




(86) Date de dépôt PCT/PCT Filing Date: 2013/06/14
(87) Date publication PCT/PCT Publication Date: 2013/12/19
(85) Entrée phase nationale/National Entry: 2014/12/09
(86) N° demande PCT/PCT Application No.: US 2013/045994
(87) N° publication PCT/PCT Publication No.: 2013/188831
(30) Priorités/Priorities: 2012/06/15 (US61/660,665);
2013/03/15 (US61/789,408)

(51) Cl.Int./Int.Cl. *C12Q 1/68* (2006.01)
(71) Demandeur/Applicant:
ADAPTIVE BIOTECHNOLOGIES CORPORATION, US
(72) Inventeur/Inventor:
ROBINS, HARLAN S., US
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : GENES DE RECEPTEUR IMMUNITAIRE ADAPTATIF REORGANISES A MARQUAGE UNIQUE DANS UN ENSEMBLE GENIQUE COMPLEXE
(54) Title: UNIQUELY TAGGED REARRANGED ADAPTIVE IMMUNE RECEPTOR GENES IN A COMPLEX GENE SET



(57) **Abrégé/Abstract:**

Compositions and methods are disclosed for uniquely tagging each rearranged gene segment that encodes a T cell receptor (TCR) and/or an immunoglobulin (Ig), in a DNA (or mRNA or cDNA reverse transcribed therefrom) sample from lymphoid cells. These and related embodiments permit accurate, high throughput quantification of distinct TCR and/or Ig encoding sequences. Also provided are compositions and methods for quantitatively sequencing the genes that encode both chains of a TCR or Ig heterodimer in a single cell, for example, to characterize the degree of T or B cell clonality in a sample.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
19 December 2013 (19.12.2013)

WIPO | PCT

(10) International Publication Number
WO 2013/188831 A1

(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:

PCT/US2013/045994

(22) International Filing Date:

14 June 2013 (14.06.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/660,665 15 June 2012 (15.06.2012) US
61/789,408 15 March 2013 (15.03.2013) US

(71) Applicant: **ADAPTIVE BIOTECHNOLOGIES CORPORATION** [US/US]; 1551 Eastlake Avenue East, Suite 200, Seattle, WA 98102 (US).

(72) Inventor: **ROBINS, Harlan, S.**; c/o Adaptive Biotechnologies Corporation, 1551 Eastlake Avenue East, Suite 200, Seattle, WA 98102 (US).

(74) Agents: **LEE, Michelle** et al.; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Mountain View, CA 94041 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))


Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: UNIQUELY TAGGED REARRANGED ADAPTIVE IMMUNE RECEPTOR GENES IN A COMPLEX GENE SET



(57) Abstract: Compositions and methods are disclosed for uniquely tagging each rearranged gene segment that encodes a T cell receptor (TCR) and/or an immunoglobulin (Ig), in a DNA (or mRNA or cDNA reverse transcribed therefrom) sample from lymphoid cells. These and related embodiments permit accurate, high throughput quantification of distinct TCR and/or Ig encoding sequences. Also provided are compositions and methods for quantitatively sequencing the genes that encode both chains of a TCR or Ig heterodimer in a single cell, for example, to characterize the degree of T or B cell clonality in a sample.

[0001] Uniquely Tagged Rearranged Adaptive Immune Receptor Genes In A Complex Gene Set

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on April 12, 2013, is named 23496PCT_CRF_sequencelisting.txt, and is 4,382,702 bytes in size.

BACKGROUND OF THE INVENTION

Technical Field

[0003] The present disclosure relates generally to quantitative high-throughput sequencing of adaptive immune receptor encoding DNA or RNA (*e.g.*, DNA or RNA encoding T cell receptors and immunoglobulins) in multiplexed nucleic acid amplification reactions. In particular, the compositions and methods described herein permit quantitative sequencing of DNA sequences encoding both chains of an adaptive immune receptor heterodimer in a single cell. Also disclosed herein are embodiments that overcome undesirable distortions in the quantification of adaptive immune receptor encoding sequences that can result from biased over-utilization and/or under-utilization of specific oligonucleotide primers in multiplexed DNA amplification.

Description of the Related Art

[0004] The adaptive immune system employs several strategies to generate a repertoire of T- and B-cell antigen receptors, *i.e.*, adaptive immune receptors, with sufficient diversity to recognize the universe of potential pathogens. The ability of T cells to recognize the universe of antigens associated with various cancers or infectious organisms is conferred by its T cell antigen receptor (TCR), which is a heterodimer of an α (alpha) chain from the TCRA locus and a β (beta) chain from the TCRB locus, or a heterodimer of a γ (gamma) chain from the TCRG locus and a δ (delta) chain from the TCRD locus. The proteins which make up these chains are encoded by DNA, which in lymphoid cells employs a unique rearrangement mechanism for generating the tremendous diversity of the TCR. This multi-subunit immune recognition receptor associates with the CD3 complex and binds to peptides presented by the major histocompatibility complex (MHC) class I and II proteins on the surface of antigen-presenting cells (APCs). Binding of TCR to the antigenic peptide on the APC is the central event in T cell activation, which occurs at an immunological synapse at the point of contact between the T cell and the APC.

[0005] Each TCR peptide contains variable complementarity determining regions (CDRs), as well as framework regions (FRs) and a constant region. The sequence diversity of $\alpha\beta$ T cells is largely determined by the amino acid sequence of the third complementarity-determining region (CDR3) loops of the α and β chain variable domains, which diversity is a result of recombination between variable (V_β), diversity (D_β), and joining (J_β) gene segments in the β chain locus, and between analogous V_α and J_α gene segments in the α chain locus, respectively. The existence of multiple such gene segments in the TCR α and β chain loci allows for a large number of distinct CDR3 sequences to be encoded. CDR3 sequence diversity is further increased by independent addition and deletion of nucleotides at the V_β - D_β , D_β - J_β , and V_α - J_α junctions during the process of TCR gene rearrangement. In this respect, immunocompetence is reflected in the diversity of TCRs.

[0006] The $\gamma\delta$ TCR is distinctive from the $\alpha\beta$ TCR in that it encodes a receptor that interacts closely with the innate immune system, and recognizes antigen in a non-HLA-dependent manner. TCR $\gamma\delta$ is expressed early in development, and has specialized anatomical distribution, unique pathogen and small-molecule specificities, and a broad spectrum of innate and adaptive cellular interactions. A biased pattern of TCR γ V and J segment expression is established early in ontogeny. Consequently, the diverse TCR γ repertoire in adult tissues is the result of extensive peripheral expansion following stimulation by environmental exposure to pathogens and toxic molecules.

[0007] Immunoglobulins (Igs or IG) expressed by B cells, also referred to herein as B cell receptors (BCR), are proteins consisting of four polypeptide chains, two heavy chains (H chains) from the IGH locus and two light chains (L chains) from either the IGK (kappa) or the IGL (lambda) locus, forming an H_2L_2 structure. Both H and L chains contain complementarity determining regions (CDR) involved in antigen recognition, and a constant domain. The H chains of Igs are initially expressed as membrane-bound isoforms using either the IgM or IgD constant region isoform, but after antigen recognition the H chain constant region can class switch to several additional isotypes, including IgG, IgE and IgA. As with TCR, the diversity of naïve Igs within an individual is mainly determined by the hypervariable complementarity determining regions (CDR). Similar to the TCR, the CDR3 domain of IGH chains is created by the combinatorial joining of the V_H , D_H , and J_H gene segments. Hypervariable domain sequence diversity is further increased by independent addition and deletion of nucleotides at the V_H - D_H , D_H - J_H , and V_H - J_H junctions during the process of Ig gene rearrangement. Distinct from TCR, Ig sequence diversity is further

augmented by somatic hypermutation (SHM) throughout the rearranged IG gene after a naïve B cell initially recognizes an antigen. The process of SHM is not restricted to CDR3, and therefore can introduce changes in the germline sequence in framework regions, CDR1 and CDR2, as well as in the somatically rearranged CDR3.

[0008] As the adaptive immune system functions in part by clonal expansion of cells expressing unique TCRs or BCRs, accurately measuring the changes in total abundance of each clone is important to understanding the dynamics of an adaptive immune response. For instance, a healthy human has a few million unique TCR β chains, each carried in hundreds to thousands of clonal T-cells out of the roughly trillion T cells in a healthy individual. Utilizing advances in high-throughput sequencing, a new field of molecular immunology has recently emerged to profile the vast TCR and BCR repertoires. Compositions and methods for the sequencing of rearranged adaptive immune receptor gene sequences and for adaptive immune receptor clonotype determination are described, for example, in Robins et al., 2009 *Blood* 114, 4099; Robins et al., 2010 *Sci. Translat. Med.* 2:47ra64; Robins et al., 2011 *J. Immunol. Meth.* doi:10.1016/j.jim.2011.09.001; Sherwood et al. 2011 *Sci. Translat. Med.* 3:90ra61; U.S.A.N. 13/217,126 (US Pub. No. 2012/0058902), U.S.A.N. 12/794,507 (US Pub. No. 2010/0330571), WO/2010/151416, WO/2011/106738 (PCT/US2011/026373), WO2012/027503 (PCT/US2011/049012), U.S.A.N. 61/550,311, and U.S.A.N. 61/569,118, all herein incorporated by reference.

[0009] To date, several different strategies have been employed to sequence nucleic acids encoding adaptive immune receptors quantitatively at high throughput, and these strategies may be distinguished, for example, by the approach that is used to amplify the CDR3-encoding regions, and by the choice of sequencing genomic DNA (gDNA) or messenger RNA (mRNA).

[0010] Sequencing mRNA is a potentially easier method than sequencing gDNA, because mRNA splicing events remove the intron between J and C segments. This allows for the amplification of adaptive immune receptors (*e.g.*, TCRs or Igs) having different V regions and J regions using a common 3' polymerase chain reaction (PCR) amplification primer in the C region. For each TCR β , for example, the thirteen J segments are all less than 60 base pairs (bp) long. Therefore, splicing events bring identical polynucleotide sequences encoding TCR β constant regions (regardless of which V and J sequences are used) to within less than 100 bp of the rearranged VDJ junction. The spliced mRNA can then be reverse transcribed into complementary DNA (cDNA) using poly-dT primers complementary to the poly-A tail

of the mRNA, random small primers (usually hexamers or nonamers) or C-segment-specific oligonucleotides. This reverse transcription should produce an unbiased library of TCR cDNA (because all cDNAs are primed with the same oligonucleotide, whether poly-dT, random hexamer, or C segment-specific oligo) that may then be sequenced to obtain information on the V and J segment used in each rearrangement, as well as the specific sequence of the CDR3. Such sequencing could use single, long reads spanning CDR3 (“long read”) technology, or could instead involve fractionating many copies of the longer sequences and using higher throughput shorter sequence reads.

[0011] Efforts to quantify the number of cells in a sample that express a particular rearranged TCR (or Ig) based on mRNA sequencing are difficult to interpret, however, because each cell potentially expresses different quantities of TCR mRNA. For example, T cells activated in vitro have 10-100 times as much mRNA per cell than quiescent T cells. To date, there is very limited information on the relative amount of TCR mRNA in T cells of different functional states, and therefore quantitation of mRNA in bulk does not necessarily accurately measure the number of cells carrying each clonal TCR.

[0012] Most T cells, on the other hand, have one productively rearranged TCR α and one productively rearranged TCR β gene (or two rearranged TCR γ and TCR δ), and most B cells have one productively rearranged Ig heavy-chain gene and one productively rearranged Ig light-chain gene (either IGK or IGL) so quantification in a sample of genomic DNA encoding TCRs or BCRs should directly correlate with, respectively, the number of T or B cells in the sample. Genomic sequencing of polynucleotides encoding any one or more of the adaptive immune receptor chains, for instance, using the human TCR β chain as a representative example, desirably entails amplifying with equal efficiency all of the many possible rearranged TCR β encoding sequences that are present in a sample containing DNA from lymphoid cells of a subject, followed by quantitative sequencing, such that a quantitative measure of the relative abundance of each clonotype can be obtained.

[0013] Difficulties are encountered with such approaches, however, in that equal amplification and sequencing efficiencies may not be achieved readily, for example, for each rearranged TCR β encoding clone, where each clone employs one of 54 possible germline V region-encoding genes and one of 13 possible J region-encoding genes. The specific sequences of the highly diverse V and J segments in the TCR β genomic locus vary widely among the large number of possible rearrangements that result from using different V or J genes, due to diversity-generating mechanisms such as those summarized above.

[0014] This sequence diversity yields complex DNA samples in which accurate determination of the multiple distinct sequences contained therein is hindered by technical limitations on the ability to quantify a plurality of molecular species simultaneously using multiplexed amplification and high throughput sequencing. In addition, it is difficult from existing methodologies to sequence quantitatively DNA or RNA encoding both chains of a TCR or IG heterodimer in a manner that permits determination that both chains originated from the same lymphoid cell.

[0015] One or more factors can give rise to artifacts that skew sequencing data outputs, compromising the ability to obtain reliable quantitative data from sequencing strategies that are based on multiplexed amplification of a highly diverse collection of TCR or IG gene templates. These artifacts often result from unequal use of diverse primers during the multiplexed amplification step. Such biased utilization of one or more oligonucleotide primers in a multiplexed reaction that uses diverse amplification templates may arise as a function of one or more of differences in the nucleotide base composition of templates and/or oligonucleotide primers, differences in template and/or primer length, the particular polymerase that is used, the amplification reaction temperatures (*e.g.*, annealing, elongation and/or denaturation temperatures), and/or other factors (*e.g.*, Kanagawa, 2003 *J. Biosci. Bioeng.* 96:317; Day et al., 1996 *Hum. Mol. Genet.* 5:2039; Ogino et al., 2002 *J. Mol. Diagnost.* 4:185; Barnard et al., 1998 *Biotechniques* 25:684; Aird et al., 2011 *Genome Biol.* 12:R18).

[0016] Clearly there remains a need for improved compositions and methods that will permit accurate quantification of adaptive immune receptor-encoding DNA and RNA sequence diversity in complex samples, in a manner that avoids skewed results such as misleading over- or underrepresentation of individual sequences due to biases in the utilization of one or more oligonucleotide primers in an oligonucleotide primer set used for multiplexed amplification of a complex template DNA population, and in a manner that permits determination of the coding sequences for both chains of a TCR or IG heterodimer that originate from the same lymphoid cell. The presently described embodiments address this need and provide other related advantages.

SUMMARY OF THE INVENTION

[0017] The invention provides compositions comprising an oligonucleotide amplification primer composition. The oligonucleotide amplification primer composition comprises (A) a first oligonucleotide amplification primer set comprising a plurality of forward

oligonucleotide sequences of a general formula (A): U1 - B1 - V1 (A), and a plurality of reverse oligonucleotide sequences of a general formula (B): U2 - B2 - J1 (B), wherein U1 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence, and U2 comprises an oligonucleotide sequence comprising a second universal adaptor oligonucleotide sequence. In one embodiment, B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, and B2 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, such that at least one of B1 or B2 is present. In another embodiment, V1 comprises an oligonucleotide sequence comprising at least 15 and not more than 100 contiguous nucleotides of a V region encoding gene sequence of a first adaptive immune receptor, or the complement thereof. In some embodiments, J1 comprises an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) a joining (J) region encoding gene sequence of said first adaptive immune receptor, or the complement thereof, or (ii) a constant (C) region encoding gene sequence of said first adaptive immune receptor, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1-B1-V1, V1 comprises a unique oligonucleotide sequence, and in each of the plurality of oligonucleotide sequences of general formula U2-B2-J1, J1 comprises a unique oligonucleotide sequence. The oligonucleotide amplification primer composition comprises a second oligonucleotide amplification primer set comprising a plurality of forward oligonucleotide sequences of a general formula (C): U3 - B3 - V2 (C) and a plurality of reverse oligonucleotide sequences of a general formula (D): U4 - B4 - J2 (D), wherein U3 comprises an oligonucleotide sequence identical to either U1 or U2, and U4 comprises an oligonucleotide sequence identical to either U1 or U2, whichever sequence is not identical to U3. In some embodiments, B3 comprises an oligonucleotide sequence comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B1, and B4 comprises an oligonucleotide sequence comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B2. In another embodiment, V2 comprises an oligonucleotide sequence comprising at least 15 and not more than 100 contiguous nucleotides of a V region encoding gene sequence of a second adaptive immune receptor, or the complement thereof. In another embodiment, J2 comprises an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) a joining (J) region encoding gene sequence of said second adaptive immune receptor, or the complement thereof, or (ii) a constant (C) region

encoding gene sequence of said second adaptive immune receptor, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3-B3-V2, V2 comprises a unique oligonucleotide sequence, and in each of the plurality of oligonucleotide sequences of general formula U4-B4-J2, J2 comprises a unique oligonucleotide sequence. In one embodiment, U1 is the same as U3. In another embodiment, U2 is the same as U4.

[0018] The invention provides a method for labeling individual rearranged DNA sequences encoding a plurality of adaptive immune receptors in a biological sample that comprises lymphoid cells of a subject, the method comprising: (a) amplifying said rearranged DNA sequences using a first amplification primer set comprising an oligonucleotide primer composition described herein under conditions that promote amplification to obtain double-stranded DNA products. Each double-stranded DNA product comprises (i) a sequence comprising at least two universal adaptor oligonucleotide sequences with one at each end of the product, at least one oligonucleotide barcode sequence, an X1 oligonucleotide sequence, an X2 oligonucleotide sequence, and (ii) a complementary sequence to the sequence in (i); (b) amplifying the double-stranded DNA products of (a) with a second amplification primer set comprising a plurality of first and second sequencing platform tag-containing oligonucleotides that each comprise either: (i) a first sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, or (ii) a second sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence. In some embodiments, amplifying takes place under conditions that promote amplification of both strands of the separated double-stranded DNA product of (a), to obtain a library of rearranged DNA sequences encoding a plurality of adaptive immune receptors for sequencing. The method also comprises a step (c) for sequencing the DNA library obtained in (b), wherein each of the sequences in the DNA library comprises a unique oligonucleotide barcode sequence, thereby labeling each sequence with an unique identifiable barcode sequence.

[0019] In some embodiments, a plurality of oligonucleotides in the second amplification primer set each further comprises either or both of: (i) a sample-identifying barcode oligonucleotide which comprises a third barcode oligonucleotide B5 comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides having a sequence that is distinct from B1 and B2, wherein in the first sequencing platform tag-containing oligonucleotide B5 is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence, and wherein in the second sequencing platform tag-containing oligonucleotide B3 is situated between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence; and (ii) a spacer oligonucleotide of any sequence of 1 to 20 contiguous nucleotides, wherein said spacer oligonucleotide is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence in the first sequencing platform tag-containing oligonucleotide, and between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence in the second sequencing platform tag-containing oligonucleotide.

[0020] In other embodiments, the invention provides an oligonucleotide primer composition, comprising a plurality of oligonucleotide sequences having a general formula (I): 5' - U1 - B1_n - X - 3' (I) wherein: U1 comprises an oligonucleotide sequence which comprises a first universal adaptor oligonucleotide sequence, B1 comprises an oligonucleotide sequence that comprises a first oligonucleotide barcode sequence of n contiguous nucleotides, wherein n is at least 6 nucleotides, and X comprises either (i) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (ii) an oligonucleotide comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences, X comprises a unique oligonucleotide sequence.

[0021] In some embodiments, the plurality of oligonucleotide sequences comprises up to 4ⁿ unique B1 oligonucleotide sequences. In one embodiment, n is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides. In other embodiments, X comprises an oligonucleotide sequence comprising at least 20, 30, 40 or 50 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof. In another embodiment, X comprises an oligonucleotide sequence comprising not more than 70,

60, or 55 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof. In yet another embodiment, X comprises an oligonucleotide sequence comprising at least 16-50 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof. In other embodiments, X comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof. In one embodiment, X is capable of hybridizing to a V region encoding gene sequence. In another embodiment, X is capable of hybridizing to a J region encoding gene sequence.

[0022] In other embodiments, B1 is a unique tag for identifying individual rearranged TCR or Ig encoding sequences. In another embodiment, U1 comprises SEQ ID NOs: 1710-1731. B1 can include sequences listed in Table 8. X can comprise SEQ ID NOs: 1631-1643 or 1696-1708. In some embodiments, X comprises SEQ ID NOs: 1644-1695. In other embodiments, X comprises SEQ ID NOs: 5613-5625. In some embodiments, the oligonucleotide composition comprising said plurality of oligonucleotide sequences comprising SEQ ID NOs: 5626-5685. In other embodiments, the oligonucleotide composition comprising said plurality of oligonucleotide sequences comprises SEQ ID NOs:1-1630.

[0023] In some embodiments, the composition includes a second plurality of oligonucleotide sequences comprising a general formula (II): 5'- P1 - S1 – B2 – U1 – 3' (II), wherein P1 comprises a sequencing platform-specific oligonucleotide, S1 comprises a sequencing platform tag-containing oligonucleotide sequence, wherein B2 comprises an oligonucleotide barcode sequence and wherein said oligonucleotide barcode sequence can be used to identify a sample source, and wherein U1 comprises said first universal adaptor oligonucleotide sequence. In other embodiments, the second plurality of oligonucleotide sequences comprises SEQ ID NOs: 5686-5877.

[0024] In another embodiment, the invention includes an oligonucleotide primer composition for a first amplification primer set comprising: (A) a plurality of first oligonucleotide sequences of a general formula (III): 5'- U1 - B1_n - X1 - 3' (III). In some embodiments, U1 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence, (ii) B1 comprises an oligonucleotide sequence comprising a first oligonucleotide barcode sequence of n contiguous nucleotides, wherein n is 0 or 6 to 20, and (iii) X1 comprises either (a) an oligonucleotide sequence comprising at least 15 and not more than 80

contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (b) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences X1 comprises a unique oligonucleotide sequence.

[0025] In one embodiment, the plurality of oligonucleotide sequences comprises up to 4^n unique B1 oligonucleotide sequences,

[0026] In another embodiment, the first amplification primer set also comprises: (B) a plurality of second oligonucleotide sequences of a general formula (IV): 5'-U2-B2_m-X2-3' (IV), wherein: (i) U2 comprises an oligonucleotide sequence comprising a second universal adaptor oligonucleotide sequence, (ii) B2 comprises an oligonucleotide sequence comprising a second oligonucleotide barcode sequence of m contiguous nucleotides, wherein m is 0 or 6 to 20, (iii) X2 comprises (a) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (b) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences X1 comprises a unique oligonucleotide sequence, wherein n and m are independent of each other, and in said first and second pluralities of oligonucleotides, m and n are not both zero, and wherein if X1 comprises an oligonucleotide sequence comprising an adaptive immune receptor V region encoding gene sequence, then X2 comprises an oligonucleotide sequence comprising an adaptive immune receptor J region encoding gene sequence, and if X1 comprises an oligonucleotide sequence comprising an adaptive immune receptor J region encoding gene sequence, then X2 comprises an oligonucleotide sequence comprising an adaptive immune receptor V region encoding gene sequence.

[0027] In one embodiment, the plurality of oligonucleotide sequences comprises up to 4^m unique B2 oligonucleotide sequences.

[0028] In another embodiment, X1 or X2 comprises an oligonucleotide sequence comprising at least 20, 30, 40 or 50 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof. In yet another embodiment, X1 or X2 comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said

complement thereof. In other embodiments, X1 or X2 comprises an oligonucleotide sequence comprising at least 16-50 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof. In one embodiment, X1 or X2 comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof. In another embodiment, B1 is a unique tag for identifying an individual rearranged TCR or Ig encoding sequence. In yet another embodiment, B2 is a unique tag for identifying an individual rearranged TCR or Ig encoding sequence.

[0029] In some embodiments, U1 or U2 comprises SEQ ID NOs: 1710-1731. In one embodiment, B1 or B2 comprises sequences listed in Table 8. In another embodiment, X1 or X2 comprises SEQ ID NOs: 1631-1643 or 1696-1708. In yet another embodiment, X1 or X2 comprises SEQ ID NOs: 1644-1695. X1 or X2 can comprise SEQ ID NOs: 5613-5625. In other embodiments, the plurality of first or second oligonucleotide sequences comprises SEQ ID NOs: 5626-5685. In another embodiment, the plurality of first or second oligonucleotide sequences comprise SEQ ID NOs: 1-1630.

[0030] In another embodiment, the invention comprises an oligonucleotide amplification primer composition, comprising: (A) a first oligonucleotide amplification primer set comprising a plurality of oligonucleotide sequences of a general formula (V): U1/2 - B1 - X1 (V), wherein U1/2 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence when B1 is present, or a second universal adaptor oligonucleotide sequence when B1 is nothing, and wherein B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, and wherein X1 comprises either: (1) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor V region encoding gene sequence, or the complement thereof, or (2) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence.

[0031] In some embodiments, the oligonucleotide amplification primer composition also comprises: (B) a second oligonucleotide amplification primer set comprising a plurality of oligonucleotide sequences of a general formula (VI): U3/4 - B2 - X2 (VI), wherein U3/4

comprises an oligonucleotide sequence comprising a third universal adaptor oligonucleotide sequence when B2 is present or a fourth universal adaptor oligonucleotide sequence when B2 is nothing, and wherein B2 comprises an oligonucleotide sequence comprising either nothing or a second oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B1, and wherein X2 comprises either (1) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor V region encoding gene sequence, or the complement thereof, or (2) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence. In some embodiments, U3 has the same sequence as U1 or U2. In other embodiments, U4 has the same sequence as U1 or U2.

[0032] Certain embodiments of the invention include a method for identifying individual rearranged DNA sequences encoding a plurality of adaptive immune receptors in a biological sample that comprises lymphoid cells of a subject, the method comprising: (a) amplifying said rearranged DNA sequences using a first amplification primer set comprising an oligonucleotide primer composition described herein under conditions that promote amplification to obtain double-stranded DNA products that each comprise (i) a sequence comprising at least one universal adaptor oligonucleotide sequence, at least one oligonucleotide barcode sequence, and at least one of an X, X1 or X2 oligonucleotide sequence, and (ii) a complementary sequence to the sequence in (i).

[0033] The method includes the step of (b) amplifying the double-stranded DNA products of (a) with a second amplification primer set comprising a plurality of first and second sequencing platform tag-containing oligonucleotides that each comprise either: (i) a first sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, or (ii) a second sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence, wherein amplifying takes place under conditions that promote amplification of both strands of the separated

double-stranded DNA product of (a), to obtain a library of rearranged DNA sequences encoding a plurality of adaptive immune receptors for sequencing.

[0034] The method includes the step of (c) sequencing the DNA library obtained in (b), wherein each of the sequences in the DNA library comprises a unique oligonucleotide barcode sequence, thereby labeling each sequence with a unique identifiable barcode sequence. In some embodiments, a plurality of oligonucleotides in the second amplification primer set each further comprises either or both of: (i) a sample-identifying barcode oligonucleotide which comprises a third barcode oligonucleotide B3 comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides having a sequence that is distinct from B1 and B2, wherein in the first sequencing platform tag-containing oligonucleotide B3 is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence, and wherein in the second sequencing platform tag-containing oligonucleotide B3 is situated between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence, and (ii) a spacer oligonucleotide of any sequence of 1 to 20 contiguous nucleotides, wherein said spacer oligonucleotide is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence in the first sequencing platform tag-containing oligonucleotide, and between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence in the second sequencing platform tag-containing oligonucleotide.

[0035] In some embodiments, the invention includes a method for labeling individual rearranged DNA sequences or mRNA sequences transcribed therefrom encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising: contacting (A) a first plurality of individual microdroplets that each contain a single lymphoid cell or genomic DNA isolated therefrom or complementary DNA (cDNA) that has been reverse transcribed from messenger RNA (mRNA) of a single lymphoid cell, with (B) a second plurality of individual microdroplets. The second plurality of individual microdroplets each contain: (i) a first oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a first polypeptide of an adaptive immune receptor heterodimer, and (ii) a second oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a second polypeptide of the adaptive immune receptor heterodimer. In some embodiments, the first oligonucleotide amplification primer set comprises a composition of U1/2-B1-X1 described

herein, and the second oligonucleotide amplification primer set comprises a composition of U3/4-B2-X2 described herein.

[0036] The method also includes providing conditions for a time sufficient such that a plurality of fusion events occur between one of said first microdroplets and one of said second microdroplets to produce a plurality of fused microdroplets, and providing conditions that permit amplification of the genomic DNA, or the cDNA that has been reverse transcribed from mRNA, using the first and second oligonucleotide amplification primer sets within the plurality of fused microdroplets. In some embodiments, each of one or more of said plurality of fused microdroplets comprises: a first double-stranded DNA product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and a second double-stranded DNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, thereby upon amplification of the genomic DNA, or the cDNA that has been reverse transcribed from mRNA, labeling each of the individual rearranged DNA sequences or mRNA sequences transcribed therefrom with an oligonucleotide barcode sequence.

[0037] In some embodiments, the method comprises disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of said first and second double-stranded DNA products. The method also includes contacting the mixture of the first and second double-stranded DNA products with a third amplification primer set and a fourth amplification primer set, wherein the third amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each

comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence. In some embodiments, the fourth amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence. In one embodiment, the step of contacting takes place under conditions and for a time sufficient to amplify both strands of the first and second double-stranded DNA products, to obtain a DNA library for sequencing. [0038] In another embodiment, the method includes sequencing the DNA library to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer. In some embodiments, the third and fourth amplification primer sets are the same.

[0039] In one embodiment, the invention comprises a method for labeling individual rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising: contacting (A) a first plurality of individual microdroplets that each contain complementary DNA (cDNA) that has been reverse transcribed from messenger RNA (mRNA) of a single lymphoid cell, with (B) a second plurality of individual microdroplets. The second plurality of individual microdroplets each contain (i) a first oligonucleotide amplification primer set that is capable of amplifying a first cDNA sequence encoding a first polypeptide of an adaptive immune receptor heterodimer, and (ii) a second oligonucleotide amplification primer set that is capable of amplifying a second cDNA sequence encoding a second polypeptide of the adaptive immune receptor heterodimer. The first oligonucleotide amplification primer set comprises a composition of U1/2-B1-X1 described herein, and the second oligonucleotide amplification primer set comprises a composition of U3/4-B2-X2 described herein.

[0040] In other embodiments, the method includes providing conditions for a time sufficient for a plurality of fusion events between one of said first microdroplets and one of said second microdroplets to produce a plurality of fused microdroplets and conditions that permit amplification of the cDNA that has been reverse transcribed from mRNA of a single lymphoid cell, using the first and second oligonucleotide amplification primer sets within the plurality of fused microdroplets. In some embodiments, each of one or more of said plurality of fused microdroplets comprises: a first double-stranded DNA product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and a second double-stranded DNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, thereby upon amplification of the cDNA, uniquely labeling each of the individual rearranged cDNA sequences with a unique oligonucleotide barcode sequence.

[0041] In another embodiment, the method includes disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of said first and second double-stranded DNA products. In other embodiments, the method includes contacting the mixture of first and second double-stranded DNA products with a third amplification primer set and a fourth amplification primer set. In one embodiment, the third amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor

oligonucleotide sequence. In one embodiment, the fourth amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence. In some embodiments, the step of contacting takes place under conditions and for a time sufficient to amplify both strands of the first and second double-stranded DNA products, to obtain a DNA library for sequencing.

[0042] In certain embodiments, the method includes sequencing the DNA library to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer. In another embodiment, the third amplification primer set is identical to the fourth amplification primer set.

[0043] In certain other embodiments, the method includes either or both of: (1) the first oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the first polypeptide, a rearranged DNA sequence encoding a first complementarity determining region-3 (CDR3) of the first polypeptide; and (2) the second oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the second polypeptide, a rearranged DNA sequence encoding a second complementarity determining region-3 (CDR3) of the second polypeptide.

[0044] In some embodiments, the first polypeptide of the adaptive immune receptor heterodimer is a TCR alpha (TCRA) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR beta (TCRB) chain. In other embodiments, the first polypeptide of the adaptive immune receptor heterodimer is a TCR gamma (TCRG) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR delta (TCRD) chain.

[0045] In another embodiment, the first polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin heavy (IGH) chain and the second polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin light (IGL or IGK or both IGL and IGK) chain. In some embodiments, if the first polypeptide of the adaptive immune

receptor heterodimer is an IGH chain and the second polypeptide of the adaptive immune receptor heterodimer is both IGL and IGK, then three different amplification primer sets are used comprising: a first oligonucleotide amplification primer set for IGH, a second oligonucleotide amplification primer set for IGK, and a third oligonucleotide amplification primer set for IGL.

[0046] In yet another embodiment, each of the second plurality of individual microdroplets further contains a third oligonucleotide primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule and that comprises a composition comprising a plurality of oligonucleotide sequences having a general formula (VII): U5/6 - B - X3 (VII). In one aspect, U5/6 comprises a fifth universal adaptor oligonucleotide sequence when B is present or a sixth universal adaptor oligonucleotide sequence when B is nothing. In another aspect, B comprises B1 or B2. In yet another aspect, X3 comprises an oligonucleotide that is one of (i) a forward primer of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and (ii) a reverse primer of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U5/6-B-X3, X3 comprises a unique oligonucleotide sequence.

[0047] In one embodiment, the lymphocyte status indicator molecule comprises one or more of FoxP3, CD4, CD8, CD11a, CD18, CD21, CD25, CD29, CCD30, CD38, CD44, CD45, CD45RA, CD45RO, CD49d, CD62, CD62L, CD69, CD71, CD103, CD137 (4-1BB), CD138, CD161, CD294, CCR5, CXCR4, IgG1-4 H-chain constant region, IgA H-chain constant region, IgE H-chain constant region, IgD H-chain constant region, IgM H-chain constant region, HLA-DR, IL-2, IL-5, IL-6, IL-9, IL-10, IL-12, IL-13, IL-15, IL-21, TGF- β , TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9 and TLR10.

[0048] In some embodiments, the method includes sorting the data set of sequences according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode, sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset, and clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets. The method

further includes identifying as originating from the same cell sequences that are members of an X1 and an X2 sequence cluster set that belong to the same one or more barcode sequence sets.

[0049] In another embodiment, methods of the invention include determining rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising: (1) distributing cells of a cell suspension that comprises a population of lymphoid cells of a subject, amongst a plurality of containers that are capable of containing said cells, to obtain a plurality of containers that each contain a subpopulation of the lymphoid cells that comprises one lymphoid cell or a plurality of lymphoid cells. The method also includes (2) contacting each of said plurality of containers, under conditions and for a time sufficient to promote reverse transcription of messenger RNA (mRNA) in the lymphoid cells in the plurality of containers, with a first and a second oligonucleotide reverse transcription primer set, wherein (A) the first oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of first mRNA sequences encoding a plurality of polypeptides of a first adaptive immune receptor heterodimer, and (B) the second oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of second mRNA sequences encoding a plurality of polypeptides of a second adaptive immune receptor heterodimer.

[0050] In another embodiment, the method comprises (I) the first oligonucleotide reverse transcription primer set comprising a composition of a general formula of U1/2-B1-X1 described herein, and (II) the second oligonucleotide reverse transcription primer set comprises a composition comprising a general formula U3/4-B2-X2 described herein.

[0051] In yet another embodiment, the step of contacting takes place under conditions and for a time sufficient to obtain in each of one or more of said plurality of containers: a first reverse-transcribed complementary DNA (cDNA) product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and a second reverse-transcribed cDNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second

polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer.

[0052] In one embodiment, the method includes combining the first and second reverse-transcribed cDNA products from the plurality of containers to obtain a mixture of reverse-transcribed cDNA products and contacting the mixture of first and second reverse-transcribed cDNA products of (3) with a first oligonucleotide amplification primer set and a second oligonucleotide amplification primer set. In some embodiments, the first amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence.

[0053] In another embodiment, the second oligonucleotide amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence.

[0054] In another embodiment, the step of contacting takes place under conditions and for a time sufficient to amplify both of the first and second reverse-transcribed cDNA products of (2), to obtain a DNA library for sequencing. In one embodiment, the method includes sequencing the DNA library obtained in (3) to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

[0055] In yet another embodiment, the method includes (a) sorting the data set of sequences according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode and (b) sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset. The method can further include (c) clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets.

[0056] In another embodiment, the method includes (d) identifying each first and second adaptive immune receptor heterodimer polypeptide encoding sequence based on known X1 and X2 sequences, wherein each X1 sequence and each X2 sequence is associated with one or a plurality of unique B sequences to identify the container from which each B sequence-associated X1 sequence and each B sequence-associated X2 sequence originated. In some embodiments, the method includes (e) combinatorically matching B sequence-associated X1 and X2 sequences of (d) as being of common clonal origin based on a probability of B sequences that are coincident with common first and second adaptive immune receptor heterodimer polypeptide encoding sequences, and therefrom determining that rearranged DNA sequences encoding first and second polypeptide sequences of the adaptive immune receptor heterodimer originated in a single lymphoid cell.

[0057] In one embodiment, the first oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the first polypeptide, a rearranged DNA sequence encoding a first complementarity determining region-3 (CDR3) of the first polypeptide. In another embodiment, the second oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the second polypeptide, a rearranged DNA sequence encoding a second complementarity determining region-3 (CDR3) of the second polypeptide.

[0058] In certain embodiments, the first polypeptide of the adaptive immune receptor heterodimer is a TCR alpha (TCRA) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR beta (TCRB) chain, or (b) the first polypeptide of the adaptive immune receptor heterodimer is a TCR gamma (TCRG) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR delta (TCRD) chain, or (c) the first polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin

heavy (IGH) chain and the second polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin light (IGL, IGK, or both IGL and IGK) chain.

[0059] In certain other embodiments, one or more of the containers comprises a third oligonucleotide amplification primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule and that comprises a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula (VI): U5/6 - B3 - X3 (VI). In some embodiments, U5/6 comprises an oligonucleotide which comprises a fifth universal adaptor oligonucleotide sequence when B3 is present or a sixth universal adaptor oligonucleotide sequence when B3 is nothing. In one embodiment, B3 comprises an oligonucleotide that comprises either nothing or a third oligonucleotide barcode sequence of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides that is either the same as or different than at least one of B1 or B2. In another embodiment, X3 comprises an oligonucleotide that is one of (i) a forward primer polynucleotide of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and (ii) a reverse primer polynucleotide of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U5/6-B3-X3, X3 comprises a unique oligonucleotide sequence.

[0060] In certain embodiments, the lymphocyte status indicator molecule comprises one or more of FoxP3, CD4, CD8, CD11a, CD18, CD21, CD25, CD29, CCD30, CD38, CD44, CD45, CD45RA, CD45RO, CD49d, CD62, CD62L, CD69, CD71, CD103, CD137 (4-1BB), CD138, CD161, CD294, CCR5, CXCR4, IgG1-4 H-chain constant region, IgA H-chain constant region, IgE H-chain constant region, IgD H-chain constant region, IgM H-chain constant region, HLA-DR, IL-2, IL-5, IL-6, IL-9, IL-10, IL-12, IL-13, IL-15, IL-21, TGF- β , TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9 and TLR10.

[0061] These and other aspects of the herein described invention embodiments will be evident upon reference to the following detailed description and attached drawings. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet are incorporated herein by reference in their entirety, as if each was incorporated individually. Aspects and embodiments of the invention can be modified, if necessary, to employ concepts of the various patents, applications and publications to provide yet further embodiments.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0062] Figure 1 depicts a schematic representation of certain herein described compositions and methods. U1 and U2 represent universal adaptor oligonucleotides. BC1 and BC2 represent barcode oligonucleotides. J represents an adaptive immune receptor joining (J) region gene and Jpr represents a region of such a gene to which a J-specific oligonucleotide primer specifically anneals. V represents an adaptive immune receptor variable (V) region gene and Vpr represents a region of such a gene to which a V-specific oligonucleotide primer specifically anneals. NDN represents the diversity (D) region found in some adaptive immune receptor encoding genes, flanked on either side by junctional nucleotides (N) which may include non-templated nucleotides. Adap1 and Adap2 represent sequencing platform-specific adapters. The segment shown as “n6” represents a spacer nucleotide segment of any nucleotide sequence, in this case, a spacer of six randomly selected nucleotides.

[0063] Figure 2 depicts a schematic representation of certain herein described compositions and methods in which individual first and second microdroplets are contacted to permit fusion events between single first and second microdroplets, by which fusion events DNA from individual lymphoid cells (*e.g.*, T or B cells) is introduced, within a fused microdroplet, to first and second oligonucleotide amplification primer sets that are capable of amplifying, respectively, DNA encoding sequences (*e.g.*, CDR3 encoding DNA) of first and second adaptive immune receptor polypeptide encoding genes from the same cell. Amplification and oligonucleotide barcode labeling of at least two rearranged DNA loci from the same cell are thus contemplated as described herein, *e.g.*, [IGH + IGL], [IGH + IGK], [IGH + IGK + IGL], [TCRA + TCRB], [TCRG + TCRG], etc.

[0064] Figure 3 depicts a schematic representation of certain herein described compositions and methods according to which, for example, DNA from individual lymphoid cells (*e.g.*, T or B cells), or cDNA that has been reverse transcribed from mRNA of single lymphoid cells, is introduced, within a fused microdroplet, to first and second oligonucleotide amplification primer sets that are capable of amplifying, respectively, DNA encoding sequences (*e.g.*, CDR3 encoding DNA) of first and second adaptive immune receptor polypeptide encoding genes from the same cell, after which the individual microdroplets are disrupted (*e.g.*, by chemical, physical and/or mechanical dissolution, dissociation, breakage, etc.) and the released bar-coded double-stranded DNAs are amplified with universal oligonucleotide primers and sequencing platform-specific adapters to permit large-scale multiplexed quantitative sequencing. See Brief Description of Fig. 1 for abbreviations.

[0065] Figure 4 depicts a schematic representation of labeling adaptive immune receptor polypeptide encoding cDNA during reverse transcription by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences into cDNA.

[0066] Figure 5 depicts a schematic representation of labeling adaptive immune receptor polypeptide encoding cDNA during reverse transcription by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences into cDNA.

[0067] Figure 6 presents a schematic representation of a DNA product that is amenable to sequencing following modification with Illumina sequencing adapters of amplified adaptive immune receptor polypeptide encoding cDNA that has been labeled during reverse transcription by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences.

DETAILED DESCRIPTION OF THE INVENTION

[0068] The present invention provides, in certain embodiments and as described herein, compositions and methods that are useful for reliably quantifying and determining the sequences of large and structurally diverse populations of rearranged genes encoding adaptive immune receptors, such as immunoglobulins (IG) and/or T cell receptors (TCR). These rearranged genes may be present in a biological sample containing DNA from lymphoid cells of a subject or biological source, including a human subject, and/or mRNA transcripts of these rearranged genes may be present in such a sample and used as templates for cDNA synthesis by reverse transcription.

[0069] Disclosed herein are unexpectedly advantageous approaches for uniquely and unambiguously labeling individual, sequence-distinct IG and TCR encoding gene segments or mRNA transcripts thereof, or cDNA that has been reverse transcribed from such mRNA transcripts, by performing such labeling prior to conventional steps of expanding a population of such gene segments or transcripts thereof (including reverse transcripts) through established nucleic acid amplification techniques. Without wishing to be bound by theory, by labeling individual TCR and IG encoding gene segments or transcripts thereof (including complementary DNA generated by reverse transcription) as described herein, prior to commonly practiced amplification steps which are employed to generate DNA copies in sufficient quantities for sequencing, the present embodiments offer unprecedented sensitivity in the detection and quantification of diverse TCR and IG encoding sequences, while at the

same time avoiding misleading, inaccurate or incomplete results that may occur due to biases in oligonucleotide primer utilization during multiple rounds of nucleic acid amplification from an original sample, using a sequence-diverse set of amplification primers.

[0070] Also described herein in certain embodiments are unprecedented compositions and methods that permit quantitative determination of the sequences encoding both polypeptides in an adaptive immune receptor heterodimer from a single cell, such as both TCRA and TCRB from a T cell, or both IgH and IgL from a B cell. By providing the ability to obtain such information from a complex sample such as a sample containing a heterogeneous mixture of T and/or B cells from a subject, these and related embodiments permit more accurate determination of the relative representation in a sample of particular T and/or B cell clonal populations than has previously been possible.

[0071] Certain embodiments contemplate modifications as described herein to oligonucleotide primer sets that are used in multiplexed nucleic acid amplification reactions to generate a population of amplified rearranged DNA molecules from a biological sample containing rearranged genes encoding adaptive immune receptors, prior to quantitative high throughput sequencing of such amplified products. Multiplexed amplification and high throughput sequencing of rearranged TCR and BCR encoding DNA sequences are described, for example, in Robins et al., 2009 *Blood* 114:4099; Robins et al., 2010 *Sci. Translat. Med.* 2:47ra64; Robins et al., 2011 *J. Immunol. Meth.* doi:10.1016/j.jim.2011.09.001; Sherwood et al. 2011 *Sci. Translat. Med.* 3:90ra61; U.S.A.N. 13/217,126 (US Pub. No. 2012/0058902), U.S.A.N. 12/794,507 (US Pub. No. 2010/0330571), WO/2010/151416, WO/2011/106738 (PCT/US2011/026373), WO2012/027503 (PCT/US2011/049012), U.S.A.N. 61/550,311, and U.S.A.N. 61/569,118; accordingly these disclosures are incorporated by reference and may be adapted for use according to the embodiments described herein.

[0072] According to certain embodiments, in a sample containing a plurality of sequence-diverse TCR or IG encoding gene segments, such as a sample comprising DNA (or mRNA transcribed therefrom or cDNA reverse-transcribed from such mRNA) from lymphoid cells in which DNA rearrangements have taken place to encode functional TCR and/or IG heterodimers (or in which non-functional TCR or IG pseudogenes have been involved in DNA rearrangements), a plurality of individual TCR or IG encoding sequences may each be uniquely tagged with a specific oligonucleotide barcode sequence as described herein, through a single round of nucleic acid amplification (*e.g.*, polymerase chain reaction PCR). The population of tagged polynucleotides can then be amplified to obtain a library of tagged

molecules, which can then be quantitatively sequenced by existing procedures such as those described, for example, in U.S.A.N. 13/217,126 (US Pub. No. 2012/0058902), U.S.A.N. 12/794,507 (US Pub. No. 2010/0330571), WO/2010/151416, WO/2011/106738 (PCT/US2011/026373), WO2012/027503 (PCT/US2011/049012), U.S.A.N. 61/550,311, and U.S.A.N. 61/569,118.

[0073] In the course of these sequence reads, the incorporated barcode tag sequence is sequenced and can be used as an identifier in the course of compiling and analyzing the sequence data so obtained. In certain embodiments, it is contemplated that for each barcode tag sequence, a consensus sequence for the associated TCR or IG sequences may be determined. A clustering algorithm can then be applied to identify molecules generated from the same original clonal cell population. By such an approach, sequence data of high quality can be obtained in a manner that overcomes inaccuracies associated with sequencing artifacts.

[0074] An exemplary embodiment is depicted in Figure 1, according to which from a starting template population of genomic DNA or cDNA from a lymphoid cell-containing population, two or more cycles of PCR are performed using an oligonucleotide primer composition that contains primers having the general formula U1-B1_n-X as described herein. As shown in Figure (Fig.) 1, the J-specific primer 110a contains a J primer sequence 100 that is complementary to a portion of the J segment, a barcode tag (BC1) 101 in Fig. 1, or B1_n in the generic formula) and also includes a first external universal adaptor sequence (U1) 102, while the V-specific primer 110b includes a V primer sequence 103 that is complementary to a portion of the V segment and a second external universal adaptor sequence (U2) 104. The invention need not be so limited, however, and also contemplates related embodiments, such as those where the barcode may instead or may in addition be present as part of the V-specific primer and is situated between the V-sequence and the second universal adaptor. It will be appreciated that based on the present disclosure, those skilled in the art can design other suitable primers by which to introduce the herein described barcode tags to uniquely label individual TCR and/or IG encoding gene segments.

[0075] As described herein, a large number (up to 4ⁿ, where n is the length of the barcode sequence) of different barcode sequences are present in the oligonucleotide primer composition that contains primers having the general formula U1-B1_n-X as described herein, such that the PCR products of the large number of different amplification events following specific annealing of appropriate V- and J-specific primers are differentially labeled. In some

embodiments, the number of barcode sequences is up to or smaller than 4^n . In one embodiment, a set of 192 different barcode sequences are used based on a barcode of length $n=8$. The length of the barcode “n” determines the possible number of barcodes (4^n as described herein), but in some embodiments, a smaller subset is used to avoid closely related barcodes or barcodes with different annealing temperatures. In other embodiments, as described herein, sets of m and n barcode sequences are used in subsequent amplification steps (e.g., to individually label each rearranged TCR or IG sequence and then to uniformly label (“tailing”) a set of sequences obtained from the same source, or sample. In preferred embodiments, the V and J primers 100 and 103 are capable of promoting the amplification of a TCR or Ig encoding sequence that includes the CDR3 encoding sequence, which in Fig. 1 includes the NDN region 111. As also indicated in Fig. 1, following no more than two amplification cycles, the first amplification primer set 110a, 110b is separated from the double-stranded DNA product. By such a step, it is believed according to non-limiting theory that contamination of the product preparation by subsequent rounds of amplification is avoided, where contaminants could otherwise be produced by amplifying newly formed double-stranded DNA molecules with amplification primers that are present in the complex reaction but which are primers other than those used to generate the double-stranded DNA in the first one or two amplification cycles. A variety of chemical and biochemical techniques are known in the art for separating double-stranded DNA from oligonucleotide amplification primers.

[0076] Once the first amplification primer set 110a, 110b is removed, by which the unique barcode tag sequences have been introduced, the tagged double-stranded DNA (dsDNA) products can be amplified using a second amplification primer set 120a, 120b as described herein and depicted in Fig. 1, to obtain a DNA library suitable for sequencing. The second amplification primer set advantageously exploits the introduction, during the preceding step, of the universal adaptor sequences 102, 104 (e.g., U1 and U2 in Fig. 1) into the dsDNA products. Accordingly, because these universal adaptor sequences have been situated external to the unique barcode tags (BC1) 101 in Fig. 1, the amplification products that comprise the DNA library to be sequenced retain the unique barcode identifier sequences linked to each particular rearranged V-J gene segment combination, whilst being amenable to amplification via the universal adaptors. An exemplary set of such a second primer set, also known as “tailing” primers, is shown in Table 7.

[0077] In preferred embodiments and as also depicted in Fig. 1, the second amplification primer set 120a, 120b may introduce sequencing platform-specific oligonucleotide sequences (Adap1 105 and Adap2 106 in Fig. 1), however these are not necessary in certain other related embodiments. The second amplification primer set 120a, 120b may also optionally introduce a second oligonucleotide barcode identifier tag (BC2 107 in Fig. 1), such as a single barcode sequence that may desirably identify all products of the amplification from a particular sample (*e.g.*, as a source subject-identifying code) and ease multiplexing multiple samples to allow for higher throughput. The barcode (BC2; 107 in Fig. 1) is a modification that increases the throughput of the assay (*e.g.*, allows samples to be multiplexed on the sequencer), but is not required. Alternatively, a universal primer without adaptors can be used to amplify the tagged molecules. After amplification, the molecules can be additionally tagged with platform specific oligonucleotide sequences. Such inclusion of a second, sample-identifying barcode, may beneficially aid in the identification of sample origins when samples from several different subjects are mixed, or in the identification of inadvertent contamination of one sample preparation with material from another sample preparation. The second amplification primer set may also, as shown in Fig. 1, optionally include a spacer nucleotide (“n6”; 108 in Fig. 1), which may facilitate the operation of the sequencing platform-specific sequences. The spacer improves the quality of the sequencing data, but is not required or present in certain embodiments. The spacer is specifically added to increase the number of random base pairs during the first 12 cycles of the sequencing step of the method. By increasing the diversity of the first 12 cycles, cluster definition and basecalling is improved. The spacer nucleotide 108 may be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11-20, 21-30 or more nucleotides of any sequence, typically a randomly generated sequence. Where it may be of concern that the presence of such random sequences will result in uneven annealing rates amongst the oligonucleotide primers containing such sequences, it may be preferred to perform a relatively small number of amplification cycles, typically three, four or five cycles, or optionally 1-6 or no more than eight cycles, to reduce the potential for unevenness in amplification that could skew downstream results.

[0078] The resulting DNA library can then be sequenced according to standard methodologies and using available instrumentation as provided herein and known in the art. Where a second, sample-identifying barcode (BC2 107 in Fig. 1) is present, sequencing that includes reading both such barcodes is performed, with the sequence information (V-J junction including CDR3 encoding sequence, along with the first oligonucleotide barcode

BC1 101 that uniquely tags each distinct sequence) between the two occurrences of the sample-identifying barcode 107 also being read. Sequencing primers may include, for instance, and with reference to Fig. 1, the universal primer 102 on the J side of NDN 111 for the first read, followed by a barcode sequence BC1 101, a J primer sequence 100 and CDR3 sequences. The second set of amplification primers include a forward primer comprising the platform-specific primer (Adap1 105) on the J side, a spacer sequence comprising random nucleotides (labeled “n6”; 108 in Fig. 1), and BC2 sample-identifying barcodes 107. The reverse primer in the second set of amplification primers includes the universal primer 104 on the V side of NDN 111, a spacer sequence 108 comprising random nucleotides, and a BC2 sample-identifying barcode sequence 107, and optionally a paired-end read using the reverse second sequencing platform-specific primer (Adap2 106). The second sequencing platform-specific primer (Adap2 106) is used to sequence and “read” the spacer sequence 108, the sample-identifying barcode sequence BC2 107, the universal adaptor sequence 104, the V sequence 103, and NDN 111. To capture the CDR3 sequence, one can use J amplification primers, C amplification primers or the V amplification primers.

[0079] Sequence data may be sorted using the BC2 sample-identifying barcodes 107 and then further sorted according to sequences that contain a common first barcode BC1 101. Within such sorted sequences, CDR3 sequences may be clustered to determine whether more than one sequence cluster is present using any of a known variety of algorithms for clustering (e.g., BLASTClust, UCLUST, CD-HIT, or others, or as described in Robins et al., 2009 *Blood* 114:4099). Additionally or alternatively, sequence data may be sorted and selected on the basis of those sequences that are found at least twice. Consensus sequences may then be determined by sequence comparisons, for example, to correct for sequencing errors. Where multiple unique identifier barcode tags (BC1 101) are detected among sequences that otherwise share a common consensus sequence, the number of such barcode tags that is identified may be regarded as reflective of the number of molecules in the sample from the same T cell or B cell clone.

Identifying Both Chains Of A TCR Or IG Heterodimer From A Single Adaptive Immune Cell

[0080] As also noted above, in certain other embodiments there is provided herein a method for determining rearranged DNA sequences (or mRNA sequences transcribed therefrom or cDNA that has been reverse transcribed from such mRNA) encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell.

The method includes uniquely labeling each rearranged DNA sequence with a unique barcode sequence for identifying a particular cell and/or sample.

[0081] Briefly, and by way of illustration and not limitation, these and related embodiments comprise a method comprising steps of (1) in each of a plurality of parallel reactions, contacting first and second microdroplets and permitting them to fuse under conditions permissive for nucleic acid amplification, to generate double-stranded DNA products (or single-stranded cDNA products) that all contain an identical barcode oligonucleotide sequence and that correspond to the two chains of an adaptive immune receptor heterodimer; (2) disrupting the fused microdroplets to obtain a heterogeneous mixture of double-stranded (or single-stranded) DNA products; (3) amplifying the heterogeneous mixture of double-stranded DNA (or single-stranded) products to obtain a DNA library for sequencing; and (4) sequencing the library to obtain a data set of DNA sequences encoding the first and second polypeptides of the heterodimer.

[0082] The method comprises contacting and permitting to fuse in pairwise fashion (A) individual first microdroplets that each (or in every n^{th} droplet) contain a single lymphoid cell or genomic DNA isolated therefrom, or cDNA has been reverse transcribed from mRNA, with (B) individual second microdroplets from a plurality of second liquid microdroplets that each contain two oligonucleotide amplification primer sets, the first set for amplifying any rearranged DNA that encodes the first chain of an adaptive immune receptor heterodimer (*e.g.*, an IGH chain, or a TCRA chain), and the second set for amplifying any rearranged DNA that encodes the second chain of the heterodimer (*e.g.*, an IGL chain, or a TCRB chain). Significantly, in a given second microdroplet, all oligonucleotide amplification primers will comprise the same barcode oligonucleotide, but within different second microdroplets, the primer sets will comprise different barcode sequences. The step of contacting is controlled so that in each of a plurality of events, a single first microdroplet fuses with a single second microdroplet to obtain a fused microdroplet. The contents of each of the first and second microdroplets come into contact with one another in the fused microdroplet. Oligonucleotide amplification primer sets capable of amplifying any rearranged DNA encoding a given TCR or IG polypeptide are described elsewhere herein and in the references incorporated for such disclosure.

[0083] Those familiar with the art will be aware of any of a number of microfluidics apparatus and devices by which microdroplet compositions that have defined contents and properties (such as the ability to controllably undergo fusion) may be prepared, such as the

RainDance™ microdroplet digital PCR system (RainDance Technologies, Lexington, MA) or any of the systems described, for example, in Pekin et al., 2011 *Lab Chip* 11:2156; Miller et al., 2012 *Proc. Nat. Acad. Sci. USA* 109:378; Brouzes et al., 2009 *Proc. Nat. Acad. Sci. USA* 106:14195; Joensson et al., 2009 *Angew. Chem. Int. Ed.* 81:4813; Baret et al., 2009 *Lab Chip* 9:1850; Frenz et al., 2009 *Lab Chip* 9:1344; Kiss et al., 2008 *Anal. Chem.* 80:8975; Leamon et al., 2006 *Nat. Meths.* 3:541; which may be adapted to a particular method such as those described herein through modifications that are routine in view of the present disclosure.

[0084] As a non-limiting example, certain embodiments may exploit the properties of aqueous phase microdroplets dispersed in an oil phase using microfluidic channels. Microdroplets may be water-in-oil emulsions, oil-in-water emulsions, or similar aqueous and non-aqueous emulsion compositions. Microdroplets may also be called microdroplets or micellar microdroplets. Conventional water-in-oil (WO) emulsions have found many applications in biology, including next-generation sequencing (Margulies et al., *Nature* 2005, 437, 376-380), rare mutation detection (Diehl, F. et al. *Proc. Natl. Acad. Sci. U.S.A.* 2005, 102, 16368-16373; Li, M. et al., *Nat. Methods* 2006, 3, 95-97; Diehl, F. et al., *Nat. Med.* 2008, 14, 985-990) and quantitative detection of DNA methylation (Li, M. et al., *Nat. Biotechnol.* 2009, 27, 858-U118), but these emulsions suffer from droplet polydispersity and shearing stresses which can disrupt cells during mechanical agitation used to form the emulsions. The use of microfluidics overcomes these limitations and leads to an improved performance of biochemical and cell based assays (Zeng, Y. et al., *Anal. Chem.* 2010, 82, 3183-3190). Microfluidic chips with channel diameters of 10-100 μm are typically fabricated from quartz, silicon, glass, or polydimethylsiloxane (PDMS) using standard soft photolithography techniques (A. Manz, N. Graber and H.M. Widmer: Miniaturized total Chemical Analysis systems: A Novel Concept for Chemical Sensing, *Sensors and Actuators, B Chemical* (1990) 244-248). Droplets are typically generated at rates of ~ 1 -10Hz by flowing an aqueous solution in one channel into a stream of oil. The use of flow focusing nozzles enables generation of controlled size droplets of aqueous phase. The droplet size and rate of droplet generation are controlled by the ratio of oil and aqueous phase flow rates, for a given nozzle geometry. The chip channel surface is usually modified to be hydrophobic, for instance, by one of the many published silanization chemistries (Zeng, Y. et al., *Anal. Chem.* 2010, 82, 3183-3190). For droplets to be fully functional microvessels, the use of hydrophobic and lipophobic oils may be beneficial, since the molecular diffusion between

droplets is minimized, the oils have low solubility for biological reagents contained in the aqueous phase and have good gas solubility, which ensures viability of encapsulated cells in certain applications. In addition, surfactants may desirably, according to certain embodiments, be mixed into the oil phase, since droplets tend to coalesce. Surfactants may also inhibit adsorption of biomolecules at the microdroplet interfaces. A novel class of block copolymer surfactants, comprising perfluorinated polyethers (PFPE) coupled to polyethyleneglycol (PEG), has been described for use with fluorocarbon oils, for example, the fluorinated oil FC-40 (Sigma), a mix of perfluoro tri-n-butyl amine with di(perfluoro(n-butyl))perfluoromethyl amine (Holtze, C. et al., *Lab Chip*, 2008, DOI: 10.1039/b806706f). These compositions have led to very stable, biocompatible emulsions (Brouzes, E., et al., *PNAS* 2009, 106(34), 14195-14200).

[0085] Droplets traveling in microfluidic channels may be maintained as discrete microdroplets by means of their surface tension. Various methods have also been proposed to overcome the surface tension and allow droplets to merge when desired, thus allowing reagent mixing, *e.g.*, by microfabrication of passive, flow reducing elements in channels (Niu, X. et al., *Lab Chip* 2008, 8, 1837-1841), by the use of electrostatic charge (electrocoalescence) (Zagnoni, M. et al., *Langmuir*, 2010, 26(18), 14443-14449), or by manipulating microchannel geometry (Dolomite Merger chip; *see also* WO/2012/083225). A method of adding reagents to droplets in microfluidic channels via picoinjectors (pressurized reagent filled channels, perpendicular to the droplet channel, operated by electric fields), has recently been published (Abate, A.R. et al., *PNAS* 2010, 107(45), 19163-19166) and may also be adapted according to certain presently contemplated embodiments as described herein.

[0086] The microdroplet contents and the step of contacting are selected to be permissive for nucleic acid amplification interactions between the genomic DNA and the amplification primers. Nucleic acid amplification (*e.g.*, PCR) reagents and conditions are well known. Such amplification is permitted to proceed at least to obtain first and second double-stranded DNA products that include the nucleotide sequences of the first and second oligonucleotide amplification primers as provided herein, and the complementary sequences thereto. Thus, for example, any single fused microdroplet may contain (i) a first double-stranded DNA product that comprises at least a first universal adaptor sequence, the barcode sequence, a V region and a J or C region sequence that encode a portion of the first adaptive immune receptor polypeptide of the heterodimer, and a second universal adaptor sequence, and (ii) a second double-stranded DNA product that comprises at least a third universal adaptor

sequence, the same barcode sequence as in (i), a V region and a J or C region sequence that encode a portion of the second adaptive immune receptor polypeptide of the heterodimer, and a fourth universal adaptor sequence.

[0087] Conditions for the amplification step in the fused microdroplets are stopped prior to the next step. This can be achieved by changing the temperature of the environment in which the microdroplets are contained (e.g., in a container or well) to stop the amplification process.

[0088] In some embodiments, the method comprises disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of the first and second double-stranded products. Disruption may be selected on the basis of the chemical properties and composition of the microdroplets, and may be achieved, for instance, by chemical, biochemical and/or physical manipulations, such as the introduction of a diluent, detergent, chaotrope, surfactant, osmotic agent, or other chemical agent, or by the use of sonication, pressure, electrical field or other disruptive conditions. It will be appreciated that preferred conditions will involve the use of aqueous solvents for the included volumes within the microdroplets and/or for the heterogeneous mixture that is obtained by the step of disrupting. By using microdroplets instead of individual cells as an assay format, one can analyze data on the number of input cells in the sample. One can correct for PCR and sequencing errors, and in the case of IG molecules differentiate between non-germline sequences due to somatic hypermutation (SHM) from non-germline sequences introduced due to PCR error.

[0089] In some embodiments, the method comprises an ensuing step for contacting the mixture of first and second double-stranded DNA products with the herein described third and fourth amplification primer sets. Conditions for this step may similarly be achieved using accepted methodologies for DNA amplification to obtain a DNA library for sequencing, which may also be achieved according to any of a number of established DNA sequencing technologies. In certain related embodiments, instead of using first liquid microdroplets that each contain a single lymphoid cell or genomic DNA isolated therefrom, each of the first liquid microdroplets contains complementary DNA (cDNA) that has been reverse transcribed from the mRNA of a single lymphoid cell, such as a first cDNA that encodes the first chain of the adaptive immune receptor heterodimer and a second cDNA that encodes the second chain of the heterodimer.

[0090] In certain related embodiments, the individual second microdroplets may each contain a third oligonucleotide primer set that is capable of amplifying additional cDNA sequences that encode a lymphocyte status indicator molecule or molecules, The third primer set is

labeled with the same barcode sequence that is present in the first and second primer sets that are in the microdroplet. In such embodiments, the biological status can be determined for the single source cell from which a given TCR or IG heterodimeric sequence is identified. The biological status can be activated vs. quiescent, maturational stage, naïve vs. memory, regulatory vs. effector, etc. Exemplary lymphocyte status indicator molecules include, *e.g.*, lck, fyn, FoxP3, CD4, CD8, CD11a, CD18, CD25, CD28, CD29, CD44, CD45, CD49d, CD62, CD69, CD71, CD103, CD137 (4-1BB), HLA-DR, etc.

[0091] Certain embodiments include a third oligonucleotide primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule, where the third oligonucleotide primer set is labeled with the same barcode sequence that is present in the first and second primer sets, and where the lymphocyte status indicator molecule comprises one or more of the following: FoxP3, CD4, CD8, CD11a, CD18, CD21, CD25, CD29, CCD30, CD38, CD44, CD45, CD45RA, CD45RO, CD49d, CD62, CD62L, CD69, CD71, CD103, CD137 (4-1BB), CD138, CD161, CD294, CCR5, CXCR4, IgG1-4 H-chain constant region, IgA H-chain constant region, IgE H-chain constant region, IgD H-chain constant region, IgM H-chain constant region, HLA-DR, IL-2, IL-5, IL-6, IL-9, IL-10, IL-12, IL-13, IL-15, IL-21, TGF- β , TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9 and TLR10.

Table 1: EXEMPLARY LYMPHOCYTE STATUS INDICATORS

Gene Name	Status Marker for:	Human Transcript Sequence Accession #
FOXP3	Treg cells	NM_014009, NM_001114377
IL9	Th9 cells	NM_000590
CD21	EBV receptor on B cells	NM_001006658, NM_001877
CD30	Activated T and B cells, NK cells monocytes, and Reed-Sternburg cells (Hodgkin's Lymphoma)	NM_001243, NM_152942
CD38	Plasma cells, activated B and T cells	NM_001775
CD138	Plasma cells	NM_001006946, NM_002997

CD45RA	Naïve T cells	NM_002838, NM_080921, NM_001267798
CD45RO	Memory T cells	NM_002838, NM_080921, NM_001267798
CD62L	Homing of naïve cells to peripheral lymph nodes	NM_000655
CD294	TH2 cells	NM_004778
Helios	Thymic Treg cells	NM_001079526, NM_016260
CD161	NK cells	NM_002258
IL2	CD4 ⁺ T cells and some CD8 ⁺ T cells	NM_000586
IL5	TH2 cells	NM_000879
IL6	Macrophages, endothelial cells, and T cells	NM_000600
IL10	Macrophages and TH2 cells	NM_000572
TGF-β	T cells and macrophages	NM_000660
IL12B	Macrophages and dendritic cells	NM_002187
IL12A	Macrophages and dendritic cells	NM_000882
IL13	TH2 cells	NM_002188
IL15	Macrophages	NM_0172175, NM_000585
IL21	Activated T cells (mainly TH2, TH17, and NKT cells)	NM_021803, NM_001207006
CCR5	T cells and macrophages	NM_000579, NM_001100168
CXCR4	T cells	NM_003467, NM_001008540
IGHG1	IgG1 heavy chain constant region	AJ294730, J00228
IGHG2	IgG2 heavy chain constant region	AJ294731, J00230
IGHG3	IgG3 heavy chain constant region	D78345
IGHG4	IgG4 heavy chain constant region	AJ294733, K01316
IGHA1	IgA1 heavy chain constant region	J00220
IGHA2	IgA2 heavy chain constant region	M60192, J00221
IGHE	IgGE1 heavy chain constant region	L00022, J00222

IGHD	IgD heavy chain constant region	K02875, K02876, K02877, K02878, K02879, K02880, K02881, K02992, X57331
IGHM	IgM heavy chain constant region	J00260, K01310, X14939, X14940, X57331
TLR1	B cells	NM_003263
TLR2	T and B cells	NM_003264
TLR3	T cells	NM_003265
TLR4	T cells	NM_003266, NM_138554, NM_138557
TLR5	Treg and naïve T cells	NM_003268

[0092] These and related embodiments need not be so limited, however, such that there are also contemplated embodiments according to which, additionally or alternatively, there may be included a third oligonucleotide primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule, where the third primer set is labeled with the same barcode sequence that is present in the first and second primer sets, and where the lymphocyte status indicator molecule comprises a cell surface receptor.

[0093] Examples of cell surface receptors include the following, or the like: CD2 (e.g., GenBank Acc. Nos. Y00023, SEG_HUMCD2, M16336, M16445, SEG_MUSCD2, M14362), 4-1BB (CDw137, Kwon et al., 1989 *Proc. Nat. Acad. Sci. USA* 86:1963, 4-1BB ligand (Goodwin et al., 1993 *Eur. J. Immunol.* 23:2361; Melero et al., 1998 *Eur. J. Immunol.* 3:116), CD5 (e.g., GenBank Acc. Nos. X78985, X89405), CD10 (e.g., GenBank Acc. Nos. M81591, X76732) CD27 (e.g., GenBank Acc. Nos. M63928, L24495, L08096), CD28 (June et al., 1990 *Immunol. Today* 11:211; see also, e.g., GenBank Acc. Nos. J02988, SEG_HUMCD28, M34563), CD152/CTLA-4 (e.g., GenBank Acc. Nos. L15006, X05719, SEG_HUMIGCTL), CD40 (e.g., GenBank Acc. Nos. M83312, SEG_MUSC040A0, Y10507, X67878, X96710, U15637, L07414), interferon- γ (IFN- γ ; see, e.g., Farrar et al. 1993 *Ann. Rev. Immunol.* 11:571 and references cited therein, Gray et al. 1982 *Nature* 295:503, Rinderknecht et al. 1984 *J. Biol. Chem.* 259:6790, DeGrado et al. 1982 *Nature* 300:379), interleukin-4 (IL-4; see, e.g., 53rd *Forum in Immunology*, 1993 *Research in Immunol.* 144:553-643; Banchereau et al., 1994 in *The Cytokine Handbook*, 2nd ed., A. Thomson, ed., Academic Press, NY, p. 99; Keegan et al., 1994 *J Leukocyt. Biol.* 55:272, and references

cited therein), interleukin-17 (IL-17) (*e.g.*, GenBank Acc. Nos. U32659, U43088) and interleukin-17 receptor (IL-17R) (*e.g.*, GenBank Acc. Nos. U31993, U58917).

[0094] Additional cell surface receptors include the following or the like: CD59 (*e.g.*, GenBank Acc. Nos. SEG_HUMCD590, M95708, M34671), CD48 (*e.g.*, GenBank Acc. Nos. M59904), CD58/LFA-3 (*e.g.*, GenBank Acc. No. A25933, Y00636, E12817; see also JP 1997075090-A), CD72 (*e.g.*, GenBank Acc. Nos. AA311036, S40777, L35772), CD70 (*e.g.*, GenBank Acc. Nos. Y13636, S69339), CD80/B7.1 (Freeman et al., 1989 *J. Immunol.* 43:2714; Freeman et al., 1991 *J. Exp. Med.* 174:625; see also *e.g.*, GenBank Acc. Nos. U33208, I683379), CD86/B7.2 (Freeman et al., 1993 *J. Exp. Med.* 178:2185, Boriello et al., 1995 *J. Immunol.* 155:5490; see also, *e.g.*, GenBank Acc. Nos. AF099105, SEG_MMB72G, U39466, U04343, SEG_HSB725, L25606, L25259), B7-H1/B7-DC (*e.g.*, Genbank Acc. Nos. NM_014143, AF177937, AF317088; Dong et al., 2002 *Nat. Med.* Jun 24 [epub ahead of print], PMID 12091876; Tseng et al., 2001 *J. Exp. Med.* 193:839; Tamura et al., 2001 *Blood* 97:1809; Dong et al., 1999 *Nat. Med.* 5:1365), CD40 ligand (*e.g.*, GenBank Acc. Nos. SEG_HUMCD40L, X67878, X65453, L07414), IL-17 (*e.g.*, GenBank Acc. Nos. U32659, U43088), CD43 (*e.g.*, GenBank Acc. Nos. X52075, J04536), ICOS (*e.g.*, Genbank Acc. No. AH011568), CD3 (*e.g.*, Genbank Acc. Nos. NM_000073 (gamma subunit), NM_000733 (epsilon subunit), X73617 (delta subunit)), CD4 (*e.g.*, Genbank Acc. No. NM_000616), CD25 (*e.g.*, Genbank Acc. No. NM_000417), CD8 (*e.g.*, Genbank Acc. No. M12828), CD11b (*e.g.*, Genbank Acc. No. J03925), CD14 (*e.g.*, Genbank Acc. No. XM_039364), CD56 (*e.g.*, Genbank Acc. No. U63041), CD69 (*e.g.*, Genbank Acc. No. NM_001781) and VLA-4 ($\alpha_4\beta_7$) (*e.g.*, GenBank Acc. Nos. L12002, X16983, L20788, U97031, L24913, M68892, M95632).

[0095] The following cell surface receptors are typically associated with B cells: CD19 (*e.g.*, GenBank Acc. Nos. SEG_HUMCD19W0, M84371, SEG_MUSCD19W, M62542), CD20 (*e.g.*, GenBank Acc. Nos. SEG_HUMCD20, M62541), CD22 (*e.g.*, GenBank Acc. Nos. I680629, Y10210, X59350, U62631, X52782, L16928), CD30 (*e.g.*, Genbank Acc. Nos. M83554, D86042), CD153 (CD30 ligand, *e.g.*, GenBank Acc. Nos. L09753, M83554), CD37 (*e.g.*, GenBank Acc. Nos. SEG_MMCD37X, X14046, X53517), CD50 (ICAM-3, *e.g.*, GenBank Acc. No. NM_002162), CD106 (VCAM-1) (*e.g.*, GenBank Acc. Nos. X53051, X67783, SEG_MMVCAM1C, *see also* U.S. Patent No. 5,596,090), CD54 (ICAM-1) (*e.g.*, GenBank Acc. Nos. X84737, S82847, X06990, J03132, SEG_MUSICAM0), interleukin-12 (see, *e.g.*, Reiter et al, 1993 *Crit. Rev. Immunol.* 13:1, and references cited therein), CD134 (OX40, *e.g.*, GenBank Acc. No. AJ277151), CD137 (41BB, *e.g.*, GenBank Acc. No. L12964,

NM_001561), CD83 (*e.g.*, GenBank Acc. Nos. AF001036, AL021918), DEC-205 (*e.g.*, GenBank Acc. Nos. AF011333, U19271).

[0096] Examples of other cell surface receptors include the following, or the like: HER1 (*e.g.*, GenBank Accession Nos. U48722, SEG_HEGFREXS, KO3193), HER2 (Yoshino et al., 1994 *J. Immunol.* 152:2393; Disis et al., 1994 *Canc. Res.* 54:16; see also, *e.g.*, GenBank Acc. Nos. X03363, M17730, SEG_HUMHER20), HER3 (*e.g.*, GenBank Acc. Nos. U29339, M34309), HER4 (Plowman et al., 1993 *Nature* 366:473; see also *e.g.*, GenBank Acc. Nos. L07868, T64105), epidermal growth factor receptor (EGFR) (*e.g.*, GenBank Acc. Nos. U48722, SEG_HEGFREXS, KO3193), vascular endothelial cell growth factor (*e.g.*, GenBank No. M32977), vascular endothelial cell growth factor receptor (*e.g.*, GenBank Acc. Nos. AF022375, 1680143, U48801, X62568), insulin-like growth factor-I (*e.g.*, GenBank Acc. Nos. X00173, X56774, X56773, X06043, *see also* European Patent No. GB 2241703), insulin-like growth factor-II (*e.g.*, GenBank Acc. Nos. X03562, X00910, SEG_HUMGFIA, SEG_HUMGFI2, M17863, M17862), transferrin receptor (Trowbridge and Omary, 1981 *Proc. Nat. Acad. USA* 78:3039; see also *e.g.*, GenBank Acc. Nos. X01060, M11507), estrogen receptor (*e.g.*, GenBank Acc. Nos. M38651, X03635, X99101, U47678, M12674), progesterone receptor (*e.g.*, GenBank Acc. Nos. X51730, X69068, M15716), follicle stimulating hormone receptor (FSH-R) (*e.g.*, GenBank Acc. Nos. Z34260, M65085), retinoic acid receptor (*e.g.*, GenBank Acc. Nos. L12060, M60909, X77664, X57280, X07282, X06538), MUC-1 (Barnes et al., 1989 *Proc. Nat. Acad. Sci. USA* 86:7159; see also *e.g.*, GenBank Acc. Nos. SEG_MUSMUCIO, M65132, M64928) NY-ESO-1 (*e.g.*, GenBank Acc. Nos. AJ003149, U87459), NA 17-A (*e.g.*, European Patent No. WO 96/40039), Melan-A/MART-1 (Kawakami et al., 1994 *Proc. Nat. Acad. Sci. USA* 91:3515; see also *e.g.*, GenBank Acc. Nos. U06654, U06452), tyrosinase (Topalian et al., 1994 *Proc. Nat. Acad. Sci. USA* 91:9461; see also *e.g.*, GenBank Acc. Nos. M26729, SEG_HUMTYR0, *see also* Weber et al., *J. Clin. Invest* (1998) 102:1258), Gp-100 (Kawakami et al., 1994 *Proc. Nat. Acad. Sci. USA* 91:3515; see also *e.g.*, GenBank Acc. No. S73003, *see also* European Patent No. EP 668350; Adema et al., 1994 *J. Biol. Chem.* 269:20126), MAGE (van den Bruggen et al., 1991 *Science* 254:1643; see also *e.g.*, GenBank Acc. Nos. U93163, AF064589, U66083, D32077, D32076, D32075, U10694, U10693, U10691, U10690, U10689, U10688, U10687, U10686, U10685, L18877, U10340, U10339, L18920, U03735, M77481), BAGE (*e.g.*, GenBank Acc. No. U19180, *see also* U.S. Patent Nos. 5,683,886 and 5,571,711), GAGE (*e.g.*, GenBank Acc. Nos. AF055475, AF055474, AF055473, U19147, U19146, U19145, U19144, U19143,

U19142), any of the CTA class of receptors including in particular HOM-MEL-40 antigen encoded by the SSX2 gene (*e.g.*, GenBank Acc. Nos. X86175, U90842, U90841, X86174), carcinoembryonic antigen (CEA, Gold and Freedman, 1985 *J. Exp. Med.* 121:439; see also *e.g.*, GenBank Acc. Nos. SEG_HUMCEA, M59710, M59255, M29540), and PyLT (*e.g.*, GenBank Acc. Nos. J02289, J02038).

[0097] A lymphocyte status indicator may also include one or more apoptosis signaling polypeptides, sequences of which are known to the art, as reviewed, for example, in *When Cells Die: A Comprehensive Evaluation of Apoptosis and Programmed Cell Death* (R.A. Lockshin et al., Eds., 1998 John Wiley & Sons, New York; see also, *e.g.*, Green et al., 1998 *Science* 281:1309 and references cited therein; Ferreira et al., 2002 *Clin. Canc. Res.* 8:2024; Gurumurthy et al., 2001 *Cancer Metastas. Rev.* 20:225; Kanduc et al., 2002 *Int. J. Oncol.* 21:165). Typically, an apoptosis signaling polypeptide sequence comprises all or a portion of, or is derived from, a receptor death domain polypeptide, for instance, FADD (*e.g.*, Genbank Acc. Nos. U24231, U43184, AF009616, AF009617, NM_012115), TRADD (*e.g.*, Genbank Acc. No. NM_003789), RAIDD (*e.g.*, Genbank Acc. No. U87229), CD95 (FAS/Apo-1; *e.g.*, Genbank Acc. Nos. X89101, NM_003824, AF344850, AF344856), TNF- α -receptor-1 (TNFR1, *e.g.*, Genbank Acc. Nos. S63368, AF040257), DR5 (*e.g.*, Genbank Acc. No. AF020501, AF016268, AF012535), an ITIM domain (*e.g.*, Genbank Acc. Nos. AF081675, BC015731, NM_006840, NM_006844, NM_006847, XM_017977; see, *e.g.*, Billadeau et al., 2002 *J. Clin. Invest.* 109:161), an ITAM domain (*e.g.*, Genbank Acc. Nos. NM_005843, NM_003473, BC030586; see, *e.g.*, Billadeau et al., 2002), or other apoptosis-associated receptor death domain polypeptides known to the art, for example, TNFR2 (*e.g.*, Genbank Acc. No. L49431, L49432), caspase/procaspase-3 (*e.g.*, Genbank Acc. No. XM_54686), caspase/procaspase-8 (*e.g.*, AF380342, NM_004208, NM_001228, NM_033355, NM_033356, NM_033357, NM_033358), caspase/procaspase-2 (*e.g.*, Genbank Acc. No. AF314174, AF314175), etc. Cells in a biological sample that are suspected of undergoing apoptosis may be examined for morphological, permeability, biochemical, molecular genetic, or other changes that will be apparent to those familiar with the art.

[0098] These and related methods for the first time permit rapid determination of the rearranged DNA sequences that encode both chains of a TCR or IG heterodimer from a single cell. Such embodiments will find uses for diagnostic and prognostic purposes, by permitting high-throughput sequencing of adaptive immune receptor encoding sequences from each of a plurality of single cells, and will also usefully inform immunological

investigations into TCR or IG heterodimeric pairings and their underlying molecular mechanisms. The rapid and large-scale availability of DNA sequence information for both subunits of a large number of TCR and/or IG heterodimers will accelerate development of synthetic antibody technologies and related arts, for example, where antibodies or complete or partial TCR or IG antigen-binding regions may be usefully engineered into diagnostic, therapeutic, biomimetic, enzymatic or catalytic (*e.g.*, Abzymes) or other industrially useful compositions. By virtue of the quantitative nature of the high throughput TCR and/or IG sequencing afforded by the present disclosure, high precision in the quantitative characterization of TCR and/or IG heterodimer sequences that are present in a sample will advantageously improve the ability to determine the number of cells that belong to a specific T cell or B cell clone.

[0099] As noted above, according to these embodiments for identifying both chains of a TCR or IG heterodimer from a single adaptive immune cell, in any given second microdroplet, all oligonucleotide amplification primers will comprise the same barcode oligonucleotide, but within different second microdroplets the primer sets will comprise different barcode sequences. Accordingly, after sequencing the DNA library obtained as described above to obtain a data set of sequences, the sequences in the data set can be sorted into groups of sequences that have identical barcode sequences, and such barcode groups can be further sorted into those having X1 or X2 sequences (which include portions of V and J or C regions) that will indicate whether a given sequence reflects the amplification product of a first TCR or IG encoding chain (*e.g.*, a TCRA or IGH chain) or a second TCR or IG encoding chain (*e.g.*, a TCRB or IGL chain).

[00100] Sequences that have been so sorted by barcode and by TCR or IG chain may be further subject to cluster analysis using any of a known variety of algorithms for clustering (*e.g.*, BLASTClust, UCLUST, CD-HIT, *see also* IEEE Rev Biomed Eng. 2010;3:120-54. doi: 10.1109/RBME.2010.2083647; Clustering algorithms in biomedical research: a review, Xu R, Wunsch DC 2nd; *Mol Biotechnol.* 2005 Sep;31(1):55-80; Data clustering in life sciences. Zhao Y, Karypis G; *Methods Mol Biol.* 2010;593:81-107. doi: 10.1007/978-1-60327-194-3_5; Overview on techniques in cluster analysis. Frades I, Matthiesen R, and error correction in the case of sequences that fail to cluster with other sequences having shared barcode sequences but which instead would cluster with sequences having a barcode that differs by a single nucleotide. *See, e.g., Proc Natl Acad Sci U S A.* 2012 Jan 24;109(4):1347-52. doi: 10.1073/pnas.1118018109. Epub 2012 Jan 9. Digital RNA sequencing minimizes sequence-

dependent bias and amplification noise with optimized single-molecule barcodes. Shiroguchi K, Jia TZ, Sims PA, Xie XS; *Proc Natl Acad Sci U S A*. 2012 Sep 4;109(36):14508-13. doi: 10.1073/pnas.1208715109. Epub 2012 Aug 1. Detection of ultra-rare mutations by next-generation sequencing. Schmitt MW, Kennedy SR, Salk JJ, Fox EJ, Hiatt JB, Loeb LA.

[00101] Accordingly, certain embodiments comprise a method including steps of (a) sorting the data set of sequences (obtained as described above) according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode; (b) sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset; (c) clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets; and (d) identifying as originating from the same cell sequences that are members of an X1 and an X2 sequence cluster set that belong to the same one or more barcode sequence sets.

[00102] It will be appreciated that according to non-limiting theory, first and second adaptive immune receptor chain encoding sequences that occur with the same set of barcode sequences have an extremely high probability of having originated from the same fused microdroplet, and thus from the same source cell. For example, where 10^4 different barcodes are used in the construction of the first and second oligonucleotide amplification primers, the probability that two independent (*i.e.*, originating from different cells) double-stranded first and second products would be obtained having the same barcode sequence is one in 10^8 . Hence, if according to the methods described herein, three or more copies of a given set of first and second adaptive immune receptor polypeptide encoding sequences (*e.g.*, X1 and X2) share common barcode sequences (*e.g.*, belong to the same barcode sequence set), the probability that the sequences are of independent cellular origin approaches zero.

[00103] Similarly, it will be appreciated that analysis of the data set of sequences obtained according to the present methods may also be used to characterize the biological status of the lymphoid cell source of genomic DNA. For example, because in B cells IGH gene rearrangement is known to precede IGL gene rearrangement, barcode sequence analysis as described herein may reveal multiple single lymphoid cell genomes having the same rearranged IGH sequence but different IGL sequences, indicating origins of these sequences in immunologically naïve cells.

[00104] Alternatively, the analysis may exploit the observation that T cells express proteins that are specific to their functions, such as lymphocyte status indicator molecules as described herein. For example, regulatory T cells express the protein FOXP3. If a cDNA that has been reverse transcribed from T cell mRNA is subsequently amplified, co-amplification products may include cDNA species that reflect other mRNAs encoding phenotypic specific proteins such as FOXP3, along with cDNAs encoding the TCRB and TCRA molecules. This approach may permit identification of the adaptive immune receptors that are expressed by T cells having specific phenotypes, such as T regulatory cells or effector T cells.

[00105] Thus, there is provided herein a method for determining rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising (1) contacting (A) individual first microdroplets that each contain a single lymphoid cell or genomic DNA isolated therefrom, with (B) individual second microdroplets from a plurality of second liquid microdroplets that each contain (i) a first oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a first complementarity determining region-3 (CDR3) of a first polypeptide of an adaptive immune receptor heterodimer, and (ii) a second oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a second complementarity determining region-3 (CDR3) of a second polypeptide of the adaptive immune receptor heterodimer. The first oligonucleotide amplification primer set comprises a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula: U1/2-B1-X1, in which U1/2 comprises an oligonucleotide which comprises a first universal adaptor oligonucleotide sequence when B1 is present or a second universal adaptor oligonucleotide sequence when B1 is nothing. In some embodiments, B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides, and X1 comprises an oligonucleotide that is one of: (a) a polynucleotide comprising at least 20, 30, 40 or 50 and not more than 100, 90, 80, 70 or 60 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence, and (b) a polynucleotide comprising at least 15-30 or 31-50 and not more than 80, 70, 60 or 55

contiguous nucleotides of either (i) an adaptive immune receptor joining (J) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence. The second oligonucleotide amplification primer set can comprise a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula: U3/4-B2-X2 in which U3/4 comprises an oligonucleotide which comprises a third universal adaptor oligonucleotide sequence when B2 is present or a fourth universal adaptor oligonucleotide sequence when B2 is nothing, B2 comprises an oligonucleotide that comprises either nothing or a second oligonucleotide barcode sequence of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides that is from the same as B1, and X2 comprises an oligonucleotide that is one of: (a) a polynucleotide comprising at least 20, 30, 40 or 50 and not more than 100, 90, 80, 70 or 60 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence, and (b) a polynucleotide comprising at least 15-30 or 31-50 and not more than 80, 70, 60 or 55 contiguous nucleotides of either (i) an adaptive immune receptor joining (J) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence. The step of contacting can take place under conditions and for a time sufficient for a plurality of fusion events between one of the first microdroplets and one of the second microdroplets to produce a plurality of fused microdroplets in which nucleic acid amplification interactions occur between the genomic DNA and the first and second oligonucleotide amplification primer sets, to obtain in each of one or more of said plurality of fused microdroplets: a first double-stranded DNA product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said

first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer. The conditions also permit obtaining in each of one or more of said plurality of fused microdroplets: a second double-stranded DNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer.

[00106] The method also includes disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of said first and second double-stranded DNA products and contacting the mixture of first and second double-stranded DNA products with a third amplification primer set and a fourth amplification primer set. In some embodiments, the third amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence. In other embodiments, the fourth amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence.

The contacting step can take place under conditions and for a time sufficient to amplify both strands of the first and second double-stranded DNA products of (2), to obtain a DNA library for sequencing. The method also includes sequencing the DNA library obtained in (3) to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

[00107] Figure 2 illustrates one method by which a plurality of first microdroplets 210 that contain a single lymphoid cell or genomic DNA fuse with a plurality of individual second microdroplets 220 to form a plurality of fused microdroplets 230. The second plurality of droplets may comprise amplification primer sets, as described herein, and the fused droplets can be placed under conditions where the amplification primers can amplify the DNA found in the single lymphoid cell or the genomic DNA (or cDNA) within the microdroplet.

[00108] These and related embodiments permit high throughput sequencing of rearranged genes encoding both chains from the same cell of an adaptive immune receptor heterodimer, such as IGH plus IGL, or IGH plus IGK, or TCRA plus TCRB, or TCRG plus TCRD. Advantageously, this approach also permits quantifying the number of cells having a given TCR or IG. A schematic depiction of an exemplary embodiment is shown in Figure 3, according to which steps highly similar to those described above are carried out, significantly, however, with the step of contacting DNA from a single lymphoid cell with first and second amplification primer sets as described herein to effect the first amplification reaction by which the unique molecular-tagging barcode is incorporated taking place within a single microdroplet, such as those that are formed from emulsions for use in the RainDance™ microdroplet digital PCR system (RainDance Technologies, Lexington, MA) (e.g., Pekin et al., 2011 *Lab. Chip* 11(13):2156; Zhong et al., 2011 *Lab. Chip* 11(13):2167; Tewhey et al., 2009 *Nature Biotechnol.* 27:1025; 2010 *Nature Biotechnol.* 28:178) or other comparable systems, any of which may be adapted by the skilled person for use with the herein described compositions and methods. Subsequent to the incorporation into a plurality of distinct dsDNA products of the plurality of unique molecular-tagging barcodes, the microdroplets may be disrupted and the ensuing steps that include amplifying and introducing sequencing platform-specific oligonucleotides may be carried out as described herein and shown in Fig. 3.

[00109] In these and related embodiments, a single tagging barcode (BC1) may be shared by all J primers (or in certain embodiments by all V primers) and it may be desirable to produce such primers with a finite set of specific and pre-identified barcode sequences. Only

a single tagging barcode sequence (BC1) will be present within any given microdroplet during the first step, however. Hence, even after a large and diverse set of sequence information is obtained following the sequencing step when practiced starting with a sample that comprises a plurality of heterogeneous lymphoid cells as provided herein, analysis of such information may include determination of first and second TCR or Ig heterodimeric polypeptide chain encoding sequences that contain the same tagging barcode (BC1), from which a probabilistic basis would indicate an extremely high likelihood that both chains are the products of the same cell. Accordingly, the present disclosure for the first time provides compositions and methods for determining and quantifying the relative representation in a sample of both chains of a TCR or Ig heterodimer that are expressed in the same cell.

[00110] Clonal Heterodimer Sequence Determination Without MicroDroplets

[00111] According to certain other embodiments, determination of rearranged DNA sequences encoding first and second adaptive immune receptor heterodimer polypeptide sequences in a single cell may be achieved without first preparing separate populations of first and second microdroplets that contain, respectively, single lymphoid cell genomic DNA (or cDNA that has been reverse transcribed from mRNA therefrom) and oligonucleotide amplification primer sets.

[00112] Instead, these alternative embodiments contemplate separating the cells of a lymphoid cell-containing cell suspension (e.g., a blood cell preparation from a subject or a cell subpopulation thereof) into subpopulations by distributing the cells to a plurality of containers, such as multiple wells of a multi-well cell culture plate or assay plate (e.g., 96-, 384- or 1536-well formats). Persons familiar with the art will be aware of a number of devices and methodologies for distributing a cell suspension into such multiple containers, for instance, using fluorescence activated cell sorting (FACS) or with automated low-volume dispensing equipment or by limiting dilution, to obtain a desired number of cells per well, container, tube, compartment or the like. In certain embodiments it may be preferred to distribute substantially the same number of cells to each container, although certain other contemplated embodiments need not be so limited.

[00113] Briefly, according to these and related embodiments, separated lymphoid cell subpopulations may provide mRNA molecules that are used as templates for reverse transcription to produce cDNA molecules that are concomitantly labeled during the reverse transcription (RT) step (see Figures 4 and 5). Figure 4 depicts a schematic representation of labeling adaptive immune receptor polypeptide encoding cDNA during reverse transcription

by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences into cDNA. The cDNA strand is amplified with primers comprising a pGEX-Rev sequence, a barcode BC and N6 spacer sequence (BC-N6) and a “Cn-RC” sequence. The 3’ end of the amplified cDNA strand includes a pGEX-FRC sequence, a barcode BC-N6 spacer sequence, and a “Smarter UAll” sequence. The wells or containers of amplified cDNA are pooled, and SPRI bead purification is performed of the first cDNA strand pool. PCR amplification is performed using a tailing-pGEX F/R sequence. The amplicons are purified and selected based on size. The resulting cDNA amplicon is shown in Figure 4.

[00114] Figure 5 depicts a schematic representation of labeling adaptive immune receptor polypeptide encoding cDNA during reverse transcription by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences into cDNA. Figure 6 presents a schematic representation of a DNA product that is amenable to sequencing following modification with Illumina sequencing adapters of amplified adaptive immune receptor polypeptide encoding cDNA that has been labeled during reverse transcription by using an oligonucleotide reverse transcription primer that directs incorporation of oligonucleotide barcode and universal adaptor oligonucleotide sequences.

[00115] As provided herein, oligonucleotide RT primers in such embodiments include oligonucleotide sequences that specifically hybridize to target adaptive immune receptor encoding regions such as V, J or C region sequences, and also include oligonucleotide barcode sequences as molecular labels, along with universal adaptor oligonucleotide sequences as described herein. The process of reverse transcription from adaptive immune receptor encoding mRNA may thus be accompanied by incorporation into cDNA products of (i) oligonucleotide barcode sequences as source identifiers, and (ii) universal adaptors to facilitate automated high throughput sequencing as described herein. By way of illustration and not limitation, in certain of these embodiments all RT primers in the oligonucleotide RT primer sets that are contacted with the contents of a single particular container (e.g., one well of a multi-well plate) share a common barcode oligonucleotide sequence (B), and a different barcode oligonucleotide sequence (B) is present in each separate container (such as each well of a multi-well plate).

[00116] For instance, a cell suspension (e.g., blood cells or a fraction thereof, such as nucleated cells, lymphoid cells, etc.) may be divided by random distribution among different

wells of a multi-well plate to physically separate the cells into subsets. The subset of cells in each well may then be lysed or otherwise processed according to any of a number of conventional procedures to liberate mRNA present within the cells, which may include mRNA encoding both chains of TCR (e.g., TCRA and TCRB, or TCRG and TCRG) or IG (e.g., IGH and IGL) heterodimers expressed by the cells, and which may also include mRNA encoding one or more lymphocyte status indicator molecules.

[00117] The mRNA may then be used as a template for cDNA synthesis by modification of established reverse transcription (RT) protocols, using oligonucleotide reverse transcription primer sets as described herein that are capable of introducing into the cDNA products, in each separate well, a unique oligonucleotide barcode sequence that is linked to the TCR or IG encoding sequence or complement thereof (see, e.g., Figs. 4-5). External to the barcode (e.g., distal from the TCR or IG encoding sequence, relative to the barcode), the oligonucleotide reverse transcription primer sets may also be designed to introduce a universal adaptor oligonucleotide sequence as described herein and/or other known oligonucleotide sequence features such as those that may facilitate downstream amplification, processing and/or other manipulation steps such as those that will be compatible with automated high throughput quantitative sequencing.

[00118] Following DNA amplification of the reverse transcription cDNA products, each amplified DNA molecule within a given well of the multi-well plate will have the same oligonucleotide barcode sequence, while the barcode sequences of the amplification products in each different well will be distinct from one another. In this manner within each well, all DNA molecules that encode either chain of an adaptive immune receptor heterodimer (e.g., IGH and IGL, TCRA and TCRB, TCRG and TCRD) will have the same oligonucleotide barcode sequence.

[00119] The amplification products may be pooled and quantitatively sequenced using automated high throughput DNA sequencing as described elsewhere herein to obtain a data set of sequences, which include TCR and/or IG sequences along with associated oligonucleotide barcode sequences. As disclosed herein, in certain preferred embodiments the data set of sequences may be analyzed by a combinatorics approach, which permits matching particular pairs of adaptive immune receptor heterodimer subunit encoding sequences to identify them as having originated from the same lymphoid cell.

[00120] As a non-limiting illustrative example, a hypothetical data set of sequences may be obtained from a set of 100 wells into which a lymphoid cell suspension is distributed. In

each well, the cells' mRNA cDNA is reverse transcribed using first and second oligonucleotide reverse transcription primer sets that are specific, respectively, for portions of TCRA and TCRB encoding sequences. The oligonucleotide reverse transcription primer sets also introduce a different oligonucleotide barcode sequence into the cDNA products in each distinct well. If, hypothetically, T cells having a single, common clonal origin (e.g., T cells that express the identical TCRA/B sequences) are randomly distributed into five different wells of the 100 wells, then the sequence data set will include five separate instances in which the unique pair of TCRA and TCRB sequences occurs in DNA amplification products that share an identical barcode sequence. In other words, in each of the five separate wells, the oligonucleotide reverse transcription primer set promotes the generation of cDNAs having identical rearranged TCRA and TCRB sequences, but the cDNA products of each well include a distinct, well-specific barcode sequence. According to non-limiting theory, on a probabilistic basis the likelihood would be extremely high that the unique TCRA/TCRB sequence pair originates in the same T cell clone, members of which would have been randomly distributed into the five different wells.

[00121] According to certain embodiments, a more detailed description of this high throughput method for determining rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell is as follows:

[00122] Lymphoid cells are isolated from an anti-coagulated whole blood sample using either density gradient centrifugation (e.g., FicollPaque®, GE Healthcare Bio-Sciences, Piscataway, NJ), or by binding to antibody-coated magnetic beads, such as CD45 beads from Miltenyi Biotec (Auburn, CA). Alternatively, T lymphocytes may be purified from a whole blood sample by binding to CD3+ magnetic beads, and B lymphocytes may be purified from a whole blood sample by binding to CD19+ magnetic beads. Isolated cell populations may then be checked for viability. Dead cells may be removed from the sample with a filter, for example, using a Miltenyi Biotec Dead Cell Removal kit. Depending on the application, isolated viable lymphoid cells (e.g., as may be present in unsorted peripheral blood mononuclear cells (PBMC), or as preparations of specific cell sub-sets) may be cultured in short-term cell culture, and in certain embodiments cells may be activated by any of a number of known activation paradigms, such as by exposure to one or more of cytokines, chemokines, specific antibodies, mitogens, polyclonal activators, etc. The final cell sample may be prepared by resuspending the cells in culture media (e.g., RPMI with 10% fetal

bovine serum) or appropriate isotonic buffered solutions (*e.g.*, phosphate buffered saline, PBS), supplemented with agents which prevent cell clumping (*e.g.*, 0.1% BSA, 1% Pluronic® F-68). Alternatively, whole blood or PBMCs may be utilized without sorting. As the most general case, any set of cells present as a suspension in an aqueous solution that contains B or T cells may be used.

[00123] The cell preparation comprising a plurality of lymphoid cells is divided into a plurality of physically separated subsets, for example, by distributing the suspension of cells amongst a plurality of containers or compartments that are capable of containing the cells to obtain a plurality of containers or compartments that each contain a subpopulation of the lymphoid cells, wherein each subpopulation comprises one lymphoid cell or a plurality of lymphoid cells, and wherein each container or compartment is physically separate so that the contents are not in fluid communication with one another. Preferably the cells are distributed or divided into the plurality of containers so that each container contains a substantially equivalent number of cells, which may result in there being the same number of cells in each container, or in there being in each container a number of cells that is within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-30, 31-50, 51-70, 71-80, or 81-100 percent of the number of cells in any other container. Exemplary containers may be wells of multi-well culture or assay plates such as 6-, 12-, 24-, 48-, 96-, 384- or 1536-well multi-well plates or any other multi-well plate format; arrays of tubes, filters, microfabricated well arrays, laser-generated matrices or any other suitable containers that are capable of containing the cells are also contemplated. In certain exemplary embodiments, cells may be distributed amongst the plurality of containers by fluorescence activated cell sorting (FACS): A predetermined number of cells may be isolated, sorted, and deposited into a multi-well (*e.g.*, 96, 384 or 1536) reaction plate using FACS. Any of a number of methodologies and instrumentation may be employed using flow cytometers that are capable of preparative sorting of cells onto multi-well plates (*e.g.*, Beckton Dickinson FACS Aria® III, Beckman MoFlo™ XDP, etc.). FACS allows for specific subsets of cells to be isolated by antibody staining, viability staining or multicolor combination of specific cell staining reagents. Cell sorters may be employed to count target cells and deposit specified numbers of cells into each well of a collection multi-well plate (10-20%CV). Alternatively, automated low volume (nl to µl volumes per well) dispensers, capable of preferably non-contact dispensing of uniform cell suspensions onto high density micro-well plates (384, 1536, 3456 wells), such as Beckman Coulter BioRAPTR FRD™, LambdaJet™ IIIMT (Thermo Fisher Scientific),

CyBi™ Drop (Jena Analytik), Furukawa Perflow™, or similar instruments, may be used to deposit specified numbers of cells into each well of a collection multi-well plate with high precision and reproducibility (10-20%CV).

[00124] The adaptive immune receptor encoding polynucleotide sequences are then amplified from each well, with a unique, well-specific, barcode oligonucleotide attached to all samples. One way to do this is to convert cellular mRNA to cDNA by reverse transcription, and to add to the cDNA products a molecular label in the form of an oligonucleotide barcode during the reverse transcription step. The same barcode may be added to cDNAs that are complementary to mRNAs encoding both chains of each heterodimeric adaptive immune receptor molecule within the well, for instance, the immunoglobulin heavy and light chains, the TCRA and TCRB chains, and the TCRG and TCRD chains. In this and related embodiments, antigen receptor encoding sequences are amplified from cDNA made by reverse transcription from mRNA; genomic DNA (gDNA) is not amplified. To do this, each well of a microwell plate may contain a medium containing an RNase inhibitor, and a medium designed either to protect RNA in cells (such as Qiagen RNeasy™, Qiagen, Valencia, CA), or to lyse cells and isolate RNA (Trizol, guanidium isothiocyanate - Qiagen RNeasy™ etc.). Extracted total cellular RNA may then be transferred into another multi-well plate for the reverse transcription reaction using robotic liquid handlers. Alternatively, sorted cells may be lysed directly in a reverse-transcription reaction mix containing an RNase inhibitor. Reverse transcription reaction (RT) may be initiated by exposing cellular RNA to a reaction mix containing an appropriate buffer, dNTPs, an enzyme (reverse transcriptase) and a set of oligonucleotide reverse transcription primers. These primers will generally comprise a multiplicity of subsets of primers that may anneal to IgG, IgM, IgA, IgD, IgE, Ig kappa, Ig lambda, TCR alpha, beta, gamma and delta constant region (C-segment) gene-specific oligonucleotide sequences, as well as a universal template switching oligonucleotide (*e.g.*, Clontech Smarter™ UAII oligonucleotide, Clontech, Mountain View, CA). For instance, either the C-segment gene specific primers, or the Smarter™ UAII oligonucleotide, or both, will be uniquely tagged with a DNA barcode, which will be a unique sequence 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, ... etc. base pairs long. Each well of the RT reaction plate will contain the same multiplicity of primers, where each primer in the mix will be tagged with the same DNA barcode, but a different barcode will be used in each well. Thus, upon completion of the reverse transcription reaction, each first

strand cDNA molecule in a given well will be barcoded with an identical DNA barcode sequence.

List of BCR / TCR C-segment primers for 1st cDNA strand synthesis:

Name	Sequence	SEQ ID NO:
Ck	GATGAAGACAGATGGTGCAGC	5579
Cl-1	GGCGGGAACAGAGTGAC	5580
Cl-2	AGGGTGGGAACAGAGTGAC	5581
Cl-3	GCTTGAAGCTCCTCAGAGG	5582
Cl-4	GGCGGGAACAGAGTGAC	5583
IgA	AGGCTCAGCGGGAAGAC	5584
IgD	GAACACATCCGGAGCCTTG	5585
IgE	GGTGGCATTGGAGGGAATG	5586
IgG-1	AAGACCGATGGGCCCTTG	5587
IgG-2	CTCTCGGAGGTGCTCCTG	5588
IgM	AATTCTCACAGGAGACGAGGG	5589
TCRa	TGGTACACGGCAGGGTC	5590
TCRA_RACE_JB2	AGTCTCTCAGCTGGTACACGGCAGGGTC	5591
	5'-AGTCTCTCAGCTGGTACACGGCAGGGTC-3'	5591
TCRA_50	5'- ACA GAC TTG TCA CTG GAT TTA GAG TCT CTC AGC TGG TAC ACG GCA GGG TC -3'	5592
TCRB_50	5'- GAG ATC TCT GCT TCT GAT GGC TCA AAC ACA GCG ACC TCG GGT GGG AAC AC -3'	5593
TCRb-1	CAAACACAGCGACCTCGG	5594
TCRb-2	ATGGCTCAAACACAGCGAC	5595
TRCd-1	GATGGTTTGGTATGAGGCTGAC	5596
TCRd-2	CCTTCACCAGACAAGCGAC	5597
TCRg-1	GAAAAATAGTGGGCTTGGGGG	5598
	Primers from Bolotin et al., Eur. J. Immunol. 2012	
TCRb_BC1R	CAGTATCTGGAGTCATTGA	5599
TCRb_BC2R	TGCTTCTGATGGCTCAAACAC	5600

	Primers from Glanville et al., PNAS 2011	
IgM_RACE	5'-GATGGAGTCGGGAAGGAAGTCCTGTGCGAG-3'	5601
IgG_RACE	5'-GGGAAGACSGATGGGCCCTTGGTGG-3'	5602
IgA_RACE	5'-CAGGCAKGCGAYGACCACGTTCCCATC-3'	5603
Igκ_RACE	5'-CATCAGATGGCGGGAAGATGAAGACAGATGGTGC-3'	5604
Igλ_RACE	5'-CCTCAGAGGAGGGTGGGAACAGAGTGAC-3'	5605
TCRB_RACE	5'-GCTCAAACACAGCGACCTCGGGTGGGAACAC-3'	5606
	Clontech Smarter primers	
Smarter UAI	5'-AAGCAGTGGTATCAACGCAGAGTACrGrGrGrGrG-P-3	5607
Islam UAI	5'- AAGCAGTGGTATCAACGCAGAGTGCAGUGCUXXXXXXr GrGrG-3'	5608
Smarter CDS	5'-Bio-AAGCAGTGGTATCAACGCAGAGTACT(30)N-1N-3'	5609
Smarter IS PCR	5'-Bio-AAGCAGTGGTATCAACGCAGAGT -3'	5610
5'RACE long	5'- CTAATACGACTCACTATAGGGCAAGCAGTGGTATCAAC GCAGAGT-3'	5611
5'RACE short	5'-CTAATACGACTCACTATAGGGC-3'	5612

[00125] Accordingly, following the step of distributing cells to a plurality of containers, each of the containers is contacted, under conditions and for a time sufficient to promote reverse transcription of mRNA in the lymphoid cells in the plurality of containers, with a first and a second oligonucleotide reverse transcription primer set, wherein (A) the first oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of first mRNA sequences encoding a plurality of first polypeptides of an adaptive immune receptor heterodimer, and (B) the second oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of second mRNA sequences encoding a plurality of second polypeptides of the adaptive immune receptor heterodimer, and wherein: (I) the first oligonucleotide reverse transcription primer set comprises a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula:

U1/2-B1-X1

[00126] in which U1/2 comprises an oligonucleotide which comprises a first universal adaptor oligonucleotide sequence when B1 is present or a second universal adaptor oligonucleotide sequence when B1 is nothing, B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides, and X1 comprises an oligonucleotide that is one of: (a) a polynucleotide comprising at least 20, 30, 40 or 50 and not more than 100, 90, 80, 70 or 60 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence, and (b) a polynucleotide comprising at least 15-30 or 31-50 and not more than 80, 70, 60 or 55 contiguous nucleotides of either (i) an adaptive immune receptor joining (J) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence for said first polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence, and (II) the second oligonucleotide reverse transcription primer set comprises a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula:

[00127] U3/4-B2-X2

[00128] in which U3/4 comprises an oligonucleotide which comprises a third universal adaptor oligonucleotide sequence when B2 is present or a fourth universal adaptor oligonucleotide sequence when B2 is nothing, B2 comprises an oligonucleotide that comprises either nothing or a second oligonucleotide barcode sequence of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides that is, for each of the first and second reverse transcription primer sets that are contacted with a single one of the plurality of containers, the same as B1, and X2 comprises an oligonucleotide that is one of: (a) a polynucleotide comprising at least 20, 30, 40 or 50 and not more than 100, 90, 80, 70 or 60 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general

formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence, and (b) a polynucleotide comprising at least 15-30 or 31-50 and not more than 80, 70, 60 or 55 contiguous nucleotides of either (i) an adaptive immune receptor joining (J) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence for said second polypeptide of an adaptive immune receptor heterodimer, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence, said step of contacting taking place under conditions and for a time sufficient to obtain in each of one or more of said plurality of containers: a first reverse-transcribed complementary DNA (cDNA) product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and also to obtain in each of one or more of said plurality of containers: a second reverse-transcribed cDNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer.

[00129] After the step of contacting, there is performed a step of combining the first and second reverse-transcribed cDNA products from the plurality of containers to obtain a mixture of reverse-transcribed cDNA products.

[00130] The combining step is followed by contacting the mixture of first and second reverse-transcribed cDNA products with a first oligonucleotide amplification primer set and a second oligonucleotide amplification primer set, wherein the first amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor

oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence, and wherein the second oligonucleotide amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence, said step of contacting taking place under conditions and for a time sufficient to amplify both of the first and second reverse-transcribed cDNA products, to obtain a DNA library for sequencing.

[00131] Once the DNA library for sequencing has been so obtained, in a step which follows there takes place the sequencing of the DNA library, to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

[00132] Analysis of the data set of sequences may then proceed essentially as described elsewhere herein, to determine rearranged DNA sequences encoding first and second polypeptides of an adaptive immune receptor heterodimer that originate in a single (*i.e.*, the same) lymphoid cell. Briefly, the method may further comprise the steps of: (a) sorting the data set of sequences according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode; (b) sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset; (c) clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets; (d) identifying each first and second adaptive immune receptor

heterodimer polypeptide encoding sequence based on known X1 and X2 sequences, wherein each X1 sequence and each X2 sequence is associated with one or a plurality of unique B sequences to identify the container from which each B sequence-associated X1 sequence and each B sequence-associated X2 sequence originated; and (e) combinatorically matching B sequence-associated X1 and X2 sequences of (d) as being of common clonal origin based on a probability of B sequences that are coincident with common first and second adaptive immune receptor heterodimer polypeptide encoding sequences, and therefrom determining that rearranged DNA sequences encoding first and second polypeptide sequences of the adaptive immune receptor heterodimer originated in a single lymphoid cell.

[00133] Accordingly and in summary, in certain of the herein disclosed embodiments, sequencing adapters may be put onto each end of all reverse transcribed/ amplified TCR and/or IG encoding segments, for instance, by synthesizing universal adaptor sequences onto each end of each cDNA molecule outside of the well-specific barcode. Then, the adapters can be synthesized onto each molecule in a tailing PCR reaction. In such embodiments, fusion RT primers may be synthesized and used for the first cDNA strand synthesis. These primers will all contain the same unique DNA barcode, as well as universal (*e.g.*, pGEX) priming sites. Upon completion of the first cDNA strand synthesis by reverse transcription, the contents of all plate wells will be recovered in a quantitative manner and pooled (*e.g.*, by an inverted centrifugation onto a trough), purified and consequently split into a multiplicity of wells for PCR with universal adapter primers (pGEX) containing “tail” sequences designed to incorporate sequences to be used for amplification and sequencing using a next-generation sequence analysis system (*e.g.*, Illumina, San Diego, CA). Alternatively, the sequencing platform specific adapters can be ligated onto the ends of tagged molecules (*e.g.*, Illumina TrueSeq™ sample preparation method). The molecules from all the wells are pooled thus generating a high-complexity sequencing library of uniquely tagged BCR or TCR ds-cDNA products. The molecules are all sequenced using high-throughput sequencing.

[00134] Universal sequencing primers, complementary to the sequencing platform-specific adapters may desirably be used. This will allow sample indexing of multiple samples, where a sample specific index will be used for each pool of uniquely tagged IGH / TCR products, originating from 96, 384, 1536 etc. original RT reaction wells. Or, a multiplex PCR with a mix of a universal UAI-Forward/ multiplex V, J or C reverse primers may be used to amplify specific target fragments while preserving the original cell transcripts barcoding. If

the Illumina sequencing platform (MiSeq™) is used, a paired end sequencing of 2x 250bp would span the majority of the whole BCR / TCR heavy and light (alpha / beta; gamma / delta) chain sequences, thus allowing recovery of the whole coding sequence of each receptor domain. Alternatively, sequencing platforms with extended read length (Roche 454, Life Ion Torrent, OGT etc.) may be used to read through all library fragments in a single sequencing read in one direction. After sequencing, the reads from each sample may be demultiplexed, provided that more than one sample were in the same sequencing lane. Demultiplexing may be performed by assigning sequencing reads to one of multiple indexes used as part of the universal sequencing adapters. For each sample demultiplexed sequence reads, all reads may be divided by the well specific barcodes. Each set of reads with a specific barcode may be clustered separately to correct PCR and sequencing errors and determine the unique sequences for each barcode:

[00135] Sequences that have been so sorted by barcode and by TCR or IG chain may be further subject to cluster analysis using any of a known variety of algorithms for clustering (*e.g.*, BLASTClust, UCLUST, CD-HIT) and error correction in the case of sequences that fail to cluster with other sequences having shared barcode sequences but which instead would cluster with sequences having a barcode that differs by a single nucleotide. The unique sequences can be identified as IG heavy or light (kappa or lambda) chain, or as TCR (alpha or beta; gamma or delta) chains, by sequence match to known receptor sequences. Each heavy and light chain sequence may thus be associated with a list of barcodes corresponding to an original sample well position. The data can then be reordered by sequence. Associated to each unique sequence will be the set of multi-well plate well-specific barcodes within which set that sequence is found. For every B or T cell clone, the heavy and light chain sequences may be associated with the barcodes from all the wells for which one or more copies of the clone is present. Combinatorics may then be used to match heavy and light chains from the same clone. For example, in a 96 well plate, if particular heavy and light chain sequences are both associated with the same 12 barcodes, this particular pair of heavy and light chains may be assumed to have originated from the same clone, insofar as the probability of two sequences randomly having the exact same 12 barcodes out of 96 is infinitesimally small.

[00136] Exemplary Algorithm: It will be appreciated that according to non-limiting theory, first and second adaptive immune receptor chain encoding sequences that occur with the same set of barcode sequences have a high probability of having originated from the same

plate well, and thus from the same source cell. For example, where 10^3 different barcodes are used in the construction of the first and second oligonucleotide reverse transcription primer sets, the probability that two independent (*i.e.*, originating from different cells) double-stranded cDNA first and second products would be obtained having the same barcode sequence is one in 10^6 , if one cell per each plate well were sorted.

[00137] Hence, if according to the methods described herein, three or more copies of a given set of first and second adaptive immune receptor polypeptide encoding sequences (*e.g.*, X1 and X2) share common barcode sequences (*e.g.*, belong to the same barcode sequence set), the probability that the sequences are of independent cellular origin approaches zero.

[00138] In certain embodiments barcode oligonucleotides B (B1, B2) may optionally comprise a first and a second oligonucleotide barcode sequence, wherein the first barcode sequence is selected to identify uniquely a particular V oligonucleotide sequence and the second barcode sequence is selected to identify uniquely a particular J oligonucleotide sequence. The relative positioning of the barcode oligonucleotides B1 and B2 and universal adaptors (U) advantageously permits rapid identification and quantification of the amplification products of a given unique template oligonucleotide by short sequence reads and paired-end sequencing on automated DNA sequencers (*e.g.*, Illumina HiSeq™ or Illumina MiSeq®, or GeneAnalyzer™-2, Illumina Corp., San Diego, CA). In particular, these and related embodiments permit rapid high-throughput determination of specific combinations of a V and a J sequence that are present in an amplification product, thereby to characterize the relative representation of annealing targets for each combination of a V-specific primer and a J-specific primer that may be present in a sample such as a sample comprising rearranged TCR or BCR encoding DNA. Verification of the identities and/or quantities of the amplification products may be accomplished by longer sequence reads.

[00139] A large number of adaptive immune receptor variable (V) region and joining (J) region gene sequences are known as nucleotide and/or amino acid sequences, including non-rearranged genomic DNA sequences of TCR and Ig loci, and productively rearranged DNA sequences at such loci and their encoded products. See, *e.g.*, U.S.A.N. 13/217,126; U.S.A.N. 12/794,507; PCT/US2011/026373; PCT/US2011/049012. These and other sequences known to the art may be used according to the present disclosure for the design and production of oligonucleotides to be included in the presently provided compositions and methods.

[00140] V region-specific oligonucleotides may include a polynucleotide sequence of at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200,

210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400 or 450 and not more than 1000, 900, 800, 700, 600 or 500 contiguous nucleotides of an adaptive immune receptor (*e.g.*, TCR or BCR) variable (V) region gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences V comprises a unique oligonucleotide sequence. Genomic sequences for TCR and BCR V region genes of humans and other species are known and available from public databases such as Genbank; V region gene sequences include polynucleotide sequences that encode the products of expressed, rearranged TCR and BCR genes and also include polynucleotide sequences of pseudogenes that have been identified in the V region loci. The diverse V polynucleotide sequences that may be incorporated into the presently disclosed oligonucleotides may vary widely in length, in nucleotide composition (*e.g.*, GC content), and in actual linear polynucleotide sequence, and are known, for example, to include “hot spots” or hypervariable regions that exhibit particular sequence diversity.

[00141] The polynucleotide V may thus includes sequences to which members of oligonucleotide primer sets specific for TCR or BCR genes can specifically anneal. Primer sets that are capable of amplifying rearranged DNA encoding a plurality of TCR or BCR are described, for example, in U.S.A.N. 13/217,126; U.S.A.N. 12/794,507; PCT/US2011/026373; or PCT/US2011/049012; or the like; or as described therein may be designed to include oligonucleotide sequences that can specifically hybridize to each unique V gene and to each J gene in a particular TCR or BCR gene locus (*e.g.*, TCRA, TCRB, TCRG, TCRD, IGH, IGK or IGL). For example by way of illustration and not limitation, an oligonucleotide primer of an oligonucleotide primer amplification set that is capable of amplifying rearranged DNA encoding one or a plurality of TCR or BCR may typically include a nucleotide sequence of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 contiguous nucleotides, or more, and may specifically anneal to a complementary sequence of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 contiguous nucleotides of a V or a J polynucleotide as provided herein. In certain embodiments the primers may comprise at least 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides, and in certain embodiment the primers may comprise sequences of no more than 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 contiguous nucleotides. Primers and primer annealing sites of other lengths are also expressly contemplated, as disclosed herein.

[00142] The V polynucleotide may thus, in certain embodiments, comprise a nucleotide sequence having a length that is less than, the same or similar to that of the length of a typical V gene from its start codon to its CDR3 encoding region and may, but need not, include a nucleotide sequence that encodes the CDR3 region. In certain preferred embodiments the V polynucleotide includes all or a portion of a CDR3 encoding nucleotide sequence or the complement thereto and CDR3 sequence lengths may vary considerably and have been characterized by several different numbering schemes (*e.g.*, Lefranc, 1999 *The Immunologist* 7:132; Kabat et al., 1991 *In: Sequences of Proteins of Immunological Interest*, NIH Publication 91-3242; Chothia et al., 1987 *J. Mol. Biol.* 196:901; Chothia et al., 1989 *Nature* 342:877; Al-Lazikani et al., 1997 *J. Mol. Biol.* 273:927; see also, *e.g.*, Rock et al., 1994 *J. Exp. Med.* 179:323; Saada et al., 2007 *Immunol. Cell Biol.* 85:323).

[00143] Briefly, the CDR3 region typically spans the polypeptide portion extending from a highly conserved cysteine residue (encoded by the trinucleotide codon TGY; Y = T or C) in the V segment to a highly conserved phenylalanine residue (encoded by TTY) in the J segment of TCRs, or to a highly conserved tryptophan (encoded by TGG) in IGH. More than 90% of natural, productive rearrangements in the TCRB locus have a CDR3 encoding length by this criterion of between 24 and 54 nucleotides, corresponding to between 9 and 17 encoded amino acids. The numbering schemes for CDR3 encoding regions described above denote the positions of the conserved cysteine, phenylalanine and tryptophan codons, and these numbering schemes may also be applied to pseudogenes in which one or more codons encoding these conserved amino acids may have been replaced with a codon encoding a different amino acid. For pseudogenes which do not use these conserved amino acids, the CDR3 length may be defined relative to the corresponding position at which the conserved residue would have been observed absent the substitution, according to one of the established CDR3 sequence position numbering schemes referenced above.

[00144] The polynucleotide J may comprise a polynucleotide comprising at least 15-30, 31-50, 51-60, 61-90, 91-120, or 120-150, and not more than 600, 500, 400, 300 or 200 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences J comprises a unique oligonucleotide sequence. The polynucleotide J (or its complement) includes sequences to which members of oligonucleotide primer sets specific for TCR or BCR genes can specifically anneal. Primer sets that are capable of amplifying rearranged DNA encoding a plurality of TCR or BCR are described, for example, in

U.S.A.N. 13/217,126; U.S.A.N. 12/794,507; PCT/US2011/026373; or PCT/US2011/049012; or the like; or as described therein may be designed to include oligonucleotide sequences that can specifically hybridize to each unique V gene and to each unique J gene in a particular TCR or BCR gene locus (*e.g.*, TCR α , β , γ or δ , or IgH μ , γ , δ , α or ϵ , or IgL κ or λ).

[00145] It may be preferred in certain embodiments that the plurality of J polynucleotides that are present in the herein described primer compositions have lengths that simulate the overall lengths of known, naturally occurring J gene nucleotide sequences. The J region lengths in the herein described templates may differ from the lengths of naturally occurring J gene sequences by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 percent. The J polynucleotide may thus, in certain embodiments, comprise a nucleotide sequence having a length that is the same or similar to that of the length of a typical naturally occurring J gene and may, but need not, include a nucleotide sequence that encodes the CDR3 region, as discussed above.

[00146] Genomic sequences for TCR and BCR J region genes of humans and other species are known and available from public databases such as Genbank; J region gene sequences include polynucleotide sequences that encode the products of expressed and unexpressed rearranged TCR and BCR genes. The diverse J polynucleotide sequences that may be incorporated into the presently disclosed primers may vary widely in length, in nucleotide composition (*e.g.*, GC content), and in actual linear polynucleotide sequence.

[00147] Alternatives to the V and J sequences described herein, for use in construction of the herein described V-segment and J-segment oligonucleotide primers, may be selected by a skilled person based on the present disclosure using knowledge in the art regarding published gene sequences for the V- and J-encoding regions of the genes for each TCR and Ig subunit. Reference Genbank entries for human adaptive immune receptor sequences include: TCR α : (TCRA/D): NC_000014.8 (chr14:22090057..23021075); TCR β : (TCRB): NC_000007.13 (chr7:141998851..142510972); TCR γ : (TCRG): NC_000007.13 (chr7:38279625..38407656); immunoglobulin heavy chain, IgH (IGH): NC_000014.8 (chr14: 106032614..107288051); immunoglobulin light chain-kappa, IgL κ (IGK): NC_000002.11 (chr2: 89156874..90274235); and immunoglobulin light chain-lambda, IgL λ (IGL): NC_000022.10 (chr22: 22380474..23265085). Reference Genbank entries for mouse adaptive immune receptor loci sequences include: TCR β : (TCRB): NC_000072.5 (chr6: 40841295..41508370), and immunoglobulin heavy chain, IgH (IGH): NC_000078.5 (chr12:114496979..117248165).

[00148] Primer design analyses and target site selection considerations can be performed, for example, using the OLIGO primer analysis software and/or the BLASTN 2.0.5 algorithm software (Altschul et al., *Nucleic Acids Res.* 1997, 25(17):3389-402), or other similar programs available in the art.

[00149] Accordingly, based on the present disclosure and in view of these known adaptive immune receptor gene sequences and oligonucleotide design methodologies, for inclusion in the instant oligonucleotides those skilled in the art can design a plurality of V region-specific and J region-specific polynucleotide sequences that each independently contain oligonucleotide sequences that are unique to a given V and J gene, respectively. Similarly, from the present disclosure and in view of known adaptive immune receptor sequences, those skilled in the art can also design a primer set comprising a plurality of V region-specific and J region-specific oligonucleotide primers that are each independently capable of annealing to a specific sequence that is unique to a given V and J gene, respectively, whereby the plurality of primers is capable of amplifying substantially all V genes and substantially all J genes in a given adaptive immune receptor-encoding locus (*e.g.*, a human TCR or IGH locus). Such primer sets permit generation, in multiplexed (*e.g.*, using multiple forward and reverse primer pairs) PCR, of amplification products that have a first end that is encoded by a rearranged V region-encoding gene segment and a second end that is encoded by a J region-encoding gene segment.

[00150] Typically and in certain embodiments, such amplification products may include a CDR3-encoding sequence although the invention is not intended to be so limited and contemplates amplification products that do not include a CDR3-encoding sequence. The primers may be preferably designed to yield amplification products having sufficient portions of V and J sequences and in certain preferred embodiments also of barcode (B) sequences as described herein, such that by sequencing the products (amplicons), it is possible to identify on the basis of sequences that are unique to each gene segment (i) the particular V gene, and (ii) the particular J gene in the proximity of which the V gene underwent rearrangement to yield a rearranged adaptive immune receptor-encoding gene. Typically, and in preferred embodiments, the PCR amplification products will not be more than 600 base pairs in size, which according to non-limiting theory will exclude amplification products from non-rearranged adaptive immune receptor genes. In certain other preferred embodiments the amplification products will not be more than 500, 400, 300, 250, 200, 150, 125, 100, 90, 80,

70, 60, 50, 40, 30 or 20 base pairs in size, such as may advantageously provide rapid, high-throughput quantification of sequence-distinct amplicons by short sequence reads.

Primers

[00151] According to the present disclosure, oligonucleotide primers are provided in an oligonucleotide primer set that comprises a plurality of V-segment primers and a plurality of J-segment primers, where the primer set is capable of amplifying rearranged DNA encoding adaptive immune receptors in a biological sample that comprises lymphoid cell DNA.

Suitable primer sets are known in the art and disclosed herein, for example, the primer sets in US 2012/0058902, U.S.A.N. 13/217,126; U.S.A.N. 12/794,507; PCT/US2011/026373; or PCT/US2011/049012; or the like; or those shown in Table 1. In certain embodiments the primer set is designed to include a plurality of V sequence-specific primers that includes, for each unique V region gene (including pseudogenes) in a sample, at least one primer that can specifically anneal to a unique V region sequence; and for each unique J region gene in the sample, at least one primer that can specifically anneal to a unique J region sequence.

[00152] Primer design may be achieved by routine methodologies in view of known TCR and BCR genomic sequences. Accordingly, the primer set is preferably capable of amplifying every possible V-J combination that may result from DNA rearrangements in the TCR or BCR locus. As also described below, certain embodiments contemplate primer sets in which one or more V primers may be capable of specifically annealing to a “unique” sequence that may be shared by two or more V regions but that is not common to all V regions, and/or in which one or more J primers may be capable of specifically annealing to a “unique” sequence that may be shared by two or more J regions but that is not common to all J regions.

[00153] In particular embodiments, oligonucleotide primers for use in the compositions and methods described herein may comprise or consist of a nucleic acid of at least about 15 nucleotides long that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence of the target V- or J- segment (*i.e.*, portion of genomic polynucleotide encoding a V-region or J-region polypeptide). Longer primers, *e.g.*, those of about 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, or 50, nucleotides long that have the same sequence as, or sequence complementary to, a contiguous sequence of the target V- or J- region encoding polynucleotide segment, will also be of use in certain embodiments. All intermediate lengths of the presently described oligonucleotide primers are contemplated for use herein. As would be recognized by the

skilled person, the primers may have additional sequence added (*e.g.*, nucleotides that may not be the same as or complementary to the target V- or J-region encoding polynucleotide segment), such as restriction enzyme recognition sites, adaptor sequences for sequencing, barcode sequences, and the like (see *e.g.*, primer sequences provided in the Tables and sequence listing herein). Therefore, the length of the primers may be longer, such as about 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 80, 85, 90, 95, 100 or more nucleotides in length or more, depending on the specific use or need.

[00154] Also contemplated for use in certain embodiments are adaptive immune receptor V-segment or J-segment oligonucleotide primer variants that may share a high degree of sequence identity to the oligonucleotide primers for which nucleotide sequences are presented herein, including those set forth in the Sequence Listing. Thus, in these and related embodiments, adaptive immune receptor V-segment or J-segment oligonucleotide primer variants may have substantial identity to the adaptive immune receptor V-segment or J-segment oligonucleotide primer sequences disclosed herein, for example, such oligonucleotide primer variants may comprise at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity compared to a reference polynucleotide sequence such as the oligonucleotide primer sequences disclosed herein, using the methods described herein (*e.g.*, BLAST analysis using standard parameters). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding ability of an oligonucleotide primer variant to anneal to an adaptive immune receptor segment-encoding polynucleotide by taking into account codon degeneracy, reading frame positioning and the like.

[00155] Typically, oligonucleotide primer variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the annealing ability of the variant oligonucleotide is not substantially diminished relative to that of an adaptive immune receptor V-segment or J-segment oligonucleotide primer sequence that is specifically set forth herein.

[00156] Table 2 presents as a non-limiting example an oligonucleotide primer set that is capable of amplifying productively rearranged DNA encoding TCR β -chains (TCRB) in a biological sample that comprises DNA from lymphoid cells of a subject. In this primer set the J segment primers share substantial sequence homology, and therefore may cross-prime amongst more than one target J polynucleotide sequence, but the V segment primers are designed to anneal specifically to target sequences within the CDR2 region of V and are

therefore unique to each V segment. An exception, however, is present in the case of several V primers where the within-family sequences of the closely related target genes are identical (e.g., V6-2 and V6-3 are identical at the nucleotide level throughout the coding sequence of the V segment, and therefore may have a single primer, TRB2V6-2/3).

Table 2. Exemplary Oligonucleotide Primer Set
(hsTCRB PCR Primers)

Name	Sequence	SEQ ID NO:
TRBJ1-1	TTACCTACAACCTGTGAGTCTGGTGCCTTGTCCAAA	1631
TRBJ1-2	ACCTACAACGGTTAACCTGGTCCCCGAACCGAA	1632
TRBJ1-3	ACCTACAACAGTGAGCCAACTTCCCTCTCCAAA	1633
TRBJ1-4	CCAAGACAGAGAGCTGGGTTCCTACTGCCAAA	1634
TRBJ1-5	ACCTAGGATGGAGAGTCGAGTCCCATCACCAAAA	1635
TRBJ1-6	CTGTCACAGTGAGCCTGGTCCCGTTCCCAAAA	1636
TRBJ2-1	CGGTGAGCCGTGTCCCTGGCCCGAA	1637
TRBJ2-2	CCAGTACGGTCAGCCTAGAGCCTTCTCCAAA	1638
TRBJ2-3	ACTGTCAGCCGGGTGCCTGGGCGCAAAA	1639
TRBJ2-4	AGAGCCGGGTCCCGGCGCCGAA	1640
TRBJ2-5	GGAGCCGCGTGCCTGGCCCGAA	1641
TRBJ2-6	GTCAGCCTGCTGCCGGCCCCGAA	1642
TRBJ2-7	GTGAGCCTGGTGCCCCGGCCCCGAA	1643
TRB2V10-1	AACAAAGGAGAAGTCTCAGATGGCTACAG	1644
TRB2V10-2	GATAAAGGAGAAGTCCCCGATGGCTATGT	1645
TRB2V10-3	GACAAAGGAGAAGTCTCAGATGGCTATAG	1646
TRB2V6-2/3	GCCAAAGGAGAGGTCCCTGATGGCTACAA	1647
TRB2V6-8	CTCTAGATTAAACACAGAGGATTTCCAC	1648
TRB2V6-9	AAGGAGAAGTCCCCGATGGCTACAATGTA	1649
TRB2V6-5	AAGGAGAAGTCCCCAATGGCTACAATGTC	1650
TRB2V6-6	GACAAAGGAGAAGTCCCGAATGGCTACAAC	1651
TRB2V6-7	GTTCCAATGGCTACAATGTCTCCAGATC	1652

Name	Sequence	SEQ ID NO:
TRB2V6-1	GTCCCCAATGGCTACAATGTCTCCAGATT	1653
TRB2V6-4	GTCCCTGATGGTTATAGTGTCTCCAGAGC	1654
TRB2V24-1	ATCTCTGATGGATACAGTGTCTCTCGACA	1655
TRB2V25-1	TTTCCTCTGAGTCAACAGTCTCCAGAATA	1656
TRB2V27	TCCTGAAGGGTACAAAGTCTCTCGAAAAG	1657
TRB2V26	CTCTGAGAGGTATCATGTTTCTTGAAATA	1658
TRB2V28	TCCTGAGGGGTACAGTGTCTCTAGAGAGA	1659
TRB2V19	TATAGCTGAAGGGTACAGCGTCTCTCGGG	1660
TRB2V4-1	CTGAATGCCCCAACAGCTCTCTCTTAAAC	1661
TRB2V4-2/3	CTGAATGCCCCAACAGCTCTCACTTATTC	1662
TRB2V2P	CCTGAATGCCCTGACAGCTCTCGCTTATA	1663
TRB2V3-1	CCTAAATCTCCAGACAAAGCTCACTTAAA	1664
TRB2V3-2	CTCACCTGACTCTCCAGACAAAGCTCAT	1665
TRB2V16	TTCAGCTAAGTGCCTCCCAAATTCACCCT	1666
TRB2V23-1	GATTCTCATCTCAATGCCCCAAGAACGC	1667
TRB2V18	ATTTTCTGCTGAATTTCCCAAAGAGGGCC	1668
TRB2V17	ATTCACAGCTGAAAGACCTAACGGAACGT	1669
TRB2V14	TCTTAGCTGAAAGGACTGGAGGGACGTAT	1670
TRB2V2	TTCGATGATCAATTCTCAGTTGAAAGGCC	1671
TRB2V12-1	TTGATTCTCAGCACAGATGCCTGATGT	1672
TRB2V12-2	GCGATTCTCAGCTGAGAGGCCTGATGG	1673
TRB2V12-3/4	TCGATTCTCAGCTAAGATGCCTAATGC	1674
TRB2V12-5	TTCTCAGCAGAGATGCCTGATGCAACTTTA	1675
TRB2V7-9	GGTTCTCTGCAGAGAGGCCTAAGGGATCT	1676
TRB2V7-8	GCTGCCCAGTGATCGCTTCTTTGCAGAAA	1677
TRB2V7-4	GGCGGCCCCAGTGGTCGGTTCTCTGCAGAG	1678
TRB2V7-6/7	ATGATCGGTTCTCTGCAGAGAGGCCTGAGG	1679
TRB2V7-2	AGTGATCGCTTCTCTGCAGAGAGGACTGG	1680
TRB2V7-3	GGCTGCCCAACGATCGGTTCTTTGCAGT	1681

Name	Sequence	SEQ ID NO:
TRB2V7-1	TCCCCGTGATCGGTTCTCTGCACAGAGGT	1682
TRB2V11-123	CTAAGGATCGATTTTCTGCAGAGAGGCTC	1683
TRB2V13	CTGATCGATTCTCAGCTCAACAGTTCAGT	1684
TRB2V5-1	TGGTCGATTCTCAGGGCGCCAGTTCTCTA	1685
TRB2V5-3	TAATCGATTCTCAGGGCGCCAGTTCCATG	1686
TRB2V5-4	TCCTAGATTCTCAGGTCTCCAGTTCCCTA	1687
TRB2V5-8	GGAAACTTCCCTCCTAGATTTTCAGGTCG	1688
TRB2V5-5	AAGAGGAAACTTCCCTGATCGATTCTCAGC	1689
TRB2V5-6	GGCAACTTCCCTGATCGATTCTCAGGTCA	1690
TRB2V9	GTTCCCTGACTTGCACTCTGAACTAAAC	1691
TRB2V15	GCCGAACACTTCTTTCTGCTTTCTTGAC	1692
TRB2V30	GACCCCAGGACCGGCAGTTCATCCTGAGT	1693
TRB2V20-1	ATGCAAGCCTGACCTTGTCCTACTCTGACA	1694
TRB2V29-1	CATCAGCCGCCCAAACCTAACATTCTCAA	1695

[00157] In certain preferred embodiments, the V-segment and J-segment oligonucleotide primers as described herein are designed to include nucleotide sequences such that adequate information is present within the sequence of an amplification product of a rearranged adaptive immune receptor (TCR or Ig) gene to identify uniquely both the specific V and the specific J genes that give rise to the amplification product in the rearranged adaptive immune receptor locus (*e.g.*, at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 base pairs of sequence upstream of the V gene recombination signal sequence (RSS), preferably at least about 22, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 39 or 40 base pairs of sequence upstream of the V gene recombination signal sequence (RSS), and in certain preferred embodiments greater than 40 base pairs of sequence upstream of the V gene recombination signal sequence (RSS), and at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 base pairs downstream of the J gene RSS, preferably at least about 22, 24, 26, 28 or 30 base pairs downstream of the J gene RSS, and in certain preferred embodiments greater than 30 base pairs downstream of the J gene RSS).

[00158] This feature stands in contrast to oligonucleotide primers described in the art for amplification of TCR-encoding or Ig-encoding gene sequences, which rely primarily on the amplification reaction merely for detection of presence or absence of products of appropriate sizes for V and J segments (*e.g.*, the presence in PCR reaction products of an amplicon of a particular size indicates presence of a V or J segment but fails to provide the sequence of the amplified PCR product and hence fails to confirm its identity, such as the common practice of spectratyping).

[00159] Oligonucleotides (*e.g.*, primers) can be prepared by any suitable method, including direct chemical synthesis by a method such as the phosphotriester method of Narang et al., 1979, *Meth. Enzymol.* 68:90-99; the phosphodiester method of Brown et al., 1979, *Meth. Enzymol.* 68:109-151; the diethylphosphoramidite method of Beaucage et al., 1981, *Tetrahedron Lett.* 22:1859-1862; and the solid support method of U.S. Pat. No. 4,458,066, each incorporated herein by reference. A review of synthesis methods of conjugates of oligonucleotides and modified nucleotides is provided in Goodchild, 1990, *Bioconjugate Chemistry* 1(3): 165-187, incorporated herein by reference.

[00160] The term "primer," as used herein, refers to an oligonucleotide capable of acting as a point of initiation of DNA synthesis under suitable

[00161] conditions. Such conditions include those in which synthesis of a primer extension product complementary to a nucleic acid strand is induced in the presence of four different nucleoside triphosphates and an agent for extension (*e.g.*, a DNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature.

[00162] A primer is preferably a single-stranded DNA. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 6 to 50 nucleotides, or in certain embodiments, from 15-35 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template nucleic acid, but must be sufficiently complementary to hybridize with the template. The design of suitable primers for the amplification of a given target sequence is well known in the art and described in the literature cited herein.

[00163] As described herein, primers can incorporate additional features which allow for the detection or immobilization of the primer but do not alter the basic property of the primer, that of acting as a point of initiation of DNA synthesis. For example, primers may contain an additional nucleic acid sequence at the 5' end which does not hybridize to the target nucleic

acid, but which facilitates cloning, detection, or sequencing of the amplified product. The region of the primer which is sufficiently complementary to the template to hybridize is referred to herein as the hybridizing region.

[00164] As used herein, a primer is "specific," for a target sequence if, when used in an amplification reaction under sufficiently stringent conditions, the primer hybridizes primarily to the target nucleic acid. Typically, a primer is specific for a target sequence if the primer-target duplex stability is greater than the stability of a duplex formed between the primer and any other sequence found in the sample. One of skill in the art will recognize that various factors, such as salt conditions as well as base composition of the primer and the location of the mismatches, will affect the specificity of the primer, and that routine experimental confirmation of the primer specificity will be needed in many cases. Hybridization conditions can be chosen under which the primer can form stable duplexes only with a target sequence. Thus, the use of target-specific primers under suitably stringent amplification conditions enables the selective amplification of those target sequences which contain the target primer binding sites.

[00165] In particular embodiments, primers for use in the methods described herein comprise or consist of a nucleic acid of at least about 15 nucleotides long that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence of the target V or J segment. Longer primers, *e.g.*, those of about 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, or 50, nucleotides long that have the same sequence as, or sequence complementary to, a contiguous sequence of the target V or J segment, will also be of use in certain embodiments. All intermediate lengths of the aforementioned primers are contemplated for use herein. As would be recognized by the skilled person, the primers may have additional sequence added (*e.g.*, nucleotides that may not be the same as or complementary to the target V or J segment), such as restriction enzyme recognition sites, adaptor sequences for sequencing, barcode sequences, and the like (see *e.g.*, primer sequences provided herein and in the sequence listing). Therefore, the length of the primers may be longer, such as 55, 56, 57, 58, 59, 60, 65, 70, 75, nucleotides in length or more, depending on the specific use or need. For example, in one embodiment, the forward and reverse primers are both modified at the 5' end with the universal forward primer sequence compatible with a DNA sequencer.

[00166] Also contemplated for use in certain embodiments are adaptive immune receptor V-segment or J-segment oligonucleotide primer variants that may share a high degree of

sequence identity to the oligonucleotide primers for which nucleotide sequences are presented herein, including those set forth in the Sequence Listing. Thus, in these and related embodiments, adaptive immune receptor V-segment or J-segment oligonucleotide primer variants may have substantial identity to the adaptive immune receptor V-segment or J-segment oligonucleotide primer sequences disclosed herein, for example, such oligonucleotide primer variants may comprise at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity compared to a reference polynucleotide sequence such as the oligonucleotide primer sequences disclosed herein, using the methods described herein (*e.g.*, BLAST analysis using standard parameters). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding ability of an oligonucleotide primer variant to anneal to an adaptive immune receptor segment-encoding polynucleotide by taking into account codon degeneracy, reading frame positioning and the like.

[00167] Typically, oligonucleotide primer variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the annealing ability of the variant oligonucleotide is not substantially diminished relative to that of an adaptive immune receptor V-segment or J-segment oligonucleotide primer sequence that is specifically set forth herein. As also noted elsewhere herein, in preferred embodiments adaptive immune receptor V-segment and J-segment oligonucleotide primers are designed to be capable of amplifying a rearranged TCR or IGH sequence that includes the coding region for CDR3.

[00168] According to certain embodiments contemplated herein, the primers for use in the multiplex PCR methods of the present disclosure may be functionally blocked to prevent non-specific priming of non-T or B cell sequences. For example, the primers may be blocked with chemical modifications as described in U.S. patent application publication US2010/0167353. According to certain herein disclosed embodiments, the use of such blocked primers in the present multiplex PCR reactions involves primers that may have an inactive configuration wherein DNA replication (*i.e.*, primer extension) is blocked, and an activated configuration wherein DNA replication proceeds. The inactive configuration of the primer is present when the primer is either single-stranded, or when the primer is specifically hybridized to the target DNA sequence of interest but primer extension remains blocked by a chemical moiety that is linked at or near to the 3' end of the primer.

[00169] The activated configuration of the primer is present when the primer is hybridized to the target nucleic acid sequence of interest and is subsequently acted upon by RNase H or

another cleaving agent to remove the 3' blocking group, thereby allowing an enzyme (*e.g.*, a DNA polymerase) to catalyze primer extension in an amplification reaction. Without wishing to be bound by theory, it is believed that the kinetics of the hybridization of such primers are akin to a second order reaction, and are therefore a function of the T cell or B cell gene sequence concentration in the mixture. Blocked primers minimize non-specific reactions by requiring hybridization to the target followed by cleavage before primer extension can proceed. If a primer hybridizes incorrectly to a sequence that is related to the desired target sequence but which differs by having one or more non-complementary nucleotides that result in base-pairing mismatches, cleavage of the primer is inhibited, especially when there is a mismatch that lies at or near the cleavage site. This strategy to improve the fidelity of amplification reduces the frequency of false priming at such locations, and thereby increases the specificity of the reaction. As would be recognized by the skilled person, reaction conditions, particularly the concentration of RNase H and the time allowed for hybridization and extension in each cycle, can be optimized to maximize the difference in cleavage efficiencies between highly efficient cleavage of the primer when it is correctly hybridized to its true target sequence, and poor cleavage of the primer when there is a mismatch between the primer and the template sequence to which it may be incompletely annealed.

[00170] As described in US2010/0167353, a number of blocking groups are known in the art that can be placed at or near the 3' end of the oligonucleotide (*e.g.*, a primer) to prevent extension. A primer or other oligonucleotide may be modified at the 3'-terminal nucleotide to prevent or inhibit initiation of DNA synthesis by, for example, the addition of a 3' deoxyribonucleotide residue (*e.g.*, cordycepin), a 2',3'-dideoxyribonucleotide residue, non-nucleotide linkages or alkane-diol modifications (U.S. Pat. No. 5,554,516). Alkane diol modifications which can be used to inhibit or block primer extension have also been described by Wilk et al., (1990 *Nucleic Acids Res.* 18 (8):2065), and by Arnold et al. (U.S. Pat. No. 6,031,091). Additional examples of suitable blocking groups include 3' hydroxyl substitutions (*e.g.*, 3'-phosphate, 3'-triphosphate or 3'-phosphate diesters with alcohols such as 3-hydroxypropyl), 2'3'-cyclic phosphate, 2' hydroxyl substitutions of a terminal RNA base (*e.g.*, phosphate or sterically bulky groups such as triisopropyl silyl (TIPS) or tert-butyl dimethyl silyl (TBDMS)). 2'-alkyl silyl groups such as TIPS and TBDMS substituted at the 3'-end of an oligonucleotide are described by Laikhter et al., U.S. patent application Ser. No. 11/686,894, which is incorporated herein by reference. Bulky substituents can also be

incorporated on the base of the 3'-terminal residue of the oligonucleotide to block primer extension.

[00171] In certain embodiments, the oligonucleotide may comprise a cleavage domain that is located upstream (*e.g.*, 5' to) of the blocking group used to inhibit primer extension. As examples, the cleavage domain may be an RNase H cleavage domain, or the cleavage domain may be an RNase H2 cleavage domain comprising a single RNA residue, or the oligonucleotide may comprise replacement of the RNA base with one or more alternative nucleosides. Additional illustrative cleavage domains are described in US2010/0167353.

[00172] Thus, a multiplex PCR system may use 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, or more forward primers, wherein each forward primer is complementary to a single functional TCR or Ig V segment or a small family of functional TCR or Ig V segments, *e.g.*, a TCR V β segment, (see *e.g.*, the TCRBV primers as shown in Table 2, SEQ ID NOS:1644-1695), and, for example, thirteen reverse primers, each specific to a TCR or Ig J segment, such as TCR J β segment (see *e.g.*, TCRBJ primers in Table 2, SEQ ID NOS:1631-1643). In another embodiment, a multiplex PCR reaction may use four forward primers each specific to one or more functional TCR γ V segment and four reverse primers each specific for one or more TCR γ J segments. In another embodiment, a multiplex PCR reaction may use 84 forward primers each specific to one or more functional V segments and six reverse primers each specific for one or more J segments.

[00173] Thermal cycling conditions may follow methods of those skilled in the art. For example, using a PCR Express™ thermal cycler (Hybaid, Ashford, UK), the following cycling conditions may be used: 1 cycle at 95°C for 15 minutes, 25 to 40 cycles at 94°C for 30 seconds, 59°C for 30 seconds and 72°C for 1 minute, followed by one cycle at 72°C for 10 minutes. As will be recognized by the skilled person, thermal cycling conditions may be optimized, for example, by modifying annealing temperatures, annealing times, number of cycles and extension times. As would be recognized by the skilled person, the amount of primer and other PCR reagents used, as well as PCR parameters (*e.g.*, annealing temperature, extension times and cycle numbers), may be optimized to achieve desired PCR amplification efficiency.

[00174] Alternatively, in certain related embodiments also contemplated herein, "digital PCR" methods can be used to quantitate the number of target genomes in a sample, without the need for a standard curve. In digital PCR, the PCR reaction for a single sample is performed in a multitude of more than 100 microcells or droplets, such that each droplet

either amplifies (*e.g.*, generation of an amplification product provides evidence of the presence of at least one template molecule in the microcell or droplet) or fails to amplify (evidence that the template was not present in a given microcell or droplet). By simply counting the number of positive microcells, it is possible directly to count the number of target genomes that are present in an input sample. Digital PCR methods typically use an endpoint readout, rather than a conventional quantitative PCR signal that is measured after each cycle in the thermal cycling reaction (see, *e.g.*, Pekin et al., 2011 *Lab. Chip* 11(13):2156; Zhong et al., 2011 *Lab. Chip* 11(13):2167; Tewhey et al., 2009 *Nature Biotechnol.* 27:1025; 2010 *Nature Biotechnol.* 28:178). Accordingly, any of the herein described compositions (*e.g.*, adaptive immune receptor gene-specific oligonucleotide primer sets) and methods may be adapted for use in such digital PCR methodology, for example, the ABI QuantStudio™ 12K Flex System (Life Technologies, Carlsbad, CA), the QuantaLife™ digital PCR system (BioRad, Hercules, CA) or the RainDance™ microdroplet digital PCR system (RainDance Technologies, Lexington, MA).

Adaptors

[00175] The herein described oligonucleotides may in certain embodiments comprise first (U1) and second (U2) (and optionally third (U3) and fourth (U4)) universal adaptor oligonucleotide sequences, or may lack either or both of U1 and U2 (or U3 or U4). A universal adaptor oligonucleotide U thus may comprise either nothing or an oligonucleotide having a sequence that is selected from (i) a first universal adaptor oligonucleotide sequence, and (ii) a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to a first universal adaptor oligonucleotide sequence, and U2 may comprise either nothing or an oligonucleotide having a sequence that is selected from (i) a second universal adaptor oligonucleotide sequence, and (ii) a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to a second universal adaptor oligonucleotide sequence. A similar relationship pertains for U3 and U4.

[00176] U1 and/or U2 may, for example, comprise universal adaptor oligonucleotide sequences and/or sequencing platform-specific oligonucleotide sequences that are specific to a single-molecule sequencing technology being employed, for example the HiSeq™ or GeneAnalyzer™-2 (GA-2) systems (Illumina, Inc., San Diego, CA) or another suitable sequencing suite of instrumentation, reagents and software. Inclusion of such platform-specific adaptor sequences permits direct quantitative sequencing of the presently described dsDNA amplification products into which U has been incorporated as described herein, using

a nucleotide sequencing methodology such as the HiSeq™ or GA2 or equivalent. This feature therefore advantageously permits qualitative and quantitative characterization of the dsDNA composition.

[00177] For example, dsDNA amplification products may be generated that have universal adaptor sequences at both ends, so that the adaptor sequences can be used to further incorporate sequencing platform-specific oligonucleotides at each end of each template.

[00178] Without wishing to be bound by theory, platform-specific oligonucleotides may be added onto the ends of such dsDNA using 5' (5'-platform sequence-universal adaptor-1 sequence-3') and 3' (5'-platform sequence-universal adaptor-2 sequence-3') oligonucleotides in three cycles of denaturation, annealing and extension, so that the relative representation in the dsDNA composition of each of the component dsDNAs is not quantitatively altered.

Unique identifier sequences (*e.g.*, barcode sequences B that are associated with and thus identify individual V and/or J regions, or sample-identifier barcodes as described herein) are placed adjacent to the adaptor sequences, thus permitting quantitative sequencing in short sequence reads, in order to characterize the DNA population by the criterion of the relative amount of each unique sequence that is present.

[00179] In addition to adaptor sequences described in the Examples and included in the exemplary template sequences in the Sequence Listing (*e.g.*, at the 5' and 3' ends of SEQ ID NOS:1-1630), other oligonucleotide sequences that may be used as universal adaptor sequences will be known to those familiar with the art in view of the present disclosure. Non-limiting examples of additional adaptor sequences are shown in Table 3 and set forth in SEQ ID NOS:1710-1731.

Table 3. Exemplary Adaptor Sequences

Adaptor (primer) name	Sequence	SEQ ID NO:
T7 Promotor	AATACGACTCACTATAGG	1710
T7 Terminator	GCTAGTTATTGCTCAGCGG	1711
T3	ATTAACCCTCACTAAAGG	1712
SP6	GATTTAGGTGACACTATAG	1713
M13F(-21)	TGTAAAACGACGGCCAGT	1714
M13F(-40)	GTTTCCAGTCACGAC	1715
M13R Reverse	CAGGAAACAGCTATGACC	1716

Adaptor (primer) name	Sequence	SEQ ID NO:
AOX1 Forward	GACTGGTTCCAATTGACAAGC	1717
AOX1 Reverse	GCAAATGGCATTCTGACATCC	1718
pGEX Forward (GST 5, pGEX 5')	GGGCTGGCAAGCCACGTTTGGTG	1719
pGEX Reverse (GST 3, pGEX 3')	CCGGGAGCTGCATGTGTCAGAGG	1720
BGH Reverse	AACTAGAAGGCACAGTCGAGGC	1721
GFP (C' terminal, CFP, YFP or BFP)	CACTCTCGGCATGGACGAGC	1722
GFP Reverse	TGGTGCAGATGAACTTCAGG	1723
GAG	GTTCGACCCCGCCTCGATCC	1724
GAG Reverse	TGACACACATTCCACAGGGTC	1725
CYC1 Reverse	GCGTGAATGTAAGCGTGAC	1726
pFastBacF	5'-d(GGATTATTCATACCGTCCCA)-3'	1727
pFastBacR	5'-d(CAAATGTGGTATGGCTGATT)-3'	1728
pBAD Forward	5'-d(ATGCCATAGCATTTTATCC)-3'	1729
pBAD Reverse	5'-d(GATTTAATCTGTATCAGG)-3'	1730
CMV-Forward	5'-d(CGCAAATGGGCGGTAGGCGTG)-3'	1731

Barcodes

[00180] As described herein, certain embodiments contemplate designing oligonucleotide sequences to contain short signature sequences that permit unambiguous identification of the polynucleotide sequence into which they are incorporated, and hence of at least one primer responsible for amplifying that product, without having to sequence the entire amplification product. In the herein described oligonucleotides, such barcodes B (*e.g.*, B1, B2) are each either nothing or each comprise an oligonucleotide B that comprises an oligonucleotide barcode sequence of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50 or more contiguous nucleotides (including all integer values therebetween), wherein in each of the plurality of oligonucleotide sequences B comprises a unique oligonucleotide sequence which uniquely identifies a particular V and/or J oligonucleotide primer sequence.

[00181] Exemplary barcodes may comprise a first barcode oligonucleotide of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 nucleotides that uniquely identifies each oligonucleotide primer (*e.g.*, a V or a J primer) in the primer composition, and optionally in certain embodiments a second barcode oligonucleotide of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 nucleotides that uniquely identifies each partner primer in a primer set (*e.g.*, a J or a V primer), to provide barcodes of, respectively, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 or 32 nucleotides in length, but these and related embodiments are not intended to be so limited. Barcode oligonucleotides may comprise oligonucleotide sequences of any length, so long as a minimum barcode length is obtained that precludes occurrence of a given barcode sequence in two or more product polynucleotides having otherwise distinct sequences (*e.g.*, V and J sequences).

[00182] Thus, the minimum barcode length, to avoid such redundancy amongst the barcodes that are used to uniquely identify different V-J sequence pairings, is X nucleotides, where 4^X is greater than the number of distinct template species that are to be differentiated on the basis of having non-identical sequences. In practice, barcode oligonucleotide sequence read lengths may be limited only by the sequence read-length limits of the nucleotide sequencing instrument to be employed. For certain embodiments, different barcode oligonucleotides that will distinguish individual species of template oligonucleotides should have at least two nucleotide mismatches (*e.g.*, a minimum hamming distance of 2) when aligned to maximize the number of nucleotides that match at particular positions in the barcode oligonucleotide sequences.

[00183] The skilled artisan will be familiar with the design, synthesis, and incorporation into a larger oligonucleotide or polynucleotide construct, of oligonucleotide barcode sequences of, for instance, at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35 or more contiguous nucleotides, including all integer values therebetween. For non-limiting examples of the design and implementation of oligonucleotide barcode sequence identification strategies, see, *e.g.*, de Carcer et al., 2011 *Adv. Env. Microbiol.* 77:6310; Parameswaran et al., 2007 *Nucl. Ac. Res.* 35(19):330; Roh et al., 2010 *Trends Biotechnol.* 28:291.

[00184] Typically, barcodes are placed in oligonucleotides at locations where they are not found naturally, *i.e.*, barcodes comprise nucleotide sequences that are distinct from any naturally occurring oligonucleotide sequences that may be found in the vicinity of the sequences adjacent to which the barcodes are situated (*e.g.*, V and/or J sequences). Such

barcode sequences may be included, according to certain embodiments described herein, as elements B1 and/or B2 of the presently disclosed oligonucleotides. Accordingly, certain of the herein described oligonucleotide compositions may in certain embodiments comprise one, two or more barcodes, while in certain other embodiments some or all of these barcodes may be absent. In certain embodiments all barcode sequences will have identical or similar GC content (*e.g.*, differing in GC content by no more than 20%, or by no more than 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10%).

Sequencing

[00185] Sequencing may be performed using any of a variety of available high throughput single molecule sequencing machines and systems. Illustrative sequence systems include sequence-by-synthesis systems such as the Illumina Genome Analyzer and associated instruments (Illumina, Inc., San Diego, CA), Helicos Genetic Analysis System (Helicos BioSciences Corp., Cambridge, MA), Pacific Biosciences PacBio RS (Pacific Biosciences, Menlo Park, CA), or other systems having similar capabilities. Sequencing is achieved using a set of sequencing oligonucleotides that hybridize to a defined region within the amplified DNA molecules. The sequencing oligonucleotides are designed such that the V- and J-encoding gene segments can be uniquely identified by the sequences that are generated, based on the present disclosure and in view of known adaptive immune receptor gene sequences that appear in publicly available databases. See, *e.g.*, U.S.A.N. 13/217,126; U.S.A.N. 12/794,507; PCT/US2011/026373; or PCT/US2011/049012. Exemplary TCRB J-region sequencing primers are set forth in Table 4:

Table 4: TCRBJ Sequencing Primers

PRIMER	SEQUENCE	SEQ ID NO:
>Jseq1-1	ACAACGTGAGTCTGGTGCCTTGTCCAAAGAAA	1696
>Jseq1-2	ACAACGGTTAACCTGGTCCCCGAACCGAAGGTG	1697
>Jseq1-3	ACAACAGTGAGCCAACCTCCCTCTCCAAAATAT	1698
>Jseq1-4	AAGACAGAGAGCTGGGTTCCACTGCCAAAAAAC	1699
>Jseq1-5	AGGATGGAGAGTCGAGTCCCATCACCAAAATGC	1700
>Jseq1-6	GTCACAGTGAGCCTGGTCCCGTTCCCAAAGTGG	1701
>Jseq2-1	AGCACGGTGAGCCGTGTCCCTGGCCCCGAAGAAC	1702
>Jseq2-2	AGTACGGTCAGCCTAGAGCCTTCTCCAAAAAAC	1703
>Jseq2-3	AGCACTGTCAGCCGGGTGCCTGGGCCCCAAATAC	1704
>Jseq2-4	AGCACTGAGAGCCGGGTCCCGGCGCCGAAGTAC	1705
>Jseq2-5	AGCACCAGGAGCCGCGTGCCTGGCCCCGAAGTAC	1706
>Jseq2-6	AGCACGGTCAGCCTGCTGCCGGCCCCGAAAGTC	1707
>Jseq2-7	GTGACCGTGAGCCTGGTGCCCGGCCCCGAAGTAC	1708

[00186] The term “gene” means the segment of DNA involved in producing a polypeptide chain such as all or a portion of a TCR or Ig polypeptide (*e.g.*, a CDR3-containing polypeptide); it includes regions preceding and following the coding region “leader and trailer” as well as intervening sequences (introns) between individual coding segments (exons), and may also include regulatory elements (*e.g.*, promoters, enhancers, repressor binding sites and the like), and may also include recombination signal sequences (RSSs) as described herein.

[00187] The nucleic acids of the present embodiments, also referred to herein as polynucleotides, may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. A coding sequence which encodes a TCR or an immunoglobulin or a region thereof (*e.g.*, a V region, a D segment, a J region, a C region, etc.) for use according to the present embodiments may be identical to the coding sequence known in the art for any given TCR or immunoglobulin gene regions or polypeptide domains (*e.g.*, V-region domains, CDR3 domains, etc.), or may be a different coding sequence, which, as a result of the redundancy or

degeneracy of the genetic code, encodes the same TCR or immunoglobulin region or polypeptide.

[00188] In certain embodiments, the amplified J-region encoding gene segments may each have a unique sequence-defined identifier tag of 2, 3, 4, 5, 6, 7, 8, 9, 10 or about 15, 20 or more nucleotides, situated at a defined position relative to a RSS site. For example, a four-base tag may be used, in the J β -region encoding segment of amplified TCR β CDR3-encoding regions, at positions +11 through +14 downstream from the RSS site. However, these and related embodiments need not be so limited and also contemplate other relatively short nucleotide sequence-defined identifier tags that may be detected in J-region encoding gene segments and defined based on their positions relative to an RSS site. These may vary between different adaptive immune receptor encoding loci.

[00189] The recombination signal sequence (RSS) consists of two conserved sequences (heptamer, 5'-CACAGTG-3', and nonamer, 5'-ACAAAAACC-3'), separated by a spacer of either 12 +/- 1 bp ("12-signal") or 23 +/- 1 bp ("23-signal"). A number of nucleotide positions have been identified as important for recombination including the CA dinucleotide at position one and two of the heptamer, and a C at heptamer position three has also been shown to be strongly preferred as well as an A nucleotide at positions 5, 6, 7 of the nonamer. (Ramsden et. al 1994; Akamatsu et. al. 1994; Hesse et. al. 1989). Mutations of other nucleotides have minimal or inconsistent effects. The spacer, although more variable, also has an impact on recombination, and single-nucleotide replacements have been shown to significantly impact recombination efficiency (Fanning et. al. 1996, Larijani et. al 1999; Nadel et. al. 1998). Criteria have been described for identifying RSS polynucleotide sequences having significantly different recombination efficiencies (Ramsden et. al 1994; Akamatsu et. al. 1994; Hesse et. al. 1989 and Cowell et. al. 1994). Accordingly, the sequencing oligonucleotides may hybridize adjacent to a four base tag within the amplified J-encoding gene segments at positions +11 through +14 downstream of the RSS site. For example, sequencing oligonucleotides for TCRB may be designed to anneal to a consensus nucleotide motif observed just downstream of this "tag", so that the first four bases of a sequence read will uniquely identify the J-encoding gene segment (see, *e.g.*, WO/2012/027503).

[00190] The average length of the CDR3-encoding region, for the TCR, defined as the nucleotides encoding the TCR polypeptide between the second conserved cysteine of the V segment and the conserved phenylalanine of the J segment, is 35+/-3 nucleotides.

Accordingly and in certain embodiments, PCR amplification using V-segment oligonucleotide primers with J-segment oligonucleotide primers that start from the J segment tag of a particular TCR or IgH J region (*e.g.*, TCR J β , TCR J γ or IgH JH as described herein) will nearly always capture the complete V-D-J junction in a 50 base pair read. The average length of the IgH CDR3 region, defined as the nucleotides between the conserved cysteine in the V segment and the conserved phenylalanine in the J segment, is less constrained than at the TCR β locus, but will typically be between about 10 and about 70 nucleotides.

Accordingly and in certain embodiments, PCR amplification using V-segment oligonucleotide primers with J-segment oligonucleotide primers that start from the IgH J segment tag will capture the complete V-D-J junction in a 100 base pair read.

[00191] PCR primers that anneal to and support polynucleotide extension on mismatched template sequences are referred to as promiscuous primers. In certain embodiments, the TCR and Ig J-segment reverse PCR primers may be designed to minimize overlap with the sequencing oligonucleotides, in order to minimize promiscuous priming in the context of multiplex PCR. In one embodiment, the TCR and Ig J-segment reverse primers may be anchored at the 3' end by annealing to the consensus splice site motif, with minimal overlap of the sequencing primers. Generally, the TCR and Ig V and J-segment primers may be selected to operate in PCR at consistent annealing temperatures using known sequence/primer design and analysis programs under default parameters.

[00192] For the sequencing reaction, the exemplary IGHJ sequencing primers extend three nucleotides across the conserved CAG sequences as described in WO/2012/027503.

Samples

[00193] The subject or biological source, from which a test biological sample may be obtained, may be a human or non-human animal, or a transgenic or cloned or tissue-engineered (including through the use of stem cells) organism. In certain preferred embodiments of the invention, the subject or biological source may be known to have, or may be suspected of having or being at risk for having, a circulating or solid tumor or other malignant condition, or an autoimmune disease, or an inflammatory condition, and in certain preferred embodiments of the invention the subject or biological source may be known to be free of a risk or presence of such disease.

[00194] Certain preferred embodiments contemplate a subject or biological source that is a human subject such as a patient that has been diagnosed as having or being at risk for developing or acquiring cancer according to art-accepted clinical diagnostic criteria, such as

those of the U.S. National Cancer Institute (Bethesda, MD, USA) or as described in *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology* (2008, Lippincott, Williams and Wilkins, Philadelphia/ Ovid, New York); Pizzo and Poplack, *Principles and Practice of Pediatric Oncology* (Fourth edition, 2001, Lippincott, Williams and Wilkins, Philadelphia/ Ovid, New York); and Vogelstein and Kinzler, *The Genetic Basis of Human Cancer* (Second edition, 2002, McGraw Hill Professional, New York); certain embodiments contemplate a human subject that is known to be free of a risk for having, developing or acquiring cancer by such criteria.

[00195] Certain other embodiments contemplate a non-human subject or biological source, for example a non-human primate such as a macaque, chimpanzee, gorilla, vervet, orangutan, baboon or other non-human primate, including such non-human subjects that may be known to the art as preclinical models, including preclinical models for solid tumors and/or other cancers. Certain other embodiments contemplate a non-human subject that is a mammal, for example, a mouse, rat, rabbit, pig, sheep, horse, bovine, goat, gerbil, hamster, guinea pig or other mammal; many such mammals may be subjects that are known to the art as preclinical models for certain diseases or disorders, including circulating or solid tumors and/or other cancers (*e.g.*, Talmadge et al., 2007 *Am. J. Pathol.* 170:793; Kerbel, 2003 *Canc. Biol. Therap.* 2(4 Suppl 1):S134; Man et al., 2007 *Canc. Met. Rev.* 26:737; Cespedes et al., 2006 *Clin. Transl. Oncol.* 8:318). The range of embodiments is not intended to be so limited, however, such that there are also contemplated other embodiments in which the subject or biological source may be a non-mammalian vertebrate, for example, another higher vertebrate, or an avian, amphibian or reptilian species, or another subject or biological source.

[00196] Biological samples may be provided by obtaining a blood sample, biopsy specimen, tissue explant, organ culture, biological fluid or any other tissue or cell preparation from a subject or a biological source. Preferably the sample comprises DNA from lymphoid cells of the subject or biological source, which, by way of illustration and not limitation, may contain rearranged DNA at one or more TCR or BCR loci. In certain embodiments a test biological sample may be obtained from a solid tissue (*e.g.*, a solid tumor), for example by surgical resection, needle biopsy or other means for obtaining a test biological sample that contains a mixture of cells.

[00197] According to certain embodiments it may be desirable to isolate lymphoid cells (*e.g.*, T cells and/or B cells) according to any of a large number of established methodologies,

where isolated lymphoid cells are those that have been removed or separated from the tissue, environment or milieu in which they naturally occur. B cells and T cells can thus be obtained from a biological sample, such as from a variety of tissue and biological fluid samples including bone marrow, thymus, lymph glands, lymph nodes, peripheral tissues and blood, but peripheral blood is most easily accessed. Any peripheral tissue can be sampled for the presence of B and T cells and is therefore contemplated for use in the methods described herein. Tissues and biological fluids from which adaptive immune cells, may be obtained include, but are not limited to skin, epithelial tissues, colon, spleen, a mucosal secretion, oral mucosa, intestinal mucosa, vaginal mucosa or a vaginal secretion, cervical tissue, ganglia, saliva, cerebrospinal fluid (CSF), bone marrow, cord blood, serum, serosal fluid, plasma, lymph, urine, ascites fluid, pleural fluid, pericardial fluid, peritoneal fluid, abdominal fluid, culture medium, conditioned culture medium or lavage fluid. In certain embodiments, adaptive immune cells may be isolated from an apheresis sample. Peripheral blood samples may be obtained by phlebotomy from subjects. Peripheral blood mononuclear cells (PBMC) are isolated by techniques known to those of skill in the art, *e.g.*, by Ficoll-Hypaque[®] density gradient separation. In certain embodiments, whole PBMCs are used for analysis.

[00198] For nucleic acid extraction, total genomic DNA may be extracted from cells using methods known in the art and/or commercially available kits, *e.g.*, by using the QIAamp[®] DNA blood Mini Kit (QIAGEN[®]). The approximate mass of a single haploid genome is 3 pg. Preferably, at least 100,000 to 200,000 cells are used for analysis, *i.e.*, about 0.6 to 1.2 μ g DNA from diploid T or B cells. Using PBMCs as a source, the number of T cells can be estimated to be about 30% of total cells. The number of B cells can also be estimated to be about 30% of total cells in a PBMC preparation.

[00199] The Ig and TCR gene loci contain many different variable (V), diversity (D), and joining (J) gene segments, which are subjected to rearrangement processes during early lymphoid differentiation. Ig and TCR V, D and J gene segment sequences are known in the art and are available in public databases such as GENBANK. The V-D-J rearrangements are mediated via a recombinase enzyme complex in which the RAG1 and RAG2 proteins play a key role by recognizing and cutting the DNA at the recombination signal sequences (RSS), which are located downstream of the V gene segments, at both sides of the D gene segments, and upstream of the J gene segments. Inappropriate RSS reduce or even completely prevent rearrangement. The recombination signal sequence (RSS) consists of two conserved

sequences (heptamer, 5'-CACAGTG-3', and nonamer, 5'-ACAAAAACC-3'), separated by a spacer of either 12 +/- 1 bp ("12-signal") or 23 +/- 1 bp ("23-signal").

[00200] A number of nucleotide positions have been identified as important for recombination including the CA dinucleotide at position one and two of the heptamer, and a C at heptamer position three has also been shown to be strongly preferred as well as an A nucleotide at positions 5, 6, 7 of the nonamer. (Ramsden et al. 1994 *Nucl. Ac. Res.* 22:1785; Akamatsu et. al. 1994 *J. Immunol.* 153:4520; Hesse et. al. 1989 *Genes Dev.* 3:1053). Mutations of other nucleotides have minimal or inconsistent effects. The spacer, although more variable, also has an impact on recombination, and single-nucleotide replacements have been shown to significantly impact recombination efficiency (Fanning et. al. 1996 *Cell. Immunol. Immunopath.* 79:1, Larijani et al. 1999 *Nucl. Ac. Res.* 27:2304; Nadel et al. 1998 *J. Immunol.* 161:6068; Nadel et al., 1998 *J. Exp. Med.* 187:1495). Criteria have been described for identifying RSS polynucleotide sequences having significantly different recombination efficiencies (Ramsden et al 1994 *Nucl. Ac. Res.* 22:1785; Akamatsu et. al. 1994 *J. Immunol.* 153:4520; Hesse et al. 1989 *Genes Dev.* 3:1053, and Lee et al., 2003 *PLoS* 1(1):E1).

[00201] The rearrangement process generally starts with a D to J rearrangement followed by a V to D-J rearrangement in the case of Ig heavy chain (IgH), TCR beta (TCRB), and TCR delta (TCRD) genes or concerns direct V to J rearrangements in case of Ig kappa (IgK), Ig lambda (IgL), TCR alpha (TCRA), and TCR gamma (TCRG) genes. The sequences between rearranging gene segments are generally deleted in the form of a circular excision product, also called TCR excision circle (TREC) or B cell receptor excision circle (BREC).

[00202] The many different combinations of V, D, and J gene segments represent the so-called combinatorial repertoire, which is estimated to be $\sim 2 \times 10^6$ for Ig molecules, $\sim 3 \times 10^6$ for TCR $\alpha\beta$ and $\sim 5 \times 10^3$ for TCR $\gamma\delta$ molecules. At the junction sites of the V, D, and J gene segments, deletion and random insertion of nucleotides occurs during the rearrangement process, resulting in highly diverse junctional regions, which significantly contribute to the total repertoire of Ig and TCR molecules, estimated to be $> 10^{12}$.

[00203] Mature B-lymphocytes further extend their Ig repertoire upon antigen recognition in follicle centers via somatic hypermutation, a process, leading to affinity maturation of the Ig molecules. The somatic hypermutation process focuses on the V- (D-) J exon of IgH and Ig light chain genes and concerns single nucleotide mutations and sometimes also insertions or deletions of nucleotides. Somatic-mutated Ig genes are also found in mature B-cell malignancies of follicular or post-follicular origin.

[00204] In certain embodiments described herein, V-segment and J-segment primers may be employed in a PCR reaction to amplify rearranged TCR or BCR CDR3-encoding DNA regions in a test biological sample, wherein each functional TCR or Ig V-encoding gene segment comprises a V gene recombination signal sequence (RSS) and each functional TCR or Ig J-encoding gene segment comprises a J gene RSS. In these and related embodiments, each amplified rearranged DNA molecule may comprise (i) at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 (including all integer values therebetween) or more contiguous nucleotides of a sense strand of the TCR or Ig V-encoding gene segment, with the at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 or more contiguous nucleotides being situated 5' to the V gene RSS and/or each amplified rearranged DNA molecule may comprise (ii) at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 (including all integer values therebetween) or more contiguous nucleotides of a sense strand of the TCR or Ig J-encoding gene segment, with the at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 or more contiguous nucleotides being situated 3' to the J gene RSS.

[00205] The practice of certain embodiments of the present invention will employ, unless indicated specifically to the contrary, conventional methods in microbiology, molecular biology, biochemistry, molecular genetics, cell biology, virology and immunology techniques that are within the skill of the art, and reference to several of which is made below for the purpose of illustration. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (3rd Edition, 2001); Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); Maniatis et al., *Molecular Cloning: A Laboratory Manual* (1982); Ausubel et al., *Current Protocols in Molecular Biology* (John Wiley and Sons, updated July 2008); *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*, Greene Pub. Associates and Wiley-Interscience; Glover, *DNA Cloning: A Practical Approach*, vol. I & II (IRL Press, Oxford Univ. Press USA, 1985); *Current Protocols in Immunology* (Edited by: John E. Coligan, Ada M. Kruisbeek, David H. Margulies, Ethan M. Shevach, Warren Strober 2001 John Wiley & Sons, NY, NY); *Real-Time PCR: Current Technology and Applications*, Edited by Julie Logan, Kirstin Edwards and Nick Saunders, 2009, Caister Academic Press, Norfolk, UK; Anand, *Techniques for the Analysis of Complex Genomes*, (Academic Press, New York, 1992); Guthrie and Fink, *Guide to Yeast Genetics and Molecular Biology* (Academic Press, New York, 1991); *Oligonucleotide Synthesis* (N. Gait, Ed., 1984); *Nucleic*

Acid Hybridization (B. Hames & S. Higgins, Eds., 1985); *Transcription and Translation* (B. Hames & S. Higgins, Eds., 1984); *Animal Cell Culture* (R. Freshney, Ed., 1986); Perbal, *A Practical Guide to Molecular Cloning* (1984); *Next-Generation Genome Sequencing* (Janitz, 2008 Wiley-VCH); *PCR Protocols (Methods in Molecular Biology)* (Park, Ed., 3rd Edition, 2010 Humana Press); *Immobilized Cells And Enzymes* (IRL Press, 1986); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller and M. P. Calos eds., 1987, Cold Spring Harbor Laboratory); Harlow and Lane, *Antibodies*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1998); *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and CC Blackwell, eds., 1986); Riott, *Essential Immunology*, 6th Edition, (Blackwell Scientific Publications, Oxford, 1988); *Embryonic Stem Cells: Methods and Protocols* (Methods in Molecular Biology) (Kurstad Turksen, Ed., 2002); *Embryonic Stem Cell Protocols: Volume I: Isolation and Characterization* (Methods in Molecular Biology) (Kurstad Turksen, Ed., 2006); *Embryonic Stem Cell Protocols: Volume II: Differentiation Models* (Methods in Molecular Biology) (Kurstad Turksen, Ed., 2006); *Human Embryonic Stem Cell Protocols* (Methods in Molecular Biology) (Kursad Turksen Ed., 2006); *Mesenchymal Stem Cells: Methods and Protocols* (Methods in Molecular Biology) (Darwin J. Prockop, Donald G. Phinney, and Bruce A. Bunnell Eds., 2008); *Hematopoietic Stem Cell Protocols* (Methods in Molecular Medicine) (Christopher A. Klug, and Craig T. Jordan Eds., 2001); *Hematopoietic Stem Cell Protocols* (Methods in Molecular Biology) (Kevin D. Bunting Ed., 2008) *Neural Stem Cells: Methods and Protocols* (Methods in Molecular Biology) (Leslie P. Weiner Ed., 2008).

[00206] Unless specific definitions are provided, the nomenclature utilized in connection with, and the laboratory procedures and techniques of, molecular biology, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for recombinant technology, molecular biological, microbiological, chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[00207] The term “isolated” means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally occurring tissue, cell, nucleic acid or polypeptide present in its original milieu in a

living animal is not isolated, but the same tissue, cell, nucleic acid or polypeptide, separated from some or all of the co-existing materials in the natural system, is isolated. Such nucleic acid could be part of a vector and/or such nucleic acid or polypeptide could be part of a composition (*e.g.*, a cell lysate), and still be isolated in that such vector or composition is not part of the natural environment for the nucleic acid or polypeptide. The term “gene” means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region “leader and trailer” as well as intervening sequences (introns) between individual coding segments (exons).

[00208] Unless the context requires otherwise, throughout the present specification and claims, the word “comprise” and variations thereof, such as, “comprises” and “comprising” are to be construed in an open, inclusive sense, that is, as “including, but not limited to”. By “consisting of” is meant including, and typically limited to, whatever follows the phrase “consisting of.” By “consisting essentially of” is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase “consisting essentially of” indicates that the listed elements are required or mandatory, but that no other elements are required and may or may not be present depending upon whether or not they affect the activity or action of the listed elements.

[00209] In this specification and the appended claims, the singular forms “a,” “an” and “the” include plural references unless the content clearly dictates otherwise. As used herein, in particular embodiments, the terms “about” or “approximately” when preceding a numerical value indicates the value plus or minus a range of 5%, 6%, 7%, 8% or 9%. In other embodiments, the terms “about” or “approximately” when preceding a numerical value indicates the value plus or minus a range of 10%, 11%, 12%, 13% or 14%. In yet other embodiments, the terms “about” or “approximately” when preceding a numerical value indicates the value plus or minus a range of 15%, 16%, 17%, 18%, 19% or 20%.

[00210] Reference throughout this specification to “one embodiment” or “an embodiment” or “an aspect” means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present invention. Thus, the appearances of the phrases “in one embodiment” or “in an embodiment” in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments.

EXAMPLES

Example 1: Single Molecule Labeling.

[00211] The single molecule labeling process used a Polymerase Chain Reaction approach to tag adaptive immune receptor encoding sequences with a unique barcode and a universal primer. The PCR reaction to tag the individual barcodes used QIAGEN Multiplex PCR master mix (QIAGEN part number 206145, Qiagen, Valencia, CA), 10% Q- solution (QIAGEN), and 300 ng of template DNA. The pooled primers were added so the final reaction had an aggregate forward primer concentration of 2 uM and an aggregate reverse primer concentration of 2 uM. The forward primers were composed of nucleotide sequence portions that annealed to V genes (segments that annealed to the V genes are shown in Table 2) and at the 5' end a universal primer (pGEX f, Table 3). The aggregate primer is listed in Table 6. These primers may, for greater specificity, have a random nucleotide insertion between the 3' end of the V primer and the 5' end of the universal primer sequence. The reverse primers have a section of nucleotides that can anneal to the J gene region (Table 2), on the 5' end of the J primer an 8 bp barcode composed of random nucleotides, and on the 5' end of the 8 bp random barcode a universal primer (pGEXr, Table 3). An example of these primers is listed in Table 5. The 8 bp barcode made of random nucleotides may be shorter or longer, additional basepairs increase the number of unique barcodes.

[00212] The nucleotide tags were incorporated onto the molecules in a 7 cycle PCR reaction. The thermocycle conditions were: 95° C for 5 minutes, followed by 7 cycles of 95° for 30 sec, 68° for 90 sec, and 72° for 30 sec. Following cycling, the rxn is held for 10 minutes at 72°.

[00213] Once the antigen receptor molecules were tagged by the primers carrying a random 8 bp tag, any remaining primers were destroyed using ExoSAP-IT (Product # 78200, Affymetrix, Santa Clara, CA). ExoSAP-IT is a product from Affymetrix that uses Exonuclease I and Shrimp Alkaline Phosphatase activities; the Exonuclease I destroys single stranded DNA and SAP degrades dNTPs. For this example, 10 ul of PCR reagents and 4 ul of exoSAP-IT were used. The reaction was incubated for 15 minutes at 37°C and the ExoSAP-it was inactivated by a 15 minute incubation at 80°C. At this point, the molecules were uniquely tagged with a barcode and a universal primer. To amplify the tagged products, another PCR reaction was performed with the universal pGEX primers. This reaction used QIAGEN Multiplex PCR master mix (QIAGEN part number 206145, Qiagen, Valencia, CA), 10% Q- solution (QIAGEN), and 6 ul of cleaned PCR reaction as template. The

forward universal (pGEXf) primer was added to the mix so the final concentration was 2 uM and the reverse universal primer (pgEXr) was added to the reaction so its final concentration was 2uM. To sequence these molecules, an Illumina adapter was incorporated using the pGEX primers. The reaction conditions were the same as above, except that the primers were replaced with the tailing primers (Table 7 below (SEQ ID NOs: 5686-5877)). The Illumina adapters, which also included an 8 bp tag and a 6 bp random set of nucleotides, were incorporated onto the molecules in a 7 cycle PCR reaction. The thermocycle conditions were: 95° C for 5 minutes, followed by 7 cycles of 95° for 30 sec, 68° for 90 sec, and 72° for 30 sec. Following cycling, the reaction was held for 10 minutes at 72°.

[00214] Once the labeled molecules were “tailed” with Illumina adaptors, they were amenable to sequencing. For this example, sequencing was conducted through the 8 bp randomer into the adaptive immune receptor encoding sequence on an Illumina HISEQ™ sequencing platform. The sequenced molecules included an 8 bp random tag. Every sequenced molecule having identical CDR3 and 8 bp random tag sequences was amplified from the adaptive immune receptor encoding polynucleotide sequences of a single cell.

[00215] Table 5 shows the J primers for the single molecule sequencing (reverse primers) and Table 6 shows the V primers (forward primers). The PCR protocol is short: 1st PCR (5 cycles) with the above primers to uniquely tag each molecule, followed by a second PCR (35 cycles) with a universal primer (PGEX) to amplify the molecules. These reactions are followed by a PCR reaction to tail on the Illumina adapters.

Table 5

J Primer Name	Bases	SEQ ID NO:	Sequence
pGEXr_TCRBJ1-1_vD12	60	5613	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NGT CTT ACC TAC AAC TGT GAG TCT GGT GCC
pGEXr_TCRBJ1-2_vD12	59	5614	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCC TTA CCT ACA ACG GTT AAC CTG GTC CC
pGEXr_TCRBJ1-3	62	5615	CCG GGA GCT GCA TGT GTC

_vD12			AGA GGN NNN NNN NCT TAC TCA CCT ACA ACA GTG AGC CAA CTT CC
pGEXr_TCRBJ1-4 _vD12	57	5616	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NAT ACC CAA GAC AGA GAG CTG GGT TCC
pGEXr_TCRBJ1-5 _vD12	60	5617	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NAA CTT ACC TAG GAT GGA GAG TCG AGT CCC
pGEXr_TCRBJ1-6 _vD12	53	5618	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCT GTC ACA GTG AGC CTG GTC CC
pGEXr_TCRBJ2-1 _vD12	49	5619	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCA CGG TGA GCC GTG TCC C
pGEXr_TCRBJ2-2 _vD12	53	5620	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCC AGT ACG GTC AGC CTA GAG CC
pGEXr_TCRBJ2-3 _vD12	49	5621	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCA CTG TCA GCC GGG TGC C
pGEXr_TCRBJ2-4 _vD12	49	5622	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCA CTG AGA GCC GGG TCC C
pGEXr_TCRBJ2-5 _vD12	48	5623	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NAC CAG GAG CCG CGT GCC
pGEXr_TCRBJ2-6 _vD12	49	5624	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NCA CGG TCA GCC TGC TGC C

pGEXr_TCRBJ2-7 _vD12	49	5625	CCG GGA GCT GCA TGT GTC AGA GGN NNN NNN NGA CCG TGA GCC TGG TGC C
-------------------------	----	------	---

Table 6

Primer Name	SEQ ID NO:	Sequence
pGEXf_TCRB V01_verD10	5626	GGGCTGGCAAGCCACGTTTGGTGGGAATGCCCTGAC AGCTCTCGCTTATA
pGEXf_TCRB V02_verD10	5627	GGGCTGGCAAGCCACGTTTGGTGCTCAGAGAAGTC TGAAATATTCGATGATCAATTCTCAGTTG
pGEXf_TCRB V03-1_verD10	5628	GGGCTGGCAAGCCACGTTTGGTGCCAAATCGMTTC TCACCTAAATCTCCAGACAAAG
pGEXf_TCRB V03-2_verD10	5629	GGGCTGGCAAGCCACGTTTGGTGCACCTGACTCTCC AGACAAAGCTCAT
pGEXf_TCRB V04- 1/2/3_verD10	5630	GGGCTGGCAAGCCACGTTTGGTGCCTGAATGCCCCA ACAGCTCTC
pGEXf_TCRB V05-1_verD10	5631	GGGCTGGCAAGCCACGTTTGGTGGATTCTCAGGGC GCCAGTTCTCTA
pGEXf_TCRB V05-2_verD10	5632	GGGCTGGCAAGCCACGTTTGGTGCCTAATTGATTCT CAGCTCACACGTCCATA
pGEXf_TCRB V05-3_verD10	5633	GGGCTGGCAAGCCACGTTTGGTGTCAGGGCGCCAG TTCCATG
pGEXf_TCRB V05-4_verD10	5634	GGGCTGGCAAGCCACGTTTGGTGTCCTAGATTCTCA GGTCTCCAGTTCCCTA
pGEXf_TCRB V05-5_verD10	5635	GGGCTGGCAAGCCACGTTTGGTGGAGGAAACTTCC CTGATCGATTCTCAGC
pGEXf_TCRB V05-6_verD10	5636	GGGCTGGCAAGCCACGTTTGGTGCAACTTCCCTGAT CGATTCTCAGGTCA
pGEXf_TCRB V05-7_verD10	5637	GGGCTGGCAAGCCACGTTTGGTGAGGAAACTTCCCT GATCAATTCTCAGGTCA

pGEXf_TCRB V05-8_verD10	5638	GGGCTGGCAAGCCACGTTTGGTGGGAACTTCCCTC CTAGATTTTCAGGTCG
pGEXf_TCRB V06-1_verD10	5639	GGGCTGGCAAGCCACGTTTGGTGCCCCAATGGCTAC AATGTCTCCAGATT
pGEXf_TCRB V06- 2/3_verD10	5640	GGGCTGGCAAGCCACGTTTGGTGGGAGAGGTCCCT GATGGCTACAA
pGEXf_TCRB V06-4_verD10	5641	GGGCTGGCAAGCCACGTTTGGTGTCCCTGATGGTTA TAGTGTCTCCAGAGC
pGEXf_TCRB V06-5_verD10	5642	GGGCTGGCAAGCCACGTTTGGTGGGAGAAGTCCCC AATGGCTACAATGTC
pGEXf_TCRB V06-6_verD10	5643	GGGCTGGCAAGCCACGTTTGGTGAAAGGAGAAGTC CCGAATGGCTACAA
pGEXf_TCRB V06-7_verD10	5644	GGGCTGGCAAGCCACGTTTGGTGGTTCCCAATGGCT ACAATGTCTCCAGATC
pGEXf_TCRB V06-8_verD10	5645	GGGCTGGCAAGCCACGTTTGGTGGAAGTCCCCAAT GGCTACAATGTCTCTAGATT
pGEXf_TCRB V06-9_verD10	5646	GGGCTGGCAAGCCACGTTTGGTGGAGAAGTCCCCG ATGGCTACAATGTA
pGEXf_TCRB V07-1_verD10	5647	GGGCTGGCAAGCCACGTTTGGTGGTGATCGGTTCTC TGCACAGAGGT
pGEXf_TCRB V07-2_verD10	5648	GGGCTGGCAAGCCACGTTTGGTGCGCTTCTCTGCAG AGAGGACTGG
pGEXf_TCRB V07-3_verD10	5649	GGGCTGGCAAGCCACGTTTGGTGGGTCTTTGCAGT CAGGCCTGA
pGEXf_TCRB V07-4_verD10	5650	GGGCTGGCAAGCCACGTTTGGTGCAGTGGTCGGTTC TCTGCAGAG
pGEXf_TCRB V07-5_verD10	5651	GGGCTGGCAAGCCACGTTTGGTGGCTCAGTGATCA ATTCTCCACAGAGAGGT
pGEXf_TCRB V07- 6/7_verD10	5652	GGGCTGGCAAGCCACGTTTGGTGTTCTCTGCAGAGA GGCCTGAGG

pGEXf_TCRB V07-8_verD10	5653	GGGCTGGCAAGCCACGTTTGGTGCCCAGTGATCGCT TCTTTGCAGAAA
pGEXf_TCRB V07-9_verD10	5654	GGGCTGGCAAGCCACGTTTGGTGCTGCAGAGAGGC CTAAGGGATCT
pGEXf_TCRB V08-1_verD10	5655	GGGCTGGCAAGCCACGTTTGGTGGAAGGGTACAAT GTCTCTGGAAACAAACTCAAG
pGEXf_TCRB V08-2_verD10	5656	GGGCTGGCAAGCCACGTTTGGTGGGGGTACTGTGTT TCTTGAAACAAGCTTGAG
pGEXf_TCRB V09_verD10	5657	GGGCTGGCAAGCCACGTTTGGTGCAAGTTCCTGACT TGCACTCTGAACTAAAC
pGEXf_TCRB V10-1_verD10	5658	GGGCTGGCAAGCCACGTTTGGTGACTAACAAAGGA GAAGTCTCAGATGGCTACAG
pGEXf_TCRB V10-2_verD10	5659	GGGCTGGCAAGCCACGTTTGGTGAGATAAAGGAGA AGTCCCCGATGGCTA
pGEXf_TCRB V10-3_verD10	5660	GGGCTGGCAAGCCACGTTTGGTGGATACTGACAAA GGAGAAGTCTCAGATGGCTATAG
pGEXf_TCRB V11- 1/2/3_verD10	5661	GGGCTGGCAAGCCACGTTTGGTGCTAAGGATCGATT TTCTGCAGAGAGGCTC
pGEXf_TCRB V12-1_verD10	5662	GGGCTGGCAAGCCACGTTTGGTGTTGATTCTCAGCA CAGATGCCTGATGT
pGEXf_TCRB V12-2_verD10	5663	GGGCTGGCAAGCCACGTTTGGTGATTCTCAGCTGAG AGGCCTGATGG
pGEXf_TCRB V12- 3/4_verD10	5664	GGGCTGGCAAGCCACGTTTGGTGGGATCGATTCTCA GCTAAGATGCCTAATGC
pGEXf_TCRB V12-5_verD10	5665	GGGCTGGCAAGCCACGTTTGGTGCTCAGCAGAGAT GCCTGATGCAACTTTA
pGEXf_TCRB V13_verD10	5666	GGGCTGGCAAGCCACGTTTGGTGCTGATCGATTCTC AGCTCAACAGTTCAGT
pGEXf_TCRB V14_verD10	5667	GGGCTGGCAAGCCACGTTTGGTGTAGCTGAAAGGA CTGGAGGGACGTAT

pGEXf_TCRB V15_verD10	5668	GGGCTGGCAAGCCACGTTTGGTGCCAGGAGGCCGA ACACTTCTTTCT
pGEXf_TCRB V16_verD10	5669	GGGCTGGCAAGCCACGTTTGGTGGCTAAGTGCCTCC CAAATTCACCCT
pGEXf_TCRB V17_verD10	5670	GGGCTGGCAAGCCACGTTTGGTGACAGCTGAAAG ACCTAACGGAACGT
pGEXf_TCRB V18_verD10	5671	GGGCTGGCAAGCCACGTTTGGTGCTGCTGAATTTCC CAAAGAGGGCC
pGEXf_TCRB V19_verD10	5672	GGGCTGGCAAGCCACGTTTGGTGAGGGTACAGCGT CTCTCGGG
pGEXf_TCRB V20_verD10	5673	GGGCTGGCAAGCCACGTTTGGTGGCCTGACCTTGTC CACTCTGACA
pGEXf_TCRB V21_verD10	5674	GGGCTGGCAAGCCACGTTTGGTGATGAGCGATTTTT AGCCCAATGCTCCA
pGEXf_TCRB V22_verD10	5675	GGGCTGGCAAGCCACGTTTGGTGTGAAGGCTACGT GTCTGCCAAGAG
pGEXf_TCRB V23_verD10	5676	GGGCTGGCAAGCCACGTTTGGTGCTCATCTCAATGC CCCAAGAACGC
pGEXf_TCRB V24_verD10	5677	GGGCTGGCAAGCCACGTTTGGTGAGATCTCTGATGG ATACAGTGTCTCTCGACA
pGEXf_TCRB V25_verD10	5678	GGGCTGGCAAGCCACGTTTGGTGAGATCTTTCCTCT GAGTCAACAGTCTCCAGAATA
pGEXf_TCRB V26_verD10	5679	GGGCTGGCAAGCCACGTTTGGTGCACTGAAAAAGG AGATATCTCTGAGGGGTATCATG
pGEXf_TCRB V27_verD10	5680	GGGCTGGCAAGCCACGTTTGGTGGTTCCTGAAGGGT ACAAAGTCTCTCGAAAAG
pGEXf_TCRB V28_verD10	5681	GGGCTGGCAAGCCACGTTTGGTGCTGAGGGGTACA GTGTCTCTAGAGAGA
pGEXf_TCRB V29_verD10	5682	GGGCTGGCAAGCCACGTTTGGTGAGCCGCCCAAAC CTAACATTCTCAA
pGEXf_TCRB V30_verD10	5683	GGGCTGGCAAGCCACGTTTGGTGCCCAGGACCGGC AGTTCA

pGEXf_TCRB VA_verD10	5684	GGGCTGGCAAGCCACGTTTGGTGTTGATTAGAGAC ATATCCCTATTGAAAATATTTCTGCA
pGEXf_TCRB VB_verD10	5685	GGGCTGGCAAGCCACGTTTGGTGAGATGCCCTGAG TCAGCATAGTCATTCTAAC

Table 7. Exemplary Tailing Primers

Sequence	SEQ ID NO:
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAA GGT CAC CGG GAG CTG CAT GTG TCA GAG G	5686
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GCA TAA CTC CGG GAG CTG CAT GTG TCA GAG G	5687
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTC TGA TTC CGG GAG CTG CAT GTG TCA GAG G	5688
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAC GTA CGC CGG GAG CTG CAT GTG TCA GAG G	5689
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAC GCG TTC CGG GAG CTG CAT GTG TCA GAG G	5690
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTC AGT GAC CGG GAG CTG CAT GTG TCA GAG G	5691
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCT GAT ATC CGG GAG CTG CAT GTG TCA GAG G	5692
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAT ATG CTC CGG GAG CTG CAT GTG TCA GAG G	5693
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CGT AAT TAC CGG GAG CTG CAT GTG TCA GAG G	5694
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT	5695

GCT GAA CCG CTC TTC CGA TCT NNN NNN ACG TAC TCC CGG GAG CTG CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTT CTA AGC CGG GAG CTG CAT GTG TCA GAG G	5696
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ACT ATG ACC CGG GAG CTG CAT GTG TCA GAG G	5697
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAC GTT AAC CGG GAG CTG CAT GTG TCA GAG G	5698
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ACA AGA TAC CGG GAG CTG CAT GTG TCA GAG G	5699
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAC TAA GAC CGG GAG CTG CAT GTG TCA GAG G	5700
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTG TCT ACC CGG GAG CTG CAT GTG TCA GAG G	5701
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTC ACT AGC CGG GAG CTG CAT GTG TCA GAG G	5702
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AAT CGG ATC CGG GAG CTG CAT GTG TCA GAG G	5703
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGT ACC GAC CGG GAG CTG CAT GTG TCA GAG G	5704
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTG CCT CAC CGG GAG CTG CAT GTG TCA GAG G	5705

AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCG TTA GCC CGG GAG CTG CAT GTG TCA GAG G	5706
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAT AGT TCC CGG GAG CTG CAT GTG TCA GAG G	5707
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGG CGT ATC CGG GAG CTG CAT GTG TCA GAG G	5708
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGG ACA TGC CGG GAG CTG CAT GTG TCA GAG G	5709
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGG TTG CTC CGG GAG CTG CAT GTG TCA GAG G	5710
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATA TGC TGC CGG GAG CTG CAT GTG TCA GAG G	5711
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTA CAG TGC CGG GAG CTG CAT GTG TCA GAG G	5712
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATC CAT GGC CGG GAG CTG CAT GTG TCA GAG G	5713
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGA TGC GAC CGG GAG CTG CAT GTG TCA GAG G	5714
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTA GCA GTC CGG GAG CTG CAT GTG TCA GAG G	5715
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GGA TCA TCC CGG GAG CTG	5716

CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTG AAC GTC CGG GAG CTG CAT GTG TCA GAG G	5717
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATT AAG CGC CGG GAG CTG CAT GTG TCA GAG G	5718
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAT TGG CGC CGG GAG CTG CAT GTG TCA GAG G	5719
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CGA TTA CAC CGG GAG CTG CAT GTG TCA GAG G	5720
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGT CAT CGC CGG GAG CTG CAT GTG TCA GAG G	5721
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAT CAA GTC CGG GAG CTG CAT GTG TCA GAG G	5722
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGG CTT GAC CGG GAG CTG CAT GTG TCA GAG G	5723
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAT AAC CAC CGG GAG CTG CAT GTG TCA GAG G	5724
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AAT CCT GCC CGG GAG CTG CAT GTG TCA GAG G	5725
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTT ATA TCC CGG GAG CTG CAT GTG TCA GAG G	5726
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT	5727

GCT GAA CCG CTC TTC CGA TCT NNN NNN ACA CAC GTC CGG GAG CTG CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATA CGA CTC CGG GAG CTG CAT GTG TCA GAG G	5728
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATC TTC GTC CGG GAG CTG CAT GTG TCA GAG G	5729
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ACA TGT ATC CGG GAG CTG CAT GTG TCA GAG G	5730
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCC ACA GTC CGG GAG CTG CAT GTG TCA GAG G	5731
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAG TCT GTC CGG GAG CTG CAT GTG TCA GAG G	5732
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCC ATG TGC CGG GAG CTG CAT GTG TCA GAG G	5733
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCA CTG CAC CGG GAG CTG CAT GTG TCA GAG G	5734
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATG GTC AAC CGG GAG CTG CAT GTG TCA GAG G	5735
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAA GTC ACC CGG GAG CTG CAT GTG TCA GAG G	5736
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAG ACG GAC CGG GAG CTG CAT GTG TCA GAG G	5737

AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAG CTC TTC CGG GAG CTG CAT GTG TCA GAG G	5738
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAG CGA TAC CGG GAG CTG CAT GTG TCA GAG G	5739
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTC GAG AAC CGG GAG CTG CAT GTG TCA GAG G	5740
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATG ACA CCC CGG GAG CTG CAT GTG TCA GAG G	5741
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTT CAC GAC CGG GAG CTG CAT GTG TCA GAG G	5742
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTA TAA GGC CGG GAG CTG CAT GTG TCA GAG G	5743
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CGT AGA GTC CGG GAG CTG CAT GTG TCA GAG G	5744
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ATA GAT ACC CGG GAG CTG CAT GTG TCA GAG G	5745
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCG TCG ATC CGG GAG CTG CAT GTG TCA GAG G	5746
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAA GAA TCC CGG GAG CTG CAT GTG TCA GAG G	5747
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AAT GAC AGC CGG GAG CTG	5748

CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGC TAG TGC CGG GAG CTG CAT GTG TCA GAG G	5749
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGA GAC CTC CGG GAG CTG CAT GTG TCA GAG G	5750
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGC GTA ATC CGG GAG CTG CAT GTG TCA GAG G	5751
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TAA CCA AGC CGG GAG CTG CAT GTG TCA GAG G	5752
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAT GGC TTC CGG GAG CTG CAT GTG TCA GAG G	5753
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GCA TCT GAC CGG GAG CTG CAT GTG TCA GAG G	5754
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTC CGG TAC CGG GAG CTG CAT GTG TCA GAG G	5755
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAC ACT CTC CGG GAG CTG CAT GTG TCA GAG G	5756
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTA AGC ATC CGG GAG CTG CAT GTG TCA GAG G	5757
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGC TAC ACC CGG GAG CTG CAT GTG TCA GAG G	5758
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT	5759

GCT GAA CCG CTC TTC CGA TCT NNN NNN TCA GCT TGC CGG GAG CTG CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAT GTA GAC CGG GAG CTG CAT GTG TCA GAG G	5760
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTC GGA ACC CGG GAG CTG CAT GTG TCA GAG G	5761
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GCA ATT CGC CGG GAG CTG CAT GTG TCA GAG G	5762
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CAA GAG GTC CGG GAG CTG CAT GTG TCA GAG G	5763
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TCG ATT AAC CGG GAG CTG CAT GTG TCA GAG G	5764
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GAA TGG ACC CGG GAG CTG CAT GTG TCA GAG G	5765
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGA ATC AGC CGG GAG CTG CAT GTG TCA GAG G	5766
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AAC TGC CAC CGG GAG CTG CAT GTG TCA GAG G	5767
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AAG TAA CGC CGG GAG CTG CAT GTG TCA GAG G	5768
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN ACT CAA TGC CGG GAG CTG CAT GTG TCA GAG G	5769

AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CCT AGT AGC CGG GAG CTG CAT GTG TCA GAG G	5770
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTG ACG TTC CGG GAG CTG CAT GTG TCA GAG G	5771
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGC AGA CAC CGG GAG CTG CAT GTG TCA GAG G	5772
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGT TGA CCC CGG GAG CTG CAT GTG TCA GAG G	5773
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GTC TCC TAC CGG GAG CTG CAT GTG TCA GAG G	5774
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CTG CAA TCC CGG GAG CTG CAT GTG TCA GAG G	5775
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TGA GCG AAC CGG GAG CTG CAT GTG TCA GAG G	5776
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTG GAC TGC CGG GAG CTG CAT GTG TCA GAG G	5777
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN AGC AAT CCC CGG GAG CTG CAT GTG TCA GAG G	5778
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN CGA ACT ACC CGG GAG CTG CAT GTG TCA GAG G	5779
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN TTA ATG GCC CGG GAG CTG	5780

CAT GTG TCA GAG G	
AAT GAT ACG GCG ACC ACC GAG ATC TAC ACC GGT CTC GGC ATT CCT GCT GAA CCG CTC TTC CGA TCT NNN NNN GCT TAG TAC CGG GAG CTG CAT GTG TCA GAG G	5781
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAA GGT CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5782
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GCA TAA CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5783
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTC TGA TTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5784
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAC GTA CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5785
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAC GCG TTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5786
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTC AGT GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5787
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCT GAT ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5788
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAT ATG CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5789
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CGT AAT TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5790
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG	5791

CTC TTC CGA TCT ACG TAC TCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTT CTA AGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5792
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ACT ATG ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5793
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAC GTT AAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5794
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ACA AGA TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5795
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAC TAA GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5796
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTG TCT ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5797
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTC ACT AGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5798
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AAT CGG ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5799
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGT ACC GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5800
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTG CCT CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5801

CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCG TTA GCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5802
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAT AGT TCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5803
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGG CGT ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5804
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGG ACA TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5805
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGG TTG CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5806
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATA TGC TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5807
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTA CAG TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5808
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATC CAT GGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5809
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGA TGC GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5810
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTA GCA GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5811
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GGA TCA TCN NNN NNG GGC TGG CAA GCC ACG TTT	5812

GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTG AAC GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5813
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATT AAG CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5814
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAT TGG CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5815
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CGA TTA CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5816
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGT CAT CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5817
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAT CAA GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5818
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGG CTT GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5819
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAT AAC CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5820
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AAT CCT GCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5821
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTT ATA TCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5822
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG	5823

CTC TTC CGA TCT ACA CAC GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATA CGA CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5824
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATC TTC GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5825
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ACA TGT ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5826
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCC ACA GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5827
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAG TCT GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5828
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCC ATG TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5829
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCA CTG CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5830
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATG GTC AAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5831
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAA GTC ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5832
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAG ACG GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5833

CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAG CTC TTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5834
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAG CGA TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5835
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTC GAG AAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5836
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATG ACA CCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5837
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTT CAC GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5838
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTA TAA GGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5839
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CGT AGA GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5840
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ATA GAT ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5841
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCG TCG ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5842
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAA GAA TCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5843
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AAT GAC AGN NNN NNG GGC TGG CAA GCC ACG TTT	5844

GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGC TAG TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5845
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGA GAC CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5846
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGC GTA ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5847
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TAA CCA AGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5848
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAT GGC TTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5849
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GCA TCT GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5850
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTC CGG TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5851
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAC ACT CTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5852
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTA AGC ATN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5853
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGC TAC ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5854
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG	5855

CTC TTC CGA TCT TCA GCT TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAT GTA GAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5856
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTC GGA ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5857
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GCA ATT CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5858
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CAA GAG GTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5859
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TCG ATT AAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5860
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GAA TGG ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5861
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGA ATC AGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5862
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AAC TGC CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5863
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AAG TAA CGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5864
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT ACT CAA TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5865

CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CCT AGT AGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5866
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTG ACG TTN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5867
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGC AGA CAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5868
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGT TGA CCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5869
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GTC TCC TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5870
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CTG CAA TCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5871
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TGA GCG AAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5872
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTG GAC TGN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5873
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT AGC AAT CCN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5874
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT CGA ACT ACN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5875
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT TTA ATG GCN NNN NNG GGC TGG CAA GCC ACG TTT	5876

GGT G	
CAA GCA GAA GAC GGC ATA CGA GAT ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT GCT TAG TAN NNN NNG GGC TGG CAA GCC ACG TTT GGT G	5877

Example 2: Single Cell Labeling of Adaptive Immune Receptor Encoding Sequences

[00216] This example describes single cell labeling of immunoglobulin and T cell receptor heavy and light chain encoding sequences by RT-PCR. Freshly drawn blood from healthy human volunteers is used as a source of leukocytes. The amount of whole blood required to obtain 100,000 – 300,000 leukocytes is less than 1mL; 1-3 mL of blood are used for isolation of blood cells. Peripheral blood mononuclear cells (PBMC) are isolated from blood by density gradient centrifugation on Histopaque®-1077 (Sigma, St. Louis, MO) according to the supplier’s instructions. CD45⁺ hematopoietic cells are isolated by binding to anti-CD45 coated magnetic beads using Whole Blood CD45 Microbeads (Miltenyi Biotec, Auburn, CA) as instructed by the manufacturer and essentially as described in Koehl et al. (2003 *Leukemia* 17:232). Leukocyte cell suspensions are washed in phosphate-buffered saline solution (PBS) and adjusted to a concentration of 1 x 10⁶ cells/ mL. Aliquots of 1-3 µL (1-3 x 10³ cells) are distributed into wells of 96-well PCR multiwell plates held on ice in pre-chilled plate racks. Immediately after all plate wells are filled, the plates are sealed and placed on dry ice to freeze and lyse the cells. Plates are held on dry ice during the reverse transcription preparation steps below.

[00217] Reverse transcription is performed using the SMARTer™ Ultra Low RNA kit for Illumina sequencing (Clontech, Mountain View, CA) essentially according to the supplier’s instructions. Stock Reaction Buffer is prepared by mixing 380 µl of Dilution Buffer with 20 µl of RNase inhibitor (40U/µl). 250 µl of Reaction Buffer is then mixed with 100 µl of a 12 µM solution of the 3’ Smarter™ CDS II oligonucleotide (5’-Bio–AAGCAGTGGTATCAACGCAGAGTACT₍₃₀₎NN–3’ [SEQ ID NO: 5878], where Bio is a biotin moiety; AAGCAGTGGTATCAACGCAGAGTAC [SEQ ID NO: 5879] is a universal adapter sequence, T₍₃₀₎ (SEQ ID NO: 5880) is a 30-mer of thymine residues, and N is any nucleotide (A, C, G or T).

[00218] The first-step annealing reactions for reverse transcription are set up by adding 3.5 µl of the Reaction Buffer containing the 3’ Smarter™ CDS II oligonucleotide primer to each

well of the 96-well plate containing the lysed cells, sealing the plate and incubating it for 3 minutes at 72°C, after which it is returned to a chilled rack on ice.

[00219] Reverse Transcription Master Mix (450 µl for 100 rxns) is prepared by combining 200 µl of 5x First Strand Buffer, 25 µl of 100 mM dithithreitol (DTT), 100 µl of dNTPs (10mM), 25 µl of RNase inhibitor (40U/µl), and 100 µl of reverse transcriptase. A 96-well working plate is prepared containing 1.0 µl of a barcoded 3'-Smart™ CDSII oligonucleotide per well. The 3'-Smart CDSII oligo sequence is: 5'–

AAGCAGTGGTATCAACGCAGAGTACBBBBBBBBBrGrGrG–P-3' [SEQ ID NO: 5881]

where AAGCAGTGGTATCAACGCAGAGTAC [SEQ ID NO: 5879] is a universal adapter sequence; BBBBBBBB is an 8-nucleotide barcode (see list below for examples of barcodes); rG is riboguanine; and P is a 3' phosphate blocking moiety.

Table 8. Barcode list (96 JS barcodes):

Name	Sequence
JS01	CAAGGTCA
JS02	GCATAACT
JS03	CTCTGATT
JS04	TACGTACG
JS05	TACGCGTT
JS06	CTCAGTGA
JS07	TCTGATAT
JS08	CATATGCT
JS09	CGTAATTA
JS10	ACGTACTC
JS11	CTTCTAAG
JS12	ACTATGAC
JS13	GACGTTAA
JS14	ACAAGATA
JS15	GACTAAGA
JS16	GTGTCTAC
JS17	TTCACTAG
JS18	AATCGGAT
JS19	AGTACCGA

JS20	TTGCCTCA
JS21	TCGTTAGC
JS22	TATAGTTC
JS23	TGGCGTAT
JS24	TGGACATG
JS25	AGGTTGCT
JS26	ATATGCTG
JS27	GTACAGTG
JS28	ATCCATGG
JS29	TGATGCGA
JS30	GTAGCAGT
JS31	GGATCATC
JS32	GTGAACGT
JS33	ATTAAGCG
JS34	TATTGGCG
JS35	CGATTACA
JS36	TGTCATCG
JS37	TATCAAGT
JS38	AGGCTTGA
JS39	GATAACCA
JS40	AATCCTGC
JS41	GTTATATC
JS42	ACACACGT
JS43	ATACGACT
JS44	ATCTTCGT
JS45	ACATGTAT
JS46	TCCACAGT
JS47	CAGTCTGT
JS48	TCCATGTG
JS49	TCACTGCA
JS50	ATGGTCAA
JS51	CAAGTCAC

JS52	TAGACGGA
JS53	CAGCTCTT
JS54	GAGCGATA
JS55	CTCGAGAA
JS56	ATGACACC
JS57	CTTCACGA
JS58	CTATAAGG
JS59	CGTAGAGT
JS60	ATAGATAC
JS61	TCGTCGAT
JS62	TAAGAATC
JS63	AATGACAG
JS64	AGCTAGTG
JS65	TGAGACCT
JS66	AGCGTAAT
JS67	TAACCAAG
JS68	GATGGCTT
JS69	GCATCTGA
JS70	TTCCGGTA
JS71	GACACTCT
JS72	TTAAGCAT
JS73	TGCTACAC
JS74	TCAGCTTG
JS75	CATGTAGA
JS76	TTCGGAAC
JS77	GCAATTCG
JS78	CAAGAGGT
JS79	TCGATTAA
JS80	GAATGGAC
JS81	AGAATCAG
JS82	AACTGCCA
JS83	AAGTAACG

JS84	ACTCAATG
JS85	CCTAGTAG
JS86	CTGACGTT
JS87	TGCAGACA
JS88	AGTTGACC
JS89	GTCTCCTA
JS90	CTGCAATC
JS91	TGAGCGAA
JS92	TTGGACTG
JS93	AGCAATCC
JS94	CGAACTAC
JS95	TTAATGGC
JS96	GCTTAGTA

[00220] To each well of the 96-well working plate containing 1.0 µl of a barcoded 3'-Smart™ CDSII oligonucleotide is added 4.5 µl of the Master Mix, and following completion of the annealing reaction, 5.5µl of the Master Mix containing barcoded 3'-Smart™ CDSII oligonucleotide is transferred from each well of the 96-well working plate to the correspondingly positioned (respective) wells of the reverse transcription annealing plate. The reverse transcription annealing plate is placed onto a thermocycler and a program is run with the steps 42°C for 90 minutes followed by 70°C for 10 minutes. This temperature profile performs first cDNA strand synthesis on all poly-A mRNA transcript molecules released from leukocytes in each well. According to non-limiting theory, after the first cDNA strand synthesis, each cDNA molecule in a well contains universal adaptor sequences at both the 5' and 3' ends, and is uniquely tagged with an 8-nt barcode at the 5' end.

[00221] Optionally, the barcoded cDNA molecules from all 96 reactions can be pooled at this step, and re-aliquoted onto a PCR plate where PCR amplification of immunoglobulin or T cell receptor cDNA takes place. The combining and splitting step permit substantially all barcoded cDNA molecules to be substantially evenly represented in subsequent PCR amplification reactions with adaptive immune receptor encoding (*e.g.*, IG or TCR) C-segment gene specific primers.

[00222] The products of reverse transcription/ cDNA first strand synthesis are next isolated by Solid Phase Reversible Immobilization Purification (SPRI) by mixing the

contents of each well from the reverse transcription reaction plate with 25 μ l of a suspension of Ampure™ XP SPRI magnetic beads (Beckman-Coulter Inc., Brea, CA) and incubating for 8 minutes at room temperature, followed by bead separation using a MagnaBot™ magnetic separator (Promega, Madison, WI) at room temperature according to the suppliers' instructions.

[00223] SPRI bead-immobilized cDNA first strands are immediately added to 5'RACE (rapid amplification of cDNA ends) PCR amplification reactions using Advantage 2™ PCR reagents (Clontech) according to the manufacturer's instructions. For each reaction, 50 μ l of PCR Master Mix is added containing dNTPs, UPM primer mix, IG/TCR primer mix as described elsewhere herein, and Advantage 2™ polymerase and PCR buffer. The thermocycling conditions are: 95°C for 1 minute; 30 cycles of 95°C for 30 seconds, 63°C for 30 seconds, and 72°C for 3 minutes; 72°C for 7 minutes; and then reactions are held at 10°C prior to preparation for Illumina sequencing. PCR primer sequences are:

5'RACE UPM long	5'-CTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT-3'(SEQ ID NO: 5611)
5'RACE UPM short	5'-CTAATACGACTCACTATAGGGC-3' (SEQ ID NO: 5612)

IgM_RACE	5'-GATGGAGTCGGGAAGGAAGTCCTGTGCGAG-3' (SEQ ID NO: 5601)
IgG_RACE	5'-GGGAAGACSGATGGGCCCTTGGTGG-3' (SEQ ID NO: 5602)
IgA_RACE	5'-CAGGCAKGC GAYGACCACGTTCCCATC-3' (SEQ ID NO: 5603)
Igκ_RACE	5'-CATCAGATGGCGGGGAAGATGAAGACAGATGGTGC-3' (SEQ ID NO: 5604)
Igλ_RACE	5'-CCTCAGAGGAGGGTGGGAACAGAGTGAC-3' (SEQ ID NO: 5605)
TCRB_RACE	5'-GCTCAAACACAGCGACCTCGGGTGGGAACAC-3' (SEQ ID NO: 5606)
TCRA_RACE_JB2	5'-AGTCTCTCAGCTGGTACACGGCAGGGTC-3' (SEQ ID NO: 5591)
TCRA_50	5'- ACA GAC TTG TCA CTG GAT TTA GAG TCT CTC AGC TGG TAC ACG GCA GGG TC -3' (SEQ ID NO: 5592)
TCRB_50	5'- GAG ATC TCT GCT TCT GAT GGC TCA AAC ACA GCG ACC TCG GGT GGG AAC AC -3' (SEQ ID NO: 5593)
S	G or C
K	G or T
Y	C or T

Illumina Sequencing Library Preparation

[00224] PCR products are pooled by inverted centrifugation of the 96-well plates and the pooled products are purified to remove DNA fragments shorter than 200-300bp using Beckman Coulter Ampure™ XP beads according to the supplier's instructions. DNA purity is assessed by capillary electrophoresis using a Caliper Bioanalyzer (Perkin Elmer, Norwalk, CT) to confirm that most of the dsDNA is within a size range of 600-700 bp. dsDNA products are quantified fluorometrically or by A260 UV absorbance.

[00225] Sequencing library construction is conducted using 1 µg of purified DNA as an input for the Illumina TruSeq® sample preparation protocol (Illumina Inc., San Diego, CA) according to the Illumina TruSeq® DNA Sample Preparation Guide (Part number 15026486 Rev. C, July 2012, Illumina, Inc., San Diego, CA). This protocol generates a sequencing library that can be sequenced using the paired-end flow cell on the Illumina MiSeq®, HiSeq®2000, and HiSeq®2500 sequencers.

[00226] Illumina sequencing is conducted according to a sequencing protocol on the Illumina MiSeq® sequencer that utilizes the MiSeq® reagents kit v2, for 500 cycles. This chemistry provides kitted reagents for up to 525 cycles of sequencing on the MiSeq® instrument and provides sufficient reagents for a 251-cycle paired-end run, plus two eight-cycle indexed reads. The Illumina sequencing protocol is described in MiSeq® ReagentKit v2 ReagentPrepGuide, Part number 15034097 Rev. B, October 2012 (Illumina Inc., San Diego, CA). A schematic representation of the structure of DNA targets to be sequenced is shown in Fig. 6 (in which Ig heavy chain is used as an example).

[00227] The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and are incorporated herein by reference, in their entirety. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, applications and publications to provide yet further embodiments.

[00228] These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

CLAIMS

What is claimed is:

1. An oligonucleotide amplification primer composition, comprising:

(A) a first oligonucleotide amplification primer set comprising a plurality of forward oligonucleotide sequences of a general formula (A):

U1 - B1 - V1 (A),

and a plurality of reverse oligonucleotide sequences of a general formula (B):

U2 - B2 - J1 (B),

wherein U1 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence, and U2 comprises an oligonucleotide sequence comprising a second universal adaptor oligonucleotide sequence,

wherein B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, and B2 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, such that at least one of B1 or B2 is present,

wherein V1 comprises an oligonucleotide sequence comprising at least 15 and not more than 100 contiguous nucleotides of a V region encoding gene sequence of a first adaptive immune receptor, or the complement thereof,

wherein J1 comprises an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) a joining (J) region encoding gene sequence of said first adaptive immune receptor, or the complement thereof, or (ii) a constant (C) region encoding gene sequence of said first adaptive immune receptor, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1-B1-V1, V1 comprises a unique oligonucleotide sequence, and in each of the plurality of oligonucleotide sequences of general formula U2-B2-J1, J1 comprises a unique oligonucleotide sequence; and

(B) a second oligonucleotide amplification primer set comprising a plurality of forward oligonucleotide sequences of a general formula (C):

U3 - B3 - V2 (C),

and a plurality of reverse oligonucleotide sequences of a general formula (D):

U4 - B4 - J2 (D),

wherein U3 comprises an oligonucleotide sequence identical to either U1 or U2, and U4 comprises an oligonucleotide sequence identical to either U1 or U2, whichever sequence is not identical to U3,

wherein B3 comprises an oligonucleotide sequence comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B1, and B4 comprises an oligonucleotide sequence comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B2, and

wherein V2 comprises an oligonucleotide sequence comprising at least 15 and not more than 100 contiguous nucleotides of a V region encoding gene sequence of a second adaptive immune receptor, or the complement thereof,

wherein J2 comprises an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) a joining (J) region encoding gene sequence of said second adaptive immune receptor, or the complement thereof, or (ii) a constant (C) region encoding gene sequence of said second adaptive immune receptor, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3-B3-V2, V2 comprises a unique oligonucleotide sequence, and in each of the plurality of oligonucleotide sequences of general formula U4-B4-J2, J2 comprises a unique oligonucleotide sequence.

2. The composition of claim 1, wherein U1 is the same as U3.
3. The composition of claim 1, wherein U2 is the same as U4.
4. A method for labeling individual rearranged DNA sequences encoding a plurality of adaptive immune receptors in a biological sample that comprises lymphoid cells of a subject, the method comprising:

(a) amplifying said rearranged DNA sequences using a first amplification primer set comprising an oligonucleotide primer composition of claims 1-3 under conditions that promote amplification to obtain double-stranded DNA products that each comprise (i) a sequence comprising at least two universal adaptor oligonucleotide sequences with one at each end of the product, at least one oligonucleotide barcode sequence, an X1 oligonucleotide sequence, an X2 oligonucleotide sequence, and (ii) a complementary sequence to the sequence in (i);

(b) amplifying the double-stranded DNA products of (a) with a second amplification primer set comprising a plurality of first and second sequencing platform tag-containing oligonucleotides that each comprise either:

(i) a first sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, or

(ii) a second sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence,

wherein amplifying takes place under conditions that promote amplification of both strands of the separated double-stranded DNA product of (a), to obtain a library of rearranged DNA sequences encoding a plurality of adaptive immune receptors for sequencing; and


(c) sequencing the DNA library obtained in (b), wherein each of the sequences in the DNA library comprises a unique oligonucleotide barcode sequence, thereby labeling each sequence with an unique identifiable barcode sequence.

5. The method of claim 4, wherein a plurality of oligonucleotides in the second amplification primer set each further comprises either or both of:

(i) a sample-identifying barcode oligonucleotide which comprises a third barcode oligonucleotide B5 comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides having a sequence that is distinct from B1 and B2, wherein in the first sequencing platform tag-containing oligonucleotide B5 is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence, and wherein in the second sequencing platform tag-containing oligonucleotide B3 is situated between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence; and

(ii) a spacer oligonucleotide of any sequence of 1 to 20 contiguous nucleotides, wherein said spacer oligonucleotide is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence in the first sequencing platform tag-containing oligonucleotide, and between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence in the second sequencing platform tag-containing oligonucleotide.

6. An oligonucleotide primer composition, comprising a plurality of oligonucleotide sequences having a general formula (I):



wherein:

U1 comprises an oligonucleotide sequence which comprises a first universal adaptor oligonucleotide sequence,

B1 comprises an oligonucleotide sequence that comprises a first oligonucleotide barcode sequence of n contiguous nucleotides, wherein n is at least 6 nucleotides, and

X comprises either (i) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (ii) an oligonucleotide comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences, X comprises a unique oligonucleotide sequence.

7. The composition of claim 6, wherein the plurality of oligonucleotide sequences comprises up to 4^n unique B1 oligonucleotide sequences,

8. The composition of claim 6, wherein n is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous nucleotides.

9. The composition as in any one of claims 6-8, wherein X comprises an oligonucleotide sequence comprising at least 20, 30, 40 or 50 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof.


10. The composition as in any one of claims 6-9, wherein X comprises an oligonucleotide sequence comprising not more than 70, 60, or 55 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof.

11. The composition as in any one of claims 6-8, wherein X comprises an oligonucleotide sequence comprising at least 16-50 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof.

12. The composition as in any one of claims 6-8 or 11, wherein X comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof.

13. The composition as in any one of claims 6-12, wherein X is capable of hybridizing to a V region encoding gene sequence.

14. The composition as in any one of claims 6-12, wherein X is capable of hybridizing to a J region encoding gene sequence.
15. The composition as in any one of claims 6-14, wherein B1 is a unique tag for identifying individual rearranged TCR or Ig encoding sequences.
16. The composition of claim 15, wherein B1 is used to identify individual rearranged TCR or Ig encoding sequences derived from a single cell.
17. The composition as in any one of claims 6-16, wherein U1 comprises SEQ ID NOs: 1710-1731.
18. The composition as in any one of claims 6-17, wherein B1 comprises sequences listed in Table 8.
19. The composition as in any one of claims 6-18, wherein X comprises SEQ ID NOs: 1631-1643 or 1696-1708.
20. The composition as in any one of claims 6-18, wherein X comprises SEQ ID NOs: 1644-1695.
21. The composition as in any one of claims 6-18, wherein X comprises SEQ ID NOs: 5613-5625.
22. The composition as in any one of claims 6-16, wherein said plurality of oligonucleotide sequences comprises SEQ ID NOs: 5626-5685.
23. The composition as in any one of claims 6-16, wherein said plurality of oligonucleotide sequences comprises SEQ ID NOs: 1-1630.
24. The composition as in any one of claims 6-23, further comprising: a second plurality of oligonucleotide sequences comprising a general formula (II):



wherein P1 comprises a sequencing platform-specific oligonucleotide,

wherein S1 comprises a sequencing platform tag-containing oligonucleotide sequence;

wherein B2 comprises an oligonucleotide barcode sequence and wherein said oligonucleotide barcode sequence can be used to identify a sample source, and

wherein U1 comprises said first universal adaptor oligonucleotide sequence.

25. The composition of claim 24, wherein said second plurality of oligonucleotide sequences comprises SEQ ID NOs: 5686-5877.

26. An oligonucleotide primer composition for a first amplification primer set comprising:

(A) a plurality of first oligonucleotide sequences of a general formula

(III):

5'-U1-B1_n-X1-3' (III),

wherein:

(i) U1 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence,

(ii) B1 comprises an oligonucleotide sequence comprising a first oligonucleotide barcode sequence of n contiguous nucleotides, wherein n is 0 or 6 to 20,

(iii) X1 comprises either (a) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (b) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences X1 comprises a unique oligonucleotide sequence; and

(B) a plurality of second oligonucleotide sequences of a general formula

(IV):

5'-U2-B2_m-X2-3' (IV),

wherein:

(i) U2 comprises an oligonucleotide sequence comprising a second universal adaptor oligonucleotide sequence,

(ii) B2 comprises an oligonucleotide sequence comprising a second oligonucleotide barcode sequence of m contiguous nucleotides, wherein m is 0 or 6 to 20,

(iii) X2 comprises (a) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor variable (V) region encoding gene sequence, or the complement thereof, or (b) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the

complement thereof, and in each of the plurality of oligonucleotide sequences X1 comprises a unique oligonucleotide sequence,

wherein n and m are independent of each other, and in said first and second pluralities of oligonucleotides, m and n are not both zero, and

wherein if X1 comprises an oligonucleotide sequence comprising an adaptive immune receptor V region encoding gene sequence, then X2 comprises an oligonucleotide sequence comprising an adaptive immune receptor J region encoding gene sequence, and if X1 comprises an oligonucleotide sequence comprising an adaptive immune receptor J region encoding gene sequence, then X2 comprises an oligonucleotide sequence comprising an adaptive immune receptor V region encoding gene sequence.

27. The composition of claim 26, wherein X1 or X2 comprises an oligonucleotide sequence comprising at least 20, 30, 40 or 50 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof.

28. The composition as in any one of claims 26-27, wherein X1 or X2 comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor V region encoding gene sequence, or said complement thereof.

29. The composition of claim 26, wherein X1 or X2 comprises an oligonucleotide sequence comprising at least 16-50 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof.

30. The composition as in any one of claims 26 or 29, wherein X1 or X2 comprises an oligonucleotide sequence comprising not more than 70, 60 or 55 contiguous nucleotides of said adaptive immune receptor J region encoding gene sequence, or said complement thereof.

31. The composition of claim 26, wherein B1 is a unique tag for identifying an individual rearranged TCR or Ig encoding sequence.

32. The composition of claim 26, wherein B2 is a unique tag for identifying an individual rearranged TCR or Ig encoding sequence.

33. The composition of claim 26, wherein U1 or U2 comprises SEQ ID NOs: 1710-1731.

34. The composition of claim 26, wherein B1 or B2 comprises sequences listed in Table 8.

35. The composition of claim 26, wherein X1 or X2 comprises SEQ ID NOs: 1631-1643 or 1696-1708.

36. The composition of claim 26, wherein X1 or X2 comprises SEQ ID NOs: 1644-1695.

37. The composition of claim 26, wherein X1 or X2 comprises SEQ ID NOs: 5613-5625.

38. The composition of claim 26, wherein said plurality of first or second oligonucleotide sequences comprise SEQ ID NOs: 5626-5685.

39. The composition of claim 26, wherein said plurality of first or second oligonucleotide sequences comprise SEQ ID NOs: 1-1630.

40. The composition as in any of claims 26-39, wherein the plurality of oligonucleotide sequences comprises up to 4^n unique B1 oligonucleotide sequences.

41. The composition as in any of claims 26-40, wherein the plurality of oligonucleotide sequences comprises up to 4^m unique B2 oligonucleotide sequences.

42. An oligonucleotide amplification primer composition, comprising:

(A) a first oligonucleotide amplification primer set comprising a plurality of oligonucleotide sequences of a general formula (V):

$$U1/2 - B1 - X1 \quad (V),$$

wherein U1/2 comprises an oligonucleotide sequence comprising a first universal adaptor oligonucleotide sequence when B1 is present, or a second universal adaptor oligonucleotide sequence when B1 is nothing, and

wherein B1 comprises an oligonucleotide that comprises either nothing or a first oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides, and

wherein X1 comprises either: (1) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor V region encoding gene sequence, or the complement thereof, or (2) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of (i) an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, or (ii) an adaptive immune receptor constant (C) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U1/2-B1-X1, X1 comprises a unique oligonucleotide sequence; and

(B) a second oligonucleotide amplification primer set comprising a plurality of oligonucleotide sequences of a general formula (VI):

$$U3/4 - B2 - X2 \quad (VI),$$

wherein U3/4 comprises an oligonucleotide sequence comprising a third universal adaptor oligonucleotide sequence when B2 is present or a fourth universal adaptor oligonucleotide sequence when B2 is nothing, and

wherein B2 comprises an oligonucleotide sequence comprising either nothing or a second oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is the same as B1, and

wherein X2 comprises either (1) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor V region encoding gene sequence, or the complement thereof, or (2) an oligonucleotide sequence comprising at least 15 and not more than 80 contiguous nucleotides of an adaptive immune receptor joining (J) region encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U3/4-B2-X2, X2 comprises a unique oligonucleotide sequence.

43. The method of claim 42, wherein U3 has the same sequence as U1 or U2.

44. The method of claim 42, wherein U4 has the same sequence as U1 or U2.

45. A method for labeling individual rearranged DNA sequences encoding a plurality of adaptive immune receptors in a biological sample that comprises lymphoid cells of a subject, the method comprising:

(a) amplifying said rearranged DNA sequences using a first amplification primer set comprising an oligonucleotide primer composition of claims 1-36 under conditions that promote amplification to obtain double-stranded DNA products that each comprise (i) a sequence comprising at least one universal adaptor oligonucleotide sequence, at least one oligonucleotide barcode sequence, and at least one of an X, X1 or X2 oligonucleotide sequence, and (ii) a complementary sequence to the sequence in (i);

(b) amplifying the double-stranded DNA products of (a) with a second amplification primer set comprising a plurality of first and second sequencing platform tag-containing oligonucleotides that each comprise either:

(i) a first sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, or

(ii) a second sequencing platform tag-containing oligonucleotide comprising an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second

sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence,

wherein amplifying takes place under conditions that promote amplification of both strands of the separated double-stranded DNA product of (a), to obtain a library of rearranged DNA sequences encoding a plurality of adaptive immune receptors for sequencing; and

(c) sequencing the DNA library obtained in (b), wherein each of the sequences in the DNA library comprises a unique oligonucleotide barcode sequence, thereby labeling each sequence with an unique identifiable barcode sequence.

46. The method of claim 43, wherein a plurality of oligonucleotides in the second amplification primer set each further comprises either or both of:

(i) a sample-identifying barcode oligonucleotide which comprises a third barcode oligonucleotide B3 comprising an oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides having a sequence that is distinct from B1 and B2, wherein in the first sequencing platform tag-containing oligonucleotide B3 is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence, and wherein in the second sequencing platform tag-containing oligonucleotide B3 is situated between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence; and

(ii) a spacer oligonucleotide of any sequence of 1 to 20 contiguous nucleotides, wherein said spacer oligonucleotide is situated between the first universal adaptor oligonucleotide and the first sequencing platform-specific oligonucleotide sequence in the first sequencing platform tag-containing oligonucleotide, and between the second universal adaptor oligonucleotide and the second sequencing platform-specific oligonucleotide sequence in the second sequencing platform tag-containing oligonucleotide.

47. A method for labeling individual rearranged DNA sequences or mRNA sequences transcribed therefrom encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising:

contacting (A) a first plurality of individual microdroplets that each contain a single lymphoid cell or genomic DNA isolated therefrom or complementary DNA (cDNA) that has been reverse transcribed from messenger RNA (mRNA) of a single lymphoid cell, with (B) a second plurality of individual microdroplets that each contain:

(i) a first oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a first polypeptide of an adaptive immune receptor heterodimer, and

(ii) a second oligonucleotide amplification primer set that is capable of amplifying a rearranged DNA sequence encoding a second polypeptide of the adaptive immune receptor heterodimer,

wherein:

(I) the first oligonucleotide amplification primer set comprises a composition of U1/2-B1-X1 of claim 42, and

(II) the second oligonucleotide amplification primer set comprises a composition of U3/4-B2-X2 of claim 42;

providing conditions for a time sufficient such that a plurality of fusion events occur between one of said first microdroplets and one of said second microdroplets to produce a plurality of fused microdroplets; and

providing conditions that permit amplification of the genomic DNA, or the cDNA that has been reverse transcribed from mRNA, using the first and second oligonucleotide amplification primer sets within the plurality of fused microdroplets,

wherein each of one or more of said plurality of fused microdroplets comprises:

a first double-stranded DNA product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and

a second double-stranded DNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer,

thereby upon amplification of the genomic DNA, or the cDNA that has been reverse transcribed from mRNA, labeling each of the individual rearranged DNA sequences or mRNA sequences transcribed therefrom with an oligonucleotide barcode sequence.

48. The method of claim 47, further comprising: disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of said first and second double-stranded DNA products.

49. The method of claim 48, further comprising: contacting the mixture of the first and second double-stranded DNA products with a third amplification primer set and a fourth amplification primer set,

wherein the third amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence,

wherein the fourth amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence, and

wherein said step of contacting takes place under conditions and for a time sufficient to amplify both strands of the first and second double-stranded DNA products, to obtain a DNA library for sequencing.

50. The method of claim 49, further comprising sequencing the DNA library to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

51. The method of claim 49, wherein the third and fourth amplification primer sets are the same.

52. A method for labeling individual rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising:

contacting (A) a first plurality of individual microdroplets that each contain complementary DNA (cDNA) that has been reverse transcribed from messenger RNA (mRNA) of a single lymphoid cell, with (B) a second plurality of individual microdroplets that each contain:

(i) a first oligonucleotide amplification primer set that is capable of amplifying a first cDNA sequence encoding a first polypeptide of an adaptive immune receptor heterodimer, and

(ii) a second oligonucleotide amplification primer set that is capable of amplifying a second cDNA sequence encoding a second polypeptide of the adaptive immune receptor heterodimer,

wherein:

(I) the first oligonucleotide amplification primer set comprises a composition of U1/2-B1-X1 of claim 42;

and

(II) the second oligonucleotide amplification primer set comprises a composition of U3/4-B2-X2 of claim 42,

providing conditions for a time sufficient for a plurality of fusion events between one of said first microdroplets and one of said second microdroplets to produce a plurality of fused microdroplets and conditions that permit amplification of the cDNA that has been reverse transcribed from mRNA of a single lymphoid cell, using the first and second oligonucleotide amplification primer sets within the plurality of fused microdroplets, wherein each of one or more of said plurality of fused microdroplets comprises:

a first double-stranded DNA product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first

polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and

a second double-stranded DNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer;

thereby upon amplification of the cDNA, uniquely labeling each of the individual rearranged cDNA sequences with a unique oligonucleotide barcode sequence.

53. The method of claim 52, further comprising: disrupting the plurality of fused microdroplets to obtain a heterogeneous mixture of said first and second double-stranded DNA products.

54. The method of claim 53, further comprising: contacting the mixture of first and second double-stranded DNA products with a third amplification primer set and a fourth amplification primer set,

wherein the third amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence, and

wherein the fourth amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is

linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence,

wherein said step of contacting takes place under conditions and for a time sufficient to amplify both strands of the first and second double-stranded DNA products, to obtain a DNA library for sequencing.

55. The method of claim 54, further comprising sequencing the DNA library to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

56. The method of claim 54, wherein the third amplification primer set is identical to the fourth amplification primer set.

57. The method as in any one of claims 47 or 52, wherein either or both of:

(1) the first oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the first polypeptide, a rearranged DNA sequence encoding a first complementarity determining region-3 (CDR3) of the first polypeptide; and

(2) the second oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the second polypeptide, a rearranged DNA sequence encoding a second complementarity determining region-3 (CDR3) of the second polypeptide.

58. The method of any one of claims 47 or 52, wherein:

(a) the first polypeptide of the adaptive immune receptor heterodimer is a TCR alpha (TCRA) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR beta (TCRB) chain, or

(b) the first polypeptide of the adaptive immune receptor heterodimer is a TCR gamma (TCRG) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR delta (TCRD) chain, or

(c) the first polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin heavy (IGH) chain and the second polypeptide of the

adaptive immune receptor heterodimer is selected from an immunoglobulin light IGL or IGK chain.

59. The method of claim 58, wherein if the first polypeptide of the adaptive immune receptor heterodimer is an IGH chain and the second polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin light IGL or IGK chain, then three different amplification primer sets are used comprising: a first oligonucleotide amplification primer set for IGH, a second oligonucleotide amplification primer set for IGK, and a third oligonucleotide amplification primer set for IGL.

60. The method of claim 52, wherein the second plurality of individual microdroplets each further comprise: a third oligonucleotide primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule and that comprises a composition comprising a plurality of oligonucleotide sequences having a general formula (VII):

$$U5/6 - B - X3 \quad (VII),$$

wherein U5/6 comprises a fifth universal adaptor oligonucleotide sequence when B is present or a sixth universal adaptor oligonucleotide sequence when B is nothing,

B comprises B1 or B2, and

X3 comprises an oligonucleotide that is one of (i) a forward primer of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and (ii) a reverse primer of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U5/6-B-X3, X3 comprises a unique oligonucleotide sequence.

61. The method of claim 60, wherein the lymphocyte status indicator molecule comprises one or more of FoxP3, CD4, CD8, CD11a, CD18, CD21, CD25, CD29, CCD30, CD38, CD44, CD45, CD45RA, CD45RO, CD49d, CD62, CD62L, CD69, CD71, CD103, CD137 (4-1BB), CD138, CD161, CD294, CCR5, CXCR4, IgG1-4 H-chain constant region, IgA H-chain constant region, IgE H-chain constant region, IgD H-chain constant region, IgM H-chain constant region, HLA-DR, IL-2, IL-5, IL-6, IL-9, IL-10, IL-12, IL-13, IL-15, IL-21, TGF- β , TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9 and TLR10.

62. The method as in any one of claims 50 or 55, further comprising:

(a) sorting the data set of sequences according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode;

(b) sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset;

(c) clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets; and

(d) identifying as originating from the same cell sequences that are members of an X1 and an X2 sequence cluster set that belong to the same one or more barcode sequence sets.

63. A method for determining rearranged DNA sequences encoding first and second polypeptide sequences of an adaptive immune receptor heterodimer in a single lymphoid cell, comprising:

(1) distributing cells of a cell suspension that comprises a population of lymphoid cells of a subject, amongst a plurality of containers that are capable of containing said cells, to obtain a plurality of containers that each contain a subpopulation of the lymphoid cells that comprises one lymphoid cell or a plurality of lymphoid cells;

(2) contacting each of said plurality of containers, under conditions and for a time sufficient to promote reverse transcription of messenger RNA (mRNA) in the lymphoid cells in the plurality of containers, with a first and a second oligonucleotide reverse transcription primer set, wherein (A) the first oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of first mRNA sequences encoding a plurality of polypeptides of a first adaptive immune receptor heterodimer, and (B) the second oligonucleotide reverse transcription primer set is capable of reverse transcribing a plurality of second mRNA sequences encoding a plurality of polypeptides of a second adaptive immune receptor heterodimer,

and wherein:

(I) the first oligonucleotide reverse transcription primer set comprises a composition comprising a general formula of U1/2-B1-X1 of claim 32, and

(II) the second oligonucleotide reverse transcription primer set comprises a composition comprising a general formula U3/4-B2-X2 of claim 32, and

wherein said step of contacting taking place under conditions and for a time sufficient to obtain in each of one or more of said plurality of containers:

a first reverse-transcribed complementary DNA (cDNA) product that comprises at least one first universal adaptor oligonucleotide sequence, at least one first oligonucleotide barcode sequence, at least one X1 oligonucleotide V region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, at least one second universal adaptor oligonucleotide sequence, and at least one X1 oligonucleotide J region or C region encoding gene sequence of said first polypeptide of the adaptive immune receptor heterodimer, and

a second reverse-transcribed cDNA product that comprises at least one third universal adaptor oligonucleotide sequence, at least one second oligonucleotide barcode sequence, at least one X2 oligonucleotide V region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer, at least one fourth universal adaptor oligonucleotide sequence, and at least one X2 oligonucleotide J region or C region encoding gene sequence of said second polypeptide of the adaptive immune receptor heterodimer;

(3) combining the first and second reverse-transcribed cDNA products from the plurality of containers to obtain a mixture of reverse-transcribed cDNA products;

(4) contacting the mixture of first and second reverse-transcribed cDNA products of (3) with a first oligonucleotide amplification primer set and a second oligonucleotide amplification primer set, wherein the first amplification primer set comprises (i) a plurality of first sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the first universal adaptor oligonucleotide and a first sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the first universal adaptor oligonucleotide sequence, and (ii) a plurality of second sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the second universal adaptor oligonucleotide sequence and a second sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the second universal adaptor oligonucleotide sequence,

and wherein the second oligonucleotide amplification primer set comprises (i) a plurality of third sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the third universal adaptor oligonucleotide and a third sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the third universal adaptor oligonucleotide sequence, and (ii) a plurality of fourth sequencing platform tag-containing oligonucleotides that each comprise an oligonucleotide sequence that is capable of specifically hybridizing to the fourth universal adaptor oligonucleotide sequence and a fourth sequencing platform-specific oligonucleotide sequence that is linked to and positioned 5' to the fourth universal adaptor oligonucleotide sequence,

said step of contacting taking place under conditions and for a time sufficient to amplify both of the first and second reverse-transcribed cDNA products of (2), to obtain a DNA library for sequencing; and

(5) sequencing the DNA library obtained in (3) to obtain a data set of sequences encoding the first and second polypeptide sequences of the adaptive immune receptor heterodimer.

64. The method of claim 63, wherein step (5) further comprises:

(a) sorting the data set of sequences according to oligonucleotide barcode sequences identified therein to obtain a plurality of barcode sequence sets each having a unique barcode;

(b) sorting each barcode sequence set of (a) into an X1 sequence-containing subset and an X2 sequence-containing subset;

(c) clustering members of each of the X1 and X2 sequence-containing subsets according to X1 and X2 sequences to obtain one or a plurality of X1 sequence cluster sets and one or a plurality of X2 sequence cluster sets, respectively, and error-correcting single nucleotide barcode sequence mismatches within any one or more of said X1 and X2 sequence cluster sets;

(d) identifying each first and second adaptive immune receptor heterodimer polypeptide encoding sequence based on known X1 and X2 sequences, wherein each X1 sequence and each X2 sequence is associated with one or a plurality of unique B sequences to identify the container from which each B sequence-associated X1 sequence and each B sequence-associated X2 sequence originated; and

(e) combinatorically matching B sequence-associated X1 and X2 sequences of (d) as being of common clonal origin based on a probability of B sequences that are coincident with common first and second adaptive immune receptor heterodimer polypeptide encoding sequences, and therefrom determining that rearranged DNA sequences encoding first and second polypeptide sequences of the adaptive immune receptor heterodimer originated in a single lymphoid cell.

65. The method of claim 63, wherein either or both of:

(1) the first oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the first polypeptide, a rearranged DNA sequence encoding a first complementarity determining region-3 (CDR3) of the first polypeptide; and

(2) the second oligonucleotide amplification primer set is capable of amplifying, in the rearranged DNA sequence encoding the second polypeptide, a rearranged DNA sequence encoding a second complementarity determining region-3 (CDR3) of the second polypeptide.

66. The method of claim 65, wherein:

(a) the first polypeptide of the adaptive immune receptor heterodimer is a TCR alpha (TCRA) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR beta (TCRB) chain, or

(b) the first polypeptide of the adaptive immune receptor heterodimer is a TCR gamma (TCRG) chain and the second polypeptide of the adaptive immune receptor heterodimer is a TCR delta (TCRD) chain, or

(c) the first polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin heavy (IGH) chain and the second polypeptide of the adaptive immune receptor heterodimer is an immunoglobulin light (IGL, IGK, or both IGL and IGK) chain.

67. The method of claim 63, wherein one or more of the containers comprises a third oligonucleotide amplification primer set that is capable of amplifying a third cDNA sequence that encodes a lymphocyte status indicator molecule and that comprises a composition comprising a plurality of oligonucleotides having a plurality of oligonucleotide sequences of general formula (VIII):

U5/6 - B3 - X3 (VIII),

in which U5/6 comprises an oligonucleotide which comprises a fifth universal adaptor oligonucleotide sequence when B3 is present or a sixth universal adaptor oligonucleotide sequence when B3 is nothing,

B3 comprises an oligonucleotide that comprises either nothing or a third oligonucleotide barcode sequence of 6 to 20 contiguous nucleotides that is either the same as or different than at least one of B1 or B2, and

X3 comprises an oligonucleotide that is one of (i) a forward primer polynucleotide of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and (ii) a reverse primer polynucleotide of 15-80 contiguous nucleotides of a lymphocyte status indicator molecule encoding gene sequence, or the complement thereof, and in each of the plurality of oligonucleotide sequences of general formula U5/6-B3-X3, X3 comprises a unique oligonucleotide sequence.

68. The method of claim 67, wherein the lymphocyte status indicator molecule comprises one or more of FoxP3, CD4, CD8, CD11a, CD18, CD21, CD25, CD29, CCD30, CD38, CD44, CD45, CD45RA, CD45RO, CD49d, CD62, CD62L, CD69, CD71, CD103, CD137 (4-1BB), CD138, CD161, CD294, CCR5, CXCR4, IgG1-4 H-chain constant region, IgA H-chain constant region, IgE H-chain constant region, IgD H-chain constant region, IgM H-chain constant region, HLA-DR, IL-2, IL-5, IL-6, IL-9, IL-10, IL-12, IL-13, IL-15, IL-21, TGF- β , TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9 and TLR10.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

