

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2015231804 B2

- (54) Title
Transgenic maize event MON 87419 and methods of use thereof
- (51) International Patent Classification(s)
A01H 5/00 (2006.01) **C12N 15/82** (2006.01)
C12N 5/04 (2006.01)
- (21) Application No: **2015231804** (22) Date of Filing: **2015.03.10**
- (87) WIPO No: **WO15/142571**
- (30) Priority Data
- (31) Number **61/968,342** (32) Date **2014.03.20** (33) Country **US**
- (43) Publication Date: **2015.09.24**
(44) Accepted Journal Date: **2021.02.25**
- (71) Applicant(s)
Monsanto Technology LLC
- (72) Inventor(s)
Goley, Michael E;Burns, Wen C.;Huang, Jintai;McCann, Melinda C.;Shao, Aihua;Sparks, Oscar C.;Stoecker, Martin A.;Wei, Liping
- (74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, GPO Box 1374, Brisbane, QLD, 4001, AU
- (56) Related Art
WO 2008051633 A2
US 20120255050 A1
WO 2011066382 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 September 2015 (24.09.2015)

(10) International Publication Number
WO 2015/142571 A8

(51) International Patent Classification:
A01H 5/00 (2006.01) *C12N 5/04* (2006.01)
C12N 15/82 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2015/019663

(22) International Filing Date:
10 March 2015 (10.03.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/968,342 20 March 2014 (20.03.2014) US

(71) Applicant: **MONSANTO TECHNOLOGY LLC**
[US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US).

(72) Inventors: **BURNS, Wen, C.**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **GOLEY, Michael, E.**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **HUANG, Jintai**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **MCCANN, Melinda, C.**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **SHAO, Aihua**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **SPARKS, Oscar, C.**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **STOECKER, Martin, A.**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). **WEL, Liping**; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US).

(74) Agent: **HANSON, Robert, E.**; Dentons US LLP, 233 South Wacker Drive, Suite 7800, Chicago, IL 60606-6306 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(48) Date of publication of this corrected version:

17 December 2015

(15) Information about Correction:
see Notice of 17 December 2015

WO 2015/142571 A8

(54) Title: TRANSGENIC MAIZE EVENT MON 87419 AND METHODS OF USE THEREOF

(57) Abstract: The invention provides recombinant DNA molecules that are unique to the maize MON 87419 event and transgenic maize plants, plant parts, seeds, cells, and agricultural products containing the MON 87419 event as well as methods of using and detecting the maize MON 87419 event. Transgenic maize plants containing the MON 87419 event exhibit tolerance to dicamba and glufosinate herbicides.

TRANSGENIC MAIZE EVENT MON 87419 AND METHODS OF USE THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS**

[001] This application claims the benefit of priority to U.S. Provisional Application Serial No. 61/968,342 filed March 20, 2014.

INCORPORATION OF SEQUENCE LISTING

[002] The sequence listing that is contained in the file named “MONS362WO.txt”, which is 29.4 KB (measured in MS-Windows) and created on March 9, 2015, is filed herewith by electronic submission and incorporated herein by reference.

FIELD OF THE INVENTION

[003] The invention relates to recombinant DNA molecules that are unique to the transgenic maize event MON 87419. The invention also relates to transgenic maize plants, parts, seeds, cells, and agricultural products containing the maize MON 87419 event as well as methods of using the same. Transgenic maize plants containing the maize MON 87419 event exhibit tolerance to dicamba and glufosinate herbicides.

BACKGROUND OF THE INVENTION

[004] Maize (*Zea mays*) is an important crop in many areas of the world. The methods of biotechnology have been applied to this crop in order to produce maize with desirable traits. One such desirable trait is herbicide tolerance. Expression of a heterologous gene, also known as a transgene, for herbicide tolerance in a plant can confer herbicide tolerance on the plant. However, the expression of a transgene, and therefore its effectiveness, may be influenced by many different factors including the orientation and composition of the cassette driving expression of the individual transgene transferred to the plant chromosome and the chromosomal location and the genomic result of the transgene insertion. This is complicated further in transgenic plants with multiple molecularly-linked transgenes, each conferring a separate trait. In such a situation, proper expression of each of the molecularly-linked transgenes in the plant must result from the same transgene insertion (also called a multi-gene event). In such cases, it is

necessary to design and test multiple expression cassettes, each with a different configuration of transgenes and expression elements, and then to produce and analyze a large number of individual plant transformation events through multiple generations of plants in order to select the transgenic event having superior properties relative to the each of the desirable traits and the optimal phenotypic and agricultural characteristics necessary to make it suitable for commercial purposes. Such selection requires extensive molecular characterization as well as greenhouse and field trials over multiple years, in multiple locations, and under a variety of conditions so that a significant amount of agronomic, phenotypic, and molecular data may be collected. The resulting data and observations must then be analyzed by teams of scientists and agronomists with the goal of selecting the event suitable for commercial agricultural use across a wide range of germplasm and in a variety of field conditions. Once selected, the commercial event conferring the desirable traits may be introgressed into other genetic backgrounds using plant breeding methods, thus producing a number of different crop varieties that contain the desirable trait and are suitably adapted to specific local growing conditions.

[005] To make a transgenic plant containing a single transformation event, a portion of a recombinant DNA construct is transferred into the genome of a maize cell using plant transformation techniques. This maize cell is subsequently used to produce a unique R₀ plant, which can then be used to produce transgenic progeny plants. The genome of the progeny plants contains the unique event, and these plants can be tested for the desired trait(s) as well as for agronomic performance. The effectiveness of an event can be impacted by *cis* and/or *trans* factors relative to the integration site in the transformation event. The phenotype conferred by the event can also be impacted by the size and design of the DNA construct, which can vary by the combination of genetic elements in an expression cassette, number of transgenes, number of expression cassettes, and configuration of such elements and such cassettes. The performance of a given event can be further complicated by factors such as plant developmental, diurnal, temporal, or spatial patterns of transgene expression; or by extrinsic factors, for example, environmental plant growth conditions, water availability, nitrogen availability, heat, or stress. Thus, the ability to create an event conferring a desirable set of phenotypic traits is not readily predictable.

BRIEF SUMMARY OF THE INVENTION

[006] The invention provides a recombinant DNA molecule containing a sequence selected from the group consisting of SEQ ID NO:1-10. The invention also provides a recombinant DNA derived from a transgenic maize plant or seed containing the maize MON 87419 event, a representative sample of seed comprising the maize MON 87419 event having been deposited as ATCC Accession No. PTA-120860. The invention also provides a recombinant DNA molecule that is an amplicon diagnostic for the presence of DNA derived from the maize MON 87419 event. The invention also provides a DNA molecule that is in a maize plant, cell, seed, progeny plant, or plant part derived from transgenic maize comprising the maize MON 87419 event.

[007] The invention provides a DNA molecule having sufficient length of contiguous DNA sequence of SEQ ID NO:10 to function as a DNA probe that hybridizes under stringent hybridization conditions with a DNA molecule comprising a DNA sequence selected from the group consisting of SEQ ID NO:1-10 and does not hybridize under the stringent hybridization conditions with a DNA molecule not comprising a DNA sequence selected from the group consisting of SEQ ID NO:1-10. The invention also provides a pair of DNA molecules comprising a first DNA molecule and a second DNA molecule, wherein the first DNA molecule is a fragment of SEQ ID NO:9 and the second DNA molecule is a fragment of the maize genomic DNA of the maize MON 87419 event, and wherein the first and second DNA molecules each comprise a DNA sequence of sufficient length of contiguous nucleotides to function as DNA primers when used together in an amplification reaction with DNA containing the maize MON 87419 event to produce an amplicon diagnostic for the maize MON 87419 event in a sample.

[008] The invention provides a method of detecting the presence of the maize MON 87419 event in a sample of DNA by contacting the sample with a DNA probe, subjecting the sample and the DNA probe to stringent hybridization conditions, and detecting hybridization of the DNA probe to a DNA molecule in the sample, where the hybridization of the DNA probe to the DNA molecule indicates the presence of the maize MON 87419 event in the sample of DNA.

[009] The invention provides a method of detecting the presence of the maize MON 87419 event in a sample of DNA by contacting the sample with a pair of DNA primers,

performing an amplification reaction sufficient to produce a DNA amplicon comprising a sequence selected from the group consisting of SEQ ID NO:1-8 and SEQ ID NO:10, and detecting the presence of the DNA amplicon in the reaction, wherein the presence of the DNA amplicon in the reaction indicates the presence of the maize MON 87419 event in the sample of DNA.

[0010] The invention provides a DNA detection kit containing at least one DNA molecule comprising a DNA sequence of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as a primer or probe specific for detecting the presence of the maize MON 87419 event in a sample of DNA.

[0011] The invention provides a recombinant maize plant, seed, cell, plant part, or commodity product comprising a DNA molecule having a DNA sequence selected from the group consisting of SEQ ID NO:1-10. The invention also provides a transgenic maize plant, seed, or cell that is tolerant to glufosinate or dicamba, or glufosinate and dicamba herbicides. The invention also provides a transgenic maize plant, seed, cell, plant part, or commodity product containing the maize MON 87419 event. The invention also provides a transgenic maize plant or seed that is a hybrid having at least one parent plant that comprised the maize MON 87419 event.

[0012] The invention provides a method for controlling weeds in an area comprising planting transgenic maize comprising the maize MON 87419 event in an area and applying an effective dose of dicamba or glufosinate or dicamba and glufosinate herbicides to control the weeds in the area without injuring the transgenic maize. The invention also provides a method for controlling weeds by applying an effective dose of glufosinate herbicide of about 0.1 pounds acid equivalent per acre to about 16 pounds acid equivalent per acre of glufosinate herbicide over a growing season. The invention also provides a method for controlling weeds by applying an effective dose of glufosinate herbicide of about 0.4 pounds acid equivalent per acre to about 1.59 pounds acid equivalent per acre of glufosinate herbicide over a growing season. The invention also provides a method for controlling weeds by applying an effective dose of dicamba herbicide of about 0.1 pounds acid equivalent per acre to about 16 pounds acid equivalent per acre of dicamba herbicide over a growing season. The invention also provides a method for controlling weeds by applying an effective dose of dicamba herbicide is about 0.5 pounds acid equivalent per acre to about 2 pounds acid equivalent per acre of dicamba herbicide over a growing season.

[0013] The invention provides a method of producing a transgenic maize plant that is tolerant to glufosinate and dicamba herbicides by sexually crossing a transgenic maize plant comprising the maize MON 87419 event with a second maize plant, collecting the seed produced, growing the seed to produce progeny plants, treating the progeny plants with glufosinate or dicamba, or glufosinate and dicamba herbicides, and selecting a progeny plant that is tolerant to glufosinate and dicamba herbicides. The invention also provides a method of producing a transgenic maize plant that is tolerant to glufosinate and dicamba herbicides by selfing a transgenic maize plant comprising the maize MON 87419 event, collecting the seed produced, growing the seed to produce progeny plants, treating the progeny plants with glufosinate or dicamba, or dicamba and glufosinate herbicides, and selecting a progeny plant that is tolerant to glufosinate and dicamba herbicides.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] Figure 1: illustrates the organization of the transgene insert in the genome of a maize plant comprising maize event MON 87419. The horizontal lines correspond to the relative positions of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10; the thick arrows labeled SQ26644 and SQ26645 represent the approximate position of a pair of primers used to identify transgenic maize containing the maize MON 87419 event; the thick, short lines numbered 14 through 22 represent the relative position of unique recombinant construct sequences within the DNA insert (SEQ ID NO:9) and the number refers to the corresponding SEQ ID NO of each, respectively; the thin horizontal arrows represent the relative organization of the two separate expression cassettes of the heterologous transgene inserted DNA of the maize MON 87419 event and the boxes indicate the separate elements of the two expression cassettes; a leading 'P' represents a promoter element, a leading 'L' represents a leader element, a leading 'I' represents an intron, a leading 'TS' represents a chloroplast transit peptide, a leading 'T' represents a 3' transcription termination and polyadenylation element (3' UTR), *pat* represents the coding region for the phosphinothrin acetyl transferase (PAT) protein, and *dmo* represents the coding region for the dicamba mono-oxygenase (DMO) protein.

BRIEF DESCRIPTION OF THE SEQUENCES

[0015] SEQ ID NO:1 is a thirty nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:1 corresponds to nucleotide positions 1232 to 1261 of SEQ ID NO:10.

[0016] SEQ ID NO:2 is a thirty nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:2 corresponds to nucleotide positions 7994 to 8023 of SEQ ID NO:10.

[0017] SEQ ID NO:3 is a sixty nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:3 corresponds to nucleotide positions 1217 to 1276 of SEQ ID NO:10.

[0018] SEQ ID NO:4 is a sixty nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:4 corresponds to nucleotide positions 7979 to 8038 of SEQ ID NO:10.

[0019] SEQ ID NO:5 is a one-hundred nucleotide DNA sequence representing the 5' junction of maize genomic DNA and the transgene insert. SEQ ID NO:5 corresponds to nucleotide positions 1197 to 1296 of SEQ ID NO:10.

[0020] SEQ ID NO:6 is a one-hundred nucleotide DNA sequence representing the 3' junction of maize genomic DNA and the transgene insert. SEQ ID NO:6 corresponds to nucleotide positions 7959 to 8058 of SEQ ID NO:10.

[0021] SEQ ID NO:7 is a 1771 nucleotide DNA sequence representing 1246 nucleotides of the 5' flanking maize genomic DNA and 525 nucleotides of the 5' end of the transgene insert.

[0022] SEQ ID NO:8 is a 1767 nucleotide DNA sequence representing 516 nucleotides of the 3' end of the transgene insert and 1251 nucleotides of the 3' flanking maize genomic DNA.

[0023] SEQ ID NO:9 is a 6762 nucleotide DNA sequence corresponding to the transgene insert of the maize MON 87419 event.

[0024] SEQ ID NO:10 is a 9259 nucleotide DNA sequence corresponding to the maize MON 87419 event; the sequence contains the 5' flanking genomic DNA sequence from positions 1 to 1246, the transgenic DNA insert from positions 1247 to 8008, and the 3' flanking genomic DNA sequence from positions 8009 to 9259.

[0025] SEQ ID NO:11 is a 33 nucleotide DNA sequence corresponding to a primer referred to as SQ26644 and used to identify maize MON 87419 event DNA in a sample; it corresponds to positions 7966 to 7998 of SEQ ID NO:10.

[0026] SEQ ID NO:12 is a 24 nucleotide DNA sequence corresponding to a primer referred to as SQ26645 and used to identify maize MON 87419 event DNA in a sample; it corresponds to positions 8022 to 8045 of SEQ ID NO:10.

[0027] SEQ ID NO:13 is a 19 nucleotide DNA sequence corresponding to a probe referred to as PB11207 and used to identify maize MON 87419 event DNA in a sample; it corresponds to positions 8002 to 8020 of SEQ ID NO:10.

[0028] SEQ ID NOs:14-22 are DNA sequences corresponding to unique sequences within the transgene insert of the maize MON 87419 event.

DETAILED DESCRIPTION OF THE INVENTION

[0029] The following definitions and methods are provided to better define the invention and to guide those of ordinary skill in the art in the practice of the invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

[0030] Modern plant transformation techniques are used to generate genetically engineered plants. The term 'transgenic' may also be used to refer to genetically engineered plants. During the process of generating a transgenic plant, foreign DNA is randomly inserted into the genome of a plant cell. During the transformation procedure, many individual cells are transformed. Due to random integration, a separate and unique DNA recombination event will take place within the genome of each individual transformed plant cell. An entire transgenic plant is then generated from a single individual transgenic cell which necessarily results in every cell of the transgenic plant containing the uniquely inserted DNA as a stable part of its genome. The transgenic herbicide tolerant maize containing the maize MON 87419 event comprises a single insertion of transgenic DNA into the chromosome/genome of the maize germplasm. The maize MON 87419 event was produced by: (i) transformation of thousands of maize plant cells with a nucleic acid construct that includes the transgenes of interest, (ii) regeneration of a population of maize plants each containing a unique transgenic event, and (iii) multi-year testing, screening, and selection to select an event having the desirable agronomic properties, the maize

MON 87419 event. The maize MON 87419 event is characterized by the unique DNA sequence of the insertion of the transgene into the particular location of the maize plant's genome.

[0031] The act of inserting the transgenic DNA into the genome of the maize plant is accomplished by the act of plant transformation and results in the creation of a new transgenic genomic molecular sequence, known as an "event". This sequence is unique to and specific for the event and can be readily identified when compared to the original maize genomic sequence or other transgenic maize events. Molecular analysis of the maize MON 87419 event identified the genomic insertion site of the inserted DNA (SEQ ID NO:9) and the flanking maize genomic DNA sequence immediately adjacent to either side of the inserted DNA (SEQ ID NO:7 and SEQ ID NO:8). This arrangement of the inserted DNA in relation to the surrounding maize plant genome DNA is therefore specific and unique to the transgenic herbicide tolerant maize comprising the maize MON 87419 event. This new genomic molecular sequence (SEQ ID NO:10) is also an integral part of the chromosome of transgenic herbicide tolerant maize plants comprising the maize MON 87419 event and as such is static in the plant and may be passed on to progeny of the plant.

[0032] The present invention also provides progeny of the original transformant comprising the maize MON 87419 event. Such progeny may be produced by selfing of a maize plant comprising the maize MON 87419 event or by sexual outcross between a maize plant comprising the maize MON 87419 event and another plant that does or does not contain the maize MON 87419 event. Such other plant may be a transgenic plant comprising the same or different event(s) or a nontransgenic plant, such as one from a different variety. Even after repeated back-crossing to a recurrent parent, the maize MON 87419 event from the transformed parent is present in the progeny of the cross at the same genomic location.

[0033] As used herein, the term "maize" means *Zea mays* (also referred to as corn) and includes all plant varieties that can be bred with maize.

[0034] The invention provides a transgenic herbicide tolerant maize plant containing the maize MON 87419 event that is tolerant to dicamba (3,6-dichloro-2-methoxybenzoic acid) herbicide and glufosinate (2-amino-4-(hydroxymethylphosphinyl) butanoic acid) herbicide. Dicamba is a synthetic auxin herbicide useful for controlling broadleaf weeds. Glufosinate is an organophosphorus herbicide useful for controlling a broad spectrum of annual and perennial grass and broadleaf weeds. The maize MON 87419 event contains a demethylase (*dmo*) gene from

Stenotrophomonas maltophilia that expresses a dicamba mono-oxygenase (DMO) protein to confer tolerance to dicamba herbicide and a bialaphos resistance (*pat*) gene from *Streptomyces viridochromogenes* that expresses the phosphinothricin N-acetyltransferase (PAT) protein to confer tolerance to glufosinate herbicide.

[0035] As used herein, the term “recombinant” refers to a non-natural DNA, protein, or organism that would not normally be found in nature and was created by human intervention. As used herein, a “recombinant DNA molecule” is a DNA molecule comprising a combination of DNA molecules that would not naturally occur together and is the result of human intervention, for example, a DNA molecule that is comprised of a combination of at least two DNA molecules heterologous to each other, such as a DNA molecule that comprises a transgene and the plant genomic DNA adjacent to the transgene. An example of a recombinant DNA molecule is a DNA molecule comprising at least one sequence selected from SEQ ID NO:1-10. As used herein, a “recombinant plant” is a plant that would not normally exist in nature, is the result of human intervention, and contains a transgenic DNA molecule. As a result of such genomic alteration, the recombinant plant is something new and distinctly different from the related wild-type plant. An example of a recombinant plant is a maize plant containing the maize MON 87419 event.

[0036] As used herein, the term “transgene” refers to a DNA molecule artificially incorporated into an organism’s genome as a result of human intervention, such as by plant transformation methods. A transgene may be heterologous to the organism. The term “transgene insert” as used herein refers to the transgene inserted by plant transformation techniques into the maize genome to produce maize event MON 87419. The sequence for this transgene insert is provided as SEQ ID NO:9. The term “transgenic” refers to comprising a transgene, for example a “transgenic plant” refers to a plant comprising a transgene.

[0037] As used herein, the term “heterologous” refers to a first molecule not normally associated with a second molecule or an organism in nature. For example, a DNA molecule may be derived from a first species and inserted into the genome of a second species. The DNA molecule would thus be heterologous to the genome and the organism.

[0038] As used herein, the term “chimeric” refers to a single DNA molecule produced by fusing a first DNA molecule to a second DNA molecule, where neither first nor second DNA molecule would normally be found in that configuration fused to the other. The chimeric DNA molecule is thus a new DNA molecule not normally found in nature. An example of a chimeric

DNA molecule is a DNA molecule comprising at least one sequence selected from SEQ ID NO:1-10.

[0039] The invention provides DNA molecules and their corresponding DNA sequences. As used herein, the terms “DNA” and “DNA molecule” refer to a deoxyribonucleic acid (DNA) molecule. A DNA molecule may be of genomic or synthetic origin, and is by convention from the 5’ (upstream) end to the 3’ (downstream) end. As used herein, the term “DNA sequence” refers to the nucleotide sequence of a DNA molecule. The nomenclature used is that required by Title 37 of the United States Code of Federal Regulations § 1.822 and set forth in the tables in WIPO Standard ST.25 (1998), Appendix 2, Tables 1 and 3. By convention, the DNA sequences of the invention and fragments thereof are disclosed with reference to only one strand of the two complementary DNA sequence strands. By implication and intent, the complementary sequences of the sequences provided here (the sequences of the complementary strand), also referred to in the art as the reverse complementary sequences, are within the scope of the invention and are expressly intended to be within the scope of the subject matter claimed. Thus, as used herein references to SEQ ID NO:1-10 and SEQ ID NO:14-22 and fragments thereof include and refer to the sequence of the complementary strand and fragments thereof.

[0040] As used herein, the term “fragment” refers to a smaller piece of a whole. For example, fragments of SEQ ID NO:10 would include sequences that are at least about 20 consecutive nucleotides, at least about 25 consecutive nucleotides, at least about 30 consecutive nucleotides, at least about 35 consecutive nucleotides, at least about 40 consecutive nucleotides, at least about 45 consecutive nucleotides, at least about 50 consecutive nucleotides, at least about 60 consecutive nucleotides, at least about 70 consecutive nucleotides, at least about 80 consecutive nucleotides, at least about 90 consecutive nucleotides, or at least about 100 consecutive nucleotides of the complete sequence of SEQ ID NO:10.

[0041] The DNA sequence corresponding to the complete DNA sequence of the transgene insert and substantial segments of the maize genome DNA flanking either end of the transgene insert is provided as SEQ ID NO:10. The DNA sequences of the maize genomic DNA physically linked by phosphodiester bond linkage to and therefore flanking the 5’ end of the transgene insert is provided as SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, and SEQ ID NO:7. The DNA sequence of the maize genomic DNA physically linked by phosphodiester bond

linkage to and therefore flanking the 3' end of the transgene insert is provided as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, and SEQ ID NO:8.

[0042] Transgenic maize containing the maize MON 87419 event comprises two regions referred to as junctions. A “junction” is where one end of the transgene insert has been connected to the genomic DNA. A junction spans or extends across a portion of the transgene insert and the adjacent flanking genomic DNA and as such is the connection point of these two as one contiguous molecule. One junction is at the 5' end of the transgene insert and one is at the 3' end of the transgene insert, referred to herein as the 5' and 3' junction, respectively. A “junction sequence” refers to a DNA sequence of any length that spans the 5' or 3' junction of an event. Junction sequences of the maize MON 87419 event are readily apparent to one of skill in the art using SEQ ID NO:10. Examples of junction sequences of the maize MON 87419 event are provided as SEQ ID NO:1-8. Figure 1 illustrates the physical arrangement of SEQ ID NO:1-10 arranged from 5' to 3'. The invention thus provides a DNA molecule that contains at least one of the DNA sequences as set forth in SEQ ID NO:1-8.

[0043] The junction sequences of the maize MON 87419 event may be present as part of the genome of a transgenic maize plant, seed, or cell containing the maize MON 87419 event. The identification of any one or more of SEQ ID NO:1-8 in a sample derived from a transgenic maize plant, plant part, seed, or cell indicates that the DNA was obtained from transgenic maize containing the maize MON 87419 event and is diagnostic for the presence of the maize MON 87419 event.

[0044] The maize MON 87419 event contains sequences which are unique to the transgene insert, specifically SEQ ID NO:14-22. These sequences are unique to the specific chimeric configuration of the various promoters, introns, chloroplast targeting peptides (CTP), 3' termination signal, the *pat* and *dmo* genes within the transgene insert of the event. Figure 1 illustrates the relative position of each of these unique transgene insert sequences with respect to SEQ ID NO:9.

[0045] Provided are exemplary DNA molecules that can be used either as primers or probes for diagnosing the presence in a sample of DNA derived from the maize MON 87419 event. Such primers or probes are specific for a target nucleic acid sequence and as such are useful for the identification of the maize MON 87419 event by the methods described here.

[0046] A “primer” is a DNA molecule that is designed for use in annealing or hybridization methods that involve an amplification reaction. An amplification reaction is an *in vitro* reaction that amplifies template DNA to produce an amplicon. As used herein, an “amplicon” is a DNA molecule that has been synthesized using amplification techniques. Amplicons of the invention have a DNA sequence comprising one or more of SEQ ID NO:1-10, or fragments thereof. A pair of primers may be used with template DNA, such as a sample of maize genomic DNA, in an amplification reaction, such as polymerase chain reaction (PCR), to produce an amplicon, where the amplicon produced would have a DNA sequence corresponding to sequence of the template DNA located between the two sites where the primers hybridized to the template. A primer is typically designed to hybridize to a complementary target DNA strand to form a hybrid between the primer and the target DNA strand. The presence of a primer is a point of recognition by a polymerase to begin extension of the primer using as a template the target DNA strand. Primer pairs refer to use of two primers binding opposite strands of a double stranded nucleotide segment for the purpose of amplifying the nucleotide segment between them. Examples of primer sequences are provided as SEQ ID NO:11 and SEQ ID NO:12. The primer pair provided as SEQ ID NO:11 and SEQ ID NO:12 are useful as a first DNA molecule and a second DNA molecule, where the first DNA molecule is a fragment of SEQ ID NO:9 and the second DNA molecule is a fragment of the maize genomic DNA sequence of SEQ ID NO:10, and each are of sufficient length to function as DNA primers when used together in an amplification reaction with DNA containing the maize MON 87419 event to produce an amplicon diagnostic for the maize MON 87419 event in a sample. The maize genomic DNA sequence of the maize MON 87419 event is provided as positions 1-1246 and 8009-9259 of SEQ ID NO:10.

[0047] A “probe” is a nucleic acid molecule that is complementary to a strand of a target nucleic acid and useful in hybridization detection methods. Probes according to the invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and the detection of such binding can be useful in detecting the presence or absence of the target DNA sequence. A probe may be attached to a conventional detectable label or reporter molecule, such as a radioactive isotope, ligand, chemiluminescent agent, or enzyme. An exemplary DNA sequence useful as a probe for detecting maize MON 87419 event is provided as SEQ ID NO:13.

[0048] Methods for designing and using primers and probes are well known in the art, and DNA molecules comprising fragments of SEQ ID NO:1-10 and useful as primers and probes for detecting the maize MON 87419 event can readily be designed by one of skill in the art.

[0049] The DNA molecules and corresponding DNA sequences provided herein are therefore useful for identifying the maize MON 87419 event in transgenic maize plants, cells, seeds, or parts; selecting maize varieties or hybrids comprising the maize MON 87419 event; and detecting the presence or absence of the maize MON 87419 event in a sample.

[0050] As used herein, the term “isolated” refers to separating a molecule from other molecules normally associated with it in its native or natural state. The term “isolated” thus may refer to a DNA molecule that has been separated from other DNA molecule(s) which normally are associated with it in its native or natural state. Such a DNA molecule may be present in a recombined state, such as a recombinant DNA molecule. Thus, DNA molecules fused to regulatory or coding sequences with which they are not normally associated, for example as the result of recombinant techniques, are considered isolated, even when integrated as a transgene into the chromosome of a cell or present with other DNA molecules.

[0051] The invention provides maize plants, progeny, seeds, plant cells, and plant parts containing the maize MON 87419 event, and commodity products produced using these. A representative sample of transgenic herbicide tolerant maize seed comprising MON 87419 event has been deposited according to the Budapest Treaty with the American Type Culture Collection (ATCC®). The ATCC repository has assigned the Patent Deposit Designation PTA-120860 to the seed of the transgenic herbicide tolerant maize plant containing the maize MON 87419 event. The plants, progeny, seeds, plant cells, plant parts, and commodity products of the invention contain a detectable amount of DNA having at least one of the sequences provided as SEQ ID NO:1-10 and SEQ ID NO:14-22. Plants, progeny, seeds, plant cells, and plant parts of the invention may also contain one or more additional transgenic traits, particularly those introduced by crossing a maize plant containing the maize MON 87419 event with another plant containing the additional transgenic trait(s). Such traits include but are not limited to increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, hybrid seed production, and/or increased herbicide tolerance, in which the trait is measured with respect to a maize plant lacking such transgenic trait.

[0052] Plants of the invention may be used to produce progeny that contain the maize MON 87419 event. As used herein, “progeny” includes any plant, seed, and plant cell comprising the maize MON 87419 event inherited from an ancestor plant, indicated by the plant comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1-10. Plants, progeny, and seeds may be homozygous or heterozygous for the maize MON 87419 event. Progeny plants may be grown from seeds produced by a maize plant containing the maize MON 87419 event or from seeds produced by a maize plant fertilized with pollen containing the maize MON 87419 event.

[0053] As used herein, a “plant part” of the invention is any part derived from a transgenic maize plant containing the maize MON 87419 event. Plant parts include but are not limited to pollen, ovule, pod, flower, roots, stems, fibers, and leaves. Plant parts may be viable or nonviable.

[0054] The invention provides a commodity product that is produced from transgenic maize containing the maize MON 87419 event. Commodity products of the invention contain a detectable amount of DNA comprising a DNA sequence selected from the group consisting of SEQ ID NO:1-10. As used herein, a “commodity product” refers to any composition or product which is comprised of material derived from a transgenic maize plant, maize seed, maize plant cell, or maize plant part containing the maize MON 87419 event. Commodity products include but are not limited to processed seeds, grains, plant parts, and meal. Transgenic maize containing the maize MON 87419 event can be used to manufacture any commodity product typically acquired from maize. A commodity product of the invention will contain a detectable amount of DNA corresponding to the maize MON 87419 event. Detection of one or more of this DNA in a sample may be used for determining the content or the source of the commodity product. Any standard method of detection for DNA molecules may be used, including methods of detection disclosed herein.

[0055] The invention provides methods for controlling weeds using glufosinate or dicamba, or glufosinate and dicamba herbicides with transgenic maize containing the maize MON 87419 event. A method for controlling weeds in an area, such as a field, is provided that consists of planting transgenic maize plants containing the maize MON 87419 event in an area and applying an herbicidally effective dose of glufosinate or dicamba, or glufosinate and dicamba herbicides to the area for the purpose of controlling weeds in the area without injuring

the transgenic maize plants containing the maize MON 87419 event. Such application of glufosinate or dicamba, or glufosinate and dicamba herbicides may be pre-emergence (any time after transgenic maize seed containing the maize MON 87419 event is planted and before transgenic maize plants containing the maize MON 87419 event emerge) or post-emergence (any time after transgenic maize plants containing the maize MON 87419 event emerge). A herbicidally effective dose of glufosinate for use in the area for controlling weeds should consist of a range from about 0.1 pound acid equivalent per acre (ae/ac) to as much as about 16 pounds ae/ac of glufosinate over a growing season. A herbicidally effective dose of dicamba for use in the area for controlling weeds should consist of a range from about 0.1 pound ae/ac to as much as about 16 pounds ae/ac of dicamba over a growing season. Multiple applications of glufosinate or dicamba, or glufosinate and dicamba herbicides may be used over a growing season, for example, two applications (such as a pre-planting application and a post-emergence application or a pre-emergence application and a post-emergence application) or three applications (such as a pre-planting application, a pre-emergence application, and a post-emergence application).

[0056] As used herein, the “active ingredient” or “ai” is the component of an herbicide formulation responsible for the herbicidal activity, often measured in pounds per gallon or applied as pounds per acre. For herbicides that are acids (for example, molecules that have a carboxyl group as part of their structure), the acidic group is often converted to (may be replaced by the desired ions to form) a salt or (reacted with an alcohol to form) an ester during the formulation process. This may alter not only the chemical characteristics of a particular herbicide molecule, but also the mass. However, the corresponding acid is the herbicidally active portion of the formulation and equivalency of herbicidal activity between different active ingredients can be calculated using the acid equivalent as the standard unit of measurement. The term “acid equivalent” or “ae” means the portion of an active ingredient in a formulation that theoretically could be converted back to the corresponding acid. Herbicide application rates may be expressed as “acid equivalent per acre” (abbreviated as “ae/ac”) or as “active ingredient per acre” (abbreviated as “ai/ac”).

[0057] Methods for producing an herbicide tolerant transgenic maize plant containing the maize MON 87419 event are provided. Progeny produced by these methods may be varietal or hybrid plants; may be grown from seeds produced by a transgenic maize plant containing the maize MON 87419 event or from seeds produced by a maize plant fertilized with pollen from a

transgenic maize plant containing the maize MON 87419 event; and may be homozygous or heterozygous for the maize MON 87419 event. Plants may be self-pollinated (also known as “selfing”) or cross-pollinated (also known as “crossing”). Transgenic maize plants containing the maize MON 87419 event may be self-pollinated to generate a true breeding line of plants that are homozygous for the maize MON 87419 event. Selfing results in progeny known as “inbred” and is used to produce inbred lines that are genetically uniform. Alternatively, transgenic maize plants containing the maize MON 87419 event may be outcrossed (bred with another plant that is transgenic or nontransgenic) to produce a varietal or a hybrid seed. Seed and progeny plants made by the methods of the invention would contain the maize MON 87419 event and may then be treated with glufosinate or dicamba, or glufosinate and dicamba herbicides. Treatment with glufosinate or dicamba, or glufosinate and dicamba herbicides may be used to select progeny that are tolerant. Alternatively, these progeny plants may be analyzed using diagnostic methods to select for plants or seeds containing the maize MON 87419 event.

[0058] Plants, progeny, seeds, plant cells, and plant parts of the invention may also contain one or more additional maize transgenic traits, particularly those introduced by crossing a maize plant containing the maize MON 87419 event with another maize plant containing the additional transgenic trait(s). Such maize transgenic traits include, but are not limited to, increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, hybrid seed production, and herbicide tolerance, in which the trait is measured with respect to a maize plant lacking such transgenic trait. Such maize transgenic traits are known to one of skill in the art; for example, a list of such traits is provided the United States Department of Agriculture’s (USDA) Animal and Plant Health Inspection Service (APHIS) and can be found on their website at <http://www.aphis.usda.gov>. Two transgenic plants may thus be crossed to produce progeny that contain two or more independently segregating transgenic traits. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of breeding methods that are commonly used for different traits and crops can be found in one of several references, for example, Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison WI (1987).

[0059] The plants, seeds, cells, plant parts, and commodity products of the invention may be used for detection of DNA and protein molecules indicative of the presence of the maize

MON 87419 event. Such detection would be done using the DNA sequences provided herein and the respective DMO and PAT protein sequences encoded by the transgene insert that is provided as SEQ ID NO:9. Detection of the presence of the maize MON 87419 event may be done by using methods known in the art, such as thermal amplification of nucleic acid, nucleic acid hybridization techniques (such as northern blotting and southern analysis), protein detection techniques (such as western blotting, immuno-precipitation, and enzyme-linked immunosorbent assay-based (ELISA) techniques) or by using the methods of detection and/or the detection kits provided herein. One method provides for contacting a DNA sample with a primer pair that is capable of producing an amplicon from DNA of the transgenic maize containing the maize MON 87419 event, performing an amplification reaction and thereby producing a DNA amplicon comprising at least one of the DNA sequences provided as SEQ ID NO:1-10 and SEQ ID NO:14-22, and then detecting the presence or absence of the amplicon molecule and optionally confirming within the sequence of the amplicon a sequence comprising at least one of the sequences provided as SEQ ID NO:1-10 and SEQ ID NO:14-22. The presence of such an amplicon is diagnostic for the presence of DNA specific for the transgenic maize containing the maize MON 87419 event and thus biological material in the sample that is derived from transgenic maize containing the maize MON 87419 event. Another method provides for contacting a DNA sample with a DNA probe, subjecting the probe and the DNA sample to stringent hybridization conditions, and then detecting hybridization between the probe and the target DNA sample. Detection of hybridization is diagnostic for the presence of DNA specific for the transgenic maize containing the maize MON 87419 event in the DNA sample.

[0060] DNA detection kits for the maize MON 87419 event are provided. Variations on such kits can also be developed using the compositions and methods disclosed herein and the methods well known in the art of DNA detection. DNA detection kits can be applied to methods for breeding with transgenic maize plants containing the maize MON 87419 event. Such kits contain DNA primers or probes comprising fragments of SEQ ID NO:1-10. One example of such a kit comprises at least one DNA molecule of sufficient length of contiguous nucleotides of SEQ ID NO:10 to function as a DNA probe useful for detecting the presence and/or absence in a sample of DNA derived from transgenic herbicide tolerant maize plants containing the maize MON 87419 event. A DNA molecule sufficient for use as a DNA probe is provided that is useful for determining, detecting, or diagnosing the presence and/or absence in a sample of transgenic

herbicide tolerant maize containing the maize MON 87419 event DNA is provided as SEQ ID NO:13. Other probes may be readily designed by one of skill in the art and should comprise at least about fifteen nucleotides of SEQ ID NO:10 and be sufficiently unique to transgenic herbicide tolerant maize containing the maize MON 87419 event DNA in order to identify DNA derived from the maize MON 87419 event. Another type of kit comprises a primer pair useful for producing an amplicon useful for detecting the presence or absence in a sample of the maize MON 87419 event. Such a kit would employ a method comprising contacting a target DNA sample with a primer pair, then performing a nucleic acid amplification reaction sufficient to produce an amplicon comprising a DNA molecule having at least one sequence selected from SEQ ID NO:1-10, and then detecting the presence or absence of the amplicon. Such a method may also include sequencing the amplicon or a fragment thereof. Other primer pairs may be readily designed by one of skill in the art and should comprise at least twenty nucleotides of SEQ ID NO:10 and be sufficiently unique to the transgenic herbicide tolerant maize containing the maize MON 87419 event DNA in order to detect the maize MON 87419 event. Kits of the invention may optionally also comprise reagents for performing the detection or diagnostic reactions described herein or instructions for the use of the kit and its contents.

[0061] As used herein, the term “comprising” means “including but not limited to”.

Deposit Information

[0062] A deposit of a representative sample of transgenic maize seed comprising the maize MON 87419 event has been made according to the Budapest Treaty with the American Type Culture Collection (ATCC) having an address at 10801 University Boulevard, Manassas, Virginia USA, Zip Code 20110. The ATCC Patent Deposit Designation (accession number) for seeds comprising the maize MON 87419 event is PTA-120860 and the date of deposit was January 17, 2014. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer.

EXAMPLES

[0063] The following examples are included to more fully describe the invention. It should be appreciated by those of skill in the art that many modifications can be made in the specific examples which are disclosed and still obtain a similar result. Certain agents which are

both chemically and physiologically related may be substituted for the agents described herein while achieving the same or similar results. All such substitutions and modifications apparent to those skilled in the art are deemed to be within the scope of the invention.

Example 1: MON 87419 event production and selection

[0064] This example describes the production, analysis, and selection of transgenic maize containing the event MON 87419, an event which can provide tolerance to both dicamba and glufosinate herbicides. Summarized are the production and analysis of tens of thousands of individual plants over multiple years through the rigorous molecular, phenotypic, and field testing required for the ultimate selection of the maize MON 87419 event.

[0065] Transformation vectors containing a variety of different expression cassettes were designed and tested to confirm their utility for expressing the *dmo* and *pat* genes. Using this data, expression element combinations were selected and eight different transformation vectors were constructed and transformed into maize. These vectors tested promoters and terminators in various combinations with the two coding regions for the *pat* gene and the *dmo* gene (Table 1). The resulting plants were analyzed for protein expression and two vectors (shown as A and B in Table 1) were selected for commercial maize transformation.

Table 1. Cassette Configuration of Transformation Vectors.

Vector	cassette 1 (PAT)			cassette 2 (DMO)		
	Promoter	Gene of Interest	Terminator	Promoter	Gene of Interest	Terminator
A	AND.ge.Ubq1	PAT	Os.Ara5	PCSV/I-Act1	CTP4/DMO	Hsp17.5
B	P-1	PAT	T-1	PCSV/I-Act1	CTP4/DMO	Hsp17.5
C	AND.ge.Ubq1	PAT	Os.Ara5	P-1	CTP4/DMO	Hsp17.5
D	P-1	PAT	T-1	AND.ge.Ubq1	CTP4/DMO	Hsp17.5
E	AND.ge.Ubq1	PAT	Os.Ara5	P-2	CTP4/DMO	Hsp17.5
F	AND.ge.Ubq1	PAT	Os.Ara5	P-3	CTP4/DMO	Hsp17.5
G	P-1	PAT	T-1	P-2	CTP4/DMO	Hsp17.5
H	P-1	PAT	T-1	P-3	CTP4/DMO	Hsp17.5

[0066] Over thirteen thousand unique transformed maize plants were produced using the two vectors (A and B) that were selected. In plants there is often wide variation in the levels of expression of an introduced gene among individual events, and gene expression can directly

correlate positively or negatively with the phenotype of the plant containing the event. The expression of foreign genes in plants is known to be influenced, among other things, by their chromosomal position. For this reason, it was necessary to screen a large number of individual plants containing random insertion events through multiple years and locations to identify the optimal event. The transgenic maize MON 87419 event was created through *Agrobacterium*-mediated transformation of LH244 maize immature embryos. Methods for transforming maize are known in the art. Maize cells were transformed and regenerated into intact maize plants. Rooted plants with normal phenotypic characteristics were selected. Thousands of individual, independent events were then transferred to soil for growth and further assessment.

Table 2. Event Selection Process

	Milestone	Vector A Events Advanced	Vector B Events Advanced	Total Unique Events Advanced
R0 Evaluation	Transgenic Events Produced	5236	8413	13649
	First Pass Single Copy analysis	1300	1698	2998
	R0 Spray	642	798	1440
	Initial Detailed Molecular Analysis	85	99	184
	R0 Southern	54	58	112
Early Screens (R1 & R2)	R1 Trials & molecular analysis	22	22	44
	R2 Trials & molecular analysis	20	22	42
Advanced Field Trials & Molecular Analysis	SA Year 1	7	10	17
	US Year 1	5	6	11
	SA Year 2	2	3	5
	In-depth evaluation	2	0	2
	Final event selection	1	0	1

[0067] Throughout the event selection process, molecular analysis as well as field trials to assess phenotype, agronomics, and efficacy of the events were conducted, often concurrently (Table 2). The 13,000 individual, unique transformed maize R0 plants were analyzed first by

PCR to select for events with a single copy of the transgene insert (First Pass Single Copy). This resulted in the advancement of 2,998 events. The next selection was R0 herbicide spray tolerance conducted in a green house. The plants were tested for tolerance to both glufosinate (Ignite® 280 herbicide) and dicamba (Clarity® herbicide) by using a tank mix of glufosinate (0.9 lb ai/ac) and dicamba (2.0 lb ae/ac) sprayed at the V1/V2 growth stage. Plants that showed >15% injury were discarded, and 1,440 events were selected for further analysis. An initial detailed molecular analysis was conducted and included sequence identification and confirmation, and a second check of copy number and absence of backbone. This analysis resulted in 184 events containing only a single copy of the transgene insert selected for advancement. Southern analysis on DNA extracted from R0 events was done to further confirm transgene insert copy number and to confirm absence of transformation backbone. Based on this, 112 events were selected for advancement to R1 for further analysis. R0 plants were self-pollinated and seed was collected for R1 trials.

[0068] Concurrently with all field trials, additional molecular analysis was in progress. Northern analysis was done to detect and measure mRNA transcripts of the *pat* and *dmo* genes. N-terminal protein sequencing of the PAT and DMO proteins purified from transgenic plants containing select events was done to confirm the recombinant protein sequence. Western analysis to detect the PAT and DMO proteins was done with transgenic plant samples. Sequencing of the entire transgene and both the 5' and 3' ends of the insert was conducted and subsequently used to develop methods of detecting individual events. In depth Southern analysis was performed on R1 plants to confirm copy number and the absence of backbone.

[0069] For the R1 field trial early screens, of the 112 events selected from the R0 screening 82 events were selected based on seed return and nursery size considerations. The R1 plants were segregating, and thus were null, hemizygous, or homozygous for an event. The 82 events in R1 plants were evaluated in a field efficacy screen with high rates of herbicide application. Plant injury was assessed following treatments with various combinations of glufosinate (Ignite 280) and dicamba (Clarity) (up to 20X glufosinate and up to 16X dicamba labeled field application rates) and with application timing at various growth stages from V1/V2 to V8/V10. Injury ratings were taken 10 to 14 days after herbicide application. Standard injury ratings include scoring for percent chlorosis, malformation, and breeding. Overall averages for multiple plants containing the same event were used to select events for advancement. Herbicide

applications at 2X rates generally produced less than 10% injury, and herbicide application at the 16X and 20X rates produced injury ratings of more than 10%. In addition to the efficacy testing, agronomic scoring was collected for each plant and correlated to the event it contained. The agronomic scoring criteria that were evaluated included plant height, ear height, percent moisture, test weight, days to 50% pollen, and days to 50% silk. Based on analysis of the data collected from the R1 field trials and the molecular analysis, 44 events were selected for advancement to R2 for further analysis. R1 plants were self-pollinated and seed was collected for R2 trials.

[0070] In the R2 field trial early screens, R2 and F1 events (from an R1 outcross) were evaluated in field efficacy screens at three locations (two states and Puerto Rico). Plant injury was assessed following treatments with various combinations of glufosinate and dicamba application rates and application timing. The R2 plants are homozygous for the event, and the herbicide tolerance failure rate after herbicide application was low, confirming the R1 results. Agronomic data was collected and scored as in the R1 field trials. Additional molecular analysis, including gene expression analysis, was also used to select plants containing the best events. Based on analysis of the data collected from the R2 and F1 field trials and the molecular analysis, 42 events were selected for advancement to R3 for further analysis. R2 plants were self-pollinated and seed was collected for R3 trials.

[0071] For the advanced field trials, both hybrid and inbred efficacy and hybrid and inbred agronomic field trials were conducted. The agronomic field trials were run during the same season as the efficacy field trials. All field trials used a randomized complete block design and were conducted at multiple locations. For both efficacy and agronomic field trials, agronomic scoring was collected throughout the field trial season, and at the end of the season yield was determined (efficacy yield or agronomic yield). Efficacy field trials were conducted to assess crop injury 10 to 14 days following herbicide application, crop injury ratings, and yield. The target crop injury rating was a score of less than 10% for advancement of the event. For agronomic field trials, the plots were maintained weed free and no glufosinate or dicamba herbicide was applied during the growing season. The hybrid agronomic field trials included controls of a comparable hybrid (hybrid control) produced using the same parental maize lines used to make the transgenic hybrid cross, but not containing a transgenic event. Inbred controls were a comparable inbred to the transgenic inbred lines.

[0072] A meta-analysis was performed using the aggregate of the multi-season, multi-location field trial data. Table 3 illustrates the number of replications for which an observation was repeated for the particular field trial type for plants containing either one of the events. For each of the two events, there were 135 data points recorded for hybrid agronomic performance, 933 data points for hybrid efficacy, 179 data points for inbred agronomics, 30 data points for inbred efficacy, and 16 data points for event based pressure testing of herbicide application rates for maize containing the event.

Table 3. Field trial replications for two final events.

Description	Reps per event
Hybrid Agronomics	135 reps
Hybrid Efficacy	933 reps
Inbred Agronomics	179 reps
Inbred Efficacy	30 reps
Event based Pressure Test	16 reps
Total Replications per event	1293

[0073] Hybrid plants each containing one of 23 selected events (a subset of the 42 events from the R2 field trial) were evaluated in South America (SA) Year 1 contra-season efficacy field trials and agronomic field trials. Trials were conducted at six locations in a randomized complete block design with 4 treatments and 2 replications per treatment. In the efficacy field trials, the dicamba formulation was Banvel® 4SL herbicide and the glufosinate formulation was Liberty® 1.67SL. Herbicide treatments consisted of the following: (1) non-treated control; (2) dicamba at 2 lbs ae/acre (ac) PRE (where PRE is defined as at planting or before crop emergence) followed by dicamba at 1lb ae/ac applied at each of VE-V2 followed by V4 followed by V8; (3) glufosinate applied at 0.8 lb ai/ac at VE-V2 followed by V4 followed by V8; and (4) glufosinate applied at 0.8 lb ai/ac plus dicamba applied at 1 lb ae/ac at VE-V2 followed by V4 followed by V8. Injury ratings were taken at 10 to 14 days after herbicide application. Overall averages for multiple plants containing the same event were used to select events for advancement. The target crop injury rating was a score of less than 10%, and the observed injury rating was below 1%. Based upon hybrid injury scoring, agronomic scoring, efficacy yield, agronomic yield, and additional molecular analysis, 17 events were selected for advancement.

[0074] Inbred and hybrid plants from the 17 events advanced from the SA Year 1 contra-season field trials were then further evaluated in United States (US) Year 1 efficacy field trials and agronomic field trials. These trials were conducted in 2012, which was a season of severe drought in the United States. The hybrid efficacy field trials were conducted at 12 locations, 2 states in a randomized complete block design with 6 treatments and 3 replications per treatment. The hybrid plants containing the transgenic event derived glyphosate tolerance from the male parent in the cross. In these efficacy field trials, the glyphosate formulation was Roundup PowerMAX® 4.5SL, the dicamba formulation was Clarity 4SL, and the glufosinate formulation was Ignite 280 2.34SL. Herbicide treatments consisted of the following: (1) non-treated control; (2) glyphosate at 3 lbs ae/ac applied at V4 followed by V8; (3) glufosinate at 0.8 lb ai/ac applied at V2 followed by V4 followed by V8; (4) dicamba at 2 lbs ae/ac applied PRE and then again applied at V4 followed by V8; (5) glyphosate at 3lbs ae/ac plus dicamba at 1.5 lbs ae/ac applied at V2 followed by V4 followed by V8; (6) dicamba at 2 lbs ae/ac applied at V2 followed by glufosinate at 0.8 lbs ai/ac plus dicamba at 1 lb ae/ac applied at growth stage V4 followed by glyphosate at 3 lbs ae/ac plus dicamba at 1.5 lbs ae/ac applied at growth stage V8. Injury ratings were taken 10 to 14 days after herbicide application. Overall averages for multiple plants containing the same event were used to select events for advancement. The target crop injury rating was a score of less than 10%, and the observed injury rating was below 1%. Based on hybrid injury scoring, agronomic scoring, efficacy yield, agronomic yield, and additional molecular analysis, 11 events were selected for advancement.

[0075] South America (SA) Year 2 contra-season field trials to assess hybrid efficacy and inbred agronomic yield were then conducted with plants containing these 11 events. The hybrid plants containing the transgenic event derived glyphosate tolerance from the male parent in the cross. The hybrid efficacy field trials were conducted essentially as described for South America (SA) year 1 contra-season field trials but with the following herbicide treatments: (1) non-treated control; (2) glyphosate at 3 lbs ae/ac applied at V4 followed by V8; (3) glufosinate at 0.8 lb ai/ac applied at V4 followed by V8; (4) dicamba at 2 lbs ae/ac applied PRE and then again applied at V4 followed by V8; (5) glyphosate at 3lbs ae/ac plus dicamba at 1.5 lbs ae/ac applied at V4 followed by V8; (6) glufosinate at 0.8 lbs ai/ac plus dicamba at 1 lb ae/ac applied at V4 followed by glyphosate at 3 lbs ae/ac plus dicamba at 1.5 lbs ae/ac applied at V8. Injury ratings were taken 10 to 14 days after herbicide application. Overall averages for multiple plants containing the

same event were used to select events for advancement. Based on hybrid injury scoring, agronomic scoring, hybrid efficacy yield, inbred agronomic yield, and additional molecular analysis, 5 events were selected for advancement.

[0076] Additional molecular analysis was then completed for these 5 events. The multi-year field and molecular data for each of the 5 events, including hybrid trait efficacy field trials, hybrid and inbred yield measurements, agronomic scoring, and molecular information was then reviewed, and two events were selected for further analysis. Both of these events were produced using the same transformation vector, and therefore had the same transgene insert but not the same genomic location or flanking sequence.

[0077] United States (US) Year 2 hybrid and inbred efficacy field trials, and hybrid and inbred agronomic field trials were conducted to evaluate these two events. The hybrid efficacy trials were done similar to year 1 US field trials, but included different spray regimens. Efficacy was measured by injury ratings and hybrid efficacy yield. Additional molecular analysis for the events was also done. The hybrid plants containing the transgenic event derived glyphosate tolerance from the male parent in the cross. The hybrid Efficacy 1 and 2 field trials were conducted at twelve locations across two states and the hybrid Efficacy 3 field trials were conducted at thirty-three locations across four states. In these field trials, the glyphosate formulation was Roundup PowerMAX 4.5SL, the dicamba formulation was Clarity 4SL, and the glufosinate formulation was Ignite 280 2.34SL. Herbicide applications for the Efficacy 1 and Efficacy 2 trials (with crosses to two separate inbred lines) were: (1) non-treated control; (2) glufosinate at 0.4 lb ai/ac applied at VE-V2 followed by V6; (3) glufosinate at 0.8 lb ai/ac applied at VE-V2 followed by V6; (4) dicamba at 0.5 lbs ae/ac applied at V4 followed by V8; (5) dicamba at 1.0 lbs ae/ac applied at V4 followed by V8; and (6) glyphosate at 2.25lbs ae/ac plus dicamba at 1.0 lbs ae/ac applied at V4 followed by V8. Herbicide applications for the Efficacy 3 trial (representing hybrid from a cross with a third inbred line) were: (1) non-treated control; (2) dicamba at 0.5 lbs ae/ac applied at VE-V2 followed by glufosinate at 0.4 lb ai/ac applied at V6; and (3) dicamba at 1.0 lbs ae/ac applied at VE-V2 followed by glufosinate at 0.8 lb ai/ac applied at V6. Injury ratings were taken 10 to 14 days after herbicide application and for multiple plants containing the same event were used to select events for advancements. Yield and agronomic data was collected.

[0078] To compare the hybrid injury ratings for the two events, meta-analysis of the multiple hybrid efficacy field trials was completed. (Table 4) Injury rating was scored at V8 (where V8 analysis encompasses the cumulative injury from V2, V4, V6, and V8 herbicide applications) with a statistical least significant difference (LSD at $p<0.05$). Tester 1, Tester 2, and Tester 3 represent crosses with 3 independent inbred maize parent lines, which were used to generate the hybrid for the indicated field trial. For each of the trials, no statistical difference in injury rating was found between hybrids generated using a transgenic maize parent containing either of the two events.

Table 4. Meta-analysis of injury rating from hybrid efficacy field trials.

Field Trial	Transgenic Maize	V8 injury	LSD (P<0.05)
Year 1 combined SA and US	MON 87419	0.43	0.3
Year 1 combined SA and US	EVENT 2	0.49	0.3
SA Year 2 Hybrid Efficacy	MON 87419	2.58	3
SA Year 2 Hybrid Efficacy	EVENT 2	1.98	3
US Year 2 Hybrid Tester 1	MON 87419	0	0
US Year 2 Hybrid Tester 1	EVENT 2	0	0
US Year 2 Hybrid Tester 2	MON 87419	0	0
US Year 2 Hybrid Tester 2	EVENT 2	0	0
US Year 2 Hybrid Tester 3	MON 87419	0.31	0.42
US Year 2 Hybrid Tester 3	EVENT 2	0.12	0.42

[0079] A meta-analysis of the hybrid efficacy yield (bushels/acre) from the multiple efficacy field trials was completed comparing yield from hybrids containing each of the two events. (Table 5) For each of the trials, no statistical difference in hybrid efficacy yield was found between hybrids generated using a transgenic maize parent containing either of the two events.

Table 5. Meta-analysis of yield from hybrid efficacy field trials.

Field Trial	Hybrid Maize	Yield (Bu/ac)	LSD (p<0.05)
Year 1 combined SA and US	MON 87419	171.07	4
Year 1 combined SA and US	EVENT 2	172.90	4

SA Year 2	MON 87419	233.66	14
SA Year 2	EVENT 2	231.59	14
US Year 2 Tester 1	MON 87419	217.43	10
US Year 2 Tester 1	EVENT 2	220.06	10
US Year 2 Tester 2	MON 87419	219.68	7
US Year 2 Tester 2	EVENT 2	221.21	7
US Year 2 Tester 3	MON 87419	208.76	5.67
US Year 2 Tester 3	EVENT 2	213.02	5.67

[0080] Pressure testing field trials were also conducted with hybrid transgenic maize containing either one of the two events. In the pressure tests, either glufosinate (Ignite 280, 2.34SL) or dicamba (Clarity 4SL) herbicide was applied at non-commercially high rates. For typical field trials, the 1X rate for glufosinate was 0.4 lb ai/ac and the 1X rate for dicamba was 0.5 lb ae/ac. For the glufosinate pressure testing field trials, glufosinate was applied at VE-V2 followed by V4 followed by V8 at the following rates: (1) 1 lb ai/ac (2.5X); (2) 2 lb ai/ac (5X); (3) 4 lb ai/ac (10X); and (4) 8 lb ai/ac (20X). For the dicamba pressure testing field trials, dicamba was applied at VE-V2 followed by V4 followed by V8 at the following rates: (1) 2 lb ae/ac (4X); (2) 4 lb ae/ac (8X); (3) 8 lb ae/ac (16X); and (4) 16 lb ae/ac (32X). At the end of the season, the hybrid pressure testing field trials were harvested and yield (bushels/acre or bu/ac) was determined. An analysis of the yield data compared hybrids containing either of the two events. For each of the trials, no statistical difference in yield at any of the herbicide application rates was found between hybrids generated using a transgenic maize parent containing either of the two events.

Table 6. Yield from hybrid pressure testing efficacy field trials.

Field Trial	Transgenic Hybrid Maize	Yield (Bu/ac)	LSD (p<0.05)
Glufosinate Pressure Test 2.5-20X	MON 87419	207.31	40
Glufosinate Pressure Test 2.5-20X	EVENT 2	201.65	40
Dicamba Pressure Test 4-32X	MON 87419	239.93	40
Dicamba Pressure Test 4-32X	EVENT 2	234.60	40

[0081] Hybrid agronomic field trials were conducted in 3 sets with 15 locations per set at 21 locations across 3 states and were run during the same season with the hybrid efficacy field trials. Agronomic measures were collected through out the field trial season, and at the end of the

season agronomic yield was determined. Meta-analysis across the multi-season, multi-location hybrid agronomic field trials was used to compare the yield of the hybrid control and the hybrids containing the either one of the two events. No statistical difference in hybrid agronomic yield was found either between the transgenic hybrids or as compared to the hybrid controls (Table 7).

Table 7. Meta-analysis of yield from hybrid agronomic field trials.

Field Trial	Hybrid Maize	Yield (Bu/ac)	LSD (p<0.05)
Year 1 combined SA and US	Control	180.28	6
Year 1 combined SA and US	MON 87419	179.01	6
Year 1 combined SA and US	EVENT 2	184.88	6
SA Year 2	Control	231.70	10
SA Year 2	MON 87419	232.47	10
SA Year 2	EVENT 2	225.07	10
US Year 2 Tester 1	Control	185.80	17.40
US Year 2 Tester 1	MON 87419	187.09	17.40
US Year 2 Tester 1	EVENT 2	192.28	17.40
US Year 2 Tester 2	Control	219.12	12.83
US Year 2 Tester 2	MON 87419	219.56	12.83
US Year 2 Tester 2	EVENT 2	221.22	12.83
US Year 2 Tester 3	Control	197.62	8.66
US Year 2 Tester 3	MON 87419	191.49	8.66
US Year 2 Tester 3	EVENT 2	195.48	8.66

[0082] Inbred efficacy field trials were conducted using a randomized complete block design at 6 locations, 1 state. In these field trials, the dicamba formulation was Clarity 4SL, and the glufosinate formulation was Ignite 280 2.34SL. The herbicide application for the inbred efficacy field trials were glufosinate at 0.8 lb ai/ac and dicamba at 2.0 lbs ae/ac applied at VE-V2 followed by glufosinate at 0.8 lb ai/ac and dicamba at 2.0 lbs ae/ac applied at V8. At the end of the season, yield was measured. For each of the trials, a statistical difference in inbred efficacy yield was found when comparing inbred yield harvested from these trials from transgenic maize containing either of the two events (Table 8). These data indicate the superior performance of transgenic maize containing the maize MON 87419 event.

Table 8. Meta-analysis of yield from inbred efficacy field trials.

Field Trial	Inbred Maize	Yield (Bu/ac)	LSD (p<0.05)
year 2 US Inbred	MON 87419	110.95	7
year 2 US Inbred	EVENT 2	98.89	7

[0083] Inbred agronomic field trials were run during the same season with the US Year 1 efficacy field trials, SA Year 1 efficacy field trials, and US Year 2 efficacy field trials (11 locations, 1 state). The plots were set up in randomized complete block design conducted at multiple locations, and the trials included controls of a comparable inbred to the transgenic inbred lines. Meta-analysis across the multi-season, multi-location inbred agronomic field trials was conducted comparing yield for the paired control and the transgenic inbreds generated using either of the two events. No statistical difference in inbred agronomic yield was found between the control and transgenic maize containing the MON 87419 event (Table 9). In contrast, there was a statistically significant decrease in yield in transgenic maize containing the event 2 when compared to either control or transgenic maize containing the MON 87419 event. These data further indicated the superior performance of transgenic maize containing the maize MON 87419 event.

Table 9. Meta-analysis of yield from inbred agronomic field trials.

Field Trial	Inbred Maize	Total (Bu/ac)	LSD (p<0.05)
year 1 US Inbred	Control	65.87	31
year 1 US Inbred	MON 87419	60.33	31
year 1 US Inbred	EVENT 2	43.10	31
year 2 SA Inbred	Control	95.88	7
year 2 SA Inbred	MON 87419	98.77	7
year 2 SA Inbred	EVENT 2	75.85	7
year 2 US Inbred	Control	108.90	6
year 2 US Inbred	MON 87419	109.22	6
year 2 US Inbred	EVENT 2	96.88	6

Example 2: Characterization of the DNA sequence of the maize MON 87419 event

[0084] This example describes the extensive molecular characterization that was conducted on the maize MON 87419 event. The transgene insert of the maize MON 87419 event

contains, from the 5' to 3' orientation: (i) the promoter (P-ANDge.Ubq1), leader, and intron (L-I-ANDge.Ubq1) of the ubiquitin gene (Ubq) from *Andropogon gerardii*; operably linked to the *pat* gene from *Streptomyces viridochromogenes* (CR-STRvi.pat) that encodes a phosphinothricin N-acetyltransferase (PAT) that confers tolerance to glufosinate herbicide; operably linked to the polyadenylation signal (also known as a terminator that directs polyadenylation of mRNA) from the RA5B precursor gene of *Oryza sativa* (T-Os.Ara5) and (ii) the promoter for the full length transcript of Peanut Chlorotic Streak Virus with a duplicated enhancer region (PCISV); operably linked to the leader of the light harvesting complex b1 gene from *Triticum aestivum* (L-Ta.Lhcb1); operably linked to the first intron from the actin 1 gene from *Oryza sativa* (I-Os.Act1); operably linked to the N-terminal chloroplast transit peptide from the Petunia x hybrida 5-enolpyruvylshikimate-3-phosphate synthase gene (TS-Ph.ShkG-CTP4); operably linked to the *dmo* gene *Stenotrophomonas maltophilia* optimized for monocot expression(CR-STEma.DMO) that encodes a dicamba monooxygenase (DMO) that confers tolerance to dicamba herbicide; operably linked to the heat shock protein 17 polyadenylation signal from *Triticum aestivum* (T-Ta.Hsp17). The 5' end of the transgene insert was flanked by the Right border of *Agrobacterium tumifaciens* and the 3' end of the transgene insert was flanked by the Left border of *Agrobacterium tumifaciens*.

[0085] Southern blot analysis was conducted to confirm that transgenic maize containing the maize MON 87419 event contained a single, intact copy of the entire transgene insert without any vector backbone. Flank sequences were isolated from both the 5' and 3' ends of the insert, and the respective junctions were determined using inverse PCR and/or genome walking techniques. The chromosomal location of the insert of the maize MON 87419 event was determined using inverse PCR to amplify genomic DNA outside of the site of interest. The flank sequences for the maize MON 87419 event were mapped to the known maize genome physical assembly and the maize MON 87419 event was confirmed to not be within any known genes. This sequence information was used to design event specific endpoint TAQMAN® assays to identify the presence of the maize MON 87419 event in a sample. The insertion site sequence information was also used for bioinformatics analysis of the chromosomal location of the event. Insertion site integrity was determined by PCR across the wild-type allele using primers specific to the flanking regions of the maize MON 87419 event. The wild-type insertion site was used to map to the maize reference genome the unique site of transgene integration for the maize

MON 87419 event. To ensure that no alterations or mutations were introduced to any region of the transgene during transformation, the entire transgene insert of the maize MON 87419 even was isolated from the plant and sequenced.

[0086] N-terminal protein sequencing of the expressed PAT and DMO proteins was performed using immunopurified protein extracts from transgenic maize grain containing the maize MON 87419 event. This sequence was then used to confirm the authentic N-terminal amino acid sequence. Western analysis was conducted on protein extracts from grain of transgenic maize containing the maize MON 87419 event. This confirmed that a single expected-sized protein was being produced for PAT and for DMO, respectively. ELISAs were developed to determine protein levels in various transgenic maize tissue types (leaf, seed, roots, and pollen) for the PAT or DMO protein expressed from the maize MON 87419 event. Northern analysis was conducted on poly-A RNA isolated from grain of transgenic maize containing the maize MON 87419 event. This confirmed the transcript size and number for the *pat* and *dmo* mRNA products. RNA expression levels were also measured by real-time PCR using samples from transgenic maize containing the maize MON 87419 event.

Example 3: Event specific endpoint TAQMAN® assays

[0087] This example describes an event specific endpoint TAQMAN® thermal amplification method developed to identify transgenic maize containing the maize MON 87419 event in a sample. The DNA primers used in the endpoint assay are primers SQ26644 (SEQ ID NO:11), SQ26645 (SEQ ID NO:12), and 6-FAM™ labeled probe PB11207 (SEQ ID NO:13). 6-FAM™ is a fluorescent dye product of Applied Biosystems (Foster City, CA) attached to the DNA probe. For TAQMAN® MGB™ probes, the 5' exonuclease activity of Taq DNA polymerase cleaves the probe from the 5'-end, between the fluorophore and quencher. When hybridized to the target DNA strand, quencher and fluorophore are separated enough to produce a fluorescent signal, thus releasing fluorescence. SQ26644 and SQ26645 when used with these reaction methods and PB11207 produce a DNA amplicon that is diagnostic for the maize MON 87419 event. The controls for this analysis should include a positive control containing the maize MON 87419 event, a negative control from non-transgenic maize, and a negative control that contains no template DNA. Additionally, a control for the PCR reaction should optimally include Internal Control Primers and an Internal Control Probe, specific to a single copy gene in

the maize genome. These assays are optimized for use with either an Applied Biosystems GeneAmp® PCR System 9700 (run at maximum speed) or MJ Research DNA Engine PTC-225 thermal cycler, but other equipment may be used.

[0088] An example of conditions useful with Endpoint TAQMAN® assay method useful for detection of the maize MON 87419 event is as follows. Step 1: 18 megohm water adjusted for final volume of 10 μ l. Step 2: 5.0 μ l of 2X Universal Master Mix (dNTPs, enzyme, buffer) to a 1X final concentration. Step 3: 0.5 μ l Event Primer-1 (SQ26644) and Event Primer-2 (SQ26645). Mix (resuspended in 18 megohm water to a concentration of 20 uM for each primer) to 1.0 μ M final concentration (for example in a microcentrifuge tube, the following should be added to achieve 500 μ l at a final concentration of 20 uM: 100 μ l of Primer SQ26644 at a concentration of 100 μ M; 100 μ l of Primer SQ26645 at a concentration of 100 μ M; 300 μ l of 18 megohm water). Step 4: 0.2 μ l Event 6-FAM™ MGB Probe PB11207 (10uM) (resuspended in 18 megohm water to a concentration of 10 μ M to 0.2 μ M final concentration. Step 5: 0.5 μ l Internal Control Primer-1 and Internal Control Primer-2 Mix (resuspended in 18 megohm water to a concentration of 20 μ M for each primer) to 1.0 μ M final concentration. Step 6: 0.2 μ l Internal Control VIC™ Probe (10uM) to 0.2 μ M final concentration (resuspended in 18 megohm water to a concentration of 10 μ M). Step 7: 3.0 μ l Extracted DNA (template) for each sample with one each of the following comprising 1. Leaf Samples to be analyzed; 2. Negative control (non-transgenic DNA); 3. Negative water control (no template); 4. Positive control transgenic maize containing the maize MON 87419 event DNA. Step 8: Thermocycler Conditions as follows: One Cycle at 50°C for 2 minutes; One Cycle at 95°C for 10 minutes; Ten Cycles of 95°C for 15 seconds then 64°C for 1 minute with (-1°C/cycle); Thirty Cycles of 95°C for 15 seconds then 54°C 1 minute, optional additional 10 to 20 cycles (95°C for 15 seconds then 64°C for 1 minute (-1°C/cycle) may provide more distinct population separation during EndPoint TaqMan® analysis); final cycle of 10°C.

Example 4: Zygosity assay

[0089] A zygosity assay may be used to determine whether or not a plant comprising the maize MON 87419 event is heterozygous or homozygous for the event or the wild-type allele. An amplification reaction assay can be designed using the sequence information provided herein. For example, such a PCR assay would include design of at least three primers: primer-1, primer-

2, and primer-3, where primer-1 is specific to maize genomic DNA on the 3' flank of the maize MON 87419 event; primer-2 is specific to the maize MON 87419 event transgene insert; and primer-3 is specific to the wild-type allele. When used as a primer pair in an amplification reaction, primer-1 with primer-2 will produce a PCR amplicon specific for the maize MON 87419 event. When used as a primer pair in an amplification reaction, primer-1 with primer-3 will produce a PCR amplicon specific for wild-type allele. In a PCR reaction performed on maize genomic DNA, the respective PCR amplicons generated from (primer-1 + primer-2) and (primer-1 + primer-3) will differ in sequence and size of the amplicon. When the three primers are included in a PCR reaction with DNA extracted from a plant homozygous for the maize MON 87419 event, only the primer-1 + primer-2 amplicon will be generated. When the three primers are included in a PCR reaction with DNA extracted from a plant heterozygous for the maize MON 87419 event, both the primer-1 + primer-2 amplicon and the primer-1 + primer-3 amplicon will be generated. When the three primers are mixed together in a PCR reaction with DNA extracted from a plant that is null for the maize MON 87419 event (that is wild-type), only the primer-1 + primer-3 amplicon will be generated.

[0090] Another method to determine zygosity of a maize plant for the maize MON 87419 event is an endpoint TAQMAN® thermal amplification reaction. For this type of assay, in addition to primers as described above, the assay would include two fluorescently labeled probes. Probe-1 would be specific for the maize MON 87419 event, and probe-2 would be specific for a maize plant that is null for the maize MON 87419 event (wild-type), and where the two probes contain different fluorescent labels, for example the 6-FAM™-label or VIC™-label. When used in an endpoint TAQMAN® thermal amplification reaction, primer-1 + primer-2 + probe-1 will produce a first fluorescent signal specific for the maize MON 87419 event. When used in an endpoint TAQMAN® thermal amplification reaction, primer-1 + primer-3 + probe-2 will produce a second fluorescent signal specific for wild-type maize. When the three primers and two probes are included in an endpoint TAQMAN® thermal amplification reaction with DNA extracted from a plant homozygous for the maize MON 87419 event, only the first fluorescent signal (specific to primer-1 + primer-2 + probe-1) will be generated. When the three primers are included in an endpoint TAQMAN® thermal amplification reaction with DNA extracted from a plant heterozygous for the maize MON 87419 event, both the first fluorescent signal (specific to primer-1 + primer-2 + probe-1) and the second fluorescent signal (specific to

primer-1 + primer-3 + probe-2) will be generated. When the three primers are mixed together in an endpoint TAQMAN® thermal amplification reaction with DNA extracted from a plant which is null for the maize MON 87419 event (wild-type), only the second fluorescent signal (specific to primer-1 + primer-3 + probe-2) will be generated.

[0091] Another method to determine zygosity of a plant for the maize MON 87419 event would be Southern analysis. One of skill in art would understand how to design Southern hybridization probe(s) specific for the maize MON 87419 event and a second southern hybridization probe specific for a maize plant which is null for the maize MON 87419 event (wild-type). With Southern analysis, a signal detected only from the first Southern hybridization probe will be indicative of a plant homozygous for the maize MON 87419 event; a signal detected from both the first Southern hybridization probe and the second Southern hybridization probe will be indicative of a plant heterozygous for the maize MON 87419 event; and a signal detected only from the second Southern hybridization probe will be indicative that the DNA was extracted from a plant that is null for the maize MON 87419 event (wild-type).

What is claimed is:

Claim 1. A recombinant DNA molecule comprising a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10.

Claim 2. The recombinant DNA molecule of claim 1, wherein the DNA molecule is from a transgenic maize plant or seed comprising the maize MON 87419 event, a representative sample of seed comprising said event having been deposited as ATCC Accession No. PTA-120860.

Claim 3. The recombinant DNA molecule of claim 1, wherein the DNA molecule is an amplicon diagnostic for the presence of DNA from the maize MON 87419 event.

Claim 4. The recombinant DNA molecule of claim 1, wherein the DNA molecule is in a maize plant, cell, seed, progeny plant, or plant part from transgenic maize comprising the maize MON 87419 event, a representative sample of seed comprising said event having been deposited as ATCC PTA-120860.

Claim 5. A DNA molecule comprising a sufficient length of contiguous DNA sequence of SEQ ID NO:10 to function as a DNA probe that hybridizes under stringent hybridization conditions with a DNA sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10, wherein the DNA molecule does not hybridize under the stringent hybridization conditions with a DNA sequence not comprising a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and SEQ ID NO:10, and wherein said DNA molecule comprises SEQ ID NO:1 or SEQ ID NO:2.

Claim 6. A pair of DNA molecules comprising a first DNA molecule and a second DNA molecule, wherein the first DNA molecule is a fragment of SEQ ID NO:9 and the second DNA molecule is a fragment of the maize genomic DNA of the maize MON 87419 event, and wherein the first and second DNA molecules each comprise a DNA sequence of sufficient length of contiguous nucleotides to produce an amplicon diagnostic for the maize MON 87419 event when used together as DNA primers in an amplification reaction with DNA containing the maize MON 87419 event in a sample, wherein said amplicon comprises SEQ ID NO:1 or SEQ ID NO:2.

Claim 7. A method of detecting the presence of the maize MON 87419 event in a sample of DNA, the method comprising:

- a) contacting the sample with the DNA molecule of claim 5;
- b) subjecting the sample and the DNA molecule to stringent hybridization conditions; and
- c) detecting hybridization of the DNA molecule to a target DNA molecule in the sample, wherein the hybridization of the DNA molecule to the target DNA molecule indicates the presence of the maize MON 87419 event in the sample of DNA.

Claim 8. A method of detecting the presence of the maize MON 87419 event in a sample of DNA, the method comprising:

- a) contacting the sample with the pair of DNA molecules of claim 6;
- b) performing an amplification reaction sufficient to produce a DNA amplicon comprising
 - a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8; and
 - c) detecting the presence of the DNA amplicon in the reaction, wherein the presence of the DNA amplicon in the reaction indicates the presence of the maize MON 87419 event in the sample of DNA.

Claim 9. A DNA detection kit when used for detecting the presence of the maize MON 87419 event comprising:

- a) a DNA molecule comprising SEQ ID NO:1 or SEQ ID NO:2; or
- b) a pair of DNA molecules comprising a first DNA molecule and a second DNA molecule, wherein the first DNA molecule is a fragment of SEQ ID NO:9 and the second DNA molecule is a fragment of the maize genomic DNA of the maize MON 87419 event, and wherein the first and second DNA molecules each comprise a DNA sequence of sufficient length of contiguous nucleotides when used as DNA primers together in an amplification reaction with DNA containing the maize MON 87419 event to produce an amplicon diagnostic for the maize MON 87419 event in a sample, wherein said amplicon comprises SEQ ID NO:1 or SEQ ID NO:2.

Claim 10. A transgenic maize plant, seed, cell, plant part, or commodity product comprising a DNA molecule having a DNA sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:10.

Claim 11. The transgenic maize plant, seed, or cell of claim 10, wherein the plant, seed, or cell is tolerant to glufosinate or dicamba, or glufosinate and dicamba herbicides.

Claim 12. A transgenic maize plant, seed, cell, plant part, or commodity product comprising the maize MON 87419 event, a representative sample of seed comprising said event having been deposited as ATCC Accession No. PTA-120860.

Claim 13. The transgenic maize plant or seed of claim 12, wherein the maize plant or seed is a hybrid having at least one parent plant comprising the maize MON 87419 event.

Claim 14. A method for controlling weeds in an area comprising planting transgenic maize comprising the maize MON 87419 event in an area and applying an effective dose of dicamba, or glufosinate, or dicamba and glufosinate herbicides to control the weeds in the area without injuring the transgenic maize.

Claim 15. The method of claim 14, wherein the effective dose of glufosinate herbicide is a total of about 0.1 pounds acid equivalent per acre to about 16 pounds acid equivalent per acre of glufosinate herbicide over a growing season.

Claim 16. The method of claim 14, wherein the effective dose of glufosinate herbicide is a total of about 0.4 pounds acid equivalent per acre to about 1.59 pounds acid equivalent per acre of glufosinate herbicide over a growing season

Claim 17. The method of claim 14, wherein the effective dose of dicamba herbicide is a total of about 0.1 pounds acid equivalent per acre to about 16 pounds acid equivalent per acre of dicamba herbicide over a growing season.

Claim 18. The method of claim 14, wherein the effective dose of dicamba herbicide is a total of about 0.5 pounds acid equivalent per acre to about 2 pounds acid equivalent per acre of dicamba herbicide over a growing season.

Claim 19. A method of producing a transgenic maize plant that is tolerant to glufosinate and dicamba herbicides, the method comprising:

- a) sexually crossing a transgenic maize plant comprising the maize MON 87419 event with itself or a second maize plant;
- b) collecting the seed produced;
- c) growing the seed to produce progeny plants;

- d) treating the progeny plants with glufosinate, or dicamba, or glufosinate and dicamba herbicides; and
- e) selecting a progeny plant that is tolerant to glufosinate and dicamba herbicides.

Claim 20. A method of growing a transgenic maize plant that is tolerant to glufosinate and dicamba herbicides, the method comprising:

- a) planting a maize seed comprising event MON 87419;
- b) allowing a plant to grow from said seed; and
- c) treating the plant with glufosinate, or dicamba, or glufosinate and dicamba herbicides.

Claim 21. A method of producing a plant that tolerates application of glufosinate and dicamba herbicides, the method comprising:

- a) providing a DNA construct comprising the transgene insert comprising: (i) the promoter, leader, and intron of the ubiquitin gene from *Andropogon gerardii*; operably linked to the phosphinothricin N-acetyltransferase (PAT) coding sequence from *Streptomyces viridochromogenes*; operably linked to the polyadenylation signal from the RA5B precursor gene of *Oryza sativa* and (ii) the promoter for the full length transcript of Peanut Chlorotic Streak Virus with a duplicated enhancer region; operably linked to the leader of the light harvesting complex b1 gene from *Triticum aestivum*; operably linked to the first intron from the actin 1 gene from *Oryza sativa*; operably linked to the N-terminal chloroplast transit peptide from the *Petunia x hybrida* 5-enolpyruvylshikimate-3-phosphate synthase gene; operably linked to the dicamba monooxygenase (DMO) coding sequence from *Stenotrophomonas maltophilia* optimized for monocot expression; operably linked to the heat shock protein 17 polyadenylation signal from *Triticum aestivum*;
- b) introducing the DNA construct into a maize plant cell; and
- c) regenerating the maize plant cell to produce a maize plant such that the plant comprises a sequence selected from SEQ ID NO:1-8 and 10, wherein the plant tolerates application of glufosinate and dicamba herbicides.

Claim 22. The method of claim 21, wherein the transgene insert comprises SEQ ID NO:9.

Claim 23. A maize plant produced by the method of claim 21.

Claim 24. A progeny plant, seed, or part of the maize plant of claim 23.




Figure 1

MONS362WO. txt
SEQUENCE LISTING

<110> Monsanto Technology LLC
<120> Transgenic Maize Event MON 87419 and Methods of Use Thereof
<130> MONS: 362WO
<140> Unknown
<141> 2015-03-10
<150> 61/968, 342
<151> 2014-03-20
<160> 22
<170> PatentIn version 3.5
<210> 1
<211> 30
<212> DNA
<213> Artificial Sequence
<220>
<223> Recombinant
<400> 1
caggtattga tgtgcgccag tcagcatcat 30
<210> 2
<211> 30
<212> DNA
<213> Artificial Sequence
<220>
<223> Recombinant
<400> 2
ttcttttct ccatagcatt cgcaatacag 30
<210> 3
<211> 60
<212> DNA
<213> Artificial Sequence
<220>
<223> Recombinant
<400> 3
gcaggcggtc gctgccaggt attgatgtgc gccagtcagc atcatcacac caaaagttag 60
<210> 4
<211> 60
<212> DNA
<213> Artificial Sequence
<220>
<223> Recombinant
<400> 4
attggtaatt actctttctt tttctccata gcattcgcaa tacagttaga tgcgagtgaa 60
<210> 5
<211> 100

MONS362W0. txt

<212> DNA
<213> Artificial Sequence

<220>
<223> Recombinant

<400> 5
agggggggg tggcggggtg gcaggcggtc gctgccaggt attgatgtgc gccagtcagc 60
atcatcacac caaaagttag gcccgaatag tttgaaatta 100

<210> 6
<211> 100
<212> DNA
<213> Artificial Sequence

<220>
<223> Recombinant

<400> 6
cattttgaa ttgaaaaaaaaa attggtaatt actcttctt tttctccata gcattcgcaa 60
tacagttaga tgcgagtgaa gcacgataag tcacaaccat 100

<210> 7
<211> 1771
<212> DNA
<213> Artificial Sequence

<220>
<223> Recombinant

<400> 7
gttgatattt tgacccaaag acatgccatt ctcattgttt tcagtttat cctaccaatg 60
tagtaaccca aacagtttaa ttacaagtga atcaactgga aacaagatct acagactgca 120
cacagcacat taacaattct gcgc当地atg aatgacagag tgatttatag gcactaagct 180
ggttttcatc atgaattttg atcagagcat aaacctgtt gacaacagaa tagaatcaaa 240
caagctgctc agctcagatg gacaaaata atgaacgcgc gcacagtata ttgaaccaaa 300
atagtgaacg cacagtagaa aacagacgga gaagcttagt cacactaaaa atagaactca 360
ctgatgcatt tttaccgaa tcattaaaaa acagaccaac atgatttagac cagctatcaa 420
aagacacact gctgagaatc aacaactgct atatgtgtga tctacctgaa ataaacagat 480
gcagttttt gtccctacag accaacatgt tttatcaga atgccaatga taaatcctta 540
gagatctgga gagcaaacta attaaccact gagaatcaac aactgctata atccttagag 600
atctggagag cacactaatt aaccactgag cctctttgtt cagcatgcag aaatccccaa 660
aactatactg taattcgaaa cccataagcg cacatcaaag caactccatt gttgatggtt 720
ggaaatgtta aaaatatggt tggacacc aattgccgt aacaaaccca cagggagaac 780
atgagttatgtt acaatcaaaa ctaacatgaa taagtggatg ttcagattaa gtaagaaaa 840
tggcagatggtt attaagaaca gagcaatgtt tagcacaagt gggaaattaca aacctttttt 900
cacaatcaatgtt caatctctgtt acagcacaca gcaagatctt ggaatggatg agggatgtt 960
atcggtgaaa aaagacagca cacagcaaga tctctagaac agaacatgaa atgttggaga 1020

MONS362W0. txt

gtggggatgg	gtttagacc	taaccttgt	ccatcctttt	aatagccgtc	gctgccaggt	1080
atggaatagg	gcgcgtgag	tgccgtcgag	ggcgccgca	ggggcgcggt	ttggtgtggg	1140
cgggataccc	ggcgccgtcg	ggaaacccca	cggcggtcat	cacggcgagc	cggggaggg	1200
gggggttggc	gggtggcag	gcggtcgtg	ccaggtattg	atgtgcgcca	gtcagcatca	1260
tcacacaaa	agtaggccc	aatagttt	aaattagaaa	gctcgcaatt	gaggtctgtc	1320
gaggaggta	acctaggta	tgaattaccc	tgttatccct	aactagatat	cgaattctaa	1380
ctataacgg	cctaaggtag	cgaatcggtc	ttcttggcgc	gccaagacgc	aaactcggac	1440
cgttcaagc	cgtcaaggca	cttctatgca	accacagtca	acttgaatgc	cgcttgagtg	1500
ccttctcaag	ttttttttc	ttgaaaaat	catttctttt	ttttaaaaaaa	agtataattt	1560
ggatcgtgca	aatttcttc	taggtgtgt	tgtgactgtg	ttagtaacaa	tttctctagt	1620
tgtgcgcgac	tgctgcttac	tttggagatt	acaatatctt	tctaaaatgc	ttcgattact	1680
tatttataaa	ccgtctctaa	ggccaattgc	tcaagattca	ttcaacaattt	gaaacgtctc	1740
acatgattaa	atcatataaa	gtttctaagt	c			1771

<210> 8
 <211> 1767
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Recombinant

<400> 8	gcgagtctga	tgagacatct	ctgtattgt	tttcttccc	cagtgtttc	tgtacttgt	60
	taatcgctta	atcgccaaca	gattcggcga	tgaataatg	agaaataat	tgttctgatt	120
	ttgagtgcaa	aaaaaaagga	attagatctg	tgtgtttt	ttggatcccc	ggggcggccg	180
	cacaacaaac	gaatacgtcc	tgcttggcct	actaggccaa	cgcaggcgct	ggccgtgacg	240
	gccacgagcg	aactgatatac	gaattctagg	gataacaggg	taatccacgt	gtagctaaac	300
	gcccctcat	ctaagcccc	atttggacgt	gaatgtagac	acgtcgaaat	aaagatttcc	360
	gaattagaat	aatttgttta	ttgcttcgc	ctataaatac	gacggatcgt	aatttgcgt	420
	tttatcaaaa	tgtacttca	ttttataata	acgctgcgga	catctacatt	tttgaattga	480
	aaaaaaattt	gtaattactc	tttcttttcc	tccatagcat	tcgcaataca	gttagatgcg	540
	agtgaagcac	gataagtac	aaccataata	catactatta	gaatccggct	ctttgccag	600
	tgctttttt	gggcactcgg	caaagacttc	tttaccggca	aagtccact	ctcggttaacg	660
	accacgttta	ccgagagcag	gacgctcggt	acagggagac	actcggcaaa	gaccttttg	720
	tcgagtgc	aacgctcg	aaaggccgt	cagcagccgt	ctatagcaga	tgtttattat	780
	ctttgccag	cgccaagcgt	tggcactcgg	caaaacaat	gttgccgagt	gcttaaattt	840
	gacactcggc	aaaatataatt	tttctttttt	cttttgcaa	ccaaactttc	tgtggttgt	900
	tcctacacta	tgttagacata	catgttccat	tttggcacaa	ttataaaaat	gtttgctata	960

MONS362W0. txt

aatattagat	tttggcggtt	taattgaatt	ttctcgata	attcagattt	gaactacaag	1020
tcactcaaaa	aatggaaaac	catgaatgca	aaaatgatat	ccatgttatt	tagcacaagt	1080
tacggccat	ttcaggagta	gaccgaatt	tttggagcacc	atgctcacga	aacatgactg	1140
tgaacttgc	atccagttgt	tttaaaattt	tataaaacac	aaacaaaatc	aaaaaatcat	1200
gaaacttgc	cacatgtcat	gatatcatca	tatgttaggg	ttgtgataaa	aagttgaaaa	1260
tatTCgtga	aagttgtcca	tagttcactg	agcctcttgc	tcgagtgtca	cactcgccaa	1320
agccttgc	gagtgttca	gacactcgcc	aaataagttt	tttccagtag	tgacattaat	1380
agaaaaagag	tgcatagttc	gttcacatag	ctaactaatt	ggttcattac	aaagaaacat	1440
tagaacaagc	ataatgtga	cttggacata	atgaaataaa	atacataggt	tgagggacc	1500
aaacataatg	gaacacagaa	caaataaggg	acaacatcat	agaaaacgac	ataacttcac	1560
atgggttttta	cgcatttgc	tccacgcca	gtccataagt	gctcgatgag	atcatgttga	1620
agttgggtgt	atcattttag	actagtgtat	ttgtggtggg	agcatagcca	ctctgtacga	1680
acttcaggga	ttgtgatgag	agatccacgt	gcactccaat	tgccgaaac	tcaacattac	1740
ttgcacgtgg	gaccttgc	tcgacaa				1767

<210> 9
 <211> 6762
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Recombinant

<400> 9	gccagtgc	atcatcacac	caaaagttt	gcccgaatag	tttggaaattt	gaaagctcgc	60
aattgagg	tcgtcgagg	gttacccat	gtactgaatt	accctgttat	ccctaactag		120
atatcgaa	tttcaactataa	cggccta	gtacgtatc	gttcttcttgc	gcgcgcca		180
acgcaaactc	ggaccgg	ttcaaaagtttca	ggcacttctt	tgcaccacac	gtcaacttga		240
atgcgc	tttgccttct	caagttttttt	tttcttgc	aaatcatttc	ttttttttaa		300
aaaaagtata	atttggatcg	tgc	aaatcatttc	tctctaggtt	tgtgtgtac		360
acaatttctc	tagttgtcg	cgactgctc	tttacttttgc	gattacaata	tctttctaa		420
atgc	tttgccttct	taaaccgtct	ctaaaggccaa	ttgctcaaga	ttcattcaac		480
aattgaaac	tctcacatga	ttaaatcata	taaagtttct	aagtcttgc	tgacaagatt		540
tttttagatt	ttcatctaaa	ttggatgaaa	ctatcaaaca	ctaaattttaa	aaaatataag		600
agaagctcc	gagataaaag	gtcgtctat	tttattataag	agtaaaagtgc	tctattctct		660
tcgtcc	ccatataatt	ctaaaggcat	attgcttct	tttggacaa	aggagcatg		720
ccacaacaca	agaatgtatgt	caccgtcat	cttggatcct	tttatggtaa	agttcacct		780
tctataatct	aacaatagag	aaatcaggaa	aaaatcatgt	tttggttgtt	tttatttcta		840
acctccacaa	taacttttgtt	ttaccat	ttgtttgatt	ttgttttag	agaagcg		900

MONS362W0. txt

ataacaggac	ctaaaatctt	tttcagtagc	acagtacaac	gcagacgctc	atacacgcac	960
gcacactcac	ctctatgaac	acacgtaaga	aaaccctaca	cctttagcac	cttcgaagga	1020
ctgagccgt	aaatatacg	attctcgaa	tcactattag	cgccctcggt	tcaacggaa	1080
tgtcgcttac	cacttaaagc	ataacgcca	gaaatcccgt	aataaatcca	gtaaaatacg	1140
agcaccctgt	ccaagttgaa	tatttgaacc	cgagtggta	gattccaccg	caaaggacct	1200
aaccagatca	tttcgcaaac	aggaactaaa	atcggttagag	agcccagaca	aaagccttc	1260
ctaagagcca	ctccagtgaa	agcccctact	ttaggatataa	aatgcaatac	tagtgggct	1320
cctaaataaa	cttctatttt	tcatggcctt	ctaaaattca	ctcccaaacc	cctagctata	1380
gaagtcttt	atccatcctc	taaataaaaa	tgggagttca	ttttatttca	ccagagttga	1440
tcgtaaattt	agtctctcaa	attttataag	ttgagggtag	aggatgactg	gagttgctct	1500
aaacggacct	atcttcaagt	gacctcagt	agcccgttta	acggcgtcga	caagtttaat	1560
ctaacggaca	ccaaccagag	aagagaacca	ccgcccagcgc	cgagccaagc	gacgttgaca	1620
tcttggcgcg	gcacggcatc	tccctggcgt	ctggccccc	ctcgagactt	ccgctccacc	1680
tcccaccgg	ggcggtttcc	aagtccgttc	cgcctcctt	cacacggcac	gaaaccgtga	1740
cgggcaccgg	cagcacgggg	ggattccctt	cccaccgctc	cttcccttcc	ccttcctctc	1800
ccgcccgtat	aaatagccag	ccccatcccc	agcttcttcc	cccaacctca	tcttctctcg	1860
tgttggcgg	cacaacccga	tcgatcccc	actccctcgt	cgtctctcct	cgcgagcctc	1920
gtcgatcccc	cgcttcaagg	tacggcgatc	gattatcttc	cctctctcta	ccttctctct	1980
ctttagggc	ctgctagctc	tgttccgtt	tttccatggc	tgcgaggtac	aatagatcgg	2040
cgatccatgg	ttagggcctg	ctagttgtgt	tcctgttttt	ccatggctgc	gaggcacaat	2100
agatctgatg	gcgttatgt	ggttaacttg	tcatacttt	gcgtatctatg	gtcccttttag	2160
gagtttagga	catctattta	atttcggata	gttcgagatc	tgtgatccat	ggttagtacc	2220
ctaggcagt	gggtagatc	cgtgtgtt	tggttcgtag	atggattctg	attgctcagt	2280
aactggaaat	cctggatgg	ttctagctgg	ttcgcagata	agatcgattt	catgatatgc	2340
tatacttgt	ttgggtgccg	tggttccgtt	aatctgtct	gttatgtatct	tagtcttga	2400
taaggttcg	tcgtgttagc	tacgtcctgt	gcagcactta	attgtcaggt	cataattttt	2460
agcatgcctt	tttttattt	gtttgggttt	gtctgactgg	gctgttagata	gtttcaatct	2520
ttgtctgact	gggctgtaga	tagttcaat	ctacctgtcg	gtttattttt	ttaaatttgg	2580
atctgtatgt	gtgtcatata	tcttcatctt	ttagatataat	cgataggttt	atatgttgc	2640
gtcggttttt	tactgttcc	ttatgagata	tattcatgct	tagatacatg	aaacaacgtg	2700
ctgttacagt	ttaatagttc	ttgttatct	aataaaca	taaggatagg	tatatgctgc	2760
agttatgttt	actggactt	ttttgacat	gaaccctacgg	cttaataatt	agtcttcatc	2820
aaataaaaag	catatttttt	aattatccg	atatacttga	atgtatgtat	atgcagcatc	2880
tgtgtgaatt	tttggccctg	tcttcatatg	ctgtttattt	gtttgggact	gtttcttgg	2940

MONS362W0. txt

ttgataactc atcctgttgt ttggtgatcc ttttgcaggt gcaaccatgt ctccggagag	3000
gagaccagtt gagattaggc cagctacagc agctgatatg gccgcgggtt gtgatatcgt	3060
taaccattac attgagacgt ctacagtcaa ctttaggaca gagccacaaa caccacaaga	3120
gtggattgat gatctagaga ggttgcaga tagataccct tggttgggtt ctgagggtga	3180
gggtgttgt gctggatttg cttacgctgg gccctgaaag gcttaggaacg cttacgattt	3240
gacagttgag agtactgttt acgtgtcaca taggcatcaa agtttggcc taggatccac	3300
attgtacaca catttgctta agtctatgga ggcgcaggt ttaagtctg tggttgcgt	3360
tataggcctt ccaaacgatc catctgttag gttgcattgag gctttggat acacagcccg	3420
gggtacattt cgccgcagctg gatacaagca tggtggatgg catgtatgtt gttttggca	3480
aagggatttt gagttgccag ctcctccaag gccagttagg ccagttaccc agatctgatt	3540
aattaactag gctactgttag ctatgtgc atgtatgtgg tgtggttact aaaataatta	3600
tggtttcct tttgtttgga agcatatgtg tggtgaataa atgatgaact ccgatgttcc	3660
tctctataaa tcttgatgat tcgctagcta tccgtacgtc gttgttcttt gatttgatga	3720
ttagattgaa aaatgaaatg tcattgtcaag gagggtgccg cggccggcc gtgacggcca	3780
cgagcgaact cctgcaggac aacggagcag ctcctcagc aaatcctacc acctcattt	3840
aatagagtga ggttggatgg ctgaggtacg ggccgcgtt acaagcttct gcagaattcg	3900
tcaacgagat cttgagccaa tcaaagagga gtgatgtaga cctaaagcaa taatggagcc	3960
atgacgtaag ggcttacgcc catacgaaat aattaaaggc ttagtgcacc tgtcggtctc	4020
tcagaacctt tactttttat gttggcgtg tattttaaa tttccacggc aatgacgatg	4080
tgacccaacg agatctttag ccaatcaaag aggagtgtatg tagacctaataa gcaataatgg	4140
agccatgacg taagggctt cgcccatagc aaataattaa aggctgtatgt gacctgtcgg	4200
tctctcagaa cctttacttt ttatatttgg cgtgtatttt taaatttcca cggcaatgac	4260
gatgtgaccc gtgcattccgc tttgcctata aataagttt agttttagt gatgcacacg	4320
gtcgagaaga cacggccatt cttagaccat cttccacaca ctcaagccac actattggag	4380
aacacacagg gacaacacac cataagatcc aaggaggcc tccgcccggc cggtaacca	4440
ccccggccct ctccttttc tttctccgtt ttttttccg tctcggtctc gatctttggc	4500
cttggtagtt tgggtggcg agaggcgct tcgtgcgcgc ccagatcggt gcgcgggagg	4560
ggcgggatct cgccgcgtgg gctctcgccg gcgtggatcc gccccggatc tcgcggggaa	4620
tggggctctc ggtatgtatgt ctgcgtatccg ccgttgggttgg gggagatgtatgttta	4680
aaatttccgc cgtgctaaac aagatcagga agagggaaaa agggcactat ggtttatatt	4740
tttatatatatt tctgctgctt cgtcaggctt agatgtgcta gatctttctt tcttctttt	4800
gtgggtagaa tttgaatccc tcagcattgt tcattcgatg tttttctttt catgatttgc	4860
gacaaatgca gcctcgatcg gagctttttt gtaggtagaa gtgatcaacc atggcccaga	4920
tcaacaacat ggcccaggc atccagaccc tgaaccctaa ctctaacttc cacaagccgc	4980

MONS362W0.txt

aagtgcccaa	gtctagctcc	ttcctcggt	tcggctccaa	gaagctcaag	aatagcgcca	5040
attccatgct	ggtcctgaag	aaagactcga	tcttcatgca	gaagttctgc	tccttcgca	5100
tcagtgcttc	ggttgcgact	gcctgcatgc	tcacccctcg	taggaacgccc	tggtaacgtcg	5160
ccgctctccc	tgaggagctg	agcgagaagc	ccttgggtcg	caccatccta	gacactccgt	5220
tagccctta	ccgcccagcct	gacggcgtag	tggcggccct	gcttgcacatc	tgcccgata	5280
ggttcgctcc	gctcagcgcac	ggcatcctcg	tcaacgggca	tcttcagtc	ccgtaccacg	5340
ggcttggaaatt	tgacggcggt	gggcagtgtg	tccacaaccc	gcacggcaac	ggcgacacggc	5400
cagcttccct	caacgtttagg	tcgttccctg	ttgtcgagcg	cgacgcactg	atctggatct	5460
ggcctggcga	cccagctctg	gccgatccag	gagccattcc	cgacttcggt	tgcccgctgg	5520
acccagccta	tcggacggtc	ggcggttacg	ggcacgtcga	ttgttaactat	aagctccctg	5580
tggacaacct	tatggatttgc	ggccacgctc	agtacgtcga	ccgggctaacc	gctcagactg	5640
acgccttga	ccgtctcgaa	aggaggatca	tcgtcggcga	cgagagagatt	caggcgctga	5700
tgaagatccc	tggaggcacf	ccctctgtgc	tcatggcgaa	gtttctcaga	ggcgcaaca	5760
cgcggcgatgga	cgcctggaaac	gacatccgct	ggaataaggt	ctccgcgatg	ctgaacttca	5820
tcgcccgtgc	gcccgagggc	acacccaaag	agcagtaat	ccacagcaga	gggaccata	5880
ttcttacacc	ggaaaccgag	gctagttgcc	actacttctt	cggtcgatc	cggaatttcg	5940
ggatagacga	tccggagatg	gacgggttgc	ttcgatcttgc	gcaagcgca	gctctcgatc	6000
aggaagataa	ggtggtcgtg	gaggctatcg	agcgtaggcg	cgcctacgtt	gaggcgaacg	6060
gtattaggcc	cgcgatgctg	tcctgcgacg	aggccgcagt	taggtgtcg	cgcgagatag	6120
aaaagctgga	gcagctagag	gccgcctgag	gtaccgagct	cgtcaatcac	tagtgaattc	6180
tgcattgcgtt	tggacgtatg	ctcattcagg	ttggagccaa	tttggttgat	gtgtgtgcga	6240
gttcttgcga	gtctgatgag	acatctctgt	attgttttc	tttccccagt	gttttctgtat	6300
cttgcgtat	cggctaattcg	ccaacagatt	cggcgatgaa	taaatgagaa	ataaaattgtt	6360
ctgatttga	gtgcaaaaaaa	aaaggaatta	gatctgtgt	tgtttttgg	atccccgggg	6420
cggccgcaca	acaaacgaat	acgtcctgct	tggcctacta	ggccaacgca	ggcgctggcc	6480
gtgacggcca	cgagcgaact	gatatcgaaat	tctaggata	acagggtaat	ccacgtgtag	6540
ctaaacgcgc	cctcatctaa	gccccattt	ggacgtaaat	gtagacacgt	cgaataaaag	6600
atttccgaat	tagaataatt	tgtttattgc	tttcgcctat	aaatacgcac	gatcgtaatt	6660
tgtcgttta	tcaaaatgtat	cttcattttt	ataataacgc	tgcggacatc	tacatttttg	6720
aattgaaaaaa	aaattggtaa	ttactcttgc	tttttctcca	ta		6762

<210> 10
 <211> 9259
 <212> DNA
 <213> Artificial Sequence

<220>

MONS362W0.txt

<223> Recombinant

<400>	10					
gttgatattt	tgacccaaag	acatgccatt	ctcattgttt	tcagtttat	cctaccaatg	60
tagtaaccca	aacagttaa	ttacaagtga	atcaactgga	aacaagatct	acagactgca	120
cacagcacat	taacaattct	gcgccaatg	aatgacagag	tgatttata	gcactaagct	180
ggtttcatc	atgaattttg	atcagagcat	aaaccttgtt	gacaacagaa	tagaatcaaa	240
caagctgctc	agctcagatg	gacaaaata	atgaacgcgc	gcacagtata	ttgaaccaaa	300
atagtgaacg	cacagttagaa	aacagacgga	gaagctagt	cacactaaaa	atagaactca	360
ctgatgcatt	tttaaccgaa	tcattaaaaa	acagaccaac	atgatttagac	cagctatcaa	420
aagacacact	gctgagaatc	aacaactgct	atatgtgt	tctacctgaa	ataaacagat	480
gcagttttt	gtccctacag	accaacatgt	tttaatcaga	atgccaatga	taaatccta	540
gagatctgga	gagcaaacta	attaaccact	gagaatcaac	aactgctata	atccttagag	600
atctggagag	cacactaatt	aaccactgag	cctcttgg	cagcatgcag	aaatccccaa	660
aactatactg	taattcgaaa	cccataagcg	cacatcaaag	caactccatt	gttgatggtt	720
ggaatgtta	aaaatatggt	tgtggacacc	aattgccgt	aacaaaccca	cagggagaac	780
atgagtatga	acaatcaaaa	ctaacatgaa	taagtggat	ttcagattaa	gtaaggaaaa	840
tgggcacag	attaagaaca	gagcaatgta	tagcacaagt	ggaattaca	aactttttt	900
cacaaaatta	caatctctag	acagcacaca	gcaagatctt	ggaatggat	agggtgatga	960
atcgggtaaa	aaagacagca	cacagcaaga	tctctagaac	agaacatgaa	atgtggaga	1020
gtggggatgg	gtttagacc	taacctctgt	ccatcctttt	aatagccgtc	gctgccaggt	1080
atggaatagg	gcccgtgag	tgccgtcgag	ggcgccgc	ggggcgcggt	ttgggtgtgg	1140
cgggataccc	ggcgccgtcg	ggaaacccca	cggcggtcat	cacggcgagc	cggggaggg	1200
gggggttggc	gggtggcag	gcggtcgt	ccaggtattt	atgtgcgcca	gtcagcatca	1260
tcacaccaaa	agttaggccc	aatagttt	aaattagaaa	gctcgcaatt	gaggctgtc	1320
gaggaggta	acctaggta	tgaattaccc	tgttatccct	aactagat	cgaattctaa	1380
ctataacgg	cctaaggtag	cgaatcg	ttcttggcgc	gccaagacgc	aaactcggac	1440
cgttcaagc	cgtcaaggca	cttctatgca	accacagtca	acttgaatgc	cgttgagtg	1500
ccttctcaag	ttttttttc	ttgaaaaat	catttctttt	ttttaaaaaaa	agtataattt	1560
ggatcgtca	aatttctctc	taggtgtgt	tgtgactgt	tgagtaacaa	tttctctgt	1620
tgtgcgcac	tgctgcttac	tttggagatt	acaatatctt	tctaaaatgc	ttcgattact	1680
tatttataaa	ccgtctctaa	ggccaattgc	tcaagattca	ttcaacaatt	gaaacgtctc	1740
acatgattaa	atcatataaa	gttctta	cttgcgtac	aagattttt	tagattttca	1800
tctaaattgg	atgaaactat	caaacactaa	ttttaaaaaaa	tataagagaa	gctccggaga	1860
taaaaggctc	tctatgttat	tataagagta	aagtcgtcta	ttctcttcgt	cccaacat	1920
ataattctaa	gcatgaattt	cttctttt	ggacaaaagg	agcatgccac	aacacaagaa	1980

MONS362W0. txt

tgtatgtcacc gtcatgcttg gatccttta tggtaaagct tcaccctcta taatctaaca 2040
atagagaat cagggaaaaa tcatgttttgc ttgttttta tttctaacct ccacaataac 2100
tttggtttac catttttgc ttgattttag ttttagagaa gcgttataa caggacctaa 2160
aatctttt cagtacacag tacaacgcag acgctcatac acgcacgcac actcacctct 2220
atgaacacac gtaagaaaac cctacacctt gaggcaccc tcgaaggactga gccggtaat 2280
atagagattc tcgaagtcac tattagcgcc tcgttgtcaa cggaaatgtc gcttaccact 2340
taaagcataa cggcgagaaa tcccgtataa aatccagtaa aatacgagca cccgtgcca 2400
gttgaatatt tgaacccgag tggtagatt ccacccgaaa ggacctaacc agatcatttc 2460
gcaaaacagga actaaaatcg gtagagagcc cagacaaaag ctttcctaa gagccactcc 2520
agtggaaagcc cctacttttag gtataaaaatg caatactagt ggggctccta aataaaacttc 2580
tattttcat ggccttctaa aattcactcc caaaccctta gctatagaag tctcttatcc 2640
atcctctaaa taaaaatggg agtctatTTT atttcaccag agttgatcgt aaatTTtagtc 2700
tctcaaattt tataagttga gggtagagga tgactggagt tgctctaaac ggacctatct 2760
tcaagtgacc tcagtgagcc cggttaacgg cgtagacaag tttaatctaa cggacaccaa 2820
ccagagaaga gaaccaccgc cagcgccgag ccaagcgacg ttgacatctt ggcgcggcac 2880
ggcatctccc tggcgtctgg cccctctcg agactccgc tccacccccc accggtggcg 2940
gtttccaagt ccgttccgccc tcctctcaca cggcacgaaa ccgtgacggg caccggcagc 3000
acggggggat tccttccca ccgtcccttc ctttccctt cctctccgc cgctataaat 3060
agccagcccc atccccagct tctttccca acctcatctt ctctcgtgtt gttcggcaca 3120
acccgatcga tcccccaactc cctcgctc tctcccgcg agccctcgat atcccccgct 3180
tcaaggtagc gcgatcgatt atcttccctc tctctacctt ctctctctta tagggcctgc 3240
tagctctgtt cctgttttc catggctcg aggtacaata gatggcgat ccattgttag 3300
ggcctgttag ttgtgttccat ggtgtcgagg cacaatagat ctgtggcgt 3360
tatgtatggtt aacttgtcat actcttgca tctatggtcc cttaggagt ttaggacatc 3420
tatttaattt cgatagttc gagatctgtg atccatgtt agtaccctag gcagtggggt 3480
tagatccgtg ctgttatggt tcgttagatgg attctgttgc ctctgtact ggaaatccgt 3540
ggatggttct agctggttcg cagataagat cgatttcgt atatgtata tcttgggtgg 3600
ttgccgtggt tccgttaaat ctgtctgtta tgatcttagt ctgtataag gttcggtcg 3660
gctagctacg tcctgtgcag cacttaattt tcaggtcata attttagca tgccttttt 3720
ttattggttt ggtttgtct gactgggtcg tagatgtttt caatcttgcgt ctgactggc 3780
tgttagatgt ttcaatctac ctgtcggtt attttattaa atttggatct gtatgtgtgt 3840
catatatctt catcttttag atatatcgat aggtttatat gttgtgtcg gttttttact 3900
gttcctttat gagatatatt catgcttaga tacatgaaac aacgtgtgt tacagttaa 3960
tagttctgtt ttatctaata aacaataag gataggtata tgctgcgtt agttttactg 4020

MONS362WO. txt

MONS362W0. txt

aatccctcag	cattgttcat	cggtagttt	tctttcatg	atttgtgaca	aatgcagcct	6120
cgtgcggagc	ttttttag	gtagaagtga	tcaaccatgg	cccagatcaa	caacatggcc	6180
cagggcatcc	agaccctgaa	ccctaactct	aacttccaca	agccgcaagt	gcccaagtct	6240
agctccttcc	tcgtgttcgg	ctccaagaag	ctcaagaata	gccccaaattc	catgctggtc	6300
ctgaagaaag	actcgatctt	catcgagaag	ttctgctcct	ttcgcatcag	tgcttcggtt	6360
gcgactgcct	gcatgctcac	cttcgttagg	aacgcctggt	acgtcgccgc	tctccctgag	6420
gagctgagcg	agaagccctt	gggtcgacc	atcctagaca	ctccgttagc	ccttaccgc	6480
cagcctgacg	gcgttagtggc	ggccctgctt	gacatctgcc	cgcatacggtt	cgctccgctc	6540
agcgacggca	tcctcgtaa	cggcatctt	cagtcccgt	accacgggct	ggaatttgc	6600
ggcgggtggc	agtgtgtcca	caacccgcac	ggcaacggcg	cacggccagc	ttccctcaac	6660
gttaggtcgt	tccctgttgt	cgagcgcac	gcactgatct	ggatctggcc	tggcgacc	6720
gctctggccg	atccaggagc	cattccgac	ttcggtgcc	gcgtggaccc	agcctatcgg	6780
acggtcggcg	gttacgggca	cgtcgattgt	aactataagc	tccttggtt	caacctttag	6840
gatttggcc	acgctcagta	cgtgcaccgg	gctaacgctc	agactgacgc	ctttgaccgt	6900
ctcgaaggg	aggtcatcgt	cggcgacgga	gagattcagg	cgcgtatgaa	gatccctgga	6960
ggcacgccc	ctgtgctcat	ggcgaagttt	ctcagaggcg	cgaacacgccc	cgtggacgccc	7020
tggaaacgaca	tccgctggaa	taaggtctcc	gcgtatgtt	acttcatcgc	cgttgccccc	7080
gagggcacac	ccaaagagca	gtcaatccac	agcagaggga	cccatattct	tacaccggaa	7140
accgaggcta	gttgcacta	cttcttcggc	tcgtcacgga	atttcggat	agacgatccg	7200
gagatggacg	gtgttcttcg	atcttggcaa	gcgcagctc	tcgtcaagga	agataagg	7260
gtcgtggagg	ctatcgacg	taggcgcgc	tacgttgggg	cgaacggat	taggcccgc	7320
atgctgtcct	gcgacgaggc	cgcagttt	gtgtcgccgc	agatagaaaa	gctggagcag	7380
ctagaggccg	cctgaggta	cgagctcg	aatcactat	gaattctgca	tgcgttgga	7440
cgtatgctca	ttcagggtgg	agccaattt	gtttagtgtt	gtgcgagtt	ttgcgagtct	7500
gatgagacat	ctctgtattt	tgtttttt	cccagtgtt	tctgtactt	tgtatcgcc	7560
taatcgccaa	cagattccgc	gatgaataaa	tgagaataaa	attgttctga	ttttgagtgc	7620
aaaaaaaaaaag	gaatttagatc	tgtgtgtgtt	ttttggatcc	ccggggcgcc	cgcacaacaa	7680
acgaataacgt	cctgcttggc	ctactaggcc	aacgcaggcg	ctggccgt	cggccacgag	7740
cgaactgata	tcgaattcta	gggataacag	ggtatccac	gtgttagctaa	acgcgccctc	7800
atctaagccc	ccatttggac	gtaatgtt	acacgtcgaa	ataaagattt	ccgaattt	7860
ataatttgg	tattgcttcc	gcctataaaat	acgacggatc	gtatgttgc	gttttatcaa	7920
aatgtacttt	catttataaa	taacgctgcg	gacatctaca	ttttgaattt	aaaaaaaat	7980
tggttaattac	tctttctttt	tctccatagc	attcgcaata	cagttatgt	cgagtgaagc	8040
acgataagtc	acaaccataa	tacatactat	tagaatccgg	ctctttccgc	agtgcgtttt	8100

MONS362W0. txt

ttgggcactc ggcaaagact tctttaccgg caaagtccctt ctctcgtaa cgaccacgtt 8160
taccgagagc aggacgctcg gtacagggag acactcggca aagaccttt tgtcgagtgc 8220
caaacgctcg gcaaagggcc gtcagcagcc gtctatagca gatgtttatt atcttgcgg 8280
agcgccaagc gttggcactc ggcaaaacaa gtgttgcga gtgcttaat tggacactcg 8340
gcaaaatata ttttctttt ttcttttgc aaccaaactt tctgtggttt gttcctacac 8400
tatgttagaca tacatgttcc attttggcac aattataaaa gtgttgcta taaatattag 8460
attttggcg ttttaattgaa ttttctcgaa taattcagat ttgaactaca agtcactcaa 8520
aaaatggaaa accatgaatg caaaaatgat atccatgtt ttttagcaca gttacggccg 8580
atttcaggag tagacccgaa ttttgagca ccatgctcac gaaacatgac tgtgaacttg 8640
tcatccagtt gttttaaaat tgtataaaaac acaaacaaaa tcaaaaaatc atgaaacttg 8700
tccacatgtc atgatatcat catatgtaga gttgtgata aaaagttgaa aatatttcgt 8760
gaaagtgtc catagttcac tgagcccttt tgtcgagtgt cacactcggc aaagcccttg 8820
tcgagtgtctt cagacactcg gcaaataagt ttttccagt agtgacatta atagaaaaag 8880
agtgcatagt tcgttcacat agctaactaa ttggttcatt acaaagaaac attagaacaa 8940
gcataatgtat gacttggaca taatgaaata aaatacatag gttgagggaa ccaaacataa 9000
tggaaacacag aacaaataga ggacaacatc atagaaaacg acataacttc acatgggtt 9060
tacgccaatt gctccacgccc aagtccataa gtgctcgatg agatcatgtt gaagttggtt 9120
gtatcatttgc agactagtga ttgtgtggtg ggagcatagc cactctgtac gaacttcagg 9180
gattgtgtatg agagatccac gtgcactcca attgcggcaa actcaacatt acttgcacgt 9240
gggaccttgc cctcgacaa 9259

<210> 11
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Recombinant

<400> 11
gaattgaaaa aaaattggta attactcttt ctt

33

<210> 12
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Recombinant

<400> 12
atcgtgcttc actcgcatct aact

24

<210> 13
<211> 19

MONS362W0. txt

<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 13		19
ctccatagca ttcgcaata		
<210> 14		
<211> 81		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 14		60
attagaaagc tcgcaattga ggtctgtcga ggaggtaac ctaggtactg aattaccctg		
ttatccctaa ctagatatcg a		81
<210> 15		
<211> 61		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 15		60
tggcgcgcca agacgcaaac tcggaccggt tcaagccgtc aaggcacttc tatgcaacca		
c		61
<210> 16		
<211> 59		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 16		59
atcctgttgc ttgggtatcc ttttcaggt gcaaccatgt ctccggagag gagaccagt		
<210> 17		
<211> 56		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 17		56
ccagtttaggc cagttaccca gatctgatta attaactagg ctactgttagc tagctg		
<210> 18		
<211> 99		
<212> DNA		
<213> Artificial Sequence		
<220>		

MONS362W0.txt

<223> Recombinant		
<400> 18		
aaaaatggaa tgtcatgcta aggagggtgc cgccggccgg ccgtgacggc cacgagcgaa		60
ctcctgcagg acaacggagc agcctcctca gcaaattcct		99
<210> 19		
<211> 56		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 19		
tgagggttgc ttgctgaggt agcggccgct ttaacaagct tctgcagaat tcgtca		56
<210> 20		
<211> 55		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 20		
aggtagaagt gatcaaccat ggcccgatc aacaacatgg cccagggcat ccaga		55
<210> 21		
<211> 91		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 21		
cgcgagatag aaaagctgga gcagctagag gccgcctgag gtaccgagct cgtcaatcac		60
tagtgaattc tgcattgcgtt tggacgtatgc		91
<210> 22		
<211> 67		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> Recombinant		
<400> 22		
ctaggataa caggtaatc cacgtgtac taaacgcgcc ctcatctaag ccccatgg		60
gacgtga		67