



(51) International Patent Classification:

C07D 471/04 (2006.01) A61K 31/519 (2006.01)

C07D 487/04 (2006.01) A61K 31/53 (2006.01)

A61P 35/00 (2006.01)

(21) International Application Number:

PCT/IB2018/060015

(22) International Filing Date:

13 December 2018 (13.12.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

201721044886 13 December 2017 (13.12.2017) IN

201821024634 02 July 2018 (02.07.2018) IN

201821040029 23 October 2018 (23.10.2018) IN

(71) **Applicant:** LUPIN LIMITED [IN/IN]; Kalpataru Inspire, 3rd Floor, Off Western Express Highway, Santacruz (East), Maharashtra, Mumbai 400 055 (IN).

(72) **Inventors:** NAIR, Prathap, Sreedharan; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **GUDEDE, Ganesh, Bhausaheb**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **SETHI, Sachin**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **LAGAD, Dipak, Raychand**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **PAWAR, Chetan, Sanjay**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **TRYAMBAKE, Mahadeo, Bhaskar**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **KULKARNI, Chaitanya, Prabhakar**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **HAJARE, Anil, Kashiram**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **GORE, Balasa-**


heb, Arjun; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **KULKARNI, Sanjeev, Anant**; K52, B801, Karve Nagar, Maharashtra, Pune 411052 (IN). **SINDKHEDKAR, Milind, Dattatraya**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **PALLE, Venkata, P.**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN). **KAMBOJ, Rajender, Kumar**; Lupin Limited (Research Park), 46A / 47A, Village Nande, Taluka Mulshi, Maharashtra, Pune 412115 (IN).

(74) **Agent:** MAJUMDAR, Subhotosh et al.; S. Majumdar & Co., 5 Harish Mukherjee Road, State of West Bengal, Kolkata 700 025 (IN).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) **Title:** SUBSTITUTED BICYCLIC HETEROCYCLIC COMPOUNDS AS PRMT5 INHIBITORS



(57) **Abstract:** The invention relates to substituted bicyclic heterocyclic compounds of formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions for treating diseases, disorders or conditions associated with the overexpression of PRMT5 enzyme. The invention also relates to methods of treating diseases, disorders or conditions associated with the overexpression of PRMT5 enzyme.



Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE*

**SUBSTITUTED BICYCLIC HETEROCYCLIC COMPOUNDS AS PRMT5
INHIBITORS**

Field of the Invention

5 The invention relates to substituted bicyclic heterocyclic compounds of formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions for treating diseases, disorders or conditions associated with the overexpression of PRMT5 enzyme. The invention also relates to methods of treating diseases, disorders or conditions associated with the overexpression of PRMT5 enzyme.

Cross-reference to related applications

10 The present application claims the benefit of Indian Provisional Patent Applications Nos. IN 201721044886 filed on December 13, 2017, IN 201821040029 filed on October 23, 2018, and IN 201821024634 filed on July 02, 2018, the disclosures of which are incorporated herein by reference in their entirety for all purposes.

Background to the invention

15 Methylation of proteins is a common post-translational modification that affects the protein's activity and its interaction with other biological molecules. N-methylation typically occurs on the nitrogen atoms of arginine, lysine and histidine residues and there are different families of enzymes that catalyze the methylation reaction, each being specific to the amino acid residue that will be methylated.

20 A family of 9 enzymes, called *Protein Arginine N-Methyl Transferases* (PRMTs), are responsible for the methylation of the guanidinium group of arginine. The guanidinium group of arginine bears 2 terminal nitrogen atoms that undergo monomethylation or dimethylation. Depending on the type of dimethylation, the enzymes are further classified as type I or type II. Type I PRMTs catalyse the monomethylation or
25 the asymmetric dimethylation whereas type II enzymes catalyse the symmetric dimethylation. Some of the substrates that undergo methylation are histones, Sm ribonucleoproteins, MRE11 and p53 binding protein 1.

The methylation of arginine side-chains has an important role to play in various cell functions that include transcription activation as well as transcription repression, mRNA translation, pre-mRNA splicing, protein trafficking and signal transduction. It also occurs on myriad substrates. The enzymatic activity of the PRMTs hence affects cellular processes like cell proliferation, repair of damaged DNA as well as cell cycle and cell death. It has been shown that PRMT enzyme-mediated hypermethylation leads to certain disease conditions like cancer (Nature Reviews Cancer 2013, 13, p37; Cellular and Molecular Life Sciences 2015, 72, p2041; Trends in Biochemical Sciences 2011, 36, p633).


At present, the most studied type II enzyme is PRMT5, which is conserved across the eukaryotic organisms. Overexpression of PRMT5 is linked with carcinogenesis and decreased patient survival in several human malignancies (Cell Mol Life Sci., 2015, 72, p2041). PRMT5 directly interacts with proteins often dysregulated or mutated in cancers, hence a putative oncogene (Mol Cell Biol, 2008, 28, p6262). PRMT5 mediated transcriptional repression of tumor suppressor genes like p53, RB-1, ST7, or upregulation of Cyclin D1, CDK4, CDK6, eLF4E, MITF, FGFR3 associate with the oncogenesis in both solid tumors and hemaological malignancies. PRMT5 is located in the nucleus as well as the cytoplasm and its overexpression has been linked to a wide range of cancers including, but not limited to, glioblastoma multiforme (Oncogene, 2017, 36, p263), prostate cancer (Oncogene, 2017, 36, p1223), and pancreatic cancer (Science, 2016, 351, p1214), mantle cell lymphoma (Nature Chemical Biology, 2015, 11, p432), non-Hodgkin's lymphomas and diffuse large B-cell lymphoma (Journal of Biological Chemistry, 2013, 288, p35534), acute myeloid leukemia (Leukemia, 2018, 32, p499), acute lymphoblastic leukemia (AACR; Cancer Research 2017;77(13 Suppl):Abstract nr 1128), multiple myeloma (Leukemia, 2018, 32, p996), non-small cell lung cancer (The Biochemical Journal, 2012, 446, p235), small cell lung cancer (AACR; Cancer Research 2017;77(13 Suppl):Abstract nr DDT02-04), breast cancer (Cell Reports, 2017, 21, p3498), triple negative breast cancer (AACR; Cancer Res 2015;75(15 Suppl):Abstract nr 4786), gastric cancer (International Journal of Oncology, 2016, 49, p1195), colorectal cancer (Oncotarget, 2015, 6, p22799), ovarian cancer (J Histochem Cytochem 2013, 61, p206), bladder cancer (Clinical Cancer Research, 2018, CCR-18-1270), hepatocellular

cancer (Oncology Reports, 2018, 40, p536), melanoma (PLoS One, 2013, 8, e74710; J Clin Invest. 2018, 128, p517), sarcoma (Oncology Letters, 2018, 16, p2161), oropharyngeal squamous cell carcinoma (Oncotarget, 2017, 8, p14847), chronic myelogenous leukemia (J Clin Invest, 2016, 126, p3961), epidermal squamous cell carcinoma (Carcinogenesis, 2017, 38, p827), nasopharyngeal carcinoma (Oncology Reports, 2016, 35, p1703), neuroblastoma (Molecular Oncology, 2015, 9, p617), endometrial carcinoma (Gynecol Oncol., 2016, 140, p145), cervical cancer (Pharmazie, 2018, 73, p269).. These findings have led to further research which show that inhibiting PRMT5 reduces cell proliferation (Molecular and Cellular Biology 2008, 28, p6262, The Journal of Biological Chemistry 2013, 288, p35534).

Inhibitors of arginine methyl transferases were first disclosed in 2004 by Cheng *et al* in the Journal of Biological Chemistry – Vol. 279 (23), p.23892. Since then, various other compounds and substances having greater selectivity towards either type I or type II arginine methyl transferases have been disclosed. Other publications that disclose small molecules as inhibitors in relation to PRMT5 are: WO2011077133, WO2011079236, WO2014100695, WO2014100716, WO2014100719, WO2014100730, WO2014100734, WO2014128465, WO2014145214, WO2015200677, WO2015200680, WO2015198229, WO2016022605, WO2016034671, WO2016034673, WO2016034675, WO2016038550, WO2016135582, WO2016145150, WO2016178870, WO2017032840 and ACS Medicinal Chemistry Letters 2015, 6, p408.

Summary of the Invention

In accordance with one aspect, the invention provides compound of general formula (I), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,




wherein,

L_1 is selected from $-CR^aR^b-$, $-NR^a-$, S, and O;

$Z=CH$ or N;

- 5 R^a and R^b are independently selected at each occurrence from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

ring A is selected from,



- 10 R^c and R^d are selected from substituted or unsubstituted alkyl or together with the carbon atoms to which they are attached form a C_3 - C_6 cycloalkyl ring;

R is selected from $-NR^4R^5$, hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted heteroaryl and substituted or unsubstituted cycloalkyl;

- 15 R^1 and R^2 together with the carbon atoms to which they are attached form a bond in order to form a $-C=C-$; or R^1 and R^2 together with the carbon atoms to which they are attached form a cyclopropane ring;

$R^{2'}$ and R^{2a} which may be same or different and are independently selected from hydrogen and substituted or unsubstituted alkyl;

- 20 R^3 is independently selected at each occurrence from halogen, cyano, nitro, substituted or unsubstituted alkyl, $-OR^6$, $-NR^7R^8$, substituted or unsubstituted cycloalkyl, $-C(O)OH$, $-C(O)O$ -alkyl, $-C(O)R^9$, $-C(O)NR^7R^8$, $-NR^7C(O)R^9$, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted heterocyclyl;

R^4 and R^5 are independently selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

5

R⁶ is selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

R⁷ and R⁸ are independently selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

- 5 R⁹ is selected from substituted or unsubstituted alkyl and substituted or unsubstituted cycloalkyl;

R¹⁰ is selected from hydrogen, halogen, and substituted or unsubstituted alkyl;

‘n’ is an integer ranging from 0 to 4, both inclusive;

- 10 when an alkyl group is substituted, it is substituted with 1 to 4 substituents independently selected from oxo (=O), halogen, cyano, cycloalkyl, aryl, heteroaryl, heterocyclyl, -OR^{7a}, -C(=O)OH, -C(=O)O(alkyl), -NR^{8a}R^{8b}, -NR^{8a}C(=O)R^{9a}, and -C(=O)NR^{8a}R^{8b};

- when the heteroaryl group is substituted, it is substituted with 1 to 4 substituents independently selected from halogen, nitro, cyano, alkyl, haloalkyl, perhaloalkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, -OR^{7a}, -NR^{8a}R^{8b}, -NR^{7a}C(=O)R^{9a}, -C(=O)R^{9a}, -C(=O)NR^{8a}R^{8b}, -SO₂-alkyl, -C(=O)OH, and -C(=O)O-alkyl;
- 15

- when the heterocycle group is substituted, it is substituted either on a ring carbon atom or on a ring hetero atom, and when it is substituted on a ring carbon atom, it is substituted with 1 to 4 substituents independently selected from oxo (=O), halogen, cyano, alkyl, cycloalkyl, perhaloalkyl, -OR^{7a}, -C(=O)NR^{8a}R^{8b}, -C(=O)OH, -C(=O)O-alkyl, -N(H)C(=O)(alkyl), -N(H)R^{8a}, and -N(alkyl)₂; and when the heterocycle group is substituted on a ring nitrogen, it is substituted with substituents independently selected from alkyl, cycloalkyl, aryl, heteroaryl, -SO₂(alkyl), -C(=O)R^{9a}, and -C(=O)O(alkyl);
- 20 when the heterocycle group is substituted on a ring sulfur, it is substituted with 1 or 2 oxo (=O) group(s);


- 25 R^{7a} is selected from hydrogen, alkyl, perhaloalkyl, and cycloalkyl;

R^{8a} and R^{8b} are each independently selected from hydrogen, alkyl, and cycloalkyl; and

R^{9a} is selected from alkyl and cycloalkyl.

The details of one or more embodiments of the invention set forth in below are only illustrative in nature and not intended to limit to the scope of the invention. Other features, objects and advantages of the inventions will be apparent from the description and claims.


- 5 According to one embodiment, the invention provides compounds having the structure of Formula (II), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,



10 wherein,

Ring A, L_1 , Z, R^a , R^b , $R^{2'}$, R, R^{2a} , R^3 , R^{10} and 'n' are as defined herein above.


According to another embodiment, the invention provides compounds having the structure of Formula (III), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,



wherein,

Ring A, L₁, Z, R^a, R^b, R^{2'}, R, R^{2a}, R³, R¹⁰ and 'n' are as defined herein above.

According to one embodiment, the invention provides compounds having the structure of Formula (IV), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,



wherein,

X² is Br or Cl;

L₁, R^a, R^b, R¹, R^{2'}, R², R, R^{2a} and R¹⁰ are as defined herein above.

In accordance with an embodiment of the invention, R^c and R^d are independently selected from substituted or unsubstituted alkyl or R^c and R^d together with the carbon atom to which they are attached form a cyclobutyl ring.


In certain embodiment, R^c and R^d are independently selected from methyl or R^c and R^d together with the carbon atom to which they are attached form a cyclobutyl ring.

In any of above embodiments, R^a and R^b are independently selected from hydrogen, methyl, and cyclopropyl.

In any of above embodiments, R³ is selected from halogen, cyano, -OR⁶, -NR⁷R⁸, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl.

In certain embodiments, R³ is independently selected from -F, Cl, Br, CN, -NH₂, -NH(CH₃), -NHCH(CH₃)₂, -CH₃, cyclopropyl, -CH(CH₃)₂, -CF₂CH₃, -OCH₃,


8



In any of above embodiments, $R^{2'}$ and R^{2a} are independently selected from hydrogen and methyl.

- In any of above embodiments, R is selected from hydrogen, halogen, $-NR^4R^5$, and substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl and substituted or unsubstituted heteroaryl.

In certain embodiments, R is selected from hydrogen, $-NH_2$, methyl, Chloro, cyclopropyl,



and

In any of above embodiments, R^4 and R^5 are independently selected from hydrogen.

- In any of above embodiments, R^6 is selected from substituted and unsubstituted alkyl.

In certain embodiments, R^6 is selected from methyl.


In any of above embodiments, R^7 and R^8 are independently selected from hydrogen, methyl, $-CH(CH_3)_2$, $-CH_2$ -cyclopropyl, cyclopropyl, and cyclobutyl.

In any of above embodiments, R^{10} is selected from hydrogen, $-F$, and methyl.

- In any of above embodiments, n is selected from 1 to 3.


According to another embodiment, there are provided compounds having the structure of Formula (I) wherein L_1 is selected from $-CH_2-$, $-CH(CH_3)-$, $-NH-$, $-N(CH_3)-$, S, and O;

According to another embodiment, there are provided compounds having the




structure of Formula (I) wherein ring A is


According to another embodiment, there are provided compounds having the



According to another embodiment, there are provided compounds having the





- 5 According to another embodiment, there are provided compounds having the structure of Formula (I), wherein ring A is selected from-



L1 is selected from $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{NH}-$, $-\text{N}(\text{CH}_3)-$, S, and O; R^3 is selected from F, Cl, Br, CN, $-\text{NH}_2$, $-\text{NH}(\text{CH}_3)$, $-\text{NHCH}(\text{CH}_3)_2$, $-\text{CH}_3$, cyclopropyl, $-\text{CH}(\text{CH}_3)_2$, $-\text{CF}_2\text{CH}_3$, $-\text{OCH}_3$, CF_3 ,

10



selected from hydrogen, $-\text{NH}_2$, Cl, $-\text{CH}(\text{CH}_3)_2$, methyl, ethyl, cyclopropyl and  ;

R^a and R^b are independently selected from hydrogen, methyl, and cyclopropyl; $\text{R}^{2'}$ and R^{2a} are independently selected from hydrogen and methyl; R^{10} is selected from hydrogen,

- 15 $-\text{F}$, and methyl.

The examples 1 to 84 given herein are representative compounds, which are only illustrative in nature and are not intended to limit to the scope of the invention.

It should be understood that formula (I) structurally encompasses all tautomers, stereoisomers and isotopes wherever applicable and pharmaceutically acceptable salts that may be contemplated from the chemical structures generally described herein.

20

According to one embodiment, there are provided compounds of formula (I) to (IV) wherein the compound is in the form of the free base or is a pharmaceutically acceptable salt thereof.

In another aspect of the invention, there are provided compounds of formula (I) to (IV) or a pharmaceutically acceptable salt thereof for treating the diseases, disorders, syndromes or conditions associated with PRMT5 enzyme.

In one embodiment of the present invention, there are provided compounds of formula (I) to (IV), or a pharmaceutically acceptable salt thereof for treating diseases, disorders, syndromes or conditions by inhibition of PRMT5 enzyme.

In another aspect of the invention, there are provided compounds of formula (I) to (IV) or a pharmaceutically acceptable salt thereof for use as a medicament.

In another aspect of the invention, there are provided compounds of formula (I) to (IV) or a pharmaceutically acceptable salt thereof for use in treating the diseases, disorders, syndromes or conditions associated with PRMT5.

In one embodiment of the present invention, there are provided compounds of formula (I) to (IV) or a pharmaceutically acceptable salt thereof for use in treating diseases, disorders, syndromes or conditions by the inhibition of PRMT5.

In another aspect of the invention, there is provided a method of inhibiting PRMT5 by using a compound selected from formula (I) to (IV) or a pharmaceutically acceptable salt thereof.

In another aspect of the invention, there is provided a method of treating diseases, disorders or conditions associated with PRMT5 by using a compound selected from formula (I) to (IV).

In another aspect of the present invention, a method of treating diseases, disorders or conditions is selected from glioblastoma multiforme, prostate cancer, and pancreatic cancer, mantle cell lymphoma, non-Hodgkin's lymphomas and diffuse large B-cell lymphoma, acute myeloid leukemia, acute lymphoblastic leukemia, multiple myeloma, non-small cell lung cancer, small cell lung cancer, breast cancer, triple negative breast

cancer, gastric cancer, colorectal cancer, ovarian cancer, bladder cancer, hepatocellular cancer, melanoma, sarcoma, oropharyngeal squamous cell carcinoma, chronic myelogenous leukemia, epidermal squamous cell carcinoma, nasopharyngeal carcinoma, neuroblastoma, endometrial carcinoma, and cervical cancer by using a compound selected from formula (I) to (IV) is provided.

In another aspect of the invention, there is provided a use of a compound selected from formula (I) to (IV) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for treating, the diseases, disorders or conditions associated with PRMT5.

In another aspect, the invention provides a pharmaceutical composition comprising at least one compound of formula (I) to (IV) or a pharmaceutically acceptable salt thereof and at least one pharmaceutically acceptable excipient.

In another aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of compound of formula (I) to (IV) or a pharmaceutically acceptable salt thereof, for use in treating, the diseases, disorders or conditions associated with PRMT5 by administering to the subject in need thereof.

In another aspect of the present invention, wherein the use of compounds of formula (I) to (IV) or a pharmaceutically acceptable salt thereof for the diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 are selected from the group consisting of glioblastoma multiforme, prostate cancer, and pancreatic cancer, mantle cell lymphoma, non-Hodgkin's lymphomas and diffuse large B-cell lymphoma, acute myeloid leukemia, acute lymphoblastic leukemia, multiple myeloma, non-small cell lung cancer, small cell lung cancer, breast cancer, triple negative breast cancer, gastric cancer, colorectal cancer, ovarian cancer, bladder cancer, hepatocellular cancer, melanoma, sarcoma, oropharyngeal squamous cell carcinoma, chronic myelogenous leukemia, epidermal squamous cell carcinoma, nasopharyngeal carcinoma, neuroblastoma, endometrial carcinoma, and cervical cancer.

In another aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of compound of formula (I) to (IV) or a

pharmaceutically acceptable salt thereof, for treating the diseases, disorders or conditions associated with PRMT5 by administering to the subject in need thereof.

In another embodiment of the invention the compounds, their stereoisomers or pharmaceutically acceptable salts thereof are:

- 5 (1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-1);
- (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-2);
- (1S,2R,5R)-3-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-5-fluoro-7H-
- 10 pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-3);
- (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)thio)methyl)cyclopent-3-ene-1,2-diol (Compound-4);
- (1S,2R,5R)-3-(((2-amino-3-chloroquinolin-7-yl)thio)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-5);
- 15 (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)(methyl)amino)methyl)cyclopent-3-ene-1,2-diol (Compound-6);
- (1S,2R,5R)-3-(1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-7a and 7b);
- (1S,2R,5R)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(1-((2-(methylamino)quinolin-7-yl)oxy)ethyl)cyclopent-3-ene-1,2-diol (Compound-8a and 8b);
- 20 (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-9);
- (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2-(methylamino)quinolin-7-yl)ethyl)cyclopent-3-ene-1,2-diol (Compound-10);
- 25 (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)cyclopent-3-ene-1,2-diol (Compound-11);
- (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-4-methylcyclopent-3-ene-1,2-diol (Compound-12);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-13);
- 30

- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol hydrochloride (Compound-14);
(1S,2R,5R)-3-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol (Compound-15);
5 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-16);
(1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)amino)methyl)cyclopent-3-ene-1,2-diol (Compound-17);
(1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-18);
10 (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-(methylamino)quinolin-7-yl)oxy)methyl)cyclopent-3-ene-1,2-diol (Compound-19);
(1S,2R,5R)-3-(1-((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-20a and 20b);
15 (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-21);
(1S,2R,5R)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2-(cyclobutylamino)quinolin-7-yl)ethyl)cyclopent-3-ene-1,2-diol (Compound-22);
(1S,2R,5R)-3-(2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-23);
20 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-24);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-25);
25 (1S,2R,5R)-3-(2-(2-amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-26);
(1S,2R,5R)-3-(2-(2-amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-27);
(1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2'-aminospiro[cyclobutane-1,3'-indol]-6'-yl)ethyl)cyclopent-3-ene-1,2-diol (Compound-28);
30

- (1S,2R,5R)-3-(2-(2-amino-3,5-dichloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-29);
(1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(2-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-30);
5 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-31);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-32);
10 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-33a and 33b);
(1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-34a and 34b);
(1S,2R,5R)-3-(1-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-35a and 35b);
15 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-2-methyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-36a and 36b);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-37);
20 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-38);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-2-methyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-39);
25 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-ethylcyclopent-3-ene-1,2-diol (Compound-40);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-ethyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-41);
(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-42);
30 (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-43);

- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-44);
- (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoro quinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-45a and 45b);
- 5 (1S,2R,5R)-3-(2-(2-Amino-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-46);
- (1S,2R,5R)-3-(2-(2-amino-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-47);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-48);
- 10 (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(methylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0] hexane-2,3-diol (Compound-49);
- (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(isopropylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol (Compound-50);
- 15 (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(cyclobutylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol (Compound-51);
- (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-((cyclopropylmethyl)amino) quinolin-7-yl) ethyl)bicyclo[3.1.0]hexane-2,3-diol (Compound-52);
- 20 (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-amino-8-fluoroquinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol (Compound-53);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol (Compound-54);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-isopropyl quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-55);
- 25 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(1,1-difluoroethyl) quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d] pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-56);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-cyclopropylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol(Compound-57);
- 30 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methoxyquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-58);

- 2-amino-7-(2-((1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,3-dihydroxybicyclo [3.1.0]hexan-1-yl)ethyl)quinoline-3-carbonitrile(Compound-59);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-60);
5 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-61);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol(Compound-62);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-63);
10 (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-64);
(1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl) bicyclo[3.1.0]hexane-2,3-diol (Compound-65);
15 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol(Compound-66);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-6-methyl-7H-pyrrolo[2,3-d] pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol
20 (Compound-67);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-6-methyl-7H-pyrrolo[2,3-d] pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol
(Compound-68);
(1R,2R,3S,4R,5S)-1-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol (Compound-69);
25 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo [3.1.0]hexane-2,3-diol (Compound-70);
(1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2'-aminospiro[cyclobutane-1,3'-indol]-6'-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol
30 (Compound-71);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-72);

- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-73);
- (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinolin-7-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol (Compound-74);
- 5 (1R,2R,3S,4R,5S)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinazolin-7-yl)ethyl)bicyclo[3.1.0] hexane-2,3-diol (Compound-75);
- (1S,2R,3S,4R,5S)-1-((S)-1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-76a and 76b);
- 10 (1S,2R,3S,4R,5S)-1-((S)-2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)-1-cyclopropylethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-77a and 77b);
- (1S,2R,3S,4R,5S)-1-(1-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-
- 15 78a and 78b);
- (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinolin-7-yl)propyl) bicyclo[3.1.0]hexane-2,3-diol (Compound-79a and 79b);
- (1R,2R,3S,4R,5S)-1-(((2-Amino-3-bromoquinolin-7-yl)oxy)methyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-80);
- 20 (1S,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(((2-aminoquinolin-7-yl)thio)methyl) bicyclo [3.1.0] hexane-2,3-diol (Compound-81);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(4-fluorophenyl)quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol_ (Compound-82);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(pyridin-3-yl)quinolin-7-yl)ethyl)-4-(4-amino-
- 25 7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-83); and
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(3-methyl isoxazol-4-yl)quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol
- (Compound-84).
- 30 In another embodiment of the invention the compounds, their stereoisomer thereof, or a pharmaceutically acceptable salt thereof, are selected from:

- (1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-1);
- (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-2);
- 5 (1S,2R,5R)-3-(1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-7a and 7b);
- (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-9);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-13);
- 10 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol hydrochloride (Compound-14);
- (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-18);
- 15 (1S,2R,5R)-3-(1-((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-20a and 20b);
- (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-21);
- (1S,2R,5R)-3-(2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-23);
- 20 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-24);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-25);
- 25 (1S,2R,5R)-3-(2-(2-amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-26);
- (1S,2R,5R)-3-(2-(2-amino-3,5-dichloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-29);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-31);
- 30

- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-32);
- 5 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-33a and 33b);
- (1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-34a and 34b);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-37);
- 10 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-38);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-ethylcyclopent-3-ene-1,2-diol (Compound-40);
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-ethyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-41);
- 15 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-42);
- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-43);
- 20 (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-44);
- (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoro quinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-45a and 45b);
- (1S,2R,5R)-3-(2-(2-Amino-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-46);
- 25 (1S,2R,5R)-3-(2-(2-amino-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-47);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-48);
- 30 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol (Compound-54);

(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-61);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol(Compound-62);
5 (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-64);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-72); and
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-73).
10

In another embodiment of the invention the compounds, their stereoisomer thereof, or a pharmaceutically acceptable salt thereof, are selected from:

(1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-13);
15 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-24);
(1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol(Compound-33a and 33b);
20 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-37);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-48);
(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol(Compound-62);
25 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-73);
(1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-1);
30 (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-25);

- (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-18);
- (1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-34a and 34b);
- 5 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol (Compound-38);
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol(Compound-61);
- 10 (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-43);
- (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoro quinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-45a);
- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-44);
- 15 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-72); and
- (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-64).

20 Detailed description of the invention

Definitions and Abbreviations

Unless otherwise stated, the following terms used in the specification and claims have the meanings given below.

- 25 For purposes of interpreting the specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa.

The terms "halogen" or "halo" means fluorine, chlorine, bromine, or iodine.

The term "alkyl" refers to an alkane derived hydrocarbon radical that includes solely carbon and hydrogen atoms in the backbone, contains no unsaturation, has from one to six carbon atoms, and is attached to the remainder of the molecule by a single bond, for example (C₁-C₆)alkyl or (C₁-C₄)alkyl, representative groups include *e.g.*,
5 methyl, ethyl, n-propyl, 1-methylethyl (isopropyl), n-butyl, n-pentyl and the like. Unless set forth or recited to the contrary, all alkyl groups described or claimed herein may be straight chain or branched.

The term "alkenyl" refers to a hydrocarbon radical containing from 2 to 10 carbon atoms and including at least one carbon-carbon double bond. Non-limiting Examples of
10 alkenyl groups include, for example (C₂-C₆)alkenyl, (C₂-C₄)alkenyl, ethenyl, 1-propenyl, 2-propenyl (allyl), *iso*-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like. Unless set forth or recited to the contrary, all alkenyl groups described or claimed herein may be straight chain or branched.

The term "alkynyl" refers to a hydrocarbon radical containing 2 to 10 carbon
15 atoms and including at least one carbon-carbon triple bond. Non-limiting Examples of alkynyl groups include, for example (C₂-C₆)alkynyl, (C₂-C₄)alkynyl, ethynyl, propynyl, butynyl and the like. Unless set forth or recited to the contrary, all alkynyl groups described or claimed herein may be straight chain or branched.

The term "haloalkyl" refers to an alkyl group as defined above that is substituted
20 by one or more halogen atoms as defined above. For example (C₁-C₆)haloalkyl or (C₁-C₄)haloalkyl. Suitably, the haloalkyl may be monohaloalkyl, dihaloalkyl or polyhaloalkyl including perhaloalkyl. A monohaloalkyl can have one iodine, bromine, chlorine or fluorine atom. Dihalalkyl and polyhaloalkyl groups can be substituted with two or more of the same halogen atoms or a combination of different halogen atoms. Suitably, a
25 polyhaloalkyl is substituted with up to 12 halogen atoms. Non-limiting Examples of a haloalkyl include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl and the like. A perhaloalkyl refers to an alkyl having all hydrogen atoms

replaced with halogen atoms. Unless set forth or recited to the contrary, all haloalkyl groups described or claimed herein may be straight chain or branched.

The term "alkoxy" denotes an alkyl group attached via an oxygen linkage to the rest of the molecule. Representative examples of such groups are -OCH₃ and -OC₂H₅.

- 5 Unless set forth or recited to the contrary, all alkoxy groups described or claimed herein may be straight chain or branched.

The term "alkoxyalkyl" refers to an alkoxy group as defined above directly bonded to an alkyl group as defined above, *e.g.*, -CH₂-O-CH₃, -CH₂-O-CH₂CH₃, -CH₂CH₂-O-CH₃ and the like.

- 10 The term "cycloalkyl" refers to a non-aromatic mono or multicyclic ring system having 3 to 12 carbon atoms, such as (C₃-C₁₀)cycloalkyl, (C₃-C₆)cycloalkyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like. Examples of multicyclic cycloalkyl groups include, but are not limited to, perhydronaphthyl, adamantyl and norbornyl groups, bridged cyclic groups or spirobicyclic groups, *e.g.*, spiro(4,4)non-2-yl and the like.
- 15

The term "aryl" refers to an aromatic radical having 6- to 14- carbon atoms, including monocyclic, bicyclic and tricyclic aromatic systems, such as phenyl, naphthyl, tetrahydronaphthyl, indanyl, and biphenyl and the like.

- 20 The term "heterocyclic ring" or "heterocyclyl ring" or "heterocyclyl", unless otherwise specified, refers to substituted or unsubstituted non-aromatic 3- to 15-membered ring which consists of carbon atoms and with one or more heteroatom(s) independently selected from N, O or S. The heterocyclic ring may be a mono-, bi- or tricyclic ring system, which may include fused, bridged or spiro ring systems and the nitrogen, carbon, oxygen or sulfur atoms in the heterocyclic ring may be optionally
- 25 oxidized to various oxidation states. In addition, the nitrogen atom may be optionally quaternized, the heterocyclic ring or heterocyclyl may optionally contain one or more olefinic bond(S), and one or two carbon atoms(S) in the heterocyclic ring or heterocyclyl may be interrupted with -CF₂-, -C(O)-, -S(O)-, S(O)₂ etc. In addition heterocyclic ring may also be fused with aromatic ring. Non-limiting Examples of heterocyclic rings

include azetidiny, benzopyranyl, chromanyl, decahydroisoquinolyl, indoliny, isoindoliny, isochromanyl, isothiazolidiny, isoxazolidiny, morpholiny, oxazoliny, oxazolidiny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolidiny, 2-oxoazepiny, octahydroindolyl, octahydroisoindolyl, perhydroazepiny, piperaziny, 4-piperidony, 5 pyrrolidiny, piperidiny, phenothiaziny, phenoxaziny, quinuclidiny, tetrahydroisoquinolyl, tetrahydrofuryl, tetrahydropyranyl, thiazoliny, thiazolidiny, thiamorpholiny, thiamorpholinylsulfoxide, thiamorpholinylsulfoneindoline, benzodioxole, tetrahydroquinoline, tetrahydrobenzopyran and the like. The heterocyclic ring may be attached by any atom of the heterocyclic ring that results in the creation of a 10 stable structure.

The term "heteroaryl" unless otherwise specified, refers to a substituted or unsubstituted 5- to 14- membered aromatic heterocyclic ring with one or more heteroatom(S) independently selected from N, O or S. The heteroaryl may be a mono-, bi- or tricyclic ring system. The heteroaryl ring may be attached by any atom of the 15 heteroaryl ring that results in the creation of a stable structure. Non-limiting Examples of a heteroaryl ring include oxazolyl, isoxazolyl, imidazolyl, furyl, indolyl, isoindolyl, pyrrolyl, triazolyl, triazinyl, tetrazolyl, thienyl, thiazolyl, isothiazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, benzothienyl, carbazolyl, quinoliny, isoquinoliny, quinazoliny, 20 cinnoliny, naphthyridiny, pteridiny, puriny, quinoxaliny, quinolyl, isoquinolyl, thiadiazolyl, indoliziny, acridiny, phenaziny, phthalaziny and the like.

The compounds of the present invention may have one or more chiral centers. The absolute stereochemistry at each chiral center may be 'R' or 'S'. The compounds of the invention include all diastereomers and enantiomers and mixtures thereof. Unless 25 specifically mentioned otherwise, reference to one stereoisomer applies to any of the possible stereoisomers. Whenever the stereoisomeric composition is unspecified, it is to be understood that all possible stereoisomers are included.

The term "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures which are not 30 interchangeable. The three-dimensional structures are called configurations. As used

herein, the term "enantiomer" refers to two stereoisomers whose molecules are non-superimposable mirror images of one another. The term "chiral center" refers to a carbon atom to which four different groups are attached. As used herein, the term "diastereomers" refers to stereoisomers which are not enantiomers. The terms "racemate" or "racemic mixture" refer to a mixture of equal parts of enantiomers.

A "tautomer" refers to a compound that undergoes rapid proton shifts from one atom of the compound to another atom of the compound. Some of the compounds described herein may exist as tautomers with different points of attachment of hydrogen. The individual tautomers as well as mixture thereof are encompassed with compounds of formula (I).

The term "treating" or "treatment" of a state, disorder or condition includes: (a) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a subject that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition; (b) inhibiting the state, disorder or condition, i.e., arresting or reducing the development of the disease or at least one clinical or subclinical symptom thereof; c) lessening the disease, disorder or condition or at least one of its clinical or subclinical symptoms or (d) relieving the disease, i.e., causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

The term "inhibitor" refers to a molecule that binds to an enzyme to inhibit the activity of the said enzyme either partially or completely.

The term "subject" includes mammals (especially humans) and other animals, such as domestic animals (e.g., household pets including cats and dogs) and non-domestic animals (such as wildlife).

A "therapeutically effective amount" means the amount of a compound that, when administered to a subject for treating a disease, disorder or condition, is sufficient to cause the effect in the subject, which is the purpose of the administration. The "therapeutically

effective amount" will vary depending on the compound, the disease and its severity and the age, weight, physical condition and responsiveness of the subject to be treated.

Pharmaceutically Acceptable Salts

The compounds of the invention may form salts with acid or base. The
5 compounds of invention may be sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Non-limiting Examples of pharmaceutically acceptable salts are inorganic, organic acid addition salts formed by addition of acids including hydrochloride salts. Non-limiting Examples of pharmaceutically acceptable salts are inorganic, organic
10 base addition salts formed by addition of bases. The compounds of the invention may also form salts with amino acids. Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting sufficiently basic compound such as an amine with a suitable acid.

Screening of the compounds of invention for PRMT5 inhibitory activity can be
15 achieved by using various in vitro and in vivo protocols mentioned herein below or methods known in the art.

Pharmaceutical Compositions

The invention relates to pharmaceutical compositions containing the compounds of the formula (I), or pharmaceutically acceptable salts thereof disclosed herein. In
20 particular, pharmaceutical compositions containing a therapeutically effective amount of at least one compound of formula (I) described herein and at least one pharmaceutically acceptable excipient (such as a carrier or diluent). Preferably, the contemplated pharmaceutical compositions include the compound(s) described herein in an amount sufficient to inhibit PRMT5 to treat the diseases described herein when administered to a
25 subject.

The subjects contemplated include, for example, a living cell and a mammal, including human. The compound of the invention may be associated with a pharmaceutically acceptable excipient (such as a carrier or a diluent) or be diluted by a carrier, or enclosed within a carrier which can be in the form of a capsule, sachet, paper or

other container. The pharmaceutically acceptable excipient includes pharmaceutical agent that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity.

Examples of suitable carriers or excipients include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, peanut oil, olive oil, gelatin, lactose, terra alba, sucrose, dextrin, magnesium carbonate, sugar, cyclodextrin, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, salicylic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerytritol fatty acid esters, polyoxyethylene, hydroxymethylcellulose and polyvinylpyrrolidone.

The pharmaceutical composition may also include one or more pharmaceutically acceptable auxiliary agents, wetting agents, emulsifying agents, suspending agents, preserving agents, salts for influencing osmotic pressure, buffers, sweetening agents, flavoring agents, colorants, or any combination of the foregoing. The pharmaceutical composition of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the subject by employing procedures known in the art.

The pharmaceutical compositions described herein may be prepared by conventional techniques known in the art. For example, the active compound can be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier, which may be in the form of an ampoule, capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material that acts as a vehicle, excipient, or medium for the active compound. The active compound can be adsorbed on a granular solid container, for Example, in a sachet.

The pharmaceutical compositions may be in conventional forms, for example, capsules, tablets, caplets, orally disintegrating tablets, aerosols, solutions, suspensions or products for topical application.

The route of administration may be any route which effectively transports the active compound of the invention to the appropriate or desired site of action. Suitable

routes of administration include, but are not limited to, oral, oral inhalation, nasal, pulmonary, buccal, subdermal, intradermal, transdermal, parenteral, rectal, depot, subcutaneous, intravenous, intraurethral, intramuscular, intranasal, ophthalmic (such as with an ophthalmic solution) or topical (such as with a topical ointment).

5 Solid oral formulations include, but are not limited to, tablets, caplets, capsules (soft or hard gelatin), orally disintegrating tablets, dragees (containing the active ingredient in powder or pellet form), troches and lozenges. Tablets, dragees, or capsules having talc and/or a carbohydrate carrier or binder or the like are particularly suitable for oral application. Liquid formulations include, but are not limited to, syrups, emulsions, 10 suspensions, solutions, soft gelatin and sterile injectable liquids, such as aqueous or non-aqueous liquid suspensions or solutions. For parenteral application, particularly suitable are injectable solutions or suspensions, preferably aqueous solutions with the active compound dissolved in polyhydroxylated castor oil.

 The pharmaceutical preparation is preferably in unit dosage form. In such form 15 the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as pocketed tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, caplet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged 20 form.

 For administration to subject patients, the total daily dose of the compounds of the invention depends, of course, on the mode of administration. For example, oral administration may require a higher total daily dose, than an intravenous (direct into blood). The quantity of active component in a unit dose preparation may be varied or 25 adjusted from 0.1 mg to 1000 mg by oral administration and 1 μ g to 5000 μ g by inhalation according to the potency of the active component or mode of administration.

 Those skilled in the relevant art can determine suitable doses of the compounds for use in treating the diseases and disorders described herein. Therapeutic doses are generally identified through a dose ranging study in subject based on preliminary 30 evidence derived from the animal studies. Doses must be sufficient to result in a desired

therapeutic benefit without causing unwanted side effects for the patient. For example, the daily dosage of the PRMT5 inhibitor can range from about 0.1 to about 30.0 mg/kg by oral administration. Mode of administration, dosage forms, suitable pharmaceutical excipients, diluents or carriers can also be well used and adjusted by those skilled in the art. All changes and modifications envisioned are within the scope of the invention.

Methods of Treatment

The invention provides compound of formula (I) and pharmaceutical compositions thereof as protein arginine methyl transferase-5 (PRMT5) inhibitors for treating the diseases, disorders or conditions associated with overexpression of PRMT5.

10 The invention further provides a method of treating diseases, disorders or conditions associated with overexpression of PRMT5 in a subject in need thereof by administering to the subject a therapeutically effective amount of a compound or a pharmaceutical composition of the invention.

In another aspect, the invention relates to a method of treating diseases, disorders or conditions associated with the overexpression of PRMT5. In this method, a subject in need of such treatment is administered a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as described herein.

In one embodiment of the present invention, the diseases, disorders, or conditions associated with the overexpression of PRMT5 are cancer.

20 In another embodiment, the invention provides a method of treating cancers, particularly, glioblastoma multiforme, prostate cancer, pancreatic cancer, mantle cell lymphoma, non-Hodgkin's lymphomas and diffuse large B-cell lymphoma, acute myeloid leukemia, acute lymphoblastic leukemia, multiple myeloma, non-small cell lung cancer, small cell lung cancer, breast cancer, triple negative breast cancer, gastric cancer, colorectal cancer, ovarian cancer, bladder cancer, hepatocellular cancer, melanoma, sarcoma, oropharyngeal squamous cell carcinoma, chronic myelogenous leukemia, epidermal squamous cell carcinoma, nasopharyngeal carcinoma, neuroblastoma, endometrial carcinoma, and cervical cancer.


It is to be understood that the invention encompasses the compounds of formula (I) or pharmaceutically acceptable salts thereof for use in the treatment of a disease or disorder mentioned herein.

It is to be understood that the invention encompasses the compounds of formula (I) or pharmaceutically acceptable salts thereof in the manufacture of a medicament for treating a disease or disorder mentioned herein.

General Methods of Preparation

The compound of formula described herein may be prepared by techniques known in the art. In addition, the compound of formula described herein may be prepared by following the reaction sequence as depicted in Scheme-1 to Scheme-27. Further, in the following schemes, where specific bases, acids, reagents, solvents, coupling agents, etc., are mentioned, it is understood that other bases, acids, reagents, solvents, coupling agents etc., known in the art may also be used and are therefore included within the scope of the present invention. Variations in reaction conditions, for example, temperature and/or duration of the reaction, which may be used as known in the art, are also within the scope of the present invention. All the isomers of the compound of formula in described in these schemes, unless otherwise specified, are also encompassed within the scope of this invention.


Scheme-1:



Scheme-1 illustrates the synthesis of compound of formula 10 (when Q¹ is N, CH or CX,

where X can be Cl or Br). Compound of formula 1 (where PG = Protecting group), is prepared by following the procedure reported in Kenneth A. Jacobson et.al; Purinergic Signalling (2015) 11:371–387. Mitsunobu reaction of compound of formula 1 with compound of formula 2 (X = -Cl, -Br) using various azo dicarboxylate reagents such as but not limited to DEAD or DIAD in presence of phosphine such as but not limited to PPh₃ to form the compound of formula 3. Compound of formula 4 is formed upon treatment of compound of formula 3 with fluoride ions such as but not limited to TBAF. Oxidation of compound of formula 4 with oxidising agents such as but not limited to Dess-Martin periodinane can furnish the compound of formula 5. Reagents such as but not limited to methyltriphenylphosphonium bromide in presence of a base such as but not limited to KO^tBu, NaO^tBu, LiHMDS, NaHMDS, or KHMDS when treated with compound of formula 5 affords compound of formula 6. Compound of formula 6a (where R⁴ and R⁵ are defined herein above) upon treatment with compound of formula 6 affords compound of formula 7. Compound of formula 9 can be achieved by hydroboration of compound of formula 7 with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to K₃PO₄ or Cs₂CO₃, in presence of Pd catalyst such as but not limited to Pd(dppf)Cl₂ or Pd(PPh₃)₂Cl₂ and compound of formula 8 (Y = -Br, -I, which can be prepared by following the procedure reported J. Med. Chem., 2017, 60 (9), 3958-3978). Acetonide deprotection of compound of formula 9 with acids such as but not limited to HCl or TFA affords compound of formula 10.


Scheme-2:



Scheme-2 illustrates the synthesis of compound of formula 16. This can be achieved by Mitsunobu reaction of compound of formula 1a, which is prepared by following the procedure reported in Kenneth A. Jacobson et.al; Purinergic Signalling (2015) 11:371–387,

with compound of formula **2** using various azo dicarboxylate reagents such as but not limited to DIAD in presence of phosphine such as but not limited to PPh_3 to form the compound of formula **11**, which can be further converted to compound of formula **12** upon treatment with fluoride ions such as but not limited to TBAF. Mitsunobu reaction of compound of formula **12** with compound of formula **13** (commercially available or synthesized as per known literature, where $\text{Q}^1 = \text{C}$ or N , R^3 , R^7 and R^8 are defined herein above) using various azo dicarboxylate reagents such as but not limited to DIAD in presence of phosphine such as but not limited to PPh_3 affords the compound of formula **14**. Compound of formula **6a** upon treatment with compound of formula **14** affords compound of formula **15**. Acetonide deprotection of compound of formula **15** with acids such as but not limited to HCl or TFA affords compound of formula **16**.


Scheme-3:



Scheme-3 illustrates the synthesis of compound of formula **21**. Compound of formula **17** is formed upon treatment of compound of formula **12** with oxidising agents such as but not limited to Dess-Martin periodinane. Reagents such as but not limited to methyltriphenylphosphonium bromide in presence of a base such as but not limited to KO^tBu , NaO^tBu , LiHMDS , NaHMDS , or KHMDS when treated with compound of formula **17** affords compound of formula **18**. Compound of formula **19** can be synthesized by hydroboration of compound of formula **18** with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to tripotassium phosphate or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $\text{Pd}(\text{dppf})\text{Cl}_2$ or $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and compound of formula **8** ($\text{Y} = -\text{Br}$, $-\text{I}$), which was synthesized by following the procedure reported in WO2012002577 A1, followed by N-oxide formation, chlorination with phosphoroxychloride and nucleophilic substitution with PMB-NH_2 or J .

Med. Chem, 2017, 60 (9), 3958-3978). Compound of formula **6a** (where R^4 and R^5 are defined herein above) upon treatment with compound of formula **19** affords compound of formula **20**. Acetonide deprotection of compound of formula **20** with acids such as but not limited to HCl or TFA affords compound of formula **21**.

5 **Scheme-4:**




Scheme-4 illustrates the synthesis of compound of formula **27**. Compound of formula **22** is formed upon treatment of compound of formula **5** with Grignard reagent such as but not limited to methylmagnesium bromide, ethylmagnesium bromide, cyclopropylmagnesium bromide etc. Compound of formula **22** on oxidation with oxidising agents such as but not limited to Dess-Martin periodinane gives compound of formula **23**. Reagents such as but not limited to methyltriphenylphosphonium bromide in presence of base such as but not limited to KO^tBu , NaO^tBu , LiHMDS, NaHMDS, or KHMDS when treated with compound of formula **23** affords compound of formula **24**. Compound of formula **25** can be achieved by hydroboration of compound of formula **24** with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to tripotassium phosphate or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $\text{Pd}(\text{dppf})\text{Cl}_2$ or $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and compound of formula **8** ($Y = -\text{Br}, -\text{I}$), which was synthesized by following the procedure reported in WO2012002577 A1, followed by N-oxide formation, chlorination with Phosphoroxychloride, and nucleophilic substitution with PMB-NH_2 or J.Med.Chem, 2017, 60 (9), 3958-3978). Compound of formula **6a**

(where R_4 and R_5 are defined herein above) upon treatment with compound of formula **25** affords compound of formula **26**. Acetonide deprotection of compound of formula **26** with acids such as but not limited to HCl or TFA affords compound of formula **27**.

Scheme-5:


5



Scheme-5 illustrates the synthesis of compound of formula **32**. Compound of formula **28** is formed upon treatment of compound of formula **5** with Wittig reagents such as but not limited to (1-iodoethyl) triphenylphosphonium bromide. Compound of formula **29** can be achieved by Suzuki coupling of compound of formula **28** with suitable inorganic base such as but not limited to tripotassium phosphate or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}(\text{dppf})\text{Cl}_2$ or $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and compound of formula **8a** ($Y = -\text{Br}, -\text{I}$). Compound of formula **6a** (where R_4 and R_5 are defined herein above) upon treatment with compound of formula **29** affords compound of formula **30**. Hydrogenation of compound of formula **30** affords compound of formula **31**. Acetonide deprotection of compound of formula **31** with acids such as but not limited to HCl or TFA affords compound of formula **32**.

Scheme-6:


35



Compound of formula-33 (can be synthesized by following the protocol as mentioned in Med. Chem. Comm.; vol. 4; nb. 5; (2013); p. 822 - 826) upon treatment with iodine in presence of pyridine affords Compound of formula-34, which can be reduced by cerium chloride and sodium borohydride to get compound of formula-35. Mitsunobu reaction using compound of formula-2 with compound of formula-35 provides the compound of formula-36. Halogen of compound of formula-36 on substitution with PMB amine, followed by Suzuki reaction affords the compound of formula-38. TBDMS deprotection of compound of formula-38 followed by tosylation of compound of formula-39 provides compound of formula-40. The tosyl of compound of formula-40 is replaced with compound of formula-13 gives the compound of formula-41, which is further deprotected in acidic condition to get the final compound of formula -42.


Scheme-7:

36




Reaction of compound of formula-43 [can be prepared by method reported in *J. Org. Chem.* 2014, 79, 8059–8066] with methyl magnesium bromide gives compound of formula-44, with only one stereoisomer with good yield. This when subjected to acetone shuffling affords compound of formula-46. Mitsunobu reaction using compound of formula 2 with compound of formula-46 yields the compound of formula-47, which can be converted to compound of formula-48 on treating with TBAF. Compound of formula-48 can be oxidized with DMP to afford Compound of formula-49, which can undergo Wittig reaction to give compound of formula-50. Compound of formula 51 can be achieved by hydroboration of compound of formula 50 with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to K_3PO_4 or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $Pd(dppf)Cl_2$ or $Pd(PPh_3)_2Cl_2$ and compound of formula 8. Compound of formula-51 on treating with aq. ammonia followed by treatment with trifluoroacetic acid affords compound of formula-53.

15 Scheme-8:




Compound of formula-48 (compound of formula 48a can be synthesized from 48 by oxidation and Grignard reaction) when condensed with compound of formula-13 using Mitsunobu reaction affords compound of formula-54, which can be reacted with ammonia followed by treatment with trifluoroacetic acid to provide a compound of formula 56.

5 **Scheme-9:**



6-Halo-oxindole on treatment with a base, and methyl iodide or diiodopropane or appropriately substituted dihalide provides compound of formula-57. Compound of formula 57 when treated with Lawesson's reagent in hydrocarbon solvent such as but not limited to toluene at 100°C for 3h gives compound of formula-58, which is then treated with sodium hydride followed by methyl iodide in THF to yield compound of formula 59. Compound of formula 59 on treating with 7N ammonia in methanol at 100°C for 16h provides compound of formula-60.


Scheme-10:



Hydroboration of compound of formula-18 with 9-BBN followed Suzuki coupling with compound of formula-59 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h provide compound of formula-61, which is then treated with compound of formula-6a followed by TFA to give compound of formula 63.


20 **Scheme-11:**

38



Hydroboration of compound of formula-7 with 9-BBN followed Suzuki coupling with compound of formula-59 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-64, which is then treated with compound of formula-6a followed by TFA to afford compound of formula 66.


Scheme-12:



Hydroboration of compound of formula-18 with 9-BBN followed Suzuki coupling with compound of formula-67 in presence of Pd-118 in THF/H₂O at 50°C for 5-16h affords compound of formula-68, which is when treated with compound of formula-6a followed by TFA affords compound of formula-70. Hydroboration of compound of formula-7 with 9-BBN followed Suzuki coupling with compound of formula-67 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-71, which is when treated with TFA affords compound of formula-72.


Scheme-13:

39



Hydroboration of compound of formula-6 with 9-BBN followed by Suzuki coupling of compound of formula 8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula 73, which is when treated with compound of formula-6a followed by treatment with TFA affords compound of formula 75.


Scheme-14:




Compound of formula-17 when treated with methylmagnesium bromide in THF gives compound of formula-76, which can be oxidized by DMP in methylene chloride at 0°C to afford compound of formula-77. Addition of methylmagnesium bromide on compound of formula-77 provides compound of formula-78. Dehydration of compound of formula-78 with Martin Sulfurane gives compound of formula-79, which then treated with 9-BBN followed by Suzuki coupling with compound of formula-8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-80. Treating compound of formula-80 with compound of formula 6a followed by treatment with TFA affords compound of formula-82.

Scheme-15:


40




Compound of formula-11 on treatment with trimethylboroxine in presence of tripotassium phosphate, and Pd-118 or PdCl₂dppf gives mixture of compound of formula-83 and compound of formula-84. The PG of compound of formula-83 can be deprotected to afford compound of formula-85, which on oxidation with DMP gives compound of formula-86. Wittig reaction of compound of formula-86 provides compound of formula-87. Hydroboration of compound of formula-87 with 9-BBN followed by Suzuki coupling with compound of formula-8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-88, which when treated with TFA affords compound of formula-89.

Scheme-16:

The protecting group of compound of formula-84 can be deprotected to afford compound of formula-90, which on oxidation with DMP gives compound of formula-91. Wittig reaction of compound of formula-91 provides compound of formula-92. Hydroboration of compound of formula-92 with 9-BBN followed by Suzuki coupling with compound of formula-8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-93, which is then treated with TFA to afford compound of formula 94.


Scheme-17:

10 Reductive amination of compound of formula-17 with compound of formula-95 affords compound of formula-96, which is then treated with compound of formula-6a followed by treatment with TFA to afford compound of formula-98.

Scheme-18:

15 Compound of formula-12 is treated with TsCl/ MsCl in presence of a base to give a compound of formula -99, which is then reacted with compound of formula-100 to yield compound of formula -101. Reaction of compound of formula-101 with compound of formula-6a followed by treatment with TFA affords compound of formula-103.


Scheme-19:



Compound of formula-99 is reacted with compound of formula-104 in presence of a base such as but not limited to cesium carbonate to afford a compound of formula-105.

- 5 Substitution reaction of compound of formula-105 with compound of formula-6a followed by treatment with TFA affords compound of formula-107.


Scheme-20:



- 10 The compound of formula-109 can be synthesized by Mitsunobu reaction of compound of formula 1a with compound of formula 108 using various azo dicarboxylate reagents such as but not limited to DIAD in presence of phosphine such as but not limited to PPh_3 . Compound of formula-109 can be further converted to compound of formula 110 upon treatment with fluoride ions such as but not limited to TBAF. Oxidation followed by Wittig
- 15 reaction on compound of formula 110 gives compound of formula 112. The aromatic halogen of compound of formula 112 can be converted to alkyl groups such as a methyl group using Pd-H18 or PdCl_2dppf with trimethylboroxine to afford compound of formula 113. Hydroboration of compound of formula-113 with 9-BBN followed by Suzuki coupling

with compound of formula-8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-114, which when treated with TFA or HCl/MeOH affords compound of formula-115.


Scheme-21:



Mitsunobu reaction of compound of formula 4 (can be prepared from compound of formula 1 as described in scheme 1) with compound of formula 13 using various azo dicarboxylate reagents such as but not limited to DIAD in presence of phosphine such as but not limited to PPh₃ provides compound of formula 116. Compound of formula-116 on treating compound of formula 6a followed by treatment with trifluoroacetic acid affords compound of formula-118.

10


Scheme-22:




Scheme-22 illustrates the synthesis of compound of formula 126. Compound of formula 119 (where PG = Protecting group), is prepared by Simmons Smith reaction. Mitsunobu reaction of compound of formula 119 with compound of formula 2 (X = -Cl, -Br) using

15

various azo dicarboxylate reagents such as but not limited to DEAD or DIAD in presence of phosphine such as but not limited to PPh_3 to form the compound of formula 120. Compound of formula 121 is formed upon treatment of compound of formula 120 with fluoride ions such as but not limited to TBAF. Oxidation of compound of formula 121 with oxidising agents such as but not limited to Dess-Martin periodinane can furnish the compound of formula 122. Reagents such as but not limited to methyltriphenylphosphonium bromide in presence of a base such as but not limited to KOtBu, NaOtBu, LiHMDS, NaHMDS, or KHMDS when treated with compound of formula 122 affords compound of formula 123. Amine of formula 6a (where R_4 and R_5 are defined herein above) upon treatment with compound of formula 123 affords compound of formula 124. Compound of formula 125 can be achieved by hydroboration of compound of formula 124 with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to K_3PO_4 or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $\text{Pd}(\text{dppf})\text{Cl}_2$ or $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and compound of formula 8 ($\text{Y} = -\text{Br}, -\text{I}$, which can be prepared by following the procedure reported J. Med. Chem., 2017, 60 (9), 3958-3978). Acetonide deprotection of compound of formula 125 with acids such as but not limited to HCl or TFA affords compound of formula 126.


Scheme-23:

Suzuki coupling of compound of formula 10 with aryl or heteroaryl boronic acid or ester in presence of $\text{Pd}(\text{PPh}_3)_4$ or Pd-118 or PdCl_2dppf in dioxane at 80-100°C for 5-16h affords compound of formula-127.

Scheme-24:


Compound of formula-18 when treated with compound of formula-128 using Suzuki coupling affords compound of formula-129. Compound of formula 130 can be achieved by hydroboration of compound of formula 129 with suitable boranes such as but not limited to 9-BBN followed by addition of inorganic base such as but not limited to K_3PO_4 or Cs_2CO_3 , in presence of Pd catalyst such as but not limited to $Pd(dppf)Cl_2$ or $Pd(PPh_3)_2Cl_2$ and compound of formula 8 ($Y = -Br, -I$, which can be prepared by following the procedure reported J. Med. Chem., 2017, 60 (9), 3958-3978). Acetonide deprotection of compound of formula 130 with acids such as but not limited to HCl or TFA affords compound of formula 131.

10 **Scheme-25:**



Compound of formula-76 when condensed with compound of formula-13 using Mitsunobu reaction affords compound of formula-132, which can be reacted with compound of formula-6a followed by treatment with trifluoroacetic acid to provide a compound of formula-134.

15 **Scheme-26:**




Compound of formula-48a when treated DMP in methylene chloride at $0^\circ C$ to afford compound of formula -135, which when react with methylmagnesium bromide in THF

gives compound of formula-136. Dehydration of compound of formula-136 with Martin Sulfurane gives compound of formula-137, which then treated with 9-BBN followed by Suzuki coupling with compound of formula-8 in presence of Pd-118 or PdCl₂dppf in THF/H₂O at 50-70°C for 5-16h affords compound of formula-138. Treating compound of

5 formula-138 with compound of formula 6a followed by treatment with TFA affords compound of formula-140.

Scheme-27:



Compound of formula-4 is treated with TsCl/ MsCl in presence of a base to give a

10 compound of formula 141. Compound of formula-141 is reacted with compound of formula-104 in presence of a base such as but not limited to cesium carbonate to afford a compound of formula -142. Substitution reaction of compound of formula-142 with compound of formula-6a followed by treatment with TFA affords compound of formula-

15 144.

Abbreviations

The following abbreviations may be used herein:

20

AcOH = Acetic acid

Aq.= aqueous

ca = about or approximately

NH₄Cl = Ammonium chloride

25

9-BBN = 9-Borabicyclononane

BINAP = 2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl

Boc = *tert*-Butoxycarbonyl

t-Bu or *t*Bu = *tert*-Butyl

	Cs_2CO_3 = Cesium Carbonate
	CHCl_3 = Chloroform
	CDCl_3 = Deuterated chloroform
	DAST = Diethylaminosulphur trifluoride
5	dba = Dibenzylideneacetone
	CH_2Cl_2 or DCM = Dichloromethane
	DMP = Dess Martin Periodinane
	DEAD = Diethyl azodicarboxylate
	DIAD = Diisopropyl azodicarboxylate
10	DIPEA = Diisopropylethylamine
	DMAP = 4-Dimethylaminopyridine
	DMF = <i>N,N</i> -Dimethylformamide
	DMSO = Dimethylsulphoxide
	DMSO- d_6 = Deuterated dimethylsulphoxide
15	Et = ethyl
	EtOH = Ethanol
	EtOAc = Ethyl acetate
	g = gram
	H_2O_2 = Hydrogen peroxide
20	H_2SO_4 = Sulphuric acid
	K_2CO_3 = Potassium carbonate
	KOH = Potassium hydroxide
	KO ^t Bu = Potassium tert-butoxide
	K_3PO_4 = Potassium phosphate
25	KHMDS = Potassium bis(trimethylsilyl)amide
	LDA = Lithium diisopropylamide
	LHMDS = Lithium bis(trimethylsilyl)amide
	LCMS = Liquid chromatography mass spectrometry
	<i>m</i> -CPBA = <i>meta</i> -chloroperoxybenzoic acid
30	mg = milligram
	Me = Methyl
	MeOH = Methanol

	MeOD = Deuterated methanol
	MS = Molecular sieves
	MsCl = Methanesulphonyl chloride
	MgSO ₄ = Magnesium sulphate
5	NaH = Sodium hydride
	NaO ^t Bu = Sodium tert-butoxide
	NaHCO ₃ = Sodium bicarbonate
	Na ₂ SO ₄ = Sodium sulphate
	Na ₂ S ₂ O ₃ = Sodium thiosulphate
10	Na ₂ SO ₃ = Sodium sulphite
	NaHMDS = Sodium bis(trimethylsilyl)amide
	NMP = N-Methyl-2-pyrrolidone
	NBS = <i>N</i> -Bromosuccinimide
	NCS = <i>N</i> -Chlorosuccinimide
15	NIS = <i>N</i> -Iodosuccinimide
	NMO = N-Methylmorpholine-N-oxide
	NMR = Nuclear magnetic resonance
	Ph = phenyl
	PDC = Pyridinium dichlorochromate
20	Pd(OAc) ₂ = Palladium acetate
	Pd/C = Palladium on carbon
	Pd-118 = [1,1'-Bis(di-tert-butylphosphino)ferrocene]dichloropalladium(II)
	Pd(PPh ₃) ₄ = Tetrakis(triphenylphosphine)palladium(0)
	POCl ₃ = Phosphorous oxychloride
25	PdCl ₂ (dppf) = [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II)
	Pd(PPh ₃) ₂ Cl ₂ = Bis(triphenylphosphine)palladium(II) dichloride
	PCC = Pyridinium chlorochromate
	PMB = p-Methoxybenzyl
	PTSA = p-Toluenesulphonic acid
30	Rt = Retention time
	rt = room temperature
	Sat.= saturated

SFC = Supercritical fluid chromatography

TLC = Thin layer chromatography

TBAF = Tetrabutylammonium fluoride

TsCl = p-Toluenesulphonyl chloride

5 TBDMS = tert-Butyldimethylsilyl

TBDPS = tert-Butyldiphenylsilyl

Et₃N or NEt₃ or TEA = Triethylamine

TFA = Trifluoroacetic acid

THF = Tetrahydrofuran


10 Ts = p-Toluenesulphonyl

p-TsOH = p-Toluenesulphonic acid

15


INTERMEDIATES

7-Bromoquinolin-2-amine




20 The title compound was prepared by following an analogous reaction protocol as described in Cinelli, Maris A et al, Journal of Medicinal Chemistry, 2017, vol. 60, # 9, p. 3958 - 3978.

7-Bromo-N-(4-methoxybenzyl)quinolin-2-amine



The title compound was prepared by following an analogous reaction protocol as described
25 in Arnould, Jean-Claude et al, WO 2007/141473 A1.


7-Bromo-N-methylquinolin-2-amine




The title compound was prepared by following same reaction protocol as described in
Arnould, Jean-Claude et al, WO 2007/141473 A1.

30 **7-Bromo-N-isopropylquinolin-2-amine**

50




The title compound was prepared by following an analogous reaction protocol as described in Arnould, Jean-Claude et al, WO 2007/141473 A1.

7-Bromo-N-cyclobutylquinolin-2-amine


5

The title compound was prepared by following an analogous reaction protocol as described in Arnould, Jean-Claude et al, WO 2007/141473 A1.

7-Bromo-N-(cyclopropylmethyl)quinolin-2-amine


10

The title compound was prepared by following analogous reaction protocol as described in Arnould, Jean-Claude et al, WO 2007/141473 A1.

7-Bromo-2-chloro-8-fluoroquinoline


The title compound was prepared by following same reaction protocol as described in Aciro, Caroline et al, WO2013/185103 A1.

15

7-Bromo-2-chloro-3-isopropylquinoline

The title compound was prepared by following same reaction protocol as described in Vialard, Jorge Eduardo et al, WO2008/107478 A1; LCMS m/z= 284.1, 286.1 (M⁺, M+2; 100%).


20

2-Amino-4-bromo-6-fluorobenzaldehyde

To a stirred solution of 4-bromo-2-fluoro-6-nitrobenzaldehyde (prepared by following same reaction protocol as described in Li, Liansheng et al, WO 2015/054572 A1; 4.15g,

16.73 mmol) in ethanol (20ml) & acetic acid (20ml) was added iron powder (2.80 g, 50.2 mmol) at 0°C and stirred the reaction mixture for 1h. The reaction mixture was diluted with ethyl acetate (70ml) and netralized with aq. sat. NaHCO₃ (100 ml). The resulting emulsion was filtered through celite. Layers were separated, organic layer was washed with brine (100 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to afford (3.36g, 92%) as a light green solid which was used for next step without purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.10 (s, 1H), 7.78 – 7.54 (m, 2H), 6.84 (t, *J* = 1.5 Hz, 1H), 6.64 (dd, *J* = 11.1, 1.8 Hz, 1H).

1-Chloro-5-iodo-2-methyl-3-nitrobenzene



A solution of iodine (10.43g, 41.1mmol), potassium iodate (1.247g, 5.83mmol) in conc. H₂SO₄ (51.4g, 525mmol) was added to a solution of 1-chloro-2-methyl-3-nitrobenzene (5g, 29.1mmol) in conc. H₂SO₄ (51.4g, 525mmol) at 25°C. The resulting mixture was stirred at 25°C for 6h. The reaction mixture was added slowly onto crushed ice and extracted the product with ethyl acetate (75ml). The organic layer was washed with aq.sat.NaHCO₃ (75ml), aq.sat.Na₂S₂O₃ (75ml) and brine (75ml) successively. Dried the organic layer over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give 9g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with elution of petroleum ether to afford the title compound (8.5g, 98%) as a pale yellow oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.02 (d, *J* = 1.4 Hz, 1H), 7.96 (d, *J* = 1.7 Hz, 1H), 2.50 (s, 3H).


2-(Bromomethyl)-1-chloro-5-iodo-3-nitrobenzene



To a stirred solution of 1-chloro-5-iodo-2-methyl-3-nitrobenzene (27.5g, 92mmol) in CCl₄ (280 ml) was added NBS (19.74g, 111mmol) and benzoyl peroxide (2.99g, 9.24 mmol) at 25°C. The resulting mixture was stirred at 80°C for 15h. The solvent was evaporated *in vacuo* and this residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution of (0 to 1%) of ethyl acetate in petroleum ether


to afford the title compound (12g, 34.5 %) as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.37 – 8.33 (m, 2H), 4.72 (s, 2H).

2-Chloro-4-iodo-6-nitrobenzaldehyde




- 5 To a stirred solution of 2-(bromomethyl)-1-chloro-5-iodo-3-nitrobenzene (12 g, 31.9 mmol) in acetonitrile (150ml) was added 4-methylmorpholine-N-oxide (9.19 g, 78 mmol) and molecular sieves 4A° (30g) at 25°C. The resulting mixture was stirred at 25°C for 1.5h. Water (75ml) was added, pH was adjusted to 6 by adding 1N HCl. Extracted the product with ethyl acetate (75ml). Layers were separated, organic layer was washed with brine
- 10 (20ml) and dried over anhydrous Na_2SO_4 . The organic layer was filtered and concentrated *in vacuo* to give 16.5g of crude compound. This residue was purified by combiflash (R_f 200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 4%) of ethyl acetate in petroleum ether to afford the title compound (7g, 70.5%) as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.22 (s, 1H), 8.46 (s, 2H).

15 **2-Amino-6-chloro-4-iodobenzaldehyde**



- The title compound was prepared by following an analogous reaction protocol as described in the preparation of 2-amino-4-bromo-6-fluorobenzaldehyde. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.26 (s, 1H), 7.80 – 7.55 (m, 2H), 7.23 (d, J = 1.4 Hz, 1H), 6.99 (d, J = 1.5
- 20 Hz, 1H).


1-Fluoro-2-iodo-5-methyl-4-nitrobenzene



- To a stirred solution of 2-fluoro-4-methyl-5-nitroaniline (2.0 g, 11.75 mmol) in conc.HCl (6.15 ml, 73.8 mmol) was added a solution of sodium nitrite (0.884 g, 12.81 mmol) in water (4ml) in a dropwise manner at 0°C. After stirring for 15mins, the mixture was filtered
- 25 through a cotton pad and slowly poured into a stirred solution of potassium iodide (6.83 g, 41.1mmol) in water (25ml) at 0°C. The resulting mixture was stirred at 25°C for 16h. The


reaction mixture was diluted with ethyl acetate (20ml) and washed with 10% aq.NaOH (50ml), aq.sat.NaHCO₃ (50ml) successively. Layers were separated, organic layer was washed with brine (50ml) and was dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 3.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with petroleum ether as a eluent to afford the title compound (1.7g, 51.5%) of the title compound as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.45 (d, *J* = 5.7 Hz, 1H), 7.06 (d, 1H), 2.62 (d, *J* = 0.7 Hz, 3H); GCMS *m/z* = 281.03 (M⁺, 50%).

1-(Bromomethyl)-5-fluoro-4-iodo-2-nitrobenzene




The title compound was prepared by following an analogous reaction protocol as described in the preparation of 2-(bromomethyl)-1-chloro-5-iodo-3-nitrobenzene. ¹H NMR (400 MHz, Chloroform-d) δ 8.53 (d, *J* = 5.6 Hz, 1H), 7.31 (d, *J* = 7.8 Hz, 1H), 4.80 (s, 2H).

5-Fluoro-4-iodo-2-nitrobenzaldehyde




The title compound was prepared by following an analogous reaction protocol as described in the preparation of 2-chloro-4-iodo-6-nitrobenzaldehyde. ¹H NMR (400 MHz, Chloroform-d) δ 10.42 (d, *J* = 2.3 Hz, 1H), 8.62 (d, *J* = 5.1 Hz, 1H), 7.60 (d, *J* = 7.2 Hz, 1H).

2-Amino-5-fluoro-4-iodobenzaldehyde




The title compound was prepared by following an analogous reaction protocol as described in the preparation of 2-amino-4-bromo-6-fluorobenzaldehyde. ¹H NMR (400 MHz, Chloroform-*d*) δ 9.78 (d, *J* = 0.6 Hz, 1H), 7.17 – 7.13 (m, 2H), 5.98 (s, 2H).

7-Bromo-3-cyclopropylquinoline



To a stirred mixture of 2-amino-4-bromobenzaldehyde (2 g, 10.00 mmol) and 2-cyclopropylacetaldehyde (0.841 g, 10.00 mmol) in absolute ethanol (12ml) was added a solution of KOH (0.191 g, 3.40 mmol) in ethanol (12ml) in a dropwise manner under N₂ atmosphere. The resulting mixture was stirred at 95°C for 5h. The volatiles were evaporated
5 *in vacuo* and the residue was dissolved in dichloromethane (60 ml) and washed with water (40 ml). Layers were separated, organic layer was washed with brine (50 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 2.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 6%) of ethyl acetate in
10 petroleum ether to afford 7-bromo-3-cyclopropylquinoline (1g, 40.3%) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (d, *J* = 2.3 Hz, 1H), 8.17 (d, *J* = 2.1 Hz, 1H), 8.00 (d, *J* = 2.4 Hz, 1H), 7.89 – 7.80 (m, 1H), 7.70 (dd, *J* = 8.8, 2.1 Hz, 1H), 2.15 (tt, *J* = 8.1, 5.1 Hz, 1H), 1.14 – 1.03 (m, 2H), 0.95 – 0.81 (m, 2H); LCMS *m/z* = 247.83, 249.83 (M⁺, M+2, 100%).


15 **7-Bromo-3-(1,1-difluoroethyl)quinoline**



A mixture of 1-(7-bromoquinolin-3-yl)ethan-1-one (synthesized by following same reaction protocol as described in Alam, Muzaffar et al, US20120230951 A1; 2.4 g, 9.60 mmol) in diethylaminosulfur trifluoride (2.5 ml, 18.92 mmol) was stirred at 70°C for 16h.
20 The resulting mixture was slowly poured into aq. sat. sodium bicarbonate (50ml) and extracted with dichloromethane (50ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 3.2g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient
25 elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (1.7g, 65.1 %) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.13 (d, *J* = 2.4 Hz, 1H), 8.71 (s, 1H), 8.34 (d, *J* = 2.1 Hz, 1H), 8.11 (d, *J* = 8.7 Hz, 1H), 7.88 (dd, *J* = 8.7, 2.0 Hz, 1H), 2.14 (t, *J* = 19.2 Hz, 3H), LCMS *m/z* = 271.90, 273.90 (M+1; 100%).

7-Bromo-3-chloro-8-fluoroquinoline


55



To a stirred solution of 7-bromo-8-fluoroquinoline (synthesized by following same reaction protocol as described in Ghergurovich, Jonathan Michael et al, WO2013028447 A1, 3.4 g, 15.04 mmol) in DMF (10ml) was added N-chlorosuccinimide (4.02g, 30.1mmol) at 25°C.

- 5 The reaction mixture was stirred at 100°C for 16h. The reaction mixture was cooled to 25°C and poured onto ice cold water (100ml) and stirred for 30 minutes. The precipitate was collected by filtration and washed with water. The precipitate was dried *in vacuo* to afford the title compound (2g, 51%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.90 (d, *J* = 2.3 Hz, 1H), 8.18 (t, *J* = 1.9 Hz, 1H), 7.72 (dd, *J* = 8.9, 6.1 Hz, 1H), 7.48 (dd, *J* = 8.8, 1.5 Hz, 1H); LCMS *m/z* = 261.71 (M⁺; 100%).
- 10





7-Bromo-3-chloro-5-fluoroquinoline




To a stirred solution of 2-amino-4-bromo-6-fluorobenzaldehyde (3.36 g, 15.41 mmol) in toluene (35 ml) was added 2-chloro-1,1-dimethoxyethane (2.304 g, 18.49 mmol) followed by *p*-toluene sulfonic acid monohydrate (2.93 g, 15.41 mmol) at 25°C. The resulting mixture was stirred at 110°C using Dean Stark apparatus for 4h under N₂ atmosphere. The reaction mixture was diluted with ethyl acetate (100ml) and basified with aq.sat.NaHCO₃ (75ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 2%) of ethyl acetate in petroleum ether to afford the title compound (1.64g, 40.9%) as a light green solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.88 (d, *J* = 2.4 Hz, 1H), 8.38 (d, *J* = 2.4 Hz, 1H), 8.15 (s, 1H), 7.45 (ddd, *J* = 9.0, 3.5, 1.8 Hz, 1H); LCMS *m/z* = 261.76 (M+1, 100%).

- 25 Intermediates in table-1 were synthesized by following an analogous reaction protocol as was used for the preparation of 7-bromo-3-chloro-5-fluoroquinoline using the appropriate starting materials.

Table-1:

Intermediate's Structure	Starting materials used	¹ H NMR and LCMS data
 7-Bromo-3-chloro-6-fluoroquinoline	2-Amino-4-bromo-5-fluorobenzaldehyde, which was synthesized as per US2014/200216 A1.	¹ H NMR (400 MHz, Chloroform-d) δ 8.81 (s, 1H), 8.43 – 8.38 (m, 1H), 8.12 – 8.08 (m, 1H), 7.47 (d, J = 8.4 Hz, 1H); LCMS m/z= 261.83 (M+1; 100%).
 3,5-Dichloro-7-iodoquinoline	2-Amino-6-chloro-4-iodobenzaldehyde	¹ H NMR (400 MHz, Chloroform-d) δ 8.85 (d, J = 2.3 Hz, 1H), 8.50 (dd, J = 2.3, 0.9 Hz, 1H), 8.48 – 8.45 (m, 1H), 7.96 (d, J = 1.6 Hz, 1H).
 3,7-Dibromo-5-fluoroquinoline	2-Amino-4-bromo-6-fluorobenzaldehyde	Crude was taken as such for the next step
 3-Bromo-6-fluoro-7-iodoquinoline	2-Amino-5-fluoro-4-iodobenzaldehyde	¹ H NMR (400MHz, Chloroform-d) δ 8.88 (d, J = 2.2 Hz, 1H), 8.61 (d, J = 6.0 Hz, 1H), 8.27 (d, J = 2.3 Hz, 1H), 7.39 (d, J = 7.8 Hz, 1H); LCMS m/z= 351.03, 353.03 (M-1, M+1, 60%).

7-Bromo-3-chloro-5-fluoroquinoline 1-oxide



A mixture of 7-bromo-3-chloro-5-fluoroquinoline (1.64 g, 6.30 mmol) and m-CPBA (2.90 g, 12.59 mmol) in CHCl₃ (30ml) was heated at 50°C for 16h. The reaction mixture was






5 diluted with chloroform (50ml) and washed with aq.sat. Na₂SO₃ (50ml) followed by



aq. sat. NaHCO₃ (50ml). Layers were separated, organic layer was washed with brine (50ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.35g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (542mg, 31.1%) as a light brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.97 (d, *J* = 1.8 Hz, 1H), 8.46 (s, 1H), 8.17 – 8.10 (m, 1H), 8.03 (dd, *J* = 9.5, 1.9 Hz, 1H); LCMS *m/z* = 275.83, 277.83 (M⁺, M+2; 100%).


Intermediates in table-2 were synthesized by following an analogous reaction protocol as was used for the preparation of 7-bromo-3-chloro-5-fluoroquinoline 1-oxide using the appropriate starting materials.

Table-2:

Intermediate's Structure	Starting materials used	¹ H NMR and LCMS data
 7-Bromo-3-methylquinoline 1-oxide	7-Bromo-3-methylquinoline	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.60 (dd, <i>J</i> = 12.2, 1.7 Hz, 2H), 7.97 (d, <i>J</i> = 8.7 Hz, 1H), 7.85 (dd, <i>J</i> = 8.7, 2.0 Hz, 1H), 7.77 (q, <i>J</i> = 1.1 Hz, 1H), 2.40 (d, <i>J</i> = 1.0 Hz, 3H); LCMS <i>m/z</i> = 237.9, 239.9 (M ⁺ , M+2, 100%).
 7-Bromo-3-cyclopropylquinoline 1-oxide	7-Bromo-3-cyclopropylquinoline	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.58 (dd, <i>J</i> = 13.9, 1.7 Hz, 2H), 7.93 (d, <i>J</i> = 8.7 Hz, 1H), 7.84 (dd, <i>J</i> = 8.7, 2.1 Hz, 1H), 7.66 (s, 1H), 2.08 (tt, <i>J</i> = 8.4, 5.0 Hz, 1H), 1.12 – 1.04 (m, 2H), 0.95 – 0.87 (m, 2H); LCMS <i>m/z</i> = 263.83, 265.83 (M ⁺ , M+2, 100%).

 7-Bromo-3-(1,1-difluoroethyl)quinoline 1-oxide	7-Bromo-3-(1,1-difluoroethyl) quinoline	LCMS m/z= 289.96 (M+1; 100%).
 7-Bromo-3-methoxyquinoline 1-oxide	7-Bromo-3-methoxy quinoline, which was synthesized as per Adams, Nicholas David et al, WO2014/008223 A1.	Crude was taken as such for the next step
 7-Bromo-3-chloro-6-fluoroquinoline 1-oxide	7-Bromo-3-chloro-6-fluoroquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.87 (d, J = 1.8 Hz, 1H), 8.75 (d, J = 6.6 Hz, 1H), 8.14 – 8.03 (m, 2H); LCMS m/z= 275.77 (M+1; 60%).
 7-Bromo-3-chloro-8-fluoroquinoline 1-oxide	7-Bromo-3-chloro-8-fluoroquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.78 (d, J = 1.5 Hz, 1H), 8.17 (s, 1H), 7.99 (dd, J = 8.8, 5.7 Hz, 1H), 7.80 (dd, J = 8.8, 1.6 Hz, 1H); LCMS m/z= 277.83 (M+1; 100%).
 3,5-Dichloro-7-iodoquinoline 1-oxide	3,5-Dichloro-7-iodoquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.95 (d, J = 1.7 Hz, 1H), 8.80 (t, J = 1.3 Hz, 1H), 8.33 (d, J = 1.6 Hz, 1H), 8.08 (t, J = 1.3 Hz, 1H).







 3,7-Dibromo-5-fluoroquinoline 1-oxide	3,7-Dibromo-5-fluoroquinoline	LCMS m/z= 322.0 (M+1; 100%).
 3-Bromo-6-fluoro-7-iodoquinoline 1-oxide	3-Bromo-6-fluoro-7-iodoquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.91 (d, J = 6.0 Hz, 1H), 8.87 (d, J = 1.5 Hz, 1H), 8.22 (t, J = 1.1 Hz, 1H), 7.92 (d, J = 8.3 Hz, 1H).




7-Bromo-2,3-dichloro-5-fluoroquinoline

To a stirred solution of 7-bromo-3-chloro-5-fluoroquinoline 1-oxide (542 mg, 1.960 mmol) in CHCl₃ (10 ml) was added POCl₃ (1.867 ml, 20.03 mmol) at 25°C. The resulting mixture was stirred at 65°C for 2h under N₂ atmosphere. The reaction mixture was poured onto ice cold water (50ml), carefully basified with solid NaHCO₃ and extracted the product with dichloromethane (50ml). Layers were separated, organic layer was washed with brine (50ml) and was dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.2g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redise[®] R_f column with gradient elution (0 to 10%) of ethyl acetate in petroleum ether to afford the title compound (410mg, 70.9%) as a white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.49 – 8.43 (m, 1H), 8.04 (dt, J = 1.9, 1.0 Hz, 1H), 7.45 (dd, J = 8.9, 1.7 Hz, 1H); LCMS m/z= 296.19 (M+1; 100%). Intermediates in table-3 were synthesized by following an analogous reaction protocol as was used for the preparation of 7-bromo-2,3-dichloro-5-fluoroquinoline using the appropriate starting materials.


Table-3:

Intermediate's Structure	Starting materials used	¹ H NMR and LCMS data
--------------------------	-------------------------	----------------------------------

 7-Bromo-2-chloro-3-methylquinoline	7-Bromo-3-methylquinoline 1-oxide	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.40 (t, J = 1.0 Hz, 1H), 8.17 (d, J = 2.0 Hz, 1H), 7.94 (d, J = 8.7 Hz, 1H), 7.78 (dd, J = 8.7, 2.0 Hz, 1H), 2.48 (d, J = 1.0 Hz, 3H); LCMS m/z= 256, 258 (M ⁺ , M+2, 100%).
 7-Bromo-2-chloro-3-cyclopropyl quinoline	7-Bromo-3-cyclopropylquinoline 1-oxide	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (d, J = 3.0 Hz, 2H), 7.92 (d, J = 8.7 Hz, 1H), 7.77 (dd, J = 8.7, 2.0 Hz, 1H), 2.21 (tt, J = 8.4, 5.3 Hz, 1H), 1.10 (dt, J = 8.5, 3.2 Hz, 2H), 0.92 – 0.81 (m, 2H); LCMS m/z= 281.90, 283.90 (M ⁺ , M+2, 100%).
 7-Bromo-2-chloro-3-(1,1-difluoro ethyl)quinoline	7-Bromo-3-(1,1-difluoroethyl) quinoline 1-oxide	LCMS m/z= 306, 308 (M ⁺ , M+2; 100%).
 7-Bromo-2-chloro-3-methoxy quinoline	7-Bromo-3-methoxyquinoline 1-oxide	LCMS m/z= 271.69, 273.69
 7-Bromo-2,3-dichloro-6-fluoro quinoline	7-Bromo-3-chloro-6-fluoroquinoline 1-oxide	¹ H NMR (400 MHz, Chloroform-d) δ 8.31 (d, J = 6.3 Hz, 1H), 8.23 – 8.18 (m, 1H), 7.48 (d, J = 8.2 Hz, 1H).
	7-Bromo-3-chloro-8-fluoroquinoline 1-oxide	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.94 (d, J = 1.6 Hz, 1H),

7-Bromo-2,3-dichloro-8-fluoro quinoline		7.97 (dd, J = 8.9, 6.3 Hz, 1H), 7.83 (dd, J = 8.9, 1.3 Hz, 1H), LCMS m/z= 295.65 (M+1; 100%).
 2,3,5-Trichloro-7-iodoquinoline	3,5-Dichloro-7-iodoquinoline 1-oxide	¹ H NMR (400 MHz, Chloroform-d) δ 8.59 (d, J = 0.8 Hz, 1H), 8.38 (dd, J = 1.6, 0.8 Hz, 1H), 7.96 (d, J = 1.5 Hz, 1H).
 3,7-Dibromo-2-chloro-5-fluoroquinoline	3,7-Dibromo-5-fluoroquinoline 1-oxide	Crude was taken as such for the next step
 3-Bromo-2-chloro-6-fluoro-7-iodoquinoline	3-Bromo-6-fluoro-7-iodoquinoline 1-oxide	¹ H NMR (400 MHz, Chloroform-d) δ 8.54 (dd, J = 5.9, 0.7 Hz, 1H), 8.38 (s, 1H), 7.40 (d, J = 7.6 Hz, 1H).

7-Bromo-3-chloro-5-fluoroquinolin-2-amine







A mixture of 7-bromo-2,3-dichloro-5-fluoroquinoline (410mg, 1.390mmol), aqueous ammonia (9.74 ml, 250 mmol) in dioxane (10ml) was heated at 120°C in a steel bomb for 24h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.15g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (297 mg, 78%) as a white solid. ¹H NMR (400







MHz, DMSO-*d*₆) δ 8.23 (s, 1H), 7.51 (d, *J* = 1.9 Hz, 1H), 7.32 (dd, *J* = 9.5, 1.9 Hz, 1H), 7.25 (s, 2H); LCMS *m/z* = 276.83 (*M*+1; 100%).


Intermediates in table-4 were synthesized by following an analogous reaction protocol as was used for the preparation of 7-bromo-3-chloro-5-fluoroquinolin-2-amine using the

5 appropriate starting materials.


Table-4:

Intermediate's Structure	Starting materials used	¹ H NMR and LCMS data
 7-Bromo-8-fluoroquinolin-2-amine	7-Bromo-2-chloro-8-fluoroquinoline	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.95 (dd, <i>J</i> = 8.9, 1.8 Hz, 1H), 7.42 (dd, <i>J</i> = 8.7, 1.4 Hz, 1H), 7.35 – 7.27 (m, 1H), 6.93 (s, 2H), 6.88 – 6.79 (m, 1H); LCMS <i>m/z</i> = 240.8 (<i>M</i> +; 100%)
 7-Bromo-3-methylquinolin-2-amine	7-Bromo-2-chloro-3-methylquinoline	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.75 (t, <i>J</i> = 1.0 Hz, 1H), 7.60 (d, <i>J</i> = 2.0 Hz, 1H), 7.54 (d, <i>J</i> = 8.4 Hz, 1H), 7.25 (dd, <i>J</i> = 8.5, 2.0 Hz, 1H), 6.51 (s, 2H), 2.19 (d, <i>J</i> = 1.1 Hz, 3H); LCMS <i>m/z</i> = 237, 239 (<i>M</i> +, <i>M</i> +2, 100%).
 7-Bromo-3-isopropylquinolin-2-amine	7-Bromo-2-chloro-3-isopropyl quinoline	LCMS <i>m/z</i> = 265.1, 267.1 (<i>M</i> +, <i>M</i> +2; 100%).
 7-Bromo-3-cyclopropylquinolin-2-amine	7-Bromo-2-chloro-3-cyclopropyl quinoline	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.63 (s, 1H), 7.59 (d, <i>J</i> = 2.0 Hz, 1H), 7.55 (d, <i>J</i> = 8.5 Hz, 1H), 7.24 (dd, <i>J</i> = 8.5, 2.0 Hz, 1H), 6.65 (s, 2H), 1.80 (tdd, <i>J</i> = 8.2, 4.6, 2.9 Hz, 1H), 1.01 – 0.92 (m, 2H), 0.69 –


		0.62 (m, 2H); LCMS m/z= 262.83, 264.83 (M ⁺ , M+2, 100%).
 7-Bromo-3-(1,1-difluoroethyl)quinolin-2-amine	7-Bromo-2-chloro-3-(1,1-difluoroethyl)quinoline	LCMS m/z= 287.96 (M ⁺ , 100%).
 7-Bromo-3-methoxyquinolin-2-amine	7-Bromo-2-chloro-3-methoxyquinoline	LCMS m/z= 253, 255 (M ⁺ , M+2; 100%).
 7-Bromo-3-chloro-6-fluoroquinolin-2-amine	7-Bromo-2,3-dichloro-6-fluoroquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.23 (s, 1H), 7.79 (d, J = 6.7 Hz, 1H), 7.67 (dd, J = 9.3, 2.0 Hz, 1H), 6.97 (s, 2H); LCMS m/z= 276.83 (M ⁺ ; 100%).
 7-Bromo-3-chloro-8-fluoroquinolin-2-amine	7-Bromo-2,3-dichloro-8-fluoroquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.32 (d, J = 1.7 Hz, 1H), 7.30 (s, 2H), 7.35 – 7.16 (m, 2H), LCMS m/z= 276.86 (M ⁺ ; 100%).
 3,5-Dichloro-7-iodoquinolin-2-amine	2,3,5-Trichloro-7-iodoquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.22 (s, 1H), 7.94 – 7.81 (m, 1H), 7.67 (d, J = 1.6 Hz, 1H), 7.24 (s, 2H).
 3,7-Dibromo-5-fluoroquinolin-2-amine	3,7-Dibromo-2-chloro-5-fluoroquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.39 (d, J = 0.7 Hz, 1H), 7.50 (d, J = 1.7 Hz, 1H), 7.31 (dd, J = 9.5, 1.8 Hz, 1H), 7.14 (s, 2H); LCMS

		m/z= 318.96, 320.34, 322.34 (M-1, M+, M+2; 100%).
 3-Bromo-6-fluoro-7-iodoquinolin-2-amine	3-Bromo-2-chloro-6-fluoro-7-iodoquinoline	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.39 (s, 1H), 7.97 (d, J = 5.9 Hz, 1H), 7.55 (d, J = 8.5 Hz, 1H), 6.82 (s, 2H).

7-Bromo-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine



- 5 The title compound was prepared by following an analogous reaction protocol as described in Banka, Anna Lindsey et al, WO2012/037108 A1 using appropriate starting materials. ¹H NMR (400 MHz, DMSO-d₆) δ 8.22 (d, J = 0.8 Hz, 1H), 7.96 (t, J = 6.1 Hz, 1H), 7.58 (dd, J = 1.8, 1.0 Hz, 1H), 7.39 – 7.28 (m, 3H), 6.91 – 6.82 (m, 2H), 4.62 (d, J = 6.1 Hz, 2H), 3.71 (s, 3H); LCMS m/z= 397 (M+1; 100%).




10

- A mixture of 3,7-dibromo-5-fluoroquinolin-2-amine (2.05g, 6.41mmol), Et₃N (2.68ml, 19.22mmol), DMAP (0.078g, 0.641mmol) and Boc anhydride (3.12 ml, 13.45 mmol) in THF (25ml) was stirred at 25°C for 16h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 3.9g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 5%) of ethyl acetate in petroleum ether to afford the di-boc compound (2.5g, 75%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.65 (d, J = 0.8 Hz, 1H), 8.08 (q, J = 1.2 Hz, 1H), 7.46 (dd, J = 8.9, 1.7 Hz, 1H), 1.42 (s, 18H).
- 15
- 20


2-Amino-7-bromoquinoline-3-carbonitrile.

65




A mixture of 2-amino-4-bromobenzaldehyde (0.448 g, 2.240 mmol), malononitrile (0.222 g, 3.36 mmol) and piperidine (0.111ml, 1.120mmol) in ethanol (10ml) was stirred at 100 °C for 16h. The volatiles were evaporated *in vacuo* and the residue was purified by
 5 combiflash (R_f200, Teledyne/Isco) instrument onto a redise[®] R_f column with gradient elution (0 to 10%) of ethyl acetate in petroleum ether to afford the title compound (0.343g, 61.7%) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.71 (s, 1H), 7.74 – 7.65 (m, 2H), 7.42 (dd, J = 8.6, 2.0 Hz, 1H), 7.20 (s, 2H); LCMS m/z= 248, 250 (M⁺, M+2, 100%).

10 **7-Bromo-N-(4-methoxybenzyl)quinazolin-2-amine**




The title compound was prepared by following an analogous reaction protocol as described in Li, Liansheng et al, WO2017/087528 A1; LCMS m/z= 344.1 (M⁺, 100%).

7-Bromo-3-fluoroquinolin-2-amine




15 The title compound was prepared by an following analogous reaction protocol as described in Banka, Anna Lindsey et al, WO2012/037108 A1.

7-Bromo-3-chloroquinolin-2-amine



20 The title compound was prepared by following an analogous reaction protocol as described in Banka, Anna Lindsey et al, WO2012/037108 A1.


7-Bromo-3-chloro-N-(4-methoxybenzyl)quinolin-2-amine



The title compound was prepared by following an analogous reaction protocol as described
 25 in Banka, Anna Lindsey et al, WO2012/037108 A1.


7-Bromo-3-chloro-N,N-bis(4-methoxybenzyl)quinolin-2-amine

66




To a stirred suspension of 7-bromo-3-chloroquinolin-2-amine (2.0 g, 7.77 mmol) in DMF (20 ml) was added NaH (0.932 g, 23.30 mmol) at 0°C. The resulting mixture was stirred at 0°C for 15 min. Then 1-(chloromethyl)-4-methoxybenzene (3.65 g, 23.30 mmol) was added dropwise under N₂ atmosphere. The reaction mixture was then stirred for 16h at 25°C. The reaction mixture was poured into ice water (150 mL) and extracted with ethyl acetate (150 ml). Layers were separated, organic layer was washed with brine (100ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 3.87g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (2.2g, 56.9%) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.45 (s, 1H), 7.89 (d, J = 1.9 Hz, 1H), 7.77 (d, J = 8.6 Hz, 1H), 7.56 (dd, J = 8.6, 2.0 Hz, 1H), 7.31 – 7.22 (m, 4H), 6.93 – 6.82 (m, 4H), 4.54 (s, 4H), 3.70 (s, 6H); LCMS m/z = 498.97 (M+1; 100%).

15 3-Bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine



The title compound was prepared by following an analogous reaction protocol as described in Banka, Anna Lindsey et al, WO2012/037108 A1.

6-Bromo-3,3-dimethylindoline-2-thione.




20

The suspension of 6-bromo-3,3-dimethylindolin-2-one (3.5 g, 14.58 mmol), which was synthesized by following an analogous reaction protocol as was reported in WO2015/177110, A1 and Lawesson's reagent (7.66 g, 18.95 mmol) in toluene (15 ml) was heated at 100°C for 3h under N₂ atmosphere. Solvent was evaporated *in vacuo* and this residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0-20%) of ethyl acetate in petroleum ether to afford the title compound (3.3g, 88 %) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 12.67 (s,

25


1H), 7.40 (d, J = 7.9 Hz, 1H), 7.31 (dd, J = 7.9, 1.8 Hz, 1H), 7.15 (d, J = 1.7 Hz, 1H), 1.30 (s, 6H); LCMS m/z= 256.89 (M+1; 30%).

6-Bromo-3,3-dimethyl-2-(methylthio)-3H-indole.




- 5 To a stirred suspension of 6-bromo-3,3-dimethylindoline-2-thione (3 g, 11.71 mmol) in THF (40ml) was added NaH (0.703g, 17.57mmol) at 0°C. The resulting mixture was stirred for 15 min at 0°C. Methyl iodide (1.098 ml, 17.57 mmol) was added and stirred the reaction mixture for 1h at 0°C under N₂ atmosphere. The reaction mixture was diluted with ethyl acetate (50 ml) and washed with water (50 ml). Layers were separated, organic layer was
- 10 washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 3.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisept® R_f column with gradient elution (0-5%) of ethyl acetate in petroleum ether to afford the title compound (3.1g, 98%) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.60 (d, J = 1.7 Hz, 1H),
- 15 7.41 (d, J = 7.9 Hz, 1H), 7.32 (dd, J = 7.9, 1.8 Hz, 1H), 2.60 (s, 3H), 1.29 (s, 6H); LCMS m/z= 269.90 (M+; 40%).

6'-Bromospiro[cyclobutane-1,3'-indoline]-2'-thione.



- The title compound was prepared by following an analogous reaction protocol as was
- 20 described in the preparation of 6-bromo-3,3-dimethylindoline-2-thione. LCMS m/z= 267.65 (M+; 20%).


6'-Bromo-2'-(methylthio)spiro[cyclobutane-1,3'-indole]



- The title compound was prepared by following an analogous reaction protocol as was
- 25 described in the preparation of 6-bromo-3,3-dimethyl-2-(methylthio)-3H-indole. ¹H NMR

(400 MHz, Chloroform-d) δ 7.63 (d, J = 1.7 Hz, 1H), 7.44 (d, J = 7.9 Hz, 1H), 7.32 (dd, J = 7.9, 1.7 Hz, 1H), 2.73 (s, 3H), 2.61 – 2.46 (m, 4H), 2.37 – 2.30 (m, 2H); LCMS m/z = 281.78, 283.78 (M^+ , $M+2$; 100%).

6'-Bromospiro[cyclobutane-1,3'-indol]-2'-amine.




5

To stirred solution of 6'-bromo-2'-(methylthio)spiro[cyclobutane-1,3'-indole] (1.2 g, 4.25 mmol) in 7N ammonia in methanol (15ml, 371mmol) was heated at 100°C for 16 h. Solvent was evaporated *in vacuo* and this residue was purified by combiflash (R_f 200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0-10%) of methanol in dichloromethane to afford the title compound (650mg, 60.9%) as white solid.

^1H NMR (400 MHz, DMSO- d_6) δ 7.69 (s, 2H), 7.46 (d, J = 8.2 Hz, 1H), 7.01 (h, J = 1.8 Hz, 2H), 2.63 – 2.53 (m, 2H), 2.43 – 2.30 (m, 1H), 2.23 – 2.09 (m, 3H); LCMS m/z = 250.87 (M^+ ; 100%).

10


7-Bromo-3-methylimidazo[1,2-a]pyridine



15

The title compound was prepared by following an analogous reaction protocol as was described in Dubois., Laurent et al, WO 2009/112679 A1.

2-(Methylamino)quinolin-7-ol




20

The title compound was prepared by following an analogous reaction protocol as was described in Doherty, Elizabeth M. et.al, Journal of Medicinal Chemistry, 2007, vol. 50, # 15, p. 3515 – 3527.


3-Bromo-N-(4-methoxybenzyl)-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2-amine

69




A mixture of 3-bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine (1.5g, 3.20mmol), bispinacolatodiboron (0.974g, 3.84mmol), [1,1'-bis(diphenylphosphino) ferrocene] dichloropalladium(II), complex with dichloromethane (0.261g, 0.320mmol) and potassium acetate (0.533g, 5.44mmol) in DMSO (50ml) was heated at 80°C for 15min in a preheated oil bath. The reaction mixture was allowed to cool to 25°C, diluted with ethyl acetate (50ml) and poured onto ice-cold water (100ml). Layers were separated, organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 2.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-7%) of ethyl acetate in petroleum ether to afford the title compound (1.35g, 90%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.42 (d, *J* = 0.7 Hz, 1H), 7.86 (d, *J* = 1.1 Hz, 1H), 7.65 (d, *J* = 7.9 Hz, 1H), 7.43 (dd, *J* = 7.9, 1.1 Hz, 1H), 7.40 – 7.32 (m, 3H), 6.99 – 6.74 (m, 2H), 4.62 (d, *J* = 6.0 Hz, 2H), 3.71 (s, 3H), 1.32 (s, 12H); LCMS *m/z* = 468.89 (M⁺; 100%).

3-Bromo-2-((4-methoxybenzyl)amino)quinolin-7-ol




To a stirred solution of 3-bromo-N-(4-methoxybenzyl)-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2-amine (0.35g, 0.746mmol) in THF (20 ml) was added AcOH (0.064ml, 1.119mmol) dropwise at 0°C and stirred for 1h. Aq. H₂O₂ (0.5ml, 1.492mmol) was added slowly at 0°C. The resulting mixture was stirred at 25°C for 5h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with aq.sat.Na₂SO₃ (20ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 0.41g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-25%) of ethyl acetate in petroleum ether to afford the title compound (0.13g, 48.5%) as an off-white solid. LCMS *m/z* = 359.22 (M⁺; 100%).

3-Chloro-5-fluoro-N-(4-methoxybenzyl)-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2-amine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 3-bromo-N-(4-methoxybenzyl)-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2-amine. ¹H NMR (400 MHz, Chloroform-d) δ 8.17 (d, J = 0.9 Hz, 1H), 8.07 (s, 1H), 7.41 – 7.36 (m, 2H), 7.29 (dd, J = 10.0, 0.8 Hz, 1H), 6.94 – 6.89 (m, 2H), 5.69 (s, 1H), 4.77 (d, J = 5.4 Hz, 2H), 3.83 (s, 3H), 1.29 (s, 12H); LCMS m/z = 443.05 (M⁺; 100%).

3-Chloro-5-fluoro-2-((4-methoxybenzyl)amino)quinolin-7-ol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 3-bromo-2-((4-methoxybenzyl)amino)quinolin-7-ol. ¹H NMR (400 MHz, DMSO-d₆) δ 10.19 (d, J = 1.0 Hz, 1H), 8.02 (s, 1H), 7.55 (t, J = 6.2 Hz, 1H), 7.43 – 7.28 (m, 2H), 6.91 – 6.82 (m, 2H), 6.67 (d, J = 2.1 Hz, 1H), 6.57 (dd, J = 11.7, 2.2 Hz, 1H), 4.60 (d, J = 6.0 Hz, 2H), 3.71 (s, 3H); LCMS m/z = 333.15, 335.15 (M⁺, M+2; 30%).


N2-(4-Methoxybenzyl)quinoline-2,7-diamine.



- 20 A mixture of 7-bromo-N-(4-methoxybenzyl)quinolin-2-amine (0.6 g, 1.748 mmol), copper(I) iodide (0.033g, 0.175mmol), and N1,N1-dimethylethane-1,2-diamine (0.171ml, 1.748mmol) in AMMONIA (0.757 ml, 35.0 mmol) and DMSO (1ml) at 130°C for 15h. The resulting suspension was cooled to 25°C and saturated aqueous sodium sulphate solution (5mL) was added. The resulting mixture was extracted with ethyl acetate (20ml X
- 25 3). The organic layer was separated, dried over MgSO₄, filtered and concentrated. This


residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-5%) of methanol in dichloromethane to afford the title compound (0.18g, 36.9%) as a brown solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.55 (d, J = 8.7 Hz, 1H), 7.34 – 7.29 (m, 2H), 7.25 (d, J = 8.4 Hz, 1H), 7.07 (t, J = 5.8 Hz, 1H), 6.91 – 6.83 (m, 2H), 6.56 (d, J = 2.2 Hz, 1H), 6.52 (dd, J = 8.4, 2.2 Hz, 1H), 6.40 (d, J = 8.7 Hz, 1H), 5.32 (s, 2H), 4.49 (d, J = 5.8 Hz, 2H), 3.72 (s, 3H); LCMS m/z= 280.2 (M+1, 100%).

N2-(4-Methoxybenzyl)-N7-methylquinoline-2,7-diamine.




A mixture of 7-bromo-N-(4-methoxybenzyl)quinolin-2-amine (0.5 g, 1.457 mmol), methanamine (7.54 ml, 7.28 mmol) and copper (0.046 g, 0.728 mmol) was stirred at 110°C in a sealed tube for 12h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). The organic layer was separated, dried over MgSO₄, filtered and concentrated. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-5%) of methanol in dichloromethane to afford the title compound (0.4g, 94%) as a brown liquid. LCMS m/z= 294.09 (M+1; 100%).


S-(2-((4-Methoxybenzyl)amino)quinolin-7-yl) ethanethioate.




To a stirred solution of 7-bromo-N-(4-methoxybenzyl)quinolin-2-amine (2g, 5.83mmol), potassium thioacetate (1.331g, 11.65mmol), DIEA (2.035ml, 11.65mmol) and xantphos (0.337g, 0.583mmol) in 1,4-dioxane (20ml) was added Pd₂(dba)₃ (0.534g, 0.583mmol) at 25°C under N₂ atmosphere. The resulting mixture was stirred at 110°C for 1.5h under microwave condition. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). The organic layer was separated, dried over MgSO₄, filtered and concentrated. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-20%) of ethyl acetate in petroleum ether to afford the title compound (0.67g, 34%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 7.87 – 7.80 (m, 2H), 7.63 (d, J = 8.3 Hz, 1H), 7.39 – 7.32 (m, 2H), 7.28 – 7.23 (m, 1H), 6.95 – 6.87 (m, 2H), 6.67 (d, J = 9.0 Hz, 1H), 5.21 (s, 1H), 4.67 (d, J = 5.4 Hz, 2H), 3.82 (s, 3H), 2.47 (s, 3H); LCMS m/z= 339.22 (M+1; 100%).

2-((4-Methoxybenzyl)amino)quinoline-7-thiol.

To a stirred solution of S-(2-((4-methoxybenzyl)amino)quinolin-7-yl) ethanethioate (0.67g, 1.98 mmol) in 20 mL of ethanol was added KOH (0.33 g, 5.94 mmol) at 25°C. The resulting mixture was stirred at 25°C for 1h. The reaction mixture was quenched with sat. aqueous NH₄Cl (20ml) and extracted with ethyl acetate (20ml). The organic layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo* to give the title compound (0.56g, 95%) as a brown solid. ¹H NMR (400 MHz, Chloroform-d) δ 7.77 (d, J = 8.9 Hz, 1H), 7.64 (d, J = 1.8 Hz, 1H), 7.46 (d, J = 8.3 Hz, 1H), 7.37 – 7.32 (m, 2H), 7.10 (dd, J = 8.3, 1.9 Hz, 1H), 6.93 – 6.87 (m, 2H), 6.58 (d, J = 8.9 Hz, 1H), 5.58 (s, 1H), 4.64 (d, J = 5.3 Hz, 2H), 3.82 (s, 3H); LCMS m/z = 297.09 (M⁺; 100%).

S-(2-(Bis(4-methoxybenzyl)amino)-3-chloroquinolin-7-yl) ethanethioate


The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of S-(2-((4-methoxybenzyl)amino)quinolin-7-yl) ethanethioate. LCMS m/z = 493.30 (M⁺; 100%).

2-(Bis(4-methoxybenzyl)amino)-3-chloroquinoline-7-thiol

To a solution of solution of S-(2-(bis(4-methoxybenzyl)amino)-3-chloroquinolin-7-yl) ethanethioate (300 mg, 0.608 mmol) in ethanol (9 ml) was added KOH (51.2 mg, 0.913 mmol) at 25°C. The mixture was heated at 50°C for 2h. The reaction solution was allowed to cool to 25°C, adjusted pH to 4 with HCl (1N) and concentrated *in vacuo* to get 0.4g of crude compound, which was triturated with diethyl ether (20ml) to get the title compound (200 mg, 72.9%) as an off-white solid. LCMS m/z = 450.42 (M⁺; 100%).


((3aR,6R,6aS)-6-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol

73




The title compound was prepared by following the same reaction protocol as was described in Kenneth A. Jacobson et.al; Purinergic Signalling (2015) 11:371–387.

5 ((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl 4-methylbenzenesulfonate




To a stirred solution of ((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol (0.5 g, 1.554 mmol) in dichloromethane (10ml) was added TEA (0.651 ml, 4.66 mmol), DMAP (0.038 g, 0.311 mmol), followed by a slow addition of p-TsCl (0.355 g, 1.865 mmol) at 0°C and stirred for 10mins. The resulting mixture was stirred at 25°C for 2h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). The organic layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo* to give 0.71g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redise[®] R_f column with gradient elution (0-10%) of ethyl acetate in petroleum ether to afford the title compound (0.282g, 38.1%) as a pale yellow liquid. ¹H NMR (400 MHz, Chloroform-d) δ 8.69 (t, J = 2.8 Hz, 1H), 7.89 – 7.81 (m, 2H), 7.42 – 7.34 (m, 2H), 7.12 (s, 1H), 6.64 (d, J = 3.8 Hz, 1H), 5.86 (d, J = 12.4 Hz, 2H), 5.36 (d, J = 5.6 Hz, 1H), 4.90 – 4.77 (m, 2H), 4.63 (t, J = 4.5 Hz, 1H), 2.52 – 2.45 (m, 3H), 1.45 (s, 3H), 1.34 (s, 3H); LCMS m/z = 476.17 (M⁺; 100%).

(3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde



To a stirred solution of ((3a*S*,4*R*,6a*R*)-4-(4-Chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxol-6-yl)methanol (2.50g, 7.77mmol) in CH₂Cl₂ (40ml) at 0°C, was added Dess-Martin Periodinane (3.95 g, 9.32 mmol) portion-wise and stirred for 1h. The reaction mixture was diluted with dichloromethane (50ml) and washed with water (50ml). The organic layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo* to give 2.71g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-30%) of ethyl acetate in petroleum ether to afford the title compound (2.32g, 93%) as a colorless oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 10.00 (s, 1H), 8.67 (s, 1H), 7.12 (d, *J* = 3.6 Hz, 1H), 6.78 (dd, *J* = 2.6, 0.9 Hz, 1H), 6.69 (d, *J* = 3.7 Hz, 1H), 5.97 (dt, *J* = 2.8, 1.4 Hz, 1H), 5.76 (dd, *J* = 5.9, 1.5 Hz, 1H), 4.88 (dt, *J* = 5.9, 1.1 Hz, 1H), 1.54 (s, 3H), 1.40 (s, 3H); LCMS *m/z* = 320.2 (*M*+1, 100%).

4-Chloro-7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine




15

To a cooled suspension of methyl(triphenyl)phosphonium bromide (5.03g, 14.07mmol) in THF (30mL) at 0°C, was added 1M KHMDS in THF (14.07mL, 14.07mmol) slowly and stirred for 5 min. The reaction mixture was allowed to warm to 25°C and stirred for 10min. Cooled the reaction mixture to 0°C and slowly added a solution of (3a*S*,4*R*,6a*R*)-4-(4-Chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxole-6-carbaldehyde (1.8g, 5.63mmol) in THF (1ml). Stirred the reaction mixture at 25°C for 10min. The reaction mixture was quenched with sat. aqueous NH₄Cl (50ml) and extracted with ethyl acetate (50ml). The organic layer was separated, dried over Na₂SO₄, filtered and concentrated *in vacuo* to give 2.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0-7%) of ethyl acetate in petroleum ether to afford the title compound (0.81g, 45.3%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.72 (s, 1H), 7.11 (d, *J* = 3.6 Hz, 1H), 6.66 – 6.57 (m, 2H), 5.94 (d, *J* = 2.6 Hz, 1H), 5.81 – 5.75 (m, 2H),

25


5.57 (dd, $J = 6.0, 1.5$ Hz, 1H), 5.49 (d, $J = 10.9$ Hz, 1H), 4.66 (dt, $J = 6.0, 1.0$ Hz, 1H), 1.52 (s, 3H), 1.40 (s, 3H); LCMS $m/z = 318.09$ ($M+1$, 100%).

7-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta [d] [1,3] dioxol-4-yl)-4-ethyl-7H-pyrrolo[2,3-d]pyrimidine



5 To a stirred solution of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (500mg, 1.573mmol) in THF (1ml) and NMP (0.2ml) was added ferric acetylacetonate (55.6 mg, 0.157mmol) at 25°C. 2M Ethylmagnesium chloride in THF (1.573ml, 3.15mmol) was added dropwise and the
 10 reaction mixture was stirred for 4h. The reaction mixture was quenched with sat. aqueous NH_4Cl (10ml) and extracted with ethyl acetate (10ml). The organic layer was separated, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to give 0.6g of crude compound. This residue was purified by combiflash (R_f 200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0-90%) of ethyl acetate in petroleum ether to afford the
 15 title compound (0.28g, 57.1%) as an off-white solid. ^1H NMR (400 MHz, DMSO-d_6) δ 8.71 (s, 1H), 7.35 (d, $J = 3.6$ Hz, 1H), 6.73 (d, $J = 3.6$ Hz, 1H), 6.60 (dd, $J = 17.7, 10.8$ Hz, 1H), 5.88 (d, $J = 2.7$ Hz, 1H), 5.83 – 5.74 (m, 1H), 5.66 – 5.56 (m, 2H), 5.45 – 5.37 (m, 1H), 4.67 (dd, $J = 6.0, 1.1$ Hz, 1H), 2.99 (q, $J = 7.6$ Hz, 2H), 1.40 (s, 3H), 1.30 (s, 6H); LCMS $m/z = 312.21$ ($M+1$, 100%).


20 **7-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d] [1,3]dioxol-4-yl)-4-isopropyl-7H-pyrrolo[2,3-d]pyrimidine**



A solution of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (1g, 3.15mmol), 2M
 25 isopropyl magnesium bromide in THF (5.51 ml, 11.01 mmol) and $\text{PdCl}_2(\text{dppf})$ (0.230 g, 0.315 mmol) in toluene (10 ml) was heated at 100°C for 30min. After completion of the


reaction, the reaction mixture was quenched with methanol and concentrated. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0-2%) of methanol in dichloromethane to afford the title compound (150mg, 14.65%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.98 (s, 1H), 7.48 (d, *J* = 4.0 Hz, 1H), 7.11 (t, *J* = 1.6 Hz, 1H), 6.86 (d, *J* = 4.0 Hz, 1H), 3.98 (dt, *J* = 2.5, 1.2 Hz, 2H), 3.50 – 3.43 (m, 1H), 2.66 (s, 1H), 2.20 (d, *J* = 2.4 Hz, 1H), 2.00 (dd, *J* = 7.2, 1.0 Hz, 2H), 1.44 (dd, *J* = 6.9, 3.1 Hz, 6H), 1.28 (s, 6H).

4-Cyclopropyl-7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d] [1,3]dioxol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine.



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta [d] [1,3] dioxol-4-yl)-4-ethyl-7H-pyrrolo[2,3-*d*]pyrimidine. LCMS *m/z* = 323.90 (*M*⁺; 100%).


1-((3a*S*,4*R*,6a*R*)-4-(4-Chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-ol.



To a stirred solution of (3a*S*,4*R*,6a*R*)-4-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde (2.30 g, 7.19 mmol) in THF (120ml) at -78°C, was added methylmagnesium bromide (4.80 ml, 14.39 mmol) and the reaction mixture was stirred at same temperature for 3h. The reaction mixture was quenched with a sat. aqueous NH₄Cl (50 ml) and extracted with ethyl acetate (50 ml). Layers were separated, the organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 2.2g of crude compound and this crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 20%) of


ethyl acetate in petroleum ether to afford the title compound (1.82g, 75%) as a off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.70 (s, 1H), 7.11 (d, J = 3.7 Hz, 1H), 6.63 (d, J = 3.7 Hz, 1H), 5.88 – 5.81 (m, 1H), 5.80 – 5.71 (m, 1H), 5.59 – 5.45 (m, 1H), 4.76 – 4.64 (m, 2H), 1.59 – 1.49 (m, 6H), 1.38 (d, J = 1.0 Hz, 3H), 1.32 – 1.23 (m, 1H); LCMS m/z = 336.2 (M⁺; 100%).

1-((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-one.




To a stirred solution of 1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-ol (4.75 g, 14.15 mmol) in dichloromethane (45 ml) at 0 °C, was added Dess-Martin Periodinane (7.20 g, 16.97 mmol) portion-wise and stirred for 30 min. Water (50ml) was added and filtered the reaction mixture through a celite bed, washed with dichloromethane (25ml X 2). Separated the layers and the organic layer was washed with brine (50ml), dried over anhydrous sodium sulphate and concentrated *in vacuo* to give 4.8g crude compound. This crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a rediseq® Rf column with gradient elution (0 to 50%) of ethyl acetate in petroleum ether to afford the title compound (3.8g, 80%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.68 (s, 1H), 7.10 (d, J = 3.7 Hz, 1H), 6.68 (d, J = 3.6 Hz, 1H), 6.63 (dt, J = 2.7, 0.7 Hz, 1H), 5.96 (dt, J = 2.8, 1.5 Hz, 1H), 5.75 (dd, J = 5.9, 1.6 Hz, 1H), 4.82 (dt, J = 5.9, 1.1 Hz, 1H), 2.47 (s, 3H), 1.52 (s, 3H), 1.39(s, 3H); LCMS m/z = 334.09 (M⁺; 100%).

2-((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propan-2-ol.



To a stirred solution of 1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-one (1.0 g, 3.00 mmol) in THF (10 ml) at -20 °C, was added dropwise methyl magnesium bromide (1.498 ml, 4.49 mmol) and stirred the reaction mixture for 30 min at the same temperature. The reaction mixture was quenched with a sat. aqueous NH₄Cl (50 ml) and extracted with ethyl acetate (50 ml). Layers were separated, the organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated in vacuo to give 1.5g of crude compound and this crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisepp® Rf column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (850 mg, 81%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.69 (s, 1H), 7.09 (d, J = 3.6 Hz, 1H), 6.71 – 6.60 (m, 1H), 5.82 (s, 1H), 5.78 – 5.69 (m, 1H), 5.65 – 5.58 (m, 1H), 4.69 (dt, J = 5.8, 1.0 Hz, 1H), 1.59 – 1.53 (m, 9H), 1.38 (s, 3H); LCMS m/z = 350.2 (M⁺; 100%).


4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine.



To a stirred solution of 2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propan-2-ol (8.2 g, 23.44 mmol) in dichloromethane (80ml) at 0°C was added Martin's Sulfurane (17.34 g, 25.8 mmol) and the stirred the reaction mixture for 45 min at 25°C. The reaction mixture was quenched with an sat.aq. sodium bicarbonate (100ml), extracted with dichloromethane (100ml). Layers were separated, the organic layer was washed with brine (50 ml) and was dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 8.5g of crude compound. This crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisepp® Rf column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (4.2g, 54.0%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.71 (s, 1H), 7.11 (d, J = 3.6 Hz, 1H), 6.63 (d, J = 3.6 Hz, 1H), 6.00 – 5.94 (m, 1H), 5.76 (d, J = 2.7 Hz, 1H), 5.62 – 5.52 (m, 2H),

5.33 (d, $J = 1.7$ Hz, 1H), 4.67 (dt, $J = 6.0, 1.0$ Hz, 1H), 2.03 (t, $J = 1.0$ Hz, 3H), 1.52 (s, 3H), 1.40 (s, 3H); LCMS $m/z = 332.28$ (M^+ ; 100%).

7-((3aS,4R,6aR)-2,2-Dimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine



5



To a degassed solution of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (1.000 g, 3.01 mmol) in dioxane (10 ml) and water (1 ml), was added potassium phosphate, tribasic (1.575 g, 9.04 mmol), dichloro[1,1'-bis(di-*t*-butylphosphino)ferrocene]palladium(II) (0.196 g, 0.301 mmol) and 2,4,6-trimethyl-1,3,5,2,4,6-trioxatriborinane (4.21 ml, 30.1 mmol) at 25°C. The resulting mixture was stirred at 100°C for 50min under microwave condition. Solvent was removed and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisepp® Rf column with gradient elution (0 to 50%) of ethyl acetate in petroleum ether to afford the title compound (0.84g, 90%) as an off-white solid. LCMS $m/z = 312.28$ (M^+ ; 100%).

15

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine (A).

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (B).

20




To a degassed solution of 7-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-7H-pyrrolo[2,3-d]pyrimidine (5g, 8.93 mmol) in dioxane (80 ml) and water (10ml), was added potassium phosphate tribasic (4.66 g, 26.8 mmol), dichloro[1,1'-bis(di-*t*-butylphosphino)ferrocene]palladium(II) (0.582 g, 0.893 mmol) and 2,4,6-trimethyl-1,3,5,2,4,6-

25

trioxatriborinane (12.48 ml, 89 mmol) at 25°C. The reaction mixture was heated at 80°C for 8h. The reaction mixture was diluted with ethyl acetate (50 ml) and washed with water (50 ml). Layers were separated and the organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 4.3g of crude compound. This crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound, A (3.2g, 66%) and B (0.75g, 15.98%) as an off-white solids. ¹H NMR of A (400 MHz, Chloroform-d) δ 8.83 (s, 1H), 7.71 (tt, *J* = 6.6, 1.5 Hz, 4H), 7.48 – 7.37 (m, 6H), 6.91 (d, *J* = 3.6 Hz, 1H), 6.57 (d, *J* = 3.6 Hz, 1H), 5.88 (s, 2H), 5.25 (d, *J* = 5.7 Hz, 1H), 4.60 (d, *J* = 5.7 Hz, 1H), 4.55 – 4.45 (m, 2H), 2.77 (s, 3H), 1.45 (s, 3H), 1.32 (s, 3H), 1.11 (s, 9H); LCMS *m/z* = 540.4 (*M*+1; 100%); ¹H NMR of B (400 MHz, Chloroform-d) δ 7.71 (tt, *J* = 6.6, 1.5 Hz, 4H), 7.53 – 7.35 (m, 6H), 6.98 (d, *J* = 3.6 Hz, 1H), 6.57 (d, *J* = 3.6 Hz, 1H), 5.90 (d, *J* = 14.5 Hz, 2H), 5.26 (d, *J* = 5.7 Hz, 1H), 4.61 (d, *J* = 5.7 Hz, 1H), 4.51 (d, *J* = 9.3 Hz, 2H), 1.46 (s, 3H), 1.33 (s, 3H), 1.28 (s, 2H), 1.11 (s, 9H); LCMS *m/z* = 526.44 (*M*+1; 100%).


((3a*S*,4*R*,6a*R*)-2,2-dimethyl-4-(4-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3]dioxol-6-yl)methanol



To a stirred solution of 7-((3a*S*,4*R*,6a*R*)-6-(((tert-butyl)diphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3]dioxol-4-yl)-4-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidine (3.20 g, 5.93 mmol) in THF (20 ml), was slowly added TBAF (8.89 ml, 8.89 mmol) at 25°C and stirred the reaction mixture at 25°C for 15h. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 100%) of ethyl acetate in petroleum ether to afford the title compound (1.5g, 84%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.81 (s, 1H), 7.06 (d, *J* = 3.6 Hz, 1H), 6.55 (d, *J* = 3.6 Hz, 1H), 5.90 – 5.78 (m, 2H), 5.41 (ddd, *J* = 5.8, 1.7, 0.9 Hz, 1H), 4.65 (dt, *J* = 5.8, 0.9 Hz, 1H), 4.56 –

4.42 (m, 2H), 3.35 (d, $J = 8.2$ Hz, 1H), 2.74 (s, 3H), 1.53 (s, 3H), 1.37 (s, 3H); LCMS $m/z = 302.21$ ($M+1$; 100%) .


(3aS,4R,6aR)-2,2-Dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde.



5


To a stirred solution of 7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (1.50 g, 4.98 mmol) in dichloromethane (100 ml) at 0°C, was added Dess-Martin Periodinane (2.53 g, 5.97 mmol) portion-wise and stirred for 1h. The reaction mixture was diluted with methylene chloride (50 ml) and washed with water (50 ml). Layers were separated, the organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated in vacuo to give a crude compound and this crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (0.95g, 63.8%) as an off-white solid. ¹H NMR (400 MHz, Chloroform-d) δ 10.00 (s, 1H), 8.80 (s, 1H), 7.02 (d, $J = 3.6$ Hz, 1H), 6.82 – 6.76 (m, 1H), 6.63 (d, $J = 3.6$ Hz, 1H), 6.00 (dt, $J = 2.7, 1.4$ Hz, 1H), 5.76 (dd, $J = 5.9, 1.5$ Hz, 1H), 4.87 (dt, $J = 5.9, 1.1$ Hz, 1H), 2.77 (s, 3H), 1.53 (s, 3H), 1.38 (s, 3H); LCMS $m/z = 300.15$ ($M+1$; 100%)

7-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine.




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-d) δ 8.83 (s, 1H), 7.01 (d, J = 3.6 Hz, 1H), 6.68 – 6.53 (m, 2H), 5.95 (d, J = 2.5 Hz, 1H), 5.80 – 5.71 (m, 2H), 5.56 (dd, J = 6.0, 1.4 Hz, 1H), 5.47 (d, J = 10.8 Hz, 1H), 4.65 (d, J = 5.8 Hz, 1H), 2.75 (s, 3H), 1.52 (s, 3H), 1.40 (s, 3H); LCMS m/z = 298.5 (M+1; 100%).

((3aS,4R,6aR)-2,2-Dimethyl-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol.




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3aS,4R,6aR)-6-(((tert-butyl)diphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. LCMS m/z = 287.90 (M+; 100%).

(3aS,4R,6aR)-2,2-Dimethyl-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde.




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-Dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-d) δ 10.01 (s, 1H), 9.04 (s, 1H), 8.93 (s, 1H), 7.11 (d, J = 3.6 Hz, 1H), 6.83 – 6.77 (m, 1H), 6.67 (d, J = 3.7 Hz, 1H), 6.03 (dt, J = 2.8, 1.4 Hz, 1H), 5.77 (dd, J = 5.9, 1.5 Hz, 1H), 4.89 (dt, J = 6.0, 1.2 Hz, 1H), 1.54 (s, 3H), 1.41 (s, 3H); LCMS m/z = 286.09 (M+; 100%).

7-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d] [1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine.




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-d) δ 8.97 (d, *J* = 15.8 Hz, 2H), 7.09 (d, *J* = 3.7 Hz, 1H), 6.68 – 6.56 (m, 2H), 5.98 (d, *J* = 2.7 Hz, 1H), 5.82 – 5.72 (m, 2H), 5.60 – 5.54 (m, 1H), 5.52 – 5.44 (m, 1H), 4.66 (dt, *J* = 5.8, 1.0 Hz, 1H), 1.52 (s, 3H), 1.40 (s, 3H); LCMS *m/z* = 284.04 (*M*+1; 100%).

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-5-fluoro-7H-pyrrolo[2,3-d] pyrimidine




To a stirred solution of (3aS,4S,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (3.5 g, 8.24 mmol) in THF (50 ml) was added 4-chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidine (3.54 g, 20.61 mmol), which was synthesized by following the same reaction protocol as was described in WO2005/16878, A2), triphenylphosphine (5.40 g, 20.61 mmol), DIAD (4.01 ml, 20.61 mmol) at 0°C and stirred for 30 min. The resulting reaction mixture was stirred at 25°C for 16h. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 10%) of ethyl acetate in petroleum ether to afford the title compound (2.7g, 56.7%) as an off-white solid. LCMS *m/z* = 577.94 (*M*+; 100%).

((3aR,6R,6aS)-6-(4-Chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3aS,4R,6aR)-6-(((tert-butylidiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. LCMS m/z = 340.03 (M^+ ; 100%).

(3aR,6R,6aS)-6-(4-Chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxole-4-carbaldehyde




10

The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-Dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ^1H NMR (400 MHz, DMSO- d_6) δ 9.90 (s, 1H), 8.73 (s, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.10 – 7.03 (m, 1H), 6.01 (dq, J = 2.7, 1.4 Hz, 1H), 5.57 (dd, J = 6.0, 1.5 Hz, 1H), 4.80 (dt, J = 6.0, 1.2 Hz, 1H), 1.39 (s, 3H), 1.28 (s, 3H); LCMS m/z = 338.03 (M^+ ; 100%).

15

4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-5-fluoro-7H-pyrrolo[2,3-d]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-

20


dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.71 (s, 1H), 7.66 (d, *J* = 2.0 Hz, 1H), 6.66 – 6.52 (m, 1H), 5.90 – 5.78 (m, 2H), 5.67 – 5.55 (m, 2H), 5.47 – 5.36 (m, 1H), 4.71 (d, *J* = 6.0 Hz, 1H), 1.38 (s, 3H), 1.31 (s, 3H); LCMS *m/z* = 336.03 (M⁺; 100%).

5 **7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-2-chloro-7H-pyrrolo[2,3-d]pyrimidine**



To a stirred solution of (3aS,4S,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (1.2g, 2.83 mmol) in THF
10 (15ml) was added 2-chloro-7H-pyrrolo[2,3-d]pyrimidine (0.738 g, 4.80 mmol), triphenylphosphine (2.59 g, 9.89 mmol) and DIAD (1.923 ml, 9.89 mmol) slowly at 0°C and stirred for 5mins. The reaction mixture was brought to 25°C and stirred for 1h. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisept® Rf column with gradient elution (0 to 15%) of
15 ethyl acetate in petroleum ether to afford the title compound (1.1g, 69.5%) as an off- white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.84 (s, 1H), 7.70 (ddt, *J* = 6.6, 5.0, 1.5 Hz, 4H), 7.50 – 7.34 (m, 6H), 6.94 (d, *J* = 3.6 Hz, 1H), 6.59 (d, *J* = 3.6 Hz, 1H), 5.84 (dt, *J* = 19.8, 2.2 Hz, 2H), 5.30 (d, *J* = 5.7 Hz, 1H), 4.63 (d, *J* = 5.6 Hz, 1H), 4.57 – 4.42 (m, 2H), 1.43 (s, 3H), 1.33 (s, 3H), 1.10 (s, 9H); LCMS *m/z* = 560.3 (M⁺; 100).


20 **((3aR,6R,6aS)-6-(2-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol**



To a stirred solution of 7-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-2-chloro-7H-pyrrolo[2,3-
25 d]pyrimidine (500 mg, 0.893 mmol) in THF (5 ml) at 0 °C, was added TBAF (1.250 ml, 1.250 mmol) slowly and stirred the reaction mixture at the same temperature for 10 min.


The reaction mixture was brought to 25°C and stirred for 30mins. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 40%) of ethyl acetate in petroleum ether to afford the title compound (0.27g, 94%) as a colourless oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.81 (s, 1H), 7.28 (s, 1H), 7.08 (d, *J* = 3.7 Hz, 1H), 6.57 (d, *J* = 3.7 Hz, 1H), 5.92 – 5.62 (m, 2H), 5.47 (d, *J* = 5.7 Hz, 1H), 4.69 (dt, *J* = 5.6, 0.9 Hz, 1H), 4.59 – 4.37 (m, 2H), 1.52 (s, 3H), 1.38 (s, 3H); LCMS *m/z* = 321.09 (M⁺; 100).

(3aR,6R,6aS)-6-(2-Chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[*d*][1,3]dioxole-4-carbaldehyde




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-Dimethyl-4-(4-methyl-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-*d*) δ 9.99 (s, 1H), 8.86 (s, 1H), 7.04 (d, *J* = 3.7 Hz, 1H), 6.75 (dd, *J* = 2.6, 0.9 Hz, 1H), 6.65 (d, *J* = 3.7 Hz, 1H), 6.00 (dt, *J* = 2.7, 1.4 Hz, 1H), 5.77 (dd, *J* = 5.9, 1.5 Hz, 1H), 4.88 (dd, *J* = 5.9, 1.2 Hz, 1H), 1.53 (s, 3H), 1.40 (s, 3H); LCMS *m/z* = 319.90 (M⁺; 100).

2-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[*d*][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.82 (s, 1H), 7.03 (d, *J* = 3.6 Hz, 1H), 6.64 – 6.53 (m, 2H), 5.93 (d, *J* = 2.6 Hz, 1H), 5.81 – 5.69 (m, 2H), 5.60 (dd, *J* = 5.8, 1.4 Hz, 1H), 5.53 – 5.44 (m, 1H), 4.68 (dd, *J* = 5.8, 1.1 Hz, 1H), 1.45 (s, 6H); LCMS *m/z* = 318.15 (M⁺; 100).

1-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-c]pyridine



To a stirred solution of (3aS,4S,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (0.5 g, 1.178 mmol) in THF (7 ml) at 0°C was added 4-chloro-1H-pyrrolo[3,2-c]pyridine (0.305 g, 2.002 mmol), triphenylphosphine (1.081 g, 4.12 mmol) and DIAD (0.801 ml, 4.12 mmol) slowly and stirred for 5min. The reaction mixture was stirred at 25°C for 16h. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 10%) of ethyl acetate in petroleum ether to afford the title compound (0.3g, 45.6%) as an off-white solid. LCMS m/z = 559.23 (M^+ ; 100).


((3aR,6R,6aS)-6-(4-Chloro-1H-pyrrolo[3,2-c]pyridin-1-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol



To a stirred solution of 1-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-c]pyridine (1.3 g, 2.325 mmol) in THF (13ml) was added TBAF (3.25 ml, 3.25 mmol) at 0°C and stirred for 5mins. The reaction mixture was brought to 25°C and stirred for 1h. Volatiles were removed *in vacuo* and the crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 45%) of ethyl acetate in petroleum ether to afford the title compound (0.5g, 67%) as an off-white solid. LCMS m/z = 319.71 ($M-1$; 100).


(3aR,6R,6aS)-6-(4-Chloro-1H-pyrrolo[3,2-c]pyridin-1-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxole-4-carbaldehyde

88




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-Dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-*d*) δ 10.05 (s, 1H), 8.19 (d, *J* = 5.8 Hz, 1H), 7.34 (dd, *J* = 5.9, 0.9 Hz, 1H), 6.99 (d, *J* = 3.3 Hz, 1H), 6.94 – 6.89 (m, 1H), 6.74 (dd, *J* = 3.3, 0.9 Hz, 1H), 5.73 – 5.58 (m, 2H), 4.67 (dt, *J* = 5.9, 1.2 Hz, 1H), 1.55 (s, 3H), 1.39 (s, 3H); LCMS *m/z* = 319.05 (M⁺; 100).

4-Chloro-1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-1H-pyrrolo[3,2-c]pyridine



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.14 (d, *J* = 5.8 Hz, 1H), 7.37 (dd, *J* = 5.9, 0.9 Hz, 1H), 7.06 (d, *J* = 3.3 Hz, 1H), 6.70 – 6.59 (m, 2H), 5.91 – 5.72 (m, 2H), 5.55 – 5.48 (m, 3H), 4.59 – 4.47 (m, 1H), 1.53 (s, 3H), 1.40 (s, 3H); LCMS *m/z* = 317.15 (M⁺; 100).


1-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-1H-pyrrolo[3,2-c]pyridine



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine.


LCMS m/z = 297.21 (M^+ ; 100). Des chloro compound i.e. 1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-1H-pyrrolo[3,2-c]pyridine was also formed, which was separated after Suzuki coupling step (Table-6) by reverse phase preparative HPLC.

5 **(3aR,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one**



To a stirred solution of (3aR,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (20 g, 47.3 mmol) and iodine (14.41 g, 56.8 mmol) in dichloromethane (250 ml), was added pyridine (3.45 ml, 42.6 mmol) under nitrogen atmosphere at 0°C and stirred at 25°C for 4h. The reaction mixture was diluted with methylene chloride (100 ml) and washed with a saturated aqueous sodium thiosulfate (100ml). Layers were separated, the organic layer was washed with brine (100 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 18g of crude compound. This crude residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a rediseq® Rf column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (15 g, 57.8%) as a colorless oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.74 (ddt, J = 17.2, 6.5, 1.6 Hz, 4H), 7.58 – 7.36 (m, 6H), 5.19 (d, J = 5.8 Hz, 1H), 4.79 (t, J = 5.6 Hz, 1H), 4.53 – 4.33 (m, 2H), 1.45 (s, 3H), 1.35 (s, 3H), 1.08 (s, 9H).


(3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol



To a stirred solution of (3aR,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (13.7g, 24.98mmol) in


methanol (130 ml), was added cerium (III) chloride heptahydrate (10.24 g, 27.5 mmol) at 0°C and the reaction mixture was stirred for 30min. Sodium borohydride (0.992 g, 26.2 mmol) was added portion-wise and the resulting mixture was stirred at 0°C for 2h under N₂ atmosphere. The reaction mass was quenched with water (150 ml) and extracted with ethyl acetate (150 ml x 2). The combined organic layer was washed with brine (100 ml), dried over sodium sulfate & concentrated to give 8.5g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisepp® Rf column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (6 g, 43.6%) as a colorless oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.84 – 7.62 (m, 4H), 7.48 – 7.39 (m, 6H), 5.19 (d, *J* = 5.8 Hz, 1H), 4.86 – 4.71 (m, 1H), 4.46 – 4.31 (m, 3H), 2.84 (d, *J* = 10.3 Hz, 1H), 1.45 (s, 3H), 1.38 – 1.33 (s, 3H), 1.08 (s 9H).

7-((3a*S*,4*S*,6a*R*)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-7H-pyrrolo[2,3-*d*]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3a*S*,4*R*,6a*R*)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-*c*]pyridine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.64 (s, 1H), 7.75 (ddt, *J* = 20.3, 6.8, 1.5 Hz, 4H), 7.53 – 7.31 (m, 6H), 6.96 (d, *J* = 3.7 Hz, 1H), 6.67 (d, *J* = 3.6 Hz, 1H), 5.74 – 5.62 (m, 2H), 4.86 – 4.80 (m, 1H), 4.54 – 4.42 (m, 2H), 1.48 (s, 3H), 1.42 (s, 3H), 1.13 (s, 9H). LCMS *m/z* = 686.33 (*M*⁺; 100).

7-((3a*S*,4*S*,6a*R*)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-N-(4-methoxybenzyl)-7H-pyrrolo[2,3-*d*]pyrimidin-4-amine




A mixture of 7-((3aS,4S,6aR)-6-(((tert-butylidiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-7H-pyrrolo[2,3-d]pyrimidine (1.7 g, 2.478 mmol) and (4-methoxyphenyl)methanamine (3.40 g, 24.78 mmol) in ethanol (1.7 ml) was refluxed for 4h. The reaction mixture was quenched with water (150 ml) and extracted with ethyl acetate (150 ml). The combined organic layer was washed with brine (100 ml), dried over sodium sulfate and concentrated to give 8.5g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 40%) of ethyl acetate in petroleum ether to afford the title compound (1.8 g, 92%) as an off white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.37 (d, J = 6.4 Hz, 1H), 7.76 (ddt, J = 19.4, 6.8, 1.4 Hz, 4H), 7.52 – 7.32 (m, 8H), 7.00 – 6.88 (m, 2H), 6.67 (d, J = 3.7 Hz, 1H), 6.42 (d, J = 3.6 Hz, 1H), 5.71 (s, 1H), 5.61 (d, J = 6.2 Hz, 1H), 4.80 (dd, J = 12.7, 5.8 Hz, 3H), 4.60 – 4.37 (m, 2H), 3.84 (s, 3H), 3.77 (s, 1H), 1.46 (s, 3H), 1.41 (s, 3H), 1.13 (s, 9H); LCMS m/z = 786.41 (M⁺; 100).

7-((3aS,4R,6aR)-6-(((tert-Butylidiphenylsilyl)oxy)methyl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-N-(4-methoxybenzyl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine




A mixture of 7-((3aS,4S,6aR)-6-(((tert-butylidiphenylsilyl)oxy)methyl)-5-iodo-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-N-(4-methoxybenzyl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine (3.2 g, 4.07 mmol), 2,4,6-trimethyl-1,3,5,2,4,6-trioxatriborinane (1.021 g, 8.13 mmol) and potassium Carbonate (2.53 g, 18.30 mmol) in DMF (10 ml) was purged with nitrogen for 10 min in a seal tube. Pd(PPh₃)₄ (0.470 g, 0.407 mmol) was added to reaction mixture and stirred at 80°C for 8h. The reaction mixture was diluted with water (50 ml) and extracted with ethyl acetate (50 ml X 2). The combined organic layer was washed with brine (50 ml), dried over sodium sulfate, concentrated *in vacuo* to give 3.1g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 40%) of ethyl acetate in petroleum ether to afford the title compound (2.7 g, 98 %) as an off-white solid. LCMS m/z = 675.60 (M⁺; 100).

((3aS,4R,6aR)-4-(4-((4-Methoxybenzyl)amino)-7H-pyrrolo[2,3-d] pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol



To a stirred solution of 7-((3aS,4R,6aR)-6-(((tert-butyl)diphenylsilyl)oxy)methyl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-N-(4-methoxybenzyl)-7H-pyrrolo [2,3-d]pyrimidin-4-amine (2.7 g, 4.00 mmol) in THF (40 ml) at 0 °C, was added TBAF (4.80 ml, 4.80 mmol) slowly and stirred for 2h at 25°C. The reaction mixture was diluted with water (50ml) and extracted with ethyl acetate (50 ml x 2). The combined organic layer was washed with brine (50ml), dried over sodium sulfate, filtered and concentrated *in vacuo* to give 2.2g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redisep® Rf column with gradient elution (0 to 100%) of ethyl acetate in petroleum ether to afford the title compound (1.7 g., 97 %) as a colorless oil. ¹H NMR (400 MHz, Chloroform-d) δ 7.69 (ddd, J = 12.0, 8.3, 1.4 Hz, 1H), 7.62 – 7.41 (m, 1H), 7.41 – 7.30 (m, 2H), 6.99 – 6.80 (m, 2H), 6.70 (d, J = 3.6 Hz, 1H), 6.35 (d, J = 3.5 Hz, 1H), 5.67 (s, 1H), 5.51 (d, J = 5.9 Hz, 1H), 4.79 (d, J = 5.5 Hz, 2H), 4.60 (dd, J = 5.9, 0.9 Hz, 1H), 4.45 (s, 2H), 3.83 (s, 3H), 1.61 (t, J = 1.1 Hz, 3H), 1.51 (s, 3H), 1.37 (s, 3H). LCMS m/z = 437.17 (M⁺, 100%).


((3aS,4R,6aR)-4-(4-((4-Methoxybenzyl)amino)-7H-pyrrolo[2,3-d] pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl 4-methylbenzeneulfonate



To a mixture of ((3aS,4R,6aR)-4-(4-((4-methoxybenzyl)amino)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol (100mg, 0.229mmol), DMAP (5.60 mg, 0.046 mmol) and TEA (0.096 ml, 0.687 mmol) in CH₂Cl₂ (10 ml) at 0°C, was added a solution of p-toluene sulphonyl chloride (65.5 mg, 0.344 mmol) in CH₂Cl₂ (1ml) and stirred for 1h at 25°C. The reaction

mixture was diluted with water (5 ml) and extracted with ethyl acetate (10 ml x 2). The combined organic layer was washed with brine (10 ml), dried over sodium sulfate, filtered and concentrated *in vacuo* to give 0.2g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redise[®] Rf column with gradient elution (0 to 40%) of ethyl acetate in petroleum ether to afford the title compound (0.08 g, 59.1 %) as a colorless oil. Obtained product was used as such for next step without characterization.

(3aS,4R,6aS)-5-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,4-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol



10


To a stirred solution of (3aS,6aS)-5-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (10.0 g, 23.66 mmol) [synthesized by following the same reaction protocol as was described in *J. Org. Chem.* **2014**, 79, 8059–8066] in THF (100ml), was added 3M methyl magnesium bromide in THF (11.85 ml, 35.5 mmol) at 0°C. The reaction mixture was stirred at 0°C for 1h. The reaction mixture was quenched with a saturated aqueous ammonium chloride (100ml) and extracted with ethyl acetate (200ml x 2). The combined organic layer was washed with brine (50 ml), dried over sodium sulfate, filtered and concentrated *in vacuo* to give 9.8g of crude compound. This residue was purified by combiflash (Rf200, Teledyne/Isco) instrument onto a redise[®] Rf column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (8.50g, 82%) as a colorless oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.72 – 7.67 (m, 4H), 7.47 – 7.37 (m, 6H), 5.84 (q, *J* = 2.0 Hz, 1H), 5.11 – 5.00 (m, 1H), 4.46 – 4.28 (m, 3H), 3.51 (s, 1H), 1.47 (s, 3H), 1.42 (s, 3H), 1.26 (s, 3H), 1.09 (s, 9H).

15

20

(3aS,4S,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol


25



To a stirred solution of (3aS,4R,6aS)-5-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2,4-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (8.0 g, 18.24 mmol) in


anhydrous acetone (80ml), was added p-TsOH·H₂O (0.69 g, 3.65 mmol) at 25°C and stirred for 18h. TLC showed 50% conversion. The reaction mixture was then concentrated *in vacuo*, diluted with methylene chloride (100 ml) and washed with saturated NaHCO₃ (50 ml). Layers were separated, the organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 4.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (3.5g, 43%) as an oil. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.74 – 7.65 (m, 4H), 7.48 – 7.35 (m, 6H), 5.81 (p, *J* = 1.6 Hz, 1H), 4.62 – 4.55 (m, 1H), 4.44 – 4.24 (m, 3H), 1.39 (s, 3H), 1.34 (s, 3H), 1.29 (s, 3H), 1.10 (s, 9H).

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-7H-pyrrolo[2,3-*d*]pyrimidine.



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-*c*] pyridine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.74 (s, 1H), 7.78 – 7.67 (m, 4H), 7.52 – 7.39 (m, 6H), 6.94 (d, *J* = 3.6 Hz, 1H), 6.61 (d, *J* = 3.6 Hz, 1H), 5.94 – 5.86 (m, 1H), 5.80 (q, *J* = 2.4 Hz, 1H), 4.62 – 4.48 (m, 2H), 4.13 (d, *J* = 1.2 Hz, 1H), 1.39 (s, 3H), 1.38 (s, 3H), 1.36 (s, 3H), 1.13 (s, 9H), LCMS *m/z* = 574.44 (*M*⁺; 100%).


((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of ((3aR,6R,6aS)-6-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol.


¹H NMR (400 MHz, Chloroform-d) δ 8.73 (s, 1H), 7.16 (d, J = 3.6 Hz, 1H), 6.62 (d, J = 3.6 Hz, 1H), 5.87 – 5.78 (m, 2H), 4.62 – 4.44 (m, 2H), 4.25 – 4.20 (m, 1H), 1.56 (s, 3H), 1.49 (s, 3H), 1.42 (s, 3H); LCMS m/z = 336.15 (M⁺; 100%).

(3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-d) δ 10.01 (s, 1H), 8.73 (s, 1H), 7.09 (d, J = 3.6 Hz, 1H), 6.78 (dd, J = 2.8, 1.2 Hz, 1H), 6.69 (d, J = 3.6 Hz, 1H), 5.99 (dd, J = 2.8, 0.8 Hz, 1H), 4.34 (t, J = 0.8 Hz, 1H), 1.78 (s, 3H), 1.47 (s, 3H), 1.44 (s, 3H), LCMS m/z = 334.22 (M+1; 100%).

4-Chloro-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-d) δ 8.74 (s, 1H), 7.16 (d, J = 3.6 Hz, 1H), 6.62 (d, J = 3.6 Hz, 1H), 6.49 (dd, J = 17.6, 11.2 Hz, 1H), 5.94 – 5.87 (m, 1H), 5.85 – 5.76 (m, 2H), 5.47 (dd, J = 11.2, 1.2 Hz, 1H), 4.15 (d, J = 1.1 Hz, 1H), 1.60 (s, 3H), 1.44 (s, 6H), LCMS m/z = 332.22 (M+1; 100%).

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.86 (s, 1H), 7.73 (ddt, *J* = 15.2, 6.7, 1.5 Hz, 4H), 7.52 – 7.39 (m, 6H), 6.90 (d, *J* = 3.6 Hz, 1H), 6.55 (dd, *J* = 3.7, 2.3 Hz, 1H), 5.90 (dt, *J* = 2.9, 1.5 Hz, 1H), 5.82 (q, *J* = 2.4 Hz, 1H), 4.54 (dq, *J* = 6.2, 1.8 Hz, 2H), 4.14 (dd, *J* = 6.7, 2.1 Hz, 1H), 1.64 (s, 6H), 1.39 (s, 3H), 1.36 (d, *J* = 2.1 Hz, 3H), 1.12 (s, 9H); LCMS *m/z* = 554.32 (M⁺, 100%).


((3aS,4R,6aR)-2,2,6a-Trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methanol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of ((3aR,6R,6aS)-6-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol.


¹H NMR (400 MHz, Chloroform-*d*) δ 8.86 (s, 1H), 7.08 (d, *J* = 3.7 Hz, 1H), 6.57 (dd, *J* = 3.6, 1.1 Hz, 1H), 5.86 – 5.78 (m, 2H), 4.59 – 4.47 (m, 2H), 4.23 (dt, *J* = 6.9, 0.8 Hz, 1H), 2.77 (s, 3H), 1.56 (s, 3H), 1.50 (s, 3H), 1.42 (s, 3H); LCMS *m/z* = 316.21 (M⁺, 100%).

(3aS,4R,6aR)-2,2,6a-Trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3a*S*,4*R*,6a*R*)-2,2-dimethyl-4-(4-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-*d*) δ 10.01 (s, 1H), 8.87 (s, 1H), 7.03 (d, *J* = 3.7 Hz, 1H), 6.79 (td, *J* = 2.8, 1.1 Hz, 1H), 6.67 (d, *J* = 3.7 Hz, 1H), 6.02 (dd, *J* = 2.9, 0.7 Hz, 1H), 4.33 (d, *J* = 0.9 Hz, 1H), 2.82 (s, 3H), 1.78 (s, 3H), 1.47 (s, 3H), 1.44 (s, 3H); LCMS *m/z*=314.28 (M⁺, 100%).

(4-Methyl-7-((3a*S*,4*R*,6a*R*)-2,2,6a-trimethyl-6-vinyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3] dioxol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3] dioxol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.87 (s, 1H), 7.09 (d, *J* = 3.7 Hz, 1H), 6.60 – 6.44 (m, 2H), 5.94 – 5.75 (m, 3H), 5.45 (dd, *J* = 11.3, 1.5 Hz, 1H), 4.21 – 4.07 (m, 1H), 2.78 (s, 3H), 2.64 (s, 3H), 1.44 (s, 6H); LCMS *m/z*=312.28 (M⁺, 100%).

1-((3a*R*,6*R*,6a*S*)-6-(4-Chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2,3a-trimethyl-6,6a-di hydro-3a*H*-cyclopenta[*d*][1,3]dioxol-4-yl)ethanol




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3a*S*,4*R*,6a*R*)-4-(4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3]dioxol-6-yl)ethan-1-ol. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.73 (d, *J* = 2.0 Hz, 1H), 7.12 (dd, *J* = 31.8, 3.6 Hz, 1H), 6.62 (t, *J* = 3.3 Hz, 1H), 5.87 – 5.76 (m, 2H), 4.78 – 4.68 (m, 2H), 4.26 – 4.20 (m, 1H), 1.61 (d, *J* = 7.1 Hz, 3H), 1.54 (s, 6H), 1.43 (d, *J* = 3.5 Hz, 3H); LCMS *m/z*=350.22 (M⁺, 100%).

1-((3aR,6R,6aS)-6-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,3a-trimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)ethanone




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-one. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.74 (s, 1H), 7.10 (d, *J* = 3.7 Hz, 1H), 6.74 – 6.56 (m, 2H), 5.95 (d, *J* = 2.9 Hz, 1H), 4.27 (t, *J* = 0.9 Hz, 1H), 2.49 (s, 3H), 1.76 (s, 3H), 1.48 (s, 6H); LCMS *m/z*=348.16 (M⁺, 100%).

2-((3aR,6R,6aS)-6-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,3a-trimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)propan-2-ol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propan-2-ol. LCMS *m/z*=364.22 (M⁺, 100%).


4-Chloro-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H

NMR (400 MHz, Chloroform-*d*) δ 8.74 (s, 1H), 7.15 (d, J = 3.6 Hz, 1H), 6.62 (d, J = 3.7 Hz, 1H), 5.84 (d, J = 3.0 Hz, 1H), 5.76 (ddd, J = 12.2, 2.4, 1.0 Hz, 2H), 5.31 (t, J = 1.6 Hz, 1H), 4.15 – 4.13 (m, 1H), 2.05 (t, J = 1.0 Hz, 3H), 1.61 (s, 3H), 1.43 (s, 6H); LCMS m/z = 346.02 (M^+ , 100%).

5 **4-Methyl-7-((3a*S*,4*R*,6a*R*)-2,2,6a-trimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine**




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-*d*]pyrimidine. ¹H
 10 NMR (400 MHz, Chloroform-*d*) δ 8.87 (s, 1H), 7.06 (d, J = 3.6 Hz, 1H), 6.56 (d, J = 3.6 Hz, 1H), 5.89 – 5.67 (m, 3H), 5.29 (t, J = 1.6 Hz, 1H), 4.14 (d, J = 1.3 Hz, 1H), 2.76 (s, 3H), 2.05 (t, J = 1.0 Hz, 3H), 1.61 (s, 3H), 1.43 (s, 6H); LCMS m/z = 326.28 (M^+ , 100%).
 7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidine
 15



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3a*S*,4*R*,6a*R*)-2,2-dimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-*d*]pyrimidine.
 20 LCMS m/z = 364.22 (M^+ , 100%).


(3a*S*,4*R*,6a*S*)-5-(((tert-Butyldiphenylsilyl)oxy)methyl)-4-ethyl-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-ol



100


The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aS)-5-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,4-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.76 – 7.74 (m, 2H), 7.73 – 7.68 (m, 4H), 7.44 – 7.39 (m, 8H), 5.95 (d, *J* = 1.9 Hz, 1H), 5.01 (ddt, *J* = 5.5, 2.9, 1.5 Hz, 1H), 4.43 – 4.36 (m, 2H), 1.49 (s, 3H), 1.43 (s, 3H), 1.10 (s, 9H), 0.74 (t, *J* = 7.5 Hz, 3H).

(3aS,4S,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-6a-ethyl-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-ol




10 The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4S,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.72 – 7.67 (m, 4H), 7.41 (dddt, *J* = 15.6, 7.9, 6.7, 2.1 Hz, 6H), 5.92 (p, *J* = 1.7 Hz, 1H), 4.52 (s, 1H), 4.43 – 4.35 (m, 2H), 4.22 (ddd, *J* = 15.6, 3.0, 2.0 Hz, 1H), 2.71 (s, 1H), 1.79 – 1.64 (m, 2H), 1.39 (s, 3H), 1.32 (s, 3H), 1.10 (s, 9H), 0.74 (t, *J* = 7.6 Hz, 3H).

7-((3aS,4R,6aR)-6-(((tert-Butyldiphenylsilyl)oxy)methyl)-6a-ethyl-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-7H-pyrrolo[2,3-*d*]pyrimidine




20 The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3aS,4R,6aR)-6-(((tert-butylidiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-*c*] pyridine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.76 (s, 1H), 7.79 – 7.67 (m, 4H), 7.55 – 7.37 (m, 6H), 6.99 (d, *J* = 3.6 Hz, 1H), 6.60 (d, *J* = 3.6 Hz, 1H), 6.03 – 5.96 (m, 1H), 5.78 (q, *J* = 2.5 Hz, 1H), 4.61 – 4.43 (m, 2H), 4.25 (d, *J* = 0.9 Hz, 1H), 1.75 (q, *J* = 7.3 Hz, 1H), 1.55 (dq, *J* = 14.8, 7.4 Hz, 1H), 1.41 (s, 6H), 1.13 (s, 9H), 0.68 (t, *J* = 7.4 Hz, 3H); LCMS *m/z* = 588.21 (*M*⁺, 100%).

((3aR,6R,6aS)-6-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a-ethyl-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of ((3aR,6R,6aS)-6-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.74 (s, 1H), 7.17 (d, *J* = 3.7 Hz, 1H), 6.60 (d, *J* = 3.6 Hz, 1H), 5.93 (dq, *J* = 2.7, 1.5 Hz, 1H), 5.79 (q, *J* = 2.2 Hz, 1H), 4.60 – 4.43 (m, 2H), 4.32 (t, *J* = 0.8 Hz, 1H), 1.94 (dq, *J* = 14.8, 7.4 Hz, 1H), 1.74 (dq, *J* = 14.7, 7.5 Hz, 1H), 1.51 (s, 3H), 1.41 (s, 3H), 0.83 (t, *J* = 7.5 Hz, 3H); LCMS *m/z* = 350.22 (*M*⁺, 100%).

(3aR,6R,6aS)-6-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a-ethyl-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxole-4-carbaldehyde



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ¹H NMR (400 MHz, Chloroform-*d*) δ 10.02 (s, 1H), 8.75 (s, 1H), 7.09 (d, *J* = 3.7 Hz, 1H), 6.88 (dd, *J* = 2.8, 1.1 Hz, 1H), 6.68 (d, *J* = 3.6 Hz, 1H), 5.98 (dd, *J* = 2.8, 1.0 Hz, 1H), 4.43 (d, *J* = 1.0 Hz, 1H), 2.24 (dq, *J* = 14.9, 7.5 Hz, 1H), 1.97 (dq, *J* = 14.8, 7.4 Hz, 1H), 1.47 (s, 6H), 0.89 (t, *J* = 7.5 Hz, 3H); LCMS *m/z* = 348.22 (*M*⁺, 100%).


4-Chloro-7-((3aS,4R,6aR)-6a-ethyl-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine



102


The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.75 (s, 1H), 7.16 (d, *J* = 3.7 Hz, 1H), 6.60 (d, *J* = 3.7 Hz, 1H), 6.56 – 6.36 (m, 1H), 5.99 – 5.69 (m, 3H), 5.44 (dd, *J* = 11.3, 1.5 Hz, 1H), 4.25 (d, *J* = 0.9 Hz, 1H), 2.15 – 1.99 (m, 1H), 1.76 (dq, *J* = 14.8, 7.5 Hz, 1H), 1.46 (s, 6H), 0.81 (t, *J* = 7.5 Hz, 3H). LCLCMS *m/z* = 346.22 (*M*⁺, 100%).

((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl hexa hydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)methanol



The title compound was prepared by an analogous reaction protocol as described in WO2006/091905 A1.


((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl hexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)methyl 4-methyl benzenesulfonate



To a stirred solution of ((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methanol (2 g, 5.96 mmol) in CH₂Cl₂ (40 ml) at 0°C, was added TEA (2.494 ml, 17.87 mmol), DMAP (0.146 g, 1.191 mmol) and followed by p-TsCl (1.363 g, 7.15 mmol) slowly and stirred for 10 min. The reaction mixture was stirred at 25°C for 16h. The reaction mixture was diluted with methylene chloride (100 ml) and washed with water (100 ml). Layers were separated, organic layer was washed with brine (100 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 12g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 7%) of ethyl acetate in

petroleum ether to afford the title (0.267g, 9.15%) as an off-white solid. LCMS m/z = 490.17 (M^+ , 100%).

(3aR,3bS,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxole-3b-carbaldehyde




5

The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. ^1H NMR (400 MHz, Chloroform- d) δ 9.34 (s, 1H), 8.63 (s, 1H), 7.18 (d, J = 3.6 Hz, 1H), 6.66 (d, J = 3.6 Hz, 1H), 5.91 (dd, J = 7.1, 1.2 Hz, 1H), 5.11 (s, 1H), 4.82 (dd, J = 7.1, 1.6 Hz, 1H), 2.34 (ddd, J = 9.4, 6.1, 1.6 Hz, 1H), 1.89 – 1.77 (m, 2H), 1.58 (s, 3H), 1.30 (s, 3H); LCMS m/z = 333.9 (M^+ , 100%).

10

4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydro cyclopropa [3,4] cyclo penta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine.



15


To a stirred suspension of methyltriphenylphosphonium bromide (24.62 g, 68.9 mmol) in THF (200 ml), was added 1M KHMDS in THF (68.9 ml, 68.9 mmol) at 25°C and stirred for 10 min. The resulting yellow suspension was cooled to 0°C and a solution of (3aR,3bS,4aS, 5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxole-3b-carbaldehyde (9.2 g, 27.6 mmol) in THF (80 ml) was added slowly. The reaction mixture was stirred at the same temperature for 1h. The reaction mixture was quenched with a saturated aq. NH_4Cl (200ml) and extracted with ethyl acetate (200 ml). Layers were separated, organic layer was washed with brine (250 ml) and dried over anhydrous Na_2SO_4 . The organic layer was filtered and concentrated *in vacuo* to give 11g of crude compound. This residue was purified by combiflash (R_f 200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient

20

25

elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (7g, 77%) as a white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.68 (s, 1H), 7.23 (d, *J* = 3.7 Hz, 1H), 6.67 (d, *J* = 3.6 Hz, 1H), 5.86 (dd, *J* = 17.3, 10.6 Hz, 1H), 5.39 – 5.32 (m, 2H), 5.29 (s, 1H), 5.18 (dd, *J* = 10.6, 0.9 Hz, 1H), 4.59 (dd, *J* = 7.1, 1.6 Hz, 1H), 1.77 (ddd, *J* = 9.3, 4.9, 1.6 Hz, 1H), 1.63 (s, 3H), 1.49 (t, *J* = 5.3 Hz, 1H), 1.27 (s, 3H), 1.18 (ddd, *J* = 9.3, 5.6, 1.6 Hz, 1H); LCMS *m/z* = 332.28 (M⁺, 50%).

7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta [1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine.



10


A mixture of 4-chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa [3,4] cyclo penta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine (3g, 9.04 mmol) and aq. ammonia (19.57 ml, 904 mmol) in dioxane (6ml) stirred at 130°C in a steel bomb for 16 h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 4.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 3%) of methanol in dichloromethane to afford the title compound (2.45g, 87%) as a white solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.07 (s, 1H), 7.02 (s, 2H), 6.96 (d, *J* = 3.5 Hz, 1H), 6.62 (d, *J* = 3.5 Hz, 1H), 5.86 (dd, *J* = 17.4, 10.7 Hz, 1H), 5.33 (dd, *J* = 7.2, 1.3 Hz, 1H), 5.23 (dd, *J* = 17.4, 1.3 Hz, 1H), 5.10 – 5.01 (m, 2H), 4.50 (dd, *J* = 7.1, 1.6 Hz, 1H), 1.70 (ddd, *J* = 9.3, 4.8, 1.6 Hz, 1H), 1.46 (s, 3H), 1.29 – 1.22 (m, 1H), 1.19 (s, 3H), 1.10 (ddd, *J* = 9.1, 5.1, 1.5 Hz, 1H); LCMS *m/z* = 313 (M+1, 100%).

25

7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta [1,2-d][1,3]dioxol-5-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine


105



To a degassed mixture dioxane (8ml) and water (1ml) in a microwave vial was added 4-chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine (1.00 g, 3.01 mmol, potassium phosphate, tribasic (1.575 g, 9.04 mmol), dichloro[1,1'-bis(di-t-butylphosphino)ferrocene] palladium(II) (0.196 g, 0.301 mmol) and 2,4,6-trimethyl-1,3,5,2,4,6-trioxatriborinane (2.107 ml, 15.07 mmol) at 25°C. Stirred the reaction mixture at 100°C for 1h. The reaction mixture was diluted with ethyl acetate (20ml) and washed with water (20ml). Layers were separated, organic layer was washed with brine (20ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (0.66g, 70.3%) as an off white solid. ¹H NMR (400 MHz, Chloroform-d) δ 8.81 (s, 1H), 7.16 (d, *J* = 3.6 Hz, 1H), 6.63 (d, *J* = 3.6 Hz, 1H), 5.87 (dd, *J* = 17.3, 10.6 Hz, 1H), 5.37 – 5.31 (m, 3H), 5.17 (dd, *J* = 10.6, 0.9 Hz, 1H), 4.58 (dd, *J* = 7.1, 1.6 Hz, 1H), 2.78 (s, 3H), 1.78 (ddt, *J* = 9.3, 4.9, 1.8 Hz, 1H), 1.59 (d, *J* = 1.8 Hz, 3H), 1.50 (td, *J* = 5.2, 2.7 Hz, 1H), 1.27 (d, *J* = 2.2 Hz, 3H), 1.19 – 1.14 (m, 1H); LCMS *m/z* = 312.21 (M+1, 100%).

Des chloro compound i.e. 7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine was also formed, which was separated at final step by reverse phase preparative HPLC.


-(S)-1-((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethan-1-ol



To a solution of (3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxole-3b(3aH)-carbaldehyde (2.0 g, 5.99 mmol) in tetrahydrofuran (20 ml), was added 1M methyl magnesium bromide in THF (3.00 ml, 8.99 mmol) at 0°C. The reaction mixture was then stirred at 0°C for 2h.


- 5 The reaction mixture was quenched with a sat. aq. ammonium chloride (50ml) and extracted with ethyl acetate (50ml X 2). The combined organic layer was washed with brine (50ml), dried over sodium sulfate, filtered and concentrated *in vacuo* to give 2.2g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (1.85g, 88%) as an off-white solid. LCMS m/z = 350.03 (M+1; 100%).

1-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethan-1-one



- 15 The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. LCMS m/z = 348.03 (M⁺; 100%).


4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydro cyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine.



- To a stirred suspension of methyltriphenylphosphonium bromide (3.85 g, 10.78 mmol) in THF (40 ml) was added 1M KHMDS in THF (10.78 ml, 10.78 mmol) at 0°C and stirred for 10min. To this yellow suspension was added a solution of 1-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetra hydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethan-1-one (1.5 g, 4.31 mmol) in THF (20 ml)


was added slowly at 0°C. The reaction mixture was stirred at 25°C for 1h. The reaction mixture was quenched with a saturated aqueous ammonium chloride (50ml) and extracted with ethyl acetate (50 ml). Layers were separated, organic layer was washed with brine (50 ml) and dried over anhydrous sodium sulphate. The organic layer was filtered and concentrated *in vacuo* to give 2.2g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisept[®] R_f column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (0.7g, 46.9%) as a white solid. LCMS m/z = 346.03 (M+1; 100%).

(R)-((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl) (cyclopropyl)methanol



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (S)-1-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethan-1-ol. LCMS m/z = 375.91 (M+; 100%).


((3aR,3bS,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl) (cyclopropyl)methanone



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethan-1-one. ¹H NMR (400 MHz, DMSO-d₆) δ 8.58 (s, 1H), 7.71 (d, *J* = 3.7 Hz, 1H), 6.72 (d, *J* = 3.6 Hz, 1H), 5.84 (dd, *J* = 7.3, 1.2 Hz, 1H), 5.21 (s, 1H), 4.79 (dd, *J*


= 7.3, 1.5 Hz, 1H), 2.09 (s, 1H), 2.07 – 2.01 (m, 1H), 1.79 (ddd, $J = 9.5, 5.3, 1.4$ Hz, 1H), 1.51 (t, $J = 5.5$ Hz, 1H), 1.48 (s, 3H), 1.21 (s, 3H), 0.96 – 0.74 (m, 4H); LCMS $m/z = 373.97$ (M^+ ; 100%).

**4-Chloro-7-((3aR,3bR,4aS,5R,5aS)-3b-(1-cyclopropylvinyl)-2,2-dimethylhexahydro
5 cyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine**



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydro cyclopropa[3,4] cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine. ^1H NMR (400 MHz, Chloroform- d) δ 8.67 (s, 1H), 7.29 (d, $J = 3.6$ Hz, 1H), 6.68 (d, $J = 3.6$ Hz, 1H), 5.37 – 5.35 (m, 2H), 4.98 (d, $J = 0.8$ Hz, 1H), 4.80 (s, 1H), 4.59 (dd, $J = 7.5, 1.6$ Hz, 1H), 2.04 (ddd, $J = 9.3, 4.7, 1.6$ Hz, 1H), 1.67 (s, 2H), 1.42 (t, $J = 5.1$ Hz, 1H), 1.27 (s, 6H), 0.75 – 0.69 (m, 2H), 0.56 – 0.52 (m, 2H); LCMS $m/z = 372.2$ (M^+ ; 100%).


**4-Chloro-7-((3aR,3bR,4aS,5R,5aS)-3b-((E)-2-iodoprop-1-en-1-yl)-2,2-dimethylhexa
15 hydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine.**



To a suspension of (1-iodoethyl)triphenylphosphonium bromide (4.08 g, 7.49 mmol, synthesized by following same reaction protocol as was described in WO2004/9574, A1) in THF (20 ml)), was added NaHMDS (7.49 ml, 7.49 mmol) at -25°C . The resulting red coloured solution was stirred at -25°C for 10min. A solution of (3aR,3bS,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydro cyclopropa[3,4]cyclopenta [1,2-d][1,3]dioxole-3b(3aH)-carbaldehyde (1.00 g, 3.00 mmol) in THF (20 ml) was added at -30°C and stirred for 30mins. The reaction mixture was quenched with sat


aq.NH₄Cl (20 ml) and extracted with ethyl acetate (20 ml). Layers were separated, organic layer was washed with brine (20 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.04g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with
 5 gradient elution (0 to 10%) of ethyl acetate in petroleum ether to afford the title compound (0.79g, 55.9%) as an off-white solid. LCMS m/z = 471.80 (M⁺; 100%).

7-((3aR,3bR,4aS,5R,5aS)-3b-(((tert-Butyldiphenylsilyl)oxy)methyl)-2,2-dimethylhexahydro cyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-4-chloro-6-methyl-7H-pyrrolo[2,3-d]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 1-((3aS,4R,6aR)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-2,2-dimethyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-chloro-1H-pyrrolo[3,2-c] pyridine. LCMS m/z = 587.82(M⁺; 100%).

15 ((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methanol




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of ((3aR,6R,6aS)-6-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol. ¹H NMR (400 MHz, DMSO-d₆) δ 8.57 (s, 1H), 6.48 (d, *J* = 1.1 Hz, 1H), 5.34 – 5.25 (m, 1H), 4.98 (s, 1H), 4.90 (dt, *J* = 7.4, 1.3 Hz, 1H), 4.66 (t, *J* = 5.7 Hz, 1H), 3.65 (q, *J* = 5.8, 5.3 Hz, 2H), 2.57 (s, 3H), 1.54 – 1.48 (m, 1H), 1.46 (s, 3H), 1.19 (s, 3H), 0.92 (ddd, *J* = 9.0, 5.0, 1.3 Hz, 1H), 0.85 (q, *J* = 4.7, 4.3 Hz, 1H); LCMS m/z = 350.03 (M⁺; 100%).

(3aR,3bS,4aS,5R,5aS)-5-(4-Chloro-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxole-3b(3aH)-carbaldehyde




5 The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of (3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxole-6-carbaldehyde. LCMS $m/z = 348.03$ (M^+ ; 100%).

10 4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydro cyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-6-methyl-7H-pyrrolo[2,3-d]pyrimidine




The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 4-chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydro cyclopropa [3,4] cyclo penta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine. ^1H NMR (400 MHz, Chloroform- d) δ 8.57 (s, 1H), 6.38 (q, $J = 1.1$ Hz, 1H), 5.88 (dd, $J = 17.3, 10.6$ Hz, 1H), 5.64 (dd, $J = 7.2, 1.4$ Hz, 1H), 5.34 – 5.23 (m, 1H), 5.10 (dd, $J = 10.6, 1.1$ Hz, 1H), 5.02 – 4.96 (m, 1H), 4.87 (s, 1H), 2.55 (s, 3H), 1.59 (s, 3H), 1.56 – 1.51 (m, 1H), 1.30 (s, 3H), 1.29 – 1.26 (m, 1H), 1.19 – 1.12 (m, 1H); LCMS $m/z = 346.03$ (M^+ ; 100%).

20 7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta [1,2-d][1,3]dioxol-5-yl)-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine



The title compound was synthesized by following an analogous reaction protocol as was described in the preparation of 7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta [1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine. ¹H NMR (400 MHz, DMSO-d₆) δ 7.99 (s, 1H), 6.83 (s, 2H), 6.29 (d, *J* = 1.2 Hz, 1H), 5.87 – 5.74 (m, 2H), 5.44 (d, *J* = 7.3 Hz, 1H), 5.19 (dd, *J* = 17.4, 1.4 Hz, 1H), 5.02 (dd, *J* = 10.7, 1.3 Hz, 1H), 4.80 (d, *J* = 7.3 Hz, 1H), 2.36 (s, 3H), 1.61 (dd, *J* = 9.0, 5.3 Hz, 1H), 1.46 (s, 3H), 1.21 (s, 3H), 1.15 – 1.08 (m, 2H); LCMS *m/z* = 327.1 (M+1; 100%).




3-Bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine







4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (0.25g, 0.787mmol) in 9-BBN (0.5 molar, 6.29ml, 3.15mmol) was heated at 70°C for 2h under N₂ atmosphere. The reaction mixture was cooled to 25°C, then potassium phosphate tribasic (0.685g, 3.93mmol) in water (0.5ml) was added and stirred for 20mins. A solution of 3-bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine (0.369g, 0.787mmol) in THF (1ml) was added, followed by PdCl₂(dppf) (0.058g, 0.079mmol). The resulting mixture was stirred at 70°C for 2h. The reaction mixture was diluted with ethyl acetate (10 ml) and washed with water (10 ml). Layers were separated, organic layer was washed with brine (10 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated in vacuo to give 0.35g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 15%) of ethyl acetate in petroleum ether to afford the title compound (0.25g, 48.1%) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.65 (s, 1H), 8.37 (s, 1H), 7.60 (d, *J* = 8.2 Hz, 1H), 7.45 (s, 1H), 7.33 (d, *J* = 8.5 Hz, 2H), 7.28 – 7.14 (m, 2H), 6.98 (d, *J* = 3.7 Hz, 1H), 6.87 – 6.75 (m, 2H), 6.36 (d, *J* = 3.6 Hz, 1H), 5.66 (s, 1H), 5.52 (s, 1H), 5.35 (d, *J* = 5.7 Hz, 1H), 4.63 (d, *J* = 6.0 Hz, 2H), 4.49 (d, *J* = 5.7 Hz, 1H), 3.66 (s, 3H), 3.07 – 2.97 (m, 2H), 2.65 – 2.59 (m, 2H), 1.39 (s, 3H), 1.28 (s, 3H). LCMS *m/z* = 658.89, 661.64 (M-2, M+1, 100%).

Intermediates in table-5 were synthesized by an analogous reaction protocol as was used for the preparation of 3-bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-methoxy benzyl)quinolin-2-amine using the appropriate starting materials.





5 Table-5:



Structure & IUPAC name-	Intermediates used	¹ H NMR & LCMS data
 7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-methylquinolin-2-amine	7-Bromo-N-methyl quinolin-2-amine and 4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	LCMS m/z = 476.05 (M ⁺ , 100%)
 3-Bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3] dioxol-4-yl)-5-fluoro-7H-pyrrolo[2,3-d]pyrimidine and 3-Bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z = 679.85 (M ⁺ , 100%)
 Chemical Formula: C₂₅H₂₃ClFN₅O₂ 7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-	7-Bromo-3-fluoro quinolin-2-amine and 4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d] pyrimidine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.65 (s, 1H), 7.81 (d, <i>J</i> = 11.9 Hz, 1H), 7.60 (d, <i>J</i> = 8.2 Hz, 1H), 7.41 (d, <i>J</i> = 1.6 Hz, 1H), 7.19 (dd, <i>J</i> = 8.2, 1.6 Hz, 1H), 6.95

dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-fluoroquinolin-2-amine		(d, $J = 3.7$ Hz, 1H), 6.84 – 6.67 (m, 2H), 6.41 (d, $J = 3.7$ Hz, 1H), 5.66 (s, 1H), 5.52 (s, 1H), 5.34 (d, $J = 5.7$ Hz, 1H), 4.51 (dd, $J = 15.1, 5.6$ Hz, 1H), 3.18 – 2.98 (m, 2H), 2.80 – 2.56 (m, 2H), 1.39 (s, 3H), 1.28 (s, 3H). LCMS $m/z = 480.2$ (M^+, 100%)
 3-Bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-N-(4-methoxybenzyl)quinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 3-Bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine	LCMS $m/z = 676.47$ ($M+1$, 100%)
 7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-cyclobutylquinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-N-cyclobutylquinolin-2-amine	^1H NMR (400 MHz, Chloroform-d) δ 8.65 (s, 1H), 7.89 (d, $J = 8.9$ Hz, 1H), 7.60 – 7.51 (m, 2H), 7.16 (dd, $J = 8.1, 1.7$ Hz, 1H), 6.65 (d, $J = 8.9$ Hz, 1H), 6.46 (d, $J = 3.7$ Hz, 1H), 6.25 (d, $J = 3.7$ Hz, 1H), 5.77 (s, 1H), 5.44 (t, $J = 1.7$ Hz,


		1H), 5.32 – 5.28 (m, 2H), 4.48 – 4.41 (m, 2H), 3.22 – 3.02 (m, 2H), 2.83 – 2.77 (m, 2H), 2.58 – 2.46 (m, 2H), 2.07 – 1.96 (m, 2H), 1.85 (ddd, $J=10.6$, 9.0, 6.4 Hz, 2H), 1.51 (s, 3H), 1.38 (s, 3H); LCMS $m/z = 515.19$ (M-1, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine & 7-bromo-3-chloroquinolin-2-amine	LCMS $m/z = 476.36$ (M+, 60%)
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)quinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-bromo-3-chloroquinolin-2-amine	LCMS $m/z = 510.2$ (M+, 100%)

115

	 and 4-chloro-7-((3a<i>S</i>,4<i>R</i>,6a<i>R</i>)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4<i>H</i>-cyclopenta[<i>d</i>] [1,3]dioxol-4-yl)-7<i>H</i>-pyrrolo[2,3-<i>d</i>]pyrimidine	¹H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.76 – 8.62 (m, 2H), 7.76 (s, 1H), 7.25 (dd, <i>J</i> = 10.4, 1.5 Hz, 1H), 6.84 (d, <i>J</i> = 3.7 Hz, 1H), 6.54 (d, <i>J</i> = 3.6 Hz, 1H), 5.81 (s, 1H), 5.55 (s, 1H), 5.37 – 5.29 (m, 1H), 4.58 (d, <i>J</i> = 5.7 Hz, 1H), 3.25 – 3.12 (m, 2H), 2.89 – 2.72 (m, 2H), 1.51 (s, 3H), 1.45 (s, 9H), 1.43 (s, 9H), 1.38 (s, 3H); LCMS <i>m/z</i> = 760.41 (<i>M</i>+1, 40%).
	 and 7-((3a<i>S</i>,4<i>R</i>,6a<i>R</i>)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4<i>H</i>-cyclopenta[<i>d</i>][1,3]dioxol-4-yl)-4-methyl-7<i>H</i>-pyrrolo[2,3-<i>d</i>]pyrimidine	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.88 (s, 1H), 8.64 (s, 1H), 7.85 (s, 1H), 7.63 (dd, <i>J</i> = 11.0, 1.5 Hz, 1H), 6.80 (d, <i>J</i> = 3.6 Hz, 1H), 6.48 (d, <i>J</i> = 3.7 Hz, 1H), 5.65 (s, 1H), 5.56 (s, 1H), 5.37 (d, <i>J</i> = 5.6 Hz, 1H), 4.44 (d, <i>J</i> = 5.7 Hz, 1H), 3.22 – 3.13 (m, 2H), 2.78 – 2.66 (m, 2H), 2.61 (s, 3H), 1.38 (s, 9H), 1.35 (s, 3H), 1.33 (s, 9H), 1.27 (s, 3H);

		LCMS m/z = 738.61 (M ⁺ , 90%), 740.61 (M+2, 100%).
	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and	LCMS m/z = 796.20 (M+23, 60%).
	4-Chloro-1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-1H-pyrrolo[3,2-c]pyridine & 7-Bromo-3-methylimidazo[1,2-a]pyridine	LCMS m/z =450.04 (M ⁺ , 100%)

3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine








- 5 4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (0.242g, 0.762mmol) in 9-BBN (0.5 molar, 4.36ml,



2.178mmol) was heated at 50°C for 1h under N₂ atmosphere. The reaction mixture was cooled to 25°C, then potassium phosphate tribasic (0.578g, 2.72mmol) in water (0.5ml) was added and stirred for 20 mins. A solution of 7-bromo-3-chloro-5-fluoroquinolin-2-amine (0.150g, 0.544mmol) in THF (0.5ml) was added, followed by dichloro[1,1'-bis(di-
5 butylphosphino)ferrocene]palladium(II) (0.035g, 0.054mmol). The resulting mixture was stirred at 50°C for 6h. The reaction mixture was diluted with water (10 ml) and extracted with ethyl acetate (10 ml). Layers were separated, organic layer was washed with brine (10 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 0.345g of crude compound. This residue was purified by combiflash (R_f200,
10 Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 35%) of ethyl acetate in petroleum ether to afford the title compound (0.16g, 57.5%) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.64 (s, 1H), 8.18 (s, 1H), 7.25 (s, 1H), 7.09 – 6.96 (m, 4H), 6.44 (d, J = 3.6 Hz, 1H), 5.67 (s, 1H), 5.56 – 5.51 (m, 1H), 5.35 (d, J = 5.7 Hz, 1H), 4.52 (d, J = 5.7 Hz, 1H), 3.05 – 2.98 (m, 2H), 2.74 – 2.56 (m, 2H), 1.38 (s, 3H), 1.28
15 (s, 3H); LCMS m/z = 514.2 (M⁺, 100%).



Intermediates in table-6 were synthesized by an analogous reaction protocol as was used for the preparation of 3-chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine using the appropriate starting materials.



Table-6:



Structure & IUPAC name	Intermediates used 118	¹ H NMR & LCMS data
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-6-fluoroquinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-6-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.65 (s, 1H), 8.18 (s, 1H), 7.56 – 7.44 (m, 2H), 7.14 (d, <i>J</i> = 3.7 Hz, 1H), 6.73 (s, 2H), 6.50 (d, <i>J</i> = 3.7 Hz, 1H), 5.69 (s, 1H), 5.60 – 5.52 (m, 1H), 5.42 – 5.31 (m, 1H), 4.53 (dd, <i>J</i> = 5.7, 2.8 Hz, 1H), 3.08 – 3.01 (m, 2H), 2.66 – 2.62 (m, 2H), 1.39 (s, 3H), 1.28 (s, 3H); LCMS <i>m/z</i> = 515 (M+1, 100%).
 3-(3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine.	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.65 (s, 1H), 8.19 (s, 1H), 7.61 (d, <i>J</i> = 8.2 Hz, 1H), 7.40 (s, 1H), 7.19 (dd, <i>J</i> = 8.1, 1.7 Hz, 1H), 6.96 (d, <i>J</i> = 3.6 Hz, 1H), 6.72 (s, 2H), 6.41 (d, <i>J</i> = 3.6 Hz, 1H), 5.66 (s, 1H), 5.52 (s, 1H), 5.34 (d, <i>J</i> = 5.7 Hz, 1H), 4.49 (d, <i>J</i> = 5.6 Hz, 1H), 3.10 – 2.96 (m, 2H), 2.79 – 2.60 (m, 2H), 1.38 (s, 3H), 1.28 (s, 3H); LCMS <i>m/z</i> = 496.05 (M+, 100%)




 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-5-fluoroquinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.64 (d, <i>J</i> = 6.1 Hz, 1H), 8.22 – 8.16 (m, 1H), 7.29 – 7.14 (m, 2H), 7.00 – 6.95 (m, 3H), 6.56 (d, <i>J</i> = 3.6 Hz, 1H), 5.67 (d, <i>J</i> = 14.7 Hz, 1H), 5.57 (s, 1H), 5.40 – 5.34 (m, 1H), 4.54 (dd, <i>J</i> = 15.1, 5.7 Hz, 1H), 3.04 (q, <i>J</i> = 9.6, 7.8 Hz, 1H), 2.87 (s, 2H), 1.30 (d, <i>J</i> = 9.7 Hz, 3H), 1.19 (d, 6H); LCMS <i>m/z</i> = 528.32 (M⁺, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	7-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.83 (s, 1H), 8.22 (d, <i>J</i> = 0.8 Hz, 1H), 7.34 (s, 1H), 6.92 (dd, <i>J</i> = 10.6, 1.5 Hz, 1H), 6.68 (d, <i>J</i> = 3.6 Hz, 1H), 6.38 (d, <i>J</i> = 3.6 Hz, 1H), 5.80 (s, 1H), 5.52 – 5.47 (m, 2H), 4.55 (d, <i>J</i> = 5.7 Hz, 1H), 3.12 – 3.06 (m, 2H), 2.77 – 2.74 (m, 2H), 2.74 (s, 3H), 1.51 (s, 3H), 1.38 (s, 3H); LCMS <i>m/z</i> = 494.3 (M⁺, 80%)
 2-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-	2-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-	¹H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.76 (s, 1H), 7.98 (s, 1H), 7.55 (d, <i>J</i> = 8.2 Hz, 2H), 7.22 (dd, <i>J</i> =



3-Chloro-7-(2-((3aS,4R,6aR)-4-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloroquinolin-2-amine	8.1, 1.7 Hz, 1H), 6.52 (d, J = 3.6 Hz, 1H), 6.28 (d, J = 3.7 Hz, 1H), 5.75 (s, 1H), 5.42 (d, J = 2.4 Hz, 1H), 5.34 (d, J = 10.9 Hz, 3H), 4.52 (d, J = 5.6 Hz, 1H), 3.18 – 3.07 (m, 2H), 2.81 – 2.76 (m, 2H), 1.49 (s, 3H), 1.38 (s, 3H); LCMS m/z = 496.24 (M^+, 60%)
 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-5-fluoroquinolin-2-amine	7-((3aS,4R,6aR)-2,2-Dimethyl-6-(prop-1-en-2-yl)-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z = 508.31 (M^+, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-	^1H NMR (400 MHz, Chloroform-d) δ 8.93 (d, J = 10.4 Hz, 2H), 8.24 (s, 1H), 7.37 (s, 1H), 6.94 (dd, J = 10.8, 1.4 Hz, 1H), 6.71 (d, J = 3.6 Hz, 1H), 6.37 (d, J = 3.6 Hz, 1H), 5.83 (s, 1H), 5.72 (s, 2H), 5.50 (s, 1H), 5.31 (d, J = 5.7 Hz, 1H), 4.56 (d, J = 5.7 Hz, 1H), 3.16 –



	fluoroquinolin-2-amine	3.04 (m, 2H), 2.78 – 2.76 (m, 2H), 1.52 (s, 3H), 1.38 (s, 3H); LCMS m/z =480.2 (M^+ , 70%)
 3-Chloro-5-fluoro-7-(2-((3aS,4R,6aR)-4-(4-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-isopropyl-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z = 522.32 (M^+ , 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-ethyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-ethyl-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	^1H NMR (400 MHz, DMSO- d_6) δ 8.68 (s, 1H), 8.18 (s, 1H), 7.25 (s, 1H), 7.05 (dd, J = 11.0, 1.4 Hz, 1H), 6.96 (s, 2H), 6.85 (d, J = 3.6 Hz, 1H), 6.49 (d, J = 3.6 Hz, 1H), 5.66 (s, 1H), 5.52 (s, 1H), 5.34 (d, J = 5.7 Hz, 1H), 4.46 (d, J = 5.6 Hz, 1H), 3.05 – 2.93 (m, 4H), 2.66 – 2.59 (m, 2H), 1.38 (s, 3H), 1.31 – 1.24 (m, 6H); LCMS m/z =508.31 (M^+ , 100%)



 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.70 (s, 1H), 8.60 (s, 1H), 8.23 (s, 1H), 8.19 (s, 1H), 7.27 (s, 1H), 7.10 – 6.95 (m, 4H), 6.79 (d, J = 3.7 Hz, 1H), 5.70 (s, 1H), 5.55 (s, 1H), 5.36 (d, J = 5.6 Hz, 1H), 4.50 (d, J = 5.7 Hz, 1H), 3.97 (s, 3H), 3.10 – 2.94 (m, 2H), 2.71 – 2.60 (m, 2H), 1.39 (s, 3H), 1.29 (s, 3H); LCMS m/z = 560.33 (M⁺, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-8-fluoroquinolin-2-amine	4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-8-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.65 (s, 1H), 8.26 (d, J = 1.6 Hz, 1H), 7.62 (s, 1H), 7.45 (d, J = 8.3 Hz, 1H), 7.27 – 7.15 (m, 1H), 7.03 (s, 2H), 6.51 (d, J = 3.7 Hz, 1H), 5.69 (s, 1H), 5.55 (s, 1H), 5.36 (d, J = 5.7 Hz, 1H), 4.54 (d, J = 5.7 Hz, 1H), 3.12 – 3.0 (m, 2H), 2.66 – 2.58 (m, 2H), 1.29 (s, 3H), 1.24 (s, 3H); LCMS m/z = 515.57 (M+1, 40%)


 4-Chloro-7-((3aS,4R,6aR)-6-(2-(3,3-dimethyl-2-(methylthio)-3H-indol-6-yl)ethyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 6-Bromo-3,3-dimethyl-2-(methylthio)-3H-indole	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.67 (s, 1H), 7.42 – 7.28 (m, 2H), 7.17 – 6.99 (m, 2H), 6.51 (d, <i>J</i> = 3.6 Hz, 1H), 5.68 (s, 1H), 5.53 (s, 1H), 5.36 (d, <i>J</i> = 5.7 Hz, 1H), 4.52 (d, <i>J</i> = 5.6 Hz, 1H), 3.07 – 2.85 (m, 2H), 2.71 – 2.56 (m, 5H), 1.38 (s, 3H), 1.29 (d, <i>J</i> = 3.4 Hz, 9H); LCMS <i>m/z</i> = 510.31 (M+1, 60%)
 6'-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-2'-(methylthio)spiro[cyclobutane-1,3'-indole]	4-Chloro-1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-1H-pyrrolo[3,2-c]pyridine & 6'-Bromo-2'-(methylthio)spiro[cyclobutane-1,3'-indole]	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.66 (s, 1H), 7.61 (d, <i>J</i> = 7.5 Hz, 1H), 7.31 (d, <i>J</i> = 1.4 Hz, 1H), 7.09 (dd, <i>J</i> = 7.6, 1.5 Hz, 1H), 7.01 (d, <i>J</i> = 3.7 Hz, 1H), 6.46 (d, <i>J</i> = 3.6 Hz, 1H), 5.67 (s, 1H), 5.53 (s, 1H), 5.35 (d, <i>J</i> = 5.6 Hz, 1H), 4.52 (d, <i>J</i> = 5.6 Hz, 1H), 3.02 – 2.86 (m, 2H), 2.69 – 2.56 (m, 5H), 2.48 – 2.39 (m, 4H), 2.39 – 2.18 (m, 2H), 1.39 (s, 3H), 1.29 (s, 3H); LCMS <i>m/z</i> = 521.19 (M+, 30%)

 3-Chloro-5-fluoro-7-(2-((3aS,4R,6aR)-2,2,6a-trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)quinolin-2-amine	4-methyl-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z =522.32 (M^+, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	4-Cyclopropyl-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z=520.14 (M^+, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	4-Chloro-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-bromo-3-chloro-5-fluoroquinolin-2-amine	^1H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.72 (s, 1H), 8.22 (d, J = 0.8 Hz, 1H), 7.37 (s, 1H), 6.99 (d, J = 3.6 Hz, 1H), 6.96 – 6.89 (m, 1H), 6.55 (d, J = 3.7 Hz, 1H)


d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	¹ H), 5.76 (s, 1H), 5.58 – 5.46 (m, 3H), 4.18 – 4.13 (m, 1H), 3.20 – 3.05 (m, 2H), 2.77 – 2.62 (m, 2H), 1.50 (s, 3H), 1.44 (s, 3H), 1.42 (s, 3H); LCMS m/z =528.19 (M ⁺ , 100%)
 3-Chloro-5-fluoro-7-(2-((3aS,4R,6aR)-2,2,6a-trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	4-Methyl-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.68 (s, 1H), 8.18 (s, 1H), 7.27 (s, 1H), 7.12 – 7.03 (m, 2H), 6.96 (s, 2H), 6.66 – 6.51 (m, 1H), 5.64 (s, 1H), 5.58 (s, 1H), 4.04 (s, 1H), 3.14 – 3.02 (m, 2H), 2.74 – 2.59 (m, 5H), 1.39 (s, 3H), 1.30 (d, <i>J</i> = 3.5 Hz, 6H) ; LCMS m/z =508.31 (M ⁺ , 100%)
 3,5-Dichloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-	4-chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 3,5-	LCMS m/z=532.19 (M ⁺ , 90%)

cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	Dichloro-7-iodoquinolin-2-amine	
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-5-fluoroquinolin-2-amine	4-chloro-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-bromo-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.66 (s, 1H), 8.14 (s, 1H), 7.25 (s, 1H), 7.07 (dd, <i>J</i> = 11.1, 1.3 Hz, 1H), 7.01 (s, 2H), 6.69 (d, <i>J</i> = 3.7 Hz, 1H), 6.35 (d, <i>J</i> = 3.7 Hz, 1H), 5.77 (d, <i>J</i> = 2.7 Hz, 1H), 5.60 (d, <i>J</i> = 2.7 Hz, 1H), 4.01 (s, 1H), 3.10 – 2.93 (m, 2H), 2.92 – 2.80 (m, 1H), 1.37 (s, 3H), 1.27 (s, 3H), 1.25 (d, <i>J</i> = 6.0 Hz, 3H), 1.11 (s, 3H). LCMS <i>m/z</i> = 542.20 (M⁺, 100%)
 3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	1-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-1H-pyrrolo[3,2-c]pyridine, 1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dio	LCMS <i>m/z</i> = 493.36 (M⁺, 100%)

	xol-4-yl)-1H-pyrrolo[3,2-c]pyridine and 7-bromo-3-chloro-5-fluoroquinolin-2-amine	
	1-((3aS,4R,6aR)-2,2-Dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-4-methyl-1H-pyrrolo[3,2-c]pyridine, 1-((3aS,4R,6aR)-2,2-dimethyl-6-vinyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-1H-pyrrolo[3,2-c]pyridine and 7-bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z =479.30 (M⁺, 100%)
 3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-6a-ethyl-2,2-	4-Chloro-7-((3aS,4R,6aR)-6a-ethyl-2,2-dimethyl-6-vinyl-4,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.70 (s, 1H), 8.18 (s, 1H), 7.34 (d, <i>J</i> = 3.7 Hz, 1H), 7.28 (s, 1H), 7.07 (dd, <i>J</i> = 11.1, 1.4 Hz, 1H), 6.96 (s, 2H), 6.60 (d, <i>J</i> = 3.7 Hz, 1H), 5.79 (s, 1H), 5.58 (d, <i>J</i> = 2.3 Hz, 1H), 4.23 (s,

dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3] dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	Bromo-3-chloro-5-fluoroquinolin-2-amine	1H), 3.18 – 2.98 (m, 2H), 2.67 – 2.53 (m, 2H), 1.68 – 1.52 (m, 2H), 1.31 (s, 3H), 1.28 (s, 3H), 0.70 – 0.61 (m, 3H); LCMS m/z = 542.3 (M+, 100%)
 Chemical Formula: C₂₈H₂₉ClFN₅O₂ 3-chloro-5-fluoro-7-(2-((3aS,4R,6aR)-2,2,6a-trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)quinolin-2-amine	4-methyl-7-((3aS,4R,6aR)-2,2,6a-trimethyl-6-(prop-1-en-2-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS m/z = 522.32 (M+, 100%).

7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine




- 5 7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine (1g, 3.20mmol) in 9-BBN (0.5 molar, 25.6 ml, 12.81 mmol) was heated at 60°C for 1h under N₂ atmosphere. The reaction mixture was cooled to 25°C, then



potassium phosphate tribasic (3.40g, 16.01mmol) in water (2 ml) was added and stirred for 30 mins. A solution of 7-bromo-3-chloro-5-fluoroquinolin-2-amine (0.882g, 3.20mmol) in THF (12 ml) was added, followed by PdCl₂(dppf) (0.234g, 0.320mmol). The resulting mixture was stirred at 50°C for 6h. The reaction mixture was diluted with water (10 ml) and extracted with ethyl acetate (10 ml). Layers were separated, organic layer was washed with brine (10 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 5%) of methanol in dichloromethane to afford the title compound (1.3g, 80%) as a pale yellow semisolid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.16 (s, 1H), 8.08 (s, 1H), 7.22 (s, 1H), 7.12 (d, *J* = 3.5 Hz, 1H), 7.05 (d, *J* = 1.4 Hz, 1H), 7.02 (s, 2H), 6.94 (s, 2H), 6.61 (d, *J* = 3.5 Hz, 1H), 5.20 (d, *J* = 7.2 Hz, 1H), 5.01 (s, 1H), 4.52 (dd, *J* = 7.3, 1.5 Hz, 1H), 2.85 – 2.80 (m, 2H), 2.32 – 2.26 (m, 1H), 1.72 – 1.56 (m, 1H), 1.48 (s, 3H), 1.46 – 1.41 (m, 1H), 1.19 (s, 3H), 0.94 – 0.92 (m, 1H), 0.77 – 0.68 (m, 1H); LCMS *m/z* = 509.06 (M⁺, 20%).

Intermediates in table-7 were synthesized by an analogous reaction protocol as was used for the preparation of 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine using the appropriate starting materials and at suitable temperature.


Table-7:




25


Structure & IUPAC name-	Intermediates used	¹ H NMR & LCMS data
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.16 (s, 1H), 8.07 (s, 1H), 7.50 – 7.40 (m, 2H), 7.14 (d, <i>J</i> = 3.5 Hz, 1H), 7.09 – 7.01 (m, 3H),



yl)-2,2-dimethyltetra hydrocyclopropa[3,4]cyclopenta[1,2- d][1,3]dioxol-3b(3aH)-yl)ethyl)-3- chloro-6-fluoroquinolin-2-amine	d][1,3]dioxol-5-yl)-7H- pyrrolo[2,3-d]pyrimidin- 4-amine and 7-Bromo-3- chloro-6-fluoroquinolin- 2-amine	6.67 (s, 1H), 6.62 (dd, J = 6.0, 3.6 Hz, 1H), 5.20 (d, J = 7.2 Hz, 1H), 5.01 (s, 1H), 4.54 (d, J = 7.2 Hz, 1H), 2.97 – 2.76 (m, 2H), 2.30 – 2.20(m, 1H), 1.68 – 1.60 (m, 1H), 1.48 (s, 3H), 1.24 (s, 3H), 1.20 – 1.14 (m, 1H), 0.97 – 0.94 (m, 1H), 0.76 – 0.70 (m, 1H); LCMS m/z =509.06 (M^+ , 30%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4- amino-7H-pyrrolo[2,3-d]pyrimidin-7- yl)-2,2-dimethyltetra hydrocyclopropa[3,4]cyclopenta[1,2- d][1,3]dioxol-3b(3aH)-yl)ethyl)-3- chloroquinolin-2-amine	7- ((3aR,3bS,4aS,5R,5aS)- 2,2-Dimethyl-3b- vinylhexahydrocyclopro pa [3,4]cyclopenta[1,2- d][1,3]dioxol-5-yl)-7H- pyrrolo[2,3-d]pyrimidin- 4-amine and 7-Bromo-3- chloroquinolin-2-amine	^1H NMR (400 MHz, DMSO- d_6) δ 8.15 (s, 1H), 8.08 (s, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.37 (d, J = 1.6 Hz, 1H), 7.20 – 7.10 (m, 2H), 7.01 (s, 2H), 6.73 – 6.57 (m, 3H), 5.21 (dd, J = 7.1, 1.3 Hz, 1H), 5.01 (s, 1H), 4.52 (dd, J = 7.3, 1.5 Hz, 1H), 2.86– 2.81 (m, 2H), 2.32 – 2.21 (m, 1H), 1.69 – 1.65 (m, 1H), 1.48 (s, 3H), 1.47 – 1.42 (m, 1H), 1.20 (s, 3H), 0.97 – 0.91 (m, 1H), 0.80 – 0.72 (m, 1H); LCMS m/z =491.06 (M^+ , 100%)
	7- ((3aR,3bS,4aS,5R,5aS)- 2,2-Dimethyl-3b-	^1H NMR (400 MHz, DMSO- d_6) δ 8.08 (s, 1H), 7.50 (d, J = 8.1 Hz, 1H),




131

7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-methoxyquinolin-2-amine	vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-methoxyquinolin-2-amine	7.40 – 7.24 (m, 2H), 7.18 – 7.05 (m, 2H), 7.01 (s, 2H), 6.61 (d, $J = 3.5$ Hz, 1H), 6.29 (s, 2H), 5.21 (dd, $J = 7.2, 1.3$ Hz, 1H), 5.02 (s, 1H), 4.52 (dd, $J = 7.3, 1.5$ Hz, 1H), 3.89 (s, 3H), 2.88 – 2.74 (m, 2H), 2.30 – 2.18 (m, 1H), 1.72 – 1.60 (m, 1H), 1.48 (s, 3H), 1.46 – 1.41 (m, 1H), 1.20 (s, 3H), 0.98 – 0.91 (m, 1H), 0.79 – 0.73 (m, 1H); LCMS $m/z = 487.2$ (M+1, 50%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-isopropylquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-isopropylquinolin-2-amine	^1H NMR (400 MHz, DMSO-d_6) δ 8.08 (s, 1H), 7.83 (s, 1H), 7.60 (d, $J = 8.1$ Hz, 1H), 7.34 (d, $J = 1.5$ Hz, 1H), 7.14 (dd, $J = 7.6, 2.5$ Hz, 2H), 7.02 (s, 2H), 6.69 – 6.59 (m, 3H), 5.27 – 5.16 (m, 1H), 5.01 (s, 1H), 4.53 (dd, $J = 7.3, 1.5$ Hz, 1H), 3.10 – 3.0 (m, 1H), 2.91 – 2.76 (m, 2H), 2.32 – 2.19 (m, 1H), 1.72 – 1.60 (m, 1H), 1.48 (s, 3H), 1.45 – 1.41 (m, 1H), 1.25 – 1.23 (m, 6H), 1.20 (s, 3H), 0.96 – 0.91 (m, 1H), 0.77 – 0.73



		(m, 1H); LCMS m/z = 499.3 (M+1, 75%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-8-fluoroquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-chloro-8-fluoroquinolin-2-amine	LCMS m/z =509.12 (M+, 60%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-methylquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-methylquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.08 (s, 1H), 7.73 (s, 1H), 7.51 (d, <i>J</i> = 8.1 Hz, 1H), 7.34 (d, <i>J</i> = 1.6 Hz, 1H), 7.13 – 7.07 (m, 2H), 7.01 (s, 2H), 6.61 (d, <i>J</i> = 3.5 Hz, 1H), 6.40 (s, 2H), 5.21 (d, <i>J</i> = 7.2 Hz, 1H), 5.01 (s, 1H), 4.53 (dd, <i>J</i> = 7.2, 1.4 Hz, 1H), 2.90 – 2.76 (m, 2H), 2.31 – 2.20 (m, 1H), 2.21 (s, 3H), 1.72 – 1.61 (m, 1H), 1.48 (s, 3H), 1.47 – 1.41 (m, 1H), 1.23 (s, 3H), 0.97 – 0.91 (m, 1H), 0.80 – 0.73 (m, 1H); LCMS m/z =471.3 (M+1, 90%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-methylquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-methylquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.07 (s, 1H),

7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3] dioxol-3b(3aH)-yl)ethyl)-3-cyclopropylquinolin-2-amine	2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-cyclopropylquinolin-2-amine	7.59 (s, 1H), 7.52 (d, $J = 8.1$ Hz, 1H), 7.32 (d, $J = 1.6$ Hz, 1H), 7.17 – 7.05 (m, 2H), 7.02 (s, 1H), 6.61 (d, $J = 3.5$ Hz, 1H), 6.46 (s, 2H), 5.21 (d, $J = 7.2$ Hz, 1H), 5.01 (s, 1H), 4.52 (dd, $J = 7.3, 1.5$ Hz, 1H), 2.90 – 2.73 (m, 2H), 2.30 – 2.18 (m, 1H), 1.86 – 1.74 (m, 1H), 1.73 – 1.60 (m, 1H), 1.51 – 1.41 (m, 4H), 1.26 – 1.22 (m, 4H), 1.00 – 0.90 (m, 3H), 0.79 – 0.71 (m, 1H), 0.68 – 0.60 (m, 2H); LCMS $m/z = 497.11$ (M+1, 15%)
 2-Amino-7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3] dioxol-3b(3aH)-yl)ethyl)quinoline-3-carbonitrile	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 2-Amino-7-bromoquinoline-3-carbonitrile	^1H NMR (400 MHz, DMSO-d_6) δ 8.61 (s, 1H), 8.07 (s, 1H), 7.66 (d, $J = 8.3$ Hz, 1H), 7.38 (s, 1H), 7.29 – 7.19 (m, 1H), 7.12 (d, $J = 3.5$ Hz, 1H), 7.01 (s, 2H), 6.89 (s, 2H), 6.61 (dd, $J = 8.7, 3.5$ Hz, 1H), 5.21 (d, $J = 7.1$ Hz, 1H), 5.01 (s, 1H), 4.52 (dd, $J = 7.0, 1.6$ Hz, 1H), 2.89 – 2.83 (m, 2H), 2.31 – 2.22 (m, 1H), 1.72 – 1.60 (m, 1H), 1.48 (s, 3H), 1.45 – 1.40 (m, 1H), 1.20 (s, 3H), 0.97 – 0.90 (m, 1H),


		0.80 – 0.60 (m, 1H); LCMS m/z =482.36 (M+1, 15%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-fluoroquinolin-2-amine	7- ((3aR,3bS,4aS,5R,5aS)- 2,2-Dimethyl-3b- vinylhexahydrocyclopro pa [3,4]cyclopenta[1,2- d][1,3]dioxol-5-yl)-7H- pyrrolo[2,3-d]pyrimidin- 4-amine and 7-Bromo-3- fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.07 (s, 1H), 7.77 (d, <i>J</i> = 11.8 Hz, 1H), 7.57 (d, <i>J</i> = 8.2 Hz, 1H), 7.37 (d, <i>J</i> = 1.6 Hz, 1H), 7.20 – 7.09 (m, 2H), 7.01 (s, 2H), 6.71 (s, 2H), 6.60 (d, <i>J</i> = 3.5 Hz, 1H), 5.21 (dd, <i>J</i> = 7.2, 1.3 Hz, 1H), 5.02 (s, 1H), 4.52 (dd, <i>J</i> = 7.4, 1.5 Hz, 1H), 2.87 – 2.77 (m, 2H), 2.31 – 2.20 (m, 1H), 1.70 – 1.60 (m, 1H), 1.48 (s, 3H), 1.46 – 1.43 (m, 1H), 1.20 (s, 3H), 0.96 – 0.92 (m, 1H), 0.82 – 0.69 (m, 1H); LCMS m/z =475.3 (M+1, 100%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	7- ((3aR,3bS,4aS,5R,5aS)- 2,2-Dimethyl-3b- vinylhexahydrocyclopro pa [3,4]cyclopenta[1,2- d][1,3]dioxol-5-yl)-6- methyl-7H-pyrrolo[2,3- d]pyrimidin-4-amine and 7-Bromo-3-chloro- 5-fluoroquinolin-2- amine	LCMS m/z =522.94 (M+, 15%)

 7-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-6-fluoroquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine & 7-Bromo-3-chloro-6-fluoroquinolin-2-amine	LCMS m/z =522.94 (M+, 10%)
 7-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-3b-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)hexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-methylimidazo[1,2-a]pyridine	LCMS m/z=445.3 (M+, 70%)
 7-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-bromo-6-fluoroquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 3-Bromo-6-fluoro-7-iodoquinolin-2-amine	LCMS m/z= 553.20 (M+, 100%)

136

	 7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	LCMS m/z = 755.59 (M+2, 100%).
---	---	--------------------------------

7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-isopropyl quinolin-2-amine



5 7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine (150mg, 0.480mmol) in 9-BBN (0.5molar, 3.84ml, 1.921mmol) was heated at 50°C for 1h under N₂ atmosphere. The reaction mixture was cooled to 25°C, then



10 potassium phosphate tribasic (510 mg, 2.401 mmol) in water (0.5ml) was added and stirred for 30 mins. A solution of 7-bromo-N-isopropylquinolin-2-amine (0.127g, 0.480mmol) in THF (3ml) was added, followed by dichloro[1,1'-bis(di-*t*-butylphosphino)ferrocene]palladium(II) (6.26 mg, 9.60 μmol). The resulting mixture was stirred at 50°C for 6h. The reaction mixture was diluted with water (10 ml) and extracted



15 with ethyl acetate (10 ml). Layers were separated, organic layer was washed with brine (10 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 1.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 10%) of 50% 7N NH₃/MeOH in dichloromethane to afford the title compound (0.12g, 50.1%) as a


20 pale yellow semisolid. LCMS m/z=499.2 (M+1, 40%).

Intermediates in table-8 were synthesized by an analogous reaction protocol as was used for the preparation of 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-isopropylquinolin-2-amine using the appropriate starting materials and at suitable temperature.

Table-8:


Structure & IUPAC name	Intermediates used	¹ H NMR & LCMS data
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-cyclobutylquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-N-cyclobutylquinolin-2-amine	LCMS m/z=511.3 (M+1, 40%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-(1,1-difluoroethyl)quinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-3-(1,1-difluoroethyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.15 (s, 1H), 8.08 (d, <i>J</i> = 1.4 Hz, 1H), 7.69 (d, <i>J</i> = 8.2 Hz, 1H), 7.38 (s, 1H), 7.22 – 7.11 (m, 2H), 7.01 (s, 2H), 6.61 (dd, <i>J</i> = 3.6, 1.4 Hz, 1H), 6.23 (s, 2H), 5.22 (d, <i>J</i> = 7.1 Hz, 1H), 5.02 (s, 1H), 4.53 (d, <i>J</i> = 7.1 Hz, 1H), 2.92 – 2.78 (m, 2H), 2.32 – 2.22 (m, 1H), 2.08 (t, <i>J</i> = 19.1 Hz,

		3H), 1.74 – 1.60 (m, 1H), 1.48 (s, 3H), 1.46 – 1.41 (m, 1H), 1.24 (s, 3H), 0.97 – 0.92 (m, 1H), 0.79 – 0.71 (m, 1H); LCMS m/z = 520.94 (M ⁺ , 50%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-N-(cyclopropylmethyl)quinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 7-Bromo-N-(cyclopropylmethyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 7.78 (d, J = 8.9 Hz, 1H), 7.51 (d, J = 8.1 Hz, 1H), 7.39 (d, J = 1.6 Hz, 1H), 7.19 – 7.05 (m, 3H), 7.02 (s, 2H), 6.73 (d, J = 8.9 Hz, 1H), 6.61 (d, J = 3.6 Hz, 1H), 5.21 (dd, J = 7.2, 1.3 Hz, 1H), 5.02 (s, 1H), 4.52 (dd, J = 7.3, 1.5 Hz, 1H), 3.26 (t, J = 6.1 Hz, 2H), 2.90 – 2.74 (m, 2H), 2.34 – 2.22 (m, 1H), 1.75 – 1.59 (m, 2H), 1.49 (s, 3H), 1.48 – 1.44 (m, 1H), 1.20 (s, 3H), 0.94 (t, J = 4.6 Hz, 1H), 0.77 – 0.70 (m, 1H), 0.51 – 0.43 (m, 2H), 0.29 – 0.22 (m, 2H); LCMS m/z = 511.12 (M+1, 100%)
	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-	¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 7.34 – 7.26 (m, 2H),

7-((3aR,3bR,4aS,5R,5aS)-3b-(2-(3,3-dimethyl-2-(methylthio)-3H-indol-6-yl)ethyl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	vinylhexahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 6-Bromo-3,3-dimethyl-2-(methylthio)-3H-indole	7.12 (d, $J = 3.5$ Hz, 1H), 7.08 – 6.94 (m, 3H), 6.61 (d, $J = 3.5$ Hz, 1H), 5.22 – 5.15 (m, 1H), 5.02 (s, 1H), 4.52 (dd, $J = 7.2$, 1.5 Hz, 1H), 2.81 – 2.69 (m, 2H), 2.59 (s, 3H), 2.29 – 2.16 (m, 1H), 1.68 – 1.56 (m, 1H), 1.48 (s, 3H), 1.45 – 1.41 (m, 1H), 1.27 (s, 6H), 1.20 (s, 3H), 0.94 (t, $J = 4.7$ Hz, 1H), 0.80 – 0.71 (m, 1H); LCMS $m/z = 504.31$ (M+1, 100%)
 6'-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)spiro[cyclobutane-1,3'-indol]-2'-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa [3,4]cyclopenta [1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine and 6'-Bromospiro [cyclobutane-1,3'-indol]-2'-amine.	LCMS $m/z = 485.3$ (M+1, 70%)

3-Bromo-7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine

140






4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine (2.9g, 8.74mmol) in 9-BBN (0.5 molar, 87ml, 43.7mmol) was stirred at 70°C for 1h. The reaction mixture was cooled to 25°C, then potassium phosphate tribasic (5.57g, 26.2mmol) in water (45ml) was added and stirred for 30 mins. A solution of 3-bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine (4.72g, 10.05mmol) in THF (60ml) was added, followed by PdCl₂(dppf)-CH₂Cl₂ (0.357g, 0.437mmol). The resulting mixture was stirred at 70°C for 3h. The reaction mixture was diluted with water (50 ml) and extracted with ethyl acetate (50 ml). Layers were separated, organic layer was washed with brine (50 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 3.5g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 30%) of ethyl acetate in petroleum ether to afford the title compound (3.1 g, 52.5%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.68 (s, 1H), 8.33 (s, 1H), 7.73 (d, *J* = 3.7 Hz, 1H), 7.56 (d, *J* = 8.2 Hz, 1H), 7.43 (d, *J* = 1.6 Hz, 1H), 7.37 – 7.31 (m, 2H), 7.25 – 7.12 (m, 1H), 7.15 (dd, *J* = 8.2, 1.6 Hz, 1H), 6.90 – 6.83 (m, 2H), 6.70 (d, *J* = 3.7 Hz, 1H), 5.27 (dd, *J* = 7.2, 1.3 Hz, 1H), 5.12 (s, 1H), 4.71 – 4.56 (m, 3H), 3.70 (s, 3H), 2.90 – 2.75 (m, 2H), 2.38 – 2.26 (m, 1H), 1.63 – 1.58 (m, 1H), 1.55 – 1.51 (m, 1H), 1.49 (s, 3H), 1.20 (s, 3H), 0.95 (t, *J* = 4.7 Hz, 1H), 0.79 – 0.72 (m, 1H); LCMS *m/z* = 674.1 (M-1, 100%).

Intermediates in table-9 were synthesized by an analogous reaction protocol as was used for the preparation of 3-bromo-7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine using the appropriate starting materials.


Table-9:

Structure & IUPAC name	Intermediates used	¹ H NMR & LCMS data
------------------------	--------------------	--------------------------------

 3-Chloro-7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	4-chloro-7-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-3b-(prop-1-en-2-yl)hexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z =662.10 (M⁺, 100%)
 3-Chloro-7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)-2-cyclopropylethyl)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	4-Chloro-7-((3aR,3bR,4aS,5R,5aS)-3b-(1-cyclopropylvinyl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 8.62 (s, 1H), 8.16 (d, <i>J</i> = 3.6 Hz, 1H), 7.78 – 7.71 (m, 1H), 7.61 (dd, <i>J</i> = 10.1, 5.1 Hz, 1H), 7.40 – 7.30 (m, 3H), 7.09 – 7.03 (m, 1H), 6.86 (dd, <i>J</i> = 8.6, 3.2 Hz, 2H), 6.78 – 6.74 (m, 1H), 5.53 (t, <i>J</i> = 7.2 Hz, 2H), 5.34 (d, <i>J</i> = 7.0 Hz, 1H), 5.16 (s, 1H), 4.63 (d, <i>J</i> = 6.5 Hz, 1H), 3.70 (s, 3H), 2.98 – 2.88 (m, 2H), 1.73 – 1.68 (m, 1H), 1.53 (d, <i>J</i> = 7.4 Hz, 4H), 1.26 – 1.22 (m, 4H), 0.91 –

		0.86 (m, 1H), 0.63 – 0.55 (m, 1H), 0.43 – 0.33 (m, 2H), 0.22 – 0.12 (m, 2H).
 3-Bromo-7-(2-(((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-N-(4-methoxybenzyl)quinolin-2-amine	4-chloro-7-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-3b-(prop-1-en-2-yl)hexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine & 3-Bromo-7-iodo-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z = 687.98 (M-1, 80%); 689.98 (M+1, 100%)

7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-8-fluoroquinolin-2-amine




5 4-chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4] cyclo
penta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine (0.38g, 1.145mmol) in 9-BBN
(0.5 molar, 9.16 ml, 4.58 mmol) was stirred at 70°C for 8h. The reaction mixture was cooled
to 25°C, then potassium phosphate tribasic (0.729g, 3.44mmol) in water (6ml) was added
10 and stirred for 30mins. A solution of 7-bromo-8-fluoroquinolin-2-amine (0.304g,
1.26mmol) in THF (30ml) was added, followed by [1,1'-bis(di-tert-butylphosphino)
ferrocene] dichloropalladium(II) (0.022g, 0.034mmol). The resulting mixture was stirred
at 70°C for 7h. The reaction mixture was diluted with water (20 ml) and extracted with
ethyl acetate (20 ml). Layers were separated, organic layer was washed with brine (20 ml)
15 and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo*



to give 0.5g of crude compound. This residue was purified by combiflash (R_f 200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 50%) of ethyl acetate in petroleum ether to afford the title compound (0.12g, 21.21%) as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.67 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.75 (d, J = 3.6 Hz, 1H), 7.36 (d, J = 8.1 Hz, 1H), 7.14 – 7.05 (m, 1H), 6.79 – 6.70 (m, 2H), 6.64 (s, 2H), 5.27 (d, J = 7.2 Hz, 1H), 5.13 (s, 1H), 4.69 (d, J = 7.2 Hz, 1H), 2.96 – 2.77 (m, 2H), 2.31 – 2.22 (m, 1H), 1.70 – 1.60 (m, 1H), 1.59 – 1.54 (m, 1H), 1.50 (s, 3H), 1.21 (s, 3H), 0.99 (t, J = 4.7 Hz, 1H), 0.83 – 0.77 (m, 1H); LCMS m/z = 494 (M^+ , 100%)


Intermediates in table-10 were synthesized by an analogous reaction protocol as was used for the preparation of 7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-8-fluoroquinolin-2-amine using the appropriate starting materials and at suitable temperature.

Table-10:


15

Structure & IUPAC name	Intermediates used	^1H NMR & LCMS data
 7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-N-(4-	^1H NMR (400 MHz, Chloroform- d) δ 8.68 (s, 1H), 7.82 (d, J = 8.9 Hz, 1H), 7.60 (d, J = 1.5 Hz, 1H), 7.54 (d, J = 8.1 Hz, 1H), 7.41 – 7.33 (m, 2H), 7.21 – 7.14 (m, 2H), 6.97 – 6.84 (m, 2H), 6.65 – 6.57 (m, 2H), 5.23 (dd, J = 7.3, 1.4 Hz, 1H), 5.15 (s, 1H), 4.70 – 4.62 (m, 3H), 3.83 (s, 3H), 3.11 – 2.86 (m, 2H), 2.48 – 2.37 (m,

	methoxybenzyl)quinolin-2-amine	¹ H), 1.83 – 1.76 (m, 1H), 1.55 (s, 3H), 1.50 (dd, <i>J</i> = 4.8, 1.5 Hz, 1H), 1.28 (s, 3H), 1.16 (t, <i>J</i> = 5.0 Hz, 1H), 0.86 - 0.78 (m, 1H); LCMS <i>m/z</i> = 596.20 (M ⁺ , 100%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-methylquinolin-2-amine	4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-N-methylquinolin-2-amine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.69 (s, 1H), 7.77 (d, <i>J</i> = 8.9 Hz, 1H), 7.74 (d, <i>J</i> = 3.7 Hz, 1H), 7.51 (d, <i>J</i> = 8.1 Hz, 1H), 7.41 (d, <i>J</i> = 1.6 Hz, 1H), 7.07 (dd, <i>J</i> = 8.1, 1.7 Hz, 1H), 6.93 (d, <i>J</i> = 5.9 Hz, 1H), 6.71 (d, <i>J</i> = 3.6 Hz, 1H), 6.67 (d, <i>J</i> = 8.9 Hz, 1H), 5.27 (dd, <i>J</i> = 7.2, 1.3 Hz, 1H), 5.13 (s, 1H), 4.67 (dd, <i>J</i> = 7.3, 1.5 Hz, 1H), 2.88 (d, <i>J</i> = 4.7 Hz, 3H), 2.85 – 2.74 (m, 2H), 2.36 – 2.30 (m, 1H), 1.71 – 1.65 (m, 1H), 1.57 – 1.52 (m, 1H), 1.50 (s, 3H), 1.21 (s, 3H), 0.96 (t, <i>J</i> = 4.7 Hz, 1H), 0.82 – 0.76 (m, 1H); LCMS <i>m/z</i> = 490.24 (M ⁺ , 60%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxyphenyl)quinolin-2-amine	4-Chloro-7-((3aR,3bS,4aS,5R,5aS)-2,2-dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-N-(4-methoxyphenyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.04 (s, 1H), 8.68 (s, 1H), 7.79 (s, 1H), 7.74 (d, <i>J</i> = 3.7 Hz, 1H), 7.69 (d, <i>J</i> = 8.2 Hz, 1H), 7.35 (s, 1H), 7.30 (d, <i>J</i> = 8.2 Hz, 2H), 7.17 (dd, <i>J</i> = 8.3, 1.6 Hz, 1H), 6.89 – 6.83 (m, 2H), 6.71 (d, <i>J</i> =

d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinazolin-2-amine	7H-pyrrolo[2,3-d]pyrimidine & 7-Bromo-N-(4-methoxybenzyl)quinazolin-2-amine	3.7 Hz, 1H), 5.27 (d, $J = 7.1$ Hz, 1H), 5.13 (s, 1H), 4.71 – 4.64 (m, 1H), 4.52 (d, $J = 6.3$ Hz, 2H), 3.71 (s, 3H), 2.90 – 2.80 (m, 2H), 2.37 – 2.27 (m, 1H), 1.67 – 1.62 (m, 1H), 1.56 – 1.52 (m, 1H), 1.49 (s, 3H), 1.20 (s, 3H), 0.95 (t, $J = 4.6$ Hz, 1H), 0.79 – 0.74 (m, 1H); LCMS $m/z = 597.3$ (M^+ , 100%)
 3-Chloro-7-(2-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-5-fluoroquinolin-2-amine	7-((3aR,3bS,4aS,5R,5aS)-2,2-Dimethyl-3b-vinylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-4-methyl-7H-pyrrolo[2,3-d]pyrimidine and 7-Bromo-3-chloro-5-fluoroquinolin-2-amine	LCMS $m/z = 508.19$ (M^+ , 80%)


7-(1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-N-methylquinolin-2-amine



- 5 To a stirred solution of 1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-ol (0.2g, 0.596


mmol), 2-(methylamino)quinolin-7-ol (0.145g, 0.834mmol) and triphenylphosphine (0.469g, 1.787mmol) at 0°C was added DEAD (0.283ml, 1.787mmol) slowly and stirred for 30min. The resulting mixture was stirred at 25°C for 1h. The solvent was evaporated under reduced pressure and the residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (0.1g, 34.1%) as an off-white solid. LCMS m/z = 492.2 (M⁺, 100%).

7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-N-methylquinolin-2-amine



To a stirred solution of ((3aR,6R,6aS)-6-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methanol (0.25g, 0.777mmol), 2-(methylamino)quinolin-7-ol (0.338g, 1.942mmol) and triphenylphosphine (0.611g, 2.331mmol) in THF (20ml) was added DEAD (0.369ml, 2.331mmol) dropwise at 0°C. The resulting mixture was stirred at 25°C for 20h. The solvent was evaporated under reduced pressure and the residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 70%) of ethyl acetate in petroleum ether (0.2 g, 53.9 %) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.66 (s, 1H), 7.74 (d, J = 8.8 Hz, 1H), 7.53 (d, J = 1.7 Hz, 1H), 7.51 (d, J = 3.5 Hz, 1H), 7.07 (d, J = 2.5 Hz, 1H), 6.95 (d, J = 5.0 Hz, 1H), 6.85 (dd, J = 8.7, 2.5 Hz, 1H), 6.69 – 6.51 (m, 2H), 5.87 (s, 1H), 5.78 (s, 1H), 5.55 – 5.42 (m, 1H), 5.04 – 4.85 (m, 2H), 4.74 – 4.63 (m, 1H), 2.89 (d, J = 4.7 Hz, 3H), 1.49 (s, 3H), 1.24 (s, 3H); LCMS m/z = 479.3 (M+1, 100%).


3-Bromo-7-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methoxy)-N-(4-methoxybenzyl)quinolin-2-amine



To a stirred solution of ((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-

yl)methanol (0.09g, 0.268mmol), 3-bromo-2-((4-methoxybenzyl)amino)quinolin-7-ol (0.106g, 0.295mmol) and triphenylphosphine (0.176g, 0.670mmol) in THF (8 ml), was added DEAD (0.106ml, 0.670mmol) dropwise at 0°C and stirred for 30 min. The resulting mixture stirred for at 25°C for 14h. The solvent was evaporated under reduced pressure and the residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 60%) of ethyl acetate in petroleum ether (0.12g, 66.1%) as an off-white solid. LCMS m/z= 677.97 (M+1, 100%).


3-Bromo-7-((E)-1-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)prop-1-en-2-yl)-N-(4-methoxybenzyl)quinolin-2-amine



A mixture of 4-chloro-7-((3aR,3bR,4aS,5R,5aS)-3b-((E)-2-iodoprop-1-en-1-yl)-2,2-dimethylhexahydro cyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidine (0.12g, 0.254mmol), triphenylphosphine (6.67mg, 0.025mmol), 3-bromo-N-(4-methoxybenzyl)-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-2-amine (0.179g, 0.382mmol), sodium carbonate (0.054g, 0.509mmol) and Pd(OAc)₂ (2.86mg, 0.013mmol) in DMF (4ml) and water (1 ml) was stirred at 25°C for 16h. The reaction mixture was diluted with water (20 ml) and extracted with ethyl acetate (20 ml). Layers were separated, organic layer was washed with brine (20 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 0.3g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 15%) of ethyl acetate in petroleum ether to afford the title compound (0.13g, 74.4%) as an off-white solid. LCMS m/z = 687.98 (M+1, 100%).


N7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)quinoline-2,7-diamine

148



To a stirred solution of (3a*S*,4*R*,6a*R*)-4-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxole-6-carbaldehyde (0.18g, 0.563mmol) and N2-(4-methoxybenzyl)quinoline-2,7-diamine (0.157g, 0.563mmol) in methanol (5ml) was added acetic acid (0.1mL) at 25°C and stirred for 2h. Sodium cyanoborohydride (0.106g, 1.689mmol) was added stirred for 16h. The reaction was quenched by addition of sat. NH₄Cl (15mL). The reaction mixture was concentrated under reduced pressure and the residue was diluted with ethyl acetate (40mL). The organic phase was washed with water (30ml), saturated aqueous sodium bicarbonate (30ml) and brine (30ml) successively. Dried the organic layer over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give 0.2g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisept® R_f column with gradient elution (0 to 15%) of methanol in dichloromethane to afford the title compound (0.15g, 45.7%) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.66 (s, 1H), 7.59 (d, *J* = 8.8 Hz, 1H), 7.48 (d, *J* = 3.7 Hz, 1H), 7.30 (m, 3H), 7.10 (t, *J* = 5.8 Hz, 1H), 6.92 – 6.86 (m, 2H), 6.65 (dd, *J* = 8.6, 2.3 Hz, 1H), 6.57 (d, *J* = 2.2 Hz, 1H), 6.51 (d, *J* = 3.7 Hz, 1H), 6.45 (d, *J* = 8.7 Hz, 1H), 5.72 (s, 1H), 5.69 (s, 1H), 5.38 (d, *J* = 5.7 Hz, 1H), 4.65 – 4.52 (m, 1H), 4.55 (d, *J* = 5.8 Hz, 2H), 4.13 – 3.93 (m, 3H), 3.72 (s, 3H), 1.48 (s, 3H), 1.30 (s, 3H); LCMS *m/z* = 583.4 (M⁺, 100%).


N7-(((3a*S*,4*R*,6a*R*)-4-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)-N7-methylquinoline-2,7-diamine



To a stirred solution of N2-(4-methoxybenzyl)-N7-methylquinoline-2,7-diamine in DMF (2 ml) was added K₂CO₃ (0.058g, 0.420mmol) at 0°C slowly followed by ((3a*S*,4*R*,6a*R*)-4-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[*d*][1,3] dioxol-6-yl)methyl 4-methylbenzenesulfonate (0.1g, 0.210mmol) and


stirred the reaction mixture for 10 mins. The resulting mixture was stirred at 25°C for 16h. The reaction mixture was diluted with water (20ml) and extracted with ethyl acetate (20ml). Layers were separated, organic layer was washed with brine (20 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 0.3g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 7%) of ethyl acetate in petroleum ether to afford the title compound (0.025g, 19.93%) as a colourless oil. LCMS *m/z* = 597.29 (M⁺, 80%).

7-(((3a*S*,4*R*,6a*R*)-4-(4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3]dioxol-6-yl)methylthio)-*N*-(4-methoxybenzyl)quinolin-2-amine




To a stirred solution of 2-((4-methoxybenzyl)amino)quinoline-7-thiol (125mg, 0.422mmol) in DMSO (3 ml) at 0°C was added Cs₂CO₃ (302mg, 0.928mmol) and ((3a*S*,4*R*,6a*R*)-4-(4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4*H*-cyclopenta[*d*][1,3] dioxol-6-yl)methyl 4-methylbenzenesulfonate (201mg, 0.422mmol). The resulting mixture was stirred at 25°C for 3h. The reaction mixture was diluted with water (20ml) and extracted with ethyl acetate (20ml). Layers were separated, organic layer was washed with brine (20 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 0.3g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 20%) of ethyl acetate in petroleum ether to afford the title compound (0.112g, 44.3%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.60 (s, 1H), 7.85 (d, *J* = 8.9 Hz, 1H), 7.60 – 7.48 (m, 3H), 7.33 – 7.26 (m, 2H), 7.19 (dd, *J* = 8.3, 1.9 Hz, 1H), 6.85 – 6.79 (m, 3H), 6.48 (d, *J* = 3.7 Hz, 1H), 6.06 (d, *J* = 3.6 Hz, 1H), 5.72 (s, 1H), 5.65 (s, 1H), 5.43 (d, *J* = 5.7 Hz, 1H), 4.54 (d, *J* = 5.7 Hz, 2H), 4.48 (d, *J* = 5.7 Hz, 1H), 4.22 (d, *J* = 15.3 Hz, 1H), 3.77 – 3.70 (m, 1H), 3.68 (s, 3H), 1.43 (s, 3H), 1.28 (s, 3H); LCMS *m/z* = 600.21 (M⁺, 100%).

3-Chloro-7-(1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine



- 5 To a stirred solution of 1-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethan-1-ol (0.7g, 2.085mmol), 3-chloro-5-fluoro-2-((4-methoxybenzyl)amino)quinolin-7-ol (0.416g, 1.251mmol) and triphenylphosphine (1.640g, 6.25mmol) in THF (10ml), was added DEAD (0.990 ml, 6.25 mmol) dropwise at 0°C and stirred for 30mins. The resulting mixture stirred
10 at 25°C for 16h. The solvent was evaporated under reduced pressure and this residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisept® R_f column with gradient elution (0 to 15%) of ethyl acetate in petroleum ether to afford the title compound (0.5g, 36.9 % yield) as an off-white solid. LCMS m/z = 650.3 (M⁺, 100%).

- 3-chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine**





- Cesium carbonate (558 mg, 1.713 mmol) was added to a solution of ((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d]
20 [1,3]dioxol-6-yl)methyl 4-methylbenzenesulfonate (272 mg, 0.571 mmol) and 3-chloro-5-fluoro-2-((4-methoxybenzyl)amino)quinolin-7-ol (190 mg, 0.571 mmol) in DMF (4 ml) at 0°C and stirred at 25°C for 1h. Ice cold water (20ml) was added to the reaction mixture and stirred for 10 minutes, precipitated solid was filtered, washed with water and dried under vacuum to afford the title compound (300 mg, 83%) as a brown solid. ¹H NMR (400
25 MHz, DMSO-d₆) δ 8.64 (s, 1H), 8.11 (s, 1H), 7.65 (t, J = 6.1 Hz, 1H), 7.50 (d, J = 3.7 Hz, 1H), 7.36 – 7.30 (m, 2H), 6.94 (d, J = 2.3 Hz, 1H), 6.89 – 6.79 (m, 3H), 6.60 (d, J = 3.7 Hz,


1H), 5.87 (s, 1H), 5.78 (s, 1H), 5.48 (d, J = 5.7 Hz, 1H), 5.05 – 4.89 (m, 2H), 4.70 (d, J = 5.7 Hz, 1H), 4.64 (d, J = 6.1 Hz, 2H), 3.71 (s, 3H), 1.45 (s, 3H), 1.30 (s, 3H); LCMS m/z = 636.34 (M⁺, 100%).

Intermediates in table-11 were synthesized by an analogous reaction protocol as was used

- 5 for the preparation of 3-chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine using the appropriate starting materials and at suitable temperature.


10 Table-11:

Structure & IUPAC name	Intermediates used	¹ H NMR & LCMS data
 3-Chloro-5-fluoro-N-(4-methoxybenzyl)-7-(((3aS,4R,6aR)-4-(4-((4-methoxybenzyl)amino)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)quinolin-2-amine	((3aS,4R,6aR)-4-(4-((4-methoxybenzyl)amino)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl 4-methylbenzenesulfonate & 3-chloro-5-fluoro-2-((4-methoxybenzyl)amino)quinolin-7-ol	¹H NMR & LCMS data LCMS m/z=751.61 (M⁺, 100%).
 3-Chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-	((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl 4-	¹H NMR (400 MHz, DMSO-d₆) δ 8.53 (s, 1H), 8.41 (s, 1H), 7.74 (d, J = 8.6 Hz, 1H), 7.67 (d, J = 1.8 Hz, 1H), 7.43 (dd, J = 8.5, 1.9 Hz, 1H), 7.26 – 7.20 (m,

yl)methyl)thio)-N,N-bis(4-methoxybenzyl)quinolin-2-amine	methylbenzenesulfonate & 2-(bis(4-methoxybenzyl)amino)-3-chloroquinoline-7-thiol	4H), 6.81 – 6.75 (m, 4H), 6.38 (d, J = 3.7 Hz, 1H), 5.98 (d, J = 3.6 Hz, 1H), 5.72 (s, 1H), 5.63 (s, 1H), 5.44 (d, J = 5.7 Hz, 1H), 4.57 – 4.42 (m, 5H), 4.28 (d, J = 15.4 Hz, 1H), 3.82 – 3.76 (m, 1H), 3.64 (s, 6H), 1.42 (s, 3H), 1.28 (s, 3H); LCMS m/z = 754.49 (M ⁺ , 100%)
 7-(((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methyl)thio)-N-(4-methoxybenzyl)quinolin-2-amine	((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methyl 4-methylbenzenesulfonate & 2-((4-methoxybenzyl)amino)quinoline-7-thiol	¹ H NMR (400 MHz, Chloroform-d) δ 8.68 (s, 1H), 7.76 (d, J = 8.9 Hz, 1H), 7.64 (d, J = 1.8 Hz, 1H), 7.48 (d, J = 8.3 Hz, 1H), 7.39 (d, J = 3.6 Hz, 1H), 7.37 – 7.32 (m, 2H), 7.20 (dd, J = 8.4, 1.9 Hz, 1H), 6.93 – 6.87 (m, 2H), 6.59 (d, J = 8.8 Hz, 1H), 6.55 (d, J = 3.6 Hz, 1H), 5.39 (dd, J = 7.2, 1.5 Hz, 1H), 5.21 (s, 1H), 4.65 (d, J = 5.3 Hz, 2H), 4.64 – 4.60 (m, 1H), 3.82 (s, 3H), 3.62 (d, J = 13.3 Hz, 1H), 3.43 (d, J = 13.3 Hz, 1H), 1.76 – 1.73 (m,


		1H), 1.60 (s, 3H), 1.33 – 1.30 (m, 1H), 1.26 (s, 3H), 1.12 – 1.06 (m, 1H); LCMS m/z = 614.21 (M ⁺ , 100%)
--	--	--

3-Chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine



5 To a stirred solution of ((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl) methanol (250 mg, 0.745 mmol) and 3-chloro-5-fluoro-2-((4-methoxybenzyl)amino)quinolin-7-ol (297 mg, 0.893 mmol) in toluene (2 ml) was added triphenyl phosphine (234 mg, 0.893 mmol) and
10 stirred for 5 minutes then added DEAD (0.118 ml, 0.745 mmol) dropwise at 25°C and reaction mixture was stirred at 90°C for 2 hours. The volatiles were removed *in vacuo* and obtained residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 50%) of ethyl acetate in petroleum ether to afford the title compound (420mg, 87%) as a off-white solid. ¹H NMR (400 MHz, DMSO-
15 d₆) δ 8.70 (s, 1H), 8.12 (s, 1H), 7.69 (t, J = 6.1 Hz, 1H), 7.48 (d, J = 3.7 Hz, 1H), 7.42 – 7.36 (m, 1H), 7.36 – 7.30 (m, 2H), 6.97 (d, J = 2.2 Hz, 1H), 6.88 – 6.86 (m, 2H), 6.62 (d, J = 3.6 Hz, 1H), 5.86 (d, J = 2.8 Hz, 1H), 5.68 (d, J = 2.5 Hz, 1H), 5.08 (d, J = 15.5 Hz, 1H), 4.89 (d, J = 15.5 Hz, 1H), 4.65 (d, J = 6.0 Hz, 2H), 4.25 (s, 1H), 3.70 (s, 3H), 1.55 (s, 3H), 1.41 (s, 3H), 1.34 (s, 3H); LCMS m/z = 650.34 (M⁺, 100%).


20 **7-(2-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-methylquinolin-2-amine**









A mixture of 7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-methylquinolin-2-amine (0.100 g, 0.210 mmol) in dioxane (3 ml) was added aq. ammonia (0.227 ml, 10.50 mmol) at 25°C and stirred the reaction mixture at 130°C for 16h. The reaction mixture was diluted with brine (20ml) and extracted with ethyl acetate (20ml). Layers were separated, the organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give 0.15g of crude compound. The obtained residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a rediseq® R_f column with gradient elution (0 to 5%) of methanol in dichloromethane to afford the title compound (0.07g, 73%) as an off-white solid. LCMS m/z= 457.2 (M+1; 40%).




Intermediates in table-12 were synthesized by an analogous reaction protocol as was used for the preparation of 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-methylquinolin-2-amine using the appropriate starting materials (Instead of aq.NH₃, 7N NH₃ in MeOH could also be used).




Table-12




Structure & IUPAC name	Intermediate used	¹ H NMR & LCMS data
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-cyclobutylquinolin-2-amine	7-(2-((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-cyclobutylquinolin-2-amine	¹ H NMR (400 MHz, Chloroform-d) δ 8.35 (s, 1H), 7.86 (d, J = 8.9 Hz, 1H), 7.58 – 7.53 (m, 2H), 7.15 (dd, J = 8.2, 1.7 Hz, 1H), 6.62 (d, J = 8.9 Hz, 1H), 6.31 (d, J = 3.6 Hz, 1H), 6.09 (d, J = 3.6 Hz, 1H), 5.75 (s, 1H), 5.51 – 5.45 (m, 2H), 5.26 (d, J = 6.0 Hz, 1H), 5.15 (s, 2H), 4.47 (d, J = 5.7 Hz, 1H), 4.45 – 4.37 (m, 1H), 3.19 – 3.01 (m, 2H), 2.87 – 2.68 (m, 2H), 2.57 – 2.46 (m, 2H), 2.02 – 1.93 (m, 2H), 1.89 – 1.79 (m, 2H), 1.50 (s, 3H),



		1.37 (s, 3H); LCMS m/z= 497.3 (M+1; 70%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-fluoroquinolin-2-amine	7-(2-((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.05 (s, 1H), 7.79 (d, J = 11.8 Hz, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.41 (s, 1H), 7.19 (d, J = 8.2 Hz, 1H), 6.96 (s, 2H), 6.72 (s, 2H), 6.35 – 6.31 (m, 2H), 5.54 (s, 1H), 5.50 (s, 1H), 5.29 (d, J = 5.9 Hz, 1H), 4.35 (d, J = 5.7 Hz, 1H), 3.10 – 2.93 (m, 2H), 2.70 – 2.60 (m, 2H), 1.37 (s, 3H), 1.24 (s, 3H); LCMS m/z= 461.3 (M+1; 60%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.17 (s, 1H), 8.05 (s, 1H), 7.61 (d, J = 8.3 Hz, 1H), 7.41 (d, J = 1.6 Hz, 1H), 7.19 (dd, J = 8.2, 1.7 Hz, 1H), 6.98 (s, 2H), 6.69 (s, 2H), 6.41 (d, J = 3.6 Hz, 1H), 6.35 (d, J = 3.5 Hz, 1H), 5.55 (s, 1H), 5.51 (s, 1H), 5.29 (d, J = 5.7 Hz, 1H), 4.36 (d, J = 5.7 Hz, 1H), 3.09 – 2.94 (m, 2H), 2.72 – 2.56 (m, 2H), 1.37 (s, 3H), 1.28 (s, 3H); LCMS m/z= 477.05 (M ⁺ ; 20%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromoquinolin-2-amine	3-Bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.35 (s, 1H), 8.07 (s, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.45 (s, 1H), 7.38 – 7.31 (m, 2H), 7.22 – 7.16 (m, 2H), 7.01 (s, 2H), 6.88 – 6.82 (m, 2H), 6.52 (d, J = 3.5 Hz, 1H), 6.40




cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	(d, J = 3.6 Hz, 1H), 5.56 (s, 1H), 5.52 (s, 1H), 5.30 (d, J = 5.7 Hz, 1H), 4.66 – 4.61 (m, 2H), 4.38 (d, J = 5.6 Hz, 1H), 3.69 (s, 3H), 3.06 – 2.93 (m, 2H), 2.70 – 2.59 (m, 2H), 1.38 (s, 3H), 1.27 (s, 3H); LCMS m/z = 641.97 (M ⁺ ; 60%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.17 (d, J = 0.7 Hz, 1H), 8.05 (s, 1H), 7.26 (s, 1H), 7.05 (dd, J = 11.1, 1.4 Hz, 1H), 7.01 – 6.91 (m, 4H), 6.41 (d, J = 3.5 Hz, 1H), 6.35 (d, J = 3.5 Hz, 1H), 5.54 (s, 1H), 5.51 (s, 1H), 5.30 (d, J = 5.7 Hz, 1H), 4.37 (d, J = 5.7 Hz, 1H), 3.07 – 2.95 (m, 2H), 2.71 – 2.56 (m, 2H), 1.37 (s, 3H), 1.27 (s, 3H); LCMS m/z = 495.05 (M ⁺ ; 40%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-5-fluoroquinolin-2-amine		¹ H NMR (400 MHz, DMSO-d ₆) δ 8.34 (s, 1H), 8.05 (s, 1H), 7.25 (s, 1H), 7.04 (dd, J = 11.0, 1.4 Hz, 1H), 6.97 (s, 2H), 6.86 (s, 2H), 6.43 (d, J = 3.5 Hz, 1H), 6.37 (d, J = 3.5 Hz, 1H), 5.55 (s, 1H), 5.51 (s, 1H), 5.30 (d, J = 5.7 Hz, 1H), 4.38 (d, J = 5.7 Hz, 1H), 3.08 – 2.95 (m, 2H), 2.71 – 2.58 (m, 2H), 1.37 (s, 3H), 1.28 (s, 3H); LCMS m/z = 541.20 (M ⁺ ; 100%)


 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-6-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-6-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.06 (s, 1H), 7.53 – 7.44 (m, 2H), 6.99 (s, 2H), 6.71 (s, 2H), 6.52 (d, J = 3.5 Hz, 1H), 6.40 (d, J = 3.5 Hz, 1H), 5.57 (s, 1H), 5.55 (s, 1H), 5.32 (d, J = 5.7 Hz, 1H), 4.40 (d, J = 5.7 Hz, 1H), 3.12 – 2.99 (m, 2H), 2.70 – 2.56 (m, 2H), 1.36 (s, 3H), 1.28 (s, 3H); LCMS m/z= 494.99 (M⁺; 20%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-8-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-8-fluoroquinolin-2-amine	LCMS m/z= 495.30 (M⁺; 70%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3,5-dichloroquinolin-2-amine	3,5-Dichloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	LCMS m/z= 511.2 (M⁺; 100%).


 7-((3aS,4R,6aR)-2,2-Dimethyl-6-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	4-Chloro-7-((3aS,4R,6aR)-2,2-dimethyl-6-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	¹H NMR (400 MHz, DMSO-d₆) δ 8.19 (d, J = 7.0 Hz, 1H), 8.04 (s, 1H), 7.43 (s, 1H), 7.31 (d, J = 1.1 Hz, 1H), 6.98 (s, 2H), 6.91 (dd, J = 7.1, 1.7 Hz, 1H), 6.44 (d, J = 3.5 Hz, 1H), 6.35 (d, J = 3.5 Hz, 1H), 5.55 (s, 1H), 5.50 (s, 1H), 5.32 (d, J = 5.6 Hz, 1H), 4.38 (d, J = 5.7 Hz, 1H), 3.02 – 2.91 (m, 2H), 2.71 – 2.57 (m, 2H), 2.45 (d, J = 1.0 Hz, 3H), 1.38 (s, 3H), 1.28 (s, 3H); LCMS m/z= 430.98 (M⁺; 30%).
 7-((3aS,4R,6aR)-6-(2-(2-Amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	4-chloro-7-((3aS,4R,6aR)-6-(2-(3,3-dimethyl-2-(methylthio)-3H-indol-6-yl)ethyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	¹H NMR (400 MHz, DMSO-d₆) δ 9.42 (s, 2H), 8.06 (s, 1H), 7.32 (d, J = 7.6 Hz, 1H), 7.10 – 6.95 (m, 4H), 6.45 (s, 2H), 5.56 (s, 1H), 5.51 (s, 1H), 5.28 (d, J = 5.6 Hz, 1H), 4.37 (d, J = 5.7 Hz, 1H), 3.02 – 2.84 (m, 2H), 2.65 – 2.54 (m, 2H), 1.46 (s, 6H), 1.36 (s, 3H), 1.28 (s, 3H); LCMS m/z= 459.36 (M+1; 20%).
 6'-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-2'-	4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-2'-	¹H NMR (400 MHz, DMSO-d₆) δ 8.06 (s, 1H), 7.64 (d, J = 7.6 Hz, 1H), 7.09 – 6.97 (m, 4H), 6.47 – 6.40 (m, 2H), 5.55 (s, 1H), 5.50 (s, 1H), 5.28 (d, J = 5.7 Hz, 1H), 4.37 (d, J = 5.6 Hz, 1H), 2.99 – 2.87 (m, 2H), 2.83 – 2.71 (m, 2H), 2.64

yl)ethyl)spiro [cyclobutane-1,3'-indol]-2'-amine	(methylthio)spiro[cyclobutane-1,3'-indole]	– 2.53 (m, 2H), 2.36 – 2.29 (m, 2H), 2.27 – 2.16 (m, 2H), 1.37 (s, 3H), 1.28 (s, 3H); LCMS m/z= 472.3 (M+2; 100%).
 7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-N-methylquinolin-2-amine	7-(((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-N-methylquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.08 (s, 1H), 7.80 (d, J = 8.9 Hz, 1H), 7.57 (d, J = 8.7 Hz, 1H), 7.15 (s, 1H), 7.06 (s, 2H), 6.90 (dd, J = 8.6, 2.4 Hz, 1H), 6.86 (d, J = 3.6 Hz, 1H), 6.64 (d, J = 8.9 Hz, 1H), 6.52 (d, J = 3.6 Hz, 1H), 5.84 (s, 1H), 5.65 (s, 1H), 5.42 (d, J = 5.6 Hz, 1H), 4.93 (s, 2H), 4.53 (d, J = 5.7 Hz, 1H), 2.93 (d, J = 4.8 Hz, 3H), 1.46 (s, 3H), 1.30 (s, 3H); LCMS m/z= 459.40 (M+1; 30%).
 7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	3-Chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.11 (s, 1H), 8.07 (s, 1H), 7.67 (t, J = 6.2 Hz, 1H), 7.36 – 7.31 (m, 2H), 7.01 (s, 2H), 6.94 (d, J = 2.2 Hz, 1H), 6.90 – 6.81 (m, 4H), 6.51 (d, J = 3.6 Hz, 1H), 5.83 (s, 1H), 5.65 (s, 1H), 5.42 (d, J = 5.7 Hz, 1H), 4.94 (q, J = 15.3 Hz, 2H), 4.65 (d, J = 6.1 Hz, 2H), 4.55 (d, J = 5.7 Hz, 1H), 3.70 (s, 3H), 1.43 (s, 3H), 1.29 (s, 3H); LCMS m/z= 617.34 (M+; 100%).
 N7-(((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-N-(4-methoxybenzyl)quinolin-2-amine	N7-(((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z= 564.40 (M+1; 100%).




N7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)quinoline-2,7-diamine	dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)quinoline-2,7-diamine	
 N7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)-N7-methylquinoline-2,7-diamine	N7-(((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)-N7-methylquinoline-2,7-diamine	LCMS m/z= 576.95 (M-1; 100%).
 7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methylthio)-N-(4-methoxybenzyl)quinolin-2-amine	7-(((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methylthio)-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.03 (s, 1H), 7.84 (d, J = 8.9 Hz, 1H), 7.57 (d, J = 8.3 Hz, 1H), 7.49 (s, 2H), 7.36 – 7.26 (m, 2H), 7.18 (d, J = 8.4 Hz, 1H), 6.96 (s, 2H), 6.84 (d, J = 8.2 Hz, 2H), 6.82 – 6.76 (m, 1H), 6.16 (s, 1H), 6.01 (s, 1H), 5.76 (s, 2H), 5.71 (s, 1H), 5.54 (s, 1H), 5.38 (s, 1H), 4.55 (s, 1H), 4.34 (d, J = 5.5 Hz, 1H), 4.19 (d, J = 15.2 Hz, 1H), 3.70 (s, 3H), 1.42 (s, 3H), 1.27 (s, 3H); LCMS m/z= 580.83 (M⁺; 100%).



 7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methylthio)-3-chloro-N,N-bis(4-methoxybenzyl)quinolin-2-amine	3-Chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methylthio)-N,N-bis(4-methoxybenzyl)quinolin-2-amine	LCMS m/z= 735.61 (M⁺; 80%).
 7-(2-(((3aS,4R,6aR)-4-(2-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloroquinolin-2-amine	3-Chloro-7-(2-(((3aS,4R,6aR)-4-(2-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.43 (s, 1H), 8.17 (s, 1H), 7.61 (d, J = 8.3 Hz, 1H), 7.40 (d, J = 1.5 Hz, 1H), 7.28 (s, 2H), 7.19 (dd, J = 8.3, 1.7 Hz, 1H), 6.69 (s, 2H), 6.23 (d, J = 3.7 Hz, 1H), 6.03 (d, J = 3.6 Hz, 1H), 5.46 (s, 1H), 5.43 (s, 1H), 5.33 (d, J = 5.7 Hz, 1H), 4.36 (d, J = 5.6 Hz, 1H), 3.10 – 2.93 (m, 2H), 2.67 – 2.56 (m, 2H), 1.36 (s, 3H), 1.28 (s, 3H); LCMS m/z= 476.98 (M⁺; 100%).
 7-(2-(((3aS,4R,6aR)-4-(4-amino-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-bromo-5-fluorophenyl)-3-chloroquinolin-2-amine	3-Bromo-7-(2-(((3aS,4R,6aR)-4-(4-chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-(4-bromo-5-fluorophenyl)-3-chloroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.35 (s, 1H), 8.06 (s, 1H), 7.60 (d, J = 8.3 Hz, 1H), 7.45 (s, 1H), 7.36 – 7.31 (m, 2H), 7.21 – 7.16 (m, 2H), 6.95 (s, 2H), 6.86 – 6.82 (m, 2H), 6.35 (d, J = 2.1 Hz, 1H), 5.58 (s, 1H), 5.48 (s, 1H), 5.28 (d, J = 5.6 Hz, 1H), 4.37 (d, J = 5.6 Hz, 1H), 3.10 – 2.93 (m, 2H), 2.67 – 2.56 (m, 2H), 1.36 (s, 3H), 1.28 (s, 3H); LCMS m/z= 554.12 (M⁺; 100%).




yl)ethyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	methoxybenzyl)quinolin-2-amine	1H), 4.27 (d, J = 4.1 Hz, 2H), 3.68 (s, 3H), 3.05 – 2.97 (m, 2H), 2.67 – 2.57 (m, 2H), 1.36 (s, 3H), 1.27 (s, 3H); LCMS m/z= 660.97 (M+1; 25%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-chloroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)quinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. Run Time (min): 10.00, Injection Volume: 5.00 μL Wavelength: 225nm HEX_0.1%DEA_IPA_DCM_60_40_A_C_1.2ML_10MIN Flow Rate: 1.2 ml/min. Column : CHIRALPAK IA CRL-027 OLD Column Temp: 25.0 $^{\circ}$C Mobile Phase A: HEX_0.1%DEA Mobile Phase C: IPA:DCM Mobile Phase B: NA Mobile Phase D: NA Diastereomer-1 ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.17 (s, 1H), 8.04 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 7.38 (s, 1H), 7.16 (dd, J = 8.1, 1.6 Hz, 1H), 6.98 (s, 2H), 6.67 (s, 2H), 6.32 (d, J = 3.5 Hz, 1H), 6.29 (d, J = 3.6 Hz, 1H), 5.55 (s, 1H), 5.52 (s, 1H), 5.41 (d, J = 5.8 Hz, 1H), 4.35 (d, J = 5.8 Hz, 1H), 3.12 – 3.03 (m, 1H), 2.96 – 2.90 (m, 1H), 2.88 –



		2.81 (m, 1H), 1.31 (s, 3H), 1.27 (s, 3H), 1.17 (d, J = 6.8 Hz, 3H); LCMS m/z= 491.36 (M⁺; 80%). Diastereomer-2 ¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.06 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 7.36 (s, 1H), 7.15 (dd, J = 8.3, 1.6 Hz, 1H), 6.97 (s, 2H), 6.67 (s, 2H), 6.56 (d, J = 3.6 Hz, 1H), 6.44 (d, J = 3.6 Hz, 1H), 5.60 – 5.56 (m, 1H), 5.52 (d, J = 2.5 Hz, 1H), 5.34 (d, J = 5.7 Hz, 1H), 4.40 (d, J = 5.7 Hz, 1H), 3.13 – 3.04 (m, 1H), 2.93 (q, J = 7.2 Hz, 1H), 2.83 (d, J = 10.0 Hz, 1H), 1.40 (s, 3H), 1.28 (s, 3H), 1.14 (d, J = 6.8 Hz, 3H); LCMS m/z= 491.36 (M⁺; 80%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	3-Bromo-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-N-(4-methoxybenzyl)quinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. Run Time (min): 10.00 Injection Volume: 5.00 μL, Wavelength: 251nm, HEX_0.1%DEA_IPA-DCM_50_50_A_C_1.0ML_10M IN_251nm Flow Rate: 1.0 ml/min. Column : CHIRALPAK IG CRL-071 Column Temp: 25.0 °C. Mobile Phase A: HEX_0.1%DEA Mobile Phase C:



		IPA-DCM_1-1, Mobile Phase B: NA Mobile Phase D: NA Diastereomer-1: ^1H NMR (400 MHz, DMSO-d_6) δ 8.36 (s, 1H), 8.07 (s, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.41 (d, J = 1.4 Hz, 1H), 7.37 – 7.33 (m, 2H), 7.19 – 7.13 (m, 2H), 6.99 (s, 2H), 6.87 – 6.83 (m, 2H), 6.63 (d, J = 3.6 Hz, 1H), 6.46 (d, J = 3.5 Hz, 1H), 5.59 (s, 1H), 5.53 (s, 1H), 5.32 (d, J = 5.7 Hz, 1H), 4.65 (d, J = 6.0 Hz, 2H), 4.40 (d, J = 5.8 Hz, 1H), 3.69 (s, 3H), 3.11 – 3.04 (m, 1H), 2.86 – 2.76 (m, 2H), 1.41 (s, 3H), 1.28 (s, 3H), 1.14 (d, J = 6.5 Hz, 3H); LCMS m/z = 657.47 ($M+2$; 45%). Diastereomer-2: ^1H NMR (400 MHz, DMSO-d_6) δ 8.35 (s, 1H), 8.05 (s, 1H), 7.60 (d, J = 8.1 Hz, 1H), 7.42 (s, 1H), 7.33 (d, J = 8.5 Hz, 2H), 7.18 – 7.13 (m, 2H), 7.00 (s, 2H), 6.84 (d, J = 8.6 Hz, 2H), 6.43 (d, J = 3.6 Hz, 1H), 6.38 (d, J = 3.5 Hz, 1H), 5.55 (s, 1H), 5.52 (s, 1H), 5.42 (d, J = 5.7 Hz, 1H), 4.63 (d, J = 6.0 Hz, 2H), 4.38 (d, J = 5.7 Hz, 1H), 3.69 (s, 3H), 3.10 – 3.05 (m, 1H), 2.96 – 2.90 (m, 2H), 1.32 (s, 3H), 1.27 (s, 3H),
--	--	---



		1.16 (d, J = 6.8 Hz, 3H); LCMS m/z= 655.47 (M ⁺ ; 35%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-chloro-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-5-fluoroquinolin-2-amine	LCMS m/z= 509.3 (M ⁺ ; 60%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-bromo-5-fluoroquinolin-2-amine		Diastereomeric mixture was separated by chiral preparative HPLC. Run Time (min): 15.00, Injection Volume: 10.00 μL, Wavelength: 254nm, Flow Rate: 1.50 ml/min, Column Temp: 30.0 $^{\circ}$C, Instrument Method: ACN_DEA_100_B_1.5ML_15M IN_254NM, Column : CHIRALPAK OX-H CRL-061, Mobile Phase A: NA, Mobile Phase B: ACN_0.1%DEA Diastereomer-1: LCMS m/z = 553.20, 555.20 (M⁺;M+2; 100%) Diastereomer-2: ¹H NMR (400 MHz, DMSO-d₆) δ 8.35 (s, 1H), 8.06 (s, 1H), 7.20 (s, 1H), 7.00 –



		6.96 (m, 3H), 6.86 (s, 2H), 6.57 (d, J = 3.5 Hz, 1H), 6.45 (d, J = 3.5 Hz, 1H), 5.57 (s, 1H), 5.53 (s, 1H), 5.34 (d, J = 5.8 Hz, 1H), 4.41 (d, J = 5.7 Hz, 1H), 3.09 – 3.01 (m, 2H), 2.85 – 2.83 (m, 1H), 1.40 (s, 3H), 1.28 (s, 3H), 1.15 (d, J = 6.0 Hz, 3H); LCMS m/z = 555.20 (M+2; 100%)
 7-(1-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-N-methylquinolin-2-amine	7-(1-((3aS,4R,6aR)-4-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-N-methylquinolin-2-amine	LCMS m/z = 473.11 (M+1; 20%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.17 (d, J = 0.7 Hz, 1H), 8.09 (s, 1H), 7.27 (s, 1H), 7.09 – 7.01 (m, 3H), 6.95 (s, 2H), 6.65 (d, J = 3.5 Hz, 1H), 6.47 (d, J = 3.5 Hz, 1H), 5.63 – 5.58 (m, 1H), 5.49 – 5.44 (m, 1H), 4.00 (d, J = 0.9 Hz, 1H), 3.06 (t, J = 7.6 Hz, 2H), 2.70 – 2.59 (m, 1H), 2.59 – 2.53 (m, 1H), 1.39 (s, 3H), 1.30 (s, 3H), 1.28 (s, 3H); LCMS m/z = 509.2 (M+; 90%).



 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-6a-ethyl-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-6a-ethyl-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.18 (s, 1H), 8.09 (s, 1H), 7.28 (s, 1H), 7.06 (dd, J = 11.1, 1.4 Hz, 1H), 6.99 (s, 2H), 6.95 (s, 2H), 6.68 (d, J = 3.5 Hz, 1H), 6.45 (d, J = 3.5 Hz, 1H), 5.73 (d, J = 2.3 Hz, 1H), 5.47 – 5.43 (m, 1H), 4.09 (s, 1H), 3.07 (t, J = 7.6 Hz, 2H), 2.70 – 2.54 (m, 2H), 1.84 (dq, J = 14.9, 7.4 Hz, 1H), 1.59 (dq, J = 14.7, 7.4 Hz, 1H), 1.31 (s, 3H), 1.27 (s, 3H), 0.68 (t, J = 7.4 Hz, 3H); LCMS m/z= 523.44 (M⁺; 95%).
 7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-chloro-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-5-fluoroquinolin-2-amine	Diastereomer-1: ¹H NMR (400 MHz, DMSO-d₆) δ 8.20 (s, 1H), 8.06 (s, 1H), 7.26 (s, 1H), 7.06 (dd, J = 11.1, 1.4 Hz, 1H), 6.96 (s, 4H), 6.21 (d, J = 3.5 Hz, 1H), 5.97 (d, J = 3.6 Hz, 1H), 5.69 (d, J = 2.6 Hz, 1H), 5.45 (d, J = 2.7 Hz, 1H), 3.86 (s, 1H), 3.10 – 2.97 (m, 2H), 2.93 – 2.84 (m, 1H), 1.36 (s, 3H), 1.27 (s, 3H), 1.24 (d, J = 3.0 Hz, 3H), 1.19 (s, 3H); LCMS m/z= 523.32 (M⁺; 50%). Note- Only desired isomer was isolated
	3-Chloro-7-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-	LCMS m/z= 631.30 (M⁺; 60%).


7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	
 7-(1-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	3-Chloro-7-(1-(((3aS,4R,6aR)-4-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	Diastereomer-1 LCMS m/z= 631.34 (M⁺; 100%). Diastereomer-2 LCMS m/z= 631.34 (M⁺; 100%).
 7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	3-bromo-7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.33 (s, 1H), 8.08 (s, 1H), 7.57 (d, J = 8.2 Hz, 1H), 7.43 (d, J = 1.6 Hz, 1H), 7.37 – 7.31 (m, 2H), 7.19 (t, J = 6.1 Hz, 1H), 7.16 (dd, J = 8.2, 1.6 Hz, 1H), 7.14 (d, J = 3.5 Hz, 1H), 7.03 (s, 2H), 6.90 – 6.84 (m, 2H), 6.61 (d, J = 3.5 Hz, 1H), 5.21 (d, J = 7.3 Hz, 1H), 5.01 (s, 1H), 4.62 (d, J = 6.1 Hz, 2H), 4.51 (dd, J = 7.4, 1.5 Hz, 1H), 3.71 (s, 3H), 2.88 – 2.79 (m, 2H), 2.36 – 2.24 (m, 1H), 1.67 –



		1.58 (m, 1H), 1.48 (s, 3H), 1.46 – 1.41 (m, 1H), 1.19 (s, 3H), 0.92 (t, J = 4.7 Hz, 1H), 0.74 – 0.68 (m, 1H); LCMS m/z = 656.2 (M+1; 40%).
 7-((E)-1-((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)prop-1-en-2-yl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	3-bromo-7-((E)-1-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)prop-1-en-2-yl)-N-(4-methoxybenzyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.38 (s, 1H), 8.08 (s, 1H), 7.64 (s, 1H), 7.58 (d, J = 8.3 Hz, 1H), 7.45 – 7.40 (m, 1H), 7.33 (d, J = 8.4 Hz, 2H), 7.16 (d, J = 3.6 Hz, 1H), 7.12 (t, J = 6.0 Hz, 1H), 7.01 (s, 2H), 6.87 (d, J = 8.6 Hz, 2H), 6.63 (d, J = 3.5 Hz, 1H), 6.17 (s, 1H), 5.26 (d, J = 6.9 Hz, 1H), 4.94 (s, 1H), 4.70 (dd, J = 14.7, 6.0 Hz, 1H), 4.59 (dd, J = 14.7, 6.0 Hz, 1H), 4.48 (d, J = 6.9 Hz, 1H), 3.71 (s, 3H), 2.12 (s, 3H), 1.51 (s, 3H), 1.46 – 1.39 (m, 1H), 1.19 (s, 3H), 0.90 (t, J = 5.1 Hz, 1H), 0.26 – 0.18 (m, 1H); LCMS m/z = 668.97 (M+1; 30%).
 7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	3-chloro-7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. <i>Wavelength:</i> 225nm, <i>Instrument Method:</i> HEX-0.1%DEA_IPA-DCM_50_50_A_C_1.2ML_8MIN <i>Flow Rate:</i> 1.2 ml/min, <i>Column:</i> CHIRALPAK IG CRL-071

fluoro-N-(4-methoxybenzyl)quinolin-2-amine	methoxybenzyl)quinolin-2-amine	Column Temp: 25°C, Mobile Phase A: HEX_0.1%DEA Mobile Phase C: IPA-DCM, Mobile Phase B: NA Mobile Phase D: NA First Diastereomer: LCMS m/z = 643.09(M⁺; 30%). Second Diastereomer: LCMS m/z = 643.09(M⁺; 30%).
 7-(2-(((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)-2-cyclopropylethyl)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	3-chloro-7-(2-(((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)-2-cyclopropylethyl)-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z = 669.10(M ⁺ ; 20%).
 7-(2-(((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-N-(4-bromobenzyl)quinolin-2-amine	3-Bromo-7-(2-(((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propyl)-N-(4-methoxybenzyl)quinolin-2-amine	LCMS m/z = 670.97(M ⁺ ; 20%).

3b(3aH)-yl)propyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	methoxybenzyl)quinolin-2-amine	
 7-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinazolin-2-amine	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinazolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.04 (s, 1H), 8.07 (s, 1H), 7.79 (t, J = 6.3 Hz, 1H), 7.69 (d, J = 8.2 Hz, 1H), 7.35 (s, 1H), 7.30 (d, J = 8.2 Hz, 2H), 7.17 (dd, J = 8.2, 1.6 Hz, 1H), 7.13 (d, J = 3.5 Hz, 1H), 7.01 (s, 2H), 6.90 – 6.84 (m, 2H), 6.61 (d, J = 3.5 Hz, 1H), 5.21 (d, J = 7.1 Hz, 1H), 5.01 (s, 1H), 4.57 – 4.48 (m, 3H), 3.71 (s, 1H), 2.91 – 2.80 (m, 2H), 2.35 – 2.24 (m, 1H), 1.69 – 1.58 (m, 1H), 1.48 (s, 3H), 1.47 – 1.42 (m, 1H), 1.20 (s, 3H), 0.93 (t, J = 4.7 Hz, 1H), 0.75 – 0.69 (m, 1H); LCMS m/z = 578.3 (M+1; 90%).
 7-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinazolin-2-amine	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(4-methoxybenzyl)quinazolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.07 (s, 1H), 7.88 (d, J = 8.9 Hz, 1H), 7.36 (d, J = 8.2 Hz, 1H), 7.13 (d, J = 3.7 Hz, 1H), 7.09 (t, J = 7.4 Hz, 1H), 7.00 (s, 2H), 6.75 (d, J = 8.9 Hz, 1H), 6.65 (s, 2H), 6.61 (d, J = 3.5 Hz, 1H),


cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-8-fluoroquinolin-2-amine	-d][1,3]dioxol-3b(3aH)-yl)ethyl)-8-fluoroquinolin-2-amine	5.21 (d, J = 7.2 Hz, 1H), 5.01 (s, 1H), 4.55 (d, J = 7.2 Hz, 1H), 2.93 – 2.79 (m, 2H), 2.29 – 2.20 (m, 1H), 1.68 – 1.58 (m, 1H), 1.49 (s, 3H), 1.47 – 1.43 (m, 1H), 1.21 (s, 3H), 0.96 (t, J = 4.8 Hz, 1H), 0.75 (dd, J = 9.0, 5.0 Hz, 1H); LCMS m/z = 475.2 (M+1; 30%).
 7-((2-(((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-methylquinolin-2-amine	7-(2-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-methylquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 7.84 (d, J = 8.9 Hz, 1H), 7.55 (d, J = 8.1 Hz, 1H), 7.46 (s, 1H), 7.16 – 7.10 (m, 2H), 7.06 (s, 2H), 6.72 (d, J = 8.9 Hz, 1H), 6.62 (d, J = 3.5 Hz, 1H), 5.22 (d, J = 7.2 Hz, 1H), 5.02 (s, 1H), 4.52 (dd, J = 7.3, 1.5 Hz, 1H), 2.91 (d, J = 4.7 Hz, 3H), 2.88 – 2.77 (m, 2H), 2.32 – 2.24 (m, 1H), 1.70 – 1.61 (m, 1H), 1.49 (s, 3H), 1.48 – 1.44 (m, 1H), 1.20 (s, 3H), 0.94 (t, J = 4.7 Hz, 1H), 0.74 (dd, J = 9.2, 5.0 Hz, 1H); LCMS m/z = 471.23 (M+1; 15%).
 7-(((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-3-bromo-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-methylquinolin-2-amine	3-bromo-7-(((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-methylquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.30 (s, 1H), 8.09 (s, 1H), 7.64 – 7.59 (m, 1H), 7.38 (d, J = 3.6 Hz, 1H), 7.37 – 7.31 (m, 2H), 7.16 (t, J = 5.9 Hz, 1H), 7.04 (s,

7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methoxy)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methoxy)-N-(4-methoxybenzyl)quinolin-2-amine	2H), 7.00 (d, J = 2.5 Hz, 1H), 6.97 – 6.92 (m, 1H), 6.91 – 6.84 (m, 2H), 6.65 (d, J = 3.6 Hz, 1H), 5.30 (d, J = 7.1 Hz, 1H), 5.14 (s, 1H), 4.63 (d, J = 6.0 Hz, 2H), 4.50 (d, J = 7.1 Hz, 1H), 4.35 (d, J = 10.5 Hz, 1H), 4.21 (d, J = 10.5 Hz, 1H), 3.71 (s, 3H), 1.78 – 1.71 (m, 1H), 1.48 (s, 3H), 1.18 (s, 3H), 1.11 – 1.06 (m, 1H), 0.88 – 0.83 (m, 1H); LCMS m/z = 659.3 (M+2; 100%).
 7-((((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methylthio)-N-(4-methoxybenzyl)quinolin-2-amine	7-((((3aR,3bS,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methylthio)-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (s, 1H), 7.80 (d, J = 8.9 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 7.46 (t, J = 5.7 Hz, 1H), 7.41 (d, J = 1.8 Hz, 1H), 7.35 – 7.30 (m, 2H), 7.23 (d, J = 3.5 Hz, 1H), 7.12 (dd, J = 8.3, 1.9 Hz, 1H), 7.01 (s, 2H), 6.91 – 6.86 (m, 2H), 6.75 (d, J = 8.9 Hz, 1H), 6.58 (d, J = 3.5 Hz, 1H), 5.23 (dd, J = 7.3, 1.4 Hz, 1H), 5.03 (s, 1H), 4.55 (d, J = 5.7 Hz, 2H), 4.51 (dd, J = 7.4, 1.5 Hz, 1H), 3.72 (s, 3H), 3.67 (d, J = 12.6 Hz, 1H), 3.32 (d, J = 12.7 Hz, 1H), 1.71 – 1.63 (m, 1H), 1.47 (s, 3H), 1.17 (s, 3H), 1.06 (t, J = 4.8 Hz, 1H), 1.01 –

		0.95 (m, 1H); LCMS m/z = 595.21 (M ⁺ ; 70%)
 7-((3aR,3bR,4aS,5R,5aS)-3b-(2-(2-amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	7-((3aR,3bR,4aS,5R,5aS)-3b-(2-(3,3-dimethyl-2-(methylthio)-3H-indol-6-yl)ethyl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.07 (s, 1H), 7.18 (d, J = 7.5 Hz, 1H), 7.11 (d, J = 3.6 Hz, 1H), 7.01 (s, 2H), 6.94 (s, 1H), 6.88 (d, J = 7.6 Hz, 1H), 6.61 (d, J = 3.5 Hz, 1H), 5.18 (d, J = 7.2 Hz, 1H), 5.00 (s, 1H), 4.52 (d, J = 7.1 Hz, 1H), 2.77 – 2.65 (m, 2H), 2.22 – 2.12 (m, 1H), 1.66 – 1.55 (m, 1H), 1.47 (s, 3H), 1.44 (dd, J = 9.2, 4.3 Hz, 1H), 1.36 (s, 3H), 1.24 (s, 3H), 1.19 (s, 3H), 0.93 (t, J = 4.7 Hz, 1H), 0.75 (dd, J = 9.1, 5.0 Hz, 1H); LCMS m/z = 473.4 (M ⁺ ; 80%)
 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-(4-methoxybenzyl)quinolin-2-amine	¹ H NMR (400 MHz, Chloroform-d) δ 8.34 (s, 1H), 7.87 (d, J = 8.9 Hz, 1H), 7.63 (s, 1H), 7.55 (d, J = 8.0 Hz, 1H), 7.35 (d, J = 8.5 Hz, 2H), 7.22 (d, J = 8.1 Hz, 1H), 6.97 (d, J = 3.6 Hz, 1H), 6.91 (d, J = 8.7 Hz, 2H), 6.65 (d, J = 9.0 Hz, 1H), 6.39 (d, J = 3.6 Hz, 1H), 5.25 (s, 2H), 5.19 (d, J = 7.1 Hz, 1H), 5.14 (s, 1H), 4.67 – 4.62 (m, 3H), 3.82 (s, 3H), 3.06 – 2.93 (m, 2H), 2.41 – 2.36 (m, 1H), 2.06 – 2.01 (m,

		1H), 1.60 (s, 3H), 1.51 (dd, J = 8.7, 4.2 Hz, 1H), 1.28 (s, 3H), 1.15 (t, J = 4.8 Hz, 1H), 0.78 (dd, J = 5.5, 3.2 Hz, 1H); LCMS m/z = 577.5 (M+1; 60%)
--	--	--

7-(1-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)propan-2-yl)-N-(4-methoxybenzyl)quinolin-2-amine




- 5 A mixture of 7-((E)-1-((3aR,3bS,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)prop-1-en-2-yl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine (0.05g, 0.075 mmol), ammonium formate (0.331 g, 5.24 mmol) and Pd/C (0.024 g, 0.225 mmol) in EtOH (15 ml) was heated at 75°C for 8h. The reaction mixture was cooled 25°C, filtered through
- 10 celite and concentrated *in vacuo* to get 0.06g of crude compound. The obtained residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisept[®] R_f column with gradient elution (0 to 5%) of methanol in dichloromethane to afford the title compound (0.042g, 95%) as an off-white solid. LCMS m/z = 591.29 (M+1, 100%).

15

Examples

Example-1: (1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-1)






- 20 The mixture of 7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine (0.220 g, 0.343 mmol) in TFA (3.96 ml, 51.4 mmol) was stirred at 50°C for 1.5h. The resulting mixture was concentrated *in vacuo* and obtained residue




was dissolved in MeOH (5ml). K₂CO₃ (0.142 g, 1.029 mmol) was added and stirred the reaction mixture at 50°C for 1.5h. Reaction mixture was filtered and filtrate was concentrated under reduced pressure to get 0.27g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisepp® R_f column with gradient elution (0 to 7%) of methanol in dichloromethane to afford the title compound (0.03g, 18%) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.35 (s, 1H), 8.06 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 7.36 (d, J = 1.6 Hz, 1H), 7.16 (dd, J = 8.2, 1.7 Hz, 1H), 7.08 (s, 2H), 6.67 (d, J = 3.5 Hz, 1H), 6.61 (s, 2H), 6.45 (d, J = 3.5 Hz, 1H), 5.50 (d, J = 4.4 Hz, 1H), 5.44 (t, J = 1.7 Hz, 1H), 4.98 (d, J = 6.4 Hz, 2H), 4.45 (t, J = 5.9 Hz, 1H), 3.97 (d, J = 5.0 Hz, 1H), 3.04 – 2.87 (m, 2H), 2.61 – 2.53 (m, 2H); LCMS m/z = 483.01 (M+2, 90%).


Examples in table-13 were synthesized by following an analogous reaction protocol as was used for the preparation of (1S,2R,5R)-3-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol using the appropriate starting materials (Instead of TFA, 3N HCl/MeOH could also be used).

Table-13



Structure & IUPAC name	Intermediate used	¹ H NMR & LCMS data
 (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-N-(4-methoxybenzyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.17 (s, 1H), 8.09 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 7.37 (s, 1H), 7.29 (s, 2H), 7.16 (dd, J = 8.3, 1.7 Hz, 1H), 6.75 – 6.66 (m, 3H), 6.49 (d, J = 3.6 Hz, 1H), 5.50 (d, J = 4.4 Hz, 1H), 5.43 (d, J = 1.9 Hz, 1H), 5.01 (d, J = 6.7 Hz, 2H), 4.45 (t, J = 5.9 Hz, 1H), 3.96 (d, J = 5.0 Hz, 1H), 2.94 (ddt, J = 21.4, 14.1, 7.2 Hz, 2H), 2.58 – 2.55 (m, 2H); LCMS m/z = 436.92 (M ⁺ , 20%)




Compound-3  (1S,2R,5R)-3-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-1,2-diol	7-(2-(((3aS,4R,6aR)-4-(4-amino-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.39 (s, 1H), 8.08 (s, 1H), 7.63 (d, J = 8.2 Hz, 1H), 7.37 (s, 1H), 7.18 (dd, J = 8.2, 1.6 Hz, 3H), 6.72 (d, J = 27.4 Hz, 2H), 6.55 (d, J = 2.1 Hz, 1H), 5.54 (s, 1H), 5.40 (q, J = 1.7 Hz, 1H), 5.10 - 4.91 (m, 2H), 4.43 (d, J = 5.7 Hz, 1H), 3.92 (q, J = 5.2 Hz, 1H), 3.04 - 2.87 (m, 2H) 2.65 - 2.52 (m, 2H); LCMS m/z = 501 (M+1, 30%)
Compound-4  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)thio)methyl)cyclopent-1,2-diol	7-((((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)thio)-N-(4-methoxybenzyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.00 (s, 1H), 7.86 (d, J = 8.8 Hz, 1H), 7.56 (d, J = 8.3 Hz, 1H), 7.40 (d, J = 1.8 Hz, 1H), 7.15 (dd, J = 8.3, 1.9 Hz, 1H), 6.89 (s, 2H), 6.72 (d, J = 8.8 Hz, 1H), 6.48 (s, 2H), 6.26 (d, J = 3.5 Hz, 1H), 6.21 (d, J = 3.6 Hz, 1H), 5.57 (s, 1H), 5.49 (s, 1H), 5.11 (d, J = 5.5 Hz, 1H), 5.04 (d, J = 6.4 Hz, 1H), 4.58 (s, 1H), 4.01 (d, J = 14.7 Hz, 1H), 3.88 - 3.81 (m, 1H), 3.75 - 3.66 (m, 1H); LCMS m/z = 420.85 (M⁺, 100%)
Compound-5	7-((((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-	¹H NMR (400 MHz, DMSO-d₆) δ 8.19 (s, 1H), 8.00 (s, 1H), 7.62 (d, J = 8.4 Hz, 1H), 7.43 (s, 1H), 7.23 (dd, J = 8.4, 1.9 Hz, 1H), 6.94 (s, 2H), 6.82 (s, 2H), 6.28 (q, J = 3.5



 (1S,2R,5R)-3-(((2-amino-3-chloroquinolin-7-yl)thio)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	cyclopenta[d][1,3]dioxol-6-yl)methyl)thio)-3-chloro-N,N-bis(4-methoxybenzyl)quinolin-2-amine	Hz, 2H), 5.61 (s, 1H), 5.50 (s, 1H), 5.09 (dd, J = 24.6, 6.5 Hz, 2H), 4.57 (t, J = 6.0 Hz, 1H), 4.04 (d, J = 14.9 Hz, 1H), 3.87 (q, J = 5.7 Hz, 1H), 3.74 (d, J = 15.0 Hz, 1H); LCMS m/z = 455.50 (M ⁺ , 50%)
Compound-6  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)(methyl)amino)methyl)cyclopent-3-ene-1,2-diol	N7-(((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)-N7-methylquinoline-2,7-diamine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.02 (s, 1H), 7.65 (d, J = 8.7 Hz, 1H), 7.39 (d, J = 8.8 Hz, 1H), 6.95 – 6.86 (m, 3H), 6.81 (dd, J = 8.9, 2.6 Hz, 1H), 6.61 (d, J = 2.5 Hz, 1H), 6.49 (d, J = 3.5 Hz, 1H), 6.41 (d, J = 8.7 Hz, 1H), 6.09 (s, 2H), 5.51 (s, 1H), 5.41 (d, J = 1.9 Hz, 1H), 5.06 (dd, J = 6.5, 2.5 Hz, 2H), 4.44 (t, J = 6.0 Hz, 1H), 4.17 (d, J = 4.9 Hz, 2H), 4.11 – 4.04 (m, 1H), 3.05 (s, 3H); LCMS m/z = 417.10 (M ⁺ , 100%)
Compound-7a and 7b  (1S,2R,5R)-3-(1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-(((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-bromo-N-(4-methoxybenzyl)quinolin-2-amine	First Diastereomer(Compound-7a): ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.35 (s, 1H), 8.05 (s, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.33 (s, 1H), 7.13 (d, J = 8.4 Hz, 1H), 6.93 (s, 2H), 6.78 (d, J = 3.6 Hz, 1H), 6.59 (s, 2H), 6.49 (d, J = 3.6 Hz, 1H), 5.53 (d, J = 4.5 Hz, 1H), 5.42 (s, 1H), 5.01 – 4.88 (m, 2H), 4.46 (t, J = 5.8 Hz, 1H), 3.99 (t, J = 7.8




		Hz, 1H), 3.03 (dd, J = 11.9, 4.4 Hz, 1H), 2.73 (q, J = 10.6, 8.9 Hz, 2H), 1.24 (s, 3H); LCMS m/z = 497.30 (M+2, 100%) Second Diastereomer(Compound-7b): ¹H NMR (400 MHz, DMSO-d6) δ 8.36 (s, 1H), 8.05 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 7.33 (s, 1H), 7.21 – 7.02 (m, 3H), 6.61 (s, 2H), 6.50 (d, J = 3.5 Hz, 1H), 6.43 (d, J = 3.6 Hz, 1H), 5.52 (d, J = 5.0 Hz, 1H), 5.35 (s, 1H), 5.00 (s, 1H), 4.91 (d, J = 6.4 Hz, 1H), 4.56 (s, 1H), 3.90 (d, J = 5.8 Hz, 1H), 3.07 – 2.99 (m, 1H), 2.83 – 2.74 (m, 2H) 1.09 (d, J = 5.8 Hz, 3H); LCMS m/z = 495.24 (M+, 40%)
(Compound-8a and 8b)  (1S,2R,5R)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(1-((2-(methylamino)quinolin-7-yl)oxy)ethyl)cyclopent-3-ene-1,2-diol	7-(1-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta [d] [1,3] dioxol-6-yl)ethoxy)-N-methyl quinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. <i>Wavelength:</i> 225 nm, <i>Instrument Method:</i> IPA_0.1%DEA_MeOH_0.1%DEA_50_50_0.7ML_12MIN, <i>Flow Rate:</i> 0.70 ml/min <i>Column :</i> CHIRALPAK IB CRL-043 OLD, <i>Column Temp:</i> 30°C, <i>Mobile Phase A:</i> IPA_0.1%DEA, <i>Mobile Phase B:</i> MeOH_0.1%DEA



		First Diastereomer (Compound-8a): ^1H NMR (400 MHz, DMSO-d_6) δ 8.04 (s, 1H), 7.73 (d, J = 8.8 Hz, 1H), 7.49 (d, J = 8.7 Hz, 1H), 7.00 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 4.2 Hz, 2H), 6.84 – 6.80 (m, 2H), 6.57 (d, J = 8.8 Hz, 1H), 6.44 (d, J = 3.5 Hz, 1H), 5.70 (t, J = 1.7 Hz, 1H), 5.60 – 5.56 (m, 1H), 5.21 (d, J = 6.7 Hz, 1H), 5.14 – 5.08 (m, 1H), 5.04 – 4.99 (m, 1H), 4.62 – 4.58 (m, 1H), 4.15 – 4.11 (m, 1H), 3.19 – 3.16 (m, 1H), 2.90 (d, J = 4.7 Hz, 3H), 1.54 (d, J = 6.3 Hz, 3H); LCMS m/z = 433.04 (M+1, 30%) Second Diastereomer(Compound-8b): ^1H NMR (400 MHz, DMSO-d_6) δ 8.02 (s, 1H), 7.73 (d, J = 8.9 Hz, 1H), 7.49 (d, J = 8.7 Hz, 1H), 7.04 (d, J = 2.5 Hz, 1H), 6.94 (d, J = 3.5 Hz, 3H), 6.81 (dd, J = 8.7, 2.5 Hz, 1H), 6.57 (d, J = 8.8 Hz, 1H), 6.54 (d, J = 3.6 Hz, 1H), 5.72 (t, J = 1.5 Hz, 1H), 5.57 (d, J = 5.0 Hz, 1H), 5.26 (d, J = 6.7 Hz, 1H), 5.15 (s, 1H), 5.10 (s, 1H), 4.48 (s, 1H), 4.36 (d, J = 4.2 Hz, 1H),
--	--	---



		3.94 (d, $J = 5.8$ Hz, 1H), 2.91 (d, $J = 4.7$ Hz, 3H), 1.57 (d, $J = 6.5$ Hz, 3H); LCMS (m/z) = 433.04 (M^+ , 30%).
Compound-9  (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol	7-(((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	^1H NMR (400 MHz, DMSO- d_6) δ 8.37 (s, 1H), 8.14 (s, 1H), 7.71 (d, $J = 3.7$ Hz, 1H), 7.07 (s, 2H), 6.97 (d, $J = 3.6$ Hz, 1H), 6.90 (d, $J = 2.2$ Hz, 1H), 6.79 (dd, $J = 11.5, 2.2$ Hz, 1H), 5.61 (s, 2H), 4.89 – 4.77 (m, 2H), 4.76 – 4.63 (m, 3H), 1.82 (s, 3H); LCMS m/z = 471.07 (M^+ , 100%)
Compound-10  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2-(methylamino)quinolin-7-yl)ethyl)cyclopent-3-ene-1,2-diol	7-(2-(((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-N-methylquinolin-2-amine	^1H NMR (400 MHz, DMSO- d_6) δ 8.03 (s, 1H), 7.79 (d, $J = 8.9$ Hz, 1H), 7.54 (d, $J = 8.0$ Hz, 1H), 7.39 (s, 1H), 7.07 (dd, $J = 8.1, 1.6$ Hz, 1H), 6.93 (s, 2H), 6.72 – 6.64 (m, 2H), 6.41 (d, $J = 3.5$ Hz, 1H), 5.51 (s, 1H), 5.43 (d, $J = 1.9$ Hz, 1H), 4.95 (dd, $J = 10.7, 6.5$ Hz, 2H), 4.45 (t, $J = 6.0$ Hz, 1H), 4.01 – 3.90 (m, 1H), 3.12 – 3.08 (m, 2H), 2.90 (d, $J = 4.7$ Hz, 3H), 2.60 – 2.52 (m, 2H); LCMS m/z = 416.48 (M^+ , 60%)
Compound-11	7-(((3aS,4R,6aR)-2,2-dimethyl-6-(2-(3-	^1H NMR (400 MHz, DMSO- d_6) δ 8.20 (d, $J = 7.0$ Hz, 1H), 8.03

 (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)cyclopent-3-ene-1,2-diol	methylimidazo[1,2-a]pyridin-7-yl)ethyl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	(s, 1H), 7.39 (s, 1H), 7.32 (s, 1H), 7.00 – 6.86 (m, 3H), 6.66 (d, J = 3.5 Hz, 1H), 6.42 (d, J = 3.5 Hz, 1H), 5.53 – 5.41 (m, 2H), 4.98 (d, J = 6.4 Hz, 2H), 4.46 (t, J = 6.0 Hz, 1H), 3.98 (q, J = 5.5 Hz, 1H), 3.40 (d, J = 7.0 Hz, 2H), 2.99 – 2.82 (m, 2H), 2.45 (s, 3H); LCMS m/z = 390.91 (M⁺, 90%)
Compound-12  (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-4-methylcyclopent-3-ene-1,2-diol	3-chloro-5-fluoro-N-(4-methoxybenzyl)-7-(((3aS,4R,6aR)-4-(4-((4-methoxybenzyl)amino)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,5-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)quinolin-2-amine	LCMS m/z = 471.23 (M⁺, 60%)
Compound-13  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-4-methylcyclopent-3-ene-1,2-diol	3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.62 (s, 1H), 8.18 (s, 1H), 7.22 (s, 1H), 7.13 – 6.90 (m, 4H), 6.55 (d, J = 3.6 Hz, 1H), 5.61 (d, J = 4.4 Hz, 1H), 5.46 (t, J = 1.7 Hz, 1H), 5.02 (dd, J = 6.5, 4.3 Hz, 2H), 4.47 (t, J = 6.0 Hz, 1H),

yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	6-yl)ethyl)-5-fluoroquinolin-2-amine	4.08 – 3.98 (m, 1H), 3.04 – 2.85 (m, 2H), 2.63 (s, 3H), 2.60 – 2.54 (m, 2H); LCMS m/z = 454.17 (M ⁺ , 100%)
Compound-14  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol hydrochloride	3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 14.95 (s, 1H), 8.41 (s, 1H), 8.26 (t, J = 5.9 Hz, 1H), 7.94 (d, J = 6.9 Hz, 1H), 7.53 (d, J = 3.5 Hz, 1H), 7.43 – 7.29 (m, 2H), 7.29 – 7.03 (m, 3H), 5.64 (s, 1H), 5.58 (d, J = 5.3 Hz, 1H), 4.41 (d, J = 5.6 Hz, 1H), 3.88 (t, J = 5.6 Hz, 1H), 3.10 – 2.99 (m, 2H), 2.93 (s, 3H), 2.62-2.50 (m, 2H); LCMS m/z= 453.11 (M+1, 100%)
Compound-15  (1S,2R,5R)-3-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol	3-Chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(1H-pyrrolo[3,2-c]pyridin-1-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.91 (s, 1H), 8.20 (s, 1H), 8.09 (d, J = 6.2 Hz, 1H), 7.49 (d, J = 6.1 Hz, 1H), 7.24 (s, 1H), 7.11 (d, J = 3.3 Hz, 1H), 7.07 – 7.01 (m, 1H), 6.98 (s, 2H), 6.60 (d, J = 3.3 Hz, 1H), 5.58 (d, J = 1.8 Hz, 1H), 5.39 (s, 1H), 5.18 (dd, J = 17.2, 6.7 Hz, 2H), 4.41 (t, J = 5.4 Hz, 1H), 3.84 (q, J = 5.9 Hz, 1H), 3.11 – 2.91 (m, 2H), 2.68 – 2.58 (m, 2H); LCMS (m/z) = 439.23 (M ⁺ , 100%).
Compound-16	3-chloro-7-(2-((3aS,4R,6aR)-2,2-	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.98 (s, 1H), 8.76 (s, 1H), 8.18


 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-1,2-diol	dimethyl-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	(s, 1H), 7.22 (s, 1H), 7.11 (d, J = 3.6 Hz, 1H), 7.02 (dd, J = 11.1, 1.4 Hz, 1H), 6.96 (s, 2H), 6.49 (d, J = 3.6 Hz, 1H), 5.65 (s, 1H), 5.47 (d, J = 1.9 Hz, 1H), 5.03 (d, J = 6.4 Hz, 2H), 4.48 (t, J = 6.1 Hz, 1H), 4.06 – 4.02 (m, 1H), 3.02 – 2.93 (m, 2H), 2.60 – 2.55 (m, 2H) ; LCMS m/z= 440.17 (M⁺, 60%)
Compound-17  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-aminoquinolin-7-yl)amino)methyl)cyclopent-1,2-diol	N7-(((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methyl)-N2-(4-methoxybenzyl)quinoline-2,7-diamine	¹H NMR (400 MHz, DMSO-d₆) δ 8.04 (s, 1H), 7.60 (d, J = 8.6 Hz, 1H), 7.29 (d, J = 8.6 Hz, 1H), 6.96 - 6.91 (m, 3H), 6.64 (dd, J = 8.7, 2.3 Hz, 1H), 6.51 (d, J = 3.5 Hz, 1H), 6.48 (d, J = 2.2 Hz, 1H), 6.37 (d, J = 8.6 Hz, 1H), 6.20 (t, J = 5.6 Hz, 1H), 6.08 (s, 2H), 5.61 (q, J = 1.8 Hz, 1H), 5.55 (s, 1H), 5.09 - 5.04 (m, 2H), 4.52 (t, J = 5.9 Hz, 1H), 4.11 (q, J = 5.8 Hz, 1H), 3.90 (s, 2H); LCMS m/z= 404.16 (M⁺, 100%)
Compound-18  (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-	7-(((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)methoxy)-3-chloro-5-fluoro-N-(4-	¹H NMR (400 MHz, DMSO-d₆) δ 8.58 (bs, 2H), 8.32 (s, 1H), 8.11 (s, 1H), 7.32 (d, J = 3.6 Hz, 1H), 6.99 (bs, 2H), 6.89 – 6.83 (m, 2H), 6.81 (d, J = 2.3 Hz, 1H), 5.82 (q, J = 1.8 Hz, 1H), 5.63 (d, J = 5.2 Hz, 1H), 5.27

yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	methoxybenzyl)quinolin-2-amine	(bs, 2H), 4.93 – 4.80 (m, 2H), 4.56 (d, J = 5.7 Hz, 1H), 4.21 (t, J = 5.6 Hz, 1H); LCMS m/z= 457.17 (M ⁺ , 70%)
Compound-19  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(((2-(methylanino)quinolin-7-yl)oxy)methyl)cyclopent-3-ene-1,2-diol	7-(((3aR,6R,6aS)-6-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-6,6a-dihydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)methoxy)-N-methylquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (s, 1H), 8.00 (d, J = 8.9 Hz, 1H), 7.69 (d, J = 8.8 Hz, 1H), 7.40 (d, J = 13.8 Hz, 3H), 7.12 (d, J = 3.7 Hz, 1H), 7.03 (d, J = 8.9 Hz, 1H), 6.80 (d, J = 9.0 Hz, 1H), 6.64 (d, J = 3.5 Hz, 1H), 5.82 (d, J = 2.0 Hz, 1H), 5.61 (s, 1H), 5.30 – 5.08 (m, 2H), 4.86 (s, 2H), 4.60 (t, J = 5.6 Hz, 1H), 4.28 – 4.04 (m, 2H) 3.05 (d, J = 4.6 Hz, 3H); LCMS m/z= 419.16 (M ⁺ , 100%)
Compound-20a and 20b  (1S,2R,5R)-3-(1-((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(1-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethoxy)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. First Diastereomer (Compound-20a): ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (s, 1H), 7.92 (s, 1H), 7.36 (d, J = 3.6 Hz, 1H), 7.16 (d, J = 3.5 Hz, 1H), 6.83 (s, 2H), 6.65 – 6.35 (m, 3H), 5.96 (t, J = 9.7 Hz, 1H), 5.82 – 5.54 (m, 2H), 5.07 – 4.91 (m, 2H), 4.61 – 4.36 (m, 2H), 4.18 (dd, J = 9.6, 4.8 Hz, 1H), 1.65 (ddd, J

		= 26.6, 6.9, 2.5 Hz, 3H); LCMS m/z= 471.23 (M⁺, 90%) Second Diastereomer (Compound-20b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 8.01 (s, 1H), 6.93 (s, 5H), 6.87 – 6.75 (m, 2H), 6.54 (d, J = 3.6 Hz, 1H), 5.62 (d, J = 1.6 Hz, 1H), 5.53 – 5.49 (m, 1H), 5.26 (d, J = 5.9 Hz, 1H), 5.23 – 5.13 (m, 2H), 4.55 (t, J = 5.7 Hz, 1H), 4.15 – 4.00 (m, 1H), 1.57 (d, J = 6.5 Hz, 3H); LCMS m/z= 471.23 (M⁺, 80%)
Compound-21  (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-1,2-diol	3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)quinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.62 (s, 1H), 8.17 (s, 1H), 7.61 (d, J = 8.3 Hz, 1H), 7.44 (s, 1H), 7.37 (s, 1H), 7.17 (d, J = 8.2 Hz, 1H), 7.02 (d, J = 3.6 Hz, 1H), 6.70 (s, 2H), 5.62 (s, 1H), 5.45 (s, 1H), 5.04 (s, 2H), 4.47 (d, J = 5.5 Hz, 1H), 4.14 – 3.91 (m, 1H), 3.06 – 2.87 (m, 2H) 2.58 (d, J = 32.3 Hz, 5H); LCMS m/z= 436.23 (M⁺, 90%)
Compound-22  Chemical Formula: C₂₆H₂₈N₆O₂ (1S,2R,5R)-5-(4-Amino-7H-	7-(2-((3aS,4R,6aR)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.05 (d, J = 9.1 Hz, 1H), 7.77 – 7.51 (m, 4H), 7.24 (d, J = 8.1 Hz, 1H), 6.82 (dd, J = 17.2, 6.4 Hz, 2H), 6.57 (d, J = 3.6 Hz, 1H), 5.53 (s, 1H),

pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2-(cyclobutylamino) quinolin-7-yl)ethyl)cyclopent-3-ene-1,2-diol	dioxol-6-yl)ethyl)-N-cyclobutylquinolin-2-amine	5.45 (s, 1H), 5.03 (s, 2H), 4.55 – 4.43 (m, 2H), 4.01 (t, J = 5.2 Hz, 1H), 3.06 – 2.87 (m, 2H), 2.59 – 2.54 (m, 2H), 2.41 (d, J = 8.7 Hz, 2H), 2.01 (q, J = 9.8 Hz, 2H), 1.76 (q, J = 9.4 Hz, 2H); LCMS m/z= 457.3 (M ⁺ , 90%)
--	---	---

Example-2: (1S,2R,5R)-3-(2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-23)



5 The mixture of 7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-fluoroquinolin-2-amine (0.060 g, 0.130 mmol) in TFA (1.004 ml, 13.03 mmol) was stirred at 25°C for 6h under N₂ atmosphere. The reaction mixture was basified with ice cold solution of

10 aq.sat.NaHCO₃ (20ml) and extracted with ethyl acetate (20 ml). Layers were separated, organic layer was washed with brine (20 ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 0.12g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redise[®] R_f column with gradient elution (0 to 10%) of methanol in dichloromethane to afford the title




15 compound (25 mg, 45.6 % yield) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.05 (d, J = 7.3 Hz, 1H), 7.79 (d, J = 11.8 Hz, 1H), 7.59 (d, J = 8.2 Hz, 1H), 7.37 (s, 1H), 7.16 (d, J = 8.3 Hz, 1H), 6.99 (s, 2H), 6.74 (s, 2H), 6.58 (d, J = 3.5 Hz, 1H), 6.40 (d, J = 3.5 Hz, 1H), 5.50 (d, J = 9.5 Hz, 1H), 5.42 (d, J = 1.9 Hz, 1H), 4.97 (dd, J = 6.6, 3.1 Hz, 2H), 4.45 (t, J = 6.0 Hz, 1H), 3.95 (q, J = 5.5 Hz, 1H), 3.04 – 2.85 (m, 2H), 2.60 – 2.53 (m, 2H);




20 LCMS m/z= 420.92 (M⁺, 100%).



Examples in table-14 were synthesized by following an analogous reaction protocol as was used for the preparation of (1S,2R,5R)-3-(2-(2-amino-3-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol using the appropriate



starting materials (Instead of TFA, aq. TFA or FeCl₃. DCM could also be used at appropriate temperature).


Table-14



Structure & IUPAC name	Intermediate used	¹ H NMR & LCMS data
Compound-24  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (d, J = 0.8 Hz, 1H), 8.03 (s, 1H), 7.22 (s, 1H), 7.01 (dd, J = 11.0, 1.4 Hz, 1H), 6.95 (s, 2H), 6.92 (s, 2H), 6.63 (d, J = 3.5 Hz, 1H), 6.42 (d, J = 3.5 Hz, 1H), 5.50 (t, J = 3.2 Hz, 1H), 5.45 (t, J = 1.7 Hz, 1H), 4.96 (dd, J = 6.3, 3.0 Hz, 2H), 4.45 (t, J = 5.8 Hz, 1H), 3.97 (q, J = 5.5 Hz, 1H), 3.03 – 2.87 (m, 2H), 2.56 (t, J = 7.0 Hz, 2H); LCMS m/z = 454.98 (M⁺, 40%)
Compound-25  (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-6-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.03 (s, 1H), 7.53 – 7.30 (m, 2H), 6.97 (s, 2H), 6.75 – 6.55 (m, 3H), 6.44 (d, J = 3.6 Hz, 1H), 5.57 – 5.31 (m, 2H), 4.98 (dd, J = 6.4, 1.8 Hz, 2H), 4.45 (t, J = 6.0 Hz, 1H), 3.98 (q, J = 5.6 Hz, 1H), 2.98 (q, J = 6.7 Hz, 2H), 2.59 – 2.52 (m, 2H); LCMS m/z = 455.10 (M⁺, 90%)
Compound-26  (1S,2R,5R)-3-(2-(2-amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-8-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.24 (d, J = 1.6 Hz, 1H), 8.04 (s, 1H), 7.44 (d, J = 8.3 Hz, 1H), 7.17 (dd, J = 8.3, 6.5 Hz, 1H), 7.00 (bs, 4H), 6.66 (d, J = 3.6 Hz, 1H), 6.44 (d, J = 3.5 Hz, 1H), 5.51 (d, J = 4.5 Hz, 1H), 5.42 (d, J = 1.8 Hz, 1H),



yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)-3-chloro-8-fluoroquinolin-2-amine	4.96 (dd, J = 6.6, 4.8 Hz, 2H), 4.44 (t, J = 5.9 Hz, 1H), 3.98 (q, J = 5.6 Hz, 1H), 2.98 (d, J = 8.3 Hz, 2H), 2.49 – 2.40 (m, 2H); LCMS m/z= 455.05 (M ⁺ , 100%)
Compound-27  (1S,2R,5R)-3-(2-(2-amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-((3aS,4R,6aR)-6-(2-(2-amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.04 (s, 1H), 7.02 (d, J = 7.3 Hz, 1H), 6.97 – 6.85 (m, 4H), 6.81 (d, J = 1.4 Hz, 1H), 6.67 (dd, J = 7.4, 1.5 Hz, 1H), 6.63 (d, J = 3.5 Hz, 1H), 6.46 (d, J = 3.5 Hz, 1H), 5.55 – 5.46 (m, 1H), 5.42 (t, J = 1.7 Hz, 1H), 4.95 (s, 2H), 4.44 (d, J = 5.5 Hz, 1H), 3.93 (q, J = 4.8 Hz, 1H), 2.87 – 2.62 (m, 2H), 2.49 – 2.41 (m, 2H), 1.30 – 1.17 (m, 6H); LCMS m/z= 418.41 (M ⁺ , 100%)
Compound-28  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2'-aminospiro[cyclobutane-1,3'-indol]-6'-yl)ethyl)cyclopent-3-ene-1,2-diol	6'-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)spiro[cyclobutane-1,3'-indol]-2'-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.04 (s, 1H), 7.39 (d, J = 7.3 Hz, 1H), 7.07 (s, 2H), 6.92 (s, 2H), 6.77 (d, J = 1.4 Hz, 1H), 6.71 (dd, J = 7.5, 1.5 Hz, 1H), 6.62 (d, J = 3.5 Hz, 1H), 6.45 (d, J = 3.5 Hz, 1H), 5.49 (d, J = 4.3 Hz, 1H), 5.41 (d, J = 1.9 Hz, 1H), 4.93 (d, J = 5.9 Hz, 2H), 4.45 (d, J = 5.5 Hz, 1H), 3.93 (q, J = 5.3 Hz, 1H), 2.87 – 2.64 (m, 2H), 2.62 – 2.49 (m, 4H), 2.40 – 2.30 (m, 1H), 2.24 – 2.13 (m, 3H); LCMS m/z= 431.23 (M ⁺ , 50%)
Compound-29  (1S,2R,5R)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3-(2-(2'-aminospiro[cyclobutane-1,3'-indol]-6'-yl)ethyl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)spiro[cyclobutane-1,3'-indol]-2'-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.20 (s, 1H), 8.03 (s, 1H), 7.35 (d, J = 8.7 Hz, 2H), 7.14 – 6.88 (m, 4H), 6.62 (d, J = 3.5 Hz, 1H), 6.43 (d, J = 3.6 Hz, 1H), 5.49 (s, 1H), 5.43 (s, 1H), 4.98 (dd, J = 11.2, 6.4



(1S,2R,5R)-3-(2-(2-amino-3,5-dichloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)-3,5-dichloroquinolin-2-amine	Hz, 2H), 4.46 (t, $J = 5.9$ Hz, 1H), 3.96 (q, $J = 5.5$ Hz, 1H), 3.04 – 2.85 (m, 2H), 2.62 – 2.56 (m, 2H); LCMS $m/z = 471.30$ (M^+ , 50%)
Compound-30 	7-(2-((3aS,4R,6aR)-4-(2-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)-3-chloroquinolin-2-amine	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.69 (s, 1H), 8.37 (s, 1H), 7.69 (d, $J = 8.2$ Hz, 1H), 7.42 (s, 1H), 7.26 (d, $J = 8.3$ Hz, 1H), 6.95 (d, $J = 3.9$ Hz, 1H), 6.46 (d, $J = 3.8$ Hz, 1H), 5.42 (dd, $J = 10.5, 3.3$ Hz, 2H), 5.30 – 4.81 (m, 2H), 4.47 (d, $J = 5.5$ Hz, 1H), 4.22 – 3.88 (m, 1H), 3.09 – 2.87 (m, 2H) 2.57 (d, $J = 7.7$ Hz, 2H); LCMS $m/z = 437.21$ ($M+1$, 70%)
Compound-31 	3-chloro-5-fluoro-7-(2-((3aS,4R,6aR)-4-(4-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)quinolin-2-amine	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.68 (s, 1H), 8.18 (s, 1H), 7.22 (s, 1H), 7.05 (d, $J = 3.7$ Hz, 1H), 7.02 (dd, $J = 11.0, 1.4$ Hz, 1H), 6.96 (s, 2H), 6.57 (d, $J = 3.6$ Hz, 1H), 5.63 (d, $J = 4.5$ Hz, 1H), 5.46 (d, $J = 1.8$ Hz, 1H), 5.01 (d, $J = 6.5$ Hz, 2H), 4.47 (t, $J = 6.1$ Hz, 1H), 4.10 - 3.98 (m, 1H), 3.40 (dt, $J = 13.9, 6.9$ Hz, 1H), 3.04 - 2.89 (m, 2H), 2.57 (d, $J = 8.3$ Hz, 2H), 1.32 (d, $J = 2.1$ Hz, 3H), 1.30 (d, $J = 2.1$ Hz, 3H); LCMS $m/z = 483.92$ (M^+ , 30%)
Compound-32	3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-(1-methyl-1H-pyrazol-4-	^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.68 (s, 1H), 8.62 (s, 1H), 8.21 (d, $J = 19.9$ Hz, 2H), 7.26 - 7.17 (m, 2H), 7.03 (dd, $J = 11.0, 1.4$ Hz, 2H), 6.98 (s, 1H), 6.87 (d, J




 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)-5-fluoroquinolin-2-amine	= 3.7 Hz, 1H), 5.66 (d, J = 4.6 Hz, 1H), 5.50 (t, J = 1.7 Hz, 1H), 5.04 (d, J = 6.1 Hz, 2H), 4.48 (s, 1H), 4.08 (d, J = 5.4 Hz, 1H), 3.97 (s, 3H), 3.06 – 2.88 (m, 2H), 2.60 – 2.55 (m, 2H); LCMS m/z= 520.32 (M⁺, 100%)
Compound-33a and 33b  (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)propyl)-3-chloro-5-fluoroquinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. First Diastereomer(Compound-33a) : ¹H NMR (400 MHz, DMSO-d₆) δ 8.18 (s, 1H), 8.03 (s, 1H), 7.18 (s, 1H), 7.14 – 6.83 (m, 5H), 6.50 (d, J = 3.5 Hz, 1H), 6.42 (d, J = 3.5 Hz, 1H), 5.51 (d, J = 5.1 Hz, 1H), 5.36 (d, J = 1.8 Hz, 1H), 5.00 (d, J = 6.8 Hz, 1H), 4.89 (d, J = 6.2 Hz, 1H), 4.56 (t, J = 5.8 Hz, 1H), 3.91 (d, J = 5.5 Hz, 1H), 3.08 – 2.94 (m, 1H), 2.79 (d, J = 9.3 Hz, 2H), 1.09 (d, J = 6.7 Hz, 3H); LCMS m/z= 469.36 (M⁺, 30%) Second Diastereomer (Compound-33b) : ¹H NMR (400 MHz, DMSO-d₆) δ 8.18 (s, 1H), 8.05 (s, 1H), 7.19 (s, 1H), 7.08 – 6.86 (m, 5H), 6.78 (d, J = 3.5 Hz, 1H), 6.50 (d, J = 3.5 Hz, 1H), 5.56 – 5.47 (m, 1H), 5.43 (t, J = 1.4 Hz, 1H), 4.97 (t, J = 5.5 Hz, 2H), 4.51 – 4.42 (m, 1H), 3.98 (q, J = 6.1, 5.3 Hz, 1H), 3.09 – 2.96 (m, 1H),




		2.75 (s, 2H), 1.08 (d, J = 6.3 Hz, 3H); LCMS m/z= 469.36 (M ⁺ , 30%)
Compound-34a and 34b  (1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)propyl)-3-chloroquinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. First Diastereomer (Compound-34a) : ¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.03 (s, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.33 (d, J = 1.5 Hz, 1H), 7.13 (dd, J = 8.2, 1.6 Hz, 1H), 6.94 (s, 2H), 6.67 (s, 2H), 6.47 (d, J = 3.5 Hz, 1H), 6.39 (d, J = 3.5 Hz, 1H), 5.56 – 5.46 (m, 1H), 5.35 (d, J = 1.8 Hz, 1H), 4.98 (d, J = 6.7 Hz, 1H), 4.88 (d, J = 6.3 Hz, 1H), 4.57 (t, J = 5.9 Hz, 1H), 3.90 (q, J = 5.7 Hz, 1H), 3.10 – 2.96 (m, 1H), 2.84 – 2.73 (m, 2H), 1.13 – 1.09 (m, 3H); LCMS m/z= 451.2 (M⁺, 100%) Second Diastereomer(Compound-34b) : ¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.05 (s, 1H), 7.60 (d, J = 8.2 Hz, 1H), 7.34 (d, J = 1.6 Hz, 1H), 7.13 (dd, J = 8.2, 1.6 Hz, 1H), 6.94 (s, 2H), 6.77 (d, J = 3.6 Hz, 1H), 6.67 (s, 2H), 6.49 (d, J = 3.5 Hz, 1H), 5.59 – 5.49 (m, 1H), 5.47 – 5.38 (m, 1H), 4.95 (t, J = 7.2 Hz, 2H), 4.47 (t, J = 6.0 Hz, 1H), 3.98 (q, J = 5.5 Hz, 1H), 3.04 (dd, J = 12.0, 4.5 Hz, 1H), 2.82 – 2.64 (m, 2H), 1.07 (d, J = 6.3 Hz, 3H); LCMS m/z= 452.98 (M⁺, 80%)

Compound-35a and 35b  (1S,2R,5R)-3-(1-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	3-chloro-7-(2-((3aS,4R,6aR)-2,2-dimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)propyl)-5-fluoroquinolin-2-amine	Diastereomeric mixture was separated by chiral preparative HPLC. First Diastereomer (Compound-35a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.63 (s, 1H), 8.18 (s, 1H), 7.27 – 7.12 (m, 2H), 6.96 (s, 3H), 6.64 (d, J = 3.6 Hz, 1H), 5.64 (d, J = 4.9 Hz, 1H), 5.46 (d, J = 2.0 Hz, 1H), 5.01 (dd, J = 6.5, 3.4 Hz, 2H), 4.46 (t, J = 5.9 Hz, 1H), 4.03 (q, J = 5.7 Hz, 1H), 3.02 (q, J = 9.1 Hz, 1H), 2.86 – 2.68 (m, 2H), 2.64 (s, 3H), 1.09 (d, J = 7.0 Hz, 3H); LCMS m/z = 468.36 (M⁺, 80%) Second Diastereomer (Compound-35b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.60 (s, 1H), 8.19 (s, 1H), 7.19 (s, 1H), 7.04 – 6.89 (m, 4H), 6.52 (d, J = 3.6 Hz, 1H), 5.63 (d, J = 5.0 Hz, 1H), 5.39 (d, J = 1.8 Hz, 1H), 5.08 – 5.00 (m, 1H), 4.95 (d, J = 6.2 Hz, 1H), 4.58 (t, J = 5.9 Hz, 1H), 3.98 (dt, J = 6.8, 5.4 Hz, 1H), 3.03 (q, J = 10.1 Hz, 1H), 2.81 (s, 2H), 2.62 (s, 3H), 1.11 (d, J = 5.3 Hz, 3H); LCMS m/z = 468.36 (M⁺, 80%)
Compound-36a and 36b  (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-2-	3-chloro-5-fluoro-7-(2-((3aS,4R,6aR)-2,2,6a-trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo	Diastereomeric mixture was separated by chiral preparative HPLC. First Diastereomer(Compound 36a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.63 (s, 1H), 8.19 (s, 1H), 7.22 – 7.12 (m, 2H), 7.03 – 6.90 (m, 3H), 6.66 (d, J = 3.7 Hz, 1H), 5.64 – 5.54 (m, 2H), 5.05 (d, J = 7.1

methyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	1-6-yl)propyl)quinolin-2-amine	Hz, 1H), 4.49 (s, 1H), 3.66 (t, J = 6.5 Hz, 1H), 2.96 (dd, J = 13.2, 7.1 Hz, 1H), 2.85 – 2.68 (m, 2H), 2.65 (s, 3H), 1.24 (s, 3H), 1.13 (d, J = 6.5 Hz, 3H); LCMS m/z= 482.2 (M⁺, 40%) Second Diastereomer (Compound 36b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.64 (s, 1H), 8.17 (s, 1H), 7.43 (d, J = 3.6 Hz, 1H), 7.21 (s, 1H), 7.00 (dd, J = 11.1, 1.3 Hz, 1H), 6.95 (s, 2H), 6.69 (d, J = 3.6 Hz, 1H), 5.68 – 5.61 (m, 1H), 5.46 (d, J = 1.4 Hz, 1H), 5.18 (d, J = 7.3 Hz, 1H), 4.63 (s, 1H), 3.90 (t, J = 6.9 Hz, 1H), 3.23 – 3.16 (m, 1H), 2.73 (s, 2H), 2.65 (s, 3H), 1.42 (s, 3H), 0.99 (d, J = 6.6 Hz, 3H); LCMS m/z= 482.2 (M⁺, 40%).
Compound-37  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.12 (s, 1H), 7.34 (s, 2H), 7.22 (s, 1H), 7.07 – 6.92 (m, 4H), 6.65 – 6.57 (m, 1H), 5.56 – 5.41 (m, 2H), 5.15 – 4.99 (m, 1H), 4.62 (s, 1H), 3.83 (d, J = 6.0 Hz, 1H), 2.99 – 2.86 (m, 2H), 2.46 – 2.31 (m, 2H), 1.27 (s, 3H); LCMS m/z= 469.30 (M⁺, 40%)
Compound-38 	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-	First Diastereomer: ¹H NMR (400 MHz, DMSO-d₆) δ 8.18 (s, 1H), 8.04 (s, 1H), 7.18 (s, 1H), 7.01 – 6.90 (m, 5H), 6.70 (d, J = 3.6 Hz, 1H), 6.47 (d, J = 3.5 Hz, 1H),


(1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol	2,2,6a-trimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)propyl)-3-chloro-5-fluoroquinolin-2-amine	5.55 (d, J = 1.7 Hz, 1H), 5.47 (d, J = 5.9 Hz, 1H), 5.06 (d, J = 7.0 Hz, 1H), 4.44 (s, 1H), 3.63 (dd, J = 7.0, 5.8 Hz, 1H), 2.97 (dd, J = 12.9, 6.7 Hz, 1H), 2.82 - 2.68 (m, 2H), 1.24 (s, 3H), 1.11 (s, 3H); LCMS m/z= 482.67 (M ⁺ , 80%)
Compound-39  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-2-methyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	3-chloro-5-fluoro-7-(2-((3aS,4R,6aR)-2,2,6a-trimethyl-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)quinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.63 (s, 1H), 8.17 (s, 1H), 7.38 (d, J = 3.7 Hz, 1H), 7.22 (s, 1H), 7.02 (dd, J = 11.0, 1.5 Hz, 1H), 6.96 (s, 2H), 6.67 (d, J = 3.5 Hz, 1H), 5.64 – 5.57 (m, 1H), 5.50 – 5.45 (m, 1H), 5.07 (d, J = 7.3 Hz, 1H), 4.65 (s, 1H), 3.89 (dd, J = 7.4, 6.1 Hz, 1H), 2.94 (t, J = 7.8 Hz, 2H), 2.64 (s, 3H), 2.57 – 2.53 (m, 2H), 1.28 (s, 3H); LCMS m/z= 468.36 (M ⁺ , 60%)
Compound-40  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-ethylcyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-6a-ethyl-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxo 1-6-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (s, 1H), 8.05 (s, 1H), 7.22 (s, 1H), 7.07 – 6.89 (m, 6H), 6.56 (d, J = 3.6 Hz, 1H), 5.57 – 5.43 (m, 2H), 5.04 (d, J = 7.6 Hz, 1H), 4.56 (s, 1H), 3.95 (t, J = 6.4 Hz, 1H), 2.94 (t, J = 8.0 Hz, 2H), 2.48 – 2.42 (m, 2H), 1.69 (dq, J = 14.4, 7.2 Hz, 1H), 1.56 (dq, J = 14.4, 7.2 Hz, 1H), 0.85 (t, J = 7.2 Hz, 3H); LCMS m/z= 482.67 (M ⁺ , 80%)
Compound-41	3-chloro-7-(2-((3aS,4R,6aR)-4-(4-ethyl-7H-pyrrolo[2,3-	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.66 (s, 1H), 8.18 (s, 1H), 7.22 (s, 1H), 7.11 – 6.91 (m, 4H), 6.56 (d, J = 3.6 Hz, 1H),

 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-ethyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	5.63 (s, 1H), 5.46 (d, J = 1.9 Hz, 1H), 5.02 (dd, J = 6.5, 4.3 Hz, 2H), 4.47 (t, J = 6.0 Hz, 1H), 4.08 – 4.00 (m, 1H), 2.96 (q, J = 7.7 Hz, 4H), 2.57 (d, J = 7.2 Hz, 2H), 1.31 (d, J = 7.6 Hz, 3H); LCMS m/z= 468.08 (M⁺, 100%)
Compound-42  (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	3-Chloro-7-(2-((3aS,4R,6aR)-4-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.63 (s, 1H), 8.27 (s, 1H), 7.30 (d, J = 36.5 Hz, 3H), 7.18 (d, J = 3.7 Hz, 1H), 7.08 (d, J = 10.9 Hz, 1H), 6.76 (d, J = 3.7 Hz, 1H), 5.62 (d, J = 3.7 Hz, 1H), 5.47 (d, J = 1.9 Hz, 1H), 4.47 (d, J = 5.5 Hz, 1H), 4.06 (t, J = 5.2 Hz, 1H), 3.07 – 2.88 (m, 2H), 2.62 – 2.54 (m, 3H) 1.27 – 1.14 (m, 4H); LCMS m/z= 479.86 (M⁺, 100%)
Compound-43  (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-5-fluoroquinolin-2-amine	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-6-yl)ethyl)-3-bromo-5-fluoroquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.34 (s, 1H), 8.04 (s, 1H), 7.21 (s, 1H), 7.06 – 6.94 (m, 3H), 6.86 (s, 2H), 6.65 (d, J = 3.5 Hz, 1H), 6.43 (d, J = 3.5 Hz, 1H), 5.49 (d, J = 4.4 Hz, 1H), 5.45 (t, J = 1.7 Hz, 1H), 4.96 (dd, J = 6.5, 2.1 Hz, 2H), 4.45 (t, J = 6.0 Hz, 1H), 3.97 (q, J = 5.5 Hz, 1H), 3.04 – 2.86 (m, J = 7.4 Hz, 2H), 2.55 (d, J = 6.4 Hz, 2H); LCMS m/z= 499.30, 501.24 (M⁺, M+2, 100%)

Compound-44  (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol		¹H NMR (400 MHz, DMSO-d₆) δ 8.61 (s, 1H), 8.34 (s, 1H), 7.21 (s, 1H), 7.06 (d, J = 3.6 Hz, 1H), 7.02 – 6.98 (m, 1H), 6.87 (s, 2H), 6.54 (d, J = 3.6 Hz, 1H), 5.61 (t, J = 3.3 Hz, 1H), 5.46 (q, J = 1.6 Hz, 1H), 5.01 (dd, J = 6.5, 5.2 Hz, 2H), 4.47 (t, J = 6.0 Hz, 1H), 4.08 – 3.99 (m, 1H), 3.07 – 2.86 (m, 2H), 2.62 (s, 3H), 2.57 (t, J = 8.3 Hz, 2H); LCMS m/z= 498.24, 500.24 (M⁺, M+2, 100%).
Compound-45a and 45b  (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol	7-(2-((3aS,4R,6aR)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl-3a,6a-dihydrocyclopenta[d][1,3]dioxo-1-yl)propyl)-3-bromo-5-fluoroquinolin-2-amine	First Diastereomer(Compound 45a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.34 (s, 1H), 8.03 (s, 1H), 7.18 (s, 1H), 7.11 – 6.68 (m, 5H), 6.52 (d, J = 3.7 Hz, 1H), 6.41 (d, J = 3.4 Hz, 1H), 5.51 (d, J = 4.8 Hz, 1H), 5.37 (s, 1H), 5.00 (d, J = 7.1 Hz, 1H), 4.89 (d, J = 6.2 Hz, 1H), 4.56 (s, 1H), 3.91 (d, J = 6.2 Hz, 1H), 3.01 (d, J = 7.4 Hz, 1H), 2.88 – 2.71 (m, 2H), 1.10 (d, J = 7.1, Hz, 3H). LCMS m/z= 514.19 (M+1, 100%) Second Diastereomer (Compound 45b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.34 (s, 1H), 8.04 (s, 1H), 7.18 (s, 1H), 7.02 – 6.92 (m, 3H), 6.86 (s, 2H), 6.78 (d, J = 3.5 Hz, 1H), 6.50 (d, J = 3.5 Hz, 1H), 5.52 (d, J = 4.8 Hz, 1H), 5.43 (s, 1H), 4.97 (dd, J = 6.5, 3.6 Hz, 2H), 4.46 (t, J = 6.1 Hz, 1H), 3.98 (q, J = 5.6 Hz, 1H), 3.01 (d, J = 9.0 Hz, 1H), 2.84 – 2.65 (m,

		2H), 1.07 (d, J = 5.9 Hz, 3H). LCMS m/z= 514.19 (M+1, 100%)
--	--	--

Example-3: (1S,2R,5R)-3-(2-(2-Amino-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-46)



5


The mixture of (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (311mg, 0.684mmol) and palladium hydroxide (168mg, 0.239mmol) in ethanol (40ml) was stirred at 25°C for 8h under hydrogen atmosphere (60 psi). The resulting mixture was filtered through celite and filtrate was concentrated *in vacuo* to give 0.32g of crude compound. Obtained residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 10%) of methanolic ammonia in dichloromethane to afford the title compound (7mg, 2.4% yield) as an off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.03 (s, 1H), 7.85 (d, J = 8.8 Hz, 1H), 7.40 (t, J = 9.3 Hz, 2H), 6.91 (s, 2H), 6.75 (d, J = 8.8 Hz, 1H), 6.65 (d, J = 3.6 Hz, 1H), 6.43 (d, J = 3.6 Hz, 1H), 6.36 (s, 2H), 5.51 (s, 1H), 5.45 (s, 1H), 4.97 (d, J = 6.3 Hz, 2H), 4.46 (s, 1H), 3.97 (d, J = 5.7 Hz, 1H), 3.05 – 2.88 (m, 2H), 2.48 – 2.26 (m, 2H).

15

LCMS m/z= 420.98 (M⁺, 50%).

Example-4: (1S,2R,5R)-3-(2-(2-amino-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (Compound-47)

20




The mixture of (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol (30 mg, 0.066 mmol), Pd-C (1.755 mg, 1.649 μmol) and ammonium formate (16.63mg, 0.264mmol) in MeOH

25

(2 ml) was stirred at 78°C for 8h. The resulting mixture was filtered through celite and filtrate was concentrated *in vacuo* to give 0.32g of crude compound. Obtained residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 10%) of methanolic ammonia in dichloromethane to afford the title compound (9mg, 32.5%) as an off-white solid. ¹H NMR (400MHz, DMSO-d₆) δ 8.03 (s, 1H), 7.97 (d, J = 9.0 Hz, 1H), 7.15 (s, 1H), 6.94 – 6.87 (m, 3H), 6.77 (d, J = 9.0 Hz, 1H), 6.64 – 6.59 (m, 3H), 6.41 (d, J = 3.5 Hz, 1H), 5.49 (d, J = 3.9 Hz, 1H), 5.44 (t, J = 1.7 Hz, 1H), 4.97 (d, J = 5.5 Hz, 2H), 4.46 (s, 1H), 3.96 (q, J = 5.3 Hz, 1H), 3.00 – 2.86 (m, 2H), 2.54 (d, J = 10.1 Hz, 2H); LCMS m/z= 420.98 (M⁺, 90%).



Example-5: (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-48)





TFA (44.3 ml, 575 mmol) was added to 7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyl tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-5-fluoro quinolin-2-amine (4.5g, 8.84 mmol) at 0°C. The resulting mixture was stirred at 25°C for 16h under N₂ atmosphere. The solvent was removed *in vacuo* at 30°C. The obtained residue was dissolved with ethyl acetate (100ml) and basified with aq. sat. NaHCO₃. Layers were separated, organic layer was washed with brine (50ml) and dried over anhydrous Na₂SO₄. The organic layer was filtered and concentrated *in vacuo* to give 5.1g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 10%) of methanol in dichloromethane to afford the title compound (2.8g, 67.5%) as a light Brown solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.16 (s, 1H), 8.07 (s, 1H), 7.21 (s, 1H), 7.04 (d, J = 3.6 Hz, 1H), 7.03 – 6.99 (m, 1H), 6.99 – 6.90 (m, 4H), 6.58 (d, J = 3.5 Hz, 1H), 5.12 (d, J = 3.8 Hz, 1H), 4.90 (s, 1H), 4.52 (d, J = 4.1 Hz, 2H), 3.72 (s, 1H), 2.96 – 2.80 (m, 2H), 2.14 (ddd, J = 16.1, 11.3, 5.3 Hz, 1H), 1.84 (ddd, J = 13.7, 11.2, 5.6 Hz, 1H), 1.24 (d, J = 5.3 Hz, 2H), 0.57 (q, J = 5.9 Hz, 1H); LCMS m/z= 469.23 (M+1, 50%).


Examples in table-15 were synthesized by following an analogous reaction protocol as was used for the preparation of (1R,2R,3S,4R,5S)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinolin-7-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol using the appropriate starting materials.



5 Table-:15



Structure & IUPAC name	Intermediate used	¹ H NMR & LCMS data
Compound-49  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(methylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0] hexane-2,3-diol	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropano [3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-N-methylquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (s, 1H), 7.91 (d, J = 9.0 Hz, 1H), 7.60 (d, J = 8.1 Hz, 1H), 7.52(s, 1H), 7.16 (d, J = 8.2 Hz, 1H), 7.12 – 7.03 (m, 2H), 6.77 (d, J = 9.0 Hz, 1H), 6.61 (d, J = 3.5 Hz, 1H), 5.12 (d, J = 4.5 Hz, 1H), 4.91 (s, 1H), 4.53 (d, J = 7.0 Hz, 2H), 3.74 (t, J = 5.0 Hz, 1H), 2.95 (d, J = 4.8 Hz, 3H), 2.87 – 2.84 (m, 2H), 2.21 – 2.05 (m, 1H), 1.88 (ddd, J = 13.7, 11.5, 5.5 Hz, 1H), 1.32 – 1.19 (m, 2H), 0.61 – 0.58 (m, 1H); ; LCMS m/z =431.06 (M+1; 60 %).
Compound-50  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(isopropylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0] hexane-2,3-diol	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropano [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-	¹H NMR (400 MHz, DMSO-d₆) δ 8.13 (s, 1H), 7.99 (s, 1H), 7.70 - 7.53 (m, 2H), 7.27 (d, J = 38.6 Hz, 3H), 7.10 (d, J = 3.6 Hz, 1H), 6.83 (d, J = 9.0 Hz, 1H), 6.66 (d, J = 3.5 Hz, 1H), 5.14 (d, J = 4.4 Hz, 1H), 4.91



yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol.	yl)ethyl)-N-isopropylquinolin-2-amine.	(d, J = 1.3 Hz, 1H), 4.54 (d, J = 3.2 Hz, 2H), 4.31 – 4.18 (m, 1H) 3.75 (s, 1H), 3.03 - 2.80 (m, 2H), 2.14 (ddd, J = 17.0, 11.9, 6.1 Hz, 1H), 1.94 - 1.79 (m, 1H), 1.27 - 1.23 (m, 8H), 0.64 - 0.54 (m, 1H); LCMS m/z = 459.3 (M+1; 100%).
Compound-51  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(cyclobutylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropano [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-cyclobutylquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.13 (s, 1H), 7.96 (d, J = 9.2 Hz, 1H), 7.62 (d, J = 8.1 Hz, 1H), 7.52 (s, 1H), 7.29 (s, 1H), 7.19 (d, J = 8.4 Hz, 1H), 7.10 (d, J = 3.6 Hz, 1H), 6.77 (d, J = 9.1 Hz, 1H), 6.65 (d, J = 3.5 Hz, 1H), 5.13 (d, J = 4.5 Hz, 1H), 4.91 (d, J = 1.3 Hz, 1H), 4.57 - 4.45 (m, 3H), 3.75 (s, 1H), 3.17 (s, 1H), 2.95 - 2.85 (m, 2H), 2.45 - 2.33 (m, 2H), 2.13 (ddd, J = 13.4, 11.3, 5.2 Hz, 1H), 2.06 - 1.81 (m, 2H), 1.74 (ddd, J = 15.3, 10.1, 7.1 Hz, 2H), 1.33 - 1.21 (m, 2H), 0.64 - 0.55 (m, 1H); LCMS m/z = 471.3 (M+1; 100%).
Compound-52  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-(cyclopropylamino)quinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropano [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-cyclopropylquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.10 (s, 1H), 7.89 (d, J = 9.0 Hz, 1H), 7.58 (d, J = 8.1 Hz, 1H), 7.46 (s, 1H), 7.18 – 7.04 (m, 4H), 6.81 (d, J = 9.0 Hz,



202



(2-((cyclopropylmethyl)amino)quinolin-7-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol	opa [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-N-(cyclopropylmethyl)quinolin-2-amine.	¹H), 6.61 (d, J = 3.5 Hz, 1H), 5.12 (d, J = 4.4 Hz, 1H), 4.91 (d, J = 1.2 Hz, 1H), 4.53 (d, J = 5.6 Hz, 2H), 3.74 (t, J = 4.8 Hz, 1H), 3.31 – 3.28 (m, 1H), 2.94 – 2.83 (m, 2H), 2.12 (ddd, J = 13.7, 11.5, 5.2 Hz, 1H), 1.88 (ddd, J = 13.7, 11.5, 5.4 Hz, 1H), 1.28 – 1.23 (m, 3H), 1.19 – 1.08 (m, 1H), 0.59 (q, J = 5.9 Hz, 1H), 0.53 – 0.45 (m, 2H), 0.33 – 0.22 (m, 2H); LCMS m/z = 471.05(M+1; 90%).
Compound-53  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-amino-8-fluoroquinolin-7-yl)ethyl)bicyclo [3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa [3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-8-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.10 (s, 1H), 7.90 (dd, J = 9.0, 1.6 Hz, 1H), 7.38 (d, J = 8.2 Hz, 1H), 7.30 – 7.00 (m, 4H), 6.77 (d, J = 9.1 Hz, 3H), 6.62 (d, J = 3.6 Hz, 1H), 5.15 (d, J = 4.5 Hz, 1H), 4.91 (s, 1H), 4.53 (d, J = 3.0 Hz, 2H), 3.75 (s, 1H), 2.97 (td, J = 12.6, 5.1 Hz, 1H), 2.84 (td, J = 12.6, 5.3 Hz, 1H), 2.19 – 2.01 (m, 1H), 1.82 (td, J = 12.7, 5.2 Hz, 1H), 1.28 (d, J = 3.5 Hz, 2H), 0.63 – 0.54 (m, 1H); LCMS m/z = 435.04 (M+1; 40 %).
Compound-54	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-	¹H NMR (400 MHz, DMSO-d₆) δ 8.07 (s, 1H), 7.71 (s, 1H), 7.51 (d, J = 8.2 Hz, 1H), 7.33



 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol.	d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopenta[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-methylquinolin-2-amine.	(d, J = 1.6 Hz, 1H), 7.08 (dd, J = 8.2, 1.7 Hz, 1H), 7.03 (d, J = 3.6 Hz, 1H), 6.98 (s, 2H), 6.58 (d, J = 3.5 Hz, 1H), 6.36 (s, 2H), 5.12 (d, J = 4.5 Hz, 1H), 4.90 (d, J = 1.2 Hz, 1H), 4.59 – 4.46 (m, 2H), 3.72 (t, J = 5.2 Hz, 1H), 2.92 – 2.83 (m, 2H), 2.20 (d, J = 1.1 Hz, 3H), 2.10 (ddd, J = 12.7, 11.0, 5.2 Hz, 1H), 1.88 (ddd, J = 13.8, 11.5, 5.5 Hz, 1H), 1.23 (d, J = 3.5 Hz, 2H), 0.58 (q, J = 5.5 Hz, 1H); LCMS m/z = 431.08 (M+1; 100%).
Compound-55  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-isopropyl quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopenta[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-isopropylquinolin-2-amine	¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (d, J = 6.3 Hz, 2H), 7.74 (d, J = 8.2 Hz, 3H), 7.45 (s, 1H), 7.30 (dd, J = 8.1, 1.5 Hz, 1H), 7.07 (dd, J = 9.1, 3.6 Hz, 3H), 6.61 (d, J = 3.5 Hz, 1H), 5.16 (d, J = 4.5 Hz, 1H), 4.90 (d, J = 1.2 Hz, 1H), 4.53 (d, J = 7.5 Hz, 2H), 3.74 (t, J = 5.1 Hz, 1H), 3.10 (p, J = 6.7 Hz, 1H), 3.01 – 2.83 (m, 2H), 2.14 (ddd, J = 15.8, 12.1, 6.4 Hz, 1H), 1.85 (ddd, J = 13.6, 11.1, 5.9 Hz, 1H), 1.26 (d, J = 6.6 Hz, 8H), 0.56 (q, J = 6.0 Hz, 1H); LCMS m/z = 459.3 (M⁺; 40%).



Compound-56  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(1,1-difluoroethyl) quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-(1,1-difluoroethyl)quinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.09 (s, 1H), 7.70 (d, J = 8.1 Hz, 1H), 7.38 (s, 1H), 7.21 – 7.15 (m, 1H), 7.10 – 7.00 (m, 3H), 6.61 (d, J = 3.6 Hz, 1H), 6.25 (s, 2H), 5.11 (d, J = 4.5 Hz, 1H), 4.91 (s, 1H), 4.53 (d, J = 7.5 Hz, 2H), 3.74 (t, J = 5.2 Hz, 1H), 2.83 – 2.91 (m, 2H), 2.18 – 1.98 (m, 4H), 1.88 (td, J = 12.4, 5.5 Hz, 1H), 1.28 – 1.25 (m, 2H), 0.58 (q, J = 5.9 Hz, 1H); LCMS m/z = 481.2 (M+1; 90%).
Compound-57  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-cyclopropylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-cyclopropylquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (s, 1H), 7.91 (s, 1H), 7.80 (s, 2H), 7.67 (d, J = 8.2 Hz, 1H), 7.44 (s, 1H), 7.27 (dd, J = 8.2, 1.5 Hz, 1H), 7.11 (s, 2H), 7.06 (d, J = 3.5 Hz, 1H), 6.61 (d, J = 3.5 Hz, 1H), 5.17 (d, J = 4.5 Hz, 1H), 4.90 (d, J = 1.2 Hz, 1H), 4.53 (d, J = 4.3 Hz, 2H), 3.73 (s, 1H), 2.95 – 2.87 (m, 2H), 2.12 (ddd, J = 13.8, 11.0, 5.4 Hz, 1H), 1.90 – 1.74 (m, 2H), 1.23 (d, J = 3.4 Hz, 2H), 1.08 – 0.96 (m, 2H), 0.78 – 0.68 (m, 2H), 0.55 (q, J = 5.8 Hz, 1H); LCMS m/z = 457.13 (M+; 50%).



Compound-58  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methoxyquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-methoxyquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 7.54 (d, J = 8.1 Hz, 1H), 7.40 – 7.31 (m, 2H), 7.15 – 7.05 (m, 3H), 7.04 (d, J = 3.5 Hz, 1H), 6.67 (d, J = 18.5 Hz, 2H), 6.60 (d, J = 3.5 Hz, 1H), 5.12 (d, J = 4.5 Hz, 1H), 4.90 (d, J = 1.3 Hz, 1H), 4.52 (d, J = 5.8 Hz, 2H), 3.91 (s, 3H), 3.79 – 3.66 (m, 1H), 2.90 – 2.80 (m, 2H), 2.09 (ddd, J = 13.7, 11.2, 5.4 Hz, 1H), 1.88 (ddd, J = 13.8, 11.5, 5.7 Hz, 1H), 1.24 (d, J = 5.2 Hz, 2H), 0.58 (q, J = 5.8 Hz, 1H).); LCMS m/z = 447.01 (M+1; 90%).
Compound-59  2-amino-7-(2-((1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,3-dihydroxybicyclo[3.1.0]hexan-1-yl)ethyl)quinoline-3-carbonitrile.	2-amino-7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)quinoline-3-carbonitrile.	¹H NMR (400 MHz, DMSO-d₆) δ 8.62 (s, 1H), 8.06 (d, J = 2.3 Hz, 1H), 7.66 (d, J = 8.3 Hz, 1H), 7.37(s, 1H), 7.22 (dd, J = 8.3, 1.6 Hz, 1H), 7.03 (d, J = 3.6 Hz, 1H), 6.97 (s, 2H), 6.89 (s, 2H), 6.58 (d, J = 3.5 Hz, 1H), 5.11 (d, J = 4.5 Hz, 1H), 4.90 (s, 1H), 4.57 – 4.48 (m, 2H), 3.73 (t, J = 5.3 Hz, 1H), 3.0 - 2.81 (m, 2H), 2.21 – 2.04 (m, 1H), 1.93 – 1.81 (m, 1H), 1.26 – 1.23 (m, 2H), 0.57 (q, J = 5.9 Hz,




		1H); LCMS m/z = 442.23 (M+1; 80%).
Compound-60  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 7.78 (d, J = 11.8 Hz, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.49 (s, 2H), 7.37 (d, J = 1.6 Hz, 1H), 7.15 (dd, J = 8.2, 1.6 Hz, 1H), 7.11 (d, J = 3.6 Hz, 1H), 6.77 (s, 2H), 6.67 (d, J = 3.6 Hz, 1H), 5.14 (d, J = 4.3 Hz, 1H), 4.90 (d, J = 1.3 Hz, 1H), 4.54 (d, J = 8.9 Hz, 2H), 3.74 (d, J = 5.5 Hz, 1H), 2.92 – 2.82 (m, 2H), 2.10 (ddd, J = 13.8, 11.2, 5.4 Hz, 1H), 1.87 (ddd, J = 13.8, 11.6, 5.6 Hz, 1H), 1.27 – 1.23 (m, 2H), 0.63 – 0.55 (m, 1H).); LCMS m/z = 435.3 (M+; 80%).
Compound-61  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.07 (s, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.36 (s, 1H), 7.15 (dd, J = 8.2, 1.6 Hz, 1H), 7.08 – 6.89 (m, 3H), 6.66 (s, 2H), 6.58 (d, J = 3.5 Hz, 1H), 5.11 (d, J = 4.5 Hz, 1H), 4.90 (s, 1H), 4.55 – 4.50 (m, 2H), 3.73 (t, J = 5.3 Hz, 1H), 2.96 – 2.80 (m, 2H), 2.11 (ddd, J = 13.7, 11.3, 5.3 Hz, 1H), 1.87 (ddd, J = 13.9, 11.6, 5.6 Hz, 1H), 1.26


		– 1.23 (m, 2H), 0.58 (q, J = 5.9 Hz, 1H); LCMS m/z = 451.3 (M+1; 80%).
Compound-62  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-6-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (d, J = 6.2 Hz, 2H), 7.56 – 7.35 (m, 4H), 7.13 (d, J = 3.6 Hz, 1H), 6.68 (d, J = 3.1 Hz, 3H), 5.16 (s, 1H), 4.91 (s, 1H), 4.53 (d, J = 7.2 Hz, 2H), 3.75 (s, 1H), 2.98 (t, J = 12.0 Hz, 1H), 2.91 – 2.78 (m, 1H), 2.13 (q, J = 8.2, 4.7 Hz, 1H), 1.89 – 1.76 (m, 1H), 1.25 (d, J = 11.4 Hz, 2H), 0.59 (q, J = 6.0 Hz, 1H); LCMS m/z = 451.3 (M+1; 80%).
Compound-63  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-8-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.23 (s, 1H), 8.07 (s, 1H), 7.42 (d, J = 8.3 Hz, 1H), 7.17 (t, J = 7.4 Hz, 1H), 7.04 (d, J = 3.6 Hz, 1H), 7.01 – 6.92 (m, 4H), 6.58 (d, J = 3.5 Hz, 1H), 5.16 (d, J = 4.6 Hz, 1H), 4.90 (s, 1H), 4.59 – 4.44 (m, 2H), 3.73 (t, J = 5.4 Hz, 1H), 3.09 – 2.92 (m, 1H), 2.92 – 2.77 (m, 1H), 2.13 (t, J = 7.2 Hz, 1H), 1.87-1.74 (m, 1H), 1.27 (t, J = 6.5 Hz, 2H), 0.58 (d, J = 5.6 Hz, 1H);

		LCMS m/z = 468.68 (M+; 80%).
Compound-64  (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-bromo-6-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.35 (s, 1H), 8.24 (s, 1H), 7.96 (d, J = 30.9 Hz, 2H), 7.45 (t, J = 8.4Hz, 1H), 7.24 (d, J = 3.6 Hz, 1H), 6.80 (d, J = 3.5 Hz, 1H), 6.60 (s, 1H), 5.17 (s, 1H), 4.91 (d, J = 4.0 Hz, 1H), 4.53 (s, 1H), 4.40 (d, J = 6.8 Hz, 1H), 3.78 (d, J = 6.4 Hz, 1H), 3.04 – 2.80(m, 2H), 2.15 (s, 1H), 1.82 (s, 1H), 1.30 – 1.27 (m, 2H), 0.60 – 0.55 (m, 1H); LCMS m/z = 514.19 (M+1; 80%).
Compound-65  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl) bicyclo[3.1.0]hexane-2,3-diol.	7-((3aR,3bR,4aS,5R,5aS)-2,2-Dimethyl-3b-(2-(3-methylimidazo[1,2-a]pyridin-7-yl)ethyl)hexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.16 (d, J = 6.9 Hz, 1H), 8.07 (s, 1H), 7.36 (s, 1H), 7.28 (s, 1H), 7.08 – 6.85 (m, 4H), 6.58 (d, J = 3.4 Hz, 1H), 5.12 (d, J = 4.5 Hz, 1H), 4.90 (s, 1H), 4.52 (s, 2H), 3.72 (s, 1H), 3.29 (s, 1H), 2.95-2.75 (m, 2H), 2.61 – 2.53 (m, 1H), 2.44 (s, 3H), 2.21 – 2.03 (m, 1H), 1.97 – 1.74 (m, 1H), 0.61 – 0.47 (m, 1H); LCMS m/z = 405.16 (M+1; 100%).
Compound-66	7-((3aR,3bR,4aS,5R,5aS)-3b-(2-(2-Amino-3,3-dimethyl-3H-indol-6-	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.06 (s, 1H), 7.09 (d, J = 7.4 Hz, 1H), 7.01 (d, J = 3.6 Hz,


 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3,3-dimethyl-3H-indol-6-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	yl)ethyl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine.	¹H), 6.98 (s, 2H), 6.87 (d, J = 1.4 Hz, 1H), 6.79 – 6.74 (m, 1H), 6.58 (d, J = 3.5 Hz, 1H), 5.11 (s, 1H), 4.90 (d, J = 1.2 Hz, 1H), 4.49 (s, 2H), 3.71 (s, 1H), 2.82 – 2.59 (m, 2H), 2.09-1.96 (m, 1H), 1.81 (td, J = 12.7, 5.3 Hz, 1H), 1.30 (s, 6H), 1.26 – 1.19 (m, 2H), 0.63 – 0.51 (m, 1H); LCMS m/z = 433.40 (M+1; 80%).
Compound-67  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-5-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.03 (s, 1H), 7.18 (s, 1H), 7.06 – 6.87 (m, 5H), 6.36 – 6.29 (m, 1H), 4.93 (d, J = 5.4 Hz, 1H), 4.74 (t, J = 6.7 Hz, 1H), 4.51 (d, J = 6.6 Hz, 1H), 4.41 (d, J = 3.4 Hz, 1H), 4.28 (s, 1H), 2.95 (td, J = 12.6, 11.3, 4.9 Hz, 1H), 2.82 (td, J = 12.8, 12.1, 5.7 Hz, 1H), 2.38 (s, 3H), 2.14 – 1.98 (m, 1H), 1.79 (ddd, J = 13.6, 11.3, 5.8 Hz, 1H), 0.99 – 0.79 (m, 2H), 0.58 (dd, J = 8.4, 4.7 Hz, 1H); LCMS m/z = 485.05 (M+2; 40%).
Compound-68	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-	¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.01 (s, 1H), 7.46 – 7.39 (m, 2H), 6.94 (s, 2H), 6.67 (s, 2H), 6.32 (d, J =

 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-3-chloro-6-fluoroquinolin-2-amine.	1.2 Hz, 1H), 4.98 – 4.91 (m, 1H), 4.76 (t, J = 6.7 Hz, 1H), 4.53 (d, J = 6.6 Hz, 1H), 4.42 (d, J = 3.4 Hz, 1H), 4.28 (s, 1H), 3.08 – 2.94 (m, 1H), 2.84 (td, J = 13.3, 12.7, 5.3 Hz, 1H), 2.43 – 2.37 (m, 3H), 2.04 (dq, J = 19.8, 7.7, 6.3 Hz, 1H), 1.86 – 1.72 (m, 1H), 1.32 – 1.26 (m, 2H), 0.59 (dd, J = 8.3, 4.7 Hz, 1H); LCMS m/z = 482.30 (M; 80%).
Compound-69  (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	3-Chloro-7-(2-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-5-fluoroquinolin-2-amine.	A mixture of two compound was separated by reverse phase HPLC to afford compound X and compound Y as shown below.¹H NMR (400 MHz, DMSO-d₆) δ 8.66 (s, 1H), 8.16 (s, 1H), 7.43 (d, J = 3.7 Hz, 1H), 7.21 (s, 1H), 7.00 (dd, J = 11.1, 1.4 Hz, 1H), 6.93 (s, 2H), 6.70 (d, J = 3.6 Hz, 1H), 5.16 (s, 1H), 4.99 (d, J = 1.4 Hz, 1H), 4.56 (d, J = 6.5 Hz, 2H), 3.80 (d, J = 5.8 Hz, 1H), 2.97 – 2.80 (m, 2H), 2.65 (s, 3H), 2.22 – 2.09 (m, 1H), 1.91 – 1.81 (m, 1H), 1.31 – 1.25 (m, 2H), 0.60 (q, J = 5.8 Hz, 1H); LCMS m/z = 468.2 (M+1; 80%).

Compound-70  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	3-Chloro-7-(2-((3aR,3bR,4aS,5R,5aS)-2,2-dimethyl-5-(7H-pyrrolo[2,3-d]pyrimidin-7-yl)tetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)-5-fluoroquinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 9.02 (s, 1H), 8.81 (s, 1H), 8.16 (s, 1H), 7.51 (d, J = 3.7 Hz, 1H), 7.22 (s, 1H), 7.03 – 6.98 (m, 1H), 6.92 (bs, 2H), 6.67 (d, J = 3.6 Hz, 1H), 5.02 (d, J = 1.5 Hz, 1H), 4.56 (d, J = 6.7 Hz, 1H), 3.81 (d, J = 6.4 Hz, 1H), 2.87 (ddd, J = 19.0, 14.0, 8.5 Hz, 2H), 2.20 – 2.10 (m, 1H), 1.88 (d, J = 10.5 Hz, 1H), 1.32 – 1.26 (m, 2H), 0.64 – 0.58 (m, 1H); LCMS m/z = 454.11 (M+1; 40%).
Compound-71  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2'-aminospiro[cyclobutane-1,3'-indol]-6'-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol.	6'-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)ethyl)spiro[cyclobutane-1,3'-indol]-2'-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.07 (s, 1H), 7.68 (d, J = 7.7 Hz, 1H), 7.22 – 7.07 (m, 4H), 7.07 – 6.96 (m, 3H), 6.59 (d, J = 3.5 Hz, 1H), 5.14 (d, J = 4.6 Hz, 1H), 4.89 (s, 1H), 4.51 (d, J = 7.2 Hz, 1H), 3.74 (t, J = 5.4 Hz, 1H), 2.79 (dd, J = 24.0, 14.1 Hz, 3H), 2.35 (d, J = 12.3 Hz, 2H), 2.20 (d, J = 9.9 Hz, 1H), 2.04 (d, J = 17.8 Hz, 1H), 1.87 – 1.74 (m, 1H), 1.26 – 1.18 (m, 5H), 0.60 – 0.50 (m, 1H); LCMS m/z = 445.03 (M+1; 90%).
Compound-72		¹H NMR (400 MHz, DMSO-d₆) δ 8.33 (s, 1H), 8.08 (s, 1H),

 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol		7.20 (s, 2H), 7.17 – 6.96 (m, 3H), 6.84 (s, 2H), 6.60 (d, J = 3.5 Hz, 1H), 5.11 (d, J = 4.5 Hz, 1H), 4.90 (d, J = 1.2 Hz, 1H), 4.52 (d, J = 2.7 Hz, 2H), 3.72 (s, 1H), 3.00 – 2.74 (m, 2H), 2.14 (s, 1H), 1.90 – 1.76 (m, 1H), 1.42 – 1.28 (m, 1H), 0.91 – 0.75 (m, 1H), 0.57 (q, J = 5.9 Hz, 1H); LCMS m/z = 514.19, 516.19 (M⁺, M+2, 100%).
---	--	---

Example-6: (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-73)



5 7-((2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)-3-bromo-N-(4-methoxy benzyl)quinolin-2-amine (2.6g, 3.97mmol) in TFA (55.0 ml, 714 mmol) was stirred at 50°C for 1h under N₂ atmosphere. The resulting mixture was concentrated *in vacuo* and obtained residue was dissolved in MeOH (50ml). K₂CO₃ (0.982 g, 7.10mmol) was added and stirred the reaction mixture at 60°C for 1h. The reaction mixture was filtered, and filtrate was concentrated under reduced pressure to get 2.7g of crude compound. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redisep® R_f column with gradient elution (0 to 9%) of methanol in dichloromethane to afford the title compound (1.35g, 77%) as a white solid. ¹H NMR (400 MHz, DMSO-d₆): δ 8.34 (s, 1H), 8.12 (s, 1H), 7.58 (d, J = 8.3 Hz, 1H), 7.36 (d, J = 1.6 Hz, 1H), 7.26 (s, 2H), 7.15 (dd, J = 8.2, 1.7 Hz, 1H), 7.09 (d, J = 3.5 Hz, 1H), 6.64 (d, J = 3.5 Hz, 1H), 6.60 (s, 2H), 5.13 (d, J = 4.6 Hz, 1H), 4.97 – 4.83 (m, 1H), 4.53 (d, J = 3.9 Hz, 2H), 3.74 (s, 1H), 2.94 – 2.81 (m,

10



15


2H), 2.11 (dt, $J = 11.2, 6.5$ Hz, 1H), 1.87 (ddd, $J = 13.9, 11.5, 5.6$ Hz, 1H), 1.33 – 1.12 (m, 2H); 0.60 – 0.57 (m, 1H). LCMS $m/z = 494.99, 496.99$ (M^+ , $M+2$; 100%).


Examples in table-16 were synthesized by following an analogous reaction protocol as was used for the preparation of (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-


- 5 4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol using the appropriate starting materials.


Table-16:

Structure & IUPAC name	Intermediate used	^1H NMR & LCMS data
Compound-74  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinolin-7-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol	7-(2-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethylhexahydrocyclopropano[3,4]cyclopenta[1,2-d][1,3]dioxol-3b-yl)ethyl)quinolin-2-amine	^1H NMR (400 MHz, DMSO-d_6) δ 8.07 (s, 1H), 7.82 (d, $J = 8.8$ Hz, 1H), 7.52 (d, $J = 8.2$ Hz, 1H), 7.30 (d, $J = 1.6$ Hz, 1H), 7.08 - 7.02 (m, 2H), 6.97 (s, 2H), 6.68 (d, $J = 8.8$ Hz, 1H), 6.58 (d, $J = 3.5$ Hz, 1H), 6.34 (s, 2H), 5.11 (d, $J = 4.5$ Hz, 1H), 4.91 (d, $J = 1.2$ Hz, 1H), 4.55 - 4.49 (m, 2H), 3.73 (t, $J = 5.2$ Hz, 1H), 2.91 - 2.78 (m, 2H), 2.10 (ddd, $J = 13.6, 11.3, 5.3$ Hz, 1H), 1.92 - 1.85 (m, 1H), 1.28 - 1.24 (m, 2H), 0.59 (td, $J = 6.6, 3.2$ Hz, 1H); LCMS $m/z = 418.17$ ($M+2$; 40%).
Compound-75  (1R,2R,3S,4R,5S)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinazolin-7-yl)ethyl)bicyclo[3.1.0]hexane-2,3-diol.	N-(7-(2-((1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,3-dihydroxybicyclo[3.1.0]hexan-1-yl)ethyl)quinazolin-2-yl)-2,2,2-trifluoroacetamide.	^1H NMR (400 MHz, DMSO-d_6) δ 9.04 (s, 1H), 8.16 (s, 1H), 7.71 (d, $J = 8.2$ Hz, 1H), 7.53 (s, 2H), 7.28 (s, 1H), 7.15 (dd, $J = 11.1, 5.8$ Hz, 2H), 6.83 (s, 2H), 6.69 (d, $J = 3.6$ Hz, 1H), 5.14 (s, 1H), 4.90 (s, 1H), 4.55 (s, 2H), 3.75 (s, 1H), 2.99 – 2.79 (m, 2H), 2.11 (d, $J = 15.9$ Hz, 1H), 1.87 (q, $J = 10.9, 8.1$ Hz, 1H), 1.29 – 1.23



		(m, 2H), 0.61 – 0.56 (m, 1H); LCMS m/z = 418.10 (M+1; 80%).
Compound-76a and 76b  (1S,2R,3S,4R,5S)-1-((S)-1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3] dioxol-3b(3aH)-yl)propyl)-3-bromo-N-(4-methoxybenzyl) quinolin-2-amine.	Diastereomeric mixture was separated by chiral preparative HPLC. <i>Wavelength:</i> 225nm, <i>Instrument Method:</i> HEX-0.1%DEA_IPA-DCM_A_C_40_60_1.2ML_10MIN <i>Flow Rate:</i> 1.2 ml/min, <i>Column :</i> CHIRALPAK IA CRL-025 <i>Column Temp:</i> 25°C, <i>Mobile Phase A:</i> HEX_0.1%DEA <i>Mobile Phase C:</i> IPA-DCM_1-1, <i>Mobile Phase B:</i> NA <i>Mobile Phase D:</i> NA First Diastereomer (Compound 76a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.33 (s, 1H), 8.06 (s, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.33 (s, 1H), 7.12 (d, J = 8.4 Hz, 1H), 7.01 – 6.87 (m, 3H), 6.64 – 6.47 (m, 3H), 5.13 (d, J = 5.2 Hz, 1H), 4.80 (d, J = 2.9 Hz, 1H), 4.61 (t, J = 6.7 Hz, 1H), 4.47 (d, J = 7.2 Hz, 1H), 3.96 – 3.85 (m, 1H), 3.22 (dd, J = 13.2, 3.9 Hz, 1H), 2.94 – 2.88 (m, 1H), 1.97 – 1.83 (m, 1H), 1.31 – 1.21 (m, 2H), 0.79 (d, J = 6.8 Hz, 3H), 0.67 – 0.59 (m, 1H); LCMS m/z = 510.94 (M+2; 40%) Second Diastereomer (Compound 76b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.32 (s, 1H), 8.06 (s, 1H), 7.57 (d, J = 8.2 Hz, 1H), 7.32 (s, 1H), 7.11

		(dd, $J = 8.2, 1.6$ Hz, 1H), 7.07 (d, $J = 3.6$ Hz, 1H), 6.97 (s, 2H), 6.58 (d, $J = 3.5$ Hz, 1H), 6.55 (s, 2H), 5.15 (d, $J = 5.1$ Hz, 1H), 4.80 (d, $J = 2.6$ Hz, 1H), 4.64 (t, $J = 7.0$ Hz, 1H), 4.43 (d, $J = 7.3$ Hz, 1H), 3.97 – 3.88 (m, 1H), 3.07 – 2.97 (m, 1H), 2.97 – 2.88 (m, 1H), 1.70 – 1.55 (m, 1H), 1.31 – 1.21 (m, 2H), 0.98 (d, $J = 6.8$ Hz, 3H), 0.61 – 0.52 (m, 1H); LCMS $m/z = 510.94$ (M+2; 40%)
Compound-77a and 77b  (1S,2R,3S,4R,5S)-1-((S)-2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)-1-cyclopropylethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	7-(2-((3aR,3bS,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3] dioxol-3b(3aH)-yl)-2-cyclopropyl ethyl)-3-chloro-5-fluoro-N-(4-methoxybenzyl)quinolin-2-amine.	Diastereomeric mixture was separated by chiral preparative HPLC. <i>Wavelength:</i> 225 nm, <i>Instrument Method:</i> MeOH_0.1%DEA_100_1.0ML_12MIN <i>Flow Rate:</i> 1.00 ml/min <i>Column :</i> CHIRALPAK IE CRL-042 <i>Column Temp:</i> 30°C, <i>Mobile Phase A:</i> MEOH_0.1%DEA, <i>Mobile Phase B:</i> NA First Diastereomer (Compound 77a): ^1H NMR (400 MHz, DMSO-d_6) δ 8.39 (s, 1H), 8.16 (s, 1H), 8.04 (s, 1H), 7.28 (s, 1H), 7.05 (dd, $J = 11.3, 1.3$ Hz, 1H), 6.93 (dd, $J = 11.6, 8.0$ Hz, 4H), 6.52 (d, $J = 3.5$ Hz, 1H), 5.28 – 5.08 (m, 1H), 4.82 (d, $J = 2.8$ Hz, 1H), 4.61 (d, $J = 6.7$ Hz, 1H), 3.86 (s, 1H), 2.90 (dd, $J = 13.3, 8.5$ Hz, 1H), 2.77 (q, $J = 7.2$ Hz, 2H), 2.68 (q, $J = 1.8$ Hz, 1H),

		1.37 (dd, J = 8.8, 4.0 Hz, 1H), 1.27 – 1.01 (m, 3H), 0.66 (dd, J = 8.6, 4.9 Hz, 1H), 0.46 (s, 1H), 0.26 – 0.17 (m, 1H), 0.16 – 0.01 (m, 1H), LCMS m/z = 509.4 (M+1; 50%) Second Diastereomer (Compound 77b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.05 (s, 1H), 7.31 – 7.17 (m, 2H), 7.08 – 6.81 (m, 5H), 6.59 (d, J = 3.5 Hz, 1H), 5.18 (s, 1H), 4.82 (dd, J = 9.3, 4.6 Hz, 2H), 4.54 (s, 1H), 3.92 (d, J = 6.7 Hz, 1H), 3.16 (q, J = 6.5 Hz, 1H), 2.99 – 2.89 (m, 1H), 1.22 – 1.06 (m, 4H), 0.57 (q, J = 8.5, 7.9 Hz, 1H), 0.50 – 0.34 (m, 1H), 0.26 (ddd, J = 13.1, 8.4, 4.7 Hz, 2H), 0.06 (dd, J = 9.4, 4.7 Hz, 1H); LCMS m/z = 509.4 (M+1; 50%)
Compound-78a and 78b  (1S,2R,3S,4R,5S)-1-(1-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	N-(7-(2-((1S,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,3-dihydroxybicyclo[3.1.0]hexan-1-yl)propyl)-3-chloro-5-fluoroquinolin-2-yl)-2,2,2-trifluoroacetamide.	First Diastereomer(Compound-78a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.15 (s, 2H), 8.06 (s, 1H), 7.17 (s, 1H), 7.07 (d, J = 3.6 Hz, 1H), 7.04 – 6.90 (m, 4H), 6.58 (d, J = 3.5 Hz, 1H), 5.16 (d, J = 5.1 Hz, 1H), 4.79 (d, J = 2.6 Hz, 1H), 4.63 (t, J = 7.1 Hz, 1H), 4.45 (d, J = 7.3 Hz, 1H), 3.93 (d, J = 6.9 Hz, 1H), 2.99 (dd, J = 13.3, 4.9 Hz, 1H), 2.61 (d, J = 11.2 Hz, 1H), 1.63 (s, 1H), 1.17 (d, J = 5.8 Hz, 2H), 0.99 (d, J =


		6.7 Hz, 3H), 0.57 (d, J = 4.3 Hz, 1H); LCMS m/z = 485.03 (M+2; 40%) Second Diastereomer (Compound-78b): ¹H NMR (400 MHz, DMSO-d₆) δ 8.16 (s, 1H), 8.08 (s, 1H), 7.19 (s, 1H), 7.10 (s, 2H), 6.98 (dd, J = 10.0, 2.2 Hz, 2H), 6.92 (s, 2H), 6.56 (d, J = 3.6 Hz, 1H), 5.13 (d, J = 5.3 Hz, 1H), 4.80 (d, J = 2.9 Hz, 1H), 4.61 (t, J = 6.7 Hz, 1H), 4.48 (d, J = 7.1 Hz, 1H), 3.90 (s, 1H), 3.21 (dd, J = 13.3, 4.1 Hz, 1H), 2.68 (p, J = 1.9 Hz, 1H), 2.36 – 2.32 (m, 1H), 1.17 (t, J = 4.3 Hz, 2H), 0.79 (d, J = 6.8 Hz, 3H), 0.63 (dd, J = 8.6, 4.7 Hz, 1H); LCMS m/z = 483.02 (M+; 90%)
Compound-79a and 79b  (1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(2-(2-aminoquinolin-7-yl)propyl)bicyclo[3.1.0]hexane-2,3-diol.	7-(1-((3aR,3bR,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3] dioxol-3b(3aH)-yl)propan-2-yl)-N-(4-methoxybenzyl)quinolin-2-amine.	Diastereomeric mixture was separated by chiral preparative HPLC. Wavelength: 225 nm, Instrument Method: MeOH_0.1%DEA_A_1.0ML_10MIN Flow Rate: 1.00 ml/min, Column : CHIRALPAK IB CRL-043 Column Temp: 30°C, Mobile Phase A: MeOH_0.1%DEA, Mobile Phase B: NA First Diastereomer(Compound 79a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.13 – 8.03 (m, 2H), 7.66 (d, J = 8.2 Hz, 1H), 7.53 – 7.43 (m, 3H) 7.31 (d,

		J= 8.3 Hz, 1H), 7.03 (s, 2H), 6.93 (d, J = 3.5 Hz, 1H), 6.84 (d, J = 9.1 Hz, 1H), 6.56 (d, J = 3.5 Hz, 1H), 5.15 (d, J = 4.7 Hz, 1H), 4.76 (d, J = 1.4 Hz, 1H), 4.49 – 4.32 (m, 2H), 3.70 (t, J = 5.5 Hz, 1H), 3.30 – 3.18 (m, 1H), 2.25 (dd, J = 14.1, 4.8 Hz, 1H), 1.82 (dd, J = 14.2, 9.1 Hz, 1H), 1.31 – 1.22 (m, 4H), 1.02 – 0.91 (m, 1H), 0.64 (dd, J = 8.5, 4.7 Hz, 1H); LCMS m/z = 430.92 (M+1; 90%). Second Diastereomer(Compound 79a): ¹H NMR (400 MHz, DMSO-d₆) δ 8.02 (d, J = 13.7 Hz, 2H), 7.64 (d, J = 8.2 Hz, 1H), 7.37 (d, J = 1.6 Hz, 1H), 7.21 (dd, J = 8.2, 1.6 Hz, 3H), 6.98 (s, 2H), 6.81 (d, J = 9.0 Hz, 1H), 6.69 (d, J = 3.6 Hz, 1H), 6.42 (d, J = 3.5 Hz, 1H), 5.16 (d, J = 4.8 Hz, 1H), 4.83 (d, J = 1.7 Hz, 1H), 4.47 (d, J = 2.4 Hz, 2H), 3.68 (s, 1H), 2.37 – 2.27 (m, 1H), 1.68 – 1.57 (m, 1H), 1.39 – 1.26 (m, 5H), 1.04 (dd, J = 8.6, 3.8 Hz, 1H), 0.69 – 0.60 (m, 1H); LCMS m/z = 430.98 (M+1; 90%)
--	--	--

Compound-80  (1R,2R,3S,4R,5S)-1-(((2-Amino-3-bromoquinolin-7-yl)oxy)methyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	N-(7-(((1R,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,3-dihydroxybicyclo[3.1.0]hexan-1-yl)methoxy)-3-bromoquinolin-2-yl)-2,2,2-trifluoroacetamide.	¹H NMR (400 MHz, DMSO-d₆) δ 9.39 (bs, 2H), 8.45 – 8.36 (m, 3H), 7.83 – 7.62 (m, 2H), 7.03-7.14 (m, 3H), 6.99 (d, J = 3.6 Hz, 1H), 5.28 (s, 1H), 5.05 (s, 1H), 4.86 (s, 1H), 4.71 – 4.57 (m, 2H), 3.93 (d, J = 10.4 Hz, 1H), 3.72 (d, J = 6.4 Hz, 1H), 1.62 (d, J = 8.6 Hz, 1H), 1.53 (t, J = 4.4 Hz, 1H), 0.90– 0.82 (m, 1H); LCMS m/z = 499.2 (M+2; 40%).
Compound-81  (1S,2R,3S,4R,5S)-4-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-1-(((2-aminoquinolin-7-yl)thio)methyl)bicyclo[3.1.0]hexane-2,3-diol.	7-(((3aR,3bS,4aS,5R,5aS)-5-(4-Amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2,2-dimethyltetrahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-3b(3aH)-yl)methyl)thio)-N-(4-methoxybenzyl)quinolin-2-amine.	¹H NMR (400 MHz, DMSO-d₆) δ 8.05 (s, 1H), 7.82 (d, J = 8.8 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 7.38(d, J = 1.8 Hz, 1H), 7.16 (d, J = 3.5 Hz, 1H), 7.13 (dd, J = 8.4, 1.9 Hz, 1H), 6.96 (s, 2H), 6.67 (d, J = 8.8Hz, 1H), 6.52 (d, J = 3.5 Hz, 1H), 6.44 (s, 2H), 5.21 – 5.11 (m, 1H), 4.90 (s, 1H), 4.68 (d, J = 7.1 Hz, 1H), 4.57 (s, 1H), 3.71 – 3.60 (m, 2H), 3.38 (d, J = 13.5 Hz, 1H), 1.43 – 1.38 (m, 2H), 0.84 – 0.77 (m, 1H); LCMS m/z = 435.10 (M+2; 90%).

Example-7: (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(4-fluorophenyl)quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (Compound-82)

5





In a sealed tube, the mixture of (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol (100

mg, 0.202 mmol), K₂CO₃ (84mg, 0.606mmol), (4-fluorophenyl)boronic acid (42.4mg, 0.303mmol) in dioxane (10ml) was degassed for 10min with nitrogen at 25°C. PdCl₂(dppf)-CH₂Cl₂ adduct (16.49mg, 0.020mmol) was added and stirred the reaction mixture at 100°C for 16h. The resulting mixture was filtered through celite and filtrate was concentrated *in vacuo* to get 0.15g of crude compound.. This residue was purified by combiflash (R_f200, Teledyne/Isco) instrument onto a redise[®] R_f column with gradient elution (0 to 8%) of methanol in dichloromethane to afford the title compound (0.025g, 93.81%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.07 (s, 1H), 7.78 (s, 1H), 7.68 – 7.50 (m, 3H), 7.43 – 7.24 (m, 3H), 7.13 (dd, *J* = 8.2, 1.7 Hz, 1H), 7.05 (d, *J* = 3.6 Hz, 1H), 6.97 (s, 2H), 6.59 (d, *J* = 3.5 Hz, 1H), 5.96 (s, 2H), 5.11 (d, *J* = 4.5 Hz, 1H), 4.97 – 4.81 (m, 1H), 4.58 – 4.39 (m, 2H), 3.74 (t, *J* = 5.5 Hz, 1H), 3.04 – 2.72 (m, 2H), 2.18 – 2.09 (m, 1H), 1.90 (td, *J* = 12.5, 5.6 Hz, 1H), 1.34 – 1.16 (m, 2H), 0.64 – 0.54 (m, 1H); LCMS *m/z* = 511.09 (M+1; 90%).

Examples in table-17 were synthesized by following an analogous reaction protocol as was used for the preparation of (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-*d*]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol using the appropriate starting materials.

Table-17

Srtucture & IUPAC name	Intermediate used	¹ H NMR & LCMS data
Compound-83  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(pyridin-3-yl)quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-<i>d</i>]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.07 (s, 1H), 7.78 (s, 1H), 7.68 – 7.50 (m, 3H), 7.43 – 7.24 (m, 3H), 7.13 (dd, <i>J</i> = 8.2, 1.7 Hz, 1H), 7.05 (d, <i>J</i> = 3.6 Hz, 1H), 6.97 (s, 2H), 6.59 (d, <i>J</i> = 3.5 Hz, 1H), 5.96 (s, 2H), 5.11 (d, <i>J</i> = 4.5 Hz, 1H), 4.97 – 4.81 (m, 1H), 4.58 – 4.39 (m, 2H), 3.74 (t, <i>J</i> = 5.5 Hz, 1H), 3.04 – 2.72 (m, 2H), 2.18 – 2.09 (m, 1H), 1.90 (td, <i>J</i> =

		12.5, 5.6 Hz, 1H), 1.34 – 1.16 (m, 2H), 0.64 – 0.54 (m, 1H); LCMS m/z= 494.2(M+1;30%)
Compound-84  (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-(3-methyl isoxazol-4-yl)quinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.	(1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.01 (s, 1H), 8.07 (s, 1H), 7.85 (s, 1H), 7.59 (d, J = 8.2 Hz, 1H), 7.37 (d, J = 1.5 Hz, 1H), 7.13 (dd, J = 8.2, 1.6 Hz, 1H), 7.04 (d, J = 3.6 Hz, 1H), 6.97 (s, 2H), 6.58 (d, J = 3.5 Hz, 1H), 6.15 (s, 2H), 5.12 (d, J = 4.1 Hz, 1H), 4.59 – 4.47 (m, 2H), 3.74 (d, J = 5.2 Hz, 1H), 3.00 – 2.77 (m, 3H), 2.22 (s, 3H), 1.89 (ddd, J = 13.9, 11.4, 5.7 Hz, 1H), 1.30 – 1.25 (m, 2H), 0.90 – 0.80 (m, 1H), 0.59 (q, J = 5.8 Hz, 1H); 498.07 (M+1;100%)

Biological Examples

Biochemical assay protocol 1

Inhibitory effect of compounds on PRMT5 was assessed using HTRF detection technology in biochemical assay. Biotinylated H4R3 (residues 1–21) was used as a substrate. Compounds were pre-incubated with 15-25 ng PRMT5:MEP50 per well of a 384-well plate for 30 min at room temperature in the assay buffer containing 20 mM Bicine, pH 7.6, 25 mM NaCl, 2 mM DTT, 0.01% Chicken albumin and 0.01% Tween-20. Reaction was initiated by adding 1 μM of SAM and 50 nM biotinylated H4R3. Total assay volume was 15 μL. Reaction was continued for 120 min at room temperature. Then detection solution containing Streptavidin-Eu cryptate, anti-rabbit IgG-XL-665, Histone H4R3

Dimethyl Symmetric (H4R3me2s) Polyclonal Antibody, all prepared in HTRF detection buffer was added and further incubated for 30 min at room temperature. HTRF signal was recorded in PHERAStar microplate reader. Ratio of signal obtained at 665 nm and 620 nm was used to compute the percent inhibition of compound as follows

5 % Inhibition = $100 - ((\text{Test Ratio} - \text{Negative control Ratio}) / (\text{Positive control Ratio} - \text{Negative control Ratio}) * 100)$ where

Positive control = PRMT5 + SAM + H4R3

Negative control = PRMT5 + H4R3

Biochemical assay protocol 2

10 Inhibitory effect of compounds on PRMT5 was assessed using HTRF detection technology in biochemical assay. Biotinylated H4R3 (residues 1–21) was used as a substrate. Compounds were pre-incubated with 2.5 ng PRMT5:MEP50 per well of a 384-well plate for 30 min at room temperature in the assay buffer containing 20 mM Bicine, pH 7.6, 25 mM NaCl, 2 mM DTT, 0.01% Chicken albumin and 0.01% Tween-20. Reaction

15 was initiated by adding 1 μM of SAM and 50 nM biotinylated H4R3. Total assay volume was 15 μL . Reaction was continued for 4 h at room temperature. Then detection solution containing Streptavidin-Eu cryptate, anti-rabbit IgG-XL-665, Histone H4R3 Dimethyl Symmetric (H4R3me2s) Polyclonal Antibody, all prepared in HTRF detection buffer was added and further incubated for 30 min at room temperature. HTRF signal was recorded in

20 PHERAStar microplate reader. Ratio of signal obtained at 665 nm and 620 nm was used to compute the percent inhibition of compound as follows

% Inhibition = $100 - ((\text{Test Ratio} - \text{Negative control Ratio}) / (\text{Positive control Ratio} - \text{Negative control Ratio}) * 100)$ where

Positive control = PRMT5 + SAM + H4R3

25 Negative control = PRMT5 + H4R3

Activity Range	Compound numbers
IC ₅₀ 300pM to 950pM	43, 26, 35b, 47, 21, 38, 34b, 34a, 7a, 13, 37, 18, 33b, 33a, 76a, 23, 25, 24, 48, 1, 61, 73, 44, 2, 15, 36a, 45a, 45b, 35a

SDMA inhibition assay

Protocol

Z-138 cells (ATCC, CRL-3001TM) were seeded at a density of 1 million cells/well in transparent, flat bottomed tissue culture grade 48-well plates. Cells were treated with various concentration of test compounds for a period of 48 h. Cell lysate was prepared using 1X CST Lysis buffer (Cell Signaling Technology, USA) and 500 ng/well/50 µL of lysate in pH 9.6 carbonate buffer was coated on 96-well Maxisorb plate and incubated overnight at 4°C. The plate was washed twice in 1 x PBS containing 0.05% Tween 20 and blocked in 1% BSA for 1 h at ambient temperature. Further, the plate was incubated first with primary antibody (anti-SDMA antibody; CST#13222s) at ambient temperature for 2 h and then with HRP-conjugated secondary antibody at ambient temperature for 1 h with 2 intermittent washing steps in between.

For luminiscence based detection, HRP substrates (substrate A + substrate B in a 1:1 proportion) were added followed by luminiscence reading after 30 min in SynergyTM 2 reader (Biotek, USA).

For absorbance based detection, TMB substrate was added followed by addition of STOP solution (2N H₂SO₄) post colour development and absorbance (excitation 450 nm and emission 540 nm) was measured in SynergyTM 2 reader (Biotek, USA).

% inhibition of SDMA was calculated relative to the vehicle control samples containing media with 0.1% DMSO alone as per the formula below.

$$\frac{(\text{Avg. of Untreated Control} - \text{Avg. of Test})}{\text{Avg. of Untreated control}} \times 100$$

Avg. of Untreated control

The IC₅₀ values of individual compounds were calculated with Non Linear Regression Analysis using Graph Pad Prism (Graph Pad software, Inc, USA).

Activity Range	Compound numbers
IC ₅₀ 1pM to 1nM	7b, 13, 37, 33b, 46, 76b, 23, 24, 2, 48, 1, 61, 73, 47, 38, 34b, 30, 41, 31, 72, 29, 34a, 21, 35a, 26, 43.
IC ₅₀ 1.1nM to 50nM	32, 64, 33a, 76a, 25, 62, 35b.

Anticancer activity assay

Z-138 cells were seeded at a density of 2000-3000 cells per well in culture media (IMDM + 10% FBS). PANC-1 (ATCC, CRL-1469TM) and MIA PaCa-2 (ATCC, CRL-1420TM) cells were seeded at a density of 200-300 cells per well in culture media (DMEM + 10% FBS). Cells were seeded in opaque, flat bottomed tissue culture grade 96-well plates and Z-138 cells (suspension) were seeded and treated on the same day with various concentrations of test compounds. PANC-1 and MIA PaCa-2 cells, being adherent, were kept for overnight settlement at standard cell culture conditions (37°C, 5% CO₂). On the following day, cells were treated with various concentrations of test compounds. Cells were treated with test compounds for a period of 96 h, 7 days and 10 days, for Z-138 cells, PANC-1 cells and MIA PaCa-2 cells, respectively. Cell viability was assessed using CellTiterGloTM (Promega, USA) as per manufacturer's instructions. Relative Light Units (RLU) were read in SynergyTM 2 reader (Biotek, USA). The assay measures cellular ATP as an indicator of cell viability. RLU is proportional to the number of viable cells in the respective well.

% inhibition of cell viability was calculated relative to the vehicle control samples containing media with 0.1% DMSO alone as per the formula below.

$$\frac{(\text{Avg. of Untreated Control} - \text{Avg. of Test})}{\text{Avg. of Untreated control}} \times 100$$

Avg. of Untreated control

The IC₅₀ values of individual compounds were calculated with Non Linear Regression Analysis using Graph Pad Prism (Graph Pad software, Inc, USA).

Anti-cancer Assay (Z-138)

Activity Range	Compound numbers
IC ₅₀ 0.1pM to 100pM	43, 35a, 47, 21, 40, 38, 34a, 13, 37, 18, 33b, 46, 76b, 24, 2, 48, 1, 54, 61, 73, 34b, 7a, 25, 33a, 44, 72.

IC ₅₀ 101pM to 1nM	26 , 64, 39, 7b, 76a, 62, 20b
-------------------------------	-------------------------------

Anti-cancer assay (Panc-1)

Activity Range	Compound numbers
IC ₅₀ 300pM to 20nM	41, 47, 64, 40, 21, 38, 34b, 34a, 7b, 13, 37, 33b , 46 , 25 , 24, 48 , 61, 73, 18, 30, 7a, 2, 1, 54, 35a, 43
IC ₅₀ 20nM to 100nM	39, 29, 33a, 76b, 62

5

Anti-cancer assay (MiaPaCa-2)

Activity Range	Compound numbers
IC ₅₀ 1pM to 40nM	21, 38, 34b, 34a, 13, 37, 33b, 25, 24, 2, 48, 1, 61, 73, 18, 30, 40 , 64 , 62, 33a, 35a, 43 , 26

In vivo efficacy experiments

- 10 Tumor xenograft for mantle cell lymphoma was established by injection of cells into the right flank of female NOD.CB17-Prkdc^{scid}/J mice with an age between 7- 11 weeks purchased from The Jackson Laboratory, USA. All animal study proposals were reviewed and approved by the Institutional Animal Ethics Committee (IAEC) prior to initiation of experimentation.

15 Z-138 xenograft

For Z-138 xenograft mouse model, Z-138 cells (ATCC® CRL-3001™) were grown in IMDM medium supplemented with 10% FBS. Cells were incubated under standard conditions at 37°C and 5% CO₂. For generating tumors, Z-138 cells in IMDM medium were mixed with Matrigel (Corning® Matrigel® Basement Membrane Matrix) in a ratio of 1:1.

- 20 10 x 10⁶ cells) in a volume of 200 µL were injected subcutaneously in each mouse to establish tumors. Mice were randomized into treatment groups of 8-10 mice, once tumors reached an average volume between 100 to 120 mm³. Treatment was initiated on day of randomization and continued until end of the study. The Vehicle and test compound treatment groups were administered respective treatments orally, using gavage tubing, at
- 25 an application volume of 10 mL/kg per mouse twice a day.

Mice were housed in individually ventilated cages (IVC) at room temperature of 22±3°C, humidity 50±20% and 12/12 h light/dark cycle. All the experimental activities were carried-out inside the biosafety cabinets to ensure sterility.

Tumor size was measured with Digimatic Vernier caliper (Mitutoyo, Japan) when the tumors became palpable. Tumor volume (T. V.) is calculated by using the formula:

$$\text{Tumor volume (mm}^3\text{)} = (L \times W^2)/2$$

Where, L: Length of tumor, W: Width of tumor in millimeter

Percent tumor growth inhibition (% TGI) is calculated using the formula:

$$\% \text{ TGI} = [1 - (T_f - T_i)/(C_f - C_i)] \times 100$$

Where, T_f and T_i , are the final and initial tumor volumes (test compound), and C_f and C_i are the final and initial mean tumor volumes (vehicle group), respectively.

Percent tumor regression is calculated as:

$$\% \text{ TR} = (T_i - T_f)/(T_i) \times 100$$

Where, T_f and T_i , are the final and initial tumor volumes, respectively.


The compounds 24, 33b, and 13 were tested for tumor growth inhibition in Z-138 xenograft model using assay procedure given above; the % of tumor growth inhibition after 38 days at 1 mg/kg dose was found to be 100% and tumor regression was 67-74%. The compound-48 was tested at 5 mg/kg dose, it showed 100% tumor growth inhibition and 63% tumor regression.

20

25

CLAIMS

1. A compound having the general formula (I), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,




5 wherein,

L_1 is selected from $-CR^aR^b-$, $-NR^a-$, S, and O;

Z is selected from CH and N;

R^a and R^b are independently selected at each occurrence from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

10 ring A is selected from,



R^c and R^d are selected from substituted or unsubstituted alkyl or together with the carbon atoms to which they are attached form a C_3 - C_6 cycloalkyl ring;

15 R is selected from $-NR^4R^5$, hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted heteroaryl and substituted or unsubstituted cycloalkyl;

R^1 and R^2 together with the carbon atoms to which they are attached form a bond in order to form a $-C=C-$; or R^1 and R^2 together with the carbon atoms to which they are attached form a cyclopropane ring;

$R^{2'}$ and R^{2a} which may be same or different and are independently selected from hydrogen and substituted or unsubstituted alkyl;

R^3 is independently selected at each occurrence from halogen, cyano, nitro, substituted or unsubstituted alkyl, $-OR^6$, $-NR^7R^8$, substituted or unsubstituted cycloalkyl, $-C(O)OH$, $-C(O)O$ -alkyl, $-C(O)R^9$, $-C(O)NR^7R^8$, $-NR^7C(O)R^9$, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted heterocyclyl;

R^4 and R^5 are independently selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

R^6 is selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

R^7 and R^8 are independently selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted cycloalkyl;

R^9 is selected from substituted or unsubstituted alkyl and substituted or unsubstituted cycloalkyl;

R^{10} is selected from hydrogen, halogen, and substituted or unsubstituted alkyl;

'n' is an integer ranging from 0 to 4, both inclusive;

when an alkyl group is substituted, it is substituted with 1 to 4 substituents independently selected from oxo ($=O$), halogen, cyano, cycloalkyl, aryl, heteroaryl, heterocyclyl, $-OR^{7a}$, $-C(=O)OH$, $-C(=O)O$ (alkyl), $-NR^{8a}R^{8b}$, $-NR^{8a}C(=O)R^{9a}$, and $-C(=O)NR^{8a}R^{8b}$;

when the heteroaryl group is substituted, it is substituted with 1 to 4 substituents independently selected from halogen, nitro, cyano, alkyl, haloalkyl, perhaloalkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, $-OR^{7a}$, $-NR^{8a}R^{8b}$, $-NR^{7a}C(=O)R^{9a}$, $-C(=O)R^{9a}$, $-C(=O)NR^{8a}R^{8b}$, $-SO_2$ -alkyl, $-C(=O)OH$, and $-C(=O)O$ -alkyl;

when the heterocycle group is substituted, it is substituted either on a ring carbon atom or on a ring hetero atom, and when it is substituted on a ring carbon atom, it


is substituted with 1 to 4 substituents independently selected from oxo (=O), halogen, cyano, alkyl, cycloalkyl, perhaloalkyl, $-OR^{7a}$, $-C(=O)NR^{8a}R^{8b}$, $-C(=O)OH$, $-C(=O)O$ -alkyl, $-N(H)C(=O)(alkyl)$, $-N(H)R^{8a}$, and $-N(alkyl)_2$; and when the heterocycle group is substituted on a ring nitrogen, it is substituted with substituents independently selected from alkyl, cycloalkyl, aryl, heteroaryl, $-SO_2(alkyl)$, $-C(=O)R^{9a}$, and $-C(=O)O(alkyl)$; when the heterocycle group is substituted on a ring sulfur, it is substituted with 1 or 2 oxo (=O) group(s);

R^{7a} is selected from hydrogen, alkyl, perhaloalkyl, and cycloalkyl;

R^{8a} and R^{8b} are each independently selected from hydrogen, alkyl, and cycloalkyl; and

R^{9a} is selected from alkyl and cycloalkyl.

2. The compound of claim 1 having the structure of Formula (II), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,




15 wherein,

Ring A, Z, L_1 , R^a , R^b , $R^{2'}$, R, R^{2a} , R^3 , R^{10} and 'n' are as defined herein above.

3. The compound of claim 1 having the structure of Formula (III), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,


230



wherein,

Ring A, Z, L₁, R^a, R^b, R^{2'}, R, R^{2a}, R³, R¹⁰ and 'n' are as defined herein above.

4. The compound of claim 1 having the structure of Formula (IV), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof,




wherein,

X² is Br or Cl;

L₁, R^a, R^b, R¹, R^{2'}, R², R, R^{2a} and R¹⁰ are as defined herein above.


5. The compound of claim 1 to 3, wherein ring A is selected from-




6. The compound of claim 1 to 4, wherein L₁ is selected from -CH₂-, -CH(CH₃)-, -NH-, -N(CH₃)-, S, and O.

231

7. The compound of claim 1 to 3, wherein R^3 is selected from F, Cl, Br, CN, $-NH_2$, $-NH(CH_3)$, $-NHCH(CH_3)_2$, $-CH_3$, cyclopropyl, $-CH(CH_3)_2$, $-CF_2CH_3$, $-OCH_3$, CF_3 ,



- 5 8. The compound of claim 1 to 4, wherein R is selected from hydrogen, $-NH_2$, Cl, $-CH(CH_3)_2$, methyl, ethyl, cyclopropyl and




9. The compound of claim 1 to 4, wherein R^a and R^b are independently selected from hydrogen, methyl, and cyclopropyl.


- 10 10. The compound of claim 1 to 4, wherein $R^{2'}$ and R^{2a} are independently selected from hydrogen and methyl.


11. The compound of claim 1 to 4, wherein R^{10} is selected from hydrogen, $-F$, and methyl.


12. The compound of claim 1, wherein ring A is selected from-



- 15 L1 is selected from $-CH_2-$, $-CH(CH_3)-$, $-NH-$, $-N(CH_3)-$, S, and O; R^3 is selected from F, Cl, Br, CN, $-NH_2$, $-NH(CH_3)$, $-NHCH(CH_3)_2$, $-CH_3$, cyclopropyl, $-CH(CH_3)_2$, $-CF_2CH_3$, $-OCH_3$, CF_3 ,





cyclopropyl and ; R^a and R^b are independently selected from hydrogen, methyl, and cyclopropyl; R^{2'} and R^{2a} are independently selected from hydrogen and methyl; R¹⁰ is selected from hydrogen, -F, and methyl.

13. The compound of formula (I), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof, as claimed in any one of preceding claims, wherein the compound is selected from:
 - 5 (1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (1S,2R,5R)-3-(((2-amino-3-chloroquinolin-7-yl)thio)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - 10 (1S,2R,5R)-3-(1-(2-Amino-3-bromoquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol
(Compound-9);
 - 15 (1S,2R,5R)-3-(1-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - 20 (1S,2R,5R)-3-(1-((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (1S,2R,5R)-3-(2-(2-amino-3-chloroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (2-(2-Amino-3-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - 25 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
 - (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;

- (1S,2R,5R)-3-(2-(2-amino-3-chloro-8-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-amino-3,5-dichloroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 5 (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 10 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 15 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-ethylcyclopent-3-ene-1,2-diol;
- 20 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-ethyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-cyclopropyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 25 (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoro quinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 30

- (1S,2R,5R)-3-(2-(2-Amino-3-chloro-5-fluoro quinolin-7-yl)ethyl)-5-(4-methyl-1H-pyrrolo[3,2-c]pyridin-1-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-Amino-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 5 (1S,2R,5R)-3-(2-(2-amino-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-methylquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol;
- 10 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol;
- 15 (1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol; and
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo [3.1.0]hexane-2,3-diol.
- 20
14. The compound of formula (I), a stereoisomer thereof, or a pharmaceutically acceptable salt thereof, as claimed in any one of preceding claims, wherein the compound is selected from:
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 25 (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;

- (1S,2R,5R)-3-(2-(2-amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol;
- 5 (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl) bicyclo[3.1.0]hexane-2,3-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo [2,3-d]pyrimidin-7-yl)bicyclo [3.1.0]hexane-2,3-diol;
- 10 (1S,2R,5R)-3-(2-(2-Amino-3-bromoquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-amino-3-chloro-6-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(((2-amino-3-chloro-5-fluoroquinolin-7-yl)oxy)methyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 15 (1S,2R,5R)-3-(1-(2-amino-3-chloroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 20 (1S,2R,5R)-3-(1-(2-amino-3-chloro-5-fluoroquinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylcyclopent-3-ene-1,2-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-chloroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol;
- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- 25 (1S,2R,5R)-3-(1-(2-Amino-3-bromo-5-fluoro quinolin-7-yl)propan-2-yl)-5-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1S,2R,5R)-3-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-5-(4-methyl-7H-pyrrolo[2,3-d]pyrimidin-7-yl)cyclopent-3-ene-1,2-diol;
- (1R,2R,3S,4R,5S)-1-(2-(2-Amino-3-bromo-5-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol; and
- 30

(1R,2R,3S,4R,5S)-1-(2-(2-amino-3-bromo-6-fluoroquinolin-7-yl)ethyl)-4-(4-amino-7H-pyrrolo[2,3-d]pyrimidin-7-yl)bicyclo[3.1.0]hexane-2,3-diol.

15. A pharmaceutical composition comprising at least one compound of any one of the claims 1 to 14, a stereoisomer thereof, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
16. A method for treating the diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 enzyme to a subject in need thereof, comprising administering to the subject, an effective amount of compound as claimed in claim 1 to 14, or their pharmaceutically acceptable salt thereof.
17. A method as claimed in claim 16, wherein the said diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 enzyme is glioblastoma multiforme, prostate cancer, pancreatic cancer, mantle cell lymphoma, non-Hodgkin's lymphomas and diffuse large B-cell lymphoma, acute myeloid leukemia, acute lymphoblastic leukemia, multiple myeloma, non-small cell lung cancer, small cell lung cancer, breast cancer, triple negative breast cancer, gastric cancer, colorectal cancer, ovarian cancer, bladder cancer, hepatocellular cancer, melanoma, sarcoma, oropharyngeal squamous cell carcinoma, chronic myelogenous leukemia, epidermal squamous cell carcinoma, nasopharyngeal carcinoma, neuroblastoma, endometrial carcinoma, and cervical cancer.
18. A method as claimed in claim 16, wherein the said diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 enzyme is cancer
19. Use of a compound, of any one of claim 1 to 14, in the preparation of medicament for treating the diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 in a subject in need thereof.
20. The use as claimed in claim 19, wherein the diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 are selected from the group consisting of glioblastoma multiforme, prostate cancer, pancreatic cancer, mantle cell lymphoma, non-Hodgkin's lymphomas and diffuse large B-cell lymphoma, acute myeloid leukemia, acute lymphoblastic leukemia, multiple myeloma, non-small cell lung cancer, small cell lung cancer, breast cancer, triple negative breast cancer, gastric cancer, colorectal cancer, ovarian cancer, bladder cancer,

hepatocellular cancer, melanoma, sarcoma, oropharyngeal squamous cell carcinoma, chronic myelogenous leukemia, epidermal squamous cell carcinoma, nasopharyngeal carcinoma, neuroblastoma, endometrial carcinoma, and cervical cancer.

- 5 21. The use as claimed in claim 19, wherein the diseases, disorders, syndromes or conditions associated by inhibition of PRMT5 is cancer.

10

15

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2018/060015

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D471/04 C07D487/04 A61P35/00 A61K31/519 A61K31/53
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2017/032840 A1 (JANSSEN PHARMACEUTICA NV [BE]) 2 March 2017 (2017-03-02) cited in the application examples; claims; p. 1, lines 5-8; -----	1-21
Y	WO 2014/100695 A1 (EPIZYME INC [US]) 26 June 2014 (2014-06-26) cited in the application examples; claims; p. 2. paragraph [0006]; -----	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 February 2019

Date of mailing of the international search report

27/02/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Wolf, Claudia

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2018/060015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2017032840	A1	02-03-2017	AR 105820 A1 15-11-2017
		AU 2016311295 A1 08-02-2018	
		BR 112018003595 A2 25-09-2018	
		CA 2992688 A1 02-03-2017	
		CL 2018000510 A1 06-07-2018	
		CN 107922413 A 17-04-2018	
		CO 2018002063 A2 19-07-2018	
		EA 201890573 A1 31-07-2018	
		EP 3341368 A1 04-07-2018	
		JP 2018528946 A 04-10-2018	
		KR 20180041222 A 23-04-2018	
		PE 9292018 A1 08-06-2018	
		PH 12018500425 A1 10-09-2018	
		TW 201718575 A 01-06-2017	
		US 2018243328 A1 30-08-2018	
		WO 2017032840 A1 02-03-2017	

WO 2014100695	A1	26-06-2014	CA 2894126 A1 26-06-2014
		EP 2935240 A1 28-10-2015	
		JP 2016505000 A 18-02-2016	
		US 2014228343 A1 14-08-2014	
		US 2015133427 A1 14-05-2015	
		US 2016214985 A1 28-07-2016	
		US 2017114061 A1 27-04-2017	
		US 2018186798 A1 05-07-2018	
		US 2018298010 A1 18-10-2018	
		WO 2014100695 A1 26-06-2014	
