



(86) Date de dépôt PCT/PCT Filing Date: 2001/01/12
(87) Date publication PCT/PCT Publication Date: 2001/07/19
(45) Date de délivrance/Issue Date: 2011/03/15
(85) Entrée phase nationale/National Entry: 2002/07/12
(86) N° demande PCT/PCT Application No.: US 2001/000981
(87) N° publication PCT/PCT Publication No.: 2001/051490
(30) Priorité/Priority: 2000/01/14 (US60/176,373)

(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/505 (2006.01), *A61K 31/52* (2006.01),
A61P 29/00 (2006.01), *A61P 35/00* (2006.01),
C07D 235/00 (2006.01), *C07D 239/00* (2006.01),
C07D 239/54 (2006.01), *C07D 239/56* (2006.01),
C07D 239/58 (2006.01), *C07D 473/34* (2006.01),
C07D 487/04 (2006.01)

(72) Inventeurs/Inventors:
JACOBSON, KENNETH A., US;
MARQUEZ, VICTOR E., US

(73) Propriétaire/Owner:

(54) Titre : ANALOGUES NUCLEOSIDIQUES DE METHANOCARBA CYCLOALKYLE
(54) Title: METHANOCARBA CYCLOALKYL NUCLEOSIDE ANALOGUES

(57) **Abrégé/Abstract:**

The present invention provides novel nucleoside and nucleotide derivatives that are useful agonists or antagonists of P1 or P2 receptors. For example, the present invention provides a compound of formula A-M, wherein A is modified adenine or uracil and M is a constrained cycloalkyl group. The adenine or uracil is bonded to the constrained cycloalkyl group. The compounds of the present invention are useful in the treatment or prevention of various diseases including airway diseases (through A_{2B}, A₃, P2Y₂ receptors), cancer (through A₃, P2 receptors), cardiac arrhythmias (through A₁ receptors), cardiac ischemia (through A₁, A₃ receptors), epilepsy (through A₁, P2X receptors), and Huntington's Disease (through A_{2A} receptors).



(73) **Propriétaires(suite)/Owners(continued):**

THE GOVERNMENT OF THE UNITED STATES OF AMERICA, REPRESENTED BY THE SECRETARY, DEPARTMENT
OF HEALTH AND HUMAN SERVICES, US

(74) **Agent:** R. WILLIAM WRAY & ASSOCIATES

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2001 (19.07.2001)

PCT

(10) International Publication Number
WO 01/51490 A1

(51) International Patent Classification⁷: **C07D 471/04**,
487/04, 239/54, 239/56, 239/58, A61P 29/00, 35/00,
A61K 31/505, 31/52 // (C07D 487/04, 239:00, 235:00)

(74) Agent: **PILLAI, Xavier**; Leydig, Voit & Mayer, Ltd., 700
Thirteenth Street, N.W., Suite 300, Washington, DC 20005
(US).

(21) International Application Number: PCT/US01/00981

(22) International Filing Date: 12 January 2001 (12.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/176,373 14 January 2000 (14.01.2000) US

(71) Applicant (for all designated States except US): **THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES** [US/US]; National Institutes of Health, Office of Technology Transfer, 6011 Executive Boulevard, Suite 325, Rockville, MD 20852 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JACOBSON, Kenneth, A.** [US/US]; 1111 University Building W, Apartment 504, Silver Spring, MD 20902 (US). **MARQUEZ, Victor, E.** [US/US]; 220 Doolittle Street, Montgomery Village, MD 20886 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHANOCARBA CYCLOALKYL NUCLEOSIDE ANALOGUES

(57) Abstract: The present invention provides novel nucleoside and nucleotide derivatives that are useful agonists or antagonists of P1 or P2 receptors. For example, the present invention provides a compound of formula A-M, wherein A is modified adenine or uracil and M is a constrained cycloalkyl group. The adenine or uracil is bonded to the constrained cycloalkyl group. The compounds of the present invention are useful in the treatment or prevention of various diseases including airway diseases (through A_{2B}, A₃, P2Y₂ receptors), cancer (through A₃, P2 receptors), cardiac arrhythmias (through A₁ receptors), cardiac ischemia (through A₁, A₃ receptors), epilepsy (through A₁, P2X receptors), and Huntington's Disease (through A_{2A} receptors).



WO 01/51490 A1

instance, the prevention of reperfusion injury after cardiac ischemia or stroke, and treatment of hypertension and epilepsy (Jacobson, et al., J. Med. Chem., 35, 407-422 (1992)).

Adenosine itself has recently been approved for the treatment
5 of paroxysmal supra ventricular tachycardia (Pantely, et al., Circulation, 82, 1854 (1990)). Adenosine receptor agonists also find use as anti-arrhythmics, antinociceptives, anti-lipolytics, cerebroprotectives, and antipsychotics.

P2 receptors, are present in heart, skeletal, various
10 smooth muscles, prostate, ovary, and brain and have been implicated in certain aggregation processes associated with thrombosis and as anti-hypertensive and anti-diabetic agents. Agonists that bind the P2 receptor induce activation of phospholipase C, which leads to the generation of inositol
15 phosphates and diacyl glycerol with a subsequent rise in intracellular calcium concentration and muscle relaxation. P2 receptor antagonists block ADP-promoted aggregation in platelets and thereby exert an anti-thrombotic effect.

All P1 and P2 receptor nucleoside ligands suffer from
20 chemical instability that is caused by the labile glycosidic linkage in the sugar moiety of the nucleoside. However, it has been found that relatively few ribose modifications are tolerated by the presently known agonists and antagonists of P1 and P2 receptors.

25 New compositions are needed that have improved chemical stability and that do not destroy the activity of such compounds.

The invention provides such compositions and methods of using them in the treatment of disease. These and other
30 advantages of the present invention, as well as additional

inventive features, will be apparent from the description of the invention provided herein.

The following documents disclosed certain analogues
5 containing adenine, thymidine, or uracil: Siddiqui et al.,
Nucleosides & Nucleotides, 15, 235-250 (1996); Katagiri et
al., Tetr. Lett., 40, 9069-9072 (1999); Dyatkina et al.,
Bioorg. & Med. Chem., 6, 2639-2642 (1996); WO 95 08541 A
(1995); WO 95 03304 A (1995); EP-A-0 577 558 (1994); US-A-5
10 840 728 (1998); WO 98 05662 A (1998); US-A-5 629 454 (1997);
Laks et al., Tetr. Lett., 37, 2353-2356 (1996); Marquez et
al., J. Med. Chem., 39, 3739-3747 (1996); Marquez et al.,
JACS, 120, 2780-2789 (1998); Ezzitouni et al., JCS, Perkin
Trans., 1, 1073-1078 (1996); Marquez et al., Helv. Chim. Acta,
15 82, 2119-2129 (2000); Shin et al., JOC, 65, 2172-2178 (2000);
H.R. Moon, JOC, 64, 4733-4741 (1999); A. Ezzitouni, JOC, 62,
4870-4873 (1997); Altmann et al., Tetr. Lett., 35, 2331-2334
(1994); Ezzitouni et al., JCS, Chem. Comm., 1345-1346 (1995);
Theil et al., JCS, Perkin Trans., 1, 255-258 (1996); Rodriguez
20 et al., Tetr. Lett., 34, 6233-6236 (1993); Marquez et al.,
Nucleosides & Nucleotides, 16, 1431-1434 (1997); V. E.
Marquez, Nucleosides & Nucleotides, 18, 521-530 (1999); Jeong
et al., Nucleosides & Nucleotides, 16, 1059-1062 (1997);
Altmann et al., Tetr. Lett., 35, 7625-7628 (1994); Rodriguez
25 et al., J. Med. Chem., 37, 3389-3399 (1994); US-A-4 954 504
(1990); US-A-5 063 233 (1991); Jacobson et al., J. Med. Chem.,
35, 407-422 (1992).

BRIEF SUMMARY OF THE INVENTION

30 The present invention provides novel nucleoside and
nucleotide derivatives that are useful agonists or antagonists
of P1 or P2 receptors. The invention is premised upon the
novel combination of adenine and uracil and their derivatives
with a constrained cycloalkyl group, typically a cyclopentyl
35 group. The constraint on the cycloalkyl group is introduced

3a

by fusion to a second cycloalkyl group. In the case of cyclopentane, the fusion is typically with cyclopropane. The present compounds retain a surprising binding affinity despite the substitution for the ribose group. Moreover, the absence of the glycosidic bond in the compounds assists in improving the chemical stability of the compounds and aids in overcoming the stability problem associated with the glycosidic bond in previously known P1 and P2 receptor ligands.

The compounds of the present invention are useful in the treatment or prevention of various airway diseases (through A_{2B}, A₃, P2Y₂ receptors), cancer (through A₃, P2 receptors), cardiac arrhythmias (through A₁ receptors), cardiac ischemia (through A₁, A₃ receptors), epilepsy (through A₁, P2X receptors), Huntington's Disease (through A_{2A} receptors), immunodeficient disorders (through A₂, A₃ receptors), inflammatory disorders (through A₃, P₂ receptors), neonatal hypoxia (through A₁ receptors), neurodegenerative diseases (through A₁, A₃, P2 receptors), pain (through A₁, A₃, P2X₃ receptors), Parkinson's Disease (through A_{2A} receptors), renal failure (through A₁ receptors), schizophrenia (through A_{2A} receptors), sleep disorders (through A₁ receptors), stroke (through A₁, A₃,

P2 receptors), thrombosis (through P2Y₁, P2Y_{AC} receptors), urinary incontinence (through P2X₁ receptors), diabetes (through A₁ receptors), psoriasis (through P2X receptors), septic shock (through P2 receptors), brain trauma (through A₁ receptors), glaucoma (through A₃ receptors) and congestive heart failure (through P2 receptors).

The invention may best be understood with reference to the accompanying drawings and in the following detailed description of the preferred embodiments.

10

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a new class of nucleoside and nucleotide analogs that serve as selective agonists or antagonists for P1 and P2 receptors.

15

Generally, the compounds of the present invention comprise two basic chemical components designated "A" and "M" which are covalently bonded to one another. Component A comprises adenine or uracil, and component M includes a constrained cycloalkyl group. Preferably the adenine and uracil are chemically modified or substituted with moieties that allow the compound to bind to a P1 or P2 receptor. To that end any of a wide variety of chemical groups can be used to modify adenine and uracil. Those groups are well known to those of skill in the receptor art. Preferably, when A is purine or a purine derivative, the linkage between A and M is a chemical bond between the N9 purine nitrogen and the C1 carbon of the cycloalkyl group. Where A is pyrimidine or a pyrimidine derivative, the bond is between N1 pyrimidine nitrogen and the C1 carbon of the cycloalkyl group. The


20

25


compounds of the present invention have improved stability and surprising receptor binding affinity.

While not wishing to be bound to any particular theory, it is believed that the constrained cycloalkyl group assists
5 in improving chemical stability and receptor affinity. Preferably the cycloalkyl groups are capable of adopting a conformation such that the compound can bind to P1 or P2 receptors. As a result, preferred cycloalkyl groups are those that tend to form energetically favorable interactions with P1
10 and P2 receptors and avoid energetically unfavorable ones, such as unfavorable ionic and/or steric interactions. Further, the cycloalkyl group is derivatized with a bridging group. The constraint restricts the cycloalkyl group to certain conformations that are believed to be beneficial to
15 binding affinity. The preferred cycloalkyl group is a cyclopentyl group. With cyclopentyl groups the preferred method for introducing a conformational constraint is by derivatizing with a fused cyclopropane bridge. With this modification the cyclopentane ring is believed to be
20 constrained to mimic the conformation of a rigid furanose ring.

Compounds of the present invention include the compounds shown below in Formulae I and II.



Formula 1



Formula II

Formulae I and II show compounds in which a derivatized or underivatized adenine base is joined to a constrained cyclopentyl group. For purposes of reference, the carbon atom of the cyclopentyl group, M, that is joined to adenine, A, is the C1 carbon and the adenine is joined to M through its N9 nitrogen. In the compounds of Formulae I and II the constrained cyclopentyl group is derivatized with a fused cyclopropane bridge. In Formula I the cyclopropyl group bridges carbon atoms C4 and C6. In Formula II the cyclopropyl group bridges carbon atoms C6 and C1. These distinct bridging patterns constrain the cyclopentyl group into distinct conformations, specifically the N- (northern) conformation as in Formula I and the S- (southern) conformation as in Formula II. These two conformations are thought to mimic the two

biologically active conformations of furanose groups for P1 and P2 receptor binding pockets.

The compounds described by Formulae I and II can be further defined by a
5 variety of suitable modifications to the adenine group. As discussed above, any of a wide variety of chemical groups can be used to form suitable adenine derivatives that comprise the novel compounds of the present invention, provided that the resulting compound is capable of binding to a P1 or P2 receptor. These chemical groups are well known in the art and have been described, for example in U.S. Patent Nos.
10 5,284,834; 5,498,605; 5,620,676; 5,688,774; and Jacobson and Van Rhee, PURINERGIC APPROACHES IN EXPERIMENTAL THERAPEUTICS, Chapter 6, p. 101 (Jacobson and Jarvis eds., 1997); and Jacobson et al., THE P2 NUCLEOTIDE RECEPTORS, p. 81-107, in THE RECEPTORS (Turner et al., eds. 1998). The combination of the chemically modified adenine and the constrained cycloalkyl group
15 provides a surprising improvement in both chemical stability and binding affinity.

By way of example and not in limitation of the present invention in the compounds of Formulae I and II, R₁ is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, sulfonyl, arylsulfonyl, thiazolyl or bicyclic alkyl; R₂ is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, thioether, cyano,
20 alkylthio or arylalkylthio; R₃, R₄, and R₅, are each hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and they can be the same or different; R₆ is hydrogen, alkyl, alkenyl, alkynyl, or aminoalkyl. R₇ is a methylene,

dihalomethyl, carbonyl, or sulfoxide group. R_8 is carbon or nitrogen. At least one of R_1 , R_2 , and R_6 is not hydrogen. It can be appreciated that various combinations of the above groups are also within the invention provided that they retain
5 agonist or antagonist activity with a P1 or P2 type receptor.

Where an alkyl, alkenyl, alkynyl group is referenced by itself or as part of another group, the reference is to an uninterrupted carbon chain consisting of no more than 20 carbon atoms. Aryl and cycloalkyl groups contain no more than
10 8 carbons in the ring.

Reference to alkyl groups is further meant to include straight or branched chain alkyls, arylalkyl, aminoalkyl, haloalkyl, alkylthio or arylalkylthio groups. Alkyls specifically include methyl through dodecyl. Where alkyl
15 groups are present at position R_6 in adenine, it is preferred that the chain length be no longer than 6 carbons. Arylalkyl groups include, phenylisopropyl, phenylethyl. Aminoalkyl groups can be any suitable alkyl group also containing an amine. Similarly, haloalkyl groups can be any suitable alkyl
20 group that contains a halo substituent, such as bromo, chloro, fluoro, iodo. Alkylthio includes such moieties as thiomethyl, thiopentyl, thiohexyl, thioheptyl, thiooctyl, thiodecyl, thioundecyl, ethylthioethyl, or 6-cyanoethylthio groups. Alkylthio also is meant to include arylalkylthio such as 2-(p-
25 nitrophenyl)ethylthio, 2-aminophenylethylthio, 2-(p-nitrophenyl)ethylthio, or 2-aminophenylethylthio.

Cycloalkyls for example cyclopentyl, cyclohexyl, hydroxycyclopentyl.

Alkoxys include for example methoxy groups.

Cycloalkoxys can include cyclopentoxy.

Aryl moieties can be arylalkyl, arylalkylthio, arylsulfonyl, arylamino, aryloxy, heteroaryl, haloaryl, arylurea, arylcarboxamido, heteroarylamino or sulfoaryl.

5 Benzyl groups are one species of aryl group. In addition, the arylalkyls include R-phenylisopropyl or phenylethyl.

Aryloxides can be phenyl, R-phenylisopropyl, phenylethyl, 3,5-dimethoxyphenyl-2-(2-methylphenyl)ethyl and sulfophenyl.

10 Haloaryl can be iodobenzyl among other halogenated aryl groups. Additionally, the heteroaryls include, for example, furans such as tetrahydrofuran.

Acyl groups include carbonyls.

Alkenyl groups are analogous to alkyl groups but include at least one carbon-carbon double bond. When present at the
15 R₆ group of adenine it is preferred that the carbon chain length be from 2 to 6 carbons.

Similarly, alkynyls are analogous to alkenyl groups but contain at least one triple carbon-carbon bond. As with other groups, when present at the R₆ position of adenine it is
20 preferred that they are not longer than 6 carbons.

Phosphoryl groups include diphosphoryl, triphosphoryl, thiophosphoryl, thiodiphosphoryl, thiotriphosphoryl, imidodiphosphate, imidotriphosphate, methylene diphosphate, methylenetriphosphate, halomethylene diphosphate,
25 halomethylene triphosphate, boranophosphate, boranodiphosphate, boranotriphosphate, or phosphorothioate-2-thioether for example.

Thio groups include alkylthio, arylalkylthio, alkenylthio, or arylthios. Alkylthio includes such groups as

thiomethyl, thiopentyl, thiohexyl, thioheptyl, thiooctyl, thiodecyl, thioundecyl, ethylthioethyl, or 6-cyanoethylthio. Alkenylthio includes 5-hexenylthio. Arylthios include 2-(p-nitrophenyl)ethylthio, 2-aminophenylethylthio, 2-(p-nitrophenyl)ethylthio, or 2-aminophenylethylthio.

One example of a suitable thiazolyl is (benzothiazolyl)thio-2-propyl.

Examples of bicycloalkyls include s-endonorbornyl, or carbamethylcyclopentane.


Halo groups include such elements as fluoro, bromo, chloro, or iodo.

It will also be appreciated that any group that may be further substituted can be, and still be within the scope of the invention. For example, all of the R_1 groups except hydrogen can be further substituted. By way of illustration, when R_1 is not hydrogen, it can be further modified by substitutions with any of the following chemical substituents including amino, cyano, alkoxyl, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxy, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, sulfonamido, carboxyl, and carboxamido groups. Similarly, for $R_2 - R_{10}$ all of the groups other than hydrogen can be substituted further. Multiple substitutions are also contemplated.


In a preferred embodiment R_1 can be either methyl, cyclopentyl, cyclohexyl, phenyl, R-phenylisopropyl, benzyl, or phenylethyl; R_2 is chloride; and R_6 can be a C_1-C_6 alkylamino, C_1-C_6 alkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl group.

Other compounds of the present invention include the compounds shown below in Formulae III and IV. The Formulae

show compounds in which a derivatized or underivatized uracil base is joined to a constrained cyclopentyl group.



Formula III



Formula IV

- 5 The compounds defined by formulae III and IV can be further defined by a variety of suitable modifications. For example R₁ can be hydrogen, or an alkyl group; R₂ can be hydrogen, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, or a C₁-C₆ aminoalkyl group; R₃, R₄, R₅, can each independently be the same as discussed
- 10 previously with respect to Formulae I and Formulae II. R₆ and R₇ are each independently either sulfur or oxygen.

Certain compounds of the present invention are ligands of P2 receptors. A variety of P2 receptors are known in the art and the present compounds act at one or more of these, which

15 include for example, P2X and P2Y receptors. These Receptor ligands are compounds that bind receptors, preferably in the binding pocket. In certain embodiments the compound can be a

P2 receptor agonist. In other embodiments the compound can be a P2 receptor antagonist.

Certain compounds of the present invention are ligands for the P1 receptor. A variety of subclasses of P1 receptors are known and various of present compounds act at one or more
5 these species, which include for example A₁, A₂, and A₃ receptors. Certain compounds act as P1 receptor agonists while others appear to act as antagonists.

The compounds of the present invention are useful in the
10 treatment or prevention of various airway diseases (through A_{2B}, A₃, P2Y₂ receptors), cancer (through A₃, P2 receptors), cardiac arrhythmias (through A₁ receptors), cardiac ischemia (through A₁, A₃ receptors), epilepsy (through A₁, P2X receptors), Huntington's Disease (through A_{2A} receptors),
15 Immunodeficient disorders (through A₂, A₃ receptors), inflammatory disorders (through A₃, P₂ receptors), neonatal hypoxia (through A₁ receptors), neurodegenerative (through A₁, A₃, P2 receptors), pain (through A₁, A₃, P2X₃ receptors), Parkinson's Disease (through A_{2A} receptors), renal failure
20 (through A₁ receptors), schizophrenia (through A_{2A} receptors), sleep disorders (through A₁ receptors), stroke (through A₁, A₃, P2 receptors), thrombosis (through P2Y₁, P2Y_{Ac} receptors), urinary incontinence (through P2X₁ receptors), diabetes (through A₁ receptors), psoriasis (through P2X receptors),
25 septic shock (through P2 receptors), brain trauma (through A₁ receptors), glaucoma (through A₃ receptors), and congestive heart failure (through P2 receptors).

The present invention is further directed to a pharmaceutical composition comprising a pharmaceutically

acceptable carrier and at least one compound selected from the group consisting of the presently described compounds.

The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or
5 diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one that is chemically inert to the active compounds and one that has no detrimental side effects or toxicity under the conditions of
10 use.

The choice of excipient will be determined in part by the particular compound of the present invention chosen, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable
15 formulations of the pharmaceutical composition of the present invention. The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

20 One skilled in the art will appreciate that suitable methods of utilizing a compound and administering it to a mammal for the treatment of disease states, which would be useful in the method of the present invention, are available. Although more than one route can be used to administer a
25 particular compound, a particular route can provide a more immediate and more effective reaction than another route. Accordingly, the described methods are merely exemplary and are in no way limiting.

The dose administered to an animal, particularly human
30 and other mammals, in accordance with the present invention

should be sufficient to effect the desired response. Such responses include reversal or prevention of the bad effects of the disease for which treatment is desired or to elicit the desired benefit. One skilled in the art will recognize that dosage will depend upon a variety of factors, including the age, species, condition or disease state, and body weight of the animal, as well as the source and extent of the disease condition in the animal. The size of the dose will also be determined by the route, timing and frequency of administration as well as the existence, nature, and extent of any adverse side-effects that might accompany the administration of a particular compound and the desired physiological effect. It will be appreciated by one of skill in the art that various conditions or disease states may require prolonged treatment involving multiple administrations.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages that are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. The present inventive method typically will involve the administration of about 0.1 to about 300 mg of one or more of the compounds described above per kg body weight of the individual.

The following examples further illustrate the present invention but, of course, should not be construed as in any way limiting its scope. In the examples, unless otherwise noted, compounds were characterized and resonances assigned by 300

MHz proton nuclear magnetic resonance mass spectroscopy using a Varian GEMINI-300 FT-NMR spectrometer. Also, unless noted otherwise, chemical shifts are expressed as ppm downfield from tetramethylsilane. Synthetic intermediates were characterized
5 by chemical ionization mass spectrometry (NH_3) and adenosine derivatives by fast atom bombardment mass spectrometry (positive ions in a noba or m-bullet matrix) on a JEOL SX102 mass spectrometer. Low resolution CI- NH_3 (chemical
10 ionization) mass spectra were carried out with Finnigan 4600 mass spectrometer and high-resolution EI (electron impact) mass spectrometry with a VG7070F mass spectrometry at 6kV. Elemental analysis was performed by Atlantic Microlab Inc. (Norcross, GA). NMR and mass spectra were consistent with the assigned structure.

15


EXAMPLE 1

In all of the potent adenosine agonists previously developed, the ribose moiety is present, and consequently, these agonists are subject to deglycosylation and other pathways of metabolic degradation *in vivo*. In order to design
20 non-glycosyl adenosine agonists and thereby increase biological stability and potential receptor selectivity, carbocyclic modifications of the ribose moiety have been introduced. In previous studies of adenosine analogues it was found that if adenosine derivatives having carbocyclic
25 modifications of the ribose ring (compounds 1-4, below)

bind to adenosine receptors it is only with greatly reduced affinity.

In the present study we have incorporated a complex
30 carbocyclic modification of ribose for use with adenosine

agonists. This modification, wherein only one isomeric form retains high affinity and receptor selectivity, is the "methanocarba" ring. In this modification a fused cyclopropane ring constrains the accompanying cyclopentane moiety to mimic the conformation of a rigid furanose ring. The furanose ring of nucleosides and nucleotides in solution is known to exist in a rapid, dynamic equilibrium between a range of Northern and opposing Southern conformations as defined in the pseudorotational cycle. For methanocarba analogues, the bicyclo[3.1.0]hexane ring can constrain the cyclopentane ring into a N-, 2'-exo envelope pucker, and a S-, 3' exo form.



These two extreme forms of ring pucker usually define biologically active conformations. This example shows that nucleoside binding to P1- (adenosine) receptors, is favored when the fixed ring-twist conformation is in the N-conformation.

Nucleosides and synthetic reagents were purchased from Sigma Chemical Co. (St. Louis, MO) and Aldrich (St. Louis, MO).

2,6-Dichloropurine was obtained from Sigma. *m*-iodobenzyl

bromide was purchased from Aldrich (St. Louis, MO). 4-(6-


5 Aminopurin-9-yl)-1-hydroxymethyl-bicyclo[3.1.0]hexane-2,3-diol

(1) and compounds 5c and 5d were obtained from Dr. Victor

Marquez. Compounds 7a and 9a were synthesized in our

laboratory.

The synthetic strategy used in this example is shown below. The synthesis of N6-substituted N-methanocarba adenosine derivatives optimized for interaction with A1 (CP = cyclopentyl) or A3 (IB = 3-iodobenzyl) receptors. Reagents: a) DEAD, Ph_3P ; b) MeOH , rt; c) BCl_3 ; d) H_2/Pd ; e) 3-iodobenzyl bromide, 50°C , DMF , 2 days; f) NH_4OH , MeOH , 80°C , 3 days.



(1'R, 2R, 3'R, 4'R, 1'aR)-2,3-(dihydroxy)-4-(hydroxymethyl)-1-(6-cyclopentylaminopurine-9-yl)bicyclo(3.1.0)hexane) (6c):

A solution of 8c (4 mg, 0.01 mmol) in methanol (0.5 ml) was hydrogenated at atmospheric pressure over 10% Pd/C (1 mg) to furnish the product 6c (83% yield). ¹H NMR (CD₃OD): δ 0.7-0.8 (m, 1H, 6'-CHH), 1.46-1.88 (m, 1OH, 6'CHH, 1'aH, 4CH₂), 2.01-2.20 (m, 1H, NCH), 3.34 (d, 1H, J = 9.77 Hz, 5'CHH), 3.88 (d, 1H, J = 6.84 Hz, 3'CH), 4.26 (d, 1H, J = 9.77 Hz, 5'CHH), 4.66-4.98 (m, 2H, 2'CH, 1'CH), 8.28 (s, 1H, 2CH), 8.5 (s, 1H, 8CH). HRMS (FAB): Cal: 346.1879 Found: 346.1879

(1'R, 2'R, 3'R, 4'R)-2,3-(dihydroxy)-4-(hydroxymethyl)-1-(6-(3-iodobenzylamino)purine-9-yl)cyclopentane (7b):

A mixture of aristeromycin (3.5 mg, 0.013 mmol) and 3-iodobenzylbromide (12 mg, 0.039 mmol) in anhydrous DMF was heated for 3 days, and solvent was removed under vacuum. The excess 3-iodobenzylamine was removed from the reaction mixture by adding ether to the reaction mixture, and stirring was continued for 5 min. followed by decantation of the supernatant ether phase. The residue was dried, suspended in methanol (1 ml) and ammonium hydroxide (0.5 ml), and heated at 80 °C in a closed tube for 1 h. Solvent was removed under vacuum, and the residue obtained was purified by flash column chromatography using 7/3 chloroform/methanol to furnish 3.0 mg (47%) of the product.

¹H NMR (CD₃OD) δ 1.86-1.96 (m, 1H, 1'CHH), 2.14-2.30 (m, 1H, 1'CHH), 2.38-2.48 (m, 1H, 4'CH), 3.3-3.38 (m, 1H, 5'CHH), 3.67 (d, 1H, J = 6.84 Hz, 5'CHH), 3.96-4.06 (m, 1H, 3'CH), 4.43-4.48 (m, 1H, 2'CH), 4.73-4.82 (m, 1H, 1'CH), 5.26 (s, 2H, ArCH₂), 7.12 (t, 1H, J = 7.82 Hz, ArH), 7.32 (d, 1H, J =

7.82Hz, ArH), 7.66 (d, 1H, $J = 7.82$ Hz, ArH), 7.73 (s, 1H, ArH), 8.06 (s, 1H, 2CH). 8.08 (s, 1H, 8CH).

Preparation of 4-[6-(3-iodobenzylamino)-purin-9-yl]-1-hydroxymethyl-bicyclo[3.1.0]hexane-2,3-diol (7c, (N)-

5 Methanocarpa- N^6 -(3-iodobenzyl)adenosine) by Dimroth rearrangement:¹

To a solution of 4-(6-amino-purin-9-yl)-1-hydroxymethyl-bicyclo[3.1.0]hexane-2,3-diol (5c, 20 mg, 0.0721 mmol) in DMF (0.5 mL) was added *m*-iodobenzyl bromide (64 mg, 0.216 mmol),
10 and the mixture was stirred at 50 °C for 2 days. DMF was then removed under a stream of N_2 . To the resulting syrup 0.5 mL of acetone and 1 mL of ether were added and the syrup solidified. The solvents were removed by decantation, and again ether was added and removed. The solid was dried and
15 dissolved in 1 mL MeOH. NH_4OH (1.5 mL) was added and the mixture was stirred at 80 °C for 3 days. After cooling down to room temperature, the solvents were removed under reduced pressure and the residue was purified by preparative TLC (silica 60; 1 000 μm ; Analtech, Newark, DE; ethyl acetate-*i*-
20 $PrOH-H_2O$ (8:2:1)) to give 26 mg of the product (7c), yield: 73%. ^1H NMR ($CDCl_3$): δ 0.82 (t, $J = 6.0$ Hz, 1 H), 1.41 (t, $J = 4.8$ Hz, 1 H), 1.72 (dd, $J = 8.5, 6.0$ Hz, 1H), 3.36 (d, $J = 10.8$ Hz, 1 H), 4.05 (d, $J = 6.9$ Hz, 1 H), 4.33 (m, 1 H), 4.80-4.88 (m, 3 H), 5.21 (d, $J = 6.9$ Hz, 1 H), 6.25 (m, br, 1),
25 7.07 (t, $J = 7.8$ Hz, 1 H), 7.35 (d, $J = 7.8$ Hz, 1 H), 7.61 (d, $J = 7.8$ Hz, 1 H), 7.74 (s, 1), 7.93 (s, 1 H), 8.33 (s, 1 H). MS(FAB): m/z 494 ($M^+ + I$).

(1'R, 2'R, 3'R, 4'R, 1'aR,)-2,3-(dihydroxy)-4-(hydroxymethyl)-
1-(2-chloro-6-cyclopentylaminopurine-9-
yl)bicyclo(3.1.0)hexane) (8c):

To a solution of 15 (36 mg, 0.076 mmol) in anhydrous
5 dichloromethane was added BCl₃ (1M solution in
dichloromethane, 0.23 ml, 0.23 mmol) at 0 °C. The reaction
mixture was warmed to room temperature and stirred for 10 min.
To this mixture was added methanol (1 ml) followed by ammonium
hydroxide (0.5 ml). The mixture was concentrated under
10 vacuum, and the residue obtained was purified by flash column
chromatography using 9/1 chloroform-1/methanol as eluent to
furnish 14 mg of the product 8c (48% yield) as a solid.

¹H NMR(CDCl₃): δ 0.65-0.9 (m, 1H, 6'CHH), 1.1-1.4 (m, 2H, 6'CHH,
1'aH), 1.4-1.9 (m, 8H, 4CH₂), 2.0-2.2 (m, 1H, N⁶CH), 3.34 (d,
15 1H, J = 7.2 Hz, 5'CHH), 3.97 (d, 1H, J = 4.6 Hz, 3'CH), 4.25
(d, 1H, J = 7.2 Hz, 5'CHH), 4.687 (s, 1H, 1'CH), 5.11 (d, 1H,
J 4.6, 2'CH), 7.85 (s, 1H, 8CH). HRMS(FAB): Cal: 380.1489
found: 380.1498

(1'R, 2'R, 3'R, 4'R, 1'aR)-2,3-(dihydroxy)-4-(hydroxymethyl)-
20 1-(2-chloro-6-(3-idobenzylamino)purine-9-
yl)bicyclo(3.1.0)hexane) (9c) was synthesized by the same
method as 8c in 53% yield.

¹H NMR(CD₃OD): δ 0.70-0.78 (m, 1H, 6'CHH), 1.50-1.63 (m, 2H,
6', CHH, 1'aH), 3.33 (d, 1H, J = 11.72 Hz, 5'CHH), 3.88 (d, 1H,
25 J = 6.84 Hz, 3'CH), 4.26 (d, 1H, J = 11.72 Hz, 5'CHH), 4.71-
4.83 (m, 2H, 1'CH, 2'CH), 7.1 (t, 1H, J = 7.82 Hz, ArH), 7.40
(d, 1H, J = 7.82 Hz, ArH), 7.61 (d, 1H, 7.82 Hz, ArH), 7.78
(s, 1H, ArH), 8.54 (s, 1H, 8CH). HRMS(FAB): Cal: 528.0299
Found: 528.0295

(2R, 3R, 4R, 1'aR, 1S)-2,3-(O-isopropylidene)-4-(methylenebenzyloxy)-1-(2,6dichloropurine-9-yl)bicyclo(3.1.0)hexane) (12):

To a solution of triphenyl phosphine (260 mg, 1 mmol) in anhydrous THF (2 ml) was added DEAD (0.16 ml, 1 mmol) dropwise at 0 °C, and stirring was continued for 20 min. To this solution was added a solution of 2,6-dichloropurine in THF (4 ml) followed by the addition of 11 (145 mg, 0.5 mmol) in THF (4 ml). The reaction mixture was warmed to room temperature, and stirring was continued for 6 h. Solvent was evaporated under vacuum, and the residue obtained was purified by flash chromatography using 7/3 petroleum ether/ethylacetate as eluent to furnish 141 mg of the product (12) (70% yield) as a gum.

^1H NMR (CDCl_3): δ 1.0 (m, 1H, 6'CHH), 1.24 (s, 3H, CH_3), 1.27-1.38 (m, 1H, 6'CHH), 1.55 (s, 3H, CH_3), 1.62 (dd, 1H, $J = 4.88$, 9.77 Hz, 1'aH), 3.34 (d, 1H, $J = 9.77$ Hz, 5'CHH), 3.97 (d, 1H, $J = 9.77$ Hz, 5'CHH), 4.50 (d, 1H, $J = 6.84$ Hz, 3'CH), 4.57-4.68 (qAB, 2H, $J = 12.7$ Hz, ArCH_2), 5.17 (s, 1H, 1'CH), 5.32 (d, 1H, $J = 6.84$ Hz, 2'H), 7.27-7.4 (m, 5H, Ar), 8.63 (s, 1H, 8CH).

(2R, 3R, 4R, 1'aR, 1S)-2,3-(O-isopropylidene)-4-(methylenebenzyloxy)-1-(2-chloro-6-cyclopentylaminopurine-9-yl)bicyclo(3.1.0)hexane) (15):

To a solution of 12 (42 mg, 0.105 mmol) in methanol (2 ml) was added cyclopentylamine at room temperature, and stirring was continued for 6 hr for complete reaction. Solvent was removed under vacuum, and the residue obtained was purified by flash column chromatography using 7/3 petroleum ether/ethylacetate as eluent to furnish 45 mg of the product 15 (90% yield) as a gum.

^1H NMR(CDCl_3): δ 0.92-0.96 (m, 1H, 6' $\underline{\text{CHH}}$), 1.14-1.01 (m, 1H, 6' $\underline{\text{CHH}}$), 1.23 (s, 3H, CH_3), 1.42-1.81 (m, 9H, 1'aH, 4 CH_2), 1.54 (s, 3H, CH_3), 2.08-2.21 (m, 1H, N⁶ $\underline{\text{CH}}$), 3.44 (d, 1H, J = 9.76 Hz, 5' $\underline{\text{CHH}}$), 3.90 (d, 1H, J = 9.76 Hz, 5' $\underline{\text{CHH}}$), 4.51 (d, 1H, J = 6.84 Hz, 3' $\underline{\text{CH}}$), 4.57-4.67 (qAB, 2H, J = 12.7 Hz, Ar CH_2), 5.04 (s, 1H, 1' $\underline{\text{CH}}$), 5.32 (d, 1H, J = 6.84 Hz, 2' $\underline{\text{CH}}$), 7.2-7.4 (m, 5H, Ar), 8.18 (s, 1H, 8CH).

(1'R, 2'R, 3'R, 4'R, 1'aR)-2,3-(O-isopropylidene)-4-(methylenebenzyloxy)-1-(2-chloro-6-(3-iodobenzylamino)purine-9-yl)bicyclo(3.1.0)hexane (16) was synthesized in 70% yield by the same method as 15, except using 3-iodobenzylamine hydrochloride and two equivalents of triethylamine.

^1H NMR(CDCl_3): δ 0.87-0.91 (m, 1H, 6' $\underline{\text{CHH}}$), 1.10-1.29 (m, 1H, 6' $\underline{\text{CHH}}$), 1.17 (s, 3H, CH_3), 1.42-1.56 (m, 1H, 1'aH), 1.47 (s, 3H, CH_3), 3.37 (d, 1H, J = 9.77 Hz, 5' $\underline{\text{CHH}}$), 3.84 (d, 1H, J = 9.77 Hz, 5' $\underline{\text{CHH}}$), 4.44 (d, 1H, J = 6.84 Hz, 3' $\underline{\text{CH}}$), 4.50-4.60 (qAB, 2H, J = 11.72 Hz, Ar CH_2), 4.70 (bs, 1H, NH), 4.98 (s, 1H, 1' $\underline{\text{CH}}$), 5.24 (d, 1H, J = 6.84 Hz, 2' $\underline{\text{CH}}$), 7.0 (t, 1H, J = 7.82 Hz, ArH), 7.2-7.34 (m, 6H, ArH), 7.55 (d, 1H, J = 7.82, ArH), 7.65 (s, 1H, ArH), 8.08 (s, 1H, 8CH).

Pharmacological Analyses.

Materials

F-12 (Ham's) medium, fetal bovine serum (FBS) and penicillin/streptomycin were from Gibco BRL (Gaithersburg, MD). [^{125}I]AB-MECA (1000 Ci/mmol) and [^{35}S]guanosine 5'-(γ -thio)triphosphate (1000-1500 Ci/mmol) were from DuPont NEN (Boston, MA). Adenosine deaminase (ADA) was from Boehringer Mannheim (Indianapolis, IN). All other materials were from

standard local sources and of the highest grade commercially available.

Cell culture and membrane preparation

5 CHO cells stably transfected with either human A₁ or A₃ receptors (gift of Dr. Gary Stiles and Dr. Mark Olah, Duke University Medical Center) were cultured as monolayers in medium supplemented with 10% fetal bovine serum. Cells were washed twice with 10 ml of ice-cold phosphate buffered saline,
10 lysed in lysis buffer (10 mM Tris.HCl buffer, pH 7.4, containing 2 mM MgCl₂ and 0.5 mM EDTA), and homogenized in a Polytron homogenizer in the presence of 0.2 U/ml adenosine deaminase. The crude membranes were prepared by centrifuging the homogenate at 1000 x g for 10 min followed by
15 centrifugation of the supernatant at 40,000 x g for 15 min. The pellet was washed once with the lysis buffer and recentrifuged at 40,000 x g for 15 min. The final pellets were resuspended in 50 mM Tris.HCl buffer, pH 7.4, containing 10 mM MgCl₂ and 0.1 mM EDTA and stored at -70 °C.

20

Radioreceptor Binding

Determination of binding to adenosine A₁, A_{2A} and A_{2B} receptors was carried out as reported. Determination of A₃ adenosine receptor binding was carried out using [¹²⁵I]AB-MECA. Briefly,
25 aliquots of crude transfected CHO cell membranes (approximately 40 µg protein/tube) were incubated with 0.5 nM [¹²⁵I]AB-MECA, 10 mM MgCl₂, 2 units/ml adenosine deaminase, 50 mM Tris.HCl (pH 7.4) at 37 °C for 60 min. The total volume of the reaction mixture was 125 µl. Bound and free ligands were
30 separated by rapid filtration of the reaction mixture through

Whatman GF/B glass filters. The filters were immediately washed with two 5 ml-portions of ice-cold 50 mM Tris.HCl buffer (pH 7.4). The radioactivity bound to the filters was determined in a Beckman gamma counter. Specific binding was defined as the amount of the radioligand bound in the absence of competing ligand minus the amount of that bound in the presence of 100 μ M NECA. K_i -values were calculated using the K_d for [^{125} I]AB-MECA binding of 0.56 nM.

10 Determination of [^{3} S]GTP γ S Binding

[^{35} S]GTP γ S binding was determined by the method of Lorenzen et al. The incubation mixture contained in a total volume of 125 μ l, 50 mM Tris.HCl (pH 7.4), 1 mM EDTA, 10 mM MgCl₂, 10 μ M guanosine 5'-diphosphate, 1 mM dithiothreitol, 100 mM NaCl, 0.2 units/ml adenosine deaminase, 0.16 nM [^{35} S] GTP γ S (about 50,000 cpm) and 0.5% BSA. The CHO cell membranes expressing A₁ or A₃ receptors were preincubated with the above-mentioned assay mixture at 37 °C for 1 h and further incubated for 1 hr after the addition of [^{35} S]GTP γ S. Incubations were terminated by rapid filtration of the samples through glass fiber filters (Whatman GF/B), followed by two 5 ml washes of the same buffer. After transferring the filters into a vial containing 3 ml of scintillation cocktail, the radioactivity was determined in a scintillation counter.

Data analysis. Analyses of saturation binding assays and concentration-response curves were carried out using the GraphPad Prism (GraphPad Software Inc., San Diego, CA). Comparisons between groups were carried out using the unpaired Student's test.

RESULTS

Chemical synthesis

The methanocarbo-cyclic 2'-deoxyadenosine analogues, shown
5 below in Table 1, in which a fused cyclopropane ring
constrains the cyclopentane ring into a rigid envelope
configuration of either a *N*- or *S*- conformation, were
synthesized in a manner similar as shown above. The *N*-
methanocarba analogues of various N^6 -substituted adenosine
10 derivatives, including cyclopentyl and iodobenzyl, in which
the parent compounds are potent and selective agonists at
either A_1 or A_3 receptors, respectively, were prepared. 2,6-
Dichloropurine, 10, was condensed with the cyclopentyl
derivative, 11, using the Mitsunobu reaction, followed by
15 substitution at the 6-position and deprotection to give 8c or
9e. The 2-chloro substitution of compound 8c was removed by
catalytic reduction to give 6c. This allowed the
incorporation in the *N*-configuration series of the 2-chloro
modification of adenine, which was of interest for its effect
20 on adenosine receptor affinity. An N^6 -(3-iodobenzyl) group
could also be introduced in either aristeromycin, 5b, or *N*-
methanocarba-adenosine, 5c, by the Dimroth rearrangement, to
give 7b and 7c.

25 Biological activity

A pair of methanocarba analogues of adenosine, 5c and 5d,
corresponding to *N*- and *S*- conformations of ribose, were
tested in binding assays, the results of which are shown in
Table 1 below, at four subtypes of adenosine receptors. The
30 more synthetically challenging *S*-isomer (5d) was available

only as the racemate and therefore was tested as such. At rat A1, rat A2A, and human A3 subtypes, the N-analogue proved to be of much higher affinity than the S-analogue. At the human A2B receptor, binding was carried out using [3H]ZM 241,385, however the affinity was too weak to establish selectivity for a specific isomer. Affinity of N-methanocarba-adenosine, 5c, vs. adenosine, 5a, was particularly enhanced at the A3 receptor subtype, for which the ratio of affinities of N-to S-analogues was 150-fold. Although a poor substrate for adenosine deaminase (ADA), the binding curve for 5c was shifted in the presence of ADA, therefore the affinity values for 5c and 5d obtained in the absence of ADA are entered in Table 1, below. The South conformer, 5d, is even a worse substrate of ADA (100-fold less) which explains why the curves in the presence and absence of ADA for 5d are virtually the same. Aristeromycin, 5b, bound weakly to adenosine receptors, with slight selectivity for the A_{2A} subtype. Compound 5c was more potent than aristeromycin, 5b, in binding to A1 (4-fold) and A3 (4500-fold) adenosine receptors.

Compounds 6c and 8c are patterned after A1 receptor-selective agonists, while compounds 7c and 9c are patterned after A3 receptor-selective agonists. Compounds 6 and 7 are unsubstituted at the 2-position, while compounds 8 and 9 contain the potency enhancing 2-chloro substituent. The N6-cyclopentyl N-methanocarba derivative, 6c, based on CPA, 6a, maintained high selectivity for A1 receptors, although the affinity of 6c at rat A1 receptors was 3-fold less than for 6a. In one series it was possible to compare ribose, cyclopentyl, and N-methanocarba derivatives having the same N6-substitution. The N6-(3-iodobenzyl) derivative, 7c, based on a 5'-hydroxy analogue, 7a, of IB-MECA, with a K_i value of

4.1 nM was 2.3-fold more potent at A3 receptors than the ribose-containing parent. Thus, the selectivity of 7c for human A3 versus rat A1 receptors was 17-fold. The aristeromycin analogue, 7b, was relatively weak in binding to
5 adenosine receptors.



Among 2-chloro-substituted derivatives, the N-methanocarba analogue, 8c, was less potent at A1 and A2A receptors than its parent 2-chloro-N6-cyclopentyladenosine, 8a, and roughly equipotent at A3 receptors. Thus, 8c was 53-
10 fold selective in binding to rat A1 vs. human A3 receptors. The N-methanocarba analogue, 9c, of 2-chloro-N6-(3iodobenzyl)adenosine, 9a, had Ki values (nM) of 141, 732, and 2.2 at A1, A2A, and A3 receptors, respectively. Thus, the 2-chloro group slightly enhanced affinity at A3 receptors,
15 while reducing affinity at A1 receptors.


The receptor binding affinity upon replacement of ribose with the N-methanocarba moiety was best preserved for the A3 subtype, at which differences were small. At A1 receptors the loss of affinity for structures 6 - 9 was between 3- and 8-
20 fold. At A2A receptors the loss of affinity was between 6- and 34-fold.

The agonist-induced stimulation of binding of guanine nucleotides to activated G-proteins has been used as a functional assay for a variety of receptors, including
25 adenosine receptors. Binding of [³⁵S]GTP-γ-S was studied in membranes prepared from CHO cells stably expressing human A1 or A3 receptors (Table 2). The non-selective adenosine agonist NECA (5'-N-ethyluronamidoadenosine) caused a concentration-dependent increase in the level of the guanine
30 nucleotide bound. Compound 6c was highly selective and a full agonist at human A1 but not rat A1 receptors. Both 7c and 9c

stimulated the binding of [³⁵S]GTP-γ-S, however the maximal stimulation was significantly less than that produced by either NECA or N6(3-iodobenzyl)adenosine, 7a, both being full A3 agonists. Compounds 7c and 9c resulted in relative stimulation of [³⁵S]GTP-γ-S binding of only 45% and 22%, respectively, indicating that the efficacy of the N-methanocarba analogue at A3 receptors was further reduced upon 2-chloro modification. The potency of compounds 7c and 9c, indicated by the EC50 values in this functional assay, was greater than the potencies of either NECA or compound 7a (Table 2). Thus, the N-methanocarba N6-(3-iodobenzyl) analogues appear to be highly potent and selective partial agonists at human A3 receptors.

Table I
Affinities of Adenosine Derivatives¹

 1a-4a		 1b-4b R'' = OH 1c-4c R'' = H	<u>K_i (nM) or % displacement</u>			
Compound	R'	R	RA ₁ ^a	RA _{2A} ^b	hA ₃ ^c	
1a	H	cyclopentyl	0.59	462	274±20	
CPA					240 (r)	
1B	H	cyclopentyl	5.06±0.51	6800±1800	170±51	
1781						
1c	H	cyclopentyl	5110±790	15% at 10 μM		
1783						
2a	H	3-iodobenzyl	20.0±8.5	17.5±0.5	9.5±1.4 (r)	
IBO-ADO, 541						
2b	H	3-iodobenzyl	69.2±9.8	601±236	4.13±1.76	



1743									
3a	C1	cyclopentyl	0.6	950	237 (r)				
CCPA									
3b	C1	cyclopentyl	8.76±0.81	3390±520	466±58				
1761									
3c	C1	cyclopentyl	3600±780	45±5% at 100 μM					
1782									
4a	C1	3-iodobenzyl	18.5±4.7	38.5±2.0	1.41±0.17 (r)				
542									
4b	C1	3-iodobenzyl	141±22	732±207	2.24±1.45				
1760									
4c	C1	3-iodobenzyl	8730±370	25,400±3800					
1784									

Compound	R ₂	$\frac{rA_1^a}{rA_2^b}$	$\frac{rA_2^b}{rA_3^b}$	$\frac{hA_2^b}{hA_3^b}$	$\frac{A_1/A_2}{A_1/A_3}$
5a	H	Estd. 10 ^d	estd. 30 ^d	<10% at 100 μM	estd. 1000(r) d,e
5b	H	6260±730	2150±950	47,300±10,600	20,000±7900(r) e
5c	H	1680±80	22,500±100 (h) ^{e,f}	35±2% at 50 μM ^f	404±70 ^f

5d	H	15% at 100 μ M	>100,000 (h) ^{e,f}	20 $\pm$ 4% at 50 μ M ^f	62,500 $\pm$ 2900 ^f	>1
(racemic)						
6a	CP	1.50 $\pm$ 0.51	857 $\pm$ 163	21,200 $\pm$ 4300	274 $\pm$ 20, 240 (r) ^e	0.0055
6c	CP	5.06 $\pm$ 0.51	6800 $\pm$ 1800	139k $\pm$ 19k	170 $\pm$ 51	0.030
7a	IB	20.0 $\pm$ 8.5	17.5 $\pm$ 0.5	3570 $\pm$ 100	9.5 $\pm$ 1.4 (r) ^e	2.1
7b	IB	25,900 $\pm$ 1600	<10% 100 μ M	n.d.	1960 $\pm$ 370	13
7c	IB	69.2 $\pm$ 9.8	601 $\pm$ 236	12,100 $\pm$ 1300	4.13 $\pm$ 1.76	17
8a	CP	1.33 $\pm$ 0.19	605 $\pm$ 154	20,400 $\pm$ 1200	237 (r) ^e	0.0056
8c	CP	8.76 $\pm$ 0.81	3390 $\pm$ 520	27 $\pm$ 7% at 100 μ M	466 $\pm$ 58	0.019
9a	IB	18.5 $\pm$ 4.7	38.5 $\pm$ 2.0	5010 $\pm$ 1400	1.41 $\pm$ 0.17 (r) ^e	13
9c	IB	141 $\pm$ 22	732 $\pm$ 207	41,000 $\pm$ 700	2.24 $\pm$ 1.45	63

31

¹ (a) simple carbocyclic, (b) and methanocarpa-adenosine (N)-conformation, (c) and S-conformation, (d) derivatives in radioligand binding assays at rat A₁,^a rat A_{2A},^b human A_{2B},^b and human A₃ receptors,^c unless noted.^e

TABLE II

Effect of ligands to stimulate [^{35} S]GTP γ S binding to membranes of cells expressing the cloned hA₁AR or hA₃AR or in rat cerebral cortical membranes containing the A₁AR

Ligand	cloned hA ₁ AR EC ₅₀ (nM) ^a	% Maximal Stimulation ^c	hA ₁ AR EC ₅₀ (nM) ^a	% Maximal Stimulation ^c	cloned hA ₃ AR EC ₅₀ (nM) ^a	% Maximal Stimulation ^c
NECA	n.d.		n.d.		155±15	100
6a	4.15 ± 0.90	100	20.3 ± 13.1	100	7980 ± 60	100
6c	21.5 ± 2.3	102 ± 1	100 ± 17	75 ± 6	>10,000	14±2% at 10 μ M
7a	43.1 ± 10.4	91 ± 1	340 ± 98	95 ± 4	5.16 ± 0.71	100
7b	>10,000	5±2% at 10 μ M	n.d.		>10,000	15±5% at 10 μ M
7c	218 ± 18	86 ± 2	940 ± 114	55 ± 5	0.70 ± 0.16	45.3 ± 6.8
8c	31.2 ± 3.3	97 ± 1	145 ± 35	96 ± 2	n.d.	
9c	142 ± 24	91 ± 1	684 ± 75	48 ± 3	0.67 ± 0.19	22.0 ± 2.8

^a EC₅₀ for stimulation of basal [^{35} S]GTP- γ S binding by agonists in membranes from transfected CHO cells ($\pm$ S.E.M.), n = 3.

n.d. not determined.

5

Discussion

Nearly all of the thousands of known adenosine agonists are derivatives of adenosine. Although molecular modeling of adenosine agonists has been carried out, there has been no direct evidence from this for a conformational preference of the ribose ring in the receptor binding site. In the present study, methanocarba-adenosine analogues have defined the role of sugar puckering in stabilizing the active receptor-bound conformation. The S-methanocarba analogue of adenosine, 5d, was only weakly active, presumably because of a disfavored conformation that decreases receptor binding. In contrast, the methanocarba analogues constrained in the N-conformation, e.g. 5c - 9c, displayed high receptor affinity, particularly at the A₃ receptor. In binding assays at A₁, A_{2A}, and A₃ receptors, N-methanocarba-adenosine proved to be of higher

20

affinity than the S-analogue, with an N:S-affinity ratio of 150 at the human A3 receptor. Thus, the biological potency and efficacy of this series of nucleosides appears to be highly dependent on ring puckering, which in turn would influence the orientation of the hydroxyl groups within the receptor binding site.

The structure activity relationship (SAR) of adenosine agonists indicates that the ribose ring oxygen may be substituted with carbon, as in 5b and 7b, however much affinity is lost. As demonstrated with the aristeromycin derivative, 7b, simple carbocyclic substitution of the ribose moiety of otherwise potent, N6-substituted adenosine agonists greatly diminishes affinity, even in comparison to aristeromycin, 5b.

In comparison to the ribose analogues, the N-methanocarba N6-substituted adenosine agonists were of comparable affinity at A3 receptors, but less potent at A1, A2A, and A2B receptors. The N-methanocarba N6-cyclopentyl derivatives were A1 receptor-selective and maintained high efficacy at human recombinant but not rat brain A1 receptors, as indicated by stimulation of binding of [³⁵S]GTPγS. This may be related to either species differences or heterogeneity of G proteins, since the degree of agonist efficacy of a given compound may be highly dependent on the receptor-associated G protein. N-Methanocarba N6-(3-iodobenzyl)adenosine and the 2-chloro derivative had K_i values of 4.1 and 2.2 nM at A3 receptors, respectively, and were selective partial agonists. As for the ribose parents, additional 2-chloro substitution was favorable for receptor selectivity. However, unlike the ribose forms, efficacy was reduced in N6-(3-iodobenzyl) analogues, such that partial A3 receptor agonists 7c and 9c were produced.

Partial agonists are possibly more desirable than full agonists as therapeutic agents due to potentially reduced side effects in the former. Partial agonists may display *in vivo* specificity for sites at which spare receptors are present, and the drug would therefore behave with apparent "full" efficacy. Thus, for compounds 7c and 9c, partial agonism combined with unprecedented functional potency at A3 receptors (<1 nM) may give rise to tissue selectivity.

Thus, at least three of the four adenosine receptors favor the N-conformation. For another member of the GPCR superfamily, the P2Y1 receptor, we recently reported that the ribose N-conformation of adenine nucleotides also appears to be preferred at the receptor binding site. Thus, the P1 and at least one of the P2 purinoceptors share the preference for the N-conformation. This may suggest a common motif of binding of nucleoside moieties among these GPCRS. The insights of this conformational preference may be utilized in simulated docking of adenosine agonists in a putative receptor binding site and to design even more potent and selective agents.

At the binding site of ADA, the N-isomer is also preferred, although the carbocyclic adenosine analogues are relatively poor substrates (relative rates of deamination are: 5a, 100; 5b, 0.99; 5c, 0.58; 5d, 0.010. N6-substituted analogues, such as 6c - 9c, would not be expected to be substrates for ADA. Other enzymes, such as HIV reverse transcriptase and Herpes thymidine kinase (HSV-1 TK) are also able to discriminate between the two antipodal conformations of restricted methanocarba thymidine analogues.

In conclusion, we have found that the introduction of a methano-carbocyclic modification of the ribose ring of purine

agonists represents a general approach for the enhancement of pharmacodynamic and because of the absence of the glycosyl bond, potentially of pharmacokinetic properties. This approach could therefore be applied to the development of cardioprotective, cerebroprotective, and anti-inflammatory agents.

EXAMPLE 2

Introduction

P2 receptors, which are activated by purine and/or pyrimidine nucleotides, consist of two families: G protein-coupled receptors termed P2Y, of which 5 mammalian subtypes have been cloned, and ligand-gated cation channels termed P2X, of which 7 mammalian subtypes have been cloned. The P2Y₁ receptor, which is present in the heart, skeletal and various smooth muscles, prostate, ovary, and brain, was the first P2 subtype to be cloned. The nomenclature of P2 receptors and their various ligand specificities is well established.

Nucleotide agonists binding at P2Y₁ receptors induce activation of phospholipase C (PLC), which generates inositol phosphates and diacylglycerol from phosphatidyl inositol-(4,5)-bisphosphate, leading to a rise in intracellular calcium. A P2Y₁ receptor antagonist may have potential as an anti-thrombotic agent, while a selective P2Y₁ receptor agonist may have potential as an anti-hypertensive or anti-diabetic agent.

Recently, progress in the synthesis of selective P2 receptor antagonists has occurred. Adenosine 3',5'- and 2',5'-bisphosphates were recently shown to be selective antagonists or partial agonists at P2Y₁ receptors, and other

classes of P2 antagonists include pyridoxal phosphate derivatives, isoquinolines, large aromatic sulfonates related to the trypanocidal drug suramin and various dyestuffs, and 2',3'-nitrophenylnucleotide derivatives. Synthesis of analogues of adenosine bisphosphates has resulted in N⁶-methyl-2'-deoxyadenosine-3',5'-bisphosphate (1a, MRS 2179), a competitive antagonist at human and turkey P2Y₁ receptors, with a KB value of approximately 100 Nm. The presence of an N⁶-methyl group and the absence of a 2'-hydroxyl group both enhanced affinity and decreased agonist efficacy, thus resulting in a pure antagonist at both turkey and human P2Y₁ receptor. The corresponding 2-Cl analogue (1b, MRS 2216) was slightly more potent than 1a as an antagonist at turkey P2Y₁ receptors, with an IC₅₀ value of 0.22 μM in blocking the effects of 10 nm 2-methylthioadenosine-5'diphosphate (2-MeSADP). MRS2179 (compound 1a) was inactive at P2Y₂, P2Y₄, and P2Y₆ subtypes, at the adenylyl cyclase-linked P2Y receptor in C6 glioma cells and at a novel avian P2Y receptor that inhibits adenylyl cyclase. However, the selectivity of this series of nucleotides for the P2Y₁ receptor is not absolute, since 1a also displayed considerable activity at P2X₁ receptors (EC₅₀ 1.2 μM), but not at P2Y₂₋₄ receptors.


In order to move away from the nucleotide structure of 1a and thereby increase biological stability and selectivity for the receptors in the present study, further structural modifications of the ribose moiety have been carried out. We have explored the SAR of these two series and introduced major modifications of the ribose moiety. These modifications include fixing the ring pucker conformation in the carbocyclic

series using a bridging cyclopropane ring, ring enlargement with introduction of a nitrogen atom, and ring contraction.

Results


Chemical synthesis

5 The methanocarborocyclic 2'-deoxyadenosine analogues in which the fused cyclopropane ring fixes the conformation of the carbocyclic nucleoside into a rigid northern or southern envelope conformation, as defined in the pseudorotational cycle, were synthesized as precursors of nucleotides 4 and 5
10 by the general approach of Marquez and coworkers. Again, the N⁶-methyl group was introduced by the Dimroth rearrangement, as shown below.



2-

Position adenine modifications were further introduced in the
15 N-configuration series as shown below.



Biological activity

Adenine nucleotides markedly stimulate inositol lipid hydrolysis by phospholipase C in turkey erythrocyte membranes, through activation of a P2Y_1 receptor. The agonist used in screening these analogues, 2-MeSADP, has a higher potency than the corresponding triphosphate for stimulation of inositol phosphate accumulation in membranes isolated from [^3H]inositol-labeled turkey erythrocytes.

The deoxyadenosine bisphosphate nucleotide analogues prepared in the present study were tested separately for agonist and antagonist activity in the PLC assay at the P2Y_1 receptor in turkey erythrocyte membranes, and the results are reported in Table 3. Concentration-response curves were determined for each compound alone and in combination with 10nM 2-MeSADP.

Marquez and coworkers have introduced the concept of ring-constrained carbocyclic nucleoside analogues, based on cyclopentane rings constrained in the *N*-(Northern) and *S*-(Southern) conformations by fusion with a cyclopropane

(methanocarba) ring. In the presnet studies the series of ring-constrained N-methanocarba derivatives, the 6-NH₂ analogues, 4a was a pure agonist of EC₅₀ 152 nM and 88-fold more potent than the corresponding *S*-isomer, 5, also an agonist.

5. Thus, the ribose ring *N*-conformation appeared to be favored in recognition at P2Y₁ receptors. The N⁶-methy- and 2-chloro-N⁶-methyl-N-methanocarba analogues, 4b and 4c, were antagonists having IC₅₀ values of 276 and 53 nM, respectively.

10 Molecular modeling.

- To better understand the role of the sugar puckering on the human P2Y₁ agonist and antagonists activities, we carried out a molecular modeling study of this new generation of ribose-modified ligands. Such modifications include
- 15 cyclopentyl rings constrained in the *N*- and *S*- conformations with cyclopropyl (methanocarba) groups, six-membered rings (morpholino and anhydrohexitol analogues), and cyclobutyl nucleotides. We have recently developed a model of the human P2Y₁ receptor, using rhodopsin as a template, by adapting a
- 20 facile method to simulate the reorganization of the native receptor structure induced by the ligand coordination (cross-docking procedure). Details of the model building are given in the Experimental Section. We have also reported the hypothetical molecular basis for recognition by human P2Y₁
- 25 receptors of the natural ligand ATP and the new potent, competitive antagonist 2'-deoxy-N⁶-methyladenosine-3',5'-bisphosphate. Both ATP and 1a are present in the hypothetical binding site with a *N*-sugar ring conformation. In the present work, the sterically constrained *N*- and *S*-methanocarba agonist

analogues, 4a and 5, respectively, were docked into the putative binding site of our previously reported P2Y₁ receptor model. According to their structural similarity, the cross-docking procedure demonstrated that the receptor architecture found for binding the ATP and 1a was energetically appropriate also for the binding of both 4a and 5. However, N-methanocarba/ P2Y₁ complex appeared more stable by approximately 20 kcal/mol than S-methanocarba/P2Y₁ complex.

In the lowest energy docked complex of N-methanocarba agonist in the proposed ligand binding cavity the side chain of Gln307 is within hydrogen bonding distance of the N⁶ atom at 1.8Å, and the side chain of Ser314 is positioned at 2.0Å from the N¹ atom and at 3.4Å from the N⁶ of the purine ring. As already reported, another three amino acids are important for the coordination of the phosphate groups in the antagonist: Arg128, Lys280 and Arg310. Lys280 may interact directly with both 3'-5'-phosphates (1.7Å, O3' and 1.7Å, O5'), whereas Arg128 and Arg310 are within ionic coupling range to both the O2 and O3 atoms of the 5'-phosphate. In molecular modeling studies poor superimposition (rms=1.447) between the N- and S-methanocarba agonist analogues has been found inside the receptor binding domain. In particular, the adenine moiety and 5' phosphate of the S-methanocarba derivative are shifted out position relative to with the N-methanocarba isomer, decreasing the stability of the S-methanocarba/PSY₁ complex. This fact might be correlated with the difference of their biological activity as seen in Table 4 below.

Using the information that a common binding site could be hypothesized among these deoxyadenosine bisphosphate analogues, a superimposition analysis of the energy-minimized

of the more potent antagonists has been performed. In this analysis we have used 1a as a reference compound, and we have defined three matching pairs of atoms, corresponding to N¹ atom of the purine ring and the P atoms of both 3' and 5' phosphate groups, to carry out the superimposition analysis. As reported in Table 4, acceptable RMS values have been obtained for all the antagonists compared with the 1a structure. As shown in Fig 4A, this superimposition study suggested that the two phosphate groups may occupy a common receptor regions, and a general pharmacophore model for bisphosphate antagonists binding to the human PSY₁ receptor can be extrapolated.

Discussion

In conclusion the present study has identified new pharmacological probes of PSY₁ receptors, including full agonists, partial agonists, and antagonists. The SAR of 1a indicates that the ribose ring oxygen may be readily substituted with carbon. Furthermore, analogues of constrained conformation, e.g. the methanocarba analogues, display enhanced receptor affinity. Additional 2-chloro and N⁶-methyl substitution is favorable for affinity at PSY₁ receptors, and nearly pure antagonism is maintained provided that the N⁶-methyl group is present.

Thus, the biological potency and efficacy of this series of bisphosphates appears to be highly dependent on subtle conformational factors, which would influence the orientation of the phosphate groups within the receptor binding site.

The sugar moiety of nucleosides and nucleotides in solution is known to exist in a rapid, dynamic equilibrium

between extreme 2-*exo*/3'-*endo* (*N*-) and 2'-*endo*/3'-*exo* (*S*-) conformations as defined in the pseudorotational cycle. While the energy gap between *N*- and *S*-conformation is in the neighborhood of 4kcal/mol, such a disparity can explain the difference between micromolar and nanomolar binding affinities. Using a molecular modeling approach, we have analyzed the sugar conformational requirements for a new class of bisphosphate ligands binding to the human PSY₁ receptor. As experimentally shown, the ribose ring Northern conformation appeared to be favored in recognition at human PSY₁ receptor (see Table 4). We have found new support to our recently presented hypothesis in which three important recognition regions are present in the bisphosphate molecular structures; The N¹ atom of the purine ring and the P atoms of both 3' and 5' phosphate groups. The *N*-conformation seems to be essential to maximize the electrostatic interactions between the negatively charged phosphates and the positively charged amino acids present in the receptor binding cleft, as well Arg128, Lys280, and Arg310.

Interestingly, the electrostatic contacts also appear to be crucial for the recognition of bisphosphate antagonists. Using superimposition analysis, a general pharmacophore model for the bisphosphate antagonists binding to the PSY₁ receptor has been proposed. According to the pharmacophore map, recognition of the bisphosphates antagonists at a common region inside the receptor binding site and, consequently, a common electrostatic potential profile is possible. As well for the agonists, the Northern conformation seems to be essential to maximize the electrostatic interactions between the negatively charged phosphates and the positively charged

amino acids presents in the receptor binding cleft. As we predicted using the previously reported PSY₁ receptor model, sugar moiety does not seem to be crucial for the ligand recognition process.

5 As already described, the simple addition of the N⁶-methyl group in several cases converted pure agonists to antagonists. From a pharmacological point of view, this is really a unique situation. With the addition of the N⁶-methyl group it is not possible to have a double hydrogen-bonding
10 interaction and, consequently, the activation pathway is blocked. However, for all the N⁶-methyl antagonists the possibility to participate in at least one of the two possible hydrogen bonds appears to be very important for the increase in affinity at the PSY₁ receptor.

15

Chemical Synthesis

Nucleosides and synthetic reagents were purchased from Sigma Chemical Co. (St. Louis, MO) and Aldrich (St. Louis, MO). 6-Chloro-2'-deoxypurine riboside was obtained from
20 Sigma. Several 2'-deoxynucleosides, including an anhydrohexitol-adenine nucleoside and 2'-deoxyaristeromycin were also synthesized.

Purity of compounds was checked using a Hewlett-Packard 1090 HPLC apparatus equipped with an SMT OD-5-60 RP-C18
25 analytical column (250 x 4.6 mm; Separation Methods Technologies, Inc., Newark, DE) in two solvent systems. System A: Linear gradient solvent system: 0.1 M TEAA/CH₃CN from 95/5 to 40/60 in 20 min and the flow rate was of 1 mL/min. System B: linear gradient solvent system: 5 mM
30 TBAP/CH₃CN from 80/20 to 40/60 in 20 min and the flow rate was

of 1 mL/min. Peaks were detected by UV absorption using a diode array detector. All derivatives showed more than 95% purity in the HPLC systems.

Purification of most of the nucleotide analogues for biological testing was carried out on DEAE-A25 Sephadex columns as described above. However, compounds 7b and 8a - c required HPLC purification (system a, semi-preparative C18 column) of the reaction mixtures.

General procedure of phosphorylation.

10 Method A: The nucleoside (0.1 mmol) and Proton Sponge[®] (107 mg, 0.5 mmol) were dried for several h in high vacuum at room temperature and then suspended in 2 mL of trimethyl phosphate. Phosphorous oxychloride (Aldrich, 37 μ L, 0.4 mmol) was added, and the mixture was stirred for 1 h at 0° C. The reaction was
15 monitored by analytical HPLC (eluting with a gradient consisting of buffer: CH₃CN in the ratio 95 : 5 to 40 : 60, in which the buffer was 0.1 M triethylammonium acetate (TEAA); elution time was 20 min; flow rate was 1 mL/min; column was SMT OD -5-60 RP-C18; detector was by UV in the E_{max} range of
20 260-300 nm). The reaction was quenched by adding 2 mL of triethylammonium bicarbonate buffer and 3 mL of water. The mixture was subsequently frozen and lyophilized. Purification was performed on an ion-exchange column packed with Sephadex-DEAE A-25 resin, a linear gradient (0.01 to 0.5 M) of 0.5 M
25 ammonium bicarbonate was applied as the mobile phase, and UV and HPLC were used to monitor the elution. All nucleotide bisphosphates were collected, frozen and lyophilized as the ammonium salts. All synthesized compounds gave correct molecular mass (high resolution FAB) and showed more than 95%
30 purity (HPLC, retention times are reported in Table 4).

Method B: Nucleoside (0.1 mmol) dried for several h in high vacuum at room temperature was dissolved in 2 mL of dry THF. Lithium diisopropylamide solution (Aldrich, 2.0 M in THF, 0.4 mmol) was added slowly at -78 °C. After 15 min tetrabenzyl pyrophosphate (Aldrich, 0.4 mmol) was added and the mixture was stirred for 30-60 min at -78°C. The reaction mixture was warmed to 0°C - rt and stirred for an addition period ranging from 2h to 24 h. Chromatographic purification (pTLC, CHCl₃:CH₃OH(10:1) gave the tetrabenzyl phosphorylated compound. This compound (20 mg) was dissolved in a mixture of methanol (2 mL) and water (1 mL) and hydrogenated over a 10 % Pd-on-C catalyst (10 mg) at rt for 62 h. The catalyst was removed by filtration and the methanol was evaporated. The residue was treated with ammonium bicarbonate solution and subsequently frozen and lyophilized. Purification, if necessary, was by the same procedure as in method A.

(*N*-Methanocarpa-2'-deoxyadenosine-3',5'-bis(diammonium phosphate) (4a) [(1R,2S,4S,5S)-1-[(phosphato)methyl]-4-(6-aminopurin-9-yl) bicyclo [3.1.0.] -hexane-2-phosphate tetraammonium salt]

Starting from 16 mg (0.06 mmol) of (*N*)-methanocarpa-2'-deoxyadenosine and following the general phosphorylation procedure A we obtained 1.8 mg (0.0037 mmol, 5.5 % yield) of the desired compound.

¹H-NMR (D₂O) δ 0.90 (1H, m, CH₂-6), 1.10 (1H, m, CH₂6'), 1.82 (1H, m, CH-5), 1.91 (1H, m, CH₂-3') 2.23 (1H, m, CH₂-3'), 3.49 (1H, d, J = 11.7 Hz, CH₂-OH), 4.16 (1H, d, J = 6.9 Hz, CH₂-2'), 8.39 (1H, s, H-2), 8.54 (1H, s, H-8).

³¹P-NMR (D₂O) δ 0.43 (s, 5'P); -0.19 (s, 3'P).

(N)-Methanocarba-N⁶-methyl-2'-deoxyadenosine-3',5'-
bis(diammonium phosphate) (4b)

(1R,2S,4S,5S)-1-[(phosphato)methyl]-4-(6-methylaminopurin-9-yl) bicyclo [3.1.0.]-hexane-2-phosphate tetraammonium salt]

5 13.5 mg (0.0170 mmol) of compound 18 was converted to the corresponding phosphoric acid analog using hydrogenation following the general procedure B. Purification was performed on an ion-exchange column packed with Sephadex-DEAE A-25 resin, linear gradient (0.01 to 0.5 M) of 0.5 M ammonium
10 bicarbonate was applied as the elan to give 3.0 mg (0.0060 mmol, 35.3 % yield) of the desired compound.

¹H-NMR (D₂O) δ 0.93-0.98 (1H, m, CH₂-6'), 1.17 (1H, m, CH₂-6'), 1.86-1.88 (1H, m, CH₅'), 1.94-1.98 (1H, m, CH₂-3'), 2.23-2.31 (1H, m, CH₂-3'), 3.09 (3H, bs, N⁶-CH₃), 3.61-3.64 (1H, m, CH₂OH), 4.51-4.55 (1H, m, CH₂OH), 5.01-5.03 (1H, m, CH-4'), 5.19-5.21 (1H, m, CH-2'), 8.22 (1H, s, H-2), 8.51 (1H, s, H-8). 31P-NMR (D₂O) δ 1.26, 1.92 (2s, 3'-P, 5'-P).

(N)-Methanocarba-N⁶-methyl-2-chloro-2'-deoxyadenosine-3',5'-bis(diammonium phosphate) (4c)

20 [(1R,2S,4S,5S)-1[(phosphato)methyl]-4-(2-chloro-6-aminopurin-9-yl) bicyclo [3.1.0]-hexane-2-phosphate tetraammonium salt]

The nucleoside, compound 23, reacted with tetrabenzyl pyrophosphate, as in Method B, followed by an alternative deprotection procedure. Starting from 10 mg (0.0323 mmol) of
25 (N)-methanorcarba-N⁶-methyl-2-chloro-2'-deoxyadenosine and following the general phosphorylation procedure (Method B) we obtained 9.5 mg (0.0114 mmol, 35.3 % yield) of the desired

compound, (N)-methanocarba-N⁶-methyl-2-chloro-2'-deoxyadenosine-3',5'-bis(dibenzyl phosphate).

¹H-NMR (CDCl₃) δ 0.75-0.81 (H, m, CH₂-6'), 1.03-1.08 (1H, m, CH₂-6'), 1.49-1.51 (1H, m, CH-5'), 1.84-1.94 (1H, m, CH₂-3'), 1.99-2.10 (1H, m, CH₂-3'), 3.12 (3H, bs, N⁶-CH₃), 4.11-4.20 (1H, m, CH₂OH), 4.50-4.55 (H, m, CH₂OH), 4.90-4.98, (8H, m, -OCH₂), 4.99-5.01 (1H, m, CH-4'), 5.23-5.30 (1H, m, CH-2'), 5.90 (1H, BS, NH), 7.20-7.29 (20H, m, C₆H₅), 7.82 (1H, s, H-8)

³¹P-NMR (D₂O) δ -0.58 (s, 5'P); -1.06 (s, 3'P).

10 MS(CI-NH₃) (M+1) 830 HRMS (FAB-) (M+Cs) Calcd. 962.1252; Found 962, 1252

9.5 mg (0.0114 mmol) of the tetrabenzyl-protected intermediate added to dry CH₂Cl₂ (1.0 mL) was cooled to -78 °C under argon and treated with 100 µL of boron trichloride solution (1M in CH₂Cl₂) and 100 µL of anisole. The reaction mixture was stirred for 12 hr at 0 °C to rt and extracted with triethylamine solution. Purification was performed on an ion-exchange column packed with Sephadex-DEAE A-25 resin, linear gradient (0.01 to 0.5 M) of 0.5 M ammonium bicarbonate was applied as the eluent to give 0.4 mg (0.0007 mmol, 6.52 yield) of the desired compound 4c.

¹H-NMR (D₂O) δ 0.91-0.96 (1H, m, CH₂-6'), 1.12-1.16 (1H, m, CH₂-6'), 1.80-1.84 (1H, m, CH-5'), 1.85-1.98 (1H, m, CH₂-3'), 2.20-2.50 (1H, m, CH₂-3'), 3.08 (3H, bs, N⁶-CH₃), 3.57-3.60 (1H, m, CH₂OH), 4.52-4.67 (1H, m, CH₂OH), 4.94-4.96 (1H, m, CH-4'), 5.18-5.21 (1H, m, CH-2'), 8.52 (1H, s, H-8)

³¹P-NMR (D₂O) δ 1.82, 2.52 (2s, 3'-P, 5'P)

(S)-Methanocarba-2', deoxyadeonosine-3',5'-bis(diammonium phosphate (5) [(1S,3S,4R,5S)-4-[(phosphato)methyl]-1-(6-aminopurin-9-yl) bicyclo [3.1.0]-hexane-3-phosphate tetraammonium salt]

5 Starting from 16 mg (0.06 mmol) of (S)-methanocarba-2'-deoxyadenosine and following the general phosphorylation procedure A, we obtained 2.1 mg (0.0043 mmol, 7.55 yield) of the desired compound 5.

¹H-NMR (D₂O) δ 1.36 (1H, m, CH₂-6'), 1.53 (1H, t, J = 4.8 Hz, CH₂-6'), 2.05 (1H, m, CH₂-5'), 2.30 (1H, m, CH-4'), 2.46 (2H, m, CH₂-2'), 3.97 (2H, m, CH₂OH), 4.45 (1H, d, J = 6.6 Hz, CH-3'), 8.16 (1H, s, H-2), 8.30 (1H, s, H-8). ³¹P-NMR (D₂O) δ 0.85 (bs, 5'P); 0.31 (bs, 3'P).

15 [(1S,3S,4R,5S)-1-[(Hydroxy)methyl]-2-hydroxy-4-(6-methylaminopurin-9-yl) bicyclo [3.1.0]-hexane (17b)

The Dimroth rearrangement (Scheme 2) was carried out on (N)-methanocarba-2'-deoxyadenosine. Specifically, the (N)-methanorcarba-2'-deoxyadenosine (17a, 50.0 mg, 0.191 mmol) was heated at 40° C with methyl iodide (71.5 μL, 1.15 mmol) in dry DMF (2.0 mL) for 48 h. The solvent was evaporated under reduced pressure, and the residue was heated at 90 °C with ammonium hydroxide (4.0 mL) for 4 h. The water was evaporated, and the residue was purified by pTLC using MeOH; CHCl₃ (1:9) to afford compound 17b as a colorless solid (40 mg, 0.15 mmol, 76%).

¹H-NMR (CD₃OD) δ 0.77.-0.81 (1H, m, CH₂-6'), 1.03-1.07 (1H, m, CH₂-6'), 1.68-1.72 (1H, m, CH-5'), 1.79-1.89 (1H, m, CH₂-3'), 2.00-2.07 (1H, m, CH₂-3'), 3.12 (3H, bs, N⁶-CH₃), 3-33 (1H, d, J = CH₂OH), 4.29 (1H, d, J = 11.7 Hz, CH₂OH), 4.89-4.92 (1H, m,

CH-4'), 5.02 (1H, d, $J = 6.9$ Hz, CH-2'), 8.24 (1H, s, H-2), 8.49 (1H, s, H-8).

MS(CI-NH₃) : 276 (M+1) 830 HRMS(FAB-) (M+Cs) Calcd. 275.1382; Found 275.1389.

5 (N)-Methanocarba-N⁶-methyl-2'-deoxyadenosine-3',5'-

bis(dibenzylphosphate) (18)

[(1S,2S,4Z,5S)-1-[(dibenzylphosphato)methyl]-4-(6-methylaminopurin-9-yl) bicyclo [3.1.0]-hexane-2-dibenzylphosphate]

10 Starting from 20.0 mg (0.0726 mmol) of N-methanocarba-N⁶-methyl-2'-deoxyadenosine 17b and following the general phosphorylation procedure (Method B we obtained 13.5 mg (0.0170 mmol, 23.4 % yield) of the desired protected intermediate, 18 as shown in Scheme 2.

15 ¹H-NMR (CDCl₃) δ 0.73-0.78 (1H, m, CH₂-6'), 0.94-.0.98 (1H, m, CH₂-6'), 1.53-1.54 (1H, m, CH-5'), 1.81-1.91 (1H, m, CH₂-3'), 2.05-2.13 (1H, m, CH₂-3'), 3.15 (3H, bs, N⁶-CH₃), 3.70-3.83 (1H, m, CH₂OP), 4.49-4.55 (1H, m, CH₂OP), 4.89-5.00 (8H, m, OCH-2), 5.02-5.06 (1H, m, CH-4'), 5.27-5.32 (1H, m, CH-2'), 5.86 (1H, bs, NH), 7.21-7.23 (20H, m, C₆H₅), 7.86 (1H, s, H-2), 8.31 (1H, s, H-8). ³¹P-NMR (D₂O) δ -0.56, -1.05 (2s, 3'-P, 5'-P) HRMS (FAB-) (M-Cs) Calcd. 928.1641; Found 928.1700.

[(1S,2S,4Z,5S)-1-[(Benzyloxy)methyl]-2-benzyloxy-4-(2-6-dichloropurin-9-yl) bicyclo [3.1.0]-hexane (21)

25 To an ice cold solution of triphenylphosphine (278 mg, 1.06 mmol) in dry THF (2 mL) was added diethylazadicarboxylate (170 μ L, 1.06 mmol) dropwise under a nitrogen atmosphere, and the mixture was stirred for 20 min until the solution turned red orange (Scheme 3). This mixture was added dropwise to a cold

stirred mixture of the starting alcohol (135 mg, 0.417 mmol) and 2,6-dichloropurine (157 mg, 0.883 mmol) under a nitrogen atmosphere. The reaction mixture was stirred in an ice bath for 30 min and then allowed to warm to room temperature, and stirring continued for 12 h. Solvent was removed by nitrogen purge, and the residue was purified by pTLC using EtOAc : petroleum ether (1 : 1) to afford a thick liquid (132 mg, 0.263 mmol, 64%).

¹H NMR: (CD₃OD) δ 0.85 (m, 1H), 1.13 (m, 1H), 1.59 (m, 1H), 1.68 (m, 1H), 2.06 (m, 1H), 3.17 (d, J = 10.8 Hz, 1H), 4.11-4.57 (m, 5H), 5.20 (d, J = 6.9 Hz, 1H), 6.6 (bs, 1H), 7.23-7.37 (m, 10H), 8.98 (s, 1H).

MS : (EI) 494 (M+).

[(1R,2S,4S,5S)-1-[(Benzyloxy)methyl]-2-benzyloxy-4-(2-chloro-6-methylaminopurin-9-yl) bicyclo [3.1.0]-hexane (22)

Compound 21 (100 mg, 0.202 mmol) was dissolved in methylamine in methanol (30 % solution, 3mL) and was stirred at rt for 12 h under a nitrogen atmosphere. The solvent was evaporated, and the crude product was purified by pTLC using EtOAc : petroleum ether (6 : 4) to afford 22 as a light yellow solid (86 mg, 0.176 mmol, 88 %).

¹H NMR: (CD₃OD) δ 0.70 (m, 1H), 1.06 (m, 1H), 1.50 (m, 1H), 1.76 (m, 1H), 1.96 (m, 1H), 3.01 (s, 3H), 3.08 (m, 2H), 4.03 (m, 4H), 4.45 (bs, 1H), 5.02 (bs, 1H), 8.38 (s, 1H).

MS : (CI) : 490 (M+1).

[(1R,2S,4S,5S)-1-[(Hydroxy)methyl]-2-hydroxy-4-(2-chloro-6-methylaminopurin-9-yl) bicyclo [3.1.0]-hexane (23)

Compound 22 (40 mg 0.0816 mmol) was dissolved in dry CH₂Cl₂ (1.0 mL), and hydrogenated using BCl₃ (1M in CH₂Cl₂, 175 μL) for 50 min at -78 °C under argon. The solvent was evaporated,

and the crude product was purified by pTLC using CHCl_3 : MeOH (10 : 1) to afford 23 as a light yellow solid (10.0 mg, 0.0323 mmol, 39.6 %).

^1H NMR: (CD_3OD) δ 0.77-0.81 (1H, m, CH_2 -6'), 1.02-1.05 (1H, m, CH_2 -6'), 1.65-1.68 (1H, m, CH-5'), 1.78-1.91 (1H, m, CH_2 -3'), 1.99-2.07 (1H, m, CH_2 -3'), 3.08 (3H, bs, N^6 - CH_3), 3.37 (1H, d, J = 11.7 Hz, CH_2OH), 4.27 (1H, d, J = 11.7 Hz, CH_2OH), 4.89-4.91 (1H, m, CH-4), 4.97 (1H, d, J = 6.8 Hz, CH-2'), 8.46 (1H, s, H-8).

MS : (CI-NH_3): 310 ($\text{M}+1$), HRMS (FAB-): Calcd 309.0992, Found 309.0991.

Pharmacological Analyses.

P2Y_1 receptor promoted stimulation of inositol phosphate formation by adenine nucleotide analogues was measured in turkey erythrocyte membranes as previously described. The $\text{K}_{0.5}$ values were averaged from 3-8 independently determined concentration-effect curves for each compound. Briefly, 1 mL of washed turkey erythrocytes was incubated in inositol-free medium (DMEM; Gibco, Gaithersburg MD) with 0.5 mCi of 2- $[\text{^3H}]$ myo-inositol (20Ci/mmol: American Radiolabelled Chemicals, Inc., St. Louis MO) for 18-24 h in a humidified atmosphere of 95% air/5% CO_2 at 37 °C. Erythrocyte ghosts were prepared by rapid lysis in hypotonic buffer (5 mM sodium phosphate, pH 7.4, 5 mM MgCl_2 , 1mM EGTA) as described. Phospholipase C activity was measured in 25 μL of $[\text{^3H}]$ inositol-labeled ghosts (approximately 175 μg of protein, 200-500,000 cpm/assay) in a medium containing 424 μM CaCl_2 , 0.91 mM MgSO_4 , 2mM EGTA, 115 mM KCl, 5 mM KH_2PO_4 , and 10 mM Hepes pH 7.0. Assays (200 μL

final volume) contained 1 μ M GTP γ S and the indicated concentrations of nucleotide analogues. Ghosts were incubated at 30 °C for 5 min, and total [^{3} H]inositol phosphates were quantitated by anion exchange chromatography as previously
5 described.^{7,36}

Data analysis.

Agonist potencies were calculated using a four-parameter logistic equation and the GraphPad software package (GraphPad, San Diego, CA). EC₅₀ values (mean $\pm$ standard error) represent
10 the concentration at which 50 % of the maximal effect is achieved. Relative efficacy (%) was determined by comparison with the effect produced by a maximal effective concentration of 2-MeSADP in the same experiment.

Antagonist IC₅₀ values (mean $\pm$ standard error) represent
15 the concentration needed to inhibit by 50 % the effect elicited by 10 nM 2-MeSADP. The percent of maximal inhibition is equal to 100 minus the residual fraction of stimulation at the highest antagonist concentration.

All concentration-effect curves were repeated in at least
20 three separate experiments carried out with different membrane preparations using duplicate or triplicate assays.

Table 3

Stimulation of PLC at turkey erythrocyte P2Y₁ receptors (agonist effect) and the inhibition of PLC stimulation elicited by 10 nM 2-MeSADP (antagonist effect), for at least two separate determinations.

Compound	Agonist Effect, % of maximal increase ^a	EC ₅₀ , μ M ^a	Antagonist Effect, % of maximal inhibition ^b	IC ₅₀ , μ M ^b (n)
1a ^{c,e} (MRS 2179)	NE		99 $\pm$ 1	0.331 $\pm$ 0.059 (5)
1b ^e	NE		95 $\pm$ 1	0.206 $\pm$ 0.053
1c ^e	4	d	96 $\pm$ 2	1.85 $\pm$ 0.74
1d ^e	6 $\pm$ 2	d	94 $\pm$ 2	0.362 $\pm$ 0.119
4a	92 $\pm$ 5	0.155 $\pm$ 0.021	NE	
4b	NE		100	0.157 $\pm$ 0.060
4c	NE		100	0.0516 $\pm$ 0.0008
5	41 $\pm$ 13	13.3	34% at 100 μ M	small decrease

- ^a Agonist potencies were calculated using a four-parameter logistic equation and the GraphPad software package (GraphPad, San Diego, CA). EC₅₀ values (mean $\pm$ standard error) represent the concentration at which 50% of the maximal effect is achieved. Relative efficacies (%) were determined by comparison with the effect produced by a maximal effective concentration of 2-MeSADP in the same experiment. Small increase refers to <10% at 100 μ M.
- ^b Antagonist IC₅₀ values (mean $\pm$ standard error) represent the concentration needed to inhibit by 50% the effect elicited by 10 nM 2-MeSADP. The percent of maximal inhibition is equal to 100 minus the residual fraction of stimulation at the highest antagonist concentration.
- ^c 1a, MRS 2179; 4c, MRS 2279.
- ^d EC₅₀ was not calculated for increases of 10% at 100 μ M.
- ^e values from refs. 17, 19.
- NE no effect at 100 μ M.

TABLE 4

Synthetic data for nucleotide derivatives, including structural verification using high resolution mass spectroscopy and purity verification using HPLC.

No	Formula	FAB (M-H ⁺)		HPLC (rt; min) ^a		Method, Yield (%) ^b
		Calcd	Found	System A	System B	
2	C ₁₀ H ₁₅ O ₉ N ₅ P ₂	410.0267	410.0269	3.53	10.72	B, 21.7
3b	C ₁₂ H ₁₉ O ₈ N ₅ P ₂	422.0631	422.0664	3.41	8.21	B, 8.0
4a	C ₁₂ H ₁₇ O ₈ N ₅ P ₂	420.0474	420.0482	3.92	7.30	A, 5.5
4b	C ₁₃ H ₁₉ O ₈ N ₅ P ₂	434.0631	434.0622	5.91	7.83	B, 8.3
4c	C ₁₃ H ₁₈ O ₈ N ₅ P ₂ Cl	468.0241	468.0239	8.05	8.54	B, 2.3
5	C ₁₂ H ₁₇ O ₈ N ₅ P ₂	420.0474	420.0481	4.02	6.84	A, 7.5
6	C ₁₁ H ₁₆ O ₈ N ₅ P ₂ Cl	442.0084	442.0070	6.67	6.82	A, 24.3
7b	C ₁₂ H ₂₀ O ₁₂ N ₅ P ₃	518.0237	518.0243	4.98	12.74	A, 1.8
7c	C ₁₂ H ₁₉ O ₉ N ₅ P ₂	438.0580	438.0580	4.63	9.36	B, 50.1
7d	C ₁₂ H ₁₈ O ₉ N ₅ P ₂ Cl	472.0201	472.0190	5.67	9.97	B, 31.3
8a	C ₁₂ H ₂₀ O ₈ N ₆ P ₂	437.0740	437.0721	2.37	8.78	8.0
8b	C ₁₂ H ₂₁ O ₁₁ N ₆ P ₃	517.0403	517.0404	2.42	9.23	7.2
8c	C ₁₂ H ₂₂ O ₁₄ N ₆ P ₄	597.0066	597.0053	2.96	10.02	4.0

^a Purity of each derivative was 95%, as determined using HPLC with two different mobile phases. System A: gradient of 0.1M TEAA/CH₃CN from 95/5 to 40/60 and System B: gradient of 5mM TBAP/ CH₃CN from 80/20 to 40/60.

^b Phosphorylation methods: Method A refers to use of phosphorous oxychloride, and Method B refers to use of tetrabenzyl pyrophosphate/lithium diisopropylamide followed by hydrogenation. The percent yields refer to overall yield for each phosphorylation sequence. For the method of synthesis of 8 refer to Experimental Section.

Abbreviations

AIBN, 2,2' -azobisisobutyronitrile;

ATP, adenosine 5'-triphosphate;

DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene;


- DCTIDS, 1,3-dichlorotetraisopropyl-1,1,3,3, -disiloxane;
DEAD, diethylazadicarboxylate;
DEAE, diethylaminoethyl;
DMAP, 4-dimethylaminopyridine;
5 DMF, dimethylformamide;
DMSO, dimethylsulfoxide;
FAB, fast atom bombardment (mass spectroscopy);
HPLC, high pressure liquid chromatography;
MS, mass spectroscopy;
10 HRMS, high resolution mass spectroscopy;
LDA, lithium diisopropylamide;
2-MeSADP, 2-methylthioadenosine-5'-diphosphate;
TBAP, tetrabutylammonium phosphate;
TEAA, triethylammonium acetate;
15 THF, tetrahydrofuran;

While this invention has been described with an emphasis upon preferred embodiments, it will be obvious to those of ordinary skill in the art that variations of the preferred embodiments may be used and that it is intended that the invention may be practiced otherwise than as specifically described herein. Accordingly, this
20 invention includes all modifications encompassed within the scope of the invention as defined by the following claims.

CLAIMS

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A compound of the formula



wherein

R₁ is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, sulfonyl, arylsulfonyl, thiazolyl, (benzothiazolyl) thio-2-propyl, or bicyclic alkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, phosphoryl, halo, sulfonyl, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido;

R₂ is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, thiol, cyano, alkylthio, or arylalkylthio;

R₃, R₄, and R₅, are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, diphosphoryl, triphosphoryl, imidodiphosphato, imidotriphosphato, methylene diphosphato, methylene triphosphato, halomethylene diphosphato, halomethylene triphosphato, borono phosphato, borono diphosphato, borono triphosphato, phosphorothioato, phosphonyl, boronyl, thiophosphoryl, thiodiphosphoryl, thiotriphosphoryl or vanadyl, and can be the same or different;

R₆ is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;

R₇ is methylene, dihalomethylene, carbonyl, sulfoxide;


and at least one of R₁, R₂, and R₆ is other than hydrogen; and

R₈ is CH or nitrogen;

wherein said alkyl is a C₁-C₂₀ alkyl; said alkenyl is a C₂-C₂₀ alkenyl; said alkynyl is a C₂-C₂₀ alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring;

wherein any of R₂ - R₈ may be further substituted with a substituent selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; or a salt of said compound.

2. A compound of the formula



wherein

R₁ is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, sulfonyl, arylsulfonyl, thiazolyl, (benzothiazolyl) thio-2-propyl, or bicyclic alkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, halo, sulfonyl, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, sulfonamido, carboxyl, and carboxamido;

R₂ is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, thiol, cyano, alkylthio, or arylalkylthio;

R₃ is hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, diphosphoryl, triphosphoryl, phosphonyl, boronyl, thiophosphoryl, thiodiphosphoryl, thiotriphosphoryl or vanadyl;

R₄ and R₅, are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, diphosphoryl, triphosphoryl, imidodiphosphato, imidotriphosphato, methylene diphosphato, methylene triphosphato, halomethylene diphosphato, halomethylene triphosphato, borono phosphato, borono diphosphato, borono triphosphato, phosphorothioate-2-thioether, phosphonyl, boronyl, thiophosphoryl, thiodiphosphoryl, thiotriphosphoryl or vanadyl, and can be the same or different;

R₆ is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;

R₇ is methylene, dihalomethyl, carbonyl, or sulfoxide;

and at least one of R₁, R₂, and R₆ is other than hydrogen; and

R₈ is CH or nitrogen;

wherein said alkyl is a C₁-C₂₀ alkyl; said alkenyl is a C₂-C₂₀ alkenyl; said alkynyl is a C₂-C₂₀ alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring;

wherein any of R₂ - R₈ may be further substituted with a substituent selected from the group consisting of amino, cyano, alkoxyl, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; or a salt of said compound.

3. The compound or salt of claim 1 or 2, wherein R₁ is alkyl, cycloalkyl, alkoxy, aryl, arylalkyl, bicycloalkyl, or sulfonyl.

4. The compound or salt of claim 1 or 2, wherein R₁ is methyl, cyclopentyl, cyclohexyl, phenyl, *R*-phenylisopropyl, benzyl, or phenylethyl;

R₂ is chloro;

and R₆ is C₁-C₆ alkylamino, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

5. The compound or salt of claim 1 or 2, wherein R₁ is methyl group and R₂ is chloro, alkylthio, or arylalkylthio.


6. The compound or salt of claim 1 or 2, wherein R₆ is methyl and R₂ is chloro, alkylthio, arylalkylthio or hydrogen.

7. The compound or salt of claim 1 or 2, wherein R₆ is halo and R₂ is chloro, alkylthio, arylalkylthio or hydrogen.

8. The compound or salt of claim 1 or 2, wherein R₂ is chloro.


9. The compound or salt of claim 1 or 2, wherein R_1 is methyl, R_2 is chloro, and R_3 is hydrogen.

10. The compound or salt of claim 1, wherein the compound has the formula




wherein R_1 is iodobenzyl, or cyclopentyl and R_2 is hydrogen or chloro.

11. A compound of the formula




or a salt thereof.

12. The compound or salt of claim 1, wherein the compound has the formula



13. A compound of the formula:




wherein R_1 , R_2 , and R_9 are each independently hydrogen, alkyl, alkenyl, alkynyl, or aminoalkyl, R_3 and R_4 are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and can be the same or different; R_5 is hydrogen, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and can be the same or different; R_8 and R_7 are each independently sulfur or oxygen; and R_{10} is methylene, dihalomethylene, carbonyl, or sulfoxide;

wherein said alkyl is a C_1 - C_{20} alkyl; said alkenyl is a C_2 - C_{20} alkenyl; said alkynyl is a C_2 - C_{20} alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring;

wherein any of R_2 - R_5 and R_9 - R_{10} other than hydrogen may be further substituted with a substituent selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; or a salt of said compound.

14. A compound of the formula:



wherein R_1 , R_2 , and R_9 are each independently hydrogen, alkyl, alkenyl, alkynyl, or aminoalkyl;

R_3 , R_4 , and R_5 are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and can be the same or different;


R_8 and R_7 are each independently sulfur or oxygen;

and R_{10} is dihalomethylene, carbonyl, or sulfoxide;

wherein said alkyl is a C_1 - C_{20} alkyl; said alkenyl is a C_2 - C_{20} alkenyl; said alkynyl is a C_2 - C_{20} alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring;

wherein any of $R_1 - R_5$ and $R_9 - R_{10}$ other than hydrogen may be further substituted with a substituent selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; or a salt of said compound.

15. A compound of the formula:



wherein

R_1 , R_2 , and R_9 are each independently hydrogen, alkyl, alkenyl, alkynyl, aminoalkyl;

R_3 is hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and can be the same or different;

R_4 and R_5 are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, phosphonyl, boronyl, or vanadyl, and can be the same or different;


R_8 and R_7 are each independently sulfur or oxygen;

and R_{10} is methylene, dihalomethylene, carbonyl, or sulfoxide; wherein said alkyl is a C_1 - C_{20} alkyl; said alkenyl is a C_2 - C_{20} alkenyl; said alkynyl is a C_2 - C_{20} alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring;

wherein any of R_1 - R_5 and R_9 other than hydrogen may be further substituted with a substituent selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, phosphoryl, sulfonyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; or a salt of said compound.

16. The compound or salt of any one of claims 13-15, wherein R_1 is methyl.

17. A compound of the formula



wherein

R_1 is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, sulfonyl, arylsulfonyl, thiazolyl, (benzothiazolyl) thio-2-propyl, or bicyclic alkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, phosphoryl, halo, sulfonyl, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, sulfonamido, carboxyl, thiohydroxyl, and carboxamido; R_2 is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, thiol, cyano, alkylthio, or arylalkylthio; R_3 , R_4 , and R_5 are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, diphosphoryl, triphosphoryl, imidodiphosphato, imidotriphosphato, methylene diphosphato, methylene triphosphato, halomethylene diphosphato, halomethylene triphosphato, borono phosphato, borono diphosphato, borono triphosphato, phosphorothioate-2-thioether, phosphonyl, boronyl, thiophosphoryl, thiodiphosphoryl, thiotriphosphoryl or vanadyl, and can be the same or different;

R_6 is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;

R_7 is methylene, dihalomethylene, carbonyl, or sulfoxide;

and at least one of R_1 , R_2 , and R_6 is other than hydrogen; and

R_8 is CH or nitrogen;


wherein said alkyl is a C₁-C₂₀ alkyl; said alkenyl is a C₂-C₂₀ alkenyl; said alkynyl is a C₂-C₂₀ alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring; or a salt of said compound.

18. The compound or salt of claim 1 or 17, wherein R₂ is hydrogen or halo.

19. The compound or salt of claim 1 or 17, wherein R₂ is alkyl, aryl, arylamino, aryloxy, alkynyl, or alkenyl.

20. The compound or salt of claim 1 or 17, wherein R₂ is thiol, alkylthio, arylalkylthio, or cyano.

21. A compound of the formula



wherein

R₁ is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, arylsulfonyl, thiazolyl or bicycloalkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, dihydrogen phosphato, halo, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, sulfonamido, carboxyl, and carboxamido;

R₂ is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, mercapto, cyano, alkylthio, or arylalkylthio;

R₄ and R₅ are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, dihydrogen phosphato, trihydrogen diphosphato, tetrahydrogen triphosphato, dihydrogen imidodiphosphato, trihydrogen imidotriphosphato, dihydrogen methylenediphosphato, or dihydrogen halomethylene diphosphato, and can be the same or different;

R₃ is hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, dihydrogen phosphato, dihydrogen imidodiphosphato, dihydrogen methylenediphosphato, or dihydrogen halomethylene diphosphato;

R₆ is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;

R₇ is methylene, dihalomethylene, carbonyl, or sulfoxide;

and at least one of R₁, R₂, and R₆ is other than hydrogen; and

R₈ is nitrogen;

wherein said alkyl is a C₁-C₂₀ alkyl; said alkenyl is a C₂-C₂₀ alkenyl; said alkynyl is a C₂-C₂₀ alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring; wherein any of R₂-R₇ other than hydrogen may be further substituted with one or more substituents selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, dihydrogen phosphato, sulfonamido, carboxyl, mercapto, and carboxamido, or a salt of said compound.

22. The compound of claim 21, wherein R₁ is alkyl, cycloalkyl, alkoxy, aryl, arylalkyl, or bicycloalkyl.

23. The compound of claim 21, wherein R₁ is methyl, cyclopentyl, cyclohexyl, phenyl, (*R*)-phenylisopropyl, benzyl, or phenylethyl;

R₂ is chloro; and

R₆ is C₁-C₆ alkylamino, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

24. The compound of claim 21, wherein R₆ is methyl and R₂ is chloro, alkylthio, arylalkylthio or hydrogen.


25. The compound of claim 21, wherein R₆ is halo and R₂ is chloro, alkylthio, arylalkylthio or hydrogen.

26. The compound of claim 21, wherein R₂ is chloro.

27. The compound of claim 21, wherein R₁ is methyl, R₂ is chloro and R₃ is hydrogen.


28. The compound of claim 21, wherein the compound has the formula

65



wherein R_1 is iodobenzyl, or cyclopentyl and R_2 is hydrogen or chloro.

29. A compound of the formula



wherein

R_1 is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, arylsulfonyl, thiazolyl or bicycloalkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, halo, dihydrogen phosphato, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, sulfonamido, carboxyl, thiol, and carboxamido;

R_2 is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, mercapto, cyano, alkylthio, or arylalkylthio;

R_4 and R_5 are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, or dihydrogen phosphato, and can be the same or different;

R_3 is hydroxyl, alkoxy, alkyl, alkenyl, aryl, acyl, alkylamino, arylamino, or dihydrogen phosphato;

R_6 is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;


R_7 is methylene, dihalomethylene, carbonyl, or sulfoxide;

and at least one of R_1 , R_2 , and R_6 is other than hydrogen; and

R_8 is nitrogen;

wherein said alkyl is a C_1 - C_{20} alkyl; said alkenyl is a C_2 - C_{20} alkenyl; said alkynyl is a C_2 - C_{20} alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring; or a salt of said compound.

30. A compound of the formula:



wherein

R_1 is hydrogen, alkyl, cycloalkyl, alkoxy, cycloalkoxy, aryl, arylalkyl, acyl, arylsulfonyl, thiazolyl or bicycloalkyl, each of which, other than hydrogen, may be further substituted with a member selected from the group consisting of hydroxyl, dihydrogen phosphato, halo, amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, sulfonamido, carboxyl, and carboxamido;

R_2 is hydrogen, halo, alkyl, aryl, arylamino, aryloxy, alkynyl, alkenyl, mercapto, cyano, alkylthio, or arylalkylthio;

R_3 , R_4 , and R_5 are each independently alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, dihydrogen phosphato, trihydrogen diphosphato, tetrahydrogen triphosphato, dihydrogen imidodiphosphato, trihydrogen imidotriphosphato, dihydrogen methylene diphosphato, or dihydrogen halomethylene diphosphato, and can be the same or different;

R_6 is hydrogen, alkyl, alkenyl, alkynyl, heteroaryl or aminoalkyl;

R_7 is methylene, dihalomethylene, carbonyl, or sulfoxide;

and at least one of R_1 , R_2 , and R_6 is other than hydrogen; and

R_8 is nitrogen;

wherein said alkyl is a C₁-C₂₀ alkyl; said alkenyl is a C₂-C₂₀ alkenyl; said alkynyl is a C₂-C₂₀ alkynyl; and said aryl has no more than 8 carbon atoms in an aromatic ring; wherein any of R₂-R₇ other than hydrogen may be further substituted with one or more substituents selected from the group consisting of amino, cyano, alkoxy, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, acyl, halo, hydroxyl, dihydrogen phosphato, sulfonamido, carboxyl, mercapto, and carboxamido, or a salt of said compound.

31. The compound of claim 30, wherein R₄ and R₅ are each independently dihydrogen phosphato, trihydrogen diphosphato, tetrahydrogen triphosphato, dihydrogen imidodiphosphato, trihydrogen imidotriphosphato, dihydrogen methylene diphosphato, or dihydrogen halomethylene diphosphato, and can be the same or different.

32. The compound of claim 31, wherein R₄ and R₅ are each independently dihydrogen phosphato, trihydrogen diphosphato, or tetrahydrogen triphosphato.

33. The compound of claim 32, wherein R₄ and R₅ are dihydrogen phosphato.


34. A compound selected from the group consisting of (*N*)-methanocarba-2'-deoxyadenosine-3',5'-bis(diammonium phosphate), (*N*)-methanocarba-N⁶-methyl-2'-deoxyadenosine-3',5'-bis(diammonium phosphate), and (*N*)-methanocarba-N⁶-methyl-2-chloro-2'-deoxyadenosine-3',5'-bis(diammonium phosphate).

35. A compound selected from the group consisting of (*N*)-methanocarba-N⁶-methyl-2'-deoxyadenosine-3',5'-bis(diammonium phosphate) and (*N*)-methanocarba-N⁶-methyl-2-chloro-2'-deoxyadenosine-3',5'-bis(diammonium phosphate).

36. The compound of claim 21, wherein R₁ is iodobenzyl, R₂ is hydrogen or chloro, R₃ and R₄ are hydroxyl, R₅ is methylamino, R₆ is hydrogen, and R₇ is carbonyl.


37. A compound of the formula:

68



wherein R_1 is methyl; R_2 is chloro or iodo; R_3 is hydrogen; R_4 and R_5 are dihydrogen phosphato; R_6 is hydrogen; and R_7 is methylene.

38. A compound of the formula:



wherein R_1 is hydrogen, alkenyl, alkynyl, or aminoalkyl;

R_2 and R_9 are independently hydrogen, alkyl, alkenyl, alkynyl, or aminoalkyl;

R_3 , R_4 , and R_5 , are each independently hydrogen, hydroxyl, alkoxy, alkyl, alkenyl, alkynyl, aryl, acyl, alkylamino, arylamino, phosphoryl, or phosphonyl;

R_8 and R_7 are each independently sulfur or oxygen;

and R_{10} is methylene, dihalomethyl, carbonyl, or sulfoxide;

or a salt of said compound;

with the proviso that when R_1 is hydrogen, R_2 is methyl, R_4 and R_5 are hydroxyl, R_7 and R_8 are O, R_9 is hydrogen, and R_{10} is methylene, R_3 is not hydrogen.

39. The compound or salt of claim 38, wherein the phosphoryl group is a diphosphoryl group.

40. The compound or salt of claim 39, wherein R_1 , R_2 , and R_9 are hydrogen, R_7 and R_8 are oxygen, R_{10} is methylene, R_5 is diphosphoryl, and R_3 and R_4 are hydrogen or hydroxyl.
41. The compound or salt of claim 40, wherein R_3 and R_4 are hydroxyl.
42. The compound or salt of claim 40, wherein R_3 is hydrogen and R_4 is hydroxyl.
43. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound or salt of any one of claims 1-42.
44. Use of the compound or salt of any one of claims 1-42 in the preparation of a medicament for the treatment of an airway disease, cancer, cardiac arrhythmia, cardiac ischemia, epilepsy, Huntington's Disease, an immunodeficient disorder, an inflammatory disorder, neonatal hypoxia, neurodegenerative disease, pain, Parkinson's Disease, renal failure, schizophrenia, a sleep disorder, stroke, thrombosis, urinary incontinence, diabetes, psoriasis, septic shock, brain trauma, glaucoma, or congestive heart failure in a mammal.
45. Use of a compound or salt of any one of claims 1-42 as an agonist or antagonist of an adenosine, ATP, or UTP receptor.
46. Use of a compound or salt of any one of claims 30-37 as an agonist or antagonist of a P2Y1 receptor.