




(86) Date de dépôt PCT/PCT Filing Date: 2009/08/19
(87) Date publication PCT/PCT Publication Date: 2010/02/25
(85) Entrée phase nationale/National Entry: 2011/02/16
(86) N° demande PCT/PCT Application No.: US 2009/054268
(87) N° publication PCT/PCT Publication No.: 2010/022125
(30) Priorité/Priority: 2008/08/20 (US61/090,478)

(51) Cl.Int./Int.Cl. *C07D 239/46* (2006.01),
A61K 31/505 (2006.01), *A61K 31/506* (2006.01),
C07D 401/06 (2006.01), *C07D 401/14* (2006.01),
C07D 403/06 (2006.01), *C07D 405/06* (2006.01),
C07D 409/06 (2006.01), *C07D 413/06* (2006.01),
C07D 417/06 (2006.01)

(71) Demandeurs/Applicants:
SOUTHERN RESEARCH INSTITUTE, US;
SCHERING CORPORATION, US

(72) Inventeurs/Inventors:
NJOROGÉ, F. GEORGE, US;
ARASAPPAN, ASHOK, US;
BENNETT, FRANK, US; ...

(54) Titre : DERIVES DE PYRIDINE ET DE PYRIMIDINE A SUBSTITUTION ETHYLYLE ET LEUR UTILISATION DANS
LE TRAITEMENT D'INFECTIONS VIRALES
(54) Title: ETHYNYL-SUBSTITUTED PYRIDINE AND PYRIMIDINE DERIVATIVES AND THEIR USE IN TREATING
VIRAL INFECTIONS



(57) Abrégé/Abstract:

The present invention provides compounds of Formula (I): (Chemical formula should be inserted here as it appears on abstract in paper form) (I) and tautomers, isomers, and esters of said compounds, and pharmaceutically acceptable salts, solvates, and



(72) **Inventeurs(suite)/Inventors(continued)**: PIWINSKI, JOHN J., US; SHIH, NENG-YANG, US; KWONG, CECIL D., US; ANANTHAN, SUBRAMANIAM, US; CLARK, JEREMY, US; FOWLER, ANITA T., US; GENG, FENG, US; KEZAR, HOLLIS S., III, US; MADDRY, JOSEPH A., US; REYNOLDS, ROBERT C., US; ROYCHOWDHURY, ABHIJIT, US; SECRIST, JOHN A., III, US

(74) **Agent**: OGILVY RENAULT LLP/S.E.N.C.R.L., S.R.L.

(57) **Abrégé(suite)/Abstract(continued)**:

prodrugs of said compounds, wherein wherein each of R, R¹, X, Y, Z, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁸, R¹⁹ and n is selected independently and as defined herein. Compositions comprising such compounds are also provided. The compounds of the invention are effective as inhibitors of HCV, and are useful, alone and together with other therapeutic agents, in treating or preventing diseases or disorders such as viral infections and virus-related disorders.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
25 February 2010 (25.02.2010)

PCT

(10) International Publication Number
WO 2010/022125 A1

(51) International Patent Classification:

C07D 239/46 (2006.01) **C07D 409/06** (2006.01)
C07D 401/06 (2006.01) **C07D 413/06** (2006.01)
C07D 403/06 (2006.01) **C07D 417/06** (2006.01)
C07D 405/06 (2006.01) **A61K 31/505** (2006.01)
C07D 401/14 (2006.01) **A61K 31/506** (2006.01)

(21) International Application Number:

PCT/US2009/054268

(22) International Filing Date:

19 August 2009 (19.08.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/090,478 20 August 2008 (20.08.2008) US

(71) Applicants (for all designated States except US):

SCHERING CORPORATION [US/US]; 2000 Gallop-
 ing Hill Road, Kenilworth, New Jersey 07033-0530 (US).
SOUTHERN RESEARCH INSTITUTE [US/US];
 2000 9th Avenue South, P.O.Box 55305, Birmingham,
 Alabama 35205-5305 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **NJOROGE, F.,**
George [US/US]; 11 Softwood Way, Warren, New Jersey
 07059 (US). **ARASAPPAN, Ashok** [IN/US]; 18 Larsen

Court, Bridgewater, New Jersey 08807 (US). **BENNETT,**
Frank [GB/US]; 38 Keith Jeffries Avenue, Cranford,
 New Jersey 07016 (US). **PIWINSKI, John, J.** [US/US];
 6 Saddle Ridge Drive, Lebanon, New Jersey 08833 (US).
SHIH, Neng-Yang [US/US]; 2 Tyler Road, Lexington,
 Massachusetts 02420 (US). **KWONG, Cecil, D.**
 [US/US]; 1734 Woodbrine Drive, Birmingham, Alabama
 35216 (US). **ANANTHAN, Subramaniam** [US/US];
 1318 Atkins-Trim Boulevard, Birmingham, Alabama
 35226 (US). **CLARK, Jeremy** [US/US]; 349 S. Burbank
 Drive, Hoover, Alabama 35244 (US). **FOWLER, Anita,**
T. [US/US]; 5703 Kacie Circle, Irondale, Alabama 35210
 (US). **GENG, Feng** [CN/US]; 17 Snowdrift Drive, Pis-
 cataway, New Jersey 08854 (US). **KEZAR, III, Hollis, S.**
 [US/US]; 4566 Lavelley Drive, Hoover, Alabama 35244
 (US). **MADDRY, Joseph, A.** [US/US]; 565 Rutherford
 Circle, Birmingham, Alabama 35206 (US). **REYNOLDS,**
Robert, C. [US/US]; 4300 Fair Oaks Drive, Birmingham,
 Alabama 35213 (US). **ROYCHOWDHURY, Abhijit**
 [IN/US]; 3460 Manor Lane, # 105, Homewood, Alabama
 35209 (US). **SECRIST, III, John, A.** [US/US]; 5564
 Surrey Lane, Birmingham, Alabama 35242 (US).

(74) Agent: **MACMILLAN, Keith, D.**; Schering-plough Cor-
 poration, Patent Department, K-6-1 1990, 2000 Galloping
 Hill Road, Kenilworth, New Jersey 07033-0530 (US).

[Continued on next page]

(54) Title: ETHYNYL-SUBSTITUTED PYRIDINE AND PYRIMIDINE DERIVATIVES AND THEIR USE IN TREATING VIRAL INFECTIONS



(57) Abstract: The present invention provides com-
 pounds of Formula (I): (Chemical formula should be in-
 serted here as it appears on abstract in paper form) (I)
 and tautomers, isomers, and esters of said compounds,
 and pharmaceutically acceptable salts, solvates, and pro-
 drugs of said compounds, wherein wherein each of R,
 R¹, X, Y, Z, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁸, R¹⁹ and n
 is selected independently and as defined herein. Compo-
 sitions comprising such compounds are also provided.
 The compounds of the invention are effective as in-
 hibitors of HCV, and are useful, alone and together with
 other therapeutic agents, in treating or preventing dieas-
 es or disorders such as viral infections and virus-related
 disorders.

WO 2010/022125 A1



(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

-1-

**ETHYNYL-SUBSTITUTED PYRIDINE AND PYRIMIDINE DERIVATIVES AND
THEIR USE IN TREATING VIRAL INFECTIONS**

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to provisional application U.S. Serial No. 61/090,478,
5 filed August 20, 2008, incorporated by reference.

FIELD OF THE INVENTION

The present invention relates to certain ethynyl-substituted pyridine and pyrimidine
derivatives, to compositions comprising them, and to methods for their use as inhibitors of
HCV and in treating or preventing viral infections or virus-related disorders.

BACKGROUND OF THE INVENTION

HCV is a (+)-sense single-stranded RNA virus that has been implicated as the major
causative agent in non-A, non-B hepatitis (NANBH). NANBH is distinguished from other
types of viral-induced liver disease, such as hepatitis A virus (HAV), hepatitis B virus
15 (HBV), hepatitis delta virus (HDV), as well as from other forms of liver disease such as
alcoholism and primary biliary cirrhosis.

Hepatitis C virus is a member of the hepacivirus genus in the family Flaviviridae. It
is the major causative agent of non-A, non-B viral hepatitis and is the major cause of
transfusion-associated hepatitis and accounts for a significant proportion of hepatitis cases
20 worldwide. Although acute HCV infection is often asymptomatic, nearly 80% of cases
resolve to chronic hepatitis. About 60% of patients develop liver disease with various
clinical outcomes ranging from an asymptomatic carrier state to chronic active hepatitis and
liver cirrhosis (occurring in about 20% of patients), which is strongly associated with the
development of hepatocellular carcinoma (occurring in about 1-5% of patients). The World
25 Health Organization estimates that 170 million people are chronically infected with HCV,
with an estimated 4 million living in the United States.

HCV has been implicated in cirrhosis of the liver and in induction of hepatocellular
carcinoma. The prognosis for patients suffering from HCV infection remains poor as HCV
infection is more difficult to treat than other forms of hepatitis. Current data indicates a four-
30 year survival rate of below 50% for patients suffering from cirrhosis and a five-year survival
rate of below 30% for patients diagnosed with localized resectable hepatocellular carcinoma.

Patients diagnosed with localized unresectable hepatocellular carcinoma fare even worse, having a five-year survival rate of less than 1%.

HCV is an enveloped RNA virus containing a single-stranded positive-sense RNA genome approximately 9.5 kb in length. The RNA genome contains a 5'-nontranslated region (5' NTR) of 341 nucleotides, a large open reading frame (ORF) encoding a single polypeptide of 3,010 to 3,040 amino acids, and a 3'-nontranslated region (3'-NTR) of variable length of about 230 nucleotides. HCV is similar in amino acid sequence and genome organization to flaviviruses and pestiviruses, and therefore HCV has been classified as a third genus of the family Flaviviridae.

The 5' NTR, one of the most conserved regions of the viral genome, contains an internal ribosome entry site (IRES) which plays a pivotal role in the initiation of translation of the viral polyprotein. A single long open reading frame encodes a polyprotein, which is co- or post-translationally processed into structural (core, E1, E2 and p7) and nonstructural (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) viral proteins by either cellular or viral proteinases. The 3' NTR consists of three distinct regions: a variable region of about 38 nucleotides following the stop codon of the polyprotein, a polyuridine tract of variable length with interspersed substitutions of cytidines, and 98 nucleotides (nt) at the very 3' end which are highly conserved among various HCV isolates. By analogy to other plus-strand RNA viruses, the 3'-NTR is thought to play an important role in viral RNA synthesis. The order of the genes within the genome is: NH₂-C-E1-E2-p7-NS2-NS3-NS4A-NS4B-NS5A-NS5B-COOH.

Processing of the structural proteins core (C), envelope protein 1 and (E1, E2), and the p7 region is mediated by host signal peptidases. In contrast, maturation of the nonstructural (NS) region is accomplished by two viral enzymes. The HCV polyprotein is first cleaved by a host signal peptidase generating the structural proteins C/E1, E1/E2, E2/p7, and p7/NS2. The NS2-3 proteinase, which is a metalloprotease, then cleaves at the NS2/NS3 junction. The NS3/4A proteinase complex (NS3 being a serine protease and NS4A acting as a cofactor of the NS3 protease), is then responsible for processing all the remaining cleavage junctions. RNA helicase and NTPase activities have also been identified in the NS3 protein. One-third of the NS3 protein functions as a protease, and the remaining two-thirds of the molecule acts as the helicase/ATPase that is thought to be involved in HCV replication. NS5A may be phosphorylated and acts as a putative cofactor of NS5B. The fourth viral

enzyme, NS5B, is a membrane-associated RNA-dependent RNA polymerase (RdRp) and a key component responsible for replication of the viral RNA genome. NS5B contains the "GDD" sequence motif, which is highly conserved among all RdRps characterized to date.

Replication of HCV is thought to occur in membrane-associated replication

5 complexes. Within these, the genomic plus-strand RNA is transcribed into minus-strand RNA, which in turn can be used as a template for synthesis of progeny genomic plus-strands. At least two viral enzymes appear to be involved in this reaction: the NS3 helicase/NTPase, and the NS5B RNA-dependent RNA polymerase. While the role of NS3 in RNA replication is less clear, NS5B is the key enzyme responsible for synthesis of progeny RNA strands.

10 Using recombinant baculoviruses to express NS5B in insect cells and a synthetic nonviral RNA as a substrate, two enzymatic activities have been identified as being associated with it: a primer-dependent RdRp and a terminal transferase (TNTase) activity. It was subsequently confirmed and further characterized through the use of the HCV RNA genome as a substrate. Other studies have shown that NS5B with a C-terminal 21 amino-acid truncation expressed in

15 Escherichia coli is also active for *in vitro* RNA synthesis. On certain RNA templates, NS5B has been shown to catalyze RNA synthesis via a *de novo* initiation mechanism, which has been postulated to be the mode of viral replication *in vivo*. Templates with single-stranded 3' termini, especially those containing a 3'-terminal cytidylate moiety, have been found to direct *de novo* synthesis efficiently. There has also been evidence for NS5B to utilize di- or tri-

20 nucleotides as short primers to initiate replication.

It is well-established that persistent infection of HCV is related to chronic hepatitis, and as such, inhibition of HCV replication is a viable strategy for the prevention of hepatocellular carcinoma. Present treatment approaches for HCV infection suffer from poor efficacy and unfavorable side-effects and there is currently a strong effort directed to the

25 discovery of HCV replication inhibitors that are useful for the treatment and prevention of HCV related disorders. New approaches currently under investigation include the development of prophylactic and therapeutic vaccines, the identification of interferons with improved pharmacokinetic characteristics, and the discovery of agents designed to inhibit the function of three major viral proteins: protease, helicase and polymerase. In addition, the

30 HCV RNA genome itself, particularly the IRES element, is being actively exploited as an antiviral target using antisense molecules and catalytic ribozymes.

Particular therapies for HCV infection include α -interferon monotherapy and combination therapy comprising α -interferon and ribavirin. These therapies have been shown to be effective in some patients with chronic HCV infection. The use of antisense oligonucleotides for treatment of HCV infection has also been proposed as has the use of free bile acids, such as ursodeoxycholic acid and chenodeoxycholic acid, and conjugated bile acids, such as tauroursodeoxycholic acid. Phosphonoformic acid esters have also been proposed as potentially for the treatment of various viral infections including HCV. Vaccine development, however, has been hampered by the high degree of viral strain heterogeneity and immune evasion and the lack of protection against reinfection, even with the same inoculum.

The development of small-molecule inhibitors directed against specific viral targets has become a major focus of anti-HCV research. The determination of crystal structures for NS3 protease, NS3 RNA helicase, and NS5B polymerase has provided important structural insights that should assist in the rational design of specific inhibitors.


NS5B, the RNA-dependent RNA polymerase, is an important and attractive target for small-molecule inhibitors. Studies with pestiviruses have shown that the small molecule compound VP32947 (3-[[[(2-dipropylamino)ethyl]thio]-5H-1,2,4-triazino[5,6-b]indole) is a potent inhibitor of pestivirus replication and most likely inhibits the NS5B enzyme since resistant strains are mutated in this gene. Inhibition of RdRp activity by (-) β -L-2',3'-dideoxy-3'-thiacytidine 5'-triphosphate (3TC; lamivudine triphosphate) and phosphonoacetic acid also has been observed.

Despite the intensive effort directed at the treatment and prevention of HCV and related viral infections, there exists a need in the art for non-peptide, small-molecule compounds having desirable or improved physicochemical properties that are useful for inhibiting viruses and treating viral infections and virus-related disorders.

SUMMARY OF THE INVENTION

The present invention provides certain ethynyl-substituted pyridine and pyrimidine derivatives (collectively referred to herein as “compounds of the invention”), compositions comprising such compounds, and methods for their use as HCV inhibitors and for treating viral infections and disorders related thereto.

In one embodiment, the compounds of the invention have a general structure shown in Formula (I):



(I)

and include tautomers, isomers, and esters of said compounds, and pharmaceutically acceptable salts, solvates, and prodrugs of said compounds, tautomers, isomers, and esters, wherein each of R, R¹, X, Y, Z, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁸, R¹⁹ and n are selected independently and wherein:

R is selected from H, alkyl, aryl, heteroaryl, cycloalkyl, aryl-fused cycloalkyl, heteroaryl-fused cycloalkyl, cycloalkenyl, aryl-fused cycloalkenyl, heteroaryl-fused cycloalkenyl, heterocycloalkyl, aryl-fused heterocycloalkyl, and heteroaryl-fused heterocycloalkyl,

wherein each of said alkyl, said aryl, said heteroaryl, said cycloalkyl, said aryl-fused cycloalkyl, said heteroaryl-fused cycloalkyl, said cycloalkenyl, said aryl-fused cycloalkenyl, said heteroaryl-fused cycloalkenyl, said heterocycloalkyl, said aryl-fused heterocycloalkyl, and said heteroaryl-fused heterocycloalkyl, is unsubstituted or optionally independently substituted with from one to five substituents, which are the same or different, each substituent being independently selected from halo, -OH, -CN, oxo, alkyl, cycloalkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heterohaloalkyl,

-alkyl-OH, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, heteroarylalkyl-, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, heterocycloalkylalkyl-, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)-cycloalkyl, -C(O)-heterocycloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -C(O)O-haloalkyl, -C(O)O-cycloalkyl, -C(O)O-heterocycloalkyl, -OC(O)-alkyl, -OC(O)-haloalkyl, -OC(O)-cycloalkyl, -OC(O)-heterocycloalkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -OC(O)NH₂, -CO(O)NHR¹⁰, -CO(O)NR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NH₂, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -S(O)₂R¹⁰, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to five substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl;

X and Y are each independently selected from N and CH, with the proviso that at least one of X or Y is N;

Z = H, halo, -OH, -SH, -CN, alkyl, alkenyl, alkynyl, heteroalkyl, haloalkyl, heterohaloalkyl, -S-alkyl, -O-alkyl, -O-aryl, -O-heteroaryl, cycloalkyl, aryl, heteroaryl, -NH₂, -NHR¹², and -NR¹²R¹³;

R¹ is selected from H, halo, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, heteroaryl, -OH, -O-alkyl, -O-aryl, -O-heteroalkyl, -O-heteroaryl, -SH, -S-alkyl, -S-aryl, -S-heteroalkyl, -S-heteroaryl, -NH₂, -NHR¹⁴, -NR¹⁴R¹⁵, -NO₂, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰;

R² (when R² is not joined with R⁹) is selected from H and alkyl;

n = 0, 1, or 2;

R³ is selected from H, -alkyl, -alkenyl, alkynyl, aryl, heteroaryl, and cycloalkyl,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, and said cycloalkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^4 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^5 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl,

-OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R^4 and R^5 are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

R^6 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R^5 and R^6 are taken together to form a double bond;

R^7 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl,

-OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R⁶ and R⁷ are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

R⁸ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R¹⁰)R¹¹, -O(C)O-NHR¹¹, -O(C)O-NH₂, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰, -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R⁹ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl,

10

-O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R¹⁰)R¹¹, -O(C)O-NHR¹¹, -O(C)O-NH₂,
 -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl,
 -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰,
 -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰,
 5 -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and
 heterocycloalkenyl;

or, alternatively, R⁸ and R⁹ are taken together with the carbon atom to which they are
 shown attached to form a 3- to 7-membered, saturated or partially unsaturated,
 spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N,
 10 and S;

each R¹⁸ (when present) is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl,
 heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -
 NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰,
 -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

15 wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said
 -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or
 optionally independently substituted with from one to three substituents, which can be
 the same or different, each substituent being independently selected from halo, -OH,
 alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl,
 20 -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl,
 -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl,
 -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and
 heterocycloalkenyl;

each R¹⁹ (when present) is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl,
 25 heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -
 NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰,
 -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,


wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said
 -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or
 30 optionally independently substituted with from one to three substituents, which can be
 the same or different, each substituent being independently selected from halo, -OH,

11

alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, n is 1 and R^{18} and R^{19} are taken together with the carbon atom to which they are attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1C):




(1C)

wherein R^{20} and R^{21} are each independently selected from H, alkyl, and heteroalkyl and

wherein R^5 and R^6 are defined above, with the proviso that when R^4 and R^7 form a moiety (1C), then R^5 and R^6 are not taken together to form a double bond;

or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1D):




(1D),

wherein R^5 and R^6 are as defined above;

12

or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1E):



(1E),

5 wherein R^5 and R^6 are as defined above;

each R^{10} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

10 each R^{11} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{10} and R^{11} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl;

15 each R^{12} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

each R^{13} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

20 or, alternatively, R^{12} and R^{13} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R^{14} is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, alkylamino, alkylthio, heteroalkenyl, haloalkenyl, $-S(O)_2$ -alkyl, -alkyl-OH,

-alkyl-O-Acyl, -C(O)Oalkyl, -C(O)alkyl, cycloalkyl, cycloalkyl-alkyl-, heterocycloalkyl, heterocycloalkyl-alkyl-, heterocycloalkenyl, heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-, heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said alkylthio, each said heteroalkenyl, each said haloalkenyl, each said -S(O)₂-alkyl, each said -alkyl-OH, each said -alkyl-O-Acyl, each said -C(O)Oalkyl, each said -C(O)alkyl, each said cycloalkyl, each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said

heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl, and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted with from one to five substituent, which can be the same or different, each substituent being independently selected from -OH, halo, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)₂alkyl, -S(O)₂aryl, alkyl, alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, heterohaloalkyl, aryl, cycloalkyl, and heterocycloalkyl;

each R¹⁵ is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, alkylamino, alkylthio, heteroalkenyl, haloalkenyl, -S(O)₂-alkyl, -alkyl-OH, -alkyl-O-Acyl, -C(O)Oalkyl, -C(O)alkyl, cycloalkyl, cycloalkyl-alkyl-, heterocycloalkyl, heterocycloalkyl-alkyl-, heterocycloalkenyl, heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-, heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said alkylthio, each said heteroalkenyl, each said haloalkenyl, each said -S(O)₂-alkyl, each said -alkyl-OH, each said -alkyl-O-Acyl, each said -C(O)Oalkyl, each said -C(O)alkyl, each said cycloalkyl, each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl, and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted with from one to five substituent, which can be the same or different, each substituent being independently selected from -OH, halo, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH,

-C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)₂alkyl, -S(O)₂aryl, alkyl, alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, heterohaloalkyl, aryl, cycloalkyl, and heterocycloalkyl;

or, alternatively, R¹⁴ and R¹⁵ are linked together with the nitrogen to which they are
5 attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R¹⁶ is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl, -C(O)N(alkyl)₂, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl; and

each R¹⁷ is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl,
10 heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl, -C(O)N(alkyl)₂, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R¹⁶ and R¹⁷ are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl.

In another embodiment, the invention provides compositions, including
15 pharmaceutical compositions, comprising one or more compounds of the invention (e.g., one compound of the invention), or a pharmaceutically acceptable salt, solvate, ester, or prodrug thereof, and a pharmaceutically acceptable carrier or diluent. In one embodiment, said compound or compounds of the invention are present in the composition in an amount effective for inhibiting HCV, and/or for treating or preventing a viral infection or a virus-
20 related disorder in a patient in need thereof.

In another embodiment, the invention provides a pharmaceutical composition comprising one or more compounds of the invention, or a pharmaceutically acceptable salt, solvate, ester, or prodrug thereof, together with one or more additional therapeutic agents, optionally further comprising a pharmaceutically effective carrier or diluent. Non-limiting
25 examples of such additional therapeutic agents include one or more of any of the following: HCV polymerase inhibitors, HCV protease inhibitors, HCV replicase inhibitors, nucleosides, Interferon, and/or ribavirin (or Levovirin or Viramidine). Non-limiting examples of interferon include PEG-interferon, PEG interferon alpha conjugate, alpha-interferon, and pegylated interferon. These and other examples are known to those of ordinary skill in the
30 art.

In another embodiment, the present invention provides for the use of one or more compounds of the invention, or a pharmaceutically acceptable salt, solvate, ester, and/or prodrug thereof, alone or in combination with one or more additional therapeutic agents such as those described above, for inhibiting HCV and/or for treating or preventing a viral infection or a virus-related disorder in a patient in need thereof.

In another embodiment, the invention provides a method of inhibiting HCV *in vivo*, *ex vivo*, or *in vitro*, comprising exposing a population of cells comprising HCV to an effective amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester, or prodrug thereof, alone or in combination with one or more additional therapeutic agents such as those described above. In one such embodiment, the compound or compounds of the invention are used as the neat chemical. In another such embodiment, the compounds of the invention are used in the form of a pharmaceutically acceptable composition.

In another embodiment, the invention provides a method for treating or preventing a viral infection or a virus-related disorder in a patient, comprising administering to the patient an effective amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester, or prodrug thereof, alone or in combination with one or more additional therapeutic agents such as those described above. In one such embodiment, the compound or compounds of the invention are used as the neat chemical. In another such embodiment, the compounds of the invention are used in the form of a pharmaceutically acceptable composition.

The details of the invention are set forth in the accompanying detailed description below. Although any methods and materials similar to those described herein can be used in the practice or testing of the present invention, illustrative methods and materials are described herein. Other features, objects, and advantages of the invention will be apparent from the description and the claims. All patents and publications cited in this specification are incorporated herein by reference.

DETAILED DESCRIPTION OF THE INVENTION

In one embodiment, the compounds of the invention have the structural Formula (I) as described above, and include pharmaceutically acceptable salts, esters, prodrugs, tautomers, and isomers of said compounds.

In one embodiment, in Formula (I), each of R³, R⁵, R⁶, and R⁸ is H.

In one embodiment, in Formula (I), n is 1; each of R^2 , R^3 , R^5 , R^6 , R^8 , R^{18} and R^{19} is H; R^4 and R^7 are OH; and R^9 is alkyl, wherein said alkyl is unsubstituted or substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl.

In one embodiment, in Formula (I), n is 1; each of R^2 , R^3 , R^5 , R^6 , R^8 , R^{18} and R^{19} is H; R^4 and R^7 are OH; and R^9 is alkyl, wherein said alkyl is unsubstituted or substituted with from one to five groups independently selected from -OH, halo, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -Ocycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

In one embodiment, in Formula (I), n is 1; each R^2 , R^3 , R^5 , R^6 , R^8 , R^{18} and R^{19} is H; R^4 and R^7 are OH; and R^9 is methyl, wherein said methyl is unsubstituted or substituted with from one to three groups independently selected from -OH, halo, alkyl, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -Ocycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

In some embodiments, R^9 is -alkyl-NHS(O)₂R¹⁰, wherein R^{10} is selected from methyl, ethyl, and cyclopropyl.

In some embodiments, R^9 is selected from -alkyl -N(CH₃)S(O)₂R¹⁰ and -alkyl-N(CH₂CH₃)S(O)₂R¹⁰, wherein R^{10} is selected from methyl, ethyl, and cyclopropyl.

In some embodiments, R^9 is -alkyl-O(C)O-NHR¹⁰, wherein R^{10} is selected from methyl, ethyl, and cyclopropyl.

In some embodiments, R^9 is selected from R^9 -alkyl-O(C)O-N(CH₃)R¹⁰ and -O(C)O-N(CH₂CH₃)R¹⁰, wherein R^{10} is selected from methyl, ethyl, and cyclopropyl.

17

In one embodiment, in Formula (I), n is 1; each of R^2 , R^3 , R^5 , R^6 , R^8 , R^{18} and R^{19} is H; R^4 and R^7 are OH; and R^9 is selected from $-\text{CH}_2\text{-O-alkyl}$, $-\text{CH}_2\text{-OH}$, $-\text{CH}_3$, H, $-\text{CH}_2\text{-CH}_3$, $-\text{CH}_2\text{-OC(O)CF}_3$, $-\text{CH}_2\text{-NH}_2$, $-\text{CH}_2\text{-NHR}^{16}$, and $-\text{CH}_2\text{-NR}^{16}\text{R}^{17}$.

In one embodiment, in Formula (I), each of R^3 , R^5 , R^6 , and R^8 is H and each of R^4 and R^7 is -OH.


In one embodiment, in Formula (I), each of R^3 , R^5 , R^6 , and R^8 is H; each of R^4 and R^7 is -OH; and R^9 is -O-alkyl.

In one embodiment, in Formula (I), each of R^3 , R^5 , R^6 , and R^8 is H; each of R^4 and R^7 is -OH; and R^9 is -O-CH₃.

In one embodiment, in Formula (I), each of R^3 , R^5 , R^6 , and R^8 is H; each of R^4 and R^7 is -OH; R^9 is -O-CH₃, and n is 1.

In one embodiment, in Formula (I), each of R^3 , R^5 , R^6 , and R^8 is H; each of R^4 and R^7 is -OH; R^9 is -O-CH₃, and n is 2.

In one embodiment, the compounds of the invention have the structural Formula (I.A):



(I.A)

and includes tautomers, isomers, and esters of such compounds, and pharmaceutically acceptable salts, solvates, and prodrugs of said compounds, tautomers, isomers, and esters, wherein each of R , R^1 , X , Y , Z , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{18} , R^{19} and n are selected independently and wherein:

R , R^1 , R^2 , X , Y , Z , and n are as defined in Formula (I);

R³ is selected from H, -alkyl, -alkenyl, alkynyl, aryl, heteroaryl, and cycloalkyl,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, and said cycloalkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

10 R⁴ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

20 R⁵ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

30 R⁶ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH,
 5 alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^7 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH,
 15 alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R^6 and R^7 are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

R^8 is selected from is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and
 25 -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH,
 30 alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R^{10}) R^{11} , -O(C)O-NHR¹¹, -O(C)O-NH₂,

-OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰, -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R⁹ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R¹⁰)R¹¹, -O(C)O-NHR¹¹, -O(C)O-NH₂, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰, -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

each R¹⁸ (when present) is independently selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

each R^{19} (when present) is independently selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NHalkyl, and -N(alkyl)₂,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said
 5 -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl,
 10 -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

each R^{10} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

15 each R^{11} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl, -C(O)(alkyl)₂, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{10} and R^{11} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl;

20 each R^{12} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl, -C(O)(alkyl)₂, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

each R^{13} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, -S(O)₂-alkyl, -alkyl-OH, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl,
 25 -C(O)(alkyl)₂, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{12} and R^{13} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R^{14} is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, alkylamino, heteroalkenyl, haloalkenyl, -S(O)₂-alkyl, -alkyl-OH,
 30 -alkyl-O-Acyl, -C(O)Oalkyl, -C(O)alkyl, -C(O)NHalkyl, -C(O)(alkyl)₂, cycloalkyl,

cycloalkyl-alkyl-, heterocycloalkyl, heterocycloalkyl-alkyl-, heterocycloalkenyl,
heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-, heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each
said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said

5 heteroalkenyl, each said haloalkenyl, each said -S(O)₂-alkyl, each said -alkyl-OH, each
said -alkyl-O-Acyl, each said -C(O)Oalkyl, each said -C(O)alkyl, each said cycloalkyl,
each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said
heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said
heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl,
10 and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted
with from one to three substituent, which can be the same or different, each substituent
being independently selected from halo, -OH, -NH₂, -NHalkyl, -N(alkyl)₂, alkyl,
alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, and heterohaloalkyl;

each R¹⁵ is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl,
15 heterohaloalkyl, alkylamino, heteroalkenyl, haloalkenyl, -S(O)₂-alkyl, -alkyl-OH,
-alkyl-O-Acyl, -C(O)Oalkyl, -C(O)alkyl, cycloalkyl, cycloalkyl-alkyl-, heterocycloalkyl,
heterocycloalkyl-alkyl-, heterocycloalkenyl, heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-,
heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each
20 said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said
heteroalkenyl, each said haloalkenyl, each said -S(O)₂-alkyl, each said -alkyl-OH, each
said -alkyl-O-Acyl, each said -C(O)Oalkyl, each said -C(O)alkyl, each said cycloalkyl,
each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said
heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said
25 heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl,
and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted
with from one to three substituent, which can be the same or different, each substituent
being independently selected from halo, -OH, -NH₂, -NHalkyl, -N(alkyl)₂, alkyl,
alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, and heterohaloalkyl;


30 or, alternatively, R¹⁴ and R¹⁵ are linked together with the nitrogen to which they are
attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R^{16} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl; and

each R^{17} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{16} and R^{17} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl.

In one embodiment, the compounds of the invention have the structural Formula (I.a):



(I.a)

and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R , R^1 , X , Y , Z , R^2 , R^4 , R^5 , R^7 , and R^9 is selected independently and defined in Formula (I).

In one embodiment, in Formula (I.a), n is 1; R^2 is H; R^4 and R^7 are each independently selected from H and OH; R^5 is selected from H, halo, and alkyl; and R^9 is selected from H, $-COOH$, $-C(O)O$ -alkyl, $-OC(O)$ -alkyl, $-C(O)O$ -aryl, $-OC(O)$ -aryl, $-C(O)O$ -alkyl-aryl, $-OC(O)$ -alkyl-aryl, $-C(O)O$ -alkyl-heteroaryl, $-OC(O)$ -alkyl-heteroaryl, alkyl, $-O$ -alkyl, heteroalkyl, haloalkyl, heterohaloalkyl, $-O$ -heteroalkyl, $-O$ -haloalkyl, $-O$ -heterohaloalkyl, $-alkyl-OH$, $-alkyl-OC(O)$ -alkyl, $-alkyl-OC(O)$ -haloalkyl, $-alkyl-NH_2$, $-alkyl-NHR^{16}$, and $-alkyl-NR^{16}R^{17}$.

In one embodiment, in Formula (I.a), n is 1; each of R^2 , R^3 , R^5 , R^6 , R^8 , R^{18} and R^{19} is H; R^4 and R^7 are OH; and R^9 is alkyl, wherein said alkyl is unsubstituted or substituted with

from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -O(C)O-NHR¹⁰,
 5 -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl.

In one embodiment, in Formula (I.a), n is 1; each of R², R³, R⁵, R⁶, R⁸, R¹⁸ and R¹⁹ is H;
 10 R⁴ and R⁷ are OH; and R⁹ is alkyl, wherein said alkyl is unsubstituted or substituted with from one to five groups independently selected from -OH, halo, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -O-cycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

In one embodiment, in Formula (I.a), n is 1; each R², R³, R⁵, R⁶, R⁸, R¹⁸ and R¹⁹ is H;
 15 R⁴ and R⁷ are OH; and R⁹ is methyl, wherein said methyl is unsubstituted or substituted with from one to three groups independently selected from -OH, halo, alkyl, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -O-cycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰,
 20 -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

In some embodiments, R⁹ is -alkyl-NHS(O)₂R¹⁰, wherein R¹⁰ is selected from methyl, ethyl, and cyclopropyl.

In some embodiments, R⁹ is selected from -alkyl -N(CH₃)S(O)₂R¹⁰ and -alkyl-N(CH₂CH₃)S(O)₂R¹⁰, wherein R¹⁰ is selected from methyl, ethyl, and cyclopropyl.

25 In some embodiments, R⁹ is -alkyl-O(C)O-NHR¹⁰, wherein R¹⁰ is selected from methyl, ethyl, and cyclopropyl.

In some embodiments, R⁹ is selected from R⁹ -alkyl-O(C)O-N(CH₃)R¹⁰ and -O(C)O-N(CH₂CH₃)R¹⁰, wherein R¹⁰ is selected from methyl, ethyl, and cyclopropyl.

30 In one embodiment, in Formula (I.a), n is 1; R² is H; R⁴ and R⁷ are each independently selected from H and OH; R⁵ is selected from H, halo, and alkyl; and R⁹ is selected from H,

-CH₃, -CH₂-CH₃, -CH₂-OH, -CH₂-O-alkyl, -CH₂-OC(O)-alkyl, -CH₂-OC(O)-haloalkyl, -CH₂-NH₂, -CH₂-NHR¹⁶, and -CH₂-NR¹⁶R¹⁷.

In one embodiment, in Formula (I.a), n is 1, R² is H, R⁵ is -CH₃, and R⁹ is selected from H, -CH₃, -CH₂-CH₃, -CH₂-OH, -CH₂-O-alkyl, -CH₂-OC(O)CF₃, -CH₂-NH₂,
5 -CH₂-NHR¹⁶, and -CH₂-NR¹⁶R¹⁷.

In one embodiment, in Formula (I.a), n is 1; R² is H; R⁴ and R⁷ are each -OH, R⁵ is -CH₃, and R⁹ is H.


In one embodiment, in Formula (I.a), n is 1; R² is H; R⁴ and R⁷ are each -OH, R⁵ is selected from H and -CH₃, and R⁹ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl,
10 azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or
15 optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl,
20 -C(O)O-haloalkenyl, -S(O)₂alkyl, -S-alkyl, -CN, -NH₂, -NHR¹⁶, and -N(alkyl)₂, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl.

In one embodiment, in Formula (I.a), X is N, Y is N, n is 1; R² is H; R⁴ and R⁷ are each -OH, R⁵ is selected from H and -CH₃, and R⁹ is selected from H, -alkyl, -alkyl-OH,
25 -alkyl-S(O)₂alkyl, -alkyl-S-alkyl, haloalkyl, heteroalkyl, -alkyl-CN, -alkyl-NH₂, -alkyl-NHR¹⁶, and -alkyl-N(alkyl)₂. In one such embodiment, each said alkyl is selected from straight or branched lower alkyl.


26

In one embodiment, the compounds of the invention have the structural Formula (I.a.1):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural
 Formula (I.a.1.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in
 Formula (I).


15

In one embodiment, the compounds of the invention have the structural Formula (I.a.2):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural
 Formula (I.a.2.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).

28


In one embodiment, the compounds of the invention have the structural Formula (I.a.3):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently and defined
 in Formula (I).

In one embodiment, the compounds of the invention have the structural
 Formula (I.a.3.i)


10 :



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in
 15 Formula (I).


29

In one embodiment, the compounds of the invention have the structural Formula (I.a.4):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R , R^1 , X , Y , Z , R^2 , and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural
 Formula (I.a.4.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R , R^1 , X , Y , Z , and R^2 is selected independently and defined in Formula (I).


30

In one embodiment, the compounds of the invention have the structural Formula (I.a.5):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural
 Formula (I.a.5.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).


31

In one embodiment, the compounds of the invention have the structural Formula (I.a.6):




and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently defined in
 Formula (I).

In one embodiment, the compounds of the invention have the structural
 Formula (I.a.6.i):




and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in
 Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.a.7):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural
 Formula (I.a.7.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).

33


In one embodiment, the compounds of the invention have the structural Formula (I.a.8):



(I.a.8)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently defined in
 Formula (I).

In one embodiment, the compounds of the invention have the structural
 Formula (I.a.8.i):




(I.a.8.i)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in
 Formula (I).


34

In one embodiment, the compounds of the invention have the structural Formula (I.a.9):




and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², n, and R¹⁶ is selected independently defined
 in Formula (I).

In one embodiment, the compounds of the invention have the structural
 Formula (I.a.9.i):




and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, R², and R¹⁶ is selected independently and
 defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.a.10):




- 5 and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², n, R¹⁶, and R¹⁷ is selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.a.10.i):




- and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², R¹⁶, and R¹⁷ is selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.a.10.j):



(I.a.10.j)

- 5 and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and is selected independently and defined in Formula (I), and wherein CB is a moiety selected from the group consisting of:




wherein each R¹⁰ is independently selected from the group consisting of methyl, ethyl, and cyclopropyl.

In one embodiment, in Formula (I.a.10.j):

X is N; Y is N; R² is H; and Z is selected from the group consisting of H, methyl, and chloro; and R and R¹ are each as defined in Formula (I). In other such embodiments, R and R¹ are each independently as defined in any of the various embodiments described herein, or in the examples.

5 In one embodiment, the compounds of the invention have the structural Formula (I.B):




(I.B)

and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², R³, R⁴, R⁷, R⁸, R⁹, R¹⁸, R¹⁹, and n is selected independently and defined in Formula (I).

10

In one embodiment, the compounds of the invention have the structural Formula (I.b):



(I.b)


and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², and R⁹ is selected independently and defined in Formula (I).

15

In one embodiment, in Formula (I.b), R^2 is H, and R^9 is selected from H, $-CH_3$, $-CH_2-O$ -alkyl, $-CH_2-OH$, $-CH_2-OC(O)$ -alkyl, $-CH_2-OC(O)$ -haloalkyl, $-CH_2-CH_3'-CH_2-NH_2$, $-CH_2-NHR^{16}$, and $-CH_2-NR^{16}R^{17}$.


- 5 In one embodiment, in Formula (I.b), R^2 is H, and R^9 is selected from $-CH_2-O$ -alkyl, and $-CH_2-OH$.

In one embodiment, the compounds of the invention have the structural Formula (I.b.1):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R^1 , X, Y, Z, and R^2 is selected independently and defined in Formula (I).


- 15 In one embodiment, the compounds of the invention have the structural Formula (I.b.1.i):



39


and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).

5 In one embodiment, the compounds of the invention have the structural Formula (I.b.2):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
10 compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the Formula (I.b.2.i):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
15 compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).

40


In one embodiment, the compounds of the invention have the structural Formula (I.C):



(I.C)

and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², R³, R⁵, R⁶, R⁸, R⁹, R¹⁸, R¹⁹, R²⁰, R²¹, and n is selected independently and defined in Formula (I), with the proviso that R⁵ and R⁶ are not taken together to form a double bond.

In one embodiment, the compounds of the invention have the structural Formula (I.c):



(I.c)


and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 10 compounds, wherein each of R, R¹, X, Y, Z, R², R⁹, R²⁰, and R²¹ is selected independently and defined in Formula (I).

41

In one embodiment, in Formula (I.c), R^2 is H; R^9 is selected from H, $-CH_3$, $-CH_2-O$ -alkyl, $-CH_2-OH$, $-CH_2-OC(O)$ -alkyl, $-CH_2-OC(O)$ -haloalkyl, $-CH_2-CH_3$, $-CH_2-NH_2$, $-CH_2-NHR^{16}$, and $-CH_2-NR^{16}R^{17}$; and each of R^{20} and R^{21} is independently selected from H and $-CH_3$.


5

In one embodiment, the compounds of the invention have the structural Formula (I.c.1):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R^1 , X, Y, Z, and R^2 is selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.c.1.i):




15

42

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural Formula (I.c.2):



(I.c.2)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural Formula (I.c.2.i):



(I.c.2.i)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently defined in Formula (I).


5 In one embodiment, the compounds of the invention have the structural Formula (I.D):



(I.D)

and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², R³, R⁵, R⁶, R⁸, R⁹, R¹⁸, and R¹⁹, and n is
 10 selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (I.d):



(I.d)


and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said
 5 compounds, wherein each of R, R¹, X, Y, Z, R², R⁹, and n is selected independently and defined in Formula (I).

In one embodiment, in Formula (I.d), n is 1 and R⁹ is selected from H, -CH₃,
 -CH₂-O-alkyl, -CH₂-OH, -CH₂-OC(O)-alkyl, -CH₂-OC(O)-haloalkyl, -CH₂-CH₃, -CH₂-NH₂,
 -CH₂-NHR¹⁶, and -CH₂-NR¹⁶R¹⁷.

10 In one embodiment, in Formula (I.d), n is 1 and R⁹ is selected from -CH₂-O-alkyl, and
 -CH₂-OH.

45

In one embodiment, the compounds of the invention have the structural Formula (I.d.1):




5

(I.d.1)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural

10 Formula (I.d.1.i):




(I.d.1.i)

and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, and R² is selected independently and defined in

15 Formula (I).


In one embodiment, the compounds of the invention have the structural Formula (I.E):



(I.E)

- 5 and includes pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R², R³, R⁵, R⁶, R⁸, R⁹, R¹⁸, and R¹⁹, and n is selected independently and defined in Formula (I).


In one embodiment, the compounds of the invention have the structural Formula (II):



(II)


and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, Z, R³, R⁴, R⁵, R⁶, R⁷, and R⁸ is selected independently and defined in Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (II):



and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, and Z is selected independently and wherein R, R¹, X, Y, and Z are defined in Formula (I).

10 In one embodiment, in Formula (I), the compounds of the invention have the structural Formula (II.a.1):



15 and include pharmaceutically acceptable salts, esters, prodrugs, or isomers of said compounds, wherein each of R, R¹, X, Y, and Z is selected independently and wherein R, R¹, X, Y, and Z are defined in Formula (I).

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 5 (II.A.1), X is N and Y is N.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 10 (II.A.1), X is N and Y is CH.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 15 (II.A.1), X is CH and Y is N.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 20 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is selected from aryl, heteroaryl, benzo-fused heteroaryl, cycloalkyl, cycloalkenyl, benzo-fused cycloalkyl, benzo-fused cycloalkenyl, heterocycloalkyl, and benzo-fused heterocycloalkyl,

wherein each of said alkyl, said aryl, said heteroaryl, said benzo-fused heteroaryl, said
 25 cycloalkyl, said cycloalkenyl, said heterocycloalkyl, said heterocycloalkenyl, and said benzo-fused heterocycloalkyl is unsubstituted or optionally independently substituted with from one to three substituents, which are the same or different, each substituent being independently selected from halo, -OH, -CN, alkyl, cycloalkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heterohaloalkyl, -alkyl-OH, -O-alkyl, -O-haloalkyl,
 30 -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl,

-C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains
 5 from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted benzo-fused heteroaryl, each of said substituents being independently selected from the group consisting
 15 of alkyl and -O-alkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 20 (II.A.1), R is selected from substituted alkynyl, substituted alkynyl, unsubstituted aryl, substituted aryl, unsubstituted cycloalkyl, substituted cycloalkyl, unsubstituted benzo-fused cycloalkyl, substituted benzo-fused cycloalkyl, unsubstituted cycloalkenyl, and substituted cycloalkenyl, which substituents, when present, are as defined in Formula (I).

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted aryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 30

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted aryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted benzo-fused cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 15 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted benzo-fused cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 20 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 25 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted cycloalkenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 30 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted cycloalkenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 5 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), R is selected from unsubstituted heteroaryl, substituted heteroaryl, unsubstituted
 benzo-fused heteroaryl, substituted benzo-fused heteroaryl, unsubstituted heterocycloalkyl,
 10 substituted heterocycloalkyl, unsubstituted benzo-fused heterocycloalkyl, substituted benzo-
 fused heterocycloalkyl, unsubstituted heterocycloalkenyl, and unsubstituted
 heterocycloalkenyl, which substituents, when present, are as defined in Formula (I).

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 15 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), R is unsubstituted heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 20 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), R is substituted heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 25 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), R is unsubstituted benzo-fused heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 30 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted benzo-fused heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted heterocycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted heterocycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 15 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted benzo-fused heterocycloalkyl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 20 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted benzo-fused heterocycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 25 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is unsubstituted heterocycloalkenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 30 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is substituted heterocycloalkenyl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R is an unsubstituted or substituted monocyclic aryl moiety or an unsubstituted or substituted heteroaryl moiety. Non-limiting examples of such unsubstituted or substituted monocyclic aryl moiety or unsubstituted or substituted heteroaryl moiety include:



wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl,

-C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰,
 -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹,
 -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and
 said substituted heteroaryl independently contains from one to three substituents, which may
 5 be the same or different, each substituent being independently selected from halo, alkyl,
 -O-alkyl, and -C(O)Oalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 10 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), R is an unsubstituted or an substituted bicyclic heteroaryl moiety. Non-limiting
 examples of such unsubstituted or substituted bicyclic heteroaryl moieties include:



wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, cycloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is selected from the group consisting of halo, alkyl, haloalkyl, cycloalkyl, and -NH₂. Non-limiting examples of Z when Z is cycloalkyl include cyclopropyl. Non-limiting examples of Z when Z is haloalkyl include fluoroalkyl (up to perfluoroalkyl).

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is selected from the group consisting of halo, alkyl, and cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is H.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), Z is halo.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), Z is -Cl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), Z is -F.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), Z is -OH.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
25 (II.A.1), Z is -SH.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
30 (II.A.1), Z is -Salkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), Z is $-S-CH_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), Z is $-alkyl$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), Z is $-CH_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), Z is $-CH_2CH_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
25 (II.A.1), Z is $-Oalkyl$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
30 (II.A.1), Z is $-OCH_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), Z is -haloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), Z is -CF₃.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), Z is -CHF₂.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), Z is -CH₂F.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
25 (II.A.1), Z is cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
30 (II.A.1), Z is cyclopropyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), Z is aryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), Z is phenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), Z is heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), Z is 2-thiophenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
25 (II.A.1), Z is 3-thiophenyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
30 (II.A.1), Z is 2-thiazolyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), Z is 2-oxazolyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), Z is 2-pyrimidinyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), Z is 2-pyridyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), Z is 2-pyrazinyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
25 (II.A.1), Z is 2-imidazolyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
30 (II.A.1), Z is $-\text{NH}_2$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 5 (II.A.1), Z is -NHR^{12} .

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 10 (II.A.1), Z is $\text{-NR}^{12}\text{R}^{13}$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 15 (II.A.1), Z is selected from the group consisting of Cl and methyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 20 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is selected from the group consisting of -NH_2 , -NHR^{14} , and $\text{-NR}^{14}\text{R}^{15}$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 25 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is selected from the group consisting of -NH_2 and -NHR^{14} .

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 30 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is H.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 5 (II.A.1), R¹ is halo.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 10 (II.A.1), R¹ is Cl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 15 (II.A.1), R¹ is F.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 20 (II.A.1), R¹ is alkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-\text{CH}_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-\text{CH}_2\text{CH}_3$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is heteroalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 15 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 20 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-\text{OH}$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
5 (II.A.1), R¹ is -O-alkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
10 (II.A.1), R¹ is -O-aryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
15 (II.A.1), R¹ is -O-heteroalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
20 (II.A.1), R¹ is -O-heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-SH$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-S$ -alkyl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-S$ -aryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 15 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-S$ -heteroalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 20 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-S$ -heteroaryl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-NH_2$.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-NHR^{14}$. Non-limiting examples of R^1 when R^1 is $-NHR^{14}$ include:




wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each R_{aa} is independently selected from haloalkyl (non-limiting examples of which include $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, etc.), R_{ab} is selected from OH, OAc, and $-\text{O-alkyl}$ (non-limiting examples of which include $-\text{O-Me}$, $-\text{O-Et}$, $-\text{O-}n\text{-Pr}$, $-\text{O-}i\text{-Pr}$, $-\text{O-}n\text{-Bu}$, $-\text{O-}i\text{-Bu}$, and $-\text{O-}t\text{-Bu}$), $-\text{O-haloalkyl}$ (non-limiting examples of which include $-\text{O-CH}_2\text{F}$, $-\text{O-CHF}_2$, and $-\text{O-CF}_3$), $-\text{NH}_2$, $-\text{NHalkyl}$, and $-\text{N(alkyl)}_2$.

Additional non-limiting examples of R^1 when R^1 is $-\text{NHR}^{14}$ include:




wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein R_{af} is selected from H and acetyl. It shall be understood that positional isomers of the heteroatoms shown in the moieties above are also contemplated. Such positional isomers



include symmetric positional isomers such as


Additional non-limiting examples of R^1 when R^1 is $-\text{NHR}^{14}$ include:



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, cycloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl,

-O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected from alkyl and haloalkyl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-NR^{14}R^{15}$. Non-limiting examples of R^1 when R^1 is $-NR^{14}R^{15}$ include:




wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each R_{aa} is independently selected from haloalkyl (non-limiting examples of which include $-CH_2F$, $-CHF_2$, $-CF_3$, etc.), R_{ab} is selected from OH, OAc, and -O-alkyl (non-limiting examples of which include -O-Me, -O-Et, -O-*n*-Pr, -O-*i*-Pr, -O-*n*-Bu, -O-*i*-Bu, and -O-*t*-Bu), -O-haloalkyl (non-limiting examples of which include -O- CH_2F , -O- CHF_2 , and -O- CF_3), -NH₂, -NHalkyl, and -N(alkyl)₂.

Additional non-limiting examples of R^1 when R^1 is $-NR^{14}R^{15}$ include:


70



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein R_{af} is selected from H and acetyl. It shall be understood that positional isomers of the heteroatoms shown in the moieties above are also contemplated. Such positional isomers



Additional non-limiting examples of R^1 when R^1 is $-NR^{14}R^{15}$ include:




wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN,

- 5 alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹,
- 10 -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl,

-O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected from alkyl and haloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), R^1 is $-NR^{14}R^{15}$, wherein R^{14} and R^{15} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl.

Non-limiting examples of R^1 when R^1 is $-NR^{14}R^{15}$ and R^{14} and R^{15} are so linked include:



In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1):

R is selected from aryl, heteroaryl, benzo-fused heteroaryl, cycloalkyl, cycloalkenyl, benzo-fused cycloalkyl, benzo-fused cycloalkenyl, heterocycloalkyl, and benzo-fused heterocycloalkyl,

wherein each of said alkyl, said aryl, said heteroaryl, said benzo-fused heteroaryl, said cycloalkyl, said cycloalkenyl, said heterocycloalkyl, said heterocycloalkenyl, and said benzo-fused heterocycloalkyl is unsubstituted or optionally independently substituted with from one to three substituents, which are the same or different, each substituent being independently selected from halo, -OH, -CN, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heterohaloalkyl, -alkyl-OH, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl,

-C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂,
 -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂,
 -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of
 5 said substituted aryl and said substituted heteroaryl independently contains from one to
 three substituents, which may be the same or different, each substituent being
 independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl;

R¹ is selected from -NH₂, -NHR¹⁴, and -NR¹⁴R¹⁵; and

Z is selected from H, halo, alkyl, -OH, haloalkyl, and cycloalkyl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 10 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), Z is halo; R¹ is selected from -NH₂, -NHR¹⁴, and -NR¹⁴R¹⁵; and R is as defined in
 claim 1.

15 In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and
 (II.A.1), Z is heteroaryl; R¹ is selected from -NH₂, -NHR¹⁴, and -NR¹⁴R¹⁵; and R is as
 20 defined in claim 1.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and

(II.A.1), Z is H; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is as defined in claim 1.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is alkyl; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is as defined in claim 1.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 10 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is halo; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2),
 15 (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is heteroaryl; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is heteroaryl.

20 In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),

(I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is H; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 5 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), Z is alkyl; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R is heteroaryl.

In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i),
 10 (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i), (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1), X and Y are each N; R is selected from unsubstituted aryl, substituted aryl, unsubstituted heteroaryl, and substituted heteroaryl, wherein said substituents, when present, are defined in Formula (I); Z is selected from halo, $-OH$, $-SH$, alkyl, $-NH_2$, $-NHR^{12}$, and
 15 $-NR^{12}R^{13}$; R^1 is selected from $-NH_2$, $-NHR^{14}$, and $-NR^{14}R^{15}$; and R^2 is selected from H and alkyl.


In other embodiments, in each of Formulas (I), (I.A), (I.a), (I.a.1), (I.a.1.i), (I.a.2), (I.a.2.i), (I.a.3), (I.a.3.i), (I.a.4), (I.a.4.i), (I.a.5), (I.a.5.i), (I.a.6), (I.a.6.i), (I.a.7), (I.a.7.i), (I.a.8), (I.a.8.i), (I.a.10), (I.a.10.i), (I.a.10.j), (I.B), (I.b), (I.b.1), (I.b.1.i), (I.b.2), (I.b.2.i),
 20 (I.C), (I.c), (I.c.1), (I.c.1.i), (I.c.2), (I.c.2.i), (I.D), (I.d), (I.d.1), (I.d.1.i), (I.E), (II), (II.A), and (II.A.1):

X is N;

Y is N;


R is selected from the group consisting of:

(a) an unsubstituted or substituted monocyclic aryl moiety or an unsubstituted or substituted heteroaryl moiety selected from the group consisting of:



- 5 wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a, R_b, R_c, R_d, and R_e, is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl,
- 10 -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl,
- 15 -O-alkyl, and -C(O)Oalkyl, and

(b) an unsubstituted or an substituted bicyclic heteroaryl moiety selected from the group consisting of:




- 5 wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl,
- 10 -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may

be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl;

R^1 is selected from the group consisting of:


(a) $-NH_2$,



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each R_{aa} is independently selected from haloalkyl (non-limiting examples of which include $-CH_2F$, $-CHF_2$, $-CF_3$, etc.), R_{ab} is selected from OH, OAc, and -O-alkyl (non-limiting examples of which include $-O-Me$, $-O-Et$, $-O-n-Pr$, $-O-i-Pr$, $-O-n-Bu$, $-O-i-Bu$, and $-O-t-Bu$), -O-haloalkyl (non-limiting examples of which include $-O-CH_2F$, $-O-CHF_2$, and $-O-CF_3$), $-NH_2$, $-NHalkyl$, and $-N(alkyl)_2$,


10

79



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein R_{af} is selected from H and acetyl,


(d)



5


wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN,

alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰,
 5 -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹,
 -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and
 said substituted heteroaryl independently contains from one to three substituents, which may
 be the same or different, each substituent being independently selected from halo, alkyl,
 -O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected
 10 from alkyl and haloalkyl,



wherein the wavy line represents the point of attachment of R¹ to the rest of the molecule, and
 wherein each R_{aa} is independently selected from haloalkyl (non-limiting examples of which
 15 include -CH₂F, -CHF₂, -CF₃, etc.), R_{ab} is selected from OH, OAc, and -O-alkyl (non-limiting
 examples of which include -O-Me, -O-Et, -O-*n*-Pr, -O-*i*-Pr, -O-*n*-Bu, -O-*i*-Bu, and -O-*t*-


Bu), -O-haloalkyl (non-limiting examples of which include -O-CH₂F, -O-CHF₂, and -O-CF₃), -NH₂, -NHalkyl, and -N(alkyl)₂,




wherein the wavy line represents the point of attachment of R¹ to the rest of the molecule, and

5 wherein R_{af} is selected from H and acetyl,

(g)



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected from alkyl and haloalkyl, and



Z is selected from the group consisting of H, halo, -OH, -SH, -CN, alkyl, alkenyl, alkynyl, heteroalkyl, haloalkyl, heterohaloalkyl, -S-alkyl, -O-alkyl, -O-aryl, -O-heteroaryl, cycloalkyl, aryl, heteroaryl, -NH₂, -NHR¹², and -NR¹²R¹³.

In other embodiments, the compounds of the invention have a structural formula as depicted in Table I below and include tautomers, and pharmaceutically acceptable salts, esters, prodrugs, isomers, and solvates of such compounds and such tautomers.

DEFINITIONS

The terms used herein have their ordinary meaning and the meaning of such terms is independent at each occurrence thereof. That notwithstanding and except where stated otherwise, the following definitions apply throughout the specification and claims.

5 Chemical names, common names and chemical structures may be used interchangeably to describe that same structure. These definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Hence the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portion of "hydroxyalkyl", "haloalkyl", arylalkyl-, alkylaryl-, "alkoxy" etc.

10 "At least one" means one or more than one, for example, 1, 2, or 3, or in another example, 1 or 2, or in another example 1.

"One or more" means one or more than one, for example, 1, 2, or 3, or in another example, 1 or 2, or in another example 1.

"Patient" includes both human and non-human animals. Non-human animals include
15 research animals, farm animals, and companion animals such as mice, primates, monkeys, great apes, cows, sheep, horse, canine (*e.g.*, dogs), and feline (*e.g.*, house cats), etc.

"Composition" includes "pharmaceutical composition" and other compositions not suitable for pharmaceutical use but which may be suitable for other uses such as research or other uses.

20 "Pharmaceutical composition" (or "pharmaceutically acceptable composition") means a composition suitable for administration to a patient. Such compositions may contain the neat compound (or compounds) of the invention or mixtures thereof, or salts, solvates, prodrugs, isomers, or tautomers thereof, or they may contain one or more pharmaceutically acceptable carriers or diluents. The term "pharmaceutical composition" is also intended to
25 encompass both the bulk composition and individual dosage units comprised of more than one (*e.g.*, two) pharmaceutically active agents such as, for example, a compound of the present invention and an additional agent selected from the lists of the additional agents described herein, along with any pharmaceutically inactive excipients. The bulk composition and each individual dosage unit can contain fixed amounts of the afore-said "more than one
30 pharmaceutically active agents". The bulk composition is material that has not yet been formed into individual dosage units. An illustrative dosage unit is an oral dosage unit such as tablets, pills and the like. Similarly, the herein-described method of treating a patient by

administering a pharmaceutical composition of the present invention is also intended to encompass the administration of the afore-said bulk composition and individual dosage units.

“Halogen” means fluorine, chlorine, bromine, or iodine. Preferred are fluorine, chlorine and bromine.

5 “Alkyl” means an aliphatic hydrocarbon group which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. “Lower alkyl” means a group
10 having about 1 to about 6 carbon atoms in the chain which may be straight or branched. “Alkyl” may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being as described herein or independently selected from the group consisting of halo, alkyl, haloalkyl, spirocycloalkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -O-C(O)-
15 alkyl, -O-C(O)-aryl, -O-C(O)-cycloalkyl, carboxy and -C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl and t-butyl.

“Haloalkyl” means an alkyl as defined above wherein one or more hydrogen atoms on the alkyl is replaced by a halo group defined above.

20 “Aminoalkyl” means an alkyl which has been substituted at one or more available carbon atoms by one or more amino group(s). Non-limiting examples of such amino groups include those described herein, such as -NH₂, -NHR¹², -NR¹²R¹³, -NHR¹⁴, and -NHR¹⁵.

 “Heteroalkyl” means an alkyl moiety as defined above, having one or more carbon atoms, for example one, two or three carbon atoms, including a terminal carbon atom, replaced with one or more heteroatoms, which may be the same or different, where the point
25 of attachment to the remainder of the molecule is through a carbon atom of the heteroalkyl radical. Suitable such heteroatoms include O, S, S(O), S(O)₂, -NH-, -N(alkyl)-, and -N(alkyl)₂. Non-limiting examples include ethers, thioethers, amines, hydroxymethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 2-methoxyethyl, 2-aminoethyl, 2-dimethylaminoethyl, and the like. Additional non-limiting examples include -alkyl-NHalkyl and -alkyl-N(alkyl)₂.
30 A non-limiting example of heteroalkyl wherein a terminal carbon atom is replaced with a heteroatom includes -alkyl-NH₂.

"Heterohaloalkyl" means an haloalkyl moiety as defined above, having one or more, for example one, two, or three carbon atoms, including a terminal carbon atom, replaced with one or more heteroatoms, which may be the same or different, where the point of attachment to the remainder of the molecule is through a carbon atom of the heterohaloalkyl radical.

5 "Alkenyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkenyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are
10 attached to a linear alkenyl chain. "Lower alkenyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. "Alkenyl" may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, alkoxy and -S(alkyl). Non-limiting examples of suitable alkenyl groups include ethenyl,
15 propenyl, n-butenyl, 3-methylbut-2-enyl, n-pentenyl, octenyl and decenyl.

"Alkylene" means a difunctional group obtained by removal of a hydrogen atom from an alkyl group that is defined above. Non-limiting examples of alkylene include methylene, ethylene and propylene. More generally, the suffix "ene" on alkyl, aryl, heterocycloalkyl, etc.

indicates a divalent moiety, e.g., -CH₂CH₂- is ethylene, and  is para-phenylene.

20 "Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear
25 alkynyl chain. "Lower alkynyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butyne and 3-methylbutynyl. "Alkynyl" may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and
30 cycloalkyl.

"Alkenylene" means a difunctional group obtained by removal of a hydrogen from an alkenyl group that is defined above. Non-limiting examples of alkenylene include $-\text{CH}=\text{CH}-$, $-\text{C}(\text{CH}_3)=\text{CH}-$, and $-\text{CH}=\text{CHCH}_2-$.


"Aryl" means an aromatic monocyclic or multicyclic ring system comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a heteroaryl can be optionally oxidized to the corresponding N-oxide. "Heteroaryl" may also include a heteroaryl as defined above fused to an aryl as defined above. Non-limiting examples of suitable heteroaryls include pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridone (including N-substituted pyridones), isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxaliny, phthalazinyl, oxindolyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinolinyl, imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinolinyl, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like. The term "heteroaryl" also refers to partially saturated heteroaryl moieties such as, for example, tetrahydroisoquinolyl, tetrahydroquinolyl and the like.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable monocyclic cycloalkyls include

cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like.

Further non-limiting examples of cycloalkyl include the following:




"Spirocycloalkyl" means a cycloalkyl moiety in which two available hydrogen atoms attached to the same carbon atom are replacted to form a cycloalkyl group.

10 "Cycloalkenyl" means a non-aromatic mono or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms which contains at least one carbon-carbon double bond. Preferred cycloalkenyl rings contain about


5 to about 7 ring atoms. The cycloalkenyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkenyls include cyclopentenyl, cyclohexenyl, cyclohepta-1,3-dienyl, and the like. Non-limiting example of a suitable
5 multicyclic cycloalkenyl is norbornylenyl.

"Heterocycloalkyl" (or "heterocyclyl") means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no
10 adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. Any -NH in a heterocyclyl ring may exist protected such as, for example, as an -N(Boc), -N(CBz), -N(Tos) group and the like; such protections are also considered part of
15 this invention. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Thus, the term "oxide," when it appears in a definition of a variable in a general structure described herein, refers to the corresponding N-oxide, S-oxide, or S,S-
20 dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, lactam, lactone, and the like. "Heterocyclyl" also includes rings wherein =O replaces two available hydrogens on the same carbon atom (i.e., heterocyclyl includes rings having a carbonyl group in the ring). Such =O groups may be
25 referred to herein as "oxo." An example of such a moiety is pyrrolidinone (or pyrrolidone):




"Heterocycloalkenyl" (or "heterocyclenyl") means a non-aromatic monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to

about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur atom, alone or in combination, and which contains at least one carbon-carbon double bond or carbon-nitrogen double bond. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclenyl rings contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclenyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclenyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. The nitrogen or sulfur atom of the heterocyclenyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable heterocyclenyl groups include 1,2,3,4- tetrahydropyridinyl, 1,2-dihydropyridinyl, 1,4-dihydropyridinyl, 1,2,3,6- tetrahydropyridinyl, 1,4,5,6-tetrahydropyrimidinyl, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazoliny, 2-pyrazoliny, dihydroimidazolyl, dihydrooxazolyl, dihydrooxadiazolyl, dihydrothiazolyl, 3,4-dihydro-2H-pyranyl, dihydrofuranyl, fluorodihydrofuranyl, 7-oxabicyclo[2.2.1]heptenyl, dihydrothiophenyl, dihydrothiopyranyl, and the like. "Heterocyclenyl" also includes rings wherein =O replaces two available hydrogens on the same carbon atom (i.e., heterocyclenyl includes rings having a carbonyl group in the ring). Example of such moiety is pyrrolidenone (or pyrrolone):



It should be noted that in hetero-atom containing ring systems of this invention, there are no hydroxyl groups on carbon atoms adjacent to a N, O or S, as well as there are no N or S groups on carbon adjacent to another heteroatom. Thus, for example, in the ring:




there is no -OH attached directly to carbons marked 2 and 5.

It should also be noted that tautomeric forms of the compounds of the invention are also contemplated as being within the scope of the invention.

“Arylcycloalkyl” (or “arylfused cycloalkyl”) means a group derived from a fused aryl and cycloalkyl as defined herein. Preferred arylcycloalkyls are those wherein aryl is phenyl (which may be referred to as “benzofused”) and cycloalkyl consists of about 5 to about 6 ring atoms. The arylcycloalkyl can be optionally substituted as described herein. Non-limiting examples of suitable arylcycloalkyls include indanyl (a benzofused cycloalkyl) and 1,2,3,4-tetrahydronaphthyl and the like. The bond to the parent moiety is through a non-aromatic carbon atom.

“Arylheterocycloalkyl” (or “arylfused heterocycloalkyl”) means a group derived from a fused aryl and heterocycloalkyl as defined herein. Preferred arylheterocycloalkyls are those wherein aryl is phenyl (which may be referred to as “benzofused”) and heterocycloalkyl consists of about 5 to about 6 ring atoms. The arylheterocycloalkyl can be optionally substituted, and/or contain the oxide or oxo, as described herein. Non-limiting examples of suitable arylfused heterocycloalkyls include:



The bond to the parent moiety is through a non-aromatic carbon atom.

It is also understood that the terms “arylfused aryl”, “arylfused cycloalkyl”, “arylfused cycloalkenyl”, “arylfused heterocycloalkyl”, “arylfused heterocycloalkenyl”, “arylfused heteroaryl”, “cycloalkylfused aryl”, “cycloalkylfused cycloalkyl”, “cycloalkylfused cycloalkenyl”, “cycloalkylfused heterocycloalkyl”, “cycloalkylfused heterocycloalkenyl”, “cycloalkylfused heteroaryl”, “cycloalkenylfused aryl”, “cycloalkenylfused cycloalkyl”, “cycloalkenylfused cycloalkenyl”, “cycloalkenylfused heterocycloalkyl”, “cycloalkenylfused heterocycloalkenyl”, “cycloalkenylfused heteroaryl”, “heterocycloalkylfused aryl”, “heterocycloalkylfused cycloalkyl”, “heterocycloalkylfused cycloalkenyl”, “heterocycloalkylfused heterocycloalkyl”, “heterocycloalkylfused heterocycloalkenyl”, “heterocycloalkylfused heteroaryl”, “heterocycloalkenylfused aryl”, “heterocycloalkenylfused cycloalkyl”, “heterocycloalkenylfused cycloalkenyl”, “heterocycloalkenylfused heterocycloalkyl”, “heterocycloalkenylfused heterocycloalkenyl”, “heterocycloalkenylfused heteroaryl”, “heteroaryl-fused aryl”, “heteroaryl-fused cycloalkyl”,

“heteroaryl-fused cycloalkenyl”, “heteroaryl-fused heterocycloalkyl”, “heteroaryl-fused heterocycloalkenyl”, and “heteroaryl-fused heteroaryl” are similarly represented by the combination of the groups aryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, and heteroaryl, as previously described. Any such groups may be unsubstituted or substituted with one or more ring system substituents at any available position as described herein.

“Aralkyl” or “arylalkyl” means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the parent moiety is through the alkyl. The term (and similar terms) may be written as “arylalkyl-” to indicate the point of attachment to the parent moiety.

Similarly, “heteroarylalkyl”, “cycloalkylalkyl”, “cycloalkenylalkyl”, “heterocycloalkylalkyl”, “heterocycloalkenylalkyl”, etc., mean a heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, etc. as described herein bound to a parent moiety through an alkyl group. Preferred groups contain a lower alkyl group. Such alkyl groups may be straight or branched, unsubstituted and/or substituted as described herein.

Similarly, “aryl-fused arylalkyl-”, aryl-fused cycloalkylalkyl-, etc., means an aryl-fused aryl group, aryl-fused cycloalkyl group, etc. linked to a parent moiety through an alkyl group. Preferred groups contain a lower alkyl group. Such alkyl groups may be straight or branched, unsubstituted and/or substituted as described herein.

“Alkylaryl” means an alkyl-aryl- group in which the alkyl and aryl are as previously described. Preferred alkylaryls comprise a lower alkyl group. Non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl.

“Cycloalkylether” means a non-aromatic ring of 3 to 7 members comprising an oxygen atom and 2 to 7 carbon atoms. Ring carbon atoms can be substituted, provided that substituents adjacent to the ring oxygen do not include halo or substituents joined to the ring through an oxygen, nitrogen or sulfur atom.

“Cycloalkylalkyl” means a cycloalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkylalkyls include cyclohexylmethyl, adamantylmethyl, adamantylpropyl, and the like.

“Cycloalkenylalkyl” means a cycloalkenyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkenylalkyls include cyclopentenylmethyl, cyclohexenylmethyl and the like.

5 “Heteroarylalkyl” means a heteroaryl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable heteroaryls include 2-pyridinylmethyl, quinolinylmethyl and the like.

10 “Heterocyclalkyl” (or “heterocycloalkyl”) means a heterocyclalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable heterocyclalkyls include piperidinylmethyl, piperazinylmethyl and the like.

“Heterocyclenylalkyl” means a heterocyclenyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core.

15 “Alkynylalkyl” means an alkynyl-alkyl- group in which the alkynyl and alkyl are as previously described. Preferred alkynylalkyls contain a lower alkynyl and a lower alkyl group. The bond to the parent moiety is through the alkyl. Non-limiting examples of suitable alkynylalkyl groups include propargylmethyl.

20 “Heteroaralkyl” means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

“Hydroxyalkyl” means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

25 “Cyanoalkyl” means a NC-alkyl- group in which alkyl is as previously defined. Preferred cyanoalkyls contain lower alkyl. Non-limiting examples of suitable cyanoalkyl groups include cyanomethyl and 2-cyanoethyl.

30 “Acyl” means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1-naphthoyl.

5 "Heteroaroyl" means an heteroaryl-C(O)- group in which the heteroaryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include pyridoyl.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, *n*-propoxy, isopropoxy and *n*-butoxy. The bond to the parent moiety is through the ether
10 oxygen.

"Alkyoxyalkyl" means a group derived from an alkoxy and alkyl as defined herein. The bond to the parent moiety is through the alkyl.

"Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond
15 to the parent moiety is through the ether oxygen.

"Aralkyloxy" (or "arylalkyloxy") means an aralkyl-O- group (an arylalkyl-O- group) in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is through the ether oxygen.

20 "Arylalkenyl" means a group derived from an aryl and alkenyl as defined herein. Preferred arylalkenyls are those wherein aryl is phenyl and the alkenyl consists of about 3 to about 6 atoms. The arylalkenyl can be optionally substituted by one or more substituents. The bond to the parent moiety is through a non-aromatic carbon atom.

"Arylalkynyl" means a group derived from a aryl and alkynyl as defined herein.
25 Preferred arylalkynyls are those wherein aryl is phenyl and the alkynyl consists of about 3 to about 6 atoms. The arylalkynyl can be optionally substituted by one or more substituents. The bond to the parent moiety is through a non-aromatic carbon atom.

"Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkylthio groups include methylthio and
30 ethylthio. The bond to the parent moiety is through the sulfur.

"Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

5 "Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

"Alkoxycarbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl. The bond to the parent moiety is through the carbonyl.

10 "Aryloxycarbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aralkoxycarbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxycarbonyl group is benzyloxycarbonyl. The bond to the parent moiety is
15 through the carbonyl.

"Alkylsulfonyl" means an alkyl-S(O₂)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

"Arylsulfonyl" means an aryl-S(O₂)- group. The bond to the parent moiety is through the sulfonyl.

20 "Spriocycloalkyl" means a cycloalkyl group attached to a parent moiety at a single carbon atom. Non-limiting examples of spirocycloalkyl wherein the parent moiety is a cycloalkyl include spiro [2.5] octane, spiro [2.4] heptane, etc. Non-limiting examples of spriocycloalkyl wherein the parent moiety is an The alkyl moiety linking fused ring systems (such as the alkyl moiety in heteroaryl fused heteroarylalkyl-) may optionally be substituted
25 with spirocycloalkyl or other groups as described herein. Non-limiting spirocycloalkyl groups include spirocyclopropyl, spriorcyclobutyl, spirocycloheptyl, and spirocyclohexyl.

The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution
30 results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful

degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term “optionally substituted” means optional substitution with the specified groups, radicals or moieties.

5 Substitution on a cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylfused cycloalkylalkyl- moiety or the like includes substitution on any ring portion and/or on the alkyl portion of the group.


When a variable appears more than once in a group, e.g., R^8 in $-N(R^8)_2$, or a variable appears more than once in a structure presented herein such as Formula (I), the variables can
10 be the same or different.

With reference to the number of moieties (e.g., substituents, groups or rings) in a compound, unless otherwise defined, the phrases “one or more” and “at least one” mean that there can be as many moieties as chemically permitted, and the determination of the maximum number of such moieties is well within the knowledge of those skilled in the
15 art. With respect to the compositions and methods comprising the use of “at least one compound of the invention, e.g., of Formula (I),” one to three compounds of the invention, e.g., of Formula (I) can be administered at the same time, preferably one.

Compounds of the invention may contain one or more rings having one or more ring system substituents. “Ring system substituent” means a substituent attached to an aromatic or
20 non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being as described herein or independently selected from the group consisting of alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, aryl, heteroaryl, aralkyl, alkylaryl, heteroaralkyl, heteroarylalkenyl, heteroarylalkynyl, alkylheteroaryl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl,
25 aroyl, halo, nitro, cyano, carboxy, alkoxycarbonyl, aryloxycarbonyl, aralkoxycarbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, heterocyclyl, $-O-C(O)-$ alkyl, $-O-C(O)-$ aryl, $-O-C(O)-$ cycloalkyl, $-C(=N-CN)-NH_2$, $-C(=NH)-NH_2$, $-C(=NH)-NH(alkyl)$, Y_1Y_2N- , $Y_1Y_2N-alkyl-$, $Y_1Y_2NC(O)-$, $Y_1Y_2NSO_2-$ and $-SO_2NY_1Y_2$, wherein Y_1 and Y_2 can be the same or different
30 and are independently selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl, and aralkyl. “Ring system substituent” may also mean a single moiety which simultaneously replaces two available hydrogens on two adjacent carbon atoms (one H on each carbon) on a

ring system. Examples of such moieties are rings such as heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl rings. Additional non-limiting examples include methylene dioxy, ethylenedioxy, $-\text{C}(\text{CH}_3)_2-$ and the like which form moieties such as, for example:


5



As used herein, the term “composition” is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.


10

The line ----, as a bond generally indicates a mixture of, or either of, the possible isomers, e.g., containing (R)- and (S)- stereochemistry. For example:




15

The wavy line ~~~, as used herein, indicates a point of attachment to the rest of the compound. For example, each wavy line in the following structure:



indicates a point of attachment to the core structure, as described herein.

Lines drawn into the ring systems, such as, for example:




20

indicate that the indicated line (bond) may be attached to any of the substitutable ring carbon atoms.

“Oxo” is defined as a oxygen atom that is double bonded to a ring carbon in a cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, or other ring described herein, e.g.,


97



In this specification, where there are multiple oxygen and/or sulfur atoms in a ring system, there cannot be any adjacent oxygen and/or sulfur present in said ring system.

It is noted that the carbon atoms for compounds of the invention may be replaced with 1 to 3 silicon atoms so long as all valency requirements are satisfied.

As well known in the art, a bond drawn from a particular atom wherein no moiety is depicted at the terminal end of the bond indicates a methyl group bound through that bond to the atom, unless stated otherwise. For example:



The term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of said compound after being isolated from a synthetic process (e.g. from a reaction mixture), or natural source or combination thereof. Thus, the term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of said compound after being obtained from a purification process or processes described herein or well known to the skilled artisan (e.g., chromatography, recrystallization and the like), in sufficient purity to be characterizable by standard analytical techniques described herein or well known to the skilled artisan.

It should also be noted that any carbon (or other atom or heteroatom) with unsatisfied valences in the text, schemes, examples and Tables herein is assumed to have the sufficient number of hydrogen atom(s) to satisfy the valences.

When a functional group in a compound is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for

example, T. W. Greene *et al*, *Protective Groups in Organic Synthesis* (1991), Wiley, New York.

As used herein, the term “composition” is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. A discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems* (1987) 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press. The term “prodrug” means a compound (e.g, a drug precursor) that is transformed *in vivo* to yield a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate of the compound. The transformation may occur by various mechanisms (e.g., by metabolic or chemical processes), such as, for example, through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, “Pro-drugs as Novel Delivery Systems,” Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

For example, if a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate of the compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as, for example, (C₁–C₈)alkyl, (C₂–C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁–C₂)alkylamino(C₂–C₃)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C₁–C₂)alkyl, N,N-di(C₁–C₂)alkylcarbamoyl-(C₁–C₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂–C₃)alkyl, and the like.

Similarly, if a compound of the invention contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as, for example, (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl, (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α -amino(C₁-C₄)alkanyl, arylacyl and α -aminoacyl, or α -aminoacyl- α -aminoacyl, where each α -aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate), and the like.

If a compound of the invention incorporates an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the amine group with a group such as, for example, R-carbonyl, RO-carbonyl, NRR'-carbonyl where R and R' are each independently (C₁-C₁₀)alkyl, (C₃-C₇) cycloalkyl, benzyl, or R-carbonyl is a natural α -aminoacyl or natural α -aminoacyl, —C(OH)C(O)OY¹ wherein Y¹ is H, (C₁-C₆)alkyl or benzyl, —C(OY²)Y³ wherein Y² is (C₁-C₄) alkyl and Y³ is (C₁-C₆)alkyl, carboxy (C₁-C₆)alkyl, amino(C₁-C₄)alkyl or mono-N— or di-N,N-(C₁-C₆)alkylaminoalkyl, —C(Y⁴)Y⁵ wherein Y⁴ is H or methyl and Y⁵ is mono-N— or di-N,N-(C₁-C₆)alkylamino morpholino, piperidin-1-yl or pyrrolidin-1-yl, and the like.

One or more compounds of the invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. "Solvate" means a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanlates, methanlates, and the like. "Hydrate" is a solvate wherein the solvent molecule is H₂O.

One or more compounds of the invention may optionally be converted to a solvate. Preparation of solvates is generally known. Thus, for example, M. Caira *et al*, *J. Pharmaceutical Sci.*, 93(3), 601-611 (2004) describe the preparation of the solvates of the antifungal fluconazole in ethyl acetate as well as from water. Similar preparations of solvates,

hemisolvate, hydrates and the like are described by E. C. van Tonder *et al*, *AAPS PharmSciTech.*, 5(1), article 12 (2004); and A. L. Bingham *et al*, *Chem. Commun.*, 603-604 (2001). A typical, non-limiting, process involves dissolving the inventive compound in desired amounts of the desired solvent (organic or water or mixtures thereof) at a higher than ambient temperature, and cooling the solution at a rate sufficient to form crystals which are then isolated by standard methods. Analytical techniques such as, for example IR spectroscopy, show the presence of the solvent (or water) in the crystals as a solvate (or hydrate).

"Effective amount" or "therapeutically effective amount" is meant to describe an amount of compound or a composition of the present invention effective in inhibiting the above-noted diseases and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect.

The compounds of the invention can form salts which are also within the scope of this invention. Reference to a compound of the invention herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of the invention contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of the invention may be formed, for example, by reacting a compound of the invention with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) and the like. Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are

discussed, for example, by P. Stahl *et al*, Camille G. (eds.) *Handbook of Pharmaceutical Salts. Properties, Selection and Use*. (2002) Zurich: Wiley-VCH; S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

Pharmaceutically acceptable esters of the present compounds include the following groups: (1) carboxylic acid esters obtained by esterification of the hydroxy groups, in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, acetyl, n-propyl, t-butyl, or n-butyl), alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted with, for example, halogen, C₁₋₄alkyl, or C₁₋₄alkoxy or amino); (2) sulfonate esters, such as alkyl- or aralkylsulfonyl (for example, methanesulfonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); (4) phosphonate esters and (5) mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a C₁₋₂₀ alcohol or reactive derivative thereof, or by a 2,3-di (C₆₋₂₄)acyl glycerol.

Compounds of the invention, and salts, solvates, esters and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

The compounds of the invention may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the compounds of the invention as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention embraces all geometric and positional isomers. For example, if a compound of the invention incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the invention.

Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of the invention may be atropisomers (e.g., substituted biaryls) and are considered as part of this invention. Enantiomers can also be separated by use of chiral HPLC column.

It is also possible that the compounds of the invention may exist in different tautomeric forms, and all such forms are embraced within the scope of the invention. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the invention.

All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this invention, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). (For example, if a compound of the invention incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the invention. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the invention.).

Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the *IUPAC* 1974 Recommendations. The use of the
5 terms "salt", "solvate", "ester", "prodrug" and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

The present invention also embraces isotopically-labelled compounds of the present invention which are identical to those recited herein, but for the fact that one or more atoms
10 are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively.

15 Certain isotopically-labelled compounds of the invention (e.g., those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (i.e., ^3H) and carbon-14 (i.e., ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g.,
20 increased in vivo half-life or reduced dosage requirements) and hence may be preferred in some circumstances. Isotopically labelled compounds of the invention can generally be prepared by following procedures analogous to those disclosed in the Schemes and/or in the Examples hereinbelow, by substituting an appropriate isotopically labelled reagent for a non-isotopically labelled reagent.

25 Polymorphic forms of the compounds of the invention, and of the salts, solvates, esters and prodrugs of the compounds of the invention, are intended to be included in the present invention.

Suitable doses for administering compounds of the invention to patients may readily be determined by those skilled in the art, e.g., by an attending physician, pharmacist, or other
30 skilled worker, and may vary according to patient health, age, weight, frequency of administration, use with other active ingredients, and/or indication for which the compounds are administered. Doses may range from about 0.001 to 500 mg/kg of body weight/day of the

compound of the invention. In one embodiment, the dosage is from about 0.01 to about 25 mg/kg of body weight/day of a compound of the invention, or a pharmaceutically acceptable salt or solvate of said compound. In another embodiment, the quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 100 mg, preferably from about 1 mg to about 50 mg, more preferably from about 1 mg to about 25 mg, according to the particular application. In another embodiment, a typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 500 mg/day, preferably 1 mg/day to 200 mg/day, in two to four divided doses.

As discussed above, the amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated.

When used in combination with one or more additional therapeutic agents, the compounds of this invention may be administered together or sequentially. When administered sequentially, compounds of the invention may be administered before or after the one or more additional therapeutic agents, as determined by those skilled in the art or patient preference.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically active agent or treatment within its dosage range.

Accordingly, in an aspect, this invention includes combinations comprising an amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof, and an effective amount of one or more additional agents described above.

The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological assays. Certain assays are exemplified elsewhere in this document.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium

carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

The compounds of this invention may also be delivered subcutaneously.

In one embodiment, the compound is administered orally.

In some embodiments, it may be advantageous for the pharmaceutical preparation comprising one or more compounds of the invention be prepared in a unit dosage form. In such forms, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

PREPARATIVE EXAMPLES

Compounds of the invention can be made using procedures known in the art. The following reaction schemes show typical procedures, but those skilled in the art will recognize that other procedures can also be suitable.

Techniques, solvents and reagents may be referred to by their following abbreviations:

Thin layer chromatography: TLC

High performance liquid chromatography: HPLC

ethyl acetate: AcOEt or EtOAc

methanol: MeOH

5 ether: Et₂O

tetrahydrofuran: THF

Acetonitrile: MeCN

1,2-dimethoxyethane: DME

Trifluoroacetic acid: TFA

10 Dimethylacetamide: DMA

Dimethylformamide: DMF

Dimethylsulfoxide: DMSO

triethylamine: Et₃N or TEA

tert-Butoxycarbonyl: t-Boc or Boc

15 2-(Trimethylsilyl)ethoxycarbonyl: Teoc

nuclear magnetic resonance spectroscopy: NMR

liquid chromatography mass spectrometry: LCMS

high resolution mass spectrometry: HRMS

milliliters: mL

20 millimoles: mmol

microliters: μ l

grams: g

milligrams: mg

centimeters: cm

25 room temperature (ambient, about 25°C): rt

Retention time: tR

N-bromosuccinimide: NBS

N-chlorosuccinimide: NCS

Methyl magnesium bromide: MeMgBr

30 iron(III) acetylacetonate: Fe(acac)₃

Diphenylphosphoryl azide: DPPA

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride: EDCI

Diisopropylethylamine: DIEA or *i*-Pr₂NEt or DIPEA

Diisopropylamine: *i*-Pr₂NH

2-(Trimethylsilyl)ethanol: TMSethanol

3-Chloroperoxybenzoic acid: mCPBA

5 n-Butyllithium: nBuLi

lithium diisopropylamide: LDA


[1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II): PdCl₂dppf

Palladium(II) acetate: Pd(OAc)₂

Methanesulfonyl chloride : MeSO₂Cl

10 Triphenyl phosphine: TPP or Ph₃P

General Method 1:



wherein X, Y, Z, R, R¹, R², R³ to R⁹, R¹⁸, R¹⁹, and n are as defined herein.

To a stirred mixture of 2-amino-4,6-dichloropyrimidine (**1**, X, Y = N, R¹ = -NH₂, Z = Cl, 5.0 g, 30.5 mmol) in glacial acetic acid (120 mL) was added dropwise a solution of ICl (5.01 mL, 100 mmol) in glacial acetic acid (120 mL). After 5 h, the mixture was filtered, and the collected solids were washed with glacial acetic acid and then azeotroped with toluene (2X), giving 2.78 g of **2** as a white solid. After 7 days, more solid was visible in the filtrate, and thus, it was again filtered, the collected solids washed with glacial acetic acid and azeotroped

with toluene (2X), giving another 4.22 g of **2**, with TLC and MS data that matched the first batch (79% total yield).

MS m/z (M+H)⁺ 289.93 (2 Cl pattern);

A mixture of **2** (58.0 g, 0.20 mol), the cyclopentylamine sugar (1R, 2S, 3R, 4R)-2,3-dihydroxy-4-(hydroxymethyl)-1-aminocyclopentane hydrochloride, 40.4 g, 0.22 mol) in ethanol (800 ml) and triethylamine (92 ml, 0.66 mol) was refluxed for 18 h, during which time complete dissolution occurred. After concentrating and adsorbing the residue onto silica gel, the crude was purified by chromatography, eluting with a gradient of EtOAc/MeOH (97.5/2.5 → 95/5). The desired product **3** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) was obtained as a white solid (67 g, 83% yield).

MS m/z (M+H)⁺ 401.00 (Cl pattern);

¹H NMR (DMSO-*d*₆): δ 6.61 (s, 2H, D₂O exchangeable), 6.22 (d, 1H, *J* = 7.7 Hz, D₂O exchangeable), 4.75 (dd, 1H, *J* = 4.8, 4.8 Hz, D₂O exchangeable), 4.60 (d, 1H, *J* = 5.2 Hz, D₂O exchangeable), 4.41 (d, 1H, *J* = 4.5 Hz, D₂O exchangeable), 4.30 – 4.18 (m, 1H; upon D₂O exchange collapses to 4.22, dd, *J* = 7.4, 12.9 Hz), 3.77 – 3.70 (m, 2H), 3.39 (dd, 2H, *J* = 5.1, 5.1 Hz; upon D₂O exchange collapses to 3.43, d, *J* = 5.3 Hz), 2.24 – 2.14 (m, 1H), 1.93 – 1.83 (m, 1H), 1.15 – 1.06 (m, 1H).

Analysis calculated for C₁₆H₁₈Cl₂N₆O₃: C, 46.50; H, 4.39; N, 20.34. Found: C, 46.25; H, 4.26; N, 20.09

To a degassed solution of **3** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, 5.0 g, 12.5 mmol) in DMF (50 ml) was added (with protection from light) triethylamine (7.0 mL, 50 mmol) dropwise over 10 min, followed by CuI (952 mg, 5.0 mmol) and then tetrakis(triphenylphosphine)-palladium (2.9 g, 2.5 mmol). After degassing with Ar for 10 min, the addition of TMS acetylene (5.3 mL, 37.5 mmol) was followed by sealing the reaction vessel with a rubber septum. Then, while still protecting the sealed flask from light, the reaction mixture was heated in an oil bath at 55°C for 18 h. After concentrating, the methanol extract was filtered, and the filtrate concentrated onto silica. Chromatography on silica (eluting gradient of CHCl₃/MeOH (95/5 → 90/10) gave 2.62 g of a brown foam that contained triethylamine salts. A second chromatography (eluting with EtOAc/MeOH, 95/5)

gave 2.55 g of **4** (55% yield, X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) as a reddish brown solid. (Even after the 2 chromatographic isolations, the obtained **4** still contained impurities, by TLC. Since these impurities would not interfere with the next reaction and since they could be easily removed after the next reaction step (the conversion of **4** to **5**), the slightly impure **4** fractions could also be used as is in the next reaction.)

¹H NMR (DMSO-*d*₆): δ 6.94 (s, 2H), 6.10 (d, 1H, *J* = 7.7 Hz), 4.68 – 4.62 (m, 2H), 4.45 (d, 1H, *J* = 4.5 Hz), 4.27 – 4.17 (m, 1H), 3.76 – 3.65 (m, 2H), 3.40 (dd, 2H, *J* = 5.1, 5.1 Hz), 2.26 – 2.16 (m, 1H), 1.93 – 1.83 (m, 1H), 1.16 – 1.07 (m, 1H), 0.23 (s, 3H).

To a solution of **4** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) 4.17 g, 11.2 mmol) in acetonitrile (100 ml) was added tetraethylammonium fluoride dihydrate (1.04 g, 5.62 mmol). After 2 h, MeOH was added to dissolve precipitated material, and the resulting solution was concentrated onto silica gel. Chromatography on silica (eluting gradient of CHCl₃/MeOH, 95/5 → 92.5/7.5) followed by chromatography eluting with EtOAc/MeOH (95/5) resulted in the recovery of 3.06 g of **5** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) as a light tan solid (94% yield).

¹H NMR (DMSO-*d*₆): δ 1.10 (m, 1H), 1.89 (m, 1H), 2.15 (m, 1H), 3.38 (t, 2H, *J* = 5.1 Hz), 3.73 (m, 2H), 4.28 (m, 1H), 4.40 (d, 1H, *J* = 3.9 Hz), 4.58 (d, 1H, *J* = 5.1 Hz), 4.71 (t, 1H, *J* = 5.1 Hz), 6.58 (d, 1H, *J* = 8.1 Hz), 6.87 (s, 2H)

MS *m/z* (M+H)⁺: 299.15 (Cl pattern);

Compound 20, (1R,2S,3R,5R)-3-[[2-Amino-6-chloro-5-[2-(2-methoxypyridin-5-yl)ethynyl]-4-pyrimidinyl]amino]-5-(hydroxymethyl)-1,2-cyclopentanediol (**6**, R = 2-methoxypyridin-5-yl).

To a solution of **5** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, 100 mg, 0.335 mmol) in DMF (4 mL) in a small microwave reaction tube was added 5-bromo-2-methoxypyridine (65 μL, 0.487 mmol) and triethylamine (200 μL, 1.43 mmol). The solution was degassed with argon before and during the addition of cuprous iodide (6.4 mg, 0.033 mmol), followed by tetrakis(triphenylphosphine)palladium(O) (20 mg, 0.017 mmol). The tightly capped tube was placed in a microwave reactor at 90°C for 10 min at which time the reaction was complete. Evaporation of solvent in vacuo gave a brown residue

which was purified by silica gel flash chromatography (0 – 10% MeOH in EtOAc). The off-white solid obtained (112 mg) was crystallized from acetone to give pure **6** (R = 2-methoxyxypyridin-5-yl, X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, 76 mg, 56%).

5

¹H NMR (DMSO-d₆): δ 1.20 (m, 1H), 1.87 (m, 1H), 2.17 (m, 1H), 3.41 (m, 2H), 3.78 (m, 2H), 3.88 (s, 3H), 4.32 (m, 1H), 4.41 (d, 1H, J = 4.6 Hz), 4.60 (d, 1H, J = 4.9 Hz), 4.77 (m, 1H), 6.64 (d, 1H, J = 8 Hz), 6.89 (m, 3H), 7.83 (dd, 1H, J = 2.2 Hz, 8.7 Hz), 8.37 (m, 1H).

10 MS *m/z* (M+H)⁺ 406.19 (Cl pattern);

Method 1A, Same as Method 1 except for the last step (Conversion of 5 to 6)

The conversion of **5** to **6** was also done under non-microwave conditions using 1 eq of **5**, 3 eq of the aryl halide, 4 eq of Et₃N, 0.4 eq CuI, and 0.2 eq


15 tetrakis(triphenylphosphine)palladium(O). The addition of components was the same as above. The reaction was run at 50°C overnight, and the desired products were isolated by the same procedure as above. Product yields were consistently in the 25-30% range.

Method 1B (Conversion of Z = Cl (e.g., Compound 21) to Z = Me (e.g., Compound 1)

20 A mixture of **6** (R = 2-ethoxyxypyridin-5-yl, X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, 56 mg, 0.133 mmol), methyl boronic acid (25 mg, 0.4 mmol), potassium carbonate (93 mg, 0.667 mmol) and dichloro(bis-triphenylphosphine)palladium II (19 mg, 0.027 mmol) in dioxane (3 ml) and water (3 ml) was heated to 100-110°C for 1 h. Since a TLC of the dark brown reaction mixture indicated that only 10% conversion had
25 occurred, additional aliquots of the methyl boronic acid, potassium carbonate, and palladium catalyst were added and heating was continued for another 24 h, monitoring progress by TLC. After 24 h, ~60% conversion had occurred, and so the reaction solution was diluted with more 1:1 dioxane/water (2 mL), degassed with argon, and the whole was treated with another aliquot of the methyl boronic acid, potassium carbonate, and palladium catalyst.
30 After 18 h more heating, TLC indicated <10% starting material, and so the mixture was filtered through a celite pad, the pad was washed with dioxane, and the filtrate and wash were evaporated to a dark solid. This material was triturated with chloroform, and the insolubles were collected and washed with more chloroform. Since the chloroform soluble material was found not to contain any desired product, the insoluble solid (product plus a brown tar solid)
35 was dissolved in methanol, and then filtered through silica. The methanol soluble material was evaporated to dryness (93.7 mg) and was then applied onto 2 Analtech GF, 10X 20 cm,

1,000 u plates. Development in 5:1 chloroform/MeOH gave 23.2 mg of impure product which was further purified on prep TLC plates as before to give 15 mg (28% yield) of the desired product Compound 1 shown in Table I (Compound **6**, as shown in General Method I, wherein R = 2-ethoxyxyridin-5-yl, X, Y = N, Z = Me, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, and R⁹ = CH₂OH) as an orange solid.

Method 2 (Initial Synthesis)



wherein X, Y, Z, R, R¹, R², R³ to R⁹, R¹⁸, R¹⁹, and n are as defined herein.

Compound 25, (1R,2S,3R,5R)-3-[[2-Amino-6-chloro-5-[2-phenylethynyl]-4-pyrimidinyl]amino]-5-(hydroxymethyl)-1,2-cyclopentanediol (**6**, R = phenyl).


To a stirred solution of compound **3** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, synthesized as presented above, 0.15 g, 0.375 mmol) and phenyl acetylene (0.1 mL, 0.93 mmol) in anhydrous DMF (5.0 mL) under N₂ at room temperature was added tetrakis(triphenylphosphine)palladium (0) (0.048 g, 0.037 mmol), TEA (0.1 mL, 0.75 mmol) and CuI (0.014 g, 0.075 mmol). The reaction mixture was then stirred at room temperature for 18 h, and then, MeOH (0.5 mL) was added and the whole stirred for and additional 15 min at room temperature. The solvents were removed *in vacuo*, and the obtained residue was dissolved in EtOAc (20.0 mL) and washed first with saturated solution of NaHCO₃ (3 x 20 mL) and then with saturated solution of NaCl (30 mL). The organic layer was dried over Na₂SO₄ and evaporated *in vacuo*, and the obtained residue was purified by column chromatography (silica gel, Flash), eluting with 95:5; CHCl₃ / MeOH (v/v) to give the pure product **6** (R = phenyl, X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) as a white solid (0.1 g, 78% yield).

¹H NMR (DMSO-d₆): δ 1.18 (m, 1H), 1.89 (m, 1H), 2.18 (m, 1H), 3.45 (m, 2H), 3.78 (m, 2H), 4.34 (m, 1H), 4.41 (d, 1H, J = 4.6 Hz), 4.62 (d, 1H, J = 4.9 Hz), 4.74 (t, 1H, J = 5.1 Hz), 6.60 (d, 1H, J = 8.1 Hz), 6.91 (bs, 2H), 7.41 (m, 3H), 7.54 (m, 2H)

MS m/z ($M+H$)⁺ : 375.12 (Cl pattern);

Analysis: calculated for C₁₈H₁₉ClN₄O₃: C 57.67; H 5.11; N 14.94 Found: C 57.37; H 4.96; N 14.91

Method 3 (Microwave-Assisted Synthesis)




wherein X, Y, Z, R, R¹, R², R³ to R⁹, R¹⁸, R¹⁹, and n are as defined herein.

Compound 26, (1R,2S,3R,5R)-3-[[2-Amino-6-chloro-5-[2-pyridinylethynyl]-4-pyrimidinyl]amino]-5-(hydroxymethyl)-1,2-cyclopentanediol (**6**, R = 2-pyridinyl).

To a stirred solution of compound **3** (X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH, synthesized as presented above, 0.2 g, 0.5 mmol) under N₂ at room temperature and 2-ethynylpyridine (0.13 mL, 1.25 mmol) in anhydrous DMF (5.0 mL), were added tetrakis(triphenylphosphine) palladium (0) (0.06 g, 0.05 mmol), Et₃N (0.14 mL) and CuI (0.04 g, 0.1 mmol). The reaction mixture was irradiated at 45°C for 40 min in a microwave reactor. Then, MeOH (0.5 mL) was added and the whole stirred for 15 min at room temperature. The solvents were removed *in vacuo*, and the obtained residue was purified by flash silica gel column chromatography, eluting with 96:4 CHCl₃/MeOH (v/v), giving **6** (R = 2-pyridinyl, X, Y = N, Z = Cl, R¹ = NH₂, R², R³, R⁴, R⁷, R⁸ = H, R⁵, R⁶ = OH, R⁹ = CH₂OH) as a white solid (0.11 g, 63% yield)

¹H NMR (DMSO-d₆): δ 1.22 (m, 1H), 1.90 (m, 1H), 2.21 (m, 1H), 3.43 (t, 2H, J = 4.8 Hz), 3.77 (m, 2H), 4.34 (m, 1H), 4.42 (d, 1H, J = 4.8 Hz), 4.94 (t, 1H, J = 4.5 Hz), 6.84 (d, 1H, J = 7.8 Hz), 7.02 (bs, 2H), 7.37 (m, 1H), 7.63 (m, 1H), 7.83 (m, 1H), 8.85 (m, 1H)

MS m/z ($M+H$)⁺ : 376.19 (Cl pattern)

Method 4: (2-Substituted Amino Pyrimidines 5)

5

Step 1: To trichloropyrimidine (2.9 ml, 25 mmol) in THF (25) at -15°C was added a solution of trifluoroethylamine (3.74 ml, 47.5 mmol) in THF (25 ml) over 1 hr. The reaction was warmed to 10° C over 4 hrs and stored at 8° C for approximately 48 h. Then water (150 ml) was added and extracted with EtOAc (2 x 150 ml). The combined organic layer was washed with brine (150 ml), dried (Na₂SO₄), filtered and concentrated. The residue (white solid, ~7g, **1** and **1A**) was stirred in heptane (100 ml) for 30 min at room temperature. The solid was filtered off, the filtrate was concentrated and purified by silica gel (120 g prepacked cartridge) using 0/100 to 50/50 EtOAc/hexanes to provide only **1** (1.04 g, white solid).

10

15

Step 2: To **1** (1 g, 4.07 mmol) in acetic acid (10 ml) was added a solution of ICl (1.98 g, 12.2 mmol) in glacial acetic acid (10 ml) over 30 minutes at room temperature. The mixture was stirred at room temperature, overnight. Added more ICl (2 x ~2 g) in glacial acetic acid (5 ml) for complete conversion of starting material. The reaction was quenched by dropwise addition of ice cold saturated sodium bicarbonate solution (200 ml). Then added EtOAc (150

114


ml) and the mixture was stirred overnight. The organic layer was separated and the aqueous layer was extracted with EtOAc (100 ml). The combined organic layer was washed with saturated sodium bicarbonate (200 ml), 10% aq. sodium bisulfite (2 x 200 ml), brine (150 ml), dried (Na_2SO_4), filtered and concentrated to afford **2** (1.68 g) as a white solid.

5

Step 3: The above obtained intermediate **2** (1.65 g, 4.44 mmol) in EtOH (25 ml) was treated with the carbasugar (4.40 mmol) and triethylamine (2.2 ml, 15.54 mmol). The mixture was refluxed, overnight. Then solvent was evaporated, and the residue was washed with water several times to afford **3** (1.88 g) as a solid.

10

115

Method 5 (Synthesis of 4-Me, 2-Alkylamino, 5-Arylethynyl Compounds)

5 2-Methylthio-4-methyl-6-hydroxypyrimidine **7** was prepared according to the method in (*J. Med. Chem.*, **2007**, 50, 1146-57).

2-Methylthio-4-methyl-5-iodo-6-hydroxypyrimidine **8** was prepared according to the method in *Chem. Pharm. Bull.*, **1986**, 34, 2719.

10

2-Methylthio-4-methyl-5-iodo-6-chloropyrimidine **9** was prepared according to the method in *Chem. Pharm. Bull.*, **1986**, 34, 2719.

Step 1: To a stirred mixture of the chloropyrimidine **9** (2-methylthio-4-Me-5-iodo-6-chloropyrimidine, 51.6 g, 0.172 mol) and the cyclopentylamine carbasugar (34.6 g, 0.189 mol) in EtOH (all under argon) was added diisopropylethylamine 100 mL, 0.566 mol). The resulting mixture was refluxed overnight, becoming a solution after ~1 h of heating. After
5 TLC showed that a small amount of the starting chloropyrimidine was still present, another 0.1 eq of the cyclopentylamine carbasugar (3.46 g) and more diisopropylethylamine (10 mL) were added, and again, while heating, the mixture became a solution. After refluxing overnight, the reaction was allowed to cool to room temperature and to set for ~2 h. The resulting precipitated solid was filtered and collected, washed with EtOH, and dried under
10 high vacuum to give a 79% yield of the desired adduct as an off-white solid. The filtrate was then concentrated onto silica gel and chromatographed using Flash I 65, eluting with a chloroform/methanol gradient (0 to 10% MeOH) to give more of the desired adduct **10** as a slightly impure solid product (total yield was ess. quantitative).

Step 2: Compound **10** (6.0 g, 0.0145 mol) was dissolved in DMF (80.0 mL) with stirring while a gentle stream of argon was passed through the solution for ~5 min. With a continuous stream of argon, TMS acetylene (6.2 mL, 0.043 mol) was added all at once, followed by triethylamine (8.0 mL, 0.0058 mol). After ~5 min., (Ph₃P)₄Pd (3.3 g, 0.0029 mol) (Note 1), and CuI (1.1 g, 0.0058 mol) were added in this order. The reaction mixture
20 was sealed with a rubber septum and stirred at ~45 °C with stirring under Argon for ~20 h. The resulting mixture was concentrated to dryness *in vacuo*, reconstituted in methanol-acetone (~1:1) and pre-adsorbed unto silica gel. Flash chromatography eluting with a stepwise gradient of 0 to 10% MeOH in EtOAc afforded compound **11** (4.5 g, 80.8%). [Note 1: This Pd catalyst should be a nice yellow color that is easily handled (e.g., free flowing
25 solid). A good, reliable source of this and other such catalysts is Strem Chemicals.]

Step 3: A solution of *m*CPBA (1.1 g, ~0.006 mol) was dissolved in CH₂Cl₂ (10.0 mL) and washed with brine (1 x 5.0 mL), dried (MgSO₄), filtered, and transferred to a round bottom flask. Under argon with stirring, this *m*CPBA solution is cooled to -5 °C and treated with a
30 solution of **11** (1.0 g, 0.00262 mol) in CH₂Cl₂-DMF (1:1, 6.0 mL). The bath was removed and stirring was continued overnight. The resulting mixture was concentrated to dryness *in vacuo*, reconstituted in methanol-acetone (~1:1) and pre-adsorbed unto silica gel. Flash

chromatography eluting with a stepwise gradient of 0 to 10% MeOH in EtOAc afforded compound **12** that was contaminated with a small amount of the sulfoxide **13** (0.6 g).

Step 4: To a stirred solution of 5-TMS-ethynyl **12** (200.2 mg, 0.484 mmol) in MeCN at room temperature was added Et₄NF (46 mg, 0.241 mmol), all in one portion. (The yellow solution turned dark orange upon addition.) The solution was stirred for 1h at room temp, and then a TLC showed that the reaction was complete. The solution was applied to 2 Analtech GF, 20X20 cm, 2000 μ plates, air-dried overnight, and developed 2X in 9:1 CHCl₃/MeOH. The product band was extracted with warm MeOH, and the extract evaporated to give the desired product **14** as a purple glass (83 mg, 50% yield).

Step 4a: 2-Methylsulfoxide **13** was also a suitable intermediate in the preparation of target **6** compounds by the same reaction Steps 5 and 6 shown below. Its use also required its initial deprotection to the acetylene by the same procedure described above in Step 4. Thus, a solution of **13** (1.1 g, 2.62 mmol) was treated with Et₄NF (249 mg, 1.3 mmol). Purification by flash column chromatography (25 g silica gel 230-400 mesh) eluting with CHCl₃/MeOH (0 to 20% MeOH) gave the desired product **16** as purple foam (763 mg, 89% yield).

Step 5: The solution of 2-methylsulfonylpyrimidine **14** (83.6 mg, 0.243 mmol in 2.9 mL DMF) was added to a small microwave reaction tube, and the whole was treated with the 5-bromo-2-methoxypyridine (50 uL, 0.375 mmol) followed by Et₃N (172 uL, 1.23 mmol). The solution was degassed with argon and the argon flow was continued during the addition of CuI (4.8 mg, 0.025 mmol) followed by Pd(PPh₃)₄ (14 mg, 0.012 mmol). The reaction tube was tightly capped, placed in the microwave reactor, and the mixture reacted at 90°C, 150 watts, for 10 min. After TLC of an aliquot showed that the reaction was complete, the solution was evaporated in vacuo to a light brown residue. This material in 1:1 CHCl₃/MeOH was applied to two Analtech GF (10 X 20 cm, 1000 micron) plates and then developed 3 X in 9:1 CHCl₃/MeOH. The product band was extracted with warm MeOH, and the extract was evaporated to give the desired product **15** as a white solid product (86.6 mg, 79% yield).

Step 6: To a suspension of 5-arylethynylpyrimidine **15** (83.6 mg, 0.186 mmol) in acetonitrile (10 mL) was added 3-hydroxypropylamine, 143 μ L, 1.86 mmol). The resulting cloudy reaction mixture was then placed in a preheated 95°C oil bath, and the solution cleared while refluxing. After 3 h, TLC indicated that only <5% of the sulfone starting material remained, and the heating bath was removed (3.5 h total). As the slightly yellow reaction solution cooled to RT, a solid began precipitating, and so the mixture was kept at RT for another ~2 h and then refrigerated overnight, resulting in an ivory-colored solid which was collected, washed with cold MeCN, and dried in vacuo to give the desired product. The product was then heated in boiling EtOH, filtered hot to remove insoluble material, and the filtrate concentrated to give a 2nd crop of the desired product as a slightly off-white solid. Both solid products were the desired product **6**, and the total yield was essentially quantitative.

In other embodiments, the compounds of the invention have a structural formula as depicted in Table I below and include tautomers, and pharmaceutically acceptable salts, esters, prodrugs, isomers, and solvates of such compounds and such tautomers.


Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
1		A	(DMSO- <i>d</i> ₆): δ 1.16 (dt, 1H, <i>J</i> =7.5, 13.0 Hz), 1.32 (t, 3H, <i>J</i> =7.0 Hz), 1.84-1.95 (m 1H), 2.15-2.27 (m, 1H), 2.25 (s, 3H), 3.41 (app t, 2H, <i>J</i> =4.5 Hz), 3.70-3.81 (m, 2H), 4.32 (q, 2H, <i>J</i> =7.0 Hz), 4.23-4.40 (m, 1H), 4.39-4.45 (m, 1H), 4.61-4.68 (m, 1H), 4.73-4.81 (m, 1H), 6.23 (d, 1H, <i>J</i> =7.8 Hz, NH), 6.39 (br s, 1H, Ar), 6.83 (dd, 1H, <i>J</i> =0.7, 8.6 Hz), 7.83 (dd, 1H, <i>J</i> =2.4, 8.6 Hz), 8.34 (dd, 1H, <i>J</i> =0.7, 2.3 Hz).	400.2	1, 1B

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
2		A	(DMSO- <i>d</i> ₆): δ. 1.15 (dt, 1H, <i>J</i> =7.5, 13.0 Hz), 1.31 (t, 3H, <i>J</i> =7.0 Hz), 1.83-1.95 (m 1H), 2.11-2.23 (m, 1H), 2.56 (s, 3H), 3.41 (app t, 2H, <i>J</i> =5.1 Hz), 3.72-3.82 (m, 2H), 4.24-4.37 (m, 1H), 4.31 (q, 2H, <i>J</i> =7.0 Hz), 4.44 (br d, 1H, <i>J</i> =3.5 Hz), 4.62 (br d, 1H, <i>J</i> =4.4 Hz), 4.76 (app t, 1H, <i>J</i> =4.8 Hz), 6.58 (d, 1H, <i>J</i> =8.1 Hz, NH), 6.67 (d, 1H, <i>J</i> =8.4 Hz, Ar), 6.91 (br s, 2H, NH ₂), 7.82 (d, 1H, <i>J</i> =8.4 Hz, Ar).	434.1 (Cl pattern)	1
3		A	(DMSO- <i>d</i> ₆): δ 1.17 (dt, 1H, <i>J</i> =7.5, 13.1 Hz), 1.84-1.95 (m 1H), 2.12-2.23 (m, 1H), 2.57 (s, 3H), 3.41 (app t, 2H, <i>J</i> =4.8 Hz), 3.72-3.83 (m, 2H), 3.87 (s, 3H), 4.26- 4.38 (m, 1H), 4.41-4.49 (m, 1H), 4.60-4.68 (m, 1H), 4.72-4.80 (m, 1H), 6.58 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.70 (dd, 1H, <i>J</i> =0.4, 8.6 Hz, Ar), 6.91 (br s, 2H, NH ₂), 7.84 (d, 1H, <i>J</i> =8.5 Hz, Ar).	420.2 (Cl pattern)	1



Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
4		B	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.6, 13.1 Hz), 1.84-1.96 (m 1H), 2.12-2.24 (m, 1H), 2.45 (s, 3H), 3.41 (app t, 2H, <i>J</i> =5.1 Hz), 3.73-3.84 (m, 2H), 4.28-4.39 (m, 1H), 4.38-4.53 (br s, 1H, OH), 4.55-4.70 (br s, 1H, OH), 4.72-4.84 (br s, 1H, OH), 6.65 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.99 (br s, 2H, NH ₂), 7.34 (br d, 1H, <i>J</i> =4.4 Hz), 8.37-8.42 (br s, 1H), 8.68-8.72 (br s, 1H).	390.2 (Cl pattern)	1A
5		B	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.6, 13.0 Hz), 1.82-1.96 (m 1H), 2.11-2.22 (m, 1H), 2.51 (s, 3H), 3.41 (app t, 2H, <i>J</i> =5.1 Hz), 3.73-3.84 (m, 2H), 4.28-4.39 (m, 1H), 4.43 (d, 1H, <i>J</i> =4.5 Hz), 4.63 (d, 1H, <i>J</i> =5.1 Hz), 4.79 (t, 1H, <i>J</i> =5.2 Hz), 6.69 (d, 1H, <i>J</i> =8.1 Hz, NH), 6.95 (br s, 2H, NH ₂), 7.30 (d, 1H, <i>J</i> =8.0 Hz), 7.81 (dd, 1H, <i>J</i> =2.2, 8.0 Hz), 8.60 (d, 1H, <i>J</i> =1.7 Hz).	390.2 (Cl pattern)	1 ^a

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
9		C	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.3, 13.0 Hz), 1.83-1.95 (m, 1H), 2.11-2.22 (m, 1H), 3.41 (app, 2H, <i>J</i> =5.0 Hz), 3.74-3.86 (m, 2H), 4.28-4.40 (m, 1H), 4.45 (br d, 1H, <i>J</i> =4.1 Hz, OH), 4.63 (br d, 1H, <i>J</i> =4.8 Hz, OH), 4.81 (app t, 1H, <i>J</i> =4.9 Hz, OH), 6.77 (d, 1H, <i>J</i> =8.2 Hz, NH), 7.02 (br s, 2H, NH ₂), 7.58 (dd, 1H, <i>J</i> =0.7, 8.3 Hz), 7.99 (dd, 1H, <i>J</i> =2.4, 8.3 Hz), 8.57 (dd, 1H, <i>J</i> =0.7, 2.4 Hz).	410.1 (2 Cl pattern)	1 ^a
10		B	(DMSO- <i>d</i> ₆): δ 1.15 (dt, 1H, <i>J</i> =7.8, 13.1 Hz), 1.83-1.96 (m, 1H), 2.16-2.28 (m, 1H), 3.40 (app t, 2H, <i>J</i> =5.1 Hz), 3.72-3.80 (m, 2H), 4.28-4.40 (m, 1H), 4.46 (br d, 1H, <i>J</i> =3.7 Hz, OH), 4.65 (br d, 1H, <i>J</i> =5.0 Hz, OH), 4.74 (app t, 1H, <i>J</i> =5.1 Hz, OH), 6.55 (d, 1H, <i>J</i> =7.8 Hz, NH), 7.10 (br s, 2H, NH ₂), 7.49 (dd, 1H, <i>J</i> =4.8, 7.8 Hz), 8.11 (dd, 1H, <i>J</i> =1.9, 7.7 Hz), 8.38 (dd, 1H, <i>J</i> =1.9, 4.7 Hz).	410.1 (2 Cl pattern)	1 ^a

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
11		B	(DMSO- <i>d</i> ₆): δ 1.16 (dt, 1H, <i>J</i> =7.5, 13.3 Hz), 1.82-1.95 (m, 1H), 2.10-2.22 (m, 1H), 3.40 (br t, 2H, <i>J</i> =5.0 Hz), 3.70-3.80 (m, 2H), 4.25-4.37 (m, 1H), 4.41 (br d, 1H, <i>J</i> =4.1 Hz, OH), 4.60 (br d, 1H, <i>J</i> = 4.9 Hz, OH), 4.72 (app t, 1H, <i>J</i> =5.1 Hz, OH), 6.56 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.67-6.69 (m, 1H), 6.88 (br s, 2H, NH ₂), 7.75 (app t, 1H, <i>J</i> =1.8 Hz), 8.05 (s, 1H).	365.1 (Cl pattern)	1A
12		B	(DMSO- <i>d</i> ₆): δ 1.13 (dt, 1H, <i>J</i> =8.4, 12.8 Hz), 1.86-1.97 (m, 1H), 2.14-2.26 (m, 1H), 3.40 (app t, 2H, <i>J</i> =5.3 Hz), 3.69-3.79 (m, 2H), 3.96 (s, 3H), 4.28-4.38 (m, 1H), 4.46 (br d, 1H, <i>J</i> =4.1 Hz, OH), 4.66-4.71 (m, 2H), 6.33 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.99 (br s, 2H, NH ₂), 7.04 (dd, 1H, <i>J</i> =5.1, 7.5 Hz), 7.86 (dd, 1H, <i>J</i> =1.9, 7.4 Hz), 8.16 (dd, 1H, <i>J</i> =1.9, 4.9 Hz).	406.1 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
13		B	(DMSO- <i>d</i> ₆): δ 1.17 (dt, 1H, <i>J</i> =7.4, 13.0 Hz), 1.83-1.96 (m, 1H), 2.10-2.23 (m, 1H), 3.43 (br d, 2H, <i>J</i> =4.9 Hz), 3.72-3.81 (m, 2H), 4.28-4.40 (m, 1H), 4.44 (br d, 1H, <i>J</i> =3.6 Hz, OH), 4.58 (br d, 1H, <i>J</i> =4.2 Hz, OH), 5.0-5.13 (br s, 1H, OH), 6.90 (d, 1H, <i>J</i> =8.1 Hz, NH), 6.99 (s, 2H), 7.14 (br s, 2H), 12.7 (br s, 1H).	365.2 (Cl pattern)	1
14		B	(DMSO- <i>d</i> ₆): δ 1.13 (dt, 1H, <i>J</i> =7.7, 13.0 Hz), 1.32 (d, 6H, <i>J</i> =6.2 Hz), 1.85-1.97 (m, 1H), 2.11-2.25 (m, 1H), 3.40 (app t, 2H, <i>J</i> =5.2 Hz), 3.71-3.81 (m, 2H), 4.28-4.38 (m, 1H), 4.45 (br d, 1H, <i>J</i> =3.4 Hz, OH), 4.65 (br d, 1H, <i>J</i> =5.0 Hz), 4.74 (app t, 1H, 2 <i>J</i> =5.1 Hz), 5.33 (quint, 1H, <i>J</i> =6.2 Hz), 6.38 (d, 1H, <i>J</i> =8.1 Hz, NH), 6.96 (br s, 2H, NH ₂), 6.98 (dd, 1H, <i>J</i> =5.0, 7.4 Hz), 7.88 (dd, 1H, <i>J</i> =2.0, 7.5 Hz), 8.12 (dd, 1H, <i>J</i> =2.0, 4.9 Hz).	434.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
15		B	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.8, 13.0 Hz), 1.85-1.96 (m, 1H), 2.13-2.25 (m, 1H), 2.64 (s, 3H), 3.41 (app t, 2H, <i>J</i> =5.1 Hz), 3.71-3.84 (m, 2H), 4.28-4.39 (m, 1H), 4.43 (br d, 1H, <i>J</i> =4.3 Hz, OH), 4.61 (br d, 1H, <i>J</i> =4.9 Hz), 4.76 (app t, 1H, <i>J</i> =5.0 Hz), 6.64 (d, 1H, <i>J</i> =7.9 Hz, NH), 6.98 (br s, 2H, NH ₂), 7.27 (dd, 1H, <i>J</i> =5.0, 7.8 Hz), 7.93 (dd, 1H, <i>J</i> =1.8, 7.8 Hz), 8.40 (dd, 1H, <i>J</i> =1.8, 4.9 Hz).	390.2 (Cl pattern)	2
16		A	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.4, 13.2 Hz), 1.83-1.95 (m, 1H), 2.12-2.25 (m, 1H), 2.45 (s, 3H), 2.60 (s, 3H), 3.41 (app t, 2H, <i>J</i> =5.2 Hz), 3.71-3.84 (m, 2H), 4.28-4.39 (m, 1H), 4.44 (br d, 1H, <i>J</i> =4.2 Hz, OH), 4.63 (br d, 1H, <i>J</i> =5.1 Hz), 4.77 (app t, 1H, <i>J</i> =5.0 Hz), 6.61 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.97 (br s, 2H, NH ₂), 7.13 (d, 1H, <i>J</i> =7.9 Hz), 7.81 (d, 1H, <i>J</i> =7.9 Hz).	404.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
17		B	(DMSO- <i>d</i> ₆): δ 1.17 (dt, 1H, <i>J</i> =7.7, 13.0 Hz), 1.24 (t, 3H, <i>J</i> =7.6 Hz), 1.84-1.96 (m 1H), 2.12-2.25 (m, 1H), 2.47 (s, 3H), 2.95 (q, 2H, <i>J</i> =7.5 Hz), 3.41 (app t, 2H, <i>J</i> =5.1 Hz), 3.72-3.84 (m, 2H), 4.27-4.39 (m, 1H), 4.44 (br d, 1H, <i>J</i> =3.9 Hz, OH), 4.63 (br d, 1H, <i>J</i> =4.7 Hz), 4.76 (t, 1H, <i>J</i> =5.0 Hz), 6.57 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.96 (br s, 2H, NH ₂), 7.13 (d, 1H, <i>J</i> =8.0 Hz), 7.83 (d, 1H, <i>J</i> =7.9 Hz).	418.2 (Cl pattern)	1
18		A	(DMSO- <i>d</i> ₆): δ 1.20 (dt, 1H, <i>J</i> =7.5, 12.9 Hz), 1.85-1.96 (m 1H), 2.15-2.26 (m, 1H), 2.65 (s, 3H), 3.43 (app t, 2H, <i>J</i> =5.0 Hz), 3.75-3.85 (m, 2H), 4.29-4.40 (m, 1H), 4.45 (br d, 1H, <i>J</i> =4.2 Hz, OH), 4.63 (br d, 1H, <i>J</i> =4.8 Hz), 4.80 (app t, 1H, <i>J</i> =5.0 Hz), 6.68 (d, 1H, <i>J</i> =7.9 Hz, NH), 6.99 (br s, 2H, NH ₂), 7.17 (dd, 1H, <i>J</i> =3.7, 5.0 Hz), 7.67 (dd, 1H, <i>J</i> =1.0, 5.1 Hz), 7.78-7.86 (m, 2H), 7.96 (d, 1H, <i>J</i> =8.2 Hz).	472.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
19		A	(DMSO- <i>d</i> ₆): δ 1.14-35 (m, 1H), 1.85-1.96 (m 1H), 2.13-2.35 (m, 1H), 2.72-2.85 (m, 3H), 3.39-3.47 (m, 2H), 3.74-3.84 (m, 2H), 3.88 (s, 3H), 4.22-4.35 (m, 1H), 4.43 (br d, 1H, <i>J</i> =4.6 Hz, OH), 4.58-4.68 (m, 1H), 4.76-4.84 (m, 1H), 6.58-6.78 (m, 1H, NH), 6.88 (d, 1H, <i>J</i> =8.6 Hz), 7.25-7.45 (m, 1H, NH), 7.84 (dd, 1H, <i>J</i> =2.2, 8.8 Hz), 8.36 (d, 1H, <i>J</i> =1.8 Hz).	420.2 (Cl pattern)	4
20		A	(DMSO- <i>d</i> ₆): δ 1.20 (m, 1H), 1.87 (m, 1H), 2.17 (m, 1H), 3.41 (m, 2H), 3.78 (m, 2H), 3.88 (s, 3H), 4.32 (m, 1H), 4.41 (d, 1H, <i>J</i> =4.6 Hz), 4.60 (d, 1H, <i>J</i> =4.9 Hz), 4.77 (m, 1H), 6.64 (d, 1H, <i>J</i> =8 Hz), 6.89 (m, 3H), 7.83 (dd, 1H, <i>J</i> =2.2Hz, 8.7 Hz), 8.37 (m, 1H).	406.2 (Cl pattern)	1

Table I

Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
21		A	(DMSO- <i>d</i> ₆): δ 1.19 (m, 1H), 1.32 (t, 3H, <i>J</i> =6.9 Hz), 1.89 (m, 1H), 2.18 (m, 1H), 3.41 (m, 2H), 3.76 (m, 2H), 4.30 (q, 3H, <i>J</i> =7.0, 14.0 Hz), 4.41 (d, 1H, <i>J</i> =4.4 Hz), 4.63 (d, 1H, <i>J</i> =5.0 Hz), 4.78 (t, 1H, <i>J</i> =5.0 Hz), 6.74 (d, 1H, <i>J</i> =6.0 Hz, NH), 6.84 (dd, 1H, <i>J</i> =0.4, 8.6 Hz, Ar), 6.92 (br s, 2H, NH ₂), 7.44 (d, 1H, Ar), 7.83 (dd, 1H, <i>J</i> =2.4, 8.7 Hz, Ar), 8.34 (dd, 1H, <i>J</i> =0.6, 2.3 Hz).	420.2 (Cl pattern)	1
22		A	(DMSO- <i>d</i> ₆): δ 1.11 (m, 1H), 1.90 (m, 1H), 2.20 (m, 1H), 3.39 (m, 2H), 3.73 (m, 2H), 4.31 (m, 1H), 4.47 (d, 1H, <i>J</i> =3.8 Hz), 4.68 (m, 1H), 6.30 (d, 1H, <i>J</i> =7.9 Hz, NH), 6.44 (d, 1H, <i>J</i> =8.1 Hz, Ar), 6.92 (br s, 2H, NH ₂), 7.77 (d, 1H, <i>J</i> =8.2 Hz).	436.1 (Cl pattern)	1
23		C	(DMSO- <i>d</i> ₆): δ 1.25 (m, 1H), 1.32 (t, 3H, <i>J</i> =6.9 Hz), 1.93 (m, 1H), 2.23 (m, 1H), 3.10 (s, 6H), 3.70 (q, 1H, <i>J</i> =7.0, 12.1 Hz), 3.84 (q, 1H, <i>J</i> =5.2, 10.3 Hz), 4.24 (m, 1H), 4.34 (q, 2H, <i>J</i> =7.0, 14.2 Hz), 4.43 (d, 1H,	448.2 (Cl pattern)	4



Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			<i>J</i> =5.2 Hz), 4.63 (d, 1H, <i>J</i> =5.2 Hz), 4.81 (t, 1H, <i>J</i> =5.0 Hz), 6.71 (d, 1H, <i>J</i> =7.5 Hz, NH), 6.84 (dd, 1H, <i>J</i> =0.7, 8.6 Hz, Ar), 7.83 (dd, 1H, <i>J</i> =2.3, 8.6 Hz, Ar), 8.35 (dd, 1H, <i>J</i> =0.55, 2.3 Hz).		
24		C	(DMSO- <i>d</i> ₆): δ 1.25 (m, 1H), 1.93 (m, 1H), 2.23 (m, 1H), 3.10 (br s, 6H), 3.42 (t, 2H, <i>J</i> =4.6 Hz), 3.77 (q, 1H, <i>J</i> =4.7, 9.9 Hz), 3.84 (q, 1H, <i>J</i> =5.2, 10.4 Hz), 4.24 (m, 1H), 4.44 (d, 1H, <i>J</i> =5.3 Hz), 4.64 (d, 1H, <i>J</i> =5.2 Hz), 4.82 (t, 1H, <i>J</i> =4.90 Hz), 6.73 (d, 1H, <i>J</i> =7.6 Hz, NH), 6.88 (d, 1H, <i>J</i> =8.7 Hz, Ar), 7.84 (dd, 1H, <i>J</i> =2.3, 8.6 Hz, Ar), 8.37 (m, 1H).	434.2 (Cl pattern)	4
25		B	(DMSO- <i>d</i> ₆): δ 1.18 (m, 1H), 1.89 (m, 1H), 2.18 (m, 1H), 3.45 (m, 2H), 3.78 (m, 2H), 4.34 (m, 1H), 4.41 (d, 1H, <i>J</i> =4.6 Hz), 4.62 (d, 1H, <i>J</i> =4.9 Hz), 4.74 (t, 1H, <i>J</i> =5.1 Hz), 6.60 (d, 1H, <i>J</i> =8.1 Hz), 6.91 (bs, 2H), 7.41 (m, 3H), 7.54 (m, 2H)	375.1 (Cl pattern)	2

Table I




Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
26		B	(DMSO- <i>d</i> ₆) δ 1.22 (m, 1H), 1.90 (m, 1H), 2.21 (m, 1H), 3.43 (t, 2H, <i>J</i> =4.8 Hz), 3.77 (m, 2H), 4.34 (m, 1H), 4.42 (d, 1H, <i>J</i> =4.8 Hz), 4.94 (t, 1H, <i>J</i> =4.5 Hz), 6.84 (d, 1H, <i>J</i> =7.8 Hz), 7.02 (bs, 2H), 7.37 (m, 1H), 7.63 (m, 1H), 7.83 (m, 1H), 8.85 (m, 1H)	376.2 (Cl pattern)	3
27		B	(CD ₃ OD) δ 1.28–1.40 (m, 1H), 2.05–2.18 (m, 1H), 2.38–2.48 (m, 1H), 3.58–3.69 (m, 2H), 3.89 (t, 1H, <i>J</i> =5.3 Hz), 3.97 (t, 1H, <i>J</i> =5.3 Hz), 4.35–4.43 (m (app q), 1H), 7.37 (app d, 2H, <i>J</i> =8.7 Hz), 7.51 (app d, 2H, <i>J</i> =8.7 Hz).	409.1 (2 Cl pattern)	2
28		B	(CD ₃ OD) δ 1.35 (dt, 1H, <i>J</i> =7.3, 14.3 Hz), 2.05–2.16 (m, 1H), 2.37–2.47 (m, 1H), 3.56–3.70 (m, 2H), 3.88 (t, 1H, <i>J</i> =5.3 Hz), 3.96 (t, 1H, <i>J</i> =5.2 Hz), 4.35–4.42 (m, 1H), 7.21 (dd, 1H, <i>J</i> =5.06, 1.1 Hz), 7.42 (dd, 1H, <i>J</i> =5.1, 3.0 Hz), 7.65 (dd, 1H, <i>J</i> =1.1, 3.0 Hz)	381.2 (Cl pattern)	2

Table I




Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
29		B	(CD ₃ OD) δ 1.28–1.40 (m, 1H), 2.07–2.15 (m, 1H), 2.38–2.49 (m, 1H), 2.46 (s, 2H), 3.55–3.67 (m, 2H), 3.80 (s, 3H), 3.87 (t, 1H, <i>J</i> =5.6 Hz), 3.95 (t, 1H, <i>J</i> =5.2 Hz), 4.32–4.41 (m, 1H), 6.75 (dd, 1H, <i>J</i> =8.5, 2.6 Hz), 6.82 (app d, 1H, <i>J</i> =2.6 Hz), 7.44 (d, 1H, <i>J</i> =8.5 Hz).	419.2 (Cl pattern)	2
30		C	(CD ₃ OD) δ 1.28–1.41 (m, 1H), 2.07–2.15 (m, 1H), 2.36–2.47 (m, 1H), 3.54–3.67 (m, 2H), 3.76 (s, 3H), 3.88 (t, 1H, <i>J</i> =5.4 Hz), 3.96 (t, 1H, <i>J</i> =5.2 Hz), 4.37–4.44 (m, 1H), 7.27 (br s, 1H), 7.71 (br s, 1H).	379.2 (Cl pattern)	2
31		A	(DMSO- <i>d</i> ₆): δ 1.18–1.20 (m, 1H), 1.52–1.61 (m, 2H, -CH ₂ pip.), 1.68–1.78 (m, 2H, -CH ₂ pip.), 1.95–2.09 (m, 2H, -CH ₂ pip.), 1.80–1.91 (m, 1H), 2.10–2.18 (m, 1H), 2.91–2.99 (m, 1H, -CH ₂ pip), 3.25–3.99 (m, 1H, -CH ₂ pip) 3.37–3.44 (m, 2H), 3.72 (q, 1H, <i>J</i> =5.1 Hz), 3.78 (t, 1H, <i>J</i> =5.1 Hz), 4.29–4.36 (m, 1H), 4.41 (d, 1H, <i>J</i> =6 Hz), 4.55 (d, 1H, <i>J</i> =5.7 Hz), 4.69 (t,	464.2 (Cl pattern)	1




Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			1H, <i>J</i> =4.8 Hz), 6.67 (d, 1H, <i>J</i> =8.4 Hz), 6.97 (bs, 2H), 7.14-7.12 (m, 3H), 7.29 (d, 2H, <i>J</i> =6 Hz).		
32		A	(DMSO- <i>d</i> ₆): δ 1.19-1.23 (m, 1H), 1.83-1.91 (m, 1H), 2.07-2.14 (m, 1H), 3.33-3.41 (m, 2H), 3.71 (q, 2H, <i>J</i> =5.0 Hz), 3.80 (s, 3H), 3.62-3.80 (m, 1H), 4.40 (d, 1H, <i>J</i> =4 Hz), 4.45-4.50 (m, 1H), 4.58 (t, 1H, <i>J</i> =5.1 Hz), 4.61 (d, 1H, <i>J</i> =6 Hz), 7.05 (bs, 2H), 7.05 (bs, 2H), 7.08 (m, 1H), 7.67 (m, 2H).	455.1 (Cl pattern)	1
33		A	(DMSO- <i>d</i> ₆): δ 1.13-1.22 (m, 1H), 1.73-1.81 (m, 1H), 2.12-2.19 (m, 2H), 3.38-3.42 (m, 2H), 3.70-3.75 (m, 2H), 3.78 (s, 3H), 4.25-4.33 (m, 1H), 4.41 (d, 1H, <i>J</i> =4.2 Hz), 4.61 (d, 1H, <i>J</i> =5.1 Hz), 4.73 (t, 1H, <i>J</i> =5.1 Hz), 6.54 (d, 1H, <i>J</i> =8.1 Hz), 6.85 (bs, 2H), 6.97 (d, 1H, <i>J</i> =8.7 Hz), 7.46-7.49 (m, 2H).	405.8 (Cl pattern)	2
34		A	(DMSO- <i>d</i> ₆): δ 1.18-1.22 (m, 1H), 1.84-1.90 (m, 1H), 2.10-2.20 (m, 1H), 2.37-3.43 (m, 2H), 3.71-3.83 (m, 2H), 3.31-3.37 (m, 2H), 4.63	458.1 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			(d, 1H, $J=5.4$ Hz), 4.73 (t, 1H, $J=5.1$ Hz), 6.74 (d, 1H, $J=8.1$ Hz), 7.0 (bs, 2H), 7.29-7.33 (m, 1H), 7.38 (d, 1H, $J=3.9$ Hz), 7.78 (d, 1H, $J=3.9$ Hz), 7.96 (m, 1H), 8.52-8.54 (m, 1H).		
35		A	(DMSO- d_6): δ 1.10-1.18 (m, 1H), 1.83-1.86 (m, 1H), 2.09-2.19 (m, 1H), 2.71 (s, 3H), 3.34-3.43 (m, 2H), 3.70-3.78 (m, 2H), 4.29-4.37 (m, 2H), 4.41 (d, 1H, $J = 4.2$ Hz), 4.60 (d, 1H, $J = 5.1$ Hz), 4.68 (t, 1H, $J = 4.8$ Hz), 6.73-6.76 (m, 1H), 7.04 (bs, 2H), 7.35-7.40 (m, 1H), 7.43-7.48 (m, 2H), 8.20-8.24 (m, 2H).	472.1 (Cl pattern)	1
36		A	(DMSO- d_6): δ 1.10-1.20 (m, 1H), 1.82-1.89 (m, 1H), 2.08-2.18 (m, 1H), 3.43-3.37 (m, 2H), 3.70-3.80 (m, 2H), 4.27-4.36 (m, 2H), 4.40 (d, 1H, $J=4.5$ Hz), 4.60 (d, 1H, $J=5.4$ Hz), 4.69 (t, 1H, $J=4.1$ Hz), 6.67-6.68 (m, 1H), 7.04 (bs, 2H), 7.09-7.12 (m, 1H), 7.36-7.38 (m, 1H), 7.61-7.64 (m, 1H).	381.2 (Cl pattern)	1

Table I




Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
37		B	(DMSO- <i>d</i> ₆): δ 1.11-1.23 (m, 1H), 1.82-1.91 (m, 1H), 2.09-2.18 (m, 1H), 3.39 (t, 2H, <i>J</i> =5.1 Hz), 3.71-3.82 (m, 2H), 4.28-4.35 (m, 1H), 4.41 (d, 1H, <i>J</i> =4.5 Hz), 4.56 (d, 1H, <i>J</i> =5.1 Hz), 4.72 (t, 1H, <i>J</i> =4.8 Hz), 6.78 (d, 1H, <i>J</i> =4.1 Hz), 7.02 (bs, 2H), 8.12 (s, 1H), 9.12 (s, 1H).	382.2 (Cl pattern)	1
38		B	(DMSO- <i>d</i> ₆): δ 1.15-1.21 (m, 1H), 1.85-1.91 (m, 1H), 2.15-2.21 (m, 1H), 3.41 (t, 2H, <i>J</i> =4.8 Hz), 3.73-3.83 (m, 2H), 3.99-4.06 (m, 1H), 4.42 (d, 1H, <i>J</i> =4.5 Hz), 4.61 (d, 1H, <i>J</i> =5.1 Hz), 4.78 (t, 1H, <i>J</i> =5.1 Hz), 6.72 (d, 1H, <i>J</i> =8.1 Hz), 6.97 (bs, 2H), 7.43-7.46 (m, 1H), 7.90-7.94 (m, 1H), 8.53 (bs, 1H), 8.72 (bs, 1H).	376.1 (Cl pattern)	2
39		B	(DMSO- <i>d</i> ₆): δ 1.12-1.21 (m, 1H), 1.82-1.91 (m, 1H), 2.10-2.21 (m, 1H), 2.69 (s, 3H), 3.40 (t, 2H, <i>J</i> =5.7 Hz), 3.71-3.83 (m, 2H), 4.30-4.38 (m, 1H), 4.42 (d, 1H, <i>J</i> =4.1 Hz), 4.62 (d, 1H, <i>J</i> =5.4 Hz), 4.72 (t, 1H, <i>J</i> =5.1 Hz), 6.71 (d, 1H, <i>J</i> =8.1 Hz), 6.97	478.0 (Cl pattern)	1

Table I




Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			(bs, 2H), 7.33 (d, 1H, $J=3.6$ Hz), 7.51 (t, 1H, $J=3.6$ Hz), 7.89 (s, 1H).		
40		B	(DMSO- d_6): δ 1.18-1.21 (m, 1H), 1.85-1.92 (m, 1H), 2.08-2.20 (m, 1H), 3.38-3.43 (m, 2H), 3.70-3.82 (m, 2H), 4.30-4.38 (m, 1H), 4.42 (d, 1H, $J=4.1$ Hz), 4.59-4.61 (m, 1H), 4.80 (t, 1H, $J=5.1$ Hz), 6.96 (bs, 1H), 7.18 (bs, 1H), 7.52 (d, 1H, $J=4.2$ Hz), 8.08-8.10 (m, 1H).	477.1 (Cl pattern)	1
41		B	(DMSO- d_6): δ 1.10-1.25 (m, 1H), 1.82-1.92 (m, 1H), 2.08-2.18 (m, 1H), 3.37-3.40 (m, 2H), 3.70-3.82 (m, 2H), 4.30-4.48 (m, 1H), 4.41 (d, 1H, $J=4.5$ Hz), 4.62 (d, 1H, $J=5.4$ Hz), 4.71 (t, 1H, $J=5.1$ Hz), 6.95 (d, 1H, $J=8.1$ Hz), 7.10 (bs, 2H), 7.86 (d, 1H, $J=3.3$ Hz), 7.91 (d, 1H, $J=3.3$ Hz).	382.1 (Cl pattern)	1
42		B	(DMSO- d_6): δ 1.12-1.22 (m, 1H), 1.82-1.91 (m, 1H), 2.08-2.19 (m, 1H), 3.40 (t, 1H, $J=5.4$ Hz), 3.72-3.83 (m, 2H), 4.31-4.36 (m, 1H), 4.41 (d, 1H, $J=4.5$ Hz), 4.62 (d, 1H, $J=5.4$ Hz), 4.73 (t,	368.1 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			1H, <i>J</i> =5.1 Hz), 6.74 (d, 1H, <i>J</i> =8.1 Hz), 7.00 (bs, 2H), 7.43 (d, 1H, <i>J</i> =3.9 Hz), 7.44-7.49 (m, 1H), 7.64 (d, 1H, <i>J</i> =3.3 Hz), 8.06-8.10 (m, 1H), 8.52-8.54 (m, 1H), 8.92 (bs, 1H).		
43		B	(DMSO- <i>d</i> ₆): δ 1.10-1.22 (m, 1H), 1.82-1.91 (m, 1H), 2.08-2.19 (m, 1H), 3.41-3.51 (m, 1H), 3.70-3.85 (m, 2H), 3.89-4.41 (m, 1H), 4.41 (d, 1H, <i>J</i> =4.5 Hz), 4.62 (d, 1H, <i>J</i> =5.4 Hz), 4.75 (t, 1H, <i>J</i> =5.1 Hz), 6.80 (d, 1H, <i>J</i> =8.1 Hz), 6.96 (d, 1H, <i>J</i> =2.1 Hz), 7.03 (bs, 2H), 7.43 (d, 1H, <i>J</i> =3.9 Hz), 7.68 (d, 1H, <i>J</i> =3.9 Hz).	448.1 (Cl pattern)	1
44		B	(DMSO- <i>d</i> ₆): δ 1.15-1.25 (m, 1H), 1.85-1.95 (m, 1H), 2.08-2.19 (m, 1H), 3.42 (t, 2H, <i>J</i> =5.1 Hz), 3.74-3.84 (m, 2H), 4.29-4.39 (m, 1H), 4.42 (d, 1H, <i>J</i> =4.5 Hz), 4.63 (d, 1H, <i>J</i> =5.4 Hz), 4.77 (t, 1H, <i>J</i> =5.1 Hz), 6.64 (d, 1H, <i>J</i> =8.1 Hz), 6.93 (bs, 2H), 7.35-7.40 (m, 1H), 7.45-7.52 (m, 2H), 7.60-7.66 (m, 2H), 7.70-7.74 (m, 4H).	451.2 (Cl pattern)	2

Table I

Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
45		B	(DMSO- <i>d</i> ₆): δ 1.15-1.25 (m, 1H), 1.83-1.95 (m, 1H), 2.10-2.22 (m, 1H), 3.42 (t, 2H, <i>J</i> =5.1 Hz), 3.74-3.83 (m, 2H), 4.28-4.37 (m, 1H), 4.45 (d, 1H, <i>J</i> =4.5 Hz), 4.62 (d, 1H, <i>J</i> =5.4 Hz), 4.84 (t, 1H, <i>J</i> =5.1 Hz), 6.80 (d, 1H, <i>J</i> =8.1 Hz), 7.06 (bs, 2H), 8.95 (bs, 2H), 9.14 (bs, 1H).	377.2 (Cl pattern)	2
46		C	(DMSO- <i>d</i> ₆): δ 1.10-1.21 (m, 1H), 1.82-1.91 (m, 1H), 2.18-2.28 (m, 1H), 2.39 (s, 3H), 3.38-3.44 (m, 2H), 3.70-3.85 (m, 2H), 4.32-4.39 (m, 2H), 4.61 (d, 1H, <i>J</i> =5.4 Hz), 4.71 (t, 1H, <i>J</i> =5.1 Hz), 6.91 (d, 1H, <i>J</i> =8.1 Hz), 7.09 (bs, 2H), 7.41 (bs, 1H).	396.2 (Cl pattern)	1
47		C	(DMSO- <i>d</i> ₆): δ 1.03-1.31 (m, 1H), 1.80-1.93 (m, 1H), 2.09-2.18 (m, 1H), 3.37-3.41 (m, 2H), 3.70-3.83 (m, 2H), 4.30-4.42 (m, 2H), 4.61 (d, 1H, <i>J</i> =5.4 Hz), 4.68-4.71 (m, 3H), 5.68 (t, 1H, <i>J</i> =5.7 Hz), 6.91 (d, 1H, <i>J</i> =8.1 Hz), 7.09 (bs, 2H), 7.73 (bs, 1H).	412.2 (Cl pattern)	1

Table I




Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
48		C	(DMSO- <i>d</i> ₆): δ 1.18-1.27 (m, 1H), 1.86-1.92 (m, 1H), 2.15-2.26 (m, 1H), 3.39-3.42 (m, 2H), 3.71-3.83 (m, 2H), 4.19-4.28 (m, 1H), 4.40 (d, 1H, <i>J</i> =5.4 Hz), 4.60 (m, 1H), 4.71 (t, 1H, <i>J</i> =5.7 Hz), 6.73 (d, 1H, <i>J</i> =7.5 Hz), 7.09-7.12 (bs, 1H), 7.37-7.38 (m, 1H), 7.61-7.63 (m, 1H).	409.1 (Cl pattern)	4
49		B	(CD ₃ OD) δ 1.53 (dt, 1H, <i>J</i> =8.9, 13.2 Hz), 2.07-2.19 (m, 1H), 2.37 (dt, 1H, <i>J</i> =8.6, 13.2 Hz), 3.60-3.71 (m, 2H), 3.96 (t, 1H, <i>J</i> =5.1 Hz), 4.06-4.13 (m, 2H), 4.40-4.50 (m (app dt), 1H), 7.39 (dt, 1H, <i>J</i> =7.6, 1.4 Hz), 7.49 (dt, 1H, <i>J</i> =1.3, 7.4 Hz), 7.56 (dd, 1H, <i>J</i> =1.3, 7.7 Hz), 7.67 (dd, 1H, <i>J</i> =7.6, 1.0 Hz).	418.2 (Cl pattern).	1
50		B	(DMSO- <i>d</i> ₆): δ 1.14-1.31 (m, 1H), 1.34 (d, 3H, <i>J</i> =4.62 Hz), 1.36 (d, 3H, <i>J</i> =4.7 Hz), 1.87-2.04 (m, 1H), 2.05-2.32 (m, 1H), 2.75-2.85 (m, 3H), 3.30-3.48 (m, 2H), 3.70-3.79 (m, 1H), 3.85-3.99 (m, 1H), 4.40-4.55 (m, 2H), 4.59 (t, 1H, =5.1 Hz),	475.3 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			4.64 (d, 1H, <i>J</i> =5.9 Hz), 5.21 (septet, 1H, <i>J</i> =6.2 Hz), 7.17-7.32 (m, 1H), 7.40-7.56 (m, 2H), 7.56-7.70 (m, 2H), 8.03 (d, 1H, <i>J</i> =7.3 Hz).		
51		**	(DMSO- <i>d</i> ₆): δ 1.22 (dt, 1H, <i>J</i> =9.7, 12.7 Hz), 1.35 (t, 3H, <i>J</i> = 7.1 Hz), 1.87-1.99 (m, 1H), 2.07-2.17 (m, 1H), 3.32-3.45 (m, 2H), 3.71-3.78 (m, 1H), 3.86-3.95 (m, 1H), 4.38 (q, 2H, <i>J</i> =7.1 Hz), 4.42-4.56 (m, 2H), 4.60 (t, 1H, <i>J</i> =5.2 Hz, D ₂ O exchangeable), 4.67 (d, 1H, <i>J</i> =6.2 Hz, D ₂ O exchangeable), 7.07 (br s, 2H, D ₂ O exchangeable), 7.23 (d, 1H, <i>J</i> =8.5 Hz), 7.49 (dt, 1H, <i>J</i> =1.4, 7.5 Hz), 7.55-7.70 (m, 2H), 8.06 (dd, 1H, <i>J</i> = 1.1, 7.9 Hz).	447.2 (Cl pattern)	1
52		A	((DMSO- <i>d</i> ₆): δ 1.23 (d, 1H, <i>J</i> =9.8, 12.9 Hz), 1.35 (t, 3H, <i>J</i> =7.0 Hz), 1.87-2.00 (m, 1H), 2.12 (dt, 1H, <i>J</i> =8.2, 12.8 Hz), 3.32-3.45 (m, 2H), 3.74 (dd, 1H, <i>J</i> =1.3, 3.7 Hz), 3.89 (dd, 1H, <i>J</i> =5.4, 8.1 Hz), 4.38 (q, 2H, <i>J</i> =7.1 Hz), 4.42-4.55 (m, 1H), 7.25	447.1 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			(d, 1H, $J=8.5$ Hz), 7.49 (app dt, 1H, $J=1.1, 8.2$ Hz), 7.58-7.70 (m, 2H), 8.06 (d, 1H, $J=7.9$ Hz).		
53		B	(DMSO- d_6): δ 1.16-1.29 (m, 1H), 1.34 (d, 3H, $J=5.6$ Hz), 1.36 (d, 3H, $J=5.6$ Hz), 1.86-1.99 (m, 1H), 2.11 (dt, 1H, $J=7.8, 12.6$ Hz), 3.34-3.48 (m, 2H), 3.69-3.78 (m, 1H), 3.86-3.92 (m, 1H), 4.44 (d, 1H, $J=4.0$ Hz), 4.45-4.57 (m, 1H), 4.60 (app t, 1H, $J=5.0$ Hz, D ₂ O exchangeable), 4.66 (d, 1H, $J=6.1$ Hz, D ₂ O exchangeable), 5.21 (quint, 1H, $J=6.2$ Hz), 7.06 (br s, 2H, D ₂ O exchangeable), 7.23 (d, 1H, $J=8.5$ Hz), 7.51 (dt, 1H, $J=1.1, 7.8$ Hz), 7.56-7.70 (m, 2H), 8.03 (d, 1H, $J=7.8$ Hz).	461.2 (Cl pattern)	1
54		A	(DMSO- d_6): δ 1.22 (dt, 1H, $J=9.7, 12.9$ Hz), 1.35 (t, 3H, $J=7.1$ Hz), 1.86-1.99 (m, 1H), 2.13 (dt, 1H, $J=8.4, 12.5$ Hz), 3.39-3.45 (m, 2H), 3.72-3.79 (m, 1H), 3.86-3.94 (m, 1H), 4.40 (ap q, 2H, $J=7.1$ Hz), 4.43-	448.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			4.52 (m, 2H), 4.61 (t, 1H, <i>J</i> =5.1 Hz), 4.66 (d, 1H, <i>J</i> =5.8 Hz, D ₂ O exchangeable), 7.05 (d, 1H, <i>J</i> =8.5 Hz), 7.12 (br s, 2H, D ₂ O exchangeable), 7.67 (dd, 1H, <i>J</i> =4.6, 8.0 Hz), 8.02 (dd, 1H, <i>J</i> =1.6, 8.0 Hz), 8.63 (dd, 1H, <i>J</i> =4.6, 1.6 Hz).		
55		A	(DMSO- <i>d</i> ₆): δ 1.23 (dt, 1H, <i>J</i> =9.0, 12.9 Hz), 1.87–1.98 (m, 1H), 2.15 (dt, 1H, <i>J</i> =8.2, 12.5 Hz), 3.35–3.44 (m, 2H), 3.73–3.79 (m, 1H), 3.88–3.97 (m, 1H), 3.93 (s, 3H), 4.43 (d, 1H, <i>J</i> =4.1 Hz), 4.42–4.49 (m, 1H), 4.61 (t, 1H, <i>J</i> =5.2 Hz), 4.65 (d, 1H, <i>J</i> =5.9 Hz), 7.04 (d, 1H, <i>J</i> =8.5 Hz), 7.12 (br s, 2H, D ₂ O exchangeable), 7.67 (dd, 1H, <i>J</i> =8.0, 4.6 Hz), 8.03 (dd, 1H, <i>J</i> =8.0, 1.6 Hz), 8.63 (dd, 1H, <i>J</i> =4.6, 1.6 Hz).	434.1 (Cl pattern)	1
56		B	(DMSO- <i>d</i> ₆): δ 1.12 (t, 3H, <i>J</i> = 7.2 Hz), 1.24 (dt, 1H, <i>J</i> =9.9, 13.1 Hz), 1.85–1.97 (m, 1H), 2.09 (dt, 1H, <i>J</i> =8.0, 12.6 Hz), 3.26–3.49 (m, 2H), 3.70–3.76 (m, 1H), 3.97–4.05 (m, 1H),	446.2 (Cl pattern)	1

Table I


Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
			4.39 (d, 1H, <i>J</i> =3.9 Hz), 4.47 (app quint, 1H, <i>J</i> =8.4 Hz), 4.57 (app t, 1H, <i>J</i> =5.2 Hz), 4.59 (d, 1H, <i>J</i> =5.9 Hz), 6.95 (br s, 2H, D ₂ O exchangeable), 7.37 (d, 1H, <i>J</i> =8.5 Hz), 7.40–7.48 (m, 1H), 7.51 (d, 1H, <i>J</i> =4.0 Hz), 7.64 (d, 1H, <i>J</i> =7.6 Hz), 8.65 (app t, 1H, <i>J</i> =5.5 Hz).		
57		B	(DMSO- <i>d</i> ₆): δ 1.25 (dt, 1H, <i>J</i> =9.7, 12.6 Hz), 1.84–1.96 (m, 1H), 2.10 (dt, 1H, <i>J</i> =8.3, 12.7 Hz), 2.80 (d, 3H, <i>J</i> =4.6 Hz), 3.30–3.47 (m, 2H), 3.70–3.78 (m, 1H), 4.05 (dt, 1H, <i>J</i> =5.8, 8.0 Hz), 4.39 (d, 1H, <i>J</i> =4.1 Hz), 4.48 (app quint, 1H, <i>J</i> =8.4 Hz), 4.57 (app t, 1H, <i>J</i> =5.2 Hz), 4.61 (d, 1H, <i>J</i> =5.9 Hz), 6.95 (br s, 2H, D ₂ O exchangeable), 7.35–7.50 (m, 3H), 7.51 (d, 1H, <i>J</i> =4.0 Hz), 7.64 (d, 1H, <i>J</i> =7.6 Hz), 8.59–8.67 (m, 1H).	432.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
58		B	(DMSO- <i>d</i> ₆): δ 1.04-1.15 (m, 1H), 1.85-1.97 (m, 1H), 2.06-2.17 (m, 1H), 3.35-3.48 (m, 2H), 3.69-3.75 (m, 1H), 3.78-3.87 (m, 1H), 4.38 (d, 1H, <i>J</i> =4.1 Hz, D ₂ O exchangeable), 4.45 (app quint, 1H, <i>J</i> =8.6 Hz), 4.53-4.60 (m, 2H, D ₂ O exchangeable), 6.83 (d, 1H, <i>J</i> =8.7 Hz, D ₂ O exchangeable), 7.02 (br s, 2H, D ₂ O exchangeable), 7.50-7.66 (m, 5H including 4 aryls), 7.92 (d, 1H, <i>J</i> =7.7 Hz).	454.1 (Cl pattern)	1
59		B	(DMSO- <i>d</i> ₆): δ 1.05 (dt, 1H, <i>J</i> =8.7, 13.1 Hz), 1.85-1.97 (m, 1H), 2.16 (dt, 1H, <i>J</i> =8.5, 13.1 Hz), 3.69-3.76 (m, 1H), 3.81-3.90 (m, 1H), 4.38-4.50 (m, 1H), 4.45 (d, 1H, <i>J</i> =4.0 Hz), 4.63 (t, 1H, <i>J</i> =5.1 Hz, D ₂ O exchangeable), 4.65 (d, 1H, <i>J</i> =6.1 Hz, D ₂ O exchangeable), 6.80 (d, 1H, <i>J</i> =8.4 Hz), 7.14 (br s, 2H), 7.60-7.68 (m, 1H), 7.70-7.80 (m, 2H), 8.01 (dd, 1H, <i>J</i> =0.9, 7.91 Hz)	453.1 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
60		B	(DMSO- <i>d</i> ₆): δ 1.13 (dt, 1H, <i>J</i> =8.2, 13.2 Hz), 1.85-1.97 (m, 1H), 2.21 (dt, 1H, <i>J</i> =8.2, 13.2 Hz), 3.39 (t, 2H, <i>J</i> =5.3 Hz), 3.69-3.78 (m, 2H), 4.27-4.38 (m, 1H), 4.45 (d, 1H, <i>J</i> =4.0 Hz), 4.65-4.72 (m, 2H), 6.43 (d, 1H, <i>J</i> =7.9 Hz), 7.04 (br s, 2H), 7.36-7.43 (m, 2H), 7.55-7.61 (m, 1H), 7.65-7.71 (m, 1H).	409 (2 Cl pattern)	2
61		B	(DMSO- <i>d</i> ₆): δ 1.14 (dt, 1H, <i>J</i> =8.6, 12.9 Hz), 1.92-2.05 (m, 1H), 2.15 (dt, 1H, <i>J</i> =8.7, 12.4 Hz), 3.21 (s, 3H), 3.22-3.40 (m, 2H), 3.64-3.71 (m, 1H), 3.78-3.86 (m, 1H), 4.32 (quint, 1H, <i>J</i> =7.8 Hz), 4.49 (d, 1H, <i>J</i> =4.5 Hz, D ₂ O exchangeable), 4.67 (d, 1H, <i>J</i> =5.7 Hz, D ₂ O exchangeable), 6.51 (d, 1H, <i>J</i> =8.0 Hz, D ₂ O exchangeable), 6.90 (br s, 2H), 7.25 (dd, <i>J</i> =1.2, 5.1 Hz), 7.63 (dd, 1H, <i>J</i> =3.0, 4.9 Hz), 7.83 (dd, 1H, <i>J</i> =1.1, 2.9 Hz).	395.1 (Cl pattern)	2

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
62		B	(DMSO- <i>d</i> ₆): δ 1.28 (dt, 1H, <i>J</i> =8.6, 13.0 Hz), 1.91-2.04 (m, 1H), 2.25 (dt, 1H, <i>J</i> =8.6, 13.1 Hz), 3.04-3.19 (br s, 6H), 3.30-3.51 (m, 2H), 3.75-3.82 (m, 1H), 3.92-4.02 (m, 1H), 3.94 (s, 3H), 4.32-4.46 (m, 1H), 4.51 (d, 1H, <i>J</i> =4.7 Hz), 4.58-4.65 (m, 2H), 7.07 (d, 1H, <i>J</i> =7.7 Hz), 7.67 (dd, 1H, <i>J</i> =4.6, 8.0 Hz), 8.03 (dd, 1H, <i>J</i> =1.5, 8.0 Hz), 8.63 (dd, 1H, <i>J</i> =1.3, 4.5 Hz).	462.2(Cl pattern)	1
63		A	(DMSO- <i>d</i> ₆): δ 1.21 (dt, 1H, <i>J</i> =7.0, 12.9 Hz), 1.85-1.98 (m, 1H), 2.13-2.26 (m, 1H), 2.71 (s, 3H), 3.39-3.47 (m, 2H), 3.74-3.87 (m, 2H), 4.35 (quint, 1H, <i>J</i> =7.5 Hz), 4.45 (d, 1H, <i>J</i> =4.4 Hz, OH), 4.64 (d, 1H, <i>J</i> =5.2 Hz), 4.79 (app t, 1H, <i>J</i> =5.0 Hz), 6.75 (d, 1H, <i>J</i> =8.0 Hz, NH), 7.05 (br s, 2H, NH ₂), 7.90 (d, 1H, <i>J</i> =3.2 Hz), 7.99-8.03 (m, 2H), 8.03 (d, 1H, <i>J</i> =8.2 Hz).	473.2 (Cl pattern)	1

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
64		A	(DMSO- <i>d</i> ₆): δ 1.18 (dt, 1H, <i>J</i> =7.5, 13.0 Hz), 1.84-1.95 (m, 1H), 2.12-2.24 (m, 1H), 2.51 (s, 3H), 2.60 (s, 3H), 3.39-3.45 (m, 2H), 3.72-3.83 (m, 2H), 4.26-4.38 (m, 1H), 4.41-4.49 (m, 1H), 4.61-4.67 (m, 1H), 4.75-4.81 (m, 1H), 6.63 (d, 1H, <i>J</i> =8.0 Hz, NH), 6.96 (br s, 2H, NH ₂), 7.18 (d, 1H, <i>J</i> =8.1 Hz, Ar), 7.79 (d, 1H, <i>J</i> =8.2 Hz, Ar).	436.2 (Cl pattern)	1
65		A	(DMSO- <i>d</i> ₆): δ 1.11-1.33 (m, 1H), 1.86-1.96 (m, 1H), 2.13-2.32 (m, 1H), 2.57 (s, 3H), 2.72-2.84 (m, 3H, N-Me), 3.39-3.45 (m, 2H), 3.74-3.83 (m, 2H), 3.87 (s, 3H), 4.24-4.35 (m, 1H), 4.42-4.49 (m, 1H), 4.60-4.69 (m, 1H), 4.73-4.81 (m, 1H), 6.50-6.75 (m, 1H, NH), 6.70 (d, 1H, <i>J</i> =8.5 Hz, Ar), 7.25-7.42 (m, 1H, NH), 7.84 (d, 1H, <i>J</i> =8.5 Hz, Ar).	434.2 (Cl pattern)	4

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
66		A	(DMSO- <i>d</i> ₆): δ 1.11-1.33 (m, 1H), 1.86-1.98 (m, 1H), 2.12-2.32 (m, 1H), 2.52 (s, 3H), 2.60 (s, 3H), 2.72-2.84 (m, 3H, N-Me), 3.38-3.45 (m, 2H), 3.73-3.90 (m, 2H), 4.24- 4.35 (m, 1H), 4.42-4.49 (m, 1H), 4.59-4.70 (m, 1H), 4.75-4.83 (m, 1H), 6.57-6.78 (m, 1H, NH), 7.18 (d, 1H, <i>J</i> =8.2 Hz), 7.29-7.46 (m, 1H, NH), 7.79 (d, 1H, <i>J</i> =8.4 Hz, Ar).	450.1 (Cl pattern)	4
67		A	(DMSO- <i>d</i> ₆): δ 1.13-1.31 (m, 1H), 1.86-1.97 (m, 1H), 2.14-2.32 (m, 1H), 2.57 (s, 3H), 3.26 (s, 3H), 3.33-3.50 (m, 6H), 3.72-3.92 (m, 2H), 3.87 (s, 3H), 4.18- 4.34 (m, 1H), 4.41-4.49 (m, 1H), 4.62 (d, 1H, <i>J</i> =4.7 Hz), 4.72-4.83 (m, 1H), 6.53-6.75 (m, 1H, NH), 6.71 (d, 1H, <i>J</i> =8.5 Hz, Ar), 7.25-7.41 (m, 1H, NH), 7.84 (d, 1H, <i>J</i> =8.7 Hz, Ar).	478.2 (Cl pattern)	4

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
68		A	(DMSO- <i>d</i> ₆): δ 1.17-1.32 (m, 1H), 1.86-1.97 (m, 1H), 2.15-2.30 (m, 1H), 2.58 (s, 3H), 3.37-3.48 (m, 2H), 3.72-3.92 (m, 2H), 3.87 (s, 3H), 3.98-4.39 (m, 3H), 4.43-4.51 (m, 1H), 4.60-4.67 (m, 1H), 4.76-4.85 (m, 1H), 6.72 (d, 1H, <i>J</i> =8.5 Hz, Ar), 6.78-6.92 (m, 1H, NH), 7.86 (d, 1H, <i>J</i> =8.5 Hz, Ar), 7.90-8.10 (m, 1H, NH).	502.2	4
69		B	(DMSO- <i>d</i> ₆): δ 1.14-1.32 (m, 1H), 1.85-1.97 (m, 1H), 2.13-2.30 (m, 1H), 2.59 (s, 3H), 3.26 (s, 3H), 3.35-3.50 (m, 6H), 3.72-3.89 (m, 2H), 4.19-4.35 (m, 1H), 4.41-4.51 (m, 1H), 4.59-4.67 (m, 1H), 4.75-4.85 (m, 1H), 5.01 (q, 2H, <i>J</i> =9.1 Hz), 6.60-6.79 (m, 1H, NH), 6.88 (d, 1H, <i>J</i> =8.6 Hz, Ar), 7.30-7.55 (m, 1H, NH), 7.94 (d, 1H, <i>J</i> =8.5 Hz, Ar).	546.2	4

Table I

Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
70		B	(DMSO- <i>d</i> ₆): δ 1.17-1.33 (m, 1H), 1.86-1.98 (m 1H), 2.14-2.30 (m, 1H), 2.60 (s, 3H), 3.38-3.48 (m, 2H), 3.72-3.81 (m, 1H), 3.78-3.89 (m, 1H), 3.97-4.40 (m, 3H), 4.41-4.51 (m, 1H), 4.60-4.67 (m, 1H), 4.75-4.87 (m, 1H), 5.01 (q, 2H, <i>J</i> =9.0 Hz), 6.81-6.97 (m, 1H, NH), 6.89 (d, 1H, <i>J</i> =8.6 Hz, Ar), 7.97 (d, 1H, <i>J</i> =8.5 Hz, Ar), 8.0-8.11 (m, 1H, NH).	570.1	4
71		A	(DMSO- <i>d</i> ₆): δ 1.11-1.34 (m, 1H), 1.31 (t, 3H, <i>J</i> =7.0 Hz), 1.86-1.97 (m 1H), 2.14-2.30 (m, 1H), 2.56 (s, 3H), 3.26 (s, 3H), 3.37-3.49 (m, 6H), 3.72-3.89 (m, 2H), 4.17-4.38 (m, 2H), 4.31 (q, 2H, <i>J</i> =7.0 Hz), 4.40-4.48 (m, 1H), 4.60-4.66 (m, 1H), 4.73-4.82 (m, 1H), 6.54-6.63 (m, 1H, NH), 6.67 (d, 1H, <i>J</i> =8.4 Hz, Ar), 7.38-7.50 (m, 1H, NH), 7.82 (d, 1H, <i>J</i> =8.4 Hz, Ar).	492.2 (Cl pattern)	4

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
72		A	(DMSO- <i>d</i> ₆): δ 1.17-1.33 (m, 1H), 1.32 (t, 3H, <i>J</i> =7.0 Hz), 1.86-1.98 (m, 1H), 2.14-2.30 (m, 1H), 2.56 (s, 3H), 3.37-3.49 (m, 2H), 3.72-3.80 (m, 1H), 3.79-3.89 (m, 1H), 3.97-4.34 (m, 3H), 4.32 (q, 2H, <i>J</i> =7.0 Hz), 4.42-4.49 (m, 1H), 4.62 (app d, 1H, <i>J</i> =5.1 Hz), 4.74-4.83 (m, 1H), 6.68 (d, 1H, <i>J</i> =8.5 Hz, Ar), 6.77-6.90 (m, 1H, NH), 7.85 (d, 1H, <i>J</i> =8.5 Hz, Ar), 7.90-8.08 (m, 1H, NH).	516.2 (Cl pattern)	4
73		A	(DMSO- <i>d</i> ₆): δ 1.12-1.35 (m, 1H), 1.32 (t, 3H, <i>J</i> =7.0 Hz), 1.84-1.97 (m, 1H), 2.13-2.30 (m, 1H), 3.26 (s, 3H), 3.36-3.50 (m, 6H), 3.72-3.89 (m, 2H), 4.19-4.33 (m, 1H), 4.33 (q, 2H, <i>J</i> =7.0 Hz), 4.40-4.48 (m, 1H), 4.62 (app d, 1H, <i>J</i> =4.9 Hz), 4.74-4.84 (m, 1H), 6.60-6.80 (m, 1H, NH), 6.84 (dd, 1H, <i>J</i> =8.6, 0.6 Hz, Ar), 7.08-7.52 (m, 1H, NH), 7.83 (dd, 1H, <i>J</i> =8.6, 2.5 Hz), 8.34 (d, 1H, <i>J</i> =2.1 Hz, Ar).	478.2 (Cl pattern)	4

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
74		A	(DMSO- <i>d</i> ₆): δ 1.18-1.33 (m, 1H), 1.34 (t, 3H, <i>J</i> =7.0 Hz), 1.85-1.98 (m, 1H), 2.12-2.31 (m, 1H), 3.38-3.48 (m, 2H), 3.72-3.82 (m, 1H), 3.79-3.89 (m, 1H), 3.97-4.30 (m, 3H), 4.34 (q, 2H, <i>J</i> =7.0 Hz), 4.45 (app d, 1H, <i>J</i> =4.9 Hz), 4.62 (app d, 1H, <i>J</i> =5.2 Hz), 4.75-4.87 (m, 1H), 6.86 (d, 1H, <i>J</i> =8.6 Hz, Ar), 6.85-6.97 (m, 1H, NH), 7.84 (dd, 1H, <i>J</i> =8.6, 2.3 Hz), 7.90-8.09 (m, 1H, NH), 8.36 (d, 1H, <i>J</i> =2.1 Hz, Ar).	502.2	4
75		A	(DMSO- <i>d</i> ₆): δ 1.14-1.32 (m, 1H), 1.84-1.98 (m, 1H), 2.12-2.30 (m, 1H), 3.27 s, 3H), 3.38-3.50 (m, 6H), 3.72-3.90 (m, 2H), 3.88 (s, 3H), 4.19-4.36 (m, 1H), 4.40-4.48 (m, 1H), 4.62 (app d, 1H, <i>J</i> =5.0 Hz), 4.75-4.85 (m, 1H), 6.62-6.80 (m, 1H, NH), 6.88 (d, 1H, <i>J</i> =8.7 Hz, Ar), 7.27-7.50 (m, 1H, NH), 7.84 (dd, 1H, <i>J</i> =8.6, 2.2 Hz), 8.36 (d, 1H, <i>J</i> =2.0 Hz, Ar).	464.2 (Cl pattern)	4



Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
76		A	(DMSO- <i>d</i> ₆): δ 1.14-1.32 (m, 1H), 1.85-1.98 (m 1H), 2.12-2.30 (m, 1H), 3.38-3.47 (m, 2H), 3.73-3.80 (m, 1H), 3.80-3.90 (m, 1H), 3.89 (s, 3H), 3.96-4.15 (m, 2H), 4.14 (m, 1H), 4.45 (app d, 1H, <i>J</i> =5.1 Hz), 4.63 (d, 1H, <i>J</i> =5.3 Hz), 4.75-4.87 (m, 1H), 6.84-6.97 (m, 1H, NH), 6.90 (d, 1H, <i>J</i> =8.6 Hz, Ar), 7.86 (dd, 1H, <i>J</i> =8.6, 2.3 Hz), 7.90-8.10 (m, 1H, NH), 8.38 (d, 1H, <i>J</i> =2.1 Hz, Ar).	488.1	4
77		A	(DMSO- <i>d</i> ₆): δ 1.13-1.28 (m, 1H), 1.85-1.97 (m 1H), 2.17-2.35 (m, 1H), 2.27 (br s, 3H), 3.26 (s, 3H), 3.30-3.52 (m, 6H), 3.73-3.82 (m, 2H), 3.88 (s, 3H), 4.17-4.32 (m, 1H), 4.38-4.43 (m, 1H), 4.60 (app d, 1H, <i>J</i> =4.4 Hz), 4.75-4.83 (m, 1H), 6.22-6.38 (br s, 1H, NH), 6.65-6.88 (m, 1H), 6.86 (d, 1H, <i>J</i> =8.6 Hz, Ar), 7.84 (dd, 1H, <i>J</i> =8.7, 2.3 Hz), 8.36 (d, 1H, <i>J</i> =1.9 Hz, Ar).	444.2	5

Table I



Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
78		A	(DMSO- <i>d</i> ₆): δ 1.13-1.28 (m, 1H), 1.60-1.72 (m, 2H), 1.85-1.97 (m 1H), 2.27 (br s, 3H), 3.26-3.39 (m, 2H), 3.39-3.50 (m, 2H), 3.74-3.84 (m, 2H), 3.88 (s, 3H), 4.18-4.32 (m, 1H), 4.38-4.43 (m, 1H), 4.40-4.50 (m, 1H), 4.51-4.55 (m, 1 H), 4.75-4.83 (m, 1H), 6.22-6.33 (m, 1H, NH), 6.70-6.98 (m, 1H), 6.86 (app d, 1H, <i>J</i> =8.7 Hz, Ar), 7.82 (dd, 1H, <i>J</i> =8.6, 2.4 Hz), 8.35 (d, 1H, <i>J</i> =1.8 Hz, Ar).	444.3	5
79		A	¹H NMR (DMSO-<i>d</i>₆) δ 1.12-1.24 (m, 1H), 1.62-1.73 (m, 2H), 1.84-1.92 (m, 1H), 2.10-2.20 (m, 1H), 2.68-2.77 (m, 2H), 3.38-3.95 (m, 2H), 3.72-3.82 (m, 2H), 4.28-4.37 (m, 1H), 4.43 (d, OH, <i>J</i> =4.6 Hz), 4.61 (d, OH, <i>J</i> =5.0 Hz), 4.78 (t, OH, <i>J</i> =5.0Hz), 6.94 (br. s, 1 NH), 7.30 (d, 1 H, <i>J</i> =7.9Hz), 7.78-7.84 (m, 1H) and 8.64 (s, 1 H).	418.2	


Table I					
Compd #	Structure	EC90 Range (uM)	¹ H NMR data	MS	Synth. Method
80		B	¹H NMR (DMSO-d₆) 1.22-1.29 (m, 1H), 1.85-1.92 (m, 1H), 2.13-2.21 (m, 1H), 2.27 (s, 3H, <u>CH₃</u>), @3.40 (m, 2H), 3.72-3.80 (m, 2H), 4.28-4.36 (m, 1H), 4.40 (br. s, 1 <u>OH</u>), 4.63 (d, 1 <u>OH</u> , <i>J</i> =4.0 Hz), 4.71 (t, 1 <u>OH</u> , <i>J</i> =4.0 Hz), 6.45 (d, 1 H, <i>J</i> =7.9 Hz), 6.60 (br. s, <u>NH₂</u>), 7.83 (d, 1 H, <i>J</i> = 3.2Hz), and 7.89 (d, 1H, <i>J</i> = 3.2 Hz).	362.2	

Table I, Continued



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
81		A	(DMSO- <i>d</i> ₆): δ 1.13-1.28 (m, 1H), 1.32 (t, 3H, <i>J</i> = 7.1 Hz), 1.85-1.97 (m, 1H), 2.17-2.35 (m, 1H), 2.27 (br s, 3H), 3.26 (s, 3H), 3.35-3.49 (m, 6H), 3.73-3.82 (m, 2H), 4.17-4.30 (m, 1H), 4.32 (q, 2H, <i>J</i> = 7.0 Hz), 4.38-4.45 (m, 1H), 4.61 (app d, 1H, <i>J</i> = 3.6 Hz), 4.76-4.83 (app t, 1H), 6.22-6.39 (br s, 1H, NH), 6.68-6.95 (m, 1H), 6.83 (dd, 1H, <i>J</i> = 8.6, 0.8 Hz, Ar), 7.83 (dd, 1H, <i>J</i> = 8.7, 2.4 Hz), 8.36 (dd, 1H, <i>J</i> = 0.7, 2.3 Hz, Ar).	458.3	5
82		A	(DMSO- <i>d</i> ₆): δ 1.11-1.26 (m, 1H), 1.85-1.97 (m, 1H), 2.16-2.35 (m, 1H), 2.29 (br s, 3H), 2.56 (s, 3H), 3.26 (s, 3H), 3.37-3.49 (m, 6H), 3.72-3.82 (m, 2H), 3.86 (s, 3H), 4.17-4.28 (m, 1H), 4.38-4.44 (m, 1H), 4.59-4.64 (m, 1H), 4.70-4.78 (m, 1H), 6.12-6.28 (m, 1H), 6.54-6.93 (m, 1H, NH), 6.69 (d, 1H, <i>J</i> = 8.6 Hz, Ar), 7.82 (d, 1H, <i>J</i> = 8.6 Hz, Ar).	458.3	5

Table I, Continued

Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			Ar).		
83		B	(DMSO- <i>d</i> ₆): δ 1.21 (s, 3H), 1.40 (s, 3H), 1.59-1.73 (m 2H), 2.08-2.19 (m, 1H), 2.22-2.38 (m, 1H), 2.27 (br s, 3H), 3.24-3.40 (m, 2H), 3.39-3.52 (m, 2H), 3.50-3.61 (m, 1H), 3.88 (s, 3H), 4.33-4.60 (m, 1H), 4.47 (br s, 1H), 5.00-5.18 (m, 1H), 6.74-6.99 (br s, 2H, NHs), 6.85 (d, 1H, <i>J</i> =8.7 Hz, Ar), 7.83 (dd, 1H, <i>J</i> =2.4, 8.6 Hz, Ar), 8.35 (d, 1H, <i>J</i> = 2.1 Hz).	484.3	5
84		A	(DMSO- <i>d</i> ₆): δ 1.11-1.26 (m, 1H), 1.86-1.97 (m 1H), 2.10 (s, 3H), 2.16-2.35 (m, 1H), 2.27 (br s, 3H), 2.62 (app t, 2H, <i>J</i> = 8.0 Hz), 3.37-3.49 (m, 4H), 3.72-3.85 (m, 2H), 3.88 (s, 3H), 4.19-4.32 (m, 1H), 4.38-4.46 (m, 1H), 4.59 (app d, 1H, <i>J</i> = 4.7 Hz), 4.78 (app t, 1H, <i>J</i> = 5.0 Hz), 6.22-6.38 (m, 1H), 6.80-6.91 (m, 1H), 7.03-7.13 (m, 1H), 7.84 (dd, 1H, <i>J</i> = 2.3, 8.7 Hz), 8.35 (d, 1H, <i>J</i> = 2.3 Hz).	460.2	5



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
85		A	(DMSO- <i>d</i> ₆): δ 0.88 (s, 9H), 1.14-1.30 (m, 1H), 1.32 (t, 3H, <i>J</i> = 7.0 Hz), 1.85-1.97 (m, 1H), 2.17-2.32 (m, 1H), 2.26 (br s, 3H), 3.08-3.23 (m, 2H), 3.38-3.49 (m, 2H), 3.73-3.83 (m, 2H), 4.17-4.30 (m, 1H), 4.32 (q, 2H, <i>J</i> = 7.0 Hz), 4.39-4.46 (m, 1H), 4.51-4.61 (m, 1H), 4.76-4.83 (m, 1H), 6.12-6.34 (m, 1H, NH), 6.83 (dd, 1H, <i>J</i> = 0.6, 8.6 Hz, Ar), 6.91-7.02 (m, 1H), 7.81 (dd, 1H, <i>J</i> = 8.6, 2.3 Hz), 8.33 (dd, 1H, <i>J</i> = 0.6, 2.3 Hz, Ar).	470.3	5
86		A	(DMSO- <i>d</i> ₆): δ 1.14-1.30 (m, 1H), 1.32 (t, 3H, <i>J</i> = 7.0 Hz), 1.84-1.98 (m, 1H), 2.13-2.38 (m, 1H), 2.30 (br s, 3H), 3.38-3.48 (m, 2H), 3.72-3.84 (m, 2H), 3.96-4.30 (m, 3H), 4.32 (q, 2H, <i>J</i> = 7.0 Hz), 4.40-4.48 (m, 1H), 4.58-4.64 (m, 1H), 4.78-4.86 (m, 1H), 6.42-6.52 (m, 1H, NH), 6.83 (dd, 1H, <i>J</i> = 0.8, 8.7 Hz, Ar), 7.41-7.61 (m, 1H),	482.2	5

Table I, Continued



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			7.85 (dd, 1H, $J=8.6, 2.3$ Hz), 8.36 (dd, 1H, $J=0.6, 2.3$ Hz, Ar).		
87		A	(DMSO- d_6): δ 0.87 (t, 3H, $J = 7.3$ Hz), 1.12-1.30 (m, 1H), 1.32 (t, 3H, $J = 7.0$ Hz), 1.51 (hex, 2H, $J = 7.3$ Hz), 1.84-1.97 (m 1H), 2.15-2.32 (m 1H), 2.26 (br s, 3H), 3.11-3.28 (m, 2H), 3.42 (app t, 2H, $J = 5.0$ Hz), 3.72-3.83 (m, 2H), 4.18-4.35 (m, 1H), 4.32 (q, 2H, $J= 7.0$ Hz), 4.37-4.43 (m, 1H), 4.60 (app d, 1H, $J = 4.7$ Hz), 4.77 (app t, 1H, $J = 5.0$ Hz), 6.14-6.32 (m, 1H, NH), 6.82 (dd, 1H, $J=0.7, 8.6$ Hz, Ar), 6.86-7.04 (m, 1H), 7.82 (dd, 1H, $J=8.6, 2.4$ Hz), 8.33 (dd, 1H, $J=0.6, 2.2$ Hz, Ar).	442.3	5
88		A	(DMSO- d_6): δ 1.10 (t, 3H, $J = 7.1$ Hz), 1.12-1.30 (m, 1H), 1.32 (t, 3H, $J = 7.0$ Hz), 1.84-2.05 (m 1H), 2.16-2.32 (m, 1H), 2.27 (br s, 3H), 3.18-3.48 (m, 4H), 3.74-3.85 (m, 2H), 4.18-4.35 (m, 1H), 4.32 (q, 2H, $J= 7.0$ Hz), 4.37-4.43 (m,	428.3	5


Table I, Continued					
Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			1H), 4.57-4.67 (m, 1H), 4.73-4.83 (m, 1H), 6.08-6.32 (m, 1H, NH), 6.70-7.01 (m, 1H), 6.82 (dd, 1H, $J=0.7, 8.6$ Hz, Ar), 7.82 (dd, 1H, $J=8.6, 2.4$ Hz), 8.33 (dd, 1H, $J=0.7, 2.3$ Hz, Ar).		
89		B	(DMSO- d_6): δ 1.14-1.30 (m, 1H), 1.32 (t, 3H, $J = 7.0$ Hz), 1.85-1.98 (m 1H), 2.16-2.32 (m, 1H), 2.28 (br s, 3H), 3.20-3.45 (m, 2H), 3.51 (app q, 1H), 3.59 (app q, 1H), 3.77 (br s, 2H), 4.18-4.35 (m, 1H), 4.32 (q, 1H, $J = 7.0$ Hz), 4.39-4.51 (m, 1H), 4.44 (app t, 1H, $J = 5.1$ Hz), 4.56-4.71 (m, 1H), 4.60 (t, 1H, $J = 5.1$ Hz), 4.76-4.85 (m, 1H), 6.28-6.40 (m, 1H, NH), 6.82 (dd, 1H, $J=0.7, 8.6$ Hz, Ar), 6.92-7.19 (m, 1H), 7.83 (dd, 1H, $J=8.6, 2.3$ Hz), 8.34 (dd, 1H, $J=0.8, 2.4$ Hz, Ar).	446.3	5

Table I, Continued



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
90		B	(DMSO- <i>d</i> ₆): δ 1.14-1.30 (m, 1H), 1.86-1.98 (m, 1H), 2.15-2.35 (m, 1H), 2.31 (br s, 3H), 3.38-3.49 (m, 2H), 3.74-3.85 (m, 2H), 3.88 (s, 3H), 3.94-4.32 (m, 3H), 4.41 (app d, 1H, <i>J</i> = 4.1 Hz), 4.58 (app d, 1H, <i>J</i> = 4.5 Hz), 4.76-4.85 (m, 1H), 6.41-6.52 (m, 1H, NH), 6.86 (d, 1H, <i>J</i> = 0.7, 8.6 Hz, Ar), 7.39-7.57 (m, 1H), 7.86 (dd, 1H, <i>J</i> = 8.6, 2.2 Hz), 8.38 (d, 1H, <i>J</i> = 1.5 Hz, Ar).	468.2	5
91		A	(DMSO- <i>d</i> ₆): δ 1.10 (t, 3H, <i>J</i> = 7.1 Hz), 1.13-1.29 (m, 1H), 1.86-1.98 (m, 1H), 2.17-2.33 (m, 1H), 2.27 (br s, 3H), 3.22-3.35 (m, 2H), 3.43 (app t, 2H, <i>J</i> = 5.0 Hz), 3.75-3.84 (m, 1H), 3.88 (s, 3H), 4.19-4.32 (m, 1H), 4.40 (app d, 1H, <i>J</i> = 4.3 Hz), 4.61 (app d, 1H, <i>J</i> = 4.7 Hz), 4.77 (app t, 1H, <i>J</i> = 5.0 Hz), 6.19-6.32 (m, 1H, NH), 6.80-6.97 (m, 1H), 6.85 (dd, 1H, <i>J</i> = 0.7, 8.6 Hz, Ar), 7.83 (dd, 1H, <i>J</i> = 8.7, 2.3 Hz), 8.35 (dd, 1H,	414.2	5

Table I, Continued



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			<i>J</i> =0.7, 2.3 Hz, Ar).		
92		A	(DMSO- <i>d</i> ₆): δ 0.18-0.23 (m, 2H), 0.36-0.43 (m, 2H), 0.98-1.13 (m, 1H), 1.13-1.29 (m, 1H), 1.85-1.98 (m, 1H), 2.17-2.32 (m, 1H), 2.27 (br s, 3H), 3.10-3.21 (m, 2H), 3.43 (app t, 2H, <i>J</i> = 4.9 Hz), 3.75-3.84 (m, 2H), 3.88 (s, 3H), 4.19-4.32 (m, 1H), 4.38-4.43 (m, 1H), 4.60 (app d, 1H, <i>J</i> = 4.7 Hz), 4.77 (app t, 1H, <i>J</i> = 5.1 Hz), 6.18-6.33 (m, 1H, NH), 6.82-7.08 (m, 1H), 6.85 (dd, 1H, <i>J</i> =0.5, 8.6 Hz, Ar), 7.83 (dd, 1H, <i>J</i> =8.6, 2.4 Hz), 8.35 (dd, 1H, <i>J</i> =0.4, 1.9 Hz, Ar).	440.2	5
93		A	(DMSO- <i>d</i> ₆): δ 1.10-1.21 (m, 1H), 1.84-1.96 (m, 1H), 2.09-2.28 (m, 1H), 2.27 (br s, 3H), 3.37-3.46 (m, 2H), 3.71 (s, 6H), 3.75-3.82 (m, 2H), 3.88 (s, 3H), 4.19-4.32 (m, 1H), 4.30-4.50 (m, 3H), 4.58 (app d, 1H, <i>J</i> = 3.9 Hz), 4.76 (app t, 1H, <i>J</i> = 5.0 Hz), 6.22-6.36 (m, 2H), 6.51 (br s, 2H, Ph), 6.85 (dd,	536.1	5

Table I, Continued

Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			1H, <i>J</i> =0.7, 8.7 Hz, Ar), 7.20-7.57 (m, 1H), 7.83 (dd, 1H, <i>J</i> =8.6, 2.3 Hz), 8.35 (dd, 1H, <i>J</i> =0.5, 2.2 Hz, Ar).		
94		A	(DMSO- <i>d</i> ₆): δ 1.08-1.21 (m, 1H), 1.85-1.99 (m, 1H), 2.18-2.35 (m, 1H), 2.27 (br s, 3H), 3.26 (s, 3H), 3.33 (s, 6H), 3.37-3.49 (m, 4H), 3.69-3.79 (m, 2H), 3.89 (s, 3H), 3.97 (s, 3H), 4.18-4.29 (m, 1H), 4.43-4.52 (m, 1H), 4.63-4.72 (m, 2H), 5.99-6.13 (m, 1H), 6.43 (d, 1H, <i>J</i> = 8.2 Hz), 6.62-6.94 (m, 1H), 7.77 (d, 1H, <i>J</i> = 8.1 Hz).	474.2	5
95		A	(DMSO- <i>d</i> ₆): δ 0.18-0.23 (m, 2H), 0.36-0.43 (m, 2H), 0.91-1.21 (m, 2H), 1.87-2.00 (m, 1H), 2.18-2.32 (m, 1H), 2.27 (br s, 3H), 3.03-3.23 m, 2H), 3.37-3.47 (m, 2H), 3.68-3.80 (m, 2H), 3.89 (s, 3H), 3.97 (s, 3H), 4.18-4.29 (m, 1H), 4.43-4.50 (m, 1H), 4.63-4.72 (m, 2H), 5.96-6.10 (m, 1H), 6.43 (d, 1H, <i>J</i> = 8.1 Hz), 6.80-7.06 (m, 1H),	470.2	5

Table I, Continued



Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			7.77 (d, 1H, <i>J</i> = 8.2 Hz).		
96		A	(DMSO- <i>d</i> ₆): δ 1.10-1.20 (m, 1H), 1.85-1.97 (m, 1H), 2.12 (s, 3H, CH ₃), 2.17-2.20 (m, 1H), 2.21-2.32 (bs, 4H, CH ₂ & CH ₃), 2.50-2.65 (m, 1H), 3.72-3.85 (m, 2H), 4.23-4.32 (m, 1H), 4.34-4.41 (m, 1H), 4.59 (d, <i>J</i> = 5.1 Hz, 1H), 4.71 (t, 1H, <i>J</i> = 4.5 Hz), 6.39 (bs, 1H), 7.14 (bs, 1H), 7.32 (t, 1H, <i>J</i> = 4.2 Hz), 7.31 (d, 1H, <i>J</i> = 3.6 Hz), 7.75 (d, 1H, <i>J</i> = 3.9 Hz), 7.87 - 7.84 (m, 1H), 7.94 (d, 1H, <i>J</i> = 7.8 Hz), 8.52 (d, 1H, <i>J</i> = 4.5 Hz).	512.2	5
97		A	(DMSO- <i>d</i> ₆): δ 1.16-1.21 (m, 1H), 1.85-1.95 (m, 1H), 2.15-2.21 (m, 1H), 3.38-3.46 (m, 2H), 3.73-3.80 (m, 1H), 3.81-3.89 (m, 1H), 4.01-4.12 (m, 2H), 4.16-4.28 (m, 1H), 4.40-4.49 (m, 1H, 3'-OH), 4.61 (d, 1H, <i>J</i> = 5.4 Hz, 2'-OH), 4.77 (bs, 1H, 5'-OH), 7.02 (bs, 1H, NH-4), 7.30-7.33 (m, 1H), 7.40 (d, 1H, <i>J</i> = 3.9 Hz), 7.80 (d, 1H, <i>J</i> = 3.9 Hz).	540.1 (Cl pattern)	5



Table I, Continued					
Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			Hz), 7.88-7.83 (m, 1H), 7.97 (d, 1H, <i>J</i> = 7.8 Hz), 8.10 (bs, 1H), 8.52 (d, 1H, <i>J</i> = 4.8 Hz).		
98		**	(DMSO- <i>d</i> ₆): δ 1.16-1.27 (m, 1H), 1.56-1.68 (m, 1H), 1.75-2.10 (m, 6H, pip), 2.10-2.21 (m, 1H), 2.96-3.18 (m, 1H), 3.81-3.89 (m, 1H), 3.20-3.28 (m, 1H, pip), 3.40-3.45 (m, 2H), 3.73 (q, 1H, <i>J</i> = 4.8 Hz), 3.79-3.88 (m, 1H), 3.90-4.30 (m, 3H, 1'CH + CH ₂ -CF ₃), 4.61 (bs, 1H, 3'-OH), 4.58 (d, 1H, <i>J</i> = 5.4 Hz, 2'-OH), 4.72 (bs, 1H, 5'-OH), 7.98 (bs, 1H, NH-4), 7.20 (d, 1H, <i>J</i> = 3.9 Hz), 7.33 (d, 1H, <i>J</i> = 3.9 Hz), 8.08 (bs, 1H), 8.69 (bs, 1H, NH-pip).	273.6 (Cl pattern)	5
99		B	(DMSO- <i>d</i> ₆): δ 1.30 (s, 3H), 1.50 (s, 3H), 1.55-1.84 (m, 3H), 2.10 (bs, 1H, 5'-OH), 2.35-2.44 (m, 4H), 2.59-2.61 (m, 1H), 3.48-3.72 (bs, 4H, CH ₂), 3.74-3.77 (m, 1H), 3.85-3.94 (m, 1H), 4.46 (d, <i>J</i> = 5.1 Hz, 1H), 4.60-4.69 (m, 2H), 5.20 (bs, 2H), 6.88 (bs, 1H), 7.14-7.2 (m, 2H), 7.43	459.3	5




Table I, Continued					
Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			(d, 1H, J = 3.9 Hz), 7.60 -7.75 (m, 2H), 8.53 (d, 1H, J = 4.8 Hz).		
100		B	(DMSO- <i>d</i> ₆): δ 1.20 (s, 3H), 1.39 (s, 3H), 1.59-1.70 (m, 3H), 2.06-2.15 (m, 1H), 2.23 (s, 3H), 2.24-2.30 (m, 2H), 3.20-3.20 (m, 2H), 3.39-3.50 (m, 6H, 2xCH ₂ , 2xCH), 4.41-4.58 (m, 2H), 4.95 (bs, 1H, 5'-OH), 6.86 (bs, 2H), 7.05-7.10 (m, 1H), 7.31 (q, 1H, J = 3.4 Hz), 7.56 (q, 1H, J = 5.1 Hz).	459.3	5
101		B	(DMSO- <i>d</i> ₆): δ 1.20-1.34 (m, 1H), 2.05-2.13 (m, 1H), 2.36-2.45 (m, 1H), 2.72 (s, 3H), 3.57-3.64 (m, 2H), 3.80-3.89 (m, 1H), 3.91 (t, 1H, J = 5.1 Hz), 4.02-4.25 (m, 2H), 4.38 (q, 1H, J = 6 Hz, 1'-CH), 7.35-7.49 (m, 3H, arom), 8.16-8.21 (m, 2H, arom).	554.2 (Cl pattern)	5
102		A	(DMSO- <i>d</i> ₆): δ 1.10-1.20 (m, 1H), 1.85-1.97 (m, 1H), 2.12 (s, 3H, CH ₃), 2.17-2.20 (m, 1H), 2.21-2.32 (bs, 4H, CH ₂ & CH ₃), 2.50-2.65 (m, 1H), 3.72-3.85 (m, 2H), 4.23-4.32 (m, 1H), 4.34-	452.2	5




Table I, Continued					
Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			4.41 (m, 1H), 4.59 (d, <i>J</i> = 5.1 Hz, 1H), 4.71 (t, 1H, <i>J</i> = 4.5 Hz), 6.39 (bs, 1H), 7.14 (bs, 1H), 7.32 (t, 1H, <i>J</i> = 4.2 Hz), 7.31 (d, 1H, <i>J</i> = 3.6 Hz), 7.75 (d, 1H, <i>J</i> = 3.9 Hz), 7.87 - 7.84 (m, 1H), 7.94 (d, 1H, <i>J</i> = 7.8 Hz), 8.52 (d, 1H, <i>J</i> = 4.5 Hz).		
103		A	(DMSO- <i>d</i> ₆): δ 1.10 (t, <i>J</i> = 6.9 Hz, 3H), 1.14 -1.22 (m, 1H), 1.86-1.92 (m, 1H), 2.18-2.23 (m, 1H), 2.25 (s, 3H, CH ₃), 3.34-3.51 (m, 8H, CH ₂ x3 & OCH ₃), 3.72-3.81 (m, 2H), 4.19-4.28 (m, 1H), 4.38-4.42 (m, 1H), 4.60 (d, <i>J</i> = 5.1 Hz, 1H), 4.74 (t, 1H, <i>J</i> = 4.5 Hz), 6.25 (bs, 1H), 6.80 (bs, 1H), 7.24 (dd, 1H, <i>J</i> = 12 Hz), 7.60 (q, 1H, <i>J</i> = 3 Hz), 7.74-7.76 (m, 1H).	433.2	5
104		B	(DMSO- <i>d</i> ₆): δ 0.18-0.24 (m, 2H), 0.35-0.44 (m, 2H), 1.02-1.08 (m, 1H), 1.14-1.21 (m, 1H), 1.87-1.98 (m, 1H), 2.15-2.22 (m, 1H), 2.26 (s, 3H), 3.10-3.21 (m, 2H), 3.38-3.45 (bs, 2H, CH ₂), 3.74-3.81 (m, 2H),	492.2	5

Table I, Continued					
Compd #	Structure	EC90 (uM)	¹ H NMR data	MS (M+H)	Synth. Method
			4.20-4.31 (m, 1H), 4.37-4.44 (m, 1H, 3'-OH), 4.62 (d, 1H, <i>J</i> = 4.8 Hz), 4.73 (t, 1H, <i>J</i> = 5.1 Hz), 6.24-6.38 (m, 1H), 7.13 (bs, 1H), 7.29-7.31 (m, 1H), 7.33 (d, 1H, <i>J</i> = 3.6 Hz), 7.75 (d, 1H, <i>J</i> = 3.9 Hz), 7.84-7.91 (m, 1H), 7.93-7.96 (m, 1H), 8.51-8.53 (m, 1H).		
105		B	(DMSO- <i>d</i> ₆): δ 1.12-1.23 (m, 1H), 1.85-1.94 (m, 1H), 1.87-1.98 (m, 1H), 2.18-2.25 (m, 1H), 2.26 (s, 3H), 3.26 (s, 3H, OCH ₃), 3.10-3.21 (m, 2H), 3.39-3.49 (m, 6H), 3.73-3.82 (m, 2H), 4.24 (bs, 1H), 4.40 (m, 1H, 3'-OH), 4.62 (d, 1H, <i>J</i> = 4.8 Hz), 4.74 (t, 1H, <i>J</i> = 5.1 Hz), 6.37 (bs, 1H), 7.00 (bs, 1H), 7.29-7.31 (m, 1H), 7.34 (d, 1H, <i>J</i> = 3.6 Hz), 7.76 (d, 1H, <i>J</i> = 3.9 Hz), 7.95 (d, 1H, <i>J</i> = 8.1 Hz), 8.50-8.58 (m, 1H).	496.2	5

ASSAYS

Cell-based HCV Replicon Assay

To measure cell-based anti-HCV activity of the compounds of the present invention, replicon cells were seeded at 5000 cells/well in 96-well collagen I-coated Nunc plates in the presence of the compound of the invention. Various concentrations of a compound of the invention, typically in 10 serial 2-fold dilutions, were added to the assay mixture, the starting concentration of the compound ranging from 25 μ M to 1 μ M. The final concentration of DMSO was 0.5%, fetal bovine serum was 10%, in the assay media. Cells were harvested on day 3 by the addition of 1x cell lysis buffer (Ambion cat #8721).

The replicon RNA level was measured using real time PCR (Taqman assay). The amplicon was located in 5B. The PCR primers were: 5B.2F, ATGGACAGGCGCCCTGA; 5B.2R, TTGATGGGCAGCTTGGTTTC; the probe sequence was FAM-labeled CACGCCATGCGCTGCGG. GAPDH RNA was used as endogenous control and was amplified in the same reaction as NS5B (multiplex PCR) using primers and VIC-labeled probe recommended by the manufacturer (PE Applied Biosystem). The real-time RT-PCR reactions were run on ABI PRISM 7900HT Sequence Detection System using the following program: 48°C for 30 min, 95°C for 10 min, 40 cycles of 95°C for 15 sec, 60°C for 1 min. The Δ CT values ($CT_{5B} - CT_{GAPDH}$) were plotted against the concentration of test compound and fitted to the sigmoid dose-response model using GraphPad PRISM software. EC_{50} was defined as the concentration of inhibitor necessary to achieve Δ CT=1 over the projected baseline; EC_{90} the concentration necessary to achieve Δ CT=3.2 over the baseline. Alternatively, to quantitate the absolute amount of replicon RNA, a standard curve was established by including serially diluted T7 transcripts of replicon RNA in the Taqman assay. All Taqman reagents were from PE Applied Biosystems. Such an assay procedure was described in detail in e.g. Malcolm *et al.*, *Antimicrobial Agents and Chemotherapy* 50: 1013-1020 (2006).

HCV Replicon assay data for compounds of the invention that were tested was obtained using the above method. Calculated EC_{90} values are reported for each compound in Table I as a falling within the following ranges:

30 “A” - less than or equal to about 0.5 μ M

 “B” - greater than about 0.5 μ M to less than or equal to about 5.0 μ M

“C” - greater than about 5.0 μ M

METHODS OF USE

The compounds of the invention are useful in human and veterinary medicine for
5 treating or preventing a viral infection or a virus-related disorder in a patient. In
accordance with the invention, the compounds of the invention can be administered to a
patient in need of treatment or prevention of a viral infection or a virus-related disorder.

Accordingly, in one embodiment, the invention provides methods for treating a
viral infection in a patient comprising administering to the patient an effective amount of
10 at least one compounds of the invention or a pharmaceutically acceptable salt, ester,
prodrug, isomer, tautomer, or solvate thereof. In another embodiment, the invention
provides methods for treating a virus-related disorder in a patient comprising
administering to the patient an effective amount of at least one compounds of the
invention or a pharmaceutically acceptable salt, ester, prodrug, isomer, tautomer, or
15 solvate thereof.

Treatment or Prevention of a Viral Infection

The compounds of the invention can be used to treat or prevent a viral infection.
In one embodiment, the compounds of the invention can be used to inhibit viral
20 replication. In a specific embodiment, the compounds of the invention can be inhibitors
of HCV replication. Accordingly, the compounds of the invention are useful for treating
viral diseases and disorders related to the activity of a virus, such as HCV polymerase.

Such uses as are described herein may be performed in a patient in need thereof,
although *in vitro* and *ex vivo* uses, such as in diagnostic and research contexts, are also
25 contemplated. References made herein to the use of compounds of the invention also
refers to uses of compositions comprising compounds of the invention.

Examples of viral infections that can be treated or prevented using the present
methods, include but are not limited to, hepatitis A infection, hepatitis B infection and
hepatitis C infection.

30 In one embodiment, the viral infection is hepatitis C infection.

In one embodiment, the hepatitis C infection is acute hepatitis C. In another
embodiment, the hepatitis C infection is chronic hepatitis C.

The compositions and combinations of the present invention can be useful for treating a patient suffering from infection related to any HCV genotype. HCV types and subtypes may differ in their antigenicity, level of viremia, severity of disease produced, and response to interferon therapy as described in Holland *et al.*, *Pathology*, 30(2):192-195 (1998). The nomenclature set forth in Simmonds *et al.*, *J Gen Virol*, 74(Pt11):2391-2399 (1993) is widely used and classifies isolates into six major genotypes, 1 through 6, with two or more related subtypes, e.g., 1a, 1b. Additional genotypes 7-10 and 11 have been proposed, however the phylogenetic basis on which this classification is based has been questioned, and thus types 7, 8, 9 and 11 isolates have been reassigned as type 6, and type 10 isolates as type 3 (see Lamballerie *et al.*, *J Gen Virol*, 78(Pt1):45-51 (1997)). The major genotypes have been defined as having sequence similarities of between 55 and 72% (mean 64.5%), and subtypes within types as having 75%-86% similarity (mean 80%) when sequenced in the NS-5 region (see Simmonds *et al.*, *J Gen Virol*, 75(Pt 5):1053-1061 (1994)).

15

Treatment or Prevention of a Virus-Related Disorder

The compounds of the invention can be used to treat or prevent a virus-related disorder. Accordingly, the compounds of the invention are useful for treating disorders related to the activity of a virus, such as liver inflammation or cirrhosis. Virus-related disorders include, but are not limited to, RNA-dependent polymerase-related disorders and disorders related to HCV infection.

20

Treatment or Prevention of a RNA-Dependent Polymerase-Related Disorder

The compounds of the invention are useful for treating or preventing a RNA dependent polymerase (RdRp) related disorder in a patient. Such disorders include viral infections wherein the infective virus contain a RdRp enzyme.

25

Accordingly, in one embodiment, the present invention provides a method for treating a RNA dependent polymerase-related disorder in a patient, comprising administering to the patient an effective amount of at least one compounds of the invention or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof.

30

Treatment or Prevention of a Disorder Related to HCV Infection

The compounds of the invention can also be useful for treating or preventing a disorder related to an HCV infection. Examples of such disorders include, but are not limited to, cirrhosis, portal hypertension, ascites, bone pain, varices, jaundice, hepatic
5 encephalopathy, thyroiditis, porphyria cutanea tarda, cryoglobulinemia, glomerulonephritis, sicca syndrome, thrombocytopenia, lichen planus and diabetes mellitus.

Accordingly, in one embodiment, the invention provides methods for treating an HCV-related disorder in a patient, wherein the method comprises administering to the
10 patient a therapeutically effective amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof.

Combination Therapy

In another embodiment, the present methods for treating or preventing a viral
15 infection can further comprise the administration of one or more additional therapeutic agents. In one embodiment, such one or more additional therapeutic agent may be one or more additional compounds of the invention. In another embodiment, such one or more additional therapeutic agent is an agent other than a compound of the invention.

In one embodiment, the additional therapeutic agent is an antiviral agent. Non-
20 limiting examples of antiviral agents are as described herein and include, e.g., interferon.

In another embodiment, the additional therapeutic agent is an immunomodulatory agent, such as an immunosuppressive agent.

Accordingly, in one embodiment, the present invention provides methods for treating a viral infection in a patient, the method comprising administering to the patient:
25 (i) at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof, and (ii) at least one antiviral agent other than a compound of the invention, wherein the amounts administered are together effective to treat or prevent a viral infection.

When administering such a combination to a patient, the therapeutic agents in the
30 combination, or a pharmaceutical composition or compositions comprising the therapeutic agents, may be administered in any order such as, for example, sequentially, concurrently, together, simultaneously and the like. The amounts of the various actives in

such combination therapy may be different amounts (different dosage amounts) or same amounts (same dosage amounts). Thus, for non-limiting illustration purposes, a compound of the invention and an additional therapeutic agent may be present in fixed amounts (dosage amounts) in a single dosage unit (e.g., a capsule, a tablet and the like).

5 (A commercial example of such single dosage unit containing fixed amounts of two different active compounds is VYTORIN[®] (available from Merck Schering-Plough Pharmaceuticals, Kenilworth, New Jersey)).

In one embodiment, the at least one compound of the invention is administered at time when the additional antiviral agent(s) exert their prophylactic or therapeutic effect,
10 or *vice versa*.

In another embodiment, the at least one compound of the invention and the additional antiviral agent(s) are administered in doses commonly employed when such agents are used as monotherapy for treating a viral infection.

In another embodiment, the at least one compound of the invention and the
15 additional antiviral agent(s) are administered in doses lower than the doses commonly employed when such agents are used as monotherapy for treating a viral infection.

In another embodiment, the at least one compound of the invention and the additional antiviral agent(s) act synergistically and are administered in doses lower than the doses commonly employed when such agents are used as monotherapy for treating a
20 viral infection.

In one embodiment, the at least one compound of the invention and the additional antiviral agent(s) are present in the same composition. In one embodiment, this composition is suitable for oral administration. In another embodiment, this composition is suitable for intravenous administration.

25 Viral infections and virus-related disorders that can be treated or prevented using the combination therapy methods of the present invention include, but are not limited to, those listed above.

In one embodiment, the viral infection is HCV infection.

The at least one compound of the invention and the additional antiviral agent(s)
30 can act additively or synergistically. A synergistic combination may allow the use of lower dosages of one or more agents and/or less frequent administration of one or more agents of a combination therapy. A lower dosage or less frequent administration of one

or more agents may lower toxicity of the therapy without reducing the efficacy of the therapy.

In one embodiment, the administration of at least one compound of the invention and the additional antiviral agent(s) may inhibit the resistance of a viral infection to these
5 agents.

Non-limiting examples of other therapeutic agents useful in the present compositions and methods include an an viral (e.g., HCV) polymerase inhibitor, a viral (e.g., HCV) protease inhibitor, an interferon, a viral replication inhibitor, an antisense agent, a therapeutic vaccine, a viral protease inhibitor, a virion production inhibitor, an
10 immunosuppressive agent, an antiviral antibody, a CYP-450 inhibitor, an antiviral booster, and an antiviral sensitizer, and any agent useful for treating an RNA-dependent polymerase-related disorder.

In one embodiment, the at least one additional antiviral agent is a viral polymerase inhibitor.

15 In another embodiment, the at least one additional antiviral agent is an HCV polymerase inhibitor.

In one embodiment, the at least one additional antiviral agent is a viral protease inhibitor.

20 In another embodiment, the at least one additional antiviral agent is an HCV protease inhibitor.

In another embodiment, the at least one additional antiviral agent is an interferon.

In still another embodiment, the at least one additional antiviral agent is a viral replication inhibitor.

25 In another embodiment, the at least one additional antiviral agent is an antisense agent.

In another embodiment, the at least one additional antiviral agent is a therapeutic vaccine.

In a further embodiment, the at least one additional antiviral agent is an virion production inhibitor.

30 In another embodiment, the at least one additional antiviral agent is an antibody.

In another embodiment, the at least one additional antiviral agents comprise a protease inhibitor and a polymerase inhibitor.

In still another embodiment, the at least one additional antiviral agents comprise a protease inhibitor and an immunosuppressive agent.

In yet another embodiment, the at least one additional antiviral agents comprise a polymerase inhibitor and an immunosuppressive agent.

5 In a further embodiment, the at least one additional antiviral agents comprise a protease inhibitor, a polymerase inhibitor and an immunosuppressive agent.

In another embodiment the at least one additional agent is ribavirin, Levovirin, or Viramidine.

10 In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention and a CYP-450 inhibitor. Non-limiting examples of suitable CYP-450 inhibitors include ritonavir.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention and an interferon. Non-limiting examples of such interferon are as described herein and include alpha interferon, 15 pegylated interferon and conjugates thereof. Additional non-limiting examples of interferon include PEG-intron™ brand pegylated interferon, Pegasys™ brand pegylated interferon, Infergen™ brand interferon, and Alferon™ brand pegylated interferon.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention and an interferon. Further comprising 20 ribavirin, Levovirin, or Viramidine.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention and a protease inhibitor.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention, a protease inhibitor, and an interferon.

25 In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention, a protease inhibitor, an interferon, and ribavirin.

In other embodiments, pharmaceutical compositions according to the invention 30 comprise at least one compound of the invention, a polymerase inhibitor, and an interferon.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention, a polymerase inhibitor, an interferon, and ribavirin.

5 In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention, a protease inhibitor, polymerase inhibitor, and an interferon.

In other embodiments, pharmaceutical compositions according to the invention comprise at least one compound of the invention, a protease inhibitor, a polymerase
10 inhibitor, an interferon, and ribavirin.

HCV polymerase inhibitors useful in the present methods and compositions include, but are not limited to VP-19744 (Wyeth/ViroPharma), HCV-796 (Wyeth/ViroPharma), NM-283 (Idenix/Novartis), R-1626 (Roche), MK-0608 (Merck), A848837 (Abbott), GSK-71185 (Glaxo SmithKline), XTL-2125 (XTL
15 Biopharmaceuticals), and those disclosed in Ni *et al.*, *Current Opinion in Drug Discovery and Development*, 7(4):446 (2004); Tan *et al.*, *Nature Reviews*, 1:867 (2002); and Beaulieu *et al.*, *Current Opinion in Investigational Drugs*, 5:838 (2004).

Additional non-limiting examples of HCV polymerase inhibitors useful in the present methods and compositions include: MK00608, NM283, HCV796, R1626,
20 A848837, GSK71185, R7128, VCH759, GS9190, VP19744, and XTL2125.

Additional non-limiting examples of HCV polymerase inhibitors and HCV protease inhibitors useful in the present methods and compositions include: ANA598 (Anadys Pharmaceuticals), ABT-333, (Abbott), VCH-916, (Virochem), MK7009, (Merck), PF-00868554, (Pfizer) VX-500, (Vertex) GS9190, (Gilead) GSK625433,
25 (GlaxoSmithKline) ITMN-191 (R-7227), (Intermune), R7128, (Pharmasset/Roche), VCH-759 (Virochem), R1626, (Roche), TMC435350, (Medivir/Tibotec), SCH 503034 (Boceprevir, Schering), SCH900519 (Schering), and VX 950 (telaprevir) (Vertex).
Additional non-limiting examples of HCV polymerase inhibitors include MK-3281 (Merck), PSI-7851 (Pharmasset), IDX184 (Indenix), ANA598 (Anadys), ABT-333
30 (Abbott), VCH-916 (Vertex), PF-0086554 (Pfizer), R7128 (Pharmasset/Roche), GS 9190 (Gilead), and VCH-759 (Vertex).

Additional non-limiting examples of agents useful in the present methods and compositions include: SPC3649 (LNA-antimiRTM-122), microRNA, Santaris Pharma, CF102, (A3AR AGONISTS) (CAN-FITE), IMO-2125, TLR9 agonist, (Idera Pharmaceuticals), PYN17, Botanical, (Phynova), Bavituximab (formerly Tarvacin), Anti-
 5 Phospholipid Therapy, (Peregrine), A-831 and/or A-832 (each of which are listed as NS5A Inhibitors from ArrowTherapeutics Ltd.), BMS-790052 (NS5A inhibitors from BMS), NOV-205, Immunomodulator, (Novelos Therapeutics), CTS-1027, Anti-inflammatory, (Conatus), Oglufanide disodium, Immunomodulator, (Implicit Bioscience), Alinia (nitazoxanide), Thiazolides, (Romark Laboratories), SCV-07, Broad spectrum
 10 immune stimulator, (SciClone), MitoQ (mitoquinone), Inflammation/Fibrosis Inhibitor, (Antipodean Pharmaceuticals), DEBIO-025, Cyclophilin inhibitor, (Debio Pharm Group), SCY-635, cyclophilin inhibitor (SCYNEXIS), PF-03491390 (Formerly IDN-6556), Pancaspase Inhibitor, (Pfizer Pharmaceuticals), Civacir, HCV Immune Globulin, NABI, MX-3253 (celgosivir), Glucosidase I Inhibitor, (MIGENIX), VGX-410C (Mifepristone),
 15 IRES Inhibitor, (VGX Pharmaceuticals), Viramidine (Taribavirin), Nucleoside Analogue, (Valeant Pharmaceuticals), and ZADAXIN® (thymalfasin or thymosin alpha 1), Immunomodulator, (SciClone/Sigma-Tau).

Additional non-limiting examples of agents useful in the present methods and compositions include: TLR agonists (e.g., ANA773, Anadys Pharmaceuticals),
 20 immunomodulators (e.g., CYT107, Cytheris; oglufanide disodium, Implicit Bioscience), microRNA (e.g., SPC3649 (LNA-antimiRTM-122, Santaris Pharma), A3AR agonists (e.g., CF102, CAN-FITE), TLR9 agonists (e.g., Idera Pharmaceuticals), anti-phospholipid therapeutics (e.g., bavituximab (formerly Tarvacin), Peregrine), immunomodulators (e.g., NOV-205, Novelos Therapeutics), caspase inhibitors (e.g., GS-9450, Gilead), anti-
 25 inflammatories (e.g., CTS-1027, Conatus), thiazolides (e.g., alinia (nitazoxanide), Romark Laboratories), broad spectrim immune stimulators (e.g., SCV-07, SciClone), inflammation/fibrosis inhibitors (e.g., MitoQ (mitoquinone), Antipodean Pharmaceuticals, cyclophilin inhibitors (e.g., DEBIO-025, Debio Pharm Group), pancaspase inhibitors (e.g., PF-03491390 (formerly IDN-6556, Pfizer Pharmaceuticals),
 30 and nucleoside analogues (e.g., Viramidine (Taribavirin), Valeant Pharmaceuticals).

Interferons useful in the present methods and compositions include, but are not limited to, interferon alfa-2a, interferon alfa-2b, interferon alfacon-1 and PEG-interferon

alpha conjugates. "PEG-interferon alpha conjugates" are interferon alpha molecules covalently attached to a PEG molecule. Illustrative PEG-interferon alpha conjugates include interferon alpha-2a (RoferonTM, Hoffman La-Roche, Nutley, New Jersey) in the form of pegylated interferon alpha-2a (*e.g.*, as sold under the trade name PegasysTM),
5 interferon alpha-2b (IntronTM, from Schering-Plough Corporation) in the form of pegylated interferon alpha-2b (*e.g.*, as sold under the trade name PEG-IntronTM), interferon alpha-2c (Berofer AlphaTM, Boehringer Ingelheim, Ingelheim, Germany), interferon alpha fusion polypeptides, or consensus interferon as defined by determination of a consensus sequence of naturally occurring interferon alphas (InfergenTM, Amgen,
10 Thousand Oaks, California).

Additional examples of Interferons useful in the present methods and compositions include, but are not limited to: IL-29 (PEG-Interferon Lambda), Long acting Interferon, ZymoGenetics, Oral Interferon alpha, Oral Interferon, (Amarillo Biosciences), Belerofon (oral), Oral interferon, (Nautilus Biotech), BLX-883 (Locteron),
15 Long Acting Interferon, (Biolex Therapeutics / OctoPlus), Omega Interferon, Interferon, (Intarcia Therapeutics), Albuferon, Long Acting Interferon (injections every two weeks), (Human Genome Sciences), Consensus interferon (Infergen), and Interferon, (Three Rivers Pharma).

Antiviral antibodies (antibody therapy agents) useful in the present methods and
20 compositions include, but are not limited to, antibodies specific to IL-10 (such as those disclosed in US Patent Publication No. US2005/0101770, humanized 12G8, a humanized monoclonal antibody against human IL-10, plasmids containing the nucleic acids encoding the humanized 12G8 light and heavy chains were deposited with the American Type Culture Collection (ATCC) as deposit numbers PTA-5923 and PTA-5922,
25 respectively), and the like). Viral protease inhibitors useful in the present methods and compositions include, but are not limited to, NS3 serine protease inhibitors (including, but are not limited to, those disclosed in U.S. Patent Nos. 7,012,066, 6,914,122, 6,911,428, 6,846,802, 6,838,475, 6,800,434, 5,017,380, 4,933,443, 4,812,561 and 4,634,697; and U.S. Patent Publication Nos. US20020160962, US20050176648 and
30 US20050249702), HCV protease inhibitors (*e.g.*, SCH503034 (Schering-Plough), VX-950 (Vertex), GS-9132 (Gilead/Achillion), ITMN-191 (InterMune/Roche)), and HIV

protease inhibitors (*e.g.*, amprenavir, atazanavir, fosemprenavir, indinavir, lopinavir, ritonavir, nelfinavir, saquinavir, tipranavir and TMC114).

Viral replication inhibitors useful in the present methods and compositions include, but are not limited to, NS3 helicase inhibitors, NS5A inhibitors, ribavirin, 5 viramidine, A-831 (Arrow Therapeutics); an antisense agent or a therapeutic vaccine.

In one embodiment, viral replication inhibitors useful in the present methods and compositions include, but are not limited to, NS3 helicase inhibitors or NS5A inhibitors.

Examples of protease inhibitors useful in the present methods include, but are not limited to, an HCV protease inhibitor and a NS-3 serine protease inhibitor.

10 Examples of NS-3 serine protease inhibitors include, but are not limited to, SCH 503034 (Boceprevir, Schering), SCH900519 (Schering), Telaprevir (VX950), ITMN-191, TMC435350, GS9132, MK7009, and BILN2061.

Examples of HCV protease inhibitors useful in the present methods include, but are not limited to, those disclosed in Landro *et al.*, *Biochemistry*, 36(31):9340-9348 15 (1997); Ingallinella *et al.*, *Biochemistry*, 37(25):8906-8914 (1998); Llinàs-Brunet *et al.*, *Bioorg Med Chem Lett*, 8(13):1713-1718 (1998); Martin *et al.*, *Biochemistry*, 37(33):11459-11468 (1998); Dimasi *et al.*, *J Virol*, 71(10):7461-7469 (1997); Martin *et al.*, *Protein Eng*, 10(5):607-614 (1997); Elzouki *et al.*, *J Hepat*, 27(1):42-48 (1997); *BioWorld Today*, 9(217):4 (November 10, 1998); and International Publication Nos. WO 20 98/14181; WO 98/17679, WO 98/17679, WO 98/22496 and WO 99/07734. Additional non-limiting examples of protease inhibitors include ACH-1625 (Achillion), ABT-450 (Abbott/Enanta), BI201335 (Boehringer Ingelheim Pharma), VX-813 (Vertex), PHX1766 (Phenomix), VX-500 (Vertex), ITMN-191 (R-7227) (InterMune), MK7009 (Merck), BI 207127 (Boehringer Ingelheim), SCH900518 (Schering/Merck), TMC435 25 (Medivir/Tibotec), SCH 503034 (Boceprevir, Schering), SCH900519 (Schering), Telaprevir (VX950) and (Vertex), XTL-2125 (XTL Biopharmaceuticals).

Additional examples of other therapeutic agents useful in the present methods and compositions include vaccines. Non-limiting examples of antiviral vaccines include: ChronVac-C, DNA-based Therapeutic Vaccine, (Inovio / Tripep), TG4040, Therapeutic 30 Vaccine, (Transgene), PeviPROTM, Therapeutic vaccine, (Pevion Biotect), HCV/MF59, Vaccine(s), (Chiron/Novartis), GI-5005, Therapeutic Vaccine, (Globe Immune), IC41,

Therapeutic Vaccine, (InterCell), HCV/MF59 (Chiron/Novartis), GI-5005 (Globe Immune), and Civacir (NABI).

Additional examples of other therapeutic agents useful in the present methods and compositions include anti-cancer agents. Non-limiting examples of antiviral anti-cancer agents include: ZIO-101, Anti-Liver Cancer (Arsenic), (Ziopharm Oncology), GV1001 (Heptovax), Anti-Liver Cancer, (Pharmexa), PI-88, Anti-liver cancer, (Progen Industries), Nexavar (sorafenib), Anti-liver cancer, (Onyx Pharmaceuticals), and ThermoDox (doxorubicin), Anti-liver cancer, (Celsion). Additional non-limiting examples of viral anticancer agents include CF102 (Can-Fite BioPharma), ZIO-101 (Ziopharm Oncology), GV1001 (Heptovax) (Pharmexa), PI-88 (Progen Industries), ThermoDox (doxorubicin) (Celsion), and Nexavar (sorafenib) (Onyx Pharmaceuticals).

Additional examples of other therapeutic agents useful in the present compositions and methods include, but are not limited to, LevovirinTM (ICN Pharmaceuticals, Costa Mesa, California), VP 50406TM (Viropharma, Incorporated, Exton, Pennsylvania), ISIS 14803TM (ISIS Pharmaceuticals, Carlsbad, California), HeptazymeTM (Ribozyme Pharmaceuticals, Boulder, Colorado), VX-950TM (Vertex Pharmaceuticals, Cambridge, Massachusetts), ThymosinTM (SciClone Pharmaceuticals, San Mateo, California), MaxamineTM (Maxim Pharmaceuticals, San Diego, California), NKB-122 (JenKen Bioscience Inc., North Carolina), mycophenolate mofetil (Hoffman-LaRoche, Nutley, New Jersey).

Additional examples of other therapeutic agents useful in the present methods and compositions include adjunct therapeutics such as thrombopoietin receptor antagonists (e.g., LGD-4665, Ligand Pharmaceuticals Inc., and eltromobopag (Promacta), GlaxoSmithKline).

Additional examples of other therapeutic agents useful in the present compositions and methods include, but are not limited to: HCV/MF59, Oral Interferon alpha, Viramidine, Infergen/Consensus, JBK-122, Bavituximab (Tarvacin), Civacir, Albuferon, IL-29 (PEG-Interferon lambda), Omega Interferon, ZADAXIN® (thymalfasin or thymosin alpha 1), NOV-205, PF-03491390 (formerly IDN-6556), Nexavar, ITMN-191, IC41, VX 950 (telaprevir), R1656, MX-3253 (Celgosivir), SCH 503034 (Boceprevir, Schering), SCH900519 (Schering), Belerofon (oral), VGX-410C, ThermoDox (doxorubicin), R7128, R1626, A-831, DEBIO-025, PeviPROTM, GV1001, PYN17, PI-

88, TG4040, BLX-883 (Locteron), ChronVac-R, MitoQ, GSK625433, SOV-07, IMO-2125, Alinia (nitazoxanide), LGD-4665, Z10-101, CF102, VCH-759, VCH-916, Oglufanide disodium, VX-500, TMC435350, PF-00868554, GGI-5005 (Tarmogen), SPC3649 (LNA-antimiRTM-122), CTS-1027, ABT-333, Eltrombopag, and ANA598.

5 Additional examples of other therapeutic agents useful in the present compositions and methods include, but are not limited to adjunct therapeutics. Non-limiting examples include: LGD-4665, Thrombopoietin Receptor Agonist, (Ligand Pharmaceuticals Inc.), and Eltrombopag (Promacta), Thrombopoietin Receptor Agonist, (GlaxcoSmithKline).

The doses and dosage regimen of the other agents used in the combination
10 therapies of the present invention for the treatment or prevention of a viral infection can be determined by the attending clinician, taking into consideration the the approved doses and dosage regimen in the package insert; the age, sex and general health of the patient; and the type and severity of the viral infection or related disease or disorder. When
administered in combination, the compound(s) of the invention and the other agent(s) for
15 treating diseases or conditions listed above can be administered simultaneously (*i.e.*, in the same composition or in separate compositions one right after the other) or sequentially. This is particularly useful when the components of the combination are given on different dosing schedules, *e.g.*, one component is administered once daily and another every six hours, or when the preferred pharmaceutical compositions are different,
20 *e.g.* one is a tablet and one is a capsule. A kit comprising the separate dosage forms is therefore advantageous.

Generally, a total daily dosage of the at least one compound of the invention and the additional antiviral agent(s), when administered as combination therapy, can range from about 0.1 to about 2000 mg per day, although variations will necessarily occur
25 depending on the target of the therapy, the patient and the route of administration. In one embodiment, the dosage is from about 10 to about 500 mg/day, administered in a single dose or in 2-4 divided doses. In another embodiment, the dosage is from about 1 to about 200 mg/day, administered in a single dose or in 2-4 divided doses. In still another embodiment, the dosage is from about 1 to about 100 mg/day, administered in a single
30 dose or in 2-4 divided doses. In yet another embodiment, the dosage is from about 1 to about 50 mg/day, administered in a single dose or in 2-4 divided doses. In a further embodiment, the dosage is from about 1 to about 20 mg/day, administered in a single

dose or in 2-4 divided doses. In another embodiment, the dosage is from about 500 to about 1500 mg/day, administered in a single dose or in 2-4 divided doses. In still another embodiment, the dosage is from about 500 to about 1000 mg/day, administered in a single dose or in 2-4 divided doses. In yet another embodiment, the dosage is from about 100 to
5 about 500 mg/day, administered in a single dose or in 2-4 divided doses.

In one embodiment, when the other therapeutic agent is INTRON-A interferon alpha 2b (commercially available from Schering-Plough Corp.), this agent is administered by subcutaneous injection at 3MIU(12 mcg)/0.5mL/TIW is for 24 weeks or 48 weeks for first time treatment.

10 In another embodiment, when the other therapeutic agent is PEG-INTRON interferon alpha 2b pegylated (commercially available from Schering-Plough Corp.), this agent is administered by subcutaneous injection at 1.5 mcg/kg/week, within a range of 40 to 150 mcg/week, for at least 24 weeks.

In another embodiment, when the other therapeutic agent is ROFERON A
15 inteferon alpha 2a (commercially available from Hoffmann-La Roche), this agent is administered by subcutaneous or intramuscular injection at 3MIU(11.1 mcg/mL)/TIW for at least 48 to 52 weeks, or alternatively 6MIU/TIW for 12 weeks followed by 3MIU/TIW for 36 weeks.

In another embodiment, when the other therapeutic agent is PEGASUS interferon
20 alpha 2a pegylated (commercially available from Hoffmann-La Roche), this agent is administered by subcutaneous injection at 180mcg/1mL or 180mcg/0.5mL, once a week for at least 24 weeks.

In another embodiment, when the other therapeutic agent is INFERGEN
interferon alphacon-1 (commercially available from Amgen), this agent is administered
25 by subcutaneous injection at 9mcg/TIW is 24 weeks for first time treatment and up to 15 mcg/TIW for 24 weeks for non-responsive or relapse treatment.

In another embodiment, when the other therapeutic agent is Ribavirin
(commercially available as REBETOL ribavirin from Schering-Plough or COPEGUS
ribavirin from Hoffmann-La Roche), this agent is administered at a daily dosage of from
30 about 600 to about 1400 mg/day for at least 24 weeks.

Compositions and Administration

The compounds of the invention may be used as the neat chemical or as part of a composition, such as a pharmaceutical composition. For example, when administered to a patient, the compounds of the invention can be administered as a component of a composition that comprises a pharmaceutically acceptable carrier or vehicle. The present invention provides pharmaceutical compositions comprising an effective amount of at least one compound of the invention and a pharmaceutically acceptable carrier. In the pharmaceutical compositions and methods of the present invention, the active ingredients will typically be administered in admixture with suitable carrier materials suitably selected with respect to the intended form of administration, i.e. oral tablets, capsules (either solid-filled, semi-solid filled or liquid filled), powders for constitution, oral gels, elixirs, dispersible granules, syrups, suspensions, and the like, and consistent with conventional pharmaceutical practices. For example, for oral administration in the form of tablets or capsules, the active drug component may be combined with any oral non-toxic pharmaceutically acceptable inert carrier, such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol, ethyl alcohol (liquid forms) and the like. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. Powders and tablets may be comprised of from about 5 to about 95 percent inventive composition. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration.

Moreover, when desired or needed, suitable binders, lubricants, disintegrating agents and coloring agents may also be incorporated in the mixture. Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethylcellulose, polyethylene glycol and waxes. Among the lubricants there may be mentioned for use in these dosage forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrants include starch, methylcellulose, guar gum and the like. Sweetening and flavoring agents and preservatives may also be included where appropriate.

Liquid form preparations include solutions, suspensions and emulsions and may include water or water-propylene glycol solutions for parenteral injection.

Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas.

Also included are solid form preparations which are intended to be converted,
5 shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides or cocoa butter is first melted, and the active ingredient is dispersed
homogeneously therein as by stirring. The molten homogeneous mixture is then poured
10 into convenient sized molds, allowed to cool and thereby solidify.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

15 Additionally, the compositions of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects, i.e. anti-inflammatory activity and the like. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled
20 release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

In one embodiment, the one or more compounds of the invention are in a form suitable for oral administration.

25 In another embodiment, the one or more compounds of the invention are in a form suitable for intravenous administration.

In another embodiment, the one or more compounds of the invention are in a form suitable for topical administration.

In another embodiment, the one or more compounds of the invention are in a form
30 suitable for sublingual administration.

In one embodiment, a pharmaceutical preparation comprising at least one compound of the invention is formulated in a unit dosage form. In such form, the

preparation is subdivided into unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

Compositions can be prepared according to conventional mixing, granulating or coating methods, respectively, and the present compositions can contain, in one
5 embodiment, from about 0.1% to about 99% of the compound(s) of the invention by weight or volume. In various embodiments, the the present compositions can contain, in one embodiment, from about 1% to about 70% or from about 5% to about 60% of the compound(s) of the invention by weight or volume.

The quantity of compound(s) of the invention in a unit dose of preparation may be
10 varied or adjusted from about 0.1 mg to about 2000 mg. In various embodiment, the quantity is from about 1 mg to about 2000 mg, 100 mg to about 200 mg, 500 mg to about 2000 mg, 100 mg to about 1000 mg, and 1 mg to about 500 mg.

For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In one embodiment, the daily dosage is administered
15 in one portion. In another embodiment, the total daily dosage is administered in two divided doses over a 24 hour period. In another embodiment, the total daily dosage is administered in three divided doses over a 24 hour period. In still another embodiment, the total daily dosage is administered in four divided doses over a 24 hour period.

The amount and frequency of administration of the compound(s) of the invention
20 will be determined according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. Generally, a total daily dosage of the compound(s) of the invention range from about 0.1 to about 2000 mg per day, although variations will necessarily occur depending on the target of the therapy, the patient and the route of administration. In one
25 embodiment, the dosage is from about 1 to about 200 mg/day, administered in a single dose or in 2-4 divided doses. In another embodiment, the dosage is from about 10 to about 2000 mg/day, administered in a single dose or in 2-4 divided doses. In another embodiment, the dosage is from about 100 to about 2000 mg/day, administered in a single dose or in 2-4 divided doses. In still another embodiment, the dosage is from about 500
30 to about 2000 mg/day, administered in a single dose or in 2-4 divided doses.

The compositions of the invention can further comprise one or more additional therapeutic agents, selected from those described above. Accordingly, in one

embodiment, the present invention provides compositions comprising: (i) at least one compound of the invention or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof; (ii) one or more additional therapeutic agents that are not a compound of the invention; and (iii) a pharmaceutically acceptable carrier, wherein the amounts in the
5 composition are together effective to treat a viral infection or a virus-related disorder.

Kits

In another embodiment, the present invention provides a kit comprising a therapeutically effective amount of at least one compound of the invention, or a
10 pharmaceutically acceptable salt, solvate, ester, isomer, tautomer, or prodrug of said compound and a pharmaceutically acceptable carrier, vehicle or diluent.

In another aspect the present invention provides a kit comprising an amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester, isomer, tautomer, or prodrug of said compound and an amount of at least one additional
15 therapeutic agent listed above, wherein the amounts of the two or more ingredients result in a desired therapeutic effect.


The present invention is not to be limited by the specific embodiments disclosed in the examples that are intended as illustrations of a few aspects of the invention and any embodiments that are functionally equivalent are within the scope of this invention.
20 Indeed, various modifications of the invention in addition to those shown and described herein will become apparant to those skilled in the art and are intended to fall within the scope of the appended claims.

A number of references have been cited herein. The entire disclosures of such references are incorporated herein by reference.

CLAIMS

THEREFORE, WE CLAIM:

1. A compound, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I):



(I)

wherein each of R, R¹, X, Y, Z, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁸, R¹⁹ and n are selected independently and wherein:

R is selected from H, alkyl, aryl, heteroaryl, cycloalkyl, aryl-fused cycloalkyl, heteroaryl-fused cycloalkyl, cycloalkenyl, aryl-fused cycloalkenyl, heteroaryl-fused cycloalkenyl, heterocycloalkyl, aryl-fused heterocycloalkyl, and heteroaryl-fused heterocycloalkyl,

wherein each of said alkyl, said aryl, said heteroaryl, said cycloalkyl, said aryl-fused cycloalkyl, said heteroaryl-fused cycloalkyl, said cycloalkenyl, said aryl-fused cycloalkenyl, said heteroaryl-fused cycloalkenyl, said heterocycloalkyl, said aryl-fused heterocycloalkyl, and said heteroaryl-fused heterocycloalkyl, is unsubstituted or optionally independently substituted with from one to five substituents, which are the same or different, each substituent being independently selected from halo, -OH, -CN, oxo, alkyl, cycloalkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heterohaloalkyl,

11/0007/0

187

-alkyl-OH, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, heteroarylalkyl-, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, heterocycloalkylalkyl-, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)-cycloalkyl, -C(O)-heterocycloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -C(O)O-haloalkyl, -C(O)O-cycloalkyl, -C(O)O-heterocycloalkyl, -OC(O)-alkyl, -OC(O)-haloalkyl, -OC(O)-cycloalkyl, -OC(O)-heterocycloalkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -OC(O)NH₂, -CO(O)NHR¹⁰, -CO(O)NR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NH₂, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -S(O)₂R¹⁰, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to five substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl;

X and Y are each independently selected from N and CH, with the proviso that at least one of X or Y is N;

Z = H, halo, -OH, -SH, -CN, alkyl, alkenyl, alkynyl, heteroalkyl, haloalkyl, heterohaloalkyl, -S-alkyl, -O-alkyl, -O-aryl, -O-heteroaryl, cycloalkyl, aryl, heteroaryl, -NH₂, -NHR¹², and -NR¹²R¹³;

R¹ is selected from H, halo, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, heteroaryl, -OH, -O-alkyl, -O-aryl, -O-heteroalkyl, -O-heteroaryl, -SH, -S-alkyl, -S-aryl, -S-heteroalkyl, -S-heteroaryl, -NH₂, -NHR¹⁴, -NR¹⁴R¹⁵, -NO₂, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰;

R² (when R² is not joined with R⁹) is selected from H and alkyl;

n= 0, 1, or 2;

R³ is selected from H, -alkyl, -alkenyl, alkynyl, aryl, heteroaryl, and cycloalkyl,

11/0007/0

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, and said cycloalkyl, is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^4 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^5 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl,

-O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R⁴ and R⁵ are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

R⁶ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R⁵ and R⁶ are taken together to form a double bond;

R⁷ is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH,

110000 / 0

190

alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R^6 and R^7 are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

R^8 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R¹⁰)R¹¹, -O(C)O-NHR¹¹, -O(C)O-NH₂, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰, -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

R^9 is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or

optionally independently substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -O(C)O-N(R¹⁰)R¹¹, -O(C)O-NHR¹¹, -O(C)O-NH₂, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -N(R¹⁰)S(O)₂R¹⁰, -NHS(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, R⁸ and R⁹ are taken together with the carbon atom to which they are shown attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

each R¹⁸ (when present) is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

each R¹⁹ (when present) is selected from H, -OH, halo, -alkyl, -alkenyl, alkynyl, azido, aryl, heteroaryl, -O-alkyl, -O-alkenyl, -OC(O)-alkyl, -SH, -S-alkyl, -NH₂, -NO₂, -NHR¹⁰, -NR¹⁰R¹¹, -C(O)OH, -C(O)OR¹⁰, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -S(O)NHR¹⁰, -S(O)NR¹⁰R¹¹, -S(O)R¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, and -S(O)₂R¹⁰,


11/0000/0

192

wherein each of said -alkyl, said -alkenyl, said alkynyl, said aryl, said heteroaryl, said -O-alkyl, said -O-alkenyl, said -OC(O)-alkyl, and said -S-alkyl is unsubstituted or optionally independently substituted with from one to three substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl;

or, alternatively, n is 1 and R^{18} and R^{19} are taken together with the carbon atom to which they are attached to form a 3- to 7-membered, saturated or partially unsaturated, spirocycloalkyl ring containing from 0 to 3 spiro ring heteroatoms selected from O, N, and S;

or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1C):




wherein R^{20} and R^{21} are each independently selected from H, alkyl, and heteroalkyl and wherein R^5 and R^6 are defined above, with the proviso that when R^4 and R^7 form a moiety (1C), then R^5 and R^6 are not taken together to form a double bond;

11/0000/0

193


or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1D):



(1D),

wherein R^5 and R^6 are as defined above;

or, alternatively, R^4 and R^7 , together with the carbon atoms to which they are shown attached, form a moiety (1E):



(1E),

wherein R^5 and R^6 are as defined above;

each R^{10} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

each R^{11} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{10} and R^{11} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl;

each R^{12} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

each R^{13} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, -alkyl-OH, $-C(O)O$ alkyl, $-C(O)$ alkyl, $-C(O)NH$ alkyl, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{12} and R^{13} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R^{14} is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, alkylamino, alkylthio, heteroalkenyl, haloalkenyl, $-S(O)_2$ -alkyl, -alkyl-OH, -alkyl-O-Acyl, $-C(O)O$ alkyl, $-C(O)$ alkyl, cycloalkyl, cycloalkyl-alkyl-, heterocycloalkyl, heterocycloalkyl-alkyl-, heterocycloalkenyl, heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-, heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said alkylthio, each said heteroalkenyl, each said haloalkenyl, each said $-S(O)_2$ -alkyl, each said -alkyl-OH, each said -alkyl-O-Acyl, each said $-C(O)O$ alkyl, each said $-C(O)$ alkyl, each said cycloalkyl, each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl, and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted with from one to five substituent, which can be the same or different, each substituent being independently selected from -OH, halo, $-NH_2$, $-NHR^{10}$, $-NR^{10}R^{11}$, $-C(O)OH$, $-C(O)OR^{10}$, $-C(O)NH_2$, $-C(O)NHR^{10}$, $-C(O)NR^{10}R^{11}$, $-S(O)_2$ alkyl, $-S(O)_2$ aryl, alkyl, alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, heterohaloalkyl, aryl, cycloalkyl, and heterocycloalkyl;

each R^{15} is independently selected from alkyl, alkoxy, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, alkylamino, alkylthio, heteroalkenyl, haloalkenyl, $-S(O)_2$ -alkyl, -alkyl-OH, -alkyl-O-Acyl, $-C(O)O$ alkyl, $-C(O)$ alkyl, cycloalkyl, cycloalkyl-alkyl-, heterocycloalkyl,

heterocycloalkyl-alkyl-, heterocycloalkenyl, heterocycloalkenyl-alkyl-, aryl, aryl-alkyl-, heteroaryl, and heteroaryl-alkyl-,

wherein each said alkyl, each said alkoxy, each said alkenyl, each said haloalkyl, each said heteroalkyl, each said heterohaloalkyl, each said alkylamino, each said alkylthio, each said heteroalkenyl, each said haloalkenyl, each said $-S(O)_2$ -alkyl, each said $-alkyl-OH$, each said $-alkyl-O-Acyl$, each said $-C(O)Oalkyl$, each said $-C(O)alkyl$, each said cycloalkyl, each said cycloalkyl-alkyl-, each said heterocycloalkyl, each said heterocycloalkyl-alkyl-, each said heterocycloalkenyl, each said heterocycloalkenyl-alkyl-, each said aryl, each said aryl-alkyl-, each said heteroaryl, and each said heteroaryl-alkyl-, is unsubstituted or optionally independently substituted with from one to five substituent, which can be the same or different, each substituent being independently selected from $-OH$, halo, $-NH_2$, $-NHR^{10}$, $-NR^{10}R^{11}$, $-C(O)OH$, $-C(O)OR^{10}$, $-C(O)NH_2$, $-C(O)NHR^{10}$, $-C(O)NR^{10}R^{11}$, $-S(O)_2alkyl$, $-S(O)_2aryl$, alkyl, alkoxy, haloalkyl, haloalkoxy, heteroaryl, heteroalkyl, heterohaloalkyl, aryl, cycloalkyl, and heterocycloalkyl;

or, alternatively, R^{14} and R^{15} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- to 6-membered heterocycloalkyl;

each R^{16} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, $-alkyl-OH$, $-C(O)Oalkyl$, $-C(O)alkyl$, $-C(O)NHalkyl$, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl; and

each R^{17} is independently selected from alkyl, alkenyl, haloalkyl, heteroalkyl, heterohaloalkyl, $-S(O)_2$ -alkyl, $-alkyl-OH$, $-C(O)Oalkyl$, $-C(O)alkyl$, $-C(O)NHalkyl$, $-C(O)N(alkyl)_2$, cycloalkyl, heterocycloalkyl, heterocycloalkenyl, aryl, and heteroaryl;

or, alternatively, R^{16} and R^{17} are linked together with the nitrogen to which they are attached to form an unsubstituted or substituted 4- or 6-membered heterocycloalkyl.

2. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

110000/0

R is selected from aryl, heteroaryl, benzo-fused heteroaryl, cycloalkyl, cycloalkenyl, benzo-fused cycloalkyl, benzo-fused cycloalkenyl, heterocycloalkyl, and benzo-fused heterocycloalkyl,

wherein each of said alkyl, said aryl, said heteroaryl, said benzo-fused heteroaryl, said cycloalkyl, said cycloalkenyl, said heterocycloalkyl, said heterocycloalkenyl, and said benzo-fused heterocycloalkyl is unsubstituted or optionally independently substituted with from one to three substituents, which are the same or different, each substituent being independently selected from halo, -OH, -CN, alkyl, cycloalkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heterohaloalkyl, -alkyl-OH, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl;

X is N;

Y is N;

R² is H;

R¹ is selected from -NH₂, -NHR¹⁴, and -NR¹⁴R¹⁵; and

Z is selected from H, halo, alkyl, -OH, haloalkyl, and cycloalkyl.

3. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

X is N;


110000/0

197

Y is N;

R is selected from the group consisting of:

(a) an unsubstituted or substituted monocyclic aryl moiety or an unsubstituted or substituted heteroaryl moiety selected from the group consisting of:




wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a, R_b, R_c, R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and

110000/0

said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl, and

(b) an unsubstituted or an substituted bicyclic heteroaryl moiety selected from the group consisting of:



wherein the wavy line represents the point of attachment of R to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, cycloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰,

11000070

199

$-\text{C}(\text{O})\text{NR}^{10}\text{R}^{11}$, $-\text{C}(\text{O})\text{ONH}_2$, $-\text{C}(\text{O})\text{ONHR}^{10}$, $-\text{C}(\text{O})\text{ONR}^{10}\text{R}^{11}$, $-\text{NH}_2$, $-\text{NHR}^{10}$, $-\text{NR}^{10}\text{R}^{11}$, $-\text{NO}_2$, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, $-\text{O}$ -alkyl, and $-\text{C}(\text{O})\text{Oalkyl}$;

R^1 is selected from the group consisting of:

(a) $-\text{NH}_2$,





FIGURE 10

200

Bu), -O-haloalkyl (non-limiting examples of which include -O-CH₂F, -O-CHF₂, and -O-CF₃), -NH₂, -NHalkyl, and -N(alkyl)₂,




wherein the wavy line represents the point of attachment of R¹ to the rest of the molecule, and wherein R_{af} is selected from H and acetyl,

119000 / 0

201

(d)




wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may

110000 / 0

202


be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected from alkyl and haloalkyl,



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each R_{aa} is independently selected from haloalkyl (non-limiting examples of which include -CH₂F, -CHF₂, -CF₃, etc.), R_{ab} is selected from OH, OAc, and -O-alkyl (non-limiting examples of which include -O-Me, -O-Et, -O-*n*-Pr, -O-*i*-Pr, -O-*n*-Bu, -O-*i*-Bu, and -O-*t*-Bu), -O-haloalkyl (non-limiting examples of which include -O-CH₂F, -O-CHF₂, and -O-CF₃), -NH₂, -NHalkyl, and -N(alkyl)₂,


FIGURE 10

203



wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein R_{af} is selected from H and acetyl,


(g)



11/0000/0


204

wherein the wavy line represents the point of attachment of R^1 to the rest of the molecule, and wherein each of R_a , R_b , R_c , R_d , and R_e , is independently selected from H, halo, -OH, -CN, alkyl, haloalkyl, -alkyl-OH, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-haloalkyl, -O-alkyl-OH, aryl, -O-aryl, -S-aryl, -O-alkyl-aryl, -S-alkyl-aryl, heteroaryl, -O-heteroaryl, -S-heteroaryl, -O-alkyl-heteroaryl, -S-alkyl-heteroaryl, heterocycloalkyl, -C(O)-alkyl, -C(O)-haloalkyl, -C(O)H, -C(O)OH, -C(O)O-alkyl, -OC(O)-alkyl, -C(O)NH₂, -C(O)NHR¹⁰, -C(O)NR¹⁰R¹¹, -C(O)ONH₂, -C(O)ONHR¹⁰, -C(O)ONR¹⁰R¹¹, -NH₂, -NHR¹⁰, -NR¹⁰R¹¹, -NO₂, substituted aryl, and substituted heteroaryl, wherein each of said substituted aryl and said substituted heteroaryl independently contains from one to three substituents, which may be the same or different, each substituent being independently selected from halo, alkyl, -O-alkyl, and -C(O)Oalkyl, and wherein each R_{ad} and each R_{ae} is independently selected from alkyl and haloalkyl, and



Z is selected from the group consisting of H, halo, -OH, -SH, -CN, alkyl, alkenyl, alkynyl, heteroalkyl, haloalkyl, heterohaloalkyl, -S-alkyl, -O-alkyl, -O-aryl, -O-heteroaryl, cycloalkyl, aryl, heteroaryl, -NH₂, -NHR¹², and -NR¹²R¹³.

4. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I.a):



(I.a)

wherein each of R, R¹, X, Y, Z, R², R⁴, R⁵, R⁷, and R⁹ is selected independently and defined in Claim 1.

5. A compound of Claim 4, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

n is 1;

R² is H;

R⁴ and R⁷ are each independently selected from H and OH;

R⁵ is selected from H, halo, and alkyl; and

R⁹ is alkyl, wherein said alkyl is unsubstituted or substituted with from one to five substituents, which can be the same or different, each substituent being independently selected from halo, -OH, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, -O-alkyl, -O-cycloalkyl, -O-alkenyl, -O-haloalkyl, -O-haloalkenyl, -OC(O)-alkyl, -OC(O)-alkenyl, -OC(O)-haloalkyl, -OC(O)-haloalkenyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -C(O)O-alkenyl, -C(O)O-haloalkyl, -C(O)O-haloalkenyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, -S(O)₂NR¹⁰R¹¹, -CN, -NH₂, -NHR¹⁶, and -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl.

6. A compound of Claim 4, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

n is 1;

R² is H;

R⁴ is OH;

R⁷ is OH;

R⁵ is H; and

R⁹ is alkyl, wherein said alkyl is unsubstituted or substituted with from one to five groups independently selected from -OH, halo, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰,

110000/0

206

-N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -Ocycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

7. A compound of Claim 4, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

n is 1,

R² is H,

R⁴ is OH;

R⁷ is OH;

R⁵ is -CH₃, and

R⁹ is methyl, wherein said methyl is unsubstituted or substituted with from one to three groups independently selected from -OH, halo, alkyl, -CN, -NH₂, -NHR¹⁶, -NR¹⁶R¹⁷, -NHS(O)₂R¹⁰, -N(R¹⁰)S(O)₂R¹⁰, -Oalkyl, -Ocycloalkyl, -O-alkyl-cycloalkyl, -OC(O)-alkyl, -O(C)O-NHR¹⁰, -O(C)O-N(R¹⁰)R¹¹, -C(O)O-alkyl, -S(O)₂R¹⁰, -SR¹⁰, -S(O)₂NHR¹⁰, and -S(O)₂NR¹⁰R¹¹.

8. A compound of Claim 4, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

X and Y are each N;

R is selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted benzo-fused heteroaryl, each of said substituents being independently selected from the group consisting of alkyl, -O-alkyl, and cycloalkyl;

Z is selected from H, halo, alkyl, fluoroalkyl, cyclopropyl, and -NH₂;


R¹ is selected from -NH₂, -NHR¹⁴, and -NR¹⁴R¹⁵; and

R² is selected from H.

110000 / 0


207

9. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I.a.10.j):



(I.a.10.j)

wherein CB is a moiety selected from the group consisting of:



wherein each R¹⁰ is independently selected from the group consisting of methyl, ethyl, and cyclopropyl.


11/0000/0

208

10. A compound of claim 9, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, wherein:

X is N; Y is N; R^2 is H; and Z is selected from the group consisting of H, methyl, and chloro.

11. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I.C):




(I.C)

wherein each of R, R¹, X, Y, Z, R², R³, R⁵, R⁶, R⁸, R⁹, R¹⁸, R¹⁹, R²⁰, R²¹, and n is selected independently and defined in Claim 1, with the proviso that R⁵ and R⁶ are not taken together to form a double bond.

110000 / 0

209

12. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I.D):




(I.D)

wherein each of R, R¹, X, Y, Z, R², R³, R⁵, R⁶, R⁸, R⁹, R¹⁸, and R¹⁹, and n is selected independently and defined in Claim 1.


13. A compound of Claim 1, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (I.E):

210



(I.E)

14. A compound, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound having the general structure shown in Formula (II):







(II)

wherein each of R, R¹, X, Y, Z, R³, R⁴, R⁵, R⁶, R⁷, and R⁸ is selected independently and defined in Claim 1.

11/0000/0






211

15. A compound, or an isomer, ester, or tautomer of such compound, or a pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, said compound selected from the group consisting of:

Compd #	Structure
1	
2	
3	
4	






11/0000/0

212

Compd #	Structure
5	
7	
8	
9	
10	





11/0000/0

213

Compd #	Structure
11	
12	
13	
14	
15	

1180008 / U

214

Compd #	Structure
16	
17	
18	
19	






11/0000/0

215

Compd #	Structure
20	 <chem>COc1cc(C#CC2=NC(Cl)=NC(N)=N2NC[C@H]3C[C@@H](O)[C@H](O)C3)cnc1</chem>
21	 <chem>CCOc1cc(C#CC2=NC(Cl)=NC(N)=N2NC[C@H]3C[C@@H](O)[C@H](O)C3)cnc1</chem>
22	 <chem>COc1cc(C#CC2=NC(Cl)=NC(N)=N2NC[C@H]3C[C@@H](O)[C@H](O)C3)cnc1OC</chem>
23	 <chem>CCOc1cc(C#CC2=NC(Cl)=NC(N(C)C)=N2NC[C@H]3C[C@@H](O)[C@H](O)C3)cnc1</chem>
24	 <chem>COc1cc(C#CC2=NC(Cl)=NC(N(C)C)=N2NC[C@H]3C[C@@H](O)[C@H](O)C3)cnc1</chem>

11/0000/0

216

Compd #	Structure
25	
26	
27	
28	
29	






110000/0

217

Compd #	Structure
30	 <chem>CN1C=CN(C1)C#CC2=NC(Cl)=NC(N)=C2N[C@@H]3[C@H](CO)[C@H](O)[C@H]3O</chem>
31	 <chem>C1=CC=C(C=C1S)C2=NC(Cl)=NC(N)=C2N[C@@H]3[C@H](CO)[C@H](O)[C@H]3O</chem>
32	 <chem>COC(=O)C1=NC(N)=NC=C1C#CC2=NC(Cl)=NC(N)=C2N[C@@H]3[C@H](CO)[C@H](O)[C@H]3O</chem>
33	 <chem>COC1=CC=C(C=C1)C#CC2=NC(Cl)=NC(N)=C2N[C@@H]3[C@H](CO)[C@H](O)[C@H]3O</chem>
34	 <chem>C1=CC=C(C=C1S)C2=NC(Cl)=NC(N)=C2N[C@@H]3[C@H](CO)[C@H](O)[C@H]3O</chem>






11/0000/0

218

Compd #	Structure
35	
36	
37	
38	
39	






110000/0

219

Compd #	Structure
40	
41	
42	
43	
44	





11/0000/0

220

Compd #	Structure
45	
46	
47	
48	
49	





110006 / 0

221

Compd #	Structure
50	
51	
52	 HCl
53	

11/0007/0

222

Compd #	Structure
54	
55	
56	
57	

11/0008/0

223

Compd #	Structure
58	 <chem>NC1=NC=C(NC1C#CC2=CC=CC=C2S(=O)(=O)N)N=C(N)N=C2C</chem>
59	 <chem>CC(=O)S(=O)(=O)C1=CC=CC=C1C#CC2=CC=CC=C2N=C(N)N=C2N</chem>
60	 <chem>NC1=NC=C(NC1C#CC2=CC=CC=C2Cl)N=C(N)N=C2N</chem>
61	 <chem>CO[C@@H]1C[C@H](O)[C@H](O)[C@H]1NC2=NC=C(N)N=C2C#CC3=CC=CS3</chem>
62	 <chem>CN(C)C1=NC=C(NC1C#CC2=CC=CC=C2C(=O)OC3=CC=CC=N3)N=C(N)N=C2N</chem>






11/000/0

224

Compd #	Structure
63	 <chem>Cc1cc(ccc1c2nc(N)nc(Cl)c2N[C@@H]3C[C@H](CO)[C@@H](O)[C@H]3O)c4cc5ccsc5n4</chem>
64	 <chem>Cc1cc(ccc1c2nc(N)nc(Cl)c2N[C@@H]3C[C@H](CO)[C@@H](O)[C@H]3O)c4cc(CS)cc4</chem>
65	 <chem>CN(C)c1nc(NC[C@@H]2C[C@H](CO)[C@@H](O)[C@H]2O)c(Cl)c(NC3=CC=C(C)C(OC)=N3)c1</chem>
66	 <chem>CN(C)c1nc(NC[C@@H]2C[C@H](CO)[C@@H](O)[C@H]2O)c(Cl)c(NC3=CC=C(C)C(S)=N3)c1</chem>

110008/0

225

Compd #	Structure
67	
68	
69	
70	
71	





11/0000/0

226

Compd #	Structure
72	
73	
74	
75	



11/000/0

227

Compd #	Structure
76	
77	
78	
79	 , and

11/0000/0

228



11/000/0

229

Compd #	Structure
84	 <chem>COc1ccc(cc1)C#CC2=CN(C(=N2)NCCS)N[C@@H]3[C@H](O)[C@H](O)[C@@H](O)CO3</chem>
85	 <chem>CCOC1=CC=CC=C1C#CC2=CN(C(=N2)NCC(C)(C)C)N[C@@H]3[C@H](O)[C@H](O)[C@@H](O)CO3</chem>
86	 <chem>CCOC1=CC=CC=C1C#CC2=CN(C(=N2)NCC(F)(F)F)N[C@@H]3[C@H](O)[C@H](O)[C@@H](O)CO3</chem>
87	 <chem>CCOC1=CC=CC=C1C#CC2=CN(C(=N2)NCCC)N[C@@H]3[C@H](O)[C@H](O)[C@@H](O)CO3</chem>





11/000/0

230

Compd #	Structure
88	 <chem>CCOC1=CC=C(C#CC2=C(C)N(C(=N2)NCC)NC[C@H]3O[C@@H](CO)[C@H](O)[C@H]3O)C=C1</chem>
89	 <chem>CCOC1=CC=C(C#CC2=C(C)N(C(=N2)NCC(F)C)NC[C@H]3O[C@@H](CO)[C@H](O)[C@H]3O)C=C1</chem>
90	 <chem>COc1ccc(C#CC2=C(C)N(C(=N2)NCC(F)(F)F)NC[C@H]3O[C@@H](CO)[C@H](O)[C@H]3O)cc1</chem>
91	 <chem>COc1ccc(C#CC2=C(C)N(C(=N2)NCC)NC[C@H]3O[C@@H](CO)[C@H](O)[C@H]3O)cc1</chem>






1180006/0

231

Compd #	Structure
92	
93	
94	
95	

11/0000/0

232

Compd #	Structure
96	
97	
98	
99	
100	

11/0008/0

233

Compd #	Structure
101	
102	
103	
104	
105	

16. A pharmaceutical composition comprising at least one compound according to any one of claims 1-15, or an isomer, ester, or tautomer of such compound, or a


pharmaceutically acceptable salt, solvate, or prodrug of said compound, isomer, ester, or tautomer, and a pharmaceutically acceptable carrier.

17. A pharmaceutical composition of Claim 16, further comprising at least one additional therapeutic agent selected from the group consisting of: an HCV polymerase inhibitor, an interferon, a viral replication inhibitor, an antisense agent, a therapeutic vaccine, a viral protease inhibitor, a virion production inhibitor, an immunosuppressive agent, an antiviral antibody, a CYP-450 inhibitor, an antiviral booster, and an antiviral sensitizer.

18. A method of treating a viral infection or a virus-related disorder comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one compound according to any one of Claims 1-15.

19. A method of Claim 18, wherein said viral infection or virus-related disorder is an HCV infection.

20. A method of Claim 19, further comprising administering a therapeutically effective amount of at least one additional therapeutic agent useful for treating a viral infection or a virus-related disorder, wherein said at least one additional therapeutic agent is selected from the group consisting of: an HCV polymerase inhibitor, an interferon, a viral replication inhibitor, an antisense agent, a therapeutic vaccine, a viral protease inhibitor, a virion production inhibitor, an immunosuppressive agent, an antiviral antibody, a CYP-450 inhibitor, an antiviral booster, and an antiviral sensitizer.



(I)