



US 20130295022A1

(19) **United States**(12) **Patent Application Publication**
Friedman et al.(10) **Pub. No.: US 2013/0295022 A1**(43) **Pub. Date: Nov. 7, 2013**(54) **MOISTURIZING FOAM CONTAINING
LANOLIN**(71) Applicant: **Foamix Ltd.**, Rehovot, IL (US)(72) Inventors: **Doron Friedman**, Karmei Yosef (IL);
Alex Besonov, Rehovot (IL); **Dov
Tamarkin**, Macabim (IL); **Meir Eini**,
Ness Ziona (IL)(21) Appl. No.: **13/911,470**(22) Filed: **Jun. 6, 2013****Related U.S. Application Data**(63) Continuation of application No. 11/099,942, filed on
Apr. 6, 2005, now Pat. No. 8,486,376, which is a con-
tinuation-in-part of application No. PCT/IB0300/
005527, filed on Oct. 24, 2003, said application No.
11/099,942 is a continuation-in-part of application No.
10/911,367, filed on Aug. 4, 2004.(60) Provisional application No. 60/492,546, filed on Aug.
4, 2003, provisional application No. 60/492,385, filed
on Aug. 4, 2003.(30) **Foreign Application Priority Data**

Oct. 25, 2002 (IL) 152486

Publication Classification(51) **Int. Cl.****A61K 9/12** (2006.01)**A61K 45/06** (2006.01)**A61K 31/575** (2006.01)(52) **U.S. Cl.**CPC **A61K 9/122** (2013.01); **A61K 31/575**
(2013.01); **A61K 45/06** (2013.01)USPC . **424/43**; 514/170; 424/62; 514/1.1; 424/617;
424/642; 424/646; 424/683; 424/665(57) **ABSTRACT**

The present invention provides a highly tolerable, hydrating lanolin containing foamable composition for administration to the skin, body surface, body cavity or mucosal surface, e.g., the mucosa of the nose, mouth, eye, ear, respiratory system, vagina or rectum. The foamable composition includes lanolin, a surface-active agent, about 0.01% to about 5% by weight of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent and a phase change agent, water, and a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the total composition.



Figure 1

Figure 2

MOISTURIZING FOAM CONTAINING LANOLIN

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part application of co-pending International Patent Application No. IB03/005527 designating the United States and filed on Oct. 24, 2003, which claims the benefit of priority under 35 U.S.C. §119(e) to U.S. Patent Application Ser. No. 60/492,546, filed on Nov. 29, 2002, both entitled “Cosmetic and Pharmaceutical Foam,” and which also claims the benefit of priority under 35 USC §119(a) to Israeli Patent Appl. No. 152486, filed Oct. 25, 2002, all of which are hereby incorporated in their entirety by reference.

[0002] This application is a continuation-in-part application of co-pending U.S. patent application Ser. No. 10/911,367, filed on Aug. 4, 2004, which claims the benefit of priority under 35 U.S.C. §119(e) to U.S. Patent Application Ser. No. 60/492,385, filed on Aug. 4, 2003, both entitled “Foam Carrier Containing Amphiphilic Copolymer Gelling Agent” and both hereby incorporated in their entirety by reference.

BACKGROUND OF THE INVENTION

[0003] Foams and, in particular, foam emulsions are complex dispersion systems which do not form under all circumstances. Slight shifts in foam emulsion composition, such as by the addition of active ingredients, may destabilize the foam.

[0004] There have been a few attempts to create foams containing lanolin.

[0005] U.S. Pat. No. 5,679,324 pertains to an aerosol foamable fragrance composition, translucent in its pre-dispensed state, which forms a fast-breaking foam. The composition contains a surfactant selected from the group consisting of ethoxylated lanolin oil derivatives, propoxylated lanolin oil derivatives, and mixtures thereof, a propellant, a fragrance, a thickener, and a cosmetic vehicle (preferably water). Emollients may be included, however, being translucent, the composition cannot comprise significant oil concentrations (which would make it opaque). Apparently the foam breaks spontaneously upon discharging from an aerosol container (with no need of any rubbing or sheer force application), thus making it impractical for certain applications such as intravaginal applications.

SUMMARY OF THE INVENTION

[0006] The present invention provides a highly tolerable, hydrating lanolin containing foamable composition product for administration to the skin, a body surface, a body cavity or mucosal surface, e.g., the mucosa of the nose, mouth, eye, ear, respiratory system, vagina or rectum (the “target site”). The foamable composition includes lanolin, a surface-active agent, about 0.01% to about 5% by weight of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent and a phase change agent, water, and a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the total composition. The hydrating lanolin foam composition absorbs and stores water from the environment to thereby hydrate the target site.

[0007] According to further embodiments of the present invention, the foamable composition further contains at least

one organic carrier selected from the group consisting of a hydrophobic organic carrier, a polar solvent, an emollient and mixtures thereof, wherein the at least one organic carrier is present in an amount selected from the group consisting of about 2% to about 5%, about 5% to about 10%, about 10% to about 20%, and about 20% to about 50%.

[0008] According to further embodiments of the present invention, lanolin is selected from the group consisting of lanolin, wool fat, lanolin wax and hydrophobic derivatives thereof.

[0009] According to further embodiments of the present invention, there is provided a method of treating, alleviating or preventing a disorders of the skin, a body cavity or mucosal surface, wherein the disorder involves insufficient hydration as one of its etiological factors. The method includes administering topically to a subject having the disorder, a foamed composition including lanolin, about 0.1% to about 5% by weight of a surface-active agent, about 0.01% to about 5% by weight of a polymeric additive selected from a bioadhesive agent, a gelling agent, a film forming agent and a phase change agent, and water.

[0010] According to still further embodiments of method according to the present invention, the composition further includes at least one additional therapeutic agent.

[0011] According to yet further embodiments of method according to the present invention, the organic carrier includes at least one therapeutically active oil.

BRIEF DESCRIPTION OF THE DRAWING

[0012] The invention is described with reference to the figures, which are presented for the purpose of illustration and are not intended to be limiting of the invention.

[0013] FIG. 1 is a plot of hydration value vs. time for two foamed lanolin compositions, illustrating the skin hydrating effect of two lanolin-containing foams.

[0014] FIG. 2 is a bar graph illustrating the difference in skin hydration between two lanolin-containing foams and “no treatment”.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention provides a highly tolerable, hydrating lanolin-containing foamable composition product.

[0016] All % values are provided on a weight (w/w) basis.

[0017] According to one or more embodiments of the present invention, the foamable lanolin containing composition for administration to the skin, a body surface, a body cavity or mucosal surface, e.g., the mucosa of the nose, mouth, eye, ear, respiratory system, vagina or rectum (severally and interchangeably termed herein “target site”) includes:

[0018] (1) lanolin;

[0019] (2) about 0.1% to about 5% by weight of a surface-active agent;

[0020] (3) about 0.01% to about 5% by weight of at least one polymeric agent selected from a bioadhesive agent, a gelling agent, a film forming agent and a phase change agent; and

[0021] (4) a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the total composition.

[0022] Water and optional ingredients are added to complete the total mass to 100%. Upon release from an aerosol

container, the foamable composition forms an expanded foam suitable for topical administration.

[0023] Lanolin, also termed “wool fat” and “wool fat alcohols” is a hydrophobic greasy, substance extracted from wool, produced from the oil glands of sheep. Lanolin wax is a high melting wax fraction separated from lanolin, and white lanolin is obtained by further purifications to obtain desired lanolin fractions or derivatives.

[0024] Natural lanolin is produced, among other methods, by extraction from the wool by kneading with water, with which it readily forms an emulsion. On heating, it separates as a distinct layer at the surface of the liquid. Purification is effected by repeated treatment with water in a centrifugal machine, or by some other suitable process.

[0025] The constituents of lanolin include cholesterol and ischolesterol, together with various esters of cholesterol and/or ischolesterol with one or more fatty acids, including, but not limited to lanoceric acid, lanopalmitic acid, carnaubic acid, myristic acid, oleic acid, cerotic acid and palmitic acid.

[0026] In one or more embodiments of the present invention, lanolin is a synthetic lanolin or a synthetic lanolin derivative. In the context of the present invention, a synthetic lanolin or a synthetic lanolin derivative is a hydrophobic derivative of lanolin, which is the result of a chemical or biochemical modification of natural lanolin. Synthetic lanolin or a synthetic lanolin derivative may consist of cholesterol or ischolesterol covalently linked with an organic moiety. By way of non-limiting examples, synthetic lanolin derivatives include organic alcohols derived from the hydrolysis of lanolin, esters of lanolin with an aliphatic acid (e.g., acetylated lanolin), lanolin acid (a mixture of organic acids obtained from the hydrolysis of lanolin), esters of lanolin acid with aliphatic alcohols (e.g., isopropyl lanolate and cetyl lanolate).

[0027] The presence of lanolin may be determined by detection of cholesterol, a compound making up lanolin. To test, one gram of a fat is dissolved in 3 or 4 mL of acetic anhydride and 0.3 mL of sulfuric acid, resulting in a pink coloration, further changing to green and blue.

[0028] Optionally, the composition further contains at least one organic carrier selected from a hydrophobic organic carrier, a polar solvent, an emollient and mixtures thereof, at a concentration of about 2% to about 5%, or about 5% to about 10%, or about 10% to about 20%, or about 20% to about 50% by weight.

[0029] A “hydrophobic organic carrier” as used herein refers to a material having solubility in distilled water at ambient temperature of less than about 1 gm per 100 mL, more preferable less than about 0.5 gm per 100 mL, and most preferably less than about 0.1 gm per 100 mL. It is liquid at ambient temperature. The identification of a hydrophobic organic carrier or “hydrophobic solvent”, as used herein, is not intended to characterize the solubilization capabilities of the solvent for any specific active agent or any other component of the foamable composition. Rather, such information is provided to aid in the identification of materials suitable for use as a hydrophobic carrier in the foamable compositions described herein.

[0030] In one or more embodiments, the hydrophobic organic carrier is an oil, such as mineral oil. Mineral oil (Chemical Abstracts Service Registry number 8012-95-1) is a mixture of aliphatic, naphthalenic, and aromatic liquid hydrocarbons that derive from petroleum. It is typically liquid; its viscosity is in the range of between about 35 CST and about

100 CST (at 40° C.), and its pour point (the lowest temperature at which an oil can be handled without excessive amounts of wax crystals forming so preventing flow) is below 0° C. In one or more embodiments, the term hydrophobic organic carrier does not include thick or semi-solid materials, such as white petrolatum, also termed “Vaseline”, which, in certain compositions is disadvantageous due to its waxy nature and semi-solid texture.

[0031] According to one or more embodiments, hydrophobic solvents are liquid oils originating from vegetable, marine or animal sources. Suitable liquid oil includes saturated, unsaturated or polyunsaturated oils. By way of example, the unsaturated oil may be olive oil, corn oil, soybean oil, canola oil, cottonseed oil, coconut oil, sesame oil, sunflower oil, borage seed oil, syzigium aromaticum oil, hempseed oil, herring oil, cod-liver oil, salmon oil, flaxseed oil, wheat germ oil, evening primrose oils or mixtures thereof, in any proportion.

[0032] Suitable hydrophobic solvents also include polyunsaturated oils containing poly-unsaturated fatty acids. In one or more embodiments, said unsaturated fatty acids are selected from the group of omega-3 and omega-6 fatty acids. Examples of such polyunsaturated fatty acids are linoleic and linolenic acid, gamma-linoleic acid (GLA), eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Such unsaturated fatty acids are known for their skin-conditioning effect, which contribute to the therapeutic benefit of the present foamable composition. Thus, the hydrophobic solvent can include at least 6% of an oil selected from omega-3 oil, omega-6 oil, and mixtures thereof. In the context of the present invention, oils that possess therapeutically beneficial properties are termed “therapeutically active oil”.

[0033] Another class of hydrophobic solvents is the essential oils, which are also considered therapeutically active oil, which contain active biologically occurring molecules and, upon topical application, exert a therapeutic effect, which is conceivably synergistic to the beneficial effect of the NSAID in the composition.

[0034] Another class of therapeutically active oils includes liquid hydrophobic plant-derived oils, which are known to possess therapeutic benefits when applied topically.

[0035] Silicone oils also may be used and are desirable due to their known skin protective and occlusive properties. Suitable silicone oils include non-volatile silicones, such as polyalkyl siloxanes, polyaryl siloxanes, polyalkylaryl siloxanes and polyether siloxane copolymers, polydimethylsiloxanes (dimethicones) and poly(dimethylsiloxane)-(diphenylsiloxane) copolymers. These are chosen from cyclic or linear polydimethylsiloxanes containing from about 3 to about 9, preferably from about 4 to about 5, silicon atoms. Volatile silicones such as cyclomethicones can also be used. Silicone oils are also considered therapeutically active oil, due to their barrier retaining and protective properties.

[0036] In one or more embodiments, the hydrophobic carrier includes at least 2% by weight silicone oil or at least 5% by weight.

[0037] The solvent may be a mixture of two or more of the above hydrophobic solvents in any proportion.

[0038] A further class of solvents includes “emollients” that have a softening or soothing effect, especially when applied to body areas, such as the skin and mucosal surfaces. Emollients are not necessarily hydrophobic. Examples of suitable emollients include hexyleneglycol, propylene glycol, isostearic acid derivatives, isopropyl palmitate, isopropyl

isostearate, diisopropyl adipate, diisopropyl dimerate, maleated soybean oil, octyl palmitate, cetyl lactate, cetyl ricinoleate, tocopheryl acetate, acetylated lanolin alcohol, cetyl acetate, phenyl trimethicone, glyceryl oleate, tocopheryl linoleate, wheat germ glycerides, arachidyl propionate, myristyl lactate, decyl oleate, propylene glycol ricinoleate, isopropyl lanolate, pentaerythrityl tetrastearate, neopentylglycol dicaprylate/dicaprate, isononyl isononanoate, isotridecyl isononanoate, myristyl myristate, triisocetyl citrate, octyl dodecanol, sucrose esters of fatty acids, octyl hydroxystearate and mixtures thereof.

[0039] According to one or more embodiments of the present invention, the hydrophobic organic carrier includes a mixture of a hydrophobic solvent and an emollient. According to one or more embodiments, the foamable composition is a mixture of mineral oil and an emollient in a ratio between 2:8 and 8:2 on a weight basis.

[0040] A "polar solvent" is an organic solvent, typically soluble in both water and oil. Examples of polar solvents include polyols, such as glycerol (glycerin), propylene glycol, hexylene glycol, diethylene glycol, propylene glycol n-alkanols, terpenes, di-terpenes, tri-terpenes, terpen-ols, limonene, terpene-ol, 1-menthol, dioxolane, ethylene glycol, other glycols, sulfoxides, such as dimethylsulfoxide (DMSO), dimethylformamide, methyl dodecyl sulfoxide, dimethylacetamide, monooleate of ethoxylated glycerides (with 8 to 10 ethylene oxide units), azone (1-dodecylazacycloheptan-2-one), 2-(n-nonyl)-1,3-dioxolane, esters, such as isopropyl myristate/palmitate, ethyl acetate, butyl acetate, methyl propionate, capric/caprylic triglycerides, octylmyristate, dodecyl-myristate; myristyl alcohol, lauryl alcohol, lauric acid, lauryl lactate ketones; amides, such as acetamide oleates such as triolein; various alkanolic acids such as caprylic acid; lactam compounds, such as azone; alkanols, such as dialkylamino acetates, and admixtures thereof.

[0041] According to one or more embodiments, the polar solvent is a polyethylene glycol (PEG) or PEG derivative that is liquid at ambient temperature, including PEG200 (MW (molecular weight) about 190-210 kD), PEG300 (MW about 285-315 kD), PEG400 (MW about 380-420 kD), PEG600 (MW about 570-630 kD) and higher MW PEGs such as PEG 4000, PEG 6000 and PEG 10000 and mixtures thereof.

[0042] According to one or more embodiments, the foamable composition is substantially alcohol-free, i.e., free of short chain alcohols. Short chain alcohols, having up to 5 carbon atoms in their carbon chain skeleton and one hydroxyl group, such as ethanol, propanol, isopropanol, butanol, isobutanol, t-butanol and pentanol, are considered less desirable solvents or polar solvents due to their skin-irritating effect. Thus, the composition is substantially alcohol-free and includes less than about 5% final concentration of lower alcohols, preferably less than about 2%, more preferably less than about 1%.

[0043] The polymeric agent serves to stabilize the foam composition and to control drug residence in the target organ. Exemplary polymeric agents, are classified below in a non-limiting manner. In certain cases, a given polymer can belong to more than one of the classes provided below.

[0044] In one or more embodiments, the composition of the present invention includes at least one gelling agent. A gelling agent controls the residence of a therapeutic composition in the target site of treatment by increasing the viscosity of the

composition, thereby limiting the rate of its clearance from the site. Many gelling agents are known in the art to possess mucoadhesive properties.

[0045] The gelling agent can be a natural gelling agent, a synthetic gelling agent and an inorganic gelling agent. Exemplary gelling agents that can be used in accordance with one or more embodiments of the present invention include, for example, naturally-occurring polymeric materials, such as locust bean gum, sodium alginate, sodium caseinate, egg albumin, gelatin agar, carrageenin gum, sodium alginate, xanthan gum, quince seed extract, tragacanth gum, guar gum, starch, chemically modified starches and the like, semi-synthetic polymeric materials such as cellulose ethers (e.g. hydroxyethyl cellulose, methyl cellulose, carboxymethyl cellulose, hydroxy propylmethyl cellulose), guar gum, hydroxypropyl guar gum, soluble starch, cationic celluloses, cationic guar, and the like, and synthetic polymeric materials, such as carboxyvinyl polymers, polyvinylpyrrolidone, polyvinyl alcohol, polyacrylic acid polymers, polymethacrylic acid polymers, polyvinyl acetate polymers, polyvinyl chloride polymers, polyvinylidene chloride polymers and the like. Mixtures of the above compounds are contemplated.

[0046] Further exemplary gelling agents include the acrylic acid/ethyl acrylate copolymers and the carboxyvinyl polymers sold, for example, by the B.F. Goodrich Company under the trademark of Carbopol® resins. These resins consist essentially of a colloidal water-soluble polyalkenyl polyether crosslinked polymer of acrylic acid crosslinked with from 0.75% to 2% of a crosslinking agent such as polyallyl sucrose or polyallyl pentaerythritol. Examples include Carbopol® 934, Carbopol® 940, Carbopol® 950, Carbopol® 980, Carbopol® 951 and Carbopol® 981. Carbopol® 934 is a water-soluble polymer of acrylic acid crosslinked with about 1% of a polyallyl ether of sucrose having an average of about 5.8 allyl groups for each sucrose molecule.

[0047] In one or more embodiment, the composition of the present invention includes at least one polymeric agent, which is a water-soluble cellulose ether. Preferably, the water-soluble cellulose ether is selected from the group consisting of methylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose (Methocel), hydroxyethyl cellulose, methylhydroxyethylcellulose, methylhydroxypropylcellulose, hydroxyethylcarboxymethylcellulose, carboxymethylcellulose and carboxymethylhydroxyethylcellulose. More preferably, the water-soluble cellulose ether is selected from the group consisting of methylcellulose, hydroxypropyl cellulose and hydroxypropyl methylcellulose (Methocel). In one or more embodiments, the composition includes a combination of a water-soluble cellulose ether; and a naturally-occurring polymeric materials, selected from the group including xanthan gum, guar gum, carrageenan gum, locust bean gum and tragacanth gum.

[0048] Yet, in other embodiments, the gelling agent includes inorganic gelling agents, such as silicone dioxide (fumed silica).

[0049] Mucoadhesion/bioadhesion is defined as the attachment of synthetic or biological macromolecules to a biological tissue. Mucoadhesive agents are a class of polymeric biomaterials that exhibit the basic characteristic of a hydrogel, i.e. swell by absorbing water and interacting by means of adhesion with the mucous that covers epithelia. Compositions of the present invention may contain a mucoadhesive macromolecule or polymer in an amount sufficient to confer bioadhesive properties. The bioadhesive macromolecule

enhances the delivery of biologically active agents on or through the target surface. The mucoadhesive macromolecule may be selected from acidic synthetic polymers, preferably having at least one acidic group per four repeating or monomeric subunit moieties, such as poly(acrylic)- and/or poly(methacrylic) acid (e.g., Carbopol®, Carbomer®), poly(methylvinyl ether/maleic anhydride) copolymer, and their mixtures and copolymers; acidic synthetically modified natural polymers, such as carboxymethylcellulose (CMC); neutral synthetically modified natural polymers, such as (hydroxypropyl)methylcellulose; basic amine-bearing polymers such as chitosan; acidic polymers obtainable from natural sources, such as alginic acid, hyaluronic acid, pectin, gum tragacanth, and karaya gum; and neutral synthetic polymers, such as polyvinyl alcohol or their mixtures. An additional group of mucoadhesive polymers includes natural and chemically modified cyclodextrin, especially hydroxypropyl- β -cyclodextrin. Such polymers may be present as free acids, bases, or salts, usually in a final concentration of about 0.01% to about 0.5% by weight.

[0050] A suitable bioadhesive macromolecule is the family of acrylic acid polymers and copolymers, (e.g., Carbopol®). These polymers contain the general structure $-\text{[CH}_2-\text{CH}(\text{COOH})-\text{]}_n$. Hyaluronic acid and other biologically-derived polymers may be used.

[0051] Exemplary bioadhesive or mucoadhesive macromolecules have a molecular weight of at least 50 kDa, or at least 300 kDa, or at least 1,000 kDa. Favored polymeric ionizable macromolecules have not less than 2 mole percent acidic groups (e.g., COOH, SO₃H) or basic groups (NH₂, NRH, NR₂), relative to the number of monomeric units. The acidic or basic groups can constitute at least 5 mole percent, or at least 10 mole percent, or at least 25, at least 50 more percent, or even up to 100 mole percent relative to the number of monomeric units of the macromolecule.

[0052] Yet, another group of mucoadhesive agent includes inorganic gelling agents such as silicon dioxide (fumed silica), including but not limited to, AEROSIL 200 (DE-GUSSA).

[0053] Many mucoadhesive agents are known in the art to also possess gelling properties.

[0054] The foam composition may contain a film forming component. The film forming component may include at least one water-insoluble alkyl cellulose or hydroxyalkyl cellulose. Exemplary alkyl cellulose or hydroxyalkyl cellulose polymers include ethyl cellulose, propyl cellulose, butyl cellulose, cellulose acetate, hydroxypropyl cellulose, hydroxybutyl cellulose, and ethylhydroxyethyl cellulose, alone or in combination. In addition, a plasticizer or a cross linking agent may be used to modify the polymer's characteristics. For example, esters such as dibutyl or diethyl phthalate, amides such as diethyldiphenyl urea, vegetable oils, fatty acids and alcohols such as oleic and myristic acid may be used in combination with the cellulose derivative.

[0055] In one or more embodiments, the composition of the present invention includes a phase change polymer, which alters the composition behavior from fluid-like prior to administration to solid-like upon contact with the target mucosal surface. Such phase change results from external stimuli, such as changes in temperature or pH and exposure to specific ions (e.g., Ca²⁺).

[0056] Non-limiting examples of phase change polymers include poly(N-isopropylamide), Poloxamer 407® and Smart-Gel® (Poloxamer PAA)

[0057] The polymeric agent is present in an amount in the range of about 0.01% to about 5.0% by weight of the foam composition. In one or more embodiments, it is typically less than about 1 wt % of the foamable composition.

[0058] Surface-active agents (also termed "surfactants") include any agent linking oil and water in the composition, in the form of emulsion. A surfactant's hydrophilic/lipophilic balance (HLB) describes the emulsifier's affinity toward water or oil. The HLB scale ranges from 1 (totally lipophilic) to 20 (totally hydrophilic), with 10 representing an equal balance of both characteristics. Lipophilic emulsifiers form water-in-oil (w/o) emulsions; hydrophilic surfactants form oil-in-water (o/w) emulsions. The HLB of a blend of two emulsifiers equals the weight fraction of emulsifier A times its HLB value plus the weight fraction of emulsifier B times its HLB value (weighted average).

[0059] According to one or more embodiments of the present invention, the surface-active agent has a hydrophilic lipophilic balance (HLB) between about 9 and about 14, which is the required HLB (the HLB required to stabilize an O/W emulsion of a given oil) of most oils and hydrophobic solvents. Thus, in one or more embodiments, the composition contains a single surface active agent having an HLB value between about 9 and 14, and in one or more embodiments, the composition contains more than one surface active agent and the weighted average of their HLB values is between about 9 and about 14. Yet, in other embodiments, when a water in oil emulsion is desirable, the composition contains one or more surface active agents, having an HLB value between about 2 and about 9.

[0060] The surface-active agent is selected from anionic, cationic, nonionic, zwitterionic, amphoteric and ampholytic surfactants, as well as mixtures of these surfactants. Such surfactants are well known to those skilled in the therapeutic and cosmetic formulation art. Nonlimiting examples of possible surfactants include polysorbates, such as polyoxyethylene (20) sorbitan monostearate (Tween 60) and poly(oxyethylene) (20) sorbitan monooleate (Tween 80); poly(oxyethylene) (POE) fatty acid esters, such as Myrj 45, Myrj 49, Myrj 52 and Myrj 59; poly(oxyethylene)alkyl ethers, such as poly(oxyethylene)cetyl ether, poly(oxyethylene) palmityl ether, polyethylene oxide hexadecyl ether, polyethylene glycol cetyl ether, brij 38, brij 52, brij 56 and brij W1; sucrose esters, partial esters of sorbitol and its anhydrides, such as sorbitan monolaurate and sorbitan monolaurate; mono or diglycerides, isoceteth-20, sodium methyl cocoyl taurate, sodium methyl oleoyl taurate, sodium lauryl sulfate, triethanolamine lauryl sulfate and betaines.

[0061] In one or more embodiments of the present invention, the surface-active agent includes at least one non-ionic surfactant. Ionic surfactants are known to be irritants. Therefore, non-ionic surfactants are preferred in applications including sensitive tissue such as found in most mucosal tissues, especially when they are infected or inflamed. We have surprisingly found that non-ionic surfactants alone provide foams of excellent quality, i.e. a score of "E" according to the grading scale discussed herein below.

[0062] In one or more embodiments, the surface active agent includes a mixture of at least one non-ionic surfactant and at least one ionic surfactant in a ratio in the range of about 100:1 to 6:1. In one or more embodiments, the non-ionic to ionic surfactant ratio is greater than about 6:1, or greater than about 8:1; or greater than about 14:1, or greater than about 16:1, or greater than about 20:1.

[0063] In one or more embodiments of the present invention, a combination of a non-ionic surfactant and an ionic surfactant (such as sodium lauryl sulphate and cocamidopropylbetaine) is employed, at a ratio of between 1:1 and 20:1, or at a ratio of 4:1 to 10:1. The resultant foam has a low specific gravity, e.g., less than 0.1 g/ml.

[0064] It has been surprisingly discovered that the stability of the composition is especially pronounced when a combination of at least one non-ionic surfactant having HLB of less than 9 and at least one non-ionic surfactant having HLB of equal or more than 9 is employed. The ratio between the at least one non-ionic surfactant having HLB of less than 9 and the at least one non-ionic surfactant having HLB of equal or more than 9, is between 1:8 and 8:1, or at a ratio of 4:1 to 1:4. The resultant HLB of such a blend of at least two emulsifiers is between about 9 and about 14.

[0065] Thus, in an exemplary embodiment, a combination of at least one non-ionic surfactant having HLB of less than 9 and at least one non-ionic surfactant having HLB of equal or more than 9 is employed, at a ratio of between 1:8 and 8:1, or at a ratio of 4:1 to 1:4, wherein the HLB of the combination of emulsifiers is between about 9 and about 14.

[0066] In one or more embodiments of the present invention, the surface-active agent includes mono-, di- and triesters of sucrose with fatty acids (sucrose esters), prepared from sucrose and esters of fatty acids or by extraction from sucro-glycerides. Suitable sucrose esters include those having high monoester content, which have higher HLB values.

[0067] The total surface active agent is in the range of about 0.1 to about 5% of the foamable composition, and is typically less than about 2% or less than about 1%.

[0068] Preferably, a therapeutically effective foam adjuvant is included in the foamable compositions of the present invention to increase the foaming capacity of surfactants and/or to stabilize the foam. In one or more embodiments of the present invention, the foam adjuvant agent includes fatty alcohols having 15 or more carbons in their carbon chain, such as cetyl alcohol and stearyl alcohol (or mixtures thereof). Other examples of fatty alcohols are arachidyl alcohol (C20), behenyl alcohol (C22), 1-triacontanol (C30), as well as alcohols with longer carbon chains (up to C50). Fatty alcohols, derived from beeswax and including a mixture of alcohols, a majority of which has at least 20 carbon atoms in their carbon chain, are especially well suited as foam adjuvant agents. The amount of the fatty alcohol required to support the foam system is inversely related to the length of its carbon chains. Foam adjuvants, as defined herein are also useful in facilitating improved spreadability and absorption of the composition.

[0069] In one or more embodiments of the present invention, the foam adjuvant agent includes fatty acids having 16 or more carbons in their carbon chain, such as hexadecanoic acid (C16) stearic acid (C18), arachidic acid (C20), behenic acid (C22), octacosanoic acid (C28), as well as fatty acids with longer carbon chains (up to C50), or mixtures thereof. As for fatty alcohols, the amount of fatty acids required to support the foam system is inversely related to the length of its carbon chain.

[0070] Optionally, the carbon atom chain of the fatty alcohol or the fatty acid may have at least one double bond. A further class of foam adjuvant agent includes a branched fatty alcohol or fatty acid. The carbon chain of the fatty acid or fatty alcohol also can be substituted with a hydroxyl group, such as 12-hydroxy stearic acid.

[0071] An important property of the fatty alcohols and fatty acids used in context of the composition of the present invention is related to their therapeutic properties per se. Long chain saturated and mono unsaturated fatty alcohols, e.g., stearyl alcohol, erucyl alcohol, arachidyl alcohol and behenyl alcohol (docosanol) have been reported to possess antiviral, antiinfective, antiproliferative and antiinflammatory properties (see, for example, U.S. Pat. No. 4,874,794). Longer chain fatty alcohols, e.g., tetracosanol, hexacosanol, heptacosanol, octacosanol, triacontanol, etc., are also known for their metabolism modifying properties and tissue energizing properties. Long chain fatty acids have also been reported to possess anti-infective characteristics.

[0072] Thus, in preferred embodiments of the present invention, a combined and enhanced therapeutic effect is attained by including both a nonsteroidal immunomodulating agent and a therapeutically effective foam adjuvant in the same composition, thus providing a simultaneous anti-inflammatory and antiinfective effect from both components. Furthermore, in a further preferred embodiment, the composition concurrently comprises a nonsteroidal immunomodulating agent, a therapeutically effective foam adjuvant and a therapeutically active oil, as detailed above. Such combination provides an even more enhanced therapeutic benefit. Thus, the foamable carrier, containing the foam adjuvant provides an extra therapeutic benefit in comparison with currently used vehicles, which are inert and non-active.

[0073] The foam adjuvant according to preferred embodiments of the present invention includes a mixture of fatty alcohols, fatty acids and hydroxy fatty acids and derivatives thereof in any proportion, providing that the total amount is 0.1% to 5% (w/w) of the carrier mass. More preferably, the total amount is 0.4%-2.5% (w/w) of the carrier mass.

[0074] The therapeutic foam of the present invention may further optionally include a variety of formulation excipients, which are added in order to fine-tune the consistency of the formulation, protect the formulation components from degradation and oxidation and modify their consistency. Such excipients may be selected, for example, from stabilizing agents, antioxidants, humectants, preservatives, colorant and odorant agents and other formulation components, used in the art of formulation.

[0075] Aerosol propellants are used to generate and administer the foamable composition as a foam. The total composition including propellant, foamable compositions and optional ingredients is referred to as the foamable carrier. The propellant makes up about 3% to about 25 wt % of the foamable carrier. Examples of suitable propellants include volatile hydrocarbons such as butane, propane, isobutane or mixtures thereof, and fluorocarbon gases.

[0076] Several disorders of the skin, body cavity or mucosal surface (e.g., the mucosa of the nose, mouth, eye, ear, vagina or rectum), involve a combination of dry and scaly skin, which can be treated with lanolin as the hydrating component of the composition and additional therapeutic modalities that address other etiological factors of the disorder. For example, psoriasis involves dry and scaly skin, as well as inflammation, excessive cell proliferation and inadequate cell differentiation, which can be alleviated by treatment with lanolin and an additional active agent, such as a steroid, a retinoid or an alpha or beta hydroxy acid. Likewise, atopic dermatitis involves skin dryness and inflammation, which may be treated with lanolin and a steroid or an immunomodulating agent.

[0077] Thus, in one or more embodiments, the composition of the present invention is a carrier of a cosmetically or pharmaceutically active agents. Exemplary, non binding and cosmetically or pharmaceutically active agents include, but are not limited to an anti-infective, an antibiotic, an antibacterial agent, an antifungal agent, an antiviral agent, an antiparasitic agent, an steroidal antiinflammatory agent, an immunosuppressive agent, an immunomodulator, an immunoregulating agent, a hormonal agent, vitamin A, a vitamin A derivative, vitamin B, a vitamin B derivative, vitamin C, a vitamin C derivative, vitamin D, a vitamin D derivative, vitamin E, a vitamin E derivative, vitamin F, a vitamin F derivative, vitamin K, a vitamin K derivative, a wound healing agent, a disinfectant, an anesthetic, an antiallergic agent, an alpha hydroxyl acid, lactic acid, glycolic acid, a beta-hydroxy acid, a protein, a peptide, a neuropeptide, a allergen, an immunogenic substance, a haptene, an oxidizing agent, an antioxidant, a dicarboxylic acid, azelaic acid, sebacic acid, adipic acid, fumaric acid, a retinoid, an antiproliferative agent, an anticancer agent, a photodynamic therapy agent, an anti-wrinkle agent, a radical scavenger, a metal oxide (e.g., titanium dioxide, zinc oxide, zirconium oxide, iron oxide), silicone oxide, an anti wrinkle agent, a skin whitening agent, a skin protective agent, a masking agent, an anti-wart agent, a refatting agent, a lubricating agent and mixtures thereof.

Composition and Foam Physical Characteristics

[0078] A pharmaceutical or cosmetic composition manufactured using the foam carrier according to one or more embodiments of the present invention is very easy to use. When applied onto the afflicted body surface of mammals, i.e., humans or animals, it is in a foam state, allowing free application without spillage. Upon further application of a mechanical force, e.g., by rubbing the composition onto the body surface, it freely spreads on the surface and is rapidly absorbed.

[0079] The foam composition of the present invention creates a stable emulsion having an acceptable shelf-life of at least one year, or at least two years at ambient temperature. A feature of a product for cosmetic or medical use is long term stability. Propellants, which are a mixture of low molecular weight hydrocarbons, tend to impair the stability of emulsions. It has been observed, however, that foam compositions according to the present invention are surprisingly stable. Following accelerated stability studies, they demonstrate desirable texture; they form fine bubble structures that do not break immediately upon contact with a surface, spread easily on the treated area and absorb quickly.

[0080] The composition should also be free flowing, to allow it to flow through the aperture of the container, e.g., and aerosol container, and create an acceptable foam. Compositions containing semi-solid hydrophobic solvents, e.g., white petrolatum, as the main ingredients of the oil phase of the emulsion, exhibit high viscosity and poor flowability and are inappropriate candidates for a foamable composition.

[0081] Foam quality can be graded as follows:

[0082] Grade E (excellent): very rich and creamy in appearance, does not show any bubble structure or shows a very fine (small) bubble structure; does not rapidly become dull; upon spreading on the skin, the foam retains the creaminess property and does not appear watery.

[0083] Grade G (good): rich and creamy in appearance, very small bubble size, "dulls" more rapidly than an excellent foam, retains creaminess upon spreading on the skin, and does not become watery.

[0084] Grade FG (fairly good): a moderate amount of creaminess noticeable, bubble structure is noticeable; upon spreading on the skin the product dulls rapidly and becomes somewhat lower in apparent viscosity.

[0085] Grade F (fair): very little creaminess noticeable, larger bubble structure than a "fairly good" foam, upon spreading on the skin it becomes thin in appearance and watery.

[0086] Grade P (poor): no creaminess noticeable, large bubble structure, and when spread on the skin it becomes very thin and watery in appearance.

[0087] Grade VP (very poor): dry foam, large very dull bubbles, difficult to spread on the skin.

[0088] Typically administratable foams are typically of quality grade E or G, when released from the aerosol container. Smaller bubbles are indicative of more stable foam, which does not collapse spontaneously immediately upon discharge from the container. The finer foam structure looks and feels smoother, thus increasing its usability and appeal.

[0089] As further aspect of the foam is breakability. The breakable foam is thermally stable, yet breaks under sheer force. Sheer-force breakability of the foam is clearly advantageous over thermally-induced breakability. Thermally sensitive foams immediately collapse upon exposure to skin temperature and, therefore, cannot be applied on the hand and afterwards delivered to the afflicted area.

[0090] Another property of the foam is specific gravity, as measured upon release from the aerosol can. Typically, foams have specific gravity of less than 0.1 g/mL or less than 0.05 g/mL.

Fields of Pharmaceutical Applications

[0091] By including lanolin and optional active agents in the compositions of the present invention, the composition are useful in treating an animal or a human patient having any one of a variety of dermatological disorders that include dry and/or scaly skin as one or their etiological factors (also termed "dermatoses"), such as classified in a non-limiting exemplary manner according to the following groups:

[0092] Dermatitis including contact dermatitis, atopic dermatitis, seborrheic dermatitis, nummular dermatitis, chronic dermatitis of the hands and feet, generalized exfoliative dermatitis, stasis dermatitis; lichen simplex chronicus; diaper rash;

[0093] Bacterial infections including cellulitis, acute lymphangitis, lymphadenitis, erysipelas, cutaneous abscesses, necrotizing subcutaneous infections, staphylococcal scalded skin syndrome, folliculitis, furuncles, hidradenitis suppurativa, carbuncles, paronychia infections, erythrasma;

[0094] Fungal Infections including dermatophyte infections, yeast Infections; parasitic Infections including scabies, pediculosis, creeping eruption;

[0095] Viral Infections;

[0096] Disorders of hair follicles and sebaceous glands including acne, rosacea, perioral dermatitis, hypertrichosis (hirsutism), alopecia, including male pattern baldness, alopecia areata, alopecia universalis and alopecia totalis; pseudo-folliculitis barbae, keratinous cyst;

[0097] Scaling papular diseases including psoriasis, pityriasis rosea, lichen planus, pityriasis rubra pilaris;

[0098] Benign tumors including moles, dysplastic nevi, skin tags, lipomas, angiomas, pyogenic granuloma, seborrheic keratoses, dermatofibroma, keratoacanthoma, keloid; [0099] Malignant tumors including basal cell carcinoma, squamous cell carcinoma, malignant melanoma, paget's disease of the nipples, kaposi's sarcoma;

[0100] Reactions to sunlight including sunburn, chronic effects of sunlight, photosensitivity;

[0101] Bullous diseases including pemphigus, bullous pemphigoid, dermatitis herpetiformis, linear immunoglobulin A disease;

[0102] Pigmentation disorders including hypopigmentation such as vitiligo, albinism and postinflammatory hypopigmentation and hyperpigmentation such as melasma (chloasma), drug-induced hyperpigmentation, postinflammatory hyperpigmentation;

[0103] Disorders of cornification including ichthyosis, keratosis pilaris, calluses and corns, actinic keratosis;

[0104] Pressure sores;

[0105] Disorders of sweating; and

[0106] Inflammatory reactions including drug eruptions, toxic epidermal necrolysis; erythema multiforme, erythema nodosum, granuloma annulare.

[0107] According to one or more embodiments of the present invention, the compositions are also useful in the therapy of non-dermatological disorders by providing transdermal delivery of an active nonsteroidal immunomodulating agent that is effective against non-dermatological disorders.

[0108] The same advantage is expected when the composition is topically applied to a body cavity or mucosal surface (e.g., the mucosa of the nose, mouth, eye, ear, vagina or rectum) to treat conditions such as chlamydia infection, gonorrhea infection, hepatitis B, herpes, HIV/AIDS, human papillomavirus (HPV), genital warts, bacterial vaginosis, candidiasis, chancroid, granuloma Inguinale, lymphogranuloma venereum, mucopurulent cervicitis (MPC), molluscum contagiosum, nongonococcal urethritis (NGU), trichomoniasis, vulvar disorders, vulvodynia, vulvar pain, yeast infection, vulvar dystrophy, vulvar intraepithelial neoplasia (VIN), contact dermatitis, pelvic inflammation, endometritis, salpingitis, oophoritis, genital cancer, cancer of the cervix, cancer of the vulva, cancer of the vagina, vaginal dryness, dyspareunia, anal and rectal disease, anal abscess/fistula, anal cancer, anal fissure, anal warts, Crohn's disease, hemorrhoids, anal itch, pruritus ani, fecal incontinence, constipation, polyps of the colon and rectum.

[0109] The following examples exemplify the therapeutic kits and pharmacological compositions and methods described herein. The examples are for the purposes of illustration only and are not intended to be limiting of the invention.

Example 1

Lanolin Based Foamable Compositions

[0110]

Ingredient	Composition No:			
	LAN1	LAN2	LAN3	LAN4
	%			
Lanolin	5.00	15.00	30.00	40.00
Mineral oil light	20.00	—	—	—

-continued

Ingredient	Composition No:			
	LAN1	LAN2	LAN3	LAN4
	%			
MCT oil	—	10.00	—	—
Glyceryl monostearate	0.45	0.45	0.45	1.00
Stearyl alcohol	0.95	—	—	—
Xanthan gum	0.28	—	—	—
Methocel K100M	0.28	—	—	—
Polysorbate 80	0.95	0.95	0.95	0.95
PEG-40 Stearate	2.80	2.80	2.80	2.80
Preservative	0.25	0.25	0.25	0.25
Propellant	8.00	8.00	8.00	8.00
Water	To 100	To 100	To 100	To 100
Composition Properties				
Emulsion color	White	White	White	White
Foam Density	0.06	0.06	0.08	0.08

Example 2

Additional Lanolin Based Foamable Compositions

[0111]

Ingredient	Composition No:			
	LAN5	LAN6	LAN7	LAN8
	%			
Lanolin	5.00	5.00	5.00	5.00
Mineral oil	20.00	20.00	20.00	—
IPP	10.00	—	—	—
IPM	—	10.00	10.00	—
Beeswax	2.00	2.00	2.00	2.00
Glyceryl oleate	0.80	0.80	0.80	0.80
Glyceryl monostearate SE	0.80	0.80	0.80	0.80
Arlacel P135	0.40	0.40	0.40	0.40
Ceteth-2 (Lipocol C2)	6.00	5.00	5.00	5.00
Arlatone 2121	0.50	0.50	0.50	0.50
Sucrose ester SP30	0.80	0.80	0.80	0.80
Sucrose ester SP10	0.80	0.80	0.80	0.80
<i>Aloe vera</i> extract	0.10	0.10	0.10	0.10
Xanthan gum	0.40	0.40	0.40	0.40
Magnesium sulfate	—	0.20	0.20	0.20
D-Pantenol	4.00	4.00	4.00	4.00
Disodium EDTA	0.20	0.40	0.40	0.40
Alpha Bisabolol	—	0.20	0.20	0.20
Benzalkonium chloride (50% sol)	0.20	0.20	0.20	0.20
Zinc oxide	10.00	10.00	—	—
Water	To 100	To 100	To 100	To 100
Composition Properties				
Emulsion color	White	White	White	White
Foam Density	0.072	0.086	0.088	0.070

Example 6

Controlled Study, to Assess the Skin Hydration of Two Lanolin Based Foam Compositions

[0112] Six subjects were treated with foam compositions LAN1 and LAN2, on their arm. Skin hydration was measured

using Corneometer CM825 for 240 minutes (6 hours) at the treatment sites and at an untreated site. The skin hydration values for the two lanolin based foams vs. “no treatment” is presented in Table 1 below and is shown in FIG. 1. Table 2 provides the “delta” values, i.e., the difference between the hydration effect of LAN1 and LAN2, and the hydration value of the non-treated sites, which is illustrated in FIG. 2. As demonstrated in Tables 1 and 2, LAN1 and LAN2 foams provided significant hydration for at least 6 hours. The improvement rendered by LAN1 was in the range of 16.7%-22.4%, and for LAN2, there was a 24.9%-19.6% improvement. For comparison purposes, in a parallel group, which was treated with a commercial 12% ammonium lactate lotion, the skin moisture was improved by 20%-28%. Notable, while 12% ammonium lactate is a prescription drug with known side effects, the lanolin foam compositions can be generally considered as safe and can be used daily without safety concerns.

TABLE 1

Moisturizing Test: Two Lanolin Based Foams vs. No Treatment, using Corneometer CM825			
Time (min)	Corneometer CM825 Values		
	No Treatment	LAN1	LAN2
0	41	40	38
60	41	48	47
120	40	47	49
180	40	47	49
240	42	49	49

TABLE 2

Moisturizing Test: The Absolute (Incremental) Skin Hydrating Effect of Two Lanolin, Using Corneometer CM825		
Time (min)	Corneometer CM825 Values	
	LAN1	LAN2
60	7.75 (+19.4%)	9.46 (+24.9%)
120	6.71 (+16.7%)	11.04 (+29.1%)
180	6.75 (+16.9%)	11.25 (+29.6%)
240	8.96 (+22.4%)	11.21 (+29.5%)

What is claimed is:

1.-20. (canceled)

21. A hydrating foamable composition comprising an emulsion, the emulsion comprising:

- i) a hydrating component comprising an emollient, a humectant, or a mixture thereof;
- ii) about 0.1% to about 5% of a surface-active agent by weight of the emulsion;
- iii) about 0.01% to about 5% by weight of the emulsion of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent, a phase change agent, a water-soluble cellulose ether, a naturally-occurring polymeric material, and mixtures of two or more thereof;
- iv) water; and
- v) optionally at least one therapeutic agent; and
 - a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the hydrating foamable composition;

wherein upon release from a pressurized container the composition expands to form a foam that collapses upon application of shear force but does not immediately collapse upon exposure to skin temperature.

22. The foamable composition of claim **21**, further comprising at least one organic carrier selected from the group consisting of a hydrophobic organic carrier, a polar solvent, and mixtures of two or more thereof, wherein the at least one organic carrier is present in an amount selected from the group consisting of (i) about 2% to about 5% by weight of the emulsion; (ii) about 5% to about 10% by weight of the emulsion; (iii) about 10% to about 20% by weight of the emulsion; and (iv) about 20% to about 50% by weight of the emulsion.

23. The foamable composition of claim **21**, further comprising about 0.1% to about 5% by weight of the emulsion of a foam adjuvant selected from the group consisting of a fatty alcohol having 15 or more carbons in its carbon chain; a fatty acid having 16 or more carbons in its carbon chain; fatty alcohols derived from beeswax and including a mixture of alcohols, a majority of which has at least 20 carbon atoms in their carbon chain; a fatty alcohol having at least one double bond; a fatty acid having at least one double bond; a branched fatty alcohol; a branched fatty acid; and a fatty acid substituted with a hydroxyl group; and mixtures of any two or more thereof.

24. The foamable composition of claim **21**, wherein the foamable composition is substantially free of short chain alcohols.

25. The foamable composition of claim **21**, wherein the hydrating component comprises a lanolin selected from the group consisting of:

- i) lanolin, wool fat, lanolin wax, and a synthetic lanolin;
- ii) a synthetic lanolin comprising at least one of cholesterol, ischolesterol, and analogs, derivatives, salts, and esters thereof;
- iii) an oil-soluble hydrophobic derivative of lanolin;
- iv) a synthetic lanolin comprising at least one of cholesterol and ischolesterol covalently linked with an organic moiety;
- v) a synthetic lanolin comprising at least one of cholesterol and ischolesterol covalently linked with an aliphatic acid;
- vi) a lanolin acid;
- vii) an ester of lanolin acid with an aliphatic alcohol;
- viii) a compound selected from the group consisting of acetylated lanolin, isopropyl lanolate, and cetyl lanolate; and
- ix) mixtures of any two or more thereof.

26. The foamable composition of claim **25**, wherein the concentration range of lanolin is selected from the group of (i) between about 5% and about 12% by weight of the emulsion; and (ii) between about 12% and about 24% by weight of the emulsion.

27. The foamable composition of claim **21**, wherein the surface active agent is non ionic and the concentration of the non-ionic surface active agent is less than about 2% by weight of the emulsion.

28. The foamable composition of claim **21**, wherein the surface-active agent comprises a combination of at least one non-ionic surfactant having HLB of less than 9 and at least one non-ionic surfactant having HLB of equal or more than 9, wherein the ratio between the at least one non-ionic surfactant

having HLB of less than 9 and the at least one non-ionic surfactant having HLB of equal or more than 9, is between 1:8 and 8:1.

29. The foamable composition of claim **21**, wherein the polymeric agent is selected from the group consisting of methylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hydroxyethyl cellulose, methylhydroxyethylcellulose, methylhydroxypropylcellulose, hydroxyethylcarboxymethylcellulose, carboxymethylcellulose, carboxymethylhydroxyethylcellulose, xanthan gum, guar gum, carrageenan gum, locust bean gum, tragacanth gum, and mixtures of any two or more thereof.

30. The foamable composition of claim **21**, comprising a therapeutic agent selected from the group consisting of an anti-infective, an antibiotic, an antibacterial agent, an antifungal agent, an antiviral agent, an antiparasitic agent, a steroid antiinflammatory agent, an immunosuppressive agent, an immunomodulator, an immunoregulating agent, a hormonal agent, a vitamin A, a vitamin A derivative, a vitamin B, a vitamin B derivative, a vitamin C, a vitamin C derivative, a vitamin D, a vitamin D derivative, a vitamin E, a vitamin E derivative, a vitamin F, a vitamin F derivative, a vitamin K, a vitamin K derivative, a wound healing agent, a disinfectant, an anesthetic, an antiallergic agent, an alpha hydroxyl acid, lactic acid, glycolic acid, a beta-hydroxy acid, a protein, a peptide, a neuropeptide, an allergen, an immunogenic substance, a haptene, an oxidizing agent, an antioxidant, a dicarboxylic acid, azelaic acid, sebacic acid, adipic acid, fumaric acid, a retinoid, an antiproliferative agent, an anticancer agent, a photodynamic therapy agent, benzoyl chloride, calcium hypochlorite, magnesium hypochlorite, an anti-wrinkle agent, a radical scavenger, a metal, silver, a metal oxide, titanium dioxide, zinc oxide, zirconium oxide, iron oxide, silicone oxide, talc, carbon, a skin whitening agent, a skin protective agent, a masking agent, an anti-wart agent, a refatting agent, a lubricating agent, and mixtures of any two or more thereof.

31. The composition of claim **21**, wherein the hydrating component is selected from the group consisting of a lanolin, a beeswax, a mineral oil, a MCT oil, isopropyl myristate (IPM), and isopropyl palmitate (IPP).

32. The composition of claim **25**, wherein the hydrating component is at least about 5% by weight of the emulsion and comprises a combination of a lanolin and a component selected from the group consisting of a beeswax, a mineral oil, a MCT oil, isopropyl myristate (IPM) isopropyl palmitate (IPP), and mixtures of any two or more thereof.

33. A hydrating foamable composition comprising an emulsion, the emulsion comprising:

- i) a hydrating component comprising a humectant;
- ii) about 0.1% to about 5% of a surface-active agent by weight of the emulsion;
- iii) about 0.01% to about 5% by weight of the emulsion of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent, a phase change agent, a water-soluble cellulose ether, a naturally-occurring polymeric material, and mixtures of two or more thereof;
- iv) water; and
- v) optionally at least one therapeutic agent; and
 - a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the hydrating foamable composition;

wherein upon release from a pressurized container the composition expands to form a foam that collapses upon application of shear force but does not immediately collapse upon exposure to skin temperature.

34. The foamable composition of claim **33**, further comprising at least one organic carrier selected from the group consisting of a hydrophobic organic carrier, a polar solvent, an emollient, and mixtures of two or more thereof, wherein the at least one organic carrier is present in an amount selected from the group consisting of (i) about 2% to about 5% by weight of the emulsion; (ii) about 5% to about 10% by weight of the emulsion; (iii) about 10% to about 20% by weight of the emulsion; and (iv) about 20% to about 50% by weight of the emulsion.

35. A method of treating or alleviating insufficient hydration of skin, a body cavity or a mucosal surface, comprising: administering topically to skin, a body cavity or a mucosal surface a foam that collapses upon application of shear force but does not immediately collapse upon exposure to skin temperature, wherein the foam is obtained by releasing from a pressurized container a foamable composition comprising an emulsion comprising:

- i) a hydrating component comprising an emollient, a humectant, or mixture thereof;
- ii) about 0.1% to about 5% of a non-ionic surface-active agent by weight of the emulsion;
- iii) about 0.01% to about 5% by weight of the emulsion of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent, a phase change agent, a water-soluble cellulose ether, a naturally-occurring polymeric material, and mixtures of two or more thereof; and
- iv) water; and

a liquefied or compressed gas propellant at a concentration of about 3% to about 25% by weight of the hydrating foamable composition;

wherein the non-ionic surface-active agent comprises a combination of at least one non-ionic surfactant having HLB of less than 9 and at least one non-ionic surfactant having HLB of equal or more than 9, wherein the ratio between the at least one non-ionic surfactant having HLB of less than 9 and the at least one non-ionic surfactant having HLB of equal or more than 9, is between 1:8 and 8:1.

36. The method of claim **35**, wherein the hydrating component comprises a lanolin selected from the group consisting of:

- i) lanolin, wool fat, lanolin wax, and a synthetic lanolin;
- ii) a synthetic lanolin comprising at least one of cholesterol, ischolesterol, and analogs, derivatives, salts, and esters thereof;
- iii) an oil-soluble hydrophobic derivative of lanolin;
- iv) a synthetic lanolin comprising at least one of cholesterol and ischolesterol covalently linked with an organic moiety;
- v) a synthetic lanolin comprising at least one of cholesterol and ischolesterol covalently linked with an aliphatic acid;
- vi) a lanolin acid;
- vii) an ester of lanolin acid with an aliphatic alcohol;
- viii) a compound selected from the group consisting of acetylated lanolin, isopropyl lanolate, and cetyl lanolate; and
- ix) mixtures of any two or more thereof.

37. The method of claim **35**, wherein the composition further comprises at least about 5% by weight of the emulsion of a lanolin and about 0.1% to about 5% by weight of the emulsion of a foam adjuvant selected from the group consisting of a fatty alcohol having 15 or more carbons in its carbon chain; a fatty acid having 16 or more carbons in its carbon chain; fatty alcohols, derived from beeswax and including a mixture of alcohols, a majority of which has at least 20 carbon atoms in their carbon chain; a fatty alcohol having at least one double bond; a fatty acid having at least one double bond; a branched fatty alcohol; a branched fatty acid; a fatty acid substituted with a hydroxyl group; and mixtures of any two or more thereof.

38. The method of claim **35**, wherein the composition further comprises an additional active agent effective to treat a disorder of skin, a body cavity or a mucosal surface.

39. The method of claim **35**, wherein the foamable composition further comprises an organic carrier consisting of at least one therapeutically active oil.

40. The method of claim **38**, wherein the additional active agent is selected from the group consisting of an anti-infective, an antibiotic, an antibacterial agent, an antifungal agent, an antiviral agent, an antiparasitic agent, a steroidal anti-inflammatory agent, an immunosuppressive agent, an immunomodulator, an immunoregulating agent, a hormonal agent, a vitamin A, a vitamin A derivative, a vitamin B, a vitamin B derivative, a vitamin C, a vitamin C derivative, a vitamin D, a

vitamin D derivative, a vitamin E, a vitamin E derivative, a vitamin F, a vitamin F derivative, a vitamin K, a vitamin K derivative, a wound healing agent, a disinfectant, an anesthetic, an antiallergic agent, an alpha hydroxyl acid, lactic acid, glycolic acid, a beta-hydroxy acid, a protein, a peptide, a neuropeptide, an allergen, an immunogenic substance, a hapten, an oxidizing agent, an antioxidant, a dicarboxylic acid, azelaic acid, sebacic acid, adipic acid, fumaric acid, a retinoid, an antiproliferative agent, an anticancer agent, a photodynamic therapy agent, benzoyl chloride, calcium hypochlorite, magnesium hypochlorite, an anti-wrinkle agent, a radical scavenger, a metal, silver, a metal oxide, titanium dioxide, zinc oxide, zirconium oxide, iron oxide, silicone oxide, talc, carbon, a skin whitening agent, a skin protective agent, a masking agent, an anti-wart agent, a refatting agent, a lubricating agent, and mixtures of any two or more thereof.

41. The method of claim **35**, wherein the surface active agent is non ionic and the concentration of the nonionic surface active agent is less than about 2% by weight of the emulsion.

42. The method of claim **35**, wherein the hydrating component is selected from the group consisting of a lanolin, beeswax, mineral oil, MCT oil, isopropyl myristate (IPM) isopropyl palmitate (IPP), and mixtures of any two or more thereof.

* * * * *