

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau

(43) International Publication Date
25 April 2013 (25.04.2013)(10) International Publication Number
WO 2013/057320 A1

(51) International Patent Classification:

$C07D\ 205/04$ (2006.01) $C07D\ 221/20$ (2006.01)
 $C07D\ 207/09$ (2006.01) $C07D\ 223/12$ (2006.01)
 $C07D\ 207/14$ (2006.01) $C07D\ 309/04$ (2006.01)
 $C07D\ 209/96$ (2006.01) $C07D\ 309/14$ (2006.01)
 $C07D\ 211/26$ (2006.01) $C07D\ 401/04$ (2006.01)
 $C07D\ 211/56$ (2006.01) $C07D\ 401/12$ (2006.01)
 $C07D\ 211/58$ (2006.01) $C07D\ 405/12$ (2006.01)
 $C07D\ 215/42$ (2006.01)

Genomics, S.A., C/Sant Ferran, 74, E-08940 Cornellà de Llobregat (ES). **CASTRO PALOMINO LARIA, Julio Cesar**; Oryzon Genomics, S.A., C/Sant Ferran, 74, E-08940 Cornellà de Llobregat (ES).

(74) Agent: VOSSIUS & PARTNER; Siebertstraße 4, 81675 München (DE).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: (HETERO)ARYL CYCLOPROPYLAMINE COMPOUNDS AS LSD1 INHIBITORS



(57) Abstract: The invention relates to (hetero)aryl cyclopropylamine compounds, including particularly the compounds of formula I as described and defined herein, and their use in therapy, including, e.g., in the treatment or prevention of cancer, a neurological disease or condition, or a viral infection.

5

(HETERO)ARYL CYCLOPROPYLAMINE COMPOUNDS AS LSD1 INHIBITORS

The invention relates to (hetero)aryl cyclopropylamine compounds, particularly compounds of formula I, Ia, Ia-1, Ib and Ic, more particularly compounds of formula I and Ia, as described and defined herein, and their use in 10 therapy, including e.g., in the treatment or prevention of cancer, a neurological disease, or a viral infection.

Aberrant gene expression in affected tissue as compared to normal tissue is a common characteristic of many 15 human diseases. This is true for cancer and many neurological diseases which are characterized by changes in gene expression patterns. Gene expression patterns are controlled at multiple levels in the cell. Control of gene expression can occur through modifications of DNA: DNA promoter methylation is associated with suppression of gene expression. Several inhibitors of DNA methylation are approved for clinical use including the blockbuster Vidaza™. Another class of modifications involve histones which form the protein scaffold that DNA is normally associated with (coiled around) in eukaryotic cells. Histones play a crucial role in organizing 20 DNA and the regulated coiling and uncoiling of DNA around the histones is critical in controlling gene expression – coiled DNA is typically not accessible for gene transcription. A number of histone modifications have been discovered including histone acetylation, histone lysine methylation, histone arginine methylation, 25 histone ubiquinylation, and histone sumoylation, many of which modify accessibility to the associated DNA by the cells transcriptional machinery. These histone marks serve to recruit various protein complexes involved in transcription and repression. An increasing number of studies are painting an intricate picture of how various combinations of histone marks control gene expression in cell-type specific manner and a new term has been coined to capture this concept: the histone code.

The prototypical histone mark is histone acetylation. Histone acetyl transferase and histone deacetylases are 30 the catalytic machines involved in modulation of this histone mark although typically these enzymes are parts of multiprotein complexes containing other proteins involved in reading and modifying histone marks. The components of these protein complexes are typically cell-type specific and typically comprise transcriptional regulators, repressors, co-repressors, receptors associated with gene expression modulation (e.g., estrogen or androgen receptor). Histone deacetylase inhibitors alter the histone acetylation profile of chromatin. Accordingly, histone deacetylase inhibitors like Vorinostat (SAHA), Trichostatin A (TSA), and many others have 35 been shown to alter gene expression in various *in vitro* and *in vivo* animal models. Clinically, histone deacetylase inhibitors have demonstrated activity in the cancer setting and are being investigated for oncology indications as well as for neurological conditions and other diseases.

Another modification that is involved in regulating gene expression is histone methylation including lysine and arginine methylation. The methylation status of histone lysines has recently been shown to be important in dynamically regulating gene expression.

5 A group of enzymes known as histone lysine methyl transferases and histone lysine demethylases are involved in histone lysine modifications. One particular human histone lysine demethylase enzyme called Lysine Specific Demethylase-1 (LSD1) was recently discovered (Shi *et al.* (2004) *Cell* 119:941) to be involved in this crucial histone modification. LSD1 has a fair degree of structural similarity, and amino acid identity/homology to polyamine oxidases and monoamine oxidases, all of which (*i.e.*, MAO-A, MAO-B and LSD1) are flavin dependent amine oxidases which catalyze the oxidation of nitrogen-hydrogen bonds and/or nitrogen carbon bonds. LSD1 has been recognized as an interesting target for the development of new drugs to treat cancer, 10 neurological diseases and other conditions.

15 Cyclopropylamine containing compounds are known to inhibit a number of medically important targets including amine oxidases like Monoamine Oxidase A (MAO-A; or MAOA), Monoamine Oxidase B (MAO-B; or MAOB), and Lysine Specific Demethylase-1 (LSD1). Tranylcypromine (also known as 2-phenylcyclopropylamine), which is the active ingredient of Parnate® and one of the best known examples of a cyclopropylamine, is known to inhibit all of these enzymes. Since MAO-A inhibition may cause undesired side effects, it would be desirable to identify cyclopropylamine derivatives that exhibit potent LSD1 inhibitory activity while being devoid of or having 20 substantially reduced MAO-A inhibitory activity.

In view of the lack of adequate treatments for conditions such as cancer and neurodegeneration, there is a desperate need for disease modifying drugs and drugs that work by inhibiting novel targets. There is thus a need for the development of LSD1 inhibitors, particularly those which selectively inhibit LSD1.

25


SUMMARY OF THE INVENTION

The present invention relates to the identification of compounds and their use in treating or preventing diseases. The invention provides (hetero)aryl cyclopropylamine compounds, including the compounds of 30 Formula I, Ia, Ia-1, Ib and Ic, and particularly the compounds I, Ia and Ia-1, as described and defined herein. The present invention particularly provides a compound of Formula I, Ia, Ia-1, Ib and Ic, and particularly a compound of Formula I, Ia and Ia-1, pharmaceutical compositions comprising a compound of Formula I, Ia, Ia-1, Ib or Ic, and particularly a compound of Formula I, Ia or Ia-1, and a pharmaceutically acceptable carrier, and their uses for treating diseases. One use of the compound of Formula I, Ia, Ia-1, Ib and Ic is for treating or 35 preventing cancer. Another use for the compound of Formula I, Ia, Ia-1, Ib and Ic is to inhibit LSD1. The invention thus relates to a compound of Formula I, Ia, Ia-1, Ib or Ic, and particularly a compound of Formula I,

la or la-1, for use in treating or preventing human disease. The present invention further relates to a compound of Formula I, la, la-1, Ib and Ic, and particularly a compound of Formula I, la or la-1, for use in treating or preventing cancer. The present invention further relates to a compound of Formula I, la, la-1, Ib and Ic, and particularly a compound of Formula I, la or la-1, for use in treating or preventing a neurological disease. The present invention further relates to a compound of Formula I, la, la-1, Ib and Ic, and particularly a compound of Formula I, la or la-1, for use in treating or preventing a viral infection.

Accordingly, the present invention provides a compound of Formula I and relates, in particular, to the compound of Formula I for use as a medicament:

10



wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

15

B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

20

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- 25 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

30 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;
or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- 5 (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

10 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;


15 each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

20 each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

25 each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl.

30 In another embodiment, the present invention provides a compound of Formula I wherein R^w, R^x, R^y and R^z are each hydrogen, i.e. a compound of formula Ia, and relates, in particular, to the compound of Formula Ia for use as a medicament:



35 wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

B is H, R¹ or -L¹-E;

5 E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

10 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

15 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula Ia through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

20 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

25 (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula Ia through a ring C atom,

30 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and


wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;


5 each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

10 each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea.

15 In another embodiment, the invention provides a compound of formula Ia as defined above wherein the substituents of the cyclopropyl moiety -A-B and -NH-L²-D are in the *trans*- configuration, i.e. a compound of formula Ia-1:



wherein the groups and variables of formula Ia-1, including A, B, D, E, L¹, L², R¹, R² and R³, are as defined above in relation to a compound of formula Ia. The above chemical representation for a compound of formula Ia-1 does not intend to indicate absolute stereochemistry of the two chiral centers on the cyclopropyl ring, but only their relative stereochemistry (which is *trans*). Thus a compound of formula Ia-1 could likewise be represented as



25 A compound of formula Ia-1 therefore relates to the individual optically active *trans* isomers as well as any mixtures thereof.

In another embodiment, the invention provides a compound of formula I as defined above wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl with the proviso that at least one of R^w, R^x, R^y and R^z is not hydrogen; that compound is referred to as a compound of formula Ib in the following. In a more preferred embodiment, R^w is selected from halo and C₁₋₄ alkyl, preferably from fluoro and methyl, and each R^x, R^y and R^z is hydrogen. A compound of formula I wherein R^w is selected from halo and C₁₋₄ alkyl,

preferably from fluoro and methyl, and each R^x , R^y and R^z is hydrogen is referred to as a compound of formula Ic in the following.

Also included within the scope of the invention are all isomers, including all stereoisomers and mixtures thereof, of the compounds of formula I, Ia, Ia-1, Ib and Ic. All salts and all solvates, preferably pharmaceutically acceptable salts and solvates, of the compounds of formula I, Ia, Ia-1, Ib and Ic are also encompassed within the scope of the invention. Furthermore, all physical forms (including amorphous and crystalline forms) of any such compounds are also encompassed within the scope of the invention. Any reference to a compound of formula I, Ia, Ia-1, Ib or Ic should be construed, unless otherwise indicated, as a reference to a compound of formula I, Ia, Ia-1, Ib or Ic (respectively), any isomer thereof (including any stereoisomer thereof or any mixtures thereof), any salt thereof (including any pharmaceutically acceptable salt thereof), any solvate thereof (including any pharmaceutically acceptable solvate thereof), and any physical form thereof.

The compounds of formula Ia, including also the compounds of formula Ia-1, are particularly preferred compounds according to the present invention. The most preferred compounds of the invention are the compounds of formula Ia-1.

Any chemical drawing or formula given herein is intended to represent unlabeled forms as well as isotopically labeled forms of the compounds of the invention. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine, chlorine, and iodine, such as 2H , 3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , and ^{125}I , respectively. Such isotopically labelled compounds are useful in metabolic studies (preferably with ^{14}C), reaction kinetic studies (with, for example 2H or 3H), detection or imaging techniques [such as positron emission tomography (PET) or single- photon emission computed tomography (SPECT)] including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F or ^{11}C labeled compound may be particularly preferred for PET or SPECT studies. Further, substitution with heavier isotopes such as deuterium (i.e., 2H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements. Particularly preferred are the deuterated forms of the compounds of the invention, i.e. a compound of formula I, Ia, Ia-1, Ib and Ic above wherein one or more hydrogen atoms has been replaced with deuterium. Isotopically labeled compounds of the invention can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non- isotopically labeled reagent. In addition to the unlabeled form, all isotopically labeled forms of the compounds of formula I, Ia, Ia-1, Ib and Ic are included within the scope of the invention.

In a compound of formula I, Ia, Ib or Ic the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are preferably in the *trans*-configuration.

5 The compounds of formula I, Ia, Ia-1, Ib and Ic are potent inhibitors of LSD1 and therefore can be useful for the treatment or prevention of any disease associated with LSD1.

10 The invention thus provides a pharmaceutical composition comprising a compound of Formula I, Ia, Ia-1, Ib or Ic and a pharmaceutically acceptable carrier. Preferred embodiments of the compound of Formula I, Ia, Ia-1, Ib and Ic, e. g. for use in the compositions of the invention, are defined and described herein below in more detail.

15 In another aspect, the invention provides a method of treating or preventing a disease comprising administering, to a patient (preferably a human) in need of such treatment or prevention, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic (as described above or as defined in the embodiments thereof described below) effective to treat or prevent said disease. In one embodiment, such disease is a disease associated with LSD1.

20 In a related aspect, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (as described above or as defined in the embodiments thereof as described below) for use as a medicament. In a more specific embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic for use in the treatment or prevention of a disease associated with LSD1.

25 In yet another aspect, the invention provides a method of inhibiting LSD1 activity comprising administering, to a patient in need of treatment, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic sufficient to inhibit LSD1 activity. Preferably the patient is a human. In a related aspect, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic as herein defined for use as a LSD1 inhibitor. Preferred embodiments of the compounds of Formula I, Ia, Ia-1, Ib or Ic for use herein are as described in more detail below.

30 In another aspect, the invention provides a method of treating or preventing cancer comprising administering, to a patient (preferably a human) in need of such treatment or prevention, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or as defined in the embodiments described in more detail herein) sufficient to treat or prevent such cancer. In a related aspect, the invention provides a method of treating or preventing a cancer wherein said cancer is chosen from breast cancer, lung cancer, prostate cancer, colorectal cancer, brain cancer, skin cancer, blood cancer (e.g., leukemia, including, for example, acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia), lymphoma

and myeloma, comprising administering to a patient (preferably a human) in need of such treatment or prevention, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or as defined in the embodiments described in more detail herein) sufficient to treat or prevent such cancer. In an even more specific aspect, said cancer is chosen from prostate, brain, colorectal, lung, breast, skin, and blood cancer. In 5 one specific aspect, the cancer is prostate cancer. In one specific aspect, the cancer is lung cancer. In one specific aspect, the cancer is brain cancer. In one specific aspect, the cancer is blood cancer (e.g., leukemia, including, for example, acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia). In one specific aspect, the cancer is breast cancer. In one specific 10 aspect, the cancer is colorectal cancer. In one specific aspect, the cancer is lymphoma. In one specific aspect, the cancer is myeloma. In a preferred embodiment, the method comprises administering a therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic sufficient for treating or preventing said cancer. In a preferred aspect, the therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic is an amount sufficient to inhibit LSD1. In another preferred aspect, the therapeutically effective amount is an amount 15 sufficient to modulate histone methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-4 methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-9 methylation levels. While the present invention relates to both the treatment and the prevention of cancer, the treatment of cancer 20 is particularly preferred.

20 In a related aspect, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or as defined in the embodiments described in more detail herein), for use in the treatment or prevention of cancer. In another related aspect, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic for use in the treatment or prevention of a cancer wherein said cancer is chosen from breast cancer, lung cancer, prostate 25 cancer, colorectal cancer, brain cancer, skin cancer, blood cancer (e.g., leukemia, including, for example, acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia), lymphoma and myeloma. In a more specific aspect, said cancer is chosen from prostate, brain, colorectal, lung, breast, skin, and blood cancer. In one specific aspect, the cancer is prostate cancer. In one specific aspect, the cancer is lung cancer. In one specific aspect, the cancer is brain cancer. In one specific aspect, the cancer is blood cancer (e.g., leukemia, including, for example, acute myelogenous leukemia (AML), 30 chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia). In one specific aspect, the cancer is breast cancer. In one specific aspect, the cancer is colorectal cancer. In one specific aspect, the cancer is lymphoma. In one specific aspect, the cancer is myeloma. In preferred embodiment, a therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic sufficient for treating or preventing 35

5 said cancer is administered. In a preferred aspect, the therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic is an amount sufficient to inhibit LSD1. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-4 methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-9 methylation levels.

10 In another aspect, the invention provides a method of treating or preventing a neurological disease (e.g., a neurodegenerative disease) comprising administering, to a patient in need of such treatment or prevention, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or in the embodiments described in more detail herein) sufficient to treat or prevent said neurological disease. In a related aspect, the invention provides a method of treating or preventing a neurological disease wherein said neurological disease is selected from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, Dementia with Lewy Bodies, or Frontotemporal Dementia, particularly from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, or Dementia with Lewy Bodies, comprising administering to 15 a patient (preferably a human) in need of such treatment or prevention, an amount of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or as defined in the embodiments described in more detail herein) sufficient to treat or prevent such neurological disease. In a preferred embodiment, the method comprises administering a therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic sufficient for treating or preventing said neurological disease. In a preferred aspect, the therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic is an amount sufficient to inhibit LSD1. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-4 methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-9 methylation levels.

20 In another related aspect, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or in the embodiments described in more detail herein) for use in the treatment or prevention of a neurological disease (e.g., a neurodegenerative disease). In one embodiment, said neurological disease is selected from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, Dementia with Lewy Bodies and Frontotemporal Dementia, particularly from depression, Alzheimer's disease, Huntington disease, Parkinson's disease and Dementia with Lewy Bodies. In a preferred embodiment, a 25 therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic sufficient for treating or preventing said neurological disease is administered. In a preferred aspect, the therapeutically effective amount of a compound of Formula I, Ia, Ia-1, Ib or Ic is an amount sufficient to inhibit LSD1. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone methylation levels. In another

preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-4 methylation levels. In another preferred aspect, the therapeutically effective amount is an amount sufficient to modulate histone-3 lysine-9 methylation levels.

5 In another aspect, the invention provides a method of treating or preventing a viral infection comprising administering to a patient in need thereof (preferably a human) an amount of a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein) sufficient to treat or prevent said viral infection. In a related aspect, the invention also provides a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein) for use in treating or preventing a 10 viral infection. In one specific embodiment, the viral infection is a herpesvirus infection. In a more specific embodiment, the herpesvirus infection is caused by and/or associated with a herpesvirus chosen from HSV-1, HSV-2, and Epstein-Barr virus. In another embodiment, the viral infection is caused by and/or associated with HIV. In another embodiment, the viral infection is caused by and/or associated with a Hepadnavirus (i.e., a 15 virus of the *Hepadnaviridae* family), particularly Hepatitis B virus (HBV). In another embodiment, the viral infection is caused by and/or associated with a Flavivirus (i.e., a virus of the *Flaviviridae* family), particularly Hepatitis C virus (HCV), yellow fever virus, West Nile virus, Dengue virus or Japanese encephalitis virus, and more preferably HCV. In an even more specific embodiment, the invention provides a method for treating or preventing viral reactivation after latency, the method comprising administering to an individual (preferably a 20 human) a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein). Accordingly, the invention also provides a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein) for use in treating or preventing viral reactivation after latency. In a specific embodiment, the virus that is reactivating is a herpesvirus. In a more specific embodiment, the herpesvirus that is reactivating is chosen from HSV-1, HSV-2, and Epstein-Barr virus. In an even more specific embodiment, the virus that is reactivating is HSV. In a further specific embodiment, the virus 25 that is reactivating is HIV.

In still another aspect, the invention provides the use of a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein) for the manufacture of a medicament for the treatment or prevention of cancer. In a preferred embodiment, said cancer is chosen from breast cancer, lung 30 cancer, prostate cancer, colorectal cancer, brain cancer, skin cancer, blood cancer (e.g., leukemia, including, for example, acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia), lymphoma and myeloma.

35 In still another aspect, the invention provides the use of a compound of Formula I, Ia, Ia-1, I_b or I_c (as defined above or in the embodiments described in more detail herein) for the manufacture of a medicament for the


treatment or prevention of a neurological disease (e.g., a neurodegenerative disease). In a preferred embodiment said neurological disease is selected from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, Dementia with Lewy Bodies, or Frontotemporal Dementia, particularly from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, and Dementia with Lewy Bodies.

5 In still another aspect, the invention provides the use of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or in the embodiments described in more detail herein) for the manufacture of a medicament for the treatment or prevention of a viral infection. In a preferred embodiment, said viral infection is a herpesvirus 10 infection (e.g., a herpesvirus infection caused by and/or associated with a herpesvirus chosen from HSV-1, HSV-2, and Epstein-Barr virus) or a viral infection caused by and/or associated with HIV. In another preferred embodiment, said viral infection is caused by and/or associated with a Hepadnavirus, particularly Hepatitis B virus (HBV). In another embodiment, said viral infection is caused by and/or associated with a Flavivirus, particularly Hepatitis C virus (HCV), yellow fever virus, West Nile virus, Dengue virus or Japanese encephalitis 15 virus, and more preferably HCV.


In still another aspect, the invention provides the use of a compound of Formula I, Ia, Ia-1, Ib or Ic (as defined above or in the embodiments described in more detail herein) for the manufacture of a medicament for the treatment or prevention of viral reactivation after latency. In a preferred embodiment, the virus that is 20 reactivating is a herpesvirus (e.g., HSV-1, HSV-2, or Epstein-Barr virus), HSV, or HIV.

In still another aspect, the invention provides a method for identifying a compound which is a selective inhibitor 25 of LSD1, the method comprising selecting or providing a compound of Formula I, Ia, Ia-1, Ib or Ic as defined herein, and determining the ability of the compound to inhibit LSD1 and MAO-A and/or MAO-B, wherein a compound that inhibits LSD1 to a greater extent than MAO-A and/or MAO-B is identified as a LSD1 selective inhibitor. The compound of this aspect that is an LSD1 inhibitor can be used to treat disease, particularly 30 human disease.


In another aspect, the invention provides a process for the preparation of a compound of formula I, or a salt 30 thereof, which comprises reacting a compound of formula II




wherein A, B, R^w, R^x, R^y, R^z have the meaning disclosed above in relation to a compound of formula I, with a compound of formula IIIa, IIIb or IIIc



IIIa



IIIb



IIIc

wherein D has the meaning disclosed above in relation to a compound of formula I, and wherein any amino group that may be present in D is optionally protected with a protecting group,
5 in the presence of a reducing agent, followed by the removal of any protecting group that may be present. The reducing agent may be, e.g., a borohydride, such as sodium borohydride or sodium triacetoxyborohydride.


Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

15 Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.


DETAILED DESCRIPTION OF THE INVENTION

20 The present invention relates to the identification of compounds and their use in treating and preventing diseases. The present invention provides compounds of Formula I, Ia, Ia-1, Ib and Ic, pharmaceutical compositions comprising a compound of Formula I, Ia, Ia-1, Ib or Ic and a pharmaceutically acceptable carrier, and their use for treating diseases. One use of the compounds of Formula I, Ia, Ia-1, Ib and Ic is for treating cancer.

25 The present invention provides a compound of Formula I and relates, in particular, to the compound of Formula I for use as a medicament.



In a compound of formula I, each R^w , R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl. In one embodiment, each R^w , R^x , R^y and R^z is independently selected from hydrogen, fluoro and C_{1-4} alkyl, preferably from hydrogen, fluoro and methyl. In another embodiment, each R^w , R^x , R^y and R^z is independently selected from hydrogen and fluoro. In another embodiment, R^w is fluoro and each R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl; preferably, R^w is fluoro and each R^x , R^y and R^z is hydrogen. In another embodiment, R^z is fluoro and each R^w , R^x and R^y is independently selected from hydrogen, halo and C_{1-4} alkyl; preferably, R^z is fluoro and each R^w , R^x and R^y is hydrogen. In another embodiment, R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C_{1-4} alkyl; preferably, R^w and R^z are fluoro and each R^x and R^y is hydrogen. In a preferred embodiment, R^w is selected from hydrogen, halo and C_{1-4} alkyl, preferably from hydrogen, fluoro and methyl, and each R^x , R^y and R^z is hydrogen. In a more preferred embodiment, each R^w , R^x , R^y and R^z is hydrogen. A compound of formula I wherein each R^w , R^x , R^y and R^z is hydrogen is a compound of formula Ia, which can be depicted as follows:



Ia


15

In another embodiment, in a compound of formula I each R^w , R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl with the proviso that at least one of R^w , R^x , R^y and R^z is not hydrogen, this is a compound of formula Ib. In a more preferred embodiment, R^w is selected from halo and C_{1-4} alkyl, preferably fluoro and methyl, and each R^x , R^y and R^z is hydrogen. A compound of formula I wherein R^w is selected from halo and C_{1-4} alkyl, preferably fluoro and methyl, and each R^x , R^y and R^z is hydrogen is a compound of formula Ic. Preferably, in a compound of formula Ic R^w is methyl.

The invention also relates to a compound of formula Ia, Ia-1, Ib and Ic as defined above, and in particular to a compound of formula Ia, Ia-1, Ib and Ic for use as a medicament.

In a compound of formula I, Ia, Ia-1, Ib or Ic, the group A is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R^1 . In one embodiment, A is aryl (preferably phenyl or naphthyl) optionally substituted with one or more R^1 . In a specific embodiment, A is phenyl optionally substituted with one or more R^1 . In another specific embodiment, A is naphthyl (preferably 2-naphthyl) optionally substituted with one or more R^1 . In another embodiment, A is heteroaryl optionally substituted with one or more R^1 . In a more specific embodiment, A is monocyclic heteroaryl optionally substituted with one or more R^1 . In a preferred embodiment, A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R^1 . Preferably, A is monocyclic aryl (i.e. phenyl) or monocyclic heteroaryl, wherein said monocyclic aryl or said monocyclic heteroaryl is

optionally substituted with one or more R¹. More preferably, A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein A (i.e. said phenyl, said pyridyl, said thiophenyl, said pyrrolyl, said furanyl, or said thiazolyl) is optionally substituted with one or more R¹. Still more preferably, A is phenyl, pyridyl or thiazolyl, wherein A is optionally substituted with one or more R¹. Even more preferably, A is phenyl, 3-pyridyl or 5-thiazolyl, as shown below:



wherein A is optionally substituted with one or more R¹. In one embodiment, A is phenyl or pyridyl, preferably phenyl or 3-pyridyl. In another embodiment, A is phenyl. In another embodiment, A is pyridyl, preferably 3-pyridyl. In another embodiment, A is thiazolyl, preferably 5-thiazolyl. In one embodiment, A has 0, 1 or 2 substituents R¹. In a further embodiment, A has 0 or 1 substituent R¹. In a further embodiment, A has 0 substituent R¹. In a further embodiment, A has 1 or 2 substituents R¹. In a further embodiment, A has 1 substituent R¹. In the aforementioned embodiments, in which A has 0, 1 or 2 substituents R¹, the total number of substituents R¹ is defined, including the possibility that B may be R¹. Accordingly, if A has 0 substituents R¹, then B is not R¹.

15

In a compound of formula I, Ia, Ia-1, Ib or Ic, B is hydrogen, R¹ or -L¹-E. In one embodiment, B is -L¹-E. In another embodiment, B is hydrogen or R¹. In a further embodiment, B is hydrogen. In another embodiment, B is R¹.

20 In a compound of formula I, Ia, Ia-1, Ib or Ic, E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R². In one embodiment, E is an aryl group (e.g., phenyl, naphthyl or anthracenyl) optionally substituted with one or more R². In another embodiment, E is a heteroaryl group (e.g., pyridinyl, thiophenyl, pyrrolyl, furanyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolyl, indolyl, pyrazolyl, indazolyl, imidazolyl or benzimidazolyl) optionally substituted with one or more R². In another embodiment, E is pyridinyl, thiophenyl, pyrrolyl, furanyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolyl, indolyl, indazolyl, imidazolyl or benzimidazolyl, wherein said pyridinyl, said thiophenyl, said pyrrolyl, said furanyl, said thiazolyl, said oxazolyl, said isoxazolyl, said oxadiazolyl, said thiadiazolyl, said triazinyl, said pyridazinyl, said pyrazinyl, said pyrimidinyl, said quinolyl, said indolyl, said indazolyl, said imidazolyl or said benzimidazolyl is optionally substituted with one or more R². Preferably, E is monocyclic aryl (i.e. phenyl) or monocyclic heteroaryl, wherein said monocyclic aryl or said monocyclic heteroaryl is optionally substituted with one or more R². In one preferred embodiment, E is phenyl or pyridinyl, wherein said phenyl or said pyridinyl is optionally

substituted with one or more R². In one embodiment, E has 0, 1, 2 or 3 substituents R². In another embodiment, E has 0, 1 or 2 substituents R². In another embodiment, E has 0 or 1 substituents R². In another embodiment, E has 0 substituents R². In another embodiment, E has 1 substituent R². Preferably, E is phenyl optionally substituted with one or more R². In one embodiment, E is phenyl optionally substituted with one, two or three R². In another embodiment, E is phenyl optionally substituted with one or two R². In a further embodiment, E is phenyl optionally substituted with one R². In another embodiment, E is phenyl. In another embodiment, E is phenyl substituted with one, two or three, preferably one or two, R². In another preferred embodiment, E is heteroaryl, preferably monocyclic heteroaryl, optionally substituted with one or more (preferably one, two or three) R². In one embodiment, E is heteroaryl, preferably monocyclic heteroaryl. In another embodiment, E is heteroaryl (preferably monocyclic heteroaryl) substituted with one, two or three, preferably one or two, R².

10 In a compound of formula I, Ia, Ia-1, Ib or Ic, L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄ alkylene. Preferably said heteroC₁₋₄ alkylene is -(CH₂)_x-NH-, or -(CH₂)_x-O-, wherein x is 1, 2, 3 or 4; still more preferably, said -(CH₂)_x-NH- or -(CH₂)_x-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -(CH₂)_x group. More preferably, said heteroC₁₋₄ alkylene is -CH₂-NH- or -CH₂-O-, wherein said -CH₂-NH- and -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group.

15 In one embodiment, L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, -CH₂-, CH₂-CH₂-, -CH₂-NH-, or -CH₂-O-. In another embodiment, L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, -CH₂-NH-, or -CH₂-O-. In another embodiment, L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-. In another embodiment, L¹ is a bond, -O-, -NH-, or -CH₂-NH-. In another embodiment, L¹ is a bond or -CH₂-O-. In another embodiment, L¹ is -NH- or -CH₂-NH-. In a further embodiment, L¹ is a bond. In another embodiment, L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄ alkylene; preferably, L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, -CH₂-, CH₂-CH₂-, -CH₂-NH-, or -CH₂-O-; more preferably L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, -CH₂-NH-, or -CH₂-O-; even more preferably L¹ -O-, -NH-, -CH₂-NH-, or -CH₂-O-; and particularly preferably L¹ is -CH₂-O-. In another embodiment, L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, -(CH₂)_x-NH-, -SO₂N(CH₃)-, wherein x is 1, 2, 3 or 4; preferably L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-, and more preferably L¹ is -O-, -NH- or -CH₂-NH-. Preferably, in all these embodiments, said -(CH₂)_x-NH-, -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group.

20 In one embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein the groups -CH₂-NH- and -CH₂-O- are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is phenyl, pyridinyl, thiophenyl, pyrrolyl, furanyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolyl, indolyl,

indazolyl, imidazolyl or benzimidazolyl, wherein said phenyl, said pyridinyl, said thiophenyl, said pyrrolyl, said furanyl, said thiazolyl, said oxazolyl, said isoxazolyl, said oxadiazolyl, said thiadiazolyl, said triazinyl, said pyridazinyl, said pyrazinyl, said pyrimidinyl, said quinolyl, said indolyl, said indazolyl, said imidazolyl or said benzimidazolyl is optionally substituted with one or more R²; and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein the groups -CH₂-NH- and -CH₂-O- are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is phenyl optionally substituted with one or more R²; and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein the groups -CH₂-NH- and -CH₂-O- are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and L¹ is a bond or -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is phenyl optionally substituted with one or more R²; and L¹ is a bond or -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and L¹ is a bond. In another embodiment, B is -L¹-E; E is aryl, pyridinyl, thiophenyl, pyrrolyl, furanyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolyl, indolyl, indazolyl, imidazolyl or benzimidazolyl, wherein said aryl, said pyridinyl, said thiophenyl, said pyrrolyl, said furanyl, said thiazolyl, said oxazolyl, said isoxazolyl, said oxadiazolyl, said thiadiazolyl, said triazinyl, said pyridazinyl, said pyrazinyl, said pyrimidinyl, said quinolyl, said indolyl, said indazolyl, said imidazolyl or said benzimidazolyl is optionally substituted with one or more R²; and L¹ is a bond. In another embodiment, B is -L¹-E; E is aryl, wherein said aryl is optionally substituted with one or more R²; and L¹ is a bond. In another embodiment, B is -L¹-E; E is phenyl optionally substituted with one or more R²; and L¹ is a bond. In another embodiment, B is -L¹-E; E is pyridinyl optionally substituted with one or more R²; and L¹ is a bond. In another embodiment, B is -L¹-E; E is aryl or heteroaryl; and L¹ is a bond. In another embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and L¹ is -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is heteroaryl (preferably monocyclic heteroaryl, more preferably pyridinyl), wherein said heteroaryl is optionally substituted with one or more R²; and L¹ is -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is phenyl optionally substituted with one or more R²; and L¹ is -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, -(CH₂)_x-NH-, -SO₂NH- or -SO₂N(CH₃)-, wherein x is 1, 2, 3 or 4, preferably L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-, and more preferably L¹ is -O-, -NH- or -CH₂-NH- wherein the group -CH₂-NH- is linked to ring A through the N atom and to ring E through the -CH₂- group. In another embodiment, B is -L¹-E; E is aryl or heteroaryl, wherein said aryl or said heteroaryl is

optionally substituted with one or more R²; and L¹ is -NH- or -CH₂-NH-, wherein the group -CH₂-NH- is linked to ring A through the N atom and to ring E through the -CH₂- group.

In a compound of formula I, Ia, Ia-1, Ib or Ic, each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea. When there is more than one R¹ as a substituent on ring A, they can be the same or different. In one embodiment, each R¹ is independently selected from C₁₋₈ alkyl, cyclyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another embodiment, each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another embodiment, each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In a compound of formula I, Ia, Ia-1, Ib or Ic, each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea. When there is more than one R² as a substituent on ring E, they can be the same or different. In one embodiment, each R² is independently selected from C₁₋₈ alkyl, cyclyl, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide and C₁₋₈ alkoxy. In another embodiment, each R² is independently selected from C₁₋₈ alkyl, cyclyl, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, N-sulfonamido and C₁₋₈ alkoxy. In another embodiment, each R² is independently selected from hydroxyl, halo (for example fluoro or chloro), haloC₁₋₈ alkyl (for example trifluoromethyl) and sulfonamide (preferably N-sulfonamido). In another embodiment, each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido. In another embodiment, each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and -NR'SO₂R (wherein R and R' are as defined herein below; preferably R' is H and R is C₁₋₈ alkyl (for example, methyl, ethyl or isopropyl) or R' is H and R is optionally substituted phenyl). In another embodiment, each R² is independently selected from hydroxyl, halo and haloC₁₋₈ alkyl. In another embodiment, each R² is independently selected from hydroxyl, halo and haloC₁₋₄ alkyl. In another embodiment, each R² is independently selected from hydroxyl, chloro, fluoro or trifluoromethyl. In a further embodiment, ring E is substituted with one R² and said R² is N-sulfonamido, i.e. -NR'SO₂R (wherein R and R' are as defined herein below; preferably R' is H and R is C₁₋₈ alkyl (for example, methyl, ethyl or isopropyl) or R' is H and R is optionally substituted phenyl).

In a compound of formula I, Ia, Ia-1, Ib or Ic, L² and D have the following meanings:

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- 5 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I (i.e., of formula I, Ia, Ia-1, Ib or Ic, respectively) through a ring C atom,

10 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

15 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) 20 a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I (i.e., of formula I, Ia, Ia-1, Ib or Ic, respectively) through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

25 wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

In one embodiment, L² is a bond and D is a cyclic group selected from:

- (i) 30 a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

35 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I, Ia, Ia-1, Ib or Ic through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

5 D can thus be a monocyclic ring (group i) or polycyclic ring (group ii). "Polycyclic ring system" as used herein means a ring system containing 2 or more rings, preferably containing two (i.e. bicyclic) or three (i.e. tricyclic) rings, which rings can be fused, bridged or spiro rings, or any combination thereof. Thus, tricyclic rings include for example three fused rings, or two rings fused together, one of which is a spiro bicyclic ring. A fused ring as used herein means that two rings are linked to each other through two adjacent ring atoms common to both 10 rings. A bridged ring as used herein means that a ring comprises a linker group -(C(R^a)₂)_p- linking together any two non-adjacent carbon or nitrogen atoms of the ring, wherein p is 1 or 2 and each R^a independently is hydrogen or C₁₋₄ alkyl. A spiro ring as used herein means two rings linked together via a single carbon atom common to both rings.

15 Ring D can contain 1 or 2 heteroatoms independently selected from N, O and S in case of cyclic group (i) or from 1 to 4 heteroatoms independently selected from N, O and S in case of cyclic group (ii). In one embodiment, D contains 1 or 2 heteroatoms independently selected from N, O and S in case of cyclic group (i) or from 1 to 4 heteroatoms independently selected from N, O and S in case of cyclic group (ii), wherein ring D contains at least 1 N atom. In another embodiment, cyclic group (i) contains 1 heteroatom selected from N, O and S. In another embodiment, cyclic group (i) contains 1 N atom. In another embodiment, cyclic group (ii) 20 contains 1 or 2 heteroatoms independently selected from N, O and S. In another embodiment, cyclic group (ii) contains 1 or 2 N atoms.

25 Cyclic group (ii) as defined above refers to a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring. One ring forming part of cyclic group (ii) must thus be a saturated heterocyclic ring, while the other ring(s) can be saturated, partially unsaturated or aromatic carbocyclic or heterocyclic rings. It is to be understood that, in the polycyclic ring group (ii), a saturated heterocyclic ring which is fused to an aromatic ring is still to be considered as a saturated heterocyclic ring, even if the two adjacent ring atoms common to both rings (i.e., common to the saturated heterocyclic ring and to the aromatic ring) are connected by a double bond. Accordingly, the cyclic group (ii) may also be, e.g., a saturated heterocyclic ring which is fused to an aromatic carbocyclic or heterocyclic ring. In one embodiment, cyclic group (ii) is a 30 saturated polycyclic ring system, i.e. all rings forming cyclic group (ii) are saturated.

D (i.e. groups (i) or (ii)) is linked to the remainder of the compound of Formula I through a ring C atom. In the case of cyclic group (ii), the C atom linking D to the remainder of the compound of formula I can be in any of the rings and thus can be in the saturated heterocyclic ring or any other ring forming part of cyclic group (ii).

35 D, i.e. cyclic group (i) or (ii), may be optionally substituted with one or more R³, which can be the same or different and may be placed at any available carbon or nitrogen atom of cyclic group (i) or of any of the rings of

cyclic group (ii). In one embodiment, the substituent(s) R³, if present, are placed on ring C atom(s). In another embodiment, D is not substituted with any R³.

In one embodiment, L² is a bond and D is a cyclic group selected from:

5 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

10 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I (i.e., the remainder of the compound of Formula I, Ia, Ia-1, Ib or Ic, respectively) through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

15 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

In another embodiment, L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

20 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

25 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

30 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

In another embodiment, L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

35 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,
wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,
5 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

In another embodiment, L² is a bond and D is a cyclic group selected from:

10 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,
wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,
15 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and
wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

In another embodiment, L² is a bond and D is a cyclic group selected from:

20 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
(ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 heteroatoms independently selected from N, O and S,
wherein the cyclic group (i) or (ii) contains at least one N atom,
25 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,
wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
30 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

In another embodiment, L² is a bond and D is a cyclic group selected from:

35 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
(ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

5

In another embodiment, L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, and
- (ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 N atom,

10 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

15 In another embodiment, L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

25 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³, wherein said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

30 In another embodiment, L² is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

35 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³.

In another embodiment, L^2 is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

5 wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO_2 groups, and

wherein ring D is optionally substituted with one or more R^3 .

10 In another embodiment, L^2 is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one S atom in ring D, if present, is optionally oxidized to form independently SO group or SO_2 group,

15 and

wherein ring D is optionally substituted with one or more R^3 .

In another embodiment, L^2 is a bond and D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S,

20 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one S atom in ring D, if present, is optionally oxidized to form independently SO group or SO_2 group, and

wherein ring D is optionally substituted with one or more R^3 .

25

In another embodiment, L^2 is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and

30 wherein ring D is optionally substituted with one or more R^3 .

In another embodiment, L^2 is a bond and D is piperidinyl,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and

35 wherein ring D is optionally substituted with one or more R^3 .


In another embodiment, L² is a bond and D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and wherein said piperidinyl is optionally substituted with one or more R³.

5 In another embodiment, L² is a bond and D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and wherein said piperidinyl is optionally substituted with one or more R³, wherein said substituent(s) R³, if present, are placed on C atom(s) of said piperidine ring.

10 In another embodiment, L² is a bond and D is 4-piperidinyl optionally substituted with one or more R³. In another embodiment, L² is a bond and D is 4-piperidinyl optionally substituted with one or more R³, wherein said substituent(s) R³, if present, are placed on C atom(s) of said piperidine ring.

15 In a specific embodiment, L² is a bond and D is 4-piperidinyl.

In another specific embodiment, L² is a bond and D is a group of the following formula:




20 In another embodiment, L² is a bond and D is a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.

30 In another embodiment, L² is a bond and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
 wherein one or more ring C atoms in D are optionally oxidized to form CO groups,
 wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
 5 wherein D is optionally substituted with one or more R³.

In a preferred embodiment, L² is a bond and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 or 2 N atoms,
 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
 10 wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and
 wherein D is optionally substituted with one or more R³.


In a specific embodiment, L² is a bond and D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)




(c)



(d)

15 wherein D is optionally substituted with one or more R³. Said substituent(s) R³, if present, can be placed on any available position, including on the N atom.


In another specific embodiment, L² is a bond and D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b) and (c)



(a)



(b)



(c)

wherein D is optionally substituted with one or more R³. Said substituent(s) R³, if present, can be placed on any available position, including on the N atom.

5 In a further embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

10 wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I, Ia, Ia-1, Ib or Ic through a ring C atom,

15 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

20 D can thus be a monocyclic saturated ring (group iii) or polycyclic saturated ring (group iv). "Polycyclic saturated ring system" as used herein means a ring system containing 2 or more saturated rings, preferably containing two (i.e. bicyclic) or three (i.e. tricyclic) saturated rings, which rings can be fused, bridged or spiro rings, or any combination thereof. Thus, tricyclic rings include for example three fused rings, or two rings fused together, one of which is a spiro bicyclic ring. A fused ring as used herein means that two rings are linked to each other through two adjacent ring atoms common to both rings. A bridged ring as used herein means that a ring comprises a linker group -(C(R^a)₂)_p- linking together any two non-adjacent carbon or nitrogen atoms of the ring, wherein p is 1 or 2 and each R^a independently is hydrogen or C₁₋₄ alkyl. A spiro ring as used herein means 25 two rings linked together via a single carbon atom common to both rings.

Ring D can contain 1 or 2 heteroatoms independently selected from N, O and S in case of cyclic group (iii) or from 1 to 4 heteroatoms independently selected from N, O and S in case of cyclic group (iv). In one embodiment, D contains 1 or 2 heteroatoms independently selected from N, O and S in case of cyclic group (iii) or from 1 to 4 heteroatoms independently selected from N, O and S in case of cyclic group (iv), wherein ring D 30 contains at least 1 N atom. In another embodiment, cyclic group (iii) contains 1 heteroatom selected from N, O and S. In another embodiment, cyclic group (iv) contains 1 N atom. In another embodiment, cyclic group (iii) contains 1 or 2 heteroatoms independently selected from N, O and S. In another embodiment, cyclic group (iv) contains 1 or 2 N atoms.

35 Cyclic group (iv) as defined above refers to a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring. One ring forming part of cyclic group (iv) must thus be a saturated heterocyclic ring, while the other ring(s) can be saturated carbocyclic or heterocyclic rings.

D (i.e. groups (iii) or (iv)) is linked to the remainder of the compound of Formula I through a ring C atom. In the case of cyclic group (iv), the C atom linking D to the remainder of the compound of formula I can be in any of the rings and thus can be in the saturated heterocyclic ring or any other (carbocyclic or heterocyclic) saturated ring forming part of cyclic group (iv).

5 D, i.e. cyclic group (iii) or (iv), may be optionally substituted with one or more R³, which can be the same or different and may be placed at any available carbon or nitrogen atom of cyclic group (iii) or of any of the rings of cyclic group (iv). In one embodiment, the substituent(s) R³, if present, are placed on ring C atom(s). In another embodiment, D is not substituted with any R³.

10 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

15 wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I (i.e., the remainder of the compound of Formula I, Ia, Ia-1, Ib or Ic, respectively) through a ring C atom,

20 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

25 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 2 heteroatoms independently selected from N, O and S,

30 wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

35 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

5 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 2 N atoms,

10 wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

15 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 N atom,

20 wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

25

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered saturated polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

35 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³, wherein said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

5

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

10 wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,
wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³.

15 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

20 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein ring D is optionally substituted with one or more R³.

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, 25 O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

30 wherein one S atom in ring D, if present, is optionally oxidized to form independently SO group or SO₂ group, and

wherein ring D is optionally substituted with one or more R³.

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and 35 S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,
wherein one S atom in ring D, if present, is optionally oxidized to form independently SO group or SO₂ group,
and
wherein ring D is optionally substituted with one or more R³.

5

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and
10 wherein ring D is optionally substituted with one or more R³.

10

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
and D is piperidinyl,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
15 wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and
wherein ring D is optionally substituted with one or more R³.

15

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
and D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a
20 ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R³.

20

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
and D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a
25 ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R³, wherein said substituent(s) R³, if present,
are placed on C atom(s) of said piperidine ring.

25

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
30 and D is 4-piperidinyl optionally substituted with one or more R³.


30

In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-)
and D is 4-piperidinyl optionally substituted with one or more R³, wherein said substituent(s) R³, if present, are
placed on C atom(s) of said piperidine ring.

35

In a specific embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is 4-piperidinyl.

5 In another specific embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a group of the following formula:



10 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is azetidinyl, wherein said azetidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said azetidinyl is optionally substituted with one or more R³.


15 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is 3-azetidinyl optionally substituted with one or more R³.

20 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

25 wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.

30 In another embodiment, L² is C₁₋₄ alkylene (preferably linear C₁₋₄ alkylene, more preferably -CH₂- or -CH₂CH₂-) and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 or 2 N atoms, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and wherein D is optionally substituted with one or more R³.


In another embodiment, L^2 is C_{1-4} alkylene (preferably linear C_{1-4} alkylene, more preferably $-CH_2-$ or $-CH_2CH_2-$) and D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)



(c)




(d)


5

wherein D is optionally substituted with one or more R^3 . Said substituent(s) R^3 , if present, can be placed on any available position, including on the N atom.


In another embodiment, L^2 is C_{1-4} alkylene (preferably linear C_{1-4} alkylene, more preferably $-CH_2-$ or $-CH_2CH_2-$) and D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b) and (c)



(a)



(b)



(c)


wherein D is optionally substituted with one or more R^3 . Said substituent(s) R^3 , if present, can be placed on any available position, including on the N atom.

In a compound of formula I, Ia, Ia-1, Ib or Ic, each R^3 is independently selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, cyclol, amino, amido, hydroxyl, nitro, halo, halo C_{1-8} alkyl, halo C_{1-8} alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C_{1-8} alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea. In one embodiment, each R^3 is independently selected from C_{1-8} alkyl, amino, amido, hydroxyl, nitro, halo, halo C_{1-8} alkyl, halo C_{1-8} alkoxy, cyano, sulfonyl, sulfonamide, C_{1-8} alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea. In another embodiment, each R^3 is independently selected from C_{1-8} alkyl, halo, C_{1-8} alkoxy, hydroxyl, amino, and amido.

20

In another embodiment, each R^3 is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

5 The substituents of the cyclopropyl moiety $-A-B$ and $-NH-L^2-D$ in a compound of formula I, **Ia**, **Ib** or **Ic** are preferably in the *trans*- configuration. Thus, in one embodiment, the invention provides a compound of formula I (including a compound **Ia**, **Ib** or **Ic**) wherein the groups $-A-B$ and $-NH-L^2-D$ are in *trans* configuration. In a preferred embodiment, the invention provides a compound of formula **Ia** wherein the groups $-A-B$ and $-NH-L^2-D$ are in *trans* configuration, which is a compound of formula **Ia-1**:

**Ia-1**

10

wherein the groups and variables of formula **Ia-1**, including A, B, D, E, L^1 , L^2 , R^1 , R^2 and R^3 , are as defined above in relation to a compound of formula I and **Ia** and in the various preferred embodiments for a compound of formula I and **Ia** described above. The above chemical representation for a compound of formula **Ia-1** does not intend to indicate absolute stereochemistry of the two chiral centers on the cyclopropyl ring, but only their relative stereochemistry (which is *trans*). Thus a compound of formula **Ia-1** therefore relates to individual optically active *trans* isomers as well as mixtures of *trans*-isomers.

15

In one embodiment, the invention provides a compound of formula I wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, fluoro and C_{1-4} alkyl, preferably from hydrogen, fluoro and methyl.

20

In another embodiment, the invention provides a compound of formula I wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen and fluoro.

25

In another embodiment, the invention provides a compound of formula I wherein R^w is fluoro and each R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl; preferably, R^w is fluoro and each R^x , R^y and R^z is hydrogen.


30

In another embodiment, the invention provides a compound of formula I wherein R^z is fluoro and each R^w , R^x and R^y is independently selected from hydrogen, halo and C_{1-4} alkyl; preferably, R^z is fluoro and each R^w , R^x and R^y is hydrogen.

In another embodiment, the invention provides a compound of formula I wherein R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl; preferably, R^w and R^z are fluoro and each R^x and R^y is hydrogen.

5 In a preferred embodiment, the invention provides a compound of formula I wherein R^w is selected from hydrogen, halo and C₁₋₄ alkyl, preferably from hydrogen, fluoro and methyl, and each R^x, R^y and R^z is hydrogen.

In a more preferred embodiment, the invention provides a compound of formula I wherein each R^w, R^x, R^y and R^z is hydrogen, i.e. a compound of formula Ia:



In another embodiment, the invention provides a compound of formula I wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl with the proviso that at least one is not hydrogen, i.e. 15 a compound of formula Ib.

In another embodiment, the invention provides a compound of formula I wherein R^w is selected from halo and C₁₋₄ alkyl, preferably from fluoro and methyl, and each R^x, R^y and R^z is hydrogen, i.e. a compound of formula Ic. Preferably, in a compound of formula Ic R^w is methyl.

20 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein L² is a bond or linear C₁₋₄ alkylene, preferably L² is a bond or linear C₁₋₂ alkylene.

25 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein L² is a bond.

30 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein L² is CH₂.

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein L² is CH₂CH₂.

10 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond; and

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

15 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

20 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

25 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

30 L² is a bond; and

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

5 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

10 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, , Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond; and

D is a cyclic group selected from:

15 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

20 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

25 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond; and

D is a cyclic group selected from:

30 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
(ii) a 7- to 15-membered saturated polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and

35 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

In another embodiment, the invention provides a compound of formula **I**, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein:

5 L^2 is a bond; and
D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,
10 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein ring D is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

15 In another embodiment, the invention provides a compound of formula **I**, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein:
 L^2 is a bond; and
D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and
wherein ring D is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25 In another embodiment, the invention provides a compound of formula **I**, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein:
 L^2 is a bond; and
D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is a bond; and

5 D is a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

10 wherein D is optionally substituted with one or more R³.

In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is a bond; and


D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 or 2 N atoms, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and

20 wherein D is optionally substituted with one or more R³.

In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is a bond; and


D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)



(c)



(d)

wherein D is optionally substituted with one or more R³.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

5 L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene; and

D is a cyclic group selected from:

- 10 (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

15 wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

20 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³. In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (iii) or (iv).

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and
D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 2 N atoms,

5 wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, and
wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R^3 . In a more specific embodiment,
10 said substituent(s) R^3 , if present, are placed on ring C atom(s) of cyclic group (iii) or (iv).

In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

15 L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and
D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

20 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO_2 groups, and

wherein ring D is optionally substituted with one or more R^3 . In a more specific embodiment, said substituent(s) R^3 , if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R^3 , if present, is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

25 In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

30 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and
wherein ring D is optionally substituted with one or more R^3 . In a more specific embodiment, said substituent(s) R^3 , if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R^3 , if present, is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

5 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula **I** through a ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R^3 . In a more specific embodiment, said substituent(s) R^3 , if present, are placed on ring C atom(s) of ring D . In a further specific embodiment, each R^3 , if present, is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

10 In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is CH_2 ; and

15 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula **I** through a ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R^3 . In a more specific embodiment, said substituent(s) R^3 , if present, are placed on ring C atom(s) of ring D . In a further specific embodiment, each R^3 , if present, is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

20 In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is CH_2CH_2 ; and

25 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula **I** through a ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R^3 . In a more specific embodiment, said substituent(s) R^3 , if present, are placed on ring C atom(s) of ring D . In a further specific embodiment, each R^3 , if present, is independently selected from C_{1-4} alkyl, halo, C_{1-4} alkoxy, hydroxyl, amino, and amido.

30 In another embodiment, the invention provides a compound of formula **I**, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula **I**, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:


L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

35 D is 4-piperidinyl.

In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

5 D is a group of formula:



In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

D is azetidinyl, wherein said azetidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said azetidinyl is optionally substituted with one or more R^3 .

15

In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

20 D is 3-azetidinyl optionally substituted with one or more R^3 .

In another embodiment, the invention provides a compound of formula I, **Ia**, **Ia-1**, **Ib** or **Ic** (preferably a compound of formula I, **Ia**, or **Ia-1**, more preferably a compound of formula **Ia** or **Ia-1**, and most preferably a compound of formula **Ia-1**) wherein:

25 L^2 is C_{1-4} alkylene, preferably linear C_{1-4} alkylene, more preferably linear C_{1-2} alkylene; and

D is a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

30 wherein one or more ring C atoms in D are optionally oxidized to form CO groups,

wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein D is optionally substituted with one or more R³.

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene; and

D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, 10 wherein the polycyclic saturated ring system contains 1 or 2 N atoms,


wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and

wherein D is optionally substituted with one or more R³.


15 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene; and


D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d), 20 preferably from a group of formula (a), (b) and (c):




(a)



(b)



(c)



(d)

wherein D is optionally substituted with one or more R³.

25 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a

compound of formula **la-1**) wherein A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, naphthyl or monocyclic heteroaryl is optionally substituted with one or more (e.g., one or two) R¹. In a more preferred embodiment, A is phenyl, naphthyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein A is optionally substituted with one or more R¹. More preferably, A is phenyl, naphthyl, pyridyl or thiazolyl, wherein A is optionally substituted with one or more R¹. Still more preferably, A is phenyl, 2-naphthyl, 3-pyridyl or 5-thiazolyl, wherein A is optionally substituted with one or more R¹. In one embodiment, A is phenyl optionally substituted with one or more R¹. In another embodiment, A is naphthyl, preferably 2-naphthyl, optionally substituted with one or more R¹. In another embodiment, A is pyridyl, preferably 3-pyridyl, optionally substituted with one or more R¹. In another embodiment, A is thiazolyl, preferably 5-thiazolyl, optionally substituted with one or more R¹.

10 In another embodiment, the invention provides a compound of Formula I, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹. In a more preferred embodiment, A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein A is optionally substituted with one or more R¹. More preferably, A is phenyl, pyridyl or thiazolyl, wherein A is optionally substituted with one or more R¹. In one embodiment, A is phenyl. In another embodiment, A is monocyclic heteroaryl. In another embodiment, A is pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl. In another embodiment, A is 3-pyridyl. In another embodiment, A is 5-thiazolyl.

15 In another embodiment, the invention provides a compound of Formula I, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein A is phenyl optionally substituted with one or more R¹.

20 In another embodiment, the invention provides a compound of Formula I, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein A is naphthyl (e.g. 2-naphthyl) optionally substituted with one or more R¹.

25 In another embodiment, the invention provides a compound of Formula I, **la**, **la-1**, **lb** or **lc** (preferably a compound of formula **I**, **la**, or **la-1**, more preferably a compound of formula **la** or **la-1**, and most preferably a compound of formula **la-1**) wherein A is heteroaryl optionally substituted with one or more R¹. Preferably, A is monocyclic heteroaryl optionally substituted with one or more R¹.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

5 B is hydrogen, R¹ or -L¹-E;

E is phenyl optionally substituted with one or more R²; and

10 L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

15 A is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

25 A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

30 In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is heteroaryl, wherein said heteroaryl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is monocyclic heteroaryl, wherein said monocyclic heteroaryl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl, pyridyl or thiazolyl (preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

5 B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In

10 another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

15 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In

20 another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

25 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is naphthyl, wherein said naphthyl is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In the above embodiment, preferably each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, carbamate, and urea. In

30 another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl (e.g. methyl), haloC₁₋₄ alkyl (e.g. trifluoromethyl), C₁₋₄ alkoxy (e.g. methoxy) and C₃₋₆ cycloalkyl (e.g. cyclopropyl). In another preferred embodiment, each R¹ is independently selected from halo, C₁₋₄ alkyl and C₃₋₆ cycloalkyl.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl, naphthyl or monocyclic heteroaryl; and

5 B is hydrogen.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

10 A is phenyl, pyridyl or thiazolyl (preferably phenyl, 3-pyridyl or 5-thiazolyl); and

B is hydrogen.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

15 A is phenyl; and

B is hydrogen.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

20 A is naphthyl, preferably 2-naphthyl; and

B is hydrogen.

25 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is heteroaryl, preferably monocyclic heteroaryl; and

30 B is hydrogen.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

35 A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, and more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹; and

B is $-L^1-E$.

In the above embodiment preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

10 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹; and

15 B is $-L^1-E$.

In the above embodiment preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

20 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is phenyl; and

25 B is $-L^1-E$.

In the above embodiment preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

A is 3-pyridyl; and

5 B is $-L^1-E$.

In the above embodiment preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through 10 the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

15 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is $-L^1-E$;

20 L^1 is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, -(CH₂)_x-NH-, -SO₂NH- or -SO₂N(CH₃)-, wherein x is 1, 2, 3 or 4; preferably L^1 is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH, and more preferably L^1 is -O-, -NH- or -CH₂-NH-; and

25 E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

30 B is $-L^1-E$;

L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein the groups -CH₂-NH- and -CH₂-O- are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group; and

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

35 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is $-L^1-E$;

40 L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein the groups -CH₂-NH- and -CH₂-O- are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group; and

E is phenyl optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

5 B is -L¹-E;

L¹ is a bond or -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group; and

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

10 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is -L¹-E;

15 L¹ is a bond; and

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

20 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is -L¹-E;

L¹ is a bond; and

25 E is aryl, pyridinyl, thiophenyl, pyrrolyl, furanyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolyl, indolyl, indazolyl, imidazolyl or benzimidazolyl, wherein said aryl, said pyridinyl, said thiophenyl, said pyrrolyl, said furanyl, said thiazolyl, said oxazolyl, said isoxazolyl, said oxadiazolyl, said thiadiazolyl, said triazinyl, said pyridazinyl, said pyrazinyl, said pyrimidinyl, said quinolyl, said indolyl, said indazolyl, said imidazolyl or said benzimidazolyl is optionally substituted with one or more R².

30 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is -L¹-E;

L¹ is a bond; and

E is aryl (preferably phenyl), wherein said aryl is optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is $-L^1-E$;

5 L¹ is a bond; and

E is pyridinyl optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

10 B is $-L^1-E$;

L¹ is a bond; and

E is aryl or heteroaryl.

15 In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is $-L^1-E$;

L¹ is a bond;

20 E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²; and each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

25 B is $-L^1-E$;

L¹ is $-CH_2-O-$, wherein the group $-CH_2-O-$ is linked to ring A through the O atom and to ring E through the $-CH_2-$ group; and

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

30

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is $-L^1-E$;

35 L¹ is $-CH_2-O-$, wherein the group $-CH_2-O-$ is linked to ring A through the O atom and to ring E through the $-CH_2-$ group; and

E is heteroaryl (preferably monocyclic heteroaryl, more preferably pyridinyl), wherein said heteroaryl is optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is -L¹-E;

L¹ is -CH₂-O-, wherein the group -CH₂-O- is linked to ring A through the O atom and to ring E through the -CH₂- group; and

E is phenyl optionally substituted with one or more R².

In another embodiment, the invention provides a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

B is -L¹-E;

L¹ is -NH- or -CH₂-NH-, wherein the group -CH₂-NH- is linked to ring A through the N atom and to ring E through the -CH₂- group; and

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a cyclic group selected from:

- 25 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

30 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

35 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is phenyl, pyridyl or thiazolyl (preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

B is hydrogen, R¹ or -L¹-E;

E is phenyl optionally substituted with one or more R²; and

5 L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

10 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

15 L² is a bond;

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

20 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

25 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is aryl or heteroaryl, wherein A is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

30

In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

35 D is a cyclic group selected from:

5 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

10 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, and more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

15 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

20 L² is a bond;

D is a cyclic group selected from:

25 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

30 wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is phenyl optionally substituted with one or more R¹; and

35 B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- 10 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

15 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

20 A is naphthyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

25 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- 30 (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

35 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

5 A is phenyl; and

B is hydrogen.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

10 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a cyclic group selected from:

15 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

20 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

25 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R¹; and

B is -L¹-E.

In the above embodiment, preferably L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

35 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

5 L² is a bond;

D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

10 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

15 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹; and

20 B is -L¹-E.

In the above embodiment, preferably L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

25 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of cyclic group (i) or (ii).

30 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

5 wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 10 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

B is hydrogen, R¹ or -L¹-E;

E is phenyl optionally substituted with one or more R²; and

15 L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

20 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

25 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

30 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

35 B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is phenyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

30 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

35 A is naphthyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

10 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

15 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is phenyl; and

B is hydrogen.

20 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

30 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, heteroatoms independently selected from N, O and S, preferably containing 1 heteroatom selected from N, O and S, and more preferably containing 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

35 A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹; and

B is $-L^1-E$.

In the above embodiment, preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In a more specific embodiment of the above embodiments, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L^2 is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

B is hydrogen, R¹ or $-L^1-E$;

E is phenyl optionally substituted with one or more R²; and

L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L^2 is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

10

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

15

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

20

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

30

wherein said piperidinyl is optionally substituted with one or more R³;

A is naphthyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

35

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

5 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl; and

B is hydrogen.

10 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

20 wherein said piperidinyl is optionally substituted with one or more R³;

A is naphthyl; and

B is hydrogen.

25 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

30 L² is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R¹; and

35 B is -L¹-E.

In the above embodiment, preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In a more specific embodiment of the above embodiments, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

15 L^2 is a bond;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹; and

20 B is -L¹-E.

In the above embodiment, preferably L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L^1 is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L^1 is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

In a more specific embodiment of the above embodiments, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

35 L^2 is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

5 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

10 B is hydrogen, R¹ or -L¹-E;

E is phenyl optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

15 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

20 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

25 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

30 wherein ring D is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

35 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

5 L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;
D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,
10 wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein ring D is optionally substituted with one or more R³;
A is phenyl optionally substituted with one or more R¹; and
B is hydrogen or R¹.
15 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

20 L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;
D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,
25 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,
wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein ring D is optionally substituted with one or more R³;
30 A is naphthyl optionally substituted with one or more R¹; and
B is hydrogen or R¹.
In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

5 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

10 wherein ring D is optionally substituted with one or more R³;

A is phenyl; and

B is hydrogen.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

25 wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³;

A is naphthyl; and

30 B is hydrogen.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

5 D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2, preferably 1, heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

10 wherein ring D is optionally substituted with one or more R³;

A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹; and

15 B is -L¹-E.

In the above embodiment, preferably L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

20 In a more specific embodiment of the above embodiments, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

30 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹;

35 B is hydrogen, R¹ or -L¹-E;

E is phenyl optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

5 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

10 In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

15 wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or monocyclic heteroaryl (preferably phenyl, pyridyl or thiazolyl, more preferably phenyl, 3-pyridyl or 5-thiazolyl), wherein A is optionally substituted with one or more R¹; and

B is hydrogen or R¹.

20 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25 In another embodiment, the invention provides a compound of formula I, Ia,la-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

29 wherein said piperidinyl is optionally substituted with one or more R³;

30 A is phenyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

5 D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is naphthyl optionally substituted with one or more R¹; and

B is hydrogen or R¹.

10 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

15 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

20 wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl; and

B is hydrogen.

25 In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

30 L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is naphthyl; and

35 B is hydrogen.

In a more specific embodiment, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

5 In another embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) wherein:

L² is C₁₋₄ alkylene, preferably linear C₁₋₄ alkylene, more preferably linear C₁₋₂ alkylene;

D is piperidinyl, preferably 4-piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of

10 Formula I through a ring C atom, and

wherein said piperidinyl is optionally substituted with one or more R³;

A is phenyl or pyridyl (preferably phenyl or 3-pyridyl), wherein A is optionally substituted with one or more R¹;

and

B is -L¹-E.

15 In the above embodiment, preferably L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. In a more specific embodiment, E is phenyl optionally substituted with one or more R² and L¹ is a bond, -O-, -NH-, -CH₂-NH-, or -CH₂-O-, wherein said -CH₂-NH- or -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group. Preferably, L¹ is a bond or -CH₂-O-, wherein said -CH₂-O- group is linked to ring A through the O atom and to ring E through the -CH₂- group.

20 In a more specific embodiment of the above embodiments, said substituent(s) R³, if present, are placed on ring C atom(s) of ring D. In a further specific embodiment, each R³, if present, is independently selected from C₁₋₄ alkyl, halo, C₁₋₄ alkoxy, hydroxyl, amino, and amido.

25 In a further embodiment, the invention provides a compound of formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) selected from:

N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;

30 N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;

(Trans)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;

(Trans)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

35 N-((trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine;

(Trans)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;

(Trans)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;
N-((trans)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
5 (Trans)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(Trans)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(Trans)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(Trans)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
10 (Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(Trans)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
N-((trans)-2-phenylcyclopropyl)piperidin-3-amine;
15 N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-phenylcyclopropyl)azetidin-3-amine;
20 N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((trans)-2-phenylcyclopropyl)azepan-3-amine;
N-((trans)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((trans)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
25 N-((trans)-2-phenylcyclopropyl)decahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((trans)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine;
N-((trans)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine;
30 N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
35 N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
(1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
5 (1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine ;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
10 (1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
15 (1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
20 (1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
N-((*trans*)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(2-(*o*-tolyl)cyclopropyl)piperidin-4-amine;
N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
25 N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-(6-methoxy-4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
30 N-(4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(methylsulfonyl)-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-(4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((*trans*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
35 2,2,6,6-tetramethyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;

1-isopropyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;
5 N-((*trans*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
10 N-((*trans*)-2-(*o*-tolyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
(*Trans*)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine;
15 (*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(*Trans*)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
20 (*Trans*)-2-(4-(1*H*-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
30 (*trans*)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine; as well as salts and solvates thereof (including also pharmaceutically acceptable salts and solvates thereof).
25 Moreover, the invention provides the following compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1), in which L² is a bond:
N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
30 N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
35 N-((*trans*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-phenylcyclopropyl)piperidin-3-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-phenylcyclopropyl)pyrrolidin-3-amine;
5 N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-phenylcyclopropyl)azetidin-3-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
10 N-((trans)-2-phenylcyclopropyl)azepan-3-amine;
N-((trans)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((trans)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((trans)-2-phenylcyclopropyl)decahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine;
15 N-((trans)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((trans)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine;
N-((trans)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine;
N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine ;
20 N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
25 N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
30 N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-(6-methoxy-4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
35 1-(methylsulfonyl)-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;
1-(4-((trans)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;

4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((*trans*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
2,2,6,6-tetramethyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
5 1-isopropyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;
N-((*trans*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
10 N-((1*S*,2*S*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(*o*-tolyl)cyclopropyl)piperidin-4-amine;
15 N-((*trans*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine; as well as salts and solvates thereof (including
also pharmaceutically acceptable salts and solvates).

20 The invention further provides the following compounds of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound
of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of
formula Ia-1), in which L² is C₁₋₄ alkylene:
(*Trans*)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
25 (*Trans*)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
30 (*Trans*)-N-(piperidin-4-ylmethyl)-2-(3'-trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
35 (*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1*S*,2*R*)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1*R*,2*S*)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;

(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine ;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
5 (1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
10 (1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
15 (1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
20 (1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(Trans)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine;
(Trans)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
25 (Trans)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(Trans)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(Trans)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(Trans)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
30 (trans)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(trans)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine; as well as salts and solvates thereof (including also pharmaceutically acceptable salts and solvates).
In a further embodiment, the invention provides a compound of formula I selected from:
N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
35 N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine ;
(S)-N-((1S,2R)-2-phenylcyclopropyl)piperidin-3-amine;

(*R*)-N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-3-amine;
(*S*)-N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-3-amine;
(*R*)-N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-3-amine;
(*S*)-N-((1*S*,2*R*)-2-phenylcyclopropyl)pyrrolidin-3-amine;
5 (*R*)-N-((1*S*,2*R*)-2-phenylcyclopropyl)pyrrolidin-3-amine;
(*S*)-N-((1*R*,2*S*)-2-phenylcyclopropyl)pyrrolidin-3-amine;
(*R*)-N-((1*R*,2*S*)-2-phenylcyclopropyl)pyrrolidin-3-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)azetidin-3-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)azetidin-3-amine;
10 N-((1*S*,2*R*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((1*R*,2*S*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((1*R*,2*S*)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((1*S*,2*R*)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
15 N-((1*S*,2*R*)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
20 N-((1*S*,2*R*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
25 N-((1*R*,2*S*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(6-methoxy-4'-((1*R*,2*S*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(6-methoxy-4'-((1*S*,2*R*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
30 N-(4'-((1*R*,2*S*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
N-(4'-((1*S*,2*R*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(4-(((1*R*,2*S*)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
1-(4-(((1*S*,2*R*)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-(((1*R*,2*S*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
35 4-(((1*S*,2*R*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((1*R*,2*S*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2*H*-pyran-4-amine;

N-((1S,2R)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine;

2,2,6,6-tetramethyl-N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;

2,2,6,6-tetramethyl-N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;

1-methyl-N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;

5 1-methyl-N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;

1-isopropyl-N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;

1-isopropyl-N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;

N-((1S,2R)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;

10 4-(((1R,2S)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;

4-(((1S,2R)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;

N-((1S,2S)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2R)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;

15 N-((1S,2R)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;

N-((1S,2R)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;

N-((1S,2R)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;

20 N-((1R,2S)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;

N-((1S,2R)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;

N-((1S,2R)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;

as well as salts and solvates thereof (including also pharmaceutically acceptable salts and solvates).

25

The present invention furthermore provides a compound of formula I selected from:

(1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;

(1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;

(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

30 (1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

35 (1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
5 (1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
10 (1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1S,2R)-2-phenyl-N-((S)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1S,2R)-2-phenyl-N-((R)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-((S)-pyrrolidin-3-ylmethyl)cyclopropanamine;
15 (1R,2S)-2-phenyl-N-((R)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
20 (1S,2R)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(1R,2S)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
25 (1S,2R)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(1R,2S)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
as well as salts and solvates thereof (including also pharmaceutically acceptable salts and solvates).
30

The invention also relates to any one or any subgroup of the compounds listed above. The invention likewise relates to a pharmaceutically acceptable salt, preferably a hydrochloride salt (such as, e.g., a monohydrochloride salt, a dihydrochloride salt or, where applicable, a trihydrochloride salt), of any of the compounds listed above.

It is preferred that the compound of Formula I, Ia or Ia-1 is not 1-methyl-N-(2-phenylcyclopropyl)-4-piperidinamine or a salt or solvate thereof. Accordingly, in all of the embodiments described herein, the compound Formula I, Ia or Ia-1 for use as a medicament as well as for use in the treatment or prevention of cancer, a neurological disease or a viral infection is preferably not 1-methyl-N-(2-phenylcyclopropyl)-4-piperidinamine or a salt or solvate thereof.

Preferred embodiments of the compounds of Formula I, Ia, Ia-1, Ib and Ic for use in the compositions and methods of the invention are as defined herein above.

The present invention also provides compounds of Formula I, Ia or Ia-1, as defined and described herein above, including also compounds as defined in any of the preferred embodiments of Formula I, Ia or Ia-1 described herein, with the proviso that the following compounds are excluded:

1-(1-methylethyl)-N-(2-phenylcyclopropyl)- 3-Pyrrolidinamine;
4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid 1,1-dioxide;

15 N-(2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)- 4-Piperidinamine;

1-(3-methyl-2-buten-1-yl)-N-(2-phenylcyclopropyl)- 4-Piperidinamine;

4-[(2-phenylcyclopropyl)amino]-1-Piperidinecarboxylic acid ethyl ester;

1-[4-[(2-phenylcyclopropyl)amino]-1-piperidyl]- ethanone;

hexahydro-3-[(2-phenylcyclopropyl)amino]- 2H-azepin-2-one;

20 1-cyclopropyl-3-[(2-phenylcyclopropyl)amino]- 2,5-pyrrolidinedione;

3-[(2-phenylcyclopropyl)amino]-1-propyl-2,5-pyrrolidinedione;

1-(1-methylethyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;

1-(1-methylpropyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;

1,2,5-trimethyl-N-(2-phenylcyclopropyl)- 4-Piperidinamine;

25 3-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;

4-[(2-phenylcyclopropyl)amino]- 1-piperidinecarboxamide;

3-hydroxy-4-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;

tetrahydro-4-[(2-phenylcyclopropyl)amino]- 2H-pyran-4-carboxylic acid;

4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;

30 N-(2-phenylcyclopropyl)-1-(2-propyn-1-yl)- 4-piperidinamine;

1-ethyl-N-(2-phenylcyclopropyl)- 4-piperidinamine;

1-ethyl-3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;

4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid;

N-(2-phenylcyclopropyl)-1-propyl-4-piperidinamine;

35 3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;

tetrahydro-3-[(2-phenylcyclopropyl)amino]- 3-Thiophenecarboxylic acid;

tetrahydro-N-(2-phenylcyclopropyl)-2H-Thiopyran-4-amine 1-oxide;
1-(1-methylethyl)-N-(2-phenylcyclopropyl)-4-piperidinamine;
1-methyl-3-[(2-phenylcyclopropyl)amino]-2,5-pyrrolidinedione;
tetrahydro-3-[(2-phenylcyclopropyl)amino]-2H-thiopyran-3-carboxylic acid;
5 tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-amine;
N-(2-phenylcyclopropyl)-3-piperidinamine;
tetrahydro-4-[(2-phenylcyclopropyl)amino]-3-furanol;
N-(2-phenylcyclopropyl)-4-piperidinamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-thiopyran-4-amine;
10 tetrahydro-N-(2-phenylcyclopropyl)-3-thiophenamine;
1-methyl-N-(2-phenylcyclopropyl)-4-piperidinamine;
3-methoxy-N-[[5-[[4-[(1S,2R)-2-phenylcyclopropyl]amino]-1-piperidinyl]sulfonyl]-2-thienyl]methyl]-benzamide;
N-(2-phenylcyclopropyl)-1,4-dioxaspiro[4.5]decan-8-amine;
8-methyl-N-(2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
15 N-(2-phenylcyclopropyl)-1-azabicyclo[2.2.2]octan-3-amine;
6-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
8-fluoro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
5-amino-1,3-dihydro-6-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
20 3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
6-bromo-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
2,3-dihydro-3-[(2-phenylcyclopropyl)amino]-6-benzofuranol;
2,3-dihydro-N-(2-phenylcyclopropyl)-benzo[b]thiophen-3-amine 1,1-dioxide;
25 2,3-dihydro-N6-(2-phenylcyclopropyl)-1,4-benzodioxin-6,7-diamine;
3,4-dihydro-N-(2-phenylcyclopropyl)-1H-2-benzothiopyran-4-amine;
7-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-1-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-7-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
30 6-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
7-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
35 4-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
4-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;

1,3-dihydro-5-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
3-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-thieno[3,2-b]pyrrol-5-amine;
5 8-methoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-methyl-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-ethoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-fluoro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
10 8-chloro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione;
1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
15 a-methyl-N-(2-phenylcyclopropyl)-1-propyl-3-piperidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-2-methanamine;
octahydro-N-(2-phenylcyclopropyl)-1H-indole-2-methanamine;
a-methyl-N-(2-phenylcyclopropyl)-1-propyl-4-piperidinemethanamine;
3-methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
20 N-(2-phenylcyclopropyl)-2-piperidinemethanamine;
N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
a,1-dimethyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
4-methoxy-N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
25 N-(2-phenylcyclopropyl)-3-propyl-3-pyrrolidinemethanamine;
N-(2-phenylcyclopropyl)-2-morpholinemethanamine;
4-ethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
4-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
a,1-dimethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
30 1-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-methanamine;
3-methyl-N-(2-phenylcyclopropyl)-3-pyrrolidinemethanamine;
N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-3-furanmethanamine;
35 tetrahydro-N-(2-phenylcyclopropyl)-2-furanmethanamine;
N-(2-phenylcyclopropyl)-4-piperidineethanamine;

N-(2-phenylcyclopropyl)-2-pyrrolidineethanamine;
1-methyl-N-(2-phenylcyclopropyl)-2-piperidineethanamine; and
N-(2-phenylcyclopropyl)-3-piperidinepropanamine.

5 Accordingly, the invention relates to a compound of Formula I wherein:
A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;
B is H, R¹ or -L¹-E;
E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;
L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;
10 L² is a bond and D is a cyclic group selected from:
(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,
15 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,
20 wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;
or L² is C₁₋₄ alkylene and D is a cyclic group selected from:
25 (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,
30 wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,
wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
35 wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;
or a salt or solvate thereof;

with the proviso that the following compounds are excluded:

1-(1-methylethyl)-N-(2-phenylcyclopropyl)- 3-Pyrrolidinamine;

4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid 1,1-dioxide;

15 N-(2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)- 4-Piperidinamine;

1-(3-methyl-2-buten-1-yl)-N-(2-phenylcyclopropyl)- 4-Piperidinamine;

4-[(2-phenylcyclopropyl)amino]-1-Piperidinecarboxylic acid ethyl ester;

1-[4-[(2-phenylcyclopropyl)amino]-1-piperidinyl]- ethanone;

hexahydro-3-[(2-phenylcyclopropyl)amino]- 2H-azepin-2-one;

20 1-cyclopropyl-3-[(2-phenylcyclopropyl)amino]- 2,5-pyrrolidinedione;

3-[(2-phenylcyclopropyl)amino]-1-propyl-2,5-pyrrolidinedione;

1-(1-methylethyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;

1-(1-methylpropyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;

1,2,5-trimethyl-N-(2-phenylcyclopropyl)- 4-Piperidinamine;

25 3-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;

4-[(2-phenylcyclopropyl)amino]- 1-piperidinecarboxamide;

3-hydroxy-4-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;

tetrahydro-4-[(2-phenylcyclopropyl)amino]- 2H-pyran-4-carboxylic acid;

4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;

30 N-(2-phenylcyclopropyl)-1-(2-propyn-1-yl)- 4-piperidinamine;

1-ethyl-N-(2-phenylcyclopropyl)- 4-piperidinamine;

1-ethyl-3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;

4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid;

N-(2-phenylcyclopropyl)-1-propyl-4-piperidinamine;

35 3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;

tetrahydro-3-[(2-phenylcyclopropyl)amino]- 3-Thiophenecarboxylic acid;

tetrahydro-N-(2-phenylcyclopropyl)- 2H-Thiopyran-4-amine 1-oxide;
1-(1-methylethyl)-N-(2-phenylcyclopropyl)-4-piperidinamine;
1-methyl-3-[(2-phenylcyclopropyl)amino]-2,5-pyrrolidinedione;
tetrahydro-3-[(2-phenylcyclopropyl)amino]-2H-thiopyran-3-carboxylic acid;
5 tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-amine;
N-(2-phenylcyclopropyl)-3-piperidinamine;
tetrahydro-4-[(2-phenylcyclopropyl)amino]-3-furanol;
N-(2-phenylcyclopropyl)-4-piperidinamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-thiopyran-4-amine;
10 tetrahydro-N-(2-phenylcyclopropyl)-3-thiophenamine;
1-methyl-N-(2-phenylcyclopropyl)-4-piperidinamine;
3-methoxy-N-[[5-[[4-[(1S,2R)-2-phenylcyclopropyl]amino]-1-piperidinyl]sulfonyl]-2-thienyl]methyl]-benzamide;
N-(2-phenylcyclopropyl)-1,4-dioxaspiro[4.5]decan-8-amine;
8-methyl-N-(2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
15 N-(2-phenylcyclopropyl)-1-azabicyclo[2.2.2]octan-3-amine;
6-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
8-fluoro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
5-amino-1,3-dihydro-6-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
20 3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
6-bromo-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
2,3-dihydro-3-[(2-phenylcyclopropyl)amino]-6-benzofuranol;
2,3-dihydro-N-(2-phenylcyclopropyl)-benzo[b]thiophen-3-amine 1,1-dioxide;
25 2,3-dihydro-N6-(2-phenylcyclopropyl)-1,4-benzodioxin-6,7-diamine;
3,4-dihydro-N-(2-phenylcyclopropyl)-1H-2-benzothiopyran-4-amine;
7-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-1-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-7-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
30 6-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
7-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
35 4-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
4-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;

1,3-dihydro-5-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
3-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-thieno[3,2-b]pyrrol-5-amine;
5 8-methoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-methyl-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-ethoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-fluoro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
10 8-chloro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione;
1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
15 a-methyl-N-(2-phenylcyclopropyl)-1-propyl-3-piperidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-2-methanamine;
octahydro-N-(2-phenylcyclopropyl)-1H-indole-2-methanamine;
a-methyl-N-(2-phenylcyclopropyl)-1-propyl-4-piperidinemethanamine;
3-methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
20 N-(2-phenylcyclopropyl)-2-piperidinemethanamine;
N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
a,1-dimethyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
4-methoxy-N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
25 N-(2-phenylcyclopropyl)-3-propyl-3-pyrrolidinemethanamine;
N-(2-phenylcyclopropyl)-2-morpholinemethanamine;
4-ethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
4-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
a,1-dimethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
30 1-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-methanamine;
3-methyl-N-(2-phenylcyclopropyl)-3-pyrrolidinemethanamine;
N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
tetrahydro-N-(2-phenylcyclopropyl)-3-furanmethanamine;
35 tetrahydro-N-(2-phenylcyclopropyl)-2-furanmethanamine;
N-(2-phenylcyclopropyl)-4-piperidineethanamine;

N-(2-phenylcyclopropyl)-2-pyrrolidineethanamine;
1-methyl-N-(2-phenylcyclopropyl)-2-piperidineethanamine; and
N-(2-phenylcyclopropyl)-3-piperidinopropanamine.

5 Accordingly, the invention relates to a compound of Formula I, Ia or Ia-1, as defined and described herein above, including also compounds as defined in any of the preferred embodiments of Formula I, Ia or Ia-1 described herein, with the proviso that the following compounds are excluded:

3-Pyrrolidinamine, 1-(1-methylethyl)-N-(2-phenylcyclopropyl)-;
2H-Thiopyran-4-carboxylic acid, tetrahydro-4-[(2-phenylcyclopropyl)amino]-, 1,1-dioxide;

10 4-Piperidinamine, N-(2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)-;
4-Piperidinamine, 1-(3-methyl-2-buten-1-yl)-N-(2-phenylcyclopropyl)-;
1-Piperidinecarboxylic acid, 4-[(2-phenylcyclopropyl)amino]-, ethyl ester;
Ethanone, 1-[4-[(2-phenylcyclopropyl)amino]-1-piperidinyl]-;
2H-Azepin-2-one, hexahydro-3-[(2-phenylcyclopropyl)amino]-;

15 2,5-Pyrrolidinedione, 1-cyclopropyl-3-[(2-phenylcyclopropyl)amino]-;
2,5-Pyrrolidinedione, 3-[(2-phenylcyclopropyl)amino]-1-propyl-;
2,5-Pyrrolidinedione, 1-(1-methylethyl)-3-[(2-phenylcyclopropyl)amino]-;
2,5-Pyrrolidinedione, 1-(1-methylpropyl)-3-[(2-phenylcyclopropyl)amino]-;
4-Piperidinamine, 1,2,5-trimethyl-N-(2-phenylcyclopropyl)-;

20 3-Thiophenamine, tetrahydro-N-(2-phenylcyclopropyl)-, 1,1-dioxide;
1-Piperidinecarboxamide, 4-[(2-phenylcyclopropyl)amino]-;
Thiophene-3-ol, tetrahydro-4-[(2-phenylcyclopropyl)amino]-, 1,1-dioxide; 2H-Pyran-4-carboxylic acid, tetrahydro-4-[(2-phenylcyclopropyl)amino]-;
2H-Thiopyran-4-amine, tetrahydro-N-(2-phenylcyclopropyl)-, 1,1-dioxide;

25 4-Piperidinamine, N-(2-phenylcyclopropyl)-1-(2-propyn-1-yl)-;
4-Piperidinamine, 1-ethyl-N-(2-phenylcyclopropyl)-;
2,5-Pyrrolidinedione, 1-ethyl-3-[(2-phenylcyclopropyl)amino]-;
2H-Thiopyran-4-carboxylic acid, tetrahydro-4-[(2-phenylcyclopropyl)amino]-;
4-Piperidinamine, N-(2-phenylcyclopropyl)-1-propyl-;

30 2,5-Pyrrolidinedione, 3-[(2-phenylcyclopropyl)amino]-;
3-Thiophenecarboxylic acid, tetrahydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Thiopyran-4-amine, tetrahydro-N-(2-phenylcyclopropyl)-, 1-oxide;
4-Piperidinamine, 1-(1-methylethyl)-N-(2-phenylcyclopropyl)-;

35 2,5-Pyrrolidinedione, 1-methyl-3-[(2-phenylcyclopropyl)amino]-;
2H-Thiopyran-3-carboxylic acid, tetrahydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Pyran-4-amine, tetrahydro-N-(2-phenylcyclopropyl)-;

3-Piperidinamine, N-(2-phenylcyclopropyl)-;
3-Furanol, tetrahydro-4-[(2-phenylcyclopropyl)amino]-;
4-Piperidinamine, N-(2-phenylcyclopropyl)-;
2H-Thiopyran-4-amine, tetrahydro-N-(2-phenylcyclopropyl)-;
5 3-Thiophenamine, tetrahydro-N-(2-phenylcyclopropyl)-;
4-Piperidinamine, 1-methyl-N-(2-phenylcyclopropyl)-;
Benzamide, 3-methoxy-N-[[5-[[4-[(1S,2R)-2-phenylcyclopropyl]amino]-1-piperidinyl]sulfonyl]-2-thienyl]methyl]-;
1,4-Dioxaspiro[4.5]decan-8-amine, N-(2-phenylcyclopropyl)-;
8-Azabicyclo[3.2.1]octan-3-amine, 8-methyl-N-(2-phenylcyclopropyl)-;
10 1-Azabicyclo[2.2.2]octan-3-amine, N-(2-phenylcyclopropyl)-;
2H-1-Benzothiopyran-4-amine, 6-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-;
2H-1-Benzothiopyran-4-amine, 8-fluoro-3,4-dihydro-N-(2-phenylcyclopropyl)-;
2H-1-Benzothiopyran-4-amine, 3,4-dihydro-N-(2-phenylcyclopropyl)-;
2H-Indol-2-one, 5-amino-1,3-dihydro-6-[(2-phenylcyclopropyl)amino]-;
15 2H-1-Benzopyran-4-amine, 3,4-dihydro-N-(2-phenylcyclopropyl)-;
2H-1-Benzopyran-4-amine, 6-bromo-3,4-dihydro-N-(2-phenylcyclopropyl)-;
-Benzofuranol, 2,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
Benzo[b]thiophen-3-amine, 2,3-dihydro-N-(2-phenylcyclopropyl)-, 1,1-dioxide;
1,4-Benzodioxin-6,7-diamine, 2,3-dihydro-N6-(2-phenylcyclopropyl)-;
20 1H-2-Benzothiopyran-4-amine, 3,4-dihydro-N-(2-phenylcyclopropyl)-;
2H-Indol-2-one, 7-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 1,3-dihydro-1-methyl-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 1,3-dihydro-7-methyl-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 6-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
25 2H-Indol-2-one, 5-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 7-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 6-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 4-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
30 2H-Indol-2-one, 4-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 5-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 1,3-dihydro-5-methyl-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 5-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
2H-Indol-2-one, 6-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-;
35 2H-Thieno[3,2-b]pyrrol-5-amine, 3-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-;
5H-Pyrimido[5,4-b]indol-4-amine, 8-methoxy-N-(2-phenylcyclopropyl)-;

5H-Pyrimido[5,4-b]indol-4-amine, N-(2-phenylcyclopropyl)-;
5H-Pyrimido[5,4-b]indol-4-amine, 8-methyl-N-(2-phenylcyclopropyl)-;
5H-Pyrimido[5,4-b]indol-4-amine, 8-ethoxy-N-(2-phenylcyclopropyl)-;
5H-Pyrimido[5,4-b]indol-4-amine, 8-fluoro-N-(2-phenylcyclopropyl)-;
5H-Pyrimido[5,4-b]indol-4-amine, 8-chloro-N-(2-phenylcyclopropyl)-;
5 Benzo[g]pteridine-2,4(3H,4aH)-dione, 5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-;
3-Piperidinemethanamine, 1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, 1-ethyl-a-methyl-N-(2-phenylcyclopropyl)-;
10 3-Piperidinemethanamine, a-methyl-N-(2-phenylcyclopropyl)-1-propyl-;
2H-Pyran-2-methanamine, tetrahydro-N-(2-phenylcyclopropyl)-;
1H-Indole-2-methanamine, octahydro-N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, a-methyl-N-(2-phenylcyclopropyl)-1-propyl-;
3-Piperidinemethanamine, 3-methyl-N-(2-phenylcyclopropyl)-;
15 2-Piperidinemethanamine, N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, N-(2-phenylcyclopropyl)-;
3-Piperidinemethanamine, a,1-dimethyl-N-(2-phenylcyclopropyl)-;
3-Piperidinemethanamine, N-(2-phenylcyclopropyl)-;
2-Pyrrolidinemethanamine, 4-methoxy-N-(2-phenylcyclopropyl)-;
20 3-Pyrrolidinemethanamine, N-(2-phenylcyclopropyl)-3-propyl-;
2-Morpholinemethanamine, N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, 4-ethyl-N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, 4-methyl-N-(2-phenylcyclopropyl)-;
4-Piperidinemethanamine, a,1-dimethyl-N-(2-phenylcyclopropyl)-;
25 4-Piperidinemethanamine, 1-methyl-N-(2-phenylcyclopropyl)-;
2H-Pyran-4-methanamine, tetrahydro-N-(2-phenylcyclopropyl)-;
3-Pyrrolidinemethanamine, 3-methyl-N-(2-phenylcyclopropyl)-;
2-Pyrrolidinemethanamine, N-(2-phenylcyclopropyl)-;
3-Furanmethanamine, tetrahydro-N-(2-phenylcyclopropyl)-;
30 2-Furanmethanamine, tetrahydro-N-(2-phenylcyclopropyl)-;
4-Piperidineethanamine, N-(2-phenylcyclopropyl)-;
2-Pyrrolidineethanamine, N-(2-phenylcyclopropyl)-;
2-Piperidineethanamine, 1-methyl-N-(2-phenylcyclopropyl)-; and
3-Piperidinepropanamine, N-(2-phenylcyclopropyl)-.

The present invention further provides compounds of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1), as defined and described herein above, including also compounds as defined in any of the preferred embodiments of Formula I, Ia, Ia-1, Ib or Ic described herein, in which (i) A is aryl optionally substituted with one or more R¹ and B is -L¹-E, or (ii) A is heteroaryl optionally substituted with one or more R¹ and B is H, R¹ or -L¹-E.

Accordingly, the invention provides a compound of Formula I wherein:

A is aryl optionally substituted with one or more R¹, and B is -L¹-E, or

A is heteroaryl optionally substituted with one or more R¹, and B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl; or a salt or solvate thereof.

The present invention furthermore provides compounds of Formula I, Ia, Ib or Ic (preferably a compound of formula I or Ia, more preferably a compound of formula Ia), as defined and described herein above, including also compounds as defined in any of the preferred embodiments of Formula I, Ia, Ia-1, Ib or Ic described herein, in which the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration. For these compounds, it is preferred that the compound 5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione is excluded.

Accordingly, the invention provides a compound of Formula I wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

25 B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

5 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

10 (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

15 wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, 20 halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

25 each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;

or a salt or solvate thereof;

30 wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration,

and further wherein it is preferred that the compound 5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione is excluded.

Accordingly, for the compounds of Formula I, **Ia** and **Ia-1**, in which the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration, it is preferred that the compound 5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(trans-2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione is excluded.

5 The present invention also provides compounds of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1), as defined and described herein above, including also compounds as defined in any of the preferred embodiments of Formula I, Ia, Ia-1, Ib or Ic described herein, in which the compound of formula I, Ia, Ia-1, Ib or Ic is an optically active stereoisomer.

Accordingly, the invention provides a compound of Formula I wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

10 B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

15 (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

20 wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

25 wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

30 (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl; or a salt or solvate thereof;

wherein the compound of formula I is an optically active stereoisomer.

The invention relates to compounds which are cyclopropylamino derivatives, so that cyclopropylimino compounds are not covered. Accordingly, compounds such as, e.g., (S,Z)-N-(2,3-dichlorophenyl)-3-oxo-1-(((1S,2R)-2-phenylcyclopropyl)imino)tetrahydro-1H-oxazolo[3,4-a]pyrazine-7(3H)-carboxamide, (S,Z)-tert-butyl 3-oxo-1-(((1S,2R)-2-phenylcyclopropyl)imino)tetrahydro-1H-oxazolo[3,4-a]pyrazine-7(3H)-carboxylate or 3H-Oxazolo[3,4-a]pyrazin-3-one, hexahydro-1-[(1S,2R)-2-phenylcyclopropyl]imino]-, (8aS)- are not covered by the definition of a compound of formula I, Ia, Ia-1, Ib or Ic.

In a further aspect, the invention provides a method for identifying a compound which is a selective inhibitor of LSD1, the method comprising selecting or providing a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) and determining the ability of the said compound to inhibit LSD1 and MAO-A and/or MAO-B using assays such as the ones disclosed in more detail later on, wherein a compound that inhibits LSD1 to a greater extent than MAO-A and/or MAO-B is identified as a LSD1 selective inhibitor. LSD1 selective inhibitors have IC₅₀ values for LSD1 which are lower than the IC₅₀ value for MAO-A and/or MAO-B. Preferably, the IC₅₀ values for LSD1 are two-fold lower than for MAO-A and/or MAO-B. In one aspect of this embodiment, the LSD1 IC₅₀ value is at least 5-fold lower than the IC₅₀ value for MAO-A and/or MAO-B. In one aspect of this embodiment, the LSD1 IC₅₀ value is at least 10-fold lower than the IC₅₀ value for MAO-A and/or MAO-B. Preferably, a selective LSD1 inhibitor exhibits an IC₅₀ value for LSD1 that is > 50-fold, preferably >100-fold lower than the IC₅₀ value for MAO-A and/or MAO-B.

Asymmetric centers exist in the compounds of formula I, Ia, Ia-1, Ib and Ic disclosed herein. It should be understood that the invention encompasses all individual stereochemical isomeric forms of a compound of formula I, Ia, Ia-1, Ib and Ic, including diastereomeric, enantiomeric, and epimeric forms, as well as d-isomers and l-isomers ((+)-isomers and (-)-isomers), and any mixtures thereof, including wholly or partially equilibrated mixtures. Individual stereoisomers of compounds of the invention can be prepared synthetically from commercially available chiral starting materials or by separation from mixtures of stereoisomers, as also shown in the Examples. Methods of separation of enantiomeric and diastereomeric mixtures are well known to one skilled in the art. For example, mixtures of diastereomers can be separated by conventional separation techniques such as recrystallization or chromatography. Mixtures of enantiomeric products can be separated by conversion to a mixture of diastereomers followed by separation using recrystallization or chromatographic techniques, direct separation of enantiomers on chiral chromatographic columns, or any other appropriate method of chiral resolution known in the art. Starting compounds of particular stereochemistry are either commercially available or can be made and resolved by techniques known in the art.

15 Additionally, the compounds disclosed herein may exist as geometric isomers. The present invention includes all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the mixtures thereof.

20 Additionally, compounds may exist as tautomers; all tautomeric isomers are provided by this invention.

25 The compounds of the invention contain one or more basic nitrogen atoms and may therefore form salts with organic or inorganic acids. The compounds of the invention may also contain one or more acidic protons and therefore they may also form salts with bases. There is no limitation on the type of salt that can be used provided that these are pharmaceutically acceptable when used for therapeutic purposes. The salts of a compound of the invention can be obtained during the final isolation and purification of the compounds of the invention or can be obtained by treating a compound of formula I, Ia, Ia-1, Ib or Ic with a sufficient amount of the desired acid or base to give the corresponding salt in a conventional manner. All salts of the compounds of formula I, Ia, Ia-1, Ib or Ic, including pharmaceutically acceptable salts, are included within the scope of the invention. In one embodiment, a compound of formula I, Ia, Ia-1, Ib or Ic is provided in the form of a salt. In a more preferred embodiment, a compound of formula I, Ia, Ia-1, Ib or Ic is provided in the form of a pharmaceutically acceptable salt. In one embodiment, such pharmaceutically acceptable salt is a hydrochloride, for example a monohydrochloride, a dihydrochloride or a trihydrochloride.

30 Additionally, the compounds disclosed herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. Solvated forms with water are known as hydrates. In general, the solvated forms are considered equivalent to the unsolvated forms. The

invention thus relates to the unsolvated and solvated forms of the compounds of formula I, Ia, Ia-1, Ib or Ic (or of any salt thereof).

The compounds of formula I, Ia, Ia-1, Ib and Ic may exist in different physical forms, i.e. amorphous and crystalline forms. Moreover, the compounds of the invention may have the ability to crystallize in more than one form, a characteristic which is known as polymorphism. Polymorphs can be distinguished by various physical properties well known in the art such as X-ray diffraction pattern, melting point or solubility. All physical forms of the compounds of the invention, including all polymorphic forms (also known as polymorphs) thereof, are included within the scope of the invention.

10 The present invention further covers all unlabeled and isotopically labeled forms of the compounds of formula I, Ia, Ia-1, Ib and Ic. In one embodiment, the invention relates to deuterated forms of the compounds of formula I, Ia and Ia-1.

15 The invention also relates to a compound of Formula I, Ia, Ia-1, Ib or Ic (preferably a compound of formula I, Ia, or Ia-1, more preferably a compound of formula Ia or Ia-1, and most preferably a compound of formula Ia-1) as described and defined herein, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration and further wherein the compound is optically active. As used herein, the term "optically active," refers to the ability of a compound to rotate plane polarized light.

20 The invention, in another aspect, relates to a substantially pure, optically active stereoisomer of a compound of Formula I, Ia, Ia-1, Ib or Ic as described and defined herein, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration, or a pharmaceutically acceptable salt or solvate thereof, as well as its use as a medicament. As used herein, "substantially pure" means that there is 90 mole-% or greater of the desired stereoisomer and 10 mole-% or less of any other stereoisomer, preferably that there is 95 mole-% or greater of the desired stereoisomer and 5 mole-% or less of any other stereoisomer, more preferably, that there is 98 mole-% or greater of the desired stereoisomer and 2 mole-% or less of any other stereoisomer, still more preferably, that there is 99 mole-% or greater of the desired stereoisomer and 1 mole-% or less of any other stereoisomer, and even more preferably that there is 99.5 mole-% or greater of the desired stereoisomer and 0.5 mole-% or less of any other stereoisomer. The substantially pure, optically active stereoisomer of a compound of Formula I, Ia, Ia-1, Ib or Ic as described and defined herein, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration, is useful in treating or preventing a disease or disorder, particularly cancer, a neurological disease, or a viral infection.

Definitions:

Any definition herein may be used in combination with any other definition to describe a composite structural group. By convention, the trailing element of any such definition is that which attaches to the parent moiety.

5

As used herein, the term "acyl" refers to a carbonyl attached to an alkenyl, alkyl, aryl, cycloalkyl, heteroaryl, heterocyclyl, or any other moiety where the atom attached to the carbonyl is carbon. Preferably, the term "acyl" refers to a group of formula $-C(=O)R'$, wherein R' represents alkenyl, alkyl, aryl, cycloalkyl, heteroaryl or heterocyclyl. An "acetyl" group refers to a $-C(=O)CH_3$ group. An "alkylcarbonyl" or "alkanoyl" group refers to an alkyl group attached to the parent molecular moiety through a carbonyl group. Examples of such groups include, but are not limited to, methylcarbonyl or ethylcarbonyl. Examples of acyl groups include, but are not limited to, formyl, alkanoyl or aroyl.

10

As used herein, the term "alkenyl" refers to a straight-chain or branched-chain hydrocarbon group having one or more double bonds and containing from 2 to 20 carbon atoms. A C_{2-8} alkenyl is an alkenyl group having from 2 to 8 carbon atoms.

15

As used herein, the term "alkoxy" refers to an alkyl ether group (i.e. a group of formula $alkyl-O-$), wherein the term alkyl is as defined below. Examples of suitable alkyl ether groups include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, or n-pentoxy. The term C_{1-z} alkoxy refers to an alkoxy group wherein the alkyl moiety has from 1 to z carbon atoms; for example a C_{1-8} alkoxy is an alkoxy group wherein the alkyl moiety is C_{1-8} alkyl, i.e. a group of formula $C_{1-8} alkyl-O-$.

20

As used herein, the term "alkyl" refers to a straight-chain or branched-chain alkyl group containing from 1 to 20 carbon atoms. A C_{1-z} alkyl is an alkyl from 1 to z carbon atoms; thus, a C_{1-8} alkyl has from 1 to 8 carbon atoms, a C_{1-4} alkyl has from 1 to 4 carbon atoms and a C_{1-2} alkyl has from 1 to 2 carbon atoms. Examples of alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neo-pentyl, iso-amyl, hexyl, heptyl, octyl, or nonyl.

25

As used herein, the term " C_{1-4} alkylene" refers to an C_{1-4} alkyl group attached at two positions, i.e. an alkanediyl group. Examples include, but are not limited to, methylene, ethylene, propylene and butylene. Accordingly, the term " C_{1-4} alkylene" may refer to a straight-chain or branched-chain alkylene group having from 1 to 4 carbon atoms. A "linear C_{1-4} alkylene" refers to a straight chain alkylene group having from 1 to 4 carbon atoms, i.e. a $-(CH_2)_y-$ group wherein y is 1, 2, 3 or 4.

30

35

As used herein, the term "alkynyl" refers to a straight-chain or branched-chain hydrocarbon group having one or more triple bonds and containing from 2 to 20 carbon atoms. A C₂₋₈ alkynyl has from 2 to 8 carbon atoms. Examples of alkynyl groups include, but are not limited to, ethynyl, propynyl, hydroxypropynyl, butyn-1-yl, butyn-2-yl, pentyn-1-yl, 3-methylbutyn-1-yl, or hexyn-2-yl.

5

As used herein, the term "amido" refers to an amino group as described below attached to the parent molecular moiety through a carbonyl group (e.g., -C(=O)NRR') , or vice versa (-N(R)C(=O)R'). "Amido" encompasses "C-amido" and "N-amido" as defined herein. R and R' are as defined herein.

10

As used herein, the term "C-amido" refers to a -C(=O)NRR' group with R and R' as defined herein.

As used herein, the term "N-amido" refers to a -N(R)C(=O)R' group with R and R' as defined herein.

As used herein, the term "amino" refers to -NRR', wherein R and R' are independently selected from the group 15 consisting of hydrogen, alkyl, heteroalkyl, aryl, carbocyclyl, and heterocyclyl. Additionally, R and R' may be combined to form a heterocyclyl. Exemplary "amino" groups include, without being limited thereto, -NH₂, -NH(C₁₋₄ alkyl) and -N(C₁₋₄ alkyl)(C₁₋₄ alkyl).

As used herein, the term "aryl" refers to a carbocyclic aromatic system containing one ring, or two or three rings 20 fused together where in the ring atoms are all carbon. The term "aryl" groups includes, but is not limited to groups such as phenyl, naphthyl, or anthracenyl. The term "monocyclic aryl" refers to phenyl.

As used herein, the term "aryloxy" refers to an aryl group attached to the parent molecular moiety through an oxy (-O-).

25

As used herein, the term "carbamate" refers to an O-carbamyl or N-carbamyl group as defined herein. An N-carbamyl group refers to -NR-COOR', wherein R and R' are as defined herein. An O-carbamyl group refers to -OCO-NRR', wherein R and R' are as defined herein.

As used herein, the term "carbonyl" when alone includes formyl -C(=O)H and in combination is a -C(=O)- group.

As used herein, the term "carboxyl" or "carboxy" refers to -C(=O)OH or the corresponding "carboxylate" anion, such as is in a carboxylic acid salt.

35

An "O-carboxy" group refers to a RC(=O)O- group, where R is as defined herein.

A "C-carboxy" group refers to a -C(=O)OR groups where R is as defined herein.

As used herein, the term "cyano" refers to -CN.

5

As used herein, the term "carbocycll" refers to a saturated or partially saturated monocyclic or a fused bicyclic or tricyclic group wherein the ring atoms of the cyclic system are all carbon and wherein each cyclic moiety contains from 3 to 12 carbon atom ring members. "Carbocycll" encompasses benzo fused to a carbocycll ring system. One group of carbocyclls have from 5 to 7 carbon atoms. Examples of carbocycll groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, indanyl, 10 octahydronaphthyl, 2,3-dihydro-1H-indenyl, or adamantly.

10

As used herein, the term "cycloalkyl" refers to a saturated monocyclic, bicyclic or tricyclic group wherein the ring atoms of the cyclic system are all carbon and wherein each cyclic moiety contains from 3 to 12 carbon atom ring members. A C₃₋₆ cycloalkyl is a cycloalkyl that has from 3 to 6 carbon atoms, i.e. cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, or adamantly.

15

As used herein, the term "cycll" refers to an aryl, heterocycll, or carbocycll group as defined herein.

20

As used herein, the term "halo" or "halogen" refers to fluorine, chlorine, bromine, or iodine.

25

As used herein, the term "haloalkoxy" refers to a haloalkyl group (as defined below) attached to the parent molecular moiety through an oxygen atom. A haloC₁₋₈ alkoxy group refers to a haloalkoxy group wherein the haloalkyl moiety has from 1 to 8 C atoms. Examples of haloalkoxy groups include, but are not limited to, trifluoromethoxy, 2-fluoroethoxy, pentafluoroethoxy, or 3-chloropropoxy.

30

As used herein, the term "haloalkyl" refers to an alkyl group having the meaning as defined above wherein one or more hydrogens are replaced with a halogen. A haloC₁₋₈ alkyl group refers to a haloalkyl group wherein the alkyl moiety has from 1 to 8 C atoms. Specifically embraced are monohaloalkyl, dihaloalkyl or polyhaloalkyl groups. A monohaloalkyl group, for one example, may have an iodo, bromo, chloro or fluoro atom within the group. Dihalo or polyhaloalkyl groups may have two or more of the same halo atoms or a combination of different halo groups. Examples of haloalkyl groups include, but are not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, 35 difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl or dichloropropyl.

As used herein, the term "heteroalkyl" refers to a straight or branched alkyl chain, wherein one, two, or three carbons forming the alkyl chain are each replaced by a heteroatom independently selected from the group consisting of O, N, and S, and wherein the nitrogen and/or sulfur heteroatom(s) (if present) may optionally be oxidized and the nitrogen heteroatom(s) (if present) may optionally be quaternized. The heteroatom(s) O, N and S may, for example, be placed at the end(s) or at an interior position of the heteroalkyl group, i.e., the heteroalkyl may be bound to the remainder of the molecule via a heteroatom or a carbon atom. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃. Accordingly, a further example for a "heteroalkyl" group is a straight or branched alkyl group, in which two consecutive carbon atoms are replaced by the heteroatoms S and N, respectively, and the sulfur heteroatom is furthermore oxidized, resulting in 10 moieties such as, e.g., -S(=O)₂-NH₂, -S(=O)₂-NH(alkyl) or -S(=O)₂-N(alkyl)(alkyl).

As used herein, the term "heteroC₁₋₄ alkylene" refers to a straight or branched C₁₋₄ alkylene group (i.e., a straight or branched C₁₋₄ alkanediyl group) linked to one heteroatom selected from O, N and S and also refers to a straight or branched C₁₋₄ alkylene group wherein one or more (e.g., 1, 2 (if present) or 3 (if present)) of the 15 carbon atoms of said alkylene group are each replaced by a heteroatom independently selected from O, N or S. The nitrogen and/or sulfur heteroatom(s) (if present) may optionally be oxidized and the nitrogen heteroatom(s) (if present) may optionally be quaternized. The heteroatom(s) O, N and S may be placed at the end(s) and/or at an interior position of the heteroC₁₋₄ alkylene group. It is to be understood that the presence of 20 hydrogen atoms will depend on the valence of the heteroatom replacing the respective carbon atom. If, for example, the carbon atom in a -CH₂- group is replaced by O or S, the resulting group will be -O- or -S-, respectively, while it will be -N(H)- when the carbon atom is replaced by N. Likewise, if the central carbon atom in a group -CH₂-CH(-CH₃)-CH₂- is replaced by N, the resulting group will be -CH₂-N(-CH₃)-CH₂- An example for 25 a "heteroC₁₋₄alkylene" group is a straight or branched C₁₋₄ alkylene group, in which two consecutive carbon atoms are replaced by the heteroatoms S and N, respectively, and the sulfur heteroatom is furthermore oxidized, resulting in moieties such as, e.g., -S(=O)₂-N(H)- or -S(=O)₂-N(CH₃)-.

As used herein, the term "heteroaryl" refers to a 5 to 6 membered unsaturated monocyclic ring, or a fused 30 bicyclic or tricyclic ring system in which the rings are aromatic and in which at least one ring contains at least one heteroatom selected from the group consisting of O, S, and N. Preferred heteroaryl groups are 5- to 6-membered monocyclic or 9- to 10-membered bicyclic heteroaryl groups. Examples of heteroaryl groups include, but are not limited to, pyridinyl, imidazolyl, imidazopyridinyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, 35 tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxadiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indolizinyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, or furopyridinyl.

As used herein, the term "heterocyclyl" or "heterocycle" or "heterocyclic ring" each refer to a saturated, partially unsaturated, or fully unsaturated monocyclic, bicyclic, or tricyclic heterocyclic group containing at least one heteroatom as a ring member, wherein each of said at least one heteroatoms may be independently selected from the group consisting of nitrogen, oxygen, and sulfur wherein the nitrogen or sulfur atoms may be oxidized (e.g., $-N=O$, $-S(=O)$ -, or $-S(=O)_2$ -). Additionally, 1, 2, or 3 of the carbon atoms of the heterocyclyl may be optionally oxidized (e.g., to give an oxo group or $=O$). One group of heterocyclyls has from 1 to 4 heteroatoms as ring members. Another group of heterocyclyls has from 1 to 2 heteroatoms as ring members. One group of heterocyclyls has from 3 to 8 ring members in each ring. Yet another group of heterocyclyls has from 3 to 7 ring members in each ring. Again another group of heterocyclyls has from 5 to 6 ring members in each ring. "Heterocyclyl" is intended to encompass a heterocyclyl group fused to a carbocyclyl or benzo ring systems. Examples of heterocyclyl groups include, but are not limited to, pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, homopiperazinyl, azetidinyl, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 2-pyrrolinyl, 3-pyrrolinyl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolinyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinylimidazolinyl, or imidazolidinyl. Examples of heteroaryls that are heterocyclyls include, but are not limited to, pyridinyl, imidazolyl, imidazopyridinyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxadiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indolizinyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, or furopyridinyl.

As used herein, the term "heterocycloalkyl" refers to a heterocyclyl group that is not fully unsaturated e.g., one or more of the rings systems of a heterocycloalkyl is not aromatic. Examples of heterocycloalkyls include piperazinyl, morpholinyl, piperidinyl, or pyrrolidinyl.

As used herein, the term "hydroxyl" or "hydroxy" refers to $-OH$.

As used herein, the term "lower" where not otherwise specifically defined, means containing from 1 to and including 6 carbon atoms.

As used herein, the term "nitro" refers to $-NO_2$.

As used herein, the term "saturated" in relation to a ring means that the ring does not contain any unsaturation.

As used herein, the terms "sulfonate" "sulfonic acid" and "sulfonic" refer to the $-\text{SO}_3\text{H}$ group and its anion as the sulfonic acid is used in salt formation.

As used herein, the term "sulfinyl" refers to $-\text{S}(=\text{O})(\text{R})$, with R as defined herein.

5

As used herein, the term "sulfonyl" refers to $-\text{S}(=\text{O})_2\text{R}$, with R as defined herein.

As used herein, the term "sulfonamide" refers to an N-sulfonamido or S-sulfonamido group as defined herein.

10 As used herein, the term "N-sulfonamido" refers to a $\text{RS}(=\text{O})_2\text{N}(\text{R}')$ - group with R and R' as defined herein. Exemplary, non-limiting N-sulfonamido groups are $-\text{NHSO}_2\text{alkyl}$ such as $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CH}_2\text{CH}_3$ or $-\text{NHSO}_2\text{(isopropyl)}$, and $-\text{NHSO}_2\text{(optionally substituted aryl)}$ such as $-\text{NHSO}_2\text{phenyl}$ $-\text{NHSO}_2\text{-(2-cyanophenyl)}$, $-\text{NHSO}_2\text{-(3-cyanophenyl)}$, $-\text{NHSO}_2\text{-(4-cyanophenyl)}$, $-\text{NHSO}_2\text{-(2-aminophenyl)}$, $-\text{NHSO}_2\text{-(3-aminophenyl)}$ or $-\text{NHSO}_2\text{-(4-aminophenyl)}$.

15

As used herein, the term "S-sulfonamido" refers to a $-\text{S}(=\text{O})_2\text{NRR}'$, group, with R and R' as defined herein.

As used herein, the term "urea" refers to a $-\text{N}(\text{R})\text{C}(=\text{O})\text{N}(\text{R})(\text{R}')$ group wherein R and R' are as defined herein.

20 The term R or the term R', appearing by itself and without a number designation, unless otherwise defined, refers to a moiety selected from the group consisting of hydrogen, alkyl, cycloalkyl, heteroalkyl, aryl, heteroaryl and heterocycloalkyl.

25 Whether an R group has a number designation or not, every R group, including R, R' and R^z where $z=(1, 2, 3, \dots, z)$, every substituent, and every term should be understood to be independent of every other in terms of selection from a group. Should any variable, substituent, or term (e.g., aryl, heterocycle, R, etc.) occur more than one time in a formula or generic structure, its definition at each occurrence is independent of the definition at every other occurrence. Those of skill in the art will further recognize that certain groups may be attached to a parent molecule or may occupy a position in a chain of elements from either end as written. Thus, by way of example only, an unsymmetrical group such as $-\text{C}(=\text{O})\text{N}(\text{R})-$ may be attached to the parent moiety at either the carbon or the nitrogen.

30 As used herein, the term "optionally substituted" means the preceding or anteceding group may be substituted or unsubstituted. When substituted and unless otherwise specified, the substituents of an "optionally substituted" group may include, without limitation, one or more substituents independently selected from the following groups or a particular designated set of groups, alone or in combination: lower alkyl, lower alkenyl,

lower alkynyl, lower alkanoyl, lower heteroalkyl, lower heterocycloalkyl, lower haloalkyl, lower cycloalkyl, phenyl, aryl, heteroaryl, pyridyl, aryloxy, lower alkoxy, lower haloalkoxy, oxo, lower acyloxy, carbonyl, carboxyl, lower alkylcarbonyl, lower carboxyester, lower carboxamido, cyano, halogen, hydroxyl, amino, amido, nitro, thiol, lower alkylthio, lower haloalkylthio, lower perhaloalkylthio, arylthio, sulfonate, sulfonic acid, trisubstituted silyl, N₃, SH, SCH₃, C(O)CH₃, CO₂CH₃, CO₂H, carbamate, and urea. Two substituents may be joined together to form a fused five-, six-, or seven-membered carbocyclic or heterocyclic ring consisting of zero to three heteroatoms, for example forming methylenedioxy or ethylenedioxy. An optionally substituted group may be unsubstituted (e.g., -CH₂CH₃), fully substituted (e.g., -CF₃CF₃), monosubstituted (e.g., -CH₂CH₂F) or substituted at a level anywhere in-between fully substituted and monosubstituted (e.g., -CH₂CF₃). Where substituents are recited without qualification as to substitution, both substituted and unsubstituted forms are encompassed. Where a substituent is qualified as "substituted," the substituted form is specifically intended. Additionally, different sets of optional substituents to a particular moiety may be defined as needed; in these cases, the optional substitution will be as defined, often immediately following the phrase, "optionally substituted with." In one specific definition, the optional substituents are chosen from hydroxyl, halo, alkyl, alkoxy, haloalkyl, haloalkoxy, -N(C₁₋₃ alkyl)₂, -NH(C₁₋₃ alkyl), -NHC(=O)(C₁₋₃ alkyl), -C(=O)OH, -C(=O)O(C₁₋₃ alkyl), -C(=O)(C₁₋₃ alkyl), -C(=O)NH₂, -C(=O)NH(C₁₋₃ alkyl), -C(=O)NH(cycloalkyl), -C(=O)N(C₁₋₃ alkyl)₂, -S(=O)₂(C₁₋₃ alkyl), -S(=O)₂NH₂, -S(=O)₂N(C₁₋₃ alkyl)₂, -S(=O)₂NH(C₁₋₃ alkyl), -CHF₂, -OCF₃, -OCHF₂, -SCF₃, -CF₃, -CN, -NH₂, -NO₂, or tetrazolyl.

As used herein, the term "optional substituent" denotes that the corresponding substituent may be present or may be absent. Accordingly, a compound having 1, 2 or 3 optional substituents may be unsubstituted or may be substituted with 1, 2 or 3 substituents.

As used herein, the term "treating a disease" refers to a slowing of or a reversal of the progress of the disease. Treating a disease includes treating a symptom and/or reducing the symptoms of the disease.

As used herein, the term "preventing a disease" refers to a slowing of the disease or of the onset of the disease or the symptoms thereof. Preventing a disease or disorder can include stopping the onset of the disease or symptoms thereof.

As used herein, the term "dosage unit" refers to a physically discrete unit, such as a capsule or tablet suitable as a unitary dosage for a human patient. Each unit contains a predetermined quantity of a compound of Formula I which was discovered or believed to produce the desired pharmacokinetic profile which yields the desired therapeutic effect. The dosage unit is composed of a compound of Formula I in association with at least one pharmaceutically acceptable carrier, salt, excipient, or combination thereof.

As used herein, the term "subject" or "patient" or "individual", such as the subject in need of treatment or prevention, may be a eukaryote, an animal, a vertebrate animal, a mammal, a rodent (e.g., a guinea pig, a hamster, a rat, a mouse), a murine (e.g., a mouse), a canine (e.g., a dog), a feline (e.g., a cat), an equine (e.g. a horse), a primate, a simian (e.g., a monkey or ape), a monkey (e.g., a marmoset, a baboon), an ape (e.g., a gorilla, chimpanzee, orangutang, gibbon), or a human. The meaning of the terms "eukaryote", "animal", "mammal", etc. is well known in the art and can, for example, be deduced from Wehner und Gehring (1995; Thieme Verlag). In the context of this invention, it is particularly envisaged that animals are to be treated which are economically, agronomically or scientifically important. Scientifically important organisms include, but are not limited to, mice, rats, and rabbits. Lower organisms such as, e.g., fruit flies like *Drosophila melanogaster* and nematodes like *Caenorhabditis elegans* may also be used in scientific approaches. Non-limiting examples of agronomically important animals are sheep, cattle and pig, while, for example, cats and dogs may be considered as economically important animals. Preferably, the subject/patient/individual is a mammal; more preferably, the subject/patient/individual is a human or a non-human mammal (such as, e.g., a guinea pig, a hamster, a rat, a mouse, a rabbit, a dog, a cat, a horse, a monkey, an ape, a marmoset, a baboon, a gorilla, a chimpanzee, an orangutang, a gibbon, a sheep, cattle, or a pig); even more preferably, the subject/patient/individual is a human.

As used herein, the term "dose" or "dosage" refers to the amount of active ingredient that an individual takes or is administered at one time. For example, a 40 mg dose of a compound of Formula I refers to, in the case of a twice-daily dosage regimen, a situation where the individual takes 40 mg of a compound of Formula I twice a day, e.g., 40 mg in the morning and 40 mg in the evening. The 40 mg of a compound of Formula I dose can be divided into two or more dosage units, e.g., two 20 mg dosage units of a compound of Formula I in tablet form or two 20 mg dosage units of a compound of Formula I in capsule form.

As used herein, the term "therapeutically effective amount", such as the therapeutically effective amount of a compound of the present invention, refers to the amount sufficient to produce a desired biological effect (e.g., a therapeutic effect) in a subject. Accordingly, a therapeutically effective amount of a compound may be an amount which is sufficient to treat or prevent a disease or disorder, and/or delay the onset or progression of a disease or disorder, and/or alleviate one or more symptoms of the disease or disorder, when administered to a subject suffering from or susceptible to that disease or disorder.

As used herein, a "pharmaceutically acceptable prodrug" is a compound that may be converted under physiological conditions or by solvolysis to the specified compound or to a pharmaceutically acceptable salt of such compound.

As used herein, a "pharmaceutically acceptable salt" is intended to mean a salt that retains the biological effectiveness of the free acids and bases of the specified compound and that is not biologically or otherwise undesirable. A compound for use in the invention may possess a sufficiently acidic, a sufficiently basic, or both functional groups, and accordingly react with any of a number of inorganic or organic bases, and inorganic and 5 organic acids, to form a pharmaceutically acceptable salt. Exemplary pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a mineral or organic acid, such as hydrochlorides, hydrobromides, sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrophosphates, dihydrophosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, nitrates, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrates, caproates, 10 heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4 dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, 15 phenylpropionates, phenylbutyrate, citrates, lactates, gamma-hydroxybutyrate, glycollates, tartrates, methane-sulfonates, ethane-sulfonates, propanesulfonates, benzenesulfonates, toluenesulfonates, trifluoromethansulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, or mandelates, pyruvates, 20 stearates, ascorbates, or salicylates. When the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands such as ammonia, alkylamines, hydroxyalkylamines, lysine, arginine, N-methylglucamine, procaine and the like.

Pharmaceutically acceptable salts are well known in the art.

As used herein, a "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" refers to a non-API (API refers to Active Pharmaceutical Ingredient) substances such as disintegrators, binders, fillers, and 25 lubricants used in formulating pharmaceutical products. They are generally safe for administering to humans according to established governmental standards, including those promulgated by the United States Food and Drug Administration and the European Medical Agency. Pharmaceutically acceptable carriers or excipients are well known to those skilled in the art.

In the chemical drawings used herein, such as in the drawings for specific D rings disclosed above, the wavy 30 line represents the point of attachment to the remainder of the compound.

As is understood by the skilled artisan, certain variables in the list of substituents are repetitive (different name for the same substituent), generic to other terms in the list, and/or partially overlap in content with other terms. In the compounds of the invention, the skilled artisan recognizes that substituents may be attached to the 35 remainder of the molecule via a number of positions and the preferred positions are as illustrated in the Examples.

The compounds of the invention are unexpectedly potent and selective inhibitors of LSD1. Avoiding inhibition of "off-targets" can avoid unwanted or undesirable side-effects like the cheese effect associated with MAO-A. The compounds of the invention are thus useful for the treatment or prevention of any disease or disorder associated with LSD1. This includes cancer, neurological diseases and viral infections, among others.

5 Preferably, the compounds of formula I, including compounds of formula Ia, Ia-1, Ib and Ic, as well as any salts and solvates thereof, are used for the treatment or prevention of cancer, and most preferably for the treatment of cancer. Cancers that may be treated (or prevented) with the compounds of the invention include, but are not limited to cancers such as:

10 Hematologic cancers (also designated herein as blood cancers), including cancers of the blood, bone marrow and lymph nodes such as leukemias (e.g. acute myelogenous leukemia (AML), acute promyelocytic leukemia (APL), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia), myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome, and lymphomas (e.g. Hodgkin's disease, non-Hodgkin's lymphoma (malignant lymphoma));

15 Breast cancer, including invasive ductal carcinoma, in situ ductal carcinoma, lobular carcinoma, and mixed ductal and lobular carcinoma;

20 Lung cancer such as bronchogenic carcinoma (e.g. squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, and mesothelioma;

25 Gastrointestinal cancers such as esophagus (e.g. squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (e.g. carcinoma, lymphoma, leiomyosarcoma), pancreas (e.g. ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (e.g. adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), and large bowel (e.g. adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma);

30 Genitourinary tract cancers such as kidney (e.g. adenocarcinoma, Wilm's tumor (nephroblastoma), lymphoma, leukemia), bladder and urethra (e.g. squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (e.g. adenocarcinoma, sarcoma), and testis (e.g. seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma);

35 Liver cancer such as hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, and hemangioma;

Bone cancer such as osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma,

malignant giant cell tumor chordoma, osteochronfroma (osteocartilaginous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors;

Nervous system cancers such as skull (e.g. osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (e.g. meningioma, meningiosarcoma, gliomatosis), brain (e.g. astrocytoma, medulloblastoma, 5 glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendrogioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, and sarcoma;

Gynecological cancers such as uterus (e.g. endometrial carcinoma), cervix (e.g. cervical carcinoma, pre-tumor cervical dysplasia), ovaries (e.g. ovarian carcinoma (serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma), granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, 10 dysgerminoma, malignant teratoma), vulva (e.g. squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (e.g. clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma)), and fallopian tubes (carcinoma);

Cardiac cancer such as sarcoma (e.g. angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma;

15 Skin cancer such as malignant melanoma, basal cell carcinoma, squamous cell carcinoma, and Kaposi's sarcoma; and

Adrenal glands cancer such as neuroblastoma.

Accordingly, in one embodiment, the compounds of the invention are used for the treatment or prevention of 20 cancer, particularly for the treatment of cancer, wherein said cancer is chosen from blood cancer, leukemia, lymphoma, breast cancer, lung cancer, prostate cancer, colorectal cancer, brain cancer, neuroblastoma, bladder cancer, liver cancer, sarcoma, myeloma and skin cancer. In another embodiment, the compounds of the invention are used for the treatment or prevention (particularly for the treatment) of blood cancers (also known as hematological cancers), including leukemias (for example, acute myelogenous leukemia (AML), 25 chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), or hairy cell leukemia), lymphomas and myelomas, prostate cancer, breast cancer, lung cancer, colorectal cancer, brain cancer or skin cancer. In a preferred embodiment, the compounds of formula I, including compounds of formula Ia, Ia-1, Ib and Ic, are used for the treatment of a blood cancer. More preferably, the compounds of formula I, including compounds of 30 formula Ia, Ia-1, Ib and Ic, are used for the treatment of leukemia, including acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), and hairy cell leukemia.

Typically, compounds according to Formula I, Ia, Ia-1, Ib or Ic can be effective at an amount of from about 0.01 35 $\mu\text{g}/\text{kg}$ to about 100 mg/kg per day based on total body weight. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at predetermined intervals of time.

5 The suitable dosage unit for each administration can be, e.g., from about 1 μ g to about 2000 mg, preferably from about 5 μ g to about 1000 mg. Even more preferably, the amount of active ingredient administered is from about 5 μ g to about 100 mg per day. These doses will depend on the pharmacokinetic parameters of the particular compound and other ADME properties as well as the efficacy of the compound in a particular disease setting.

10 It should be understood that the dosage ranges set forth above are exemplary only and are not intended to limit the scope of this invention. The therapeutically effective amount for each active compound can vary with factors including but not limited to the activity of the compound used, stability of the active compound in the patient's body, the severity of the conditions to be alleviated, the total weight of the patient treated, the route of administration, the ease of absorption, distribution, and excretion of the active compound by the body, the age and sensitivity of the patient to be treated, and the like, as will be apparent to a skilled artisan. The amount of administration can be adjusted as the various factors change over time.

15 While it is possible that a compound of the invention may be administered for use in therapy directly as such, it is typically administered in the form of a pharmaceutical composition, which comprises said compound as active pharmaceutical ingredient together with one or more pharmaceutically acceptable excipients or carriers.

20 The compounds of the invention may be administered by any means that accomplish their intended purpose. Examples include administration by the oral, parenteral, intravenous, subcutaneous or topical routes.

25 For oral delivery, the active compounds can be incorporated into a formulation that includes pharmaceutically acceptable carriers such as binders (e.g., gelatin, cellulose, gum tragacanth), excipients (e.g., starch, lactose), lubricants (e.g., magnesium stearate, silicon dioxide), disintegrating agents (e.g., alginate, Primogel, and corn starch), and sweetening or flavoring agents (e.g., glucose, sucrose, saccharin, methyl salicylate, and peppermint). The formulation can be orally delivered in the form of enclosed gelatin capsules or compressed tablets. Capsules and tablets can be prepared in any conventional techniques. The capsules and tablets can also be coated with various coatings known in the art to modify the flavors, tastes, colors, and shapes of the capsules and tablets. In addition, liquid carriers such as fatty oil can also be included in capsules.

30 Suitable oral formulations can also be in the form of suspension, syrup, chewing gum, wafer, elixir, and the like. If desired, conventional agents for modifying flavors, tastes, colors, and shapes of the special forms can also be included. In addition, for convenient administration by enteral feeding tube in patients unable to swallow, the active compounds can be dissolved in an acceptable lipophilic vegetable oil vehicle such as olive oil, corn oil and safflower oil.

The active compounds can also be administered parenterally in the form of solution or suspension, or in lyophilized form capable of conversion into a solution or suspension form before use. In such formulations, diluents or pharmaceutically acceptable carriers such as sterile water and physiological saline buffer can be used. Other conventional solvents, pH buffers, stabilizers, anti-bacteria agents, surfactants, and antioxidants can all be included. For example, useful components include sodium chloride, acetates, citrates or phosphates buffers, glycerin, dextrose, fixed oils, methyl parabens, polyethylene glycol, propylene glycol, sodium bisulfate, benzyl alcohol, ascorbic acid, and the like. The parenteral formulations can be stored in any conventional containers such as vials and ampoules.

Routes of topical administration include nasal, bucal, mucosal, rectal, or vaginal applications. For topical administration, the active compounds can be formulated into lotions, creams, ointments, gels, powders, pastes, sprays, suspensions, drops and aerosols. Thus, one or more thickening agents, humectants, and stabilizing agents can be included in the formulations. Examples of such agents include, but are not limited to, polyethylene glycol, sorbitol, xanthan gum, petrolatum, beeswax, or mineral oil, lanolin, squalene, and the like. A special form of topical administration is delivery by a transdermal patch. Methods for preparing transdermal patches are disclosed, e.g., in Brown, *et al.* (1988) *Ann. Rev. Med.* 39:221-229 which is incorporated herein by reference.

Subcutaneous implantation for sustained release of the active compounds may also be a suitable route of administration. This entails surgical procedures for implanting an active compound in any suitable formulation into a subcutaneous space, e.g., beneath the anterior abdominal wall. See, e.g., Wilson *et al.* (1984) *J. Clin. Psych.* 45:242-247. Hydrogels can be used as a carrier for the sustained release of the active compounds. Hydrogels are generally known in the art. They are typically made by crosslinking high molecular weight biocompatible polymers into a network, which swells in water to form a gel like material. Preferably, hydrogels are biodegradable or biosorbable. For purposes of this invention, hydrogels made of polyethylene glycols, collagen, or poly(glycolic-co-L-lactic acid) may be useful. See, e.g., Phillips *et al.* (1984) *J. Pharmaceut. Sci.*, 73: 1718-1720.

The active compounds can also be conjugated, to a water soluble non- immunogenic non-peptidic high molecular weight polymer to form a polymer conjugate. For example, an active compound is covalently linked to polyethylene glycol to form a conjugate. Typically, such a conjugate exhibits improved solubility, stability, and reduced toxicity and immunogenicity. Thus, when administered to a patient, the active compound in the conjugate can have a longer half-life in the body, and exhibit better efficacy. See generally, Burnham (1994) *Am. J. Hosp. Pharm.* 15:210-218. PEGylated proteins are currently being used in protein replacement therapies and for other therapeutic uses. For example, PEGylated interferon (PEG-INTRON A®) is clinically used for treating Hepatitis B. PEGylated adenosine deaminase (ADAGEN®) is being used to treat severe

combined immunodeficiency disease (SCIDS). PEGylated L-asparaginase (ONCAPSPAR®) is being used to treat acute lymphoblastic leukemia (ALL). It is preferred that the covalent linkage between the polymer and the active compound and/or the polymer itself is hydrolytically degradable under physiological conditions. Such conjugates known as "prodrugs" can readily release the active compound inside the body. Controlled release of an active compound can also be achieved by incorporating the active ingredient into microcapsules, nanocapsules, or hydrogels generally known in the art. Other pharmaceutically acceptable prodrugs of the compounds of this invention include, but are not limited to, esters, carbonates, thiocarbonates, N-acyl derivatives, N-acyloxyalkyl derivatives, quaternary derivatives of tertiary amines, N-Mannich bases, Schiff bases, amino acid conjugates, phosphate esters, metal salts and sulfonate esters.

10 Liposomes can also be used as carriers for the active compounds of the present invention. Liposomes are micelles made of various lipids such as cholesterol, phospholipids, fatty acids, and derivatives thereof. Various modified lipids can also be used. Liposomes can reduce the toxicity of the active compounds, and increase their stability. Methods for preparing liposomal suspensions containing active ingredients therein are generally known in the art. See, e.g., U.S. Patent No. 4,522,811 ; Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N. Y. (1976).

20 The active compounds can also be administered in combination with another active agent that synergistically treats or prevents the same symptoms or is effective for another disease or symptom in the patient treated so long as the other active agent does not interfere with or adversely affect the effects of the active compounds of this invention. Such other active agents include but are not limited to anti-inflammatory agents, antiviral agents, antibiotics, antifungal agents, antithrombotic agents, cardiovascular drugs, cholesterol lowering agents, anti-cancer drugs, hypertension drugs, and the like.

25 Combination therapy includes administration of a single pharmaceutical dosage formulation which contains a compound of the invention and one or more additional active agents, as well as administration of the compound of the invention and each additional active agent in its own separate pharmaceutical dosage formulation. If administered separately, the administration can be simultaneous, sequential or separate, and the compound of the invention and the additional therapeutic agent(s) can be administered via the same administration route or 30 using different administration routes, for example one compound can be administered orally and the other intravenously.

35 In particular, when a compound of formula I, Ia, Ia-1, Ib or Ic is used for the treatment or prevention of cancer, said compound can be administered in combination with one or more further agents known to be useful in the treatment or prevention of cancer, including chemotherapy or radiotherapy.

Typically, for combination therapy with a compound of the invention any antineoplastic agent that has activity versus a cancer being treated or prevented with a compound of the invention may be used. Examples of antineoplastic agents that can be used in combination with the compounds and methods of the present invention include, in general, and as appropriate, alkylating agents, anti-metabolites, epidophyllotoxins, 5 antineoplastic enzymes, topoisomerase inhibitors, procarbazines, mitoxantrones, platinum coordination complexes, biological response modifiers and growth inhibitors, hormonal/anti-hormonal therapeutic agents and haematopoietic growth factors. Exemplary classes of antineoplastic agents include the anthracyclines, vinca drugs, mitomycins, bleomycins, cytotoxic nucleosides, epothilones, discodermolides, pteridines, diyenes and podophyllotoxins. Particularly useful members of those classes include, for example, carminomycin, 10 daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podo-phyllotoxin derivatives such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosidine, vindesine, leurosine, paclitaxel and the like. Other useful antineoplastic agents include 15 estramustine, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabin, idarubicin, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins.

Thus, a compound of formula I, Ia, Ia-1, Ib or Ic according to the present invention can be used for the 20 treatment or prevention of cancer, wherein said compound is to be administered in combination with one or more antineoplastic agents. The antineoplastic agents to be administered for combination therapy may be selected, as appropriate, from: a tumor angiogenesis inhibitor (for example, a protease inhibitor, an epidermal growth factor receptor kinase inhibitor, or a vascular endothelial growth factor receptor kinase inhibitor); a 25 cytotoxic drug (for example, an antimetabolite, such as purine and pyrimidine analog antimetabolites); an antimitotic agent (for example, a microtubule stabilizing drug or an antimitotic alkaloid); a platinum coordination complex; an anti-tumor antibiotic; an alkylating agent (for example, a nitrogen mustard or a nitrosourea); an endocrine agent (for example, an adrenocorticosteroid, an androgen, an anti-androgen, an estrogen, an anti-estrogen, an aromatase inhibitor, a gonadotropin-releasing hormone agonist, or a somatostatin analog); or a 30 compound that targets an enzyme or receptor that is overexpressed and/or otherwise involved in a specific metabolic pathway that is misregulated in the tumor cell (for example, ATP and GTP phosphodiesterase inhibitors, histone deacetylase inhibitors, protein kinase inhibitors (such as serine, threonine and tyrosine kinase inhibitors (for example, Abelson protein tyrosine kinase)) and the various growth factors, their receptors and kinase inhibitors therefor (such as epidermal growth factor receptor kinase inhibitors, vascular endothelial growth factor receptor kinase inhibitors, fibroblast growth factor inhibitors, insulin-like growth factor receptor inhibitors and platelet-derived growth factor receptor kinase inhibitors)); aminopeptidase inhibitors; proteasome 35 inhibitors; cyclooxygenase inhibitors (for example, cyclooxygenase-1 or cyclooxygenase-2 inhibitors);

topoisomerase inhibitors (for example, topoisomerase I inhibitors or topoisomerase II inhibitors); or retinoid agents.

An alkylating agent which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, a nitrogen mustard (such as cyclophosphamide, mechlorethamine (chlormethine), uramustine, melphalan, chlorambucil, ifosfamide, bendamustine, or trofosfamide), a nitrosourea (such as carmustine, streptozocin, fotemustine, lomustine, nimustine, prednimustine, ranimustine, or semustine), an alkyl sulfonate (such as busulfan, mannosulfan, or treosulfan), an aziridine (such as hexamethylmelamine (altretamine), triethylenemelamine, ThioTEPA (N,N'N'-triethylenethiophosphoramide), carboquone, or triaziquone), a hydrazine (such as procarbazine), a triazene (such as dacarbazine), or an imidazotetrazines (such as temozolomide).

A platinum coordination complex which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, cisplatin, carboplatin, nedaplatin, oxaliplatin, satraplatin, or triplatin tetranitrate.

A cytotoxic drug which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, an antimetabolite, including folic acid analog antimetabolites (such as aminopterin, methotrexate, pemetrexed, or raltitrexed), purine analog antimetabolites (such as cladribine, clofarabine, fludarabine, 6-mercaptopurine (including its prodrug form azathioprine), pentostatin, or 6-thioguanine), and pyrimidine analog antimetabolites (such as cytarabine, decitabine, azacytidine, 5-fluorouracil (including its prodrug forms capecitabine and tegafur), floxuridine, gemcitabine, enocitabine, or sapacitabine).

An antimitotic agent which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, a taxane (such as docetaxel, larotaxel, ortataxel, paclitaxel/taxol, or tesetaxel), a Vinca alkaloid (such as vinblastine, vincristine, vinflunine, vindesine, vinzolidine, or vinorelbine), an epothilone (such as epothilone A, epothilone B, epothilone C, epothilone D, epothilone E, or epothilone F) or an epothilone B analog (such as ixabepilone/azaepothilone B).

An anti-tumor antibiotic which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, an anthracycline (such as aclarubicin, daunorubicin, doxorubicin, epirubicin, idarubicin, amrubicin, pirarubicin, valrubicin, or zorubicin), an anthracenedione (such as mitoxantrone, or pixantrone) or an anti-tumor antibiotic isolated from *Streptomyces* (such as actinomycin (including actinomycin D), bleomycin, mitomycin (including mitomycin C), or plicamycin).

A tyrosine kinase inhibitor which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, axitinib, bosutinib, cediranib, dasatinib, erlotinib, gefitinib, imatinib, lapatinib, lestaurtinib, nilotinib, semaxanib, sorafenib, sunitinib, or vandetanib.

5 A topoisomerase-inhibitor which can be used as an antineoplastic agent in combination with a compound of the present invention may be, for example, a topoisomerase I inhibitor (such as irinotecan, topotecan, camptothecin, belotecan, rubitecan, or lamellarin D) or a topoisomerase II inhibitor (such as amsacrine, etoposide, etoposide phosphate, teniposide, or doxorubicin).

10 Further antineoplastic agents may be used in combination with a compound of the present invention. The antineoplastic agents may include biological or chemical molecules, such as TNF-related apoptosis-inducing ligand (TRAIL), tamoxifen, toremifene, fluoxymesterol, raloxifene, diethylstibestrol, bicalutamide, nilutamide, flutamide, aminoglutethimide, anastrozole, tetrazole, luteinizing hormone release hormone (LHRH) analogues, ketoconazole, goserelin acetate, leuprolide, megestrol acetate, prednisone, mifepristone, amsacrine, 15 bexarotene, estramustine, irofulven, trabectedin, cetuximab, panitumumab, tositumomab, alemtuzumab, bevacizumab, edrecolomab, gemtuzumab, alvocidib, seliciclib, aminolevulinic acid, methyl aminolevulinate, efaproxiral, porfimer sodium, talaporfin, temoporfin, verteporfin, anagrelide, arsenic trioxide, atrasentan, bortezomib, carmofur, celecoxib, demecolcine, elesclomol, elsamitrucin, etoglucid, Isonidamine, lucanthone, masoprocol, mitobronitol, mitoguazone, mitotane, oblimersen, omacetaxine, sitimagene, ceradenovect, tegafur, 20 testolactone, tiazofurine, tipifarnib, and vorinostat.

Examples of retinoid agents include all natural, recombinant, and synthetic derivatives or mimetics of vitamin A, for example, retinyl palmitate, retinyl-beta-glucuronide (vitamin A1 beta-glucuronide), retinyl phosphate (vitamin A1 phosphate), retinyl esters, 4-oxoretinol, 4-oxoretinaldehyde, 3-dehydroretinol (vitamin A2), 11-cis-retinal (11-cis-retinaldehyde, 11-cis or neo b vitamin A1 aldehyde), 5,6-epoxyretinol (5,6-epoxy vitamin A1 alcohol), anhydroretinol (anhydro vitamin A1) and 4-ketoretinol (4-keto-vitamin A1 alcohol), all-trans retinoic acid (ATRA; Tretinoin; vitamin A acid; 3,7-dimethyl-9-(2,6,6,-trimethyl-1-cyclohexen-1-yl)-2,4,6,8-nonatetraenoic acid [CAS No. 302-79-4]), lipid formulations of all-trans retinoic acid (e.g., ATRA-IV), 9-cis retinoic acid (9-cis-RA; Alitretinoin; PanretinTM; LGD1057), 13-cis retinoic acid (Isotretinoin), (E)-4-[2-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthalenyl)-1-propenyl]-benzoic acid, 3-methyl-(E)-4-[2-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthalenyl)-1-propenyl]-benzoic acid, Fenretinide (N-(4-hydroxyphenyl)retinamide; 4-HPR), Etretinate ((all-E)-9-(4-methoxy-2,3,6-trimethylphenyl)-3,7-dimethyl-2,4,6,8-nonatetraenoic acid ethyl ester; Tegison), Acitretin ((all-E)-9-(4-methoxy-2,3,6-trimethylphenyl)-3,7-dimethyl-2,4,6,8-nonatetraenoic acid; Ro 10-1670; Soriatane; Neotigason), Tazarotene (ethyl 6-[2-(4,4-dimethylthiochroman-6-yl)-ethynyl] nicotinate; 25 Tazorac; Avage; Zorac), Tocoretinate (9-cis-tretinoin; Tocoferil), Adapalene (6-[3-(1-adamantyl)-4-methoxyphenyl]-2-naphthoic acid; Differin), Motretinide (trimethylmethoxyphenyl-N-ethyl retinamide;

Trasmaderm), retinaldehyde (Retinal), CD437 (6-[3-(1-adamantyl)-4-hydroxyphenyl]-2-naphthalene carboxylic acid; AHPN), CD2325, ST1926 ([E-3-(4'-hydroxy-3'-adamantylbiphenyl-4-yl)acrylic acid), ST1878 (methyl 2-[3-[2-[3-(2-methoxy-1,1-dimethyl-2-oxoethoxy)phenoxy]ethoxy]phenoxy]isobutyrate), ST2307, ST1898, ST2306, ST2474, MM11453, MM002 (3-CI-AHPC), MX2870-1, MX3350-1, MX84, and MX90-1, docosahexaenoic acid (DHA), phytanic acid (3,7,11,15-tetramethyl hexadecanoic acid), MS6682 (methoprene acid), LG100268 (LG268), LG100324, SR11203 ([2-(4-carboxyphenyl)-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1,3-dithiane), SR11217 (4-(2-methyl-1-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)propenyl)benzoic acid), SR11234, SR11236 (2-(4-carboxyphenyl)-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1,3-dioxane), SR11246, AGN194204, derivatives of 9-cis-RA such as LGD1069 (3-methyl TTNEB; Bexarotene; Targretin®; 4-[1-(5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl]benzoic acid).

Examples of histone deacetylase inhibitors include, without limitation, MS-275 (SNDX-275; Entinostat), FK228 (FR901228; depsipeptide; Romidepsin), CI-994 (Acetylinaline; Tacedinaline), Apicidin (cyclo[(2S)-2-amino-8-oxodecanoyl-1-methoxy-L-tryptophyl-L-isoleucyl-(2R)-2-piperidinexcarbonyl]), A-161906 (7-[4-(4-cyanophenyl)phenoxy]-heptanohydroxamic acid), Scriptaid (6-(1,3-Dioxo-1H,3H-benzo[de]isoquinolin-2-yl)-hexanoic acid hydroxyamide), PXD-101 (Belinostat), CHAP (cyclic hydroxamic acid-containing peptide), LAQ-824 (Dacinostat), BML-EI319 (Depudecin), O3139 (Oxamflatin), NSC 696085 (Pyroxamide), MW2796; MW2996, T2580 (Trapoxin A), AN-9 (Pivanex), W222305 (Tributyrin) Trichostatin A, Trichostatin C, Butyric acid, Valproic acid (VPA), Suberoylanilide hydroxamic acid (SAHA; Vorinostat), m-Carboxycinnamic acid bishydroxamide (CBHA), Salicylbishydroxamic acid (S607; SHA; SHAM); Suberoyl bishydroxamic acid (SBHA); Azelaic bishydroxamic acid (ABHA); Azelaic-1-hydroxamate-9-anilide (AAHA); 3CI-UCHA (6-(3-chlorophenylureido) caproic hydroxamic acid); and sodium butyrate, 4-phenylbutyrate, phenylacetate, valerate, isovalerate, butyramide, isobutyramide, 3-bromopropionate, and valproate.

Also biological drugs, like antibodies, antibody fragments, antibody constructs (for example, single-chain constructs), and/or modified antibodies (like CDR-grafted antibodies, humanized antibodies, "full humanized" antibodies, etc.) directed against cancer or tumor markers/factors/cytokines involved in cancer can be employed in cotherapeutic approaches with the compounds of the invention. Examples of such biological molecules are alemtuzumab, apolizumab, aselizumab, atlizumab, bapineuzumab, bevacizumab, bivatuzumab mertansine, cantuzumab mertansine, cedelizumab, certolizumab pegol, cidefusituzumab, cideftuzumab, daclizumab, eculizumab, efalizumab, epratuzumab, erlizumab, felizumab, fontolizumab, gemtuzumab ozogamicin, inotuzumab ozogamicin, ipilimumab, labetuzumab, lintuzumab, matuzumab, mepolizumab, motavizumab, motovizumab, natalizumab, nimotuzumab, nolovizumab, numavizumab, ocrelizumab, omalizumab, palivizumab, pascolizumab, pecfusituzumab, pectuzumab, pertuzumab, pexelizumab, ralivizumab, ranibizumab, reslivizumab, reslizumab, resyvizumab, rituximab, rovelizumab, rolizumab,

sibrotuzumab, siplizumab, sontuzumab, tacatuzumab tetraxetan, tadocizumab, talizumab, tefibazumab, tocilizumab, toralizumab, trastuzumab, tucotuzumab celmoleukin, tucusituzumab, umavizumab, urtoxazumab, and visilizumab.

5 Other biologic agents include, but are not limited to, immunomodulating proteins such as cytokines (such as interleukin-2 (IL-2, Aldesleukin), Epoietin-alpha.; EPO), granulocyte-CSF (G-CSF; Filgrastin), and granulocyte-macrophage-CSF (GM-CSF; Sargramostim) and interferons, (e.g., interferon-alpha, interferon-beta and interferon-gamma), bacillus Calmette-Guerin, levamisole, and octreotide, endostatin, tumor suppressor genes (e.g., DPC4, NF- 1, NF-2, RB, p53, WT1, BRCA1, and BRCA2), and cancer vaccines (e.g., tumor associated 10 antigens such as gangliosides (GM2), prostate specific antigen (PSA), alpha-fetoprotein (AFP), carcinoembryonic antigen (CEA) (produced by colon cancers and other adenocarcinomas, e.g., breast, lung, gastric, and pancreatic cancers), melanoma-associated antigens (MART-1, gap100, MAGE 1,3 tyrosinase), papillomavirus E6 and E7 fragments, whole cells or portions/lysates of autologous tumor cells and allogeneic tumor cells.


15

General Synthetic Route Description

Compounds of Formula I can be synthesized in accordance with or in analogy to the general routes described 20 below. Unless otherwise stated, in the methods described below the meanings of the different substituents in each synthetic intermediate and compound of formula I are the meanings described above with regard to a compound of formula I. Other routes known by the ordinary skilled artisan, as well as other reactants and intermediates, can also be used to arrive at the compounds of Formula I. The reaction schemes described 25 below are only meant to represent examples of the invention and are in no way meant to be a limit of the invention. In some of the processes described below it may be necessary or advisable to protect reactive or labile groups with conventional protecting groups. Both the nature of these protecting groups and the procedures for their introduction and removal are well known in the art (see for example Greene TW and Wuts PGM "Greene's Protecting Groups in Organic Synthesis", 4th edition, Wiley, 2006). Whenever a protecting group is present, a subsequent deprotection step will be required, which can be performed under standard conditions well known in the art, such as those described in the above reference.

30

In general, the compounds of formula I can be prepared by reductive alkylation of a cyclopropylamino derivative of formula II with a ketone of formula IIIa or and aldehyde of formula IIIb or IIIc, as shown below in Scheme 1:



SCHEME 1

Wherein A, B, D, R^W , R^X , R^Y , R^Z and L^2 have the meaning disclosed above in relation to a compound of formula I.


5

Said reductive alkylation can be performed under standard conditions for reductive alkylations, well known in the art. For example, a suitable set of conditions is reacting II with IIIa, IIIb or IIIc using a reducing agent such as a borohydride (e.g. sodium triacetoxyborohydride or sodium borohydride) in a suitable solvent such as dichloroethane or methanol, optionally in the presence of an acid such as acetic acid. In order to conduct the reaction, it is necessary that any other amino group that may be present either in II or in IIIa, IIIb or IIIc be protected using conventional amino-protecting groups to avoid any side reactions; a subsequent deprotection step will be required then if such amino protecting group is present, in order to obtain a compound of formula I. Any suitable amino-protecting group may be used, such as for example a tert-butoxycarbonyl (Boc) group. If Boc is used, deprotection can be carried out under standard conditions, for example under acidic conditions using HCl in an organic solvent such as diethyl ether or 1,4-dioxane, or trifluoroacetic acid (TFA) in dichloromethane. When HCl is used in the last step of the synthesis, compounds of formula I are obtained as a hydrochloride salt. Likewise, if TFA is used, the compounds will be obtained as a trifluoroacetate.

The cyclopropylamino derivatives of formula II and the ketones of formula IIIa and the aldehydes of formula IIIb or IIIc are commercially available or can be prepared following methods disclosed in the literature.

More detailed methods to obtain compounds of formula I are described below.

The compounds of Formula I wherein R^w , R^x , R^y and $R^z = H$ can be synthesized, for example, by the general route described in Scheme 2. This route is particularly suitable for compounds wherein $B = H$ or R^1 since the corresponding aldehyde (1) is either commercially available or can be readily obtained. In Scheme 2 below, for schematic purposes "B" has been omitted.



SCHEME 2: DCE (Dichloroethane), DMF (N,N-dimethylformamide), DMSO (Dimethyl sulfoxide), THF (Tetrahydrofuran). aq = aqueous.


Aldehydes of Formula (1) are subjected to a Horner-Wadsworth-Emmons reaction using triethyl phosphono acetate and a base preferably potassium tert-butoxide in a suitable solvent such as tetrahydrofuran to get the ethyl acrylate derivatives of formula (2) which are then subjected to cyclopropanation reaction using trimethylsulfoxonium iodide and sodium hydride in dimethyl sulfoxide as a solvent leading to (trans)-ethyl cyclopropanecarboxylate derivatives of formula (3) (obtained as a trans ((1S, 2R) and (1R, 2S)) racemic mixture). Hydrolysis to the corresponding (trans)-cyclopropanecarboxylic acid derivatives of formula (4) can be performed under basic conditions using for example NaOH in a suitable solvent such as MeOH. The subsequent reaction of compound (4), first with ethyl chloroformate and triethylamine in acetone and later with sodium azide in water leads to the formation of (trans)-cyclopropanecarbonyl azide derivatives of formula (5). Reaction with tert-butanol results in the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (6). Alkylation with alkyl halides of formula (7) under basic conditions using for example NaH in a suitable solvent such as DMF leads to the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (8). Deprotection of the Boc-group in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or using HCl in 1,4-dioxane leads to the formation of the (trans)-cyclopropanamine derivatives of formula (9), which corresponds to a compound of formula I, and particularly Ia, wherein B is H or R¹.

Alternatively, the (trans)-cyclopropanamine derivatives of formula (12) can be synthesized by reaction of aldehydes of formula (1) with nitromethane and ammonium acetate using tetrahydrofuran as a solvent, leading to the formation of nitrostyrene of formula (10). Later cyclopropanation reaction using trimethylsulfoxonium iodide and potassium tert-butoxide results in the formation of trans nitrocyclopropyl derivatives of formula (11) (obtained as a trans ((1S, 2R), (1R, 2S)) racemic mixture) and final reduction using zinc in hydrochloric acid affords the (trans)-cyclopropanamine derivatives of formula (12).

Reductive alkylation of the derivatives of formula (12) with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under standard conditions, for example using sodium triacetoxyborohydride or sodium borohydride as reducing agent in a suitable solvent such as dichloroethane or methanol leads to the formation of (trans)-cyclopropylamino derivatives of formula (14a), (14b) or (14c) respectively, which corresponds to a compound of formula I, and particularly Ia, wherein R^w, R^x, R^y and R^z = H. In case the ketones of formula (13a) or aldehydes of formula (13b) or (13c) containing a protected amino group, for example a Boc-protected amine (Boc: tert-Butyloxycarbonyl), an additional deprotection reaction step will be required to render a compound (14a), (14b) or (14c) respectively, which can be performed in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether, or using HCl in 1,4-dioxane.

Aldehydes of formula (1), alkyl halides of formula (7), ketones of formula (13a) and aldehydes of formula (13b) and (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

5 The compounds of Formula I wherein B = -L¹-E, and R^w, R^x, R^y and R^z = H and L¹ = -(CH₂)_x-O- (wherein x is as defined previously) can be synthesized, for example, by the general route described in Scheme 3



SCHEME 3: DCE (Dichloroethane), DMF (N,N -dimethylformamide), DMSO (Dimethyl sulfoxide), THF (Tetrahydrofuran)


The alkylation of aldehydes of formula 1 (where $R^1 = -OH$) using bromo derivatives of formula (15) (other halo derivatives could also be used) and a base, preferably potassium carbonate in a suitable solvent such as N,N-dimethylformamide leads to the formation of the aldehyde derivatives of formula (16). These are subjected to a Horner-Wadsworth-Emmons reaction under the same conditions disclosed in Scheme 2 to get the ethyl acrylate derivatives of formula (17) which are then subjected to cyclopropanation reaction under the same conditions disclosed in Scheme 2 to give the (trans)-ethyl cyclopropanecarboxylate derivatives of formula (18). Following the same conditions disclosed for the conversion of a compound (3) to a compound (6) in scheme 2, a compound (18) is converted into the (trans)-cyclopropanamine derivative of formula (21). Later alkylation with alkyl halides of formula (7) under basic conditions using for example NaH in a suitable solvent such as DMF leads to the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (22). Deprotection of the Boc-group in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether leads to the formation of the (trans)-cyclopropanamine derivatives of formula (23), which corresponds to a compound of formula I wherein $B = -L^1-E$, R^w , R^x , R^y and $R^z = H$ and $L^1 = -(CH_2)_x-O-$.

Alternatively, the (trans)-cyclopropanamine derivatives of formula (26) can be synthesized from aldehydes of formula (16) by conversion into a nitrostyrene of formula (24), subsequent cyclopropanation to give a compound (25) and reduction of the nitro group under the same conditions disclosed in scheme 2 for the conversion of a compound (1) into a compound (12) via compounds (10) and (11).

Reductive alkylation of the derivatives of formula (26) with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under the conditions disclosed in scheme 1 or 2 yields compounds (27a), (27b) or (27c) respectively, which corresponds to a compound of formula I wherein $B = -L^1-E$, and R^w , R^x , R^y and $R^z = H$ and $L^1 = -(CH_2)_x-O-$. In case the ketones of formula (13a) or aldehydes of formula (13b) or (13c) containing a protected amino group, for example a Boc-protected amine (Boc: tert-butoxycarbonyl), an additional deprotection reaction step will be required to render compounds (27a), (27b) or (27c) respectively, which can be performed in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or using HCl in 1,4-dioxane.

Aldehydes of formula (1, where $R^1 = -OH$), bromo derivatives of formula (15) and ketones of formula (13a) or aldehydes of formula (13b) or (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

The compounds of Formula I where $B = -L^1-E$, and R^w , R^x , R^y and $R^z = H$ and $L^1 = -O-$ can be synthesized, for example, by the general route described in Scheme 4



SCHEME 4: DCE (Dichloroethane), DMF (N,N-dimethylformamide), DMSO (Dimethyl sulfoxide), Pd₂(dba)₃ (Tris(dibenzylideneacetone)dipalladium(0)), THF (Tetrahydrofuran), Xantphos (4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene).

5


Aldehydes of Formula (1, where R¹ = Br) are subjected to a Horner-Wadsworth-Emmons reaction under the conditions disclosed in Scheme 2 to get the ethyl acrylate derivatives of formula (2, where R¹ = Br) which are then subjected to cyclopropanation reaction under the same conditions disclosed in Scheme 2 for converting a

compound (2) into (3), leading to the (trans)-ethyl cyclopropanecarboxylate derivatives of formula (3, where R¹ = Br). Compounds of formula (3) (where R¹= Br) are converted into the corresponding (trans)-cyclopropanecarboxylic acid derivatives of formula (4, where R¹ = Br), which are then converted into the (trans)-cyclopropanecarbonyl azide derivatives of formula (5, where R¹ = Br) and then into the tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (6, where R¹ = Br) following the same conditions disclosed in Scheme 2. The reaction of the compounds (6, where R¹=Br) with hydroxy-derivatives of formula (28) using a palladium catalyst such as Tris(dibenzylideneacetone)dipalladium(0), Xantphos and a base such as sodium tert-butoxide in a suitable solvent such as dioxane leads to the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (29). Alkylation with alkyl halides of formula (7) under basic conditions using for example NaH in a suitable solvent such as DMF leads to the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (30). Deprotection of the Boc-group in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane leads to the formation of the (trans)-cyclopropanamine derivatives of formula (31), which correspond to a compound of formula I wherein B= -L¹-E, and R^w, R^x, R^y and R^z = H and L¹ is O.

Alternatively, deprotection of the Boc-group of derivatives of formula (29) in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane leads to the formation of the (trans)-cyclopropanamine derivatives of formula (32). Reductive alkylation with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under the conditions disclosed in Scheme 1 or 2 leads to the formation of (trans)-cyclopropylamino derivatives of formula (33a), (33b) or (33c) respectively, which correspond to a compound of formula I wherein B= -L¹-E, R^w, R^x, R^y and R^z = H and L¹ is O. In case the ketones of formula (13a) or aldehydes of formula (13b) or (13c) containing a protected amino group, for example a Boc-protected amine (Boc:tert-butoxycarbonyl), an additional deprotection reaction step will be required to render compounds (33a), (33b) or (33c) respectively, which can be performed in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane.

Aldehydes of formula (1, where R¹ = Br), hydroxy-derivatives of formula (28) and ketones of formula (13a) or aldehydes of formula (13b) or (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

The compounds of Formula I wherein B = -L¹-E, and R^w, R^x, R^y and R^z = H and L¹ =-NH- or -(CH₂)_x-NH- can be synthesized, for example, by the general route described in Scheme 5.



SCHEME 5: Boc_2O (Di-*tert*-butyl dicarbonate), DCE (Dichloroethane), DMF (*N,N*-dimethylformamide), DMSO (Dimethyl sulfoxide), $\text{Pd}_2(\text{dba})_3$ (Tris(dibenzylideneacetone)dipalladium(0)), THF (Tetrahydrofuran), Xantphos (4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene).

5 Tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (6, where R¹ = Br), obtained following the same procedure disclosed in Scheme 4, are converted into the (trans)-cyclopropanamine derivatives of formula (34) by alkylation with alkyl halides of formula (7) under basic conditions using for example NaH in a suitable solvent such as DMF leads to the formation of tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (34), which are then reacted with amino-derivatives of formula (35) using a palladium catalyst such as Tris(dibenzylideneacetone)dipalladium(0), Xantphos and a base such as sodium tert-butoxide in a suitable solvent such as dioxane to give the tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (36). Deprotection of the Boc-group of a compound (36) in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or using HCl in 1,4-dioxane leads to the formation of the (trans)-cyclopropanamine derivatives of formula (37), which correspond to compound of formula I wherein B = -L¹-E, R^w, R^x, R^y and R^z = H and L¹ = -NH- or -(CH₂)_x-NH-.


10

15 Alternatively, tert-butyl (trans)- cyclopropylcarbamate derivatives of formula (6, where R¹ = Br), obtained following the same procedure disclosed in Scheme 4, are converted into the (trans)-cyclopropanamine derivatives of formula (12, where R¹ = Br) by deprotection of the Boc-group in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane. Reductive alkylation of compounds (12, R¹= Br) with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under the conditions disclosed in Scheme 1 leads to the formation of (trans)-cyclopropylamino derivatives of formula (38a), (38b) or (38c) respectively. Later reaction with di-*t*-butyl dicarbonate under basic conditions 20 using for example triethylamine in a suitable solvent such tetrahydrofuran leads to the Boc-protected derivatives of formula (39a), (39b) or (39c) respectively. Reaction with amino-derivatives of formula (35) using a palladium catalyst such as Tris(dibenzylideneacetone)dipalladium(0), Xantphos and a base such as sodium tert-butoxide in a suitable solvent such as dioxane to give the tert-butyl (trans)-cyclopropylcarbamate derivatives of formula (40a), (40b) or (40c) respectively. Deprotection of the Boc-group in acidic conditions, for 25 example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane leads to the formation of the (trans)-cyclopropanamine derivatives of formula (41a), (41b) or (41c) respectively, which correspond to compound of formula I wherein B = -L¹-E, R^w, R^x, R^y and R^z = H and L¹ = -NH- or -(CH₂)_x-NH-.

30 Aldehydes of formula (1, where R¹ = Br), amino-derivatives of formula (35) and ketones of formula (13a) or aldehydes of formula (13b) or (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

35 The compounds of Formula I wherein R^w = F can be synthesized, for example, by the general route described in Scheme 6. This method is useful to obtain compounds having either a trans- or cis-configuration at the cyclopropyl ring (i.e. wherein the B-A- and -NH-D groups are in trans or cis configuration), or mixtures thereof, since the cyclopropanation reaction used yields a mixture of cis/trans isomers, as represented by the wavy line

in Scheme 6, which can be used as such to obtain compounds of the invention as cis/trans mixtures, or can be separated if desired to yield at the end of the synthesis the desired cis or trans products.




5 **SCHEME 6:** Boc₂O (Di-*tert*-butyl dicarbonate), DCE (Dichloroethane), DPPA (Diphenylphosphoryl azide), Cu(acac)₂ (Copper(II) acetylacetone), NBS (N-Bromosuccinimide),

Bromofluorination of derivatives of formula (42) using N-Bromosuccinimide and triethylamine trihydrofluoride in a suitable solvent such as dichloromethane leads to the formation of fluoro-derivatives of formula (43). Elimination reaction using a base, as for example potassium tert-butoxide in a suitable solvent, as for example pentane leads to fluoro-derivatives of formula (44). Cyclopropanation using ethyl diazoacetate and copper (II) 5 acetylacetone, as catalyst, in a suitable solvent such as dichloromethane leads to a 1:1 mixtures of cis- and trans- derivatives of formula (45). The diastereomers can be separated at this point either chromatographically or, after saponification (performed under basic conditions using for example NaOH in a suitable solvent such as MeOH), by recrystallisation of the corresponding carboxylic acids of formula (46). Curtius degradation to Boc-protected cyclopropylamines of formula (47) can be performed by using a base, as for example, triethylamine, 10 diphenylphosphoryl azide and di-*tert*-butyl dicarbonate in a suitable solvent, as for example, *tert*-butanol. Deprotection of the Boc-group in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or HCl in 1,4-dioxane leads to the formation of the cyclopropanamine derivatives 15 of formula (48). Reductive alkylation of the derivatives of formula (48) with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under standard conditions, for example using sodium triacetoxyborohydride as reducing agent in a suitable solvent such as dichloroethane leads to the formation of (trans)-cyclopropylamino derivatives of formula (49a), (49b) or (49c) respectively, which corresponds to a compound of 20 formula I, wherein R^w = F. In case the ketones of formula (13a) or aldehydes of formula (13b) or (13c) containing a protected amino group, for example a Boc-protected amine (Boc: *tert*-Butyloxycarbonyl), an additional deprotection reaction step will be required to render a compound (49a), (49b) or (49c) respectively, 25 which can be performed in acidic conditions, for example using HCl 2M in diethyl ether in a suitable solvent such as diethyl ether or using HCl in 1,4-dioxane.

Compounds of formula (42) and ketones of formula (13a) or aldehydes of formula (13b) or (13c) are 25 commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

Compounds of Formula I wherein R^w is H, fluoro or C₁₋₄ alkyl and R^x, R^y, R^z = H can be synthesized, for example, by the general route described in Scheme 7 below. This method is useful to obtain compounds 30 wherein R^w is different from hydrogen having either a trans- or cis-configuration at the cyclopropyl ring (i.e. wherein the B-A- and -NH-D groups are in trans or cis configuration), as well as compounds of formula I wherein R^w, R^x, R^y, R^z = H (i.e. a compound of formula Ia) having a cis configuration, since the cyclopropanation reaction used yields a mixture of cis/trans isomers, as represented by the wavy line in Scheme 7, which can be separated to yield the desired cis or trans compounds of the invention.



SCHEME 7: DCE (Dichloroethane)


5 Derivatives of formula (50) are subjected to cyclopropanation using ethyl diazoacetate and copper (I) chloride, as catalyst, in a suitable solvent as chloroform affording to a 1:1 mixtures of *cis*- and *trans*- derivatives of formula (51). Alternatively, the copper catalyst disclosed in Scheme 6 can be used. The diastereomers can be separated at this point either chromatographically or, after saponification (performed under basic conditions using for example NaOH in a suitable solvent such as MeOH), by recrystallisation of the corresponding carboxylic acids of formula (52). Curtius degradation to Boc-protected cyclopropylamines of formula (54) can be performed, first by using ethyl chloroformate and a base, as for example, triethylamine in a suitable solvent, as for example, acetone, and subsequent reaction with sodium azide in water leading to cyclopropanecarbonyl azide derivatives of formula (53). Reaction with *tert*-butanol results in the formation of Boc-protected cyclopropylamines of formula (54). Deprotection of the Boc-group in acidic conditions, for example using HCl in

10

1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent leads to the formation of the cyclopropanamine derivatives of formula (55). Reductive alkylation with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under the same conditions disclosed in Scheme 2 leads to the formation of cyclopropylamino derivatives of the invention, designated as compounds of formula (56a), (56b) or (56c) in the above scheme, which corresponds to a compound of formula I, wherein R^w is H, fluoro or C₁₋₄ alkyl and R^x, R^y, R^z = H. In case the ketone of formula (13a) or the aldehydes of formula (13b) or (13c) contain a protected amino group, for example a Boc-protected amine (Boc: *tert*-butoxycarbonyl), an additional deprotection reaction step will be required to render a compound (56a), (56b) or (56c), which can be performed in acidic conditions, for example using HCl in 1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent.

Compounds of formula (50), ketones of formula (13a) and the aldehydes of formula (13b) or (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

The compounds of Formula I wherein B = -L¹-E, and R^w, R^x, R^y, R^z = H and L¹ = bond can be synthesized, for example, by the general route described in Scheme 8.



Scheme 8: ACN (Acetonitrile), DCE (Dichloroethane), DMF (N,N-dimethylformamide), DMSO (Dimethyl sulfoxide).

5 *Tert*-butyl (*trans*)-cyclopropylcarbamate derivatives of formula (6, where R¹ = Br), obtained following the same procedure disclosed in Scheme 4, are converted into the (*trans*)-cyclopropanamine derivatives of formula (60) by reaction with boronic acid or ester derivatives of formula (59) using a suitable solvent such as acetonitrile and water, a base, such as for example potassium carbonate, and a palladium catalyst such as tetrakis(triphenylphosphine)palladium (0). Deprotection of the Boc-group in acidic conditions, for example using HCl 1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent leads to the formation of the (*trans*)-cyclopropanamine derivatives of formula (61). Reductive alkylation with ketones of formula (13a) or aldehydes of formula (13b) or (13c) under the same conditions disclosed in Scheme 2 leads to the formation of (*trans*)-cyclopropylamino derivatives of formula (62a), (62b) or (62c) which correspond to compounds of formula I wherein B = -L¹-E, R^w, R^x, R^y, R^z = H and L¹ = bond. When the ketones of formula (13a) or aldehydes of formula (13b) or (13c) contain a protected amino group, for example a Boc-protected amine (Boc: *tert*-butoxycarbonyl), an additional deprotection reaction step will be required to render a compound (62a), (62b) or (62c). The deprotection can be performed in acidic conditions, for example using HCl 1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent.

10

15

20 Alternatively, (*trans*)-cyclopropylamino derivatives of formula (62a), (62b) or (62c) can be synthesized by reaction of Boc-protected derivatives of formula (39a), (39b) or (39c) with boronic acid or ester derivatives of formula (59) using a suitable solvent such as acetonitrile and water, a base, such as for example potassium carbonate, and a palladium catalyst such as tetrakis(triphenylphosphine)palladium (0) affording (*trans*)-cyclopropanamine derivatives of formula (63a), (63b) or (63c). Deprotection of the Boc-group in acidic conditions, for example using HCl 1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent leads to the formation of the (*trans*)-cyclopropylamino derivatives of formula (62a), (62b) or (62c), which correspond to a compound of formula I, wherein B = -L¹-E, R^w, R^x, R^y, R^z = H and L¹ = bond.

25

30 Alternatively, alkylation of *tert*-butyl (*trans*)-cyclopropylcarbamate derivatives of formula (6, where R¹ = Br) with alkyl halides of formula (7) under basic conditions using for example NaH in a suitable solvent such as DMF leads to the formation of *tert*-butyl (*trans*)-cyclopropylcarbamate derivatives of formula (34). Reaction with boronic acid or ester derivatives of formula (59) using a suitable solvent such as acetonitrile and water, a base, such as for example potassium carbonate, and a palladium catalyst such as tetrakis(triphenylphosphine)palladium (0) leads to the formation of compounds of formula (57). Deprotection of the Boc-group in acidic conditions, for example using HCl 1,4-dioxane in a suitable solvent such as 1,4-dioxane or HCl in Et₂O using Et₂O as solvent leads to the formation of the (*trans*)-cyclopropanamine derivatives of

35

formula (58), which also correspond to a compound of formula I, wherein B = -L¹-E, R^w, R^x, R^y, R^z = H and L¹ = bond.

5 Aldehydes of formula (1), boronic acid or ester derivatives of formula (59) and ketones of formula (13a) or aldehydes of formula (13b) or (13c) are commercially available or can be prepared using well known synthetic procedures starting from readily available starting materials.

10 Furthermore, some compounds of the invention can be obtained from other compounds of formula I by appropriate interconversion reactions of functional groups present in a compound of formula I in one or several steps, using well known reactions in organic synthesis under standard experimental conditions. Said transformations can be carried out upon R¹, R² and/or R³ and may include, for example, the substitution of a primary or secondary amine or of an alcohol by treatment with an alkylating agent, the conversion of an amine into an amide, sulfonamide, carbamate or urea, the palladium-catalyzed cross-coupling of aryl halides with diboranes, boronic acids or esters or with amines, etc. Such interconversion reactions can be performed upon a 15 compound of formula I as well as upon any suitable synthetic intermediate described in the above Schemes.

20 The salts of a compound of formula I can be obtained during the final isolation and purification of the compounds of the invention or can be prepared by treating a compound of formula I with a sufficient amount of the desired acid (or base) to give the salt in a conventional manner.

25 In the above schemes 2 to 5 the cyclopropanation reaction under the conditions disclosed always leads to a racemic mixture of the trans-isomers of compounds (3), (11), (18) and (25). If the synthetic procedures are continued using the trans racemic mixture thus obtained, the corresponding compounds of formula I are obtained as mixtures of trans-isomers. Likewise, in scheme 6 and 7 the cyclopropanation reaction under the conditions disclosed leads to a mixture of cis/trans isomers of compounds (45) and (51). If the synthetic procedure is continued using said isomer mixture, the corresponding compounds of formula I are obtained as mixtures of cis/trans isomers. As used herein, cis and trans refers to the disposition of groups -A-B versus -NH-L²-D on the cyclopropyl ring.

30 Where the processes for the preparation of the compounds of the invention give rise to mixtures of stereoisomers, individual stereoisomers of a compound of formula I can be obtained by separation from a compound of formula I obtained as a mixture of stereoisomers, using well known methods such as formation of diastereomeric pairs by salt formation with an optically active acid followed by fractional crystallization and regeneration of the free base, or by chiral preparative chromatography. Alternatively, it is possible to obtain 35 optically pure or enantiomerically enriched synthetic intermediates, which can then be used as such in subsequent steps, at various stages of the synthetic procedures described above, using any known method for

chiral resolution. Preferably, the chiral separation is performed upon trans-cyclopropylamines of formula (12), (26), (32), (48), (55) and (61). Separation can also be performed at other stages of the procedure, for example upon a compound of formula (45). A suitable method to obtain the enantiomers of the trans cyclopropylamines (12), (26), (32), (48), (55) and (61) comprises contacting a trans-substituted cyclopropylamine with a chiral 5 recrystallization agent in a solvent (particularly under conditions that are sufficient for the crystallization of the salt of the chiral recrystallization agent and the trans substituted cyclopropylamine); and isolating the crystallized salt of the chiral recrystallization agent and the trans substituted cyclopropylamine, thereby preparing an enantiomer of a trans N-substituted cyclopropylamine. A suitable chiral recrystallization agent is S (+) mandelic acid, D (-) tartaric acid, L (+) tartaric acid, L (-) di-p-toluoyl tartaric acid, or R (-) mandelic acid. 10 Suitable solvents are tetrahydrofuran, ethanol or mixtures thereof with H₂O.

Alternatively, it is possible for a person skilled in the art to obtain optically pure or enantiomerically enriched final compounds (or synthetic intermediates) by using chiral chromatography.


Examples

15 Unless stated otherwise, in the compounds of all Examples of the present specification the stereochemical configuration is defined by the chemical name indicated for the respective compound, even though the drawn structure may represent a more specific configuration. Nevertheless, the invention relates to all stereoisomers of the compounds described and defined herein. Accordingly, the invention encompasses the compounds 20 described in the Examples as defined by their chemical names and, in addition thereto, also the corresponding compounds having the absolute configuration shown in the respective drawn structures.

The following abbreviations have been used:


ACN: acetonitrile, AcOH: acetic acid, aq: aqueous, Boc: tert-butyloxycarbonyl, (Boc)₂O: di-tert-butyl 25 dicarbonate, brm: broad multiplet, brs: broad singlet, Cu(acac)₂: copper(II) acetylacetone, d: doublet, DCE: 1,2-dichloroethane, DCM: dichloromethane, DMF: N,N-dimethylformamide, DMSO: dimethylsulfoxide, DPPA: 30 diphenylphosphoryl azide, Et₂O: diethyl ether, EtOAc: ethyl acetate, HPLC: high performance liquid chromatography, m: multiplet, MEM: methoxy methyl ether, MeOH: methanol, NBS: N-bromosuccinimide, NMR: nuclear magnetic resonance, Pd₂(dba)₃: tris(dibenzylideneacetone)dipalladium(0), Pet ether: petroleum ether, q: quadruplet, Rf: retention factor, RT: room temperature, s: singlet, sat.: saturated, t: triplet, TEA: triethylamine, THF: tetrahydrofuran, TLC: thin layer chromatography, vbrs: very broad singlet, Xantphos: 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene.

Intermediate A: 1-(benzyloxy)-4-[(trans)-2-nitrocyclopropyl]benzene



Trimethylsulfoxonium iodide (0.62 g, 2.82 mmol) was added in portions to a solution of *t*-BuOK (0.32 g, 2.82 mmol) in dry DMSO (5 mL). After 10 min a solution of 1-(benzyloxy)-4-[(E)-2-nitrovinyl]benzene (0.60 g, 2.35 mmol) in DMSO (5 mL) was transferred via canula and the mixture was stirred at room temperature for 6 h. The reaction was poured over water (10 mL) and extracted with Et₂O (3x10 mL); the organic layers were washed with brine (2x15 mL), dried over anhydrous Na₂SO₄ and filtered. After removal of the solvent, the residual orange oil was purified by column chromatography on silica gel (5% EtOAc/hexanes) affording 0.16 g of 1-(benzyloxy)-4-[(trans)-2-nitrocyclopropyl]benzene [R_f= 0.5 (20% EtOAc/hexanes), white solid, 26% yield].


10 **Intermediate B: Trans-2-[4-(benzyloxy)phenyl]cyclopropanamine**



Zn dust (1.97 g, 30 mol) was added in small portions, over a period of 30 min, to a vigorously stirred solution of 1-(benzyloxy)-4-[(trans)-2-nitrocyclopropyl]benzene (Intermediate A, 0.81 g, 3.0 mmol) in *i*-PrOH (25 mL) and HCl (11 mL of aqueous solution 2.7 N, 30 mmol). After 17 h the mixture was filtered through a pad of celite, that was washed with 10 mL of methanol. The filtrate was concentrated and 10 mL of water were added, washing with CH₂Cl₂ (3x15 mL). The organic layers were dried over anhydrous Na₂SO₄ and filtered. After removal of the solvent, the crude product was purified by column chromatography on silica gel (10% MeOH/CH₂Cl₂) affording 0.50 g of (trans)-2-[4-(benzyloxy)phenyl]cyclopropanamine [R_f= 0.2 (10% MeOH/CH₂Cl₂), white solid, 70% yield].

20 ¹H-NMR (MeOH, 250 MHz, δ): 7.45-7.27 (m, 5H, ArH); 6.96 (d, J = 8.5 Hz, 2H, ArH); 6.86 (d, J = 8.5 Hz, 2H, ArH); 5.03 (s, 2H, CH₂); 2.41-2.34 (m, 1H, CH); 1.86-1.76 (m, 1H, CH); 0.98-0.85 (m, 2H, CH₂).


Intermediate C: 4-(benzyloxy)benzaldehyde



Potassium Carbonate (678 g, 4.91 mol) was added to a solution of 4-hydroxybenzaldehyde (200 g, 1.63 mol) in DMF (2 L) followed to the addition of benzyl bromide (214 mL, 1.80 mol) at 0° C and stirred for 18 h at RT. After completion, the reaction mixture was poured into ice water (3 L), filtered the solid and dried to get 4-(benzyloxy)benzaldehyde (230 g, 66 %).

5


Intermediate D: (E)-ethyl 3-(4-(benzyloxy)phenyl)acrylate



Triethyl phosphonoacetate (259 mL, 1.3 mol) was added slowly dropwise to a solution of Potassium-*tert*-butoxide (145 g, 1.29 mol) in dry THF (2 L) at -5 °C and stirred for 30-45 mins. Then a solution of 4-(benzyloxy)benzaldehyde (Intermediate C, 230 g, 1.08 mol) in dry THF (1.5 L) was added slowly dropwise at -10 °C over a period of 15 mins and stirred for 30 mins. After completion, the reaction mixture was poured into ice water (1 L) and extracted with EtOAc (2 x 1.5 L). The combined organic extracts were washed with sat NaHCO₃ solution (1 L), water (1 L), brine (1 L), dried over *anhydrous* Na₂SO₄, filtered and evaporated to get crude (E)-ethyl 3-(4-(benzyloxy)phenyl)acrylate (290 g, 95 %). The crude was carried to next step without further purification.

15

Intermediate E: (Trans)-ethyl 2-(4-(benzyloxy)phenyl)cyclopropanecarboxylate




Trimethyl sulfoxonium iodide (224 g, 1.02 mol) was added portion wise to a suspension of NaH (40.8 g, 1.02 mol) in dry DMSO (2 L) at RT over a period of 20 min and stirred for 1 h till the formation of a clear solution. A solution of (E)-ethyl 3-(4-(benzyloxy)phenyl)acrylate (Intermediate D, 240 g, 0.85 mol) in dry DMSO (2 L) was added dropwise and stirred at RT for 30 mins. After completion, the reaction mixture was poured into ice water (2 L), extracted with EtOAc (2 x 1 L). Combined organic extracts were washed with ice water (1 L), brine (1 L), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (Trans)-ethyl 2-(4-(benzyloxy)phenyl)cyclopropanecarboxylate (142 g, 58.6 %) as an off white solid. The crude was carried to next step without further purification.

20


25

Intermediate F: (Trans)-2-(4-(benzyloxy)phenyl)cyclopropanecarboxylic acid




4N NaOH solution (4 L) was added to a solution of (trans)-ethyl 2-(4-(benzyloxy)phenyl)cyclopropanecarboxylate (Intermediate E, 250 g, 0.844 mol) in Methanol (1.2 L) at 0 °C and stirred at RT for 4 h. After completion, the solvent was evaporated, the residue was diluted with water (1 L), acidified with 4 N HCl solution, extracted with EtOAc (2 x 2 L). Combined organic extracts were washed with water (1 L), brine (1 L), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (trans)-2-(4-(benzyloxy)phenyl)cyclopropanecarboxylic acid (190 g, 84 %) as off white solid. The crude was carried to next step without further purification.

Intermediate G: (Trans)-2-(4-(benzyloxy)phenyl)cyclopropanecarbonyl azide




Ethyl chloroformate (143 mL, 1.48 mol) was added to a solution of (trans)- 2-(4-(benzyloxy) phenyl) cyclopropanecarboxylic acid (Intermediate F, 190 g, 0.70 mol), Triethyl amine (229 mL, 1.63 mol) in acetone (2.8 L) at -20 °C and stirred for 1 h, then a solution of NaN₃ (138 g, 2.1 mol) in water (200 mL) was added and stirred at RT for 30 mins. After completion, the solvent was evaporated, residue was dissolved in EtOAc (2 L), washed with water (2 L), brine (1 L), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (trans)-2-(4-(benzyloxy)phenyl)cyclopropanecarbonyl azide (178 g, 85.9 %).

Intermediate H: Tert-butyl ((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)carbamate




A solution of (trans)-2-(4-(benzyloxy)phenyl)cyclopropanecarbonyl azide (Intermediate G, 178 g, 0.64 mol) in *tert*-butanol (2.6 L) was heated at 90 °C for 16 h. After completion, the solvent was evaporated and the crude residue was purified by column chromatography by using (SiO₂) EtOAc: Pet ether (4: 96) to get tert-butyl ((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)carbamate (78 g, 37.8 %) as off-white solid.

Intermediate I: (E)-ethyl 3-(6-bromopyridin-3-yl)acrylate




5 Triethyl phosphonoacetate (26.6g, 118.8 mmol) was added slowly dropwise to a mixture of Potassium-*tert*-butoxide (14.5g, 129.6 mmol) in dry THF (200 mL) at -5 °C, stirred for 20 min and then a solution of 6-bromopyridine-3-carboxaldehyde (20 g, 108 mmol) in dry THF (100 mL) was added slowly dropwise at -5 °C and stirred for 30 min. After completion, the reaction mixture was poured into ice water (350 mL) and extracted with EtOAc (2 x 300 mL). The combined organic extracts were washed with saturated NaHCO₃ solution (250 mL), water (250 mL) and brine (250 mL) and dried over *anhydrous* Na₂SO₄, filtered and evaporated to get (E)-ethyl 3-(6-bromopyridin-3-yl) acrylate (20 g, 72.9 %) as brown color liquid. This is carried to next step without further purification.

10 **Intermediate J: (Trans)-ethyl-2-(6-bromopyridin-3-yl)cyclopropanecarboxylate**




15 Trimethyl sulfoxonium iodide (20.8g, 94.7 mmol) was added in small portions to a suspension of sodium hydride (4g, 170.6 mmol) in dry DMSO (400 mL) at rt., stirred for 1 h until clear solution was obtained. A solution of (E)-ethyl 3-(6-bromopyridin-3-yl) acrylate (Intermediate I, 20 g, 78.7 mmol) in dry DMSO (20 mL) was added and stirred for 4 h. After completion, the reaction mixture was poured into ice water (700 mL), extracted with EtOAc (2 x 350 mL). The combined organic extracts were washed with water (250 mL), brine (250 mL) and dried over *anhydrous* Na₂SO₄, filtered and evaporated to give (trans)-ethyl-2-(6-bromopyridin-3-yl)cyclopropanecarboxylate (10g, 47 %) as brown liquid.

20 **Intermediate K: (Trans)-2-(6-bromopyridin-3-yl)cyclopropanecarboxylic acid hydrochloride**




25 NaOH 4N solution (60 mL) was added to a solution of (trans)-ethyl-2-(6-bromopyridin-3-yl)cyclopropanecarboxylate (Intermediate J, 10 g, 37.1 mmol) in methanol (100 mL) and the reaction mixture was stirred at RT for 4 h. After completion, the solvent was evaporated and the residue was diluted with ice water (250 mL) and acidified with 4 N HCl solution, the aqueous layer was extracted with EtOAc (2 x 350 mL). The combined organic extracts were washed with water (250 mL), brine (250 mL) and dried over *anhydrous* Na₂SO₄, filtered and evaporated to give (trans)-2-(6-bromopyridin-3-yl)cyclopropanecarboxylic acid hydrochloride (5g, 55.8 %) as a light brown color solid.

30 **Intermediate L: (Trans)-2-(6-bromopyridin-3-yl)cyclopropanecarbonyl azide**



Ethyl chloroformate (5.8 mL, 62 mmol) was added to a solution of (trans)-2-(6-bromopyridin-3-yl)cyclopropanecarboxylic acid hydrochloride (Intermediate K, 5 g, 20.7 mmol) and Et₃N (14.2 mL, 103.7 mmol) in Acetone (100 mL) at -5 °C, then reaction mixture was stirred at -5 °C for 1 h, then a solution of NaN₃ (2.7 g, 41.4 mmol) in water (10 mL) was added and stirred for 30 mins at RT. After completion the solvent was evaporated under vacuum. The crude residue was dissolved in ethyl acetate (200 mL), washed with water (80 mL), brine (80 mL), dried over anhydrous Na₂SO₄, filtered and evaporated to get (trans)-2-(6-bromopyridin-3-yl)cyclopropanecarbonyl azide (2.5 g, 45.5 %) as a brown color gummy liquid.


10 **Intermediate M: tert-butyl (trans)-2-(6-bromopyridin-3-yl)cyclopropylcarbamate**



A solution of (trans)-2-(6-bromopyridin-3-yl)cyclopropanecarbonyl azide (Intermediate L, 2.5 g, 9.36 mmol) in *tert*-butanol (80 mL) was heated at 90 °C for 16 h. After completion, the solvent was evaporated under vacuum and the residue was taken in water (100 mL) and extracted with EtOAc (2 x 100 mL). The combined organic extracts were washed with water (100 mL), brine (100 mL) and dried over anhydrous Na₂SO₄, filtered and evaporated. The crude residue was purified by flash column chromatography (SiO₂) by eluting with EtOAc: Hexane (2: 8) to get tert-butyl (trans)-2-(6-bromopyridin-3-yl)cyclopropylcarbamate (1.1 g, 37.5 %) as a light yellow solid.

20 ¹H-NMR (CDCl₃) δ (ppm): 1.16 (q, 1H), 1.23 (quin, 1H), 1.45 (s, 9H), 2.01 (m, 1H), 2.69 (m, 1H), 4.88 (br, 1H), 7.36 (s, 2H), 8.20 (s, 1H).

Intermediate N: (E)-ethyl 3-(4-bromophenyl)acrylate




A solution of triethyl phosphonoacetate (13.1 g, 0.0589 mol) was added slowly (dropwise) to a solution of Potassium-*tert*-butoxide (6.59 g, 0.0589 mol), in dry THF (150 mL) at -5 °C, stirred for 30-45 mins at the same temperature, then a solution of 4-Bromo benzaldehyde (10 g, 0.054 mol), in dry THF (50 mL) was slowly added dropwise at -5 °C over a period of 15 mins, stirred the reaction mixture for 30 mins at the same temperature.

After completion of reaction by TLC, the reaction mixture was poured into ice water (300 mL), extracted with EtOAc (2 x 200 mL). The combined organic extracts were washed with sat NaHCO₃ solution (200 mL), water (200 mL), brine (200 mL) and dried over anhydrous Na₂SO₄, filtered and evaporated to get crude (E)-ethyl 3-(4-bromophenyl) acrylate (10 g, 72 %) as pale green liquid. This is carried to next step without further purification.

5


Intermediate O: (Trans)-ethyl 2-(4-bromophenyl)cyclopropanecarboxylate



Trimethyl sulfoxonium iodide (5.19 g, 0.0236 mol) was added slowly in small portions over a period of 20 min. to a suspension of sodium hydride (0.44 g, 0.0236 mol) in dry DMSO (80 mL) at rt, stirred for 1 h, till the formation of clear solution. Then a solution of (E)-ethyl 3-(4-bromophenyl) acrylate (Intermediate N, 5 g, 0.01968), in dry DMSO (20 mL) was added slowly dropwise, stirred at rt for 30 mins. After completion of reaction, checked by TLC, the reaction mixture was poured into ice water (200 mL), extracted with EtOAc (2 x 150 mL). Combined organic extracts were washed with ice water (2 x 150 mL), brine (150 mL), dried over anhydrous Na₂SO₄, filtered and evaporated to get (trans)-ethyl 2-(4-bromophenyl)cyclopropanecarboxylate (4 g, 75.9 %) as a green liquid. The crude is carried to next step without further purification.

10


Intermediate P: (Trans)-2-(4-bromophenyl)cyclopropanecarboxylic acid



NaOH 4N (20 mL) was added to a solution of (trans)-ethyl 2-(4-bromophenyl)cyclopropanecarboxylate (Intermediate O, 4 g, 0.0149 mol), in Methanol (40 mL) and stirred at rt for 2 h. After completion of reaction, checked by TLC, the solvent was evaporated and the residue was diluted with water (50 mL), acidified with HCl 4 N solution, the solid formed was filtered and dried to get (trans)-2-(4-bromophenyl)cyclopropanecarboxylic acid (2.59 g, 72 %), as a white solid.

20


Intermediate Q: (Trans)-2-(4-bromophenyl)cyclopropanecarbonyl azide



Ethyl chloroformate (1.9 mL) was added to a solution of (trans)-2-(4-bromophenyl) cyclopropanecarboxylic acid (Intermediate P, 4 g, 0.0165 mol) and Et₃N (2.51 mL, 0.0199 mol) in acetone (60 mL) at -20 °C, stirred at same

temperature for 1 h, then a solution of NaN_3 (1.3 g, 0.0199 mol) in water (5 mL), was added and stirred for 30 mins at rt. After completion of reaction, checked by TLC, the solvent was evaporated and crude residue was dissolved in ethyl acetate (100 mL), washed with water (40 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated to get (*trans*)-2-(4-bromophenyl)cyclopropanecarbonyl azide (4 g). The crude residue is carried to next step without further purification.


5 **Intermediate R: tert-butyl (*trans*)-2-(4-bromophenyl)cyclopropylcarbamate**



10 A solution of (*trans*)-2-(4-bromophenyl) cyclopropanecarbonyl azide (Intermediate Q, 4 g) in tert-Butanol (40 mL) was heated at 90 °C for 16 h. After completion of reaction, checked by TLC, the solvent was evaporated residue was poured into water (50 mL), extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated. The crude residue was purified by column chromatography (SiO_2) by eluting with EtOAc : Petroleum ether (2: 98), to get tert-butyl (*trans*)-2-(4-bromophenyl)cyclopropylcarbamate (2.5 g, 48 % overall 2 steps) as a white solid.


15 $^1\text{H-NMR}$ (CDCl_3 , 250 MHz) δ : 1.07-1.19 (m, 2H), 1.44 (s, 9H); 2.05-1.94 (m, 1H); 2.72-2.62 (m, 1H); 4.85 (br, 1H,); 7.09-6.96 (m, 2H); 7.44-7.33 (m, 2H).

20 **Intermediate S: (*E*)-ethyl 3-(pyridin-3-yl)acrylate**



25 A solution of triethyl phosphonoacetate (66.75 mL, 336.44 mmol) was added dropwise to a solution of Potassium-*tert*-butoxide (37.7 g, 280.37 mmol) in dry THF (300 mL) at -5 °C over a period of 10 mins and stirred at 0 °C for 30 mins. Then a solution of nicotinaldehyde (30 g, 280.37 mmol) in dry THF (50 mL) was added dropwise at 0 °C over a period of 15 mins and stirred at RT for 2 h. After completion, the reaction mixture was poured into ice water (150 ml) and extracted with EtOAc (2 x 300 mL). The combined extracts were washed with sat NaHCO_3 solution (200 mL), water (200 mL), brine (200 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated to afford crude liquid (*E*)-ethyl 3-(pyridin-3-yl) acrylate (42 g, 84.67 %). The crude was carried to next step without further purification.

Intermediate T: (Trans)-ethyl 2-(pyridin-3-yl)cyclopropanecarboxylate




Trimethyl sulfoxonium iodide (14.90 g, 67.76 mmol) was added portion wise to a suspension of NaH (2.71 g, 67.76 mmol) in dry DMSO (100 mL) at RT over a period of 20 min. and stirred for 1 h till the formation of a clear solution. A solution of (E)-ethyl 3-(pyridin-3-yl) acrylate (Intermediate S, 10 g, 56.47 mmol) in dry DMSO (50 mL) was added dropwise and stirred at RT for 20 min. After completion, the reaction mixture was poured into ice water (200 mL), extracted with EtOAc (2 x 200 mL). Combined organic extracts were washed with ice water (150 mL), brine (150 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (*trans*)-ethyl 2-(pyridin-3-yl)cyclopropanecarboxylate (4 g, 37.07 %) as pale brown liquid. The crude was carried to next step without further purification.

5

10

Intermediate U: (*Trans*)-2-(pyridin-3-yl)cyclopropanecarboxylic acid




15

20

A solution of NaOH (7.116 g in 45 mL of H₂O, 177.92 mmol) was added to a solution of (*trans*)-ethyl 2-(pyridin-3-yl) cyclopropanecarboxylate (Intermediate T, 17 g, 88.96 mmol) in Methanol (170 mL) at 0 °C and stirred at RT for 16 h. After completion, the solvent was evaporated, the residue was diluted with water (50 mL), neutralized with Acetic acid and extracted with EtOAc (4 x 100 mL). The combined extracts were washed with water (100 mL), brine (100 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (*trans*)-2-(pyridin-3-yl)cyclopropanecarboxylic acid (9 g, 62.06 %) as off white solid. The crude was carried to next step without further purification.


25

Intermediate V: (*Trans*)-2-(pyridin-3-yl)cyclopropanecarbonyl azide




25


Ethyl chloroformate (6.89 mL, 71.15 mmol) was added to a solution of (*trans*)-2-(pyridin-3-yl)cyclopropanecarboxylic acid (Intermediate U, 9 g, 55.194 mmol) and triethyl amine (11.03 mL, 82.79 mmol) in acetone (90 mL) at -20 °C and stirred for 1 h, then a solution of NaN₃ (5.38 g, 82.79 mmol) in water (25 mL) was added and stirred at RT for 30 mins. After completion, the solvent was evaporated, residue was dissolved in EtOAc (100 mL), washed with water (2 x 50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (*trans*)-2-(pyridin-3-yl)cyclopropanecarbonyl azide (8.4 g, 81 %). The crude was carried to next step without further purification.

Intermediate W: *tert*-butyl ((*trans*)-2-(pyridin-3-yl)cyclopropyl)carbamate


A solution of (*trans*)-2-(pyridin-3-yl)cyclopropanecarbonyl azide (Intermediate V, 8.4 g, 44.66 mmol) in *tert*-butanol (85 mL) was heated at 90 °C for 16 h. After completion, the solvent was evaporated and crude residue was purified by column chromatography (SiO₂) using EtOAc: Petroleum ether (25: 75) to afford *tert*-butyl (*trans*)-2-(pyridin-3-yl) cyclopropylcarbamate (3.9 g, 37.32 %) as colourless liquid.

Intermediate X: (*Trans*)-2-(pyridin-3-yl)cyclopropanamine hydrochloride

HCl in Dioxane (10 mL) was added to a solution of *tert*-butyl (*trans*)-2-(pyridin-3-yl)cyclopropylcarbamate (Intermediate W, 2 g, 8.541 mmol) in 1,4-dioxane (10 mL) at 0 °C and stirred at RT for 12 h. After completion, the solvent was evaporated and the residue was triturated with diethyl ether (20 mL) followed by hexane (20 mL) to get (*trans*)-2-(pyridin-3-yl)cyclopropanamine hydrochloride (1.2 g, 82.7 %).


Intermediate Y: (*E*)-ethyl 3-(thiazol-5-yl)acrylate

A solution of triethyl phosphonoacetate (11.88 g, 53.03 mmol) was added dropwise to a solution of Potassium-*tert*-butoxide (5.94 g, 53.03 mmol) in dry THF (100 mL) at -5 °C and stirred for 30 mins. A solution of thiazole-5-carbaldehyde (5 g, 44.19 mmol) in dry THF (25 mL) was then added dropwise at -5 °C over a period of 15 mins and stirred for 30 mins. After completion, the reaction mixture was poured into ice water (150 mL), extracted with EtOAc (2 x 100 mL). The combined extracts were washed with sat NaHCO₃ solution (100 mL), water (100 mL), brine (100 mL), dried over anhydrous Na₂SO₄, filtered and evaporated to afford crude (*E*)-ethyl 3-(thiazol-5-yl)acrylate (10 g, 82.3%) as a white solid. The crude was carried to next step without further purification

Intermediate Z: (*Trans*)-ethyl 2-(thiazol-5-yl)cyclopropanecarboxylate


Trimethyl sulfoxonium iodide (14.40 g, 65.49 mmol) was added portionwise to a suspension of NaH (2.61 g, 108.75 mmol) in dry DMSO (200 mL) at RT over a period of 20 min and stirred for 1 h till the formation of clear solution. A solution of (E)-ethyl 3-(thiazol-5-yl)acrylate (Intermediate Y, 10 g, 54.57 mmol) in dry DMSO (50 mL) was then added dropwise and stirred at RT for 30 mins. After completion, the reaction mixture was poured into ice water (100 mL) and extracted with EtOAc (2 x 100 mL). Combined organic extracts were washed with water (2 x 50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford *trans*-ethyl 2-(thiazol-5-yl)cyclopropanecarboxylate (8 g, 61.9 %) as a reddish brown liquid. The crude was carried to next step without further purification.

10 **Intermediate AA: (Trans)-2-(thiazol-5-yl)cyclopropanecarboxylic acid**




A 4N NaOH solution (40 mL) was added to a solution of *trans*-ethyl 2-(thiazol-5-yl)cyclopropanecarboxylate (Intermediate Z, 8 g, 40.55 mmol) in methanol (80 mL) and stirred at RT for 4 h. After completion, the solvent was evaporated, the residue was diluted with water (50 mL), acidified with Acetic acid and extracted with EtOAc (2 x 75mL). The combined extracts were washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford *trans*-2-(thiazol-5-yl)cyclopropanecarboxylic acid (4 g, 58.30 %). The crude was carried to next step without further purification.

15 **Intermediate AB: (Trans)-2-(thiazol-5-yl)cyclopropanecarbonyl azide**




20 Ethyl chloroformate (3.34 g, 30.76 mmol) was added to a solution of *trans*-2-(thiazol-5-yl)cyclopropanecarboxylic acid (Intermediate AA, 4 g, 26.3 mmol) and triethylamine (3.62 g, 35.50 mmol) in acetone (40 mL) at -20 °C, stirred at same temperature for 1 h. A solution of NaN₃ (2.84 g, 47.33 mmol) in water (10 mL) was then added and stirred at RT for 30 mins. After completion, the solvent was evaporated, the crude residue was dissolved in EtOAc (100 mL), washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford *trans*-2-(thiazol-5-yl)cyclopropanecarbonyl azide (3 g, 58.7 %) as brown liquid. The crude was carried to next step without further purification.

25 **Intermediate AC: tert-butyl ((trans)-2-(thiazol-5-yl)cyclopropyl)carbamate**




A solution of *trans*-2-(thiazol-5-yl)cyclopropanecarbonyl azide (Intermediate AB, 3 g, 15.44 mmol) in *tert*-butanol (60 mL) was heated at 90 °C for 16 h. After completion, the solvent was evaporated and the residue was taken in water (50 mL), extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude residue was purified by column chromatography (SiO₂) by using EtOAc: Petroleum ether (20:80) to get *tert*-butyl *trans*-2-(thiazol-5-yl)cyclopropylcarbamate (1.1 g, 29.64 %) as a pale yellow liquid.

Intermediate AD: (Trans)-2-(thiazol-5-yl)cyclopropanamine hydrochloride



HCl in dioxane (10 mL) was added to a solution of *tert*-butyl *trans*-2-(thiazol-5-yl)cyclopropylcarbamate (Intermediate AC, 1.1 g, 45.83 mmol) in dioxane (10 mL) at 15 °C and stirred at RT for 3 h. After completion, the solvent was evaporated, the residue was triturated with EtOAc to afford *trans*-2-(thiazol-5-yl)cyclopropanamine hydrochloride (600 mg, 74.8 %) as pale yellow solid.


Intermediate AE: *tert*-butyl ((*trans*)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)carbamate



A solution of *tert*-butyl (*trans*)-2-(6-bromopyridin-3-yl)cyclopropylcarbamate (Intermediate M, 100 mg, 0.32 mmol), potassium carbonate (132 mg, 0.96 mmol) and 3-trifluoromethylbenzeneboronic acid (72 mg, 0.38 mmol) in CH₃CN:H₂O (4: 1) (10 mL) was degassed for 30 mins. Tetrakis triphenylphosphine palladium (37mg, 0.032 mmol) was added and degassed for 10 mins and the reaction mixture was heated at reflux temperature for 2 h. After completion, the reaction mixture was poured in ice water (100 mL), extracted with ethyl acetate (5 x 40 mL). The combined extract was washed with water (70 mL), brine (70 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude residue was purified by column chromatography (SiO₂), by using EtOAc:Petroleum ether (1:9) to get *tert*-butyl (*trans*)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropylcarbamate (70 mg, 58.3%) as a white solid.

¹H-NMR (CDCl₃) δ (ppm): 1.26 (m, 2H), 1.46 (s, 9H), 2.10 (m, 1H), 2.78 (m, 1H), 4.86 (br, 1H), 7.55 (m, 2H), 7.65 (t, 2H), 8.14 (d, 1H), 8.24 (s, 1H), 8.54 (s, 1H). **MS (M+H):** 379.1.


Intermediate AF: (Trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropanamine dihydrochloride



5 HCl in diethyl ether (5 mL) was added to a solution of tert-butyl (trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropylcarbamate (Intermediate AE, 70 mg, 0.185mmol) in diethyl ether (10 mL) at 0 °C slowly dropwise over a period of 10 mins and then stirred for 2 h. After completion, the reaction mixture was filtered under inert atmosphere and washed with hexane (10 mL), EtOAC (5 mL), and dried under reduced pressure to get (trans)-
10 2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropanamine dihydrochloride (50 mg, 86.2%) as a pale yellow powder.

¹H-NMR (D₂O) δ (ppm): 1.52 (q, 1H), 1.63 (quin, 1H), 2.66 (m, 1H), 3.08 (m, 1H), 7.72 (t, 1H), 7.89 (d, 1H), 7.98 (d, 1H), 8.09 (s, 1H), 8.14 (d, 1H), 8.27 (d, 1H), 8.61 (s, 1H). **MS (M+H) :** 279.1.


15 **Intermediate AG: tert-butyl ((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate**



20 A solution of tert-butyl (trans)-2-(4-bromophenyl)cyclopropylcarbamate (Intermediate R, 1 g, 3.2 mmol), potassium carbonate (1.31 g, 9.6 mmol) and 3-(trifluoromethyl) phenylboronic acid (0.73 g, 3.8 mmol) in acetonitrile:water (4: 1) was degassed for 30 mins. Tetrakis triphenylphosphine palladium (36 mg, 0.032 mmol) was then added, degassed again for 10 mins and the reaction mixture was heated at reflux temperature for 5 h. After completion, the reaction mixture was poured in ice water (50 mL) and extracted with ethyl acetate (2 x 50 mL). Combined extracts were washed with water (70 mL), brine (70 mL), dried over anhydrous Na₂SO₄ and then filtered and evaporated. The crude residue was purified by column chromatography (SiO₂), by using EtOAc: Petroleum ether (2:8) to get tert-butyl ((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (0.8 g, 66 %) as a white solid.
25

Intermediate AH: (Trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine hydrochloride


150



HCl in diethyl ether (3 mL) was added slowly dropwise to a solution of tert-butyl ((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (Intermediate AG, 200 mg, 0.53 mmol) in diethyl ether (5 mL) at 10 °C over a period of 10 min and then stirred for 4 h. After completion, the solvent was evaporated and the residue was triturated with hexane (5 mL), diethyl ether (5 mL) and dried under reduced pressure to get (trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine hydrochloride (140 mg, 77.8 %) as a white solid.


5 **¹H-NMR (DMSO-d6) δ (ppm):** 1.27 (q, 1H), 1.46 (quin, 1H), 2.41 (m, 1H), 2.86 (m, 1H), 7.29 (d, 2H), 7.69 (m, 4H), 7.96 (m, 2H), 8.53 (s, 1H), 8.61 (br, 2H). **MS (M+H) :** 278.3

10 **Intermediate AI: (1R, 2S)-2-phenylcyclopropanamine**



L (+) tartaric acid (15.7 g, 0.105 mmol) was added to a solution of (trans)-2-phenylcyclopropanamine (14 g, 0.105 mol) in EtOH (150 mL) at 0 °C and stirred at RT for 1 h. After completion, solid was filtered and dried to afford tartarate salt (28 g, 94.2 %). The salt was taken in Isopropanol: water (3:1) (260 mL) and stirred at 70 °C for 2 h. The clear solution was allowed to cool to RT. The solid separated was collected by filtration, taken in water (100 mL), basified with NaHCO3 solution, and extracted with EtOAc (2 x 150 mL). The combined extracts were washed with water (150 mL), brine (150 mL), dried over *anhydrous* Na2SO4, filtered and evaporated to afford (1R, 2S)-2-phenylcyclopropanamine (4.7 g, 67.1 %).


20 **Intermediate AJ: (1S, 2R)-2-phenylcyclopropanamine**



D (-) tartaric acid (6.7 g, 45.1 mmol) was added to a solution of (1S, 2R)-2-phenylcyclopropanamine (crude of the reaction to get Intermediate AI)(6 g, 45.1 mmol) in EtOH (60 mL), at 0 °C and stirred at RT for 1 h. After completion, solid was filtered and dried to afford tartarate salt (12 g, 94.4 %). The salt was taken in Isopropanol: water (3: 1) (120 mL) and stirred at 70 °C for 2 h. The clear solution was allowed to cool to RT. The solid separated was collected by filtration, taken in water (75 mL), basified with NaHCO3 solution, and extracted with


EtOAc (2 x 75 mL). The combined extracts were washed with water (100 mL), brine (100 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated to afford (1*S*, 2*R*)-2-phenylcyclopropanamine (3 g, 50 %).

Intermediate AK: 4-((2-methoxyethoxy)methoxy)benzaldehyde




4-hydroxybenzaldehyde (50 g, 409 mmol) in THF (50 mL) was added dropwise and slowly over a period of 30 min to a suspension of sodium hydride (19.6 g, 817 mmol) in THF (750 mL) at 0 °C and stirred for 15 min, followed by addition of 1-(chloromethoxy)-2-methoxyethane (MEM chloride, 61.10 g, 490 mmol) at 0 °C. The reaction mixture was stirred at RT for 30 min and, after completion, poured into ice water (500 mL) and extracted with EtOAc (2 x 750 mL). The combined organic extracts were washed with ice water (500 mL), brine (500 mL), dried over *anhydrous* Na₂SO₄, filtered and concentrated affording 4-((2-methoxyethoxy)methoxy)benzaldehyde (52 g, 60 %) as a pale yellow liquid. The crude was used in the next step without further purification.

15 **Intermediate AL: (E)-ethyl 3-(4-((2-methoxyethoxy)methoxy)phenyl)acrylate**



20 Triethyl phosphonoacetate (66.53 g, 296.8 mmol) was added to a solution of potassium-*tert*-butoxide (33.3 g, 296.8 mmol) in dry THF (500 mL) at -5 °C and stirred for 30-45 min at the same temperature. Then, a solution of 4-((2-methoxyethoxy)methoxy)benzaldehyde (Intermediate AK, 52 g, 247 mmol) in dry THF (100 mL) was added dropwise over a period of 15 min at -5 °C and stirred for 30 min at the same temperature. After completion, the reaction mixture was poured into ice water (600 mL) and extracted with EtOAc (2 x 600 mL). The combined extracts were washed with saturated aq. NaHCO₃ (300 mL), water (300 mL), brine (300 mL) and dried over *anhydrous* Na₂SO₄, filtered and concentrated yielding crude (E)-ethyl 3-(4-((2-methoxyethoxy)methoxy)phenyl)acrylate (58 g, 83 %) as a thick viscous liquid. The crude product was used in the next step without further purification.


25 **Intermediate AM: (*Trans*)-ethyl 2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylate**



30 Trimethyl sulfoxonium iodide (53.7 g, 244 mmol) was added portionwise over a period of 20 min to a suspension of sodium hydride (9.58 g, 399 mmol) in dry DMSO (143 mL) and stirred at RT for 1 h, until a clear solution was formed. A solution of (E)-ethyl 3-(4-((2-methoxyethoxy)methoxy)phenyl)acrylate (Intermediate AL,


58 g (2 x 29 g, 207 mmol) in dry DMSO (20 mL) was added dropwise and stirred at RT for 30 min. After completion, the reaction mixture was poured into ice water (500 mL) and extracted with EtOAc (2 x 500 mL). The combined extracts were washed with ice water (2 x 250 mL), brine (250 mL), dried over *anhydrous* Na₂SO₄, filtered and concentrated yielding (trans)-ethyl 2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylate (35 g, 57 %) as a brown liquid. The crude product was used in the next step without further purification.

Intermediate AN: (Trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylic acid




10 4N NaOH solution (175 mL) was added to a solution of (trans)-ethyl 2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylate (Intermediate AM, 35 g, 118.9 mmol) in methanol (350 mL) and stirred at RT for 6 h. After completion, the solvent was evaporated and the residue was taken up in water (150 mL), acidified with 4N HCl and extracted with EtOAc (2 x 400 mL). The combined extracts were washed with ice water (200 mL), brine (200 mL), dried over *anhydrous* Na₂SO₄, filtered and concentrated to render (trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylic acid (30 g, 95 %) as a brown thick viscous liquid. The crude product was used in the next step without further purification.

Intermediate AO: (Trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarbonyl azide



20 Ethyl chloroformate (14.73 g, 135.73 mmol) was added to a solution of (trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarboxylic acid (Intermediate AN, 30 g, 112.6 mmol) and TEA (23.14 mL, 168.9 mmol) in acetone (300 mL) at -20 °C and stirred at same temperature for 1 h. Then, a solution of NaN₃ (14.64 g, 225.2 mmol) in water (15 mL) was added and stirred for 30 min at RT. After completion, the solvent was evaporated and the crude residue was dissolved in ethyl acetate (2 x 300 mL), washed with water (200 mL), brine (200 mL), dried over *anhydrous* Na₂SO₄, filtered and concentrated to yield (trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarbonyl azide (30.5 g, 93 %). The crude product was used in the next step without further purification.


Intermediate AP: *Tert*-butyl ((trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropyl)carbamate



30 A solution of (trans)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropanecarbonyl azide (Intermediate AO, 30.5 g, 104.7 mmol) in *tert*-butanol (305 mL) was heated at 90 °C for 16 h. After completion, the solvent was


evaporated and the residue was taken up in water (300 mL) and extracted with EtOAc (2 x 300 mL). The combined extracts were washed with water (200 mL), brine (200 mL), dried over *anhydrous* Na₂SO₄, filtered and concentrated. The crude residue was purified by column chromatography (SiO₂, EtOAc/petroleum ether 1:9) affording *tert*-butyl ((*trans*)-2-(4-((2-methoxyethoxy)methoxy)phenyl)cyclopropyl) carbamate (25 g, 71 %) as a white solid.

Intermediate AQ: 4-((*trans*)-2-aminocyclopropyl)phenol hydrochloride




HCl in 1,4-dioxane (125 mL) was added to a solution of *tert*-butyl ((*trans*)-2-((2-methoxyethoxy)methoxy)phenyl)cyclopropyl carbamate (Intermediate AP, 25 mg, 74.18 mmol) in 1,4-dioxane (250 mL) at 10 °C and stirred at RT for 16 h. After completion, the solvent was evaporated and the solid residue was triturated with Et₂O, filtered off and dried affording 4-((*trans*)-2-aminocyclopropyl)phenol hydrochloride (13 g, 95 %) as a brown solid.

15 Intermediate AR: Tert-butyl ((*trans*)-2-(4-hydroxyphenyl)cyclopropyl)carbamate



20 K_2CO_3 (20.36 g, 147.56 mmol) and $(\text{Boc})_2\text{O}$ (16.8 mL, 70.27 mmol) was added to a solution of 4-((*trans*)-2-aminocyclopropyl)phenol hydrochloride (Intermediate AQ, 13 g, 70.27 mmol) in 1,4- dioxane (78 mL) and water (195 mL) and stirred at RT for 16 h. After completion, the reaction mixture was poured into water (300 mL) and extracted with EtOAc (2 x 200 mL). The combined extracts were washed with water (75 mL), brine (75 mL), dried over *anhydrous* Na_2SO_4 , filtered and concentrated. The crude residue was purified by column chromatography (SiO_2 , EtOAc/petroleum ether 3:7) affording *tert*-butyl ((*trans*)-2-(4-hydroxyphenyl)cyclopropyl)carbamate (14 g, 80 %) as a brown thick viscous liquid.


25 Intermediate AS: *Tert*-butyl ((*trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)cyclopropyl)carbamate



2-fluorobenzyl bromide (5.83 mL, 48.13 mmol) was added to a solution of *tert*-butyl ((*trans*)-2-(4-hydroxyphenyl)cyclopropyl)carbamate (Intermediate AR, 10.0 g, 40.11 mmol) and K₂CO₃ (11.07 g, 80.22 mmol) in dry DMF (100 mL) at 0 °C and stirred at RT for 18 h. After completion, the reaction mixture was poured into ice water (350 mL) and extracted with EtOAc (2 x 200 mL). The combined extracts were washed with water (2 x 100 mL), brine (100 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude was purified by


column chromatography (SiO_2) using EtOAc: petroleum ether (1: 3) to afford *tert*-butyl ((*trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)cyclopropyl)carbamate (7.3 g, 51.01 %) as an off white solid.

5 **Intermediate AT: (*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)cyclopropanamine**




HCl in 1,4-dioxane (50 mL) was added dropwise to a solution of *tert*-butyl ((*trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)cyclopropyl)carbamate (Intermediate AS, 7.3 g, 20.16 mmol) in 1,4-dioxane (100 mL) at 0 °C and stirred at RT for 12 h. After completion, the solvent was evaporated and the residue was taken in water, basified with NaHCO_3 and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated to afford (*trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)cyclopropanamine (4.3 g, 81.90 %) as an off white solid.

15 **Intermediate AU: (*Trans*)-2-(4-bromophenyl)cyclopropanamine**



To a solution of *tert*-butyl *trans*-2-(4-bromophenyl)cyclopropylcarbamate (Intermediate R, 10 g, 32.05 mmol) in 1,4-dioxane (100 mL) at 10 °C was added HCl in dioxane (50 mL) and stirred at RT for 20 h. After completion, the solvent was evaporated and the residue was taken up in ice water, basified with saturated aq. NaHCO_3 and extracted with EtOAc (2 x 100 mL). The combined extracts were washed with water, brine, dried over *anhydrous* Na_2SO_4 , filtered and concentrated to afford (*trans*)-2-(4-bromophenyl)cyclopropanamine (6.2 g, 91.5%). The crude was used in the next step without further purification.


25 **Intermediate AV: (*Tert*-butyl 4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)methyl)piperidine-1-carboxylate**



Tert-butyl 4-formylpiperidine-1-carboxylate (602 mg, 2.83 mmol) and AcOH (169.8 mg, 2.83 mmol) were added to a solution of *trans* 2-(4-bromophenyl)cyclopropanamine (Intermediate AU, 600 mg, 2.83 mmol) in DCE (12 mL) and stirred for 5 min.. The reaction mixture was then cooled to 0 °C and sodium triacetoxy borohydride (1.07 g, 5.09 mmol) was added and stirred at RT for 5 h. After completion, the reaction mixture was poured into ice water and extracted with DCM (2 x 50 mL). The combined extracts were washed with water (50 mL), brine (50 mL) and dried over *anhydrous* Na_2SO_4 , filtered and evaporated. The crude was purified by column


chromatography (SiO_2) using EtOAc: petroleum ether (2: 8) to afford *tert*-butyl 4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)methyl)piperidine-1-carboxylate (630 mg, 54.45 %) as gummy liquid.

5 **Intermediate** AW: **(*Tert*-butyl 4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)(*tert*-butoxycarbonyl)amino)methyl)piperidine-1-carboxylate**




NaOH (246 mg, 6.17 mmol) and Boc_2O (839 mg, 3.85 mmol) were added to a solution of *tert*-butyl 4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)methyl)piperidine-1-carboxylate (Intermediate AV, 630 mg, 1.54 mmol) in 1,4-dioxane (13 mL) and stirred at RT for 16 h. After completion, the reaction mixture was poured into water, 10 and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with water (50 mL), brine (50 mL) and dried over *anhydrous* Na_2SO_4 , filtered and evaporated. The crude was purified by column chromatography (SiO_2) using EtOAc: petroleum ether (1:9) to afford (*Tert*-butyl 4-(((*trans*)-2-(4-bromophenyl)cyclopropyl)(*tert*-butoxycarbonyl)amino)methyl)piperidine-1-carboxylate (600 mg, 76.53 %) as gummy liquid.

15 **Intermediate AX: *Tert*-butyl 4-methylenepiperidine-1-carboxylate**



t-BuOK (4.2 g, 37.63 mmol) was added portionwise to a solution of methyltriphenylphosphonium bromide (12.5 g, 35.13 mmol) in dry Et_2O (40 mL) at 0 °C and stirred at 0 °C for 30 mins. Then a solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (5 g, 25.09 mmol) in dry Et_2O (25 mL) was added and stirred at RT for 16 h. After 20 completion, the reaction mixture was poured into saturated NH_4Cl solution (25 mL) and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with water (30 mL), brine (30 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated. The crude residue was purified by column chromatography (SiO_2) using EtOAc : petroleum ether (5: 95) to get *tert*-butyl 4-methylenepiperidine-1-carboxylate (3.8 g, 76.76%) as a white solid.

25 **Intermediate AY: *Tert*-butyl 1,1-dichloro-2-oxo-7-azaspiro[3.5]nonane-7-carboxylate**




Zn-Cu couple (15.6 g, 121.74 mmol) was added to a solution of *tert*-butyl 4-methylenepiperidine-1-carboxylate (Intermediate AX, 2 g, 10.14 mmol) in Diethyl ether (20 mL) followed by the addition of trichloroacetyl chloride (10.14 g, 55.77 mmol) in Dimethoxyethane (10 mL) at RT and stirred for 16 h. After completion the reaction 30

mixture was poured into NaHCO_3 solution, filtered through celite and extracted with EtOAc ($2 \times 30 \text{ mL}$). The combined extracts were washed with water (20 mL), brine (20 mL), dried over *anhydrous* Na_2SO_4 , filtered and evaporated to get *tert*-butyl 1,1-dichloro-2-oxo-7-azaspiro [3.5]nonane-7-carboxylate (3 g crude, 96%). The crude was carried to next step without further purification.

5


Intermediate AZ: *Tert*-butyl 2-oxo-7-azaspiro[3.5]nonane-7-carboxylate



Zinc powder (4.5 g, 1.5 weight equiv) was added portionwise to a solution of *tert*-butyl 1,1-dichloro-2-oxo-7-azaspiro[3.5]nonane-7-carboxylate (Intermediate AY, 3 g, 9.77 mmol) in saturated NH_4Cl (1.81 g, 34.20 mmol) in Methanol 30 ml) at RT and stirred for 4 h. After completion, the reaction mixture was filtered through a pad of celite and the solvent was evaporated. The crude residue was purified by column chromatography (SiO_2) using EtOAc : petroleum ether (10:90) as the eluent to get *tert*-butyl 2-oxo-7-azaspiro[3.5]nonane-7-carboxylate (750 mg, 32.18 %) as a white solid

10


Intermediate BA: *N*-(3-bromo-4-methoxyphenyl)methanesulfonamide



Methanesulphonyl chloride (55.82 mL, 0.494 mmol) was added to a solution of 3-bromo-4-methoxyaniline (100 mg, 0.494 mmol) in pyridine (1 mL) at 0°C and stirred at RT for 2 h. After completion, reaction mixture was poured into ice water (10 mL) and extracted with EtOAc ($2 \times 15 \text{ mL}$). The combined extracts were washed with water ($3 \times 10 \text{ mL}$), brine (15 mL) and dried over *anhydrous* Na_2SO_4 filtered, and evaporated. The crude residue was purified by column chromatography (SiO_2) by using EtOAc :Hexane (3:7) to afford *N*-(3-bromo-4-methoxyphenyl)methanesulfonamide (137 mg, 99.2 %) as a white solid

20

Intermediate BB: *N*-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanesulfonamide




25

A solution of *N*-(3-bromo-4-methoxyphenyl)methanesulfonamide (Intermediate BA, 136 mg, 0.485 mmol), bispinacolborane (147 mg, 0.582 mmol) and KOAc (87.3 mg, 0.89 mmol) in dioxane (5.5 mL) was degassed for 30 min, then $\text{PdCl}_2(\text{dppf})_2$ (17.7 mg, 0.020 mmol) was added and the reaction mixture was heated at 100°C for 3 h. After completion, the reaction mixture was poured into water (10 mL), extracted with EtOAc ($2 \times 15 \text{ mL}$).

The combined extracts were washed with water (10 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The residue was purified by column chromatography (SiO₂) EtOAc: petroleum ether (1:9) to afford *N*-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) phenyl)methanesulfonamide (100 mg, 63.2 %) as a white solid.


5

Intermediate BC: *Tert*-butyl ((*trans*)-2-(2'-methoxy-5'-(methylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate




K₂CO₃ (246 mg, 1.923 mmol) was added to a solution of *tert*-butyl (*trans*)-2-(4-bromophenyl)cyclopropylcarbamate (Intermediate R, 200 mg, 0.641 mmol) and *N*-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) phenyl)methanesulfonamide (251 mg, 0.769 mmol) in ACN:water (8:2) (10 mL). The reaction mixture was degassed with argon for 20 min. and tetrakis(triphenylphosphine)palladium (0) (Pd(PPh₃)₄) (37 mg, 0.032 mmol) was then added to the reaction mixture and heated to reflux at 90 °C for 4 h. After completion, reaction mixture was poured into ice water (15 mL) and extracted with EtOAc (2 x 20 mL). The combined organic extracts were washed with water (15 mL), brine (15 mL) and dried over *anhydrous* Na₂SO₄ filtered, and evaporated under reduced pressure. The crude residue was purified by column chromatography (SiO₂) by using EtOAc: Hexane (20: 80) to get *tert*-butyl ((*trans*)-2-(2'-methoxy-5'-(methylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (80 mg, 28.9 %) as an off white solid.

Intermediate BD: *N*-(4'-((*trans*)-2-aminocyclopropyl)-6-methoxy-[1,1'-biphenyl]-3-yl)methanesulfonamide hydrochloride




HCl in 1,4-dioxane (0.8 ml) was added to a solution *tert*-butyl ((*trans*)-2-(2'-methoxy-5'-(methylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (Intermediate BC, 80 mg, 0.185 mmol) in 1,4-dioxane (1.6 mL) at 0 °C and stirred at RT for 6 h. After completion, the solvent was evaporated and the residue was triturated with diethyl ether (5 mL) followed by n-hexane (5 mL) to afford *N*-(4'-((*trans*)-2-aminocyclopropyl)-6-methoxy-[1,1'-biphenyl]-3-yl)methanesulfonamide hydrochloride (55 mg, 80.8 %), a yellow solid.

Intermediate BE: *Tert*-butyl ((*trans*)-2-(3'-amino-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate



K_2CO_3 (331 mg, 2.403 mmol) was added to a solution of *tert*-butyl (*trans*)-2-(4-bromophenyl)cyclopropylcarbamate (Intermediate R, 250 mg, 0.801 mmol) and 3-Amino phenyl boronic acid (130.7 mg, 0.961 mmol) in ACN:water (8:2 mL), the reaction mixture was degassed with argon for 20 min. and tetrakis(triphenylphosphine)palladium (0) ($\text{Pd}(\text{PPh}_3)_4$) (9.2 mg, 0.008 mmol) was then added to the reaction mixture and heated to reflux at 90 °C for 4 h. After completion, reaction mixture was poured into ice water (15 mL) and extracted with EtOAc (2 x 20 mL). The combined extracts were washed with water (15 mL), brine (15 mL) and dried over *anhydrous* Na_2SO_4 filtered, and evaporated. The crude residue was purified by column chromatography (SiO_2) using EtOAc: Hexane (3:7) to get *tert*-butyl ((*trans*)-2-(3'-amino-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (100 mg, 38.6 %) as a light yellow solid.


Intermediate BF: *Tert*-butyl ((*trans*)-2-(3'-(1-methylethylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate



Propane-2-sulfonyl chloride (43.9 mg, 0.308 mmol) was added to a solution of *tert*-butyl ((*trans*)-2-(3'-amino-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (Intermediate BE, 100 mg, 0.308 mmol) in pyridine (1 mL) at 0 °C stirred at RT for 4 h. After completion, the reaction mixture was poured into ice water and extracted with EtOAc. The combined extracts were washed with water, brine, dried over *anhydrous* Na_2SO_4 , filtered and evaporated. The crude residue was purified by column chromatography (SiO_2) using EtOAc: petroleum ether (3:7) to afford *tert*-butyl ((*trans*)-2-(3'-(1-methylethylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (100 mg, 75.7 %) as a light brown solid.


Intermediate BG: N-(4'-((*trans*)-2-aminocyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide hydrochloride

159



HCl in 1,4-dioxane (0.5 mL) was added to a solution *tert*-butyl ((*trans*)-2-(3'-(1-methylethylsulfonamido)-[1,1'-biphenyl]-4-yl)cyclopropyl)carbamate (Intermediate BF, 100 mg, 0.232 mmol) in 1,4-dioxane (1 mL) and stirred at RT for 16 h. After completion, the solvent was evaporated. The solid residue was triturated with Et₂O and dried to afford N-(4'-((*trans*)-2-aminocyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide hydrochloride (60 mg, 70.58 %), as a pale yellow powder.

Example 1: N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine hydrochloride



10 **Step 1:**


tert-butyl 4-oxopiperidine-1-carboxylate (890 mg, 4.511 mmol) and sodium triacetoxy borohydride (2.2 g, 11.25 mmol) was added to a solution of *trans*-2-phenylcyclopropanamine (500 mg, 3.75 mmol) in DCE (10 mL) at 0 °C and stirred at RT for 16 h. After completion, solvent was evaporated, the crude residue was taken in water (25 mL) and extracted with EtOAc (2 x 25 mL). The combined extracts were washed with NaHCO₃, water (25 mL), brine (25 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude was purified by column chromatography using (SiO₂) MeOH: CHCl₃ (2: 98) to afford *tert*-butyl 4-((*trans*)-2-phenylcyclopropyl)amino)piperidine-1-carboxylate (350 mg, 25.4 %).

15 **Step 2:**

HCl in dioxane was added dropwise to a solution of *tert*-butyl *tert*-butyl 4-((*trans*)-2-phenylcyclopropyl)amino)piperidine-1-carboxylate (150 mg, 0.0473 mol) in 1, 4 dioxane at 10 °C and stirred at RT for 16 h. After completion, the solvent was evaporated, the solid was triturated with Et₂O and dried to afford N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine hydrochloride (80 mg, 84.3 %) as off white solid.

¹HNMR (400 MHz, DMSO D6) δ: 9.95 (brs, 2H), 9.09 (brs, 1H), 8.88 (brs, 1H), 7.33-7.28 (m, 2H), 7.24-7.17 (m, 3H), 3.49 (s, 1H), 3.40-3.36 (m, 3H), 2.95-2.91 (m, 3H), 2.57 (s, 1H), 2.23 (d, *J* = 12 Hz, 2H), 1.94-1.90 (m, 2H), 1.61-1.56 (m, 1H), 1.31-1.26 (m, 1H); **Mass (M+H):** 217.3

Example 1 describes a racemic, which was the combination of the two different conformations regarding to the cyclopropyl ring ((1*R*,2*S* and 1*S*, 2*R*). The synthesis of each of these isomers was performed as follows:


Example 2: N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine hydrochloride**Step 1:**

5 *Tert*-butyl 4-oxopiperidine-1-carboxylate (1.49 g, 7.51 mmol), acetic acid (450 mg 7.51 mmol) and sodium triacetoxy borohydride (856 mg, 22.53 mmol) was added to a solution of (1S, 2R)-2-phenylcyclopropanamine (Intermediate AJ, 1 g, 7.51 mmol) in DCE (20 mL) at 0 °C, and stirred at RT for 3 h. After completion, solvent was evaporated, crude residue was taken in water (50 mL), treated with NaHCO₃, and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with water (50 mL), brine (50 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude was purified by column chromatography using SiO₂ by eluting 10 EtOAc: pet ether (3: 7) to afford *tert*-butyl 4-((1S, 2R)-2-phenylcyclopropylamino)piperidine-1-carboxylate (1.2 g, 52 %).

Step 2:

15 HCl in 1, 4 dioxane (10 mL) was added to a solution of *tert*-butyl 4-((1S, 2R)-2-phenylcyclopropylamino)piperidine-1-carboxylate (1.2 g, 3.78 mmol) in dioxane (12 mL) at 15 °C and stirred at RT for 16 h. After completion, the solvent was evaporated. The solid residue was triturated with Et₂O and dried to afford N-((1S, 2R)-2-phenylcyclopropyl)piperidin-4-amine hydrochloride (750 mg, 91.4 %) as off white solid
 1H NMR (400 MHz, DMSO d6) δ: 9.86 (brs, 2H), 9.09 (brs, 1H), 8.80 (brs, 1H), 7.33-7.29 (m, 2H), 7.24-7.17 (m, 3H), 3.49 (s, 1H), 3.40-3.36 (m, 3H), 2.96-2.91 (m, 3H), 2.67 (s, 1H), 2.22 (d, J = 12 Hz, 2H), 1.87 (brs, 2H), 1.55 (brs, 1H), 1.31-1.29 (m, 1H); Mass (M+H): 217.21; [α]_D^{27.6}: +72.11 (C=0.5 % in DMSO)


20

Example 3: N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine hydrochloride


This compound was synthesized following the same procedure described in example 2 but using the intermediate AI in Step 1, to afford 735 mg (89 %) as off-white solid

25 1H NMR (400 MHz, DMSO d6) δ: 9.91 (brs, 2H), 9.03 (brs, 1H), 8.84 (brs, 1H), 7.33-7.29 (m, 2H), 7.24-7.17 (m, 3H), 3.49 (s, 1H), 3.40-3.36 (m, 3H), 2.94-2.88 (m, 3H), 2.55 (s, 1H), 2.23 (d, J = 12 Hz, 2H), 1.90-1.87 (m, 2H), 1.58-1.56 (m, 1H), 1.30-1.27 (m, 1H); Mass (M+H): 217.21; [α]_D^{27.4}: -63.82 (C=0.5 % in DMSO).


30 The following compounds can be synthesized following the methodology described for example 1 by using the corresponding intermediates. The Step 2 was only performed in case the intermediate used in the reductive alkylation (Step 1) contained a Boc (tert-butyloxycarbonyl) protecting group

Example 4: N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine hydrochloride


¹HNMR (400 MHz, DMSO D6) δ : 9.80 (brs, 2H), 8.97 (brs, 1H), 8.81 (brs, 1H), 7.44-7.37 (m, 3H), 7.34-7.30 (m, 2H), 7.11 (d, J = 8 Hz, 2H), 6.96 (d, J = 8 Hz, 2H), 5.08 (s, 2H), 3.47-3.35 (m, 2H), 2.93-2.88 (m, 3H), 2.50 (brs, 1H), 2.23-2.20 (d, J = 8 Hz, 2H), 1.86-1.84 (m, 2H), 1.49 (brs, 1H), 1.25-1.20 (m, 1H); Mass (M+H): 323.4

Example 5: (Trans)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine hydrochloride

¹HNMR (400 MHz, DMSO D6) δ : 9.56 (brs, 2H), 8.91 (brs, 1H), 8.73 (brs, 1H), 7.33-7.28 (m, 2H), 7.24-7.17 (m, 3H), 3.27 (d, J = 12 Hz, 1H), 3.00-2.83 (m, 5H), 2.59 (brs, 1H), 2.05 (brs, 1H), 1.96 (d, J = 11 Hz, 2H), 1.63-1.59 (m, 1H), 1.46-1.37 (m, 1H), 1.28-1.23 (m, 1H); Mass (M+H): 231.27


Example 6: (Trans)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine hydrochloride

¹HNMR (400 MHz, CD3OD) δ : 7.39-7.31 (m, 2H), 7.23-7.21 (m, 1H), 7.21 (d, J = 8 Hz, 2H), 3.38 (d, J = 4 Hz, 2H), 3.38 (t, J = 6 Hz, 2H), 3.29-3.21 (m, 2H), 3.00-2.92 (m, 3H), 2.57-2.46 (m, 1H), 1.98 (d, J = 6 Hz, 2H), 1.57-1.51 (m, 1H), 1.49-1.42 (m, 1H), 1.41-1.31 (m, 3H); Mass: (M+H): 244.9.

Example 7: N-((trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine hydrochloride

¹HNMR (400 MHz, DMSO D6) δ : 8.66 (brs, 1H), 8.66 (brs, 1H), 8.41 (brs, 1H), 8-38 (d, J = 8 Hz, 1H), 8.10 (d, J = 8 Hz, 2H), 7.72-7.82 (m, 3H), 3.93 (d, J = 11 Hz, 1H), 3.469-3.46 (m, 1H), 3.32 (t, J = 4 Hz, 2H), 3.12 (s, 1H), 2.70 (s, 1H), 2.09-1.98 (m, 2H), 1.70-1.74 (m, 3H); 1.49-1.46 (m, 1H); Mass (M+H): 363.1

5 Example 8: (Trans)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine hydrochloride



Step 1:

NaH (108 mg, 2.69 mmol) was added to a solution of tert-butyl ((trans)-2-phenylcyclopropyl)carbamate (300 mg, 1.28 mmol) in dry DMF (12 mL) and stirred at RT for 4 h, then 4-(2-bromoethyl)tetrahydro-2H-pyran (281 mg, 1.41 mmol) was added and stirred at RT for 2 h. After completion, the reaction mixture was evaporated, then diluted with CH₂Cl₂ (25 mL) and washed with a saturated NH₄Cl solution. The organic layer was washed with brine (10 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The residue was purified by column chromatography (SiO₂) using Hexane: Methyl tert-butylether (60:40) to get tert-butyl ((trans)-2-phenylcyclopropyl)(2-(tetrahydro-2H-pyran-4-yl)ethyl)carbamate (314 mg, 71.2 %) as a colorless oil


15 Step 2:

HCl in diethyl ether (5 mL) was added to a solution of *tert*-butyl ((trans)-2-phenylcyclopropyl)(2-(tetrahydro-2H-pyran-4-yl)ethyl)carbamate (314 g, 0.91 mmol) in diethyl ether (6 mL) and stirred at RT for 16 h. After completion, the solvent was evaporated. The solid residue was triturated with Et₂O and dried to afford (Trans)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine hydrochloride (151 mg, 41.8 %) as off white solid

¹HNMR (400 MHz, CD₃OD) δ : 7.32-7.29 (m, 2H), 7.23-7.21 (m, 1H), 7.17 (d, J = 8Hz, 2H), 3.92 (dm, J = 4 Hz, 2H), 3.41 (t, J = 6 Hz, 2H), 3.21 (t, J = 4 Hz, 2H), 2.97-2.90 (m, 1H), 2.52 (brs, 1H), 1.74-1.60 (m, 5 H), 1.57-1.49 (m, 1H), 1.39-1.24 (m, 3H); Mass: (M+H): 246.3.

25 The following compound can be synthesized following the methodology described for example 8 by using the corresponding intermediates.


Example 9: (Trans)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine hydrochloride



¹HNMR (400 MHz, DMSO D6) δ : 9.2-9.3 (brs, 2H), 7.67 (d, J = 8 Hz, 2H), 7.61 (d, J = 8 Hz, 2H), 7.50 (d, J = 8 Hz, 2H), 7.28 (d, J = 8 Hz, 2H), 3.82 (d, J = 8 Hz, 2H), 3.25 (m, J = 11 Hz, 2H), 3.08 (brs, 2H), 2.99 (brs, 1H), 2.55 (s, 1H), 1.61-1.57 (m, 6H), 1.33-1.30 (m, 1H), 1.12-1.19 (m, 1H); Mass: (M+H): 356.3.

5 The following compounds can be synthesized following the methodology described for example 1 by using the corresponding appropriate intermediates or commercially available reagents. Step 2 was only performed when the intermediate used in the reductive alkylation (Step 1) contained a Boc (tert-butyloxycarbonyl) protecting group.


10 Example 10: N-((*trans*)-2-phenylcyclopropyl)piperidin-3-amine



¹HNMR (400 MHz, D₂O) δ : 7.52 (brs, 2H), 7.48-7.42 (m, 1H), 7.30 (brs, 2H), 3.89-3.81 (brm, 2H), 3.56-3.53 (m, 1H), 3.21 (brs, 1H), 3.12-3.02 (m, 2H), 2.61 (brs, 1H), 2.46 (brs, 1H), 2.23-2.17 (m, 1H), 1.93-1.78 (brm, 2H), 1.60 (brs, 2H); Mass (M+H): 217.28. This compound was obtained as a hydrochloride salt.


15

Example 11: N-((*trans*)-2-phenylcyclopropyl)pyrrolidin-3-amine




19 ¹HNMR (400 MHz, D₂O) δ : 7.57-7.46 (m, 2H), 7.43-7.39 (m, 1H), 7.36-7.31 (m, 2H), 4.39-4.32 (m, 1H), 3.97-3.89 (m, 1H), 3.73-3.51 (m, 3H), 3.10 (s, 1H), 2.68 (brs, 2H), 2.39-2.31 (m, 1H), 1.63 (brs, 1H), 1.57 (brs, 1H); Mass (M+H): 203.2. This compound was obtained as a hydrochloride salt.

Example 12: N-((*trans*)-2-phenylcyclopropyl)azetidin-3-amine



20 ¹HNMR (400 MHz, D₂O) δ : 7.57-7.46 (m, 2H), 7.43-7.39 (m, 1H), 7.36-7.31 (m, 2H), 4.79-4.72 (m, 1H), 4.55 (brs, 4H), 3.01 (brs, 1H), 2.60 (brs, 1H), 1.63 (brs, 1H), 1.52-1.43 (m, 1H); Mass (M+H): 189.20. This compound was obtained as hydrochloride salt


Example 13: N-((*trans*)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine



¹HNMR (400 MHz, D₂O) δ: 7.55-7.44 (brm, 3H), 7.37-7.34 (m, 2H), 4.31 (s, 2H), 3.94 (s, 1H), 3.23 (s, 1H), 2.76 (brs, 3H), 2.45-2.34 (brm, 4H), 2.22-2.18 (m, 2H), 1.78 (brs, 1H), 1.72-1.60 (m, 1H); Mass (M+H): 243.30. This compound was obtained as a hydrochloride salt.


5

Example 14: N-((trans)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine




¹HNMR (400 MHz, DMSO d₆) δ: 9.37 (brs, 2H), 8.53 (brs, 2H), 7.38-7.29 (m, 2H), 7.22-7.18 (m, 3H), 3.18 (brs, 1H), 3.02 (brs, 5H), 2.92 (brs, 1H), 1.92-1.83 (m, 2H), 1.79-1.71 (m, 2H), 1.62-1.42 (m, 7H), 1.38-1.30 (m, 1H), 1.20-1.05 (m, 2H); Mass (M+H): 285.32. This compound was obtained as a hydrochloride salt.

Example 15: N-((trans)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine




¹HNMR (400 MHz, D₂O) δ: 7.45-7.39 (m, 2H), 7.38-7.32 (m, 1H), 7.27-7.21 (m, 2H), 4.11-4.04 (m, 1H), 3.26-3.18 (m, 4H), 2.96-2.94 (m, 1H), 2.57-2.43 (m, 3H), 2.20-2.15 (m, 2H), 1.98-1.85 (m, 4H), 1.42 (brs, 1H), 1.42-1.37 (m, 1H); Mass (M+H): 257.32. This compound was obtained as a hydrochloride salt.

Example 16: N-((trans)-2-phenylcyclopropyl)decahydroquinolin-4-amine



¹HNMR (400 MHz, DMSO-d₆) δ: 10.24-10.01 (m, 1H), 9.40-9.02 (m, 2H), 7.38-7.28 (m, 2H), 7.22-7.18 (m, 3H), 3.81-3.58 (m, 2H), 3.30-3.22 (m, 1H), 2.98-2.74 (m, 2H), 2.33-2.20 (m, 1H), 2.18-1.92 (m, 3H), 1.85-1.64 (m, 4H), 1.48-1.30 (m, 3H), 1.28-1-18 (m, 2H); Mass (M+H): 271.31. This compound was obtained as a hydrochloride salt.


25 **Example 17: N-((trans)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine**



¹HNMR (300 MHz, D₂O) δ: 8.78-8.62 (m, 2H), 8.32 (d, *J* = 8.2 Hz, 1H), 8.08-8.01 (m, 1H), 3.82-3.77 (m, 1H), 3.66-3.62 (m, 1H), 3.36-3.30 (m, 2H), 3.11-3.02 (m, 2H), 2.86-2.82 (m, 1H), 2.46-2.31 (m, 2H), 2.03-1.65 (m, 4H); Mass(M+H): 218.00. This compound was obtained as a hydrochloride salt.

5

Example 18: N-((trans)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, D₂O) δ: 9.24 (s, 1H), 7.86 (s, 1H), 3.75-3.65 (m, 1H), 3.58-3.52 (m, 2H), 3.19-3.04 (brm, 3H), 2.83 (brs, 1H), 2.43-2.39 (m, 2H), 1.97-1.85 (brm, 2H), 1.77-1.72 (m, 1H), 1.59-1.52 (m, 1H); Mass (M+H): 224.21. This compound was obtained as a hydrochloride salt.

10


Example 19: N-(2-(o-tolyl)cyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, DMSO-d₆) δ: 10.13 (s, 1H), 9.99 (s, 1H), 9.26 (s, 1H), 8.99 (s, 1H), 7.19-7.12 (m, 3H), 7.01 (s, 1H), 3.49 (s, 1H), 3.34 (s, 2H), 2.95 (s, 3H), 2.72-2.69 (m, 1H), 2.50 (s, 3H), 2.29-2.25 (m, 2H), 1.99-1.91 (m, 2H), 1.57 (q, *J* = 5.8 Hz, 1H), 1.80 (q, *J* = 6.8 Hz, 1H); Mass (M+H): 231.3. This compound was obtained as a hydrochloride salt.

15

Example 20: N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine




20

¹HNMR (400 MHz, DMSO-d₆) δ: 10.04 (s, 2H), 9.09 (s, 1H), 8.87 (s, 1H), 7.32-7.25 (m, 1H), 7.22-7.11 (m, 3H), 3.32 (s, 3H), 3.04 (s, 1H), 2.93 (s, 2H), 2.71 (s, 1H), 2.23 (s, 2H), 1.89 (s, 2H), 1.60 (s, 1H), 1.32 (s, 1H); Mass (M+H): 235.3. This compound was obtained as a hydrochloride salt.

25


Example 21: N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, DMSO-d₆) δ: 9.41 (s, 1H), 8.74 (s, 1H), 8.54 (s, 1H), 7.39-7.22 (m, 2H), 7.06 (brs, 1H), 3.36 (brs, 3H), 2.93 (brs, 2H), 2.57-2.53 (m, 1H), 2.14 (brs, 2H), 1.70 (brs, 2H), 1.43 (s, 1H), 1.33 (s, 1H); **Mass (M+H):** 253.19. This compound was obtained as a hydrochloride salt.


5

Example 22: N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine




¹HNMR (400 MHz, DMSO-d₆) δ: 9.83 (s, 1H), 8.99 (s, 1H), 8.89 (s, 1H), 7.11 (d, *J* = 8.8 Hz, 2H), 6.87 (d, *J* = 8.8 Hz, 2H), 3.72 (s, 3H), 3.38 (brs, 1H), 2.93-2.85 (m, 3H), 2.33-2.30 (m, 2H), 1.87 (brs, 2H), 1.50 (s, 1H), 1.24-1.21 (m, 1H); **Mass (M+H):** 247.28. This compound was obtained as a hydrochloride salt.

Example 23: 4-(2-(piperidin-4-ylamino)cyclopropyl)phenol




¹HNMR (400 MHz, D₂O) δ: 7.10 (d, *J* = 8.8 Hz, 2H), 6.86 (d, *J* = 8.8 Hz, 2H), 3.74-3.67 (m, 1H), 3.61-3.57 (m, 2H), 3.15-3.07 (m, 2H), 2.94-2.90 (m, 1H), 2.49-2.41 (m, 3H), 1.94-1.88 (m, 2H), 1.50-1.38 (brm, 2H); **Mass (M+H):** 233.19. This compound was obtained as a hydrochloride salt.

Example 24: N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, D₂O) δ: 7.94-7.88 (m, 3H), 7.72 (s, 2H), 7.57-7.54 (m, 1H), 7.34 (d, *J* = 8.8 Hz, 1H), 3.74-3.73 (m, 1H), 3.61-3.57 (m, 2H), 3.12-3.09 (m, 3H), 2.70-2.68 (m, 1H), 2.46-2.44 (m, 2H), 1.97-1.91 (m, 2H), 1.64-1.60 (m, 1H); **Mass (M+H):** 267.22. This compound was obtained as a hydrochloride salt.


Example 25: N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, DMSO-d₆) δ: 9.97 (s, 1H), 9.49 (s, 1H), 8.98 (s, 1H), 8.87 (s, 1H), 7.34 (s, 4H), 7.23 (s, 1H), 3.55 (s, 1H), 3.29-3.24 (m, 1H), 2.95 (s, 2H), 2.76 (s, 1H), 2.38-2.29 (m, 2H), 1.97 (brs, 2H), 1.62 (s, 3H), 1.44-1.40 (m, 1H), 1.34-1.31 (m, 1H); Mass (M+H): 231.3. This compound was obtained as a hydrochloride salt.


5

Example 26: (Trans)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine




¹HNMR (400 MHz, D₂O) δ: 7.46-7.38 (m, 2H), 7.37-7.29 (m, 1H), 7.22-7.18 (m, 2H), 3.64-3.59 (m, 1H), 3.51-3.23 (brm, 4H), 3.07-3.02 (brm, 2H), 2.80-2.76 (m, 1H), 2.57 (brs, 1H), 2.36-2.32 (m, 1H), 1.84-1.78 (m, 1H), 1.58 (brs, 1H), 1.47-1.41 (brm, 1H); Mass (M+H): 217.3. This compound was obtained as a hydrochloride salt.

Example 27: (Trans)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine



¹HNMR (400 MHz, DMSO d₆) δ: 9.37 (s, 2H), 8.80 (s, 1H), 8.60 (s, 1H), 7.54 (t, J = 8.0 Hz, 1H), 7.46-7.39 (m, 1H), 7.25 (q, J = 8.0 Hz, 2H), 7.13 (d, J = 8.0 Hz, 2H), 6.97 (d, J = 8.0 Hz, 2H), 5.11 (s, 2H), 3.27-3.21 (m, 2H), 3.02 (s, 2H), 2.91-2.80 (m, 3H), 2.52-2.48 (m, 1H), 2.02 (s, 1H), 1.94-1.99 (m, 2H), 1.52 (s, 1H), 1.41-1.32 (s, 2H), 1.22-1.18 (s, 1H); Mass (M+H): 416.18. This compound was obtained as a hydrochloride salt.


Example 28: (Trans)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine



¹HNMR (400 MHz, DMSO d₆) δ: 9.52-9.23 (brm, 3H), 8.78 (brs, 2H), 8.45 (brs, 2H), 7.98-7.94 (m, 2H), 7.70-7.63 (m, 4H), 7.31 (d, J = 8.0 Hz, 2H), 3.3 (brs, 2H), 3.05 (s, 3H), 2.91-2.83 (m, 2H), 2.62-2.54 (m, 1H), 2.01-1.92 (m, 3H), 1.6 (brs, 1H), 1.42-1.25 (m, 3H); Mass (M+H): 375.30. This compound was obtained as a hydrochloride salt.


25

Example 29: N-(6-methoxy-4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide




5 ^1H NMR (400 MHz, DMSO d_6) δ : 10.01 (s, 2H), 9.48 (s, 1H), 9.12 (s, 1H), 8.91 (s, 1H), 7.39 (d, J = 8.0 Hz, 2H), 7.23 (d, J = 8.0 Hz, 2H), 7.19-7.08 (m, 3H), 3.73 (s, 3H), 3.51 (brs, 1H), 3.08-2.95 (m, 6H), 2.82 (s, 1H), 2.28-2.21 (m, 2H), 1.97 (s, 2H); Mass (M+H): 416.18. This compound was obtained as a hydrochloride salt.

10 Example 30: N-(4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide



15 ^1H NMR (400 MHz, DMSO d_6) δ : 7.63 (d, J = 8.0 Hz, 2H), 7.52-7.47 (m, 3H), 7.30 (d, J = 8.0 Hz, 3H), 3.76-3.70 (m, 1H), 3.62-3.58 (m, 2H), 3.49-3.43 (m, 1H), 3.15-3.03 (m, 3H), 2.58-2.55 (m, 1H), 2.46-2.43 (m, 2H), 1.97-1.94 (m, 2H), 1.62-1.51 (m, 2H), 1.39-1.29 (2s, 6H); Mass (M+H): 414.22. This compound was obtained as a hydrochloride salt.


Example 31: (*Trans*)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine



20 ^1H NMR (400 MHz, D₂O) δ : 7.48-7.41 (m, 2H), 7.40-7.35 (m, 1H), 7.27-7.21 (m, 2H), 4.28 (brs, 2H), 4.09-4.04 (m, 2H), 3.60-3.58 (m, 2H), 3.46-3.42 (m, 1H), 2.98 (s, 1H), 2.54 (s, 1H), 1.55 (brs, 1H), 1.46-1.40 (brm, 1H); Mass (M+H): 203.17.

Step 2 (Boc-deprotection) was performed with TFA in dichloromethane and the compound was obtained as a trifluoroacetate salt.


25 Example 32: 1-(methylsulfonyl)-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine



¹HNMR (400 MHz, CDCl₃) δ: 7.28-7.23 (m, 2H), 7.18-7.14 (m, 1H), 7.01 (d, *J* = 7.2 Hz, 2H), 3.70-3.67 (m, 2H), 2.84-2.76 (m, 3H), 2.77 (s, 3H), 2.33-2.29 (m, 1H), 2.02-1.97 (m, 2H), 1.88-1.84 (m, 1H), 1.54-1.48 (m, 2H), 1.09-0.99 (m, 2H); Mass (M+H): 295.12.


5 This compound was obtained as free amine.

Example 33: 1-((trans)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone




¹HNMR (400 MHz, DMSO-d₆) δ: 7.39 (d, *J* = 8.8 Hz, 2H), 6.99 (d, *J* = 8.8 Hz, 2H), 4.11-4.07 (m, 1H), 3.71-3.67 (m, 1H), 3.03 (t, *J* = 11.2 Hz, 1H), 2.76-2.67 (m, 2H), 2.20 (s, 1H), 1.96 (brs, 2H), 1.76-1.71 (m, 3H), 1.21-0.95 (m, 4H); Mass (M+H): 337.05. This compound was obtained as free amine

Example 34: 4-((trans)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide




15 ¹HNMR (400 MHz, DMSO-d₆) δ: 7.39 (d, *J* = 8.8 Hz, 2H), 6.99 (d, *J* = 8.8 Hz, 2H), 5.83 (s, 2H), 3.77 (d, *J* = 13.6 Hz, 2H), 2.73-2.62 (m, 4H), 2.32-2.18 (m, 1H), 1.77-1.67 (m, 3H), 1.14-1.12 (m, 2H), 0.98-0.93 (m, 2H); Mass (M+H): 337.99. This compound was obtained as free amine


Example 35: N-((trans)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine




20 ¹HNMR (400 MHz, CDCl₃) δ: 7.35 (d, *J* = 8.8 Hz, 2H), 6.88 (d, *J* = 8.8 Hz, 2H), 3.96 (dm, *J* = 11.6 Hz, 2H), 3.42-3.35 (m, 2H), 2.86-2.79 (m, 1H), 2.32-2.28 (m, 1H), 1.86-1.81 (m, 3H), 1.48-1.37 (m, 2H), 1.12-1.07 (m, 1H), 0.99-0.94 (m, 1H); Mass (M+H): 296.10. This compound was obtained as free amine

Example 36: 2,2,6,6-tetramethyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine


¹HNMR (400 MHz, DMSO d₆) δ: 8.70 (vbrs, 1H), 7.70 (vbrs, 1H), 7.25-7.21 (m, 2H), 7.18-7.09 (m, 1H), 7.08-7.03 (m, 3H), 3.08 (m, 1H), 2.24 (s, 1H), 1.92-1.79 (brm, 3H), 1.33-1.14 (brm, 13H), 0.98-0.94 (s, 2H); Mass (M+H): 273.30. This compound was obtained as free amine. The reagent used in Step 1 (reductive alkylation) was directly 2,2,6,6-tetramethylpiperidin-4-one.

Example 37: 1-methyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine

¹HNMR (400 MHz, DMSO d₆) δ: 10.60 (brs, 1H), 9.98 (brs, 2H), 7.38-7.30 (m, 2H), 7.23-7.18 (m, 3H), 3.56-3.38 (m, 3H), 3.02-2.96 (m, 3H), 2.78 (s, 3H), 2.57 (brs, 1H), 2.28 (brs, 2H), 2.05 (brs, 2H), 1.58 (brs, 1H), 1.28 (m, 1H); Mass (M+H): 231.22. After Step 1 (reductive alkylation), the compound was treated with HCl in 1,4-dioxane to afford the corresponding hydrochloride salt.

Example 38: 1-isopropyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine


¹HNMR (400 MHz, D₂O) δ: 7.44-7.41 (m, 2H), 7.39-7.35 (m, 1H), 7.26-7.21 (m, 2H), 3.75-3.57 (brm, 4H), 3.21-3.13 (m, 2H), 3.03 (s, 1H), 2.58-2.51 (m, 3H), 2.07-1.97 (m, 2H), 1.58-1.51 (m, 2H), 1.36 (2s, 6H); Mass (M+H): 259.30. After Step 1 (reductive alkylation), the compound was treated with HCl in 1,4-dioxane to afford the corresponding hydrochloride salt.

Example 39: N-((*trans*)-2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)piperidin-4-amine

¹HNMR (400 MHz, D₂O) δ: 7.44-7.41 (m, 2H), 7.39-7.35 (m, 1H), 7.26-7.21 (m, 2H), 4.05-3.96 (brm, 2H), 3.82-3.68 (brm, 3H), 3.23 (brs, 2H), 3.06-3.02 (m, 1H), 2.61 (brs, 1H), 2.51 (brs, 2H), 2.10 (brs, 2H), 1.61-1.45 (m,

2H); **Mass (M+H):** 299.30. After Step 1 (reductive alkylation), the compound was treated with HCl in 1,4-dioxane to afford the corresponding hydrochloride salt.

Example 40: N-((trans)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine




5

¹HNMR (400 MHz, DMSO-d₆) δ: 9.71 (brs, 2H), 8.27 (d, *J* = 8.4 Hz, 2H), 7.32 (t, *J* = 8.0 Hz, 2H), 7.24 (d, *J* = 8.0 Hz, 2H), 7.19 (d, *J* = 7.2 Hz, 2H), 4.34 (d, *J* = 13.6 Hz, 2H), 3.62 (brs, 1H), 3.28-3.18 (m, 2H), 2.98 (s, 1H), 2.67-2.55 (m, 2H), 2.50-2.49 (m, 2H), 2.32-2.18 (m, 2H), 1.69-1.74 (m, 2H), 1.56 (s, 1H), 1.35-1.30 (m, 1H);

Mass (M+H): 294.3. After Step 1 (reductive alkylation), the compound was treated with HCl in 1,4-dioxane to afford the corresponding hydrochloride salt.

10

Example 41: 4-((trans)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide




15

¹HNMR (400 MHz, D₂O) δ: 7.53 (d, *J* = 8.8 Hz, 2H), 7.11 (d, *J* = 8.8 Hz, 2H), 3.75-3.68 (m, 1H), 3.44-3.35 (m, 4H), 3.01 (s, 1H), 2.62-2.58 (m, 2H), 5.53-2.48 (m, 1H), 2.31-2.21 (m, 2H), 1.58-1.41 (m, 2H); **Mass (M+H):** 344.05. After Step 1 (reductive alkylation), the compound was treated with HCl in 1,4-dioxane to afford the corresponding hydrochloride salt.

20

Example 42: (Trans)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine



Step 1:

25

A solution of *tert*-butyl 4-(((trans)-2-(4-bromophenyl)cyclopropyl)(*tert*-butoxycarbonyl)amino)methyl)piperidine-1-carboxylate (Intermediate AW, 500 mg, 0.98 mmol), cyclopropyl boronic acid (101 mg, 1.17 mmol) and K₂CO₃ (405 mg, 2.94 mmol) in ACN:H₂O (4:1) (10 mL) was degassed for 30 minutes, then Pd (PPh₃)₄ (56 mg, 0.049 mmol) was added and heated at reflux for 16 h. After completion, the reaction mixture was poured into water and extracted with EtOAc (2 x 25 mL). The combined extracts were washed with water (25 mL), brine (25 mL), dried over *anhydrous* Na₂SO₄, filtered and evaporated. The crude was purified by Preparative HPLC to

afford *tert*-butyl 4-(((*tert*-butoxycarbonyl)((*trans*)-2-(4-cyclopropylphenyl)cyclopropyl)amino)methyl)piperidine-1-carboxylate (120 mg) as a white solid.


Step 2:

HCl in 1,4-dioxane (1 mL) was added to a solution of *tert*-butyl 4-(((*tert*-butoxycarbonyl)((*trans*)-2-(4-cyclopropylphenyl)cyclopropyl)amino)methyl)piperidine-1-carboxylate (120 mg, 0.23 mmol) in 1,4-dioxane (2 mL) at 10 °C and stirred at RT for 16 h. After completion, the solvent was evaporated. The solid was triturated with Et₂O and dried to afford (*trans*)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine as a hydrochloride salt (80 mg), a white solid.

¹H NMR (400 MHz, DMSO d₆) δ: 9.49 (s, 1H), 8.89 (s, 1H), 8.70 (s, 1H), 7.06-6.98 (m, 4H), 3.40-3.34 (m, 2H), 2.98 (s, 2H), 2.89-2.80 (m, 3H), 2.08-1.82 (m, 4H), 1.59-1.53 (m, 1H), 1.47-1.38 (m, 2H), 1.18-1.22 (m, 1H), 0.95-0.84 (m, 2H), 0.75-0.62 (m, 2H); Mass (M+H): 271.29.


The following compounds can be synthesized following the methodology described for example 42 by using the corresponding intermediates or commercially available reagents.

Example 43: (*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine



¹H NMR (400 MHz, DMSO d₆) δ: 9.16 (s, 1H), 8.89 (d, J = 8.4 Hz, 2H), 8.82-8.80 (m, 2H), 8.16-8.12 (m, 1H), 7.80 (d, J = 8.4 Hz, 2H), 7.45 (d, J = 8.4 Hz, 2H), 3.45 (d, J = 13.2 Hz, 2H), 3.23-3.01 (m, 5H), 2.76-2.68 (m, 1H), 2.19 (s, 1H), 2.08 (brd, J = 12.8 Hz, 2H), 1.74-1.68 (m, 1H), 1.62-1.43 (m, 1H); Mass (M+H): 308.24. This compound was obtained as hydrochloride salt


Example 44: (*Trans*)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine




¹H NMR (400 MHz, D₂O) δ: 7.71 (brs, 3H), 7.23 (brs, 2H), 6.69 (brs, 1H), 3.44 (brs, 2H), 3.16 (brs, 1H), 2.96 (brs, 3H), 2.52 (brs, 1H), 2.28-2.05 (m, 4H), 1.54-1.42 (m, 4H); Mass (M+H): 297.25. This compound was obtained as hydrochloride salt

The following compounds can be synthesized following the general methods disclosed under the General Synthetic Route Description Section, including the methodologies described in Schemes 1,2,3, 4, 5,6, 7 and 8 and the examples above.


5 **Example 45: N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine**



Example 46: (Trans)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine




Example 47: (Trans)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine




10

Example 48: (Trans)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 49: (Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine




15


Example 50: (Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine




Example 51: (Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine



Example 52: (Trans)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine




Example 53: N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine




5

Example 54: N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine




Example 55: N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine




10


Example 56: N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine




Example 57: N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine




Example 58: N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine




Example 59: N-((trans)-2-phenylcyclopropyl)azepan-3-amine




Example 60: N-((trans)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine




10 Example 61: N-((trans)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine




Example 62: N-((trans)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine




Example 63: N-((trans)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine




Example 64: N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine




Example 65: N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine




Example 66: (1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine




10 Example 67: (1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 68: (1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine




Example 69: (1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine




Example 70: N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine




Example 71: N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine




Example 72: N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine




Example 73: N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine




Example 74: N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine




Example 75: N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine




Example 76: (1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine




Example 77: (1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine




Example 78: (1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine




Example 79: (1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine




Example 80: (1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine




Example 81: (1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine




Example 82: (1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 83: (1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 84: (1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine




Example 85: (1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine




Example 86: (1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine



Example 87: (1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine




10 Example 88: (1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine




Example 89: (1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine


180




Example 90: (1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine




Example 91: (1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine




Example 92: N-((trans)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine




Example 93: N-((1S,2S)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine




Example 94: N-((1R,2R)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine




Example 95: N-((trans)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine




Example 96: (Trans)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 97: N-((trans)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine




Example 98: 2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 99: (trans)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine




Example 100: (trans)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine




Example 101: N-((trans)-2-(o-tolyl)cyclopropyl)piperidin-4-amine




Example 102: N-((trans)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine



Example 103: N-((trans)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine



Example 104: N-((trans)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine

**Example 105: Biological Assays - Inhibition of LSD1**

5 The compounds of the invention can be tested for their ability to inhibit LSD1. The ability of the compounds of the invention to inhibit LSD1 can be tested as follows. Human recombinant LSD1 protein was purchased from BPS Bioscience Inc (catalog reference number 50100: human recombinant LSD1, GenBank accession no. NM_015013, amino acids 158-end with N-terminal GST tag, MW: 103 kDa). In order to monitor LSD1 enzymatic activity and/or its inhibition rate by our inhibitor(s) of interest, di-methylated H3-K4 peptide (Anaspec) 10 was chosen as a substrate. The demethylase activity was estimated, under aerobic conditions, by measuring the release of H₂O₂ produced during the catalytic process, using the Amplex® Red hydrogen peroxide/peroxidase assay kit (Invitrogen).

15 Briefly, a fixed amount of LSD1 was incubated on ice for 15 minutes, in the absence and/or in the presence of at least eight 3-fold serial dilutions of the respective test compound (e.g., from 0 to 75 μ M, depending on the inhibitor strength). Tranylcypromine (Biomol International) was used as a control for inhibition. Within the experiment, each concentration of inhibitor was tested in duplicate. After leaving the enzyme interacting with the inhibitor, K_M of di-methylated H3-K4 peptide was added to each reaction and the experiment was left for 30 20 minutes at 37°C in the dark. The enzymatic reactions were set up in a 50 mM sodium phosphate, pH 7.4 buffer. At the end of the incubation, Amplex® Red reagent and horseradish peroxidase (HPR) solution were added to the reaction according to the recommendations provided by the supplier (Invitrogen), and left to incubate for 5 extra minutes at room temperature in the dark. A 1 μ M H₂O₂ solution was used as a control of the assay, was monitored by fluorescence (excitation at 540 nm, emission at 590 nm) using a microplate reader 25 (Infinite 200, Tecan). Arbitrary units were used to measure level of H₂O₂ produced in the absence and/or in the presence of inhibitor.

30 The maximum demethylase activity of LSD1 was obtained in the absence of inhibitor and corrected for background fluorescence in the absence of LSD1. The IC₅₀ value of each inhibitor was calculated with GraphPad Prism Software.

The results presented in Table 1 below show the results of the LSD1 inhibition studies for a number of the Example compounds. In Table 2 the IC₅₀ values for all examples tested in this assay are shown. Parnate

(tranylcypromine; i.e., 2-trans phenylcyclopropylamine) was found to have a IC50 value of 35±10 micromolar. The studies show that the compounds of the invention have unexpectedly potent LSD1 inhibition.

5 **Example 106: Biological Assays - Monoamine Oxidase Assays for determining the selectivity of the compounds of the invention for LSD1**

Human recombinant monoamine oxidase proteins MAO-A and MAO-B were purchased from Sigma Aldrich. MAOs catalyze the oxidative deamination of primary, secondary and tertiary amines. In order to monitor MAO enzymatic activities and/or their inhibition rate by inhibitor(s) of interest, a fluorescence-based (inhibitor)-screening assay was set up. 3-(2-Aminophenyl)-3-oxopropanamine (kynuramine dihydrobromide, Sigma Aldrich), a non fluorescent compound was chosen as a substrate. Kynuramine is a non-specific substrate for both MAO-A and MAO-B activities. While undergoing oxidative deamination by MAO activities, kynuramine is converted into 4-hydroxyquinoline (4-HQ), a resulting fluorescent product.

15 The monoamine oxidase activity was estimated by measuring the conversion of kynuramine into 4-hydroxyquinoline. Assays were conducted in 96-well black plates with clear bottom (Corning) in a final volume of 100 μ L. The assay buffer was 100 mM HEPES, pH 7.5. Each experiment was performed in duplicate within the same experiment.

20 Briefly, a fixed amount of MAO (0.25 μ g for MAO-A and 0.5 μ g for MAO-B) was incubated on ice for 15 minutes in the reaction buffer, in the absence and/or in the presence of at least eight 3-fold serial dilutions each. Clorgyline and Deprenyl (Sigma Aldrich) was used as a control for specific inhibition of MAO-A and MAO-B respectively.

25 After leaving the enzyme(s) interacting with the inhibitor, K_m of kynuramine was added to each reaction for MAO-B and MAO-A assay respectively, and the reaction was left for 1 hour at 37°C in the dark. The oxidative deamination of the substrate was stopped by adding 50 μ L of NaOH 2N. The conversion of kynuramine to 4-hydroxyquinoline, was monitored by fluorescence (excitation at 320 nm, emission at 360 nm) using a microplate reader (Infinite 200, Tecan). Arbitrary units were used to measure levels of fluorescence produced 30 in the absence and/or in the presence of inhibitor.

35 The maximum of oxidative deamination activity was obtained by measuring the amount of 4-hydroxyquinoline formed from kynuramine deamination in the absence of inhibitor and corrected for background fluorescence in the absence of MAO enzymes. The IC50 values of each inhibitor were calculated with GraphPad Prism Software.

The results obtained with compounds of the invention in the biological assays of examples 105 and 106 are shown below.

Example No.	MAO-A (Ki)	MAO-B (Ki)	LSD1 (Ki)
1	I	I	V
2	I	I	IV
3	I	I	V
4	II	III	V
5	I	II	V
6	II	II	V
7	II	III	IV
8	II	II	III
9	II	III	IV

5 Table 1: Summary of Data from MAO-A, MAO-B, and LSD1 Inhibition Studies

The ranges for the Ki value reported in Table 1 are for MAO-A, MAO-B and LSD1: I = higher than 100 μ M ; II = between 10 μ M and 100 μ M; III = between 1 μ M and 10 μ M; IV = between 0.1 μ M and 1 μ M; V = between 0.001 and 0.1 μ M. The term Ki value is used herein as a designation for the IC50 value, i.e. the concentration required for a half-maximal (50%) inhibition of the corresponding target (MAO-A, MAO-B, or LSD1).

Generally compounds of the Examples were found to have particularly low IC50 values for LSD1, as compared to MAO-A and MAO-B. For some of the compounds of the Examples, IC50 values for LSD1 were lower than 0.1 μ M.

15 The specific IC50 values obtained for the compounds disclosed in the Examples when tested in the assays of Examples 105 (LSD1 inhibition) and 106 (MAO-A and B inhibition) are provided in Table 2 below:

Example no.	MAO-A (IC ₅₀ - μ M)	MAO-B (IC ₅₀ - μ M)	LSD1 (IC ₅₀ - μ M)
1	>100	>100	0.056
2	>100	>100	0.491
3	>100	>100	0.042
4	39.86	3.18	0.019
5	>100	35.37	0.022
6	38.32	10.12	0.038
7	33.45	0.70	0.111
8	>100	>100	2.18
9	14.61	5.60	0.296
10	>100	26.47	0.119
11	37.70	46.78	0.908
12	23.75	>100	1.942
13	>100	>100	0.069
14	>100	54.33	0.071
15	>100	≈100	0.048
16	>100	>100	0.139
17	>100	>100	0.107

18	>100	>100	0.081
19	>100	>100	0.105
20	>100	>100	0.038
21	>100	≈100	0.127
22	≈100	≈100	0.111
23	>100	>100	≈100
24	≈50	15.05	0.047
25	≈100	>100	0.097
26	>100	20.79	0.035
27	3.58	5.01	0.008
28	11.57	1.02	0.084
29	≈100	≈100	0.023
30	>100	>100	0.111
31	>100	25.50	0.042
32	>100	>100	1.677
33	≈50	≈50	0.212
34	34.12	32.40	0.104
35	45.41	18.69	0.549

36	>100	>100	0.068
37	>100	>100	0.153
38	>100	>100	0.108
39	>100	>100	4.499
40	>100	35.96	0.188
41	>50	16.93	0.177
42	≈100	3.79	0.037
43	43.10	≈100	0.032
44	≈100	>100	0.072

As the data in the above table show, the compounds of the invention are very potent LSD1 inhibitors, with IC50 values in many cases below 100 nM or even below 50 nM. In addition, the compounds exhibit high selectivity versus MAO-A and MAO-B, with IC50 values for LSD1 in general \geq 100-fold more potent than the corresponding IC50 values for MAO-A and MAO-B.

5

Example 107: Cellular assay – Induction of differentiation of THP-1 leukemia cells

Acute Myeloid Leukemia (AML) is characterized by the presence of leukemic cells with a maturation arrest that divide rapidly. With the induction of terminal differentiation, AML cells lose the ability to proliferate and end up dying without the need of a direct cytotoxic effect.

10 By analyzing the induction of CD11b membrane expression on THP-1 cells we are assessing the ability of LSD1 inhibitors to induce terminal monocytic differentiation of a MLL-AF9 AML cell line.

15

The assay was performed as follows:

THP-1 cells were established from the peripheral blood of a 1-year-old boy with acute monocytic leukemia at relapse in 1978. They carry t(9;11)(p21;q23) leading to MLL-MLLT3 (MLL-AF9) fusion gene. This cell line can undergo monocytic differentiation upon treatment with the appropriate stimulus. THP-1 were purchased from

DSMZ GmbH (Deutsche Sammlung von Mikroorganismen und Zellkulturen) and cultured in RPMI 1640 medium containing 10% of fetal bovine serum.

In this assay, 150,000 THP-1 cells were seeded in 1 ml of complete culture medium in 6-well tissue culture plates. Serial dilutions of the compounds were prepared in DMSO and then further diluted in complete medium to generate solutions of concentrations that are double of the final concentration at which the cells will be exposed. 1ml of these 2x concentrated solutions was added to the cells. DMSO final content must be the same in all the wells and must be kept below 0.1% v/v (usually 0.01-0.02% v/v), since higher DMSO content can induce differentiation of THP-1 cells.

10

Cells were kept in the presence of compounds for 96h at a 5% CO₂ atmosphere at 37°C. After this treatment period, cells were harvested, washed twice with PBS buffer and placed in a V-bottom 96-well plate. Each treated sample was split in two. One was stained with a phycoerythrin-labeled anti-CD11b antibody (clone ICRF44, purchased from eBiosciences) and the other with the relevant phycoerythrin-labeled isotype control antibody (mouse IgG₁, purchased from eBiosciences). Samples were incubated in the dark at 4°C for 30-60 minutes and washed three times in PBS buffer containing 1% fetal bovine serum.

15

Samples were analyzed in a flow cytometer equipped with a blue laser (488nm). Emitted fluorescence was detected and quantified with a 575/30nm filter. Percentage of CD11b positive cells, as an indicator of monocytic differentiation, was determined compared to isotype control antibody stained cells. EC₅₀ values were calculated by non-linear regression analysis.

20

The results obtained with compounds of the invention in this test are shown in Table 3 below.

Example No	EC ₅₀ (nM)
1	2.1
3	0.8
4	1.9
5	7.4
15	2.1

25

These results show that compounds of the invention exhibit very potent activity in inducing differentiation of leukemia THP-1 cells, with EC₅₀ values in the low nanomolar range, which indicates that these compounds are particularly useful for the treatment or prevention of leukemias.

Previous reports of LSD1 have found that it is involved in cell proliferation and growth. Some studies have implicated LSD1 as a therapeutic target for cancer. Huang *et al.* (2007) *PNAS* 104:8023-8028 found that polyamine inhibitors of LSD1 modestly cause the reexpression of genes aberrantly silenced in cancer cells and particularly colorectal cancer (Huang *et al.* *Clin Cancer Res.* (2009) Dec 1;15(23):7217-28. Epub 2009 Nov 24. PMID: 19934284). Scoumanne *et al.* ((2007) *J. Biol. Chem.* May 25;282(21):15471-5) found that deficiency in LSD1 leads to a partial cell cycle arrest in G2/M and sensitizes cells to growth suppression induced by DNA damage. Kahl *et al.* ((2006) *Cancer Res.* 66(23):11341-7.) found that LSD1 expression is correlated with prostate cancer aggressiveness. Metzger *et al.* ((2005) *Nature*, 437 (7057), 436-439) reported that LSD1 modulation by siRNA and pargyline regulates androgen receptor (AR) and may have therapeutic potential in cancers where AR plays a role, like prostate, testis, and brain cancers. Lee *et al.* ((2006) *Chem. Biol.* 13:563-567) reported that tranylcypromine derepresses Egr-1 gene expression in some cancer lines. A body of evidence is accumulating that Egr-1 is a tumor suppressor gene in many contexts (see e.g., Calogero *et al.* (2004) *Cancer Cell International* 4:1 exogenous expression of EGR-1 resulted in growth arrest and eventual cell death in primary cancer cell lines; Lucerna *et al.* (2006) *Cancer Research* 66 (13), 6708-6713 show that sustained expression of Egr-1 causes antiangiogenic effects and inhibits tumor growth in some models; Ferraro *et al.* ((2005) *J. Clin. Oncol.* Mar 20;23(9):1921-6) reported that Egr-1 is downregulated in lung cancer patients with a higher risk of recurrence and may be more resistant to therapy. Thus, increasing Egr-1 expression via inhibition of LSD1 is a therapeutic approach for some cancers. Recent studies have also implicated LSD1 in brain cancer (Schulte *et al.* (2009) *Cancer Res.* Mar 1;69(5):2065-71). Other studies have implicated LSD1 in breast cancer (Lim *et al.* *Carcinogenesis*, (2010), 31(3): 512-20, Epub 2009 Dec 30. [Epub ahead of print] PMID: 20042638), lung, bladder and colorectal cancers (Hayami *et al* (2011), *Int J Cancer*, 128(3): 574-86, PMID:20333681) and leukemia (Binda *et al* (2010), *J Am Chem Soc*, 132(19): 6827-33, PMID:20415477).

Thus, a body of evidence has implicated LSD1 in a number of cancers, which suggests that LSD1 is a therapeutic target for cancer. The instant inventors have discovered a class of LSD1 inhibitors that can be used to treat diseases where LSD1 is implicated as a therapeutic target like cancer. Accordingly, the (hetero)aryl cyclopropylamine compounds of the invention can be used to treat such diseases.

Recent studies have also implicated LSD1 in viral infection and reactivation. In particular it was shown that pharmacological inhibitors of LSD1 like parnate and siRNA knock down of LSD1 caused reduced viral infectivity and reduced reactivation after latency (Liang *et al.* (2009) *Nat. Med.* 15(11):1312-1317). Therefore it is believed that the compounds of the invention can be used for treating or preventing viral infection. Furthermore, it is believed that the compounds of the invention can treat or prevent viral reactivation after latency.

Thus, without being bound by theory, the inventors have identified a new class of cyclopropanamine-based LSD1 inhibitors with unexpected potency and selectivity for LSD1, a biologically relevant target in oncology and other diseases.

5


All publications and patent applications mentioned in the specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The mere mentioning of the publications and patent applications does not necessarily constitute an admission that they are prior art to the instant application.

10

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

CLAIMS

1. A compound of Formula I



wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

B is H, R¹ or —L¹—E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, —O—, —NH—, —N(C₁₋₄ alkyl)—, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,
wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,
wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;

or a salt or solvate thereof

for use as a medicament.

2. The compound of claim 1 wherein L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

3. The compound of claim 2, wherein D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

4. The compound of claim 2, wherein D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

5. The compound of claim 2, wherein D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

6. The compound of claim 2, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³.

7. The compound of claim 2, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein ring D is optionally substituted with one or more R³.

8. The compound of claim 2, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups,

wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and

wherein ring D is optionally substituted with one or more R³.

9. The compound of claim 2, wherein D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

10. The compound of claim 2, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and wherein ring D is optionally substituted with one or more R³.

11. The compound of claim 2, wherein D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and wherein said piperidinyl is optionally substituted with one or more R³.

12. The compound of claim 2, wherein D is 4-piperidinyl optionally substituted with one or more R³.

13. The compound of claim 2, wherein D is a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.


14. The compound of claim 2, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.

15. The compound of claim 2, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 or 2 N atoms,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and wherein D is optionally substituted with one or more R³.

16. The compound of claim 2, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 N atom, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and wherein D is optionally substituted with one or more R³.

17. The compound of claim 2, wherein D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)



wherein D is optionally substituted with one or more R³.

18. The compound of any of claims 2 to 17, wherein said substituent(s) R³, if present, are placed on ring C atom(s) of D.

19. The compound of any of claims 2 to 18, wherein each R³ is independently selected from C₁₋₈ alkyl, halo, C₁₋₈ alkoxy, hydroxyl, amino, and amido.

20. The compound of any of claims 2 to 17, wherein D does not comprise any substituents R³.

21. The compound of any of claims 2 to 20, wherein A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

22. The compound of any of claims 2 to 20, wherein A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.
23. The compound of any of claims 2 to 20, wherein A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said phenyl, said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.
24. The compound of any of claims 2 to 20, wherein A is phenyl, pyridyl or thiazolyl, wherein said phenyl, said pyridyl or said thiazolyl is optionally substituted with one or more R¹.
25. The compound of any of claims 2 to 20, wherein A is phenyl, 3-pyridyl or 5-thiazolyl, wherein said phenyl, said 3-pyridyl or said 5-thiazolyl is optionally substituted with one or more R¹.
26. The compound any of claims 2 to 20, wherein A is phenyl or 3-pyridyl, wherein said phenyl or said 3-pyridyl is optionally substituted with one or more R¹.
27. The compound any of claims 2 to 20, wherein A is phenyl optionally substituted with one or more R¹.
28. The compound any of claims 2 to 20, wherein A is naphthyl optionally substituted with one or more R¹.
29. The compound any of claims 2 to 20, wherein A is heteroaryl optionally substituted with one or more R¹.
30. The compound any of claims 2 to 20, wherein A is monocyclic heteroaryl optionally substituted with one or more R¹.
31. The compound any of claims 2 to 20, wherein A is 3-pyridyl optionally substituted with one or more R¹.
32. The compound any of claims 2 to 20, wherein A is 5-thiazolyl optionally substituted with one or more R¹.
33. The compound of any of claims 2 to 32, wherein B is hydrogen or R¹.
34. The compound of claim 33, wherein B is hydrogen.

35. The compound of any of claims 2 to 32, wherein B is $-L^1-E$.

36. The compound of any of claims 2 to 35, wherein each R^1 is independently selected from C_{1-8} alkyl, amino, amido, hydroxyl, halo, $haloC_{1-8}$ alkyl, $haloC_{1-8}$ alkoxy, cyano, sulfonamide, C_{1-8} alkoxy, acyl, carboxyl, carbamate, and urea.

37. The compound of any of claims 2 to 35, wherein each R^1 is independently selected from halo, C_{1-4} alkyl, $haloC_{1-4}$ alkyl, C_{1-4} alkoxy and C_{3-6} cycloalkyl.

38. The compound of claim 34 or 35, wherein A is not substituted with any R^1 .

39. The compound of any of claims 2 to 32 and 35 to 38, wherein L^1 is a bond, $-O-$, $-NH-$, $-CH_2-NH-$, or $-CH_2-O-$, wherein said $-CH_2-NH-$ and $-CH_2-O-$ groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-NH-$ and $-CH_2-O-$ groups.

40. The compound of claim 39, wherein L^1 is a bond.

41. The compound of claim 39, wherein L^1 is $-CH_2-O-$ linked to ring A through the O atom comprised in said $-CH_2-O-$ and linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-O-$.

42. The compound of any of claims 2 to 32 and 35 to 38, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4}$ alkyl) $-$, C_{1-4} alkylene, or $-CH_2-NH-$.

43. The compound of any of claims 2 to 32 and 35 to 38, wherein L^1 is $-NH-$ or $-CH_2-NH-$, wherein said $-CH_2-NH-$ is linked to ring A through the N atom comprised in said $-CH_2-NH-$ and is linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-NH-$.

44. The compound of any of claims 2 to 32 and 35 to 43, wherein E is phenyl which is optionally substituted with one or more R^2 .

45. The compound of any of claims 2 to 32 and 35 to 43, wherein E is phenyl which is substituted with one R^2 .

46. The compound of any of claims 2 to 32 and 35 to 43, wherein E is heteroaryl which is optionally substituted with one or more R^2 .

47. The compound of claim 46, wherein E is monocyclic heteroaryl which is optionally substituted with one or more R².

48. The compound of claim 46, wherein E is pyridinyl which is optionally substituted with one or more R².

49. The compound of any of claims 2 to 32 and 35 to 48, wherein each R² is independently selected from C₁₋₈ alkyl, cyclol, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, N-sulfonamido and C₁₋₈ alkoxy.

50. The compound of any of claims 2 to 32 and 35 to 48, wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

51. The compound of claim 50, wherein each R² is independently selected from hydroxyl, halo and haloC₁₋₈ alkyl.

52. The compound of any of claims 2 to 32 and 35 to 48, wherein ring E is substituted with one R² and said R² is N-sulfonamido.

53. The compound of any of claims 2 to 32 and 35 to 43, wherein E is phenyl.

54. The compound of any of claims 2 to 32 and 35 to 43, wherein E is heteroaryl.

55. The compound of any of claims 2 to 32, wherein B is hydrogen and A is not substituted with any R¹.

56. The compound of any of claims 2 to 32, wherein B is hydrogen and each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.

57. The compound of any of claims 2 to 32, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is aryl which is optionally substituted with one or more R².

58. The compound of any of claims 2 to 32, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R².

59. The compound of any of claims 2 to 32, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R², wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

60. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

61. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is aryl or heteroaryl.

62. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, C_{1-4} alkylene, or $-CH_2-NH-$, and wherein E is phenyl which is optionally substituted with one or more R^2 .

63. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, C_{1-4} alkylene, or $-CH_2-NH-$, and wherein E is monocyclic heteroaryl which is optionally substituted with one or more R^2 .

64. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, C_{1-4} alkylene, or $-CH_2-NH-$, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

65. The compound of any of claims 2 to 32, wherein B is $-L^1-E$, wherein L^1 is $-NH-$ or $-NH-CH_2-$, wherein the group $-NH-CH_2-$ is linked to ring A through the N atom and to ring E through the $-CH_2-$ group, and wherein E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R^2 .

66. The compound of any of claims 2 to 65, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, fluoro and C_{1-4} alkyl.

67. The compound of any of claims 2 to 65, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen and fluoro.

68. The compound of any of claims 2 to 65, wherein R^w is selected from hydrogen, halo and C_{1-4} alkyl and each R^x , R^y and R^z is hydrogen.

69. The compound of any of claims 2 to 65, wherein each R^w , R^x , R^y and R^z is hydrogen.

70. The compound of any of claims 2 to 65, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl with the proviso that at least one of R^w , R^x , R^y and R^z is not hydrogen.

71. The compound of any of claims 2 to 65, wherein R^w is selected from halo and C₁₋₄ alkyl and each R^x, R^y and R^z is hydrogen.

72. The compound of claim 71, wherein R^w is methyl and each R^x, R^y and R^z is hydrogen.

73. The compound of any of claims 2 to 65, wherein R^w is fluoro and each R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl.

74. The compound of any of claims 2 to 65, wherein R^z is fluoro and each R^w, R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

75. The compound of any of claims 2 to 65, wherein R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

76. The compound of any of claims 2 to 75, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration.

77. The compound of claim 1 or 2, wherein said compound is selected from:

N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;

N-((1S,2R)-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-phenylcyclopropyl)piperidin-4-amine;

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine;

N-((trans)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-phenylcyclopropyl)piperidin-3-amine;

N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;

N-((trans)-2-phenylcyclopropyl)pyrrolidin-3-amine;

N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;

N-((trans)-2-phenylcyclopropyl)azetidin-3-amine;

N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;

N-((trans)-2-phenylcyclopropyl)azepan-3-amine;

N-((trans)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((trans)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((trans)-2-phenylcyclopropyl)decahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((trans)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine;
N-((trans)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine;
N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine ;
N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-(6-methoxy-4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(methylsulfonyl)-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;
1-(4-(((trans)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-(((trans)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((trans)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
2,2,6,6-tetramethyl-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;
1-isopropyl-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)piperidin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-(((trans)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;
N-((trans)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;

N-((1R,2R)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((trans)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine; N-((trans)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
or a salt or solvate thereof.

78. The compound of claim 1, wherein L^2 is C_{1-4} alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO_2 groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R^3 .

79. The compound of claim 78, wherein L^2 is linear C_{1-4} alkylene.

80. The compound of claim 78, wherein L^2 is $-CH_2-$.

81. The compound of claim 78, wherein L^2 is $-CH_2CH_2-$.

82. The compound of any of claims 78 to 81, wherein D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

83. The compound of any of claims 78 to 81, wherein D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 2 N atoms,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

84. The compound of any of claims 78 to 81, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

85. The compound of any of claims 78 to 81, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

86. The compound of any of claims 78 to 81, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

87. The compound of any of claims 78 to 81, wherein D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

88. The compound of any of claims 78 to 81, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and wherein ring D is optionally substituted with one or more R³.

89. The compound of any of claims 78 to 81, wherein D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and wherein said piperidinyl is optionally substituted with one or more R³.

90. The compound of any of claims 78 to 81, wherein D is 4-piperidinyl optionally substituted with one or more R³.

91. The compound of any of claims 78 to 81, wherein D is 3-azetidinyl optionally substituted with one or more R³.

92. The compound of any of claims 78 to 81, wherein D is a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.


93. The compound of any of claims 78 to 81, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 or 2 N atoms,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and wherein D is optionally substituted with one or more R³.

94. The compound of any of claims 78 to 81, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 N atom,

wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and wherein D is optionally substituted with one or more R³.


95. The compound of any of claims 78 to 81, wherein D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)



(c)



(d)

wherein D is optionally substituted with one or more R³.

96. The compound of any of claims 78 to 95, wherein said substituent(s) R³, if present, are placed on ring C atom(s) of D.

97. The compound of any of claims 78 to 96, wherein each R³ is independently selected from C₁₋₈ alkyl, halo, C₁₋₈ alkoxy, hydroxyl, amino, and amido.

98. The compound of any of claims 78 to 95, wherein D does not comprise any substituents R³.

99. The compound of any of claims 78 to 98, wherein A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

100. The compound of any of claims 78 to 98, wherein A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

101. The compound of any of claims 78 to 98, wherein A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said phenyl, said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.

102. The compound of any of claims 78 to 98, wherein A is phenyl, pyridyl or thiazolyl, wherein said phenyl, said pyridyl or said thiazolyl is optionally substituted with one or more R¹.

103. The compound of any of claims 78 to 98, wherein A is phenyl, 3-pyridyl or 5-thiazolyl, wherein said phenyl, said 3-pyridyl or said 5-thiazolyl is optionally substituted with one or more R¹.

104. The compound any of claims 78 to 98, wherein A is phenyl or 3-pyridyl, wherein said phenyl or said 3-pyridyl is optionally substituted with one or more R¹.

105. The compound any of claims 78 to 98, wherein A is phenyl optionally substituted with one or more R¹.

106. The compound any of claims 78 to 98, wherein A is naphthyl optionally substituted with one or more R¹.

107. The compound any of claims 78 to 98, wherein A is heteroaryl optionally substituted with one or more R¹.

108. The compound any of claims 78 to 98, wherein A is monocyclic heteroaryl optionally substituted with one or more R¹.

109. The compound any of claims 78 to 98, wherein A is 3-pyridyl optionally substituted with one or more R¹.

110. The compound any of claims 78 to 98, wherein A is 5-thiazolyl optionally substituted with one or more R¹.

111. The compound of any of claims 78 to 110, wherein B is hydrogen or R¹.
112. The compound of claim 111, wherein B is hydrogen.
113. The compound of any of claims 78 to 110, wherein B is –L¹-E.
114. The compound of any of claims 78 to 113, wherein each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈alkoxy, cyano, sulfonamide, C₁₋₈alkoxy, acyl, carboxyl, carbamate, and urea.
115. The compound of any of claims 78 to 113, wherein each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.
116. The compound of claim 112 or 113, wherein A is not substituted with any R¹.
117. The compound of any of claims 78 to 110 and 113 to 116, wherein L¹ is a bond, -O-, -NH-, -CH₂NH-, or -CH₂O-, wherein said -CH₂NH- and -CH₂O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the –CH₂- group comprised in said -CH₂NH- and -CH₂O- groups.
118. The compound of claim 117, wherein L¹ is a bond.
119. The compound of claim 117, wherein L¹ is –CH₂O- linked to ring A through the O atom comprised in said -CH₂O- and linked to ring E through the –CH₂- group comprised in said -CH₂O-.
120. The compound of any of claims 78 to 110 and 113 to 116, wherein L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or –CH₂NH-.
121. The compound of any of claims 78 to 110 and 113 to 116, wherein L¹ is –NH- or –CH₂NH-, wherein said –CH₂NH- is linked to ring A through the N atom comprised in said –CH₂NH- and is linked to ring E through the –CH₂- group comprised in said –CH₂NH-.
122. The compound of any of claims 78 to 110 and 113 to 121, wherein E is phenyl which is optionally substituted with one or more R².
123. The compound of any of claims 78 to 110 and 113 to 121, wherein E is phenyl which is substituted with one R².

124. The compound of any of claims 78 to 110 and 113 to 121, wherein E is heteroaryl which is optionally substituted with one or more R².

125. The compound of claim 124, wherein E is monocyclic heteroaryl which is optionally substituted with one or more R².

126. The compound of claim 124, wherein E is pyridinyl which is optionally substituted with one or more R².

127. The compound of any of claims 78 to 110 and 113 to 126, wherein each R² is independently selected from C₁₋₈ alkyl, cyclyl, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, N-sulfonamido and C₁₋₈ alkoxy

128. The compound of any of claims 78 to 110 and 113 to 126, wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

129. The compound of claim 128, wherein each R² is independently selected from hydroxyl, halo and haloC₁₋₈ alkyl.

130. The compound of any of claims 78 to 110 and 113 to 126, wherein ring E is substituted with one R² and said R² is N-sulfonamido.

131. The compound of any of claims 78 to 110 and 113 to 121, wherein E is phenyl.

132. The compound of any of claims 78 to 110 and 113 to 121, wherein E is heteroaryl.

133. The compound of any of claims 78 to 110, wherein B is hydrogen and A is not substituted with any R¹.

134. The compound of any of claims 78 to 110, wherein B is hydrogen and each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.

135. The compound of any of claims 78 to 110, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is aryl which is optionally substituted with one or more R².

136. The compound of any of claims 78 to 110, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R².

137. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is phenyl which is optionally substituted with one or more R^2 , wherein each R^2 is independently selected from hydroxyl, halo, halo C_{1-8} alkyl and N-sulfonamido.

138. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

139. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is aryl or heteroaryl.

140. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is phenyl which is optionally substituted with one or more R^2 .

141. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is monocyclic heteroaryl which is optionally substituted with one or more R^2 .

142. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

143. The compound of any of claims 78 to 110, wherein B is $-L^1-E$, wherein L^1 is $-NH-$ or $-NH-CH_2-$, wherein the group $-NH-CH_2-$ is linked to ring A through the N atom and to ring E through the $-CH_2-$ group, and wherein E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R^2 .

144. The compound of any of claims 78 to 143, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, fluoro and C_{1-4} alkyl.

145. The compound of any of claims 78 to 143, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen and fluoro.

146. The compound of any of claims 78 to 143 wherein R^w is selected from hydrogen, halo and C_{1-4} alkyl and each R^x , R^y and R^z is hydrogen.

147. The compound of any of claims 78 to 143, wherein each R^w , R^x , R^y and R^z is hydrogen.

148. The compound of any of claims 78 to 143 wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl with the proviso that at least one of R^w, R^x, R^y and R^z is not hydrogen.

149. The compound of any of claims 78 to 143 wherein R^w is selected from halo and C₁₋₄ alkyl, preferably fluoro and methyl, and each R^x, R^y and R^z is hydrogen.

150. The compound of claim 149 wherein R^w is methyl and each R^x, R^y and R^z is hydrogen.

151. The compound of any of claims 78 to 143, wherein R^w is fluoro and each R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl.

152. The compound of any of claims 78 to 143, wherein R^z is fluoro and each R^w, R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

153. The compound of any of claims 78 to 143, wherein R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

154. The compound of any of claims 78 to 153, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration.

155. The compound of claim 1 or 78, wherein said compound is selected from:

(Trans)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;

(Trans)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

(Trans)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;

(Trans)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(Trans)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(Trans)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

(1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;

(1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;

(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

(1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine ;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine;
(*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(*Trans*)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(*Trans*)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine; or a salt or solvate thereof.

156. The compound of any of claims 1 to 155, wherein said compound is an optically active stereoisomer.

157. A pharmaceutical composition comprising the compound of any of claims 1 to 156 and a pharmaceutically acceptable carrier.

158. The compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157 for use in the treatment or prevention of cancer.

159. The compound of claim 158 or the pharmaceutical composition of claim 158, wherein said cancer is chosen from breast cancer, lung cancer, prostate cancer, colorectal cancer, brain cancer, skin cancer, blood cancer, leukemia, lymphoma and myeloma.

160. The compound of claim 159 or the pharmaceutical composition of claim 159, wherein said cancer is blood cancer.

161. The compound of claim 159 or the pharmaceutical composition of claim 159, wherein said cancer is leukemia.

162. The compound of claim 159 or 161 or the pharmaceutical composition of claim 159 or 161, wherein said leukemia is chosen from acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), and hairy cell leukemia.

163. The compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157 for use in the treatment or prevention of a neurological disease.

164. The compound of claim 163 or the pharmaceutical composition of claim 163, wherein said neurological disease is selected from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, Dementia with Lewy Bodies, or Frontotemporal Dementia.

165. The compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157 for use in the treatment or prevention of a viral infection.

166. The compound of claim 165 or the pharmaceutical composition of claim 165, wherein said viral infection is a herpesvirus infection.

167. The compound of claim 166 or the pharmaceutical composition of claim 166, wherein said herpesvirus infection is caused by and/or associated with a herpesvirus chosen from HSV-1, HSV-2 and Epstein-Barr virus.

168. The compound of claim 165 or the pharmaceutical composition of claim 165, wherein said viral infection is caused by and/or associated with HIV.

169. The compound of claim 165 or the pharmaceutical composition of claim 165, wherein said viral infection is caused by and/or associated with a *Hepadnavirus*.

170. The compound of claim 169 or the pharmaceutical composition of claim 169, wherein said *Hepadnavirus* is Hepatitis B virus.

171. The compound of claim 165 or the pharmaceutical composition of claim 165, wherein said viral infection is caused by and/or associated with a *Flavivirus*.

172. The compound of claim 171 or the pharmaceutical composition of claim 171, wherein said *Flavivirus* is chosen from Hepatitis C virus (HCV), yellow fever virus, West Nile virus, Dengue virus and Japanese encephalitis virus.

173. The compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157 for use in the treatment or prevention of viral reactivation after latency.

174. The compound of claim 173 or the pharmaceutical composition of claim 173, wherein the virus that is reactivating is a herpesvirus.

175. The compound of claim 174 or the pharmaceutical composition of claim 174, wherein said herpesvirus is chosen from HSV-1, HSV-2 and Epstein-Barr virus.

176. The compound of claim 173 or the pharmaceutical composition of claim 173, wherein the virus that is reactivating is HIV.

177. A method of treating or preventing a disease, the method comprising administering, to a subject in need of such treatment or prevention, the compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157.

178. A method of treating or preventing cancer, the method comprising administering, to a subject in need of such treatment or prevention, the compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157.

179. The method of claim 178, wherein said cancer is chosen from breast cancer, lung cancer, prostate cancer, colorectal cancer, brain cancer, skin cancer, blood cancer, leukemia, lymphoma and myeloma.

180. The method of claim 179, wherein said cancer is blood cancer.
181. The method of claim 179, wherein said cancer is leukemia.
182. The method of claim 179 or 181, wherein said leukemia is chosen from acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, chronic eosinophilic leukemia, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), and hairy cell leukemia.
183. A method of treating or preventing a neurological disease, the method comprising administering, to a subject in need of such treatment or prevention, the compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157.
184. The method of claim 183, wherein said neurological disease is selected from depression, Alzheimer's disease, Huntington disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, Dementia with Lewy Bodies, or Frontotemporal Dementia.
185. A method of treating or preventing a viral infection, the method comprising administering, to a subject in need of such treatment or prevention, the compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157.
186. The method of claim 185, wherein said viral infection is a herpesvirus infection.
187. The method of claim 186, wherein said herpesvirus infection is caused by and/or associated with a herpesvirus chosen from HSV-1, HSV-2 and Epstein-Barr virus.
188. The method of claim 185, wherein said viral infection is caused by and/or associated with HIV.
189. The method of claim 185, wherein said viral infection is caused by and/or associated with a *Hepadnavirus*.
190. The method of claim 189, wherein said *Hepadnavirus* is Hepatitis B virus.
191. The method of claim 185, wherein said viral infection is caused by and/or associated with a *Flavivirus*.
192. The method of claim 191, wherein said *Flavivirus* is chosen from Hepatitis C virus (HCV), yellow fever virus, West Nile virus, Dengue virus and Japanese encephalitis virus.

193. A method for treating or preventing viral reactivation after latency, the method comprising administering, to a subject in need of such treatment or prevention, the compound of any of claims 1 to 156 or the pharmaceutical composition of claim 157.


194. The method of claim 193, wherein the virus that is reactivating is a herpesvirus.

195. The method of claim 194, wherein said herpesvirus is chosen from HSV-1, HSV-2 and Epstein-Barr virus.

196. The method of claim 193, wherein the virus that is reactivating is HIV.

197. The method of any of claims 177 to 196, wherein said subject is a human.

198. A compound of Formula I



wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

B is H, R¹ or —L¹—E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, —O—, —NH—, —N(C₁₋₄ alkyl)—, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;

or a salt or solvate thereof;


with the proviso that the following compounds are excluded:

1-(1-methylethyl)-N-(2-phenylcyclopropyl)- 3-Pyrrolidinamine;
4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid 1,1-dioxide;
N-(2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)- 4-Piperidinamine;
1-(3-methyl-2-buten-1-yl)-N-(2-phenylcyclopropyl)- 4-Piperidinamine;
4-[(2-phenylcyclopropyl)amino]-1-Piperidinecarboxylic acid ethyl ester;
1-[4-[(2-phenylcyclopropyl)amino]-1-piperidinyl]- ethanone;
hexahydro-3-[(2-phenylcyclopropyl)amino]- 2H-azepin-2-one;
1-cyclopropyl-3-[(2-phenylcyclopropyl)amino]- 2,5-pyrrolidinedione;
3-[(2-phenylcyclopropyl)amino]-1-propyl-2,5-pyrrolidinedione;
1-(1-methylethyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;
1-(1-methylpropyl)-3-[(2-phenylcyclopropyl)amino]- 2,5-Pyrrolidinedione;
1,2,5-trimethyl-N-(2-phenylcyclopropyl)- 4-Piperidinamine;
3-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;
4-[(2-phenylcyclopropyl)amino]- 1-piperidinecarboxamide;
3-hydroxy-4-((2-phenylcyclopropyl)amino)tetrahydrothiophene 1,1-dioxide;
tetrahydro-4-[(2-phenylcyclopropyl)amino]- 2H-pyran-4-carboxylic acid;
4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;
N-(2-phenylcyclopropyl)-1-(2-propyn-1-yl)- 4-piperidinamine;
1-ethyl-N-(2-phenylcyclopropyl)- 4-piperidinamine;
1-ethyl-3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;
4-((2-phenylcyclopropyl)amino)tetrahydro-2H-thiopyran-4-carboxylic acid;
N-(2-phenylcyclopropyl)-1-propyl-4-piperidinamine;
3-[(2-phenylcyclopropyl)amino]-2,5-Pyrrolidinedione;
tetrahydro-3-[(2-phenylcyclopropyl)amino]- 3-Thiophenecarboxylic acid;
tetrahydro-N-(2-phenylcyclopropyl)- 2H-Thiopyran-4-amine 1-oxide;
1-(1-methylethyl)-N-(2-phenylcyclopropyl)-4-piperidinamine;
1-methyl-3-[(2-phenylcyclopropyl)amino]-2,5-pyrrolidinedione;
tetrahydro-3-[(2-phenylcyclopropyl)amino]-2H-thiopyran-3-carboxylic acid;
tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-amine;
N-(2-phenylcyclopropyl)-3-piperidinamine;
tetrahydro-4-[(2-phenylcyclopropyl)amino]-3-furanol;
N-(2-phenylcyclopropyl)-4-piperidinamine;
tetrahydro-N-(2-phenylcyclopropyl)-2H-thiopyran-4-amine;
tetrahydro-N-(2-phenylcyclopropyl)-3-thiophenamine;

1-methyl-N-(2-phenylcyclopropyl)-4-piperidinamine;
N-(2-phenylcyclopropyl)-1,4-dioxaspiro[4.5]decan-8-amine;
8-methyl-N-(2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-(2-phenylcyclopropyl)-1-azabicyclo[2.2.2]octan-3-amine;
6-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
8-fluoro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzothiopyran-4-amine;
5-amino-1,3-dihydro-6-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
6-bromo-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-1-benzopyran-4-amine;
2,3-dihydro-3-[(2-phenylcyclopropyl)amino]-6-benzofuranol;
2,3-dihydro-N-(2-phenylcyclopropyl)-benzo[b]thiophen-3-amine 1,1-dioxide;
2,3-dihydro-N6-(2-phenylcyclopropyl)-1,4-benzodioxin-6,7-diamine;
3,4-dihydro-N-(2-phenylcyclopropyl)-1H-2-benzothiopyran-4-amine;
7-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-1-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-7-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
7-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
4-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
4-bromo-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
1,3-dihydro-5-methyl-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
5-fluoro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
6-chloro-1,3-dihydro-3-[(2-phenylcyclopropyl)amino]-2H-indol-2-one;
3-chloro-3,4-dihydro-N-(2-phenylcyclopropyl)-2H-thieno[3,2-b]pyrrol-5-amine;
8-methoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-methyl-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-ethoxy-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-fluoro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;
8-chloro-N-(2-phenylcyclopropyl)-5H-pyrimido[5,4-b]indol-4-amine;

5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione;
 1-ethyl- α -methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
 1-ethyl- α -methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 α -methyl-N-(2-phenylcyclopropyl)-1-propyl-3-piperidinemethanamine;
 tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-2-methanamine;
 octahydro-N-(2-phenylcyclopropyl)-1H-indole-2-methanamine;
 α -methyl-N-(2-phenylcyclopropyl)-1-propyl-4-piperidinemethanamine;
 3-methyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
 N-(2-phenylcyclopropyl)-2-piperidinemethanamine;
 N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 $\alpha,1$ -dimethyl-N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
 N-(2-phenylcyclopropyl)-3-piperidinemethanamine;
 4-methoxy-N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
 N-(2-phenylcyclopropyl)-3-propyl-3-pyrrolidinemethanamine;
 N-(2-phenylcyclopropyl)-2-morpholinemethanamine;
 4-ethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 4-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 $\alpha,1$ -dimethyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 1-methyl-N-(2-phenylcyclopropyl)-4-piperidinemethanamine;
 tetrahydro-N-(2-phenylcyclopropyl)-2H-pyran-4-methanamine;
 3-methyl-N-(2-phenylcyclopropyl)-3-pyrrolidinemethanamine;
 N-(2-phenylcyclopropyl)-2-pyrrolidinemethanamine;
 tetrahydro-N-(2-phenylcyclopropyl)-3-furanmethanamine;
 tetrahydro-N-(2-phenylcyclopropyl)-2-furanmethanamine;
 N-(2-phenylcyclopropyl)-4-piperidineethanamine;
 N-(2-phenylcyclopropyl)-2-pyrrolidineethanamine;
 1-methyl-N-(2-phenylcyclopropyl)-2-piperidineethanamine; and
 N-(2-phenylcyclopropyl)-3-piperidinepropanamine.

199. A compound of Formula I



wherein:

A is aryl optionally substituted with one or more R¹, and B is -L¹-E, or

A is heteroaryl optionally substituted with one or more R¹, and B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;


each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cycll, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;

or a salt or solvate thereof.

200. A compound of Formula I



wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R¹;

B is H, R¹ or -L¹-E;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R²;

L¹ is a bond, -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene or heteroC₁₋₄alkylene;

L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and


each R^w , R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl;

or a salt or solvate thereof;

wherein the substituents $-A-B$ and $-NH-L^2-D$ on the cyclopropyl moiety are in trans-configuration;

with the proviso that the compound 5-ethyl-5,10-dihydro-3,7,8,10-tetramethyl-4a-[(2-phenylcyclopropyl)amino]-benzo[g]pteridine-2,4(3H,4aH)-dione is excluded.

201. A compound of Formula I



wherein:

A is aryl or heteroaryl, wherein said aryl or heteroaryl is optionally substituted with one or more R^1 ;

B is H, R^1 or $-L^1-E$;

E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R^2 ;

L^1 is a bond, $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, C_{1-4} alkylene or hetero C_{1-4} alkylene;

L^2 is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO_2 groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³;

or L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and
- (iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups,

wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³;

each R¹ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R² is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea;

each R³ is independently selected from C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyclyl, amino, amido, hydroxyl, nitro, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, sulfinyl, sulfonyl, sulfonamide, C₁₋₈ alkoxy, acyl, carboxyl, O-carboxy, C-carboxy, carbamate and urea; and

each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl;

or a salt or solvate thereof;

wherein the compound of formula I is an optically active stereoisomer.

202. The compound of any of claims 198 to 201, wherein L² is a bond and D is a cyclic group selected from:

- (i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

203. The compound of any of claims 198 to 201, wherein L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) contains at least one N atom,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

204. The compound of any of claims 198 to 201, wherein L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and

(ii) a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 2 N atoms,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, and wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

205. The compound of any of claims 198 to 201, wherein L² is a bond and D is a cyclic group selected from:

(i) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(ii) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (i) or (ii) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (i) or (ii) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (i) or (ii), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (i) or (ii) is optionally substituted with one or more R³.

206. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

207. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

208. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

209. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

210. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and wherein ring D is optionally substituted with one or more R³.

211. The compound of any of claims 198 to 201, wherein L² is a bond and D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and wherein said piperidinyl is optionally substituted with one or more R³.

212. The compound of any of claims 198 to 201, wherein L² is a bond and D is 4-piperidinyl optionally substituted with one or more R³.


213. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 7- to 15-membered polycyclic ring system which comprises at least one saturated heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.

214. The compound of any of claims 198 to 201, wherein L² is a bond and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains from 1 to 4 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in D are optionally oxidized to form CO groups, wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein D is optionally substituted with one or more R³.


215. The compound of any of claims 198 to 201, wherein L^2 is a bond and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 or 2 N atoms,
 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
 wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and
 wherein D is optionally substituted with one or more R^3 .

216. The compound of any of claims 198 to 201, wherein L^2 is a bond and D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic ring system contains 1 N atom,
 wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
 wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and
 wherein D is optionally substituted with one or more R^3 .


217. The compound of any of claims 198 to 201, wherein L^2 is a bond and D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)



(c)



(d)

wherein D is optionally substituted with one or more R^3 .

218. The compound of any of claims 198 to 217, wherein said substituent(s) R^3 , if present, are placed on ring C atom(s) of D.

219. The compound of any of claims 198 to 218, wherein each R^3 is independently selected from C_{1-8} alkyl, halo, C_{1-8} alkoxy, hydroxyl, amino, and amido.

220. The compound of any of claims 198 to 217, wherein D does not comprise any substituents R³.

221. The compound of any of claims 198 to 220, wherein A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

222. The compound of any of claims 198 to 221, wherein A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

223. The compound of any of claims 198 to 222, wherein A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said phenyl, said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.

224. The compound of any of claims 198 to 222, wherein A is phenyl, pyridyl or thiazolyl, wherein said phenyl, said pyridyl or said thiazolyl is optionally substituted with one or more R¹.

225. The compound of any of claims 198 to 222, wherein A is phenyl, 3-pyridyl or 5-thiazolyl, wherein said phenyl, said 3-pyridyl or said 5-thiazolyl is optionally substituted with one or more R¹.

226. The compound of any of claims 198 to 222, wherein A is phenyl or 3-pyridyl, wherein said phenyl or said 3-pyridyl is optionally substituted with one or more R¹.

227. The compound of any of claims 198 to 222, wherein A is phenyl optionally substituted with one or more R¹.

228. The compound of any of claims 198 to 221, wherein A is naphthyl optionally substituted with one or more R¹.

229. The compound of any of claims 198 to 220, wherein A is heteroaryl optionally substituted with one or more R¹.

230. The compound of any of claims 198 to 221, wherein A is monocyclic heteroaryl optionally substituted with one or more R¹.

231. The compound of any of claims 198 to 222, wherein A is pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.

232. The compound of any of claims 198 to 222, wherein A is pyridyl or thiazolyl, wherein said pyridyl or said thiazolyl is optionally substituted with one or more R¹.

233. The compound of any of claims 198 to 222, wherein A is 3-pyridyl optionally substituted with one or more R¹.

234. The compound any of claims 198 to 222, wherein A is 5-thiazolyl optionally substituted with one or more R¹.

235. The compound of any of claims 198 to 234, wherein B is hydrogen or R¹.

236. The compound of claim 235, wherein B is hydrogen.

237. The compound of any of claims 198 to 234, wherein B is -L¹-E.

238. The compound of any of claims 198 to 237, wherein each R¹ is independently selected from C₁₋₈ alkyl, amino, amido, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈alkoxy, cyano, sulfonamide, C₁₋₈alkoxy, acyl, carboxyl, carbamate, and urea.

239. The compound of any of claims 198 to 237, wherein each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.

240. The compound of claim 236 or 237, wherein A is not substituted with any R¹.

241. The compound of any of claims 198 to 234 and 237 to 240, wherein L¹ is a bond, -O-, -NH-, -CH₂NH-, or -CH₂O-, wherein said -CH₂-NH- and -CH₂-O- groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the -CH₂- group comprised in said -CH₂-NH- and -CH₂-O- groups.

242. The compound of claim 241, wherein L¹ is a bond.

243. The compound of claim 241, wherein L¹ is -CH₂-O- linked to ring A through the O atom comprised in said -CH₂-O- and linked to ring E through the -CH₂- group comprised in said -CH₂-O-.

244. The compound of any of claims 198 to 234 and 237 to 240, wherein L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-.

245. The compound of any of claims 198 to 234 and 237 to 240, wherein L¹ is –NH- or –CH₂–NH–, wherein said –CH₂–NH– is linked to ring A through the N atom comprised in said –CH₂–NH– and is linked to ring E through the –CH₂– group comprised in said –CH₂–NH–.

246. The compound of any of claims 198 to 234 and 237 to 245, wherein E is phenyl which is optionally substituted with one or more R².

247. The compound of any of claims 198 to 234 and 237 to 245, wherein E is phenyl which is substituted with one R².

248. The compound of any of claims 198 to 234 and 237 to 245, wherein E is heteroaryl which is optionally substituted with one or more R².

249. The compound of claim 248, wherein E is monocyclic heteroaryl which is optionally substituted with one or more R².

250. The compound of claim 248, wherein E is pyridinyl which is optionally substituted with one or more R².

251. The compound of any of claims 198 to 234 and 237 to 250, wherein each R² is independently selected from C₁₋₈ alkyl, cyclol, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, N-sulfonamido and C₁₋₈ alkoxy.

252. The compound of any of claims 198 to 234 and 237 to 250, wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

253. The compound of claim 252, wherein each R² is independently selected from hydroxyl, halo and haloC₁₋₈ alkyl.

254. The compound of any of claims 198 to 234 and 237 to 250, wherein ring E is substituted with one R² and said R² is N-sulfonamido.

255. The compound of any of claims 198 to 234 and 237 to 245, wherein E is phenyl.

256. The compound of any of claims 198 to 234 and 237 to 245, wherein E is heteroaryl.

257. The compound of any of claims 198 to 234, wherein B is hydrogen and A is not substituted with any R¹.

258. The compound of any of claims 198 to 234, wherein B is hydrogen and each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.

259. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is aryl which is optionally substituted with one or more R².

260. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R².

261. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R², wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

262. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is pyridinyl which is optionally substituted with one or more R².

263. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is aryl or heteroaryl.

264. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-, and wherein E is phenyl which is optionally substituted with one or more R².

265. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-, and wherein E is monocyclic heteroaryl which is optionally substituted with one or more R².

266. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is -O-, -NH-, -N(C₁₋₄ alkyl)-, C₁₋₄ alkylene, or -CH₂-NH-, and wherein E is pyridinyl which is optionally substituted with one or more R².

267. The compound of any of claims 198 to 234, wherein B is -L¹-E, wherein L¹ is -NH- or -NH-CH₂-, wherein the group -NH-CH₂- is linked to ring A through the N atom and to ring E through the -CH₂- group, and wherein E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R².

268. The compound of any of claims 198 to 267, wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen, fluoro and C₁₋₄ alkyl.

269. The compound of any of claims 198 to 267, wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen and fluoro.

270. The compound of any of claims 198 to 267, wherein R^w is selected from hydrogen, halo and C₁₋₄ alkyl and each R^x, R^y and R^z is hydrogen.

271. The compound of any of claims 198 to 267, wherein each R^w, R^x, R^y and R^z is hydrogen.

272. The compound of any of claims 198 to 267, wherein each R^w, R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl with the proviso that at least one is not hydrogen.

273. The compound of any of claims 198 to 267, wherein R^w is selected from halo and C₁₋₄ alkyl, preferably fluoro and methyl, and each R^x, R^y and R^z is hydrogen.

274. The compound of claim 273, wherein R^w is methyl and each R^x, R^y and R^z is hydrogen.

275. The compound of any of claims 198 to 267, wherein R^w is fluoro and each R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl.

276. The compound of any of claims 198 to 267, wherein R^z is fluoro and each R^w, R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

277. The compound of any of claims 198 to 267 wherein R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

278. The compound of any of claims 198 to 267, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration.

279. The compound of claim 198 or 202, wherein said compound is selected from:

N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine;

N-((trans)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;

N-((trans)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;
N-((trans)-2-phenylcyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;
N-((trans)-2-phenylcyclopropyl)azetidin-3-amine;
N-((trans)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((trans)-2-phenylcyclopropyl)azepan-3-amine;
N-((trans)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((trans)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((trans)-2-phenylcyclopropyl)decahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine;
N-((trans)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((trans)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine;
N-((trans)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine;
N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((trans)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-(6-methoxy-4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((trans)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(methylsulfonyl)-N-((trans)-2-phenylcyclopropyl)piperidin-4-amine;

1-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((*trans*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
2,2,6,6-tetramethyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2H-thiopyran 1,1-dioxide;
N-((*trans*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine;N-((*trans*)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
or a salt or solvate thereof.

280. The compound of claim 199 or 202, wherein said compound is selected from:

N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2H-pyran-4-amine;
N-((*trans*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((1*S*,2*R*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(3'-(trifluoromethyl)-[1',1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;

N-(6-methoxy-4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
or a salt or solvate thereof.

281. The compound of claim 200 or 202, wherein said compound is selected from:

N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(6-(3-(trifluoromethyl)phenyl)pyridin-3-yl)cyclopropyl)tetrahydro-2*H*-pyran-4-amine;
N-((*trans*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)piperidin-3-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-3-amine;
N-((*trans*)-2-phenylcyclopropyl)pyrrolidin-3-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)pyrrolidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)pyrrolidin-3-amine;
N-((*trans*)-2-phenylcyclopropyl)azetidin-3-amine;
N-((*trans*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((*trans*)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((*trans*)-2-phenylcyclopropyl)azepan-3-amine;
N-((*trans*)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((*trans*)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((*trans*)-2-phenylcyclopropyl)decahydroquinolin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1,2,3,4-tetrahydroquinolin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((*trans*)-2-phenylcyclopropyl)-2-azaspiro[4.5]decan-8-amine;
N-((*trans*)-2-phenylcyclopropyl)-2,3-dihydrospiro[indene-1,4'-piperidin]-3-amine;
N-((1*S*,2*R*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;

N-((1*S*,2*R*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(2-(*o*-tolyl)cyclopropyl)piperidin-4-amine;
N-(2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-(2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-(2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-(2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-(6-methoxy-4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((*trans*)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(methylsulfonyl)-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-(4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((*trans*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2*H*-pyran-4-amine;
2,2,6,6-tetramethyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
1-isopropyl-N-((*trans*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(2,2,2-trifluoroethyl)piperidin-4-amine;
N-((*trans*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-((*trans*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2*H*-thiopyran 1,1-dioxide;
N-((*trans*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*S*,2*S*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-methyl-2-phenylcyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(*o*-tolyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((*trans*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
or a salt or solvate thereof.

282. The compound of claim 201 or 202, wherein said compound is selected from:

N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine;

N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)piperidin-4-amine ;
(S)-N-((1S,2R)-2-phenylcyclopropyl)piperidin-3-amine;
(R)-N-((1S,2R)-2-phenylcyclopropyl)piperidin-3-amine;
(S)-N-((1R,2S)-2-phenylcyclopropyl)piperidin-3-amine;
(R)-N-((1R,2S)-2-phenylcyclopropyl)piperidin-3-amine;
(S)-N-((1S,2R)-2-phenylcyclopropyl)pyrrolidin-3-amine;
(R)-N-((1S,2R)-2-phenylcyclopropyl)pyrrolidin-3-amine;
(S)-N-((1R,2S)-2-phenylcyclopropyl)pyrrolidin-3-amine;
(R)-N-((1R,2S)-2-phenylcyclopropyl)pyrrolidin-3-amine;
N-((1S,2R)-2-phenylcyclopropyl)azetidin-3-amine;
N-((1R,2S)-2-phenylcyclopropyl)azetidin-3-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)azetidin-3-amine;
N-((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((1S,2R)-2-(4-(benzyloxy)phenyl)cyclopropyl)azetidin-3-amine;
N-((1R,2S)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((1S,2R)-2-phenylcyclopropyl)-8-azabicyclo[3.2.1]octan-3-amine;
N-((1R,2S)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((1S,2R)-2-phenylcyclopropyl)-3-azabicyclo[3.2.1]octan-8-amine;
N-((1R,2S)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((1S,2R)-2-phenylcyclopropyl)-3-azaspiro[5.5]undecan-9-amine;
N-((1S,2R)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(pyridin-3-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2S)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2R)-2-(thiazol-5-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1R,2S)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropyl)piperidin-4-amine;
N-((1S,2R)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-((1R,2S)-2-phenylcyclopropyl)-7-azaspiro[3.5]nonan-2-amine;
N-(6-methoxy-4'-((1R,2S)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(6-methoxy-4'-((1S,2R)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)methanesulfonamide;
N-(4'-((1R,2S)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
N-(4'-((1S,2R)-2-(piperidin-4-ylamino)cyclopropyl)-[1,1'-biphenyl]-3-yl)propane-2-sulfonamide;
1-(4-(((1R,2S)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
1-(4-(((1S,2R)-2-(4-bromophenyl)cyclopropyl)amino)piperidin-1-yl)ethanone;
4-(((1R,2S)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;

4-(((1*S*,2*R*)-2-(4-bromophenyl)cyclopropyl)amino)piperidine-1-carboxamide;
N-((1*R*,2*S*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2*H*-pyran-4-amine;
N-((1*S*,2*R*)-2-(4-bromophenyl)cyclopropyl)tetrahydro-2*H*-pyran-4-amine;
2,2,6,6-tetramethyl-N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
2,2,6,6-tetramethyl-N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
1-methyl-N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
1-isopropyl-N-((1*R*,2*S*)-2-phenylcyclopropyl)piperidin-4-amine;
1-isopropyl-N-((1*S*,2*R*)-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
N-((1*S*,2*R*)-2-phenylcyclopropyl)-1-(pyridin-4-yl)piperidin-4-amine;
4-(((1*R*,2*S*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2*H*-thiopyran 1,1-dioxide;
4-(((1*S*,2*R*)-2-(4-bromophenyl)cyclopropyl)amino)tetrahydro-2*H*-thiopyran 1,1-dioxide;
N-((1*S*,2*S*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*R*)-2-fluoro-2-phenylcyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(naphthalen-2-yl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(o-tolyl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(2-fluorophenyl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(3,4-difluorophenyl)cyclopropyl)piperidin-4-amine;
N-((1*R*,2*S*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;
N-((1*S*,2*R*)-2-(4-methoxyphenyl)cyclopropyl)piperidin-4-amine;

or a salt or solvate thereof.

283. The compound of any of claims 198 to 282, wherein said compound is an optically active stereoisomer.

284. The compound of any of claims 198 to 201, wherein L² is C₁₋₄ alkylene and D is a cyclic group selected from:

- (iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

285. The compound of claim 284, wherein L² is linear C₁₋₄ alkylene.

286. The compound of claim 284, wherein L² is -CH₂-.

287. The compound of claim 284, wherein L² is -CH₂CH₂-.

288. The compound of any of claims 284 to 287, wherein D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,

wherein the cyclic group (iii) or (iv) contains at least one N atom,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, wherein one or more S atoms in the cyclic group (iii) or (iv), if present, are optionally oxidized to form independently SO groups or SO₂ groups, and

wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

289. The compound of any of claims 284 to 287, wherein D is a cyclic group selected from:

(iii) a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 N atoms, and

(iv) a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 2 N atoms,

wherein the cyclic group (iii) or (iv) is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in the cyclic group (iii) or (iv) are optionally oxidized to form CO groups, and wherein the cyclic group (iii) or (iv) is optionally substituted with one or more R³.

290. The compound of any of claims 284 to 287, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

291. The compound of any of claims 284 to 287, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one or more S atoms in ring D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and wherein ring D is optionally substituted with one or more R³.

292. The compound of any of claims 284 to 287, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

293. The compound of any of claims 284 to 287, wherein D is a 6-membered monocyclic saturated heterocyclic ring containing 1 heteroatom selected from N, O and S, wherein D is linked to the remainder of the compound of Formula I through a ring C atom, wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, wherein one S atom in ring D, if present, is optionally oxidized to form an SO group or an SO₂ group, and wherein ring D is optionally substituted with one or more R³.

294. The compound of any of claims 284 to 287, wherein D is a 3- to 7-membered monocyclic saturated heterocyclic ring containing 1 N atom, wherein D is linked to the remainder of the compound of Formula I through a ring C atom,

wherein one or more ring C atoms in ring D are optionally oxidized to form CO groups, and
wherein ring D is optionally substituted with one or more R³.

295. The compound of any of claims 284 to 287, wherein D is piperidinyl, wherein said piperidinyl is linked to the remainder of the compound of Formula I through a ring C atom, and
wherein said piperidinyl is optionally substituted with one or more R³.

296. The compound of any of claims 284 to 287, wherein D is 4-piperidinyl optionally substituted with one or more R³.


297. The compound of any of claims 284 to 287, wherein D is 3-azetidinyl optionally substituted with one or more R³.

298. The compound of any of claims 284 to 287, wherein D is a 7- to 15-membered polycyclic saturated ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains from 1 to 4 heteroatoms independently selected from N, O and S,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in D are optionally oxidized to form CO groups,
wherein one or more S atoms in D, if present, are optionally oxidized to form independently SO groups or SO₂ groups, and
wherein D is optionally substituted with one or more R³.


299. The compound of any of claims 284 to 287, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 or 2 N atoms,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and
wherein D is optionally substituted with one or more R³.

300. The compound of any of claims 284 to 287, wherein D is a 7- to 15-membered saturated polycyclic ring system which comprises at least one heterocyclic ring, wherein the polycyclic saturated ring system contains 1 N atom,
wherein D is linked to the remainder of the compound of Formula I through a ring C atom,
wherein one or more ring C atoms in D are optionally oxidized to form CO groups, and
wherein D is optionally substituted with one or more R³.


301. The compound of any of claims 284 to 287, wherein D is a 7- to 15-membered saturated polycyclic ring system selected from a group of formula (a), (b), (c) and (d)




(a)



(b)



(c)



(d)

wherein D is optionally substituted with one or more R³.

302. The compound of any of claims 284 to 301, wherein said substituent(s) R³, if present, are placed on ring C atom(s) of D.

303. The compound of any of claims 284 to 302, wherein each R³ is independently selected from C₁₋₈ alkyl, halo, C₁₋₈ alkoxy, hydroxyl, amino, and amido.

304. The compound of any of claims 284 to 301, wherein D does not comprise any substituents R³.

305. The compound of any of claims 284 to 304, wherein A is phenyl, naphthyl or monocyclic heteroaryl, wherein said phenyl, said naphthyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

306. The compound of any of claims 284 to 305, wherein A is phenyl or monocyclic heteroaryl, wherein said phenyl or said monocyclic heteroaryl is optionally substituted with one or more R¹.

307. The compound of any of claims 284 to 306, wherein A is phenyl, pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said phenyl, said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.

308. The compound of any of claims 284 to 306, wherein A is phenyl, pyridyl or thiazolyl, wherein said phenyl, said pyridyl or said thiazolyl is optionally substituted with one or more R¹.

309. The compound of any of claims 284 to 306, wherein A is phenyl, 3-pyridyl or 5-thiazolyl, wherein said phenyl, said 3-pyridyl or said 5-thiazolyl is optionally substituted with one or more R¹.

310. The compound of any of claims 284 to 306, wherein A is phenyl or 3-pyridyl, wherein said phenyl or said 3-pyridyl is optionally substituted with one or more R¹.

311. The compound of any of claims 284 to 306, wherein A is phenyl optionally substituted with one or more R¹.

312. The compound of any of claims 284 to 305, wherein A is naphthyl optionally substituted with one or more R¹.

313. The compound of any of claims 284 to 304, wherein A is heteroaryl optionally substituted with one or more R¹.

314. The compound of any of claims 284 to 306, wherein A is monocyclic heteroaryl optionally substituted with one or more R¹.

315. The compound of any of claims 284 to 306, wherein A is pyridyl, thiophenyl, pyrrolyl, furanyl, or thiazolyl, wherein said pyridyl, said thiophenyl, said pyrrolyl, said furanyl or said thiazolyl is optionally substituted with one or more R¹.

316. The compound of any of claims 284 to 306, wherein A is pyridyl or thiazolyl, wherein said pyridyl or said thiazolyl is optionally substituted with one or more R¹.

317. The compound of any of claims 284 to 306, wherein A is 3-pyridyl optionally substituted with one or more R¹.

318. The compound of any of claims 284 to 306, wherein A is 5-thiazolyl optionally substituted with one or more R¹.

319. The compound of any of claims 284 to 318, wherein B is hydrogen or R¹.

320. The compound of claim 319, wherein B is hydrogen.

321. The compound of any of claims 284 to 318, wherein B is $-L^1-E$.

322. The compound of any of claims 284 to 321, wherein each R^1 is independently selected from C_{1-8} alkyl, amino, amido, hydroxyl, halo, $haloC_{1-8}$ alkyl, $haloC_{1-8}$ alkoxy, cyano, sulfonamide, C_{1-8} alkoxy, acyl, carboxyl, carbamate, and urea.

323. The compound of any of claims 284 to 321, wherein each R^1 is independently selected from halo, C_{1-4} alkyl, $haloC_{1-4}$ alkyl, C_{1-4} alkoxy and C_{3-6} cycloalkyl.

324. The compound of claim 320 or 321, wherein A is not substituted with any R^1 .

325. The compound of any of claims 284 to 318 and 321 to 324, wherein L^1 is a bond, $-O-$, $-NH-$, $-CH_2-NH-$, or $-CH_2-O-$, wherein said $-CH_2-NH-$ and $-CH_2-O-$ groups are linked to ring A through the N or O atom, respectively, and are linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-NH-$ and $-CH_2-O-$ groups.

326. The compound of claim 325, wherein L^1 is a bond.

327. The compound of claim 325, wherein L^1 is $-CH_2-O-$ linked to ring A through the O atom comprised in said $-CH_2-O-$ and linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-O-$.

328. The compound of any of claims 284 to 318 and 321 to 324, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4}$ alkyl) $-$, C_{1-4} alkylene, or $-CH_2-NH-$.

329. The compound of any of claims 284 to 318 and 321 to 324, wherein L^1 is $-NH-$ or $-CH_2-NH-$, wherein said $-CH_2-NH-$ is linked to ring A through the N atom comprised in said $-CH_2-NH-$ and is linked to ring E through the $-CH_2-$ group comprised in said $-CH_2-NH-$.

330. The compound of any of claims 284 to 318 and 321 to 329, wherein E is phenyl which is optionally substituted with one or more R^2 .

331. The compound of any of claims 284 to 318 and 321 to 329, wherein E is phenyl which is substituted with one R^2 .

332. The compound of any of claims 284 to 318 and 321 to 329, wherein E is heteroaryl which is optionally substituted with one or more R^2 .

333. The compound of claim 332, wherein E is monocyclic heteroaryl which is optionally substituted with one or more R².

334. The compound of claim 332, wherein E is pyridinyl which is optionally substituted with one or more R².

335. The compound of any of claims 284 to 318 and 321 to 334 wherein each R² is independently selected from C₁₋₈ alkyl, cyclyl, hydroxyl, halo, haloC₁₋₈ alkyl, haloC₁₋₈ alkoxy, cyano, N-sulfonamido and C₁₋₈ alkoxy.

336. The compound of any of claims 284 to 318 and 321 to 334, wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

337. The compound of claim 336, wherein each R² is independently selected from hydroxyl, halo and haloC₁₋₈ alkyl.

338. The compound of any of claims 284 to 318 and 321 to 334, wherein ring E is substituted with one R² and said R² is N-sulfonamido.

339. The compound of any of claims 284 to 318 and 321 to 329, wherein E is phenyl.

340. The compound of any of claims 284 to 318 and 321 to 329, wherein E is heteroaryl.

341. The compound of any of claims 284 to 318, wherein B is hydrogen and A is not substituted with any R¹.

342. The compound of any of claims 284 to 318, wherein B is hydrogen and each R¹ is independently selected from halo, C₁₋₄ alkyl, haloC₁₋₄ alkyl, C₁₋₄ alkoxy and C₃₋₆ cycloalkyl.

343. The compound of any of claims 284 to 318, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is aryl which is optionally substituted with one or more R².

344. The compound of any of claims 284 to 318, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R².

345. The compound of any of claims 284 to 318, wherein B is -L¹-E, wherein L¹ is a bond, and wherein E is phenyl which is optionally substituted with one or more R², wherein each R² is independently selected from hydroxyl, halo, haloC₁₋₈ alkyl and N-sulfonamido.

346. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

347. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is a bond, and wherein E is aryl or heteroaryl.

348. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is phenyl which is optionally substituted with one or more R^2 .

349. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is monocyclic heteroaryl which is optionally substituted with one or more R^2 .

350. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is $-O-$, $-NH-$, $-N(C_{1-4} \text{ alkyl})-$, $C_{1-4} \text{ alkylene}$, or $-CH_2-NH-$, and wherein E is pyridinyl which is optionally substituted with one or more R^2 .

351. The compound of any of claims 284 to 318, wherein B is $-L^1-E$, wherein L^1 is $-NH-$ or $-NH-CH_2-$, wherein the group $-NH-CH_2-$ is linked to ring A through the N atom and to ring E through the $-CH_2-$ group, and wherein E is aryl or heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with one or more R^2 .

352. The compound of any of claims 284 to 351, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, fluoro and C_{1-4} alkyl.

353. The compound of any of claims 284 to 351, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen and fluoro.

354. The compound of any of claims 284 to 351, wherein R^w is selected from hydrogen, halo and C_{1-4} alkyl and each R^x , R^y and R^z is hydrogen.

355. The compound of any of claims 284 to 351, wherein each R^w , R^x , R^y and R^z is hydrogen.

356. The compound of any of claims 284 to 351, wherein each R^w , R^x , R^y and R^z is independently selected from hydrogen, halo and C_{1-4} alkyl with the proviso that at least one is not hydrogen.

357. The compound of any of claims 284 to 351, wherein R^w is selected from halo and C₁₋₄ alkyl, preferably fluoro and methyl, and each R^x, R^y and R^z is hydrogen.

358. The compound of claim 357, wherein R^w is methyl and each R^x, R^y and R^z is hydrogen.

359. The compound of any of claims 284 to 351, wherein R^w is fluoro and each R^x, R^y and R^z is independently selected from hydrogen, halo and C₁₋₄ alkyl.

360. The compound of any of claims 284 to 351, wherein R^z is fluoro and each R^w, R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

361. The compound of any of claims 284 to 351, wherein R^w and R^z are fluoro and each R^x and R^y is independently selected from hydrogen, halo and C₁₋₄ alkyl.

362. The compound of any of claims 284 to 361, wherein the substituents -A-B and -NH-L²-D on the cyclopropyl moiety are in trans-configuration.

363. The compound of claim 198 or 284, wherein said compound is selected from:

(Trans)-2-phenyl-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;

(Trans)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

(Trans)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(Trans)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;

(Trans)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(Trans)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;

(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;

(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;

(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine;
(*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(*Trans*)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(*Trans*)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine;
or a salt or solvate thereof.

364. The compound of claim 199 or 284, wherein said compound is selected from:

(*Trans*)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2H-pyran-4-yl)ethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(*Trans*)-2-(4-(1*H*-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine;
or a salt or solvate thereof.

365. The compound of claim 200 or 284, wherein said compound is selected from:

(*Trans*)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(2-(tetrahydro-2*H*-pyran-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-(4'-chloro-[1,1'-biphenyl]-4-yl)-N-(2-(tetrahydro-2*H*-pyran-4-yl)ethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(*Trans*)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(*Trans*)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine ;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;

(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(*Trans*)-2-phenyl-N-(pyrrolidin-3-ylmethyl)cyclopropanamine;
(*Trans*)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(*Trans*)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(*Trans*)-2-(4-(1*H*-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*Trans*)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-methyl-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine;
(*trans*)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine;
or a salt or solvate thereof.


366. The compound of claim 201 or 284, wherein said compound is selected from:

(1S,2R)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1R,2S)-2-phenyl-N-(piperidin-4-ylmethyl)cyclopropanamine ;
(1S,2R)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine ;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(piperidin-4-ylmethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;

(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(pyridin-3-yl)cyclopropanamine;
(1S,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1R,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(thiazol-5-yl)cyclopropanamine;
(1S,2R)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1R,2S)-N-(2-(piperidin-4-yl)ethyl)-2-(3'-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-(2-(piperidin-4-yl)ethyl)cyclopropanamine;
(1S,2R)-2-phenyl-N-((S)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1S,2R)-2-phenyl-N-((R)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-((S)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1R,2S)-2-phenyl-N-((R)-pyrrolidin-3-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-((2-fluorobenzyl)oxy)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(1S,2R)-N-(azetidin-3-ylmethyl)-2-phenylcyclopropanamine;
(1R,2S)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-cyclopropylphenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(1S,2R)-N-(piperidin-4-ylmethyl)-2-(4-(pyridin-3-yl)phenyl)cyclopropanamine;
(1R,2S)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(4-(1H-pyrazol-5-yl)phenyl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1S,2R)-2-(naphthalen-2-yl)-N-(piperidin-4-ylmethyl)cyclopropanamine;
(1R,2S)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine;
(1S,2R)-2-(4-(benzyloxy)phenyl)-N-((1-methylpiperidin-4-yl)methyl)cyclopropanamine;
or a salt or solvate thereof.


367. The compound of any of claims 198 to 201 and 284 to 366, wherein said compound is an optically active stereoisomer.

368. A process for the preparation of a compound of formula I according to claim 198, or a salt thereof, which comprises reacting a compound of formula II




II


wherein A, B, R^w , R^x , R^y , and R^z have the meaning defined in claim 198, with a compound of formula IIIa, IIIb or IIIc



IIIa



IIIb



IIIc

wherein D has the meaning defined in claim 198, and wherein any amino group that may be present in D is optionally protected with a protecting group,

in the presence of a reducing agent, followed by the removal of any protecting group that may be present.

369. The process according to claim 368, wherein said reducing agent is a borohydride.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/070898

A. CLASSIFICATION OF SUBJECT MATTER				
INV.	C07D205/04	C07D207/09	C07D207/14	C07D209/96
	C07D211/56	C07D211/58	C07D215/42	C07D221/20
	C07D309/04	C07D309/14	C07D401/04	C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2011/035941 A1 (ORYZON GENOMICS) 31 March 2011 (2011-03-31) claims; examples -----	1-369
A	US 2010/324147 A1 (MCCAFFERTY ET. AL.) 23 December 2010 (2010-12-23) page 11, paragraph 146 - page 12, paragraph 163; claims; examples -----	1-369
A	WO 2011/106105 A2 (ORYZON GENOMICS) 1 September 2011 (2011-09-01) page 1, line 14 - page 7, line 2; claims; examples ----- -/-	1-369

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
19 February 2013	14/03/2013
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3046	Authorized officer Helps, Ian

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/070898

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	S. MINASU ET. AL.: "Structurally Designed trans-2-Phenylcyclopropylamine Derivatives Potently Inhibit Histone Demethylase LSD1/KDM1.", BIOCHEMISTRY, vol. 49, 22 June 2010 (2010-06-22), pages 6494-6503, XP002692366, figures 1-5 ----- H. BENELKEBIR ET. AL.: "Enantioselective Synthesis of Tranylcypromine Analogues as Lysine Demethylase (LSD1) Inhibitors.", BIOORGANIC AND MEDICINAL CHEMISTRY, vol. 19, 13 February 2011 (2011-02-13), pages 3709-3716, XP002659903, figures 1-7 ----- D. M. GOODEN ET. AL.: "Facile Synthesis of Substituted trans-2-Arylcyclopropylamine Inhibitors of the Human Histone Demethylase LSD1 and Monoamine Oxidases A and B", BIOORGANIC AND MEDICINAL CHEMISTRY LETTERS, vol. 18, 8 January 2008 (2008-01-08), pages 3047-3051, XP002692367, tables 1-3 ----- WO 2011/131697 A1 (ORYZON GENOMICS) 27 October 2011 (2011-10-27) page 1, line 13 - page 5, line 24; claims; examples ----- WO 2012/013728 A1 (ORYZON GENOMICS) 2 February 2012 (2012-02-02) page 1, line 8 - page 6, line 13; claims; examples ----- WO 2011/131576 A1 (UNIVERSITA DEGLI STUDI DI ROMA) 27 October 2011 (2011-10-27) page 1, line 15 - page 3, line 18; claims; examples -----	1-369 1-369 1-369 1-369 1-369 1-369 1-369 1-369 1-369
A,P		
Y,P		
Y,P		
A,P		

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2012/070898

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
WO 2011035941	A1 31-03-2011	AU CN EP KR US WO	2010297557 A1 102639496 A 2480528 A1 20120104525 A 2012283266 A1 2011035941 A1		17-05-2012 15-08-2012 01-08-2012 21-09-2012 08-11-2012 31-03-2011
US 2010324147	A1 23-12-2010		NONE		
WO 2011106105	A2 01-09-2011	WO WO	2011106105 A2 2011106574 A2		01-09-2011 01-09-2011
WO 2011131697	A1 27-10-2011	AU CA EP WO	2011244325 A1 2796726 A1 2560947 A1 2011131697 A1		06-12-2012 27-10-2011 27-02-2013 27-10-2011
WO 2012013728	A1 02-02-2012		NONE		
WO 2011131576	A1 27-10-2011	AU CA EP US WO	2011244478 A1 2797011 A1 2560949 A1 2013035377 A1 2011131576 A1		01-11-2012 27-10-2011 27-02-2013 07-02-2013 27-10-2011