

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2018232886 B2**

(54) Title
Pharmaceutical combinations comprising an anti-LY75 antibody

(51) International Patent Classification(s)
C07K 16/28 (2006.01) **A61P 35/00** (2006.01)
A61K 31/505 (2006.01) **A61P 35/02** (2006.01)
A61K 39/395 (2006.01) **C07K 16/46** (2006.01)
A61K 47/68 (2017.01)

(21) Application No: **2018232886** (22) Date of Filing: **2018.03.09**

(87) WIPO No: **WO18/162727**

(30) Priority Data

(31) Number	(32) Date	(33) Country
1703876.1	2017.03.10	GB

(43) Publication Date: **2018.09.13**

(44) Accepted Journal Date: **2024.09.19**

(71) Applicant(s)
Berlin-Chemie AG

(72) Inventor(s)
Binaschi, Monica;Bigioni, Mario;Merlino, Giuseppe;Simonelli, Cecilia;Bertoni, Francesco;Pellacani, Andrea

(74) Agent / Attorney
Griffith Hack, Level 15 376-390 Collins St, MELBOURNE, VIC, 3000, AU

(56) Related Art
WO 2015/052537 A1



(51) International Patent Classification:

C07K 16/28 (2006.01) A61K 47/68 (2017.01)

A61P 35/00 (2006.01) A61K 31/505 (2006.01)

A61P 35/02 (2006.01) C07K 16/46 (2006.01)

A61K 39/395 (2006.01)

(21) International Application Number:

PCT/EP2018/055939

(22) International Filing Date:

09 March 2018 (09.03.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1703876.1 10 March 2017 (10.03.2017) GB

(71) Applicant: BERLIN-CHEMIE AG [DE/DE]; Glienicke Weg 125-127, 12489 Berlin (DE).

(72) Inventors: BINASCHI, Monica; Menarini Ricerche, Via Tito Speri 10 Pomezia, Rome (IT). BIGIONI, Mario; Menarini Ricerche, Via Tito Speri 10 Pomezia, Rome (IT). MERLINO, Giuseppe; Menarini Ricerche, Via Tito Speri 10 Pomezia, Rome (IT). SIMONELLI, Cecilia; Menarini Ricerche, Via Sette Santi 3, Firenze (IT). BERTONI, Francesco; Institute of Oncology Research Fondazione

per la Ricerca e la Cura sui Linfomi, Via Vela 6, 6500 Bellinzona (CH). PELLACANI, Andrea; Menarini Ricerche, Via Sette Santi 3, Firenze (IT).


(74) Agent: DEHNS; St Bride's House, 10 Salisbury Square, London, Greater London EC4Y 8JD (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: PHARMACEUTICAL COMBINATIONS COMPRISING AN ANTI-LY75 ANTIBODY

Figure 9



(57) Abstract: The present disclosure relates generally to the fields of immunology and molecular biology. More specifically, provided herein are pharmaceutical combinations comprising (A) antibodies, or antigen-binding portions thereof, directed against LY75, and (B) a second anti-cancer entity; nucleic acids encoding antibody combinations; methods for preparing antibody combinations; and methods for the treatment of diseases, such as cancers mediated by LY75 expression or activity.



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

PHARMACEUTICAL COMBINATIONS COMPRISING AN ANTI-LY75 ANTIBODY

INTRODUCTION

5 The present disclosure relates generally to the fields of immunology and molecular biology. More specifically, provided herein are pharmaceutical combinations comprising (A) antibodies, or antigen-binding portions thereof, directed against LY75, and (B) a second anti-cancer entity; nucleic acids encoding antibody combinations; methods for preparing antibody combinations; and methods for the treatment of diseases, such as cancers mediated by LY75 expression or
10 activity.

BACKGROUND

Leukemias and lymphomas belong to a broad group of tumours that affect the blood, bone marrow, and lymphoid system; these are known as tumors of the hematopoietic and
15 lymphoid tissues.

Lymphoma is a group of blood cell tumours that develop from lymphocytes. Signs and symptoms may include enlarged lymph nodes, fever, drenching sweats, unintended weight loss, itching and constantly feeling tired. There are a number of subtypes of lymphomas: the two main categories of lymphomas are Hodgkin's lymphomas (HL) and the non-Hodgkin lymphomas
20 (NHL). The World Health Organization (WHO) includes two other categories as types of lymphoma: multiple myeloma and immunoproliferative diseases. About 90% of lymphomas are non-Hodgkin lymphomas.

Leukemia is a group of cancers that usually begin in the bone marrow and result in high numbers of abnormal white blood cells. Symptoms may include bleeding and bruising problems,
25 feeling tired, fever and an increased risk of infections. These symptoms occur due to a lack of normal blood cells. Diagnosis is typically made by blood tests or bone marrow biopsy. There are four main types of leukemia: acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL) and chronic myeloid leukemia (CML), as well as a number of less common types.

30 Treatment of leukemias and lymphomas may involve one or more of chemotherapy, radiation therapy, targeted therapy and surgery (and bone marrow transplant in the case of leukemias). The success of leukemia treatment depends on the type of leukemia and the age of the person. The lymphoma treatment outcome depends on the subtype with some being curable and treatment prolonging survival in most.

35 A number of chemotherapeutic agents have previously been used for the treatment of leukemias including prednisone, vincristine, anthracyclines, L-asparaginase, cyclophosphamide, methotrexate, 6-mercaptopurine, fludarabine, pentostatin and cladribine. Chemotherapeutic

agents for the treatment of lymphomas include cyclophosphamide, hydroxydaunorubicin (also known as doxorubicin or adriamycin), oncovin (vincristine), prednisone, prednisolone, bleomycin, dacarbazine, etoposide and procarbazine.

Combination chemotherapy involves treating a patient with a two or more different drugs simultaneously. The drugs may differ in their mechanism and side-effects. The biggest advantage of this is minimising the chances of resistance developing to any one agent.

Furthermore, the drugs can often be used at lower doses, reducing toxicity. Combination therapies for the treatment of Hodgkin's Disease include MOPP (mustine, vincristine, procarbazine, prednisolone) and ABVD (Doxorubicin, bleomycin, vinblastine, dacarbazine).

Combination therapies for the treatment of Non-Hodgkin's lymphoma include CHOP (cyclophosphamide, doxorubicin, vincristine, prednisolone). Given the number of drugs that are known for the treatment of leukemias and lymphomas, the number of permutations and combinations of possible drug therapies is clearly large. Furthermore, the aforementioned combination therapies do not include antibodies.

There remains, however, a need for new treatments of leukemias and lymphomas, and particularly for efficacious combination therapies.

Lymphocyte antigen 75 acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment and is thought to cause a reduction in proliferation of B-lymphocytes. Expression of Lymphocyte antigen 75 has been observed in pancreatic, ovarian, breast, colorectal, esophageal, skin, thyroid and lung (non-small-cell) cancers as well as Multiple Myeloma and many different subtypes of lymphomas and leukaemias. WO2009/061996 discloses isolated monoclonal antibodies which bind to human DEC-205 (LY75) and related antibody based compositions and molecules. Also disclosed are pharmaceutical compositions comprising the antibodies, as well as therapeutic and diagnostic methods for using the antibodies. WO2008/104806 discloses affinity reagents capable of binding to LY75 for use in the treatment or prophylaxis of cancer. WO2015/052537 discloses specific isolated antibodies capable of binding to LY75 and their use in the treatment various cancers.

Rituximab is a monoclonal antibody against the protein CD20 which is widely expressed on B cells (Oncogene (2003 Oct 20), Smith MR, "Rituximab (monoclonal anti-CD20 antibody):

mechanisms of action and resistance", 22(47): 7359-68). Rituximab destroys both normal and malignant B cells that have CD20 on their surfaces and is therefore used to treat diseases which are characterized by having too many B cells, overactive B cells, or dysfunctional B cells.

Rituximab has previously been used to treat a number of autoimmune diseases and some types of cancer, including rheumatoid arthritis, idiopathic thrombocytopenic purpura, pemphigus vulgaris, multiple sclerosis, systemic lupus erythematosus, non-Hodgkin's lymphoma, chronic inflammatory demyelinating polyneuropathy, chronic lymphocytic leukemia and autoimmune anemias.

Ibrutinib (Imbruvica) is a small molecule drug that binds permanently to Bruton's tyrosine kinase (BTK), which is important in B cells. Ibrutinib has previously been used to treat B cell cancers like mantle cell lymphoma, chronic lymphocytic leukemia, and Waldenström's macroglobulinemia.

5 It has now been found that combinations of (i) certain anti-LY75 antibodies with Rituximab and (ii) certain anti-LY75 antibodies with Ibrutinib demonstrate synergistic results in the treatment of lymphomas.

BRIEF SUMMARY OF THE INVENTION

10 In one aspect, the invention provides a pharmaceutical combination comprising:

(A) an anti-LY75 antibody, or an antigen-binding portion thereof, which competes for binding to LY75 with an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2; or

15 an anti-LY75 antibody, or an antigen-binding portion thereof, said antibody comprising:

a) a heavy chain variable region comprising:

i) a first vhCDR comprising SEQ ID NO: 5;

ii) a second vhCDR comprising SEQ ID NO: 6; and

iii) a third vhCDR comprising SEQ ID NO: 7; and

20 b) a light chain variable region comprising:

i) a first vlCDR comprising SEQ ID NO: 8;

ii) a second vlCDR comprising SEQ ID NO: 9; and

iii) a third vlCDR comprising SEQ ID NO: 10;

optionally wherein any one or more of the above SEQ ID NOs independently comprise

25 one, two, three, four or five amino acid substitutions, additions or deletions;

and

(B) an anti-CD20 antibody, or an antigen-binding portion thereof, said antibody comprising:

a) a heavy chain variable region comprising:

i) a first vhCDR comprising SEQ ID NO: 40;

30 ii) a second vhCDR comprising SEQ ID NO: 41; and

iii) a third vhCDR comprising SEQ ID NO: 42; and

b) a light chain variable region comprising:

i) a first vlCDR comprising SEQ ID NO: 43;

ii) a second vlCDR comprising SEQ ID NO: 44; and

35 iii) a third vlCDR comprising SEQ ID NO: 45;

optionally wherein any one or more of the above SEQ ID NOs independently comprise one, two, three, four or five amino acid substitutions, additions or deletions;

wherein the pharmaceutical combination is in the form of a combined preparation for simultaneous, separate or sequential use.

In a second aspect, the invention provides a pharmaceutical combination comprising:

(A) an anti-LY75 antibody, or an antigen-binding portion thereof, which competes for binding to LY75 with an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2;

or

an anti-LY75 antibody, or an antigen-binding portion thereof, said antibody comprising:

a) a heavy chain variable region comprising:

- i) a first vhCDR comprising SEQ ID NO: 5;
- ii) a second vhCDR comprising SEQ ID NO: 6; and
- iii) a third vhCDR comprising SEQ ID NO: 7; and

b) a light chain variable region comprising:

- i) a first vlCDR comprising SEQ ID NO: 8;
- ii) a second vlCDR comprising SEQ ID NO: 9; and
- iii) a third vlCDR comprising SEQ ID NO: 10;

optionally wherein any one or more of the above SEQ ID NOs independently comprise one, two, three, four or five amino acid substitutions, additions or deletions;

and

(B) ibrutinib or a pharmaceutically-acceptable salt thereof,

wherein the pharmaceutical combination is in the form of a combined preparation for simultaneous, separate or sequential use.

In one embodiment, the anti-LY75 antibody or antigen-binding portion thereof comprises a heavy chain variable region comprising 1, 2 or 3 CDRs selected from the group consisting of CDRs comprising SEQ ID NOs: 5, 6, and 7, and/or a light chain variable region comprising 1, 2 or 3 CDRs selected from the group consisting of CDRs comprising SEQ ID NOs: 8, 9 and 10.

In some embodiments, the anti-LY75 antibodies bind to LY75 (SEQ ID NO: 15) and are capable of being internalized by a cell expressing LY75.

In another embodiment, the anti-LY75 antibody comprises the heavy and/or light chain complementarity determining regions (CDRs) or variable regions (VRs) of the particular antibody described herein (e.g., referred to herein as "LY75_A1"). Accordingly, in one embodiment, the anti-LY75 antibody comprises the CDR1, CDR2, and CDR3 domains of the heavy chain variable (VH) region of antibody LY75_A1 having the sequence shown in SEQ ID NO:1, and/or the

CDR1, CDR2 and CDR3 domains of the light chain variable (VL) region of LY75_A1 having the sequence shown in SEQ ID NO:2.

In another embodiment, the anti-LY75 antibodies bind to human LY75 and include a heavy chain variable region comprising SEQ ID NO: 1, and/or conservative sequence modifications thereof. The antibody may further include a light chain variable region comprising SEQ ID NO:2, and/or conservative sequence modifications thereof.

In a further embodiment, the anti-LY75 antibodies bind to human LY75 and include a heavy chain variable region and a light chain variable region including the amino acid sequences set forth in SEQ ID NOs: 1 and/or 2, respectively, and conservative sequence modifications thereof.

Antibodies which include heavy and light chain variable regions having at least 80%, or at least 85%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98%, or at least 99%, or more sequence identity to any of the above sequences are also included in the present invention. Ranges intermediate to the above-recited values, e.g., heavy and light chain variable regions having at least 80-85%, 85-90%, 90-95% or 95-100% sequence identity to any of the above sequences are also intended to be encompassed by the present invention.

In one embodiment, the anti-LY75 antibody comprises a heavy chain variable region comprising SEQ ID NO:1 or a sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to SEQ ID NO: 1. In another embodiment, the anti-LY75 antibody comprises a light chain variable region comprising SEQ ID NO:2 or a sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to SEQ ID NO: 2. In another embodiment, the anti-LY75 antibody comprises a heavy chain framework region comprising an amino acid sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to the framework of the heavy chain variable region of SEQ ID NO: 1 as shown in SEQ ID NOS: 16, 17, 18 and 19. In another embodiment, the anti-LY75 antibody comprises a light chain framework region comprising an amino acid sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to the framework of the light chain variable region of SEQ ID NO:2 as shown in SEQ ID NOS: 20, 21, 22 and 23.

In one embodiment, the anti-LY75 antibody competes for binding to LY75 with an antibody comprising heavy and/or light chain variable regions comprising the amino acid sequences set forth in SEQ ID NOs:1 and 2, respectively, or amino acid sequences at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%,

at least 97%, at least 98%, at least 99% identical thereto. In another embodiment, the anti-LY75 antibody competes for binding to LY75 with an antibody comprising heavy and/or light chain variable regions comprising the amino acid sequences set forth in SEQ ID NOs:1 and 2 (LY75_A1).

5 Other antibodies of the invention bind to the same epitope or an epitope on LY75 recognized by the antibodies described herein. In another particular embodiment, the antibody binds to an epitope on LY75 recognized by an antibody comprising heavy and/or light chain variable regions comprising the amino acid sequences set forth in SEQ ID NOs:1 and 2, respectively, or amino acid sequences at least 80% identical thereto. In another embodiment, 10 the antibody binds to an epitope on LY75 recognized by an antibody comprising heavy and/or light chain variable regions comprising the amino acid sequences set forth in SEQ ID NOs:1 and 2 (LY75_A1).

In a further embodiment, the anti-LY75 antibodies bind specifically to one or more, for example, 2, 3, 4, 5, 6, 7, 8, 9 or 10, peptide(s) selected from the group comprising SEQ ID NOs: 15 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36 or 37 or fragments thereof, wherein said fragments comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 contiguous amino acids. In a further embodiment, the epitope recognized by the anti-LY75 antibodies comprises one or more peptides, two or more or three or more peptides selected from the group consisting of SEQ ID NOs: 27, 29, 30, 34, 35, 36 or 37 or 20 fragments thereof wherein said fragments comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 contiguous amino acids. In a further embodiment, the epitope recognized by the anti-LY75 antibodies comprises one or more peptides, for example, two or three peptides selected from the group consisting of SEQ ID NOs: 30, 36 and 37 or fragments thereof wherein said fragments comprises at least 2, at least 3, at 25 least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 contiguous amino acids.

In a further embodiment, the anti-LY75 antibodies comprise variable CDRs as compared to the parent antibodies described herein. Thus, the variant antibodies comprising variant variable regions of a parent antibody, wherein the parent antibody comprises a first vhCDR comprising 30 SEQ ID NO:5, a second vhCDR comprising SEQ ID NO: 6, a third vhCDR comprising SEQ ID NO:7, a first vlCDR comprising SEQ ID NO:8, a second vlCDR comprising SEQ ID NO:9 and a third vlCDR comprising a SEQ ID NO:10, and wherein the variant antibody has 1, 2, 3, 4, 5 or 6 amino acid substitutions collectively in the set of the first vhCDR, the second vhCDR, the third vhCDR, the first vlCDR, the second vlCDR and the third vlCDR, with from 1 to 4, 1 to 3 or 1 to 2 35 substitutions of particular use, and wherein the antibody retains specific binding to LY75.

All of the antibodies disclosed herein can be full-length, for example, any of the following isotypes: IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD, and IgE. Alternatively, the

antibodies can be fragments such as an antigen-binding portion or a single chain antibody (e.g., a Fab, F(ab')₂, Fv, a single chain Fv fragment, an isolated complementarity determining region (CDR) or a combination of two or more isolated CDRs). The antibodies can be any kind of antibody, including, but not limited to, human, humanized, and chimeric antibodies.

5 In other embodiments, the anti-LY75 antibodies are in the form of an immunoconjugate (i.e., further include a covalently-attached moiety). In a particular embodiment, the moiety is a drug, such as a maytansinoid, a dolastatin, an auristatin, a trichothecene, a calicheamicin, CC1065 or derivatives thereof. In a preferred embodiment, the drug moiety is DM1 or DM4.

10 In one embodiment, the anti-LY75 antibody comprises a heavy chain variable region and a light chain variable region encoded by nucleic acid sequences comprising SEQ ID NOs: 3 and 4, respectively, or nucleic acid sequences having at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to the aforementioned nucleic acid sequences or sequences which differ from SEQ ID NOs: 3 and 4 due to degeneracy of the genetic code.

15 In one embodiment, anti-CD20 antibody is a mouse/human chimeric antibody, a humanised antibody or a human antibody. Preferably the anti-CD20 antibody is Rituximab.

In another aspect of the present invention there are provided kits expression vectors comprising nucleic acids encoding heavy and light chain variable regions of the antibodies described herein operably linked to one or more regulatory elements.

20 In a preferred embodiment the host cell comprises expression vectors comprising nucleic acids encoding:

- (i) the heavy chain of the anti-LY75 antibody or antigen-binding portion thereof;
- (ii) the light chain of the anti-LY75 antibody or antigen-binding portion thereof;
- (iii) the heavy chain of the anti-CD20 antibody or antigen-binding portion thereof; and
- 25 (iv) the light chain of the anti-CD20 antibody or antigen-binding portion thereof.

In a further aspect there is provided a method of treating cancer in a patient comprising simultaneously, sequentially or separately administering to a patient in need thereof therapeutically-effective amounts of components (A) and (B) of a pharmaceutical combination of the invention.

30 In a further aspect of the present invention, there is provided a pharmaceutical combination of the invention for use in the treatment of cancer.

Also provided is the use of components (A) and (B) as defined herein in the manufacture of a pharmaceutical combination for simultaneous, separate or sequential use for the treatment of cancer. In one embodiment the cancer is preferably leukemia or lymphoma.

35 In some embodiments, the cancer is selected from the group consisting of non-Hodgkin's lymphoma, diffuse large B-cell lymphoma (DLBCL), B-Cell Lymphoma, Follicular Lymphoma, Mantle Cell Lymphoma, Lymphoma of Mucosa-Associated Lymphoid Tissue (MALT), T-

Cell/Histiocyte-Rich B-Cell Lymphoma, Burkitt's Lymphoma, Lymphoplasmacytic Lymphoma, Small Lymphocytic Lymphoma, Marginal Zone Lymphoma, T Cell Lymphoma, Peripheral T-Cell Lymphoma, Anaplastic Large Cell Lymphoma and AngioImmunoblastic T-Cell Lymphoma, acute myeloid leukaemia, and chronic lymphocytic leukaemia. More preferably, the cancer is DLBCL or non-Hodgkin's lymphoma.

Also within the scope of the invention are kits comprising a pharmaceutical combination of the invention and, optionally, instructions for use. The kit can further contain a least one additional reagent or one or more additional antibodies.

Other features and advantages of the instant invention will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the alignment of LY75_A1 heavy chain (SEQ ID NO:1), the human VH 3-15 Germline (SEQ ID NO:11) and the human JH4 Germline (SEQ ID NO:12). The CDR regions of LY75_A1 heavy chain are underlined.

Figure 2 depicts the alignment of LY75_A1 light chain (SEQ ID NO:2), the human VK O12 Germline (SEQ ID NO:13) and the human JK4 Germline (SEQ ID NO:14). The CDR regions of LY75_A1 light chain are underlined.

Figure 3a depicts cytotoxic activity of anti-LY75 monoclonal antibodies conjugated with DM1 in HT-29 and shows while most antibodies bind to LY75 only a few display efficacy.

Figure 3b depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in HT-29.

Figure 3c depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in RAJI cells.

Figure 3d depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in Namalwa cells.

Figure 3e depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in Karpas 299 cells.

Figure 3f depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in BxPC3 cells.

Figure 3g depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in HupT4 cells.

Figure 3h depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in HPAFFII cells.

Figure 3i depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in EHEB cells.

Figure 3j depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in Mec-1 cells.

Figure 3k depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in AML-193 cells.

5 Figure 3l depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in HCC 70 cells.

Figure 3m depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in HCC 1806 cells.

10 Figure 3n depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in MDA-MB-468 cells.

Figure 3o depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in RT4 cells.

Figure 3p depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in 5637 cells.

15 Figure 3q depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in SW780 cells.

Figure 3r depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in SCC-9 cells.

20 Figure 3s depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in OE 19 cells.

Figure 3t depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in OVCAR-3 cells.

Figure 3u depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in SK-OV-3 cells.

25 Figure 3v depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in MOLP-8 cells.

Figure 3w depicts cytotoxic activity of anti-LY75 antibodies conjugated to either DM1 or DM4 in RPMI8226 cells.

30 Figure 4a depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in Raji Burkitt's lymphoma SCID mouse xenograft model.

Figure 4b depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in Namalwa Burkitt's lymphoma SCID mouse xenograft model.

Figure 4c depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in HPAFII pancreatic adenocarcinoma athymic nude mouse xenograft model.

35 Figure 4d depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in SW780 human bladder carcinoma SCID mouse xenograft model.

Figure 4e depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in MDA-MB-468 athymic nude mouse xenograft model.

Figure 4f depicts the efficacy of anti-LY75 antibodies conjugated to either DM1 or DM4 in COLO205 colorectal adenocarcinoma athymic nude mouse xenograft model.

Figure 5a shows competitive binding of anti-LY75-mAb and an anti-LY75-mAb conjugated to MCC-DM1.

Figure 5b shows non-competitive binding of LY75_A1 and an anti-LY75-mAb conjugated to MCC-DM1.

Figure 6a-6j show graphical representations of the binding of antibody LY75_A1 to LY75 peptides on a peptide microarray.

Figure 7 shows an amino acid alignment of peptides bound by antibody LY75_A1 in both the peptide microarray assay and the peptide pull down assay. Peptides highlighted are those likely to form the epitope recognized by antibody LY75_A1.

Figures 8A and 8B show the anti-proliferative effect of different (nM) doses of LY75_DM4 as a single treatment or in combination with Rituximab on two different ABC-DLBCL cell lines (TMD8 and HBL1). (CI = Chou-Talalay Combination Index). Figure 8A: median CI = 0.27. Figure 8B: median CI = 0.57.

Figure 9 shows the anti-proliferative effect of different (nM) doses of LY75_DM4 as a single treatment or in combination with Ibrutinib on HBL-1 cell lines (ABC-DLBCL). (CI = Chou-Talalay Combination Index). Median CI = 0.24.

DETAILED DESCRIPTION OF THE INVENTION

The present disclosure relates to pharmaceutical combinations comprising components (A) and (B) as defined herein, wherein the pharmaceutical combination is in the form of a combined preparation for simultaneous, separate or sequential use. Component (A) relates to an anti-LY75 antibody as defined herein. Component (B) relates either to (i) an anti-CD20 antibody as defined herein or (ii) to ibrutinib or a pharmaceutically-acceptable salt thereof.

One example of the LY75 protein is given in SEQ ID NO: 15 herein. The terms "anti-LY75 antibodies" and "LY75 antibodies" are used interchangeably herein.

The LY75 antibodies disclosed herein may be internalized when contacted with cells expressing the LY75 receptor. As discussed herein, the LY75 receptor is overexpressed and/or differentially expressed on certain cancer cells, including but not limited to, leukaemia preferably acute myeloid leukaemia or chronic lymphocytic leukaemia, lymphoma, preferably DLBCL B-Cell Lymphoma, Follicular Lymphoma, Mantle Cell Lymphoma, Lymphoma of Mucosa-Associated Lymphoid Tissue (MALT), T-Cell/Histiocyte-Rich B-Cell Lymphoma, Burkitt's Lymphoma, Lymphoplasmacytic Lymphoma, Small Lymphocytic Lymphoma, Marginal Zone Lymphoma, T

Cell Lymphoma, Peripheral T-Cell Lymphoma, Anaplastic Large Cell Lymphoma and Angioimmunoblastic T-Cell Lymphoma.

As such, when the LY75 antibodies disclosed herein are conjugated to drugs (sometimes referred to herein as “antibody-drug conjugates” or “ADCs”), the internalization of these ADC molecules into cancer cells results in cell death and thus tumor treatment.

The anti-LY75 antibodies possess particular structural features such as CDR regions with particular amino acid sequences. Described herein are a set of CDRs which can form an affinity reagent, e.g. an antibody, which exhibits binding to LY75.

Thus, the disclosure provides antibodies, preferably isolated antibodies (which, as outlined below, includes a wide variety of well-known antibody structures, derivatives, mimetics and conjugates), nucleic acids encoding antibody combinations, host cells used to make the antibody combinations, methods of making the antibody combinations, and pharmaceutical combinations comprising the antibodies and optionally a pharmaceutical carrier, methods of treatment comprising the use of the pharmaceutical combinations and the use of the pharmaceutical combinations for the treatment of cancers.

Lymphocyte antigen 75 acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment and is thought to cause a reduction in proliferation of B-lymphocytes.

According to SWISS-PROT, Lymphocyte antigen 75 is expressed in spleen, thymus, colon and peripheral blood lymphocytes. It has been detected in myeloid and B lymphoid cell lines. Isoforms designated herein OGTA076b and OGTA076c are expressed in malignant Hodgkin's lymphoma cells called Hodgkin's and Reed-Sternberg (HRS) cells. LY75 acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment. It causes reduced proliferation of B-lymphocytes.

Expression of LY75 has been observed in pancreatic, bladder, ovarian, breast (including triple negative), colorectal, esophageal, skin, thyroid and lung (non-small-cell) cancers as well as Multiple Myeloma and many different subtypes of lymphomas (including DLBCL) and leukaemias.

The anti-LY75 antibody may, in certain cases, cross-react with the LY75 from species other than human. For example, to facilitate clinical testing, the anti-LY75 antibodies may cross react with murine or primate LY75 molecules. Alternatively, in certain embodiments, the antibodies may be completely specific for human LY75 and may not exhibit species or other types of non-human cross-reactivity.

The present invention relates to anti-LY75 antibodies and, in some embodiments anti-CD20 antibodies, generally therapeutic antibodies as described herein. Antibodies that find use in the present invention can take on a number of formats as described herein, including traditional antibodies as well as antibody derivatives, fragments and mimetics, described below.

In one embodiment, the invention provides antibody structures that contain a set of 6 CDRs as defined herein (including small numbers of amino acid changes as described below).

"Antibody" as used herein includes a wide variety of structures, as will be appreciated by those in the art, that in some embodiments contain at a minimum a set of 6 CDRs as defined
5 herein; including, but not limited to traditional antibodies (including both monoclonal and polyclonal antibodies), humanized and/or chimeric antibodies, antibody fragments, engineered antibodies (e.g. with amino acid modifications as outlined below), multispecific antibodies (including bispecific antibodies), and other analogs known in the art.

Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically
10 composed of two identical pairs of polypeptide chains, each pair having one "light" (typically having a molecular weight of about 25 kDa) and one "heavy" chain (typically having a molecular weight of about 50-70 kDa). Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but
15 not limited to IgG1, IgG2, IgG3, and IgG4. IgM has subclasses, including, but not limited to, IgM1 and IgM2. Thus, "isotype" as used herein is meant any of the subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. The known human immunoglobulin isotypes are IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgM1, IgM2, IgD, and IgE. It should be understood that therapeutic antibodies can also comprise hybrids of any
20 combination of isotypes and/or subclasses.

In many embodiments, IgG isotypes are used in the present invention, with IgG1 finding particular use in a number of applications.

The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. In the variable region, three loops
25 are gathered for each of the V domains of the heavy chain and light chain to form an antigen-binding site. Each of the loops is referred to as a complementarity-determining region (hereinafter referred to as a "CDR"), in which the variation in the amino acid sequence is most significant. "Variable" refers to the fact that certain segments of the variable region differ extensively in sequence among antibodies. Variability within the variable region is not evenly
30 distributed. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called "hypervariable regions" that are each 9-15 amino acids long or longer.

Each VH and VL is composed of three hypervariable regions ("complementary determining regions," "CDRs") and four FRs, arranged from amino-terminus to carboxy-terminus in the
35 following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4.

The hypervariable region generally encompasses amino acid residues from about amino acid residues 24-34 (LCDR1; "L" denotes light chain), 50-56 (LCDR2) and 89-97 (LCDR3) in the

light chain variable region and around about 31-35B (HCDR1; "H" denotes heavy chain), 50-65 (HCDR2), and 95-102 (HCDR3) in the heavy chain variable region; Kabat et al., SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991) and/or those residues forming a hypervariable loop (e.g. residues 26-32 (LCDR1), 50-52 (LCDR2) and 91-96 (LCDR3) in the light chain variable region and 26-32 (HCDR1), 53-55 (HCDR2) and 96-101 (HCDR3) in the heavy chain variable region; Chothia and Lesk (1987) J. Mol. Biol. 196:901-917. Specific CDRs of the invention are described below.

Throughout the present specification, the Kabat numbering system is generally used when referring to a residue in the variable domain (approximately, residues 1-107 of the light chain variable region and residues 1-113 of the heavy chain variable region) (e.g, Kabat et al., supra (1991)).

The CDRs contribute to the formation of the antigen-binding, or more specifically, epitope binding site of antibodies. The term "epitope" or "antigenic determinant" refers to a site on an antigen to which an immunoglobulin or antibody specifically binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids in a unique spatial conformation. As described herein, methods for determining what epitopes are bound by a given antibody (*i.e.*, epitope mapping) are well known in the art and include, for example, immunoblotting and immunoprecipitation assays, wherein overlapping or contiguous peptides from LY75 are tested for reactivity with the given anti-LY75 antibody. Methods of determining spatial conformation of epitopes include techniques in the art and those described herein, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance (see, *e.g.*, *Epitope Mapping Protocols in Methods in Molecular Biology*, Vol. 66, G. E. Morris, Ed. (1996)). The term "epitope mapping" refers to the process of identification of the molecular determinants for antibody-antigen recognition.

The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Kabat *et al.* collected numerous primary sequences of the variable regions of heavy chains and light chains. Based on the degree of conservation of the sequences, they classified individual primary sequences into the CDR and the framework and made a list thereof (see SEQUENCES OF IMMUNOLOGICAL INTEREST, 5th edition, NIH publication, No. 91-3242, E.A. Kabat *et al.*).

In the IgG subclass of immunoglobulins, there are several immunoglobulin domains in the heavy chain. By "immunoglobulin (Ig) domain" herein is meant a region of an immunoglobulin having a distinct tertiary structure. Of interest in the present invention are the heavy chain

domains, including, the constant heavy (CH) domains and the hinge domains. In the context of IgG antibodies, the IgG isotypes each have three CH regions. Accordingly, "CH" domains in the context of IgG are as follows: "CH1" refers to positions 118-220 according to the EU index as in Kabat. "CH2" refers to positions 237-340 according to the EU index as in Kabat, and "CH3" refers to positions 341-447 according to the EU index as in Kabat.

Another type of Ig domain of the heavy chain is the hinge region. By "hinge" or "hinge region" or "antibody hinge region" or "immunoglobulin hinge region" herein is meant the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 236 (G236 in IgG1), wherein the numbering is according to the EU index as in Kabat. In some embodiments, for example in the context of an Fc region, the lower hinge is included, with the "lower hinge" generally referring to positions 226 or 230.

Of particular interest in the present invention are the Fc regions. By "Fc" or "Fc region" or "Fc domain" as used herein is meant the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, the Fc domain comprises immunoglobulin domains Cy2 and Cy3 (Cy2 and Cy3) and the lower hinge region between Cy1 (Cy1) and Cy2 (Cy2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. In some embodiments, as is more fully described below, amino acid modifications are made to the Fc region, for example to alter binding to one or more FcγR receptors or to the FcRn receptor.

In some embodiments, the antibodies are full length. By "full length antibody" herein is meant the structure that constitutes the natural biological form of an antibody, including variable and constant regions, including one or more modifications as outlined herein.

Alternatively, the antibodies can be a variety of structures, including, but not limited to, antibody fragments, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as "antibody mimetics"), chimeric antibodies, humanized antibodies, antibody fusions (sometimes referred to as "antibody conjugates"), and fragments of each, respectively. Structures that rely on the use of a set of CDRs are included within the definition of "antibody".

In one embodiment, the antibody is an antibody fragment. Specific antibody fragments include, but are not limited to, (i) the Fab fragment consisting of VL, VH, CL and CH1 domains, (ii) the Fd fragment consisting of the VH and CH1 domains, (iii) the Fv fragment consisting of the

VL and VH domains of a single antibody; (iv) the dAb fragment (Ward *et al.*, 1989, Nature 341:544-546, entirely incorporated by reference) which consists of a single variable region, (v) isolated CDR regions, (vi) F(ab')₂ fragments, a bivalent fragment comprising two linked Fab fragments (vii) single chain Fv molecules (scFv), wherein a VH domain and a VL domain are
5 linked by a peptide linker which allows the two domains to associate to form an antigen binding site (Bird *et al.*, 1988, Science 242:423-426, Huston *et al.*, 1988, Proc. Natl. Acad. Sci. U.S.A. 85:5879-5883, entirely incorporated by reference), (viii) bispecific single chain Fv (WO 03/11161, hereby incorporated by reference) and (ix) "diabodies" or "triabodies", multivalent or multispecific fragments constructed by gene fusion (Tomlinson *et al.*, 2000, Methods Enzymol. 326:461-479;
10 WO94/13804; Holliger *et al.*, 1993, Proc. Natl. Acad. Sci. U.S.A. 90:6444-6448, all entirely incorporated by reference).

In some embodiments, the antibody can be a mixture from different species, e.g. a chimeric antibody and/or a humanized antibody. That is, in the present invention, the CDR sets can be used with framework and constant regions other than those specifically described by
15 sequence herein.

In general, both "chimeric antibodies" and "humanized antibodies" refer to antibodies that combine regions from more than one species. For example, "chimeric antibodies" traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human. "Humanized antibodies" generally refer to non-human antibodies that have had the
20 variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an
25 antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018, Jones, 1986, Nature 321:522-525, Verhoeyen *et al.*, 1988, Science 239:1534-1536, all entirely incorporated by reference. "Backmutation" of selected acceptor framework residues to the corresponding donor residues is often required to regain affinity that is lost in the initial grafted construct (US 5530101; US 5585089; US 5693761;
30 US 5693762; US 6180370; US 5859205; US 5821337; US 6054297; US 6407213, all entirely incorporated by reference). The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin, and thus will typically comprise a human Fc region. Humanized antibodies can also be generated using mice with a genetically engineered immune system. Roque *et al.*, 2004, Biotechnol. Prog.
35 20:639-654, entirely incorporated by reference. A variety of techniques and methods for humanizing and reshaping non-human antibodies are well known in the art (See Tsurushita & Vasquez, 2004, Humanization of Monoclonal Antibodies, Molecular Biology of B Cells, 533-545,

Elsevier Science (USA), and references cited therein, all entirely incorporated by reference).

Humanization methods include but are not limited to methods described in Jones et al., 1986,

Nature 321:522-525; Riechmann et al., 1988; Nature 332:323-329; Verhoeyen et al., 1988,

Science, 239:1534-1536; Queen et al., 1989, Proc Natl Acad Sci, USA 86:10029-33; He et al.,

5 1998, J. Immunol. 160: 1029-1035; Carter et al., 1992, Proc Natl Acad Sci USA 89:4285-9,

Presta et al., 1997, Cancer Res. 57(20):4593-9; Gorman et al., 1991, Proc. Natl. Acad. Sci. USA

88:4181-4185; O'Connor et al., 1998, Protein Eng 11:321-8, all entirely incorporated by

reference. Humanization or other methods of reducing the immunogenicity of nonhuman

antibody variable regions may include resurfacing methods, as described for example in

10 Roguska et al., 1994, Proc. Natl. Acad. Sci. USA 91:969-973, entirely incorporated by reference.

In one embodiment, the parent antibody has been affinity matured, as is known in the art.

Structure-based methods may be employed for humanization and affinity maturation, for example

as described in USSN 11/004,590. Selection based methods may be employed to humanize

and/or affinity mature antibody variable regions, including but not limited to methods described in

15 Wu et al., 1999, J. Mol. Biol. 294:151-162; Baca et al., 1997, J. Biol. Chem. 272(16):10678-

10684; Rosok et al., 1996, J. Biol. Chem. 271(37): 22611-22618; Rader et al., 1998, Proc. Natl.

Acad. Sci. USA 95: 8910-8915; Krauss et al., 2003, Protein Engineering 16(10):753-759, all

entirely incorporated by reference. Other humanization methods may involve the grafting of only

parts of the CDRs, including but not limited to methods described in USSN 09/810,510; Tan et

20 al., 2002, J. Immunol. 169:1119-1125; De Pascalis et al., 2002, J. Immunol. 169:3076-3084, all

entirely incorporated by reference.

The antibodies disclosed herein may be isolated or recombinant. "Isolated," when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a cell or cell culture from which it was expressed. Thus an

25 isolated antibody is intended to refer to an antibody that is substantially free of other antibodies

having different antigenic specificities (e.g. an isolated antibody that specifically binds to the

LY75 is substantially free of antibodies that specifically bind antigens other than the LY75). Thus,

an "isolated" antibody is one found in a form not normally found in nature (e.g. non-naturally

occurring). An isolated antibody as defined herein may, in one embodiment, include at least one

30 amino acid which does not occur in the "naturally" occurring antibody. This amino acid may be

introduced by way of an addition or a substitution. It will be understood that the introduced amino

acid may be a naturally occurring or non-naturally occurring amino acid. In some embodiments,

the antibodies of the invention are recombinant proteins, isolated proteins or substantially pure

proteins. An "isolated" protein is unaccompanied by at least some of the material with which it is

35 normally associated in its natural state, for example constituting at least about 5%, or at least

about 50% by weight of the total protein in a given sample. It is understood that the isolated

protein may constitute from 5 to 99.9% by weight of the total protein content depending on the

circumstances. For example, the protein may be made at a significantly higher concentration through the use of an inducible promoter or high expression promoter, such that the protein is made at increased concentration levels. In the case of recombinant proteins, the definition includes the production of an antibody in a wide variety of organisms and/or host cells that are known in the art in which it is not naturally produced. Ordinarily, an isolated polypeptide will be prepared by at least one purification step. An "isolated antibody," refers to an antibody which is substantially free of other antibodies having different antigenic specificities. For instance, an isolated antibody that specifically binds to LY75 is substantially free of antibodies that specifically bind antigens other than LY75, with the exception of antibodies which bind to CD20.

Isolated monoclonal antibodies, having different specificities, can be combined in a well-defined composition. Thus for example, the antibody of the invention can optionally and individually be included or excluded in a formulation, as is further discussed below.

The anti-LY75 antibodies of the present invention specifically bind LY75 (e.g. SEQ ID NO: 15). The anti-CD20 antibodies of the present invention specifically bind CD20. "Specific binding" or "specifically binds to" or is "specific for" a particular antigen or an epitope means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target.

Specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a K_D for an antigen or epitope of at least about 10^{-4} M, at least about 10^{-5} M, at least about 10^{-6} M, at least about 10^{-7} M, at least about 10^{-8} M, at least about 10^{-9} M, alternatively at least about 10^{-10} M, at least about 10^{-11} M, at least about 10^{-12} M, or greater, where K_D refers to a dissociation rate of a particular antibody-antigen interaction. Typically, an antibody that specifically binds an antigen will have a K_D that is 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for a control molecule relative to the antigen or epitope. However, in the present invention, when administering ADCs of the LY75 antibodies of the invention, what is important is that the K_D is sufficient to allow internalization and thus cell death without significant side effects.

Also, specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a K_A or K_a for an antigen or epitope of at least 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for the epitope relative to a control, where K_A or K_a refers to an association rate of a particular antibody-antigen interaction.

Standard assays to evaluate the binding ability of the antibodies toward LY75 or CD20 can be done on the protein or cellular level and are known in the art, including for example, ELISAs, Western blots, RIAs, BIAcore® assays and flow cytometry analysis. Suitable assays are

described in detail in the Examples. The binding kinetics (e.g. binding affinity) of the antibodies also can be assessed by standard assays known in the art, such as by Biacore® system analysis. To assess binding to Raji or Daudi B cell tumor cells, Raji (ATCC Deposit No. CCL-86) or Daudi (ATCC Deposit No. CCL-213) cells can be obtained from publicly available sources, such as the American Type Culture Collection, and used in standard assays, such as flow cytometric analysis.

The LY75 antibodies that bind to LY75 (SEQ ID NO: 15) maybe internalized when contacted with cells expressing LY75 on the cell surface. These antibodies are referred to herein either as “anti-LY75” antibodies or, for ease of description, “LY75 antibodies”. Both terms are used interchangeably herein.

The LY75 antibodies are internalized upon contact with cells, particularly tumor cells, which express LY75 on the surface. That is, LY75 antibodies as defined herein that also comprise drug conjugates are internalized by tumor cells, resulting in the release of the drug and subsequent cell death, allowing for treatment of cancers that exhibit LY75 expression.

Internalization in this context can be measured in several ways. In one embodiment, the LY75 antibodies are contacted with cells, such as a cell line as outlined herein, using standard assays such as MAbZap. It would be clear to the skilled person that the MabZap assay is representative of the effect that would be expected to be seen with an antibody-drug conjugate (ADC). In the latter case, the ADC would be internalized, thus taking the drug into the cell. A toxic drug would have the capacity to kill the cell, i.e. to kill the targeted cancer cell. Data from MabZap assays are readily accepted by persons of skill in the art to be representative of ADC assays (Kohls, M and Lappi, D., [2000] Biotechniques, vol. 28, no. 1, 162-165).

In these *in vitro* assay embodiments, the LY75 antibodies are added, along with an anti-LY75 antibody comprising a toxin; for example, the LY75 antibody may be murine or humanized and the anti-LY75 antibody can be anti-murine or anti-humanized and contain a toxin such as saporin. Upon formation of the [LY75 antibody]-[anti-LY75 antibody-drug conjugate] complex, the complex is internalized and the drug (e.g. saporin) is released, resulting in cell death. Only upon internalization does the drug get released, and thus cells remain viable in the absence of internalization. As outlined below, without being bound by theory, in therapeutic applications, the anti-LY75 antibody contains the toxin, and upon internalization, the bond between the antibody and the toxin is cleaved, releasing the toxin and killing the cell.

In one embodiment, the anti-LY75 antibody comprises the heavy and light chain complementarity determining regions (CDRs) or variable regions (VRs) of the particular antibody described herein (e.g., referred to herein as “LY75_A1”). Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the heavy chain variable (VH) region of antibody LY75_A1 having the sequence shown in SEQ ID NO:1, and the CDR1, CDR2

and CDR3 domains of the light chain variable (VL) region of antibody LY75_A1 having the sequence shown in SEQ ID NO:2.

In another embodiment, the anti-LY75 antibody comprises a heavy chain variable region comprising a first vhCDR comprising SEQ ID NO: 5; a second vhCDR comprising SEQ ID NO: 6; and a third vhCDR comprising SEQ ID NO:7; and a light chain variable region comprising a first vlCDR comprising SEQ ID NO:8; a second vlCDR comprising SEQ ID NO: 9; and a third vlCDR comprising SEQ ID NO:10.

In another embodiment, the anti-LY75 antibodies bind to human LY75 and include a heavy chain variable region comprising an amino acid sequence comprising SEQ ID NO:1, and conservative sequence modifications thereof. The antibody may further include a light chain variable region comprising an amino acid sequence comprising SEQ ID NO:2, and conservative sequence modifications thereof.

In a further embodiment, the anti-LY75 antibodies bind to human LY75 and include a heavy chain variable region and a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs:1 and/or 2, respectively, and conservative sequence modifications thereof. As used herein, the term conservative sequence modification refers to, for example, the substitution of an amino acid with an amino acid having similar characteristics. It is common general knowledge for one skilled in the art what such substitutions may be considered conservative. Other modifications which can be considered to be conservative sequence modifications include, for example, glycosylation.

Optionally, one or more of SEQ ID NOs: 5-10 independently comprise one, two, three, four or five conservative amino acid substitutions; optionally, one or more SEQ ID NOs: 5-10 independently comprise one or two conservative amino acid substitutions.

Preferably, the term "conservative sequence modifications" is intended to include amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody containing the amino acid sequence. Such conservative modifications include amino acid substitutions, additions and deletions. Modifications can be introduced into an antibody of the invention by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues within the CDR regions of an antibody of the

invention can be replaced with other amino acid residues from the same side chain family and the altered antibody can be tested for retained function using the functional assays described herein.

Isolated antibodies which include heavy and light chain variable regions having at least 80%, or at least 85%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98%, or at least 99%, or more sequence identity to any of the above sequences are also included in the present invention. Ranges intermediate to the above-recited values, e.g., heavy and light chain variable regions having at least 80-85%, 85-90%, 90-95% or 95-100% sequence identity to any of the above sequences are also intended to be encompassed by the present invention. In one embodiment, the anti-LY75 antibody comprises a heavy chain variable region comprising SEQ ID NO:1 or a sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to SEQ ID NO: 1. In another embodiment, the anti-LY75 antibody comprises a light chain variable region comprising SEQ ID NO:2 or a sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to SEQ ID NO: 2. In another embodiment, the anti-LY75 antibody comprises a heavy chain framework region comprising an amino acid sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to the framework of the heavy chain variable region of SEQ ID NO: 1 comprising SEQ ID NOs: 16, 17 and 18. In another embodiment, the anti-LY75 antibody comprises a light chain framework region comprising an amino acid sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to the framework of the light chain variable region of SEQ ID NO:2 comprising SEQ ID NOs:19, 20 and 21.

In one embodiment, the anti-LY75 antibody is referred to herein as "LY75_A1 antibody" comprising the following CDRs, as well as variants containing a limited number of amino acid variants:

A1	SEQ ID NOs
variable heavy CDR1	5
variable heavy CDR2	6
variable heavy CDR3	7
variable light CDR1	8
variable light CDR2	9
variable light CDR3	10

Disclosed herein are also variable heavy and light chains that comprise the CDR sets of the invention, as well as full length heavy and light chains (e.g. comprising constant regions as well). As will be appreciated by those in the art, the CDR sets of the anti-LY75 antibody can be incorporated into murine, humanized or human constant regions (including framework regions).

Accordingly, the present disclosure provides variable heavy and light chains that are at least about 90%-99% identical to the SEQ IDs disclosed herein, with 90, 91, 92, 93, 94, 95, 96, 97, 98 and 99% all finding use in the present invention.

In some embodiments, the anti-LY75 antibody is one which competes for binding to human LY75 with an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2. Antibodies that compete for binding can be identified using routine techniques. Such techniques include, for example, an immunoassay, which shows the ability of one antibody to block the binding of another antibody to a target antigen, *i.e.*, a competitive binding assay. Competitive binding is determined in an assay in which the immunoglobulin under test inhibits specific binding of a reference antibody to a common antigen, such as LY75. Numerous types of competitive binding assays are known, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahl *et al.*, *Methods in Enzymology* 9:242 (1983)); solid phase direct biotin-avidin EIA (see Kirkland *et al.*, *J. Immunol.* 137:3614 (1986)); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press (1988)); solid phase direct label RIA using I-125 label (see Morel *et al.*, *Mol. Immunol.* 25(1):7 (1988)); solid phase direct biotin-avidin EIA (Cheung *et al.*, *Virology* 176:546 (1990)); and direct labeled RIA. (Moldenhauer *et al.*, *Scand. J. Immunol.* 32:77 (1990)). Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 50-55%, 55-60%, 60-65%, 65-70% 70-75% 75-80% 80-85% 85-90% 90-95% 95-99% or more.

Monoclonal antibodies can be characterized for binding to LY75 or CD20 using a variety of known techniques. Generally, the antibodies are initially characterized by ELISA. Briefly, microtiter plates can be coated with purified LY75 or CD20 in PBS, and then blocked with irrelevant proteins such as bovine serum albumin (BSA) diluted in PBS. Dilutions of plasma from LY75-immunized mice or CD20-immunized mice are added to each well and incubated for 1-2 hours at 37°C. The plates are washed with PBS/Tween 20 and then incubated with a goat-anti-

human IgG Fc-specific polyclonal reagent conjugated to alkaline phosphatase for 1 hour at 37°C. After washing, the plates are developed with ABTS substrate, and analyzed at OD of 405. Preferably, mice which develop the highest titers will be used for fusions.

An ELISA assay as described above can be used to screen for antibodies and, thus, hybridomas that produce antibodies that show positive reactivity with the LY75/CD20 immunogen. Hybridomas that bind, preferably with high affinity, to LY75/CD20 can then be sub-cloned and further characterized. One clone from each hybridoma, which retains the reactivity of the parent cells (by ELISA), can then be chosen for making a cell bank, and for antibody purification.

To purify anti-LY75 or anti-CD20 antibodies, selected hybridomas can be grown in roller bottles, two-liter spinner-flasks or other culture systems. Supernatants can be filtered and concentrated before affinity chromatography with protein A-Sepharose (Pharmacia, Piscataway, NJ) to purify the protein. After buffer exchange to PBS, the concentration can be determined by OD₂₈₀ using 1.43 extinction coefficient or preferably by nephelometric analysis. IgG can be checked by gel electrophoresis and by antigen specific method.

To determine if the selected anti-LY75 or anti-CD20 monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (Pierce, Rockford, IL). Biotinylated MAb binding can be detected with a streptavidin labeled probe. To determine the isotype of purified antibodies, isotype ELISAs can be performed using art recognized techniques. For example, wells of microtiter plates can be coated with 10 µg/ml of anti- Ig overnight at 4°C. After blocking with 5% BSA, the plates are reacted with 10 µg/ml of monoclonal antibodies or purified isotype controls, at ambient temperature for two hours. The wells can then be reacted with either IgG1 or other isotype specific conjugated probes. Plates are developed and analyzed as described above.

To test the binding of monoclonal antibodies to live cells expressing LY75 or CD20, flow cytometry can be used. Briefly, cell lines and/or human PBMCs expressing membrane-bound LY75 or CD20 (grown under standard growth conditions) are mixed with various concentrations of monoclonal antibodies in PBS containing 0.1% BSA at 4°C for 1 hour. After washing, the cells are reacted with Fluorescein-labeled anti- IgG antibody under the same conditions as the primary antibody staining. The samples can be analyzed by FACScan instrument using light and side scatter properties to gate on single cells and binding of the labeled antibodies is determined. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells can be stained exactly as described above and examined by fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

Anti-LY75 IgGs can be further tested for reactivity with the LY75 antigen by Western blotting; the same may be done with anti-CD20 antibodies. Briefly, cell extracts from cells

expressing LY75/CD20 can be prepared and subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens will be transferred to nitrocellulose membranes, blocked with 20% mouse serum, and probed with the monoclonal antibodies to be tested. IgG binding can be detected using anti- IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, MO).

Methods for analyzing binding affinity, cross-reactivity, and binding kinetics of various anti-LY75 antibodies include standard assays known in the art, for example, Biacore™ surface plasmon resonance (SPR) analysis using a Biacore™ 2000 SPR instrument (Biacore AB, Uppsala, Sweden).

In one embodiment, the antibody specifically binds to human LY75 comprising SEQ ID NO: 15) Preferably, the anti-LY75 antibody binds to human LY75 with high affinity.

Preferably, the anti-LY75 antibody binds to a LY75 protein with a K_D of 5×10^{-8} M or less, binds to a LY75 protein with a K_D of 2×10^{-8} M or less, binds to a LY75 protein with a K_D of 5×10^{-9} M or less, binds to a LY75 protein with a K_D of 4×10^{-9} M or less, binds to a LY75 protein with a K_D of 3×10^{-9} M or less, binds to a LY75 protein with a K_D of 2×10^{-9} M or less, binds to a LY75 protein with a K_D of 1×10^{-9} M or less, binds to a LY75 protein with a K_D of 5×10^{-10} M or less, or binds to a LY75 protein with a K_D of 1×10^{-10} M or less.

In one embodiment, anti-LY75 antibodies compete (e.g., cross-compete) for binding to LY75 with the particular anti-LY75 antibodies described herein (e.g., LY75_A1). Such competing antibodies can be identified based on their ability to competitively inhibit binding to LY75 of one or more of mAbs in standard LY75 binding assays. For example, standard ELISA assays can be used in which a recombinant human LY75 protein is immobilized on the plate, one of the antibodies is fluorescently labeled and the ability of non-labeled antibodies to compete off the binding of the labeled antibody is evaluated. Additionally or alternatively, BIAcore analysis can be used to assess the ability of the antibodies to cross-compete. The ability of a test antibody to inhibit the binding of an anti-LY75 antibody of the invention to human LY75 demonstrates that the test antibody can compete with the antibody for binding to human LY75.

In one embodiment, the competing antibody is an antibody that binds to the same epitope on human LY75 as the particular anti-LY75 monoclonal antibodies described herein (e.g., LY75_A1). Standard epitope mapping techniques, such as x-ray crystallography and 2-dimensional nuclear magnetic resonance, can be used to determine whether an antibody binds to the same epitope as a reference antibody (see, e.g., Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66, G. E. Morris, Ed. (1996)).

In one embodiment, the antibody that competes for binding to LY75 and/or binds to the same epitope on human LY75 is a human antibody.

Once a single, archetypal anti-LY75 mAb has been isolated that has the desired properties described herein, other mAbs with similar properties, e.g., having the same epitope

may be generated. For example, mice may be immunized with LY75 as described herein, hybridomas produced, and the resulting mAbs screened for the ability to compete with the archetypal mAb for binding to LY75. Mice can also be immunized with a smaller fragment of LY75 containing the epitope to which the archetypal mAb binds. The epitope can be localized by, e.g., screening for binding to a series of overlapping peptides spanning LY75. Alternatively, the method of Jespers et al., *Biotechnology* 12:899, 1994 may be used to guide the selection of mAbs having the same epitope and therefore similar properties to the archetypal mAb. Using phage display, first the heavy chain of the archetypal antibody is paired with a repertoire of (preferably human) light chains to select a LY75-binding mAb, and then the new light chain is paired with a repertoire of (preferably human) heavy chains to select a (preferably human) LY75-binding mAb having the same epitope as the archetypal mAb. Alternatively variants of the archetypal mAb can be obtained by mutagenesis of cDNA encoding the heavy and light chains of the antibody.

To assess the level of competition between two antibodies, for example, radio-immunoassays or assays using other labels for the antibodies, can be used. For example, a LY75 antigen can be incubated with a saturating amount of a first anti-LY75 antibody or antigen-binding fragment thereof conjugated to a labeled compound (e.g., ^3H , ^{125}I , biotin, or rubidium) in the presence the same amount of a second unlabeled anti-LY75 antibody. The amount of labeled antibody that is bound to the antigen in the presence of the unlabeled blocking antibody is then assessed and compared to binding in the absence of the unlabeled blocking antibody. Competition is determined by the percentage change in binding signals in the presence of the unlabeled blocking antibody compared to the absence of the blocking antibody. Thus, if there is a 50% inhibition of binding of the labeled antibody in the presence of the blocking antibody compared to binding in the absence of the blocking antibody, then there is competition between the two antibodies of 50%. Thus, reference to competition between a first and second antibody of 50% or more, 60% or more, 70% or more, such as 70%, 71%, 72%, 73%, 74%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more, means that the first antibody inhibits binding of the second antibody (or vice versa) to the antigen by 50%, 60%, 70%, 71%, 72%, 73%, 74%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more (compared to binding of the antigen by the second antibody in the absence of the first antibody). Thus, inhibition of binding of a first antibody to an antigen by a second antibody of 50%, 60%, 70%, 71%, 72%, 73%, 74%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more indicates that the two antibodies bind to the same epitope.

In some embodiments, component (B) of the pharmaceutical combination is an anti-CD20 antibody as defined herein.

B-lymphocyte antigen CD20 or CD20 is an activated-glycosylated phosphoprotein expressed on the surface of all B-cells beginning at the pro-B phase (CD45R+, CD117+) and

progressively increasing in concentration until maturity (Hardy, Richard (2008). "Chapter 7: B Lymphocyte Development and Biology". In Paul, William. *Fundamental Immunology* (Book) (6th ed.). Philadelphia: Lippincott Williams & Wilkins. pp. 237–269. ISBN 0-7817-6519-6.)

CD20 is expressed on all stages of B-cell development except the first and last; it is present from late pro-B cells through memory cells, but not on either early pro-B cells or plasma blasts and plasma cells. It is found on B-cell lymphomas, hairy cell leukemia, B-cell chronic lymphocytic leukemia, and melanoma cancer stem cells. The expression of CD20 is regulated by the chemokine signalling through the CXCR4/SDF1 axis.

Immunohistochemistry can be used to determine the presence of CD20 on cells in histological tissue sections.

In humans, CD20 is encoded by the MS4A1 gene. This gene encodes a member of the membrane-spanning 4A gene family.

In one embodiment, component (B) of the pharmaceutical combination comprises an anti-CD20 antibody, or an antigen-binding portion thereof, said antibody comprising:

- a) a heavy chain variable region comprising:
 - i) a first vhCDR comprising SEQ ID NO: 40;
 - ii) a second vhCDR comprising SEQ ID NO: 41; and
 - iii) a third vhCDR comprising SEQ ID NO: 42; and
- b) a light chain variable region comprising:
 - i) a first vlCDR comprising SEQ ID NO: 43;
 - ii) a second vlCDR comprising SEQ ID NO: 44; and
 - iii) a third vlCDR comprising SEQ ID NO: 45;

optionally wherein any one or more of the above SEQ ID NOs independently comprise one, two, three, four or five amino acid substitutions, additions or deletions.

Optionally, one or more of the above SEQ ID NOs: 40-45 independently comprise one, two, three, four or five conservative amino acid substitutions; more preferably, any one or more of the above SEQ ID NOs independently comprise one or two conservative amino acid substitutions.

Preferably, the anti-CD-20 antibody is a monoclonal antibody, more preferably a human, monoclonal antibody, and even more preferably a fully human IgG1 monoclonal antibody. In a particularly preferred embodiment of the invention, the anti-CD20 antibody is Rituximab. Rituximab is a monoclonal antibody against the protein CD20 which is widely expressed on B cells (Oncogene (2003 Oct 20), Smith MR, "Rituximab (monoclonal anti-CD20 antibody): mechanisms of action and resistance", 22(47): 7359-68). Rituximab destroys both normal and malignant B cells that have CD20 on their surfaces and is therefore used to treat diseases which

are characterized by having too many B cells, overactive B cells, or dysfunctional B cells.

Rituximab has previously been used to treat a number of autoimmune diseases and some types of cancer, including rheumatoid arthritis, idiopathic thrombocytopenic purpura, pemphigus vulgaris, multiple sclerosis, systemic lupus erythematosus, non-Hodgkin's lymphoma, chronic inflammatory demyelinating polyneuropathy, chronic lymphocytic leukemia and autoimmune anemias. Rituximab is sold under the tradename Rituxan® by Roche.

Preferably, the heavy and/or light chain of the rituximab antibody comprises or consists of the amino acid sequence given in SEQ ID NO: 38 and 39, respectively.

The present invention encompasses variant antibodies, sometimes referred to as “antibody derivatives” or “antibody analogs” as well. That is, there are a number of modifications that can be made to the antibodies disclosed herein, including, but not limited to, amino acid modifications in the CDRs (affinity maturation), amino acid modifications in the framework regions, amino acid modifications in the Fc region, glycosylation variants, covalent modifications of other types (e.g. for attachment of drug conjugates, etc.).

By “variant” herein is meant a polypeptide sequence that differs from that of a parent polypeptide by virtue of at least one amino acid modification. In this case, the parent polypeptide is either the full length variable heavy or light chains, e.g. as listed in SEQ ID NOs: 1 or 2, respectively or the CDR regions or the framework regions of the heavy and light chains listed in SEQ ID NOs 5-10 and 16-21 for LY75. Amino acid modifications can include substitutions, insertions and deletions, with the former being preferred in many cases. It will be understood that an amino acid substitution may be a conservative or non-conservative substitution with conservative substitutions being preferred. Further said substitution may be a substitution with either a naturally or non-naturally occurring amino acid.

In general, variants can include any number of modifications, as long as the function of the antibody is still present, as described herein. That is, LY75_A1, for example, the antibody should still specifically bind to human LY75. Similarly, anti-CD20 antibodies should still specifically bind to human CD20. If amino acid variants are generated with the Fc region, for example, the variant antibodies should maintain the required receptor binding functions for the particular application or indication of the antibody.

“Variants” in this case can be made in either the listed CDR sequences, the framework or Fc regions of the antibody.

However, in general, from 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acid substitutions are generally utilized as often the goal is to alter function with a minimal number of modifications. In some cases, there are from 1 to 5 modifications (e.g. individual amino acid substitutions, insertions or deletions), with from 1-2, 1-3 and 1-4 also finding use in many embodiments. The number of modifications can depend on the size of the region being modified; for example, in general, fewer modifications are desired in CDR regions. It will be understood by the skilled

person that even within the CDR regions the location of the modification can significantly alter the effect. In one embodiment, the modifications can be made in any of CDR1, CDR2 or CDR3 of the heavy and/or light chains. In a further embodiment, the modifications are made in any of CDR1 or CDR2 of the heavy and/or light chains. In a still further embodiment, the modifications are located in CDR1 of the heavy and/or light chains.

It should be noted that the number of amino acid modifications may be within functional domains: for example, it may be desirable to have from 1-5 modifications in the Fc region of wild-type or engineered proteins, as well as from 1 to 5 modifications in the Fv region, for example. A variant polypeptide sequence will preferably possess at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to the parent sequences (e.g. the variable regions, the constant regions, and/or the heavy and light chain sequences and/or the CDRs of LY75_A1 or Rituximab). It should be noted that depending on the size of the sequence, the percent identity will depend on the number of amino acids.

By "amino acid substitution" or "substitution" herein is meant the replacement of an amino acid at a particular position in a parent polypeptide sequence with another amino acid which may be a natural or non-naturally occurring amino acid. For example, the substitution S100A refers to a variant polypeptide in which the serine at position 100 is replaced with alanine. By "amino acid insertion" or "insertion" as used herein is meant the addition of an amino acid at a particular position in a parent polypeptide sequence. By "amino acid deletion" or "deletion" as used herein is meant the removal of an amino acid at a particular position in a parent polypeptide sequence.

By "parent polypeptide", "parent protein", "precursor polypeptide", or "precursor protein" as used herein is meant an unmodified polypeptide that is subsequently modified to generate a variant. In general, the parent polypeptides herein are LY75_A1 and Rituximab. Accordingly, by "parent antibody" as used herein is meant an antibody that is modified to generate a variant antibody.

By "wild type" or "WT" or "native" herein is meant an amino acid sequence or a nucleotide sequence that is found in nature, including allelic variations. A WT protein, polypeptide, antibody, immunoglobulin, IgG, etc. has an amino acid sequence or a nucleotide sequence that has not been intentionally modified.

By "variant Fc region" herein is meant an Fc sequence that differs from that of a wild-type Fc sequence by virtue of at least one amino acid modification. Fc variant may refer to the Fc polypeptide itself, compositions comprising the Fc variant polypeptide, or the amino acid sequence.

In some embodiments, one or more amino acid modifications are made in one or more of the CDRs of LY75_A1 or Rituximab. In general, only 1 or 2 or 3 amino acids are substituted in any single CDR, and generally no more than from 4, 5, 6, 7, 8, 9 or 10 changes are made within a set of 6 CDRs. However, it should be appreciated that any combination of no substitutions, 1, 2

or 3 substitutions in any CDR can be independently and optionally combined with any other substitution. It will be apparent that substitutions can be made in any of the 6 CDRs. In one embodiment, substitutions are made in CDR1 of the heavy and/or light chains.

In some cases, amino acid modifications in the CDRs are referred to as "affinity maturation". An "affinity matured" antibody is one having one or more alteration(s) in one or more CDRs which results in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In some cases, although rare, it may be desirable to decrease the affinity of an antibody to its antigen, but this is generally not preferred.

Affinity maturation can be done to increase the binding affinity of the antibody for the antigen by at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 140%, about 150% or more, or 1, 2, 3, 4 to 5 fold as compared to the "parent" antibody. Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by known procedures. See, for example, Marks *et al.*, 1992, Biotechnology 10:779-783 that describes affinity maturation by variable heavy chain (VH) and variable light chain (VL) domain shuffling. Random mutagenesis of CDR and/or framework residues is described in: Barbas, *et al.* 1994, Proc. Nat. Acad. Sci, USA 91:3809-3813; Shier *et al.*, 1995, Gene 169:147-155; Yelton *et al.*, 1995, J. Immunol. 155:1994-2004; Jackson *et al.*, 1995, J. Immunol. 154(7):3310-9; and Hawkins *et al.*, 1992, J. Mol. Biol. 226:889-896, for example.

Alternatively, amino acid modifications can be made in one or more of the CDRs of the antibodies of the invention that are "silent", e.g. that do not significantly alter the affinity of the antibody for the antigen. These can be made for a number of reasons, including optimizing expression (as can be done for the nucleic acids encoding the antibodies of the invention).

Thus, included within the definition of the CDRs and antibodies disclosed herein are variant CDRs and antibodies; that is, the antibodies can include amino acid modifications in one or more of the CDRs of LY75_A1 or Rituximab. In addition, as outlined below, amino acid modifications can also independently and optionally be made in any region outside the CDRs, including framework and constant regions as described herein.

In some embodiments, the anti-LY75 antibodies and/or anti-CD20 antibodies disclosed herein are composed of a variant Fc domain. As is known in the art, the Fc region of an antibody interacts with a number of Fc receptors and ligands, imparting an array of important functional capabilities referred to as effector functions. These Fc receptors include, but are not limited to, (in humans) FcγRI (CD64) including isoforms FcγRIa, FcγRIb, and FcγRIc; FcγRII (CD32), including isoforms FcγRIIa (including allotypes H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and FcγRIIc; and FcγRIII (CD16), including isoforms FcγRIIIa (including

allotypes V158 and F158, correlated to antibody-dependent cell cytotoxicity (ADCC)) and FcγRIIIb (including allotypes FcγRIIIb-NA1 and FcγRIIIb-NA2), FcRn (the neonatal receptor), C1q (complement protein involved in complement dependent cytotoxicity (CDC)) and FcRn (the neonatal receptor involved in serum half-life). Suitable modifications can be made at one or more positions as is generally outlined, for example in US Patent Application 11/841,654 and references cited therein, US 2004/013210, US 2005/0054832, US 2006/0024298, US 2006/0121032, US 2006/0235208, US 2007/0148170, USSN 12/341,769, US Patent No. 6,737,056, US Patent No. 7,670,600, US Patent No. 6,086,875 all of which are expressly incorporated by reference in their entirety, and in particular for specific amino acid substitutions that increase binding to Fc receptors.

In addition to the modifications outlined above, other modifications can be made. For example, the molecules may be stabilized by the incorporation of disulphide bridges linking the VH and VL domains (Reiter et al., 1996, Nature Biotech. 14:1239-1245, entirely incorporated by reference).

In addition, modifications at cysteines are particularly useful in antibody-drug conjugate (ADC) applications, further described below. In some embodiments, the constant region of the antibodies can be engineered to contain one or more cysteines that are particularly "thiol reactive", so as to allow more specific and controlled placement of the drug moiety. See for example US Patent No. 7,521,541, incorporated by reference in its entirety herein.

In addition, there are a variety of covalent modifications of antibodies that can be made as outlined below.

Covalent modifications of antibodies are included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antibody are introduced into the molecule by reacting specific amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteinyl residues most commonly are reacted with α-haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues may also be derivatized by reaction with bromotrifluoroacetone, α-bromo-β-(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole and the like.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl

residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ¹²⁵I or ¹³¹I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R'—N=C=N—R'), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking antibodies to a water-insoluble support matrix or surface for use in a variety of methods, in addition to methods described below. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis (succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propionimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cynomolgusogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440, all entirely incorporated by reference, are employed for protein immobilization.

Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α-amino groups of lysine, arginine, and

histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 [1983], entirely incorporated by reference), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

In addition, as will be appreciated by those in the art, labels (including fluorescent, enzymatic, magnetic, radioactive, etc. can all be added to the antibodies (as well as the other compositions of the invention).

Another type of covalent modification is alterations in glycosylation. In some embodiments, the antibodies disclosed herein can be fully or partially aglycosylated, e.g. afucosylated.

Another type of covalent modification of the antibody comprises linking the antibody to various non-proteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in, for example, 2005-2006 PEG Catalog from Nektar Therapeutics (available at the Nektar website) US Patents 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337, all entirely incorporated by reference. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody to facilitate the addition of polymers such as PEG. See for example, U.S. Publication No. 2005/0114037A1, entirely incorporated by reference.

In additional embodiments, the antibodies may comprise a label. By "labeled" herein is meant that a compound has at least one element, isotope or chemical compound attached to enable the detection of the compound. In general, labels fall into three classes: a) isotopic labels, which may be radioactive or heavy isotopes; b) magnetic, electrical, thermal; and c) colored or luminescent dyes; although labels include enzymes and particles such as magnetic particles as well. Preferred labels include, but are not limited to, fluorescent lanthanide complexes (including those of Europium and Terbium), and fluorescent labels including, but not limited to, quantum dots, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade Blue, Texas Red, the Alexa dyes, the Cy dyes, and others described in the 6th Edition of the *Molecular Probes Handbook* by Richard P. Haugland, hereby expressly incorporated by reference.

Antibody-Drug Conjugates

In some embodiments, the anti-LY75 antibodies disclosed herein are conjugated with drugs to form antibody-drug conjugates (ADCs). In general, ADCs are used in oncology applications, where the use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents allows for the targeted delivery of the drug moiety to tumors, which can allow higher efficacy, lower toxicity, etc. An overview of this technology is provided in Ducry et al., *Bioconjugate Chem.*, 21:5-13 (2010), Carter et al., *Cancer J.* 14(3):154 (2008) and Senter, *Current Opin. Chem. Biol.* 13:235-244 (2009), all of which are hereby incorporated by reference in their entirety.

Thus the invention provides pharmaceutical combinations comprising, *inter alia*, anti-LY75 antibodies conjugated to drugs. Generally, conjugation is done by covalent attachment to the antibody, as further described below, and generally relies on a linker, often a peptide linkage (which, as described below, may be designed to be sensitive to cleavage by proteases at the target site or not). In addition, as described above, linkage of the linker-drug unit (LU-D) can be done by attachment to cysteines within the antibody. As will be appreciated by those in the art, the number of drug moieties per antibody can change, depending on the conditions of the reaction, and can vary from 1:1 to 10:1 drug:antibody. As will be appreciated by those in the art, the actual number is an average.

Thus the anti-LY75 antibodies may be conjugated to drugs. As described below, the drug of the ADC can be any number of agents, including but not limited to cytotoxic agents such as chemotherapeutic agents, growth inhibitory agents, toxins (for example, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (that is, a radioconjugate) are provided. In other embodiments, the invention further provides methods of using the ADCs.

Drugs for use in the present invention include cytotoxic drugs, particularly those which are used for cancer therapy. Such drugs include, in general, DNA damaging agents, anti-metabolites, natural products and their analogs. Exemplary classes of cytotoxic agents include the enzyme inhibitors such as dihydrofolate reductase inhibitors, and thymidylate synthase inhibitors, DNA intercalators, DNA cleavers, topoisomerase inhibitors, the anthracycline family of drugs, the vinca drugs, the mitomycins, the bleomycins, the cytotoxic nucleosides, the pteridine family of drugs, diynenes, the podophyllotoxins, dolastatins, maytansinoids, differentiation inducers, and taxols.

Members of these classes include, for example, taxol, methotrexate, methopterin, dichloromethotrexate, 5-fluorouracil, 6-mercaptopurine, cytosine arabinoside, melphalan, leurosine, leurosideine, actinomycin, daunorubicin, doxorubicin, mitomycin C, mitomycin A, caminomycin, aminopterin, tallysomylin, podophyllotoxin and podophyllotoxin derivatives such as etoposide or etoposide phosphate, vinblastine, vincristine, vindesine, taxanes including taxol, taxotere, retinoic acid, butyric acid, N8-acetyl spermidine, camptothecin, calicheamicin, esperamicin, ene-diynes, duocarmycin A, duocarmycin SA, calicheamicin, camptothecin, hemiasterlins, maytansinoids (including DM1), monomethylauristatin E (MMAE), monomethylauristatin F (MMAF), and maytansinoids (DM4) and their analogues.

Toxins may be used as antibody-toxin conjugates and include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) J. Nat. Cancer Inst. 92(19):1573-1581; Mandler et al (2000) Bioorganic & Med. Chem. Letters 10:1025-1028; Mandler et al (2002) Bioconjugate Chem. 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) Proc. Natl. Acad. Sci. USA 93:8618-8623), and

calicheamicin (Lode et al (1998) Cancer Res. 58:2928; Hinman et al (1993) Cancer Res. 53:3336-3342), hemiasterlins (WO2004/026293; Zask *et al.*, (2004) J. Med. Chem, 47: 4774-4786). Toxins may exert their cytotoxic and cytostatic effects by mechanisms including tubulin binding, DNA binding, or topoisomerase inhibition.

5 Conjugates of an anti-LY75 antibody and one or more small molecule toxins, such as a maytansinoids, dolastatins, auristatins, a trichothecene, calicheamicin, duocarmycins, pyrrolobenzodiazepines and CC1065, and the derivatives of these toxins that have toxin activity, may also be used.

10 Preferably, the anti-LY75 antibody is conjugated to DM1 or DM4, most preferably to DM4. Maytansine compounds suitable for use as maytansinoid drug moieties are well known in the art, and can be isolated from natural sources according to known methods, produced using genetic engineering techniques (see Yu et al (2002) PNAS 99:7968-7973), or maytansinol and maytansinol analogues prepared synthetically according to known methods. As described below, drugs may be modified by the incorporation of a functionally active group such as a thiol or amine group for conjugation to the antibody.

15 Exemplary maytansinoid drug moieties include those having a modified aromatic ring, such as: C-19-dechloro (U.S. Pat. No. 4,256,746) (prepared by lithium aluminum hydride reduction of ansamycin P2); C-20-hydroxy (or C-20-demethyl) +/-C-19-dechloro (U.S. Pat. Nos. 4,361,650 and 4,307,016) (prepared by demethylation using *Streptomyces* or *Actinomyces* or dechlorination using LAH); and C-20-demethoxy, C-20-acyloxy (--OCOR), +/-dechloro (U.S. Pat. No. 4,294,757) (prepared by acylation using acyl chlorides) and those having modifications at other positions.

25 Exemplary maytansinoid drug moieties also include those having modifications such as: C-9-SH (U.S. Pat. No. 4,424,219) (prepared by the reaction of maytansinol with H₂S or P₂S₅); C-14-alkoxymethyl(demethoxy/CH₂OR) (U.S. Pat. No. 4,331,598); C-14-hydroxymethyl or acyloxymethyl (CH₂OH or CH₂OAc) (U.S. Pat. No. 4,450,254) (prepared from *Nocardia*); C-15-hydroxy/acyloxy (U.S. Pat. No. 4,364,866) (prepared by the conversion of maytansinol by *Streptomyces*); C-15-methoxy (U.S. Pat. Nos. 4,313,946 and 4,315,929) (isolated from *Trewia nudiflora*); C-18-N-demethyl (U.S. Pat. Nos. 4,362,663 and 4,322,348) (prepared by the demethylation of maytansinol by *Streptomyces*); and 4,5-deoxy (U.S. Pat. No. 4,371,533) (prepared by the titanium trichloride/LAH reduction of maytansinol).

30 Of particular use are DM1 (disclosed in US Patent No. 5,208,020, incorporated by reference) and DM4 (disclosed in US Patent No. 7,276,497, incorporated by reference). See also a number of additional maytansinoid derivatives and methods in 5,416,064, WO/01/24763, 7,303,749, 7,601,354, USSN 12/631,508, WO02/098883, 6,441,163, 7,368,565, WO02/16368 and WO04/1033272, all of which are expressly incorporated by reference in their entirety.

ADCs containing maytansinoids, methods of making same, and their therapeutic use are disclosed, for example, in U.S. Pat. Nos. 5,208,020; 5,416,064; 6,441,163 and European Patent EP 0 425 235 B1, the disclosures of which are hereby expressly incorporated by reference. Liu et al., Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described ADCs comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an in vivo tumor growth assay.

Chari et al., Cancer Research 52:127-131 (1992) describe ADCs in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an antigen on human colon cancer cell lines, or to another murine monoclonal antibody TA.1 that binds the HER-2/neu oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested in vitro on the human breast cancer cell line SK-BR-3, which expresses 3x10⁵ HER-2 surface antigens per cell. The drug conjugate achieved a degree of cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

For compositions comprising a plurality of antibodies, the drug loading is represented by p, the average number of drug molecules per Antibody. Drug loading may range from 1 to 20 drugs (D) per Antibody. The average number of drugs per antibody in preparation of conjugation reactions may be characterized by conventional means such as mass spectroscopy, ELISA assay, and HPLC. The quantitative distribution of Antibody-Drug-Conjugates in terms of p may also be determined.

In some instances, separation, purification, and characterization of homogeneous Antibody-Drug-conjugates where p is a certain value from Antibody-Drug-Conjugates with other drug loadings may be achieved by means such as reverse phase HPLC or electrophoresis. In exemplary embodiments, p is 2, 3, 4, 5, 6, 7, or 8 or a fraction thereof.

The generation of Antibody-drug conjugate compounds can be accomplished by any technique known to the skilled artisan. Briefly, the Antibody-drug conjugate compounds can include an anti-LY75 antibody as the Antibody unit, a drug, and optionally a linker that joins the drug and the binding agent.

A number of different reactions are available for covalent attachment of drugs and/or linkers to binding agents. This can be accomplished by reaction of the amino acid residues of the binding agent, for example, antibody molecule, including the amine groups of lysine, the free carboxylic acid groups of glutamic and aspartic acid, the sulfhydryl groups of cysteine and the various moieties of the aromatic amino acids. A commonly used non-specific method of covalent attachment is the carbodiimide reaction to link a carboxy (or amino) group of a compound to amino (or carboxy) groups of the antibody. Additionally, bifunctional agents such as

dialdehydes or imidoesters have been used to link the amino group of a compound to amino groups of an antibody molecule.

Also available for attachment of drugs to binding agents is the Schiff base reaction. This method involves the periodate oxidation of a drug that contains glycol or hydroxy groups, thus forming an aldehyde which is then reacted with the binding agent. Attachment occurs via formation of a Schiff base with amino groups of the binding agent. Isothiocyanates can also be used as coupling agents for covalently attaching drugs to binding agents. Other techniques are known to the skilled artisan and within the scope of the present invention.

In some embodiments, an intermediate, which is the precursor of the linker, is reacted with the drug under appropriate conditions. In other embodiments, reactive groups are used on the drug and/or the intermediate. The product of the reaction between the drug and the intermediate, or the derivatized drug, is subsequently reacted with an anti-LY75 antibody of the invention under appropriate conditions.

It will be understood that chemical modifications may also be made to the desired compound in order to make reactions of that compound more convenient for purposes of preparing conjugates of the invention. For example a functional group e.g. amine, hydroxyl, or sulfhydryl, may be appended to the drug at a position which has minimal or an acceptable effect on the activity or other properties of the drug.

Typically, the antibody-drug conjugate compounds comprise a Linker unit between the drug unit and the antibody unit. In some embodiments, the linker is cleavable under intracellular or extracellular conditions, such that cleavage of the linker releases the drug unit from the antibody in the appropriate environment. For example, solid tumors that secrete certain proteases may serve as the target of the cleavable linker; in other embodiments, it is the intracellular proteases that are utilized. In yet other embodiments, the linker unit is not cleavable and the drug is released, for example, by antibody degradation in lysosomes.

In some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (for example, within a lysosome or endosome or caveolea). The linker can be, for example, a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. In some embodiments, the peptidyl linker is at least two amino acids long or at least three amino acids long or more.

Cleaving agents can include, without limitation, cathepsins B and D and plasmin, all of which are known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells (see, e.g., Dubowchik and Walker, 1999, Pharm. Therapeutics 83:67-123). Peptidyl linkers that are cleavable by enzymes that are present in LY75-expressing cells. For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a Phe-Leu or a Gly-Phe-Leu-Gly linker

(SEQ ID NO: 52)). Other examples of such linkers are described, e.g., in U.S. Pat. No. 6,214,345, incorporated herein by reference in its entirety and for all purposes.

In some embodiments, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker or a Phe-Lys linker (see, e.g., U.S. Pat. No. 6,214,345, which describes the synthesis of doxorubicin with the val-cit linker).

In other embodiments, the cleavable linker is pH-sensitive, that is, sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acid-labile linker that is hydrolyzable in the lysosome (for example, a hydrazone, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like) may be used. (See, e.g., U.S. Pat. Nos. 5,122,368; 5,824,805; 5,622,929; Dubowchik and Walker, 1999, Pharm. Therapeutics 83:67-123; Neville et al., 1989, Biol. Chem. 264:14653-14661.) Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome. In certain embodiments, the hydrolyzable linker is a thioether linker (such as, e.g., a thioether attached to the therapeutic agent via an acylhydrazone bond (see, e.g., U.S. Pat. No. 5,622,929).

In yet other embodiments, the linker is cleavable under reducing conditions (for example, a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-5-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), SPDB (N-succinimidyl-3-(2-pyridyldithio)butyrate) and SMPT (N-succinimidyl-oxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio)toluene)-, SPDB and SMPT. (See, e.g., Thorpe et al., 1987, Cancer Res. 47:5924-5931; Wawrzynczak et al., In Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer (C. W. Vogel ed., Oxford U. Press, 1987. See also U.S. Pat. No. 4,880,935).

In other embodiments, the linker is a malonate linker (Johnson et al., 1995, Anticancer Res. 15:1387-93), a maleimidobenzoyl linker (Lau et al., 1995, Bioorg-Med-Chem. 3(10):1299-1304), or a 3'-N-amide analog (Lau et al., 1995, Bioorg-Med-Chem. 3(10):1305-12).

In yet other embodiments, the linker unit is not cleavable and the drug is released by antibody degradation. (See U.S. Publication No. 2005/0238649 incorporated by reference herein in its entirety and for all purposes).

In many embodiments, the linker is self-immolative. As used herein, the term "self-immolative Spacer" refers to a bifunctional chemical moiety that is capable of covalently linking together two spaced chemical moieties into a stable tripartite molecule. It will spontaneously separate from the second chemical moiety if its bond to the first moiety is cleaved. See for example, WO 2007/059404A2, WO06/110476A2, WO05/112919A2, WO2010/062171, WO09/017394, WO07/089149, WO 07/018431, WO04/043493 and WO02/083180, which are directed to drug-cleavable substrate conjugates where the drug and cleavable substrate are

optionally linked through a self-immolative linker and which are all expressly incorporated by reference.

Often the linker is not substantially sensitive to the extracellular environment. As used herein, "not substantially sensitive to the extracellular environment," in the context of a linker, means that no more than about 20%, 15%, 10%, 5%, 3%, or no more than about 1% of the linkers, in a sample of antibody-drug conjugate compound, are cleaved when the antibody-drug conjugate compound presents in an extracellular environment (for example, in plasma).

Whether a linker is not substantially sensitive to the extracellular environment can be determined, for example, by incubating with plasma the antibody-drug conjugate compound for a predetermined time period (for example, 2, 4, 8, 16, or 24 hours) and then quantitating the amount of free drug present in the plasma.

In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (that is, in the milieu of the linker-therapeutic agent moiety of the antibody-drug conjugate compound as described herein). In yet other embodiments, the linker promotes cellular internalization when conjugated to both the auristatin compound and the anti-LY75 antibodies of the invention.

A variety of exemplary linkers that can be used with the present compositions and methods are described in WO 2004/010957, U.S. Publication No. 2006/0074008, U.S. Publication No. 20050238649, and U.S. Publication No. 2006/0024317 (each of which is incorporated by reference herein in its entirety and for all purposes). Preferably, the linker is SPDB (N-succinimidyl-3-(2-pyridyldithio)butyrate).

Drug loading is represented by p and is the average number of Drug moieties per antibody in a molecule. Drug loading (" p ") may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more moieties (D) per antibody, although frequently the average number is a fraction or a decimal. Generally, drug loading of from 1 to 4 is frequently useful, and from 1 to 2 is also useful. ADCs of the invention include collections of antibodies conjugated with a range of drug moieties, from 1 to 20, for example, 1-15, 1-10, 2-9, 3-8, 4-7, 5-6. The average number of drug moieties per antibody in preparations of ADC from conjugation reactions may be characterized by conventional means such as mass spectroscopy and, ELISA assay.

The quantitative distribution of ADC in terms of p may also be determined. In some instances, separation, purification, and characterization of homogeneous ADC where p is a certain value from ADC with other drug loadings may be achieved by means such as electrophoresis.

For some antibody-drug conjugates, p may be limited by the number of attachment sites on the antibody. For example, where the attachment is a cysteine thiol, as in the exemplary embodiments above, an antibody may have only one or several cysteine thiol groups, or may

have only one or several sufficiently reactive thiol groups through which a linker may be attached. In certain embodiments, higher drug loading, e.g. $p > 5$, may cause aggregation, insolubility, toxicity, or loss of cellular permeability of certain antibody-drug conjugates. In certain embodiments, the drug loading for an ADC of the invention ranges from 1 to about 8; from about 2 to about 6; from about 3 to about 5; from about 3 to about 4; from about 3.1 to about 3.9; from about 3.2 to about 3.8; from about 3.2 to about 3.7; from about 3.2 to about 3.6; from about 3.3 to about 3.8; or from about 3.3 to about 3.7. Indeed, it has been shown that for certain ADCs, the optimal ratio of drug moieties per antibody may be less than 8, and may be about 2 to about 5. See US 2005/0238649 A1 (herein incorporated by reference in its entirety).

In certain embodiments, fewer than the theoretical maximum of drug moieties are conjugated to an antibody during a conjugation reaction. An antibody may contain, for example, lysine residues that do not react with the drug-linker intermediate or linker reagent, as discussed below. Generally, antibodies do not contain many free and reactive cysteine thiol groups which may be linked to a drug moiety; indeed most cysteine thiol residues in antibodies exist as disulfide bridges. In certain embodiments, an antibody may be reduced with a reducing agent such as dithiothreitol (DTT) or tricarboylethylphosphine (TCEP), under partial or total reducing conditions, to generate reactive cysteine thiol groups. In certain embodiments, an antibody is subjected to denaturing conditions to reveal reactive nucleophilic groups such as lysine or cysteine.

The loading (drug/antibody ratio) of an ADC may be controlled in different ways, e.g., by: (i) limiting the molar excess of drug-linker intermediate or linker reagent relative to antibody, (ii) limiting the conjugation reaction time or temperature, (iii) partial or limiting reductive conditions for cysteine thiol modification, (iv) engineering by recombinant techniques the amino acid sequence of the antibody such that the number and position of cysteine residues is modified for control of the number and/or position of linker-drug attachments (such as thioMab or thioFab prepared as disclosed herein and in WO2006/034488 (herein incorporated by reference in its entirety)).

It is to be understood that where more than one nucleophilic group reacts with a drug-linker intermediate or linker reagent followed by drug moiety reagent, then the resulting product is a mixture of ADC compounds with a distribution of one or more drug moieties attached to an antibody. The average number of drugs per antibody may be calculated from the mixture by a dual ELISA antibody assay, which is specific for antibody and specific for the drug. Individual ADC molecules may be identified in the mixture by mass spectroscopy and separated by HPLC, e.g. hydrophobic interaction chromatography.

In some embodiments, a homogeneous ADC with a single loading value may be isolated from the conjugation mixture by electrophoresis or chromatography.

Methods of Determining Cytotoxic Effect of ADCs

Methods of determining whether a Drug or Antibody-Drug conjugate exerts a cytostatic and/or cytotoxic effect on a cell are known. Generally, the cytotoxic or cytostatic activity of an Antibody Drug conjugate can be measured by: exposing mammalian cells expressing a target
5 protein of the Antibody Drug conjugate in a cell culture medium; culturing the cells for a period from about 6 hours to about 5 days; and measuring cell viability. Cell-based in vitro assays can be used to measure viability (proliferation), cytotoxicity, and induction of apoptosis (caspase activation) of the Antibody Drug conjugate.

For determining whether an Antibody Drug conjugate exerts a cytostatic effect, a thymidine
10 incorporation assay may be used. For example, cancer cells expressing a target antigen at a density of 5,000 cells/well of a 96-well plated can be cultured for a 72-hour period and exposed to 0.5 μ Ci of ^{3} H-thymidine during the final 8 hours of the 72-hour period. The incorporation of ^{3} H-thymidine into cells of the culture is measured in the presence and absence of the Antibody Drug conjugate.

For determining cytotoxicity, necrosis or apoptosis (programmed cell death) can be
15 measured. Necrosis is typically accompanied by increased permeability of the plasma membrane; swelling of the cell, and rupture of the plasma membrane. Apoptosis is typically characterized by membrane blebbing, condensation of cytoplasm, and the activation of endogenous endonucleases. Determination of any of these effects on cancer cells indicates that
20 an Antibody Drug conjugate is useful in the treatment of cancers.

Cell viability can be measured by determining in a cell the uptake of a dye such as neutral red, trypan blue, or ALAMAR™ blue (see, e.g., Page *et al.*, 1993, Intl. J. Oncology 3:473-476). In such an assay, the cells are incubated in media containing the dye, the cells are washed, and the
25 remaining dye, reflecting cellular uptake of the dye, is measured spectrophotometrically. The protein-binding dye sulforhodamine B (SRB) can also be used to measure cytotoxicity (Skehan *et al.*, 1990, J. Natl. Cancer Inst. 82:1107-12).

Alternatively, a tetrazolium salt, such as MTT, is used in a quantitative colorimetric assay for mammalian cell survival and proliferation by detecting living, but not dead, cells (see, e.g.,
Mosmann, 1983, J. Immunol. Methods 65:55-63).

Apoptosis can be quantitated by measuring, for example, DNA fragmentation. Commercial
30 photometric methods for the quantitative in vitro determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucleotides in fragmented DNA) and ELISA-based assays, are described in Biochemica, 1999, no. 2, pp. 34-37 (Roche Molecular Biochemicals).

Apoptosis can also be determined by measuring morphological changes in a cell. For
35 example, as with necrosis, loss of plasma membrane integrity can be determined by measuring uptake of certain dyes (e.g., a fluorescent dye such as, for example, acridine orange or ethidium

bromide). A method for measuring apoptotic cell number has been described by Duke and Cohen, Current Protocols in Immunology (Coligan et al. eds., 1992, pp. 3.17.1-3.17.16). Cells also can be labeled with a DNA dye (e.g., acridine orange, ethidium bromide, or propidium iodide) and the cells observed for chromatin condensation and margination along the inner
5 nuclear membrane. Other morphological changes that can be measured to determine apoptosis include, e.g., cytoplasmic condensation, increased membrane blebbing, and cellular shrinkage.

The presence of apoptotic cells can be measured in both the attached and "floating" compartments of the cultures. For example, both compartments can be collected by removing the supernatant, trypsinizing the attached cells, combining the preparations following a
10 centrifugation wash step (e.g., 10 minutes at 2000 rpm), and detecting apoptosis (e.g., by measuring DNA fragmentation). (See, e.g., Piazza *et al.*, 1995, Cancer Research 55:3110-16).

In vivo, the effect of a therapeutic composition of the anti-LY75 antibody of the invention can be evaluated in a suitable animal model. For example, xenogenic cancer models can be used, wherein cancer explants or passaged xenograft tissues are introduced into immune
15 compromised animals, such as nude or SCID mice (Klein *et al.*, 1997, Nature Medicine 3: 402-408). Efficacy can be measured using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like.

The therapeutic compositions used in the practice of the foregoing methods can be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery
20 method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is generally non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, Remington's Pharmaceutical Sciences 16th
25 Edition, A. Osal., Ed., 1980).

Methods for producing antibodies

The antibodies disclosed herein may be made by any suitable method. These methods include culturing a host cell containing isolated nucleic acid(s) encoding the antibodies. As will
30 be appreciated by those in the art, this can be done in a variety of ways, depending on the nature of the antibody. In the case where the antibodies are full length traditional antibodies, for example, a heavy chain variable region and a light chain variable region under conditions such that an antibody is produced and can be isolated.

The variable heavy and light chains of LY75_A1 are disclosed herein (both protein and
35 nucleic acid sequences); as will be appreciated in the art, these can be easily augmented to produce full length heavy and light chains. That is, having provided the DNA fragments encoding V_H and V_K segments as outlined herein, these DNA fragments can be further manipulated by

standard recombinant DNA techniques, for example, to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes, or to a scFv gene. In these manipulations, a V_K - or V_H -encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked", as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

The isolated DNA encoding the V_H region can be converted to a full-length heavy chain gene by operatively linking the V_H -encoding DNA to another DNA molecule encoding heavy chain constant regions (C_{H1} , C_{H2} and C_{H3}). The sequences of murine heavy chain constant region genes are known in the art [see e.g. Kabat, E. A., *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, US Department of Health and Human Services, NIH Publication No. 91-3242] and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, but most preferably is an IgG1 or IgG4 constant region. For a Fab fragment heavy chain gene, the V_H -encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain C_{H1} constant region.

The isolated DNA encoding the V_L / V_K region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the V_L -encoding DNA to another DNA molecule encoding the light chain constant region, C_L . The sequences of murine light chain constant region genes are known in the art [see, e.g. Kabat, E. A., *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, US Department of Health and Human Services, NIH Publication No. 91-3242] and DNA fragments encompassing these regions can be obtained by standard PCR amplification. In preferred embodiments, the light chain constant region can be a kappa or lambda constant region.

To create a scFv gene, the V_H - and V_L / V_K -encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g. encoding the amino acid sequence $(Gly_4-Ser)_3$, such that the V_H and V_L / V_K sequences can be expressed as a contiguous single-chain protein, with the V_L / V_K and V_H regions joined by the flexible linker [see e.g. Bird *et al.* (1988) *Science* 242:423-426; Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; McCafferty *et al.*, (1990) *Nature* 348:552-554].

Nucleic acids are provided which encode the antibodies disclosed herein. Such polynucleotides encode for both the variable and constant regions of each of the heavy and light chains, although other combinations are also contemplated in accordance with the compositions described herein.

The polynucleotides can be in the form of RNA or DNA. Polynucleotides in the form of DNA, cDNA, genomic DNA, nucleic acid analogs, and synthetic DNA are also usable. The DNA may be double-stranded or single-stranded, and if single stranded, may be the coding (sense)

strand or non-coding (anti-sense) strand. The coding sequence that encodes the polypeptide may be identical to the coding sequence provided herein or may be a different coding sequence, which sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same polypeptides as the DNA provided herein.

5 In some embodiments, nucleic acid(s) encoding the antibodies disclosed herein are incorporated into expression vectors, which can be extrachromosomal or designed to integrate into the genome of the host cell into which it is introduced. Expression vectors can contain any number of appropriate regulatory sequences (including, but not limited to, transcriptional and translational control sequences, promoters, ribosomal binding sites, enhancers, origins of
10 replication, etc.) or other components (selection genes, etc.), all of which are operably linked as is well known in the art. In some cases two nucleic acids are used and each put into a different expression vector (e.g. heavy chain in a first expression vector, light chain in a second expression vector), or alternatively they can be put in the same expression vector. It will be appreciated by those skilled in the art that the design of the expression vector(s), including the
15 selection of regulatory sequences may depend on such factors as the choice of the host cell, the level of expression of protein desired, etc.

In general, the nucleic acids and/or expression can be introduced into a suitable host cell to create a recombinant host cell using any method appropriate to the host cell selected (e.g., transformation, transfection, electroporation, infection), such that the nucleic acid molecule(s) are
20 operably linked to one or more expression control elements (e.g., in a vector, in a construct created by processes in the cell, integrated into the host cell genome). The resulting recombinant host cell can be maintained under conditions suitable for expression (e.g. in the presence of an inducer, in a suitable non-human animal, in suitable culture media supplemented with appropriate salts, growth factors, antibiotics, nutritional supplements, etc.), whereby the
25 encoded polypeptide(s) are produced. In some cases, the heavy chains are produced in one cell and the light chain in another.


Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), Manassas, VA including but not limited to Chinese hamster ovary (CHO) cells, HEK 293 cells,
30 NSO cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. Non-mammalian cells including but not limited to bacterial, yeast, insect, and plants can also be used to express recombinant antibodies. In some embodiments, the antibodies can be produced in transgenic animals such as cows or chickens.

35 General methods for antibody molecular biology, expression, purification, and screening are well known, for example, see US Patent Nos. 4,816,567, 4,816,397, 6,331,415 and 7,923,221, as well as Antibody Engineering, edited by Kontermann & Dubel, Springer,

Heidelberg, 2001 and 2010 Hayhurst & Georgiou, 2001, Curr Opin Chem Biol 5:683-689; Maynard & Georgiou, 2000, Annu Rev Biomed Eng 2:339-76; and Morrison, S. (1985) Science 229:1202.

In a further embodiment of the invention, component (B) of the pharmaceutical combination is ibrutinib or a pharmaceutically-acceptable salt thereof. Ibrutinib (Imbruvica®) is a small molecule drug that binds permanently to Bruton's tyrosine kinase (BTK). It has previously been used to treat B-cell cancers such as mantle cell lymphoma, chronic lymphocytic leukemia and Waldenström's macroglobulinemia (a form of non-Hodgkin's lymphoma). The structural and chemical formulae of ibrutinib are given below:

(R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one.



Ibrutinib is sold under the tradename Imbruvica®. It is available from Pharmacyclics, Inc. (US).

Pharmaceutical Compositions

The pharmaceutical combination of the invention is in the form of a combined preparation for simultaneous, separate or sequential use. Similarly, in the methods of the invention, components (A) and (B) of the pharmaceutical combination may be administered to a patient simultaneously, separately or sequentially.

The term "combined preparation" includes both fixed combinations and non-fixed combinations.

The term "fixed combination" means that the active ingredients (e.g. components (A) and (B)) are in the form of a single entity or dosage. In other words, the active ingredients are present in a single composition or formulation.

The term "non-fixed combination" means that the active ingredients (e.g. components (A) and (B)) are present in different entities or dosages (e.g. as separate compositions or formulations), for example as a kit of parts. The independent components (A) and (B) (in their desired compositions or formulations) can then be administered separately or sequentially, at the same time point or at different time points.

Where the administration is sequential, the delay in administering the second component should not be such as to lose the benefit of the effect arising from use of the combination.

Therefore, in one embodiment sequential treatment involves administration of each component of the combination within a period of 11 days. In another embodiment this period is 10 days. In another embodiment this period is 9 days. In another embodiment this period is 8 days. In another embodiment this period is 7 days. In another embodiment this period is within 6 days. In another embodiment this period is within 5 days. In another embodiment this period is within 4 days. In another embodiment this period is within 3 days. In another embodiment this period is within 2 days. In another embodiment this period is within 24 hours. In another embodiment this period is within 12 hours.

Components (A) and (B) may be administered in any order, e.g. component (A) first and then component (B); or component (B) first and then component (A).

The ratio of the total amounts of component (A) to component (B) to be administered in the combined preparation can be varied, e.g. in order to cope with the needs of a patient sub-population to be treated or the needs of the single patient which different needs can be due to age, sex, body weight, etc. of the patients.

Components (A) and (B), whether present in a single composition or in separate compositions, may independently be formulated with one or more pharmaceutically-acceptable carriers. The pharmaceutical combinations of the invention may also include at least one other anti-tumor agent, or an anti-inflammatory or immunosuppressant agent. Examples of therapeutic agents that can be used in combination therapy are described in greater detail below in the section on uses of the antibodies disclosed herein.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g. by injection or infusion). Depending on the route of administration, the active compound, i.e. antibody, immunoconjugate, or bispecific molecule, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

Components (A) and/or (B) may be in the form of one or more pharmaceutically acceptable salts. A "pharmaceutically acceptable salt" refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects [see, e.g. Berge, S.M., *et al.* (1977) *J. Pharm. Sci.* 66:1-19]. Examples of such salts include acid addition salts and base addition salts. Acid addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous and the like, as well as from nontoxic organic acids such as aliphatic mono- and dicarboxylic

acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-dibenzylethylenediamine, N-methylglucamine, chloroprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

A pharmaceutical combination of the invention or part thereof also may include a pharmaceutically acceptable anti-oxidant. Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Examples of suitable aqueous and non-aqueous carriers that may be employed in the pharmaceutical combinations of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These combinations or parts thereof may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the invention is contemplated. Supplementary active compounds can also be incorporated into the compositions.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures

thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged
5 absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared
10 by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the subject being treated, and the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the composition which produces a therapeutic effect. Generally, out of 100 per cent, this amount will range from about 0.01 per
20 cent to about 99 per cent of active ingredient, preferably from about 0.1 per cent to about 70 per cent, most preferably from about 1 per cent to about 30 per cent of active ingredient in combination with a pharmaceutically acceptable carrier.

Dosage regimens are adjusted to provide the optimum desired response (e.g. a synergistic combination, a therapeutic response). For example, a single bolus may be administered, several
25 divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a
30 predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in
35 individuals.

For administration of the anti-LY75 or anti-CD20 antibody, the dosage ranges from about 0.0001 to 100 mg/kg, for example, 0.001 to 50mg/kg, 0.005 to 20mg/kg, 0.01 to 10mg/kg and

more usually 0.01 to 5 mg/kg, of the host body weight. For example dosages can be 0.05 mg/kg body weight, 0.1 mg/kg body weight, 0.3 mg/kg body weight, 0.3 mg/kg body weight, 0.5 mg/kg body weight, 1 mg/kg body weight, 2 mg/kg body weight, 3 mg/kg body weight, 4 mg/kg body weight, 5 mg/kg body weight 6 mg/kg body weight, 7 mg/kg body weight, 8 mg/kg body weight, 9 mg/kg body weight, 10 mg/kg body weight, 12 mg/kg body weight, 15 mg/kg body weight, 20 mg/kg body weight, 25 mg/kg body weight, 30 mg/kg body weight, or within the range of 0.1-20 mg/kg, 0.5-15 mg/kg, 1-10 mg/kg, 2-8 mg/kg, 3-7 mg/kg, 4-6 mg/kg. An exemplary treatment regime entails administration once per day, once every 2 days, once per week, once every two weeks, once every three weeks, once every four weeks, once a month, once every 6 weeks, once every 3 months or once every three to 6 months. Preferred dosage regimens for an anti-LY75 antibody of the invention include 1 mg/kg body weight or 3 mg/kg body weight via intravenous administration, with the antibody being given using one of the following dosing schedules: (i) every four weeks for six dosages, then every three months; (ii) every three weeks; (iii) 3 mg/kg body weight once followed by 1 mg/kg body weight every three weeks.

In some embodiments, the anti-LY75 antibody (e.g. LY75-DM4) dosage is adjusted to achieve a plasma antibody concentration of 0.01 to 1.5 nM or 0.018 to 1.2 nM (e.g. about 0.018, 0.037, 0.075, 0.15, 0.3, 0.6 or 1.2 nM). Preferably, the anti-LY75 antibody is adjusted to achieve a plasma antibody concentration of 0.03 nM to 0.30 nM.

In some embodiments, the anti-CD20 antibody is formulated in a 10mg/ml solution. In some embodiments, the anti-CD20 antibody is administered i.v. at a dose of 350-400mg/m², preferably once weekly. In some embodiments, the anti-CD20 antibody (e.g. Rituximab) dosage is adjusted to achieve a plasma antibody concentration of about 2.34 to 150 nM (e.g. about 2.34, 4.68, 9.37, 18.75, 37.5, 75 or 150 nM). The anti-CD20 antibody (e.g. Rituximab) may be administered at a dose of about 1400 mg/23,400 units subcutaneously. In some embodiments, the dose is 200 mg, 400 mg, 600 mg, 800 mg, 1000 mg or 1200 mg. The anti-CD20 antibody (e.g. Rituximab) may be administered with one or more of cyclophosphamide, hydroxydaunorubicin, oncovin and predisone or prednisolone (i.e. CHOP therapy).

In some embodiments, the Ibrutinib dosage is adjusted to achieve a plasma concentration of about 15.6 to 1000 nM (e.g. about 15.6, 31.2, 62.5, 125, 250, 500 or 1000 nM). In other embodiments, the Ibrutinib dosage is adjusted to achieve a plasma concentration of about 1.56 to 100 nM (e.g. about 1.5, 3.1, 6.2, 12.5, 25, 50 or 100 nM). The Ibrutinib dose may be about 560 mg (e.g. four 140 mg capsules). It may be orally administered. In some embodiments, the Ibrutinib dose is about 100 mg, 200 mg, 300 mg, 400 mg or 500 mg.

Preferably, the combination of components (A) and (B) is a synergistic combination. The skilled person will understand that a synergistic combination is one wherein the effect of the combination is greater than the sum of the effects of the individual components. Synergism may be quantified using the Chou-Talalay combination index (CI) (see

“Evaluation of combination chemotherapy: integration of nonlinear regression, curve shift, isobologram, and combination index analyses”, Zhao L, *et al.* Clin Cancer Res. (2004) Dec 1;10(23):7994-8004; and “Computerized quantitation of synergism and antagonism of taxol, topotecan, and cisplatin against human teratocarcinoma cell growth: a rational approach to clinical protocol design”, Chou TC, Motzer RJ, Tong Y, Bosl GJ., J. Natl. Cancer Inst. (1994) Oct 19;86(20):1517-24). This combination index (CI) method is based on the multiple drug effect equation derived from the median-effect principle of the mass-action law. This provides a quantitative definition for strong synergism ($CI < 0.3$), synergism ($CI = 0.3-0.9$), additive effect ($CI = 0.9-1.1$) or antagonism/no benefit ($CI > 1.1$), and it provides the algorithm for computer software for automated simulation for drug combinations. It takes into account both the potency (the $D(m)$ value) and the shape of the dose-effect curve (the m value) of each drug alone and their combination. The Chou-Talalay combination index (CI) may be estimated using the Synergy R package (see “Preclinical versus Clinical Drugs Combination Studies”, Chou TC. Leuk. Lymphoma. (2008);49(11):2059-2080, and references therein, all of which are specifically incorporated herein by reference). The CI of the combination may be tested in a suitable cell-line, e.g. in ABC-DLBLC cell lines (such as TMD8 or HBL1), e.g. under the conditions used in Example 26.

Preferably, the pharmaceutical combination of the invention is a synergistic combination wherein the Chou-Talalay combination index (CI) is less than 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3 or 0.2. Preferably, the CI is 0.1-0.5, 0.1-0.3 or 0.1-0.2.

In particular, there is provided a method of treating cancer in a patient comprising simultaneously, sequentially or separately administering to a patient in need thereof therapeutically-effective synergistic amounts of components (A) and (B) of a pharmaceutical combination of the invention. Also provided is a pharmaceutical combination of the invention for use in the treatment of cancer, wherein synergistic amounts of components (A) and (B) are simultaneously, separately or sequentially administered to the patient for the treatment of the cancer. Preferably, amounts of components (A) and (B) are administered to the patient in order to provide the plasma concentrations disclosed above.

Also provided is the use of synergistic amounts of components (A) and (B) of the pharmaceutical combination of the invention in the manufacture of a pharmaceutical combination for simultaneous, separate or sequential use for the treatment of cancer. Also provided is a synergistic pharmaceutical combination of the invention for use in therapy or for use as a medicament.

In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. Antibody is usually administered on multiple occasions. Intervals

between single dosages can be, for example, daily, twice weekly, weekly, monthly, every three months, every six months, or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to the target antigen in the patient. In some methods, dosage is adjusted to achieve a plasma antibody concentration of about 1-1000 µg /ml, 5-750 µg /ml, 10-600 µg /ml, 15-500 µg /ml, 20-400 µg /ml and in some methods about 25-300 µg /ml.

Alternatively, the anti-LY75 and or anti-CD20 antibodies can be administered as sustained release formulations, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half-life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

Actual dosage levels of the active ingredients in the pharmaceutical combinations of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A "therapeutically effective dosage" of an anti-LY75 antibody or anti-CD20 antibody preferably results in a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, or a prevention of impairment or disability due to the disease affliction. For example, for the treatment of the LY75 or CD20 mediated tumors, a "therapeutically effective dosage" preferably inhibits cell growth or tumor growth by at least about 20%, at least about 30%, more preferably by at least about 40%, at least about 50% even more preferably by at least about 60%, at least about 70% and still more preferably by at least about 80% or at least about 90%, relative to untreated subjects. The ability of a compound to inhibit tumor growth can be evaluated in an animal model system predictive of efficacy in human tumors. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit cell growth, such inhibition can be measured *in vitro* by assays known to

the skilled practitioner. A therapeutically effective amount of a therapeutic compound can decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

A pharmaceutical combination of the present invention can be administered via one or more routes of administration using one or more of a variety of methods known in the art. Components (A) and (B) may be administered by the same route or by different routes. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Preferred routes of administration for antibodies of the invention include intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other parenteral routes of administration, for example by injection or infusion. The phrase "parenteral administration" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

Alternatively, the anti-LY75 or anti-CD20 antibody can be administered via a non-parenteral route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically.

The active compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art [see, e.g. *Sustained and Controlled Release Drug Delivery Systems* (1978) J.R. Robinson, ed., Marcel Dekker, Inc., N.Y].

Therapeutic compositions can be administered with medical devices known in the art. For example, in a preferred embodiment, component (A) and/or (B) can be administered with a needleless hypodermic injection device, such as the devices disclosed in US Patent Nos. 5,399,163; 5,383,851; 5,312,335; 5,064,413; 4,941,880; 4,790,824; or 4,596,556. Examples of well-known implants and modules useful in the present invention include: US Patent No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; US Patent No. 4,486,194, which discloses a therapeutic device for administering medicaments through the skin; US Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; US Patent No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; US Patent

No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and US Patent No. 4,475,196, which discloses an osmotic drug delivery system. These patents are incorporated herein by reference. Many other such implants, delivery systems, and modules are known to those skilled in the art.

5 In certain embodiments, the anti-LY75 and/or anti-CD20 antibodies can be formulated to ensure proper distribution *in vivo*. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g. US Patents 4,522,811; 5,374,548; and 5,399,331. The liposomes may
10 comprise one or more moieties which are selectively transported into specific cells or organs, thus enhance targeted drug delivery [see, e.g. V.V. Ranade (1989) *J. Clin. Pharmacol.* 29:685]. Exemplary targeting moieties include folate or biotin (see, e.g. US Patent 5,416,016.); mannosides [Umezawa *et al.* (1988) *Biochem. Biophys. Res. Commun.* 153:1038]; antibodies [P.G. Bloeman *et al.* (1995) *FEBS Lett.* 357:140; M. Owais *et al.* (1995) *Antimicrob. Agents
15 Chemother.* 39:180]; surfactant protein A receptor [Briscoe *et al.* (1995) *Am. J. Physiol.* 1233:134]; p120 [Schreier *et al.* (1994) *J. Biol. Chem.* 269:9090]; see also K. Keinänen; M.L. Laukkanen (1994) *FEBS Lett.* 346:123; J.J. Killian; I.J. Fidler (1994) *Immunomethods* 4:273.

Uses and Methods

20 As used herein, the term "subject" is intended to include human and non-human animals. Non-human animals include all vertebrates, e.g. mammals and non-mammals, such as non-human primates, sheep, dogs, cats, cows, horses, chickens, amphibians, and reptiles. Preferred subjects include human patients having disorders mediated by LY75 activity and/or CD20 activity.

25 The methods are particularly suitable for treating human patients having a disorder associated with the aberrant LY75 expression and/or CD20 expression. Given the expression of LY75 on tumor cells, the combinations and methods of the present invention can be used to treat a subject with a tumorigenic disorder, e.g. a disorder characterized by the presence of tumor cells expressing LY75 or in the manufacture of a medicament for the treatment of such a disorder
30 including, for example leukaemia, including chronic lymphocytic leukaemia and acute myeloid leukaemia, non-Hodgkin's lymphoma, including DLBCL, B-Cell Lymphoma, Follicular Lymphoma, Mantle Cell Lymphoma, Lymphoma of Mucosa-Associated Lymphoid Tissue (MALT), T-Cell/Histiocyte-Rich B-Cell Lymphoma, Burkitt's Lymphoma, Lymphoplasmacytic Lymphoma, Small Lymphocytic Lymphoma, Marginal Zone Lymphoma, T Cell Lymphoma, Peripheral T-Cell
35 Lymphoma, Anaplastic Large Cell Lymphoma and AngioImmunoblastic T-Cell Lymphoma. LY75 has been demonstrated to be internalised on antibody binding as illustrated in Examples 5 and 7

below, thus enabling the anti-LY75 antibodies to be used in any payload mechanism of action e.g. an ADC approach, radioimmunoconjugate, or ADEPT approach.

The anti-LY75 antibodies, generally administered as ADCs, can be used to inhibit or block LY75 function which, in turn, can be linked to the prevention or amelioration of certain disease symptoms, thereby implicating the LY75 as a mediator of the disease. This can be achieved by contacting a sample and a control sample with the anti-LY75 antibody under conditions that allow for the formation of a complex between the antibody and LY75. Any complexes formed between the antibody and the LY75 are detected and compared in the sample and the control.

Suitable routes of administering the antibody compositions (e.g. monoclonal antibodies, and immunoconjugates) *in vivo* and *in vitro* are well known in the art and can be selected by those of ordinary skill. For example, the antibody compositions can be administered by injection (e.g. intravenous or subcutaneous). Suitable dosages of the molecules used will depend on the age and weight of the subject and the concentration and/or formulation of the antibody composition.

As previously described, the anti-LY75 and/or anti-CD20 antibodies can be co-administered with one or other more therapeutic agents, e.g. a cytotoxic agent, a radiotoxic agent or an immunosuppressive agent. The antibody can be linked to the agent (as an immunocomplex) or can be administered separate from the agent. In the latter case (separate administration), the antibody can be administered before, after or concurrently with the agent or can be co-administered with other known therapies, e.g. an anti-cancer therapy, e.g. radiation. Such therapeutic agents include, among others, anti-neoplastic agents such as doxorubicin (adriamycin), cisplatin bleomycin sulfate, carmustine, chlorambucil, and cyclophosphamide hydroxyurea which, by themselves, are only effective at levels which are toxic or subtoxic to a patient. Cisplatin is intravenously administered as a 100 mg/kg dose once every four weeks and adriamycin is intravenously administered as a 60-75 mg/ml dose once every 21 days. Other agents suitable for co-administration with the antibodies of the invention include other agents used for the treatment of cancers, e.g. gastric cancer, colorectal cancer, prostate cancer, breast cancer, ovarian cancer or lung cancer, such as Avastin[®], 5FU and gemcitabine. Co-administration of the anti-LY75 antibodies or antigen binding fragments thereof, of the present invention with chemotherapeutic agents provides two anti-cancer agents which operate via different mechanisms which yield a cytotoxic effect to human tumor cells. Such co-administration can solve problems due to development of resistance to drugs or a change in the antigenicity of the tumor cells which would render them unreactive with the antibody.

The pharmaceutical combinations of the invention can also be administered together with serum and/or complement. These compositions can be advantageous when the complement is located in close proximity to the antibodies. Alternatively, the antibodies, and the complement or serum can be administered separately.

2018232886 24 May 2024

Also within the scope of the present invention are kits comprising components (A) and (B), together with instructions for use. The kit can further contain one or more additional reagents, such as an immunosuppressive reagent, a cytotoxic agent or a radiotoxic agent, or one or more additional antibodies (e.g. an antibody having a complementary activity which binds to an epitope in the LY75 antigen distinct from the first antibody).

Accordingly, patients treated with pharmaceutical combinations of the invention can be additionally administered (prior to, simultaneously with, or following administration of an antibody disclosed herein) with another therapeutic agent, such as a cytotoxic or radiotoxic agent, which enhances or augments the therapeutic effect of the antibodies.

In other embodiments, the subject can be additionally treated with an agent that modulates, e.g. enhances or inhibits, the expression or activity of Fcγ or Fcγ receptors by, for example, treating the subject with a cytokine. Preferred cytokines for administration during treatment with the multispecific molecule include of granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor (GM-CSF), interferon-γ (IFN-γ), and tumor necrosis factor (TNF).

All references cited in this specification, including without limitation all papers, publications, patents, patent applications, presentations, texts, reports, manuscripts, brochures, books, internet postings, journal articles, periodicals, product fact sheets, and the like, one hereby incorporated by reference into this specification in their entireties. The discussion of the references herein is intended to merely summarize the assertions made by their authors and no admission is made that any reference constitutes prior art and Applicants' reserve the right to challenge the accuracy and pertinence of the cited references. However, it is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the dependent claims.

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

The present invention is further illustrated by the following examples which should not be construed as further limiting.

Example 1: Generation of Human Monoclonal Antibodies Against LY75-Antigen

Following standard procedures, mice (xenomouse IgG1) were immunized with CHO cells transfected with full length LY75.

The specificity of antibodies raised against the LY75 was tested by flow cytometry on HEK293 cells transfected with LY75 and subsequently on LY75-expressing HT29 cells. To test the ability of the antibodies to bind to the cell surface LY75 protein, the antibodies were incubated with the LY75-expressing cells. Cells were washed in FACS buffer (DPBS, 2% FBS),

2018232886 24 May 2024

centrifuged and resuspended in 100µl of the diluted primary LY75 antibody (also diluted in FACS buffer). The antibody-cell line complex was incubated on ice for 60 min and then washed twice with FACS buffer as described above. The cell-antibody pellet was resuspended in 100µl of the diluted secondary antibody (also diluted in FACS buffer) and incubated on ice for 60 min on ice.

- 5 The pellet was washed as before and resuspended in 200µl FACS buffer. The samples were loaded onto the BD FACScanto II flow cytometer and the data analyzed using the BD FACSDiva software (results not shown).

Example 2: Structural Characterization of Monoclonal Antibodies to LY75

- 10 The cDNA sequences encoding the heavy and light chain variable regions of the LY75_A1 monoclonal antibody were obtained using standard PCR techniques and were sequenced using standard DNA sequencing techniques.

The antibody sequences may be mutagenized to revert back to germline residues at one or more residues.

- 15 The nucleotide and amino acid sequences of the heavy chain variable region of LY75_A1 are shown in SEQ ID NO: 3 and 1, respectively.

The nucleotide and amino acid sequences of the light chain variable region of LY75_A1 are shown in SEQ ID NO: 4 and 2, respectively.

- 20 Comparison of the LY75_A1 heavy chain immunoglobulin sequence to the known human germline immunoglobulin heavy chain sequences demonstrated that the LY75_A1 heavy chain utilizes a V_H segment from human germline V_H 3-15 and a J_H segment from human germline J_H JH4. Further analysis of the LY75_A1 V_H sequence using the Kabat system of CDR region determination led to the delineation of the heavy chain CDR1, CDR2 and CDR3 regions as shown in SEQ ID NOs: 5, 6 and 7, respectively. The alignments of the LY75_A1 CDR1, CDR2
25 and CDR3 V_H sequences to the germline V_H 3-15 and germline J_H JH4 sequence are shown in Figure 1.

- Comparison of the LY75_A1 light chain immunoglobulin sequence to the known human germline immunoglobulin light chain sequences demonstrated that the LY75_A1 light chain utilizes a V_K segment from human germline V_K O12 and a J_K segment from human germline J_K
30 JK4. Further analysis of the LY75_A1 V_K sequence using the Kabat system of CDR region determination led to the delineation of the light chain CDR1, CDR2 and CDR3 regions as shown in SEQ ID NOs: 8, 9 and 10, respectively. The alignments of the LY75_A1 CDR1, CDR2 and CDR3 V_K sequences to the germline V_K O12 and germline J_K JK4 sequences are shown in Figure 2.

Example 3: Immunohistochemistry Using Monoclonal Antibody to LY75

Using the human monoclonal antibodies specific to LY75, immunohistochemistry was performed on FFPE HT-29 and A549 cell pellets, FFPE non-Hodgkin's lymphoma and pancreatic cancer arrays, and fresh frozen lymphoma/leukaemia tumours, ovarian cancer, pancreatic cancer, and breast cancer sections and a normal tissue array.

*Materials and methods**Materials*

Xylenes (X5P-1gal) from Fisher Scientific, PA, USA.

10 Histoprep 100% ethanol (HC-800-1GAL) from Fisher Scientific, PA, USA.

10x Citrate buffer for heat induced epitope retrieval (AP9003125) from Thermo Scientific, MA, USA.

Thermo Scientific* Pierce* Peroxidase Suppressor (35000) from Thermo Scientific, MA, USA.

Serum free protein block (X0909) from Dako, CA, USA

15 Secondary antibody: goat anti-human IgG Fab-FITC conjugated (109-097-003) from Jackson ImmunoResearch, PA, USA

Chrome pure Human IgG, whole molecule (09-000-003) from Jackson ImmunoResearch, PA, USA

Tertiary antibody: mouse anti-FITC (ab10257) from Abcam, MA, USA

20 Purified human IgG isotype control (1-001A) from R&D Systems, MN, USA

Tween-20 (BP337-100) from Fisher Scientific, PA, USA

Acetone (BP2403-4) from Fisher Scientific, PA, USA

Dual Link EnVision+ HRP-conjugated polymer, Mouse and Rabbit (K4063) from Dako, CA, USA.

DAB 2-solution kit (882014) from Invitrogen, NY, USA.

25 Harris Hematoxylin (23-245-677) from Fisher Scientific, PA, USA.

Faramount mounting media (S302580) from Dako, CA, USA.

Tissue sections and arrays were purchased from US Biomax Inc., MD, USA or Origene, MD, USA.

30

Preparation of FFPE slides: Deparaffinisation and Rehydration

FFPE slides were deparaffinised in xylene (2 x 3 minutes) then rehydrated through 1:1 xylene: 100% ethanol (1 x 3 minutes), 100% ethanol (2 x 3 minutes), 95% ethanol (1 x 3 minutes), 70% ethanol (1 x 3 minutes), 50% ethanol (1 x 3 minutes), and tap water (1 x 3 minutes).

35

Preparation of FFPE slides: Antigen Retrieval (Microwave).

The LY75 antigen was retrieved using microwave heat, high power until boiling then low power for 10 minutes in 50 mL 1x citrate buffer in a Coplin jar. Slides were then left to cool to room temperature for a further 15 min, then washed in tap water, 3 minutes. Circles were drawn around each tissue section/TMA with a hydrophobic barrier pen and slides were then washed 3 times in PBS, 3 minutes each wash.

Preparation of FF slides

Slides were removed from storage at -80°C and allowed to dry at room temperature in the fume hood for 20-30 minutes. The slides were fixed for 10 min in ice cold acetone at -20°C, then allowed to dry for 20 min in the fume hood at room temperature. Slides were washed and rehydrated in PBS, 3 washes for 3 min each. Sections were outlined with a hydrophobic barrier pen.

Preparation of antibody complexes

The primary anti-LY75 antibody was diluted in serum free protein block (SFPB) to obtain a solution with a concentration 20-fold greater than the final desired concentration (20 µg/mL for 1 µg/mL final). The secondary antibody, goat anti-human immunoglobulin G (IgG) antigen-binding fragment (Fab), was prepared similarly in SFPB to create a solution of equal concentration.

Equal volumes of primary and secondary antibodies were combined in a labelled tube, gently mixed, and incubated for 3 minutes at room temperature, resulting in a primary antibody concentration 10-fold greater than the desired final concentration (10 µg/mL for 1 µg/mL final). This mixture was diluted 1:5 with SFPB, gently mixed, and incubated for 30 minutes at room temperature, resulting in a primary antibody concentration twice that of the desired final concentration (2 µg/mL for 1 µg/mL final).

To produce the final staining complexes, a 1% (10 µg/µL) solution of human IgG was prepared in SFPB and equal volume added to the primary/secondary antibody mixture. This combination was gently mixed and incubated at room temperature for 30 minutes, diluting by half the primary antibody concentration of the primary/secondary antibody mixture and resulting in the desired final primary antibody concentration (1 µg/mL).

Immunostaining

Meanwhile, endogenous tissue peroxidase activity was blocked by incubating tissues with peroxidase suppressor for 5-10 minutes at RT in a humidified chamber. Slides were then washed in PBS 3 x 3 minutes each wash. Tissues were incubated in SFPB for 30 minutes at room temperature in a humidified chamber. Final staining complexes were applied to each tissue section and/or microarray, and the slides were incubated for 30 min at room temperature in a humidified chamber. Slides were then washed once in PBS and once in PBST (PBS+0.125% Tween-20), 3 minutes each wash. The tertiary antibody mouse anti-FITC, was

applied at 2 µg/mL concentration for 30 min, room temperature, in a humidified chamber. Sections were then washed once in PBS and once in PBST, 3 min each wash. Dual Link EnVision+ anti-mouse/rabbit-HRP-conjugated polymer was then applied to the tissues and the slides were incubated for 30 min at room temperature in a humidified chamber. Slides were then washed once in PBS, once in PBST, 3 minutes each wash. Tissues were incubated in DAB solution prepared according to the manufacturer's instructions at room temperature for 10 min. Slides were then washed once in running tap water for 2 minutes and once in PBS for 3 minutes. The slides were counterstained with Hematoxylin for 30 seconds at room temperature, and washed with running tap water. The slides were dried at room temperature for 30 minutes and coverslips were then mounted onto the slides using Faramount mounting media.

Results

LY75_A1 showed positivity in FFPE Triple Negative breast cancer samples, where 77% of the sections showed positive staining and 55% exhibited robust (+++) staining.

Staining for LY75 in FF normal tissues was generally absent to low. Ductal epithelium of the breast, salivary gland, and pancreas exhibited marked low to moderate staining, and the spleen stained low positive. Therefore antibodies directed to LY75 may have utility as therapeutics and diagnostics in some of the tested cancers and possibly other cancer types showing expression of LY75.

Example 4: Efficacy of DM1-Conjugated Anti-LY75 Monoclonal Antibodies in HT-29 Cells

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5e3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10e3 cells/well). The pellet was resuspended in culture media to a concentration of 1e5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

The results depicted in Figure 3a show a subpopulation of antibodies, know to bind to LY75, which can induce cell kill of HT-29 cells. This suggests while antibodies can bind to LY75 only a few display efficacy when conjugated to DM1. Antibodies where then chosen from the subpopulation for further cytotoxic activity analysis.

Example 5: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Colorectal Cancer Cells

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3b shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards HT-29 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration and other anti-LY75 antibodies conjugated to a toxin (selected from Example 1).

Example 6: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Lymphoma Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

50 μ l/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50 μ l/well) were added to the appropriate wells. Excess media (200 μ l/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37°C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100 μ l/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100 μ l/well PBS and 100 μ l Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3c shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards RAJI cells. Figure 3d shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards Namalwa cells. Figure 3e shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards Karpas 299 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration and other anti-LY75 antibodies conjugated to DM1 and DM4 (selected from Example 1).

Example 7: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Pancreatic Cancer Cell Lines

Materials

- 5 Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

10 *Method*

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

- 15 $50 \mu\text{l}$ /well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) ($50 \mu\text{l}$ /well) were added to the appropriate wells. Excess media ($200 \mu\text{l}$ /well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was
20 incubated for 72h at 37°C.

- The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with $100 \mu\text{l}$ /well PBS (for suspension cells, plate is centrifuged first to pellet cells). $100 \mu\text{l}$ /well PBS and $100 \mu\text{l}$ Cell titer glo was added to each well and triturated to mix. The plate was incubated in
25 the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

- Figure 3f shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and
30 DM4 towards BxPC3 cells. Figure 3g shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards HupT4 cells. Figure 3h shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards HPAAFFII cells. These results demonstrates an increase in cytotoxic activity proportional to antibody concentration and other anti-LY75 antibodies conjugated to DM1 and DM4 (selected from Example 1).

Example 8: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Chronic Lymphocytic Leukaemia Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

5 PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

Cell Titer Glo (G7572) from Promega, WI, USA.

Method

10 Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed
15 plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

20 The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell
25 lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3i shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards EHEB cells. Figure 3j shows the cytotoxic activity of anti-LY75 antibodies
30 conjugated to DM1 and DM4 towards Mec-1 cells. These results demonstrates an increase in cytotoxic activity proportional to antibody concentration and other anti-LY75 antibodies conjugated to DM1 and DM4 (selected from Example 1).

Example 9: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Acute Monocytic Leukaemia Cell Lines

35

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

Cell Titer Glo (G7572) from Promega, WI, USA.

5 *Method*

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

10 50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was
15 incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in
20 the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3k shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and
25 DM4 towards AML-193 cells. These results demonstrates an increase in cytotoxic activity proportional to antibody concentration and other anti-LY75 antibodies conjugated to DM1 and DM4 (selected from Example 1).

Example 10: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal

30 Antibodies in Breast Cancer Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

35 Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5e3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10e3 cells/well). The pellet was resuspended in culture media to a concentration of 1e5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3l shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards HCC 70 (ER negative, PR negative and Her2 negative) cells. Figure 3m shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards HCC 1806 (ER negative, PR negative and Her2 negative) cells. Figure 3n shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards MDA-MB-468 cells. These results demonstrates an increase in cytotoxic activity proportional to antibody concentration.

Example 11: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Bladder Cancer Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5e3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells,

such as 10^3 cells/well). The pellet was resuspended in culture media to a concentration of 10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3o shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards RT4 cells. Figure 3p shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards 5637 cells. Figure 3q shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards SW780 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration.

Example 12: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Head and Neck Cancer Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.

PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.

RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.

Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10^3 cells/well). The pellet was resuspended in culture media to a concentration of 10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to

concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

5 The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell
10 lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3r shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards SCC-9 cells. These results demonstrate an increase in cytotoxic activity
15 proportional to antibody concentration.

Example 13: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Oesophageal Cancer Cell Lines

Materials

20 Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

25 Method

Cells were dissociated using cell stripper and counted. 5e3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10e3 cells/well). The pellet was resuspended in culture media to a concentration of 1e5 cells/mL.

30 50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was
35 incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x

with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3s shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards OE 19 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration.

Example 14: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Ovarian Cancer Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3t shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards OVCAR-3 cells. Figure 3u shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards SK-OV-3 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration.

Example 15: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Multiple Myeloma Cell Lines

Materials

Cell stripper (Non-enzymatic cell dissociation) (MT-25-056CI) from Fisher Scientific, PA, USA.
PBS pH 7.4 (1X) (SH30028LS) from Fisher Scientific, PA, USA.
RPMI 1640 Media (MT-10-041-CM) from Fisher Scientific, PA, USA.
Cell Titer Glo (G7572) from Promega, WI, USA.

Method

Cells were dissociated using cell stripper and counted. 5×10^3 cells/well were spun down into a pellet (for suspension cells, more can be used depending on the doubling time of the cells, such as 10×10^3 cells/well). The pellet was resuspended in culture media to a concentration of 1×10^5 cells/mL.

50ul/well cell suspension was added to wells of a 96-well white sided, clear bottomed plate. Antibodies were diluted and titrated to 8 points (3-fold titrations) corresponding to concentrations between 0-20 nM (twice the test concentrations). Diluted antibodies or media (for untreated samples) (50ul/well) were added to the appropriate wells. Excess media (200ul/well) was added to the outside rows and columns of the plate to prevent evaporation. The plate was incubated for 72h at 37C.

The plate was removed from the incubator and incubated at room temperature for 30 minutes. Meanwhile Cell Titer Glo solution was thawed. The plate was flicked and washed 1x with 100ul/well PBS (for suspension cells, plate is centrifuged first to pellet cells). 100ul/well PBS and 100ul Cell titer glo was added to each well and triturated to mix. The plate was incubated in the dark at room temperature for 15 minutes and visualized by microscopy to ensure efficient cell lysis occurred. The plate was then read on a Glomax luminometer.

Results

Figure 3v shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards MOLP-8 cells. Figure 3w shows the cytotoxic activity of anti-LY75 antibodies conjugated to DM1 and DM4 towards RPMI8226 cells. These results demonstrate an increase in cytotoxic activity proportional to antibody concentration.

Example 16: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Raji Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in subcutaneous Raji Burkitt's lymphoma SCID mouse xenograft model.

Immunodeficient SCID mice were inoculated subcutaneously with Raji (human Burkitt's lymphoma) tumour cells. Tumours were allowed to establish and mice were sorted into five treatment groups of 3-6 mice per group. When the mean tumour volume reached an average size of 129-132 mm³ per group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; phosphate buffered saline (PBS)); Group 2 (LY75_DM1; 10 mg/kg), Group 3 (Isotype control-DM1; 10 mg/kg), Group 4 (LY75_DM4; 5 mg/kg), Group 5 (isotype control-SPBDDM4; 5 mg/kg). A second dose was administered one week later. Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumours were measured twice weekly. Mice were euthanized when their tumours reached the tumour volume endpoint of 2000 mm³ or after 60 days, whichever came first. Efficacy was determined from tumour growth delay (TGD), the increase in median time-to-endpoint (TTE) and from logrank analysis of differences in Kaplan Meier survival curves in ADC-treated versus PBS-treat mice. The first five vehicle-treated control mice to reach endpoint were sampled for tumours that were processed by formalin fixation and paraffin embedded.

Results

Figure 4a shows LY75_DM1 and LY75_DM4 each demonstrated significant anti-tumour activity and significantly extended survival in the Raji Burkitt's lymphoma SCID mouse xenograft model compared to controls; however, the 5 mg/kg LY75_DM4 doses were significantly more effective than the 10 mg/kg doses of LY75_DM1, resulting in 5 of 6 mice with complete but transient tumour regression. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human non-Hodgkin lymphoma cancer patients.

Example 17: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Namalwa Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in subcutaneous Namalwa Burkitt's lymphoma SCID mouse xenograft model.

Immunodeficient SCID mice were inoculated subcutaneously with Namalwa (human Burkitt's lymphoma) tumour cells. Tumours were allowed to establish and mice were sorted into five treatment groups of 6 mice per group. When the mean tumour volume reached an average

size of 114 mm³ per group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; phosphate buffered saline (PBS)); Group 2 (LY75_DM1; 10 mg/kg), Group 3 (Isotype control-DM1; 10 mg/kg), Group 4 (LY75_DM4; 5 mg/kg), Group 5 (isotype control-SPBDDM4; 5 mg/kg). Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumours were measured twice weekly. Mice were euthanized when their tumours reached the tumour volume endpoint of 2000 mm³ or after 60 days, whichever came first. Efficacy was determined from tumour growth delay (TGD), the increase in median time-to-endpoint (TTE), and from log rank analysis of differences in Kaplan Meier survival curves in ADC-treated versus PBS-treated mice. The first five vehicle-treated control mice to reach endpoint were sampled for tumours that were processed by formalin fixation and paraffin embedded.

Results

Figure 4b shows LY75_DM1 and LY75_DM4 each demonstrated significant anti-tumour activity and survival extension in the Namalwa Burkitt's lymphoma SCID mouse xenograft model compared to controls; however, the 5 mg/kg LY75_DM4 dose was significantly more effective than the 10 mg/kg dose of LY75_DM1, causing a brief reduction in tumour volume. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human non-Hodgkin lymphoma cancer patients.

Example 18: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Pancreatic Cancer Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in the subcutaneous HPAFII pancreatic adenocarcinoma athymic nude mouse xenograft model.

Immunodeficient athymic nude mice were inoculated subcutaneously with HPAFII (human pancreatic adenocarcinoma) tumor cells. Tumors were allowed to establish and mice were sorted into five treatment groups of 6 mice per group. When the mean tumor volume reached an average size of ~114 mm³/group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; phosphate buffered saline (PBS)); Group 2 (LY75_DM1; 10 mg/kg), Group 3 (Isotype control-DM1; 10 mg/kg), Group 4 (LY75_DM4; 5 mg/kg), Group 5 (isotype control-SPBDDM4; 5 mg/kg). Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumors were measured thrice weekly. Mice were euthanized when their tumors reached the tumor volume endpoint of 2000 mm³ or after 90 days, whichever came first. Efficacy was determined from the effect of treatment on tumor volume and from log rank analysis of differences in Kaplan-Meier survival curves in ADC-treated or PBS-treated mice. The tumors

were sampled from vehicle-treated control mice and processed by formalin fixation and paraffin embedded.

Results

Figure 4c shows LY75_DM1 and LY75_DM4 displayed significant and similarly potent anti-tumor activity and survival extension in the HPAFII nude mouse xenograft model compared to controls. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human pancreatic cancer patients.

Example 19: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Bladder Cancer Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in the subcutaneous SW780 human bladder carcinoma SCID mouse xenograft model.

Immunodeficient athymic nude mice were inoculated subcutaneously with HPAFII (human pancreatic adenocarcinoma) tumor cells. Tumors were allowed to establish and mice were sorted into five treatment groups of 6 mice per group. When the mean tumor volume reached an average size of ~114 mm³/group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; phosphate buffered saline (PBS)); Group 2 (LY75_DM1; 1 mg/kg), Group 3 (LY75_DM1; 2.5 mg/kg), Group 4 (LY75_DM1; 5 mg/kg), Group 5 (LY75_DM4; 1 mg/kg), Group 6 (LY75_DM4; 2.5 mg/kg), Group 7 (LY75_DM4; 5 mg/kg), Group 8 (isotype control-SPBDDM4; 5 mg/kg). Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumors were measured thrice weekly. Mice were euthanized when their tumors reached the tumor volume endpoint of 2000 mm³ or after 90 days, whichever came first. Efficacy was determined from the effect of treatment on tumor volume and from log rank analysis of differences in Kaplan-Meier survival curves in ADC-treated or PBS-treated mice. The tumors were sampled from vehicle-treated control mice and processed by formalin fixation and paraffin embedded.

Results

Figure 4d shows LY75_DM1 and LY75_DM4 displayed significant and similarly potent anti-tumor activity and survival extension in the SW780 nude mouse xenograft model compared to controls. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human bladder cancer patients.

Example 20: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Breast Cancer Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in the subcutaneous MDA-MB-468 athymic nude mouse xenograft model.

Immunodeficient athymic nude mice were inoculated subcutaneously with MDA-MB-468 (human triple negative breast adenocarcinoma) tumour cells. Tumours were allowed to establish and mice were sorted into seven treatment groups of 10 mice per group. When the mean tumour volume reached an average size of 167 mm³ per group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; 20 mM sodium succinate, pH 5.0, 6% trehalose, 0.04% polysorbate); Group 2 (LY75_DM1; 5 mg/kg), Group 3 (LY75_DM1; 10 mg/kg), Group 4 (LY75_DM4; 5 mg/kg), Group 5 (LY75_DM4; 2.5 mg/kg), Group 6 (LY75_DM4; 1 mg/kg), Group 7 (Isotype control-DM4; 5 mg/kg). Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumours were measured twice weekly. Mice were euthanized 82 days after tumour inoculation. Efficacy was determined from anti-tumour activity (mean tumour size in treatment group/mean tumour size in control group x 100) and the increase in mean time-to-endpoint (TTE) in ADC-treated versus PBS-treated mice. The five largest tumours in vehicle-treated control mice on day 71 post inoculation were sampled processed by formalin fixation and paraffin embedded.

Results

Figure 4e shows LY75_DM1 and LY75_DM4 each demonstrated dramatic anti-tumour activity in the MDA-MB-468 nude mouse xenograft model compared to controls. Dose dependent activity was observed with LY75_DM4, where 2.5 and 5 mg/kg were much more potent than 1 mg/kg. At 5 mg/kg, LY75_DM1 and LY75_DM4 were similarly effective. Sustained regressions in mean tumour volume were observed for LY75_DM1 at 10 and 5 mg/kg and LY75_DM4 at 5 and 2.5 mg/kg. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human triple negative breast cancer patients.

Example 21: Efficacy of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Colorectal Cancer Xenograft Models

The efficacy of LY75_DM1 and LY75_DM4 were tested in the subcutaneous COLO205 colorectal adenocarcinoma athymic nude mouse xenograft model.

Immunodeficient athymic nude mice were inoculated subcutaneously with COLO205 (human colorectal adenocarcinoma) tumor cells. Tumors were allowed to establish and mice were sorted into five treatment groups of 6 mice per group. When the mean tumor volume

reached an average size of 117 mm³ per group, each group was treated with one of the following compounds, administered intravenously at the indicated dosages: Group 1 (Vehicle; phosphate buffered saline (PBS)); Group 2 LY75_DM1; 10 mg/kg), Group 3 (Isotype control-DM1; 10mg/kg), Group 4 (LY75_DM4; 5 mg/kg), Group 5 (Isotype control-DM4; 5 mg/kg). A second dose was administered twelve days after the first. Body weights (BW) were monitored, the mice were examined frequently for health and adverse side effects, and tumors were measured twice weekly. Mice were euthanized when their tumors reached the tumor volume endpoint of 1000 mm³ or after 60 days, whichever came first. Efficacy was determined from tumor growth delay (TGD), the increase in median time-to-endpoint (TTE) and from log rank analysis of differences in Kaplan Meier survival curves in ADC-treated versus PBS-treated mice. The first five vehicle-treated control mice to reach endpoint were sampled for tumors that were processed by formalin fixation and paraffin embedded.

Results

Figure 4f shows LY75_DM1 and LY75_DM4 exhibited similar modest anti-tumor activity and survival extension in the COLO205 colorectal adenocarcinoma nude mouse xenograft model compared to controls. All treatments were well-tolerated and no clinical signs of toxicity were observed. These data suggest the potential for ADCs directed towards LY75, for example LY75_DM1 and LY75_DM4, to provide clinical benefit in the treatment of human colorectal cancer patients.

Example 22: Toxicity of DM1-Conjugated and DM4-Conjugated Anti-LY75 Monoclonal Antibodies in Cynomolgus Monkeys

Six male monkeys were assigned to the study with 2 monkeys/group. Either vehicle (PBS), LY75_DM4 (cleavable) or LY75_DM1 (non-cleavable) was administered twice (on Day 1 and Day 29) by a 15-minute intravenous infusion at 0 mg/kg/dose (PBS, vehicle), 5 mg/kg/dose (LY75_DM4, cleavable) or 10 mg/kg/dose (LY75_DM1, non-cleavable). Blood samples were collected for toxicokinetic evaluations prior to dose initiation (Day 1), and 1, 2, 3, 7, 14, 21 and 28 days post each dose. Blood samples for clinical pathology analyses were collected prior to dose initiation (Day 1), and 1, 3, 7, 14, 21 and 28 days post each dose (28 days post the 1st dose was also served as the pre-dose time point for the 2nd dose). All study animals were euthanized and necropsied following the final blood collection on Day 57. The plasma separated from each blood draw was isolated, frozen and shipped to Oxford BioTherapeutics, Inc. to be analyzed for ADC concentration by ELISA.

Treatment-related clinical pathology findings included a mild regenerative anemia and transient decreases in the blood leukocyte profile most notably in neutrophils counts. Anemia was observed in both animals treated with 5 mg/kg LY75_DM4 and in one of the two animals

5 treated with 10 mg/kg LY75_DM1. Severe neutropenia with a nadir at one-week post dose and a rapid recovery in counts was observed in all animals; the nadir in absolute neutrophil count was lower in LY75_DM4 treated animals. There were no test article-related effects on the APTT and PT coagulation parameters. Serum chemistry changes included transient increases in AST, CK, LDH (in 1 of 2 animals in each treatment group) and globulin following administration of 5 mg/kg LY75_DM4 and 10 mg/kg LY75_DM1. In addition, a transient increase in the liver specific enzyme ALT was observed only in the LY75_DM4 treated animals. The short duration of and/or the magnitude of the increases in serum chemistry parameters suggest they were not adverse. There were no test-article related urinalysis findings. Upon examination at necropsy following a 4-week recovery period there were no treatment related gross pathology findings or changes in absolute and relative organ weights. Histopathology findings only in the thyroid gland (an alteration in the colloid morphology in follicles) and kidney (dilated tubules in the outer cortex), were graded as minimal severity; not associated with changes in other study parameters; and, not adverse and of minimal toxicological significance. Conclusion: Repeated dose treatment with two doses of 5 mg/kg LY75_DM4 or 10 mg/kg LY75_DM1 was well tolerated in cynomolgus monkeys. All treatment-related toxicity findings were reversible following a 4-week recovery period.

Example 23: Epitope characterisation of LY75_A1 by competitive Fluorescence Activated Cell Sorting (FACS) binding analysis

Method

COLO205 cells (ATCC, catalog # CCL-222) were detached from tissue culture flasks with Cell Stripper (Cellgro, catalog # MT-25-056CI). Cells were washed and resuspended in FACS buffer (PBS + 2%FBS), neutralized with growth media, and counted. Cells were plated at 50,000 cells per well in a V Bottom 96-well plate. Cells were washed once with FACS buffer (PBS (Fisher, catalog # SH30028-03) + 2% FBS). An anti-LY75-mAb (Selected from Example 1) or LY75_A1 was added to wells starting at 250 nM and diluted serially 3 fold and applied to the relevant wells for 45 minutes on ice. Test wells that required single or multiple staining steps were left in FACS buffer as appropriate to ensure the final staining was completed simultaneously for all conditions tested. Two wells were left unstained in FACS buffer as controls.

After the incubation with blocking antibody, cells were washed twice in FACS buffer. The cells were resuspended in FACS buffer containing the anti-LY75-mAb conjugated to MCC-DM1 (1 nM) and incubated on ice for 45 minutes. The cells were washed as above and resuspended in FACS buffer plus 1 ug/ml mouse anti-maytansine antibody and incubated in ice for 45 minutes. The cells were washed as above and resuspended in FACS buffer containing 2 ug/ml goat anti-mouse kappa RPE. The cells were incubated on ice for 45 minutes then washed as above. The cells were resuspended in FACS buffer at 200 ul per well. Mean fluorescence intensity of each

sample was determined using a Guava EasyCyte Plus HT Flow Cytometer (96 well plate formats) and the raw data was analyzed using the Guava Cytosoft.

Results

Figure 5a shows blocking with the anti-LY75-mAb-MCC-DM1 reduced the binding of anti-LY75-mAb. Analysis of the binding of LY75_A1 to COLO205 cells showed that LY75_A1 is unable to block binding of anti-LY75-mAb-MCC-DM1 (see Figure 5b). It can therefore be determined that the anti-LY75-mAb and LY75_A1 are non-competing antibodies and LY75_A1 recognizes a different and unique epitope of LY75 to that of other anti-LY75 antibodies.

Example 24 Epitope characterization of LY75_A1 by peptide micro array assay.

Method

The peptide microarray analysis was performed by LC Sciences, Houston TX, in brief the method comprised the following steps:- Contiguous 8mer peptides of LY75 protein having one amino acid overlap spanning residues 216 to 1666 of the full length LY75 protein were synthesized and immobilized on a microarray chip. The chip comprised three panels such that the experiment was performed in triplicate. The microarray was addressed with LY75_A1 to identify the peptides to which the antibody bound. The binding assay was performed under the following conditions:-

The microarray comprising the contiguous peptides in triplicate was washed with 1 mL of binding buffer at 4 °C for 20 min. It was then incubated with 1 µg/mL LY75_A1 in binding buffer (pH 7.0) at 4 °C for 2 hrs. The array was again washed with 0.5 mL of washing buffer at 4 °C for 30 min then incubated with 25 ng/mL anti-human IgG Alexa 647 conjugate in binding buffer (pH 7.0) at 4 °C for 1 hr. The array was again washed with 0.5 mL of washing buffer at 4 °C for 30 min.

The array was then Scanned at 635 nm and PMT 500 and the signal Intensity was recorded. The peptide was classed as detectable if it was present in at least 2/3 legal duplicates. The average signal intensity of the replicates was reported as the final signal intensity.

Results

As can be seen from Figure 6 antibody LY75_A1 showed specific binding to a number of peptides located on the array. The maximum signal seen for LY75_A1 binding was 25000 (scale 1-65535), with the average signal for all spots on the array being about 885. A signal intensity of 3000 was set as the background cut off point for non-specific binding. Based on the level of antibody binding signal intensity seen potential sequences forming the epitope for LY75_A1 were identified. These regions are shown in Figures 6a – 6j and as SEQ ID NOs: 22-31.

Example 25 LY75 A1 Peptide pull down assay

Method

1.1 Pull Down Assay

5 Recombinant LY75 protein was digested by on-bead tryptic proteolysis (Promega, US). The resulting digest peptides were recovered using a C18 capture column (Thermo Fisher Scientific). Purified peptides were then incubated with 200 µl of protein A beads cross-linked with LY75A1 antibody overnight at 4°C. Next day the unbound peptides were collected and the
10 beads were washed with 1 ml of PBS twice. The antibody bound peptides were eluted from beads by heating them at 90°C in 100 µl of PBS for 5 minutes. This elution step was repeated.

1.2 Mass spectrometry

15 Samples were analysed by liquid chromatography-mass spectrometry using a Waters nanoACQUITY UPLC System fitted with a nanoACQUITY UPLC BEH 130 C18 column, 75 µm x 250mm (186003545) and a LTQ Orbitrap Velos (Thermo Fisher Scientific). Peptides were eluted with a 300nl/min gradient increasing from 3% to 35% acetonitrile over 120 min. Full-scan mass spectra were acquired at 60000 resolving power between 400-2000 m/z mass range in the Orbitrap. In each cycle, the twenty most intense peptides were selected for CID MS/MS scans in
20 the linear ion trap with nanospray ion source fitted on the instrument.

1.3 Amino acid sequence analysis of peptide

25 The raw data generated from the LTQ Orbitrap Velos was processed through the Mascot software (Matrix Science) which uses the Mowse algorithm (Curr Biol. 1993 Jun 1;3(6):327-3) to infer amino acids sequences from the peak lists by searching against a sequence database consisting of Ensembl (<http://www.ensembl.org/index.html>), IPI (www.ebi.ac.uk/IPI/IPIhuman.html) and SwissProt (<http://www.uniprot.org>) along with contaminant protein sequences. Criteria for peptide identification included trypsin digestion, up to 2 missed cleavage sites and various biological and chemical modifications (oxidized
30 methionine, cysteine modification by MMTS or iodoacetamide and phosphorylation of serine, threonine and tyrosine). Peptides ranked 1 with an expectation value of 0.05% or less, an ion score of 28 or higher were loaded into the OGAP database.

1.4 Discrimination of LY75 associated peptides

35 The process to identify LY75 used the peptide sequences obtained experimentally by mass spectrometry, as described above, of naturally occurring human proteins to identify and organize coding exons in the published human genome sequence. These experimentally

determined sequences were compared with the OGAP® database which was compiled by processing and integration of peptide masses, peptide signatures, ESTs and Public Domain Genomic Sequence Data as described in International Patent Application WO2009/087462.

5 Results

The results of the peptide pull down assay using antibody LY75_A1 are shown in Table 1 below and in Figure 7. Peptides which were identified in both peptide elutions 1a and 1b in the pull down assay and in the microarray assay were considered to be the most likely candidates for forming the epitope.

Table 1: Comparison of peptide microarray and peptide pull down experiments

SEQ ID NO:	Peptide Identified by Microarray Assay	Peptide Identified by Pull Down Assay
	Region 1 (aa609-618)	-
	Region 2 (aa651-662)	-
46	Region 3 (aa761-780)	GWHFYDDR (765-772)
47	Region 4 (aa883-901)	ISEWPIDDHFTYSR(877 to 890)
48		FPVTFGEECLYMSAK(896-910)
49	Region 5 (aa1029-1040)	ELTYSNFHPLLVSGR(1030-1044)
50	Region 6 (aa1077-1093)	HFVSLCQK (1084-1091)
51	Region 7 (aa1107-1118)	QTLQNASETVK (1099-1109)
	Region 8 (aa1368-1378)	-
	Region 9 (aa1518-1528)	-
	Region 10 (aa1535-1554)	-

Table 1 shows that a number of overlapping LY75 peptide regions were identified in both the Peptide Microarray assay and in both elutions 1a and 1b the Peptide pull down assay. These regions are considered to be the most likely to contain the epitope recognized by antibody LY75_A1 as they are bound by LY75_A1 tested by both techniques employed.

Example 26

A number of activated B-cells Diffuse Large B-cell Lymphoma (ABC-DLBCL) cell lines (i.e. TMD8 and HBL1 cell lines) were exposed for 72 hours to increasing doses of LY75_DM4 (i.e. 0.018 – 0.037 – 0.075 – 0.15 – 0.3 – 0.6 – 1.2 nM) either alone or in combination with increasing doses of Rituximab (2.34 – 4.68 – 9.37 – 18.75 – 37.5 – 75 – 150) or Ibrutinib (15.6 –

31.2 – 62.5 – 125 – 250 – 500 – 1000 nM). This was followed by an MTT [3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazoliumbromide] assay.

The Chou-Talalay combination index (CI) was estimated using the Synergy R package (Preclinical versus Clinical Drugs Combination Studies. Chou TC. Leuk. Lymphoma.

5 2008;49(11):2059-2080). This provides a quantitative definition for strong synergism (<0.3), synergism (0.3-0.9), additive effect (0.9-1.1) or antagonism/no benefit (> 1.1).

Figures 8A-B show the synergistic effects seen between LY75_DM4 and Rituximab. Figure 9 shows the synergistic effects seen between LY75_DM4 and Ibrutinib. Supporting data is shown in Tables 2, 3, and 4 below.

10

Table 2: Chou-Talalay Combination Index (CI) of different doses of LY75_DM4 in combination with Rituximab on three TMD8 ABC-DLBCL cell lines.

N°	Rituximab (nM)	LY75_DM4 (nM)	CI
1	2.3438	0.01875	17.124
2	2.3438	0.075	0.225
3	2.3438	0.15	0.346
4	2.3438	0.3	0.583
5	2.3438	0.6	1.165
6	2.3438	1.2	2.514
7	2.3438	0.0375	0.145
8	4.6875	0.01875	1.499
9	4.6875	0.0375	0.147
10	4.6875	0.075	0.216
11	4.6875	0.15	0.346
12	4.6875	0.3	0.628
13	4.6875	0.6	1.165
14	4.6875	1.2	2.514
15	9.375	0.01875	2.918
16	9.375	0.0375	0.144
17	9.375	0.075	0.219
18	9.375	0.15	0.332
19	9.375	0.3	0.583
20	9.375	0.6	1.025
21	9.375	1.2	2.514
22	18.75	0.01875	0.8
23	18.75	0.0375	0.145
24	18.75	0.075	0.213
25	18.75	0.15	0.346
26	18.75	0.3	0.628
27	18.75	0.6	1.165

28	18.75	1.2	2.514
29	37.5	0.01875	11.427
30	37.5	0.075	0.216
31	37.5	0.15	0.332
32	37.5	0.3	0.583
33	37.5	0.6	1.165
34	37.5	1.2	2.655
35	37.5	0.0375	0.151
36	75	0.01875	0.216
37	75	0.0375	0.147
38	75	0.075	0.209
39	75	0.15	0.332
40	75	0.3	0.628
41	75	0.6	1.165
42	75	1.2	2.514
43	150	0.01875	45.466
44	150	0.0375	0.152
45	150	0.075	0.209
46	150	0.15	0.332
47	150	0.3	0.628
48	150	0.6	1.257
49	150	1.2	2.770

Table 3: Chou-Talalay Combination Index (CI) of different doses of LY75_DM4 in combination with Rituximab on three HBL-1 ABC-DLBCL cell lines.

N°	Rituximab (nM)	LY75_DM4 (nM)	CI
1	2.3438	0.0188	0.534
2	2.3438	0.0750	0.958
3	2.3438	0.1500	0.603
4	2.3438	0.3000	0.509
5	2.3438	0.6000	0.454
6	2.3438	1.2000	0.817
7	2.3438	0.0375	0.413
8	4.6875	0.0188	0.568
9	4.6875	0.0375	0.837
10	4.6875	0.0750	0.919
11	4.6875	0.1500	0.635
12	4.6875	0.3000	0.580
13	4.6875	0.6000	0.501
14	4.6875	1.2000	0.908
15	9.375	0.0188	0.249
16	9.375	0.0375	1.576

17	9.375	0.0750	0.717
18	9.375	0.1500	0.439
19	9.375	0.3000	0.618
20	9.375	0.6000	0.501
21	9.375	1.2000	0.908
22	18.75	0.0188	0.787
23	18.75	0.0375	0.497
24	18.75	0.0750	0.562
25	18.75	0.1500	0.545
26	18.75	0.3000	0.510
27	18.75	0.6000	0.366
28	18.75	1.2000	0.817
29	37.50	0.0375	0.734
30	37.50	0.0750	0.548
31	37.50	0.0188	0.543
32	37.50	0.1500	0.465
33	37.50	0.3000	0.476
34	37.50	0.6000	0.324
35	37.50	1.2000	0.648
36	75	0.0188	0.442
37	75	0.0375	1.578
38	75	0.0750	0.432
39	75	0.1500	0.493
40	75	0.3000	0.383
41	75	0.6000	0.409
42	75	1.2000	0.908
43	150	0.0188	0.680
44	150	0.0375	0.640
45	150	0.0750	0.911
46	150	0.1500	0.498
47	150	0.3000	0.355
48	150	0.6000	0.366
49	150	1.2000	0.817

Table 4: Chou-Talalay Combination Index (CI) of different doses of LY75_DM4 in combination with Ibrutinib on three HBL-1 ABC-DLBCL cell lines.

N°	Ibrutinib (nM)	LY75_DM4 (nM)	CI
1	1.5625	0.01875	0.481
2	1.5625	0.0375	0.627
3	1.5625	0.075	0.862
4	1.5625	0.15	0.360

5	1.5625	0.3	0.287
6	1.5625	0.6	0.270
7	1.5625	1.2	0.343
8	3.125	0.01875	0.411
9	3.125	0.0375	0.358
10	3.125	0.075	0.461
11	3.125	0.15	0.287
12	3.125	0.3	0.251
13	3.125	0.6	0.246
14	3.125	1.2	0.394
15	6.25	0.01875	0.204
16	6.25	0.0375	0.219
17	6.25	0.075	0.275
18	6.25	0.15	0.226
19	6.25	0.3	0.216
20	6.25	0.6	0.222
21	6.25	1.2	0.343
22	12.5	0.01875	0.205
23	12.5	0.0375	0.152
24	12.5	0.075	0.287
25	12.5	0.15	0.217
26	12.5	0.3	0.228
27	12.5	0.6	0.222
28	12.5	1.2	0.343
29	25	0.01875	0.152
30	25	0.0375	0.183
31	25	0.075	0.202
32	25	0.15	0.199
33	25	0.3	0.193
34	25	0.6	0.222
35	25	1.2	0.343
36	50	0.01875	0.374
37	50	0.0375	0.473
38	50	0.075	0.287
39	50	0.15	0.192
40	50	0.3	0.194
41	50	0.6	0.246
42	50	1.2	0.394
43	100	0.01875	0.715
44	100	0.0375	0.748
45	100	0.075	0.457
46	100	0.15	0.187
47	100	0.3	0.195

48	100	0.6	0.222
49	100	1.2	0.394

SEQUENCES:

SEQ ID No	Description	Sequence
1	A1_VH aa	EVQLVESGGGLVKPGGSLRLSCAASGFTYSNAWMSWVRQAPG KGLEWVGRIKSKTDGGTTDYAAPVQGRFTISRDDSKNTLYLQMN SLKTEDTAVYYCTIFGVVSFDYWGQGTLVTVSS
2	A1_VL aa	DVQMTQSPSSLSASVGDRVITICRASQSIDYLSWYQQRPGKAP NLLIYAASNLKTGVPSRFSGSGSGTDFTLTISTLQPEDFATYYCQ QSYRSPWTFGQGTKVEIKR
3	A1_VH nt	gagggtgcagctggtggagctgggggaggcttggtaaagccgggggggtagactct cctgtgcagcctctggcttacttacagtaacgcctggatgagctgggtccgccagggtcca gggaaggggctggagtggttggccgtattaaaagcaaaactgatggtgggacaacaga ctacgtgcacccgtgcaaggcagattcaccatctcaagagatgattcaaaaaacacgctg tatctgcaaatgaacagcctgaaaaccgaggacacagccgtgtattactgtacgatttttga gtggtagctttgactactggggccagggaaccctggcaccgtctcctca
4	A1_VL nt	gacgtccagatgaccagctcctcatcctccctgtctgcatctgttgagacagagtcaccatc acttgccgggcaagtgcagagcattagcgactatttaagttggtatcagcagagaccaggga aagcccctaacctcctgatctatgctgcatccaatttaaagactgggggtcccatcaagggtca gtggcagtggtatctgggacagattcactctcaccatcagcactctgcaacctgaagattttg caacgtactactgtcaacagagttacagggtcccgtggacgttcggccaagggaaccaagg tggaatcaaacga
5	A1_VH_CDR 1 aa	NAWMS
6	A1_VH_CDR 2 aa	RIKSKTDGGTTDYAAPVQG
7	A1_VH_CDR 3 aa	FGVVSFDY
8	A1_VL_CDR1 aa	RASQSIDYLS
9	A1_VL_CDR2 aa	AASNLKT
10	A1_VL_CDR3 aa	QSYRSPWT
11	VH3 3- 15/D4 411	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPG KGLEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMN SLKTEDTAVYYCTTTT
12	JH4	YFDYWGQGTLVTVSS

13	O12	DIQMTQSPSSLSASVGDRVITICRASQSISSYLNWYQQKPGKAP KLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQ QSYS
14	JK1	WTFGQGGTKVEIKR
15	LY75 (DEC- 205)	MRTGWATPRRPAGLLMLLFWFFDLAEPSGRAANDPFTIVHGNT GKCIKPVYGWIVADDCDEDEDKLWKWVSQHRFLHLSQKCLGL DITKSVNELRMFSCDSSAMLWWKCEHHSYGAARYRLALKDGH GTAISNASDVWKKGGSEESLCDQPYHEIYTRDGNSYGRPCEFPF LIDGTWHHDCILDEDHSGPWCATTLNIEYDRKWGICLKPENGCE DNWEKNEQFGSCYQFNTQTALSWKEAYVSCQNQGADLLSINSA AELTYLKEKEGIAKIFWIGLNQLYSARGWEWSDHKPLNFLNWD DRPSAPTIGGSSCARMDAESGLWQSFSCEAQLPYVCRKPLNNT VELTDVWTYSOTRCDAGWLPNNGFCYLLVNESNSWDKAHAKCK AFSSDLISIHSLADVEVVVTKLHNEDIKEEVWIGLKNINIPTLFQWS DGTEVTLTYWDENEPNPYPNKTNPNCVSYLGELGQWKVQSCEEK LKYVCKRKGEKLNDASSDKMCPPEDEGWKRHGETCYKIYEDEV FGTNCNLTITSRFEQEYLNDLMKKYDKSLRKYFWTGLRDVDSG EYNWATVGGRRRAVTFNSWNFLPASPGGCAMSTGKSVGKW EVKDCRSFKALSICKKMSGPLGPEEASPKPDDPCPEGWQSFP SLSCYKVFHAERIVRKRNWEEAERFCQALGAHLSSFSHVDEIKEF LHFLTDQFSGQHWLWIGLNKRSPDLQGSWQWSDRTPVSTIIMP NEFQQDYDIRDCAAVKVFHRPWRRGWHFYDDREFIYLRPFACD TKLEWVCQIPKGRTPKTPDWYNPDRAIGHGPPLIEGSEYWFVAD LHLNIEEAVLYCASNHSFLATITSFVGLKAIKNKIANISGDGQKWW IRISEWPIDDHFTYSRYPWHRFPVTFGEECLYMSAKTWLIDLGP TDCSTKLFPICEKYNVSSLEKYSPPSAKVQCSEQWIPFQNKCF KIKPVSLTFSQASDTCHSYGGTLPSVLSQIEQDFITSLLPDMEATL WIGLRWTAYEKINKWTDNRELTYSNFHPLLVSGRLRIPENFFEEE SRYHCALILNLQKSPFTGTWNFTSCSERHFVSLCQKYSEVKS TLQNASETVKYLNLYKIIPKTLTWHSAREKLSNMQLVSITDPY QQAFLSVQALLHNSSLWIGLFSQDDELNFGWSDGKRLHFSRWA ETNGQLEDVLDLTDGFWKTVDNDNQPAGIYYSNETEKEV KPVDSVKCPSPVLNTPWIPFQNCYNFIITKNRHMATTQDEVHTK CQKLNPKSHLSIRDEKENNFVLEQLLYFNIMASWVMLGITYRNK SLMWFDKTPLSYTHWRAGRPTIKNEKFLAGLSTDGFWDIQTFKVI EEAVYFHQHSILACKIEMVDYKEEYNTTLPQFMPYEDGIYSVIQKK VTWYEALNMCSQSGGHLASVHNQNGQLFLEDIVKRDGFPLWVG

		LSSHDGSESSFEWSDGSTFDYIPWKGQTSPGNCVLLDPKGTWK HEKCNSVKDGAICYKPTKSKKLSRLTYSSRCPAAKENGSRWIIQY KGHCYKSDQALHSFSEAKKLC SKHDHSATIVSIKDEDENKFVSRL MRENNNITMRVWLGLSQHSVDQSWSWLDGSEVTFVKWENKSK SGVGRCSMLIASNETWKKVECEHGFGRVVCKVPLGPDYTAIIIV ATLSILVLMGGLIWFLFQRHRLHLAGFSSVRYAQGVNEDEIMLPS FHD
16	A1_VH_FR1	EVQLVESGGGLVKPGGSLRLSCAASGFTYS
17	A1_VH_FR2	WVRQAPGKGLEWVG
18	A1_VH_FR3	RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTI
19	A1_VH_FR4	WGQGTLVTVSS
20	A1_VL_FR1	DVQMTQSPSSLSASVGDRVITC
21	A1_VL_FR2	WYQQRPGKAPNLLIY
22	A1_VL_FR3	GVPSRFSGSGSGTDFTLTISTLQPEDFATYYC
23	A1_VL_FR4	FGQGTKVEIKR
24	LY75 609-618	WEVKDCRSFK
25	LY75 651-662	PASLSCYKVFHA
26	LY75 761-780	PWRRGWHFYDDREFIYLRPF
27	LY75 883-901	DDHFTYSRYPWHRFPVTFG
28	LY75 1029-1040	RELTYSNFHPLL
29	LY75 1077-1093	FTSCSERHFVSLCQKYS
30	LY75 1107-1118	TVKYLNNLYKII
31	LY75 1368-1378	EAVYFHQHSIL
32	LY75 1518-1528	KKLSRLTYSSC
33	LY75 1535-1554	NGSRWIIQYKGHCYKSDQALH
34	LY75 877-901	ISEWPIDDHFTYSRYPWHRFPVTFG
35	LY75 1099-1118	QTLQNASETVKYLNNLYKII
36	LY75 883-892	DDHFTYSRYP
37	LY75 1077-	FTSCSERHFVSLCQK

	1091	
38	Rituximab VH	QVQLQQPGAELVKPGASVKMSCKASGYTFTSYNMHWVKQTPGRG LEWIGAIYPGNGDTSYNQKFKGKATLTADKSSSTAYMQLSSLTSEDS AVYYCARSTYYGGDWYFNVWGAGTTVTVSAASTKGPSVFPLAPSS KSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKAEPKSCDKT HTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED PEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLN GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRD ELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGD SFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
39	Rituximab VL	QIVLSQSPAILSASPGEKVTMTCRASSSVSYIHWFQQKPGSSPKP WIYATSNLASGVPVRFSGSGSGTSYSLTISRVEAEDAATYYCQQ WTSNPPTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCL LNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYLSST LTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
40	Rituximab VH_CDR1	SYNMH
41	Rituximab VH_CDR2	AIYPGNGDTSYNQKFKG
42	Rituximab VH_CDR3	STYYGGDWYFNV
43	Rituximab VL_CDR1	RASSSVSYIH
44	Rituximab VL_CDR2	ATSNLAS
45	Rituximab VL_CDR3	QQWTSNPPTF
46	(765-772)	GWHFYDDR
47	(877 to 890)	ISEWPIDDHFTYSR
48	(896-910)	FPVTFGEECLYMSAK
49	(1030-1044)	ELTYSNHFPLLVSGR
50	(1084-1091)	HFVSLCQK
51	(1099-1109)	QTLQNASETVK
52	Linker	Gly-Phe-Leu-Gly

Sequence Listing Free Text:

SEQ ID NO: 38 <223> Rituximab VH sequence
SEQ ID NO: 39 <223> Rituximab VL sequence
5 SEQ ID NO: 40 <223> Rituximab VH_CDR1
SEQ ID NO: 41 <223> Rituximab VH_CDR2
SEQ ID NO: 42 <223> Rituximab VH_CDR3
SEQ ID NO: 43 <223> Rituximab VL_CDR1
SEQ ID NO: 44 <223> Rituximab VL_CDR2
10 SEQ ID NO: 45 <223> Rituximab VL_CDR3
SEQ ID NO: 52 <223> Linker
SEQ ID NOs: 53-163 <223> Peptide
SEQ ID NOs: 165-207 <223> Peptide

CLAIMS

1. A method of treating diffuse large B-cell lymphoma (DLBCL) in a patient comprising simultaneously, sequentially or separately administering to a patient in need thereof therapeutically-effective amounts of a pharmaceutical combination comprising:

(A) an anti-LY75 antibody, or an antigen-binding portion thereof, said antibody or portion thereof comprising:

a) a heavy chain variable region comprising:

- i) a first vhCDR comprising SEQ ID NO: 5;
- ii) a second vhCDR comprising SEQ ID NO: 6; and
- iii) a third vhCDR comprising SEQ ID NO: 7; and

b) a light chain variable region comprising:

- i) a first vlCDR comprising SEQ ID NO: 8;
- ii) a second vlCDR comprising SEQ ID NO: 9; and
- iii) a third vlCDR comprising SEQ ID NO: 10;

wherein the anti-LY75 antibody or antigen-binding portion thereof further comprises a covalently-attached moiety, and wherein said moiety is a maytansinoid;

and

(B) Rituximab.

2. Use of components (A) and (B) of the pharmaceutical combination as defined in claim 1 in the manufacture of a pharmaceutical combination for simultaneous, separate or sequential use in the treatment of diffuse large B-cell lymphoma (DLBCL).

3. Use of components (A) and (B) of the pharmaceutical combination as defined in claim 1 in the treatment of diffuse large B-cell lymphoma (DLBCL).

4. Use of component (A) of the pharmaceutical combination as defined in claim 1 in the manufacture of a pharmaceutical combination for simultaneous, separate or sequential use with component (B) in the treatment of diffuse large B-cell lymphoma (DLBCL).

5. Use of component (B) of the pharmaceutical combination as defined in claim 1 in the manufacture of a pharmaceutical combination for simultaneous, separate or sequential use with component (A) in the treatment of diffuse large B-cell lymphoma (DLBCL).

```

SEQ ID No: 11      EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIKSKTDGGTT
SEQ ID No: 1       EVQLVESGGGLVKPGGSLRLSCAASGFTYSNAWMSWVRQAPGKGLEWVGRIKSKTDGGTT
SEQ ID No: 12      -----
                    *****:*****
                    :
SEQ ID No: 11      DYAAPVKGRETISRDDSKNTLYLQMNSLKTEDTAVYYCTTTTVT-----
SEQ ID No: 1       DYAAPVQGRETISRDDSKNTLYLQMNSLKTEDTAVYYCTIFGVVSFDYWGQGLVTVSS
SEQ ID No: 12      -----YFDYWGQGLVTVSS
                    *****:*****
                    :

```

FIGURE 1

```

SEQ ID No: 2       DVQMTQSPSSLSASVGDRVTITCRASQSIDYLSWYQQRPGKAPNLLIYAASNLKTGVPS
SEQ ID No: 13      DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPS
SEQ ID No: 14      -----
                    *:***** ** *:*****:***** *: ****
                    :
SEQ ID No: 2       RFGSGSGTDFTLTISTLQPEDFATYYCQQSYRSPWTFGQGTKVEIKR
SEQ ID No: 13      RFGSGSGTDFTLTISLQPEDFATYYCQQSYS-----
SEQ ID No: 14      -----WTFGQGTKVEIKR
                    *****:*****

```

FIGURE 2




Figure 3a




Figure 3b




Figure 3c




Figure 3d




Figure 3e




Figure 3f




Figure 3g




Figure 3h




Figure 3i




Figure 3j




Figure 3k




Figure 3l




Figure 3m




Figure 3n




Figure 3o




Figure 3p




Figure 3q




Figure 3r




Figure 3s




Figure 3t




Figure 3u




Figure 3v




Figure 3w




Figure 4a




Figure 4b




Figure 4c




Figure 4d




Figure 4e




Figure 4f




Figure 5a




Figure 5b




Figure 6a




Figure 6b




Figure 6c




Figure 6d




Figure 6e




Figure 6f




Figure 6g




Figure 6h




Figure 6i




Figure 6j

Figure 7

```
sp|O60449|LY75_HUMAN      MRTGWATPRRPAGLLMLLFWFDDLAEPSGRAANDPFTIVHGNTGKCIKPVYGWIVADDCD  60
MS-Peptides_Elutela      -----CIKPVYGWIVADDCD
MS-Peptides_Elutelb      -----CIKPVYGWIVADDCD
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      ETEDKLWKWVSQHRLFHLHSQKCLGLDITKSVNELRMFSCDSSAMLWWKCEHHSLYGAAR 120
MS-Peptides_Elutela      ETEDKLWK-----CEHHSLYGAAR
MS-Peptides_Elutelb      ETEDKLWK-----
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      YRLALKDGHGTAISNASDVWKKGGSEESLCDQPYHEIYTRDGN SYGRPCEFPFLIDGTWH 180
MS-Peptides_Elutela      -----DGHGTAISNASDVWKKGGSEESLCDQPYHEIYTR-----
MS-Peptides_Elutelb      -----KGGSEESLCDQPYHEIYTR-----
MS-Peptides_Elute2a      -----GGSEESLCDQPYHEIYTR-----
MS-Peptides_Elute2b      -----GGSEESLCDQPYHEIYTR-----
LC-Peptides               -----DGHGTAISNASDVWK-----

sp|O60449|LY75_HUMAN      HDCILDEDHSGPWCATTLNIEYDRKWGICLKPEGNCEDNWEKNEQFGSCYQFNTQTALSW 240
MS-Peptides_Elutela      -----
MS-Peptides_Elutelb      -----WGICLKPEGNCEDNWEK-----
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      KEAYVSCQNQGADLLSINSAAELTYLKEKEGIKIFWIGLNQLYSARGWEWS DHKPLNFL 300
MS-Peptides_Elutela      -----IFWIGLNQLYSAR-----
MS-Peptides_Elutelb      -----
MS-Peptides_Elute2a      -----IFWIGLNQLYSAR-----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      NWDPRPSAPTIGSSCARMDAESGLWQSFSCQAQLPYVCRKPLNNTVELTDVW TYS DTR 360
MS-Peptides_Elutela      -----
MS-Peptides_Elutelb      -----
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      CDAGWLPNNGFCYLLVNESNSWDKAHAKCKAFSSDLISIHSLADVEVVVTKLHNEDIKEE 420
MS-Peptides_Elutela      -----LHNEDIKEE
MS-Peptides_Elutelb      -----LHNEDIKEE
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----

sp|O60449|LY75_HUMAN      VWIGLKNINIPTLFQWSDGTEVTLT YWDENEPNVPYKTPNCVSYLGELGQWKVQSCEEK 480
MS-Peptides_Elutela      VWIGLK-----TPNCVSYLGELGQWK-----
MS-Peptides_Elutelb      VWIGLK-----TPNCVSYLGELGQWK-----
MS-Peptides_Elute2a      -----
MS-Peptides_Elute2b      -----
LC-Peptides               -----
```

sp|O60449|LY75_HUMAN LKYVCKRKGEKLNDASSDKMCPDEGWKRHGETCYKIYEDEVFPGTNCNLITITSRFEQY 540
MS-Peptides_Elutela -----
MS-Peptides_Elutelb -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN LNDLMKKYDKSLRKYFWTGLRDVDSCGEYNWATVGRRRAVTFSNWNFLEPASPGGCVM 600
MS-Peptides_Elutela -----YFWTGLRDVDSCGEYNWATVGGR-----
MS-Peptides_Elutelb -----YFWTGLRDVDSCGEYNWATVGGR-----
MS-Peptides_Elute2a -----YFWTGLRDVDSCGEYNWATVGGR-----
MS-Peptides_Elute2b -----YFWTGLRDVDSCGEYNWATVGGR-----
LC-Peptides -----

sp|O60449|LY75_HUMAN STGKSVGKWEVKDCRSFKALSICKKMSGPLGPPEEASPKPDDPCPEGWQSFPASLSCYKVF 660
MS-Peptides_Elutela ----SVGKWEVKDCR---ALSICKK-----
MS-Peptides_Elutelb -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----WEVKDCRSFK-----PASLSCYKVF

sp|O60449|LY75_HUMAN HAERIVRKRNWEEAERFCQALGAHLSSFSHVDEIKEFLHFLTDQFSGQHWLWIGLNKRSP 720
MS-Peptides_Elutela -----RNWEEAER-----RSP
MS-Peptides_Elutelb -----RNWEEAER-----RSP
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides HA-----

sp|O60449|LY75_HUMAN DLQGSWQWSDRTPVSTIIMPNEFQQDYDIRDCAAVKVFHRPWRRGWHFYDDREFIYLRF 780
MS-Peptides_Elutela -----TPVSTIIMPNEFQQDYDIR-----
MS-Peptides_Elutelb DLQGSWQWSDRTPVSTIIMPNEFQQDYDIR-----
MS-Peptides_Elute2a -----GWHFYDDR-----
MS-Peptides_Elute2b -----
LC-Peptides -----PWRRGWHFYDDREFIYLRF

sp|O60449|LY75_HUMAN ACDTKLEWVCQIPKGRTPKTPDWYNPDRAGIHGPPLIIEGSEYWFWADLHLNYYEEAVLYC 840
MS-Peptides_Elutela -----
MS-Peptides_Elutelb -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN ASNHSFLATITSFVGLKAIKNKIANISGDGQKWWIRISEWPI **DDHFTYSRYF**WHRFPVTF 900
MS-Peptides_Elutela -----ISEWPI **DDHFTYSR**-----FPVTF
MS-Peptides_Elutelb -----ISEWPI **DDHFTYSR**-----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----**DDHFTYSRYF**WHRFPVTF

sp|O60449|LY75_HUMAN GEECLYMSAKTWLIDL GKPTDCSTKL PFI CEKYNVSSLEK YSPDSAAKVQCSEQWIPFQN 960
MS-Peptides_Elutela GEECLYMSAKTWLIDL GKPTDCSTK-----YNVSSLEK-----VQCSEQWIPFQN
MS-Peptides_Elutelb -----TWLIDL GKPTDCSTK-----YNVSSLEK-----VQCSEQWIPFQN
MS-Peptides_Elute2a -----TWLIDL GKPTDCSTK-----
MS-Peptides_Elute2b -----TWLIDL GKPTDCSTK-----
LC-Peptides G-----

sp|O60449|LY75_HUMAN KCFLKIKPVSLTFSQASDTHCHSYGGTLPVSLSQIEQDFITSLLPDMEATLWIGLRWTAYE 1020
MS-Peptides_Elutela -----

```
MS-Peptides_Elute1b -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN KINKWTDNRELTYSNFHPLLVSGLRIPENFFEEESRYHCALILNLQKSPFTGTWN[FTSC] 1080
MS-Peptides_Elute1a K-----ELTYSNFHPLLVSGLRIPENFFEEESRYHCALILNLQK-----
MS-Peptides_Elute1b K-----YHCALILNLQK-----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----RELTYSNFHPLL-----[FTSC]

sp|O60449|LY75_HUMAN [SERHVFVSLCQK]YSEVKSRQTLQNASE[TVKYLNNLYKII]PKTLTWHS AKRECLKSNMQLVS 1140
MS-Peptides_Elute1a --[HFVSLCQK]-----QTLQNASE[TVK]-----TLTWHS AK-----
MS-Peptides_Elute1b --[HFVSLCQK]-----QTLQNASE[TVK]-----TLTWHS AK-----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides [SERHVFVSLCQKYS]-----[TVKYLNNLYKII]-----
***** ***

sp|O60449|LY75_HUMAN ITDPYQQAFLSVQALLHNSSLWIGLFSQDDELNFGWSDGKRLHFSRWAETNGQLEDCVVL 1200
MS-Peptides_Elute1a -----
MS-Peptides_Elute1b -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN DTDGFWKTVDCNDNQPGAICYSGNETEKEVKPVD SVKCPSPVLNTPWIPFQNCYNFII 1260
MS-Peptides_Elute1a -----
MS-Peptides_Elute1b -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN TKNRHMAT TQDEVHTKCQKLNPKSHILSIRDEKENNFVLEQLLYFN YMASWVMLGITYRN 1320
MS-Peptides_Elute1a -----SHILSIR-----
MS-Peptides_Elute1b --NRHMAT TQDEVHTK-----SHILSIR-----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN KSLMWFDK TPLSYTHWRAGRPTIKNEKFLAGLSTDGFWDIQTFKVIEEAVYFH QHSILAC 1380
MS-Peptides_Elute1a -SLMWFDK TPLSYTHWR-----
MS-Peptides_Elute1b -SLMWFDK-----
MS-Peptides_Elute2a -SLMWFDK-----
MS-Peptides_Elute2b -----
LC-Peptides -----EAVYFH QHSIL--

sp|O60449|LY75_HUMAN KIEMVDYKEEYNTLPQFMPYEDGIYSVIQKKVTWY EALNMCSQSGGHLASVHNQNGQLF 1440
MS-Peptides_Elute1a -----
MS-Peptides_Elute1b -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----

sp|O60449|LY75_HUMAN LEDIVKRDGFPLWVGLSSH DGSSESSF EWSDGSTFDYIPWK GQ TSPGNCVLLDPKGTWKHE 1500
MS-Peptides_Elute1a -----
MS-Peptides_Elute1b -----
MS-Peptides_Elute2a -----
MS-Peptides_Elute2b -----
LC-Peptides -----
```

sp O60449 LY75_HUMAN	KCNSVKDGAICYKPTKSKKLSRLTYSSRCFAAKENGSRWIQYKGHCYKSDQALHSFSEAK	1560
MS-Peptides_Elutela	-----	
MS-Peptides_Elute1b	-----	
MS-Peptides_Elute2a	-----	
MS-Peptides_Elute2b	-----	
LC-Peptides	-----KKLSRLTYSS-C-----NGSRWIQYKGHCYKSDQALH-----	

sp O60449 LY75_HUMAN	KLCSKHDHSATIVSIKDEDEDENKFVSRMLRENNITMRVWLGLSQHSVDQSWSLDGSEVT	1620
MS-Peptides_Elutela	-----HDHSATIVSIKDEDEDENKFVSR-----	
MS-Peptides_Elute1b	-----HDHSATIVSIKDEDEDENKFVSR-----	
MS-Peptides_Elute2a	-----HDHSATIVSIKDEDEDENKFVSR-----	
MS-Peptides_Elute2b	-----	
LC-Peptides	-----	

sp O60449 LY75_HUMAN	FVKWENKSKSGVGRCSMLIASNETWKKVECEHGFGRVVCKVPLGPDYTAIAIIVATLSIL	1680
MS-Peptides_Elutela	-----VECEHGFGR-----	
MS-Peptides_Elute1b	-----VECEHGFGR-----	
MS-Peptides_Elute2a	-----	
MS-Peptides_Elute2b	-----	
LC-Peptides	-----	

sp O60449 LY75_HUMAN	VLMGGLIWFLFQRHRLHLAGFSSVRYAQGVNEDEIMLPSEFHD	1722
MS-Peptides_Elutela	-----	
MS-Peptides_Elute1b	-----	
MS-Peptides_Elute2a	-----	
MS-Peptides_Elute2b	-----	
LC-Peptides	-----	

Figure 8A




Figure 8B





Figure 9



eolf-seql.txt
SEQUENCE LISTING

<110> Berlin-Chemie AG

<120> Pharmaceutical combinations

<130> 489.131183/01

<160> 207

<170> PatentIn version 3.5

<210> 1

<211> 119

<212> PRT

<213> Human

<400> 1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Tyr Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
50 55 60

Pro Val Gln Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95

Tyr Cys Thr Ile Phe Gly Val Val Ser Phe Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser
115

<210> 2

eolf-seql.txt

<211> 108
<212> PRT
<213> Human

<400> 2

Asp Val Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Asp Tyr
20 25 30

Leu Ser Trp Tyr Gln Gln Arg Pro Gly Lys Ala Pro Asn Leu Leu Ile
35 40 45

Tyr Ala Ala Ser Asn Leu Lys Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Thr Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Arg Ser Pro Trp
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 3
<211> 357
<212> DNA
<213> Human

<400> 3

gaggtgcagc tgggtggagtc tggggggaggc ttggtaaagc cggggggggtc ccttagactc	60
tcctgtgcag cctctggctt cacttacagt aacgcctgga tgagctgggt ccgccaggct	120
ccagggaagg ggctggagtg ggttggccgt attaaaagca aaactgatgg tgggacaaca	180
gactacgctg caccctgca aggcagattc accatctcaa gagatgattc aaaaaacacg	240
ctgtatctgc aaatgaacag cctgaaaacc gaggacacag ccgtgtatta ctgtacgatt	300
tttggagtgg ttagctttga ctactggggc cagggaaccc tggtcaccgt ctcctca	357

eolf-seql.txt

<210> 4
 <211> 324
 <212> DNA
 <213> Human

<400> 4
 gacgtccaga tgaccagtc tccatcctcc ctgtctgcat ctgttggaga cagagtcacc 60
 atcacttgcc gggcaagtca gagcattagc gactatttaa gttggtatca gcagagacca 120
 gggaaagccc ctaacctcct gatctatgct gcatccaatt taaagactgg ggtcccatca 180
 aggttcagtg gcagtggatc tgggacagat ttcactctca ccatcagcac tctgcaacct 240
 gaagattttg caacgtacta ctgtcaacag agttacaggt ccccgaggac gttcggccaa 300
 gggaccaagg tggaaatcaa acga 324

<210> 5
 <211> 5
 <212> PRT
 <213> Human

<400> 5

Asn Ala Trp Met Ser
 1 5

<210> 6
 <211> 19
 <212> PRT
 <213> Human

<400> 6

Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala Pro
 1 5 10 15

Val Gln Gly

<210> 7
 <211> 8
 <212> PRT
 <213> Human

<400> 7

eolf-seql.txt

Phe Gly Val Val Ser Phe Asp Tyr
1 5

<210> 8
<211> 11
<212> PRT
<213> Human

<400> 8

Arg Ala Ser Gln Ser Ile Ser Asp Tyr Leu Ser
1 5 10

<210> 9
<211> 7
<212> PRT
<213> Human

<400> 9

Ala Ala Ser Asn Leu Lys Thr
1 5

<210> 10
<211> 9
<212> PRT
<213> Human

<400> 10

Gln Gln Ser Tyr Arg Ser Pro Trp Thr
1 5

<210> 11
<211> 104
<212> PRT
<213> Human

<400> 11

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

eolf-seql.txt

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60

Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
 65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Thr Thr Thr Thr Val Thr
 100

<210> 12
 <211> 15
 <212> PRT
 <213> Human

<400> 12

Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 1 5 10 15

<210> 13
 <211> 93
 <212> PRT
 <213> Human

<400> 13

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
 20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly

eolf-seql.txt

50

55

60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser
85 90

<210> 14
<211> 13
<212> PRT
<213> Human

<400> 14

Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
1 5 10

<210> 15
<211> 1722
<212> PRT
<213> Human

<400> 15

Met Arg Thr Gly Trp Ala Thr Pro Arg Arg Pro Ala Gly Leu Leu Met
1 5 10 15

Leu Leu Phe Trp Phe Phe Asp Leu Ala Glu Pro Ser Gly Arg Ala Ala
20 25 30

Asn Asp Pro Phe Thr Ile Val His Gly Asn Thr Gly Lys Cys Ile Lys
35 40 45

Pro Val Tyr Gly Trp Ile Val Ala Asp Asp Cys Asp Glu Thr Glu Asp
50 55 60

Lys Leu Trp Lys Trp Val Ser Gln His Arg Leu Phe His Leu His Ser
65 70 75 80

Gln Lys Cys Leu Gly Leu Asp Ile Thr Lys Ser Val Asn Glu Leu Arg
85 90 95

eolf-seql.txt

Met Phe Ser Cys Asp Ser Ser Ala Met Leu Trp Trp Lys Cys Glu His
100 105 110

His Ser Leu Tyr Gly Ala Ala Arg Tyr Arg Leu Ala Leu Lys Asp Gly
115 120 125

His Gly Thr Ala Ile Ser Asn Ala Ser Asp Val Trp Lys Lys Gly Gly
130 135 140

Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile Tyr Thr Arg
145 150 155 160

Asp Gly Asn Ser Tyr Gly Arg Pro Cys Glu Phe Pro Phe Leu Ile Asp
165 170 175

Gly Thr Trp His His Asp Cys Ile Leu Asp Glu Asp His Ser Gly Pro
180 185 190

Trp Cys Ala Thr Thr Leu Asn Tyr Glu Tyr Asp Arg Lys Trp Gly Ile
195 200 205

Cys Leu Lys Pro Glu Asn Gly Cys Glu Asp Asn Trp Glu Lys Asn Glu
210 215 220

Gln Phe Gly Ser Cys Tyr Gln Phe Asn Thr Gln Thr Ala Leu Ser Trp
225 230 235 240

Lys Glu Ala Tyr Val Ser Cys Gln Asn Gln Gly Ala Asp Leu Leu Ser
245 250 255

Ile Asn Ser Ala Ala Glu Leu Thr Tyr Leu Lys Glu Lys Glu Gly Ile
260 265 270

Ala Lys Ile Phe Trp Ile Gly Leu Asn Gln Leu Tyr Ser Ala Arg Gly
275 280 285

Trp Glu Trp Ser Asp His Lys Pro Leu Asn Phe Leu Asn Trp Asp Pro
290 295 300

eolf-seql.txt

Asp Arg Pro Ser Ala Pro Thr Ile Gly Gly Ser Ser Cys Ala Arg Met
305 310 315 320

Asp Ala Glu Ser Gly Leu Trp Gln Ser Phe Ser Cys Glu Ala Gln Leu
325 330 335

Pro Tyr Val Cys Arg Lys Pro Leu Asn Asn Thr Val Glu Leu Thr Asp
340 345 350

Val Trp Thr Tyr Ser Asp Thr Arg Cys Asp Ala Gly Trp Leu Pro Asn
355 360 365

Asn Gly Phe Cys Tyr Leu Leu Val Asn Glu Ser Asn Ser Trp Asp Lys
370 375 380

Ala His Ala Lys Cys Lys Ala Phe Ser Ser Asp Leu Ile Ser Ile His
385 390 395 400

Ser Leu Ala Asp Val Glu Val Val Val Thr Lys Leu His Asn Glu Asp
405 410 415

Ile Lys Glu Glu Val Trp Ile Gly Leu Lys Asn Ile Asn Ile Pro Thr
420 425 430

Leu Phe Gln Trp Ser Asp Gly Thr Glu Val Thr Leu Thr Tyr Trp Asp
435 440 445

Glu Asn Glu Pro Asn Val Pro Tyr Asn Lys Thr Pro Asn Cys Val Ser
450 455 460

Tyr Leu Gly Glu Leu Gly Gln Trp Lys Val Gln Ser Cys Glu Glu Lys
465 470 475 480

Leu Lys Tyr Val Cys Lys Arg Lys Gly Glu Lys Leu Asn Asp Ala Ser
485 490 495

Ser Asp Lys Met Cys Pro Pro Asp Glu Gly Trp Lys Arg His Gly Glu
500 505 510

eolf-seql.txt

Thr Cys Tyr Lys Ile Tyr Glu Asp Glu Val Pro Phe Gly Thr Asn Cys
515 520 525

Asn Leu Thr Ile Thr Ser Arg Phe Glu Gln Glu Tyr Leu Asn Asp Leu
530 535 540

Met Lys Lys Tyr Asp Lys Ser Leu Arg Lys Tyr Phe Trp Thr Gly Leu
545 550 555 560

Arg Asp Val Asp Ser Cys Gly Glu Tyr Asn Trp Ala Thr Val Gly Gly
565 570 575

Arg Arg Arg Ala Val Thr Phe Ser Asn Trp Asn Phe Leu Glu Pro Ala
580 585 590

Ser Pro Gly Gly Cys Val Ala Met Ser Thr Gly Lys Ser Val Gly Lys
595 600 605

Trp Glu Val Lys Asp Cys Arg Ser Phe Lys Ala Leu Ser Ile Cys Lys
610 615 620

Lys Met Ser Gly Pro Leu Gly Pro Glu Glu Ala Ser Pro Lys Pro Asp
625 630 635 640

Asp Pro Cys Pro Glu Gly Trp Gln Ser Phe Pro Ala Ser Leu Ser Cys
645 650 655

Tyr Lys Val Phe His Ala Glu Arg Ile Val Arg Lys Arg Asn Trp Glu
660 665 670

Glu Ala Glu Arg Phe Cys Gln Ala Leu Gly Ala His Leu Ser Ser Phe
675 680 685

Ser His Val Asp Glu Ile Lys Glu Phe Leu His Phe Leu Thr Asp Gln
690 695 700

Phe Ser Gly Gln His Trp Leu Trp Ile Gly Leu Asn Lys Arg Ser Pro
705 710 715 720

eolf-seql.txt

Asp Leu Gln Gly Ser Trp Gln Trp Ser Asp Arg Thr Pro Val Ser Thr
725 730 735

Ile Ile Met Pro Asn Glu Phe Gln Gln Asp Tyr Asp Ile Arg Asp Cys
740 745 750

Ala Ala Val Lys Val Phe His Arg Pro Trp Arg Arg Gly Trp His Phe
755 760 765

Tyr Asp Asp Arg Glu Phe Ile Tyr Leu Arg Pro Phe Ala Cys Asp Thr
770 775 780

Lys Leu Glu Trp Val Cys Gln Ile Pro Lys Gly Arg Thr Pro Lys Thr
785 790 795 800

Pro Asp Trp Tyr Asn Pro Asp Arg Ala Gly Ile His Gly Pro Pro Leu
805 810 815

Ile Ile Glu Gly Ser Glu Tyr Trp Phe Val Ala Asp Leu His Leu Asn
820 825 830

Tyr Glu Glu Ala Val Leu Tyr Cys Ala Ser Asn His Ser Phe Leu Ala
835 840 845

Thr Ile Thr Ser Phe Val Gly Leu Lys Ala Ile Lys Asn Lys Ile Ala
850 855 860

Asn Ile Ser Gly Asp Gly Gln Lys Trp Trp Ile Arg Ile Ser Glu Trp
865 870 875 880

Pro Ile Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro Trp His Arg Phe
885 890 895

Pro Val Thr Phe Gly Glu Glu Cys Leu Tyr Met Ser Ala Lys Thr Trp
900 905 910

Leu Ile Asp Leu Gly Lys Pro Thr Asp Cys Ser Thr Lys Leu Pro Phe
915 920 925

eolf-seql.txt

Ile Cys Glu Lys Tyr Asn Val Ser Ser Leu Glu Lys Tyr Ser Pro Asp
 930 935 940

Ser Ala Ala Lys Val Gln Cys Ser Glu Gln Trp Ile Pro Phe Gln Asn
 945 950 955 960

Lys Cys Phe Leu Lys Ile Lys Pro Val Ser Leu Thr Phe Ser Gln Ala
 965 970 975

Ser Asp Thr Cys His Ser Tyr Gly Gly Thr Leu Pro Ser Val Leu Ser
 980 985 990

Gln Ile Glu Gln Asp Phe Ile Thr Ser Leu Leu Pro Asp Met Glu Ala
 995 1000 1005

Thr Leu Trp Ile Gly Leu Arg Trp Thr Ala Tyr Glu Lys Ile Asn
 1010 1015 1020

Lys Trp Thr Asp Asn Arg Glu Leu Thr Tyr Ser Asn Phe His Pro
 1025 1030 1035

Leu Leu Val Ser Gly Arg Leu Arg Ile Pro Glu Asn Phe Phe Glu
 1040 1045 1050

Glu Glu Ser Arg Tyr His Cys Ala Leu Ile Leu Asn Leu Gln Lys
 1055 1060 1065

Ser Pro Phe Thr Gly Thr Trp Asn Phe Thr Ser Cys Ser Glu Arg
 1070 1075 1080

His Phe Val Ser Leu Cys Gln Lys Tyr Ser Glu Val Lys Ser Arg
 1085 1090 1095

Gln Thr Leu Gln Asn Ala Ser Glu Thr Val Lys Tyr Leu Asn Asn
 1100 1105 1110

Leu Tyr Lys Ile Ile Pro Lys Thr Leu Thr Trp His Ser Ala Lys
 1115 1120 1125

eolf-seql.txt

Arg	Glu	Cys	Leu	Lys	Ser	Asn	Met	Gln	Leu	Val	Ser	Ile	Thr	Asp
	1130					1135					1140			
Pro	Tyr	Gln	Gln	Ala	Phe	Leu	Ser	Val	Gln	Ala	Leu	Leu	His	Asn
	1145					1150					1155			
Ser	Ser	Leu	Trp	Ile	Gly	Leu	Phe	Ser	Gln	Asp	Asp	Glu	Leu	Asn
	1160					1165					1170			
Phe	Gly	Trp	Ser	Asp	Gly	Lys	Arg	Leu	His	Phe	Ser	Arg	Trp	Ala
	1175					1180					1185			
Glu	Thr	Asn	Gly	Gln	Leu	Glu	Asp	Cys	Val	Val	Leu	Asp	Thr	Asp
	1190					1195					1200			
Gly	Phe	Trp	Lys	Thr	Val	Asp	Cys	Asn	Asp	Asn	Gln	Pro	Gly	Ala
	1205					1210					1215			
Ile	Cys	Tyr	Tyr	Ser	Gly	Asn	Glu	Thr	Glu	Lys	Glu	Val	Lys	Pro
	1220					1225					1230			
Val	Asp	Ser	Val	Lys	Cys	Pro	Ser	Pro	Val	Leu	Asn	Thr	Pro	Trp
	1235					1240					1245			
Ile	Pro	Phe	Gln	Asn	Cys	Cys	Tyr	Asn	Phe	Ile	Ile	Thr	Lys	Asn
	1250					1255					1260			
Arg	His	Met	Ala	Thr	Thr	Gln	Asp	Glu	Val	His	Thr	Lys	Cys	Gln
	1265					1270					1275			
Lys	Leu	Asn	Pro	Lys	Ser	His	Ile	Leu	Ser	Ile	Arg	Asp	Glu	Lys
	1280					1285					1290			
Glu	Asn	Asn	Phe	Val	Leu	Glu	Gln	Leu	Leu	Tyr	Phe	Asn	Tyr	Met
	1295					1300					1305			
Ala	Ser	Trp	Val	Met	Leu	Gly	Ile	Thr	Tyr	Arg	Asn	Lys	Ser	Leu
	1310					1315					1320			

eolf-seql.txt

Met	Trp	Phe	Asp	Lys	Thr	Pro	Leu	Ser	Tyr	Thr	His	Trp	Arg	Ala
1325						1330					1335			
Gly	Arg	Pro	Thr	Ile	Lys	Asn	Glu	Lys	Phe	Leu	Ala	Gly	Leu	Ser
1340						1345					1350			
Thr	Asp	Gly	Phe	Trp	Asp	Ile	Gln	Thr	Phe	Lys	Val	Ile	Glu	Glu
1355						1360					1365			
Ala	Val	Tyr	Phe	His	Gln	His	Ser	Ile	Leu	Ala	Cys	Lys	Ile	Glu
1370						1375					1380			
Met	Val	Asp	Tyr	Lys	Glu	Glu	Tyr	Asn	Thr	Thr	Leu	Pro	Gln	Phe
1385						1390					1395			
Met	Pro	Tyr	Glu	Asp	Gly	Ile	Tyr	Ser	Val	Ile	Gln	Lys	Lys	Val
1400						1405					1410			
Thr	Trp	Tyr	Glu	Ala	Leu	Asn	Met	Cys	Ser	Gln	Ser	Gly	Gly	His
1415						1420					1425			
Leu	Ala	Ser	Val	His	Asn	Gln	Asn	Gly	Gln	Leu	Phe	Leu	Glu	Asp
1430						1435					1440			
Ile	Val	Lys	Arg	Asp	Gly	Phe	Pro	Leu	Trp	Val	Gly	Leu	Ser	Ser
1445						1450					1455			
His	Asp	Gly	Ser	Glu	Ser	Ser	Phe	Glu	Trp	Ser	Asp	Gly	Ser	Thr
1460						1465					1470			
Phe	Asp	Tyr	Ile	Pro	Trp	Lys	Gly	Gln	Thr	Ser	Pro	Gly	Asn	Cys
1475						1480					1485			
Val	Leu	Leu	Asp	Pro	Lys	Gly	Thr	Trp	Lys	His	Glu	Lys	Cys	Asn
1490						1495					1500			
Ser	Val	Lys	Asp	Gly	Ala	Ile	Cys	Tyr	Lys	Pro	Thr	Lys	Ser	Lys
1505						1510					1515			

eolf-seql.txt

Lys	Leu	Ser	Arg	Leu	Thr	Tyr	Ser	Ser	Arg	Cys	Pro	Ala	Ala	Lys
1520						1525					1530			
Glu	Asn	Gly	Ser	Arg	Trp	Ile	Gln	Tyr	Lys	Gly	His	Cys	Tyr	Lys
1535						1540					1545			
Ser	Asp	Gln	Ala	Leu	His	Ser	Phe	Ser	Glu	Ala	Lys	Lys	Leu	Cys
1550						1555					1560			
Ser	Lys	His	Asp	His	Ser	Ala	Thr	Ile	Val	Ser	Ile	Lys	Asp	Glu
1565						1570					1575			
Asp	Glu	Asn	Lys	Phe	Val	Ser	Arg	Leu	Met	Arg	Glu	Asn	Asn	Asn
1580						1585					1590			
Ile	Thr	Met	Arg	Val	Trp	Leu	Gly	Leu	Ser	Gln	His	Ser	Val	Asp
1595						1600					1605			
Gln	Ser	Trp	Ser	Trp	Leu	Asp	Gly	Ser	Glu	Val	Thr	Phe	Val	Lys
1610						1615					1620			
Trp	Glu	Asn	Lys	Ser	Lys	Ser	Gly	Val	Gly	Arg	Cys	Ser	Met	Leu
1625						1630					1635			
Ile	Ala	Ser	Asn	Glu	Thr	Trp	Lys	Lys	Val	Glu	Cys	Glu	His	Gly
1640						1645					1650			
Phe	Gly	Arg	Val	Val	Cys	Lys	Val	Pro	Leu	Gly	Pro	Asp	Tyr	Thr
1655						1660					1665			
Ala	Ile	Ala	Ile	Ile	Val	Ala	Thr	Leu	Ser	Ile	Leu	Val	Leu	Met
1670						1675					1680			
Gly	Gly	Leu	Ile	Trp	Phe	Leu	Phe	Gln	Arg	His	Arg	Leu	His	Leu
1685						1690					1695			
Ala	Gly	Phe	Ser	Ser	Val	Arg	Tyr	Ala	Gln	Gly	Val	Asn	Glu	Asp
1700						1705					1710			

eolf-seql.txt

Glu Ile Met Leu Pro Ser Phe His Asp
1715 1720

<210> 16
<211> 30
<212> PRT
<213> Human

<400> 16

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Tyr Ser
20 25 30

<210> 17
<211> 14
<212> PRT
<213> Human

<400> 17

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly
1 5 10

<210> 18
<211> 32
<212> PRT
<213> Human

<400> 18

Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Leu Tyr Leu Gln
1 5 10 15

Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys Thr Ile
20 25 30

<210> 19
<211> 11
<212> PRT
<213> Human

<400> 19

eolf-seql.txt

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
1 5 10

<210> 20
<211> 23
<212> PRT
<213> Human

<400> 20

Asp Val Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys
20

<210> 21
<211> 15
<212> PRT
<213> Human

<400> 21

Trp Tyr Gln Gln Arg Pro Gly Lys Ala Pro Asn Leu Leu Ile Tyr
1 5 10 15

<210> 22
<211> 32
<212> PRT
<213> Human

<400> 22

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Thr Ile Ser Thr Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
20 25 30

<210> 23
<211> 11
<212> PRT
<213> Human

<400> 23

eolf-seql.txt

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
1 5 10

<210> 24
<211> 10
<212> PRT
<213> Human

<400> 24

Trp Glu Val Lys Asp Cys Arg Ser Phe Lys
1 5 10

<210> 25
<211> 12
<212> PRT
<213> Human

<400> 25

Pro Ala Ser Leu Ser Cys Tyr Lys Val Phe His Ala
1 5 10

<210> 26
<211> 20
<212> PRT
<213> Human

<400> 26

Pro Trp Arg Arg Gly Trp His Phe Tyr Asp Asp Arg Glu Phe Ile Tyr
1 5 10 15

Leu Arg Pro Phe
20

<210> 27
<211> 19
<212> PRT
<213> Human

<400> 27

Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro Trp His Arg Phe Pro Val
1 5 10 15

eolf-seql.txt

Thr Phe Gly

<210> 28
 <211> 12
 <212> PRT
 <213> Human

<400> 28

Arg Glu Leu Thr Tyr Ser Asn Phe His Pro Leu Leu
 1 5 10

<210> 29
 <211> 17
 <212> PRT
 <213> Human

<400> 29

Phe Thr Ser Cys Ser Glu Arg His Phe Val Ser Leu Cys Gln Lys Tyr
 1 5 10 15

Ser

<210> 30
 <211> 12
 <212> PRT
 <213> Human

<400> 30

Thr Val Lys Tyr Leu Asn Asn Leu Tyr Lys Ile Ile
 1 5 10

<210> 31
 <211> 11
 <212> PRT
 <213> Human

<400> 31

Glu Ala Val Tyr Phe His Gln His Ser Ile Leu
 1 5 10

eolf-seql.txt

<210> 32
 <211> 11
 <212> PRT
 <213> Human

<400> 32

Lys Lys Leu Ser Arg Leu Thr Tyr Ser Ser Cys
 1 5 10

<210> 33
 <211> 20
 <212> PRT
 <213> Human

<400> 33

Asn Gly Ser Arg Trp Ile Gln Tyr Lys Gly His Cys Tyr Lys Ser Asp
 1 5 10 15

Gln Ala Leu His
 20

<210> 34
 <211> 25
 <212> PRT
 <213> Human

<400> 34

Ile Ser Glu Trp Pro Ile Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro
 1 5 10 15

Trp His Arg Phe Pro Val Thr Phe Gly
 20 25

<210> 35
 <211> 20
 <212> PRT
 <213> Human

<400> 35

Gln Thr Leu Gln Asn Ala Ser Glu Thr Val Lys Tyr Leu Asn Asn Leu
 1 5 10 15

eolf-seql.txt

Tyr Lys Ile Ile
20

<210> 36
<211> 10
<212> PRT
<213> Human

<400> 36

Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro
1 5 10

<210> 37
<211> 15
<212> PRT
<213> Human

<400> 37

Phe Thr Ser Cys Ser Glu Arg His Phe Val Ser Leu Cys Gln Lys
1 5 10 15

<210> 38
<211> 451
<212> PRT
<213> Artificial Sequence

<220>
<223> Rituximab VH sequence

<400> 38

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Asn Met His Trp Val Lys Gln Thr Pro Gly Arg Gly Leu Glu Trp Ile
35 40 45

Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr Asn Gln Lys Phe
50 55 60

eolf-seql.txt

Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Thr Tyr Tyr Gly Gly Asp Trp Tyr Phe Asn Val Trp Gly
100 105 110

Ala Gly Thr Thr Val Thr Val Ser Ala Ala Ser Thr Lys Gly Pro Ser
115 120 125

Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
130 135 140

Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
145 150 155 160

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
165 170 175

Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
180 185 190

Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
195 200 205

Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Ala Glu Pro Lys Ser Cys
210 215 220

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
225 230 235 240

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
245 250 255

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
260 265 270

eolf-seql.txt

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
275 280 285

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
290 295 300

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
305 310 315 320

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
325 330 335

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
340 345 350

Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser
355 360 365

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
370 375 380

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
385 390 395 400

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
405 410 415

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
420 425 430

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
435 440 445

Pro Gly Lys
450

<210> 39
<211> 213
<212> PRT

<213> Artificial Sequence

<220>

<223> Rituximab VL sequence

<400> 39

Gln Ile Val Leu Ser Gln Ser Pro Ala Ile Leu Ser Ala Ser Pro Gly
1 5 10 15

Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser Tyr Ile
20 25 30

His Trp Phe Gln Gln Lys Pro Gly Ser Ser Pro Lys Pro Trp Ile Tyr
35 40 45

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Val Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Arg Val Glu Ala Glu
65 70 75 80

Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Thr Ser Asn Pro Pro Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala

180

185

190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
 195 200 205

Asn Arg Gly Glu Cys
 210

<210> 40
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Rituximab VH_CDR1

<400> 40

Ser Tyr Asn Met His
 1 5

<210> 41
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Rituximab VH_CDR2

<400> 41

Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr Asn Gln Lys Phe Lys
 1 5 10 15

Gly

<210> 42
 <211> 12
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Rituximab VH_CDR3

<400> 42

eolf-seql.txt

Ser Thr Tyr Tyr Gly Gly Asp Trp Tyr Phe Asn Val
1 5 10

<210> 43
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Rituximab VL_CDR1

<400> 43

Arg Ala Ser Ser Ser Val Ser Tyr Ile His
1 5 10

<210> 44
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Rituximab VL_CDR2

<400> 44

Ala Thr Ser Asn Leu Ala Ser
1 5

<210> 45
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Rituximab VL_CDR3

<400> 45

Gln Gln Trp Thr Ser Asn Pro Pro Thr Phe
1 5 10

<210> 46
<211> 8
<212> PRT
<213> Homo sapiens

<400> 46

Gly Trp His Phe Tyr Asp Asp Arg
1 5

<210> 47

<211> 14

<212> PRT

<213> Homo sapiens

<400> 47

Ile Ser Glu Trp Pro Ile Asp Asp His Phe Thr Tyr Ser Arg
1 5 10

<210> 48

<211> 15

<212> PRT

<213> Homo sapiens

<400> 48

Phe Pro Val Thr Phe Gly Glu Glu Cys Leu Tyr Met Ser Ala Lys
1 5 10 15

<210> 49

<211> 15

<212> PRT

<213> Homo sapiens

<400> 49

Glu Leu Thr Tyr Ser Asn Phe His Pro Leu Leu Val Ser Gly Arg
1 5 10 15

<210> 50

<211> 8

<212> PRT

<213> Homo sapiens

<400> 50

His Phe Val Ser Leu Cys Gln Lys
1 5

<210> 51

<211> 11

<212> PRT
 <213> Homo sapiens

<400> 51

Gln Thr Leu Gln Asn Ala Ser Glu Thr Val Lys
 1 5 10

<210> 52
 <211> 4
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Linker

<400> 52

Gly Phe Leu Gly
 1

<210> 53
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 53

Trp Glu Val Lys Asp Cys Arg Ser
 1 5

<210> 54
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 54

Glu Val Lys Asp Cys Arg Ser Phe
 1 5

<210> 55

<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 55

Val Lys Asp Cys Arg Ser Phe Lys
1 5

<210> 56
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 56

Lys Asp Cys Arg Ser Phe Lys Ala
1 5

<210> 57
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 57

Asp Cys Arg Ser Phe Lys Ala Leu
1 5

<210> 58
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 58

Cys Arg Ser Phe Lys Ala Leu Ser

1 5

<210> 59
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 59

Arg Ser Phe Lys Ala Leu Ser Ile
1 5

<210> 60
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 60

Pro Ala Ser Leu Ser Cys Tyr Lys
1 5

<210> 61
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 61

Ala Ser Leu Ser Cys Tyr Lys Val
1 5

<210> 62
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 62

Ser Leu Ser Cys Tyr Lys Val Phe
1 5

<210> 63

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 63

Leu Ser Cys Tyr Lys Val Phe His
1 5

<210> 64

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 64

Ser Cys Tyr Lys Val Phe His Ala
1 5

<210> 65

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 65

Cys Tyr Lys Val Phe His Ala Glu
1 5

<210> 66

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 66

Tyr Lys Val Phe His Ala Glu Arg
1 5

<210> 67

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 67

Lys Val Phe His Ala Glu Arg Ile
1 5

<210> 68

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 68

Val Phe His Ala Glu Arg Ile Val
1 5

<210> 69

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 69

Phe His Ala Glu Arg Ile Val Arg
1 5

<210> 70
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 70

Asp Cys Ala Ala Val Lys Val Phe
 1 5

<210> 71
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 71

Ala Ala Val Lys Val Phe His Arg
 1 5

<210> 72
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 72

Val Lys Val Phe His Arg Pro Trp
 1 5

<210> 73
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 73

Val Phe His Arg Pro Trp Arg Arg
1 5

<210> 74
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 74

His Arg Pro Trp Arg Arg Gly Trp
1 5

<210> 75
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 75

Pro Trp Arg Arg Gly Trp His Phe
1 5

<210> 76
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 76

Arg Arg Gly Trp His Phe Tyr Asp
1 5

<210> 77
<211> 8
<212> PRT
<213> Artificial Sequence

<220>

<223> Peptide

<400> 77

Gly Trp His Phe Tyr Asp Asp Arg
1 5

<210> 78

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 78

His Phe Tyr Asp Asp Arg Glu Phe
1 5

<210> 79

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 79

Tyr Asp Asp Arg Glu Phe Ile Tyr
1 5

<210> 80

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 80

Asp Arg Glu Phe Ile Tyr Leu Arg
1 5

<210> 81

<211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 81

Glu Phe Ile Tyr Leu Arg Pro Phe
 1 5

<210> 82
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 82

Pro Ile Asp Asp His Phe Thr Tyr
 1 5

<210> 83
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 83

Ile Asp Asp His Phe Thr Tyr Ser
 1 5

<210> 84
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 84

Asp Asp His Phe Thr Tyr Ser Arg

1

5

<210> 85
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 85

Asp His Phe Thr Tyr Ser Arg Tyr
1 5

<210> 86
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 86

His Phe Thr Tyr Ser Arg Tyr Pro
1 5

<210> 87
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 87

Phe Thr Tyr Ser Arg Tyr Pro Trp
1 5

<210> 88
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 88

Thr Tyr Ser Arg Tyr Pro Trp His
1 5

<210> 89

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 89

Tyr Ser Arg Tyr Pro Trp His Arg
1 5

<210> 90

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 90

Ser Arg Tyr Pro Trp His Arg Phe
1 5

<210> 91

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 91

Arg Tyr Pro Trp His Arg Phe Pro
1 5

<210> 92

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 92

Tyr Pro Trp His Arg Phe Pro Val
1 5

<210> 93

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 93

Pro Trp His Arg Phe Pro Val Thr
1 5

<210> 94

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 94

Trp His Arg Phe Pro Val Thr Phe
1 5

<210> 95

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 95

His Arg Phe Pro Val Thr Phe Gly
1 5

<210> 96
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 96

Arg Glu Leu Thr Tyr Ser Asn Phe
1 5

<210> 97
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 97

Glu Leu Thr Tyr Ser Asn Phe His
1 5

<210> 98
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 98

Leu Thr Tyr Ser Asn Phe His Pro
1 5

<210> 99
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 99

Thr Tyr Ser Asn Phe His Pro Leu
1 5

<210> 100
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 100

Tyr Ser Asn Phe His Pro Leu Leu
1 5

<210> 101
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 101

Ser Asn Phe His Pro Leu Leu Val
1 5

<210> 102
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 102

Asn Phe His Pro Leu Leu Val Ser
1 5

<210> 103
<211> 8
<212> PRT
<213> Artificial Sequence

<220>

<223> Peptide

<400> 103

Phe His Pro Leu Leu Val Ser Gly
1 5

<210> 104

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 104

His Pro Leu Leu Val Ser Gly Arg
1 5

<210> 105

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 105

Pro Leu Leu Val Ser Gly Arg Leu
1 5

<210> 106

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 106

Leu Leu Val Ser Gly Arg Leu Arg
1 5

<210> 107

<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 107

Leu Val Ser Gly Arg Leu Arg Ile
1 5

<210> 108
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 108

Val Ser Gly Arg Leu Arg Ile Pro
1 5

<210> 109
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 109

Phe Thr Ser Cys Ser Glu Arg His
1 5

<210> 110
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 110

Thr Ser Cys Ser Glu Arg His Phe

1

5

<210> 111
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 111

Ser Cys Ser Glu Arg His Phe Val
1 5

<210> 112
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 112

Cys Ser Glu Arg His Phe Val Ser
1 5

<210> 113
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 113

Ser Glu Arg His Phe Val Ser Leu
1 5

<210> 114
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 114

Glu Arg His Phe Val Ser Leu Cys
1 5

<210> 115

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 115

Arg His Phe Val Ser Leu Cys Gln
1 5

<210> 116

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 116

His Phe Val Ser Leu Cys Gln Lys
1 5

<210> 117

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 117

Phe Val Ser Leu Cys Gln Lys Tyr
1 5

<210> 118

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 118

Val Ser Leu Cys Gln Lys Tyr Ser
1 5

<210> 119

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 119

Ser Leu Cys Gln Lys Tyr Ser Glu
1 5

<210> 120

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 120

Leu Cys Gln Lys Tyr Ser Glu Val
1 5

<210> 121

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 121

Cys Gln Lys Tyr Ser Glu Val Lys
1 5

<210> 122
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 122

Gln Lys Tyr Ser Glu Val Lys Ser
 1 5

<210> 123
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 123

Thr Val Lys Tyr Leu Asn Asn Leu
 1 5

<210> 124
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 124

Val Lys Tyr Leu Asn Asn Leu Tyr
 1 5

<210> 125
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 125

Lys Tyr Leu Asn Asn Leu Tyr Lys
1 5

<210> 126
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 126

Tyr Leu Asn Asn Leu Tyr Lys Ile
1 5

<210> 127
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 127

Leu Asn Asn Leu Tyr Lys Ile Ile
1 5

<210> 128
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 128

Asn Asn Leu Tyr Lys Ile Ile Pro
1 5

<210> 129
<211> 8
<212> PRT
<213> Artificial Sequence

<220>

<223> Peptide

<400> 129

Asn Leu Tyr Lys Ile Ile Pro Lys
1 5

<210> 130

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 130

Leu Tyr Lys Ile Ile Pro Lys Thr
1 5

<210> 131

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 131

Tyr Lys Ile Ile Pro Lys Thr Leu
1 5

<210> 132

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 132

Lys Ile Ile Pro Lys Thr Leu Thr
1 5

<210> 133

<211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 133

Ile Ile Pro Lys Thr Leu Thr Trp
 1 5

<210> 134
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 134

Ile Pro Lys Thr Leu Thr Trp His
 1 5

<210> 135
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 135

Pro Lys Thr Leu Thr Trp His Ser
 1 5

<210> 136
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 136

Lys Thr Leu Thr Trp His Ser Ala

1

5

<210> 137
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 137

Thr Leu Thr Trp His Ser Ala Lys
1 5

<210> 138
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 138

Glu Ala Val Tyr Phe His Gln His
1 5

<210> 139
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 139

Ala Val Tyr Phe His Gln His Ser
1 5

<210> 140
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 140

Val Tyr Phe His Gln His Ser Ile
1 5

<210> 141

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 141

Tyr Phe His Gln His Ser Ile Leu
1 5

<210> 142

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 142

Phe His Gln His Ser Ile Leu Ala
1 5

<210> 143

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 143

Lys Lys Leu Ser Arg Leu Thr Tyr
1 5

<210> 144

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 144

Lys Leu Ser Arg Leu Thr Tyr Ser
1 5

<210> 145

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 145

Leu Ser Arg Leu Thr Tyr Ser Ser
1 5

<210> 146

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 146

Ser Arg Leu Thr Tyr Ser Ser Arg
1 5

<210> 147

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 147

Arg Leu Thr Tyr Ser Ser Arg Cys
1 5

<210> 148
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 148

Asn Gly Ser Arg Trp Ile Gln Tyr
 1 5

<210> 149
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 149

Gly Ser Arg Trp Ile Gln Tyr Lys
 1 5

<210> 150
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 150

Ser Arg Trp Ile Gln Tyr Lys Gly
 1 5

<210> 151
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 151

Arg Trp Ile Gln Tyr Lys Gly His
1 5

<210> 152
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 152

Trp Ile Gln Tyr Lys Gly His Cys
1 5

<210> 153
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 153

Ile Gln Tyr Lys Gly His Cys Tyr
1 5

<210> 154
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 154

Gln Tyr Lys Gly His Cys Tyr Lys
1 5

<210> 155
<211> 8
<212> PRT
<213> Artificial Sequence

<220>

<223> Peptide

<400> 155

Tyr Lys Gly His Cys Tyr Lys Ser
1 5

<210> 156

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 156

Lys Gly His Cys Tyr Lys Ser Asp
1 5

<210> 157

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 157

Gly His Cys Tyr Lys Ser Asp Gln
1 5

<210> 158

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 158

His Cys Tyr Lys Ser Asp Gln Ala
1 5

<210> 159

<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 159

Cys Tyr Lys Ser Asp Gln Ala Leu
1 5

<210> 160
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 160

Tyr Lys Ser Asp Gln Ala Leu His
1 5

<210> 161
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 161

Lys Ser Asp Gln Ala Leu His Ser
1 5

<210> 162
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide

<400> 162

Ser Asp Gln Ala Leu His Ser Phe

1

5

<210> 163
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 163

Asp Gln Ala Leu His Ser Phe Ser
 1 5

<210> 164
 <211> 1722
 <212> PRT
 <213> Homo sapiens

<400> 164

Met Arg Thr Gly Trp Ala Thr Pro Arg Arg Pro Ala Gly Leu Leu Met
 1 5 10 15

Leu Leu Phe Trp Phe Phe Asp Leu Ala Glu Pro Ser Gly Arg Ala Ala
 20 25 30

Asn Asp Pro Phe Thr Ile Val His Gly Asn Thr Gly Lys Cys Ile Lys
 35 40 45

Pro Val Tyr Gly Trp Ile Val Ala Asp Asp Cys Asp Glu Thr Glu Asp
 50 55 60

Lys Leu Trp Lys Trp Val Ser Gln His Arg Leu Phe His Leu His Ser
 65 70 75 80

Gln Lys Cys Leu Gly Leu Asp Ile Thr Lys Ser Val Asn Glu Leu Arg
 85 90 95

Met Phe Ser Cys Asp Ser Ser Ala Met Leu Trp Trp Lys Cys Glu His
 100 105 110

eolf-seql.txt

His Ser Leu Tyr Gly Ala Ala Arg Tyr Arg Leu Ala Leu Lys Asp Gly
115 120 125

His Gly Thr Ala Ile Ser Asn Ala Ser Asp Val Trp Lys Lys Gly Gly
130 135 140

Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile Tyr Thr Arg
145 150 155 160

Asp Gly Asn Ser Tyr Gly Arg Pro Cys Glu Phe Pro Phe Leu Ile Asp
165 170 175

Gly Thr Trp His His Asp Cys Ile Leu Asp Glu Asp His Ser Gly Pro
180 185 190

Trp Cys Ala Thr Thr Leu Asn Tyr Glu Tyr Asp Arg Lys Trp Gly Ile
195 200 205

Cys Leu Lys Pro Glu Asn Gly Cys Glu Asp Asn Trp Glu Lys Asn Glu
210 215 220

Gln Phe Gly Ser Cys Tyr Gln Phe Asn Thr Gln Thr Ala Leu Ser Trp
225 230 235 240

Lys Glu Ala Tyr Val Ser Cys Gln Asn Gln Gly Ala Asp Leu Leu Ser
245 250 255

Ile Asn Ser Ala Ala Glu Leu Thr Tyr Leu Lys Glu Lys Glu Gly Ile
260 265 270

Ala Lys Ile Phe Trp Ile Gly Leu Asn Gln Leu Tyr Ser Ala Arg Gly
275 280 285

Trp Glu Trp Ser Asp His Lys Pro Leu Asn Phe Leu Asn Trp Asp Pro
290 295 300

Asp Arg Pro Ser Ala Pro Thr Ile Gly Gly Ser Ser Cys Ala Arg Met
305 310 315 320

eolf-seql.txt

Asp Ala Glu Ser Gly Leu Trp Gln Ser Phe Ser Cys Glu Ala Gln Leu
325 330 335

Pro Tyr Val Cys Arg Lys Pro Leu Asn Asn Thr Val Glu Leu Thr Asp
340 345 350

Val Trp Thr Tyr Ser Asp Thr Arg Cys Asp Ala Gly Trp Leu Pro Asn
355 360 365

Asn Gly Phe Cys Tyr Leu Leu Val Asn Glu Ser Asn Ser Trp Asp Lys
370 375 380

Ala His Ala Lys Cys Lys Ala Phe Ser Ser Asp Leu Ile Ser Ile His
385 390 395 400

Ser Leu Ala Asp Val Glu Val Val Val Thr Lys Leu His Asn Glu Asp
405 410 415

Ile Lys Glu Glu Val Trp Ile Gly Leu Lys Asn Ile Asn Ile Pro Thr
420 425 430

Leu Phe Gln Trp Ser Asp Gly Thr Glu Val Thr Leu Thr Tyr Trp Asp
435 440 445

Glu Asn Glu Pro Asn Val Pro Tyr Asn Lys Thr Pro Asn Cys Val Ser
450 455 460

Tyr Leu Gly Glu Leu Gly Gln Trp Lys Val Gln Ser Cys Glu Glu Lys
465 470 475 480

Leu Lys Tyr Val Cys Lys Arg Lys Gly Glu Lys Leu Asn Asp Ala Ser
485 490 495

Ser Asp Lys Met Cys Pro Pro Asp Glu Gly Trp Lys Arg His Gly Glu
500 505 510

Thr Cys Tyr Lys Ile Tyr Glu Asp Glu Val Pro Phe Gly Thr Asn Cys
515 520 525

eolf-seql.txt

Asn Leu Thr Ile Thr Ser Arg Phe Glu Gln Glu Tyr Leu Asn Asp Leu
530 535 540

Met Lys Lys Tyr Asp Lys Ser Leu Arg Lys Tyr Phe Trp Thr Gly Leu
545 550 555 560

Arg Asp Val Asp Ser Cys Gly Glu Tyr Asn Trp Ala Thr Val Gly Gly
565 570 575

Arg Arg Arg Ala Val Thr Phe Ser Asn Trp Asn Phe Leu Glu Pro Ala
580 585 590

Ser Pro Gly Gly Cys Val Ala Met Ser Thr Gly Lys Ser Val Gly Lys
595 600 605

Trp Glu Val Lys Asp Cys Arg Ser Phe Lys Ala Leu Ser Ile Cys Lys
610 615 620

Lys Met Ser Gly Pro Leu Gly Pro Glu Glu Ala Ser Pro Lys Pro Asp
625 630 635 640

Asp Pro Cys Pro Glu Gly Trp Gln Ser Phe Pro Ala Ser Leu Ser Cys
645 650 655

Tyr Lys Val Phe His Ala Glu Arg Ile Val Arg Lys Arg Asn Trp Glu
660 665 670

Glu Ala Glu Arg Phe Cys Gln Ala Leu Gly Ala His Leu Ser Ser Phe
675 680 685

Ser His Val Asp Glu Ile Lys Glu Phe Leu His Phe Leu Thr Asp Gln
690 695 700

Phe Ser Gly Gln His Trp Leu Trp Ile Gly Leu Asn Lys Arg Ser Pro
705 710 715 720

Asp Leu Gln Gly Ser Trp Gln Trp Ser Asp Arg Thr Pro Val Ser Thr
725 730 735

eolf-seql.txt

Ile Ile Met Pro Asn Glu Phe Gln Gln Asp Tyr Asp Ile Arg Asp Cys
740 745 750

Ala Ala Val Lys Val Phe His Arg Pro Trp Arg Arg Gly Trp His Phe
755 760 765

Tyr Asp Asp Arg Glu Phe Ile Tyr Leu Arg Pro Phe Ala Cys Asp Thr
770 775 780

Lys Leu Glu Trp Val Cys Gln Ile Pro Lys Gly Arg Thr Pro Lys Thr
785 790 795 800

Pro Asp Trp Tyr Asn Pro Asp Arg Ala Gly Ile His Gly Pro Pro Leu
805 810 815

Ile Ile Glu Gly Ser Glu Tyr Trp Phe Val Ala Asp Leu His Leu Asn
820 825 830

Tyr Glu Glu Ala Val Leu Tyr Cys Ala Ser Asn His Ser Phe Leu Ala
835 840 845

Thr Ile Thr Ser Phe Val Gly Leu Lys Ala Ile Lys Asn Lys Ile Ala
850 855 860

Asn Ile Ser Gly Asp Gly Gln Lys Trp Trp Ile Arg Ile Ser Glu Trp
865 870 875 880

Pro Ile Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro Trp His Arg Phe
885 890 895

Pro Val Thr Phe Gly Glu Glu Cys Leu Tyr Met Ser Ala Lys Thr Trp
900 905 910

Leu Ile Asp Leu Gly Lys Pro Thr Asp Cys Ser Thr Lys Leu Pro Phe
915 920 925

Ile Cys Glu Lys Tyr Asn Val Ser Ser Leu Glu Lys Tyr Ser Pro Asp
930 935 940

eolf-seql.txt

Ser Ala Ala Lys Val Gln Cys Ser Glu Gln Trp Ile Pro Phe Gln Asn
945 950 955 960

Lys Cys Phe Leu Lys Ile Lys Pro Val Ser Leu Thr Phe Ser Gln Ala
965 970 975

Ser Asp Thr Cys His Ser Tyr Gly Gly Thr Leu Pro Ser Val Leu Ser
980 985 990

Gln Ile Glu Gln Asp Phe Ile Thr Ser Leu Leu Pro Asp Met Glu Ala
995 1000 1005

Thr Leu Trp Ile Gly Leu Arg Trp Thr Ala Tyr Glu Lys Ile Asn
1010 1015 1020

Lys Trp Thr Asp Asn Arg Glu Leu Thr Tyr Ser Asn Phe His Pro
1025 1030 1035

Leu Leu Val Ser Gly Arg Leu Arg Ile Pro Glu Asn Phe Phe Glu
1040 1045 1050

Glu Glu Ser Arg Tyr His Cys Ala Leu Ile Leu Asn Leu Gln Lys
1055 1060 1065

Ser Pro Phe Thr Gly Thr Trp Asn Phe Thr Ser Cys Ser Glu Arg
1070 1075 1080

His Phe Val Ser Leu Cys Gln Lys Tyr Ser Glu Val Lys Ser Arg
1085 1090 1095

Gln Thr Leu Gln Asn Ala Ser Glu Thr Val Lys Tyr Leu Asn Asn
1100 1105 1110

Leu Tyr Lys Ile Ile Pro Lys Thr Leu Thr Trp His Ser Ala Lys
1115 1120 1125

Arg Glu Cys Leu Lys Ser Asn Met Gln Leu Val Ser Ile Thr Asp
1130 1135 1140

eolf-seql.txt

Pro	Tyr	Gln	Gln	Ala	Phe	Leu	Ser	Val	Gln	Ala	Leu	Leu	His	Asn
1145						1150					1155			
Ser	Ser	Leu	Trp	Ile	Gly	Leu	Phe	Ser	Gln	Asp	Asp	Glu	Leu	Asn
1160						1165					1170			
Phe	Gly	Trp	Ser	Asp	Gly	Lys	Arg	Leu	His	Phe	Ser	Arg	Trp	Ala
1175						1180					1185			
Glu	Thr	Asn	Gly	Gln	Leu	Glu	Asp	Cys	Val	Val	Leu	Asp	Thr	Asp
1190						1195					1200			
Gly	Phe	Trp	Lys	Thr	Val	Asp	Cys	Asn	Asp	Asn	Gln	Pro	Gly	Ala
1205						1210					1215			
Ile	Cys	Tyr	Tyr	Ser	Gly	Asn	Glu	Thr	Glu	Lys	Glu	Val	Lys	Pro
1220						1225					1230			
Val	Asp	Ser	Val	Lys	Cys	Pro	Ser	Pro	Val	Leu	Asn	Thr	Pro	Trp
1235						1240					1245			
Ile	Pro	Phe	Gln	Asn	Cys	Cys	Tyr	Asn	Phe	Ile	Ile	Thr	Lys	Asn
1250						1255					1260			
Arg	His	Met	Ala	Thr	Thr	Gln	Asp	Glu	Val	His	Thr	Lys	Cys	Gln
1265						1270					1275			
Lys	Leu	Asn	Pro	Lys	Ser	His	Ile	Leu	Ser	Ile	Arg	Asp	Glu	Lys
1280						1285					1290			
Glu	Asn	Asn	Phe	Val	Leu	Glu	Gln	Leu	Leu	Tyr	Phe	Asn	Tyr	Met
1295						1300					1305			
Ala	Ser	Trp	Val	Met	Leu	Gly	Ile	Thr	Tyr	Arg	Asn	Lys	Ser	Leu
1310						1315					1320			
Met	Trp	Phe	Asp	Lys	Thr	Pro	Leu	Ser	Tyr	Thr	His	Trp	Arg	Ala
1325						1330					1335			

eolf-seql.txt

Gly	Arg	Pro	Thr	Ile	Lys	Asn	Glu	Lys	Phe	Leu	Ala	Gly	Leu	Ser
1340						1345					1350			
Thr	Asp	Gly	Phe	Trp	Asp	Ile	Gln	Thr	Phe	Lys	Val	Ile	Glu	Glu
1355						1360					1365			
Ala	Val	Tyr	Phe	His	Gln	His	Ser	Ile	Leu	Ala	Cys	Lys	Ile	Glu
1370						1375					1380			
Met	Val	Asp	Tyr	Lys	Glu	Glu	Tyr	Asn	Thr	Thr	Leu	Pro	Gln	Phe
1385						1390					1395			
Met	Pro	Tyr	Glu	Asp	Gly	Ile	Tyr	Ser	Val	Ile	Gln	Lys	Lys	Val
1400						1405					1410			
Thr	Trp	Tyr	Glu	Ala	Leu	Asn	Met	Cys	Ser	Gln	Ser	Gly	Gly	His
1415						1420					1425			
Leu	Ala	Ser	Val	His	Asn	Gln	Asn	Gly	Gln	Leu	Phe	Leu	Glu	Asp
1430						1435					1440			
Ile	Val	Lys	Arg	Asp	Gly	Phe	Pro	Leu	Trp	Val	Gly	Leu	Ser	Ser
1445						1450					1455			
His	Asp	Gly	Ser	Glu	Ser	Ser	Phe	Glu	Trp	Ser	Asp	Gly	Ser	Thr
1460						1465					1470			
Phe	Asp	Tyr	Ile	Pro	Trp	Lys	Gly	Gln	Thr	Ser	Pro	Gly	Asn	Cys
1475						1480					1485			
Val	Leu	Leu	Asp	Pro	Lys	Gly	Thr	Trp	Lys	His	Glu	Lys	Cys	Asn
1490						1495					1500			
Ser	Val	Lys	Asp	Gly	Ala	Ile	Cys	Tyr	Lys	Pro	Thr	Lys	Ser	Lys
1505						1510					1515			
Lys	Leu	Ser	Arg	Leu	Thr	Tyr	Ser	Ser	Arg	Cys	Pro	Ala	Ala	Lys
1520						1525					1530			

eolf-seql.txt

Glu Asn Gly Ser Arg Trp Ile Gln Tyr Lys Gly His Cys Tyr Lys
1535 1540 1545

Ser Asp Gln Ala Leu His Ser Phe Ser Glu Ala Lys Lys Leu Cys
1550 1555 1560

Ser Lys His Asp His Ser Ala Thr Ile Val Ser Ile Lys Asp Glu
1565 1570 1575

Asp Glu Asn Lys Phe Val Ser Arg Leu Met Arg Glu Asn Asn Asn
1580 1585 1590

Ile Thr Met Arg Val Trp Leu Gly Leu Ser Gln His Ser Val Asp
1595 1600 1605

Gln Ser Trp Ser Trp Leu Asp Gly Ser Glu Val Thr Phe Val Lys
1610 1615 1620

Trp Glu Asn Lys Ser Lys Ser Gly Val Gly Arg Cys Ser Met Leu
1625 1630 1635

Ile Ala Ser Asn Glu Thr Trp Lys Lys Val Glu Cys Glu His Gly
1640 1645 1650

Phe Gly Arg Val Val Cys Lys Val Pro Leu Gly Pro Asp Tyr Thr
1655 1660 1665

Ala Ile Ala Ile Ile Val Ala Thr Leu Ser Ile Leu Val Leu Met
1670 1675 1680

Gly Gly Leu Ile Trp Phe Leu Phe Gln Arg His Arg Leu His Leu
1685 1690 1695

Ala Gly Phe Ser Ser Val Arg Tyr Ala Gln Gly Val Asn Glu Asp
1700 1705 1710

Glu Ile Met Leu Pro Ser Phe His Asp
1715 1720

eolf-seql.txt

<210> 165
 <211> 23
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 165

Cys Ile Lys Pro Val Tyr Gly Trp Ile Val Ala Asp Asp Cys Asp Glu
 1 5 10 15

Thr Glu Asp Lys Leu Trp Lys
 20

<210> 166
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 166

Cys Glu His His Ser Leu Tyr Gly Ala Ala Arg
 1 5 10

<210> 167
 <211> 34
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 167

Asp Gly His Gly Thr Ala Ile Ser Asn Ala Ser Asp Val Trp Lys Lys
 1 5 10 15

Gly Gly Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile Tyr
 20 25 30

Thr Arg

eolf-seql.txt

<210> 168
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 168

Lys Gly Gly Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile
 1 5 10 15

Tyr Thr Arg

<210> 169
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 169

Gly Gly Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile Tyr
 1 5 10 15

Thr Arg

<210> 170
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 170

Asp Gly His Gly Thr Ala Ile Ser Asn Ala Ser Asp Val Trp Lys
 1 5 10 15

eolf-seql.txt

<210> 171
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 171

Trp	Gly	Ile	Cys	Leu	Lys	Pro	Glu	Asn	Gly	Cys	Glu	Asp	Asn	Trp	Glu
1				5					10					15	

Lys

<210> 172
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 172

Ile	Phe	Trp	Ile	Gly	Leu	Asn	Gln	Leu	Tyr	Ser	Ala	Arg
1				5				10				

<210> 173
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 173

Leu	His	Asn	Glu	Asp	Ile	Lys	Glu	Glu	Val	Trp	Ile	Gly	Leu	Lys
1				5					10				15	

<210> 174
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Peptide

<400> 174

Thr	Pro	Asn	Cys	Val	Ser	Tyr	Leu	Gly	Glu	Leu	Gly	Gln	Trp	Lys
1				5					10					15

<210> 175

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 175

Tyr	Phe	Trp	Thr	Gly	Leu	Arg	Asp	Val	Asp	Ser	Cys	Gly	Glu	Tyr	Asn
1				5					10					15	

Trp	Ala	Thr	Val	Gly	Gly	Arg
			20			

<210> 176

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 176

Ser	Val	Gly	Lys	Trp	Glu	Val	Lys	Asp	Cys	Arg
1			5					10		

<210> 177

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 177

Trp	Glu	Val	Lys	Asp	Cys	Arg	Ser	Phe	Lys
1			5						10

<210> 178
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 178

Ala Leu Ser Ile Cys Lys Lys
 1 5

<210> 179
 <211> 12
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 179

Pro Ala Ser Leu Ser Cys Tyr Lys Val Phe His Ala
 1 5 10

<210> 180
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 180

Arg Asn Trp Glu Glu Ala Glu Arg
 1 5

<210> 181
 <211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

eolf-seql.txt

<400> 181

Arg	Ser	Pro	Asp	Leu	Gln	Gly	Ser	Trp	Gln	Trp	Ser	Asp	Arg	Thr	Pro
1				5					10					15	

Val	Ser	Thr	Ile	Ile	Met	Pro	Asn	Glu	Phe	Gln	Gln	Asp	Tyr	Asp	Ile
			20				25						30		

Arg

<210> 182

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 182

Thr	Pro	Val	Ser	Thr	Ile	Ile	Met	Pro	Asn	Glu	Phe	Gln	Gln	Asp	Tyr
1				5					10					15	

Asp Ile Arg

<210> 183

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 183

Pro	Trp	Arg	Arg	Gly	Trp	His	Phe	Tyr	Asp	Asp	Arg	Glu	Phe	Ile	Tyr
1				5					10					15	

Leu	Arg	Pro	Phe
			20

<210> 184

<211> 8

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 184

Gly Trp His Phe Tyr Asp Asp Arg
 1 5

<210> 185
 <211> 14
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 185

Ile Ser Glu Trp Pro Ile Asp Asp His Phe Thr Tyr Ser Arg
 1 5 10

<210> 186
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 186

Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro Trp His Arg Phe Pro Val
 1 5 10 15

Thr Phe Gly

<210> 187
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

eolf-seql.txt

<400> 187

Phe	Pro	Val	Thr	Phe	Gly	Glu	Glu	Cys	Leu	Tyr	Met	Ser	Ala	Lys	Thr
1				5					10					15	

Trp	Leu	Ile	Asp	Leu	Gly	Lys	Pro	Thr	Asp	Cys	Ser	Thr	Lys
			20					25					30

<210> 188

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 188

Thr	Trp	Leu	Ile	Asp	Leu	Gly	Lys	Pro	Thr	Asp	Cys	Ser	Thr	Lys
1				5					10					15

<210> 189

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 189

Tyr	Asn	Val	Ser	Ser	Leu	Glu	Lys
1				5			

<210> 190

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 190

Val	Gln	Cys	Ser	Glu	Gln	Trp	Ile	Pro	Phe	Gln	Asn
1				5					10		

<210> 191
 <211> 12
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 191

Arg Glu Leu Thr Tyr Ser Asn Phe His Pro Leu Leu
 1 5 10

<210> 192
 <211> 39
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 192

Glu Leu Thr Tyr Ser Asn Phe His Pro Leu Leu Val Ser Gly Arg Leu
 1 5 10 15

Arg Ile Pro Glu Asn Phe Phe Glu Glu Glu Ser Arg Tyr His Cys Ala
 20 25 30

Leu Ile Leu Asn Leu Gln Lys
 35

<210> 193
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 193

Tyr His Cys Ala Leu Ile Leu Asn Leu Gln Lys
 1 5 10

<210> 194
 <211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 194

Phe	Thr	Ser	Cys	Ser	Glu	Arg	His	Phe	Val	Ser	Leu	Cys	Gln	Lys	Tyr
1				5					10					15	

Ser

<210> 195
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 195

His	Phe	Val	Ser	Leu	Cys	Gln	Lys
1				5			

<210> 196
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 196

Gln	Thr	Leu	Gln	Asn	Ala	Ser	Glu	Thr	Val	Lys
1				5					10	

<210> 197
 <211> 12
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 197

Thr	Val	Lys	Tyr	Leu	Asn	Asn	Leu	Tyr	Lys	Ile	Ile
1				5					10		

<210> 198

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 198

Thr	Leu	Thr	Trp	His	Ser	Ala	Lys
1				5			

<210> 199

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 199

Asn	Arg	His	Met	Ala	Thr	Thr	Gln	Asp	Glu	Val	His	Thr	Lys
1				5					10				

<210> 200

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 200

Ser	His	Ile	Leu	Ser	Ile	Arg
1				5		

<210> 201

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 201

Ser	Leu	Met	Trp	Phe	Asp	Lys	Thr	Pro	Leu	Ser	Tyr	Thr	His	Trp	Arg
1				5					10					15	

<210> 202

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 202

Ser	Leu	Met	Trp	Phe	Asp	Lys
1				5		

<210> 203

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 203

Glu	Ala	Val	Tyr	Phe	His	Gln	His	Ser	Ile	Leu
1				5					10	

<210> 204

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Peptide

<400> 204

Lys	Lys	Leu	Ser	Arg	Leu	Thr	Tyr	Ser	Ser
1				5					10

eolf-seql.txt

<210> 205
 <211> 20
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 205

Asn	Gly	Ser	Arg	Trp	Ile	Gln	Tyr	Lys	Gly	His	Cys	Tyr	Lys	Ser	Asp
1				5					10					15	

Gln	Ala	Leu	His
			20

<210> 206
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 206

His	Asp	His	Ser	Ala	Thr	Ile	Val	Ser	Ile	Lys	Asp	Glu	Asp	Glu	Asn
1				5					10					15	

Lys	Phe	Val	Ser	Arg
				20

<210> 207
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide

<400> 207

Val	Glu	Cys	Glu	His	Gly	Phe	Gly	Arg
1				5				