

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46367 A1

(51) International Patent Classification⁷: **C11D 3/12, 3/40**

(74) Agent: **DICKSON, Elizabeth, Anne**; Reckitt Benckiser plc, Group Patents Department, Dansom Lane, Hull HU8 7DS (GB).

(21) International Application Number: PCT/GB00/04948

(22) International Filing Date:
21 December 2000 (21.12.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9930253.1 22 December 1999 (22.12.1999) GB
9930248.1 22 December 1999 (22.12.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
RECKITT BENCKISER (UK) LIMITED [GB/GB]; 67 Alma Road, Windsor, Berkshire SL4 3HD (GB).

Published:

— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FOX, Rod, Thomas** [GB/GB]; 30 South Street, Cottingham, Hull HU16 4AS (GB). **HARPER, Duncan, Roger** [GB/GB]; Flat 2, 78 The Boulevard, Hull HU3 2TA (GB).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHOTOCATALYTIC COMPOSITIONS AND METHODS

(57) Abstract: Cleaning compositions including a photocatalytic material and a sensitizer employ a photocatalytic material and a sensitizer at a locus, for example on a surface. The residue combats soils and/or undesired microorganisms at the locus.



WO 01/46367 A1

PHOTOCATALYTIC COMPOSITIONS AND METHODS

The present invention relates to photocatalytic compositions and in particular, but not exclusively, to photocatalytic cleaning compositions, intended to reduce the frequency and/or effort of cleaning; and to methods employing such compositions. References will be made herein to cleaning compositions and/or to compositions which are effective in combating malodours and/or soils and/or microorganisms, these being preferred compositions, but descriptions and definitions which follow are applicable also to compositions intended for other purposes.

Cleaning compositions of the invention are of particular interest for cleaning surfaces such as ceramic tiles, sinks, baths, washbasins, toilets, worksurfaces, ovens, hobs, carpets, fabrics, floors, painted woodwork, metalwork, laminates, glass surfaces and the like.

Cleaning compositions intended for general and for specific uses are well known in the art. Such compositions, when liquid, will commonly comprise one or more surfactants, to loosen and/or disperse oily deposits and to dissolve water soluble materials. These cleaning compositions may include one or more of solvents (including water), thickening agents, abrasive particles, bleaching agents, disinfectants/antibacterial agents, perfumes, waxes or other polishing agents, preservatives, colouring agents and like additives. The liquid formulation provides a vehicle for the removal of insoluble

particulate matter and builders and suspending agents are often included in the compositions to facilitate this process. These prior art compositions are, to a greater or lesser extent, effective in removing
5 soils, usually organic soils, from surfaces and in preventing their redeposition during the cleaning process. However, re-soiling of the surfaces after cleaning is an inevitable and continuous process.

10 Thus, domestic and other surfaces are continually dirtied or soiled by various means including, for example, soiling resulting from the preparation of food, contact by people and domestic pets, deposition of oily deposits and of airborne materials. Not only
15 are these and like soils aesthetically displeasing, they may also have deleterious effects on health. The soils may contain allergenic material such as pollen, dust mites, dust mite droppings, cat and other animal allergens and furthermore may include
20 harmful or toxic materials derived from adjacent or nearby industrial, horticultural or agricultural processes. Deposited soils may also harbour and give sustenance to pathogenic microorganisms or might include residues of human or animal faeces or urine.
25 It is therefore important that these and like deposited soils are removed from surfaces efficiently and frequently.

Cleaning of surfaces is therefore a frequent and
30 often time consuming requirement and is inevitably regarded as an unpleasant chore. There is a need for means to reduce the frequency of cleaning, and desirably also to facilitate the removal of soils

deposited on surfaces. It will be appreciated that known, conventional, cleaning compositions have no effect on soils deposited on the surfaces after the cleaning process until such time as the cleaning process is undertaken again. The present invention therefore seeks to provide cleaning compositions which, after the cleaning process, are effective to reduce the required frequency of cleaning and/or to facilitate the removal of deposited soils.

It is an object of embodiments of the invention to provide a composition showing improved photocatalytic action.

In accordance with a first aspect of the present invention there is provided a composition which comprises in admixture a photocatalytic material or a precursor to a photocatalytic material, and a sensitiser which acts to absorb visible or ultra violet or infra-red radiation and enhance the photocatalytic action of the photocatalytic material.

It is an object of embodiments of the invention to provide a cleaning composition which, in addition to combating existing malodours and/or soils and/or undesired microorganisms when applied to a locus, for example a surface, combats further malodour compounds and/or soils and/or undesired microorganisms, after its application to a locus.

In accordance with a second aspect of the present invention there is provided a composition comprising a photocatalytic material able to combat malodours

and/or soils and/or undesired microorganisms at a locus, or a precursor to such a photocatalytic material, and a sensitiser which acts to absorb visible or ultra-violet or infra-red radiation and
5 improve the efficacy of the photocatalytic material in combating malodours and/or soils and/or undesired microorganisms at the locus.

By "combat" we mean that the composition of the
10 second aspect can be used to remove and/or break down malodour compounds and/or soils and/or microorganisms at the locus and/or it can prevent malodours and/or soils and/or microorganisms from building up at the locus. The term "microorganism" is used in this
15 specification to denote any microscopic organism which is combatted; but especially a bacterium. Also of interest, however, as microorganisms which are prospectively combatted by compositions of the invention, are viruses and fungi, in particular
20 yeasts. One pathogenic microorganism which is of particular interest as demonstrating the efficacy of the compositions of the present invention is the bacterium *Staphylococcus aureus*.

25 Said composition of the second aspect includes deodorising compositions and anti-allergenic compositions. For example the compositions may have a deodorising effect, by breaking down odoriferous compounds, as deposits and/or as airborne compounds.
30 For such uses the compositions may be applied to surfaces in the appropriate location or may be used in room sprays.

By means of the present invention a residue or layer of photocatalytic material can be provided at a locus, for example on a surface whereby soils and/or undesired microorganisms deposited on the residue or layer or soils or undesired microorganisms which are present on the surface prior to deposition of the residue or layer are subject to a photocatalytic or other photochemical oxidation, reduction, free radical or other photochemical reaction effective to break down, "burn away" or otherwise decompose the soils or undesired microorganisms or at least major components thereof; and/or to weaken their contact with the surface. Consequently it may be said that the cleaning process continues after the conventional act of removal of the soil or undesired microorganisms is completed.

As noted above soils may contain allergenic material which is decomposed or otherwise degraded by means of the present invention. Of particular interest is the use of the compositions of the invention in combating allergenic soils associated with house dust mites.

It is believed that the faeces of two particular house dust mite species, *Dermatophagoides farinae* (known as Der-f) and *Dermatophagoides pteronyssinus* (known as Der-p) trigger the immune responses of the body, thereby giving rise to well known allergenic responses.

A review of this is given in Experimental and


Applied Acarology, 10 (1991) p. 167-186 in an article entitled "House dust-mite allergen", a review by L. G. Arlian.

5 Both the Der-f and Der-p species are found throughout the world. In some areas, Der-f will be the sole *Dermatophagoides* species. In other areas Der-p will be the sole species. In still other areas, the two species are both present through,
10 generally, one or the other will predominate.

Using the photocatalytic material, a decomposition reaction undergone by a malodour compound or a soil may involve photo-induced oxidation and/or photo-
15 induced reduction reactions with organic or inorganic components of the malodour compound or soil. These reactions may in turn result in the production of free radicals which are effective in breaking down organic matter in the malodour compound or soil.
20 These reactions may also provide an ongoing benefit after the initial deodorising or cleaning process has been completed.

One suitable photocatalytic material is titania and
25 a possible mode of action using titania is now described, and shown schematically below. Whilst we are not bound by any scientific theory, in this suggested mode of action, incident light of appropriate energy can promote an electron from a valence band of the titania to a conductance band.
30 There is then an electron (e^-) in the conductance band and a hole (h^+) in the valence band. Both the

electron and the hole may migrate to the surface of the titania particle and interact with oxygen and water to produce radical species. These radical species may then generate free radical decomposition reactions in the organic soil which may ultimately generate carbon dioxide if the free radical reaction continues to its conclusion. It is believed that the sensitiser is able to absorb light from the visible or ultra violet or infra-red (preferably the visible) region which causes an excitation of the sensitiser. Electrons are then emitted as the sensitiser decays or decomposes from the excited state, and these electrons are transferred to the conductance band of the photocatalytic material, such as titania.



The photocatalytic material in the compositions of the present invention preferably includes titania, zinc oxide or a combination of the two, and is preferably present in an amount of from 0.01% to 20%, especially 0.2% to 3%, and most preferably 0.3 to 1%, by weight of the composition. Titania is preferred as

the sole photocatalytic material. Most preferred is titania in anatase form, although the rutile form may be highly effective.

5 Preferably the photocatalytic material is imperceptible or almost imperceptible to the user after application. Preferably, the photocatalytic material used in the present invention is of a microscopic particle size. The microscopic particle
10 size also assists in achieving a uniform dispersion throughout the formulation and in maximising the efficiency of the photocatalytic reaction. Suitably the photocatalytic material has a mean particle size (diameter) of at least 5 nm, preferably at least 10
15 nm, most preferably at least 15 nm. Suitably the photocatalytic material has a mean particle size of less than 200 nm, preferably less than 100 nm. One especially preferred class of titania particles, made using the Woodhead process described later, has a
20 mean particle size in the range 5-30 nm. Another preferred class, being titania commercially available from Millenium Inorganic Chemicals, has a mean particle size in the range 30-100 nm.

25 The photocatalytic material may be doped with an additional element which has the effect of reducing the energy required to promote an electron of the photocatalytic material to the conductance band, leaving the corresponding hole in the valence band.


30

Preferably, the sensitiser is present in an amount up to 1%, more preferably up to 0.1%, still more preferably up to 0.02%, and yet more preferably up to

0.01%. Preferably it is present in an amount from 0.00001%, more preferably from 0.0001%.

In this invention the sensitiser preferably absorbs radiation of wavelength which is in the band 200-1200 nm, preferably 400-800 nm. Its absorbency peak within these bands may be narrow. Thus, it may typically absorb within a sub-band 50-200 nm in width.

There are many sensitisers which will improve the efficacy of the photocatalytic material. Examples may include cationic, anionic, nonionic and amphoteric dyes. Cationic dyes are one preferred class. Examples include the sensitisers described in US 5,200,292. Thus, suitable sensitisers include cationic dye/anionic borate dye complexes represented by the general formula (I):



wherein D^+ represents a cationic dye; and R_1 , R_2 , R_3 and R_4 , which may be the same or different, each represents an unsubstituted or substituted alkyl group, an unsubstituted or substituted aryl group, an unsubstituted or substituted aralkyl group, an unsubstituted or substituted alkenyl group, an unsubstituted or substituted alkynyl group, an unsubstituted or substituted alicyclic group, or an unsubstituted or substituted heterocyclic group.

Examples of the cationic sensitisers which can be used in the present invention include cyanine dyes and dyes comprising a cation portion such as a quaternary ammonium ion covalently bonded to other neutral sensitizer structures via a bonding group.

Cationic dye/borate anion complexes are known in the art. Examples of methods for the preparation of these complexes and the use of these complexes in an image formation system are described in U.S. Pat. Nos. 3,567,453, 4,307,182, 4,343,891, 4,447,521, 4,450,227, and 5,200,292.

For example the cationic dye/borate anion complex which can be used in the present invention can be prepared by allowing a borate salt and a sensitizer to react in a known counter ion exchange process. This process is further disclosed in Hishiki, Y., *Repts. Sci-Research Inst.* (1953), 29, pp 72 to 79. Examples of useful borate salts include sodium salts such as sodium tetraphenyl borate, sodium triphenyl butyl borate and sodium trianisyl butyl borate, and ammonium salts such as tetraethyl ammonium tetraphenyl borate.

25

Examples of useful cationic dyes to be used in the present invention include photo-reducible cationic dyes capable of forming a complex which is stable in a dark place with a borate anion, such as cationic methine, polymethine, triarylmethane, indoline, thiazine, xanthene, oxazine and acridine dyes. More particularly, these dyes are cationic, carbocyanine, hemicyanine, rhodamine and azomethine dyes.


30

Cationic cyanine dyes disclosed in US-A-3,495,987 and US-A-5,200,292 are believed to be useful in the present invention.

5

Specific examples of dyes believed useful include methylene blue, safarine O, malachite green, cyanine dyes of the general formula (II) below and rhodamine dyes of the general formula (III) below (e.g., Rhodamine B or Rhodamine 6G).

10



wherein n represents 0 or an integer of 1 to 3; R represents an alkyl group; and Y represents CH-CH, N-CH₃, C(CH₃)₂, O, S or Se.

15


In the general formula (II), R is preferably a lower alkyl group (preferably having 1 to 8 carbon atoms) or an alkyl group (preferably having 1 to 5 carbon atoms) substituted by at least one of a carboxyl group, a sulfo group (itself optionally substituted by, for example, a hydroxy group or a halogen atom), a hydroxyl group, a halogen atom, an alkoxy group having 1 to 4 carbon atoms (itself optionally substituted by, for example, one or more alkoxy groups having 1 to 4 carbon atoms or

20

25

sulfoalkoxy groups having 1 to 4 carbon atoms), a phenyl group or a substituted phenyl, for example, β -sulfoethyl, γ -sulfopropyl, γ -sulfobutyl, δ -sulfobutyl, 2-[2-(3-sulfopropoxy)ethoxy]ethyl,
 5 2-hydroxysulfopropyl, 2-chlorosulfopropyl, 2-methoxyethyl, 2-hydroxyethyl, carboxymethyl, 2-carboxyethyl, 2,2,3,3'-tetrafluoropropyl, 3,3,3-trifluoropropyl or trifluoroethyl.

10



wherein R' and R'' each represents a hydrogen atom, an alkyl group (preferably having 1 to 6 carbon atoms),
 15 an aryl group or combination thereof, for example, methyl, ethyl, propyl, butyl, pentyl, hexyl, phenyl or benzyl.

The borate anion used in the present invention is
 20 so designed that a borate radical produced by the transfer of an electron to a sensitiser upon exposure to light easily dissociates into a radical as follows:



For example, triphenylbutyl borate anion and trianisylbutyl borate anion easily dissociate into triphenyl boran or trianisyl boran and a butyl radical. Thus, these anions are particularly preferred anions. On the other hand, tetrabutyl borate anion does not easily dissociate probably because a tetrabutyl borate radical produced therefrom is so unstable that it accepts an electron from a sensitiser. Similarly, tetraphenyl borate anion functions poorly because it cannot easily produce a phenyl radical.

In the borate anion represented by the general formula (I), one or two of R_1 , R_2 , R_3 and R_4 are preferably alkyl groups. R_1 , R_2 , R_3 and R_4 each may contain 20 or less carbon atoms, preferably 1 to 7 carbon atoms. A preferable combination of R_1 , R_2 , R_3 and R_4 is one or more alkyl groups and one or more aryl groups, or one or more alkyl groups and one or more aralkyl groups. Particularly, a combination of three aryl groups and one alkyl group is most preferred.

Typical examples of alkyl groups represented by R_1 , R_2 , R_3 and R_4 include methyl, ethyl, propyl, butyl, pentyl, hexyl, octyl and stearyl groups. Such an alkyl group may be substituted by one or more halogen atoms, one or more cyano, acyloxy, acyl, alkoxy or hydroxy groups.

Typical examples of aryl groups represented by R_1 , R_2 , R_3 and R_4 include phenyl, naphthyl, and substituted aryl groups such as anisyl, and alkaryl such as methyl phenyl and dimethyl phenyl.

5

Typical examples of aralkyl groups represented by R_1 , R_2 , R_3 and R_4 include benzyl and phenethyl groups. Typical examples of alicyclic groups represented by R_1 , R_2 , R_3 and R_4 include cyclobutyl, cyclopentyl and cyclohexyl groups. Examples of unsubstituted alkynyl groups represented by R_1 , R_2 , R_3 and R_4 include propynyl and ethynyl groups. Examples of substituted alkynyl groups represented by R_1 , R_2 , R_3 and R_4 include a 3-chloropropynyl group. Examples of unsubstituted alkenyl groups represented by R_1 , R_2 , R_3 and R_4 include propenyl and vinyl groups. Examples of substituted alkenyl groups represented by R_1 , R_2 , R_3 and R_4 include 3-chloropropenyl and 2-chloroethenyl groups. Examples of unsubstituted heterocyclic groups represented by R_1 , R_2 , R_3 and R_4 include 3-thiophenyl and 4-pyridinyl groups. Examples of substituted heterocyclic groups represented by R_1 , R_2 , R_3 and R_4 include a 4-methyl-3-thiophenyl group.

Useful cationic dye/borate anion complexes may empirically confirmed. A combination of a cationic dye and a borate anion having a useful possibility can be fixed by Weller's equation (Rehm, D. and Weller, A., Isr. J. Chem., (1970), 8, pages 259 to 271). The equation can be simplified as follows:

$$\Delta G = E_{\text{ox}} - E_{\text{red}} - E_{\text{kv}}$$

wherein ΔG represents the change in Gibbs' free energy; E_{ox} represents the oxidation potential of borate anion BR_4''' -; E_{red} represents the reduction potential of an anionic sensitiser; and E_{kv} represents the energy of light used for the excitation of the sensitiser.





It is believed that a useful complex has a negative free energy change. Similarly, the difference in the reduction potential of the sensitiser and the oxidation potential of borate must be negative with respect to a complex which is stable in a dark place.

Namely,


$$E_{ox} - E_{red} > 0$$

As previously mentioned, this is a simplified equation and thus does not absolutely predict if a complex is useful in the present invention. There are many other factors which affect such a decision. One of these factors is the effect of the use of a monomer on a complex. It is known that if Weller's equation gives an excessive negative value, there can be some deviation from the equation. Thus, this equation is only a first approximation.


Particular examples of cationic dye/borate anion complexes believed useful in the present invention will be shown hereafter together with their peak absorbency wavelength, in nanometers (λ_{max}).

Complex	Structure	$\lambda_{\max}$
1	 $\text{Ph}_3\text{B}^- - n\text{-C}_4\text{H}_9$	552nm
2	 $\text{Ph}_3\text{B}^- - n\text{-C}_4\text{H}_9$	568nm
3	 $\text{Ph}_3\text{B}^- - n\text{-C}_4\text{H}_9$	492nm
4	 $\text{Ph}_3\text{B}^- - n\text{-C}_4\text{H}_9$	428nm


5




658nm



6




528nm




5

7




450nm



10


No.	R ¹	Ar
7A	n-Butyl	Phenyl
7B	n-Hexyl	Phenyl
7C	n-Butyl	Anisyl





550nm

 $\text{Ar}_3\text{B}^-\text{R}^1$


No.	R ²	R ¹	Ar
8A	Methyl	n-Butyl	Phenyl
8B	Methyl	n-Hexyl	Phenyl
8C	n-Butyl	n-Butyl	Phenyl
8D	n-Butyl	n-Hexyl	Phenyl
8E	n-Heptyl	n-Butyl	Phenyl
8F	n-Heptyl	n-Hexyl	Phenyl
8G	Ethyl	n-Butyl	Phenyl




570nm




590nm




11



640nm



No.	R ³	R ¹	Ar
11A	Methyl	n-Butyl	Phenyl
11B	Methyl	n-Hexyl	Phenyl
11C	n-Butyl	n-Butyl	Phenyl
11D	n-Butyl	n-Hexyl	Phenyl
11E	n-Pentyl	n-Butyl	Phenyl
11F	n-Pentyl	n-Hexyl	Phenyl
11G	n-Heptyl	n-Butyl	Phenyl
11H	n-Heptyl	n-Heptyl	Phenyl
11I	Methyl	n-Butyl	Anisyl



5

Other preferred sensitizers include the ruthenium sensitizers described in J.Am.Chem. Soc., Vol. 122, No. 12, 2000, pp. 2840-2849. These have three pairs of carboxylated bipyridyl groups complexed to a ruthenium (II) or ruthenium (III) atom. Two such complexes may be coupled together to make a polypyridine dyad, preferably an Ru(II) - Ru(III) polypyridine dyad.

10

Examples of preferred ruthenium sensitisers thus include the compounds:

5 ruthenium (III) bis-(4,4'-dicarboxyl-2,2'-bipyridine)-
(1,2-bis[4-(4'-methyl-2,2'-bipyridyl)]ethane)-
ruthenium(II) bis-(4,7-dimethyl-1,10-phenanthroline)



10 ruthenium (III) bis-(4,4'-dicarboxyl-2,2'-bipyridine)-
(1,2-bis[4-(4'-methyl-2,2'-bipyridyl)]ethane)-
ruthenium(II) bis-(2,2'-bipyridine)

ruthenium (II) bis-(4,4'-dicarboxyl-2,2'-bipyridine)-
(4,4'-dimethyl-2,2'-bipyridine)

15 ruthenium (II) tris-(4,4' -dicarboxyl-2,2'-bipyridine).

Other classes of sensitisers of interest for use with a photocatalytic material in the present invention the materials described in GB 1408144.
20 They include eosin, rose bengal, fluorescein, chlorophyll, metal-free porphyrin, sulphonated phthalocyanine and sulphonated zinc phthalocyanine.


Other classes of sensitisers of interest for use
25 with a photocatalytic material in the present invention include organosilicon (IV) phthalocyanines and naphthocyanines having Q-band absorption maxima at wavelengths greater than 660 nm, having the formula




wherein R^1 , R^2 , R^3 , R^4 , R^5 and R^6 units are each
5 independently selected from the group consisting of:

- a) hydrogen;
 b) halogen;
 c) hydroxyl;
 d) cyano;
 5 e) nitrilo;
 f) oximino;
 g) C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;
 h) halogen substituted C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;
 10 i) polyhydroxyl substituted C₃-C₂₂ alkyl;
 j) C₁-C₂₂ alkoxy;
 k) branched alkoxy having the formula:

15




or




wherein B is hydrogen, hydroxyl, C₁-C₃₀ alkyl, C₁-C₃₀ alkoxy, -CO₂H, -CH₂CO₂H, -SO₃⁻M⁺, -OSO₃⁻M⁺, -PO₃²⁻M, -OPO₃²⁻M, or mixtures thereof, M is a water soluble cation in sufficient amount to satisfy charge balance; x is 0 or 1, each y independently has the value from 0 to 6, each z independently has the value from 0 to 100;

- 1) substituted or unsubstituted aryl;
m) substituted or unsubstituted alkylenearyl;
n) substituted or unsubstituted aryloxy;
o) substituted or unsubstituted oxyalkylenearyl;
5 p) substituted or unsubstituted alkyleneoxyaryl;
q) C₁-C₂₂ thioalkyl, C₄-C₂₂ branched thioalkyl, or mixtures thereof;
r) an ester of the formula -CO₂R¹⁰ wherein R¹⁰ comprises:
- 10 i) C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;
ii) halogen substituted C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;
15 iii) polyhydroxyl substituted C₃-C₂₂ alkyl;
iv) C₃-C₂₂ glycol;
20 v) C₁-C₂₂ alkoxy;
vi) C₄-C₂₂ branched alkoxy;
vii) substituted or unsubstituted aryl;
viii) substituted or unsubstituted alkylaryl;
25 ix) substituted or unsubstituted aryloxy;
x) substituted or unsubstituted alkoxyaryl;
xi) substituted or unsubstituted alkyleneoxyaryl; or mixtures thereof,
30 s) an amino unit of the formula:



- wherein R^{11} and R^{12} comprises $\text{C}_1\text{-C}_{22}$ alkyl, $\text{C}_4\text{-C}_{22}$ branched alkyl, $\text{C}_2\text{-C}_{22}$ alkenyl, $\text{C}_4\text{-C}_{22}$ branched alkenyl, or mixtures thereof;
- 5 t) an alkylethyleneoxy unit of the formula:



- 10 wherein Z comprises:

- i) hydrogen;
- ii) hydroxyl;
- iii) $-\text{CO}_2\text{H}$;
- 15 iv) $-\text{SO}_3^-\text{M}^+$;
- v) $-\text{OSO}_3^-\text{M}^+$;
- vi) $\text{C}_3\text{-C}_6$ alkoxy;
- vii) substituted or unsubstituted aryl;
- viii) substituted or unsubstituted aryloxy;
- 20 ix) alkyleneamino; or mixtures thereof;

A units comprise nitrogen or oxygen, M is a water soluble cation, v is 0 or 1, x is from 0 to 100, y is from 0 to 12;

- u) and mixtures thereof;
- 25 axial R units wherein each R is independently selected from the group consisting of:


- a) hydrogen;
- b) cyano;
- c) nitrilo;
- 30 d) oximino;
- e) $\text{C}_1\text{-C}_{22}$ alkyl, $\text{C}_4\text{-C}_{22}$ branched alkyl, $\text{C}_2\text{-C}_{22}$ alkenyl, $\text{C}_4\text{-C}_{22}$ branched alkenyl, or mixtures thereof;
- f) halogen substituted $\text{C}_1\text{-C}_{22}$ alkyl, $\text{C}_4\text{-C}_{22}$ branched

alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;


g) polyhydroxyl substituted C₃-C₂₂ alkyl;

h) branched alkoxy having the formula:

5




or



wherein R is hydrogen, hydroxyl, C₁-C₃₀ alkyl, C₁-C₃₀ alkoxy, -CO₂H, -CH₂-CO₂H, -SO₃⁻M⁺, -OSO₃⁻M⁺, -PO₃²⁻M, -OPO₃²⁻M, or mixtures thereof; M is a water soluble cation in sufficient amount to satisfy charge balance; x is 0 or 1, each y independently has the value from 0 to 6, each z independently has the value from 0 to 100;

15

i) an alkyleneamino unit of the formula:



20

wherein R¹¹ and R¹² comprises C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;

R¹⁶ comprises:

- i) hydrogen;
- ii) C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;


5 A units comprise nitrogen or oxygen; X comprises chlorine, bromine, iodine or other water soluble anion, v is 0 or 1, u is from 0 to 22;

- j) an amino unit is of the formula:



wherein R¹¹ and R¹² comprises C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;

- 15 k) carboxylate of the formula:



wherein R¹⁰ comprises:

- 20 i) C₄-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₁-C₂₂ branched alkenyl, or mixtures thereof;
- ii) halogen substituted C₁-C₂₂ alkyl, C₄-C₂₂ branched alkyl, C₂-C₂₂ alkenyl, C₄-C₂₂ branched alkenyl, or mixtures thereof;
- 25 iii) poly-hydroxyl substituted C₃-C₂₂ alkyl;
- iv) C₃-C₂₂ glycol;
- v) C₁-C₂₂ alkoxy;
- vi) C₄-C₂₂ branched alkoxy;
- vii) substituted or unsubstituted aryl;
- 30 viii) substituted or unsubstituted alkylaryl;
- ix) substituted or unsubstituted aryloxy;

- x) substituted or unsubstituted alkoxyaryl;
 - xi) substituted or unsubstituted alkyleneoxyaryl;
 - xii) alkyleneamino, or mixtures thereof;
- l) and mixtures thereof.


5

Preferably in phthalocyanines (I above) each moiety R_1-R_4 is independently selected from hydrogen and C_{1-4} alkoxy, for example methoxy.


- 10 Preferably in the naphthocyanine (II above) each moiety R_1-R_6 is independently selected from hydrogen and halogen.

- 15 Preferably in phthalocyanine and naphthocyanine compounds the moieties R bonded to the central silicon atoms are polyhydroxyl substituted C_{3-22} alkylene moieties, preferably polyglycols of formula $-(CHOH)_nCH_2OH$, where n is 2-21, preferably 2-6, or branched alkoxy groups having the formula

20




or




where B is hydrogen, hydroxyl, C₁-C₃₀ alkyl, C₁-C₃₀ alkoxy, -CO₂H, -CH₂CO₂H, -SO₃⁻M⁺, -OSO₃⁻M⁺, -PO₃²⁻M, -OPO₃²⁻M, and mixtures thereof; M is a water soluble cation in sufficient amount to satisfy charge balance; x is 0 or 1, each y independently has the value from 0 to 6, preferably from 0 to 6; each z independently has the value from 0 to 100, preferably from 0 to about 10, more preferably from 0 to about 3.

Further information on these sensitisers may be found in US 5,916,481, the contents of which are incorporated herein by reference.

Further information about useful sensitisers is found in WO 98/32829. The sensitisers described therein could be used in the present invention, and the descriptions thereof are preferably incorporated herein by reference. Thus, they may suitably have the formula:



or the formula:



- 5 wherein M is a photoactive metal or non-metal having a valence greater than 3, rings A, B, C and D are aromatic rings, each of said rings being independently selected from the group consisting of benzene, 1,2-naphthalene, 2,3-naphthalene,
 10 anthracene, phenanthrene, and mixtures thereof.

Suitably rings A, B, C, and D are each independently:

- 15 i) a benzene ring unit, a 2,3-naphthylene ring unit, a 1,2-naphthylene ring unit, an anthracene ring unit or a phenanthrene ring unit, each such ring unit being fused to the pyrrole ring shown above, and each ring unit being substituted, possible substituents
 20 being as defined in WO 98/32829, or, preferably, unsubstituted.

Disclosures of similar sensitisers are given in WO 98/32826, the contents of which are also incorporated herein by reference.

5 In a third aspect of the present invention there is also provided a composition which comprises:

- a) a photocatalytic material able to combat malodours and/or soils and/or undesired microorganisms or a precursor to such a
10 photocatalytic material; and
- b) a sensitiser which is capable of absorbing radiation of a first wavelength from visible light, for example room lighting or daylight, consequently adopting an excited state, and relaxing from that
15 excited state by ejecting an electron, thereby enhancing the efficacy of the photocatalytic material against the malodours and/or soils and/or undesired microorganisms.

20 The compositions of the present invention may be provided in any appropriate dry or wet form such as, for example, a liquid, cream, mousse, emulsion, microemulsion, gel, powder or block. They may be dispensed in conventional manner directly from a
25 bottle or by means of, for example, a pump or a trigger spray or roller or an aerosol or, in the case of a powder, by a puffer or sprinkler. Also, they could be applied to a surface by a brush, dispensing stick, impregnated woven or non-woven cloth, or
30 sponge.

Liquid compositions are especially preferred, especially aqueous liquid compositions. Aqueous


liquid compositions can be emulsions, including microemulsions, and/or may contain solvents which solubilise those sensitisers which do not dissolve in a water phase. Liquid compositions could be supplied
5 ready-for-use or dilutable.

Whilst the person skilled in the art will be able to prepare aqueous and non-aqueous liquid formulations tailored to the above dispensing forms,
10 the compositions of the present invention generally comprise not more than 99.7%, preferably 75% to 95% water, and cationic, anionic, nonionic or amphoteric surfactants, or compatible combinations thereof, in an amount of 0.05% to 80%, typically 0.5% to 10%.
15 Surfactants should be selected having regard to the nature of the composition, in particular the photocatalytic agent or the precursor therefor, to ensure in-pack stability. In general, anionic surfactants are not suitable for incorporation in
20 acidic compositions, especially those containing titania. In general cationic surfactants are not suitable for incorporation in alkaline compositions, especially those containing titania. Nonionic surfactants are especially preferred in compositions
25 of the present invention.

Examples of nonionic surfactants which may be employed in the composition include those which are water soluble or water miscible and include but are
30 not limited to one or more of the following: amine oxides, block copolymers, alkoxylated alkanolamides, alkoxylated alcohols, alkoxylated alkyl phenols, and sorbitan esters, for example sorbitan mono oleate.

In each case the respective alkyl group is preferably a fatty alkyl group, suitably having from 7 to 24 carbon atoms, preferably 8 to 16, and may be branched or, more preferably, linear. Alkoxylate chains may be propoxylate chains, mixed ethoxylate/propoxylate chains or, most preferably, ethoxylate chains. Good examples include linear fatty alcohol ethoxylates (e.g. NEODOL, from Shell) and secondary fatty alcohol ethoxylates (e.g. TERGITOL, from Union Carbide). Other examples include alkoxylated octyl and nonyl phenols (e.g. IGEPAL, from Rhône-Poulenc).

Examples of cationic surfactants which may be used in the present invention include quaternary ammonium compounds and salts thereof, including quaternary ammonium compounds which also have germicidal activity and which may be characterized by the general structural formula:




when at least one of R_1 , R_2 , R_3 and R_4 is a hydrophobic, aliphatic, aryl aliphatic or aliphatic aryl group containing from 6 to 26 carbon atoms, and the entire cationic portion of the molecule has a molecular weight of at least 165. The hydrophobic groups may be long-chain alkyl, long-chain alkoxy

aryl, long-chain alkyl aryl, halogen-substituted long-chain alkyl aryl, long-chain alkyl phenoxy alkyl or aryl alkyl. The remaining groups on the nitrogen atoms, other than the hydrophobic radicals, are generally hydrocarbon groups usually containing a total of no more than 12 carbon atoms. The radicals R_1 , R_2 , R_3 and R_4 may be straight chain or may be branched, but are preferably straight chain, and may include one or more amide or ester linkages. The radical X may be any salt-forming anionic radical.

Examples of quaternary ammonium salts within the above description include the alkyl ammonium halides such as cetyl trimethyl ammonium bromide, alkyl aryl ammonium halides such as octadecyl dimethyl benzyl ammonium bromide, and N-alkyl pyridinium halides such as N-cetyl pyridinium bromide. Other suitable types of quaternary ammonium salts include those in which the molecule contains either amide or ester linkages, such as octyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride and N-laurylcocoaminoformylmethyl)-pyridinium chloride. Other effective types of quaternary ammonium compounds which are useful as germicides includes those in which the hydrophobic radical is characterized by a substituted aromatic nucleus as in the case of lauryloxyphenyltrimethyl ammonium chloride, cetylaminophenyltrimethyl ammonium methosulphate, dodecylphenyltrimethyl ammonium methosulphate, dodecylphenyltrimethyl ammonium chloride and chlorinated dodecylphenyltrimethyl ammonium chloride.

Preferred quaternary ammonium compounds which act as germicides and which are useful in the present invention include those which have the structural formula:

5



10

wherein R_2 and R_3 are the same or different C_8 - C_{12} alkyl, or R_2 is C_{12} - C_{16} alkyl, C_8 - C_{18} alkylethoxy, C_8 - C_{18} alkyl-phenoethoxy and R_2 is benzyl, and X is a halide, for example chloride, bromide or iodide, or methosulphate. The alkyl groups R_2 and R_3 may be straight chain or branched, but are preferably substantially linear.

15

20

A mixture of two or more surface active agents may also be used. Other known surface active agents not particularised above may also be used in some compositions; especially when one of them is a nonionic surfactant. Surface active agents in general are described in McCutcheon's Detergents and Emulsifiers, North American Edition, 1982; Kirk-Othmer, Encyclopaedia of Chemical Technology, 3rd Ed., Vol. 22, pp 346-387.

25

30

Grease cutting, adhesion promoting or other solvents may also be included generally in amounts of not more than 99%, typically not more than 50%.

Examples include glycols and glycol ethers.

Preferred ingredients of the composition are C₁₋₆ alkanols, for example ethanol and isopropanol. These may aid adhesion, promote soil removal and appear, surprisingly, to enhance photocatalytic activity. When present they preferably constitute 1-10%, preferably 2-5% of the composition, by weight.

Other ingredients of the compositions may include dispersing agents, suspending agents, colorants, fragrances, polishes, sequestrants, fabric softening agents, optical brighteners, laundry anti-fade agents, enzymes, thickeners, preservatives, bleaches, bleach activators, waxes, stabilising agents, propellants and further material(s) to combat undesired microorganisms. In particular variations of liquid compositions of the invention, some or all of the ingredients may be of high volatility whereby a residue of photocatalytic material can be left behind on a surface in a controlled manner.

Suitable dispersing agents may include hydroxyethyl cellulose, polyvinyl alcohol, polyvinyl acetate and ethylene oxide-propylene oxide block copolymers. Such agents may aid in-pack stability and promote good surface contact, on application.

Suitable adhesion promoters may include materials selected from polyvinyl alcohols, polyacrylic acids, ethylene oxide-propylene oxide block copolymers, hydroxyethyl celluloses, protein polymers and polysaccharide polymers. Preferred adhesion

promoters may include polyvinyl alcohols, alginates, gum arabic, and pectin.

5 Liquid compositions of the invention, ready for use, may be of pH in the range 1 to 13, preferably 2 to 12, most preferably 3 to 11. The pH may not be the same as that of as-supplied liquid compositions, because the latter may be diluted.

10 In one embodiment the composition is a bleaching composition containing a peroxygen compound, for example hydrogen peroxide or a generator thereof, or peracetic acid or persuccinic acid.

15 The components of the composition should be selected, and/or the composition formulated, such that the composition is stable for a sufficient period, without components being degraded or rendered unstable by the photocatalytic material and the
20 sensitiser. Alternatively, certain components could be kept apart from other components, for example in a twin pack formulation. Preferably the compositions are packaged for sale in containers which shield the
25 compositions from electromagnetic radiation of wavelength which would promote its photocatalytic action. All such measures are within the ordinary competence of persons skilled in the art.

30 Liquid compositions preferably have suitable rheology to suspend particles and/or to inhibit run off from upright surfaces, on application. To this end liquid compositions may be thixotropic, and preferably exhibit shear thinning with a suitable,

preferably low, yield point.

Preferred compositions of the invention are colloidal suspensions of photocatalytic particles, more preferably transition metal oxide particles, and most preferably titania particles.

Preferred colloidal suspensions of titania particles for use in the present invention are prepared by steps of hydrolysis of titanium tetrachloride in ammonium hydroxide, washing the precipitate thus formed, decreasing the pH to 3.3 by addition of a mineral acid, preferably nitric acid, washing until the conductivity drops below 500 μ S, and peptisation by addition of a mineral acid, preferably nitric acid, either at room temperature for 7 days or at 60-70°C for 30-90 minutes. The resultant colloidal suspension of titania typically has a titania concentration of about 10g/l and a mean particle size of about 20 nm. This method is known as the Woodhead method, after the inventor and patentee thereof.

Alternative colloidal suspensions of titania particles for use in the present invention may be prepared by the "isopropoxide" method. This method involves the steps of hydrolysis of titanium isopropoxide, suitably in ammonium hydroxide, washing the precipitate thus formed, filtration, and peptisation by addition of a mineral acid, preferably nitric acid, either at room temperature for 7 days or at 60-70°C for 30-90 minutes. The resultant colloidal suspension of titania typically has a

titania concentration of 25-30g/l and a mean particle size of about 20 nm, when the peptisation is at ambient temperature. When the peptisation is at the elevated temperature, the resultant colloidal suspension typically has a titania concentration in excess of 100g/l and a mean particle size of about 90-100 nm, but with a wide particle size distribution.

Further alternative colloidal suspensions of titania particles for use in the present invention may be prepared by the Kormann method. In this method titanium tetrachloride is hydrolysed at 0°C under a nitrogen blanket. Dialysis is carried out for 3-12 hours to remove undesired by-products of the hydrolysis. The resulting titania suspension is dried using a rotary evaporator, aided by a water bath held at 30°C. The resulting solid is re-suspended in deionised water. No peptisation step is required. The resulting colloidal suspension of titania typically has a titania concentration of about 1g/l and a mean particle size in the range 30-70 nm.

In accordance with a fourth aspect of the present invention there is provided a method of cleaning or sanitising a surface, the method comprising the steps of contacting the surface with a composition of the invention as defined above thereby depositing a residue of the photocatalytic material on the surface, and allowing the photocatalytic material to combat soils or undesired microorganisms present on or subsequently deposited on the surface.

In the case of soils the combating may be by catalysing or effecting an oxidation, reduction or other decomposition of the soils.

5

The method is suitably carried out with the surface and the composition at ambient temperature and without any subsequent heat treatment.

10

The method is suitably carried out under visible light of intensity at least 5,000 lux. Preferably the method is carried out under ambient light conditions, for example daylight and/or under room lighting.

15

Acidic conditions may be favoured for methods of cleaning or sanitising bathrooms and lavatories.

20

Alkaline conditions may be favoured for methods of cleaning or sanitising laundry and kitchen environments.

25

Neutral or near-neutral conditions may be favoured for methods of treating delicate fabrics and surfaces (for example marble, and certain painted surfaces).

30

The skilled person may consult readily available zeta potential plots for chosen photocatalytic materials in order to ascertain available and optimal ranges of surfactants. Furthermore, the skilled person may use dispersing agents to allow co-formulation of materials which may otherwise be incompatible.

The colloidal and interfacial nature of the photocatalytic material will determine the nature of the sensitisers, surfactants and other materials which can be employed to good effect, having regard to in-pack stability, surface coverage and adhesion and photocatalytic activity. In the case of any doubt, of course, trial and error can be used. However, by way of guidance we can make the following general statements.

Preferred acidic titania-containing compositions include a cationic and/or a nonionic surfactant; and preferably no anionic surfactant. A nonionic surfactant is in all cases a preferred constituent.

Preferred alkaline titania-containing compositions include an anionic and/or a nonionic surfactant; and preferably no cationic surfactant (in contrast, with certain mildly alkaline compositions containing zinc oxide cationic surfactants may also be used). A nonionic surfactant is in all cases a preferred constituent.

Neutral or near-neutral compositions may contain a surfactant of any type, and preferably include a nonionic surfactant.

The surfaces treated in the method may be hard surfaces, for example surfaces of wooden objects, tiles, sanitaryware, painted objects, panels, kitchen surfaces, worktops, walls, floors, windows, mirrors, shower cubicles and shower curtains, and cars. The

hard surfaces may be the surfaces of outdoor garden structures, for example greenhouses, outdoor furniture, patios and paths.

5 The surfaces treated in the method may be fibrous surfaces, for example clothes, furnishing fabrics and carpets.

10 As mentioned above and as is evidenced from the foregoing description and following examples our main interest is in providing a consumable surface cleaning composition which has, to paraphrase, a keep-clean or self-clean action. However, other compositions having a photocatalytic material or a
15 precursor to the photocatalytic material and a sensitiser in admixture are included in the scope of the invention. Such compositions may, for example, be permanently secured to the surface of a substrate, for example of ceramic, glass or plastics.
20 Securement may be by chemical bonding and/or a quasi-mechanical process, such as sputtering; or may be incorporated in an article, for example of ceramic, glass or plastics, during its manufacture. For example, the composition could be compounded with a
25 plastics material prior to its moulding or extrusion. Also covered are compositions to be added to water, to sanitise and/or decolorise it and/or to combat soils and/or microorganisms on surfaces in contact with the water.

30

 The following examples are illustrative of compositions according to the invention in the form of a liquid. They may all contain sensitisers,

colorants, fragrances and preservatives, preferably at concentrations not more than 1% each, with the balance of the formulations being titania and water.

5 All percentages in this specification are expressed in weight of component per total weight of composition (that is w/w) unless otherwise stated.

10 The invention will now be further described by way of example, with reference to the following non-limiting embodiments.

Unless otherwise stated the examples now described employ the sensitiser ruthenium (II) tris-(4,4' -
15 dicarboxyl-2,2'-bipyridine) mentioned above, hereinafter called "Sensitiser A", and having the CAS number CAS 97333-46-5. When acid blue colorant is mentioned it is the water soluble colorant known as acid blue F.Y.D.

20

EXAMPLE 1

Preparation of photocatalytic materials

25 In a series of "in-vitro" experiments the effects of pH as well as the addition of a surfactant were investigated for a particular titania/sensitiser colloidal suspension.

30 The colloidal suspension was obtained by precipitation of titanium isopropoxide via hydrolysis. The precipitate was then washed several times and filtered. The wet solid was peptised with

concentrated nitric acid (1M) and deionised water for one hour at 70°C to produce the suspension. The concentration of the resultant acidic titania (mainly anatase) suspension was greater than 100g/l. The
5 titania had a mean particle size of 95 nm.

The suspension was diluted to 1g/l for the experiment.

10 A non-acidic titania colloidal suspension containing polyvinyl alcohol (PVA, MW 15,000) was prepared as follows. PVA (0.10g, MW 15,000) was diluted in hot water then allowed to cool to room temperature. A known amount of the concentrated
15 titania colloidal suspension was mixed with the PVA/water with vigorous stirring, then rendered alkaline using sodium hydroxide, to a pH of 9.5-10.5. The volume was made up to 100 ml with deionised water. The final titania concentration was 1g/l.

20

Sensitiser preparation

Sensitiser A was dissolved in deionised water with the help of sonication, to make up a solution of
25 concentration 3.5×10^{-5} M.

Activity test

Acid system, no PVA

30

A mixture of the acidic titania colloidal suspension (1 ml) and Sensitiser A (4 ml) was stirred for about 1 minute using a rotary mixer. Gentian

violet (0.08 ml, 0.03%) was added to the mixture. pH was in the range 1.98.

Alkaline system, no PVA

5

A mixture of an alkaline titania colloidal suspension (10 ml) (described as above but without PVA) and Sensitiser A (4 ml) was mixed using a stirrer hotplate. The pH was increased with sodium hydroxide (0.1 M) to pH 10. Gentian violet (0.08 ml, 0.03%) was added to the mixture.

10

Acid system, with PVA

15

A mixture of the titania colloidal suspension containing PVA (1 ml) described above (but not made alkaline) and Sensitiser A (4 ml) was stirred for about 1 minute using a rotary mixer. Gentian violet (0.08 ml, 0.03%) was added to the mixture. The pH was 1.98.

20

Alkaline system, with PVA, (1)

An acidic mixture of titania colloidal suspension containing PVA (10 ml) and Sensitiser A (4 ml) was stirred using a stirrer hotplate. The pH was then increased with sodium hydroxide (0.1 M) to pH 10. Gentian violet (0.08 ml, 0.03%) was added to the mixture.

25

30

Alkaline system, with PVA (2)

5 A mixture of titania colloidal suspension
containing PVA at pH 10.03 ml (1 ml) and Sensitiser A
(4 ml) at pH 10.1 was stirred for about 1 minute
using a rotamixer. The pH (9.85) was slightly
increased with a sodium hydroxide solution (0.1 M) in
order to reach pH 10. Gentian violet (0.08 ml,
10 0.03%) was added to the mixture.

All samples were placed onto an overhead projector
(but at 2 cm above it in order to reduce heat).
UV/visible spectra were taken over a period of time.

15

Summary of Results

The acid system with no PVA decolorised the target
colorant gentian violet (0.08 ml, 0.03%) within 40
20 minutes. The presence of PVA slowed down the
decolorisation to 80 minutes. Both acid systems had
a precipitate. However, the target colorant was
still decolorised.

25 Activity in each of the alkaline systems was also
present but was found to be slower than in the acid
systems. After 4 hours there was still a minor
presence of the target colorant in each case.
However, there was no difference in activity between
30 PVA-containing systems (1) and (2), (1) being mixed
at acid pH and then having its pH raised to 10, and
(2) being wholly mixed under alkaline conditions.
Again, addition of PVA seemed to increase

decolorisation time.

Thus, preliminary indications are that acidic systems may offer the prospect of more rapid combating of soils and undesired microorganisms. Alkaline systems may offer the prospect of an extended period of activity. PVA, aside from benefits it may bring in surface deposition of the composition, may give prolonged activity, in both alkaline and acidic systems.

EXAMPLE 2 - colorant degradation in solution

A solution was made of 0.002% acid blue colorant in deionised water (1:4, v:v). A colloidal suspension of titania/sensitiser was added, such that the titania comprised 0.5% of the aqueous solution, and the sensitiser (Sensitiser A as used in Example 1) had a concentration of 6×10^{-6} M. The solution was mixed and poured into three glass vials. The pH of these was rendered, respectively, 2.5, 7.0 and 10.0 using sodium hydroxide as required. The samples were subjected to a light cabinet having D65 class bulbs to mimic daylight conditions at an illumination level of 2,500-3,000 lux. Colour intensity was measured by UV/visible spectrophotometry.

During the assessment the solution within each vial was subjected to magnetic stirring.

The sample at pH 2.5 was completely decolorised after 26 minutes.

After four hours the sample at pH 7.0 was put in a dark place for storage overnight. Some particulate matter settled overnight and the resulting clear solution was assessed by the UV/vis method, giving a
5 reading of 0.39.

After four hours the sample at pH 10.0 was also put in a dark place for storage overnight. In the morning it was slightly turbid so it was filtered.
10 The clear filtrate was assessed by the UV/vis method, giving a reading of 0.42.

A control sample with the same concentration of colorant but without the titania/sensitiser gave a
15 UV/vis reading of 0.6 after the same illumination/storage regime.

The titania used in this experiment was not prepared by the isopropoxide route described in
20 Example 1. The route used was the Woodhead route described earlier, involving hydrolysis of titanium tetrachloride, acidification, washing and peptisation. The concentration of titania in the resulting material was about 10 g/l and the mean
25 particle size was about 20 nm.

EXAMPLE 3

An anti-microbial evaluation was made of a
30 composition of the invention on surfaces against *Staphylococcus aureus*.

The following formulations were evaluated for antibacterial activity on surfaces.

Cleaning composition: 0.93% Titania/ 1.31×10^{-5} M,
Sensitiser A prepared as described in Example 2
above.

Control composition: As cleaning composition above
but titania/Sensitiser A.

Sterile deionised water: (inert control).

20 mm x 20 mm pieces of glazed tile were
sterilised. 0.1 ml of the respective composition was
applied to each tile and spread over the entire
surface. To allow the samples to dry, tiles were
incubated at 50°C for 15 minutes, re-spreading the
composition over the entire tile surface every 3
minutes. 5 tiles were required for each formulation,
one for each contact time (0, 1, 2, 3, 6 and 23
hours).

Treated, dried tiles were inoculated by spreading
10 μ L of a suspension of *Staphylococcus aureus*
(containing approx. 10^8 cfu/ml). After spreading the
innoculum over the 6 tiles, one tile was immediately
removed into a flask containing glass beads and
neutralising fluid, to allow bacterial recovery ($t =$
0). The number of viable bacteria recovered from the
tile was determined using pour plates prepared using
the neutralising fluid and suitable dilutions
thereof.

The remaining 4 tiles were placed in a sterile petri dish and incubated at approx. 25°C under a 50 W halogen lamp at a light intensity of 9,000-12,000 lux. The numbers of surviving bacteria on the test surface at each subsequent time-point was determined as described above. The microbiocidal effect (ME value) of each treatment was calculated as:

Log (surviving bacteria on water treated tile at t=0) minus log (surviving bacteria on test tile at a given contact time).

Table 1 - Antimicrobial activity on surfaces against *Staphylococcus aureus*

Sample	Mean ME values (n=3) at each contact time (hours)				
	0	1	2	4	6
Deionised Water	0	0.6	0.6	1.5	2.1
Control composition	0.1	0.5	1.5	2.0	2.3
Cleaning composition	0.1	5.4	5.5	5.5	5.5

Table 1 shows the means ME values for each treatment (n = 3). The cleaning composition of the invention shows benefits at all contact times.

When the antimicrobial evaluation was repeated with the inclusion of an organic soil, 5% (v/v) heat-inactivated horse serum, to represent "dirty"

conditions, a similar level of antimicrobial activity was demonstrated.

5 In similar experiments but using a lesser application of the composition of the invention and with the tiles kept in darkness, the antimicrobial activity was less.

10 **EXAMPLE 4 - Assessment of soap scum removal by QCM (quartz crystal microbalance)**

15 In this assessment the vibration frequency of a quartz crystal is altered as a function of soil removal from its surface. The technique allows soil removal to be sensitively monitored.

20 A quartz crystal was cleaned by dipping it in a hot solution of 50% chloroform/50% ethanol. Then the crystal was rinsed with ethanol and air dried. This cleaning procedure was then repeated 3 times. The crystal was then dipped once in a colloidal suspension made in accordance with the method of Example 2, having 0.5 wt% titania and 6×10^{-6} M of Sensitiser A, in deionised water. The crystal was 25 then oven dried at 50°C. A 0.045 wt% lime soap soil was then placed in a 450ml trigger spray bottle and sprayed once onto both sides of the crystal surface. The spray was held approx. 30cm above the crystal surface, and the spray was angled downwards at the 30 crystal surface. After being sprayed the crystal was again oven dried at 50°C.

The soap scum had been prepared using the following ingredients:

- Bar soap (Stearic acid based) 3.9%
- 5 Shampoo (moderate-cleaning type, without conditioner)
0.35%
- China clay powder 0.06%
- Artificial sebum 0.15%
- Hydrochloric acid (0.1M) to pH 3.3, and
- 10 Hard water (to 100%)

The hard water was made by mixing 100g calcium chloride dihydrate, 50g magnesium chloride hexahydrate and 4850g deionised water in a 5-litre screw top container.

15

The quartz crystal was illuminated by a halogen lamp giving a light intensity of 9,000 to 12,000 lux.

20 The frequency of vibration increased quickly over a period of 600 seconds and continued to increase more slowly thereafter, to a maximum value reached after 1200 seconds.

25 **EXAMPLE 5 - Action against colorant plus soil**

This procedure was created to monitor and evaluate the effects seen by a composition decomposing a lime soap soil and colorant on a hard surface (i.e. glass, ceramic tile, etc).

30

Preparation of the test surface

Using a tile cutter, tiles (approx. area 165 x 15cm)
5 were cut in half to make two pieces (15 x 7.5cm).
Each of these two tiles was then cut again widthwise
to produce smaller strips of area 7.5 x 1.5cm. Each
tile strip was then cleaned once with deionised
water, once with acetone, and then again with
10 deionised water. The tile strips were then dried in
the oven for 10 minutes, then wiped with a clean
paper tissue.

Reconstitutable soil preparation

15

The following ingredients were used to make the
soil:

Bar soap	3.9%
20 Shampoo	0.35%
China clay powder	0.06%
Artificial sebum	0.15%
Hard water	95.54%
(all as described in Example 4).	

25

The bar of soap was first shaved into a suitable
sized beaker. The remaining ingredients were added
to the soap in the above order, and stirred with a
four-blade propelled mixer. The mixture was then
30 warmed to 45-50°C and mixed until a smooth, lump free
suspension was achieved. This took about 2 hours.
The suspension was then filtered using a Buchner
funnel fitted with a Whatman No 1 filter paper. The

filtrate cake produced by filtration was then re-suspended in deionised water, using the same amount of water used in making the soil, and filtered again. The filtrate cake was uniformly dried in an oven overnight at 45°C. The dry cake was then pulverised, stored in a sealed container, and kept at 4-5°C until needed. This lime soap parent soil can be kept for up to 6 months at this temperature.

Reconstituted soil was then produced when needed in the following percentages:

Parent soil	0.045%
Hard water	9.00%
Hydrochloric acid 0.1M	0.77%
Deionised water	90.19%

The above ingredients were combined in a beaker. First the parent soil, hard water and deionised water were added together. The solution was mixed until uniform, then the hydrochloric acid was added. The suspension was next homogenised with a high shear mixer for 5 minutes then placed on a magnetic stirrer until needed.

The reconstituted soil was then placed in a 450ml trigger spray bottle and shaken well. The spray bottle was then held about 30 cm above the respective tile and the spray angled downwards. The tile surface was sprayed with the reconstituted soil 3 times, or more times if a thicker coat of soil was required. The tiles were then air dried on a flat

surface. Once the soil layer had completely dried, the surface was then treated with a colorant.

Application of a water-soluble blue colorant to the tile surface

0.06% of acid blue colorant was placed in a glass vial (approx. 2cm in width and 6cm in height), and filled until about one-third full. Each of the tiles was dipped once only into the vial of colorant, making sure that the bottom of the tile touched the base of the vial. The colorant should partly cover the bottom half of the tile. All of the treated tiles were immediately placed at a 45° angle (face of tile facing the side of the oven), into a 50°C oven for 10 minutes. Once the colorant had dried, excess colorant was washed off with a squirt of deionised water. The tiles were then placed back in the oven to dry.

Application of a photocatalytic composition to the tile surface

An acidic photocatalytic composition prepared by the method of Example 2 and also having 0.5% titania and 6×10^{-6} M of Sensitiser A was applied to the tile surfaces by wiping. 0.1 ml of the composition was placed in the centre of each tile. Carefully the composition was spread so that it covered approximately 2/3 of the tile area (approx. 5 x 1.5 cm). A folded tissue (approx. 5 x 2 cm) was used to wipe over the area with the composition, so that a film of liquid was left behind. The tiles were then

air dried for 5 minutes. This process was then repeated twice so that the tiles had 3 applications of the composition.

5 **Irradiation and measurement**

All of the tiles were photographed using a Digital Video Camcorder. A set of the treated tiles was left in a dark place as a control. All the other tiles
10 were then placed under an overhead projector (50,000 lux). Photographs of the tiles were taken at intervals until all of the blue colorant had disappeared and the time taken was noted.

15 **Results**

After approximately 2.5 hours under the 50,000 lux light the lime soap soil with the acid blue colorant on top had totally decomposed. It was noted that
20 after the blue colorant had disappeared a yellow residue was left behind. However after a further period of 4 hours under the 50,000 LUX source the yellow residue has also disappeared.

25 **EXAMPLE 6**

Tiles were prepared as described above in Example 5, bearing acid blue colorant, but not soil. They were then treated with the following compositions:

30

Composition A - 0.5% titania sol, containing 6×10^{-6} M of Dye A. The titania starting material was a 260 g/l colloidal titania suspension, mean particle

size about 95 nm, supplied by Millenium Inorganics, of Belgium. Composition A was diluted to a 10 g/l colloidal suspension, and was of pH 3.3, being adjusted thereto by 1M nitric acid.

5

Composition B - 0.5% titania colloidal solution, containing 6×10^{-6} M of Sensitiser A, made as described in Example 2 above.

10

Tiles were treated with Compositions A and B and were decolorised under illumination under an overhead projector (50,000 lux). They were then re-colored as before. After they were again decolorised they were again re-colored; and so on. The results are as follows:

15

Time (mins)	Composition A	Composition B
Time for 1 st decolorisation	84	35
Time for 2 nd decolorisation	29	12
Time for 3 rd decolorisation	21	10
Time for 4 th decolorisation	23	10
Time for 5 th decolorisation	22	6
Time for 6 th decolorisation	20	6

EXAMPLE 7 - Varying concentrations of sensitiser

The test described in Example 2, using the 50,000 lux illumination, was used for an acidic (pH 2.5) colloidal suspension of titania made as described in Example 2, and with 0.5% titania and concentration of the sensitiser up to 1×10^{-5} M. Measurements were taken using UV/visible spectrophotometry. The results are set out in the table below and show that activity is present at all sensitiser concentrations, with excellent activity at and above 3×10^{-5} M.

0.5 wt% titania measured after 4 minutes.

Sensitiser concentration ($\times 10^{-5}$ M)	Colorant decomposed (%)
0	5.6
0.065	27.3
0.300	64.0
0.600	82.1
0.800	91.1
1.000	93.9

0.5 wt% titania. Measured after 8 minutes

Sensitiser concentration ($\times 10^{-5}$ M)	Colorant decomposed (%)
0	6.5
0.065	47.4
0.300	98.9
0.600	100.0
0.800	99.7
1.000	100.0

5

EXAMPLE 8 - Varying concentrations of titania

Corresponding tests to those of Example 7 were carried out, but with the concentration of Sensitiser A fixed at 1.3×10^{-5} M, and with the titania concentration varying between 0.04% and 1.0%. The titania was prepared by the method described in Example 2, but the method employed hydrochloric acid. The results are set out in the table below.

15

Titania concentration %wt	Colorant decomposed (%)	
0.04	After 15 mins	8
	After 30 mins	13
	After 45 mins	32
	After 60 mins	46
0.2	After 15 mins	27
	After 30 mins	56
	After 45 mins	75
	After 60 mins	88
0.4	After 15 mins	27
	After 30 mins	56
	After 45 mins	75
	After 60 mins	89
1.0	After 15 mins	21
	After 30 mins	40
	After 45 mins	55
	After 60 mins	74

EXAMPLE 9 - removal of carpet stains

5

In these tests square carpet tiles (10 cm x 10 cm) of beige colour, polypropylene fibres and looped pile construction were stained with square blue stains of diameter approx 8 cm x 8 cm, sprayed through a mask onto the carpet until the stain was pronounced. The stains were applied using an aqueous solution of 0.06% acid blue colorant.

10

After drying, the tile was similarly treated with a colloidal suspension of 0.73% titania, prepared as

15

described in Example 2 above, and 8.5×10^{-6} M Sensitiser A, such that a central circular portion of the coloured area, 6 cm in diameter, had been thoroughly subjected to the spray. The tile was then
5 subjected to illumination from a halogen lamp (9,000 lux). Within 4 hours no trace of the stain could be seen.

Further work was carried out with the same titania
10 concentration but different amounts of Sensitiser A. Good results were also achieved with sensitiser concentrations of 5.7×10^{-6} M and 1.3×10^{-5} M. By visual assessment, weaker activity was achieved at sensitiser concentrations of 2.8×10^{-6} M and,
15 especially, 5.7×10^{-7} M. Using a control formulation with 0.73% titania and no sensitiser no observable stain removal was observed by eye.

20 **EXAMPLES 10-13**

Further compositions, which would employ a commercially available surfactant-stabilized colloidal aqueous solution of titania available as
25 260 g/l sols from Millenium Inorganics, may be blended at ambient temperature with the other materials described below, in water.

EXAMPLE 10

Ethoxylated alcohol	1%
Disodium cocoamphodiacetate	0.2%
POLYTERGENT noninonic surfactant	1%
Propylene glycol n-butyl ether	3%
Xanthan gum	1%
Titania (mainly anatase form)	0.5%
Sensitiser - cyanate	0.001%
dye/borate anion complex No. 1 defined above	

EXAMPLE 11

Nonionic surfactant	2%
Amphoteric surfactant	0.2%
Propylene glycol n-butyl ether	3%
Bentonite clay	1%
Zinc oxide	1.0%
Sensitiser - cyanate	0.002%
dye/borate anion complex No. 8E defined above	

5

The following example is illustrative of a composition according to the invention in the form of a cream. It may contain a colorant, fragrance and preservative at concentrations not more than 1% each with the balance of the formulation being water.

10

EXAMPLE 12

Chalk	9%
Titania	1%
Sodium lauryl sulphate (28% active content)	2%
Monoethanolamine	0.4%
Cyclodimethicone/dimethicone	9%
Poly dimethylsiloxane	0.5%
Sensitiser - cyanate dye/borate anion complex No. 10 defined above	0.0005%

5 The following example is illustrative of a composition according to the invention in the form of a mousse from an aerosol. It may contain a colorant, fragrance and preservative at a maximum concentration of 1% each with the balance of the formulation being water.

10

EXAMPLE 13

Aminomethylpropanol	0.5%
Ammonium hydroxide	0.5%
Butane	10%
Benzalkonium chloride	0.0-0.5%
Ethanol	2%
Morpholine	0.5%
PP3 methyl ether	8%
Trideceth 7	1%
Titania (mainly in anatase form)	1%
Sensitiser - cyanate dye/borate anion complex No. 8E defined above	0.001%

CLAIMS

1. A composition which comprises in admixture a photocatalytic material or a precursor to a photocatalytic material, and a sensitiser which acts to absorb visible or ultra violet or infra-red radiation and enhance the photocatalytic action of the photocatalytic material
2. A composition which includes a photocatalytic material able to combat soils and/or undesired microorganisms at a locus, or a precursor to such a photocatalytic material, and a sensitiser which acts to absorb visible or ultra violet or infra-red radiation and improve the efficacy of the photocatalytic material in combating soils and/or undesired microorganisms at the locus.
3. A composition according to claim 1 or 2, wherein the photocatalytic material is titania and/or zinc oxide.
4. A composition according to claim 3, wherein the photocatalytic material is titania.
5. A composition according to any preceding claim, wherein the photocatalytic material comprises particles having a size imperceptible or almost imperceptible to the eye when deposited on the surface.
6. A composition according to claim 5, wherein the particles have a mean size of from 5 nm to 100 nm.

7. A composition as claimed in any preceding claim, wherein the sensitiser constitutes up to 1% of the weight of the composition.
- 5
8. A composition according to any preceding claim, wherein the composition is a liquid composition which in use is applied to a surface and dries to leave the photocatalytic material and the sensitiser as a residue.
- 10
9. A composition according to claim 8, which composition comprises one or more surfactants.
- 15
10. A composition as claimed in claim 7 or 8, which composition comprises one or more ingredients selected from solvents, antimicrobial agents, suspending agents, stabilising agents, fragrances, colorants, bleaching agents, waxes, thickeners, preservatives and adhesion promoting agents.
- 20
11. A composition as claimed in any of claims 8 to 10, which composition contains a material selected from PVA, polyacrylic acid, an EO-PO block copolymer, a hydroxyethyl cellulose, a protein polymer or a polysaccharide polymer.
- 25
12. A composition which comprises:
- 30
- a) a photocatalytic material able to combat soils and/or undesired microorganisms or a precursor to such a photocatalytic material; and

b) a sensitiser which is capable of absorbing radiation of a first wavelength from visible light and consequently emitting radiation of a second wavelength which enhances the efficacy of the photocatalytic material against the soils and/or undesired microorganisms.

13. A method of cleaning and/or sanitising a surface comprising the steps of contacting the surface with a composition as claimed in any preceding claim thereby depositing a residue of the photocatalytic material on the surface, and allowing the photocatalytic material to combat soils and/or undesired microorganisms present or subsequently deposited on the surface.

14. A photocatalytic composition or method substantially as hereinbefore described with particular reference to the examples.

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/GB 00/04948

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D3/12 C11D3/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 376 288 A (FALHOLT PER ET AL) 27 December 1994 (1994-12-27) column 1, line 51-54 column 3, line 43,44 claims 1,7,9; example 10 ---	1-4,12, 14
X	WO 99 00477 A (PROCTER & GAMBLE (US)) 7 January 1999 (1999-01-07) page 21, line 11-28; examples ---	1-5, 7-12,14
X	US 4 174 292 A (MICHELOTTI FRANCIS W ET AL) 13 November 1979 (1979-11-13) column 3, line 17-65 --- -/--	1-4, 7-10,12, 14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

06/04/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pentek, E

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/GB 00/04948

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 209 863 A (DIXIT NAGARAJ S ET AL) 11 May 1993 (1993-05-11) column 11, line 21 -column 12, line 13; claims 1,10,12 -----	1-12,14
X	FR 2 729 673 A (RHONE POULENC CHIMIE) 26 July 1996 (1996-07-26) page 1, line 6 -page 10, line 34 -----	1-14
Y	-----	1-14
Y	WO 98 32829 A (PROCTER & GAMBLE (US)) 30 July 1998 (1998-07-30) cited in the application page 1, line 5 -page 2, line 10 -----	1-14
Y	GB 1 372 035 A (PROCTER & GAMBLE LTD) 30 October 1974 (1974-10-30) page 1, line 38 -page 2, line 23 -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/04948

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5376288	A	27-12-1994	AT 98295 T	15-12-1993
			AU 641672 B	30-09-1993
			AU 5960190 A	08-01-1991
			BR 9007470 A	16-06-1992
			DE 69005112 D	20-01-1994
			DE 69005112 T	14-04-1994
			WO 9015856 A	27-12-1990
			DK 478684 T	18-04-1994
			EG 19045 A	30-12-1994
			EP 0478684 A	08-04-1992
			ES 2046788 T	01-02-1994
			JP 4506229 T	29-10-1992
			MX 21242 A	01-11-1993
			NZ 234159 A	29-01-1992
			PT 94439 A, B	08-02-1991
			TR 27368 A	17-01-1995
WO 9900477	A	07-01-1999	BR 9810482 A	12-09-2000
			EP 0991748 A	12-04-2000
US 4174292	A	13-11-1979	BE 874870 A	02-07-1979
			CA 1120818 A	30-03-1982
			DE 2908736 A	27-09-1979
			FR 2420563 A	19-10-1979
			GB 2016508 A, B	26-09-1979
			IT 1114740 B	27-01-1986
			JP 54125191 A	28-09-1979
			MX 149508 A	16-11-1983
			NL 7901642 A	24-09-1979
US 5209863	A	11-05-1993	US 4968446 A	06-11-1990
			AU 662903 B	21-09-1995
			AU 1623692 A	13-05-1993
			CA 2069766 A	09-05-1993
			EP 0541202 A	12-05-1993
			FI 922509 A	09-05-1993
			GR 1001250 B	30-06-1993
			IE 921737 A	19-05-1993
			NO 922069 A	10-05-1993
			NZ 242844 A	27-04-1995
			PT 100541 A	29-10-1993
			TR 26040 A	01-11-1993
			AU 2461888 A	11-05-1989
			BR 8805765 A	25-07-1989
			CA 1326803 A	08-02-1994
			DK 618988 A	06-05-1989
			EP 0315024 A	10-05-1989
			FI 885113 A	06-05-1989
			IL 88166 A	21-06-1992
			JP 1230699 A	14-09-1989
			NO 173612 C	05-01-1994
			NZ 226709 A	26-10-1990
			PT 88929 B	29-01-1993
			PT 88929 A	01-12-1988
			US 5084198 A	28-01-1992
			US 5205954 A	27-04-1993
			ZA 8808078 A	27-06-1990

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/GB 00/04948

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2729673 A	26-07-1996	AU 4543796 A WO 9623051 A	14-08-1996 01-08-1996
WO 9832829 A	30-07-1998	BR 9807507 A CN 1250472 T EP 0970176 A ZA 9800529 A	21-03-2000 12-04-2000 12-01-2000 29-07-1998
GB 1372035 A	30-10-1974	AT 323303 B AU 460165 B AU 4216372 A BE 783317 A CA 998966 A CH 572974 A DE 2222829 A DE 2265710 C ES 402667 A FI 54339 B FR 2137741 A IE 36376 B IT 961167 B JP 52018724 B NL 7206405 A,B, SE 377816 B ZA 7203212 A	10-07-1975 17-04-1975 15-11-1973 10-11-1972 26-10-1976 27-02-1976 23-11-1972 01-07-1982 01-10-1975 31-07-1978 29-12-1972 27-10-1976 10-12-1973 24-05-1977 14-11-1972 28-07-1975 28-02-1973