



(86) Date de dépôt PCT/PCT Filing Date: 2002/02/28
(87) Date publication PCT/PCT Publication Date: 2002/10/24
(45) Date de délivrance/Issue Date: 2010/11/09
(85) Entrée phase nationale/National Entry: 2003/08/12
(86) N° demande PCT/PCT Application No.: US 2002/006240
(87) N° publication PCT/PCT Publication No.: 2002/083079
(30) Priorités/Priorities: 2001/04/17 (US60/284,606);
2001/06/22 (US09/887,281)

(51) Cl.Int./Int.Cl. *A61K 31/167* (2006.01),
A61K 31/135 (2006.01), *A61K 31/16* (2006.01),
A61K 31/165 (2006.01), *A61K 31/35* (2006.01),
A61K 31/375 (2006.01), *A61K 31/444* (2006.01),
A61K 31/46 (2006.01), *A61K 31/56* (2006.01),
A61K 31/58 (2006.01), *A61K 39/395* (2006.01),
A61K 45/00 (2006.01), *A61K 47/04* (2006.01),
A61K 47/10 (2006.01), *A61K 47/12* (2006.01),
A61K 47/16 (2006.01), *A61K 47/18* (2006.01),
A61K 47/22 (2006.01), *A61K 47/26* (2006.01),
A61K 47/38 (2006.01), *A61K 9/12* (2006.01),
A61L 9/04 (2006.01), *A61P 11/06* (2006.01),
A61P 11/08 (2006.01)

(72) Inventeurs/Inventors:

(54) Titre : COMPOSITIONS BRONCHODILATATRICES ET METHODES ASSOCIEES
(54) Title: AEROSOL COMPOSITIONS CONTAINING FORMOTEROL FOR DELIVERY TO THE LUNGS VIA
NEBULIZATION

(57) Abrégé/Abstract:

Bronchodilating compositions and methods are provided. The compositions are intended for administration as a nebulized aerosol. In certain embodiments, the compositions contain formoterol, or a derivative thereof. Methods for treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders using the compositions provided herein are also provided.



(72) Inventeurs(suite)/Inventors(continued): BANERJEE, PARTHA S., US; PHAM, STEPHEN, US; CHAUDRY, IMTIAZ A., US

(73) Propriétaires(suite)/Owners(continued): DEY, L.P., US

(74) Agent: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 October 2002 (24.10.2002)

PCT

(10) International Publication Number
WO 02/083079 A3

(51) International Patent Classification⁷: **A61K 31/167**,
31/58, 31/56, A61P 11/08

SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/US02/06240

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 28 February 2002 (28.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/284,606	17 April 2001 (17.04.2001)	US
09/887,281	22 June 2001 (22.06.2001)	US

Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/284,606 (CON)
Filed on	17 April 2001 (17.04.2001)
US	09/887,281 (CON)
Filed on	22 June 2001 (22.06.2001)

(71) Applicant (*for all designated States except US*): **DEY, L.P.** [US/US]; 2751 Napa Valley Corporate Drive, Napa, CA 94558 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BANERJEE, Partha, S.** [IN/US]; 108 Rosario Court, San Ramon, CA 94583 (US). **PHAM, Stephen** [US/US]; 7950 Pocket Road, #40, Sacramento, CA 95831 (US). **AKAPO, Samuel, O.** [NG/US]; 293 Shasta Drive, #115, Vacaville, CA 95687 (US). **CHAUDRY, Imtiaz, A.** [US/US]; 76 Golden Gate Circle, Napa, CA 94558 (US).

(74) Agents: **SEIDMAN, Stephanie, L.** et al.; Heller Ehrman White & McAuliffe LLP, 4350 La Jolla Village Drive, 6th floor, San Diego, CA 92122-1246 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— with international search report

[Continued on next page]

(54) Title: AEROSOL COMPOSITIONS CONTAINING FORMOTEROL FOR DELIVERY TO THE LUNGS VIA NEBULIZATION

(57) Abstract: Bronchodilating compositions and methods are provided. The compositions are intended for administration as a nebulized aerosol. In certain embodiments, the compositions contain formoterol, or a derivative thereof. Methods for treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders using the compositions provided herein are also provided.



WO 02/083079 A3

WO 02/083079 A3



(88) Date of publication of the international search report:
13 February 2003

(15) Information about Correction:
see PCT Gazette No. 15/2003 of 10 April 2003, Section II

(48) Date of publication of this corrected version:
10 April 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

50728-5

-1-

AEROSOL COMPOSITIONS CONTAINING FORMOTEROL FOR DELIVERY TO THE LUNGS VIA NEBULIZATION

FIELD OF THE INVENTION

Compositions and methods are provided relating to treatment, prevention, or amelioration of one or more symptoms of broncho-
5 constrictive disorders. In particular, the compositions and methods herein include formoterol, and/or derivatives thereof. The compositions are propellant-free, sterile unit dose or multidose inhalation solutions intended for administration via nebulization.

BACKGROUND OF THE INVENTION

10 Bronchoconstrictive disorders affect millions worldwide. Such disorders include asthma (including bronchial asthma, allergic asthma and intrinsic asthma, e.g., late asthma and airway hyper-responsiveness), chronic bronchitis and other chronic obstructive pulmonary diseases. Compounds having β_2 -adrenoreceptor agonist activity have been
15 developed to treat these conditions. Such compounds include, but are not limited to, Albuterol (α^1 -(((1,1-dimethylethyl)amino)methyl)-4-hydroxy-1,3-benzenedimethanol); Bambuterol (dimethylcarbamic acid 5-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-phenylene ester); Bitolterol (4-methylbenzoic acid 4-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,2-
20 phenylene ester); Broxaterol (3-bromo- α -(((1,1-dimethylethyl)amino)-methyl)-5-isoxazolemethanol); Isoproterenol (4-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,2-benzenediol); Trimetoquinol (1,2,3,4-tetrahydro-1-((3,4,5-trimethoxyphenyl)methyl)-6,7-isoquinolinediol); Clenbuterol (4-

-2-

amino-3,5-dichloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); Fenoterol (5-(1-hydroxy-2-((2-(4-hydroxyphenyl)-1-methylethyl)-amino)ethyl)-1,3-benzenediol); Formoterol (2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide);

5 (R,R)-Formoterol; Desformoterol ((R,R) or (S,S)-3-amino-4-hydroxy- α -(((2-(4-methoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol); Hexoprenaline (4,4'-(1,6-hexanediyl)-bis(imino(1-hydroxy-2,1-ethanediyl)))bis-1,2-benzenediol); Isoetharine (4-(1-hydroxy-2-((1-methylethyl)amino)-butyl)-1,2-benzenediol); Isoprenaline (4-(1-hydroxy-2-((1-methylethyl)-

10 amino)ethyl)-1,2-benzenediol); Metaproterenol (5-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,3-benzenediol); Picumeterol (4-amino-3,5-dichloro- α -(((6-(2-(2-pyridinyl)ethoxy)hexyl)amino)methyl)benzenemethanol); Pirbuterol (α^6 -(((1,1-dimethylethyl)amino)methyl)-3-hydroxy-2,6-pyridine-methanol); Procaterol (((R*,S*)-($\pm$)-8-hydroxy-5-(1-hydroxy-2-((1-

15 methylethyl)amino)butyl)-2(1H)-quinolinone); Reproterol ((7-(3-((2-(3,5-dihydroxyphenyl)-2-hydroxyethyl)amino)propyl)-3,7-dihydro-1,3-dimethyl-1H-purine-2,6-dione); Rimiterol (4-(hydroxy-2-piperidinylmethyl)-1,2-benzenediol); Salbutamol (($\pm$)- α^1 -(((1,1-dimethylethyl)amino)methyl)-4-

20 hydroxy-1,3-benzenedimethanol); (R)-Salbutamol; Salmeterol (($\pm$)-4-hydroxy- α^1 -(((6-(4-phenylbutoxy)hexyl)amino)methyl)-1,3-benzenedimethanol); (R)-Salmeterol; Terbutaline (5-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-benzenediol); Tulobuterol (2-chloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); and TA-2005 (8-hydroxy-5-((1R)-

25 1-hydroxy-2-(N-((1R)-2-(4-methoxyphenyl)-1-methylethyl)amino)ethyl)-carbostyryl hydrochloride).

These compounds are typically formulated for inhalation therapy. Aqueous or liquid formulations are preferred over solid formulations. Powdered formulations are more difficult to administer, particularly to the young and elderly who are most often the patients in need of such

-3-

therapy. Compounds, such as formoterol, which has many desirable properties, are not adequately stable in aqueous solutions to be formulated as liquids. Hence there is a need for formulations of compounds, such as formoterol, in a form that can be conveniently
5 administered and that are stable for extended periods of time. Therefore, it is an object herein to provide liquid formulations of β_2 -adrenoreceptor agonist compounds. It is also an object herein to provide more stable formulations of others of these compounds.

SUMMARY OF THE INVENTION

10 Compositions and methods for treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders are provided. The compositions provided herein are stable solutions of a bronchodilating agent, or a derivative thereof, in a pharmacologically suitable fluid that contains water, that are stable during long term
15 storage. The compositions are suitable for direct administration to a subject in need thereof. Pharmacologically suitable fluids include, but are not limited to, polar fluids, including protic fluids. In certain embodiments herein, the compositions are aqueous solutions.

The compositions provided herein possess an estimated shelf-life of
20 greater than 1, 2 or 3 months usage time at 25 °C and greater than or equal to 1, 2 or 3 years storage time at 5 °C. In certain of these embodiments, using Arrhenius kinetics, >80% or >85% or >90% or >95% estimated bronchodilating agent remains after such storage. These compositions are particularly useful for administration via
25 nebulization. In certain embodiments herein, the subject is a mammal. In other embodiments, the subject is a human.

The compositions provided herein are formulated to remain stable over a relatively long period of time. For example, the compositions provided herein are stored between -15 °C and 25 °C, or between 2 °C

-4-

and 8 °C, and remain stable for the desired time. In one embodiment, the compositions are stored at 5 °C.

Among the bronchodilating agents for use herein are Albuterol (α^1 -(((1,1-dimethylethyl)amino)methyl)-4-hydroxy-1,3-benzenedimethanol);

5 Bambuterol (dimethylcarbamic acid 5-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-phenylene ester); Bitolterol (4-methylbenzoic acid 4-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,2-phenylene ester); Broxaterol (3-bromo- α -(((1,1-dimethylethyl)amino)methyl)-5-isoxazolemethanol);

10 Isoproterenol (4-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,2-benzenediol); Trimetoquinol (1,2,3,4-tetrahydro-1-((3,4,5-trimethoxyphenyl)-methyl)-6,7-isoquinolinediol); Clenbuterol (4-amino-3,5-dichloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); Fenoterol (5-(1-hydroxy-2-((2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1,3-benzenediol); Formoterol (2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide); (R,R)-Formoterol; Desformoterol

15 ((R,R) or (S,S)-3-amino-4-hydroxy- α -(((2-(4-methoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol); Hexoprenaline (4,4'-(1,6-hexanediyl)-bis(imino(1-hydroxy-2,1-ethanediyl)))bis-1,2-benzenediol); Isoetharine (4-(1-hydroxy-2-((1-methylethyl)amino)butyl)-1,2-benzenediol); Isoprenaline


20 (4-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,2-benzenediol); Metaproterenol (5-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,3-benzenediol); Picumeterol (4-amino-3,5-dichloro- α -(((6-(2-(2-pyridinyl)ethoxy)hexyl)-amino)methyl)benzenemethanol); Pirbuterol (α^6 -(((1,1-dimethylethyl)-amino)methyl)-3-hydroxy-2,6-pyridinemethanol); Procaterol (((R*,S*)-(±)-8-hydroxy-5-(1-hydroxy-2-((1-methylethyl)amino)butyl)-2(1H)-quinolin-

25 one); Reproterol ((7-(3-((2-(3,5-dihydroxyphenyl)-2-hydroxyethyl)amino)-propyl)-3,7-dihydro-1,3-dimethyl-1H-purine-2,6-dione); Rimiterol (4-(hydroxy-2-piperidinylmethyl)-1,2-benzenediol); Salbutamol ((±)- α^1 -(((1,1-dimethylethyl)amino)methyl)-4-hydroxy-1,3-benzenedimethanol); (R)-

-5-

Salbutamol; Salmeterol (($\pm$)-4-hydroxy- α^1 -(((6-(4-phenylbutoxy)hexyl)-amino)methyl)-1,3-benzenedimethanol); (R)-Salmeterol; Terbutaline (5-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-benzenediol); Tulobuterol (2-chloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); and TA-
 5 2005 (8-hydroxy-5-((1R)-1-hydroxy-2-(N-((1R)-2-(4-methoxyphenyl)-1-methylethyl)amino)ethyl)carbostyryl hydrochloride).

Of particular interest herein is formoterol, having the formula:



20

Formoterol for use in the compositions and methods provided herein includes 2-hydroxy-5-((1RS)-1-hydroxy-2-(((1RS)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide; or a stereoisomer thereof; and also includes the single enantiomers 2-hydroxy-5-((1S)-1-hydroxy-2-
 25 (((1S)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide and 2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)-amino)ethyl)formanilide.

In certain embodiments, the compositions are administered via nebulization. Administration of a nebulized aerosol is preferred over the
 30 use of dry powders for inhalation in certain subject populations, including pediatric and geriatric groups.

In one embodiment, the compositions for use in the methods provided herein contain a pharmaceutically acceptable derivative of formoterol. In another embodiment, the compositions for use in the

50728-5

-6-

methods provided herein contain a pharmaceutically acceptable salt of formoterol. Pharmaceutically acceptable salts include, but are not limited to, salts of mineral acids, such as but not limited to hydrochlorides and sulfates; and salts of organic acids, such as but not limited to acetates, lactates, malates, tartrates, citrates, ascorbates, succinates, butyrates, valerates and fumarates. In one embodiment, the compositions for use in the methods provided herein contain formoterol fumarate or formoterol fumarate dihydrate. In another embodiment, the compositions for use in the methods provided herein contain formoterol tartrate.

According to yet another aspect of the present invention, there is provided a pharmaceutical composition, comprising formoterol, or a salt, ester, enol-ester, acid, base, solvate or prodrug thereof, in a pharmacologically acceptable fluid, wherein the composition is stable during long term storage, the fluid comprises water, and the composition is formulated at a concentration suitable for direct administration to a subject in need thereof.

Also provided herein are combinations containing a composition provided herein and a nebulizer. The combinations can be packaged as kits, which optionally contain other components, including instructions for use of the nebulizer. Any nebulizer is contemplated for use in the kits and methods provided herein. In particular, the nebulizers for use herein nebulize liquid formulations, including the compositions provided herein, containing no propellant. The nebulizer may produce the nebulized mist

50728-5

-6a-

by any method known to those of skill in the art, including, but not limited to, compressed air, ultrasonic waves, or vibration. The nebulizer may further have an internal baffle. The internal baffle, together with the housing of the nebulizer, selectively removes large droplets from the mist
5 by impaction and allows the droplets to return to the reservoir. The fine aerosol droplets thus produced are entrained into the lung by the inhaling air/oxygen.

Methods for the treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders, including, but not
10 limited to, asthma, including, but not limited to, bronchial asthma, allergic asthma and intrinsic asthma, e.g., late asthma and airway hyper-responsiveness; chronic bronchitis; and other chronic obstructive pulmonary diseases are provided. The methods involve administering an

-7-

effective amount of a pharmaceutical composition provided herein to a subject in need of such treatment.

Articles of manufacture, containing packaging material, a composition provided herein, which is useful for treatment, prevention or amelioration of one or more symptoms of diseases or disorders associated with undesired and/or uncontrolled bronchoconstriction, and a label that indicates that the composition is used for treatment, prevention or amelioration of one or more symptoms of diseases or disorders associated with undesired and/or uncontrolled bronchoconstriction, are also provided.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published applications and other publications are incorporated by reference in their entirety. In the event that there are a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

As used herein, formoterol refers to 2-hydroxy-5-((1RS)-1-hydroxy-2-(((1RS)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide; or a stereoisomer thereof. The term formoterol also refers to the single enantiomers 2-hydroxy-5-((1S)-1-hydroxy-2-(((1S)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide and 2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide.

As used herein, formoterol fumarate refers to a salt of formoterol having the formula (formoterol) • ½ fumarate.

As used herein, formoterol free base refers to the neutral, anhydrous form of formoterol. Thus, a recitation that a composition

-8-

contains, e.g., 59 $\mu\text{g/mL}$ of formoterol free base means that the composition contains 59 $\mu\text{g/mL}$ of neutral, anhydrous formoterol. Such compositions may be prepared using a derivative of formoterol.

As used herein, an aerosol is liquid or particulate matter dispersed
5 in air. Aerosols include dispersions of liquids, including aqueous and other solutions, and solids, including powders, in air.

As used herein, a nebulized solution refers to a solution that is dispersed in air to form an aerosol. Thus, a nebulized solution is a particular form of an aerosol.

10 As used herein, a nebulizer is an instrument that is capable of generating very fine liquid droplets for inhalation into the lung. Within this instrument, the nebulizing liquid or solution is atomized into a mist of droplets with a broad size distribution by methods known to those of skill in the art, including, but not limited to, compressed air, ultrasonic waves,
15 or a vibrating orifice. Nebulizers may further contain, e.g., a baffle which, along with the housing of the instrument, selectively removes large droplets from the mist by impaction. Thus, the mist inhaled into the lung contains fine aerosol droplets.

As used herein, a pharmacologically suitable fluid is a solvent
20 suitable for pharmaceutical use which is not a liquified propellant gas. Exemplary pharmacologically suitable fluids include polar fluids, including protic fluids such as water.

As used herein, a combination refers to any association between two or among more items.

25 As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

-9-

As used herein, a mixture is a mutual incorporation of two or more substances, without chemical union, the physical characteristics of each of the components being retained.

As used herein, the stability of a composition provided herein refers
5 to the length of time at a given temperature that is greater than 80%,
85%, 90% or 95% of the initial amount of active ingredient, e.g.,
formoterol, is present in the composition. Thus, for example, a
composition that is stable for 30 days at 25 °C would have greater than
80%, 85%, 90% or 95% of the initial amount of active ingredient present
10 in the composition at 30 days following storage at 25 °C.

As used herein, pharmaceutically acceptable derivatives of a
compound include salts, esters, enol ethers, enol esters, acids, bases,
solvates, hydrates or prodrugs thereof. Such derivatives may be readily
prepared by those of skill in this art using known methods for such
15 derivatization. The compounds produced may be administered to animals
or humans without substantial toxic effects and either are
pharmaceutically active or are prodrugs. Pharmaceutically acceptable
salts include, but are not limited to, amine salts, such as but not limited
to N,N'-dibenzylethylenediamine, chlorprocaine, choline, ammonia,
20 diethanolamine and other hydroxyalkylamines, ethylenediamine, N-
methylglucamine, procaine, N-benzylphenethylamine, 1-para-chlorobenzyl-
2-pyrrolidin-1'-ylmethylbenzimidazole, diethylamine and other alkylamines,
piperazine and tris(hydroxymethyl)aminomethane; alkali metal salts, such
as but not limited to lithium, potassium and sodium; alkali earth metal
25 salts, such as but not limited to barium, calcium and magnesium;
transition metal salts, such as but not limited to zinc; and other metal
salts, such as but not limited to sodium hydrogen phosphate and
disodium phosphate; and also including, but not limited to, salts of
mineral acids, such as but not limited to hydrochlorides and sulfates; and

-10-

salts of organic acids, such as but not limited to acetates, lactates, malates, tartrates, citrates, ascorbates, succinates, butyrates, valerates and fumarates. Pharmaceutically acceptable esters include, but are not limited to, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, heteroaralkyl, 5 cycloalkyl and heterocyclyl esters of acidic groups, including, but not limited to, carboxylic acids, phosphoric acids, phosphinic acids, sulfonic acids, sulfinic acids and boronic acids. Pharmaceutically acceptable enol ethers include, but are not limited to, derivatives of formula $C=C(OR)$ where R is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, 10 heteroaralkyl, cycloalkyl and heterocyclyl. Pharmaceutically acceptable enol esters include, but are not limited to, derivatives of formula $C=C(OC(O)R)$ where R is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cycloalkyl and heterocyclyl. Pharmaceutically acceptable solvates and hydrates are complexes of a 15 compound with one or more solvent or water molecule, in certain embodiments 1 to about 100, in other embodiments 1 to about 10, in further embodiments one to about 2, 3 or 4, solvent or water molecules. Formoterol salts and hydrates are used in certain embodiments herein.

As used herein, treatment means any manner in which one or more 20 of the symptoms of a condition, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein, such as use for treating cancer.

As used herein, amelioration of the symptoms of a particular 25 disorder by administration of a particular pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

-11-

As used herein, a prodrug is a compound that, upon in vivo administration, is metabolized or otherwise converted to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a prodrug, the pharmaceutically active compound is modified
5 such that the active compound will be regenerated by metabolic processes. The prodrug may be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes
10 and drug metabolism in vivo, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) Medicinal Chemistry A Biochemical Approach, Oxford University Press, New York, pages 388-392).

It is to be understood that the compounds for use in the
15 compositions and methods provided herein may contain chiral centers. Such chiral centers may be of either the (R) or (S) configuration, or may be a mixture thereof. Thus, the compounds for use in the compositions provided herein may be enantiomerically pure, or be stereoisomeric or diastereomeric mixtures. It is to be understood that the chiral centers of
20 the compounds provided herein may undergo epimerization in vivo. Thus, one of skill in the art will recognize that administration of a compound in its (R) form is equivalent, for compounds that undergo epimerization in vivo, to administration of the compound in its (S) form.

As used herein, bronchoconstriction refers to a reduction in the
25 caliber of a bronchus or bronchi.

As used herein, undesired and/or uncontrolled bronchoconstriction refers to bronchoconstriction that results in or from a pathological symptom or condition. Pathological conditions include, but are not

-12-

limited to, asthma and chronic obstructive pulmonary disease (COPD).
Pathological symptoms include, but are not limited to, asthma and COPD.

As used herein, the statement that a composition is stable during
"long term storage" means that the composition is suitable for
5 administration to a subject in need thereof when it has an estimated
shelf-life of greater than 1, 2 or 3 months usage time at 25 °C and
greater than or equal to 1, 2 or 3 years storage time at 5 °C. In certain
embodiments herein, using Arrhenius kinetics; >80% or >85% or
>90% or >95% estimated bronchodilating agent remains after such
10 storage.

A. Formoterol

Formoterol (2-hydroxy-5-((1RS)-1-hydroxy-2-(((1RS)-2-(p-methoxy-
phenyl)-1-methylethyl)amino)ethyl)formanilide) is derived from adrenaline
and, as noted above, is used as a β_2 -stimulator in inhalation therapy of
15 respiratory diseases, particularly for the treatment of bronchial asthma. It
has been reported that in patients with reversible obstructive respiratory
diseases, formoterol has a bronchodilatory effect. This effect has a
relatively rapid onset (approximately 1-3 minutes) and a relatively long
duration (greater than 12 hours). Formoterol inhibits the release of
20 leukotrienes and other messenger substances involved with inflammation,
such as histamines. In addition, formoterol may bring about a
hyperglycaemic activity.

To date, formoterol has been formulated as a dry powder and
administered via devices such as the Turbuhaler® and the Aerolizer®.
25 See, e.g., Seberova et al. (2000) Respir. Med. 94(6):607-611; Lotvall et
al. (1999) Can. Respir. J. 6(5):412-416; Campbell et al. (1999) Respir.
Med. 93(4):236-244; Nightingale et al. (1999) Am. J. Respir. Crit. Care
Med. 159(6):1786-1790; Lecaillon et al. (1999) Eur. J. Clin. Pharmacol.
55(2):131-138; Bartow et al. (1998) Drugs 55(2):303-322; Ekstrom et

-13-

al. (1998) Respir. Med. 92(8):1040-1045; Ringdal et al. (1998) Respir. Med. 92(8):1017-1021; Totterman et al. (1998) Eur. Respir. J. 12(3):573-579; Palmqvist et al. (1997) Eur. Respir. J. 10(11):2484-2489; Nielsen et al. (1997) Eur. Respir. J. 10(9):2105-2109; Ullman et
5 al. (1996) Allergy 51(10):745-748; Selroos et al. (1996) Clin. Immunother. 6:273-299; and Schreurs et al. (1996) Eur. Respir. J. 9(8):1678-1683.

Formoterol is also available as a tablet and a dry syrup in certain areas of the world (e.g., Atock®, marketed by Yamanouchi
10 Pharmaceutical Co. Ltd., Japan). Formoterol formulations are also available in other areas (e.g., Europe and U.S.) for propellant-based metered dose inhalers and dry powder inhalers (e.g., Turbuhaler®, Aerolizer® and Foradil Aerolizer®). None of these formulations are water
15 based. Sterile, stable, aqueous based inhalation solutions of formoterol for nebulization are not available, nor have they been reported.

Compositions containing formoterol in combination with other active ingredients have been disclosed. See, e.g., U.S. Patent Nos. 6,004,537, 5,972,919 and 5,674,860 (formoterol and budesonide), 5,668,110, 5,683,983, 5,677,280 and 5,654,276 (formoterol and IL-5
20 inhibitors), 6,136,603 (formoterol and antisense modulators of IL-5), 5,602,110 (formoterol and millrinone), 5,525,623 (formoterol and a tryptase inhibitor), 5,691,336, 5,877,191, 5,929,094, 5,750,549 and 5,780,467 (formoterol and a tachykinin receptor antagonist); and International Patent Application Publication Nos. WO 99/00134
25 (formoterol and rofleponide) and WO 99/36095 (formoterol and a dopamine D₂ receptor agonist).

Other compositions containing formoterol have been disclosed in U.S. Patent Nos. 5,677,809, 6,126,919, 5,733,526, 6,071,971,

-14-

6,068,833, 5,795,564, 6,040,344, 6,041,777, 5,874,481, 5,965,622 and 6,161,536.

U.S. Patent No. 6,150,418 discloses a "liquid active substance concentrate" containing formoterol in the form of its free base or in the
5 form of one of the pharmacologically acceptable salts or addition products (adducts) thereof as active substance. This "liquid active substance concentrate" is reported to be a concentrated (i.e., greater than 10 mg/mL, preferably 75 to 500 mg/mL) solution or suspension that is stable for a period of several months possibly up to several years
10 without any deterioration in the pharmaceutical quality. This patent teaches that it is the high concentration that allows for the stability of the concentrate. The "liquid active substance concentrate" is not suitable for direct administration to a patient.

U.S. Patent No. 6,040,344 discloses an aqueous aerosol formula-
15 tion of formoterol tartrate for use in a nebulizer. This patent states that the formulation disclosed therein is not attractive for long term storage.

B. Compositions for use in treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders

Pharmaceutical compositions containing a β_2 -adrenoreceptor
20 agonist for administration via nebulization are provided. The compositions are sterile filtered and filled in vials, including unit dose vials providing sterile unit dose formulations which are used in a nebulizer and suitably nebulized. Each unit dose vial is sterile and is suitably nebulized without contaminating other vials or the next dose.

25 The unit dose vials are formed in a form-fill-seal machine or by any other suitable method known to those of skill in the art. The vials may be made of plastic materials that are suitably used in these processes. For example, plastic materials for preparing the unit dose vials include, but are not limited to, low density polyethylene, high density polyethylene,

-15-

polypropylene and polyesters. In one embodiment, the plastic material is low density polyethylene.

In one embodiment, the β_2 -adrenoreceptor agonist is formoterol, or a pharmaceutically acceptable derivative thereof. In other embodiments, the formoterol for use in the compositions provided herein is formoterol fumarate. Formoterol refers to 2-hydroxy-5-((1RS)-1-hydroxy-2-(((1RS)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide; or a stereo-isomer thereof. The term formoterol also refers herein to the single enantiomers 2-hydroxy-5-((1S)-1-hydroxy-2-(((1S)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide and 2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide.

In one embodiment, the compositions contain formoterol free base at a concentration of about 5 $\mu\text{g/mL}$ to about 2 mg/mL. In other embodiments, the maximum concentration of formoterol free base in the compositions is 1.5 mg/mL. In further embodiments, the concentration of formoterol free base in the compositions is about 10 $\mu\text{g/mL}$ to about 1 mg/mL, or about 50 $\mu\text{g/mL}$ to about 200 $\mu\text{g/mL}$. In other embodiments, the compositions contain formoterol fumarate at a concentration of about 80 $\mu\text{g/mL}$ up to about 175 to 200 $\mu\text{g/mL}$. In further embodiments, the compositions contain formoterol fumarate at a concentration of about 90 $\mu\text{g/mL}$ up to about 125 to 150 $\mu\text{g/mL}$. The formoterol fumarate is formulated, in certain compositions provided herein, at a concentration of about 100 $\mu\text{g/mL}$. The formoterol fumarate is formulated, in other compositions provided herein, at a concentration of about 85 $\mu\text{g/mL}$ or about 170 $\mu\text{g/mL}$. In one embodiment, the formoterol fumarate is formulated for single dosage administration via nebulization at a concentration of about 100 $\mu\text{g/mL}$. In another embodiment, the compositions contain formoterol free base at a concentration of about 40 to about 150 $\mu\text{g/mL}$, particularly about 59 or about 118 $\mu\text{g/mL}$.

-16-

The compositions containing the β_2 -adrenoreceptor agonist, including formoterol, are formulated with a pharmacologically suitable fluid. Pharmacologically suitable fluids include, but are not limited to, polar solvents, including, but not limited to, compounds that contain
5 hydroxyl groups or other polar groups. Such solvents include, but are not limited to, water or alcohols, such as ethanol, isopropanol, and glycols including propylene glycol, polyethylene glycol, polypropylene glycol, glycol ether, glycerol and polyoxyethylene alcohols.

Polar solvents also include protic solvents, including, but not
10 limited to, water, aqueous saline solutions with one or more pharmaceutically acceptable salt(s), alcohols, glycols or a mixture thereof. For a saline solution as the solvent or as a component thereof, particularly suitable salts are those which display no or only negligible pharmacological activity after administration.

15 In the embodiments herein, the compositions have a pH of about 2.0 to about 8.0. In other embodiments, the compositions have a pH of about 4.0 to about 6.0, or about 4.5 to about 5.5. In certain of the above embodiments, the compositions are formulated at a pH of about 4, 4.4 or 4.6 up to about 5.5, 5.7 or 6. In other embodiments, the pH is
20 about 5.0. It has been found herein that the rate constant for decomposition of an aqueous solution of formoterol is dependent on pH. The rate constant (k_{obs}) at 60 °C at a pH of 3, 4, 5 and 7 is approximately 0.62, 0.11, 0.044 and 0.55 day⁻¹, respectively. Therefore, the decomposition of formoterol in aqueous solution at 60 °C
25 at a buffer concentration of 5 mM and an ionic strength of 0.05 is slowest at a pH of about 5.0.

The solubility of formoterol in aqueous solution has been found herein to be dependent on pH. Thus, at a pH of between about 5 and about 7, the aqueous solubility of formoterol at ambient temperature is

-17-

approximately 2.2 mg/mL. At a pH of about 4, the aqueous solubility of formoterol at ambient temperature is approximately 3 mg/mL, while at a pH of about 3, the aqueous solubility of formoterol at ambient temperature is about 4.8 mg/mL. The solubility of formoterol in pure water, for example, high performance liquid chromatography (HPLC) water, at ambient temperature is approximately 2 mg/mL.

In other of the above embodiments, the compositions further contain a buffer, including, but not limited to, citric acid/phosphate, acetate, barbital, borate, Britton-Robinson, cacodylate, citrate, collidine, formate, maleate, McIlvaine, phosphate, Prideaux-Ward, succinate, citrate-phosphate-borate (Teorell-Stanhagen), veronal acetate, MES (2-(N-morpholino)ethanesulfonic acid), BIS-TRIS (bis(2-hydroxyethyl)iminotris(hydroxymethyl)methane), ADA (N-(2-acetamido)-2-iminodiacetic acid), ACES (N-(carbamoylmethyl)-2-aminoethanesulfonic acid), PIPES (piperazine-N,N'-bis(2-ethanesulfonic acid)), MOPSO (3-(N-morpholino)-2-hydroxypropanesulfonic acid), BIS-TRIS PROPANE (1,3-bis(tris(hydroxymethyl)methylamino)propane), BES (N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid), MOPS (3-(N-morpholino)propanesulfonic acid), TES (N-tris(hydroxymethyl)methyl-2-aminoethanesulfonic acid), HEPES (N-(2-hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid)), DIPSO (3-(N,N-bis(2-hydroxyethyl)amino)-2-hydroxypropanesulfonic acid), MOBS (4-(N-morpholino)butanesulfonic acid), TAPSO (3-(N-tris(hydroxymethyl)methylamino)-2-hydroxypropanesulfonic acid), TRIZMA® (tris(hydroxymethyl)aminomethane), HEPPSO (N-(2-hydroxyethyl)piperazine-N'-(2-hydroxypropanesulfonic acid)), POPSO (piperazine-N,N'-bis(2-hydroxypropanesulfonic acid)), TEA (triethanolamine), EPPS (N-(2-hydroxyethyl)piperazine-N'-(3-propanesulfonic acid)), TRICINE (N-tris(hydroxymethyl)methylglycine), GLY-GLY (glycylglycine), BICINE (N,N-bis(2-hydroxyethyl)glycine), HEPBS (N-(2-hydroxyethyl)piperazine-N'-(4-

-18-

butanesulfonic acid)), TAPS (N-tris(hydroxymethyl)methyl-3-amino-
propanesulfonic acid), AMPD (2-amino-2-methyl-1,3-propanediol), and/or
any other buffers known to those of skill in the art. In one embodiment,
the buffer is citric acid/phosphate buffer, acetate buffer, citrate buffer or
5 phosphate buffer. In another embodiment, the buffer is a citrate buffer
(citric acid/sodium citrate). The buffer concentration has been found
herein to affect the stability of the composition. Buffer concentrations
for use herein include from about 0 or 0.01 mM to about 150 mM, or
about 1 mM to about 20 mM. In one embodiment, the buffer
10 concentration is about 5 mM. In other embodiments, the buffer
concentration is about 1 mM to about 50 mM. In one embodiment, the
buffer concentration is about 20 mM. The kinetic-pH profile of formoterol
is dependent on buffer concentration. At low and approximately neutral
conditions, increasing the buffer concentration from 5 mM to 20 mM
15 increased the rate constant of decomposition significantly. However, no
noticeable differences in rate constant were observed in the pH region of
about 4.5 to about 5.5 with increasing buffer concentration from 5 mM to
20 mM. The particular buffer and buffer concentration of a given
composition for long term storage provided herein may be determined
20 empirically using standard stability assays well known to those of skill in
the art (see, e.g., the Examples).

The ionic strength of the compositions provided herein also has
been found herein to affect the stability of the composition. Ionic
strengths of the compositions provided herein are from about 0 to about
25 0.4, or from about 0.05 to about 0.16. Compositions having a lower
ionic strength exhibit improved stability over formulations having higher
ionic strength. The rate constant of decomposition was essentially the
same at ionic strength 0.05 to 0.1, but increased to some extent at ionic
strength of 0.2. The particular ionic strength of a given composition for

-19-

long term storage provided herein may be determined empirically using standard stability assays well known to those of skill in the art (see, e.g., the Examples).

In embodiments where the pharmacologically suitable fluid is a saline solution, tonicity adjusting agents may be added to provide the desired ionic strength. Tonicity adjusting agents for use herein include those which display no or only negligible pharmacological activity after administration. Both inorganic and organic tonicity adjusting agents may be used in the compositions provided herein. Tonicity adjusting agents include, but are not limited to, ammonium carbonate, ammonium chloride, ammonium lactate, ammonium nitrate, ammonium phosphate, ammonium sulfate, ascorbic acid, bismuth sodium tartrate, boric acid, calcium chloride, calcium disodium edetate, calcium gluconate, calcium lactate, citric acid, dextrose, diethanolamine, dimethylsulfoxide, edetate disodium, edetate trisodium monohydrate, fluorescein sodium, fructose, galactose, glycerin, lactic acid, lactose, magnesium chloride, magnesium sulfate, mannitol, polyethylene glycol, potassium acetate, potassium chlorate, potassium chloride, potassium iodide, potassium nitrate, potassium phosphate, potassium sulfate, propylene glycol, silver nitrate, sodium acetate, sodium bicarbonate, sodium biphosphate, sodium bisulfite, sodium borate, sodium bromide, sodium cacodylate, sodium carbonate, sodium chloride, sodium citrate, sodium iodide, sodium lactate, sodium metabisulfite, sodium nitrate, sodium nitrite, sodium phosphate, sodium propionate, sodium succinate, sodium sulfate, sodium sulfite, sodium tartrate, sodium thiosulfate, sorbitol, sucrose, tartaric acid, triethanolamine, urea, urethan, uridine and zinc sulfate. In certain embodiments, the tonicity adjusting agent is sodium chloride. In these embodiments, the pharmacologically suitable fluid is aqueous saline.

-20-

The storage temperature of the compositions provided herein also has been found herein to affect the stability of the composition.

Compositions stored at a lower temperature exhibit improved stability over formulations stored at higher temperatures. The effect of

5 temperature on the rate constant of decomposition at pH5, a buffer concentration of 5 mM, and an ionic strength of 0.05, was linear according to Arrhenius kinetics, i.e., when $\ln k_{\text{obs}}$ was plotted against $1/T$, where T is the temperature in degree Kelvin.

The estimated shelf-life of formoterol in the compositions provided
10 herein is significantly greater than that reported for known formoterol compositions. The estimated shelf-life of formoterol in the compositions provided herein is about 6.2 years at 5 °C and about 7.5 months at 25 °C. The estimated formoterol concentrations in the compositions provided herein as a function of storage time at 5 °C and usage time at
15 25 °C was determined. It is estimated that greater than 90% of the initial formoterol present in the composition remains after 3 months of usage time at 25 °C and 3 years of storage time at 5 °C as well as after 0.5 months of usage time at 25 °C and 1 year of storage time at 5 °C.

In one embodiment, the compositions provided herein are prepared
20 containing formoterol fumarate at a nominal concentration of 0.1 mg/mL at the indicated pH and citric acid/phosphate buffer concentrations. The solutions were stored at 60 °C. In these compositions, formoterol is relatively more stable at a pH from about 4 to about 5, and is also more stable at lower buffer concentration.

25 The compositions provided herein also may include excipients and additives. The particular excipient or additive for use in the compositions for long term storage provided herein may be determined empirically using methods well known to those of skill in the art (see, e.g., the Examples). Excipients and additives are any pharmacologically suitable and

-21-

therapeutically useful substance which is not an active substance.

Excipients and additives generally have no pharmacological activity, or at least no undesirable pharmacological activity. The excipients and additives include, but are not limited to, surfactants, stabilizers,

- 5 complexing agents, antioxidants, or preservatives which prolong the duration of use of the finished pharmaceutical formulation, flavorings, vitamins, or other additives known in the art. Complexing agents include, but are not limited to, ethylenediaminetetraacetic acid (EDTA) or a salt thereof, such as the disodium salt, citric acid, nitrilotriacetic acid and the
- 10 salts thereof. In one embodiment, the complexing agent is EDTA. Preservatives include, but are not limited to, those that protect the solution from contamination with pathogenic particles, including benzalkonium chloride or benzoic acid, or benzoates such as sodium benzoate. Antioxidants include, but are not limited to, vitamins,
- 15 provitamins, ascorbic acid, vitamin E or salts or esters thereof.

- The compositions provided herein also may include a cosolvent, which increases the solubility of additives or the active ingredient(s). The particular cosolvent for use in the compositions for long term storage provided herein may be determined empirically using methods well known
- 20 to those of skill in the art (see, e.g., the Examples). Cosolvents for use herein include, but are not limited to, hydroxylated solvents or other polar solvents, such as alcohols such as isopropyl alcohol, glycols such as propylene glycol, polyethylene glycol, polypropylene glycol, glycol ether, glycerol, and polyoxyethylene alcohols.

25 **C. Preparation of compounds for use in the compositions**

The preparation of the compounds used in the compositions provided herein is described below. Any such compound or similar compound may be synthesized according to a method discussed in

-22-

general below or by only minor modification of the methods by selecting appropriate starting materials.

Formoterol may be prepared according to the method disclosed in U.S. Patent No. 3,994,974. Briefly, 4-benzyloxy-3-nitro- α -
5 bromoacetophenone is reacted with N-benzyl-N-(1-methyl-2-p-methoxyphenylethyl)amine to form the α -aminoacetophenone. This compound was subjected to the following series of reactions: (i) reduction of the ketone with sodium borohydride; (ii) reduction of the nitro group with aqueous hydrochloric acid and iron powder; (iii) amine
10 formylation with acetic anhydride and formic acid; and (iv) catalytic reduction over 10% palladium on carbon to afford formoterol free base. Crystallization of the $\frac{1}{2}$ fumarate salt from ethanol provides (formoterol) $\cdot \frac{1}{2}$ fumarate.

The individual enantiomers of formoterol, 2-hydroxy-5-((1S)-1-
15 hydroxy-2-(((1S)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)-formanilide and 2-hydroxy-5-((1R)-1-hydroxy-2-(((1R)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide, may be prepared by the method disclosed in U.S. Patent No. 6,040,344. Briefly, reaction of optically pure 4-benzyloxy-3-formamidostyrene oxide with an optically
20 pure 4-methoxy- α -methyl-N-(phenylmethyl)benzeneethanamine, followed by debenzylolation, affords the desired enantiomer of formoterol. Debenzylolation may be accomplished by reduction with hydrogen gas in the presence of a noble metal catalyst, such as palladium on carbon.

The required optically pure 4-benzyloxy-3-formamidostyrene oxide
25 may be prepared from 4-benzyloxy-3-nitro- α -bromoacetophenone by (i) reduction with vorane in the presence of an optically pure aminoindanol, (ii) hydrogenation over platinum oxide catalyst, (iii) formylation with formic acid and acetic anhydride, and (iv) epoxide formation in the presence of potassium carbonate.

-23-

The required optically pure 4-methoxy- α -methyl-N-(phenylmethyl)-benzeneethanamine may be prepared from 4-methoxyphenylacetone by (i) reductive amination with benzylamine in the presence of hydrogen and a platinum catalyst, and (ii) crystallization of the desired optically pure
5 amine from the resulting racemic mixture as its mandelic acid salt.

D. Formulation of pharmaceutical compositions

The compositions provided herein are prepared by procedures well known to those of skill in the art. For example, a formoterol fumarate solution may be prepared by the procedure of EXAMPLE 1. Briefly, a
10 buffer solution having a pH and ionic strength of interest herein is prepared. In one embodiment, the buffer is a mixture of citric acid and sodium citrate, with sodium chloride added to achieve the desired ionic strength. Formoterol fumarate dihydrate is added to the buffer solution with agitation to produce a solution of the desired formoterol
15 concentration. Exemplary formoterol concentrations are 0.17 g formoterol fumarate dihydrate/2 L and 0.34 g formoterol fumarate dihydrate/2 L buffer.

E. Evaluation of the activity of the compositions

Standard physiological, pharmacological and biochemical
20 procedures are available for testing the compositions provided herein to identify those that possess bronchodilatory activity.

In vitro and in vivo assays that may be used to evaluate bronchodilatory activity are well known to those of skill in the art. See also, e.g., U.S. Patent Nos. 3,994,974, and 6,068,833; German Patent
25 No. 2,305,092; Kaumann et al. (1985) Naunyn-Schmied Arch. Pharmacol. 331:27-39; Lemoine et al. (1985) Naunyn-Schmied Arch. Pharmacol. 331:40-51; Tomioka et al. (1981) Arch. Int. Pharmacodyn. 250:279-292; Dellamary et al. (2000) Pharm. Res. 17(2):168-174; Rico-Mendez et al. (1999) Rev. Alerg. Mex. 46(5):130-135; Seberova et al.

-24-

- (2000) Respir. Med. 94(6):607-611; Lotvall et al. (1999) Can. Respir. J. 6(5):412-416; Campbell et al. (1999) Respir. Med. 93(4):236-244; Nightingale et al. (1999) Am. J. Respir. Crit. Care Med. 159(6):1786-1790; Lecaillon et al. (1999) Eur. J. Clin. Pharmacol. 55(2):131-138;
- 5 Bartow et al. (1998) Drugs 55(2):303-322; Ekstrom et al. (1998) Respir. Med. 92(8):1040-1045; Ringdal et al. (1998) Respir. Med. 92(8):1017-1021; Totterman et al. (1998) Eur. Respir. J. 12(3):573-579; Palmqvist et al. (1997) Eur. Respir. J. 10(11):2484-2489; Nielsen et al. (1997) Eur. Respir. J. 10(9):2105-2109; Ullman et al. (1996) Allergy 51(10):745-
- 10 748; Selroos et al. (1996) Clin. Immunother. 6:273-299; and Schreurs et al. (1996) Eur. Respir. J. 9(8):1678-1683.

F. Methods of treatment of bronchoconstrictive disorders

The compositions provided herein are used for treating, preventing, or ameliorating one or more symptoms of a bronchoconstrictive disorders

15 in a subject. In one embodiment, the method includes administering to a subject an effective amount of a composition containing a bronchodilating agent, including, but not limited to, formoterol, whereby the disease or disorder is treated or prevented. The subject treated is, in certain embodiments, a mammal. The mammal treated is, in certain

20 embodiments, a human.

In another embodiment, the method provided herein includes oral administration of a composition provided herein. In certain embodiments herein, the composition is directly administered to a subject in need of such treatment via nebulization without dilution or other modification of

25 the composition prior to administration.

The methods for treatment, prevention, or amelioration of one or more symptoms of bronchoconstrictive disorders, in another embodiment, further include administering one or more of (a), (b), (c) or (d) as follows: (a) a β_2 -adrenoreceptor agonist; (b) a dopamine (D_2) receptor agonist; (c)

-25-

a prophylactic therapeutic, such as a steroid; or (d) an anticholinergic agent; simultaneously with, prior to or subsequent to the composition provided herein.

- β_2 -Adrenoreceptor agonists for use in combination with the
- 5 compositions provided herein include, but are not limited to, Albuterol (α^1 -(((1,1-dimethylethyl)amino)methyl)-4-hydroxy-1,3-benzenedimethanol); Bambuterol (dimethylcarbamic acid 5-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-phenylene ester); Bitolterol (4-methylbenzoic acid 4-(2-((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,2-phenylene ester); Broxa-
- 10 terol (3-bromo- α -(((1,1-dimethylethyl)amino)methyl)-5-isoxazolemethanol); Isoproterenol (4-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,2-benzenediol); Trimetoquinol (1,2,3,4-tetrahydro-1-((3,4,5-trimethoxyphenyl)-methyl)-6,7-isoquinolinediol); Clenbuterol (4-amino-3,5-dichloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); Fenoterol (5-(1-hydroxy-2-
- 15 ((2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1,3-benzenediol); Formoterol (2-hydroxy-5-((1RS)-1-hydroxy-2-(((1RS)-2-(p-methoxyphenyl)-1-methylethyl)amino)ethyl)formanilide); (R,R)-Formoterol; Desformoterol ((R,R) or (S,S)-3-amino-4-hydroxy- α -(((2-(4-methoxyphenyl)-1-methylethyl)amino)methyl)benzenemethanol); Hexoprenaline (4,4'-(1,6-hexane-
- 20 diyl)-bis(imino(1-hydroxy-2,1-ethanediyl)))bis-1,2-benzenediol); Isoetharine (4-(1-hydroxy-2-((1-methylethyl)amino)butyl)-1,2-benzenediol); Isoprenaline (4-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,2-benzenediol); Metaproterenol (5-(1-hydroxy-2-((1-methylethyl)amino)ethyl)-1,3-benzenediol); Picumeterol (4-amino-3,5-dichloro- α -(((6-(2-(2-pyridinyl)ethoxy)hexyl)-
- 25 amino)methyl)benzenemethanol); Pirbuterol (α^6 -(((1,1-dimethylethyl)-amino)methyl)-3-hydroxy-2,6-pyridinemethanol); Procaterol (((R*,S*)-($\pm$))-8-hydroxy-5-(1-hydroxy-2-((1-methylethyl)amino)butyl)-2(1H)-quinolinone); Reproterol ((7-(3-((2-(3,5-dihydroxyphenyl)-2-hydroxyethyl)amino)-propyl)-3,7-dihydro-1,3-dimethyl-1H-purine-2,6-dione); Rimiterol (4-

-26-

(hydroxy-2-piperidinylmethyl)-1,2-benzenediol); Salbutamol (($\pm$)- α^1 -(((1,1-dimethylethyl)amino)methyl)-4-hydroxy-1,3-benzenedimethanol); (R)-Salbutamol; Salmeterol (($\pm$)-4-hydroxy- α^1 -(((6-(4-phenylbutoxy)hexyl)-amino)methyl)-1,3-benzenedimethanol); (R)-Salmeterol; Terbutaline (5-(2-
5 ((1,1-dimethylethyl)amino)-1-hydroxyethyl)-1,3-benzenediol); Tulobuterol (2-chloro- α -(((1,1-dimethylethyl)amino)methyl)benzenemethanol); and TA-2005 (8-hydroxy-5-((1R)-1-hydroxy-2-(N-((1R)-2-(4-methoxyphenyl)-1-methylethyl)amino)ethyl)carbostyryl hydrochloride).

Dopamine (D_2) receptor agonists include, but are not limited to,
10 Apomorphine ((r)-5,6,6a,7-tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-10,11-diol); Bromocriptine ((5' α)-2-bromo-12'-hydroxy-2'-(1-methylethyl)-5'-(2-methylpropyl)ergotaman-3',6',18-trione); Cabergoline ((8 β)-N-(3-(dimethylamino)propyl)-N-((ethylamino)carbonyl)-6-(2-propenyl)ergoline-8-carboxamide); Lisuride (N'-((8 α)-9,10-didehydro-6-methylethyl-8-yl)-N,N-
15 diethylurea); Pergolide ((8 β)-8-((methylthio)methyl)-6-propylergoline); Levodopa (3-hydroxy-L-tryrosine); Pramipexole ((s)-4,5,6,7-tetrahydro-N⁶-propyl-2,6-benzothiazolodiamine); Quinpirole hydrochloride (trans-(-)-4aR-4,4a,5,6,7,8,8a,9-octahydro-5-propyl-1H-pyrazolo[3,4-g]quinoline hydrochloride); Ropinirole (4-(2-(dipropylamino)ethyl)-1,3-dihydro-2H-
20 indol-2-one); and Talipexole (5,6,7,8-tetrahydro-6-(2-propenyl)-4H-thiazolo[4,5-d]azepin-2-amine). Other dopamine D_2 receptor agonists for use herein are disclosed in International Patent Application Publication No. WO 99/36095.

Prophylactic therapeutics for use in combination therapy herein
25 include steroidal anti-inflammatory agents, including, but not limited to, beclomethasone dipropionate (BDP), beclomethasone monopropionate (BMP), flunisolide, triamcinolone acetonide, dexamethasone, tipredane, ciclesonid, rofleponide, mometasone, mometasone furoate (Asmanex[®] Twisthaler[™], Shering-Plough Corporation, Kenilworth, NJ), RPR 106541,

-27-

fluticasone or fluticasone propionate and budesonide or by way of sodium cromoglycate or nedocromil sodium.

Anticholinergic agents for use herein include, but are not limited to, ipratropium bromide, oxitropium bromide, atropine methyl nitrate, atropine sulfate, ipratropium, belladonna extract, scopolamine, scopolamine methobromide, homatropine methobromide, hyoscyamine, isopropiuramide, orphenadrine, benzalkonium chloride, tiotropium bromide and glycopyrronium bromide. In certain embodiments, the compositions contain an anticholinergic agent, such as ipratropium bromide or tiotropium bromide, at a concentration of about 5 $\mu\text{g/mL}$ to about 5 mg/mL, or about 50 $\mu\text{g/mL}$ to about 200 $\mu\text{g/mL}$. In other embodiments, the compositions for use in the methods herein contain an anticholinergic agent, including ipratropium bromide and tiotropium bromide, at a concentration of about 83 $\mu\text{g/mL}$ or about 167 $\mu\text{g/mL}$.

Other active ingredients for use herein in combination therapy, include, but are not limited to, IL-5 inhibitors such as those disclosed in U.S. Patent Nos. 5,668,110, 5,683,983, 5,677,280 and 5,654,276; antisense modulators of IL-5 such as those disclosed in U.S. Patent No. 6,136,603; milrinone (1,6-dihydro-2-methyl-6-oxo-[3,4'-bipyridine]-5-carbonitrile); milrinone lactate; tryptase inhibitors such as those disclosed in U.S. Patent No. 5,525,623; tachykinin receptor antagonists such as those disclosed in U.S. Patent Nos. 5,691,336, 5,877,191, 5,929,094, 5,750,549 and 5,780,467; leukotriene receptor antagonists such as montelukast sodium (Singular[®], R-(E)-1-[[[1-[3-[2-(7-chloro-2-quinolinyl)ethenyl]phenyl]-3-[2-(1-hydroxy-1-methylethyl)phenyl]-propyl]thio]methyl]cyclopropaneacetic acid, monosodium salt), 5-lipoxygenase inhibitors such as zileuton (Zyflo[®], Abbott Laboratories, Abbott Park, IL), and anti-IgE antibodies such as Xolair[®] (recombinant

-28-

humanized anti-IgE monoclonal antibody (CGP 51901; IGE 025A; rhuMAb-E25), Genentech, Inc., South San Francisco, CA).

The bronchoconstrictive disorder to be treated, prevented, or whose one or more symptoms are to be ameliorated is associated with
5 asthma, including, but not limited to, bronchial asthma, allergic asthma and intrinsic asthma, e.g., late asthma and airway hyper-responsiveness; and, particularly in embodiments where an anticholinergic agent is used, other chronic obstructive pulmonary diseases (COPDs), including, but not limited to, chronic bronchitis, emphysema, and associated cor pulmonale
10 (heart disease secondary to disease of the lungs and respiratory system) with pulmonary hypertension, right ventricular hypertrophy and right heart failure. COPD is frequently associated with cigarette smoking, infections, environmental pollution and occupational dust exposure.

G. Nebulizers

15 The compositions provided herein are intended for administration to a subject in need of such treatment via nebulization. Nebulizers that nebulize liquid formulations containing no propellant are suitable for use with the compositions provided herein. Nebulizers are available from, e.g., Pari GmbH (Starnberg, Germany), DeVilbiss Healthcare (Heston,
20 Middlesex, UK), Healthdyne, Vital Signs, Baxter, Allied Health Care, Invacare, Hudson, Omron, Bmed, AirSep, Luminscope, Medisana, Siemens, Aerogen, Mountain Medical, Aerosol Medical Ltd. (Colchester, Essex, UK), AFP Medical (Rugby, Warwickshire, UK), Bard Ltd. (Sunderland, UK), Carri-Med Ltd. (Dorking, UK), Plaem Nuiva (Brescia,
25 Italy), Henleys Medical Supplies (London, UK), Intersurgical (Berkshire, UK), Lifecare Hospital Supplies (Leies, UK), Medic-Aid Ltd. (West Sussex, UK), Medix Ltd. (Essex, UK), Sinclair Medical Ltd. (Surrey, UK), and many others.

Nebulizers for use herein include, but are not limited to, jet nebulizers (optionally sold with compressors), ultrasonic nebulizers, and others. Exemplary jet nebulizers for use herein include Pari LC plus/ProNeb^{*}, Pari LC plus/ProNeb Turbo^{*}, Pari LC plus/Dura Neb 1000 & 2000^{*}, Pari LC plus/Walkhaler^{*}, Pari LC plus/Pari Master^{*}, Pari LC star^{*}, Omron CompAir XL Portable Nebulizer System (NE-C18 and JetAir Disposable nebulizer)^{*}, Omron CompAir Elite Compressor Nebulizer System (NE-C21 and Elite Air Reusable Nebulizer)^{*}, Pari LC Plus or Pari LC Star nebulizer with Proneb Ultra compressor^{*}, Pulmo-aide^{*}, Pulmo-aide LT^{*}, Pulmo-aide traveler^{*}, Invacare Passport^{*}, Inspiration Healthdyne 626^{*}, Pulmo-Neb Traverler^{*}, DeVilbiss 646^{*}, Whisper Jet^{*}, Acorn II^{*}, Misty-Neb^{*}, Allied aerosol^{*}, Schuco Home Care^{*}, Lexan Plastic Pocet Neb^{*}, SideStream Hand Held Neb^{*}, Mobil Mist^{*}, Up-Draft^{*}, Up-Draft II^{*}, T Up-Draft^{*}, ISO-NEB^{*}, AVA-NEB^{*}, Micro Mist^{*}, and PulmoMate^{*}. Exemplary ultrasonic nebulizers for use herein include MicroAir^{*}, UltraAir^{*}, Siemens Ultra Nebulizer 145^{*}, CompAir^{*}, Pulmosonic^{*}, Scout^{*}, 5003 Ultrasonic Neb^{*}, 5110 Ultrasonic Neb^{*}, 5004 Desk Ultrasonic Nebulizer^{*}, Mystique Ultrasonic^{*}, Luminscope's Ultrasonic Nebulizer^{*}, Medisana Ultrasonic Nebulizer^{*}, Microstat Ultrasonic Nebulizer^{*}, and MABISMist Hand Held Ultrasonic Nebulizer^{*}. Other nebulizers for use herein include 5000 Electromagnetic Neb^{*}, 5001 Electromagnetic Neb 5002 Rotary Piston Neb^{*}, Lumineb I Piston Nebulizer 5500^{*}, Aeroneb™ Portable Nebulizer System, Aerodose™ Inhaler, and AeroEclipse Breath Actuated Nebulizer^{*}.

H. Articles of manufacture

The compositions provided herein may be packaged as articles of manufacture containing packaging material, a composition provided herein, which is useful for treatment, prevention or amelioration of one or more symptoms of diseases or disorders associated with undesired and/or uncontrolled bronchoconstriction, and a label that indicates that the

*Trade-mark

-30-

composition is used for treatment, prevention or amelioration of one or more symptoms of diseases or disorders associated with undesired and/or uncontrolled bronchoconstriction.

The articles of manufacture provided herein contain packaging materials. Packaging materials for use in packaging pharmaceutical products are well known to those of skill in the art. See, e.g., U.S. Patent Nos. 5,323,907, 5,052,558 and 5,033,252. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, inhalers, pumps, bags, vials, containers, syringes, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment.

In one embodiment herein, the compositions are packaged with a nebulizer for direct administration of the composition to a subject in need thereof.

15

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1

Preparation of Formoterol Inhalation Solution Formulation

20 To a 5 L stainless steel vessel were added 0.68 g citric acid USP, 1.99 g sodium citrate USP, and 17.5 g sodium chloride USP. Purified water USP (2 L) was added to the stainless steel vessel and the contents were mixed with an overhead stirrer at a speed of 240 rpm for 10 minutes. Formoterol fumarate dihydrate (0.17 g for low dosage strength formulation, 0.34 g for high dosage strength formulation) was added and 25 the solution was stirred at 240 rpm for 90 minutes.

-31-

EXAMPLE 2

Preparation of Formoterol Unit Dose Formulations

Following the procedure of EXAMPLE 1, the following formoterol unit dose formulations were prepared.

5 **Low Strength (0.0085%)**

A low strength formoterol unit dose formulation was prepared using the following reagents in the amounts indicated: formoterol fumarate dihydrate (0.170 mg), citric acid monohydrate, USP (0.68 mg), sodium citrate dihydrate, USP (1.99 mg), sodium chloride, USP (17.5
10 mg), and purified water, USP (qs to 2 mL).

High Strength (0.0170%)

A high strength formoterol unit dose formulation was prepared using the following reagents in the amounts indicated: formoterol fumarate dihydrate (0.340 mg), citric acid monohydrate, USP (0.68 mg),
15 sodium citrate dihydrate, USP (1.99 mg), sodium chloride, USP (17.5 mg), and purified water, USP (qs to 2 mL).

EXAMPLE 3

Procedure for stability testing of formoterol solutions

Stability samples of the solutions prepared in EXAMPLES 1 and 2
20 were placed in scintillation vials with teflon-lined caps and stored in stability ovens at accelerated temperatures. At selected time points, aliquots of the samples were removed from the scintillation vials. The formoterol concentrations of the samples were analyzed by high performance liquid chromatography.

25

Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

50728-5

-32-

CLAIMS:

1. A pharmaceutical composition, comprising formoterol, or a salt, ester, enol-ester, acid, or base thereof, in a pharmacologically acceptable fluid, wherein the
5 composition is stable during long term storage, the fluid comprises water, and the composition is formulated at a concentration for direct administration to a subject in need thereof.
2. The pharmaceutical composition of claim 1, wherein
10 the composition has an estimated shelf-life of greater than 1 month usage time at 25°C and greater than or equal to 1 year storage time at 5°C.
3. The pharmaceutical composition of claim 1 or 2, wherein greater than about 80% of the initial formoterol is
15 present after 1 month usage time at 25°C and 1 year storage time at 5°C.
4. The pharmaceutical composition of any one of claims 1-3, wherein the pharmacologically acceptable fluid comprises a polar solvent.
- 20 5. The pharmaceutical composition of claim 4, wherein the polar solvent is a protic solvent.
6. The pharmaceutical composition of any one of claims 1-5, further comprising a tonicity adjusting agent.
7. The pharmaceutical composition of claim 6, wherein
25 the tonicity adjusting agent is ammonium carbonate, ammonium chloride, ammonium lactate, ammonium nitrate, ammonium phosphate, ammonium sulfate, ascorbic acid, bismuth sodium tartrate, boric acid, calcium chloride, calcium disodium edetate, calcium gluconate, calcium lactate, citric acid,
30 dextrose, diethanolamine, dimethylsulfoxide, edetate disodium, edetate trisodium monohydrate, fluorescein sodium,

50728-5

-33-

fructose, galactose, glycerin, lactic acid, lactose, magnesium chloride, magnesium sulfate, mannitol, polyethylene glycol, potassium acetate, potassium chlorate, potassium chloride, potassium iodide, potassium nitrate, potassium phosphate, potassium sulfate, propylene glycol, silver nitrate, sodium acetate, sodium bicarbonate, sodium biphosphate, sodium bisulfite, sodium borate, sodium bromide, sodium cacodylate, sodium carbonate, sodium chloride, sodium citrate, sodium iodide, sodium lactate, sodium metabisulfite, sodium nitrate, sodium nitrite, sodium phosphate, sodium propionate, sodium succinate, sodium sulfate, sodium sulfite, sodium tartrate, sodium thiosulfate, sorbitol, sucrose, tartaric acid, triethanolamine, urea, urethane, uridine or zinc sulfate.

8. The pharmaceutical composition of claim 6 or 7, wherein the tonicity adjusting agent is sodium chloride.

9. The pharmaceutical composition of any one of claims 1-3, wherein the pharmacologically acceptable fluid comprises a buffer.

10. The pharmaceutical composition of any one of claims 4-8, further comprising a buffer.

11. The pharmaceutical composition of claim 9 or 10, wherein the buffer is citric acid/phosphate, acetate, barbital, borate, Britton-Robinson, cacodylate, citrate, collidine, formate, maleate, McIlvaine, phosphate, Prideaux-Ward, succinate, citrate-phosphate-borate (Teorell-Stanhagen), veronal acetate, MES (2-(N-morpholino)-ethanesulfonic acid), BIS-TRIS (bis(2-hydroxyethyl)-iminotris(hydroxymethyl)methane), ADA (N-(2-acetamido)-2-iminodiacetic acid), ACES (N-(carbamoylmethyl)-2-aminoethanesulfonic acid), PIPES (piperazine-N,N'-bis(2-ethanesulfonic acid)), MOPSO (3-(N-morpholino)-

50728-5

-34-

2-hydroxypropanesulfonic acid), BIS-TRIS PROPANE
 (1,3-bis(tris(hydroxymethyl)methylamino)propane),
 BES (N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid),
 MOPS (3-(N-morpholino)propanesulfonic acid),
 5 TES (N-tris(hydroxymethyl)methyl-2-aminoethanesulfonic
 acid), HEPES (N-(2-hydroxyethyl)piperazine-
 N'-(2-ethanesulfonic acid), DIPSO (3-(N,N-bis-
 (2-hydroxyethyl)amino)-2-hydroxypropanesulfonic acid),
 MOBS (4-(N-morpholino)butanesulfonic acid),
 10 TAPSO (3-(N-tris(hydroxymethyl)methylamino)-
 2-hydroxypropanesulfonic acid), tris(hydroxymethylamino-
 methane, HEPPSO (N-(2-hydroxyethyl)piperazine-
 N'-(2-hydroxypropanesulfonic acid), POPSO (piperazine-
 N,N'-bis(2-hydroxypropanesulfonic acid)),
 15 TEA (triethanolamine), EPPS (N-(2-hydroxyethyl)piperazine-
 N'-(3-propanesulfonic acid), TRICINE (N-tris(hydroxymethyl)-
 methylglycine), GLY-GLY (glycylglycine),
 BICINE (N,N-bis(2-hydroxyethyl)glycine),
 HEPBS (N-(2-hydroxyethyl)piperazine-N'-(4-butanefulfonic
 20 acid)), TAPS (N-tris(hydroxymethyl)methyl-3-
 aminopropanesulfonic acid), or AMPD (2-amino-2-methyl-
 1,3-propanediol) buffer.

12. The pharmaceutical composition of claim 10 or 11,
 wherein the buffer is citrate buffer.

25 13. The pharmaceutical composition of any one of
 claims 10-12, wherein the buffer concentration is from about
 0.01 mM to about 150 mM.

14. The pharmaceutical composition of any one of
 claims 10-13, wherein the buffer concentration is from about
 30 1 mM to about 20 mM.

50728-5

-35-

15. The pharmaceutical composition of any one of claims 10-14, wherein the buffer concentration is about 5 mM.

16. The pharmaceutical composition of any one of claims 1-15, wherein the ionic strength of the composition is about 0 to about 0.4.

17. The pharmaceutical composition of any one of claims 1-16, wherein the ionic strength of the composition is about 0.05 to about 0.16.

18. The pharmaceutical composition of any one of claims 1-17, wherein the pH of the composition is about 2.0 to about 8.0.

19. The pharmaceutical composition of any one of claims 1-18, wherein the pH of the composition is about 4.0 to about 6.0.

20. The pharmaceutical composition of any one of claims 1-19, wherein the pH of the composition is about 4.5 to about 5.5.

21. The pharmaceutical composition of any one of claims 1-20, wherein the pH of the composition is about 5.0.

22. The pharmaceutical composition of any one of claims 1-21, wherein the formoterol free base concentration is about 5 µg/mL to about 2 mg/mL.

23. The pharmaceutical composition of any one of claims 1-22, wherein the formoterol free base concentration is about 10 µg/mL to about 1 mg/mL.

24. The pharmaceutical composition of any one of claims 1-23, wherein the formoterol free base concentration is about 50 µg/mL to about 200 µg/mL.

50728-5

-36-

25. The pharmaceutical composition of any one of claims 1-24, wherein the formoterol free base concentration is about 59 µg/mL.

26. The pharmaceutical composition of any one of
5 claims 1-24, wherein the formoterol free base concentration is about 118 µg/mL.

27. The pharmaceutical composition of any one of claims 1-26, further comprising one or more of (a) to (j) as follows: (a) a β_2 -adrenoreceptor agonist; (b) a dopamine
10 (D₂) receptor agonist; (c) an IL-5 inhibitor; (d) an antisense modulator of IL-5; (e) a tryptase inhibitor; (f) a tachykinin receptor antagonist; (g) milrinone or milrinone lactate; (h) a leukotriene receptor antagonist; (i) a 5-lipoxygenase inhibitor; or (j) an anti-IgE antibody.

15 28. The pharmaceutical composition of claim 10 or 11, wherein the buffer comprises citric acid/phosphate buffer, acetate buffer, citrate buffer or phosphate buffer.

29. The pharmaceutical composition of any one of claims 10-13, wherein the buffer concentration is from about
20 1 mM to about 50 mM.

30. The pharmaceutical composition of claim 29, wherein the buffer concentration is about 20 mM.

31. The pharmaceutical composition of any one of claims 1-30, further comprising an anticholinergic agent.

25 32. The pharmaceutical composition of claim 31, wherein the anticholinergic agent is ipratropium bromide, oxitropium bromide, atropine methyl nitrate, tiotropium bromide or glycopyrronium bromide.

50728-5

-37-

33. The pharmaceutical composition of claim 31 or 32, wherein the anticholinergic agent is ipratropium bromide.

34. The pharmaceutical composition of claim 32 or 33, wherein the ipratropium bromide is present at a
5 concentration of about 5 $\mu\text{g/mL}$ to about 5 mg/mL .

35. The pharmaceutical composition of claim 31 or 32, wherein the anticholinergic agent is tiotropium bromide.

36. The pharmaceutical composition of claim 32 or 35, wherein the tiotropium bromide is present at a concentration
10 of about 5 $\mu\text{g/mL}$ to about 5 mg/mL .

37. The pharmaceutical composition of any one of claims 1-36 that has been nebulized.

38. A nebulized solution, comprising formoterol or a derivative thereof in a pharmacologically acceptable fluid.

15 39. A kit, comprising:

(a) a pharmaceutical composition, comprising formoterol, or a derivative thereof, in a pharmacologically acceptable fluid, wherein the composition is stable during long term storage, the fluid comprising water, and the
20 composition is formulated at a concentration for direct administration to a subject in need thereof; and

(b) a nebulizer.

40. The kit of claim 39, wherein the aqueous composition comprises (a) formoterol free base at a
25 concentration of about 59 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 5 mM ;

50728-5

-38-

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

41. The kit of claim 39, wherein the composition
5 comprises (a) formoterol free base at a concentration of about 118 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 5 mM;

wherein the ionic strength of the composition is
10 about 0.05 to about 0.16; and the pH of the composition is about 5.0.

42. A kit, comprising:

(a) a composition of any one of claims 1-36; and

(b) a nebulizer.

15 43. The kit of claim 39, wherein the composition comprises (a) formoterol free base at a concentration of about 59 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 20 mM;

20 wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

44. The kit of claim 39, wherein the composition
25 comprises (a) formoterol free base at a concentration of about 118 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 20 mM;

50728-5

-39-

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

45. A combination, comprising:

5 (a) the pharmaceutical composition of any one of claims 1-36 formulated for single dosage administration; and

(b) a vial.

46. The combination of claim 45, wherein the pharmaceutical composition comprises (a) formoterol free
10 base at a concentration of about 59 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 5 mM;

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is
15 about 5.0.

47. The combination of claim 45, wherein the aqueous composition comprises (a) formoterol free base at a concentration of about 118 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a
20 concentration of about 5 mM;

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

48. The combination of claim 45, wherein the aqueous
25 composition comprises (a) formoterol free base at a concentration of about 59 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 20 mM;

50728-5

-40-

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

49. The combination of claim 45, wherein the aqueous composition comprises (a) formoterol free base at a concentration of about 118 $\mu\text{g/mL}$; (b) aqueous saline comprising sodium chloride; and (c) citrate buffer at a concentration of about 20 mM;

wherein the ionic strength of the composition is about 0.05 to about 0.16; and the pH of the composition is about 5.0.

50. Use of the pharmaceutical composition of any one of claims 1-36 for the treatment, prevention, or amelioration of one or more symptoms of a bronchoconstrictive disorder.

51. The use of claim 50, wherein the pharmaceutical composition is for simultaneous or sequential use with one or more of (a) to (j) as follows: (a) a β_2 -adrenoreceptor agonist; (b) a dopamine (D_2) receptor agonist; (c) an IL-5 inhibitor; (d) an antisense modulator of IL-5; (e) a tryptase inhibitor; (f) a tachykinin receptor antagonist; (g) milrinone or milrinone lactate; (h) a leukotriene receptor antagonist; (i) a 5-lipoxygenase inhibitor; or (j) an anti-IgE antibody.

52. An article of manufacture, comprising packaging material, an aqueous composition comprising the composition of any one of claims 1-36 formulated for single dosage administration, and a label that indicates that the composition is used for treatment, prevention or amelioration of one or more symptoms of a bronchoconstrictive disorder.

50728-5

-41-

53. The pharmaceutical composition of any one of claims 1-36, when used for the treatment, prevention, or amelioration of one or more symptoms of a bronchoconstrictive disorder.

5 54. Use of the pharmaceutical composition of any one of claims 1-36 for the preparation of a medicament for the treatment, prevention, or amelioration of one or more symptoms of a bronchoconstrictive disorder.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS