



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2019/02/07
(87) **Date publication PCT/PCT Publication Date:** 2019/08/15
(85) **Entrée phase nationale/National Entry:** 2020/08/05
(86) **N° demande PCT/PCT Application No.:** US 2019/017116
(87) **N° publication PCT/PCT Publication No.:** 2019/157224
(30) **Priorités/Priorities:** 2018/02/07 (US62/627,721);
2018/12/10 (US62/777,683)

(51) **Cl.Int./Int.Cl.** **C07K 19/00** (2006.01),
A61K 47/68 (2017.01), **A61K 48/00** (2006.01),
A61K 9/00 (2006.01), **A61P 3/00** (2006.01),
C07K 16/28 (2006.01), **C07K 16/46** (2006.01),
C12N 15/35 (2006.01), **C12N 15/62** (2006.01),
C12N 15/86 (2006.01), **C12N 15/864** (2006.01),
C12N 9/00 (2006.01), **C12N 9/24** (2006.01)

(71) **Demandeur/Applicant:**
REGENERON PHARMACEUTICALS, INC., US

(72) **Inventeurs/Inventors:**
BAIK, ANDREW, US;
CYGNAR, KATHERINE, US;
PRAGGASTIS, MARIA, US

(74) **Agent:** CPST INTELLECTUAL PROPERTY INC.

(54) **Titre : PROCÉDES ET COMPOSITIONS POUR L'ADMINISTRATION DE PROTÉINES THÉRAPEUTIQUES**

(54) **Title: METHODS AND COMPOSITIONS FOR THERAPEUTIC PROTEIN DELIVERY**

(57) **Abrégé/Abstract:**

Compositions and methods for delivering a therapeutic protein to the central nervous system (CNS), in order to treat diseases and disorders that impair the CNS, such as treating lysosomal storage diseases are disclosed. Therapeutic proteins delivered via a therapeutically effective amount of a nucleotide composition encoding the therapeutic protein conjugated to a cell surface receptor-binding protein that crosses the blood brain barrier (BBB) are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

15 August 2019 (15.08.2019)



(10) International Publication Number

WO 2019/157224 A1

(51) International Patent Classification:

A61K 47/68 (2017.01) C12N 9/24 (2006.01)

A61K 38/47 (2006.01) A61P 3/00 (2006.01)

A61K 39/395 (2006.01) A61P 43/00 (2006.01)

A61K 48/00 (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2019/017116

(22) International Filing Date:

07 February 2019 (07.02.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/627,721 07 February 2018 (07.02.2018) US

62/777,683 10 December 2018 (10.12.2018) US

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))
- with sequence listing part of description (Rule 5.2(a))

(71) Applicant: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(72) Inventors: **BAIK, Andrew**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **CYGNAR, Katherine**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **PRAGGASTIS, Maria**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(74) Agent: **WU, Rita S.**; FISHERBROYLES, LLP, 945 East Paces Ferry Road Suite 2000, Atlanta, Georgia 30326 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: METHODS AND COMPOSITIONS FOR THERAPEUTIC PROTEIN DELIVERY

(57) Abstract: Compositions and methods for delivering a therapeutic protein to the central nervous system (CNS), in order to treat diseases and disorders that impair the CNS, such as treating lysosomal storage diseases are disclosed. Therapeutic proteins delivered via a therapeutically effective amount of a nucleotide composition encoding the therapeutic protein conjugated to a cell surface receptor-binding protein that crosses the blood brain barrier (BBB) are provided.

WO 2019/157224 A1

METHODS AND COMPOSITIONS FOR THERAPEUTIC PROTEIN DELIVERY

REFERENCE TO A SEQUENCE LISTING SUBMITTED AS A TEXT FILE VIA EFS- WEB

[0001] The Sequence Listing written in file 10457WO01_ST25.txt is 33.5 kilobytes, was created on January 29, 2019, and is hereby incorporated by reference.

FIELD

[0002] This application is generally directed to compositions and methods for delivering a therapeutic protein to the central nervous system (CNS), in order to treat diseases and disorders that impair the CNS, such as treating lysosomal storage diseases. This application is directed to providing a therapeutically effective amount of a nucleotide composition encoding a therapeutic protein conjugated to one or more delivery domains that crosses the blood brain barrier (BBB).

BACKGROUND

[0003] Drug delivery approaches have been developed to overcome the blood brain barrier (BBB), such as nanocarriers, however have shortcomings. Carriers have exhibited instability in blood circulation and undesirable bio-distribution profile (Gelperina S, et al., 2005, *Am J Respir Crit Care Med.* 172(12):1487-90; which reference is incorporated herein in its entirety by reference). Targeting efficiencies have also been compromised depending on the trafficking mechanisms at the BBB and whether a CNS disease state has altered the integrity of the barrier. Proper selection of the targeting moiety or carrier must take into account neuroinflammatory conditions that effect these trafficking mechanisms.

[0004] Delivery of therapeutic proteins via DNA expression in the liver or other tissues has provided a convenient approach eliminating the need for bolus injection of protein and therefore lessening immunogenicity concerns. Therapeutic protein conjugated to a receptor binding protein, especially a cell specific receptor, solves the problem of targeting therapeutics to specific tissues. However, there is still a need to provide methods that efficiently provide therapeutics to the CNS.

SUMMARY

[0005] Applicants have discovered that therapeutic proteins, especially replacement enzymes, can be effectively delivered into the central nervous system when associated with a receptor binding protein and provided that the circulating blood levels achieve consistent levels over

time. The multidomain therapeutic protein can be delivered to the liver via a gene therapy vector harboring the coding sequence of the therapeutic protein and binding protein complex.

[0006] In one aspect, the invention provides a method of delivering a therapeutic protein to the central nervous system (CNS) of a subject, comprising administering to the subject a nucleotide composition encoding the therapeutic protein conjugated to a cell surface receptor (CSR)-binding protein (CSR-BP) (tpCSR-BP) via a liver-targeted delivery method sufficient to provide a therapeutically effective amount of the tpCSR-BP in the CNS.

[0007] In one embodiment the CSR-BP is an antibody or antigen-binding fragment thereof that binds specifically to the CSR. In another embodiment, the therapeutic protein is a lysosomal enzyme.

[0008] In one embodiment, the enzyme has hydrolase activity, such as a glycosylase, such as a glycosidase, such as an alpha-glucosidase or alpha-galactosidase A. In one embodiment, the cell surface receptor (CSR)-binding protein (CSR-BP) is an antigen-binding protein that binds to an internalization receptor. In one embodiment, the internalization receptor is a cell-surface molecule that is endocytosed and trafficked to the lysosome. In a specific embodiment, the internalization receptor is a CD63 molecule. In one embodiment, the internalization receptor is a ITGA7 molecule. In a specific embodiment, the CSR-BP is an antibody, an antibody fragment, or a single-chain variable fragment (scFv), such as an scFv that binds CD63 or ITGA7.

[0009] In some embodiments, a multidomain therapeutic protein described herein comprises one or more delivery domain(s) and an enzyme domain, wherein the one or more delivery domain(s) binds human transferrin receptor (hTfR). In some embodiments, the multidomain therapeutic protein further comprises a second delivery domain that binds to an internalizing effector. In some embodiments, the second delivery domain binds to (i) an internalizing effector selected from the group consisting of CD63, Integrin alpha-7 (ITGA7), MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), myelin and lymphocyte protein (MAL), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptor A1-5 (SCARA1-5), SCARB1-3, and CD36; (ii) an internalizing effector expressed in several tissue types, optionally selected from the group consisting of CD63, MHC-I, vacuolar-type H⁺ ATPase, IGF2R, Integrin alpha-7 (ITGA7), LRP5, LRP6,

LRP8, Kremen-2, LDL- receptor, LDL-related protein 1 receptor, amyloid precursor protein-like protein- 2 (APLP2), apelin receptor (APLNR), PRLR, MAL (myelin and lymphocyte protein (MAL)), diphtheria toxin receptors, HBEGF (heparin binding EGF like growth factor), glutathione receptors, glutamate receptors, leptin receptors, and folate receptors; (iii) an internalizing effector preferentially expressed by bone and/or cartilage, optionally selected from the group consisting of Collagen X, Integrin alpha 10 (ITGA10), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor isoform C (FGFR3C), Hyaluronan and proteoglycan link protein 1 (CRTL1), Aggrecan, Collagen II, and Kremen-1; (iv) an internalizing effector preferentially expressed by monocytes, macrophages, or microglia, optionally selected from the group consisting of scavenger receptor A1-5 (SCARA1-5), SCARB1-3, CD36, MSR1 (macrophage scavenger receptor 1), MRC1 (macrophage mannose receptor 1), VSIG4 (V-set and immunoglobulin domain-containing protein 4), CD68 (Macrosialin), and CSF1R (Macrophage colony-stimulating factor 1 receptor); (v) an internalizing effector preferentially expressed by kidney cells, optionally selected from the group consisting of CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin). In other certain embodiments, the internalization effector is a muscle specific internalizer, such as BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNAIS (L-type calcium channel subunit alpha-15), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3); (vi) an internalizing effector preferentially expressed by liver cells, optionally ASGR1 or ASGR2; (vii) an internalizing effector preferentially expressed by muscle cells, optionally selected from the group consisting of BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNAIS (L-type calcium channel subunit alpha-15), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye

domain-containing protein 3), and/or (viii) an internalizing effector protein selected from the group consisting of ITGA7, CD9, CD63, ALPL2, MSR1, ASGR1, ASGR2, or PRLR. In some embodiments, the second delivery domain binds to the internalizing effector CD63. In some embodiments, at least one of the one or more delivery domain(s) comprises an antigen-binding protein. In some embodiments, each of the one or more delivery domain(s) comprises an antigen-binding protein. In some embodiments, at least one of the one or more delivery domain(s) comprises a single-chain variable fragment (scFv). In some embodiments, at least one of the one or more delivery domain(s) comprises a half-body. In some embodiments, the delivery domain that binds hTfR is an scFv, wherein the half-body binds CD63, wherein the enzyme domain is GAA, and wherein GAA is conjugated to the carboxy terminus of the half-body that binds CD63. In some embodiments, each of the one or more delivery domain(s) comprises an scFv. In some embodiments, at least one scFv is fused to an Fc. In some embodiments, the Fc comprises a wildtype human IgG4 isotype, or derivative thereof. In some embodiments, GAA is conjugated to the carboxy terminus of the Fc. In some embodiments, the multidomain therapeutic protein comprises an anti-hTfR scFv and an anti hCD63 scFv. In some embodiments, the anti hTfR scFv and anti hCD63 scFv are both linked, at their carboxy termini, to a single GAA enzyme. In some embodiments, the delivery domain is an anti-hTfR scFv, and the enzyme domain is linked to carboxy terminus of the VL domain of the scFv. In some embodiments, the multidomain further comprises a second delivery domain linked to the N-terminus of the VH domain of the anti-hTfR scFv. In some embodiments, the second delivery domain is an anti-hCD63 scFv. In some embodiments, the enzyme domain comprises the amino acid sequence set forth as SEQ ID NO:1

[0010] Also provided are multidomain therapeutic proteins comprising at least two delivery domains and at least one enzyme domain, wherein each of the two delivery domains is independently selected from the group consisting of an antibody, a half-body, and an scFv, and wherein at least one or more of the delivery domains is associated the at least one enzyme domain, preferably wherein the one or more delivery domains is covalently linked to the at least one enzyme domain. In some embodiments, the multidomain therapeutic protein comprises no more than two delivery domains. In some embodiments, only one of the delivery domains is associated with the at least one enzyme domain. In some embodiments, each of the at least two delivery domains is covalently linked to an enzyme domain. In some embodiments, each of the at least two delivery domains is covalently linked to the same enzyme domain. In some embodiments, each of the at least two delivery domains is covalently linked to a different

enzyme domain. In some embodiments, the multidomain therapeutic protein comprises no more than two delivery domains, wherein the first delivery domain comprises a half-body, and wherein the second delivery domain comprises an scFv. In some embodiments, the scFv is fused to an Fc. In some embodiments, the half-body is covalently linked at its carboxy terminus to a first enzyme domain and/or wherein the scFv is covalently linked at its carboxy terminus to an Fc, and optionally, a second enzyme domain. In some embodiments, the multidomain therapeutic protein comprises no more than two delivery domains, wherein the first and second delivery domains each comprise an scFv. In some embodiments, both the first and second scFv are covalently linked to an enzyme domain. In some embodiments, the multidomain therapeutic protein comprises from N-terminus to C-terminus: the first scFv, the second scFv, and the enzyme domain. In some embodiments, at least one delivery domain binds a lysosomal trafficking molecule and at least one delivery domain binds a transcytosis effector. In some embodiments, the lysosomal trafficking molecule is selected from the group consisting of CD63, ITGA7, CD9, CD63, CD81, CD82, or CD151, and wherein the transcytosis effector is selected from the group consisting of an LDL receptor, an IgA receptor, a transferrin receptor, a neonatal Fc receptor, insulin receptor, CD98, and Basigin. In some embodiments, the multidomain therapeutic protein comprises a structure as depicted in Figure 1C, Figure 1D, Figure 1E, or Figure 1F.

[0011] Also provided herein are polynucleotides that encode the multidomain therapeutic proteins described herein. In some embodiments, a polynucleotide provided herein further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence. In some embodiments, the polynucleotide further comprises a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence. In some embodiments, the polynucleotide further comprises a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence, and wherein the AAV nucleic acid sequence comprises an internal terminal repeat sequence, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter. In some embodiments, the polynucleotide further comprises a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence comprising an internal terminal repeat sequence that comprises SEQ ID NO:6, SEQ ID NO:7, or both, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter. In some

embodiments, the polynucleotide further comprises a tissue specific regulatory element comprising the sequence set forth as SEQ ID NO:8 and/or SEQ ID NO:9.

[0012] In one aspect, the invention provides a gene therapy vector, such as an AAV vector, that contains a nucleic acid sequence encoding a therapeutic protein conjugated to or fused to a CSR-BP, e.g., a polynucleotide as described herein. In some embodiments, the gene therapy vector is selected from the group consisting of a viral vector, optionally wherein the viral vector is a natural virus, an engineered virus, or a chimeric virus, and a naked polynucleotide comprising a polynucleotide described herein, a polynucleotide complex, optionally wherein the polynucleotide complex is a lipid nanoparticle comprising the polynucleotide of any one of claims 20-25 and lipids, and any combination thereof. In some embodiments, the gene therapy vector is a viral vector selected from the group consisting of a retrovirus, adenovirus, herpes simplex virus, pox virus, vaccinia virus, lentivirus, or an adeno-associated virus. In some embodiments, the gene therapy vector is AAV9, Anc80, an AAV2/8 chimera and/or an AAV pseudotyped to a specific tissue, e.g., the liver or neuronal tissue.

[0013] In one embodiment, a therapeutic protein, nucleotide encoding same, and/or gene therapy vectors comprising the nucleotide encoding same is used to treat a subject in need of enzyme replacement therapy, e.g., in a method of delivering a therapeutic protein to the central nervous system (CNS) of a subject, comprising administering to the subject a nucleotide composition encoding a multidomain therapeutic protein via a liver-targeted delivery method sufficient to provide a therapeutically effective amount of the multidomain therapeutic protein in the CNS, wherein the multidomain therapeutic protein comprises a delivery domain and an enzyme domain. In some embodiments, the subject is an animal. In some embodiments, the subject is a human.

[0014] In one aspect, an AAV vector containing a polynucleotide encoding an scFv-hydrolase fusion protein is administered to a human or non-human subject. The polynucleotide subsequently integrates at a genomic locus in the liver and the encoded fusion protein is produced. In another embodiment, the polynucleotide is transcribed episomally in the liver and the encoded fusion protein is produced. In a specific embodiment, the fusion protein is an anti-CD63scFv-GAA fusion protein or an anti-ITGA7scFv-GAA fusion protein, the human or non-human subject lacks endogenous GAA activity, and the GAA activity is effectively restored in the subject.

[0015] In one aspect, the invention provides a method of treating a subject (human or non-human) with an enzyme deficiency by administering to the patient a gene therapy vector containing a gene encoding a therapeutic protein conjugated to or fused to a CSR-BP.

[0016] Described herein are methods of delivering a therapeutic protein to the central nervous system (CNS) of a subject, comprising administering to the subject a nucleotide composition encoding a multidomain therapeutic protein via a liver-targeted delivery method sufficient to provide a therapeutically effective amount of the multidomain therapeutic protein in the CNS, wherein the multidomain therapeutic protein comprises a delivery domain and an enzyme domain. In some embodiments, the delivery domain is an antibody or antigen-binding fragment thereof that binds specifically to an internalizing effector. In some embodiments, the therapeutic protein is a lysosomal enzyme. In some embodiments, the lysosomal enzyme is GAA. In some embodiments, the nucleotide composition is administered via a viral vector. In some embodiments, the viral vector is an AAV vector. In some embodiments, the nucleotide composition is administered at a dose of at least 2×10^{12} viral genomes per kilogram (vg/kg). In some embodiments, the internalizing effector is expressed on the surface of cells selected from the group consisting of: cells in the CNS, epithelial cells, and cells that cross the blood brain barrier. In some embodiments, the delivery domain binds an internalizing effector. In some embodiments, the internalizing effector is (i) selected from the group consisting of CD63, Integrin alpha-7 (ITGA7), MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), myelin and lymphocyte protein (MAL), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptor A1-5 (SCARA1-5), SCARB1-3, and CD36; (ii) expressed in several tissue types, e.g., CD63, MHC-I, vacuolar-type H⁺ ATPase, IGF2R, Integrin alpha-7 (ITGA7), LRP5, LRP6, LRP8, Kremen-2, LDL- receptor, LDL-related protein 1 receptor, amyloid precursor protein-like protein- 2 (APLP2), apelin receptor (APLNR), PRLR, MAL (myelin and lymphocyte protein (MAL), diphtheria toxin receptors, HBEGF (heparin binding EGF like growth factor), glutathione receptors, glutamate receptors, leptin receptors, and folate receptors, optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of Fabry disease, Gaucher disease, MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVB, MPS VI, MPS VII, MPS IX, Pompe disease, Lysosomal acid lipase deficiency, Metachromatic leukodystrophy, Niemann-Pick diseases types A, B, and C2, Alpha mannosidosis, Neuraminidase deficiency,

Sialidosis, Aspartylglycosaminuria, Combined saposin deficiency, Atypical Gaucher disease, Farber lipogranulomatosis, Fucosidosis, and Beta mannosidosis; (iii) preferentially expressed by bone and/or cartilage, e.g., Collagen X, Integrin alpha 10 (ITGA10), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor isoform C (FGFR3C), Hyaluronan and proteoglycan link protein 1 (CRTL1), Aggrecan, Collagen II, and Kremen-1, optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVA, MPS IVB, MPS VI, MPS VII, MPS IX, Beta mannosidosis, Gaucher disease, atypical Gaucher disease, combined Saposin deficiency, Aspartylglycosaminuria, Farber lipogranulomatosis, Sialidosis, Neuraminidase deficiency, and Alpha mannosidosis; (iv) preferentially expressed by monocytes, macrophages, or microglia, e.g., scavenger receptor A1-5 (SCARA1-5), SCARB1-3, CD36, MSR1 (macrophage scavenger receptor 1), MRC1 (macrophage mannose receptor 1), VSIG4 (V-set and immunoglobulin domain-containing protein 4), CD68 (Macrosialin), and CSF1R (Macrophage colony-stimulating factor 1 receptor), optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of lysosomal acid lipase deficiency, Gaucher disease, Atypical Gaucher disease, combined Saposin deficiency, and Farber lipogranulomatosis; (v) preferentially expressed by kidney cells, e.g., CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin), optionally wherein the subject exhibits one or more symptoms or is diagnosed with a disease selected from the group consisting of Fabry disease, Alport syndrome, polycystic kidney disease, and Thrombotic Thrombocytopenic Purpura; (vi) preferentially expressed by liver cells, e.g., ASGR1 or ASGR2, optionally wherein the subject exhibits one or more symptoms or is diagnosed with a disease selected from the group consisting of as lysosomal acid lipase deficiency, Gaucher disease, MPS VI, MPS VII, MPS II, Niemann-Pick diseases types A, B, and C2, Sialidosis, Neuraminidase deficiency, atypical Gaucher disease, combined Saposin deficiency, Farber lipogranulomatosis; (vii) preferentially expressed by muscle cells, e.g., BMPR1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-15), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing

protein 3), optionally wherein the subject exhibits one or more symptoms or is diagnosed with Pompe disease; (viii) selected from the group consisting of ITGA7, CD9, CD63, ALPL2, MSR1, ASGR1, ASGR2, or PRLR; and/or (ix) is CD63. In some embodiments, the delivery domain is a single-chain variable fragment (scFv). In some embodiments, the cell surface receptor (CSR)-binding protein (CSR-BP) comprises an amino acid sequence of SEQ ID NO:2. In some embodiments, the therapeutic protein comprises a hydrolase. In some embodiments, the therapeutic protein comprises a glycosylase. In some embodiments, the therapeutic protein comprises a glycosidase. In some embodiments, the therapeutic protein comprises an alpha-glucosidase. In some embodiments, the therapeutic protein comprises an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:13, or a fragment thereof. In some embodiments, the therapeutic protein comprises an anti-ABeta, or an anti-Tau antibody. In some embodiments, the polynucleotide comprises a nucleic acid sequence of SEQ ID NO:11. In some embodiments, the enzyme domain comprises an alpha-glucosidase, and wherein the glycogen levels in any CNS tissue in the subject is reduced for at least nine months post-treatment. In some embodiments, the subject has Pompe disease. In some embodiments, the administered nucleotide composition provides a multidomain therapeutic protein serum level of at least 1 µg/mL. In some embodiments, the therapeutic protein comprises a glycosidase, such as GAA (*e.g.*, SEQ ID NO:1) or GLA (*e.g.*, UniProtKB No. P06280, aa32-429, SEQ ID NO:13), and the patient has Pompe disease or Fabry disease. In some embodiments, the CSR-BP is an antigen-binding protein that binds to an internalization receptor, such as CD63 or ITGA7. In some embodiments, the CSR-BP is an scFv molecule that binds CD63. In some embodiments, the CSR-BP is an scFv molecule that binds ITGA7. In some embodiments, the gene therapy vector is an AAV vector comprising a polynucleotide that encodes an anti-CD63-GAA fusion therapeutic protein. In some embodiments, the gene therapy vector is an AAV vector comprising a polynucleotide that encodes an anti-ITGA7-GAA fusion therapeutic protein.

[0017] In some embodiment, the therapeutic protein comprises a GAA enzyme domain, and high serum levels of GAA are maintained in the serum of the patient for at least 12 weeks after administering the gene therapy vector. In some embodiments, the therapeutic protein comprises a GAA enzyme, and glycogen levels in CNS tissue in the patient are significantly reduced. In some embodiments, the therapeutic protein comprises a GAA enzyme, and glycogen levels are maintained at wildtype levels 3 months, 6 months, or 9 months after administration of the gene therapy vector. In some embodiments, the therapeutic protein comprises a GAA enzyme, and the muscle strength of the patient after treatment is restored to wildtype levels.

[0018] In one aspect, the invention provides a method of reducing glycogen accumulation in a tissue, particularly a CNS tissue, in a human or non-human subject by administering a gene therapy vector containing a polynucleotide that encodes a therapeutic protein fused to a CSR-BP. In some embodiments, the gene therapy vector is administered at a dose sufficient to provide a threshold serum level of the therapeutic protein fused to a CSR-BP. In some embodiments, the threshold level is at least 1 µg/mL. In some embodiments, the threshold level is at least 2 µg/mL. In some embodiments, the threshold level is at least 3 µg/mL. In some embodiments, the threshold level is at least 4 µg/mL. In some embodiments, the threshold level is at least 5 µg/mL. In some embodiments, the threshold level is at least 6 µg/mL. In some embodiments, the threshold level is at least 7 µg/mL. In some embodiments, the threshold level is at least 8 µg/mL. In some embodiments, the threshold level is at least 9 µg/mL. In some embodiments, the threshold level is at least 10 µg/mL. In some embodiments, the threshold level is at least 11 µg/mL. In some embodiments, the threshold level is at least 12 µg/mL. In some embodiments, the threshold level is at least 13 µg/mL. In some embodiments, the threshold level is at least 14 µg/mL. In some embodiments, the threshold level is at least 15 µg/mL. In one embodiment, the tissue is cerebellum, spinal cord or hippocampus. In one embodiment, the human or non-human subject has Pompe disease. In one embodiment, the therapeutic protein comprises an anti-CD63 scFv-GAA fusion protein. In another embodiment, the therapeutic protein comprises an anti-ITGA7 scFv-GAA fusion protein.

DRAWINGS

[0019] **Figure 1A** schematically represents multidomain therapeutic proteins. Panel A depicts a multidomain therapeutic protein comprising a bispecific antibody (ii) and a replacement enzyme (i). Panel B depicts an enzyme-Fc fusion polypeptide (i) associating with an internalizing effector-specific half-body (ii) to form the multidomain therapeutic protein. Panel C depicts a replacement enzyme (hexagon) covalently linked to the C-terminus of the heavy chain of an anti-internalizing effector antibody. Panel D depicts a replacement enzyme (hexagon) covalently linked to the N-terminus of the heavy chain of an anti-internalizing effector antibody. Panel E depicts a replacement enzyme (hexagon) covalently linked to the C-terminus of the light chain of an anti-internalizing effector antibody. Panel F depicts a replacement enzyme (hexagon) covalently linked to the N-terminus of the light chain of an anti-internalizing effector antibody. Panel G depicts a replacement enzyme (hexagon) covalently linked to the C-terminus of a single-chain variable fragment (scFv) containing a VH region (shaded bar) and a VL region (open bar). Panel H depicts a replacement enzyme (hexagon) covalently linked to two scFv

domains, the first scFv (i) which serves as a first delivery domain, and the second scFv (ii) which serves as a second delivery domain. Additional multidomain therapeutic proteins not depicted in Figure 1A include, but are not limited to, multidomain therapeutic proteins comprising two or more delivery domains and at least one enzyme domain. As non-limiting examples, the antibodies, half-bodies and scFv domains depicted in panels A-H of this figure may represent any type of delivery domain, and additional delivery domains or replacement enzymes can be also associated to make a multidomain therapeutic protein. Non-limiting examples of multidomain therapeutic proteins comprising two or more delivery domains are further depicted in Figures 1C, 1D, and 1F, which include a replacement enzyme (depicted as, but not limited to, GAA) covalently linked to a first internalizing effector-specific half-body, which associates with a second internalizing effector-specific scFv-Fc fusion, which may or may not also be covalently linked to a replacement enzyme (depicted as, but not limited to, GAA), to form the multidomain therapeutic protein (Figures 1C and 1D), a replacement enzyme (depicted as, but not limited to, GAA) covalently linked to the C-terminus of each of an anti-internalizing effector-specific half-body, which serves as a first delivery domain, and an internalizing effector-specific scFv-Fc fusion, which serves as a second delivery domain, where both the anti-internalizing effector-specific half-body and associate together to form the multidomain therapeutic protein (Figure 1D), and a replacement enzyme covalently linked to a first scFv, which is linked, e.g., via a linker, to a second scFv (Figure 1F).

[0020] Figure 1B provides non-limiting exemplary illustrations of AAV gene therapy vectors that each encode a multidomain therapeutic protein represented in panel G of Figure 1A, wherein the scFv is an anti-human CD63 scFv and the replacement enzyme is GAA (e.g., anti-hCD63scFv:hGAA; see, e.g., the amino acid sequence set forth as SEQ ID NO:10). Amino acids 1-117 of SEQ ID NO:10 provide the amino acid sequence of the heavy chain variable domain (V_H) of the H4H12450N antibody; amino acids 118-132 of SEQ ID NO:10 provide an amino acid linker sequence between the heavy and light chain variable domains of H4H12450N; amino acids 133-240 of SEQ ID NO:364 provide the amino acid sequence of the light chain variable domain (V_L) of the H4H12450N antibody; amino acids 241-245 of SEQ ID NO:10 provides an amino acid linker sequence between the anti-hCD63scFv and GAA; and amino acids 246-1128 of SEQ ID NO:10 provides the amino acid sequence of the replacement enzyme GAA, or biologically active portion thereof. Exemplary 5' ITR and 3' ITR sequences are respectively set forth as SEQ ID NO:6 and SEQ ID NO:7. Panel A of this Figure provides an exemplary vector for liver specific expression comprising an exemplary liver specific

enhancer (e.g., but not limited to, *Serpina1*; set forth as SEQ ID NO:9), an exemplary liver specific promoter (e.g., but not limited to, *TTR*; set forth as SEQ ID NO:8), an exemplary signal peptide ; a nucleic acid sequence encoding the anti-hCD63scFv::hGAA multitherapeutic domain (SEQ ID NO:10), and a poly-A tail. Panel B of this figure provides an exemplary vector similar to that shown in Panel A with an exemplary ubiquitous promoter in place of the liver specific enhancer and liver specific promoter sequences. Panel C of this figure provides an exemplary vector similar to that shown in Panel A with an exemplary neuron specific promoter in place of the liver specific enhancer (e.g., *Serpina1*) and promoter (e.g., *TTR*). Panel D of this figure provides an exemplary vector similar to that shown in Panel A with an exemplary neuron specific promoter in combination with a liver specific (e.g., *Serpina1*) enhancer and promoter (e.g., *TTR*).

[0021] Figure 1C provides non-limiting exemplary illustrations of expression vectors that each encode a multidomain therapeutic protein as depicted, wherein the half-body is an anti-CD63 antibody, the scFv is an anti-human transferrin receptor scFv, and wherein the replacement enzyme is GAA (e.g., anti-hTfRscFv::hGAA).

[0022] Figure 1D provides non-limiting exemplary illustrations of expression vectors that each encode a multidomain therapeutic protein as depicted, wherein the half-body is an anti-CD63 antibody, wherein the scFv is an anti-human transferrin receptor (TfR) scFv and the Fc domain is a human IgG4 Fc, and wherein the replacement enzyme is GAA (e.g., anti-hTfRscFv::hGAA).

[0023] Figure 1E provides non-limiting exemplary illustrations of expression vectors that each encode a multidomain therapeutic protein represented in Panel H of Figure 1A, wherein one of the two scFv is an anti-human CD63 scFv, the other of the two scFv is an anti-human transferrin receptor (TfR) scFv, and the replacement enzyme is GAA (e.g., anti-hCD63scFv::hGAA::anti-TfRscFv),

[0024] Figure 1F provides non-limiting exemplary illustrations of expression vectors that each encode a multidomain therapeutic protein as depicted, wherein one of the two scFv is an anti-human CD63 scFv, the other of the two scFv is an anti-human transferrin receptor (TfR) scFv, and the replacement enzyme is GAA (e.g., anti-hCD63scFv::anti-TfRscFv::GAA or anti-TfRscFv::anti-hCD63scFv::GAA).

[0025] Figure 1G provides non-limiting exemplary illustrations of expression vectors that each encode a multidomain therapeutic protein as depicted in panel G of Figure 1A, wherein the scFv

is an anti-human transferrin receptor (TfR) scFv and the replacement enzyme is GAA (e.g., anti-TfRscFv::GAA).

[0026] Figure 2 is a bar graph depicting the amount of stored glycogen in micrograms per milligram of tissue as a function of delivered enzyme. The X-axis depicts tissues from a CD63^{hu/hu}; GAA^{-/-} mouse from left to right: heart, quadriceps, gastrocnemius, diaphragm, soleus, and extensor digitorum longus (EDL) muscle. Lane 1 boxes depict the amount of stored glycogen in an untreated mouse Pompe disease model. Lane 6 boxes depict the amount of stored glycogen in an untreated wildtype mouse model. Lane 2 boxes depict the amount of stored glycogen in a mouse Pompe disease model treated with AAV-hGAA (adeno-associated virus vector containing gene encoding human GAA) at a dose of 10^{10} vg. Lane 3 boxes depict the amount of stored glycogen in a mouse Pompe disease model treated with AAV-hGAA at a dose of 10^{11} vg. Lane 4 boxes depict the amount of stored glycogen in a mouse Pompe disease model treated with AAV-anti-hCD63scFv::hGAA (adeno-associated virus vector containing gene encoding an anti-human CD63 scFv domain linked to human GAA) at a dose of 10^{10} vg. Lane 5 boxes depict the amount of stored glycogen in a mouse Pompe disease model treated with AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0027] Figure 3 is a graph depicting the average glycogen measured ($\mu\text{g}/\text{mg}$) in skeletal muscle tissue in each mouse at 3 months post-AAV injection. Each measurement is plotted as a function of GAA exposure (*i.e.*, serum levels) per mouse treated with a particular enzyme construct at a particular dosage. Filled squares represent AAV-hGAA at a dose of 10^{10} vg. Filled pyramids represent AAV-hGAA at a dose of 10^{11} vg. Filled inverse pyramids represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{10} vg. Filled diamonds represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0028] Figure 4 is a dot plot depicting the average cardiac muscle glycogen measured ($\mu\text{g}/\text{mg}$) in heart tissue at 3 months post-AAV injection as a function of GAA exposure (*i.e.*, serum levels), per mouse treated with a particular enzyme construct at a particular dosage. Filled squares represent AAV-hGAA at a dose of 10^{10} vg. Filled pyramids represent AAV-hGAA at a dose of 10^{11} vg. Filled inverse pyramids represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{10} vg. Filled diamonds represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0029] Figure 5 is a dot plot depicting anti-GAA antibody titers at 3 months post-AAV injection as a function of GAA exposure (*i.e.* serum levels), per mouse treated with a particular enzyme construct at a particular dosage. Open squares represent AAV-hGAA at a dose of 10^{10}

vg. Open circles represent AAV-hGAA at a dose of 10^{11} vg. Open diamonds represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{10} vg. Hexagons represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0030] Figure 6 is a dot plot depicting anti-GAA antibody titers at 3 months post-AAV injection as a function of enzyme construct and dose. Circles represent control mice receiving empty AAV vector. Squares represent AAV-hGAA at a dose of 10^{10} vg. Pyramids represent AAV-hGAA at a dose of 10^{11} vg. Inverse pyramids represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{10} vg. Diamonds represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0031] Figure 7A is a line graph depicting serum levels of GAA (arbitrary units “a.u.”; y-axis) as a function of time in weeks after gene therapy vector injection. Squares (bottom line) represent AAV-hGAA at a dose of 10^{10} vg. Pyramids (second from the top line) represent AAV-hGAA at a dose of 10^{11} vg. Inverse pyramids (third from the top line) represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{10} vg. Diamonds (top line) represent AAV-anti-hCD63scFv::hGAA at a dose of 10^{11} vg.

[0032] Figure 7B is a bar graph depicting mRNA ratios (hGAA mRNA relative to mGADPH mRNA) following administration of AAV constructs in CD63 HumIn GAA KO mice (GAA^{-/-}, CD63hu/hu mice) or GAA^{+/+}, CD63hu/hu mice, as such: (1) untreated control, (2) AAV-liver-specific promoter-hGAA (1×10^{10} vg), (3) AAV-liver-specific promoter-hGAA (1×10^{11} vg), (4) AAV-liver-specific promoter-anti-hCD63::hGAA (1×10^{10} vg), (5) AAV-liver-specific promoter-anti-hCD63::hGAA (1×10^{11} vg), or (6) untreated control (GAA^{+/+}, CD63hu/hu). Liver expression of GAA was detected for all injections of AAV construct.

[0033] Figure 7C is a plot graph comparing serum GAA level to RNA expression level of GAA for mice receiving the AAV encoding the fusion protein (squares) and mice receiving the AAV encoding GAA (both constructs provided a liver-specific promoter (LSP) to drive expression).

[0034] Figure 7D is a bar graph showing Huh-7 human hepatocytes transiently transfected with liver-specific promoter driven constructs encoding for hGAA, anti-hCD63 scFv::GAA (fusion construct), or a non-binding fusion construct scFv::GAA control. Both scFv::GAA fusion constructs had a higher ratio of protein in the secreted supernatant than hGAA alone 3 days after transfection. Addition of M6P into the supernatant during the experimental period to mitigate CI-MPR-mediated uptake did not affect the ratio. (* = $p < 0.05$, $n=3$).

[0035] **Figure 7E** is a bar graph depicting the amount of serum titer of GAA as a function of delivered vector. The X-axis depicts serum from KO mice humanized for CD63 (GAA^{-/-};CD63hu/hu) that were given plasmids encoding GAA or ScFv-GAA fusion from left to right: 1) Control (no treatment); 2) AAV-LSP-hGAA treatment (1e10vg/mouse); 3) AAV-LSP-hGAA treatment (1e11vg/mouse); 4) AAV-LSP-anti-CD63::hGAA treatment (1e10vg/mouse); and 5) AAV-LSP-anti-CD63::hGAA treatment (1e11vg/mouse).

[0036] **Figure 8** are fluorescent micrographs depicting the lamp1-stained lysosomes in mouse muscle fibers counter-stained with DAPI to reveal nuclei. Panels A and A1 depict quadriceps cells derived from an untreated wildtype (GAA^{+/+}) mouse and stained for lamp1 (panel A), and nuclei (panel A1). Panels B and B1 depict quadriceps cells derived from an untreated GAA null (GAA^{-/-}) mouse and stained for lamp1 (panel B), and nuclei (panel B1). Panels C and C1 depict quadriceps cells derived from a GAA^{-/-} mouse treated with an AAV-hGAA construct and stained for lamp1 (panel C), and nuclei (panel C1). Panels D and D1 depict quadriceps cells derived from a GAA^{-/-} mouse treated with an AAV-hCD63scFv::hGAA construct and stained for lamp1 (panel D), and nuclei (panel D1).

[0037] **Figure 9** depicts line graphs showing grip strength and Rotarod test performance of mice treated with either AAV-LSP hGAA or AAV-LSP anti-hCD63::hGAA. Accelerating Rotarod measurements (**A**) and forelimb grip strength measurements (**B**) of wild-type GAA mice (inverted triangle), untreated control (square), AAV-LSP-hGAA treatment (1e11vg/mouse) (triangle) or AAV-LSP-anti-hCD63::hGAA treatment (1e11vg/mouse) (circle) were taken at monthly intervals for 6 months. Error bars are +/- SD. N=8-10 for all groups.

[0038] **Figure 10A and Figure 10B** depict the use of other membrane proteins as guides, such as anti-ITGA7 (Integrin alpha-7) scFv fusion proteins to guide GAA. **Figure 10A** shows GAA activity (y-axis) of C2C12 mouse myoblasts incubated overnight with anti-mouse CD63-GAA or anti-mouse ITGA7-GAA with or without the presence of 5mM M6P. **Figure 10B** shows GAA KO mice humanized for CD63 (GAA^{-/-};CD63hu/hu) that were given plasmids encoding an scFv::GAA format of anti-hCD63::GAA (2) or a full-length IgG4::GAA format of anti-mouse integrin alpha-7 (3) by hydrodynamic delivery (HDD), then tissue glycogen levels were measured 3 weeks post-HDD. Untreated control mice, GAA^{-/-};CD63hu/hu (1) and untreated wild-type GAA control mice, GAA^{+/+};CD63hu/hu (4) were also tested for glycogen levels in the same tissues.

[0039] **Figure 11** illustrates serum clearance of a full-length anti-CD63 antibody fused to GAA (i.e. full-length IgG4 antibody) in CD63hu/hu mice compared to WT CD63^{+/+} mice. Plasmids expressing the heavy and light chain of an anti-CD63 antibody fused to GAA were injected via the tail vein of each mouse. The serum pK of anti-CD63(full length IgG4)::GAA was observed to clear from serum within 24 hours.

[0040] **Figure 12A** is a dot plot depicting serum levels of GAA (arbitrary units “a.u.”; y-axis) at one-month post-AAV injection as a function of as a function of enzyme construct and dose. Squares represent AAV-LSP-Δ8GAA. Pyramids represent AAV-anti-hCD63scFv::GAA. Both constructs provided a liver-specific promoter (LSP) to drive expression). Dose is provided as viral genome (vg) per kilogram (kg) of the mouse.

[0041] **Figure 12B** provides dot blots depicting the levels of glycogen in micrograms per milligram of tissue (heart, quadricep, diaphragm, or tricep) as a function of GAA serum levels. Squares represent AAV-LSP-Δ8GAA. Pyramids represent AAV-anti-hCD63scFv::GAA. Both constructs provided a liver-specific promoter (LSP) to drive expression).

[0042] **Figure 13** is a bar graph depicting the amount of stored glycogen in micrograms per milligram of CNS tissue as a function of delivered enzyme/vector (9 month study). The X-axis depicts each CNS tissue (spinal cord, cerebellum or hippocampus) sampled from wild-type mice comparators (each Lane 4) or KO mice (GAA^{-/-}) that were untreated (each Lane 1), or given plasmids encoding GAA (each Lane 2) or ScFv-GAA fusion (each Lane 3). It was an unexpected finding that the mice treated with fusion protein exhibited a more robust decrease in glycogen stores in CNS tissue compared to delivery of vector encoding GAA without any fused binding domain.

[0043] **Figure 14A** is a graph depicting the amount of stored glycogen per mg of spinal cord tissue (ug glycogen/mg tissue; y-axis) 3 months after AAV delivery of GAA in a specific vector construct at a specific dose (vg= viral genomes). Glycogen levels are provided for wild-type mice (Lane 1; x-axis) compared to KO mice (GAA^{-/-}) that were untreated (Lane 2; x-axis), or given plasmids encoding GAA (each Lane 3; x-axis) or varying doses of anti-CD63ScFv-GAA fusion (Lanes 4-6; x-axis). Stored glycogen levels in spinal cord decrease with increasing dose of ScFv-GAA fusion, with the 1e11vg dose providing greater benefit to the subject than an equivalent dose of GAA alone (no fusion).

[0044] **Figure 14B** is a graph depicting the amount of stored glycogen per mg of brain tissue (ug glycogen/mg tissue; y-axis) 3 months after AAV delivery of GAA in a specific vector

constructs at a specific dose (vg= viral genomes). Glycogen levels are provided for wild-type mice (Lane 1; x-axis) compared to KO mice (GAA^{-/-}) that were untreated (Lane 2; x-axis), or given plasmids encoding GAA (each Lane 3; x-axis) or varying doses of anti-CD63ScFv-GAA fusion (Lanes 4-6; x-axis). Stored glycogen levels in brain decrease with increasing dose of ScFv-GAA fusion, with the 1e11vg dose providing greater benefit to the subject than an equivalent dose of GAA alone (no fusion).

DESCRIPTION

[0045] This invention is not limited to particular embodiments, compositions, methods and experimental conditions described, as such embodiments, compositions, methods and conditions may vary. The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0046] Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, some preferred methods and materials are now described. All publications cited herein are incorporated herein by reference to describe in their entirety. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0047] “Blood Brain Barrier” refers to the semipermeable membrane barrier that separates the blood from the brain and extracellular fluid in the central nervous system. The barrier blocking the passage of, or selectively transports, certain substances to the brain and spinal cord. The blood–brain barrier is formed by brain endothelial cells.

[0048] “Therapeutically effective amount” refers to an amount or dosage of the vector delivered to a subject such that the subject achieves a consistent blood level (serum/plasma level) of the encoded therapeutic protein. Generally, concentrations of from about 1×10^9 to about 1×10^{16} genomes vector may be utilized in the method. The dosage for delivery to liver may be about 1×10^{10} to 5×10^{13} AAV genomes per 1 kg. The dosage will be adjusted to balance the therapeutic benefit of crossing the blood brain barrier to achieve the desired effect of the molecule against any side effects and such dosages may vary depending upon the recombinant vector that is employed. The levels of expression of the transgene can be monitored in the blood circulation by extraction of serum or plasma to determine the frequency of dosage of vectors that will

achieve a steady state of circulating protein. One skilled in the art can determine specific values for an effective amount by, for example, performing experiments to determine consistent blood levels of therapeutic protein over consecutive days, weeks or months following vector delivery. Suitable experiments to test for circulating therapeutic protein are known in the art, including but not limited to western blot, ELISA, LC-MS, etc. In one example, a therapeutically effective amount of scFv-GAA fusion protein in the CNS is an amount of viral vector that produces sufficient amounts of scFv-GAA fusion protein to reduce stored glycogen in CNS tissue, for example in spinal cord, cerebellum or hippocampus tissue.

CNS Disorders

[0049] Various brain disorders may benefit from the mode of delivery of therapeutic proteins described herein. CNS disorders and disorders with neurological symptoms amenable to protein therapies include, but are not limited to: Alzheimer's, brain cancer, Behcet's Disease, cerebral Lupus, Creutzfeldt-Jakob Disease, dementia, epilepsy, encephalitis, Friedreich's Ataxia, Guillain-Barre Syndrome, Gaucher Disease, headache, hydrocephalus, Huntington's disease, intracranial hypertension, leukodystrophy, migraine, myasthenia gravis, muscular dystrophy, multiple sclerosis, narcolepsy, neuropathy, Prader-Willi Syndrome, Parkinson's disease, Rett Syndrome, restless leg syndrome, sleep disorders, subarachnoid haemorrhage, stroke, traumatic brain injury, trigeminal neuralgia, transient ischaemic attack, and Von Hippel-Lindau Syndrome (angiomatosis).

[0050] Anti-CD63-fusion delivery of a therapeutic protein to the CNS may be particularly beneficial due to its ubiquitous expression, its role as a membrane protein of extracellular vesicles (EVs; *e.g.* exosomes) and association with integrins. Other cell-surface receptors with similar properties as internalizing effectors include: ITGA7, CD9, CD63, CD81, CD82, and CD151, and may be tissue or cell-type specific in order to enhance the desired location of the uptake, as discussed throughout the specification.

[0051] Anti-transferrin-fusion delivery of a therapeutic protein to the CNS has also been shown to be particularly beneficial. Trafficking and delivery of therapeutic proteins therefore will be enhanced by the use of delivery mechanisms, such as anti-receptor fusion to particular blood-brain-barrier (BBB) targets. Some BBB targets have been shown as beneficial for CNS uptake (Zuchero, et al., 6 January 2016, Neuron, 89(1): 70-82; Boado, RJ et al, Mol Pharm. 2014 Aug 4; 11(8): 2928–2934; each of which reference is incorporated herein in its entirety by

reference). Other cell-surface receptors with similar BBB uptake properties to transferrin receptors include, but are not limited to: insulin receptor, CD98, and Basigin (Bsg).

[0052] In some embodiments, the targeted delivery of the therapeutic protein to CNS tissue (e.g. brain) is employed by use of anti-transferrin receptor or anti-insulin receptor, or anti-CD98 or anti-Bsg. The therapeutic protein may also be fused to an internalizing effector antibody, as discussed herein throughout the specification. In some embodiments, the targeted delivery of the therapeutic protein to CNS and to peripheral tissues is employed by use of an anti-insulin receptor delivery domain, for example to enhance both brain uptake as well as peripheral uptake (Boado, RJ et al, 2014, supra; Yu et al., 25 May 2011, Science Transl Med. 3:84ra44, each of which reference is incorporated herein in its entirety by reference).

[0053] Exemplary anti-transferrin receptor, anti-insulin receptor, anti-CD98, and anti-Bsg antibodies and portions thereof that may be useful as part of a multidomain therapeutic protein as described herein are well-known in the art (see, e.g., US20170174778; US20150196663; US9629801; US20180002433; WO2016081643; US20180134797; WO2014189973; US20150110791; US9708406; US20170260292; WO2016081640; US20180057604; US9611323; WO2012075037; WO2018210898, US20180344869, US20180282408, US20170051071, WO2016207240, WO2015101588, US20160324984; US20180222993; WO2017055542; US20180222992; WO2017055540; Cabezon, I., et al. Mol Pharm. 2015 Nov 2;12(11):4137-45; Yu YJ, et al. Sci Transl Med (2014) 6:261ra154; Couch, et al. Sci Transl Med. 2013 May 1;5(183):183ra57, 1-12; and Yu et al., 2011, supra, for non-limiting exemplary anti-transferrin receptor antibodies; see, e.g., WO2017214456, WO2017214458, WO2017214462; WO2008017828; WO2015146132; WO2016094566; WO2013078377; WO2017026497; Hayes GM et al. Int. J. Cancer (2015) 137:710-20; and Bixby, et al. American Society Hematology 2015 Meeting, Abstract# 3809 for non-limiting exemplary anti-CD98 antibodies; see, e.g., WO2011112566; US20110223176; US20140079711; WO2010036460; US8618264; WO2005092381; WO2018165619; and WO2017186182 for non-limiting exemplary anti-BSG antibodies; see, e.g., US8974791; WO2013081706; US20160152719; US20160208006; US20170114152; Pardridge, WM et al. BioDrugs. 2018 Apr;32(2):169-176; Boado RJ et al. Mol. Pharm. (2016) 13:3241-6; Cieniewicz AM, et al. Diabetes (2017) 66:206-217; Bexwada, P. et al. J Pharmacol Exp Ther. 2016 Feb;356(2):466-73; and Bedinger, DH, et al. J Pharmacol Exp Ther. 2015 Apr;353(1):35-43 for non-limiting exemplary anti-insulin receptor antibodies; each of which reference is incorporated in its entirety by reference). A skilled artisan could readily link these well-known antibodies, or antigen binding portions

thereof (e.g., scFv derived therefrom) to a therapeutic protein as described herein to make and use a multidomain therapeutic protein as described herein.

Lysosomal Storage Diseases

[0054] “Enzyme-deficiency diseases” include non-lysosomal storage disease such as Krabbe disease (galactosylceramidase), phenylketonuria, galactosemia, maple syrup urine disease, mitochondrial disorders, Friedreich ataxia, Zellweger syndrome, adrenoleukodystrophy, Wilson disease, hemochromatosis, ornithine transcarbamylase deficiency, methylmalonic academia, propionic academia, and lysosomal storage diseases. “Lysosomal storage diseases” include any disorder resulting from a defect in lysosome function. Currently, approximately 50 lysosomal storage disorders have been identified, the most well-known of which include Tay-Sachs, Gaucher, and Niemann-Pick disease. The pathogeneses of the diseases are ascribed to the buildup of incomplete degradation products in the lysosome, usually due to loss of protein function. Lysosomal storage diseases are caused by loss-of-function or attenuating variants in the proteins whose normal function is to degrade or coordinate degradation of lysosomal contents. The proteins affiliated with lysosomal storage diseases include enzymes, receptors and other transmembrane proteins (*e.g.*, NPC1), post-translational modifying proteins (*e.g.*, sulfatase), membrane transport proteins, and non-enzymatic cofactors and other soluble proteins (*e.g.*, GM2 ganglioside activator). Thus, lysosomal storage diseases encompass more than those disorders caused by defective enzymes *per se*, and include any disorder caused by any molecular defect. Thus, as used herein, the term “enzyme” is meant to encompass those other proteins associated with lysosomal storage diseases.

[0055] The nature of the molecular lesion affects the severity of the disease in many cases, *i.e.* complete loss-of-function tends to be associated with pre-natal or neo-natal onset, and involves severe symptoms; partial loss-of-function is associated with milder (relatively) and later-onset disease. Generally, only a small percentage of activity needs to be restored to have to correct metabolic defects in deficient cells. Table 1 lists some of the more common lysosomal storage diseases and their associated loss-of-function proteins. Lysosomal storage diseases are generally described in Desnick and Schuchman, 2012.

[0056] Lysosomal storage diseases are a class of rare diseases that affect the degradation of myriad substrates in the lysosome. Those substrates include sphingolipids, mucopolysaccharides, glycoproteins, glycogen, and oligosaccharides, which can accumulate in the cells of those with disease leading to cell death. Organs affected by lysosomal storage

diseases include the central nervous system (CNS), the peripheral nervous system (PNS), lungs, liver, bone, skeletal and cardiac muscle, and the reticuloendothelial system.

[0057] Options for the treatment of lysosomal storage diseases include enzyme replacement therapy (ERT), substrate reduction therapy, pharmacological chaperone-mediated therapy, hematopoietic stem cell transplant therapy, and gene therapy. An example of substrate reduction therapy includes the use of Miglustat or Eliglustat to treat Gaucher Type 1. These drugs act by blocking synthase activity, which reduces subsequent substrate production. Hematopoietic stem cell therapy (HSCT), for example, is used to ameliorate and slow-down the negative central nervous system phenotype in patients with some forms of MPS. See R.M. Boustany, “Lysosomal storage diseases--the horizon expands,” 9(10) Nat. Rev. Neurol. 583-98, Oct. 2013; which reference is incorporated herein in its entirety by reference. Table 1 lists some lysosomal storage diseases and their associated enzymes or other proteins.

Table 1: Lysosomal Storage Diseases

Class	Disease	Involved Enzyme/Protein
Sphingolipidoses	Fabry disease	α -Galactosidase A
	Farber lipogranulomatosis	Ceramidase
	Gaucher disease type I	β -Glucosidase
	Gaucher disease types II and III	Saposin-C activator
	Niemann-Pick diseases types A and B	Sphingomyelinase
	GM1-gangliosidosis	β -Galactosidase
	GM2-gangliosidosis (Sandhoff)	β -Hexosaminidase A and B
	GM2-gangliosidosis (Tay-Sachs)	β -Hexosaminidase A
	GM2-gangliosidosis (GM2-activator deficiency)	GM2-activator protein
	GM3-gangliosidosis	GM3 synthase
	Metachromatic leukodystrophy	Arylsulfatase A
	Sphingolipid-activator deficiency	Sphingolipid activator
Mucopoly-saccharidoses	MPS I (Scheie, Hurler-Scheie, and Hurler disease)	α -Iduronidase
	MPS II (Hunter)	Iduronidase-2-sulphatase
	MPS IIIA (Sanfilippo A)	Heparan <i>N</i> -sulphatase
	MPS IIIB (Sanfilippo B)	<i>N</i> -acetyl- α -glucosaminidase
	MPS IIIC (Sanfilippo C)	Acetyl-CoA; α -glucosamide <i>N</i> -acetyltransferase
	MPS IIID (Sanfilippo D)	<i>N</i> -acetylglucosamine-6-sulphatase
	MPS IVA (Morquio syndrome A)	<i>N</i> -acetylgalactosamine-6-sulphate sulphatase
	MPS IVB (Morquio syndrome B)	β -Galactosidase
	MPS VI (Maroteaux-Lamy)	<i>N</i> -acetylgalactosamine-4-sulphatase (arylsulphatase B)

Class	Disease	Involved Enzyme/Protein
	MPS VII (Sly disease)	β -Glucuronidase
	MPS IX	Hyaluronidase
	Pompe (glycogen storage disease type II)	α -Glucosidase 2
Glycogen storage disease		
Lipid metabolism	Lysosomal acid lipase deficiency (LAL-D; Wolman disease)	Lysosomal acid lipase

[0058] Two of the most common LSDs are Pompe disease and Fabry disease. Pompe disease, which has an estimated incidence of 1 in 10,000, is caused by defective lysosomal enzyme alpha-glucosidase (GAA), which results in the deficient processing of lysosomal glycogen. Accumulation of lysosomal glycogen occurs predominantly in skeletal, cardiac, and hepatic tissues. Infantile onset Pompe causes cardiomegaly, hypotonia, hepatomegaly, and death due to cardiorespiratory failure, usually before 2 years of age. Adult onset Pompe occurs as late as the second to sixth decade and usually involves only skeletal muscle. Treatments currently available include Genzyme's MYOZYME®/LUMIZYME® (alglucosidase alfa), which is a recombinant human alpha-glucosidase produced in CHO cells and administered by intravenous infusion.

[0059] Fabry disease, which has including mild late onset cases an overall estimated incidence of 1 in 3,000, is caused by defective lysosomal enzyme alpha-galactosidase A (GLA), which results in the accumulation of globotriaosylceramide within the blood vessels and other tissues and organs. Symptoms associated with Fabry disease include pain from nerve damage and/or small vascular obstruction, renal insufficiency and eventual failure, cardiac complications such as high blood pressure and cardiomyopathy, dermatological symptoms such as formation of angiokeratomas, anhidrosis or hyperhidrosis, and ocular problems such as cornea verticillata, spoke-like cataract, and conjunctival and retinal vascular abnormalities. Treatments currently available include Genzyme's FABRAZYME® (agalsidase beta), which is a recombinant human alpha-galactosidase A produced in CHO cells and administered by intravenous infusion; Shire's REPLAGAL™ (agalsidase alfa), which is a recombinant human alpha-galactosidase A produced in human fibroblast cells and administered by intravenous infusion; and Amicus's GALAFOLD™ (migalastat or 1-deoxygalactonojirimycin) an orally administered small molecule chaperone that shifts the folding of abnormal alpha-galactosidase A to a functional conformation.

[0060] Current treatments for lysosomal storage diseases are less than optimal. For example, ERT generally must be administered at a high frequency and a high dose, such as biweekly and

up to 40 mg/kg. Also, some replaced enzymes can be immunologically cross-reactive (CRIM), stimulating production of IgG in the subject and thus hindering delivery of the enzyme to the lysosome via the mannose-6-phosphate (M6P) receptor. The IgGs might shield the M6P residues of the replacement enzyme, and the antigen-IgG-antibody complex may be taken up into cellular lysosomes via the Fc receptor, thereby shunting the replacement enzyme preferentially to macrophages.

[0061] Delivery of replacement enzymes to the appropriate affected tissues is also inefficient (see Table 2 and Desnick & Schuchman, “Enzyme replacement therapy for lysosomal diseases: lessons from 20 years of experience and remaining challenges,” 13 Annu. Rev. Genomics Hum. Genet. 307-35, 2012), which reference is incorporated herein in its entirety by reference. For example, patients undergoing long-term enzyme replacement therapy for Infantile Pompe can still suffer from hypernasal speech, residual muscle weakness, ptosis, osteopenia, hearing loss, risk for aspiration, dysphagia, cardiac arrhythmia, and difficulty swallowing. Doses of replacement enzyme oftentimes must be increased over time to 40 mg/kg weekly or biweekly.

Table 2: Inefficient tissue targeting of ERT

Disease	Subtype(s)	Easy to reach tissue	Hard to reach tissue
Gaucher disease	Type 1	Spleen, liver, bone marrow	Bone
	Types 2 and 3	Spleen, liver, bone marrow	Bone, brain
Fabry disease	Classic and late onset	Vascular endothelium	Kidney, heart
Mucopolysaccharidoses	All	Spleen, liver, bone marrow	Bone, brain, cartilage
α -Mannosidosis	---	Spleen, liver, bone marrow	Bone, brain
Niemann-Pick disease	Type B	Spleen, liver, bone marrow	Alveolar macrophages
Pompe disease	Infantile	---	Heart, smooth and skeletal muscle
	Later onset	---	Smooth muscle and respiratory skeletal muscle

[0062] Endogenous mannose-6 phosphate receptor (MPR) mediates the transport of most recombinant enzymes to the lysosome. Two complementary forms of MPR exist: cation-independent (CI-MPR), and cation dependent (CD-MPR). Knock-outs of either form have missorted lysosomal enzymes. Lysosomal hydrolases are synthesized in the endoplasmic

reticulum and move to the cis-Golgi network, where they are covalently modified by the addition of mannose-6-phosphate (M6P) groups. The formation of this marker depends on the sequential effect of two lysosomal enzymes: UDP-*N*-acetylglucosamine-1-phosphotransferase (GlcNAc-phosphotransferase) and *N*-acetylglucosamine-1-phosphodiester- α -*N*-acetylglucosaminidase (uncovering enzyme). GlcNAc-phosphotransferase catalyzes the transfer of a GlcNAc-1-phosphate residue from UDP-GlcNAc to C6 positions of selected mannoses in high-mannose type oligosaccharides of the hydrolases. Then, the uncovering enzyme removes the terminal GlcNAc, exposing the M6P recognition signal. At the trans-Golgi network, the M6P signal allows the segregation of lysosomal hydrolases from all other types of proteins through selective binding to the M6P receptors. The clathrin-coated vesicles produced bud off from the trans-Golgi network and fuse with late endosomes. At the low pH of the late endosome, the hydrolases dissociate from the M6P receptors and the empty receptors are recycled to the Golgi apparatus for further rounds of transport.

[0063] With the exception of β -glucocerebrosidase, which is delivered via the mannose receptor, recombinant lysosomal enzymes comprise M6P glycosylation and are delivered to the lysosome primarily via CI-MPR/IGF2R. Glycosylation/CI-MPR-mediated enzyme replacement delivery however does not reach all clinically relevant tissues (Table 2). Improvement to enzyme replacement therapy have centered on improving CI-MPR delivery by (i) increasing surface expression of CI-MPR using the β 2-agonist clenbuterol (Koeberl *et al.*, “Enhanced efficacy of enzyme replacement therapy in Pompe disease through mannose-6-phosphate receptor expression in skeletal muscle,” 103(2) Mol. Genet. Metab. 107-12, 2011, which reference is incorporated herein in its entirety by reference), (ii) increasing the amount of M6P residues on enzyme (Zhu *et al.*, “Conjugation of mannose-6-phosphate-containing oligosaccharides to acid alpha-glucosidase improves the clearance of glycogen in Pompe mice,” 279(48) J. Biol. Chem. 50336-41, 2004, which reference is incorporated herein in its entirety by reference), or (iii) fusing an IGF-II domain to the enzyme (Maga *et al.*, “Glycosylation-independent lysosomal targeting of acid alpha-glucosidase enhances muscle glycogen clearance in Pompe mice,” 288(3) J. Biol. Chem. 1428-38, 2013, which reference is incorporated herein in its entirety by reference).

[0064] A large number of lysosomal storage diseases are inadequately treated by enzyme replacement therapy or gene therapy mainly due to poor targeting of the replacement enzyme to the relevant tissue or organ, negative immunological reactions in the recipient host, and low serum half-life. A need exists for improved enzyme replacement therapies that enhance and

promote better tissue biodistribution and lysosomal uptake of the enzyme, especially in the brain and spinal cord without undesirable intrathecal injections. Applicants have developed an improved enzyme replacement therapy using CI-MPR independent binding protein-guided delivery of enzymes and liver expression to provide enzyme to the lysosome of target affected tissues, particularly CNS tissues.

[0065] Lysosomal storage diseases can be categorized according to the type of product that accumulates within the defective lysosome. Sphingolipidoses are a class of diseases that affect the metabolism of sphingolipids, which are lipids containing fatty acids linked to aliphatic amino alcohols (reviewed in S. Hakomori, “Glycosphingolipids in Cellular Interaction, Differentiation, and Oncogenesis,” 50 Annual Review of Biochemistry 733-764, July 1981; which reference is incorporated herein in its entirety by reference). The accumulated products of sphingolipidoses include gangliosides (*e.g.*, Tay-Sachs disease), glycolipids (*e.g.*, Fabry's disease), and glucocerebrosides (*e.g.*, Gaucher's disease).

[0066] Mucopolysaccharidoses are a group of diseases that affect the metabolism of glycosaminoglycans (GAGS or mucopolysaccharides), which are long unbranched chains of repeating disaccharides that help build bone, cartilage, tendons, corneas, skin and connective tissue (reviewed in J. Muenzer, “Early initiation of enzyme replacement therapy for the mucopolysaccharidoses,” 111(2) Mol. Genet. Metab. 63-72 (Feb. 2014); Sasisekharan *et al.*, “Glycomics approach to structure-function relationships of glycosaminoglycans,” 8(1) Ann. Rev. Biomed. Eng. 181-231 (Dec. 2014); each of which reference is incorporated herein in its entirety by reference). The accumulated products of mucopolysaccharidoses include heparan sulfate, dermatan sulfate, keratin sulfate, various forms of chondroitin sulfate, and hyaluronic acid. For example, Morquio syndrome A is due to a defect in the lysosomal enzyme galactose-6-sulfate sulfatase, which results in the lysosomal accumulation of keratin sulfate and chondroitin 6-sulfate.

[0067] Glycogen storage diseases (a.k.a., glycogenesis) result from a cell's inability to metabolize (make or break-down) glycogen. Glycogen metabolism is moderated by various enzymes or other proteins including glucose-6-phosphatase, acid alpha-glucosidase, glycogen de-branching enzyme, glycogen branching enzyme, muscle glycogen phosphorylase, liver glycogen phosphorylase, muscle phosphofructokinase, phosphorylase kinase, glucose transporter, aldolase A, beta-enolase, and glycogen synthase. An exemplar lysosomal storage/glycogen storage disease is Pompe's disease, in which defective acid alpha-glucosidase causes glycogen to accumulate in lysosomes. Symptoms include hepatomegaly, muscle

weakness, heart failure, and in the case of the infantile variant, death by age 2 (see DiMauro and Spiegel, “Progress and problems in muscle glycogenosis,” 30(2) *Acta Myol.* 96-102 (Oct. 2011); which reference is incorporated herein in its entirety by reference).

[0068] “Multidomain therapeutic protein” includes (i) a single protein that contains more than one functional domain, (ii) a protein that contains more than one polypeptide chain, and (iii) a mixture of more than one protein or more than one polypeptide. The term polypeptide is generally taken to mean a single chain of amino acids linked together via peptide bonds. The term protein encompasses the term polypeptide, but also includes more complex structures. That is, a single polypeptide is a protein, and a protein can contain one or more polypeptides associated in a higher order structure. For example, hemoglobin is a protein containing four polypeptides: two alpha globin polypeptides and two beta globin polypeptides. Myoglobin is also a protein, but it contains only a single myoglobin polypeptide.

[0069] The multidomain therapeutic protein comprises one or more polypeptide(s) and at least two domains providing two functions. One of those domains is the “enzyme domain” which provides the replacement of a defective protein activity associated with an enzyme deficiency disease. The other of those domains is the “delivery domain” which provides binding to an internalizing effector. Thus, a single polypeptide that provides an enzyme replacement activity and the ability to bind to an internalizing effector (a.k.a. internalizing effector-binding protein (delivery domain activity) is a multidomain therapeutic protein. Also, a mixture of proteins, wherein one protein provides the enzyme function, and another protein provides the internalizing effector binding activity, is a multidomain therapeutic protein. Figure 1A depicts various exemplars of multidomain therapeutic proteins. In one example (Figure 1A, panel A), the multidomain therapeutic protein contains an enzyme (represented by the hexagon) and a bispecific antibody (the IE-BP) that binds the enzyme (hashed lines) and an internalizing effector (solid lines). Here, one arm of the bispecific antibody binds non-covalently to the enzyme, and the other arm binds non-covalently to the internalizing effector, thereby enabling the internalization of the replacement enzyme into the cell or subcellular compartment. In another example (panel B), the multidomain therapeutic protein comprises a single protein containing two polypeptides, one polypeptide having enzyme function and the other having delivery domain function. Here, the enzyme is fused to an immunoglobulin Fc domain or heavy chain constant region, which associates with the Fc domain of the enzyme half-antibody to form the bifunctional multidomain therapeutic protein. The embodiment depicted in panel B is similar to that in panel A, except that the enzyme is covalently attached to one of the half-

antibodies, rather than through antigen-antibody interaction at the immunoglobulin variable domain of the half-antibody.

[0070] In other examples, the multidomain therapeutic protein consists of the enzyme covalently linked (directly or indirectly through a linker) to the delivery domain. In one embodiment, the enzyme is attached to the C-terminus of an immunoglobulin molecule (*e.g.*, the heavy chain or alternatively the light chain). In another embodiment, the enzyme is attached to the N-terminus of the immunoglobulin molecule (*e.g.*, the heavy chain or alternatively the light chain). In these exemplars, the immunoglobulin molecule is the delivery domain. In yet another embodiment, the enzyme is attached to the C-terminus of a scFv molecule that binds the internalizing effector.

[0071] In one embodiment, the multidomain therapeutic protein comprises at least two, and in some embodiments no more than two, delivery domains, each of which is directed toward a distinct epitope, either on the same antigen or on two different antigens. In one embodiment, the first delivery domain binds to a lysosomal trafficking molecule or other internalizing effector (*e.g.*, CD63) or other similar cell-surface receptor, such as ITGA7, CD9, CD63, CD81, CD82, or CD151. In another embodiment, the second delivery domain binds to a transcytosis effector to facilitate transcellular transport of the multidomain therapeutic protein. In one embodiment, the transcytosis effector is *inter alia* an LDL receptor, an IgA receptor, a transferrin receptor, or a neonatal Fc receptor (FcRn). In a specific embodiment, the transcytosis delivery domain comprises a molecule that binds to a transferrin receptor, such as *e.g.*, an anti-transferrin receptor antibody or an anti-transferrin receptor scFv molecule. Tuma and Hubbard, “Transcytosis: Crossing Cellular Barriers,” *Physiological Reviews*, 83(3): 871-935 (1 July 2003) is incorporated herein by reference for cell surface receptors that mediate transcytosis that are useful in the practice of the subject invention. In one embodiment, a second delivery domain binds to a transferrin receptor, or other similar cell-surface protein, such as an insulin receptor, CD98, or Basigin (Bsg). Each multidomain therapeutic protein comprising at least two delivery domains also comprises at least one enzyme domain, *e.g.*, each of the at least two delivery domains may or may not be independently associated an enzyme domain in a manner described herein (*e.g.*, via an antigen-antibody interaction, via a direct covalent link, via an indirect covalent link, etc.), wherein at least one of the at least two delivery domains is associated with the enzyme domain. Additionally, each of the at least two delivery domains may independently comprise an antibody, a half-body, or an scFv (*e.g.*, an scFv fused with an Fc).

[0072] “Enzyme domain” or “enzyme” denotes any protein associated with the etiology or physiological effect of an enzyme deficiency disease. An enzyme includes the actual enzyme, transport protein, receptor, or other protein that is defective and which is attributed as the molecular lesion that caused the disease. An enzyme also includes any protein that can provide a similar or sufficient biochemical or physiological activity that replaces or circumvents the molecular lesion of the disease. For example, an “isozyme” may be used as an enzyme. Examples of lysosomal storage disease-related proteins include those listed in Table 1 as “Involved Enzyme/Protein” and any known or later discovered protein or other molecule that circumvents the molecular defect of the enzyme-deficiency disease.

[0073] In some embodiments, the enzyme is a hydrolase, including esterases, glycosylases, hydrolases that act on ether bonds, peptidases, linear amidases, diphosphatases, ketone hydrolases, halogenases, phosphoamidases, sulfohydrolases, sulfinases, desulfinases, and the like. In some embodiments, the enzyme is a glycosylase, including glycosidases and *N*-glycosylases. In some embodiments, the enzyme is a glycosidase, including alpha-amylase, beta-amylase, glucan 1,4-alpha-glucosidase, cellulose, endo-1,3(4)-beta-glucanase, inulinase, endo-1,4-beta-xylanase, endo-1,4-b-xylanase, dextranase, chitinase, polygalacturonidase, lysozyme, exo-alpha-sialidase, alpha-glucosidase, beta-glucosidase, alpha-galactosidase, beta-galactosidase, alpha-mannosidase, beta-mannosidase, beta-fructofuranosidase, alpha,alpha-trehalose, beta-glucuronidase, xylan endo-1,3-beta-xylosidase, amylo-alpha-1,6-glucosidase, hyaluronoglucosaminidase, hyaluronoglucuronidase, and the like.

[0074] In the case of Pompe disease, in which the molecular defect is a defect in α -glucosidase activity, enzymes include human alpha-glucosidase, and “isozymes” such as other alpha-glucosidases, engineered recombinant alpha-glucosidase, other glucosidases, recombinant glucosidases, any protein engineered to hydrolyze a terminal non-reducing 1-4 linked alpha-glucose residue to release a single alpha-glucose molecule, any EC 3.2.1.20 enzyme, natural or recombinant low pH carbohydrate hydrolases for glycogen or starches, and glucosyl hydrolases such as sucrase isomaltase, maltase glucoamylase, glucosidase II, and neutral alpha-glucosidase.

[0075] An “internalizing effector” includes a protein, in some cases a receptor protein, that is capable of being internalized into a cell or that otherwise participates in or contributes to retrograde membrane trafficking. Internalization effector, internalizing effector, internalization receptor, and internalizing receptor are used interchangeably herein. In some instances, the internalizing effector is a protein that undergoes transcytosis; that is, the protein is internalized on one side of a cell and transported to the other side of the cell (*e.g.*, apical-to-basal). In many

embodiments, the internalizing effector protein is a cell surface-expressed protein or a soluble extracellular protein. However, the present invention also contemplates embodiments in which the internalizing effector protein is expressed within an intracellular compartment such as the endosome, endoplasmic reticulum, Golgi, lysosome, *etc.* For example, proteins involved in retrograde membrane trafficking (*e.g.*, pathways from early/recycling endosomes to the trans-Golgi network) may serve as internalizing effector proteins in various embodiments of the present invention. In any event, the binding of the delivery domain to an internalizing effector protein causes the entire multidomain therapeutic protein, and any molecules associated therewith (*e.g.*, enzyme), to also become internalized into the cell. As explained below, internalizing effector proteins include proteins that are directly internalized into a cell, as well as proteins that are indirectly internalized into a cell.

[0076] Internalizing effector proteins that are directly internalized into a cell include membrane-associated molecules with at least one extracellular domain (*e.g.*, transmembrane proteins, GPI-anchored proteins, *etc.*), which undergo cellular internalization, and are preferably processed via an intracellular degradative and/or recycling pathway. Specific non-limiting examples of internalizing effector proteins that are directly internalized into a cell include, *e.g.*, CD63, MHC-I (*e.g.*, HLA-B27), Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), MAL (Myelin And Lymphocyte protein, a.k.a. VIP17), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptors (*e.g.*, SCARA1-5, SCARB1-3, CD36), and the like.

[0077] In one embodiment, the internalizing effector is expressed in several tissue types and is useful in treatment where targeting of both the CNS and a peripheral cell type is desired. Internalizing effectors useful in trafficking to both CNS and peripheral cell types include, but are not limited to CD63, MHC-I, vacuolar-type H⁺ ATPase, IGF2R, Integrin alpha-7 (ITGA7), LRP5, LRP6, LRP8, Kremen-2, LDL- receptor, LDL-related protein 1 receptor, amyloid precursor protein-like protein- 2 (APLP2), apelin receptor (APLNR), PRLR, MAL (myelin and lymphocyte protein (MAL), diphtheria toxin receptors, HBEGF (heparin binding EGF like growth factor), glutathione receptors, glutamate receptors, leptin receptors, and folate receptors. In certain embodiments, the internalizing effector is prolactin receptor (PRLR). It was discovered that PRLR is, not only a target for certain therapeutic applications, but also an effective internalizing effector protein on the basis of its high rate of internalization and turn-

over. The potential for PRLR as an internalizing effector protein, for example, is illustrated in W02015/026907, where it is demonstrated, *inter alia*, that anti-PRLR antibodies are effectively internalized by PRLR-expressing cells in vitro.

[0078] Targeting internalizing effectors expressed by several cell types may be useful where targeting of both the CNS and a peripheral cell type is desired, e.g., in treating disease such as Fabry disease, Gaucher disease, MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVB, MPS VI, MPS VII, MPS IX, Pompe disease, Lysosomal acid lipase deficiency, Metachromatic leukodystrophy, Niemann-Pick diseases types A, B, and C2, Alpha mannosidosis, Neuraminidase deficiency, Sialidosis, Aspartylglycosaminuria, Combined saposin deficiency, Atypical Gaucher disease, Farber lipogranulomatosis, Fucosidosis, and Beta mannosidosis.

[0079] In another embodiment, the internalizing effector is expressed in a few tissue types. In one example, the internalizing effector may target bone and cartilage preferentially. Effectors useful in trafficking to CNS, and to either or both bone and cartilage include, but are not limited to Collagen X, Integrin alpha 10 (ITGA10), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor isoform C (FGFR3C), Hyaluronan and proteoglycan link protein 1 (CRTL1), Aggrecan, Collagen II, and Kremen-1. Such effectors are useful in treatment where targeting of both the CNS and the skeleton and cartilage is desired.

[0080] Targeting internalizing effectors preferentially expressed by bone and cartilage may be useful where targeting both the CNS and the skeleton and cartilage is desired, e.g., in treating disease such as MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVA, MPS IVB, MPS VI, MPS VII, MPS IX, Beta mannosidosis, Gaucher disease, atypical Gaucher disease, combined Saposin deficiency, Aspartylglycosaminuria, Farber lipogranulomatosis, Sialidosis, Neuraminidase deficiency, Mucopolysaccharidoses, and Alpha mannosidosis.

[0081] In yet another embodiment, the internalizing effector is expressed preferentially in a particular tissue or cell type, such as macrophages, monocytes, and microglia. Effectors useful in trafficking to CNS, and to macrophages include, but are not limited to scavenger receptor A1-5 (SCARA1-5), SCARB1-3, CD36, MSR1 (macrophage scavenger receptor 1), MRC1 (macrophage mannose receptor 1), VSIG4 (V-set and immunoglobulin domain-containing protein 4), CD68 (Macrosialin), and CSF1R (Macrophage colony-stimulating factor 1 receptor). Such effectors are useful in treatment where targeting of both the CNS and macrophages is desired. CNS macrophages may be referred to as microglia.

[0082] Targeting internalizing effectors expressed preferentially by macrophages (monocytes or microglia) may be useful where targeting both CNS and macrophages (or microglia) is desired, e.g., in treating disease such as lysosomal acid lipase deficiency, Gaucher disease, Atypical Gaucher disease, combined Saposin deficiency, and Farber lipogranulomatosis.

[0083] In certain embodiments, the internalizing effector is a kidney specific internalizing effector, such as CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin).

[0084] Targeting internalizing effectors preferentially expressed in the kidney may be useful where targeting both the CNS and the kidney is desired, e.g., in treating disease such as Fabry disease, Alport syndrome, polycystic kidney disease, and Thrombotic Thrombocytopenic Purpura.

[0085] In yet another embodiment, the internalizing effector is expressed preferentially in a particular tissue or cell type, such as the liver. Effectors useful in trafficking to CNS, and to liver include, but are not limited to ASGR1 and ASGR2. Such effectors are useful in treatment where targeting of both the CNS and liver is desired.

[0086] Targeting internalizing effectors expressed preferentially in the liver may be useful where targeting both CNS and liver is desired, e.g., in treating disease such as lysosomal acid lipase deficiency, Gaucher disease, MPS VI, MPS VII, MPS II, Niemann-Pick diseases types A, B, and C2, Sialidosis, Neuraminidase deficiency, atypical Gaucher disease, combined Saposin deficiency, Farber lipogranulomatosis.

[0087] In some embodiments, the internalizing effector is a muscle specific internalizer, such as BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-1S), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3).

[0088] Targeting internalizing effectors expressed preferentially by muscle may be useful where targeting both the CNS and muscle tissue is desired, e.g., in treating disease such as Pompe disease.

[0089] In some embodiments, the internalizing effector is ITGA7, ITGA10, CD9, CD63, ALPL2, MSR1, ASGR1, ASGR2, or PRLR. Antibodies to ITGA7, ITGA10, CD9, CD63, APLP2, MSR1, ASGR1, ASGR2 or PRLR are well-known in the art (see, e.g., R & D Systems “Integrin alpha 7: Products” for exemplary non-limiting anti-IGTA7 antibodies; see, e.g., US8048991, US20120034625, US8563255, US20140099716, US20160319023, WO2018138322, and US10087253 for exemplary non-limiting examples of anti-ITGA10 antibodies; see, e.g., WO201102982; WO2014185908; US20160115229; WO2017134197; US9738717; and de Goeij BE et al. Mol. Cancer Ther. (2016) Nov;15(11):2688-2697 for exemplary non-limiting anti-CD63 antibodies; see, e.g., US543153A; US5441931A; US5677146A; and US5935854A for exemplary non-limiting anti-APLP2 antibodies; see, e.g., R&D Systems product sheet for MAB27081; R&D Systems product sheet for AF2708; AbCam Product Sheet ab1515707; Yu X, et al., J. Biol. Chem., 2011;286(21):18795-806; and N Nishijima, et al., Front Immunol, 2017;8(0):379 for exemplary non-limiting examples of anti-MSR1 antibodies; see, e.g., WO2017058944 for exemplary non-limiting examples of anti-ASGR1 antibody; see, e.g., Origene Antibodies for ASGR2 available at www.origene.com/category/antibodies?q=ASGR2&sub_category=Primary+Antibodies&reactivities=Human for non-limiting exemplary anti-ASGR2 antibodies; see, e.g., US9649374; US9302015; US20130171147; US10106616; WO2015026907; US9777063; WO2011069795; US20130272968; US20140141003; WO2011069799; US20120315276; WO2012163932; US9241989; WO2012136519; WO2011069798; WO2019011719; US8883979; US20140065158; WO2015187596; US9353186; WO2018102304; US20130022606; US9688764; US20130129739; US20160002342; US20150056222; US20150056221; US20170008965; US20150093393; WO2011069797; US20160251442; US20180094066; WO2014036076; US20180185504; US20140271659; WO2011069794; WO2014143909; US20160319029; US9545451; US9023357; US20150252116; WO2011069796 Kelly MP, et al., Mol. Cancer Ther. (2017) Jul;16(7):1299-1311; Otto C. et al. Endocrinology (2015) 156:4365-73 2017-01-20; and Andreev J et al., Mol. Cancer Ther. (2017) Apr;16(4):681-693 for exemplary non-limiting anti-PRLR antibodies; each of which references is incorporated herein in its entirety by reference). A skilled artisan could readily link these well-known antibodies, or

antigen binding portions thereof (e.g., scFv derived therefrom) to a therapeutic protein as described herein to make and use a multidomain therapeutic protein as described herein.

[0090] In those embodiments in which the internalizing effector (IE) is directly internalized into a cell, the delivery domain can be, *e.g.*, an antibody or antigen-binding fragment of an antibody that specifically binds the IE, or a ligand or portion of a ligand that specifically interacts with the IE. For example, if the IE is Kremen-1 or Kremen-2, the delivery domain can comprise or consist of a Kremen ligand (*e.g.*, DKK1) or Kremen-binding portion thereof. As another example, if the IE is a receptor molecule such as ASGR1, the delivery domain can comprise or consist of a ligand specific for the receptor (*e.g.*, asialoorosomucoid [ASOR] or Beta-GalNAc) or a receptor-binding portion thereof.

[0091] Internalizing effector proteins that are indirectly internalized into a cell include proteins and polypeptides that do not internalize on their own but become internalized into a cell after binding to or otherwise associating with a second protein or polypeptide that is directly internalized into the cell. Proteins that are indirectly internalized into a cell include, *e.g.*, soluble ligands that are capable of binding to an internalizing cell surface-expressed receptor molecule. A non-limiting example of a soluble ligand that is (indirectly) internalized into a cell via its interaction with an internalizing cell surface-expressed receptor molecule is transferrin. In embodiments, wherein the IE is transferrin (or another indirectly internalized protein), the binding of the delivery domain to the IE, and the interaction of IE with transferrin receptor (or another internalizing cell-surface expressed receptor molecule), causes the entire delivery domain, and any molecules associated therewith (*e.g.*, the enzyme), to become internalized into the cell concurrent with the internalization of the IE and its binding partner.

[0092] In those embodiments in which the IE is indirectly internalized into a cell, the delivery domain can be, *e.g.*, an antibody, antigen-binding fragment of an antibody, or an scFv that specifically binds IE, or a receptor or portion of a receptor that specifically interacts with the soluble effector protein. For example, if the IE is a cytokine, the delivery domain can comprise or consist of the corresponding cytokine receptor or ligand-binding portion thereof.

[0093] An exemplar IE is CD63, which is a member of the tetraspanin superfamily of cell surface proteins that span the cell membrane four times. CD63 is expressed in virtually all tissues and is thought to be involved in forming and stabilizing signaling complexes. CD63 localizes to the cell membrane, lysosomal membrane, and late endosomal membrane. CD63 is known to associate with integrins and may be involved in epithelial-mesenchymal transitioning.

See H. Maecker *et al.*, “The tetraspanin superfamily: molecular facilitators,” 11(6) FASEB J. 428-42, May 1997; and M. Metzelaar *et al.*, “CD63 antigen. A novel lysosomal membrane glycoprotein, cloned by a screening procedure for intracellular antigens in eukaryotic cells,” 266 J. Biol. Chem. 3239-3245, 1991; each of which references is incorporated herein in its entirety by reference.

[0094] Another exemplar IE is amyloid beta (A4) precursor-like protein 2 (“APLP2”), a ubiquitously expressed member of the APP (amyloid precursor protein) family. APLP2 is a membrane-bound protein known to interact with major histocompatibility complex (MHC) class I molecules (*e.g.*, Kd). It binds Kd at the cell surface and is internalized in a clathrin-dependent manner with Kd in tow. See Tuli *et al.*, “Mechanism for amyloid precursor-like protein 2 enhancement of major histocompatibility complex class I molecule degradation,” 284 The Journal of Biological Chemistry 34296 -34307 (2009); which reference is incorporated herein in its entirety by reference.

[0095] Another IE exemplar is the prolactin receptor (PRLR). The prolactin receptor is a member of the type I cytokine receptor family and upon ligand binding and subsequent dimerization activates “the tyrosine kinases Jak2, Fyn and Tec, the phosphatase SHP-2, the guanine nucleotide exchange factor Vav, and the signaling suppressor SOCS,” (see Clevenger and Kline, “Prolactin receptor signal transduction,” 10(10) Lupus 706-18 (2001), abstract; each of which reference is incorporated herein in its entirety by reference). The prolactin receptor undergoes endocytotic recycling and can be found in lysosomal fractions. See Genty *et al.*, “Endocytosis and degradation of prolactin and its receptor in Chinese hamster ovary cells stably transfected with prolactin receptor cDNA,” 99(2) Mol. Cell Endocrinol. 221-8 (1994); and Ferland *et al.*, “The effect of chloroquine on lysosomal prolactin receptors in rat liver,” 115(5) Endocrinology 1842-9 (1984), which reference is incorporated herein in its entirety by reference.

[0096] As used herein, “immunological reaction” generally means a patient's immunological response to an outside or “non-self” protein. This immunological response includes an allergic reaction and the development of antibodies that interfere with the effectiveness of the replacement enzyme. Some patients may not produce any of the non-functioning protein, thus rendering the replacement enzyme a “foreign” protein. For example, repeated injection of recombinant GLA (rGLA) to those Fabry patients who lack GLA frequently results in an allergic reaction. In other patients, the production of antibodies against rGLA has been shown to decrease the effectiveness of the replacement enzyme in treating the disease. See for example

Tajima *et al.* (“Use of a Modified α -N-Acetylglactosaminidase (NAGA) in the Development of Enzyme Replacement Therapy for Fabry Disease,” 85(5) *Am. J. Hum. Genet.* 569-580 (2009)), which reference is incorporated herein in its entirety by reference, which discusses the use of modified NAGA as the “isozyme” to replace GLA. The modified NAGA has no immunological cross-reactivity with GLA, and “did not react to serum from a patient with Fabry disease recurrently treated with a recombinant GLA.” *Id.*, abstract.

[0097] An “immunosuppressive agent” includes drugs and/or proteins that result in general immunosuppression and may be used to prevent cross-reactive immunological materials (CRIM) against replacement enzymes, e.g., GAA or GLA respectively in a patient with Pompe or Fabry’s disease. Non-limiting examples of an immunosuppressive agent include methotrexate, mycophenolate mofetil, cyclophosphamide, rapamycin DNA alkylating agents, anti-CD20 antibody, anti-BAFF antibody, anti-CD3 antibody, anti-CD4 antibody, and any combination thereof.

[0098] Regulatory elements, e.g., promoters, that are specific to a tissue, e.g., liver, enhance expression of nucleic acid sequences, e.g., genes, under the control of such regulatory element in the tissue for which the regulatory element is specific. Non-limiting examples of a liver specific regulatory element, e.g., liver specific promoters, may be found in Chuah *et al.* (2014) *Mol. Ther.* 22:1605-13, which reference is incorporated herein in its entirety by reference.

[0099] The term “protein” means any amino acid polymer having more than about 20 amino acids covalently linked via amide bonds. Proteins contain one or more amino acid polymer chains, generally known in the art as “polypeptides”. Thus, a polypeptide may be a protein, and a protein may contain multiple polypeptides to form a single functioning biomolecule. Disulfide bridges (*i.e.*, between cysteine residues to form cystine) may be present in some proteins. These covalent links may be within a single polypeptide chain, or between two individual polypeptide chains. For example, disulfide bridges are essential to proper structure and function of insulin, immunoglobulins, protamine, and the like. For a recent review of disulfide bond formation, see Oka and Bulleid, “Forming disulfides in the endoplasmic reticulum,” 1833(11) *Biochim Biophys Acta* 2425-9 (2013), which reference is incorporated herein in its entirety by reference.

[0100] As used herein, “protein” includes biotherapeutic proteins, recombinant proteins used in research or therapy, trap proteins and other Fc-fusion proteins, chimeric proteins, antibodies, monoclonal antibodies, human antibodies, bispecific antibodies, antibody fragments, nanobodies, recombinant antibody chimeras, scFv fusion proteins, cytokines, chemokines,

peptide hormones, and the like. Proteins may be produced using recombinant cell-based production systems, such as the insect baculovirus system, yeast systems (*e.g.*, *Pichia* sp.), mammalian systems (*e.g.*, CHO cells and CHO derivatives like CHO-K1 cells). For a recent review discussing biotherapeutic proteins and their production, see Ghaderi *et al.*, “Production platforms for biotherapeutic glycoproteins. Occurrence, impact, and challenges of non-human sialylation,” 28 Biotechnol Genet Eng Rev. 147-75 (2012), which reference is incorporated herein in its entirety by reference.

[0101] The term “antibody”, as used herein, includes immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region comprises one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4 (heavy chain CDRs may be abbreviated as HCDR1, HCDR2 and HCDR3; light chain CDRs may be abbreviated as LCDR1, LCDR2 and LCDR3. The term “high affinity” antibody refers to those antibodies having a binding affinity to their target of at least 10^{-9} M, at least 10^{-10} M; at least 10^{-11} M; or at least 10^{-12} M, as measured by surface plasmon resonance, *e.g.*, BIACORE™ or solution-affinity ELISA. The term “antibody” may encompass any type of antibody, such as *e.g.* monoclonal or polyclonal. Moreover, the antibody may be of any origin, such as *e.g.* mammalian or non-mammalian. In one embodiment, the antibody may be mammalian or avian. In a further embodiment, the antibody may be of human origin and may further be a human monoclonal antibody.

[0102] The phrase “bispecific antibody” includes an antibody capable of selectively binding two or more epitopes. Bispecific antibodies generally comprise two different heavy chains, with each heavy chain specifically binding a different epitope—either on two different molecules (*e.g.*, antigens) or on the same molecule (*e.g.*, on the same antigen). If a bispecific antibody is capable of selectively binding two different epitopes (a first epitope and a second epitope), the affinity of the first heavy chain for the first epitope will generally be at least one to two or three

or four orders of magnitude lower than the affinity of the first heavy chain for the second epitope, and vice versa. The epitopes recognized by the bispecific antibody can be on the same or a different target (*e.g.*, on the same or a different protein). Bispecific antibodies can be made, for example, by combining heavy chains that recognize different epitopes of the same antigen. For example, nucleic acid sequences encoding heavy chain variable sequences that recognize different epitopes of the same antigen can be fused to nucleic acid sequences encoding different heavy chain constant regions, and such sequences can be expressed in a cell that expresses an immunoglobulin light chain. A typical bispecific antibody has two heavy chains each having three heavy chain CDRs, followed by (N-terminal to C-terminal) a CH1 domain, a hinge, a CH2 domain, and a CH3 domain, and an immunoglobulin light chain that either does not confer antigen-binding specificity but that can associate with each heavy chain, or that can associate with each heavy chain and that can bind one or more of the epitopes bound by the heavy chain antigen-binding regions, or that can associate with each heavy chain and enable binding of one or both of the heavy chains to one or both epitopes.

[0103] The phrase “heavy chain,” or “immunoglobulin heavy chain” includes an immunoglobulin heavy chain constant region sequence from any organism, and unless otherwise specified includes a heavy chain variable domain. Heavy chain variable domains include three heavy chain CDRs and four FR regions, unless otherwise specified. Fragments of heavy chains include CDRs, CDRs and FRs, and combinations thereof. A typical heavy chain has, following the variable domain (from N-terminal to C-terminal), a CH1 domain, a hinge, a CH2 domain, and a CH3 domain. A functional fragment of a heavy chain includes a fragment that is capable of specifically recognizing an antigen (*e.g.*, recognizing the antigen with a KD in the micromolar, nanomolar, or picomolar range), that is capable of expressing and secreting from a cell, and that comprises at least one CDR.

[0104] The phrase “light chain” includes an immunoglobulin light chain constant region sequence from any organism, and unless otherwise specified includes human kappa and lambda light chains. Light chain variable (VL) domains typically include three light chain CDRs and four framework (FR) regions, unless otherwise specified. Generally, a full-length light chain includes, from amino terminus to carboxyl terminus, a VL domain that includes FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4, and a light chain constant domain. Light chains that can be used with this invention include *e.g.*, those, that do not selectively bind either the first or second antigen selectively bound by the antigen-binding protein. Suitable light chains include those that can be identified by screening for the most commonly employed light chains in existing

antibody libraries (wet libraries or *in silico*), where the light chains do not substantially interfere with the affinity and/or selectivity of the antigen-binding domains of the antigen-binding proteins. Suitable light chains include those that can bind one or both epitopes that are bound by the antigen-binding regions of the antigen-binding protein.

[0105] The phrase “variable domain” includes an amino acid sequence of an immunoglobulin light or heavy chain (modified as desired) that comprises the following amino acid regions, in sequence from N-terminal to C-terminal (unless otherwise indicated): FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. A “variable domain” includes an amino acid sequence capable of folding into a canonical domain (VH or VL) having a dual beta sheet structure wherein the beta sheets are connected by a disulfide bond between a residue of a first beta sheet and a second beta sheet.

[0106] The phrase “complementarity determining region,” or the term “CDR,” includes an amino acid sequence encoded by a nucleic acid sequence of an organism's immunoglobulin genes that normally (*i.e.*, in a wildtype animal) appears between two framework regions in a variable region of a light or a heavy chain of an immunoglobulin molecule (*e.g.*, an antibody or a T cell receptor). A CDR can be encoded by, for example, a germline sequence or a rearranged or unrearranged sequence, and, for example, by a naive or a mature B cell or a T cell. In some circumstances (*e.g.*, for a CDR3), CDRs can be encoded by two or more sequences (*e.g.*, germline sequences) that are not contiguous (*e.g.*, in an unrearranged nucleic acid sequence) but are contiguous in a B cell nucleic acid sequence, *e.g.*, as the result of splicing or connecting the sequences (*e.g.*, V-D-J recombination to form a heavy chain CDR3).

[0107] The term “antibody fragment”, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen. Examples of binding fragments encompassed within the term “antibody fragment” include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.* (1989) *Nature* 241:544-546, which reference is incorporated herein in its entirety by reference), which consists of a VH domain, (vi) an isolated CDR, and (vii) an scFv, which consists of the two domains of the Fv fragment, VL and VH, joined by a synthetic linker to form a single protein chain in which the VL and VH regions pair to form monovalent molecules. Other forms of single chain antibodies, such as diabodies are also encompassed under the term “antibody” (see *e.g.*,

Holliger *et al.* (1993) PNAS USA 90:6444-6448; Poljak *et al.* (1994) Structure 2:1121-1123, which reference is incorporated herein in its entirety by reference).

[0108] The phrase “Fc-containing protein” includes antibodies, bispecific antibodies, immunoadhesins, and other binding proteins that comprise at least a functional portion of an immunoglobulin CH2 and CH3 region. A “functional portion” refers to a CH2 and CH3 region that can bind a Fc receptor (*e.g.*, an FcγR; or an FcRn, *i.e.*, a neonatal Fc receptor), and/or that can participate in the activation of complement. If the CH2 and CH3 region contains deletions, substitutions, and/or insertions or other modifications that render it unable to bind any Fc receptor and also unable to activate complement, the CH2 and CH3 region is not functional.

[0109] Fc-containing proteins can comprise modifications in immunoglobulin domains, including where the modifications affect one or more effector function of the binding protein (*e.g.*, modifications that affect FcγR binding, FcRn binding and thus half-life, and/or CDC activity). Such modifications include, but are not limited to, the following modifications and combinations thereof, with reference to EU numbering of an immunoglobulin constant region: 238, 239, 248, 249, 250, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 297, 298, 301, 303, 305, 307, 308, 309, 311, 312, 315, 318, 320, 322, 324, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 337, 338, 339, 340, 342, 344, 356, 358, 359, 360, 361, 362, 373, 375, 376, 378, 380, 382, 383, 384, 386, 388, 389, 398, 414, 416, 419, 428, 430, 433, 434, 435, 437, 438, and 439.

[0110] For example, and not by way of limitation, the binding protein is an Fc-containing protein and exhibits enhanced serum half-life (as compared with the same Fc-containing protein without the recited modification(s)) and have a modification at position 250 (*e.g.*, E or Q); 250 and 428 (*e.g.*, L or F); 252 (*e.g.*, L/Y/F/W or T), 254 (*e.g.*, S or T), and 256 (*e.g.*, S/R/Q/E/D or T); or a modification at 428 and/or 433 (*e.g.*, L/R/SI/P/Q or K) and/or 434 (*e.g.*, H/F or Y); or a modification at 250 and/or 428; or a modification at 307 or 308 (*e.g.*, 308F, V308F), and 434. In another example, the modification can comprise a 428L (*e.g.*, M428L) and 434S (*e.g.*, N434S) modification; a 428L, 259I (*e.g.*, V259I), and a 308F (*e.g.*, V308F) modification; a 433K (*e.g.*, H433K) and a 434 (*e.g.*, 434Y) modification; a 252, 254, and 256 (*e.g.*, 252Y, 254T, and 256E) modification; a 250Q and 428L modification (*e.g.*, T250Q and M428L); a 307 and/or 308 modification (*e.g.*, 308F or 308P).

[0111] The term “antigen-binding protein,” as used herein, refers to a polypeptide or protein (one or more polypeptides complexed in a functional unit) that specifically recognizes an

epitope on an antigen, such as a cell-specific antigen and/or a target antigen of the present invention. An antigen-binding protein may be multi-specific. The term “multi-specific” with reference to an antigen-binding protein means that the protein recognizes different epitopes, either on the same antigen or on different antigens. A multi-specific antigen-binding protein of the present invention can be a single multifunctional polypeptide, or it can be a multimeric complex of two or more polypeptides that are covalently or non-covalently associated with one another. The term “antigen-binding protein” includes antibodies or fragments thereof of the present invention that may be linked to or co-expressed with another functional molecule, *e.g.*, another peptide or protein. For example, an antibody or fragment thereof can be functionally linked (*e.g.*, by chemical coupling, genetic fusion, non-covalent association or otherwise) to one or more other molecular entities, such as a protein or fragment thereof to produce a bispecific or a multi-specific antigen-binding molecule with a second binding specificity.

[0112] As used herein, the term “epitope” refers to the portion of the antigen which is recognized by the multi-specific antigen-binding polypeptide. A single antigen (such as an antigenic polypeptide) may have more than one epitope. Epitopes may be defined as structural or functional. Functional epitopes are generally a subset of structural epitopes and are defined as those residues that directly contribute to the affinity of the interaction between the antigen-binding polypeptide and the antigen. Epitopes may also be conformational, that is, composed of non-linear amino acids. In certain embodiments, epitopes may include determinants that are chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl groups, or sulfonyl groups, and, in certain embodiments, may have specific three-dimensional structural characteristics, and/or specific charge characteristics. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents.

[0113] The term “domain” refers to any part of a protein or polypeptide having a particular function or structure. Preferably, domains of the present invention bind to cell-specific or target antigens. Cell-specific antigen- or target antigen-binding domains, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen.

[0114] The term “half-body” or “half-antibody”, which are used interchangeably, refers to half of an antibody, which essentially contains one heavy chain and one light chain. Antibody heavy chains can form dimers, thus the heavy chain of one half-body can associate with heavy chain associated with a different molecule (*e.g.*, another half-body) or another Fc-containing

polypeptide. Two slightly different Fc-domains may “heterodimerize” as in the formation of bispecific antibodies or other heterodimers, -trimers, -tetramers, and the like. See Vincent and Murini, “Current strategies in antibody engineering: Fc engineering and pH-dependent antigen binding, bispecific antibodies and antibody drug conjugates,” 7 Biotechnol. J. 1444-1450 (2012); and Shimamoto *et al.*, “Peptibodies: A flexible alternative format to antibodies,” 4(5) MAbs 586-91 (2012), each of which references is incorporated herein in its entirety by reference.

[0115] In one embodiment, the half-body variable domain specifically recognizes the internalizing effector and the half body Fc-domain dimerizes with an Fc-fusion protein that comprises a replacement enzyme (*e.g.*, a peptibody) Id, 586.

[0116] The term “single-chain variable fragment” or “scFv” includes a single chain fusion polypeptide containing an immunoglobulin heavy chain variable region (VH) and an immunoglobulin light chain variable region (VL). In some embodiments, the VH and VL are connect by a linker sequence of 10 to 25 amino acids. ScFv polypeptides may also include other amino acid sequences, such as CL or CH1 regions. ScFv molecules can be manufactured by phage display or made by directly subcloning the heavy and light chains from a hybridoma or B-cell. Ahmad *et al.*, Clinical and Developmental Immunology, volume 2012, article ID 98025 is incorporated herein by reference for methods of making scFv fragments by phage display and antibody domain cloning.

[0117] “Alpha-glucosidase” (or “ α -glucosidase”), “ α -glucosidase activity”, “GAA”, and “GAA activity” are used interchangeably and refer to any protein that facilitates the hydrolysis of 1,4-alpha bonds of glycogen and starch into glucose. GAA is also known *inter alia* as EC 3.2.1.20, maltase, glucoinvertase, glucosidosucrase, maltase-glucoamylase, alpha-glucopyranosidase, glucosidoinvertase, alpha-D-glucosidase, alpha-glucoside hydrolase, alpha-1,4-glucosidase, and alpha-D-glucoside glucohydrolase. GAA can be found in the lysosome and at the brush border of the small intestine. Patients who suffer from Pompe disease lack functioning lysosomal α -glucosidase. See S. Chiba, “Molecular mechanism in alpha-glucosidase and glucoamylase,” 61(8) Biosci. Biotechnol. Biochem. 1233-9 (1997); and Hesselink *et al.*, “Lysosomal dysfunction in muscle with special reference to glycogen storage disease type II,” 1637(2) Biochim. Biophys. Acta. 164-70 (2003), which reference is incorporated herein in its entirety by reference.

[0118] “Alpha-galactosidase A”(or “ α -galactosidase A”), “ α -galactosidase A activity”, “ α -galactosidase”, “ α -galactosidase activity”, “GLA”, and “GLA activity” are used interchangeably and refer to any protein that facilitates the hydrolysis of terminal α -galactosyl moieties from glycolipids and glycoproteins, and also hydrolyses α -D-fucosides. GLA is also known *inter alia* as EC 3.2.1.22, melibiase, α -D-galactosidase, α -galactosidase A, α -galactoside galactohydrolase, α -D-galactoside galactohydrolase. GLA is a lysosomal enzyme encoded by the X-linked GLA gene. Defects in GLA can lead to Fabry Disease, in which the glycolipid known as globotriaosylceramide (a.k.a. Gb3, GL-3, or ceramide trihexoside) accumulates within blood vessels (*i.e.*, prominent vasculopathy), resulting in pain and impairment in the function of kidney, heart, skin, and/or cerebrovascular tissues. and other tissues, and organs. See for example Prabakaran *et al.* “Mannose 6-phosphate receptor and sortilin mediated endocytosis of α -galactosidase A in kidney endothelial cells,” 7(6) PLoS One e39975 pp. 1-9 (2012), which reference is incorporated herein in its entirety by reference.

[0119] In one aspect, the invention provides a method of treating a patient (or subject) suffering from a lysosomal storage disease by administering to the patient a “multidomain therapeutic protein”. The multidomain therapeutic protein enters the cells of the patient and delivers to the lysosomes an enzyme or enzymatic activity that (*i.e.*, “replacement enzyme”) that replaces the enzyme (*i.e.*, “endogenous enzyme”) or enzymatic activity that is associated with the LSD. In one embodiment, the multidomain therapeutic protein is delivered to the patient via a gene therapy vector that contains a polynucleotide that encodes the multidomain therapeutic protein.

[0120] LSDs include sphingolipidoses, a mucopolysaccharidoses, and glycogen storage diseases. In some embodiments, the LSD is any one or more of Fabry disease, Gaucher disease type I, Gaucher disease type II, Gaucher disease type III, Niemann-Pick disease type A, Niemann-Pick disease type BGM1-gangliosidosis, Sandhoff disease, Tay-Sachs disease, GM2-activator deficiency, GM3-gangliosidosis, metachromatic leukodystrophy, sphingolipid-activator deficiency, Scheie disease, Hurler-Scheie disease, Hurler disease, Hunter disease, Sanfilippo A, Sanfilippo B, Sanfilippo C, Sanfilippo D, Morquio syndrome A, Morquio syndrome B, Maroteaux-Lamy disease, Sly disease, MPS IX, and Pompe disease. In a specific embodiment, the LSD is Fabry disease. In another embodiment, the LSD is Pompe disease.

[0121] In some embodiments, the multidomain therapeutic protein comprises (a) the replacement enzyme, and (b) a molecular entity that binds an internalizing effector (delivery domain). In some cases, the replacement enzyme is any one or more of α -galactosidase, β -galactosidase, α -glucosidase, β -glucosidase, saposin-C activator, ceramidase, sphingomyelinase,

β -hexosaminidase, GM2 activator, GM3 synthase, arylsulfatase, sphingolipid activator, α -iduronidase, iduronidase-2-sulfatase, heparin *N*-sulfatase, *N*-acetyl- α -glucosaminidase, α -glucosamide *N*-acetyltransferase, *N*-acetylglucosamine-6-sulfatase, *N*-acetylgalactosamine-6-sulfate sulfatase, *N*-acetylgalactosamine-4-sulfatase, β -glucuronidase, and hyaluronidase.

[0122] In some cases, the patient may not make sufficient protein such that a replacement enzyme is recognized by the patient as “non-self” and an immunological reaction ensues after administering a replacement enzyme. This is not desirable. Therefore, in some embodiments, the replacement enzyme is designed or produced in such a way as to avoid inducing an immunological reaction in the subject. One such solution is to use an “isozyme” as a replacement enzyme. An isozyme is sufficiently close to a “self” protein of the patient but has the replacement enzyme activity sufficient to ameliorate the symptoms of the LSD.

[0123] In one particular embodiment, in which the LSD is Pompe disease and the endogenous enzyme is α -glucosidase (GAA), the isozyme can be any one of acid α -glucosidase, sucrase-isomaltase (SI), maltase-glucoamylase (MGAM), glucosidase II (GANAB), and neutral α -glucosidase (C GNAC). In another particular embodiment, in which the LSD is Fabry disease and the endogenous enzyme is α -galactosidase A (GLA), the isozyme can be an α -*N*-acetylgalactosaminidase engineered to have GLA activity.

[0124] Provided herein are methods, other than to use an isozyme, to reduce cross-reactive immunological materials (CRIM) against the replacement enzyme. As demonstrated in Figures 5 and 6, administration of a multidomain therapeutic protein (e.g., via a gene therapy vector) comprising an internalizing effector binding domain and the enzyme domain reduces the level of CRIM against the replacement enzyme comprised to administration of a control therapeutic protein (lacking the internalizing effector domain and comprising an enzyme domain). As such, in one embodiment or reducing CRIM against an enzyme in a patient with a deficiency in the enzyme comprises administering to the patient a multidomain therapeutic protein (or nucleic acid encoding same, e.g., a gene therapy vector containing a gene encoding the multidomain therapeutic protein, wherein the multidomain therapeutic protein comprises a delivery domain (e.g., internalizing effector binding protein) and an enzyme domain.

[0125] The multidomain therapeutic protein has an internalizing effector binding protein component that enables the uptake of the replacement enzyme into the cell. Thus, in some embodiments, the internalizing effector can be CD63, MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1,

ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), PRLR (prolactin receptor), MAL (Myelin And Lymphocyte protein, a.k.a. VIP17), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptor, scavenger receptor, SCARA1-5, SCARB1-3, and CD36. In certain embodiments, the internalizing effector is a kidney specific internalizer, such as CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin). In other certain embodiments, the internalizing effector is a muscle specific internalizer, such as BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-1S), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3). In some specific embodiments, the internalizing effector is ITGA7, CD9, CD63, APLP2, ASGR1, ASGR2, or PRLR.

[0126] In some embodiments, the internalizing effector-binding protein comprises an antigen-binding protein, which includes for example a receptor-fusion molecule, a trap molecule, a receptor-Fc fusion molecule, an antibody, an Fab fragment, an F(ab')₂ fragment, an Fd fragment, an Fv fragment, a single-chain Fv (scFv) molecule, a dAb fragment, an isolated complementarity determining region (CDR), a CDR3 peptide, a constrained FR3-CDR3-FR4 peptide, a domain-specific antibody, a single domain antibody, a domain-deleted antibody, a chimeric antibody, a CDR-grafted antibody, a diabody, a triabody, a tetrabody, a minibody, a nanobody, a monovalent nanobody, a bivalent nanobody, a small modular immunopharmaceutical (SMIP), a camelid antibody (VHH heavy chain homodimeric antibody), and a shark variable IgNAR domain.

[0127] In one embodiment, the molecular entity that binds the internalizing effector is an antibody, an antibody fragment, or other antigen-binding protein. For example, the molecular entity can be a bispecific antibody, in which one arm binds the internalizing effector (*e.g.*, ITGA7, CD9, CD63, PRLR, APLP2, ASGR1, ASGR2), and the other arm binds the replacement enzyme. Here, the multidomain therapeutic protein comprises the bispecific antibody and the replacement enzyme (Fig. 1A). In a specific embodiment, the disease treated is

Fabry disease, and the multidomain therapeutic protein comprises GLA and a bispecific antibody that binds GLA and CD63. In a specific embodiment, the disease treated is Fabry disease, and the multidomain therapeutic protein comprises GLA and a bispecific antibody that binds GLA and ITGA7. In another specific embodiment, the disease treated is Pompe disease, and the multidomain therapeutic protein comprises GAA and a bispecific antibody that binds GAA and CD63. In another specific embodiment, the disease treated is Pompe disease, and the multidomain therapeutic protein comprises GAA and a bispecific antibody that binds GAA and ITGA7.

[0128] In another embodiment, the molecular entity that binds the internalizing effector comprises a half-antibody, and the replacement enzyme contains an Fc domain (enzyme-Fc fusion polypeptide). In one embodiment, the Fc domain of the enzyme-Fc fusion polypeptide associates with the Fc domain of the internalizing effector-specific half-body to form the multidomain therapeutic protein (Fig. 1B).

[0129] In other embodiments, the replacement enzyme is covalently linked to internalizing effector-binding protein. The enzyme-Fc fusion:half-body embodiment described in the previous paragraph (see also Fig. 1B) falls into this class, since the Fc dimer can be secured via one or more disulfide bridges. The covalent linkage between the enzyme activity domain or polypeptide and the internalization-binding domain or polypeptide may be any type of covalent bond, *i.e.*, any bond that involved sharing of electrons. In some cases, the covalent bond is a peptide bond between two amino acids, such that the replacement enzyme and the internalizing effector-binding protein in whole or in part form a continuous polypeptide chain, as in a fusion protein. In some cases, the replacement enzyme portion and the internalizing effector-binding protein are directly linked. In other cases, a linker is used to tether the two portions. See Chen *et al.*, "Fusion protein linkers: property, design and functionality," 65(10) Adv Drug Deliv Rev. 1357-69 (2013).

[0130] The term "linker" or "spacer" refers to a short (e.g., 2 to 25 amino acids) polypeptide that typically allow for proper folding of one or more linked components of the fusion protein, e.g., a VH linked to a VL of an scFv, a therapeutic protein (e.g., replacement enzyme) linked to a delivery domain (e.g., an anti-internalizing effector antibody) of a multidomain therapeutic protein as described herein. The linker provides a flexible junction region of the component of the fusion protein, allowing the two ends of the molecule to move independently, and may play an important role in retaining each of the two moieties' appropriate functions. Therefore, the junction region acts in some cases as both a linker, which combines the two parts together, and

as a spacer, which allows each of the two parts to form its own biological structure and not interfere with the other part. Furthermore, the junction region should create an epitope that will not be recognized by the subject's immune system as foreign, in other words, will not be considered immunogenic. Linker selection may also have an effect on binding activity of the fusion molecule. (See Huston, et al, 1988, *PNAS*, 85:16:5879-83; Robinson & Bates, 1998, *PNAS* 95(11):5929-34; Arai, et al. 2001, *PEDS*, 14(8):529-32; and Chen, X. et al., 2013, *Advanced Drug Delivery Reviews* 65:1357-1369.) In one embodiment, the delivery domain is connected to the therapeutic polypeptide, or fragment thereof, via one or more peptide linkers. In another embodiment, the variable regions of an scFv antibody are connected to each other, or a fragment thereof, via one or more peptide linkers.

[0131] The length of the linker chain may be 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 15 or more amino acid residues, but typically is between 5 and 25 residues. Examples of linkers include polyGlycine linkers, such as Gly-Gly, Gly-Gly-Gly (3Gly), 4Gly, 5Gly, 6Gly, 7Gly, 8Gly or 9Gly. Examples of linkers also include Gly-Ser peptide linkers such as Ser-Gly, Gly-Ser, Gly-Gly-Ser, Ser-Gly-Gly, Gly-Gly-Gly-Ser, Ser-Gly-Gly-Gly, Gly-Gly-Gly-Gly-Ser, Ser-Gly-Gly-Gly-Gly, Gly-Gly-Gly-Gly-Gly-Ser, Ser-Gly-Gly-Gly-Gly-Gly, Gly-Gly-Gly-Gly-Gly-Gly-Ser, Ser-Gly-Gly-Gly-Gly-Gly-Gly, (Gly-Gly-Gly-Gly-Ser) n , and (Ser-Gly-Gly-Gly-Gly) n , wherein $n = 1$ to 10. (Gly-Gly-Gly-Gly-Ser) n and (Ser-Gly-Gly-Gly-Gly) n are also known as (G4S) n and (S4G) n , respectively.

[0132] In some embodiment, the therapeutic protein, e.g., replacement enzyme, is covalently linked to the C- terminus of the heavy chain of an anti-internalizing effector antibody (see Fig. 1C) or to the C-terminus of the light chain (Fig. 1E). In some embodiments, the replacement enzyme is covalently linked to the N-terminus of the heavy chain of an anti-internalizing effector antibody (see Fig. 1D) or to the N-terminus of the light chain (Fig. 1F). In some embodiments, the enzyme is linked to the C-terminus of an anti-internalizing effector scFv domain (Fig. 1G).

[0133] In some cases, especially where the therapeutic protein, e.g., replacement enzyme, is not normally proteolytically processed in the lysosome, a cleavable linker is added to those embodiments of the multidomain therapeutic protein that comprise an antibody-enzyme fusion. In some embodiments, a cathepsin cleavable linker is inserted between the antibody and the replacement enzyme to facilitate removal of the antibody in the lysosome in order to a) possibly help preserve enzymatic activity by removing the sterically large antibody and b) possibly increase lysosomal half-life of the enzyme.

[0134] In one particular embodiment, the multidomain therapeutic protein is delivered to the patient or cell in a gene therapy vector that contains a polynucleotide that encodes the multidomain therapeutic protein. In one embodiment, the multidomain therapeutic protein comprises a delivery domain and an enzyme domain. In a specific embodiment, the delivery domain binds to an internalizing effector, such as CD63, MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), MAL (myelin and lymphocyte protein (MAL)), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptor A1-5 (SCARA1-5), SCARB1-3, or CD36. In one embodiment, the delivery domain is a single-chain variable fragment (scFv) that binds to CD63 (*i.e.*, anti-CD63 scFv). In another embodiment, the delivery domain is a single-chain variable fragment (scFv) that binds to ITGA7 (*i.e.*, anti-ITGA7 scFv).

[0135] In one particular embodiment, the enzyme domain of the multidomain therapeutic protein comprises a hydrolase. In a specific embodiment, the enzyme domain comprises a hydrolase that is a glycosylase. In a more specific embodiment, the enzyme domain comprises a glycosylase that is a glycosidase. In a more specific embodiment, the enzyme domain is a glycosidase that is alpha-glucosidase.

[0136] Generally, disclosed herein are compositions comprising and use of polynucleotides, e.g., (m)RNA, DNA, and modified forms thereof, that encode a multidomain therapeutic protein comprising an internalizing effector domain and an enzyme domain in the treatment of lysosomal storage diseases, e.g., for the reduction of glycogen and/or the enhancement of immune tolerance for GAA in a patient with Pompe disease.

[0137] The term "polynucleotide" includes a polymer of nucleotides (e.g., RNA or DNA) that encodes at least one polypeptide, including fusion polypeptides, e.g., a multidomain therapeutic polypeptide comprising an internalizing effector domain and an enzyme domain. Polynucleotide as used herein encompasses polymers comprising both modified and unmodified nucleotides. A polynucleotide may contain one or more coding and non-coding regions. A polynucleotide can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, etc. Where appropriate, e.g., in the case of chemically synthesized molecules, A polynucleotide can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, backbone modifications, etc. A polynucleotide sequence is presented in the 5' to 3' direction unless otherwise indicated. In some

embodiments, a polynucleotide is or comprises natural nucleosides (e.g., adenosine, guanosine, cytidine, uridine); nucleoside analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (e.g., methylated bases); intercalated bases; modified sugars (e.g., 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (e.g., phosphorothioates and 5'-N-phosphoramidite linkages).

[0138] In some embodiments, a polynucleotide comprises one or more nonstandard nucleotide residues. The nonstandard nucleotide residues may include, e.g., 5-methyl-cytidine ("5mC"), pseudouridine ("ψU"), and/or 2-thio-uridine ("2sU"). See, e.g., U.S. Pat. No. 8,278,036 or WO2011012316, each of which is incorporated in its entirety by reference for a discussion of such residues and their incorporation into a polynucleotide. The presence of nonstandard nucleotide residues may render a polynucleotide more stable and/or less immunogenic than a control a polynucleotide with the same sequence but containing only standard residues. In further embodiments, a polynucleotide may comprise one or more nonstandard nucleotide residues chosen from isocytosine, pseudoisocytosine, 5-bromouracil, 5-propynyluracil, 6-aminopurine, 2-aminopurine, inosine, diaminopurine and 2-chloro-6-aminopurine cytosine, as well as combinations of these modifications and other nucleobase modifications. Certain embodiments may further include additional modifications to the furanose ring or nucleobase. Additional modifications may include, for example, sugar modifications or substitutions (e.g., one or more of a 2'-O-alkyl modification, a locked nucleic acid (*LNA*)). In some embodiments, the polynucleotide may be complexed or hybridized with additional polynucleotides and/or peptide polynucleotides (*PNA*). In embodiments where the sugar modification is a 2'-O-alkyl modification, such modification may include, but are not limited to a 2'-deoxy-2'-fluoro modification, a 2'-O-methyl modification, a 2'-O-methoxyethyl modification and a 2'-deoxy modification. In certain embodiments, any of these modifications may be present in 0-100% of the nucleotides--for example, more than 0%, 1%, 10%, 25%, 50%, 75%, 85%, 90%, 95%, or 100% of the constituent nucleotides individually or in combination. In some embodiments, a polynucleotide comprises messenger RNA (mRNA) molecules, which may or may not be modified, e.g., which may or may not comprise a modified nucleotide, by well-known methods

to increase their stability and/or decrease their immunogenicity. In some embodiments, a polynucleotide comprises DNA molecules, which may or may not be modified, e.g., which may or may not comprise a modified nucleotide, by well-known methods to increase their stability and/or decrease their immunogenicity.

[0139] In some embodiments, the polynucleotide also includes a “locus-targeting nucleic acid sequence”. The locus targeting sequence enables the integration of the multidomain therapeutic protein-encoding polynucleotide into the genome of the recipient host cell. In some embodiments, the locus targeting sequence include flanking homology arms to enable homologous recombination. In some embodiments, the locus targeting sequence includes guide RNA sequences and a type II Cas enzyme to facilitate integration (*i.e.*, the CRISPR-Cas9 method). In some embodiments, the locus targeting sequence includes guide zinc-finger nuclease (ZFN) recognition sequences to facilitate integration. In some embodiments, the locus targeting sequence includes transcription activator-like effector nuclease (TALEN) recognition sequences to facilitate integration. In still other embodiments, the locus targeting sequence includes a single residue-to-nucleotide code used by BuD-derived nucleases to facilitate integration.

[0140] In some embodiments, the genomic locus into which the multidomain therapeutic protein-encoding polynucleotide is integrated is a “safe harbor locus”. In one embodiment, a “safe harbor locus” enables high expression of the multidomain therapeutic protein, while not interfering with the expression of essential genes or promoting the expression of oncogenes or other deleterious genes. In one embodiment, the genomic locus is at or proximal to the liver-expressed albumin (Alb) locus, a EESYR locus, a SARS locus, position 188,083,272 of human chromosome 1 or its non-human mammalian orthologue, position 3,046,320 of human chromosome 10 or its non-human mammalian orthologue, position 67,328,980 of human chromosome 17 or its non-human mammalian orthologue, an adeno-associated virus site 1 (AAVS1) on chromosome, a naturally occurring site of integration of AAV virus on human chromosome 19 or its non-human mammalian orthologue, a chemokine receptor 5 (CCR5) gene, a chemokine receptor gene encoding an HIV-1 coreceptor, or a mouse Rosa26 locus or its non-murine mammalian orthologue. In one embodiment, the genomic locus is an adeno-associated virus site. In one embodiment, the genomic locus for integration is selected according to the method of Papapetrou and Schambach, J. Molecular Therapy, vol. 24 (4):678-684, April 2016, which is herein incorporated by reference for the step-wise selection of a safe harbor genomic locus for gene therapy vector integration; see also Barzel et al. Nature, vol.

517:360-364, incorporated herein by reference in its entirety, for the promoterless gene targeting into the liver-expressed albumin (*Alb*) locus.

[0141] In some embodiments, the polynucleotide, e.g., DNA, also contains a promoter operably linked to the multidomain therapeutic protein-encoding nucleic acid sequence. In a specific embodiment, the promoter is a tissue-specific promoter that drives gene expression in a particular tissue. In one embodiment, the tissue specific promoter is a liver-specific enhancer/promoter derived from serpinA1 (e.g., SEQ ID NO:9) and/or is a TTR promoter (SEQ ID NO:8). In other embodiments, the promoter is a CMV promoter. In other embodiments, the promoter is a ubiquitin C promoter

[0142] In one embodiment, the multidomain therapeutic protein-encoding “gene therapy vector” is any vector capable of delivering the polynucleotide encoding the multidomain therapeutic protein to a host, e.g., a patient. In some embodiments the gene therapy vector targets a specific host cell or organ, e.g., for local delivery, e.g., tissue specific delivery. Typically, local delivery requires a protein (e.g., a multidomain therapeutic protein) encoded by mRNAs be translated and expressed mainly in and/or by an organ, e.g., a liver, whereby thereby forming a depot, e.g., a liver depot for production (and secretion) of the protein. In some embodiments, a gene therapy vector delivers a multidomain therapeutic protein polynucleotide to the liver in a patient to form a liver depot. See, e.g., DeRosa et al. *Gene Therapy*, vol. 10:699-707, incorporated herein by reference in its entirety. In some embodiments, a gene therapy vector delivers a polynucleotide encoding a multidomain therapeutic protein to muscle tissue in a patient. In some embodiments, a gene therapy vector delivers a polynucleotide encoding a multidomain therapeutic protein to the brain of a patient.

[0143] Any now-known or future-developed gene therapy delivery vector, natural or engineered, can be used in the practice of this invention. In some embodiments, the gene therapy vector is a viral vector, e.g., comprises a virus, viral capsid, viral genome etc. In some embodiments, the gene therapy vector is a naked polynucleotide, e.g., an episome. In some embodiments, the gene therapy vector comprises a polynucleotide complex. Exemplary non-limiting polynucleotide complexes for use as a gene therapy vector include lipoplexes, polymersomes, polyplexes, dendrimers, inorganic nanoparticles (e.g., polynucleotide coated gold, silica, iron oxide, calcium phosphate, etc.). In some embodiments, a gene therapy vector as described herein comprises a combination of a viral vector, naked polynucleotides, and polynucleotide complexes.

[0144] In one embodiment, the gene therapy vector is a virus, including a retrovirus, adenovirus, herpes simplex virus, pox virus, vaccinia virus, lentivirus, or an adeno-associated virus. In one embodiment, the gene therapy vector is an adeno-associated virus (AAV), including serotypes AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, and AAV11, or engineered or naturally selected variants thereof.

[0145] In one embodiment, the polynucleotide also contains adeno-associated virus (AAV) nucleic acid sequence. In one embodiment, the gene therapy vector is a chimeric adeno-associated virus containing genetic elements from two or more serotypes. For example, an AAV vector with rep genes from AAV1 and cap genes from AAV2 (designated as AAV1/2 or AAV RC1/2) may be used as a gene therapy vector to deliver the multidomain therapeutic protein polynucleotide to a cell or a cell of a patient in need. In one embodiment, the gene therapy vector is an AAV1/2, AAV1/3, AAV1/4, AAV1/5, AAV1/6, AAV1/7, AAV1/8, AAV1/9, AAV1/10, AAV1/11, AAV2/1, AAV2/3, AAV2/4, AAV2/5, AAV2/6, AAV2/7, AAV2/8, AAV2/9, AAV2/10, AAV2/11, AAV3/1, AAV3/2, AAV3/4, AAV3/5, AAV3/6, AAV3/7, AAV3/8, AAV3/9, AAV3/10, AAV3/10, AAV4/1, AAV4/2, AAV4/3, AAV4/5, AAV4/6, AAV4/7, AAV4/8, AAV4/9, AAV4/10, AAV4/11, AAV5/1, AAV5/2, AAV5/3, AAV5/4, AAV5/6, AAV5/7, AAV5/8, AAV5/9, AAV5/10, AAV5/11, AAV6/1, AAV6/2, AAV6/3, AAV6/4, AAV6/5, AAV6/7, AAV6/8, AAV6/9, AAV6/10, AAV6/10, AAV7/1, AAV7/2, AAV7/3, AAV7/4, AAV7/5, AAV7/6, AAV7/8, AAV7/9, AAV7/10, AAV7/11, AAV8/1, AAV8/2, AAV8/3, AAV8/4, AAV8/5, AAV8/6, AAV8/7, AAV8/9, AAV8/10, AAV8/11, AAV9/1, AAV9/2, AAV9/3, AAV9/4, AAV9/5, AAV9/6, AAV9/7, AAV9/8, AAV9/10, AAV9/11, AAV10/1, AAV10/2, AAV10/3, AAV10/4, AAV10/5, AAV10/6, AAV10/7, AAV10/8, AAV10/9, AAV10/11, AAV11/1, AAV11/2, AAV11/3, AAV11/4, AAV11/5, AAV11/6, AAV11/7, AAV11/8, AAV11/9, AAV11/10, chimeric viral vector or derivatives thereof. Gao *et al.*, “Novel adeno-associated viruses from rhesus monkeys as vectors for human gene therapy,” PNAS 99(18): 11854–11859, Sep. 3, 2002, is incorporated herein by reference for AAV vectors and chimeric viral vectors useful as gene therapy vectors, and their construction and use.

[0146] In a more specific embodiment, the gene therapy vector is a chimeric AAV vector with a serotype 2 rep gene sequence and a serotype 8 cap sequence (“AAV2/8” or “AAV RC2/8”).

[0147] In some embodiments, the gene therapy vector is a viral vector that has been pseudotyped (e.g., engineered) to target a specific cell, e.g., a hepatocyte. Many of the advances in targeted gene therapy using viral vectors may be summarized as non-recombinatorial (non-

genetic) or recombinatorial (genetic) modification of the viral vector, which result in the pseudotyping, expanding, and/or retargeting of the natural tropism of the viral vector. (Reviewed in Nicklin and Baker (2002) *Curr. Gene Ther.* 2:273-93; Verheij and Rottier (2012) *Advances Virol* 2012:1-15; each of which references is incorporated herein in its entirety by reference). Non-genetic approaches typically utilize an adaptor, which recognizes both a wildtype (non-modified) virus surface protein and a target cell. Soluble pseudo-receptors (for the wildtype virus), polymers such as polyethylene glycol, and antibodies or portions thereof, have been used as the virus binding domain of the adaptors, while natural peptide or vitamin ligands, and antibodies and portions thereof have been used for the cell binding domain of the adaptors described above. For example, retargeting of the viral vector to a target cell may be accomplished upon binding of the vector:adaptor complex to a protein expressed on the surface of the target cell, e.g., a cell surface protein. Such approach has been used for AAV (Bartlett et al. (1999) *Nat. Biotechnol.* 17: 2777-2785), adenoviruses (Hemminki et al. (2001) *Cancer Res.* 61: 6377-81; van Beusechem et al. (2003) *Gene Therapy* 10:1982-1991; Einfeld, et al. (2001) *J. Virol.* 75:11284-91; Glasgow et al. (2009) *PLOS One* 4:e8355), herpesviruses (Nakano et al. (2005) *Mol. Ther.* 11:617-24), and paramyxoviruses (Bian et al. (2005) *Cancer Gene Ther.* 12:295-303; Bian et al. (2005) *Int. J. Oncol.* 29:1359-69), Coronaviruses (Haijema et al. (2003) *J. Virol.* 77:4528-4538; Wurdinger et al. (2005) *Gene Therapy* 12:1394-1404; each of which references is incorporated herein in its entirety by reference).

[0148] A more popular approach has been the recombinatorial genetic modification of viral capsid proteins, and thus, the surface of the viral capsid. In indirect recombinatorial approaches, a viral capsid is modified with a heterologous “scaffold”, which then links to an adaptor. The adaptor binds to the scaffold and the target cell. (Arnold et al. (2006) *Mol. Ther.* 5:125-132; Ponnazhagen et al. (2002) *J. Virol.* 76:12900-907; see also WO 97/05266 each of which references is incorporated herein in its entirety by reference) Scaffolds such as (1) Fc binding molecules (e.g., Fc receptors, Protein A, etc.), which bind to the Fc of antibody adaptors, (2) (strept)avidin, which binds to biotinylated adaptors, (3) biotin, which binds to adaptors fused with (strept)avidin, and (4) protein:protein binding pairs that form isometric peptide bonds such as SpyCatcher, which binds a SpyTagged adaptor, have been incorporated into Ad (Pereboeva et al. (2007) *Gene Therapy* 14: 627-637; Park et al. (2008) *Biochemical and Biophysical Research Communications* 366: 769-774; Henning et al. (2002) *Human Gene Therapy* 13:1427-1439; Banerjee et al. (2011) *Bioorganic and Medicinal Chemistry Letters* 21:4985-4988), AAV (Gigout et al. (2005) *Molecular Therapy* 11:856-865; Stachler et al.

(2008) *Molecular Therapy* 16:1467–1473), and togavirus (Quetglas et al. (2010) *Virus Research* 153:179–196; Ohno et al. (1997) *Nature Biotechnology* 15:763–767; Klimstra et al. (2005) *Virology* 338:9–21; each of which references is incorporated herein in its entirety by reference).

[0149] In a direct recombinatorial targeting approach, a targeting ligand is directly inserted into, or coupled to, a viral capsid, i.e., protein viral capsids are modified to express a heterologous ligand. The ligand then redirects, e.g., binds, a receptor or marker preferentially or exclusively expressed on a target cell. (Stachler et al. (2006) *Gene Ther.* 13:926–931; White et al. (2004) *Circulation* 109:513–519; each of which references is incorporated herein in its entirety by reference). Direct recombinatorial approaches have been used in AAV (Park et al., (2007) *Frontiers in Bioscience* 13:2653–59; Girod et al. (1999) *Nature Medicine* 5:1052–56; Grifman et al. (2001) *Molecular Therapy* 3:964–75; Shi et al. (2001) *Human Gene Therapy* 12:1697–1711; Shi and Bartlett (2003) *Molecular Therapy* 7:515–525, each of which references is incorporated herein in its entirety by reference), retrovirus (Dalba et al. *Current Gene Therapy* 5:655–667; Tai and Kasahara (2008) *Frontiers in Bioscience* 13:3083–3095; Russell and Cosset (1999) *Journal of Gene Medicine* 1:300–311; Erlwein et al. (2002) *Virology* 302:333–341; Chadwick et al. (1999) *Journal of Molecular Biology* 285:485–494; Pizzato et al. (2001) *Gene Therapy* 8:1088–1096), poxvirus (Guse et al. (2011) *Expert Opinion on Biological Therapy* 11:595–608; Galmiche et al. (1997) *Journal of General Virology* 78:3019–3027; Paul et al. (2007) *Viral Immunology* 20:664–671), paramyxovirus (Nakamura and Russell (2004) *Expert Opinion on Biological Therapy* 4:1685–1692; Hammond et al. (2001) *Journal of Virology* 75:2087–2096; Galanis (2010) *Clinical Pharmacology and Therapeutics* 88:620–625; Blechacz and Russell (2008) *Current Gene Therapy* 8:162–175; Russell and Peng (2009) *Current Topics in Microbiology and Immunology* 330:213–241), and herpesvirus (Shah and Breakefield (2006) *Current Gene Therapy* 6:361–370; Campadelli-Fiume et al. (2011) *Reviews in Medical Virology* 21:213–226; each of which references is incorporated herein in its entirety by reference).

[0150] In some embodiments, a gene therapy vector as described herein is pseudotyped to those tissues that are particularly suited for generating a regulatory response, e.g., tolerance toward, e.g., the replacement enzyme. Such tissues include, but are not limited to mucosal tissue, e.g., gut-associated lymphoid tissue (GALT), hematopoietic stem cells, and the liver. In some embodiments, the gene therapy vector, or gene encoding a multidomain therapeutic protein as described herein is expressed under the control of promoters specific for those tissues, e.g., a liver specific promoter.

[0151] In some embodiments, a gene therapy vector as described herein comprises a naked polynucleotide. For example, in some embodiments, a polynucleotide encoding a multidomain therapeutic polypeptide may be injected, e.g., intramuscularly, directly into an organ for the formation of a depot, intravenously, etc. Additional methods well-known for the enhanced delivery of naked polynucleotides include but are not limited to electroporation, sonoporation, use of a gene gun to shoot polynucleotides coated gold particles, magnetofection, and hydrodynamic delivery.

[0152] In some embodiments, a gene therapy vector as described herein comprises polynucleotide complexes, such as, but not limited to, nanoparticles (e.g., polynucleotide self-assembled nanoparticles, polymer-based self-assembled nanoparticles, inorganic nanoparticles, lipid nanoparticles, semiconductive/metallic nanoparticles), gels and hydrogels, polynucleotide complexes with cations and anions, microparticles, and any combination thereof.

[0153] In some embodiments, the polynucleotides disclosed herein may be formulated as self-assembled nanoparticles. As a non-limiting example, polynucleotides may be used to make nanoparticles which may be used in a delivery system for the polynucleotides (See e.g., International Pub. No. WO2012125987; herein incorporated by reference in its entirety). In some embodiments, the polynucleotide self-assembled nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

[0154] In some embodiment, these self-assembled nanoparticles may be microsponges formed of long polymers of polynucleotide hairpins which form into crystalline 'pleated' sheets before self-assembling into microsponges. These microsponges are densely-packed sponge like microparticles which may function as an efficient carrier and may be able to deliver cargo to a cell. The microsponges may be from 1 μm to 300 nm in diameter. The microsponges may be complexed with other agents known in the art to form larger microsponges. As a non-limiting example, the microsphere may be complexed with an agent to form an outer layer to promote cellular uptake such as polycation polyethyleneimine (PEI). This complex can form a 250-nm diameter particle that can remain stable at high temperatures (150 °C.) (Grabow and Jaegar, Nature Materials 2012, 11:269-269; herein incorporated by reference in its entirety). Additionally, these microsponges may be able to exhibit an extraordinary degree of protection from degradation by ribonucleases. In another embodiment, the polymer-based self-assembled nanoparticles such as, but not limited to, microsponges, may be fully programmable

nanoparticles. The geometry, size and stoichiometry of the nanoparticle may be precisely controlled to create the optimal nanoparticle for delivery of cargo such as, but not limited to, polynucleotides.

[0155] In some embodiments, polynucleotides may be formulated in inorganic nanoparticles (U.S. Pat. No. 8,257,745, herein incorporated by reference in its entirety). The inorganic nanoparticles may include, but are not limited to, clay substances that are water swellable. As a non-limiting example, the inorganic nanoparticle may include synthetic smectite clays which are made from simple silicates (See e.g., U.S. Pat. Nos. 5,585,108 and 8,257,745 each of which are herein incorporated by reference in their entirety).

[0156] In some embodiments, a polynucleotide may be formulated in water-dispersible nanoparticle comprising a semiconductive or metallic material (U.S. Pub. No. 20120228565; herein incorporated by reference in its entirety) or formed in a magnetic nanoparticle (U.S. Pub. No. 20120265001 and 20120283503; each of which is herein incorporated by reference in its entirety). The water-dispersible nanoparticles may be hydrophobic nanoparticles or hydrophilic nanoparticles.

[0157] In some embodiments, the polynucleotides disclosed herein may be encapsulated into any hydrogel known in the art which may form a gel when injected into a subject. Hydrogels are a network of polymer chains that are hydrophilic, and are sometimes found as a colloidal gel in which water is the dispersion medium. Hydrogels are highly absorbent (they can contain over 99% water) natural or synthetic polymers. Hydrogels also possess a degree of flexibility very similar to natural tissue, due to their significant water content. The hydrogel described herein may be used to encapsulate lipid nanoparticles which are biocompatible, biodegradable and/or porous.

[0158] As a non-limiting example, the hydrogel may be an aptamer-functionalized hydrogel. The aptamer-functionalized hydrogel may be programmed to release one or more polynucleotides using polynucleotide hybridization. (Battig et al., J. Am. Chem. Society. 2012 134:12410-12413; herein incorporated by reference in its entirety). In some embodiment, the polynucleotide may be encapsulated in a lipid nanoparticle and then the lipid nanoparticle may be encapsulated into a hydrogel.

[0159] In some embodiments, the polynucleotides disclosed herein may be encapsulated into a fibrin gel, fibrin hydrogel or fibrin glue. In another embodiment, the polynucleotides may be formulated in a lipid nanoparticle or a rapidly eliminated lipid nanoparticle prior to being

encapsulated into a fibrin gel, fibrin hydrogel or a fibrin glue. In yet another embodiment, the polynucleotides may be formulated as a lipoplex prior to being encapsulated into a fibrin gel, hydrogel or a fibrin glue. Fibrin gels, hydrogels and glues comprise two components, a fibrinogen solution and a thrombin solution which is rich in calcium (See e.g., Spicer and Mikos, Journal of Controlled Release 2010. 148: 49-55; Kidd et al. Journal of Controlled Release 2012. 157:80-85; each of which is herein incorporated by reference in its entirety). The concentration of the components of the fibrin gel, hydrogel and/or glue can be altered to change the characteristics, the network mesh size, and/or the degradation characteristics of the gel, hydrogel and/or glue such as, but not limited to changing the release characteristics of the fibrin gel, hydrogel and/or glue. (See e.g., Spicer and Mikos, Journal of Controlled Release 2010. 148: 49-55; Kidd et al. Journal of Controlled Release 2012. 157:80-85; Catelas et al. Tissue Engineering 2008. 14:119-128; each of which is herein incorporated by reference in its entirety). This feature may be advantageous when used to deliver the polynucleotide disclosed herein. (See e.g., Kidd et al. Journal of Controlled Release 2012. 157:80-85; Catelas et al. Tissue Engineering 2008. 14:119-128; each of which is herein incorporated by reference in its entirety).

[0160] In some embodiments, a polynucleotide disclosed herein may include cations or anions. In one embodiment, the formulations include metal cations such as, but not limited to, Zn^{2+} , Ca^{2+} , Cu^{2+} , Mg^{+} and combinations thereof. As a non-limiting example, formulations may include polymers and a polynucleotide complexed with a metal cation (See e.g., U.S. Pat. Nos. 6,265,389 and 6,555,525, each of which is herein incorporated by reference in its entirety).

[0161] In some embodiments, a polynucleotide may be formulated in nanoparticles and/or microparticles. These nanoparticles and/or microparticles may be molded into any size shape and chemistry. As an example, the nanoparticles and/or microparticles may be made using the PRINT® technology by LIQUIDA TECHNOLOGIES.RTM. (Morrisville, N.C.) (See e.g., International Pub. No. WO2007024323; herein incorporated by reference in its entirety).

[0162] In some embodiments, the polynucleotides disclosed herein may be formulated in NanoJackets and NanoLiposomes by Keystone Nano (State College, Pa.). NanoJackets are made of compounds that are naturally found in the body including calcium, phosphate and may also include a small amount of silicates. Nanojackets may range in size from 5 to 50 nm and may be used to deliver hydrophilic and hydrophobic compounds such as, but not limited to, polynucleotides, primary constructs and/or polynucleotide. NanoLiposomes are made of lipids such as, but not limited to, lipids which naturally occur in the body. NanoLiposomes may range

in size from 60-80 nm and may be used to deliver hydrophilic and hydrophobic compounds such as, but not limited to, polynucleotides, primary constructs and/or polynucleotide. In one aspect, the polynucleotides disclosed herein are formulated in a NanoLiposome such as, but not limited to, Ceramide NanoLiposomes.

[0163] In one embodiment, the multidomain therapeutic protein is an anti-CD63 scFv-GAA fusion protein or an anti-ITGA7 scFv-GAA fusion protein. The administration of the anti-CD63 scFv-GAA fusion protein or the anti-ITGA7 scFv-GAA fusion protein via AAV-delivery provides long term stable production of GAA in the serum of the patient after administration of the multidomain therapeutic protein-harboring gene therapy vector. In one embodiment, the level of GAA in the serum of the recipient patient is ≥ 1.5 fold to 100 fold, ≥ 1.5 fold to 10 fold, ≥ 2.5 fold, 2.5 fold – 3 fold, 2.5 fold, 2.6 fold, 2.7 fold, 2.8 fold, 2.9 fold, 3.0 fold, 3.1 fold, 3.2 fold, 3.3 fold, 3.4 fold, 3.5 fold, 3.6 fold, 3.7 fold, 3.8 fold, 3.9 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, or 10 fold greater than the serum levels of a patient receiving GAA not linked to a delivery domain after 1 month, 3 months, 4 months, 5 months, or 6 months after administration of the multidomain therapeutic protein-harboring gene therapy vector.

[0164] In one embodiment, the administration of the anti-CD63 scFv-GAA fusion protein or the anti-ITGA7 scFv-GAA fusion protein via AAV-delivery provides long term stable reduction in stored glycogen levels in patients with Pompe disease. In one embodiment, the glycogen levels in heart, skeletal muscle, and liver tissue in the patient are reduced to wildtype (non-disease) levels. In one embodiment, the glycogen levels in heart, skeletal muscle, and liver tissue in the patient are maintained at wildtype levels 1 month, 2 months, 3 months, 4 months, 5 months, or 6 months after administration of the multidomain therapeutic protein-harboring gene therapy vector.

[0165] In one embodiment, the administration of the anti-CD63 scFv-GAA fusion protein or the anti-ITGA7 scFv-GAA fusion protein via AAV-delivery provides long term restoration of muscle strength in patients with Pompe disease. In one embodiment, the strength of the patient as measured by grip strength is restored to normal (*i.e.*, non-disease normal levels) 1 month, 2 months, 3 months, 4 months, 5 months, or 6 months after administration of the multidomain therapeutic protein-harboring gene therapy vector.

[0166] In another aspect, the invention provides a composition comprising an enzyme activity and an antigen-binding protein, wherein the enzyme is associated with an enzyme-deficiency disease (LSD) and internalizing effector-binding protein. Enzymes (which include proteins that

are not *per se* catalytic) associated with lysosomal storage diseases include for example any and all hydrolases, α -galactosidase, β -galactosidase, α -glucosidase, β -glucosidase, saposin-C activator, ceramidase, sphingomyelinase, β -hexosaminidase, GM2 activator, GM3 synthase, arylsulfatase, sphingolipid activator, α -iduronidase, iduronidase-2-sulfatase, heparin *N*-sulfatase, *N*-acetyl- α -glucosaminidase, α -glucosamide *N*-acetyltransferase, *N*-acetylglucosamine-6-sulfatase, *N*-acetylgalactosamine-6-sulfate sulfatase, *N*-acetylgalactosamine-4-sulfatase, β -glucuronidase, hyaluronidase, and the like.

[0167] Internalizing effector-binding proteins for example include a receptor-fusion molecule, a trap molecule, a receptor-Fc fusion molecule, an antibody, an Fab fragment, an F(ab')₂ fragment, an Fd fragment, an Fv fragment, a single-chain Fv (scFv) molecule, a dAb fragment, an isolated complementarity determining region (CDR), a CDR3 peptide, a constrained FR3-CDR3-FR4 peptide, a domain-specific antibody, a single domain antibody, a domain-deleted antibody, a chimeric antibody, a CDR-grafted antibody, a diabody, a triabody, a tetrabody, a minibody, a nanobody, a monovalent nanobody, a bivalent nanobody, a small modular immunopharmaceutical (SMIP), a camelid antibody (VHH heavy chain homodimeric antibody), a shark variable IgNAR domain, other antigen-binding proteins, and the like.

[0168] Internalizing effectors include for example CD63, MHC-I, Kremen-1, Kremen- 2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), PRLR (prolactin receptor), MAL (Myelin And Lymphocyte protein, a.k.a. VIP17), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptor, scavenger receptor, SCARA1-5, SCARB1-3, and CD36. In certain embodiments, the internalizing effector is a kidney specific internalizer, such as CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin). In other certain embodiments, the internalizing effector is a muscle specific internalizer, such as BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-1S), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1),

and POPDC3 (Popeye domain-containing protein 3). In some specific embodiments, the internalizing effector is ITGA7, CD9, CD63, ALPL2, ASGR1, ASGR2 or PRLR.

[0169] In some embodiments, the enzyme is covalently linked (*i.e.*, electrons shared across atoms) to the antigen-binding protein. In one particular embodiment, the internalizing effector-binding protein consists of or contains a half-body; the enzyme is fused to an Fc-fusion domain (*e.g.*, at the C-terminus); and the Fc-domain that is covalently linked to the enzyme associates with the Fc-domain of the antigen-binding protein, such that the association contains one or more disulfide bridges. This particular embodiment is schematically depicted in Figure 1A, panel B.

[0170] In another particular embodiment, the internalizing effector-binding protein (delivery domain) consists of or contains an antibody or an antibody fragment, and the enzyme is covalently linked to the antibody or antibody fragment. In a specific embodiment, the delivery domain is an antibody, and the enzyme is covalently linked (directly through a peptide bond, or indirectly via a linker) to the C-terminus of the heavy chain or the light chain of the antibody (Figure 1A, panels C or E, respectively). In another specific embodiment, the delivery domain is an antibody, and the enzyme is covalently linked (directly through a peptide bond, or indirectly via a linker) to the N-terminus of the heavy chain or the light chain of the antibody (Figure 1A, panels D or F, respectively).

[0171] In some embodiments, the enzyme and delivery domain are not covalently linked, but are combined in an admixture. The delivery domain and the enzyme can associate through non-covalent forces to form a complex. For example, in one particular embodiment, the delivery domain is a bispecific antibody in which one arm of the antibody binds the internalizing effector and the other arm binds the enzyme. This embodiment is schematically depicted in Figure 1A, panel A.

[0172] In some embodiments, the enzyme is GAA or comprises GAA activity (*e.g.*, an isozyme with GAA activity), and the internalizing effector is ITGA7, CDH15, CD9, CD63, APLP2, ASGR1, ASGR2 or PRLR. In a particular embodiment, the enzyme is GAA or comprises GAA activity, the internalization domain is CD63, and the delivery domain is a bispecific antibody with specificity for CD63 and GAA. In a particular embodiment, the enzyme is GAA or comprises GAA activity, the internalization domain is ITGA7, and the delivery domain is a bispecific antibody with specificity for ITGA7 and GAA.

[0173] In some embodiments, the enzyme is GLA or comprises GLA activity (*e.g.*, an isozyme with GAA activity), and the internalizing effector is ITGA7, CD9, CD63, APLP2, ASGR1, ASGR2, or PRLR. In a particular embodiment, the enzyme is GLA or comprises GLA activity, the internalization domain is CD63, and the delivery domain is a bispecific antibody with specificity for CD63 and GLA. In a particular embodiment, the enzyme is GLA or comprises GLA activity, the internalization domain is ITGA7, and the delivery domain is a bispecific antibody with specificity for ITGA7 and GLA.

Pharmaceutical Compositions and Administration thereof

[0174] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's The Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro (Lippincott, Williams & Wilkins, Baltimore, Md., 2006; incorporated herein by reference in its entirety) discloses various excipients used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional excipient medium is incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention.

[0175] In some embodiments, a pharmaceutically acceptable excipient is at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% pure. In some embodiments, an excipient is approved for use in humans and for veterinary use. In some embodiments, an excipient is approved by United States Food and Drug Administration. In some embodiments, an excipient is pharmaceutical grade. In some embodiments, an excipient meets the standards of the United States Pharmacopoeia (USP), the European Pharmacopoeia (EP), the British Pharmacopoeia, and/or the International Pharmacopoeia.

[0176] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in pharmaceutical compositions.

[0177] Exemplary diluents include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, etc., and/or combinations thereof.

[0178] Exemplary granulating and/or dispersing agents include, but are not limited to, potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (VEEGUM®), sodium lauryl sulfate, quaternary ammonium compounds, etc., and/or combinations thereof.

[0179] Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (e.g. bentonite [aluminum silicate] and VEEGUM® [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g. carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (e.g. carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (e.g. polyoxyethylene sorbitan monolaurate [TWEEN® 20], polyoxyethylene sorbitan [TWEEN® 60], polyoxyethylene sorbitan monooleate [TWEEN® 80], sorbitan monopalmitate [SPAN® 40], sorbitan monostearate [SPAN® 60], sorbitan tristearate [SPAN® 65], glyceryl monooleate, sorbitan monooleate [SPAN® 80]), polyoxyethylene esters (e.g. polyoxyethylene monostearate [MYRJ® 45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and SOLUTOL®), sucrose fatty acid esters, polyethylene glycol fatty acid esters (e.g. CREMOPHOR®), polyoxyethylene ethers, (e.g. polyoxyethylene lauryl ether [BRIJ® 30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid,

ethyl laurate, sodium lauryl sulfate, PLUORINC® F 68, POLOXAMER® 188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, etc. and/or combinations thereof.

[0180] Exemplary binding agents include, but are not limited to, starch (e.g. cornstarch and starch paste); gelatin; sugars (e.g. sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol,); natural and synthetic gums (e.g. acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum®), and larch arabogalactan); alginates; polyethylene oxide; polyethylene glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; etc.; and combinations thereof.

[0181] Exemplary preservatives may include, but are not limited to, antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and/or other preservatives. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfate, sodium metabisulfite, and/or sodium sulfite. Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate, edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and/or trisodium edetate. Exemplary antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and/or thimerosal. Exemplary antifungal preservatives include, but are not limited to, butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and/or sorbic acid. Exemplary alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and/or phenylethyl alcohol. Exemplary acidic preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and/or phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate,

deteroxime mesylate, cetrimide, butylated hydroxyanisol (BHA), butylated hydroxytoluened (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, GLYDANT PLUS®, PHENONIP®, methylparaben, GERMALL® 115, GERMABEN® II, NEOLONE.TM., KATHON.TM., and/or EUXYL®.

[0182] Exemplary buffering agents include, but are not limited to, citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, etc., and/or combinations thereof.

[0183] Exemplary lubricating agents include, but are not limited to, magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behenate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, etc., and combinations thereof.

[0184] Exemplary oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, chamomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macadamia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and/or combinations thereof.

[0185] Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and/or perfuming agents can be present in the composition, according to the judgment of the formulator.

[0186] Delivery

[0187] The present disclosure encompasses the delivery of the gene therapy vector (e.g., the polynucleotides) by any appropriate route taking into consideration likely advances in the sciences of drug delivery. Delivery may be naked or formulated.

[0188] Naked Delivery

[0189] The polynucleotides of the present invention may be delivered to a cell naked. As used herein in, "naked" refers to delivering polynucleotides free from agents which promote transfection. For example, the polynucleotides delivered to the cell may contain no modifications. The naked polynucleotides may be delivered to the cell using routes of administration known in the art and described herein.

[0190] Formulated Delivery

[0191] The polynucleotides may be formulated, using the methods described herein. The formulations may contain polynucleotides and may further include, but are not limited to, cell penetration agents, a pharmaceutically acceptable carrier, a delivery agent, a bioerodible or biocompatible polymer, a solvent, and a sustained-release delivery depot. The formulated polynucleotides mRNA may be delivered to the cell using routes of administration known in the art and described herein.

[0192] Administration

[0193] The polynucleotides of the present invention may be administered by any route which results in a therapeutically effective outcome. These include, but are not limited to enteral, gastroenteral, epidural, oral, transdermal, epidural (peridural), intracerebral (into the cerebrum), intracerebroventricular (into the cerebral ventricles), epicutaneous (application onto the skin), intradermal, (into the skin itself), subcutaneous (under the skin), nasal administration (through the nose), intravenous (into a vein), intraarterial (into an artery), intramuscular (into a muscle), intracardiac (into the heart), intraosseous infusion (into the bone marrow), intrathecal (into the spinal canal), intraperitoneal, (infusion or injection into the peritoneum), intravesical infusion, intravitreal, (through the eye), intracavernous injection, (into the base of the penis), intravaginal administration, intrauterine, extra-amniotic administration, transdermal (diffusion through the

intact skin for systemic distribution), transmucosal (diffusion through a mucous membrane), insufflation (snorting), sublingual, sublabial, enema, eye drops (onto the conjunctiva), or in ear drops. In specific embodiments, compositions may be administered in a way which allows them to cross the blood-brain barrier, vascular barrier, or other epithelial barrier. Non-limiting routes of administration for the polynucleotides, primary constructs or mRNA of the present invention are described below.

[0194] Parenteral and Injectable Administration

[0195] Liquid dosage forms for parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as CREMOPHOR®, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

[0196] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

[0197] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid

compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0198] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[0199] Depot Administration

[0200] As described herein, in some embodiments, the composition is formulated in depots for extended release. Generally, a specific organ or tissue (a "target tissue") is targeted for administration.

[0201] In some aspects of the invention, the polynucleotides are spatially retained within or proximal to a target tissue. Provided are method of providing a composition to a target tissue of a mammalian subject by contacting the target tissue (which contains one or more target cells) with the composition under conditions such that the composition, in particular the nucleic acid component(s) of the composition, is substantially retained in the target tissue, meaning that at least 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the composition is retained in the target tissue. Advantageously, retention is determined by measuring the amount of the nucleic acid present in the composition that enters one or more target cells. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the nucleic acids administered to the subject are present intracellularly at a period of time following administration. For example, intramuscular injection to a mammalian subject is performed using an aqueous composition containing a

polynucleotide and a transfection reagent, and retention of the composition is determined by measuring the amount of the ribonucleic acid present in the muscle cells.

[0202] Aspects of the invention are directed to methods of providing a composition to a target tissue of a mammalian subject, by contacting the target tissue (containing one or more target cells) with the composition under conditions such that the composition is substantially retained in the target tissue. The composition contains an effective amount of a polynucleotide such that the polypeptide of interest is produced in at least one target cell. The compositions generally contain a cell penetration agent, although "naked" nucleic acid (such as nucleic acids without a cell penetration agent or other agent) is also contemplated, and a pharmaceutically acceptable carrier.

[0203] In some circumstances, the amount of a protein produced by cells in a tissue is desirably increased. Preferably, this increase in protein production is spatially restricted to cells within the target tissue. Thus, provided are methods of increasing production of a protein of interest in a tissue of a mammalian subject. A composition is provided that contains polynucleotides characterized in that a unit quantity of composition has been determined to produce the polypeptide of interest in a substantial percentage of cells contained within a predetermined volume of the target tissue.

[0204] In some embodiments, the composition includes a plurality of different polynucleotides, where one or more than one of the polynucleotides encodes a polypeptide of interest. Optionally, the composition also contains a cell penetration agent to assist in the intracellular delivery of the composition. A determination is made of the dose of the composition required to produce the polypeptide of interest in a substantial percentage of cells contained within the predetermined volume of the target tissue (generally, without inducing significant production of the polypeptide of interest in tissue adjacent to the predetermined volume, or distally to the target tissue). Subsequent to this determination, the determined dose is introduced directly into the tissue of the mammalian subject.

[0205] In one embodiment, the invention provides for the polynucleotides to be delivered in more than one injection or by split dose injections.

[0206] In one embodiment, the invention may be retained near target tissue using a small disposable drug reservoir, patch pump or osmotic pump. Non-limiting examples of patch pumps include those manufactured and/or sold by BD® (Franklin Lakes, N.J.), Insulet Corporation (Bedford, Mass.), SteadyMed Therapeutics (San Francisco, Calif.), Medtronic (Minneapolis,

Minn.) (e.g., MiniMed), UniLife (York, Pa.), Valeritas (Bridgewater, N.J.), and SpringLeaf Therapeutics (Boston, Mass.). A non-limiting example of an osmotic pump include those manufactured by DURECT® (Cupertino, Calif.) (e.g., DUROS® and ALZET®).

[0207] Dosing

[0208] The present invention provides methods comprising administering a gene therapy vector comprising polynucleotide encoding a multidomain therapeutic polypeptide, and optionally subsequently the multidomain therapeutic polypeptide to a subject in need thereof. In some embodiments, a method comprises administering a gene therapy vector comprising polynucleotide encoding a multidomain therapeutic polypeptide in a therapeutically effective amount to a patient in need thereof, wherein the therapeutically effective amount is sufficient to obviate the subsequent administration of the multidomain therapeutic polypeptide.

Accordingly, in some embodiments, a method of treating a patient in need thereof lacking an enzyme, e.g., reducing glycogen levels and/or reducing CRIM to GAA in a patient with Pompe disease, comprises administering to the patient a gene therapy vector comprising a polynucleotide encoding a multidomain therapeutic protein comprising the replacement enzyme, e.g., an anti-CD63 scFv::GAA fusion protein, e.g., a multidomain therapeutic protein comprising the sequence set forth as SEQ ID NO:11, in a therapeutically effective amount, wherein the therapeutically effective amount negates the need for subsequent administration to the patient of the replacement enzyme, e.g., GAA or derivatives thereof. In some embodiments, a method of treating a patient lacking an enzyme and in need thereof, e.g., reducing glycogen levels and/or reducing CRIM to GAA in a patient with Pompe disease, comprises administering to the patient a gene therapy vector comprising a polynucleotide encoding a multidomain therapeutic protein comprising a replacement enzyme, e.g., an anti-CD63 scFv::GAA fusion protein, e.g., a multidomain therapeutic protein comprising the sequence set forth as SEQ ID NO:11, in a therapeutically effective amount, and further comprises administering to the patient a therapeutically effective amount of the replacement enzyme. Nucleic acids, proteins or complexes, or pharmaceutical, imaging, diagnostic, or prophylactic compositions thereof, may be administered to a subject using any amount and any route of administration effective for preventing, treating, diagnosing, or imaging a disease, disorder, and/or condition (e.g., a disease, disorder, and/or condition relating to working memory deficits).

[0209] The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease, the particular composition, its mode of administration, its mode of activity, and the like.

[0210] The dose of AAV viral vectors, e.g., the units of dose in vector genomes/per kilogram of body weight (vg/kg), required to achieve a desired effect or "therapeutic effect" (e.g., a certain serum concentration of a replacement enzyme) will vary based on several factors including, but not limited to: the route of AAV administration, the level of expression required to achieve a therapeutic effect, the specific disease or disorder being treated, and the stability of the expression multidomain therapeutic protein. One of skill in the art can readily determine a AAV virion dose range to treat a subject having a particular disease or disorder based on the aforementioned factors, as well as other factors that are well known in the art, see, e.g., CDER "Guidance for Industry Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers," July 2005, incorporated herein in its entirety by reference. An effective amount of the AAV is generally in the range of from about 10 μ l to about 100 ml of solution containing from about 10^9 to 10^{16} genome copies per subject. Other volumes of solution may be used. The volume used will typically depend, among other things, on the size of the subject, the dose of the AAV, and the route of administration. In some embodiments, a dosage between about 10^{10} to 10^{12} AAV viral genome per subject is appropriate. In some embodiments the AAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , or 10^{15} genome copies per subject. In some embodiments the AAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} viral genomes per kg. In some embodiments, at least 2×10^{12} viral genomes per kilogram (vg/kg) is administered. In some embodiments, the dose administered provides a threshold multidomain therapeutic protein serum level. In some embodiments the threshold therapeutic protein serum level is at least 1 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 2 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 3 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 4 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 5 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 6 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 7 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 8 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 9 μ g/mL. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 10 μ g/mL. In some embodiments, the dose administered provides a multidomain

therapeutic protein serum level of greater than 11 $\mu\text{g/mL}$. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 12 $\mu\text{g/mL}$. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 13 $\mu\text{g/mL}$. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 14 $\mu\text{g/mL}$. In some embodiments, the dose administered provides a multidomain therapeutic protein serum level of greater than 15 $\mu\text{g/mL}$.

[0211] Compositions in accordance with the invention are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention may be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or appropriate imaging dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[0212] Nonlimiting and exemplary embodiments are listed below.

Embodiment 1. A method of delivering a therapeutic protein to the central nervous system (CNS) of a subject, comprising administering to the subject a nucleotide composition encoding a multidomain therapeutic protein via a liver-targeted delivery method sufficient to provide a therapeutically effective amount of the multidomain therapeutic protein in the CNS, wherein the multidomain therapeutic protein comprises a delivery domain and an enzyme domain.

Embodiment 2. The method of embodiment 1, wherein the delivery domain is an antibody or antigen-binding fragment thereof that binds specifically to an internalizing effector.

Embodiment 3. The method of embodiment 1 or embodiment 2, wherein the therapeutic protein is a lysosomal enzyme.

Embodiment 4. The method of embodiment 3, wherein the lysosomal enzyme is GAA.

Embodiment 5. The method of any one of embodiments 1-4, wherein the nucleotide composition is administered via a viral vector, optionally wherein the nucleotide composition is administered at a dose of at least 2×10^{12} viral genomes per kilogram (vg/kg).

Embodiment 6. The method of embodiment 5, wherein the viral vector is an AAV vector.

Embodiment 7. The method of any one of embodiments 1-6, wherein the internalizing effector is expressed on the surface of cells selected from the group consisting of: cells in the CNS, epithelial cells, and cells that cross the blood brain barrier.

Embodiment 8. The method of any one of embodiments 1-7, wherein the delivery domain binds an internalizing effector

(i) selected from the group consisting of CD63, Integrin alpha-7 (ITGA7), MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), myelin and lymphocyte protein (MAL), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptor A1-5 (SCARA1-5), SCARB1-3, and CD36;

(ii) expressed in several tissue types, optionally selected from the group consisting of CD63, MHC-I, vacuolar-type H⁺ ATPase, IGF2R, Integrin alpha-7 (ITGA7), LRP5, LRP6, LRP8, Kremen-2, LDL- receptor, LDL-related protein 1 receptor, amyloid precursor protein-like protein- 2 (APLP2), apelin receptor (APLNR), PRLR, MAL (myelin and lymphocyte protein (MAL), diphtheria toxin receptors, HBEGF (heparin binding EGF like growth factor), glutathione receptors, glutamate receptors, leptin receptors, and folate receptors,

optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of Fabry disease, Gaucher disease, MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVB, MPS VI, MPS VII, MPS IX, Pompe disease, Lysosomal acid lipase deficiency, Metachromatic leukodystrophy, Niemann-Pick diseases types A, B, and C2, Alpha mannosidosis, Neuraminidase deficiency, Sialidosis, Aspartylglycosaminuria, Combined saposin deficiency, Atypical Gaucher disease, Farber lipogranulomatosis, Fucosidosis, and Beta mannosidosis;

(iii) preferentially expressed by bone and/or cartilage, optionally selected from the group consisting of Collagen X, Integrin alpha 10 (ITGA10), Fibroblast growth factor receptor 3

(FGFR3), Fibroblast growth factor receptor isoform C (FGFR3C), Hyaluronan and proteoglycan link protein 1 (CRTL1), Aggrecan, Collagen II, and Kremen-1,

optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of MPS I, MPS II, MPS IIIA, MPS IIIB, MPS IIID, MPS IVA, MPS IVB, MPS VI, MPS VII, MPS IX, Beta mannosidosis, Gaucher disease, atypical Gaucher disease, combined Saposin deficiency, Aspartylglycosaminuria, Farber lipogranulomatosis, Sialidosis, Neuraminidase deficiency, and Alpha mannosidosis;

(iv) preferentially expressed by monocytes, macrophages, or microglia, optionally selected from the group consisting of scavenger receptor A1-5 (SCARA1-5), SCARB1-3, CD36, MSR1 (macrophage scavenger receptor 1), MRC1 (macrophage mannose receptor 1), VSIG4 (V-set and immunoglobulin domain-containing protein 4), CD68 (Macrosialin), and CSF1R (Macrophage colony-stimulating factor 1 receptor),

optionally wherein the subject exhibits one or more symptoms of a disease selected from the group consisting of lysosomal acid lipase deficiency, Gaucher disease, Atypical Gaucher disease, combined Saposin deficiency, and Farber lipogranulomatosis;

(v) preferentially expressed by kidney cells, optionally selected from the group consisting of CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin),

optionally wherein the subject exhibits one or more symptoms or is diagnosed with a disease selected from the group consisting of Fabry disease, Alport syndrome, polycystic kidney disease, and Thrombotic Thrombocytopenic Purpura;

(vi) preferentially expressed by liver cells, optionally ASGR1 or ASGR2,

optionally wherein the subject exhibits one or more symptoms or is diagnosed with a disease selected from the group consisting of as lysosomal acid lipase deficiency, Gaucher disease, MPS VI, MPS VII, MPS II, Niemann-Pick diseases types A, B, and C2, Sialidosis, Neuraminidase deficiency, atypical Gaucher disease, combined Saposin deficiency, Farber lipogranulomatosis;

(vii) preferentially expressed by muscle cells, optionally selected from the group consisting of BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic

receptor (nicotinic) alpha 1, CDH15 (Cadheri-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-15), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3),

optionally wherein the subject exhibits one or more symptoms or is diagnosed with Pompe disease;

(viii) selected from the group consisting of ITGA7, CD9, CD63, ALPL2, MSR1, ASGR1, ASGR2, or PRLR; or

(ix) that is CD63.

Embodiment 9. The method of any one of embodiments 1-8, wherein the delivery domain is a single-chain variable fragment (scFv).

Embodiment 10. The method of any one of embodiments 1-9, wherein the cell surface receptor (CSR)-binding protein (CSR-BP) comprises an amino acid sequence of SEQ ID NO:2.

Embodiment 11. The method of any one of embodiments 1-10, wherein the therapeutic protein comprises a hydrolase.

Embodiment 12. The method of any one of embodiments 1-11, wherein the therapeutic protein comprises a glycosylase.

Embodiment 13. The method of any one of embodiments 1-12, wherein the therapeutic protein comprises a glycosidase.

Embodiment 14. The method of any one of embodiments 1-13, wherein the therapeutic protein comprises an alpha-glucosidase.

Embodiment 15. The method of any one of embodiments 1-14, wherein the therapeutic protein comprises an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:13, or a fragment thereof.

Embodiment 16. The method of any one of embodiments 1-15, wherein the therapeutic protein comprises an anti-ABeta, or an anti-Tau antibody.

Embodiment 17. The method of any one of embodiments 1-16, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO:11.

Embodiment 18. The method of any one of embodiments 1-17, wherein the enzyme domain comprises an alpha-glucosidase, and wherein the glycogen levels in any CNS tissue in the subject is reduced for at least nine months post-treatment.

Embodiment 19. The method of any one of embodiments 1-18, wherein the subject has Pompe disease.

Embodiment 20. The method of any one of embodiments 1-19, wherein the administered nucleotide composition provides a multidomain therapeutic protein serum level of at least 1 µg/mL.

Embodiment 21. A multidomain therapeutic protein comprising one or more delivery domain(s) and an enzyme domain, wherein the one or more delivery domain(s) binds human transferrin receptor (hTfR).

Embodiment 22. The multidomain therapeutic protein of embodiment 21, further comprising a second delivery domain that binds to an internalizing effector.

Embodiment 23. The multidomain therapeutic protein of embodiment 22, wherein the second delivery domain binds to

(i) an internalizing effector selected from the group consisting of CD63, Integrin alpha-7 (ITGA7), MHC-I, Kremen-1, Kremen-2, LRP5, LRP6, LRP8, transferrin receptor, LDL-receptor, LDL-related protein 1 receptor, ASGR1, ASGR2, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), myelin and lymphocyte protein (MAL), IGF2R, vacuolar-type H⁺ ATPase, diphtheria toxin receptor, folate receptor, glutamate receptors, glutathione receptor, leptin receptors, scavenger receptor A1-5 (SCARA1-5), SCARB1-3, and CD36;

(ii) an internalizing effector expressed in several tissue types, optionally selected from the group consisting of CD63, MHC-I, vacuolar-type H⁺ ATPase, IGF2R, Integrin alpha-7 (ITGA7), LRP5, LRP6, LRP8, Kremen-2, LDL-receptor, LDL-related protein 1 receptor, amyloid precursor protein-like protein-2 (APLP2), apelin receptor (APLNR), PRLR, MAL (myelin and lymphocyte protein (MAL), diphtheria toxin receptors, HBEGF (heparin binding EGF like growth factor), glutathione receptors, glutamate receptors, leptin receptors, and folate receptors;

(iii) an internalizing effector preferentially expressed by bone and/or cartilage, optionally selected from the group consisting of Collagen X, Integrin alpha 10 (ITGA10), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor isoform C (FGFR3C), Hyaluronan and proteoglycan link protein 1 (CRTL1), Aggrecan, Collagen II, and Kremen-1;

(iv) an internalizing effector preferentially expressed by monocytes, macrophages, or microglia, optionally selected from the group consisting of scavenger receptor A1-5 (SCARA1-5), SCARB1-3, CD36, MSR1 (macrophage scavenger receptor 1), MRC1 (macrophage mannose receptor 1), VSIG4 (V-set and immunoglobulin domain-containing protein 4), CD68 (Macrosialin), and CSF1R (Macrophage colony-stimulating factor 1 receptor);

(v) an internalizing effector preferentially expressed by kidney cells, optionally selected from the group consisting of CDH16 (Cadherin-16), CLDN16 (Claudin-16), KL (Klotho), PTH1R (parathyroid hormone receptor), SLC22A13 (Solute carrier family 22 member 13), SLC5A2 (Sodium/glucose cotransporter 2), and UMOD (Uromodulin). In other certain embodiments, the internalization effector is a muscle specific internalizer, such as BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-1S), CACNG6 (L-type calcium channel subunit gamma-6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3);

(vi) an internalizing effector preferentially expressed by liver cells, optionally ASGR1 or ASGR2;

(vii) an internalizing effector preferentially expressed by muscle cells, optionally selected from the group consisting of BMPRI1A (Bone morphogenetic protein receptor 1A), m-cadherin, CD9, MuSK (muscle-specific kinase), LGR4/GPR48 (G protein-coupled receptor 48), cholinergic receptor (nicotinic) alpha 1, CDH15 (Cadherin-15), ITGA7 (Integrin alpha-7), CACNG1 (L-type calcium channel subunit gamma-1), CACNA1S (L-type calcium channel subunit alpha-1S), CACNG6 (L-type calcium channel subunit gamma-6),

6), SCN1B (Sodium channel subunit beta-1), CHRNA1 (ACh receptor subunit alpha), CHRND (ACh receptor subunit delta), LRRC14B (Leucine-rich repeat-containing protein 14B), dystroglycan (DAG1), and POPDC3 (Popeye domain-containing protein 3), or

(viii) an internalizing effector protein selected from the group consisting of ITGA7, CD9, CD63, ALPL2, MSR1, ASGR1, ASGR2, or PRLR.

Embodiment 24. The multidomain therapeutic protein of any one of embodiments 21-23, wherein the second delivery domain binds to the internalizing effector CD63.

Embodiment 25. The multidomain therapeutic protein of any one of embodiments 21-24, wherein at least one of the one or more delivery domain(s) comprises an antigen-binding protein.

Embodiment 26. The multidomain therapeutic protein of embodiment 25, wherein each of the one or more delivery domain(s) comprises an antigen-binding protein.

Embodiment 27. The multidomain therapeutic protein of any one of embodiments 21-26, wherein at least one of the one or more delivery domain(s) comprises a single-chain variable fragment (scFv).

Embodiment 28. The multidomain therapeutic protein of any one of embodiments 21-27, wherein at least one of the one or more delivery domain(s) comprises a half-body.

Embodiment 29. The multidomain therapeutic protein of embodiment 28, wherein the delivery domain that binds hTfR is an scFv, wherein the half-body binds CD63, wherein the enzyme domain is GAA, and wherein GAA is conjugated to the carboxy terminus of the half-body that binds CD63.

Embodiment 30. The multidomain therapeutic protein of embodiment 27, wherein each of the one or more delivery domain(s) comprises an scFv.

Embodiment 31. The multidomain therapeutic protein of any one of embodiments 27-30, wherein at least one scFv is fused to an Fc.

Embodiment 32. The multidomain therapeutic protein of embodiment 31, wherein the Fc comprises a wildtype human IgG4 isotype, or derivative thereof.

Embodiment 33. The multidomain therapeutic protein of any one of embodiments 31-32, wherein GAA is conjugated to the carboxy terminus of the Fc.

Embodiment 34. The multidomain therapeutic protein of embodiment 30, comprising an anti-hTfR scFv, an anti-hCD63 scFv.

Embodiment 35. The multidomain therapeutic protein of embodiment 34, wherein the anti-hTfR scFv and anti-hCD63 scFv are both linked, at their carboxy termini, to a single GAA enzyme.

Embodiment 36. The multidomain therapeutic protein of any one of embodiments 21-27, wherein the delivery domain is an anti-hTfR scFv, and the enzyme domain is linked to carboxy terminus of the VL domain of the scFv.

Embodiment 37. The multidomain therapeutic protein of embodiment 36, further comprising a second delivery domain linked to the N-terminus of the VH domain of the anti-hTfR scFv.

Embodiment 38. The multidomain therapeutic protein of embodiment 37, wherein the second delivery domain is an anti-hCD63 scFv.

Embodiment 39. The multidomain therapeutic protein of any one of the preceding embodiments, wherein the enzyme domain comprises the amino acid sequence set forth as SEQ ID NO:1

Embodiment 40. A polynucleotide encoding the multidomain therapeutic protein of any one of embodiments 21-39 or embodiments 50-64.

Embodiment 41. The polynucleotide of embodiment 40, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence.

Embodiment 42. The polynucleotide of embodiment 40 or embodiment 41, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence.

Embodiment 43. The polynucleotide of any one of embodiments 40-42, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence, and wherein the AAV nucleic acid sequence comprises an internal terminal repeat sequence, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter.

Embodiment 44. The polynucleotide of any one of embodiments 40-43, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence comprising an internal terminal repeat sequence that comprises SEQ ID NO:6, SEQ ID NO:7, or both, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter.

Embodiment 45. The polynucleotide of any one of embodiments 40-44 further comprising a tissue specific regulatory element comprising the sequence set forth as SEQ ID NO:8 and/or SEQ ID NO:9.

Embodiment 46. A gene therapy vector comprising a polynucleotide of any one of embodiments 40-45.

Embodiment 47. The gene therapy vector of embodiment 46, wherein the gene therapy vector is selected from the group consisting of

- a viral vector, optionally wherein the viral vector is a natural virus, an engineered virus, or a chimeric virus,

- a naked polynucleotide comprising the polynucleotide of any one of embodiments 20-25,

- a polynucleotide complex, optionally wherein the polynucleotide complex is a lipid nanoparticle comprising the polynucleotide of any one of embodiments 20-25 and lipids, and

- any combination thereof.

Embodiment 48. The gene therapy vector of embodiment 46 or embodiment 47, wherein the gene therapy vector is a viral vector selected from the group consisting of a retrovirus, adenovirus, herpes simplex virus, pox virus, vaccinia virus, lentivirus, or an adeno-associated virus.

Embodiment 49. The gene therapy vector of embodiment 47 or embodiment 48, wherein the gene therapy vector is AAV9, Anc80, an AAV2/8 chimera and/or an AAV pseudotyped to a specific tissue, e.g., the liver or neuronal tissue.

Embodiment 50. A multidomain therapeutic protein comprising at least two delivery domains and at least one enzyme domain, wherein each of the two delivery domains is independently selected from the group consisting of an antibody, a half-body, and an scFv,

and wherein at least one or more of the delivery domains is associated the at least one enzyme domain, preferably wherein the one or more delivery domains is covalently linked to the at least one enzyme domain.

Embodiment 51. The multidomain therapeutic protein of embodiment 50, comprising no more than two delivery domains.

Embodiment 52. The multidomain therapeutic protein of embodiment 50 or embodiment 51, wherein only one of the delivery domains is associated with the at least one enzyme domain.

Embodiment 53. The multidomain therapeutic protein of any one of embodiments 50-52, wherein each of the at least two delivery domains is covalently linked to an enzyme domain.

Embodiment 54. The multidomain therapeutic protein of embodiment 53, wherein each of the at least two delivery domains is covalently linked to the same enzyme domain.

Embodiment 55. The multidomain therapeutic protein of embodiment 53, wherein each of the at least two delivery domains is covalently linked to a different enzyme domain.

Embodiment 56. The multidomain therapeutic protein of any one of embodiments 50-55, wherein the multidomain therapeutic protein comprises no more than two delivery domains, wherein the first delivery domain comprises a half-body, and wherein the second delivery domain comprises an scFv.

Embodiment 57. The multidomain therapeutic protein of embodiment 56, wherein the scFv is fused to an Fc.

Embodiment 58. The multidomain therapeutic protein of embodiment 56 or embodiment 57, wherein the half-body is covalently linked at its carboxy terminus to a first enzyme domain and/or wherein the scFv is covalently linked at its carboxy terminus to an Fc, and optionally, a second enzyme domain.

Embodiment 59. The multidomain therapeutic protein of any one of embodiments 50-55, wherein the multidomain therapeutic protein comprises no more than two delivery domains, wherein the first and second delivery domains each comprise an scFv.

Embodiment 60. The multidomain therapeutic protein of embodiment 59, wherein both the first and second scFv are covalently linked to an enzyme domain.

Embodiment 61. The multidomain therapeutic protein of embodiment 59, comprising from N-terminus to C-terminus: the first scFv, the second scFv, and the enzyme domain.

Embodiment 62. The multidomain therapeutic protein of any one of embodiments 50-61, wherein at least one delivery domain binds a lysosomal trafficking molecule and at least one delivery domain binds a transcytosis effector.

Embodiment 63. The multidomain therapeutic protein of embodiment 62, wherein the lysosomal trafficking molecule is selected from the group consisting of CD63, ITGA7, CD9, CD63, CD81, CD82, or CD151, and wherein the transcytosis effector is selected from the group consisting of an LDL receptor, an IgA receptor, a transferrin receptor, a neonatal Fc receptor, insulin receptor, CD98, and Basigin.

Embodiment 64. The multidomain therapeutic protein of any one of embodiments 50-63, comprising a structure as depicted in Figure 1C, Figure 1D, Figure 1E, or Figure 1F.

Embodiment 65. Use of a nucleotide that encodes the multidomain therapeutic protein of any one of embodiments 21-39 and 50-64, the polynucleotide of any one of embodiments 40-45, or the gene therapy vector of any one of embodiments 46-49 in the method of any one of embodiments 1-20.

[0213] The following examples are provided to further illustrate the methods of the present invention. These examples are illustrative only and are not intended to limit the scope of the invention in any way.

EXAMPLES

Example 1: Construction of Anti-hCD63 ScFv::GAA Polynucleotide and Gene Therapy Vector

[0214] AAV2/8 viruses encoding for the expression of human GAA (hGAA; SEQ ID NO: 1; nucleic acid sequence represented by SEQ ID NO:12) or an anti-human CD63 single chain variable fragment (ScFv) fused on its C-terminus to human GAA (anti-hCD63 ScFv-hGAA; SEQ ID NO: 10; nucleic acid represented by SEQ ID NO:11) were generated using a standard triple transfection protocol (Gray et al. 2011; see also “Production of recombinant adeno-associated viral vectors and use *in vitro* and *in vivo* administration”, Current Protocols in Neuroscience, John Wiley & Sons, New York (1999), pp. 4.17.1–4.17.25, Vol 1). For the production, 1×10^7 HEK293 cells were plated onto 15 cm plates. The following day the cells

were transfected with (A) either 8 µg of a control pAAV vector comprising a liver specific serpin1 enhancer (SEQ ID NO:9) and encoding TTR driven human GAA or test pAAV comprising a liver specific serpin1 enhancer (SEQ ID NO:9) and encoding a TTR driven hCD63 ScFv-hGAA (see **Figure 1B**) and (B) pAAV RC2/8-derived vector (Gao, 2002) and 16 µg of pHelper (Agilent, Cat #240074) using PEIpro (Polyplus transfection, New York, NY catalog# 115-100)-mediated transfection at ratio of 1:1 (1ul PEIpro : 1µg DNA). Seventy-two hours after transfection, the cells were collected and lysed in a buffer comprised of 20mM Tris-HCl, 1mM MgCl₂, 2.5mM KCl, 100mM NaCl using a standard freeze-thaw method. Next, benzonase (Sigma, Cat# E1014-25KU) was added to the samples at a final concentration of 0.5 U/µL, and this was then incubated at 37°C for 60 minutes. Viruses were then purified using iodixanol gradient ultracentrifugation as described in (Zolotukhin et al., 1999, *Gene Ther* 1999;6:973–985) and were subsequently titrated by qPCR.

[0215] AAV samples were treated with DNaseI (ThermoFisher Scientific, Cat #EN0525) at 37°C for one hour and lysed using DNA extract All Reagents (ThermoFisher Scientific Cat# 4403319). Encapsidated viral genomes were quantified using an QuantStudio 3 Real-Time PCR System (ThermoFisher Scientific) using primers directed to the AAV2 ITRs. The sequences of the AAV2 ITRs primers are 5'-GGAACCCCTAGTGATGGAGTT-3' (fwd ITR; SEQ ID NO:3) and 5'-CGGCCTCAGTGAGCGA-3' (rev ITR; SEQ ID NO:4) (Aurnhammer *et al.*, 2012), derived the left internal inverted repeat (ITR) sequence from of the AAV (SEQ ID NO:6) and the right internal inverted repeat (ITR) sequence from of the AAV (SEQ ID NO:7), respectively. The sequence of the AAV2 ITRs probe is 5'-6-FAM-CACTCCCTCTCTGCGCGCTCG-TAMRA-3' (SEQ ID NO:5) (Aurnhammer C., Haase M., Muether N., et al., 2012, *Hum. Gene Ther. Methods* 23, 18–28). After a 95°C activation step for 10 min, a two-step PCR cycle was performed at 95°C for 15 seconds and 60°C for 30 seconds for 40 cycles. The TaqMan Universal PCR Master Mix (ThermoFisher Scientific, Cat #4304437) was used in the qPCR. DNA plasmid (Agilent, Cat #240074) was used as standard to determine absolute titers.

[0216] Anti-human CD63 antibodies and their fusions used the H5C6 mouse anti-human CD63 variable domains (amino acids 1-119 of SEQ ID NO:10 provide the amino acid sequence of the heavy chain variable domain (V_H) of the H5C6 antibody and amino acids 135-245 of SEQ ID NO:10 provide the amino acid sequence of the light chain variable domain (V_L) of the H5C6 antibody). The anti-hCD63 ScFv used here (SEQ ID NO:2) was derived from the H5C6 clone, which is mouse-anti-hCD63 monoclonal IgG1, kappa light chain antibody (H5C6 was deposited

to the Developmental Studies Hybridoma Bank at the University of Iowa by August, J.T. / Hildreth, J.E.K. (DSHB Hybridoma Product H5C6; DSHB Cat# h5c6, RRID:AB 528158). ScFv versions of the antibodies were cloned with variable domains in heavy-light order with a glycine-serine linker in between (5'-VH-Gly-Ser-VL-3')

Example 2: Glycogen content in Murine Pompe Model post-AAV

[0217] To determine the effect of AAV delivered anti-hCD63 ScFv-GAA fusion versus AAV delivered GAA, in a relevant glycogen storage *in vivo* model, both therapies were delivered to a Pompe disease mouse model where mice were homozygous for the deletion of the mouse GAA gene and were homozygous for the expression of human CD63 in place of mouse CD63 with a strain background of 75% C57BL/6; 25% 129SvJ. These mice are herein referred to as CD63 HumIn GAA KO mice or alternatively as CD63hu/hu; GAA^{-/-} mice.

[0218] For the experiment, 2-month-old CD63 HumIn GAA KO mice were administered via tail vein injection with either AAV2/8 virus containing a genome with either the TTR liver specific promoter driving human GAA (AAV-hGAA; described in Example 1) or the TTR liver specific promoter driving anti-human CD63 ScFv fused at its C-terminus with human GAA (AAV-anti-hCD63 ScFv-hGAA; described in Example 1). Both AAV2/8 viruses were delivered at either one of two doses, 1e10 vg/mouse or 1e11 vg/mouse. As controls, untreated CD63 HumIn GAA KO mice and untreated CD63 HumIn with the mouse GAA gene intact were included in the assay. Mice were housed for 3 months after treatment and bled incrementally (monthly) during this period for serum measurements of GAA levels and anti-GAA antibodies. After 3 months, all mice were sacrificed and individual tissues were harvested for glycogen measurements, PAS-H staining, quantification of central nuclei, measurement of lysosomal proliferation, and measurement of LC3b expression. Experimental dosing and treatment protocol for groups of mice are shown in Table 3.

Table 3: Experimental dosing and treatment protocol for groups of mice

Group	Mice	Number of Mice	Treatment	Dosage
1	CD63 HumIn GAA KO	4	None	N/A
2	CD63 HumIn GAA KO	4	AAV-hGAA	1e10 vg/mouse
3	CD63 HumIn GAA KO	4	AAV-hGAA	1e11 vg/mouse
4	CD63 HumIn GAA KO	5	AAV-anti-hCD63 ScFv-hGAA	1e10 vg/mouse

Group	Mice	Number of Mice	Treatment	Dosage
5	CD63 HumIn GAA KO	4	AAV-anti-hCD63 ScFv-hGAA	1e11 vg/mouse
6	CD63 HumIn GAA WT	2	None	N/A

[0219] The results are also depicted in Figure 2, which shows that anti-hCD63scFv::GAA brings glycogen down to wildtype levels in skeletal muscle, unlike GAA alone. Treatment with the two-domain anti-hCD63scFv::GAA multidomain therapeutic protein resulted in much greater reduction in stored glycogen compared to the single-domain GAA replacement enzyme. By plotting quadriceps glycogen levels (Figure 3) or heart glycogen levels (Figure 4) against the total serum expression of GAA or scfv-GAA over three months for individual mice, it was observed that the anti-hCD63scFv::GAA fusion protein removes more glycogen than the GAA enzyme alone, even at similar serum levels (Figures 3 and 4).

Example 3: Immunological Response to GAA

[0220] To measure anti-human GAA antibody serum levels, serum from all the treatment groups was separated from the blood collected during the terminal bleed using serum separator tubes (BD Biosciences, Cat#365967) as per the manufacturer's specifications. Separately, 96-well high protein binding plates (ThermoFisher, Cat#15041) were coated with 20µg of hGAA (R&D Systems, Cat#8329-GH-025) diluted in PBS overnight. Plates were washed with PBS + 0.05% Tween (PBS-T) 3 times. Plates were blocked with 0.5% BSA in PBS-T, and serial dilutions of mouse serum ranging from 1:300 to 1:5.1e7 were added to the plate overnight. Total anti-mouse IgG (subclasses 1 + 2a + 2b + 3) was measured using a HRP conjugated goat anti-mouse IgG antibody (Jackson Immuno Research, Cat# 115-035-164) and the BD Opt EIA substrate kit. The colorimetric reactions were stopped using 1 N phosphoric acid. Absorbance was then read at 450nm on a Spectramax i3 plate reader (Molecular Devices). Dilution curves were fit to sigmoidal curves, and titers were calculated from the curves. The titers expressed as mean total IgG titer +/- SD are shown in Table 4.

[0221] As shown in Table 4, mice that did not receive treatment showed an average background titer with mean levels of 1.1E+03. Mice treated with the low dose of virus (1e10 vg/mouse) of either AAV-anti-hCD63 ScFv-hGAA or AAV-hGAA demonstrated high titers, whereas in mice treated with the high dose (1e11 vg/mouse), titers were lower. Mice treated with 1e11 vg of AAV-anti-hCD63 ScFv-hGAA that had the highest levels of GAA in serum had titers within the range of untreated mice.

Table 4: Serum anti-GAA antibody levels

	Total IgG anti-GAA titer				
	CD63 HumIn GAA KO + no treatment	CD63 HumIn GAA KO + AAV-hGAA (1e10vg)	CD63 HumIn GAA KO + AAV-hGAA (1e11vg)	CD63 HumIn GAA KO + AAV-anti-hCD63 ScFv-hGAA (1e10vg)	CD63 HumIn GAA KO + AAV-anti-hCD63 ScFv-hGAA (1e11vg)
Mean	1.1E+03	7.6E+06	2.4E+04	5.5E+04	4.0E+03
SD	1.3E+03	1.0E+07	2.5E+04	1.9E+04	4.9E+03

[0222] Higher levels of GAA or anti-hCD63scFv::GAA after AAV administration correspond with lower anti-GAA titers. The serum of GAA null mice treated with high or low titers of AAV-anti-hCD63scFv::GAA or AAV-GAA were assessed for anti-GAA antibodies over the course of the three months post-injection. **Figure 5** depicts serum anti-GAA antibody titers vs GAA exposure (*i.e.*, the total serum expression over 3 months of GAA or scfv-GAA) for individual mice, which demonstrates a negative correlation between antibody titer and serum exposure to GAA, demonstrating that mice with high GAA exposure were tolerized to GAA. Likewise, **Figure 6**, which plots anti-GAA antibody titers for various groups infected with AAV encoding GAA or an anti-hCD63scFv::GAA protein, demonstrates that higher doses of construct led to lower titers of anti-GAAs.

Example 4: Serum GAA

[0223] To measure human GAA serum levels over the course of the experiment, samples were collected at monthly time points via tail bleed. Serum was separated from the blood using serum separator tubes (BD Biosciences, Cat# 365967) as per the manufacturer's specifications. 1μL of isolated serum was then loaded onto a 4-20% Novex wedgewell pre-cast gel, run at 220V for 45 minutes and transferred to nitrocellulose membrane at 200mA for 1 hour using standard procedures. The nitrocellulose membrane was then probed with an anti-GAA primary antibody (Abcam, #ab137068) used at a dilution of 1:2000 and an anti-GAPDH antibody (Abcam, #AB9484) used at a dilution of 1:1000 in 12mL and incubated overnight at 4 °C. After primary antibody incubation, the membrane was washed three times with 1 x TBST for 5 minutes per wash. Anti-rabbit IgG (LiCor, 926-32211) and anti-mouse IgG (LiCor, 925-68070) (LiCor, Lincoln, NE) secondary antibodies at a dilution of 1:15000 in 12mL were then added to the membrane and incubated for 1 hour at room temperature. After secondary antibody incubation,

the membrane was washed two times with 1 x TBST for 5 minutes per wash and one time with 1 x TBS for 5 minutes. The membrane was then imaged and quantified using a LiCor Odyssey instrument (LI-COR Biotechnology). Serum levels of GAA expressed as mean +/- standard deviation (SD) in arbitrary units are shown in Table 5.

[0224] As shown in Table 5, CD63 HumIn GAA KO mice treated with the high dose (10^{11} vg/mouse) of AAV-anti-hCD63 ScFv-hGAA or AAV-hGAA tested demonstrated sustained levels of GAA in the serum over the course of the experiment, with serum levels of GAA somewhat higher in AAV-anti-hCD63 ScFv-hGAA treated mice than in the AAV-hGAA treated mice. In mice treated with the treated with the low dose (10^{10} vg/mouse) of either AAV-anti-hCD63 ScFv-hGAA or AAV-hGAA, the levels of GAA dropped over the course of the experiment, approaching negligible levels in some mice by the 12 week time point.

Table 5: Serum GAA levels

	AAV-anti-hCD63 ScFv-hGAA (10^{10} vg)		AAV-hGAA (10^{10} vg)		AAV-anti-hCD63 ScFv-hGAA (10^{11} vg)		AAV-hGAA (10^{11} vg)	
Week	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	0.21	0.17	0.04	0.03	2.36	1.78	0.77	0.53
2	0.19	0.16	0.07	0.07	2.02	1.18	1.15	0.56
4	0.17	0.15	0.01	0.02	2.80	1.16	1.22	0.75
8	0.29	0.33	0.03	0.05	2.58	1.18	0.62	0.60
12	0.12	0.19	0.00	0.01	2.61	1.53	0.77	0.86
Area under the curve	2.27	2.23	0.27	0.36	28.13	13.73	9.80	7.22

[0225] Expression of GAA or anti-hCD63scfv::GAA was maintained over time in mice receiving the high dose of AAV (10^{11} vg/mouse), but fell off in mice receiving the lower dose (10^{10} vg/mouse). Figure 7A depicts a graph plotting serum levels of GAA, as probed by western blot, over time for various groups infected with AAV encoding GAA or an anti-hCD63 scfv fusion to GAA. The fusion protein (scFv::GAA) demonstrated consistently higher levels (e.g., 2.5 to 3-fold) of serum GAA than the GAA enzyme without the delivery domain (Figure 7A).

[0226] Real-time PCR quantifications of expression in liver, heart, and quadriceps lysates 3 months after injection are shown in Figure 7B. Liver expression was detected for all injections of AAV construct, with highest levels for the 1×10^{11} vg/mouse injections for both AAV-hGAA

and AAV-anti-hCD63::hGAA (both driven by liver-specific promoter, LSP). A comparison of serum GAA level to RNA expression level of GAA was also made (Figure 7C) and the results show that mice receiving the AAV encoding the fusion protein presented lower GAA RNA expression localized to the liver at 3 months however GAA serum levels were high in that particular mouse. AAV-LSP-hGAA injections did not present high serum levels of GAA when RNA levels were low in the liver. See Figure 7C. This data suggests that the AAV encoding the fusion protein (and expression is driven by a liver-specific promoter) attains an improved secretion profile for GAA.

[0227] A higher secreted to intracellular ratio of antibody::hGAA versus hGAA alone in Huh-7 hepatocytes was also observed. In one experiment, Huh-7 human hepatocytes were transiently transfected with liver-specific promoter driven constructs encoding for hGAA, anti-hCD63 scFv::GAA fusion, or a non-binding scFv::GAA fusion control. Both scFv::GAA fusion constructs had a higher ratio of protein in the secreted supernatant than hGAA alone 3 days after transfection (statistically significant to $p < 0.05$, $n=3$). Addition of M6P into the supernatant during the experimental period to mitigate CI-MPR-mediated uptake did not affect the ratio.

Example 5: Tissue Measurement of Glycogen and Histological Characterization of Muscle Tissue

[0228] **Tissue measurements of glycogen:** To measure the glycogen content in individual tissues, heart, quadriceps, gastrocnemius, diaphragm, soleus, and EDL tissue were dissected from mice from all groups immediately after CO₂ asphyxiation, and were then snap frozen in liquid nitrogen, and stored at -80°C. ~50mg of each tissue was lysed on a benchtop homogenizer with stainless steel beads in distilled water at a ratio of 1mg to 25μL water for glycogen measurements. Glycogen analysis lysates were heated at 105 ° for 15 minutes and centrifuged at 21000 x g to clear debris. Glycogen measurements were performed using a Glycogen Assay Kit (Sigma-Aldrich, #MAK016) according to manufacturer's instructions for fluorometric assays. The fluorescence of each sample was measured at 535nm excitation and 587nm emission on a fluorescence plate reader (Molecular Devices, Spectramax i3). The calculated amount of glycogen was calculated using the following formula provided by the manufacturer. The calculated amount of glycogen from each tissue in each treatment group was then averaged and is expressed as mean +/- standard deviation (SD) in Table 6.

[0229] As shown in Table 6, loss of *Gaa* causes a large increase in mean glycogen levels across all tissues measured, as compared to GAA WT mice. Treatment with AAV-anti-hCD63 ScFv-hGAA at 10¹¹ vg/mouse reduced glycogen to WT- or near-WT levels in all tissues tested, unlike

treatment with AAV-GAA which only partially reduced stored glycogen. The low doses of either virus also reduced glycogen, but to a lesser extent than the high doses. The 10^{10} vg/mouse dose of AAV-anti-hCD63 ScFv-hGAA reduced glycogen levels in a similar manner as the 10^{11} vg/mouse dose of AAV-GAA.

Table 6: Mean +/- SD glycogen level measured in heart, quadriceps, gastrocnemius, diaphragm, soleus, and EDL

	no treatment		AAV-hGAA (10^{10} vg)		AAV-hGAA (10^{11} vg)	
	Mean	SD	Mean	SD	Mean	SD
Heart	27.798	3.013	17.246	4.375	1.770	2.279
Quadricep	14.650	1.783	11.012	0.528	5.878	3.504
Gastrocnemius	14.295	0.480	10.990	0.868	6.073	3.080
Diaphragm	15.463	1.173	11.446	1.237	3.995	3.395
Soleus	17.260	2.262	13.684	2.506	6.533	5.201
EDL	13.588	0.498	11.178	1.760	6.275	3.159
	AAV-anti-hCD63 ScFv-hGAA (10^{10} vg)		AAV-anti- hCD63 ScFv- hGAA (10^{11} vg)		CD63 HumIn GAA WT mice (control)	
	Mean	SD	Mean	SD	Mean	SD
Heart	2.190	2.678	0.058	0.010	0.085	0.007
Quadricep	4.485	3.147	0.798	0.251	0.440	0.042
Gastrocnemius	5.198	2.516	0.825	0.461	0.790	0.014
Diaphragm	3.083	2.968	0.388	0.121	0.385	0.007
Soleus	5.268	2.786	1.040	0.896	0.545	0.049
EDL	2.495	1.750	0.313	0.099	0.260	0.141

[0230] Quadricep harvest for histopathology and quantification: Quadricep tissue samples from mice from each group besides the low dose (1×10^9 vg/mouse) treatment group were either snap frozen immediately after dissection in liquid nitrogen and stored at -80°C for quantification of LC3b expression or were placed onto blocks containing O.C.T medium (Tissue-Tek, #4583).

[0231] Tissues samples in O.C.T medium were sent to Histoserv, Inc. (Germantown, MD) for sectioning and periodic acid Schiff (PAS) staining to detect polysaccharides. Additional sections were prepared and returned for staining of central nuclei and lysosomal proliferation.

[0232] PAS staining: PAS stain sections were imaged using a Leica slide scanner at 20x magnification. The resulting images from representative mice for each treatment group are shown in Figure 8.

[0233] As shown in Figure 8, CD63 HumIn GAA KO that were treated with AAV-anti-hCD63 ScFv-hGAA at 3 months demonstrated a marked decrease in staining as compared to both the CD63 HumIn GAA KO mice with no treatment and the CD63 HumIn GAA KO mice treated with AAV-hGAA, which displayed high levels of PAS staining. This further indicates that the treatment with AAV-anti-hCD63 ScFv-hGAA can reduce polysaccharides accumulation in CD63 HumIn GAA KO mice and can do so in a uniform manner across muscle fibers.

[0234] **Quantification of central nuclei and lysosomal proliferation:** Unstained sections from Histoserv were removed from the freezer and then fixed with 4% paraformaldehyde in PBS for 15 minutes in a staining chamber. The fixed slides were then washed twice for 5 minutes in PBS and subsequently incubated with blocking buffer (eBiosciences, 00-4953-54) for 1 hour at room temperature. Slides were then either stained with either a rat anti-Lamp-1 antibody (Abcam, #AB25245) at a dilution of 1:50 in blocking buffer, a rabbit anti-Laminin antibody (Sigma, #L9393) at a dilution of 1:1000 in blocking buffer or blocking buffer with no added antibody while in a humidified staining chamber and then transferred to 4°C for overnight incubation. The following day, slides were then washed twice for 5 minutes in PBS and subsequently stained with either goat anti-rabbit IgG (H+L) superclonal secondary antibody conjugated with Alexa Fluor 647 (Life Tech Thermo, #A27040) or goat anti-rat IgG (H+L) cross-adsorbed secondary antibody conjugated with Alexa Fluor 555 (Life Tech Thermo, #A21434) in a staining chamber then allowed to incubate for 1 hour at room temperature. Stained slides were then washed twice for 5 minutes in PBS before they were mounted with Fluoromount-G with DAPI (Life Tech Thermo, #00-4959-52) and imaged on a Zeiss LSM710 instrument (Carl Zeiss Microscopy GmbH). Number of centralized nuclei was quantified using Halo software (Indica Labs, NM) and is expressed as percentage of fibers showing central nuclei +/- standard deviation are shown in Table 7. Lysosomal proliferation is depicted in Figure 8.

Table 7: Quantification of central nuclei

	no treatment		AAV-hGAA (1e11vg)		AAV-anti- hCD63 ScFv-hGAA (1e11vg)		CD63 HumIn GAA WT mice (control)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
% of Fiber with central nuclei	22.00	10.10	33.00	4.36	12.75	6.29	8.50	7.78

[0235] Quantification of LC3b expression: For quantification of LC3b expression, snap frozen samples were thawed, homogenized and then lysed in RIPA buffer at a 1mg tissue to 25 μ L RIPA buffer ratio (150 mM NaCl, 1.0% IGEPAL[®] CA-630, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0, Sigma Aldrich, R0278) by bead impaction for 45 seconds (MP Biomedical). Lysates were cleared of insoluble material by centrifugation at 21,000 x g and then 300 μ g of lysate in RIPA buffer was loaded on a 4-20% Novex wedgewell pre-cast gel, transferred to a nitrocellulose membrane and analyzed by western blot using a similar protocol as previously described for the analysis of serum GAA levels, substituting the use of primary antibody that recognizes mouse LC3b-I and LC3b-II (Sigma, #L7543) in place of the primary antibody against GAA. The membrane was then imaged and quantified using a LiCor Odyssey instrument (LI-COR Biotechnology). LC3b-I and LC3b-II levels expressed as (mean +/- standard deviation) in arbitrary units are shown in Table 8.

[0236] As shown in Table 8, there was a significant increase in both mean LC3b-I and LC3b-II levels in mice lacking GAA as compared to CD63 HumIn GAA WT mice. Treatment with AAV-anti-hCD63 ScFv-hGAA decreased mean LC3b-I and LC3b-II levels in CD63 HumIn GAA KO to the WT- or near-WT levels. CD63 HumIn GAA KO treated with AAV-hGAA demonstrated slightly decreased mean LC3b-I and LC3b-II levels as compared to CD63 HumIn GAA KO mice, but this decrease was not as pronounced as with AAV-anti-hCD63 ScFv-hGAA treatment.

Table 8: LC3b-I and LC3b-II levels in the quadriceps of mice

LC3b-I levels (arbitrary units)				
treatment	untreated	AAV-hGAA	AAV-anti-hCD63 ScFv-hGAA	CD63 HumIn GAA WT (control)
average	833	628	403	282
SD	109	139	33	49
LC3b-II levels (arbitrary units)				
treatment	untreated	AAV-hGAA	AAV-anti-hCD63 ScFv-hGAA	CD63 HumIn GAA WT (control)
average	3308	2888	445	369
SD	582	1282	398	33

Example 6: AAV anti-hCD63::GAA treatment leads to significant gains in tests of muscle strength and coordination

[0237] Grip strength and Rotarod test performance of mice treated (see above) with either AAV-LSP hGAA or AAV-LSP anti-hCD63::hGAA. Accelerating Rotarod measurements (Figure 9A) and forelimb grip strength measurements (Figure 9B) of wild-type GAA mice, untreated control, AAV-LSP-hGAA (1e11vg/mouse) or AAV-LSP-anti-hCD63:: hGAA treatment (1e11vg/mouse) were taken at monthly intervals for 6 months. Error bars are +/- SD. N=8-10 for all groups.

Example 7: Other membrane proteins as “guides” directing GAA to tissues

[0238] Other membrane proteins were tested, such as anti-ITGA7 (Integrin alpha-7) fusion proteins, to guide GAA to tissues to replace GAA in enzyme-deficient mice. C2C12 mouse myoblasts were incubated overnight with anti-mCD63-GAA or anti-ITGA7-GAA with or without the presence of 5mM M6P. Active GAA enzyme was detected in myoblast lysates over time for both fusion proteins (Figure 10A). In further experiments, GAA KO mice humanized for CD63 (GAA^{-/-};CD63hu/hu) were given plasmids encoding an scFv::GAA format of anti-hCD63::GAA or a full-length IgG4::GAA format of anti-integrin alpha-7 by hydrodynamic delivery (HDD), and mice were sacrificed 3 weeks post-HDD. Tissue glycogen levels were measured in heart, quadriceps, gastrocnemius and diaphragm. Untreated control mice, GAA^{-/-} xCD63hu/hu and untreated wild-type GAA control mice, GAA^{+/+};CD63hu/hu (4) were also tested under the same conditions. Glycogen levels were at very low levels in both anti-

hCD63::GAA treated mice and anti-ITGA7::GAA treated mice groups, as in the wild-type mice. **See Figure 10B.**

Example 8: At comparable serum levels, AAV anti-CD63::GAA treatment is more effective than AAV with optimized GAA construct

[0239] CD63 HumIn GAA KO mice ($GAA^{-/-}$ x $CD63^{hu/hu}$) infected with AAVs containing a liver specific enhancer (serpina 1; SEQ ID NO:9) and a liver specific promoter (LSP; TTR; SEQ ID NO:8) driving the expression of an anti-hCD63::GAA multidomain therapeutic (SEQ ID NO:10), which uses a chymotrypsinogen B2 signal peptide (SP7) and contains amino acids 36-952 of human GAA ($\Delta 8GAA$) exhibited significant gains in tests of muscle strength and coordination. Three different doses were given for each virus: 5×10^{11} vg/kg, 2×10^{12} vg/kg, and 4×10^{12} vg/kg. Serum was collected by submandibular bleeds on a regular basis. One month post-AAV infection, mice were sacrificed. Cardiac and skeletal muscle tissue samples were collected and snap frozen in liquid nitrogen and kept at $-80^{\circ}C$ for storage. Glycogen in tissues were measured by homogenizing tissues by bead impaction in distilled water. Samples were boiled and centrifuged, and the supernatants were used in a commercial glycogen assay kit. Serum was quantified using western blot with an antibody against human GAA as described in previous examples. For each mouse, the glycogen level in each tissue was plotted against the serum level of the construct at 1 month. 4-parameter curve fits were used to determine the EC50 of the two treatments in each tissue.

[0240] Infection with AAVs containing a liver specific promoter (LSP) encoding either anti-hCD63::GAA or sp7- $\Delta 8GAA$ provided comparable serum levels of GAA at each infection dose. **Figure 11.** However, in every muscle tissue assayed, an ~ 2.2 fold reduction in EC50 was observed when using anti-hCD63::GAA vs. sp7- $\Delta 8GAA$, demonstrating that at equivalent serum levels, anti-CD63::GAA clears glycogen more efficiently than a modified GAA expression construct that is not fused to an antibody. **See Figure 12.**

Example 9: Glycogen content in CNS of Murine Pompe Model post-AAV Treatment with Various GAA Constructs and Doses

[0241] Two-month-old CD63 HumIn GAA KO mice were administered via tail vein injection with either AAV2/8 virus containing a genome with either the TTR liver specific promoter driving human GAA (AAV-hGAA; described in Example 1) or the TTR liver specific promoter driving anti-human CD63 ScFv fused at its C-terminus with human GAA (AAV-anti-hCD63 ScFv-hGAA; described in Example 1). Both AAV2/8 viruses were delivered at 1×10^{11} vg/mouse.

As controls, untreated CD63 HumIn GAA KO (*Gaa*^{-/-}) mice and untreated CD63 HumIn with the mouse GAA gene intact (wildtype mice) were included in the assay. Mice were housed for 9 months after treatment, after which all mice were sacrificed and individual tissues were harvested for glycogen measurements. CNS tissues) were dissected on ice and quickly flash frozen 9 months after AAV transduction. Spinal cord, cerebellum, and hippocampus tissues were homogenized in distilled deionized water using bead impaction, and glycogen was measured in the tissue lysate supernatants using a commercial fluorometric glycogen assay kit (**FIGURE 13**).

[0242] In another analogous experiment, knockout mice were treated with AAV constructs encoding GAA (1e11 vg) or anti-CD63ScFv-GAA fusion (doses 1e10vg, 5e10 vg, or 1e11 vg). Wild-type mice and the amount of stored glycogen per mg of CNS tissue (spinal cord: **Figure 14A**; brain: **Figure 14B**) was examined 3 months after AAV delivery. Wild-type mice and KO mice (*GAA*^{-/-}) that were untreated were used as comparators for their stored glycogen levels in CNS tissue for the same duration (3 months).

[0243] ScFv-GAA fusion constructs were more effective than GAA alone at reducing stored glycogen levels in diseased mice at a dose of 1e11 vg/mouse (**Figures 13-14**). A dose of 5e10 vg ScFv-GAA fusion provided decreased glycogen storage levels equivalent to the higher dosed GAA alone construct (**Figures 14A-14B**). These levels of reduced glycogen storage were shown to be effective by Hordeaux et al 2017 (e.g. a decrease in glycogen and improved muscle strength in mice injected intrathecally with AAV GAA). The correlative serum level of ScFv-GAA fusion protein is detectable at 5e10 vg and more than 15 ug/mL in the mice receiving 1e11 vg (Table 9).

Table 9: Correlation of dose by weight and level of anti-CD63::GAA in the serum.

Experimental dose (viral genomes/mouse)	Approximate dose by weight (viral genomes/kg)	Level of anti-CD63::GAA in the serum (micrograms / mL)
1e10	4e11	undetectable
5e10	2e12	1.97 +/- 1.18
1e11	4e12	15.84 +/- 13.37

[0244] Without being bound by any one theory, detectable serum levels, e.g., serum levels higher than 1 ug/mL of GAA linked to a delivery domain that crosses the blood-brain-barrier are considered therapeutic.

Example 10: Expression of multidomain therapeutic proteins comprising at least two delivery domains

[0245] CHO cells were transfected with an expression constructs depicted in Figures 1C-1G, and expression of each of the constructs was confirmed (data not shown). Additionally, binding of some of the multidomain therapeutic proteins encoded by 4W1 and 4M1 (Figure 1F) to CD63 was confirmed by ELISA (data not shown).

REPLACEMENT CLAIMS

1. A method of delivering a therapeutic protein to the central nervous system (CNS) of a patient, comprising delivering a nucleotide composition encoding a multidomain therapeutic protein to the liver in the patient to form a liver depot for the production and secretion of the multidomain therapeutic at a consistent serum level of at least 1 µg/mL over consecutive days, weeks or months following delivery to provide a therapeutically effective amount of the multidomain therapeutic protein in the CNS,

wherein the multidomain therapeutic protein comprises an antibody or antigen-binding portion thereof that binds CD63 or ITGA7 and an enzyme domain.

2. The method of claim 1, wherein the therapeutic protein is a lysosomal enzyme.

3. The method of claim 2, wherein the lysosomal enzyme is GAA.

4. The method of any one of claims 1-3, wherein the nucleotide composition is administered via a viral vector.

5. The method of claim 4, wherein the viral vector is an AAV vector, optionally wherein the nucleotide composition is administered at a dose of at least 2×10^{12} viral genomes per kilogram (vg/kg).

6. The method of any one of claims 1-5, wherein the antibody or antigen-binding protein thereof comprises a single-chain variable fragment (scFv).

7. The method of any one of claims 1-6, wherein the antibody or antigen-binding protein thereof comprises an amino acid sequence of SEQ ID NO:2.

8. The method of any one of claims 1-7, wherein the therapeutic protein comprises a hydrolase.

9. The method of any one of claims 1-8, wherein the therapeutic protein comprises a glycosylase.

10. The method of any one of claims 1-9, wherein the therapeutic protein comprises a glycosidase.

11. The method of any one of claims 1-10, wherein the therapeutic protein comprises an alpha-glucosidase.

12. The method of any one of claims 1-11, wherein the therapeutic protein comprises an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:13, or a fragment thereof.
13. The method of any one of claims 1-12, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO:11.
14. The method of any one of claims 1-13, wherein the enzyme domain comprises an alpha-glucosidase, and wherein the glycogen levels in any CNS tissue in the subject is reduced for at least nine months post-treatment.
15. The method of any one of claims 1-14, wherein the subject has Pompe disease.
16. The method of any one of claims 1-15, wherein the serum level is at least 2 µg/mL.
17. The method of any one of claims 1-16, wherein the antibody or antigen-binding portion thereof binds an extracellular domain of CD63 or ITGA7.
18. A multidomain therapeutic protein comprising one or more delivery domain(s) and an enzyme domain, wherein one of the one or more delivery domain(s) binds human transferrin receptor (hTfR), and wherein an other of the one or more delivery domain(s) comprises an antibody or antigen-binding protein thereof that binds CD63 or ITGA7.
19. The multidomain therapeutic protein of claim 18, wherein each of the one or more delivery domain(s) comprises an antigen-binding protein.
20. The multidomain therapeutic protein of claim 18 or claim 19, wherein at least one of the one or more delivery domain(s) comprises a single-chain variable fragment (scFv).
21. The multidomain therapeutic protein of any one of claims 18-20, wherein at least one of the one or more delivery domain(s) comprises a half-body.
22. The multidomain therapeutic protein of claim 21, wherein the delivery domain that binds hTfR is an scFv, wherein the half-body binds CD63, wherein the enzyme domain is GAA, and wherein GAA is conjugated to the carboxy terminus of the half-body that binds CD63.
23. The multidomain therapeutic protein of claim 22, wherein each of the one or more delivery domain(s) comprises an scFv.
24. The multidomain therapeutic protein of any one of claims 20-23, wherein at least one scFv is fused to an Fc.

25. The multidomain therapeutic protein of claim 24, wherein the Fc comprises a wildtype human IgG4 isotype, or derivative thereof.
26. The multidomain therapeutic protein of any one of claims 24-25, wherein GAA is conjugated to the carboxy terminus of the Fc.
27. The multidomain therapeutic protein of claim 23, comprising an anti-hTfR scFv, an anti-hCD63 scFv.
28. The multidomain therapeutic protein of claim 27, wherein the anti-hTfR scFv and anti-hCD63 scFv are both linked, at their carboxy termini, to a single GAA enzyme.
29. The multidomain therapeutic protein of any one of claims 18-28, wherein the delivery domain is an anti-hTfR scFv, and the enzyme domain is linked to carboxy terminus of the VL domain of the scFv.
30. The multidomain therapeutic protein of claim 29, comprising the other delivery domain linked to the N-terminus of the VH domain of the anti-hTfR scFv.
31. The multidomain therapeutic protein of claim 30, wherein the second delivery domain is an anti-hCD63 scFv.
32. The multidomain therapeutic protein of any one claims 18-31, wherein the enzyme domain comprises the amino acid sequence set forth as SEQ ID NO:1
33. A multidomain therapeutic protein comprising at least two delivery domains and at least one enzyme domain, wherein each of the two delivery domains is independently selected from the group consisting of an antibody, a half-body, and an scFv, and wherein at least one or more of the delivery domains is associated the at least one enzyme domain, preferably wherein the one or more delivery domains is covalently linked to the at least one enzyme domain.
34. The multidomain therapeutic protein of claim 33, comprising a structure as depicted in Figure 1C, Figure 1D, Figure 1E, or Figure 1F.
35. A polynucleotide encoding the multidomain therapeutic protein of any one of claims 18-34.
36. The polynucleotide of claim 35, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence.

37. The polynucleotide of claim 35 or claim 36, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence.

38. The polynucleotide of any one of claims 35-37, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence, and wherein the AAV nucleic acid sequence comprises an internal terminal repeat sequence, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter.

39. The polynucleotide of any one of claims 35-38, further comprising a virus nucleic acid sequence and a locus-targeting nucleic acid sequence, wherein the virus nucleic acid sequence is an adeno-associated virus (AAV) nucleic acid sequence comprising an internal terminal repeat sequence that comprises SEQ ID NO:6, SEQ ID NO:7, or both, and optionally, a tissue specific regulatory element such as a liver specific promoter or a neuronal specific promoter.

40. The polynucleotide of any one of claims 35-39, further comprising a tissue specific regulatory element comprising the sequence set forth as SEQ ID NO:8 and/or SEQ ID NO:9.

41. A gene therapy vector comprising a polynucleotide of any one of claims 35-40.

42. The gene therapy vector of claim 41, wherein the gene therapy vector is selected from the group consisting of

a viral vector, optionally wherein the viral vector is a natural virus, an engineered virus, or a chimeric virus,

a naked polynucleotide comprising the polynucleotide of any one of claims 35-40,

a polynucleotide complex, optionally wherein the polynucleotide complex is a lipid nanoparticle comprising the polynucleotide of any one of claims 34-39 and lipids, and

any combination thereof.

43. The gene therapy vector of claim 41 or claim 42, wherein the gene therapy vector is a viral vector selected from the group consisting of a retrovirus, adenovirus, herpes simplex virus, pox virus, vaccinia virus, lentivirus, or an adeno-associated virus.

44. The gene therapy vector of claim 42 or claim 43, wherein the gene therapy vector is AAV9, Anc80, an AAV2/8 chimera and/or an AAV pseudotyped to a specific tissue, e.g., the liver or neuronal tissue.

45. Use of a nucleotide that encodes the multidomain therapeutic protein of any one of claims 18-34, the polynucleotide of any one of claims 35-40, or the gene therapy vector of any one of claims 41-44 in the method of any one of claims 1-17.

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 1D




FIGURE 1E




FIGURE 1F






FIGURE 1G




8/28

FIGURE 2




9/28

FIGURE 3




10/28

FIGURE 4




11/28

FIGURE 5




12/28

FIGURE 6




13/28

FIGURE 7A




14/28

FIGURE 7B




15/28

FIGURE 7C




16/28

FIGURE 7D




17/28

FIGURE 7E




18/28

FIGURE 8



19/28


FIGURE 8 (Cont'd)




20/28

FIGURE 9

A




B




21/28

FIGURE 10A




22/28

FIGURE 10B



23/28

FIGURE 11



24/28

FIGURE 12A




FIGURE 12B






FIGURE 13



27/28

FIGURE 14A

3 months post AAV spinal cord



28/28

FIGURE 14B

