

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number
WO 02/22829 A2

(51) International Patent Classification⁷: **C12N 15/54**,
1/21, C12P 13/08

(21) International Application Number: PCT/EP01/09456

(22) International Filing Date: 16 August 2001 (16.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
100 45 497.6 13 September 2000 (13.09.2000) DE

(71) Applicant: **DEGUSSA AG** [DE/DE]; Bennigsenplatz 1,
40474 Düsseldorf (DE).

(72) Inventors: **MÖCKEL, Bettina**; Benrodestrasse 35,
40597 Düsseldorf (DE). **MARX, Achim**; Altenbrede 4B,
33613 Bielefeld (DE). **BASTUCK, Christine**; Sieker-
wall 17, 33602 Bielefeld (DE). **BUCHHOLZ, Michael**;
Wellensiek 30, 33619 Bielefeld (DE). **PFEFFERLE,**
Walter; Jahnstrasse 33, 33790 Halle (Westf.) (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: NUCLEOTIDE SEQUENCES WHICH CODE FOR THE PPSA GENE

(57) Abstract: The invention relates to an isolated polynucleotide comprising a polynucleotide sequence chosen from the group consisting of a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2, b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2, c) polynucleotide which is complementary to the polynucleotides of a) or b), and d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c), and a process for the fermentative preparation of L-amino acids using coryneform bacteria in which at least the ppsA gene is present in enhanced form, and the use of polynucleotides which comprise the sequences according to the invention as hybridization probes.



WO 02/22829 A2

Nucleotide sequences which code for the ppsA gene

Field of the Invention

The invention provides nucleotide sequences from coryneform bacteria which code for the ppsA gene and a process for the fermentative preparation of amino acids using bacteria in which the ppsA gene is enhanced.

Prior Art

L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and produce amino acids are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acid, by

amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

Object of the Invention

The inventors had the object of providing new measures for
5 improved fermentative preparation of amino acids.

Summary of the Invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-
10 asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Lysine is particularly preferred.

15 When L-lysine or lysine are mentioned in the following, not only the bases but also the salts, such as e.g. lysine monohydrochloride or lysine sulfate, are meant by this.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence
20 which codes for the ppsA gene, chosen from the group consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of
25 SEQ ID No. 2,
- b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,
- 30 c) polynucleotide which is complementary to the polynucleotides of a) or b), and

d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of phosphoenol pyruvate synthase.

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- 10 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii),
- 15 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides

a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as shown in SEQ ID No. 1;

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing the polynucleotide according to the invention, in particular a shuttle vector or plasmid vector, and

coryneform bacteria which contain the vector or in which the ppsA gene is enhanced.

The invention also provides polynucleotides, which substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which
5 comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide according to the invention according to SEQ ID No.1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

10 Detailed Description of the Invention

Polynucleotides which comprise the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for
15 phosphoenol pyruvate synthase or to isolate those nucleic acids or polynucleotides or genes which have a high similarity of sequence with that of the ppsA gene. They are also suitable for incorporation into so-called "arrays", "micro arrays" or "DNA chips" in order to detect and
20 determine the corresponding polynucleotides.

Polynucleotides which comprise the sequences according to the invention are furthermore suitable as primers with the aid of which DNA of genes which code for phosphoenol pyruvate synthase can be prepared by the polymerase chain
25 reaction (PCR).

Such oligonucleotides which serve as probes or primers comprise at least 25, 26, 27, 28, 29 or 30, preferably at least 20, 21, 22, 23 or 24, very particularly preferably at least 15, 16, 17, 18 or 19 successive nucleotides.
30 Oligonucleotides with a length of at least 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40, or at least 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50 nucleotides are also suitable. Oligonucleotides with a length of at least 100, 150, 200, 250 or 300 nucleotides are optionally also suitable.

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

5 The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70% to 80%, preferably at least 81% to 85%, particularly preferably at least 86% to 90%, and very particularly
10 preferably at least 91%, 93%, 95%, 97% or 99% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via
15 peptide bonds.

The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of phosphoenol pyruvate synthase, and also those which are at least 70% to 80%,
20 preferably at least 81% to 85%, particularly preferably at least 86% to 90%, and very particularly preferably at least 91%, 93%, 95%, 97% or 99% identical to the polypeptide according to SEQ ID No. 2 and have the activity mentioned.

The invention furthermore relates to a process for the
25 fermentative preparation of amino acids chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-
30 arginine using coryneform bacteria which in particular already produce amino acids and in which the nucleotide sequences which code for the ppsA gene are enhanced, in particular over-expressed.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes in a microorganism which are coded by the corresponding DNA, for example by increasing the number of
5 copies of the gene or genes, using a potent promoter or using a gene which codes for a corresponding enzyme having a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is
10 in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the activity or concentration of the protein in the starting microorganism.

15 The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus
20 *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in
25 particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

Corynebacterium glutamicum ATCC13032
Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
30 *Corynebacterium thermoaminogenes* FERM BP-1539
Corynebacterium melassecola ATCC17965
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869 and
Brevibacterium divaricatum ATCC14020

and L-amino acid-producing mutants or strains prepared therefrom.

The new ppsA gene from *C. glutamicum* which codes for the enzyme phosphoenol pyruvate synthase (EC 2.7.9.2) has been
5 isolated.

To isolate the ppsA gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an
10 example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ vectors by Kohara et al. (*Cell* 50, 495 -508 (1987)). Bathe et al. (*Molecular and General Genetics*, 252:255-265, 1996) describe a gene
15 library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, *Proceedings of the National Academy of Sciences USA*, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, *Nucleic Acids Research* 16:1563-1575).
20

Börmann et al. (*Molecular Microbiology* 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pHC79 (Hohn and Collins, *Gene* 11, 291-298 (1980)).
25

To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, *Life Sciences*, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, *Gene*, 19:259-268). Suitable hosts are, in particular, those
30 *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 α mc r , which has been described by Grant et al. (*Proceedings of*

the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by
5 Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as
10 e.g. that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of *C. glutamicum* which codes for the
15 ppsA gene and which, as SEQ ID No. 1, is a constituent of the present invention has been found. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequence of the
20 ppsA gene product is shown in SEQ ID No. 2.

Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a
25 constituent of the invention. Conservative amino acid exchanges, such as e.g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the
30 protein, i.e. are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of
35 Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene

77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a
5 corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Finally, DNA sequences which are prepared by the
10 polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of
15 hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260). The
20 hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i. e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of the hybridization, including the washing
25 steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington,
30 UK, 1996).

A 5x SSC buffer at a temperature of approx. 50°C - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with
polynucleotides which are less than 70% identical to the
35 sequence of the probe. Such hybrids are less stable and are

removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50°C - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50°C to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonucleotide Synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce amino acids in an improved manner after over-expression of the ppsA gene.

To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative amino acid production. The expression is likewise improved by measures to prolong the life of the mRNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes

or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be
5 achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)),
10 Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in EP 0 472 869, in US 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132
15 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in WO 96/15246, in Malumbres et al. (Gene 134, 15 - 24 (1993)), in JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)), in Makrides (Microbiological Reviews 60:512-538
20 (1996)) and in known textbooks of genetics and molecular biology.

For enhancement, by way of example the ppsA gene according to the invention was over-expressed with the aid of episomal plasmids. Suitable plasmids are those which are
25 replicated in coryneform bacteria. Numerous known plasmid vectors, such as e.g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554), pEKEx1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic
30 plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, such as e.g. those based on pCG4 (US-A 4,489,160), or pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)), or pAG1 (US-A 5,158,891), can be used in the same manner.

Plasmid vectors which are furthermore suitable are also those with the aid of which the process of gene amplification by integration into the chromosome can be used, as has been described, for example, by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for duplication or amplification of the *hom-thrB* operon. In this method, the complete gene is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US-A 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)), pEM1 (Schrumpf et al, 1991; Journal of Bacteriology 173:4510-4516) or pBGS8 (Spratt et al., 1986, Gene 41: 337-342). The plasmid vector which contains the gene to be amplified is then transferred into the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene in question.

In addition, it may be advantageous for the production of L-amino acids to enhance, in particular over-express one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle, of the pentose phosphate cycle, of amino acid export and

optionally regulatory proteins, in addition to the ppsA gene.

Thus, for the preparation of L-amino acids, in addition to enhancement of the ppsA gene, one or more genes chosen from
5 the group consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology
10 174:6076-6086),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pgk gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- 15 • the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661),
- the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
- the mqo gene which codes for malate-quinone
20 oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
- the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512),
- the lysE gene which codes for lysine export
25 (DE-A-195 48 222),
- the hom gene which codes for homoserine dehydrogenase (EP-A 0131171),
- the ilvA gene which codes for threonine dehydratase (Möckel et al., Journal of Bacteriology (1992) 8065-

8072)) or the *ilvA*(Fbr) allele which codes for a "feed back resistant" threonine dehydratase (Möckel et al., (1994) Molecular Microbiology 13: 833-842),

• the *ilvBN* gene which codes for acetohydroxy-acid synthase
5 (EP-B 0356739),

• the *ilvD* gene which codes for dihydroxy-acid dehydratase (Sahm and Eggeling (1999) Applied and Environmental Microbiology 65: 1973-1979),

• the *zwa1* gene which codes for the Zwa1 protein (DE:
10 19959328.0, DSM 13115),

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of L-amino acids, in addition to the enhancement of the *ppsA* gene, for one or more genes chosen from the group
15 consisting of

• the *pck* gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),

• the *pgi* gene which codes for glucose 6-phosphate isomerase (US 09/396,478; DSM 12969),

20 • the *poxB* gene which codes for pyruvate oxidase (DE: 1995 1975.7; DSM 13114),

• the *zwa2* gene which codes for the Zwa2 protein (DE: 19959327.2, DSM 13113)

to be attenuated, in particular for the expression thereof
25 to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak

promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

- 5 By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the
10 starting microorganism.

In addition to over-expression of the ppsA gene it may furthermore be advantageous for the production of amino acids to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in:
15 Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch
20 culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of amino acids. A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die
25 Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of
30 the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as e.g. antibiotics, can be added to the medium to maintain the

stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.

Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by ion exchange chromatography with subsequent ninhydrin derivation, or it can be carried out by reversed phase HPLC, for example as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The process according to the invention is used for fermentative preparation of amino acids.

The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA).

Methods for transformation of Escherichia coli are also described in this handbook.

The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

Example 1

Preparation of a genomic cosmid gene library from
Corynebacterium glutamicum ATCC 13032

- Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032
5 was isolated as described by Tauch et al. (1995, Plasmid
33:168-179) and partly cleaved with the restriction enzyme
Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product
Description Sau3AI, Code no. 27-0913-02). The DNA fragments
were dephosphorylated with shrimp alkaline phosphatase
10 (Roche Diagnostics GmbH, Mannheim, Germany, Product
Description SAP, Code no. 1758250). The DNA of the cosmid
vector SuperCos1 (Wahl et al. (1987) Proceedings of the
National Academy of Sciences USA 84:2160-2164), obtained
from Stratagene (La Jolla, USA, Product Description
15 SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved
with the restriction enzyme XbaI (Amersham Pharmacia,
Freiburg, Germany, Product Description XbaI, Code no. 27-
0948-02) and likewise dephosphorylated with shrimp alkaline
phosphatase.
- 20 The cosmid DNA was then cleaved with the restriction enzyme
BamHI (Amersham Pharmacia, Freiburg, Germany, Product
Description BamHI, Code no. 27-0868-04). The cosmid DNA
treated in this manner was mixed with the treated ATCC13032
DNA and the batch was treated with T4 DNA ligase (Amersham
25 Pharmacia, Freiburg, Germany, Product Description T4-DNA-
Ligase, Code no. 27-0870-04). The ligation mixture was then
packed in phages with the aid of Gigapack II XL Packing
Extract (Stratagene, La Jolla, USA, Product Description
Gigapack II XL Packing Extract, Code no. 200217).
- 30 For infection of the *E. coli* strain NM554 (Raleigh et al.
1988, Nucleic Acid Research 16:1563-1575) the cells were
taken up in 10 mM MgSO₄ and mixed with an aliquot of the
phage suspension. The infection and titering of the cosmid
library were carried out as described by Sambrook et al.

(1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones
5 were selected.

Example 2

Isolation and sequencing of the ppsA gene

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen,
10 Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline
15 phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen,
20 Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, Holland, Product Description Zero Background Cloning Kit, Product No. K2500-01), was cleaved with the restriction enzyme BamHI (Amersham Pharmacia,
25 Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being
30 incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A.,

87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l zeocin.

The plasmid preparation of the recombinant clones was carried out with the Biorobot 9600 (Product No. 900200, 5 Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). 10 The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) 15 (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 20 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231).

25 The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1095 base pairs, which was called the ppsA gene. The ppsA gene codes for a protein of 364 amino acids.

What is claimed is:

1. Isolated polynucleotide from coryneform bacteria,
comprising a polynucleotide sequence which codes for
the ppsA gene, chosen from the group consisting of
 - 5 a) polynucleotide which is identical to the extent of
at least 70% to a polynucleotide which codes for a
polypeptide which comprises the amino acid sequence
of SEQ ID No. 2,
 - 10 b) polynucleotide which codes for a polypeptide which
comprises an amino acid sequence which is identical
to the extent of at least 70% to the amino acid
sequence of SEQ ID No. 2,
 - c) polynucleotide which is complementary to the
polynucleotides of a) or b), and
 - 15 d) polynucleotide comprising at least 15 successive
nucleotides of the polynucleotide sequence of a),
b) or c)

the polypeptide preferably having the activity of
phosphoenol pyruvate synthase.
- 20 2. Polynucleotide according to claim 1, wherein the
polynucleotide is a preferably recombinant DNA which is
capable of replication in coryneform bacteria.
3. Polynucleotide according to claim 1, wherein the
polynucleotide is an RNA.
- 25 4. Polynucleotide according to claim 2, comprising the
nucleic acid sequence as shown in SEQ ID No. 1.
5. DNA according to claim 2 which is capable of
replication, comprising

- (i) the nucleotide sequence shown in SEQ ID No. 1,
or
- (ii) at least one sequence which corresponds to
sequence (i) within the range of the
degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the
sequence complementary to sequence (i) or (ii),
and optionally
- (iv) sense mutations of neutral function in (i).
6. DNA according to claim 5 which is capable of
replication, wherein the hybridization is carried out
under a stringency corresponding to at most 2x SSC.
7. Polynucleotide sequence according to claim 1, which
codes for a polypeptide which comprises the amino acid
sequence shown in SEQ ID No. 2.
8. Coryneform bacteria in which the ppsA gene is enhanced,
in particular over-expressed.
9. Process for the fermentative preparation of L-amino
acids, in particular L-lysine, wherein the following
steps are carried out:
- a) fermentation of the coryneform bacteria which
produce the desired L-amino acid and in which at
least the ppsA gene or nucleotide sequences which
code for it are enhanced, in particular over-
expressed;
- b) concentration of the L-amino acid in the medium or
in the cells of the bacteria, and
- c) isolation of the L-amino acid.

10. Process according to claim 9, wherein bacteria in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
- 5 11. Process according to claim 9, wherein bacteria in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partly eliminated are employed.
- 10 12. Process according to claim 9, wherein a strain transformed with a plasmid vector is employed, and the plasmid vector carries the nucleotide sequence which codes for the ppsA gene.
- 15 13. Process according to claim 9, wherein the expression of the polynucleotide(s) which code(s) for the ppsA gene is enhanced, in particular over-expressed.
14. Process according to claim 9, wherein the catalytic properties of the polypeptide (enzyme protein) for which the polynucleotide ppsA codes are increased.
- 20 15. Process according to claim 9, wherein for the preparation of L-amino acids, coryneform microorganisms in which at the same time one or more of the genes chosen from the group consisting of
 - 15.1 the dapA gene which codes for dihydrodipicolinate synthase,
 - 25 15.2 the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase,
 - 15.3 the tpi gene which codes for triose phosphate isomerase,
 - 15.4 the pgk gene which codes for 3-phosphoglycerate kinase,
- 30

- 15.5 the zwf gene which codes for glucose 6-phosphate dehydrogenase,
- 15.6 the pyc gene which codes for pyruvate carboxylase,
- 5 15.7 the mgo gene which codes for malate-quinone oxidoreductase,
- 15.8 the lysC gene which codes for a feed-back resistant aspartate kinase,
- 15.9 the lysE gene which codes for lysine export,
- 10 15.10 the hom gene which codes for homoserine dehydrogenase
- 15.11 the ilvA gene which codes for threonine dehydratase or the ilvA(Fbr) allele which codes for a feed back resistant threonine dehydratase,
- 15 15.12 the ilvBN gene which codes for acetohydroxy-acid synthase,
- 15.13 the ilvD gene which codes for dihydroxy-acid dehydratase,
- 20 15.14 the zwal gene which codes for the Zwal protein, is or are enhanced or over-expressed are fermented.
16. Process according to claim 9, wherein for the preparation of L-amino acids, coryneform microorganisms in which at the same time one or more of the genes
- 25 chosen from the group consisting of
- 16.1 the pck gene which codes for phosphoenol pyruvate carboxykinase,

SEQUENCE LISTING

<110> Degussa AG

5 <120> Nucleotide sequences which code for the ppsA gene

<130> 000175 BT

<140>

10 <141>

<160> 2

15 <170> PatentIn Ver. 2.1

<210> 1

<211> 1494

<212> DNA

20 <213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (200)..(1291)

25 <223> ppsA gene

<400> 1

cggcattccac tattgcctag gaaacatgct ggccaaactt caagccaaga tctgtctcga 60

30 ggaagtcacc aggcttggtc ctccctgca cttggttgcg gacaaagcta tcgggttccg 120

ggagaacctc tccttcgcg tccccacttc tgttcccggtg acttggaacg cttaacgctt 180

tattaaataa ggagacacc atg acc aac agt ttg aac atc ccg ttt gtc cag 232
 Met Thr Asn Ser Leu Asn Ile Pro Phe Val Gln
 35 1 5 10

cgc ttc gat gaa ggc ctg gat cct gtt cta gaa gta ctc ggt ggc aag 280
 Arg Phe Asp Glu Gly Leu Asp Pro Val Leu Glu Val Leu Gly Gly Lys
 40 15 20 25

ggc gct tca cta gtc acc atg aca gat gct gga atg ccc gtt cca cct 328
 Gly Ala Ser Leu Val Thr Met Thr Asp Ala Gly Met Pro Val Pro Pro
 50 30 35 40

gga ttt gtg gtc act act gcc agc ttt gat gaa ttc atc cgt gaa gca 376
 Gly Phe Val Val Thr Thr Ala Ser Phe Asp Glu Phe Ile Arg Glu Ala
 45 50 55

ggg gtt gct gaa cac atc gat aaa ttc cta aac gat ctc gat gca gaa 424
 Gly Val Ala Glu His Ile Asp Lys Phe Leu Asn Asp Leu Asp Ala Glu
 60 65 70 75

gat gtt aag gaa gtg gat cga gtt tct gcg atc atc cgc gat gag ctg 472
 Asp Val Lys Glu Val Asp Arg Val Ser Ala Ile Ile Arg Asp Glu Leu
 55 80 85 90

tgc agt ctt gac gtt cca gag aat gct cgt ttc gca gtg cac cag gct 520
 Cys Ser Leu Asp Val Pro Glu Asn Ala Arg Phe Ala Val His Gln Ala
 95 100 105

	tat cgc gat ctc atg gaa cga tgc ggt ggc gac gtc ccg gtt gct gtc	568
	Tyr Arg Asp Leu Met Glu Arg Cys Gly Gly Asp Val Pro Val Ala Val	
	110 115 120	
5	cgg tca tcg gcc act gcc gaa gat ctg ccc gat gct tcc ttc gca ggg	616
	Arg Ser Ser Ala Thr Ala Glu Asp Leu Pro Asp Ala Ser Phe Ala Gly	
	125 130 135	
10	caa cag gac acc tat ctg tgg caa gtc ggt ttg agc gct gtc act gaa	664
	Gln Gln Asp Thr Tyr Leu Trp Gln Val Gly Leu Ser Ala Val Thr Glu	
	140 145 150 155	
15	cac atc cgt aaa tgc tgg gct tcg ctg ttc act tcc cgt gcc att atc	712
	His Ile Arg Lys Cys Trp Ala Ser Leu Phe Thr Ser Arg Ala Ile Ile	
	160 165 170	
20	tac cgt ctg aaa aac aac atc ccc aat gag ggc ctc tcc atg gcg gta	760
	Tyr Arg Leu Lys Asn Asn Ile Pro Asn Glu Gly Leu Ser Met Ala Val	
	175 180 185	
25	gtt gtt caa aaa atg gtc aac tct cgt gtc gca ggc gtg gca atc act	808
	Val Val Gln Lys Met Val Asn Ser Arg Val Ala Gly Val Ala Ile Thr	
	190 195 200	
30	atg aat cct tcc aac ggc gac cgc tcg aag atc acc atc gat tcc tca	856
	Met Asn Pro Ser Asn Gly Asp Arg Ser Lys Ile Thr Ile Asp Ser Ser	
	205 210 215	
35	tgg ggt gtt ggt gaa atg gtg gtc tca ggt gaa gtg aca cca gac aat	904
	Trp Gly Val Gly Glu Met Val Val Ser Gly Glu Val Thr Pro Asp Asn	
	220 225 230 235	
40	atc ttg ctg gac aag atc acg ctg cag gtt gtc tcc gaa cac att gga	952
	Ile Leu Leu Asp Lys Ile Thr Leu Gln Val Val Ser Glu His Ile Gly	
	240 245 250	
45	agc aaa cac gct gaa ctc atc ccc gat gcc acc agt gga agc ctc gtg	1000
	Ser Lys His Ala Glu Leu Ile Pro Asp Ala Thr Ser Gly Ser Leu Val	
	255 260 265	
50	gaa aag ccc gtt gat gaa gaa cgc gca aac cgc cgc agt ctg act gat	1048
	Glu Lys Pro Val Asp Glu Glu Arg Ala Asn Arg Arg Ser Leu Thr Asp	
	270 275 280	
55	gag gaa atg ctc gct gtg gca caa atg gct aag cgt gca gaa aaa cac	1096
	Glu Glu Met Leu Ala Val Ala Gln Met Ala Lys Arg Ala Glu Lys His	
	285 290 295	
60	tac aag tgc cca caa gat atc gaa tgg gcg ctg gac gct gat ctg cca	1144
	Tyr Lys Cys Pro Gln Asp Ile Glu Trp Ala Leu Asp Ala Asp Leu Pro	
	300 305 310 315	
65	gat gga gaa aac ctt ctg tta ttg caa tcc cgc ccg gaa act atc cac	1192
	Asp Gly Glu Asn Leu Leu Leu Leu Gln Ser Arg Pro Glu Thr Ile His	
	320 325 330	
70	tcc aac ggt gtg aag aag gaa acc cca act ccg cag gct gcc aaa acc	1240
	Ser Asn Gly Val Lys Lys Glu Thr Pro Thr Pro Gln Ala Ala Lys Thr	
	335 340 345	

ata ggc acc ttc gat ttc agc tca atc acc gtc gca atg acc ggc acg 1288
 Ile Gly Thr Phe Asp Phe Ser Ser Ile Thr Val Ala Met Thr Gly Thr
 350 355 360
 5 aag taaaaccacc gcattttttc gtcgaaaagc atctaaaagg agtttgacca 1341
 Lys
 10 tggctaataa atctttcccc aagccctccg atcttccagt gccaagggc gctgaagggtt 1401
 gggaagatct gtaccgtac tacctcggtt tccaagacaa gctcatggat caagagaatg 1461
 agaaattctg gttctgcgat tcacagcact ggc 1494
 15 <210> 2
 <211> 364
 <212> PRT
 <213> Corynebacterium glutamicum
 20 <400> 2
 Met Thr Asn Ser Leu Asn Ile Pro Phe Val Gln Arg Phe Asp Glu Gly
 1 5 10 15
 25 Leu Asp Pro Val Leu Glu Val Leu Gly Gly Lys Gly Ala Ser Leu Val
 20 25 30
 30 Thr Met Thr Asp Ala Gly Met Pro Val Pro Pro Gly Phe Val Val Thr
 35 40 45
 35 Thr Ala Ser Phe Asp Glu Phe Ile Arg Glu Ala Gly Val Ala Glu His
 50 55 60
 Ile Asp Lys Phe Leu Asn Asp Leu Asp Ala Glu Asp Val Lys Glu Val
 65 70 75 80
 Asp Arg Val Ser Ala Ile Ile Arg Asp Glu Leu Cys Ser Leu Asp Val
 85 90 95
 40 Pro Glu Asn Ala Arg Phe Ala Val His Gln Ala Tyr Arg Asp Leu Met
 100 105 110
 45 Glu Arg Cys Gly Gly Asp Val Pro Val Ala Val Arg Ser Ser Ala Thr
 115 120 125
 Ala Glu Asp Leu Pro Asp Ala Ser Phe Ala Gly Gln Gln Asp Thr Tyr
 130 135 140
 50 Leu Trp Gln Val Gly Leu Ser Ala Val Thr Glu His Ile Arg Lys Cys
 145 150 155 160
 Trp Ala Ser Leu Phe Thr Ser Arg Ala Ile Ile Tyr Arg Leu Lys Asn
 165 170 175
 55 Asn Ile Pro Asn Glu Gly Leu Ser Met Ala Val Val Val Gln Lys Met
 180 185 190
 Val Asn Ser Arg Val Ala Gly Val Ala Ile Thr Met Asn Pro Ser Asn
 195 200 205

