

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 March 2011 (10.03.2011)

(10) International Publication Number
WO 2011/028121 A1

PCT

(51) International Patent Classification:

A01H 1/04 (2006.01) *A01H 5/08* (2006.01)
A01H 5/00 (2006.01)

(21) International Application Number:

PCT/NL2010/050562

(22) International Filing Date:

6 September 2010 (06.09.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09169555.1 4 September 2009 (04.09.2009) EP

(71) Applicant (for all designated States except US): **MON-SANTO INVEST N.V.** [NL/NL]; Leeuwenhoekweg 52, NL-2661 CZ Bergschenhoek (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MILLENAAR, Franciscus Frank** [NL/NL]; Achterstraat 5, NL-4064 EH Varik (NL). **VAN POPPEL, Petrus Marinus Johannes Abraham** [NL/NL]; Lombardi 60, NL-6708 LT Wageningen (NL).

(74) Agent: **JANSEN, C.M.**; Vereenigde, Johan de Wittlaan 7, NL-2517 JR Den Haag (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: TOMATO PLANTS EXHIBITING TOLERANCE TO CONTINUOUS LIGHT

(57) Abstract: The present invention relates to a method of selecting a tomato plant comprising exposing a tomato plant and determining whether said plant exhibits tolerance to exposure to continuous light, wherein said tolerance is indicated by decreased level of chlorosis upon said exposure, and further determining whether said plant exhibits a significant increase in yield when exposed to continuous light. The invention further relates to a method for increasing the yield of a tomato plant line, said method comprising selecting a tomato plant by the method of the invention; crossing the thus selected plant with a plant of a said tomato plant line to produce offspring plants; optionally backcrossing or selfing said offspring plant to produce further offspring plants, and selecting from said offspring plants a plant having tolerance to continuous light and having increased yield as compared to said plant of the original tomato plant line.



WO 2011/028121 A1

Title: Tomato plants exhibiting tolerance to continuous light

FIELD OF THE INVENTION

The present invention relates to plant breeding and molecular biology. More specifically, the present invention relates to a method for detecting a gene associated with tolerance to continuous light in tomato, to a method of producing a continuous
5 light-tolerant tomato plant therewith and to continuous light-tolerant tomato plants thus obtained and parts thereof.

BACKGROUND OF THE INVENTION

Tomato (*Solanum lycopersicum*, formerly known as *Lycopersicon esculentum*)
10 is a plant of the Solanaceae or nightshade family. It is a short-lived perennial plant, grown as an annual plant, and a close relative of the potato. The fruit (i.e. the tomato) is an edible, brightly coloured (usually red, from the pigment lycopene) berry, 1-2 cm diameter in wild plants, commonly much larger in cultivated forms. The plant is now grown worldwide for its edible fruits, which are major fresh market vegetables
15 worldwide.

During the last decades, tomato breeding was mainly focused on yield, disease tolerance, and fruit quality aspects such as uniform ripening and taste. Yield improvements have been achieved due to new production methods, improved pest management and varieties that are better suited for new production methods, but the
20 gains in yield become smaller. New varieties with 5 or 15 fruits more per plant gave a yield increase of 2-4%.

As the world population continues to grow, the demand for fresh vegetables, such as tomatoes, is ever increasing worldwide. Thus, a continuing need exists for means and methods for improving yield of tomatoes.

25

SUMMARY OF THE INVENTION

The inventors have surprisingly found that the yield of some tomato plants can be increased by the use of longer photoperiods or even continuous photoperiods. This is surprising, because commercially used varieties, when grown in continuous

light, show a strong increase in chloroplastic starch, leaf chlorosis, necrosis and consequently very poor biomass production. In effect, continuous light is often lethal for the crop. However, the inventors have found that tomato plants exist which are not only tolerant to continuous light, but in fact display an increase in yield when
5 exposed to continuous light when compared to 16 hours of light.

The present inventors have successfully identified a quantitative trait locus (gene) which is responsible for tolerance to continuous light present in the genome of several wild relatives of tomato, i.e. in *S. neorickii*, *S. habrochaites*, *S. pennellii*, *S. peruvianum*, *S. chilense* and *S. chemielewskii*, and in *S. lycopersicum*, e.g. var. Sub
10 Arctic Plenty.

The inventors were subsequently able to produce continuous light tolerant tomato plants by crossing these continuous light tolerant wild (donor) tomato lines with non-tolerant recipient tomato plants. These plants thus produced exhibited more tolerance to continuous light than any cultivated tomato plant produced thus far. In
15 addition, several plants also produced more biomass when exposed to continuous light than plants from the original recipient line when exposed to 16 hours of light. Therefore, the present invention enables more efficient culturing and higher yields.

The present invention provides in a first aspect a method of selecting a tomato plant comprising exposing a tomato plant and determining whether said plant
20 exhibits tolerance to continuous light, wherein said tolerance is indicated by decreased level of chlorosis upon said exposure, and further determining whether said plant exhibits a significant increase in yield.

The invention further provides a method for producing a first tomato plant by a method according to the invention, crossing said selected plant with a second
25 tomato plant to produce offspring plants and selecting from said offspring plants a plant having tolerance to continuous light and having increased yield as compared to said second tomato plant.

The invention further provides a method of producing a tomato plant, tolerant to exposure to continuous light, said method comprising steps of:

30 a) providing a tomato donor plant tolerant to exposure to continuous light, preferably of the species *S. neorickii*, *S. habrochaites*, *S. pennellii*, *S. lycopersicum*, *S. peruvianum*, *S. chilense* or *S. chemielewskii*;

b) transferring nucleic acid from said donor plant to at least one recipient tomato plant which is not tolerant to exposure to continuous light, wherein said transfer results in the introduction of genomic material from said donor plant in the corresponding region of the genome of said at least one recipient plant;

- 5 c) selecting from said at least one recipient plant a plant that comprises within its genome a gene for continuous light tolerance derived from said donor plant, wherein said selection comprises detecting in chromosome 7 of said recipient tomato plant at least one genetic marker linked to said gene for light tolerance.

10 In preferred embodiments, the location of said gene on chromosome 7 of said plant is indicated by a genomic region comprising the genetic markers "7-20-1", "7-20-2" on chromosome 7 of *S. neorickii*, "7-10", "7-25" on chromosome 7 of *S. neorickii*, "7-17", "7-25" on chromosome 7 of *S. chilense*, "7-19", "7-25" on chromosome 7 of *S. pennellii*.

15 In another preferred embodiment, said transfer of nucleic acid comprising said gene for tolerance to continuous light, or a light tolerance conferring part thereof, is performed by crossing said donor tomato plant tolerant to exposure to continuous light with a recipient tomato plant which is not tolerant to exposure to continuous light, to produce offspring plants comprising said gene as an introgression, and wherein step c) is performed on at least one offspring plant.

20 In another preferred embodiment, said step c) is performed by detecting said genetic marker in DNA isolated from said at least one recipient tomato plant.

 In another preferred embodiment, said step c) further comprises subjecting said plant to bioassay for measuring the capability of growing in continuous light of said plant.

25 In another aspect the invention provides a continuous light tolerant tomato plant or a part thereof, obtainable by a method according to the invention.

 The invention further provides a gene for continuous light tolerance, wherein said gene is associated with tolerance to continuous light and wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region comprising
30 the genetic markers "7-20-1", "7-20-2" on chromosome 7 of *S. neorickii*, "7-10", "7-25" on chromosome 7 of *S. neorickii*, "7-17", "7-25" on chromosome 7 of *S. chilense*, "7-19", "7-25" on chromosome 7 of *S. pennellii*. The alleles on the positions of the genome indicated by this gene is an aspect of the present invention.

A gene of the present invention may be in the form of an isolated, preferably double stranded nucleic acid sequence comprising said gene or a tolerance-conferring part thereof. Very suitably, the size of the nucleic acid sequence, which may for instance be isolated from the chromosome of a suitable donor plant, may represent a genetic distance of 1-100 cM, preferably 10-80 cM on said chromosome. Preferably said genetic distance is between 64 and 78 cM on said chromosome, even more preferably between 72 and 75 cM. Said nucleic acid may comprise at least 50, more preferably at least 500, even more preferably at least 1000, still more preferably at least 5000 base pairs. One or more nucleic acid sequences comprising a gene or a tolerance-conferring part thereof according to the invention may in turn be comprised in a nucleic acid construct, said construct may further comprise regions that flank said one or more nucleic acid sequences and which regions are capable of being integrated into a suitable vector for transfer of said one or more nucleic acid sequences into a suitable continuous light intolerant recipient tomato plant. The vector may further comprise suitable promoter regions or other regulatory sequences. Said gene may also be in a form present within the genome of a tomato plant. The gene of the present invention preferably comprises at least one marker, preferably two markers associated with continuous light-tolerance selected from the group consisting of the markers of Figure 10 linked to said gene.

The invention further provides a method for detecting a gene for tolerance to continuous light, comprising detecting at least one genetic marker linked to a gene for tolerance to continuous light derived on chromosome 7 of a suspected continuous light tolerant tomato plant, wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region comprising the genetic markers "7-20-1", "7-20-2" on chromosome 7 of *S. neorickii*, "7-10", "7-25" on chromosome 7 of *S. neorickii*, "7-17", "7-25" on chromosome 7 of *S. chilense*, "7-19", "7-25" on chromosome 7 of *S. pennellii*.

In another aspect the invention provides a continuous light-tolerant tomato plant, or a part thereof, comprising within its genome a gene for continuous light tolerance, or a continuous light-tolerance-conferring part thereof, wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region comprising the genetic markers "7-20-1", "7-20-2" on chromosome 7 of *S. neorickii*, "7-10", "7-25" on chromosome 7 of *S. neorickii*, "7-17", "7-25" on chromosome 7 of *S. chilense*, "7-19", "7-25" on chromosome 7 of *S. pennellii*, wherein said gene or said

continuous light-tolerance-conferring part thereof is not in its natural genetic background.

In another aspect the invention provides a method of producing a continuous light-tolerant inbred tomato plant, comprising

- 5 a) producing a continuous light-tolerant tomato plant according to the invention;
- b) crossing said continuous light-tolerant tomato plant with itself or another tomato plant to yield progeny tomato seed;
- c) growing said progeny tomato seed of step b) to yield an additional continuous light-tolerant tomato plant;
- 10 d) repeating the crossing and growing steps from 0 to 7 times to generate a continuous light-tolerant inbred tomato plant.

Preferably said step c) further comprises the step of identifying plants that exhibit continuous light tolerance and possess commercially desirable characteristics.

- In another preferred embodiment said method further comprises the step of
- 15 selecting homozygote inbred tomato plants, preferably wherein said inbred plants are homozygous for said continuous light allele.

In a further aspect, the invention provides a continuous light-tolerant inbred tomato plant, or parts thereof, obtainable the method according to the invention.

- In a further aspect, the invention provides a hybrid tomato plant, or a part
- 20 thereof, that exhibits tolerance to continuous light, wherein said hybrid tomato plant is obtainable by crossing a continuous light-tolerant inbred tomato plant obtainable by a method according to the invention with an inbred tomato plant that exhibits commercially desirable characteristics. Such characteristics include *inter alia* a high yield (in excess of 50, 60 kg/m² of tomatoes and/or resistance to tobacco mosaic virus (TMV), blight and/or *Botrytis*.
 - 25

In another aspect, the invention further provides the use of a genetic marker selected from the group consisting of the genetic markers of Figure 10 for the detection of genes for continuous light-tolerance, and/or for the detection of continuous light-tolerant tomato plants.

- 30 In another aspect, the invention provides an indeterminate tomato plant, tolerant to exposure to continuous light or a part thereof.

In a further aspect, the invention further provides a tissue culture of regenerable cells of the tomato plants according to the invention, preferably said

regenerable cells comprise cells or protoplasts isolated from a tissue selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, flowers, fruits, and stems and seeds.

The continuous light-tolerant donor tomato plant used in methods of the present invention is preferably selected from the group consisting of *S. neorickii*, *S. habrochaites*, *S. pennellii*, *S. lycopersicum*, *S. peruvianum*, *S. chemielewskii*, more preferably, a wild tomato accession is used as the donor plant. Highly preferred donor plants are *S. pennellii* LA716, *S. peruvianum* LA1708, *S. habrochaites* LA1777, *S. habrochaites* G1560, *S. habrochaites* Lyc4/78, *S. chilense* LA1959, *S. chmielewskii* LA1840, *S. neorickii* LA2133 and *S. lycopersicum* variety "Sub arctic plenty".

The continuous light-tolerant recipient tomato plant used in methods of the present invention is preferably a plant of the species *Solanum lycopersicum*, more preferably an *S. lycopersicum* cultivar that possess commercially desirable characteristics, or another commercial tomato line.

DESCRIPTION OF THE DRAWINGS

Figure 1: Species and varieties tolerant to continuous light. Plants shown on the left were exposed to 16 hr and plants on the right are exposed to continuous light. Clearly the typical symptoms of light intolerance (leaf chlorosis, necrosis and poor biomass production) are not expressed in these tolerant lines, while the symptoms are clearly visible in non tolerant lines under the same conditions.

Figure 2: typical symptoms caused by exposure to continuous light in non-tolerant lines. The first visible symptoms of non tolerant lines are light green leaves with inter-vine chlorosis, followed by severe chlorosis. In severe non tolerant plants chlorosis is followed by necrosis, decreased leaf development decreased stem development and even plant mortality.

Figure 3: Degreening and chlorosis. Quantification of a part of the visible symptoms (degreening and chlorosis) by measuring the chlorophyll concentration by a SPAD meter was performed with plants grown at $131 \mu\text{mol m}^{-2} \text{s}^{-1}$ (ML) or $230 \mu\text{mol m}^{-2} \text{s}^{-1}$ (HL). The resolution to detect differences between continuous and non-continuous light was very difficult or impossible at very low light intensities ($16 \mu\text{mol m}^{-2} \text{s}^{-1}$ =LL). The visible symptoms developed earlier in time at $230 \mu\text{mol m}^{-2} \text{s}^{-1}$ compared to $131 \mu\text{mol m}^{-2} \text{s}^{-1}$. Left: averages with standard deviations, number of

replicates is 20 to 40. Right: Data extracted from the left graph, plotted is the differences between 16 and 24 hours of light per light intensity. Significance level is shown by ns= not significant, *= $p < 0.05$, **= $p < 0.01$, ***= $p < 0.001$ and ****= $p < 0.00001$.

5 **Figure 4: Stem diameter and plant height is decreased in non tolerant lines.** Therefore stem diameter and plant height are also parameters to be able to quantify continuous light symptoms. Averages with standard errors, number of replicates is 4 to 8. Significance level is shown by += $p < 0.1$, *= $p < 0.05$, **= $p < 0.01$, ***= $p < 0.001$.

10 **Figure 5: Chlorophyll concentration of IL lines from a *neorickii* * a sensitive *esculentum* (e.g. Moneyberg) population where 9 days old seedlings exposed for 33 days to continuous light.** Temperature 21°C for 24 hours and, with an RH of 70% and a light intensity of 110-120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR) from fluorescent tubes. Averages with standard errors, number of replicates was 3 to 6 plants per line.

15 **Figure 6: One highly tolerant line (42) from a *neorickii* * a sensitive *esculentum* (left).** Tolerant lines are exposed to continuous light for 90 days. The first 33 days under fluorescent tubes, followed by illumination under High pressure sodium lamp (SON-T). The leaves stay green, leaf and stem development is normal compared to non tolerant lines.

20 **Figure 7: Chlorophyll concentration of IL lines from a *chilense* LA1959 * Moneyberg GT population where 15 days old seedlings exposed for 35 (CLT1), 42 (CLT2), 49 (CLT3) and 60 (CLT4) days to continuous light.** Temperature 21°C/20 °C for 16/8 hours and, with an RH of 70% and a light intensity of 110-120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR) from fluorescent tubes during the first 15 days followed
25 by illumination under High pressure sodium lamp (SON-T) for 24 Hr. Averages with standard errors, number of replicates was 3 to 6 plants per line.

Figure 8: One highly tolerant line (5) from a *chilense* LA1959 * a sensitive Moneyberg GT. Tolerant lines are exposed to continuous light for 90 days. The leaves stay green, leaf and stem development is normal compared to non tolerant
30 lines.

Figure 9: Chlorophyll concentration of IL lines from a *pennellii* * M82 population with an introgression on chromosome 7. Chlorophyll concentration of IL lines from a *pennellii* * M82 population where 14 days old seedlings were

exposed for 21 (CLT1), 28 (CLT2), 35 (CLT3), 42 (CLT4) and 53 (CLT5) days to continuous light. Temperature 21°C/20 °C for 16/8 hours and, with an RH of 70% and a light intensity of 110-120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR) from fluorescent tubes the first 14 days after that under SON-T for 24 Hr. Averages with standard errors, number of replicates was 4 to 6 plants per line. Two tolerant lines were found (4 and 5), which have been in continuous light for 53 days.

Figure 10: Genotypic information of the different mapping populations, based on Illumina SNP markers. Combining the phenotypic and genotypic data from the three populations, the continuous light tolerant loci is located on chromosome 7 between marker “7-19” and “7-23” that is between 64 and 99,6 cM based on the EXPEN2000 linkage map. Wild donor alleles are shown in light grey, recurrent parent alleles in dark grey, different alleles in black and no call in white. Some SNPs are non polymorphic in some wild donor x recurrent parent combinations, in such cases a gap is present. SNPs markers were genotyped using GoldenGate assays (Illumina).

Figure 11: Sequences of genetic markers. The markers indicate the polymorphic sequences between the tolerant and sensitive genotype. The polymorphic positions as indicated in these sequences are SNPs. An example for use of these SNPs as markers is provided in Example 7 below. The skilled artisan will understand that the markers as indicated in this figure refer specifically to the polymorphic positions in the sequences given. **Marker 1:** Sequence of the marker “7-1” derived from *S. neorickii* LA2133 located at 0,4 cM of chromosome 7 ([T/G] indicates the polymorphic position). **Marker 2:** Sequence of the marker “7-2” derived from *S. pennellii* 0716 located at 1,2 cM of chromosome 7 ([T/G] indicates the polymorphic position). **Marker 3:** Sequence of the marker “7-3” derived from *S. neorickii* LA2133 located at 2,0 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 4:** Sequence of the marker “7-4” derived from *S. neorickii* LA2133 located at 3,5 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 5:** Sequence of the marker “7-5” derived from *S. neorickii* LA2133 located at 21,4 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 6:** Sequence of the marker “7-6” derived from *S. neorickii* LA2133 located at 22,3 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 7:** Sequence of the marker “7-7” derived from *S. neorickii* LA2133 located at 24,8 cM of chromosome 7 ([T/C] indicates the polymorphic position).

Marker 8: Sequence of the marker “7-8” derived from *S. neorickii* LA2133 located at 27,5 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 9:** Sequence of the marker “7-9” derived from *S. pennellii* 0716 located at 28,1 cM of chromosome 7 ([A/G] indicates the polymorphic position). **Marker 10:** Sequence of the marker “7-10” derived from *S. neorickii* LA2133 located at 38,0 cM of chromosome 7 ([A/G] indicates the polymorphic position). **Marker 11:** Sequence of the marker “7-11” derived from *S. neorickii* LA2133 located at 38,8 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 12:** Sequence of the marker “7-12” derived from *S. pennellii* 0716 located at 40,0 cM of chromosome 7 ([T/G] indicates the polymorphic position). **Marker 13:** Sequence of the marker “7-13” derived from *S. neorickii* LA2133 located at 43,0 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 14:** Sequence of the marker “7-14” derived from *S. neorickii* LA2133 located at 45,6 cM of chromosome 7 ([A/G] indicates the polymorphic position). **Marker 15:** Sequence of the marker “7-15” derived from *S. neorickii* LA2133 located at 48,1 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 16:** Sequence of the marker “7-16” derived from *S. neorickii* LA2133 located at 49,2 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 17:** Sequence of the marker “7-17” derived from *S. neorickii* LA2133 located at 54,0 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 18:** Sequence of the marker “7-18” derived from *S. neorickii* LA2133 located at 61,0 cM of chromosome 7 ([T/G] indicates the polymorphic position). **Marker 19:** Sequence of the marker “7-19” derived from *S. pennellii* 0716 located at 64,0 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 20:** Sequence of the marker “7-20” derived from *S. neorickii* LA2133 located at 46,18 cM of chromosome 7 based on mapping in the 2133-42 population and at 73,0 cM based on mapping using the EXPEN 2000 linkage map of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 21:** Sequence of the marker “7-21” derived from *S. neorickii* LA2133 located 46,74 cM of chromosome 7 based on mapping in the 2133-42 population and at 78,0 cM based on mapping using the EXPEN 2000 linkage map of chromosome 7 ([A/G] indicates the polymorphic position). **Marker 22:** Sequence of the marker “7-22” derived from *S. neorickii* LA2133 located at 80,0 cM of chromosome 7 ([T/C] indicates the polymorphic position). **Marker 23:** Sequence of the marker “7-23” derived from *S. neorickii* LA2133 located at 99,6 cM of chromosome 7

([T/C] indicates the polymorphic position). **Marker 24:** Sequence of the marker “7-24” derived from *S. pennellii* 0716 located at 100,4 cM of chromosome 7 ([A/T] indicates the polymorphic position). **Marker 25:** Sequence of the marker “7-25” derived from *S. neorickii* LA2133 located at 104,0 cM of chromosome 7 ([A/G] indicates the polymorphic position). **Marker 26** Sequence of the marker “7-20-1” derived from *S. neorickii* LA2133 located at 43.68 cM of chromosome 7 based on mapping in the LA2133-42 population and at 75 cM based on mapping using the EXPEN 2000 linkage map ([T/G] indicates the polymorphic position). **Marker 27** Sequence of the marker “7-20-2” derived from *S. neorickii* LA2133 located at 44.35 cM of chromosome 7 based on mapping in the LA2133-42 population and at 72 cM based on mapping using the EXPEN 2000 linkage map ([C/A] indicates the polymorphic position). **Marker 28** Sequence of the marker “7-20-3” derived from *S. neorickii* LA2133 located at 44.46 cM of chromosome 7 based on mapping in the LA2133-42 population and at 73 cM based on mapping using the EXPEN 2000 linkage map ([G/C] indicates the polymorphic position).

Figure 12: the CTL locus. The CTL locus is positioned between markers 7-17 and 7-20. Presented genotypic data (the genotypes include “homozygous for donor” or “recurrent parent” and “heterozygous”) is generated from F2 plants, while phenotypic scoring (the phenotypes include “Tolerant to continuous light”, “susceptible for continuous light” and “segregating”) is performed on F3 plants. Light grey blocks indicate genomic segments originating from the tolerant parent. Dark grey blocks indicate segments that are originating from the sensitive parent. White blocks indicate heterozygous segments. Markers are ordered according to map LA2133-42.

Figure 13. The CTL locus is positioned between markers 7-20-1 and 7-20-2. Presented genotypic data (the genotypes include “homozygous for donor” or “recurrent parent” and “heterozygous”) is generated from F2 plants, while phenotypic scoring (the phenotypes include “tolerant to continuous light”, “susceptible for continuous light” and “segregating”) is performed on F3 plants. Dark grey blocks indicate genomic segments originating from the sensitive parent. Light grey blocks indicate segments that are originating from the tolerant parent. White blocks indicate heterozygous segments. Markers are ordered according to map LA2133-42.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The term "light" as used herein means light of a suitable wavelength for growing plants at an average light intensity which is higher than the light energy known as the "light compensation point". This is the light intensity at which the amount of carbon dioxide released in respiration equals the amount used in photosynthesis, and the amount of oxygen released in photosynthesis equals the amount used in respiration.

The compensation point is the amount of light intensity on the light curve where the rate of photosynthesis exactly matches the rate of respiration. At this point, the uptake of CO₂ through photosynthetic pathways is exactly matched to the respiratory release of carbon dioxide, and the uptake of O₂ by respiration is exactly matched to the photosynthetic release of oxygen. Methods to establish the light compensation point of a plant are well known in the art. Preferably, said light has a light intensity of at least 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$, more preferably at least 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$, still more preferably at least 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PAR (photosynthetic active radiation). Preferably said light has a light intensity of less than 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$, preferably less than 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

As used herein, the term "continuous light" means a photoperiod of more than 20 hours light per day, preferably more than 21 hours, preferably more than 22 hours, more preferably more than 23 hours and most preferably 24 hours. Short interruptions of a few minutes of this photoperiod may be comprised in continuous lighting. The total duration of the exposure to said light per day must be more than 20 hours.

As used herein, the term "continuous light tolerance" refers to a plant which, when grown under continuous light, exhibits a reduction in severity of at least one symptom of the complex of symptoms typically associated with continuous light intolerance as observed in normal non tolerant (tomato) plants of for instance the variety Moneyberg within 20 days after the first exposure to continuous light. Preferably, a plant which is tolerant to continuous light does not exhibit said at least one symptom after at least 3 weeks of exposure to continuous light. Even more preferably, said at least 3 weeks is at least four weeks. More preferably, it is at least 35, 42, 63 or 90 days. Preferably, said plant which is tolerant to continuous light has a

higher chlorophyll concentration in its chlorophyll containing cells when exposed to continuous light than a non tolerant control plant which is exposed to the same conditions. Preferably, said plant which is tolerant to continuous light has a chlorophyll concentration of at least 35 (relative scale from 0-100, 100 is deep green),
5 more preferably 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60 as determined by a SPAD-502 (Konica Minolta) meter, wherein said plant has been exposed for at least, 33, 35, 42, 49, 53, 90 days to continuous light with a light intensity of $130 \mu\text{mol m}^{-2} \text{s}^{-1}$ from fluorescent tubes and grown at a temperature of 21°C and a relative humidity between 60-80%, preferably 70%. In
10 another preferred embodiment, said plant which is tolerant to continuous light has a chlorophyll concentration which is at least 5%, more preferably 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50% higher than its sensitive parent plant exposed to the same conditions, as determined by using a SPAD-502 (Konica Minolta) meter, wherein said plant has been exposed for at least, 33, 35, 42, 49, 53, 90 days to continuous light with
15 a light intensity of $130 \mu\text{mol m}^{-2} \text{s}^{-1}$ from fluorescent tubes and grown at a temperature of 21°C and a relative humidity between 60-80%, preferably 70%.

Said complex of symptoms includes but is not limited to leaf chlorosis, necrosis, decreased leaf development, increase in chloroplastic starch, decreased stem development, reduced photosynthesis, degreening and poor biomass production.
20 Preferably all symptoms are absent in a plant tolerant to continuous light.

The term "leaf chlorosis" as used herein refers to a yellowing of leaf tissue due to absence of chlorophyll. The absence of chlorophyll can be visually observed as yellow spots on the leaf tissue.

The term "necrosis" as used herein refers to the premature death of cells and
25 living tissue due to external factors. Necrosis can be visually observed as light to dark brown spots or absence of cells in tissue, preferably leaf tissue.

The term "decreased leaf development" as used herein refers to a decreased total surface area of the combined leaves of a plant which has been exposed to continuous light compared to the total surface area of the combined leaves of a
30 comparable control plant which has not been exposed to continuous light. Total surface area of the combined leaves of a plant can be determined by measuring the total leaf surface area of a plant and/or by weighing the mass of the leaves of a plant.

The term “increase in chloroplastic starch” refers to an increase in the amount of starch in a sample comprising chloroplasts from a plant which has been exposed to continuous light compared to the amount of starch from a comparable sample of a comparable control plant which has not been exposed to continuous light.

5 Alternatively, the increase of chloroplastic starch is determined by comparing the chloroplastic starch content of a plant or part thereof which contains chloroplasts with the chloroplastic starch content of a comparable control plant or chloroplast containing part thereof which has not been exposed to continuous light. The amount of starch of a plant is preferably determined by allowing a sample from said plant or
10 part thereof which contains chloroplasts to gelatinize, incubating said sample with amyloglucosidase and subsequently determining the amount of glucose released. Preferably, said sample comprises isolated chloroplasts from said plant. The chloroplastic starch content is determined by calculating the amount of chloroplastic starch per kg of the plant.

15 The term “decreased stem development” as used herein refers to a decreased length and/or diameter of a stem of a plant or a part thereof which has been exposed to continuous light compared to the length and/or diameter of a comparable stem of a comparable control plant which has not been exposed to continuous light.

The term “reduced photosynthesis” as used herein refers to a reduced level of
20 activity of photosynthesis in a plant or in a part thereof which has been exposed to continuous light compared to level of activity of photosynthesis in a comparable control plant which has not been exposed to continuous light. The term “photosynthesis” as used herein refers to the process in which the plant converts carbon dioxide into organic compounds using sunlight or artificial light as energy. The
25 level of activity of photosynthesis can be determined by measuring the carbon dioxide uptake or oxygen production. A decrease in the carbon oxide uptake of a plant and/or in the oxygen production in comparison to a control plant indicates a decrease in the activity of photosynthesis.

The term “degreening” as used herein refers to the decrease in the content of
30 chlorophyll in leaf tissue of a plant or in a part thereof which has been exposed to continuous light compared to a comparable control plant which has not been exposed to continuous light. Degreening can be visually observed as light green or whitened leaves. Alternatively, the content of chlorophyll in leaf tissue can be determined by

performing a SPAD measurement well known in the art and compare the chlorophyll measurements performed in leaf tissue of said control plant and establish whether the content of chlorophyll is decreased.

5 The term "poor biomass production" as used herein refers to a lower (e.g. at least 5%, 10%, 15% lower) amount of biomass of a plant which has been exposed to continuous light compared to a comparable control plant which has not been exposed to continuous light. The biomass of a plant can be determined by measuring the fresh and/or dry weight of a plant.

10 As used herein, the term "allele(s)" means any of one or more alternative forms of a gene, all of which alleles relate to at least one trait or characteristic. In a diploid cell or organism, the two alleles of a given gene occupy corresponding loci on a pair of homologous chromosomes. Since the present invention relates to genes, i.e. genomic regions that may comprise one or more genes, but also regulatory sequences, it is in some instances more accurate to refer to "haplotype" (i.e. an allele of a
15 chromosomal segment) in stead of "allele", however, in those instances, the term "allele" should be understood to comprise the term "haplotype".

A "gene" is defined herein as a hereditary unit consisting of a sequence of DNA that occupies a specific location on a chromosome and that contains the genetic instruction for a particular characteristics or trait in an organism.

20 A "locus" is defined herein as the position that a given gene occupies on a chromosome of a given species.

As used herein, the term "heterozygous" means a genetic condition existing when different alleles reside at corresponding loci on homologous chromosomes.

25 As used herein, the term "homozygous" means a genetic condition existing when identical alleles reside at corresponding loci on homologous chromosomes.

As used herein, the term "hybrid" means any offspring of a cross between two genetically unlike individuals, including but not limited to the cross between two inbred lines.

30 As used herein, the term "inbred" means a substantially homozygous individual or line

In this application a "recombination event" is understood to mean a meiotic crossing-over.

As used herein, the terms "introgression", "introgressed" and "introgressing" refer to both a natural and artificial process whereby genes of one species, variety or cultivar are moved into the genome of another species, variety or cultivar, by crossing those species, varieties or cultivars. The process may optionally be completed by

5 backcrossing to the recurrent parent.

"Genetic engineering", "transformation" and "genetic modification" are all used herein as synonyms for the transfer of isolated and cloned genes into the DNA, usually the chromosomal DNA or genome, of another organism.

As used herein, the term "plant part" indicates a part of the tomato plant, including single cells and cell tissues such as plant cells that are intact in plants, cell clumps and tissue cultures from which tomato plants can be regenerated. Examples of plant parts include, but are not limited to, single cells and tissues from pollen, ovules, leaves, embryos, roots, root tips, anthers, flowers, fruits, stems shoots, and seeds; as well as pollen, ovules, leaves, embryos, roots, root tips, anthers, flowers, fruits, stems, shoots, scions, rootstocks, seeds, protoplasts, calli, and the like.

As used herein, the term "population" means a genetically heterogeneous collection of plants sharing a common genetic derivation.

As used herein, the term "tomato" means any plant, line or population formerly known under the genus name of *Lycopersicon* including but not limited to

20 *Lycopersicon cerasiforme*, *Lycopersicon cheesmanii*, *Lycopersicon chilense*, *Lycopersicon chmielewskii*, *Lycopersicon esculentum* (now *Solanum lycopersicum*), *Lycopersicon hirsutum*, *Lycopersicon parviflorum*, *Lycopersicon pennellii*, *Lycopersicon peruvianum*, *Lycopersicon pimpinellifolium*, or *Solanum lycopersicoides*. The newly proposed scientific name for *Lycopersicon esculentum* is *Solanum*

25 *lycopersicum*. Similarly, the names of the wild species has been altered. *L. pennellii* has become *Solanum pennellii*, *L. hirsutum* has become *S. habrochaites*, *L. peruvianum* has be split into *S. 'N peruvianum'* and *S. 'Callejon de Huayles'*, *S. peruvianum*, and *S. corneliomuelleri*, *L. parviflorum* has become *S. neorickii*, *L. chmielewskii* has become *S. chmielewskii*, *L. chilense* has become *S. chilense*, *L. cheesmaniae* has become *S. cheesmaniae* or *S. galapagense*, and *L. pimpinellifolium*

30 has become *S. pimpinellifolium* (Solanacea Genome Network (2005) Spooner and Knapp; http://www.sgn.cornell.edu/help/about/solanum_nomenclature.html).

The term "*S. lycopersicum*", as used herein, refers to any variety or cultivar of the garden tomato.

It is especially noted that *S. habrochaites* can be defined as a tomato species that carries hairy fruits, while *S. lycopersicum* is a tomato species carrying hairless
5 fruits.

As used herein, the term "variety" or "cultivar" means a group of similar plants that by structural or genetic features and/or performance can be distinguished from other varieties within the same species.

The term "gene associated with tolerance to continuous light" as well as the
10 shorter term "gene for continuous light tolerance" refer to a region located on a particular chromosome of tomato that is associated with at least one gene that encodes for continuous light-tolerance or at least a regulatory region, i.e. a region of a chromosome that controls the expression of one or more genes involved in tolerance to continuous light. The phenotypic expression of that gene may for instance be observed
15 as a reduced degree of chlorosis or an improved biomass production upon exposure to continuous light. A gene may for instance comprise one or more genes of which the products confer the genetic continuous light tolerance. Alternatively, a gene may for instance comprise regulatory genes or sequences of which the products influence the expression of genes on other loci in the genome of the plant thereby conferring the
20 tolerance to continuous light. The gene of the present invention may be defined by indicating their genetic location in the genome of the respective wild tomato accession using one or more molecular genomic markers. One or more markers, in turn, indicate a specific locus. Distances between loci are usually measured by frequency of crossing-over between loci on the same chromosome. The farther apart two loci are, the more
25 likely that a crossover will occur between them. Conversely, if two loci are close together, a crossover is less likely to occur between them. As a rule, one centimorgan (cM) is equal to 1% recombination between loci (markers). When a gene can be indicated by multiple markers the genetic distance between the end-point markers is indicative of the size of the gene. It is to be understood that the genes identified
30 herein as residing in the CLT locus (see Tables 2 and 3) may be used as markers for the continuous light resistance trait in aspects of this invention. In particular, polymorphisms in these genes between the tolerant and non-tolerant phenotype are envisioned.

The term "natural genetic background" is used herein to indicate the original genetic background of a gene. Such a background may for instance be the genome of a continuous light-tolerance wild accession of tomato. For instance, the gene of the present invention was found at specific locations on chromosomes 7 of *Solanum pennellii* 0716. As an example, the *Solanum pennellii* 0716 represents the natural genetic background of the genes on chromosomes 7 of *Solanum pennellii* 0716. Conversely, a method that involves the transfer of DNA comprising the gene, or a tolerance-conferring part thereof, from chromosome 7 of *Solanum pennellii* 0716 to the same position on chromosome 7 of another tomato species, will result in that gene, or said tolerance-conferring part thereof, not being in its natural genetic background.

The terms "determinate" and "indeterminate" as used herein refers to the type of growth habit of tomato plants, which is commonly classified as determinate or indeterminate. This classification depends preferably on the capacity of the shoot system for continued sympodial growth. The terms are used in their art-recognized meaning.

The term "standard practice conditions", "standard greenhouse conditions" and "standard conditions" refer to the conditions of light, humidity, temperature, etc. where under plants are grown or incubated, for instance for the purpose of phenotypic characterization, as being standard. For greenhouses or climate chambers for instance, this refers to 8-16-h day, 15°C-25°C. More in general, the terms refer to standard and reference growth conditions with a photoperiod of 8 to 24 h (photosynthetic photon flux (PPF) 25 to 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$), preferably a light regime of 16 hours light and 8 hours dark, an air temperature of about 21°C during the day and 18°C at night, a water vapour pressure deficit of about 4.4 g m^{-3} corresponding to a relative humidity (RH) of about 60%-85%, at 400-900 ppm CO_2 and atmospheric O_2 concentration and at atmospheric air pressure (generally 1008 hPa). Water and nutrients may be given drop wise near the stem, or in the form of spray or mist or by ebb and flow.

With the term "bioassay for measuring the capability of growing in continuous light" as used herein, is meant any bioassay which is suitable for determining differences in said symptoms. Preferably, chlorosis is determined using fluorescence measurements of the chlorophyll concentration using a SPAD measuring device Marquard and Tipton 1987, HortScience 22: 1327. Standard bioassay

experimentation conditions, such as stem diameter and plant height measurements, chlorophyll concentration measurements and assays for determining tolerance to continuous light are further specified in the Examples below.

The term “marker-assisted selection”, as used herein, refers to the diagnostic process of identifying, optionally followed by selecting a plant from a group of plants using the presence of a molecular marker as the diagnostic characteristic or selection criterion. The process usually involves detecting the presence of a certain nucleic acid sequence or polymorphism in the genome of a plant.

The term “marker”, as used herein, refers to an indicator that is used in methods for visualizing differences in characteristics of nucleic acid sequences. Examples of such indicators are restriction fragment length polymorphism (RFLP) markers, amplified fragment length polymorphism (AFLP) markers, single nucleotide polymorphisms (SNPs), microsatellite markers (e.g. SSRs), sequence-characterized amplified region (SCAR) markers, cleaved amplified polymorphic sequence (CAPS) markers or isozyme markers or combinations of the markers described herein which defines a specific genetic and chromosomal location.

The term “linked”, as used herein, with reference to markers linked to a trait, refers to a marker the presence of which in the genome of the plant coincides with the presence of the trait. Usually the term refers to a genetic marker that falls within the physical boundaries of a genomic region spanned by at least two markers. For RIL (recombinant inbred lines), such markers having established LOD scores above a certain threshold thereby indicating that no or very little recombination between these markers and the trait locus occurs in crosses; as well as any marker in linkage disequilibrium to the trait locus; as well as markers that represent the actual causal mutations within the trait locus. The term “linked” is used in its broadest sense and indicates that the marker and the gene are located within a continuous DNA sequence of several centiMorgan. The term is used herein with reference to the linkage between markers and phenotype and refers to a distance of preferably less than 20 cM, preferably less than 10 cM, still more preferably less than 6, 5, 4, 3, 2, or 1 cM.

The term “gene”, as used herein, refers to a hereditary unit consisting of a sequence of DNA that occupies a specific location on a chromosome and that contains the genetic instruction for a particular characteristics or trait in an organism. The

term “gene” thus refers to a nucleic acid (for example, DNA or RNA) sequence that comprises coding sequences necessary for the production of an RNA, or a polypeptide or its precursor. A functional polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence as long as the desired activity or functional properties (for example, enzymatic activity, ligand binding, signal transduction, etc.) of the polypeptide are retained. The term “gene” encompasses both cDNA and genomic forms of a gene.

The term ‘yield’ as used herein, refers to the biomass of a tomato plant. Any parameter which is related to the biomass of said plant can be used to determine the yield of a plant. Preferably, said parameter comprises the total leaf area, the shoot dry matter, the shoot fresh mass, the dry mass or the shoot dry mass of said plant.

For optimal growth, fruit vegetables like tomato need 6 hours of darkness per day. This implies that no photosynthesis occurs for 33% per day. If these vegetables could be grown under continuous light, a substantial increase in production would be expected. However, continuous light causes severe problems in these crops: tomato plants, grown in continuous light show a strong increase in chloroplastic starch, leaf chlorosis, necrosis and consequently very poor biomass production; continuous light can even be lethal for the crop.

The present inventors have now discovered tolerant lines in several wild species and in an old cultivar. However, tolerant lines could not be found in any of the modern tomato varieties like Encore, Campari, DRS540, Tourance, Westland, etc.

There are no published reports of tomato varieties exhibiting tolerance to continuous light, save a congress abstract of a miniature tomato cultivar. There is no report on the sensitivity of other tomato species to continuous light, except for the paper of Daskaloff and Ognjanova (Zeitschrift für Pflanzenzuchtung 1965 54: 169-181) where it is claimed that *S. habrochaites*, *S. peruvianum* and *S. racemigerum* (= *pimpinellifolium*) are tolerant to continuous light. It is remarkable that in this publication the focus is on experiments at low light intensities as it is described that the continuous light symptoms are more profound at these lower light intensities. *S. pimpinellifolium* was found to be tolerant by Daskaloff and Ognjanova. In contrast, the present inventors have now found that it is just the opposite and that the continuous light symptoms are more profound at higher light intensities, and *S. pimpinellifolium* was not found to be tolerant.

In the experiments as conducted natural variation in the continuous light symptoms was sought. Several tomato species are clearly tolerant. The symptoms are quantified by measuring the chlorophyll concentration (using e.g a SPAD meter). For some symptomless lines biomass production increased linearly with the increase of available light. Moreover, in addition to a linear increase some lines showed even a further increase in biomass production relative to the light available, i.e. higher than was predicted. The wild tomato donors pennellii LA716, *S. peruvianum* LA1708, *S. habrochaites* LA1777, G1560, Lyc4, *S. chmielewskii* LA1840, *S. neorickii* LA2133, *S. chilense* LA1959 are tolerant to continuous light the same is true for an old variety “Sub arctic plenty”. Tolerance lines do not show deleterious symptoms.

The IL lines, 4 and 5 *S. pennellii* * M82, 42 and 43 *S. neorickii* (LA2133), 5 *S. chilense* (LA1959) are tolerant to continuous light.

The chromosomal location of the tolerance is chromosome 7 for the *S. pennellii*, *S. neorickii*, and *S. chilense* source.

Producing plants with tolerance to continuous light

Plant breeders and in particular seed companies employ elite breeding lines, generally referred to as “elite lines” to provide a constant quality product. The elite lines are the result of many years of inbreeding and combine multiple superior characteristics such as high yield, fruit quality, and resistance to pests, disease, or tolerance to abiotic stress. The average yield of these elite lines is generally much higher than the original wild (landrace) accessions from which many of the modern tomato varieties are descendants. The elite lines can be used directly as crop plant, but are typically used to produce so-called F1 or single-cross hybrids, produced by a cross between two (homozygous or inbred) elite lines. The F1 hybrids thus combine the genetic properties of the two parents into a single plant. An add-on benefit of hybrids is that they express hybrid vigour or heterosis, the poorly understood phenomenon that hybrid plants grow better than either (inbred) parent and show higher yields.

Backcross or pedigree selection is one method by which breeders add desirable agronomic traits to their elite breeding lines. The method involves crossing the breeding line with a line that expresses the desirable trait followed by backcrossing offspring plants expressing the trait to the recurrent parent. As a result,

the selection of an individual as a parent in a breeding program is based on the performance of its forebears. Such methods are most effective in breeding for qualitatively-inherited traits, i.e traits which are present or absent.

Recurrent selection is an alternative breeding method for improving breeding
5 lines and involves systematic testing and selection of desirable progeny followed by recombination of the selected individuals to form a new population. Recurrent selection has proven effective for improving quantitative traits in crop plants. Recurrent selection, however, decreases the rate of broadening genetic basis underlying the various traits in a breeding program, and its potential is therefore
10 limited.

The present inventors discovered that the yield of a tomato plant may be increased by introgressing into an elite breeding line the trait of tolerance to continuous light.

A first method would comprise introgressing the trait from a tomato plant
15 exhibiting tolerance to continuous light, such as a plant of the wild tomato species *S. pennellii*, such as *S. pennellii* LA716, or an offspring plant thereof having said tolerance to continuous light, into a plant of a tomato line of interest. This may for instance be achieved by crossing a plant of a recipient breeding line of *S. lycopersicum* capable of producing commercially valuable fruits, with a plant of a donor line of a
20 tomato species, having tolerance to continuous light. This will result in a situation wherein the gene for tolerance to continuous light is in the genetic background of the tomato line of interest. The establishment of the proper introgression in offspring plants may be monitored by using specific markers as defined herein.

Recombination is the exchange of information between two homologous
25 chromosomes during meiosis. In a recombinant plant, DNA that is originally present on a specific location within the chromosome is exchanged for DNA from another plant (i.e. maternal for paternal or *vice versa*). In order to exchange only the required material, and maintain the valuable original information on the chromosome as much as possible, will usually require two crossover events. The normal way to find such a
30 recombinant is to screen a population of F₂-plants. This population must be of sufficient size in order to detect the rare (low frequency) double recombinants. The frequency of recombination can be expressed in a genetic distance. For instance, a

single recombinant in a 10 cM area can be found with a frequency of 10% (1 centimorgan is defined as 1% recombinant progeny in a testcross).

The present invention also provides methods of producing the plants of the invention using marker assisted selection (MAS). The invention therefore relates to methods of plant breeding and to methods to select plants, in particular tomato plants, particularly cultivated tomato plants as breeder plants for use in breeding programs or cultivated tomato plants for having desired genotypic or potential phenotypic properties, in particular related to producing quantities of valuable tomato fruits, also referred herein to as agronomically desirable plants. Herein, a cultivated plant is defined as a plant being purposely selected or having been derived from a plant having been purposely selected in agricultural or horticultural practice for having desired genotypic or potential phenotypic properties, in particular a plant obtained by inbreeding.

Since the gene can only be properly identified phenotypically when the plant remains healthy when subjected to continuous light, it is of particular advantage that the establishment of the proper introgression in offspring plants may be monitored by using the gene-specific markers as provided herein, either in cis or in trans coupling as explained below. By using marker assisted selection (MAS) or marker assisted breeding (MAB) methods, the skilled person is therefore provided with methods for selecting plants carrying the desired genotype loci and discarding plants lacking the potential of producing progeny tolerant to continuous light.

The present invention thus also provides methods for selecting a tomato plant exhibiting tolerance to continuous light, comprising detecting in said plant the presence of the gene for tolerance to continuous light as defined herein. In a preferred method of the invention for selecting such a plant the method comprises:

- a) providing a sample of genomic DNA from a tomato plant;
- b) detecting in said sample of genomic DNA at least one molecular marker linked to the gene for tolerance to continuous light.

The step of providing a sample of genomic DNA from a tomato plant may be performed by standard DNA isolation methods well known in the art.

The step of detecting a molecular marker (step b) may, in a preferred embodiment, comprise the use of CAPS markers, which constitute a set of bi-directional primers in combination with a restriction enzyme. This allows for the

detection of specific SNPs linked to the trait. Bi-directional means that the orientation of the primers is such that one functions as the forward and one as the reverse primer in an amplification reaction of nucleic acid.

Alternatively, the step of detecting a molecular marker (step b) may in another preferred embodiment, comprise the use of a nucleic acid probe having a base sequence which is substantially complementary to the nucleic acid sequence defining said molecular marker (e.g. said SNP) and which nucleic acid probe specifically hybridizes under stringent conditions with a nucleic acid sequence defining said molecular marker. A suitable nucleic acid probe may for instance be a single strand oligonucleotide of the amplification product corresponding to the marker.

The step of detecting a molecular marker (step b) may also comprise the performance of a unique nucleic acid amplification reaction on said genomic DNA to detect said gene. This can suitably be done by performing a PCR reaction using a pair of marker-specific primers based on the internal or adjacent (up to 500 kilo base) sequence. In a preferred embodiment, said step b) comprises the use of at least one pair of primers defining a marker for said gene (e.g. being complementary to said marker or hybridizing specifically to said marker or allowing polymerase chain extension to occur when bound to said marker), or a pair of primers which specifically hybridize under stringent conditions with the nucleic acid sequence of a marker for said gene.

The step of detecting an amplified DNA fragment having a certain predicted length or a certain predicted nucleic acid sequence may be performed such that the amplified DNA fragment has a length that corresponds (plus or minus a few bases, e.g. a length of one, two or three bases more or less) to the expected length as based on the nucleotide sequence of the genes and markers identified herein. The skilled person is aware that markers that are absent in plants having the introgression as defined herein (donor plants), while they are present in the plants receiving the introgression (recipient plants) (so-called trans-markers), may also be useful in assays for detecting the introgression among offspring plants, although detecting the presence of a specific introgression is not optimally demonstrated by the absence of a marker.

The step of detecting an amplified DNA fragment having the predicted length or the predicted nucleic acid sequence may be performed by standard gel-

electrophoresis techniques, real time PCR, or by using DNA sequencers. The methods need not be described here as they are well known to the skilled person. It should be noted that a marker is usually defined based on its nucleotide sequences in combination with its position relative to other markers on a linkage map.

5

Molecular Markers and genes

Molecular markers are used for the visualisation of differences in nucleic acid sequences. This visualisation is possible due to DNA-DNA hybridisation techniques after digestion with a restriction enzyme (RFLP) and/or due to techniques using the polymerase chain reaction (e.g. STS, microsatellites, AFLP). All differences between two parental genotypes will segregate in a mapping population (e.g., BC₁, F₂) based on the cross of these parental genotypes. The segregation of the different markers may be compared and recombination frequencies can be calculated. The recombination frequencies of molecular markers on different chromosomes is generally 50%.

Between molecular markers located on the same chromosome the recombination frequency depends on the distance between the markers. A low recombination frequency corresponds to a short genetic distance between markers on a chromosome. Comparing all recombination frequencies will result in the most logical order of the molecular markers on the chromosomes. This most logical order can be depicted in a linkage map. A group of adjacent or contiguous markers on the linkage map that is associated with tolerance to continuous light, pinpoints the position of a gene associated with tolerance to continuous light.

The markers identified herein may be used in various aspects of the invention as will now be illustrated. Aspects of the invention are not limited to the use of the markers identified herein. It is stressed that the aspects may also make use of markers not explicitly disclosed herein or even yet to be identified.

In the present invention amplified fragment length polymorphism (AFLP) markers, single nucleotide polymorphisms (SNPs), and insertion deletions (INDELs), microsatellite markers, restriction fragment length polymorphism (RFLP) markers, sequence-characterized amplified region (SCAR) markers, cleaved amplified polymorphic sequence (CAPS) markers or isozyme markers or combinations of these markers might be used.

In general, a gene may span a region of several hundreds to thousands of bases. The plants that have the genetic potential for exhibiting a particular phenotypic trait (tolerance to continuous light) may be traced amongst a population of offspring plants through the observed correlation between the presence of a (string of
5 contiguous) genomic marker(s) and the presence of the phenotypic trait. By providing a non-limiting list of markers, the present invention thus provides for the effective utility of the gene in a breeding program.

It is further important to note that the contiguous genomic markers can also be used to indicate the presence of the gene (and thus of the phenotype) in an individual
10 plant, i.e. they can be used in marker assisted selection (MAS) procedures. In principle, the number of potentially useful markers is limited but may be very large, and the skilled person may easily identify additional markers to those mentioned in the present application. Any marker that is linked to the gene, e.g. falling within the physical boundaries of the genomic region spanned by the markers, wherein between
15 said markers and the gene no or very little recombination occurs in crosses; as well as any marker in linkage disequilibrium to the gene; as well as markers that represent the actual causal mutations within the gene, may be used in MAS procedures.

This means that the markers identified herein, are mere examples of markers suitable for use in MAS procedures. Moreover, when the gene, or the specific trait-
20 conferring part thereof, is introgressed into another genetic background (i.e. into the genome of another plant line), then some markers may no longer be found in the offspring although the trait is present therein, indicating that such markers are outside the genomic region that represents the specific trait-conferring part of the gene in the original parent line only and that the new genetic background has a
25 different genomic organisation. Such markers of which the absence indicates the successful introduction of the genetic element in the offspring are called "trans markers" and may be equally suitable in MAS procedures under the present invention.

Upon the identification of the gene, the gene effect (tolerance to continuous
30 light) is confirmed by determining the tolerance to continuous light of progenies respectively recombinant or segregating for the genes under investigation. Preferably, detecting the presence of a gene of the invention is performed with at least one of the markers for a gene as defined herein. The present invention therefore also relates to a

method for detecting the presence of a gene for tolerance to continuous light as defined herein in tomato by the use of the said markers.

The nucleotide sequence of the genes of the present invention may be resolved by determining the nucleotide sequence of one or more markers associated with said
5 gene and designing internal primers for said marker sequences that may then be used to further determine the sequence the gene adjacent to said marker sequences. For instance the nucleotide sequence of CAPS markers may be obtained by isolating said markers from the electrophoresis gel used in the determination of the presence of said markers in the genome of a subject plant, and determining the nucleotide sequence of
10 said markers by for instance Sanger or pyro sequencing methods, well known in the art.

In embodiments of methods for detecting the presence of a gene in a tomato plant, the method may also comprise the steps of providing an oligonucleotide or polynucleotide capable of hybridizing under stringent hybridization conditions to a
15 nucleic acid sequence of a marker linked to said gene, contacting said oligonucleotide or polynucleotide with nucleic acid of a tomato plant, and determining the presence of specific hybridization of said oligonucleotide or polynucleotide to said nucleic acid.

Preferably said method is performed on a nucleic acid sample obtained (isolated) from said tomato plant, although *in situ* hybridization methods may also be
20 employed. Alternatively, and in a more preferred embodiment, the skilled person may, once the nucleotide sequence of the gene has been determined, design specific hybridization probes or oligonucleotides capable of hybridizing under stringent hybridization conditions to the nucleic acid sequence of said gene and may use such hybridization probes in methods for detecting the presence of a gene of the invention
25 in a tomato plant.

Production of tomato plants exhibiting tolerance to continuous light by transgenic methods

According to another aspect of the present invention, a nucleic acid (preferably
30 DNA) sequence comprising one or more of the genes as defined herein may be used for the production of a tomato plant exhibiting tolerance to continuous light. In this aspect, the invention provides for the use of genes as defined herein or tolerance to continuous light-conferring parts thereof, for producing a tomato plant tolerant to

continuous light as defined herein, which use involves the introduction of a nucleic acid sequence comprising said gene in a suitable recipient plant. As stated, said nucleic acid sequence may be derived from a suitable donor plant. A suitable source according to the present invention for the gene for tolerance to continuous light is

- 5 tomato line *S. pennellii* LA716 (PI 246502 available from the Agricultural Research Service (ARS-GRIN) of the US Department of Agriculture, Washington DC, USA), *S. neorickii*, *S. habrochaites*, *S. lycopersicum*, *S. peruvianum* or *S. chemielewskii*.

Preferably, said gene is selected from the genes present listed in Table 2. The gene sequences thereof are provided in for Table 3.

- 10 Preferably said gene is SL1.00sc07408_313.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_312.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_311.1.1.

- 15 In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_310.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_309.1.1.

In an alternative or additional preferred embodiment, said gene is

- 20 SL1.00sc07408_308.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_307.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_306.1.1.

- 25 In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_305.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_304.1.1.

In an alternative or additional preferred embodiment, said gene is

- 30 SL1.00sc07408_303.1.1.

In an alternative or additional preferred embodiment, said gene is SL1.00sc07408_302.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_301.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_300.1.1.

5 In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_299.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_298.1.1.

In an alternative or additional preferred embodiment, said gene is
10 SL1.00sc07408_297.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_296.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_295.1.1.

15 In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_294.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_293.1.1.

In an alternative or additional preferred embodiment, said gene is
20 SL1.00sc07408_292.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_291.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_290.1.1.

25 In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_289.1.1.

In an alternative or additional preferred embodiment, said gene is
SL1.00sc07408_288.1.1.

In an alternative or additional preferred embodiment, said gene is
30 SL1.00sc07408_287.1.1

In a preferred embodiment, said gene is a gene located between markers 7-20-1
and 7-20-2 as listed in Table 2. Most preferably, said gene is SL1.00sc07408_301.1.1.

The nucleic acid sequence that comprises a gene for tolerance to continuous light, or a tolerance to continuous light-conferring part thereof, may be transferred to a suitable recipient plant by any method available. For instance, the said nucleic acid sequence may be transferred by crossing a plant of line PI 246502 with a selected
5 breeding line which is not tolerant to continuous light or of which the tolerance is to be improved, i.e. by introgression, by transformation, by protoplast fusion, by a doubled haploid technique or by embryo rescue or by any other nucleic acid transfer system, optionally followed by selection of offspring plants comprising the gene for tolerance to continuous light (as assessed by markers) and/or exhibiting tolerance to
10 continuous light. For transgenic methods of transfer a nucleic acid sequence comprising a gene for tolerance to continuous light may be isolated from said donor plant by using methods known in the art and the thus isolated nucleic acid sequence may be transferred to the recipient plant by transgenic methods for plant transformation, for instance by means of a vector, in a gamete, or in any other
15 suitable transfer element, such as a bombardment with a particle coated with said nucleic acid sequence.

Plant transformation generally involves the construction of a vector with an expression cassette that will function in plant cells. In the present invention, such a vector consists of a nucleic acid sequence that comprises a gene for tolerance to
20 continuous light, which vector may comprise such a gene that is under control of or operatively linked to a regulatory element, such as a promoter. The expression vector may contain one or more such operably linked gene/regulatory element combinations, provided that at least one of the genes contained in the combinations confers tolerance to continuous light. The vector(s) may be in the form of a plasmid, and can be used,
25 alone or in combination with other plasmids, to provide transgenic plants that exhibit tolerance to continuous light, using transformation methods known in the art, such as the *Agrobacterium* transformation system.

Expression vectors can include at least one marker gene, operably linked to a regulatory element (such as a promoter) that allows transformed cells containing the
30 marker to be either recovered by negative selection (by inhibiting the growth of cells that do not contain the selectable marker gene), or by positive selection (by screening for the product encoded by the marker gene). Many commonly used selectable marker genes for plant transformation are known in the art, and include, for example, genes

that code for enzymes that metabolically detoxify a selective chemical agent which may be an antibiotic or a herbicide, or genes that encode an altered target which is insensitive to the inhibitor. Several positive selection methods are known in the art, such as mannose selection. Alternatively, marker-less transformation can be used to
5 obtain plants without mentioned marker genes, the techniques for which are known in the art.

One method for introducing an expression vector into a plant is based on the natural transformation system of *Agrobacterium* (See *e.g.* Horsch *et al.*, 1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria that genetically
10 transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*, respectively, carry genes responsible for genetic transformation of the plant. Methods of introducing expression vectors into plant tissue include the direct infection or co-cultivation of plant cells with *Agrobacterium tumefaciens*. Descriptions of
Agrobacterium vectors systems and methods for *Agrobacterium*-mediated gene
15 transfer are provided in US Pat. No. 5,591,616. General descriptions of plant expression vectors and reporter genes and transformation protocols and descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer can be found in Gruber and Crosby, 1993. General methods of culturing plant tissues are provided for example by Miki *et al.*, 1993 and by Phillips, *et al.*, 1988. A
20 proper reference handbook for molecular cloning techniques and suitable expression vectors is Sambrook and Russell, 2001.

Another method for introducing an expression vector into a plant is based on microprojectile-mediated transformation (particle bombardment) wherein DNA is carried on the surface of microprojectiles. The expression vector is introduced into
25 plant tissues with a biolistic device that accelerates the microprojectiles to speeds of 300 to 600 m/s which is sufficient to penetrate plant cell walls and membranes. Another method for introducing DNA to plants is via the sonication of target cells. Alternatively, liposome or spheroplast fusion has been used to introduce expression vectors into plants. Direct uptake of DNA into protoplasts using CaCl_2 precipitation,
30 polyvinyl alcohol or poly-L-ornithine has also been reported. Electroporation of protoplasts and whole cells and tissues has also been described.

Other well known techniques such as the use of BACs, wherein parts of the tomato genome are introduced into bacterial artificial chromosomes (BACs), *i.e.*

vectors used to clone DNA fragments (100- to 300-kb insert size; average, 150 kb) in *Escherichia coli* cells, based on naturally occurring F-factor plasmid found in the bacterium *E. coli* may for instance be employed in combination with the BIBAC system to produce transgenic plants.

5 Following transformation of tomato target tissues, expression of the above described selectable marker genes allows for preferential selection of transformed cells, tissues and/or plants, using regeneration and selection methods now well known in the art.

10 Production of tomato plants exhibiting tolerance to continuous light by non-transgenic methods

 In an alternative embodiment for producing a tomato plant exhibiting tolerance to continuous light, protoplast fusion can be used for the transfer of nucleic acids from a donor plant to a recipient plant. Protoplast fusion is an induced or
15 spontaneous union, such as a somatic hybridization, between two or more protoplasts (cells of which the cell walls are removed by enzymatic treatment) to produce a single bi- or multi-nucleate cell. The fused cell, that may even be obtained with plant species that cannot be interbred in nature, is tissue cultured into a hybrid plant exhibiting the desirable combination of traits. More specifically, a first protoplast can be
20 obtained from a tomato plant of accession PI 246502. A second protoplast can be obtained from a second tomato plant variety, preferably a tomato line that comprises commercially valuable characteristics, such as, but not limited to disease resistance, insect resistance, valuable fruit characteristics, etc. The protoplasts are then fused using traditional protoplast fusion procedures, which are known in the art.

25 Alternatively, embryo rescue may be employed in the transfer of a nucleic acid comprising the gene as described herein from a donor plant to a recipient plant. Embryo rescue can be used as a procedure to isolate embryo's from crosses wherein plants fail to produce viable seed. In this process, the fertilized ovary or immature seed of a plant is tissue cultured to create new plants.

30 The present invention also relates to a method for improving the tolerance to continuous light of a plant of a tomato breeding line, comprising the steps of:

a) crossing a plant of a tomato breeding line with a plant of tomato line PI 246502 or an offspring plant thereof harbouring the gene for tolerance to continuous light as described herein;

5 b) selecting a progeny tomato plant resulting from said crossing having an introgression from tomato accession PI 246502 or an offspring plant thereof associated with tolerance to continuous light;

c) selfing and/or backcrossing said progeny tomato plant selected in step (b) using said tomato breeding line as a recurrent parent;

10 d) selecting a progeny tomato plant resulting from the selfing or backcrossing in step (c) having an introgression from tomato accession PI 246502 or an offspring plant thereof associated with tolerance to continuous light,

e) repeating said steps of selfing and/or backcrossing and selection of steps (c) and (d) to provide a plant of a tomato breeding line essentially homozygous for said introgression,

15 wherein preferably at least one selection as performed in steps (b) or (d) is performed by marker-assisted selection.

In a preferred embodiment of such a method, said tomato breeding line is an elite line.

20 In an alternative preferred embodiment of the above method, the marker-assisted selection procedure comprises the selection for at least one marker as exemplified in the Examples below.

The introgression of the nucleic acid sequence comprising the gene for tolerance to continuous light as described herein may suitably be accomplished by using traditional breeding techniques. The gene is preferably introgressed into commercial tomato varieties by using marker-assisted selection (MAS) or marker-assisted breeding (MAB). MAS and MAB involves the use of one or more of the molecular markers for the identification and selection of those offspring plants that contain one or more of the genes that encode for the desired trait. In the present instance, such identification and selection is based on selection of the gene of the present invention or markers associated therewith. MAS can also be used to develop near-isogenic lines (NIL) harboring the gene of interest, or the generation of gene isogenic recombinants (QIRs), allowing a more detailed study of each gene effect and is also an effective method for development of backcross inbred line (BIL) populations.

Tomato plants developed according to this embodiment can advantageously derive a majority of their traits from the recipient plant, and derive tolerance to continuous light from the donor plant.

Crossing can be achieved by mechanically pollinating the female flower of one parent plant with pollen obtained from male flowers of another parent plant.

As discussed briefly above, traditional breeding techniques can be used to introgress a nucleic acid sequence encoding a gene for tolerance to continuous light into a recipient tomato plant requiring tolerance to continuous light. In one method, which is referred to as pedigree breeding, a donor tomato plant that exhibits tolerance to continuous light and comprising a nucleic acid sequence encoding for the gene associated with tolerance to continuous light as defined herein is crossed with a recipient tomato plant (preferably a plant of an elite line) that exhibits agronomically desirable characteristics, such as, but not limited to, disease (e.g. TMV) resistance, insect resistance, valuable fruit characteristics, etc., but which is not tolerance to continuous light, or which requires improvement of such tolerance. The resulting plant population (representing the F_1 hybrids) is then self-pollinated and set seeds (F_2 seeds). The F_2 plants grown from the F_2 seeds are then screened for tolerance to continuous light. The population can be screened in a number of different ways.

First, the population can be screened using a bioassay wherein the plant is grown under continuous light for a prolonged period of time and its health or productivity (phenotype) is inspected. Second, marker-assisted selection can be performed using one or more of the hereinbefore-described molecular markers to identify those progeny that comprise a nucleic acid sequence encoding a gene for tolerance to continuous light as defined herein. Other methods, described above by methods for detecting the presence of a gene may be used. Also, marker-assisted selection can be used to confirm the results obtained from the phenotype scores, and therefore, several methods may also be used in combination.

Inbred tomato plant lines exhibiting tolerance to continuous light can be developed using the techniques of recurrent selection and backcrossing, selfing and/or dihaploids or any other technique used to make parental lines. In a method of recurrent selection and backcrossing, the tolerance to continuous light-conferring genetic element as disclosed herein can be introgressed into a target recipient plant (the recurrent parent) by crossing the recurrent parent with a first donor plant, which

differs from the recurrent parent and is referred to herein as the "non-recurrent parent". The recurrent parent is a plant of which the tolerance is to be improved and possesses agronomically desirable characteristics, such as, but not limited to disease resistance, insect resistance, valuable fruit characteristics, etc. The non-recurrent, or donor, parent may suitably be a plant of suitable donor line as described herein which comprises a nucleic acid sequence that encodes for a gene that confers tolerance to continuous light. Alternatively, the donor parent can be any plant variety or inbred line that is cross-fertile with the recurrent parent and has acquired the gene for tolerance to continuous light in an earlier cross with a plant of the said donor line.

10 The progeny resulting from a cross between the recurrent parent and non-recurrent parent is backcrossed to the recurrent parent. The resulting plant population is then screened for the desired characteristics, which screening may occur in a number of different ways. For instance, the population can be screened using phenotypic screens as described herein. As an alternative to phenotypic assays, marker-assisted selection

15 (MAS) can be performed using one or more of the hereinbefore described molecular markers, hybridization probes or polynucleotides to identify progeny that comprise a nucleic acid sequence encoding the gene responsible for tolerance to continuous light.

Following screening, the F₁ hybrid plants that exhibit a tolerant phenotype or, more preferably, genotype and thus comprise the requisite nucleic acid sequence encoding for a gene conferring tolerance to continuous light are then selected and

20 backcrossed to the recurrent parent for a number of generations in order to allow for the tomato plant to become increasingly elite. This process can be performed for two to five or more, such as 6, 7 or 8 generations. In principle the progeny resulting from the process of crossing the recurrent parent with the non-recurrent parent are

25 heterozygous for one or more genes that encode for tolerance to continuous light.

In a preferred embodiment, a method of introducing a desired trait into a hybrid tomato variety comprises the steps of:

(a) crossing an inbred tomato parent with another tomato plant that comprises one or more desired traits, to produce F₁ progeny plants, wherein the desired trait is tolerance to continuous light as conferred by the gene from any suitable donor plant of the said trait as defined herein, or an offspring plant thereof;

30

(b) selecting said F₁ progeny plants that have the desired trait to produce selected F₁ progeny plants, preferably using molecular markers as defined herein;

(c) backcrossing the selected progeny plants with said inbred tomato parent plant to produce backcross progeny plants;

(d) selecting for backcross progeny plants that have the desired trait (tolerance to continuous light) and morphological and physiological characteristics of said inbred tomato parent plant, wherein said selection preferably comprises the isolation of genomic DNA and testing said DNA for the presence of at least one molecular marker for the gene as defined above;

(e) repeating steps (c) and (d) two or more times in succession to produce selected third or higher backcross progeny plants;

(f) optionally selfing selected backcross progeny in order to identify homozygous plants;

(g) crossing at least one of said backcross progeny and/or selfed plants with another inbred tomato parent plant to generate a hybrid tomato variety with the desired trait and all of the morphological and physiological characteristics of hybrid tomato variety when grown in the same environmental conditions.

As indicated, the last backcross generation may be selfed in order to provide for homozygous pure breeding (inbred) progeny exhibiting tolerance to continuous light. Thus, the result of recurrent selection, backcrossing and selfing is the generation of lines that are genetically homozygous for the genes associated with tolerance to continuous light as well as other genes associated with traits of commercial interest.

It should be noted that heterozygous plants having the gene for tolerance to continuous light may also be of interest as intermediate products, and such plants are therefore also an aspect of the present invention.

Tomato plants and seeds

The goal of plant breeding is to combine various desirable traits in a single variety or hybrid. For commercial crops, these traits may include resistance to diseases and insects, tolerance to heat and drought, reducing the time to crop maturity, greater yield, and better agronomic quality. Uniformity of plant characteristics such as germination, growth rate, maturity, and plant height may also be of importance.

Commercial crops are bred through techniques that take advantage of the plant's method of pollination. A plant is self-pollinated if pollen from one flower is transferred to the same or another flower of the same plant. A plant is sibling mated when individuals within the same family or line are used for pollination. A plant is
5 cross-pollinated if the pollen comes from a flower on a different plant from a different family or line.

Plants that have been self-pollinated and selected for type for many generations become homozygous at almost all gene loci and produce a uniform population of true-bred progeny. A cross between two different homozygous lines
10 produces a uniform population of hybrid plants that may be heterozygous for many gene loci. A cross of two plants each heterozygous at a number of gene loci will produce a population of heterogeneous plants that differ genetically and will not be uniform.

The development of a hybrid tomato variety in a tomato plant breeding
15 program involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, individually breed true and are highly uniform; and (3) crossing a selected inbred line with an unrelated inbred line to produce the hybrid progeny (F1). After a sufficient amount of
20 inbreeding successive filial generations will merely serve to increase seed of the developed inbred. Preferably, an inbred line should comprise homozygous alleles at about 80% or more of its loci.

An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid created by crossing a defined pair of inbreds will always be the
25 same. Once the inbreds that create a superior hybrid have been identified, a continual supply of the hybrid seed can be produced using these inbred parents and the hybrid tomato plants can then be generated from this hybrid seed supply.

Using the methods as described above, the skilled person will be able to produce the required inbred lines and from those produce the commercial (F1) hybrid
30 seeds by crossing said inbred lines.

The present invention will now be explained in more detail by way of the following non-limiting Examples.

EXAMPLES

Example 1 tolerant donors

Different accessions of wild species (lines) were sown in a growth chamber with 16 hours of light per day. Two weeks after sowing, the seedlings of each line were randomly divided in two groups, one group continued to grow in 16 hours of light per day while the other was grown in another growth chamber under continuous light. After 3-4 weeks symptoms of the sensitive lines were clearly visible, while absent in the tolerant lines (Figure 1). The typical symptoms of a sensitive line are leaf chlorosis, necrosis and poor biomass production which was absent in the tolerant genotypes shown in figure 1.

Example 2 quantifying the symptoms and optimise protocol

In the same experiment as experiment 1 the sensitive lines showed at first light green leaves with inter-vein chlorosis, followed by severe chlorosis (Figure 2). In severe non tolerant plants chlorosis is followed by necrosis, decreased leaf development decreased stem development and even plant mortality. Besides the clear visible symptoms of the sensitive line in continuous light the symptoms can also be quantified by measuring the chlorophyll concentration. The chlorophyll concentration can be measured chemically but also spectroscopically with a SPAD meter (SPAD-502, Konica Minolta, Nieuwegein, Netherlands). Two sensitive lines (Moneyberg and Westland) were grown as described in experiment 1, except that also the light intensity was a experimental factor from two weeks after sowing. The plants were placed in low light intensities ($16 \mu\text{mol m}^{-2} \text{s}^{-1}$ =LL) moderate light $131 \mu\text{mol m}^{-2} \text{s}^{-1}$ (ML) or relative high light $230 \mu\text{mol m}^{-2} \text{s}^{-1}$ (HL). Quantification of a part of the visible symptoms (degreening and chlorosis) by measuring the chlorophyll concentration by a SPAD meter was performed with plants grown at $131 \mu\text{mol m}^{-2} \text{s}^{-1}$ (ML) or $230 \mu\text{mol m}^{-2} \text{s}^{-1}$ (HL). The resolution to detect differences between continuous and non-continuous light was very difficult or impossible at very low light intensities ($16 \mu\text{mol m}^{-2} \text{s}^{-1}$ =LL) (Figure 3). The visible symptoms developed earlier in time at $230 \mu\text{mol m}^{-2} \text{s}^{-1}$ compared to $131 \mu\text{mol m}^{-2} \text{s}^{-1}$. In the same experiment also the stem diameter and plant height was decreased in these non tolerant lines (Figure 4). Not only a higher light intensity resulted in a faster induction of symptoms also a prolonged exposure to continuous light did increase the symptoms in sensitive lines

(Figure 7, 9). In several lines the symptoms did appear later than after 3 to 4 weeks in continuous light.

In conclusion the continuous light symptoms can be quantified by measuring the chlorophyll concentration in the leaves but also by measuring the stem diameter or height. The quantitative tolerance trait can be measured qualitative by increasing the light intensity or increasing the duration of the continuous light. This makes it easier to map the tolerance.

Example 3 tolerant lines from *neorickii* (LA2133)

10 A segregating population between *neorickii* and a sensitive *esculentum* (e.g. Moneyberg) showed large natural variation in the chlorophyll concentration as measured with a SPAD meter (Figure 5). In figure 6 two tolerant lines (42 and 43) are shown, which have been in continuous light for 90 days.

15 Example 4 tolerant lines from *chilense* (LA1959)

A segregating population between *chilense* and a sensitive *esculentum* (e.g. Moneyberg GT) showed large natural variation in the chlorophyll concentration as measured with a SPAD meter (Figure 7). In figure 8 one tolerant line (5) is shown, which have been in continuous light for 60 days.

20

Example 5 tolerant lines from *pennellii* (LA716*M82)

A segregating population between *pennellii* and M82 was used and only lines with a chromosomal introgression from *pennelli* into M82 on chromosome 7 were used. Natural variation in the chlorophyll concentration as measured with a SPAD meter was identified (Figure 9). Two tolerant lines (4 and 5) are found, which have been in continuous light for 53 days.

25

Example 6 Continuous light tolerance loci on chromosome 7

The genetic information from the 6 different mapping population it is clear that the loci for continuous light tolerance is located on chromosome 7 (Figure 10). Moreover, by combining the information of all populations it is shown that the loci is located between marker “7-19” and “7-23” that is between 64 and 99,6 cM. The

30

sequence in formation from these 2 markers and 23 additional ones which are also linked to the trait are shown in figure 11.

Example 7 Detection of marker 7-20-1

- 5 To detect the G/T polymorphism of marker 7-20-1, a TaqMan assay (Applied biosystems) was used. The assay employs two universal primers (a forward primer having the sequence 5'-GAATTGGCTTTATGTATTTGGAATCCTTGT-3' and a reverse primer having the sequence 5'-CCCATCTAGGACCCCTGCAT-3') to amplify a fragment of 76 bp and two allele specific probes. A probe having the sequence (5'-
- 10 TTTACTGTATTTTCTTTTCTCTCCA-3' (*S. lycopersicum* (recurrent)) specifically detects the *S. lycopersicum* allele (The "T" SNP) and carries a VIC reporter fluorescent dye. A second probe having the sequence 5'-TTTACTGTATTTTCTTTTCTCTCCA-3' (*S. neorickii* LA2133 (donor)) detects the allele from *S. neorickii* LA 2133 (the "G" SNP) and carries a 6-carboxyfluorescein (FAM) dye. Amplification of one of the two or both alleles can
- 15 be monitored using a real-time PCR (polymerase chain reaction) machine which measures the level of fluorescence of the respective dyes. DNA samples that only contain the *S. lycopersicum* allele will emit a VIC signal while samples that contain the *S. neorickii* allele will emit the FAM signal. Samples that contain both alleles will emit both signals.
- 20 The assay was validated on DNA samples of elite sensitive lines and CLT donor LA2133. All other CLT donor lines had the same allele. The assay was also validated on the 46 introgression lines that were derived from LA2133 and "Moneyberg". The marker showed presence of the "G" allele in LA2133 and introgression lines LA2133-42 and LA2133-43. All other tested lines showed presence
- 25 of the "T" allele

Example 8: Fine mapping of continuous light tolerance trait.

- To identify closer linked markers for the continuous light tolerance trait fine mapping in the LA2133 background was performed. Lines LA2133-42 and LA2133-43
- 30 were backcrossed with their sensitive recurrent parent. Both F1s were selfed and the F2 offspring was genotyped for recombination with flanking markers 7-10 and 7-25. In total 770 and 616 plants were genotyped for LA2133-42 and LA2133-43 respectively. In total 40 lines were selected to generate a F3 from which 15 lines were

controls. 22 Seeds per F3 family were sown in 2 blocks of 11 seeds. A total of 18 F3 families were screened. From several F3 families the germination was too low to yield sufficient (≥ 13) plants for phenotyping the continuous light tolerance trait.

Mapping indicated that the position of the continuous light locus is located
 5 between markers 7-17 and 7-20 (figure 12). Marker 7-19 was monomorphic in the tested LA2133 populations.

New markers were developed between marker 7-17 and 7-20 and are named 7-20-1, 7-20-2 and 7-20-3 (figure 11). Using the new markers on the same phenotyped populations revealed that the position of the continuous light locus is located between
 10 markers 7-20-1 and 7-20-2 (figure 13).

The three newly developed markers were tested on DNA samples of complete F2 families to generate a new linkage map of the CLT locus (continuous light trait locus). The linkage map EXPEN2000 and the generated linkage map of LA2133-42 showed some differences on the location of the new SNP markers (See Table 1 and
 15 Fig. 10). The LA2133-42 map shows the preferred order.

Table 1

Marker	LA2133-42	EXPEN2000
7-20-1	43,68 cM	75 cM
7-20-2	44,35 cM	72 cM
7-20-3	44,46 cM	73 cM
7-20	46,18 cM	73 cM
7-21	46,74 cM	78 cM

Based on the mapping in LA2133-42 highly preferred markers linked to the
 20 continuous light trait, span a locus of about 0.67 cM.

Example 9: Genes at the CLT locus.

A BLAST search to the tomato genome (version 1.0; released December 2009) revealed that the markers of the CLT locus are preferably located on scaffold 07408
 25 (SL1.00sc07408 position:1975288 through 2058583; see <http://solgenomics.net/>). The region between the markers 7-20-1 and 7-20-2 spans 83296 bp and contains 15 genes

(ITAG1 annotation). Table 2 lists all genes at this locus and the available TaqMan markers. Table 3 lists all sequences of these genes. One of the genes

(SL1.00sc07408_301.1.1 (<http://mips.helmholtz-muenchen.de/plant/tomato/searchisp/index.jsp>)) is the LHCb3 Light-Harvesting

- 5 Chlorophyll B-Binding Protein 3 structural molecule". This gene encodes a protein that - in one preferred embodiment of the invention - plays a crucial role in the continuous light trait. Plants of the present invention therefore in a preferred embodiment of the invention comprise this gene. The other genes in this locus can also be used in aspects of this invention.

Table 2. Genes in the CLT locus

Name	Type	Description	Size	Start	Stop	Strand
7-18			100	2878606	2878726	forward
<u>SL1.00sc07408.313.1.1</u>	transcript_ITAG	late embryogenesis abundant protein-related / LEA protein-related (AHRD V1 ***NG AT5G54370.1); contains Interpro domain(s) IPR009646 Root cap	573	2093160	2093732	forward
<u>SL1.00sc07408.312.1.1</u>	transcript_ITAG	LDL2 LSD1-LIKE2 amine oxidase/ electron carrier/ oxidoreductase (AHRD V1 **-.NG AT3G13682.1); contains Interpro domain(s) IPR002937 Amine oxidase	6585	2093055	2086471	reverse
<u>SL1.00sc07408.311.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1); contains Interpro domain(s) IPR009651 Aluminium resistance	7561	2082437	2074877	reverse
<u>SL1.00sc07408.310.1.1</u>	transcript_ITAG	Transmembrane protein 184C (AHRD V1 ***NG Q5RET6); contains Interpro domain(s) IPR005178 Protein of unknown function DUF300	3624	2073719	2070096	reverse
<u>SL1.00sc07408.309.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1)	4836	2069193	2064358	reverse
<u>SL1.00sc07408.308.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1)	2919	2061327	2064245	forward
<u>SL1.00sc07408.307.1.1</u>	transcript_ITAG	AGD5 ARF-GAP domain 5 ARF GTPase activator/ DNA binding / zinc ion binding (AHRD V1 *. *G AT5G54310.1); contains Interpro domain(s) IPR001164 Arf GTPase activating protein	6889	2060607	2053719	reverse
7-20-1		Flanking marker CLT	100	2057927	2058583	reverse
<u>SL1.00sc07408.306.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1); contains Interpro domain(s) IPR008480 Protein of unknown function DUF761, plant	1950	2050327	2048378	reverse
<u>SL1.00sc07408.305.1.1</u>	transcript_ITAG	cytochrome c biogenesis protein family (AHRD V1 *. *NG AT5G54290.1); contains Interpro domain(s) IPR003834 Cytochrome c assembly protein, transmembrane region	6176	2040973	2047148	forward
<u>SL1.00sc07408.304.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1)	2874	2034960	2037833	forward
<u>SL1.00sc07408.303.1.1</u>	transcript_ITAG	Myosin-Va (AHRD V1 ***NG Q02440); contains Interpro domain(s) IPR001609 Myosin head, motor region	8186	2030449	2022264	reverse
<u>SL1.00sc07408.302.1.1</u>	transcript_ITAG	ATM2 ARABIDOPSIS THALIANA MYOSIN 2 motor (AHRD V1 ***NG AT5G54280.2); contains Interpro domain(s) IPR000048 IQ calmodulin-binding region	1832	2022005	2020174	reverse
<u>SL1.00sc07408.301.1.1</u>	transcript_ITAG	LHCB3 LIGHT-HARVESTING	1633	2018017	2019649	forward

		CHLOROPHYLL B-BINDING PROTEIN 3 structural molecule (AHRD V1 ***G AT5G54270.1); contains Interpro domain(s) IPR001344 Chlorophyll A-B binding protein					
<u>SL1.00sc07408_300.1.1</u>	transcript_ITAG	Dynein light chain 1 cytoplasmic-like protein (AHRD V1 *.*NG B0X7V2_CULQU); contains Interpro domain(s) IPR001372 Dynein light chain, type 1 and 2	1195	2015289	2016483	forward	
<u>SL1.00sc07408_299.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1); contains Interpro domain(s) IPR010634 Protein of unknown function DUF1223	2477	2013512	2011036	reverse	
<u>SL1.00sc07408_298.1.1</u>	transcript_ITAG	ATMAMI ARABIDOPSIS THALIANA MEMBRANE-ASSOCIATED MANNITOL-INDUCED structural molecule (AHRD V1 *.*G AT5G54110.1); contains Interpro domain(s) IPR008962 PapD-like	3392	2006924	2010315	forward	
<u>SL1.00sc07408_297.1.1</u>	transcript_ITAG	Wound induced protein (AHRD V1 ***NG B6SKC8_MAIZE)	461	2001724	2002184	forward	
<u>SL1.00sc07408_296.1.1</u>	transcript_ITAG	Ubiquitin carboxyl-terminal hydrolase (AHRD V1 ***G A7P1D3_VITVI); contains Interpro domain(s) IPR016652 Ubiquitinyl hydrolase	9792	1988575	1998366	forward	
<u>SL1.00sc07408_295.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1)	1254	1987084	1985831	reverse	
<u>SL1.00sc07408_294.1.1</u>	transcript_ITAG	Unknown Protein (AHRD V1)	1308	1979129	1977822	reverse	
<u>SL1.00sc07408_293.1.1</u>	transcript_ITAG	universal stress protein USP family protein (AHRD V1 *.*NG AT4G27320.1); contains Interpro domain(s) IPR006016 UspA	3278	1975876	1972599	reverse	
7-20-2		Flanking marker CLT	100	1975288	1975845	forward	
<u>SL1.00sc07408_292.1.1</u>	transcript_ITAG	PBP1 PINOID-BINDING PROTEIN 1 calcium ion binding / protein binding (AHRD V1 *.*G AT5G54490.1); contains Interpro domain(s) IPR011992 EF-Hand type	668	1969824	1970491	forward	
<u>SL1.00sc07408_291.1.1</u>	transcript_ITAG	Serine/threonine kinase-like protein (AHRD V1 ***G Q0WPA4_ARATH); contains Interpro domain(s) IPR002290 Serine/threonine protein kinase	3701	1962690	1966390	forward	
<u>SL1.00sc07408_290.1.1</u>	transcript_ITAG	ARK3 A. THALIANA RECEPTOR KINASE inase/ transmembrane receptor protein serine/threonine kinase (AHRD V1 *.*G AT4G21380.1); contains Interpro domain(s) IPR002290 Serine/threonine	3742	1957638	1961379	forward	

		protein kinase				
<u>SL1.00sc07408_289.1.1</u>	transcript_ITAG	protein kinase family protein (AHRD V1 *-*NG AT4G11470.1); contains Interpro domain(s) IPR002290 Serine/threonine protein kinase	7051	1953626	1946576	reverse
<u>SL1.00sc07408_288.1.1</u>	transcript_ITAG	ferredoxin-related (AHRD V1 ***NG AT1G02180.1)	2040	1945028	1942989	reverse
7-20-3			100	1943149	1943396	reverse
<u>SL1.00sc07408_287.1.1</u>	transcript_ITAG	protein kinase family protein (AHRD V1 ***G AT4G23150.1); contains Interpro domain(s) IPR002290 Serine/threonine protein kinase	1459	1940509	1941967	forward
7-20			100	1686186	1686306	reverse
7-21			100	1480300	1480420	reverse
7-25			100	464589	464709	reverse

Table 3

5	<u>SL1.00sc07408_313.1.1</u>					
10	TGCATGCTT TCTCAGTGGG GGCTACCAAG GCAGAGTCCT GSGACGACGA AGTGTATCAT CTGAAATCTT AITATAAAGG AAAAGAGTTA GGCTIACCAG AAGGATAIAC ATCCATATGG GAATCTTCIG AAAGCGGCTAT CAAGGTACGA AGAAGTGCAG ACAAGACCGG TGCCTTATC ACACIACCAG AGGTAGCAGA AATTICAGTA AATGAGIAC CIATACCAAA GGAAGATGAC AGGATCCACA ACIATCAGIT ACCITCIGAT GACIGCTTCG CTCACCTTGA GGTGCAGTTC AGATCTATG GCCTCTCAAC GAAGGTTGAA GCGTTCCTG GCGGTACGTA CCAGCCAGAC TTCAAAAATC CAGCAAAACC AGTGTGTGCA GTGCTGTAG TGGAGGTTGA AAGATGTAC AGAAGTTCAA CCGTCTTGTG TCCAAAAGT AACCTCTGTA TCTTCTCTOC AGCTGGAGTT TCTGAAGAA CAGACCCCTT AGTCATGGAT TATGGGACTT TAGATTGTAC TGGAGGATCA AGCGGTGGCC ACGGAATAAT TTGCAGAAAA TGA					
15	<u>SL1.00sc07408_312.1.1</u>					
20	ATGGAAATCC CAATTCAGG GGTTCAGTC TCAGGTAGAC CTGTGAGAAG GAGGTTCGCT TCTAGGAATT ATGATCAGAC TGTGATGGAT AAAATCATAG ATGAGCAGIT GGTAGTCCG GTGGGGAAGA AGAATTGAAC GAAGAAGAT TTGGAGAAAG AACTGAAA AGAGGCCCTG AITGCTCTT CTTTGGGCTT CCAATTGAT GACCTTCTTG AAGAAGAAA AAAGGCTTGA GTTGTAAAGT AATGTGATGG GAAGAGACAA AACGATTACA TCGTGTGAG AAATCATATT CTGCAAAAT GGAGGGAGAA TGTCTATATT TGGCTGAACA AAGGAAGAT AAGGGAATTT ATAGTGTGCT AGTATGACA ITTGGTAGCC ATAGCAIATG AITTTCTTTT AAGTAACGGG TATATAAATT TCGGGGTTTC ATCATCATTT GATCTAATC TTCCTGAGGA ACCTAGAGAA GGGTCTGTAA TTAICATTTG TCGTGGACTC GCTGGTTTGG CIGCAGCAAG GCAACTGATG GCTTGTGGAT TTAAGGTAA TGTCTTGA AAAGAGGAAC GACCTGGAGG GAGAGTTTAT TCTGAAGAAA TGGGATGGAA GGGAAAGTTT GGTGCTGTGG AICTTGTGG CAGCGTTATA ACTGGTATCC ATGCCAATCC TTGGGAGATT TTGGCTAGAC AACCTTCCAT TCGGCTTCAC AGTGTTAGAG ATAGTGTTC ITTATACAA COTGATGTTG CICTGTTGA TTCAGTAGTT GATTCACAG TTGAACTCAT TTCAATAAG CTACTGACAA AAGTTGCTGA ACTACGAAA ATCGTAAGTG GATTGGCTAC TGAATGTCTG TTAGGCTCCG TTTTGGAGAC ACTTAGACGA TTATATTGTG TGGCTAAAC TAAAGAGGAG AAGCAACTTC TGCATGGCA TTTTGCAAAC ITGGAAATAG CAATGTCTGG AIGCTCTCTG GAATCTCTG CTGCTACTG GATCAGGAC GATCCTTATG AAATGGATGG TGAATCTGT ITTCTGTG TGSGAAATCG AGCIATGAIC AGGSCATTTT GTAAAGGAGT TCCATATATC TATGAAAGA CTGTTCAGAC AATTAAGTAT GGAATGAAAG GAGTTGAGGT CATTCCTGG GACCAACTTT TCCGGCAGA CATGGTCCIA TGTACTGTTC CTTTGGGWT ACTAAAGAA AGATCAATTA GATTGAAAC AGATTAACCT GAGAAGAAGC TTGAAGCTAT TGATAGGCTA GGATTTGCTG TCTGAATTA AGTGGCAAG GTATTTCCTC AIAITTTTITG GGGCGAGAC TTGATACCT TCGGTTCGCT CAACCATCAT AGCCACAGAC GAGCAGAGTA CTCTTATTT TACGTTTACC ATACTGTTC TGGGGTCCCA GTACTTATG CACTGTGTC TGGTACGCTT GCTCACTTT TCGAAAGCAC AGATCCGTC ACCTTAAATA ATCGAGTAT TAACATCTC AAAGCATTT ATGAGCAAAA GGGATTAAGC GGTGCTGATC CIATACATC CATATGCAAC AATGGGGAA GTGATCCCTT TTGCTTTGGC TCATATTCAC ATGTTCTGTG TCAGTCACTT GGCAGTAT ATGACATACT TCCAGAAA CTGCGAGCTC GGTGTTTCTT TGTGGAGAG GCTACCATC GACAACATCC AGCCACCATG CATGAGGCT ATTCTAGTGG CTTAAGAGAA GCCTCTCACA TTTCCCAATC CATGAAGCG AGGCAAAACA ATCCAGGAG AACTGTATCA AAGAATCTTC GACCAAGCAA TGATACATG GAAGAGTTGT TTGAAGAGCC AGATCTAGCA ITGCGGAGAT TGTATTTGT ATTGAATCC CACACTGTG AITCIAATC TTTAGGACTG ATGAGATTA CTTTGGGAAA ATCCAACGAC GAGCTTAATT CAGAAGAGGT AGACAATAAG CCTCAACATT TATTAATATC					
45						

5	ATCAGTCGAG	CTTTATACAG	TTGTCCTCG	TGAGCAAGCA	CATGAGCIGC	AGTTGGTGAA	GGAGGGAAAC
	AAATGGCAAT	TGTTAGATTT	GGTTGAAGGT	CTTGGGTTAA	AGTTAGTGGG	AGCGAATGGA	CTAGGAGTTT
	AAGGCCATAC	TTTGGCTGCT	AAAATTTTAA	AAGGIAGAAG	GAGTCGAAGC	TATAGTCCCA	AGCAGAAAGC
	AGGCGAAAGT	AGCAGTCAGT	ATAATTTGTC	CATCTGGCAG	CAGAGTGCTA	AAAAGGATC	GATGAAAGCC
	CGTTTTCGAT	CTGGTTTGGG	AGTTTCAAGC	CTTCTTACIC	TGTCGCTGTC	TTTATGTTGT	GGAAATTTT
	CTAATAGCTT	GTGCTTTGTA	CTCCTAGATA	GTCGAAACTT	TCTTGTAGGC	TAA	
10	SL1.00sc07408_311.1.1						
	AAGGCGGCTT	TATGCTGCGG	CACCTCTGCT	TATCTTACCG	ATCTCTGAG	AGTAACTTCT	GCTAAAGCTG
	CAGTGGCTTC	GAGCTCTCGA	GTTCAGTTTC	CICAGCTGCA	TCATTCGGAT	TCCCGTTTTC	TTCCTCAGGT
	TAAATAGGCT	GTTCATCTTT	TCTCAGAAAG	GTTCAGAGAA	GTTCAGCAAT	TAGTGGCAGC	CAALACTTGG
	CGAGTTTTCG	GAGCTTTCGA	AAGGGTCAAG	GTGGGGTCTC	ATCAGTTTTC	TGGTAGCACT	GGCTAAGGTC
	ATGAAGAAGC	TGGTGGGCTT	GAGGCTTTTC	ACCAGGCTTT	TCTAGAAAT	GTTCGTGGCG	AGTCTGCAAT
15	TGTTGCAICA	CAGTTCTTCT	CAGGTACICA	TGCAATCACT	TGTCCTTAT	TGCTTTTCTT	AAGGCCAGGG
	GATGAGTTGT	TAGCGATAGC	TGTCGACCTT	TATGATACTC	TGGAGGAAAT	AAATGGAAAT	AGGGACTCTG
	GTCGATTCGG	TTCCTTGAAA	GATTTTGGAG	TAGAAACCG	GGAAATCCCA	CTTCGAGAGG	ATGGCGGGCT
	TGATTTGGAT	GCACCTTAAA	CCTCTATAAG	ACCTCACACT	AAGTGTGCAC	TCAATCAGAG	ATCATGTGGT
	TATCTTTGGC	GTGCGCTTTT	GAGTGTCACT	GAGATAGGTC	GAGCAATGTA	TATATCAAG	AIGCAGAACCC
20	CAGGCTGCAI	GGTCATGCTA	GATAACTGCT	ATGGTGAAT	TGTTGAGCAC	ATCGAACCTC	CTAGGGTGGG
	TGCTGACCTA	ATTGCCGGA	GTTCATTAAT	AAATCCGGAT	GGAAACGATT	CACATGTCG	TGGAATGTT
	GGGGAAGGCA	AAAATGCGT	AGAAAGCAGC	GGTGGCGCT	TCTCGGCTCC	AGGACTTGA	GTTCATTTGG
	GTTCATCCCT	TGGTGAATTA	ATGAGAACTT	TATTTTCAAG	TATATCTCTC	TCACTCTCAA	TGGTTGGTGA
25	AGCAATAAAG	GGAAAGTTTC	TGATAGCTGA	AGTCATGGCG	GCTAAAGGCT	ATAAATGCTA	GGCACTTTGC
	CGGATCAGGC	GTCAAGACAC	AGTGCAGGCT	GTGCAACTTC	GAAATCGTGA	GAATCTACTT	TGCTTTTGGC
	AGGCTGTTCG	GAGAAATCTT	CCTGTGAGCT	CCTTTTACAG	GGCTGTGCTA	GGTGCAGCTG	CIGGCTATGG
	AICTGAGGTA	ATTTTCTGCT	ATGGAACTCT	CAATGATGGA	AGTACATGCG	AGCTTCTCAG	TGAGGAGCTA
	CTAGCAGAGC	CTTCTCTGCT	TCTCTGTCAG	GGTGGCAGCT	ALTGGACGCA	GIGGGGACTA	GTTCCTGGGG
30	AGAAACTGGG	GGCGGCAACT	GTGGGCGCTC	CGAGTAAATC	GGATCGGCAC	GTGCATGAGC	ATGCTGTTCC
	GTGGCAGCTG	TAAATGTTTC	TCTAGATAT	TGGGCGCACT	ATCTCCAGGC	TGAGCCAAAT	G
35	SL1.00sc07408_310.1.1						
	ATGGAAATGG	ATCTGGATCG	TGGGCAAGTC	ACATCTGATG	CATCCAGTAT	ATGCTTTATG	CTGACTTTCG
	ATTTGAGCAT	ACAGCTAGTG	ACAGAAACAT	TTACGCTCAG	GAAGAAGCCT	GAAGAGCAAA	AGGCCATAAT
	TATATATGCT	CTCAAGGCAC	CCTTGTATGC	TATTTGCTCC	TTCATTTGGT	TGGTGCATTT	CAAGGCAAGC
	AAACCCCTTT	TCATCTTCTT	GGAACTGCTC	AAAGAATGTT	ATGAAGCAAT	TGTGATGGCT	AAGTCTCTGG
	GGTTGATGTA	CACCTTACTT	AAATATATCA	TAGCAAAATA	CAATGCTCTT	GATGAAATTA	AGGCAAGACA
40	ATCTCAGCAT	TCATCTTCTT	TGACATCTCT	CGAGCTTCTC	ACTGCTCAT	TGAACCACTA	TGCAATGAAAT
	CTTCTCAGCA	ACCTGATGAT	GGAGTTTCTT	GTGATCTGCT	CTGTATGCTC	TAATTTAAAT	ATTTGTTTAC
	AAATGTTTGG	AGTGTATGCT	AGTTGGGTTA	CTGTGATCTT	TACCATCAT	TTAAACATAT	CGCTTTCACT
	GGCATTTGAT	TCTCTTGTGG	TCTCTTACCA	CGTGTCTTGG	AAGGAGTGG	CGCTCTCAAA	GGCACTAGCC
	AGTTCCTGCT	GTGTCAAAGG	AAATGCTCTT	TCTGCTTCTT	GGCAGGGCAT	TCTGCTTACT	GTCTCTGTTT
	CATAGGCGAT	AAATAAATCT	CATATTTCTT	GGCTTGAAGT	GGAGCGGCTT	CAGGAAGGTA	TGCAGAAATG
45	ACTAGTATCT	CTGGAGATGG	TTTCTCTGCT	TATCTTATG	CGTCAAGCAT	ACAGTGGAGC	GGCATATGCT
	GCAGAAAGCT	TTACAACTAC	TTCAGAAAGT	GCTACTTCTG	GAGATAAATA	GAATGAGTGA	
50	SL1.00sc07408_309.1.1						
	ATGGACAGCA	TCCTCTCAAT	TCTCATTAAT	TATACACCTT	TAAAAACAGC	ACATGATGCT	TGTATAGAAA
	AACCGAAGCT	TCTTGGGCTG	AAATTTACAG	GAGCGAGTTC	TTGTTTCACT	AAATTAAGGG	CAAGAAAAGT
	GTCTATTTGT	AGTATGTTGT	ATAAATTAAG	TGCTTTTAAAT	GAGAAATCT	CAATCTTTGAT	AGATAAATAT
	GGGATTTTAA	TGAAGGAGGA	GAGATGGGCT	TGTGAGAAAT	GAAATGTTT	TGTGAAATTT	AAACAGGCTT
	TGGATTTGGA	TGGGATTTGG	GATGGTGGGG	GTGGTGGGAG	GATTAATAGT	GAGACTGTGA	GAGTGTGAG
	TAATCTTGT	TTAGCATTTC	GGCTTACTTA	TCTTACTATG	ACTGCTCAGC	TGGGTTGGAT	CTTGGATGCT
	ATGTTTCTG	TTTGGCTGCT	TTCAGTTCTA	CTTCTTAAT	TAGGTTTGGG	AGCTTTTAT	TGGTGGGCTG
55	GACGAGATAT	TGTTCAAGAG	GCTTGGCCAA	ACTGCGGGAA	TGAATTTCAA	ATTTCAAGT	CTACTTAAAT
	TGATGAGGTT	CAGCTTGGCT	CTTCTGTTAC	TCAGGCATTC	TCAGTTGTTG	GCAATAAGTT	CGTGCAGAGC
	CGAGTAAGCT	TCTCAAGACA	GTCAGCTTCA	TTTGGTCAAG	CAATTTGTTG	TTTCACTACT	CGTTCTAAGA
	AAGGTAAGAA	TCTCTGCTG	GGAATTTGTT	ATAAGAGAGC	AGAAGTTAAG	GAGCGGGACT	AG
60	SL1.00sc07408_308.1.1						
	ATGGCGGGCT	GGAGCGGGCT	GGGAGGGCAA	GCAGCGGAGC	TATATCGAAT	CTCAGCTTCT	AAATCAGTTA
	GCTCAACGAA	GCAAGGGGCT	TTCATTAATC	AGCGGGGCGG	CCTTGGCGGCT	GGTGGTGAAT	ATCATGGAGC
	TCTAAAGGCT	AAATTTTGGC	AGGATCGGAT	GAGTCTGCTT	AAATGGAAG	AGGAGCATTT	TGTGATGGCT
	TCTCTTACTG	GTGGGGGCTT	GGCTTCTTAT	GGAGGTTTAA	AGCTCTTCTC	AAAGGGAAAG	AAAGAGGAGA
65	AGGAGAGAG	AGTTGGTGA	GGATCCCACT	AA			
70	SL1.00sc07408_307.1.1						
	ATGAATGAGA	AGGCTAATGT	TACTAAGGAG	CTTAATGCGA	AGCATAGAA	GATACTAGAA	GGACTTCTTA
	AGTTGCCCGA	GAAAGGGGAA	TGTCAGAGAT	GCAAGGCCAA	GGGTCTTAGA	TGGGCTAGCG	TGAACCTGGG
	AAATTTTATA	TGATGCAAT	GCTCTGGGAT	CCAGAGGAGC	CTGGGGGCTC	ACATATCAAT	GGTCTGATCA
	GCTACAGAGT	ACACATGGCT	TCTGAGAGAA	CTTGTATTTA	TCCATCTAAT	GGGAAATGAG	AGGTCATAAT
	GTCTTTGGGA	AGCAGAGCTG	CTTCTCAAT	ATGATAGAGT	TGGTATGAG	AAATTTCAAT	GAGCAATAAT

	TGAAGAGAAG AGGTGGGTCC CTAAGGATGG GATTCAAAAA TCACCTTCCA GGGTTCAGGA AGAGAGGGGT TCGGTGCAAT GGCACAAAAA CAATGATAGA AGTGGGOTIA TACATGCAGC TAGCTCAGGA LGTGCATCTG ACGAGAGGAA AATATTTCAA GCTCGAAAG TGAAGCAAGA TGTACCTGCT GCTAGAGTCA GTATACCAGT GCGTCCCTAGA GGACCCAGAGC ATGTAACCTC AGGTACAGGT GGTAAACCAG CAGTCCAGAA AGCAGAGCCA 5 GTECAGTICA CTGAACAGCT TAAAGAGGT CCGGAAGCTG CAGTCCCTCC TAAAGTGTAT TATGCTATCTG ATCTAATTTGA CATGCTTTCT ATGGATTTGT CACTGACAA TGGCTCAGAA GCAGCTTTCTA CCGATGATAA CTCCTGGGCA GGTTCAGCT CTGCTCAAGA AGCAACAAA GCAGAAAAA CTGGGGTTAC TAACTCGGT GATCAGAGAA AGTCTCAATC TGCTGCTGCT TCTGGAAITG AAGATTTAT CAAGGATTIA CCTCAATIG 10 TGGCTTCTGC CTCTTCGAG AAGCCACAGA AAGATGCTAN AAGCATATA ATGAGCCCTT TCGACAGTTC CAATATTGTT TGACCTTTTG CIATGCAICA ACAAACACTT GCTATGCTTG CGCAACAACA GTCTTACIG ATGGCTGCGG CTGCTGCAAG TGTTGCTGTA AGACTTCCTG TAAAGCACA ACAAGTACT AATGGACCCA AATGTTTAAA TCAGAACCTG CCAAAITTAG GCIATCAGIT CCGTGGAGTG TAAATGCCAG CAGCTGTAA GACTCAGCTG GAGAAATATA TGCAGGTAG CAATATGGA CCAGCACATP TAGTGGAAA CTCTGTGCCA 15 ATTCAGCAT CCAGCTTTTA CTCAAITGGA CAGAACACTT CCAGCAAGCG TATGTTGCGG CAGAGCCAAA GCAAGCAGAG AGCTACCTCG ATATCATCAA GCCTCACACA GTCAACAAA GATTCGACT TTCTACCTT GACACAGGGT ATGTTCAACA AACGCTGA SL1.00sc07408_306.1.1 ATGGCGATTT TGTTTCAAAG TCAAGTAGC TCCATGTTAT CTAICAAAGT TTTCTGATT TCGACTACTG 20 TTTTATCTGG TGCTATTATG TTAAGGIGT CTGCTCCTGT TGTACTGAA TTCGGGTCA GTGAAGTTCC GTCGAICTGG AACGGTGTGG TTTGCTGGCT TAAACCTCG TATCTAATCC TGTATATCA CIGCTTTATC AITACGAITG TAGCTTTTC TAACTGGAG AACAGCTTG ATGAGAACTC ATCTCCGGTG CCGGCGGTIG ITTCGCTGGA GAACCTGCTC CAGTTTCACC CGATTAAGGA TGTAAAGGCG GTTACAGACT ATTATACACC 25 GGTCTTTCT CACTTAAAG GCTCCGTGCT GAAGATCAA GCAGTGGAG CCGAGGCTAG ACCGATGATT TACAGATATC CIACIGTIG TSTTTATGAT GCAAGGCTG ABAAACTTCC AGTATTAAT CCGTACATAT CCGAGAAAGG TACTCTGTTC AACACCTATC CCGAGCTTAN CGATGTTCTT GCTGAAAAGG AIGATTTCGI AATCTCGAA TCTTCTTGGG CCGCGGTGAT GAGACAGGAC TCTATHTGAT ATTCAATTC AGGCACTCTG GCTGAGAAAC CTCTGCTTTC TGCCAGATTC GCTCACCGGA GAAAGTCAA ATCCACTCTT GAGGTGGAA 30 AGGGAGCAIT CAGAGTATCA AAACTTAAAC GGCAGACAC GCTGGAGAGT ACATGTAAGA CGATACCGA GGGCGGTGCA ATGCCACTAA CGAGGCACCT GAGGAAATCG GACACGTGGG AGACATACCG TGGTGGAAAC CCGCTTACAC CACCAACGCA GAAGATGAAG AAACTGTAGN CGTCAATTA CCGACTTCTT CCGACTTCTT CCGCTATTGCT GACTCCCTCC CCGGCTGGT CAGGAAACTT TAAGAAAGAG CCRCTGCTTA GTCAGAGTGA 35 GCTGAACAGG CGAGTTGAAG CGTATATTA GAAGTTTAAI GAAGATAIGA GGTTCAGAG GTCAGAGTGA ATGCAACAGT ACATCTAGAT GAICATGCA GGTCTCACT AG SL1.00sc07408_305.1.1 ATGCGCTTCC AGCACTGCCC TATGCGACA ATTACTTCTT CAGCTCAAT GAAGTGGGT GTTAGGAGTT 40 TGAGGTCGTA CGAGCACTTC CGTCTATCA CTAICTCAAG CTTCAAGCT TTTCCACGA GAATTCAGGG TGCGCATAGT AGTCTGGAGC AAGGGAAGT GCCTAGCAGA GGTATTTTGA CAAGCAAGA TAAATCGAAG GATCTCCATG AAGAIGTTTT TICTTTGCCA ATGGCTTGCA CTAGTGGCCC AATTAGATTC ACAATGCTCT CAACTGCTGT CATGGCACA AATTGGTIG CAACACTTAC CGCAAAAGCT TCGATATGG AAGACATGAT GGATTTCTCT AGCGCTGCTC ATACATTAGC TGATGGAAGC ATTGGAGATC TTTTGGGTGG CCTTCIGTAT 45 ICTCTGTTGCT AACAGCTTAA TGAAGCTTGT CAGGGCCAGT TGACCTGCTT TAGTATTACT AGTTTGGCCA CTATTTTGG TGCGGGGCTT GTACCTAGC TCTGCGCTTG TACCTCAGC TACCTGCTTC TGACCTTCTG TTAATTTGGG GCTTTTGGTT CCGGAAAGAG CCGAGTAGCG GTTGTGGAG ATTCAATTGC ATTTCAGTCTG GGATGGGCAA CCACACTAGC ATATTTGGGT ATTGGGCTTG CATTTGCTTG AAGGCTATAT GGACAAATAG GACAAAGGAT GCGCGTGGCT GGTCTCTTCT TAGCTATGTT TAIGGGGCTA AACCTGTTAG AGGTAATAGA 50 TGTGCAACTT CCTTCATTTT TACAACTT TGAICTCTGC TCAGCGCTG CTAAGCTTTC GTCAGTGT CAAGCTTAT TGGCGGTCT TACATTTGCA TTAGTCTGCT CACCAAGCAG LACACCTAGT CTGCAACCT TGTCTGGCTA TGTGCTGCT TCTCGGATC CAGTATTTGG GGGCAGCTTG CTAATGACAT ACACACCTG CTAGTTTACT CCGTTACTTC TGTGCTGCT TTTGCTTGA TTTGCTTGA GCAITGCAGA ATATCTTTC AITCGGCAAG 55 TTTCTAGCAT GGATCAATCC AATCAGTGGT GCGCTACIAC TAGGAGGGGG TGTCTATACC TTTCTGACA AGCTTTCTCC GGTGACGATG GCTATGAG SL1.00sc07408_304.1.1 ATGGCGAAGA ACAGAAACAA GAACAAAAAC GGCCTAGCTG CCAAGGATGT CTCCACTGAC CAGACGGTCA 60 TGAAGTCCCA AGCGATGGAT ACTTCAGAT CAGCTGCTCC AAAACCAAT ATAGGTGGAT CACTTAGAAA GACGAAGGGA GTACAAATGA AAAGGACGAA GAACTTTAGG AAAAGGAGG CCAATGGCAA GGCTATTTCA AAAAGTGGA AATTGGAGCA AAGAACTACT AGGAGTGAAA GUAAGATAGA GAGAACTATA AATGCCAAAC AGTATACGA ATGA SL1.00sc07408_303.1.1 ATGGAATTTG ATGCTCAGGG GGBAAATATG TCTACTATAG TCGTAGTTC GTTAGAGGAA AIGCTGAT 65 CTCTCCGGCA AAGGATGAA AATGAGAGC CAAAGACCTT GCGCGCGGCT ITGCTGCTC GGCCTAAGCT TACTTCGAGG ACTAGGCTTC CTACACAGG CCAACCGTTG AGCAAAAGGT TAAAGCAAGG TGAATTTGAA TTGGAGAAAG GTAAAGAGAA GAGGAGTIG AAGTGTTTGA AAAGGATGT ITTTGGTCT ATGAGGTGGA AAGGAAATCA AGATAGCAAA TCGCCATATG CAATGCTTTC AGTGAAGAAA AACAGCACAG GGAGATTGCG GGAAGTAAAT GGTGGGAGG ITAGAGAAATG GCGTAGTGA GGTGAATGG ACGATCGCTT GATATTTT 70 GTTAAGAGCA AGCTTCTCAT CTGGTGTCT CTGGGGAAG GGGCAIGGT ATCAGGACTT ATTCAATCAA CTCTCAGGAG GAAAGCTAAG GTTCTCTCT CAGAGGCGAG TGTGTGACA GGTGCTGCT GAGAGTAT ACCTTCTAT CCGGATGTT ITGAGGCTT GAGTATCTC ATGCAGTIA GTTATTTGAA CGAGCTATC GTTCTACA ACCTTCAACA CAGATATGG CGATATAGA TATAAGCAA GGCAGGGGCT GTATTAAAG
--	---

	CAATCAATCC	GTTCAAGAAZ	ATCCAAITGI	AIGGAAACGA	AITTGTTACA	GCTTACAGAC	AGAAGCTCTT
	GAGTGATCCT	CAATGTTACT	CTATTGCTGA	TTCTGCCTAC	GATCGAATGA	IGGAAGATGA	GATAAGTCAA
	TCTATTATCA	TAAGTGGGGA	AAGTGGATCT	GGGAAGACGG	AAACAGCAAA	ATTTGCAATG	GAATACTTGG
5	CTATGCTTGG	IGGAGGTAGT	AATGGGATAG	AGAAGGAGGT	TTTGCAAACT	AGCTACATAI	TGGAGGCTTT
	TGGGATGCTC	AAAAGTTCCA	GGAACAAABA	CTCCAGTCTA	TTTGGAAGT	GGGTGAAAT	TCATTTAGT
	CCAGCAGGAA	GAATATGTGG	TGGLAABATA	CAAACCTGTA	AGTGAATATG	TCCTTTGTCG	AGAGTGGTTC
	AGCTGCTTGA	TGGAGAGAGG	TCCTAICATA	TTTTTACCA	ACIATGTGCC	GGGGCTCCAC	CTACTTTGAG
	AGATAAARCT	AAGTTAAAAG	GTCATTCAGA	ATACAAAIAT	CTCAACACGA	GTGGCTGCTT	GGTGAICCAT
	GATGTIGATG	ATGCTGAGGA	ATTTTGLAAG	CTTATGGAAG	CTTAAATAC	TGTTAGGATT	TCTGAAAGGG
10	AICAGAGACA	TGCTTTTCAG	ATGATTCCTT	CAGTCTATAG	GCTGGGAAC	ATACATTCCT	AAGTAATTGA
	CGATGAAAGT	CGTGCTGAAG	TTTGLCBAAG	TGAAGCTGTT	ACAAATGCTG	CTAGCTTGAT	TGGCTGTACT
	GTAATATGAC	TCATACTAGC	TTTGTCAACA	TGCCAATATC	GAGCTGGCAA	GGATAAGATC	GCCAGAGATT
	TAATGTAGAT	GCAGGCAACT	GATAGAAGAG	ATGCAATGGC	AAAGTTCATT	TAIGCAAACT	TGTTTGACTG
	GATAGTAGAT	CAAAATGAACA	GAAACCTTGC	AATGGACAAA	GAACAGATGG	GTAGATCCAT	AAATATTCTA
15	GACATTTAGC	GTTTIGAAATC	ATTTCAAGGA	AACCTCATTTG	AACAATTTCT	GAIAAABCTAT	GCAAAIGAGA
	GGTCCAGCAA	GCAIGCCAAAC	AGACATCTAT	TGAAGCTTCA	GCAAGAGGAA	TATGAATTEG	AIGGAATTGA
	TTGGTCAAAA	GTAGATTTCG	AAGACAACCA	AGAGTGGCTG	AACCTTTTIG	AAAGGTAATG	TCTCCTTTTG
	GTCTTTTTC	CTAATTTATA	GGCGTTAGGA	GCTTATATAA	GCGGCAACT	TTCTTCTGCT	CTGGATAGCA
	TGTTTACAT	CAAGCCAAIT	GGCTTATAT	CTTTGTGTAA	TGAAGAAATCA	AATTCCTTTA	CAGGCCACAGA
20	TTTACCTTTT	GTATGTAAAC	TTAAGCAGCA	CAICAAAICT	AGCCCTTCTT	TTAAAGTGA	AAGAGAAAGAA
	TTTTGATCTC	GTCATTTATG	TGAGAGGCTC	AGTTTCTCTT	AZAAATGCAI	CGTCTAGT	CTGTGCAAT
	TTCTACGGAT	ATTATAGTA	TTTTCACIAT	TTTGTATAG	AGACTTGATA	GCITTCACAT	CCCAAGTCT
	AGACTTACCT	TCATTTGTTG	GGAGAGTTTT	TCCTTCTTAT	CATGGTTTAT	TTACAGTAAAC	TTATGATGCA
	ACCTGCTTCT	TAGCAAAAGA	CAGAGATGIG	TTGCACTCTG	ACATTAOTCA	GCTTCTCTCA	TCAGGTGACA
25	GTCACCTGCC	TGAAGATAAR	AAATATTCOA	TTCCATCAAC	TGATGCAAGA	GTGCTAGAIT	TTGAGAGACA
	AAATGTTTGA	ACTAAGTTTA	AGGATANTTT	GTTCAAAATG	ATGCAAGCAI	TGGAAATATC	CTTACCCAT
	TTCAATGTTT	GCTTAAACCC	AAATATTAAG	CAGCTTCCIG	GCTATCTTGA	CAAAATCTTT	GTCATAGAAC
	AACTCAGATG	CTTGGGTTCT	CTTGAAGTTC	TTAGAAIATC	AGGATCTGGC	TATCCTACTA	GGTTAACACA
	TCAGAAATTC	ACAAGCAGGT	ACGCTTTCCT	TCTGCCAAAG	GATAGTGCAT	GCCAAATATC	TTTAAATATG
30	TCAGTTGCCA	TCTTTCATCA	ATTTGGTATT	CTTCCGGAAC	TGTACCAAGT	TGGGATATCC	AAGTTATATT
	TCCGATCAGG	ACAGATTGCT	TCATTTGAGG	ATGCAAGGAA	CCAAGTTCTG	CBAAGTACTC	TTGAGTTGCA
	GAATGCTTTC	CGTGGTCTAT	GTGCTCGTCT	GCACTTCCAT	GAACTGAAG	GAGGAGTAT	CATACITCTA
	TCATGTAAG	CAGATCCCTT	TGCAACCTCT	CCTTTTCTTT	CCCTTTTCTT	GTTGAATATA	ATAGAAGATG
35	ATATTCATGA	AGATTGTTCT	GTCTAICAGG	TTGGAIGA			
	SL1.00sc07408_302.1.1						
	ATGCTCTAAG	CGAAGTTCCT	TGCTGAAGGA	AGTCAAGAGC	AACTGGTGGC	TGCTGGTGGC	ATACAAATCAG
	CTATTCGTGG	TTGGTTGGCT	AGAAGGGGTC	TTCCGAAACI	GCGAAATCTA	AAAAGTTTAA	AIGTAGACAA
40	ACGAGATACA	GGCAGAAAGA	CGGAGGTCAA	GGAGTTGGCT	CGAGAAATCC	LACCATCTGT	LGTAGAGAGC
	CTCCAAAGAC	GGGTTCGAAA	GGCGAGGGCA	ACAACTGAAC	AAAAGGAAAA	GGAATAATGCT	GGCCTGAAGG
	AACTAAGTAA	CCAAATTCAG	ACCAAGAGCT	TAGAAATATG	GCTCAAGATG	AGGTCAATGG	AGGAGATGTC
	GCAAAAGCAA	AIGACATCAT	TSCAGGTTAG	TCIAGCTGCA	GCCAGGAACA	GTCTTACCCG	TGCTGACACT
	ACTGGTTCAG	CTGGAAGGCT	TGAAGGTTCC	CCATCTCCTC	AGTATTAAGA	TTCTGATGAT	GCAACATCTA
45	TGACATCTCC	TGCGGAGGCT	ACTCCAGTTA	GCITTCCTTA	CAACAGCTTG	GGTGTGTAG	CTAATAGATA
	GGTTAAATGT	GGTTTATCTT	TAACTAGGCA	CCTTTCATTA	GAATTTGAGC	AAAGGAGACA	AAATTTTGAC
	AAAGAAAGCA	TGGCAATGCT	TCATTTGAAG	CCAGGAGAGT	TACAGTCTAC	TAAATAATCT	GCAGATGAGT
	AICGAAAGCT	GAAACACAGG	TTTGAAGGAT	GGAATAAGGA	GTACAGGCTT	CGGTTAAGG	AGCAAAAGTC
	AAAGTACAC	AAGCTTCTTC	ATCTAAGAGC	AGGGAAGAGT	GCTAGAAAT	GGTGGGTTAA	GAAGAGCAAA
50	TGA						
	SL1.00sc07408_301.1.1						
	AIGGACAGCA	CAGGTAAGCT	AGCCACAGTT	GTIAGAGCAA	CTCCATTTT	GGGCCAGACC	AAATAIGCTA
	ACCCCTTAAG	GGATATAGTT	CCTATGGGCT	CCTGACAGAT	CACCATGAGT	AATGATTTGT	GGTAAGGACC
55	TGACCGTGTG	AGTACTTGG	GAACATTTTC	TGCTCAAACT	CTTCTACTAT	TGACTGGAGA	ATTCCCTGCT
	GATTAAGGAT	GGGATAGTGC	TGGTTATCTT	GCTGATCCCG	AGGCTTTTGC	TAAAGACAGA	GCCTTGGAGG
	TTATCCATGG	CAGATGGGCT	ATGCTTGGAG	CTTTTGGTTC	CATTACACCA	GAAGTCTCTG	AAAAATGGGT
	AAAAGTGGAC	TICAAAGAAC	CAGTATGGTT	CABAAGCTGA	GCCAGATCT	TCAGTGAAGG	TGSGCTGGAC
	TAATTTGGGA	ACCCAAACCT	IGTCCAGGCT	CAGAGCATTC	TAGCAGTATT	GGGCTTCCAA	GTTGTAATAA
60	TGSGCCTTGT	AGAAGGTTTC	AGAAATTAAG	GGCTTCTCTG	AGTTGGTGA	GGCAACAATC	TCTACCCAGG
	TGAGCAGTAC	TTTGACCCAT	TGGGCTTAGC	AGATGACCCA	ACAACTTTTC	CGGAAGTCAA	GGTCAAGGAA
	ATCAAGAAAG	GAAGATAGC	TATGTTCTCT	ATGTTGGGAT	TCTTCTGCTA	AGCTATTGTC	ACCGGCAAG
	GCCCACTTGA	AAATCTATTG	GATCACTTTC	ACAAACCTCT	TGCTAAGCAAT	GCTTGGGTTT	ACGCAACTAA
	GTTTGTCTCT	GGATCTTAA					
65	SL1.00sc07408_300.1.1						
	AIGTGGAGAG	GGAAGGAGT	AAITGGAGAT	ACAGAIATGT	TGGGAACCT	GCAACAGAT	GCAITAGATT
	TAGCTGCAAA	GGCACTTAC	TTCCTTGAAT	TCACTGAGGC	CACCTGAAT	GCAAGTTTTC	TTAAAGAGGA
70	ATTTGATACA	ATGATAGGAC	CAGGGTGGCA	ATGCAATGTA	GGAACAGATT	TGGTTTCTT	TGTAACACAT
	TGTTATGGIT	GTTTCAATCA	TTCTACAT	GGCAGGCTTC	CTATTTTCT	CTTCAAGGGC	TCTGCTGCCC
	TAGAGGACCC	GAAAGCCGAG	GCTGAAGCTG	ACCGATTTTC	CACCTCTGAG	GAAATAGCAT	GA

5	SL1.00sc07408_299.1.1							
	ATGGGGGGCC	GTCTCTTCAC	CTGCTTCGGC	AAAGGCGGTT	CTTCTCATTC	TTCTTCAAAA	GATCCCGGCT	
	CGAATACAA	GGACAGTCCG	ACGGCGGATT	TGACGGCGGA	GGAGCAAAAA	CGGTGCGGGC	CGGTGCTGGT	
	GGAGTATATC	TCATCGCAAG	GCTCGGCCAC	CTCACCTIGAA	GCGGAGCTGT	TGTTTTCGAG	GATTGGGAGA	
	GGCGATTITA	ACCTAGAAAT	GTCGGTGATT	TTGTTGGCCT	ATCATGTGGA	TTATTGGGAT	TATATGGGTT	
10	GGAGGATCC	GTTTGGGTCC	AGTTTAIGGA	CGGTAAACAC	AAAAGCGTAT	GTGGAGACCT	TAAATCTAGA	
	TACCATGTTT	ACGCTCAAAA	TTGTGGTTCA	GGGAGAGGCT	CAATGCTGTG	GGATGAACA	AGAIGCGGTT	
	TTCTCTTGTA	TCAAACTCTC	CCCCAGATT	GCTGCTCCTT	CCTTCCAGGC	AACAITCGAG	AGGCGAACAC	
	CAGAGTCAIT	GCAAGTATCT	CTATTAGGAT	CTCTAAGGAG	TAAGGTGGAC	ATGATGGTG	CGAACCTGAT	
	GATTGCTCTG	TACGAAAGTG	GCTGCTGATC	TGATATCGCT	GCTAGGAGCA	ACAAAGGAAA	AATGCTTGGC	
15	AATGACTAIG	TTGTCAGGAG	GCTGGAAAAG	CITTTGCTATG	TAAAGGATAT	TACTGCCAAG	AAGACAATCT	
	CAGGAAGTGT	CAATTTCTCT	CTTTGGGATG	GCCTCATATG	CAGCAAAATG	GGCGTAGCGC	TCCTTGTGGA	
	AICTGGCTCT	CATCAAAATAT	GTCGATCACA	AAACITTTAA	TTCGCAGAAA	ATCTCTGA		
20	SL1.00sc07408_298.1.1							
	ATGGCAATCG	CCGACCAACA	CAAGTCACAT	AACCTCAGACG	GAAAGCTCTG	GAAGCTTTGT	CTCTATGGC	
	AAACAGGAGG	TACGCTCTCT	TCCTCGTCTG	CTACACAAAA	TCCTCACTCT	CAGAAATACA	GTCAACAAAA	
	CGCGCTCGGA	TCCTAGAGCT	CTCGTCTCTC	TACGCTCTGT	AGCTCCGCTG	CTAGATCACT	CGCTCCGGCT	
	AGAGCTAGGC	TTCCGGCTCGA	TCAGGCTAAC	AGTCTCTACT	TCCTTTATGA	ACCGGTAAG	CAGGTGAAGA	
25	GCTGCTGAAA	GATTAAAGAC	ACTAGCAAA	CTTAAGTTGC	ATTAAAGTTT	CAAGCGACIG	CACCAAGAG	
	CTGCTACATG	CGACCTCCCG	GAGGCAITCT	CGAACCCGGT	GAAAGTGTTA	TTCGACCTGT	CTCTAGTTT	
	GCTGCTGCTC	CTGAGAACAA	CGAAAGGCT	GTGGACCAAA	AGAGCAAACT	TAAGTTCAG	ATCATAGCT	
	TCAGCTGAAA	AGAAAGTCTA	GATACGTAAC	CTGAGTTGTT	TGAAGAACAA	AAGGATCACT	TGACTATAGA	
	ACGTAICTTA	CGGGTGGTGT	TCCTGGACCC	AGAACGACCT	TCCTCAGTGC	TGGATAACT	AAAGCTCAG	
30	TTGCTGGAAG	CTGAGGCTAG	ATTAGAACT	CGCAAGAAAC	CTCCAGTTGA	AACTGGACCT	AAGTGTGTAG	
	GAGAGGCTCT	AGTAAATAGT	GAAAGGAGG	AACGAAGGGA	GAAATATCTC	GCTCGGACG	AGGTGAGGCT	
	TGTTGATTCA	GTGTA						
35	SL1.00sc07408_297.1.1							
	ATGAATGCAG	CAGCAGCAAA	GGGATCGGCA	TGAATGCTAG	CAGCAAGTAT	TGAGACAGTA	GAAGCATTTAA	
	AAGATCAAGG	ATTTCGCCAGA	TCGAAATACG	CTTTAAGATC	GATTCATCAC	TTTGCCAAAT	CTAAITTAAT	
	TGCTCTCTAGT	ATACTCTCGG	CTCGGAGATT	CTCGACGGCT	TGGGGCGCTG	CGGCTCTCTC	TCGGGCAATC	
	GTCTCCGGTG	AGAAGCTGAG	GAAACCGGAA	GAGAGCTTGA	GLAAGTTAT	AGATCTGAAC	TGTTGGGGTC	
40	CAAGCACTGT	CAGATTCTAG						
45	SL1.00sc07408_296.1.1							
	ATGAGGCTTC	TCCGATCAAA	CTTCTGCTGT	GTTCGGATTG	CAGAGCCGAC	TACTGCTAIC	TACAGACAG	
	AGTCTGCTAT	TTCTTTGGAT	ACTCCGAAAT	CCGATGGCGG	GCTGTTTGT	GATAAGAGTA	CTTTCTTCTG	
	ATTTCGAAGG	GATTGCTGTG	ATTGGGACTA	TGAGAGGACT	GGGAGCCGAG	TTTATTTGCA	LATAAGCAG	
	ACAAGGAAGG	CAGATGCTGA	AGATAGACCG	TCCAAAAAC	CCACTCTCTT	GCTATAGGT	TTAGACGGAG	
50	GGTTTGATAA	CAGTGAACCC	CAGTACGAAG	AATCTATGTA	AATAGTTATA	TGCTGTGATA	AIGTCACTCT	
	TCCTTTCCCA	TCCGTTGAAAT	TSCCTGAGAA	GGTTAGATTG	GCTGTGATG	CTATTTTACT	AGCTGAAGGT	
	GCTGAGAGGA	AAGAGCAACT	TGCTTCTCTG	ACTGCTGACA	AGAAGCTTGT	CAGTAAATAT	GCTACGGATC	
	TGCAACAGCT	TGCAATGGGT	GTTCCTGCTC	CAGCTGTGGG	TTGGAATGT	CGGAAATGAT	ACAAGCTGA	
	CAATCTTTGG	CTGAATCTAA	CTGATGTAAC	TATCTATGTT	GGTAGGAAAA	ATTGGGATGG	ACTGCTGGGT	
55	AATGAACCAIG	CAGTTAAACA	TTACAAAGAA	ACTGTTTATC	CATCTGCTGT	AAAGCTTGGG	ACCGTAACTG	
	CIGATTGGA	GGGGCCAGAT	GTTTACTCTT	ATCCAGAGGA	TGAAAGTGT	GTGACCCAC	TTTATGAGCA	
	TCACCTGGCA	CATTTTGGTA	TTTACTTCTC	ATCCTTGCAG	AAGACTGAAA	TGACGACCTG	TGAGAGAGAA	
	CTGAGCCAAA	ATTTTTAACTT	TGATTGGAAC	CGGATTCAGG	AGACTGGTGA	GGAGCTTGA	CCACTTTTGT	
	GACCTGGTTA	CAGTGGATTA	GTCAATCTTG	GLAACAGTTG	CTACTTGGCT	GCTACAAATG	AGGTATGTT	
60	CTCAACGGCT	TCATTTTGGT	CAAGATACIA	CTTTGATCAA	CCTCTGAAAG	AAGCTTTTAC	TACGGCTCTC	
	GCTGATCCGA	CCTGAGACCT	TAACTGCAAG	CTAACAAAGC	TGGCTCATGG	TTTGTCTTCT	GGTAAATATT	
	CGGGTCTCTGT	TCCTGAGAAAG	GATAATCTG	CTAATGCTGT	AAGCTCACAG	AAACAGGAGG	GTATCCGCTC	
	TGCAATGCTC	AGTCAAGTAA	TAGCTGCTAG	TCACTCTGAA	TTTTCACAAA	TGAGACACAA	CGATCGCTTA	
	GAGTTCTCTC	TGCAATTTAT	TGATCAAGTT	GAACGAGATA	ACTCTGGGAC	ACCTAATTTT	GAATCATCAA	
65	GGAGCTTCAA	GTTTGGTATT	GAAGAAAGCC	TCCATGTTTC	CTCGGGCAAA	GTCACTTACA	ACAGAGGCAA	
	TGATTATATT	CCTGTCTCTA	ATATTCCTTT	GGAGAGGGCT	ATAAATAAAA	AAGAGCTAGC	AGATTTTCAA	
	AAATTTGAAGG	CTGAGAGAGC	TCAGGAGGGA	AAAGAAGTGT	CTGCTGATGA	ATTGTTTGGC	CTAGAGTAT	
	CATTGAAGGA	TTCCTAGAT	TGCTTTTCAAG	CTCCTGAGGA	GTTCCTATGT	TTCTACAGCA	CAGCTTTAAC	
	AGCTAGGACT	ACAGCAATCA	AAACTGCAAG	TTTCTCTCTC	TTTCCAGATT	AATGCTTTT	GTCAATGGGT	
70	AAATTTGTTA	TGGAGGAAGG	TTGGGTGCCA	AAAGAGCTCG	ATGCTCTACAT	AGAATGCTCT	GAAACCTTGT	
	ACAATAAGTAG	CATGCGAAGT	AATGGTATTC	AAACGAGAGA	AGAGCTGTTG	CTGACAGTGT	CTGACGGGGA	
	TGGTGGAGCG	TCATTAAGGC	TTCTGGCTGA	CGATGATATT	GTTCGACAAAC	TTGTTTCAAT	GGGATTTAAT	
	CTACTTCTATT	GTACAGAGGC	TGCTATCTAT	ACTTCTACCA	GTGGAGTAGA	GGGAGCAATG	GATGCTTAC	
	TTAATCTATA	GAAATGATCA	GATATTGATG	CTCTTATATC	AGAAACCTGT	CAAAATCTGT	ATATGATCAA	

	SL1.00sc07408_295.1.1					
	ATGGCAAAAT TAGCGAAAAA TCGAGGAAGG AATAGATTTT TTAATAGTIG ITATAAGCCT CTACATTTCA					
	ACGATGATAT AICTAATACA AATAAACGAG GTCAITTTGTA TAAATCAACA GATAATCTTA ATTTGAAGAG					
5	AGTAAATATCA CAAGAAAAAT TGEATTTCAAT GTTGGGAAAA GATTTGACCA AAAATTTTCG ACCAAATOGA					
	AAGGAGAREG COTTIGGGAAG GAATTTTCTT CAIGGACTCA AAAATGTGTT TTTPGATACA TCAITGGGGA					
	AGAAAGGCCA GAGGAAGGAG CATAGATATT CATTTGGATC AAGTAAABAG TATCAACAA AATTTCAGAA					
	AAATATTTAT TCTACAAAAG AAAAAAGATT TTCTTCATCA AAGGATTTAC CAAAAATTAC AAGLAGAATG					
	TCCACAGGTG CAAGCAITGG TGAATTTTCC TTATTCACATA CTTCTACATC CTCTTTATTT TCTTCTCTCT					
10	CTTCTCTCTG TCGTCTTGT TCAZCACAGA GAACGTAAAT TPCATCTTTT CACAGGTCAA AATCAGAAAA					
	CAATATGCAA GGTATGATA AGGTAAAGA AABAGATATC GCGCTATGTT ACACAGAA ACGTTGGGACG					
	IATTCICTTT TAAITTTGTC ATTGGTATG GTTTTITGTT GGAAAGTTT TGCATCTATT TGCATTTCAA					
	CAZGGTTCTA TTTTGTCTCT CATGTCTTCA AAGGTATAGA CTGGGATGAG TACAGAAAA ATGATGGATT					
	GTATATTTTA GTACAGAAAA AAAAICTAAC TTTTATG					
15	SL1.00sc07408_294.1.1					
	ATGGCTACCT CTCTTCAGTT TCTTTCAGAT CATCACCCCTA TCCCTCAGGA AATCATCTAA ACACAGAAC					
	AGACCTCAAC GGGAGGAGCT AGAAGGTGGA GTAAAAATGG ACAGAGAGAG AAGAAGCAAC CACAAAGAGG					
	AATGGGAGTT GAACAACCTG AACGTCTTGG AGTACAAGAT CAGAGAGAAA ACAGTACAT CCATGGCGTT					
20	CATCAIAAIC ATCAGTACIA CTCTAATAAC AATTTCOCTA KATTAACCTC TGTTCATCA TTTACCGCGG					
	GIGTAGIGG TAGTGTGAT CCGGGAATTT ATGATTAATC TATTTTGAA CTTCTACCCAG TACTTCAGTT					
	CCCCAAATIG TGTGCACTGA GCGCTAATGA TTTTITATG CAACAABAG TGTGAAATC TGGGTTTATT					
	GSATCTAGTA GTACAAICA GTTGATGAT TCTTCTCAG ATCATCTATC GTTTCATCT CAGATGAATC					
	ICTAGCGATT TGCACCTCT AACGCCAGTA CTGAGAAATC AAGGGAGCTG TATCCAAAGC CAAATCTGTT					
25	TAGCAGCAAC AACTCTGTT TCTCCGATCG CTGCCGATCA TCGAACAAAA AGAAACGCAT GATCAATGGA					
	GAAGAAATTA GCGTTCATAT GSAGGACATG ATCAGAGAAA AGGAAGATTC TGGAAACAA CCTTTGCTTC					
	ACTCAIACAG TTTACCTAGC CATCAACAAA AGGGGATAGG GATTTGTGCTA AICCATAGAA AGGGAATTC					
	AATCGCGTIG TCTTCGATG AAGGAGCAGT AATGATGGAG TAIGATTTT TPCAGAAA AATCAGCAGC					
	AAGTACACIA ATACTTACAA AAGTTGTTTC GAGATGGAAG CAACGATGAT GAGTGTCTAT AATTCACAG					
30	AATCTTCTTC ATTTGCTGCT GCAGCAGCAG CAGCTGGAAA TATTATTAAAT GGTGAAGCTT CTCTCTTAC					
	TACAATATCT TGGGCTGAG ATACTACTAC TACTTACCT ACCAGTTCCA TTGAICTTTC ACTGAAGCTT					
	TCUTGTTAG					
	SL1.00sc07408_293.1.1					
35	ATGAGGGAGA GGTGTGCTT TCAAGTTGAG AGGTTAGGGC TTAGTGTGCT TATTATGGGG AGTCAGGAT					
	TGGAGCTAC GAAGAGGGGG AGTATGGA GACTTGGAG TGTAGTGAT TATTGTGTTA GGCACGTGT					
	GTTGCTGCTT GTGGTTGTIA GGTATCCCGA TGAIAAGGAT GGTGGAATG CAGTGTGAGA GCGCGTGGTT					
	TCTGTGCTT CAGCTGTGTA AGAAGACGAG GAGGAAGCTG AATACCATGA TGTCTCTGAG GATCGAAAAG					
	ATTCATAA					
40	SL1.00sc07408_292.1.1					
	ATGGAAGATC CTTATGGAIT TCAAGATCAT TCCCTTICAN TGAITGGAGAG AITAGGCGCG GAAGGATTCA					
	TGAGGGAGCT TIGCAATGGA TTTGTGTTAC TTATGGAIGT GAGTAAAGGG CTAATACAT ATGAAAGTTT					
	GAAGAGAAAC ACTATGAAIC TTGGTTTAAH TGAITTAAGA GATGATGAGT TGATTTACAT GTTGGCTGAA					
	GSAGATTGG TGGTGAATG AGCACITAAI CAAATGGAAT TTGTATTCT CATGTTAGA TTAGTCTCTG					
45	GTTAATIGGA TGGATCTAAG CAATACAIGG AATGATGAG GCTTATTAT TTTCAACCTT AA					
	SL1.00sc07408_291.1.1					
	ATGAAAGGCC TATTTCTTTT ATGCATTTGC TACCAAITTC TOTTCATTT ACTAACTTCT GCAGCATIAG					
50	ACACAAATCAC TACAGATAAA TCCATTAAG ATGGTGACAC AATTTGTTCA GCTGGAGGGG TTTATGAGCT					
	TGGATTTTTC AGCCCTGGAA ATTCGAAGAA TCAITACGTT GGCALATGGI ACAAGAAAAI ATCAAAATGGA					
	ACTGTGTCTT CCGTTGCAAA CAGAAGCAIT CCACTTAATG ACACCTCAGG AGTGTAAACA CTAAATGCCA					
	AIGGATTTCT TGTACTTGTI GATAAATCCA ATGTCTCAAT TTGGTCAATC AACCTCATCA GATTTGTTAA					
	GAATCCAAAA GCACGGTTAC TGEATTCAGG GAACCTTGTI GTCAGTGTG GAAALGAAAG AGGCCTGGAA					
55	AAATATTTCT CCGTGGCAGAG TTTPGACIAT CCGGGAARIA CTITGTTACC TGGTATGAG CTAGGAAAG					
	ATTTTGTAC GGGAAATGAA TGGCATTTAA CGTCATGGAA GAGGACAGAT GATCCTACTC CTGGTGATTA					
	TGTAGATCTG TTGATTTAC ATGCAATATCC ACAATTTGTT GTGTGAAAAA ATTCATCTAT AGTATTAGC					
	TCAGGGCCAT GGAATGTTAT TSCATTTAGT GGTAGTCCCA ATAAATAAAC AAATACATAT TACAGTTTCT					
	AGTTTGTATT TAAACAGCAG GAAATTTACT ACACATATAC AATTAAGAAI GACTCCATAC CCACAGGGT					
	GSTGTCTCAT CCGTCTGGTG TSCATGAACA CCLAACATGG ATAGAGCCCA GTGAGAGCTG GTTCTCTTAC					
60	TTGACAGCAC AATTTGATAA TTGGAATGGI TTGGTTTAT GGGGACCTTA TCAAGTTTC AACATCAATA					
	ACGCCCTCTC ATGTGACGT TGAAGGGIT TTGAGCTAG GTATCCCAA GATTCGTCAA GATCTGTCTC					
	TAGTGGTTTC AIAAGGAGAA CTCTTTTGGG TTGTACCCAT TATGGTTTTC TTAATTTTTC AGGTATCAAA					
	ATGCCIGAIT CTAGAAATCT CTGGTAAAT GACAGCATGA ACCTTGAAGA TTGCGAGAAA AIGTCTTGG					
65	CTGATTCGAA TTGTACAGCC TACTCAGATC TTGATGTIAG AATGCGCGGA AGTGTGTCT TACATGTTT					
	TGGTGAATCT ATAGATATAC GCGGGTTAC GCGAAATGAA CAAAACCTGT AIGTGAGAT AIGTATCTCA					
	GAATLAGACA GGAAGGGGAG GAGAAGGAG GCAGCCCTGA ITGGCGTCTT TCGAGCAGT GTAGCAACAT					
	TTATCTTCAG CTTTTTACCT TGGTTTATC TCGGAGAAAG GAAAAGAGCA AGAGGATTT AAGTTGAAAA					
	TGAGGACARG GAGCTTCCAT TGTGTGATT AGTACCTGTT ACTACTGCTA CAGATAATT CICTCTGCT					
	AATGATGATG GAGAGGTAGG CTTPGAGCC GTTACAGAG GTATCTTACC AAATGGACCA GATATTGCTG					
70	TAATAAGACT ATCGAAGCAT TCTGGACAAG GCTTTCAGAA GTTAAABAAI GAAACGCTC TCATTTCCAA					
	CTTGCACACT AGAAACCTTG TCAAGCTATT GGGTTGCTGC CTTGAAGGAG AAGAAAGGAT CCTAATCTAT					
	GAGTCAATGC CCAATGCTAG CTGGACIAT TTCAATTTT ATTCAAGTAG AAAAGCAICA GTTCTATGGA					

5	AGAACCGTTC	TGAATATGCT	ATGGGAATAT	CACGAGGTCT	TCTTACCTT	CACGAGGTAT	CAAGATTACG
	AATATTTCAC	AGAGATCTCA	AGACTAGCAA	CAITTTATTA	GATACATGAC	IGATGCCCCA	AATTTCCGAC
	TTTGCCCTTG	CCAAAATTTT	TGGTGGAGAT	CRAGTGGAA	GAAAAACIAA	AAGAGTAATA	GCGACATAIG
	GATATATGIC	CCCGGATAT	GCTGTTGATG	GGAAATATTC	AGTAAATCTA	GATGTAATCA	GCATTGCGGT
	AATTTATCTT	GAATATCTCA	GTGGCAGAAA	GARCAGAAA	TTTCCTCAIT	TGGACATCTA	TCACAACTTC
	TTGGGACATG	TAAGCACAAA	ACTTCAITTC	CITTTATCAC	TCTTAGCTTC	CTATGTGTAT	TACAACTGTA
	TGACTGTGTT	GTTCGTGTTG	TAA				
10	SL1.00sc07408_290.1.1						
	AUGSEATTGA	TTTTGTTTTT	TGCTGACGCT	ATGTTGAGGC	TCTTCATTTG	CTGCTCATTT	CTCTTCATGT
	IATIAACTTC	TCCTGCTATTA	GACACAACTA	CIACAAATAA	ACCTATTAGA	GATGGTAATA	CAATTGTTTC
	AGCTGGAGGC	GTTTATGAGC	TTGGAITTTT	CAGCCCTGGA	MAITCGAAGA	ATGCTTACCT	GGCATATGG
	TACAAGAGAA	TATCACCTAC	AACGTGTTGT	TGGTGTGCAA	ACAGAGACAT	TCCACTTAAT	GACACTTCAG
	GAGTGTTAAC	ACTTAATGCC	AATGGAAATC	TTGTACTTGT	TGAIAAAATC	AATGTCTCAA	TTTGGCTATC
15	AAATCTCAICA	AGATTGTIAA	AGAAICCAAA	AGCAAGGCTC	CTGGATACCG	CAAACTTTGT	TGTTAGTGGC
	GGAAATGAIA	GAGATCAGGG	AATTAATTTT	GCGTGGCAGA	GTTTTGATTA	TCCAGGAAAT	ACCTTATTTAC
	CTGGAATGAA	CGTAGGAATA	GATTGGTITA	CGGGTATGGA	TAGGTATGTA	ACGTGCTGGA	AGAGCAGAGA
	TGATCCTACT	CCTGGTGATG	AAGTAGATCG	TGTTGATTTA	CATGGATACC	CGCAATTTGT	CTGTGCGAGA
	AATTCATCTG	TAGTGTCTAG	CTCAGGGCCA	TGGACTGGTG	CTGCATTTTC	TAGTAGTCTT	AGTAATTAAC
20	CATCTTTGTA	TIATACCTTC	GAGTTTGTTA	TCATACAGAA	GGAATTTTAC	TTCAATTAIG	AGCTTAAGAG
	TGACTCTTTG	CCCACCAAGG	TGCTGCTCAA	CCCGGATGGA	GTGALACAA	ACCTAATATG	GATTGAGCAT
	ACTTCAGAGC	GGTTTCTCTA	CTTGACAGCA	CACCTTGATA	ATGTGTATCG	TTTGGCTTTA	TCCGGACCTT
	ATTCAAGTTG	CAACATCAAT	AACCTCCCTC	CATGTGACTG	TTTGAAGGHI	TTTGGAGCTA	GGTATCTTCA
	AGAACTCTGA	CCAGACTGGT	CTAGTGGTGG	CGTAGGAGAA	ACTTCTTTAA	ATTGTACCCA	TGAAGTTTTC
25	CITTAATTTA	CGCGTATCA	GATGCTGTAT	TCIAGAAACT	CTTGGTATTA	TGAGAGAAIG	AACCTTGAAG
	ATTTCGAGAA	AATGTGTTTA	GCTGATGCTA	ATTGTACAGC	CTACTCAGAT	CTGTATGTTA	GAAATGCGCG
	AAGTGAAGC	TTCATATGCT	TTGGAGAACT	CATAGTATTA	CGAGAAATCA	GCGAAATGTA	GCATAATCTA
	IATGTGAGAG	IAGCTGCTTC	AGAATTAGGC	GATGTATAT	TGACAGGTTT	AAAAGTTGAA	AAAGAGGACA
	TGGAGCTTCC	ATTGTTTGAT	TTAGTTACTG	TTACAACTTC	CATCTGAAAC	TTCTCTCTCT	CTAATGTGAT
30	TGGGCAAGGC	GGATTTGAC	CGGTCTACAG	GGGTATCTTA	CCAAAGTGGC	AAGAGATAGC	AGTAAAGAGG
	CTATCGAAGT	ATTCTGAGCA	AGGCATCTCA	GAGTATATAA	ATGAAATCTT	TCTCATTTTC	ARGCTGCAAC
	AIRGGAACCT	TGTCRAATTA	TTGGGTTGCT	GTCTTGAGGC	AGAAAGAGCG	ATGCTATAT	ATGAGTCTAT
	GCCCAACCTT	AGCTTGGACT	ATTCAITTTT	TGATCCAGGC	AGAAAGGCTT	CATCTGGGTT	GAGGAATCTG
	TTTGAATTTG	CTATGGGAAT	ATCTGCTGCT	CTTCTTTTAC	TTCCACCAGG	TTCCAGATTT	AGAATTCTTC
35	ACCGAGATCT	CAAGACAGGC	AACATTTTAT	TAGATACCTA	CATGAATGCT	AAAATTTCTG	ACTTTGGGCT
	TGCGAAAAT	TTTGGTGGTG	ACCAAGAGGA	AGGAAABACT	AAGAGAGTAA	TGGGACATA	TGCAATATG
	TCCCTGGAGT	ATGCTGTGTA	TGGGAAATAC	TGCTTAATAT	CAGATGTAT	CAGCACTGCT	GTATCTATCT
	TTGAAATAGT	TAGTGGCAGA	AGAAACAGAA	ATTCTGCTCA	TTTGAACAT	CTTCAATCTA	CTTGGAGACA
	TGCAATGTTA	CTTTGGATTT	AAGCCACAGC	GTCTGACCTG	ATAGACGAAT	GTATCAAGAA	ATCTTTTCTA
40	GAACTCTCAAG	TGCTGAGATG	CATCCAGGTT	GGTTTGTAT	GCGTCCAAA	ACTCCCCGAG	GATAGGCTTA
	CAATGGGCAIC	AAGTGGTTTC	TGCTTAGGCA	ATGAAAGGCT	GCTTCTCTCT	CAACCAAGGC	AACCTGTTTT
	TTTCTATAGAG	AGGAATTCAT	TGCAATCAAC	AGAAICAICA	ACTGAAGAG	TATATGTAG	TAGCAGCGTG
	TCGATAACAG	TTCTAGAGCC	AGGATAG				
45	SL1.00sc07408_289.1.1						
	ATGAAAGGGA	AACATTTTTT	ATTTCTTTGC	TCATTTTTC	TTCCAGTCTT	ACTAATTTCC	ACTGCAITAG
	ACACATATAC	AKCAGAGAA	CCMAITAGAG	ATGGTACAC	AAITATTTCA	CGTGGAGGGG	TTTTTGAGCT
	IGGATTTTTC	AGCCCTGGGA	ATTCCAGGAA	TCGTATCGTT	GGCAZATGHI	TTAAGAAAGT	AGCACTATGA
50	ACTGTTGCT	CGGTTGGCAA	CAGAACTTTC	CCACTGAAG	ACAACTCAGG	AGTGTATCTA	CTCAATCCCA
	ATGGAATTTT	IGTACTCTTC	CGTAATTTCA	AIGCTCAAT	TGGTCTTCA	AACCTCATCA	GATTTGTTAG
	GAACTCCAAA	CCATGGCTCC	TGGATTTCTG	TAACCTTTGT	GTGACTGATG	GAAATGTATG	TGAATCCAGAA
	GTATATTTTC	CGTGGCAGAG	TTTGTATAT	CCAGGAGATA	CTTTACTACC	TGGGAIGAG	CTTGGAGCTA
	ATCTGGCTAC	GGCATGGAT	TGCTACATAG	AGTCATGCAA	GAGCATGTAT	GATCTCTGCC	CTTGGAGATA
	TATAGAACCT	CTTGATTTCT	ATGGATACCT	ACATTTTTC	GTGTGGCAGA	ATTCACTTAT	AGTATATAGC
55	ACAGGCTCAI	GGAAATGTAI	CACATTTAGC	AGTAGTCCAA	AAAALCAACC	AGCTATATAT	TATGCTTTCT
	AGTTGTTTAT	TAAACAGAA	GAGATTTACT	TTAAATACGA	GCTAAACGAG	TCCCTGCCCA	CCAGGCTAGT
	GATCATATCA	GCTGGAAIGT	TGAACACCT	AACATGGATT	GAGCGAAATC	AGGAGTGGAT	AGCTATATGA
	TCACACAAAT	CTGATATTTG	TCATGTTTTT	GCTTATATGT	GTCCCTATGC	AGGTTGCCAA	ATCAATTAAT
	CTCTCTCTAT	TGACCTGTTG	CAGGCTTCTG	AGCCTAGGTA	TGCTGATGTA	TGCTATGCTG	TGGAGCTGTC
60	TAATGTTTGT	ATAAGGAAAT	CTCTTTTCTG	TTGTAACCAA	GATGGTTTTT	TTAAATTTAT	GATATATCAAG
	ATGCTGGAT	CTAGACATCT	CTGCTATAT	GTAAGCATGA	ATCTTGAGAA	ATGCAAGAAA	AATGCTTTGG
	CTGATTTGAA	TTGTACAGCC	TACTCAATTC	TTGATATAG	AAATGGCCGA	AGTGGATGTT	TACTATGGTT
	CGGTGAGCTC	ATTGATATTA	GAGAGTCTCA	CAAAAATGAG	CAACGCTGCT	TTGTGAGAGT	TGCTGCTTCA
	GATATAGATC	CAGTCAAGAC	TTGGAGGGGA	AAGTGGCCAG	CTCTGATTTG	GGTCAITTTA	GCCTAGCTAG
65	CACTTTTAT	CTCTCTTTT	GTAGCTTTTC	TTACTTTTCA	AAGAAGGATC	AAAAGGATC	AGTATATCAAG
	TGGAGGTTCA	GAGGTTGGAA	AGGATGAGCT	TGAGTGGCCA	TTGTTTGAAT	TAGTACTGTT	TACTATCTCT
	ACTGAAAGTT	TCTCTCTGCT	GATGTGTTAT	GGTGGGCTG	GCTTTGAGCA	GCTTTACAG	GGTATCTTAT
	CAGATGGACA	AGAGATAGCA	GTABAGAGC	TATCGAGGTA	TTCTGAGCAA	GGCTTTCAAG	AGTTTAAAAA
	TGAATTTGTT	TTCATTTTCA	AGCTGCAACA	TAGAAACCTT	GTCAAGCTTT	TGGGTTGCTG	CCTTGAGAGGA
70	GAGAGAAAGA	TCATATCTTA	TGAGTTTATG	CCCAATCTTA	GCTTGGAGCT	TTTCTATTTT	GATCCAGGCA
	GAAAGCTTTC	ACTTACATGG	AAGAACTGCT	TTGAAATTTG	TTGGGGAATA	TCCTGAGGCT	TTCTATATCT
	TGATCAGGAG	TCAAGATTTA	GATTTATCTA	TAGAGAGCTC	AAGACCAACA	ACTTTTACT	AGATGCTCAAC

	ATGAATGGCA	AAATTCGTGA	CTTGGCCCTT	GCCAAAATTT	TGGCGGAGA	ACAAGTGGAA	GGAATACTA
	AGAGASTGAT	AGGGACATAT	GGATAIATGT	CACCTGAGTA	TCCTGTTGAT	GGGAAATATT	CAATAAATC
	CGATGTATTC	AGTATTGGCG	TCATCATICT	AGAAGTAGTT	AGTGGCAGAA	GGAACAGGAA	ATTTCGTAT
5	TTGGAAACAIC	ATCACATATCT	TTTGGGACAT	GCAIGGTTAC	TTTGGACTGA	AGACAAAGCG	TIGGAACTGA
	TGGACGARTG	TTTGAAGAA	TCATTTGGCG	AAICTCAAGT	GTGAGATGC	ATCCAGGTTC	GTTTGCTGTC
	CGGCCAAAAA	CACCCCTGAGG	ATAGGCCCTAC	AATGGCATCA	GTAGTTTCT	GGTTGGGAAA	TGAAGGCTTC
	GTECTTCCTC	AACCAAGCA	GCCTGGATT	TTTATCGAAA	GGAATTCAT	GGATCAACA	GAATCAGCTC
	AGTITATAAA	CACGATCACA	ACAGATAGAT	CCATTAGAGA	CGGTGACACA	ATTGTTTCAG	CTGGTGGGAT
	TTATGAGCIT	GGATTTTCA	GTCTGGAA	TCGGAAGAA	CGTTACGTAG	GCATATGGTA	CAAGAGATA
10	TCAACTCAAA	CTGTGTCTG	GSTAGCAAA	AGAGATAATC	CACCTTATGA	CACATCAGGA	GTGTTAATC
	TCAGACCCAA	IGGAATTCCT	GTACTTGTTC	ATAATCCAA	TACAICATTT	TGGTCATCAA	ACTCATCAAG
	ACCGTTAAAG	GATCCAAAAG	CACGGATCCT	GGATTCCGGG	ACCTTCTGTC	ICAAATGATGG	AAATGAAAGA
	GACCTGGAAA	TAACTTCCG	AIGGCAGAGT	TTTGATTATC	CAGGAAATAC	TTTTATACCT	GGAATGAAAC
	TTGGAAGTAA	TTTGGTCACG	GGCATGGATT	GGTATATGTC	GTCTTGGAG	AGCATTTGAT	AICCTTCTCC
15	TGGTGAAIAT	ATAAACCGTC	TIGAITCTCA	TGGATACCCG	CABTTGTTTG	TGIGGAAAA	TTCAACTAIA
	GTGCTTAGCT	CAGGGATAIG	GAAAGGTAAT	GCAITTTACTG	TGAGTGTCAA	CAGTAGACCA	AATACATTT
	ACACCTTCGA	GTITATAATT	AATCAGCAGG	AAATTICATA	CCAATTCAG	CTTAAAGAAC	AGTCATGCTC
	CAGCAGGAGT	GTGCTCAACC	CGAAGGGGCT	GATAGAACAC	CTAACATGGA	TTCAGAGCAG	TTCAAGCTGG
	TTTCTGTACT	CAACAGTACA	GTGTGATAGC	TGTGGTGGTT	TTGCTTTATC	CGGTCTTAT	TCAGTTGCA
20	ACATCAATIA	CTCCCTCTCC	TGTAAGTCTT	TGCAAGGTTT	CABTCTTAGG	GTCTCTCAAC	AGTCTGACAG
	AGATTTGGTCT	CTCTGGTGGG	TGAGGAGCAC	TCTTTGGAT	TGTAAACAA	ACGGTTTCT	TAAATTTACA
	GGCATCABGA	TGCCCTGATC	TAGAACTCC	TGCTTAAIA	AGAGCATTTA	CCCTGAAAGA	TCTGAGAAAT
	IATGCTTAGC	TAATTCGAAC	TGTACAGGCT	ACTCAAACTC	TGATGTGAGA	AATGGCGGAA	GTGGATGCTT
	ACATATGGTTC	GGAGATCTCA	TGATATTTCG	AGAGTTGAGC	CAAAATGAGC	AAACCTTGTT	TGTGAGGTTT
25	GCTGCTTCAG	AAATGACAG	GAGGCAAGG	AGAAAGAIGT	CAGTCTGTAT	TGGGTCTCAT	TCAGCAGTGG
	TAGCAACAT	TATCTCTAGC	TTTTTAGCTT	GGTTTACTT	CCAAGGAAAG	AAAGAGAGAA	TAGGCTCAGA
	ATGTGAARAT	GAGGCAAGG	AGCTTCCAT	GTITGATTTA	GTACCTGTA	CTACTGCTCC	TGGGACTTTC
	CTGCTATGTA	ATGTAAGCG	GAGGAGGCTG	TTTGACCCGG	TTTACAAGGG	TATGCTACCA	AAAGGACAA
	AGATAGCAGT	CAGAGGTTA	ICAAAGCAAT	CTGGACAAAG	CTTACGACAG	TTAAAAAATG	AATTCCTTCT
30	CATITCCAA	CTGCAACACA	GGAACCTTGT	CAGCTTTTTC	GGTTGCTGCC	TTGAAAGAGA	AGAACGGAIG
	CTCATCTAIG	AGTTTATGCC	CAATGCTAGC	TTGGACTAAT	TCATTTTGA	TCCAGCAGA	AAAACCTCAC
	TTTCTATGGA	GACCCGCTTT	GAAATIGCTA	TAGGAATATC	TAGAAGTCTT	CTTTACCTTC	ACCAGGACTC
	AGATTAAGTA	ATTATTCACA	GAGATCTCAC	GACCAAGCAAC	ALCTATATAG	ATACIGACAT	GAAATGCCAA
	ATTTCTGACT	TGGGCTTGGC	CAGGATATTT	GGTGGAGATC	AGTGGGAAG	AGAACTAAG	AGTATGTAG
35	GGACATAAGG	ATATATGCTC	CCGGAGTATG	TTGTGATG	GAAATATICA	GTAAATTCG	ATGTATTCAG
	CATTTGGTGA	ATTATTTCTG	AAATAGTTAG	TCGCAAGAA	AACAGGAAT	TTCCTCATTT	GGAACATCAT
	CACATCTCT	TGGGACAAGC	ATGCTTACTT	TGGATGAA	GGAAGGCTTT	GGAATTCATG	ATGTAAGCTT
	TGAAGAGAT	ATTTCAGAA	TCTCAAGTGT	TGAGATGCAI	CCAGGTCTGG	TGTTATGCG	TCCAGAAACT
	CCAGAGGAC	AGGCCATAAA	TGGCAICAGT	GGTTTCTTGG	TTGGGAAAG	AAGGTCTGGT	CTCTCTCAA
40	CCAGGACATC	CAGTTTCTTT	CACAGAGAGG	AATCCAAATG	AATCTACTGA	TGAAGAAATG	CTAAGTAAAC
	ACGGAGCCTT	AACTGTTCTT	GAGCCAGGAT	AG			
45	SL1.00sc07408_288.1.1						
	ATGGCAATGC	TTCAAAGGTT	TCCTCTCTCT	TGCATTTTCT	TCATCTCAGC	TTCTGCTGCT	AAAGTTCAAA
	CCAGGGTAA	TGATAATCTT	GGTATGAGC	TGGTATCTGC	CCTTAATAGT	AACAGAACTG	GGAATTAAT
	ATCTCTCTTA	TACAGTAAAC	CTGCTTTGGC	ATGCTTGGCT	CTGCAATATA	TAAAGCATTA	CGGAGGTGAT
	TGTAAAGTAG	TIGGAGGCTC	AGATGGAAAG	AAAGCTGGCT	AAICTGAAT	CGCCCAAGAA	TTTCCCCCA
	ACTGTGGTGT	GCAGGCATCA	TCCTTTGCTC	AAATAAGTGG	AGGATTTCTC	GCATGTCAAT	CTAAGTATGC
50	GGAACCTTCT	GAGCATTCAC	ATGATATTTCT	TAAAGAAAT	ACCAAGAGTT	IGGATATTTCT	CTAAGTATGC
	AATCACACATG	AGGTTGGTGC	TGCTGTGAGT	GGCTCCGGTG	GIGGIGGCCC	CTATTTCTGG	TGCTGATCT
	TCAGTAAATGG	CAAAACCAAG	AGTAGCTTTT	CCACAGGTGG	AGTTGAGGCT	AAGGTAAAGT	GACCTGGATG
	CTTCTAGTGT	TCTAATGACC	AATGCAAGTG	TGCTAATACT	TTGTCTCAAA	CCATAIATCT	CTGGACAAIC
	ACTGTAGGAG	CTTCCGTTGC	ACTGCTTTAT	GCTTACGAG	TATGA		
55	SL1.00sc07408_287.1.1						
	ATGGGAAIAT	CAAGAGGCTT	TCCTTATCTT	CACCAGGATT	CCAGATTAAG	GATCATACAT	AGAGATCTAA
	AGACGAGCAA	CATTTTACTG	GACAGTGAAT	GGAATCCCAA	GAITTCAGAC	TTTGGACTGG	CTAGGATTAAT
	EGGCTGTGAC	CAAAACGAAG	CAAGAACAAA	AAGAGTAATA	GGGACATATG	GATATAIGTC	TCCAGAAATC
	GCATTTGATG	GGAATTTTTC	ATGGAATACA	GAATTTTCTA	GTCTTGGTGT	TCCTTCTGTA	GAAATAGTAA
60	GTGGAAGAAA	GACAGGAAAC	TTTCTGTCATC	CAGATCACCA	CCATAGTCTT	ATAGGAGCTG	CTTGGTTATT
	ATGGAACGTA	GGAAGGCTCT	TGGAACATA	CGATGATTTG	TTAAAGAAAT	CGTTTGTGTA	AICCCAGCTG
	CTAAGAGTGT	TTCAATGTGC	ACTGTTATGT	GTTCAGGAC	TAAAGAGTGA	AAGACCAACA	ATGTCTATCAG
	EGGTATTCAT	GTTAAGCTAT	GAAGAAAGTG	CATTAACCTC	AACAAAGGAG	CTTGGTTTCT	TCATGAGAG
	ARGTATAGCT	GAAACAGATG	ATTCAAATGA	GAAAGGCTGT	ATCAGTACAC	ATGTTTTCAG	ATTAACAAAT
65	CTTCAACCAA	GATAG					

Claims

1. A method of selecting a tomato plant, comprising exposing a tomato plant to continuous light and
- (a) determining whether said exposed plant exhibits tolerance to exposure to continuous light, wherein said tolerance is indicated by decreased level of chlorosis
- 5 upon said exposure as compared to a control plant,
- (b) determining whether said exposed plant exhibits a significant increase in yield upon exposure to continuous light as compared to a control plant, and
- (c) selecting said exposed plant in case it exhibits tolerance to exposure to continuous light and a significant increase in yield upon exposure to continuous light.
- 10
2. A method for increasing the yield of a tomato plant line, said method comprising:
- (a) selecting a tomato plant by the method according to claim 1;
- (b) crossing the thus selected plant with a plant of said tomato plant line the
- 15 yield of which is to be increased to produce offspring plants;
- (c) optionally backcrossing or selfing said offspring plant to produce further offspring plants, and
- (d) selecting from said offspring plants of step (b) or (c) a plant having tolerance to continuous light and having increased yield as compared to said plant of
- 20 said tomato plant line used in step (b).
3. Method according to claim 2, wherein said selected plant is not a plant of the variety "Subarctic plenty".
- 25 4. Method according to claim 1, wherein said determination of tolerance to continuous light in step (a) is performed by using a molecular marker.
5. Method according to claim 4, wherein said molecular marker is selected from the group consisting of markers 1-28 of Figure 11 or a polymorphism linked to
- 30 tolerance to continuous light in a gene listed in Table 3.

6. A method for producing a tomato plant that is tolerant to exposure to continuous light, said method comprising steps of:

- a) providing a donor tomato plant tolerant to exposure to continuous light,
5 preferably of the species *S. neorickii*, *S. habrochaites*, *S. pennellii*, *S. lycopersicum*, *S. peruvianum*, *S. chilense*, *S. chemielewskii*;
- b) transferring nucleic acid from said donor tomato plant to at least one recipient tomato plant which is not tolerant to exposure to continuous light, wherein said transfer results in the introduction of genomic material from said donor tomato
10 plant in the corresponding region of the genome of said at least one recipient tomato plant; and
- c) selecting from said at least one recipient tomato plant a plant that comprises within its genome a gene for tolerance to continuous light derived from said donor tomato plant, wherein said selection comprises detecting in chromosome 7 of
15 said recipient tomato plant at least one genetic marker linked to said gene for tolerance to continuous light.

7. Method according to claim 6, wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region linked to the genetic
20 markers "7-20-1" and/or "7-20-2" on chromosome 7 of *S. neorickii*, "7-10", "7-25" on chromosome 7 of *S. neorickii*, "7-17", "7-25" on chromosome 7 of *S. chilense*, "7-19", "7-25" on chromosome 7 of *S. pennellii*.

8. Method according to claim 6 or 7, wherein said transfer of nucleic acid
25 comprising at least one gene for tolerance to continuous light, or a tolerance-conferring part thereof, is performed by crossing said donor tomato plant tolerant to exposure to continuous light with a recipient tomato plant which is not tolerant to exposure to continuous light, to produce offspring plants comprising said gene as an introgression, and wherein step c) is performed on at least one offspring plant.

30

9. Method according to any one of claims 6 to 8, wherein step c) is performed by detecting said genetic marker in DNA isolated from said at least one recipient tomato plant.

10. Method according to any one of claims 6 to 9, wherein said step c) comprises subjecting said plant to a bioassay for determining its tolerance to continuous light or wherein said step c) comprises the detection of a molecular marker selected from the group consisting of markers 1-28 of Figure 11 or a polymorphism linked to tolerance to continuous light in a gene listed in Table 3.
11. A tomato plant tolerant to continuous light or a part thereof, obtainable by a method according to any one of claims 1-10.
12. An isolated gene that confers tolerance to continuous light to a tomato plant, wherein the location of said gene on chromosome 7 of a tomato plant is indicated by a genomic region linked to the genetic markers "7-20-1" and/or "7-20-2" on chromosome 7 of *S. neorickii*, "7-10" and/or "7-25" on chromosome 7 of *S. neorickii*, "7-17" and/or "7-25" on chromosome 7 of *S. chilense*, "7-19" and/or "7-25" on chromosome 7 of *S. pennellii*.
13. An isolated DNA sample obtained from a tomato plant comprising a gene according to claim 12.
14. A method for detecting a gene for tolerance to continuous light, comprising detecting at least one genetic marker linked to a gene for tolerance to continuous light on chromosome 7 of a plant suspected of harbouring said gene, wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region linked to the genetic markers "7-20-1" and/or "7-20-2" on chromosome 7 of *S. neorickii*, "7-10" and/or "7-25" on chromosome 7 of *S. neorickii*, "7-17" and/or "7-25" on chromosome 7 of *S. chilense*, "7-19" and/or "7-25" on chromosome 7 of *S. pennellii*.
15. A continuous light-tolerant tomato plant, or a part thereof, comprising within its genome a gene for tolerance to continuous light, or a tolerance-conferring part thereof, wherein the location of said gene on chromosome 7 of said plant is indicated by a genomic region linked to the genetic markers "7-20-1" and/or "7-20-2" on chromosome 7 of *S. neorickii*, "7-10" and/or "7-25" on chromosome 7 of *S. neorickii*, "7-

17" and/or "7-25" on chromosome 7 of *S. chilense*, "7-19" and/or "7-25" on chromosome 7 of *S. pennellii* wherein said gene or said continuous light-tolerance-conferring part thereof is not in its natural genetic background.

- 5 16. A method of producing a continuous light-tolerant inbred tomato plant, comprising the steps of :
- a) producing a continuous light-tolerant tomato plant according to any one of claims 1-4;
- b) crossing said continuous light-tolerant tomato plant with itself or another
10 tomato plant to yield progeny tomato seed;
- c) growing said progeny tomato seed of step b) to yield an additional continuous light-tolerant tomato plant;
- d) repeating the crossing and growing steps from 0 to 7 times to generate a continuous light-tolerant inbred tomato plant.
- 15
17. Method according to claim 16, wherein step c) further comprises the step of identifying plants that exhibit continuous light tolerance and possess commercially desirable characteristics.
- 20 18. Method according to claim 16 or 17, wherein said method further comprises the step of selecting homozygote inbred tomato plants.
19. A continuous light-tolerant inbred tomato plant, or parts thereof, obtainable the method according to any one of claims 16-18.
- 25
20. A hybrid tomato plant, or a part thereof, that exhibits tolerance to continuous light, wherein said hybrid tomato plant is obtainable by crossing a continuous light-tolerant inbred tomato plant obtainable by a method according to any one of claims 11-13 with an inbred tomato plant that exhibits commercially
30 desirable characteristics.
21. Use of a genetic marker selected from the group consisting of the genetic markers 1-28 of Figure 11 or a polymorphism linked to tolerance to continuous light

in a gene listed in Table 3 for the detection of genes for continuous light-tolerance, and/or for the detection of continuous light-tolerant tomato plants.

22. An indeterminate tomato plant, tolerant to exposure to continuous light or a
5 part thereof.

23. A tissue culture of regenerable cells of the tomato plants of any one of claims
11, 15, 19, 20 or 22, preferably said regenerable cells comprise cells or protoplasts
isolated from a tissue selected from the group consisting of leaves, pollen, embryos,
10 roots, root tips, anthers, flowers, fruits, and stems and seeds.

Figure 1




Figure 1 Continued




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6



Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

MARKER 1:

GATGAGATTGTTACATGATTTTGTGCAGATCTAGCACAAAGCCTTAGCCAAATTAAAGGAG [T/G] CCTCTCCATATT
CGGTCCCTCTGGTCATTTTCGGTCACATGCACAAACAGCTTGCTTCTG

MARKER 2:

TTTTCATTTTGC GTTTTCTGCAGGAGAGTTCAAATTGAACATCGAGTGCAACCATGCCAC [T/G] TTGGCTGGTCAC
AGGTTTGTGTGTTTTCAGTGCAAATTA ACTATGTCCGTCTTTTGCACC

MARKER 3:

GTCATTACTATGTCCACAGATGCCACAGATCTAGATCA [A/T] CCGTTACTATAGCCACAGATGCCACAGATCTAGA
TCATATTATCCTTCTTG CATTTTGTG

MARKER 4:

TAGATTGCGTCTCCTGCGCTCCGTACTGGCTTTACTTCGTGAATGGGCAC TTTCTGTCTT [T/C] GCACAGGGAATG
GGTACAAGTGTTACTGGAGCCTCTCTGATTCTTGGTGGAAC TTTATCA

MARKER 5:

TTATGACTTAGCCAACGGGAAAACATTTAAANTGGATGTTTTTGTATATTTACCTCCAAT [A/T] AAACCCTAAATG
AATTT CATGACTAATAAATGGTTCAAANCAATGATATTTTCATCTTTA

MARKER 6:

TGTTATACCGTGATTTATANCAAATGATTGGACAATAAGGCCTTTCACCAT TCTTGCTCT [T/C] GATGTCAGTGTC
ACTGAAAGTAACAGTAGTTGGAGTACCCATAATATTTGATCGGATGGA

MARKER 7:

AGGTCCCTCATTGTCTGTGTTGCTCCGCCCTGCAATTGTCCTAAATTTAACTGAACTAATG [T/C] ATGGGGAAAAAT
GAAATGGGGATTTCGTTTGTGCAGGGAGGATTTGAATCTGGTG CAGTGG

MARKER 8:

TATTTTAGCAGCTATAATGGTGGATACAATTGGTCGAAAATTCAGTGTGGCACTTATGTG [T/C] GGTTTAAGCTTC
CTGTTCC TTTTACCGCTTCTTGACCTCAACTCCCTGCTTTNACTACT

MARKER 9:

AAAGGTCAGGCTATTGTTTCTTTTGTAGTTTGTAGTAGCACAAATTAGGAACCAAATGC [A/G] TTGTGCTACAGT
AGTCTTAGTACAAGCTGGTGCAAATATACCTGCTACCGTTATGGTCCC

MARKER 10:

CGAGATGAAGAGAACCGGTGTGGCTCCTAATGTTGTCACCTATAATACGCTGATAAACGC [A/G] TATAGTCAGGTT
GGTAATTCTGAAATGGGAAGTAGGCTTTTCGAGGAGATGGCAAACAAT

MARKER 11:

TTNGAGAATCTTCCTTCTCCAGATACTCCTGCAGTATATGTGTCCAACCATCAGAGCTTT [T/C] TGGACATATATA
CATTACTTACTCTTGGGAGAACTTCAAGTTCATC

MARKER 12:

TTTCAAGAACCTTAGTGATGTTAGCCCTATTAATACCTTGGCTGGAGGNAACTTATACTC [T/G] TTGAACCTCACC
GATGACTCTGGGACCGTTCATCTTAACCTCAGGATGGTCTAGGACTAAA

MARKER 13:

GGCTGGCTTCGCCAGCCATGACGGAGTTGGAAATGGTAGTCATGGACTGGCTTGCTAATA [T/C] GTTGAAATTACC
AAAACTTTTCATGTTTTCTGGCACGGGTGGTGGTGTACTACAAAGTAC

MARKER 14:

ACTTGTGTTTAAATTGCAGATGTCATCAGATGGACCAACCCACAATGAGTTTGATTTTGA [A/G] TTTTATAGGCAAT
ACAACTGGTGAACCATATACAGTACAAACAAATGTGTATGTCAATGGT

MARKER 15:

TCAACTAAAGAAAGAACTGCAACTGAACTCCCTTCAAACTGACACGATTAAATGCCTGCT [A/T] AANAAAGTTCAC
AAAGACGAAACAGGTATAAAATGTTTTGTTATGTGAGAATTCCCTATAC

MARKER 16:

CAGCATGGGAAGGACTAGAATGTGTCTGACATGTCCTTCAAGGTATTTTTTCAGTTGTTT [A/T] TTGCAGGTTTTTA
TTACTGTACAGAGCATGATGATACATAATTAGAAGATTGGTTNTATCT

MARKER 17:

ATTCATAGCGTAATCTGTTAATCCGTATGCCCATCAATGTTGTTTGTAGTTCAGGATGTT [T/C] GGATGGATACTC
ACAGTCTTTGGCCTATTTGNTATTATTGTACTTGGAAGCTTGTTTATA

MARKER 18:

AGAACTATTGAAGAAATCAGTACATCTCCACATCCCTTTTTTGAAGAAAAAATACACC [T/G] AACAGTTTAAAA
ATGAAAATGGAAGGCATGGAGGAATTCCTAGCTAAAATCTTTAGTTAT

MARKER 19:

TTTGTCGTTTGATACTTGTATGCAATCAGCTTCAGTAGCCAGCTTCGTATATCTTTCAGA [A/T] GAAAAANATCCT
GATTTGTTACATAGTATTTCCGGTATGCACCTGTGACTGATCAGTTTAC

MARKER 20:

TGTCCTNTGATTTCTATCTCCCTGNTTTGTGTAATAAAGAAGTTTGTGTATTTGGCAGAT [T/C] GAGGCAGTAGTG
ACTAAGGCTGAGCTCAAGTATCTTGCTTTCTTGTGCAAGTCTGAGGTT

MARKER 21:

GAAGCTGAAGAAGATTGGCTTTGCATGCTAAGGAAGAGTTTCTGATTATTACCAGGTACA [A/G] CAACTTCTTTCT
TCATGAGTTTCTCTTGCGTAAAGATACAGAAATTGGAATATTGTCAT

MARKER 22:

GGAAAGCCTACTGTTGTGGAATTCTATGCCGATTGGTGTGAAGTTTGTGCGAGAATTAGCT [T/C] CAGATGTCTATA
AAGTTGAACAGCAGTACAAGTAAATCTCTTTTTTGTATTGACTCCTG

MARKER 23:

TATTACTGGTTANTCGCCATCTATTTGGTTTCAAATGCTTGGCTGCTTAGTTATTACTAA [T/C] TTTTGGTTTTGA
TTACACATAGTGGACTAACTCCTACATGAAGATGAATAGAACAGTTAT

MARKER 24:

GTTGGNTAAAAAAGGCATCCTNTTTTGCCTCCGTTTAGTATCTAAATCTGGAATCATTAT [A/T] ATATCAATTCTA
AAGTTGAGAATTTTATGTTTGGACTTCTCAAGAAAATATTGATTTTA

MARKER 25:

ATCAAGGGTTTATAACAATAAAAAAGGTAATCATCAGAAAATGATGTATAGTTGGAAAAAA [A/G] AACCTTCCAAGA
TGGTGAATCAAAGCATAAAAAATAGCTTCCTTGAATTCGTCTTTGTA

MARKER 26:

GAATTGGCTTTATGTATTTGGAATCCTTGTATTTACTGTATTTTCTTTT [T/G] CTCCAATGCAGGGGTCCTAGAT
GG

MARKER 27:

CACTCCCAAGTCTTCCATCACTCCCCCTCTT [C/A] GTAGCTCCAAATCCTCGACTCCCCATAATAACAGCACTAAG
CCCTAACCTC


MARKER 28:

ACTGGAAGATTTCTCGCATGTCAATCTAAGTATGC [G/C] GAACCTTCTGAAGCATTCAATGATATTCTTATAAGAA
ATACCAA [G/A] AGTTTGGATATTCTCTATAGCAAGAATCACACTGAGGT

Figure 12

7-14	7-16	7-17	CLT Locus	7-20	7-21	7-25	Phenotype of CLT
				LA2133-42-1			tolerant
	LA2133-42-2						segregating
				LA2133-43-1			segregating
	LA2133-42-3						segregating
	LA2133-43-3						susceptible
	LA2133-42-4						tolerant

Figure 13



INTERNATIONAL SEARCH REPORT

International application No
PCT/NL2010/050562

A. CLASSIFICATION OF SUBJECT MATTER

INV. A01H1/04 A01H5/00 A01H5/08
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MUELLER, LUKAS A. ET AL: "A snapshot of the emerging tomato genome sequence", PLANT GENOME (MADISON, WI, UNITED STATES), 2(1), 78-92 CODEN: PGMWAT; ISSN: 1940-3372, 18 March 2009 (2009-03-18), XP9128007, page 80, column 2, paragraph 4 - page 82, column 2, paragraph 5 page 86, column 2, paragraph 4 - page 87, column 1, paragraph 4; figure 1; table 1 ----- -/--	12,13

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 December 2010

Date of mailing of the international search report

16/12/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Burkhardt, Peter

INTERNATIONAL SEARCH REPORT

International application No

PCT/NL2010/050562

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TANKSLEY ET AL: "High density molecular linkage maps of the tomato and potato genomes", GENETICS, GENETICS SOCIETY OF AMERICA, AUSTIN, TX, US, vol. 132, 1 December 1992 (1992-12-01), pages 1141-1160, XP002097980, ISSN: 0016-6731 page 1142, column 1, paragraph 3 - page 1145, column 1, paragraph 1 chromosome 7; page 1150 page 1156, column 1, paragraph 2; figures 1,5	12,13
X	CHR DASKALOFF UND A OGNJANOVA: "Das Verhalten von Lycopersicon esculentum Mill, L. racemigerum Lange und L. hirsutum Humb, et Bonpl. gegenüber Dauerbelichtung", ZEITSCHRIFT FUER PFLANZENZUECHTUNG, XX, XX, vol. 54, no. 2, 1 January 1965 (1965-01-01), pages 169-181, XP009127688, ISSN: 0044-3298 cited in the application	1-11, 15-20, 22,23
Y	page 169, paragraph 1 page 173, last paragraph - page 176, paragraph 3; figures; tables	1-10,14, 16-18,21
X	NUEZ F ET AL: "GENETICS OF THE PARTHENO-CARPY FOR TOMATO VARIETIES 'SUB-ARTIC PLENTY', '75/59' AND 'SEVERIANIN'", ZEITSCHRIFT FUER PFLANZENZUECHTUNG, vol. 96, no. 3, 1 January 1986 (1986-01-01), pages 200-206, XP002043982, ISSN: 0044-3298 page 201, paragraph 3	11,15, 19,20, 22,23
X	OHYAMA KATSUMI ET AL: "Potential use of a 24-hour photoperiod (continuous light) with alternating air temperature for production of tomato plug transplants in a closed system", HORTSCIENCE, vol. 40, no. 2, April 2005 (2005-04), pages 374-377, XP9127917, ISSN: 0018-5345	11,15, 19,20, 22,23
Y	page 375, column 3, paragraph 1 - page 376, column 1, paragraph 1; table 1	1-10,14, 16-18,21
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/NL2010/050562

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/066866 A2 (METAPONTUM AGROBIOS S C R L [IT]; D AMBROSIO CATERINA [IT]; FESTA GIUS) 14 August 2003 (2003-08-14) page 8, line 18 - page 9, line 17; claims; example 6 -----	11,15, 19,20, 22,23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/NL2010/050562

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 03066866	A2	14-08-2003	AU	2003202560 A1	02-09-2003
			IT	MI20020213 A1	06-08-2003