



HU000034962T2

(19) **HU**(11) Lajstromszám: **E 034 962**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM

SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 12 178778**
(22) A bejelentés napja: **2006. 03. 23.**
- (96) Az európai bejelentés bejelentési száma:
EP 20060178778
- (97) Az európai bejelentés közzétételi adatai:
EP 2567976 A2 **2013. 03. 13.**
- (97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2567976 B1 **2017. 07. 19.**
- (51) Int. Cl.: **C07K 16/28** (2006.01)
A61P 35/00 (2006.01)
A61P 37/00 (2006.01)
G01N 33/50 (2006.01)
C12N 5/10 (2006.01)
A61K 39/395 (2006.01)
C07K 14/705 (2006.01)

- | | |
|---|--|
| (30) Elsőbbségi adatok:
200500429 2005. 03. 23. DK
667579 P 2005. 04. 01. US
696163 P 2005. 07. 01. US
728561 P 2005. 10. 20. US | (73) Jogosult(ak):
Genmab A/S, 1560 Copenhagen V (DK) |
| (72) Feltaláló(k):
Graus, Yvo, 3984 PR Odijk (NL)
Parren, Paul, 3984 PR Odijk (NL)
van de Winkel, Jan, 3707 CK Zeist (NL)
de Weers, Michel, 3991 WR Houten (NL)
Oprins, Judith, 3584 CM Utrecht (NL)
van Vugt, Martine, 3994 KR Houten (NL) | (74) Képviselő:
PINTZ ÉS TÁRSAI Szabadalmi, Védjegy és Jogi Iroda Kft., Budapest |

(54) **CD38-ellenes antitestek a mielóma multiplex kezelésére**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 567 976 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

19.07.2017 Bulletin 2017/29

(51) Int Cl.:

C07K 16/28 ^(2006.01) **C07K 14/705** ^(2006.01)
A61K 39/395 ^(2006.01) **G01N 33/50** ^(2006.01)
C12N 5/10 ^(2006.01) **A61P 35/00** ^(2006.01)
A61P 37/00 ^(2006.01)

(21) Application number: **12178778.2**

(22) Date of filing: **23.03.2006**

(54) **Antibodies against CD38 for treatment of multiple myeloma**

Antikörper gegen CD38 zur Behandlung von multiplem Myelom

Anticorps dirigés contre CD38 pour le traitement du myelome multiple

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

(74) Representative: **J A Kemp**

**14 South Square
Gray's Inn
London WC1R 5JJ (GB)**

(30) Priority: **23.03.2005 DK 200500429**

01.04.2005 US 667579 P

01.07.2005 US 696163 P

20.10.2005 US 728561 P

(56) References cited:

- **ELLIS JONATHAN H ET AL:** "Engineered anti-CD38 monoclonal antibodies for immunotherapy of multiple myeloma", JOURNAL OF IMMUNOLOGY, THE WILLIAMS AND WILKINS CO. BALTIMORE, US, vol. 155, no. 2, 1995, pages 925-937, XP002146232, ISSN: 0022-1767
- **AUSIELLO C METAL:** "Functional topography of discrete domains of human CD38.", TISSUE ANTIGENS. DEC 2000, vol. 56, no. 6, December 2000 (2000-12), pages 539-547, XP002389874, ISSN: 0001-2815
- **HOSHINO S ET AL:** "Mapping of the catalytic and epitopic sites of human CD38/NAD+ glycohydrolase to a functional domain in the carboxyl terminus.", JOURNAL OF IMMUNOLOGY (BALTIMORE, MD. : 1950) 15 JAN 1997, vol. 158, no. 2, 15 January 1997 (1997-01-15), pages 741-747, XP002389875, ISSN: 0022-1767
- **MCKAY BROWN ET AL:** "Tolerance to single, but not multiple, amino acid replacements in antibody V-H CDR2: A means of minimizing B cell wastage from somatic hypermutation?", THE JOURNAL OF IMMUNOLOGY, THE AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US, vol. 156, no. 9, 1 January 1996 (1996-01-01), pages 3285-3291, XP002649029, ISSN: 0022-1767

(43) Date of publication of application:

13.03.2013 Bulletin 2013/11

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

06722861.9 / 1 866 338

(73) Proprietor: **Genmab A/S**

1260 Copenhagen K (DK)

(72) Inventors:

- **Graus, Yvo**
3984 PR Odijk (NL)
- **Parren, Paul**
3984 PR Odijk (NL)
- **van de Winkel, Jan**
3707 CK Zeist (NL)
- **de Weers, Michel**
3991 WR Houten (NL)
- **Oprins, Judith**
3584 CM Utrecht (NL)
- **van Vugt, Martine**
3994 KR Houten (NL)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 567 976 B1

- | | |
|---|--|
| <ul style="list-style-type: none"> • HOLT L J ET AL: "Domain antibodies: proteins for therapy", TRENDS IN BIOTECHNOLOGY, ELSEVIER PUBLICATIONS, CAMBRIDGE, GB, vol. 21, no. 11, 1 November 2003 (2003-11-01), pages 484-490, XP004467495, ISSN: 0167-7799, DOI: 10.1016/J.TIBTECH.2003.08.007 • DAVIES J ET AL: "Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding", IMMUNOTECHNOLOGY, ELSEVIER SCIENCE PUBLISHERS BV, NL, vol. 2, no. 3, 1 September 1996 (1996-09-01), pages 169-179, XP004070292, ISSN: 1380-2933, DOI: 10.1016/S1380-2933(96)00045-0 | <ul style="list-style-type: none"> • RADER C ET AL: "A PHAGE DISPLAY APPROACH FOR RAPID ANTIBODY HUMANIZATION: DESIGNED COMBINATORIAL V GENE LIBRARIES", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, NATIONAL ACADEMY OF SCIENCES, US, vol. 95, 21 July 1998 (1998-07-21), pages 8910-8915, XP000941357, ISSN: 0027-8424, DOI: 10.1073/PNAS.95.15.8910 |
|---|--|

Description**FIELD OF THE INVENTION**

5 **[0001]** The present invention relates to antibodies binding CD38, which antibodies have specific characteristics and which are useful for treating inter alia multiple myeloma.

BACKGROUND

10 **[0002]** Multiple myeloma is a B cell malignancy characterized by the latent accumulation in bone marrow of secretory plasma cells with a low proliferative index and an extended life span. The disease ultimately attacks bones and bone marrow, resulting in multiple tumors and lesions throughout the skeletal system.

15 **[0003]** Approximately 1% of all cancers, and slightly more than 10% of all hematologic malignancies, can be attributed to multiple myeloma (MM). Incidence of MM increases in the aging population, with the median age at time of diagnosis being about 61 years.

20 **[0004]** Currently available therapies for multiple myeloma include chemotherapy, stem cell transplantation, Thalomid® (thalidomide), Velcade® (bortezomib), Aredia® (pamidronate), and Zometa® (zoledronic acid). Current treatment protocols, which include a combination of chemotherapeutic agents such as vincristine, BCNU, melphalan, cyclophosphamide, adriamycin, and prednisone or dexamethasone, yield a complete remission rate of only about 5%, and median survival is approximately 36-48 months from the time of diagnosis. Recent advances using high dose chemotherapy followed by autologous bone marrow or peripheral blood mononuclear cell transplantation have increased the complete remission rate and remission duration. Yet overall survival has only been slightly prolonged, and no evidence for a cure has been obtained. Ultimately, all MM patients relapse, even under maintenance therapy with interferon-alpha (IFN- α) alone or in combination with steroids.

25 **[0005]** Efficacy of the available chemotherapeutic treatment regimens for MM is limited by the low cell proliferation rate and development of multi-drug resistance. For more than 90% of MM patients, the disease becomes chemoresistant. As a result, alternative treatment regimens aimed at adoptive immunotherapy targeting surface antigens on plasma cells are being sought.

30 **[0006]** CD38 is an example of an antigen expressed on such malignant plasma cells, and is expressed in a variety of malignant hematological diseases, including but not restricted to multiple myeloma, B-cell chronic lymphocytic leukemia, B-cell acute lymphocytic leukemia, Waldenström macroglobulinemia, primary systemic amyloidosis, mantle-cell lymphoma, pro-lymphocytic/myelocytic leukemia, acute myeloid leukemia, chronic myeloid leukemia, follicular lymphoma, NK-cell leukemia and plasma-cell leukemia. Expression of CD38 has been described on epithelial/endothelial cells of different origin, including glandular epithelium in prostate, islet cells in pancreas, ductal epithelium in glands, including parotid gland, bronchial epithelial cells, cells in testis and ovary and tumor epithelium in colorectal adenocarcinoma. Diseases, where CD38 expression could be involved, include but is not restricted to broncho-epithelial carcinomas of the lung, breast cancer (evolving from malignant proliferation of epithelial lining in ducts and lobules of the breast), pancreatic tumors, evolving from the b-cells (insulinomas), tumors evolving from epithelium in the gut (e.g. adenocarcinoma and squamous cell carcinoma) In CNS, neuroblastomas express CD38. Other diseases include carcinoma in the prostate gland, seminomas in testis and ovarian cancers.

35 **[0007]** Normally, CD38 is expressed by hemopoietic cells, and in solid tissues. With regard to hemopoietic cells, the majority of medullary thymocytes are CD38⁺, resting and circulating T- and B-cells are CD38⁻, and activated cells are CD38⁺. CD38 is also expressed on approximately 80% of resting NK cells and monocytes, and on lymph node germinal center lymphoblasts, plasma B cells and some intrafollicular cells. CD38 can also be expressed by dendritic cells. A significant proportion of normal bone marrow cells, particular precursor cells, express CD38. In addition, 50-80% of umbilical cord blood cells is CD38⁺ and remains so in human blood for the first two to three years of life. In addition to lymphoid precursor cells, CD38 is also expressed on erythrocytes and on platelets.

40 **[0008]** With regard to solid tissues, CD38 is expressed in the gut by intra-epithelial cells and lamina propria lymphocytes, by Purkinje cells and neurofibrillary tangles in the brain, by epithelial cells in the prostate, β -cells in the pancreas, osteoclasts in the bone, retinal cells in the eye, and sarcolemma of smooth and striated muscle.

45 **[0009]** Functions ascribed to CD38 include both receptor mediation in adhesion and signaling events and (ecto-) enzymatic activity. As an ectoenzyme, CD38 uses NAD⁺ as substrate for the formation of cyclic ADP-ribose (cADPR) and ADPR, but also of nicotinamide and nicotinic acid-adenine dinucleotide phosphate (NAADP). cADPR has been shown to act as second messenger for Ca²⁺ mobilization from the endoplasmatic reticulum. In addition to signaling via Ca²⁺, CD38 signaling occurs via cross-talk with antigen-receptor complexes on T and B cells or other types of receptor complexes, e.g. MHC molecules, and is in this way involved in several cellular responses, but also in switching and secretion of IgG1.

50 **[0010]** Anti-CD38 antibodies are described in the literature, for instance in Lande R, et al., Cell Immunol. 220(1), 30-8

(2002), Ausiello CM, et al., Tissue Antigens. 56(6), 539-47 (2000), and Cotner T, et al., Int J Immunopharmacol. 3(3), 255-68 (1981). Ellis, J. H. et al (J. Immunol., 1995, 155: 925-937) describes a mouse anti-CD38 mAb designated AT13/5 and a humanised form of AT13/5. CD38 has a number of functions, which may or may not be activated by a molecule binding to CD38. For instance the mouse anti-CD38 antibody IB4 has agonistic properties in relation to CD38 (Ausiello CM, et al., supra). IB4 is shown to induce T cell activation as indicated by Ca^{2+} mobilization in Jurkat cells (Zubiaur M, et al., J Immunol. 159(1), 193-205 (1997), to induce significant proliferation of peripheral blood mononuclear cells (PB-MCs), to induce release of significant IL-6 levels and to induce release of detectable IFN- γ levels (Lande, Zubiaur Morra, Ansiello supra).

SUMMARY OF THE INVENTION

[0011] The present invention provides an antibody binding to human CD38, wherein said antibody comprises human light chain and human heavy chain variable regions, wherein the light chain variable region comprises a VL CDR1 having the sequence as set forth in SEQ ID No:13, a VL CDR2 having the sequence as set forth in SEQ ID No:14 and a VL CDR3 having the sequence as set forth in SEQ ID No:15, and the heavy chain variable region comprises a VH CDR1 having the sequence as set forth in SEQ ID No:18, a VH CDR2 having the sequence as set forth in SEQ ID No:19 and a VH CDR3 having the sequence as set forth in SEQ ID No:20.

[0012] Any antibody of the invention may be a human monoclonal antibody.

[0013] Any antibody of the invention may be characterized in that it is a full length IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody. The antibody may be a full length IgG1, κ antibody or a full length IgM, κ antibody.

[0014] An antibody of the invention may comprise a V_H region having the amino acid sequence as set forth in SEQ ID No:17.

[0015] An antibody of the invention may comprise a V_H region comprising the amino acid sequence spanning the V_H CDR1 - V_H CDR3 region of SEQ ID No:17.

[0016] An antibody of the invention may comprise a V_L region having the amino acid sequence as set forth in SEQ ID NO:12.

[0017] An antibody of the invention may be encoded by:

(i) human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions as set forth in SEQ ID No:11 and SEQ ID No:16, respectively; or

(ii) human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions, which are conservative sequence modifications of the sequences as set forth in (i).

[0018] An antibody of the invention may be an antibody fragment or a single chain antibody.

[0019] An antibody of the invention may be in a substantially isolated form.

[0020] An antibody of the invention may further comprise a chelator linker for attaching a radioisotope.

[0021] The present invention also provides an isolated nucleic acid encoding an antibody of the invention.

[0022] The present invention also provides an expression vector comprising a nucleic acid sequence encoding an antibody of the invention.

[0023] The present invention also provides a eukaryotic or prokaryotic host cell which produces an antibody of the invention.

[0024] The present invention also provides an immunoconjugate comprising any of the above-described antibodies linked to a cytotoxic agent, a radioisotope, or a drug.

[0025] The present invention also provides a bispecific or multispecific molecule comprising an antibody of the invention and a binding specificity for a human effector cell.

[0026] The present invention also provides a pharmaceutical composition comprising antibodies, immunoconjugates, or bispecific or multispecific molecules of the invention and a pharmaceutically acceptable carrier.

[0027] The present invention also provides antibodies, immunoconjugates, bispecific or multispecific molecules or expression vectors of the invention for use as a medicament.

[0028] The present invention also provides antibodies, immunoconjugates, pharmaceutical compositions, bispecific or multispecific molecules or expression vectors of the invention for use in a method of treating or preventing a disease or disorder involving cells expressing CD38, wherein the disease or disorder is selected from the group consisting of B cell lymphoma/leukemias including precursor B cell lymphoblastic leukemia/lymphoma and B cell non-Hodgkin's lymphomas; acute lymphoblastic leukemia; mature B cell neoplasms including B cell chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL); B cell acute lymphocytic leukemia; B cell prolymphocytic leukemia; lymphoplasmacytic lymphoma; mantle cell lymphoma (MCL); follicular lymphoma (FL) including low-grade, intermediate-grade and high-grade FL; cutaneous follicle center lymphoma; marginal zone B cell lymphoma including MALT type, nodal and splenic type; hairy cell leukemia; diffuse large B cell lymphoma; Burkitt's lymphoma; plasmacytoma; plasma cell myeloma;

plasma cell leukemia; post-transplant lymphoproliferative disorder; Waldenström's macroglobulinemia; plasma cell leukemias; anaplastic large-cell lymphoma (ALCL); multiple myeloma; Hodgkin's lymphoma; mature T cell and NK cell neoplasms including T cell prolymphocytic leukemia; T cell large granular lymphocytic leukemia; aggressive NK cell leukemia; adult T cell leukemia/lymphoma; extranodal NK/T cell lymphoma; nasal type, enteropathy-type T cell lymphoma; hepatosplenic T cell lymphoma; subcutaneous panniculitis-like T cell lymphoma; blastic NK cell lymphoma; Mycosis Fungoides/Sézary Syndrome; primary cutaneous CD30 positive T cell lymphoproliferative disorders including cutaneous anaplastic large cell lymphoma C-ALCL, lymphomatoid papulosis and borderline lesions; angioimmunoblastic T cell lymphoma, peripheral T cell lymphoma unspecified; anaplastic large cell lymphoma; acute myeloid leukemia including acute promyelocytic leukemia; and chronic myeloproliferative diseases including chronic myeloid leukemia. The disease may be B cell non-Hodgkin's lymphoma, multiple myeloma, acute lymphoblastic leukemia or B cell chronic lymphocytic leukemia. Any such method may comprise administration of one or more further therapeutic agents to the subject, such as a chemotherapeutic agent, an anti-inflammatory agent, or an immunosuppressive and/or immunomodulatory agent. The one or more further therapeutic agents may be selected from the group consisting of cisplatin, gefitinib, cetuximab, rituximab, bevacizumab, erlotinib, bortezomib, thalidomide, pamidronate, zoledronic acid, clodronate, risendronate, ibandronate, etidronate, alendronate, tiludronate, arsenic trioxide, lenalidomide, filgrastim, pegfilgrastim, sargramostim, suberoylanilide hydroxamic acid, and SCIO-469.

[0029] The invention also provides an *in vitro* method for detecting the presence of CD38 antigen, or a cell expressing CD38, in a sample comprising:

- a) contacting the sample an antibody of the invention under conditions that allow for formation of a complex between the peptide and CD38; and
- b) detecting the formation of a complex.

[0030] The invention also provides an anti-idiotypic antibody binding to an antibody of the invention.

[0031] The antibody binding to human CD38 may comprise a V_L region having at least about 90%, such as at least about 95% amino acid sequence identity to the sequence according to SEQ ID No:12.

[0032] The antibody binding to human CD38 may comprise a V_H region having at least about 90%, such as at least about 95% amino acid sequence identity to the sequence as set forth in SEQ ID No:17.

[0033] The antibody binding to human CD38 may comprise a V_H region having 1-5, such as 1-3 amino acid substitutions, deletions or additions compared to the sequence as set forth in SEQ ID No:17.

[0034] The antibody binding to human CD38 may comprising a V_L region as defined above and a V_H region as defined above.

[0035] The antibody binding to human CD38 may be one which binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), to the same degree that it binds to human CD38 (SEQ ID No:31).

[0036] The EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), may be less than 50%, such as less than 10%, less than 5% or less than 1% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

[0037] The antibody binding to human CD38 may be one which binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), to the same degree that it binds to human CD38 (SEQ ID No:31).

[0038] The EC_{50} of the binding of the antibody to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), may be less than 50%, such as less than 10%, less than 5% or less than 1% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

[0039] Said antibody may be one which binds to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32), to the same degree that it binds to human CD38 (SEQ ID No:31).

[0040] The EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) may correspond to 75% - 125% of the EC_{50} of the binding of the peptide to human CD38 (SEQ ID No:31).

[0041] Described herein is an antibody which binds to human CD38 (SEQ ID No:31) may be one which possesses the following binding characteristics: (i) binds to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), to the same degree that it binds to human CD38 (SEQ ID No:31), (ii) binds to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), to the same degree that it binds to human CD38 (SEQ ID No:31), and (iii) binds to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32), to the same degree that it binds to human CD38 (SEQ ID No:31).

[0042] The antibody which binds to human CD38 (SEQ ID No:31) may be one which possesses the following binding

characteristics: (i) does not bind to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), to the same degree that it binds to human CD38 (SEQ ID No:31), (ii) does not bind to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), to the same degree that it binds to human CD38 (SEQ ID No:31), (iii) binds to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32), to the same degree that it binds to human CD38 (SEQ ID No:31).

[0043] The EC₅₀ may be determined by use of an ELISA as described in Example 17 of the specification.

[0044] The antibody may specifically binds to the region SKRNIQFSCCKNIYR and the region EKVQTLEAWVIHGG of human CD38 (SEQ ID No:31).

[0045] The antibody binding to human CD38 may possess one or more of the following characteristics:

(i) acts as an antagonist of CD38;

(ii) does not induce significant proliferation of peripheral blood mononuclear cells as determined by the method described in Example 18 of the specification;

(iii) does not induce release of significant IL-6 by human monocytes or peripheral blood mononuclear cells as determined by the method described in Example 19 of the specification;

(iv) does not induce release of detectable IFN- γ by human T cells or peripheral blood mononuclear cells as determined by the method described in Example 20 of the specification;

(v) is internalized by CD38 expressing cells; such as internalized by CHO-CD38 cells within 5 to 15 minutes at 37°C by the method as described in Example 12 of the specification;

(vi) induces ADCC; such as with an EC₅₀ value of below 15 ng/ml, such as below 10 ng/ml in Daudi-luc cells and with an EC₅₀ value of below 75 ng/ml, such as below 50 ng/ml, 30 ng/ml or 10 ng/ml in MM cells as determined by the method described in example 5 of the specification;

(vii) induces CDC in the presence of complement; such as with an EC₅₀ value of below 5 μ g/ml, such as below 1 μ g/ml in daudi-luc or CD38-CHO cells by the method described in Example 6 of the specification;

(viii) inhibits the synthesis of cGDPR;

(ix) inhibits the synthesis of cADPR;

(x) binds to human CD38 with an affinity (K_o) of below 10⁻⁸ M, such as in the range of from 10⁻⁸ M to 10⁻¹¹ M, for example in the range of from 7 x 10⁻⁹ M to 10⁻¹⁰ M, as determined by surface plasmon resonance as described in Example 20 of the specification.

[0046] The antibody may inhibit the synthesis of cGDPR by at least 25%, such as at least 30% after 90 minutes at a concentration of 3 μ g/ml as determined by spectrophotometric method described in Example 23 of the specification.

[0047] The antibody may inhibit the synthesis of cADPR by at least 25%, such as at least 30% after 90 minutes at a concentration of 3 μ g/ml as determined by the HPLC method described in Munshi et al., J. Biol. Chem. 275, 21566-21571 (2000).

[0048] The human monoclonal antibody may comprise

(i) a heavy chain variable region amino acid sequence derived from a human Hv1263/3M28 (V_HI) germline sequence and a light chain variable region amino acid sequence derived from a human L15 (V_KI) germline sequence; or

(ii) a heavy chain variable region amino acid sequence derived from a human V_H3-DP-47/V3-23 (V_HIII) germline sequence and a light chain variable region amino acid sequence derived from a human L6 (V_KI) germline sequence.

[0049] The antibody may be glycosylated in a eukaryotic cell.

[0050] The expression vector may comprise a V_L nucleotide sequence of SEQ ID No:11 and a V_H nucleotide sequence of SEQ ID No:16. The eukaryotic host cell may comprise such an expression vector, or an expression vector comprising a V_L nucleotide sequence of SEQ ID No:11 and an expression vector encoding a V_H nucleotide sequence of SEQ ID No:16.

[0051] The expression vectors may further comprise a nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody. The nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody may encode an IgG1 antibody.

[0052] The bispecific or multispecific molecule may comprise an antibody of the invention and a binding specificity for CD3, CD4, CD138, IL-15R, membrane bound or receptor bound TNF- α , a human Fc receptor, or membrane bound or receptor bound IL-15.

[0053] The pharmaceutical composition may comprise one or more further therapeutic agents.

[0054] The one or more further therapeutic agents may be selected from a chemotherapeutic agent, an anti-inflammatory agent, or an immunosuppressive and/or immunomodulatory agent.

[0055] The one or more further therapeutic agents may be selected from a group consisting of cisplatin, gefitinib, cetuximab, rituximab, bevacizumab, erlotinib, bortezomib, thalidomide, pamidronate, zoledronic acid, clodronate, risen-

dronate, ibandronate, etidronate, alendronate, tiludronate, arsenic trioxide, lenalidomide, filgrastim, pegfilgrastim, sargramostim, suberoylanilide hydroxamic acid, and SCIO-469.

[0056] The antibody may be comprised in a kit for detecting the presence of CD38 antigen, or a cell expressing CD38, in a sample.

[0057] The anti-idiotypic antibody may be used for detecting the level of an antibody of the invention in a sample.

[0058] The anti anti-idiotypic may be used for detecting the level of human monoclonal antibody against CD38 in a sample.

BRIEF DESCRIPTION OF THE FIGURES

[0059]

Figure 1A shows the binding of -003, -005 and the isotype control antibody HuMab-KLH to CD38-transfected CHO (CHO-CD38) cells as measured by flow cytometry. The experimental setup is described in Example 4.

Figure 1 B shows the binding of -024 and HuMab-KLH to CD38-transfected CHO (CHO-CD38) cells as measured by flow cytometry. The experimental setup is described in Example 4.

Figure 2A shows the binding of -003, -005 and HuMab-KLH to Daudi cells as measured by flow cytometry. The experimental setup is described in Example 4.

Figure 2B shows the binding of -024 and HuMab-KLH to Daudi cells as measured by flow cytometry. The experimental setup is described in Example 4.

Figure 3 shows the binding of -003, -005, -024 and HuMab-KLH to multiple myeloma cells. The experimental setup is described in Example 4.

Figure 4A shows the ability of -003 and -005 to induce lysis of Daudi cells by ADCC as compared to rituximab and HuMab-KLH. The experimental setup is described in Example 5.

Figure 4B shows the ability of -024 to induce lysis of Daudi cells by ADCC as compared to HuMab-KLH. The experimental setup is described in Example 5.

Figure 5A shows the ability of -003, -005 and -024 to induce lysis of fresh multiple myeloma tumor cells by ADCC as compared to HuMab-KLH. The experimental setup is described in Example 5.

Figure 5B shows the ability of -003, -005 and -024 to induce lysis of fresh plasma cell leukemia tumor cells by ADCC as compared to HuMab-KLH. The experimental setup is described in Example 5.

Figure 6 shows the ability of -003 and -005 to induce lysis of JK6L (a multiple myeloma cell line) by ADCC as compared to HuMab-KLH. The experimental setup is described in Example 5.

Figure 7 shows the ability of -003 and -005 to induce lysis of AMO-1 (a multiple myeloma cell line) by ADCC as compared to HuMab-KLH. The experimental setup is described in Example 5.

Figure 8 shows the CDC-mediated lysis of Daudi-luc cells induced by -003 and -005 compared to HuMab-KLH. The experimental setup is described in Example 6.

Figure 9A shows the CDC-mediated lysis of CHO-CD38 cells induced by -003 and -005 compared to HuMab-KLH. The experimental setup is described in Example 6.

Figure 9B shows the CDC-mediated lysis of CHO-CD38 cells induced by -024 compared with HuMab-KLH. The experimental setup is described in Example 6.

Figure 10A shows the CDC-mediated lysis of 3% refractory tumor cells in the presence of -003, -005 and HuMab-KLH. The experimental setup is described in Example 6.

Figure 10B shows the CDC-mediated lysis of 9% refractory tumor cells in the presence of -003, -005 and HuMab-KLH. The experimental setup is described in Example 6.

Figure 10C shows the CDC-mediated lysis of 30-40% tumor cells in the presence of -003, -005 and HuMab-KLH. The experimental setup is described in Example 6.

Figure 10D shows the CDC-mediated lysis of 70% tumor cells in the presence of -003, -005 and HuMab-KLH. The experimental setup is described in Example 6.

Figure 10E shows the CDC-mediated lysis of multiple myeloma cells in the presence of -024 and HuMab-KLH. The experimental setup is described in Example 6.

Figure 11 shows that -003 and -005 do not cross-block binding to CD38. The experimental setup is described in Example 7.

Figure 12A shows the immunohistological staining of macrophages, lymphocytes and plasma B cells with -003. The experimental setup is described in Example 10.

Figure 12B shows the immunohistological staining of bronchial epithelium with -003. The experimental setup is described in Example 10.

Figure 12C shows the immunohistological staining of myocytes with -003. The experimental setup is described in Example 10.

Figure 12D shows the immunohistological staining of cynomolgus lymphoid tissue with -003. The experimental setup is described in Example 10.

Figure 13A shows the immunohistological staining of macrophages, lymphocytes and plasma B cells with -005. The experimental setup is described in Example 10.

Figure 13B shows the immunohistological staining of bronchial epithelium with -005. The experimental setup is described in Example 10.

Figure 13C shows the immunohistological staining of myocytes with -005. The experimental setup is described in Example 10.

Figure 13D shows the immunohistological staining of cynomolgus lymphoid tissue with -005. The experimental setup is described in Example 10.

Figure 14A shows immunohistological staining of liver endothelium with CD31. The experimental setup is described in Example 10.

Figure 14B shows immunohistological staining of liver endothelium with vWF. The experimental setup is described in Example 10.

Figure 14C shows immunohistological staining of liver endothelium with anti-KLH. The experimental setup is described in Example 10.

Figure 14D shows immunohistological staining of liver endothelium with -003. The experimental setup is described in Example 10.

Figure 14E shows immunohistological staining of liver endothelium with -005. The experimental setup is described in Example 10.

Figure 15A shows the cross-reactivity of -003 and -005 compared to HuMab-KLH on cynomolgus lymphocytes as measured by flow cytometry. The experimental setup is described in Example 11.

Figure 15B shows the cross-reactivity of -003 and -005 compared to HuMab-KLH on cynomolgus monocytes as measured by flow cytometry. The experimental setup is described in Example 11.

Figure 15C shows the cross-reactivity of -003 and -005 compared to HuMab-KLH on rhesus monkey PBMCs as measured by flow cytometry. The experimental setup is described in Example 11.

Figure 16A shows the internalization of -003 as measured by EtBr-quenching. The experimental setup is described in Example 12.

Figure 16B shows the internalization of -005 as measured by EtBr-quenching. The experimental setup is described in Example 12.

Figure 17A shows the inhibition caused by -003 and -005 compared to an anti-CD20 monoclonal antibody (rituximab) and HuMab-KLH of the growth of tumor cells in a preventive setting as measured by *in vivo* SCID luciferase imaging. The experimental setup is described in Example 13.

Figure 17B shows the inhibition caused by -003 and -005 compared to an anti-CD20 monoclonal antibody (rituximab) and HuMab-KLH of the growth of tumor cells in therapeutic setting I as measured by *in vivo* SCID luciferase imaging. The experimental setup is described in Example 13.

Figure 17C shows the inhibition caused by -003 and -005 compared to an anti-CD20 monoclonal antibody (rituximab) and HuMab-KLH of the growth of tumor cells in therapeutic setting II as measured by *in vivo* SCID luciferase imaging. The experimental setup is described in Example 13.

Figure 17D shows the inhibition of tumor cell growth by -003 and -024 compared to HuMab-KLH in therapeutic setting III as measured by *in vivo* SCID luciferase imaging. The experimental set up is described in Example 13.

Figure 18 shows the induction of apoptosis by -003 and -005 compared to an anti-CD20 monoclonal antibody (rituximab) and HuMab-KLH without or with crosslinking. The experimental setup is described in Example 14.

Figure 19 shows the histological score for CD38-positive cells in implanted RA-SCID mouse xenografts on day 14, after treatment with anti-KLH (HuMab-KLH) or -005. Methods are described in Example 15.

Figure 20 shows the histological score for CD138-positive cells in implanted RA-SCID mouse xenografts on day 14, after treatment with anti-KLH or -005. Methods are described in Example 15.

Figure 21 shows CD38 staining of B cells in xenografts before implantation (A), or after treatment with anti-KLH (B), or -005 (C). Methods are described in Example 15.

Figure 22 shows CD138 staining of B cells in xenografts before implantation (A), or after treatment with anti-KLH (B), or -005 (C). Methods are described in Example 15.

Figure 23 shows the binding of -003 and -005 to wild type and mutant human CD38 as measured by ELISA. 23A: Binding of -003 and -005 to T237A mutant human CD38. 23B: Binding of -003 and -005 to Q272R mutant human CD38. 23C: Binding of -003 and -005 to S274F mutant human CD38. Methods are described in Example 17.

Figure 24 shows the effect of -003 and -005 compared to HuMab-KLH on proliferation (A), IL-6 production (B) and IFN- γ production (C) of human PBMCs. Methods are described in Examples 18, 19 and 20, respectively.

Figure 25 shows the enzymatic production of cGDPribose in the presence of various concentrations of -003 (B), -005 (C), -024 (D) or anti-KLH (A). Methods are described in Example 23.

Figure 26 shows the comparison between -003, -005 and Morphosys antibody TH-3079 in CDC of CHO-CD38 cells (26A), CDC of Daudi cells (26B), and ADCC of Daudi cells (26C).

SEQ ID No:1 is the nucleotide sequence of the V_L region of the antibody -003.
 SEQ ID No:2 is the amino acid sequence of the V_L region of the antibody -003.
 SEQ ID No:3 is the amino acid sequence of the V_L CDR1 of the antibody -003 comprising aa 24-34 of SEQ ID No:2.
 SEQ ID No:4 is the amino acid sequence of the V_L CDR2 of the antibody -003 comprising aa 50-56 of SEQ ID No:2.
 SEQ ID No:5 is the amino acid sequence of the V_L CDR3 of the antibody -003 comprising aa 89-97 of SEQ ID No:2.
 SEQ ID No:6 is the nucleotide sequence of the V_H region of the antibody -003.
 SEQ ID No:7 is the amino acid sequence of the V_H region of the antibody -003.
 SEQ ID No:8 is the amino acid sequence of the V_H CDR1 of the antibody -003 comprising aa 31-35 of SEQ ID No:7.
 SEQ ID No:9 is the amino acid sequence of the V_H CDR2 of the antibody -003 comprising aa 50-66 of SEQ ID No:7.
 SEQ ID No:10 is the amino acid sequence of the V_H CDR3 of the antibody -003 comprising aa 99-109 of SEQ ID No:7.
 SEQ ID No:11 is the nucleotide sequence of the V_L region of the antibody -005.
 SEQ ID No:12 is the amino acid sequence of the V_L region of the antibody -005.
 SEQ ID No:13 is the amino acid sequence of the V_L CDR1 of the antibody -005 comprising aa 24-34 of SEQ ID No: 12.
 SEQ ID No:14 is the amino acid sequence of the V_L CDR2 of the antibody -005 comprising aa 50-56 of SEQ ID No:12.
 SEQ ID No:15 is the amino acid sequence of the V_L CDR3 of the antibody -005 comprising aa 89-97 of SEQ ID No:12.
 SEQ ID No: 16 is the nucleotide sequence of the V_H region of the antibody -005.
 SEQ ID No:17 is the amino acid sequence of the V_H region of the antibody -005.
 SEQ ID No:18 is the amino acid sequence of the V_H CDR1 of the antibody -005 comprising aa 31-35 of SEQ ID No:17.
 SEQ ID No:19 is the amino acid sequence of the V_H CDR2 of the antibody -005 comprising aa 50-66 of SEQ ID No:17.
 SEQ ID No:20 is the amino acid sequence of the V_H CDR3 of the antibody -005 comprising aa 99-111 of SEQ ID No:17.
 SEQ ID No:21 is the nucleotide sequence of the V_L region of the antibody -024.
 SEQ ID No:22 is the amino acid sequence of the V_L region of the antibody -024.
 SEQ ID No:23 is the amino acid sequence of the V_L CDR1 of the antibody -024 comprising aa 24-34 of SEQ ID No:22.
 SEQ ID No:24 is the amino acid sequence of the V_L CDR2 of the antibody -024 comprising aa 50-56 of SEQ ID No:22.
 SEQ ID No:25 is the amino acid sequence of the V_L CDR3 of the antibody -024 comprising aa 89-97 of SEQ ID No:22.
 SEQ ID No:26 is the nucleotide sequence of the V_H region of the antibody -024.
 SEQ ID No:27 is the amino acid sequence of the V_H region of the antibody -024.
 SEQ ID No:28 is the amino acid sequence of the V_H CDR1 of the antibody -024 comprising aa 31-35 of SEQ ID No:27.
 SEQ ID No:29 is the amino acid sequence of the V_H CDR2 of the antibody -024 comprising aa 50-66 of SEQ ID No:27.
 SEQ ID No:30 is the amino acid sequence of the V_H CDR3 of the antibody -024 comprising aa 99-111 of SEQ ID No:27.
 SEQ ID No:31 is the sequence of human CD38.
 SEQ ID No:32 is the sequence of a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue.
 SEQ ID No:33 is the sequence of a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue.
 SEQ ID No:34 is the sequence of a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue.

DETAILED DESCRIPTION

[0060] The present invention provides anti-CD38 antibodies, which may be useful in the treatment, diagnosis and prevention of a variety of disorders involving cells expressing CD38, such as multiple myeloma. One embodiment in accordance with the invention is the antibody -005. -005 is a human monoclonal IgG1 antibody having a V_L region consisting of the sequence of SEQ ID No:12; and a V_H region consisting of the sequence of SEQ ID No:17.

[0061] Also described herein are CD38 binding peptides ("CD38BPs"), which may be useful in the treatment, diagnosis and prevention of a variety of disorders involving cells expressing CD38, such as multiple myeloma.

[0062] Other CD38BPs include the antibody -003 and the antibody -024. -003 is a human monoclonal IgG1 antibody having a V_L region consisting of the sequence of SEQ ID No:2 and a V_H region consisting of the sequence of SEQ ID No:7. -024 is a human monoclonal IgG1 antibody having a V_L region consisting of the sequence of SEQ ID No:22 and a V_H region consisting of the sequence of SEQ ID No:27.

[0063] Antibodies interact with target antigens primarily through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see for instance Riechmann, L. et al., Nature 332, 323-327 (1998), Jones, P. et al., Nature 321, 522-525 (1986) and Queen, C. et al., PNAS USA 86, 10029-10033 (1989)).

[0064] It is well known in the art that antibody heavy chain CDR3 domains play a particularly important role in the binding specificity/affinity of an antibody for an antigen (Ditzel HJ, et al., J Immunol. 157(2), 739-49 (1996), Barbas SM et al., J. Am. Chem. Soc. 116, 2161-2162 (1994), and Barbas SM et al., Proc Natl Acad Sci USA 92(7), 2529-33 (1995).

[0065] Described herein are CD38BPs that comprise the heavy chain CDR3s of -003, -005 or -024, the heavy and light chain CDR3s of -003, -005 or -024, the CDR2s of -003, -005 and -024, and/or the CDR1s of -003, -005 and -024.

[0066] Also described herein are CD38BPs, which compete with -003, -005 or -024 for binding to CD38.

[0067] The competition may be determined by use of an ELISA or a FACS as described in the Examples section.

[0068] Described herein are a CD38BP that specifically binds to a CD38 epitope, which epitope is also specifically bound by -003 or -005 or -024 and a CD38BP having substantially the same specific binding characteristics for binding human CD38 as -003 or -005 or -024.

[0069] The antibodies of the invention comprise a V_L CDR3 having the sequence set forth in SEQ ID No:15 and a V_L CDR2 having the sequence set forth in SEQ ID No:14 and a V_L CDR1 having the sequence set forth in SEQ ID No:13.

[0070] The antibodies of the invention comprise a V_H CDR3 having the sequence set forth in SEQ ID No:20 and a V_H CDR2 having the sequence set forth in SEQ ID No:19 and a V_H CDR1 having the sequence set forth in SEQ ID No:18.

[0071] The antibodies of the invention comprise

(a) a first V_L region comprising three V_L CDRs, which have the sequences set forth in SEQ ID No:13, SEQ ID No:14, and SEQ ID No:15; and

(b) a first V_H region comprising three V_H CDRs, which have the sequences set forth in SEQ ID No:18, SEQ ID No:19, and SEQ ID No:20.

[0072] In one embodiment, the V_L region and the V_H region are present on the same chain in the antibody.

[0073] In a further embodiment, the V_L region and the V_H region are separated by a flexible linker.

[0074] In one embodiment, the V_L region and the V_H region are present on the separate chains in the antibody.

[0075] In a further embodiment, the V_L region and the V_H region are present on the separate chains in the antibody in the context of an immunoglobulin fold protein.

[0076] In one embodiment, the first V_L region and the first V_H region are oriented such that the three CDRs in the V_L region and the three CDRs in the V_H region cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on human CD38.

[0077] In a further embodiment, the antibody comprises a second V_L region identical to the first V_L region and a second V_H region identical to the first V_H region, where the second V_L region and the second V_H region cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on human CD38.

[0078] The antibody of the invention may comprise a V_L region that is a functional variant of the V_L region of -005.

[0079] In one embodiment, the V_L region of the antibody consists essentially of a sequence having at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:12. The antibody of the invention may comprise a V_H region that is a functional variant of the V_H region of -005.

[0080] In one embodiment, the V_H region of the antibody consists essentially of a sequence having at least about

50%, at least 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:17.

[0081] The CD38BPs may bind to human CD38 with greater affinity than -003 or -005 or -024 or may have substantially the same specific CD38 binding characteristics as -003 or -005 or -024.

[0082] The CD38BP may be substantially free of other CD38 binding peptides.

[0083] The antibodies of the present invention may be human antibodies, humanized antibodies or chimeric antibodies.

[0084] In one embodiment, the antibody of the present invention is a monoclonal antibody.

[0085] In one embodiment, the antibody of the present invention is an IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody. In a further embodiment, the antibody is an IgG1 antibody. In a further embodiment, the antibody is a IgG1, κ antibody. In another further embodiment, the antibody is an IgM antibody. In a further embodiment, the antibody is an IgM, κ antibody.

[0086] In one embodiment, the antibody of the present invention is an antibody fragment or a single chain antibody.

[0087] In one embodiment, the antibody is glycosylated in a eukaryotic cell.

[0088] In one embodiment, the antibody further comprises a chelator linker for attaching a radioisotope.

[0089] In one embodiment, the antibody is in a substantially isolated form.

[0090] The present invention provides an isolated nucleic acid encoding a CD38 antibody of the present invention.

[0091] The present invention provides an expression vector comprising a nucleic acid sequence encoding a CD38 antibody of the present invention.

[0092] Also disclosed herein is an expression vector which comprises a V_L nucleotide sequence of SEQ ID No:1, a V_H nucleotide sequence of SEQ ID No:6, or a V_L nucleotide sequence of SEQ ID No:1 and a V_H nucleotide sequence of SEQ ID No:6.

[0093] An antibody of the invention may be produced using an expression vector which comprises a V_L nucleotide sequence of SEQ ID No:11 and/or an expression vector which comprises a V_H nucleotide sequence of SEQ ID No:16. In one embodiment, an expression vector of the invention comprises a V_L nucleotide sequence of SEQ ID No:11 and a V_H nucleotide sequence of SEQ ID No:16.

[0094] Also described herein is an expression vector which comprises a V_L nucleotide sequence of SEQ ID No:21, a V_H nucleotide sequence of SEQ ID No:26, or a V_L nucleotide sequence of SEQ ID No:21 and a V_H nucleotide sequence of SEQ ID No:26.

[0095] In a further embodiment, the expression vector further comprises a nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody.

[0096] In a further embodiment, the nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody encodes a IgG1 antibody.

[0097] Described herein is a hybridoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in the variable light chain region as set forth in SEQ ID No:1, or conservative sequence modifications thereof, and nucleotide sequences in the variable heavy chain region as set forth in SEQ ID No:6, or conservative sequence modifications thereof. In one embodiment, the human light chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:1, and the human heavy chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:6.

[0098] Described herein is a hybridoma which produces a human monoclonal anti-CD38 antibody having human heavy chain and light chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:2, or conservative sequence modifications thereof, and the human light chain variable amino sequence as set forth in SEQ ID No:7, or conservative sequence modifications thereof. In one embodiment, the human light chain variable region comprises an amino acid sequence as set forth in SEQ ID No:2, and the human heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID No:7.

[0099] Described herein is a transfectoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain variable nucleic acids as set forth in SEQ ID No:1, or conservative sequence modifications thereof, and human heavy chain nucleic acids as set forth SEQ ID No:6, or conservative sequence modifications thereof. In one embodiment, the human monoclonal anti-CD38 antibody is encoded by human light chain variable nucleic acids as set forth in SEQ ID No:1, and human heavy chain nucleic acids as set forth SEQ ID No:6.

[0100] Described herein is a transfectoma which produces a human monoclonal anti-CD38 antibody having human light chain and heavy chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:2, or conservative sequence modifications thereof, and the human heavy chain variable amino sequence as set forth in SEQ ID No:7, or conservative sequence modifications thereof. In one embodiment, the human light chain comprises the human light chain variable amino acid sequence as set forth in SEQ ID No:2, and the human heavy chain comprises the human heavy chain variable amino sequence as set forth in SEQ ID No:7.

[0101] The antibody of the invention may be produced by a hybridoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in the variable light chain region as set forth in SEQ ID No:11, or conservative sequence modifications thereof, and

nucleotide sequences in the variable heavy chain region as set forth in SEQ ID No:16, or conservative sequence modifications thereof. In one embodiment, the human light chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:11, and the human heavy chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:16.

[0102] The antibody of the invention may be produced by a hybridoma which produces a human monoclonal anti-CD38 antibody having human heavy chain and light chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:12, or conservative sequence modifications thereof, and the human light chain variable amino sequence as set forth in SEQ ID No:17, or conservative sequence modifications thereof. In one embodiment, the human light chain variable region comprises an amino acid sequence as set forth in SEQ ID No:12, and the human heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID No:17.

[0103] The antibody of the invention may be produced by a transfectoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain variable nucleic acids as set forth in SEQ ID No:11, or conservative sequence modifications thereof, and human heavy chain nucleic acids as set forth SEQ ID No:16, or conservative sequence modifications thereof. In one embodiment, the human monoclonal anti-CD38 antibody is encoded by human light chain variable nucleic acids as set forth in SEQ ID No:11, and human heavy chain nucleic acids as set forth SEQ ID No:16.

[0104] The antibody of the invention may be produced by a transfectoma which produces a human monoclonal anti-CD38 antibody having human light chain and heavy chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:12, or conservative sequence modifications thereof, and the human heavy chain variable amino sequence as set forth in SEQ ID No:17, or conservative sequence modifications thereof. In one embodiment, the human light chain comprises the human light chain variable amino acid sequence as set forth in SEQ ID No:12, and the human heavy chain comprises the human heavy chain variable amino sequence as set forth in SEQ ID No:17.

[0105] Described herein is a hybridoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in the variable light chain region as set forth in SEQ ID No:21, or conservative sequence modifications thereof, and nucleotide sequences in the variable heavy chain region as set forth in SEQ ID No:26, or conservative sequence modifications thereof. In one embodiment, the human light chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:21, and the human heavy chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:26.

[0106] Described herein is a hybridoma which produces a human monoclonal anti-CD38 antibody having human heavy chain and light chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:22, or conservative sequence modifications thereof, and the human light chain variable amino sequence as set forth in SEQ ID No:27, or conservative sequence modifications thereof. In one embodiment, the human light chain variable region comprises an amino acid sequence as set forth in SEQ ID No:22, and the human heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID No:27.

[0107] Described herein is a transfectoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain variable nucleic acids as set forth in SEQ ID No:21, or conservative sequence modifications thereof, and human heavy chain nucleic acids as set forth SEQ ID No:26, or conservative sequence modifications thereof. In one embodiment, the human monoclonal anti-CD38 antibody is encoded by human light chain variable nucleic acids as set forth in SEQ ID No:21, and human heavy chain nucleic acids as set forth SEQ ID No:26.

[0108] Described herein is a transfectoma which produces a human monoclonal anti-CD38 antibody having human light chain and heavy chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:22, or conservative sequence modifications thereof, and the human heavy chain variable amino sequence as set forth in SEQ ID No:27, or conservative sequence modifications thereof. In one embodiment, the human light chain comprises the human light chain variable amino acid sequence as set forth in SEQ ID No:22, and the human heavy chain comprises the human heavy chain variable amino sequence as set forth in SEQ ID No:27.

[0109] The present invention provides a eukaryotic or prokaryotic host cell which produces an anti-CD38 antibody of the present invention.

[0110] The present invention provides a eukaryotic or prokaryotic host cell containing an expression vector of the present invention.

[0111] Described herein is a transgenic non-human animal or plant comprising nucleic acids encoding a human heavy chain and a human light chain, wherein the animal or plant produces a detectable amount of a CD38BP as described herein.

[0112] The present invention provides an immunoconjugate comprising an anti-CD38 antibody of the present invention linked to a cytotoxic agent, a radioisotope, or a drug. In one embodiment, the antibody is a monomeric IgM antibody linked to a cytotoxic agent, a radioisotope, or a drug.

[0113] The present invention provides a bispecific or multispecific molecule comprising an anti-CD38 antibody of the present invention and a binding specificity for a human effector cell. In one embodiment, the binding specificity for a

human effector cell is a binding specificity for CD3, CD4, CD138, IL-15R, membrane bound or receptor bound TNF- α , a human Fc receptor, or membrane bound or receptor bound IL-15.

[0114] The present invention provides an anti-idiotypic antibody binding to an anti-CD38 antibody of the present invention.

[0115] The anti-idiotypic antibody of the present invention is useful for detecting the level of human monoclonal antibody against CD38 in a sample.

[0116] The following is a list of selected embodiments.

Embodiment 1: An antibody binding to human CD38 encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions as set forth in SEQ ID No: 11 and SEQ ID No:16, respectively, or conservative sequence modifications thereof.

Embodiment 2: An antibody binding to human CD38 encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions as set forth in SEQ ID No: 11 and SEQ ID No:16, respectively.

Embodiment 3: An antibody comprising

(a) a first V_L region comprising three V_L CDRs, which independently of each other consist essentially of SEQ ID No:13, SEQ ID No:14, and SEQ ID No:15; and

(b) a first V_H region comprising three V_H CDRs, which independently of each other consist essentially of SEQ ID No:18, SEQ ID No:19, and SEQ ID No:20.

Embodiment 4: An antibody according to embodiment 3, wherein the V_L region and the V_H region are present on the same chain in the antibody.

Embodiment 5: An antibody according to embodiment 4, wherein the V_L region and the V_H region are separated by a flexible linker.

Embodiment 6: An antibody according to embodiment 3, wherein the V_L region and the V_H region are present on the separate chains in the antibody.

Embodiment 7: An antibody according to embodiment 6, wherein the V_L region and the V_H region are present on the separate chains in the peptide in the context of an immunoglobulin fold protein.

Embodiment 8: An antibody according to any of embodiments 3 to 7, wherein the first V_L region and the first V_H region are oriented such that the three CDRs in the V_L region and the three CDRs in the V_H region cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on human CD38.

Embodiment 9: An antibody according to embodiment 8, wherein the antibody comprises a second V_L region identical to the first V_L region and a second V_H region identical to the first V_H region, where the second V_L region and the second V_H region cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on human CD38.

Embodiment 10: An antibody according to embodiment 3 comprising a V_L region that is a functional variant of the V_L region of an antibody of embodiment 2.

Embodiment 11: An antibody according to embodiment 10, wherein the V_L region of the peptide consists essentially of a sequence having at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:12.

Embodiment 12: An antibody according to embodiment 3 comprising a V_H region that is a functional variant of the V_H region of an antibody of embodiment 2.

Embodiment 13: An antibody according to embodiment 12, wherein the V_H region of the antibody consists essentially of a sequence having at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:17.

Embodiment 14: An antibody according to any of embodiments 3 to 13, which antibody specifically binds human CD38.

Embodiment 15: An antibody according to any of embodiments 3 to 14, which binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34) to the same degree that it binds to human CD38 (SEQ ID No:31).

Embodiment 16: An antibody according to embodiment 15, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 50% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 17: An antibody according to embodiment 16, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue

(SEQ ID No:34), is less than 10% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 18: An antibody according to embodiment 17, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 5% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 19: An antibody according to embodiment 18, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 1 % of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 20: An antibody which binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33) to the same degree that it binds to human CD38 (SEQ ID No:31).

Embodiment 21: An antibody according to embodiment 20, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 50% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 22: An antibody according to embodiment 21, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 10% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 23: An antibody according to any of embodiments 15 to 19 which does not bind to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33) to the same degree that it binds to human CD38 (SEQ ID No:31).

Embodiment 24: An antibody according to embodiment 23, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 50% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 25: An antibody according to embodiment 24, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 10% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 26: An antibody according to any of embodiments 15 to 25, wherein said antibody binds to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with a alanine residue (SEQ ID No:32) to the same degree that it binds to human CD38 (SEQ ID No:31).

Embodiment 27: An antibody according to embodiment 26, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with a alanine residue (SEQ ID No:32) is more than 75% of the EC_{50} of the antibody of the peptide to human CD38 (SEQ ID No:31).

Embodiment 28: An antibody according to embodiment 27 wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with a alanine residue (SEQ ID No:32) is more than 85% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 29: An antibody according to embodiment 28, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with a alanine residue (SEQ ID No:32) is more than 90% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 30: An antibody according to embodiment 29, wherein the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with a alanine residue (SEQ ID No:32) is more than 95% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

Embodiment 31: An antibody according to any of embodiments 1 to 30, wherein the antibody is not an agonist of CD38.

Embodiment 32: An antibody according to any of embodiments 1 to 31, wherein the antibody does not induce significant proliferation of peripheral blood mononuclear cells.

Embodiment 33: An antibody according to any of embodiments 1 to 32, wherein the antibody does not induce release of significant IL-6 by human monocytes or peripheral blood mononuclear cells.

Embodiment 34: An antibody according to any of embodiments 1 to 33, wherein the antibody does not induce release of detectable IFN- γ by human T cells or peripheral blood mononuclear cells.

Embodiment 35: An antibody according to any of embodiments 1 to 34, which antibody is a human antibody.

Embodiment 36: An antibody according to any of embodiments 1 to 34, which antibody is a humanized or chimeric antibody.

Embodiment 37: An antibody according to any of embodiments 1 to 36, which antibody is a monoclonal antibody.

Embodiment 38: An antibody according to any of embodiments 1 to 37, characterized in that it is an IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody.

Embodiment 39: An antibody according to embodiment 38, characterized in that it is an IgG1 antibody.

Embodiment 40: An antibody according to embodiment 39, wherein the antibody is a IgG1, κ antibody.

Embodiment 41: An antibody according to embodiment 38, characterized in that it is an IgM antibody.

Embodiment 42: An antibody according to embodiment 41, wherein the antibody is a IgM, κ antibody.

Embodiment 43: An antibody according to any of embodiments 1 to 42, wherein the antibody is glycosylated in a

eukaryotic cell.

Embodiment 44: An antibody according to any of embodiments 1 to 43, which is an antibody fragment or a single chain antibody.

Embodiment 45: An antibody according to any of embodiments 1 to 44, further comprising a chelator linker for attaching a radioisotope.

Embodiment 46: An antibody according to any of embodiments 1 to 45, which is in a substantially isolated form.

Embodiment 47: An isolated nucleic acid encoding an antibody according to any of embodiments 1 to 45.

Embodiment 48: An expression vector comprising a nucleic acid sequence encoding an antibody according to any of embodiments 1 to 45.

Embodiment 49: An expression vector comprising a V_L nucleotide sequence of SEQ ID No:11, a V_H nucleotide sequence of SEQ ID No:16, or a V_L nucleotide sequence of SEQ ID No:11 and a V_H nucleotide sequence of SEQ ID No:16.

Embodiment 50: An expression vector according to embodiment 49, further comprising a nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody.

Embodiment 51: An expression vector according to embodiment 50, wherein the nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of a human antibody encodes a IgG1 antibody.

Embodiment 52: A hybridoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain and human heavy chain nucleic acids comprising nucleotide sequences in the variable light chain region as set forth in SEQ ID No:11, or conservative sequence modifications thereof, and nucleotide sequences in the variable heavy chain region as set forth in SEQ ID No:16, or conservative sequence modifications thereof.

Embodiment 53: A hybridoma according to embodiment 54, wherein the human light chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:11, and the human heavy chain nucleic acids comprises a nucleotide sequence as set forth in SEQ ID No:16.

Embodiment 54: A hybridoma which produces a human monoclonal anti-CD38 antibody having human heavy chain and light chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:12, or conservative sequence modifications thereof, and the human heavy chain variable amino sequence as set forth in SEQ ID No:17, or conservative sequence modifications thereof.

Embodiment 55: A hybridoma according to embodiment 54, wherein the human light chain variable region comprises an amino acid sequence as set forth in SEQ ID No:12, and the human heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID No: 17.

Embodiment 56: A transfectoma which produces a human monoclonal anti-CD38 antibody encoded by human light chain variable nucleic acids as set forth in SEQ ID No:11, or conservative sequence modifications thereof, and human heavy chain nucleic acids as set forth SEQ ID No:16, or conservative sequence modifications thereof.

Embodiment 57: A transfectoma according to embodiment 56, wherein the human monoclonal anti-CD38 antibody is encoded by human light chain variable nucleic acids as set forth in SEQ ID No:11, and human heavy chain nucleic acids as set forth SEQ ID No:16.

Embodiment 58: A transfectoma which produces a human monoclonal anti-CD38 antibody having human light chain and heavy chain variable regions which comprise the human light chain variable amino acid sequence as set forth in SEQ ID No:12, or conservative sequence modifications thereof, and the human heavy chain variable amino sequence as set forth in SEQ ID No:17, or conservative sequence modifications thereof.

Embodiment 59: A transfectoma according to embodiment 58, wherein the human light chain comprises the human light chain variable amino acid sequence as set forth in SEQ ID No:12, and the human heavy chain comprises the human heavy chain variable amino sequence as set forth in SEQ ID No:17.

Embodiment 60: A eukaryotic or prokaryotic host cell which produces a peptide according to any of embodiments 1 to 45.

Embodiment 61: A eukaryotic or prokaryotic host cell containing an expression vector according to embodiment 48.

Embodiment 62: A transgenic non-human animal or plant comprising nucleic acids encoding a human heavy chain and a human light chain, wherein the animal or plant produces a detectable amount of an antibody according to any of embodiments 1 to 45.

Embodiment 63: An immunoconjugate comprising an antibody according to any of embodiments 1 to 44 linked to a cytotoxic agent, a radioisotope, or a drug.

Embodiment 64: An immunoconjugate comprising an antibody according to any of embodiments 1 to 38 or embodiments 41 to 44, wherein the antibody is a monomeric IgM antibody linked to a cytotoxic agent, a radioisotope, or a drug.

Embodiment 65: A bispecific or multispecific molecule comprising an antibody according to any of embodiments 1 to 45 and a binding specificity for a human effector cell.

Embodiment 66: A bispecific or multispecific molecule comprising an antibody according to any of embodiments 1

to 45 and a binding specificity for CD3, CD4, IL-15R, membrane bound or receptor bound TNF- α , a human Fc receptor, or membrane bound or receptor bound IL-15.

Embodiment 67: A pharmaceutical composition comprising an antibody according to any of embodiments 1 to 46 or an immunoconjugate according to any of embodiments 63 to 66 and a pharmaceutically acceptable carrier.

Embodiment 68: A pharmaceutical composition according to embodiment 67 comprising one or more further therapeutic agents.

Embodiment 69: An *in vitro* method of inhibiting growth and/or proliferation of a cell expressing CD38, comprising administration of an antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 50 to 53, such that the growth and/or proliferation of the cell is inhibited.

Embodiment 70: An antibody according to any of embodiments 1 to 48, an immunoconjugate according to any of embodiments 65 to 68, a pharmaceutical composition according to embodiment 69 or 70, or an expression vector according to any of embodiments 50 to 53 for use in a method of treating a disease or disorder involving cells expressing CD38.

Embodiment 71: An antibody according to any of embodiments 1 to 48, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use in a method of preventing a disease or disorder involving cells expressing CD38 in a subject.

Embodiment 72: An antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use according to embodiment 70 or embodiment 71, wherein the disease or disorder is rheumatoid arthritis.

Embodiment 73: An antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use according to embodiment 70 or embodiment 71, wherein the disease or disorder is multiple myeloma.

Embodiment 74: An antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use according to any of embodiments 70 to 73, wherein the method comprises administration of one or more further therapeutic agents to the subject.

Embodiment 75: An antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use according to embodiment 74, wherein the one or more further therapeutic agents are selected from a chemotherapeutic agent, an anti-inflammatory agent, or an immunosuppressive and/or immunomodulatory agent.

Embodiment 76: An antibody according to any of embodiments 1 to 46, an immunoconjugate according to any of embodiments 63 to 66, a pharmaceutical composition according to embodiment 67 or 68, or an expression vector according to any of embodiments 48 to 51 for use according to embodiment 74, wherein the one or more further therapeutic agents are selected from a group consisting of cisplatin, gefitinib, cetuximab, rituximab, bevacizumab, erlotinib, bortezomib, thalidomide, pamidronate, zoledronic acid, clodronate, risendronate, ibandronate, etidronate, alendronate, tiludronate, arsenic trioxide, lenalidomide, filgrastim, pegfilgrastim, sargramostim, suberoylanilide hydroxamic acid, and SCIO-469.

Embodiment 77: An *in vitro* method for detecting the presence of CD38 antigen, or a cell expressing CD38, in a sample comprising:

- a) contacting the sample with an antibody according to any of embodiments 1 to 46 under conditions that allow for formation of a complex between the antibody and CD38; and
- b) detecting the formation of a complex.

Embodiment 78: A kit for detecting the presence of CD38 antigen, or a cell expressing CD38, in a sample comprising an antibody according to any of embodiments 1 to 46.

Embodiment 79: An anti-idiotypic antibody binding to an antibody according to any of embodiments 1 to 44.

Embodiment 80: Use of an anti-idiotypic antibody according to embodiment 79 for detecting the level of an antibody according to any of embodiments 1 to 44 in a sample.

[0117] The terms "CD38" and "CD38 antigen" are used interchangeably herein, and include any variants, isoforms and species homologs of human CD38, which are naturally expressed by cells or are expressed on cells transfected with the CD38 gene. Synonyms of CD38, as recognized in the art, include ADP ribosyl cyclase 1, cADPr hydrolase 1,

Cd38-rs1, Cyclic ADP-ribose hydrolase 1, I-19, NIM-R5 antigen.

[0118] The term peptide with respect to both CD38-binding peptides and non-CD38 peptides described herein includes any suitable peptide and can be used synonymously with the terms polypeptide and protein, unless otherwise stated or contradicted by context; provided that the reader recognize that each type of respective amino acid polymer-containing molecule can be associated with significant differences and thereby form individual embodiments (for example, a peptide such as an antibody, which is composed of multiple polypeptide chains, is significantly different from, for example, a single chain antibody, a peptide immunoadhesin, or single chain immunogenic peptide). Therefore, the term peptide herein should generally be understood as referring to any suitable peptide of any suitable size and composition (with respect to the number of amino acids and number of associated chains in a protein molecule). Moreover, peptides in the context of the inventive methods and compositions described herein may comprise non-naturally occurring and/or non-L amino acid residues, unless otherwise stated or contradicted by context.

[0119] As will be discussed further herein, unless otherwise stated or contradicted by context, the term peptide (and if discussed as individual embodiments of the term(s) polypeptide and/or protein) also encompasses derivatized peptide molecules. Briefly, a derivative is a peptide in which one or more of the amino acid residues of the peptide have been chemically modified (for instance by alkylation, acylation, ester formation, or amide formation) or associated with one or more non-amino acid organic and/or inorganic atomic or molecular substituents (for instance a polyethylene glycol (PEG) group, a lipophilic substituent (which optionally may be linked to the amino acid sequence of the peptide by a spacer residue or group such as β -alanine, γ -aminobutyric acid (GABA), L/D-glutamic acid, succinic acid, and the like), a fluorophore, biotin, a radionuclide, etc.) and may also or alternatively comprise non-essential, non-naturally occurring, and/or non-L amino acid residues, unless otherwise stated or contradicted by context (however, it should again be recognized that such derivatives may, in and of themselves, be considered independent features and inclusion of such molecules within the meaning of peptide is done for the sake of convenience in describing the present disclosure rather than to imply any sort of equivalence between naked peptides and such derivatives). Non-limiting examples of such amino acid residues include for instance 2-amino adipic acid, 3-amino adipic acid, β -alanine, β -aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid, 2-aminopimelic acid, 2,4-diaminobutyric acid, desmosine, 2,2'-diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, hydroxylysine, alcohdroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, alloseucine, N-methylglycine, N-methylisoleucine, 6-N-methyllysine, N-methylvaline, norvaline, norleucine, ornithine, and statine halogenated amino acids.

[0120] Antigen binding peptides refers to any peptide that specifically binds to a portion of a given antigen under cellular and/or physiological conditions for an amount of time sufficient to induce, promote, enhance, and/or otherwise modulate a physiological effect associated with the antigen; to allow detection by ELISA, Western blot, or other similarly suitable protein binding technique described herein and/or known in the art and/or to otherwise be detectably bound thereto after a relevant period of time (for instance at least about 15 minutes, at least about 30 minutes, at least about 45 minutes, at least about 1 hour, at least about 2 hours, at least about 4 hours, at least about 6 hours, at least about 12 hours, about 1-24 hours, about 1-36 hours, about 1-48 hours, about 1-72 hours, about one week, or longer).

[0121] A CD38 binding peptide, or CD38BP, is an antigen binding peptide that specifically binds to the antigen CD38. Binding of a CD38BP to CD38 may be measured by use of the method described in Example 4.

[0122] The term immunoglobulin refers to a class of structurally related glycoproteins consisting of two pairs of polypeptide chains, one pair of light (L) low molecular weight chains and one pair of heavy (H) chains, all four inter-connected by disulfide bonds. The structure of immunoglobulins has been well characterized. See for instance Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)). Briefly, each heavy chain typically is comprised of a heavy chain variable region (abbreviated herein as V_H) and a heavy chain constant region. The heavy chain constant region typically is comprised of three domains, C_H1 , C_H2 , and C_H3 . Each light chain typically is comprised of a light chain variable region (abbreviated herein as V_L) and a light chain constant region. The light chain constant region typically is comprised of one domain, C_L . The V_H and V_L regions can be further subdivided into regions of hypervariability (or hypervariable regions which can be hypervariable in sequence and/or form of structurally defined loops), also termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FRs).

[0123] Each V_H and V_L is typically composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4 (see also Chothia and Lesk J. Mol. Biol. 196, 901-917 (1987)). Typically, the numbering of amino acid residues in this region is performed by the method described in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991) (phrases such as variable domain residue numbering as in Kabat or according to Kabat herein refer to this numbering system for heavy chain variable domains or light chain variable domains). Using this numbering system, the actual linear amino acid sequence of a peptide may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of V_H CDR2

and inserted residues (for instance residues 82a, 82b, and 82c, etc. according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a "standard" Kabat numbered sequence.

[0124] The term antibody (Ab) in the context of the present invention refers to an immunoglobulin molecule, a fragment of an immunoglobulin molecule, or a derivative of either thereof, which has the ability to specifically bind to an antigen under typical physiological conditions for significant periods of time such as at least about 30 minutes, at least about 45 minutes, at least about one hour, at least about two hours, at least about four hours, at least about 8 hours, at least about 12 hours, about 24 hours or more, about 48 hours or more, about 3, 4, 5, 6, 7 or more days, etc., or any other relevant functionally-defined period (such as a time sufficient to induce, promote, enhance, and/or modulate a physiological response associated with antibody binding to the antigen).

[0125] The variable regions of the heavy and light chains of the immunoglobulin molecule contain a binding domain that interacts with an antigen. The constant regions of the antibodies (Abs) may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (such as effector cells) and the first component (C1q) of the classical complement system.

[0126] An anti-CD38 antibody may be a bispecific antibody, diabody, or similar molecule (see for instance PNAS USA 90(14), 6444-8 (1993) for a description of diabodies). Indeed, bispecific antibodies, diabodies, and the like, provided by the present invention may bind any suitable target in addition to a portion of CD38.

[0127] As indicated above, the term antibody herein, unless otherwise stated or clearly contradicted by context, includes fragments of an antibody that retain the ability to specifically bind to an antigen. It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antibody" include (i) a Fab fragment, a monovalent fragment consisting of the V_L , V_H , C_L and C_H1 domains; (ii) $F(ab)_2$ and $F(ab')_2$ fragments, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting essentially of the V_H and C_H1 domains; (iv) a Fv fragment consisting essentially of the V_L and V_H domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., Nature 341, 544-546 (1989)), which consists essentially of a V_H domain; (vi) an isolated complementarity determining region (CDR), and (vii) a combination of two or more isolated CDRs which may optionally be joined by a synthetic linker. Furthermore, although the two domains of the Fv fragment, V_L and V_H , are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the V_L and V_H regions pair to form monovalent molecules (known as single chain antibodies or single chain Fv (scFv), see for instance Bird et al., Science 242, 423-426 (1988) and Huston et al., PNAS USA 85, 5879-5883 (1988)). Such single chain antibodies are encompassed within the term antibody unless otherwise noted or clearly indicated by context. Other forms of single chain antibodies, such as diabodies are included within the term antibody. Although such fragments are generally included within the meaning of antibody, they collectively and each independently are unique features of the present invention, exhibiting different biological properties and utility. These and other useful antibody fragments in the context of the present invention are discussed further herein.

[0128] It also should be understood that the term antibody also generally includes polyclonal antibodies, monoclonal antibodies (mAbs), antibody-like polypeptides, such as chimeric antibodies and humanized antibodies, anti-idiotypic (anti-Id) antibodies to antibodies, and antibody fragments retaining the ability to specifically bind to the antigen (antigen-binding fragments) provided by any known technique, such as enzymatic cleavage, peptide synthesis, and recombinant techniques. An antibody as generated can possess any isotype.

[0129] The terms "anti-CD38 antibody" and "CD38 antibody" are used interchangeably herein. An anti-CD38 antibody, or a CD38 antibody, is an antibody as described above, which binds specifically to the antigen CD38.

[0130] The term "epitope" means a protein determinant capable of specific binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents. The epitope may comprise amino acid residues directly involved in the binding (also called immunodominant component of the epitope) and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen binding peptide (in other words, the amino acid residue is within the footprint of the specifically antigen binding peptide).

[0131] The term "bispecific molecule" is intended to include any agent, such as a protein, peptide, or protein or peptide complex, which has two different binding specificities. For example, the molecule may bind to, or interact with, (a) a cell surface antigen and (b) an Fc receptor on the surface of an effector cell. The term "multispecific molecule" is intended to include any agent, for instance a protein, peptide, or protein or peptide complex, which has more than two different binding specificities. For example, the molecule may bind to, or interact with, (a) a cell surface antigen, (b) an Fc receptor on the surface of an effector cell, and (c) at least one other component. Accordingly, the present invention includes, but is not limited to, bispecific, trispecific, tetraspecific, and other multispecific molecules which are directed to CD38, and to other cell surface antigens or targets, such as Fc receptors on effector cells.

[0132] The term "bispecific antibodies" is intended to include any anti-CD38 antibody, which is a bispecific molecule. The term "bispecific antibodies" also includes diabodies. Diabodies are bivalent, bispecific antibodies in which the V_H and V_L domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see for instance Holliger, P. et al., PNAS USA 90, 6444-6448 (1993), Poljak, R.J. et al., Structure 2, 1121-1123 (1994)).

[0133] As used herein, the term "effector cell" refers to an immune cell which is involved in the effector phase of an immune response, as opposed to the cognitive and activation phases of an immune response. Exemplary immune cells include a cell of a myeloid or lymphoid origin, for instance lymphocytes (such as B cells and T cells including cytolytic T cells (CTLs)), killer cells, natural killer cells, macrophages, monocytes, eosinophils, neutrophils, polymorphonuclear cells, granulocytes, mast cells, and basophils. Some effector cells express specific Fc receptors and carry out specific immune functions. In some embodiments, an effector cell is capable of inducing antibody-dependent cellular cytotoxicity (ADCC), such as a neutrophil capable of inducing ADCC. For example, monocytes, macrophages, which express FcR are involved in specific killing of target cells and presenting antigens to other components of the immune system, or binding to cells that present antigens. In some embodiments, an effector cell may phagocytose a target antigen, target cell, or microorganism. The expression of a particular FcR on an effector cell can be regulated by humoral factors such as cytokines. For example, expression of Fc γ RI has been found to be up-regulated by interferon γ (IFN- γ) and/or G-CSF. This enhanced expression increases the cytotoxic activity of Fc γ RI-bearing cells against targets. An effector cell can phagocytose or lyse a target antigen or a target cell.

[0134] The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the present invention may include amino acid residues not encoded by human germline immunoglobulin sequences (for instance mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[0135] As used herein, a human antibody is "derived from" a particular germline sequence if the antibody is obtained from a system using human immunoglobulin sequences, for instance by immunizing a transgenic mouse carrying human immunoglobulin genes or by screening a human immunoglobulin gene library, and wherein the selected human antibody is at least 90%, such as at least 95%, for instance at least 96%, such as at least 97%, for instance at least 98%, or such as at least 99% identical in amino acid sequence to the amino acid sequence encoded by the germline V_H or V_L variable region gene segment. Typically, a human antibody derived from a particular human germline V_H or V_L variable region gene segment sequence will display no more than 10 amino acid differences, such as no more than 5, for instance no more than 4, 3, 2, or 1 amino acid difference from the amino acid sequence encoded by the germline immunoglobulin gene.

[0136] A chimeric antibody is an antibody that contains one or more regions from one antibody and one or more regions from one or more other antibodies derived from another species. A monovalent chimeric antibody is a dimer (HL) formed by a chimeric H chain associated through disulfide bridges with a chimeric L chain. A divalent chimeric antibody is tetramer (H_2L_2) formed by two HL dimers associated through at least one disulfide bridge. A polyvalent chimeric antibody may also be produced, for example, by employing a CH region that oligomerizes (for instance from an IgM H chain, or μ chain). Typically, a chimeric antibody refers to an antibody in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see for instance US 4,816,567 and Morrison et al., PNAS USA 81, 6851-6855 (1984)). Chimeric antibodies are produced by recombinant processes well known in the art (see for instance Cabilly et al., PNAS USA 81, 3273-3277 (1984), Morrison et al., PNAS USA 81, 6851-6855 (1984), Boulianne et al., Nature 312, 643-646 (1984), EP125023, Neuberger et al., Nature 314, 268-270 (1985), EP171496, EP173494, WO86/01533, EP184187, Sahagan et al., J. Immunol. 137, 1066-1074 (1986), WO87/02671, Liu et al., PNAS USA 84, 3439-3443 (1987), Sun et al., PNAS USA 84, 214-218 (1987), Better et al., Science 240, 1041-1043 (1988) and Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988)).

[0137] A humanized antibody is an antibody that is derived from a non-human species, in which certain amino acids in the framework and constant domains of the heavy and light chains have been mutated so as to avoid or abrogate an immune response in humans. Humanized forms of non-human (for instance murine) antibodies are chimeric antibodies which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or non-human primate having the desired antigen-binding characteristics such as specificity, and affinity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human

residues. Furthermore, humanized antibodies may comprise residues which are not found in the recipient antibody or in the donor antibody. These modifications are made to further optimize antibody performance. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. A humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., Nature 321, 522-525 (1986), Riechmann et al., Nature 332, 323-329 (1988) and Presta, Curr. Op. Struct. Biol. 2, 593-596 (1992).

[0138] The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope. Accordingly, the term "human monoclonal antibody" refers to antibodies displaying a single binding specificity which have variable and constant regions derived from human germline immunoglobulin sequences. The human monoclonal antibodies may be generated by a hybridoma which includes a B cell obtained from a transgenic or transchromosomal non-human animal, such as a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene, fused to an immortalized cell. A monoclonal antibody may be abbreviated as mAb.

[0139] The term "recombinant human antibody", as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (such as a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom (described further elsewhere herein), (b) antibodies isolated from a host cell transformed to express the antibody, such as from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial human antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies may be subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the V_H and V_L regions of the recombinant antibodies are sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

[0140] As used herein, a "heterologous antibody" is defined in relation to the transgenic non-human organism producing such an antibody. This term refers to an antibody having an amino acid sequence corresponding to that found in an organism not consisting of the non-human animal, and generally from a species other than that of the transgenic non-human animal.

[0141] An "isolated antibody," as used herein, is intended to refer to an antibody which is substantially free of other antibodies having different antigenic specificities (for instance an isolated antibody that specifically binds to CD38 is substantially free of antibodies that specifically bind antigens other than CD38). An isolated antibody that specifically binds to an epitope, isoform or variant of human CD38 may, however, have cross-reactivity to other related antigens, for instance from other species (such as CD38 species homologs). Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals. In one embodiment of the present invention, a combination of "isolated" monoclonal antibodies having different specificities are combined in a well defined composition.

[0142] As used herein, "specific binding" refers to an antigen binding peptide, such as an antibody, binding to a predetermined antigen. Typically, the antigen binding peptide, such as an antibody, binds with an affinity corresponding to a K_D of about 10^{-7} M or less, such as about 10^{-8} M or less, such as about 10^{-9} M or less, about 10^{-10} M or less, or about 10^{-11} M or even less when determined by surface plasmon resonance (SPR) technology in a BIAcore 3000 instrument using recombinant CD38 as the ligand and the antibody as the analyte. The antigen binding peptide may bind to the predetermined antigen with an affinity corresponding to a K_D that is at least ten-fold lower, such as at least 100 fold lower, for instance at least 1000 fold lower, such as at least 10,000 fold lower, for instance at least 100,000 fold lower than its affinity for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen. The amount with which the affinity is lower is dependent on the K_D of the antigen binding peptide, so that when the K_D of the antigen binding peptide is very low (that is, the antigen binding peptide is highly specific), then the amount with which the affinity for the antigen is lower than the affinity for a non-specific antigen may be at least 10,000 fold. The phrases "an antigen binding peptide recognizing an antigen" and "an antigen binding peptide specific for an antigen" are used interchangeably herein with the term "an antigen binding peptide which binds specifically to an antigen". Likewise, the phrases "an antibody recognizing an antigen" and "an antibody specific for an antigen" are used interchangeably herein with the term "an antibody which binds specifically to an antigen".

[0143] The term " k_d " (sec^{-1}), as used herein, is intended to refer to the dissociation equilibrium rate constant of a particular antibody-antigen interaction. Said value is also referred to as the k_{off} value.

[0144] The term " k_a " ($\text{M}^{-1} \times \text{sec}^{-1}$), as used herein, is intended to refer to the association equilibrium rate constant of a particular antibody-antigen interaction.

[0145] The term " K_D " (M), as used herein, is intended to refer to the dissociation equilibrium constant of a particular antibody-antigen interaction.

[0146] The term " K_A " (M^{-1}), as used herein, is intended to refer to the association equilibrium constant of a particular antibody-antigen interaction and is obtained by dividing the k_a by the k_d .

[0147] As used herein, "isotype" refers to the antibody class (for instance IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM) that is encoded by heavy chain constant region genes.

[0148] As used herein, "isotype switching" refers to the phenomenon by which the class, or isotype, of an antibody changes from one immunoglobulin class to one of the other immunoglobulin classes.

[0149] As used herein, "nonswitched isotype" refers to the isotypic class of heavy chain that is produced when no isotype switching has taken place; the CH gene encoding the nonswitched isotype is typically the first CH gene immediately downstream from the functionally rearranged VDJ gene. Isotype switching has been classified as classical or non-classical isotype switching. Classical isotype switching occurs by recombination events which involve at least one switch sequence region in the transgene. Non-classical isotype switching may occur by, for example, homologous recombination between human $\sigma\mu$ and human $\Sigma\mu$ (δ -associated δ deletion). Alternative non-classical switching mechanisms, such as intertransgene and/or interchromosomal recombination, among others, may occur and effectuate isotype switching.

[0150] As used herein, the term "switch sequence" refers to those DNA sequences responsible for switch recombination. A "switch donor" sequence, typically a μ switch region, will be 5' (i.e., upstream) of the construct region to be deleted during the switch recombination. The "switch acceptor" region will be between the construct region to be deleted and the replacement constant region (for instance γ , ϵ , etc.). As there is no specific site where recombination always occurs, the final gene sequence will typically not be predictable from the construct.

[0151] As used herein, "glycosylation pattern" is defined as the pattern of carbohydrate units that are covalently attached to a protein, more specifically to an immunoglobulin (antibody) protein. A glycosylation pattern of a heterologous antibody may be characterized as being substantially similar to glycosylation patterns which occur naturally on antibodies produced by the species of the non-human transgenic animal, when one of ordinary skill in the art would recognize the glycosylation pattern of the heterologous antibody as being more similar to said pattern of glycosylation in the species of the non-human transgenic animal than to the species from which the CH genes of the transgene were derived.

[0152] The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

[0153] The term "rearranged" as used herein refers to a configuration of a heavy chain or light chain immunoglobulin locus wherein a V segment is positioned immediately adjacent to a D-J or J segment in a conformation encoding essentially a complete V_H or V_L domain, respectively. A rearranged immunoglobulin (antibody) gene locus can be identified by comparison to germline DNA; a rearranged locus will have at least one recombined heptamer/nonamer homology element.

[0154] The term "unrearranged" or "germline configuration" as used herein in reference to a V segment refers to the configuration wherein the V segment is not recombined so as to be immediately adjacent to a D or J segment.

[0155] The term "nucleic acid molecule", as used herein, is intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but is preferably double-stranded DNA. The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is "isolated" or "rendered substantially pure" when purified away from other cellular components or other contaminants, such as other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel et al., ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley InterScience New York (1987).

[0156] A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence. With respect to transcription of regulatory sequences, operably linked means that the DNA sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame. For switch sequences, operably linked indicates that the sequences are capable of effecting switch recombination.

[0157] As used herein, the term "inhibits growth" (for instance when referring to cells) is intended to include any measurable decrease in the cell growth when contacted with a CD38BP, such as an anti-CD38 antibody, as compared to the growth of the same cells not in contact with a CD38BP, such as an anti-CD38 antibody, for instance an inhibition of growth of a cell culture by at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 99%, or 100%.

[0158] As used herein, the terms "inhibits binding" and "blocks binding" (for instance when referring to inhibition/blocking of binding of a CD38 binding partner to CD38) are used interchangeably and encompass both partial and complete inhibition/blocking. The inhibition/blocking of binding of a CD38 binding partner to CD38 may reduce or alter the normal level or type of cell signaling that occurs when a CD38 binding partner binds to CD38 without inhibition or blocking. Inhibition and blocking are also intended to include any measurable decrease in the binding affinity of a CD38 binding

partner to CD38 when in contact with a CD38BP, such as an anti-CD38 antibody, as compared to the ligand not in contact with a CD38BP, such as an anti-CD38 antibody, for instance a blocking of binding of a CD38 binding partner to CD38 by at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 99%, or 100%.

[0159] "Target cell" shall mean any undesirable cell in a subject (for instance a human or animal) that can be targeted by a composition (comprising for instance a CD38BP, such as a human monoclonal anti-CD38 antibody, and/or a bispecific or a multispecific molecule directed against CD38) of the present invention. In some embodiments, the target cell is a cell expressing or overexpressing CD38. Cells expressing CD38 typically include hemopoietic cells, such as medullary thymocytes, activated T and B cells, 80% of resting NK cells and monocytes, lymph node germinal center lymphoblasts, plasma B cells and some intrafollicular cells, dendritic cells, normal bone marrow cells, particular precursor cells, 50-80% of umbilical cord blood cells, erythrocytes and platelets. CD38 can also be expressed by non-hemopoietic cells, such as intra-epithelial cells and lamina propria lymphocytes in the gut, by Purkinje cells and neurofibrillary tangles in the brain, by epithelial cells in the prostate, 3-cells in the pancreas, osteoclasts in the bone, retinal cells in the eye, and sarcolemma of smooth and striated muscle. On malignant cells, CD38 is expressed in a variety of malignant hematological diseases, including but not restricted to multiple myeloma, primary or secondary plasma cell leukemia, B-cell chronic lymphocytic leukemia, B-cell acute lymphocytic leukemia, Waldenström macroglobulinemia, primary systemic amyloidosis, mantle-cell lymphoma, polymphocytic/myelocytic leukemia, acute myeloid leukemia, chronic myeloid leukemia, follicular lymphoma, and NK-cell leukemia.

[0160] The term "vector," as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (for instance bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (such as non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the present invention is intended to include such other forms of expression vectors, such as viral vectors (such as replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0161] The term "recombinant host cell" (or simply "host cell"), as used herein, is intended to refer to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein. Recombinant host cells include, for example, transfectomas, such as CHO cells, NS/O cells, and lymphocytic cells.

[0162] The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (for instance polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel, Gene Expression Technology. Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Examples of regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV), Simian Virus 40 (SV40), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)) and polyoma. Alternatively, nonviral regulatory sequences may be used, such as the ubiquitin promoter or β -globin promoter.

[0163] As used herein, the term "subject" includes any human or non-human animal. The term "non-human animal" includes all vertebrates, for instance mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc.

[0164] The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, e.g., electroporation, calcium-phosphate precipitation, DEAE-dextran transfection, lipofectin transfection and the like.

[0165] The term "transfectoma", as used herein, includes recombinant eukaryotic host cells expressing the antibody, such as CHO cells, NS/O cells, HEK293 cells, plant cells, or fungi, including yeast cells.

[0166] The term "non-human animal" includes all vertebrates, for instance, mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc. The term "non-human animal" includes all vertebrates, for instance, mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc. The term "non-human animal" includes all vertebrates, for instance, mammals and non-mammals,

such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc.

[0167] The terms "transgenic, non-human animal" refers to a non-human animal having a genome comprising one or more human heavy and/or light chain transgenes or transchromosomes (either integrated or non-integrated into the animal's natural genomic DNA) and which is capable of expressing fully human antibodies. For example, a transgenic mouse can have a human light chain transgene and either a human heavy chain transgene or human heavy chain transchromosome, such that the mouse produces human anti-CD38 antibodies when immunized with CD38 antigen and/or cells expressing CD38. The human heavy chain transgene can be integrated into the chromosomal DNA of the mouse, as is the case for transgenic mice, for instance HuMAb mice, such as HCo7 or HCo12 mice, or the human heavy chain transgene can be maintained extrachromosomally, as is the case for transchromosomal KM mice as described in WO02/43478. Such transgenic and transchromosomal mice (collectively referred to herein as "transgenic mice") are capable of producing multiple isotypes of human monoclonal antibodies to a given antigen (such as IgG, IgA, IgM, IgD and/or IgE) by undergoing V-D-J recombination and isotype switching. Transgenic, non-human animal can also be used for production of antibodies against a specific antigen by introducing genes encoding such specific antibody, for example by operatively linking the genes to a gene which is expressed in the milk of the animal.

[0168] The term specificity herein refers to the ability of a CD38 binding peptide, such as an anti-CD38 antibody, to recognize an epitope within CD38, while only having little or no detectable reactivity with other portions of CD38 (including other epitopes that are bound by other CD38BPs, such as anti-CD38 antibodies). Specificity can be relatively determined by competition assays as described herein. Specificity can more particularly be determined by any of the epitope identification/characterization techniques described herein or their equivalents known in the art.

[0169] An antibody specific for a particular antigenic determinant may nonetheless cross-react with other biomolecules that may be present in some biological context with CD38. More typically, a CD38BP, such as an anti-CD38 antibody, may cross-react with CD38 homologues from other species. In either or both contexts, typically such cross-reactive antibodies are selective for human CD38 with respect to relevant structure and/or environmental factors.

[0170] The term selectivity herein refers to the preferential binding of a CD38BP, such as an anti-CD38 antibody, for a particular region, target, or peptide; typically a region or epitope in CD38, as opposed to one or more other biological molecules, structures, cells, tissues, etc. In one embodiment, a CD38BP, such as an anti-CD38 antibody of the present invention, is selective for a portion of CD38 in the context of colon cancer cells (i.e., the anti-CD38 antibody will selectively bind to the portion of CD38 over other components of a colon cancer cell).

[0171] The CD38 antibodies of the present invention are typically used in and provided in an at least substantially isolated form. A substantially isolated molecule is a molecule that is the predominant species in the composition wherein it is found with respect to the class of molecules to which it belongs (i.e., it makes up at least about 50% of the type of molecule in the composition and typically will make up at least about 70%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more of the species of molecule, e.g., peptide, in the composition (e.g., the composition will exhibit at least about 98%, 98%, or 99% homogeneity for the CD38BP in the context of all present peptide species)).

[0172] An isolated molecule refers to a molecule that is not associated with significant levels (such as more than about 1%, more than about 2%, more than about 3%, or more than about 5%) of any extraneous and undesirable physiological factors, such as non-CD38 binding biomolecules (or CD38 binding molecules that may interfere with the binding and/or activity of a CD38 antibody of the present invention) contained within a cell or animal in which the antibody is produced. An isolated molecule also refers to any molecule that has passed through such a stage of purity due to human intervention (whether automatic, manual, or both). In many of the various compositions provided by the present invention, such as in a composition comprising one or more pharmaceutically acceptable carriers, the CD38 antibody may be present in relatively small amounts in terms of numbers of total molecular species in the composition (for instance in the case of a composition comprising a large amount of a pharmaceutically acceptable carrier, stabilizer, and/or preservative). In some cases additional peptides, such as BSA, may be included in such a composition with a previously purified antibody. However, provided that such additional constituents of the composition are acceptable for the intended application of the antibody, such a composition can still be described as comprising an isolated antibody.

[0173] The CD38 antibodies of the present invention are typically substantially free of other CD38BPs, such as CD38BPs having different antigenic specificities. However, the present invention does also provide a composition comprising a number of CD38BP antibodies with different specificities and characteristics (for instance the present invention provides a "cocktail" of CD38BPs having different specificity and/or selectivity characteristics).

[0174] "Treatment" means the administration of an effective amount of a therapeutically active compound of the present invention with the purpose of easing, ameliorating, or eradicating (curing) symptoms or disease states.

[0175] Described herein are:

- a CD38BP comprising a V_L region consisting essentially of the sequence of SEQ ID No:2;
- a CD38BP comprising a V_H region consisting essentially of the sequence of SEQ ID No:6;
- a CD38BP comprising a V_L region consisting essentially of the sequence of SEQ ID No:2 and a V_H region consisting

essentially of the sequence of SEQ ID No:6;

- a CD38BP comprising a V_L CDR1 consisting essentially of the sequence of SEQ ID No:3;
- a CD38BP comprising a V_L CDR2 consisting essentially of the sequence of SEQ ID No:4;
- a CD38BP comprising a V_L CDR3 consisting essentially of the sequence of SEQ ID No:5;
- a CD38BP comprising a V_H CDR1 consisting essentially of the sequence of SEQ ID No:8;
- a CD38BP comprising a V_H CDR2 consisting essentially of the sequence of SEQ ID No:9;
- a CD38BP comprising a V_H CDR3 consisting essentially of the sequence of SEQ ID No:10;
- a CD38BP comprising V_L CDRs (V_L CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No: 3, SEQ ID No:4 and SEQ ID No:5, respectively;
- a CD38BP that comprises V_H CDRs (V_H CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No:8, SEQ ID No:9 and SEQ ID No:10, respectively;
- a CD38BP that comprises (a) three V_L CDRs, which independently consist essentially of SEQ ID No:3, SEQ ID No:4 and SEQ ID No:5 in close proximity to one another (e.g., near the spacing of V_L CDRs in a wild-type anti-CD38 antibody) in the CD38BP and (b) three V_H CDRs which independently consist essentially of SEQ ID No:8, SEQ ID No:9 and SEQ ID No:10 in close proximity to one another (e.g., near the spacing of V_H CDRs in a wild-type anti-CD38 antibody) in the CD38BP;
- a CD38BP that comprises a flexible linker positioned between the V_L region and V_H region of the CD38BP;
- a CD38BP, wherein the V_L and V_H regions are presented on separate chains in the context of an immunoglobulin fold protein and oriented such that the V_L CDR1, CDR2, CDR3 and V_H CDR1, CDR2, and CDR3 cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on CD38;
- a CD38BP that comprises two sets of variable domains (sets of associated V_L and V_H domains on associated separate chains), such that the CD38BP comprises two identical antigenic determinant binding sites.

[0176] Any of such CD38BPs described in this paragraph are expected to, at least in part, have similar epitope specificity, selectivity, and other characteristics as an antibody having V_L region comprising the sequence of SEQ ID No:2 and a V_H region comprising the sequence of SEQ ID No:7, and, accordingly, may be useful in the treatment of multiple myeloma.

[0177] In one embodiment, the CD38 antibody of the invention comprises a V_L region consisting essentially of the sequence of SEQ ID No:12.

[0178] In one embodiment, the CD38 antibody of the invention comprises a V_H region consisting essentially of the sequence of SEQ ID No:17.

[0179] In one embodiment, the CD38 antibody of the invention comprises a V_L region consisting essentially of the sequence of SEQ ID No:12 and a V_H region consisting essentially of the sequence of SEQ ID No:17.

[0180] Described herein are:

- a CD38BP comprising a V_L CDR consisting essentially of the sequence of SEQ ID No:13;
- a CD38BP comprising a V_L CDR2 consisting essentially of the sequence of SEQ ID No:14;
- a CD38BP comprising a V_L CDR3 consisting essentially of the sequence of SEQ ID No:15;
- a CD38BP comprising a V_H CDR1 consisting essentially of the sequence of SEQ ID No:18;
- a CD38BP comprising a V_H CDR2 consisting essentially of the sequence of SEQ ID No:19;
- a CD38BP comprising a V_H CDR3 consisting essentially of the sequence of SEQ ID No:20;
- a CD38BP comprising V_L CDRs (V_L CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No:13, SEQ ID No:14 and SEQ ID No:15, respectively; and
- a CD38BP that comprises V_H CDRs (V_H CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No:18, SEQ ID No:19 and SEQ ID No:20, respectively.

[0181] In one embodiment, the CD38 antibody of the invention comprises (a) three V_L CDRs, which independently consist essentially of SEQ ID No:13, SEQ ID No:14 and SEQ ID No:15 in close proximity to one another (e.g., near the spacing of V_L CDRs in a wild-type anti-CD38 antibody) in the antibody and (b) three V_H CDRs which independently consist essentially of SEQ ID No:18, SEQ ID No:19 and SEQ ID No:20 in close proximity to one another (e.g., near the spacing of V_H CDRs in a wild-type anti-CD38 antibody) in the antibody.

[0182] CD38 antibodies of the invention may comprise a flexible linker positioned between the V_L region and V_H region of the CD38BP. The V_L and V_H regions may be presented on separate chains in the context of an immunoglobulin fold protein and oriented such that the V_L CDR1, CDR2, CDR3 and V_H CDR1, CDR2, and CDR3 cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on CD38. The antibody may comprise two sets of variable domains (sets of associated V_L and V_H domains on associated separate chains), such that the antibody comprises two identical antigenic determinant binding sites.

[0183] Any of such CD38BPs and antibodies described in this paragraph are expected to, at least in part, have similar

epitope specificity, selectivity, and other characteristics as an antibody having V_L region comprising the sequence of SEQ ID No:12 and a V_H region comprising the sequence of SEQ ID No:17, and, accordingly, may be useful in the treatment of multiple myeloma.

[0184] Described herein are:

- a CD38BP comprising a V_L region consisting essentially of the sequence of SEQ ID No:22;
- a CD38BP comprising a V_H region consisting essentially of the sequence of SEQ ID No:27; a CD38BP comprising a V_L region consisting essentially of the sequence of SEQ ID No:22 and a V_H region consisting essentially of the sequence of SEQ ID No:27;
- a CD38BP comprising a V_L CDR1 consisting essentially of the sequence of SEQ ID No:23;
- a CD38BP comprising a V_L CDR2 consisting essentially of the sequence of SEQ ID No:24;
- a CD38BP comprising a V_L CDR3 consisting essentially of the sequence of SEQ ID No:25;
- a CD38BP comprising a V_H CDR1 consisting essentially of the sequence of SEQ ID No:28;
- a CD38BP comprising a V_H CDR2 consisting essentially of the sequence of SEQ ID No:29;
- a CD38BP comprising a V_H CDR3 consisting essentially of the sequence of SEQ ID No:30;
- a CD38BP comprising V_L CDRs (V_L CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No:23, SEQ ID No:24 and SEQ ID No:25, respectively;
- a CD38BP that comprises V_H CDRs (V_H CDR1, CDR2, and CDR3) consisting essentially of SEQ ID No:28, SEQ ID No:29 and SEQ ID No:30, respectively;
- a CD38BP that comprises (a) three V_L CDRs, which independently consist essentially of SEQ ID No:23, SEQ ID No:24 and SEQ ID No:25 in close proximity to one another (e.g., near the spacing of V_L CDRs in a wild-type anti-CD38 antibody) in the CD38BP and (b) three V_H CDRs which independently consist essentially of SEQ ID No:28, SEQ ID No:29 and SEQ ID No:30 in close proximity to one another (e.g., near the spacing of V_H CDRs in a wild-type anti-CD38 antibody) in the CD38BP;
- a CD38BP that comprises a flexible linker positioned between the V_L region and V_H region of the CD38BP;
- a CD38BP, wherein the V_L and V_H regions are presented on separate chains in the context of an immunoglobulin fold protein and oriented such that the V_L CDR1, CDR2, CDR3 and V_H CDR1, CDR2, and CDR3 cooperatively associate to contribute in selectively and/or specifically bind an antigenic determinant on CD38;
- a CD38BP that comprises two sets of variable domains (sets of associated V_L and V_H domains on associated separate chains), such that the CD38BP comprises two identical antigenic determinant binding sites.

[0185] Any of such CD38BPs described in this paragraph are expected to, at least in part, have similar epitope specificity, selectivity, and other characteristics as an antibody having V_L region comprising the sequence of SEQ ID No:22 and a V_H region comprising the sequence of SEQ ID No:27, and, accordingly, may be useful in the treatment of multiple myeloma.

[0186] Described herein are:

- a CD38BP comprising a V_L CDR1 consisting essentially of a sequence according to SEQ ID No:3 or SEQ ID No:13 or SEQ ID No:23, wherein the N-terminal residue and/or one, two, or three of the C-terminal amino acid residues are missing;
- a CD38BP comprising a V_L CDR2 consisting essentially of a sequence according to SEQ ID No:4 or SEQ ID No:14 or SEQ ID No:24, wherein one or two of the N-terminal residues and/or one, two, or three of the C-terminal residues are missing;
- a CD38BP comprising a V_L CDR3 consisting essentially of a sequence according to SEQ ID No:5 or SEQ ID No:15 or SEQ ID No:25, wherein the N-terminal residue and/or one, two, three, or four of the C-terminal residues are missing;
- a V_H CDR1 consisting essentially of a sequence according to SEQ ID No:8 or SEQ ID No:18 or SEQ ID No:28, wherein one, two, three, or four of the N-terminal residues and/or one, two, three, or four C-terminal residues are missing;
- a CD38BP comprising a V_H CDR2 consisting essentially of a sequence according to SEQ ID No:9 or SEQ ID No:19 or SEQ ID No:29, wherein one, two, three, four, or five of the N-terminal amino acids thereof and/or one, two, three, four, five, or six of the C-terminal amino acids thereof are missing;
- a CD38BP comprising a V_H CDR3 consisting essentially of a sequence according to SEQ ID No:10 or SEQ ID No:20 or SEQ ID No:30, wherein the N-terminal one, two, or three amino acid residues and/or the C-terminal one, two, three, or four amino acid residues are missing.

[0187] Also described are CD38BPs wherein these "truncated" CDR sequences are combined with each other and/or other CDR sequences described herein.

[0188] Also described herein are CD38BPs comprising functional variants of the V_L region, V_H region, or one or more CDRs of the antibodies of the examples. A functional variant of a V_L, V_H, or CDR used in the context of a CD38BP still allows the CD38BP to retain at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the affinity/avidity and/or specificity/selectivity of the parent antibody and in some cases such a CD38BP may be

[0189] In one embodiment, provided herein is an antibody binding to human CD38 comprising a variant V_L consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:12, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_L sequence of SEQ ID No:12, such as an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17.

[0190] Described herein is a CD38BP comprising a variant V_L CDR1 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID No:3 or SEQ ID No:13 or SEQ ID No:23, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_L CDR1 sequence of SEQ ID No:3 or SEQ ID No:13 or SEQ ID No:23, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 or SEQ ID No:12 or SEQ ID No:22, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, respectively.

[0191] Described herein is a CD38BP comprising a variant V_L CDR2 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID Nos:4 or 14, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_L CDR2 sequence of SEQ ID No:4 or SEQ ID No:14 or SEQ ID No:24, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 or SEQ ID No:12 or SEQ ID No:22, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, respectively.

[0192] Described herein is a CD38BP comprising a variant V_L CDR3 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID No:5 or SEQ ID No:15 or SEQ ID No:25, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_L CDR3 sequence of SEQ ID No:5 or SEQ ID No:15 or SEQ ID No:25, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 or SEQ ID No:12 or SEQ ID No:22, respectively, such as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, respectively.

[0193] In one embodiment, provided herein is an antibody binding to human CD38 comprising a variant V_H consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to SEQ ID No:17, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_H sequence of SEQ ID No:17, such as an antibody having a V_H sequence of SEQ ID No:17 and a V_L sequence of SEQ ID No:12.

[0194] Described herein is a CD38BP comprising a variant V_H CDR1 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID No:8 or SEQ ID No:18 or SEQ ID No:28, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_H CDR1 sequence of SEQ ID No:8 or SEQ ID No:18 or SEQ ID No:28, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 or SEQ ID No:17 or SEQ ID No:27, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 and a V_L sequence of SEQ ID No:2, or an antibody having a V_H sequence of SEQ ID No:17 and a V_L sequence of SEQ ID No:12, or an antibody having a V_H sequence of SEQ

ID No:27 and a V_L sequence of SEQ ID No:22, respectively.

[0195] Described herein is a CD38BP comprising a variant V_H CDR2 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID No:9 or SEQ ID No:19 or SEQ ID No:29, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_H CDR2 sequence of SEQ ID No:9 or SEQ ID No:19 or SEQ ID No:29, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 or SEQ ID No:17 or SEQ ID No:27, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 and a V_L sequence of SEQ ID No:2, or an antibody having a V_H sequence of SEQ ID No:17 and a V_L sequence of SEQ ID No:12, or an antibody having a V_H sequence of SEQ ID No:27 and a V_L sequence of SEQ ID No:22, respectively.

[0196] Described herein is a CD38BP comprising a variant V_H CDR3 consisting essentially of a sequence having at least about 50%, such as at least 60%, for instance at least about 70%, such as at least about 75%, for instance at least about 80%, such as at least about 85%, for instance at least about 90%, such as at least about 95% amino acid sequence identity to a sequence according to any one of SEQ ID No: 10 or SEQ ID No:20 or SEQ ID No:30, wherein the CD38BP has at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the epitope binding characteristics of an antibody having a variant V_H CDR3 sequence of SEQ ID No: 10 or SEQ ID No:20 or SEQ ID No:30, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 or SEQ ID No:17 or SEQ ID No:27, respectively, such as an antibody having a V_H sequence of SEQ ID No:7 and a V_L sequence of SEQ ID No:2, or an antibody having a V_H sequence of SEQ ID No:17 and a V_L sequence of SEQ ID No:12, or an antibody having a V_H sequence of SEQ ID No:27 and a V_L sequence of SEQ ID No:22, respectively.

[0197] The percent identity between two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences may be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

[0198] The percent identity between two nucleotide sequences may be determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent identity between two nucleotide or amino acid sequences may also be determined using the algorithm of E. Meyers and W. Miller, Comput. Appl. Biosci 4, 11-17 (1988)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences may be determined using the Needleman and Wunsch, J. Mol. Biol. 48, 444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

[0199] The nucleic acid and protein sequences as described herein may further be used as a "query sequence" to perform a search against public databases to, for example, identify related sequences. Such searches may be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul et al., J. Mol. Biol. 215, 403-10 (1990). BLAST nucleotide searches may be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the present invention. BLAST protein searches may be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the protein molecules of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST may be utilized as described in Altschul et al., Nucleic Acids Res. 25(17), 3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) may be used. See <http://www.ncbi.nlm.nih.gov>.

[0200] The sequence of CDR variants may differ from the sequence of the CDR of the parent antibody sequences through mostly conservative substitutions; for instance at least about 35%, about 50% or more, about 60% or more, about 70% or more, about 75% or more, about 80% or more, about 85% or more, about 90% or more, about 95% or more (e.g., about 65-99%) of the substitutions in the variant are conservative amino acid residue replacements. In the context of the present invention, conservative substitutions may be defined by substitutions within the classes of amino acids reflected in one or more of the following three tables:

Amino acid residue classes for conservative substitutions

Acidic Residues	Asp and Glu
Basic Residues	Lys, Arg, and His
Hydrophilic Uncharged Residues	Ser, Thr, Asn, and Gln

(continued)

Aliphatic Uncharged Residues	Gly, Ala, Val, Leu, and Ile
Non-polar Uncharged Residues	Cys, Met, and Pro
Aromatic Residues	Phe, Tyr, and Trp

Alternative conservative amino acid residue substitution classes

1	Ala (A)	Ser (S)	Thr (T)
2	Asp (D)	Glu (E)	
3	Asp (N)	Gln (Q)	
4	Arg (R)	Lys (K)	
5	Ile (I)	Leu (L)	Met (M)
6	Phe (F)	Tyr (Y)	Trp (W)

Alternative Physical and Functional Classifications of Amino Acid Residues

Alcohol group-containing residues	S and T
Aliphatic residues	I, L, V, and M
Cycloalkenyl-associated residues	F, H, W, and Y
Hydrophobic residues	A, C, F, G, H, I, L, M, R, T, V, W, and Y
Negatively charged residues	D and E
Polar residues	C, D, E, H, K, N, Q, R, S, and T
Positively charged residues	H, K, and R
Small residues	A, C, D, G, N, P, S, T, and V
Very small residues	A, G, and S
Residues involved in turn formation	A, C, D, E, G, H, K, N, Q, R, S, P, and T
Flexible residues	Q, T, K, S, G, P, D, E, and R

[0201] More conservative substitutions groupings include: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Additional groups of amino acids may also be formulated using the principles described in, e.g., Creighton (1984) *Proteins: Structure and Molecular Properties* (2d Ed. 1993), W.H. Freeman and Company.

[0202] In one embodiment conservation in terms of hydropathic/hydrophilic properties and residue weight/size also is substantially retained in a variant CDR as compared to a CDR of an antibody of the examples (e.g., the weight class, hydropathic score, or both of the sequences are at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more (e.g., about 65-99%) retained). For example, conservative residue substitutions may also or alternatively be based on the replacement of strong or weak based weight based conservation groups, which are known in the art.

[0203] The retention of similar residues may also or alternatively be measured by a similarity score, as determined by use of a BLAST program (e.g., BLAST 2.2.8 available through the NCBI). Suitable variants typically exhibit at least about 45%, such as at least about 55%, at least about 65%, at least about 75%, at least about 85%, at least about 90%, at least about 95%, or more (e.g., about 70-99%) similarity to the parent peptide.

[0204] Substantial changes in function may be made by selecting substitutions that are less conservative than those shown in the defined groups, above. For example, nonconservative substitutions may be made which more significantly affect the structure of the peptide in the area of the alteration, for example, the alpha-helical, or beta-sheet structure; the charge or hydrophobicity of the molecule at the target site; or the bulk of the side chain. The substitutions which generally are expected to produce the greatest changes in the peptide's properties are those where 1) a hydrophilic

residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl, or alanyl; 2) a cysteine or proline is substituted for (or by) any other residue; 3) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or 4) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) a residue that does not have a side chain, e.g., glycine. Accordingly, these and other nonconservative substitutions may be introduced into peptide variants where significant changes in function/structure is desired and such changes avoided where conservation of structure/function is desired.

[0205] A convenient way for generating substitution variants is affinity maturation using phage using methods known in the art. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis may also be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues are likely suitable candidates for substitution.

[0206] Where hypervariable region insertions are made to generate a variant antibody, the typical range of lengths of the hypervariable region in question in known antibodies should be taken into consideration. For example, for the first hypervariable region of a light chain variable domain, insertions may be introduced into the V_L CDR1 sequence of a parent antibody while retaining a substantially similar and thereby expected appropriate size, which according to Kabat et al., supra, e.g., typically has an overall of about 9-20 (e.g., about 10-17) residues. Similarly, V_L CDR2 typically has an overall length from about 5-10 residues; V_L CDR3 typically has a length of about 7-20 residues; V_H CDR1 typically has a length of about 10-15 residues; V_H CDR2 typically has a length of about 15-20 residues; and V_H CDR3 typically has a length of about 6-30 residues (e.g., 3-25 residues). Insertions in the V_H region typically are made in V_H CDR3 and typically near the C-terminal of the domain, such as about residues 97-102 of the parent V_H CDR3 (for instance adjacent to, or C-terminal in sequence to, residue number 100 of the parent V_H CDR3 sequence) using the alignment and numbering as described in Kabat. Antibody variants with inserted amino acid residue(s) in a hypervariable region thereof may be prepared randomly, especially where the starting binding affinity of the parent antibody for the target antigen is such that randomly produced antibody variants may be readily screened. For example, phage display provides a convenient method of screening such random variants.

[0207] In the design, construction, and/or evaluation of CDR variants attention may be paid to the fact that CDR regions may be altered to enable a better binding to the epitope. Antibody CDRs typically operate by providing a complementary surface, possibly including fingers which can protrude into the protein surface of the antigen, or other paratope structure, onto which the epitope fits. If the epitope is not fitting tightly, the antibody may not offer the best affinity. However, as with epitopes, there often are a few key residues in a paratope structure that account for most of this binding. Thus, CDR sequences may vary in length and composition significantly between antibodies for the same peptide. The skilled artisan will recognize that certain residues, such as tyrosine residues (e.g., in the context of V_H CDR3 sequences), that are often significant contributors to such epitope binding, are typically retained in a CDR variant.

[0208] Variants of the CDR region may also increase the amino acid contacts between the antigen and an antibody variant, as compared to the amino acid contacts between the antigen and the parent antibody, by introducing one or more amino acid residues (either by substitution or insertions) which increase the contacts or energetically favorable interactions between one or more amino acid residues present in an antigen and one or more amino acid residues present in the antibody. The amino acid interactions of interest may be selected from hydrogen bonding interactions, van der Waals interactions, and ionic interactions.

[0209] Those skilled in the art will be aware of additional principles useful in the design and selection of CD38BP comprising CDR variants of the antibodies of the present invention.

[0210] In the context of CDR variants, which are variants of the CDRs of the antibodies of the examples, particularly in the context of variant CDR in anti-CD38 antibodies or fragments thereof, residues required to support and/or orientate the CDR structural loop structure(s) may typically be retained; residues which fall within about 10 angstroms of a CDR structural loop (but optionally only residues in this area that also possess a water solvent accessible surface of about 5 angstroms² or greater) may typically be unmodified or modified only by conservative amino acid residue substitutions; and/or the amino acid sequence may typically be subject to only a limited number of insertions and/or deletions (if any), such that CDR structural loop-like structures are retained in the variant (a description of related techniques and relevant principles is provided in for instance Schiweck et al., J Mol Biol. 268(5), 934-51 (1997), Morea, Biophys Chem. 68(1-3), 9-16 (1997), Shirai et al., FEBS Lett. 399(1-2), 1-8 (1996), Shirai et al., FEBS Lett. 455(1-2), 188-97 (1999), Reckzo et al., Protein Eng. 8(4), 389-95 (1995) and Eigenbrot et al., J Mol Biol. 229(4), 969-95 (1993). See also WO 03/048185, WO 03/070747 and WO 03/027246.

[0211] Additional techniques that may be used to generate variant antibodies include the directed evolution and other variant generation techniques described in for instance US 20040009498, Marks et al., Methods Mol Biol. 248, 327-43 (2004), Azriel-Rosenfeld et al., J Mol Biol. 335(1), 177-92 (2004), Park et al., Biochem Biophys Res Commun. 275(2), 553-7 (2000), Kang et al., Proc Natl Acad Sci USA. 88(24), 11120-3 (1991), Zahnd et al., J Biol Chem. 279(18), 18870-7

(2004), Xu et al., Chem Biol. 9(8), 933-42 (2002), Border et al., Proc Natl Acad Sci USA. 97(20), 10701-5 (2000), Crameri et al., Nat Med. 2(1), 100-2 (1996) and as more generally described in for instance WO 03/048185.

[0212] Generated antibody variants may be subjected to any suitable screening technique and antibodies with suitable and desirably superior properties in one or more relevant assays may be selected for further development.

[0213] CD38BPs comprising CDR sequences as described above may comprise any suitable number and combination of such V_L and V_H CDRs while retaining at least a substantial proportion (at least about 50%, 60%, 70%, 80%, 90%, 95% or more) of the affinity/avidity and/or specificity/selectivity of an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, and/or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 and/or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27,, but optionally differing in other characteristics, such as immunogenicity in a human patient, affinity for the epitope, increased half-life, etc. In some cases such a CD38BP may be associated with greater affinity, selectivity, and/or specificity than the parent antibody. In one embodiment, less than a full set of V_L CDRs and/or V_H CDRs is present in a CD38BP. In the antibodies of the present invention all of the V_L CDRs and V_H CDRs are present.

[0214] Examples of other functional properties of antibodies, which may be altered or retained in variant CD38BPs as compared to -003 and -005 and -024, are:

- (1) high affinity binding to CD38;
- (2) low dissociation rate from CD38
- (3) inhibition or blocking of CD38-binding to CD38 target;
- (4) elimination of T cells or B cells expressing CD38;
- (5) induction of a high level of CDC of either CD55/59 negative or CD55/59 positive cells;
- (6) translocation into lipid rafts upon binding to CD38;
- (7) tolerization of T cells;
- (8) inhibition of proliferation of Tor B cells cells expressing CD38;
- (9) internalization of CD38;
- (10) inhibition or induction of CD38 enzymatic activity;
- (11) inhibition or induction of CD38-induced signal transduction;
- (12) induction or inhibition of cytokine production;
- (13) induction or blocking of T cell or B cell differentiation;
- (14) induction of or rescue from apoptosis;
- (15) attenuation or augmentation of lysis induction by NK cells;
- (16) induction or inhibition of insulin production by β cells in pancreas;
- (17) prolonged survival of a subject having tumor cells which express CD38; and/or
- (18) induction of ADCC of CD38 targets when mixed with appropriate effector cells.

[0215] The present disclosure also provides CD38BPs which are characterized with respect to their ability to compete (competitively inhibit) or cross-compete (i.e., relatively partially inhibit epitope binding) with an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as antibody -003), or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 (such as antibody -005) or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, (such as antibody -024), for binding to CD38.

[0216] Such a CD38BP may be, for instance, a Fab fragment, derived from an antibody that binds to an epitope identical to or overlapping with an epitope bound by an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27. Such a Fab fragment, due to its relatively small size compared to the mAb molecules, may not significantly compete with said antibodies for binding to CD38 although the antibody from which it derived does. Nonetheless, such a CD38BP may be useful in similarly targeting nearby regions of CD38 (e.g., in the context of targeting a cytotoxin, radionuclide, or the like in the context of an immunoconjugate CD38BP). Therefore, such CD38BPs may be useful in the context of the methods as described herein.

[0217] Competition for binding to CD38 or a portion of CD38 by two or more CD38BPs may be determined by any suitable technique. In one embodiment, competition is determined for example as described in Example 7, 8 and 9.

[0218] Competition refers to any detectably significant reduction in the propensity for a particular molecule to bind a particular binding partner in the presence of another molecule that binds the binding partner. Typically, competition means an at least about 10% reduction, such as an at least about 15%, or an at least about 20% reduction in binding between a CD38BP and

- (a) a form of CD38 (e.g. "processed", "mature", "unprocessed", "not processed" or "immature" CD38);
- (b) a form of free CD38 (e.g., a CD38 fragment produced by *in vivo* processing);

- (c) a heterodimeric peptide composed of another peptide associated with CD38, such as a CD31, and CD38;
- (d) a complex of CD38 and one or more substrates, such as cAMP, NAD⁺ and/or cADPR;
- (e) a dimerized, associated and/or processed dimer of CD38 with a soluble ligand, such as CD31; or
- (f) a portion of CD38,

caused by the presence of another CD38BP as determined by, e.g., ELISA analysis or FACS analysis (as described in the examples section) using sufficient amounts of the two or more competing CD38BPs and CD38 molecule. It may also be the case that competition may exist between CD38BPs with respect to more than one of CD38, and/or a portion of CD38, e.g. in a context where the antibody-binding properties of a particular region of CD38 are retained in fragments thereof, such as in the case of a well-presented linear epitope located in various tested fragments or a conformational epitope that is presented in sufficiently large CD38 fragments as well as in CD38.

[0219] Assessing competition typically involves an evaluation of relative inhibitory binding using a first amount of a first molecule; a second amount of a second molecule; and a third amount of a third molecule (or a standard determined by binding studies that may be reasonably compared to new binding data with respect to the first and second molecules as a surrogate for actual contemporaneous data), wherein the first, second, and third amounts all are sufficient to make a comparison that imparts information about the selectivity and/or specificity of the molecules at issue with respect to the other present molecules. The first, second, and third amounts may vary with the nature of the CD38BP and potential targets therefore at issue. For instance, for ELISA assessments, similar to those described in the Examples section, about 5-50 μ g (e.g., about 10-50 μ g, about 20-50 μ g, about 5-20 μ g, about 10-20 μ g, etc.) of CD38BP and/or CD38 targets are required to assess whether competition exists. Conditions also should be suitable for binding. Typically, physiological or near-physiological conditions (e.g., temperatures of about 20-40°C, pH of about 7-8, etc.) are suitable for CD38BP:CD38 binding.

[0220] Often competition is marked by a significantly greater relative inhibition than about 5% as determined by ELISA and/or FACS analysis. It may be desirable to set a higher threshold of relative inhibition as a criteria/determinant of what is a suitable level of competition in a particular context (e.g., where the competition analysis is used to select or screen for new antibodies designed with the intended function of blocking the binding of another peptide or molecule binding to CD38 (e.g., the natural binding partners of CD38 such as CD31, also called CD31 antigen, EndoCAM, GPIIA', PECAM-1, platelet/endothelial cell adhesion molecule or naturally occurring anti-CD38 antibody)). Thus, for example, it is possible to set a criteria for competitiveness wherein at least about 10% relative inhibition is detected; at least about 15% relative inhibition is detected; or at least about 20% relative inhibition is detected before an antibody is considered sufficiently competitive. In cases where epitopes belonging to competing antibodies are closely located in an antigen, competition may be marked by greater than about 40% relative inhibition of CD38 binding (e.g., at least about 45% inhibition, such as at least about 50% inhibition, for instance at least about 55% inhibition, such as at least about 60% inhibition, for instance at least about 65% inhibition, such as at least about 70% inhibition, for instance at least about 75% inhibition, such as at least about 80% inhibition, for instance at least about 85% inhibition, such as at least about 90% inhibition, for instance at least about 95% inhibition, or higher level of relative inhibition).

[0221] Competition may be considered the inverse of cross-reactivity between a molecule and two potential binding partners. In certain embodiments, a CD38BP antibody of the present invention may specifically bind to one or more residues or regions in CD38 but also does not cross-react with other peptides, peptide regions, or molecules, e.g., an anti-CD38 antibody in accordance with the present invention may not cross-react with proteins with homology to CD38, such as BST-1 (bone marrow stromal cell antigen-1) and Mo5, also called CD157; or anti-CD38 antibodies may not cross-react with CD38 in the context of normal tissue, such as tissues not involved in multiple myeloma. Typically, a lack of cross-reactivity means less than about 5% relative competitive inhibition between the molecules when assessed by ELISA and/or FACS analysis using sufficient amounts of the molecules under suitable assay conditions.

[0222] Described herein are:

- a CD38BP that competes with an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, such as the antibody -003, for binding to CD38 or a portion thereof;
- a CD38BP that competes with an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, such as the antibody -005, for binding to CD38 or a portion thereof; and
- a CD38BP that competes with an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, such as the antibody -024, for binding to CD38 or a portion thereof.

[0223] As discussed elsewhere herein, unless otherwise stated or clearly contradicted by context, references to binding of a CD38BP to CD38 are intended to refer to binding in any suitable context, such as in a conformational context where the structure of CD38 is present; or in a linear epitope context. Of course, binding in a limited subset of such context(s) may be an important characteristic with respect to any CD38BP.

[0224] Additional methods for determining CD38BP specificity by competitive inhibition may be found in for instance

Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1988), Colligan et al., eds., *Current Protocols in Immunology*, Greene Publishing Assoc. and Wiley InterScience N.Y., (1992, 1993), and Muller, *Meth. Enzymol.* 92, 589-601 (1983)).

[0225] Human CD38 comprises a number of different epitopes, which may include (1) peptide antigenic determinants that are comprised within single peptide chains within human CD38; (2) conformational antigenic determinants which consist one or more noncontiguous amino acids on a particular chain and/or amino acids present on spatially contiguous but separate peptide chains (typically where the respective amino acid sequences of the chains are located disjointedly along the human CD38 polypeptide sequence); (3) post-translational antigenic determinants which consist, either in whole or part, of molecular structures covalently attached to human CD38, such as carbohydrate groups; or (4) combinations of (1)-(3).

[0226] An epitope in the context of the present invention includes any peptide or peptide-derivative determinant capable of specific binding to an immunoglobulin. An epitope may comprise any suitable number of amino acids, in any suitable position (with respect to the linear sequence of CD38) orientation (with respect to folded CD38, or a fragment thereof), amino acid composition (and consequently, at least in part, charge). Thus, for example, an epitope may be composed of about 3-10 amino acids, typically 3-8 amino acids, in one or more contiguous or noncontiguous locations with respect to the primary sequence of CD38 (for instance an epitope may consist essentially of 2, 3, 4, 5, 6, 7, or 8 amino acid residues distributed in 1, 2, 3, 4, or 5 noncontiguous locations in CD38). Alternatively, for example, an epitope may be considered to be defined by a region of about 5-40 contiguous amino acid residues (e.g., about 7-30 amino acid residues, about 5-20 amino acid residues, or about 3-15 amino acid residues) in CD38 (solely or in combination with a portion of an adjacent CD38 domain). In some epitopes it may be the case that just one amino acid residue or only a few amino acid residues are critical to CDR or CDR(s) recognition (and thereby most important to CD38BP:CD38 antigen affinity and avidity). As such, an epitope may be characterized on the basis of one or more of such critical residues, with the recognition that other residues may also make some lesser contribution to the epitope. In the case of an epitope defined by a region of amino acids, it may be that one or more amino acids in the region make only a minor contribution or even negligible contribution to antibody binding, such that the residue may be subject to substitution with an appropriate different residue without resulting in "a loss" of the epitope to at least some CD38BPs specific for it.

[0227] Described herein is a CD38BP, such as an anti-CD38 antibody, that specifically binds to a CD38 epitope that also is specifically bound by an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as antibody -003), or an antibody in accordance with the invention having a V_L sequence of SEQ ID No: 12 and a V_H sequence of SEQ ID No:17 (such as antibody -005) or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 (such as antibody -024). It is possible that CD38BPs having one or more CDRs that differ from the CDRs of an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, or the CDRs of an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, or the CDRs of an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, may still be specific for the same epitope as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, and an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 and an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27, respectively. In some such cases, the CD38BP in question may recognize or be more specific/selective for particular structures or regions of the epitope than the antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7, and the antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17, and the antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 respectively.

[0228] A CD38 epitope bound by an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as the antibody -003), or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 (such as the antibody -005) or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 (such as antibody -024), may be identified via standard mapping and characterization techniques, further refinement of which may be identified by any suitable technique, numerous examples of which are available to the skilled artisan. These techniques may also be used to identify and/or characterize epitopes for CD38BPs generally. As one example of such mapping/characterization methods, an epitope for an anti-CD38 antibody may be determined by epitope "foot-printing" using chemical modification of the exposed amines/carboxyls in the CD38 protein. One specific example of such a foot-printing technique is the use of HXMS (hydrogen-deuterium exchange detected by mass spectrometry) wherein a hydrogen/deuterium exchange of receptor and ligand protein amide protons, binding, and back exchange occurs, wherein the backbone amide groups participating in protein binding are protected from back exchange and therefore will remain deuterated. Relevant regions may be identified at this point by peptic proteolysis, fast microbore high-performance liquid chromatography separation, and/or electrospray ionization mass spectrometry. See, e.g., Ehring H, *Analytical Biochemistry*, 267(2) 252-259 (1999) and/or Engen, J.R. and Smith, D.L. (2001) *Anal. Chem.* 73, 256A-265A. Another example of a suitable epitope identification technique is nuclear magnetic resonance epitope mapping (NMR), where typically the position of the signals in two-dimensional NMR spectra of the free antigen and the antigen complexed with the antigen binding peptide, such as an antibody, are compared. The antigen typically is selectively

isotopically labeled with ^{15}N so that only signals corresponding to the antigen and no signals from the antigen binding peptide are seen in the NMR-spectrum. Antigen signals originating from amino acids involved in the interaction with the antigen binding peptide typically will shift position in the spectra of the complex compared to the spectra of the free antigen, and the amino acids involved in the binding may be identified that way. See for instance Ernst Schering Res Found Workshop. (44), 149-67 (2004), Huang et al., Journal of Molecular Biology 281(1), 61-67 (1998) and Saito and Patterson, Methods. 9(3), 516-24 (1996).

[0229] Epitope mapping/characterization may also be performed using mass spectrometry methods. See for instance Downard, J Mass Spectrom. 35(4), 493-503 (2000) and Kiselar and Downard, Anal Chem. 71(9), 1792-801 (1999).

[0230] Protease digestion techniques may also be useful in the context of epitope mapping and identification. Antigenic determinant-relevant regions/sequences may be determined by protease digestion, e.g. by using trypsin in a ratio of about 1:50 to CD38 overnight (O/N) digestion at 37°C and pH 7-8, followed by mass spectrometry (MS) analysis for peptide identification. The peptides protected from trypsin cleavage by the CD38BP may subsequently be identified by comparison of samples subjected to trypsin digestion and samples incubated with CD38BP and then subjected to digestion by e.g. trypsin (thereby revealing a foot print for the binder). Other enzymes like chymotrypsin, pepsin, etc. may also or alternatively be used in a similar epitope characterization method. A CD38BP which gives the significantly same result as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as the antibody -003), or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 (such as the antibody -005) or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 (such as antibody -024) in these measurements are deemed to be an antibody that bind the same epitope as an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as the antibody -003), or an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 (such as the antibody -005) or an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 (such as antibody -024), respectively. See for instance Manca, Ann Ist Super Sanita. 27(1), 15-9 (1991) for a discussion of similar techniques.

[0231] Epitope mapping by competitive binding to CD38 with two antibodies where one is biotinylated is another method for identifying relevant antigenic determinant regions.

[0232] The binding of antibodies to linear and looped peptides of CD38 by a PEPSCAN-based enzyme-linked immuno assay is another method for identifying relevant antigenic determinant regions, see for instance Sloodstra-JW et al. Mol-Divers. 1, 87-96 (1996).

[0233] Site directed mutagenesis is another method for identifying relevant antigenic determinant regions, see for instance Polyak and Deans, Blood 99, 3956-3962 (2002).

[0234] Various phage display techniques may also be used to identify epitopes. See for instance Wang and Yu, Curr Drug Targets. 5(1), 1-15 (2004), Burton, Immunotechnology. 1(2), 87-94 (1995 Aug), Cortese et al., Immunotechnology. 1(2), 87-94 (1995) and Irving et al., Curr Opin Chem Biol. 5(3), 314-24 (2001). Consensus epitopes may also be identified through modified phage display-related techniques (see, <http://www.cs.montana.edu/~mumey/papers/jcb03.pdf>) for discussion.

[0235] Other methods potentially helpful in mapping epitopes include crystallography techniques, X-ray diffraction techniques (such as the X-ray diffraction/sequence study techniques developed by Poljak and others in the 1970s-1980s), and the application of Multipin Peptide Synthesis Technology. Computer-based methods such as sequence analysis and three dimensional structure analysis and docking may also be used to identify antigenic determinants. For example, an epitope may also be determined by molecular modeling using a structure of CD38 with docking of the structure of the Fab fragment of the individual monoclonal antibody. These and other mapping methods are discussed in Epitope Mapping A Practical Approach (Westwood and Hay Eds.) 2001 Oxford University Press.

[0236] Described herein is a CD38BP having substantially the same specific CD38-binding characteristics of one or more mAbs selected from an antibody having a V_L sequence of SEQ ID No:2 and a V_H sequence of SEQ ID No:7 (such as the antibody -003), an antibody having a V_L sequence of SEQ ID No:12 and a V_H sequence of SEQ ID No:17 (such as antibody -005), and an antibody having a V_L sequence of SEQ ID No:22 and a V_H sequence of SEQ ID No:27 (such as antibody -024).

[0237] Mapping studies have indicated that several monoclonal antibodies raised against human CD38 bind to epitopes in the C-terminal region of CD38 (220-296) (Hoshino et al. J. Immunol., 1997, 158: 741-747, and Ferrero et al., BMC Immunology, 2004, 5: 21). Within this region three amino acid differences have been found between the human and the cynomolgus CD38 sequence: T237, Q272 and S274 in humans correspond to A238, R273 and F275 in cynomolgus. -005 does not bind to cynomolgus tissue (shown in examples 10 and 11). A limited number of amino acid differences exist between the human and the monkey CD38 sequence, for instance in the carboxyterminal part to the protein, for instance the following three amino acid differences between the human and the cynomolgus CD38 sequence: T237, Q272 and S274 in human CD38s correspond to A238, R273 and F275 in cynomolgus monkey CD38 (compare SEQ ID No.21 and SEQ ID No.22). -005 does not bind to a mutant huCD38 protein, wherein the glutamine residue at position 272 of SEQ ID No:31 has been substituted with an arginine residue (Q272R), or to a mutant huCD38 protein, wherein the serine residue of position 274 of SEQ ID No:31 has been substituted with a phenylalanine residue (S274F) (shown

in Example 17) to the same degree that it binds to wild type human CD38. Binding of -005 is particularly abrogated by the amino acid substitution at position S274F.

[0238] Consequently, an antibody of the present invention may be an antibody which binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33) to the same degree that it binds to human CD38 (SEQ ID No:31).

[0239] An antibody of the present invention may be an antibody which also binds to human CD38 (SEQ ID No:31), and which does not bind to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34) to the same degree that it binds to human CD38 (SEQ ID No:31).

[0240] The term "to the same degree" should be interpreted so that the binding of the peptide to the mutant human CD38 is significantly lower than the binding of the peptide to the wild type human CD38. The binding of a peptide to the CD38 molecules (wild type and mutant) may be determined in a number of ways and it is within the common general knowledge of a person skilled in the art to determine whether the binding to the mutant is "significantly lower" than the binding to the wildtype. A large number of different techniques for determining the binding of a peptide to another peptide are available to the person skilled in the art, for example ELISA, radioimmunoassay, BIAcore or flow cytometry.

[0241] One method of determining the binding is by determining the EC_{50} of the binding of the peptide to the mutant protein and to the wild type protein and then comparing the values obtained. Another method of determining the binding is by examining the magnitude of binding at saturating concentration (for instance the plateau of binding signal), or by determining kinetic rate constants k_{on} and k_{off} for example by BIAcore.

[0242] In one embodiment, the binding of the antibody in question to the CD38 proteins (mutant or wild type) is by use of an ELISA as described in Example 17.

[0243] In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 50% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 10% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 5% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the serine residue in position 274 has been substituted with a phenylalanine residue (SEQ ID No:34), is less than 1% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

[0244] In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), is less than 50% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the glutamine residue in position 272 has been substituted with an arginine residue (SEQ ID No:33), is less than 10% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

[0245] In one embodiment, an antibody of the invention may be an antibody which also binds to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) to the same degree that it binds to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) is more than 75% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) is more than 85% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) is more than 90% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31). In one embodiment, the EC_{50} of the binding of the antibody to a mutant human CD38, wherein the threonine residue in position 237 has been substituted with an alanine residue (SEQ ID No:32) is more than 95% of the EC_{50} of the binding of the antibody to human CD38 (SEQ ID No:31).

[0246] To identify more specific likely antigenic determinant regions in CD38, various predictive analytical methods may be applied. In a first analytical approach, CD38 may be analyzed for (1) highly hydrophobic regions (using the Kyte-Doolittle method); (2) antigenicity as measured by the Protrusion Index method; (3) antigenicity as determined by the Parker method; (4) antigenicity as determined by the Hopp/Woods method; and (5) hydrophilicity as measured by the methods of Goldman, Engleman, and Steitz. Sequences ranging from 10-40 amino acids in length may be selected based on exhibiting one or more of these properties. The rationale for this approach is the general consensus that many ideal B cell epitopes are hydrophilic, surface-oriented, and flexible sequences of about 8-10 amino acids in length.

[0247] Provided herein are CD38BPs specific for CD38-regions of CD38 identified in such a manner. Moreover, the termini of these sequences may be compared to predicted antigenic determinant regions located through the other analyses described herein to provide additional specific likely antigenic-determinant containing regions. Other similar

comparisons may readily be made to provide additional likely antigenic determinant regions, where CD38BPs binding to these antigenic determinant regions may be considered another feature of the present disclosure.

[0248] CD38BPs of the present invention are antibodies. Non-limiting examples of CD38 binding antibodies provided by the present invention include (a) a complete functional, immunoglobulin molecule comprising: (i) two identical chimeric heavy chains comprising a variable region with a human B cell surface antigen specificity and human constant region and (ii) two identical all (i.e. non-chimeric) human light chains; (b) a complete, functional, immunoglobulin molecule comprising: (i) two identical chimeric heavy chains comprising a variable region as indicated, and a human constant region, and (ii) two identical all (i.e. non-chimeric) non-human light chains; (c) a monovalent antibody, i.e., a complete, functional immunoglobulin molecule comprising: (i) two identical chimeric heavy chains comprising a variable region as indicated, and a human constant region, and (ii) two different light chains, only one of which has the same specificity as the variable region of the heavy chains. The resulting antibody molecule binds only to one end thereof and is therefore incapable of divalent binding. As another illustration, antibodies provided by the present invention may be said to include the following: (a) a whole immunoglobulin molecule; (b) an scFv; (c) a monoclonal antibody; (d) a human antibody; (e) a chimeric antibody; (f) a humanized antibody; (g) a Fab fragment; (h) an Fab' fragment; (i) an F(ab')₂ fragment; (j) an Fv molecule; and (k) a disulfide-linked Fv molecule.

[0249] In one embodiment, the antibody of the present invention is a polyclonal antibody. In one embodiment, the antibody of the present invention is a monoclonal antibody. In a further embodiment, the antibody of the present invention is a human monoclonal antibody. In another further embodiment, the antibody of the present invention is a humanized antibody. In another further embodiment, the antibody of the present invention is a chimeric antibody. In another further embodiment, the antibody of the present invention is a monoclonal antibody originating entirely from a mammalian species different from humans. In a further embodiment, the antibody of the present invention is a fully murine monoclonal antibody.

[0250] A monoclonal antibody refers to a composition comprising a homogeneous antibody population having a uniform structure and specificity. Typically a monoclonal antibody is an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific and each monoclonal antibody is typically directed against a single epitope, which is in contrast to polyclonal antibody preparations which typically include different antibodies directed against different epitopes. That an antibody is monoclonal is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies of the present invention may be produced by the hybridoma method first described by Kohler et al., *Nature* 256, 495 (1975), or may be produced by recombinant DNA methods. Monoclonal antibodies may also be isolated from phage antibody libraries using the techniques described in, for example, Clackson et al., *Nature* 352, 624-628 (1991) and Marks et al., *J. Mol. Biol.* 222, 581-597 (1991).

[0251] Monoclonal antibodies may be obtained from any suitable source. Thus, for example, monoclonal antibodies may be obtained from hybridomas prepared from murine splenic B cells obtained from mice immunized with an antigen of interest, for instance in form of cells expressing the antigen on the surface, or a nucleic acid encoding an antigen of interest. Monoclonal antibodies may also be obtained from hybridomas derived from antibody-expressing cells of immunized humans or non-human mammals such as rats, dogs, primates, etc.

[0252] Alternatively, the cloned antibody genes can be expressed in other expression systems, including prokaryotic cells, such as microorganisms, such as *E. coli*, for the production of single chain Fv antibodies, *algae*, as well as insect cells. Furthermore, the antibodies can be produced in transgenic non-human animals, such as in milk from sheep and rabbits or in eggs from hens, or in transgenic plants. See for instance Verma, R., et al., *J. Immunol. Meth.* 216, 165-181 (1998); Pollock, et al., *J. Immunol. Meth.* 231, 147-157 (1999); and Fischer, R., et al., *Biol. Chem.* 380, 825-839 (1999).

[0253] In one embodiment, human monoclonal antibodies directed against CD38 may be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system. Such transgenic and transchromosomal mice include mice referred to herein as HuMAb mice and KM mice, respectively, and are collectively referred to herein as "transgenic mice". A human monoclonal antibody generated in such mice may be abbreviated as HuMAb.

[0254] The HuMAb mouse contains a human immunoglobulin gene miniloci that encodes unrearranged human heavy (μ and γ) and κ light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous μ and κ chain loci (Lonberg, N. et al., *Nature* 368, 856-859 (1994)). Accordingly, the mice exhibit reduced expression of mouse IgM or κ and in response to immunization, the introduced human heavy and light chain transgenes, undergo class switching and somatic mutation to generate high affinity human IgG, κ monoclonal antibodies (Lonberg, N. et al. (1994), *supra*; reviewed in Lonberg, N. *Handbook of Experimental Pharmacology* 113, 49-101 (1994), Lonberg, N. and Huszar, D., *Intern. Rev. Immunol.* Vol. 13 65-93 (1995) and Harding, F. and Lonberg, N. *Ann. N.Y. Acad. Sci.* 764 536-546 (1995)). The preparation of HuMAb mice is described in detail in Taylor, L. et al., *Nucleic Acids Research* 20, 6287-6295 (1992), Chen, J. et al., *International Immunology* 5, 647-656 (1993), Tuailon et al., *J. Immunol.* 152, 2912-2920 (1994), Taylor, L. et al., *International Immunology* 6, 579-591 (1994), Fishwild, D. et al., *Nature Biotechnology* 14, 845-851

(1996). See also US 5,545,806, US 5,569,825, US 5,625,126, US 5,633,425, US 5,789,650, US 5,877,397, US 5,661,016, US 5,814,318, US 5,874,299, US 5,770,429, US 5,545,807, WO 98/24884, WO 94/25585, WO 93/1227, WO 92/22645, WO 92/03918 and WO 01/09187.

[0255] The HCo7 mice have a JKD disruption in their endogenous light chain (kappa) genes (as described in Chen et al., EMBO J. 12, 821-830 (1993)), a CMD disruption in their endogenous heavy chain genes (as described in Example 1 of WO 01/14424), a KCo5 human kappa light chain transgene (as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996)), and a HCo7 human heavy chain transgene (as described in US 5,770,429).

[0256] The HCo12 mice have a JKD disruption in their endogenous light chain (kappa) genes (as described in Chen et al., EMBO J. 12, 821-830 (1993)), a CMD disruption in their endogenous heavy chain genes (as described in Example 1 of WO 01/14424), a KCo5 human kappa light chain transgene (as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996)), and a HCo12 human heavy chain transgene (as described in Example 2 of WO 01/14424). In the KM mouse strain, the endogenous mouse kappa light chain gene has been homozygously disrupted as described in Chen et al., EMBO J. 12, 811-820 (1993) and the endogenous mouse heavy chain gene has been homozygously disrupted as described in Example 1 of WO 01/09187. This mouse strain carries a human kappa light chain transgene, KCo5, as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996). This mouse strain also carries a human heavy chain transchromosome composed of chromosome 14 fragment hCF (SC20) as described in WO 02/43478.

[0257] The KM mouse contains a human heavy chain transchromosome and a human kappa light chain transgene. The endogenous mouse heavy and light chain genes also have been disrupted in the KM mice such that immunization of the mice leads to production of human immunoglobulins rather than mouse immunoglobulins. Construction of KM mice and their use to raise human immunoglobulins is described in detail in WO 02/43478.

[0258] Splenocytes from these transgenic mice may be used to generate hybridomas that secrete human monoclonal antibodies according to well known techniques. Such transgenic mammals, mammals comprising an operable nucleic acid sequence coding for expression of a CD38BP, such as an antibody of the present invention, mammals stably transfected with one or more CD38-encoding nucleic acid sequences, and the like, are additional features involving antibodies of the present invention.

[0259] Human monoclonal or polyclonal antibodies of the present invention, or antibodies of the present invention originating from other species may also be generated transgenically through the generation of another non-human mammal or plant that is transgenic for the immunoglobulin heavy and light chain sequences of interest and production of the antibody in a recoverable form therefrom. In connection with the transgenic production in mammals, antibodies may be produced in, and recovered from, the milk of goats, cows, or other mammals. See for instance US 5,827,690, US 5,756,687, US 5,750,172 and US 5,741,957.

[0260] Further, human antibodies of the present invention or antibodies of the present invention from other species may be generated through display-type technologies, including, without limitation, phage display, retroviral display, ribosomal display, and other techniques, using techniques well known in the art and the resulting molecules may be subjected to additional maturation, such as affinity maturation, as such techniques are well known in the art (see for instance Hoogenboom et al., J. Mol. Biol. 227, 381 (1991) (phage display), Vaughan et al., Nature Biotech 14, 309 (1996) (phage display), Hanes and Plutchau, PNAS USA 94, 4937-4942 (1997) (ribosomal display), Parmley and Smith, Gene 73, 305-318 (1988) (phage display), Scott TIBS 17, 241-245 (1992), Cwirla et al., PNAS USA 87, 6378-6382 (1990), Russel et al., Nucl. Acids Research 21, 1081-1085 (1993), Hoogenboom et al., Immunol. Reviews 130, 43-68 (1992), Chiswell and McCafferty TIBTECH 10, 80-84 (1992), and US 5,733,743). If display technologies are utilized to produce antibodies that are not human, such antibodies may be humanized, for instance as described elsewhere herein.

[0261] Humanized monoclonal antibodies of the present invention may be generated by fusing the constant domains from a human antibody to the variable domains of a non-human species. Examples of how to make humanized antibodies may be found in for instance US 6,054,297, US 5,886,152 and US 5,877,293. A humanized antibody is designed to have greater homology to a human immunoglobulin than animal-derived monoclonal antibodies. Non-human amino acid residues from an "import" (animal) variable domain typically are transfected into a human "backbone". Humanization may essentially be performed following the method of Winter and co-workers (Jones et al., Nature 321, 522-525 (1986), Riechmann et al., Nature 332, 323-327 (1988), Verhoeyen et al., Science 239, 1534-1536 (1988)), by substituting rodent complementarity determining regions ("CDRs") or CDR sequences for the corresponding sequences of a human antibody. Accordingly, in such "humanized" antibodies, the CDR portions of the human variable domain have been substituted by the corresponding sequence from a non-human species. Thus, humanized antibodies are typically human antibodies in which some CDR residues and possibly some framework residues are substituted by residues from analogous sites in rodent antibodies. The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims et al., J. Immunol. 151, 2296 (1993), Chothia et al., J. Mol. Biol. 196, 901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light

or heavy chains. The same framework may be used for several different humanized antibodies (Carter et al., PNAS USA 89, 4285 (1992), Presta et al., J. Immunol. 151, 2623 (1993)).

[0262] It is typically also important that humanized antibodies retain high affinity for the antigen and other favorable biological properties. To achieve this goal, humanized antibodies may be prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of certain residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues may be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is maximized, although it is the CDR residues that directly and most substantially influence antigen binding.

[0263] Murine antibodies or antibodies from other species may be humanized or primatized using any suitable techniques, a number of suitable techniques being already well known in the art (see for instance Winter and Harris Immunol Today 14, 43-46 (1993) and Wright et al., Crit. Reviews in Immunol. 125-168 (1992)). The antibody of interest may be engineered by recombinant DNA techniques to substitute the C_H1, C_H2, C_H3, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190 and US 5,530,101, US 5,585,089, US 5,693,761, US 5,693,792, US 5,714,350, and US 5,777,085).

[0264] Humanization of antibodies may also be performed following the method of Winter and co-workers (Jones et al., Nature 321, 522-525 (1986), Riechmann et al., Nature 332, 323-327 (1988), Verhoeven et al., Science 239, 1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are, in a sense, chimeric antibodies (US 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0265] Also, the use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (see for instance Liu et al., PNAS USA 84, 3439 (1987) and J. Immunol. 139, 3521 (1987)). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA: The cDNA of interest may be amplified by the polymerase chain reaction using specific primers (US 4,683,195 and US 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. Sequences of human constant regions (as well as variable regions) may be found in Kabat et al., (1991) Sequences of Proteins of Immunological Interest, N.I.H. publication no. 91-3242 and more recent and related data can be accessed at <http://www.biochem.ucl.ac.uk/~martin/abs/GeneralInfo.html>. The choice of isotype typically will be guided by the desired effector functions, such as complement fixation, or activity in antibody-dependent cellular cytotoxicity. Exemplary isotypes are IgG1, IgG2, IgG3, and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody may then be expressed by conventional methods.

[0266] Antibodies of the present invention may be in any suitable form with respect to multimerization. Anti-CD38 antibodies and antibody fragments may be at least in heterotrimeric form if not in higher multimeric forms such as those associated with IgM antibodies. In other embodiments, the antibody may be presented as a dimer or monomer. Monomeric antibodies of the present invention may be, for example, modified by any suitable technique so as to form multimeric compositions.

[0267] If desired, the class of an anti-CD38 antibody of the present invention may be switched by known methods. For example, an antibody of the present invention that was originally IgM may be class switched to an IgG antibody of the present invention. Further, class switching techniques may be used to convert one IgG subclass to another, for instance from IgG1 to IgG2. Thus, the effector function of the antibodies of the present invention may be changed by isotype switching to, e.g., an IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody for various therapeutic uses.

[0268] In one embodiment an antibody of the present invention is an IgG1 antibody, for instance an IgG1,K or IgG1, λ isotype. In another embodiment an antibody of the present invention is an IgG3 antibody, for instance an IgG3, κ or IgG3, λ isotype. In another embodiment an antibody of the present invention is an IgG4 antibody, for instance an IgG4, κ or IgG4, λ isotype. In another embodiment an antibody of the present invention is an IgA1 or IgA2 antibody. In another embodiment an antibody of the present invention is an IgM antibody.

[0269] Anti-CD38 antibodies may be recovered from recombinant combinatorial antibody libraries, such as a scFv phage display library, which may be made with human V_L and V_H cDNAs prepared from mRNA derived from human lymphocytes. Methods for preparing and screening such libraries are known in the art. There are a number of commercially available kits for generating phage display libraries. There are also other methods and reagents that may be used in generating and screening antibody display libraries (see for instance US 5,223,409, WO 92/18619, WO 91/17271, WO 92/20791, WO 92/15679, WO 93/01288, WO 92/01047, WO 92/09690, Fuchs et al., Bio/Technology 9, 1370-1372 (1991),

Hay et al., Hum. Antibod. Hybridomas 3, 81-85 (1992), Huse et al., Science 246,1275-1281 *(1989), McCafferty et al., Nature 348, 552-554 (1990), Griffiths et al., EMBO J 12, 725-734 (1993), Hawkins et al., J. Mol. Biol. 226, 889-896 (1992), Clackson et al., Nature 352, 624-628 (1991), Gram et al., PNAS USA 89, 3576-3580 (1992), Garrad et al., Bio/Technology 9, 1373-1377 (1991), Hoogenboom et al., Nuc Acid Res 19, 4133-4137 (1991) and Barbas et al., PNAS USA 88, 7978-7982 (1991)). Suitable V_L and V_H nucleic acid sequences may be selected using any appropriate method. For example, V_L and V_H nucleic acids may be selected by employing the epitope imprinting methods described in WO 93/06213. Antibody libraries, such as scFv libraries may be prepared and screened using known and suitable methods (with human CD38-containing peptides as antigen(s)), such as those described in for instance WO92/01047, McCafferty et al., Nature 348, 552-554 (1990) and Griffiths et al., EMBO J 12, 725-734 (1993). Such antibody libraries and other combinations of CD38BPs (libraries, pools, etc.) are features involving antibodies of the present invention that may be used therapeutically to provide a more comprehensive immune response; as tools in screening methods for immunogenic peptides, small molecules, other anti-CD38 antibodies (e.g., by way of competition assays), and the like; and/or in diagnostic methods and compositions (e.g., an immunoassay chip comprising a panel of such antibodies optionally in association with other antibodies may be prepared by standard techniques). Once initial human V_L and V_H segments are selected, "mix and match" experiments, in which different pairs of the initially selected V_L and V_H segments are screened for CD38-containing peptide binding, may be performed to select desirable V_L/V_H pair combinations. For example, reactivity of the peptides may be determined by ELISA or other suitable epitope analysis methods (see for instance Scott, J. K. and Smith, G. P. Science 249, 386-390 (1990), Cwirla et al., PNAS USA 87, 6378-6382 (1990), Felici et al., J. Mol. Biol. 222, 301-310 (1991) and Kuwabara et al., Nature Biotechnology 15, 74-78 (1997) for discussion of such techniques and principles). Antibodies may be selected by their affinity for antigen and/or by their kinetics of dissociation (off-rate) from antigen (see for instance Hawkins et al., J. Mol. Biol. 226, 889-896 (1992)).

[0270] To further improve the quality and/or diversity of anti-CD38 antibodies, the V_L and V_H segments of V_L/N_H pair(s) may be randomly mutated, for instance within the CDR3 region of V_H and/or V_L , in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during a natural immune response. This *in vitro* affinity maturation may be accomplished by amplifying V_H and V_L regions using PCR primers complimentary to the V_H CDR3 or V_L CDR3, respectively, which primers typically are "spiked" with a random mixture of the four nucleotide bases at certain positions, such that the resultant PCR products encode V_H and V_L segments into which random mutations have been introduced into the V_H and/or V_L CDR3 regions. These randomly mutated V_H and V_L segments may be re-screened for binding to CD38-containing peptides.

[0271] Following screening, nucleic acid encoding a selected antibody may be recovered from the display package (e.g., from the phage genome) and subcloned into an appropriate vector by standard recombinant DNA techniques. If desired, such an antibody-encoding nucleic acid may be further manipulated to create other antibody forms or CD38BPs. To express a recombinant antibody isolated by screening of a combinatorial library, typically a nucleic acid comprising a sequence encoding the antibody is cloned into a recombinant expression vector and introduced into appropriate host cells (mammalian cells, yeast cells, etc.) under conditions suitable for expression of the nucleic acid and production of the antibody.

[0272] High-affinity antibody peptides, such as human single-chain Fv (scFv) and Fab antibody fragments, may also be isolated from such libraries using a panning technique in which the antigen of interest is immobilized on a solid surface, such as microtiter plates or beads (see for instance Barbas and Burton, Trends. Biotechnol. 14, 230-234 (1996) and Aujame et al., Hum. Antibodies 8, 155-68 (1997). Phage display of large naïve libraries also makes it possible to isolate human antibodies directly without immunization (see for instance de Haard et al., J. Biol. Chem. 274(26), 18218-18230 (1999)).

[0273] A "variant" anti-CD38 antibody is an antibody that differs from a parent antibody (typically generated by immunization) by one or more suitable amino acid residue alterations, that is substitutions, deletions, insertions, or terminal sequence additions, in the CDRs or other V_H and/or V_L sequences (provided that at least a substantial amount of the epitope binding characteristics of the parent antibody are retained, if not improved upon, by such changes).

[0274] Variations in an antibody variant may be made in each of the framework regions, the constant domain, and/or the variable regions (or any one or more CDRs thereof) in a single variant antibody. Alternatively, variations may be made in only one of the framework regions, the variable regions (or single CDR thereof), or the constant domain in an antibody. Alanine scanning mutagenesis techniques, such as described by Cunningham and Wells, Science 244, 1081-1085 (1989), may be used to identify suitable residues for substitution or deletion in generating CD38BPs comprising variant V_L , V_H , or particular CDR sequences, although other suitable mutagenesis techniques also may be applied. Multiple amino acid substitutions may also be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer, Science 241, 53-57 (1988) or Bowie and Sauer, PNAS USA 86, 2152-2156 (1989).

[0275] Thus, for example, in an antibody variant one or more amino acid residues may be introduced or inserted in or adjacent to one or more of the hypervariable regions of a parent antibody, such as in or adjacent to one or more CDRs. An anti-CD38 antibody variant may comprise any number of inserted amino acid residues, provided again that

at least a substantial amount of the epitope binding characteristics of the parent antibody are retained. An anti-CD38 antibody variant may for example comprise from about 1-30 inserted amino acid residues, for instance from about 1-10, such as for instance from about 2-10, for instance from 2-5 or such as from about 1-5 inserted amino acid residues. Likewise, an anti-CD38 antibody variant may for example comprise from about 1-30 deleted amino acid residues, for instance from about 1-10, such as for instance from about 2-10, for instance from 2-5 or such as from about 1-5 deleted amino acid residues. Likewise, an anti-CD38 antibody variant may for example comprise from about 1-30 substituted amino acid residues, for instance from about 1-10, such as for instance from about 2-10, for instance from 2-5 or such as from about 1-5 substituted amino acid residues. Likewise, an anti-CD38 antibody variant may for example comprise from about 1-30 terminal sequence amino acid residue additions, for instance from about 1-10, such as for instance from about 2-10, for instance from 2-5 or such as from about 1-5 terminal sequence amino acid residue additions. An antibody variant may also comprise a combination of two or more of such insertions, deletions, substitutions and terminal sequence amino acid residue additions, provided that the variant possesses at least a substantial proportion of the parent antibodies affinity, specificity, and/or selectivity with respect to one or more CD38 epitopes.

[0276] Considerations in the selection of antibody variants (e.g., conservation of amino acid residue functional characteristics, conservation of amino acid residues based on hydrophobic characteristics, and/or conservation of amino acid residues on the basis of weight/size), are described elsewhere herein. Typically, amino acid sequence alterations, such as conservative substitution variations, desirably do not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to disrupt secondary structure that characterizes the function of the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in, e.g., *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)), *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)) and Thornton et al., *Nature* 354, 105 (1991). Additional principles relevant to the design and construction of peptide variants is discussed in for instance Collinet et al., *J Biol Chem* 275(23), 17428-33 (2000).

[0277] Amino acid sequence variants of an antibody may be obtained by introducing appropriate nucleotide changes into the antibody-encoding nucleic acid (e.g., by site directed mutagenesis) or by chemical peptide synthesis. Such variants include, for example, deletions from, and/or insertions into and/or substitutions of and/or terminal sequence additions of residues within the amino acid sequences of the antibodies of the examples herein. Any combination of deletions, insertions, and substitutions may be made to arrive at a desired variant, provided that the variant possesses at least a substantial proportion of epitope binding characteristics of the parent antibody. Amino acid sequence changes, with respect to a parent antibody, also may alter post-translational processes of the variant antibody with respect to a parent antibody, such as by changing the number or position of glycosylation sites.

[0278] Variant antibodies may comprise alterations in the hypervariable region, such as in the CDRs. Examples of CD38BPs comprising such CDR variants are described elsewhere herein, and, as described above, such CD38BPs may be antibodies.

[0279] Variant antibodies of the present invention may comprise framework (FR) alterations, that is outside the hypervariable region, for instance in the Fc region, which alterations may be associated with advantageous properties, such as changing the functional or pharmacokinetic properties of the antibodies. For example, a substitution or other modification (insertion, deletion, terminal sequence additions or combination of any thereof) in a framework region or constant domain may be associated with an increase in the half-life of the variant antibody with respect to the parent antibody, or may be made to alter the immunogenicity of the variant antibody with respect to the parent antibody, to provide a site for covalent or non-covalent binding to another molecule, or to alter such properties as complement fixation, for instance resulting in a decrease or increase of C1q binding and CDC or of FcγR binding and antibody-dependent cellular cytotoxicity (ADCC). Substitutions may for example be made in one or more of the amino acid residues 234, 235, 236, 237, 297, 318, 320, and 322 of the heavy chain constant region, thereby causing an alteration in an effector function while retaining binding to antigen as compared with the unmodified antibody, cf. US 5,624,821 and US 5,648,260. Further reference may be had to WO 00/42072 disclosing antibodies with altered Fc regions that increase ADCC, and WO 94/29351 disclosing antibodies having mutations in the N-terminal region of the C_H2 domain that alter the ability of the antibodies to bind to FcRI and thereby decreases the ability of the antibodies to bind to C1 q which in turn decreases the ability of the antibodies to fix complement. Furthermore, Shields et al., *J. Biol. Chem.* 276, 6591-6604 (2001) teaches combination variants, that improve FcγRIII binding, for instance T256A/S298A, S298A/E333A, and S298A/E333A/K334A,.

[0280] The *in vivo* half-life of the antibodies may also be improved by modifying the salvage receptor epitope of the Ig constant domain or an Ig-like constant domain such that the molecule does not comprise an intact C_H2 domain or an intact Ig Fc region, cf. US 6,121,022 and US 6,194,551. The *in vivo* half-life may furthermore be increased by making mutations in the Fc region, e.g. by substituting threonine for leucine at position 252, threonine for serine at position 254, or threonine for phenylalanine at position 256, cf. US 6,277,375.

[0281] Anti-CD38 antibodies of the invention may be variant anti-CD38 antibodies wherein potential T cell epitopes in the antibody have been reduced or eliminated through rationale design. Thus, for example, in one embodiment the

present invention provides a "deimmunized" anti-CD38 antibody in which the potential T cell epitopes have been eliminated. The design and construction of deimmunized anti-CD38 antibodies may be accomplished by any suitable known technique (see for instance WO9852976 with respect to methods for preparing deimmunized antibodies). Immunogenicity in humans is expected to be eliminated or substantially reduced when such CD38BPs (e.g., anti-CD38 variant antibodies) are administered.

[0282] Other framework mutations may include sequence changes which may reduce susceptibility to proteolysis, reduce susceptibility to oxidation, and/or confer or modify other physicochemical or functional properties on the associated variant antibody.

[0283] Amino acid sequence variations in the framework may also result in an altered glycosylation pattern in the variant antibody with respect to a parent antibody. By altering is meant deleting one or more carbohydrate moieties found in the parent antibody, and/or adding one or more glycosylation sites that are not present in the parent antibody. Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are common recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide may create a potential glycosylation site. O-linked glycosylation refers to the attachment of sugars such as N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used. Addition of glycosylation sites to the antibody may be conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[0284] The antibodies may also be expressed in a transfectoma which does not add the fucose unit normally attached to the carbohydrate attached to Asn at position 297 of Fc in order to enhance the affinity of Fc for Fc γ RIII which in turn will result in an increased ADCC of the antibodies in the presence of NK cells, cf. Shield et al., J. Biol. Chem. 277, 26733 (2002). Other methods of modifying the glycosylation with focus on the fucosylation is described in WO 00/61739 to Kyowa. Furthermore, modification of galactosylation may be made in order to modify CDC. Further reference may be had to WO 99/54342 and Umana et al., Nat. Biotechnol. 17, 176 (1999) disclosing a CHO cell line engineered to express GntIII resulting in the expression of monoclonal antibodies with altered glycoforms and improved ADCC activity.

[0285] Other potentially suitable techniques for preparing novel anti-CD38 antibodies include CDR walking mutagenesis, antibody chain shuffling, "parsimonious mutagenesis" (Balint and Larrick, Gene 137, 109-118 (1993)), and other affinity maturation techniques (see for instance Wu et al., PNAS USA 95, 6037-42 (1998)). Repertoire cloning procedures may also be useful in the production of variant antibodies (see for instance WO 96/33279).

[0286] There are a number of techniques known for generating CDR variants, any suitable technique or combination of which may be used in the context of the present invention for generating CDR variants of the CDRs of the antibodies of the examples. Examples of such techniques include the removal of nonessential residues as described in Studnicka et al., Protein Engineering 7, 805-814 (1994) (see also Soderlind et al., Immunotechnology. 4(3-4), 279-85 (1999), CDR walking mutagenesis and other artificial affinity maturation techniques (see for instance Yang et al., Journal of Molecular Biology 254(3), 392-403 (1995), CDR shuffling techniques wherein typically CDRs are amplified from a diverse set of gene templates optionally comprising synthetic oligonucleotides, the constant regions of the V_L, V_H, and/or CDRs are amplified, and the various fragments mixed (in single-stranded or double-stranded format) and assembled by polymerase chain reaction (PCR) to produce a set of antibody-fragment encoding gene products carrying shuffled CDR introduced into the master framework, which is amplified using external primers annealing to sites beyond inserted restriction sites to ensure production of full-length products, which are inserted into a vector of choice and used to expressed variant CDR-containing proteins. Appropriate structure may be determined by superimposition of the variant/mimetic structures and those of the parent sequences, e.g., by comparison of NMR solution structures. Useful methods for rational design of CDR sequence variants are described in for instance WO 91/09967 and WO 93/16184. Additional examples of such methods are provided elsewhere herein.

[0287] The antibodies of the present invention include fragments of antibodies (including variant antibodies) of the present invention, which fragments has the ability to bind to CD38 (CD38 binding fragments). Antibodies of the invention thus include antibody-like molecules that comprise less than the full tetrameric structure associated with naturally-occurring antibodies. An antibody fragment may be any peptide that comprises a portion of a full length antibody, generally the antigen binding or variable region thereof (this includes, for example, fragments of humanized antibodies comprising the CDRs from an antibody of the present invention). In one embodiment, an antibody fragment refers to a peptide that consists essentially or consists only of a portion of an antibody molecule. In one embodiment, the present invention provides an antibody fragment comprising at least a portion of a heavy chain variable domain containing the three V_H CDRs of an antibody of the present invention and also a light chain-variable domain comprising the three V_L CDRs of an antibody of the present invention, wherein the heavy chain variable domain, and optionally the light chain variable domain, optionally is (are) fused to an additional moiety, such as an immunoglobulin constant domain. Constant domain

sequences may be added to the heavy chain and/or light chain sequence(s) to form species with partial length heavy and/or light chain(s). Constant regions, or portions thereof, of any antibody isotype may be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions.

[0288] Examples of CD38-binding antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments. An antibody fragment in the context of the present invention may also include a peptide comprising the CDRs of an antibody of the invention, and the like. In one embodiment, the present invention provides an antibody fragment comprising a first polypeptide chain that comprises the heavy chain CDRs of an antibody of the invention and a second polypeptide chain that comprises the light chain CDRs of an antibody of the invention, wherein the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In one embodiment, the present invention provides a two-chain antibody fragment having such features wherein the antibody fragment is selected from Fab, Fab', Fab'-SH, Fv, and/or F(ab')₂ fragments.

[0289] Antibodies may be fragmented using conventional techniques, and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')₂ fragments may be generated by treating antibody with pepsin. The resulting F(ab')₂ fragment may be treated to reduce disulfide bridges to produce Fab' fragments. Fab fragments may be obtained by treating an IgG antibody with papain; Fab' fragments may be obtained with pepsin digestion of IgG antibody. A Fab' fragment may also be produced by binding Fab' described below via a thioether bond or a disulfide bond. A Fab' fragment is an antibody fragment obtained by cutting a disulfide bond of the hinge region of the F(ab')₂. A Fab' fragment may be obtained by treating a F(ab')₂ fragment with a reducing agent, such as dithiothreitol. Antibody fragment peptides may also be generated by expression of nucleic acids encoding such peptides in recombinant cells (see for instance Evans et al., J. Immunol. Meth. 184, 123-38 (1995)). For example, a chimeric gene encoding a portion of a F(ab')₂ fragment could include DNA sequences encoding the C_H1 domain and hinge region of the H chain, followed by a translational stop codon to yield such a truncated antibody fragment molecule.

[0290] Antibodies of the invention also include univalent antibodies and single chain antibodies. Single chain antibodies are peptides in which the heavy and light chain Fv regions are connected. In one embodiment, the present invention provides a single-chain Fv (scFv) wherein the heavy and light chains in the Fv of an anti-CD38 antibody of the present invention are joined with a flexible peptide linker (typically of about 10, 12, 15 or more amino acid residues) in a single peptide chain. Methods of producing such antibodies are described in for instance US 4,946,778, Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds. Springer-Verlag, New York, pp. 269-315 (1994), Bird et al., Science 242, 423-426 (1988), Huston et al., PNAS USA 85, 5879-5883 (1988) and McCafferty et al., Nature 348, 552-554 (1990). The single chain antibody may be monovalent, if only a single V_H and V_L are used, bivalent, if two V_H and V_L are used, or polyvalent, if more than two V_H and V_L are used.

[0291] In one embodiment of the present invention, the antibody may be derivatized or linked to another functional molecule, for instance another peptide or protein (such as a Fab' fragment) to generate a bispecific or multispecific molecule which binds to multiple binding sites or target epitopes. For example, an antibody of the present invention may be functionally linked (for instance by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other binding molecules, such as another antibody, peptide or binding mimetic. Accordingly, the present invention includes bispecific and multispecific molecules comprising at least one first binding specificity for CD38 and a second binding specificity for a second target epitope. In one embodiment of the present invention, the second target epitope is an Fc receptor, e.g., human FcγRI (CD64) or a human Fcα receptor (CD89), or a T cell receptor, e.g., CD3. In one embodiment, the present invention provides bispecific and multispecific molecules capable of binding both to FcγR, FcαR or FcεR expressing effector cells (e.g., monocytes, macrophages or polymorphonuclear cells (PMNs)), and to target cells expressing CD38. These bispecific and multispecific molecules target CD38 expressing cells to effector cell and trigger Fc receptor-mediated effector cell activities, such as phagocytosis of CD38 expressing cells, antibody dependent cellular cytotoxicity (ADCC), cytokine release, or generation of superoxide anion.

[0292] Bispecific and multispecific molecules of the present invention may further include a third binding specificity, in addition to an anti-Fc binding specificity and an anti-CD38 binding specificity. In one embodiment, the third binding specificity is an anti-enhancement factor (EF) portion, e.g., a molecule which binds to a surface protein involved in cytotoxic activity and thereby increases the immune response against the target cell. The "anti-enhancement factor portion" may be an antibody, functional antibody fragment or a ligand that binds to a given molecule, e.g., an antigen or a receptor, and thereby results in an enhancement of the effect of the binding determinants for the Fc receptor or target cell antigen. The "anti-enhancement factor portion" may bind an Fc receptor or a target cell antigen. Alternatively, the anti-enhancement factor portion may bind to an entity that is different from the entity to which the first and second binding specificities bind. For example, the anti-enhancement factor portion may bind a cytotoxic T cell (e.g., via CD2, CD3, CD8, CD28, CD4, CD40, ICAM-1 or other immune cell that results in an increased immune response against the target cell).

[0293] In one embodiment, the bispecific and multispecific molecules of the present invention comprise as a binding specificity at least one further antibody, including, e.g., an Fab, Fab', F(ab')₂, Fv, or a scFv. The further antibody may also be a light chain or heavy chain dimer, or any minimal fragment thereof such as a Fv or a single chain construct as

described in Ladner et al., in US 4,946,778. The antibody may also be a binding-domain immunoglobulin fusion protein as disclosed in US 2003/0118592 and US 2003/0133939.

[0294] In one embodiment, the binding specificity for an Fc receptor is provided by a human monoclonal antibody, the binding of which is not blocked by human immunoglobulin G (IgG). As used herein, the term "IgG receptor" refers to any of the eight γ -chain genes located on chromosome 1. These genes encode a total of twelve transmembrane or soluble receptor isoforms which are grouped into three Fc γ receptor classes: Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). In one embodiment, the Fc γ receptor is a human high affinity Fc γ RI. The production and characterization of these monoclonal antibodies are described by Fanger et al., in WO 88/00052 and in US 4,954,617. These antibodies bind to an epitope of Fc γ RI, Fc γ RII or Fc γ RIII at a site which is distinct from the Fc γ binding site of the receptor and, thus, their binding is not blocked substantially by physiological levels of IgG. Specific anti-Fc γ RI antibodies useful in the present invention are mAb 22, mAb 32, mAb 44, mAb 62 and mAb 197. In other embodiments, the anti-Fc γ receptor antibody is a humanized form of mAb 22 (H22). The production and characterization of the H22 antibody is described in Graziano, R.F. et al., J. Immunol. 155(10), 4996-5002 (1995) and WO 94/10332. The H22 antibody producing cell line was deposited at the American Type Culture Collection on November 4, 1992 under the designation HA022CL1 and has the accession No. CRL 11177.

[0295] In one embodiment, the binding specificity for an Fc receptor is provided by an antibody that binds to a human IgA receptor, e.g., an Fc α receptor (Fc α I (CD89)), the binding of which in one embodiment is not blocked by human immunoglobulin A (IgA). The term "IgA receptor" is intended to include the gene product of one α -gene (Fc α RI) located on chromosome 19. This gene is known to encode several alternatively spliced transmembrane isoforms of 55 to 110 kDa. Fc α RI (CD89) is constitutively expressed on monocytes/macrophages, eosinophilic and neutrophilic granulocytes, but not on non-effector cell populations. Fc α RI has medium affinity for both IgA1 and IgA2, which is increased upon exposure to cytokines such as G-CSF or GM-CSF (Morton, H.C. et al., Critical Reviews in Immunology 16, 423-440 (1996)). Four Fc α RI-specific monoclonal antibodies, identified as A3, A59, A62 and A77, which bind Fc α RI outside the IgA ligand binding domain, have been described (Monteiro, R.C. et al., J. Immunol. 148, 1764 (1992)).

[0296] Fc α RI, Fc γ RI, Fc γ RII and Fc γ RIII, especially Fc γ RII and Fc γ RIII, are examples of trigger receptors for use in the present invention because they (1) are expressed primarily on immune effector cells, e.g., monocytes, PMNs, macrophages and dendritic cells; (2) are expressed at high levels (for instance 5,000-100,000 per cell); (3) are mediators of cytotoxic activities (for instance ADCC, phagocytosis); and (4) mediate enhanced antigen presentation of antigens, including self-antigens, targeted to them.

[0297] In one embodiment, an antibody of the present invention is a multispecific anti-CD38 antibody or antibody-like molecule, a particular example of which is a bispecific antibody comprising at least one pair of V_H sequence and V_L sequence chains specific for an epitope comprised at least in part in CD38 and a second at least one pair of V_H and V_L sequence chains specific for a second epitope. The V_H and V_L sequences in a bispecific antibody may comprise complete V_H and V_L sequences corresponding to anti-CD38 V_H and V_L regions, variant V_H and/or V_L sequences, or suitable portions of V_H and/or V_L regions, such as a suitable combination of CDR sequences and other sequences sufficient to provide binding to the epitopes of interest.

[0298] Exemplary bispecific antibody molecules comprise (i) two antibodies one with a specificity to CD38 and another to a second target that are conjugated together, (ii) a single antibody that has one chain specific to CD38 and a second chain specific to a second molecule, and (iii) a single chain antibody that has specificity to CD38 and a second molecule. Typically, the second target/second molecule is a molecule other than CD38. In one embodiment, the second molecule is a cancer antigen/tumor-associated antigen such as carcinoembryonic antigen (CEA), prostate specific antigen (PSA), RAGE (renal antigen), α -fetoprotein, CAMEL (CTL-recognized antigen on melanoma), CT antigens (such as MAGE-B5, -B6, -C2, -C3, and D; Mage-12; CT10; NY-ESO-1, SSX-2, GAGE, BAGE, MAGE, and SAGE), mucin antigens (e.g., MUC1, mucin-CA125, etc.), ganglioside antigens, tyrosinase, gp75, C-myc, Mart1, MelanA, MUM-1, MUM-2, MUM-3, HLA-B7, and Ep-CAM. In one embodiment, the second molecule is a cancer-associated integrin, such as α 5 β 3 integrin. In one embodiment, the second molecule is an angiogenic factor or other cancer-associated growth factor, such as a vascular endothelial growth factor (VEGF), a fibroblast growth factor (FGF), epidermal growth factor (EGF), epidermal growth factor receptor (EGFR), angiogenin, and receptors thereof, particularly receptors associated with cancer progression (for instance one of the HER1-HER4 receptors). Other cancer progression-associated proteins discussed herein may also be suitable second molecules. In one embodiment, the second molecule is a molecule expressed on the surface of multiple myeloma cells such as CD138.

[0299] In one embodiment, a bispecific antibody of the present invention is a diabody.

[0300] Bispecific antibodies also include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in a heteroconjugate may be coupled to avidin and the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (see for instance US 4,676,980). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable peptide cross-linking agents and techniques are well known in the art, and examples of such agents and techniques are disclosed in for instance US 4,676,980.

[0301] Thus, although the discussion herein may focus on antibodies, it should be understood that the embodiments

and features of the antibodies may equally be applied to antibody fragments, such as Fab fragments, Fab' fragments, and scFv peptides, antibody-like peptides (peptides comprising a CDR), bi- and multi-specific antibodies and other CD38BPs, as appropriate, provided that the antibody of the present invention retains at least a substantial proportion of the antigen-binding properties of the corresponding complete antibody. In some instances, antibody fragments may be associated with lower antigen-binding affinity, but may offer other advantageous features that may offset for any such loss in affinity.

[0302] Anti-CD38 antibodies in accordance with the present invention may be selected based on their ability to provide the ability of complement fixation, or not. There are a number of isotypes of antibodies that are capable of complement fixation and CDC, including, without limitation, the following: murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3. Those isotypes that do not include, without limitation, human IgG2 and human IgG4. Isotype determination and other methods for modifying the complement fixation and CDC functional characteristics of antibodies are known in the art.

[0303] CD38BPs as described herein also include immunoadhesins, which are molecules wherein one or more CDRs of an anti-CD38 antibody are covalently or noncovalently associated with the molecule. An immunoadhesin may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDRs permit the immunoadhesin to specifically bind to a CD38.

[0304] Antibodies of the present invention may also be provided as CD38BP fusion proteins. CD38BP fusion proteins may comprise any suitable amino acid sequence or combination of sequences that are specific and/or selective for at least one domain that is at least partially comprised within CD38 (e.g., an anti-CD38 antibody V_H domain, V_L domain, or particular CDRs thereof) and at least one nonhomologous and typically substantially nonsimilar amino acid sequence (e.g., less than about 40%, less than about 35%, less than about 30%, less than about 25%, or less than about 20% amino acid sequence identity to the CD38-specific/selective sequence) that imparts a detectable biological function and/or characteristic to the fusion protein that cannot solely be attributed to the CD38-specific/selective sequence (e.g., increased *in vivo* half-life, fluorescence, increased targeting to a particular type of cell, etc.). Functional sequences of such a fusion protein may be separated by flexible linker(s). Secondary sequence(s) may also be derived from cytotoxic or apoptotic peptides. Secondary sequences may also confer diagnostic properties. Examples of such sequences include those derived from easily visualized enzymes such as horseradish peroxidase.

[0305] CD38BP fusion proteins may also be characterized by comprising an epitope tag. An epitope tag sequence is an amino acid sequence having enough residues to provide an epitope against which an antibody may be made, in the context of the CD38BP, yet is short enough such that it does not substantially interfere with the activity (selectivity, specificity, affinity, and/or biological activity) of the CD38BP (as compared to a parent CD38BP lacking the epitope tag). An epitope tag desirably is sufficiently unique so that the anti-epitope tag antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least about 6 amino acid residues and usually between about 8-50 amino acid residues (e.g., about 9-30 residues). Examples of epitope tags include the flu HA tag polypeptide and its antibody 12CA5 (Field et al., Mol. Cell. Biol. 8, 2159-2165 (1988)); the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto (Evan et al., Mol. Cell. Biol. 5(12), 3610-3616 (1985)) and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody (Paborsky et al., Protein Engineering 3(6), 547-553 (1990)). In certain embodiments, the epitope tag is a "salvage receptor binding epitope". As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (for instance IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

[0306] Anti-CD38 antibodies of the present invention may also be used to produce anti-CD38 antibody derivatives. An antibody derivative comprises an antibody in which one or more of the amino acid residues of the antibody have been chemically modified (e.g. by alkylation, acylation, ester formation, or amide formation) or covalently associated with one or more heterologous substituents (e.g., a lipophilic substituent, a PEG moiety, a peptide side chain linked by a suitable organic moiety linker, etc.). The antibody may also be conjugated to a therapeutic moiety, such as a cytotoxin, a chemotherapeutic drug, an immunosuppressant, or a radioisotope (a so-called immunoconjugate). In general, CD38BPs described herein, such as anti-CD38 antibodies of the invention, may be modified by inclusion of any suitable number of such modified amino acids and/or associations with such conjugated substituents. Suitability in this context general is determined by the ability to at least substantially retain CD38 selectivity and/or specificity associated with the non-derivatized parent CD38BP. The inclusion of one or more modified amino acids may be advantageous in, for example, (a) increasing polypeptide serum half-life, (b) reducing polypeptide antigenicity, or (c) increasing polypeptide storage stability. Amino acid (s) are modified, for example, co-translationally or post-translationally during recombinant production (e.g., N-linked glycosylation at N-X-S/T motifs during expression in mammalian cells) or modified by synthetic means. Non-limiting examples of a modified amino acid include a glycosylated amino acid, a sulfated amino acid, a prenylated (e.g., farnesylated, geranylgeranylated) amino acid, an acetylated amino acid, an acylated amino acid, a PEGylated amino acid, a biotinylated amino acid, a carboxylated amino acid, a phosphorylated amino acid, and the like. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature. Example protocols are found in Walker (1998) Protein Protocols On Cd-Rom, Humana Press, Towata, NJ. The modified amino

acid may be selected from a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid / conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.

[0307] Additionally, antibodies may be chemically modified by covalent conjugation to a polymer to for instance increase their circulating half-life. Exemplary polymers, and methods to attach them to peptides, are illustrated in for instance US 4,766,106, US 4,179,337, US 4,495,285 and US 4,609,546. Additional illustrative polymers include polyoxyethylated polyols and polyethylene glycol (PEG) (e.g., a PEG with a molecular weight of between about 1,000 and about 40,000, such as between about 2000 and about 20,000, e.g., about 3,000-12,000).

[0308] Antibodies of the present invention may be conjugated to a second molecule that is selected from a radionuclide, an enzyme, an enzyme substrate, a cofactor, a fluorescent marker, a chemiluminescent marker, a peptide tag, or a magnetic particle. In one embodiment, an antibody may be conjugated to one or more antibody fragments, nucleic acids (oligonucleotides), nucleases, hormones, immunomodulators, chelators, boron compounds, photoactive agents, dyes, and the like. These and other suitable agents may be coupled either directly or indirectly to antibodies of the present invention. One example of indirect coupling of a second agent is coupling by a spacer moiety. These spacers, in turn, may be either insoluble or soluble (see for instance Diener et al., Science 231, 148 (1986)) and may be selected to enable drug release from the antibody at a target site and/or under particular conditions. Additional examples of therapeutic agents that may be coupled to antibodies include lectins and fluorescent peptides.

[0309] In one embodiment, antibody derivatives comprising one or more radiolabeled amino acids are provided. A radiolabeled antibody may be used for both diagnostic and therapeutic purposes (conjugation to radiolabeled molecules is another possible feature). Nonlimiting examples of labels for polypeptides include, but are not limited to ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{125}I , ^{131}I , and ^{186}Re . Methods for preparing radiolabeled amino acids and related peptide derivatives are known in the art (see for instance Junghans et al., in Cancer Chemotherapy and Biotherapy 655-686 (2d edition, Chafner and Longo, eds., Lippincott Raven (1996)) and US 4,681,581, US 4,735,210, US 5,101,827, US 5,102,990 (US RE35,500), US 5,648,471 and US 5,697,902. For example, a radioisotope may be conjugated by a chloramine T method.

[0310] Advantageous radionuclides in diagnostic contexts are indium isotopes and in the context of therapeutic applications yttrium isotopes, which are cytotoxic. Photon-emitting radioisotopes, in general, are advantageous in diagnostic (radioimmuno-scintigraphy (RIS)) methods. Auger electrons have a very short path length (5-10 nm) and need to be internalized to be cytotoxic (see for instance Adelstein et al., Nucl. Med. Biol. 14, 165-169 (1987)). Accordingly, antibodies conjugated to such isotopes may be useful in diagnostic methods, but only antibodies that are internalized should be considered for radioisotopes that emit Auger electrons in therapeutic contexts. Alpha particles need to be close to a cell (within 3-4 cell diameters) to be effective as therapeutic agents (Vriesendorp et al., "Radioimmunoglobulin therapy," in High Dose Cancer Therapy Armitage et al., (eds). (Williams & Wilkins, Baltimore, Md. 1992)). Both Auger electrons and alpha emitters may be considered to have high selectivity because their short-range emission typically will not irradiate neighboring normal cells.

[0311] The radiometals ^{111}In and ^{90}Y are, respectively, a pure γ -emitter and a pure β -emitter. Iodine-125, the most commonly used emitter of Auger electrons, has a half-life of about 60 days and frequently is released by immunoconjugates *in vivo* (due to dehalogenation). The most commonly considered alpha emitters for clinical use, astatine-211 and bismuth-212, have relatively short half-lives (7.2 h and 1.0 h, respectively) and decay into radioactive isotopes that may not be retained by the immunoconjugate after the first alpha emission (Wilbur, Antibiot. Immunoconjug. Radiopharm. 4, 5-97 (1991)). For diagnostic applications, CD38BPs, such as anti-CD38 antibodies of the invention, labeled with indium-111 or technetium-99m may be used. Both of these isotopes emit gamma rays within the appropriate energy range for imaging, (100-250 keV). Energies below this range typically are not penetrating enough to reach an external imaging device. Higher energy levels are difficult to collimate and provide diagnostic images with poor resolution. The short-half life of ^{99}Tc typically restricts its use to immunoconjugates with rapid tumor uptake.

[0312] In one embodiment, first and second CD38 antibodies conjugated with first and second radioisotopes are provided. In another embodiment, a single CD38 antibody conjugated with two radioisotopes is provided. An advantage of using two separate radioisotopes, e.g., one for imaging and one for therapy, is that it facilitates outpatient treatment. The low amount of radioactivity used diagnostically does not represent a radiation hazard, while the radiation emitted by a therapeutic isotope, such as a pure β -emitter, typically will largely be absorbed in the vicinity of the targeted cells.

[0313] Radioisotopes may be attached directly or indirectly to a CD38BP, such as an anti-CD38 antibody of the invention. The radioisotopes ^{125}I , ^{131}I , ^{99}Tc , ^{186}Re , and ^{188}Re may be, for example, covalently bound to proteins (including antibodies) through amino acid functional groups. For radioactive iodine it is usually through the phenolic group found on tyrosine. There are numerous methods to accomplish this: chloramine-T (see for instance Greenwood et al., Biochem J. 89, 114-123 (1963) and Iodogen (Salacinski et al., Anal. Biochem. 117, 136-146 (1981)). Tc and Re isotopes may be covalently bound through the sulfhydryl group of cysteine (see for instance Griffiths et al., Cancer Res. 51, 4594-4602 (1991)). However, such compositions may be relatively better suited for diagnostic purposes as the body often can break these covalent bonds, releasing the radioisotopes to the circulatory system.

[0314] A CD38BP, such as an anti-CD38 antibody of the invention, may also be labeled with enzymes that are useful

for detection, such as horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase, glucose oxidase, and the like. A CD38BP, such as an anti-CD38 antibody of the invention, may also be labeled with biotin, and accordingly detected through indirect measurement of avidin or streptavidin binding. A CD38BP, such as an anti-CD38 antibody of the invention, may also be labeled with a predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.). Additional examples of enzyme conjugate candidates include malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, α -glycerophosphate dehydrogenase, triose phosphate isomerase, asparaginase, glucose oxidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholinesterase.

[0315] Additional exemplary labeling moieties generally include, but are not limited to spin-labeled molecules and other labeling moieties of diagnostic value.

[0316] In one embodiment, the present invention provides crosslinked CD38 antibody derivatives. For example, such a CD38 antibody derivative may be produced by crosslinking two or more antibodies, at least one of which is an antibody of the invention (of the same type or of different types, e.g., to create bispecific antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (e.g., m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (e.g., disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

[0317] CD38 antibodies may also be conjugated with any suitable type of chemical group, such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These and other suitable conjugated groups may be used to improve the biological characteristics of the antibody, e.g., to increase serum half-life, solubility, and/or tissue binding.

[0318] CD38 antibody derivatives may be produced by chemically conjugating a radioisotope, protein, or other agent/moiety/compound to (a) the N-terminal side or C-terminal side of the antibody or subunit thereof (e.g., an anti-CD38 antibody H chain, L chain, or anti-CD38 specific/selective fragment thereof) an appropriate substituent group or side chain or (b) a sugar chain in the CD38BP (see, e.g., Antibody Engineering Handbook, edited by Osamu Kanemitsu, published by Chijin Shokan (1994)). Derivatives may also be generated by conjugation at internal residues or sugars, where appropriate.

[0319] CD38 antibodies may also be derivatized with a detection agents, for instance fluorescent compounds, including fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, lanthanide phosphors, and the like. Additional examples of suitable fluorescent labels include a ^{125}Eu label, an isothiocyanate label, a phycoerythrin label, a phycocyanin label, an allophycocyanin label, an o-phthaldehyde label, a fluorescamine label, etc. Examples of chemiluminescent labels include luminal labels, isoluminal labels, aromatic acridinium ester labels, imidazole labels, acridinium salt labels, oxalate ester labels, a luciferin labels, luciferase labels, aequorin labels, etc.

[0320] In one embodiment, a CD38 antibody derivative comprises a conjugated nucleic acid or nucleic acid-associated molecule. In one such facet of the invention the conjugated nucleic acid is a cytotoxic ribonuclease. In one embodiment, the conjugated nucleic acid is an antisense nucleic acid (for instance a S100A10 targeted antisense molecule, which may also be an independent component in a combination composition or combination administration method as described herein - see for instance Zhang et al., J Biol Chem. 279(3), 2053-62 (2004)). In one embodiment, the conjugated nucleic acid is an inhibitory RNA molecule (e.g., a siRNA molecule). In one embodiment, the conjugated nucleic acid is an immunostimulatory nucleic acid (e.g., an immunostimulatory CpG motif-containing DNA molecule). In one embodiment, the conjugated nucleic acid is an expression cassette coding for expression of a tumor suppressor gene, anti-cancer vaccine, anti-cancer cytokine, or apoptotic agent. Such derivatives also may comprise conjugation of a nucleic acid coding for expression of one or more cytotoxic proteins, such as plant and bacterial toxins.

[0321] In one embodiment, the antibody is conjugated to a functional nucleic acid molecule. Functional nucleic acids include antisense molecules, interfering nucleic acid molecules (e.g., siRNA molecules), aptamers, ribozymes, triplex forming molecules, and external guide sequences. The functional nucleic acid molecules may act as effectors, inhibitors, modulators, and stimulators of a specific activity possessed by a target molecule, or the functional nucleic acid molecules may possess a de novo activity independent of any other molecules. A representative sample of methods and techniques which aid in the design and use of antisense molecules can be found in the following non-limiting list of US patents: US 5,135,917, US 5,294,533, US 5,627,158, US 5,641,754, US 5,691,317, US 5,780,607, US 5,786,138, US 5,849,903, US 5,856,103, US 5,919,772, US 5,955,590, US 5,990,088, US 5,994,320, US 5,998,602, US 6,005,095, US 6,007,995, US 6,013,522, US 6,017,898, US 6,018,042, US 6,025,198, US 6,033,910, US 6,040,296, US 6,046,004, US 6,046,319 and US 6,057,437.

[0322] In one embodiment, the antibody is conjugated to an aptamer. Aptamers are molecules that interact with a target molecule, for instance in a specific way. Typically aptamers are small nucleic acids ranging from 15-50 bases in length that fold into defined secondary and tertiary structures, such as stem-loops or G-quartets. Aptamers can bind small molecules, such as ATP (US 5,631,146) and theophylline (US 5,580,737), as well as large molecules, such as reverse transcriptase (US 5,786,462) and thrombin (US 5,543,293). Representative examples of how to make and use aptamers to bind a variety of different target molecules can be found in the following non-limiting list of US patents: US

5,476,766, US 5,503,978, US 5,631,146, US 5,731,424, US 5,780,228, US 5,792,613, US 5,795,721, US 5,846,713, US 5,858,660, US 5,861,254, US 5,864,026, US 5,869,641, US 5,958,691, US 6,001,988, US 6,011,020, US 6,013,443, US 6,020,130, US 6,028,186, US 6,030,776 and US 6,051,698.

[0323] In one embodiment, the present invention provides a CD38 antibody which is conjugated to a ribozyme. Ribozymes are nucleic acid molecules that are capable of catalyzing a chemical reaction, either intramolecularly or intermolecularly. Ribozymes are thus catalytic nucleic acids. There are a number of different types of ribozymes that catalyze nuclease or nucleic acid polymerase type reactions which are based on ribozymes found in natural systems, such as (a) hammerhead ribozymes, (described in for example US 5,334,711, US 5,436,330, US 5,616,466, US 5,633,133, US 5,646,020, US 5,652,094, US 5,712,384, US 5,770,715, US 5,856,463, US 5,861,288, US 5,891,683, US 5,891,684, US 5,985,621, US 5,989,908, US 5,998,193, US 5,998,203, WO 9858058, WO 9858057 and WO 9718312), (b) hairpin ribozymes (described in for instance US 5,631,115, US 5,646,031, US 5,683,902, US 5,712,384, US 5,856,188, US 5,866,701, US 5,869,339 and US 6,022,962), and (c) tetrahymena ribozymes (described in for instance US 5,595,873 and US 5,652,107). There are also a number of ribozymes that are not found in natural systems, but which have been engineered to catalyze specific reactions de novo (examples of which are described in for instance US 5,580,967, US 5,688,670, US 5,807,718 and US 5,910,408). Ribozymes typically cleave RNA or DNA substrates, and more commonly cleave RNA substrates. Ribozymes typically cleave nucleic acid substrates through recognition and binding of the target substrate with subsequent cleavage. This recognition is often based mostly on canonical or non-canonical base pair interactions. This property makes ribozymes particularly good candidates for target specific cleavage of nucleic acids because recognition of the target substrate is based on the target substrates sequence. Representative examples of how to make and use ribozymes to catalyze a variety of different reactions can be found in the following non-limiting list of US patents: US 5,646,042, US 5,693,535, US 5,731,295, US 5,811,300, US 5,837,855, US 5,869,253, US 5,877,021, US 5,877,022, US 5,972,699, US 5,972,704, US 5,989,906 and US 6,017,756.

[0324] In one embodiment, the present invention provides a CD38 antibody that is conjugated to a triplex forming function nucleic acid. Such nucleic acid molecules can interact with either double-stranded or single-stranded nucleic acid. When triplex molecules interact with a target region, a structure called a triplex is formed, in which three strands of DNA form a complex dependent on both Watson-Crick and Hoogsteen base-pairing. Triplex molecules can bind target regions with high affinity and specificity. Representative examples of how to make and use triplex forming molecules to bind a variety of different target molecules can be found in the following non-limiting list of US patents: US 5,176,996, US 5,645,985, US 5,650,316, US 5,683,874, US 5,693,773, US 5,834,185, US 5,869,246, US 5,874,566 and US 5,962,426.

[0325] In one embodiment, a CD38 antibody is conjugated to an external guide sequence. External guide sequences (EGSs) are molecules that bind a target nucleic acid molecule forming a complex that is recognized by RNase P, which cleaves the target molecule. EGSs may be designed to specifically target a RNA molecule of choice. RNase P aids in processing transfer RNA (tRNA) within a cell. Bacterial RNase P can be recruited to cleave virtually any RNA sequence by using an EGS that causes the target RNA:EGS complex to mimic the natural tRNA substrate. (see for instance WO 92/03566 and Forster and Altman, Science 238, 407-409 (1990) for discussion). Representative examples of how to make and use EGS molecules to facilitate cleavage of a variety of different target molecules are provided in the following non-limiting list of US patents: US 5,168,053, US 5,624,824, US 5,683,873, US 5,728,521, US 5,869,248 and US 5,877,162.

[0326] In one embodiment, a CD38 antibody is conjugated to an interfering nucleic acid molecule, such as a siRNA or other RNAi molecule (e.g., an inhibitory double stranded (ds) RNA molecule of about 20-25 nucleotides), which is targeted to interfere with the action of a target gene expression product, such as a gene expression product involved in a CD38 mediated disease or condition. Methods for the production and use of interfering nucleic acid molecules are provided in for instance Nishikura, Cell. 107(4), 415-8 (2001), Fjose et al., Biotechnol Annu Rev. 7, 31-57 (2001), Hanon, Nature 418, 244-51 (2002), Brantl, Biochim Biophys Acta. 1575(1-3), 15-25 (2002), Tuschl, Chembiochem. 2(4), 239-45 (2001), Caplen, Expert Opin Biol Ther. 3(4), 575-86 (2003), Lu et al., Curr Opin Mol Ther. 5(3), 225-34 (2003), Shuey et al., Drug Discov Today. 7(20), 1040-6 (2002), Shi, Trends Genet. 19(1), 9-12 (2003), Kovar et al., Semin Cancer Biol. 13(4), 275-81 (2003), Lavrey et al., Curr Opin Drug Discov Devel. 6(4), 561-9 (2003), Clewey, Commun Dis Public Health. 6(2), 162-3 (2003), Duxbury et al., J Surg Res. 117(2), 339-44 (2004), Caplen et al., Ann N Y Acad Sci. 1002, 56-62 (2003), WO 01/75164, US 6,506,559, US 20040086884, US 20040077574, US 20040063654, US 20040033602, US 20030167490, US 20030157030, US 20030114409, US 20030108923, US 20040014113 and US 20020132788.

[0327] In one embodiment, a CD38 antibody is conjugated to a tumor targeting domain peptide or molecule. In one embodiment, a CD38 antibody is conjugated to a tumor targeting factor VII sequence. Any method known in the art for conjugating the CD38BPantibody to the conjugated molecule(s), such as those described above, may be employed, including those methods described by Hunter et al., Nature 144, 945 (1962), David et al., Biochemistry 13, 1014 (1974), Pain et al., J. Immunol. Meth. 40, 219 (1981) and Nygren, J. Histochem. and Cytochem. 30, 407 (1982). Linkage/conjugation may be accomplished in any suitable way. For example, a covalent linkage may take the form of a disulfide bond (if necessary and suitable, a CD38BP, such as an anti-CD38 antibody of the invention, could be engineered to

contain an extra cysteine codon, which desirably does not interfere with the CD38 binding activity of the molecule. A toxin molecule, derivatized with a sulfhydryl group reactive with the cysteine of the modified CD38BP, may form an immunoconjugate with the CD38BP. Alternatively, a sulfhydryl group may be introduced directly to a CD38BP, such as an anti-CD38 antibody of the invention, using solid phase polypeptide techniques. For example, the introduction of sulfhydryl groups into peptides is described by Hiskey, Peptides 3, 137 (1981). The introduction of sulfhydryl groups into proteins is described in Maasen et al., Eur. J. Biochem. 134, 32 (1983). Once the correct sulfhydryl groups are present, the cytotoxin and CD38BP may be purified, both sulfur groups reduced; cytotoxin and ligand mixed (for instance in a ratio of about 1:5 to 1:20); and disulfide bond formation allowed to proceed to completion (generally about 20 to 30 minutes) at room temperature. The mixture may then be dialyzed against phosphate buffered saline or chromatographed in a resin such as Sephadex to remove unreacted ligand and toxin molecules.

[0328] Numerous types of cytotoxic compounds may be joined to proteins through the use of a reactive group on the cytotoxic compound or through the use of a cross-linking agent. A common reactive group that will form a stable covalent bond *in vivo* with an amine is isothiocyanate (Means et al., Chemical modifications of proteins (Holden-Day, San Francisco 1971) pp. 105-110). This group preferentially reacts with the ϵ -amine group of lysine. Maleimide is a commonly used reactive group to form a stable *in vivo* covalent bond with the sulfhydryl group on cysteine (Ji., Methods Enzymol 91, 580-609 (1983)). Monoclonal antibodies typically are incapable of forming covalent bonds with radiometal ions, but they may be attached to the antibody indirectly through the use of chelating agents that are covalently linked to the antibodies. Chelating agents may be attached through amines (Meares et al., Anal. Biochem. 142, 68-78 (1984)) and sulfhydryl groups (Koyama, Chem. Abstr. 120, 217262t (1994)) of amino acid residues and also through carbohydrate groups (Rodwell et al., PNAS USA 83, 2632-2636 (1986), Quadri et al., Nucl. Med. Biol. 20, 559-570 (1993)). Since these chelating agents contain two types of functional groups, one to bind metal ions and the other to joining the chelate to the antibody, they are commonly referred as bifunctional chelating agents (Sundberg et al., Nature 250, 587-588 (1974)).

[0329] Crosslinking agents that have two reactive functional groups are classified as being homo or heterobifunctional. Examples of homobifunctional crosslinking agents include bismaleimido-hexane (BMH) which is reactive with sulfhydryl groups (Chen et al., J Biol Chem 266, 18237-18243 (1991)) and ethylene glycolbis[succinimidylsuccinate] (EGS) which is reactive with amino groups (Browning et al., J. Immunol. 143, 1859-1867 (1989)). An example of a heterobifunctional crosslinker is m-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) (Myers et al., J. Immunol. Meth. 121, 129-142 (1989)). These methodologies are simple and are commonly employed.

[0330] A therapeutic or diagnostic agent may also or alternatively be attached at the hinge region of a reduced antibody component via disulfide bond formation. As an alternative, such peptides may be attached to the antibody component using a heterobifunctional cross-linker, such as N-succinyl 3-(2-pyridyldithio)propionate (SPDP). Yu et al., Int. J. Cancer 56, 244 (1994). General techniques for such conjugation are well known in the art. See, for example, Wong, Chemistry Of Protein Conjugation And Cross-Linking (CRC Press 1991), Upešlaciš et al., "Modification of Antibodies by Chemical Methods," In Monoclonal Antibodies: Principles And Applications, Birch et al., (eds.) (Wiley-Liss, Inc. 1995), Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in Monoclonal Antibodies: Production, Engineering And Clinical Application, Ritter et al., (eds.) (Cambridge University Press 1995).

[0331] In some embodiments, labels or other conjugated substituents are attached to the amino acid sequence of the anti-CD38 antibody by spacer arms of various lengths to reduce potential steric hindrance.

[0332] Unlabeled CD38BP(s), such as antibodies of the invention, may be used in combination with other labeled antibodies (second antibodies) that are reactive with the CD38BP(s), such as antibodies specific for human immunoglobulin constant regions that bind to anti-CD38 mAbs. Alternatively, a CD38BP, such as an antibody of the invention, may be directly labeled. A wide variety of labels may be employed for direct or indirect labeling of CD38BPs, such as labeling with radionuclides, fluors, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, ligands (particularly haptens), etc.

[0333] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue or the antibody fused to an epitope tag. Other insertion variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody of an enzyme or a polypeptide or PEG which increases the serum half-life of the antibody. Such anti-CD38 antibody fusion proteins comprising an antibody of the invention are another feature of the present invention.

[0334] In one embodiment, the present invention provides molecules comprising a CD38 antibody, such as a human anti-CD38 antibody, of the present invention conjugated to a therapeutic moiety, such as a cytotoxin, a chemotherapeutic drug, an immunosuppressant, or a radioisotope. Such conjugates are referred to herein as "immunoconjugates". Immunoconjugates which include one or more cytotoxins are referred to as "immunotoxins".

[0335] A cytotoxin or cytotoxic agent includes any agent that is detrimental to (e.g., kills) cells. For a description of these classes of drugs which are well known in the art, and their mechanisms of action, see Goodman et al., Goodman and Gilman's The Pharmacological Basis Of Therapeutics, 8th Ed., Macmillan Publishing Co., 1990. Additional techniques relevant to the preparation of antibody immunotoxins are provided in for instance Vitetta, Immunol. Today 14, 252 (1993)

and US 5,194,594.

[0336] Suitable therapeutic agents for forming immunoconjugates of the present invention include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, actinomycin D, 1-dehydro-testosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin, antimetabolites (such as methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, fludarabine, 5-fluorouracil, decarbazine, hydroxyurea, asparaginase, gemcitabine, cladribine), alkylating agents (such as mechlorethamine, thioepa, chlorambucil, melphalan, carmustine (BSNU), lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, dacarbazine (DTIC), procarbazine, mitomycin C, cisplatin and other platinum derivatives, such as carboplatin), antibiotics (such as dactinomycin (formerly actinomycin), bleomycin, daunorubicin (formerly daunomycin), doxorubicin, idarubicin, mithramycin, calicheamicin, mitomycin, mitoxantrone, plitacmycin, anthramycin (AMC)), diphtheria toxin and related molecules (such as diphtheria A chain and active fragments thereof and hybrid molecules), ricin toxin (such as ricin A or a deglycosylated ricin A chain toxin), cholera toxin, a Shiga-like toxin (SLT-I, SLT-II, SLT-III), LT toxin, C3 toxin, Shiga toxin, pertussis toxin, tetanus toxin, soybean Bowman-Birk protease inhibitor, *Pseudomonas* exotoxin, alorin, saporin, modeccin, gelanin, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, *sapaonaria officinalis* inhibitor, gelonin, mitogellin, restrictocin, phenomycin, and enomycin toxins. Therapeutic agents, which may be administered in combination with an anti-CD38 antibody of the present invention as described elsewhere herein, may also be candidates for therapeutic moieties useful for conjugation to an antibody of the present invention. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an enzymatically active toxin, or active fragment thereof, such as abrin, ricin A, *pseudomonas* exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor or interferon- γ ; or, biological response modifiers such as, for example, lymphokines, interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), or other growth factors and apoptotic inducing protein isolated from mitochondria.

[0337] In one embodiment, the cytotoxic agent is calicheamicin, ^{90}Y , ^{125}I and ^{131}I .

[0338] Other examples of therapeutic cytotoxins that may be conjugated to a CD38 antibody of the present invention include calicheamicins and duocarmycins. As indicated above, the drug moiety need not be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an agent active at the cell surface, such as phospholipase enzymes, e.g. phospholipase C.

[0339] The lysing portion of a toxin typically may be readily joined to the Fab fragment of an antibody or antibody fragment of the present invention. Other suitable conjugated molecules include ribonuclease (RNase), DNase I, *Staphylococcal* enterotoxin-A, pokeweed antiviral protein, diphtherin toxin, and *Pseudomonas* endotoxin. See, for example, Pastan et al., *Cell* 47, 641 (1986) and Goldenberg, *Calif. A Cancer Journal for Clinicians* 44, 43 (1994). Additional toxins suitable for use in the present invention are known to those of skill in the art (see for instance US 6,077,499).

[0340] Conjugates of CD38BPs, such as antibodies of the invention, and such cytotoxic moieties may be made using a variety of bifunctional protein coupling agents. Examples of such reagents include SPDP, IT, bifunctional derivatives of imidoesters such as dimethyl adipimidate HCl, active esters such as disuccinimidyl suberate, aldehydes such as glutaraldehyde, bis-azido compounds such as bis (p-azidobenzoyl) hexanediamine, bis-diazonium derivatives such as bis-(p-diazoniumbenzoyl)-ethylenediamine, diisocyanates such as tolylene 2,6-diisocyanate, and bis-active fluorine compounds such as 1,5-difluoro-2,4-dinitrobenzene and anti-mitotic agents (e.g., vincristine, vinblastine, docetaxel, paclitaxel and vinorelbine).

[0341] Techniques for conjugating such therapeutic moieties to CD38BPs, such as antibodies, are well known, see for instance Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al., (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985), Hellstrom et al., "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson et al., (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987), Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al., (eds.), pp. 475-506 (1985), "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al., (eds.), pp. 303-16 (Academic Press 1985) and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62, 119-58 (1982).

[0342] In one embodiment, the present invention provides a CD38 antibody that is conjugated to a mixed toxin. A mixed toxin molecule is a molecule derived from two different (typically polypeptide) toxins. Generally, peptide toxins comprise one or more domains responsible for generalized eukaryotic cell binding, at least one enzymatically active domain, and at least one translocation domain. The binding and translocation domains are required for cell recognition and toxin entry respectively. Naturally-occurring proteins which are known to have a translocation domain include diphtheria toxin, *Pseudomonas* exotoxin A, and possibly other peptide toxins. The translocation domains of diphtheria toxin and *Pseudomonas* exotoxin A are well characterized (see for instance Hoch et al., *PNAS USA* 82, 1692 (1985), Colombatti

et al., J. Biol. Chem. 261, 3030 (1986) and Deleers et al., FEBS Lett. 160, 82 (1983)), and the existence and location of such a domain in other molecules may be determined by methods such as those employed by Hwang et al., Cell 48, 129 (1987) and Gray et al., PNAS USA 81 2645 (1984). In view of these techniques, a useful mixed toxin hybrid molecule may be formed, for example, by fusing the enzymatically active A subunit of E. coli Shiga-like toxin (Calderwood et al., PNAS USA 84, 4364 (1987)) to the translocation domain (amino acid residues 202 through 460) of diphtheria toxin, and to a molecule targeting a particular cell type, as described in US 5,906,820. The targeting portion of the three-part hybrid can cause the molecule to attach specifically to the targeted cells, and the diphtheria toxin translocation portion can act to insert the enzymatically active A subunit of the Shiga-like toxin into a targeted cell. The enzymatically active portion of Shiga-like toxin, like diphtheria toxin, acts on the protein synthesis machinery of the cell to prevent protein synthesis, thus killing the targeted cell.

[0343] Immunoconjugates according to the present invention may also comprise a radioisotope, e.g., iodine-131, yttrium-90 or indium-111, to generate cytotoxic radiopharmaceuticals for treating a CD38-related disorder, such as multiple myeloma.

[0344] In one embodiment, the CD38 antibodies, such as the human antibodies of the present invention are attached to a linker-chelator, e.g., tiuxetan, which allows for the antibody to be conjugated to a radioisotope.

[0345] Additionally useful conjugate substituents include anti-cancer retinoids. Taxane conjugates (see for instance Jaime et al., Anticancer Res. 21(2A), 1119-28 (2001), cisplatin conjugates, thapsigargin conjugates, linoleic acid conjugates, calicheamicin conjugates (see for instance Damle et al., Curr Opin Pharmacol. 3(4), 386-90 (2003), doxorubicin conjugates, geldanamycin conjugates, and the like, also may be useful in promoting the treatment of cancer (see, generally, Trail et al., Cancer Immunol Immunother. 52(5), 328-37 (2003)).

[0346] In one embodiment, the present invention provides anti-idiotypic antibodies raised against anti-CD38 antibodies of the present invention. Secondary antibodies raised against anti-CD38 antibodies of the present invention are also described. A secondary antibody refers to an antibody specific for, and typically raised against, an anti-CD38 antibody. An anti-idiotypic (Id) antibody is an antibody which recognizes unique determinants generally associated with the antigen-binding site of an antibody. An Id antibody may be prepared by immunizing an animal of the same species and genetic type as the source of an anti-CD38 mAb with the mAb to which an anti-Id is being prepared. The immunized animal typically can recognize and respond to the idiotype determinants of the immunizing antibody by producing an antibody to these idiotype determinants (the anti-Id antibody). Such antibodies are described in for instance US 4,699,880. Such antibodies are further features of the present invention.

[0347] An anti-Id antibody may also be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody. An anti-anti-Id may be epitopically identical to the original mAb, which induced the anti-Id. Thus, by using antibodies to the idiotype determinants of a mAb, it is possible to identify other clones expressing antibodies of identical specificity. Anti-Id antibodies may be varied (thereby producing anti-Id antibody variants) and/or derivatized by any suitable technique, such as those described elsewhere herein with respect to anti-CD38 antibodies and other CD38BPs of the present disclosure. For example, anti-Id mAbs may be coupled to a carrier such as keyhole limpet hemocyanin (KLH) and used to immunize BALB/c mice. Sera from these mice typically will contain anti-anti-Id antibodies that have the binding properties similar if not identical to an original/parent CD38 antibody.

[0348] In one embodiment, the present invention provides a nucleic acid encoding a CD38 antibody of the invention. A CD38 antibody-encoding nucleic acid of the invention may have any suitable characteristics and comprise any suitable features or combination thereof. Thus, for example, a CD38 antibody-encoding nucleic acid may be in the form of DNA, RNA, or a hybrid thereof, and may include nonnaturally-occurring bases, a modified backbone (e.g., a phosphothioate backbone that promotes stability of the nucleic acid), or both. The nucleic acid advantageously comprises features that promote desired expression in target host cell(s), replication, and/or selection. Examples of such features include an origin of replication component, a selection gene component, a promoter component, an enhancer element component, a polyadenylation sequence component, a termination component, and the like.

[0349] In one embodiment, the present invention provides a vector comprising an anti-CD38 antibody -encoding nucleic acid of the invention. A vector refers to a delivery vehicle that promotes the expression of a CD38 antibody-encoding nucleic acid, the production of a CD38 antibody, the transfection/transformation of target cells, the replication of the CD38 antibody-encoding nucleic acid, promotes stability of the nucleic acid, promotes detection of the nucleic acid and/or transformed/transfected cells, or otherwise imparts advantageous biological function to the CD38 antibody-encoding nucleic acid. A vector in the context of the present invention may be any suitable vector, including chromosomal, non-chromosomal, and synthetic nucleic acid vectors (a nucleic acid sequence comprising a suitable set of expression control elements). Examples of such vectors include derivatives of SV40, bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, and viral nucleic acid (RNA or DNA) vectors. In one embodiment, a CD38BP-encoding nucleic acid is comprised in a naked DNA or RNA vector, including, for example, a linear expression element (as described in for instance Sykes and Johnston, Nat Biotech 17, 355-59 (1997)), a compacted nucleic acid vector (as described in for instance US 6,077, 835 and/or WO 00/70087), a plasmid vector such as pBR322, pUC 19/18, or pUC 118/119, a "midge" minimally-sized nucleic acid vector (as described in for instance

Schakowski et al., *Mol Ther* 3, 793-800 (2001)), or as a precipitated nucleic acid vector construct, such as a CaP04-precipitated construct (as described in for instance WO 00/46147, Benvenisty and Reshef, *PNAS USA* 83, 9551-55 (1986), Wigler et al., *Cell* 14, 725 (1978), and Coraro and Pearson, *Somatic Cell Genetics* 7, 603 (1981)). Such nucleic acid vectors and the usage thereof are well known in the art (see for instance US 5,589,466 and US 5,973,972).

[0350] In one embodiment, the vector is suitable for expression of the CD38 antibody in a bacterial cell. Examples of such vectors include, for example, vectors which direct high level expression of fusion proteins that are readily purified (for instance multifunctional *E. coli* cloning and expression vectors such as BlueScript (Stratagene), pIN vectors (Van Heeke & Schuster, *J Biol Chem* 264, 5503-5509 (1989), pET vectors (Novagen, Madison WI) and the like).

[0351] An expression vector may also or alternatively be a vector suitable for expression in a yeast system. Any vector suitable for expression in a yeast system may be employed. Suitable vectors for use in for instance *Saccharomyces cerevisiae* include, for example, vectors comprising constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH (reviewed in: F. Ausubel et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley InterScience New York (1987), and Grant et al., *Methods in Enzymol* 153, 516-544 (1987)).

[0352] A nucleic acid and/or vector may also comprises a nucleic acid sequence encoding a secretion/ localization sequence, which can target a polypeptide, such as a nascent polypeptide chain, to a desired cellular compartment, membrane, or organelle, or which directs polypeptide secretion to periplasmic space or into cell culture media. Such sequences are known in the art, and include secretion leader or signal peptides, organelle targeting sequences (e. g., nuclear localization sequences, ER retention signals, mitochondrial transit sequences, chloroplast transit sequences), membrane localization/anchor sequences (e. g., stop transfer sequences, GPI anchor sequences), and the like.

[0353] CD38BP antibody-encoding nucleic acids may comprise or be associated with any suitable promoter, enhancer, and other expression-facilitating elements. Examples of such elements include strong expression promoters (e. g., human CMV IE promoter/enhancer as well as RSV, SV40, SL3-3, MMTV, and HIV LTR promoters), effective poly (A) termination sequences, an origin of replication for plasmid product in *E. coli*, an antibiotic resistance gene as selectable marker, and/or a convenient cloning site (e.g., a polylinker). Nucleic acids may also comprise an inducible promoter as opposed to a constitutive promoter such as CMV IE (the skilled artisan will recognize that such terms are actually descriptors of a degree of gene expression under certain conditions).

[0354] In one embodiment, the nucleic acid may be positioned in and/or delivered to the host cell or host animal via a viral vector. Any suitable viral vector may be used in this respect, and several are known in the art. A viral vector may comprise any number of viral polynucleotides, alone or in combination with one or more viral proteins, which facilitate delivery, replication, and/or expression of the nucleic acid of the present invention in a desired host cell. The viral vector may be a polynucleotide comprising all or part of a viral genome, a viral protein/nucleic acid conjugate, a virus-like particle (VLP), a vector similar to those described in US 5,849, 586 and WO 97/04748, or an intact virus particle comprising viral nucleic acids and the nucleic acid of the present invention. A viral particle viral vector may comprise a wild-type viral particle or a modified viral particle. The viral vector may be a vector which requires the presence of another vector or wild-type virus for replication and/or expression (i.e., it may be a helper-dependent virus), such as an adenoviral vector amplicon. Typically, such viral vectors consist essentially of a wild-type viral particle, or a viral particle modified in its protein and/or nucleic acid content to increase transgene capacity or aid in transfection and/or expression of the nucleic acid (examples of such vectors include the herpes virus/AAV amplicons). Typically, a viral vector is similar to and/or derived from a virus that normally infects humans. Suitable viral vector particles in this respect, include, for example, adenoviral vector particles (including any virus of or derived from a virus of the adenoviridae), adeno-associated viral vector particles (AAV vector particles) or other parvoviruses and parvoviral vector particles, papillomaviral vector particles, flaviviral vectors, alphaviral vectors, herpes viral vectors, pox virus vectors, retroviral vectors, including lentiviral vectors. Examples of such viruses and viral vectors are in for instance Fields et al., eds., *Virology* Raven Press, Ltd., New York (3rd ed., 1996 and 4th ed., 2001), *Encyclopedia of Virology*, R. G. Webster et al., eds., Academic Press (2nd ed., 1999), *Fundamental Virology*, Fields et al., eds., Lippincott-Raven (3rd ed., 1995), Levine, "Viruses," *Scientific American Library* No. 37 (1992), *Medical Virology*, D. O. White et al., eds., Acad. Press (2nd ed. 1994), and *Introduction to Modern Virology*, Dimock, N. J. et al., eds., Blackwell Scientific Publications, Ltd. (1994).

[0355] Viral vectors that may be employed with polynucleotides of the present invention and the methods described herein include adenovirus and adeno-associated vectors, as in for instance Carter, *Curr Opin Biotech* 3, 533-539 (1992) and Muzyczka, *Curr Top Microbiol Immunol* 158, 97-129 (1992). Additional types and aspects of AAV vectors are described in for instance Carter, *Contrib. Microbiol.* 4, 85-86 (2000), Smith-Arica, *Curr. Cardiol. Rep.* 3(1), 41-49 (2001), Taj, *J. Biomed. Sci.* 7(4), 279- 91 (2000), Vigna et al., *J. Gene Med.* 2(5), 308-16 (2000), Klimatcheva et al., *Front. Biosci.* 4, D481-96 (1999), Lever et al., *Biochem. Soc. Trans.* 27(6), 841-47 (1999), Snyder, *J Gene Med.* 1(3), 166-75 (1999), Gerich et al., *Knee Surg. Sports Traumatol. Arthrosc.* 5(2), 118-23 (1998), and During, *Adv. Drug Deliv. Review* 27(1), 83-94 (1997) and US 4,797,368, US 5,139,941, US 5,173,414, US 5,614,404, US 5,658,785, US 5,858,775 and US 5,994,136). Adeno-associated viral vectors may be constructed and/or purified using the methods set forth, for example, in US 4,797,368 and Laughlin et al., *Gene* 23, 65-73 (1983).

[0356] Another type of viral vector that may be employed with polynucleotides and methods of the present invention

is a papillomaviral vector. Suitable papillomaviral vectors are known in the art and described in, e. g., Hewson, *Mol Med Today* 5(1), 8 (1999), Stephens, *Biochem J.* 248(1), 1-11 (1987) and US 5,719, 054. Examples of papilloma viral vectors are provided in for instance WO 99/21979. Alphavirus vectors may be gene delivery vectors in other contexts. Alphavirus vectors are known in the art and described in for instance Carter, *Curr Opin Biotech* 3, 533-539 (1992), Muzyczka, *Curr Top Microbiol Immunol.* 158, 97-129 (1992), Schlesinger, *Expert Opin Biol Ther.* 1(2), 177-91 (2001), Polo et al., *Dev Biol (Basel).* 104, 181-5 (2000), Wahlfors et al., *Gene Ther.* 7(6), 472-80 (2000), Colombage et al., *Virology.* 250(1), 151-63 (1998) and WO 01/81609, WO 00/39318, WO 01/81553, WO 95/07994 and WO 92/10578.

[0357] Another group of viral vectors are herpes viral vectors. Examples of herpes viral vectors are described in for instance Lachmann et al., *Curr Opin Mol Ther* 1(5), 622-32 (1999), Fraefel et al., *Adv Virus Res.* 55, 425-51 (2000), Huard et al., *Neuromuscul* 7(5), 299-313 (1997), Glorioso et al., *Annu Rev Microbiol.* 49, 675-710 (1995), Latchman, *Mol Biotechnol.* 2(2), 179-95 (1994), and Frenkel et al., *Gene Ther.* 1(Suppl 1), S40-6 (1994), as well as US 6,261,552 and US 5,599,691.

[0358] Retroviral vectors, including lentiviral vectors, also may be advantageous gene delivery vehicles in particular contexts. There are numerous retroviral vectors known in the art. Examples of retroviral vectors are described in for instance Miller, *Curr Top Microbiol Immunol* 158, 1-24 (1992), Salmons and Gunzburg, *Human Gene Therapy* 4, 129-141 (1993), Miller et al., *Methods in Enzymology* 217, 581-599 (1994), Weber et al., *Curr Opin Mol Ther.* 3(5), 439-53 (2001), Hu et al., *Pharmacol Rev.* 52(4), 493-511 (2000), Kim et al., *Adv Virus Res.* 55, 545-63 (2000), Palu et al., *Rev Med Virol.* 10(3), 185-202 (2000) and Takeuchi et al., *Adv Exp Med Biol.* 465, 23-35 (2000), as well as US 6,326,195, US 5,888,502, US 5,580,766, and US 5,672, 510.

[0359] Adenoviral vectors may also be suitable viral vectors for gene transfer. Adenoviral vectors are well known in the art and described in for instance Graham et al, *Mol Biotechnol* 3(3), 207-220 (1995), Stephenson, *Clin Diagn Virol* 10(2-3), 187-94 (1998), Jacobs, *Clin Sci (Lond).* 85(2), 117-22 (1993), US 5,922, 576, US 5,965,358 and US 6,168, 941 and WO98/22588, WO98/56937, WO99/15686, WO99/54441, and WO00/32754. Adenoviral vectors, herpes viral vectors and Sindbis viral vectors, useful in the practice of the present invention, are described in for instance Jolly *Cancer Gene Therapy* 1, 51-64 (1994), Latchman *Molec Biotechnol* 2, 179-195 (1994) and Johanning et al., *Nucl Acids Res* 23, 1495-1501 (1995).

[0360] Other suitable viral vectors include pox viral vectors. Examples of such vectors are discussed in for instance Berencsi et al., *J Infect Dis* 183(8), 1171-9 (2001), Rosenwirth et al., *Vaccine* 19(13-14), 1661-70 (2001), Kittlesen et al., *J Immunol* 164(8), 4204-11 (2000), Brown et al., *Gene Ther* 7(19), 1680-9 (2000), Kanasa-athan et al., *Vaccine* 19(4- 5), 483-91 (2000), Sten, *Drugs* 60(2), 249-71 (2000). Vaccinia virus vectors may be pox virus vectors. Examples of such vectors and uses thereof are provided in for instance Venugopal et al., *Res Vet Sci* 57(2), 188-193 (1994), Moss *Dev Biol Stand* 82, 55-63 (1994), Weisz et al., *Mol Cell Biol* 43, 137-159 (1994), Mahr and Payne, *Immunobiology* 184(2-3), 126-146 (1992), Hruby, *Clin Microbiol Rev* 3(2), 153-170 (1990) and WO92/07944, WO98/13500, and WO89/08716.

[0361] Other features of the present invention include recombinant cells, such as yeast, bacterial, and mammalian cells (e.g., immortalized mammalian cells) comprising a nucleic acid of the invention, a vector of the invention, or combinations of either or both thereof. For example, one embodiment is a cell comprising a nucleic acid stably integrated into the cellular genome that comprises a sequence coding for expression of a CD38 antibody of the present invention. One embodiment is a cell comprising a non-integrated nucleic acid, such as a plasmid, cosmid, phagemid, or linear expression element, which comprises a sequence coding for expression of a CD38 antibody.

[0362] Also described herein are immunogenic peptides comprising any of the above-described antigenic determinant portions of CD38 specific for the CD38BPs as described herein such as the antigenic determinant portions of CD38 specific for -003 and -005 and -024. Such immunogens may be used to elicit a direct immune response in a method comprising an active immunotherapy regimen. The present disclosure further provides a fusion protein comprising such a CD38 immunogen and a fusion partner sequence that improves the half-life of the fusion protein (e.g., by inclusion of an immunoglobulin domain sequence); facilitates detection and/or purification of the fusion protein (by comprising, e.g., a fluorescent peptide sequence, a reporter enzyme sequence, an epitope tag, a hexa-histidine sequence, or the like); promotes the targeting of the fusion protein (e.g., by comprising a ligand or portion of a ligand specific for a receptor on a target cell); promotes induction of a distinct immune response (e.g., corresponds to a cancer antigen or an immunogenic fragment thereof); is a cytotoxic agent; or achieves any combination thereof (e.g., a heat shock fusion protein partner can increase an immune response generated against a non-similar, heterologous antigen portion of a fusion protein, while also increasing the *in vivo* half-life of a fusion protein). Fusion proteins may also comprise one or more cleavage sites, particularly between domains.

[0363] Variants of such peptides, and derivatives of such immunogenic peptides or immunogenic peptide variants are additional features as described herein (e.g., such CD38 immunogenic peptide derivatives may be modified by chemical coupling, genetic fusion, non-covalent association, and the like, to other molecular entities such as antibodies, toxins, radioisotope, cytotoxic agents, or cytostatic agents). Peptide mimetopes, comprising CD38 epitope sequences may also, for example, be useful as vaccine candidates. Such peptides may also be useful in the purification of anti-CD38 antibodies.

In addition to the B-cell epitope sequences described herein, such peptides may be engineered or selected to also or alternatively comprise one or more anti-CD38 T cell epitopes. Such epitopes may be identified by any suitable technique known in the art (e.g., by T cell epitope prediction software applications).

[0364] Described herein is a nucleic acid encoding such an immunogenic peptide. Such a nucleic acid may be delivered to a host in a suitable vector, such as a replication-deficient targeted vector (e.g., a targeted nucleic acid vector or a replication-deficient, targeted adenovirus vector). The present disclosure also provides compositions of one or more of such immunogenic peptides and/or immunogenic peptide-encoding nucleic acids.

[0365] CD38 antibodies of the present invention include "neutralizing" antibodies. The term "neutralizing antibody" refers to a CD38 antibody that is capable of substantially inhibiting or eliminating a biological activity of a CD38-associated peptide. Typically, a neutralizing anti-CD38 antibody, may inhibit, directly or indirectly, the function of CD38, such as enzymatic activity, signal transduction, induction of cytokine expression, induction of proliferation or differentiation, or induction of lysis, in a degree that is about equal or greater than the inhibition of such cells due to administration of an approximately equal amount of -003 or -005 or -024.

[0366] A CD38 antibody of the present invention may have any suitable affinity and/or avidity for one or more epitopes contained at least partially in CD38. Affinity refers to the strength of binding of the CD38 antibody to such an epitope. Typically, affinity is measured by dissociation constant K_d , defined as $[Ab] \times [Ag] / [Ab-Ag]$ where $[Ab-Ag]$ is the molar concentration of the antibody-antigen complex (or the CD38BP-antigen complex), $[Ab]$ is the molar concentration of the unbound antibody (or CD38BP) and $[Ag]$ is the molar concentration of the unbound antigen. The affinity constant K_a is defined by $1/K_d$. Suitable methods for determining specificity and affinity by competitive inhibition can be found in for instance Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988), Colligan et al., eds., *Current Protocols in Immunology*, Greene Publishing Assoc. and Wiley InterScience N.Y., (1992, 1993) and Muller, *Meth. Enzymol.* 92, 589-601 (1983).

[0367] A CD38BP, and particularly anti-CD38 antibodies of the present invention, may have an affinity for at least one epitope at least partially comprised in CD38 in the range of about 10^4 to about 10^{10} M^{-1} . The term immunoreact herein typically refers to binding of a CD38BP to a CD38 epitope with a dissociation constant K_d lower than about 10^{-4} M .

[0368] A CD38BP may have an affinity that is at least as great for CD38 as -003 and -005 and -024, and in some embodiments have an affinity that is at least about as great as -003 and -005 and -024. In particular anti-CD38 antibodies of the present invention may have an affinity that is at least as great for CD38 as -005, and in some embodiments have an affinity that is at least about as great as -005. Affinity may be determined by any of the methods described elsewhere herein or their known equivalents in the art. An example of one method that may be used to determine affinity is provided in Scatchard analysis of Munson & Pollard, *Anal. Biochem.* 107, 220 (1980). Binding affinity also may be determined by equilibrium methods (for instance enzyme-linked immunoabsorbent assay (ELISA) or radioimmunoassay (RIA)) or kinetics analysis (for instance BIACORE™ analysis).

[0369] Typically, the disassociation constant for CD38BPs, such as anti-CD38 antibodies, of the present invention is less than about 100 nM, less than about 50 nM, less than about 10 nM, about 5 nM or less, about 1 nM or less, about 0.5 nM or less, about 0.1 nM or less, about 0.01 nM or less, or even about 0.001 nM or less.

[0370] CD38BPs, such as anti-CD38 antibodies of the present invention, may exhibit similar functional characteristics as -003 and -005 and -024, such as may be determined by antibody-dependent cellular cytotoxicity (ADCC) and complement-mediated cytotoxicity (CDC) assays (see for instance US 5500362).

[0371] In one embodiment, an anti-CD38 antibody according to the present invention does not act as an agonist of CD38, but as an antagonist of CD38. An agonist of CD38 is a molecule, which activates one or more of the functions ascribed to CD38. Such functions may include receptor mediation in adhesion and signaling events and (ecto-) enzymatic activity. Furthermore, as an ectoenzyme, CD38 uses NAD^+ as substrate for the formation of cyclic ADP-ribose (cADPR) and ADPR, but also of nicotinamide and nicotinic acid-adenine dinucleotide phosphate (NAADP). cADPR has been shown to act as second messenger for Ca^{2+} mobilization from the endoplasmatic reticulum. In addition to signaling via Ca^{2+} , CD38 signaling occurs via cross-talk with antigen-receptor complexes on T and B cells or other types of receptor complexes, e.g. MHC molecules, and is in this way involved in several cellular responses, but also in switching and secretion of IgG1.

[0372] In one embodiment, an anti-CD38 antibody according to the present invention does not induce significant proliferation of PBMCs. In one embodiment, an anti-CD38 antibody according to the present invention does not induce release of significant IL-6 levels. In one embodiment, an anti-CD38 antibody according to the present invention does not induce release of detectable IFN- γ levels. Such assays may be measured as described in Ausiello et al., *Tissue antigens* 56, 538-547 (2000).

[0373] Anti-CD38 antibodies of the present invention, as well as other CD38BPs as described herein may be prepared by recombinant expression in any suitable type of cells or animals.

[0374] Recombinant CD38BPs, such as recombinant antibodies, such as recombinant human antibodies, include CD38BPs, such as antibodies, such as human antibodies that are prepared, expressed, created or isolated by recombinant means, such as CD38BPs, such as antibodies, such as human antibodies expressed using a recombinant ex-

pression vector transfected into a host cell.

[0375] Recombinant antibodies, such as recombinant human antibodies also include antibodies isolated from a recombinant, combinatorial human antibody library, antibodies isolated from an animal, such as a transgenic animal, or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin-encoding nucleic acid sequences to other nucleic acid sequences exogenous to the human immunoglobulin-encoding nucleic acids and human immunoglobulin-encoding genes. Recombinant human antibodies typically have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and, thus, the amino acid sequences of the V_H and V_L regions of the recombinant antibodies may be sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*. Both types of human antibodies are provided in accordance with the present invention.

[0376] Suitable methods for recombinant protein production are known in the art, see for instance (Sambrook and Russell (eds.), Molecular cloning, third edition, 2001, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.

[0377] Likewise, suitable methods for antibody production are known in the art and include those described in for instance Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988), Harlow and Lane: Using Antibodies: A Laboratory Manual (Cold Spring Harbor Laboratory Press (1999)), US 4,376,110 and Ausubel et al., eds., Current Protocols In Molecular Biology, Greene Publishing Assoc. and Wiley InterScience N.Y., (1987, 1992). Monoclonal antibodies may be made using the hybridoma method first described by Kohler et al., Nature 256, 495 (1975), or by other well-known, subsequently-developed methods (see, e.g., Goding, Monoclonal Antibodies: Principles and Practice, pp.59-103 (Academic Press, 1986)). Hybridomas useful in the production of anti-CD38 antibodies of the present invention are also provided by the present invention. Such hybridomas may be formed by chemical fusion, electrical fusion, or any other suitable technique, with any suitable type of myeloma, heteromyeloma, phoblastoid cell, plasmacytoma or other equivalent thereof and any suitable type of antibody-expressing cell. Transformed immortalized B cells may also be used to efficiently produce antibodies of the present invention and are also provided in accordance with the present invention. Such cells may be produced by standard techniques, such as transformation with an Epstein Barr Virus, or a transforming gene. (See, e.g., "Continuously Proliferating Human Cell Lines Synthesizing Antibody of Predetermined Specificity," Zurawaki, V. R. et al., in Monoclonal Antibodies, ed. by Kennett R. H. et al., Plenum Press, N.Y. 1980, pp 19-33.). Thus, stable and continuous and/or immortalized anti-CD38 antibody expressing cells and cell lines are a feature in accordance with the present invention. Eukaryotic and prokaryotic cells (e.g., yeast cells, continuous and/or immortalized mammalian cell lines (e.g., lymphoid antibody-producing cell derived cell lines), plant cells, insect cells, and bacterial cells such as E. coli cells, etc.) comprising CD38BP-encoding or CD38BP-fragment-encoding nucleic acids are provided in accordance with the present invention. Transgenic animals, such as non-human primates, rodents (e.g., hamsters, guinea pigs, and rats - including modified strains thereof such as severe combined immunodeficient (SCID) mice and other immunocompromised animal strains), dogs, etc., expressing human anti-CD38 antibodies of the present invention also are provided in accordance with the present invention.

[0378] Recombinant cells comprising exogenous nucleic acids encoding CD38BPs may be prepared by any suitable technique (e.g., transfection/transformation with a naked DNA plasmid vector, viral vector, invasive bacterial cell vector or other whole cell vector, etc., comprising a CD38BP-encoding sequence (or sequences) delivered into the cell by calcium phosphate-precipitation facilitated transfection, receptor-mediated targeting and transfection, biolistic delivery, electroporation, dextran-mediated transfection, liposome-mediated transformation, protoplast fusion, direct microinjection, etc.). Methods of transforming/transfecting cells are well known in the art (see, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press (2d Edition, 1989 and 3rd Edition, 2001) and F. Ausubel et al., ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley InterScience New York (1987). Such recombinant cells which express an antibody of the invention are a feature of the present invention.

[0379] Cell lines available as hosts for recombinant protein expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, inter alia, Chinese hamster ovary (CHO) cells, NSO, SP2 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), A549 cells, and a number of other cell lines. Other cell lines that may be used are insect cell lines, such as Sf9 cells. When nucleic acids (or nucleic acid-containing vectors) encoding proteins, such as CD38BPs (including anti-CD38 antibodies), are introduced into mammalian host cells, proteins, such as CD38BPs, may be produced by culturing the host cells for a period of time sufficient to allow for expression of the protein, such as a CD38BP, in the host cells or by secretion of the protein, such as a CD38BP, into the culture medium in which the host cells are grown. CD38BPs may be recovered from the culture medium using standard protein purification methods. CD38BPs may also be recovered from host cell lysates when directly expressed without a secretory signal.

[0380] CD38BPs, such as anti-CD38 antibodies, may also be produced in bacterial cells and eukaryotic unicellular microorganisms, such as yeast. Bacterial cell produced CD38BPs, such as anti-CD38 antibodies, typically lack normal

glycosylation and bacterial cell produced anti-CD38 antibodies may thus be more or less deficient in terms of ADCC functions and other aspects of the immune response associated with anti-CD38 antibodies produced in mammalian cells and/or animals (e.g., the recruitment of NK cells). Yeast cell produced CD38BPs, such as anti-CD38 antibodies normally exhibit different types of glycosylation patterns than antibodies produced in mammalian cells. However, methods for producing antibodies with effective glycosylation in yeast are currently being developed by companies such as Glycofi, Inc. (Lebanon, NH, USA). See also Wildt S et al., Nat Rev Microbiol. 3(2), 119-28 (2005).

[0381] When recombinant expression vectors encoding CD38BP genes (including anti-CD38 antibody genes) are introduced into mammalian host cells, the CD38BPs are produced by culturing the host cells for a period of time sufficient to allow for expression of the CD38BP in the host cells or for secretion of the antibody into the culture medium in which the host cells are grown. The purification of antibodies and other CD38BPs from cell cultures, cell lysates, and animals (e.g., from the ascites fluid of a transgenic animal producing anti-CD38 antibodies) may be achieved by application of any number of suitable techniques known in the art including, e.g., immunoaffinity column purification; sulfate precipitation; chromatofocusing; preparative SDS-PAGE, and the like.

[0382] Human monoclonal antibodies of the present invention may also be produced by a variety of other techniques, including conventional monoclonal antibody methodology, e.g., the standard somatic cell hybridization technique of Kohler and Milstein, Nature 256, 495 (1975). Other techniques for producing monoclonal antibody may also be employed, e.g. phage display techniques using libraries of human antibody genes. In one embodiment, anti-CD38 antibodies of the present invention produced by use of hybridomas generated in a murine system. Hybridoma production in the mouse is a very well established procedure. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known.

[0383] To generate fully human monoclonal antibodies to CD38, transgenic or transchromosomal mice containing human immunoglobulin genes (e.g., HCo12, HCo7 or KM mice) may be immunized with an enriched preparation of CD38 antigen and/or cells expressing CD38, as described, for example, by Lonberg et al., (1994), supra, Fishwild et al., (1996), supra, and WO 98/24884. Alternatively, mice may be immunized with DNA encoding human CD38. The mice may be 6-16 weeks of age upon the first infusion. For example, an enriched preparation (5-50 μ g) of the CD38 antigen may be used to immunize the HuMAb mice intraperitoneally. In the event that immunizations using a purified or enriched preparation of the CD38 antigen do not result in antibodies, mice may also be immunized with cells expressing CD38, e.g., a cell line, to promote immune responses.

[0384] Cumulative experience with various antigens has shown that the HuMAb transgenic mice respond best when initially immunized intraperitoneally (i.p.) or subcutaneously (s.c.) with CD38 expressing cells in complete Freund's adjuvant, followed by every other week i.p. immunizations (up to a total of 10) with CD38 expressing cells in PBS. The immune response may be monitored over the course of the immunization protocol with plasma samples being obtained by retroorbital bleeds. The plasma may be screened by FACS analysis, and mice with sufficient titers of anti-CD38 human immunoglobulin may be used for fusions. Mice may be boosted intravenously with CD38 expressing cells for Examples 4 and 3 days before sacrifice and removal of the spleen.

[0385] To generate hybridomas producing human monoclonal antibodies to human CD38, splenocytes and lymph node cells from immunized mice may be isolated and fused to an appropriate immortalized cell line, such as a mouse myeloma cell line. The resulting hybridomas may then be screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic lymphocytes from immunized mice may be fused to SP2/0 nonsecreting mouse myeloma cells (ATCC, CRL 1581) with 50% PEG (w/v). Cells may be plated at approximately 1×10^5 per well in flat bottom microtiter plate, followed by a two week incubation in selective medium containing besides usual reagents 10% fetal Clone Serum, 5-10% origin hybridoma cloning factor (IGEN) and 1X HAT (Sigma). After approximately two weeks, cells may be cultured in medium in which the HAT is replaced with HT. Individual wells may then be screened by ELISA for human kappa-light chain containing antibodies and by FACS analysis using CD38 expressing cells for CD38 specificity. Once extensive hybridoma growth occurs, medium may be observed usually after 10-14 days. The antibody secreting hybridomas may be replated, screened again, and if still positive for human IgG, anti-CD38 monoclonal antibodies may be subcloned at least twice by limiting dilution. The stable subclones may then be cultured *in vitro* to generate antibody in tissue culture medium for characterization.

[0386] Human antibodies of the present invention may also be produced in a host cell transfectoma using, for example, a combination of recombinant DNA techniques and gene transfection methods as is well known in the art, see for instance Morrison, S., Science 229, 1202 (1985).

[0387] For example, to express the antibodies, or antibody fragments thereof, DNAs encoding partial or full-length light and heavy chains, may be obtained by standard molecular biology techniques (for instance PCR amplification, site directed mutagenesis) and may be inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term "operatively linked" is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain

gene and the antibody heavy chain gene may be inserted into separate vectors or, more typically, both genes are inserted into the same expression vector. The antibody genes may be inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present). The light and heavy chain variable regions of the antibodies described herein may be used to create full-length antibody genes of any antibody isotype by inserting them into expression vectors already encoding heavy chain constant and light chain constant regions of the desired isotype such that the V_H segment is operatively linked to the CH segment(s) within the vector and the V_L segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector may encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide may be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

[0388] In addition to the antibody chain genes, the recombinant expression vectors of the present invention carry regulatory sequences that allows and control the expression of the antibody chain genes in a host cell.

[0389] In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the present invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see for instance US 4,399,216, US 4,634,665 and US 5,179,017). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Examples of selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr-host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

[0390] For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The host cells may be prokaryotic or eukaryotic, such as mammalian, host cells. For instance antigen binding fragments may be expressed in prokaryotic host cells and full-length antibodies may be expressed in eukaryotic host cells.

[0391] In one embodiment the antibodies are expressed in eukaryotic cells, such as mammalian host cells. Examples of mammalian host cells for expressing the recombinant antibodies of the present invention include CHO cells (including dhfr-CHO cells, described in Urlaub and Chasin, PNAS USA 77, 4216-4220 (1980), used with a DHFR selectable marker, for instance as described in R. J. Kaufman and P. A. Sharp, Mol. Biol. 159, 601-621 (1982)), NS/0 myeloma cells, COS cells, HEK293 cells and SP2.0 cells. In particular for use with NS/0 myeloma cells, another example of a expression system is the GS (glutamine synthetase) gene expression system disclosed in WO87/04462, WO89/01036 and EP338 841.

[0392] The CD38BP genes may be expressed in other expression systems, including prokaryotic cells, such as microorganisms, e.g. E. coli for the production of scFv antibodies, algae, as well as insect cells. Furthermore, the CD38BPs may be produced in transgenic non-human animals, such as in milk from sheep and rabbits or eggs from hens, or in transgenic plants. See for instance Verma, R. et al., J. Immunol. Meth. 216, 165-181 (1998), Pollock et al., J. Immunol. Meth. 231, 147-157 (1999) and Fischer, R. et al., Biol. Chem. 380, 825-839 (1999).

[0393] Bispecific and multispecific CD38BPs, such as antibodies of the present invention, may be made using chemical techniques (see for instance D. M. Kranz et al., PNAS USA 78, 5807 (1981)), "polydome" techniques (See US 4,474,893) or recombinant DNA techniques.

[0394] Bispecific antibodies of the present invention may be produced by a variety of known methods including fusion of hybridomas or linking of Fab' fragments (see for instance Songsivilai & Lachmann, Clin. Exp. Immunol. 79, 315-321 (1990) and Kostelny et al., J. Immunol. 148, 1547-1553 (1992)). Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (see for instance Milstein and Cuellar, Nature 305, 537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Similar procedures are disclosed in WO 93/08829 and Trautnecker et al., EMBO J. 10, 3655 (1991).

[0395] According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences by recombinant or synthetic methods. The variable domain sequence is typically fused to an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, C_H2 , and C_H3 regions. Also typically, a first heavy-chain constant region (C_H1), containing the site necessary for light chain binding, also is present in at least one of the fusion peptides. In a more specific example of this type of approach, a bispecific antibody is produced comprising a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. Such an asymmetric structure can facilitate the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations (such an approach is described in WO 94/04690). For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology 121, 210 (1986).

[0396] In another approach, the interface between a pair of antibody molecules may be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture so as to form a population of bispecific antibody molecules. Typically, such an interface comprises at least a part of the C_H3 domain of an antibody constant region. Normally in such a method, one or more amino acid residues with smaller side chains from the interface of the first antibody molecule are replaced with amino acid residues with larger side chains (such as tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain amino acid residue(s) are created on the interface of the second antibody molecule by replacing large amino acid side chain residues with smaller ones (such as alanine or threonine). This may provide a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0397] Bispecific and multispecific molecules of the present invention may be prepared by conjugating the constituent binding specificities, e.g., the anti-FcR and anti-CD38 binding specificities, using methods known in the art. For example, each binding specificity of the bispecific and multispecific molecule may be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents may be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-S-acetyl-thioacetate (SATA), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP) and sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (sulfo-SMCC), see for instance Karpovsky et al., J. Exp. Med. 160, 1686 (1984), Liu, M. A. et al., PNAS USA 82, 8648 (1985). In another example, Brennan et al., Science 229, 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated may then be converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives may then be reconverted to the Fab'-thiol by reduction with mercaptoethylamine and mixed with an equimolar amount of the other Fab'-TNB derivative to form a bispecific antibody. Shalaby et al., J. Exp. Med. 175, 217-225 (1992) describes the production of a fully humanized bispecific antibody F(ab')₂ molecule, according to a related technique. Other methods include those described by Paulus (Behring Ins. Mitt. No. 78, 118-132 (1985)) and Glennie et al., J. Immunol. 139, 2367-2375 (1987). Examples of conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, IL).

[0398] When the binding specificities are antibodies, they may be conjugated via sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In one embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, for instance one, prior to conjugation.

[0399] Alternatively, both binding specificities may be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific and multispecific molecule is a mAb x mAb, mAb x Fab, Fab x F(ab')₂ or ligand x Fab fusion protein. A bispecific and multispecific molecule of the present invention, e.g., a bispecific molecule may be a single chain molecule, such as a single chain bispecific antibody, a single chain bispecific molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific and multispecific molecules may also be single chain molecules or may comprise at least two single chain molecules. Methods for preparing bi- and multispecific molecules are described for example in US 5,260,203, US 5,455,030, US 4,881,175, US 5,132,405, US 5,091,513, US 5,476,786, US 5,013,653, US 5,258,498 and US 5,482,858.

[0400] Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers (see for instance Kostelny et al., J. Immunol. 148(5), 1547-1553 (1992)). Leucine zipper peptides from the Fos and Jun proteins can be linked to the Fab' portions of two different antibodies by gene fusion and the resulting antibody homodimers reduced at the hinge region to form monomers that can be re-oxidized to form the antibody heterodimers. The "diabody" technology described by Hollinger et al., PNAS USA 90, 6444-6448 (1993) also has provided an alternative mechanism for making bispecific antibody fragments. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See for instance Gruber et al., J. Immunol. 152, 5368 (1994).

[0401] In addition, bispecific antibodies may be formed as "diabodies" (Holliger et al., PNAS USA, 90, 6444-6448 (1993)) or "Janusins" (Traunecker et al., EMBO J 10, 3655-3659 (1991) and Traunecker et al., Int J Cancer Suppl 7, 51-52 (1992)). Bispecific antibodies, by definition, do not exist in the form of fragments having a single binding site (e.g., Fab, Fab', and Fv fragments, which also are provided by the present invention).

[0402] Binding of the bispecific and multispecific molecules to their specific targets may be confirmed by enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay (RIA), FACS analysis, a bioassay (e.g., growth inhibition), or a Western Blot Assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody) specific for the complex of interest. For example, the FcR-antibody complexes may be detected using e.g., an enzyme-linked antibody or antibody fragment which recognizes and specifically binds to the antibody-FcR complexes. Alternatively, the complexes may be detected using any of a variety of other immunoassays. For example, the antibody may be radioactively labeled and used in a radioimmunoassay (RIA)

(see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986). The radioactive isotope may be detected by such means as the use of a γ counter or a scintillation counter or by autoradiography.

[0403] As stated earlier, antibodies interact with target antigens primarily through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). Described herein are antibodies having CDR regions identical to or otherwise derived from the CDR regions of -003 or -024. Such antibodies may be generated by constructing expression vectors that include CDR sequences from -003 or -024 grafted onto framework sequences from a different antibody with different properties.

[0404] Antibodies of the invention have CDR regions identical to or otherwise derived from the CDR regions of -005. Such antibodies may be generated by constructing expression vectors that include CDR sequences from -005 grafted onto framework sequences from a different antibody with different properties.

[0405] Such framework sequences can be obtained from public DNA databases that include germline antibody gene sequences. These germline sequences will differ from mature antibody gene sequences because they will not include completely assembled variable genes, which are formed by V(D)J joining during B cell maturation. Germline gene sequences will also differ from the sequences of a high affinity secondary repertoire antibody which contains mutations throughout the variable gene but typically clustered in the CDRs. For example, somatic mutations are relatively infrequent in the amino terminal portion of framework region 1 and in the carboxy-terminal portion of framework region 4. For this reason, it is not necessary to obtain the entire DNA sequence of a particular antibody in order to recreate an intact recombinant antibody having binding properties similar to those of the original antibody (see WO 99/45962). Partial heavy and light chain sequence spanning the CDR regions is typically sufficient for this purpose. The partial sequence is used to determine which germline variable and joining gene segments contributed to the recombined antibody variable genes. The germline sequence is then used to fill in missing portions of the variable regions. Heavy and light chain leader sequences are cleaved during protein maturation and do not contribute to the properties of the final antibody. To add missing sequences, cloned cDNA sequences may be combined with synthetic oligonucleotides by ligation or PCR amplification. Alternatively, the entire variable region may be synthesized as a set of short, overlapping, oligonucleotides and combined by PCR amplification to create an entirely synthetic variable region clone. This process has certain advantages such as elimination or inclusion of particular restriction sites, or optimization of particular codons.

[0406] The nucleotide sequences of heavy and light chain transcripts from hybridomas are used to design an overlapping set of synthetic oligonucleotides to create synthetic V sequences with identical amino acid coding capacities as the natural sequences. The synthetic heavy and kappa chain sequences can differ from the natural sequences in three ways: strings of repeated nucleotide bases are interrupted to facilitate oligonucleotide synthesis and PCR amplification; optimal translation initiation sites are incorporated according to Kozak's rules (Kozak, J. Biol. Chem. 266, 19867-19870 (1991); and *HindIII* sites are engineered upstream of the translation initiation sites.

[0407] For both the heavy and light chain variable regions, the optimized coding and corresponding non-coding, strand sequences are broken down into 30-50 nucleotides approximately at the midpoint of the corresponding non-coding oligonucleotide. Thus, for each chain, the oligonucleotides may be assembled into overlapping double stranded sets that span segments of 150-400 nucleotides. The pools are then used as templates to produce PCR amplification products of 150-400 nucleotides. Typically, a single variable region oligonucleotide set will be broken down into two pools which are separately amplified to generate two overlapping PCR products. These overlapping products are then combined by PCR amplification to form the complete variable region. It may also be desirable to include an overlapping fragment of the heavy or light chain constant region (including the *BbsI* site of the kappa light chain, or the *AgeI* site of the gamma heavy chain) in the PCR amplification to generate fragments that can easily be cloned into the expression vector constructs.

[0408] The reconstructed heavy and light chain variable regions are then combined with cloned promoter, leader, translation initiation, constant region, 3' untranslated, polyadenylation, and transcription termination, sequences to form expression vector constructs. The heavy and light chain expression constructs may be combined into a single vector, co-transfected, serially transfected, or separately transfected into host cells which are then fused to form a host cell expressing both chains.

[0409] A similar procedure may be followed to graft novel antigen-specificity into an existing mature antibody. Typically, an acceptor antibody is chosen which originates from the same variable germ-line gene as the CDR-donor antibody, but other acceptor antibodies may also be chosen. One or more CDRs from the donor antibody are then transferred using the techniques described above.

[0410] The structural features of -003 and -005 and -024 may be used to create structurally related anti-CD38 antibodies, for instance human anti-CD38 antibodies, that retain at least one functional property of -003 and -005 and -024, namely binding to CD38. More specifically, one or more CDR regions of -003 or -005 and -024 may be combined recombinantly with known human framework regions and CDRs to create additional, recombinantly-engineered, human anti-CD38 antibodies.

[0411] Exemplary plasmids for use in construction of expression vectors for human IgG κ are described below. The

plasmids were constructed so that PCR amplified V kappa heavy and V kappa light chain cDNA sequences could be used to reconstruct complete heavy and light chain minigenes. These plasmids may be used to express completely human IgG1, κ or IgG4, κ antibodies. Similar plasmids may be constructed for expression of other heavy chain isotypes, or for expression of antibodies comprising lambda light chains.

[0412] CD38 antibodies of the present invention, such as human anti-CD38 antibodies of the present invention, may be isolated and characterized in a number of different ways. For example, selected hybridomas may be grown in suitable flasks for monoclonal antibody purification. Supernatants may then be filtered and concentrated before affinity chromatography with protein A-sepharose (for IgG1 isotype antibodies) (Pharmacia, Piscataway, NJ) or anti-human IgG coated sepharose or protein G-sepharose in case of IgG3 isotype antibodies. Eluted IgG may be checked by gel electrophoresis and high performance liquid chromatography to ensure purity. The buffer solution may be exchanged into PBS, and the concentration may be determined by OD₂₈₀ using 1.43 extinction coefficient. The monoclonal antibodies may be aliquoted and stored at -80°C.

[0413] To determine if the selected CD38BPs, such as human anti-CD38 monoclonal antibodies, bind to unique epitopes, site-directed or multi-site directed mutagenesis may be used.

[0414] To determine the isotype of purified antibodies, isotype ELISAs may be performed. Wells of microtiter plates may be coated with 10 μ g/ml of anti-human Ig overnight at 4°C. After blocking with 5% BSA (bovine serum albumin), the plates are reacted with 10 μ g/ml of monoclonal antibodies or purified isotype controls, at ambient temperature for two hours. The wells may then be reacted with either human IgG1, IgG2, IgG3 or IgG4, IgE, IgA1, IgA2, or human IgM-specific alkaline phosphatase-conjugated probes. After washing, the plates are developed with pNPP substrate (1 mg/ml) and analyzed by OD at 405 nm.

[0415] In order to demonstrate the presence of anti-CD38 antibodies in sera of immunized mice or the binding of CD38BPs (including anti-CD38 antibodies) to live cells expressing the CD38, flow cytometry may be used. Briefly, cell lines expressing CD38 (grown under standard growth conditions) are mixed with various concentrations of CD38BP in PBS containing 0.1% BSA and 0.02% sodium-azide, and incubated at 4°C for 30 minutes. After washing, the cells are reacted with fluorescein-labeled anti-human IgG antibody under the same conditions as the primary antibody staining. The samples may be analyzed by flow cytometry with a flow cytometer (e.g., Becton Dickinson FACS instrument) using light and side scatter properties to gate on single, living cells. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells may be stained exactly as described above and examined by fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

[0416] CD38BPs, such as anti-CD38 human IgGs, may be further tested for reactivity with CD38 antigen by Western blotting. Briefly, cell extracts from cells expressing CD38 may be prepared and subjected to sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens will be transferred to nitrocellulose membranes, blocked with 20% non-fat milk, and probed with the CD38BPs to be tested. Human IgG binding may be detected using anti-human IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, MO), but detecting agents directed at other specific portions of the CD38BP may also be used.

[0417] In addition to binding specifically to CD38, CD38BPs (including human anti-CD38 antibodies) may be tested for their ability to inhibit various activities of cells expressing CD38, such as but not restricted to insulin production, Ca²⁺ release, cytokine production, lysis induction, differentiation, and proliferation.

[0418] Described herein are transgenic and transchromosomal non-human animals, such as transgenic or transchromosomal mice, which are capable of expressing human antibodies that specifically bind to CD38. In particular, described herein is a transgenic or transchromosomal mouse having a genome comprising a human heavy chain transgene, such that the mouse produces human anti-CD38 antibodies when immunized with cells expressing CD38. The human heavy chain transgene may be integrated into the chromosomal DNA of the mouse, as is the case for transgenic, e.g., HuMab mice, as described in detail herein. Alternatively, the human heavy chain transgene may be maintained extrachromosomally, as is the case for transchromosomal (e.g., KM) mice as described in WO 02/43478. Such transgenic and transchromosomal animals are capable of producing multiple isotypes of human monoclonal antibodies to CD38 (e.g., IgG, IgA and/or IgE) by undergoing V-D-J/V-J recombination and isotype switching. The design of a transgenic or transchromosomal non-human animal that responds to foreign antigen stimulation with a heterologous antibody repertoire, requires that the heterologous immunoglobulin transgenes contained within the transgenic animal function correctly throughout the pathway of B cell development. This includes, for example, isotype switching of the heterologous heavy chain transgene. Accordingly, transgenes are constructed so that isotype switching may be induced and one or more of the following characteristics of antibody genes: (1) high level and cell-type specific expression, (2) functional gene rearrangement, (3) activation of and response to allelic exclusion, (4) expression of a sufficient primary repertoire, (5) signal transduction, (6) somatic hypermutation, and (7) domination of the transgene antibody locus during the immune response.

[0419] Not all of the foregoing criteria need be met. For example, where the endogenous immunoglobulin loci of the transgenic animal are functionally disrupted, the transgene need not activate allelic exclusion. Further, wherein the

transgene comprises a functionally rearranged heavy and/or light chain immunoglobulin gene, the second criteria of functional gene rearrangement is unnecessary, at least for that transgene which is already rearranged. For background on molecular immunology, see, Fundamental Immunology, 2nd edition (1989), Paul William E., ed. Raven Press, N.Y.

[0420] In certain embodiments, the transgenic or transchromosomal non-human animals used to generate the human monoclonal antibodies of the present invention contain rearranged, unrearranged or a combination of rearranged and unrearranged heterologous immunoglobulin heavy and light chain transgenes in the germline of the transgenic animal. Each of the heavy chain transgenes comprises at least one C_H gene. In addition, the heavy chain transgene may contain functional isotype switch sequences, which are capable of supporting isotype switching of a heterologous transgene encoding multiple C_H genes in the B cells of the transgenic animal. Such switch sequences may be those which occur naturally in the germline immunoglobulin locus from the species that serves as the source of the transgene C_H genes, or such switch sequences may be derived from those which occur in the species that is to receive the transgene construct (the transgenic animal). For example, a human transgene construct that is used to produce a transgenic mouse may produce a higher frequency of isotype switching events if it incorporates switch sequences similar to those that occur naturally in the mouse heavy chain locus, as presumably the mouse switch sequences are optimized to function with the mouse switch recombinase enzyme system, whereas the human switch sequences are not. Switch sequences may be isolated and cloned by conventional cloning methods, or may be synthesized *de novo* from overlapping synthetic oligonucleotides designed on the basis of published sequence information relating to immunoglobulin switch region sequences (Mills et al., Nucl. Acids Res. 15, 7305-7316 (1991) Sideras et al., Intl. Immunol. 1, 631-642 (1989)). For each of the foregoing transgenic animals, functionally rearranged heterologous heavy and light chain immunoglobulin transgenes are found in a significant fraction of the B cells of the transgenic animal (at least 10%).

[0421] The transgenes used to generate the transgenic non-human animals described herein include a heavy chain transgene comprising DNA encoding at least one variable gene segment, one diversity gene segment, one joining gene segment and at least one constant region gene segment. The immunoglobulin light chain transgene comprises DNA encoding at least one variable gene segment, one joining gene segment and at least one constant region gene segment. The gene segments encoding the light and heavy chain gene segments are heterologous to the transgenic animal in that they are derived from, or correspond to, DNA encoding immunoglobulin heavy and light chain gene segments from a species not consisting of the transgenic non-human animal. The transgene may be constructed such that the individual gene segments are unrearranged, i.e., not rearranged so as to encode a functional immunoglobulin light or heavy chain. Such unrearranged transgenes support recombination of the V, D, and J gene segments (functional rearrangement) and may support incorporation of all or a portion of a D region gene segment in the resultant rearranged immunoglobulin heavy chain within the transgenic animal when exposed to CD38 antigen.

[0422] In an alternate embodiment, the transgenes comprise an unrearranged "minilocus". Such transgenes typically comprise a substantial portion of the C, D, and J segments as well as a subset of the V gene segments. In such transgene constructs, the various regulatory sequences, e.g. promoters, enhancers, class switch regions, splice-donor and splice-acceptor sequences for RNA processing, recombination signals and the like, comprise corresponding sequences derived from the heterologous DNA. Such regulatory sequences may be incorporated into the transgene from the same or a related species of the non-human animal used. For example, human immunoglobulin gene segments may be combined in a transgene with a rodent immunoglobulin enhancer sequence for use in a transgenic mouse. Alternatively, synthetic regulatory sequences may be incorporated into the transgene, wherein such synthetic regulatory sequences are not homologous to a functional DNA sequence that is known to occur naturally in the genomes of mammals. Synthetic regulatory sequences are designed according to consensus rules, such as, for example, those specifying the permissible sequences of a splice-acceptor site or a promoter/enhancer motif. For example, a minilocus comprises a portion of the genomic immunoglobulin locus having at least one internal (i.e., not at a terminus of the portion) deletion of a non-essential DNA portion (e.g., intervening sequence; intron or portion thereof) as compared to the naturally-occurring germline Ig locus.

[0423] Examples of transgenic and transchromosomal non-human animals, such as mice, will exhibit immunoglobulin production with a significant repertoire, ideally substantially similar to that of a human after adjusting for volume.

[0424] The repertoire will ideally approximate that shown in a human when adjusted for volume, usually with a diversity at least about 10% as great, such as 25 to 50% or more. Generally, at least about a thousand different immunoglobulins (ideally IgG), such as 10^4 to 10^6 or more, will be produced, depending on the number of different V, J and D regions introduced into the mouse genome and driven by the additional diversity generated by V(-D-)J gene segment rearrangements and random nucleotide additions at the joining regions. Typically, the immunoglobulins will exhibit an affinity (K_D) for preselected antigens of below 10^{-8} M, such as of below 10^{-9} M, 10^{-10} M or 10^{-11} M or even lower. Transgenic and transchromosomal non-human animals, e.g., mice, as described above, may be immunized with, for example, cells expressing CD38. Alternatively, the transgenic animals may be immunized with DNA encoding human CD38. The animals will then produce B cells which undergo class-switching via switch recombination (cis-switching) and express immunoglobulins reactive with CD38. The immunoglobulins will be human antibodies (also referred to as "human sequence antibodies"), wherein the heavy and light chain polypeptides are encoded by human transgene sequences, which may

include sequences derived by somatic mutation and V region recombinatorial joints, as well as germline-encoded sequences; these human antibodies may be referred to as being substantially identical to a polypeptide sequence encoded by a human V_L and J_L or V_H , D_H and J_H gene segments, even though other non-germline sequences may be present as a result of somatic mutation and differential V-J and V-D-J recombination joints. The variable regions of each antibody chain are typically at least 80 percent similar to human germline V, and J gene segments, and, in the case of heavy chains, human germline V, D, and J gene segments; frequently at least 85 percent similar to human germline sequences present on the transgene; often 90 or 95 percent or more similar to human germline sequences present on the transgene. However, since non-germline sequences are introduced by somatic mutation and VJ and VDJ joining, the human sequence antibodies will frequently have some variable region sequences which are not encoded by human V, D, or J gene segments as found in the human transgene(s) in the germline of the mice. Typically, such non-germline sequences (or individual nucleotide positions) will cluster in or near CDRs, or in regions where somatic mutations are known to cluster.

[0425] The present disclosure also describes B cells derived from transgenic or transchromosomal non-human animals as described herein. The B cells may be used to generate hybridomas expressing human monoclonal antibodies which bind with high affinity (for instance with a dissociation equilibrium constant (K_D) of lower than 10^{-8} M) to human CD38. Thus, in one embodiment, provided herein is a hybridoma which produces a human antibody having an affinity (K_D) of below 10^{-8} M, such as of below 10^{-9} M, 10^{-10} M or 10^{-11} M or even lower when determined by scatchard analysis of CD38 expressing cells using a radio-actively labeled monoclonal antibody or by determination of the half-maximal binding concentration using FACS analysis, or by analysis using surface plasmon resonance as measured on a BIAcore instrument.

[0426] An anti-CD38 antibody of the present invention may comprise a human sequence light chain composed of (1) a light chain variable region having a polypeptide sequence which is substantially identical to a polypeptide sequence encoded by a human V_L gene segment and a human J_L segment, and (2) a light chain constant region encoded by a human C_L gene segment; and a human sequence heavy chain composed of a (1) a heavy chain variable region having a polypeptide sequence which is substantially identical to a polypeptide sequence encoded by a human V_H gene segment, a D region, and a human J_H segment, and (2) a constant region encoded by a human C_H gene segment. It should be noted that human D genes may be substantially altered by recombination and somatic mutation events such that the original human germ-line sequence may not be readily recognized.

[0427] The development of high affinity human monoclonal antibodies against CD38 can be facilitated by a method for expanding the repertoire of human variable region gene segments in a transgenic non-human animal having a genome comprising an integrated human immunoglobulin transgene, said method comprising introducing into the genome a V gene transgene comprising V region gene segments which are not present in said integrated human immunoglobulin transgene. Often, the V region transgene is a yeast artificial chromosome (YAC) comprising a portion of a human V_H or V_L (V_K) gene segment array, as may naturally occur in a human genome or as may be spliced together separately by recombinant methods, which may include out-of-order or omitted V gene segments. Often at least five or more functional V gene segments are contained on the YAC. In this variation, it is possible to make a transgenic animal produced by the V repertoire expansion method, wherein the animal expresses an immunoglobulin chain comprising a variable region sequence encoded by a V region gene segment present on the V region transgene and a C region encoded on the human Ig transgene. By means of the V repertoire expansion method, transgenic animals having at least 5 distinct V genes can be generated; as can animals containing at least about 24 V genes or more. Some V gene segments may be non-functional (e.g., pseudogenes and the like); these segments may be retained or may be selectively deleted by recombinant methods available to the skilled artisan, if desired.

[0428] Once the mouse germline has been engineered to contain a functional YAC having an expanded V segment repertoire, substantially not present in the human Ig transgene containing the J and C gene segments, the trait can be propagated and bred into other genetic backgrounds, including backgrounds where the functional YAC having an expanded V segment repertoire is bred into a non-human animal germline having a different human Ig transgene. Multiple functional YACs having an expanded V segment repertoire may be bred into a germline to work with a human Ig transgene (or multiple human Ig transgenes). Although referred to herein as YAC transgenes, such transgenes when integrated into the genome may substantially lack yeast sequences, such as sequences required for autonomous replication in yeast; such sequences may optionally be removed by genetic engineering (e.g., restriction digestion and pulsed-field gel electrophoresis or other suitable method) after replication in yeast is no longer necessary (i.e., prior to introduction into a mouse ES cell or mouse prozygote). Methods of propagating the trait of human sequence immunoglobulin expression, include breeding a transgenic animal having the human Ig transgene(s), and optionally also having a functional YAC having an expanded V segment repertoire. Both V_H and V_L gene segments may be present on the YAC. The transgenic animal may be bred into any background desired by the practitioner, including backgrounds harboring other human transgenes, including human Ig transgenes and/or transgenes encoding other human lymphocyte proteins. The present disclosure also describes a high affinity human sequence immunoglobulin produced by a transgenic mouse having an expanded V region repertoire YAC transgene. Although the foregoing describes a specific embodiment of the transgenic animal, other embodiments are contemplated which have been classified in three categories:

- I. Transgenic animals containing an unrearranged heavy and rearranged light chain immunoglobulin transgene;
- II. Transgenic animals containing an unrearranged heavy and unrearranged light chain immunoglobulin transgene;
- and
- III. Transgenic animal containing rearranged heavy and an unrearranged light chain immunoglobulin transgene.

[0429] In one embodiment, the present invention provides a pharmaceutical composition comprising a therapeutically effective amount of a CD38 antibody of the present invention. The pharmaceutical compositions may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

[0430] The pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients should be suitable for the chosen compound of the present invention and the chosen mode of administration. Suitability for carriers and other components of pharmaceutical compositions is determined based on the lack of significant negative impact on the desired biological properties of the chosen compound or pharmaceutical composition of the present invention (e.g., less than a substantial impact (10% or less relative inhibition, 5% or less relative inhibition, etc.) on antigen binding.

[0431] A pharmaceutical composition of the present inventions may also include diluents, fillers, salts, buffers, detergents (e. g., a nonionic detergent, such as Tween-80), stabilizers, stabilizers (e. g., sugars or protein-free amino acids), preservatives, tissue fixatives, solubilizers, and/or other materials suitable for inclusion in a pharmaceutical composition.

[0432] The actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0433] The pharmaceutical composition may be administered by any suitable route and mode. Suitable routes of administering a compound of the present invention *in vivo* and *in vitro* are well known in the art and can be selected by those of ordinary skill in the art.

[0434] The compounds of the present invention may be administered via any suitable route, such as an oral, nasal, inhalable, topical (including buccal, transdermal and sublingual), rectal, vaginal and/or parenteral route

[0435] In one embodiment, a pharmaceutical composition of the present invention is administered orally, for example, with an inert diluent or an assimilable edible carrier. The active ingredient may be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. Pharmaceutical compositions of the present invention which are suitable for oral administration include ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like containing such carriers as are known in the art to be appropriate. To administer a compound of the present invention oral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

[0436] In one embodiment, a pharmaceutical composition of the present invention is administered nasally. Pharmaceutical compositions of the present invention which are suitable for nasal administration are known in the art and typically include sprays, nose drops and inhalants.

[0437] In one embodiment, a pharmaceutical composition of the present invention is administered topically. Pharmaceutical compositions of the present invention which are suitable for topical or transdermal administration include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants containing such carriers as are known in the art to be appropriate.

[0438] In one embodiment, a pharmaceutical composition of the present invention is administered rectally. Pharmaceutical compositions of the present invention which are suitable for rectal administration are known in the art and include gels, pastes, spray formulations, suppositories.

[0439] In one embodiment, a pharmaceutical composition of the present invention is administered vaginally. Pharmaceutical compositions of the present invention which are suitable for vaginal administration include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

[0440] In one embodiment, a pharmaceutical composition of the present invention is administered parenterally.

[0441] The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and include epidermal, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, intratendinous, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, intracranial, intrathoracic, epidural and intrasternal injection and infusion.

[0442] In one embodiment that pharmaceutical composition is administered by intravenous or subcutaneous injection or infusion.

[0443] In one embodiment the compounds of the present invention are administered in crystalline form by subcutaneous injection, cf. Yang et al., PNAS USA 100(12), 6934-6939 (2003).

[0444] The pharmaceutical compositions may be administered with medical devices known in the art. For example, in one embodiment, a pharmaceutical composition of the present invention may be administered with a needleless hypodermic injection device, such as the devices disclosed in US 5,399,163, US 5,383,851, US 5,312,335, US 5,064,413, US 4,941,880, US 4,790,824, or US 4,596,556. Examples of well-known implants and modules useful in the present invention include: US 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; US 4,486,194, which discloses a therapeutic device for administering medicants through the skin; US 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; US 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; US 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and US 4,475,196, which discloses an osmotic drug delivery system. Many other such implants, delivery systems, and modules are known to those skilled in the art.

[0445] Pharmaceutical compositions of the present invention may be formulated for particular routes of administration, such as oral, nasal, topical (including buccal, transdermal and sublingual), rectal, vaginal and/or parenteral administration. The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any methods known in the art of pharmacy. The amount of active ingredient which may be combined with a carrier material to produce a single dosage form will vary depending upon the subject being treated, and the particular mode of administration. The amount of active ingredient which may be combined with a carrier material to produce a single dosage form will generally be that amount of the composition which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 0.01% to about 99% of active ingredient, such as from about 0.1 % to about 70%, for instance from about 1% to about 30%.

[0446] Regardless of the route of administration selected, the compounds of the present invention, which may be used in the form of a pharmaceutically acceptable salt or in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. A "pharmaceutically acceptable salt" refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects (see for instance Berge, S.M. et al., J. Pharm. Sci. 66, 1-19 (1977)). Examples of such salts include acid addition salts and base addition salts. Acid addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous acids and the like, as well as from nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanic acids, hydroxy alkanic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-dibenzylethylenediamine, N-methylglucamine, chlorprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

[0447] Pharmaceutically acceptable carriers include any and all suitable solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonicity agents, antioxidants and absorption delaying agents, and the like that are physiologically compatible with a compound of the present invention.

[0448] Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the present invention include water, saline, phosphate buffered saline, ethanol, dextrose, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, corn oil, peanut oil, cottonseed oil, and sesame oil, carboxymethyl cellulose colloidal solutions, tragacanth gum and injectable organic esters, such as ethyl oleate, and/or various buffers. Other carriers are well known in the pharmaceutical arts.

[0449] Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the present invention is contemplated.

[0450] Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0451] Pharmaceutical compositions of the present invention may also comprise pharmaceutically acceptable antioxidants for instance (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

[0452] Pharmaceutical compositions of the present invention may also comprise isotonicity agents, such as sugars,

polyalcohols such as mannitol, sorbitol, glycerol or sodium chloride in the compositions

[0453] Pharmaceutically acceptable diluents include saline and aqueous buffer solutions.

[0454] The pharmaceutical compositions of the present invention may also contain one or more adjuvants appropriate for the chosen route of administration such as preservatives, wetting agents, emulsifying agents, dispersing agents, preservatives or buffers, which may enhance the shelf life or effectiveness of the pharmaceutical composition. Compounds of the present invention may for instance be admixed with lactose, sucrose, powders (e.g., starch powder), cellulose esters of alkanolic acids, stearic acid, talc, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulphuric acids, acacia, gelatin, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol. Other examples of adjuvants are QS21, GM-CSF, SRL-172, histamine dihydrochloride, thymocartin, Tio-TEPA, monophosphoryl-lipid A/micobacteria compositions, alum, incomplete Freund's adjuvant, montanide ISA, ribi adjuvant system, TiterMax adjuvant, syntex adjuvant formulations, immune-stimulating complexes (ISCOMs), gerbu adjuvant, CpG oligodeoxynucleotides, lipopolysaccharide, and polyinosinic:polycytidylic acid.

[0455] Prevention of presence of microorganisms may be ensured both by sterilization procedures and by the inclusion of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0456] Pharmaceutical compositions of the present invention comprising a compound of the present invention may also include a suitable salt therefor. Any suitable salt, such as an alkaline earth metal salt in any suitable form (e.g., a buffer salt), may be used in the stabilization of the compound of the present invention. Suitable salts typically include sodium chloride, sodium succinate, sodium sulfate, potassium chloride, magnesium chloride, magnesium sulfate, and calcium chloride. In one embodiment, an aluminum salt is used to stabilize a compound of the present invention in a pharmaceutical composition of the present invention, which aluminum salt also may serve as an adjuvant when such a composition is administered to a patient.

[0457] Pharmaceutical compositions according to the present invention may be in a variety of suitable forms. Such forms include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, emulsions, microemulsions, gels, creams, granules, powders, tablets, pills, powders, liposomes, dendrimers and other nanoparticles (see for instance Baek et al., *Methods Enzymol.* 362, 240-9 (2003), Nigavekar et al., *Pharm Res.* 21(3), 476-83 (2004), microparticles, and suppositories.

[0458] The optimal form depends on the chosen mode of administration, the nature of the composition, and the therapeutic application. Formulations may include, for instance, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles, DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing may be appropriate in treatments and therapies in accordance with the present invention, provided that the active ingredient in the pharmaceutical composition is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also for instance Powell et al., "Compendium of excipients for parenteral formulations" *PDA J Pharm Sci Technol.* 52, 238-311 (1998) and the citations therein for additional information related to excipients and carriers well known to pharmaceutical chemists.

[0459] The compounds of the present invention may be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Such carriers may include gelatin, glyceryl monostearate, glyceryl distearate, biodegradable, biocompatible polymers such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid alone or with a wax, or other materials well known in the art. Methods for the preparation of such formulations are generally known to those skilled in the art. See e.g., *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

[0460] To administer compositions of the present invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound of the present invention may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al., *J. Neuroimmunol.* 7, 27 (1984)).

[0461] Depending on the route of administration, the active compound may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al., *J. Neuroimmunol.* 7, 27 (1984)).

[0462] In one embodiment, the compounds of the present invention may be formulated to ensure proper distribution *in vivo*. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds of the present invention cross the BBB (if desired), they may be formulated, for example, in liposomes. For methods of manufacturing liposomes, see for instance US 4,522,811, US 5,374,548 and US 5,399,331.

The liposomes may comprise one or more moieties which are selectively transported into specific cells or organs, thus enhance targeted drug delivery (see for instance V.V. Ranade J. Clin. Pharmacol. 29, 685 (1989)). Exemplary targeting moieties include folate or biotin (see for instance US 5,416,016), mannosides (Umezawa et al., Biochem. Biophys. Res. Commun. 153, 1038 (1988)), antibodies (P.G. Bloeman et al., FEBS Lett. 357, 140 (1995), M. Owais et al., Antimicrob. Agents Chemother. 39, 180 (1995)), surfactant protein A receptor (Briscoe et al., Am. J. Physiol. 1233, 134 (1995)), different species of which may comprise the pharmaceutical compositions of the present inventions, as well as components of the invented molecules, p120 (Schreier et al., J. Biol. Chem. 269, 9090 (1994)), see also K. Keinänen, M.L. Laukkanen, FEBS Lett. 346, 123 (1994) and J.J. Killion, I.J. Fidler, Immunomethods 4, 273 (1994).

[0463] Anti-CD38 antibodies of the present invention may be formulated in liposomes. The liposomes may include a targeting moiety. The antibodies in the liposomes may be delivered by bolus injection to a site proximal to the desired area, e.g., the site of inflammation or infection, or the site of a tumor. The composition must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.

[0464] In one embodiment, the antibodies of the present invention may be formulated to prevent or reduce their transport across the placenta. This may be done by methods known in the art, e.g., by PEGylation of the compounds or by use of F(ab')₂ fragments. Further references can be made to Cunningham-Rundles C et al., J Immunol Methods. 152, 177-190 (1992) and to Landor M., Ann Allergy Asthma Immunol 74, 279-283 (1995).

[0465] Pharmaceutically acceptable carriers for parenteral administration include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the present invention is contemplated. Supplementary active compounds may also be incorporated into the compositions.

[0466] Pharmaceutical compositions for injection must typically be sterile and stable under the conditions of manufacture and storage. The composition may be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier may be an aqueous or nonaqueous solvent or dispersion medium containing for instance water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. The proper fluidity may be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as glycerol, mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions may be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin. Sterile injectable solutions may be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients e.g. as enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients e.g. from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0467] Sterile injectable solutions may be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0468] The pharmaceutical composition of the present invention may contain one compound of the present invention or a combination of compounds of the present invention. Thus, in one embodiment, a pharmaceutical composition of the present invention includes a combination of multiple (e.g., two or more) compounds of the present invention which act by different mechanisms, e.g., one compound which predominately acts by inducing CDC in combination with another compound which predominately acts by inducing apoptosis.

[0469] The anti-CD38 antibodies (and immunoconjugates, bispecific/multispecific molecules, compositions and other derivatives described herein) of the present invention have numerous *in vitro* and *in vivo* diagnostic and therapeutic utilities involving the diagnosis and treatment of disorders involving cells expressing CD38. For example, the antibodies may be administered to cells in culture, e.g., *in vitro* or *ex vivo*, or to human subjects, e.g., *in vivo*, to treat, prevent and to diagnose a variety of disorders. As used herein, the term "subject" is intended to include human and non-human animals which respond to the CD38 antibody. Subjects may for instance include human patients having disorders that may be corrected or ameliorated by inhibiting CD38 function, such as enzymatic activity, signal transduction, induction of cytokine expression, induction of proliferation or differentiation, and/or induction of lysis and/or eliminating/reducing

the number of CD38 expressing cells.

[0470] For example, the CD38 antibodies may be used to elicit *in vivo* or *in vitro* one or more of the following biological activities: inhibition CD38 function (such as enzymatic activity, signal transduction, induction of cytokine expression, induction of proliferation or differentiation, and/or induction of lysis), killing a cell expressing CD38, mediating phagocytosis or ADCC of a cell expressing CD38 in the presence of human effector cells, and by mediating CDC of a cell expressing CD38 in the presence of complement. or by killing CD38 expressing cells by apoptosis.

[0471] Any composition comprising CD38 antibodies of the present invention having complement binding sites, such as portions from IgG1, -2, or -3 or IgM which bind complement, may also be used in the presence of complement. Ex vivo treatment of a population of cells comprising target cells with a CD38 antibody of the present invention and appropriate effector cells may be supplemented by the addition of complement or serum containing complement. Phagocytosis or lysis of target cells coated with a CD38 antibody of the present invention may be improved by binding of complement proteins. Target cells coated with the CD38 antibodies of the present invention may also be lysed by complement. In one embodiment, the CD38 antibodies of the present invention do not activate complement.

[0472] The CD38 antibodies of the present invention may also be administered together with complement. Accordingly, within the scope of the present invention are compositions comprising CD38 antibodies with serum or complement. In these compositions the complement is located in close proximity to the CD38 antibodies, for instance by conjugation or may be suited for simultaneous administration. Alternatively, the CD38 antibodies and the complement or serum may be administered separately.

[0473] The CD38 antibodies of the present invention may also be used to target cells expressing Fc γ R or CD38, for example for labeling such cells. For such use, the CD38 antibody may be linked to a molecule that can be detected. Thus, the present invention provides methods for localizing *ex vivo* or *in vitro* cells expressing Fc receptors, such as Fc γ R, or CD38. The detectable label may be, e.g., a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor.

[0474] Target-specific effector cells, e.g., effector cells linked to a CD38 antibody of the present invention may also be used as therapeutic agents. Effector cells for targeting may be human leukocytes such as macrophages, neutrophils or monocytes. Other cells include eosinophils, natural killer cells and other IgG- or IgA-receptor bearing cells. If desired, effector cells may be obtained from the subject to be treated. The target-specific effector cells, may be administered as a suspension of cells in a physiologically acceptable solution. The number of cells administered may be in the order of 10^8 to 10^9 but will vary depending on the therapeutic purpose. In general, the amount will be sufficient to obtain localization at the target cell, e.g., a tumor cell expressing CD38, and to effectively kill the cell by, e.g., phagocytosis or lysis.

[0475] Therapy with target-specific effector cells may be performed in conjunction with other techniques for removal of targeted cells. For example, anti-tumor therapy using the CD38 antibodies of the present invention and/or effector cells armed with these compositions may be used in conjunction with chemotherapy. Additionally, combination immunotherapy may be used to direct two distinct cytotoxic effector populations toward tumor cell rejection. For example, CD38 antibodies linked to anti-Fc γ RI or anti-CD3 may be used in conjunction with IgG- or IgA-receptor specific binding agents. Bispecific and multispecific molecules of the present invention may also be used to modulate Fc α R or Fc γ R levels on effector cells, such as by capping and elimination of receptors on the cell surface. Mixtures of anti-Fc receptors may also be used for this purpose.

[0476] In one embodiment, the present invention provides methods for detecting the presence of CD38 antigen in a sample, or measuring the amount of CD38 antigen, comprising contacting the sample, and a control sample, with a CD38 antibody of the invention which specifically binds to CD38, under conditions that allow for formation of a complex between the CD38 antibody or portion thereof and CD38. The formation of a complex is then detected, wherein a difference in complex formation between the sample compared to the control sample is indicative of the presence of CD38 antigen in the sample. Examples of methods for detecting immunoassays include, without limitation, an ELISA, an RIA, FACS assays, plasmon resonance assays, chromatographic assays, tissue immunohistochemistry, Western blot, and/or immunoprecipitation.

[0477] In one embodiment, CD38 antibodies of the present invention may be used to detect levels of circulating CD38 or levels of cells which contain CD38 on their membrane surface, which levels can then be linked to certain disease symptoms. Alternatively, the CD38 antibodies may be used to deplete or interact with the function of CD38 expressing cells, thereby implicating these cells as important mediators of the disease. This may be achieved by contacting a sample and a control sample with the anti-CD38 antibody under conditions that allow for the formation of a complex between the antibody and CD38. Any complexes formed between the antibody and CD38 are detected and compared in the sample and the control.

[0478] CD38 antibodies of the present invention may be initially tested for binding activity associated with therapeutic or diagnostic use *in vitro*. For example, the CD38 antibodies may be tested using flow cytometric assays. Moreover, activity of the CD38 antibodies in triggering at least one effector-mediated effector cell activity may be assayed. For example, the ability of anti-CD38 antibodies of the present invention to trigger CDC and/or apoptosis may be assayed. Protocols for assaying for CDC, homotypic adhesion, molecular clustering or apoptosis are well known in the art.

[0479] In one embodiment, the present invention provides a method for detecting the presence or quantifying the amount of CD38-expressing cells *in vivo* or *in vitro*. The method comprises (i) administering to a subject a CD38 antibody of the present invention conjugated to a detectable marker; (ii) exposing the subject to a means for detecting said detectable marker to identify areas containing CD38-expressing cells.

[0480] In one embodiment, immunoconjugates of the present invention may be used to target compounds (e.g., therapeutic agents, labels, cytotoxins, immunosuppressants, etc.) to cells which have CD38 bound to their surface by using such target compounds as the therapeutic moieties in immunoconjugates of the present invention.

[0481] Anti-CD38 antibodies of the present invention may be used in methods for localizing *ex vivo* or *in vitro* cells expressing CD38 (e.g., with a detectable label, such as a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor).

[0482] Anti-CD38 antibodies of the present invention may be used in methods for killing cells which have CD38 bound to their surface by administering immunotoxins of the present invention.

[0483] Anti-CD38 antibodies of the present invention may be used in methods for treating or preventing a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of a CD38 antibody of the present invention to a subject in need thereof. Such CD38 antibodies are used to inhibit CD38 induced activities associated with certain disorders or to eliminate or reduce the number of cells expressing CD38.

[0484] Such a method involves administering to a subject a CD38 antibody composition of the present invention in an amount effective to treat or prevent the disorder. The CD38 antibody composition may be administered alone or along with another therapeutic agent, such as is described elsewhere herein which acts in conjunction with or synergistically with the CD38 antibody composition to treat or prevent the diseases involving CD38 expressing cells. Alternatively, immunoconjugates may be used to kill cells which have CD38 expressed on their surface by targeting cytotoxins or radiotoxins to CD38.

[0485] The disorder involving cells expressing CD38 which may be treated with antibodies of the invention may be a tumorigenic disorder, such as a disorder characterized by the presence of tumor cells expressing CD38 including, for example, B cell lymphoma, plasma cell malignancies, T/NK cell lymphoma and myeloid malignancies.

[0486] Examples of such tumorigenic diseases include B cell lymphoma/leukemias including precursor B cell lymphoblastic leukemia/lymphoma and B cell non-Hodgkin's lymphomas; acute promyelocytic leukemia acute lymphoblastic leukemia and mature B cell neoplasms, such as B cell chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), B cell acute lymphocytic leukemia, B cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, mantle cell lymphoma (MCL), follicular lymphoma (FL), including low-grade, intermediate-grade and high-grade FL, cutaneous follicle center lymphoma, marginal zone B cell lymphoma (MALT type, nodal and splenic type), hairy cell leukemia, diffuse large B cell lymphoma, Burkitt's lymphoma, plasmacytoma, plasma cell myeloma, plasma cell leukemia, post-transplant lymphoproliferative disorder, Waldenström's macroglobulinemia, plasma cell leukemias and anaplastic large-cell lymphoma (ALCL).

[0487] In one embodiment, the disorder involving cells expressing CD38 is multiple myeloma.

[0488] Examples of B cell non-Hodgkin's lymphomas are lymphomatoid granulomatosis, primary effusion lymphoma, intravascular large B cell lymphoma, mediastinal large B cell lymphoma, heavy chain diseases (including γ , μ , and α disease), lymphomas induced by therapy with immunosuppressive agents, such as cyclosporine-induced lymphoma, and methotrexate-induced lymphoma.

[0489] In one embodiment of the present invention, the disorder involving cells expressing CD38 may be Hodgkin's lymphoma.

[0490] Examples of a disorder involving cells expressing CD38 may be a malignancy derived from T and NK cells including: mature T cell and NK cell neoplasms including T cell prolymphocytic leukemia, T cell large granular lymphocytic leukemia, aggressive NK cell leukemia, adult T cell leukemia/lymphoma, extranodal NK/T cell lymphoma, nasal type, enteropathy-type T cell lymphoma, hepatosplenic T cell lymphoma, subcutaneous panniculitis-like T cell lymphoma, blastic NK cell lymphoma, Mycosis Fungoides/Sézary Syndrome, primary cutaneous CD30 positive T cell lymphoproliferative disorders (primary cutaneous anaplastic large cell lymphoma C-ALCL, lymphomatoid papulosis, borderline lesions), angioimmunoblastic T cell lymphoma, peripheral T cell lymphoma unspecified, and anaplastic large cell lymphoma.

[0491] Examples of malignancies derived from myeloid cells include acute myeloid leukemia, including acute promyelocytic leukemia, and chronic myeloproliferative diseases, including chronic myeloid leukemia.

[0492] In one embodiment of the present invention, the disorder involving cells expressing CD38 may be immune disorders in which CD38 expressing B cells, plasma cells, monocytes and T cells are involved

[0493] Examples of immune disorders in which CD38 expressing B cells, plasma cells, monocytes and T cells are involved include autoimmune disorders, such as psoriasis, psoriatic arthritis, dermatitis, systemic scleroderma and sclerosis, inflammatory bowel disease (IBD), Crohn's disease, ulcerative colitis, respiratory distress syndrome, meningitis, encephalitis, uveitis, glomerulonephritis, eczema, asthma, atherosclerosis, leukocyte adhesion deficiency, multiple sclerosis, Raynaud's syndrome, Sjogren's syndrome, juvenile onset diabetes, Reiter's disease, Behget's disease, im-

immune complex nephritis, IgA nephropathy, IgM polyneuropathies, immune-mediated thrombocytopenias, such as acute idiopathic thrombocytopenic purpura and chronic idiopathic thrombocytopenic purpura, hemolytic anemia, myasthenia gravis, lupus nephritis, systemic lupus erythematosus, rheumatoid arthritis (RA), atopic dermatitis, pemphigus, Graves' disease, Hashimoto's thyroiditis, Wegener's granulomatosis, Osler-Weber's syndrome, chronic renal failure, acute infectious mononucleosis, multiple sclerosis, HIV, and herpes virus associated diseases. Further examples are severe acute respiratory distress syndrome and choreoretinitis. Furthermore, other diseases and disorders are included such as those caused by or mediated by infection of B-cells with virus, such as Epstein-Barr virus (EBV).

[0494] In one embodiment, the disorder involving cells expressing CD38 is rheumatoid arthritis.

[0495] Further examples of inflammatory, immune and/or autoimmune disorders in which autoantibodies and/or excessive B and T lymphocyte activity are prominent and which may be treated according to the present invention include the following:

vasculitides and other vessel disorders, such as microscopic polyangiitis, Churg-Strauss syndrome, and other ANCA-associated vasculitides, polyarteritis nodosa, essential cryoglobulinaemic vasculitis, cutaneous leukocytoclastic angiitis, Kawasaki disease, Takayasu arteritis, giant cell arthritis, Henoch-Schonlein purpura, primary or isolated cerebral angiitis, erythema nodosum, thrombangiitis obliterans, thrombotic thrombocytopenic purpura (including hemolytic uremic syndrome), and secondary vasculitides, including cutaneous leukocytoclastic vasculitis (e.g., secondary to hepatitis B, hepatitis C, Waldenström's macroglobulinemia, B-cell neoplasias, rheumatoid arthritis, Sjögren's syndrome, or systemic lupus erythematosus); further examples are erythema nodosum, allergic vasculitis, panniculitis, Weber-Christian disease, purpura hyperglobulinaemia, and Buerger's disease;

skin disorders, such as contact dermatitis, linear IgA dermatosis, vitiligo, pyoderma gangrenosum, epidermolysis bullosa acquisita, pemphigus vulgaris (including cicatricial pemphigoid and bullous pemphigoid), alopecia areata (including alopecia universalis and alopecia totalis), dermatitis herpetiformis, erythema multiforme, and chronic autoimmune urticaria (including angioneurotic edema and urticarial vasculitis);

immune-mediated cytopenias, such as autoimmune neutropenia, and pure red cell aplasia;

connective tissue disorders, such as CNS lupus, discoid lupus erythematosus, CREST syndrome, mixed connective tissue disease, polymyositis/dermatomyositis, inclusion body myositis, secondary amyloidosis, cryoglobulinemia type I and type II, fibromyalgia, phospholipid antibody syndrome, secondary hemophilia, relapsing polychondritis, sarcoidosis, stiff man syndrome, and rheumatic fever; a further example is eosinophil fasciitis;

arthritis, such as ankylosing spondylitis, juvenile chronic arthritis, adult Still's disease, and SAPHO syndrome; further examples are sacroileitis, reactive arthritis, Still's disease, and gout;

hematologic disorders, such as aplastic anemia, primary hemolytic anemia (including cold agglutinin syndrome), hemolytic anemia secondary to CLL or systemic lupus erythematosus; POEMS syndrome, pernicious anemia, and Waldenström's purpura hyperglobulinaemia; further examples are agranulocytosis, autoimmune neutropenia, Franklin's disease, Seligmann's disease, gamma heavy chain disease, paraneoplastic syndrome secondary to thymoma and lymphomas, an, paraneoplastic syndrome secondary to thymoma and lymphomas, and factor VIII inhibitor formation;

endocrinopathies, such as polyendocrinopathy, and Addison's disease; further examples are autoimmune hypoglycemia, autoimmune hypothyroidism, autoimmune insulin syndrome, de Quervain's thyroiditis, and insulin receptor antibody-mediated insulin resistance;

hepato-gastrointestinal disorders, such as celiac disease, Whipple's disease, primary biliary cirrhosis, chronic active hepatitis, and primary sclerosing cholangitis; a further example is autoimmune gastritis;

nephropathies, such as rapid progressive glomerulonephritis, post-streptococcal nephritis, Goodpasture's syndrome, membranous glomerulonephritis, and cryoglobulinemic nephritis; a further example is minimal change disease;

neurological disorders, such as autoimmune neuropathies, mononeuritis multiplex, Lambert-Eaton's myasthenic syndrome, Sydenham's chorea, tabes dorsalis, and Guillain-Barre's syndrome; further examples are myelopathy/tropical spastic paraparesis, myasthenia gravis, acute inflammatory demyelinating polyneuropathy, and chronic inflammatory demyelinating polyneuropathy; multiple sclerosis;

cardiac and pulmonary disorders, such as COPD, fibrosing alveolitis, bronchiolitis obliterans, allergic aspergillosis, cystic fibrosis, Löfller's syndrome, myocarditis, and pericarditis; further examples are hypersensitivity pneumonitis, and paraneoplastic syndrome secondary to lung cancer;

allergic disorders, such as bronchial asthma and hyper-IgE syndrome; a further example is amaurosis fugax;

ophthalmologic disorders, such as idiopathic chorioretinitis;

infectious diseases, such as parvovirus B infection (including hands-and-socks syndrome);

gynecological-obstetrical disorders, such as recurrent abortion, recurrent fetal loss, and intrauterine growth retardation; a further example is paraneoplastic syndrome secondary to gynaecological neoplasms;

male reproductive disorders, such as paraneoplastic syndrome secondary to testicular neoplasms; and

transplantation-derived disorders, such as allograft and xenograft rejection, and graft-versus-host disease.

[0496] The antibody may also be administered prophylactically in order to reduce the risk of developing cancer, such as a tumorigenic disorder as described above, delay the onset of the occurrence of an event in such cancer progression, and/or reduce the risk of recurrence when such a cancer is in remission. This may be especially useful in patients wherein it is difficult to locate a tumor that is known to be present due to other biological factors.

[0497] Compositions of the present invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of a CD38 antibody of the invention. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result. A therapeutically effective amount of a CD38 antibody may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the CD38 antibody to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result (e.g., a reduction in the likelihood of developing a disorder, a reduction in the intensity or spread of a disorder, an increase in the likelihood of survival during an imminent disorder, a delay in the onset of a disease condition, etc.). Typically, because a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

[0498] A "therapeutically effective amount" for tumor therapy may also be measured by its ability to stabilize the progression of disease. The ability of a compound to inhibit cancer may be evaluated in an animal model system predictive of efficacy in human tumors. Alternatively, this property of a composition may be evaluated by examining the ability of the compound to inhibit cell growth or to induce apoptosis by *in vitro* assays known to the skilled practitioner. A therapeutically effective amount of a therapeutic compound may decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

[0499] A "therapeutically effective amount" for rheumatoid arthritis may result in an at least ACR₂₀ Preliminary Definition of Improvement in the patients, such as in at least an ACR₅₀ Preliminary Definition of Improvement, for instance at least an ARC₇₀ Preliminary Definition of Improvement.

[0500] ACR₂₀ Preliminary Definition of Improvement is defined as:

≥ 20% improvement in: Tender Joint Count (TJC) and Swollen Joint Count (SJC)
and ≥ 20% improvement in 3 of following 5 assessments: Patient Pain Assessment (VAS), Patient Global assessment (VAS), Physician Global Assessment (VAS), Patient Self-Assessed Disability (HAQ), Acute Phase Reactant (CRP or ESR).

[0501] ACR₅₀ and ACR₇₀ are defined in the same way with ≥ 50% and ≥ 70% improvements, respectively. For further details see Felson et al., in American College of Rheumatology Preliminary Definition of Improvement in Rheumatoid Arthritis; *Arthritis Rheumatism* 38, 727-735 (1995).

[0502] Alternatively, a therapeutically effective amount for rheumatoid arthritis can be measured by DAS (disease activity score), including DAS28 and/or DAS56, as defined by EULAR.

[0503] Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[0504] The efficient dosages and the dosage regimens for the CD38 antibodies of the present invention depend on the disease or condition to be treated and may be determined by the persons skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of a compound of the present invention is about 0.1-100 mg/kg, such as about 0.1-50 mg/kg, for example about 0.1-20 mg/kg, such as about 0.1-10 mg/kg, for instance about 0.5, about 0.3, about 1, or about 3 mg/kg.

[0505] A physician or veterinarian having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the CD38 antibodies of the present invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a composition of the present invention will be that amount of the compound which is

the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Administration may be intravenous, intramuscular, intraperitoneal, or subcutaneous, and for instance administered proximal to the site of the target. If desired, the effective daily dose of a pharmaceutical composition may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical composition as described above.

[0506] In one embodiment, the CD38 antibodies of the present invention may be administered by infusion in a weekly dosage of from 10 to 500 mg/m², such as of from 200 to 400 mg/m². Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours.

[0507] In one embodiment, the CD38 antibodies of the present invention may be administered by slow continuous infusion over a long period, such as more than 24 hours, in order to reduce toxic side effects.

[0508] In one embodiment the CD38 antibodies of the present invention may be administered in a weekly dosage of from 250 mg to 2000 mg, such as for example 300 mg, 500 mg, 700 mg, 1000 mg, 1500 mg or 2000 mg, for up to 8 times, such as from 4 to 6 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours. Such regimen may be repeated one or more times as necessary, for example, after 6 months or 12 months. The dosage may be determined or adjusted by measuring the amount of compound of the present invention in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the antigen binding region of the CD38 antibodies of the present invention.

[0509] In one embodiment, the CD38 antibodies of the present invention may be administered by maintenance therapy, such as, e.g., once a week for a period of 6 months or more.

[0510] In one embodiment, the CD38 antibodies of the present invention may be administered by a regimen including one infusion of a CD38 antibody of the present invention followed by an infusion of a CD38 antibody of the present invention conjugated to a radioisotope. The regimen may be repeated, e.g., 7 to 9 days later.

[0511] As non-limiting examples, treatment according to the present invention may be provided as a daily dosage of an antibody of the present invention in an amount of about 0.1-100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 after initiation of treatment, or any combination thereof, using single or divided doses of every 24, 12, 8, 6, 4, or 2 hours, or any combination thereof.

[0512] The pharmaceutical compositions of the present invention may also be administered in combination therapy, i.e., combined with other therapeutic agents relevant for the disease or condition to be treated. Such administration may be simultaneous, separate or sequential. For simultaneous administration the agents may be administered as one compositions or as separate compositions, as appropriate.

[0513] Accordingly, the present invention provides antibodies of the present invention for use in methods for treating a disorder involving cells expressing CD38 as described above, which methods comprise administration of a CD38BP of the present invention combined with one or more additional therapeutic agents as described below.

[0514] Antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with at least one chemotherapeutic agent for a disorder involving cells expressing CD38 as described above.

[0515] The combination therapy may include administration of a composition of the present invention together with at least one chemotherapeutic agent, at least one anti-inflammatory agent, or at least one immunosuppressive and/or immunomodulatory agent.

[0516] In one embodiment, the present invention provides an antibody of the invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and at least one chemotherapeutic agent to a subject in need thereof

[0517] In one embodiment, the present invention provides an antibody of the invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody and at least one chemotherapeutic agent to a subject in need thereof.

[0518] In one embodiment, antibodies of the present invention are used for the preparation of a pharmaceutical composition to be administered with at least one chemotherapeutic agent for treating multiple myeloma.

[0519] In one embodiment, such a chemotherapeutic agent may be selected from an antimetabolite, such as methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, fludarabine, 5-fluorouracil, decarbazine, hydroxyurea, asparaginase, gemcitabine, cladribine and similar agents.

[0520] In one embodiment, such a chemotherapeutic agent may be selected from an alkylating agent, such as mechlorethamine, thioepa, chlorambucil, melphalan, carmustine (BSNU), lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, dacarbazine (DTIC), procarbazine, mitomycin C, cisplatin and other platinum derivatives, such as carboplatin, and similar agents.

[0521] In one embodiment, such a chemotherapeutic agent may be selected from an antibiotic, such as dactinomycin (formerly actinomycin), bleomycin, daunorubicin (formerly daunomycin), doxorubicin, idarubicin, mithramycin, mitomycin, mitoxantrone, plicamycin, anthramycin (AMC) and similar agents.

[0522] In one embodiment, such a chemotherapeutic agent may be selected from an anti-mitotic agent, such as taxanes, for instance docetaxel, and paclitaxel, and vinca alkaloids, for instance vindesine, vincristine, vinblastine, and vinorelbine.

[0523] In one embodiment, such a chemotherapeutic agent may be selected from a topoisomerase inhibitor, such as topotecan.

[0524] In one embodiment, such a chemotherapeutic agent may be selected from a growth factor inhibitor, such as an inhibitor of ErbB1 (EGFR) (such as gefitinib (Iressa®), cetuximab (Erbix®), erlotinib (Tarceva®), HuMax-EGFr (2F8 disclosed in WO 2002/100348) and similar agents), an inhibitor of ErbB2 (Her2/neu) (such as trastuzumab (Herceptin®) and similar agents) and similar agents. In one embodiment, such a growth factor inhibitor may be a farnesyl transferase inhibitor, such as SCH-66336 and R115777. In one embodiment, such a growth factor inhibitor may be a vascular endothelial growth factor (VEGF) inhibitor, such as bevacizumab (Avastin®).

[0525] In one embodiment, such a chemotherapeutic agent may be a tyrosine kinase inhibitor, such as imatinib (Gleevec, Gleevec ST1571), lapatinib, PTK787/ZK222584 and similar agents.

[0526] In one embodiment, such a chemotherapeutic agent may be a histone deacetylase inhibitor. Examples of such histone deacetylase inhibitors include hydroxamic acid-based hybrid polar compounds, such as SAHA (suberoylanilide hydroxamic acid).

[0527] In one embodiment, such a chemotherapeutic agent may be a P38a MAP kinase inhibitor, such as SCIO-469.

[0528] In one embodiment, the present invention provides antibodies of the invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and at least one inhibitor of angiogenesis, neovascularization, and/or other vascularization to a subject in need thereof.

[0529] In one embodiment, the present invention provides antibodies of the invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of a CD38 antibody of the present invention and at least one inhibitor of angiogenesis, neovascularization, and/or other vascularization to a subject in need thereof.

[0530] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with at least one inhibitor of angiogenesis, neovascularization, and/or other vascularization for treating multiple myeloma.

[0531] Examples of such angiogenesis inhibitors are urokinase inhibitors, matrix metalloprotease inhibitors (such as marimastat, neovastat, BAY 12-9566, AG 3340, BMS-275291 and similar agents), inhibitors of endothelial cell migration and proliferation (such as TNP-470, squalamine, 2-methoxyestradiol, combretastatins, endostatin, angiostatin, penicillamine, SCH66336 (Schering-Plough Corp, Madison, NJ), R115777 (Janssen Pharmaceutica, Inc, Titusville, NJ) and similar agents), antagonists of angiogenic growth factors (such as such as ZD6474, SU6668, antibodies against angiogenic agents and/or their receptors (such as VEGF, bFGF, and angiopoietin-1), thalidomide (Thalomid®), thalidomide analogs (such as CC-5013 (lenalidomide, Revlimid™) and CC4047 (Actimid™), Sugen 5416, SU5402, antiangiogenic ribozyme (such as angiozyme), interferon α (such as interferon α 2a), suramin and similar agents), VEGF-R kinase inhibitors and other anti-angiogenic tyrosine kinase inhibitors (such as SU011248), inhibitors of endothelial-specific integrin/survival signaling (such as vitaxin and similar agents), copper antagonists/chelators (such as tetrathiomolybdate, captopril and similar agents), carboxyamido-triazole (CAI), ABT-627, CM101, interleukin-12 (IL-12), IM862, PNU145156E as well as nucleotide molecules inhibiting angiogenesis (such as antisense-VEGF-cDNA, cDNA coding for angiostatin, cDNA coding for p53 and cDNA coding for deficient VEGF receptor-2) and similar agents.

[0532] Other examples of such inhibitors of angiogenesis, neovascularization, and/or other vascularization are anti-angiogenic heparin derivatives and related molecules (e.g., heparinase III), temozolomide, NK4, macrophage migration inhibitory factor (MIF), cyclooxygenase-2 inhibitors, inhibitors of hypoxia-inducible factor 1, anti-angiogenic soy isoflavones, oltipraz, fumagillin and analogs thereof, somatostatin analogues, pentosan polysulfate, tecogalan sodium, dalteparin, tumstatin, thrombospondin, NM-3, combrestatin, canstatin, avastatin, antibodies against other relevant targets (such as anti- α -v/ β -3 integrin and anti-kininostatin mAbs) and similar agents.

[0533] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with thalidomide (Thalomid®), thalidomide analogs (such as CC-5013 (lenalidomide, Revlimid™) and/or CC4047 (Actimid™). In a further embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with thalidomide.

[0534] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with an anti-CD20 antibody, such as rituximab (Rituxan®, Mabthera®), a human monoclonal anti-CD20 antibody as disclosed in WO 2004/035607, such as 11 B8, 2F2 or 7D8.

[0535] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention

for treating the disorders as described above may be a proteasome inhibitor, such as bortezomib (Velcade®).

[0536] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a corticosteroid, such as prednisone, prednisolone, dexamethasone, etc.

[0537] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a corticosteroid, such as prednisone, prednisolone, dexamethasone, etc.

[0538] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an anti-cancer immunogen, such as a cancer antigen/tumor-associated antigen (e.g., epithelial cell adhesion molecule (EpCAM/TACSTD1), mucin 1 (MUC1), carcinoembryonic antigen (CEA), tumor-associated glycoprotein 72 (TAG-72), gp100, Melan-A, MART-1, KDR, RCAS1, MDA7, cancer-associated viral vaccines (e.g., human papillomavirus vaccines), tumor-derived heat shock proteins, and similar agents. A number of other suitable cancer antigens/tumor-associated antigens described elsewhere herein and similar molecules known in the art may also or alternatively be used in such embodiment. Anti-cancer immunogenic peptides also include anti-idiotypic "vaccines" such as BEC2 anti-idiotypic antibodies, Mitumomab, CeaVac and related anti-idiotypic antibodies, anti-idiotypic antibody to MG7 antibody, and other anti-cancer anti-idiotypic antibodies (see for instance Birebent et al., Vaccine. 21(15), 1601-12 (2003), Li et al., Chin Med J (Engl). 114(9), 962-6 (2001), Schmitt et al., Hybridoma. 13(5), 389-96 (1994), Maloney et al., Hybridoma. 4(3), 191-209 (1985), Raychardhuri et al., J Immunol. 137(5), 1743-9 (1986), Pohl et al., Int J Cancer. 50(6), 958-67 (1992), Bohlen et al., Cytokines Mol Ther. 2(4), 231-8 (1996) and Maruyama, J Immunol Methods. 264(1-2), 121-33 (2002)). Such anti-idiotypic Abs may optionally be conjugated to a carrier, which may be a synthetic (typically inert) molecule carrier, a protein (for instance keyhole limpet hemocyanin (KLH) (see for instance Ochi et al., Eur J Immunol. 17(11), 1645-8 (1987)), or a cell (for instance a red blood cell - see for instance Wi et al., J Immunol Methods. 122(2), 227-34 (1989)).

[0539] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a bisphosphonate. Examples of potentially suitable bisphosphonates are pamidronate (Aredia®), zoledronic acid (Zometa®), clodronate (Bonefos®), risendronate (Actonel®), ibandronate (Boniva®), etidronate (Didronel®), alendronate (Fosamax®), tiludronate (Skelid®), incadronate (Yamanouchi Pharmaceutical) and minodronate (YM529, Yamanouchi).

[0540] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a colony stimulating factor. Examples of suitable colony stimulating factors are granulocyte-colony stimulating factors (G-CSF), such as filgrastim (Neupogen®) and pegfilgrastim (Neulasta®), and granulocyte macrophage-colony stimulating factors (GM-CSF) such as sargramostim (Leukine®).

[0541] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an erythropoietic agent. Examples of suitable erythropoietic agents are erythropoietin (EPO), such as epoetin alfa (for instance Procrit®, Epogen®, and Eprex®) and epoetin beta (for instance NeoRecormon®) and erythropoiesis-stimulating proteins (for instance Aranesp®).

[0542] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an anti-cancer cytokine, chemokine, or combination thereof. Examples of suitable cytokines and growth factors include IFN γ , IL-2, IL-4, IL-6, IL-7, IL-10, IL-12, IL-13, IL-15, IL-18, IL-23, IL-24, IL-27, IL-28a, IL-28b, IL-29, KGF, IFN α (e.g., IFN α 2b), IFN β , GM-CSF, CD40L, Flt3 ligand, stem cell factor, ancestim, and TNF α . Suitable chemokines may include Glu-Leu-Arg (ELR)-negative chemokines such as IP-10, MCP-3, MIG, and SDF-1 α from the human CXC and C-C chemokine families. Suitable cytokines include cytokine derivatives, cytokine variants, cytokine fragments, and cytokine fusion proteins.

[0543] These and other methods or uses involving naturally occurring peptide-encoding nucleic acids herein may alternatively or additionally be performed by "gene activation" and homologous recombination gene upregulation techniques, such as are described in US 5,968,502, US 6,063,630 and US 6,187,305 and EP 0505500.

[0544] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an agent that modulates, e.g., enhances or inhibits, the expression or activity of Fc α or Fc γ receptors. Examples of agents suitable for this use include interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte colony-stimulating factor (G-CSF), such as filgrastim (Neupogen®) and pegfilgrastim (Neulasta®), and granulocyte macrophage-colony stimulating factors (GM-CSF) such as sargramostim (Leukine®), interferon- γ (IFN- γ), and tumor necrosis factor (TNF).

[0545] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a cell cycle control/apoptosis regulator (or "regulating agent"). A cell cycle control/apoptosis regulator may include molecules (i) that target and modulate cell cycle control/apoptosis regulators such as cdc-25 (such as NSC 663284), (ii) cyclin-dependent kinases that overstimulate the cell cycle (such as flavopiridol (L868275, HMR1275), 7-hydroxystaurosporine (UCN-01, KW-2401), and roscovitine (R-roscovitine, CYC202)), and (iii) telomerase modulators (such as BIBR1532, SOT-095, GRN163 and compositions described in for

instance US 6,440,735 and US 6,713,055). Non-limiting examples of molecules that interfere with apoptotic pathways include TNF-related apoptosis-inducing ligand (TRAIL)/apoptosis-2 ligand (Apo-2L), agents inducing NF- κ B blockade leading to inhibition of IL-6 production, antibodies that activate TRAIL receptors, IFNs, anti-sense Bcl-2, and As₂O₃ (arsenic trioxide, Trisenox®).

[0546] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a hormonal regulating agent, such as agents useful for anti-androgen and anti-estrogen therapy. Examples of such hormonal regulating agents are tamoxifen, idoxifene, fulvestrant, droloxifene, toremifene, raloxifene, diethylstilbestrol, ethinyl estradiol/estinyl, an antiandrogene (such as flutaminde/eulexin), a progestin (such as hydroxy-progesterone caproate, medroxyprogesterone/provera, megestrol acepate/megace), an adrenocorticosteroid (such as hydrocortisone, prednisone), luteinizing hormone-releasing hormone (and analogs thereof and other LHRH agonists such as buserelin and goserelin), an aromatase inhibitor (such as anastrozole/arimidex, aminoglutethimide/cytraden, exemestane), a hormone inhibitor (such as octreotide/- sandostatin) and similar agents.

[0547] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an anti-nergic agents (for instance small molecule compounds, proteins, glycoproteins, or antibodies that break tolerance to tumor and cancer antigens). Examples of such compounds are molecules that block the activity of CTLA-4, such as MDX-010 (Phan et al., PNAS USA 100, 8372 (2003)).

[0548] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be a tumor suppressor gene-containing nucleic acid or vector such as a replication-deficient adenovirus encoding human recombinant wild-type p53/SCH58500, etc.; antisense nucleic acids targeted to oncogenes, mutated, or deregulated genes; or siRNA targeted to mutated or deregulated genes. Examples of tumor suppressor targets include, for example, BRCA1, RB1, BRCA2, DPC4 (Smad4), MSH2, MLH1, and DCC.

[0549] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an anti-cancer nucleic acid, such as genasense (augmerosen/G3139), LY900003 (ISIS 3521), ISIS 2503, OGX-011 (ISIS 112989), LE-AON/LEraf-AON (liposome encapsulated c-raf antisense oligonucleotide/ISIS-5132), MG98, and other antisense nucleic acids that target PKC α , clusterin, IGFBPs, protein kinase A, cyclin D1, or Bcl-2h.

[0550] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be an anti-cancer inhibitory RNA molecule (see for instance Lin et al., Curr Cancer Drug Targets. 1(3), 241-7 (2001), Erratum in: Curr Cancer Drug Targets. 3(3), 237 (2003), Lima et al., Cancer Gene Ther. 11(5), 309-16 (2004), Grzmil et al., Int J Oncol. 4(1), 97-105 (2004), Collis et al., Int J Radiat Oncol Biol Phys. 57(2 Suppl), S144 (2003), Yang et al., Oncogene. 22(36), 5694-701 (2003) and Zhang et al., Biochem Biophys Res Commun. 303(4), 1169-78 (2003)).

[0551] Compositions and combination administration methods of the present invention also include the administration of nucleic acid vaccines, such as naked DNA vaccines encoding such cancer antigens/tumor-associated antigens (see for instance US 5,589,466, US 5,593,972, US 5,703,057, US 5,879,687, US 6,235,523, and US 6,387,888). In one embodiment, the combination administration method and/or combination composition comprises an autologous vaccine composition. In one embodiment, the combination composition and/or combination administration method comprises a whole cell vaccine or cytokine-expressing cell (for instance a recombinant IL-2 expressing fibroblast, recombinant cytokine-expressing dendritic cell, and the like) (see for instance Kowalczyk et al., Acta Biochim Pol. 50(3), 613-24 (2003), Reilly et al., Methods Mol Med. 69, 233-57 (2002) and Tirapu et al., Curr Gene Ther. 2(1), 79-89 (2002). Another example of such an autologous cell approach that may be useful in combination methods of the present invention is the MyVax® Personalized Immunotherapy method (previously referred to as GTOP-99) (Genitope Corporation - Redwood City, CA, USA).

[0552] In one embodiment, the present invention provides combination compositions and combination administration methods wherein a CD38 antibody of the invention is combined or co-administered with a virus, viral proteins, and the like. Replication-deficient viruses, that generally are capable of one or only a few rounds of replication *in vivo*, and that are targeted to tumor cells, may for instance be useful components of such compositions and methods. Such viral agents may comprise or be associated with nucleic acids encoding immunostimulants, such as GM-CSF and/or IL-2. Both naturally oncolytic and such recombinant oncolytic viruses (for instance HSV-1 viruses, reoviruses, replication-deficient and replication-sensitive adenovirus, etc.) may be useful components of such methods and compositions. Accordingly, in one embodiment, the present invention provides combination compositions and combination administration methods wherein a CD38 antibody of the invention is combined or co-administered with an oncolytic virus. Examples of such viruses include oncolytic adenoviruses and herpes viruses, which may or may not be modified viruses (see for instance Shah et al., J Neurooncol. 65(3), 203-26 (2003), Stiles et al., Surgery. 134(2), 357-64 (2003), Sunarmura et al., Pancreas. 28(3), 326-9 (2004), Teshigahara et al., J Surg Oncol. 85(1), 42-7 (2004), Varghese et al., Cancer Gene Ther. 9(12), 967-78 (2002), Wildner et al., Cancer Res. 59(2), 410-3 (1999), Yamanaka, Int J Oncol. 24(4), 919-23 (2004) and Zwiebel

et al., Semin Oncol. 28(4), 336-43 (2001).

[0553] Combination compositions and combination administration methods involving antibodies of the present invention may also involve "whole cell and "adoptive" immunotherapy methods. For instance, such methods may comprise infusion or re-infusion of immune system cells (for instance tumor-infiltrating lymphocytes (TILs), such as CD4⁺ and/or CD8⁺T cells (for instance T cells expanded with tumor-specific antigens and/or genetic enhancements), antibody-expressing B cells or other antibody producing/presenting cells, dendritic cells (e.g., anti-cytokine expressing recombinant dendritic cells, dendritic cells cultured with a DC-expanding agent such as GM-CSF and/or Flt3-L, and/or tumor-associated antigen-loaded dendritic cells), anti-tumor NK cells, so-called hybrid cells, or combinations thereof. Cell lysates may also be useful in such methods and compositions. Cellular "vaccines" in clinical trials that may be useful in such aspects include Canvaxin™, APC-8015 (Dendreon), HSPPC-96 (Antigenics), and Melacine® cell lysates. Antigens shed from cancer cells, and mixtures thereof (see for instance Bystryn et al., Clinical Cancer Research Vol. 7, 1882-1887, July 2001), optionally admixed with adjuvants such as alum, may also be components in such methods and combination compositions.

[0554] In one embodiment, a CD38 antibody of the present invention may be delivered to a patient in combination with the application of an internal vaccination method. internal vaccination refers to induced tumor or cancer cell death, such as drug-induced or radiation-induced cell death of tumor cells, in a patient, that typically leads to elicitation of an immune response directed towards (i) the tumor cells as a whole or (ii) parts of the tumor cells including (a) secreted proteins, glycoproteins or other products, (b) membrane-associated proteins or glycoproteins or other components associated with or inserted in membranes, and/or (c) intracellular proteins or other intracellular components. An internal vaccination-induced immune response may be humoral (i.e. antibody - complement-mediated) or cell-mediated (e.g., the development and/or increase of endogenous cytotoxic T lymphocytes that recognize the internally killed tumor cells or parts thereof). In addition to radiotherapy, non-limiting examples of drugs and agents that may be used to induce said tumor cell-death and internal vaccination are conventional chemotherapeutic agents, cell-cycle inhibitors, anti-angiogenesis drugs, monoclonal antibodies, apoptosis-inducing agents, and signal transduction inhibitors.

[0555] Examples of other anti-cancer agents, which may be relevant as therapeutic agents for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above are differentiation inducing agents, retinoic acid and retinoic acid analogues (such as all trans retinoic acid, 13-cis retinoic acid and similar agents), vitamin D analogues (such as seocalcitol and similar agents), inhibitors of ErbB3, ErbB4, IGF-IR, insulin receptor, PDGFR α , PDGFR β , Flk2, Flt4, FGFR1, FGFR2, FGFR3, FGFR4, TRKA, TRKC, c-met, Ron, Sea, Tie, Tie2, Eph, Ret, Ros, Alk, LTK, PTK7 and similar agents.

[0556] Examples of other anti-cancer agents, which may be relevant as therapeutic agents for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above are cathepsin B, modulators of cathepsin D dehydrogenase activity, glutathione-S-transferase (such as glutacylcysteine synthetase and lactate dehydrogenase), and similar agents.

[0557] Examples of other anti-cancer agents, which may be relevant as therapeutic agents for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above are estramustine and epirubicin.

[0558] Examples of other anti-cancer agents, which may be relevant as therapeutic agents for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above are a HSP90 inhibitor like 17-allyl amino geldanamycin, antibodies directed against a tumor antigen such as PSA, CA125, KSA, etc., integrins like integrin β 1, inhibitors of VCAM and similar agents

[0559] Examples of other anti-cancer agents, which may be relevant as therapeutic agents for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above are calcineurin-inhibitors (such as valspodar, PSC 833 and other MDR-1 or p-glycoprotein inhibitors), TOR-inhibitors (such as sirolimus, everolimus and rapamycin). and inhibitors of "lymphocyte homing" mechanisms (such as FTY720), and agents with effects on cell signaling such as adhesion molecule inhibitors (for instance anti-LFA, etc.).

[0560] In one embodiment, the present invention provides an antibody of the invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and radiotherapy to a subject in need thereof.

[0561] In one embodiment, the present invention provides an antibody of the invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody and radiotherapy to a subject in need thereof.

[0562] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with radiotherapy for treating multiple myeloma.

[0563] Radiotherapy may comprise radiation or associated administration of radiopharmaceuticals to a patient is provided. The source of radiation may be either external or internal to the patient being treated (radiation treatment may, for example, be in the form of external beam radiation therapy (EBRT), brachytherapy (BT) or skeletal targeted radiotherapy). Radioactive elements that may be used in practicing such methods include, e.g., radium, cesium-137, iridium-192, americium-241, gold-198, cobalt-57, copper-67, technetium-99, iodide-123, iodide-131, and indium-111.

[0564] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody to a subject in need thereof combined with autologous peripheral stem cell or bone marrow transplantation.

[0565] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody to a subject in need thereof combined with autologous peripheral stem cell or bone marrow transplantation.

[0566] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with autologous peripheral stem cell or bone marrow transplantation for treating multiple myeloma.

[0567] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody to a subject in need thereof combined with orthopedic intervention.

[0568] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with autologous peripheral stem cell or bone marrow transplantation for treating multiple myeloma.

[0569] Orthopedic interventions may be used in the treatment of a disorder involving cells expressing CD38, such as multiple myeloma, to help control pain or retain function or mobility. Such interventions may include physical therapy, splinting of bones to prevent or treat fractures, or surgical procedures (minor or major) to repair fractures.

[0570] In one embodiment, a CD38 antibody of the present invention may be administered in connection with the delivery of one or more agents that promote access of the antibody or combination composition to the interior of a tumor. Such methods may for example be performed in association with the delivery of a relaxin, which is capable of relaxing a tumor (see for instance US 6,719,977). In one embodiment, a CD38 antibody of the present invention may be bonded to a cell penetrating peptide (CPP). Cell penetrating peptides and related peptides (such as engineered cell penetrating antibodies) are described in for instance Zhao et al., J Immunol Methods. 254(1-2), 137-45 (2001), Hong et al., Cancer Res. 60(23), 6551-6 (2000), Lindgren et al., Biochem J. 377(Pt 1), 69-76 (2004), Buerger et al., J Cancer Res Clin Oncol. 129(12), 669-75 (2003), Pooga et al., FASEB J. 12(1), 67-77 (1998) and Tseng et al., Mol Pharmacol. 62(4), 864-72 (2002).

[0571] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and at least one anti-inflammatory agent to a subject in need thereof.

[0572] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody and at least one anti-inflammatory agent to a subject in need thereof.

[0573] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with at least one anti-inflammatory agent for treating multiple myeloma.

[0574] In one embodiment such an anti-inflammatory agent may be selected from a steroidal drug and a NSAID (nonsteroidal anti-inflammatory drug).

[0575] In one embodiment such an anti-inflammatory agent may be selected from aspirin and other salicylates, Cox-2 inhibitors (such as rofecoxib and celecoxib), NSAIDs (such as ibuprofen, fenoprofen, naproxen, sulindac, diclofenac, piroxicam, ketoprofen, diflunisal, nabumetone, etodolac, oxaprozin, and indomethacin), anti-IL6R antibodies, anti-IL8 antibodies, anti-IL15 antibodies, anti-IL15R antibodies, anti-CD4 antibodies, anti-CD11a antibodies (e.g., efalizumab), anti- α -4/ β -1 integrin (V_L4) antibodies (e.g. natalizumab), CTLA4-Ig for the treatment of inflammatory diseases, prednisolone, prednisone, disease modifying antirheumatic drugs (DMARDs) such as methotrexate, hydroxychloroquine, sulfasalazine, pyrimidine synthesis inhibitors (such as leflunomide), IL-1 receptor blocking agents (such as anakinra), TNF- α blocking agents (such as etanercept, infliximab, and adalimumab) and similar agents.

[0576] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and at least one immunosuppressive and/or immunomodulatory agent to a subject in need thereof.

[0577] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody and at least one immunosuppressive and/or immunomodulatory agent to a subject in need thereof.

[0578] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with at least one immunosuppressive and/or immunomodulatory agent for treating multiple myeloma.

[0579] In one embodiment, such an immunosuppressive and/or immunomodulatory agent may be selected from cyclosporine, azathioprine, mycophenolic acid, mycophenolate mofetil, corticosteroids such as prednisone, methotrexate, gold salts, sulfasalazine, antimalarials, brequinar, leflunomide, mizoribine, 15-deoxyspergualine, 6-mercaptopurine, cy-

clophosphamide, rapamycin, tacrolimus (FK-506), OKT3, anti-thymocyte globulin, thymopentin, thymosin- α and similar agents.

[0580] In one embodiment, such an immunosuppressive and/or immunomodulatory agent may be selected from immunosuppressive antibodies, such as antibodies binding to p75 of the IL-2 receptor, or antibodies binding to for instance MHC, CD2, CD3, CD4, CD7, CD28, B7, CD40, CD45, IFN γ , TNF- α , IL-4, IL-5, IL-6R, IL-6; IGF, IGFR1, IL-7, IL-8, IL-10, CD11a, or CD58, or antibodies binding to their ligands.

[0581] In one embodiment, such an immunosuppressive and/or immunomodulatory agent may be selected from soluble IL-15R, IL-10, B7 molecules (B7-1, B7-2, variants thereof, and fragments thereof), ICOS, and OX40, an inhibitor of a negative T cell regulator (such as an antibody against CTLA4) and similar agents.

[0582] The CD38BPs of the present invention may be administered in combination with two or more immunosuppressive and/or immunomodulatory agents, such as in combination with prednisone and cyclosporine; prednisone, cyclosporine and azathioprine; or prednisone, cyclosporine and mycophenolate mofetil.

[0583] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating a disorder involving cells expressing CD38 in a subject, which method comprises administration of a therapeutically effective amount of the antibody and an anti-C3b(i) antibody to a subject in need thereof

[0584] In one embodiment, the present invention provides an antibody of the present invention for use in a method for treating multiple myeloma, which method comprises administration of a therapeutically effective amount of the antibody and an anti-C3b(i) antibody to a subject in need thereof.

[0585] In one embodiment, antibodies of the present invention may be used for the preparation of a pharmaceutical composition to be administered with an anti-C3b(i) antibody for treating multiple myeloma.

[0586] In one embodiment, a therapeutic agent for use in combination with the CD38 antibodies of the present invention for treating the disorders as described above may be selected from histone deacetylase inhibitors (for instance phenylbutyrate) and/or DNA repair agents (for instance DNA repair enzymes and related compositions such as dimericine).

[0587] Methods involving antibodies of the present invention for treating a disorder as described above comprising administration of a therapeutically effective amount of a CD38 antibody of the present invention may also comprise anti-cancer directed photodynamic therapy (for instance anti-cancer laser therapy - which optionally may be practiced with the use of photosensitizing agent, see, for instance Zhang et al., J Control Release. 93(2), 141-50 (2003)), anti-cancer sound-wave and shock-wave therapies (see for instance Kambe et al., Hum Cell. 10(1), 87-94 (1997)), and/or anti-cancer nutraceutical therapy (see for instance Roudebush et al., Vet Clin North Am Small Anim Pract. 34(1), 249-69, viii (2004) and Rafi, Nutrition. 20(1), 78-82 (2004). Likewise, a CD38 antibody of the present invention may be used for the preparation of a pharmaceutical composition for treating a disorder as described above to be administered with anti-cancer directed photodynamic therapy (for instance anti-cancer laser therapy - which optionally may be practiced with the use of photosensitizing agent, anti-cancer sound-wave and shock-wave therapies, and/or anti-cancer nutraceutical therapy.

[0588] As described above, a pharmaceutical composition of the present invention may be administered in combination therapy, i.e., combined with one or more agents relevant for the disease or condition to be treated either as separate pharmaceutical compositions or with a compound of the present invention coformulated with one or more additional therapeutic agents as described above. Such combination therapies may require lower dosages of the compound of the present invention and/or the co-administered agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

[0589] In one embodiment, the present invention provides a CD38 antibody that is conjugated to an immunomodulator, such as an immunomodulating cytokine, stem cell growth factor, lymphotoxin (such as a TNF such as TNF α), or a hematopoietic factor. Examples of such molecules that may be useful as conjugates include IL-1, IL-2, IL-3, IL-6, IL-10, IL-12, IL-18, and IL-21, colony stimulating factors (such as granulocyte-colony stimulating factor (G-CSF) and granulocyte macrophage-colony stimulating factor (GM-CSF)), interferons (such as IFN α , IFN β , and IFN γ) the stem cell growth factor designated "S1 factor," erythropoietin, and thrombopoietin, active fragments thereof, derivatives thereof, variants thereof, or a combination of any thereof.

[0590] In one embodiment, the CD38 antibodies of the present invention may be used *in vivo* or *in vitro* for diagnosing diseases wherein activated cells expressing CD38 play an active role in the pathogenesis by detecting levels of CD38, or levels of cells which contain CD38 on their membrane surface. This may be achieved, for example, by contacting a sample to be tested, optionally along with a control sample, with the antibody under conditions that allow for formation of a complex between the antibody and CD38. Complex formation is then detected (e.g., using an ELISA). When using a control sample along with the test sample, complex is detected in both samples and any statistically significant difference in the formation of complexes between the samples is indicative of the presence of CD38 in the test sample.

[0591] More specifically, antibodies of the present invention provide methods for the identification of, and diagnosis of invasive cells and tissues, and other cells targeted by CD38 antibodies of the present invention, and for the monitoring of the progress of therapeutic treatments, status after treatment, risk of developing cancer, cancer progression, and the like.

[0592] In one example of such a diagnostic assay, antibodies of the present invention provide a method of diagnosing the level of invasive cells in a tissue comprising forming an immunocomplex between a CD38 antibody and potential CD38 containing tissues, and detecting formation of the immunocomplex, wherein the formation of the immunocomplex correlates with the presence of invasive cells in the tissue. The contacting may be performed *in vivo*, using labeled isolated antibodies and standard imaging techniques, or may be performed *in vitro* on tissue samples.

[0593] CD38BPs, such as CD38 antibodies, may be used to detect CD38-containing peptides and peptide fragments in any suitable biological sample by any suitable technique. Examples of conventional immunoassays provided by antibodies of the present invention include, without limitation, an ELISA, an RIA, FACS assays, plasmon resonance assays, chromatographic assays, tissue immunohistochemistry, Western blot, and/or immunoprecipitation using a CD38BP. Anti-CD38 antibodies of the present invention may be used to detect CD38 and CD38-fragments from humans. Suitable labels for the CD38BP and/or secondary antibodies used in such techniques include, without limitation, various enzymes, prosthetic groups, fluorescent materials, luminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , and ^3H .

[0594] CD38BPs, such as CD38 antibodies, may also be assayed in a biological sample by a competition immunoassay utilizing CD38 peptide standards labeled with a detectable substance and an unlabeled CD38BP, such as an unlabeled anti-CD38 antibody, for example. In such an assay, the biological sample, the labeled CD38 peptide standard(s) and the CD38BP are combined and the amount of labeled CD38 standard bound to the unlabeled CD38BP is determined. The amount of CD38 peptide in the biological sample is inversely proportional to the amount of labeled CD38 standard bound to the CD38BP.

[0595] The CD38 antibodies are particularly useful in the *in vivo* imaging of tumors. *In vivo* imaging of tumors associated with CD38 may be performed by any suitable technique. For example, ^{99}Tc -labeling or labeling with another gamma-ray emitting isotope may be used to label anti-CD38 antibodies in tumors or secondary labeled (e.g., FITC labeled) CD38 antibody:CD38 complexes from tumors and imaged with a gamma scintillation camera (e.g., an Elscint Apex 409ECT device), typically using low-energy, high resolution collimator or a low-energy all-purpose collimator. Stained tissues may then be assessed for radioactivity counting as an indicator of the amount of CD38-associated peptides in the tumor. The images obtained by the use of such techniques may be used to assess biodistribution of CD38 in a patient, mammal, or tissue, for example in the context of using CD38 or CD38-fragments as a biomarker for the presence of invasive cancer cells. Variations on this technique may include the use of magnetic resonance imaging (MRI) to improve imaging over gamma camera techniques. Similar immunoscintigraphy methods and principles are described in, e.g., Srivastava (ed.), Radiolabeled Monoclonal Antibodies For Imaging And Therapy (Plenum Press 1988), Chase, "Medical Applications of Radioisotopes," in Remington's Pharmaceutical Sciences, 18th Edition, Gennaro et al., (eds.), pp. 624-652 (Mack Publishing Co., 1990), and Brown, "Clinical Use of Monoclonal Antibodies," in Biotechnology And Pharmacy 227-49, Pezzuto et al., (eds.) (Chapman & Hall 1993). Such images may also be used for targeted delivery of other anti-cancer agents, examples of which are described herein (e.g., apoptotic agents, toxins, or CHOP chemotherapy compositions). Moreover, such images may also or alternatively serve as the basis for surgical techniques to remove tumors. Furthermore, such *in vivo* imaging techniques may allow for the identification and localization of a tumor in a situation where a patient is identified as having a tumor (due to the presence of other biomarkers, metastases, etc.), but the tumor cannot be identified by traditional analytical techniques. All of these methods are features in accordance with the present invention.

[0596] The *in vivo* imaging and other diagnostic methods utilizing the antibodies of the present invention are particularly useful in the detection of micrometastases in a human patient (e.g., a patient not previously diagnosed with cancer or a patient in a period of recovery/remission from a cancer). Carcinoma cancer cells, which may make up to 90% of all cancer cells, for example, have been demonstrated to stain very well with anti-CD38 antibody conjugate compositions. Detection with monoclonal anti-CD38 antibodies and other CD38BPs described herein may be indicative of the presence of carcinomas that are aggressive/invasive and also or alternatively provide an indication of the feasibility of using related monoclonal anti-CD38 antibody, CD38BP, or related composition treatments against such micrometastases. Moreover, monoclonal anti-CD38 antibodies that are associated with cancer cells are advantageously able to distinguish such cancer-associated tissues and cells from normal cells that other forms of CD38 may be associated with.

[0597] In an *in vivo* imaging method, a CD38BP, such as an anti-CD38 antibody of the present invention, may be conjugated to a detection-promoting radio-opaque agent, the conjugated antibody is administered to a host, such as by injection into the bloodstream, and the presence and location of the labeled antibody in the host is assayed. Through this technique and any other diagnostic method described herein, provided herein is a method for screening for the presence of disease-related cells in a human patient or a biological sample taken from a human patient.

[0598] For diagnostic imaging, radioisotopes may be bound to a CD38BP, such as a CD38 antibody, either directly,

or indirectly by using an intermediary functional group. Useful intermediary functional groups include chelators, such as ethylenediaminetetraacetic acid and diethylenetriaminepentaacetic acid (see for instance US 5,057,313). In such diagnostic assays involving radioisotope-conjugated CD38BPs, the dosage of conjugated peptide delivered to the patient typically is maintained at as low a level as possible through the choice of isotope for the best combination of minimum half-life, minimum retention in the body, and minimum quantity of isotope, which will permit detection and accurate measurement.

[0599] In addition to radioisotopes and radio-opaque agents, diagnostic methods may be performed using CD38BPs, such as CD38 antibodies, that are conjugated to dyes (such as with the biotin-streptavidin complex), contrast agents, fluorescent compounds or molecules and enhancing agents (e.g. paramagnetic ions) for magnetic resonance imaging (MRI) (see, e.g., US Pat. No. 6,331,175, which describes MRI techniques and the preparation of antibodies conjugated to a MRI enhancing agent). Such diagnostic/detection agents may be selected from agents for use in magnetic resonance imaging, and fluorescent compounds. In order to load a CD38BP, such as an antibody, component with radioactive metals or paramagnetic ions, it may be necessary to react it with a reagent having a long tail to which are attached a multiplicity of chelating groups for binding the ions. Such a tail may be a polymer such as a polylysine, polysaccharide, or other derivatized or derivatizable chain having pendant groups to which can be bound chelating groups such as, e.g., porphyrins, polyamines, crown ethers, bithiosemicarbazones, polyoximes, and like groups known to be useful for this purpose. Chelates may be coupled to CD38BPs using standard chemistries. A chelate is normally linked to a CD38BP, such as an anti-CD38 mAB, by a group, which enables formation of a bond to the molecule with minimal loss of immunoreactivity and minimal aggregation and/or internal cross-linking. Other, more unusual, methods and reagents for conjugating chelates to antibodies are disclosed in for instance US 4,824,659. Examples of potentially useful metal-chelate combinations include 2-benzyl-DTPA and its monomethyl and cyclohexyl analogs, used with diagnostic isotopes in the general energy range of 60 to 4,000 keV, such as ^{125}I , ^{123}I , ^{124}I , ^{62}Cu , ^{64}Cu , ^{18}F , ^{111}In , ^{67}Ga , ^{99}Tc , ^{94}Tc , ^{11}C , ^{13}N , ^{15}O , and ^{76}Br , for radio-imaging. These and similar chelates, when complexed with non-radioactive metals, such as manganese, iron, and gadolinium may be useful for MRI diagnostic methods in connection with CD38BPs. Macrocyclic chelates such as NOTA, DOTA, and TETA are of use with a variety of metals and radiometals, most particularly with radionuclides of gallium, yttrium, and copper, respectively. Such metal-chelate complexes may be made very stable by tailoring the ring size to the metal of interest. Other ring-type chelates such as macrocyclic polyethers, which are of interest for stably binding nuclides, such as ^{223}Ra for RAIT may also be suitable in diagnostic methods.

[0600] Thus, provided herein are diagnostic CD38 antibody conjugates, wherein the CD38 antibody is conjugated to a contrast agent (such as for magnetic resonance imaging, computed tomography, or ultrasound contrast-enhancing agent) or a radionuclide that may be, for example, a gamma-, beta-, alpha-, Auger electron-, or positron-emitting isotope. Additional useful conjugated CD38BPs are described elsewhere herein, which may also be useful in diagnostic methods and compositions (e.g., diagnostic kits) provided by the present disclosure.

[0601] Described herein is a kit for diagnosis of cancer comprising a container comprising a CD38BP, such as an anti-CD38 antibody, and one or more reagents for detecting binding of the CD38BP to a CD38 peptide. Reagents may include, for example, fluorescent tags, enzymatic tags, or other detectable tags. The reagents may also include secondary or tertiary antibodies or reagents for enzymatic reactions, wherein the enzymatic reactions produce a product that can be visualized. In one embodiment, provided herein is a diagnostic kit comprising one or more CD38BPs, such as anti-CD38 antibodies of the present invention, in labeled or unlabeled form in suitable container(s), reagents for the incubations for an indirect assay, and substrates or derivatizing agents for detection in such an assay, depending on the nature of the label. Control reagent(s) and instructions for use also may be included.

[0602] Diagnostic kits may also be supplied for use with a CD38BP, such as a conjugated/labeled anti-CD38 antibody, for the detection of a cellular activity or for detecting the presence of CD38 peptides in a tissue sample or host. In such diagnostic kits, as well as in kits for therapeutic uses described elsewhere herein, a CD38BP typically may be provided in a lyophilized form in a container, either alone or in conjunction with additional antibodies specific for a target cell or peptide. Typically, a pharmaceutical acceptable carrier (e.g., an inert diluent) and/or components thereof, such as a Tris, phosphate, or carbonate buffer, stabilizers, preservatives, biocides, biocides, inert proteins, e.g., serum albumin, or the like, also are included (typically in a separate container for mixing) and additional reagents (also typically in separate container(s)). In certain kits, a secondary antibody capable of binding to the anti-CD38 antibody or other CD38BP, which typically is present in a separate container, is also included. The second antibody is typically conjugated to a label and formulated in manner similar to the anti-CD38 antibody of the present invention.

[0603] Using the methods described above and elsewhere herein CD38BPs, such as antibodies, may be used to define subsets of cancer/tumor cells and characterize such cells and related tissues/growths.

[0604] In one example, a CD38BP or anti-CD38 antibody, may be added to nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles, or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled CD38 peptide or antibody. The solid phase support may then be washed with the buffer a second time to remove unbound peptide or antibody. The amount of bound label on the solid support may then be detected by known method steps.

[0605] Linked enzymes that react with an exposed substrate may be used to generate a chemical moiety which can be detected, for example, by spectrophotometric, fluorometric or by visual means, in the context of a CD38BP conjugate and/or fusion protein. Enzymes which may be used to detectably label CD38BPs such as anti-CD38 antibodies include malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholinesterase. It is also possible to label a CD38BP, such as a CD38 antibody, with a fluorescent compound. When the fluorescent labeled antibody is exposed to light of the proper wave length, its presence may be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde, and fluorescamine.

[0606] The CD38BPs, such as anti-CD38 antibodies, may also be detectably labeled using fluorescence-emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals may be attached to an anti-CD38 antibody, for example, using such metal chelating groups as diethylenetriaminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

[0607] CD38BPs, such as anti-CD38 antibodies, may also be detectably labeled by coupling to a chemiluminescent compound. The presence of the chemiluminescently labeled CD38BP is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, thionin, acridinium ester, imidazole, acridinium salt, and oxalate ester.

[0608] Likewise, a bioluminescent compound may be used to label a CD38BP. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase, and aequorin.

[0609] Detection of a labeled peptide or antibody, antibody fragment or derivative may be accomplished by a scintillation counter, for example, if the detectable label is a radioactive gamma emitter, or by a fluorometer, for example, if the label is a fluorescent material. In the case of an enzyme label, the detection may be accomplished by colorimetric methods which employ a substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

[0610] These and other diagnostic techniques may be used to screen any suitable material for CD38 peptides or CD38-fragments. Examples of materials that may be screened include, for example, blood, serum, lymph, urine, inflammatory exudate, cerebrospinal fluid, amniotic fluid, a tissue extract or homogenate, and the like. However, the present disclosure is not limited to assays using only these samples, it being possible for one of ordinary skill in the art to determine suitable conditions which allow the use of other samples.

[0611] *In situ* detection may be accomplished by removing a histological specimen from a patient, and providing the combination of labeled CD38BPs, such as anti-CD38 antibodies, of the present invention to such a specimen. The CD38BP, anti-CD38-antibody (including fragment) of the present invention may be provided by applying or by overlaying the labeled CD38BP, such as a labeled anti-CD38 antibody, of the present invention to a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of CD38 or CD38-fragment but also the distribution of such peptides in the examined tissue (e.g., in the context of assessing the spread of cancer cells). Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) may be modified in order to achieve such *in situ* detection.

[0612] The present disclosure further includes methods of promoting the sale and/or use of a CD38BP as described herein comprising distributing information (such as by printed materials that are handed out, mailed, etc., by advertising signage, by television programs and advertisements, by radio programs and advertisements, by internet site postings, by email, by telemarketing, by door-to-door or person-to-person marketing, by funding and/or hosting conferences, panels, forums, etc., by employing and/or contracting for the services of salespeople and/or medical/scientific liaisons, by funding and/or hosting scientific research and publications related to such uses, etc.) related to the use of the compound in the prevention or treatment of any condition or combination of conditions as described elsewhere herein to any persons or entities of potential interest (such as pharmaceutical chains, formulary managers, insurance companies, HMOs, hospitals and hospital chains, other health care companies, pharmacy benefit managers, potential patients, cancer patients, former cancer patients, patients in remission, primary care physicians, nurses, doctors of pharmacy, and/or key opinion leaders).

[0613] The present disclosure also includes kits comprising a pharmaceutical composition of a compound of the present invention and instructions for use. The kit may further contain one or more additional agents, such as an immunosuppressive reagent, a chemotherapeutic reagent, an anti-inflammatory agent or a radiotoxic agent as described above, or one or more additional CD38BPs as described herein (such as a CD38BP having a complementary activity). A kit as described herein may also include diagnostic agents and/or other therapeutic agents. In one embodiment, a kit includes a CD38BP as described herein and a diagnostic agent that may be used in a diagnostic method for diagnosing the state or existence of a disorder involving cells expressing CD38 in a subject. In one embodiment, the kit includes a

CD38BP as described herein in a highly stable form (such as in a lyophilized form) in combination with pharmaceutically acceptable carrier(s) that may be mixed with the highly stable composition to form an injectable composition.

[0614] All headings and sub-headings are used herein for convenience only and should not be construed as limiting the present invention in any way.

[0615] The use of the terms "a" and "an" and "the" and similar referents in the context of describing the present invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context.

[0616] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. Unless otherwise stated, all exact values provided herein are representative of corresponding approximate values (e.g., all exact exemplary values provided with respect to a particular factor or measurement can be considered to also provide a corresponding approximate measurement, modified by "about," where appropriate).

[0617] All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context.

[0618] The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended merely to better illuminate the present invention and does not pose a limitation on the scope of the present invention unless otherwise indicated. No language in the specification should be construed as indicating any element is essential to the practice of the present invention unless as much is explicitly stated.

[0619] The citation and incorporation of patent documents herein is done for convenience only and does not reflect any view of the validity, patentability, and/or enforceability of such patent documents.

[0620] The description herein of any embodiment of the present invention using terms such as "comprising", "having," "including," or "containing" with reference to an element or elements is intended to provide support for a similar embodiment of the present invention that "consists of", "consists essentially of", or "substantially comprises" that particular element or elements, unless otherwise stated or clearly contradicted by context (e.g., a composition described herein as comprising a particular element should be understood as also describing a composition consisting of that element, unless otherwise stated or clearly contradicted by context).

[0621] The present invention is further illustrated by the following examples which should not be construed as further limiting.

EXAMPLES

EXAMPLE 1

Manufacturing luciferase-transfected (Daudi-luc) cells

[0622] Culture of Daudi cells (originating from Burkitt's lymphoma) was cultured in RPMI 1640 culture medium supplemented with 10% FCS (Optimum C241, Wisent Inc., St. Bruno, QC, Canada), 2 mM L-glutamine, 100 IU/ml penicillin, 100 mg/ml streptomycin, 1 mM sodium pyruvate (all derived from Gibco BRL, Life Technologies, Paisley, Scotland). Medium was refreshed twice a week. Before transfection, cells were split and seeded out at $1-1.5 \times 10^6$ cells/ml to ensure viability and optimal growth.

Luciferase transfection

[0623] 8.2×10^6 CD38⁺ Daudi cells were taken up in 350 μ l RPMI (supplemented with 10% dFCS, Gibco BRL) and transferred to an electroporation cuvet (Biorad, Hemel Hempstead, Herts, UK). Then, 40 μ g gWIZ luciferase from GTS (Aldevron, Fargo, ND, USA) and 10 μ g pPur vector (BD Biosciences, Alphen a/d Rijn, The Netherlands), which confers puromycin resistance, were added. After resting cells on ice for 10 minutes, cells were electroporated (250 V, 950 μ F; Gene Pulser II, Biorad Laboratories GmbH, München, Germany). Cells were again rested on ice, and taken up in 40 ml RPMI (supplemented with 10% FCS). Then, cells were plated out in 96-well tissue culture plates (100 μ l per well). After 48 hours, puromycin (final concentration: 1 μ g/ml; Sigma-Aldrich Chemie BV, Zwijndrecht, The Netherlands) was added. Puromycin-resistant clones were further grown in 24-well tissue culture plates.

Determination of luciferase activity

[0624] Luciferase activity of cells was determined using the Luciferase Assay System (#E4030, Promega, Madison, WI, USA). 1×10^5 cells were centrifuged (13.500 rpm, 1 min) in an eppendorf centrifuge, and the pellet was washed in 100 μ l PBS. After centrifugation (13.500 rpm, 1 min), pellet was lysed with 20 μ l Reporter Lysis Buffer (Promega), frozen

and thawed. After centrifugation (13,500 rpm, 1 min), 20 μ l supernatant was discarded, and 100 μ l luciferase assay reagent was added (in special luminometer tubes, Promega). Luminescence was measured (10 sec) in a luminometer (LB9507, Berthold, Vilvoorde, Belgium).

EXAMPLE 2

Immunization of mice and generation of hybridomas

Immunization protocol for -003

[0625] HCo12 mice were immunized every fortnight with 20 μ g purified HA-CD38. The first immunization was performed i.p. in the presence of 100 μ l PBS, mixed with 100 μ l Complete Freund's Adjuvant (CFA). After this first immunization, subsequent boosts (13x) with purified HA-CD38 were performed in the presence of 100 μ l PBS, mixed with 100 μ l incomplete Freund's Adjuvant (IFA) alternating s.c. and i.p. After titer development, mice were boosted with 20 μ g HA-CD38 in PBS, i.v.

Immunization protocol for -005 and -024

[0626] HCo12 mice were immunized every fortnight with 20 μ g purified HA-CD38 alternating with NIH-3T3-CD38 transfected cells. The first immunization was performed with 5×10^6 cells in 100 μ l PBS, mixed with 100 μ l CFA, i.p., the second and following immunizations with HA-CD38 s.c., in the presence of 100 μ l PBS, mixed with 100 μ l IFA. The following immunizations with transfected cells were performed in the presence of 200 μ l PBS. After titer development, mice were boosted with 20 μ g HA-CD38 in PBS, i.v.

Generation of Hybridomas Producing Human Monoclonal Antibodies to CD38

[0627] The mouse splenocytes were isolated from HCo12 mice and fused with PEG to a mouse myeloma cell line based upon standard protocols. The resulting hybridomas were then screened for human antibody production by ELISA and for CD38 specificity using human CD38-transfected NS/0 cells by FACS analysis and recombinant HA-CD38 protein binding by ELISA. Three hybridoma cell lines were selected expressing the human monoclonal anti-CD38 antibodies, -003, -005 and -024, respectively.

EXAMPLE 3

Transfection of NIH cells with CD38

[0628] The vector (pCIPuroCD38) for producing NIH-3T3-CD38 cells was obtained from Prof. M. Glennie (Tenovus Research Laboratory, Southampton General Hospital, Southampton, UK). NIH-3T3 cells (DSMZ, ACC 59; 150,000 cells/well; 0.5 ml; 96-well flat-bottom plates, Greiner) were cultured in DMEM (supplemented with glucose [4.5 g/l], 10% FCS, L-glutamine, Na-pyruvate; BioWhittaker) for 24 h. Then, DNA (0.8 μ g) and lipofectamine (Invitrogen, Breda, The Netherlands) were diluted in DMEM, and mixed (20 min, RT). Thereafter, the mixture (100 μ l) was added to each well and incubated (ON, 37°C).

Screening for CD38 expression

[0629] NIH-3T3-CD38 cells were washed (in 1 ml PBS) and trypsinized (200 μ l, trypsin-EDTA, BioWhittaker). Then, 1 ml of DMEM was added and the mixture pipetted into FACS tubes. After centrifugation (1200 rpm, 5 min), cells were washed in FACS Buffer (FB; PBS, 0.05% BSA, 0.02% NaN_3) and resuspended in 1 ml FB. After centrifugation (1200 rpm, 5 min), supernatant was removed and mouse anti-human CD38-PE was added (1/50 dilution, Sanquin, Amsterdam, The Netherlands). After washing the cells twice in FB, cells were resuspended in FB for acquisition by flow cytometry.

Expansion and selection

[0630] After trypsin treatment, cells were transferred to T25 flasks (Greiner) in DMEM (supplemented with glucose 4.5 g/l, 2 mM L-glutamine, and puromycin (2 μ g/ml) BioWhittaker). Puromycin-resistant cells were tested for stable CD38 expression by flow cytometry after two weeks on puromycin-containing medium. NIH-3T3-CD38 selected cells were subcloned by limiting dilution. After expanding these cells, all 15 NIH-3T3-CD38 clones were screened for CD38 expression. CD38high NIH-3T3-CD38 cells were frozen in liquid nitrogen (-80°C) until use.

Culture of NIH-3T3-CD38 cells

[0631] Cells are cultured in DMEM (supplemented with glucose (4.5 g/l), 10% FCS, 2 mM L-glutamine, Na-pyruvate, penicillin, streptomycin). Cells are passaged twice a week by use of trypsin/EDTA and seeded in a concentration of 1×10^6 cells/T75 flask. CD38^{high} NIH-3T3-CD38 cells were frozen in liquid nitrogen (-80°C) until use.

Purification of HA-CD38 antigen

[0632] Sepharose 4B (Amersham Bioscience, Uppsala, Sweden) was coupled with anti-CD38 antibody (Serotec, Oxford, UK). Column (column tube HR5/20 was packed to 12 cm bedheight, column volume 2.4 ml; maximum flow rate 0.5 ml/min) was equilibrated with at least 5 column volumes (CV) of PBS. Sample was filtrated and loaded to the column. Column was washed with PBS until signal returned to baseline (approximately 3 CV). Elution was carried out with 0.1 M glycine at pH 2. Eluted fractions were neutralized with 1% (v/v) 2 M Tris-HCl, pH 9.

Purification of anti-CD38 antibodies

[0633] Human anti-CD38 antibodies were purified from tissue culture supernatants. First, the supernatants were filtered over 0.20 μM dead-end filter. Then, the supernatant was loaded on a 5 ml Protein A column (rProtein A FF, Amersham Bioscience) and eluted with 0.1 M citric acid-NaOH, pH 3. The eluate was immediately neutralized with 2 M Tris-HCl, pH 9 and dialyzed O/N to 12.6 mM sodium phosphate, 140 mM NaCl, pH 7.4 (B. Braun, Oss, The Netherlands). After dialysis samples were sterile filtered over 0.20 μM dead-end filter.

Purification of His-CD38 batches

[0634] The protein is present in cell culture supernatant of His-CD38-expressing cells, with a DNA construct containing the sequence for the extracellular domain of CD38. An additional poly-His-tag sequence is included in the constructs and present at the N-terminus of the protein. This tag enables purification with immobilized metal affinity chromatography. In this process, a chelator fixed onto the chromatographic resin is charged with Co^{2+} cations. Particularly, a sequence that includes 6 histidine amino acids strongly binds Co^{2+} . Therefore the His-tagged CD38 proteins bind strongly to such a column, while other proteins present in the culture supernatant will flow through the column or will be washed away. The strongly bound His-tagged CD38 proteins are then eluted with a buffer containing imidazole, which competes with the binding of His to Co^{2+} . When sufficient His-CD38 is purified, the eluent is removed from the protein by buffer exchange on a desalting column.

EXAMPLE 4Binding of -003, -005, and -024 to CD38-transfected CHO (CHO-CD38) cells, to Daudi-luc cells and to fresh multiple myeloma (MM) tumor cells

[0635] After harvesting and counting, Daudi-luc cells, CHO cells transfected with CD38 and control CHO cells were resuspended in PBS (1×10^6 cells/ml). Then, cells were put in 96-well V-bottom plates (100 μl /well) and washed twice in PBS-BSA (PBS supplemented with 0.1 % BSA and 0.02% Na-azide). Thereafter, 50 μl antibody solution in PBS-BSA was added to the cells (4°C , 30 min). After washing three times in PBS-BSA, 50 μl (1:400 dilution) of rabbit anti-human IgG-FITC in PBS-BSA was added (4°C in the dark, 30 min). Cells were washed three times and specific binding of CD38-antibodies to CHO-CD38 and Daudi-luc cells was detected by flow cytometry. HuMab-KLH (a human monoclonal antibody against KLH (keyhole limpet haemocyanin) generated by Genmab B.V., Utrecht, The Netherlands by use of the immunization protocols described elsewhere herein) was used as a control. Figures 1 and 2 show that -003, -005, and -024 bind to CHO-CD38 cells and to Daudi-luc cells, albeit with different EC_{50} (Table 1). No binding to control CHO cells is observed (data not shown).

[0636] Fresh MM tumor cells were obtained from Dr. Lokhorst (University Medical Center Utrecht, Utrecht, The Netherlands). Tumor cells were isolated from bonemarrow of multiple myeloma patients by Ficoll (Bio Whittaker; lymphocyte separation medium, cat 17-829E) gradient centrifugation. After harvesting and counting, MM cells (100,000 cells/well) were resuspended with 25 μl FITC-labeled CD38-specific antibodies and 25 μl CD138. After incubation (4°C , 30 min), cells were washed in PBS-BSA and PE-labeled goat-anti-mouse IgG (1:200; Jackson ImmunoResearch Europe Ltd. Soham, UK) was added. After incubation (4°C , 30 min) and washing of the cells in PBS-BSA, fluorescence was measured by flow cytometry.

[0637] Figure 3 shows that -003, -005 and -024 bind to MM cells.

Table 1 - EC₅₀ values of binding of anti CD38-antibodies on CHO-CD38 cells, Daudi-luc cells and fresh MM tumor cells.

CD38-specific antibodies	EC ₅₀ CHO-CD38 (μg/ml)	EC ₅₀ Daudi-luc (μg/ml)	EC ₅₀ MM cells (μg/ml)
-003	0.54	0.26	0.56
-005	0.23	0.09	0.04
-024	0.08	0.05	0.02

EXAMPLE 5Antibody-dependent cell-mediated cytotoxicity

[0638] Daudi-luc cells, fresh multiple myeloma tumor cells, fresh Plasma Cell Leukemia tumor cells and JK6L and AMO-1 multiple myeloma cells were collected (5x10⁶ cells) in RPMI⁺⁺ (RPMI 1640 culture medium supplemented with 10% cosmic calf serum (HyClone, Logan, UT, USA)), to which 100 μCi ⁵¹Cr (Chromium-51; Amersham Biosciences Europe GmbH, Roosendaal, The Netherlands) was added, and the mixture was incubated in a 37°C water bath for 1 hr. After washing of the cells (twice in PBS, 1500 rpm, 5 min), the cells were resuspended in RPMI⁺⁺ and counted by trypan blue exclusion. Cells were brought at concentration of 1x10⁵ cells/ml.

Preparation of effector cells

[0639] Fresh peripheral blood mononuclear cells (healthy volunteers, UMC Utrecht, Utrecht, The Netherlands) were isolated from 40 ml of heparin blood by Ficoll (Bio Whittaker; lymphocyte separation medium, cat 17-829E) according to the manufacturer's instructions. After resuspension of cells in RPMI⁺⁺, cells were counted by trypan blue exclusion and brought at concentration of 1x10⁷ cells/ml.

ADCC set up

[0640] 50 μl of ⁵¹Cr-labeled targets cells were pipetted into 96-well plates, and 50 μl of antibody was added, diluted in RPMI⁺⁺ (final concentrations 10, 1, 0.1, 0.01 μg/ml). Cells were incubated (RT, 15 min), and 50 μl effector cells were added, resulting in an effector to target ratio of 100:1 (for determination of maximal lysis, 100 μl 5% Triton-X100 was added instead of effector cells; for determination of spontaneous lysis, 50 μl target cells and 100 μl RPMI⁺⁺ were used). Cells were spun down (500 rpm, 5 min), and incubated (37°C, 5% CO₂, 4 hr). After spinning down cells (1500 rpm, 5 min), 100 μl of supernatant was harvested into micronic tubes, and counted in gamma counter. The percentage specific lysis was calculated as follows:

$$(\text{cpm sample} - \text{cpm target cells only}) / (\text{cpm maximal lysis} - \text{cpm target cells only}) \text{ wherein cpm is counts per minute.}$$

[0641] In Daudi-luc cells (Figure 4 and Table 2) -003, -005, and -024 induce lysis by ADCC, and -003, and -005 perform slightly better than rituximab (anti-CD20 mAb). Interestingly, also when fresh multiple myeloma tumor cells (obtained from Dr. H. Lokhorst, UMCU, The Netherlands) are used as target cells, ADCC is induced by -003, -005 and -024 (Figure 5A and Table 2).

Table 2 - EC₅₀ values of CD38-specific antibodies obtained in ADCC

CD38-specific antibodies	EC ₅₀ Daudi-luc (ng/ml)	EC ₅₀ MM cells (ng/ml)
-003	9.0	27
-005	4.5	5.7
-024	9.7	56

Enrichment of human peripheral blood mononuclear cells Erlangen

[0642] Human blood from human volunteers (university Erlangen, Erlangen, Germany) was diluted twice in RPMI 1640 and blood cells were layered on Ficoll (Lymphocyte Separation Medium 1077 g/ml, 710 g, RT, 20 min; BioWhittaker, Cambrex Bio Science Verviers, Verviers, Belgium, cat. 17-829E, lot no. 0148 32). Peripheral blood mononuclear cells (MNCs) were collected from the interphase, washed and resuspended in RPMI 1640 culture medium supplemented with 10% FCS, 2 mM L-glutamine, 5 U/ml penicillin, 50 μg/ml streptomycin (all derived from BioWhittaker) to which 25

mM HEPES (BioWhittaker) was added.

ADCC set up II

[0643] Target B-cells (fresh plasma cell leukemia tumor cells, JK6L and AMO-1 B-cell lines, obtained from Dr. T. Valerius, University of Erlangen, Erlangen, Germany) were labeled with 20 μCi ^{51}Cr (Amersham Biosciences, Uppsala, Sweden) for 2 hours. After extensive washing in RPMI-10, cells were adjusted to 1×10^5 cells/ml. MNCs (50 μl), sensitizing antibodies (50 μl), and RPMI-10 (50 μl) were added to round-bottom microtiter plates (Greiner Bio-One GmbH, Frickhausen, Germany). Assays were started by adding fresh plasma cell leukemia tumor cells, JK6L or AMO-1 cells (50 μl) giving a final volume of 200 μl . An effector to target (E:T) ratio of 40:1 was used. After incubation (3 hr, 37°C), assays were stopped by centrifugation, and ^{51}Cr release from triplicates was measured in counts per minute (cpm) in a scintillation counter. Percentage of cellular cytotoxicity was calculated using the following formula:

$$\% \text{ specific lysis} = (\text{experimental cpm} - \text{basal cpm}) / (\text{maximal cpm} - \text{basal cpm}) \times 100$$

with maximal ^{51}Cr release determined by adding perchloric acid (3% final concentration) to target cells, and basal release was measured in the absence of sensitizing antibodies and effector cells.

[0644] In both multiple myeloma cell lines (i.e. JK6L and AMO-1), lysis is induced with both -003 and -005 (Figures 6 and 7), even when CD38 expression is low (AMO-1 cell line).

[0645] -003, -005 and -024 induce ADCC of plasma cell leukemia primary tumor cells (Figure 5B).

EXAMPLE 6

Complement-dependent cytotoxicity

[0646] After harvesting and counting of Daudi-luc cells, the viability of the cells should be $\geq 90\%$. After washing (PBS), cells are resuspended at 2.0×10^6 cells/ml in RPMI-B (RPMI supplemented with 1% BSA). Thereafter, cells are put in 96-well round-bottom plates at 1×10^5 cells/well (50 μl /well). Then, 50 μl antibodies is added to the wells (final concentration range between 0-100 $\mu\text{g/ml}$ (three-fold dilutions in RPMI-B)). After incubation (RT, 15 min), 11 μl of pooled human serum (pool of 18 healthy donors) was added to each well (37°C, 45 min). Wells were resuspended once and 120 μl was transferred to FACS tubes (Greiner). Then, 10 μl propidium iodide (PI; Sigma-Aldrich Chemie B.V.) was added (10 $\mu\text{g/ml}$ solution) to this suspension. Lysis was detected by flow cytometry (FACScalibur™, Becton Dickinson, San Diego, CA, USA) by measurement of the percentage of dead cells (corresponds to PI-positive cells).

[0647] Figure 8 and Table 2 show that lysis of Daudi-luc cells is induced by -005 (~60% maximum lysis) and that lysis by -003 is only seen at very high antibody concentrations. -024 does not induce CDC in Daudi cells (data not shown). In CHO-CD38 cells, lysis is induced by both -003, -005, and -024 (Figure 9 and Table 3). Lysis by -003 is induced at higher concentrations. In tumor cells (all obtained from Dr. Lokhorst and Dr. Bloem, University Medical Center Utrecht, The Netherlands), obtained from different MM patients (A: 3% refractory tumor cells, B: 9% refractory tumor cells, C: 30-40% tumor cells, and D: 70% tumor cells), CDC-mediated lysis is observed in the presence of -005, but not in the presence of -003 (Figure 10). -024 also induced lysis of MM tumor cells (Figure 10E).

Table 3 - EC₅₀ values of CD38-specific antibodies obtained in CDC

CD38-specific antibodies	EC ₅₀ Daudi-luc ($\mu\text{g/ml}$)	EC ₅₀ CD38-CHO ($\mu\text{g/ml}$)
-003	>90	3.14
-005	0.33	0.14
-024	>90	0.24

EXAMPLE 7

Cross-block studies using FACS

[0648] CHO-CD38 cells were incubated with an excess of unlabelled CD38-specific antibody (4°C, 15 min). Then, cells were incubated with FITC-labeled CD38-specific antibodies (concentration approximates EC₉₀, 4°C, 45 min). After twice washing the cells with PBS-BSA, fluorescence was measured by flow cytometry. Figure 11 shows that unlabelled -003 blocks binding of FITC-labeled -003, whereas binding of FITC-labeled -005 is not blocked. Also unlabelled -005 blocks binding of FITC-labeled -005, whereas binding of FITC-labeled -003 is not blocked. -003 and -005 bind to different

epitopes, because they do not compete for binding.

EXAMPLE 8

Cross-blocking studies using ELISA

[0649] Soluble human CD38 is coated on the surface of an ELISA plate. Coated CD38 is incubated with an excess of unlabelled CD38 specific antibodies for about 15 minutes and subsequently biotinylated CD38-specific antibodies are added (concentration approximates EC_{90} , RT, 1 hour). After washing three times with PBS/Tween, horseradish peroxidase (HRP)-conjugated streptavidine is added and the mixture is incubated for 1 hour at RT. The complex can be detected by addition of an ABTS-solution and the HRP mediated substrate conversion is measured using an ELISA reader at OD 405 nm.

EXAMPLE 9

Cross-blocking studies using sandwich-ELISA

[0650] CD38 specific antibodies are coated on the surface of an ELISA plate. Plate-bound antibodies are incubated with biotinylated soluble CD38 in the presence of an excess of CD38 specific antibodies in fluid phase. After washing with PBS/Tween, bound biotinylated CD38 is detected with HRP-conjugated streptavidine for 1 hr at RT. This complex can be detected by addition of an ABTS-solution (after washing with PBS/Tween) and the HRP mediated substrate conversion is measured using an ELISA reader at OD 405 nm.

EXAMPLE 10

Reactivity with a panel of human tissues and cross-reactivity with cynomolgus tissue by immunohistochemistry

[0651] Sections from frozen human tissue (obtained from Dr. H. Niessen, Free University Medical Center, Amsterdam, The Netherlands) or monkey tissue (Inveresk Research, Glasgow, Scotland) were cut at 6 μ m and air-dried overnight. These cryostat sections were fixated in acetone (RT, 10 min) and air-dried (approx. 5 min). Thereafter, sections were incubated with 1x citric acid/phosphate buffer containing 0.1% H_2O_2 (pH 5.8; Sigma), to block endogenous peroxidase. After 20 min at RT, sections were washed twice with PBS and 0.05% Tween-20 (PBST, RT, 5 min; Riedel de-Haen, Germany). Then, sections were incubated with avidin (RT, 15 min; DAKO, Glostrup, Denmark), washed twice with PBST, and incubated with biotin (RT, 15 min; DAKO) to block endogenous biotin. After washing the sections twice with PBST, sections were preincubated with PBST⁺⁺ (PBST supplemented with 10% normal human serum (NHS, CLB, Amsterdam, Netherlands) and 10% normal goat serum (NGS; DAKO) (RT, 20 min). After blotting-off of the pre-incubation serum, sections were incubated with FITC-labeled primary antibody diluted in 2% PBST⁺⁺ at the indicated concentrations (RT, 60 min). Thereafter, sections were incubated with rabbit-anti-FITC (1:1000; DAKO) in 2% PBST⁺⁺ (RT, 30 min). After washing the sections with PBST, sections were incubated with goat-anti-rabbit-biotin (1:400; DAKO) in 2% PBST⁺⁺ (RT, 30 min). Then, sections were washed and incubated with SABC-HRP (1:100; DAKO) in 2% PBST⁺⁺ (RT, 30 min). After washing the sections twice in PBST, they were incubated (RT, 10 min) with amino-ethyl-carbazole (AEC)-development solution (50 mM acetate buffer, pH4.9, 0.01 % H_2O_2 ; Riedel-de-Haen). Finally, sections were washed in millipore H_2O (5 min) and counterstained with hematoxylin (DAKO). By use of glycergel (37°C), sections were fixed with cover slips, and studied by light microscopy (Axiovision-2; Zeiss, Thornwood, NY, USA).

[0652] Bronchial epithelium is stained with -003 and -005 (Figures 12B and 13B) as well as striated muscle (myocytes, Figures 12C and 13C), macrophages, lymphocytes and plasma B cells (Figures 12A and 13A). -024 has a similar staining of striated muscle and bronchial epithelium, but staining was less intense. No staining of endothelial cells is observed, neither with -003 (Fig 14D), -005 (14E) nor -024 (data not shown), whereas clear staining was observed with the positive control antibodies against endothelial cell markers CD31 (Fig 14A) and vWF (14B). Anti-KLH was used as negative control antibody (Fig 14C). -003 (Figure 12D) and -024 (data not shown) but not -005 (Figure 13D) cross-react with cynomolgus monkey lymphoid tissue.

EXAMPLE 11

Cross-reactivity with cynomolgus or rhesus monkey peripheral blood mononuclear cells (PBMCs) by flow cytometry

[0653] 5 ml of cynomolgus monkey peripheral blood (Inveresk Research) were lysed by adding 4.5 ml shock buffer (1.7 mM NH_4CL , 1 mM EDTA), 40 ml H_2O and 450 μ l 10% $KHCO_3$. After hemolysis cells were centrifuged (1200 rpm,

10 min) and washed thrice in PBS. After counting cells with trypan blue, cells were resuspended in PBS-BSA (1x10⁶ cell/ml).

[0654] 17.5 ml of rhesus monkey peripheral blood (BPRC, Rijswijk, The Netherlands) was diluted 1:1 with RPMI 1640 and layered on Ficoll (1.077 g/ml; BioWhittaker, cat. 17-829E, lot no. 0148 32). After centrifugation (710 g, RT, 20 min), the interphase was collected and washed twice in RPMI. After the last wash cells were resuspended in RPMI 1640 at a concentration of 1x10⁵ cells/50 µl.

[0655] Cells were transferred to 96-well plate (100,000 PBMCs/well), washed in FACS buffer (PBS, 0.05% BSA, 0.02% NaN₃) and incubated with the primary antibodies (4°C, 30 min). After washing in PBS-BSA, 50 µl FITC-labeled rb-anti-hlgG (DAKO, Glostrup, Denmark) was added (4°C, 30 min). Finally, cells were collected in FACS tubes in a total volume of 150 µl. Samples were measured and analyzed by use of FACScalibur™ (Becton Dickinson, San Diego, CA, USA).

[0656] With flow cytometry cross-reactivity of -003 on cynomolgus lymphocytes (Figure 15A) and monocytes (Figure 15B) was shown, but not of -005. Also in rhesus monkeys, cross-reactivity of -003 was observed on mononuclear cells, but not of -005 (Figure 15C).

EXAMPLE 12

Internalization experiments

[0657] CHO-CD38 cells were stained with a saturating concentration of FITC-labeled CD38-specific antibodies (on ice, 30 min). After washing of cells (in RPMI1640 supplemented with 10% FCS), one cell pool was warmed up to 37°C to allow internalization, and the other pool was left on ice. At several time intervals (0-120 min) cell aliquots were taken and transferred to ice-cold PBS-BSA to stop internalization. After washing samples twice with PBS-BSA, EtBr (diluted in PBS-BSA, final concentration 2 mg/ml) was added to the samples to quench membrane-bound FITC. Fluorescence was measured by flow cytometry.

[0658] Figure 16A and 16B show that -003 and -005 are internalized by CHO-CD38 cells within 5 minutes at 37°C.

EXAMPLE 13

In vivo SCID-luciferase experiments

[0659] In this model tumor cells are transfected with firefly luciferase. Upon administration of luciferin (Molecular Probes, Leiden, The Netherlands) to the mice the labeled cells can be detected in vivo by bioluminescent imaging using a highly sensitive CCD camera, cf. Wetterwald et al., American Journal of Pathology 160(3), 1143-1153 (2002).

[0660] Daudi cells were transfected with gWIZ luciferase from Gene Therapy Systems (San Diego, CA) and cultured in RPMI with 10% FCS, Pen/Strep, Sodium Pyruvate and 1 µg/ml puromycin (Sigma). Cells were analysed for luciferase expression (expressed in RLU/1 x 10⁵ cells) in a luminometer and for CD38 expression by FACS. 2.5 x 10⁶ luciferase-transfected Daudi cells/mouse were injected i.v. into SCID mice. Mice were treated with -003, -005, isotype control antibody (HuMab-KLH) or rituximab (anti-CD20 antibody). Antibodies were injected intraperitoneally. Four treatment settings were used (see Table 4). In the preventive setting, antibody (100 µg/mouse) and cells were administered simultaneously. In therapeutic setting I, antibody (300 µg/mouse) was administered 7 days after administration of cells. In therapeutic setting II, antibody (10 µg/mouse) was administered 14 days after administration of cells. In therapeutic setting III, antibody (100 µg/mouse) was administered 7 days after administration of cells. For imaging, mice were anesthetized by i.p. injection of a mixture of ketamine/xylazine /atropine. Synthetic D-Luciferin (sodium salt, Molecular Probes) was given i.p. at a dose of 25 mg/ml. Mice were then placed in a light tight box and after 3 min, imaging was started using a VersArray 1300B liquid nitrogen cooled CCD detector (Roper Scientific). Photons emitted from the luciferase were counted over an exposure period of 5 min. Under illumination black and white images were made for reference. MetaVue software (Universal Imaging Corp) was used for data collection and image analysis. Statistical significance of differences between groups was established using one-way analysis of variance with a Newman-Keuls post test using GraphPad PRISM version 3.02 (Graphpad Software Inc).

Table 4 - Treatment settings for in vivo luciferase experiments

Experimental setting	Antibody treatment (days after cell inoculation)	Antibody dose (µg/mouse)
Preventive setting	0	100
Therapeutic setting	7	300
Therapeutic setting II	14	10
Therapeutic setting III	7	100

[0661] Figure 17A and 17B show that -003 and -005 inhibit growth of tumor cells in the preventive setting and in therapeutic setting I, similar to the inhibition observed for the anti-CD20 antibody. Both antibodies perform significantly better than the isotype control antibody. Also in therapeutic setting II CD38-antibodies slow down the growth of Daudi-luc tumor cells (Figure 17C). In therapeutic setting III, -003 and -024 show a clear inhibition of Daudi-luc tumor cell growth (Figure 17D).

EXAMPLE 14

Apoptosis

[0662] Apoptosis assay was carried out according to the manufacturer's instructions (Annexin-V Apoptosis kit, BD Biosciences, Alphen a.d. Rijn, Netherlands). In short, CD38 mAbs were added to 2.5×10^5 cells (luciferase-transfected Daudi cells, in 0.5 ml RPMI⁺⁺ in a 24-wells plate), in a concentration of 5 $\mu\text{g/ml}$ -003 or -005 or an anti-CD20 antibodies alone or in the presence of cross-blocking rb-anti-hlgG (50 $\mu\text{g/ml}$).

[0663] After incubation (37°C, 5% CO₂, 20 hr), cells were harvested carefully, and washed with Binding Buffer (1200 rpm, 4°C, 5 min, BD Biosciences). Pellet was resuspended in 100 μl Binding Buffer. Then, 5 μl Annexin-V-FITC (BD Biosciences) and 10 μl PI (BD Biosciences) was added to the suspension and incubated for 15 minutes at RT. 400 μl Binding Buffer was added and the samples were measured (PI readout in FL2). For analysis of apoptotic cells, all Annexin-V-positive cells were counted by flow cytometry using a FACScalibur flow cytometer with CellQuest pro software (BD Biosciences). At least 10,000 events were collected for analysis. This population includes both PI-positive as well as PI-negative cells.

[0664] Figure 18 shows that -003 and -005 do not induce apoptosis. However, after cross-linking, apoptosis of target cells is observed. -003 induced apoptosis after cross-linking that was similar to apoptosis induced by an anti-CD20 antibody (rituximab). -005 was less able to induce apoptosis after cross-linking. Similar results were obtained with RAMOS cells as target cells (data not shown).

EXAMPLE 15

Effect of -005 on tissue graft B cells in RA-SCID mouse model

Implantation of synovial tissue

[0665] SCID-mice, strain C.B.-17/lcrCrl-SCID-bg, male / female, 4-12 weeks, purchased from Charles River Laboratories Nederland (Maastricht, the Netherlands) were kept in IVC cages under standard conditions of temperature and light, and were fed laboratory chow and water ad libitum. Prior to implantation, mice (three mice in each experimental group, day 0) were anesthetized by intraperitoneal injection of ketamine (NIMATEK, EuroVet) and xylazine (Rompun, Bayer) at ratio 1:1. A small incision of the skin was made using surgical scissors. Inflamed synovial tissue from a patient with rheumatoid arthritis undergoing joint replacement surgery was implanted subcutaneously as a cluster of six small fragments (total 2-3 mm³) on each flank of the mouse. The wound was closed using Permacol cyanoacrylate glue. On day 1 of the experiment, remaining synovial tissue was analyzed in order to check for B cells in the inflamed synovial transplants. -005 (12 mg/kg) or control antibody (anti-KLH, 30 mg/kg) was injected (i.v.), in a volume of 200 μl on day 8 of the experiment. At the end of the experiment (day 14) mice were sacrificed by CO₂ inhalation and the synovial grafts were explanted. One of the grafts was snap-frozen in OCT compound (TissueTek, Sacura Finetek Europe) for further immunohistochemical analysis, and another one was frozen by immersion in liquid nitrogen for further RNA analysis.

Immunohistochemistry

[0666] 5 μM cryosections on SuperFrost (Menzel GmbH, Braunschweig) slides were prepared using LEICA CM1900 cryostat and stored at -80°C. Thawed sections were fixed in acetone for 10 min, dried at room temperature and washed 3 x 5 min in PBS. All steps were performed at room temperature. Endogenous peroxidase activity was blocked by incubation with PBS supplemented with 0.3% hydrogen peroxide and 0.1% sodium azide for 20 min. Slides were washed 3 x 5 min in PBS and incubated with 10% normal human serum (NHS) / 10% normal rabbit serum (NRbS) in PBS / 1% BSA for 30 min. Next, primary antibody (mouse mAb) diluted in PBS supplemented with 1% BSA / 10% NHS / 10% NRbS was incubated for 60 min. After 3 x 2 min washes in PBS, HRP-conjugate (goat anti-mouse Ig-HRP; DAKO P0447) diluted 1:50 in PBS (supplemented with 1% BSA / 10% NHS / 10% NRbS) was added for 30 min. Peroxidase signal was enhanced using TSATM Biotin system (Perkin Elmer Life Sciences, NEL700). Slides were washed 3 x 2 min in PBS and incubated with biotinyl tyramide diluted 1:1600 in amplification buffer for 30 min. After 3 x 2 min washes in PBS, streptavidin-HRP diluted 1:400 in PBS (supplemented with 1% BSA) was added for 30 min. Slides were washed 3 x 2

min in PBS and incubated with DAB solution (DAKO Cytomation K3465) for 5 min. Color reaction was stopped with distilled water. Finally, slides were counterstained with hematoxyline (MERCK), washed with running water and covered with Kaiser's glycerin and cover slips.

Scoring of staining intensity

[0667] Scoring of stained synovial tissue xenografts was performed in a blinded fashion by two trained persons. First the strongest section was selected from a series of sections and this reference section was awarded the maximum score 8. The staining intensity in the other sections was then scored on a scale of 0 to 8, relative to the reference section.

Statistical analysis

[0668] Scoring of staining intensity was analyzed by Kruskal-Wallis one-way ANOVA followed by Dunn's multiple comparison test using Graph Pad Prism version 4.01 (Graph Pad software, Inc., San Diego, CA, USA).

[0669] Figure 19 and Figure 21 show that the numbers of anti-CD38-positive plasma cells are reduced after treatment with -005. Staining of plasma cells with anti-CD138 confirms that -005 results in reduced numbers of plasma cells (Figures 20 and 22).

EXAMPLE 16

Sequencing of the coding sequence of human antibodies against CD38

RNA preparation

[0670] Total RNA was prepared from 5×10^6 cells of the hybridoma cell lines expressing the monoclonal antibody -003, -005 and -024, respectively, with the RNeasy kit (Qiagen, Westburg, Leusden, Netherlands) according to the manufacturer's protocol.

cDNA preparation of -003, -005 and -024

[0671] 5'-RACE-Complementary DNA (cDNA) of RNA was prepared from 100 ng total RNA, using the SMART RACE cDNA Amplification kit (Clontech), following the manufacturer's protocol.

[0672] Oligonucleotide primers were synthesized and quantified by Isogen Bioscience (Maarssen, The Netherlands). Primers were dissolved in H_2O to 100 pmol/ μl and stored at $-20^\circ C$. A summary of all PCR and sequencing primers is tabulated (Table 5). For PCR, PfuTurbo® Hotstart DNA polymerase (Stratagene, Amsterdam, The Netherlands; product# 600322) was used according to the manufacturer's instructions. Each reaction mix contained 200 μM mixed dNTPs (Roche Diagnostics, Almere, The Netherlands; product# 1814362), 12 pmol of the reverse primer (RACEG1A1 for V_H 3003-005, RACEV_HApal for V_H 3003-003 and RACEV_LBsiWI for V_L 3003-003 and 005), 7.2 pmol UPM-Mix (UPM-Mix: 2 μM ShortUPMH3 and 0.4 μM LongUPMH3), 0.6 μl of the 5'RACE cDNA template, and 1.5 unit of PfuTurbo® Hotstart DNA polymerase in PCR reaction buffer (supplied with polymerase) in a total volume of 30 μl . PCR reactions were carried out with a TGradient Thermocycler 96 (Whatman Biometra, Goettingen, Germany; product# 050-801) using a 35-cycle program: denaturing at $95^\circ C$ for 2 min; 35 cycles of $95^\circ C$ for 30 sec, a $55^\circ C$ for 30 sec, and $72^\circ C$ for 1.5 min; final extension at $72^\circ C$ for 10 min. If appropriate, the PCR mixes were stored at $4^\circ C$ until further analysis or processing.

Table 5 - Primers

Name	Sequence
ShortUPMH3	TGAAAGCTTCTAATACGACTCACTATAGGGC
RACEV _L BsiWI	GAAGATGAAGACAGATGGTGCAGCCACCGTACG
RACEV _H Apal	GGAGGGTGCCAGGGGGAAGACCGATGGGCCCTT
RACEG1A1	GGGAGTAGAGTCCTGAGGACTG
M13reverse	GGATAACAATTTACACAGG
LongUPMH3	TGAAAGCTTCTAATACGACTCACTATAGGGCAAGCAGTGG TATCAACGCAGAGT
HCseq5	GGTCAGGGCGCCTGAGTTCCACG

(continued)

Name	Sequence
VH3003-003for	GATAAGCTTGCCGCCACCATGGACTGGACCTGGAGGTTC CTC
VH3003-5for	GATAAGCTTGCCGCCACCATGGAGTTTGGGCTGAGCTGG CTT
VL3003-5exfor	GATAAGCTTGCCGCCACCATGGAAGCCCCAGCTCAGCTT CTC
VL3003-003for	GATAAGCTTGCCGCCACCATGAGGGTCCTCGCTCAGCTC CTG
VH300324exfor	GATAAGCTTGCCGCCACCATGGGGTCAACCGCCATCCTC GCC
VL3003-24-5exfor	GATAAGCTTGCCGCCACCATGGAAGCCCCAGCTCAGCTT CTC

Cloning of -003-2F5 V_H and V_L and -005 V_L and -024 V_H and V_L in pGEMT-Vector System II

[0673] The reaction products were separated by electrophoresis on a 1% TAE agarose gel and stained with ethidium bromide. Bands of the correct size were cut from the gels and the DNA was isolated from the agarose using the Qiaexll gel extraction kit (Qiagen, cat no 20021).

[0674] Gel isolated PCR fragments were A tailed by a 10 min 72°C incubation with 200 µM dATP and 2.5 units Amplitaq (Perkin Elmer) and purified using minielute columns (Qiagen). A-tailed PCR fragments were cloned into the pGEMTeasy vector (Promega) using the pGEMT easy vector system II kit and protocol (LJ270, page 3/4). 2 µl of the ligation mixture was transformed into OneShot DH5αT1R competent E.Coli (Invitrogen) and plated on LB/Amp/IPTG/Xgal plates.

Sequencing

[0675] The V-regions -003 and -024 and the -005 V_L region were sequenced by AGOWA (Berlin, Germany) after picking respectively 20 (V_H-003), 16 (V_L-003), 15 (V_L-005) and 6 (V_H and V_L -024) white colonies, isolating plasmid and sequencing with the M13 reverse primer. The -005 V_H region was sequenced directly on the PCR product by using primer HCseq5. Sequences were analyzed using the Vector NTI advanced suite (Invitrogen).

Generation of expression vectors for antibody -003, -005, -024 and Morphosys antibody 3079

[0676] The V_H coding region of -003 was amplified by PCR from a pGemT plasmid clone containing the V_H region of -003, using the primers VH3003-003for and RACEVHApal, introducing suitable restriction sites (HindIII and Apal) for cloning into pConG1f0.4 (Lonza Biologics, Slough, UK) and an ideal Kozak sequence (GCCGCCACC). The pConG1f0.4 vector contains the heavy chain constant region of human IgG1. The V_H PCR fragment was inserted, in frame, into the pConG1f0.4 vector using HindIII and Apal. The construct was checked by sequence analysis.

[0677] The V_H coding region of -005 was amplified by PCR from a pGemT plasmid clone containing the V_H region of -005, using the primers VH3003-5for and RACEVHApal, introducing suitable restriction sites (HindIII and Apal) for cloning into pConG1f0.4 and an ideal Kozak sequence. The V_H PCR fragment was inserted, in frame, into the pConG1f0.4 vector using HindIII and Apal. The construct was checked by sequence analysis.

[0678] The V_H coding region of -024 was amplified by PCR from a pGemT plasmid clone containing the V_H region of -024, using the primers VH300324exfor and RACEVHApal, introducing suitable restriction sites (HindIII and Apal) for cloning into pConG1f0.4 and an ideal Kozak sequence. The V_H PCR fragment was inserted, in frame, into the pConG1f0.4 vector using HindIII and Apal. The construct was checked by sequence analysis.

[0679] The V_H coding region of Morphosys antibody 3079 was synthesized by GeneArt (Regensburg, Germany), based on the data published in patent WO 2005/103083 A2. The coding region was codon optimized for expression in HEK cells to enhance expression levels and suitable restriction sites (HindIII and Apal) for cloning into pConG1f0.4 and an ideal Kozak sequence were introduced. The plasmid containing the synthetic V_H region was digested with Apal and HindIII and the V_H fragment was inserted, in frame, into the pConG1f0.4 vector.

[0680] The V_L coding region of -005 was amplified by PCR from a pGemT plasmid clone containing the V_L region of -005, using the primers VL3003-5exfor and RACEVLBSiWI, introducing suitable restriction sites (HindIII and Pfl23II) for cloning into pConKappa0.4 (Lonza Biologies) and an ideal Kozak sequence. The pConKappa0.4 vector contains the kappa light chain constant region. The V_L PCR fragment was inserted, in frame, into the pConKappa0.4 vector using HindIII and Pfl23II. The construct was checked by sequence analysis.

[0681] The V_L coding region of -003 was amplified by PCR from a pGemT plasmid clone containing the V_L region of -003, using the primers VL3003-003for and RACEVLBSiWI, introducing suitable restriction sites (HindIII and Pfl23II) for cloning into pConKappa0.4 and an ideal Kozak sequence. The V_L PCR fragment was inserted, in frame, into the pConKappa0.4 vector using HindIII and Pfl23II. The construct was checked by sequence analysis.

[0682] The V_L coding region of -024 was amplified by PCR from a pGemT plasmid clone containing the V_L region of -024, using the primers VL3003-24-5exfor and RACEVLBSiWI, introducing suitable restriction sites (HindIII and Pfl23II) for cloning into pConKappa0.4 and an ideal Kozak sequence. The V_L PCR fragment was inserted, in frame, into the pConKappa0.4 vector using HindIII and Pfl23II. The construct was checked by sequence analysis.

[0683] The V_L coding region of Morphosys antibody 3079 was synthesized by GeneArt, based on the data published in WO 2005/103083. The coding region was codon optimized for expression in HEK cells; to enhance expression levels and suitable restriction sites (HindIII and Pfl23II) for cloning into pConKappa0.4 and an ideal Kozak sequence were introduced. The plasmid, containing the synthetic V_L region, was digested with Pfl23II and HindIII and the VH fragment was inserted, in frame, into the pConKappa0.4 vector.

[0684] Antibodies were transiently expressed in HEK-293F cells, as described in Example 17, by cotransfecting their heavy chain and light chain vectors.

Generation of stable cell lines in CHO-K1SV cells

[0685] For generation of stable cell lines, the heavy and light chain vectors of -003 or -005 were combined in a single double gene vector by standard cloning techniques.

[0686] The double gene vectors of -003 or -005 were linearized and transfected into CHO-K1SV (Lonza Biologies) cells, essentially as described by the manufacturer. Stable cell lines were selected by selection with 25 μM L-Methionine sulfoximine (MSX) as described by Lonza Biologies. Top producing clones were selected and propagated in CD-CHO (Invitrogen) medium and antibodies were purified from cell culture supernatant as described in Example 3.

EXAMPLE 17

Epitope mapping using site directed mutagenesis

[0687] Oligonucleotide primers were synthesized and quantified by Isogen Bioscience (Maarssen, The Netherlands). Primers were dissolved in H₂O to 100 pmol/μl and stored at -20°C. A summary of all PCR and sequencing primers is shown in Table 6. For PCR, PfuTurbo® Hotstart DNA polymerase (Stratagene, Amsterdam, The Netherlands) was used according to the manufacturer's instructions. Each reaction mix contained 200 μM mixed dNTPs (Roche Diagnostics, Almere, The Netherlands), 10 pmol of both the forward and reverse primer, 100 ng of genomic DNA or 1 ng of plasmid DNA and 1 unit of PfuTurbo® Hotstart DNA polymerase in PCR reaction buffer (supplied with polymerase) in a total volume of 20 μl. PCR reactions were carried out with a TGradient Thermocycler 96 (Whatman Biometra, Goettingen, Germany) using a 32-cycle program: denaturing at 95°C for 2 min; 30 cycles of 95°C for 30 sec, a 60-70°C gradient (or another specific annealing temperature) for 30 sec, and 72°C for 3 min; final extension at 72°C for 10 min. If appropriate, the PCR mixtures were stored at 4°C until further analysis or processing.

[0688] Agarose gel electrophoresis was performed according to Sambrook (Sambrook, Russell et al. 2000) using gels of 50 ml, in 1 x Tris Acetate EDTA buffer. DNA was visualized by the inclusion of ethidium bromide in the gel and observation under UV light. Gel images were recorded by a CCD camera and an image analysis system (GeneGnome; Syngene, via Westburg B.V., Leusden, The Netherlands).

[0689] Purification of desired PCR fragments was carried out using a MinElute PCR Purification Kit (Qiagen, via Westburg, Leusden, The Netherlands; product# 28006), according to the manufacturer's instructions. Isolated DNA was quantified by UV spectroscopy (see below) and the quality was assessed by agarose gel electrophoresis.

[0690] Alternatively, PCR or digestion products were separated by agarose gel electrophoresis (for instance when multiple fragments were present) using a 1% Tris Acetate EDTA agarose gel. The desired fragment was excised from the gel and recovered using the QIAEX II Gel Extraction Kit (Qiagen; product# 20051), according to the manufacturer's instructions.

[0691] Optical density of nucleic acids was determined using a NanoDrop ND-1000 Spectrophotometer (Isogen Life Science, Maarssen, The Netherlands) according to the manufacturer's instructions. The DNA concentration was measured by analysis of the optical density (OD) at 260 nm (one OD_{260nm} unit = 50 μg/ml). For all samples, the buffer in

which the nucleic acids were dissolved was used as a reference.

[0692] Restriction enzymes and supplements were obtained from New England Biolabs (Beverly, MA, USA) or Fermentas (Vilnius, Lithuania) and used according to the manufacturer's instructions. DNA (100 ng) was digested with 5 units of enzyme(s) in the appropriate buffer in a final volume of 10 μ l (reaction volumes were scaled up as appropriate). Digestions were incubated at the recommended temperature for a minimum of 60 min. For fragments requiring double digestions with restriction enzymes which involve incompatible buffers or temperature requirements, digestions were performed sequentially. If necessary digestion products were purified by agarose gel electrophoresis and gel extraction.

[0693] Ligations of DNA fragments were performed with the Quick Ligation Kit (New England Biolabs) according to the manufacturer's instructions. For each ligation, vector DNA was mixed with approximately three-fold molar excess of insert DNA.

[0694] Plasmid DNA (1-5 μ l of DNA solution, typically 2 μ l of DNA ligation mix) was transformed into One Shot DH5 α -T1^R E. coli cells (Invitrogen, Breda, The Netherlands; product# 12297-016) using the heat-shock method, according to the manufacturer's instructions. Next, cells were plated on Luria-Bertani (LB) agar plates containing 50 μ g/ml ampicillin. Plates were incubated for 16-18 h at 37°C until bacterial colonies became evident.

[0695] Bacterial colonies were screened for the presence of vectors containing the desired sequences via colony PCR using the ThermoStart PCR Master Mix (Abgene, via Wetsburg, Leusden, The Netherlands; product# AB-938-DC15/b) and primers pConG1seq1 and pEE13.4seqrev2 (Table 6). Selected colonies were lightly touched with a 20 μ l pipette tip and touched briefly in 2 ml LB for small scale culture, and then resuspended in the PCR mix. PCR was performed with a TGradient Thermocycler 96 using a 35-cycle program: denaturation at 95°C for 15 min; 35 cycles of 94°C for 30 sec, 55°C for 30 sec and 72°C for 2 min; followed by a final extension step of 10 min at 72°C. If appropriate, the PCR mixtures were stored at 4°C until analysis by agarose gel electrophoresis.

[0696] Plasmid DNA was isolated from E. coli cultures using the following kits from Qiagen (via Westburg, Leusden, The Netherlands), according to the manufacturer's instructions. For bulk plasmid preparation (50-150 ml culture), either a HiSpeed Plasmid Maxi Kit (product# 12663) or a HiSpeed Plasmid Midi Kit (product# 12643) was used. For small scale plasmid preparation (\pm 2 ml culture) a Qiaprep Spin Miniprep Kit (product# 27106) was used and DNA was eluted in 50 μ l elution buffer (supplied with kit).

Construction of HA-CD38 expression vector pEE13.4HACD38

[0697] The extracellular domain of human CD38 was amplified from plasmid pCIPuroCD38 (obtained from Prof. M. Glennie, Tenovus Research Laboratory, Southampton General Hospital, Southampton, UK) using primers cd38forha and cd38exrev. By this PCR reaction an HA-tag was introduced. This PCR product was used as template for a second PCR reaction with primers SPHMM38ex and cd38exrev. By this PCR reaction, signal peptide SPHMM, restriction sites and an ideal Kozak sequence (GCCGCCACC) for optimal expression were introduced. After purification, this PCR fragment was cloned into expression vector pEE13.4 (Lonza Biologics) and the complete coding sequence was confirmed by sequencing with primers pConKseq1, pEE13.4seqrev, cd38seq1for and cd38seq2rev (Table 6). This construct was named pEE13.4HACD38

Site-directed mutagenesis

[0698] Three single mutant proteins of huCD38 was constructed, in which T was mutated to A at position 237 (T237A, SEQ ID No:32), Q was mutated to R at position 272 (Q272R, SEQ ID No:33), or S was mutated to F at position 274 (S274F, SEQ ID No:34). Site-directed mutagenesis was performed using the QuickChange II XL Site-Directed Mutagenesis Kit (Stratagene, Amsterdam, The Netherlands) according to the manufacturer's instructions. This method included the introduction of a silent extra restriction site or loss of a restriction site to screen for successful mutagenesis (extra *Xba*I site for T237A mutant, extra *Bcg*I site for Q272R mutant and loss of *Ssp*I site for S274F mutant). Briefly, 5 μ l 10x reaction buffer, 1 μ l oligonucleotide HACD38T237Afor2, HACD38Q272Rfor or HACD38S274Ffor (100 pmol/ μ l), 1 μ l oligonucleotide HACD38T237Arev2, HACD38Q272Rrev or HACD38S274Frev (100 pmol/ μ l), 1 μ l dNTP mix, 3 μ l Quicksolution, 1 μ l plasmid pEE13.4HACD38 (50 ng/ μ l) and 1 μ l PfuUltra HF DNA polymerase were mixed in a total volume of 50 μ l and amplified with a TGradient Thermocycler 96 (Whatman Biometra, Goettingen, Germany; product# 050-801) using an 18-cycle program: denaturing at 95°C for 1 min; 18 cycles of 95°C for 50 sec, 60°C for 50 sec, and 68°C for 10 min. PCR mixtures were stored at 4°C until further processing. Next, PCR mixtures were incubated with 1 μ l *Dpn*I for 60 min at 37°C to digest the pEE13.4HACD38 WT vector and stored at 4°C until further processing. The reaction mixture was precipitated with 5 μ l 3 M NaAc and 125 μ l ethanol, incubated for 20 minutes at -20°C and spun down for 20 minutes at 4°C at 14000xg. The DNA pellet was washed with 70% ethanol, dried and dissolved in 4 μ l water. The total 4 μ l reaction volume was transformed in One Shot Top 10DH5 α T1^R competent E. coli cells (Invitrogen, Breda, The Netherlands) according to the manufacturer's instructions (Invitrogen). Next, cells were plated on Luria-Bertani (LB) agar plates containing 50 μ g/ml ampicillin. Plates were incubated for 16-18 h at 37°C until bacterial colonies

became evident. Colonies were screened by colony PCR using primers pConG1seq1 and pEE13.4seqrev2 (Table 5) and digested with the relevant restriction enzymes to screen for incorporation of the mutagenic oligonucleotide. 2 positive clones for each mutant were grown and plasmid DNA was isolated. The complete HACD38 coding sequence was determined using primers cd38seq1for, pConG1seq1 and pEE13.4seqrev2 to confirm the presence of the mutations and the absence of additional undesirable mutations.

DNA sequencing

[0699] Plasmid DNA samples were sent to AGOWA (Berlin, Germany) for sequence analysis. Sequences were analyzed using Vector NTI advanced software (Informax, Oxford, UK).

Transient expression in HEK-293F cells

[0700] Freestyle™ 293-F (a HEK-293 subclone adapted to suspension growth and chemically defined Freestyle medium, (HEK-293F)) cells were obtained from Invitrogen and transfected with pEE13.4HACD38 and with the three constructs carrying the mutations T237A, Q272R and S274F, according to the manufacturer's protocol using 293fectin (Invitrogen). Culture supernatants of transfected cells were used in ELISA for anti-CD38 binding studies.

Anti-CD38 antibody binding

[0701] ELISA plates (Greiner, # 655092) were coated O/N at 4°C with 1 µg anti-HA antibody (Sigma, # H-9658) and subsequently blocked with 2% chicken serum. Culture supernatants of transfected HEK293F cells were diluted, applied to the ELISA plates and incubated for 1 hr at RT. After washing, serial dilutions of HuMabs -003 and -005 were added and incubated for 1 hr at RT. Bound antibodies were detected with HRP-conjugated goat-anti-human IgG antibodies. The assay was developed with ABTS (Roche, # 1112597) and the absorbance was measured at 405 nm using a spectrophotometer.

[0702] As can be seen from Figures 23A-23C, both -003 and -005 bind to wt human CD38. The binding of -003 was not affected by the introduction of mutations T237A (Figure 23A), Q272R (Figure 23B) or S274F (Figure 23C). -005 was able to bind CD38 harboring mutation T237A (Figure 23A). Binding of -005 to CD38 with mutation Q272R was severely affected (Figure 23B), both with respect to EC₅₀ and maximum binding capacity. -005 was not able to bind to mutant CD38 wherein serine at position 274 was replaced by phenylalanine (Figure 23C).

[0703] These data shows that -003 and -005 bind to different epitopes. Furthermore these studies revealed that binding of -005 to CD38 is sensitive to mutations at positions 272 and 274. Particularly S274 is essential for -005 binding to CD38.

Table 6 - Primers

Name	Sequence
cd38forha	CTGCTGTGGCCCATGGTGTGGGCCTACCCTTACGACGTGC CTGACTACGCCAGGTGGCGCCAGACGTGGAGC
cd38exrev	AGGTCAGGTACCTCAGATCTCAGATGTGCAAG
SPHMM38ex	TATAGCCCCGGGGCCGCCACCATGTGGTGGCGCCTGTGGTG GCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGGCCCATGGTGTGG GCC
pConG1seq1	GAAGACTTAAGGCAGCGGCAGAA
pConKseq1	GTAGTCTGAGCAGTACTCGTTGC
pEE13.4seqrev	TGCATTTCATTTTATGTTTCAGGT
pEE13.4seqrev2	TCGGACATCTCATGACTTTCTTT
cd38seq1for	AGGACACGCTGCTAGGCTACCTT
cd38seq2rev	GTCCTTTCTCCAGTCTGGGCAAG
HACD38T237Arev2	TCCACCATGTATCACCCAGGCCTCTAGAGCCTGAACCTTCT CTGGTTG
HACD38T237Afor2	CAACCAGAGAAGGTTTCAGGCTCTAGAGGCCTGGGTGATACA TGGTGGA
HACD38Q272Rrev	GATATTCTTGCAGGAAAATCGAATATTCCTTTTGCTTAT
HACD38Q272Rfor	ATAAGCAAAAGGAATATTCGATTTTCCTGCAAGAATATC

(continued)

Name	Sequence
HACD38S274Frev	TCTGTAGATATTCTTGCAGAAAAATTGAATGTTCTTTTGCTT ATA
HACD38S274Ffor	TATAAGCAAAAGGAACATTCAATTTTTCTGCAAGAATATCTAC AGA

EXAMPLE 18Induction of proliferation of PBMC

[0704] -003, -005 and -024 were tested in an assay essentially as described in Ausiello et al., Tissue antigens 56, 538-547 (2000). Briefly, PBMCs from healthy donors were cultured at 1×10^5 cells/well in flat bottom 96-well plates in the presence of antibodies (final concentration: 1.1 - 3.3 - 10 - 30 $\mu\text{g/ml}$) in 200 μl RPMI⁺⁺. Stimulation of cells with IL-15 (at 333 ng/ml; Amgen Inc., Thousand Oaks, CA, USA) was used as positive control. After a 4 day incubation at 37°C, 30 μl ³H-thymidine (16.7 $\mu\text{Ci/ml}$) was added, and culture was continued O/N. ³H-thymidine incorporation was assessed using a Packard Cobra gamma counter (Packard Instruments, Meriden, DT, USA), according to the manufacturer's instructions. Data are shown as the mean cpm (\pm SEM) of PBMCs obtained from 10 donors. The results show that -003 and -005 do not induce significant proliferation of PBMCs (Figure 24A). Also -024 did not induce significant proliferation of PBMCs (data not shown).

EXAMPLE 19Induction of IL-6

[0705] -003, -005 and -024 were tested in an assay as described in Ausiello et al., Tissue antigens 56, 538-547 (2000). Briefly, PBMCs were cultured at 1×10^6 cells/well in 48-well plates in the presence of 20 $\mu\text{g/ml}$ of antibodies and 10 ng/ml LPS (Sigma-Aldrich Chemie, Zwijndrecht, The Netherlands) in 500 μl RPMI⁺⁺. After an O/N incubation at 37°C, supernatant was harvested and stored at -20°C. The IL-6 concentration was assessed by ELISA (IL-6 ELISA kit, U-CyTech Biosciences, Utrecht, The Netherlands) according to the manufacturer's instructions. Data are shown mean concentration in pg/ml (\pm SEM) from 7 donors. The results show that -003 and -005 does not induce release of significant IL-6 levels (Figure 24B). Also -024 did not induce release of significant IL-6 levels (data not shown).

EXAMPLE 20Induction of release of IFN- γ

[0706] -003, -005 and -024 were tested in an assay as described in Ausiello et al., Tissue antigens 56, 538-547 (2000). Briefly, PBMCs were cultured at 1×10^6 cells/well in 48-well plates in the presence of 20 $\mu\text{g/ml}$ of antibodies and 1 $\mu\text{g/ml}$ OKT-3 (Sanquin, Amsterdam, The Netherlands) in 500 μl RPMI⁺⁺. After an O/N incubation at 37°C, supernatant was harvested and stored at -20°C. The IFN- γ concentration was assessed by ELISA (IFN- γ ELISA kit, U-CyTech Biosciences, Utrecht, The Netherlands) according to the manufacturer's instructions. Data are shown mean concentration in pg/ml (\pm SEM) from 9 donors. The results show that -003 and -005 does not induce release of detectable IFN- γ levels (Figure 24C). Also -024 did not induce release of significant IFN- γ levels (data not shown).

EXAMPLE 21Affinity of binding of -003 and -005 to recombinant CD38

[0707] Binding of -003 and -005 to CD38 was tested using surface plasmon resonance. Briefly, purified antibodies were immobilized on a CM-5 sensor chip (Biacore, Uppsala, Sweden) via amine coupling. HA-tagged CD38 (see Example 3) was flowed over, and the binding of antigen to mAb was detected by a change in refractive index at the surface of the chip using a Biacore 3000 (Biacore). The associated and rate constants for -003 (Table 7) and -005 (Table 8) are summarized below, mean of 3 experiments \pm SD, and show that both -003 and -005 have a high affinity for CD38.

Table 7 - Association and rate constants at 25°C**-003**

k_a (1/Ms)	$2.17 \times 10^5 \pm 2.65 \times 10^4$
k_d (1/s)	$1.9 \times 10^{-4} \pm 4.51 \times 10^{-6}$
K_A (1/M)	$1.14 \times 10^9 \pm 1.58 \times 10^8$
K_D (M)	$8.85 \times 10^{-10} \pm 1.2 \times 10^{-10}$

Table 8 - Association and rate constants at 25°C**-005**

k_a (1/Ms)	$8.88 \times 10^4 \pm 1.95 \times 10^4$
k_d (1/s)	$5.22 \times 10^{-4} \pm 1.16 \times 10^{-5}$
K_A (1/M)	$1.7 \times 10^8 \pm 3.68 \times 10^7$
K_D (M)	$6.06 \times 10^{-9} \pm 1.21 \times 10^{-9}$

EXAMPLE 22Epitope mappingEpitope mapping using PEPSCAN method

[0708] According to known procedures (Geysen et al. 1984. Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid. Proc Natl Acad Sci USA 81:3998; Slootstra et al. 1996. Structural aspects of antibody-antigen interaction revealed through small random peptide libraries. Mol Divers 1:87; Puijk et al. 2001. Segment synthesis. In PCT, The Netherlands, p.1.), overlapping 20-mer linear and 15-mer looped peptides were synthesized covering 138 amino acids at the C-terminus of human CD38. Furthermore, based on the sequence at the C-terminus single-looped peptides of different size were made covering region KNIYRPDKFLQCVKNPEDSSCTSEI, region CVHNLQPEKVQTLAWVIHGG, and region CLESIIISKRNQFSKNIYRC. In addition, extra sets were designed to reconstruct double-looped regions that were composed of SKRNQFSCKNIYR and EKVQTLAWVIHGG. Native cysteines were replaced by alanines. Peptides were screened in an ELISA-assay using credit-card format mini-PEPSCAN cards.

Synthesis of peptides

[0709] The peptides were synthesized using standard Fmoc-chemistry and deprotected using TFA with scavengers. Subsequently, the deprotected peptides were reacted on the microarray with an 0.5 mM solution of 2,6-bis(bromomethyl)pyridine or 2,4,6-tris(bromomethyl)mesitylene in ammonium bicarbonate (20 mM, pH 7.9), supplemented with acetonitrile (1:1 [volume/volume]). The microarrays were gently shaken in the solution for 30-60 min, while completely covered in the solution. Finally, the microarrays were washed extensively with excess of Millipore H₂O and sonicated in disrupt-buffer containing 1% sodium dodecylsulfate, 0.1% β -mercaptoethanol, in PBS (pH 7.2) at 70°C for 30 min, followed by sonication in millipore H₂O for another 45 min.

PEPSCAN ELISA-assay

[0710] The 455-well credit card-format polyethylene cards, containing the covalently linked peptides, were incubated with serum (e.g. diluted 1:1000 in blocking solution which contains 5% horse serum [volume/volume] and 5% ovalbumin [weight/volume]) (4°C, overnight). After washing, the peptides were incubated with rabbit- anti-human Ig peroxidase (dilution 1:1000, 25°C, 1 hour), and after washing the peroxidase substrate (2,2'-azino-di-3-ethylbenzthiazoline sulfonate and 2 μ l/ml 3% H₂O₂) was added. After one hour, the color development was measured with a CCD-camera and an image processing system. The set up consists of a CCD-camera with a 55 mm lens (Sony CCD Video Camera XC-77RR, Nikon micro-nikkor 55 mm f/2.8 lens), a camera adaptor (Sony Camera adaptor DC-77RR) and the Image Processing Software package Optimas, version 6.5 (Media Cybernetics, Silver Spring, MD 20910, U.S.A.; Optimas runs on a pentium II computer system).

Method for epitope representation

[0711] Individual amino acids were identified by dipeptide motifs which represent the smallest unique units in the

human CD38 amino acid sequence. All dipeptide motifs present in each of the 1164 peptides tested were awarded the ELISA value obtained for the respective whole peptide. To rank the dipeptide motifs from strong to poor binding, a relative signal was calculated by dividing the ELISA value obtained for each individual motif by the average ELISA value from all 1164 tested linear and looped peptides, and these were sorted for decreasing values. In this manner, amino acid contributions to conformational epitopes were considered. For each of the mAb tested, all dipeptide motifs scoring above 2.5 (i.e. ELISA values of peptides containing these motifs were at least 2.5 times the average ELISA value of those obtained with all 1164 peptides) were selected. The data were de-convoluted into single amino acid contributions represented on the linear CD38 sequence by a scoring system. By walking along the linear CD38 sequence and by using the unique dipeptide units as a reference point, one point was awarded each time a CD38 amino acid was present in this set of high scoring peptides.

[0712] -003, 005 and -024 were all found to bind to the regions SKRNIQFSCKNYR and EKVQTLEAWVIHGG of human CD38. -003 especially recognized the motifs RNIQF and WVIH, -005 especially recognized the motifs KRN and VQTL.

EXAMPLE 23

Enzymatic activity

[0713] The enzymatic activity of human CD38 was measured in an assay essentially as described in Graeff et al., J. Biol. Chem. 269, 30260-30267 (1994). Briefly, substrate NGD⁺ (80 μ M) was incubated with CD38 (0.6 μ g/ml His-tagged extracellular domain of human CD38, see Example 3 regarding purification of His-CD38) in a buffer containing 20 mM Tris-HCl, pH 7.0. The production of cGDPR can be monitored spectrophotometrically at the emission wavelength of 410 nm (excitation at 300 nm). In this example an excitation filter of 340 ± 60 nm and an emission filter of 430 ± 8 nm was used.

[0714] To test the effect of -003, -005 and -024 on the enzymatic activity of CD38, recombinant His-CD38 protein was pre-incubated for 15 min at room temperature with various concentrations (30, 3, 0.3 and 0.03 μ g/ml) of the different antibodies before adding the substrate NGD⁺. The production of cyclic GDP-ribose (cGDPR) was recorded at different time points after addition of antibodies (3, 6, 9, 12, 30, 45, 60, 75 and 90 min).

[0715] Fig 25B shows that -005 has a pronounced inhibitory effect on the production of cGDPR. After 90 minutes, addition of 30 and 3 μ g/ml -005 resulted in a 32% and 34% reduced production of cGDPR (Table 9). Similar results were observed in independent experiments using different batches of -005.

[0716] No inhibitory effect on cGPDR production was observed after addition of -003 (Figure 25B, Table 9), -024 (Figure 25D, Table 9) or anti-KLH (Figure 25A, Table 9).

[0717] Based on these findings -005 is also expected to inhibit the synthesis of Cyclic ADP-ribose (cADPR) from NAD⁺. Inhibition of the synthesis of cADPR can be determined according to the HPLC method described in Munshi et al., J. Biol. Chem. 275, 21566-21571 (2000).

Table 9. cGDPRibose production in presence of CD38-specific antibodies or anti-KLH.

	Production (% of NGD control)			
	30 μ g/ml	3 μ g/ml	0.3 μ g/ml	0.03 μ g/ml
KLH	110	99	108	111
-003	99	100	107	107
-005	68	66	98	102
-024	99	100	104	105

EXAMPLE 24

Comparison of -003 and -005 with Morphosys antibody 3079.

[0718] Antibodies -003 and -005 were functionally compared to Morphosys antibody 3079 (TH-3079). Methods for cloning and expression of Morphosys antibody TH-3079 are described in Example 16. Methods for CDC are described in Example 6. Methods for ADCC are described in Example 5. Figure 26A shows that -005 and -003 and TH-3079 induce CDC-mediated lysis of CD38-transfected CHO cells, with similar maximal lysis. When EC₅₀ values are compared, -005 antibody is better than TH3079 in inducing lysis of CHO-CD38 cells, with 2-times lower EC₅₀ (see Table 10).

[0719] Figure 26B shows that -005 is superior to TH-3079 in inducing CDC-mediated lysis of Daudi-luciferase cells, with maximal lysis by -005 being 2-3 times higher than by TH3079. When EC₅₀ values are compared, -005 antibody is

EP 2 567 976 B1

similar to TH-3079 in inducing lysis of Daudi-luciferase cells (see Table 10). -003 does not induce significant CDC-mediated lysis of Daudi-luciferase cells.

[0720] Figure 26C shows that in this experiment -005, -003 and TH-3079 mediate lysis of Daudi target cells via ADCC. No difference was found in (log) EC50 and maximal lysis (Table 11, n=5).

Table 10. Maximal lysis and EC50 values of CD38-specific antibodies in CDC.

	CHO-CD38 cells (n=2)		Daudi-luc cells (n=2)	
	EC50 μ g/ml	% Max.lysis	EC50 μ g/ml	% Max.lysis
-005	0.15 \pm 0.007	76.5 \pm 3.54	0.39 \pm 0.00	70.5 \pm 7.78
TH-3079	0.31 \pm 0.021	81.5 \pm 7.78	0.34 \pm 0.26	25.5 \pm 12.02
-003	4.5 \pm 0.933	62.0 \pm 16.79	nc	12 \pm 8.49

Table 11. Maximal lysis and EC₅₀ values of CD38 specific antibodies in ADCC.

	Log EC50	STD log EC50	Maximal lysis (%)	STD max. lysis
-005	0.76	0.18	49.2	12.8
-003	1.17	0.23	64	14.2
TH3079	0.96	0.10	43.8	12.0

SEQUENCE LISTING

[0721]

<110> Genmab A/S

<120> Antibodies against CD38 for treatment of multiple myeloma

<130> N402226EP-B

<160> 34

<170> PatentIn version 3.2

<210> 1

<211> 321

<212> DNA

<213> homo sapiens

<400> 1

```

gacatccaga tgacccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtc gggattagc agctggttag cctgggtatca gcagaaacca      120
gagaaagccc ctaagtcctt gatctatgct gcttcagtt tgcaaagtgg ggtcccatca      180
aggttcagcg gcagtggatc tgggacagat ttactctca ccatcagcag cctgcagcct      240
gaagattttg caacttatta ctgccaacag tataatagtt accctcggac gttcggccaa      300
gggaccaagg tggaaatcaa a                                              321

```

<210> 2

<211> 107

<212> PRT

EP 2 567 976 B1

<213> homo sapiens

<400> 2

5 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10
10 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
20 20 25 30
15 Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
35 40 45
20 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
25 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
30 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Arg
85 90 95
35 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105

<210> 3

<211> 11

<212> PRT

<213> homo sapiens

<400> 3

35 Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

<210> 4

<211> 7

<212> PRT

<213> homo sapiens

<400> 4

45 Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 5

<211> 9

<212> PRT

<213> homo sapiens

<400> 5

55 Gln Gln Tyr Asn Ser Tyr Pro Arg Thr
1 5

EP 2 567 976 B1

<210> 6
 <211> 366
 <212> DNA
 <213> homo sapiens

5

<400> 6

10

```

cagggtccagc tgggtgcagtc tggggctgag gtgaagaagc ctgggtcctc ggtgaaggtc      60
tcctgcaagg cttctggagg caccttcagc agctatgctt tcagctgggt gcgacaggcc      120
cctggacaag gacttgagtg gatgggaagg gtcatccctt tccttggtat agcaaactcc      180
gcacagaaat tccagggcag agtcacaatt accgcggaca aatccacgag cacagcctac      240
atggacctga gcagcctgag atctgaggac acggccgtat attactgtgc gagagatgat      300
atagcagcac ttggtccttt tgactactgg ggccaggga ccttggtcac cgtctcctca      360
gcctcc                                     366
  
```

20

<210> 7
 <211> 122
 <212> PRT
 <213> homo sapiens

25

<400> 7

30

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Ser Ser Tyr
 20 25 30

35

Ala Phe Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met

40

35 40 45

Gly Arg Val Ile Pro Phe Leu Gly Ile Ala Asn Ser Ala Gln Lys Phe
 50 55 60

45

Gln Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80

50

Met Asp Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Asp Asp Ile Ala Ala Leu Gly Pro Phe Asp Tyr Trp Gly Gln
 100 105 110

55

Gly Thr Leu Val Thr Val Ser Ser Ala Ser
 115 120

<210> 8

EP 2 567 976 B1

<211> 5
<212> PRT
<213> homo sapiens

5 <400> 8

Ser Tyr Ala Phe Ser
1 5

10
<210> 9
<211> 17
<212> PRT
<213> homo sapiens

15
<400> 9

20 Arg Val Ile Pro Phe Leu Gly Ile Ala Asn Ser Ala Gln Lys Phe Gln
1 5 10 15

Gly

25
<210> 10
<211> 11
<212> PRT
<213> homo sapiens

30
<400> 10

Asp Asp Ile Ala Ala Leu Gly Pro Phe Asp Tyr
1 5 10

35
<210> 11
<211> 321
<212> DNA
<213> homo sapiens

40
<400> 11

gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60

45 ctctcctgca gggccagtca gagggttagc agctacttag cctggtacca acagaaacct 120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc 180
50 aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 240
gaagattttg cagtttatta ctgtcagcag cgtagcaact ggcctccgac gttcggccaa 300
gggaccaagg tggaaatcaa a 321

55
<210> 12
<211> 107
<212> PRT
<213> homo sapiens

EP 2 567 976 B1

<400> 12

5 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30

10 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

 Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

20 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
 85 90 95

 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

25

<210> 13

<211> 11

<212> PRT

30 <213> homo sapiens

<400> 13

35 Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
 1 5 10

<210> 14

<211> 7

<212> PRT

40 <213> homo sapiens

<400> 14

45 Asp Ala Ser Asn Arg Ala Thr
 1 5

<210> 15

<211> 10

50 <212> PRT

<213> homo sapiens

<400> 15

55 Gln Gln Arg Ser Asn Trp Pro Pro Thr Phe
 1 5 10

<210> 16

EP 2 567 976 B1

<211> 372
<212> DNA
<213> homo sapiens

5 <400> 16

gaggtgcagc tgttggagtc tgggggaggc ttggtacagc ctgggggggtc cctgagactc 60
 10 tcatgtgcag tctctggatt cacctttaac agctttgcca tgagctgggt ccgccaggct 120
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtggtggtgg cacatactac 180
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240
 15 ctgcaaataga acagcctgag agccgaggac acggccgtat atttctgtgc gaaagataag 300
 attctctggt tcggggagcc cgtctttgac tactggggcc agggaaccct ggtcaccgtc 360
 tcctcagcct cc 372

20 <210> 17
<211> 124
<212> PRT
<213> homo sapiens

25 <400> 17

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 30 Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Phe Thr Phe Asn Ser Phe
 20 25 30
 35 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 40 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 45 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Phe Cys
 85 90 95
 Ala Lys Asp Lys Ile Leu Trp Phe Gly Glu Pro Val Phe Asp Tyr Trp
 100 105 110
 50 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser

115 120

55 <210> 18
<211> 5
<212> PRT

EP 2 567 976 B1

<213> homo sapiens

<400> 18

5

Ser Phe Ala Met Ser
1 5

<210> 19

10

<211> 17

<212> PRT

<213> homo sapiens

<400> 19

15

Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

20

Gly

<210> 20

<211> 13

<212> PRT

25

<213> homo sapiens

<400> 20

30

Asp Lys Ile Leu Trp Phe Gly Glu Pro Val Phe Asp Tyr
1 5 10

<210> 21

<211> 321

35

<212> DNA

<213> homo sapiens

<400> 21

40

gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60

ctctcctgca gggccagtca gagggttagc agctacttag cctggtacca acagaaacct 120

ggccaggctc ccgggctcct catctatgat gcttccaaca gggcctctgg catcccagcc 180

45

aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 240

gaagattttg cagtttatta ctgtcagcag cgtagcaact ggcctctcac tttcggcgga 300

gggaccaagg tggagatcaa a 321

50

<210> 22

<211> 107

<212> PRT

<213> homo sapiens

55

<400> 22

EP 2 567 976 B1

1 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 5 5 10 15
 5
 20 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 20 25 30
 10
 35 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Gly Leu Leu Ile
 35 40 45
 15
 50 Tyr Asp Ala Ser Asn Arg Ala Ser Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 20
 65 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80
 20
 85 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Leu
 85 90 95
 25
 100 Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105
 30
 <210> 23
 <211> 11
 <212> PRT
 <213> homo sapiens
 30
 <400> 23
 35
 1 Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
 1 5 10
 40
 <210> 24
 <211> 7
 <212> PRT
 <213> homo sapiens
 40
 <400> 24
 45
 1 Asp Ala Ser Asn Arg Ala Ser
 1 5
 50
 <210> 25
 <211> 9
 <212> PRT
 <213> homo sapiens
 50
 <400> 25
 55
 1 Gln Gln Arg Ser Asn Trp Pro Leu Thr
 1 5
 <210> 26

EP 2 567 976 B1

<211> 366
<212> DNA
<213> homo sapiens

5 <400> 26

gaggtgcagc tgggtgcagtc tggagcagag gtgaaaaagc ccggggagtc tctgaagatc 60
tcctgtaagg gttctggata cagcttttcc aactactgga tcggctgggt gcgccagatg 120

10

cccgggaaag gcctggagtg gatggggatc atctatcctc atgactctga tgccagatac 180
agcccgtcct tccaaggcca ggtcaccttc tcagccgaca agtccatcag caccgcctac 240
15 ctgcagtgga gcagcctgaa ggcctcggac accgccatgt attactgtgc gagacatgta 300
gggtggggat cgcggtactg gtacttcgat ctctggggcc gtggcaccct ggtcactgtc 360
tcctca 366

<210> 27
<211> 122
<212> PRT
<213> homo sapiens

25 <400> 27

30 Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15
Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Ser Asn Tyr
20 25 30
35 Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45
40 Gly Ile Ile Tyr Pro His Asp Ser Asp Ala Arg Tyr Ser Pro Ser Phe
50 55 60
45 Gln Gly Gln Val Thr Phe Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80
50 Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95
Ala Arg His Val Gly Trp Gly Ser Arg Tyr Trp Tyr Phe Asp Leu Trp
100 105 110
55 Gly Arg Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 28
<211> 5
<212> PRT
<213> homo sapiens

EP 2 567 976 B1

<400> 28

5

Asn Tyr Trp Ile Gly
1 5

<210> 29

<211> 17

<212> PRT

10

<213> homo sapiens

<400> 29

15

Ile Ile Tyr Pro His Asp Ser Asp Ala Arg Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

20

<210> 30

<211> 13

<212> PRT

<213> homo sapiens

25

<400> 30

30

His Val Gly Trp Gly Ser Arg Tyr Trp Tyr Phe Asp Leu
1 5 10

<210> 31

<211> 300

<212> PRT

35

<213> homo sapiens

<400> 31

40

45

50

55

EP 2 567 976 B1

	Met	Ala	Asn	Cys	Glu	Phe	Ser	Pro	Val	Ser	Gly	Asp	Lys	Pro	Cys	Cys
	1				5					10					15	
5	Arg	Leu	Ser	Arg	Arg	Ala	Gln	Leu	Cys	Leu	Gly	Val	Ser	Ile	Leu	Val
				20					25					30		
	Leu	Ile	Leu	Val	Val	Val	Leu	Ala	Val	Val	Val	Pro	Arg	Trp	Arg	Gln
10			35					40					45			
	Gln	Trp	Ser	Gly	Pro	Gly	Thr	Thr	Lys	Arg	Phe	Pro	Glu	Thr	Val	Leu
		50					55					60				
15	Ala	Arg	Cys	Val	Lys	Tyr	Thr	Glu	Ile	His	Pro	Glu	Met	Arg	His	Val
	65					70					75				80	
	Asp	Cys	Gln	Ser	Val	Trp	Asp	Ala	Phe	Lys	Gly	Ala	Phe	Ile	Ser	Lys
20					85					90					95	
	His	Pro	Cys	Asn	Ile	Thr	Glu	Glu	Asp	Tyr	Gln	Pro	Leu	Met	Lys	Leu
				100					105					110		
25	Gly	Thr	Gln	Thr	Val	Pro	Cys	Asn	Lys	Ile	Leu	Leu	Trp	Ser	Arg	Ile
			115					120					125			
	Lys	Asp	Leu	Ala	His	Gln	Phe	Thr	Gln	Val	Gln	Arg	Asp	Met	Phe	Thr
30		130					135					140				
	Leu	Glu	Asp	Thr	Leu	Leu	Gly	Tyr	Leu	Ala	Asp	Asp	Leu	Thr	Trp	Cys
	145					150					155					160
35	Gly	Glu	Phe	Asn	Thr	Ser	Lys	Ile	Asn	Tyr	Gln	Ser	Cys	Pro	Asp	Trp
40																
45																
50																
55																

EP 2 567 976 B1

					165					170					175	
5	Arg	Lys	Asp	Cys 180	Ser	Asn	Asn	Pro	Val 185	Ser	Val	Phe	Trp	Lys 190	Thr	Val
	Ser	Arg	Arg 195	Phe	Ala	Glu	Ala	Ala 200	Cys	Asp	Val	Val	His 205	Val	Met	Leu
10	Asn	Gly 210	Ser	Arg	Ser	Lys	Ile 215	Phe	Asp	Lys	Asn	Ser 220	Thr	Phe	Gly	Ser
15	Val 225	Glu	Val	His	Asn	Leu 230	Gln	Pro	Glu	Lys	Val 235	Gln	Thr	Leu	Glu	Ala 240
	Trp	Val	Ile	His	Gly 245	Gly	Arg	Glu	Asp	Ser 250	Arg	Asp	Leu	Cys	Gln 255	Asp
20	Pro	Thr	Ile	Lys 260	Glu	Leu	Glu	Ser	Ile 265	Ile	Ser	Lys	Arg	Asn 270	Ile	Gln
25	Phe	Ser	Cys 275	Lys	Asn	Ile	Tyr	Arg 280	Pro	Asp	Lys	Phe	Leu 285	Gln	Cys	Val
	Lys	Asn 290	Pro	Glu	Asp	Ser	Ser 295	Cys	Thr	Ser	Glu	Ile 300				

<210> 32
 <211> 300
 <212> PRT
 <213> homo sapiens

<400> 32

EP 2 567 976 B1

	Met	Ala	Asn	Cys	Glu	Phe	Ser	Pro	Val	Ser	Gly	Asp	Lys	Pro	Cys	Cys
	1				5					10					15	
5	Arg	Leu	Ser	Arg	Arg	Ala	Gln	Leu	Cys	Leu	Gly	Val	Ser	Ile	Leu	Val
				20					25					30		
	Leu	Ile	Leu	Val	Val	Val	Leu	Ala	Val	Val	Val	Pro	Arg	Trp	Arg	Gln
10			35					40					45			
	Gln	Trp	Ser	Gly	Pro	Gly	Thr	Thr	Lys	Arg	Phe	Pro	Glu	Thr	Val	Leu
		50					55					60				
15	Ala	Arg	Cys	Val	Lys	Tyr	Thr	Glu	Ile	His	Pro	Glu	Met	Arg	His	Val
	65					70					75					80
	Asp	Cys	Gln	Ser	Val	Trp	Asp	Ala	Phe	Lys	Gly	Ala	Phe	Ile	Ser	Lys
20					85					90					95	
	His	Pro	Cys	Asn	Ile	Thr	Glu	Glu	Asp	Tyr	Gln	Pro	Leu	Met	Lys	Leu
				100					105					110		
25																
30																
35																
40																
45																
50																
55																

EP 2 567 976 B1

5 Gly Thr Gln Thr Val Pro Cys Asn Lys Ile Leu Leu Trp Ser Arg Ile
115 120 125

Lys Asp Leu Ala His Gln Phe Thr Gln Val Gln Arg Asp Met Phe Thr
130 135 140

10 Leu Glu Asp Thr Leu Leu Gly Tyr Leu Ala Asp Asp Leu Thr Trp Cys
145 150 155 160

Gly Glu Phe Asn Thr Ser Lys Ile Asn Tyr Gln Ser Cys Pro Asp Trp
165 170 175

15 Arg Lys Asp Cys Ser Asn Asn Pro Val Ser Val Phe Trp Lys Thr Val
180 185 190

20 Ser Arg Arg Phe Ala Glu Ala Ala Cys Asp Val Val His Val Met Leu
195 200 205

25 Asn Gly Ser Arg Ser Lys Ile Phe Asp Lys Asn Ser Thr Phe Gly Ser
210 215 220

Val Glu Val His Asn Leu Gln Pro Glu Lys Val Gln Ala Leu Glu Ala
225 230 235 240

30 Trp Val Ile His Gly Gly Arg Glu Asp Ser Arg Asp Leu Cys Gln Asp
245 250 255

35 Pro Thr Ile Lys Glu Leu Glu Ser Ile Ile Ser Lys Arg Asn Ile Gln
260 265 270

Phe Ser Cys Lys Asn Ile Tyr Arg Pro Asp Lys Phe Leu Gln Cys Val
275 280 285

40 Lys Asn Pro Glu Asp Ser Ser Cys Thr Ser Glu Ile
290 295 300

<210> 33
<211> 300
45 <212> PRT
<213> homo sapiens

<400> 33

50 Met Ala Asn Cys Glu Phe Ser Pro Val Ser Gly Asp Lys Pro Cys Cys
1 5 10

Arg Leu Ser Arg Arg Ala Gln Leu Cys Leu Gly Val Ser Ile Leu Val
20 25 30

55 Leu Ile Leu Val Val Val Leu Ala Val Val Val Pro Arg Trp Arg Gln
35 40 45

EP 2 567 976 B1

5 Gln Trp Ser Gly Pro Gly Thr Thr Lys Arg Phe Pro Glu Thr Val Leu
 50 55 60
 10 Ala Arg Cys Val Lys Tyr Thr Glu Ile His Pro Glu Met Arg His Val
 65 70 75 80
 15 Asp Cys Gln Ser Val Trp Asp Ala Phe Lys Gly Ala Phe Ile Ser Lys
 85 90 95
 20 His Pro Cys Asn Ile Thr Glu Glu Asp Tyr Gln Pro Leu Met Lys Leu
 100 105 110
 25 Gly Thr Gln Thr Val Pro Cys Asn Lys Ile Leu Leu Trp Ser Arg Ile
 115 120 125
 30 Lys Asp Leu Ala His Gln Phe Thr Gln Val Gln Arg Asp Met Phe Thr
 130 135 140
 35 Leu Glu Asp Thr Leu Leu Gly Tyr Leu Ala Asp Asp Leu Thr Trp Cys
 145 150 155 160
 40 Gly Glu Phe Asn Thr Ser Lys Ile Asn Tyr Gln Ser Cys Pro Asp Trp
 165 170 175
 45 Arg Lys Asp Cys Ser Asn Asn Pro Val Ser Val Phe Trp Lys Thr Val
 180 185 190
 50 Ser Arg Arg Phe Ala Glu Ala Ala Cys Asp Val Val His Val Met Leu
 195 200 205
 55 Asn Gly Ser Arg Ser Lys Ile Phe Asp Lys Asn Ser Thr Phe Gly Ser
 210 215 220
 60 Val Glu Val His Asn Leu Gln Pro Glu Lys Val Gln Thr Leu Glu Ala
 225 230 235 240
 65 Trp Val Ile His Gly Gly Arg Glu Asp Ser Arg Asp Leu Cys Gln Asp
 245 250 255
 70 Pro Thr Ile Lys Glu Leu Glu Ser Ile Ile Ser Lys Arg Asn Ile Arg
 260 265 270
 75 Phe Ser Cys Lys Asn Ile Tyr Arg Pro Asp Lys Phe Leu Gln Cys Val
 275 280 285
 80 Lys Asn Pro Glu Asp Ser Ser Cys Thr Ser Glu Ile
 290 295 300

<210> 34
 <211> 300
 <212> PRT

EP 2 567 976 B1

<213> homo sapiens

<400> 34

5	Met	Ala	Asn	Cys	Glu	Phe	Ser	Pro	Val	Ser	Gly	Asp	Lys	Pro	Cys	Cys
	1				5					10					15	
10	Arg	Leu	Ser	Arg	Arg	Ala	Gln	Leu	Cys	Leu	Gly	Val	Ser	Ile	Leu	Val
				20					25					30		
	Leu	Ile	Leu	Val	Val	Val	Leu	Ala	Val	Val	Val	Pro	Arg	Trp	Arg	Gln
			35					40					45			
15	Gln	Trp	Ser	Gly	Pro	Gly	Thr	Thr	Lys	Arg	Phe	Pro	Glu	Thr	Val	Leu
		50					55					60				
20	Ala	Arg	Cys	Val	Lys	Tyr	Thr	Glu	Ile	His	Pro	Glu	Met	Arg	His	Val
	65					70					75					80
	Asp	Cys	Gln	Ser	Val	Trp	Asp	Ala	Phe	Lys	Gly	Ala	Phe	Ile	Ser	Lys
					85					90					95	
25	His	Pro	Cys	Asn	Ile	Thr	Glu	Glu	Asp	Tyr	Gln	Pro	Leu	Met	Lys	Leu
				100					105					110		
30	Gly	Thr	Gln	Thr	Val	Pro	Cys	Asn	Lys	Ile	Leu	Leu	Trp	Ser	Arg	Ile
			115					120					125			
	Lys	Asp	Leu	Ala	His	Gln	Phe	Thr	Gln	Val	Gln	Arg	Asp	Met	Phe	Thr
		130					135					140				
35	Leu	Glu	Asp	Thr	Leu	Leu	Gly	Tyr	Leu	Ala	Asp	Asp	Leu	Thr	Trp	Cys
	145					150					155					160
40	Gly	Glu	Phe	Asn	Thr	Ser	Lys	Ile	Asn	Tyr	Gln	Ser	Cys	Pro	Asp	Trp
					165					170					175	
	Arg	Lys	Asp	Cys	Ser	Asn	Asn	Pro	Val	Ser	Val	Phe	Trp	Lys	Thr	Val
				180					185					190		
45	Ser	Arg	Arg	Phe	Ala	Glu	Ala	Ala	Cys	Asp	Val	Val	His	Val	Met	Leu
			195					200					205			
50	Asn	Gly	Ser	Arg	Ser	Lys	Ile	Phe	Asp	Lys	Asn	Ser	Thr	Phe	Gly	Ser
		210					215					220				
	Val	Glu	Val	His	Asn	Leu	Gln	Pro	Glu	Lys	Val	Gln	Thr	Leu	Glu	Ala
	225					230					235					240
55	Trp	Val	Ile	His	Gly	Gly	Arg	Glu	Asp	Ser	Arg	Asp	Leu	Cys	Gln	Asp
					245					250					255	

Pro Thr Ile Lys Glu Leu Glu Ser Ile Ile Ser Lys Arg Asn Ile Gln
260 265 270

5 Phe Phe Cys Lys Asn Ile Tyr Arg Pro Asp Lys Phe Leu Gln Cys Val
275 280 285

10 Lys Asn Pro Glu Asp Ser Ser Cys Thr Ser Glu Ile
290 295 300

Claims

- 15 1. An antibody binding to human CD38, wherein said antibody comprises human light chain and human heavy chain variable regions, wherein the light chain variable region comprises a V_L CDR1 having the sequence as set forth in SEQ ID No:13, a V_L CDR2 having the sequence as set forth in SEQ ID No:14 and a V_L CDR3 having the sequence as set forth in SEQ ID No:15, and the heavy chain variable region comprises a V_H CDR1 having the sequence as set forth in SEQ ID No:18, a V_H CDR2 having the sequence as set forth in SEQ ID No:19 and a V_H CDR3 having the sequence as set forth in SEQ ID No:20.
- 20 2. The antibody according to claim 1, which is a human monoclonal antibody.
3. The antibody according to any one of claims 1 and 2, **characterized in that** it is a full length IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody.
- 25 4. The antibody according to claim 3, which is an IgG1 antibody.
5. The antibody according to claim 4, which is an IgG1, κ antibody.
- 30 6. The antibody according to claim 3, which is an IgM antibody.
7. The antibody according to claim 6, which is an IgM, κ antibody.
- 35 8. The antibody according to any one of claims 1 to 7, comprising a V_H region having the amino acid sequence as set forth in SEQ ID No:17.
9. The antibody according to any one of claims 1 to 7, comprising a V_H region comprising the amino acid sequence spanning the V_H CDR1 - V_H CDR3 region of SEQ ID No:17.
- 40 10. The antibody according to any one of claims 1 to 9, comprising a V_L region having the amino acid sequence as set forth in SEQ ID NO:12.
11. The antibody according to claim 1, wherein the antibody is encoded by
 - 45 (i) human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions as set forth in SEQ ID No:11 and SEQ ID No:16, respectively; or
 - (ii) human light chain and human heavy chain nucleic acids comprising nucleotide sequences in their variable regions, which are conservative sequence modifications of the sequences as set forth in (i).
- 50 12. An antibody according to any one of the preceding claims, which is an antibody fragment or a single chain antibody.
13. An antibody according to any one of the preceding claims, which is in a substantially isolated form.
14. An antibody according to any one of the preceding claims, further comprising a chelator linker for attaching a radioisotope.
- 55 15. An isolated nucleic acid encoding an antibody according to any one of claims 1 to 12.

16. An expression vector comprising a nucleic acid sequence encoding an antibody according to any one of claims 1 to 12.
17. A eukaryotic or prokaryotic host cell which produces an antibody according to any one of claims 1 to 12.
- 5 18. An immunoconjugate comprising an antibody according to any one of claims 1 to 13 linked to a cytotoxic agent, a radioisotope, or a drug.
19. A bispecific or multispecific molecule comprising the antibody according to any one of claims 1 to 14 and a binding specificity for a human effector cell.
- 10 20. A pharmaceutical composition comprising the antibody according to any one of claims 1 to 14, the immunoconjugate according to claim 18, or the bispecific or multispecific molecule according to claim 19 and a pharmaceutically acceptable carrier.
- 15 21. An antibody according to any one of claims 1 to 14, an immunoconjugate according to claim 18, a bispecific or multispecific molecule according to claim 19 or an expression vector according to claim 16 for use as a medicament.
22. An antibody according to any one of claims 1 to 14, an immunoconjugate according to claim 18, the pharmaceutical composition according to claim 20, the bispecific or multispecific molecule according to claim 19 or an expression vector according to claim 16 for use in a method of treating or preventing a disease or disorder involving cells expressing CD38, wherein the disease or disorder is selected from the group consisting of B cell lymphoma/leukemias including precursor B cell lymphoblastic leukemia/lymphoma and B cell non-Hodgkin's lymphomas; acute lymphoblastic leukemia; mature B cell neoplasms including B cell chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL); B cell acute lymphocytic leukemia; B cell prolymphocytic leukemia; lymphoplasmacytic lymphoma; 25 mantle cell lymphoma (MCL); follicular lymphoma (FL) including low-grade, intermediate-grade and high-grade FL; cutaneous follicle center lymphoma; marginal zone B cell lymphoma including MALT type, nodal and splenic type; hairy cell leukemia; diffuse large B cell lymphoma; Burkitt's lymphoma; plasmacytoma; plasma cell myeloma; plasma cell leukemia; post-transplant lymphoproliferative disorder; Waldenström's macroglobulinemia; plasma cell leukemias; anaplastic large-cell lymphoma (ALCL); multiple myeloma; Hodgkin's lymphoma; mature T cell and NK cell neoplasms including T cell prolymphocytic leukemia; T cell large granular lymphocytic leukemia; aggressive NK cell leukemia; adult T cell leukemia/lymphoma; extranodal NK/T cell lymphoma; nasal type, enteropathy-type T cell lymphoma; hepatosplenic T cell lymphoma; subcutaneous panniculitis-like T cell lymphoma; blastic NK cell lymphoma; Mycosis Fungoides/Sézary Syndrome; primary cutaneous CD30 positive T cell lymphoproliferative disorders including cutaneous anaplastic large cell lymphoma C-ALCL, lymphomatoid papulosis and borderline lesions; angioimmunoblastic T cell lymphoma, peripheral T cell lymphoma unspecified; anaplastic large cell lymphoma; acute myeloid leukemia including acute promyelocytic leukemia; and chronic myeloproliferative diseases including chronic myeloid leukemia.
- 30 23. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for the use according to claim 22, wherein the disease is B cell non-Hodgkin's lymphoma.
- 35 24. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for the use according to claim 22, wherein the disease is multiple myeloma.
- 40 25. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for the use according to claim 22, wherein the disease is acute lymphoblastic leukemia.
- 45 26. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for the use according to claim 22, wherein the disease is B cell chronic lymphocytic leukemia.
- 50 27. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for use according to any one of claims 21 to 26, wherein the method comprises administration of one or more further therapeutic agents to the subject.
- 55 28. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for use according to claim 27, wherein the one or more further therapeutic agents are selected from a chemotherapeutic agent, an anti-inflammatory agent, or an immunosuppressive and/or immunomodulatory agent.

29. The antibody, immunoconjugate, pharmaceutical composition, bispecific or multispecific molecule or expression vector for use according to claim 27, wherein the one or more further therapeutic agents are selected from the group consisting of cisplatin, gefitinib, cetuximab, rituximab, bevacizumab, erlotinib, bortezomib, thalidomide, pamidronate, zoledronic acid, clodronate, risendronate, ibandronate, etidronate, alendronate, tiludronate, arsenic trioxide, lenalidomide, filgrastim, pegfilgrastim, sargramostim, suberoylanilide hydroxamic acid, and SCIO-469.

30. An *in vitro* method for detecting the presence of CD38 antigen, or a cell expressing CD38, in a sample comprising:

- a) contacting the sample with the antibody according to any one of claims 1 to 11 under conditions that allow for formation of a complex between the peptide and CD38; and
- b) detecting the formation of a complex.

31. An anti-idiotypic antibody binding to an antibody according to any one of claims 1 to 12.

Patentansprüche

1. Antikörper, welcher an humanes CD38 bindet, wobei der Antikörper variable Regionen der humanen leichten Kette und humanen schweren Kette umfasst, wobei die variable Region der leichten Kette eine V_L-CDR1, die die Sequenz, wie in SEQ ID NO:13 gezeigt, besitzt, eine V_L-CDR2, welche die Sequenz, wie in SEQ ID NO:14 gezeigt, besitzt, und eine V_L-CDR3, welche die Sequenz, wie in SEQ ID NO:15 gezeigt, besitzt, umfasst, und die variable Region der schweren Kette eine V_H-CDR1, welche die Sequenz, wie in SEQ ID NO:18 gezeigt, besitzt, eine V_H-CDR2, welche die Sequenz, wie in SEQ ID NO:19 gezeigt, besitzt, und eine V_H-CDR3, welche die Sequenz, wie in SEQ ID NO:20 gezeigt, besitzt, umfasst.

2. Antikörper gemäß Anspruch 1, welcher ein humaner monoklonaler Antikörper ist.

3. Antikörper gemäß einem der Ansprüche 1 und 2, dadurch charakterisiert, dass er ein Volllängen-IgG1-, -IgG2-, -IgG3-, -IgG4-, -IgD-, -IgA-, -IgE- oder -IgM-Antikörper ist.

4. Antikörper gemäß Anspruch 3, welcher ein IgG1-Antikörper ist.

5. Antikörper gemäß Anspruch 4, welcher ein IgG1,κ-Antikörper ist.

6. Antikörper gemäß Anspruch 3, welcher ein IgM-Antikörper ist.

7. Antikörper gemäß Anspruch 6, welcher ein IgM,κ-Antikörper ist.

8. Antikörper gemäß einem der Ansprüche 1 bis 7, umfassend eine V_H-Region, welche die Aminosäuresequenz, wie in SEQ ID NO:17 gezeigt, besitzt.

9. Antikörper gemäß einem der Ansprüche 1 bis 7, umfassend eine V_H-Region, umfassend die Aminosäuresequenz, welche die V_H-CDR1 - V_H-CDR3-Region von SEQ ID NO:17 umfasst.

10. Antikörper gemäß einem der Ansprüche 1 bis 9, umfassend eine V_L-Region, welche die Aminosäuresequenz, wie in SEQ ID NO 12 gezeigt, besitzt.

11. Antikörper gemäß Anspruch 1, wobei der Antikörper codiert ist durch

- (i) Nucleinsäuren der humanen leichten Kette und der humanen schweren Kette, umfassend die Nucleotidsequenzen in ihren variablen Regionen, wie es in SEQ ID NO:11 bzw. SEQ ID NO:16 gezeigt ist; oder
- (ii) Nucleinsäuren der humanen leichten Kette und der humanen schweren Kette, umfassend die Nucleotidsequenzen in ihren variablen Regionen, welche konservative Sequenzmodifikationen der Sequenzen, wie in (i) genannt, sind.

12. Antikörper gemäß einem der vorhergehenden Ansprüche, welcher ein Antikörperfragment oder ein Einzelkettenantikörper ist.

13. Antikörper gemäß einem der vorhergehenden Ansprüche, welcher sich in einer im Wesentlichen isolierten Form befindet.
14. Antikörper gemäß einem der vorhergehenden Ansprüche, zusätzlich umfassend einen Chelator-Linker zur Anbin-
5 dung eines Radioisotops.
15. Isolierte Nucleinsäure, welche einen Antikörper gemäß einem der Ansprüche 1 bis 12 codiert.
16. Expressionsvektor, umfassend eine Nucleinsäuresequenz, welche einen Antikörper gemäß einem der Ansprüche
10 1 bis 12 codiert.
17. Eukaryotische oder prokaryotische Wirtszelle, welche einen Antikörper gemäß einem der Ansprüche 1 bis 12 pro-
duziert.
18. Immunkonjugat, umfassend einen Antikörper gemäß einem der Ansprüche 1 bis 13, welcher mit einem zytotoxischen
15 Agens, einem Radioisotop oder einem Medikament verknüpft ist.
19. Bispezifisches oder multispezifisches Molekül, umfassend den Antikörper gemäß einem der Ansprüche 1 bis 14
und eine Bindungsspezifität für eine humane Effektorzelle.
20
20. Pharmazeutische Zusammensetzung, umfassend den Antikörper gemäß einem der Ansprüche 1 bis 14, das Im-
munkonjugat gemäß Anspruch 18 oder das bispezifische oder multispezifische Molekül gemäß Anspruch 19 und
einen pharmazeutisch verträglichen Trägerstoff.
21. Antikörper gemäß einem der Ansprüche 1 bis 14, Immunkonjugat gemäß Anspruch 18, bispezifisches oder multi-
25 spezifisches Molekül gemäß Anspruch 19 oder Expressionsvektor gemäß Anspruch 16 zur Verwendung als Medi-
kament.
22. Antikörper gemäß einem der Ansprüche 1 bis 14, Immunkonjugat gemäß Anspruch 18, pharmazeutische Zusam-
30 mensetzung gemäß Anspruch 20, bispezifisches oder multispezifisches Molekül gemäß Anspruch 19 oder Expres-
sionsvektor gemäß Anspruch 16 zur Verwendung in einem Verfahren zur Behandlung oder Prävention einer Krank-
heit oder Störung, die Zellen, welche CD38 exprimieren, involviert, wobei die Krankheit oder Störung ausgewählt
ist aus der Gruppe, bestehend aus B-Zell-Lymphom/ Leukämien, einschließlich Vorläufer-B-Zell-lymphoblastischer
35 Leukämie/lymphoblastischem Lymphom und B-Zell-nicht-Hodgkin-Lymphomen; akuter Lymphoblastenleukämie;
reifen B-Zell-Neoplasmen, einschließlich chronischer lymphozytärer Leukämie (CLL) des B-Zell-Typs/kleinem lym-
phozytärem Lymphom (SLL); akuter lymphozytärer Leukämie des B-Zell-Typs; prolymphozytärer Leukämie des B-
Zell-Typs; lymphoplasmozytischem Lymphom; Mantelzellenlymphom (MCL); folliculärem Lymphom (FL), ein-
schließlich niedriggradigem, zwischengradigem und hochgradigem FL; kutanem Follikelzentrumlymphom; Margi-
40 nalzonen-B-Zell-Lymphom, einschließlich des MALT-Typs, des knotigen Typs und des lienalen Typs; Haarzellen-
leukämie; diffusen großem B-Zell-Lymphom; Burkitts Lymphom; Plasmozytom; Plasmazellenmyelom; Plasmazel-
lenleukämie; lymphoproliferativer Störung nach Transplantation; Waldenströms Makroglobulinämie; Plasmazellen-
leukämien; anaplastischem großzelligem Lymphom (anaplastic large-cell lymphoma (ALCL)); multiplem Myelom;
Hodgkin-Lymphom; reifen T-Zell- und NK-Zell-Neoplasmen, einschließlich T-Zell-prolymphozytärer Leukämie; T-
45 Zellgroßer granulärer lymphozytärer Leukämie; aggressiver NK-Zell-Leukämie; adulter T-Zell-Leukämie/adultem T-
Zell-Lymphom; extranodalem NK/T-Zell-Lymphom; T-Zell-Lymphom des nasalen Typs, Enteropathie-Typs; hepa-
tosplenischem T-Zell-Lymphom; subkutanem Panniculitis-artigem T-Zell-Lymphom; blastischem NK-Zell-Lymphom;
Mycosis Fungoides/Sézary-Syndrom; primären kutanen CD30-positiven T-Zelllymphoproliferativen Störungen, ein-
50 schließlich kutanem anaplastischem großzelligem Lymphom C-ALCL, lymphomatoider Papulose und Borderline-
Läsionen; angioimmunoblastischem T-Zell-Lymphom, peripheren T-Zell-Lymphomen, die unspezifiziert sind; ana-
plastischem großzelligem Lymphom; akuter myeloider Leukämie, einschließlich akuter promyelozytischer Leukämie,
und chronischen myeloproliferativen Krankheiten, einschließlich chronischer myeloider Leukämie.
23. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül
55 oder Expressionsvektor zur Verwendung gemäß Anspruch 22, wobei die Krankheit B-Zell-nicht-Hodgkin-Lymphom
ist.
24. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül
oder Expressionsvektor zur Verwendung gemäß Anspruch 22, wobei die Krankheit multiples Myelom ist.

25. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül oder Expressionsvektor zur Verwendung gemäß Anspruch 22, wobei die Krankheit akute Lymphoblastenleukämie ist.

26. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül oder Expressionsvektor zur Verwendung gemäß Anspruch 22, wobei die Krankheit chronische lymphozytäre Leukämie des B-Zell-Typs ist.

27. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül oder Expressionsvektor zur Verwendung gemäß einem der Ansprüche 21 bis 26, wobei das Verfahren die Verabreichung eines weiteren Therapeutikums oder mehrerer weiterer Therapeutika an das Subjekt umfasst.

28. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül oder Expressionsvektor zur Verwendung gemäß Anspruch 27, wobei das eine weitere Therapeutikum oder die mehreren weiteren Therapeutika aus einem Chemotherapeutikum, einem entzündungshemmenden Mittel oder einem immunsuppressiven und/oder immunmodulatorischen Mittel ausgewählt ist/sind.

29. Antikörper, Immunkonjugat, pharmazeutische Zusammensetzung, bispezifisches oder multispezifisches Molekül oder Expressionsvektor zur Verwendung gemäß Anspruch 27, wobei das eine weitere Therapeutikum oder die mehreren weiteren Therapeutika aus der Gruppe, bestehend aus Cisplatin, Gefitinib, Cetuximab, Rituximab, Bevacizumab, Erlotinib, Bortezomib, Thalidomid, Pamidronat, Zoledronsäure, Clodronat, Risendronat, Ibandronat, Etidronat, Alendronat, Tiludronat, Arsentrioxid, Lenalidomid, Filgrastim, Pegfilgrastim, Sargramostim, Suberoylanilidhydroxamsäure und SCIO-469, ausgewählt ist/sind.

30. *In-vitro*-Verfahren zum Detektieren des Vorliegens von CD38-Antigen oder einer Zelle, die CD38 exprimiert, in einer Probe, umfassend:

a) Inkontaktbringen der Probe mit einem Antikörper gemäß einem der Ansprüche 1 bis 11 unter Bedingungen, die für die Bildung eines Komplexes zwischen dem Peptid und CD38 sorgen, und

b) Detektieren der Bildung eines Komplexes.

31. Anti-idiotypischer Antikörper, welcher an einen Antikörper gemäß einem der Ansprüche 1 bis 12 bindet.

Revendications

1. Anticorps se liant au CD38 humain, lequel anticorps comprend des régions variables de chaîne légère humaine et de chaîne lourde humaine, laquelle région variable de chaîne légère comprend un domaine CDR1 de V_L ayant la séquence donnée en tant que Séquence N° 13, un domaine CDR2 de V_L ayant la séquence donnée en tant que Séquence N° 14, et un domaine CDR3 de V_L ayant la séquence donnée en tant que Séquence N° 15, et laquelle région variable de chaîne lourde comprend un domaine CDR1 de V_H ayant la séquence donnée en tant que Séquence N° 18, un domaine CDR2 de V_H ayant la séquence donnée en tant que Séquence N° 19, et un domaine CDR3 de V_H ayant la séquence donnée en tant que Séquence N° 20.

2. Anticorps conforme à la revendication 1, qui est un anticorps monoclonal humain.

3. Anticorps conforme à l'une des revendications 1 et 2, **caractérisé en ce qu'il** s'agit d'un anticorps IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE ou IgM, complet.

4. Anticorps conforme à la revendication 3, qui est un anticorps IgG1.

5. Anticorps conforme à la revendication 4, qui est un anticorps IgG1, κ .

6. Anticorps conforme à la revendication 3, qui est un anticorps IgM.

7. Anticorps conforme à la revendication 6, qui est un anticorps IgM, κ .

8. Anticorps conforme à l'une des revendications 1 à 7, qui comporte une région V_H ayant la séquence d'acides aminés

donnée en tant que Séquence N° 17.

9. Anticorps conforme à l'une des revendications 1 à 7, qui comporte une région V_H comprenant la séquence d'acides aminés couvrant les domaines CDR1 à CDR3 de V_H de la Séquence N° 17.

10. Anticorps conforme à l'une des revendications 1 à 9, qui comporte une région V_L ayant la séquence d'acides aminés donnée en tant que Séquence N° 12.

11. Anticorps conforme à la revendication 1, lequel anticorps est codé

i) par des acides nucléiques pour chaîne légère humaine et pour chaîne lourde humaine, comprenant respectivement, pour leurs régions variables, les séquences de nucléotides données en tant que Séquence N° 11 et Séquence N° 16,

ii) ou par des acides nucléiques pour chaîne légère humaine et pour chaîne lourde humaine comprenant, pour leurs régions variables, les séquences de nucléotides qui sont des variantes conservatives des séquences indiquées en (i).

12. Anticorps conforme à l'une des revendications précédentes, qui est un fragment d'anticorps ou un anticorps à chaîne unique.

13. Anticorps conforme à l'une des revendications précédentes, qui se présente sous forme pratiquement isolée.

14. Anticorps conforme à l'une des revendications précédentes, qui comprend en outre un raccord chélateur servant à fixer un isotope radioactif.

15. Acide nucléique isolé, codant un anticorps conforme à l'une des revendications 1 à 12.

16. Vecteur d'expression comprenant une séquence d'acide nucléique codant un anticorps conforme à l'une des revendications 1 à 12.

17. Cellule hôte eucaryote ou procaryote, qui produit un anticorps conforme à l'une des revendications 1 à 12.

18. Immuno-conjugué comprenant un anticorps conforme à l'une des revendications 1 à 13, raccordé à un agent cytotoxique, un isotope radioactif ou un médicament.

19. Molécule bispécifique ou multispécifique, comprenant un anticorps conforme à l'une des revendications 1 à 14 et dotée d'une capacité de liaison spécifique à une cellule effectrice humaine.

20. Composition pharmaceutique comprenant un anticorps conforme à l'une des revendications 1 à 14, un immuno-conjugué conforme à la revendication 18 ou une molécule bispécifique ou multispécifique conforme à la revendication 19, et un véhicule pharmacologiquement admissible.

21. Anticorps conforme à l'une des revendications 1 à 14, immuno-conjugué conforme à la revendication 18, molécule bispécifique ou multispécifique conforme à la revendication 19, ou vecteur d'expression conforme à la revendication 16, pour utilisation en tant que médicament.

22. Anticorps conforme à l'une des revendications 1 à 14, immuno-conjugué conforme à la revendication 18, composition pharmaceutique conforme à la revendication 20, molécule bispécifique ou multispécifique conforme à la revendication 19, ou vecteur d'expression conforme à la revendication 16, pour utilisation dans un procédé de traitement ou de prévention d'une maladie ou d'un trouble impliquant des cellules exprimant CD38, laquelle maladie ou lequel trouble est choisi(e) dans l'ensemble formé par les suivant(e)s : lymphomes/leucémies à lymphocytes B, y compris lymphome/leucémie lymphoblastique à précurseurs de lymphocytes B et lymphomes non-Hodgkiniens à lymphocytes B ; leucémie lymphoblastique aiguë ; néoplasmes à lymphocytes B matures, y compris leucémie lymphocytaire chronique à lymphocytes B (CLL) et lymphome lymphocytaire à petites cellules (SLL) ; leucémie lymphocytaire aiguë à lymphocytes B ; leucémie prolymphocytaire à lymphocytes B ; lymphome lymphoplasmocytaire ; lymphome des cellules du manteau (MCL) ; lymphome folliculaire (FL), y compris les FL à bas grade, à grade intermédiaire et à grade élevé ; lymphome cutané des centres folliculaires ; lymphomes B de la zone marginale, y compris ceux de type MALT, ganglionnaire et splénique ; leucémie à tricholeucocytes ; lymphome diffus à grands lymphocytes B ;

lymphome de Burkitt ; plasmacytome ; myélome plasmocytaire ; leucémie plasmocytaire ; trouble lymphoprolifératif post-transplantation ; macroglobulinémie de Waldenström ; leucémies à plasmocytes ; lymphome anaplasique à grandes cellules (ALCL) ; myélome multiple ; lymphome de Hodgkin ; néoplasmes à lymphocytes T et NK matures, y compris leucémie prolymphocytaire à lymphocytes T ; leucémie lymphocytaire à grands lymphocytes T granuleux ; leucémie agressive à lymphocytes NK ; leucémie/lymphome à lymphocytes T de l'adulte ; lymphome extra-ganglionnaire à lymphocytes NK/T ; lymphome à lymphocytes T de type nasal, de type entéropathie ; lymphome hépatosplénique à lymphocytes T ; lymphome sous-cutané à lymphocytes T, de type panniculite ; lymphome blastique à lymphocytes NK ; mycosis fongoïde, syndrome de Sézary ; troubles lymphoprolifératifs cutanés primaires à lymphocytes T CD30-positifs, y compris lymphome anaplasique cutané à grandes cellules (C-ALCL), papulose lymphomatoïde et lésions borderline ; lymphome angioimmunoblastique à lymphocytes T, lymphome périphérique non spécifié à lymphocytes T ; lymphome anaplasique à grandes cellules ; leucémie myéloïde aiguë, y compris leucémie pro-myélocytaire aiguë ; et troubles myéloprolifératifs chronique, y compris leucémie myéloïde chronique.

23. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 22, dans laquelle la maladie est un lymphome non-Hodgkinien à lymphocytes B.

24. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 22, dans laquelle la maladie est un myélome multiple.

25. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 22, dans laquelle la maladie est une leucémie lymphoblastique aiguë.

26. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 22, dans laquelle la maladie est une leucémie lymphocytaire chronique à lymphocytes B.

27. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à l'une des revendications 21 à 26, dans laquelle le procédé comporte le fait d'administrer au sujet un ou plusieurs autre(s) agent(s) thérapeutique(s).

28. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 27, dans laquelle l'autre ou les autres agent(s) thérapeutique(s) est ou sont choisi(s) parmi un agent de chimiothérapie, un agent anti-inflammatoire, et un agent immunosuppresseur et/ou immunomodulateur.

29. Anticorps, immuno-conjugué, composition pharmaceutique, molécule bispécifique ou multispécifique, ou vecteur d'expression, pour utilisation conforme à la revendication 27, dans laquelle l'autre ou les autres agent(s) thérapeutique(s) est ou sont choisi(s) dans l'ensemble formé par les suivants : cisplatine, géfitinib, cétuximab, rituximab, bévacizumab, erlotinib, bortézomib, thalidomide, pamidronate, acide zolédronique, clodronate, risédronate, ibandronate, étidronate, alendronate, tiludronate, trioxyde d'arsenic, lénalidomide, filgrastim, pegfilgrastim, sargramostim, acide anilido-subéroïl-hydroxamique, et SCIO-469.

30. Procédé *in vitro* permettant de détecter la présence, dans un échantillon, de l'antigène CD38 ou d'une cellule exprimant CD38, comportant :


a) le fait de mettre l'échantillon en contact avec un anticorps conforme à l'une des revendications 1 à 11, dans des conditions qui permettent la formation d'un complexe entre ce peptide et CD38,

b) et le fait de détecter la formation d'un complexe.

31. Liaison d'un anticorps anti-idiotype à un anticorps conforme à l'une des revendications 1 à 12.

FIGURE 1

A:



B:

Binding of -024 on CHO-CD38 cells.





FIGURE 2

A:



B:




FIGURE 3





FIGURE 4



B:

ADCC on Daudi cells





FIGURE 5 (1/2)

A:



B:




FIGURE 6




FIGURE 7




FIGURE 8





FIGURE 9

A:



B:





FIGURE 10 (1/3)

A:



B:





FIGURE 10 (2/3)

C:



D:





FIGURE 10 (3/3)

E:






FIGURE 12

FIGURE 12A




FIGURE 12B




FIGURE 12C




FIGURE 12D




FIGURE 13

FIGURE 13A




FIGURE 13B



FIGURE 13C



FIGURE 13D





FIGURE 14

14A




14B



14C



14D



14E




FIGURE 15 (1/2)

FIGURE 15A




FIGURE 15B




FIGURE 15 (2/2)

FIGURE 15C




FIGURE 16

FIGURE 16A




FIGURE 16B




FIGURE 17 (1/2)

FIGURE 17A




FIGURE 17B




FIGURE 17 (2/2)

FIGURE 17C




FIGURE 17D




FIGURE 18




FIGURE 19




FIGURE 20






FIGURE 21

A



B



C






FIGURE 22

A



B



C




FIGURE 23

A:





FIGURE 23

B:





FIGURE 23

C:







FIGURE 24

A:



B:



C:





FIGURE 25 (1/2)

A



B





FIGURE 25 (2/2)

C



D





FIGURE 26 (1/2)

A



B





FIGURE 26 (2/2)

C

Daudi cells Donor B



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4816567 A [0136] [0264]
- EP 125023 A [0136]
- EP 171496 A [0136]
- EP 173494 A [0136]
- WO 8601533 A [0136]
- EP 184187 A [0136]
- WO 8702671 A [0136]
- WO 0243478 A [0167] [0256] [0257] [0418]
- WO 03048185 A [0210] [0211]
- WO 03070747 A [0210]
- WO 03027246 A [0210]
- US 20040009498 A [0211]
- US 5545806 A [0254]
- US 5569825 A [0254]
- US 5625126 A [0254]
- US 5633425 A [0254]
- US 5789650 A [0254]
- US 5877397 A [0254]
- US 5661016 A [0254]
- US 5814318 A [0254]
- US 5874299 A [0254]
- US 5770429 A [0254] [0255]
- US 5545807 A [0254]
- WO 9824884 A [0254] [0383]
- WO 9425585 A [0254]
- WO 931227 A [0254]
- WO 9222645 A [0254]
- WO 9203918 A [0254]
- WO 0109187 A [0254] [0256]
- WO 0114424 A [0255] [0256]
- US 5827690 A [0259]
- US 5756687 A [0259]
- US 5750172 A [0259]
- US 5741957 A [0259]
- US 5733743 A [0260]
- US 6054297 A [0261]
- US 5886152 A [0261]
- US 5877293 A [0261]
- WO 9202190 A [0263]
- US 5530101 A [0263]
- US 5585089 A [0263]
- US 5693761 A [0263]
- US 5693792 A [0263]
- US 5714350 A [0263]
- US 5777085 A [0263]
- US 4683195 A [0265]
- US 4683202 A [0265]
- US 5223409 A [0269]
- WO 9218619 A [0269]
- WO 9117271 A [0269]
- WO 9220791 A [0269]
- WO 9215679 A [0269]
- WO 9301288 A [0269]
- WO 9201047 A [0269]
- WO 9209690 A [0269]
- WO 9306213 A [0269]
- US 5624821 A [0279]
- US 5648260 A [0279]
- WO 0042072 A [0279]
- WO 9429351 A [0279]
- US 6121022 A [0280]
- US 6194551 B [0280]
- US 6277375 B [0280]
- WO 9852976 A [0281]
- WO 0061739 A [0284]
- WO 9954342 A [0284]
- WO 9633279 A [0285]
- WO 9109967 A [0286]
- WO 9316184 A [0286]
- US 4946778 A [0290] [0293]
- US 20030118592 A [0293]
- US 20030133939 A [0293]
- WO 8800052 A, Fanger [0294]
- US 4954617 A [0294]
- WO 9410332 A [0294]
- US 4676980 A [0300]
- US 4766106 A [0307]
- US 4179337 A [0307]
- US 4495285 A [0307]
- US 4609546 A [0307]
- US 4681581 A [0309]
- US 4735210 A [0309]
- US 5101827 A [0309]
- US 5102990 A [0309]
- US RE35500 E [0309]
- US 5648471 A [0309]
- US 5697902 A [0309]
- US 5135917 A [0321]
- US 5294533 A [0321]
- US 5627158 A [0321]
- US 5641754 A [0321]
- US 5691317 A [0321]
- US 5780607 A [0321]
- US 5786138 A [0321]
- US 5849903 A [0321]
- US 5856103 A [0321]
- US 5919772 A [0321]
- US 5955590 A [0321]

EP 2 567 976 B1

- US 5990088 A [0321]
- US 5994320 A [0321]
- US 5998602 A [0321]
- US 6005095 A [0321]
- US 6007995 A [0321]
- US 6013522 A [0321]
- US 6017898 A [0321]
- US 6018042 A [0321]
- US 6025198 A [0321]
- US 6033910 A [0321]
- US 6040296 A [0321]
- US 6046004 A [0321]
- US 6046319 A [0321]
- US 6057437 A [0321]
- US 5631146 A [0322]
- US 5580737 A [0322]
- US 5786462 A [0322]
- US 5543293 A [0322]
- US 5476766 A [0322]
- US 5503978 A [0322]
- US 5731424 A [0322]
- US 5780228 A [0322]
- US 5792613 A [0322]
- US 5795721 A [0322]
- US 5846713 A [0322]
- US 5858660 A [0322]
- US 5861254 A [0322]
- US 5864026 A [0322]
- US 5869641 A [0322]
- US 5958691 A [0322]
- US 6001988 A [0322]
- US 6011020 A [0322]
- US 6013443 A [0322]
- US 6020130 A [0322]
- US 6028186 A [0322]
- US 6030776 A [0322]
- US 6051698 A [0322]
- US 5334711 A [0323]
- US 5436330 A [0323]
- US 5616466 A [0323]
- US 5633133 A [0323]
- US 5646020 A [0323]
- US 5652094 A [0323]
- US 5712384 A [0323]
- US 5770715 A [0323]
- US 5856463 A [0323]
- US 5861288 A [0323]
- US 5891683 A [0323]
- US 5891684 A [0323]
- US 5985621 A [0323]
- US 5989908 A [0323]
- US 5998193 A [0323]
- US 5998203 A [0323]
- WO 9858058 A [0323]
- WO 9858057 A [0323]
- WO 9718312 A [0323]
- US 5631115 A [0323]
- US 5646031 A [0323]
- US 5683902 A [0323]
- US 5856188 A [0323]
- US 5866701 A [0323]
- US 5869339 A [0323]
- US 6022962 A [0323]
- US 5595873 A [0323]
- US 5652107 A [0323]
- US 5580967 A [0323]
- US 5688670 A [0323]
- US 5807718 A [0323]
- US 5910408 A [0323]
- US 5646042 A [0323]
- US 5693535 A [0323]
- US 5731295 A [0323]
- US 5811300 A [0323]
- US 5837855 A [0323]
- US 5869253 A [0323]
- US 5877021 A [0323]
- US 5877022 A [0323]
- US 5972699 A [0323]
- US 5972704 A [0323]
- US 5989906 A [0323]
- US 6017756 A [0323]
- US 5176996 A [0324]
- US 5645985 A [0324]
- US 5650316 A [0324]
- US 5683874 A [0324]
- US 5693773 A [0324]
- US 5834185 A [0324]
- US 5869246 A [0324]
- US 5874566 A [0324]
- US 5962426 A [0324]
- WO 9203566 A [0325]
- US 5168053 A [0325]
- US 5624824 A [0325]
- US 5683873 A [0325]
- US 5728521 A [0325]
- US 5869248 A [0325]
- US 5877162 A [0325]
- WO 0175164 A [0326]
- US 6506559 B [0326]
- US 20040086884 A [0326]
- US 20040077574 A [0326]
- US 20040063654 A [0326]
- US 20040033602 A [0326]
- US 20030167490 A [0326]
- US 20030157030 A [0326]
- US 20030114409 A [0326]
- US 20030108923 A [0326]
- US 20040014113 A [0326]
- US 20020132788 A [0326]
- US 5194594 A [0335]
- US 6077499 A [0339]
- US 5906820 A [0342]
- US 4699880 A [0346]
- US 6077835 A [0349]
- WO 0070087 A [0349]
- WO 0046147 A [0349]

EP 2 567 976 B1

- US 5589466 A [0349] [0551]
- US 5973972 A [0349]
- US 5849586 A [0354]
- WO 9704748 A [0354]
- US 4797368 A [0355]
- US 5139941 A [0355]
- US 5173414 A [0355]
- US 5614404 A [0355]
- US 5658785 A [0355]
- US 5858775 A [0355]
- US 5994136 A [0355]
- US 5719 A [0356]
- US 054 A [0356]
- WO 9921979 A [0356]
- WO 0181609 A [0356]
- WO 0039318 A [0356]
- WO 0181553 A [0356]
- WO 9507994 A [0356]
- WO 9210578 A [0356]
- US 6261552 B [0357]
- US 5599691 A [0357]
- US 6326195 B [0358]
- US 5888502 A [0358]
- US 5580766 A [0358]
- US 5672510 A [0358]
- US 5922576 A [0359]
- US 5965358 A [0359]
- US 6168941 A [0359]
- WO 9822588 A [0359]
- WO 9856937 A [0359]
- WO 9915686 A [0359]
- WO 9954441 A [0359]
- WO 0032754 A [0359]
- WO 9207944 A [0360]
- WO 9813500 A [0360]
- WO 8908716 A [0360]
- US 5500362 A [0370]
- US 4376110 A [0377]
- US 4399216 A [0389]
- US 4634665 A [0389]
- US 5179017 A [0389]
- WO 8704462 A [0391]
- WO 8901036 A [0391]
- EP 338841 A [0391]
- US 4474893 A [0393]
- WO 9308829 A [0394]
- WO 9404690 A [0395]
- US 5260203 A [0399]
- US 5455030 A [0399]
- US 4881175 A [0399]
- US 5132405 A [0399]
- US 5091513 A [0399]
- US 5476786 A [0399]
- US 5013653 A [0399]
- US 5258498 A [0399]
- US 5482858 A [0399]
- WO 9945962 A [0405]
- US 5399163 A [0444]
- US 5383851 A [0444]
- US 5312335 A [0444]
- US 5064413 A [0444]
- US 4941880 A [0444]
- US 4790824 A [0444]
- US 4596556 A [0444]
- US 4487603 A [0444]
- US 4486194 A [0444]
- US 4447233 A [0444]
- US 4447224 A [0444]
- US 4439196 A [0444]
- US 4475196 A [0444]
- US 4522811 A [0462]
- US 5374548 A [0462]
- US 5399331 A [0462]
- US 5416016 A [0462]
- WO 2002100348 A [0524]
- WO 2004035607 A [0534]
- US 5968502 A [0543]
- US 6063630 A [0543]
- US 6187305 B [0543]
- EP 0505500 A [0543]
- US 6440735 B [0545]
- US 6713055 B [0545]
- US 5593972 A [0551]
- US 5703057 A [0551]
- US 5879687 A [0551]
- US 6235523 B [0551]
- US 6387888 B [0551]
- US 6719977 B [0570]
- US 5057313 A [0598]
- US 6331175 B [0599]
- US 4824659 A [0599]
- WO 2005103083 A2 [0679]
- WO 2005103083 A [0683]

Non-patent literature cited in the description

- LANDE R et al. *Cell Immunol*, 2002, vol. 220 (1), 30-8 [0010]
- AUSIELLO CM et al. *Tissue Antigens*, 2000, vol. 56 (6), 539-47 [0010]
- COTNER T et al. *Int J Immunopharmacol*, 1981, vol. 3 (3), 255-68 [0010]
- ELLIS, J. H. et al. *J. Immunol.*, 1995, vol. 155, 925-937 [0010]
- ZUBIAUR M et al. *J Immunol.*, 1997, vol. 159 (1), 193-205 [0010]
- MUNSHI et al. *J. Biol. Chem.*, 2000, vol. 275, 21566-21571 [0047] [0717]

- RIECHMANN, L. et al. *Nature*, 1998, vol. 332, 323-327 [0063]
- JONES, P. et al. *Nature*, 1986, vol. 321, 522-525 [0063]
- QUEEN, C. et al. *PNAS USA*, 1989, vol. 86, 10029-10033 [0063]
- DITZEL HJ. *J Immunol.*, 1996, vol. 157 (2), 739-49 [0064]
- BARBAS SM et al. *J. Am. Chem. Soc.*, 1994, vol. 116, 2161-2162 [0064]
- BARBAS SM et al. *Proc Natl Acad Sci USA*, 1995, vol. 92 (7), 2529-33 [0064]
- Fundamental Immunology. Raven Press, 1989 [0122] [0419]
- CHOTHIA ; LESK. *J. Mol. Biol.*, 1987, vol. 196, 901-917 [0123]
- KABAT et al. Sequences of Proteins of Immunological Interest. Public Health Service, National Institutes of Health, 1991 [0123]
- *PNAS USA*, 1993, vol. 90 (14), 6444-8 [0126]
- WARD et al. *Nature*, 1989, vol. 341, 544-546 [0127]
- BIRD et al. *Science*, 1988, vol. 242, 423-426 [0127] [0290]
- HUSTON et al. *PNAS*, 1988, vol. 85, 5879-5883 [0127]
- HOLLIGER, P. et al. *PNAS USA*, 1993, vol. 90, 6444-6448 [0132]
- POLJAK, R.J. et al. *Structure*, 1994, vol. 2, 1121-1123 [0132]
- MORRISON et al. *PNAS USA*, 1984, vol. 81, 6851-6855 [0136]
- CABILLY et al. *PNAS USA*, 1984, vol. 81, 3273-3277 [0136]
- BOULIANNE et al. *Nature*, 1984, vol. 312, 643-646 [0136]
- NEUBERGER et al. *Nature*, 1985, vol. 314, 268-270 [0136]
- SAHAGAN et al. *J. Immunol.*, 1986, vol. 137, 1066-1074 [0136]
- LIU et al. *PNAS*, 1987, vol. 84, 3439-3443 [0136]
- SUN et al. *PNAS USA*, 1987, vol. 84, 214-218 [0136]
- BETTER et al. *Science*, 1988, vol. 240, 1041-1043 [0136]
- HARLOW et al. Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 [0136] [0224] [0366] [0377]
- JONES et al. *Nature*, 1986, vol. 321, 522-525 [0137] [0261] [0264]
- RIECHMANN et al. *Nature*, 1988, vol. 332, 323-329 [0137]
- PRESTA. *Curr. Op. Struct. Biol.*, 1992, vol. 2, 593-596 [0137]
- Current Protocols in Molecular Biology. Greene Publishing and Wiley InterScience, 1987 [0155] [0378]
- GOEDDEL. Gene Expression Technology. Methods in Enzymology. Academic Press, 1990, vol. 185 [0162]
- E. MEYERS ; W. MILLER. *Comput. Appl. Biosci.*, 1988, vol. 4, 11-17 [0198]
- NEEDLEMAN ; WUNSCH. *J. Mol. Biol.*, 1970, vol. 48, 444-453 [0198]
- ALTSCHUL et al. *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0199]
- ALTSCHUL et al. *Nucleic Acids Res.*, 1997, vol. 25 (17), 3389-3402 [0199]
- CREIGHTON. Proteins: Structure and Molecular Properties. W.H. Freeman and Company, 1984 [0201]
- SCHIWECK et al. *J Mol Biol.*, 1997, vol. 268 (5), 934-51 [0210]
- MOREA. *Biophys Chem.*, 1997, vol. 68 (1-3), 9-16 [0210]
- SHIRAI et al. *FEBS Lett.*, 1996, vol. 399 (1-2), 1-8 [0210]
- SHIRAI et al. *FEBS Lett.*, 1999, vol. 455 (1-2), 188-97 [0210]
- RECKZO et al. *Protein Eng.*, 1995, vol. 8 (4), 389-95 [0210]
- EIGENBROT et al. *J Mol Biol.*, 1993, vol. 229 (4), 969-95 [0210]
- MARKS et al. *Methods Mol Biol.*, 2004, vol. 248, 327-43 [0211]
- AZRIEL-ROSENFELD et al. *J Mol Biol.*, 2004, vol. 335 (1), 177-92 [0211]
- PARK et al. *Biochem Biophys Res Commun*, 2000, vol. 275 (2), 553-7 [0211]
- KANG et al. *Proc Natl Acad Sci USA.*, 1991, vol. 88 (24), 11120-3 [0211]
- ZAHND et al. *J Biol Chem.*, 2004, vol. 279 (18), 18870-7 [0211]
- XU et al. *Chem Biol.*, 2002, vol. 9 (8), 933-42 [0211]
- BORDER et al. *Proc Natl Acad Sci USA.*, 2000, vol. 97 (20), 10701-5 [0211]
- CRAMERI et al. *Nat Med.*, 1996, vol. 2 (1), 100-2 [0211]
- COLLIGAN et al. Current Protocols in Immunology. Greene Publishing Assoc. and Wiley InterScience, 1992 [0224]
- MULLER. *Meth. Enzymol.*, 1983, vol. 92, 589-601 [0224] [0366]
- EHRLING H. *Analytical Biochemistry*, 1999, vol. 267 (2), 252-259 [0228]
- ENGEN, J.R. ; SMITH, D.L. *Anal. Chem.*, 2001, vol. 73, 256A-265A [0228]
- Ernst Schering Res Found Workshop, 2004, vol. 44, 149-67 [0228]
- HUANG et al. *Journal of Molecular Biology*, 1998, vol. 281 (1), 61-67 [0228]
- SAITO ; PATTERSON. *Methods*, 1996, vol. 9 (3), 516-24 [0228]
- DOWNWARD. *J Mass Spectrom*, 2000, vol. 35 (4), 493-503 [0229]
- KISELAR ; DOWNARD. *Anal Chem.*, 1999, vol. 71 (9), 1792-801 [0229]

- **MANCA.** *Ann Ist Super Sanita*, 1991, vol. 27 (1), 15-9 [0230]
- **SLOOTSTRA-JW et al.** *Mol-Divers*, 1996, vol. 1, 87-96 [0232]
- **POLYAK ; DEANS.** *Blood*, 2002, vol. 99, 3956-3962 [0233]
- **WANG ; YU.** *Curr Drug Targets*, 2004, vol. 5 (1), 1-15 [0234]
- **BURTON.** *Immunotechnology*, August 1995, vol. 1 (2), 87-94 [0234]
- **CORTESE et al.** *Immunotechnology*, 1995, vol. 1 (2), 87-94 [0234]
- **IRVING et al.** *Curr Opin Chem Biol.*, 2001, vol. 5 (3), 314-24 [0234]
- Epitope Mapping A Practical Approach. Oxford University Press, 2001 [0235]
- **HOSHINO et al.** *J. Immunol.*, 1997, vol. 158, 741-747 [0237]
- **FERRERO et al.** *BMC Immunology*, 2004, vol. 5, 21 [0237]
- **KOHLER et al.** *Nature*, 1975, vol. 256, 495 [0250] [0377]
- **CLACKSON et al.** *Nature*, 1991, vol. 352, 624-628 [0250] [0269]
- **MARKS et al.** *J. Mol. Biol.*, 1991, vol. 222, 581-597 [0250]
- **VERMA, R. et al.** *J.Immunol.Meth.*, 1998, vol. 216, 165-181 [0252] [0392]
- **POLLOCK et al.** *J.Immunol.Meth.*, 1999, vol. 231, 147-157 [0252] [0392]
- **FISCHER, R. et al.** *Biol.Chem.*, 1999, vol. 380, 825-839 [0252] [0392]
- **LONBERG, N. et al.** *Nature*, 1994, vol. 368, 856-859 [0254]
- **LONBERG, N.** Handbook of Experimental Pharmacology. 1994, vol. 113, 49-101 [0254]
- **LONBERG, N. ; HUSZAR, D.** *Intern. Rev. Immunol.*, 1995, vol. 13, 65-93 [0254]
- **HARDING, F. ; LONBERG, N.** *Ann. N.Y. Acad. Sci.*, 1995, vol. 764, 536-546 [0254]
- **TAYLOR, L. et al.** *Nucleic Acids Research*, 1992, vol. 20, 6287-6295 [0254]
- **CHEN, J. et al.** *International Immunology*, 1993, vol. 5, 647-656 [0254]
- **TUAILLON et al.** *J. Immunol.*, 1994, vol. 152, 2912-2920 [0254]
- **TAYLOR, L. et al.** *International Immunology*, 1994, vol. 6, 579-591 [0254]
- **FISHWILD, D. et al.** *Nature Biotechnology*, 1996, vol. 14, 845-851 [0254]
- **CHEN et al.** *EMBO J.*, 1993, vol. 12, 821-830 [0255] [0256]
- **FISHWILD et al.** *Nature Biotechnology*, 1996, vol. 14, 845-851 [0255] [0256]
- **CHEN et al.** *EMBO J.*, 1993, vol. 12, 811-820 [0256]
- **VAUGHAN et al.** *Nature Biotech*, 1996, vol. 14, 309 [0260]
- **HANES ; PLUCTHAU.** *PNAS USA*, 1997, vol. 94, 4937-4942 [0260]
- **PARMLEY ; SMITH.** *Gene*, 1988, vol. 73, 305-318 [0260]
- **SCOTT.** *TIBS*, 1992, vol. 17, 241-245 [0260]
- **CWIRLA et al.** *PNAS USA*, 1990, vol. 87, 6378-6382 [0260] [0269]
- **RUSSEL et al.** *Nucl. Acids Research*, 1993, vol. 21, 1081-1085 [0260]
- **HOOGENBOOM et al.** *Immunol. Reviews*, 1992, vol. 130, 43-68 [0260]
- **CHISWELL ; MCCAFFERTY.** *TIBTECH*, 1992, vol. 10, 80-84 [0260]
- **RIECHMANN et al.** *Nature*, 1988, vol. 332, 323-327 [0261] [0264]
- **VERHOEYEN et al.** *Science*, 1988, vol. 239, 1534-1536 [0261] [0264]
- **SIMS et al.** *J. Immunol.*, 1993, vol. 151, 2296 [0261]
- **CHOTHIA et al.** *J. Mol. Biol.*, 1987, vol. 196, 901 [0261]
- **CARTER et al.** *PNAS USA*, 1992, vol. 89, 4285 [0261]
- **PRESTA et al.** *J. Immunol.*, 1993, vol. 151, 2623 [0261]
- **WINTER ; HARRIS.** *Immunol Today*, 1993, vol. 14, 43-46 [0263]
- **WRIGHT et al.** *Crit. Reviews in Immunol*, 1992, 125-168 [0263]
- **LIU et al.** *PNAS USA*, 1987, vol. 84, 3439 [0265]
- *J.Immunol.*, 1987, vol. 139, 3521 [0265]
- **KABAT et al.** Sequences of Proteins of Immunological Interest. 1991 [0265]
- **FUCHS et al.** *Biol/Technology*, 1991, vol. 9, 1370-1372 [0269]
- **HAY et al.** *Hum. Antibod. Hybridomas*, 1992, vol. 3, 81-85 [0269]
- **HUSE et al.** *Science*, 1989, vol. 246, 1275-1281 [0269]
- **MCCAFFERTY et al.** *Nature*, 1990, vol. 348, 552-554 [0269] [0290]
- **GRIFFITHS et al.** *EMBO J*, 1993, vol. 12, 725-734 [0269]
- **HAWKINS et al.** *J. Mol. Biol.*, 1992, vol. 226, 889-896 [0269]
- **GRAM et al.** *PNAS USA*, 1992, vol. 89, 3576-3580 [0269]
- **GARRAD et al.** *Biol/Technology*, 1991, vol. 9, 1373-1377 [0269]
- **HOOGENBOOM et al.** *Nuc Acid Res*, 1991, vol. 19, 4133-4137 [0269]
- **BARBAS et al.** *PNAS USA*, 1991, vol. 88, 7978-7982 [0269]
- **SCOTT, J. K. ; SMITH, G. P.** *Science*, 1990, vol. 249, 386-390 [0269]
- **FELICI et al.** *J. Mol. Biol.*, 1991, vol. 222, 301-310 [0269]
- **KUWABARA et al.** *Nature Biotechnology*, 1997, vol. 15, 74-78 [0269]

- **BARBAS ; BURTON.** *Trends. Biotechnol.*, 1996, vol. 14, 230-234 [0272]
- **AUJAME et al.** *Hum. Antibodies*, 1997, vol. 8, 155-68 [0272]
- **HAARD et al.** *J. Biol. Chem.*, 1999, vol. 274 (26), 18218-18230 [0272]
- **CUNNINGHAM ; WELLS.** *Science*, 1989, vol. 244, 1081-1085 [0274]
- **REIDHAAR-OLSON ; SAUER.** *Science*, 1988, vol. 241, 53-57 [0274]
- **BOWIE ; SAUER.** *PNAS USA*, 1989, vol. 86, 2152-2156 [0274]
- **Proteins, Structures and Molecular Principles.** W. H. Freeman and Company, 1984 [0276]
- **Introduction to Protein Structure.** Garland Publishing, 1991 [0276]
- **THORNTON.** *Nature*, 1991, vol. 354, 105 [0276]
- **COLLINET et al.** *J Biol Chem*, 2000, vol. 275 (23), 17428-33 [0276]
- **SHIELDS et al.** *J. Biol. Chem.*, 2001, vol. 276, 6591-6604 [0279]
- **SHIELD et al.** *J. Biol. Chem.*, 2002, vol. 277, 26733 [0284]
- **UMANA et al.** *Nat. Biotechnol.*, 1999, vol. 17, 176 [0284]
- **BALINT ; LARRICK.** *Gene*, 1993, vol. 137, 109-118 [0285]
- **WU et al.** *PNAS USA*, 1998, vol. 95, 6037-42 [0285]
- **STUDNICKA et al.** *Protein Engineering*, 1994, vol. 7, 805-814 [0286]
- **SODERLIND et al.** *Immunotechnology*, 1999, vol. 4 (3-4), 279-85 [0286]
- **YANG et al.** *Journal of Molecular Biology*, 1995, vol. 254 (3), 392-403 [0286]
- **EVANS et al.** *J. Immunol. Meth.*, 1995, vol. 184, 123-38 [0289]
- **PLUCKTHUN.** *The Pharmacology of Monoclonal Antibodies.* Springer-Verlag, 1994, vol. 113, 269-315 [0290]
- **HUSTON et al.** *PNAS USA*, 1988, vol. 85, 5879-5883 [0290]
- **GRAZIANO, R.F. et al.** *J. Immunol.*, 1995, vol. 155 (10), 4996-5002 [0294]
- **MORTON, H.C. et al.** *Critical Reviews in Immunology*, 1996, vol. 16, 423-440 [0295]
- **MONTEIRO, R.C. et al.** *J. Immunol.*, 1992, vol. 148, 1764 [0295]
- **FIELD et al.** *Mol. Cell. Biol.*, 1988, vol. 8, 2159-2165 [0305]
- **EVAN et al.** *Mol. Cell. Biol.*, 1985, vol. 5 (12), 3610-3616 [0305]
- **PABORSKY et al.** *Protein Engineering*, 1990, vol. 3 (6), 547-553 [0305]
- **WALKER.** *Protein Protocols On Cd-Rom.* Humana Press, 1998 [0306]
- **DIENER et al.** *Science*, 1986, vol. 231, 148 [0308]
- **JUNGHANS et al.** *Cancer Chemotherapy and Biotherapy.* Lippincott Raven, 1996, 655-686 [0309]
- **ADELSTEIN et al.** *Nucl. Med. Biol.*, 1987, vol. 14, 165-169 [0310]
- **Radioimmunoglobulin therapy.** **VRIESENDORP et al.** *High Dose Cancer Therapy Armitage.* Williams & Wilkins, 1992 [0310]
- **WILBUR.** *Antibiot. Immunoconjug. Radiopharm.*, 1991, vol. 4, 5-97 [0311]
- **GREENWOOD et al.** *Biochem J.*, 1963, vol. 89, 114-123 [0313]
- **SALACINSKI et al.** *Anal. Biochem.*, 1981, vol. 117, 136-146 [0313]
- **GRIFFITHS et al.** *Cancer Res.*, 1991, vol. 51, 4594-4602 [0313]
- **Antibody Engineering Handbook.** Chijin Shokan, 1994 [0318]
- **ZHANG et al.** *J Biol Chem.*, 2004, vol. 279 (3), 2053-62 [0320]
- **FORSTER ; ALTMAN.** *Science*, 1990, vol. 238, 407-409 [0325]
- **NISHIKURA.** *Cell*, 2001, vol. 107 (4), 415-8 [0326]
- **FJOSE et al.** *Biotechnol Annu Rev.*, 2001, vol. 7, 31-57 [0326]
- **HANON.** *Nature*, 2002, vol. 418, 244-51 [0326]
- **BRANTL.** *Biochim Biophys Acta*, 2002, vol. 1575 (1-3), 15-25 [0326]
- **TUSCHL.** *Chembiochem*, 2001, vol. 2 (4), 239-45 [0326]
- **CAPLEN.** *Expert Opin Biol Ther*, 2003, vol. 3 (4), 575-86 [0326]
- **LU et al.** *Curr Opin Mol Ther*, 2003, vol. 5 (3), 225-34 [0326]
- **SHUEY et al.** *Drug Discov Today*, 2002, vol. 7 (20), 1040-6 [0326]
- **SHI.** *Trends Genet*, 2003, vol. 19 (1), 9-12 [0326]
- **KOVAR et al.** *Semin Cancer Biol.*, 2003, vol. 13 (4), 275-81 [0326]
- **LAVREY et al.** *Curr Opin Drug Discov Devel*, 2003, vol. 6 (4), 561-9 [0326]
- **CLEWEY.** *Commun Dis Public Health*, 2003, vol. 6 (2), 162-3 [0326]
- **DUXBURY et al.** *J Surg Res.*, 2004, vol. 117 (2), 339-44 [0326]
- **CAPLEN et al.** *Ann N Y Acad Sci.*, 2003, vol. 1002, 56-62 [0326]
- **HUNTER et al.** *Nature*, 1962, vol. 144, 945 [0327]
- **DAVID et al.** *Biochemistry*, 1974, vol. 13, 1014 [0327]
- **PAIN et al.** *J. Immunol. Meth.*, 1981, vol. 40, 219 [0327]
- **NYGREN.** *J. Histochem. and Cytochem.*, 1982, vol. 30, 407 [0327]
- **HISKEY.** *Peptides*, 1981, vol. 3, 137 [0327]
- **MAASEN et al.** *Eur. J. Biochem.*, 1983, vol. 134, 32 [0327]
- **MEANS et al.** *Chemical modifications of proteins.* Holden-Day, 1971, 105-110 [0328]
- **Jl.** *Methods Enzymol*, 1983, vol. 91, 580-609 [0328]
- **MEARES et al.** *Anal. Biochem.*, 1984, vol. 142, 68-78 [0328]

- **KOYAMA.** *Chem. Abstr.*, 1994, vol. 120, 217262t [0328]
- **RODWELL et al.** *PNAS USA*, 1986, vol. 83, 2632-2636 [0328]
- **QUADRI et al.** *Nucl. Med. Biol.*, 1993, vol. 20, 559-570 [0328]
- **SUNDBERG et al.** *Nature*, 1974, vol. 250, 587-588 [0328]
- **CHEN et al.** *J Biol Chem*, 1991, vol. 266, 18237-18243 [0329]
- **BROWNING et al.** *J. Immunol.*, 1989, vol. 143, 1859-1867 [0329]
- **MYERS et al.** *J. Immunol. Meth.*, 1989, vol. 121, 129-142 [0329]
- **YU et al.** *Int. J. Cancer*, 1994, vol. 56, 244 [0330]
- **WONG.** *Chemistry Of Protein Conjugation And Cross-Linking.* CRC Press, 1991 [0330]
- *Modification of Antibodies by Chemical Methods.* **UP-ESLACIS et al.** *Monoclonal Antibodies: Principles And Applications.* Wiley-Liss, Inc, 1995 [0330]
- *Production and Characterization of Synthetic Peptide-Derived Antibodies.* **PRICE et al.** *Monoclonal Antibodies: Production, Engineering And Clinical Application.* Cambridge University Press, 1995 [0330]
- **GOODMAN et al.** *Goodman and Gilman's The Pharmacological Basis Of Therapeutics.* Macmillan Publishing Co, 1990 [0335]
- **VITETTA.** *Immunol. Today*, 1993, vol. 14, 252 [0335]
- **PASTAN et al.** *Cell*, 1986, vol. 47, 641 [0339]
- **GOLDENBERG ; CALIF.** *A Cancer Journal for Clinicians*, 1994, vol. 44, 43 [0339]
- *Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy.* **ARNON et al.** *Monoclonal Antibodies And Cancer Therapy.* Alan R. Liss, Inc, 1985, 243-56 [0341]
- *Antibodies For Drug Delivery.* **HELLSTROM et al.** *Controlled Drug Delivery.* Marcel Dekker, Inc, 1987, 623-53 [0341]
- *Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review.* **THORPE et al.** *Monoclonal Antibodies '84: Biological And Clinical Applications.* 1985, 475-506 [0341]
- *Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy.* *Monoclonal Antibodies For Cancer Detection And Therapy.* Academic Press, 1985, 303-16 [0341]
- **THORPE et al.** *The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates.* *Immunol. Rev.*, 1982, vol. 62, 119-58 [0341]
- **HOCH et al.** *PNAS USA*, 1985, vol. 82, 1692 [0342]
- **COLOMBATTI et al.** *J. Biol. Chem.*, 1986, vol. 261, 3030 [0342]
- **DELEERS et al.** *FEBS Lett.*, 1983, vol. 160, 82 [0342]
- **HWANG et al.** *Cell*, 1987, vol. 48, 129 [0342]
- **GRAY et al.** *PNAS USA*, 1984, vol. 81, 2645 [0342]
- **CALDERWOOD et al.** *PNAS USA*, 1987, vol. 84, 4364 [0342]
- **JAIME et al.** *Anticancer Res.*, 2001, vol. 21 (2A), 1119-28 [0345]
- **DAMLE et al.** *Curr Opin Pharmacol*, 2003, vol. 3 (4), 386-90 [0345]
- **TRAIL et al.** *Cancer Immunol Immunother*, 2003, vol. 52 (5), 328-37 [0345]
- **SYKES ; JOHNSTON.** *Nat Biotech*, 1997, vol. 17, 355-59 [0349]
- **SCHAKOWSKI et al.** *Mol Ther*, 2001, vol. 3, 793-800 [0349]
- **BENVENISTY ; RESHEF.** *PNAS USA*, 1986, vol. 83, 9551-55 [0349]
- **WIGLER et al.** *Cell*, 1978, vol. 14, 725 [0349]
- **CORARO ; PEARSON.** *Somatic Cell Genetics*, 1981, vol. 7, 603 [0349]
- **VAN HEEKE ; SCHUSTER.** *J Biol Chem*, 1989, vol. 264, 5503-5509 [0350]
- **F. AUSUBEL et al.** *Current Protocols in Molecular Biology.* Greene Publishing and Wiley InterScience, 1987 [0351]
- **GRANT et al.** *Methods in Enzymol*, 1987, vol. 153, 516-544 [0351]
- *Virology.* Raven Press, Ltd, 1996 [0354]
- *VIROLOGY.* 2001 [0354]
- *Encyclopedia of Virology.* Academic Press, 1999 [0354]
- *Fundamental Virology.* Lippincott- Raven, 1995 [0354]
- **LEVINE.** *Viruses.* Scientific American Library, 1992 [0354]
- *Medical Virology.* Acad. Press, 1994 [0354]
- *Introduction to Modern Virology.* Blackwell Scientific Publications, Ltd, 1994 [0354]
- **CARTER.** *Curr Opinion Biotech*, 1992, vol. 3, 533-539 [0355] [0356]
- **MUZCZYKA.** *Curr Top Microbiol Immunol*, 1992, vol. 158, 97-129 [0355] [0356]
- **CARTER.** *Contrib. Microbiol.*, 2000, vol. 4, 85-86 [0355]
- **SMITH-ARICA.** *Curr. Cardiol. Rep.*, 2001, vol. 3 (1), 41-49 [0355]
- **TAJ.** *J. Biomed. Sci.*, 2000, vol. 7 (4), 279-91 [0355]
- **VIGNA et al.** *J. Gene Med.*, 2000, vol. 2 (5), 308-16 [0355]
- **KLIMATCHEVA et al.** *Front. Biosci.*, 1999, vol. 4, D481-96 [0355]
- **LEVER et al.** *Biochem. Soc. Trans.*, 1999, vol. 27 (6), 841-47 [0355]
- **SNYDER.** *J Gene Med.*, 1999, vol. 1 (3), 166-75 [0355]
- **GERICH et al.** *Knee Surg. Sports Traumatol. Arthrosc.*, 1998, vol. 5 (2), 118-23 [0355]
- *During.* *Adv. Drug Deliv. Review*, 1997, vol. 27 (1), 83-94 [0355]
- **LAUGHLIN et al.** *Gene*, 1983, vol. 23, 65-73 [0355]
- **HEWSON.** *Mol Med Today*, 1999, vol. 5 (1), 8 [0356]
- **STEPHENS.** *Biochem J.*, 1987, vol. 248 (1), 1-11 [0356]

- **SCHLESINGER.** *Expert Opin Biol Ther.*, 2001, vol. 1 (2), 177-91 [0356]
- **POLO et al.** *Dev Biol*, 2000, vol. 104, 181-5 [0356]
- **WAHLFORS et al.** *Gene Ther.*, 2000, vol. 7 (6), 472-80 [0356]
- **COLOMBAGE et al.** *Virology*, 1998, vol. 250 (1), 151-63 [0356]
- **LACHMANN et al.** *Curr Opin Mol Ther*, 1999, vol. 1 (5), 622-32 [0357]
- **FRAEFEL et al.** *Adv Virus Res.*, 2000, vol. 55, 425-51 [0357]
- **HUARD et al.** *Neuromuscul*, 1997, vol. 7 (5), 299-313 [0357]
- **GLORIOSO et al.** *Annu Rev Microbiol*, 1995, vol. 49, 675-710 [0357]
- **LATCHMAN.** *Mol Biotechnol*, 1994, vol. 2 (2), 179-95 [0357]
- **FRENKEL et al.** *Gene Ther*, 1994, vol. 1 (1), 40-6 [0357]
- **MILLER.** *Curr Top Microbiol Immunol*, 1992, vol. 158, 1-24 [0358]
- **SALMONS ; GUNZBURG.** *Human Gene Therapy*, 1993, vol. 4, 129-141 [0358]
- **MILLER et al.** *Methods in Enzymology*, 1994, vol. 217, 581-599 [0358]
- **WEBER et al.** *Curr Opin Mol Ther*, 2001, vol. 3 (5), 439-53 [0358]
- **HU et al.** *Pharmacol Rev.*, 2000, vol. 52 (4), 493-511 [0358]
- **KIM et al.** *Adv Virus Res.*, 2000, vol. 55, 545-63 [0358]
- **PALU et al.** *Rev Med Virol*, 2000, vol. 10 (3), 185-202 [0358]
- **TAKEUCHI et al.** *Adv Exp Med Biol.*, 2000, vol. 465, 23-35 [0358]
- **GRAHAM et al.** *Mol Biotechnol*, 1995, vol. 33 (3), 207-220 [0359]
- **STEPHENSON.** *Clin Diagn Virol*, 1998, vol. 10 (2-3), 187-94 [0359]
- **JACOBS.** *Clin Sci*, 1993, vol. 85 (2), 117-22 [0359]
- **JOLLY.** *Cancer Gene Therapy*, 1994, vol. 1, 51-64 [0359]
- **LATCHMAN.** *Molec Biotechnol*, 1994, vol. 2, 179-195 [0359]
- **JOHANNING et al.** *Nucl Acids Res*, 1995, vol. 23, 1495-1501 [0359]
- **BERENCSI et al.** *J Infect Dis*, 2001, vol. 183 (8), 1171-9 [0360]
- **ROSENWIRTH et al.** *Vaccine*, 2001, vol. 19 (13-14), 1661-70 [0360]
- **KITTLESEN et al.** *J Immunol*, 2000, vol. 164 (8), 4204-11 [0360]
- **BROWN et al.** *Gene Ther*, 2000, vol. 7 (19), 1680-9 [0360]
- **KANESA-THASAN et al.** *Vaccine*, 2000, vol. 19 (4-5), 483-91 [0360]
- **STEN.** *Drua*, 2000, vol. 60 (2), 249-71 [0360]
- **VENUGOPAL et al.** *Res Vet Sci*, 1994, vol. 57 (2), 188-193 [0360]
- **MOSS.** *Dev Biol Stand*, 1994, vol. 82, 55-63 [0360]
- **WEISZ et al.** *Mol Cell Biol*, 1994, vol. 43, 137-159 [0360]
- **MAHR ; PAYNE.** *Immunobiology*, 1992, vol. 184 (2-3), 126-146 [0360]
- **HRUBY.** *Clin Microbiol Rev*, 1990, vol. 3 (2), 153-170 [0360]
- Current Protocols in Immunology. Greene Publishing Assoc. and Wiley InterScience, 1992 [0366]
- **MUNSON ; POLLARD.** *Anal. Biochem.*, 1980, vol. 107, 220 [0368]
- **AUSIELLO et al.** *Tissue antigens*, 2000, vol. 56, 538-547 [0372] [0704] [0705] [0706]
- Molecular cloning. Cold Spring Harbor Laboratory Press, 2001 [0376]
- **HARLOW ; LANE.** Using Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1999 [0377]
- Current Protocols In Molecular Biology. Greene Publishing Assoc. and Wiley InterScience, 1987 [0377]
- **GODING.** Monoclonal Antibodies: Principles and Practice. Academic Press, 1986, 59-103 [0377]
- Continuously Proliferating Human Cell Lines Synthesizing Antibody of Predetermined Specificity. **ZURAWAKI, V. R. et al.** Monoclonal Antibodies. Plenum Press, 1980, 19-33 [0377]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0378]
- MOLECULAR CLONING : A LABORATORY MANUAL. 2001 [0378]
- **WILDT S et al.** *Nat Rev Microbiol*, 2005, vol. 3 (2), 119-28 [0380]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495 [0382]
- **MORRISON, S.** *Science*, 1985, vol. 229, 1202 [0386]
- **URLAUB ; CHASIN.** *PNAS USA*, 1980, vol. 77, 4216-4220 [0391]
- **R. J. KAUFMAN ; P. A. SHARP.** *Mol. Biol.*, 1982, vol. 159, 601-621 [0391]
- **D. M. KRANZ et al.** *PNAS USA*, 1981, vol. 78, 5807 [0393]
- **SONGSIVILAI ; LACHMANN.** *Clin. Exp. Immunol.*, 1990, vol. 79, 315-321 [0394]
- **KOSTELNY et al.** *J. Immunol.*, 1992, vol. 148, 1547-1553 [0394]
- **MILSTEIN ; CUELLO.** *Nature*, 1983, vol. 305, 537 [0394]
- **TRAUNECKER et al.** *EMBO J.*, 1991, vol. 10, 3655 [0394]
- **SURESH et al.** *Methods in Enzymology*, 1986, vol. 121, 210 [0395]
- **KARPOVSKY et al.** *J. Exp. Med.*, 1984, vol. 160, 1686 [0397]
- **LIU, M. A. et al.** *PNAS USA*, 1985, vol. 82, 8648 [0397]

- **BRENNAN et al.** *Science*, 1985, vol. 229, 81 [0397]
- **SHALABY et al.** *J. Exp. Med.*, 1992, vol. 175, 217-225 [0397]
- **PAULUS.** *Behring Ins. Mitt.*, 1985, 118-132 [0397]
- **GLENNIE et al.** *J. Immunol.*, 1987, vol. 139, 2367-2375 [0397]
- **KOSTELNY et al.** *J. Immunol.*, 1992, vol. 148 (5), 1547-1553 [0400]
- **HOLLINGER et al.** *PNAS USA*, 1993, vol. 90, 6444-6448 [0400]
- **GRUBER et al.** *J. Immunol.*, 1994, vol. 152, 5368 [0400]
- **HOLLIGER et al.** *PNAS USA*, 1993, vol. 90, 6444-6448 [0401]
- **TRAUNECKER et al.** *EMBO J*, 1991, vol. 10, 3655-3659 [0401]
- **TRAUNECKER et al.** *Int J Cancer*, 1992, vol. 7, 51-52 [0401]
- **WEINTRAUB, B.** Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques. The Endocrine Society, March 1986 [0402]
- **KOZAK.** *J. Biol. Chem.*, 1991, vol. 266, 19867-19870 [0406]
- **MILLS et al.** *Nucl. Acids Res.*, 1991, vol. 15, 7305-7316 [0420]
- **SIDERAS et al.** *Intl. Immunol.*, 1989, vol. 1, 631-642 [0420]
- Remington: The Science and Practice of Pharmacy. Mack Publishing Co, 1995 [0429]
- **YANG et al.** *PNAS USA*, 2003, vol. 100 (12), 6934-6939 [0443]
- **BERGE, S.M. et al.** *J. Pharm. Sci.*, 1977, vol. 66, 1-19 [0446]
- **BAEK et al.** *Methods Enzymol.*, 2003, vol. 362, 240-9 [0457]
- **NIGAVEKAR et al.** *Pharm Res.*, 2004, vol. 21 (3), 476-83 [0457]
- **POWELL et al.** Compendium of excipients for parenteral formulations. *PDA J Pharm Sci Technol.*, 1998, vol. 52, 238-311 [0458]
- Sustained and Controlled Release Drug Delivery Systems. Marcel Dekker, Inc, 1978 [0459]
- **STREJAN et al.** *J. Neuroimmunol.*, 1984, vol. 7, 27 [0460] [0461]
- **V.V. RANADE.** *J. Clin. Pharmacol.*, 1989, vol. 29, 685 [0462]
- **UMEZAWA et al.** *Biochem. Biophys. Res. Commun.*, 1988, vol. 153, 1038 [0462]
- **P.G. BLOEMAN et al.** *FEBS Lett.*, 1995, vol. 357, 140 [0462]
- **M. OWAIS et al.** *Antimicrob. Agents Chemother.*, 1995, vol. 39, 180 [0462]
- **BRISCOE et al.** *Am. J. Physiol.*, 1995, vol. 1233, 134 [0462]
- **SCHREIER et al.** *J. Biol. Chem.*, 1994, vol. 269, 9090 [0462]
- **K. KEINANEN ; M.L. LAUKKANEN.** *FEBS Lett.*, 1994, vol. 346, 123 [0462]
- **J.J. KILLION ; I.J. FIDLER.** *Immunomethods*, 1994, vol. 4, 273 [0462]
- **CUNNINGHAM-RUNDLES C et al.** *J Immunol Methods*, 1992, vol. 152, 177-190 [0464]
- **LANDORM.** *Ann Allergy Asthma Immunol*, 1995, vol. 74, 279-283 [0464]
- **FELSON et al.** American College of Rheumatology Preliminary Definition of Improvement in Rheumatoid Arthritis. *Arthritis Rheumatism*, 1995, vol. 38, 727-735 [0501]
- **BIREBENT et al.** *Vaccine*, 2003, vol. 21 (15), 1601-12 [0538]
- **LI et al.** *Chin Med J*, 2001, vol. 114 (9), 962-6 [0538]
- **SCHMITT et al.** *Hybridoma*, 1994, vol. 13 (5), 389-96 [0538]
- **MALONEY et al.** *Hybridoma*, 1985, vol. 4 (3), 191-209 [0538]
- **RAYCHARDHURI et al.** *J Immunol.*, 1986, vol. 137 (5), 1743-9 [0538]
- **POHL et al.** *Int J Cancer*, 1992, vol. 50 (6), 958-67 [0538]
- **BOHLEN et al.** *Cytokines Mol Ther.*, 1996, vol. 2 (4), 231-8 [0538]
- **MARUYAMA.** *J Immunol Methods*, 2002, vol. 264 (1-2), 121-33 [0538]
- **OCHI et al.** *Eur J Immunol*, 1987, vol. 17 (11), 1645-8 [0538]
- **WI et al.** *J Immunol Methods*, 1989, vol. 122 (2), 227-34 [0538]
- **PHAN et al.** *PNAS USA*, 2003, vol. 100, 8372 [0547]
- **LIN et al.** *Curr Cancer Drug Targets*, 2001, vol. 1 (3), 241-7 [0550]
- *Curr Cancer Drug Targets*, 2003, vol. 3 (3), 237 [0550]
- **LIMA et al.** *Cancer Gene Ther.*, 2004, vol. 11 (5), 309-16 [0550]
- **GRZMIL et al.** *Int J Oncol*, 2004, vol. 4 (1), 97-105 [0550]
- **COLLIS et al.** *Int J Radiat Oncol Biol Phys.*, 2003, vol. 57 (2), S144 [0550]
- **YANG et al.** *Oncogene*, 2003, vol. 22 (36), 5694-701 [0550]
- **ZHANG et al.** *Biochem Biophys Res Commun*, 2003, vol. 303 (4), 1169-78 [0550]
- **KOWALCZYK et al.** *Acta Biochim Pol.*, 2003, vol. 50 (3), 613-24 [0551]
- **REILLY et al.** *Methods Mol Med.*, 2002, vol. 69, 233-57 [0551]
- **TIRAPU et al.** *Curr Gene Ther.*, 2002, vol. 2 (1), 79-89 [0551]
- **SHAH et al.** *J Neurooncol*, 2003, vol. 65 (3), 203-26 [0552]
- **STILES et al.** *Surgery*, 2003, vol. 134 (2), 357-64 [0552]
- **SUNARMURA et al.** *Pancreas*, 2004, vol. 28 (3), 326-9 [0552]

- **TESHIGAHARA et al.** *J Surg Oncol.*, 2004, vol. 85 (1), 42-7 [0552]
- **VARGHESE et al.** *Cancer Gene Ther.*, 2002, vol. 9 (12), 967-78 [0552]
- **WILDNER et al.** *Cancer Res.*, 1999, vol. 59 (2), 410-3 [0552]
- **YAMANAKA.** *Int J Oncol.*, 2004, vol. 24 (4), 919-23 [0552]
- **ZWIEBEL et al.** *Semin Oncol.*, 2001, vol. 28 (4), 336-43 [0552]
- **BYSTRYN et al.** *Clinical Cancer Research*, July 2001, vol. 7, 1882-1887 [0553]
- **ZHAO et al.** *J Immunol Methods*, 2001, vol. 254 (1-2), 137-45 [0570]
- **HONG et al.** *Cancer Res.*, 2000, vol. 60 (23), 6551-6 [0570]
- **LINDGREN et al.** *Biochem J.*, 2004, vol. 377 (1), 69-76 [0570]
- **BUERGER et al.** *J Cancer Res Clin Oncol.*, 2003, vol. 129 (12), 669-75 [0570]
- **POOGA et al.** *FASEB J.*, 1998, vol. 12 (1), 67-77 [0570]
- **TSENG et al.** *Mol Pharmacol.*, 2002, vol. 62 (4), 864-72 [0570]
- **ZHANG et al.** *J Control Release*, 2003, vol. 93 (2), 141-50 [0587]
- **KAMBE et al.** *Hum Cell*, 1997, vol. 10 (1), 87-94 [0587]
- **ROUDEBUSH et al.** *Vet Clin North Am Small Anim Pract.*, 2004, vol. 34 (1), 249-69 [0587]
- **RAFI.** *Nutrition*, 2004, vol. 20 (1), 78-82 [0587]
- Radiolabeled Monoclonal Antibodies For Imaging And Therapy. Plenum Press, 1988 [0595]
- Medical Applications of Radioisotopes. **CHASE et al.** Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990, 624-652 [0595]
- Clinical Use of Monoclonal Antibodies. **BROWN et al.** Biotechnology And Pharmacy. Chapman & Hall, 1993, 227-49 [0595]
- **WETTERWALD et al.** *American Journal of Pathology*, 2002, vol. 160 (3), 1143-1153 [0659]
- **GEYSEN et al.** Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid. *Proc Natl Acad Sci USA*, 1984, vol. 81, 3998 [0708]
- **SLOOTSTRA et al.** Structural aspects of antibody-antigen interaction revealed through small random peptide libraries. *Mol Divers*, 1996, vol. 1, 87 [0708]
- **PUIJK et al.** Segment synthesis. *PCT*, 2001, 1 [0708]
- **GRAEFF et al.** *J. Biol. Chem.*, 1994, vol. 269, 30260-30267 [0713]

CD38-ELLENES ANTITESTEK A MIELÓMA MULTIPLEX KEZELÉSÉRE
SZABADALMI IGÉNYPONTOK

1. Antitest, amely a humán CD38-hoz kötődik, ahol az említett antitest humán könnyűlánc és humán nehézlánc variábilis régiókat tartalmaz, ahol a könnyűlánc variábilis régiója tartalmaz egy V_L CDR1-et, aminek szekvenciáját a SEQ ID No:13 határozza meg, egy V_L CDR2-t, aminek szekvenciáját a SEQ ID No:14 határozza meg és egy V_L CDR3-at, aminek szekvenciáját a SEQ ID No:15 határozza meg, és a nehézlánc variábilis régiója tartalmaz egy V_H CDR1-et, aminek szekvenciáját a SEQ ID No:18 határozza meg, egy V_H CDR2-t, aminek szekvenciáját a SEQ ID No:19 határozza meg és egy V_H CDR3-at, aminek szekvenciáját a SEQ ID No:20 határozza meg.
2. Az 1. igénypont szerinti antitest, ami humán monoklonális antitest.
3. Az 1. és a 2. igénypontok bármelyike szerinti antitest **azzal jellemezve, hogy ez egy teljes hosszúságú IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE vagy IgM antitest.**
4. A 3. igénypont szerinti antitest, ami egy IgG1 antitest.
5. A 4. igénypont szerinti antitest, ami egy IgG1_K antitest.
6. A 3. igénypont szerinti antitest, ami egy IgM antitest.
7. A 6. igénypont szerinti antitest, ami egy IgM_K antitest.
8. Az 1-7. igénypontok bármelyike szerinti antitest, ami tartalmaz egy olyan V_H régiót, aminek aminosav-szekvenciáját a SEQ ID No:17 határozza meg.
9. Az 1-7. igénypontok bármelyike szerinti antitest, ami tartalmaz egy olyan V_H régiót, ami a SEQ ID No:17 V_H CDR1-V_H CDR3-ig terjedő régiójának aminosav-szekvenciáját tartalmazza.
10. Az 1-9. igénypontok bármelyike szerinti antitest, ami tartalmaz egy olyan V_L régiót, aminek aminosav-szekvenciáját a SEQ ID No:12 határozza meg.
11. Az 1. igénypont szerinti antitest, ahol az antitestet a következő kódolja:
 - (i) a humán könnyűlánc és a humán nehézlánc nukleinsavai, amik sorrendben a SEQ ID No:11 és a SEQ ID No:16 által meghatározott nukleotidszekvenciákat tartalmazzák a variábilis régiókban; vagy
 - (ii) a humán könnyűlánc és a humán nehézlánc nukleinsavai, amik a variábilis régiókban az (i) pontban meghatározott szekvenciák konzervatív szekvenciamódosulatait tartalmazzák.
12. Az előző igénypontok bármelyike szerinti antitest, ami egy antitestfragmentum vagy egy egyláncú antitest.
13. Az előző igénypontok bármelyike szerinti antitest, ami lényegében izolált formában van.
14. Az előző igénypontok bármelyike szerinti antitest, ami még egy kelátor linkert is tartalmaz radioizotóp kapcsolásához.
15. Izolált nukleinsav, amely az 1-12. igénypontok bármelyike szerinti antitestet kódolja.
16. Expressziós vektor, amely az 1-12. igénypontok bármelyike szerinti antitestet kódoló nukleinsav-szekvenciát tartalmaz.
17. Eukarióta vagy prokarióta gazdasejt, amely az 1-12. igénypontok bármelyike szerinti antitestet termel.
18. Immunkonjugátum, ami az 1-13. igénypontok bármelyike szerinti antitestet foglalja magában, ami egy citotoxikus agenshez, a radioizotóphoz vagy egy gyógyszerhez kapcsolódik.
19. Bispecifikus vagy multispecifikus molekula, ami az 1-14. igénypontok bármelyike szerinti antitestet foglalja magában és kötési specificitása van a humán effektorsejtek felé.
20. Gyógyászati készítmény, ami tartalmazza az 1-14. igénypontok bármelyike szerinti antitestet, a 18. igénypont szerinti immunkonjugátumot vagy a 19. igénypont szerinti bispecifikus vagy multispecifikus molekulát és egy gyógyászatiilag elfogadható hordozót.



SZTNH-100072037

21. Az 1-14. igénypontok bármelyike szerinti antitest, a 18. igénypont szerinti immunkonjugátum, a 19. igénypont szerinti bispecifikus vagy multispecifikus molekula vagy a 16. igénypont szerinti expressziós vektor gyógyszerként való felhasználásra.
22. Az 1-14. igénypontok bármelyike szerinti antitest, a 18. igénypont szerinti immunkonjugátum, a 20. igénypont szerinti gyógyászati készítmény, a 19. igénypont szerinti bispecifikus vagy multispecifikus molekula vagy a 16. igénypont szerinti expressziós vektor olyan betegség vagy rendellenesség kezelésében vagy megelőzésében való felhasználásra, amelyben CD38-at expresszáló sejtek vannak, ahol a betegség vagy rendellenesség az alábbi csoportból választott: B-sejtes limfóma/leukémiák beleértve a prekursor B-limfoblasztos leukémiát/limfómát és a B-sejtes non-Hodgkin limfómákat; akut limfoblasztos leukémia; érett B-sejtes neoplazmák, beleértve a B-sejtes krónikus limfocitás leukémiát (CLL)/kis limfocitás limfómát (SLL); B-sejtes akut limfocitás leukémia; B-sejtes prolimfocitás leukémia; limfoplaszmatikus limfóma; köpenysejtes limfóma (MCL); folliculáris limfóma (FL), beleértve az alacsony, közepes és magas fokozatú FL-t; cutan folliculus centrum limfóma; marginális zónájú B-sejtes limfóma, beleértve a MALT típust, a nyirokcsomó és a lép típust is; hajás sejt leukémia; diffúz nagy B-sejtes limfóma; Burkitt-limfóma; plazmacitoma; plazmasejtes mieloma; plazmasejtes leukémia; transzplantáció utáni limfoproliferatív rendellenesség; Waldenström fele makroglobulinémia; plazmasejtes leukémiák; anaplasztikus nagysejtes limfóma (ALCL); mieloma multiplex; Hodgkin-limfóma; érett T-sejtes és NK-sejtes neoplazmák, beleértve a T-sejtes prolimfocitás leukémiát is; T-sejtes nagy granuláris limfocitás leukémia; agresszív NK-sejtes leukémia; felnőtt T-sejtes leukémia/limfóma; extranodális NK/T-sejtes limfóma; nazális típusú, enteropathia típusú T-sejtes limfóma, hepatosplenikus T-sejtes limfóma; szubkután panniculitis-szerű T-sejtes limfóma; blasztos NK-sejtes limfóma; Mycosis Fungoides/Sézary szindróma; primer kután CD30 pozitív T-sejtes limfoproliferatív betegségek, beleértve a kután anaplasztikus nagysejtes limfómát C-ALCL, limfomatoid papulosis és a borderline léziókat; angioimmunoblasztikus T-sejtes limfóma, nem részletezett perifériás T-sejtes limfóma; anaplasztikus nagysejtes limfóma; akut mieloid leukémia, beleértve az akut promielotikus leukémiát is és krónikus mieloproliferatív betegségek, beleértve a krónikus mieloid leukémiát is.
23. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 22. igénypont szerinti felhasználásra, ahol a betegség a B-sejtes non-Hodgkin limfóma.
24. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 22. igénypont szerinti felhasználásra, ahol a betegség a mieloma multiplex.
25. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 22. igénypont szerinti felhasználásra, ahol a betegség az akut limfoblasztos leukémia.
26. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 22. igénypont szerinti felhasználásra, ahol a betegség a B-sejtes krónikus limfocitás leukémia.
27. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 21-26. igénypontok bármelyike szerinti felhasználásra, ahol a módszer magában foglalja egy vagy több további terápiás szer adagolását is az alany számára.
28. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 27. igénypont szerinti felhasználásra, ahol az egy vagy több további terápiás szer az alábbiak közül választott: egy kemoterápiás szer, egy gyulladásgátló szer vagy egy immunszuppresszív és/vagy immunmoduláló szer.
29. Az antitest, az immunkonjugátum, a gyógyászati készítmény, a bispecifikus vagy multispecifikus molekula vagy az expressziós vektor a 27. igénypont szerinti felhasználásra, ahol az egy vagy több további terápiás szer az alábbi csoportból választott: ciszplatín, gefitinib, celastrol, rituximab, bevacizumab, erlotinib, bortezomib, thalidomid, pamidronát,

zoledronsav, clodronát, risendronát, ibandronát, etidronát, alendronát, tiludronát, arzén-trioxid, lenalidomid, filgrastim, pegfilgrastim, sargramostim, suberoyl anilid hidroxámsav és SCIO-469.

30. *In vitro* módszer a CD38 antigén jelenlétének kimutatására vagy egy CD38-af expresszáló sejt kimutatására mintában, ahol a módszer a következőkből áll
 - a) a mintát érintkezésbe hozzuk az 1-11. igénypontok bármelyike szerinti antitesttel olyan körülmények között, amelyek lehetővé teszik, hogy komplex képződjön a peptid és a CD38 között; és
 - b) detektáljuk a komplex képződését.
31. Anti-idiotipusú antitest, amely az 1-12. igénypontok bármelyike szerinti antitesthez kötődik.