(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
12 May 2005 (12.05.2005)

PCT

(10) International Publication Number
WO 2005/041669 A1

(51) International Patent Classification[7]: A01N 57/20

(21) International Application Number:
PCT/US2004/035807

(22) International Filing Date: 27 October 2004 (27.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/515,339 28 October 2003 (28.10.2003) US
60/532,758 24 December 2003 (24.12.2003) US

(71) Applicants *(for all designated States except US)*: **WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION** [US/US]; 1610 N.E. Eastgate Boulevard, Pullman, WA 99163 (US). **THE UNITED STATES OF AMERICA, as represented by THE SECRETARY OF AGRICULTURE** [US/US]; 1400 Independence Avenue SW, Washington, DC 20250 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **BALEY, George, J.** [US/US]; 430 West Main, Pullman, WA 99163 (US). **KIDWELL, Kimberlee, K.** [US/US]; 455 N.W. Robert Street, Pullman, WA 99163 (US). **PAULITZ, Timothy, C.** [US/US]; c/o USDA-ARS, Root Disease and Biological Control Lab, Rm. 363 Johnson Hall, Washington State University, Pullman, WA 99164-6430 (US).

(74) Agent: **YOUNG, Travis**; Klarquist Sparkman, LLP, One World Trade Center, Suite 1600, 121 SW Salmon Street, Portland, OR 97204 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:
— *of inventorship (Rule 4.17(iv)) for US only*

Published:
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUPPRESSION OF FOLIAR AND SOILBORNE PATHOGENS

(57) Abstract: Disclosed herein is a method for increasing the production of crops, particularly wheat and soybean, using herbicide resistant cultivars. In one aspect of this method, the method increases crop yield by diminishing the impact of the root diseases caused by Gaeumannomyces and Rhizoctonia species by treating the crop with an herbicide, in particular glyphosate. In another aspect the method for treating crops reduces the effects foliar pathogens and diseases, particularly fungal pathogens, such as rusts, including soybean rust, stem rust, stripe rust and leaf rust.

WO 2005/041669 A1

SUPPRESSION OF FOLIAR AND SOILBORNE PATHOGENS

CROSS REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Patent Application No. 60/515,339 filed October 28, 2003, and U.S. Provisional Patent Application No. 60/532,758 filed December 24, 2003, both of which are incorporated herein by reference.

ACKNOWLEDGEMENT OF GOVERNMENT SUPPORT

The Federal government may have certain rights in this technology pursuant to USDA Biotechnology Risk Assessment Research Grant number 2001-03734.

FIELD

This disclosure concerns methods for increasing the yield of crops that subject to disease pressure, particularly from seed borne, soil borne or foliar pathogens, such as fungi.

BACKGROUND

Commercial farming is a critical part of the economy. However crops are subject to near constant attack by insects, fungi, bacteria and other pathogens. When such pathogens encounter susceptible crops, such attacks can result in lower yield quality and can even destroy entire crops. Thus, pathogens cause substantial economic harm to growers and in some areas of the world contribute to famine.

Traditionally, farmers have relied upon conventional tillage methods to disrupt the soil and thereby control weeds, pathogens and volunteer crops. However, the current trend, particularly in the Pacific Northwest region, is to use no-till or direct seed crop production methods to reduce soil erosion and the accompanying environmental degradation associated with conventionally tilled fields. No-till and direct seeding methods aim to reduce environmental degradation but generally require the use of herbicides to control weeds and volunteer crops.

Typically growers apply herbicides prior to planting to control weeds since the crop itself may be susceptible to the herbicide. However the development of

herbicide resistant wheat varieties raises the possibility of increasing wheat yields by applying the herbicide to the standing crop. Unfortunately, weeds dying in the standing crop have been demonstrated to result in carryover of fungal pathogens, which typically are unaffected by herbicides, from the dying weeds to the standing wheat crop. Indeed, U.S. Patent No. 5,972,689 to Cook *et al.* (Cook) discloses that spraying with an herbicide such as glyphosate controls weeds, but favors the development of Rhizoctonia root rot in wheat. Because wheat is particularly susceptible to fungal pathogens, this carryover or "green-bridge" is a serious problem.

This green-bridge effect often leads to yield reductions associated with increased disease pressure which are the result of increased soilborne pathogens present on dying herbicide-sensitive volunteer and/or weeds. For example, *R. solani* significantly reduced grain yields of glyphosate-sensitive barley when glyphosate was applied three days before planting. However, no significant yield depression was detected when glyphosate was applied in the fall or three weeks prior to planting (Smiley *et al. Plant Dis.* **1992,** *76,* 937–942). These results demonstrate that a fresh source of *R. solani* inoculum, from the dying volunteer and weeds treated with glyphosate three days prior to planting, acted as a green-bridge for the fungus to infect barley planted shortly after herbicide application.

Another fungal disease of concern, take-all, is caused by *Gaeumannomyces graminis* var. *tritici* (*Ggt*), which has been a persistent pathogen plaguing wheat around the world for over a hundred years. Take-all (*Ggt*) disease is present in the roots, crown, and basal stem of infected wheat plants. Severe *Ggt* infection can decrease grain yields by as much as 50%. Symptoms of infection include stunting, blackened lower stems, and white heads. Take-all will commonly put out "runner-hyphae" to neighboring plants, so that a single site of infection is sufficient to cause multiple infections. Persistence through a green-bridge effect can occur similar to that reported for *R. solani*. Studies conducted in New Zealand have shown that treatment of cereals with glyphosate increases the levels of infection with *Ggt* due to the green-bridge effect of the herbicide on couch grass (Harvey *et al. Aglink 1/3000/3/82,* **1982,** Ministries of Agriculture and Fisheries, Wellington, New

Zealand). The most successful current form of *Ggt* control is through crop rotation, which is not always satisfactory for wheat production.

Nineteen *Pythium* species have been reported to be pathogenic to wheat roots. *Pythium* inoculum will remain active within the upper soil layer for years, utilizing residues as a source of nutrients. *Pythium* is considered to be a primary colonizer and infection levels can be reduced by removing straw and debris from the field. Unfortunately, this is not an option in a no-till system.

The interaction between glyphosate treated plants and infection by *Pythium* spp. has been investigated with numerous crop species. Soilborne *Pythium* spp. were found to be the first and predominant root colonizers of glyphosate treated plants (Levesque *et al. Mycological Research* **1993,** *20,* 307–312). This is an important observation since *Pythium* damage is often overlooked by growers, even though significant yield loss resulting from *Pythium* infection can occur.

Another group of important pathogens that affect wheat include foliar fungal pathogens, such rusts. Rust pathogens are parasitic fungi that infect wheat, barley, oats, beans, corn, sorghum, and other plants. Each rust pathogen is generally specific to its host and the location on the plant where infection occurs. The stem rust pathogen (*Puccinia graminis f.* sp. *tritici*) is a fungus that principally infects the leaf sheath of wheat plants. The leaf rust pathogen (*Puccinia recondita f.* sp. *tritici*) infects wheat plants through the stomates. The stripe rust pathogen (*Puccinia striiformis*) is similar to leaf rust, but differs in that infections appear systemic due to colonization patterns on wheat leaves.

Soybean rust is another serious rust pathogen that causes crop losses. It has not yet been detected in the continental United States, but the fact that it is principally spread by wind-borne spores indicates it may eventually reach major soybean growing areas in the United States. Soybean rust is caused by two fungal species, *Phakopsora pachyrhizi* and *Phakopsora meibomiae*. It has been reported in various countries including Australia, China, Korea, India, Japan, Nepal, Taiwan, Thailand, the Philippines, Mozambique, Nigeria, Rwanda, Uganda, Zimbabwe, South Africa, Brazil, Argentina, and Paraguay. *P. meibomiae* has been reported to be a weak pathogen. However, *Phakopsora pachyrhizi* is much more aggressive and recent introductions of *P. pachyrhizi* have rapidly spread causing severe damage in

Zimbabwe, South Africa, Paraguay, and Brazil. Yield losses have been reported from 10–80%. Other important fungal pathogens that affect soybeans include root rot caused by various species of *Phytophthora*.

There are few methods for controlling fungal diseases in wheat and soybeans, and none of these methods are widely accepted as being commercially viable. For example, some root rot diseases can be controlled through of crop rotation, that is, by not growing wheat in the same field more than every third or fourth year. However, like most other enterprises, agriculture has forced farmers to specialize in order to compete. The United States grows some 150 different crops, but fewer than 15 of these crops (including wheat and barley) occupy more than 90% of U.S. cropland, with the vast majority of farms specialized in the production of a single crop year after year on the same land, or two or at most three crops grown in a rotation on any given farm. Many wheat farms in areas well-suited to cereals tend to grow wheat every year or at least every other year in the same fields. Moreover, in certain regions, such as in the Pacific Northwest, leguminous crops commonly used in rotations do not bring the same levels of financial returns as do continuous wheat cropping systems (see, Cook and Veseth *Wheat Health Management*; American Phytopathological Society: St. Paul, MN, 1991). Therefore, crop rotation is not a feasible economic solution to reduce disease pressure in a continuously cropped no-till production system.

Many diseases of wheat, barley, and other crops are controlled by breeding varieties of the crops with resistance to the pathogens. However, this approach has not worked for certain fungal diseases of wheat, particularly root diseases. No commercial wheat is available that has resistance to take-all, Rhizoctonia root rot, or Pythium root rot. Moreover, rust pathogens mutate at a relatively high rate, and therefore new rust-resistant cultivars of wheat are needed approximately every seven years.

Another method that is currently used to combat fungal infections is the topical application of fungicides. Fungicides, although effective, are prohibitively expensive for growers and typically must be applied as a preventative, even if it is not certain that the plants will be infected. Moreover many of the fungicides previously used have been withdrawn by the EPA. Compounds that are currently

used are more easily degraded and therefore are less harmful to the environment, but fungi can quickly develop resistance to these compounds. Thus, fungicide applications are even less desirable than before.

SUMMARY OF THE DISCLOSURE

Disclosed herein is a method for conferring resistance to pathogens on crops, including wheat and soybeans. In one embodiment, the method includes treating a standing crop with an herbicide. In one aspect, the method includes treating a crop with an herbicide, which increases the crop's resistance to fungal pathogens. In one aspect of this method, the method increases crop yield by diminishing the impact of the root diseases caused by *Gaeumannomyces* and *Rhizoctonia* species. In another aspect, the method increases the crop's resistance to foliar pathogens, particularly fungal pathogens such as rusts caused by various species, including stem rust, stripe rust, leaf rust and soybean rust.

Typically, the disclosed methods result in increased crop yields. Any increase in crop yield is desirable, however typically the method produces from about 5% to about 25% greater yield, and more typically from about 5% to about 15% greater yield relative to an untreated crop.

The herbicide used in embodiments of the disclosed methods typically includes at least one of a sulfamoylurea, sulfonylcarboxamide, sulfonamide, sulfonylurea or glyphosate. In one aspect a combination of herbicides, such as two or more of the herbicides listed above or one of the listed herbicides and one or more herbicide not listed, can be applied to the crop. Typically the herbicide or herbicides are selected with reference to the particular wheat cultivar to be treated, for example, an herbicide to which the particular cultivar is resistant can be selected.

In one embodiment, the resistance of crops to various pathogens is induced by applying an effective amount of an agricultural grade formulation of a glyphosate-based herbicide (such as ROUNDUP® brand herbicide produced by Monsanto, St. Louis, MO) to a glyphosate resistant crop after emergence. Typically, the effective amount of active ingredient applied is from about 0.5 kg/ha to about 3.0 kg/ha, and more typically from about 1.0 kg/ha to about 1.5 kg/ha. In one embodiment wheat is treated with formulations containing glyphosate at a density

from about 0.5 kg/ha to about 1.5 kg/ha of glyphosate. In another embodiment a soybean crop is treated with from about greater than about 1.0 kg/ha, such as from about 1.5 to about 3.0 kg/ha of glyphosate. In one aspect of the method, an amount of glyphosate less than that typically recommended for weed control is effective to confer or induce pathogen resistance to a crop.

In one embodiment a crop is treated with glyphosate at a growth stage not currently recommended for glyphosate treatment. For example, in one embodiment glyphosate is applied to wheat at or after the 3 leaf stage, such as between the 3 leaf stage and the flowering stage. In another embodiment glyphosate is applied to soybean at growth stage after emergence and before the flowering stage. In one aspect, glyphosate is applied after symptoms of pathogen infection appear.

In embodiments of the method, treatment of herbicide resistant crop with an herbicide reduces disease on the crop for an extended period of time, and particularly for longer than the herbicide is active as an herbicide after the treatment. Typically, the disease reduction extends for at least about seven days after the treatment; more typically extends for at least about 21 days, and in certain examples extends for the life of the crop.

In another embodiment of the method, a crop is treated with an herbicide prior to emergence, and treated with the same or a different herbicide after emergence. In yet another embodiment of the method a crop is treated with two or more separate applications of one or more herbicide.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a bar graph recording the number of sporulating lesions present 14 days following inoculation of two different soybean cultivars with soybean rust spores and subject to five different conditions, demonstrating the suppression of soybean rust accompanying glyphosate treatment of soybeans before and after inoculation with rust (treatments followed by the same letters are not significantly different at P = 0.05, LSD test).

Figure 2 is a bar graph recording the number of sporulating lesions present 21 days following inoculation of two different soybean cultivars with soybean rust spores and subject to five different conditions, demonstrating the suppression of

soybean rust accompanying glyphosate treatment of soybeans before and after inoculation with rust (treatments followed by the same letters are not significantly different at P = 0.05, LSD test).

Figure 3 is a bar graph recording observed rust rating following inoculation of the soybeans with rust and subject to five different treatment conditions (treatments followed by the same letter are not significantly different at P = 0.05, Kruskal-Wallis One-Way Nonparametric test).

DETAILED DESCRIPTION

I. Introduction

The present disclosure illustrates the surprising result that the treatment of herbicide resistant wheat, particularly glyphosate resistant wheat, with an herbicide, such as glyphosate, reduces disease caused by pathogens in the wheat. Moreover, the wheat exhibits persistently superior pathogen resistance over an extended period of time, after the herbicide is no longer effective as an herbicide. In one embodiment, the deterrence of the pathogen is due to the accumulation of glyphosate in the tissues the plant. In one embodiment the herbicide acts indirectly by inducing systemic disease resistance.

II. Herbicides and Herbicide Resistant Crops

Disease reduction on herbicide resistant crops can be accomplished according to the methods disclosed herein. Suitable herbicides and herbicide resistant crops are known and will be readily apparent to those of ordinary skill in the art. Moreover, methods for producing transgenic herbicide resistant crops, including, without limitation, wheat and soybeans, are disclosed by U.S. Patent Nos. 6,635,806, 6,803,501, 6,750,383, which are incorporated herein by reference.

One herbicide for which resistant crops, including resistant wheat and soybean cultivars, have been developed is *N*-phosphonomethylglycine, commonly referred to as glyphosate. Glyphosate is the active ingredient in glyphosate herbicides, such as ROUNDUP® brand herbicide produced by Monsanto (St. Louis, MO). Typically, glyphosate is formulated as a water-soluble salt such as an ammonium, alkylamine, alkali metal or trimethylsulfonium salt. One of the most

common formulations is the isopropylamine salt of glyphosate, which is the form employed in ROUNDUP® brand herbicide.

Glyphosate is conventionally applied as an aqueous solution to the foliage of plants to be killed, where it is taken up into the leaves and transported throughout the plant. Commercial formulations of glyphosate may also include one or more surfactants to facilitate penetration of the active ingredient into the plant leaves, as well as compounds to enhance rainfastness. Numerous U.S. patents disclose various formulations of glyphosate and methods for their use, including U.S. Patent Nos. 4,405,531; 5,118,338; 5,196,044; 5,639,711; 5,652,197; 5,679,621; 5,750,468; 6,207,617; and 6,455,473. Each of these patents is incorporated herein by reference.

Glyphosate inhibits the shikimic acid pathway, which is responsible for the biosynthesis of aromatic compounds including amino acids, such as tryptophan, phenylalanine and tyrosine, as well as several secondary metabolites. Specifically, glyphosate inhibits the conversion of phosphoenolpyruvic acid (PEP) and 3-phosphoshikimic acid to 5-enolpyruvyl-3-phosphoshikimic acid by inhibiting the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (hereinafter referred to as EPSP synthase or EPSPS). For purposes of the present disclosure, the term "glyphosate" is intended to include any herbicidally effective form of *N*-phosphonomethylglycine (including any salt thereof) and other forms which result in the production of the glyphosate anion in plants.

Glyphosate resistant crops, including glyphosate resistant wheat and soybeans, are well known to those of ordinary skill in the art. Examples of glyphosate resistant wheat are disclosed in U.S. Patent Nos. 5,463,175; 5,463,175; and 6,153,812. Production of lines of other plant species, including soybean, expressing a glyphosate-tolerance gene may be produced by techniques known in the art. *See*, e.g. U.S. Pat. Nos. 5,312,910; 5,310,667; 5,463,175. Each of these patents is incorporated herein by reference.

III. Methods and Examples

Because herbicide treatment, including glyphosate treatment, typically impairs certain plant defense mechanisms, dying herbicide sensitive plants can function as a green-bridge, transferring pathogens to a standing crop. For example,

glyphosate blocks the production of phenolic compounds, such as lignin precursors and phytoalexins that are components of the defense mechanisms and confer disease resistance. Because glyphosate causes the breakdown of plant defenses, glyphosate renders glyphosate sensitive plants susceptible to pathogen invasion. This supplemental effect of glyphosate treatment has been termed Glyphosate Synergistic Interaction (GSI). GSI also provides a source or inoculum of pathogens that can attack the glyphosate resistant crop, thereby worsening the green-bridge effect.

As demonstrated herein, it has been surprisingly found that herbicides can function indirectly to reduce pathogen damage to wheat. This result is particularly surprising because herbicides have previously been shown to exacerbate the problems caused by fungal pathogens. Moreover, the herbicides have little or no direct anti-fungal activity and indeed the pathogen resistance is shown to be effective well after the herbicide itself dissipates from the treatment site. Accordingly, disclosed herein are methods for increasing disease resistance in wheat, examples of which include providing an herbicide resistant wheat crop and treating the crop with an herbicide, thereby conferring disease resistance on the crop for an extended period of time.

One of the herbicides effective in the present methods, glyphosate, is currently applied to glyphosate sensitive wheat fields (prior to emergence of the crop) by producers in the fall or spring as an inexpensive yet effective way of controlling weed competition. Glyphosate has a wide spectrum of activity and controls 97% of the world's worst weed problems. However, glyphosate has not been shown to be effective against fungal pathogens, (see, Franz, J.E. Discovery Development and Chemistry of Glyphosate. In, *The Herbicide Glyphosate,* Butterworths, London, 1985, pp 3–17). Moreover, some previous studies have suggested that glyphosate encourages the growth of toxic fungi.

Disclosed embodiments of the present methods using glyphosate resistant wheat and soybean crops provide this same level of weed control during the early crop growth stages, and moreover reduce the effect of pathogenic organisms and particularly fungal pathogens on wheat and soybean crops. Moreover, embodiments of the present methods are effective for treating crops that are already infected with a pathogen, having a curative effect on such crops. Therefore, in addition to

eliminating weed competition, the present methods reduce losses to plant diseases, thereby decreasing herbicide costs, increasing grain yields, and enhancing profitability.

For example, under severe stripe rust conditions, two Bobwhite cultivars that had not been treated with RoundUp were heavily infested with stripe rust, whereas glyphosate treated glyphosate resistant Bobwhite exhibited only moderate stripe rust symptoms. Bobwhite cultivars sprayed with Buctril/Harmony Extra (Buctril is commercially available from Bayer CropScience, and Harmony Extra is commercially available from DuPont) or the untreated control displayed severe stripe rust susceptibility symptoms and matured 2–3 weeks earlier than NILs treated with RoundUp. Buctril/Harmony Extra treated glyphosate resistant Bobwhite produced significantly ($P = 0.01$) less grain than the glyphosate resistant Bobwhite treated with glyphosate, regardless of root disease treatment. Visual differences in stripe rust severity typically were not apparent until 21 days after herbicide application, which notably is well beyond the time after application that the herbicide exerts its direct herbicidal effects.

The present results in both wheat and soybean are surprising for several reasons, including that glyphosate was not previously found to be effective for suppressing disease in glyphosate resistant crops. The results presented herein also are surprising because glyphosate is directly effective as an herbicide for only a short time and is not persistent in herbicidally effective amounts. These results demonstrate that glyphosate application suppresses pathogen growth for an extended period of time. The method is particularly effective in hindering the colonization of leaf tissue by foliar pathogens. Moreover, the pathogen suppression extends to root pathogens, such as *Rhizoctonia* and *Ggt*.

In one embodiment, the methods disclosed herein can be used to treat a crop, such as wheat or soy crops, that are infected with a pathogen, such as a fungal pathogen, including seed borne, soil borne or foliar fungal diseases. For example, the treatment of infected crops with glyphosate is demonstrated herein to decrease fungal colonization of crops and to increase crop yield. This result is surprising because the fungal pathogens, such as stripe rust, soybean rust and the like, previously have not been thought to be susceptible to glyphosate. In one

embodiment the treatment of the crops includes two or more separate applications of glyphosate. Thus, according to the embodiments disclosed herein crop yields can be increased by treating a crop before symptoms of infection are evident, following the appearance of infection, or both. In exemplary embodiments, treatment of wheat and soybean both before and after the emergence of foliar fungal infection resulted in increased crop yield. In one embodiment, the crops are treated with from about 0.5 kg/ha to about 3.0 kg/ha of glyphosate in one or more applications. In one embodiment, from about 0.5 kg/ha to about 1.5 kg/ha, such as from about 1.0 kg/ha to about 1.5 kg/ha of glyphosate is applied. In one embodiment a first application of glyphosate of from about 0.5 kg/ha to about 2.0 kg/ha is made, followed by at least a second application of from about 0.5 kg/ha to about 2.0 kg/ha. In one aspect of this embodiment the crop is wheat and at least one application is made between the 3 leaf stage and the flowering stage. In another aspect of the method the crop is soybean and at least one application is made between emergence and the flowering stage.

As demonstrated in the examples below, the reduced disease susceptibility of wheat treated according to the method also was responsible for increased grain yields observed for glyphosate treated glyphosate resistant cultivars across trial growing locations, regardless of disease treatment.

The foregoing disclosure is further explained by the following non-limiting examples.

Example 1

This example describes the response of glyphosate resistant and glyphosate sensitive spring wheat to inoculation with soil pathogens as judged by wheat yield. Inoculation simulates the green-bridge effect of dying weeds and volunteer in a standing wheat crop. The soil pathogens assessed were: *Rhizoctonia solani* isolate AG-8 and *Gaeumannomyces graminis* var. *tritici*. These pathogenic strains are prevalent in the Pacific Northwest and were isolated from the region. They have been maintained by the Department of Plant Pathology and USDA researchers at Washington State University.

Near isogenic lines (glyphosate resistant and glyphosate sensitive) of Westbred 926 and Bobwhite were seeded using a no-till drill at three different sites (Lind, Davenport and Pullman, WA).

To introduce the root pathogens into the wheat seed for distribution into the field, oat seeds, autoclaved and sterilized, were inoculated for each soilborne pathogen using isolated pathogen cultures. Oat seeds that were non-inoculated, but sterilized and autoclaved, served as a control. Following two weeks of pathogen growth, the inoculated oat seeds were dried down for 2 days and screened for potential contamination. To establish a green-bridge, a mixture of 15% barley and 20% oat inoculum seeds (by weight) were planted using a small plot no-till drill. One week after planting the introduced green-bridge, the wheat isolines were planted in rows adjacent to the barley green-bridge rows in a randomized, split-plot design, replicated in five randomized blocks. Inoculum treatment served as the main plot, and response to herbicides as the subplot factor.

The results for *Ggt* and *Rhizoctonia* inoculation compared to non-inoculated controls are illustrated in Table 1. In summary, with reference to Table 1, inoculation of wheat as described above can result in depressed yields, as demonstrated in Lind in 2002. Yield losses are often more severe in areas of lower rainfall such as Lind.

Table 1

Wheat yield depression due to fungal inoculation, without herbicide treatments, 2002.

Inoculation	**Herbicide**	**Location**	**Yield (bu/acre)**
None	None	Lind	13.5
Ggt	None	Lind	12.0
Rhizoctonia	None	Lind	12.9
None	None	Davenport	21.5
Ggt	None	Davenport	24.6
Rhizoctonia	None	Davenport	23.3
None	None	Pullman	46.5
Ggt	None	Pullman	47.0
Rhizoctonia	None	Pullman	47.5

Example 2

This example describes the effect of herbicide treatment on spring wheat, averaged over all disease treatments, planted as described in Example 1. The appropriate herbicide was applied to each trial once the plants reached the 4-5 leaf stage with a custom built hand-pulled sprayer outfitted with a boom with a length of 16 feet with spray shields on the side to minimize drift. In this manner, an entire plot was treated at once with the appropriate herbicide rate. Entries that were resistant to glyphosate were treated with ROUNDUP® brand herbicide (0.63 kg/ha), available from Monsanto, St. Louis, MO. Herbicide sensitive and resistant entries were treated with a common broadleaf herbicide treatment (Buctril/Harmony) or a no-spray control. Introduced green-bridge barley and any other weeds were manually removed 7–10 days after spray treatment to simulate the time required for ROUNDUP to take effect and maintained weed-free conditions to provide an accurate comparison to commercial weed management strategies needed to assess potential herbicide/disease interactions.

The results of the herbicide treatment are recorded in Table 2. With reference to Table 2, treatment of glyphosate resistant wheat with glyphosate resulted in higher yields in each trial.

Table 2

Effect of herbicide treatment on glyphosate resistant wheat averaged over all fungal inoculation treatments, demonstrating increased yield associated with glyphosate treatment.

Herbicide Resistance	Herbicide	Location	Yield (bu/acre)
None	None	Lind	12.8
None	Buctril/Harmony	Lind	12.7
glyphosate res.	Glyphosate	Lind	14.2
None	None	Davenport	23.2
None	Buctril/Harmony	Davenport	24.2
glyphosate res.	Glyphosate	Davenport	28.9
None	None	Pullman	47.0
None	Buctril/Harmony	Pullman	45.6
glyphosate res.	Glyphosate	Pullman	56.5

Example 3

This example describes the assessment of the disease response of non-inoculated glyphosate resistant spring wheat as described in Example 1 and treated with glyphosate in comparison to inoculated near isogenic lines of glyphosate sensitive spring wheat. The results, recorded in Table 3, demonstrate increased yield accompanying glyphosate treatment of non-inoculated wheat.

Table 3

Yields of non-inoculated wheat demonstrating increased yield of glyphosate resistant wheat due to glyphosate treatment, 2002.

Inoculation	**Herbicide Resistance**	**Herbicide**	**Location**	**Yield (bu/acre)**
none	None	None	Lind	13.5
none	None	Buctril/Harmony	Lind	12.8
none	glyphosate res.	glyphosate	Lind	14.2
none	None	None	Davenport	24.6
none	None	Buctril/Harmony	Davenport	23.6
none	glyphosate res.	glyphosate	Davenport	29.1
none	None	None	Pullman	47.0
none	None	Buctril/Harmony	Pullman	45.0
none	glyphosate res.	glyphosate	Pullman	55.6

Thus treatment of glyphosate-resistant wheat lines with glyphosate increases yield, even in the absence of additional pathogen inoculum.

Example 4

This example describes the assessment of the disease response of glyphosate resistant spring wheat inoculated with *Ggt* as described in Example 1 and treated with glyphosate in comparison to inoculated near isogenic lines of glyphosate sensitive spring wheat.

The effect of herbicide treatment on the yield of glyphosate resistant wheat inoculated with *Ggt* is illustrated in Table 4. The yield for glyphosate treated, glyphosate resistant, *Ggt* inoculated wheat was significantly superior ($P \leq 0.05$) as compared to both untreated and Buctril/Harmony treated glyphosate sensitive wheat at each location.

Table 4

Yields of *Ggt* inoculated wheat demonstrating increased yield of glyphosate resistant wheat due to glyphosate treatment.

Inoculation	Herbicide Resistance	Herbicide	Location	Yield (bu/acre)
Ggt	None	None	Lind	12.0
Ggt	None	Buctril/Harmony	Lind	13.6
Ggt	glyphosate res.	glyphosate	Lind	14.4
Ggt	None	None	Davenport	21.5
Ggt	None	Buctril/Harmony	Davenport	24.2
Ggt	glyphosate res.	glyphosate	Davenport	28.7
Ggt	None	None	Pullman	46.5
Ggt	None	Buctril/Harmony	Pullman	41.0
Ggt	glyphosate res.	glyphosate	Pullman	57.0

Example 5

This example describes the assessment of the disease response of glyphosate resistant spring wheat inoculated with *Rhizoctonia solani* as described in Example 1 and treated with glyphosate in comparison to inoculated near isogenic lines of glyphosate sensitive spring wheat.

With reference to Table 5, glyphosate treatment of glyphosate resistant wheat resulted in enhanced yield at each location relative to untreated and conventionally (Buctril/Harmony) treated glyphosate sensitive wheat.

Table 5

Yields of *Rhizoctonia* inoculated wheat demonstrating increased yield of glyphosate resistant wheat due to glyphosate treatment.

Inoculation	Herbicide Resistance	Herbicide	Location	Yield (bu/acre)
Rhizoctonia	None	None	Lind	12.9
Rhizoctonia	None	Buctril/Harmony	Lind	11.7
Rhizoctonia	glyphosate res.	glyphosate	Lind	14.0
Rhizoctonia	None	None	Davenport	23.3
Rhizoctonia	None	Buctril/Harmony	Davenport	24.7
Rhizoctonia	glyphosate res.	glyphosate	Davenport	29.0
Rhizoctonia	None	None	Pullman	47.5
Rhizoctonia	None	Buctril/Harmony	Pullman	50.6
Rhizoctonia	glyphosate res.	glyphosate	Pullman	53.9

Rhizoctonia and *Ggt* naturally prevail in areas receiving low and high levels of precipitation, respectively. In trials planted in the low and high rainfall zones,

grain yields of NILs treated with Buctril/Harmony Extra were significantly (P = 0.05) lower than NILs treated with RoundUp or the no-spray control.

Regardless of disease treatment or location, glyphosate treated RoundUp Ready® (glyphosate resistant) spring wheat, produced significantly (P = 0.001) more grain than glyphosate sensitive NILs treated with Buctril/Harmony Extra or the no spray control.

This demonstrates green-bridge transmission of fungal pathogens, such as *Rhizoctonia* and *Ggt*, due to glyphosate application does not suppress yields of glyphosate resistant varieties; rather, glyphosate application increases yields demonstrating that the glyphosate treated plants not only resist the inoculated pathogens, but also other endemic pathogens, such as *Puccinia* spp.(rusts).

Example 6

This example describes the effect of herbicides on natural stripe rust infection of NILs of Bobwhite with and without glyphosate tolerance. The wheat was seeded and inoculated with *Rhizoctonia* and *Ggt* as described in Example 1. The yield results from the Pullman site are recorded in Table 6.

With reference to Table 6, the glyphosate treated glyphosate resistant wheat Bobwhite NILs produced superior yields regardless of what pathogen had been introduced. These results highlight the advantages of the disclosed method for conferring disease resistance on herbicide resistant wheat.

Table 6

Glyphosate induced yield enhancement of Bobwhite NILs in the presence of stripe rust in Pullman in 2002.

Inoculation	Cultivar	Herbicide	Yield (bu/acre)
None	Bobwhite	None	37.2
None	Bobwhite (res.)	None	36.8
Rhizoctonia	Bobwhite	None	38.7
Rhizoctonia	Bobwhite (res.)	None	33.9
Ggt	Bobwhite	None	37.7
Ggt	Bobwhite (res.)	None	32.8
None	Bobwhite	Buctril/Harmony	33.0
None	Bobwhite (res.)	Buctril/Harmony	31.8
Rhizoctonia	Bobwhite	Buctril/Harmony	39.4
Rhizoctonia	Bobwhite (res.)	Buctril/Harmony	36.3
Ggt	Bobwhite	Buctril/Harmony	37.7
Ggt	Bobwhite (res.)	Buctril/Harmony	26.4
None	Bobwhite (res.)	Glyphosate	44.9
Rhizoctonia	Bobwhite (res.)	Glyphosate	40.0
GGT	Bobwhite (res.)	Glyphosate	49.0

Example 7

This example describes the effect of glyphosate treatment of glyphosate resistant wheat in varying amounts and at different times. Eight different treatment regimens were applied to the hard white spring wheat variety Macon, which is susceptible to stripe rust. Macon plots were planted at the Palouse Conservation Farm near Pullman, Washington and subjected to one of the following eight treatment regimens:

Treatment 1: no herbicide treatment and manual weed control;

Treatment 2: a 16 oz./A (0.50 kg/ha of active ingredient) application of glyphosate, one-half the recommended rate of 32 oz./A (1.0 kg/ha of active

ingredient), was made 'on-label' to glyphosate resistant wheat at the 4–5 leaf stage.

Treatment 3: a first application of glyphosate at one-half (0.50 kg/ha of active ingredient) the recommended density of 1.0 kg/ha was made to glyphosate resistant wheat at the 4-5 leaf stage according to the manufacturer's instructions. A second application of 16 oz./A (0.50 kg/ha of active ingredient) glyphosate was made once moderate stripe rust symptoms appeared (42 days following the first glyphosate application).

Treatment 4: a 16 oz./A (0.50 kg/ha of active ingredient) application of glyphosate, one-half the recommended rate, was made to glyphosate resistant wheat once moderate stripe rust symptoms appeared.

Treatment 5: a 32 oz./A (1.0 kg/ha of active ingredient) application of glyphosate, the recommended full rate, was made 'on-label' to glyphosate resistant wheat at the 4-5 leaf stage.

Treatment 6: a first application of 32 oz./A. (1.0 kg/ha of active ingredient) glyphosate, the recommended full rate, was made 'on-label' to glyphosate resistant wheat at the 4–5 leaf stage and a second application of 32 oz./A. (1.0 kg/ha) glyphosate was made once moderate stripe rust symptoms appeared.

Treatment 7: a 32 oz./A. (1.0 kg/ha of active ingredient) application of glyphosate was made to glyphosate resistant wheat once moderate stripe rust symptoms appeared. **Treatment 8:** a 48 oz./A. (1.0 kg/ha of active ingredient) application of glyphosate, 1.5 times the recommended rate, was made to glyphosate resistant plants at the 4–5 leaf stage.

The results of the treatments listed above are recorded in Table 7.

Table 7

Yield of glyphosate-resistant wheat with variable timing of administration and amount of glyphosate.

Treatment	Stripe Rust (96 days after planting)	Stripe Rust (102 days after planting)	Stripe Rust (106 days after planting)	Yield (bu/A)
No Spray	9, 40%	7-8, 70%	8, 50%	55.4
No Spray	9, 30%	9, 50%	8, 50%	59.6
16oz (52 days)	7-8, 10%	7-8, 30%	8, 30%	60.2
16oz (52 days) and 16oz (94 days)	4-5, 10%	0-1, 10%	2, 10%	68.7
16oz (94 days)	7-8, 20%	2-3, 20%	2-3, 15%	52.7
32oz (52 days)	7-8, 20%	7-8, 40%	8, 60%	80.3
32oz (52 days) and 32oz (94 days)	9, 5%	1-2, 5%	3, 10%	74.5
32oz (94 days)	7-8, 25%	0-1, 20%	2, 5%	64.5
48oz (52 days)	7-8, 2%	4-5, 10%	8, 20%	67.8

With reference to Table 7, stripe rust severity is characterized with two numerical indications. The first is a number characterizing the stripe rust severity on a scale of 0–9, with 0 corresponding to no infection and 9 being a high level of spore activity and infection. The second number is a percentage of the plants in the plot infected with the rating described by the first number.

With continued reference to Table 7, glyphosate resistant Macon experienced high levels of stripe rust infection when left untreated by glyphosate (rows 1 and 2). When glyphosate was applied to glyphosate resistant wheat at the 4–5 leaf stage, the 48 oz./A rate of glyphosate did the best job of controlling stripe rust compared to the 16 oz./A and 32 oz./A application. The higher rate of glyphosate clearly provided improved control of stripe rust infection.

When glyphosate was applied to glyphosate resistant Macon once plants had become moderately infected with stripe rust a curative effect was observed (rows 5 and 8). These "off-label" applications made 94 days after planting resulted in the formation of necrotic stripes on the leaves of the plants on the second day following glyphosate application. These necrotic stripes had been actively sporulating stripe rust at the time of glyphosate application. The amount of time needed to control stripe rust infection was found to be similar to when fungicides are applied to wheat plants infected with stripe rust.

The best control of stripe rust was seen for treatments 3 and treatments 6 (rows 4 and 7), which were treated with "split-treatments" of glyphosate. The stripe rust activity for both of these treatments 94 days after planting was similar to treatment 2 and treatment 4, before the second application of glyphosate was made. Once the second glyphosate application was made, a curative effect against active stripe rust was observed. These treatments had the lowest levels of stripe rust infection in the trial due to the initial fungicidal activity of glyphosate from the first (4–5 leaf stage; 52 days after planting) applications and the curative effects of glyphosate from the second applications. A curative effect was also noted when glyphosate was applied only once for treatments 4 and 7. These results demonstrate that the methods disclosed herein for treating glyphosate resistant wheat provide control of glyphosate sensitive weeds and control against the negative effects associated with fungal pathogens.

Example 8

This example describes the suppression of soybean rust using glyphosate treatment of glyphosate resistant soybean varieties.

Two different Roundup Ready soybean lines were used: Line 1 (purple flowers/grey pubescence) and Line 2 (white flowers/grey pubescence). Both lines are glyphosate resistant and have been classified as maturity group 3, with white or purple flowers and grey pubescence. The following treatments were used:

Treatment 1: No glyphosate spray before or after inoculation

Treatment 2: One application of glyphosate (48 oz./A) (1.0 kg/ha of active ingredient) three days before inoculation

Treatment 3: Twice the amount of glyphosate as in Treatment 2 (96 oz./A) (2.0 kg/ha of active ingredient) applied three days before inoculation

Treatment 4: One application of glyphosate (48 oz./A) three days before inoculation, and one application (48 oz./A) (1.5 kg/ha of active ingredient) when symptoms of soybean rust first appeared.

Treatment 5: No glyphosate spray before inoculation, and one application (48 oz./A) (1.5 kg/ha of active ingredient) when symptoms of soybean rust first appeared.

The results of this example are recorded in Figures 1–3. With reference to Figures 1–3, treatments 2 and 3 demonstrate that glyphosate treatment exerts a dose-dependent effect on soy rust. Treatments 4 and 5 demonstrate that glyphosate treatment of soybean infected with soy rust has a curative effect.

It will be apparent to those skilled in the art that various modifications and variations can be made in the disclosed methods without departing from the scope or spirit of the disclosure. Other embodiments of the methods will be apparent to those skilled in the art from consideration of the specification and practice of the procedures disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

We claim:

1. A method for reducing disease on a crop infected with at least one pathogen, comprising:

providing an herbicide resistant crop, wherein the crop is selected from glyphosate resistant wheat and glyphosate resistant soybeans;

treating the crop with glyphosate at a density of greater than about 1.0 kg/ha of glyphosate, thereby reducing the effects of the pathogen on the crop.

2. The method according to claim 1, wherein treating the crop comprises at least two separate applications of glyphosate.

3. The method according to claim 2, wherein the more than two separate applications of glyphosate are applied at least about seven days apart.

4. The method according to claim 1, wherein treating the crop comprises treating the crop with from greater than about 1.0 kg/ha to about 3.0 kg/ha of glyphosate.

5. The method according to claim 1, wherein treating the crop comprises treating the crop with from greater than about 1.0 kg/ha to about 2.0 kg/ha of glyphosate.

6. The method of claim 1, wherein treating the crop comprises treating the crop with from about 1.5 kg/ha to about 2.0 kg/ha of glyphosate.

7. The method of claim 4, wherein treating the crop with glyphosate comprises at least two separate applications of glyphosate.

8. The method of claim 1, wherein the pathogen is a fungal pathogen.

9. The method of claim 1, wherein the pathogen is a foliar pathogen.

10. The method of claim 1, wherein the pathogen is a species of *Rhizoctonia, Gaeumannomyces, Phakopsora* or *Puccinia.*

11. The method of claim 1, wherein the pathogen is *Phakopsora pachyrhizi.*

12. The method of claim 11, wherein the crop is glyphosate resistant soybean.

13. The method of claim 1, wherein the crop is glyphosate resistant wheat.

14. The method of claim 1, wherein the yield is from about 5% to about 20% higher than a crop not treated with glyphosate.

15. The method of claim 1, wherein the crop is glyphosate resistant wheat and the crop is treated with glyphosate at a stage between the 3 leaf stage and the flowering stage.

16. The method of claim 1, wherein the crop is glyphosate resistant soybean and the soybeans and the crop is treated between emergence and the flowering stage.

17. The method of claim 1, wherein treating the crop with glyphosate comprises treating the crop with glyphosate prior to the display of a symptom of pathogen presence.

18. The method of claim 1, further comprising harvesting the crop thereby yielding a harvested crop.

19. A harvested crop produced by the method of claim 18.

20. A method for reducing disease on a wheat crop with at least one pathogen, comprising:

providing an herbicide resistant wheat crop; and

treating the wheat crop with an herbicide after emergence of the herbicide resistant wheat crop, thereby reducing the effects of the pathogen on the wheat crop.

21. The method according to claim 20, wherein the herbicide resistant wheat crop is glyphosate resistant.

22. The method according to claim 20, further comprising treating the wheat crop prior to emergence.

23. The method according to claim 20, wherein the herbicide is glyphosate.

24. The method according to claim 20, wherein the herbicide is a 5-enolpyruvylshikimate-3-phosphate synthase inhibitor.

25. The method according to claim 20, wherein the pathogen is a soilborne pathogen.

26. The method according to claim 20, wherein the pathogen is a fungal pathogen.

27. The method according to claim 20, wherein the pathogen is a species of *Rhizoctonia*, *Gaeumannomyces, Phakopsora* or *Puccinia*.

28. The method according to claim 27, wherein the pathogen is *Gaeumannomyces graminis var tritici.*

29. The method according to claim 20, wherein the pathogen is a foliar pathogen.

30. The method according to claim 20, wherein the pathogen causes stripe rust, stem rust or leaf rust.

31. The method according to claim 30, wherein the pathogen is *Puccinia striiformis*.

32. The method according to claim 20, wherein pathogen activity is decreased for at least 21 days after herbicide application.

33. The method of claim 21, wherein the glyphosate resistant wheat crop is treated with from about 0.5 kg/ha to about 2.0 kg/ha glyphosate, thereby increasing the yield of the wheat, wherein the yield is at least about 5% higher than a glyphosate sensitive wheat crop.

34. The method according to claim 20, wherein glyphosate is applied at a density of from about 0.5 kg/ha to about 1.5 kg/ha.

35. The method according to claim 20, wherein glyphosate is applied at a density of from about 0.5 kg/ha to about 1.0 kg/ha.

36. The method according to claim 33, wherein the yield is from about 5% to about 20% higher.



Roundup Ready Soybean - Day 14
Line 1
Line 2
Number of Sporulating Lesions
140.00
120.00
100.00
80.00
60.00
40.00
20.00
0.00
a
ab
a
ab
c
b
a
cd
a
abc
Treatment 1
Treatment 2
Treatment 3
Treatment 4
Treatment 5
Treatment

FIG. 1



Roundup Ready Soybean - Day 21
Line 1
Line 2
Number of Sporulating Lesions
180.00
160.00
140.00
120.00
100.00
80.00
60.00
40.00
20.00
0.00
Treatment 1
Treatment 2
Treatment 3
Treatment 4
Treatment 5
Treatment
ab
ab
a
a
b
b
ab
ab

FIG. 2



Roundup Ready Soybean - Rust Rating
Line 1
Line 2
A
A
B
AB
AB
5.00
4.00
3.00
2.00
1.00
0.00
Rust Rating (1-5)
1
2
3
4
5
Treatment

FIG. 3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/035807

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01N57/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	S. SANOGO, X.B. YANG & H. SCHERM: "Effects of herbcides on Fusarium solani (...) and development of SDS in glyphosate-tolerant soybean" PHYTOPATHOLGY, vol. 90, no. 1, 2000, pages 57-66, XP001182837 page 57, column 2, paragraph 2 page 60, column 1, paragraph 2 page 63, column 1 - column 2 page 64, column 2, paragraph 2 - page 65, column 1, paragraph 2 ----- -/--	1-12, 14-19

[X] Further documents are listed in the continuation of box C.

[X] Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
12 April 2005	21/04/2005

Name and mailing address of the ISA	Authorized officer
European Patent Office, P B 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx 31 651 epo nl, Fax. (+31-70) 340-3016	Klaver, J

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	C.A.BRADLEY ET AL.: "Interaction of Rhizoctonia disease of soybeans and four postemergence herbicides on four soybean cultivars in the greenhouse" PHYTOPATHOLOGY, vol. 87, no. 6(Suppl), 1997, page S11, XP001206019 the whole document	1-12, 14-18
X	R. HARIKRISHNAN & X.B.YANG: "Effects of herbicides on root rot and damping-off caused by Rhizoctonia solani in glyphosate-tolerant soybean." PLANT DISEASE, vol. 86, no. 12, 2002, pages 1369-1373, XP001206018 page 1370, column 1, paragraph 1 page 1370, column 3, paragraphs 1,2 page 1372, column 3, paragraph 2 - page 1373, column 1, paragraph 1; tables 4-7	1-12, 14-19
X	H. ZHOU ET AL.: "Field efficacy assessment of transgenic Roundup Ready wheat" CROP SCIENCE, vol. 43, May 2003 (2003-05), pages 1072-1075, XP002324017 page 1072, column 2, paragraph 4 page 1073, column 1, paragraph 3 - page 1074, column 1, paragraph 4 page 1075, column 1, paragraph 2	19
A	C.D.LEE, D. PENNER & R. HAMMERSCHMIDT: "Influence of formualted glyphosate and actoivator adjuvants ov Sclerotinia sclerotiorum in glyphosate-resistant and -susceptible Glycine max." WEED SCIENCE, vol. 48, 2000, pages 710-715, XP009045945 page 711, column 1, paragraph 3 - column 2, paragraph 2 page 713, column 1, paragraph 2 - page 714, column 2, paragraph 2; tables 1-3	1-9,12, 14-19

-/--

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	R.W.SMILEY, A.G.OGG & R.J.COOK: "Influence of glyphosate on Rhizoctonia root rot, growth and yield of barley." PLANT DISEASE, vol. 76, no. 9, 1992, pages 937-942, XP009045170 cited in the application page 937, column 2, paragraph 2 - column 3, paragraph 1 page 937, column 3, paragraph 5 - page 938, column 1, paragraph 2 page 940, column 3, paragraph 3 - page 941, column 1, paragraph 2; figure 4; table 3	1-36
A	WO 97/36488 A (MONSANTO EUROPE S.A; BRANTS, IVO; GRAHAM, WILLIAM) 9 October 1997 (1997-10-09) the whole document	1-36

INTERNATIONAL SEARCH REPORT

nformation on patent family members

Int[redacted]tional Application No
PCT/US2004/035807

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9736488 A	09-10-1997	AP 927 A	12-01-2001
		AT 219329 T	15-07-2002
		AU 712463 B2	04-11-1999
		AU 2504997 A	22-10-1997
		BG 63796 B1	31-01-2003
		BG 102804 A	30-07-1999
		BR 9708457 A	13-04-1999
		CA 2249332 A1	09-10-1997
		CN 1220579 A ,C	23-06-1999
		CZ 9802872 A3	13-01-1999
		DE 69713496 D1	25-07-2002
		DE 69713496 T2	13-02-2003
		DK 889692 T3	15-07-2002
		EA 1150 B1	30-10-2000
		EE 9800319 A	15-04-1999
		WO 9736488 A1	09-10-1997
		EP 0889692 A1	13-01-1999
		ES 2178768 T3	01-01-2003
		IL 126307 A	21-04-2002
		JP 2000507565 T	20-06-2000
		KR 2000005108 A	25-01-2000
		NZ 331764 A	27-03-2000
		OA 10888 A	18-02-2003
		PL 329125 A1	15-03-1999
		PT 889692 T	31-10-2002
		SI 889692 T1	31-10-2002
		SK 129898 A3	11-02-1999
		TR 9801934 T2	18-01-1999
		US 6083878 A	04-07-2000
		ZA 9702721 A	23-10-1997