

ORGANISATION AFRICAINE DE LA PROPRIETE INTELLECTUELLE
(O.A.P.I.)



(19)

(11) N°

12987

(51) Inter. Cl. 7

BREVET D'INVENTION

C07C 233/78 ; C07D 295/20
C07D 307/20 ; A61K 31/166
A61K 31/132 ; A61K 31/33

(21) Numéro de dépôt : 1200400227

(22) Date de dépôt : 27.02.2003

(30) Priorité(s) : US
27.02.2002 N° 60/359953

(24) Délivré le : 30.12.2005

(45) Publié le : 13 OCT 2006

(73) Titulaire(s) :

1- Elan Pharmaceuticals, Inc.
800 Gateway Boulevard
SOUTH SAN FRANCISCO
CA 94080 (US)

2- Pharmacia & Upjohn Company
301 Henrietta Street
KALAMAZOO, MI 49007 (US)

(72) Inventeur(s) :

1- BECK James P.
6839 Windermere Drive
ZIONSVILLE, IN 46077 (US)

2- GETMAN Daniel (US)
3- JAGODZINSKA Barbara (US)
4- MAILLARD Michel (US)
5- TENBRINK Ruth E. (US)
6- JOHN Varghese (US)


(74) Mandataire :

Cabinet J. EKEME
B.P. 6370, YAOUNDE (CM)

(54) Titre : Substituted hydroxyethylamines.

(57) Abrégé :

The present invention relates to prodrug compounds of the formula (Y) and pharmaceutical salts thereof:



where R_N , R_1 , R_2 , R_3 , R_4 and R_C are defined herein, which are useful in treating Alzheimer's disease and other similar diseases.

SUBSTITUTED HYDROXYETHYLAMINES

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. Provisional Application Serial No. 60/359,953, filed February 27, 2002, the disclosure of which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

1. Field of the Invention

This invention relates to prodrugs of a class of amine compounds which are useful in the treatment of Alzheimer's disease and similar diseases.

2. Description of the Related Art

Alzheimer's disease (AD) is a progressive degenerative disease of the brain primarily associated with aging. Clinical presentation of AD is characterized by loss of memory, cognition, reasoning, judgement, and orientation. As the disease progresses, motor, sensory, and linguistic abilities are also affected until there is global impairment of multiple cognitive functions. These cognitive losses occur gradually, but typically lead to severe impairment and eventual death in the range of four to twelve years.

Alzheimer's disease is characterized by two major pathologic observations in the brain: neurofibrillary tangles and beta amyloid (or neuritic) plaques, comprised predominantly of an aggregate of a peptide fragment known as A beta. Individuals with AD exhibit characteristic beta-amyloid deposits in the brain (beta amyloid plaques) and in cerebral blood vessels (beta amyloid angiopathy) as well as neurofibrillary tangles. Neurofibrillary tangles occur not only in Alzheimer's disease but also in other dementia-inducing disorders. On autopsy, large numbers of these lesions are

generally found in areas of the human brain important for memory and cognition.

Smaller numbers of these lesions in a more restricted anatomical distribution are found in the brains of most aged humans who do not have clinical AD. Amyloidogenic plaques and vascular amyloid angiopathy also characterize the brains of individuals with Trisomy 21 (Down's Syndrome), Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type (HCHWA-D), and other neurogenerative disorders. Beta-amyloid is a defining feature of AD, now believed to be a causative precursor or factor in the development of the disease. Deposition of A beta in areas of the brain responsible for cognitive activities is a major factor in the development of AD. Beta-amyloid plaques are predominantly composed of amyloid beta peptide (A beta, also sometimes designated betaA4). A beta peptide is derived by proteolysis of the amyloid precursor protein (APP) and is comprised of 39-42 amino acids. Several proteases called secretases are involved in the processing of APP.

Cleavage of APP at the N-terminus of the A beta peptide by beta-secretase and at the C-terminus by one or more gamma-secretases constitutes the beta-amyloidogenic pathway, i.e. the pathway by which A beta is formed. Cleavage of APP by alpha-secretase produces alpha-sAPP, a secreted form of APP that does not result in beta-amyloid plaque formation. This alternate pathway precludes the formation of A beta peptide. A description of the proteolytic processing fragments of APP is found, for example, in U.S. Patent Nos. 5,441,870; 5,721,130; and 5,942,400.

An aspartyl protease has been identified as the enzyme responsible for processing of APP at the beta-secretase cleavage site. The beta-secretase enzyme has been disclosed using varied nomenclature, including BACE, Asp, and Memapsin.

See, for example, Sinha et.al., 1999, *Nature* 402:537-554 (p501) and published PCT application WO00/17369.

Several lines of evidence indicate that progressive cerebral deposition of beta-amyloid peptide (A beta) plays a seminal role in the pathogenesis of AD and can precede cognitive symptoms by years or decades. See, for example, Selkoe, 1991, *Neuron* 6:487. Release of A beta from neuronal cells grown in culture and the presence of A beta in cerebrospinal fluid (CSF) of both normal individuals and AD patients has been demonstrated. See, for example, Seubert et al., 1992, *Nature* 359:325-327.

It has been proposed that A beta peptide accumulates as a result of APP processing by beta-secretase, thus inhibition of this enzyme's activity is desirable for the treatment of AD. In vivo processing of APP at the beta-secretase cleavage site is thought to be a rate-limiting step in A beta production, and is thus a therapeutic target for the treatment of AD. See for example, Sabbagh, M., et al., 1997, *Alz. Dis. Rev.* 3, 1-19.

BACE1 knockout mice fail to produce A beta, and present a normal phenotype. When crossed with transgenic mice that overexpress APP, the progeny show reduced amounts of A beta in brain extracts as compared with control animals (Luo et. al., 2001 *Nature Neuroscience* 4:231-232). This evidence further supports the proposal that inhibition of beta-secretase activity and reduction of A beta in the brain provides a therapeutic method for the treatment of AD and other beta amyloid disorders.

Published PCT application WO00/47618 entitled "Beta-Secretase Enzyme Compositions and Methods" identifies the beta-secretase enzyme and methods of its use. This publication also discloses oligopeptide inhibitors that bind the enzyme's active site and are useful in affinity column purification of the enzyme. In addition, WO00/77030 discloses tetrapeptide


inhibitors of beta-secretase activity that are based on a statine molecule.

At present there are no effective treatments for halting, preventing, or reversing the progression of Alzheimer's disease. Therefore, there is an urgent need for pharmaceutical agents capable of slowing the progression of Alzheimer's disease and/or preventing it in the first place.

Compounds that are effective inhibitors of beta-secretase, that inhibit beta-secretase-mediated cleavage of APP, that are effective inhibitors of A beta production, and/or are effective to reduce amyloid beta deposits or plaques, are needed for the treatment and prevention of disease characterized by amyloid beta deposits or plaques, such as AD.

SUMMARY OF THE INVENTION

The invention provides compounds of formula (Y):



and the pharmaceutically acceptable salts thereof,

where R_1 is:

(I) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_7$ alkyl (optionally substituted with $\text{C}_1\text{-C}_3$ alkyl and $\text{C}_1\text{-C}_3$ alkoxy), $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl, $-\text{OC}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(II) $-\text{CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)}$,

(III) $-\text{CH}_2\text{-CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)}$,

(IV) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,

(V) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,

(VI) $-(\text{CH}_2)_{n_1}\text{-(R}_{1-\text{aryl}})$ where n_1 is zero or one and where $\text{R}_{1-\text{aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthyl, tetralinyl optionally substituted with one,

two, three or four of the following substituents on the aryl ring:

(A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C≡N, -CF₃, C₁-C₃ alkoxy,

(B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(D) -F, Cl, -Br and -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(N) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$,

(VII) $-(\text{CH}_2)_{n_1}- (\text{R}_1\text{-heteroaryl})$ where n_1 is as defined above

and where $\text{R}_1\text{-heteroaryl}$ is selected from the group consisting of
 pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl,
 5 indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl,
 quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl,
 isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indoliziny, l,
 indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl,
 furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl,
 10 triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl,
 isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-
 carbolinyl, isochromanlyl, chromanyl, tetrahydroisoquinolinyl,
 isoindolinyl, isobenzotetrahydrofuranlyl,
 isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl,
 15 pyridopyridinyl, benzotetrahydrofuranlyl, benzotetrahydrothienyl,
 purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl,
 pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl,
 dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl,
 dihydrobenzisothiazinyl, benzopyranyl, benzothiopyranyl,
 20 coumarinyl, isocoumarinyl, chromonyl, chromanonyl, pyridinyl-N-
 oxide tetrahydroquinolinyl dihydroquinolinyl
 dihydroquinolinonyl dihydroisoquinolinonyl dihydrocoumarinyl
 dihydroisocoumarinyl isoindolinonyl benzodioxanyl
 benzoxazolinonyl pyrrolyl N-oxide, pyrimidinyl N-oxide,
 25 pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide,
 indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide,
 quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-
 oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide,
 thiazolyl N-oxide, indoliziny, l N-oxide, indazolyl N-oxide,
 30 benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-
 oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-
 oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide, where the $\text{R}_1\text{-heteroaryl}$ group is

bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent R_N -heteroaryl group substituted by hydrogen such that the new bond to the R_1 -heteroaryl group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or
 5 four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy,
 10

(2) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 15

(3) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 20

(4) $-F$, $-Cl$, $-Br$ and $-I$,

(6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three $-F$,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined
 25 below,

(8) $-OH$,

(9) $-C\equiv N$,

(10) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 30

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-\text{SO}_2-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(13) $-\text{CO}-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (14) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$, with the proviso that when n_1 is zero $\text{R}_{1-\text{heteroaryl}}$ is not bonded to the carbon chain by nitrogen,

(VIII) $-(\text{CH}_2)_{n_1}-(\text{R}_{1-\text{heterocycle}})$ where n_1 is as defined above and $\text{R}_{1-\text{heterocycle}}$ is selected from the group consisting of
 10 morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide, oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl
 15 dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide, and homothiomorpholinyl S-oxide where the $\text{R}_{1-\text{heterocycle}}$ group is bonded by any atom of the parent
 20 $\text{R}_{1-\text{heterocycle}}$ group substituted by hydrogen such that the new bond to the $\text{R}_{1-\text{heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) C_1-C_6 alkyl optionally substituted with
 25 one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy,

(2) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three
 30 substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or C_1-C_6 alkyl,

(3) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are
 5 -H or C_1-C_6 alkyl,

(4) -F, -Cl, -Br and -I,

(5) C_1-C_6 alkoxy,

(6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three -F,

10 (7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,

(9) $-C\equiv N$,

(10) C_3-C_7 cycloalkyl, optionally
 15 substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH
 $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

20 (12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$,

25 (15) =O, with the proviso that when n_1 is zero $R_{1-\text{heterocycle}}$ is not bonded to the carbon chain by nitrogen;

where R_2 is selected from the group consisting of:

(I) -H,

(II) C_1-C_6 alkyl, optionally substituted with one, two
 30 or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

5 (IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

15 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

20 where R₃ is selected from the group consisting of:

(I) -H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,

25 -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

(IV) C₂-C₆ alkenyl with one or two double bonds,

30 (V) C₂-C₆ alkynyl with one or two triple bonds,

- (VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, and seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, $-NR_{N-2}-$, where R_{N-2} is as defined below;

10 where R_N is:

(I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:

- (A) $-CO-$,
 (B) $-SO_2-$,
 15 (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,
 (D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above,
 20 (E) a single bond;

where R_{N-1} is selected from the group consisting of:

- (A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, dihydronaphthyl or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, optionally substituted
 25 with one, two or three of the following substituents which can be the same or different and are:
 (1) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$,
 30 $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $-OH$,

- (3) $-\text{NO}_2$,
- (4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
- (5) $-\text{CO}-\text{OH}$,
- (6) $-\text{C}\equiv\text{N}$,
- 5 (7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:
- (a) $-\text{H}$,
- (b) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted
- 10 with one substituent selected from the group consisting of:
- (i) $-\text{OH}$,
- (ii) $-\text{NH}_2$,
- (c) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one to three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
- 15 (d) $-\text{C}_3-\text{C}_7$ cycloalkyl,
- (e) $-(\text{C}_1-\text{C}_2 \text{ alkyl})-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
- (f) $-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_3 \text{ alkyl})$,
- (g) $-\text{C}_2-\text{C}_6$ alkenyl with one or two double bonds,
- 20 (h) $-\text{C}_2-\text{C}_6$ alkynyl with one or two triple bonds,
- (i) $-\text{C}_1-\text{C}_6$ alkyl chain with one double bond and one triple bond,
- (j) $-\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined
- 25 above,
- (k) $-\text{R}_1\text{-heteroaryl}$ where $\text{R}_1\text{-heteroaryl}$ is as defined above,
- (8) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
- (9) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl with one, two or three double bonds})$,
- 30 (10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl with one, two or three triple bonds})$,
- (11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(12) - $(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined above,

(13) - $(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_1\text{-heteroaryl}$ where $\text{R}_1\text{-heteroaryl}$ is as defined above,

5 (14) - $(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_1\text{-heterocycle}$ where $\text{R}_1\text{-heterocycle}$ is as defined above,

(15) - $(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, 10 homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of $\text{C}_1\text{-C}_6$ alkyl,

(16) - $(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ is selected from the group consisting of:

15 (a) $\text{C}_1\text{-C}_6$ alkyl,

(b) - $(\text{CH}_2)_{0-2}-(\text{R}_1\text{-aryl})$ where $\text{R}_1\text{-aryl}$ is as defined above,

(c) $\text{C}_2\text{-C}_6$ alkenyl containing one or two double bonds,

20 (d) $\text{C}_2\text{-C}_6$ alkynyl containing one or two triple bonds,

(e) $\text{C}_3\text{-C}_7$ cycloalkyl,

(f) - $(\text{CH}_2)_{0-2}-(\text{R}_1\text{-heteroaryl})$ where $\text{R}_1\text{-heteroaryl}$ is as defined above,

25 (17) - $(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are as defined above,

(18) - $(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1\text{-C}_8 \text{ alkyl})$,

(19) - $(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1\text{-C}_{12} \text{ alkyl})$,

(20) - $(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3\text{-C}_7 \text{ cycloalkyl})$,

30 (21) - $(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,

(22) - $(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,

- (23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,
- (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,
- 5 (25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,
- (26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,
- 10 (27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,
- (28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is $-H$ or C_1-C_4 alkyl,
- (29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,
- 15 (30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,
- (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,
- (32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,
- 20 (33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined above,
- (34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -F)$,
- 25 (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 30 (37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(38) - $(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above,

(39) - $(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,

- 5 (B) - R_{N} -heteroaryl where R_{N} -heteroaryl is selected from the group consisting of pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, 10 thiazolyl, indoliziny, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, 15 isochromanyl, chromanyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranyl, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranyl, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, henoxazinyl, phenothiazinyl, 20 pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoquinolyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisoquinolyl, benzopyranyl, benzothiopyranyl, coumarinyl, isocoumarinyl, chromonyl, chromanonyl, pyridinyl-N-oxide, tetrahydroquinolinyl dihydroquinolinyl 25 dihydroquinolinonyl dihydroisoquinolinonyl dihydrocoumarinyl dihydroisocoumarinyl isoindolinonyl benzodioxanyl benzoxazolinonyl pyrrolyl N-oxide, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, 30 quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indoliziny, indazolyl N-oxide, benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-

oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, and benzothiopyranyl S,S-dioxide, where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

(1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$, $C_1\text{-}C_3$ alkoxy, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

(2) -OH,

(3) -NO $_2$,

(4) -F, -Cl, -Br, -I,

(5) -CO-OH,

(6) -C $\equiv$ N,

(7) -(CH $_2$) $_{0-4}$ -CO-NR $_{N-2}$ R $_{N-3}$ where R $_{N-2}$ and R $_{N-3}$ are the same or different and are selected from the group consisting of:

(a) -H,

(b) - $C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH,

(ii) -NH $_2$,

(c) - $C_1\text{-}C_6$ alkyl optionally substituted with one to three -F, -Cl, -Br, -I,

(d) - $C_3\text{-}C_7$ cycloalkyl,

(e) -($C_1\text{-}C_2$ alkyl)-($C_3\text{-}C_7$ cycloalkyl),

(f) -($C_1\text{-}C_6$ alkyl)-O-($C_1\text{-}C_3$ alkyl),

(g) - $C_2\text{-}C_6$ alkenyl with one or two

double bonds,

- triple bonds,
- (h) $-C_2-C_6$ alkynyl with one or two
- (i) $-C_1-C_6$ alkyl chain with one double bond and one triple bond,
- 5 above,
- (j) $-R_1$ -aryl where R_1 -aryl is as defined
- defined above,
- (k) $-R_1$ -heteroaryl where R_1 -heteroaryl is as
- 10 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12}$ alkyl),
- (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkenyl with one, two or three double bonds),
- (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkynyl with one, two or three triple bonds),
- 15 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7$ cycloalkyl),
- (12) $-(CH_2)_{0-4}-CO-R_1$ -aryl where R_1 -aryl is as defined above,
- (13) $-(CH_2)_{0-4}-CO-R_1$ -heteroaryl where R_1 -heteroaryl is as defined above,
- 20 (14) $-(CH_2)_{0-4}-CO-R_1$ -heterocycle where R_1 -heterocycle is as defined above,
- (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,
- 25 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:
- 30 (a) C_1-C_6 alkyl,
- (b) $-(CH_2)_{0-2}-(R_1$ -aryl) where R_1 -aryl is as defined above,
- (c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple bonds,

(e) C_3-C_7 cycloalkyl,

(f) $-(CH_2)_{0-2}-(R_1\text{-heteroaryl})$ where $R_1\text{-heteroaryl}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

(21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where

R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where

R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N\text{-aryl-1}})_2$ where $R_{N\text{-aryl-1}}$ is $-H$ or C_1-C_4 alkyl,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,

(32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,

(33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$

5 (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

10 (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

15 (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above,

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

20 (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-heterocycle}}$ is defined as $\text{R}_{1-\text{heterocycle}}$ is defined above,

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

25 (H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-1-heterocycle}}$ is defined as $\text{R}_{1-\text{heterocycle}}$ is defined above,

(I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}},$

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

where W is

30 (1) $-(\text{CH}_2)_{0-4}-,$

(2) $-\text{O}-,$

(3) $-\text{S}(\text{O})_{0-2}-,$

(4) $-N(R_{N-5})-$ where R_{N-5} is as defined above, or

(5) $-CO-$;

(II) $-CO-(C_1-C_{10} \text{ alkyl})$ where alkyl is optionally substituted with one two or three substituents selected from the group consisting of:

(A) $-OH$,

(B) $-C_1-C_6$ alkoxy,

(C) $-C_1-C_6$ thioalkoxy,

(D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, C_1-C_6 alkyl or phenyl,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$,

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,

(R) $-F$, $-Cl$,

(III) $-CO-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

- (A) -OH,
- (B) -C₁-C₆ alkoxy,
- (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -
- 5 phenyl,
- (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
- (F) -CO-R_{N-4} where R_{N-4} is as defined above,
- (G) -SO₂-(C₁-C₈ alkyl),
- 10 (H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
- (I) -NH-CO-(C₁-C₆ alkyl),
- (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
- (K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
- 15 different and are as defined above,
- (L) -R_{N-4} where R_{N-4} is as defined above,
- (M) -O-CO-(C₁-C₆ alkyl),
- (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different and are as defined above,
- 20 (O) -O-(C₁-C₅ alkyl)-COOH,
- (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, -I),
- (Q) -NH-SO₂-(C₁-C₆ alkyl),
- (R) -F, -Cl,
- 25 (IV) -CO-(C₁-C₆ alkyl)-S-(C₁-C₆ alkyl) where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of :
- (A) -OH,
- (B) -C₁-C₆ alkoxy,
- (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
- (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
- 30

- (F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
 (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same
 or different and are as defined above,
 5 (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
 (K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or
 different and are as defined above,
 10 (L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where the $\text{R}_{\text{N}-8}$ s are the same or
 different and are as defined above,
 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with
 15 one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$),
 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (R) $-\text{F}$, $-\text{Cl}$,
 (V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$
 where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is
 20 selected from the group consisting of:
 (A) $-\text{H}$,
 (B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,
 (C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
 (D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,
 25 (E) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one triple bond,
 (F) $\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined above,
 (G) $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined
 above,
 (VI) $-\text{CO}-(\text{C}_3-\text{C}_8 \text{ cycloalkyl})$ where cycloalkyl is
 30 optionally substituted with one or two substituents selected
 from the group consisting of:
 (A) $-(\text{CH}_2)_{0-4}-\text{OH}$,

- (B) $-(CH_2)_{0-4}-C_1-C_6$ alkoxy,
 (C) $-(CH_2)_{0-4}-C_1-C_6$ thioalkoxy,
 (D) $-(CH_2)_{0-4}-CO-O-R_{N-8}$ where R_{N-8} is $-H$, C_1-C_6 alkyl or $-phenyl$,
 5 (E) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,
 (F) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is as defined above,
 (G) $-(CH_2)_{0-4}-SO_2-(C_1-C_8 \text{ alkyl})$,
 10 (H) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,
 (I) $-(CH_2)_{0-4}-NH-CO-(C_1-C_6 \text{ alkyl})$,
 (J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,
 (K) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the
 15 same or different and are as defined above,
 (L) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,
 (M) $-O-CO-(C_1-C_6 \text{ alkyl})$,
 (N) $-O-CO-NR_{N-8}R_{N-8}$ where the R_{N-8} s are the same or different and are as defined above,
 20 (O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,
 (P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$,
 (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,
 (R) $-F$, $-Cl$,
 25 where R_C is:
 (I) $-C_1-C_{10}$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O-phenyl$, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-OC=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-S(=O)_{0-2} R_{1-a}$ where R_{1-a} is as defined above, -

$\text{NR}_{1-a}\text{C}=\text{O NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{C}=\text{O NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, and $-\text{S}(=\text{O})_2 \text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{CO}-\text{OH}$, $-\text{CO}-\text{O}-(\text{C}_1-\text{C}_4 \text{ alkyl})$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(III) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are $-\text{H}$, C_1-C_4 alkyl optionally substituted with one or two $-\text{OH}$,

C_1-C_4 alkoxy optionally substituted with one, two, or three of $-\text{F}$,

$-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, C_2-C_6 alkenyl containing one or two double bonds, C_2-C_6 alkynyl containing one or two triple bonds, phenyl,

and where $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six and seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{NR}_{\text{N-2}}-$ and $\text{R}_{\text{C-aryl}}$ is defined as $\text{R}_{\text{N-aryl}}$ is defined above;

(IV) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$ is defined as $\text{R}_{\text{N-heteroaryl}}$ is defined above and $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(V) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}-\text{R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(VI) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}-\text{R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

- (VII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- (VIII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- 5 (IX) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-aryl}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above, and $\text{R}_{\text{C-heterocycle}}$ is defined as $\text{R}_{\text{N-heterocycle}}$ is defined above,
- (X) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- 10 (XI) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- (XII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- 15 (XIII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- (XIV) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,
- (XV) - $[\text{C}(\text{R}_{\text{C-1}})(\text{R}_{\text{C-2}})]_{1-3}\text{-CO-N}(\text{R}_{\text{C-3}})_2$ where $\text{R}_{\text{C-1}}$ and $\text{R}_{\text{C-2}}$ are the same or different and are selected from the group consisting of:
- 20 (A) -H,
- (B) - $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH,
- 25 -SH, -C $\equiv$ N, -CF $_3$, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, -NR $_{1-a}$ R $_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br,
- 30 -I, -OH, -SH, -C $\equiv$ N, -CF $_3$, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, -NR $_{1-a}$ R $_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents

selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$, C_1 - C_6 alkoxy, -O-phenyl, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

(E) - (CH $_2$) $_{1-2}$ -S(O) $_{0-2}$ -(C_1 - C_6 alkyl),

5 (F) - (CH $_2$) $_{0-4}$ - C_3 - C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$, C_1 - C_6 alkoxy, -O-phenyl, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

10 (G) - (C_1 - C_4 alkyl)-R $_{C'}$ -aryl where R $_{C'}$ -aryl is as defined for R $_{1-aryl}$,

(H) - (C_1 - C_4 alkyl)-R $_C$ -heteroaryl where R $_C$ -heteroaryl is as defined above,

(I) - (C_1 - C_4 alkyl)-R $_C$ -heterocycle where R $_C$ -heterocycle is as defined above,

15 (J) - R $_C$ -heteroaryl where R $_C$ -heteroaryl is as defined above,

(K) - R $_C$ -heterocycle where R $_C$ -heterocycle is as defined above,

20 (M) - (CH $_2$) $_{1-4}$ -R $_{C-4}$ -(CH $_2$) $_{0-4}$ -R $_{C'}$ -aryl where R $_{C-4}$ is -O-, -S- or -NR $_{C-5}$ - where R $_{C-5}$ is C_1 - C_6 alkyl, and where R $_{C'}$ -aryl is defined above,

(N) - (CH $_2$) $_{1-4}$ -R $_{C-4}$ -(CH $_2$) $_{0-4}$ -R $_C$ -heteroaryl where R $_{C-4}$ and R $_C$ -heteroaryl are as defined above,

(O) - R $_{C'}$ -aryl where R $_{C'}$ -aryl is as defined above, and where R $_{C-3}$ is the same or different and is:

(A) -H,

(B) - C_1 - C_6 alkyl optionally substituted with one, 30 two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents
 5 selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents
 10 selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from
 15 the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) -R_{C'-aryl} where R_{C'-aryl} is as defined above,

20 (G) -R_{C-heteroaryl} where R_{C-heteroaryl} is as defined above,

(H) -R_{C-heterocycle} where R_{C-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{C'-aryl} where R_{C'-aryl} is as defined above,

25 (J) -(C₁-C₄ alkyl)-R_{C-heteroaryl} where R_{C-heteroaryl} is as defined above,

(K) -(C₁-C₄ alkyl)-R_{C-heterocycle} where R_{C-heterocycle} is as defined above,

(XVI) -CH(R_{C-aryl})₂ where R_{C-aryl} are the same or
 30 different and are as defined above,

(XVII) -CH(R_{C-heteroaryl})₂ where R_{C-heteroaryl} are the same or different and are as defined above,

(XVIII) $-\text{CH}(\text{R}_{\text{C-aryl}})(\text{R}_{\text{C-heteroaryl}})$ where $\text{R}_{\text{C-aryl}}$ and $\text{R}_{\text{C-heteroaryl}}$ are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $\text{R}_{\text{C-aryl}}$ or $\text{R}_{\text{C-heteroaryl}}$ or $\text{R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-aryl}}$ or $\text{R}_{\text{C-heteroaryl}}$ or $\text{R}_{\text{C-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH , $\text{NR}_{\text{N-5}}$, O , $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $=\text{O}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) $\text{C}_2\text{-C}_{10}$ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $\text{C}_2\text{-C}_{10}$ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{C-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-aryl}}$ is as defined above and $\text{R}_{\text{C-6}}$ is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{C-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$ and $\text{R}_{\text{C-6}}$ is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{C-aryl}} \text{ or } \text{R}_{\text{C-heteroaryl}})-\text{CO-O}(\text{C}_1\text{-C}_4 \text{ alkyl})$ where $\text{R}_{\text{C-aryl}}$ and $\text{R}_{\text{C-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,


(XXV) $(\text{C}_1\text{-C}_6 \text{ alkyl})-\text{O}-(\text{C}_1\text{-C}_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; and

where R_4 is:



where x is 0 or 1 and R_5 is H or is:

- 5 (I) R_{p-1} wherein R_{p-1} is selected from the group consisting of:

(A) $-\text{Y}-R_{p\text{-aryl}}$ where Y is absent or is oxygen, and $R_{p\text{-aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, optionally
10 substituted with one, two or three of the following substituents which can be the same or different and are:

- (1) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
15 $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are independently $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,

(2) $-\text{OH}$,

(3) $-\text{NO}_2$,

(4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,

20 (5) $-\text{CO}-\text{OH}$,

(6) $-\text{C}\equiv\text{N}$,

(7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{p-2}\text{R}_{p-3}$ where R_{p-2} and R_{p-3} are the same or different and are selected from the group consisting of:

25 (a) $-\text{H}$,

(b) $-\text{C}_1\text{-C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-\text{OH}$,

(ii) $-\text{NH}_2$,

30 (c) $-\text{C}_1\text{-C}_6$ alkyl optionally substituted with one to three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,

- (d) $-C_3-C_7$ cycloalkyl,
- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
- (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
- (g) $-C_2-C_6$ alkenyl with one or two
- 5 double bonds,
- (h) $-C_2-C_6$ alkynyl with one or two
- triple bonds,
- (i) $-C_1-C_6$ alkyl chain with one double
- bond and one triple bond,
- 10 (j) $-R_P\text{-aryl}$ where $-R_P\text{-aryl}$ is defined
- above,
- (k) $-R_P\text{-heteroaryl}$ where $R_P\text{-heteroaryl}$ is
- defined below,
- (l) $-C(O)C_1-C_6$ alkyl,
- 15 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
- (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12})$ alkenyl with one,
- two or three double bonds,
- (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12})$ alkynyl with one,
- two or three triple bonds,
- 20 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
- (12) $-(CH_2)_{0-4}-CO-R_P\text{-aryl}$ where $R_P\text{-aryl}$ is as
- defined above,
- (13) $-(CH_2)_{0-4}-CO-R_P\text{-heteroaryl}$ where $R_P\text{-heteroaryl}$
- is as defined below,
- 25 (14) $-(CH_2)_{0-4}-CO-R_P\text{-heterocycle}$ where $R_P\text{-heterocycle}$
- is selected from the group consisting of:
- (A) morpholinyl,
- (B) thiomorpholinyl,
- (C) thiomorpholinyl S-oxide,
- (D) thiomorpholinyl S,S-dioxide,
- (E) piperazinyl,
- (F) homopiperazinyl,
- 30 (G) pyrrolidinyl,

- (H) pyrrolinyl,
- (I) tetrahydropyranyl,
- (J) piperidinyl,
- (K) tetrahydrofuranyl,
- (L) tetrahythiophenyl,
- (M) homopiperidinyl,
- (N) imidazolidine,
- (O) imidazolidine dione, and
- (P) dithiane,

where the $R_{P\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{P\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{P\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one thru four:

- (1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-O\text{-}CO\text{-}(C_1\text{-}C_6 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy,
- (2) $C_2\text{-}C_6$ alkenyl with one or two double bonds,
- (3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds,
- (4) $-F$, Cl , $-Br$ and $-I$,
- (5) $C_1\text{-}C_6$ alkoxy,
- (6) $-O\text{-}C_1\text{-}C_6$ alkyl optionally substituted with one thru three $-F$,
- (7) $-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are as defined above,
- (8) $-OH$,
- (9) $-C\equiv N$,
- (10) $C_3\text{-}C_7$ cycloalkyl,
- (11) $-CO\text{-}(C_1\text{-}C_4 \text{ alkyl})$,

- (12) $-\text{SO}_2-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are
as defined above,
- (13) $-\text{CO}-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are
as defined above,
- 5 (14) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$;
(15) $=\text{O}$; or
(16) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$;
(15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{N-4}$ where R_{N-4} is selected
from the group consisting of morpholinyl, thiomorpholinyl,
10 piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl,
homomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide,
pyrrolinyl and pyrrolidinyl where each group is optionally
substituted with one, two, three, or four of: $\text{C}_1-\text{C}_6 \text{ alkyl}$,
(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{N-5}$ where R_{N-5}
15 is selected from the group consisting of:
(a) $\text{C}_1-\text{C}_6 \text{ alkyl}$,
(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{P-\text{aryl}})$ where $\text{R}_{P-\text{aryl}}$ is as
defined above,
(c) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ containing one or two
20 double bonds,
(d) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ containing one or two
triple bonds,
(e) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
(f) $-(\text{CH}_2)_{0-2}-(\text{R}_{P-\text{heteroaryl}})$ where R_{P-}
25 heteroaryl is as defined below,
(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{P-2}\text{R}_{P-3}$ where R_{P-2} and R_{P-3}
are as defined above,
(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
30 (20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{N-5})-\text{CO}-\text{O}-\text{R}_{N-5}$ where
 R_{N-5} can be the same or different and is as defined above,

- (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- 5 (24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{P}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{P}-2}$ can be the same or different and are as defined above,
- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{P}-2}\text{R}_{\text{P}-3}$ where $\text{R}_{\text{P}-2}$ and $\text{R}_{\text{P}-3}$ can be the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined
- 10 above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is $-\text{H}$ or C_1-C_4 alkyl,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as
- 15 defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 20 (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of: } -\text{F})$,
- 25 (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b}
- 30 are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,

-OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{p-2} where R_{N-5} and R_{p-2} can be the same or different and are as described above,

(39) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl,

(B) -Y-R_{p-heteroaryl} where Y is absent or is oxygen, and R_{p-heteroaryl} is selected from the group consisting of:

10	(A) pyridinyl,
	(B) pyrimidinyl,
	(C) quinolinyl,
	(F) benzothienyl,
	(G) indolyl,
15	(H) indolinyl,
	(I) pyridazinyl,
	(J) pyrazinyl,
	(K) isoindolyl,
	(L) isoquinolyl,
20	(M) quinazolinyl,
	(N) quinoxalinyl,
	(O) phthalazinyl,
	(P) imidazolyl,
	(Q) isoxazolyl,
25	(R) pyrazolyl,
	(S) oxazolyl,
	(T) thiazolyl,
	(U) indolizinyl,
	(V) indazolyl,
30	(W) benzothiazolyl,
	(X) benzimidazolyl,
	(Y) benzofuranyl,
	(Z) furanyl,

	(AA) thienyl,
	(BB) pyrrolyl,
	(CC) oxadiazolyl,
	(DD) thiadiazolyl,
5	(EE) triazolyl,
	(FF) tetrazolyl,
	(II) oxazolopyridinyl,
	(JJ) imidazopyridinyl,
	(KK) isothiazolyl,
10	(LL) naphthyridinyl,
	(MM) cinnolinyl,
	(NN) carbazolyl,
	(OO) beta-carbolinyl,
	(PP) isochromanyl,
15	(QQ) chromanyl,
	(SS) tetrahydroisoquinolinyl,
	(TT) isoindolinyl,
	(UU) isobenzotetrahydrofuranyl,
	(VV) isobenzotetrahydrothienyl,
20	(WW) isobenzothienyl,
	(XX) benzoxazolyl,
	(YY) pyridopyridinyl,
	(ZZ) benzotetrahydrofuranyl,
	(AAA) benzotetrahydrothienyl,
25	(BBB) purinyl,
	(CCC) benzodioxolyl,
	(DDD) triazinyl,
	(EEE) phenoxazinyl,
	(FFF) phenothiazinyl,
30	(GGG) pteridinyl,
	(HHH) benzothiazolyl,
	(III) imidazopyridinyl,
	(JJJ) imidazothiazolyl,

- 5 (KKK) dihydrobenzisoxazinyl,
 (LLL) benzisoxazinyl,
 (MMM) benzoxazinyl,
 (NNN) dihydrobenzisothiazinyl,
 (OOO) benzopyranyl,
 (PPP) benzothiopyranyl,
 (QQQ) coumarinyl,
 (RRR) isocoumarinyl,
 (SSS) chromonyl,
 10 (TTT) chromanonyl,
 (UUU) pyridinyl-N-oxide,

where the $R_{P\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{P\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{P\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

(1) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -
 20 CF_3 , C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

- (2) -OH,
 (3) -NO₂,
 (4) -F, -Cl, -Br, -I,
 25 (5) -CO-OH,
 (6) -C $\equiv$ N,
 (7) $-(CH_2)_{0-4}-CO-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are selected from the group consisting of:

- 30 (a) -H,
 (b) $-C_1-C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH,

- (ii) -NH_2 ,
- (c) $\text{-C}_1\text{-C}_6$ alkyl optionally substituted with one to three -F , -Cl , -Br , -I ,
- (d) $\text{-C}_3\text{-C}_7$ cycloalkyl,
- 5 (e) $\text{-(C}_1\text{-C}_2 \text{ alkyl)-(C}_3\text{-C}_7 \text{ cycloalkyl)}$,
- (f) $\text{-(C}_1\text{-C}_6 \text{ alkyl)-O-(C}_1\text{-C}_3 \text{ alkyl)}$,
- (g) $\text{-C}_2\text{-C}_6$ alkenyl with one or two double bonds,
- (h) $\text{-C}_2\text{-C}_6$ alkynyl with one or two triple bonds,
- 10 (i) $\text{-C}_1\text{-C}_6$ alkyl chain with one double bond and one triple bond,
- (j) $\text{-R}_\text{P-aryl}$ where $\text{R}_\text{P-aryl}$ is as defined above,
- 15 (k) $\text{-R}_\text{P-heteroaryl}$ where $\text{R}_\text{P-heteroaryl}$ is as defined above,
- (8) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_1\text{-C}_{12} \text{ alkyl)}$,
- (9) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_2\text{-C}_{12} \text{ alkenyl with one, two or three double bonds)}$,
- 20 (10) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_2\text{-C}_{12} \text{ alkynyl with one, two or three triple bonds)}$,
- (11) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_3\text{-C}_7 \text{ cycloalkyl)}$,
- (12) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{P-aryl}$ where $\text{R}_\text{P-aryl}$ is as defined above,
- 25 (13) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{P-heteroaryl}$ where $\text{R}_\text{P-heteroaryl}$ is as defined above,
- (14) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{P-heterocycle}$ where $\text{R}_\text{P-heterocycle}$ is as defined above,
- (15) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally
- 30

substituted with one, two, three, or four of: C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5}

is selected from the group consisting of:

(a) C_1-C_6 alkyl,

5 (b) $-(CH_2)_{0-2}-(R_{P-aryl})$ where R_{P-aryl} is as defined above,

(c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two
10 triple bonds,

(e) C_3-C_7 cycloalkyl,

(f) $-(CH_2)_{0-2}-(R_{P-heteroaryl})$ where $R_{P-heteroaryl}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3}
15 are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

(21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where
20 R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where
 R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be
the same or different and is as defined above,

25 (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{P-2}$ where R_{N-5}
and R_{P-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} can
be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined
30 above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$
1 is $-H$ or C_1-C_4 alkyl,

- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 5 (31) $-(\text{CH}_2)_{0-4}-\text{O}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-\text{N}(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 10 (34) $-(\text{CH}_2)_{0-4}-\text{O}-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one, two, three, four, or five of: $-\text{F}$),
- (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double
- 15 bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple
- bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
- 20 $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{P}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{P}-2}$ can be the same or different and are as described above,
- 25 (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,
- (C) $\text{R}_{\text{P}-\text{aryl}}-\text{W}-\text{R}_{\text{P}-\text{aryl}}$,
- (D) $\text{R}_{\text{P}-\text{aryl}}-\text{W}-\text{R}_{\text{P}-\text{heteroaryl}}$,
- (E) $\text{R}_{\text{P}-\text{aryl}}-\text{W}-\text{R}_{\text{P}-\text{heterocycle}}$,
- (F) $\text{R}_{\text{P}-\text{heteroaryl}}-\text{W}-\text{R}_{\text{P}-\text{aryl}}$,
- 30 (G) $\text{R}_{\text{P}-\text{heteroaryl}}-\text{W}-\text{R}_{\text{P}-\text{heteroaryl}}$,
- (H) $\text{R}_{\text{P}-\text{heteroaryl}}-\text{W}-\text{R}_{\text{P}-\text{heterocycle}}$,
- (I) $\text{R}_{\text{P}-\text{heterocycle}}-\text{W}-\text{R}_{\text{P}-\text{aryl}}$,
- (J) $\text{R}_{\text{P}-\text{heterocycle}}-\text{W}-\text{R}_{\text{P}-\text{heteroaryl}}$,

(K) R_{P-1} -heterocycle-W- R_{P-1} -heterocycle,

where W is

(1) $-(CH_2)_{0-4}-$,

(2) $-O-$,

5 (3) $-S(O)_{0-2}-$,

(4) $-N(R_{N-5})-$ where R_{N-5} is as defined above;

(5) $-CO-$; or

(6) $-O-(CH_2)_{1-4}-$,

10 (L) $-Y-R_{P-1}$ -heterocycle where Y is absent or is oxygen, and R_{P-1} -heterocycle is defined above;

(II) $-(C_1-C_{10} \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

15 (A) $-OH$,

(B) $-C_1-C_6$ alkoxy,

(C) $-C_1-C_6$ thioalkoxy,

(D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, C_1-C_6 alkyl, phenyl or benzyl,

20 (E) $-CO-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-SO_2-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are as defined above,

25 (I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are as defined above,

30 (L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where the R_{N-8} s are the same or different and are as defined above,

- (O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,
- (P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$,
- (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,
- 5 (R) $-F$, $-Cl$,
- (III) $-X-T-(C_1-C_6 \text{ alkyl})$ where X is absent or is C_1-C_6 alkyl, T is oxygen or is NR_{P-2} , and where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:
- 10 (A) $-OH$,
- (B) $-C_1-C_6$ alkoxy,
- (C) $-C_1-C_6$ thioalkoxy,
- (D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, C_1-C_6 alkyl, phenyl, or benzyl,
- 15 (E) $-CO-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are as defined above,
- (F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,
- (G) $-SO_2-(C_1-C_8 \text{ alkyl})$,
- (H) $-SO_2-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same
- 20 or different and are as defined above,
- (I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,
- (J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,
- (K) $-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are as defined above,
- 25 (L) $-R_{N-4}$ where R_{N-4} is as defined above,
- (M) $-O-CO-(C_1-C_6 \text{ alkyl})$,
- (N) $-O-CO-NR_{N-8}R_{N-8}$ where the R_{N-8} s are the same or different and are as defined above,
- (O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,
- 30 (P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$,
- (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,
- (R) $-F$, $-Cl$,

(S) $-O-R_{p-aryl}$,

(T) $-O-R_{p-heteroaryl}$, OR

(U) $-O-R_{p-hetrocycle}$,

(IV) $-(C_1-C_6 \text{ alkyl})-Z-(C_1-C_6 \text{ alkyl})$ where Z is S or O,

5 and alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

(A) $-OH$,

(B) $-C_1-C_6$ alkoxy,

(C) $-C_1-C_6$ thioalkoxy,

10 (D) $-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(E) $-CO-NR_{p-2}R_{p-3}$ where R_{p-2} and R_{p-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

15 (H) $-SO_2-NR_{p-2}R_{p-3}$ where R_{p-2} and R_{p-3} are the same or different and are as defined above,

(I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-NR_{p-2}R_{p-3}$ where R_{p-2} and R_{p-3} are the same or
20 different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where the R_{N-8} s are the same or
different and are as defined above,

25 (O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$,

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,

(R) $-F$, $-Cl$,

30 (V) $-CH(-(CH_2)_{0-2}-O-R_{N-10})-(CH_2)_{0-2}-R_{p-aryl}/R_{p-heteroaryl}$

where R_{p-aryl} and $R_{p-heteroaryl}$ are as defined above, where R_{N-10} is selected from the group consisting of:

(A) $-H$,

- (B) C₁-C₆ alkyl,
(C) C₃-C₇ cycloalkyl,
(D) C₂-C₆ alkenyl with one double bond,
(E) C₂-C₆ alkynyl with one triple bond,
5 (F) R_{P-aryl} where R_{P-aryl} is as defined above,
(G) R_{P-heteroaryl} where R_{P-heteroaryl} is as defined
above,

(VI) -(C₃-C₈ cycloalkyl) where cycloalkyl is
optionally substituted with one or two substituents selected
10 from the group consisting of:

- (A) -(CH₂)₀₋₄-OH,
(B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
(C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
(D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆
15 alkyl, phenyl or benzyl,
(E) -(CH₂)₀₋₄-CO-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are
the same or different and are as defined above,
(F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined
above,
20 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
(H) -(CH₂)₀₋₄-SO₂-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are
the same or different and are as defined above,
(I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
25 (K) -(CH₂)₀₋₄-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the
same or different and are as defined above,
(L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
(M) -O-CO-(C₁-C₆ alkyl),
(N) -O-CO-NR_{N-8}R_{N-8} where the R_{N-8}s are the same or
30 different and are as defined above,
(O) -O-(C₁-C₅ alkyl)-COOH,

- (P) $-O-C_1-C_6$ alkyl optionally substituted with one, two, or three of: $-F$, $-Cl$, $-Br$, $-I$,
- (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,
- (R) $-F$, $-Cl$;
- 5 (VII) $-(CR_{P-x}R_{P-y})_{0-4}-R_{P-1}$ where R_{P-1} is as defined above and each of R_{P-x} and R_{P-y} are independently any combination of:
- (A) $-H$,
- (B) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group
- 10 consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-COOH$, $-COO(C_1-C_6 \text{ alkyl})$, $-CONR_{1a}R_{1b}$ and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (C) $-CH_2-S-(C_1-C_6 \text{ alkyl})$,
- 15 (D) $-CH_2-CH_2-S-(C_1-C_6 \text{ alkyl})$,
- (E) C_2-C_6 alkenyl with one or two double bonds,
- (F) C_2-C_6 alkynyl with one or two triple bonds,
- (G) $-(CH_2)_{n_1}-(R_{P-aryl})$ where n_1 is zero or one, where R_{P-aryl} is as defined above,
- 20 (H) $-(CH_2)_{n_1}-(R_{P-heteroaryl})$ where n_1 and $R_{P-heteroaryl}$ are as defined above, or
- (I) $-(CH_2)_{n_1}-(R_{P-heterocycle})$ where n_1 and $R_{P-heterocycle}$ are as defined above,
- (J) $-(CH_2)_{0-3}-(C_3-C_7)$ cycloalkyl where cycloalkyl
- 25 can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-CO-OH$, $-CO-O-(C_1-C_4 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and
- 30 R_{1-b} are as defined above, or
- (K) F , Cl , Br or I ,
- (L) $-NR_{P-2}$;

- or R_{p-x} and R_{p-y} are taken together with the carbon to which they are attached to form a carbocycle of three thru seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of
- 5 -O-, -S-, -SO₂-, and -NR_{p-2}-;
- (VIII) -(C₁-C₆ alkyl)-Z-(C₁-C₆ alkyl)-T-(C₁-C₆ alkyl)- wherein Z is S or O, T is O or NR_{p-2}, and alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:
- 10 (A) -OH,
- (B) -C₁-C₆ alkoxy,
- (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
- (E) -CO-NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same
- 15 or different and are as defined above,
- (F) -CO-R_{N-4} where R_{N-4} is as defined above,
- (G) -SO₂-(C₁-C₈ alkyl),
- (H) -SO₂-NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same or different and are as defined above,
- 20 (I) -NH-CO-(C₁-C₆ alkyl),
- (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
- (K) -NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same or different and are as defined above,
- (L) -R_{N-4} where R_{N-4} is as defined above,
- 25 (M) -O-CO-(C₁-C₆ alkyl),
- (N) -O-CO-NR_{N-8}R_{N-8} where the R_{N-8}s are the same or different and are as defined above,
- (O) -O-(C₁-C₅ alkyl)-COOH,
- (P) -O-(C₁-C₆ alkyl) optionally substituted with
- 30 one, two, or three of -F, -Cl, -Br, -I,
- (Q) -NH-SO₂-(C₁-C₆ alkyl),
- (R) -F, -Cl.

Also disclosed is a method of treating a patient who has, or in preventing a patient from getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for
5 treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating
10 cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive
15 supranuclear palsy, dementia associated with cortical basal degeneration, or diffuse Lewy body type of Alzheimer's disease and who is in need of such treatment which comprises administration of a therapeutically effective amount of a compound of formula (Y).

20 Also disclosed are methods for inhibiting beta-secretase activity, for inhibiting cleavage of amyloid precursor protein (APP), in a reaction mixture, at a site between Met596 and Asp597, numbered for the APP-695 amino acid isotype; or at a corresponding site of an isotype or mutant thereof, for
25 inhibiting production of amyloid beta peptide (A beta) in a cell, for inhibiting the production of beta-amyloid plaque in an animal, and for treating or preventing a disease characterized by beta-amyloid deposits in the brain which comprise administration of a therapeutically effective amount
30 of a compound of formula (Y), and pharmaceutically acceptable salts thereof.

Disclosed is a pharmaceutical composition which comprises a compound of formula (Y), and one or more pharmaceutically acceptable inert carriers.

Disclosed is the use of a compound of formula (Y), and
5 pharmaceutically acceptable salts thereof for the manufacture of a medicament for use in treating a patient who has, or in preventing a patient from getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for
10 treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating
15 cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive
20 supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease and who is in need of such treatment.

The present invention provides compounds, compositions, kits, and methods for inhibiting beta-secretase-mediated
25 cleavage of amyloid precursor protein (APP). More particularly, the compounds, compositions, and methods of the invention are effective to inhibit the production of A beta peptide and to treat or prevent any human or veterinary disease or condition associated with a pathological form of A beta
30 peptide.

The invention also provides methods of preparing the compounds of the invention and the intermediates used in those methods.

The compounds, compositions, and methods of the invention are useful for treating humans who have Alzheimer's Disease (AD), for helping prevent or delay the onset of AD, for treating patients with mild cognitive impairment (MCI), and
5 preventing or delaying the onset of AD in those patients who would otherwise be expected to progress from MCI to AD, for treating Down's syndrome, for treating Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type, for treating cerebral beta-amyloid angiopathy and preventing its potential
10 consequences such as single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, for treating dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia
15 associated with cortical basal degeneration, and diffuse Lewy body type AD, and for treating frontotemporal dementias with parkinsonism (FTDP).

The compounds of the invention possess beta-secretase inhibitory activity. The inhibitory activities of the
20 compounds of the invention are readily demonstrated, for example, using one or more of the assays described herein or known in the art.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the invention provides compounds of formula Y

Preferred compounds of formula Y include those of formula Y-1, i.e., compounds of formula Y wherein

R₁ is R₁-aryl, R₁-heteroaryl, R₁-heterocycle, -C₁-C₆ alkyl- R₁-aryl, -C₁-C₆ alkyl- R₁-heteroaryl, or -C₁-C₆ alkyl- R₁-heterocycle, where the ring portions of each are optionally substituted as indicated above.

More preferred compounds of formula Y-1 include those wherein

R₁ is -C₁-C₆ alkyl- R₁-aryl, -C₁-C₆ alkyl- R₁-heteroaryl, or -C₁-C₆ alkyl- R₁-heterocycle, where the ring portions of each are optionally substituted as indicated above.

Still more preferred compounds of formula Y-1 include those wherein

R₁ is -(CH₂)- R₁-aryl, -(CH₂)- R₁-heteroaryl, or -(CH₂)- R₁-heterocycle, where the ring portions of each are optionally substituted as indicated above.

Yet more preferred compounds of formula Y-1 include those wherein

R₁ is -CH₂-phenyl or -CH₂-pyridinyl where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, C₁-C₄ alkoxy, hydroxy, -NO₂, and C₁-C₄ alkyl, wherein C₁-C₄ alkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halogen, OH, SH, NH₂, NH(C₁-C₆ alkyl), N-(C₁-C₆ alkyl)(C₁-C₆ alkyl), C≡N, CF₃.

Still more preferred compounds of formula Y-1 include those wherein

R₁ is -CH₂-phenyl or -CH₂-pyridinyl where the phenyl or pyridinyl rings are each optionally substituted with 1 or 2 groups independently selected from halogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, hydroxy, -CF₃, and -NO₂.

Preferred compounds of formula Y-1 also include those wherein

R_1 is $-\text{CH}_2$ -phenyl or $-\text{CH}_2$ -pyridinyl where the phenyl or pyridinyl rings are each optionally substituted with 2 groups independently selected from halogen, C_1 - C_2 alkyl, C_1 - C_2 alkoxy, hydroxy, and $-\text{NO}_2$.

Still yet more preferred compounds of formula Y-1 include those wherein R_1 is $-\text{CH}_2$ -pyridinyl, benzyl, 3,5-difluorobenzyl, or 5-hydroxybenzyl.

Other preferred compounds of formula Y and formula Y-1 include compounds of formula Y-2, i.e., those of formula Y or Y-1 wherein

R_N is $-\text{CO}-R_{N-1}$, wherein R_{N-1} is as defined above, or R_N is $-\text{CO}-(\text{C}_1-\text{C}_{10} \text{ alkyl})$, wherein C_1-C_{10} alkyl is optionally substituted as indicated above.

Preferred compounds of formula Y-2 include compounds wherein

R_N is $-\text{CO}-R_{N-1}$, wherein R_{N-1} is R_N -aryl, R_N -heteroaryl, or R_N -heterocycle, each of which is optionally substituted as indicated above, or R_N is $-\text{CO}-(\text{C}_1-\text{C}_{10} \text{ alkyl})$, wherein C_1-C_{10} alkyl is optionally substituted as indicated above.

Preferred compounds of formula Y-2 include compounds wherein

R_N is $-\text{CO}-R_{N-1}$, wherein R_{N-1} is R_N -aryl, or R_N -heteroaryl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from $-\text{OR}$, $-\text{NO}_2$, C_1-C_5 alkyl, halogen, $-\text{C}\equiv\text{N}$, $-\text{OCF}_3$, $-\text{CF}_3$, $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{N-2}\text{R}_{N-3}$, and $-(\text{CH}_2)_{0-4}-\text{N}(\text{R}_{N-2})-\text{SO}_2-\text{R}_{N-2}$.

Preferred compounds of formula Y-2 include compounds wherein

R_N is $-\text{C}(=\text{O})$ -phenyl or $-\text{C}(=\text{O})$ -pyridinyl where each ring is independently optionally substituted with 1 or 2 groups

independently selected from C_1-C_6 alkyl, halogen, $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$, $-(CH_2)_{0-4}-N(R_{N-2})-SO_2-R_{N-2}$.

Preferred compounds of formula Y-2 include compounds wherein

- 5 R_N is $-C(=O)$ -phenyl or $-C(=O)$ -pyridinyl where each ring is independently substituted with $-CO-NR_{N-2}R_{N-3}$.

Preferred compounds of formula Y-2 include compounds wherein

- 10 R_N is $-C(=O)$ -phenyl or $-C(=O)$ -pyridinyl where each ring is independently substituted with $-CO-N(C_1-C_6 \text{ alkyl})(C_1-C_6 \text{ alkyl})$.


Other preferred compounds of formula Y, formula Y-1 and formula Y-2 include compounds of formula Y-3, i.e., those of formula Y, Y-1 or Y-2 wherein

- 15 R_4 is $-(CH_2-O)-(C=O)-R_5$, or $-(C=O)-R_5$, wherein R_5 is as defined above.


Preferred compounds of formula Y-3 include those wherein R_5 is:

- 20 $-(CR_{P-x}R_{P-y})_{0-4}-R_{P-1}$, $-(C_1-C_{10} \text{ alkyl})$, $-X-T-(C_1-C_6 \text{ alkyl})$, $-(C_1-C_6 \text{ alkyl})-Z-(C_1-C_6 \text{ alkyl})$, wherein each of these groups is as defined above.

Preferred compounds of formula Y-3 include those wherein R_4 is:



Compounds (Y) of the invention are prodrugs that are useful in treating and preventing Alzheimer's disease. Without being bound by any particular theory, it is believed that the produg compounds (Y) are converted in vivo to the active compounds by hydrolytic or enzymatic cleavage at the R_4 position, as shown in equation (I):



The prodrugs of the invention provide increased aqueous solubility, higher oral bioavailability and facile in vivo generation of the active ingredient.

The produg compounds (Y) can be made by methods well known to those skilled in the art from starting compounds known to those skilled in the art. The process chemistry is known to those skilled in the art. The most general process to prepare the compounds (Y) of the present invention is set forth in CHART A. The chemistry is straight forward and in summary involves the steps of N-protecting an amino acid (I) starting material to produce the corresponding protected amino acid (II), reaction of the protected amino acid (II) with diazomethane followed by work-up to add a carbon atom to produce the corresponding protected compound (III), reduction of the protected compound (III) to the corresponding alcohol (IV), formation of the corresponding epoxide (V), opening of the epoxide (V) with a C-terminal amine, $\text{R}_\text{C}-\text{NH}_2$ (VI) to produce the corresponding protected alcohol (VII) which then has the nitrogen protecting group removed to produce the corresponding amine (VIII), which is then reacted with an amide forming agent of the formula $(\text{R}_{\text{N}-1}-\text{X}_\text{N})_2\text{O}$ or $\text{R}_{\text{N}-1}-\text{X}_\text{N}-\text{X}_2$ or $\text{R}_{\text{N}-1}-\text{X}_\text{N}-\text{OH}$ (IX) to

produce the amine (X). The amine groups of amine (X) are then protected using, for example, t-butoxycarbonyl, to form protected amine (B). Other protecting groups can be used as is known to a person of ordinary skill in the art. Protected
5 amine (B) is converted to compound (C) by methods well known in the art (see EXAMPLES 1260-1275 as illustrative, but non-limiting examples). Compound (C) is deprotected using, for example, trifluoroacetic acid, to provide compounds (Y) of the invention. One skilled in the art will appreciate that these
10 are all well known reactions in organic chemistry. A chemist skilled in the art, knowing the chemical structure of the compounds (Y) of the invention would be able to prepare them by known methods from known starting materials without any additional information. The explanation below therefore is not
15 necessary but is deemed helpful to those skilled in the art who desire to make the compounds of the present invention.

The hydroxyethylamine moiety, -NH-CH(R)-CH(OH)- , of the compounds of the invention, can be readily prepared by methods disclosed in the literature and known to those skilled in the
20 art. For example, *J. Med. Chem.*, 36, 288-291 (1992), *Tetrahedron Letters*, 28, 5569-5572 (1987), *J. Med. Chem.*, 38, 581-584 (1994) and *Tetrahedron Letters*, 38, 619-620 (1997) all disclose processes to prepare hydroxyethylamine type compounds.

CHART A sets forth a general method used in the present
25 invention to prepare the compounds (Y). The anti-Alzheimer compounds (Y) of the present invention are prepared by starting with the corresponding amino acid (I). The amino acids (I) are well known to those skilled in the art or can be readily prepared from known compounds by methods well known to those
30 skilled in the art. The compounds (Y) of the present invention have at least two enantiomeric centers which give four enantiomers. The first of these enantiomeric centers derives from the amino acid starting material (I). It is preferred to

commercially obtain or produce the desired enantiomer (S) rather than produce an enantiomerically impure mixture and then have to separate out the desired enantiomer (S). It is preferred to start the process with enantiomerically pure (S)-
 5 amino acid (I) of the same configuration as that of the compound (Y) product. For the amino acids (I), R_1 is as defined above for compound (Y).

It is further preferred that R_1 be $-(CH_2)_{0-1}-(R_1\text{-aryl})$ or $-(CH_2)_{n1}-(R_1\text{-heteroaryl})$. It is more preferred that R_1 is $-(CH_2)-(R_1\text{-aryl})$ or $-(CH_2)-(R_1\text{-heteroaryl})$. It is further preferred that R_1 is $-(CH_2)-(R_1\text{-aryl})$ where $R_1\text{-aryl}$ is phenyl. It is even more preferred that R_1 is $-(CH_2)-(R_1\text{-aryl})$ where $R_1\text{-aryl}$ is phenyl substituted with two -F. It is additionally preferred that the -F substitution is 3,5-difluorobenzyl.

15 When R_1 is $R_1\text{-heteroaryl}$ or $R_1\text{-heterocycle}$ the bond from the $R_1\text{-heteroaryl}$ or $R_1\text{-heterocycle}$ group to the $-(CH_2)_{n1}$ - group can be from any ring atom which has an available valence provided that such bond does not result in formation of a charged species or unstable valence. This means that the $R_1\text{-heteroaryl}$ or $R_1\text{-heterocycle}$
 20 group is bonded to $-(CH_2)_{n1}$ -by any ring atom of the parent $R_1\text{-heteroaryl}$ or $R_1\text{-heterocycle}$ group which was substituted by hydrogen such that the new bond to the $R_1\text{-heteroaryl}$ or $R_1\text{-heterocycle}$ group replaces the hydrogen atom and its bond.

25 The first step of the process is to protect the free amino group of the (S)-amino acid (I) with an amino protecting group to produce the (S)-protected amino acid (II) by methods well known to those skilled in the art. Amino protecting groups are well known to those skilled in the art. See for example,
 30 "Protecting Groups in Organic Synthesis", John Wiley and sons, New York, N.Y., 1981, Chapter 7; "Protecting Groups in Organic Chemistry", Plenum Press, New York, N.Y., 1973, Chapter 2. The function of the amino protecting group is to protect the free

amino functionality (-NH₂) during subsequent reactions on the (S)-amino acid (I) which would not proceed well, either because the amino group would react and be functionalized in a way that is inconsistent with its need to be free for subsequent reactions, or the free amino group would interfere in the reaction. When the amino protecting group is no longer needed, it is removed by methods well known to those skilled in the art. By definition the amino protecting group must be readily removable as is known to those skilled in the art by methods well known to those skilled in the art. Suitable amino PROTECTING GROUP is selected from the group consisting of t-butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl, dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluy)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycabonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxyl)benzyloxycarbonyl,

isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, -CH-CH=CH₂ and phenyl-C(=N-)-H. It is preferred that the protecting group be t-butoxycarbonyl (BOC) and benzyloxycarbonyl (CBZ), it is more preferred that the protecting group be t-butoxycarbonyl. One skilled in the art will understand the preferred methods of introducing a t-butoxycarbonyl or benzyloxycarbonyl protecting group and may additionally consult T.W. Green and P.G.M. Wuts in "Protective Groups in Organic Chemistry," John Wiley and Sons, 1991 for guidance.

The (S)-protected amino acid (II) is transformed to the corresponding (S)-protected compound (III) by two different methods depending on the nature of R₂ and R₃. R₂ and R₃ are as defined above.

It is preferred that R₂ and R₃ both be -H. If R₂ and R₃ are not the same, an additional enantiomeric center is added to the molecule. If it is desired that both R₂ and R₃ are -H, then the (S)-protected amino acid (II) is reacted with diazomethane, as is well known to those skilled in the art, followed by reaction with a compound of the formula H-X₁ to produce the (S)-protected compound (III). X₁ includes -Cl, -Br, -I, -O-tosylate, -O-mesylate, -O-nosylate.; it is preferred that -X₁ be -Br or -Cl. Suitable reaction conditions include running the reaction in inert solvents, such as but not limited to ether, tetrahydrofuran and the like. The reactions from the (S)-protected amino acid (II) to the (S)-protected compound (III) are carried out for a period of time between 10 minutes and 1 day and at temperatures ranging from -78 degrees to 20-25 degrees C. It is preferred to conduct the reactions for a period of time between 1-4 hours and at temperatures between -30 degrees to -10 degrees C. This process adds one methylene group.

Alternatively, the (S)-protected compounds of formula (III) can be prepared by first converting the (S)-protected amino acid (II) to a corresponding methyl or ethyl ester, according to methods well established in the art, followed by treatment with a reagent of formula $X_1-C(R_2)(R_3)-X_1$ and a strong metal base. The base serves to affect a halogen-metal exchange, where the $-X_1$ undergoing exchange is a halogen selected from chlorine, bromine or iodine. The nucleophilic addition to the ester derivative gives directly the (S)-protected compound (III). Suitable bases include, but are not limited to the alkylolithiums including, for example, *sec*-butyllithium, *n*-butyllithium, and *t*-butyllithium. The reactions are preferably conducted at low temperature, such as -78 degrees C. Suitable reaction conditions include running the reaction in inert solvents, such as but not limited to, ether, tetrahydrofuran and the like. Where R_2 and R_3 are both hydrogen, then examples of $X_1-C(R_2)(R_3)-X_1$ include dibromomethane, diiodomethane, chloriodomethane, bromiodomethane and bromochloromethane. One skilled in the art knows the preferred conditions required to conduct this reaction. Furthermore, if R_2 and/or R_3 are not $-H$, then by the addition of $-C(R_2)(R_3)-X_1$ to esters of the (S)-protected amino acid (II) to produce the (S)-protected compound (III), an additional chiral center will be incorporated into the product, provided that R_2 and R_3 are not the same.

The (S)-protected compound (III) is then reduced by means well known to those skilled in the art for reduction of a ketone to the corresponding secondary alcohol affording the corresponding alcohol (IV). The means and reaction conditions for reducing the (S)-protected compound (III) to the corresponding alcohol (IV) include, for example, sodium borohydride, lithium borohydride, borane, diisobutylaluminum hydride, and lithium aluminium hydride. Sodium borohydride is

the preferred reducing agent. The reductions are carried out for a period of time between 1 hour and 3 days at temperatures ranging from -78 degrees C to elevated temperature up to the reflux point of the solvent employed. It is preferred to conduct the reduction between -78 degrees C and 0 degrees C. If borane is used, it may be employed as a complex, for example, borane-methyl sulfide complex, borane-piperidine complex, or borane-tetrahydrofuran complex. The preferred combination of reducing agents and reaction conditions needed are known to those skilled in the art, see for example, Larock, R.C. in Comprehensive Organic Transformations, VCH Publishers, 1989. The reduction of the (S)-protected compound (III) to the corresponding alcohol (IV) produces the second chiral center (third chiral center if R₂ and R₃ are not the same). The reduction of the (S)-protected compound (III) produces a mixture of enantiomers at the second center, (S, R/S)-alcohol (IV). This enantiomeric mixture is then separated by means known to those skilled in the art such as selective low-temperature recrystallization or chromatographic separation, for example by HPLC, employing commercially available chiral columns. The enantiomer that is used in the remainder of the process of CHART A is the (S,S)-alcohol (IV) since this enantiomer will give the desired biologically active anti-Alzheimer (S,R)-substituted compound (Y).

The (S, S)-alcohol (IV) is transformed to the corresponding epoxide (V) by means known to those skilled in the art. The stereochemistry of the (S)-(IV) center is maintained in forming the epoxide (V). A preferred means is by reaction with base, for example, but not limited to, hydroxide ion generated from sodium hydroxide, potassium hydroxide, lithium hydroxide and the like. Reaction conditions include the use of C₁-C₆ alcohol solvents; ethanol is preferred. A common co-solvent, such as for example, ethyl acetate may also

be employed. Reactions are conducted at temperatures ranging from -45 degrees C up to the reflux temperature of the alcohol employed; preferred temperature ranges are between -20 degrees C and 20-25 degrees C.

5 The epoxide (V) is then reacted with the appropriately substituted C-terminal amine, R_C-NH_2 (VI) by means known to those skilled in the art that opens the epoxide to produce the desired corresponding enantiomerically pure (S,R)-protected alcohol (VII). The substituted C-terminal amines, R_C-NH_2 (VI)
10 of this invention are commercially available or can be readily prepared from known compounds using literature procedures. R_C is as defined above.

It is preferred that R_C is:

- C₁-C₈ alkyl,
- 15 - (CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-aryl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-heteroaryl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-heterocycle,
- cyclopentyl or -cyclohexyl ring fused to R_C-aryl or R_C-
20 heteroaryl OR
R_C-heterocycle.

It is more preferred that R_C is:

- (CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-aryl,
- 25 - (CR_{C-x}R_{C-y})₀₋₄-R_C-heteroaryl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-heterocycle,
- cyclopentyl or -cyclohexyl ring fused to a R_C-aryl OR
R_C-heteroaryl OR
R_C-heterocycle.

30 It is even more preferred that R_C is:

- (CR_{C-x}R_{C-y})₀₋₄-R_C-aryl,
- (CR_{C-x}R_{C-y})₀₋₄-R_C-heteroaryl,

-cyclopentyl or -cyclohexyl ring fused to a R_C -aryl or R_C -heteroaryl or R_C -heterocycle.

It is still more preferred that R_C is selected from the group consisting of:

- ($CR_{C-x}R_{C-y}$)₀₋₄- R_C -aryl where R_C -aryl is phenyl,
- ($CR_{C-x}R_{C-y}$)₀₋₄- R_C -heteroaryl,
- cyclopentyl or -cyclohexyl ring fused to a R_C -aryl or R_C -heteroaryl or R_C -heterocycle. Further, it is preferred that when R_C is phenyl, it is substituted in the 3-position or 3,5-positions.

Suitable reaction conditions for opening the epoxide (V) include running the reaction in a wide range of common and inert solvents. C_1 - C_6 alcohol solvents are preferred and isopropyl alcohol most preferred. The reactions can be run at temperatures ranging from 20-25 degrees C up to the reflux temperature of the alcohol employed. The preferred temperature range for conducting the reaction is between 50 degrees C up to the reflux temperature of the alcohol employed. When the substituted C-terminal amine (VI) is a 1-amino-3,5-cis-dimethyl cyclohexyldicarboxylate it is preferably prepared as follows. To dimethyl-5-aminoisophthalate in acetic acid and methanol, is added rhodium in alumina in a high-pressure bottle. The bottle is saturated with hydrogen at 55 psi and shaken for one week of time. The mixture is then filtered through a layer of diatomaceous earth and rinsed with methanol three times, the solvents are removed under reduced pressure (with heat) to give a concentrate. The concentrate is triturated with ether and filtered again to give the desired C-terminal amine (VI). When the substituted C-terminal amine (VI) is 1-amino-3,5-cis-dimethoxy cyclohexane it is prepared by following the general procedure above and making non-critical variations but starting with 3,5-dimethoxyaniline. When the substituted C-terminal

amine (VI) is an aminomethyl group where the substituent on the methyl group is an aryl group, for example $\text{NH}_2\text{-CH}_2\text{-Rc-aryl}$, and $\text{NH}_2\text{-CH}_2\text{-Rc-aryl}$ is not commercially available it is preferably prepared as follows. A suitable starting material is the (appropriately substituted) aralkyl compound. The first step is bromination of the alkyl substituent via methods known to those skilled in the art, see for example R.C. Larock in Comprehensive Organic Transformations, VCH Publishers, 1989, p. 313. Next the alkyl halide is reacted with azide to produce the aryl-(alkyl)-azide. Last the azide is reduced to the corresponding amine by hydrogen/catalyst to give the C-terminal amine (VI) of formula $\text{NH}_2\text{-CH}_2\text{-Rc-aryl}$. The suitably functionalized C-terminal amines (VI) may readily be prepared by one skilled in the art via known methods in the literature, making non-significant modifications. Select literature references include 1) Calderwood, et al., *Tet. Lett.*, 1997, 38, 1241, 2) Ciganek, J. *Org. Chem.*, 1992, 57, 4521, 3) Thurkauf, et al., *J. Med. Chem.*, 1990, 33, 1452, 4) Werner, et al., *Org. Syn., Coll. Vol. 5*, 273, 5) *J. Med. Chem.*, 1999, 42, 4193, 6) *Chem. Rev.* 1995, 95, 2457, 7) *J. Am. Chem. Soc.*, 1986, 3150, 8) Felman et al., *J. Med. Chem.*, 1992, 35, 1183, 9) *J. Am. Chem. Soc.* 1970, 92, 3700, 10) *J. Med. Chem.*, 1997, 40, 2323.

CHART B discloses an alternative process for production of the enantiomerically pure (S,R)-protected alcohol (VII) from the (S)-protected compound (III). In the alternative process, the (S)-protected compound (III) is first reacted with the appropriately substituted C-terminal amine Rc-NH_2 (VI) using the preferred conditions described above to produce the corresponding (S)-protected ketone (XI) which is then reduced using the preferred conditions described above to produce the corresponding (S,R)-protected alcohol (VII).

CHART C discloses another alternative process for production of enantiomerically pure (S,R)-protected alcohol

(VII) but this time from the epoxide (V). In the process of CHART C, the epoxide (V) is reacted with azide to produce the corresponding enantiomerically pure (S,R)-protected azide (XII). Conditions to conduct the azide mediated epoxide opening are known to those skilled in the art, see for example, J. March, *Advanced Organic Chemistry*, 3rd Edition, John Wiley & Sons Publishers, 1985, p. 380. Next, the (S,R)-protected azide (XII) is reduced to the corresponding protected amine (XIII) by methods known to those skilled in the art. Preferred reducing conditions to reduce the (S,R)-protected azide (XII) in the presence of a t-butoxycarbonyl N-protecting group include catalytic hydrogenation, the conditions for which are known to those skilled in the art. Alternative reducing conditions which may be used to avoid N-deprotection with protecting groups other than t-butoxycarbonyl are known to those skilled in the art, see for example, R.C. Larock in *Comprehensive Organic Transformations*, VCH Publishers, 1989, p. 409.

The (S,R)-protected alcohol (VII) is deprotected to the corresponding (S,R)-amine (VIII) by means known to those skilled in the art for removal of amine protecting group. Suitable means for removal of the amine protecting group depends on the nature of the protecting group. Those skilled in the art, knowing the nature of a specific protecting group, know which reagent is preferable for its removal. For example, it is preferred to remove the preferred protecting group, BOC, by dissolving the (S,R)-protected alcohol (VII) in a trifluoroacetic acid/dichloromethane mixture. When complete, the solvents are removed under reduced pressure to give the corresponding (S,R)-amine (as the corresponding salt, i.e. trifluoroacetic acid salt) which is used without further purification. However, if desired, the (S,R)-amine can be purified further by means well known to those skilled in the art, such as for example, recrystallization. Further, if the

non-salt form is desired that also can be obtained by means known to those skilled in the art, such as for example, preparing the free base amine via treatment of the salt with mild basic conditions. Additional BOC deprotection conditions and deprotection conditions for other protecting groups can be found in T.W. Green and P.G.M. Wuts in "Protective Groups in Organic Chemistry," John Wiley and Sons, 1991, p. 309. Typical chemically suitable salts include trifluoroacetate, and the anion of mineral acids such as chloride, sulfate, phosphate; preferred is trifluoroacetate and chloride.

The (S,R)-amine (VIII) is then reacted with an appropriately substituted amide forming agent (IX) such as anhydride, acyl halide, or acid of the formula $(R_N-1-X_N)_2O$ or $R_N-1-X_N-X_2$ or R_N-1-X_N-OH (IX) by nitrogen-acylation means known to those skilled in the art to produce the corresponding (S,R)-substituted amine (X). Nitrogen acylation conditions for reaction of the (S,R)-amine (VIII) with an amide forming agent (IX) to produce the corresponding (S,R)-substituted amine (X) are known to those skilled in the art and can be found in R.C. Larock in Comprehensive Organic Transformations, VCH Publishers, 1989, p. 981, 979, and 972. R_N is as defined above. Substituted amine (X) is then converted to prodrug compound (Y) of the invention as shown in CHART A and as discussed above (see also examples 1260-1275 for illustrative but non-limiting examples of the conversion of compound (X) to various compounds (Y) of the invention.

In the compounds of the invention, R_s is preferably $-(CR_{P-y})_{0-4}-M$, $-O-M$, or $-NR_{P-2}-M$ where M is H, C_1-C_{12} alkyl, C_2-C_{12} alkenyl, R_P -aryl, R_P -heteroaryl or R_P -heterocycle, wherein 1-4 of the of the $-CH_2$ radicals of the alkyl or alkenyl group is optionally replaced by a heteroatom selected from O, S, $S(O)$, $S(O)_2$, or NR_{P-2} , and wherein R_{P-x} , R_{P-y} , R_{P-2} , R_P -aryl, R_P -heteroaryl, and R_P -heterocycle are as defined above.

In the compounds of the invention, it is preferred that R_N is selected from the group consisting of:

5 $R_{N-1}-X_N-$ where X_N is $-CO-$, where R_{N-1} is R_{N-aryl} or $R_{N-heteroaryl}$ where R_{N-aryl} is phenyl where the substitution on phenyl is 1,3-, and where R_{N-aryl} or $R_{N-heteroaryl}$ are substituted with one $-CO-NR_{N-2}R_{N-3}$,

10 $R_{N-1}-X_N-$ where X_N is $-CO-$, where R_{N-1} is R_{N-aryl} or $R_{N-heteroaryl}$ where R_{N-aryl} is phenyl substituted with one C_1 alkyl where the substitution on the phenyl is 1,3,5-, and where R_{N-aryl} or $R_{N-heteroaryl}$ are substituted with one $-CO-NR_{N-2}R_{N-3}$,

$R_{N-1}-X_N-$ where X_N is $-CO-$, where R_{N-1} is $R_{N-heteroaryl}$ where $R_{N-heteroaryl}$ is substituted with one $-CO-NR_{N-2}R_{N-3}$. It is further preferred that R_{N-2} and R_{N-3} are the same and are C_3 alkyl. It is further preferred that:

15 $R_{N-1}-X_N-$ where X_N is $-CO-$, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one $-CO-NR_{N-2}R_{N-3}$ where the substitution on phenyl is 1,3-,

20 $R_{N-1}-X_N-$ where X_N is $-CO-$, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one C_1 alkyl and with one $-CO-NR_{N-2}R_{N-3}$ where the substitution on the phenyl is 1,3,5-. It is preferred that X_N is (A) $-CO-$ and (B) $-SO_2-$; it is more preferred that X_N be $-CO-$. X_2 includes $-Cl$, $-Br$; it is preferred that X_2 is $-Cl$.

25 The nitrogen-acylation of primary amines to produce secondary amides is one of the oldest known reactions. The amide forming agents, $(R_{N-1}-X_N)_2O$ or $R_{N-1}-X_N-X_2$ or $R_{N-1}-X_N-OH$ (IX) are known to those skilled in the art and are commercially available or can be readily prepared from known starting materials by methods known in the literature. It is preferred to use an isophthalic acid acylating agent (IX) of the formula $R_{N-2}R_{N-3}N-CO-phenyl-CO-$ or a methylisophthalic acid acylating agent (IX) of the formula $R_{N-2}R_{N-3}N-CO-(CH_3-)phenyl-CO-$ where the substitution is 5-methyl-1,3-isophthalic acid. The more

preferred 5-methyl-1,3-isophthalic acid is 3-[(N,N-dipropylamino)carbonyl]-5-methylbenzoic acid (IX). These compounds are preferably prepared as follows. An ester, preferably the monomethyl ester of isophthalic acid or methyl 5-methyl-1,3-isophthalate is dissolved in a THF/DMF mixture. 1,1'-Carbonyldiimidazole is added at 20-25 degrees C. Next the desired amine ($H-NR_{N-2}R_{N-3}$) is added. After 3-24 hr of stirring at 20 degrees C to the reflux temperature of the solvent, the reaction mixture is partitioned between saturated aqueous ammonium chloride and a water immiscible organic solvent such as ethyl acetate. The aqueous layer is separated and extracted twice more with the organic solvent (ethyl acetate). The organic extracts are combined and then washed with saturated aqueous solutions of bicarbonate and saline and dried over anhydrous sodium sulfate or magnesium sulfate. Filtration of the drying agent and removal of solvents by reduced pressure gives the methyl ester of the desired $R_{N-2}R_{N-3}N-CO-phenyl-CO-O-CH_3$ or a methylisophthalic acid acylating agent (IX) $R_{N-2}R_{N-3}N-CO-(CH_3-phenyl-CO-O-CH_3)$. Purification of the (methyl) ester can be achieved via chromatography on silica gel eluting with ethyl acetate in hexanes. The isophthalate ester or methylisophthalate ester of the mono-alkyl or di-alkyl amide is then treated with an aqueous solution of base such as lithium hydroxide in a minimum amount of THF/methanol/water and stirred 3-24 hours at 20 degrees C to the reflux temperature of the solvent. The solvents are then removed under reduced pressure and subsequently partitioned between water and a water immiscible solvent such as ethyl acetate, for example. If emulsions prohibit separation of the two phases, a small amount of saline is added to aid in separation. The aqueous phase is separated and extracted once more with a water immiscible solvent such as ethyl acetate, for example. The aqueous phase is then acidified with concentrated acid, preferably

hydrochloric until $\text{pH} \leq 3$. The mixture obtained is then extracted three times with a water-immiscible solvent such as ethyl acetate, for example. These combined organic extracts are dried over anhydrous sodium or magnesium sulfate. The drying agent is removed by filtration and the organic solvent is removed under reduced pressure to give product. The mono- or di-alkyl amide isophthalate/methylisophthalate is used as such in the next reaction with the (S,R)-amine (VIII) to produce the (S,R)-substituted amine (X).

When it is desired to produce a primary amide, R_{N-2} and R_{N-3} are both -H, the following procedure is preferred. An ester, preferably the methyl ester of isophthalate or methyl 5-methyl-1,3-isophthalate is dissolved in a THF/DMF mixture. CDI is added at 20-25 degrees C. After five to thirty minutes, ammonia gas is bubbled into the mixture through a syringe needle for 1 hr. The mixture is cooled to 0 degrees C for the duration of the hour. The reaction is left stirring under a balloon of ammonia overnight at 20-25 degrees C, after which time the reaction mixture is partitioned between saturated aqueous ammonium chloride and a water immiscible solvent such as ethyl acetate, for example. The phases are separated and the aqueous phase is extracted twice more with ethyl acetate. The organic extracts are washed with saturated aqueous solutions of bicarbonate and saline and dried over anhydrous sodium or magnesium sulfate. Filtration of the drying agent and removal of solvents under reduced pressure gives the ester of the desired isophthalic acid or the isophthalic acid acylating agent (IX). Purification of the (methyl) ester can be achieved via chromatography on silica gel eluting with isopropanol/chloroform. The isophthalate ester or methylisophthalate ester of the primary amide is then treated with an aqueous solution of base such as lithium hydroxide in a minimum amount of THF/methanol/water and stirred overnight at

20-25 degrees C after which time the solvents are removed under reduced pressure and subsequently partitioned between water and a water immiscible solvent such as ethyl acetate, for example. If emulsions prohibit separation of the two phases, a small
5 amount of saline is added to aid in separation. The aqueous phase is separated and extracted once more with a water immiscible solvent such as ethyl acetate, for example. The aqueous phase is then acidified with concentrated acid, preferably hydrochloric until $\text{pH} \leq 3$. The mixture obtained is
10 then extracted three times with ethyl acetate. These combined organic extracts are dried over anhydrous sodium or magnesium sulfate. The drying agent is removed by filtration and the organic solvent removed under reduced pressure to give product. The amide isophthalic acid is used as such in the next reaction
15 with (VIII) to produce (X).

When it is desired that the amine be cyclized to be a group such as morpholinyl, piperazinyl, piperidinyl and pyrrolidinyl, etc the following procedure is preferably followed. An ester, preferably the methyl ester of isophthalic
20 acid or methyl 5-methyl-1,3-isophthalate is dissolved in dry methylene chloride and three drops of DMF are added. The mixture is cooled to 0 degrees C and then oxalyl chloride is added. The mixture is stirred at 0 degrees C for 30 minutes to two hours after which the solvents are removed under reduced
25 pressure. The acid chloride is left under vacuum overnight. The crude acid chloride is dissolved in dry methylene and cooled to 0 degrees C before the addition of the cyclic amine and a tertiary amine base such as N-methyl piperidine, for example. The reaction mixture is stirred at 0 degrees C for 1
30 to 6 hr before the solvents are removed under reduced pressure. The residue is diluted with water and a water immiscible solvent such as ethyl acetate, for example, and the phases are separated. The aqueous phase is extracted twice more with a

water immiscible solvent such as ethyl acetate, for example, and the combined organic extracts are washed with saturated aqueous bicarbonate and dried over anhydrous sodium or magnesium sulfate. Filtration of the drying agent and removal
5 of solvents under reduced pressure gives the product cyclic amide. The cyclic amide is then treated with an aqueous base such as lithium hydroxide in a minimum amount of THF/methanol/water and stirred overnight at 20-25 degrees C, after which time the solvents are removed under reduced
10 pressure and the residue is subsequently partitioned between water and a water immiscible solvent such as ethyl acetate, for example. The aqueous phase is extracted once more with ethyl acetate. Removal of water from the aqueous phase under reduced pressure gives the desired cyclic amide product (IX).

15 When it is desired that R_{N-1} is carbocycle, for example but not limited to, cyclohexane, one may then start with a suitably functionalized dimethyl isophthalate and via methods taught in the literature (Meyers, A.I., *Org. Syn.*, 1971, 51, 103) one may reduce the six-membered ring with reducing agents such as
20 rhodium (5%) on alumina in the presence of acetic acid and methanol under a hydrogen atmosphere to afford the corresponding dimethyl cyclohexane dicarboxylate.

CHART D sets forth an alternative process for production of the (S,R)-substituted amine (X), used in CHART A to produce
25 compound (Y), from the (S,R)-protected azide (XII), which is produced from the corresponding epoxide (V) in CHART C. The amino protecting group is removed to produce the corresponding unprotected azide (XIV) by methods previously described in CHART A for the conversion of (S,R)-protected alcohol (VII) to
30 the corresponding (S,R)-amine (VIII). The (S,R)-unprotected azide (XIV) is then acylated on nitrogen to produce the corresponding (S,R)-azide (XV). Next, the azide functionality is reduced as previously discussed for the conversion of the

(S,R)-protected azide (XII) to the corresponding (S,R)-protected amine (XIII) to give the (S,R)-free amine (XVI). Last, the (S,R)-free amine (XVI) is transformed to the corresponding (S,R)-substituted amine (X) by nitrogen
 5 alkylation with a compound of the formula R_C-X_3 to give the corresponding (S,R)-substituted amine (X). X_3 is an appropriate leaving group, such as but not limited to, -Cl, -Br, -I, -O-mesylate, -O-tosylate, O-triflate, etc. X_3 may also be an aldehyde; the corresponding coupling with (XVI) via the well
 10 known reductive amination procedure gives the (S,R)-substituted amine (X).

Carbocyclic amide forming agents (IX) are also provided for by the invention. For example, the carbocyclic amide forming agents of the formula

15
 |-----|
 $R'-CH-C(R'')(R''')-CH-X_N-OH$ (IX) are readily prepared from known starting materials by methods disclosed in the literature and known to those skilled in the art, for example, *J. Med. Chem.* 1998, 41, 1581, *J. Org. Chem.* 2000, 65, 1305. It is also
 20 understood that instead of the carboxylic acid, one may readily employ an acyl halide, where the halide is preferably chloride, or a suitable group to produce a mixed anhydride; these methods are taught by CHART A. For additional guidance on the
 25 formation of carbocycles and preferably cyclopropanes, one may consult M.P. Doyle; M.A. McKerverly; T. Ye in *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds From Cyclopropanes to Ylides*, Wiley-Interscience, 1998, pp. 163-279.

CHARTS E, F, G, and H disclose various methods to produce
 30 the R_N portion of the substituted amine (X) (which is converted to compounds (Y) of the invention, as shown in CHART A) where the phenyl ring of the R_N 1,3-disubstituted moiety,

-CO-phenyl-CO-, is further substituted in the 5-position with various groups such as amides, nitriles, halides, and amines. These compounds are prepared by methods known to those skilled in the art. The process chemistry of each reaction is known to those skilled in the art. What is novel here is the order of each process step and/or the specific reactants used. One skilled in the art knowing the desired product would know at least one method to prepare the desired product by using known starting materials. Hence, the following discussion is not necessary but is set forth to further aid those interested in preparing the compounds of the invention.

CHART E discloses alternate processes for the transformation of the aniline (XVII) or acid ester (XVIII) to the corresponding acid (IX-XXIII). One process begins with the commercially available aniline (XVII) where R_{N-a} is preferably -H, C_1-C_4 alkyl or benzyl. The aniline (XVII) is treated with a diazotizing reagent such as sodium or potassium nitrite in mineral acid, followed by a halogen source such as copper (II) halide or alkali metal halide, or by an organic diazotizing reagent such as an alkyl nitrite in a strong acid such as acetic acid or trifluoroacetic acid, followed by a halide source such as copper (II) halide or alkali metal halide to give the halo acid ester (XIX) where R_{N-b} is -Cl, -Br or -I. Alternatively, the acid ester (XVIII) is treated with N-halosuccinimide and trifluoromethanesulfonic acid to give the halo acid ester (XIX). The halo acid ester (XIX) is then converted to the ester amide (XXI) using a primary or secondary amine of the formula $H-NR_{N\alpha}R_{N\beta}$ (AMINE) where $R_{N\alpha}$ and $R_{N\beta}$ are the same or different or can be cyclized. These groups, $R_{N\alpha}$ and $R_{N\beta}$, become part of the substituted amine (X) and are included in the definition of R_N . R_N includes $R_{N-1}-X_N-$ where the linker, $-X_N-$, includes -CO- and R_{N-1} includes R_{N-aryl} . R_{N-aryl}

is defined to include phenyl (-phenyl) optionally substituted with two amides:

-CO-NR_{N-2}R_{N-3} and -CO-R_{N-4}. R_{Nalpha} and R_{Nbeta} include both the non-cyclic amides, -CO-NR_{N-2}R_{N-3} and the cyclic amides -CO-R_{N-4} where R_{N-2}, R_{N-3} and R_{N-4} are as defined in the claims. Alternatively, the halo acid ester (XIX) is converted to the dihalo ester (XX) by methods known to those skilled in the art. R_{N-c} includes -Cl and -F. The dihalo ester (XX) is treated with a primary or secondary amine of the formula H-NR_{Nalpha}R_{Nbeta} (AMINE) to give the ester amide (XXI). The ester amide (XXI) is then reacted with an AMINE in a carbon monoxide atmosphere in the presence of a palladium catalyst using methods such as those reviewed by Heck, (Palladium Reagents in Organic Synthesis, 1985 pp. 342-365). to give the diamide (XXII). Hydrolysis of the ester portion of the diamide (XXII) using methods well known to those skilled in the art gives the diamide acid (XXIII).

In CHART F, an alternate route to intermediate diamide (XXII) is shown starting from commercially available phenol (XXIV). The phenol (XXIV) is treated with a trifluoromethanesulfonating reagent such as trifluoromethanesulfonic anhydride to give triflate (XXV). The triflate (XXV) is reacted under the conditions of palladium catalysis in the presence of carbon monoxide and an amine of the formula H-NR_{Nalpha}R_{Nbeta} (AMINE) as for the conversion of the ester amide (XXI) to the corresponding diamide (XXII) in CHART E to give the diester (XXVI). The diester (XXVI) is hydrolyzed using methods known to those skilled in the art to give the monoacid (XXVII). The monoacid (XXVII) is then converted to the diamide (XXII) using conditions such as for the conversion of the halo acid ester (XIX) to the ester amide (XXI) in CHART E.

CHART G discloses another route to prepare the ester amide (XXI). The reaction starts with commercially available nitro compound (XXVIII) which is condensed with an (AMINE) using coupling methods known to those skilled in the art to give the nitro amide (XXX). The nitro amide (XXX) can also be prepared by first treating the nitro compound (XXVIII) with reagents such as thionyl chloride, or DMF and oxalyl chloride, or other methods known to those skilled in the art to give the acid chloride (XXIX), which upon treatment with the (AMINE) gives the nitro amide (XXX). Reduction of the nitro amide (XXX) using methods known to those skilled in the art (see, for example, Smith and March, Advanced Organic Chemistry, 5th ed.) gives amide aniline (XXXI). The amide aniline (XXXI) is then treated with diazotizing reagents such as sodium or potassium nitrite in mineral acid, followed by a halogen source such as copper (II) halide or alkali metal halide, or by an organic diazotizing reagent such as an alkyl nitrite in a strong acid such as acetic acid or trifluoroacetic acid, followed by a halide source such as copper (II) halide or alkali metal halide to give the ester amide (XXI).

CHART H discloses a process to prepare the diamide acid (IX-XXIII) from the ester amide (XXI), where one of the amides is unsubstituted and is $-CO-NH_2$. This process starts from either the ester or the acid, for example the ester amide (XXI) is treated with copper (I) cyanide ($CuCN$) in N-methylpyrrolidinone or DMF, preferably N-methylpyrrolidinone, to give the nitrile (XXXII). The nitrile (XXXII) is converted to the primary amide (XXXIII) using urea-hydrogen peroxide complex (see *Synth. Commun.* (1993) 3149) or the methods of *Synth. Commun.* (1990) 1445, *Synth. Commun.* (1997) 3119, *J. Org. Chem.* (1992) 2521, *Tet. Lett.* (1996) 6555, *Ind. J. Chem., Sect. B*, (1999) 974, *Tet. Lett.* (1995) 3469, *Tet. Lett.* (1998) 3005, or others. When the ester amide (XXI) is in the form of an

ester, an additional hydrolysis step using lithium hydroxide, sodium hydroxide, potassium hydroxide, barium hydroxide, or other hydrolysis methods known to those skilled in the art is used to convert the diamide ester (XXXIII) to the diamide acid
5 (IX-XXIII).

CHART I discloses an alternate synthetic route from the protected alcohol (VII) to the substituted amine (X) (used to form compounds (Y)) which uses a diprotected intermediate (XXXIV) whereby the nitrogen atom attached to the R_C
10 substituent is protected. Using the process of CHART I, the mono protected alcohol (VII) is reacted with a new protecting group to form the orthogonally protected (XXXIV). This is a common strategy employed in traditional peptide chemistry by those skilled in the art, see M. Bodansky, Principles of
15 Peptide Chemistry. When the mono protected alcohol (VII) is protected with CBZ one skilled in the art could react it with either (BOC)₂O in methylene chloride or similar organic solvent or FMOC-Cl in methylene chloride or similar organic solvent to prepare orthogonally protected (XXXIV). Then the CBZ group is
20 removed by hydrogenation in the presence of a catalytic amount of palladium on carbon in an alcoholic solvent, such as methanol, or ethyl acetate, or with catalytic palladium on carbon in alcoholic solvents in the presence of ammonium formate as is known to those skilled in the art. This gives
25 the R_C-N protected (XXXV). Similarly, when the mono protected alcohol (VII) is protected as a BOC it can be reacted with CBZ-Cl under Schotten-Bauman conditions or CBZ-OSu in THF to prepare the reversed (XXXIV). Then the BOC group can be
30 cleaved with hydrochloric acid (4 N) in methanol, ethanol or dioxane or with trifluoroacetic acid in methylene chloride or by other methods such as those described in The Peptides, Analysis, Synthesis, Biology, Vol. 3, Ed. E. Gross and J. Meienhofer (1981) to liberate the CBZ R_C-N protected (XXXV).

This functional group manipulation gives various permutations in the sequence (VII) to (XXXIV) to (XXXV) as is apparent to one skilled in the art. When the appropriately R_C -N protected compound (XXXV) is reacted with the amide forming agent (IX),
5 in acid form, under standard peptide coupling conditions, for example, EDC/HOBt in methylene chloride or DMF or a previously activated acid, $(R_N)_2O$ gives the corresponding R_N -substituted R_C -N protected (XXXVI). Simple de-protection of the R_N -substituted R_C -N protected (XXXVI) then gives the desired
10 substituted amine (X). Thus when the R_N -substituted R_C -N protected (XXXVI) is protected with BOC, treatment with hydrochloric acid (4N) in dioxane or the other reagents discussed above gives the substituted amine (X). When the R_N -substituted R_C -N protected (XXXVI) is protected with CBZ,
15 treatment with hydrogen from 10 - 50 psi in alcoholic solvents, such as methanol with a catalytic amount of palladium on carbon will give, after work-up, the desired substituted amine (X). Similarly when the R_N -substituted R_C -N protected (XXXVI) is protected with FMOC, treatment with a secondary amine,
20 preferably either piperidine (10 %) or diethylamine (10 %) in an inert solvent such as, for example, methylene chloride will give after work up the desired substituted amine (X).

CHART J discloses a process to prepare compounds where the phenyl ring of the R_N substituent of -CO-phenyl-CO- is
25 substituted with a sulfonamide group in the 5-position. The process starts with the halo amide ester (XXI, CHART E) which is reacted with sodium nitrite, sulfur dioxide, copper chloride (II) and acetic acid by the method disclosed in *J. Med. Chem.*, 42, 3797 (1999) to prepare the sulfonyl chloride (XXXVII). The
30 sulfonyl chloride (XXXVII) is then reacted with AMINE, as defined above, by methods known to those skilled in the art to produce the corresponding sulfonamide (XXXVIII). Last the sulfonamide (XXXVIII) is transformed to the corresponding

sulfonamide acid (XXXIX) by methods known to those skilled in the art such as using lithium hydroxide, sodium hydroxide, potassium hydroxide, barium hydroxide, or other hydrolysis methods known to those skilled in the art.

5 CHART K discloses how to prepare the R_N substituents where R_N is $R_{N-1}-X_N-$, where X_N is $-\text{CO}-$ and R_{N-1} is $R_{N-\text{aryl}}$ where $R_{N-\text{aryl}}$ is phenyl substituted with one alkyl group and one $-\text{CO}-\text{NR}_{N-2}\text{R}_{N-3}$ or $-\text{CO}-\text{R}_{N-4}$. See the discussion above for CHART E regarding the amine, $\text{H}-\text{NR}_{N\alpha}\text{R}_{N\beta}$ (AMINE), used to form the amide R_N substituents. The process starts with the halo amide ester (XXI) which is then reacted with an alkyl boronic acid having the desired alkyl group in the presence of a palladium catalyst such as $\text{Pd}(\text{PPh}_3)\text{Cl}_2$ using the general method described in *J. Med. Chem.*, 4288 (2000). The alkyl boronic acids are commercially available or can be prepared by the process described in *J. Am. Chem. Soc.*, 60, 105 (1938). It is preferred that R_{N-b} is bromo. This step produces the alkyl ester (XL) which is then hydrolyzed by means known to those skilled in the art to produce the desired alkyl acid (XLI).

20 CHART L discloses a process to prepare the amide forming agent (IX - XLVII) where the R_N substituent is $R_{N-1}-X_N-$, where the linker, $-X_N-$ is $-\text{CO}-$, where R_{N-1} is $R_{N-\text{aryl}}$ and where $R_{N-\text{aryl}}$ is phenyl (-phenyl) substituted with groups:

C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, and $-\text{N}(-\text{H}$ and C_1-C_3 alkyl)- $\text{CO}-\text{R}_{N-5}$. This specific amide forming agent, (IX - XLVII) is prepared by starting with the phenyl nitro compound (XLII) which is reduced to the corresponding phenyl nitro hydroxy compound (XLIII) using borane-methyl sulfide or borane in THF. The phenyl nitro hydroxy compound (XLIII) is reduced to the corresponding phenyl amino hydroxy compound (XLIV) using

hydrogen and palladium catalyst as is known to those skilled in the art. The phenyl amino hydroxy compound (XLIV) is reacted with an aldehyde in the presence of a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride to give the phenyl substituted amino hydroxy compound (XLV). The phenyl substituted amino hydroxy compound (XLV) is acylated with an acid chloride or acid anhydride by methods known to those skilled in the art to give the phenyl disubstituted amino hydroxy compound (XLVI). The phenyl disubstituted amino hydroxy compound (XLVI) is hydrolyzed using an alkali hydroxide, followed by acidification, to give the amide forming agent (IX - XLVII). The amide forming agent (XLVII) is then coupled with amine (VIII) using methods known to those skilled in the art and methods previously discussed, such as with diethyl cyanophosphonate, to give the substituted amine (X). Further treatment of the substituted amine (X) with diethyl cyanophosphonate gives the substituted amine where the hydroxyalkyl substituent on the phenyl ring has a phosphate substituent.

CHART M discloses a process to prepare amide forming agents (IX- L) where the R_N substituent is $R_{N-1}-X_N-$, where the linker, $-X_N-$ is $-CO-$, where R_{N-1} is R_{N-aryl} and where R_{N-aryl} is phenyl (-phenyl) substituted with two groups. The first substituent at what is usually identified as position "5-" can be either:

$-R_{N-aryl}$ or

$-R_{N-heteroaryl}$. The second substituent at what is usually identified as position "3-" can be either:

$-CO-NR_{N-2}R_{N-3}$ or

$-CO-R_{N-4}$.

$R_{N\alpha}$ and $R_{N\beta}$ include both the non-cyclic amides, $-CO-NR_{N-2}R_{N-3}$ and the cyclic amides $-CO-R_{N-4}$ where R_{N-2} , R_{N-3} and R_{N-4} are as defined in the claims. The process starts with the

trisubstituted phenyl compound (XLVIII) where R_{N-d} is -Cl, -Br, -I or -O-triflate. Treatment with an aryl or heteroaryl boronic acid or heteroaryl or aryl boronic acid ester such as (aryl or heteroaryl)-B(OH)₂ or (aryl or heteroaryl)-B(OR^a)(OR^b) (where R^a and R^b are lower alkyl, ie. C₁-C₆, or taken together, R^a and R^b are lower alkylene, ie. C₂-C₁₂) in the presence of a metal catalyst with or without a base in an inert solvent yields (XLIX). Metal catalysts in these transformations include, but are not limited to, salts or phosphine complexes of Cu, Pd, or Ni (eg. Cu(OAc)₂, PdCl₂(PPh₃)₂, NiCl₂(PPh₃)₂). Bases may include, but are not limited to, alkaline earth metal carbonates, alkaline earth metal bicarbonates, alkaline earth metal hydroxides, alkali metal carbonates, alkali metal bicarbonates, alkali metal hydroxides, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium diisopropylamide), alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably diisopropylethylamine or triethylamine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, acetonitrile, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylacetamides (preferably dimethylacetamide), N,N-dialkylformamides (preferably dimethylformamide), dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes (preferably methylene chloride). Preferred reaction temperatures range from room temperature up to the boiling point of the solvent employed. The reactions may be run in conventional glassware or in one of many commercially available parallel synthesizer units. Non-commercially available

boronic acids or boronic acid esters may be obtained from the corresponding optionally substituted aryl halide as described in *Tetrahedron*, 50, 979-988 (1994). Intermediate (XLIX) is then hydrolyzed using alkali metal hydroxide, for example lithium,
 5 sodium or potassium hydroxide, followed by acidification, to give aryl or heteroaryl coupled acids (IX-L). Alternatively, as described in *Tetrahedron*, 50, 979-988 (1994), one may convert the R_N -d to the corresponding boronic acid or boronic acid ester $(OH)_2B$ - or $(OR^a)(OR^b)B$ - and obtain the same products set forth
 10 above by treating with a suitable aryl or heteroaryl halide or triflate.

CHART N discloses a process to prepare amide forming agents (IX - LII) where the R_N substituent is $R_{N-1}-X_N$ - where the linker, $-X_N$ - is $-CO-$, where R_{N-1} is R_{N-aryl} and where R_{N-aryl} is phenyl ($-phenyl$) substituted with two groups. The first substituent at
 15 what is usually identified as position "5-" is $-C\equiv C-R$. The second substituent at what is usually identified as position "3-" can be either $-CO-NR_{N-2}R_{N-3}$ or $-CO-R_{N-4}$. The halo ester (XXI) is treated with a mixture of $PdCl_2(Pphenyl)_2$ and trimethylsilyl
 20 acetylene, using methods known to those skilled in the art, to give acetylene ester (LI). Acetylene ester (LI) is then hydrolyzed using alkali metal hydroxide, followed by acidification, to give acetylene acid (IX - LII).

CHARTS O and O' disclose processes to prepare amide forming
 25 agents (IX - LX) and (IX - LXIII) with an extended methylene group where the R_N substituent is $R_{N-1}-X_N$ - where the linker, $-X_N$ - is $-CO-$, where R_{N-1} is R_{N-aryl} and where R_{N-aryl} is phenyl ($-phenyl$) substituted with two groups. The substituent at what is usually identified as position "3-" can be either $-CO-NR_{N-2}R_{N-3}$ or $-CO-R_{N-4}$.
 30 4. In the process of CHART O, the substituent at the 5-position is $-CH_2CO-NH_2$ and in the process of CHART O', the substituent at the 5-position is $-CH_2C\equiv N$. The starting diester acid (LIII)

is reduced with borane in solvents such as THF to give the corresponding diester alcohol (LIV). The diester alcohol (LIV) is converted to the corresponding diester bromo compound (LV) using a brominating agent such as PBr_3 , CBr_4 , or other halogenating agent such as are known to those skilled in the art. The bromine of the diester bromo compound (LV) is then displaced with cyanide to give the corresponding nitrile (LVI). In CHART O', the nitrile (LVI) is then hydrolyzed to the corresponding cyano ester (LXI). The cyano ester (LXI) is then coupled with $H-NR_{N-1}R_{N-2}$ (AMINE), as previously described using methods known to those skilled in the art to give the corresponding cyano amide (LXII). The cyano amide (LXII) is then hydrolyzed to the corresponding cyano acid (IX-LXIII) which is in turn coupled with amine (VIII) to give the substituted amine (X). Amine (X) is converted to compound (Y) as shown in CHART A. When the substituent on the extended methyl group is $-CO-NH_2$, the process of CHART O is used. There the nitrile (LVI) is converted to the corresponding diester amine (LVII) by methods known to those skilled in the art. The next steps are the same as for CHART O' where the diester amide (LVII) is hydrolyzed to the corresponding ester amine (LVIII) which is then converted to the corresponding diamide ester (LIX) which is hydrolyzed to the corresponding diamide acid (IX - LX). The diamide acid (IX - XL) is then coupled with the appropriate amine (VIII) to produce the desired substituted amine (X). Amine (X) is readily converted to compounds (Y) as shown in CHART A.

CHART P discloses a process to prepare amide forming agents (IX - LXVII) with an extended hydroxymethylene group where the R_N substituent is $R_{N-1}-X_N-$ where the linker, $-X_N-$ is $-CO-$, where the R_{N-1} is R_{N-aryl} , where R_{N-aryl} is phenyl (-phenyl) substituted with two groups. The substituent at what is usually identified as position "3-" can be either $-CO-NR_{N-2}R_{N-3}$ or $-CO-R_{N-4}$. The

process begins with a halo amide (LXIV), preferably iodo, which is converted to the corresponding aldehyde (LXV) and then to the corresponding alcohol (LXVI) by the method described in *Synth. Commun.* 28, 4270 (1998), optionally with variations known to those skilled in the art. Hydrolysis of the alcohol (LXVI) using alkali hydroxides, followed by acidification, gives the desired hydroxy acid (IX - LXVII). The hydroxy acid (IX - LXVII) is then coupled with the appropriate amine (VIII) to give the desired substituted amine (X) which is converted to compounds (Y) as shown in CHART A.

CHART Q discloses a process to prepare amide forming agents (IX - LXXII) with an alkyl group or a halogen atom or an amino group at the 5-position where the R_N substituent is $R_{N-1}-X_N$ where the linker, $-X_N-$ is $-CO-$, where the R_{N-1} is R_{N-aryl} , where R_{N-aryl} is phenyl (-phenyl) substituted with two groups. The substituent at what is usually identified as position "3-" can be either $-CO-NR_{N-2}R_{N-3}$ or $-CO-R_{N-4}$. The process begins with an appropriately 5-substituted diacid (LXVIII) which is esterified by methods known to those skilled in the art to give the corresponding diester (LXIX). The diester (LXIX) is then hydrolyzed using alkali hydroxides, followed by acidification, to give the corresponding monoacid (LXX). Alternatively, the monoacid (LXX) can be produced directly from the diacid (LXVIII) by known methods. The monoacid (LXX) is then coupled with $H-NR_{N\alpha}R_{N\beta}$ (AMINE) to give the corresponding amide ester (LXXI). The amide ester (LXXI) is then hydrolyzed using alkali hydroxides, followed by acidification, to give the corresponding acid amide (IX - LXXII).

CHART R discloses a general process to prepare the amide forming agents (IX - LXXVII) which, for example, have an alkyl group at what is known as the 5-position and a ketone at the 3-position. These acids (IX- LXXVII) are formed by starting with the acid (LXXIII) which is converted to the corresponding acid

halide (LXXIV) using methods known to those skilled in the art. The acid halide (LXXIV) is preferably the acid chloride. The acid halide (LXXIV) in the presence of copper (I) bromide and tetrahydrofuran and at temperatures ranging from -78 degrees C to 0 degreesC is treated with a Grignard reagent (aryl-Mg-X, or alkyl-Mg-X, where X is -Cl or -Br) to give the ketone esters (LXXVI and LXXVI'). Many Grignard reagents are available for purchase; others are prepared by methods known to those skilled in the art. An alternative method for preparing the ketone esters (LXXVI, LXXVI') is to prepare the Weinreb amide (LXXV), either from the acid (LXXIII) directly or by way of acid halide (LXXIV) followed by treatment with N,O-dimethylhydroxylamine to give Weinreb amide (LXXV) and then treating the Weinreb amide (LXXV) with a Grignard reagent, by methods known to those skilled in the art. The ketone esters (LXXVI, LXXVI') are then hydrolyzed using alkali hydroxides, followed by acidification, to give the ketone acids (LXXVII, LXXVII').

CHART S discloses various methods to modify the R_N portion of the substituted amine (X) where the phenyl ring of the R_N moiety is further substituted in the 3-position with various groups such as aryl and heteroaryl. These compounds are prepared by methods known to those skilled in the art. The process chemistry of each reaction is known to those skilled in the art. What is novel here is the order of each process step and/or the specific reactants used. One skilled in the art knowing the desired product would know at least one method to prepare the desired product by using known starting materials. Hence, the following discussion is not necessary but is set forth to further aid those interested in preparing the compounds of the invention.

CHART S sets forth a general method used in the present invention to prepare the substituted amines (X) where $R_N = R_N$.

aryl- R_N -aryl- X_N or R_N -heteroaryl- R_N -aryl- X_N . Amines (X) are converted to compounds (Y) of the invention as shown in CHART A.

Treatment of the (S,R)-amine (VIII) with amide forming agents (IX) according to the methods set forth above where for
 5 CHART S, R_N is Br- R_N -aryl generates the corresponding (S,R)-substituted amine (X) where R_N is Br- N_R -aryl- X_N . Further treatment with an aryl boronic acid or aryl boronic acid ester such as (aryl or heteroaryl)-B(OH)₂ or (aryl or heteroaryl)-B(OR^a)(OR^b) (where R^a and R^b are lower alkyl, ie. C₁-C₆, or
 10 taken together, R^a and R^b are lower alkylene, ie. C₂-C₁₂) in the presence of a metal catalyst with or without a base in an inert solvent yields the (S,R)-substituted amine (X) where R_N is N_R -aryl- N_R -aryl- X_N or R_N -heteroaryl- R_N -aryl- X_N . Metal catalysts in these transformations include, but are not limited to, salts or
 15 phosphine complexes of Cu, Pd, or Ni (eg. Cu(OAc)₂, PdCl₂(PPh₃)₂, NiCl₂(PPh₃)₂). Bases may include, but are not limited to, alkaline earth metal carbonates, alkaline earth metal bicarbonates, alkaline earth metal hydroxides, alkali metal carbonates, alkali metal bicarbonates, alkali metal hydroxides, alkali metal hydrides (preferably sodium hydride),
 20 alkali metal alkoxides (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium diisopropylamide), alkali metal bis(trialkylsilyl)amides (preferably sodium
 25 bis(trimethylsilyl)amide), trialkyl amines (preferably diisopropylethylamine or triethylamine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, acetonitrile, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-
 30 dioxane), N,N-dialkylacetamides (preferably dimethylacetamide), N,N-dialkylformamides (preferably dimethylformamide),



dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes (preferably methylene chloride). Preferred reaction temperatures range from room temperature up to the boiling point of the solvent employed. The reactions may be run in conventional glassware or in one of many commercially available parallel synthesizer units. Non-commercially available boronic acids or boronic acid esters may be obtained from the corresponding optionally substituted aryl halide as described in *Tetrahedron*, 50, 979-988 (1994).

Where the above chemistry is incompatible with other functionality present in the (S,R)-substituted amine (X) where R_N is $Br-NR-aryl-X_N$, then one skilled in the art will readily understand that an alternative sequence of coupling steps is required. For example, treatment of an appropriately substituted amide forming agent (IX) $R_{N-1}-X_N-OH$ where R_{N-1} is $Br-R_N-aryl$ with a boronic acid or boronic acid ester under the conditions described above will afford the appropriately substituted amide forming agent (IX) where R_{N-1} is $NR-aryl-NR-aryl$ or $R_N-heteroaryl-R_N-aryl$. When the amide forming agent (IX) where R_{N-1} is $NR-aryl-NR-aryl$ or $R_N-heteroaryl-R_N-aryl$ is treated with the (S,R)-amine (VIII), one then obtains the same substituted amines (X) set forth in CHART S.

The above examples for CHART S are not meant to limit the scope of the chemistry. In addition to bromine, a suitable group may include iodine or triflate. Alternatively, as described in *Tetrahedron*, 50, 979-988 (1994), one may convert the $Br-R_N-aryl$ to the corresponding boronic acid or boronic acid ester $(OH)_2B-R_N-aryl$ or $(OR^a)(OR^b)B-R_N-aryl$ and obtain the same products set forth above by treating with a suitable aryl or heteroaryl halide or triflate. Additionally, each $-R_N-aryl$ and

-R_N-heteroaryl are interchangeable at each occurrence in the chemistry described above.

CHART T discloses a process to prepare amide forming agents (IX - LXXIX) where the R_N substituent is R_{N-1}-X_N-, where
 5 the linker, -X_N- is -CO-, where R_{N-1} is R_{N-aryl} and where R_{N-aryl} is phenyl substituted with -CO-NR_{Nalpha}R_{Nbeta} (AMINE) and with an amide of the formulas:



10 The process begins with the amide aniline (XXXI) which is reacted with the corresponding acid halide or sulfonyl halide, or acid anhydride or sulfonyl anhydride to produce the corresponding amide ester (LXXVIII). Suitable solvents include THF or dichloromethane at temperatures ranging from -78 degrees
 15 to 100 degrees C. The amide ester (LXXVIII) is then hydrolyzed to the corresponding amide acid (IX - LXXIX) by methods known to those skilled in the art. When the amide forming agent (IX - LXXIX) is reacted with the appropriate amine (VIII), the substituted amine (X) is obtained. Amine (X) is converted to
 20 compounds (Y) of the invention as shown in CHART A.

CHART U discloses a general method for preparing various C-terminal amines (VI) as represented by the preparation of C-terminal amine (LXXXIV). Methods to prepare amines of this type are well understood using methods known to those skilled
 25 in the art, or one may consult the references: 1) JACS, 1970, 92, 3700, and 2) US patent 4,351,842.

CHART V further discloses general methods for preparing various C-terminal amines (VI) as represented by the preparation of C-terminal amines (LXXXIX). Multiple examples
 30 of the heterocyclic carboxylic acids or acid chlorides are commercially available. Optionally, the carboxylic acid (LXXXV) may be converted to the acid chloride (LXXXVI) with reagents such as, but not limited to, thionyl chloride.

Displacement with ammonia generates the common intermediate amides (LXXXVII) which are readily reduced to amines (VI - LXXXIX) using a variety of methods detailed previously. Alternatively, other heteroaryls are commercially available as the methyl halide (LXXXVIII) which are treated with ammonia to yield the title C-terminal amines (VI - LXXXVIII).

CHART W discloses general methods for preparing thiazolyl containing C-terminal amines as represented by the preparation of C-terminal amines (LXXXXI). The synthesis of the thiazoles is outlined in CHART W; these procedures are amply taught in the literature and are modified from the procedures outlined in: Mashraqui, SH; Keehn, PM. *J. Am. Chem. Soc.* 1982, 104, 4461-4465. The synthesis of substituted 5-aminomethylthiazoles (XCI) was achieved from 5-hydroxymethylthiazole (XC) by the procedure described in: Alterman et al. *J. Med. Chem.* 1998, 41, 3782-3792. All other thiazole analogs were transformed to the hydroxymethyl derivative using CHART W, and converted to the aminomethyl derivative by the Alterman procedure without notable changes.

CHART X discloses general methods for preparing isoxazolyl containing C-terminal amines as represented by the preparation of C-terminal amines (XCII). The synthesis of isoxazole derivatives was modified from the procedure in: Felman, SW et al. *J. Med. Chem.* 1992, 35, 1183-1190 and is readily understood by those skilled in the art making non-notable changes to achieve the title compounds. The substituted hydroxylamine precursors were synthesized using the procedure taught by Bousquet, EW. *Org. Synth. Coll. Vol II*, 313-315. Commercially available propargylamine may be protected using any number of methods known in the art (see: Greene, TW; Wuts, PGM. *Protective Groups in Organic Synthesis*, 3rd Ed. New York: John Wiley, 1999. Chapter 7.), preferred is a BOC protecting group.

Substituted propargyl amines may be obtained by a number of methods commonly known in the art.

CHART Y discloses a general route to prepare hydroxyethylamines where one carbon atom of the peptide backbone, along with R₂ and R₃ form a ring. It is understood that the present invention also allows for a heteroatom to be incorporated into the ring. In summary, the synthesis of compounds where R₂ and R₃ may form a ring proceeds from a suitably protected amino acid aldehyde and cycloalkyllithium species, both of which are commercially available or where known procedures for making such compounds are known in the art. The general procedure involved is also precedent in the literature, for example, see Klumpp, et al., *J. Am. Chem. Soc.*, 1979, 101, 7065, and it is intended that making non-critical variations, one may obtain the title compounds provided for by CHART Y. Treatment of a suitably protected amino acid aldehyde and cycloalkyllithium species affords alcohol (XCIII). These reactions are carried out in an inert solvent such as, for example, tetrahydrofuran or diethyl ether. Optimally the reactions are conducted at low temperatures, for example below 0 degrees C. Carbonylation via the Klumpp procedure yields the acid (XCIV) which when exposed to Curtius, or related procedures well known to those skilled in the art, generates the primary amine (XCV). The primary amines (XCV) may be capped C-terminally via the conditions set forth in CHART C & D followed by nitrogen deprotection and capping N-terminally via the conditions set forth in CHART A.

The compounds of the invention may contain geometric or optical isomers as well as tautomers. Thus, the invention includes all tautomers and pure geometric isomers, such as the E and Z geometric isomers, as well as mixtures thereof. Furthermore, the invention includes pure enantiomers and diastereomers as well as mixtures thereof, including racemic

mixtures. The individual geometric isomers, enantiomers, or diastereomers may be prepared or isolated by methods known in the art.

Compounds of the invention with the stereochemistry designated in formula X may be included in mixtures, including racemic mixtures, with other enantiomers, diastereomers, geometric isomers or tautomers. Compounds of the invention with the stereochemistry designated in formula X are typically present in these mixtures in excess of 50 percent. Preferably, compounds of the invention with the stereochemistry designated in formula X are present in these mixtures in excess of 80 percent. Most preferably, compounds of the invention with the stereochemistry designated in formula X are present in these mixtures in excess of 90 percent.

The compounds (Y) are amines and as such form salts when reacted with acids. Pharmaceutically acceptable salts are preferred over the corresponding compounds (Y) since they produce compounds which are more water soluble, stable and/or more crystalline. Pharmaceutically acceptable salts are any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Pharmaceutically acceptable salts include salts of both inorganic and organic acids. The preferred pharmaceutically acceptable salts include salts of the following acids: acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsyllic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycollylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic,

muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, succinic, sulfamic, sulfanilic, 5 sulfonic, sulfuric, tannic, tartaric, teoclic and toluenesulfonic. For other acceptable salts, see *Int. J. Pharm.*, 33, 201-217 (1986) and *J. Pharm. Sci.*, 66(1), 1, (1977).

The present invention provides compounds, compositions, kits, and methods for inhibiting beta-secretase enzyme activity 10 and A beta peptide production. Inhibition of beta-secretase enzyme activity halts or reduces the production of A beta from APP and reduces or eliminates the formation of beta-amyloid deposits in the brain.

15 **Methods of the Invention**

The compounds of the invention, and pharmaceutically acceptable salts thereof, are useful for treating humans or animals suffering from a condition characterized by a pathological form of beta-amyloid peptide, such as beta-amyloid 20 plaques, and for helping to prevent or delay the onset of such a condition. For example, the compounds are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with MCI (mild cognitive impairment) and preventing or delaying the 25 onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and 30 recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear

palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds of the invention can either be used individually or in combination, as is best for the patient.

As used herein, the term "treating" means that the compounds of the invention can be used in humans with at least a tentative diagnosis of disease. The compounds of the invention will delay or slow the progression of the disease thereby giving the individual a more useful life span.

The term "preventing" means that the compounds of the present invention are useful when administered to a patient who has not been diagnosed as possibly having the disease at the time of administration, but who would normally be expected to develop the disease or be at increased risk for the disease. The compounds of the invention will slow the development of disease symptoms, delay the onset of the disease, or prevent the individual from developing the disease at all. Preventing also includes administration of the compounds of the invention to those individuals thought to be predisposed to the disease due to age, familial history, genetic or chromosomal abnormalities, and/or due to the presence of one or more biological markers for the disease, such as a known genetic mutation of APP or APP cleavage products in brain tissues or fluids.

In treating or preventing the above diseases, the compounds of the invention are administered in a therapeutically effective amount. The therapeutically effective amount will vary depending on the particular compound used and the route of administration, as is known to those skilled in the art.

In treating a patient displaying any of the diagnosed above conditions a physician may administer a compound of the invention immediately and continue administration indefinitely, as needed. In treating patients who are not diagnosed as
5 having Alzheimer's disease, but who are believed to be at substantial risk for Alzheimer's disease, the physician should preferably start treatment when the patient first experiences early pre-Alzheimer's symptoms such as, memory or cognitive problems associated with aging. In addition, there are some
10 patients who may be determined to be at risk for developing Alzheimer's through the detection of a genetic marker such as APOE4 or other biological indicators that are predictive for Alzheimer's disease. In these situations, even though the patient does not have symptoms of the disease, administration
15 of the compounds of the invention may be started before symptoms appear, and treatment may be continued indefinitely to prevent or delay the outset of the disease.

Dosage forms and amounts

20 The compounds of the invention can be administered orally, parenternally, (IV, IM, depo-IM, SQ, and depo SQ), sublingually, intranasally (inhalation), intrathecally, topically, or rectally. Dosage forms known to those of skill in the art are suitable for delivery of the compounds of the
25 invention.

Compositions are provided that contain therapeutically effective amounts of the compounds of the invention. The compounds are preferably formulated into suitable pharmaceutical preparations such as tablets, capsules, or
30 elixirs for oral administration or in sterile solutions or suspensions for parenternal administration. Typically the compounds described above are formulated into pharmaceutical

compositions using techniques and procedures well known in the art.

About 1 to 500 mg of a compound or mixture of compounds of the invention or a physiologically acceptable salt or ester is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in those compositions or preparations is such that a suitable dosage in the range indicated is obtained. The compositions are preferably formulated in a unit dosage form, each dosage containing from about 2 to about 100 mg, more preferably about 10 to about 30 mg of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

To prepare compositions, one or more compounds of the invention are mixed with a suitable pharmaceutically acceptable carrier. Upon mixing or addition of the compound(s), the resulting mixture may be a solution, suspension, emulsion, or the like. Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for lessening or ameliorating at least one symptom of the disease, disorder, or condition treated and may be empirically determined.

Pharmaceutical carriers or vehicles suitable for administration of the compounds provided herein include any

such carriers known to those skilled in the art to be suitable for the particular mode of administration. In addition, the active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, or have another action. The compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients.

Where the compounds exhibit insufficient solubility, methods for solubilizing may be used. Such methods are known and include, but are not limited to, using cosolvents such as dimethylsulfoxide (DMSO), using surfactants such as Tween®, and dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as salts or prodrugs may also be used in formulating effective pharmaceutical compositions.

The concentration of the compound is effective for delivery of an amount upon administration that lessens or ameliorates at least one symptom of the disorder for which the compound is administered. Typically, the compositions are formulated for single dosage administration.

The compounds of the invention may be prepared with carriers that protect them against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, microencapsulated delivery systems. The active compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. The therapeutically effective concentration may be determined empirically by testing the compounds in known *in vitro* and *in vivo* model systems for the treated disorder.

The compounds and compositions of the invention can be enclosed in multiple or single dose containers. The enclosed

compounds and compositions can be provided in kits, for example, including component parts that can be assembled for use. For example, a compound inhibitor in lyophilized form and a suitable diluent may be provided as separated components for combination prior to use. A kit may include a compound inhibitor and a second therapeutic agent for co-administration. The inhibitor and second therapeutic agent may be provided as separate component parts. A kit may include a plurality of containers, each container holding one or more unit dose of the compound of the invention. The containers are preferably adapted for the desired mode of administration, including, but not limited to tablets, gel capsules, sustained-release capsules, and the like for oral administration; depot products, pre-filled syringes, ampules, vials, and the like for parenteral administration; and patches, medipads, creams, and the like for topical administration.

The concentration of active compound in the drug composition will depend on absorption, inactivation, and excretion rates of the active compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test data. It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the

administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

If oral administration is desired, the compound should be provided in a composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an enteric coating that maintains its integrity in the stomach and releases the active compound in the intestine. The composition may also be formulated in combination with an antacid or other such ingredient.

Oral compositions will generally include an inert diluent or an edible carrier and may be compressed into tablets or enclosed in gelatin capsules. For the purpose of oral therapeutic administration, the active compound or compounds can be incorporated with excipients and used in the form of tablets, capsules, or troches. Pharmaceutically compatible binding agents and adjuvant materials can be included as part of the composition.

The tablets, pills, capsules, troches, and the like can contain any of the following ingredients or compounds of a similar nature: a binder such as, but not limited to, gum tragacanth, acacia, corn starch, or gelatin; an excipient such as microcrystalline cellulose, starch, or lactose; a disintegrating agent such as, but not limited to, alginic acid and corn starch; a lubricant such as, but not limited to, magnesium stearate; a gildant, such as, but not limited to, colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; and a flavoring agent such as peppermint, methyl salicylate, or fruit flavoring.

When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials, which modify the physical form of the

dosage unit, for example, coatings of sugar and other enteric agents. The compounds can also be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active
5 compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings, and flavors.

The active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action.

10 Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent such as water for injection, saline solution, fixed oil, a naturally occurring vegetable oil such as sesame oil, coconut oil, peanut
15 oil, cottonseed oil, and the like, or a synthetic fatty vehicle such as ethyl oleate, and the like, polyethylene glycol, glycerine, propylene glycol, or other synthetic solvent; antimicrobial agents such as benzyl alcohol and methyl parabens; antioxidants such as ascorbic acid and sodium
20 bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates, and phosphates; and agents for the adjustment of tonicity such as sodium chloride and dextrose. Parenteral preparations can be enclosed in ampoules, disposable syringes, or multiple dose
25 vials made of glass, plastic, or other suitable material. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

Where administered intravenously, suitable carriers include physiological saline, phosphate buffered saline (PBS),
30 and solutions containing thickening and solubilizing agents such as glucose, polyethylene glycol, polypropyleneglycol, and mixtures thereof. Liposomal suspensions including tissue-targeted liposomes may also be suitable as pharmaceutically

acceptable carriers. These may be prepared according to methods known for example, as described in U.S. Patent No. 4,522,811.

The active compounds may be prepared with carriers that
5 protect the compound against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers such as collagen,
10 ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid, and the like. Methods for preparation of such formulations are known to those skilled in the art.

The compounds of the invention can be administered orally,
15 parenterally (IV, IM, depo-IM, SQ, and depo-SQ), sublingually, intranasally (inhalation), intrathecally, topically, or rectally. Dosage forms known to those skilled in the art are suitable for delivery of the compounds of the invention.

Compounds of the invention may be administered enterally
20 or parenterally. When administered orally, compounds of the invention can be administered in usual dosage forms for oral administration as is well known to those skilled in the art. These dosage forms include the usual solid unit dosage forms of tablets and capsules as well as liquid dosage forms such as
25 solutions, suspensions, and elixirs. When the solid dosage forms are used, it is preferred that they be of the sustained release type so that the compounds of the invention need to be administered only once or twice daily.

The oral dosage forms are administered to the patient 1,
30 2, 3, or 4 times daily. It is preferred that the compounds of the invention be administered either three or fewer times, more preferably once or twice daily. Hence, it is preferred that the compounds of the invention be administered in oral dosage

form. It is preferred that whatever oral dosage form is used, that it be designed so as to protect the compounds of the invention from the acidic environment of the stomach. Enteric coated tablets are well known to those skilled in the art. In addition, capsules filled with small spheres each coated to protect from the acidic stomach, are also well known to those skilled in the art.

When administered orally, an administered amount therapeutically effective to inhibit beta-secretase activity, to inhibit A beta production, to inhibit A beta deposition, or to treat or prevent AD is from about 0.1 mg/day to about 1,000 mg/day. It is preferred that the oral dosage is from about 1 mg/day to about 100 mg/day. It is more preferred that the oral dosage is from about 5 mg/day to about 50 mg/day. It is understood that while a patient may be started at one dose, that dose may be varied over time as the patient's condition changes.

Compounds of the invention may also be advantageously delivered in a nano crystal dispersion formulation. Preparation of such formulations is described, for example, in U.S. Patent 5,145,684. Nano crystalline dispersions of HIV protease inhibitors and their method of use are described in US 6,045,829. The nano crystalline formulations typically afford greater bioavailability of drug compounds.

The compounds of the invention can be administered parenterally, for example, by IV, IM, depo-IM, SC, or depo-SC. When administered parenterally, a therapeutically effective amount of about 0.5 to about 100 mg/day, preferably from about 5 to about 50 mg daily should be delivered. When a depot formulation is used for injection once a month or once every two weeks, the dose should be about 0.5 mg/day to about 50 mg/day, or a monthly dose of from about 15 mg to about 1,500 mg. In part because of the forgetfulness of the patients with

Alzheimer's disease, it is preferred that the parenteral dosage form be a depo formulation.

The compounds of the invention can be administered sublingually. When given sublingually, the compounds of the invention should be given one to four times daily in the amounts described above for IM administration.

The compounds of the invention can be administered intranasally. When given by this route, the appropriate dosage forms are a nasal spray or dry powder, as is known to those skilled in the art. The dosage of the compounds of the invention for intranasal administration is the amount described above for IM administration.

The compounds of the invention can be administered intrathecally. When given by this route the appropriate dosage form can be a parenteral dosage form as is known to those skilled in the art. The dosage of the compounds of the invention for intrathecal administration is the amount described above for IM administration.

The compounds of the invention can be administered topically. When given by this route, the appropriate dosage form is a cream, ointment, or patch. Because of the amount of the compounds of the invention to be administered, the patch is preferred. When administered topically, the dosage is from about 0.5 mg/day to about 200 mg/day. Because the amount that can be delivered by a patch is limited, two or more patches may be used. The number and size of the patch is not important, what is important is that a therapeutically effective amount of the compounds of the invention be delivered as is known to those skilled in the art. The compounds of the invention can be administered rectally by suppository as is known to those skilled in the art. When administered by suppository, the therapeutically effective amount is from about 0.5 mg to about 500 mg.

The compounds of the invention can be administered by implants as is known to those skilled in the art. When administering a compound of the invention by implant, the therapeutically effective amount is the amount described above
5 for depot administration.

The invention here is the new compounds of the invention and new methods of using the compounds of the invention. Given a particular compound of the invention and a desired dosage form, one skilled in the art would know how to prepare and
10 administer the appropriate dosage form.

The compounds of the invention are used in the same manner, by the same routes of administration, using the same pharmaceutical dosage forms, and at the same dosing schedule as described above, for preventing disease or treating patients
15 with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating or preventing Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid
20 angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear
25 palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type of Alzheimer's disease.

The compounds of the invention can be used in combination, with each other or with other therapeutic agents or approaches used to treat or prevent the conditions listed above. Such
30 agents or approaches include: acetylcholine esterase inhibitors such as tacrine (tetrahydroaminoacridine, marketed as COGNEX®), donepezil hydrochloride, (marketed as Aricept® and rivastigmine (marketed as Exelon®); gamma-secretase

inhibitors; anti-inflammatory agents such as cyclooxygenase II inhibitors; anti-oxidants such as Vitamin E and ginkgolides; immunological approaches, such as, for example, immunization with A beta peptide or administration of anti-A beta peptide
5 antibodies; statins; and direct or indirect neurotropic agents such as Cerebrolysin®, AIT-082 (Emilieu, 2000, Arch. Neurol. 57:454), and other neurotropic agents of the future.

In addition, the compounds of formula (Y) can also be used with inhibitors of P-glycoprotein (P-gp). The use of P-gp
10 inhibitors is known to those skilled in the art. See for example, Cancer Research, 53, 4595-4602 (1993), Clin. Cancer Res., 2, 7-12 (1996), Cancer Research, 56, 4171-4179 (1996), International Publications WO99/64001 and WO01/10387. The important thing is that the blood level of the P-gp inhibitor
15 be such that it exerts its effect in inhibiting P-gp from decreasing brain blood levels of the compounds of formula (I). To that end the P-gp inhibitor and the compounds of formula (I) can be administered at the same time, by the same or different route of administration, or at different times. The important
20 thing is not the time of administration but having an effective blood level of the P-gp inhibitor.

Suitable P-gp inhibitors include cyclosporin A, verapamil, tamoxifen, quinidine, Vitamin E-TGPS, ritonavir, megestrol acetate, progesterone, rapamycin, 10,11-methanodibenzosuberane,
25 phenothiazines, acridine derivatives such as GF120918, FK506, VX-710, LY335979, PSC-833, GF-102,918 and other steroids. It is to be understood that additional agents will be found that do the same function and are also considered to be useful.

The P-gp inhibitors can be administered orally,
30 parenterally, (IV, IM, IM-depo, SQ, SQ-depo), topically, sublingually, rectally, intranasally, intrathecally and by implant.

The therapeutically effective amount of the P-gp inhibitors is from about 0.1 to about 300 mg/kg/day, preferably about 0.1 to about 150 mg/kg daily. It is understood that while a patient may be started on one dose, that dose may have
5 to be varied over time as the patient's condition changes.

When administered orally, the P-gp inhibitors can be administered in usual dosage forms for oral administration as is known to those skilled in the art. These dosage forms include the usual solid unit dosage forms of tablets and
10 capsules as well as liquid dosage forms such as solutions, suspensions and elixirs. When the solid dosage forms are used, it is preferred that they be of the sustained release type so that the P-gp inhibitors need to be administered only once or twice daily. The oral dosage forms are administered to the
15 patient one thru four times daily. It is preferred that the P-gp inhibitors be administered either three or fewer times a day, more preferably once or twice daily. Hence, it is preferred that the P-gp inhibitors be administered in solid dosage form and further it is preferred that the solid dosage
20 form be a sustained release form which permits once or twice daily dosing. It is preferred that what ever dosage form is used, that it be designed so as to protect the P-gp inhibitors from the acidic environment of the stomach. Enteric coated tablets are well known to those skilled in the art. In
25 addition, capsules filled with small spheres each coated to protect from the acidic stomach, are also well known to those skilled in the art.

In addition, the P-gp inhibitors can be administered parenterally. When administered parenterally they can be
30 administered IV, IM, depo-IM, SQ or depo-SQ.

The P-gp inhibitors can be given sublingually. When given sublingually, the P-gp inhibitors should be given one thru four times daily in the same amount as for IM administration.

The P-gp inhibitors can be given intranasally. When given by this route of administration, the appropriate dosage forms are a nasal spray or dry powder as is known to those skilled in the art. The dosage of the P-gp inhibitors for intranasal
5 administration is the same as for IM administration.

The P-gp inhibitors can be given intrathecally. When given by this route of administration the appropriate dosage form can be a parenteral dosage form as is known to those skilled in the art.

10 The P-gp inhibitors can be given topically. When given by this route of administration, the appropriate dosage form is a cream, ointment or patch. Because of the amount of the P-gp inhibitors needed to be administered the patch is preferred. However, the amount that can be delivered by a patch is
15 limited. Therefore, two or more patches may be required. The number and size of the patch is not important, what is important is that a therapeutically effective amount of the P-gp inhibitors be delivered as is known to those skilled in the art.

20 The P-gp inhibitors can be administered rectally by suppository as is known to those skilled in the art.

The P-gp inhibitors can be administered by implants as is known to those skilled in the art.

The exact dosage and frequency of administration will
25 depend on the particular compounds of the invention administered, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, and other medication the individual may be taking as is well known to
30 administering physicians who are skilled in this art.

Inhibition of APP Cleavage

The compounds of the invention inhibit cleavage of APP between Met595 and Asp596 numbered for the APP695 isoform, or a mutant thereof, or at a corresponding site of a different isoform, such as APP751 or APP770, or a mutant thereof
5 (sometimes referred to as the "beta secretase site"). While not wishing to be bound by a particular theory, inhibition of beta-secretase activity is thought to inhibit production of beta amyloid peptide (A beta). Inhibitory activity is demonstrated in one of a variety of inhibition assays, whereby
10 cleavage of an APP substrate in the presence of a beta-secretase enzyme is analyzed in the presence of the inhibitory compound, under conditions normally sufficient to result in cleavage at the beta-secretase cleavage site. Reduction of APP cleavage at the beta-secretase cleavage site compared with an
15 untreated or inactive control is correlated with inhibitory activity. Assay systems that can be used to demonstrate efficacy of the compound inhibitors of the invention are known. Representative assay systems are described, for example, in U.S. Patents No. 5,942,400, 5,744,346, as well as in the
20 Examples below.

The enzymatic activity of beta-secretase and the production of A beta can be analyzed *in vitro* or *in vivo*, using natural, mutated, and/or synthetic APP substrates, natural, mutated, and/or synthetic enzyme, and the test compound. The
25 analysis may involve primary or secondary cells expressing native, mutant, and/or synthetic APP and enzyme, animal models expressing native APP and enzyme, or may utilize transgenic animal models expressing the substrate and enzyme. Detection of enzymatic activity can be by analysis of one or more of the
30 cleavage products, for example, by immunoassay, flurometric or chromogenic assay, HPLC, or other means of detection. Inhibitory compounds are determined as those having the ability to decrease the amount of beta-secretase cleavage product

produced in comparison to a control, where beta-secretase mediated cleavage in the reaction system is observed and measured in the absence of inhibitory compounds.

5 Beta-secretase

Various forms of beta-secretase enzyme are known, and are available and useful for assay of enzyme activity and inhibition of enzyme activity. These include native, recombinant, and synthetic forms of the enzyme. Human beta-secretase is known as Beta Site APP Cleaving Enzyme (BACE), Asp2, and memapsin 2, and has been characterized, for example, in U.S. Patent No. 5,744,346 and published PCT patent applications WO98/22597, WO00/03819, WO01/23533, and WO00/17369, as well as in literature publications (Hussain et.al., 1999, *Mol.Cell.Neurosci.* 14:419-427; Vassar et.al., 1999, *Science* 286:735-741; Yan et.al., 1999, *Nature* 402:533-537; Sinha et.al., 1999, *Nature* 40:537-540; and Lin et.al., 2000, *PNAS USA* 97:1456-1460). Synthetic forms of the enzyme have also been described (WO98/22597 and WO00/17369). Beta-secretase can be extracted and purified from human brain tissue and can be produced in cells, for example mammalian cells expressing recombinant enzyme.

Preferred compounds inhibit about 50% of beta-secretase enzymatic activity at a concentration of less than about 50 micromolar, preferably at a concentration of about 10 micromolar or less, more preferably about 1 micromolar or less, and most preferably about 10 nanomolar or less.

APP substrate

Assays that demonstrate inhibition of beta-secretase-mediated cleavage of APP can utilize any of the known forms of APP, including the 695 amino acid "normal" isotype described by Kang et.al., 1987, *Nature* 325:733-6, the 770 amino acid isotype described by Kitaguchi et. al., 1981, *Nature* 331:530-532, and variants such as the Swedish Mutation (KM670-1NL) (APP-SW), the London Mutation (V7176F), and others. See, for example, U.S. Patent No. 5,766,846 and also Hardy, 1992, *Nature Genet.* 1:233-234, for a review of known variant mutations. Additional useful substrates include the dibasic amino acid modification, APP-KK disclosed, for example, in WO 00/17369, fragments of APP, and synthetic peptides containing the beta-secretase cleavage site, wild type (WT) or mutated form, e.g., SW, as described, for example, in U.S. Patent No 5,942,400 and WO00/03819.

The APP substrate contains the beta-secretase cleavage site of APP (KM-DA or NL-DA) for example, a complete APP peptide or variant, an APP fragment, a recombinant or synthetic APP, or a fusion peptide. Preferably, the fusion peptide includes the beta-secretase cleavage site fused to a peptide having a moiety useful for enzymatic assay, for example, having isolation and/or detection properties. A useful moiety may be an antigenic epitope for antibody binding, a label or other detection moiety, a binding substrate, and the like.

Antibodies

Products characteristic of APP cleavage can be measured by immunoassay using various antibodies, as described, for example, in Pirttila et.al., 1999, *Neuro.Lett.* 249:21-4, and in U.S. Patent No. 5,612,486. Useful antibodies to detect A beta include, for example, the monoclonal antibody 6E10 (Senetek, St. Louis, MO) that specifically recognizes an epitope on amino

acids 1-16 of the A beta peptide; antibodies 162 and 164 (New York State Institute for Basic Research, Staten Island, NY) that are specific for human A beta 1-40 and 1-42, respectively; and antibodies that recognize the junction region of beta-amyloid peptide, the site between residues 16 and 17, as described in U.S. Patent No. 5,593,846. Antibodies raised against a synthetic peptide of residues 591 to 596 of APP and SW192 antibody raised against 590-596 of the Swedish mutation are also useful in immunoassay of APP and its cleavage products, as described in U.S. Patent Nos. 5,604,102 and 5,721,130.

Assay Systems

Assays for determining APP cleavage at the beta-secretase cleavage site are well known in the art. Exemplary assays, are described, for example, in U.S. Patent Nos. 5,744,346 and 5,942,400, and described in the Examples below.

Cell free assays

Exemplary assays that can be used to demonstrate the inhibitory activity of the compounds of the invention are described, for example, in WO00/17369, WO 00/03819, and U.S. Patents No. 5,942,400 and 5,744,346. Such assays can be performed in cell-free incubations or in cellular incubations using cells expressing a beta-secretase and an APP substrate having a beta-secretase cleavage site.

An APP substrate containing the beta-secretase cleavage site of APP, for example, a complete APP or variant, an APP fragment, or a recombinant or synthetic APP substrate containing the amino acid sequence: KM-DA or NL-DA, is incubated in the presence of beta-secretase enzyme, a fragment thereof, or a synthetic or recombinant polypeptide variant having beta-secretase activity and effective to cleave the

beta-secretase cleavage site of APP, under incubation conditions suitable for the cleavage activity of the enzyme. Suitable substrates optionally include derivatives that may be fusion proteins or peptides that contain the substrate peptide and a modification useful to facilitate the purification or detection of the peptide or its beta-secretase cleavage products. Useful modifications include the insertion of a known antigenic epitope for antibody binding; the linking of a label or detectable moiety, the linking of a binding substrate, and the like.

Suitable incubation conditions for a cell-free *in vitro* assay include, for example: approximately 200 nanomolar to 10 micromolar substrate, approximately 10 to 200 picomolar enzyme, and approximately 0.1 nanomolar to 10 micromolar inhibitor compound, in aqueous solution, at an approximate pH of 4 -7, at approximately 37 degrees C, for a time period of approximately 10 minutes to 3 hours. These incubation conditions are exemplary only, and can be varied as required for the particular assay components and/or desired measurement system. Optimization of the incubation conditions for the particular assay components should account for the specific beta-secretase enzyme used and its pH optimum, any additional enzymes and/or markers that might be used in the assay, and the like. Such optimization is routine and will not require undue experimentation.

One useful assay utilizes a fusion peptide having maltose binding protein (MBP) fused to the C-terminal 125 amino acids of APP-SW. The MBP portion is captured on an assay substrate by anti-MBP capture antibody. Incubation of the captured fusion protein in the presence of beta-secretase results in cleavage of the substrate at the beta-secretase cleavage site. Analysis of the cleavage activity can be, for example, by immunoassay of cleavage products. One such immunoassay detects

a unique epitope exposed at the carboxy terminus of the cleaved fusion protein, for example, using the antibody SW192. This assay is described, for example, in U.S. Patent No 5,942,400.

5 Cellular assay

Numerous cell-based assays can be used to analyze beta-secretase activity and/or processing of APP to release A beta. Contact of an APP substrate with a beta-secretase enzyme within the cell and in the presence or absence of a compound inhibitor of the invention can be used to demonstrate beta-secretase inhibitory activity of the compound. Preferably, assay in the presence of a useful inhibitory compound provides at least about 30%, most preferably at least about 50% inhibition of the enzymatic activity, as compared with a non-inhibited control.

15 In one embodiment, cells that naturally express beta-secretase are used. Alternatively, cells are modified to express a recombinant beta-secretase or synthetic variant enzyme as discussed above. The APP substrate may be added to the culture medium and is preferably expressed in the cells. 20 Cells that naturally express APP, variant or mutant forms of APP, or cells transformed to express an isoform of APP, mutant or variant APP, recombinant or synthetic APP, APP fragment, or synthetic APP peptide or fusion protein containing the beta-secretase APP cleavage site can be used, provided that the 25 expressed APP is permitted to contact the enzyme and enzymatic cleavage activity can be analyzed.

Human cell lines that normally process A beta from APP provide a useful means to assay inhibitory activities of the compounds of the invention. Production and release of A beta 30 and/or other cleavage products into the culture medium can be measured, for example by immunoassay, such as Western blot or enzyme-linked immunoassay (EIA) such as by ELISA.

Cells expressing an APP substrate and an active beta-secretase can be incubated in the presence of a compound inhibitor to demonstrate inhibition of enzymatic activity as compared with a control. Activity of beta-secretase can be
5 measured by analysis of one or more cleavage products of the APP substrate. For example, inhibition of beta-secretase activity against the substrate APP would be expected to decrease release of specific beta-secretase induced APP cleavage products such as A beta.

10 Although both neural and non-neural cells process and release A beta, levels of endogenous beta-secretase activity are low and often difficult to detect by EIA. The use of cell types known to have enhanced beta-secretase activity, enhanced processing of APP to A beta, and/or enhanced production of A
15 beta are therefore preferred. For example, transfection of cells with the Swedish Mutant form of APP (APP-SW); with APP-KK; or with APP-SW-KK provides cells having enhanced beta-secretase activity and producing amounts of A beta that can be readily measured.

20 In such assays, for example, the cells expressing APP and beta-secretase are incubated in a culture medium under conditions suitable for beta-secretase enzymatic activity at its cleavage site on the APP substrate. On exposure of the cells to the compound inhibitor, the amount of A beta released
25 into the medium and/or the amount of CTF99 fragments of APP in the cell lysates is reduced as compared with the control. The cleavage products of APP can be analyzed, for example, by immune reactions with specific antibodies, as discussed above.

Preferred cells for analysis of beta-secretase activity
30 include primary human neuronal cells, primary transgenic animal neuronal cells where the transgene is APP, and other cells such as those of a stable 293 cell line expressing APP, for example, APP-SW.

In vivo assays: animal models

Various animal models can be used to analyze beta-secretase activity and /or processing of APP to release A beta, as described above. For example, transgenic animals expressing APP substrate and beta-secretase enzyme can be used to demonstrate inhibitory activity of the compounds of the invention. Certain transgenic animal models have been described, for example, in U.S. Patent Nos: 5,877,399; 5,612,486; 5,387,742; 5,720,936; 5,850,003; 5,877,015,, and 5,811,633, and in Ganes et.al., 1995, Nature 373:523. Preferred are animals that exhibit characteristics associated with the pathophysiology of AD. Administration of the compound inhibitors of the invention to the transgenic mice described herein provides an alternative method for demonstrating the inhibitory activity of the compounds. Administration of the compounds in a pharmaceutically effective carrier and via an administrative route that reaches the target tissue in an appropriate therapeutic amount is also preferred.

Inhibition of beta-secretase mediated cleavage of APP at the beta-secretase cleavage site and of A beta release can be analyzed in these animals by measure of cleavage fragments in the animal's body fluids such as cerebral fluid or tissues. Analysis of brain tissues for A beta deposits or plaques is preferred.

On contacting an APP substrate with a beta-secretase enzyme in the presence of an inhibitory compound of the invention and under conditions sufficient to permit enzymatic mediated cleavage of APP and/or release of A beta from the substrate, the compounds of the invention are effective to reduce beta-secretase-mediated cleavage of APP at the beta-secretase cleavage site and/or effective to reduce released amounts of A beta. Where such contacting is the administration

of the inhibitory compounds of the invention to an animal model, for example, as described above, the compounds are effective to reduce A beta deposition in brain tissues of the animal, and to reduce the number and/or size of beta amyloid plaques. Where such administration is to a human subject, the compounds are effective to inhibit or slow the progression of disease characterized by enhanced amounts of A beta, to slow the progression of AD in the, and/or to prevent onset or development of AD in a patient at risk for the disease.

Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are hereby incorporated by reference for all purposes.

DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

I. CONVENTIONS FOR FORMULAS AND DEFINITIONS OF VARIABLES

When the claims contain a fairly complex (cyclic) substituent, at the end of the phrase naming/designating that particular substituent will be a notation in (parentheses) which will correspond to the same name/designation in one of the CHARTS which will also set forth the chemical structural formula of that particular substituent.

II. DEFINITIONS

All temperatures are in degrees Celsius.

TLC refers to thin-layer chromatography.

psi refers to pounds/in².

- HPLC refers to high pressure liquid chromatography.
THF refers to tetrahydrofuran.
DMF refers to dimethylformamide.
EDC refers to ethyl-1-(3-dimethylaminopropyl)carbodiimide
5 or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride.
HOBT refers to 1-hydroxy benzotriazole hydrate.
NMM refers to N-methylmorpholine.
NBS refers to N-bromosuccinimide.
TEA refers to triethylamine.
10 BOC refers to 1,1-dimethylethoxy carbonyl or t-butoxycarbonyl, $-\text{CO}-\text{O}-\text{C}(\text{CH}_3)_3$.
CBZ refers to benzyloxycarbonyl, $-\text{CO}-\text{O}-\text{CH}_2$ -phenyl.
Fmoc refers to 9-fluorenylmethyl carbonate.
TFA refers to trifluoroacetic acid, $\text{CF}_3\text{-COOH}$.
15 CDI refers to 1,1'-carbonyldiimidazole.
Saline refers to an aqueous saturated sodium chloride solution.

Chromatography (column and flash chromatography) refers to purification/separation of compounds expressed as (support,
20 eluent). It is understood that the appropriate fractions are pooled and concentrated to give the desired compound(s).

CMR refers to C-13 magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

NMR refers to nuclear (proton) magnetic resonance
25 spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

IR refers to infrared spectroscopy.

-phenyl refers to phenyl (C_6H_5).

MS refers to mass spectrometry expressed as m/e, m/z or
30 mass/charge unit. MH^+ refers to the positive ion of a parent plus a hydrogen atom. EI refers to electron impact. CI refers to chemical ionization. FAB refers to fast atom bombardment.
HRMS refers to high resolution mass spectrometry.

Ether refers to diethyl ether.

Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the
 5 manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

When solvent pairs are used, the ratios of solvents used are volume/volume (v/v).

10 When the solubility of a solid in a solvent is used the ratio of the solid to the solvent is weight/volume (wt/v).

BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate.

TBDMSCl refers to t-butyldimethylsilyl chloride.

15 TBDSOTf refers to t-butyldimethylsilyl trifluosulfonic acid ester.

Trisomy 21 refers to Down's Syndrome.

The following terms are used (in EXAMPLEs 321 and above) for the

20 amide forming agent (IX):

"PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-phenyl-CO-OH}$ where the attachment to the - phenyl- ring is 1,3-;

"5-Me-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(CH}_3\text{-) phenyl -CO-OH}$ where the attachment to the - phenyl - ring is 1,3- for the
 25 carbonyl groups and 5- for the methyl group;

"3,5-pyridinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(pyridinyl)-CO-OH}$ where the attachment to the -pyridinyl- ring is 3,5- for the carbonyl groups;

"-SO₂-" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{CH-SO}_2\text{- phenyl -CO-OH}$ where
 30 the attachment to the - phenyl - ring is 1,3-;

"5-OMe-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(CH}_3\text{-O-) phenyl -CO-OH}$ where the attachment to the - phenyl - ring is 1,3- for the carbonyl groups and 5- for the methoxy group;

"5-Cl-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(Cl-phenyl-CO-OH)}$ where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups and 5- for the chlorine atom;

5 "5-F-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(F-phenyl-CO-OH)}$ where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups and 5- for the fluorine atom;

"thienyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-thienyl-CO-OH}$ where the attachment to the thiophene ring is -2,5;

10 "2,4-pyridinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(pyridinyl-CO-OH)}$ where the attachment to the -pyridinyl- ring is 2,4- for the carbonyl groups;

"4,6-pyrimidinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-})_2\text{N-CO-(pyrimidinyl-phenyl-CO-OH)}$ where the attachment to the -pyrimidinyl- ring is 4,6- for the carbonyl groups;

15 "morpholinyl" refers to morpholinyl-CO-phenyl-CO-OH where the attachment to the -phenyl- ring is 1,3 for the carbonyl groups.

20 APP, amyloid precursor protein, is defined as any APP polypeptide, including APP variants, mutations, and isoforms, for example, as disclosed in U.S. Patent No. 5,766,846.

A beta, amyloid beta peptide, is defined as any peptide resulting from beta-secretase mediated cleavage of APP, including peptides of 39, 40, 41, 42, and 43 amino acids, and extending from the beta-secretase cleavage site to amino acids
25 39, 40, 41, 42, or 43.

Beta-secretase (BACE1, Asp2, Memapsin 2) is an aspartyl protease that mediates cleavage of APP at the amino-terminal edge of A beta. Human beta-secretase is described, for example, in WO00/17369.

30 "Pharmaceutically acceptable" refers to those properties and/or substances that are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical

point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

A therapeutically effective amount is defined as an amount effective to reduce or lessen at least one symptom of the disease being treated or to reduce or delay onset of one or more clinical markers or symptoms of the disease.

By "alkyl" and "C₁-C₆ alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl. It is understood that in cases where an alkyl chain of a substituent (e.g. of an alkyl, alkoxy or alkenyl group) is shorter or longer than 6 carbons, it will be so indicated in the second "C" as, for example, "C₁-C₁₀" indicates a maximum of 10 carbons.

By "alkoxy" and "C₁-C₆ alkoxy" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, attached through at least one divalent oxygen atom, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, isopentoxy, neopentoxy, hexoxy, and 3-methylpentoxy.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

"Alkenyl" and "C₂-C₆ alkenyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and from one to three double bonds and includes, for example, ethenyl, propenyl, 1-but-3-enyl, 1-pent-3-enyl, 1-hex-5-enyl and the like.

"Alkynyl" and "C₂-C₆ alkynyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and one or two triple bonds and includes ethynyl, propynyl, butynyl, pentyn-2-yl and the like.

As used herein, the term "cycloalkyl" refers to saturated carbocyclic radicals having three to twelve carbon atoms. The cycloalkyl can be monocyclic, or a polycyclic fused system. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The cycloalkyl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such cycloalkyl groups may be optionally substituted with C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, amino(C₁-C₆)alkyl, mono(C₁-C₆)alkylamino(C₁-C₆)alkyl or di(C₁-C₆)alkylamino(C₁-C₆)alkyl.

By "aryl" is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl), which is optionally mono-, di-, or trisubstituted. Preferred aryl groups of the present invention are phenyl, 1-naphthyl, 2-naphthyl, indanyl, indenyl, dihydronaphthyl, tetralinyl or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl. The aryl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such aryl groups may be optionally substituted with, for example, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, amino(C₁-C₆)alkyl, mono(C₁-C₆)alkylamino(C₁-C₆)alkyl, di(C₁-C₆)alkylamino(C₁-C₆)alkyl, -COOH, -C(=O)O(C₁-C₆ alkyl), -C(=O)NH₂, -C(=O)N(mono- or di-C₁-C₆ alkyl), -S(C₁-C₆ alkyl), -SO₂(C₁-C₆ alkyl), -O-C(=O)(C₁-C₆ alkyl), -NH-C(=O)-(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)-C(=O)-(C₁-C₆ alkyl), -NH-SO₂-(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)-SO₂-(C₁-C₆ alkyl), -NH-C(=O)NH₂, -NH-C(=O)N(mono- or

di-C₁-C₆ alkyl), -NH(C₁-C₆ alkyl)-C(=O)-NH₂ or -NH(C₁-C₆ alkyl)-C(=O)-N-(mono- or di-C₁-C₆ alkyl).

By "heteroaryl" is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heteroaryl groups of the present invention include pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, 10 quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indolizinyl, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, 15 isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, isochromanyl, chromanyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranlyl, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranlyl, benzotetrahydrothienyl, 20 purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl, benzopyranyl, benzothiopyranyl, coumarinyl, isocoumarinyl, chromonyl, chromanonyl, pyridinyl-N-oxide; tetrahydroquinolinyl, dihydroquinolinyl, 25 dihydroquinolinonyl, dihydroisoquinolinonyl, dihydrocoumarinyl, dihydroisocoumarinyl, isoindolinonyl, benzodioxanyl, benzoxazolinonyl, pyrrolyl N-oxide,, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, 30 indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indolizinyl N-oxide, indazolyl N-oxide,

benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, benzothiopyranyl S,S-dioxide. The heteroaryl groups herein are

5 unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such heteroaryl groups may be optionally substituted with C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, C₂-C₆alkenyl, C₂-

10 C₆alkynyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, amino(C₁-C₆)alkyl, mono(C₁-C₆)alkylamino(C₁-C₆)alkyl or di(C₁-C₆)alkylamino(C₁-C₆)alkyl, -COOH, -C(=O)O(C₁-C₆ alkyl), -C(=O)NH₂, -C(=O)N(mono- or di-C₁-C₆ alkyl), -S(C₁-C₆ alkyl), -SO₂(C₁-C₆ alkyl), -O-C(=O)(C₁-C₆ alkyl), -NH-C(=O)-(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)-

15 C(=O)-(C₁-C₆ alkyl), -NH-SO₂-(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)-SO₂-(C₁-C₆ alkyl), -NH-C(=O)NH₂, -NH-C(=O)N(mono- or di-C₁-C₆ alkyl), -NH(C₁-C₆ alkyl)-C(=O)-NH₂ or -NH(C₁-C₆ alkyl)-C(=O)-N-(mono- or di-C₁-C₆ alkyl).

By "heterocycle", "heterocycloalkyl" or "heterocyclyl" is

20 meant one or more carbocyclic ring systems of 3-, 4-, 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heterocycles of the present invention include morpholinyl,

25 thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide,

30 oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl, dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, azepanyl, diazepanyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide and

homothiomorpholinyl S-oxide. The heterocycle groups herein maybe unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such heterocycle groups may be optionally substituted with C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, hydroxy, cyano, nitro, amino, 5 mono(C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, amino(C₁-C₆)alkyl, mono(C₁-C₆)alkylamino(C₁-C₆)alkyl, di(C₁-C₆)alkylamino(C₁-C₆)alkyl or =O.

10 Structures were named using Name Pro IUPAC Naming Software, version 4.0 or 5.09, available from Advanced Chemical Development, Inc., 90 Adelaide Street West, Toronto, Ontario, M5H 3V9, Canada.

15 All patents and publications referred to herein are hereby incorporated by reference for all purposes.

The present invention provides compounds, compositions, and methods for inhibiting beta-secretase enzyme activity and A beta peptide production. Inhibition of beta-secretase enzyme activity halts or reduces the production of A beta from APP and 20 reduces or eliminates the formation of beta-amyloid deposits in the brain.

EXAMPLES

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, practice the present invention to its fullest extent. The following detailed examples describe how to prepare the various compounds and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques.

PREPARATION 1 3-Amino-5-(methoxycarbonyl)benzoic acid (XVII)

A suspension of *mono*-methyl 5-nitro-isophthalate (22.5 g, 100 mmol) and palladium on carbon (5%, 2.00 g) in methanol (100 mL) is shaken in a hydrogenation apparatus under hydrogen (50 psi) for 3 hours. The mixture is then filtered through diatomaceous earth and concentrated to give the title compound, NMR (300 MHz, CDCl₃) delta 7.67, 7.41, 7.40 and 3.83; MS (ESI-) for C₉H₉NO₄ *m/z* (M-H)⁻ = 194.

PREPARATION 2 3-Bromo-5-(methoxycarbonyl)benzoic acid (XIX)

A mixture of copper (II) bromide (1.85 g, 8.30 mmol), *n*-butyl nitrite (1.07 g, 10.4 mmol), and acetonitrile (30 mL) is stirred in a round bottomed flask in a water bath to which a few chunks of ice has been added. 3-Amino-5-(methoxycarbonyl)benzoic acid (XVII, PREPARATION 1, 1.35 g, 6.92 mmol) is added as a slurry in warm acetonitrile (70 mL) over 15 min and the mixture is stirred at 20-25 degrees C for an additional 2 hour, at which time the mixture is partitioned between dichloromethane and hydrochloric acid (3N). The organic phase is separated and dried over sodium sulfate and concentrated to dryness. Chromatography (silica gel, 125 mL;

methanol/dichloromethane, 15/85) and concentration of the appropriate fractions gives a solid which is crystallized from methanol to give the title compound in two crops, NMR (DMSO- d_6) δ 3.90, 8.26 and 8.65.

5

PREPARATION 3 Methyl 3-bromo-5-
[(dipropylamino)carbonyl]benzoate (XXI)

Carbonyl diimidazole (3.0 g, 18 mmol) is added to a solution of 3-bromo-5-(methoxycarbonyl)benzoic acid (XIX,
10 PREPARATION 2, 3.9 g, 15 mmol) in THF (30 mL). The mixture is stirred for 0.5 hours. Dipropylamine (AMINE, 4.2 mL, 30 mmol) is added to the mixture, which is then stirred for 24 hours. The solvent is then removed under reduced pressure and the mixture is partitioned between ethyl acetate and water. The
15 organic phase is then washed with saline, dried over anhydrous magnesium sulfate, filtered, and concentrated. Column chromatography (silica gel; ethyl acetate/hexanes, 15/85) gives the title compound, IR (diffuse reflectance) 2968, 2958, 1714, 1637, 1479, 1440, 1422, 1321, 1310, 1288, 1273, 1252, 889, 772
20 and 718 cm^{-1} ; NMR (300 MHz, CDCl_3) δ 8.21, 7.96, 7.70, 3.95, 3.46, 3.15, 1.69, 1.57, 1.00 and 0.78; MS (ESI+) for $\text{C}_{15}\text{H}_{20}\text{BrNO}_3$ m/z (M+H) $^+$ = 344.1.

PREPARATION 4 3-Bromo-5-[(dipropylamino)carbonyl]benzoic acid

25 To a solution of methyl 3-bromo-5-
[(dipropylamino)carbonyl]benzoate (XXI, PREPARATION 3, 1.4 g, 4.1 mmol) in THF/water/methanol (4/2/2, 8 mL) is added to lithium hydroxide monohydrate (0.17 g, 4.05 mmol). The mixture is stirred at 20 degrees -25 degrees C for 1 hour and then
30 solvent is removed under reduced pressure. The residue is dissolved in water (50 mL) and hydrochloric acid (1 N) is added to adjust the pH to about 3. The aqueous mixture is extracted with ethyl acetate and the organic phase is separated and dried

over magnesium sulfate to give the title compound. Analytical calculated for $C_{14}H_{18}BrNO_3$: C, 51.23; H, 5.53; N, 4.27; Br, 24.35. Found: C, 51.37; H, 5.56; N, 4.28.

5 PREPARATION 5 Methyl 3-(aminocarbonyl)-5-

[(dipropylamino)carbonyl]-benzoate (XXII)

To a mixture of methyl 3-bromo-5-
[(dipropylamino)carbonyl]benzoate (XXI, PREPARATION 3, 0.5 g, 1.47 mmol) in dry N-methyl pyrrolidinone under a carbon
10 monoxide atmosphere is added palladium (II) acetate (0.017 g, 0.074 mmol), 1,3-bis(diphenylphosphino)propane (0.045 g, 0.11 mmol), hexamethyldisilazane (1.0 mL, 4.7 mmol), and diisopropylethylamine (0.38 g, 2.94 mmol). The mixture is heated at 100 degrees C for 24 hours. The mixture is cooled to
15 20-25 degrees C and partitioned between water and ethyl acetate. The layers are separated and the aqueous phase is back-washed with ethyl acetate. The organic phases are combined and washed three times with saline, dried over anhydrous magnesium sulfate, filtered and concentrated. Column
20 chromatography (silica gel, 75 mL; methanol/methylene chloride, 2.5/97.5) gives the title compound, NMR ($CDCl_3$) delta 0.77, 1.02, 1.57, 1.71, 3.17, 3.49, 3.98, 5.78, 6.34, 8.07, 8.20 and 8.48.

25 PREPARATION 6 3-(Aminocarbonyl)-5-

[(dipropylamino)carbonyl]benzoic acid (XXIII)

To a mixture of methyl 3-(aminocarbonyl)-5-
[(dipropylamino)carbonyl]benzoate (XXII, PREPARATION 5, 0.197 g, 0.64 mmol) in methanol (5.0 mL) is added sodium hydroxide
30 (1N, 3.0 mL). The mixture is stirred at 20-25 degrees C for 24 hours. The mixture is acidified to about pH 5 with hydrochloric acid (10%). Water (50 mL) is added and the mixture is washed twice with ethyl acetate (2 x 50 mL). The

organic extracts are combined and dried over anhydrous magnesium sulfate and concentrated to give the title compound, NMR (DMSO- d_6) δ 0.66, 0.930, 1.48, 1.62, 3.12, 3.35, 7.54, 7.98, 8.22 and 8.51.

5

PREPARATION 7 3-Cyano-5-[(dipropylamino)carbonyl]benzoic acid (IX/XXXII)

A mixture of 3-bromo-5-[(dipropylamino)carbonyl]benzoic acid (PREPARATION 4, 0.596 g, 1.82 mmol) and copper nitrile
10 (0.325 g, 3.63 mmol) in N-methylpyrrolidinone (1.5 mL) is stirred at 175 degrees C for 2.5 hour, at which time the mixture is cooled and partitioned between ethyl acetate and hydrochloric acid (3N). The organic layer is washed twice more with hydrochloric acid (3N) and then twice more with saline
15 which had been acidified with a small amount of hydrochloric acid (3N). The organic layer is dried over magnesium sulfate and concentrated under high vacuum to give the title compound, NMR (CDCl₃) δ 0.80, 1.02, 1.60, 1.73, 3.17, 3.51, 7.90, 8.31 and 8.41; an aliquot is crystallized from ethyl
20 ether/dichloromethane/hexane - IR (diffuse reflectance) 3017, 2970, 2937, 2898, 2877, 2473; 2432, 2350, 2318, 2236, 1721, 1608, 1588, 1206 and 1196 cm⁻¹.

PREPARATION 8 3-(Aminocarbonyl)-5-

25 [(dipropylamino)carbonyl]benzoic acid (XXXIII)

A mixture of 3-cyano-5-[(dipropylamino)carbonyl]benzoic acid (IX/XXXII, PREPARATION 7, 0.602 g, 2.19 mmol), potassium carbonate (0.212 g, 1.53 mmol), and acetone (2.5 mL) is stirred
30 at 20-25 degrees C. Water (2.5 mL) and urea-hydrogen peroxide adduct (0.825 g, 8.78 mmol) are added and the mixture is stirred for 15 hours at 20-25 degrees C, at which time additional urea-hydrogen peroxide adduct (0.204 g) is added;

after stirring for another 3 hours, an additional 0.205 g of urea-hydrogen peroxide is added. After a total of 39 hours has elapsed, the acetone is removed under reduced pressure and the residue is acidified with hydrochloric acid (3N) to pH = 2-4.

- 5 The mixture is extracted with dichloromethane, the organic layer is separated and washed with hydrochloric acid (0.5 N), and the organic phase is dried with anhydrous magnesium sulfate to a solid. The solid is crystallized from dichloromethane/hexane/methanol to give the title compound, MS
10 (ESI+) for $C_{15}H_{20}N_2O_4$ m/z $(M+H)^+ = 293.2$.

PREPARATION 9 Methyl 3-[(dipropylamino)carbonyl]-5-nitrobenzoate (XXX)

- Carbonyl diimidazole (3.90 g, 24.0 mmol) is added to a
15 mixture of mono-methyl 5-nitro-isophthalate (XXVIII, 4.50 g, 20.0 mmol) in dry THF (50 mL). The mixture is stirred for 0.5 hours. Dipropylamine (3.28 mL, 24.0 mmol) is added slowly to the mixture. The reaction mixture is then stirred for 4 hours. The solvent is removed under reduced pressure and the mixture
20 is partitioned between ethyl acetate and water. The organic phase is separated and washed with saline, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography (silica gel; ethyl acetate/hexanes, 15/85) gives the title compound, NMR (300 MHz, $CDCl_3$) δ 8.88, 8.41, 8.35,
25 4.00, 3.48, 3.15, 1.72, 1.57, 1.00 and 0.77; MS (ESI+) for $C_{15}H_{20}N_2O_5$ m/z $(M+H)^+ = 309.2$.

PREPARATION 10 Methyl 3-amino-5-

[(dipropylamino)carbonyl]benzoate (XXXI)

- 30 A suspension of methyl 3-[(dipropylamino)carbonyl]-5-nitrobenzoate (XXX, PREPARATION 9, 6.00g, 20.0 mmol) and palladium on carbon (5%, 0.600 g) in methanol (40 mL) is shaken in a hydrogenation apparatus under hydrogen (45 psi) for 3

hours. The mixture is then filtered through diatomaceous earth and concentrated to give the title compound, NMR (300 MHz, CDCl_3) delta 7.27, 6.77, 4.10, 3.82, 3.38, 3.10, 1.62, 1.46, 0.91 and 0.68.

5

PREPARATION 11 Methyl 3-(chlorosulfonyl)-5-
[(dipropylamino)carbonyl]- benzoate (XXXVII)

Methyl 3-amino-5-[(dipropylamino)carbonyl]benzoate (XXXI, PREPARATION 10, 1.11 g, 4 mmol) is added to a mixture of water (5 mL) and concentrated hydrochloric acid (1 mL). Sodium nitrite (0.276 g, 4 mmol) is added to the mixture slowly at 0 degrees C. The mixture is then added to an acetic acid solution (5 mL) of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ saturated with sulfur dioxide. The mixture is stirred for 0.5 hours and poured into ice water. The mixture is extracted with ethyl acetate. The organic phase is separated and washed with saturated sodium bicarbonate, water, and saline and dried over anhydrous sodium sulfate, filtered, and concentrated to give the title compound, NMR (300 MHz, CDCl_3) delta 8.69, 8.38, 8.20, 4.01, 3.49, 3.14, 1.72, 1.59, 1.01 and 0.79; MS (ESI+) for $\text{C}_{15}\text{H}_{20}\text{ClNO}_5\text{S}$ m/z (M+H)⁺ = 362.2

25

PREPARATION 12 Methyl 3-(aminosulfonyl)-5-
[(dipropylamino)carbonyl]- benzoate (XXXVIII)

To a solution of methyl 3-(chlorosulfonyl)-5-[(dipropylamino)carbonyl]benzoate (XXXVII, PREPARATION 11, 0.100 g, 0.300 mmol) in dry THF (3 mL) is added ammonia (7 N solution in methanol, 0.214 mL, 1.50 mmol). The mixture is stirred for 18 hours and solvent is then removed. The residue is partitioned between ethyl acetate and water. The organic phase is separate and washed with saline, dried over anhydrous sodium sulfate, filtered, and concentrated to give the title

compound, NMR (300 MHz, CDCl_3) δ 8.45, 8.07, 8.01, 6.05, 3.93, 3.44, 3.09, 1.67, 1.52, 0.96 and 0.73; MS (ESI+) for $\text{C}_{12}\text{H}_{22}\text{N}_2\text{O}_5\text{S}$ m/z (M+H)⁺ = 343.3.

5 PREPARATION 13 3-(Aminosulfonyl)-5-

[(dipropylamino)carbonyl]benzoic acid (XXXVIII)

Lithium hydroxide monohydrate (0.011 g, 0.263 mmol) is added to a solution of methyl 3-(aminosulfonyl)-5-[(dipropylamino)carbonyl]benzoate (XXXVIII, PREPARATION 12, 10 0.090 g, 0.263 mmol) in a mixture of THF/methanol/water (2/1/1, 2 mL). The mixture is stirred at 20-25 degrees C for 3 hours. The mixture is then diluted with water and hydrochloric acid (1 N) is added to bring the pH to less than 3. The aqueous solution is extracted with ethyl acetate. The organic phase is 15 separated and washed with saline, dried over anhydrous sodium sulfate, filtered and concentrated to give the title compound. ¹H NMR (300 MHz, CDCl_3) δ 10.36 (s, 1 H), 8.39 (s, 1 H), 8.09 (s, 2 H), 6.06 (s, 2 H), 3.48 (t, J = 7 Hz, 2 H), 3.15 (t, J = 7 Hz, 2 H), 1.71 (m, 2 H), 1.55 (m, 2 H), 0.97 (t, J = 7 20 Hz, 3 H), 0.74 (t, J = 7 Hz, 3 H). MS (ESI+) for $\text{C}_{11}\text{H}_{20}\text{N}_2\text{O}_5\text{S}$ m/z 329.2 (M+H)⁺.

PREPARATION 14 Methyl 3-[(dipropylamino)carbonyl]-5-(1-pyrrolidinylsulfonyl)- benzoate (XXXVIII)

25 Following the general procedure of PREPARATION 12 and making non-critical variations but using pyrrolidine (0.347 mL, 4.16 mmol), the title compound is obtained, MS (ESI+) for $\text{C}_{19}\text{H}_{28}\text{N}_2\text{O}_5\text{S}$ m/z (M+H)⁺ = 397.1.

30 PREPARATION 15 3-[(Dipropylamino)carbonyl]-5-(1-pyrrolidinylsulfonyl)benzoic acid (XXXIX)

Following the general procedure of PREPARATION 13 and making non-critical variations, the title compound is obtained, MS (ESI+) for $C_{18}H_{26}N_2O_5S$ m/z $(M+H)^+ = 383.3$.

5 PREPARATION 16 Methyl 3-[(dipropylamino)carbonyl]-5-
[(methylamino)-sulfonyl]benzoate (XXXVIII)

Following the general procedure of PREPARATION 12 and making non-critical variations but using methyl amine (2 N solution in THF, 0.692 mL, 1.38 mmol), the title compound is
10 obtained, MS (ESI+) for $C_{16}H_{24}N_2O_5S$ m/z $(M+H)^+ = 357.1$.

PREPARATION 17 3-[(Dipropylamino)carbonyl]-5-
[(methylamino)-sulfonyl]benzoic acid (XXXIX)

Following the general procedure of PREPARATION 13 and
15 making non-critical variations, the title compound is obtained, MS (ESI+) for $C_{15}H_{22}N_2O_5S$ m/z $(M+H)^+ = 343.1$.

PREPARATION 18 Methyl 3-[(dimethylamino)sulfonyl]-5-
[(dipropylamino)-carbonyl]benzoate (XXXVIII)

20 Following the general procedure of PREPARATION 12 and making non-critical variations but using dimethylamine (2 N solution in THF, 0.692 mL, 1.38 mmol), the title compound is obtained, MS (ESI+) for $C_{17}H_{26}N_2O_5S$ m/z $(M+H)^+ = 371.1$.

25 PREPARATION 19 3-[(Dimethylamino)sulfonyl]-5-
[(dipropylamino)carbonyl]-benzoic acid (XXXIX)

Following the general procedure of PREPARATION 13 and making non-critical variations, the title compound is obtained, MS (ESI+) for $C_{16}H_{24}N_2O_5S$ m/z $(M+H)^+ = 357.1$.

30 PREPARATION 20 Methyl 3-[(dipropylamino)carbonyl]-5-ethylbenzoate (IX)

Ethylboronic acid (0.800 g, 10.8 mmol),
 dichlorobis(triphenylphosphine)-palladium(II) (0.252 g, 0.360
 mmol), potassium carbonate (2.50 g, 18.0 mmol) and lithium
 chloride (0.151 g, 3.60 mmol) are added to a mixture of methyl
 5 3-bromo-5-[(dipropylamino)carbonyl]benzoate (1.23 g, 3.60 mmol)
 in dry DMF (20 mL). The mixture is heated at 100 degrees C for
 18 hours. The mixture is then partitioned between ethyl
 acetate and water. The phases are separated and the ethyl
 acetate phase is washed with saline, dried over sodium sulfate
 10 and concentrated. The concentrate is column chromatographed
 (silica gel; ethyl acetate/hexanes, 15/85) to give the title
 compound, MS (ESI+) for $C_{17}H_{25}NO_3$ m/z $(M+H)^+ = 292.2$.

PREPARATION 21 3-[(Dipropylamino)carbonyl]-5-ethylbenzoic acid
 15 (IX)

Lithium hydroxide monohydrate (0.0680 g, 1.6 mmol) is
 added to a mixture of methyl 3-[(dipropylamino)carbonyl]-5-
 ethylbenzoate (PREPARATION 20, 0.450 g, 1.6 mmol) in a mixture
 of THF/methanol/water (2/1/1, 8 mL). The mixture is stirred at
 20 20-25 degrees C for 3 hours. The mixture is then diluted with
 water (20 mL) and hydrochloric acid (1 N) is added to bring the
 pH to less than 3. The aqueous mixture is extracted with ethyl
 acetate. The organic phase is separated and washed with
 saline, dried over anhydrous magnesium sulfate, filtered and
 25 concentrated to give the title compound, MS (ESI+) for $C_{16}H_{23}NO_3$
 m/z $(M+H)^+ = 278.2$.

EXAMPLE 1 tert-Butyl (1S)-3-bromo-1-(3,5-difluorobenzyl)-
 2-oxopropylcarbamate (III)

30 N-methyl-morpholine (5.83 mL, 53 mmole, 1.05 eq.) is
 added to (2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-
 difluorophenyl)propanoic acid (II, 15 g, 50 mmole) in THF (100
 mL) and the reaction is cooled to -78 degrees C. Isobutyl

chloroformate (6.87 mL, 53 mmole, 1.05 eq.) is added rapidly. The cold bath is then removed and the mixture stirred for 1 hour. The reaction is monitored by TLC to insure completion of the reaction and the mixture is then filtered and washed with dry THF (50 mL) and kept cold in the filtered flask at -20 degrees C.

In an ice-salt bath is placed a 500 mL graduate cylinder containing ether (200 mL) and aqueous potassium hydroxide (40%, 60 mL). 1-Methyl-3-nitro-1-nitrosoguanidine (5.6 g, 106 mmole, 2.1 eq.) is added slowly with stirring and temperature kept below 0 degrees C. The mixture turned yellow and the bubbling lasted for 10 minutes. The stirring is stopped and without mixing the layers, the top diazomethane ethereal layer is transferred with non-ground tip pipette into the stirred mixed anhydride mixture at -20 degrees C. The reaction is monitored by TLC (ethyl acetate/hexane, 50/50; $R_f = 0.69$). After 1 hour nitrogen is then bubbled into the mixture. The solvent is removed under reduced pressure (with heat) and the mixture is partitioned between ether and water. The phases are separated, the organic phase is washed with bicarbonate, saline, dried over anhydrous sodium sulfate and solvent removed under reduced pressure (with heat). The residue is dissolved in ether (100 mL) and hydrobromic acid (48%, 15 mL, 135 mmole, 2.7 eq.) is added at -20 degrees C, the cold bath is removed and the mixture is stirred for another 0.5 hours. The reaction is monitored by TLC (ethyl acetate/hexane, 50/50; $R_f = 0.88$). The mixture is partitioned between ether and water, washed with bicarbonate, saline, dried over anhydrous sodium sulfate and the solvent removed. The residue is recrystallized from ethanol to give the title compound, TLC (ethyl acetate/hexane, 50/50) $R_f = 0.88$; MS (MH^+) = 379.3.

EXAMPLE 2 *tert*-Butyl (1*S*, 2*S*)-3-bromo-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (IV)

Sodium borohydride (1.32 g, 34.9 mmole, 1.1 eq.) is added to *tert*-Butyl (1*S*)-3-bromo-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate (III, EXAMPLE 1, 12 g, 31.75 mmole) dissolved in absolute alcohol (500 mL) at -78 degrees C. The reaction mixture is stirred for 0.5 hour and monitored by TLC (ethyl acetate/hexane, 20/80; R_f = 0.2). The mixture is quenched with water (10 mL) and the solvent removed under reduced pressure with heat (not exceeding 30 degrees C) to dryness. The solid is partitioned between dichloromethane and water, washed with saline, dried over anhydrous sodium sulfate. The solvent is removed under reduced pressure to give the title compound, TLC (ethyl acetate/hexane, 20/80) R_f = 0.2; MS (MH^+) = 381.2.

EXAMPLE 3 *tert*-Butyl (1*S*)-2-(3,5-difluorophenyl)-1-[(2*S*)-oxirany]ethylcarbamate (V)

tert-Butyl (1*S*, 2*S*)-3-bromo-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (IV, EXAMPLE 2) is dissolved in absolute alcohol (150 mL) and ethyl acetate (100 mL) and potassium hydroxide (2.3 g, 34.9 mmole, 1.1eq.) in ethyl alcohol (85%, 5mL) is added at -20 degrees C. The cold bath is then removed and the mixture stirred for 0.5 hour. The reaction is monitored by TLC (ethyl acetate/hexane, 20/80). When the reaction is complete, it is diluted with dichloromethane and extracted, washed with water, saline, dried over anhydrous sodium sulfate and the solvent removed under reduced pressure. The crude material is purified by flash chromatography on silica gel to give the title compound, TLC (ethyl acetate/hexane, 20/80) R_f = 0.3; MS (MH^+) = 300.4.

EXAMPLE 4 tert-Butyl (1S, 2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII)

tert-Butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3, 245 mg, 0.82 mmol) is suspended in isopropyl alcohol (6 mL) and 3-methoxybenzylamine (160 microL, 1.22 mmol) is added with stirring at 20-25 degrees C. This mixture is heated to gentle reflux (bath temp 85 degrees C) under nitrogen for 2 hours, whereupon the resulting mixture is concentrated under reduced pressure to give the title compound. The title compound is purified by flash chromatography (2-5% methanol/methylene chloride; gradient elution) to give purified title compound.

EXAMPLE 5 (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII)

tert-Butyl (1S, 2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, EXAMPLE 4, 258 mg, 0.59 mmol) is dissolved in methylene chloride (1 mL) at 20-25 degrees C, and trifluoroacetic acid (1 mL) is added with stirring under nitrogen. The reaction mixture is stirred at 20-25 degrees C for 1 hour, whereupon the reaction mixture is concentrated under reduced pressure to give the title compound. The title compound is used in the next reaction without further purification.

EXAMPLE 6 N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N³,N³-dipropylisophthalamide (X)

(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate salt (VIII, EXAMPLE 5) is dissolved in anhydrous DMF (3 mL) and cooled to 0

degrees C. Triethylamine (500 microliter, 3.6 mmol) and 5-methyl-*N, N*-dipropylisophthalamide (156 mg, 0.59 mmol) are added with stirring. The mixture is warmed to 20-25 degrees C briefly to allow for complete dissolution of the carboxylic acid, before recooling to 0 degrees C. 1-Hydroxybenzotriazole (157 mg, 1.2 mmol) is added with stirring, followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (229 mg, 1.2 mmol). The resulting mixture is stirred at 0 degrees C for 5 minutes, then warmed to 20-25 degrees C for 15 hours. The reaction mixture is then quenched with aqueous citric acid (10%), and the mixture extracted three times with ethyl acetate. The combined organic extracts are washed with saturated sodium bicarbonate, saline, dried over sodium sulfate, filtered and concentrated under reduced pressure to give the title compound in crude form. This material is purified by flash chromatography (2-10% methanol/methylene chloride gradient elution) to give purified title compound, MS (ES) $MH^+ = 582.3$.

EXAMPLES 7-9

Following the general procedure of EXAMPLE 1 and making non critical variations but starting with the protecting group of Column A and using the acid of Column B, the protected compound (III) of Column C is obtained:

EXAMPLE	Column A	Column B	Column C
7	BOC	Hydrochloric	tert-butyl (1S)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate
8	CBZ	Hydrobromic	benzyl (1S)-3-bromo-1-(3,5-

			difluorobenzyl)-2-oxopropylcarbamate
9	CBZ	Hydrochloride	benzyl (1S)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate

EXAMPLES 10-12

Following the general procedure of EXAMPLE 2 and making non critical variations but starting with the protected compound (III) of Column A, the alcohol (IV) of Column B is obtained:

EXAMPLE	Column A	Column B
10	7	Tert-butyl (1S, 2S)-3-chloro-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate
11	8	Benzyl (1S, 2S)-3-bromo-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate
12	9	Benzyl (1S, 2S)-3-chloro-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate

EXAMPLE 13 Benzyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V)

Following the general procedure of EXAMPLE 3 and making non critical variations but starting with the alcohol (IV) of EXAMPLE 12, the title compound is obtained.

EXAMPLES 14-107

Following the general procedure of EXAMPLE 4 and making non-critical variations but reacting tert-butyl (1S,2S)-1-(2-oxiranyl)-2-phenylethylcarbamate (V, commercially available) with the C-terminal amine (VI) of Column A, the protected alcohol (VII) of Column B is obtained.

EXA	Column A C-terminal amine (VI)	Column B Protected alcohol (VII)
14	$\text{H}_2\text{N}-\text{CH}_2\text{CH}_3$	tert-butyl (1S,2R) - 1-benzyl-3- (ethylamino)-2- hydroxypropylcarbama te
15	$\text{H}_2\text{N}-\text{CH}_2$ -phenyl	tert-butyl (1S,2R) - 1-benzyl-3- (benzylamino)-2- hydroxypropylcarbama te
16	$\text{H}_2\text{N}-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R) - 1-benzyl-3- (isopropylamino)-2- hydroxypropylcarbama te
17	$\text{H}_2\text{N}-\text{CH}_2$ -phenyl-4- CH_3	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(4- methylbenzyl) amino]p ropylcarbamate
18	$\text{H}_2\text{N}-(\text{CH}_2)_2$ -phenyl-4- OCH_3	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{[2-(4- methoxyphenyl) ethyl] amino}propylcarbamat e
19	$\text{H}_2\text{N}-\text{CH}_2$ -phenyl-3- OCH_3	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(3-

		methoxybenzyl) amino] propylcarbamate
20	$\text{H}_2\text{N}-\text{CH}(-\text{phenyl})-\text{CO}-\text{OC}_2\text{H}_5$	ethyl ({(2R,3S)-3- [(tert- butoxycarbonyl) amino]-2-hydroxy-4- phenylbutyl} amino) (p henyl) acetate
21	$\text{H}_2\text{N}-(\text{CH}_2)_2-\text{phenyl}$	tert-butyl (1S,2R)- 1-benzyl-2-hydroxy- 3-[(2- phenylethyl) amino]pr opylcarbamate
22	$\text{H}_2\text{N}-\text{CH}(-\text{CH}_2\text{OH})-\text{CH}(\text{OH})-\text{phenyl}-4-\text{NO}_2$	tert-butyl (1S,2R)- 1-benzyl-2-hydroxy- 3-[(1S)-2-hydroxy- 1-(hydroxymethyl)-2- (4- nitrophenyl) ethyl] am ino}propylcarbamate
23	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-2-\text{Cl}$	tert-butyl (1S,2R)- 1-benzyl-3-[(2- chlorobenzyl) amino]- 2- hydroxypropylcarbama te
24	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-4-\text{Cl}$	tert-butyl (1S,2R)- 1-benzyl-3-[(4- chlorobenzyl) amino]- 2- hydroxypropylcarbama te

25	$\text{H}_2\text{N}-(\text{CH}_2)_2-\text{O}-(\text{CH}_2)_2-\text{OH}$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{ [2-(2- hydroxyethoxy) ethyl] amino}propylcarbamate
26	H_2N -1-indanyl	tert-butyl (1S,2R) - 1-benzyl-3-(2,3- dihydro-1H-inden-1- ylamino)-2- hydroxypropylcarbama te
27	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_3$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(2- hydroxypropyl) amino] propylcarbamate
28	$\text{H}_2\text{N}-\text{CH}_2$ - tetrahydrofuranyl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(tetrahydro-2- furanylmethyl) amino] propylcarbamate
29	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}(-\text{OCH}_2\text{CH}_3)$	tert-butyl (1S,2R) - 1-benzyl-3-[(2,2- diethoxyethyl) amino] -2- hydroxypropylcarbama te
30	$\text{H}_2\text{N}-(\text{CH}_2)_4-\text{CH}_3$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-(pentylamino) propylcarbamate

31	H ₂ N-cyclohexyl	tert-butyl (1S,2R) - 1-benzyl-3- (cyclohexylamino)-2- hydroxypropylcarbama te
32	H ₂ N-CH ₂ -pyridin-2-yl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(2- pyridinylmethyl)amin ol]propylcarbamate
33	H ₂ N-CH ₂ -phenyl-2-NH ₂	tert-butyl (1S,2R) - 3-[(2- aminobenzyl) amino] - 1-benzyl-2- hydroxypropylcarbama te
34	H ₂ N-CH ₂ -pyridin-3-yl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(3- pyridinylmethyl)amin ol]propylcarbamate
35	H ₂ N-(CH ₂) ₂ - pyrrolidin-1-yl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{[2-(1- pyrrolidinyl)ethyl]a mino}propylcarbamate
36	H ₂ N-CH ₂ -CH(OH) - phenyl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(2-hydroxy-2- phenylethyl) amino]pr opylcarbamate
37	H ₂ N-(CH ₂) ₃ -O-(CH ₂) ₃ -	tert-butyl (1S,2R) -

	CH ₃	1-benzyl-3-[(3-butoxypropyl)amino]-2-hydroxypropylcarbamate
38	H ₂ N-(CH ₂) ₃ -O-CH(CH ₃) ₂	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(3-isopropoxypropyl)aminolpropylcarbamate
39	H ₂ N-(CH ₂) ₂ -CH(CH ₃) ₂	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-(isopentylamino)propylcarbamate
40	H ₂ N-(CH ₂) ₃ -phenyl	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(3-phenylpropyl)aminolpropylcarbamate
41	H ₂ N-(CH ₂) ₂ -OCH ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(2-methoxyethyl)aminolpropylcarbamate
42	H ₂ N-(CH ₂) ₂ -O-phenyl	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(2-phenoxyethyl)aminolpropylcarbamate
43	H ₂ N-(CH ₂) ₂ -O-(CH ₂) ₂ -CH ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(2-

		propoxyethyl) amino] p ropylcarbamate
44	$\text{H}_2\text{N}-(\text{CH}_2)_2-\text{C}(\text{CH}_3)_3$	tert-butyl (1S,2R) - 1-benzyl-3-[(3,3- dimethylbutyl) amino] -2- hydroxypropylcarbama te
45	$\text{H}_2\text{N}-(\text{CH}_2)_4\text{-phenyl}$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(4- phenylbutyl) amino] pr opylcarbamate
46	$\text{H}_2\text{N}-\text{CH}_2\text{-phenyl-3-I}$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(3- iodobenzyl) amino] pro pylcarbamate
47	$\text{H}_2\text{N}-\text{CH}_2\text{-phenyl-4-NO}_2$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(4- nitrobenzyl) amino] pr opylcarbamate
48	$\text{H}_2\text{N}-\text{CH}_2\text{-phenyl-3-Cl}$	tert-butyl (1S,2R) - 1-benzyl-3-[(3- chlorobenzyl) amino] - 2- hydroxypropylcarbama te
49	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-4-Cl}$	tert-butyl (1S,2R) - 1-benzyl-3-[[2-(4- chlorophenyl) ethyl] a

		mino}-2-hydroxypropylcarbamate
50	H ₂ N-(CH ₂) ₂ -pyridin-2-yl	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[2-(2-pyridinyl)ethyl]amino}propylcarbamate
51	H ₂ N-CH ₂ -pyridin-4-yl	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{(4-pyridinylmethyl)amino}propylcarbamate
52	H ₂ N-(CH ₂) ₂ -(N-methylpyrrolidin-2-yl)	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[2-(1-methyl-2-pyrrolidinyl)ethyl]amino}propylcarbamate
53	H ₂ N-CH ₂ -phenyl-2,3-dimethyl	tert-butyl (1S,2R)-1-benzyl-3-[(2,3-dimethylbenzyl)amino]-2-hydroxypropylcarbamate
54	H ₂ N-CH ₂ -phenyl-2-OCF ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[2-(trifluoromethoxy)benzyl]amino}propylcarbamate
55	H ₂ N-CH ₂ -phenyl-2-Cl-6-O-phenyl	tert-butyl (1S,2R)-1-benzyl-3-[(2-

		chloro-6- phenoxybenzyl) amino] -2- hydroxypropylcarbama te
56	H ₂ N-CH ₂ -phenyl-4-CF ₃	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{ [4- (trifluoromethyl)ben zyl] amino} propylcarb amate
57	H ₂ N-CH ₂ -phenyl-2,3- dichloro	tert-butyl (1S,2R) - 1-benzyl-3-[(2,3- dichlorobenzyl) amino]-2- hydroxypropylcarbama te
58	H ₂ N-CH ₂ -phenyl-3,5- dichloro	tert-butyl (1S,2R) - 1-benzyl-3-[(3,5- dichlorobenzyl) amino]-2- hydroxypropylcarbama te
59	H ₂ N-CH ₂ -phenyl-3,5- difluoro	tert-butyl (1S,2R) - 1-benzyl-3-[(3,5- difluorobenzyl) amino]-2- hydroxypropylcarbama te
60	H ₂ N-CH ₂ -phenyl-4- OCF ₃	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{ [4-

		(trifluoromethoxy)benzyl]amino}propylcarbamate
61	H ₂ N-(CH ₂) ₂ -phenyl-4-SO ₂ -NH ₂	tert-butyl (1S,2R)-3-{[4-(aminosulfonyl)benzyl]amino}-1-benzyl-2-hydroxypropylcarbamate
62	H ₂ N-CH ₂ -phenyl-4-OCH ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(4-methoxybenzyl)amino]propylcarbamate
63	H ₂ N-CH ₂ -phenyl-4-CH ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(4-methylbenzyl)amino]propylcarbamate
64	H ₂ N-CH ₂ -Ph-(3,4,5-trimethoxy)	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(3,4,5-trimethoxybenzyl)amino]propylcarbamate
65	H ₂ N-CH ₂ -phenyl-3-OCF ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[3-(trifluoromethoxy)benzyl]amino}propylcarbamate
66	H ₂ N-CH ₂ -phenyl-3,5-dimethoxy	tert-butyl (1S,2R)-1-benzyl-3-[(3,5-

		dimethoxybenzyl) amino]-2-hydroxypropylcarbamate
67	H ₂ N-CH ₂ -phenyl-2,4-dimethoxy	tert-butyl (1S,2R)-1-benzyl-3-[(2,4-dimethoxybenzyl) amino]-2-hydroxypropylcarbamate
68	H ₂ N-CH ₂ -phenyl-phenyl	tert-butyl (1S,2R)-1-benzyl-3-[[[1,1'-biphenyl]-3-ylmethyl) amino]-2-hydroxypropylcarbamate
69	H ₂ N-CH ₂ -phenyl-3,4-dichloro	tert-butyl (1S,2R)-1-benzyl-3-[(3,4-dichlorobenzyl) amino]-2-hydroxypropylcarbamate
70	H ₂ N-CH ₂ -phenyl-4-F	tert-butyl (1S,2R)-1-benzyl-3-[(4-fluorobenzyl) amino]-2-hydroxypropylcarbamate
71	H ₂ N-CH ₂ -phenyl-3-CF ₃	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[3-(trifluoromethyl)ben

		zyl]amino}propylcarbamate
72	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-2-\text{CH}_3$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-[(2-methylbenzyl)amino]propylcarbamate
73	$\text{H}_2\text{N}-\text{CH}((\text{R})-\text{CH}_3)-\text{phenyl}$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-{[(1R)-1-phenylethyl]amino}propylcarbamate
74	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_3)-\text{phenyl}$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-{[(1S)-1-phenylethyl]amino}propylcarbamate
75	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-3,5-(\text{bis})\text{trifluoromethyl}$	tert-butyl (1S,2R) - 1-benzyl-3-{[3,5-bis(trifluoromethyl)benzyl]amino}-2-hydroxypropylcarbamate
76	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-2-\text{CF}_3$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-{[2-(trifluoromethyl)benzyl]amino}propylcarbamate
77	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_3)-(\text{naphth-1-yl})$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-

		3-{[(1S)-1-(1-naphthyl)ethyl]amino}propyl carbamate
78	$\text{-NH}_2\text{-CH}((R)\text{-CH}_3)\text{-(naphth-1-yl)}$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[(1R)-1-(1-naphthyl)ethyl]amino}propylcarbamate
79	$\text{H}_2\text{N-CH}_2\text{-phenyl-3-OCH}_3\text{-4-OH}$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(4-hydroxy-3-methoxybenzyl)amino]propylcarbamate
80	$\text{H}_2\text{N-CH}_2\text{-phenyl-3,4-dihydroxy}$	tert-butyl (1S,2R)-1-benzyl-3-[(3,4-dihydroxybenzyl)amino]-2-hydroxypropylcarbamate
81	$\text{H}_2\text{N-(CH}_2)_3\text{-OCH}_3$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxypropyl)amino]propylcarbamate
82	$\text{H}_2\text{N-CH}((S)\text{-CH}_3)\text{-CH}_2\text{-OH}$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[(1S)-2-hydroxy-1-methylethyl]amino}propyl carbamate
83	$\text{H}_2\text{N-CH}((R)\text{-CH}_3)\text{-CH}_2\text{-OH}$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-

		3-{[(1R)-2-hydroxy-1-methylethyl]amino}propyl carbamate
84	$\text{H}_2\text{N}-\text{CH}_2-\text{C}\equiv\text{CH}$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-(2-propynylamino)propylcarbamate
85	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-2-F}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(2-fluorophenyl)ethyl]amino}-2-hydroxypropylcarbamate
86	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-3-F}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(3-fluorophenyl)ethyl]amino}-2-hydroxypropyl carbamate
87	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-4-F}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(4-fluorophenyl)ethyl]amino}-2-hydroxypropyl carbamate
88	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-4-Br}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(4-bromophenyl)ethyl]amino}-2-hydroxypropyl carbamate

89	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-3-OCH}_3$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[2-(3-methoxyphenyl)ethyl]amino}propylcarbamate
90	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-2,4-dichloro}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(2,4-dichlorophenyl)ethyl]amino}-2-hydroxypropylcarbamate
91	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-3-Cl}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(3-chlorophenyl)ethyl]amino}-2-hydroxypropylcarbamate
92	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-2,5-dimethoxy}$	tert-butyl (1S,2R)-1-benzyl-3-{[2-(2,5-dimethoxyphenyl)ethyl]amino}-2-hydroxypropylcarbamate
93	$\text{H}_2\text{N}-(\text{CH}_2)_2\text{-phenyl-4-CH}_3$	tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[2-(4-methylphenyl)ethyl]amino}propylcarbamate
94	$\text{H}_2\text{N-CH}(-(\text{R})\text{CH}_2\text{-OH})\text{-CH}_2\text{-phenyl}$	tert-butyl (1S,2R)-1-benzyl-3-{[(1R)-1-benzyl-2-

		hydroxyethyl] amino} - 2- hydroxypropylcarbama te
95	$H_2N-(CH_2)_3-(1-$ morpholinyl)	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{ [3-(4- morpholinyl)propyl] a mino}propylcarbamate
96	$H_2N-CH_2-C(CH_3)_2$	tert-butyl (1S,2R) - 1-benzyl-3-[(3,3- dimethylbutyl) amino] -2- hydroxypropylcarbama te
97	$H_2N-(CH_2)_2-(1-$ morpholinyl)	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-{ [2-(4- morpholinyl)ethyl] am ino}propylcarbamate
98	$H_2N-CH(OH)-CH_2-CH_3$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(1- hydroxypropyl) amino] propylcarbamate
99	$H_2N-(CH_2)_2-(thien-2-$ yl)	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy- 3-[(2- thienylmethyl) amino] propylcarbamate
100	$H_2N-(CH_2)_4-OH$	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-

		3-[(4-hydroxybutyl)amino]propylcarbamate
101	H ₂ N-CH(- (S) CH ₂ -OH) - phenyl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-{ [(1S)-2-hydroxy-1-phenylethyl] amino} propylcarbamate
102	H ₂ N-CH ₂ -phenyl-2,4-dichloro	tert-butyl (1S,2R) - 1-benzyl-3-[(2,4-dichlorobenzyl) amino] - 2-hydroxypropylcarbamate
103	H ₂ N-CH(- (R) CH ₂ -OH) - phenyl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-{ [(1R)-2-hydroxy-1-phenylethyl] amino} propylcarbamate
104	H ₂ N-CH ₂ -phenyl-4-C(CH ₃) ₃	tert-butyl (1S,2R) - 1-benzyl-3-[(4-tert-butylbenzyl) amino] - 2-hydroxypropylcarbamate
105	H ₂ N-CH(CH ₃) - phenyl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-3-[(1-phenylethyl) amino]propylcarbamate
106	H ₂ N- (1R,2S) - 2-hydroxyinden-1-yl	tert-butyl (1S,2R) - 1-benzyl-2-hydroxy-

		3-{[(1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino}propylcarbamate
107	H ₂ N-CH ₂ -phenyl-3,4-dimethyl	tert-butyl (1S,2R)-1-benzyl-3-[(3,4-dimethylbenzyl)amino]-2-hydroxypropylcarbamate

EXAMPLES 108-164

Following the general procedure of EXAMPLE 4 and making non-critical variations but reacting tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3) with the C-terminal amine (VI) of Column A, the protected alcohol (VII) of Column B is obtained.

EXA	Column A C-terminal amine (VI)	Column B Protected alcohol (VII)
108	H ₂ N-(CH ₂) ₆ -CO-O-CH ₃	methyl 7-{[(2R,3S)-3-[(tert-butoxycarbonyl)amino]-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}heptanoate
109	H ₂ N-CH(-CH ₃)-CO-NH-CH ₂ -CH(CH ₃) ₂ r/s	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-

		hydroxy-3-{ [2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl carbamate
110	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_3)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [(1S)-2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl carbamate
111	$\text{H}_2\text{N}-\text{C}(-\text{CH}_3)_2-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [2-(isobutylamino)-1,1-dimethyl-2-oxoethyl]amino}propyl carbamate
112	$\text{H}_2\text{N}-\text{CH}_2-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [2-(isobutylamino)-2-oxoethyl]amino}propyl carbamate
113	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2\text{CH}_3)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({ (1S)-1-

		[(isobutylamino) carbonyl]propyl}amino)propylcarbamate
114	$\text{H}_2\text{N}-\text{CH}((\text{R})-\text{CH}_2\text{CH}_3)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino) carbonyl]propyl}amino)propylcarbamate
115	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}$	tert-butyl (1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate
116	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}_3$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-(ethylamino)-2-hydroxypropylcarbamate
117	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isobutylamino)propylcarbamate
118	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{CONH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-

		(isobutylamino)-2-methyl-3-oxopropylamino}propylcarbamate
119	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-4-\text{N}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-{[4-(dimethylamino)benzyl]amino}-2-hydroxypropylcarbamate
120	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2-\text{phenyl})-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-3-{[(1S)-1-(3,5-difluorobenzyl)-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate
121	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}(\text{CH}_3)_2)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]-3-methylbutyl}amino)propylcarbamate
122	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-

		{ [2-(dimethylamino)ethyl]amino}-2-hydroxypropylcarbamate
123	$\text{H}_2\text{N}-\text{CH}_2-(\text{pyridin-3-yl})$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-pyridinylmethyl)aminol]propylcarbamate
124	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2-\text{O}-\text{CH}_2-\text{phenyl})-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-3-{[(1S)-1-[(benzyloxy)methyl]-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate
125	$\text{H}_2\text{N}-\text{C}(-\text{CH}_3)_2-\text{phenyl}$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)aminol]propylcarbamate
126	$\text{H}_2\text{N}-\text{CH}((\text{R})-\text{CH}(\text{CH}_3)_2)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino)carb

		onyl]-3-methylbutyl}amino)propylcarbamate
127	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2-\text{CH}_2-\text{CH}_3)-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carboxonyl]butyl}amino)propylcarbamate
128	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2-\text{OH})-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-(hydroxymethyl)-2-(isobutylamino)-2-oxoethyl}amino)propylcarbamate
129	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{phenyl}$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-phenylethyl)amino]propylcarbamate
130	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_3)-\text{CO}-\text{NH}-\text{CH}_2-\text{phenyl}$	tert-butyl (1S,2R)-3-{[2-(benzylamino)-1-methyl-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate

131	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{CH}_2-\text{CH}_3)-\text{phenyl}$	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-3- {[(1S)-2- (benzylamino)-1- methyl-2- oxoethyl]amino}-2- hydroxypropylcarbama te
132	$\text{H}_2\text{N}-\text{CH}_2-\text{phenyl}-3-\text{OCH}_3$	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(3- methoxybenzyl)amino] propylcarbamate
133	$\text{H}_2\text{N}-\text{CH}((\text{S})-\text{phenyl})\text{CO}-\text{NHCH}_2\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-{[(1S)-2- (isobutylamino)-2- oxo-1- phenylethyl]amino}pr opylcarbamate
134	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{CH}(\text{CH}_3)_2$	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3- (isopentylamino)prop ylcarbamate
135	$\text{H}_2\text{N}-\text{cyclohexyl}$	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-3-

		(cyclohexylamino)-2-hydroxypropylcarbamate
136	$\text{H}_2\text{N}-(\text{CH}_2)_3-\text{CH}_3$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-(butylamino)-2-hydroxypropylcarbamate
137	$\text{H}_2\text{N}-(\text{CH}_2)_3-\text{O}-\text{CH}_3$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxypropyl)amino]propylcarbamate
138	$\text{H}_2\text{N}-\text{CH}_2-\text{CH}(\text{OH})-\text{phenyl}$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-hydroxy-2-phenylethyl)amino]propylcarbamate
139	$\text{H}_2\text{N}-\text{cyclohexyl}-3,5\text{-dimethoxy}$	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3R,5S)-3,5-dimethoxycyclohexyl]amino}-2-hydroxypropylcarbamate
140	$\text{H}_2\text{N}-\text{cyclohexyl}-3,5\text{-di-}(-\text{CO}-\text{OCH}_3)$	dimethyl (1R,3S)-5-({(2R,3S)-3-[(tert-

		butoxycarbonyl) amino]-2-hydroxy-4- phenylbutyl} amino) - 1,3- cyclohexanedicarboxy late
141	H ₂ N-cyclohexyl-3,5- di-(-COOH)	(1R,3S)-5-({(2R,3S)- 3-[(tert- butoxycarbonyl) amino]-2-hydroxy-4- phenylbutyl} amino) - 1,3- cyclohexanedicarboxy lic acid
142	H ₂ N-CH((R)-CH ₂ -CH ₃)- phenyl	tert-butyl (1S,2R)- 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(1R)-1- phenylpropyl] amino)p ropylcarbamate
143	H ₂ N-CH ₂ -phenyl-3-Cl	tert-butyl (1S,2R)- 1-(3,5- difluorobenzyl)-3- [(3- chlorobenzyl) amino]- 2- hydroxypropylcarbama te
144	H ₂ N-CH ₂ -phenyl-3- OCH ₃	tert-butyl (1S,2R)- 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(3-

		methoxybenzyl) amino] propylcarbamate
145	H ₂ N-CH ₂ -phenyl- phenyl	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-3- [[[1,1'-biphenyl]-3- ylmethyl) amino]-2- hydroxypropylcarbama te
146	H ₂ N-CH ₂ -phenyl-3-I	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(3- iodobenzyl) amino]pro pylcarbamate
147	H ₂ N-CH ₂ -phenyl-3-CH ₃	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(3- methylbenzyl) amino]p ropylcarbamate
148	H ₂ N-CH ₂ -CH(-CH ₃)- phenyl	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(2- phenylpropyl) amino]p ropylcarbamate
149	H ₂ N-CH ₂ -(thiazol-5- yl)	tert-butyl (1S,2R) - 1-(3,5- difluorobenzyl)-2- hydroxy-3-[(1,3- thiazol-5-

		ylmethyl) amino] propyl carbamate
150	H ₂ N-CH ₂ -(thien-2-yl)	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-thienylmethyl) amino] propyl carbamate
151	H ₂ N-4-methoxytetralin-1-yl	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl) amino] propyl carbamate
152	H ₂ N-CH ₂ -pyrazin-2-yl	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-pyrazinylmethyl) amino] propyl carbamate
153	H ₂ N-CH ₂ -phenyl-3,5-difluoro	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-[(3,5-difluorobenzyl) amino]-2-hydroxypropyl carbamate
154	H ₂ N-CH ₂ -phenyl-3,4-methylenedioxy	tert-butyl (1S,2R)-3-[(1,3-benzodioxol-

		5-ylmethyl) amino] -1-(3,5-difluorobenzyl) -2-hydroxypropylcarbamate
155	H ₂ N-CH ₂ -phenyl-3,5-dimethoxy	tert-butyl (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3,5-dimethoxybenzyl) amino] -2-hydroxypropylcarbamate
156	H ₂ N-CH ₂ -phenyl-3-CF ₃	tert-butyl (1S,2R) -1-(3,5-difluorobenzyl) -2-hydroxy-3-{[3-(trifluoromethyl) benzyl] amino} propylcarbamate
157	H ₂ N-CH ₂ -(furan-2-yl)	tert-butyl (1S,2R) -1-(3,5-difluorobenzyl) -3-[(2-furylmethyl) amino] -2-hydroxypropylcarbamate
158	H ₂ N-(7-methoxytetralin-1-yl)	tert-butyl (1S,2R) -1-(3,5-difluorobenzyl) -2-hydroxy-3-[(7-

		methoxy-1,2,3,4-tetrahydro-1-naphthalenyl) amino]propylcarbamate
159	H ₂ N-CH ₂ -phenyl-3-O-CF ₃	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(trifluoromethoxy)benzyl] amino}propylcarbamate
160	H ₂ N-CH ₂ -phenyl-3-F	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-fluorobenzyl) amino]-2-hydroxypropylcarbamate
161	H ₂ N-CH ₂ -phenyl-3-O-CH(CH ₃) ₂	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropoxybenzyl) amino]propylcarbamate
162	H ₂ N-CH ₂ -phenyl-3-Br	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-bromobenzyl) amino]-2-hydroxypropylcarbamate

		te
163	H ₂ N-CH ₂ -(5-methylfuran-2-yl)	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-methyl-2-furyl)methyl]amino}propylcarbamate
164	H ₂ N-(5-methoxytetralin-1-yl)	tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propylcarbamate

EXAMPLE 165 tert-Butyl-(1S, 2R)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (XII)

Sodium azide (0.22 g, 4 mmole) and ammonium chloride (2
5 eq) are added to tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3, 0.6 g, 2 mmole). The reaction is heated to 75-80 degrees C and stirred for 16 hours. The reaction is monitored by TLC to insure completion. The solvent is removed under reduced pressure. The concentrate
10 is partitioned between ethyl acetate and water, the phases are separated and the organic phase is washed with bicarbonate and saline, dried over anhydrous sodium sulfate and concentrated to give the title compound, TLC (ethyl acetate/hexane) R_f = 0.45; MS (MH⁺) = 343.

15

EXAMPLE 166 (2R, 3S)-3-amino-1-azido-4-(3,5-difluorophenyl)-2-butanol (XIV)

tert-Butyl-(1S, 2R)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (XII, EXAMPLE 165, 0.48 g, 1.41 mmole) is dissolved in dichloromethane (20 ml) to which trifluoroacetic acid (5 ml) is added. The reaction is stirred at 20-25 degrees C for 16 hours and the solvent is removed under reduced pressure with heat. Ethyl acetate is added twice and evaporated twice to give the title compound as the trifluoroacetic acid salt which is used in the next reaction without further purification; MS (MH⁺) = 242.

10

EXAMPLE 167 N¹-[(1S,2R)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N³,N³-dipropylisophthalamide (XV)

To (2R, 3S)-3-amino-1-azido-4-(3,5-difluorophenyl)-2-butanol (XIV, EXAMPLE 166, 0.34 g, 1.4 mmole) in dichloromethane (20 ml) is added N,N-dipropylamidooisophthalic acid (IX, 0.53 g, 2 mmole), t-butyl alcohol (0.27 g, 2 mmole) and triethylamine (0.84 ml, 6 mmole) and ethyl-1-(3-dimethylaminopropyl)carbodiimide (0.58 g, 3 mmole). The mixture is stirred at 20-25 degrees C for 16 hours. The reaction is monitored by TLC (methanol/dichloromethane, 20/80 + ethyl acetate/hexane, 50/50; R_f = 0.76). When the reaction is complete as measured by TLC, the reaction mixture is partitioned between dichloromethane and water, washed with hydrochloric acid (0.5 N), bicarbonate, saline, dried over anhydrous sodium sulfate and the solvent is removed under reduced pressure with heat to produce a concentrate. The concentrate is column chromatographed on silica gel to give the title compound; MS (MH⁺) = 488.

30

EXAMPLE 168 N¹-[(1S,2R)-3-amino-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N³,N³-dipropylisophthalamide acetic acid salt (XVI)

N^1 -[(1S,2R)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- N^3,N^3 -dipropylisophthalamide (XV, EXAMPLE 167, 0.3 g, 0.62 mmole) in ethyl acetate (20 ml) and acetic acid (5ml) is placed in a Parr pressure bottle. Palladium on carbon (10%, 5 g) is added and the mixture shaken under hydrogen at 50 psi for 2 hours. The mixture is filtered through a diatomaceous earth and the filtrate is concentrated to give the title compound; MS (MH^+) = 462.

10 EXAMPLE 169 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(2-furylmethyl)amino]-2-hydroxypropyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

N^1 -[(1S,2R)-3-amino-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- N^3,N^3 -dipropylisophthalamide acetic acid salt (XVI, EXAMPLE 168, 76 mg, 0.146 mmol) is dissolved in absolute ethanol (2 mL). 3-Furaldehyde (20 microL, 0.231 mmol) and triethylamine (30 microL, 0.215 mmol) are added via syringe, with stirring at 20-25 degrees C. After 10 minutes, palladium on carbon 122 mg, 5 weight %) is added and the mixture placed under a hydrogen atmosphere (50 psi) and shaken for 20 minutes. The resulting mixture is then filtered through diatomaceous earth, with ethanol washings. The filtrate is purified by flash chromatography (2-10% methanol/methylene chloride) to give purified title compound, MS (MH^+) = 542.2.

25 EXAMPLE 169a tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-3-[(1S)-2-(ethylamino)-1-methyl-2-oxoethyl]amino}-2-hydroxypropylcarbamate (VII)

30 Following the general procedure of EXAMPLES 4 and 14-164 and making non-critical variations and reacting tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V,

EXAMPLE 3) with (2S)-2-amino-N-ethylpropanamide (VI), the title compound is obtained.

EXAMPLES 170-320

- 5 Following the general procedure of EXAMPLE 5 and making non-critical variations but starting with the protected alcohol (VII) of Column A, the amine (VIII) of Column B is obtained.

EXA	Column A Protecte d Alcohol (VII) Compound of EXAMPLE	Column B Amine (VIII)
170	14	(2R,3S)-3-amino-1-(ethylamino)-4-phenyl-2-butanol
171	15	(2R,3S)-3-amino-1-(benzylamino)-4-phenyl-2-butanol
172	16	(2R,3S)-3-amino-1-(isopropylamino)-4-phenyl-2-butanol
173	17	(2R,3S)-3-amino-1-[(4-methylbenzyl)amino]-4-phenyl-2-butanol
174	18	(2R,3S)-3-amino-1-{[2-(4-methoxyphenyl)ethyl]amino}-4-phenyl-2-butanol
175	19	(2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol
176	20	ethyl {[(2R,3S)-3-amino-2-

		hydroxy-4-phenylbutyl]amino}(phenyl)acetate
177	21	(2R,3S)-3-amino-4-phenyl-1-[(2-phenylethyl)amino]-2-butanol
178	22	(2S)-2-{[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]amino}-1-(4-nitorphenyl)-1,3-propanediol
179	23	(2R,3S)-3-amino-1-[(2-chlorobenzyl)amino]-4-phenyl-2-butanol
180	24	(2R,3S)-3-amino-1-[(4-chlorobenzyl)amino]-4-phenyl-2-butanol
181	25	(2R,3S)-3-amino-1-{[2-(2-hydroxyethoxy)ethyl]amino}-4-phenyl-2-butanol
182	26	(2R,3S)-3-amino-1-(2,3-dihydro-1H-inden-1-ylamino)-4-phenyl-2-butanol
183	27	(2R,3S)-3-amino-1-[(2-hydroxypropyl)amino]-4-phenyl-2-butanol
184	28	(2R,3S)-3-amino-4-phenyl-1-[(tetrahydro-2-furanylmethyl)amino]-2-butanol
185	29	(2R,3S)-3-amino-1-[(2,2-diethoxyethyl)amino]-4-phenyl-2-butanol
186	30	(2R,3S)-3-amino-1-(butylamino)-4-phenyl-2-butanol
187	31	(2R,3S)-3-amino-1-

	-	(cyclohexylamino)-4-phenyl-2-butanol
188	32	(2R,3S)-3-amino-4-phenyl-1-[(2-pyridinylmethyl)amino]-2-butanol
189	33	(2R,3S)-3-amino-1-[(2-aminobenzyl)amino]-4-phenyl-2-butanol
190	34	(2R,3S)-3-amino-4-phenyl-1-[(3-pyridinylmethyl)amino]-2-butanol
191	35	(2R,3S)-3-amino-4-phenyl-1-{[2-(1-pyrrolidinyl)ethyl]amino}-2-butanol
192	36	(2R,3S)-3-amino-1-[(2-hydroxy-2-phenylethyl)amino]-4-phenyl-2-butanol
193	37	(2R,3S)-3-amino-1-[(3-butoxypropyl)amino]-4-phenyl-2-butanol
194	38	(2R,3S)-3-amino-1-[(3-isopropoxypropyl)amino]-4-phenyl-2-butanol
195	39	(2R,3S)-3-amino-1-(isopentylamino)-4-phenyl-2-butanol
196	40	(2R,3S)-3-amino-4-phenyl-1-[(3-phenylpropyl)amino]-2-butanol
197	41	(2R,3S)-3-amino-1-[(2-methoxyethyl)amino]-4-phenyl-2-butanol
198	42	(2R,3S)-3-amino-1-[(2-phenoxyethyl)amino]-4-phenyl-2-butanol

199	43	(2R,3S)-3-amino-4-phenyl-1-[(2-propoxyethyl)amino]-2-butanol
200	44	(2R,3S)-3-amino-1-[(3,3-dimethylbutyl)amino]-4-phenyl-2-butanol
201	45	(2R,3S)-3-amino-4-phenyl-1-[(4-phenylbutyl)amino]-2-butanol
202	46	(2R,3S)-3-amino-1-[(3-iodobenzyl)amino]-4-phenyl-2-butanol
203	47	(2R,3S)-3-amino-1-[(4-nitrobenzyl)amino]-4-phenyl-2-butanol
204	48	(2R,3S)-3-amino-1-[(3-chlorobenzyl)amino]-4-phenyl-2-butanol
205	49	(2R,3S)-3-amino-1-{[2-(4-chlorophenyl)ethyl]amino}-4-phenyl-2-butanol
206	50	(2R,3S)-3-amino-4-phenyl-1-{[2-(2-pyridinyl)ethyl]amino}-2-butanol
207	51	(2R,3S)-3-amino-4-phenyl-1-[(4-pyridinylmethyl)amino]-2-butanol
208	52	(2R,3S)-3-amino-1-{[2-(1-methyl-2-pyrrolidinyl)ethyl]amino}-4-phenyl-2-butanol
209	53	(2R,3S)-3-amino-1-[(2,3-dimethylbenzyl)amino]-4-phenyl-2-butanol
210	54	(2R,3S)-3-amino-4-phenyl-1-{[2-(trifluoromethoxy)benzyl]amino}-

		2-butanol
211	55	(2R,3S)-3-amino-1-[(2-chloro-6-phenoxybenzyl)amino]-4-phenyl-2-butanol
212	56	(2R,3S)-3-amino-4-phenyl-1-{[4-(trifluoromethyl)benzyl]amino}-2-butanol
213	57	(2R,3S)-3-amino-1-[(2,3-dichlorobenzyl)amino]-4-phenyl-2-butanol
214	58	(2R,3S)-3-amino-1-[(3,5-dichlorobenzyl)amino]-4-phenyl-2-butanol
215	59	(2R,3S)-3-amino-1-[(3,5-difluorobenzyl)amino]-4-phenyl-2-butanol
216	60	(2R,3S)-3-amino-4-phenyl-1-{[4-(trifluoromethoxy)benzyl]amino}-2-butanol
217	61	4-({[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]amino}methyl)benzenesulfonamide
218	62	(2R,3S)-3-amino-1-[(4-methoxybenzyl)amino]-4-phenyl-2-butanol
219	63	(2R,3S)-3-amino-1-[(4-methylbenzyl)amino]-4-phenyl-2-butanol
220	64	(2R,3S)-3-amino-4-phenyl-1-[(3,4,5-trimethoxybenzyl)amino]-2-butanol

221	65	(2R,3S)-3-amino-4-phenyl-1-{[3-(trifluoromethoxy)benzyl]amino}-2-butanol
222	66	(2R,3S)-3-amino-1-[(3,5-dimethoxybenzyl)amino]-4-phenyl-2-butanol
223	67	(2R,3S)-3-amino-1-[(2,4-dimethoxybenzyl)amino]-4-phenyl-2-butanol
224	68	(2R,3S)-3-amino-1-[[[1,1'-biphenyl]-3-ylmethyl]amino]-4-phenyl-2-butanol
225	69	(2R,3S)-3-amino-1-[(3,4-dichlorobenzyl)amino]-4-phenyl-2-butanol
226	70	(2R,3S)-3-amino-1-[(2-fluorobenzyl)amino]-4-phenyl-2-butanol
227	71	(2R,3S)-3-amino-4-phenyl-1-{[3-(trifluoromethyl)benzyl]amino}-2-butanol
228	72	(2R,3S)-3-amino-1-[(2-methylbenzyl)amino]-4-phenyl-2-butanol
229	73	(2R,3S)-3-amino-4-phenyl-1-{[(1R)-1-phenylethyl]amino}-2-butanol
230	74	(2R,3S)-3-amino-4-phenyl-1-{[(1S)-1-phenylethyl]amino}-2-butanol
231	75	(2R,3S)-3-amino-1-{[3,5-bis(trifluoromethyl)benzyl]amino}

		}-4-phenyl-2-butanol
232	76	(2R,3S)-3-amino-4-phenyl-1-{[2-(trifluoromethyl)benzyl]amino}-2-butanol
233	77	(2R,3S)-3-amino-1-{[(1S)-1-(1-naphthyl)ethyl]amino}-4-phenyl-2-butanol
234	78	(2R,3S)-3-amino-1-{[(1R)-1-(1-naphthyl)ethyl]amino}-4-phenyl-2-butanol
235	79	4-({[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]amino}methyl)-2-methoxyphenol
236	80	4-({[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]amino}methyl)-1,2-benzenediol
237	81	(2R,3S)-3-amino-1-[(3-methoxypropyl)amino]-4-phenyl-2-butanol
238	82	(2R,3S)-3-amino-1-{[(1S)-2-hydroxy-1-methylethyl]amino}-4-phenyl-2-butanol
239	83	(2R,3S)-3-amino-1-{[(1R)-2-hydroxy-1-methylethyl]amino}-4-phenyl-2-butanol
240	84	(2R,3S)-3-amino-4-phenyl-1-(2-propynylamino)-2-butanol
241	85	(2R,3S)-3-amino-1-{[2-(2-fluorophenyl)ethyl]amino}-4-phenyl-2-butanol
242	86	(2R,3S)-3-amino-1-{[2-(3-fluorophenyl)ethyl]amino}-4-

		phenyl-2-butanol
243	87	(2R,3S)-3-amino-1-{[2-(4-fluorophenyl)ethyl]amino}-4-phenyl-2-butanol
244	88	(2R,3S)-3-amino-1-{[2-(4-bromophenyl)ethyl]amino}-4-phenyl-2-butanol
245	89	(2R,3S)-3-amino-1-{[2-(3-methoxyphenyl)ethyl]amino}-4-phenyl-2-butanol
246	90	(2R,3S)-3-amino-1-{[2-(2,4-dichlorophenyl)ethyl]amino}-4-phenyl-2-butanol
247	91	(2R,3S)-3-amino-1-{[2-(3-chlorophenyl)ethyl]amino}-4-phenyl-2-butanol
248	92	(2R,3S)-3-amino-1-{[2-(2,5-dimethoxyphenyl)ethyl]amino}-4-phenyl-2-butanol
249	93	(2R,3S)-3-amino-1-{[2-(4-methylphenyl)ethyl]amino}-4-phenyl-2-butanol
250	94	(2R,3S)-3-amino-1-{[(1R)-1-benzyl-2-hydroxyethyl]amino}-4-phenyl-2-butanol
251	95	(2R,3S)-3-amino-1-{[3-(4-morpholinyl)propyl]amino}-4-phenyl-2-butanol
252	96	(2R,3S)-3-amino-1-(isobutylamino)-4-phenyl-2-butanol
253	97	(2R,3S)-3-amino-1-{[2-(4-

		morpholinyl) ethyl] amino}-4-phenyl-2-butanol
254	98	(2R,3S)-3-amino-4-phenyl-1-[(2-hydroxybutyl) amino]-2-butanol
255	99	(2R,3S)-3-amino-4-phenyl-1-{[2-(2-thienyl) ethyl] amino}-2-butanol
256	100	4-{[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl] amino}-1-butanol
257	101	(2R,3S)-3-amino-1-{[(1S)-2-hydroxy-1-phenylethyl] amino}-4-phenyl-2-butanol
258	102	(2R,3S)-3-amino-1-[(2,4-dichlorobenzyl) amino]-4-phenyl-2-butanol
259	103	(2R,3S)-3-amino-1-{[(1R)-2-hydroxy-1-phenylethyl] amino}-4-phenyl-2-butanol
260	104	(2R,3S)-3-amino-1-[(4-tert-butylbenzyl) amino]-4-phenyl-2-butanol
261	105	(2R,3S)-3-amino-4-phenyl-1-[(1-phenylethyl) amino]-2-butanol
262	106	(1R,2S)-1-{[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl] amino}-2,3-dihydro-1H-inden-2-ol
263	107	(2R,3S)-3-amino-1-[(3,4-dimethylbenzyl) amino]-4-phenyl-2-butanol
264	108	methyl 7-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl] amino}heptanoate

265	109	2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutylpropanamide
266	110	(2S) -2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutylpropanamide
267	111	2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutyl-2-methylpropanamide
268	112	2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutylacetamide
269	113	(2S) -2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutylbutanamide
270	114	(2R) -2- { [(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -2-hydroxybutyl] amino} -N-isobutylbutanamide
271	115	(2R, 3S) -3-amino-1- (benzylamino) -4- (3, 5-difluorophenyl) -2-butanol
272	116	(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -1- (ethylamino) -2-butanol
273	117	(2R, 3S) -3-amino-4- (3, 5-difluorophenyl) -1- (isobutylamino) -2-butanol

274	118	3- { [(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-isobutyl-2-methylpropanamide
275	119	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-{[4-(dimethylamino)benzyl]amino}-2-butanol
276	120	(2S)-2-{ [(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-isobutyl-3-phenylpropanamide
277	121	(2S)-2-{ [(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-isobutyl-3-methylbutanamide
278	122	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-{[2-(dimethylamino)ethyl]amino}-2-butanol
279	123	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-{[3-pyridinylmethyl]amino}-2-butanol
280	124	(2S)-2-{ [(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-3-(benzyloxy)-N-isobutylpropanamide
281	125	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(1-methyl-1-phenylethyl)amino]-2-butanol
282	126	(2R)-2-{ [(2R,3S)-3-amino-4-(3,5-

		difluorophenyl)-2-hydroxybutyl]amino}-N-isobutyl-3-methylbutanamide
283	127	(2S)-2-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-isobutylpentanamide
284	128	(2S)-2-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-3-hydroxy-N-isobutylpropanamide
285	129	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(2-phenylethyl)amino]-2-butanol
286	130	(2S)-2-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-benzylpropanamide
287	131	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(1S)-1-phenylpropyl]amino}-2-butanol
287 a	169a	(2S)-2-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-ethylpropanamide
288	132	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol
289	133	(2S)-2-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-N-isobutyl-2-phenylethanamide

290	134	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-(isopentylamino)-2-butanol
291	135	(2R,3S)-3-amino-1-(cyclohexylamino)-4-(3,5-difluorophenyl)-2-butanol
292	136	(2R,3S)-3-amino-1-(butylamino)-4-(3,5-difluorophenyl)-2-butanol
293	137	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxypropyl)amino]-2-butanol
294	138	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(2-hydroxy-2-phenylethyl)amino]-2-butanol
295	139	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-{[(3R,5S)-3,5-dimethoxycyclohexyl]amino}-2-butanol
296	140	dimethyl (1R,3S)-5-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-1,3-cyclohexanedicarboxylate
297	141	(1R,3S)-5-{[(2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl]amino}-1,3-cyclohexanedicarboxylic acid
298	142	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-{[(1R)-1-phenylpropyl]amino}-2-butanol
299	143	(2R,3S)-3-amino-1-[(3-chlorobenzyl)amino]-4-(3,5-difluorophenyl)-2-butanol

300	144	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol
301	145	(2R,3S)-3-amino-1-[(1,1'-biphenyl)-3-ylmethyl]amino]-4-(3,5-difluorophenyl)-2-butanol
302	146	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-iodobenzyl)amino]-2-butanol
303	147	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methylbenzyl)amino]-2-butanol
304	148	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(2-phenylpropyl)amino]-2-butanol
305	149	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(1,3-thiazol-5-ylmethyl)amino]-2-butanol
306	150	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(2-thienylmethyl)amino]-2-butanol
307	151	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]-2-butanol
308	152	(2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(2-pyrazinylmethyl)amino]-2-butanol
309	153	(2R,3S)-3-amino-1-[(3,5-difluorobenzyl)amino]-4-(3,5-difluorophenyl)-2-butanol
310	154	(2R,3S)-3-amino-1-[(1,3-

		benzodioxol-5-ylmethyl) amino] -4-(3,5-difluorophenyl) -2-butanol
311	155	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(3,5-dimethoxybenzyl) amino] -2-butanol
312	156	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-{[3-(trifluoromethyl) benzyl] amino} -2-butanol
313	157	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(2-furylmethyl) amino] -2-butanol
314	158	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(7-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl) amino] -2-butanol
315	159	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-{[3-(trifluoromethoxy) benzyl] amino} -2-butanol
316	160	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(3-fluorobenzyl) amino] -2-butanol
317	161	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(3-isopropoxybenzyl) amino] -2-butanol
318	162	(2R,3S) -3-amino-1-[(3-bromobenzyl) amino] -4-(3,5-difluorophenyl) -2-butanol
319	163	(2R,3S) -3-amino-4-(3,5-difluorophenyl) -1-[(5-methyl-2-

		furylmethyl) amino] -2-butanol
320	164	(2R,3S) -3-amino-4- (3,5-difluorophenyl) -1- [(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl) amino] -2-butanol

EXAMPLES 321-473

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with the amine (VIII) of Column A and reacting it with the amide forming agent (IX) of Column B, the substituted amine (X) of Column C is obtained.

EXA	Column A Amine (VIII) Compound of EXA	Column B Amide Forming Agent (IX)	Column C Substitued amine (X)	Column D Physica l Data
321	170	"PHTH"	N ¹ - [(1S,2R) -1-benzyl-3-(ethylamino) -2-hydroxypropyl] -N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 440.7
322	171	"PHTH"	N ¹ - [(1S,2R) -1-benzyl-3-(benzylamino) -2-hydroxypropyl] -N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 502.7
323	172	"PHTH"	N ¹ - [(1S,2R) -1-benzyl-2-hydroxy-3-(isopropylamino) propyl] -N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 454.6

324	173	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-(4-toluidino)propyl]-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 502.2
325	174	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-{[2-(4-methoxyphenyl)ethyl]amino}propyl]-N ³ ,N ³ -ipropylisophthalamide	MH ⁺ = 546
326	175	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
327	176	"PHTH"	ethyl {[(3S)-3-({3-[(dipropylamino)carbonyl] benzoyl}amino)-2-hydroxy-4-phenylbutyl]amino} (phenyl) acetate	MH ⁺ = 574
329	178	"PHTH"	N ¹ -[(1S)-1-benzyl-2-hydroxy-3-{[(1S)-2-hydroxy-1-(hydroxymethyl)-2-(4-nitrophenyl)ethyl]amino}propyl]-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 607
330	179	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(2-chlorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 537
331	180	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-	MH ⁺ =

			[(4-chlorobenzyl) amino] - 2-hydroxypropyl} -N ³ ,N ³ - dipropylisophthalamide	537
332	181	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2- hydroxy-3-{[2-(2- hydroxyethoxy) ethyl] amino}propyl) -N ³ ,N ³ - dipropylisophthalamide	MH+ = 500
333	182	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-3- (2,3-dihydro-1H-inden-1- ylamino)-2- hydroxypropyl] -N ³ ,N ³ - dipropylisophthalamide	MH+ = 528
334	183	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-[(2- hydroxypropyl) amino] propy l} -N ³ ,N ³ - dipropylisophthalamide	MH+ = 470
335	184	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-[(tetrahydro-2- furanylmethyl) amino] propy l} -N ³ ,N ³ - dipropylisophthalamide	MH+ = 496
336	185	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3- [(2,2- diethoxyethyl) amino] -2- hydroxypropyl} -N ³ ,N ³ - dipropylisophthalamide	MH+ = 528
337	186	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-3- (butylamino) -2- hydroxypropyl] -N ³ ,N ³ - dipropylisophthalamide	MH+ = 468

338	187	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-3-(cyclohexylamino)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 494
339	188	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(2-pyridinylmethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 507
340	189	"PHTH"	N ¹ -{(1S,2R)-3-[(2-aminobenzyl)amino]-1-benzyl-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 517
341	190	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-pyridinylmethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 503
342	191	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-{[2-(1-pyrrolidinyl)ethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 509
343	192	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(2-hydroxy-2-phenylethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
344	193	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3-butoxypropyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -	MH ⁺ = 526

			dipropylisophthalamide	
345	194	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-isopropoxypropyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 512
346	195	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-(isopentylamino)propyl]-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 482
347	196	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-phenylpropyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 530
348	197	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(2-methoxyethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 470
349	198	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(2-phenoxyethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
350	199	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(2-propoxyethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 498
351	200	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-	MH ⁺ =

			[(3,3-dimethylbutyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	496
352	201	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(4-phenylbutyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 544
353	202	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 628
354	203	"PHTH"	N ¹ -{(1S)-1-benzyl-2-hydroxy-3-[(4-nitrobenzyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 547
355	204	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3-chlorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 537
356	205	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-{[2-(4-chlorophenyl)ethyl]amino}-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 551
357	206	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-{[2-(2-pyridinyl)ethyl]amino}propyl}-N ³ ,N ³ -	MH ⁺ = 517

			dipropylisophthalamide	
358	207	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(4-pyridinylmethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 503
359	208	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[2-(1-methyl-2-pyrrolidinyl)ethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 523
360	209	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(2,3-dimethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 580
361	210	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[2-(trifluoromethoxy)benzyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 586
362	211	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(2-chloro-6-phenoxybenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 629
363	212	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[4-(trifluoromethyl)benzyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 570

			dipropylisophthalamide	
364	213	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(2,3-dichlorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 571
365	214	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,5-dichlorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 571
366	215	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,5-difluorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 538
367	216	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[4-(trifluoromethoxy)benzyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 586
368	217	"PHTH"	N ¹ -[(1S,2R)-3-({2-[4-(aminosulfonyl)phenyl]ethyl}amino)-1-benzyl-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 595
369	218	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(4-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
370	219	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-	MH ⁺ =

			hydroxy-3-[(4-methylbenzyl) amino]propyl }-N ³ ,N ³ - dipropylisophthalamide	516
371	220	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3,4,5-trimethoxybenzyl) amino]propyl}-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 592
372	221	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-(trifluoromethoxy)benzyl) amino] propyl}-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 586
373	222	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,5-dimethoxybenzyl) amino]-2-hydroxypropyl}-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 562
374	223	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(2,4-dimethoxybenzyl) amino]-2-hydroxypropyl}-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 562
375	224	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[[[1,1'-biphenyl]-3-ylmethyl) amino]-2-hydroxypropyl}-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 578
376	225	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,4-dichlorobenzyl) amino]-2-	MH ⁺ = 571

			hydroxypropyl}-N ³ ,N ³ - dipropylisophthalamide	
377	226	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3- [(2-fluorobenzyl)amino]- 2-hydroxypropyl}-N ³ ,N ³ - dipropylisophthalamide	MH+ = 520
378	227	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-{[3- (trifluoromethyl)benzyl]a mino}propyl)-N ³ ,N ³ - dipropylisophthalamide	MH+ = 570
379	228	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-[(2- methylbenzyl)amino]propyl }-N ³ ,N ³ - dipropylisophthalamide	MH+ = 516
380	229	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-{[(1R)-1- phenylethyl]amino}propyl) -N ³ ,N ³ - dipropylisophthalamide	MH+ = 516
381	230	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-{[(1S)-1- phenylethyl]amino}propyl) -N ³ ,N ³ - dipropylisophthalamide	MH+ = 516
382	231	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3- {[3,5- bis(trifluoromethyl)benzy l]amino}-2- hydroxypropyl)-N ³ ,N ³ - dipropylisophthalamide	MH+ = 638

383	232	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[2-(trifluoromethyl)benzyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 570
384	233	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1S)-1-(1-naphthyl)ethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 566
385	234	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1R)-1-(1-naphthyl)ethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 566
386	235	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(4-hydroxy-3-methoxybenzyl)amino]propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 548
387	236	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,4-dihydroxybenzyl)amino]-2-hydroxypropyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 534
388	237	"PHTH"	N ¹ -{(1S)-1-benzyl-2-hydroxy-3-[(3-methoxypropyl)amino]propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 484
389	238	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1S)-2-	MH ⁺ = 470

			hydroxy-1-methylethyl]amino}propyl) -N ³ ,N ³ - dipropylisophthalamide	
390	239	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1R)-2-hydroxy-1-methylethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH+ = 470
391	240	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-(2-propynylamino)propyl]-N ³ ,N ³ -dipropylisophthalamide	MH+ = 450
392	241	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3-{[2-(2-fluorophenyl)ethyl]amino}-2-hydroxypropyl)-N ³ ,N ³ -dipropylisophthalamide	MH+ = 534
393	242	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3-{[2-(3-fluorophenyl)ethyl]amino}-2-hydroxypropyl)-N ³ ,N ³ -dipropylisophthalamide	MH+ = 534
394	243	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3-{[2-(4-fluorophenyl)ethyl]amino}-2-hydroxypropyl)-N ³ ,N ³ -dipropylisophthalamide	MH+ = 534
395	244	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3-{[2-(4-	MH+ = 595

			bromophenyl) ethyl] amino}- 2-hydroxypropyl)-N ³ ,N ³ - dipropylisophthalamide	
396	245	"PHTH"	N ¹ -((1S)-1-benzyl-2- hydroxy-3-{[2-(3- methoxyphenyl) ethyl] amino }propyl)-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 546
397	246	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3- {[2-(2,4- dichlorophenyl) ethyl] amin o}-2-hydroxypropyl)-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 585
398	247	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3- {[2-(3- chlorophenyl) ethyl] amino} -2-hydroxypropyl)-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 551
399	248	"PHTH"	N ¹ -((1S)-1-benzyl-3-{[2- (2,5- dimethoxyphenyl) ethyl] ami no}-2-hydroxypropyl)- N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 576
400	249	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2- hydroxy-3-{[2-(4- methylphenyl) ethyl] amino} propyl)-N ³ ,N ³ - dipropylisophthalamide	MH ⁺ = 530
401	250	"PHTH"	N ¹ -((1S,2R)-1-benzyl-3- {[(1R)-1-benzyl-2- hydroxyethyl] amino}-2-	MH ⁺ = 546

012987

			hydroxypropyl)-N ³ ,N ³ - dipropylisophthalamide	
402	251	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2- hydroxy-3-{[3-(4- morpholinyl)propyl]amino} propyl)-N ³ ,N ³ - dipropylisophthalamide	MH+ = 539
403	252	"PHTH"	N ¹ -[(1S,2R)-1-benzyl-2- hydroxy-3- (isobutylamino)propyl]- N ³ ,N ³ - dipropylisophthalamide	MH+ = 468
404	253	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2- hydroxy-3-{[2-(4- morpholinyl)ethyl]amino}p ropyl)-N ³ ,N ³ - dipropylisophthalamide	MH+ = 525
405	254	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-[(2- hydroxybutyl)amino]propyl }-N ³ ,N ³ - dipropylisophthalamide	MH+ = 484
406	255	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2- hydroxy-3-{[2-(2- thienyl)ethyl]amino}propy l)-N ³ ,N ³ - dipropylisophthalamide	MH+ = 522
407	256	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2- hydroxy-3-[(4- hydroxybutyl)amino]propyl }-N ³ ,N ³ - dipropylisophthalamide	MH+ = 484

01298 7

408	257	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1S)-2-hydroxy-1-phenylethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
409	258	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-{(2,4-dichlorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 571
410	259	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1R)-2-hydroxy-1-phenylethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 532
411	260	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-{(4-tert-butylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 558
412	261	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(1-phenylethyl)amino]propyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 516
413	262	"PHTH"	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino}propyl)-N ³ ,N ³ -	MH ⁺ = 544

			dipropylisophthalamide	
414	263	"PHTH"	N ¹ -{(1S,2R)-1-benzyl-3-[(3,4-dimethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 530
416	265	"PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 575
417	266	"PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 575
418	266	"3,5-pyridinyl"	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl)-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	MH ⁺ = 576
419	267	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-1,1-dimethyl-2-	MH ⁺ = 603

			oxoethyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
420	268	"5-Me-PH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-2-oxoethyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 575.3
421	269	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]propyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 603.8
422	270	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino)carbonyl]propyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 603.8
423	271	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 552.3
424	272	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-	MH+ = 490.7

			(ethylamino)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
425	273	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isobutylamino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 518.0
426	274	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(isobutylamino)-2-methyl-3-oxopropyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 603
427	275	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[4-(dimethylamino)benzyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 595.8
428	276	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-{[(1S)-1-benzyl-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 665.9
429	277	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-	MH+ =

			difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]-2-methylpropyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	617.9
430	278	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[2-(dimethylamino)ethyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 533.3
431	279	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-pyridinylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 553
432	280	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-{[(1S)-1-[(benzyloxy)methyl]-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 695.9
433	281	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)amino]propyl}-5-methyl-N ³ ,N ³ -	MH ⁺ = 580.8

			dipropylisophthalamide	
434	282	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino)carbonyl]-2-methylpropyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 617.9
435	283	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]butyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 617.9
436	284	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-(hydroxymethyl)-2-(isobutylamino)-2-oxoethyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 605.8
437	285	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-phenylethyl)amino]propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 566.8
438	286	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-({(1S)-2-(benzylamino)-1-methyl-2-	MH+ = 623.2

			oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
439	287	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-1-phenylpropyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 580.8
440	287a	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(1S)-2-(ethylamino)-1-methyl-2-oxoethyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 561.2
441	289	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-2-(isobutylamino)-2-oxo-1-phenylethyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 651.9
442	290	"5-Me-PHTH"	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isopentylamino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 531.69
443	291	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-	MH+ =

			(cyclohexylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	544.2
444	292	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-(butylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 518.2
445	293	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxypropyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 534.2
446	294	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-hydroxy-2-phenylethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 582
447	295	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3R,5S)-3,5-dimethoxycyclohexyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 604.2
448	296	"5-Me-PHTH"	dimethyl (1R,3S)-5-{[(2R,3S)-4-(3,5-	MH+ = 660.2

			difluorophenyl)-3-({3- [(dipropylamino) carbonyl] -5-methylbenzoyl}amino)- 2-hydroxybutyl]amino)- 1,3- cyclohexanedicarboxylate	
449	297	"5-Me-PHTH"	(1R,3S)-5-{[(2R,3S)-4- (3,5-difluorophenyl)-3- (3- [(dipropylamino) carbonyl] -5-methylbenzoyl}amino)- 2-hydroxybutyl]amino)- 1,3- cyclohexanedicarboxylic acid	MH+ = 632.2
450	298	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5- difluorobenzyl)-2- hydroxy-3-{[(1R)-1- phenylpropyl]amino}propyl) -5-methyl-N ³ ,N ³ - dipropylisophthalamide	MH+ = 580.8
451	299	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-[(3- chlorobenzyl)amino]-1- (3,5-difluorobenzyl)-2- hydroxypropyl]-5-methyl- N ³ ,N ³ - dipropylisophthalamide	MH+ = 586.3
452	300	"-SO ₂ -"	N-{(1S,2R)-1-(3,5- difluorobenzyl)-2- hydroxy-3-[(3- methoxybenzyl)amino]propyl }-3-[(2-	MH+ = 603.2

			propylpentyl) sulfonyl] benzamide	
453	301	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-[(1,1'-biphenyl)-3-ylmethyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 628.8
454	302	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 678
455	303	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methylbenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 566.8
456	304	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-phenylpropyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 694
457	305	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1,3-thiazol-5-ylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -	MH ⁺ = 559.5

01298 7

			dipropylisophthalamide	
458	306	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-thienylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 558.5
459	307	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 814
460	308	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-pyrazinylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 554.3
461	309	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3,5-difluorobenzyl)amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 588.7
462	310	"PHTH"	N ¹ -{(1S,2R)-3-[(1,3-benzodioxol-5-ylmethyl)amino]-1-benzyl-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 546
463	311	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-	MH ⁺ =

			difluorobenzyl)-3-[(3,5-dimethoxybenzyl)amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	612.4
464	312	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(trifluoromethyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 620.3
466	314	"5-Me-PHTH"	Racemic N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(7-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 814
467	315	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(trifluoromethoxy)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 636.3
468	316	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-fluorobenzyl)amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH+ = 570.3

469	317	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 610.4
470	318	"5-Me-PHTH"	N ¹ -[(1S,2R)-3-[(3-bromobenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 631.6
471	319	"5-Me-PHTH"	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-methyl-2-furyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 556
472	320	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide "isomer A"	MH ⁺ = 622
473	320	"5-Me-PHTH"	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}-5-methyl-N ³ ,N ³ -	MH ⁺ = 622

			dipropylisophthalamide "isomer B"	

EXAMPLE 474 N^1 -[(1S,2R)-1-benzyl-2-hydroxy-3-(1,2,3,4-tetrahydro-1-naphthalenylamino)propyl]- N^3,N^3 -dipropylisophthalamide (X)

5 Following the general procedures of EXAMPLES 6 and 321-473 and making non-critical variations but starting with (2R,3S)-3-amino-4-phenyl-1-(1,2,3,4-tetrahydro-1-naphthalenylamino)-2-butanol (VIII) and using "PHTH", the title compound is obtained, $MH^+ = 542.3$.

10 EXAMPLES 475-483

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with the amine (VIII) of Column A and reacting it with the amide forming agent (IX) of Column B, the substitute amine (X) of Column C is obtained.

15

EXA	Column A Amine (VIII) Compound d of EXA	Column B Amide Forming Agent (IX)	Column C Substitued amine (X)	Column D Physica l Data
475	271	"5-OMe-PHTH"	N^1 -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methoxy- N^3,N^3 -dipropylisophthalamide	$MH^+ = 568.2$
476	271	"PHTH"	N^1 -[(1S,2R)-3-(benzylamino)-1-(3,5-	$MH^+ = 538.2$

			difluorobenzyl)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	
477	271	"5-Cl-PHTH"	N ¹ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-chloro-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 572.1
478	271	"3,5-pyridinyl"	N ³ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	MH ⁺ = 539.5
479	271	"5-F-PHTH"	N ¹ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-fluoro-N ³ ,N ³ -dipropylisophthalamide	MH ⁺ = 556.3
480	271	"thienyl"	N ² -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ⁵ ,N ⁵ -dipropyl-2,5-thiophenedicarboxamide	MH ⁺ = 544.2
481	271	"2,4-pyridinyl"	N ⁴ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ² ,N ² -dipropyl-2,4-	MH ⁺ = 539.3

			pyridinedicarboxamide	
482	271	"4,6-pyrimidinyl "	N ⁴ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ⁶ ,N ⁶ dipropyl-4,6-pyrimidinedicarboxamide	MH ⁺ = 540
483	271	"morpholinyl 1"	N-[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-3-(4-morpholinylcarbonyl)benzamide	MH ⁺ = 552

EXAMPLE 484 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methylbenzyl)amino]propyl}-N³,N³-dipropylisophthalamide (X)

5 Following the general procedures of EXAMPLES 4 and 14-107 and making non-critical variations but starting with tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V, commercially available) and NH₂-CH₂-phenyl-(3-CH₃) (VI), the corresponding protected alcohol (VII) is obtained.

10 Following the general procedure of EXAMPLES 5 and 170-320 and making non-critical variations, the corresponding amine (VIII) is obtained.

Following the general procedure of EXAMPLES 6 and 321-483 and making non-critical variations but using "PTH" (IX), the
15 title compound is obtained, MH⁺ = 516.

EXAMPLE 485 N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N⁵,N⁵-dipropylpentanediamide (X)

Following the general procedure of EXAMPLES 6 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol (VIII, EXAMPLE 5) with 5-(dipropylamino)-5-oxopentanoic acid (IX), the title compound is obtained, $MH^+ = 534.2$.

EXAMPLE 486 N^1 -[(1S,2R)-3-{[(1R)-1-[(benzyloxy)methyl]-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedures of EXAMPLES 124, 280 and 432 and making non-critical variations but using the "R" C-terminal amine (VI), the title compound is obtained, $MH^+ = 695$.

EXAMPLE 487 N^1 -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1R)-1-(hydroxymethyl)-2-(isobutylamino)-2-oxoethyl]amino}propyl)-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedures of EXAMPLES 128, 284 and 436 and making non-critical variations but using the "R" C-terminal amine (VI), the title compound is obtained, $MH^+ = 719$.

EXAMPLE 488 N^1 -[(1S,2R)-1-benzyl-2-hydroxy-3-(pentylamino)propyl]- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedures of EXAMPLES 96, 252 and 403 and making non-critical variations but using $NH_2-(CH_2)_4-CH_3$ as the C-terminal amine (VI), the title compound is obtained, $MH^+ = 481$.

EXAMPLE 489 N^1 -[(1S)-3-({2-[4-(aminosulfonyl)phenyl]ethyl}amino)-1-benzyl-2-hydroxypropyl]- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedures of EXAMPLES 61, 217 and 368 and making non-critical variations but using racemic C-terminal amine (VI), the title compound is obtained, $MH^+ = 595$.

- 5 EXAMPLE 491 N^3 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1,3-thiazol-5-ylmethyl)amino]propyl}- N^5 , N^5 -dipropyl-3,5-pyridinedicarboxamide (X)

Following the general procedure of EXAMPLE 457 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(1,3-thiazol-5-ylmethyl)amino]-2-butanol (VIII, EXAMPLE 305) with "3,5-pyridinyl" as the amide forming agent (IX), the title compound is obtained.

- 15 EXAMPLE 492 3-Benzoyl-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}benzamide (X)

Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII, EXAMPLE 5) and reacting it with phenyl-CO-phenyl-CO-OH (IX) where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups, the title compound is obtained, $MH^+ = 545.2$

- 25 EXAMPLE 493 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}[1,1'-biphenyl]-3-carboxamide (X)

Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII, EXAMPLE 5) and reacting it with phenyl-phenyl-CO-OH (IX) where the attachment to the middle -phenyl-

ring is 1,3- for the carbonyl group and other -phenyl , the title compound is obtained, $MH^+ = 517.2$.

EXAMPLE 494 $N^1-\{(1S,2R)-1\text{-benzyl-2-hydroxy-3-}[(3\text{-methylbenzyl)amino}]propyl\}-N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII, EXAMPLE 5) and reacting it with (CH₃-O-CH₂-CH₂)₂N-CO--CO-OH (IX) where the attachment to the -phenyl - ring is 1,3- for the carbonyl groups, the title compound is obtained, $MH^+ = 570.3$.

EXAMPLE 495 $N^1-\{(1S,2R)-3\text{-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl}\}-N^3\text{-(2-methoxyethyl)-}N^3\text{-propylisophthalamide (X)}$

Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-1-(benzylamino)-4-(3,5-difluorophenyl)-2-butanol (VIII, EXAMPLE 271) and reacting it with (CH₃-CH₂-CH₂)-(CH₃-O-CH₂-CH₂)-N-CO-phenyl-CO-OH (IX) where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups, the title compound is obtained, $MH^+ = 554.5$.

25

EXAMPLE 496 $N-\{(1S,2R)-1\text{-(3,5-difluorobenzyl)-2-hydroxy-3-}[(3\text{-methoxybenzyl)amino}]propyl\}\text{-3-ethoxybenzamide (X)}$
Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII, EXAMPLE 5) and reacting it with CH₃-CH₂-O-phenyl-CO-OH (IX) where the attachment to the -phenyl- ring

is 1,3- for the carbonyl and ethoxy group, the title compound is obtained, $MH^+ = 485$

EXAMPLE 497 $N-\{(1S,2R)-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxy-3-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl}\}-2\text{-naphthamide (X)}$

5 Following the general procedure of EXAMPLE 452 and making non-critical variations and using (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol trifluoroacetate (VIII, EXAMPLE 5) and reacting ortho position, the title compound is obtained, $MH^+ = 491$.

10

EXAMPLE 498 $N^1-\{(1S,2R)-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxy-3-}[(1R)-1,2,3,4\text{-tetrahydro-1-naphthalenylamino}]\text{propyl}\}-5\text{-methyl-}N^3,N^3\text{-dipropylisophthalamide (X)}$

15 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 1,2,3,4-tetrahydro-1-naphthalenylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 588$.

20

EXAMPLE 499 $N^1-\{(1R)-3-\{[3,5\text{-bis(trifluoromethyl)benzyl}]\text{amino}\}-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxypropyl}\}-5\text{-methyl-}N^3,N^3\text{-dipropylisophthalamide (X)}$

25 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), $H_2N\text{-CH}_2\text{-phenyl-(3,5-ditrifluoromethyl)}$ (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 688$.

30

EXAMPLE 500 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[2\text{-fluoro-5-}$
 $(\text{trifluoromethyl})\text{benzyl}\}\text{amino})-2\text{-hydroxypropyl})-$
 $N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 5 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N-CH_2\text{-phenyl-(2-fluoro-}$
 $5\text{-trifluoromethyl)}$ (VI) and "PHTH" (IX), the title compound is
 obtained, $MH^+ = 588$.

EXAMPLE 501 $N^1-\{(1S,2R)-1\text{-benzyl-3-}\{(2,3\text{-}$
 10 $\text{difluorobenzyl})\text{amino}\}-2\text{-hydroxypropyl}\}-N^3,N^3\text{-}$
 $\text{dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N-CH_2\text{-phenyl-(2,3-}$
 15 difluoro) (VI) and "PHTH" (IX), the title compound is obtained,
 $MH^+ = 538$.

EXAMPLE 502 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[3\text{-fluoro-4-}$
 20 $(\text{trifluoromethyl})\text{benzyl}\}\text{amino})-2\text{-hydroxypropyl})-$
 $N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N-CH_2\text{-phenyl-(3-fluoro-}$
 $4\text{-trifluoromethyl)}$ (VI) and "PHTH" (IX), the title compound is
 25 obtained, $MH^+ = 588$.

EXAMPLE 503 $N^1-\{(1S,2R)-1\text{-benzyl-3-}\{(2,5\text{-}$
 30 $\text{difluorobenzyl})\text{amino}\}-2\text{-hydroxypropyl}\}-N^3,N^3\text{-}$
 $\text{dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N-CH_2\text{-phenyl-(2,5-}$

difluoro) (VI) and "PHTH" (IX), the title compound is obtained,
 $MH^+ = 538$.

EXAMPLE 504 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[3\text{-fluoro-5-}$
 5 $(\text{trifluoromethyl})\text{benzyl}\}\text{amino})\text{-2-hydroxypropyl})\text{-}$
 $N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-(3-fluoro-}$
 10 $5\text{-trifluoromethyl})$ (VI) and "PHTH" (IX), the title compound is
 obtained, $MH^+ = 588$.

EXAMPLE 505 $N^1-\{(1S,2R)-1\text{-benzyl-3-}\{(3,4\text{-}$
 $\text{difluorobenzyl})\text{amino}\}\text{-2-hydroxypropyl}\}\text{-}N^3,N^3\text{-}$
 15 $\text{dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-(3,4-}$
 difluoro) (VI) and "PHTH" (IX), the title compound is obtained,
 20 $MH^+ = 538$.

EXAMPLE 506 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[4\text{-fluoro-3-}$
 $(\text{trifluoromethyl})\text{benzyl}\}\text{amino})\text{-2-hydroxypropyl})\text{-}$
 $N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl 1-(2-
 oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-(3-}$
 trifluoromethyl-4-fluoro) (VI) and "PHTH" (IX), the title
 compound is obtained, $MH^+ = 588$.

EXAMPLE 507 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[2\text{-chloro-5-}$
 $(\text{trifluoromethyl})\text{benzyl}\}\text{amino})\text{-2-hydroxypropyl})\text{-}$
 $N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl-(2-chloro-5-trifluoromethyl) (VI) and "PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 605$.

EXAMPLE 508 N^1 -((1S,2R)-1-benzyl-3-{[4-chloro-3-(trifluoromethyl)benzyl]amino}-2-hydroxypropyl)- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl-(3-trifluoromethyl-4-chloro) (VI) and "PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 605$.

EXAMPLE 509 N^1 -[(1S,2R)-1-benzyl-3-(2,3-dihydro-1H-inden-2-ylamino)-2-hydroxypropyl]- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), H_2N -(2,3-dihydro-1H-inden-2-yl) and "PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 528$.

EXAMPLE 510 N^1 -{(1S)-1-benzyl-2-hydroxy-3-[(3-nitrobenzyl)amino]propyl}- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl-3-nitro (VI) and "PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 547$.

EXAMPLE 511 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[3\text{-(difluoromethoxy)benzyl]amino}\}-2\text{-hydroxypropyl})\text{-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-3(-O-CHF}_2\text{)}$ (VI) and "PHTH" (IX), the title compound is obtained, $MH^+ = 538$.

EXAMPLE 512 $N^1-\{(1S,2R)-1\text{-benzyl-3-}[(3\text{-ethoxybenzyl)amino}]\text{-2-hydroxypropyl}\}\text{-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-(3-O-CH}_2\text{-CH}_3\text{)}$ (VI) and "PHTH" (IX), the title compound is obtained, $MH^+ = 546$.

EXAMPLE 513 $N^1-((1S,2R)-1\text{-benzyl-2-hydroxy-3-}\{[(5\text{-methyl-2-pyrazinyl)methyl]amino}\}\text{propyl})\text{-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 5-methyl-2-methylaminoprazine (VI) and "PHTH" (IX), the title compound is obtained, $MH^+ = 518$.

EXAMPLE 514 $N^1-\{(1S,2R)-1\text{-benzyl-3-}[(3\text{-bromo-4-fluorobenzyl)amino}]\text{-2-hydroxypropyl}\}\text{-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), $H_2N\text{-CH}_2\text{-phenyl-(3-bromo-4-fluoro)}$ (VI) and "PHTH" (IX), the title compound is obtained, $MH^+ = 599$.

EXAMPLE 515 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3,5-dimethylbenzyl)amino]-2-hydroxypropyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

5 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), H_2N - CH_2 -phenyl-(3,5-dimethyl) (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 580$.

10

EXAMPLE 516 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethoxybenzyl)amino]-2-hydroxypropyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

15 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), H_2N - CH_2 -phenyl-(3-ethoxy) (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 596$.

20 EXAMPLE 517 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-phenoxyethyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

25 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), H_2N - CH_2 - CH_2 -O-phenyl (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 582$.

30 EXAMPLE 518 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isobutoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-

(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl-O-i-butyl (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 624$.

- 5 EXAMPLE 519 $\text{N}^1-((1\text{S},2\text{R})-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxy-3-}\{[(4\text{-methyl-1,3-thiazol-2-yl)methyl]amino\}propyl)-5\text{-methyl-}\text{N}^3,\text{N}^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
10 making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -4-methyl-1,3-thiazol-2-yl (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 573$.

- 15 EXAMPLE 520 $\text{N}^1-[(1\text{S},2\text{R})-3-(\text{benzylamino})-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxypropyl}]-\text{N}^3\text{-methyl-}\text{N}^3\text{-propylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
making non-critical variations but using tert-butyl (1S)-2-
20 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl (VI) and $(\text{CH}_3-\text{CH}_2-\text{CH}_2-)(\text{CH}_3-)\text{N}-\text{CO}-(\text{CH}_3-)\text{phenyl}-\text{CO}-\text{OH}$ (IX) where the attachment to the -phenyl- ring for the carbonyls is 1-,3- and 5- for the methyl group, the title compound is obtained, $\text{MH}^+ = 510$.

- 25 EXAMPLE 521 $\text{N}^2-[(1\text{S},2\text{R})-3-(\text{benzylamino})-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxypropyl}]-\text{N}^5,\text{N}^5\text{-dipropyl-2,5-furandicarboxamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and
30 making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), $\text{H}_2\text{N}-\text{CH}_2$ -phenyl (VI) and $(\text{CH}_3-\text{CH}_2-\text{CH}_2-)_2\text{N}-\text{CO}-(2,5\text{-furanyl})-\text{CO}-$ (IX), the title compound is obtained, $\text{MH}^+ = 528$.

EXAMPLE 522 $\cdot N^3$ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
 {[3-(trifluoromethyl)benzyl]amino}propyl)- N^5, N^5 -
 dipropyl-3,5-pyridinedicarboxamide (X)

5 Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl (1S)-2-
 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), H_2N -
 CH_2 -phenyl-3-trifluoromethyl (VI) and "3,5-pyridinyl" (IX), the
 title compound is obtained, $MH^+ = 607$.

10

EXAMPLE 523 N^3 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
 [(1-methyl-1-phenylethyl)amino]propyl}- N^5, N^5 -
 dipropyl-3,5-pyridinedicarboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and
 15 making non-critical variations but using tert-butyl (1S)-2-
 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), H_2N -
 C($-CH_3$)₂-phenyl (VI) and "3,5-pyridinyl" (IX), the title
 compound is obtained, $MH^+ = 567$.

20 EXAMPLE 524 N^1 -[(1S,2R)-3-amino-1-(3,5-difluorobenzyl)-2-
 hydroxypropyl]-5-methyl- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of CHARTS A, C and D and
 EXAMPLES 165-168 and making non-critical variations, but using
 25 tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-
 oxiranyl]ethylcarbamate (V) and "5-Me-PHTH" (IX), the title
 compound is obtained, $MH^+ = 462$.

EXAMPLE 525 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(1,2-
 30 diphenylethyl)amino]-2-hydroxypropyl}-5-methyl- N^3, N^3 -
 dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and
 making non-critical variations but using tert-butyl (1S)-2-

(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), $\text{NH}_2\text{-CH()-CH}_2\text{-phenyl}$ (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 642$.

- 5 EXAMPLE 526 $\text{N}^1\text{-}\{(1\text{S},2\text{R})\text{-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(7-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}\}\text{-5-methyl-N}^3\text{,N}^3\text{-dipropylisophthalamide, isomer A (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
10 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 7-methoxy-1,2,3,4-tetrahydro-1-naphthalenylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 622$.

- 15 EXAMPLE 527 $\text{N}^1\text{-}\{(1\text{S},2\text{R})\text{-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(7-methoxy-1,2,3,4-tetrahydro-1-naphthalenyl)amino]propyl}\}\text{-5-methyl-N}^3\text{,N}^3\text{-dipropylisophthalamide, isomer B (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
20 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 7-methoxy-1,2,3,4-tetrahydro-1-naphthalenylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $\text{MH}^+ = 622$.

- 25 EXAMPLE 528 Benzyl-(1S, 2R)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (XII)

Following the general procedure of EXAMPLE 165 and making non-critical variations but starting with benzyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 13), the title compound is obtained.

- 30 EXAMPLE 529 $\text{N-}\{(1\text{S},2\text{R})\text{-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}\}\text{-3-}$

(dimethylamino)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-dimethylaminobenzoic acid (IX), the title compound is obtained, $MH^+ = 448$.

EXAMPLE 530 N-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-2-methyl-1H-benzimidazole-5-carboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 2-methyl-1H-benzimidazole-5-carboxylic acid (IX), the title compound is obtained, $MH^+ = 459$.

EXAMPLE 531 3-(aminosulfonyl)-N-{(1S)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-chlorobenzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-(aminosulfonyl)-4-chlorobenzoic acid (IX), the title compound is obtained, $MH^+ = 519$.

EXAMPLE 532 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-cyanobenzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI)

and 3-cyanobenzoic acid (IX), the title compound is obtained,
 $MH^+ = 430$.

5 EXAMPLE 533 N-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-4-chloro-3-nitrobenzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI)
 10 and 4-chloro-3-nitrobenzoic acid (IX), the title compound is obtained, $MH^+ = 484$.

15 EXAMPLE 534 Methyl 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino)carbonyl]-5-nitrobenzoate (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-nitro-5-(methyl-O-CO-)phenyl-CO-OH (IX), the title
 20 compound is obtained, $MH^+ = 508$.

EXAMPLE 535 tert-butyl 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino)carbonyl]-phenylcarbamate (X)

25 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-(t-butyl-O-CO-NH)-phenyl-CO-OH (IX), the title compound is obtained, $MH^+ = 520$.

30 EXAMPLE 536 N-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-9,10-dioxo-9,10-

dihydro-2-anthourancenylcarboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 9,10-dioxo-9,10-dihydro-2-anthourancenylcarboxylic acid (IX), the title compound is obtained, $MH^+ = 535$.

EXAMPLE 537 N-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-1H-1,2,3-benzotriazole-6-carboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 1H-1,2,3-benzotriazolyl-6-carboxylic acid (IX), the title compound is obtained, $MH^+ = 446$.

EXAMPLE 538 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-(3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-(3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl)benzoic acid (IX), the title compound is obtained, $MH^+ = 501$.

EXAMPLE 539 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-1H-indole-5-carboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-

oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and indole-5-carboxylic acid (IX), the title compound is obtained, $MH^+ = 444$.

5 EXAMPLE 540 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-fluoro-5-(trifluoromethyl)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-fluoro-5-(trifluoromethyl)benzoic acid (IX), the title compound is obtained, $MH^+ = 491$.

15 EXAMPLE 541 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3-(trifluoromethyl)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-(trifluoromethyl)benzoic acid (IX), the title compound is obtained, $MH^+ = 473$.

25 EXAMPLE 542 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-4-(butylamino)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-(butylamino)benzoic acid (IX), the title compound is obtained, $MH^+ = 476$.

EXAMPLE 543 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3-(trifluoromethoxy)benzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-(trifluoromethoxy)benzoic acid (IX), the title compound is obtained, $MH^+ = 489$.

10 EXAMPLE 544 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3,5-dimethoxybenzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3,5-dimethoxybenzoic acid (IX), the title compound is obtained, $MH^+ = 465$.

EXAMPLE 545 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3,5-dimethylbenzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3,5-dimethylbenzoic acid (IX), the title compound is obtained, $MH^+ = 433$.

EXAMPLE 546 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3,5-difluorobenzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI)

and 3,5-difluorobenzoic acid (IX), the title compound is obtained, $MH^+ = 441$.

EXAMPLE 547 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-3,5-dichlorobenzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3,5-dichlorobenzoic acid (IX), the title compound is obtained, $MH^+ = 474$.

EXAMPLE 548 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-4-(benzyloxy)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-(benzyloxy)benzoic acid (IX), the title compound is obtained, $MH^+ = 511$.

EXAMPLE 549 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]-propyl}-1,3-benzodioxole-5-carboxamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 1,3-benzodioxole-5-carboxylic acid (IX), the title compound is obtained, $MH^+ = 449$.

EXAMPLE 550 3-(acetyl-amino)-N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-(acetylamino)benzoic acid (IX), the title compound is
 5 obtained, $MH^+ = 462$.

EXAMPLE 551 4-(acetylamino)-N-((1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl)benzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and
 10 making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-(acetylamino)benzoic acid (IX), the title compound is obtained, $MH^+ = 462$.

15 EXAMPLE 552 N^1 -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3,5-dimethyl-4-isoxazolyl)methyl]amino}-2-hydroxypropyl)-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and
 20 making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 3,5-dimethyl-4-isoxazolylmethylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 571.3$.

25 EXAMPLE 553 N^1 -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-phenylpropyl)amino]propyl)-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V,
 30 EXAMPLE 3), 3-phenylpropylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 580$.

EXAMPLE 554 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-furylmethyl)amino]-2-hydroxypropyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 3-furylmethylamine (VI) and "5-Me-PHTH" (IX); the title compound is obtained, $MH^+ = 542$.

10 EXAMPLE 555 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(tetrahydro-3-furanylmethyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (tetrahydro-3-furanylmethyl)amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 546$.

20 EXAMPLE 556 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-propoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (3-propoxybenzyl)amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 610$.

30 EXAMPLE 557 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-pyridinylmethyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-

(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (2-pyridinylmethyl)amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 553$.

- 5 EXAMPLE 558 N^1 -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-hydroxy- N^3,N^3 -dipropylisophthalamide (X)

10 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), benzylamine (VI) and 5-hydroxy-N,N-dipropylisophthalic acid (IX), the title compound is obtained, $MH^+ = 554$.

- 15 EXAMPLE 559 N^1 -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[1-methyl-1-(3-methylphenyl)ethyl]amino}propyl]-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

20 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), [1-methyl-1-(3-methylphenyl)ethyl]amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 594$.

- 25 EXAMPLE 560 N^1 -{[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1S)-1,2,3,4-tetrahydro-1-naphthalenylamino]propyl]-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

30 Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V,

EXAMPLE 3), (1S)-1,2,3,4-tetrahydro-1-naphthalenylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 592$.

EXAMPLE 561 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(2,5-dimethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-
5 N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
10 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 2,5-dimethylbenzylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 580$.

EXAMPLE 562 N^1 -[(1S,2R)-3-{[2-chloro-5-(trifluoromethyl)benzyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- N^3,N^3 -
15 dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
20 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 2-chloro-5-trifluorobenzylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 654$.

EXAMPLE 563 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-hydroxy-5-methylbenzyl)amino]propyl}-5-methyl- N^3,N^3 -
25 dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
30 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), 2-hydroxy-5-methylbenzylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 582$.

EXAMPLE 564 $N^1-((1S,2R)-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxy-3-} \{[(1S,2R)-2\text{-hydroxy-2,3-dihydro-1H-inden-1-yl}] \text{amino}\} \text{propyl})-5\text{-methyl-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), [(1S,2R)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 594$.

10

EXAMPLE 565 $N^1-\{(1S,2R)-1-(3,5\text{-difluorobenzyl})-3-[(1R)-2,3\text{-dihydro-1H-inden-1-ylamino}]-2\text{-hydroxypropyl}\}-5\text{-methyl-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (1R)-2,3-dihydro-1H-inden-1-ylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 578$.

20

EXAMPLE 566 $5\text{-chloro-}N^1-\{(1S,2R)-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxy-3-}[(1\text{-methyl-1-phenylethyl}) \text{amino}] \text{propyl}\}-N^3,N^3\text{-dipropylisophthalamide (X)}$

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (1-methyl-1-phenylethyl)amine (VI) and "5-Cl-PHTH" (IX), the title compound is obtained, $MH^+ = 601$.

30

EXAMPLE 567 $N^1-[(1S,2R)-3-[(1\text{-benzofuran-2-ylmethyl}) \text{amino}]-1-(3,5\text{-difluorobenzyl})-2\text{-hydroxypropyl}]-5\text{-methyl-}$

N³,N³-dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
 5 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (1-benzofuran-2-ylmethyl)amine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, MH⁺ = 592.

EXAMPLE 568 N¹-[(1S,2R)-3-{[(1R)-1-(3-
 10 bromophenyl)ethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N³,N³-dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-
 15 (3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3), (1R)-1-(3-bromophenyl)ethylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, MH⁺ = 645.

EXAMPLE 569 N¹-{(1S,2R)-1-(4-fluorobenzyl)-2-hydroxy-3-[(3-
 20 iodobenzyl)amino]propyl}-5-methyl-N³,N³-dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(4-
 25 fluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-iodobenzylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, MH⁺ = 660.

EXAMPLE 570 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-
 30 methoxybenzyl)amino]propyl}-3-[butyl(butyryl)amino]-5-methylbenzamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-

oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 3-[butyl(butyryl)amino]-5-methylbenzoic acid (IX), the title compound is obtained, $MH^+ = 560$.

- 5 EXAMPLE 571 N^1 -{1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-methyl- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-methyl-N,N-dipropylisophthalic acid (IX), the title compound is obtained, $MH^+ = 546$.

- 15 EXAMPLE 572 N^3 -{1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-methyl- N^1, N^1 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-methyl-N,N-dipropylisophthalic acid (IX), the title compound is obtained, $MH^+ = 546$.

- 25 EXAMPLE 573 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-methyl- N^3, N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-methoxybenzylamine (VI) and 4-methyl-N,N-dipropylisophthalic acid (IX), the title compound is obtained, $MH^+ = 582$.

EXAMPLE 574 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-1-butyl-1H-indole-6-carboxamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl 1-(2-oxiranyl)-2-phenylethylcarbamate (V), 3-methoxybenzylamine (VI) and 1-butyl-1H-indole-6-carboxylic acid (IX), the title compound is obtained, $MH^+ = 500$.

10 EXAMPLE 575 N^1 -[(1S,2R)-3-anilino-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), aniline (VI) and "5-Me-PHTH" (IX), the title compound is obtained, $MH^+ = 538$.

EXAMPLE 576 5-bromo- N^1 -[(1S,2R)-3-[(3-bromobenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-bromobenzylamine (VI) and 5-bromo-N,N-dipropylisophthalic acid (IX), the title compound is obtained, $MH^+ = 696$.

EXAMPLE 577 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4-methylpentanamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-

iodobenzylamine (VI) and 4-methylpentanoic acid (IX), the title compound is obtained, $MH^+ = 531$.

EXAMPLE 578 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
5 [(3-iodobenzyl)amino]propyl}-3-methylpentanamide
(X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-iodobenzylamine (VI) and $\text{CH}_3\text{-CH}_2\text{-CH}(\text{CH}_3\text{-})\text{-CH}_2\text{-CO-OH}$ (IX), the title compound is obtained, $\text{MH}^+ = 531$.

EXAMPLE 579 N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
[(3-hydroxybenzyl)amino]propyl}-5-methyl-N³,N³-
dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V), 3-hydroxybenzylamine (VI) and "5-Me-PHTH" (IX), the title
20 compound is obtained, $MH^+ = 568$.

EXAMPLE 580 *tert*-butyl (1*S*)-1-(3,5-difluorobenzyl)-3-[(3-methoxybenzyl)amino]-2-oxopropylcarbamate (XI)

tert-butyl (1*S*)-3-bromo-1-(3,5-difluorobenzyl)-2-

25 oxopropylcarbamate (III, EXAMPLE 1, 1 equivalent) is dissolved in isopropanol and treated with 3-methoxybenzylamine (VI, 5 equivalents). The reaction is heated at reflux for 2 hours and monitored by TLC for disappearance of the ketone (III). Upon completion, the reaction is concentrated to dryness under

30 reduced pressure and partitioned between equal parts water and ethyl acetate. The organic phase is extracted, washed two additional times with water, then saline, then dried over anhydrous sodium sulfate, filtered and concentrated under

reduced pressure. The crude material is purified by column chromatography on silica gel to give the title compound.

EXAMPLE 581 *tert*-butyl (1*S*,2*R*)-3-amino-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (XIII)

5 *tert*-butyl-(1*S*, 2*R*)-3-azido-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (XII, EXAMPLE 165, 1 equivalent) is dissolved in methanol and treated with 10% palladium on carbon under hydrogen (50 psi). The reaction is shaken at 20-25
10 degrees C for 2 hours, filtered through a diatomaceous earthpad, and concentrated under reduced pressure to dryness. The crude material is purified by column chromatography on silica gel to give the title compound.

15 EXAMPLE 582 Benzyl (1*S*,2*R*)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII)

A mixture of benzyl (1*S*)-1-[(2*S*)-oxiranyl]-2-phenylethylcarbamate (V, 0.22 g, 0.74 mmol), 3-methoxybenzylamine (VI, 0.13 g, 0.92 mmol), and ethanol (2 mL)
20 is stirred at reflux for 2.3 hours and then cooled and concentrated under reduced pressure. The residue is chromatographed (silica gel; methanol/dichloromethane/ammonium hydroxide, 4/96/trace) to give the title compound, MS *m/z* at (*m*
25 + *H*)⁺ = 435.2.

EXAMPLE 583 Benzyl (1*S*,2*R*)-1-benzyl-2-hydroxy-3-(*tert*-butylcarbamoyl)-3-[(3-methoxybenzyl)amino]propyl-carbamate (XXXIV)

30 A mixture of benzyl (1*S*,2*R*)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl-carbamate (VII, EXAMPLE 582, 9.00 g, 21.0 mmol), di-*tert*-butyl dicarbonate (5.00 g, 23 mmol), sodium carbonate monohydrate (3.12 g, 25 mmol), THF (200 mL), and

water (100 mL) is stirred at 20-25 degrees C for 2 hours. THF is removed under reduced pressure and the residue is partitioned between ethyl acetate and water. The organic layer is then washed with saline, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography (silica gel; methanol/dichloromethane, 1/99) gives the title compound, NMR (300 MHz, CDCl₃) δ 7.26, 6.79, 5.00, 4.42, 3.89, 3.78, 3.44, 3.28, 2.91, 1.49; MS (ESI+) for C₃₁H₃₈N₂O₆ m/z (M+H)⁺ = 535.3.

EXAMPLE 584 *tert*-Butyl (2R,3S)-3-amino-2-hydroxy-4-phenylbutyl(3-methoxybenzyl)carbamate (XXXV)

A suspension of benzyl (1S,2R)-1-benzyl-2-hydroxy-3-(*tert*-butylcarbamoyl)-3-[(3-methoxybenzyl)amino]propyl-carbamate (XXXIV, EXAMPLE 583, 10.2 g, 18.6 mmol) and palladium on carbon (5%, 1.00 g) in ethanol (100 mL) is shaken in a hydrogenation apparatus under hydrogen (50 psi) for 2 hours. Then the mixture is filtered through diatomaceous earth and concentrated. The concentrate is chromatographed (silica gel; methanol/dichloromethane, 5/95) to give the title compound, NMR (CDCl₃) δ 1.50, 2.46, 2.92, 3.04, 3.42, 3.52, 3.72, 3.82, 4.49, 6.83, 7.19 and 7.28; MS (ESI+) for C₂₃H₃₂N₂O₄ m/z (M+H)⁺ = 401.3.

EXAMPLE 585 *tert*-butyl (2R,3S)-3-({3-cyano-5-[(dipropylamino)carbonyl]benzoyl}amino)-2-hydroxy-4-phenylbutyl(3-methoxybenzyl)carbamate (XXXVI)

To a mixture of 3-cyano-5-[(dipropylamino)carbonyl]benzoic acid (IX/XXXII, PREPARATION 7, 0.086 g, 0.314 mmol) and *tert*-butyl (2R,3S)-3-amino-2-hydroxy-4-phenylbutyl(3-methoxybenzyl)carbamate (XXXV, EXAMPLE 584, 0.126 g, 0.314 mmol) in dichloromethane (0.3 mL) is added

diethylcyanophosphonate (0.062 g, 0.330 mmol) and triethylamine (0.032 g, 0.316 mmol). The mixture is stirred at 20-25 degrees for 65 hours and then partitioned between dichloromethane and saturated aqueous sodium bicarbonate. The organic phase is separated and dried over sodium sulfate and concentrated. The residue is chromatographed (silica gel, 20 mL; methanol/dichloromethane, 5/95) to give several mixed fractions, which are combined and rechromatographed (silica gel; acetone/hexane, 20/80) to give the title compound, MS (ESI+) for $C_{38}H_{48}N_4O_6$ m/z 657.6 $(M+H)^+ = 657.6$; $(M+Na) = 679.5$ and $(M-C_4H_9O_2) = 557.5$.

EXAMPLE 586 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-cyano- N^3,N^3 -dipropylisophthalamide hydrochloride (X)

A mixture of tert-butyl (2R,3S)-3-({3-cyano-5-[(dipropylamino)carbonyl]benzoyl}amino)-2-hydroxy-4-phenylbutyl(3-methoxybenzyl)carbamate (XXXVI, EXAMPLE 585, 0.080 g, 0.122 mmol), dichloromethane (1 mL), and methanol saturated with hydrochloric acid (1 mL) is stirred for 8 hours, after which time the solvents are removed under reduced pressure. A few drops of methanol, followed by ether, gives the title compound, MS (ESI+) for $C_{33}H_{40}N_4O_4$ m/z $(M+H)^+ = 557.5$

EXAMPLE 587 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^3,N^3 -dipropyl-1,3,5-benzenetricarboxamide (X)

To a mixture of 3-(aminocarbonyl)-5-[(dipropylamino)carbonyl]benzoic acid (IX, PREPARATION 6, 0.18 g, 0.616 mmol) in dry DMF (16 mL) is added EDC (0.182 g, 0.9 mmol), HOBT (0.127 g, 0.9 mmol), triethylamine (0.062 g, 0.616 mol), and (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-

2-butanol (VIII, EXAMPLE 175, 0.185 g, 0.616 mmol). The mixture is stirred at 20-25 degrees C for 3 days. The mixture is partitioned between water and ethyl acetate. The phases are separated and the organic phase is washed three times with
5 water. The organic phase is dried over anhydrous magnesium sulfate, filtered and concentrated. Column chromatography (silica gel, 75 mL; methanol/methylene chloride, 10/90) gives the title compound, IR (diffuse reflectance) 3306, 3301, 3270, 2962, 1676, 1667, 1663, 1645, 1638, 1627, 1615, 1550, 1537,
10 1450 and 1439 cm^{-1} ; NMR (CDCl_3) δ 0.645, 0.968, 1.20, 1.43, 1.67, 2.8, 2.97, 3.38, 3.47, 3.73, 3.87, 4.31, 6.78, 6.91, 7.23, 7.72, 7.87, 8.22 and 8.43.

EXAMPLE 588 1- tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-
15 hydroxy-3-[(3-iodobenzyl)amino]propylcarbamate
(VII)

tert-Butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, EXAMPLE 3, 1.75 g, 5.8 mmole) is mixed with isopropanol (30 mL). The reaction flask is charged
20 with 3-iodobenzylamine (VI). The reaction mixture is heated to reflux for 45 minutes, HPLC analysis indicates complete disappearance of the epoxide (V). The reaction mixture is concentrated under reduced pressure and the residue is partitioned between ethyl acetate (150 mL) and aqueous
25 hydrochloric acid (3%, 35 mL). The organic phase is separated and washed with aqueous hydrochloric acid (3%, 20 mL), bicarbonate, saline and dried over sodium sulfate. Concentration under reduced pressure gives the title compound,
M + H = 535.

30
EXAMPLE 589 1-9H-fluoren-9-ylmethyl (2R,3S)-3-(3-tert-butylloxycarbonyl)amino-4-(3,5-difluorophenyl)-2-

hydroxybutyl (3'-iodobenzyl) carbamate
hydrochloride (XXXIV)

1- tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
[(3-iodobenzyl)amino]propylcarbamate (VII, EXAMPLE 588, 2.5 g,
5 4.7 mmole) and triethylamine (0.72 ml, 5.1 mmole) in THF (10
ml) are mixed. The reaction is cooled to 0 degrees and treated
with Fmoc-Cl (1.2 g, 4.7 mmole) in THF (2 ml) via addition
funnel. After 15 minutes HPLC indicates complete disappearance
of starting material. The reaction is diluted with ethyl
10 acetate and washed with aqueous potassium bisulfate, saturated
aqueous bicarbonate, saline and dried over sodium sulfate.
Concentration under reduced pressure gives crude product which
is purified by flash chromatography, eluting with ethyl
acetate/hexane (20/80) followed by ethyl acetate to give the
15 title compound, M + H = 757.

EXAMPLE 590 1-9H-fluoren-9-ylmethyl (2R,3S)-3-amino-4-(3,5-
difluorophenyl)-2-hydroxybutyl (3-
iodobenzyl) carbamate hydrochloride (XXXV)

20

1-9H-fluoren-9-ylmethyl (2R,3S)-3-(3-t-
butyloxycarbonyl)amino-4-(3,5-difluorophenyl)-2-
hydroxybutyl (3'-iodobenzyl) carbamate hydrochloride (XXXIV,
EXAMPLE 589, 2.9 g) in hydrochloric acid/dioxane (4N, 10 ml).

25 The mixture is stirred 1 hour then slowly poured into rapidly
stirring ether (200 ml). The product is filtered and dried to
give the title compound, M + H = 657.

EXAMPLE 591 1-9H-fluoren-9-ylmethyl (2R,3S)-4-(3,5-
difluorophenyl)-2-hydroxy-3-{[5-oxo-5-(1-
30 piperidinyl)pentanoyl]amino}butyl (3-
iodobenzyl) carbamate (XXXVI)

HOBt (81 mg, 0.6 mmole) and EDC (105 mg, 0.55 mmole) are added to 1-carboxy-5-piperdinyglutaramide (IX, 100 mg, 0.5 mmole) in DMF (2 ml). The acid is activated 60 minutes then treated with 1-9H-fluoren-9-ylmethyl (2R,3S)-3-amino-4-(3,5-difluorophenyl)-2-hydroxybutyl (3-iodobenzyl) carbamate hydrochloride (XXXV, EXAMPLE 590, 300 mg, 0.43 mmole) and NMM (0.19 ml, 1.72 mmole). The reaction is stirred 3 hours then concentrated under reduced pressure. The residue is partitioned between ethyl acetate and saturated aqueous bicarbonate. The organic phases are washed with aqueous potassium bisulfate, saline, dried over sodium sulfate and finally concentrated under reduced pressure to give crude product. Purification via flash chromatography with ethyl acetate/hexane (50/50) then methanol/ethyl acetate (10/90) gives the title compound, M + H = 838.

EXAMPLE 592 1- N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-oxo-5-(1-piperidinyl)pentanamide trifluroacetate (X)

1- N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-oxo-5-(1-piperidinyl)pentanamide trifluroacetate (XXXVI, EXAMPLE 591, 240 mg, 0.29 mmole is dissolved in diethylamine (10%, 9 ml) in methylene chloride. The reaction is stirred at 20-25 degrees overnight. The next morning the reaction is concentrated under reduced pressure and the residue is redissolved in methylene chloride and purified by preparative reverse phase HPLC. The appropriate fractions are pooled and concentrated under reduced pressure and partitioned between ethyl acetate and saline. The organic phase is separated and dried over sodium sulfate and concentrated to give the title compound, M + H = 614.

EXAMPLE 593 5-(Aminosulfonyl)-N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N-dipropylisophthalamide (X)

O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
 5 hexafluorophosphate (HATU, 0.0928 g, 0.244 mmol) is added to a mixture of, 3-(aminosulfonyl)-5-[(dipropylamino)-carbonyl]benzoic acid (XXXIX, PREPARATION 13, 0.0800 g, 0.244 mmol) and (2R,3S)-3-amino-1-[(3-methoxybenzyl)-amino]-4-phenyl-2-butanol (VIII, EXAMPLE 175, 0.0732 g, 0.244 mmol) in dry DMF
 10 (3 mL). The mixture is stirred for 18 hours at 20-25 degrees, and then partitioned between ethyl acetate and water. The organic phase is separated and washed with saline, dried over anhydrous sodium sulfate, filtered and concentrated. The concentrate is column chromatographed (silica gel;
 15 methanol/dichloromethane, 5/95) to give the title compound, MS (ESI+) for C₃₂H₄₂N₄O₆S m/z (M+H)⁺ = 611.5; HRMS (FAB) calculated for C₃₂H₄₂N₄O₆S + H₁ = 611.2903, found = 611.2904.

EXAMPLE 594 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropyl-5-(1-pyrrolidinylsulfonyl)isophthalamide (X)

Following the general procedure of EXAMPLE 593 and making non-critical variations but using 3-[(dipropylamino)carbonyl]-5-(1-pyrrolidinylsulfonyl)benzoic acid (XXXIX, PREPARATION 15,
 25 the title compound is obtained, ES (ESI+) for C₃₆H₄₈N₄O₆S m/z (M+H)⁺ = 665.6; HRMS (FAB) calculated for C₃₆H₄₈N₄O₆S + H₁ = 665.3372, found = 665.3393.

EXAMPLE 595 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(methylamino)sulfonyl]-N³,N³-dipropylisophthalamide (X)

Following the general procedure of EXAMPLE 593 and making non-critical variations but using 3-[(dipropylamino)carbonyl]-

5-[(methylamino)-sulfonyl]benzoic acid (XXXIX, PREPARATION 17, the title compound is obtained, MS (ESI+) for $C_{33}H_{44}N_4O_6S$ m/z $(M+H)^+ = 625.5$.

- 5 EXAMPLE 596 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(dimethylamino)sulfonyl]- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLE 593 and making
10 non-critical variations but using 3-[(dimethylamino)sulfonyl]-5-[(dipropylamino)carbonyl]-benzoic acid (XXXIX, PREPARATION 19), the title compound is obtained, MS (ESI+) for $C_{34}H_{46}N_4O_6S$ m/z $(M+H)^+ = 639.5$

- 15 EXAMPLES 597-619

Following the general procedure of EXAMPLES 589 through 592 but starting with (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII, EXAMPLE 175) and using the amide forming agent (IX) of Column A, the Substituted
20 Amine (X) of Column B is obtained.

EX #	Column A Amide Forming Agent (IX)	Column B Substituted Amine (X)	MS Data M + H
597	2-methyl-3-(methylsulfonyl)propanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-methyl-3-(methylsulfonyl)propanamide hydrochloride	449
59	3-	N-{(1S,2R)-1-benzyl-	435

8	(methylsulfonyl)propanoic acid	2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-(methylsulfonyl)propanamide hydrochloride	
599	2-amino-1,3-thiazole-4-carboxylic acid hydrochloride	2-amino-N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-1,3-thiazole-4-carboxamide dihydrochloride	427
600	5-(methylsulfonyl)pentanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(methylsulfonyl)pentanamide hydrochloride	463
601	4-anilino-4-oxobutanoic acid	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ⁴ -phenylsuccinamide hydrochloride	476
602	(2R)-4-amino-2,3,3-trimethyl-4-oxobutanoic acid	(3R)-N ⁴ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2,2,3-trimethylbutanediamide hydrochloride	442
603	3-	N-{(1S,2R)-1-benzyl-	520

3	[(dipropylamino) sulfonyl] propanoic acid	2-hydroxy-3-[(3-methoxybenzyl) amino]propyl}-3-[(dipropylamino) sulfonyl]-propanamide hydrochloride	
604	5-(dipropylamino)-5-oxopentanoic acid	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino]propyl}-N ⁵ ,N ⁵ -dipropylpentanediamide hydrochloride	498
605	4-oxo-4-(1-piperidinyl)butanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino]propyl}-4-oxo-4-(1-piperidinyl)butanamide hydrochloride	468
606	4-(dipropylamino)-4-oxobutanoic acid	N-1--{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino]propyl}-N ⁴ ,N ⁴ -dipropylsuccinamide hydrochloride	484
607	5-oxo-5-(1-piperidinyl)pentanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino]propyl}-5-oxo-5-(1-piperidinyl)pentanamide hydrochloride	482

608	5-anilino-5-oxopentanoic acid	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ⁵ -phenylpentanediamide hydrochloride	490
609	3,3-dimethyl-4-oxo-4-(1-piperidinyl)butanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3,3-dimethyl-4-oxo-4-(1-piperidinyl)butanamide hydrochloride	496
610	4-(isopentylsulfonyl)butanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-(isopentylsulfonyl)butanamide hydrochloride	505
611	4-(dipropylamino)-2,2-dimethyl-4-oxobutanoic acid	N ¹ -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2,2-dimethyl-N ⁴ ,N ⁴ -dipropylsuccinamide hydrochloride	512
612	4-[(dipropylamino)sulfonyl]butanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-	534

		[(dipropylamino)sulfonyl] butanamide hydrochloride	
61 3	4-[(methylanilino)sulfonyl] butanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-[(methylanilino)sulfonyl]butanamide hydrochloride	540
61 4	3-[(methylanilino)sulfonyl]propanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-[(methylanilino)sulfonyl]propanamide	526
61 5	Acetic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}acetamide hydrochloride	343
61 6	3-(isopentylsulfonyl)propanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-(isopentylsulfonyl)propanamide hydrochloride	491
61 7	5-oxo-5-(1-piperidiny)pentanoic acid	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-	614

		iodobenzyl) amino] propyl}-5-oxo-5-(1-piperidinyl) pentanamide trifluoroacetate	
61 8	5-oxo-5-(1-piperidinyl) pentanoic acid	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl) amino] propyl}-5-oxo-5-(1-piperidinyl) pentanamide trifluoroacetate	578
61 9	3-[(dipropylamino) sulfonyl] propanoic acid	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl) amino] propyl}-3-[(dipropylamino) sulfonyl] propanamide	652

EXAMPLE 620 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-ethyl- N^3 , N^3 -dipropylisophthalamide (X)

5

Diethyl cyanophosphonate (0.132 mL, 0.870 mmol) is added to a mixture of 3-[(dipropylamino) carbonyl]-5-ethylbenzoic acid (IX, PREPARATION 21, 0.200 g, 0.720 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl) amino]-4-phenyl-2-butanol (VIII, EXAMPLE 175, 0.216 mg, 0.720 mmol), and triethylamine (0.121 mL, 0.870 mmol) in dichloromethane (3 mL). The mixture was stirred for 1 hour at 20-25 degrees C. Dichloromethane is then removed under reduced pressure. The residue is partitioned between ethyl acetate and water. The organic phase is

separated and is washed with saline, dried over anhydrous sodium sulfate, filtered and concentrated. The concentrate is column chromatographed (silica gel; methanol/dichloromethane, 5/95) to give the title compound, MS (ESI+) for $C_{34}H_{45}N_3O_4$ m/z (M+H)⁺ = 560.4; HOURMS (FAB) calculated for $C_{34}H_{45}N_3O_4 + H$ = 560.3488, found = 560.3487.

EXAMPLE 621 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-isobutyl- N^3,N^3 -dipropylisophthalamide

Following the general procedure of PREPARATIONS 20 and 21 and making non-critical variations but using isobutylboronic acid, and following the general procedure of EXAMPLE 620 but using the 5-isobutylisophthalic acid (IX), the title compound is obtained, MS (ESI+) for $C_{36}H_{49}N_3O_4$ m/z (M+H)⁺ = 588.6; HRMS (FAB) calculated for $C_{36}H_{49}N_3O_4 + H_1$ = 588.3801, found = 588.3810.

EXAMPLE 622 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-tert-butyl- N^3,N^3 -dipropylisophthalamide

Following the general procedure of PREPARATIONS 20 and 21 and making non-critical variations but using tert-butylboronic acid, and following the general procedure of EXAMPLE 620 but using the tert-butylphthalic acid (IX), the title compound is obtained, MS (ESI+) for $C_{36}H_{49}N_3O_4$ m/z (M+H)⁺ = 588.5; HRMS (FAB) calculated for $C_{36}H_{49}N_3O_4 + H_1$ = 588.3801, found = 588.3791.

EXAMPLE 623 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-cyano- N^3 -propylisophthalamide (X)

Following the general procedure of EXAMPLE 586 and making non-critical variations, but using tert-butyl (2R,3S)-3-[(3-cyano-5-[(propylamino)carbonyl]benzoyl)amino]-2-hydroxy-4-

phenylbutyl (3-methoxybenzyl) carbamate (XXXVI) the title compound is obtained, $M + H = 515.1$.

EXAMPLES 624-628

- 5 Following the general procedure of EXAMPLE 587 and making non-critical variations, but using the appropriate amines (VIII) and amide forming agents (IX), for example PREPARATIONS 6 and 19, the titled compounds are obtained.

EXAMPLE	Substituted Amine (X)	M + H =
624	$N^1 - \{ (1S, 2R) - 1 - (3, 5\text{-difluorobenzyl}) - 2\text{-hydroxy-3-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl} \} - N^3, N^3\text{-dipropyl-1, 3, 5-benzenetricarboxamide}$	611.0
625	$N^1 - \{ (1S, 2R) - 1\text{-benzyl-2-hydroxy-3-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl} \} - N^3, N^3\text{-dimethyl-}N^5, N^5\text{-dipropyl-1, 3, 5-benzenetricarboxamide}$	603.0
626	$N^1 - \{ (1S, 2R) - 3\text{-amino-1-benzyl-2-hydroxypropyl} \} - N^3, N^3\text{-dipropyl-1, 3, 5-benzenetricarboxamide}$	455.1
627	$N^1 - \{ (1S, 2R) - 1\text{-benzyl-2-hydroxy-3-(isopentylamino)propyl} \} - N^3, N^3\text{-dipropyl-1, 3, 5-benzenetricarboxamide}$	525.6
628	$N^1 - \{ (1S, 2R) - 1\text{-benzyl-2-hydroxy-3-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl} \} - N^3\text{-propyl-1, 3, 5-benzenetricarboxamide}$	533.1

- 10 EXAMPLE 629 $N - \{ (1S, 2R) - 1\text{-Benzyl-2-hydroxy-3-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl} \} - 3\text{-[butyryl(propyl)amino]-5-methylbenzamide (X)}$

Following the procedure of EXAMPLE 570 and making non-critical variations, diethyl cyanophosphonate (0.0760 mL, 0.550 mmol) is added to a mixture of 3-[butyryl(propyl)amino]-5-methylbenzoic acid (IX, 0.120 g, 0.460 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII, 0.137 g, 0.460 mmol), and triethylamine (0.0760 mL, 0.550 mmol) in dichloromethane (5 mL). The mixture is stirred for 1 hour at 20-25 degrees C. Dichloromethane is then removed under reduced pressure. The residue is partitioned between ethyl acetate and water. The organic is separated, is washed with saline, dried over anhydrous sodium sulfate, filtered and concentrated. The concentrate is column chromatographed (silica gel; methanol/dichloromethane, 5/95) to give the title compound, NMR (400 MHz, CDCl₃) δ 7.09, 4.15, 3.80, 3.79, 3.60, 3.02, 2.84, 2.36, 1.94, 1.56, 1.49, 0.87 and 0.81; MS (ESI+) for C₃₃H₄₃N₃O₄ m/z (M+H)⁺ = 546.3; HRMS (FAB) calculated for C₃₃H₄₃N₃O₄+H = 546.3331, found = 546.3331.

EXAMPLE 630 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)aminopropyl]-1-propyl-1H-indole-6-carboxamide (X)

Following the general procedure of EXAMPLE 539 and making non-critical variations, the title compound is obtained, IR (diffuse reflectance) 3330, 3314, 2960, 2952, 2931, 2873, 1621, 1599, 1525, 1499, 1467, 1353, 1283, 1253 and 786 cm⁻¹.

EXAMPLE 631 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)aminopropyl]-1-propyl-1H-indole-6-carboxamide (X)

Following the general procedure of EXAMPLE 539 and making non-critical variations, the title compound is obtained, IR (diffuse reflectance) 3289, 1627, 1621, 1595, 1531, 1525, 1520, 1507, 1466, 1458, 1450, 1349, 1317, 1252 and 1117 cm⁻¹.

EXAMPLES 633-708

Following the general procedures of CHART A as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-5 631 and making non-critical variations and using the appropriate reagents, the substituted amines (X) of EXAMPLES 633-708 are obtained.

EXAMPLE	Substituted Amine (X)	MH ⁺ =
633	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3,4-dimethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	580
634	N ¹ -[(1S,2R)-3-[(3-aminobenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	567
635	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}octanamide	559
636	N ³ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({1-methyl-1-[3-(trifluoromethyl)phenyl]ethyl}amino)propyl]-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	635
637	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({1-methyl-1-[3-(trifluoromethyl)phenyl]ethyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	648
638	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino]propyl)-5-	594

	methyl-N ³ ,N ³ -dipropylisophthalamide	
639	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(1R)-2,3-dihydro-1H-inden-1-ylamino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	578
640	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-3-methylbenzamide	551
641	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(1H-isoindol-3-ylamino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	577
642	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1R,2S,5R)-2-isopropyl-5-methylcyclohexyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	600
643	N ¹ ,N ¹ -diallyl-5-chloro-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)amino]propyl}isophthalamide	597
644	5-chloro-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)amino]propyl}-N ³ ,N ³ -bis(2-methoxyethyl)isophthalamide	633
645	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-phenylcyclopentyl)amino]propyl}-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	593
646	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	580

647	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(dimethylamino)benzyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	595
648	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(4,5-dimethyl-2-furyl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	570
649	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-phenylcyclopentyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	606
650	N ¹ -[(1S,2R)-3-(cyclopropylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	502
651	N ¹ -[(1S,2R)-3-[(cyclopropylmethyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	516
652	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-N ⁵ ,N ⁵ -dipropylpentanediamide	630
653	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(2-furylmethyl)amino]-2-hydroxypropyl}-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	529
654	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(tetrahydro-2-furanylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	546
655	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-	565

	phenylcyclopropyl) amino] propyl} -N ⁵ , N ⁵ - dipropyl-3,5-pyridinedicarboxamide	
656	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(2-oxo-3- azepanyl) amino] propyl}-5-methyl-N ³ , N ³ - dipropylisophthalamide	573
657	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(3-methyl-2- furyl) methyl] amino} propyl)-5-methyl- N ³ , N ³ -dipropylisophthalamide	556
658	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(2S)-tetrahydro-2- furanylmethyl] amino} propyl)-5-methyl- N ³ , N ³ -dipropylisophthalamide	546
659	5-chloro-N ¹ -{(1S,2R)-1-(3,5- difluorobenzyl)-2-hydroxy-3-[(1-methyl- 1-phenylethyl) amino] propyl}-N ³ , N ³ -di(2- propynyl) isophthalamide	593
660	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(3- isopropenylbenzyl) amino] propyl}-5- methyl-N ³ , N ³ -dipropylisophthalamide	592
661	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(2- propoxyethyl) amino] propyl}-5-methyl- N ³ , N ³ -dipropylisophthalamide	548
662	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- (hexylamino)-2-hydroxypropyl]-5-methyl- N ³ , N ³ -dipropylisophthalamide	546
663	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-[(3-iodobenzyl) amino] propyl}- 4-(3-methyl-5-oxo-4,5-dihydro-1H-	633

	pyrazol-1-yl)benzamide	
664	methyl 4-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino}methyl)benzoate	610
665	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-methoxyethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	520
666	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(5-isoxazolylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	543
667	(1R,2R)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide	628
668	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2S)-tetrahydro-2-furanylmethyl]amino}propyl)-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	533
669	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	582
670	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropylbenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	594
671	4-(butyrylamino)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-	622

	iodobenzyl) amino]propyl} benzamide	
672	N ¹ -[(1S,2R)-3-[(3-amino-3-oxopropyl) amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	533
673	N ³ -[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide 1-oxide	555
674	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl) amino]propyl}-5-ethynyl-N ³ ,N ³ -dipropylisophthalamide	688
675	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(7-oxabicyclo[2.2.1]hept-2-ylmethyl) amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	572
676	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl) amino]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	576
677	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-methyl-1,3-thiazol-5-yl)methyl] amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	573
678	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethyl-1,3-thiazol-5-yl)methyl] amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	587
679	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3R)-2-oxoazepanyl] amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	573

680	N ¹ -[(1S,2R)-3-(cyclobutylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	516
681	N ¹ -[(1S,2R)-3-(butylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-ethynyl-N ³ ,N ³ -dipropylisophthalamide	528
682	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-ethynyl-N ³ ,N ³ -dipropylisophthalamide	590
683	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-(5-hexynylamino)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	542
684	N ³ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-methyl-2-furyl)methyl]amino}propyl)-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide	543
685	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)amino]propyl}-N ⁵ ,N ⁵ -dipropylpentanediamide	532
686	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(2-furyl)-1-methylethyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	570
687	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(3-isobutyl-5-isoxazolyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	599
688	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-isobutyl-1,3-thiazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -	615

	dipropylisophthalamide	
689	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl]-3-[(dipropylamino)sulfonyl]propanamide	554
690	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-[(2-phenylethyl)amino]propyl]-N ³ ,N ³ -dipropylisophthalamide	516
691	N ¹ -[(1S,2R)-1-benzyl-3-{[2-(2-chlorophenyl)ethyl]amino}-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	551
692	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-{[3-(2-oxo-1-pyrrolidinyl)propyl]amino}propyl]-N ³ ,N ³ -dipropylisophthalamide	537
693	N ¹ -[(1S,2R)-1-benzyl-3-[(cyclohexylmethyl)amino]-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	508
694	N ¹ -[(1S,2R)-1-benzyl-3-(cyclopropylamino)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	452
695	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-[(2-oxo-3-azepanyl)amino]propyl]-N ³ ,N ³ -dipropylisophthalamide	523
696	N-[(1S,2R)-3-(benzylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-3-(butylsulfonyl)benzamide	531
697	N ¹ -[(1S,2R)-1-benzyl-3-[(2-ethylhexyl)oxy]ethyl]amino)-2-hydroxypropyl]-N ³ ,N ³ -	568

	dipropylisophthalamide	
698	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[(1S,2R)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	544
699	N ¹ -((1S,2R)-1-benzyl-2-hydroxy-3-{[1-(4-hydroxyphenyl)ethyl]amino}propyl)-N ³ ,N ³ -dipropylisophthalamide	532
700	N ¹ -[(1S,2R)-1-benzyl-3-(cycloheptylamino)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	508
701	N ¹ -{(1S,2R)-1-benzyl-3-[(1,1'-biphenyl)-2-ylmethyl]amino}-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	578
702	N ¹ -{(1S,2R)-1-benzyl-3-[(2-fluorobenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	520
703	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-(dimethylamino)benzamide	484
704	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-1-naphthamide	491
705	N ¹ -[(1S,2R)-1-benzyl-3-({2-[(5-{[(dimethylamino)methyl]-2-furyl)methyl]sulfanyl}ethyl]amino)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	609
706	N ¹ -[(1S,2R)-1-benzyl-3-({2-[(2-chloro-6-	615

	fluorobenzyl)sulfanyl}ethyl}amino)-2-hydroxypropyl]-N ³ ,N ³ -dipropylisophthalamide	
707	N ¹ -[(1S,2R)-3-[[[1,1'-biphenyl]-4-ylmethyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	628
708	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(1-naphthylamino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	588

EXAMPLES 709-737

Following the general procedure of CHART D and EXAMPLES 165-169 and making non-critical variations and using the appropriate reagents, the substituted amines (X) of EXAMPLES 709-737 are obtained.

EXAMPLE	Substituted Amine (X)	MH ⁺
709	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1H-imidazol-5-ylmethyl)amino]propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	542
710	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-phenyl-1H-imidazol-5-yl)methyl]amino]propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	618
711	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1H-imidazol-2-yl)methyl]amino]propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	556
712	N ¹ -[(1S,2R)-3-[[[2-butyl-4-chloro-1H-imidazol-5-yl)methyl]amino]-1-(3,5-	633

	difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
713	N ¹ -[(1S,2R)-3-{[(6-chloroimidazo[2,1-b][1,3]thiazol-5-yl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	633
714	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1-methyl-1H-benzimidazol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	606
715	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-hydroxy-1-naphthyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	618
716	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(4-oxo-4H-chouromen-3-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	620
717	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(1,5-dimethyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-4-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	662
718	N ¹ -[(1S,2R)-3-({[5-cyano-6-(methylsulfanyl)-2-pyridinyl]methyl}amino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	624
719	[5-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino)methyl)-2-	614

	furyl)methyl acetate	
720	N ¹ -[(1S,2R)-3-[(1-benzofuran-3-ylmethyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	592
721	methyl 4-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino}methyl)-1-methyl-1H-pyrrole-2-carboxylate	613
722	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({[1-(phenylsulfonyl)-1H-pyrrol-2-yl]methyl}amino)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	681
723	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1-methyl-1H-pyrrol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	555
724	N ¹ -[(1S,2R)-3-{[(4-chloro-1-methyl-1H-pyrazol-3-yl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	591
725	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	646
726	N ¹ -[(1S,2R)-3-{[(5-chloro-3-methyl-1-phenyl-1H-pyrazol-4-yl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	667
727	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-	618

	hydroxy-3-{[(3-phenyl-1H-pyrazol-4-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
728	N ¹ -[(1S,2R)-3-{[(5-chloro-2-thienyl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	593
729	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(3-phenoxy-2-thienyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	650
730	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-quinolinylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	603
731	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(2-quinolinylmethyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	603
732	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1-methyl-1H-indol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	605
733	N ¹ -[(1S,2R)-3-{[(1-benzyl-1H-indol-3-yl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	681
734	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1-methyl-1H-indol-3-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	605
735	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-	745

	hydroxy-3-[(1-[(4-methylphenyl)sulfonyl]-1H-indol-3-yl)methyl]amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	
736	N ¹ -[(1S,2R)-3-{[(2-butyl-1H-imidazol-5-yl)methyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	598
737	methyl 3-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino}methyl)-1H-indole-6-carboxylate	649

EXAMPLE 738 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino)carbonyl]-5-[butyl(butyryl)amino]benzyl diethyl phosphate (X)

5 Following the general procedure of CHART L and EXAMPLE 620 and making non-critical variations, the title compound is obtained, HRMS (FAB) = 712.3749.

EXAMPLE 739 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(cyanomethyl)-N³,N³-dipropylisophthalamide (X)

10 Step 1. A mixture of diethyl 1,3,5-benzenetricarboxylate (5.2 g) and borane methylsulfide complex (6.1 g) is stirred in THF (150 mL) at 20-25 degrees C overnight. The mixture is then
 15 treated with methanol, concentrated to dryness, and chroumatographed (silica gel) to give diethyl 5-(hydroxymethyl)isophthalate. Diethyl 5-(hydroxymethyl)isophthalate (3.4 g) is hydrolyzed in ethanol and water with lithium hydroxide monohydrate (0.57 g) at 20-25

degrees C for 3.5 hours at which time the solvents are removed under reduced pressure. Water (100 mL) is added and the mixture is acidified to pH = 4 with concentrated hydrochloric acid. The mixture is extracted with ethyl acetate and dried over magnesium sulfate, filtered, and concentrated to give 3-(ethoxycarbonyl)-5-(hydroxymethyl)benzoic acid, high resolution MS MH⁺ = 225.0769. 3-(Ethoxycarbonyl)-5-(hydroxymethyl)benzoic acid (2.3 g), EDC (3.0 g), 1-HOBT (2.1 g), diisopropylethylamine (2.7 mL), dipropyl amine (2.8 mL), and DMF (50 mL) are stirred at 20-25 degrees C overnight. The mixture is then partitioned between ethyl acetate, water, and saline. The organic phase is separated and dried over magnesium sulfate, filtered, and concentrated. Chromatography (silica gel) gives ethyl 3-[(dipropylamino)carbonyl]-5-(hydroxymethyl)benzoate, NMR (CDCl₃) δ 0.77, 1.0, 1.4, 1.6, 1.7, 3.2, 3.5, 4.4, 4.8, 7.6, 8.0 and 8.1.

Step 2. A mixture of ethyl 3-[(dipropylamino)carbonyl]-5-(hydroxymethyl)benzoate (1.5 g) and phosphorous tribromide (0.95 mL) is stirred in dichloromethane (10 mL) and heated at 50 degrees C for 4 hours and then cooled and partitioned between dichloromethane and water. The organic phase is separated and washed with aqueous sodium bicarbonate and then dried over magnesium sulfate and taken to dryness to give ethyl 3-(bromomethyl)-5-[(dipropylamino)carbonyl]benzoate, high resolution MS MH⁺ = 370.1020. Ethyl 3-(bromomethyl)-5-[(dipropylamino)carbonyl]benzoate (1.4 g) and sodium cyanide (0.2 g) are stirred in dry DMSO (25 mL) at 20-25 degrees C for 3.5 hours and the mixture is then partitioned between ethyl acetate, water and saline. The organic layer is separated and dried over magnesium sulfate and taken to dryness under reduced pressure to give ethyl 3-(cyanomethyl)-5-[(dipropylamino)carbonyl]benzoate. Ethyl 3-(cyanomethyl)-5-[(dipropylamino)carbonyl]benzoate (0.6 g) is hydrolyzed with

lithium hydroxide monohydrate (0.1 g) in ethanol and water at 20-25 degrees C overnight and then added to water (50 mL). The pH is adjusted to 4 using concentrated hydrochloric acid and the mixture is partitioned between ethyl acetate, water, and saline. The organic phase is separated and dried over magnesium sulfate and taken to dryness under reduced pressure to give 3-(cyanomethyl)-5-[(dipropylamino)carbonyl]benzoic acid, MS M+H = 287.2.

Step 3. A mixture of 3-(cyanomethyl)-5-[(dipropylamino)carbonyl]benzoic acid (IX, 0.13 g), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII, 0.14 g), HATU (0.17 g), and dichloromethane (10 mL) is stirred at 40 degrees C overnight. After cooling, the mixture is washed with water and the organic phase is separated and dried over magnesium sulfate and taken to dryness under reduced pressure. Chromatography (silica gel) gives the title compound, M + H = 571.2

EXAMPLE 740 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(hydroxymethyl)-N³,N³-dipropylisophthalamide (X)

Following the procedure of CHART P and EXAMPLE 739 and making non-critical variations but using 3-[(dipropylamino)carbonyl]-5-(hydroxymethyl)benzoic acid (IX) and (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII), the title compound is obtained, HRMS (FAB) = 615.3571.

EXAMPLE 741 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-ethynyl-N³,N³-dipropylisophthalamide (X)

Step 1: A mixture of methyl 3-bromo-5-[(dipropylamino)carbonyl]benzoate (XXI, 200 mg, 0.58 mmol),

PdCl₂(Ph₃P)₂ (16 mg, 0.03 mol %) and copper (I) iodide (6 mg, 0.05 mol %) in triethylamine (1.2 mL) is heated to reflux. (Trimethylsilyl) acetylene (100 microliter, 0.7 mmol) is added, and the mixture stirred for 3 hours, cooled to 20-25 degrees, 5 diluted with water (20 mL), and extracted with chloroform (3 x 15 mL). The combined organic extracts are washed with saline (20 mL), dried over sodium sulfate and concentrated under reduced pressure to give methyl 3-[(dipropylamino)carbonyl]-5-ethynylbenzoate (XXXII, 185.5 mg), NMR (300 MHz, CDCl₃): δ 7.95, 10 7.75, 7.43, 3.74, 3.25, 2.95, 1.49, 1.34, 0.79, 0.56 and 0.06.

Step 2: To a stirred mixture of the protected methyl 3-[(dipropylamino)carbonyl]-5-ethynylbenzoate (XXXII, Step 1, 185.3 mg, 0.49 mmol) in methanol (2.5 mL) is added a mixture of potassium hydroxide (2.9 mL of a 1 M mixture in water, 2.9 15 mmol). The reaction mixture is stirred for 4 hours diluted with chloroform (40 mL), the phases are separated and the organic phase is concentrated under reduced pressure to give 3-[(dipropylamino)carbonyl]-5-ethynylbenzoic acid, NMR (300 MHz, CDCl₃): δ 8.22, 8.05, 7.71, 3.48, 3.17, 3.16, 1.71, 1.55, 1.00 20 and 0.78.

Step 3: To a stirred mixture of 3-[(dipropylamino)carbonyl]-5-ethynylbenzoic acid (70 mg, 0.24 mmol) in DMF (2.5 mL) is added (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 25 81 mg, 0.24 mmol), HOBt (36 mg, 0.26 mmol) and diisopropylethylamine (170 microliter, 0.96 mmol). To this reaction mixture is added EDC (51mg, 0.26 mmol) and the reaction mixture is stirred overnight. The reaction mixture is diluted with ethyl acetate (30 mL), washed with water (3 x 50 30 mL), hydrochloric acid (1 N, 30 mL), saturated sodium bicarbonate (30 mL), saline (30 mL), dried over sodium sulfate and concentrated under reduced pressure. Purification by flash chromatography (silica, ethyl acetate to methanol/chloroform,

1/10) gives the title compound, IR (KBr): 3276, 2956, 2921, 1610, 1450 and 1264 cm^{-1} ; ESI-MS (m/z) $[M + H]^+ = 556$.

EXAMPLE 742 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}- N^3,N^3 -dipropyl-5-prop-1-ynylisophthalamide (X)

Following the general procedure of EXAMPLE 741 and making non-critical variations but using propyne in place of (trimethylsilyl) acetylene and using (2R,3S)-3-amino-1-[(3-iodobenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII) in place of (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII), the title compound is obtained, IR (ATR): 3305, 2930, 2872, 1613 and 1537 cm^{-1} ; ESI-MS (m/z) $[M+H]^+ = 666$.

EXAMPLE 743 N^1 -((1S,2R)-1-benzyl-2-hydroxy-3-{[3-(trifluoromethyl)benzyl]amino}propyl)-5-ethynyl- N^3,N^3 -dipropylisophthalamide (X)

Step 1: A mixture of tert-butyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V, 2.3 g, 8.7 mmol) and 3-(trifluoromethyl)benzylamine (VI, 1.9 mL, 13.1 mmol) in 2-propanol (70 mL) is heated at reflux for 4 hours. The reaction mixture is cooled to 20-25 degrees and concentrated under reduced pressure to give tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[3-(trifluoromethyl)benzyl]amino}propylcarbamate (VII, 3.1 g) as a solid, ESI-MS (m/z) $[M + H]^+ = 439$.

Step 2: A mixture of tert-butyl (1S,2R)-1-benzyl-2-hydroxy-3-{[3-(trifluoromethyl)benzyl]amino}propylcarbamate (VII, step 1, 2.5 g, 5.7 mmol) and hydrochloric acid (29 mL of a 4.0 M mixture in dioxane, 114 mmol) is stirred at 20-25 degrees. A precipitate forms and is collected by filtration, washed with ether, and dried under reduced pressure to give (2R,3S)-3-amino-4-phenyl-1-{[3-(trifluoromethyl)benzyl]amino}-

2-butanol dihydrochloride (VIII, 2.13 g), ESI-MS (m/z) $[M +]^+ = 339$.

Step 3: A mixture of 3-[(dipropylamino)carbonyl]-5-ethynylbenzoic acid (IX, 231 mg, 0.8 mmol), (2R,3S)-3-amino-4-phenyl-1-{[3-(trifluoromethyl)benzyl]amino}-2-butanol dihydrochloride (VIII, Step 2, 493.5 mg, 1.2 mmol) HOBt (162 mg, 1.2 mmol), and diisopropylethylamine (832 Micro Liter, 4.8 mmol) is stirred in methylene chloride (4 mL) for 15 minutes EDC (206 mg, 1.2 mmol) is added and the reaction mixture is stirred overnight. The reaction mixture is diluted with water, and extracted with methylene chloride (3 x 25 mL). The organic phase is washed with hydrochloric acid (1N, 25 mL), saturated sodium bicarbonate (25 mL), saline dried over sodium sulfate and concentrated under reduced pressure. Purification by flash column chromatography (silica, 100% ethyl acetate to methanol/chloroform, 1/9) gives title compound, IR (ATR): 3302, 2963, 2932 and 1615 cm^{-1} ; MS (m/z) $[M + H]^+ = 549$.

EXAMPLE 744 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-ethynyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLE 744 and making non-critical variations but using 3-iodobenzylamine hydrochloride salt (VI), the title compound is obtained, IR (ATR) 3295, 2960, 2927 and 1616 cm^{-1} , APCI-MS (m/z) $[M + H]^+ = 652$.

EXAMPLE 745 N^1 -{(1S,2R)-1-benzyl-3-[(3-fluorobenzyl)amino]-2-hydroxypropyl}-5-ethynyl- N^3,N^3 -dipropylisophthalamide (X)

Following the general procedure of EXAMPLE 744 and making non-critical variations but using 3-fluorobenzylamine (VI), the

title compound is obtained, IR (ATR): 3217, 2961, 2918 and 1615 cm^{-1} ; APCI-MS (m/z) $[M + H]^+ = 544$.

EXAMPLE 746 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^3,N^3 -dipropyl-5-(8-quinolinyl)isophthalamide (X)

Step 1: A mixture of methyl 3-[(dipropylamino)carbonyl]benzoate (XLVIII, 200 mg, 0.58 mmol), 8-quinolineboronic acid (200.6 mg, 1.2 mmol), sodium carbonate (870 Micro Liter of a 2 M mixture in water, 1.74 mmol) in toluene (6 mL) is degassed under reduced pressure for 15 minutes and purged with argon. Palladium tetrakis(triphenylphosphine) (139 mg, 0.12 mmol) is added and the reaction mixture is degassed under reduced pressure for 15 minutes and purged with argon. The reaction mixture is heated at reflux overnight, cooled to 20-25 degrees C and diluted with chloroform. The organic phase is separated and washed with water (3 x 50 mL), and saline, dried over sodium sulfate and concentrated under reduced pressure. Purification by flash column chromatography (silica, ethyl acetate/hexanes, 1.3/1) gives methyl 3-[(dipropylamino)carbonyl]-5-(8-quinolinyl)benzoate (XLIX, 176 mg), NMR (300 MHz, CDCl_3): δ 8.91, 8.42, 8.21, 8.09, 7.95, 7.86, 7.77, 7.64, 3.94, 3.49, 3.34, 1.64, 0.99 and 0.84.

Step 2: To a mixture of methyl 3-[(dipropylamino)carbonyl]-5-(8-quinolinyl)benzoate (XLIX, step 1, 175.5 mg, 0.45 mmol) in methanol (2 mL) is added lithium hydroxide (32.3 mg, 1.4 mmol) and water (500 microliter). After stirring overnight, the reaction mixture is partitioned between ethyl acetate (10 mL) and water (10 mL). The aqueous phase is separated and acidified with hydrochloric acid (1N), and extracted with chloroform (3 x 40 mL). The organic phase is washed with saline, dried (sodium sulfate) and concentrated

under reduced pressure to give 3-[(dipropylamino)carbonyl]-5-(8-quinolinyl)benzoic acid (IX - L, 130 mg), NMR (300 MHz, CD₃OD) δ 8.84, 8.39, 8.35, 8.05, 7.96, 7.90, 7.87, 7.79, 7.68, 3.50, 3.37, 1.76-1.61, 0.99 and 0.84.

5 Step 3: A mixture of 3-[(dipropylamino)carbonyl]-5-(8-quinolinyl)benzoic acid (IX - L, Step 2, 130 mg, 0.35 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 117 mg, 0.35 mmol), HOBt (70 mg, 0.52 mmol) and diisopropylethylamine (241 microliter, 1.4 mmol) in
10 methylene chloride (2 mL) is stirred for 15 minutes EDC (89 mg, 0.52 mmol) is added and the reaction mixture is stirred overnight. The reaction mixture is diluted with water and extracted with methylene chloride (3 x 25 mL). The organic phase is washed with hydrochloric acid (1N, 25 mL), saturated
15 sodium bicarbonate (25 mL), saline, dried (sodium sulfate), and concentrated under reduced pressure. Purification by flash column chromatography (silica; methanol/chloroform, 1/9) gives the title compound, IR (NaCl): 3301, 2916, 2365 and 1613 cm⁻¹; APCI-MS (m/z) [M + H]⁺ = 659.

20

EXAMPLE 747 N³-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4'-methoxy-N⁵,N⁵-dipropyl[1,1'-biphenyl]-3,5-dicarboxamide hydrochloride (X)

Step 1: A mixture of 4-methoxyphenyl boronic acid (463 mg, 3.05 mmol), 3-bromo-5-[(dipropylamino)carbonyl]benzoic acid (XLVIII, 1.02 g, 3.05 mmol), and potassium phosphate (1.29 g, 6.10 mmol) in 1,2-dimethoxyethane (10 mL) and water (5 mL) is degassed with argon for 15 minutes
25 Bis(triphenylphosphine)palladium (II) chloride (21 mg, 0.03 mmol) is added, the reaction mixture is degassed again with argon, and heated at 85 degrees C overnight. The reaction mixture is cooled to 20-25 degrees C, and passed through a plug of diatomaceous earth.
30

The filtrate is acidified to pH = 4 with hydrochloric acid (1N) and extracted with ethyl acetate. The organic phase is washed with water and saline and dried (magnesium sulfate). The product is purified by flash column chromatography (silica gel; ethyl acetate/acetic acid, 99/1) to give 5-[(dipropylamino)carbonyl]-4'-methoxy[1,1'-biphenyl]-3-carboxylic acid (IX - L, 667 mg), ESI-MS (m/z) $[M + H]^+ = 356$.

Step 2: A mixture of 5-[(dipropylamino)carbonyl]-4'-methoxy[1,1'-biphenyl]-3-carboxylic acid (IX - L, step 1, 316 mg, 0.89 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 332 mg, 0.89 mmol), HOBt (181 mg, 1.34 mmol), and *N*-methylmorpholine (0.37 g, 3.56 mmol) in methylene chloride (8 mL) and dimethylformamide (2 mL) is stirred at 20-25 degrees for 15 minutes. EDC (257 mg, 1.34 mmol) is added and the reaction mixture is stirred for 4.5 hours. The reaction mixture is partitioned between methylene chloride and water. The organic phase is washed with hydrochloric acid (1N), water, and saline, dried (magnesium sulfate), and concentrated. The concentrate is dissolved in a minimum of methanol, treated with hydrochloric acid (3 mL of a 1.0 M mixture in ether, 3 mmol), and stirred for 10 minutes. More ether is added to precipitate the rest of the product. The precipitate is collected by filtration and dried in the vacuum oven at 50 degrees C to give the title compound, mp = 205-209 degrees C; IR (ATR): 2964 and 1649 cm^{-1} ; APCI-MS (m/z) $[M + H]^+ = 638$.

EXAMPLE 748 $\text{N}^3\text{-}\{(1\text{S},2\text{R})\text{-}1\text{-}(3,5\text{-difluorobenzyl})\text{-}2\text{-hydroxy-}3\text{-}[(3\text{-methoxybenzyl})\text{amino}]\text{propyl}\}\text{-N}^5\text{-N}^5\text{-dipropyl}[1,1'\text{-biphenyl}]\text{-}3,5\text{-dicarboxamide hydrochloride (X)}$

Step 1: A mixture of tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl]ethylcarbamate (V, 500 mg, 1.67 mmol) and 3-methoxybenzylamine (VI, 0.34g, 2.51 mmol) in

2-propanol (3 mL) is heated at reflux overnight, allowed to cool to 20-25 degrees C, and concentrated under reduced pressure. The residue is crystallized from ethyl acetate/hexanes and collected by filtration to afford tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, 575 mg) as a solid:
 5 ESI-MS (m/z): 437 $[M + H]^+$.

Step 2: A mixture of tert-butyl (1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, Step 1, 535 mg, 1.23 mmol) in methanol (2 mL) is treated with hydrochloric acid (3.2 mL of a 1.0 M mixture in ether, 3.2 mmol), and stirred at 20-25 degrees C for 30 minutes. Ether is added until a precipitate formed. The precipitate is collected by filtration is
 10 (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol dihydrochloride (VIII).

Step 3: A mixture of 5-[(dipropylamino)carbonyl][1,1'-biphenyl]-3-carboxylic acid (IX, 188 mg, 0.56 mmol), (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol dihydrochloride (VIII, Step 2, 230 mg, 0.56 mmol), HOBT (114 mg, 0.84 mmol), and *N*-methylmorpholine (0.23 g, 2.24 mmol) in methylene chloride (6 mL) and dimethylformamide (1 mL) is stirred at 20-25 degrees C for 15 minutes. EDC (161 mg, 0.84 mmol) is added and the reaction mixture is stirred at 20-25
 20 degrees C overnight. The reaction mixture is washed with water, 1 N hydrochloric acid, water, and saline, dried (sodium sulfate), and concentrated under reduced pressure to give the title compound, mp 230-233 degrees C; IR (ATR): 2965, 1651, 1596 and 1267 cm^{-1} ; ESI-MS (m/z) $[M + H]^+ = 644$.
 25

30

EXAMPLE 749 N^3 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^5,N^5 -dipropyl[1,1'-biphenyl]-3,5-dicarboxamide hydrochloride (X)

Following the general procedure of EXAMPLE 748 and making non-critical variations but using (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII) in place of (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol dihydrochloride (VIII), the title compound is obtained, mp = 214-219 degrees C; IR (KBr): 3227, 2961, 1632 and 1605 cm^{-1} ; ESI-MS (m/z) $[M + H]^+ = 608$.

EXAMPLE 750 N^3 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4'-[(dimethylamino)sulfonyl]- N^5,N^5 -dipropyl-1,1'-biphenyl-3,5-dicarboxamide (X)

Step 1: A flask is charged with 1,1'-bis(diphenylphosphino)ferrocene-dichloropalladium 1:1 complex (37 mg, 0.05 mmol), potassium acetate (492 mg, 4.5 mmol) and bis(pinacolato)diboron (408 mg, 1.6 mmol) and is degassed under reduced pressure for 15 min and purged with argon. To this mixture is added a mixture of methyl-3-bromo-5-[(dipropylamino)carbonyl]benzoate (XXI, 500 mg, 1.5 mmol) in anhydrous dimethyl sulfoxide (9 mL) and the reaction mixture is stirred at 80 degrees C for 4 hours. The reaction mixture is cooled to 20-25 degrees C, diluted with toluene (50 mL), washed with water (3 x 150 mL), saline, dried (magnesium sulfate), and concentrated under reduced pressure to give methyl 3-[(dipropylamino)carbonyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate, ESI-MS (m/z) $[M + H]^+ = 390$.

Step 2: A mixture of methyl 3-[(dipropylamino)carbonyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (Step 1, 534 mg, 1.4 mmol), 4-bromobenzenedimethyl-sulfonamide (363 mg, 1.4 mmol), and sodium carbonate (2 mL of a 2 M mixture in water, 4.1 mmol) in toluene (10 mL) is degassed under reduced pressure for 15 minutes and then purged with argon. Palladium tetrakis(triphenylphosphine) (40 mg, 0.025 mmol) is added and the reaction mixture is degassed under reduced pressure for 15

minutes and then purged with argon. The reaction mixture is heated at reflux for 4 hours, cooled to 20-25 degrees C, filtered through a plug of diatomaceous earth and sodium sulfate, and the filtrate is concentrated under reduced pressure. Purification by flash column chromatography (silica; ethyl acetate/hexanes, 1/1) gives methyl 4'-

[(dimethylamino)sulfonyl]-5-[(dipropylamino)carbonyl][1,1'-biphenyl]-3-carboxylate (XXXVIII), ESI-MS (m/z) $[M + H]^+ = 447$.
 Step 3: A mixture of methyl 4'-[(dimethylamino)sulfonyl]-5-[(dipropylamino)carbonyl][1,1'-biphenyl]-3-carboxylate (XXXVIII, step 2, 555 mg, 1.24 mmol) in methanol (6 mL) and sodium hydroxide (2 mL of a 6.0 M mixture in water, 12 mmol) is stirred at 20-25 degrees C for 4 hours. The reaction mixture is partitioned between ethyl acetate (40 mL) and water (40 mL). The aqueous phase is acidified to pH = 4 with hydrochloric acid (1N), extracted with ether (3 x 100 mL), and the combined organic phases are concentrated under reduced pressure to give methyl 4'-[(dimethylamino)sulfonyl]-5-[(dipropylamino)carbonyl][1,1'-biphenyl]-3-carboxylic acid (IX - XXXIX), NMR (300 MHz, $CDCl_3$): δ 8.37, 8.12, 7.89, 7.80, 3.51, 3.22, 2.76, 1.74, 1.59, 1.02 and 0.79.

Step 4: A mixture of the acid (IX - XXXIX, Step 3, 150 mg, 0.35 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 129 mg, 0.35 mmol) HOBt (47 mg, 0.35 mmol), and *N*-methylmorpholine (122 μ L, 1.1 mmol) is stirred in methylene chloride (4 mL) for 15 minutes EDC (107 mg, 0.62 mmol) is added and the reaction mixture is stirred overnight. The reaction mixture is diluted with water, and extracted with methylene chloride (3 x 25 mL). The organic phase is washed with hydrochloric acid (1N, 25 mL), saturated sodium bicarbonate (25 mL), saline, dried (sodium sulfate), and concentrated under reduced pressure. Purification by flash column chromatography (silica; 100% ethyl acetate to

methanol/chloroform, 1/9) gives the title compound, IR (ATR): 2932, 2837 and 1593 cm^{-1} ; APCI-MS (m/z) $[M + H]^+ = 715$.

EXAMPLE 751 N^3 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4'-[(dimethylamino)sulfonyl]- N^5,N^5 -dipropyl-1,1'-biphenyl-3,5-dicarboxamide (X)

Following the general procedure of EXAMPLE 750 and making non-critical variations but using 2R,3S)-3-amino-1-[(3-iodobenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII), the title compound is obtained, IR (ATR): 3303, 2930, 2872 and 1614 cm^{-1} ; APCI-MS (m/z) $[M + H]^+ = 811$.

EXAMPLE 752 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^3,N^3 -dipropyl-5-(3-thienyl)isophthalamide hydrochloride (X)

Step 1: To an ice-cold mixture of methyl 3-amino-5-[(dipropylamino)carbonyl]benzoate (XLVIII, 1.0 g, 3.60 mmol) in aqueous hydrogen tetrafluoroborate (48% wt. in H_2O , 12.9 mmol) is added a cold mixture of aqueous sodium nitrite (0.25 g, 3.60 mmol) dropwise. The mixture is stirred for 10 min and then extracted with ethyl acetate. The organic phase is washed with water, dried over magnesium sulfate, filtered, and concentrated under reduced pressure to give a diazonium salt which is used without further purification, NMR (500 MHz, CD_3OD): δ 9.26, 8.86, 8.71, 4.03, 3.50, 3.22, 1.75, 1.60, 1.01 and 0.79.

Step 2: To a mixture of thiophene-3-boronic acid (1.0 g, 7.82 mmol) in methanol is added a concentrated aqueous mixture of potassium hydrogen difluoride (2.01 g, 25.8 mmol) dropwise. The reaction mixture is stirred for 10 minutes and concentrated under reduced pressure. The resulting solid is extracted with acetone and concentrated under reduced pressure gives crude material, which is

recrystallized from acetone/ether to give potassium trifluoro(3-thienyl)borate salt, ESI-MS (m/z) $[M + H]^+ = 151$.

Step 3: A mixture of potassium trifluoro(3-thienyl)borate salt (step 2, 0.69 g, 1.82 mmol), diazonium salt from (XLVIII, step 1, 0.42 g, 2.19 mmol), and lead acetate (0.02 g, 0.09 mmol) in the dark is purged with argon for 15 minutes. Dioxane (8 mL) is added and the reaction mixture is degassed with argon and stirred at 20-25 degrees C overnight. The reaction mixture is diluted with ether, washed with saline, dried over magnesium sulfate and concentrated under reduced pressure to give methyl 3-[(dipropylamino)carbonyl]-5-(3-thienyl)benzoate (XLIX) which is purified by flash chromatography (silica; ethyl acetate/hexanes, 1/1), ESI-MS (m/z) $[M + H]^+ = 346$.

Step 4: A mixture of methyl 3-[(dipropylamino)carbonyl]-5-(3-thienyl)benzoate (XLIX, step 3, 0.31 g, 0.88 mmol) in THF/methanol/sodium hydroxide (3/1/1, 5 mL) is stirred at 40 degrees C for 2 hours. The reaction is cooled to 20-25 degrees C, diluted with water and extracted with ethyl acetate. The aqueous phase is acidified to pH = 4 and extracted with ethyl acetate. The organic phase is washed with water and saline, dried over magnesium sulfate and concentrated under reduced pressure to give 3-[(dipropylamino)carbonyl]-5-(3-thienyl)benzoic acid (IX - L), ESI-MS (m/z) $[M + H]^+ = 332$.

Step 5: A mixture of 3-[(dipropylamino)carbonyl]-5-(3-thienyl)benzoic acid (IX - L, step 4, 0.26 g, 0.79 mmol), (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 0.26 g, 0.71 mmol), HOBt (0.16 g, 1.18 mmol), and triethylamine (0.44 mL, 3.15 mmol) in DMF (4 mL) is stirred at 20-25 degrees C for 10 minutes. EDC (0.23 g, 1.18 mmol) is added and the reaction mixture is stirred for 4 hours. The reaction mixture is diluted with water and extracted with ethyl acetate. The organic phase is washed with hydrochloric acid (1 N), water, and saline, dried over magnesium sulfate and

concentrated under reduced pressure. Recrystallization (methylene chloride/hexanes, 1/1) gives the title compound, mp = 199-201 degrees C; IR (KBr): 3278, 2961, 2874 and 2837 cm^{-1} ; ESI-MS (m/z) $[M + H]^+ = 614$.

5 EXAMPLE 753 N-{(1R,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
[(3-methoxybenzyl)amino]propyl}-3-methyl-5-pentanoylbenzamide
(X)

Step 1: To an ice-cold, stirred mixture of oxalyl chloride
(733 mg, 5.77 mmol) in methylene chloride (5 mL) is added 3
10 drops of dimethylformamide. After 10 minutes 3-
(methoxycarbonyl)-5-methylbenzoic acid (LXXIII, 560 mg, 2.89
mmol) is added. The reaction mixture is stirred for 1 hour and
concentrated under reduced pressure to provide an acid chloride
(LXXIV), which is used without further purification.

15 Step 2: To a -78 degrees C, stirred mixture of acid halide
(LXXIV, step 1, 612 mg, 2.89 mmol) and copper (I) bromide (415
mg, 2.89 mmol) in tetrahydrofuran (5 mL) is added butyl
magnesium chloride (1.44 mL of a 2.0 M mixture in
tetrahydrofuran, 2.89 mmol). The reaction mixture is warmed to
20 20-25 degrees C, quenched by addition of saturated ammonium
chloride, and diluted with ether. The organic phase is
separated, washed with saline, dried (sodium sulfate),
filtered, and concentrated under reduced pressure.
Purification by flash column chromatography (silica;
25 hexanes/ethyl acetate, 6.5/1) gives methyl 3-methyl-5-
pentanoylbenzoate (LXXVI), NMR (300 MHz, CD_3OD): δ 8.43, 8.05,
3.96, 3.01, 1.77, 1.55 and 1.22.

Step 3: A mixture of methyl 3-methyl-5-pentanoylbenzoate
(LXXVI, step 2, 133 mg, 0.605 mmol) in methanol (1 mL) is
30 stirred with tetrahydrofuran/methanol/sodium hydroxide (2 N)
(3/1/1, 3 mL) for 3 days. The reaction mixture is diluted with
ethyl acetate and washed with water. The aqueous phase is
separated and acidified with hydrochloric acid (1 N) and

extracted with methylene chloride. The organic phase is dried (sodium sulfate), filtered, and concentrated under reduced pressure to give 3-methyl-5-pentanoylbenzoic acid (IX - LXXVII), NMR (300 MHz, CD₃OD): δ 8.44, 8.03, 3.10, 2.33, 1.78, 1.64 and 1.34.

Step 4: To a mixture of 3-methyl-5-pentanoylbenzoic acid (IX - LXXVII, 112 mg, 0.589 mmol), (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol dihydrochloride (VIII, 239 mg, 0.589 mmol), HOBt (80 mg, 0.589 mmol), and N-methylmorpholine (250 mg, 2.47 mmol) in methylene chloride (3 mL) is added EDC (203 mg, 1.06 mmol). The reaction mixture is stirred overnight and then partitioned between ethyl acetate and water. The organic phase is washed with hydrochloric acid (1 N), saturated sodium bicarbonate, saline, dried (sodium sulfate), filtered, and concentrated under reduced pressure. Purification by flash column chromatography (silica; methylene chloride/methanol, 12/1) gives the title compound, IR (ATR): 3297, 2957, 1687 and 1628 cm⁻¹; APCI-MS (m/z) [M + H]⁺ = 539.

EXAMPLE 754 N¹-(4-hydroxybutyl)-N³-{(1S)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N¹-propylisophthalamide (X)

Step 1: To a mixture of methyl (2S)-3-[4-(benzyloxy)phenyl]-2-(tert-butoxycarbonyl)aminopropanoate (1.79 g, 4.65 mmol) in a THF/methanol/water (1/2/1, 16 mL) is added lithium hydroxide (340 mg, 13.9 mmol) and the mixture stirred at 20-25 degrees C for 12 hours. The mixture is quenched with citric acid (10%). The resulting mixture is extracted with ethyl acetate (3 x 15 mL). The combined organic extracts are washed three times with water, dried over sodium sulfate, filtered, and concentrated under reduced pressure to give (2S)-3-[4-(benzyloxy)phenyl]-2-[(tert-butoxycarbonyl)amino]propanoic

acid which is carried on without purification. To a -78 degrees C, stirred mixture of (2S)-3-[4-(benzyloxy)phenyl]-2-[(tert-butoxycarbonyl)amino]propanoic acid (10.0 g, 27.0 mmol) in THF (200 mL) is added NMM (3.20 mL, 29.0 mmol) and isobutyl chloroformate (3.8 mL, 29.0 mmol). The cold bath is removed, the reaction mixture is stirred for 1 hour, and then filtered. The filtrate is kept cold and used in the next step. To an ice-cold, stirred mixture of ether (110 mL) and potassium hydroxide (40%, 35 mL) is slowly added 1-methyl-3-nitro-1-nitrosoguanidine (8.40 g, 57.0 mmol). The reaction mixture is stirred until gas evolution ends. The organic phase is separated and slowly added to an ice-cold, stirred mixture of the mixed anhydride filtrate from step 2. After the reaction mixture is stirred for 1 hour, nitrogen is bubbled into the mixture for 10 minutes. The resulting mixture is concentrated under reduced pressure, diluted with ethyl acetate (200 mL), and washed with water (100 mL). The organic phase is washed with saturated sodium bicarbonate and saline, dried over sodium sulfate, filtered, and concentrated under reduced pressure to give the diazoketone, which is carried on without purification or characterization. To an ice-cold, stirred mixture of diazoketone in ether (100 mL) is added hydrobromous acid (48%, 4 mL, 73 mmol). The cold bath is removed, the reaction mixture stirred for 30 minutes, and partitioned between ether and water. The organic phase separated and washed with saturated sodium bicarbonate and saline, dried over sodium sulfate, filtered, and concentrated under reduced pressure to give tert-butyl (1S)-1-[4-(benzyloxy)benzyl]-3-bromo-2-oxopropylcarbamate (IV) which is used without further purification or characterization. To a -78 degrees C, stirred mixture of tert-butyl (1S)-1-[4-(benzyloxy)benzyl]-3-bromo-2-oxopropylcarbamate (IV) in a isopropanol/THF (2/1, 150 mL) is slowly added sodium borohydride (1.15 g, 30.0 mmol). The reaction mixture is

stirred for 30 minutes followed by the addition of water (30 mL). The resulting mixture is warmed to 20-25 degrees C and concentrated under reduced pressure in a water bath not exceeding 30 degrees C. The crude residue is dissolved in ethyl acetate and washed with water and saline. The organic phase is dried over magnesium sulfate, filtered and concentrated under reduced pressure to give the bromohydrin as a solid. To an ice-cold, stirred mixture of bromohydrin in ethanol (150 mL) and ethyl acetate (100 mL) is added a potassium hydroxyde (1 N) ethanol mixture (36 mL, 36 mmol). The cold bath is removed and the reaction mixture is stirred for 30 minutes. The resulting mixture is partitioned between ethyl acetate and water. The organic phase is separated and washed with saline, dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by flash chromatography (silica; hexanes/ethyl acetate, 5/1) gives tert-butyl (1S)-2-[4-(benzyloxy)phenyl]-1-[(2S)-oxiranyl]ethylcarbamate (V, as a 8/1 mixture of diastereomers), NMR (500 MHz, CDCl₃) δ 7.44-7.32, 7.14, 6.93, 5.07, 4.45, 3.61, 3.00-2.60 and 1.39.

Step 2: A mixture of 4-benzyloxybutyric acid (2.69 g, 13.8 mmol), propylamine (0.82 g, 13.8 mmol), HOBt (2.05 g, 15.2 mmol), N-methylmorpholine (1.68 g, 16.6 mmol) and EDC (2.91 g, 15.2 mmol) in DMF (6 mL) is stirred at 20-25 degrees C for 18 hours. The mixture is diluted with ethyl acetate (40 mL) and washed with water (10 mL), hydrochloric acid (1 N, 10 mL), saturated sodium bicarbonate (10 mL), and saline (10 mL). The organic phase is separated, dried over magnesium sulfate, filtered, and concentrated under reduced pressure to provide 4-(benzyloxy)-N-propylbutanamide (2.59 g), APCI-MS (m/z) [M + H]⁺ = 236.

Step 3: To an ice-cold, stirred mixture of 4-(benzyloxy)-N-propylbutanamide (2.59 g, 11.0 mmol) in THF (8 mL) is added

lithium aluminum hydride (0.54 g, 14.3 mmol). The reaction mixture is heated to 40-50 degrees C for 5 hours. The cooled reaction mixture is quenched with water (0.5 mL), sodium hydroxide (2 N, 1.0 mL), and saline (0.5 mL) then diluted with ether (30 mL). The precipitate that formed is filtered off, and the ether phase dried over magnesium sulfate, filtered, and concentrated under reduced pressure to give N-[4-(benzyloxy)butyl]-N-propylamine (2.41 g), APCI-MS (m/z): 222 [$M + H$]⁺.

Step 4: A mixture of N-[4-(benzyloxy)butyl]-N-propylamine (2.31 g, 10.44 mmol), 3-(ethoxycarbonyl)-5-methylbenzoic acid (2.18 g, 10.44 mmol), HOBt (1.56 g, 11.49 mmol), N-methylmorpholine (1.37 mL, 12.52 mmol), and EDC (2.20 g, 11.49 mmol) in DMF (12 mL) is stirred at 20-25 degrees C for 18 hours. The reaction mixture is diluted with ethyl acetate (80 mL) and washed with water (2 x 20 mL), hydrochloric acid (1 N, 20 mL), saturated sodium bicarbonate (20 mL) and saline (20 mL), dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by flash chromatography (silica; hexanes/ethyl acetate, 1/1) gives ethyl 3-[[[4-(benzyloxy)butyl](propyl)amino]carbonyl]-5-methylbenzoate (1.79 g), NMR (500 MHz, DMSO- d_6): δ 7.80, 7.64, 7.40, 7.38-7.16, 4.50-4.43, 4.34-4.29, 3.53-3.30, 3.20-3.06, 2.41-2.36, 1.70-1.40, 1.36-1.29, 0.94-0.84 and 0.82-0.72; APCI-MS (m/z) [$M + H$]⁺ = 412.

Step 5: To a mixture of ethyl 3-[[[4-(benzyloxy)butyl](propyl)-amino]carbonyl]-5-methylbenzoate (1.75 g, 4.25 mmol) in THF/ethanol/water (1/2/1, 30 mL) is added lithium hydroxide (0.31 g, 12.76 mmol). The reaction mixture is stirred for 2 h and then acidified to pH = 3 with concentrated hydrochloric acid (0.5 mL). The reaction mixture is extracted with ethyl acetate (2 x 30 mL), dried over magnesium sulfate, filtered, and concentrated under reduced

pressure to give 3-[[4-(benzyloxy)butyl](propyl)amino]carbonyl}-5-methylbenzoic acid (IX, 1.63 g), ESI-MS (m/z) $[M + H]^+ = 384$.

Step 6: A mixture of tert-butyl (1S)-2-[4-(benzyloxy)phenyl]-1-[(2S)-oxiranyl]ethylcarbamate (V, 1.58 g, 4.28 mmol) and 3-methoxybenzylamine (VI, 825 microliter, 6.42 mmol) in isopropanol (45 mL) is heated to 90 degrees C for 4 hours. Upon cooling to 20-25 degrees C, the reaction mixture is concentrated under reduced pressure. Purification by flash chromatography (silica; methylene chloride/methanol/ammonium hydroxide 98/1/1 to 95/4/1) gives tert-butyl (1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, 1.97 g), NMR (300 MHz, MeOH- d_4): δ 7.41-6.79, 5.05, 4.33-3.33, 3.74, 3.54, 3.03-2.46 and 1.29; ESI-MS (m/z) $[M + H]^+ = 507$.

Step 7: tert-Butyl (1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, step 6, 2.34 g, 4.62 mmol) in dioxane (10 mL) is treated with hydrochloric acid (12 mL of a 4.0 M mixture in dioxane, 48 mmol) for 2 hours. The precipitate that forms is collected by filtration, washed with ether, and dried under reduced pressure overnight to give (2R,3S)-3-amino-4-[4-(benzyloxy)phenyl]-1-[(3-methoxybenzyl)amino]-2-butanol hydrochloride (VIII), NMR (300 MHz, MeOH- d_4): δ 7.44-6.96, 5.05, 4.21, 3.83, 3.65 and 3.21-2.77; ESI-MS (m/z) $[M + H]^+ = 407$.

Step 8: To an ice-cold, stirred mixture of 3-[[4-(benzyloxy)butyl](propyl)amino]carbonyl}-5-methylbenzoic acid (IX, 310 mg, 0.809 mmol), (2R,3S)-3-amino-4-[4-(benzyloxy)phenyl]-1-[(3-methoxybenzyl)amino]-2-butanol hydrochloride (VIII, 359 mg, 0.809 mmol), and bromotripyrrolidinophosphonium hexafluorophosphate (415 mg, 0.890 mmol) in methylene chloride (10 mL) is added diisopropylethylamine (285 microL, 1.62 mmol) dropwise. The

resulting mixture is stirred at 0 degrees C for 30 minutes and then warmed to 20-25 degrees C. After 4 hours, the reaction is concentrated under reduced pressure and is partitioned between ethyl acetate and water. The aqueous phase is separated and
 5 extracted with ethyl acetate (3 x 15 mL), the combined organic phases are dried over magnesium sulfate, and concentrated under reduced pressure. The concentrate is purified by flash chromatography (silica; methylene chloride/methanol/ammonium hydroxide 96/3/0.5) to give N¹-{(1S,2R)-1-[4-(benzyloxy)benzyl]-
 10 2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³-[4-(benzyloxy)butyl]-5-methyl-N³-propylisophthalamide (X) NMR (300 MHz, Acetone-d₆): δ 7.99-6.74, 5.01 4.51-4.29, 4.36, 4.01, 3.80, 3.55-3.16, 2.98-2.82, 2.65-2.62, 2.36, 1.85-1.29, 1.01 and 0.68; ESI-MS (m/z) [M + H]⁺ = 772.

15 Step 9. A mixture of N¹-{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³-[4-(benzyloxy)butyl]-5-methyl-N³-propylisophthalamide (X, 100 mg, 0.130 mmol) and palladium on carbon (10%, 100 mg) in absolute glacial acetic acid (5 mL) is shaken under an atmosphere of
 20 hydrogen at 35 psi for 5 hours. The resulting mixture is filtered through diatomaceous earth and washed with methanol. The combined filtrates are concentrated under reduced pressure. The concentrate is purified by flash column chromatography (silica; gradient of dichloromethane/methanol/ammonium hydroxide
 25 97/3/0.05 to 93/7/0.05) to give the title compound: NMR (300 MHz, CD₃OD): δ 7.55-6.64, 4.19, 3.99-3.72, 3.63-3.36, 3.21-3.09, 2.79-2.69, 2.39, 1.90-1.40, 1.29 and 1.02-0.6; ESI-MS (m/z) [M + H]⁺ = 592.

30 EXAMPLE 755 N¹-{(1S,2R)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N³-(3-hydroxypropyl)-5-methyl-N³-propylisophthalamide (X)

Following the general procedure of EXAMPLE 754 and making non-critical variations, but using 3-[[3-

(benzyloxy)propyl] (propyl)amino]carbonyl}-5-methylbenzoic acid (IX) in place of 3-[[[4-(benzyloxy)butyl] (propyl)amino]-carbonyl]-5-methylbenzoic acid (IX), the title compound is obtained, ESI-MS (m/z) $[M + H]^+ = 578$.

5

EXAMPLE 756 N^1 -{(1S,2R)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

Step 1. To a stirred mixture of 3-

10 [(dipropylamino)carbonyl]-5-methylbenzoic acid (IX, 150 mg, 0.570 mmol), (2R,3S)-3-amino-4-[4-(benzyloxy)phenyl]-1-[(3-methoxybenzyl)amino]-2-butanol hydrochloride (VIII, 274 mg, 0.571 mmol), *N,N*-diisopropylethylamine (400 microliter, 2.28 mmol), and HOBt (116 mg, 0.857 mmol) in dichloromethane (10 mL)
15 is added EDC (165 mg, 0.857 mmol). The resulting mixture is stirred at 20-25 degrees C for 16 hours. The reaction mixture is partitioned between dichloromethane and water. The aqueous phase is separated and extracted with dichloromethane (3 x 15 mL). The combined organic phases are washed with water, dried
20 (magnesium sulfate), and concentrated under reduced pressure. Purification by flash column chromatography (silica; dichloromethane/methanol/ammonium hydroxide, 97/3/0.05) gives N^1 -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -
25 dipropylisophthalamide, ESI-MS (m/z) $[M + H]^+ = 652$.

Step 2. A mixture of N^1 -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (140 mg, 0.215 mmol) and palladium on carbon (10%, 140 mg) in absolute glacial acetic acid (5 mL) is
30 shaken under an atmosphere of hydrogen at 35 psi for 5 hours. The resulting mixture is filtered through diatomaceous earth and washed with methanol. The combined filtrates are concentrated under reduced pressure. The concentrate is

purified by flash column chromatography (silica; methylene chloride/methanol/ammonium hydroxide gradient from 97/3/0.05 to 93/7/0.05) to give the title compound, IR (KBr) 2962, 2931, 1611, 1594 and 1263 cm^{-1} ; ESI-MS (m/z) $[M + H]^+ = 562$.

5

EXAMPLE 757 $N^1-((1S,2R)-1\text{-benzyl-3-}\{[3-(2,4\text{-dimethylphenyl})\text{propyl}]\text{amino}\}-2\text{-hydroxypropyl})-5\text{-methyl-}N^3,N^3\text{-dipropylisophthalamide (X)}$

Step 1: A stirred mixture of tert-butyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V, 247 mg, 0.939 mmol), sodium carbonate (299 mg, 2.82 mmol), and 3-(2,4-dimethylphenyl)propylamine (VI, 628 mg, 2.82 mmol) is heated at reflux overnight. The reaction mixture is cooled to 20-25 degrees C and concentrated under reduced pressure.

15 Purification by flash column chromatography (silica; methylene chloride/methanol/ammonium hydroxide, 98/2/1) gives tert-butyl (1S,2R)-1-benzyl-3- $\{[3-(2,4\text{-dimethylphenyl})\text{propyl}]\text{amino}\}-2\text{-hydroxypropylcarbamate (VII)}$, NMR (300 MHz, CD_3OD): delta 7.22-7.16, 3.81, 3.18, 2.77, 2.54, 2.15, 2.13, 1.89 and 1.23.

20 Step 2: To a stirred mixture of tert-butyl (1S,2R)-1-benzyl-3- $\{[3-(2,4\text{-dimethylphenyl})\text{propyl}]\text{amino}\}-2\text{-hydroxypropylcarbamate (VII)}$, 180 mg, 0.423 mmol) in dioxane (2 mL) is added hydrochloric acid (0.32 mL of a 4 N mixture in dioxane, 1.27 mmol). The reaction mixture is stirred overnight and concentrated under reduced pressure to give (2R,3S)-3-amino-1- $\{[3-(2,4\text{-dimethylphenyl})\text{propyl}]\text{amino}\}-4\text{-phenyl-2-butanol hydrochloride (VIII)}$, NMR (300 MHz, CDCl_3): delta 7.14, 3.73, 2.70, 2.32 and 1.86.

Step 3: To a stirred mixture of (2R,3S)-3-amino-1- $\{[3-(2,4\text{-dimethylphenyl})\text{propyl}]\text{amino}\}-4\text{-phenyl-2-butanol hydrochloride (VIII)}$, 163 mg, 0.411 mmol), 3-[(dipropylamino)carbonyl]-5-methylbenzoic acid (IX, 108 mg, 0.411 mmol), HOBt (55 mg, 0.411 mmol), and N-methylmorpholine

(133 mg, 1.32 mmol) in methylene chloride (5 mL) is added EDC (142 mg, 0.740 mmol). The reaction mixture is stirred overnight and then partitioned between ethyl acetate and water. The organic phase is washed with hydrochloric acid (1 N),
 5 saturated sodium bicarbonate, saline, dried (sodium sulfate), filtered, and concentrated under reduced pressure. Purification by flash column chromatography (silica; methylene chloride/methanol/ammonium hydroxide, 95/5/1) gives the title compound, IR (ATR): 3299, 2930 and 1614 cm^{-1} ; APCI-MS (m/z) [$M + H$]⁺ = 572.
 10

EXAMPLE 758 N^1 -((1S,2R)-1-benzyl-2-hydroxy-3-{[3-(4-methylphenyl)propyl]amino}propyl)-5-methyl- N^3,N^3 -dipropylisophthalamide hydrochloride (X)

15 Following the general procedure of EXAMPLE 757 and making non-critical variations, but using 3-(4-methylphenyl)propylamine in place of 3-(2,4-dimethylphenyl)propylamine, the title compound is obtained, IR (ATR): 3282, 2929 and 1594 cm^{-1} ; APCI-MS (m/z) [$M + H$]⁺ = 558.
 20

EXAMPLE 759 N^1 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide (X)

25 Following the general procedure of EXAMPLES 4, 5, and 6 and making non-critical variations but using tert-butyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and "5-Me-PHTH" (IX), the title compound is obtained, MS [$M+H$]⁺ = 546.3.

30 EXAMPLE 760 N -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-1,3-dioxo-2-propyl-5-isindolinecarboxamide (X)

Following the general procedure of EXAMPLEs 4, 5, and 6 and making non-critical variations but using tert-butyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and 1,3-dioxo-2-propylisoindoline-5-carboxylic acid (IX), the title compound is obtained, MS $[M+H]^+$ = 516.2.

EXAMPLE 761 N-{(1R,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-bromo-5-methylbenzamide (X)

10 Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using benzyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-bromo-5-methylbenzoic acid (IX), the title compound is obtained, MS $[M+H]^+$ = 497.1, 499.1.

15

EXAMPLE 762 3-bromo-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methylbenzamide (X)

20 Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl] ethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-bromo-5-methylbenzoic acid (IX), the title compound is obtained, MS $[M+H]^+$ = 533.3, 535.3.

25

EXAMPLE 763 N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-methyl-N³,N³-dipropylisophthalamide (X)

30 Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using benzyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine

(VI) and 3-[(dipropylamino)carbonyl]-4-methylbenzoic acid (IX), the title compound is obtained, MS $[M+H]^+ = 546.4$.

EXAMPLE 764 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-
5 [(3-methoxybenzyl)amino]propyl}-4-methyl- N^3,N^3 -
dipropylisophthalamide (X)

Following the general procedure of EXAMPLES 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl] ethylcarbamate (V) and
10 3-methoxybenzylamine (VI) and 3-[(dipropylamino)carbonyl]-4-methylbenzoic acid (IX), the title compound is obtained, MS $[M+H]^+ = 582.3$.

EXAMPLE 765 N^3 -{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-
15 methoxybenzyl)amino]propyl}-4-methyl- N^1,N^1 -
dipropylisophthalamide (X)

3-Bromo-4-methylbenzoic acid (10.94 g, 43.25 mmol), copper(I) cyanide (7.75 g, 86.5 mmol) and 1-methyl-2-pyrrolidinone (75 ml) are heated to 160 degrees C overnight.
20 The mixture is cooled and vacuum distilled to give a residue which is stirred in hydrochloric acid (6N, 60 ml) for 10 minutes. The resulting solid is collected by filtration, washed with water, ether, and dried. The solid is heated to 90 degrees C in sodium hydroxide (2N, 250 ml) for 3 hours and the
25 mixture is then cooled and stirred overnight at 20-25 degrees C. The reaction is acidified to about pH 3 with concentrated hydrochloric acid which gives a precipitate. The solids are collected by filtration and washed with water, then triturated in boiling water, filtered and dried in a vacuum oven at 60
30 degrees C. The solid is dissolved in methanol (75 ml) and concentrated hydrochloric acid (5 ml) is added and the mixture is refluxed overnight. The mixture then is cooled and

concentrated under reduced pressure. Chromatography (silica gel; methanol/methylene chloride, 8/92) gives 5-(methoxycarbonyl)-2-methylbenzoic acid.

To 5-(methoxycarbonyl)-2-methylbenzoic acid (250 mg, 1.3 mmol) and triethylamine (0.72 ml, 5.2 mmol) in methylene chloride (14 ml) is added diethylcyanopyrocarbonate (90%, 0.24 ml, 1.4 mmol) with stirring. After 1 minute, (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol dihydrochloride (VIII, 485 mg, 1.3 mmol) is added and the reaction is stirred overnight. The mixture is concentrated followed by chromatography (silica gel; methanol/methylene chloride 8/92) to afford 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino)carbonyl]-4-methylbenzoate.

3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino) carbonyl]-4-methylbenzoate (200 mg, 0.42 mmol) is treated with lithium hydroxide (39 mg, 0.96 mmol) in tetrahydrofuran/methanol/water (2/1/1, 2 ml), and the mixture stirred overnight at 20-25 degrees C. The mixture is decanted and the supernatant concentrated to give 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino)carbonyl]-4-methylbenzoic acid.

3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino) carbonyl]-4-methylbenzoic acid (124 mg, 0.27 mmol) is dissolved in triethylamine (0.07 ml, 0.54 mmol) and methylene chloride (3 ml) and treated with diethylcyanopyrocarbonate (90%, 0.06 ml, 0.32 mmol) with stirring for 2 minutes. Dipropylamine (0.04 ml, 0.32 mmol) is added and stirring continued overnight. The organic phase is diluted with methylene chloride and washed with saturated sodium bicarbonate (2 X 50 ml) and saline (50 ml) then dried over anhydrous sodium sulfate, filtered and concentrated. Chromatography (silica gel; methanol/methylene chloride, 8/92) gives the title compound, MS $[M+H]^+ = 546.3$.

EXAMPLE 766 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-(2-furyl)-5-methylbenzamide (X)

5 N-{(1R,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-bromo-5-methylbenzamide (X, EXAMPLE 761, 295 mg, 0.59 mmol), 2-furanylboronic acid (133 mg, 1.19 mmol) and sodium carbonate (366 mg, 2.95 mmol) are combined in dimethylformamide (5 ml) and sparged under a flow
10 of nitrogen for 15 minutes. Tetrakis(triphenylphosphino)palladium (136 mg, 0.12 mmol) is added and the mixture heated to 100 degrees C overnight. The mixture is cooled to 20-25 degrees C, diluted with chloroform (50 ml) and extracted with water (3 x 100 ml). The organic
15 phase is separated and washed with saturated sodium bicarbonate (2 x 100 ml) and saline (100 ml), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The residue is chromatographed (silica gel; methanol/methylene chloride, 8/92) to give the title compound, MS [M+H]⁺ = 485.3.

20

EXAMPLE 767 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3',5,5'-trimethyl-1,1'-biphenyl-3-carboxamide (X)

Following the general procedure of EXAMPLE 766 and making
25 non-critical variations but using 3,5-dimethylphenylboronic acid, the title compound is obtained, MS [M+H]⁺ = 523.3.

EXAMPLE 768 3'-Acetyl-N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl[1,1'-biphenyl]-3-carboxamide (X)

30

Following the general procedure of EXAMPLE 766 and making non-critical variations but using 3-acetylphenylboronic acid, the title compound is obtained, MS $[M+H]^+ = 537.3$.

- 5 EXAMPLE 769 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3'-methoxy-5-methyl[1,1'-biphenyl]-3-carboxamide (X)

Following the general procedure of EXAMPLE 766 and making non-critical variations but using 3-methoxyphenylboronic acid,
10 the title compound is obtained,
MS $[M+H]^+ = 525.3$.

- EXAMPLE 770 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl[1,1'-
15 biphenyl]-3-carboxamide (X)

Following the general procedure of EXAMPLE 766 and making non-critical variations but using phenylboronic acid, the title compound is obtained, MS $[M+H]^+ = 495.3$.

- 20 EXAMPLE 771 3-Methyl-5-(3-thienyl)benzoic acid (IX)

Following the general procedure of EXAMPLE 766 and making non-critical variations but using 2-thiopheneboronic acid and methyl 3-bromo-5-methylbenzoate, methyl 3-methyl-5-(3-thienyl)benzoate is obtained.

- 25 Methyl 3-methyl-5-(3-thienyl)benzoate (257 mg, 1.1 mmol) is treated with lithium hydroxide (186 mg, 4.4 mmol) in tetrahydrofuran/methanol/water (8 mL, 2:1:1) and the mixture is stirred for 2 hours at 20-25 degrees C. The mixture is acidified and concentrated to give the title compound, MS
30 $[M+H]^+ = 217.0$.

EXAMPLE 772 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-5-(3-thienyl)benzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using benzyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-methyl-5-(3-thienyl)benzoic acid (IX), the title compound is obtained, MS [M+H]⁺ = 501.2.

10 EXAMPLE 773 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-5-(2-thienyl)benzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using benzyl (1S)-1-[(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-methyl-5-(2-thienyl)benzoic acid, the title compound is obtained, MS [M+H]⁺ = 501.2.

EXAMPLE 774 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-5-(3-thienyl)benzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiranyl] ethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-methyl-5-(3-thienyl)benzoic acid (IX), the title compound is obtained, MS [M+H]⁺ = 537.2.

EXAMPLE 775 N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-3-methyl-5-(3-thienyl)benzamide (X)

Following the general procedure of EXAMPLEs 4, 5 and 6 and making non-critical variations but using tert-butyl (1S)-2-

(3,5-difluorophenyl)-1-[(2S)-oxiranyl] ethylcarbamate (V) and 3-iodobenzylamine (VI) and 3-methyl-5-(3-thienyl)benzoic acid (IX), the title compound is obtained, MS $[M+H]^+ = 633.1$.

5 EXAMPLE 776 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-methyl-3-(3-thienyl)benzamide (X)

Following the general procedure of EXAMPLES 4, 5, 6 and 766 and making non-critical variations but using benzyl (1S)-1-
10 [(2S)-oxiranyl]-2-phenylethylcarbamate (V) and 3-methoxybenzylamine (VI) and 3-bromo-4-methylbenzoic acid (IX) and 3-thiopheneboronic acid, the title compound is obtained, MS $[M+H]^+ = 501.2$.

15 EXAMPLE 777 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^3,N^3,N^5,N^5 -tetrapropylbenzene-1,3,5-tricarboxamide hydrochloride (X) Following the general procedures of CHART A as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical
20 variations and using the appropriate reagents, the title compound is obtained, HRMS (FAB) = 659.4155.

EXAMPLE 778 N^1 -{(1S,2R)-1-(3,5-Difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}- N^3,N^3 -dipropylbenzene-1,3,5-
25 tricarboxamide (X)

Following the general procedures of CHART A as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS $(M+H)^+$
30 = 609.2.

EXAMPLE 779 Ethyl 3-[(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]amino) carbonyl]-5-[(dipropylamino) carbonyl] benzoate hydrochloride (X)

Following the general procedures of CHART A as well as
5 PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, HRMS (FAB) = 604.3392.

10 EXAMPLE 780 N¹-{(1S,2R)-2-Hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropylbenzene-1,3,5-tricarboxamide (X)

Following the general procedures of CHART A as well as
15 PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 591.0.

20 EXAMPLE 781 N¹-{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropyl-5-[(trifluoromethyl)sulfonyl]amino}isophthalamide (X)

Following the general procedures of CHART A and CHART T as
25 well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 679.2.

30 EXAMPLE 782 5-Amino-N¹-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropylisophthalamide (X)

WO 03/072535

Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺
5 = 547.3.

EXAMPLE 783 N¹-{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropyl-5-[(trifluoroacetyl)amino]isophthalamide (X)

10 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺
= 643.2

15

EXAMPLE 784 N¹-{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(methanesulfonyl)amino]-N³,N³-dipropylisophthalamide hydrochloride (X)

20 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺
= 625.0

25 EXAMPLE 785 N¹-{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N³,N³-dipropyl-5-[(thien-2-ylsulfonyl)amino]isophthalamide hydrochloride (X)

30 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺
= 693.1

WO 03/072535

EXAMPLE 786 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}- N^3,N^3 -dipropyl-5-[(thien-2-ylcarbonyl)amino]isophthalamide (X)

5 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 657.2.

10

EXAMPLE 787 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(methacryloylamino)- N^3,N^3 -dipropylisophthalamide (X)

15 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 615.1.

20

EXAMPLE 788 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(2,2-dimethylpropanoyl)amino]- N^3,N^3 -dipropylisophthalamide (X)

25 Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 631.3.

30

EXAMPLE 789 N^1 -{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-

WO 03/072535

[(phenylsulfonyl)amino]-N³,N³-
dipropylisophthalamide (X)

Following the general procedures of CHART A and CHART T as well as PREPARATIONS 1-13, EXAMPLES 1-13, 321-48-579, 597-622 and 624-631 and making non-critical variations and using the appropriate reagents, the title compound is obtained, MS (M+H)⁺ = 687.2.

EXAMPLE 790 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(methylthio)pentanamide

Following the general procedure of EXAMPLES 588-592 and 597-619 and using the appropriate starting materials the title compound is obtained, HRMS m/z calculated for C₂₄H₃₅N₂O₃S (M+H) = 431.2363, found = 431.2397.

EXAMPLE 791 tert-butyl (2R,3S)-3-({3-[(dipropylamino)sulfonyl]-propanoyl}amino)-2-hydroxy-4-phenylbutyl(3-methoxybenzyl)carbamate

Following the general procedure of EXAMPLES 588-592 and 597-619 and using the appropriate starting materials the title compound is obtained, HRMS calculated for M+H is 620.3369, observed = 620.3379.

EXAMPLE 7922-Butylcyclopropylamine hydrochloride (VI)

A solution of triethylphosphonoacetate (22.4 g, 0.1 mol) in 13 mL of diglyme is added to a mixture of 13 mL of diglyme and sodium hydride (60%, 5.7 g, 0.12 mol) in mineral oil. When hydrogen evolution ceased, 1,2-epoxyhexane (12 g, 0.12 mol) in diglyme (12 mL) is added. The mixture is stirred for 1 day at 25 degrees C and 3 hours at 140 degrees C. A mixture of sodium hydroxide (15 g in 25 mL of water) is added

in the cold. The mixture is refluxed 15 hours, diluted with cold water (100 mL), and washed with ether (3 x 50 mL). Acidification to pH = 2 with sulfuric acid (25%), extraction with ether (5 x 25 mL), drying the ether over anhydrous sodium sulfate, filtration and concentration gives 2-butylcyclopropanecarboxylic acid. The acid (5.0 g, 0.035 mmol) in dichloromethane (15 mL) is heated with thionyl chloride (5.1 g, 3.1 mL) for 15 hours at 60 degrees C. The reaction mixture is distilled (76 degrees C- 80 degrees C) to give the acid chloride which is dissolved in acetone (15 mL), cooled to -10 degrees C and treated with sodium azide (2.2 g, 33.8 mmol) in water (5 mL). The reaction mixture is stirred at -10 degrees C for another 1 hour and then poured onto ice/water, extracted with ether (3x10 mL), dried, and cautiously evaporated to dryness at 20-25 degrees C under reduced pressure. The residue is dissolved in toluene (15 mL) and carefully warmed to 100 degrees C while vigorously stirring for 1 hour. Concentrated hydrochloric acid (7 mL) is added and the reaction mixture is refluxed for 15 minutes. The acidic layer is evaporated to dryness to give the title compound, $MH^+ = 114.2$.

EXAMPLE 793 2-Aminomethyl-3-methylfuran (VI)

3-Methylfuroic acid (4.0 g, 32 mmol) is dissolved in DMF (10 mL) at 20-25 degrees C, and 1,1-carbonyldiimidazole (5.7 g, 35 mmol) is added. After 15 minutes, ammonia is bubbled into the mixture for approximately 2 minutes. This mixture is stirred at 20-25 degrees C for 2 hours then the mixture is concentrated under reduced pressure. The residue is partitioned between ethyl acetate and 10% aqueous citric acid. The layers are separated, and the aqueous layer extracted with additional ethyl acetate (2 x). The combined organic phases are washed with saturated sodium bicarbonate,

then saline and dried over magnesium sulfate, filtered and concentrated. Crystals formed upon standing, which are isolated by filtration and washing with a small amount of ethyl acetate/hexanes (80/20), MS(ESI): MH+: 126.1. 3-

- 5 Methylfuroic amide (317 mg, 2.5 mmol) is dissolved in dry THF (5 mL). Lithium aluminum hydride (230 mg, 6.0 mmol) is added in one portion, and the mixture heated to reflux overnight. The mixture is cooled to 0 degrees C, and quenched by addition of THF/water (50/50). The mixture is then diluted
10 with THF, and filtered through diatomaceous earth. The filtrate is concentrated to give the title compound, MS(ESI): (M-H)+: 109.1.

EXAMPLE 7944-Aminomethyl-3,5-dimethylisoxazole (VI)

- 15 4-Chloromethyl-3,5-dimethylisoxazole (700 mg, 4.8 mmol) is suspended in concentrated aqueous ammonia at 20-25 degrees C, and vigorously stirred overnight. The reaction mixture is extracted with isopropyl alcohol/chloroform (10/90, 2 x). The combined organic phases are concentrated under nitrogen flow.
20 The residue is purified by flash chromatography methanol/methylene chloride (5-20%, 1% triethylamine) to give the title compound, MR (CDCl₃, 300 MHz) delta 3.62, 2.37, 2.29, and 1.44.

25 EXAMPLE 795 5-Hydroxymethyl-2-(2-methylpropyl) thiazole (VI)

- Isovalerothioamide is synthesized according to the procedure in J. Med. Chem. 41, 602-617 (1998). Isovaleramide (10 g, 9.9 mmol) is suspended in dry ether (400 mL), then
30 phosphorous(V) sulfide (4.4 g, 0.99 mmol) is added in portions. This is vigorously stirred at 20-25 degrees C for 2 hours, then filtered. The filtrate is concentrated under

reduced pressure and the residue used without further purification: MS(ESI): MH+: 118.1.

Isovalerothioamide (6.0 g, 51 mmol) and ethyl formylchloroacetate (*Heterocycles* 32 (4), 693-701, (1991), 5.0 g, 33 mmol) are dissolved in dry DMF (20 mL), and heated to 95 degrees C for 4 hours. The reaction is subsequently cooled to 0 degrees C, and cold water (50 mL) is added. The mixture is basified to pH = 8 with solid sodium bicarbonate, then extracted with ether (3 x 35 mL). The combined organic
10 extracts are washed with water, then saline and dried over magnesium sulfate, filtered, and concentrated. The residue is purified by flash chromatography (ethyl acetate/hexanes 4-10% elution) to give the desired product. NMR (CDCl₃, 300 MHz) δ 8.27, 4.45-4.30, 3.70-3.50, 3.00-2.80, 2.30-2.10, 1.40-1.20,
15 and 1.10-0.90.

A solution of ethyl 2-(2-methylpropyl)thiazole-5-carboxylate (2.05 g, 9.6 mmol) in THF (10 mL) is added dropwise with stirring to a suspension of lithium aluminum hydride (730 mg, 19 mmol) in dry THF (50 mL) at 0 degrees C.
20 Upon complete addition, the reaction mixture is allowed to stir at 20-25 degrees C. The reaction mixture is cooled to 0 degrees C, and water (0.75 mL), aqueous sodium hydroxide (15%, 0.75 mL), and water (2.25 mL) is added in succession. This mixture is stirred at 0 degrees C for 1 hour, then
25 filtered through diatomaceous earth, (THF and chloroform). The filtrate is concentrated to give 5-hydroxymethyl-2-(2-methylpropyl)thiazole, MS(ESI): MH+: 172.1.

EXAMPLE 796 3-(2-Methylpropyl)-5-aminomethylisoxazole (VI)

30 Isovaleraldehyde (5.4 mL, 50 mmol) and hydroxylamine hydrochloride (3.5 g, 50.4 mmol) are vigorously stirred in water (6 mL). To this is added a solution of sodium carbonate (2.65 g, 25 mmol) in water (15 mL). This is vigorously stirred

overnight. The mixture is extracted with ether. The organic layer is washed with water, then dried over sodium sulfate, filtered and concentrated. This is used in subsequent reactions without further purification: MS(ESI): MH+: 102.1.

5 Propargylamine (8.0 mL, 117 mmol) is dissolved in methylene chloride (60 mL), and di-tert-butyl dicarbonate (25 g, 114 mmol) is added. This is stirred overnight, and concentrated to provide the BOC-protected propargylamine, which is used without further purification: MS(ESI): MNa+: 178.0.

10 BOC-propargylamine (6.2 g, 39.7 mmol) and isovaleroxime (3.97 g, 39.3 mmol) is dissolved in methylene chloride (60 mL), and triethylamine (0.55 mL, 3.95 mmol) is added. This is cooled to 0 degrees C, and bleach (5% aqueous solution, 59.1 g) is added dropwise with vigorous stirring. After addition is
15 complete, the mixture is allowed to warm to 20-25 degrees C over 22 hours. The layers are separated, and the aqueous layer is extracted with methylene chloride (2 x). The combined organic extracts are washed with saline, dried over magnesium sulfate, filtered and concentrated. The residue is purified by
20 chromatography (silica gel, ethyl acetate/hexanes 5-10%) to give the BOC-protected title compound, MS(ESI): MH+: 255.3.

BOC-protected 3-(2-methylpropyl)-5-aminomethylisoxazole (2.4 g, 9.3 mmol) is dissolved in methylene chloride (10 mL) and treated with trifluoroacetic acid (10 mL) at 20-25 degrees
25 C. This is stirred at 20-25 degrees C for 70 minutes, then concentrated. The product is dissolved in methylene chloride, and washed with aqueous potassium carbonate (1 M) until basic (pH = 11). The organic layer is isolated, dried over sodium sulfate, filtered and concentrated to give the title compound:
30 MS(ESI): MH+: 155.2.

EXAMPLE 797 tert-butyl (3R)-2-oxo-1-propylazepanylcarbamate (VI)

To N-t-Boc-D-Lys-OH (10 g, 41.4mmole) in DMF (4 liters) is added benzotriazol-1-yloxytripyrrolidino-phosphonium hexafluorophosphate (BOP, 18.3 g, 41.4mmole) and sodium bicarbonate (17.4 g, 206.8mmole); the reaction is stirred at 20-25 degrees C for 12 hours. The reaction is then concentrated to 50 ml volume and diluted with ethyl acetate and washed with sodium bicarbonate 3x, water, 1M potassium bisulfate and brine, dried and concentrated. Purification by chromatography on silica gel afforded 5.05 g of the tert-butyl (3R)-2-oxoazepanylcarbamate as a solid; the procedure employed is similar to that described in *J.Med.Chem.* 1999, 4193. M+H-(t-Boc) (m/e=129.2), M+Na (m/e=251.1).

To the above lactam (2 g, 8.77mmole) in dry THF (20 ml) is added n-butyllithium /hexane (2.5 M, 5.3 ml, 13.2 mmole) at -78 degrees C, the reaction is stirred for 1 hour and 1-bromopropane (3.2 ml, 35.1 mmole) is added. The reaction is stirred for 1 hour and the cold bath removed and stirring continued for another 16 hours. Tetrabutylammonium iodide (0.49 g, 2.63mmole) is added and the reaction stirred for another 16 hours. The reaction is partitioned between ethyl acetate/hydrochloric acid + ice + water, the mixture is washed with water and saline and concentrated. Purification by chromatography on silica gel afforded the title compound, MS (M+Na+) 293.3.

EXAMPLE 798 N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl}-5-methyl-N³,N³-dipropylisophthalamide (X)

Following the procedure described in *J. Am. Chem. Soc.* 1986, 3150, the trifluoroacetic acid salt of N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N³,N³-dipropylisophthalamide (92.9 mg, 0.117 mmol) is dissolved in triethylamine (0.2 M, 0.6 mL) before the addition

- of $\text{PdCl}_2(\text{PPh}_3)_2$ (3.3 mg, 0.005 mmol), and copper (I) iodide (1.1 mg, 0.006 mmol). The reaction is heated to reflux. While the reaction is refluxing, trimethylsilylacetylene (0.02 ml, 0.14 mmol) is added via syringe. The reaction is refluxed for 3 hour
- 5 under N_2 (g), and the reaction cooled to 20-25 degrees C before partitioning between aqueous sodium bicarbonate and ethyl acetate. The product is extracted with ethyl acetate (3 x), washed with saline, dried over sodium sulfate, and filtered before the removal of solvent under reduced pressure.
- 10 The TMS protected acetylene (0.117 mmol) is dissolved in methanol (0.2 M, 0.5 mL) before the addition of potassium hydroxide (1M, 0.7 mL, 0.7 mmol). The reaction is stirred at 20-25 degrees C for 6 hours, at which point the mixture is partitioned between sodium bicarbonate and ethyl acetate. The
- 15 product is extracted with ethyl acetate (3 x), washed with saline, dried over sodium sulfate, and filtered before the removal of solvent under reduced pressure. Column chromatography (silica gel; 1.5-2 % isopropanol/chloroform under basic conditions; a few drops of ammonium hydroxide per
- 20 100 mL of elution solvent) gives the title compound, MS m/z $(\text{M}+\text{H})^+ = 576.3$.

EXAMPLE 799 1-phenylcyclopropylamine (VI)

- Following the procedure described in N.W. Werner et.al.,
- 25 J. Org. Syn. Coll. Vol. 5, 273-276, sodium azide (0.915g, 14.1 mmol) is slowly added to a solution of 1-phenylcyclopropanecarboxylic acid (1.0 g, 6.1 mmol) in concentrated sulfuric acid (5 ml) and dichloromethane (10 ml). The sodium sulfate precipitated out of solution. The reaction mixture is
- 30 heated to 50 degrees C for 17 hours and then cooled to 0 degrees C. The mixture is basified to pH = 11 with sodium hydroxide (1N) and extracted with dichloromethane (2 x). The organic layers are combined, dried over sodium sulfate, filtered and

concentrated. The residue is purified by chromatography (silica gel; isopropyl alcohol/chloroform/ammonium hydroxide 4/95/1) to give the title compound, MS (ESI+) for $C_9H_{11}N$ m/z $(M+H)^+ = 134$.

5

EXAMPLE 800 7-methoxy-1,2,3,4-tetrahydro-1-naphthalenamine
(VI)

7-Methoxy-1-tetralone (2.0 g, 11.3 mmol), hydroxylamine hydrochloride (1.56 g, 22.6 mmol) and sodium acetate (1.8g, 22.6 mmol) are suspended in ethanol/water (3/1, 40 mL). The mixture is heated for 45 min. at 100 degrees C. The mixture is allowed to cool overnight and the precipitate obtained is filtered and washed with water to yield an intermediate oxime, MS (ES) $(M+H): 192.1$. The oxime is dissolved in glacial acetic acid (25 ml) and palladium/carbon (500 mg) is added and the mixture hydrogenated under 50 psi at 20-25 degrees C overnight. The catalyst is filtered over diatomaceous earth and washed with methanol. The combined filtrates are concentrated. The concentrate is triturated with ether to give the title compound, MS (CI) $(MH+): 178.2$.

10
15
20

EXAMPLE 801 3-Amino-N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-methylbutanamide dihydrochloride

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII) and reacting it with 3-[(tert-butoxycarbonyl)amino]-2-methylbutanoic acid (IX), and then treatment with HCl, the title compound is obtained, HRMS calculated = 400.2600, observed = 400.2596.

25
30

EXAMPLE 802 N-{(1S,2R)-1-Benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-ethylhexanamide hydrochloride

Following the general procedure of EXAMPLE 6 and making
5 non-critical variations but starting with (2R,3S)-3-amino-1-[(3-methoxybenzyl)amino]-4-phenyl-2-butanol (VIII) and reacting it with 2-ethylhexanoic acid (IX), the title compound is obtained, HRMS calculated = 427.2961, observed = 427.2961.

10 EXAMPLE 803 N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-3-[(isobutylsulfonyl)amino]propanamide trifluoroacetate

Following the general procedure of CHART I and making non-critical variations but starting with (2R,3S)-3-amino-1-[(3-iodobenzyl)amino]-4-phenyl-2-butanol (VIII) and reacting it
15 with N-(isobutylsulfonyl)-beta-alanine (IX), the title compound is obtained, HRMS calculated = 588.1393, observed = 588.1379.

EXAMPLE 804 N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-N³-(isobutylsulfonyl)-beta-alaninamide trifluoroacetate
20

Following the general procedure of CHART I and making non-critical variations but starting with (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-iodobenzyl)amino]-2-butanol (VIII) and
25 reacting it with N-(isobutylsulfonyl)-beta-alanine (IX), the title compound is obtained, M + H = 624.

EXAMPLE 805 5-bromo-N¹-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-N³,N³-dipropylisophthalamide hydrochloride
30

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with (2R,3S)-3-amino-4-

(3,5-difluorophenyl)-1-[(3-iodobenzyl)amino]-2-butanol (VIII) and reacting it with 3-bromo-5-[(dipropylamino)carbonyl]benzoic acid (IX), the title compound is obtained, $M + H = 745$.

- 5 EXAMPLE 806 N^1 -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-phenylcyclopropyl)amino]propyl}-5-methyl- N^3,N^3 -dipropylisophthalamide

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with (2R,3S)-3-amino-4-
10 (3,5-difluorophenyl)-1-[(1-phenylcyclopropyl)amino]-2-butanol (VIII) and reacting it with 3-[(dipropylamino)carbonyl]-5-methylbenzoic acid (IX), the substituted amine (X) is obtained. $MH^+ = 578$.

- 15 EXAMPLE 806.1 5-bromo- N^1 -[(1S,2R)-3-[(3-bromobenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]- N^3,N^3 -dipropylisophthalamide

Following the general procedure of EXAMPLE 6 and making non-critical variations but starting with (2R,3S)-3-amino-1-
20 [(3-bromobenzyl)amino]-4-(3,5-difluorophenyl)-2-butanol hydrochloride (VIII) and reacting it with 3-bromo-5-[(dipropylamino)carbonyl]benzoic acid (IX), the title compound is obtained, NMR (500 MHz, CD_3OD): δ 7.85, 7.72, 7.69, 7.67, 7.52, 7.51, 6.88, 6.86, 6.74, 4.39-4.13, 3.95-3.88, 3.53-2.96,
25 2.82, 1.70, 1.50, 0.99 and 0.71.

EXAMPLES 807-1,064

Following the general procedure of CHART A and EXAMPLES 4-
6, 321-327, 329-464, 466-527, 529-579, 597-619 and 633-708 and
30 making non-critical variations, and using the appropriate amines (VIII) and amide forming agents (IX), the following compounds are obtained.

EXAMPLE	Substituted Amine (X)
807	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-5-(2-propylpentanoyl)benzamide
808	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-(2-ethylpentanoyl)-5-methylbenzamide
809	N-{(1S,2R)-1-benzyl-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-methyl-5-(2-propylpentanoyl)benzamide
810	N-{(1S,2R)-1-benzyl-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl}-3-methyl-5-(2-propylpentanoyl)benzamide
811	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-(2-ethylbutanoyl)-5-methylbenzamide
812	N ¹ -{(1S,2R)-1-benzyl-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-(2-propylpentanoyl)isophthalamide
813	N-{(1S,2R)-1-benzyl-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-(2-ethylpentanoyl)-5-methylbenzamide
814	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-(2-propylpentanoyl)isophthalamide
815	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(2-propylpentanoyl)isophthalamide
816	N-[(1S,2R)-3-[(3-ethylbenzyl)amino]-2-hydroxy-1-(4-hydroxybenzyl)propyl]-3-methyl-5-(2-propylpentanoyl)benzamide

817	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- [(3-methoxybenzyl) amino] propyl } -3-methyl-5-(2-propylpentanoyl) benzamide
818	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } -3-methyl-5-(2-propylpentanoyl) benzamide
819	N ¹ - ((1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- { [3-(3-pyridinyl) benzyl] amino } propyl) -5-methyl-N ³ ,N ³ -dipropylisophthalamide
820	N ¹ - ((1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- { [3-(4-pyridinyl) benzyl] amino } propyl) -5-methyl-N ³ ,N ³ -dipropylisophthalamide
821	N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- [(3-methoxybenzyl) amino] propyl } - N ³ ,N ³ -dipropyl-5-(1-propynyl) isophthalamide
822	N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } - N ³ ,N ³ -dipropyl-5-(1-propynyl) isophthalamide
823	N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } - N ³ ,N ³ -dipropyl-5-(2-propynyl) isophthalamide
824	N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- [(3-methoxybenzyl) amino] propyl } - N ³ ,N ³ -dipropyl-5-(2-propynyl) isophthalamide

825	N ¹ -{(1S,2R)-1-(cyclohexylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
826	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(3-thienylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
827	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-thienylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
828	N ¹ -{(1S)-1-[(1R)-2-(benzylamino)-1-hydroxyethyl]-3-butynyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
829	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-thienylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
830	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(2-thienylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
831	N ¹ -{(1S,2R)-1-(3-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
832	N ¹ -{(1S,2R)-3-(benzylamino)-1-[4-(benzyloxy)benzyl]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
833	N ¹ -{(1S,2R)-1-(2-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-

	methyl- N ³ ,N ³ -dipropylisophthalamide
834	N ¹ -[(1S,2R)-3-(benzylamino)-1-(cyclohexylmethyl)-2-hydroxypropyl]-5-methyl- N ³ ,N ³ -dipropylisophthalamide
835	N ¹ -{(1S,2R)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N ³ ,N ³ -dipropylisophthalamide
836	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(1-naphthylmethyl)propyl]- N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
837	2,3,5-trideoxy-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-5-[(3-methoxybenzyl)amino]-1-S-phenyl-1-thio-D-erythouro-pentitol
838	N ¹ -[(1S,2R)-3-(benzylamino)-1-(3-furylmethyl)-2-hydroxypropyl]-5-methyl- N ³ ,N ³ -dipropylisophthalamide
839	N ¹ -{(1S)-1-[(1R)-1-hydroxy-2-[(3-methoxybenzyl)amino]ethyl]-3-methylbutyl}-5-methyl- N ³ ,N ³ -dipropylisophthalamide
840	N ¹ -[(1S,2R)-3-(benzylamino)-1-(4-fluorobenzyl)-2-hydroxypropyl]- N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
841	N ¹ -{(1S,2R)-1-(4-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl- N ³ ,N ³ -dipropylisophthalamide
842	N ¹ -[(1S,2R)-3-(benzylamino)-1-(2-

	furylmethyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
843	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(1-naphthylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
844	N ¹ -{(1S)-1-[(1R)-2-(benzylamino)-1-hydroxyethyl]-3-methylbutyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
845	N ¹ -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
846	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(4-hydroxybenzyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
847	N ¹ -((1S)-1-{(1R)-1-hydroxy-2-[(3-methoxybenzyl)amino]ethyl}-3-butynyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
848	N ¹ -((1S)-1-{(1R)-1-hydroxy-2-[(3-methoxybenzyl)amino]ethyl}-3-butynyl)-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
849	5-(benzylamino)-2,3,5-trideoxy-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-1-S-phenyl-1-thio-D-erythouro-pentitol
850	N ¹ -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -

	dipropyl-1,3,5-benzenetricarboxamide
851	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(4-hydroxybenzyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
852	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(1-naphthylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
853	N ¹ -{(1S)-1-[(1R)-2-(benzylamino)-1-hydroxyethyl]-3-methylbutyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
854	N ¹ -{(1S,2R)-1-(4-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
855	*N ¹ -[(1S,2R)-3-(benzylamino)-1-(3-furylmethyl)-2-hydroxypropyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
856	N ¹ -{(1S)-1-[(1R)-1-hydroxy-2-[(3-methoxybenzyl)amino]ethyl]-3-methylbutyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
857	N ¹ -[(1S,2R)-3-(benzylamino)-1-(4-fluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
858	N ¹ -[(1S,2R)-3-(benzylamino)-1-(2-furylmethyl)-2-hydroxypropyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
859	N ¹ -{(1S,2R)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -

	dipropyl-1,3,5-benzenetricarboxamide
860	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(1-naphthylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
861	N ¹ -{(1S,2R)-1-(cyclohexylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
862	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(2-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
863	N ¹ -{(1S,2R)-1-(3-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
864	N ¹ -{(1S,2R)-3-(benzylamino)-1-[4-(benzyloxy)benzyl]-2-hydroxypropyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
865	N ¹ -{(1S,2R)-1-(2-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
866	N ¹ -[(1S,2R)-3-(benzylamino)-2-hydroxy-1-(3-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
867	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
868	N ¹ -{(1S)-1-[(1R)-2-(benzylamino)-1-hydroxyethyl]-3-butynyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide

869	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
870	N ¹ -{(1S,2R)-1-(cyclohexylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
871	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
872	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-thienylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
873	N ¹ -{(1S,2R)-1-(2-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
874	N ¹ -{(1S,2R)-1-(3-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
875	N ¹ -{(1S,2R)-2-hydroxy-1-(4-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
876	N ¹ -((1S)-1-{(1R)-1-hydroxy-2-[(3-methoxybenzyl)amino]ethyl}-3-methylbutyl)-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide

877	N ¹ -{(1S,2R)-1-(4-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
878	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(1-naphthylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
879	N ¹ -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
880	N ¹ -{(1S,2R)-2-hydroxy-1-[3-(hydroxymethyl)benzyl]-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
881	N ¹ -{(1S,2R)-3-[(3-ethylbenzyl)amino]-2-hydroxy-1-[3-(hydroxymethyl)benzyl]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
882	N ¹ -{(1S,2R)-2-hydroxy-1-[3-(hydroxymethyl)benzyl]-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
883	N ¹ -{(1S,2R)-2-hydroxy-1-[4-(hydroxymethyl)benzyl]-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
884	N ¹ -{(1S,2R)-3-[(3-ethylbenzyl)amino]-2-hydroxy-1-[4-(hydroxymethyl)benzyl]propyl}-5-methyl-

	N ³ ,N ³ -dipropylisophthalamide
885	N ¹ -{(1S,2R)-2-hydroxy-1-[4-(hydroxymethyl)benzyl]-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
886	N ¹ -{(1S,2R)-1-(3-fluoro-5-hydroxybenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
887	N ¹ -[(1S,2R)-3-[(3-ethylbenzyl)amino]-1-(3-fluoro-5-hydroxybenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
888	N ¹ -{(1S,2R)-1-(3-fluoro-5-hydroxybenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
889	N ¹ -{(1S,2R)-1-[3-(benzyloxy)-5-fluorobenzyl]-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
890	N ¹ -{(1S,2R)-1-[3-(benzyloxy)-5-fluorobenzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
891	N-[(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-3-[(dipropylamino)sulfonyl]propanamide
892	N ¹ -{(1S,2R)-1-[4-(benzyloxy)benzyl]-2-hydroxy-3-[(3-

	methoxybenzyl) amino] propyl} -N ⁵ ,N ⁵ - dipropylpentanediamide
893	3-[(dipropylamino) sulfonyl] -N-[(1S,2R)-2- hydroxy-3-[(3-methoxybenzyl) amino]-1-(1- naphthylmethyl) propyl] propanamide
894	N ¹ -[(1S,2R)-2-hydroxy-3-[(3- methoxybenzyl) amino]-1-(1- naphthylmethyl) propyl] - N ⁵ ,N ⁵ - dipropylpentanediamide
895	3-[(dipropylamino) sulfonyl] -N-{(1S,2R)-1- (4-fluorobenzyl)-2-hydroxy-3-[(3- methoxybenzyl) amino] propyl} propanamide
896	N ¹ -{(1S,2R)-1-(4-fluorobenzyl)-2-hydroxy- 3-[(3-methoxybenzyl) amino] propyl} - N ⁵ ,N ⁵ - dipropylpentanediamide
897	3-[(dipropylamino) sulfonyl] -N-{(1S,2R)-2- hydroxy-1-(4-hydroxybenzyl)-3-[(3- methoxybenzyl) amino] propyl} propanamide
898	N ¹ -{(1S,2R)-2-hydroxy-1-(4- hydroxybenzyl)-3-[(3- methoxybenzyl) amino] propyl} - N ⁵ ,N ⁵ - dipropylpentanediamide
899	3-[(dipropylamino) sulfonyl] -N-{(1S,2R)-1- (3-furylmethyl)-2-hydroxy-3-[(3- methoxybenzyl) amino] propyl} propanamide
900	N ¹ -{(1S,2R)-1-(2-furylmethyl)-2-hydroxy- 3-[(3-methoxybenzyl) amino] propyl} - N ⁵ ,N ⁵ - dipropylpentanediamide
901	3-[(dipropylamino) sulfonyl] -N-{(1S,2R)-1- (2-furylmethyl)-2-hydroxy-3-[(3- methoxybenzyl) amino] propyl} propanamide

902	N ¹ -{(1S,2R)-1-(3-furylmethyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ⁵ ,N ⁵ -dipropylpentanediamide
903	3-[(dipropylamino)sulfonyl]-N-[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-thienylmethyl)propyl]propanamide
904	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-thienylmethyl)propyl]-N ⁵ ,N ⁵ -dipropylpentanediamide
905	3-[(dipropylamino)sulfonyl]-N-[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-thienylmethyl)propyl]propanamide
906	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-thienylmethyl)propyl]-N ⁵ ,N ⁵ -dipropylpentanediamide
907	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-3-[[2R]-1-ethylpyrrolidinyl]carbonyl]-5-methylbenzamide
908	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-3-[[2S]-1-ethylpyrrolidinyl]carbonyl]-5-methylbenzamide
909	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl]-3-[(1-ethyl-1H-imidazol-2-yl)carbonyl]-5-

	methylbenzamide
910	N-((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl)-3-[(1-ethyl-4-methyl-1H-imidazol-5-yl)carbonyl]-5-methylbenzamide
911	N ¹ -((1S,2S)-1-(3,5-difluorobenzyl)-2-hydroxy-2-{1-[(3-methoxybenzyl)amino]cyclopropyl}ethyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
912	N ¹ -((1S,2S)-1-(3,5-difluorobenzyl)-2-{1-[(3-ethylbenzyl)amino]cyclopropyl}-2-hydroxyethyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
913	(1R,2R,3R)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ² ,N ² -dipropyl-1,2,3-cyclopropanetricarboxamide
914	(1R,2R,3R)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-phenyl-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
915	(1R,2R,3R)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
916	(1R,2R,3S)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-methyl-

	N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
917	(1R,2R,3S)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-phenyl-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
918	(1R,2R,3S)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ² ,N ² -dipropyl-1,2,3-cyclopropanetricarboxamide
919	(1R,2R,3S)-3-(2-amino-2-oxoethyl)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
920	(1R,2R,3R)-3-(2-amino-2-oxoethyl)-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ² ,N ² -dipropyl-1,2-cyclopropanedicarboxamide
921	(1R,2R,3S)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[2-(dipropylamino)-2-oxoethyl]-3-methylcyclopropanecarboxamide
922	(1R,2R,3R)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[2-(dipropylamino)-2-oxoethyl]-3-methylcyclopropanecarboxamide

923	(1S,2R,3R)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[2-(dipropylamino)-2-oxoethyl]-3-phenylcyclopropanecarboxamide
924	(1S,2R,3S)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[2-(dipropylamino)-2-oxoethyl]-3-phenylcyclopropanecarboxamide
925	(1S,2R,3R)-N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-[2-(dipropylamino)-2-oxoethyl]-1,2-cyclopropanedicarboxamide
926	(1S,2R,3S)-N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-3-[2-(dipropylamino)-2-oxoethyl]-1,2-cyclopropanedicarboxamide
927	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-5-[[trifluoromethyl)sulfonyl]amino}isophthalamide
928	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-[[trifluoromethyl)sulfonyl]amino}isophthalamide

929	N ¹ -{(1S,2R)-1-benzyl-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-{[(trifluoromethyl)sulfonyl]amino}isophthalamide
930	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-{methyl[(trifluoromethyl)sulfonyl]amino}-N ³ ,N ³ -dipropylisophthalamide
931	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-{methyl[(trifluoromethyl)sulfonyl]amino}-N ³ ,N ³ -dipropylisophthalamide
932	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-5-{propyl[(trifluoromethyl)sulfonyl]amino}isophthalamide
933	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(methylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide
934	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-[(phenylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide
935	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-

	hydroxy-3-[(3-isopropylbenzyl)amino]propyl}-3-[(dipropylamino)sulfonyl]propanamide
936	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl]-3-[(dipropylamino)sulfonyl]propanamide
937	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(dimethylamino)benzyl]amino}-2-hydroxypropyl]-3-[(dipropylamino)sulfonyl]propanamide
938	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethyl-1,3-thiazol-5-yl)methyl]amino}-2-hydroxypropyl]-3-[(dipropylamino)sulfonyl]propanamide
939	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-isobutyl-1,3-thiazol-5-yl)methyl]amino}propyl]-3-[(dipropylamino)sulfonyl]propanamide
940	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(3-isobutyl-5-isoxazolyl)methyl]amino}propyl]-3-[(dipropylamino)sulfonyl]propanamide
941	N-[(1S,2R)-3-[(3-cyclopropylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-3-[(dipropylamino)sulfonyl]propanamide
942	N ¹ -[(1S,2R)-3-[(3-cyclopropylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
943	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(1,3-thiazol-2-

	yl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
944	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(1,3-oxazol-2-yl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
945	N ¹ -[(1S,2R)-3-[(3-acetylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
946	N ¹ -[(1S,2R)-3-[(3-acetylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
947	N ¹ -[(1S,2R)-3-[(3-acetylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-(aminosulfonyl)-N ³ ,N ³ -dipropylisophthalamide
948	N ¹ -[(1S,2R)-3-[(3-acetylbenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-(methylsulfonyl)-N ³ ,N ³ -dipropylisophthalamide
949	N ¹ -[(1S,2R)-3-{[3-(diethylamino)benzyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
950	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(4-morpholinyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
951	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-

	hydroxy-3-{ [3-(1-piperazinyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
952	N ¹ -[(1S,2R)-3-{ [3-(aminosulfonyl)benzyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
953	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-({3-[(dimethylamino)sulfonyl]benzyl}amino)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
954	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [3-(1-piperidinylsulfonyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
955	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [3-(methylsulfonyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
956	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{ [3-(isopropylsulfonyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
957	N ¹ -[(1S,2R)-3-{ [3-(aminocarbonyl)benzyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
958	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-({3-[(dimethylamino)carbonyl]benzyl}amino)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -

	dipropylisophthalamide
959	N ¹ -[(1S,2R)-3-[(3-cyanobenzyl)amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
960	3-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino}methyl)phenylcarbamate
961	3-({[(2R,3S)-4-(3,5-difluorophenyl)-3-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}amino)-2-hydroxybutyl]amino}methyl)phenyl dimethylcarbamate
962	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(1-propynyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
963	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(3-methyl-1-butyryl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
964	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(2-propynyl)benzyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
965	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(5-isobutyl-1,3,4-oxadiazol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
966	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(5-ethyl-1,3,4-oxadiazol-2-

	yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
967	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(5-ethyl-1,3,4-thiadiazol-2-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
968	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(5-isobutyl-1,3,4-thiadiazol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
969	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(3-ethyl-1,2,4-thiadiazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
970	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(3-isobutyl-1,2,4-thiadiazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
971	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(3-isobutyl-1,2,4-oxadiazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
972	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[3-(3-ethyl-1,2,4-oxadiazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
973	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethyl-1,3-oxazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide

974	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-isobutyl-1,3-oxazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
975	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-isobutyl-1,3,4-oxadiazol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
976	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-isobutyl-1,3,4-thiadiazol-2-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
977	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(5-ethyl-1,3,4-thiadiazol-2-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
978	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(5-ethyl-1,3,4-oxadiazol-2-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
979	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3-ethyl-1,2,4-oxadiazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
980	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(3-ethyl-1,2,4-thiadiazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
981	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(3-isobutyl-1,2,4-thiadiazol-

	5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
982	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(3-isobutyl-1,2,4-oxadiazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
983	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethyl-2H-tetraazol-5-yl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
984	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-isobutyl-2H-tetraazol-5-yl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
985	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethyl-4-pyrimidinyl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
986	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(2-isopropyl-4-pyrimidinyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
987	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(2-ethynyl-4-pyrimidinyl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
988	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(6-isopropyl-4-pyrimidinyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide

989	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- ({[6-(dimethylamino)-4- pyrimidinyl]methyl}amino)-2- hydroxypropyl]-5-methyl-N ³ ,N ³ - dipropylisophthalamide
990	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- ({[2-(dimethylamino)-4- pyrimidinyl]methyl}amino)-2- hydroxypropyl]-5-methyl-N ³ ,N ³ - dipropylisophthalamide
991	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- ({[4-(dimethylamino)-2- pyrimidinyl]methyl}amino)-2- hydroxypropyl]-5-methyl-N ³ ,N ³ - dipropylisophthalamide
992	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2- hydroxy-3-{[(4-isopropyl-2- pyrimidinyl)methyl]amino}propyl]-5- methyl-N ³ ,N ³ -dipropylisophthalamide
993	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- {[(4-ethyl-2-pyrimidinyl)methyl]amino}-2- hydroxypropyl]-5-methyl-N ³ ,N ³ - dipropylisophthalamide
994	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3- {[(5-ethyl-3-pyridazinyl)methyl]amino}-2- hydroxypropyl]-5-methyl-N ³ ,N ³ - dipropylisophthalamide
995	N ³ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-{[3- (dimethylamino)benzyl]amino}-2- hydroxypropyl]-N ⁵ ,N ⁵ -dipropyl-3,5-

	pyridinedicarboxamide
996	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(5-isopropyl-3-pyridazinyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
997	N ³ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(1-propynyl)benzyl]amino}propyl)-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide
998	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(6-isopropyl-4-pyridazinyl)methyl]amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
999	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl}-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide
1000	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(6-ethyl-4-pyridazinyl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1001	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropylbenzyl)amino]propyl}-N ⁵ ,N ⁵ -dipropyl-3,5-pyridinedicarboxamide
1002	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(6-ethyl-2-pyrazinyl)methyl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1003	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-

	N ⁵ , N ⁵ -dipropyl-3,5-pyridinedicarboxamide
1004	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(6-isopropyl-2-pyrazinyl)methyl]amino}propyl)-5-methyl-N ³ , N ³ -dipropylisophthalamide
1005	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3,4,5-trifluorobenzyl)propyl]-5-methyl-N ³ , N ³ -dipropylisophthalamide
1006	N ¹ -((1S,2R)-2-hydroxy-1-(3,4,5-trifluorobenzyl)-3-{[3-(trifluoromethyl)benzyl]amino}propyl)-5-methyl-N ³ , N ³ -dipropylisophthalamide
1007	N ¹ -((1S,2R)-2-hydroxy-1-(2,3,5,6-tetrafluorobenzyl)-3-{[3-(trifluoromethyl)benzyl]amino}propyl)-5-methyl-N ³ , N ³ -dipropylisophthalamide
1008	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2,3,5,6-tetrafluorobenzyl)propyl]-5-methyl-N ³ , N ³ -dipropylisophthalamide
1009	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1R,2S)-2-hydroxy-6-methoxy-2,3-dihydro-1H-inden-1-yl]amino}propyl)-5-methyl-N ³ , N ³ -dipropylisophthalamide
1010	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1R,2S)-2-hydroxy-6-methoxy-2,3-dihydro-1H-inden-1-yl]amino}propyl)-N ³ , N ³ -dipropyl-1,3,5-benzenetricarboxamide

1011	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(1R,2S)-6-ethyl-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino}-2-hydroxypropyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1012	N ¹ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[(1R,2S)-6-ethyl-2-hydroxy-2,3-dihydro-1H-inden-1-yl]amino}-2-hydroxypropyl)-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1013	N ¹ -{(1S,2R)-2-hydroxy-1-(1H-indol-5-ylmethyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1014	N ¹ -[(1S,2R)-3-[(3-ethylbenzyl)amino]-2-hydroxy-1-(1H-indol-5-ylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1015	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-methylbenzyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1016	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(3-methylbenzyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1017	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[3-(trifluoromethyl)benzyl]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1018	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[3-

	(trifluoromethyl)benzyl]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1019	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-pyridinylmethyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1020	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(2-pyridinylmethyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1021	N ¹ -{(1S,2R)-1-[3-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1022	N ¹ -{(1S,2R)-1-[3-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1023	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[3-(trifluoromethoxy)benzyl]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1024	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[3-(trifluoromethoxy)benzyl]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1025	N ¹ -{(1S,2R)-2-hydroxy-1-(3-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide

1026	N ¹ -{(1S,2R)-2-hydroxy-1-(3-hydroxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1027	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(4-methylbenzyl)propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1028	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-(4-methylbenzyl)propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1029	N ¹ -{(1S,2R)-1-(4-fluoro-3-methylbenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1030	N ¹ -{(1S,2R)-1-(4-fluoro-3-methylbenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1031	N ¹ -{(1S,2R)-1-(4-chlorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1032	N ¹ -{(1S,2R)-1-(4-chlorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1033	N ¹ -{(1S,2R)-2-hydroxy-1-(3-methoxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide

1034	N ¹ -{(1S,2R)-2-hydroxy-1-(3-methoxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1035	N ¹ -{(1S,2R)-2-hydroxy-1-(4-methoxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1036	N ¹ -{(1S,2R)-2-hydroxy-1-(4-methoxybenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1037	N ¹ -{(1S,2R)-1-(3-chloro-5-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1038	N ¹ -{(1S,2R)-1-(3-chloro-5-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1039	N ¹ -{(1S,2R)-1-(4-chloro-3-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1040	N ¹ -{(1S,2R)-1-(4-chloro-3-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1041	N ¹ -{(1S,2R)-1-(3,5-dichlorobenzyl)-2-hydroxy-3-[(3-

	methoxybenzyl) amino] propyl} -5-methyl-N ³ ,N ³ -dipropylisophthalamide
1042	N ¹ -{(1S,2R)-1-(3,5-dichlorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1043	N ¹ -{(1S,2R)-1-[4-(dimethylamino) benzyl]-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1044	N ¹ -{(1S,2R)-1-[4-(dimethylamino) benzyl]-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1045	N ¹ -{(1S,2R)-1-(3-chlorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1046	N ¹ -{(1S,2R)-1-(3-chlorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1047	N ¹ -{(1S,2R)-1-(3-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1048	N ¹ -{(1S,2R)-1-(3-fluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1049	N ¹ -{(1S,2R)-2-hydroxy-1-(4-isopropylbenzyl)-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide

1050	N ¹ -{(1S,2R)-2-hydroxy-1-(4-isopropylbenzyl)-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1051	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[(6-methoxy-2-pyridinyl)methyl]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1052	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[(6-methoxy-2-pyridinyl)methyl]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1053	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[(5-methyl-2-pyridinyl)methyl]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1054	N ¹ -{(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl)amino]-1-[(5-methyl-2-pyridinyl)methyl]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1055	N ¹ -{(1S,2R)-1-(3-fluoro-4-methylbenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1056	N ¹ -{(1S,2R)-1-(3-fluoro-4-methylbenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1057	N ¹ -{(1S,2R)-1-(3-fluoro-4-methoxybenzyl)-2-hydroxy-3-[(3-

	methoxybenzyl) amino] propyl} -5-methyl-N ³ ,N ³ -dipropylisophthalamide
1058	N ¹ -{(1S,2R)-1-(3-fluoro-4-methoxybenzyl)-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1059	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl) amino]-1-(2-methoxy-5-methylbenzyl) propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1060	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl) amino]-1-(2-methoxy-5-methylbenzyl) propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1061	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl) amino]-1-(1,3-thiazol-2-ylmethyl) propyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1062	N ¹ -[(1S,2R)-2-hydroxy-3-[(3-methoxybenzyl) amino]-1-(1,3-thiazol-2-ylmethyl) propyl]-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide
1063	N ¹ -{(1S,2R)-1-[(5-chloro-2-thienyl) methyl]-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide
1064	N ¹ -{(1S,2R)-1-[(5-chloro-2-thienyl) methyl]-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-N ³ ,N ³ -dipropyl-1,3,5-benzenetricarboxamide

EXAMPLES 1,065-1,155

Following the general procedure of CHART A and EXAMPLES 4-6, 321-327, 329-464, 466-527, 529-579, 597-619 and 633-708 and making non-critical variations, and using the appropriate amines (VIII) and amide forming agents (IX), the titled compounds are obtained.

EXAMPLE	Substituted Amine (X)
1,065	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-3-(1-pyrrolidinylcarbonyl)benzamide
1,066	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-2-[(methylsulfonyl)amino]-1,3-thiazole-4-carboxamide
1,067	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,068	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(propylsulfonyl)amino]-1,3-thiazole-4-carboxamide
1,069	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-3-(1-pyrrolidinylcarbonyl)benzamide
1,070	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[(propylsulfonyl)amino]-1,3-thiazole-4-carboxamide
1,071	N-{(1S,2R)-1-benzyl-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(methylsulfonyl)amino]-

01298 7

WO 03/072535

PCT/US03/07287

	1,3-oxazole-4-carboxamide
1,072	N-((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(3-ethylphenyl)cyclopropyl]amino}-2-hydroxypropyl)-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide

1,073	N-((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(3-ethylphenyl)-1-methylethyl]amino}-2-hydroxypropyl)-4-hydroxy-3-(1-pyrrolidinylcarbonyl)benzamide
1,074	N-((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(3-ethylphenyl)-1-methylethyl]amino}-2-hydroxypropyl)-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide
1,075	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide
1,076	N-((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(3-ethylphenyl)-1-methylethyl]amino}-2-hydroxypropyl)-5-methyl-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide
1,077	N-((1S,2R)-1-(3,5-difluorobenzyl)-3-{[1-(3-ethylphenyl)cyclopropyl]amino}-2-hydroxypropyl)-4-hydroxy-3-(1-pyrrolidinylcarbonyl)benzamide
1,078	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl}-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide
1,079	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide
1,080	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl)amino]-2-hydroxypropyl}-5-

	methyl-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,081	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-3-(1-piperidinylcarbonyl)benzamide
1,082	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,083	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide

1,084	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-4-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,085	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-3-(1-piperidinylcarbonyl)benzamide
1,086	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,087	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,088	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-4-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,089	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-3-(4-morpholinylcarbonyl)benzamide
1,090	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-[(ethylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,091	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,092	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-

	[(3-iodobenzyl) amino]propyl}-4- [(ethylsulfonyl) amino]-1,3-oxazole-2- carboxamide
1,093	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl }-4-hydroxy-3-(4-morpholinylcarbonyl) benzamide
1,094	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3-[(3-iodobenzyl) amino]propyl }-4-[(propylsulfonyl) amino]-1,3-oxazole-2-carboxamide
1,095	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3-[(3-methoxybenzyl) amino]propyl }-5-methyl-2-[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide

1,096	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4-[(methylsulfonyl)amino]-1,3-thiazole-2-carboxamide
1,097	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-3-(1-piperazinylcarbonyl)benzamide
1,098	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-[(methylsulfonyl)amino]-1,3-thiazole-2-carboxamide
1,099	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methyl-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,100	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(methylsulfonyl)amino]-1,3-oxazole-5-carboxamide
1,101	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-3-(1-piperazinylcarbonyl)benzamide
1,102	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-methyl-2-[(methylsulfonyl)amino]-1,3-oxazole-5-carboxamide
1,103	N ⁴ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(methylsulfonyl)amino]-1,3-oxazole-4,5-dicarboxamide
1,104	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-

	[(3-iodobenzyl) amino]propyl}-2- [(methanesulfonyl) amino]-1,3-oxazole-5- carboxamide
1,105	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-4-hydroxy- N ³ -methanophthalamide
1,106	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3- [(3-iodobenzyl) amino]propyl}-4-methyl-2- [(methanesulfonyl) amino]-1,3-oxazole-5- carboxamide

1,107	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(ethylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,108	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,109	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-N ³ -methylisophthalamide
1,110	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4-methyl-5-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,111	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[(ethylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,112	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-methyl-5-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,113	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ -ethyl-4-hydroxyisophthalamide
1,114	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(methylsulfonyl)amino]-1,3-oxazole-2-carboxamide
1,115	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-

	[(3-iodobenzyl) amino]propyl}-2- [(ethylsulfonyl) amino]-1,3-oxazole-4- carboxamide
1,116	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3- ethylbenzyl) amino]-2-hydroxypropyl}-5- [(methylsulfonyl) amino]-3-isoxazolecarboxamide
1,117	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3- ethylbenzyl) amino]-2-hydroxypropyl}-N ³ -ethyl- 4-hydroxyisophthalamide

1,118	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-[(methylsulfonyl)amino]-3-isoxazolecarboxamide
1,119	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(propylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,120	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-3-[(methylsulfonyl)amino]-5-isoxazolecarboxamide
1,121	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl-N ³ -ethyl-4-hydroxyisophthalamide
1,122	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(methylsulfonyl)amino]-5-isoxazolecarboxamide
1,123	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-[(propylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,124	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-(hydroxymethyl)-2-[(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,125	N ³ -(cyclopropylmethyl)-N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl-4-hydroxyisophthalamide
1,126	5-cyclopropyl-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-

	[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,127	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]-2-[(propylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,128	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl) amino]propyl]-5-isopropyl-2-[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,129	N ³ -(cyclopropylmethyl)-N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]-4-hydroxyisophthalamide
1,130	N-[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isopentylamino)propyl]-2-[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,131	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]-5-methyl-2-[(propylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,132	N-[(1S,2R)-3-(cyclopropylamino)-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-2-[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,133	N-[(1S,2R)-3-[(3-ethylbenzyl) amino]-2-hydroxy-1-(4-hydroxybenzyl)propyl]-2-[(methylsulfonyl) amino]-1,3-oxazole-4-carboxamide
1,134	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]-4-hydroxy-N ³ -isobutylisophthalamide

1,135	2- { [(cyclopropylmethyl) sulfonyl] amino } -N- { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } -1,3-oxazole-4-carboxamide
1,136	N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } -4-hydroxy-N ³ -isobutyl-N ³ -methylisophthalamide
1,137	N- { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } -2- [(isobutylsulfonyl) amino] -1,3-oxazole-4-carboxamide
1,138	N ³ - (cyclopropylmethyl) -N ¹ - { (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2-hydroxypropyl } -4-hydroxy-N ³ -methylisophthalamide
1,139	N- { (1S,2R) -1- (3,5-difluorobenzyl) -2-hydroxy-3- [(3-methoxybenzyl) amino] propyl } -2- [(isobutylsulfonyl) amino] -1,3-oxazole-4-carboxamide

1,140	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-N ³ -methyl-N ³ -propylisophthalamide
1,141	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(isobutylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,142	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-N ³ -methyl-N ³ -propylisophthalamide
1,143	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(phenylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,144	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ -ethyl-4-hydroxy-N ³ -propylisophthalamide
1,145	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(4-methylphenyl)sulfonyl]amino}-1,3-oxazole-4-carboxamide
1,146	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -ethyl-4-hydroxy-N ³ -propylisophthalamide
1,147	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(4-methylphenyl)sulfonyl]amino}-1,3-oxazole-4-carboxamide
1,148	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-

012987

	[(phenylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,149	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-4-hydroxy-N ³ , N ³ -dipropylisophthalamide

1,150	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[methyl(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,151	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-4-hydroxy-N ³ , N ³ -dipropylisophthalamide
1,152	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[methyl(methylsulfonyl)amino]-1,3-oxazole-4-carboxamide
1,153	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-4-hydroxy-N ³ , N ³ -dipropylisophthalamide
1,154	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(methylsulfonyl)amino]-1,3-thiazole-4-carboxamide
1,155	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-2-[(methylsulfonyl)amino]-1,3-thiazole-4-carboxamide

EXAMPLES 1,156-1,214

Following the general procedure of CHART A and EXAMPLES 4-
5 6, 321-327, 329-464, 466-527, 529-579, 597-619 and 633-708 and making non-critical variations, and using the appropriate amines (VIII) and amide forming agents (IX), the titled compounds are obtained.

EXAMPLE	Substituted Amine (X)	MH+
1,156	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-	532.

	methoxybenzyl) amino] propyl}-3-methyl-5-[propionyl (propyl) amino] benzamide	5
1,157	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-1-butyl-1H-indole-5-carboxamide	Anal . Foun d: C = 73.5 8; H = 7.44 ; N = 8.24 .
1,159	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-3-[butyl (propionyl) amino]-5-methylbenzamide	546. 3
1,160	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-4-methyl-1-propyl-1H-indole-6-carboxamide	500. 3
1,161	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl) amino] propyl}-1-(1-propylbutyl)-1H-indole-6-carboxamide	542. 2
1,162	N ¹ -[(1S,2R)-1-benzyl-2-hydroxy-3-[(2-oxo-2,3-dihydro-1,3-benzoxazol-6-yl) methyl] amino] propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	573. 3
1,163	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]-N ³ ,N ³ -dipropyl-5-	713. 0

	{[(trifluoromethyl)sulfonyl]amino}isophthalamide	
1,164	3-[[{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}amino]carbonyl]-5-[(dipropylamino)carbonyl]benzoic acid	576. 1
1,165	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-prop-1-ynylisophthalamide	604. 4
1,166	N-{(1S,2R)-1-benzyl-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-2-(dipropylamino)isonicotinamide	HRMS = 505. 3176
1,167	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(methylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,168	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(ethylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,169	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-[(propylsulfonyl)amino]isophthalamide	
1,170	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(isopropylsulfonyl)amino]-N ³ ,N ³ -	

	dipropylisophthalamide	
1,171	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(isobutylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,172	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-[(thien-2-ylsulfonyl)amino]isophthalamide	
1,173	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(2-furylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,174	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-[(1,3-thiazol-5-ylsulfonyl)amino]isophthalamide	
1,175	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(1,3-oxazol-5-ylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,176	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(1,3-oxazol-4-ylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,177	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-	

	hydroxypropyl}-N ³ ,N ³ -dipropyl-5-[(1,3-thiazol-4-ylsulfonyl)amino]isophthalamide	
1,178	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-{[(1-methyl-1H-imidazol-4-yl)sulfonyl]amino}-N ³ ,N ³ -dipropylisophthalamide	
1,179	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-[(phenylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,180	5-{[(5-cyanopyridin-2-yl)sulfonyl]amino}-N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropylisophthalamide	
1,181	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -dipropyl-5-({[5-(trifluoromethyl)pyridin-2-yl)sulfonyl]amino)isophthalamide	

1,182	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-{[(1-methyl-1H-imidazol-4-yl)sulfonyl]amino}benzamide	
1,183	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-({[5-(trifluoromethyl)pyridin-2-	

	yl) sulfonyl} amino) benzamide	
1,184	3-{ [(5-cyanopyridin-2-yl) sulfonyl] amino} -N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} benzamide	
1,185	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(phenylsulfonyl) amino] benzamide	
1,186	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(methylsulfonyl) amino] benzamide	
1,187	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(ethylsulfonyl) amino] benzamide	
1,188	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(propylsulfonyl) amino] benzamide	
1,189	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(isobutylsulfonyl) amino] benzamide	
1,190	N-{ (1S,2R) -1-(3,5-difluorobenzyl) -3-[(3-ethylbenzyl) amino] -2-hydroxypropyl} -3-[(isopropylsulfonyl) amino] benzamide	

1,191	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(1-ethylpropyl)sulfonyl]amino}benzamide	
1,192	3-[(cyclohexylsulfonyl)amino]-N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}benzamide	
1,193	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(1-propylbutyl)sulfonyl]amino}benzamide	
1,194	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(thien-2-ylsulfonyl)amino]benzamide	
1,195	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(2-furylsulfonyl)amino]benzamide	
1,196	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(isoxazol-5-ylsulfonyl)amino]benzamide	
1,197	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(isoxazol-3-ylsulfonyl)amino]benzamide	
1,198	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(3-furylsulfonyl)amino]benzamide	

1,199	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl]-3-[(thien-3-ylsulfonyl)amino]benzamide	
1,200	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl]-3-[(1,3-thiazol-4-ylsulfonyl)amino]benzamide	
1,201	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl]-3-[(1,3-thiazol-5-ylsulfonyl)amino]benzamide	
1,202	N-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl]-3-[(1,3-thiazol-2-ylsulfonyl)amino]benzamide	
1,203	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isopentylamino)propyl]-N ³ ,N ³ -dipropyl-5-[[trifluoromethyl)sulfonyl]amino}isophthalamide	
1,204	N ¹ -[(1S,2R)-3-amino-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-N ³ ,N ³ -dipropyl-5-[[trifluoromethyl)sulfonyl]amino}isophthalamide	
1,205	N ¹ -[(1S,2R)-3-amino-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-[(methylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	

1,206	N ¹ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isopentylamino)propyl]-5-[(methylsulfonyl)amino]-N ³ ,N ³ -dipropylisophthalamide	
1,207	N ¹ -(tert-butyl)-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}isophthalamide	
1,208	N ¹ -(tert-butyl)-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylisophthalamide	
1,209	5-bromo-N ¹ -(tert-butyl)-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}isophthalamide	
1,210	3-tert-butoxy-N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}benzamide	
1,211	3-tert-butoxy-N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylbenzamide	
1,212	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[[trifluoromethyl)sulfonyl]amino}benzamide	
1,213	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-	

	(trifluoromethoxy) benzamide	
1,214	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -3- [(3-ethylbenzyl) amino] -2- hydroxypropyl }-3-methyl-5- (trifluoromethoxy) benzamide	

EXAMPLES 1,215-1,259

Following the general procedure of CHART A and EXAMPLES 4-6, 321-327, 329-464, 466-527, 529-579, 597-619 and 633-708 and making non-critical variations, and using the appropriate amines (VIII) and amide forming agents (IX), the titled compounds are obtained.

EXAMPLE	Substituted Amine (X)	[M+H] +
1,215	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2- hydroxy-3- [(3-iodobenzyl) amino] propyl }-2-hydroxy-2- (4-methylphenyl) acetamide	581.4
1,216	N1-{ (1S,2R) -1- (3,5-difluorobenzyl) -2- hydroxy-3- [(3-iodobenzyl) amino] propyl }-4-hydroxy-N3-methylisophthalamide	610.4
1,217	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2- hydroxy-3- [(3-iodobenzyl) amino] propyl }-2-hydroxy-2- (4-methoxy-3-nitrophenyl) acetamide	642.4
1,218	5- (aminosulfonyl) -N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2- hydroxy-3- [(3-iodobenzyl) amino] propyl }-2-methoxybenzamide	646.5
1,219	N-{ (1S,2R) -1- (3,5-difluorobenzyl) -2- hydroxy-3- [(3-iodobenzyl) amino] propyl }-	650.5

	4-hydroxy-3-(pyrrolidin-1-ylcarbonyl)benzamide	
1,220	N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-2-[(methanesulfonyl)amino]-1,3-oxazole-4-carboxamide	621.4
1,221	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-(3,5-dimethylisoxazol-4-yl)-N ³ ,N ³ -dipropylisophthalamide	663.4
1,222	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-N ³ ,N ³ -dipropyl-5-(1,3-thiazol-2-yl)isophthalamide	651.4
1,223	3-(cyclohexylcarbonyl)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methylbenzamide	565.4
1,224	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N ³ -propylisophthalamide	540.4
1,225	3-[cyclohexyl(hydroxy)methyl]-N-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methylbenzamide	567.4
1,226	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-(4-methyl-1,3-oxazol-	647.5

	2-yl)-N ³ ,N ³ -dipropylisophthalamide	
1,227	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-N ⁵ ,N ⁵ -dipropylpyridine-3,5-dicarboxamide	567
1,228	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isobutyl-1,2,4-oxadiazol-5-yl)methyl] amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	600
1,229	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethynylbenzyl) amino]-2-hydroxypropyl}-N ⁵ ,N ⁵ -dipropylpyridine-3,5-dicarboxamide	563
1,230	N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropylbenzyl) amino]propyl}-N ⁵ ,N ⁵ -dipropylpyridine-3,5-dicarboxamide	581
1,231	N ¹ -[(1S,2R)-3-[(1-acetylpiperidin-4-yl) amino]-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl-N ³ ,N ³ -dipropylisophthalamide	587
1,232	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-pent-1-ynylbenzyl) amino]propyl}-5-methyl-N ³ ,N ³ -dipropylisophthalamide	618
1,233	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-(4-hydroxybut-1-ynyl)benzyl) amino]propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	620
1,234	1-{3-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl) amino] carbonyl]-5-	593

	methylbenzoyl}-L-prolinamide	
1,235	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -isopropyl-5-methylisophthalamide	538
1,236	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -ethyl-N ³ ,5-dimethylisophthalamide	538
1,237	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,5-dimethyl-N ³ -prop-2-ynylisophthalamide	548
1,238	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -isobutyl-5-methylisophthalamide	552
1,239	N ¹ -(sec-butyl)-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylisophthalamide	552
1,240	N ¹ -butyl-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylisophthalamide	552
1,241	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,N ³ -diethyl-5-methylisophthalamide	552
1,242	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ ,5-dimethyl-N ³ -	552


	propylisophthalamide	
1,243	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -isopropyl-N ³ ,5-dimethylisophthalamide	552
1,244	N ¹ -butyl-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ¹ ,5-dimethylisophthalamide	566
1,245	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -isobutyl-N ³ ,5-dimethylisophthalamide	566
1,246	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -ethyl-5-methyl-N ³ -propylisophthalamide	566
1,247	N ¹ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-N ³ -ethyl-N ³ -isopropyl-5-methylisophthalamide	566
1,248	N ¹ ,N ¹ -diallyl-N ³ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylisophthalamide	576
1,249	3-(azepan-1-ylcarbonyl)-N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-5-methylbenzamide	578
1,250	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-[(4-hydroxypiperidin-	580

	1-yl) carbonyl] -5-methylbenzamide	
1,251	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-3-[(3-hydroxypiperidin-1-yl) carbonyl] -5-methylbenzamide	580
1,252	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-N ³ ,N ³ -diisopropyl-5-methylisophthalamide	580
1,253	N ¹ -butyl-N ³ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-N ¹ -ethyl-5-methylisophthalamide	580
1,254	N ¹ -(cyclopropylmethyl)-N ³ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-5-methyl-N ¹ -propylisophthalamide	592
1,255	1-{3-[(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl]amino) carbonyl] -5-methylbenzoyl}-D-prolinamide	593
1,256	N ¹ -cyclohexyl-N ³ -(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl) amino]-2-hydroxypropyl}-N ¹ ,5-dimethylisophthalamide	592
1,257	N ¹ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[1-(3-methylphenyl) cyclopropyl] amino}propyl)-5-methyl-N ³ ,N ³ -dipropylisophthalamide	592
1,258	N ³ -(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(1,2,3,4-tetrahydronaphthalen-1-ylamino)propyl]-	579

	N ⁵ ,N ⁵ -diisopropylpyridine-3,5-dicarboxamide	
1,259	N-{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-ethylbenzyl)amino]-2-hydroxypropyl}-3-{[(trifluoromethyl)sulfonyl]amino}benzamide	586.1

Following the procedures described above and those in international patent application WO 99/33815 (Published July 8, 1999), the following prodrug compounds are made. WO 99/33815 is herein incorporated by reference in its entirety.

EXAMPLE 1260. 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)propyl 4-nitrophenyl carbonate (1260Y).




476 is first protected by methods well known in the art, as discussed above. For example, 476 is protected with t-butoxycarbonyl by treating the compound with di-tert-butyl dicarbonate in the presence of triethylamine, 4-(dimethylamino)pyridine (DMAP) and methylene chloride. A mixture of 3.7 mMol of protected 476 and 16 mMol of di-p-nitrophenyl carbonate in 10 ml of dimethylformamide is treated at 25 °C with 4 mMol of P4-phosphazene base (Fluka, 1M in

hexane). The mixture is stirred for 6h at 25 °C until all of the starting alcohol is consumed. The reaction mixture is partitioned between ethyl acetate and 1N hydrochloric acid. The organic layer is washed with 1N sodium hydroxide and brine, dried over magnesium sulfate and concentrated in vacuo.

Titration with dichloromethane gives protected 1260.

Deprotection of protected 1260 is conducted by methods well known in the art, as discussed above. For example, protected 1260 is treated with trifluoroacetic acid to yield 1260.

EXAMPLE 1261. 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)propyl 4-methylpiperazine-1-carboxylate (1261Y).




To 0.286 mM of protected 1260 dissolved in THF is added 1.14 mM of 1-methyl-piperazine and the mixture is stirred overnight at room temperature. All the solvents are then evaporated and the solid residue partitioned between EtOAc and water. The volatiles are removed and the residue is treated with 1:1 TFA/DCM over 30 min at room temperature to remove the Boc protecting groups, yielding product 1261.

EXAMPLES 1262-1267.

Following the general procedure of EXAMPLE 1261 but using the reagents of Column A instead of 1-methyl-piperazine, prodrug (Y) of Column B is obtained.

EX.	COL. A	COL. B; Compound Y
1262	N,N-dimethylamino-ethanol	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-(dimethylamino)ethyl carbonate
1263	N-acetyl-ethylenediamine	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-(acetylamino)ethylcarbamate
1264	Mono-N-Boc-piperazine	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl piperazine-1-carboxylate
1265	Mono-N-Boc-ethylenediamine	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-aminoethylcarbamate
1266	1,3-diamino-3-N-Boc-propane	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl 3-aminopropylcarbamate
1267	(3R)-(+)-3-Boc-aminopyrrolidine	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl (3R)-3-aminopyrrolidine-1-carboxylate

EXAMPLE 1268. 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)propyl O-benzyl-L-tyrosinate



476 is protected as described above. To 0.37mM of protected 476 dissolved in CH_2Cl_2 is added 0.41mM of N-CBz-O-Benzyl-L-tyrosine followed by 1.12mM of DCC (dicyclohexylcarbodiimide), followed by 0.23mM of DMAP. The reaction is stirred at rt for 24hr. Precipitates are removed by filtration. The filtrate is then concentrated in vacuo. The product is deprotected as described above and purified by methods well known in the art to give 1268.

EXAMPLES 1269-1275

Following the general procedure of EXAMPLE 1268 but using the reagents of Column A instead of N-CBz-O-Benzyl-L-tyrosine, compound (Y) of Column B is obtained.

EX.	COL. A	COL. B; Compound Y
1269	L-tyrosine	1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl L-tyrosinate
1270	3-Methoxy-	1-[(benzylamino)methyl]-3-(3,5-

	propionic acid	difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl 3-methoxypropanoate
1271	(2-Hydroxy- ethoxy)-acetic acid	1-[(benzylamino)methyl]-3-(3,5- difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl (2-hydroxyethoxy)acetate
1272	D-lysine	1-[(benzylamino)methyl]-3-(3,5- difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl D-lysinate
1273	[2-(2-Methoxy- ethoxy)- ethoxy]-acetic acid	1-[(benzylamino)methyl]-3-(3,5- difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl [2-(2- methoxyethoxy)ethoxy]acetate
1274	{2-[(2- Dimethylamino- ethyl)-methyl- amino]-ethoxy}- acetic acid	1-[(benzylamino)methyl]-3-(3,5- difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl {2-[[2-(dimethylamino)ethyl]- (methyl)amino]ethoxy}acetate
1275	(4-Methyl- piperazin-1- yl)-acetic acid	1-[(benzylamino)methyl]-3-(3,5- difluorophenyl)-2-({3- [(dipropylamino)carbonyl]benzoyl}amino)- propyl (4-methylpiperazin-1-yl)acetate

Following the general procedures outlined above and making non-critical variations, the following compounds (Y) are obtained.
 (1*R*,2*S*)-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-
 pent-1-ynylbenzyl)amino]methyl}propyl (4-methylpiperazin-
 1-yl)acetate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl 4-methylpiperazine-
1-carboxylate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl piperazine-1-
carboxylate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl {2-[[2-
(dimethylamino)ethyl](methyl)amino]ethoxy}acetate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl (2-
methoxyethoxy)acetate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl L-tyrosinate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl (3R)-3-
aminopyrrolidine-1-carboxylate,

(1R,2S)-3-(3,5-difluorophenyl)-2-({3-
[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{{(3-
pent-1-ynylbenzyl)amino}methyl}propyl tetrahydrofuran-3-

yl carbonate,

N'-[(1*S*,2*R*)-1-benzyl-3-(cyclohexylamino)-2-hydroxypropyl]-*N,N*-dipropylisophthalamide,

(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl 4-methylpiperazine-1-carboxylate,

(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl piperazine-1-carboxylate,

(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl {2-[[2-(dimethylamino)ethyl](methyl)amino]ethoxy}acetate,


(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl (2-methoxyethoxy)acetate,

(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl *L*-tyrosinate,


(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl (3*R*)-3-aminopyrrolidine-1-carboxylate,

(1*R*,2*S*)-1-[(cyclohexylamino)methyl]-2-({3-[(dipropylamino)carbonyl]benzoyl}-amino)-3-phenylpropyl tetrahydrofuran-3-yl carbonate.


CHART A




(I)




(II)




(III)




(IV)



(V)

CHART A - Continued

10



15







CHART A - Continued

(VIII)



+



(IX)



(X)



(A)



CHART A - Continued






CHART B














CHART C



(VII)

CHART D






CHART E

5




10

15




(IX - XXIII)


CHART F




(XXIV)



(XXV)




(XXVI)






(XXVII)






(XXII)

CHART G



5

10



(XXI)










CHART H

CHART I









CHART J




(XXI)



(XXXVII)






(XXXVIII)



(IX - XXXIX)


CHART K



15


5

10

CHART I


(XLII)

15




(XLIII)

20




(XLIV)


403



5



10

CHART L - Continued

15

↓








CHART M



(XLVIII)



(XLIX)



(IX - L)









CHART N

CHART O







5

CHART O - Continued


10



15







01298 7




(IX - LX)

CHART O - continued






(IX - LXIII)

CHART P


(LXIV)

5



(LXV)

10




(LXVI)

15




↓



(IX - LXVII)


CHART Q



5

10

CHART R



5

10









CHART S

(X,
 $\text{R}_\text{N} = \text{R}_\text{N-aryl}-\text{R}_\text{N-aryl}-\text{X}_\text{N}$)



(X,
 $\text{R}_\text{N} = \text{R}_\text{N-heteroaryl}-\text{R}_\text{N-aryl}-\text{X}_\text{N}$)






CHART T

CHART U


(LXXX)



(LXXXI)



(LXXXII)



(LXXXIII)



(LXXXIV)


CHART V






CHART W


CHART X

5


10

CHART Y

(XCIII)



(XCIV)



(XCV)

BIOLOGICAL EXAMPLES

Example A

Enzyme Inhibition Assay

5

The compounds of the invention are analyzed for inhibitory activity by use of the MBP-C125 assay. This assay determines the relative inhibition of beta-secretase cleavage of a model APP substrate, MBP-C125SW, by the compounds assayed as compared with an untreated control. A detailed description of the assay parameters can be found, for example, in U.S. Patent No. 5,942,400. Briefly, the substrate is a fusion peptide formed of maltose binding protein (MBP) and the carboxy terminal 125 amino acids of APP-SW, the Swedish mutation. The beta-secretase enzyme is derived from human brain tissue as described in Sinha et.al, 1999, *Nature* 40:537-540) or recombinantly produced as the full-length enzyme (amino acids 1-501), and can be prepared, for example, from 293 cells expressing the recombinant cDNA, as described in W000/47618.

20 Inhibition of the enzyme is analyzed, for example, by immunoassay of the enzyme's cleavage products. One exemplary ELISA uses an anti-MBP capture antibody that is deposited on precoated and blocked 96-well high binding plates, followed by incubation with diluted enzyme reaction supernatant, incubation with a specific reporter antibody, for example, biotinylated anti-SW192 reporter antibody, and further incubation with streptavidin/alkaline phosphatase. In the assay, cleavage of the intact MBP-C125SW fusion protein results in the generation of a truncated amino-terminal fragment, exposing a new SW-192 antibody-positive epitope at the carboxy terminus. Detection is effected by a fluorescent substrate signal on cleavage by the phosphatase. ELISA only detects cleavage following Leu 596 at the substrate's APP-SW 751 mutation site.

Specific Assay Procedure:

Compounds are diluted in a 1:1 dilution series to a six-point concentration curve (two wells per concentration) in one 96-plate row per compound tested. Each of the test compounds is prepared in DMSO to make up a 10 millimolar stock solution. The stock solution is serially diluted in DMSO to obtain a final compound concentration of 200 micromolar at the high point of a 6-point dilution curve. Ten (10) microliters of each dilution is added to each of two wells on row C of a corresponding V-bottom plate to which 190 microliters of 52 millimolar NaOAc, 7.9% DMSO, pH 4.5 are pre-added. The NaOAc diluted compound plate is spun down to pellet precipitant and 20 microliters/well is transferred to a corresponding flat-bottom plate to which 30 microliters of ice-cold enzyme-substrate mixture (2.5 microliters MBP-C125SW substrate, 0.03 microliters enzyme and 24.5 microliters ice cold 0.09% TX100 per 30 microliters) is added. The final reaction mixture of 200 micromolar compound at the highest curve point is in 5% DMSO, 20 millimolar NaAc, 0.06% TX100, at pH 4.5.

Warming the plates to 37 degrees C starts the enzyme reaction. After 90 minutes at 37 degrees C, 200 microliters/well cold specimen diluent is added to stop the reaction and 20 microliters/well is transferred to a corresponding anti-MBP antibody coated ELISA plate for capture, containing 80 microliters/well specimen diluent. This reaction is incubated overnight at 4 degrees C and the ELISA is developed the next day after a 2 hours incubation with anti-192SW antibody, followed by Streptavidin-AP conjugate and fluorescent substrate. The signal is read on a fluorescent plate reader.

Relative compound inhibition potency is determined by calculating the concentration of compound that showed a fifty percent reduction in detected signal (IC_{50}) compared to the

enzyme reaction signal in the control wells with no added compound. In this assay, the compounds of the invention exhibited an IC_{50} of less than 50 micromolar.

5 Example B

Cell Free Inhibition Assay Utilizing a Synthetic APP Substrate

A synthetic APP substrate that can be cleaved by beta-secretase and having N-terminal biotin and made fluorescent by the covalent attachment of Oregon green at the Cys residue is used
10 to assay beta-secretase activity in the presence or absence of the inhibitory compounds of the invention. Useful substrates include the following:

Biotin-SEVNL-DAEFRC[oregon green]KK [SEQ ID NO: 1]

15 Biotin-SEVKM-DAEFRC[oregon green]KK [SEQ ID NO: 2]

Biotin-GLNIKTEEISEISY-EVEFRC[oregon green]KK [SEQ ID NO: 3]

Biotin-ADRGLTTRPGSGLTNIKTEEISEVNL-DAEFRC[oregon green]KK [SEQ
ID NO:4]

Biotin-FVNQHLCoxGSHLVEALY-LVCoXGERGFFYTPKAC[oregon green]KK
20 [SEQ ID NO: 5]

The enzyme (0.1 nanomolar) and test compounds (0.001 - 100 micromolar) are incubated in pre-blocked, low affinity, black plates (384 well) at 37 degrees for 30 minutes. The reaction is initiated by addition of 150 millimolar substrate to a final
25 volume of 30 microliter per well. The final assay conditions are: 0.001 - 100 micromolar compound inhibitor; 0.1 molar sodium acetate (pH 4.5); 150 nanomolar substrate; 0.1 nanomolar soluble beta-secretase; 0.001% Tween 20, and 2% DMSO. The assay mixture is incubated for 3 hours at 37 degrees C, and the
30 reaction is terminated by the addition of a saturating concentration of immunopure streptavidin. After incubation with streptavidin at room temperature for 15 minutes, fluorescence polarization is measured, for example, using a LJL

Acquest (Ex485 nm/ Em530 nm). The activity of the beta-secretase enzyme is detected by changes in the fluorescence polarization that occur when the substrate is cleaved by the enzyme. Incubation in the presence or absence of compound
5 inhibitor demonstrates specific inhibition of beta-secretase enzymatic cleavage of its synthetic APP substrate. In this assay, preferred compounds of the invention exhibit an IC₅₀ of less than 50 micromolar.

10 **Example C**

Beta-Secretase Inhibition: P26-P4'SW Assay

Synthetic substrates containing the beta-secretase cleavage site of APP are used to assay beta-secretase activity, using the methods described, for example, in published PCT

15 application WO00/47618. The P26-P4'SW substrate is a peptide of the sequence:

(biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNLD AEF [SEQ ID NO: 6]

The P26-P1 standard has the sequence:

(biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNL [SEQ ID NO: 7].

20 Briefly, the biotin-coupled synthetic substrates are incubated at a concentration of from about 0 to about 200 micromolar in this assay. When testing inhibitory compounds, a substrate concentration of about 1.0 micromolar is preferred. Test compounds diluted in DMSO are added to the reaction mixture,
25 with a final DMSO concentration of 5%. Controls also contain a final DMSO concentration of 5%. The concentration of beta secretase enzyme in the reaction is varied, to give product concentrations with the linear range of the ELISA assay, about 125 to 2000 picomolar, after dilution.

30 The reaction mixture also includes 20 millimolar sodium acetate, pH 4.5, 0.06% Triton X100, and is incubated at 37 degrees C for about 1 to 3 hours. Samples are then diluted in assay buffer (for example, 145.4 nanomolar sodium chloride,

9.51 millimolar sodium phosphate, 7.7 millimolar sodium azide, 0.05% Triton X405, 6g/liter bovine serum albumin, pH 7.4) to quench the reaction, then diluted further for immunoassay of the cleavage products.

- 5 Cleavage products can be assayed by ELISA. Diluted samples and standards are incubated in assay plates coated with capture antibody, for example, SW192, for about 24 hours at 4 degrees C. After washing in TTBS buffer (150 millimolar sodium chloride, 25 millimolar Tris, 0.05% Tween 20, pH 7.5), the
- 10 samples are incubated with streptavidin-AP according to the manufacturer's instructions. After a one hour incubation at room temperature, the samples are washed in TTBS and incubated with fluorescent substrate solution A (31.2 g/liter 2-amino-2-methyl-1-propanol, 30 mg/liter, pH 9.5). Reaction with
- 15 streptavidin-alkaline phosphate permits detection by fluorescence. Compounds that are effective inhibitors of beta-secretase activity demonstrate reduced cleavage of the substrate as compared to a control.

20 **Example D**

Assays using Synthetic Oligopeptide-Substrates

- Synthetic oligopeptides are prepared that incorporate the known cleavage site of beta-secretase, and optionally detectable tags, such as fluorescent or chromogenic moieties. Examples of
- 25 such peptides, as well as their production and detection methods are described in U.S. Patent No: 5,942,400, herein incorporated by reference. Cleavage products can be detected using high performance liquid chromatography, or fluorescent or chromogenic detection methods appropriate to the peptide to be
- 30 detected, according to methods well known in the art. By way of example, one such peptide has the sequence SEVNLD-DAEF [SEQ ID NO: 8], and the cleavage site is between residues 5 and 6. Another preferred substrate has the sequence

ADRLTTRPGSGLTNIKTEEISEVNL-DAEF [SEQ ID NO: 9], and the cleavage site is between residues 26 and 27.

These synthetic APP substrates are incubated in the presence of beta-secretase under conditions sufficient to result in beta-secretase mediated cleavage of the substrate. Comparison of the cleavage results in the presence of the compound inhibitor to control results provides a measure of the compound's inhibitory activity.

10 Example E

Inhibition of Beta-Secretase Activity - Cellular Assay

An exemplary assay for the analysis of inhibition of beta-secretase activity utilizes the human embryonic kidney cell line HEKp293 (ATCC Accession No. CRL-1573) transfected with APP751 containing the naturally occurring double mutation Lys651Met52 to Asn651Leu652 (numbered for APP751), commonly called the Swedish mutation and shown to overproduce A beta (Citron et al., 1992, *Nature* 360:672-674), as described in U.S. Patent No. 5,604,102.

The cells are incubated in the presence/absence of the inhibitory compound (diluted in DMSO) at the desired concentration, generally up to 10 micrograms/ml. At the end of the treatment period, conditioned media is analyzed for beta-secretase activity, for example, by analysis of cleavage fragments. A beta can be analyzed by immunoassay, using specific detection antibodies. The enzymatic activity is measured in the presence and absence of the compound inhibitors to demonstrate specific inhibition of beta-secretase mediated cleavage of APP substrate.

30

Example F

Inhibition of Beta-Secretase in Animal Models of AD

Various animal models can be used to screen for inhibition of beta-secretase activity. Examples of animal models useful in the invention include, but are not limited to, mouse, guinea pig, dog, and the like. The animals used can be wild type, transgenic, or knockout models. In addition, mammalian models can express mutations in APP, such as APP695-SW and the like described herein. Examples of transgenic non-human mammalian models are described in U.S. Patent Nos. 5,604,102, 5,912,410 and 5,811,633.

- 10 PDAPP mice, prepared as described in Games et al., 1995, *Nature* 373:523-527 are useful to analyze *in vivo* suppression of A beta release in the presence of putative inhibitory compounds. As described in U.S. Patent No. 6,191,166, 4 month old PDAPP mice are administered compound formulated in vehicle, such as corn
15 oil. The mice are dosed with compound (1-30 mg/ml; preferably 1-10 mg/ml). After time, e.g., 3-10 hours, the animals are sacrificed, and brains removed for analysis.

Transgenic animals are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode
20 of administration. Control animals are untreated, treated with vehicle, or treated with an inactive compound. Administration can be acute, i.e., single dose or multiple doses in one day, or can be chronic, i.e., dosing is repeated daily for a period of days. Beginning at time 0, brain tissue or cerebral fluid
25 is obtained from selected animals and analyzed for the presence of APP cleavage peptides, including A beta, for example, by immunoassay using specific antibodies for A beta detection. At the end of the test period, animals are sacrificed and brain tissue or cerebral fluid is analyzed for the presence of A beta
30 and/or beta-amyloid plaques. The tissue is also analyzed for necrosis.

Animals administered the compound inhibitors of the invention are expected to demonstrate reduced A beta in brain

tissues or cerebral fluids and reduced beta amyloid plaques in brain tissue, as compared with non-treated controls.

Example G

5 Inhibition of A Beta Production in Human Patients

Patients suffering from Alzheimer's Disease (AD) demonstrate an increased amount of A beta in the brain. AD patients are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration.

10 Administration is repeated daily for the duration of the test period. Beginning on day 0, cognitive and memory tests are performed, for example, once per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as
15 analyzed by changes in one or more of the following disease parameters: A beta present in CSF or plasma; brain or hippocampal volume; A beta deposits in the brain; amyloid plaque in the brain; and scores for cognitive and memory function, as compared with control, non-treated patients.

20

Example H

Prevention of A Beta Production in Patients at Risk for AD

Patients predisposed or at risk for developing AD are identified either by recognition of a familial inheritance
25 pattern, for example, presence of the Swedish Mutation, and/or by monitoring diagnostic parameters. Patients identified as predisposed or at risk for developing AD are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Administration
30 is repeated daily for the duration of the test period.


Beginning on day 0, cognitive and memory tests are performed, for example, once per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as analyzed by changes in one or more of the following disease parameters: A beta present in CSF or plasma; brain or
5 hippocampal volume; amyloid plaque in the brain; and scores for cognitive and memory function, as compared with control, non-treated patients.

While this invention has been described with respect to
10 various specific examples and embodiments, it is to be understood that the invention is not limited thereby and should only be construed by interpretation of the scope of the appended claims.

What is claimed is:

1. A compound of the formula:



5

or a pharmaceutically acceptable salt thereof,

where R_1 is:

- (I) C_1 - C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, C_1 - C_7 alkyl (optionally substituted with C_1 - C_3 alkyl and C_1 - C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$, C_1 - C_3 alkoxy, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl, -OC=O NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

15

(II) -CH $_2$ -S(O) $_{0-2}$ -(C_1 - C_6 alkyl),

(III) -CH $_2$ -CH $_2$ -S(O) $_{0-2}$ -(C_1 - C_6 alkyl),

- (IV) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C $\equiv$ N, -CF $_3$, C_1 - C_3 alkoxy, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl,

20

- (V) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C $\equiv$ N, -CF $_3$, C_1 - C_3 alkoxy, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl,

25

- (VI) -(CH $_2$) n_1 -(R $_{1-aryl}$) where n_1 is zero or one and where R $_{1-aryl}$ is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthyl, tetralinyl optionally substituted with one,

two, three or four of the following substituents on the aryl ring:

(A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C≡N, -CF₃, C₁-C₃ alkoxy,

(B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(D) -F, Cl, -Br and -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(N) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$,

(VII) $-(\text{CH}_2)_{n_1}-(\text{R}_1\text{-heteroaryl})$ where n_1 is as defined above

and where $\text{R}_1\text{-heteroaryl}$ is selected from the group consisting of
 pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl,
 5 indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl,
 quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl,
 isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indoliziny, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl,
 10 triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, isochroman, chromanyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuran, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl,
 15 pyridopyridinyl, benzotetrahydrofuran, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoquinazolinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisoquinazolinyl, benzopyranyl, benzothiopyranyl,
 20 coumarinyl, isocoumarinyl, chromon, chromanonyl, pyridinyl-N-oxide tetrahydroquinolinyl dihydroquinolinyl dihydroquinolinonyl dihydroisoquinolinonyl dihydrocoumarinyl dihydroisocoumarinyl isoindolinonyl benzodioxan, benzoxazolinonyl pyrrolyl N-oxide, pyrimidinyl N-oxide,
 25 pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indoliziny, indazolyl N-oxide,
 30 benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, and benzothiopyranyl S,S-dioxide, where the $\text{R}_1\text{-heteroaryl}$ group is

bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent R_N -heteroaryl group substituted by hydrogen such that the new bond to the R_1 -heteroaryl group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or
 5 four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy,
 10

(2) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 15

(3) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 20

(4) $-F$, $-Cl$, $-Br$ and $-I$,

(6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three $-F$,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined
 25 below,

(8) $-OH$,

(9) $-C\equiv N$,

(10) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
 30

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-\text{SO}_2-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(13) $-\text{CO}-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (14) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$, with the proviso that when n_1 is zero $\text{R}_{1-\text{heteroaryl}}$ is not bonded to the carbon chain by nitrogen,

(VIII) $-(\text{CH}_2)_{n_1}-(\text{R}_{1-\text{heterocycle}})$ where n_1 is as defined above and $\text{R}_{1-\text{heterocycle}}$ is selected from the group consisting of
 10 morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-
 15 dioxide, oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl, dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide, and homothiomorpholinyl S-oxide where the $\text{R}_{1-\text{heterocycle}}$ group is bonded by any atom of the parent
 20 $\text{R}_{1-\text{heterocycle}}$ group substituted by hydrogen such that the new bond to the $\text{R}_{1-\text{heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) C_1-C_6 alkyl optionally substituted with
 25 one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy,

(2) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three
 30 substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or C_1-C_6 alkyl,

(3) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are
 5 -H or C_1-C_6 alkyl,

(4) -F, -Cl, -Br and -I,

(5) C_1-C_6 alkoxy,

(6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three -F,

10 (7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,

(9) $-C\equiv N$,

(10) C_3-C_7 cycloalkyl, optionally
 15 substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH
 $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

20 (12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$,

25 (15) =O, with the proviso that when n_1 is zero $R_{1-heterocycle}$ is not bonded to the carbon chain by nitrogen;

where R_2 is selected from the group consisting of:

(I) -H,

(II) C_1-C_6 alkyl, optionally substituted with one, two
 30 or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

5 (IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

15 (VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

20 where R₃ is selected from the group consisting of:

(I) -H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,

25 -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

(IV) C₂-C₆ alkenyl with one or two double bonds,

30 (V) C₂-C₆ alkynyl with one or two triple bonds,

(VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, and seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, $-NR_{N-2}-$, where R_{N-2} is as defined below;

10 where R_N is:

(I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:

- (A) $-CO-$,
- (B) $-SO_2-$,
- 15 (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,
- (D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above,
- 20 (E) a single bond;

where R_{N-1} is selected from the group consisting of:

- (A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, dihydronaphthyl or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, optionally substituted
- 25 with one, two or three of the following substituents which can be the same or different and are:
 - (1) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$,
 - 30 $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
 - (2) $-OH$,

- (3) $-\text{NO}_2$,
- (4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
- (5) $-\text{CO}-\text{OH}$,
- (6) $-\text{C}\equiv\text{N}$,
- 5 (7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:
- (a) $-\text{H}$,
- (b) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted
- 10 with one substituent selected from the group consisting of:
- (i) $-\text{OH}$,
- (ii) $-\text{NH}_2$,
- (c) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one to three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
- 15 (d) $-\text{C}_3-\text{C}_7$ cycloalkyl,
- (e) $-(\text{C}_1-\text{C}_2 \text{ alkyl})-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
- (f) $-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_3 \text{ alkyl})$,
- (g) $-\text{C}_2-\text{C}_6$ alkenyl with one or two double bonds,
- 20 (h) $-\text{C}_2-\text{C}_6$ alkynyl with one or two triple bonds,
- (i) $-\text{C}_1-\text{C}_6$ alkyl chain with one double bond and one triple bond,
- (j) $-\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined
- 25 above,
- (k) $-\text{R}_1\text{-heteroaryl}$ where $\text{R}_1\text{-heteroaryl}$ is as defined above,
- (8) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
- (9) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl with one, two or three double bonds})$,
- 30 (10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl with one, two or three triple bonds})$,
- (11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(12) $-(CH_2)_{0-4}-CO-R_1\text{-aryl}$ where $R_1\text{-aryl}$ is as defined above,

(13) $-(CH_2)_{0-4}-CO-R_1\text{-heteroaryl}$ where $R_1\text{-heteroaryl}$ is as defined above,

5 (14) $-(CH_2)_{0-4}-CO-R_1\text{-heterocycle}$ where $R_1\text{-heterocycle}$ is as defined above,

(15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, 10 homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:

15 (a) C_1-C_6 alkyl,

(b) $-(CH_2)_{0-2}-(R_1\text{-aryl})$ where $R_1\text{-aryl}$ is as defined above,

(c) C_2-C_6 alkenyl containing one or two double bonds,

20 (d) C_2-C_6 alkynyl containing one or two triple bonds,

(e) C_3-C_7 cycloalkyl,

(f) $-(CH_2)_{0-2}-(R_1\text{-heteroaryl})$ where $R_1\text{-heteroaryl}$ is as defined above,

25 (17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

30 (21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

- (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,
- 5 (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- 10 (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is $-\text{H}$ or C_1-C_4 alkyl,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 15 (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 20 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
- 25 (36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$
- (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$
- 30

(38) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-SO_2-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as described above,

(39) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

- 5 (B) $-R_{N-\text{heteroaryl}}$ where $R_{N-\text{heteroaryl}}$ is selected from the group consisting of pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, 10 thiazolyl, indoliziny, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, 15 isochromanly, chromanly, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranly, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranly, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, hexoxazinyl, phenothiazinyl, 20 pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl, benzopyranly, benzothiopyranly, coumarinyl, isocoumarinyl, chromonyl, chromanonyl, pyridinyl-N-oxide, tetrahydroquinolinyl dihydroquinolinyl 25 dihydroquinolinonyl dihydroisoquinolinonyl dihydrocoumarinyl dihydroisocoumarinyl isoindolinonyl benzodioxany, benzoxazolinonyl pyrrolyl N-oxide, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, 30 quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indoliziny, indazolyl N-oxide, benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-

oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, and benzothiopyranyl S,S-dioxide, where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by
 5 hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

(1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group
 10 consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $C_1\text{-}C_3$ alkoxy, $\text{-NR}_{1\text{-}a}\text{R}_{1\text{-}b}$ where $R_{1\text{-}a}$ and $R_{1\text{-}b}$ are as defined above,

(2) -OH,
 (3) -NO_2 ,
 15 (4) -F, -Cl, -Br, -I,
 (5) -CO-OH ,
 (6) $\text{-C}\equiv\text{N}$,
 (7) $\text{-(CH}_2\text{)}_{0\text{-}4}\text{-CO-NR}_{N\text{-}2}\text{R}_{N\text{-}3}$ where $R_{N\text{-}2}$ and $R_{N\text{-}3}$ are the same or different and are selected from the group
 20 consisting of:

(a) -H,
 (b) $\text{-C}_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 (i) -OH,
 25 (ii) -NH_2 ,
 (c) $\text{-C}_1\text{-}C_6$ alkyl optionally substituted with one to three -F, -Cl, -Br, -I,
 (d) $\text{-C}_3\text{-}C_7$ cycloalkyl,
 (e) $\text{-(C}_1\text{-}C_2\text{ alkyl)-(C}_3\text{-}C_7\text{ cycloalkyl)}$,
 30 (f) $\text{-(C}_1\text{-}C_6\text{ alkyl)-O-(C}_1\text{-}C_3\text{ alkyl)}$,
 (g) $\text{-C}_2\text{-}C_6$ alkenyl with one or two double bonds,

(h) $-C_2-C_6$ alkynyl with one or two triple bonds,

(i) $-C_1-C_6$ alkyl chain with one double bond and one triple bond,

5 (j) $-R_1\text{-aryl}$ where $R_1\text{-aryl}$ is as defined above,

(k) $-R_1\text{-heteroaryl}$ where $R_1\text{-heteroaryl}$ is as defined above,

(8) $-(CH_2)_{0-4}\text{-CO-}(C_1-C_{12}\text{ alkyl})$,

10 (9) $-(CH_2)_{0-4}\text{-CO-}(C_2-C_{12}\text{ alkenyl with one, two or three double bonds})$,

(10) $-(CH_2)_{0-4}\text{-CO-}(C_2-C_{12}\text{ alkynyl with one, two or three triple bonds})$,

(11) $-(CH_2)_{0-4}\text{-CO-}(C_3-C_7\text{ cycloalkyl})$,

15 (12) $-(CH_2)_{0-4}\text{-CO-}R_1\text{-aryl}$ where $R_1\text{-aryl}$ is as defined above,

(13) $-(CH_2)_{0-4}\text{-CO-}R_1\text{-heteroaryl}$ where $R_1\text{-heteroaryl}$ is as defined above,

20 (14) $-(CH_2)_{0-4}\text{-CO-}R_1\text{-heterocycle}$ where $R_1\text{-heterocycle}$ is as defined above,

(15) $-(CH_2)_{0-4}\text{-CO-}R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, 25 pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}\text{-CO-O-}R_{N-5}$ where R_{N-5} is selected from the group consisting of:

(a) C_1-C_6 alkyl,

30 (b) $-(CH_2)_{0-2}\text{-}(R_1\text{-aryl})$ where $R_1\text{-aryl}$ is as defined above,

(c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple bonds,

(e) C_3-C_7 cycloalkyl,

(f) $-(CH_2)_{0-2}-(R_1\text{-heteroaryl})$ where $R_1\text{-heteroaryl}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

(21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N\text{-aryl-1}})_2$ where $R_{N\text{-aryl-1}}$ is $-H$ or $C_1-C_4 \text{ alkyl}$,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,

(32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,

(33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$

5 (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

10 (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

(38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above,

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

20 (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-heterocycle}}$ is defined as $\text{R}_{1-\text{heterocycle}}$ is defined above,

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-1-heterocycle}}$ is defined as $\text{R}_{1-\text{heterocycle}}$ is defined above,

(I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}},$

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

where W is

30 (6) $-(\text{CH}_2)_{0-4}-,$

(7) $-\text{O}-,$

(8) $-\text{S}(\text{O})_{0-2}-,$

(9) $-N(R_{N-5})-$ where R_{N-5} is as defined above, or

(10) $-CO-$;

(II) $-CO-(C_1-C_{10} \text{ alkyl})$ where alkyl is optionally substituted with one two or three substitutents selected from the group consisting of:

(A) $-OH$,

(B) $-C_1-C_6$ alkoxy,

(C) $-C_1-C_6$ thioalkoxy,

10 (D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, C_1-C_6 alkyl or -phenyl,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

15 (G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

20 (K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

25 (N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, $-I$),

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,

30 (R) $-F$, $-Cl$,

(III) $-CO-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substitutents selected from the group consisting of :

- (A) -OH,
- (B) -C₁-C₆ alkoxy,
- (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -
- 5 phenyl,
- (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
- (F) -CO-R_{N-4} where R_{N-4} is as defined above,
- (G) -SO₂-(C₁-C₈ alkyl),
- 10 (H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
- (I) -NH-CO-(C₁-C₆ alkyl),
- (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
- (K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
- 15 different and are as defined above,
- (L) -R_{N-4} where R_{N-4} is as defined above,
- (M) -O-CO-(C₁-C₆ alkyl),
- (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different and are as defined above,
- 20 (O) -O-(C₁-C₅ alkyl)-COOH,
- (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, -I),
- (Q) -NH-SO₂-(C₁-C₆ alkyl),
- (R) -F, -Cl,
- 25 (IV) -CO-(C₁-C₆ alkyl)-S-(C₁-C₆ alkyl) where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of :
- (A) -OH,
- (B) -C₁-C₆ alkoxy,
- 30 (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
- (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

- (F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
 (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same
 or different and are as defined above,
 5 (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
 (K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or
 different and are as defined above,
 (L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 10 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where the $\text{R}_{\text{N}-8}$ s are the same or
 different and are as defined above,
 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with
 15 one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (R) $-\text{F}$, $-\text{Cl}$,
 (V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$
 where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is
 20 selected from the group consisting of:
 (A) $-\text{H}$,
 (B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,
 (C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
 (D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,
 25 (E) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one triple bond,
 (F) $\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above,
 (G) $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined
 above,
 (VI) $-\text{CO}-(\text{C}_3-\text{C}_8 \text{ cycloalkyl})$ where cycloalkyl is
 30 optionally substituted with one or two substituents selected
 from the group consisting of:
 (A) $-(\text{CH}_2)_{0-4}-\text{OH}$,

- (B) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6$ alkoxy,
 (C) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6$ thioalkoxy,
 (D) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,
 5 (E) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
 (F) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 (G) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
 10 (H) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
 (I) $-(\text{CH}_2)_{0-4}-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
 (K) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the
 15 same or different and are as defined above,
 (L) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where the $\text{R}_{\text{N}-8}$ s are the same or different and are as defined above,
 20 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (R) $-\text{F}$, $-\text{Cl}$,

25 where R_c is:

- (I) $-\text{C}_1-\text{C}_{10}$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}-\text{phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{OC}=\text{O NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{S}(=\text{O})_{0-2} \text{R}_{1-a}$ where R_{1-a} is as defined above, -

$\text{NR}_{1-a}\text{C}=\text{O}$ $\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-\text{C}=\text{O}$ $\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, and $-\text{S}(=\text{O})_2$ $\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{CO}-\text{OH}$, $-\text{CO}-\text{O}-(\text{C}_1-\text{C}_4 \text{ alkyl})$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(III) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are $-\text{H}$, C_1-C_4 alkyl optionally substituted with one or two $-\text{OH}$,

C_1-C_4 alkoxy optionally substituted with one, two, or three of $-\text{F}$,

$-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, C_2-C_6 alkenyl containing one or two double bonds, C_2-C_6 alkynyl containing one or two triple bonds, phenyl,

and where $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six and seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{NR}_{\text{N-2}}-$ and $\text{R}_{\text{C-aryl}}$ is defined as $\text{R}_{\text{N-aryl}}$ is defined above;

(IV) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$ is defined as $\text{R}_{\text{N-heteroaryl}}$ is defined above and $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(V) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}-\text{R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(VI) $-(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}-\text{R}_{\text{C-aryl}}-\text{R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(VII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(VIII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

5 (IX) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-aryl}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above, and $\text{R}_{\text{C-heterocycle}}$ is defined as $\text{R}_{\text{N-heterocycle}}$ is defined above,

(X) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heteroaryl}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

10 (XI) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-aryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(XII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-heteroaryl}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(XIII) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(XIV) - $(\text{CR}_{\text{C-x}}\text{R}_{\text{C-y}})_{0-4}\text{-R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-heterocycle}}$, $\text{R}_{\text{C-x}}$ and $\text{R}_{\text{C-y}}$ are as defined above,

(XV) - $[\text{C}(\text{R}_{\text{C-1}})(\text{R}_{\text{C-2}})]_{1-3}\text{-CO-N}(\text{R}_{\text{C-3}})_2$ where $\text{R}_{\text{C-1}}$ and $\text{R}_{\text{C-2}}$ are the same or different and are selected from the group

20 consisting of:

(A) -H,

(B) $\text{-C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH,

25 -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br,

30 -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents

selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_6 alkoxy, -O-phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{1-2}-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

5 (F) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_6 alkoxy, -O-phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

10 (G) $-(C_1-C_4 \text{ alkyl})-R_{C'}\text{-aryl}$ where $R_{C'}\text{-aryl}$ is as defined for R_{1-aryl} ,

(H) $-(C_1-C_4 \text{ alkyl})-R_{C'}\text{-heteroaryl}$ where $R_{C'}\text{-heteroaryl}$ is as defined above,

15 (I) $-(C_1-C_4 \text{ alkyl})-R_{C'}\text{-heterocycle}$ where $R_{C'}\text{-heterocycle}$ is as defined above,

(J) $-R_{C'}\text{-heteroaryl}$ where $R_{C'}\text{-heteroaryl}$ is as defined above,

(K) $-R_{C'}\text{-heterocycle}$ where $R_{C'}\text{-heterocycle}$ is as defined above,

20 (M) $-(CH_2)_{1-4}-R_{C-4}-(CH_2)_{0-4}-R_{C'}\text{-aryl}$ where R_{C-4} is -O-, -S- or $-NR_{C-5}-$ where R_{C-5} is C_1 - C_6 alkyl, and where $R_{C'}\text{-aryl}$ is defined above,

25 (N) $-(CH_2)_{1-4}-R_{C-4}-(CH_2)_{0-4}-R_{C'}\text{-heteroaryl}$ where R_{C-4} and $R_{C'}\text{-heteroaryl}$ are as defined above,

(O) $-R_{C'}\text{-aryl}$ where $R_{C'}\text{-aryl}$ is as defined above, and where R_{C-3} is the same or different and is:

(A) -H,

30 (B) $-C_1-C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents
5 selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents
10 selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from
15 the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) -R_{C'}-aryl where R_{C'}-aryl is as defined above,

20 (G) -R_C-heteroaryl where R_C-heteroaryl is as defined above,

(H) -R_C-heterocycle where R_C-heterocycle is as defined above,

(I) -(C₁-C₄ alkyl)-R_{C'}-aryl where R_{C'}-aryl is as defined above,

25 (J) -(C₁-C₄ alkyl)-R_C-heteroaryl where R_C-heteroaryl is as defined above,

(K) -(C₁-C₄ alkyl)-R_C-heterocycle where R_C-heterocycle is as defined above,

(XVI) -CH(R_C-aryl)₂ where R_C-aryl are the same or
30 different and are as defined above,

(XVII) -CH(R_C-heteroaryl)₂ where R_C-heteroaryl are the same or different and are as defined above,

(XVIII) $-\text{CH}(\text{R}_{\text{C-aryl}})(\text{R}_{\text{C-heteroaryl}})$ where $\text{R}_{\text{C-aryl}}$ and $\text{R}_{\text{C-heteroaryl}}$ are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $\text{R}_{\text{C-aryl}}$ or $\text{R}_{\text{C-heteroaryl}}$ or $\text{R}_{\text{C-heterocycle}}$ where $\text{R}_{\text{C-aryl}}$ or $\text{R}_{\text{C-heteroaryl}}$ or $\text{R}_{\text{C-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH , $\text{NR}_{\text{N-S}}$, O , $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $=\text{O}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) $\text{C}_2\text{-C}_{10}$ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $\text{C}_2\text{-C}_{10}$ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}\text{-CHR}_{\text{C-6}}-(\text{CH}_2)_{0-1}\text{-R}_{\text{C-aryl}}$ where $\text{R}_{\text{C-aryl}}$ is as defined above and $\text{R}_{\text{C-6}}$ is $-(\text{CH}_2)_{0-6}\text{-OH}$,

(XXII) $-(\text{CH}_2)_{0-1}\text{-CHR}_{\text{C-6}}-(\text{CH}_2)_{0-1}\text{-R}_{\text{C-heteroaryl}}$ where $\text{R}_{\text{C-heteroaryl}}$ and $\text{R}_{\text{C-6}}$ is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{C-aryl}} \text{ or } \text{R}_{\text{C-heteroaryl}})-\text{CO-O}(\text{C}_1\text{-C}_4 \text{ alkyl})$ where $\text{R}_{\text{C-aryl}}$ and $\text{R}_{\text{C-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2\text{-OH})-\text{CH}(-\text{OH})\text{-micro-NO}_2$,


(XXV) $(\text{C}_1\text{-C}_6 \text{ alkyl})\text{-O-}(\text{C}_1\text{-C}_6 \text{ alkyl})\text{-OH}$,

(XXVII) $-\text{CH}_2\text{-NH-CH}_2\text{-CH}(-\text{O-CH}_2\text{-CH}_3)_2$,

(XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}\text{-C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; and

where R_4 is:



where x is 0 or 1 and R_5 is H or is:

5 (I) R_{p-1} wherein R_{p-1} is selected from the group consisting of:

(A) $-Y-R_{p-aryl}$ where Y is absent or is oxygen, and R_{p-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are independently $-H$ or C_1-C_6 alkyl,

(2) $-OH$,

(3) $-NO_2$,

(4) $-F$, $-Cl$, $-Br$, $-I$,

20 (5) $-CO-OH$,

(6) $-C\equiv N$,

(7) $-(CH_2)_{0-4}-CO-NR_{p-2}R_{p-3}$ where R_{p-2} and R_{p-3} are the same or different and are selected from the group consisting of:

25 (a) $-H$,

(b) $-C_1-C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-OH$,

(ii) $-NH_2$,

30 (c) $-C_1-C_6$ alkyl optionally substituted with one to three $-F$, $-Cl$, $-Br$, $-I$,

- (d) $-C_3-C_7$ cycloalkyl,
- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
- (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
- (g) $-C_2-C_6$ alkenyl with one or two
- 5 double bonds,
- (h) $-C_2-C_6$ alkynyl with one or two
- triple bonds,
- (i) $-C_1-C_6$ alkyl chain with one double
- bond and one triple bond,
- 10 (j) $-R_{P\text{-aryl}}$ where $-R_{P\text{-aryl}}$ is defined
- above,
- (k) $-R_{P\text{-heteroaryl}}$ where $R_{P\text{-heteroaryl}}$ is
- defined below,
- (1) $-C(O)C_1-C_6$ alkyl,
- 15 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
- (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12})$ alkenyl with one,
- two or three double bonds,
- (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12})$ alkynyl with one,
- two or three triple bonds,
- 20 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
- (12) $-(CH_2)_{0-4}-CO-R_{P\text{-aryl}}$ where $R_{P\text{-aryl}}$ is as
- defined above,
- (13) $-(CH_2)_{0-4}-CO-R_{P\text{-heteroaryl}}$ where $R_{P\text{-heteroaryl}}$
- is as defined below,
- 25 (14) $-(CH_2)_{0-4}-CO-R_{P\text{-heterocycle}}$ where $R_{P\text{-heterocycle}}$
- is selected from the group consisting of:
- (A) morpholinyl,
- (B) thiomorpholinyl,
- (C) thiomorpholinyl S-oxide,
- 30 (D) thiomorpholinyl S,S-dioxide,
- (E) piperazinyl,
- (F) homopiperazinyl,
- (G) pyrrolidinyl,

- (H) pyrrolinyl,
- (I) tetrahydropyranyl,
- (J) piperidinyl,
- (K) tetrahydrofuranyl,
- (L) tetrahythiophenyl,
- (M) homopiperidinyl,
- (N) imidazolidine,
- (O) imidazolidine dione, and
- (P) dithiane,

5

where the $R_{P\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{P\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{P\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one thru four:

15

(1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -O-CO-($C_1\text{-}C_6$ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C≡N, -CF₃, $C_1\text{-}C_3$ alkoxy,

20

(2) $C_2\text{-}C_6$ alkenyl with one or two double bonds,

(3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds,

(4) -F, Cl, -Br and -I,

25

(5) $C_1\text{-}C_6$ alkoxy,

(6) -O- $C_1\text{-}C_6$ alkyl optionally substituted with one thru three -F,

(7) -NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are as defined above,

30

(8) -OH,

(9) -C≡N,

(10) $C_3\text{-}C_7$ cycloalkyl,

(11) -CO-($C_1\text{-}C_4$ alkyl),

- (12) $-\text{SO}_2-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are
as defined above,
- (13) $-\text{CO}-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are
as defined above,
- 5 (14) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$;
(15) $=\text{O}$; or
(16) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$;
(15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{N-4}$ where R_{N-4} is selected
from the group consisting of morpholinyl, thiomorpholinyl,
10 piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl,
homomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide,
pyrrolinyl and pyrrolidinyl where each group is optionally
substituted with one, two, three, or four of: $\text{C}_1-\text{C}_6 \text{ alkyl}$,
(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{N-5}$ where R_{N-5}
15 is selected from the group consisting of:
(a) $\text{C}_1-\text{C}_6 \text{ alkyl}$,
(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{P-\text{aryl}})$ where $\text{R}_{P-\text{aryl}}$ is as
defined above,
(c) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ containing one or two
20 double bonds,
(d) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ containing one or two
triple bonds,
(e) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
(f) $-(\text{CH}_2)_{0-2}-(\text{R}_{P-\text{heteroaryl}})$ where $\text{R}_{P-\text{heteroaryl}}$
25 heteroaryl is as defined below,
(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{P-2}\text{R}_{P-3}$ where R_{P-2} and R_{P-3}
are as defined above,
(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
30 (20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{N-5})-\text{CO}-\text{O}-\text{R}_{N-5}$ where
 R_{N-5} can be the same or different and is as defined above,

- (22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,
- (23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,
- 5 (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{P-2}$ where R_{N-5} and R_{P-2} can be the same or different and are as defined above,
- (25) $-(CH_2)_{0-4}-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} can be the same or different and are as defined above,
- (26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined
- 10 above,
- (27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,
- (28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is $-H$ or C_1-C_4 alkyl,
- (29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as
- 15 defined above,
- (30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,
- (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,
- 20 (32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,
- (33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined above,
- (34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally$
- 25 substituted with one, two, three, four, or five of: $-F$),
- (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b}
- 30 are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$,

-OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) - (CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{P-2} where R_{N-5} and R_{P-2} can be the same or different and are as described
5 above,

(39) - (CH₂)₀₋₄- C₃-C₇ cycloalkyl,

(B) -Y-R_{P-heteroaryl} where Y is absent or is oxygen, and R_{P-heteroaryl} is selected from the group consisting of:

10	(A) pyridinyl,
	(B) pyrimidinyl,
	(C) quinolinyl,
	(F) benzothienyl,
	(G) indolyl,
15	(H) indolinyl,
	(I) pyridazinyl,
	(J) pyrazinyl,
	(K) isoindolyl,
	(L) isoquinolyl,
20	(M) quinazolinyl,
	(N) quinoxalinyl,
	(O) phthalazinyl,
	(P) imidazolyl,
	(Q) isoxazolyl,
25	(R) pyrazolyl,
	(S) oxazolyl,
	(T) thiazolyl,
	(U) indolizinyl,
	(V) indazolyl,
30	(W) benzothiazolyl,
	(X) benzimidazolyl,
	(Y) benzofuranyl,
	(Z) furanyl,

	(AA) thienyl,
	(BB) pyrrolyl,
	(CC) oxadiazolyl,
	(DD) thiadiazolyl,
5	(EE) triazolyl,
	(FF) tetrazolyl,
	(II) oxazolopyridinyl,
	(JJ) imidazopyridinyl,
	(KK) isothiazolyl,
10	(LL) naphthyridinyl,
	(MM) cinnolinyl,
	(NN) carbazolyl,
	(OO) beta-carbolinyl,
	(PP) isochromanyl,
15	(QQ) chromanyl,
	(SS) tetrahydroisoquinolinyl,
	(TT) isoindolinyl,
	(UU) isobenzotetrahydrofuranyl,
	(VV) isobenzotetrahydrothienyl,
20	(WW) isobenzothienyl,
	(XX) benzoxazolyl,
	(YY) pyridopyridinyl,
	(ZZ) benzotetrahydrofuranyl,
	(AAA) benzotetrahydrothienyl,
25	(BBB) purinyl,
	(CCC) benzodioxolyl,
	(DDD) triazinyl,
	(EEE) phenoxazinyl,
	(FFF) phenothiazinyl,
30	(GGG) pteridinyl,
	(HHH) benzothiazolyl,
	(III) imidazopyridinyl,
	(JJJ) imidazothiazolyl,

- 5 (KKK) dihydrobenzisoxaziny1,
 (LLL) benzisoxaziny1,
 (MMM) benzoxaziny1,
 (NNN) dihydrobenzisothiaziny1,
 (OOO) benzopyrany1,
 (PPP) benzothiopyrany1,
 (QQQ) coumariny1,
 (RRR) isocoumariny1,
 (SSS) chromony1,
 10 (TTT) chromanony1,
 (UUU) pyridiny1-N-oxide,

where the $R_{P\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{P\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{P\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted
 15 with one, two, three, or four of:

(1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined
 20 above,

(2) $-OH$,
 (3) $-NO_2$,
 (4) $-F$, $-Cl$, $-Br$, $-I$,
 25 (5) $-CO-OH$,
 (6) $-C\equiv N$,
 (7) $-(CH_2)_{0-4}-CO-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the same or different and are selected from the group consisting of:

30 (a) $-H$,
 (b) $-C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-OH$,

- (ii) -NH_2 ,
- (c) $\text{-C}_1\text{-C}_6$ alkyl optionally substituted with one to three -F , -Cl , -Br , -I ,
- (d) $\text{-C}_3\text{-C}_7$ cycloalkyl,
- 5 (e) $\text{-(C}_1\text{-C}_2 \text{ alkyl)- (C}_3\text{-C}_7 \text{ cycloalkyl)}$,
- (f) $\text{-(C}_1\text{-C}_6 \text{ alkyl)-O-(C}_1\text{-C}_3 \text{ alkyl)}$,
- (g) $\text{-C}_2\text{-C}_6$ alkenyl with one or two double bonds,
- (h) $\text{-C}_2\text{-C}_6$ alkynyl with one or two
- 10 triple bonds,
- (i) $\text{-C}_1\text{-C}_6$ alkyl chain with one double bond and one triple bond,
- (j) $\text{-R}_\text{p-aryl}$ where $\text{R}_\text{p-aryl}$ is as defined above,
- 15 (k) $\text{-R}_\text{p-heteroaryl}$ where $\text{R}_\text{p-heteroaryl}$ is as defined above,
- (8) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_1\text{-C}_{12} \text{ alkyl)}$,
- (9) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_2\text{-C}_{12} \text{ alkenyl with one, two or three double bonds)}$,
- 20 (10) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_2\text{-C}_{12} \text{ alkynyl with one, two or three triple bonds)}$,
- (11) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-(C}_3\text{-C}_7 \text{ cycloalkyl)}$,
- (12) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{p-aryl}$ where $\text{R}_\text{p-aryl}$ is as defined above,
- 25 (13) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{p-heteroaryl}$ where $\text{R}_\text{p-heteroaryl}$ is as defined above,
- (14) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_\text{p-heterocycle}$ where $\text{R}_\text{p-heterocycle}$ is as defined above,
- (15) $\text{-(CH}_2\text{)}_{0-4}\text{-CO-R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is selected
- 30 from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally

substituted with one, two, three, or four of: C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5}

is selected from the group consisting of:

(a) C_1-C_6 alkyl,

5 (b) $-(CH_2)_{0-2}-(R_{P-aryl})$ where R_{P-aryl} is as defined above,

(c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two
10 triple bonds,

(e) C_3-C_7 cycloalkyl,

(f) $-(CH_2)_{0-2}-(R_{P-heteroaryl})$ where $R_{P-heteroaryl}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3}
15 are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

(21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where
20 R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where
 R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be
the same or different and is as defined above,

25 (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{P-2}$ where R_{N-5}
and R_{P-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} can
be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined
30 above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$
is $-H$ or C_1-C_4 alkyl,

- (29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,
- (30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,
- 5 (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,
- (32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,
- 10 (33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined above,
- (34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, three, four, or five of: } -F),$
- (35) $C_3-C_7 \text{ cycloalkyl},$
- (36) $C_2-C_6 \text{ alkenyl with one or two double}$
- 15 $\text{bonds optionally substituted with } C_1-C_3 \text{ alkyl, } -F, -Cl, -Br, -I, -OH, -SH, -C\equiv N, -CF_3, C_1-C_3 \text{ alkoxy, } -NR_{1-a}R_{1-b} \text{ where } R_{1-a} \text{ and } R_{1-b} \text{ are as defined above,}$
- (37) $C_2-C_6 \text{ alkynyl with one or two triple bonds optionally substituted with } C_1-C_3 \text{ alkyl, } -F, -Cl, -Br, -I, -OH, -SH, -C\equiv N, -CF_3, C_1-C_3 \text{ alkoxy, } -NR_{1-a}R_{1-b} \text{ where } R_{1-a} \text{ and } R_{1-b} \text{ are as defined above,}$
- 20 (38) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-SO_2-R_{P-2}$ where R_{N-5} and R_{P-2} can be the same or different and are as described above,
- 25 (39) $-(CH_2)_{0-4}-C_3-C_7 \text{ cycloalkyl},$
- (C) $R_{P-aryl}-W-R_{P-aryl},$
- (D) $R_{P-aryl}-W-R_{P-heteroaryl},$
- (E) $R_{P-aryl}-W-R_{P-heterocycle},$
- (F) $R_{P-heteroaryl}-W-R_{P-aryl},$
- 30 (G) $R_{P-heteroaryl}-W-R_{P-heteroaryl},$
- (H) $R_{P-heteroaryl}-W-R_{P-heterocycle},$
- (I) $R_{P-heterocycle}-W-R_{P-aryl},$
- (J) $R_{P-heterocycle}-W-R_{P-heteroaryl},$

(K) $R_{P\text{-heterocycle}}\text{-W-}R_{P\text{-heterocycle}}$,

where W is

(1) $-(\text{CH}_2)_{0-4}-$,

(2) $-\text{O}-$,

5 (3) $-\text{S}(\text{O})_{0-2}-$,

(4) $-\text{N}(\text{R}_{N-5})-$ where R_{N-5} is as defined

above;

(5) $-\text{CO}-$; or

(6) $-\text{O}-(\text{CH}_2)_{1-4}-$,

10 (L) $-\text{Y-}R_{P\text{-heterocycle}}$ where Y is absent or is oxygen,
and $R_{P\text{-heterocycle}}$ is defined above;

(II) $-(\text{C}_1\text{-C}_{10}\text{ alkyl})$ where alkyl is optionally
substituted with one, two, or three substituents selected from
the group consisting of:

15 (A) $-\text{OH}$,

(B) $-\text{C}_1\text{-C}_6\text{ alkoxy}$,

(C) $-\text{C}_1\text{-C}_6\text{ thioalkoxy}$,

(D) $-\text{CO-O-R}_{N-8}$ where R_{N-8} is $-\text{H}$, $\text{C}_1\text{-C}_6\text{ alkyl}$,
phenyl or benzyl,

20 (E) $-\text{CO-NR}_{P-2}\text{R}_{P-3}$ where R_{P-2} and R_{P-3} are the same
or different and are as defined above,

(F) $-\text{CO-R}_{N-4}$ where R_{N-4} is as defined above,

(G) $-\text{SO}_2-(\text{C}_1\text{-C}_8\text{ alkyl})$,

(H) $-\text{SO}_2\text{-NR}_{P-2}\text{R}_{P-3}$ where R_{P-2} and R_{P-3} are the same
25 or different and are as defined above,

(I) $-\text{NH-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,

(J) $-\text{NH-CO-O-R}_{N-8}$ where R_{N-8} is as defined above,

(K) $-\text{NR}_{P-2}\text{R}_{P-3}$ where R_{P-2} and R_{P-3} are the same or
different and are as defined above,

30 (L) $-\text{R}_{N-4}$ where R_{N-4} is as defined above,

(M) $-\text{O-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,

(N) $-\text{O-CO-NR}_{N-8}\text{R}_{N-8}$ where the R_{N-8} s are the same or
different and are as defined above,

- (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl) optionally substituted with
 one, two, or three of -F, -Cl, -Br, -I,
 (Q) -NH-SO₂-(C₁-C₆ alkyl),
 5 (R) -F, -Cl,
 (III) -X-T-(C₁-C₆ alkyl) where X is absent or is C₁-C₆
 alkyl, T is oxygen or is NR_{P-2}, and where alkyl is optionally
 substituted with one, two, or three substituents selected from
 the group consisting of:
 10 (A) -OH,
 (B) -C₁-C₆ alkoxy,
 (C) -C₁-C₆ thioalkoxy,
 (D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl,
 phenyl, or benzyl,
 15 (E) -CO-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same
 or different and are as defined above,
 (F) -CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -SO₂-(C₁-C₈ alkyl),
 (H) -SO₂-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same
 20 or different and are as defined above,
 (I) -NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same or
 different and are as defined above,
 25 (L) -R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where the R_{N-8}s are the same or
 different and are as defined above,
 (O) -O-(C₁-C₅ alkyl)-COOH,
 30 (P) -O-(C₁-C₆) alkyl optionally substituted with
 one, two, or three of -F, -Cl, -Br, -I,
 (Q) -NH-SO₂-(C₁-C₆ alkyl),
 (R) -F, -Cl,

- (S) -O-R_P-aryl,
 (T) -O-R_P-heteroaryl, or
 (U) -O-R_P-hetrocycle,
 (IV) -(C₁-C₆ alkyl)-Z-(C₁-C₆ alkyl) where Z is S or O,
 5 and alkyl is optionally substituted with one, two, or three
 substituents selected from the group consisting of:
- (A) -OH,
 (B) -C₁-C₆ alkoxy,
 (C) -C₁-C₆ thioalkoxy,
 10 (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
 (E) -CO-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same
 or different and are as defined above,
 (F) -CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -SO₂-(C₁-C₈ alkyl),
 15 (H) -SO₂-NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same
 or different and are as defined above,
 (I) -NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -NR_{P-2}R_{P-3} where R_{P-2} and R_{P-3} are the same or
 20 different and are as defined above,
 (L) -R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where the R_{N-8}s are the same or
 different and are as defined above,
 25 (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆) alkyl optionally substituted with
 one, two, or three of -F, -Cl, -Br, -I,
 (Q) -NH-SO₂-(C₁-C₆ alkyl),
 (R) -F, -Cl,
 30 (V) -CH(-(CH₂)₀₋₂-O-R_{N-10})-(CH₂)₀₋₂-R_P-aryl/R_P-heteroaryl)
 where R_P-aryl and R_P-heteroaryl are as defined above, where R_{N-10} is
 selected from the group consisting of:
- (A) -H,

- 5 (B) C_1-C_6 alkyl,
 (C) C_3-C_7 cycloalkyl,
 (D) C_2-C_6 alkenyl with one double bond,
 (E) C_2-C_6 alkynyl with one triple bond,
 (F) R_{P-aryl} where R_{P-aryl} is as defined above,
 (G) $R_{P-heteroaryl}$ where $R_{P-heteroaryl}$ is as defined
 above,

(VI) - (C_3-C_8 cycloalkyl) where cycloalkyl is
 optionally substituted with one or two substituents selected
 10 from the group consisting of:

- (A) - $(CH_2)_{0-4}-OH$,
 (B) - $(CH_2)_{0-4}-C_1-C_6$ alkoxy,
 (C) - $(CH_2)_{0-4}-C_1-C_6$ thioalkoxy,
 (D) - $(CH_2)_{0-4}-CO-O-R_{N-8}$ where R_{N-8} is -H, C_1-C_6
 15 alkyl, phenyl or benzyl,
 (E) - $(CH_2)_{0-4}-CO-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are
 the same or different and are as defined above,
 (F) - $(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is as defined
 above,
 20 (G) - $(CH_2)_{0-4}-SO_2-(C_1-C_8$ alkyl),
 (H) - $(CH_2)_{0-4}-SO_2-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are
 the same or different and are as defined above,
 (I) - $(CH_2)_{0-4}-NH-CO-(C_1-C_6$ alkyl),
 (J) - $NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,
 25 (K) - $(CH_2)_{0-4}-NR_{P-2}R_{P-3}$ where R_{P-2} and R_{P-3} are the
 same or different and are as defined above,
 (L) - $(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,
 (M) - $O-CO-(C_1-C_6$ alkyl),
 (N) - $O-CO-NR_{N-8}R_{N-8}$ where the R_{N-8} s are the same or
 30 different and are as defined above,
 (O) - $O-(C_1-C_5$ alkyl)- $COOH$,

(P) $-O-C_1-C_6$ alkyl optionally substituted with one, two, or three of: $-F$, $-Cl$, $-Br$, $-I$,

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$,

(R) $-F$, $-Cl$;

5 (VII) $-(CR_{P-x}R_{P-y})_{0-4}-R_{P-1}$ where R_{P-1} is as defined above and each of R_{P-x} and R_{P-y} are independently any combination of:

(A) $-H$,

10 (B) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-COOH$, $-COO(C_1-C_6 \text{ alkyl})$, $-CONR_{1a}R_{1b}$ and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(C) $-CH_2-S-(C_1-C_6 \text{ alkyl})$,

15 (D) $-CH_2-CH_2-S-(C_1-C_6 \text{ alkyl})$,

(E) C_2-C_6 alkenyl with one or two double bonds,

(F) C_2-C_6 alkynyl with one or two triple bonds,

(G) $-(CH_2)_{n_1}-(R_{P-aryl})$ where n_1 is zero or one,

where R_{P-aryl} is as defined above,

20 (H) $-(CH_2)_{n_1}-(R_{P-heteroaryl})$ where n_1 and $R_{P-heteroaryl}$ are as defined above, or

(I) $-(CH_2)_{n_1}-(R_{P-heterocycle})$ where n_1 and $R_{P-heterocycle}$ are as defined above,

25 (J) $-(CH_2)_{0-3}-(C_3-C_7)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-CO-OH$, $-CO-O-(C_1-C_4 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, or

30 (K) F , Cl , Br or I ,

(L) $-NR_{P-2}$;

or R_{p-x} and R_{p-y} are taken together with the carbon to which they are attached to form a carbocycle of three thru seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of

5 -O-, -S-, -SO₂-, and -NR_{p-2}-;

(VIII) - (C₁-C₆ alkyl)-Z-(C₁-C₆ alkyl)-T-(C₁-C₆ alkyl)- wherein Z is S or O, T is O or NR_{p-2}, and alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

10 (A) -OH,

(B) -C₁-C₆ alkoxy,

(C) -C₁-C₆ thioalkoxy,

(D) -CO-O-R_{N-8} where R_{N-8} is as defined above,

(E) -CO-NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same

15 or different and are as defined above,

(F) -CO-R_{N-4} where R_{N-4} is as defined above,

(G) -SO₂-(C₁-C₈ alkyl),

(H) -SO₂-NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same

or different and are as defined above,

20 (I) -NH-CO-(C₁-C₆ alkyl),

(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) -NR_{p-2}R_{p-3} where R_{p-2} and R_{p-3} are the same or

different and are as defined above,

(L) -R_{N-4} where R_{N-4} is as defined above,

25 (M) -O-CO-(C₁-C₆ alkyl),

(N) -O-CO-NR_{N-8}R_{N-8} where the R_{N-8}s are the same or different and are as defined above,

(O) -O-(C₁-C₅ alkyl)-COOH,

(P) -O-(C₁-C₆ alkyl) optionally substituted with

30 one, two, or three of -F, -Cl, -Br, -I,

(Q) -NH-SO₂-(C₁-C₆ alkyl),


(R) -F, -Cl.


2. A compound or salt according to claim 1 wherein R_1 is $-(CH_2)_{0-1}-(R_1\text{-aryl})$.

3. A compound or salt according to claim 1 wherein $R_2 =$
5 $R_3 = H$.

4. A compound or salt according to claim 1 wherein R_C is
(a) $-(CR_{C-x}R_{C-y})_{0-4}-R_C\text{-aryl}$,
(b) $-(CR_{C-x}R_{C-y})_{0-4}-R_C\text{-heteroaryl}$, or
10 (c) $-(CR_{C-x}R_{C-y})_{0-4}-R_C\text{-heterocycle}$.

5. A compound or salt according to claim 1 wherein R_4 is





6. A compound or salt according to claim 1 wherein R_N is $R_{N-1}-X_N$ where X_N is $-CO-$, where R_{N-1} is R_N -aryl or R_N -heteroaryl.

- 5 7. A compound or salt according to claim 1 selected from the group consisting of
- 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)propyl 4-nitrophenyl carbonate,
- 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-

[(dipropylamino)carbonyl]benzoyl}amino)propyl 4-
 methylpiperazine-1-carboxylate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-
 (dimethylamino)ethyl carbonate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-
 (acetylaminio)ethylcarbamate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl
 piperazine-1-carboxylate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl 2-
 aminoethylcarbamate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl 3-
 aminopropylcarbamate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl (3R)-3-
 aminopyrrolidine-1-carboxylate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)propyl O-benzyl-L-
 tyrosinate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl L-
 tyrosinate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl 3-
 methoxypropanoate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino)carbonyl]benzoyl}amino)-propyl (2-
 hydroxyethoxy) acetate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-

[(dipropylamino)carbonyl]benzoyl}amino)-propyl D-lysinate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl [2-(2-methoxyethoxy)ethoxy]acetate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl {2-[[2-(dimethylamino)ethyl]-(methyl)amino]ethoxy}acetate,
 1-[(benzylamino)methyl]-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]benzoyl}amino)-propyl (4-methylpiperazin-1-yl)acetate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-pent-1-ynylbenzyl)amino]methyl}propyl (4-methylpiperazin-1-yl)acetate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-pent-1-ynylbenzyl)amino]methyl}propyl 4-methylpiperazine-1-carboxylate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-pent-1-ynylbenzyl)amino]methyl}propyl piperazine-1-carboxylate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-pent-1-ynylbenzyl)amino]methyl}propyl {2-[[2-(dimethylamino)ethyl]-(methyl)amino]ethoxy}acetate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-pent-1-ynylbenzyl)amino]methyl}propyl (2-methoxyethoxy)acetate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-[(dipropylamino)carbonyl]-5-methylbenzoyl}-amino)-1-{[(3-

pent-1-ynylbenzyl) amino] methyl} propyl L-tyrosinate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino) carbonyl]-5-methylbenzoyl}-amino)-1-{{3-
 pent-1-ynylbenzyl) amino] methyl} propyl (3R)-3-
 aminopyrrolidine-1-carboxylate,
 (1R,2S)-3-(3,5-difluorophenyl)-2-({3-
 [(dipropylamino) carbonyl]-5-methylbenzoyl}-amino)-1-{{3-
 pent-1-ynylbenzyl) amino] methyl} propyl tetrahydrofuran-3-
 yl carbonate,
 N'-[(1S,2R)-1-benzyl-3-(cyclohexylamino)-2-
 hydroxypropyl]-N,N-dipropylisophthalamide,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 4-methylpiperazine-1-carboxylate,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 piperazine-1-carboxylate,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 {2-[[2-(dimethylamino) ethyl] (methyl) amino] ethoxy} acetate,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 (2-methoxyethoxy) acetate,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 L-tyrosinate,
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 (3R)-3-aminopyrrolidine-1-carboxylate, and
 (1R,2S)-1-[(cyclohexylamino) methyl]-2-({3-
 [(dipropylamino) carbonyl] benzoyl}-amino)-3-phenylpropyl
 tetrahydrofuran-3-yl carbonate.

8. A method for making a compound of claim 1.

9. The use of a compound or salt according to claim 1 for the manufacture of a medicament.

5

10. The use of a compound or salt according to claim 1 for the manufacture of a medicament for use in the treatment or prevention of Alzheimer's disease, mild cognitive impairment Down's syndrome, Hereditary Cerebral Hemorrhage with
10 Amyloidosis of the Dutch-Type, cerebral amyloid angiopathy, other degenerative dementias, dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, or
15 diffuse Lewy body type of Alzheimer's disease.