



(51) International Patent Classification:

A23L 1/29 (2006.01) A61K 35/74 (2006.01)
A23L 1/30 (2006.01) A61P 1/00 (2006.01)
A61K 31/702 (2006.01)

(21) International Application Number:

PCT/US2011/067012

(22) International Filing Date:

22 December 2011 (22.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/428,869 31 December 2010 (31.12.2010) US

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; 100 Abbott Park Road, Dept. 0377 AP6A-1, Abbott Park, Illinois 60064 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHOW, JoMay** [US/US]; 5626 Blendon View Ct., Gahanna, Ohio 43230 (US). **BUCK, Rachael** [US/US]; 3710 Peak Ridge Drive, Gahanna, Ohio 43230 (US).

(74) Agents: **ENGLE, Mark R.** et al.; Abbott Nutrition a Division of Abbott Laboratories, 3300 Stelzer Road, Dept 108140 RP3-2, Columbus, Ohio 43219 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: SYNBIOTIC COMBINATION OF PROBIOTIC AND HUMAN MILK OLIGOSACCHARIDES TO PROMOTE GROWTH OF BENEFICIAL MICROBIOTA

FIG. 1

Effect of 2&3' Fucosyllactose on Gut Motility



(57) Abstract: Disclosed are nutritional compositions including human milk oligosaccharides that can be administered to individuals including preterm infants, infants, toddlers, and children for improving gastrointestinal function and tolerance, as well as the growth of beneficial microbiota. Suitable additional methods of using the nutritional compositions including the human milk oligosaccharides are also disclosed.

WO 2012/092155 A1

**SYNBIOTIC COMBINATION OF PROBIOTIC AND HUMAN MILK
OLIGOSACCHARIDES TO PROMOTE GROWTH OF BENEFICIAL
MICROBIOTA**

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/428,869 filed on December 31, 2010, which disclosure is incorporated by reference in its entirety.

FIELD OF THE DISCLOSURE

[0002] The present disclosure relates to human milk oligosaccharides (HMOs) for improving gastrointestinal function and tolerance in infants, toddlers, and children. More particularly, the present disclosure relates to human milk fortifiers, preterm and term infant formulas, and pediatric formulas comprising HMOs that can stimulate enteric nerve cells in the gastrointestinal tract, thereby treating and/or preventing numerous gastrointestinal-related conditions and diseases and promoting intestinal barrier integrity.

BACKGROUND OF THE DISCLOSURE

[0003] During postnatal development, a newborn's intestine experiences a process of maturation that ends with the production of gastrointestinal epithelium that functions as a selective barrier (i.e., gut barrier). The gastrointestinal epithelium permits the absorption of nutrients, electrolytes and water, while preventing exposure to dietary and microbial antigens, including food allergens. Specifically, this barrier limits the passage of antigens to the systemic circulation, thereby preventing infection, inflammatory reactions, and other gastrointestinal diseases and disorders that may occur during infancy and later in life. For very young infants, and particularly, preterm infants, who have an immature immune system and intestinal tract, development of suboptimal intestinal flora may result in infection, diarrhea, allergies, and food intolerance.

[0004] Barrier formation and maintenance has been found to be affected by the diet. Breast milk contains components that not only act as pathogen receptor analogues, but also activate immune factors by infant intestinal epithelial cells and/or associated

immune cell populations to enhance development and maturation of the infant's gastrointestinal and immune systems.

[0005] Not all infants, however, are in a position to receive human breast milk. It would therefore be desirable to provide nutritional compositions, and synthetic infant formulas in particular, that can produce nutritional benefits including improved gastrointestinal growth, development, and maturation. It would additionally be beneficial if the nutritional compositions could enhance immunity against microbial infections and other gastrointestinal diseases, conditions, and disorders.

SUMMARY OF THE DISCLOSURE

[0006] The present disclosure is directed to nutritional compositions, including synthetic infant formulas, synthetic pediatric formulas, and synthetic child formulas including at least one HMO alone or in combination with other components such as prebiotic oligosaccharides and/or probiotics, for improving gut function and immunity in an infant, toddler, child, or adult, along with related methods of use. More particularly, the nutritional compositions can improve growth and maturation of the gut barrier, thereby treating and/or preventing formula intolerance or other gastrointestinal diseases and/or disorders resulting from a loss of dysfunction of the gut barrier.

[0007] One embodiment is a synthetic pediatric formula for promoting intestinal barrier integrity. The synthetic pediatric formula comprises a first oligosaccharide in a concentration of from about 1 mg/mL to about 4 mg/mL and selected from the group consisting of a galactooligosaccharide, a fructooligosaccharide, and combinations thereof; and a second oligosaccharide on a concentration of from about 0.05 mg/mL to about 0.5 mg/mL and selected from the group consisting of 2'-fucosyllactose, 3'-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, lacto-N-neotetraose, and combinations thereof.

[0008] Another embodiment is directed to a method of promoting the growth of beneficial microbiota in the gastrointestinal tract of an infant or toddler in need thereof. The method comprises administering to the infant or toddler a synthetic pediatric formula comprising a probiotic, a first oligosaccharide in a concentration of from about 1 mg/mL to about 4 mg/mL and selected from the group consisting of galactooligosaccharide, fructooligosaccharide, and combinations thereof; and a second oligosaccharide in a

concentration of from about 0.05 mg/mL to about 0.5 mg/mL and selected from the group consisting of 2'-fucosyllactose, 3'-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, lacto-N-neotetraose, and combinations thereof.

[0009] It has been discovered that HMOs that are delivered to the gut tissue stimulate the gut-brain-immune axis, and improve the immune system and enteric nervous system. Specifically, it has been found that 2'-fucosyllactose stimulates enteric nerve cells in the gastrointestinal tract such that gut function may be improved and gastrointestinal issues minimized.

[0010] Additionally, it has been found that the digestive tolerance of an infant, toddler, child, or adult can be significantly increased by administering to the infant, toddler, child or adult a select blend of carbohydrates including HMOs. Specifically, the carbohydrate blend includes a combination of fast, medium, and slowly digested carbohydrates including specific HMOs such as lacto-N-neotetraose, 2'-fucosyllactose, 3'-fucosyllactose, 3'-sialyllactose and/or 6'-sialyllactose.

[0011] Moreover, it has been found that intestinal barrier integrity of an infant, toddler, child, or adult can be significantly improved by administering to the infant, toddler, child, or adult a synbiotic composition including HMOs. Specifically, the synbiotic combination includes a probiotic, at least one of a galactooligosaccharide and a fructooligosaccharide (such as a short chain fructooligosaccharide) and at least one HMO. The synbiotic composition promotes the colonization of beneficial intestinal microbiota in order to discourage the growth of harmful bacteria.

[0012] Although the nutritional compositions and methods are primarily discussed herein in relation to preterm infants and infants in general, it should be understood that many of the benefits discussed herein may be provided to toddlers, children, and adults administered combinations of the HMOs alone, or with other components as described herein, such as prebiotic oligosaccharides and/or probiotics, for example. Particularly, in some embodiments, the incidence of gastrointestinal diseases and disorders that generally affect adults, such as Crohn's disease, irritable bowel syndrome and the like, can be reduced with the use of the nutritional compositions of the present disclosure including HMOs.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] FIG. 1 is a graph depicting the effect of 2'FL and 3'FL on gut motility as measured in Example 35.

[0014] FIG. 2 is a table setting forth the microbiological medium used in the *in vitro* experiment of Example 36.

[0015] FIG. 3 is a graph depicting the change in pH over time as affected by the various oligosaccharide substrates as tested in Example 36.

[0016] FIG. 4 is a graph depicting change in total short chain fatty acid production over time as affected by the various oligosaccharide substrates as tested in Example 36.

[0017] FIGS. 5A-5H depict growth curves of various *Bifidobacterium* spp. as evaluated in Example 37.

[0018] FIGS. 6A-6H depict growth curves of various *Bifidobacterium* spp. as evaluated in Example 37.

[0019] FIGS. 7A-7G depict growth curves of various *Bifidobacterium* spp. as evaluated in Example 37.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0020] The nutritional compositions and methods described herein utilize HMOs alone or in combination with at least one other prebiotic oligosaccharide and/or a probiotic for controlling and reducing a number of diseases, disorders and conditions related to the gut-brain-immune system. These and other features of the nutritional compositions and methods, as well as some of the many optional variations and additions, are described in detail hereafter.

[0021] The terms "retort packaging" and "retort sterilizing" are used interchangeably herein, and unless otherwise specified, refer to the common practice of filling a container, most typically a metal can or other similar package, with a nutritional

liquid and then subjecting the liquid-filled package to the necessary heat sterilization step, to form a sterilized, retort packaged, nutritional liquid product.

[0022] The term "aseptic packaging" as used herein, unless otherwise specified, refers to the manufacture of a packaged product without reliance upon the above-described retort packaging step, wherein the nutritional liquid and package are sterilized separately prior to filling, and then are combined under sterilized or aseptic processing conditions to form a sterilized, aseptically packaged, nutritional liquid product.

[0023] The terms "fat" and "oil" as used herein, unless otherwise specified, are used interchangeably to refer to lipid materials derived or processed from plants or animals. These terms also include synthetic lipid materials so long as such synthetic materials are suitable for oral administration to humans.

[0024] The term "human milk oligosaccharide" or "HMO", unless otherwise specified, refers generally to a number of complex carbohydrates found in human breast milk that can be in acidic or neutral form, and to precursors thereof. Exemplary non-limiting human milk oligosaccharides include 3'-sialyllactose, 6'-sialyllactose, 3'-fucosyllactose, 2'-fucosyllactose, and lacto-N-neo-tetraose. Exemplary human milk oligosaccharide precursors include sialic acid and/or fucose.

[0025] The term "shelf stable" as used herein, unless otherwise specified, refers to a nutritional product that remains commercially stable after being packaged and then stored at 18-24°C for at least 3 months, including from about 6 months to about 24 months, and also including from about 12 months to about 18 months.

[0026] The terms "nutritional formulation" or "nutritional composition" as used herein, are used interchangeably and, unless otherwise specified, refer to synthetic formulas including nutritional liquids, nutritional powders, nutritional semi-liquids, nutritional semi-solids, nutritional supplements, and any other nutritional food product as known in the art. The nutritional powders may be reconstituted to form a nutritional liquid, all of which comprise one or more of fat, protein and carbohydrate and are suitable for oral consumption by a human. The terms "nutritional formulation" and "nutritional composition" do not include human breast milk.

[0027] The term “nutritional liquid” as used herein, unless otherwise specified, refers to nutritional compositions in ready-to-drink liquid form, concentrated form, and nutritional liquids made by reconstituting the nutritional powders described herein prior to use.

[0028] The term “nutritional powder” as used herein, unless otherwise specified, refers to nutritional compositions in flowable or scoopable form that can be reconstituted with water or another aqueous liquid prior to consumption and includes both spraydried and drymixed/dryblended powders.

[0029] The term “nutritional semi-solid,” as used herein, unless otherwise specified, refers to nutritional products that are intermediate in properties, such as rigidity, between solids and liquids. Some semi-solids examples include puddings, gelatins, and doughs.

[0030] The term “nutritional semi-liquid,” as used herein, unless otherwise specified, refers to nutritional products that are intermediate in properties, such as flow properties, between liquids and solids. Some semi-liquids examples include thick shakes and liquid gels.

[0031] The term “infant” as used herein, unless otherwise specified, refers to a person 12 months or younger. The term “preterm infant” as used herein, refers to a person born prior to 36 weeks of gestation.

[0032] The term “toddler” as used herein, unless otherwise specified, refers to a person greater than one year of age up to three years of age.

[0033] The term “child” as used herein, unless otherwise specified, refers to a person greater than three years of age up to twelve years of age.

[0034] The term “newborn” as used herein, unless otherwise specified, refers to a person from birth up to four weeks of age.

[0035] The terms “infant formula” or “synthetic infant formula” as used herein, unless otherwise specified, are used interchangeably and refer to liquid, solid, semi-liquid, and semi-solid human milk replacements or substitutes that are suitable for consumption by

an infant. The synthetic formulas include components that are of semi-purified or purified origin. As used herein, unless otherwise specified, the terms “semi-purified” or “purified” refer to a material that has been prepared by purification of a natural material or by synthesis. The terms “infant formula” or “synthetic infant formula” do not include human breast milk.

[0036] The term “synthetic pediatric formula” as used herein, unless otherwise specified, refers to liquid, solid, semi-solid, and semi-liquid human milk replacements or substitutes that are suitable for consumption by an infant or toddler up to the age of 36 months (3 years). The synthetic formulas include components that are of semi-purified or purified origin. As used herein, unless otherwise specified, the terms “semi-purified” or “purified” refer to a material that has been prepared by purification of a natural material or by synthesis. The term “synthetic pediatric nutritional formula” does not include human breast milk.

[0037] The term “synthetic child formula” as used herein, unless otherwise specified, refers to liquid, solid, semi-liquid, and semi-solid human milk replacements or substitutes that are suitable for consumption by a child up to the age of 12 years. The synthetic formulas include components that are of semi-purified or purified origin. As used herein, unless otherwise specified, the terms “semi-purified” or “purified” refer to a material that has been prepared by purification of a natural material or by synthesis. The term “synthetic child nutritional formula” does not include human breast milk.

[0038] The term “preterm infant formula” as used herein, unless otherwise specified, refers to liquid and solid nutritional products suitable for consumption by a preterm infant.

[0039] The term “human milk fortifier” as used herein, unless otherwise specified, refers to liquid and solid nutritional products suitable for mixing with breast milk or preterm infant formula or infant formula for consumption by a preterm or term infant.

[0040] The terms “susceptible” and “at risk” as used herein, unless otherwise specified, mean having little resistance to a certain condition or disease, including being genetically predisposed, having a family history of, and/or having symptoms of the condition or disease.

[0041] The term “cognition” as used herein, unless otherwise specified, refers to an individual’s ability for learning, memory acquisition, and memory recall.

[0042] The terms “growth of a virus” or “growth of bacteria” as used herein, unless otherwise specified, refer to the production, proliferation, or replication of a virus or bacteria.

[0043] All percentages, parts and ratios as used herein, are by weight of the total composition, unless otherwise specified. All such weights, as they pertain to listed ingredients, are based on the active level and, therefore, do not include solvents or by-products that may be included in commercially available materials, unless otherwise specified.

[0044] Numerical ranges as used herein are intended to include every number and subset of numbers within that range, whether specifically disclosed or not. Further, these numerical ranges should be construed as providing support for a claim directed to any number or subset of numbers in that range. For example, a disclosure of from 1 to 10 should be construed as supporting a range of from 2 to 8, from 3 to 7, from 5 to 6, from 1 to 9, from 3.6 to 4.6, from 3.5 to 9.9, and so forth.

[0045] All references to singular characteristics or limitations of the present disclosure shall include the corresponding plural characteristic or limitation, and vice versa, unless otherwise specified or clearly implied to the contrary by the context in which the reference is made.

[0046] All combinations of method or process steps as used herein can be performed in any order, unless otherwise specified or clearly implied to the contrary by the context in which the referenced combination is made.

[0047] The nutritional compositions and methods may comprise, consist of, or consist essentially of the essential elements of the compositions and methods as described herein, as well as any additional or optional element described herein or otherwise useful in nutritional composition applications.

Product Form

[0048] The nutritional compositions of the present disclosure may be formulated and administered in any known or otherwise suitable oral product form. Any solid, liquid, semi-solid, semi-liquid or powder product form, including combinations or variations thereof, are suitable for use herein, provided that such forms allow for safe and effective oral delivery to the individual of the essential ingredients and any optional ingredients, as also defined herein.

[0049] The nutritional compositions of the present disclosure are desirably formulated as dietary product forms, which are defined herein as those embodiments comprising the ingredients of the present disclosure in a product form that then contains at least one of fat, protein, and carbohydrate, and preferably also contains vitamins, minerals, or combinations thereof. The nutritional compositions will comprise at least one HMO, and many times at least two or more HMOs, desirably in combination with at least one of protein, fat, vitamins, and minerals, to produce a nutritional combination.

[0050] The nutritional composition may be formulated with sufficient kinds and amounts of nutrients to provide a sole, primary, or supplemental source of nutrition, or to provide a specialized nutritional composition for use in individuals afflicted with specific diseases, disorders, or conditions or with a targeted nutritional benefit as described below.

[0051] Specific non-limiting examples of product forms suitable for use with the HMO-containing compositions as disclosed herein include, for example, liquid and powdered dietary supplements, liquid and powdered human milk fortifiers, liquid and powdered preterm infant formulas, liquid and powdered infant formulas, liquid and powdered elemental and semi-elemental formulas, liquid and powdered pediatric formulas, liquid and powdered toddler formulas, liquid and powdered follow-on formulas, liquid, powdered and solid adult nutritional formulas suitable for use with individuals suffering from food intolerance, allergies, immune disorders, and other gastrointestinal diseases, conditions, and/or disorders.

Nutritional Liquids

[0052] Nutritional liquids include both concentrated and ready-to-feed nutritional liquids. These nutritional liquids are most typically formulated as suspensions or emulsions, although other liquid forms are within the scope of the present disclosure.

[0053] Nutritional emulsions suitable for use may be aqueous emulsions comprising proteins, fats, and carbohydrates. These emulsions are generally flowable or drinkable liquids at from about 1°C to about 25°C and are typically in the form of oil-in-water, water-in-oil, or complex aqueous emulsions, although such emulsions are most typically in the form of oil-in-water emulsions having a continuous aqueous phase and a discontinuous oil phase.

[0054] The nutritional emulsions may be and typically are shelf stable. The nutritional emulsions typically contain up to about 95% by weight of water, including from about 50% to about 95%, also including from about 60% to about 90%, and also including from about 70% to about 85%, by weight of water. The nutritional emulsions may have a variety of product densities, but most typically have a density greater than about 1.03 g/mL, including greater than about 1.04 g/mL, including greater than about 1.055 g/mL, including from about 1.06 g/mL to about 1.12 g/mL, and also including from about 1.085 g/mL to about 1.10 g/mL.

[0055] The nutritional emulsions may have a caloric density tailored to the nutritional needs of the ultimate user, although in most instances the emulsions comprise generally at least 19 kcal/fl oz (660 kcal/liter), more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 25 kcal/fl oz (820 kcal/liter), even more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 24 kcal/fl oz (800-810 kcal/liter). Generally, the 22-24 kcal/fl oz formulas are more commonly used in preterm or low birth weight infants, and the 20-21 kcal/fl oz (675-680 to 700 kcal/liter) formulas are more often used in term infants. In some embodiments, the emulsion may have a caloric density of from about 50-100 kcal/liter to about 660 kcal/liter, including from about 150 kcal/liter to about 500 kcal/liter. In some specific embodiments, the emulsion may have a caloric density of 25, or 50, or 75, or 100 kcal/liter.

[0056] The nutritional emulsion may have a pH ranging from about 3.5 to about 8, but are most advantageously in a range of from about 4.5 to about 7.5, including from about 5.5 to about 7.3, including from about 6.2 to about 7.2.

[0057] Although the serving size for the nutritional emulsion can vary depending upon a number of variables, a typical serving size is generally at least about 1 mL, or even at least about 2 mL, or even at least about 5 mL, or even at least about 10 mL, or even at least about 25 mL, including ranges from about 2 mL to about 300 mL, including from about 4 mL to about 250 mL, and including from about 10 mL to about 240 mL.

Nutritional Solids

[0058] The nutritional solids may be in any solid form, but are typically in the form of flowable or substantially flowable particulate compositions, or at least particulate compositions. Particularly suitable nutritional solid product forms include spray dried, agglomerated and/or dryblended powder compositions. The compositions can easily be scooped and measured with a spoon or similar other device, and can easily be reconstituted by the intended user with a suitable aqueous liquid, typically water, to form a nutritional composition for immediate oral or enteral use. In this context, "immediate" use generally means within about 48 hours, most typically within about 24 hours, preferably right after reconstitution.

[0059] The nutritional powders may be reconstituted with water prior to use to a caloric density tailored to the nutritional needs of the ultimate user, although in most instances the powders are reconstituted with water to form compositions comprising at least 19 kcal/fl oz (660 kcal/liter), more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 25 kcal/fl oz (820 kcal/liter), even more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 24 kcal/fl oz (800-810 kcal/liter). Generally, the 22-24 kcal/fl oz formulas are more commonly used in preterm or low birth weight infants, and the 20-21 kcal/fl oz (675-680 to 700 kcal/liter) formulas are more often used in term infants. In some embodiments, the reconstituted powder may have a caloric density of from about 50-100 kcal/liter to about 660 kcal/liter, including from about 150 kcal/liter to about 500 kcal/liter. In some specific embodiments, the emulsion may have a caloric density of 25, or 50, or 75, or 100 kcal/liter.

Human Milk Oligosaccharides (HMOs)

[0060] The nutritional compositions of the present disclosure include at least one HMO, and in many embodiments, a combination of two or more HMOs. Oligosaccharides are one of the main components of human breast milk, which contains, on average, 10 grams per liter of neutral oligosaccharides and 1 gram per liter of acidic oligosaccharides. The compositional structure of HMOs is very complex and more than 200 different oligosaccharide-like structures are known.

[0061] The HMO or HMOs may be included in the nutritional compositions alone, or in some embodiments, in combination with other components (e.g., prebiotic oligosaccharides, probiotics, etc.) as described herein. In many embodiments, HMOs are included in the nutritional compositions with multiple additional components. The HMO or HMOs may be isolated or enriched from milk(s) secreted by mammals including, but not limited to: human, bovine, ovine, porcine, or caprine species. The HMOs may also be produced via microbial fermentation, enzymatic processes, chemical synthesis, or combinations thereof.

[0062] Suitable HMOs for use in the nutritional compositions may include neutral oligosaccharides, acidic oligosaccharides, n-acetylglucosylated oligosaccharides, and HMO precursors. Specific non-limiting examples of HMOs that may be included individually or in combination in the compositions of the present disclosure include: sialic acid (i.e., free sialic acid, lipid-bound sialic acid, protein-bound sialic acid); D-glucose (Glc); D-galactose (Gal); N-acetylglucosamine (GlcNAc); L-fucose (Fuc); fucosyl oligosaccharides (i.e., Lacto-N-fucopentaose I; Lacto-N-fucopentaose II; 2'-Fucosyllactose; 3'-Fucosyllactose; Lacto-N-fucopentaose III; Lacto-N-difucohexaose I; and Lactodifucotetraose); non-fucosylated, non-sialylated oligosaccharides (i.e., Lacto-N-tetraose and Lacto-N-neotetraose); sialyl oligosaccharides (i.e., 3'-Sialyl-3-fucosyllactose; Disialomonofucosyllacto-N-neohexaose; Monofucosylmonosialyllacto-N-octaose (sialyl Lea); Sialyllacto-N-fucohexaose II; Disialyllacto-N-fucopentaose II; Monofucosyldisialyllacto-N-tetraose); and sialyl fucosyl oligosaccharides (i.e., 2'-Sialyllactose; 2-Sialyllactosamine; 3'-Sialyllactose; 3'-Sialyllactosamine; 6'-Sialyllactose; 6'-Sialyllactosamine; Sialyllacto-N-neotetraose c; Monosialyllacto-N-hexaose; Disialyllacto-N-hexaose I; Monosialyllacto-N-neohexaose I; Monosialyllacto-N-

neohexaose II; Disialyllacto-N-neohexaose; Disialyllacto-N-tetraose; Disialyllacto-N-hexaose II; Sialyllacto-N-tetraose a; Disialyllacto-N-hexaose I; and Sialyllacto-N-tetraose b). Also useful are variants in which the glucose (Glc) at the reducing end is replaced by N-acetylglucosamine (e.g., 2'-fucosyl-N-acetylglucosamine (2'-FLNac) is such a variant to 2'-fucosyllactose). These HMOs are described more fully in U.S. Patent Application No. 2009/0098240, which is herein incorporated by reference in its entirety. Other suitable examples of HMOs that may be included in the compositions of the present disclosure include lacto-N-fucopentaose V, lacto-N-hexaose, para-lacto-N-hexaose, lacto-N-neohexaose, para-lacto-N-neohexaose, monofucosyllacto-N-hexaose II, isomeric fucosylated lacto-N-hexaose (1), isomeric fucosylated lacto-N-hexaose (3), isomeric fucosylated lacto-N-hexaose (2), difucosyl-para-lacto-N-neohexaose, difucosyl-para-lacto-N-hexaose, difucosyllacto-N-hexaose, lacto-N-neoocataose, para-lacto-N-octanose, iso-lacto-N-octaose, lacto-N-octaose, monofucosyllacto-neoocataose, monofucosyllacto-N-octaose, difucosyllacto-N-octaose I, difucosyllacto-N-octaose II, difucosyllacto-N-neoocataose II, difucosyllacto-N-neoocataose I, lacto-N-decaose, trifucosyllacto-N-neooctaose, trifucosyllacto-N-octaose, trifucosyl-iso-lacto-N-octaose, lacto-N-difuco-hexaose II, sialyl-lacto-N-tetraose a, sialyl-lacto-N-tetraose b, sialyl-lacto-N-tetraose c, sialyl-fucosyl-lacto-N-tetraose I, sialyl-fucosyl-lacto-N-tetraose II, and disialyl-lacto-N-tetraose, and combinations thereof. Particularly suitable nutritional compositions include at least one of the following HMOs or HMO precursors: sialic acid (SA); 2'-Sialyllactose (2'SL); 3'-Sialyllactose (3'SL); 6'-Sialyllactose (6'SL); 2'-Fucosyllactose (2'FL); 3'-Fucosyllactose (3'FL); and Lacto-N-tetraose and Lacto-N-neotetraose (LNnT), and in particular, combinations of 2'FL with at least one of 6'SL and 3'SL; and combinations of LNnT with at least one of 6'SL and 3'FL.

[0063] Other exemplary combinations include: SA, 3'SL, 6'SL, 3'FL, 2'FL, and LNnT; 3'SL, 6'SL, 3'FL, 2'FL, and LNnT; SA, 6'SL, 3'FL, 2'FL, and LNnT; SA, 3'SL, 3'FL, 2'FL, and LNnT; SA, 3'SL, 6'SL, 2'FL, and LNnT; SA, 3'SL, 6'SL, 3'FL, and LNnT; SA, 3'SL, 6'SL, 3'FL, and 2'FL; SA and 3'SL; SA and 6'SL; SA and 2'FL; SA and LNnT; SA, 3'SL, and 6'SL; SA, 3'SL and 3'FL; SA, 3'SL and 2'FL; SA, 3'SL and LNnT; SA, 6'SL and 3'FL; SA, 6'SL, and 2'FL; SA, 6'SL, and LNnT; SA, 3'FL, and 2'FL; SA, 3'FL, and LNnT; SA, 2'FL, and LNnT; SA, 3'SL, 6'SL, and 3'FL; SA, 3'SL, 6'SL and 2'FL; SA, 3'SL, 6'SL, and LNnT; SA, 3'SL, 3'FL, and 2'FL; SA, 3'SL, 3'FL, and LNnT;

SA, 3'SL, 2'FL, and LNnT; SA, 6'SL, 3'FL, and 2'FL; SA, 6'SL, 2'FL, and LNnT; SA, 6'SL, 3'FL, and LNnT; SA, 3'FL, 2'FL, and LNnT; SA, 6'SL, 2'FL, and LNnT; SA, 3'SL, 3'FL, 2'FL, and LNnT; SA, 6'SL, 3'FL, 2'FL, and LNnT; SA, 3'SL, 6'SL, 3'FL, and LNnT; SA, 3'SL, 3'FL, 2'FL, and LNnT; SA, 3'SL, 6'SL, 2'FL, and LNnT; 3'SL, 6'SL, 3'FL, and 2'FL; 3'SL, 6'SL, 2'FL, and LNnT; 3'SL, 3'FL, 2'FL, and LNnT; 3'SL, 6'SL, 3'FL, and LNnT; 3'SL, 6'SL, and 3'FL; 3'SL, 3'FL, and 2'FL; 3'SL, 2'FL, and LNnT; 3'SL, 6'SL, and 2'FL; 3'SL, 6'SL, and LNnT; 3'SL and 3'FL; 3'SL and 2'FL; 3'SL and LNnT; 6'SL and 3'FL; 6'SL and 2'FL; 6'SL and LNnT; 6'SL, 3'FL, and LNnT; 6'SL, 3'FL, 2'FL, and LNnT; 3'FL, 2'FL, and LNnT; 3'FL and LNnT; and 2'FL and LNnT.

[0064] The HMOs are present in the nutritional compositions in total amounts of HMO in the composition (mg of HMO per mL of composition) of at least about 0.001 mg/mL, including at least about 0.01 mg/mL, including from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to about 15 mg/mL, including from about 0.01 mg/mL to about 15 mg/mL, including from about 0.001 mg/mL to about 10 mg/mL, including from about 0.01 mg/mL to about 10 mg/mL, including from about 0.001 mg/mL to about 5 mg/mL, including from about 0.01 mg/mL to about 5 mg/mL, and including from about 0.001 mg/mL to about 1 mg/mL of total HMO in the nutritional composition, including from about 0.001 mg/mL to about 0.23 mg/mL, and including from about 0.01 mg/mL to about 0.23 mg/mL. Typically, the amount of HMO in the nutritional composition will depend on the specific HMO or HMOs present and the amounts of other components in the nutritional compositions.

[0065] In one specific embodiment when the nutritional composition is a nutritional powder, the total concentration of HMOs in the nutritional powder is from about 0.0005% to about 5%, including from about 0.01% to about 1% (by weight of the nutritional powder).

[0066] In another specific embodiment, when the nutritional composition is a ready-to-feed nutritional liquid, the total concentration of HMOs in the ready-to-feed nutritional liquid is from about 0.0001% to about 0.50%, including from about 0.001% to about 0.15%, including from about 0.01% to about 0.10%, and further including from about 0.01% to about 0.03% (by weight of the ready-to-feed nutritional liquid).

[0067] In another specific embodiment, when the nutritional composition is a concentrated nutritional liquid, the total concentration of HMOs in the concentrated liquid is from about 0.0002% to about 0.60%, including from about 0.002% to about 0.30%, including from about 0.02% to about 0.20%, and further including from about 0.02% to about 0.06% (by weight of the concentrated nutritional liquid).

[0068] In one specific embodiment, the nutritional composition includes a neutral human milk oligosaccharide in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to less than 2 mg/mL, and including from about 0.01 mg/mL to less than 2 mg/mL.

[0069] In one specific embodiment of the present disclosure, a nutritional composition includes 2'FL. The 2'FL may be the only HMO included in the nutritional composition, or other additional HMOs may also be included in the nutritional composition (e.g., the 2'FL may be combined with 3'SL and/or 6'SL in some specific embodiments). In one embodiment, the 2'FL is included in the nutritional composition in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to less than 2 mg/mL, and including from about 0.01 mg/mL to less than 2 mg/mL. In another embodiment, the 2'FL is included in the nutritional composition in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from greater than 2.5 mg/mL to about 20 mg/mL, including from greater than 2.5 mg/mL to about 15 mg/mL, and including from greater than 2.5 mg/mL to about 10 mg/mL.

[0070] In one specific embodiment, the nutritional composition includes 6'SL, alone or in combination with other HMOs, in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to less than 0.25 mg/mL, and including from about 0.01 mg/mL to less than 0.25 mg/mL. In another embodiment, the nutritional composition includes 6'SL, alone or in combination with other HMOs, in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from greater than 0.4 mg/mL to about 20 mg/mL, including from greater than 0.4 mg/mL to about 15 mg/mL, and including from greater than 0.4 mg/mL to about 10 mg/mL.

[0071] In one embodiment, when the nutritional composition includes 6'SL, the total amount of HMOs in the nutritional composition includes at least about 88% (by total weight HMOs) 6'SL, including from about 88% (by total weight HMOs) to about 96% (by total weight HMOs), including from about 88% (by total weight HMOs) to about 100% (by total weight HMOs), and including about 100% (by total weight HMOs) 6'SL.

[0072] In another embodiment, the nutritional composition includes 3'SL, alone or in combination with other HMOs, in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to less than 0.15 mg/mL, including from about 0.01 mg/mL to less than 0.15 mg/mL, including from greater than 0.25 mg/mL to about 20 mg/mL, including from greater than 0.25 mg/mL to about 15 mg/mL, and including from greater than 0.25 mg/mL to about 10 mg/mL.

[0073] In one embodiment, when the nutritional composition includes 3'SL, the total amount of HMOs in the nutritional composition includes at least about 85% (by total weight HMOs) 3'SL, including from about 85% (by total weight HMOs) to about 88% (by total weight HMOs), including from about 88% (by total weight HMOs) to about 100% (by total weight HMOs), and including about 100% (by total weight HMOs) 3'SL.

[0074] In one specific embodiment, the nutritional composition includes LNnT, alone or in combination with other HMOs, in an amount of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from about 0.001 mg/mL to less than 0.2 mg/mL, including from about 0.01 mg/mL to less than 0.2 mg/mL, including from greater than 0.32 mg/mL to about 20 mg/mL, including from greater than 0.32 mg/mL to about 15 mg/mL, and including from greater than 0.32 mg/mL to about 10 mg/mL.

Additional Prebiotic Oligosaccharides

[0075] The nutritional compositions of the present disclosure may, in addition to the HMOs described above, comprise an additional source or sources of prebiotic oligosaccharides (the total amount of oligosaccharides being referred to herein as an "oligosaccharide blend" of the nutritional composition). Suitable additional sources of prebiotic oligosaccharides for use in the nutritional compositions include any prebiotic

oligosaccharide that is suitable for use in an oral nutritional composition and is compatible with the essential elements and features of such compositions. In some embodiments, the nutritional composition includes a combination of one or more HMOs and one or more additional prebiotic oligosaccharides such that the composition provides a synergistic benefit to the end user, such as a synergistic benefit in improving feeding intolerance in infants.

[0076] In some embodiments, the combinations of HMO or HMOs with the additional prebiotic oligosaccharides to provide the synergistic effect include HMOs and additional prebiotic oligosaccharides that ferment at a rapid rate (“rapidly-fermenting oligosaccharides”), oligosaccharides that ferment at a moderate rate (“medium-fermenting oligosaccharides”), and/or oligosaccharides that ferment at a slow rate (“slowly-fermenting oligosaccharides”). Some preferred embodiments provide a nutritional composition that includes at least one HMO in combination with a rapidly-fermenting oligosaccharide, a medium-fermenting oligosaccharide, and/or a slowly-fermenting oligosaccharide.

[0077] Non-limiting examples of suitable additional prebiotic oligosaccharides for use in the nutritional compositions described herein include prebiotic oligosaccharides that have a degree of polymerization (DP) of at least 2 monose units, which are not or only partially digested in the intestine by the action of acids or digestive enzymes present in the human upper digestive tract (small intestine and stomach), but which are fermentable by the human intestinal flora. The term “monose units” refers to units having a closed ring structure, preferably hexose, e.g., the pyranose or furanose forms. Particularly preferred oligosaccharides for use in combination with the HMO or HMOs in the nutritional compositions of the present disclosure include galactooligosaccharides (GOS), fructooligosaccharides (FOS), short chain fructooligosaccharides, inulin, polydextrose (PDX), pectin hydrolysate, and gum fiber. In one specific embodiment, the gum fiber is gum arabic.

[0078] The oligosaccharide blend is present in the nutritional compositions in a total amount of at least about 1 mg/mL, including from about 1 mg/mL to about 20 mg/mL, including from about 1 mg/mL to about 15 mg/mL, including from about 1 mg/mL to about 10 mg/mL, and including from about 1 mg/mL to about 5 mg/mL. In one embodiment, the

oligosaccharide blend is present in the nutritional composition in a total amount of from about 1 mg/mL to about 10 mg/mL.

[0079] Typically, when used as an oligosaccharide blend, the nutritional compositions, in addition to the HMO or HMOs, include at least one rapidly-fermented oligosaccharide, at least one medium-fermented oligosaccharide, and, optionally, at least one slowly-fermented oligosaccharide to provide a nutritional composition that is tolerated well by preterm and term infants (i.e., reduced gassiness and/or stool frequency). Rapidly-fermented oligosaccharides generally have a fermentation rate of greater than 4,000 $\mu\text{g/g}$ of dry matter/hour; medium-fermented oligosaccharides generally have a fermentation rate of from 1,500 $\mu\text{g/g}$ of dry matter/hour to 4,000 $\mu\text{g/g}$ of dry matter/hour; and slowly-fermented oligosaccharides generally have a fermentation rate of less than 1,500 $\mu\text{g/g}$ of dry matter/hour.

[0080] By way of specific example, rapidly-fermented oligosaccharides include FOS, GOS (about 9,304 $\mu\text{g/g}$ of dry matter/hour), LNnT (about 4,488 $\mu\text{g/g}$ of dry matter/hour), 2'FL (about 4,872 $\mu\text{g/g}$ of dry matter/hour), and combinations thereof. Medium-fermented oligosaccharides include 6'SL (about 1,809 $\mu\text{g/g}$ of dry matter/hour), 3'SL, 2'FL, 3'FL, and LNnT, and combinations thereof. Slowly-fermented oligosaccharides include longer chain carbohydrates such as inulin (about 1,435 $\mu\text{g/g}$ of dry matter/hour), gum fibers (e.g., gum arabic (about 785 $\mu\text{g/g}$ of dry matter/hour)), and combinations thereof.

[0081] When used in an oligosaccharide blend, the rapidly-fermented oligosaccharides can be included in the nutritional compositions in amounts of from about 0.05 mg/mL to about 20 mg/mL, including from about 0.5 mg/mL to about 15 mg/mL, including from about 0.5 mg/mL to about 10 mg/mL, including from about 1 mg/mL to about 15 mg/mL, including from about 1 mg/mL to about 10 mg/mL, including from about 2 mg/mL to about 8 mg/mL, and also including from about 3 mg/mL to about 5 mg/mL. The medium-fermented oligosaccharides can be included in the nutritional compositions in amounts of from about 0.05 mg/mL to about 20 mg/mL, including from about 0.05 mg/mL to about 15 mg/mL, including from about 0.05 mg/mL to about 10 mg/mL, including from about 0.05 mg/mL to about 5 mg/mL, including from about 0.05 mg/mL to about 2.5 mg/mL, including from about 0.05 mg/mL to about 1 mg/mL, including from about 0.05

mg/mL to about 0.5 mg/mL, and including from about 0.05 mg/mL to about 0.25 mg/mL. The slowly-fermented oligosaccharides can be included in the nutritional compositions in amounts of from about 0.05 mg/mL to about 20 mg/mL, including from about 0.05 mg/mL to about 15 mg/mL, including from about 0.05 mg/mL to about 10 mg/mL, including from about 0.05 mg/mL to about 5 mg/mL, and also including from about 0.05 mg/mL to about 2.5 mg/mL.

[0082] In one specific embodiment, the nutritional composition includes an oligosaccharide blend including LNnT, 6'SL and inulin in a total amount of oligosaccharide blend of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL.

[0083] In another specific embodiment, the nutritional composition includes an oligosaccharide blend including 2'FL, 6'SL and inulin in a total amount of oligosaccharide blend of from about 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL.

[0084] Other exemplary combinations include: FOS, GOS, 2'FL, LNnT, 3'SL, and 6'SL; FOS, GOS, 2'FL, 3'SL, and 6'SL; FOS, GOS, LNnT, 3'SL, and 6'SL; FOS, 2'FL, LNnT, 3'SL, and 6'SL; GOS, 2'FL, LNnT, 3'SL, and 6'SL; FOS, GOS, 3'SL, and 6'SL; FOS, 2'FL, 3'SL, and 6'SL; FOS, LNnT, 3'SL, and 6'SL; GOS, 2'FL, 3'SL, and 6'SL; GOS, LNnT, 3'SL, and 6'SL; 2'FL, LNnT, 3'SL, and 6'SL; FOS, 3'SL, and 6'SL; GOS, 3'SL, and 6'SL; 2'FL, 3'SL, and 6'SL; LNnT, 3'SL, and 6'SL; FOS, GOS, 2'FL, LNnT, and 3'SL; FOS, GOS, 2'FL, and 3'SL; FOS, GOS, LNnT, and 3'SL; FOS, 2'FL, LNnT, and 3'SL; GOS, 2'FL, LNnT, and 3'SL; FOS, GOS, and 3'SL; FOS, 2'FL, and 3'SL; FOS, LNnT, and 3'SL; GOS, 2'FL, and 3'SL; GOS, LNnT, and 3'SL; 2'FL, LNnT, and 3'SL; FOS and 3'SL; GOS and 3'SL; 2'FL and 3'SL; LNnT and 3'SL; FOS, GOS, 2'FL, LNnT, and 6'SL; FOS, GOS, 2'FL, and 6'SL; FOS, GOS, LNnT, and 6'SL; FOS, 2'FL, LNnT, and 6'SL; GOS, 2'FL, LNnT, and 6'SL; FOS, GOS, and 6'SL; FOS, 2'FL, and 6'SL; FOS, LNnT, and 6'SL; GOS, 2'FL, and 6'SL; GOS, LNnT, and 6'SL; 2'FL, LNnT, and 6'SL; FOS and 6'SL; GOS and 6'SL; 2'FL and 6'SL; and LNnT and 6'SL.

[0085] Further exemplary combinations include: FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, 2'FL,

LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, 2'FL, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; GOS, 2'FL, 3'SL, 6'SL, inulin, a gum, and polydextrose; GOS, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; 2'FL, LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, 3'SL, 6'SL, inulin, a gum, and polydextrose; GOS, 3'SL, 6'SL, inulin, a gum, and polydextrose; 2'FL, 3'SL, 6'SL, inulin, a gum, and polydextrose; LNnT, 3'SL, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, 3'SL, inulin, a gum, and polydextrose; FOS, GOS, LNnT, 3'SL, inulin, a gum, and polydextrose; FOS, 2'FL, LNnT, 3'SL, inulin, a gum, and polydextrose; GOS, 2'FL, LNnT, 3'SL, inulin, a gum, and polydextrose; FOS, GOS, 3'SL, inulin, a gum, and polydextrose; FOS, GOS, 3'SL, inulin, a gum, and polydextrose; FOS, 2'FL, 3'SL, inulin, a gum, and polydextrose; FOS, LNnT, 3'SL, inulin, a gum, and polydextrose; GOS, 2'FL, 3'SL, inulin, a gum, and polydextrose; GOS, LNnT, 3'SL, inulin, a gum, and polydextrose; 2'FL, LNnT, 3'SL, inulin, a gum, and polydextrose; FOS, 3'SL, inulin, a gum, and polydextrose; GOS, 3'SL, inulin, a gum, and polydextrose; 2'FL, 3'SL, inulin, a gum, and polydextrose; LNnT, 3'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, LNnT, 6'SL, inulin, a gum, and polydextrose; FOS, 2'FL, LNnT, 6'SL, inulin, a gum, and polydextrose; GOS, 2'FL, LNnT, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 6'SL, inulin, a gum, and polydextrose; FOS, 2'FL, 6'SL, inulin, a gum, and polydextrose; FOS, LNnT, 6'SL, inulin, a gum, and polydextrose; GOS, 2'FL, 6'SL, inulin, a gum, and polydextrose; GOS, LNnT, 6'SL, inulin, a gum, and polydextrose; 2'FL, LNnT, 6'SL, inulin, a gum, and polydextrose; FOS, 6'SL, inulin, a gum, and polydextrose; GOS, 6'SL, inulin, a gum, and polydextrose; 2'FL, 6'SL, inulin, a gum, and polydextrose; LNnT, 6'SL, inulin, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and a gum; FOS, GOS, 2'FL, 3'SL, 6'SL, inulin, and a gum; FOS, GOS, LNnT, 3'SL, 6'SL, inulin, and a gum; FOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and a gum; GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and a gum; FOS, GOS, 3'SL, 6'SL, inulin, and a gum; FOS, 2'FL, 3'SL, 6'SL, inulin, and a gum; FOS, LNnT, 3'SL, 6'SL, inulin, and a gum; GOS, 2'FL, 3'SL, 6'SL, inulin, and a gum; GOS, LNnT, 3'SL, 6'SL, inulin, and a gum; 2'FL, LNnT, 3'SL, 6'SL, inulin, and a gum; FOS, 3'SL, 6'SL, inulin, and a gum; GOS, 3'SL, 6'SL, inulin, and a gum; 2'FL, 3'SL, 6'SL, inulin, and a gum; LNnT, 3'SL, 6'SL, inulin, and a gum; FOS,

GOS, 2'FL, LNnT, 3'SL, inulin, and a gum; FOS, GOS, 2'FL, 3'SL, inulin, and a gum; FOS, GOS, LNnT, 3'SL, inulin, and a gum; FOS, 2'FL, LNnT, 3'SL, inulin, and a gum; GOS, 2'FL, LNnT, 3'SL, inulin, and a gum; FOS, GOS, 3'SL, inulin, and a gum; FOS, 2'FL, 3'SL, inulin, and a gum; FOS, LNnT, 3'SL, inulin, and a gum; GOS, 2'FL, 3'SL, inulin, and a gum; GOS, LNnT, 3'SL, inulin, and a gum; 2'FL, LNnT, 3'SL, inulin, and a gum; FOS, 3'SL, inulin, and a gum; GOS, 3'SL, inulin, and a gum; 2'FL, 3'SL, inulin, and a gum; LNnT, 3'SL, inulin, and a gum; FOS, GOS, 2'FL, LNnT, 6'SL, inulin, and a gum; FOS, GOS, 2'FL, 6'SL, inulin, and a gum; FOS, GOS, LNnT, 6'SL, inulin, and a gum; FOS, 2'FL, LNnT, 6'SL, inulin, and a gum; GOS, 2'FL, LNnT, 6'SL, inulin, and a gum; FOS, GOS, 6'SL, inulin, and a gum; FOS, 2'FL, 6'SL, inulin, and a gum; FOS, LNnT, 6'SL, inulin, and a gum; GOS, 2'FL, 6'SL, inulin, and a gum; GOS, LNnT, 6'SL, inulin, and a gum; 2'FL, LNnT, 6'SL, inulin, and a gum; FOS, 6'SL, inulin, and a gum; GOS, 6'SL, inulin, and a gum; 2'FL, 6'SL, inulin, and a gum; LNnT, 6'SL, inulin, and a gum; FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and polydextrose; FOS, GOS, 2'FL, 3'SL, 6'SL, inulin, and polydextrose; FOS, GOS, LNnT, 3'SL, 6'SL, inulin, and polydextrose; FOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and polydextrose; GOS, 2'FL, LNnT, 3'SL, 6'SL, inulin, and polydextrose; FOS, GOS, 3'SL, 6'SL, inulin, and polydextrose; FOS, 2'FL, 3'SL, 6'SL, inulin, and polydextrose; FOS, LNnT, 3'SL, 6'SL, inulin, and polydextrose; GOS, 2'FL, 3'SL, 6'SL, inulin, and polydextrose; GOS, LNnT, 3'SL, 6'SL, inulin, and polydextrose; 2'FL, LNnT, 3'SL, 6'SL, inulin, and polydextrose; FOS, 3'SL, 6'SL, inulin, and polydextrose; GOS, 3'SL, 6'SL, inulin, and polydextrose; 2'FL, 3'SL, 6'SL, inulin, and polydextrose; LNnT, 3'SL, 6'SL, inulin, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, inulin, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, inulin, and polydextrose; GOS, 2'FL, LNnT, 3'SL, inulin, and polydextrose; FOS, GOS, 3'SL, inulin, and polydextrose; FOS, 2'FL, 3'SL, inulin, and polydextrose; FOS, LNnT, 3'SL, inulin, and polydextrose; GOS, 2'FL, 3'SL, inulin, and polydextrose; GOS, LNnT, 3'SL, inulin, and polydextrose; 2'FL, LNnT, 3'SL, inulin, and polydextrose; FOS, 3'SL, inulin, and polydextrose; GOS, 3'SL, inulin, and polydextrose; 2'FL, 3'SL, inulin, and polydextrose; LNnT, 3'SL, inulin, and polydextrose; FOS, GOS, 2'FL, LNnT, 6'SL, inulin, and polydextrose; FOS, GOS, 2'FL, 6'SL, inulin, and polydextrose; FOS, GOS, LNnT, 6'SL, inulin, and polydextrose; FOS, 2'FL, LNnT, 6'SL, inulin, and polydextrose; GOS, 2'FL, LNnT, 6'SL, inulin, and polydextrose; FOS, GOS, 6'SL, inulin, and polydextrose; FOS, GOS, 6'SL, inulin, and polydextrose; FOS,

2'FL, 6'SL, inulin, and polydextrose; FOS, LNnT, 6'SL, inulin, and polydextrose; GOS, 2'FL, 6'SL, inulin, and polydextrose; GOS, LNnT, 6'SL, inulin, and polydextrose; 2'FL, LNnT, 6'SL, inulin, and polydextrose; FOS, 6'SL, inulin, and polydextrose; GOS, 6'SL, inulin, and polydextrose; 2'FL, 6'SL, inulin, and polydextrose; LNnT, 6'SL, inulin, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, a gum, and polydextrose; FOS, GOS, 2'FL, 3'SL, 6'SL, a gum, and polydextrose; FOS, GOS, LNnT, 3'SL, 6'SL, a gum, and polydextrose; FOS, 2'FL, LNnT, 3'SL, 6'SL, a gum, and polydextrose; GOS, 2'FL, LNnT, 3'SL, 6'SL, a gum, and polydextrose; FOS, GOS, 3'SL, 6'SL, a gum, and polydextrose; FOS, 2'FL, 3'SL, 6'SL, a gum, and polydextrose; FOS, LNnT, 3'SL, 6'SL, a gum, and polydextrose; GOS, 2'FL, 3'SL, 6'SL, a gum, and polydextrose; GOS, LNnT, 3'SL, 6'SL, a gum, and polydextrose; 2'FL, LNnT, 3'SL, 6'SL, a gum, and polydextrose; FOS, 3'SL, 6'SL, a gum, and polydextrose; GOS, 3'SL, 6'SL, a gum, and polydextrose; 2'FL, 3'SL, 6'SL, a gum, and polydextrose; LNnT, 3'SL, 6'SL, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, a gum, and polydextrose; FOS, GOS, 2'FL, 3'SL, a gum, and polydextrose; FOS, GOS, LNnT, 3'SL, a gum, and polydextrose; FOS, 2'FL, LNnT, 3'SL, a gum, and polydextrose; GOS, 2'FL, LNnT, 3'SL, a gum, and polydextrose; FOS, GOS, 3'SL, a gum, and polydextrose; FOS, 2'FL, 3'SL, a gum, and polydextrose; FOS, LNnT, 3'SL, a gum, and polydextrose; GOS, 2'FL, 3'SL, a gum, and polydextrose; GOS, LNnT, 3'SL, a gum, and polydextrose; 2'FL, LNnT, 3'SL, a gum, and polydextrose; FOS, 3'SL, a gum, and polydextrose; GOS, 3'SL, a gum, and polydextrose; 2'FL, 3'SL, a gum, and polydextrose; LNnT, 3'SL, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 6'SL, a gum, and polydextrose; FOS, GOS, 2'FL, 6'SL, a gum, and polydextrose; FOS, GOS, LNnT, 6'SL, a gum, and polydextrose; FOS, 2'FL, LNnT, 6'SL, a gum, and polydextrose; GOS, 2'FL, LNnT, 6'SL, a gum, and polydextrose; FOS, GOS, 6'SL, a gum, and polydextrose; FOS, 2'FL, 6'SL, a gum, and polydextrose; FOS, LNnT, 6'SL, a gum, and polydextrose; GOS, 2'FL, 6'SL, a gum, and polydextrose; GOS, LNnT, 6'SL, a gum, and polydextrose; 2'FL, LNnT, 6'SL, a gum, and polydextrose; FOS, 6'SL, a gum, and polydextrose; GOS, 6'SL, a gum, and polydextrose; 2'FL, 6'SL, a gum, and polydextrose; LNnT, 6'SL, a gum, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, and inulin; FOS, GOS, 2'FL, 3'SL, 6'SL, and inulin; FOS, GOS, LNnT, 3'SL, 6'SL, and inulin; FOS, 2'FL, LNnT, 3'SL, 6'SL, and inulin; GOS, 2'FL, LNnT, 3'SL, 6'SL, and inulin; FOS, GOS, 3'SL, 6'SL, and inulin; FOS, 2'FL, 3'SL, 6'SL, and inulin; FOS, LNnT, 3'SL, 6'SL, and inulin; GOS, 2'FL, 3'SL, 6'SL, and inulin; GOS, LNnT, 3'SL, 6'SL, and inulin; 2'FL,

LNnT, 3'SL, 6'SL, and inulin; FOS, 3'SL, 6'SL, and inulin; GOS, 3'SL, 6'SL, and inulin; 2'FL, 3'SL, 6'SL, and inulin; LNnT, 3'SL, 6'SL, and inulin; FOS, GOS, 2'FL, LNnT, 3'SL, and inulin; FOS, GOS, 2'FL, 3'SL, and inulin; FOS, GOS, LNnT, 3'SL, and inulin; FOS, 2'FL, LNnT, 3'SL, and inulin; GOS, 2'FL, LNnT, 3'SL, and inulin; FOS, GOS, 3'SL, and inulin; FOS, 2'FL, 3'SL, and inulin; FOS, LNnT, 3'SL, and inulin; GOS, 2'FL, 3'SL, and inulin; GOS, LNnT, 3'SL, and inulin; 2'FL, LNnT, 3'SL, and inulin; FOS, 3'SL, and inulin; GOS, 3'SL, and inulin; 2'FL, 3'SL, and inulin; LNnT, 3'SL, and inulin; FOS, GOS, 2'FL, LNnT, 6'SL, and inulin; FOS, GOS, 2'FL, 6'SL, and inulin; FOS, GOS, LNnT, 6'SL, and inulin; FOS, 2'FL, LNnT, 6'SL, and inulin; GOS, 2'FL, LNnT, 6'SL, and inulin; FOS, GOS, 6'SL, and inulin; FOS, 2'FL, 6'SL, and inulin; FOS, LNnT, 6'SL, and inulin; GOS, 2'FL, 6'SL, and inulin; GOS, LNnT, 6'SL, and inulin; 2'FL, LNnT, 6'SL, and inulin; FOS, 6'SL, and inulin; GOS, 6'SL, and inulin; FOS, GOS, 2'FL, LNnT, 3'SL, 6'SL, and polydextrose; FOS, GOS, 2'FL, 3'SL, 6'SL, and polydextrose; FOS, GOS, LNnT, 3'SL, 6'SL, and polydextrose; FOS, 2'FL, LNnT, 3'SL, 6'SL, and polydextrose; GOS, 2'FL, LNnT, 3'SL, 6'SL, and polydextrose; FOS, GOS, 3'SL, 6'SL, and polydextrose; FOS, 2'FL, 3'SL, 6'SL, and polydextrose; FOS, LNnT, 3'SL, 6'SL, and polydextrose; GOS, 2'FL, 3'SL, 6'SL, and polydextrose; GOS, LNnT, 3'SL, 6'SL, and polydextrose; 2'FL, LNnT, 3'SL, 6'SL, and polydextrose; FOS, 3'SL, 6'SL, and polydextrose; GOS, 3'SL, 6'SL, and polydextrose; 2'FL, 3'SL, 6'SL, and polydextrose; LNnT, 3'SL, 6'SL, and polydextrose; FOS, GOS, 2'FL, LNnT, 3'SL, and polydextrose; FOS, GOS, 2'FL, 3'SL, and polydextrose; FOS, GOS, LNnT, 3'SL, and polydextrose; FOS, 2'FL, LNnT, 3'SL, and polydextrose; GOS, 2'FL, LNnT, 3'SL, and polydextrose; FOS, GOS, 3'SL, and polydextrose; FOS, 2'FL, 3'SL, and polydextrose; FOS, LNnT, 3'SL, and polydextrose; GOS, 2'FL, 3'SL, and polydextrose; GOS, LNnT, 3'SL, and polydextrose; 2'FL, LNnT, 3'SL, and polydextrose; FOS, 3'SL, and polydextrose; GOS, 3'SL, and polydextrose; 2'FL, 3'SL, and polydextrose; LNnT, 3'SL, and polydextrose; FOS, GOS, 2'FL, LNnT, 6'SL, and polydextrose; FOS, GOS, 2'FL, 6'SL, and polydextrose; FOS, GOS, LNnT, 6'SL, and polydextrose; FOS, 2'FL, LNnT, 6'SL, and polydextrose; GOS, 2'FL, LNnT, 6'SL, and polydextrose; FOS, GOS, 6'SL, and polydextrose; FOS, 2'FL, 6'SL, and polydextrose; FOS, LNnT, 6'SL, and polydextrose; GOS, 2'FL, 6'SL, and polydextrose; GOS, LNnT, 6'SL, and polydextrose; 2'FL, LNnT, 6'SL, and polydextrose; FOS, 6'SL, and polydextrose; GOS, 6'SL, and polydextrose; 2'FL, 6'SL, and polydextrose; LNnT, 6'SL, and polydextrose; FOS, GOS, 2'FL, LNnT,

3'SL, 6'SL, and a gum; FOS, GOS, 2'FL, 3'SL, 6'SL, and a gum; FOS, GOS, LNnT, 3'SL, 6'SL, and a gum; FOS, 2'FL, LNnT, 3'SL, 6'SL, and a gum; GOS, 2'FL, LNnT, 3'SL, 6'SL, and a gum; FOS, GOS, 3'SL, 6'SL, and a gum; FOS, 2'FL, 3'SL, 6'SL, and a gum; FOS, LNnT, 3'SL, 6'SL, and a gum; GOS, 2'FL, 3'SL, 6'SL, and a gum; GOS, LNnT, 3'SL, 6'SL, and a gum; 2'FL, LNnT, 3'SL, 6'SL, and a gum; FOS, 3'SL, 6'SL, and a gum; GOS, 3'SL, 6'SL, and a gum; 2'FL, 3'SL, 6'SL, and a gum; LNnT, 3'SL, 6'SL, and a gum; FOS, GOS, 2'FL, LNnT, 3'SL, and a gum; FOS, GOS, 2'FL, 3'SL, and a gum; FOS, GOS, LNnT, 3'SL, and a gum; FOS, 2'FL, LNnT, 3'SL, and a gum; GOS, 2'FL, LNnT, 3'SL, and a gum; FOS, GOS, 3'SL, and a gum; FOS, 2'FL, 3'SL, and a gum; FOS, LNnT, 3'SL, and a gum; GOS, 2'FL, 3'SL, and a gum; GOS, LNnT, 3'SL, and a gum; 2'FL, LNnT, 3'SL, and a gum; FOS, 3'SL, and a gum; GOS, 3'SL, and a gum; 2'FL, 3'SL, and a gum; LNnT, 3'SL, and a gum; FOS, GOS, 2'FL, LNnT, 6'SL, and a gum; FOS, GOS, 2'FL, 6'SL, and a gum; FOS, GOS, LNnT, 6'SL, and a gum; FOS, 2'FL, LNnT, 6'SL, and a gum; GOS, 2'FL, LNnT, 6'SL, and a gum; FOS, GOS, 6'SL, and a gum; FOS, 2'FL, 6'SL, and a gum; FOS, LNnT, 6'SL, and a gum; GOS, 2'FL, 6'SL, and a gum; GOS, LNnT, 6'SL, and a gum; 2'FL, LNnT, 6'SL, and a gum; FOS, 6'SL, and a gum; GOS, 6'SL, and a gum; 2'FL, 6'SL, and a gum; and LNnT, 6'SL, and a gum.

Probiotics

[0086] The nutritional compositions of the present disclosure may, in addition to HMOs (and, optionally, other prebiotic oligosaccharides as described above), comprise one or more probiotics. In some embodiments, the nutritional composition includes a combination of HMOs and probiotics such that the composition provides a synergistic benefit to the end user in promoting the growth of microbiota in the gastrointestinal tract of infants.

[0087] Probiotics are live microorganisms thought to be healthy for the host organism. Lactic acid bacteria (LAB) and bifidobacteria are the most common types of microbes used as probiotics. Probiotics maintain the microbial ecology of the gut and show physiological, immuno-modulatory and antimicrobial effects, such that the use of probiotics has been found to prevent and treat gastrointestinal diseases and/or disorders, pathogen-induced diarrhea and toxin-producing bacteria, urogenital infections, and atopic diseases.

[0088] In order for microbes to exhibit beneficial probiotic effects *in vivo*, the organisms should survive for extended time periods in the gastrointestinal tract. Therefore, it is important that probiotic strains be selected that possess qualities that prevent their rapid removal by gut contraction. Effective probiotic strains are able to survive gastric conditions and colonize the intestine, at least temporarily, by adhering to the intestinal epithelium.

[0089] Non-limiting examples of probiotic strains for use in the nutritional compositions herein include the genus *Lactobacillus* including *L. acidophilus*, *L. amylovorus*, *L. brevis*, *L. bulgaricus*, *L. casei* spp. *casei*, *L. casei* spp. *rhamnosus*, *L. crispatus*, *L. delbrueckii* ssp. *lactis*, *L. fermentum*, *L. helveticus*, *L. johnsonii*, *L. paracasei*, *L. pentosus*, *L. plantarum*, *L. reuteri*, and *L. sake*; the genus *Bifidobacterium* including: *B. animalis*, *B. bifidum*, *B. breve*, *B. infantis*, and *B. longum*; the genus *Pediococcus* including: *P. acidilactici*; the genus *Propionibacterium* including: *P. acidipropionici*, *P. freudenreichii*, *P. jensenii*, and *P. theonii*; and the genus *Streptococcus* including: *S. cremoris*, *S. lactis*, and *S. thermophilus*. Particularly preferred probiotics include probiotics of human infant origin such as *B. infantis* M-63, *B. infantis* ATCC 15697, *B. infantis* 35624, *B. infantis* CHCC2228, *B. infantis* BB-02, *B. infantis* DSM20088, and *B. infantis* R-0033.

[0090] The probiotic is present in the nutritional compositions in a total amount of at least about 10^3 CFU/g, including from about 10^3 CFU/g to about 10^{12} CFU/g, and including from about 10^6 CFU/g to about 10^7 CFU/g.

[0091] In some embodiments, the nutritional composition includes a probiotic in combination with a first oligosaccharide including fructooligosaccharide and/or a galactooligosaccharide further in combination with a second oligosaccharide including at least one HMO such as 2'FL, 3'FL, 3'SL, 6'SL, and/or LNnT. In these embodiments, the first oligosaccharide and the second oligosaccharide are present in the compositions in a weight ratio of first oligosaccharide:second oligosaccharide of about 10:1, or even from about 11:1 to about 8:1.

Macronutrients

[0092] The nutritional compositions including the HMO or HMOs may be formulated to include at least one of protein, fat, and carbohydrate. In many embodiments, the nutritional compositions will include the HMO or HMOs with protein, carbohydrate and fat.

[0093] Although total concentrations or amounts of the fat, protein, and carbohydrates may vary depending upon the product type (i.e., human milk fortifier, preterm infant formula, infant formula, toddler formula, pediatric formula, follow-on formula, adult nutritional, etc.), product form (i.e., nutritional solid, powder, ready-to-feed liquid, or concentrated liquid), and targeted dietary needs of the intended user, such concentrations or amounts most typically fall within one of the following embodied ranges, inclusive of any other essential fat, protein, and/or carbohydrate ingredients as described herein.

[0094] For the liquid preterm and term infant formulas, carbohydrate concentrations (including both HMOs and any other carbohydrate/oligosaccharide sources) most typically range from about 5% to about 40%, including from about 7% to about 30%, including from about 10% to about 25%, by weight of the preterm or term infant formula; fat concentrations most typically range from about 1% to about 30%, including from about 2% to about 15%, and also including from about 3% to about 10%, by weight of the preterm or term infant formula; and protein concentrations most typically range from about 0.5% to about 30%, including from about 1% to about 15%, and also including from about 2% to about 10%, by weight of the preterm or term infant formula.

[0095] For the liquid human milk fortifiers, carbohydrate concentrations (including both HMOs and any other carbohydrate/oligosaccharide sources) most typically range from about 10% to about 75%, including from about 10% to about 50%, including from about 20% to about 40%, by weight of the human milk fortifier; fat concentrations most typically range from about 10% to about 40%, including from about 15% to about 37%, and also including from about 18% to about 30%, by weight of the human milk fortifier; and protein concentrations most typically range from about 5% to about 40%, including from about 10% to about 30%, and also including from about 15% to about 25%, by weight of the human milk fortifier.

[0096] For the adult nutritional liquids, carbohydrate concentrations (including both HMOs and any other carbohydrate/oligosaccharide sources) most typically range from about 5% to about 40%, including from about 7% to about 30%, including from about 10% to about 25%, by weight of the adult nutritional; fat concentrations most typically range from about 2% to about 30%, including from about 3% to about 15%, and also including from about 5% to about 10%, by weight of the adult nutritional; and protein concentrations most typically range from about 0.5% to about 30%, including from about 1% to about 15%, and also including from about 2% to about 10%, by weight of the adult nutritional.

[0097] The amount of carbohydrates, fats, and/or proteins in any of the liquid nutritional compositions described herein may also be characterized in addition to, or in the alternative, as a percentage of total calories in the liquid nutritional composition as set forth in the following table. These macronutrients for liquid nutritional compositions of the present disclosure are most typically formulated within any of the caloric ranges (embodiments A-F) described in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment A	Embodiment B	Embodiment C
Carbohydrate	0-98	2-96	10-75
Protein	0-98	2-96	5-70
Fat	0-98	2-96	20-85
	Embodiment D	Embodiment E	Embodiment F
Carbohydrate	30-50	25-50	25-50
Protein	15-35	10-30	5-30
Fat	35-55	1-20	2-20

[0098] In one specific example, liquid infant formulas (both ready-to-feed and concentrated liquids) include those embodiments in which the protein component may comprise from about 7.5% to about 25% of the caloric content of the formula; the carbohydrate component (including both HMOs and any other carbohydrate/oligosaccharide sources) may comprise from about 35% to about 50% of the total caloric content of the infant formula; and the fat component may comprise from about

30% to about 60% of the total caloric content of the infant formula. These ranges are provided as examples only, and are not intended to be limiting. Additional suitable ranges are noted in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment G	Embodiment H	Embodiment I
Carbohydrates:	20-85	30-60	35-55
Fat:	5-70	20-60	25-50
Protein:	2-75	5-50	7-40

[0099] When the nutritional composition is a powdered preterm or term infant formula, the protein component is present in an amount of from about 5% to about 35%, including from about 8% to about 12%, and including from about 10% to about 12% by weight of the preterm or term infant formula; the fat component is present in an amount of from about 10% to about 35%, including from about 25% to about 30%, and including from about 26% to about 28% by weight of the preterm or term infant formula; and the carbohydrate component (including both HMOs and any other carbohydrate/oligosaccharide sources) is present in an amount of from about 30% to about 85%, including from about 45% to about 60%, including from about 50% to about 55% by weight of the preterm or term infant formula.

[0100] For powdered human milk fortifiers, the protein component is present in an amount of from about 1% to about 55%, including from about 10% to about 50%, and including from about 10% to about 30% by weight of the human milk fortifier; the fat component is present in an amount of from about 1% to about 30%, including from about 1% to about 25%, and including from about 1% to about 20% by weight of the human milk fortifier; and the carbohydrate component (including both HMOs and any other carbohydrate/oligosaccharide sources) is present in an amount of from about 15% to about 75%, including from about 15% to about 60%, including from about 20% to about 50% by weight of the human milk fortifier.

[0101] For powdered adult nutritionals, the protein component is present in an amount of from about 10% to about 90%, including from about 30% to about 80%, and including from about 40% to about 75% by weight of the adult nutritional; the fat component is present in an amount of from about 0.5% to about 20%, including from about

1% to about 10%, and including from about 2% to about 5% by weight of the adult nutritional; and the carbohydrate component (including both HMOs and any other carbohydrate/oligosaccharide sources) is present in an amount of from about 5% to about 40%, including from about 7% to about 30%, including from about 10% to about 25% by weight of the adult nutritional.

[0102] The total amount or concentration of fat, carbohydrate, and protein, in the powdered nutritional compositions of the present disclosure can vary considerably depending upon the selected composition and dietary or medical needs of the intended user. Additional suitable examples of macronutrient concentrations are set forth below. In this context, the total amount or concentration refers to all fat, carbohydrate, and protein sources in the powdered composition. For powdered nutritional compositions, such total amounts or concentrations are most typically and preferably formulated within any of the embodied ranges described in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment J	Embodiment K	Embodiment L
Carbohydrate	1-85	30-60	35-55
Fat	5-70	20-60	25-50
Protein	2-75	5-50	7-40

Fat

[0103] The nutritional compositions of the present disclosure may optionally comprise any source or sources of fat. Suitable sources of fat for use herein include any fat or fat source that is suitable for use in an oral nutritional composition and is compatible with the essential elements and features of such composition. For example, in one specific embodiment, the fat is derived from long chain polyunsaturated fatty acids (LCPUFAs).

[0104] Exemplary LCPUFAs for use in the nutritional compositions include, for example, ω -3 LCPUFAs and ω -6 LCPUFAs. Specific LCPUFAs include docosahexaenoic acid (DHA), eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), arachidonic acid (ARA), linoleic acid, linolenic acid (alpha linolenic acid) and gamma-linolenic acid derived from oil sources such as plant oils, marine plankton, fungal oils, and fish oils. In

one particular embodiment, the LCPUFAs are derived from fish oils such as menhaden, salmon, anchovy, cod, halibut, tuna, or herring oil. Particularly preferred LCPUFAs for use in the nutritional compositions with the HMOs include DHA, ARA, EPA, DPA, and combinations thereof.

[0105] In order to reduce potential side effects of high dosages of LCPUFAs in the nutritional compositions, the content of LCPUFAs preferably does not exceed 3% by weight of the total fat content, including below 2% by weight of the total fat content, and including below 1% by weight of the total fat content in the nutritional composition.

[0106] The LCPUFA may be provided as free fatty acids, in triglyceride form, in diglyceride form, in monoglyceride form, in phospholipid form, in esterified form or as a mixture of one or more of the above, preferably in triglyceride form. In another specific embodiment, the fat is derived from short chain fatty acids.

[0107] Additional non-limiting examples of suitable fats or sources thereof for use in the nutritional compositions described herein include coconut oil, fractionated coconut oil, soybean oil, corn oil, olive oil, safflower oil, high oleic safflower oil, oleic acids (EMERSOL 6313 OLEIC ACID, Cognis Oleochemicals, Malaysia), MCT oil (medium chain triglycerides), sunflower oil, high oleic sunflower oil, palm and palm kernel oils, palm olein, canola oil, marine oils, fish oils, fungal oils, algae oils, cottonseed oils, and combinations thereof.

Protein

[0108] The nutritional compositions of the present disclosure may optionally further comprise protein. Any protein source that is suitable for use in oral nutritional compositions and is compatible with the essential elements and features of such compositions is suitable for use in the nutritional compositions.

[0109] Non-limiting examples of suitable proteins or sources thereof for use in the nutritional compositions include hydrolyzed, partially hydrolyzed or non-hydrolyzed proteins or protein sources, which may be derived from any known or otherwise suitable source such as milk (e.g., casein, whey), animal (e.g., meat, fish), cereal (e.g., rice, corn), vegetable (e.g., soy) or combinations thereof. Non-limiting examples of such proteins

include milk protein isolates, milk protein concentrates as described herein, casein protein isolates, extensively hydrolyzed casein, whey protein, sodium or calcium caseinates, whole cow milk, partially or completely defatted milk, soy protein isolates, soy protein concentrates, and so forth. In one specific embodiment, the nutritional compositions include a protein source derived from milk proteins of human and/or bovine origin.

[0110] In one embodiment, the protein source is a hydrolyzed protein hydrolysate. In this context, the terms “hydrolyzed protein” or “protein hydrolysates” are used interchangeably herein and include extensively hydrolyzed proteins, wherein the degree of hydrolysis is most often at least about 20%, including from about 20% to about 80%, and also including from about 30% to about 80%, even more preferably from about 40% to about 60%. The degree of hydrolysis is the extent to which peptide bonds are broken by a hydrolysis method. The degree of protein hydrolysis for purposes of characterizing the extensively hydrolyzed protein component of these embodiments is easily determined by one of ordinary skill in the formulation arts by quantifying the amino nitrogen to total nitrogen ratio (AN/TN) of the protein component of the selected liquid formulation. The amino nitrogen component is quantified by USP titration methods for determining amino nitrogen content, while the total nitrogen component is determined by the Tecator Kjeldahl method, all of which are well known methods to one of ordinary skill in the analytical chemistry art.

[0111] Suitable hydrolyzed proteins may include soy protein hydrolysate, casein protein hydrolysate, whey protein hydrolysate, rice protein hydrolysate, potato protein hydrolysate, fish protein hydrolysate, egg albumen hydrolysate, gelatin protein hydrolysate, combinations of animal and vegetable protein hydrolysates, and combinations thereof. Particularly preferred protein hydrolysates include whey protein hydrolysate and hydrolyzed sodium caseinate.

[0112] When used in the nutritional compositions, the protein source may include at least about 20% (by weight total protein) protein hydrolysate, including from about 30% to 100% (by weight total protein) protein hydrolysate, and including from about 40% to about 80% (by weight total protein) protein hydrolysate, and including about 50% (by weight total protein) protein hydrolysate. In one particular embodiment, the nutritional composition includes 100% (by weight total protein) protein hydrolysate.

Carbohydrate

[0113] The nutritional compositions of the present disclosure may further optionally comprise any carbohydrates that are suitable for use in an oral nutritional composition and are compatible with the essential elements and features of such compositions.

[0114] Non-limiting examples of suitable carbohydrates or sources thereof for use in the nutritional compositions described herein may include maltodextrin, hydrolyzed or modified starch or cornstarch, glucose polymers, corn syrup, corn syrup solids, rice-derived carbohydrates, pea-derived carbohydrates, potato-derived carbohydrates, tapioca, sucrose, glucose, fructose, lactose, high fructose corn syrup, honey, sugar alcohols (e.g., maltitol, erythritol, sorbitol), artificial sweeteners (e.g., sucralose, acesulfame potassium, stevia), and combinations thereof. A particularly desirable carbohydrate is a low dextrose equivalent (DE) maltodextrin.

Other Optional Ingredients

[0115] The nutritional compositions of the present disclosure may further comprise other optional components that may modify the physical, chemical, aesthetic or processing characteristics of the compositions or serve as pharmaceutical or additional nutritional components when used in the targeted population. Many such optional ingredients are known or otherwise suitable for use in medical food or other nutritional products or pharmaceutical dosage forms and may also be used in the compositions herein, provided that such optional ingredients are safe for oral administration and are compatible with the essential and other ingredients in the selected product form.

[0116] Non-limiting examples of such optional ingredients include preservatives, emulsifying agents, buffers, pharmaceutical actives, anti-inflammatory agents, additional nutrients as described herein, colorants, flavors, thickening agents and stabilizers, emulsifying agents, lubricants, and so forth.

[0117] The nutritional compositions may further comprise a sweetening agent, preferably including at least one sugar alcohol such as maltitol, erythritol, sorbitol, xylitol, mannitol, isomalt, and lactitol, and also preferably including at least one artificial or high

potency sweetener such as acesulfame K, aspartame, sucralose, saccharin, stevia, and tagatose. These sweetening agents, especially as a combination of a sugar alcohol and an artificial sweetener, are especially useful in formulating liquid beverage embodiments of the present disclosure having a desirable favor profile. These sweetener combinations are especially effective in masking undesirable flavors sometimes associated with the addition of vegetable proteins to a liquid beverage. Optional sugar alcohol concentrations in the nutritional composition may range from at least 0.01%, including from 0.1% to about 10%, and also including from about 1% to about 6%, by weight of the nutritional composition. Optional artificial sweetener concentrations may range from about 0.01%, including from about 0.05% to about 5%, also including from about 0.1% to about 1.0%, by weight of the nutritional composition.

[0118] A flowing agent or anti-caking agent may be included in the nutritional compositions as described herein to retard clumping or caking of the powder over time and to make a powder embodiment flow easily from its container. Any known flowing or anti-caking agents that are known or otherwise suitable for use in a nutritional powder or product form are suitable for use herein, non-limiting examples of which include tricalcium phosphate, silicates, and combinations thereof. The concentration of the flowing agent or anti-caking agent in the nutritional composition varies depending upon the product form, the other selected ingredients, the desired flow properties, and so forth, but most typically range from about 0.1% to about 4%, including from about 0.5% to about 2%, by weight of the nutritional composition.

[0119] A stabilizer may also be included in the nutritional compositions. Any stabilizer that is known or otherwise suitable for use in a nutritional composition is also suitable for use herein, some non-limiting examples of which include gums such as xanthan gum. The stabilizer may represent from about 0.1% to about 5.0%, including from about 0.5% to about 3%, including from about 0.7% to about 1.5%, by weight of the nutritional composition.

[0120] Additionally, the nutritional compositions may comprise one or more antioxidants to provide nutritional support, as well as to reduce oxidative stress. Any antioxidants suitable for oral administration may be included for use in the nutritional

compositions of the present disclosure, including, for example, vitamin A, vitamin E, vitamin C, retinol, tocopherol, and carotenoids.

[0121] In one specific embodiment, the antioxidants for use in the nutritional compositions include carotenoids such as lutein, zeaxanthin, lycopene, beta-carotene, and combinations thereof, and particularly, combinations of the carotenoids lutein, lycopene, and beta-carotene. Nutritional compositions containing these combinations, as selected and defined herein, can be used to modulate inflammation and/or levels of C-reactive protein in preterm and term infants.

[0122] The nutritional compositions may further comprise any of a variety of other vitamins or related nutrients, non-limiting examples of which include vitamin D, vitamin K, thiamine, riboflavin, pyridoxine, vitamin B₁₂, niacin, folic acid, pantothenic acid, biotin, choline, inositol, salts and derivatives thereof, and combinations thereof.

[0123] The nutritional compositions may further comprise any of a variety of other additional minerals, non-limiting examples of which include calcium, phosphorus, magnesium, iron, zinc, manganese, copper, sodium, potassium, molybdenum, chromium, chloride, and combinations thereof.

[0124] The nutritional compositions of the present disclosure may additionally comprise nucleotides and/or nucleotide precursors selected from the group consisting of nucleoside, purine base, pyrimidine base, ribose and deoxyribose to further improve intestinal barrier integrity and/or maturation. The nucleotide may be in monophosphate, diphosphate, or triphosphate form. The nucleotide may be a ribonucleotide or a deoxyribonucleotide. The nucleotides may be monomeric, dimeric, or polymeric (including RNA and DNA). The nucleotide may be present in the nutritional composition as a free acid or in the form of a salt, preferably a monosodium salt.

[0125] Suitable nucleotides and/or nucleosides for use in the nutritional compositions include one or more of cytidine 5'-monophosphate, uridine 5'-monophosphate, adenosine 5'-monophosphate, guanosine 5'-1-monophosphate, and/or inosine 5'-monophosphate, more preferably cytidine 5'-monophosphate, uridine 5'-monophosphate, adenosine 5'-monophosphate, guanosine 5'-monophosphate, and inosine 5'-monophosphate.

Methods of Manufacture

[0126] The nutritional compositions of the present disclosure may be prepared by any known or otherwise effective manufacturing technique for preparing the selected product solid or liquid form. Many such techniques are known for any given product form such as nutritional liquids or powders and can easily be applied by one of ordinary skill in the art to the nutritional compositions described herein.

[0127] The nutritional compositions of the present disclosure can therefore be prepared by any of a variety of known or otherwise effective formulation or manufacturing methods. In one suitable manufacturing process, for example, at least three separate slurries are prepared, including a protein-in-fat (PIF) slurry, a carbohydrate-mineral (CHO-MIN) slurry, and a protein-in-water (PIW) slurry. The PIF slurry is formed by heating and mixing the oil (e.g., canola oil, corn oil, etc.) and then adding an emulsifier (e.g., lecithin), fat soluble vitamins, and a portion of the total protein (e.g., milk protein concentrate, etc.) with continued heat and agitation. The CHO-MIN slurry is formed by adding with heated agitation to water: minerals (e.g., potassium citrate, dipotassium phosphate, sodium citrate, etc.), trace and ultra trace minerals (TM/UTM premix), thickening or suspending agents (e.g. avicel, gellan, carrageenan). The resulting CHO-MIN slurry is held for 10 minutes with continued heat and agitation before adding additional minerals (e.g., potassium chloride, magnesium carbonate, potassium iodide, etc.), and/or carbohydrates (e.g., HMOs, fructooligosaccharide, sucrose, corn syrup, etc.). The PIW slurry is then formed by mixing with heat and agitation the remaining protein, if any.

[0128] The resulting slurries are then blended together with heated agitation and the pH adjusted to 6.6-7.0, after which the composition is subjected to high-temperature short-time (HTST) processing during which the composition is heat treated, emulsified and homogenized, and then allowed to cool. Water soluble vitamins and ascorbic acid are added, the pH is adjusted to the desired range if necessary, flavors are added, and water is added to achieve the desired total solid level. The composition is then aseptically packaged to form an aseptically packaged nutritional emulsion. This emulsion can then be further diluted, heat-treated, and packaged to form a ready-to-feed or concentrated liquid, or it can be heat-treated and subsequently processed and packaged as a reconstitutable powder, e.g., spray dried, drymixed, agglomerated.

[0129] The nutritional solid, such as a spray dried nutritional powder or drymixed nutritional powder, may be prepared by any collection of known or otherwise effective techniques, suitable for making and formulating a nutritional powder.

[0130] For example, when the nutritional powder is a spray dried nutritional powder, the spray drying step may likewise include any spray drying technique that is known for or otherwise suitable for use in the production of nutritional powders. Many different spray drying methods and techniques are known for use in the nutrition field, all of which are suitable for use in the manufacture of the spray dried nutritional powders herein.

[0131] One method of preparing the spray dried nutritional powder comprises forming and homogenizing an aqueous slurry or liquid comprising predigested fat, and optionally protein, carbohydrate, and other sources of fat, and then spray drying the slurry or liquid to produce a spray dried nutritional powder. The method may further comprise the step of spray drying, drymixing, or otherwise adding additional nutritional ingredients, including any one or more of the ingredients described herein, to the spray dried nutritional powder.

[0132] Other suitable methods for making nutritional compositions are described, for example, in U.S. Pat. No. 6,365,218 (Borschel, et al.), U.S. Patent No. 6,589,576 (Borschel, et al.), U.S. Pat. No. 6,306,908 (Carlson, et al.), U.S. Patent Application No. 20030118703 A1 (Nguyen, et al.), which descriptions are incorporated herein by reference to the extent that they are consistent herewith.

Methods of Use

[0133] The nutritional compositions as described herein can be used to address one or more of the diseases, disorders, or conditions discussed herein, or can be used to provide one or more of the benefits described herein, to preterm infants, infants, toddlers, children, and adults, including pregnant women. The preterm infant, infant, toddler, child, adult and pregnant women utilizing the nutritional compositions described herein may actually have or be afflicted with the disease or condition described, or may be susceptible to, or at risk of, getting the disease or condition (that is, may not actually yet have the disease or condition, but is at elevated risk as compared to the general population for

getting it due to certain conditions, family history, etc.) Whether the preterm infant, infant, toddler, child, adult, and pregnant women actually have the disease or condition, or is at risk or susceptible to the disease or condition, the preterm infant, infant, toddler, child, adult, and pregnant women are classified herein as “in need of” assistance in dealing with and combating the disease or condition. For example, the preterm infant, infant, toddler, child, adult and pregnant women may actually have respiratory inflammation or may be at risk of getting respiratory inflammation (susceptible to getting respiratory inflammation) due to family history or other medical conditions, for example. Whether the preterm infant, infant, toddler, child, adult, and pregnant women actually has the disease or condition, or is only at risk or susceptible to getting the disease or condition, it is within the scope of the present disclosure to assist the preterm infant, infant, toddler, child, adult and pregnant women with the nutritional compositions described herein.

[0134] Based on the foregoing, because some of the method embodiments of the present disclosure are directed to specific subsets or subclasses of identified individuals (that is, the subset or subclass of individuals “in need” of assistance in addressing one or more specific diseases or specific conditions noted herein), not all preterm infants, infants, toddlers, children, adults and pregnant women will fall within the subset or subclass of preterm infants, infants, toddlers, children, adults, and pregnant women as described herein for certain diseases or conditions.

[0135] The nutritional compositions as described herein comprise HMOs, alone or in combination with one or more additional components, to provide a nutritional source for improving at least the intestinal/gut function. Specifically, the nutritional compositions can stimulate enteric nerve cells in the gastrointestinal tract of an individual to improve intestinal/gut barrier integrity; improve feeding tolerance (e.g., reduced diarrhea, loose stools, gas, and bloating); reduce colic in infants; protect against necrotizing enterocolitis and other disorders of prematurity; address gastrointestinal diseases and disorders associated with the enteric nervous system; address gastrointestinal diseases and disorders of gut contractility and inflammation; correct effects of gut dysbiosis; and affect long-term modulation of allergic tolerance.

[0136] More particularly, in some embodiments, the nutritional compositions may be administered to an individual having, susceptible to, or at risk of, gastrointestinal diseases and disorders associated with the enteric nervous system and/or associated with gut contractility and inflammation, which may include, for example, irritable bowel syndrome, colitis (e.g., necrotizing enterocolitis, Crohn's disease, ischemic colitis, cryptosporidium enterocolitis, pseudomembranous colitis, cytomegalovirus, ulcerative colitis), food intolerance, and food allergies.

[0137] Along with improved growth and maturation of an individual's immune system as described above, the use of the nutritional compositions of the present disclosure may also function to enhance the individual's ability to resist microbial infection and to promote the growth of beneficial microbiota in the gastrointestinal tract of an infant, toddler, child, or adult.

[0138] Additionally, the nutritional compositions of the present disclosure may also be used to improve cognition in individuals, particularly in individuals susceptible to, or at risk of, neurodegenerative diseases, which may include, for example, Alzheimer's disease, Huntington's disease, Parkinson's disease, and schizophrenia, or in individuals suffering from conditions caused by impaired cognitive development or neurodevelopmental conditions, such as attention deficit hyperactivity disorder and autism.

EXAMPLES

[0139] The following examples illustrate specific embodiments and/or features of the nutritional compositions and methods of the present disclosure. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present disclosure, as many variations thereof are possible without departing from the spirit and scope of the disclosure. All exemplified amounts are weight percentages based upon the total weight of the composition, unless otherwise specified.

[0140] The exemplified compositions are shelf stable nutritional compositions prepared in accordance with the manufacturing methods described herein, such that each exemplified composition, unless otherwise specified, includes an aseptically processed embodiment and a retort packaged embodiment.

[0141] The nutritional liquid embodiments are aqueous oil-in-water emulsions that are packaged in 240 mL plastic containers and remain physically stable for 12-18 months after composition/packaging at storage temperatures ranging from 1-25°C.

EXAMPLES 1-5

[0142] Examples 1-5 illustrate ready-to-feed nutritional emulsions of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	86.64	86.64	86.64	86.64	86.64
Lactose	54.80	54.80	54.80	54.80	54.80
High oleic safflower oil	14.10	14.10	14.10	14.10	14.10
Soybean oil	10.6	10.6	10.6	10.6	10.6
Coconut oil	10.1	10.1	10.1	10.1	10.1
2' fucosyllactose (2'FL)	0.1896	0.1801	0.1706	0.1991	0.2086
Galactooligosaccharides (GOS)	8.630	8.630	8.630	8.630	8.630
Whey protein concentrate	6.40	6.40	6.40	6.40	6.40
Potassium citrate	478.9 g	478.9 g	478.9 g	478.9 g	478.9 g
Calcium carbonate	448.28 g	448.28 g	448.28 g	448.28 g	448.28 g
Soy lecithin	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
Stabilizer	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
ARA oil	368.01 g	368.01 g	368.01 g	368.01 g	368.01 g
Nucleotide/chloride premix	293.26 g	293.26 g	293.26 g	293.26 g	293.26 g
Potassium chloride	226.45 g	226.45 g	226.45 g	226.45 g	226.45 g
Ascorbic acid	445.94 g	445.94 g	445.94 g	445.94 g	445.94 g
Vitamin mineral premix	142.88 g	142.88 g	142.88 g	142.88 g	142.88 g
DHA oil	137.8 g	137.8 g	137.8 g	137.8 g	137.8 g
Carrageenan	180.0 g	180.0 g	180.0 g	180.0 g	180.0 g
Magnesium chloride	55.0 g	55.0 g	55.0 g	55.0 g	55.0 g
Ferrous sulfate	58.0 g	58.0 g	58.0 g	58.0 g	58.0 g
Choline chloride	53.9 g	53.9 g	53.9 g	53.9 g	53.9 g
Vitamin A, D ₃ , E, K ₁ premix	47.40 g	47.40 g	47.40 g	47.40 g	47.40 g
Citric acid	29.77 g	29.77 g	29.77 g	29.77 g	29.77 g
Probiotic	1.0	1.0	1.0	1.0	1.0
Mixed carotenoid premix	26.40 g	26.40 g	26.40 g	26.40 g	26.40 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	3.31 g	3.31 g	3.31 g	3.31 g	3.31 g
Tricalcium phosphate	15.65 g	15.65 g	15.65 g	15.65 g	15.65 g
Potassium phosphate monobasic	13.67 g	13.67 g	13.67 g	13.67 g	13.67 g
Riboflavin	2.42 g	2.42 g	2.42 g	2.42 g	2.42 g
Potassium hydroxide	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 6-10

[0143] Examples 6-10 illustrate ready-to-feed nutritional emulsions of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 6	Ex. 7	Ex. 8	Ex. 9	Ex. 10
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	86.64	86.64	86.64	86.64	86.64
Lactose	54.80	54.80	54.80	54.80	54.80
High oleic safflower oil	14.10	14.10	14.10	14.10	14.10
Soybean oil	10.6	10.6	10.6	10.6	10.6
Coconut oil	10.1	10.1	10.1	10.1	10.1
2' fucosyllactose (2'FL)	0.0948	0.09005	0.0853	0.0995	0.1043
Lacto-N-neotetraose (LNnT)	0.0948	0.09005	0.0853	0.0995	0.1043
Galactooligosaccharides (GOS)	8.630	8.630	8.630	8.630	8.630
Whey protein concentrate	6.40	6.40	6.40	6.40	6.40
Potassium citrate	478.9 g	478.9 g	478.9 g	478.9 g	478.9 g
Calcium carbonate	448.28 g	448.28 g	448.28 g	448.28 g	448.28 g
Soy lecithin	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
Stabilizer	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
ARA oil	368.01 g	368.01 g	368.01 g	368.01 g	368.01 g
Nucleotide/chloride premix	293.26 g	293.26 g	293.26 g	293.26 g	293.26 g
Potassium chloride	226.45 g	226.45 g	226.45 g	226.45 g	226.45 g
Ascorbic acid	445.94 g	445.94 g	445.94 g	445.94 g	445.94 g
Vitamin mineral premix	142.88 g	142.88 g	142.88 g	142.88 g	142.88 g
DHA oil	137.8 g	137.8 g	137.8 g	137.8 g	137.8 g
Carrageenan	180.0 g	180.0 g	180.0 g	180.0 g	180.0 g
Magnesium chloride	55.0 g	55.0 g	55.0 g	55.0 g	55.0 g
Ferrous sulfate	58.0 g	58.0 g	58.0 g	58.0 g	58.0 g
Choline chloride	53.9 g	53.9 g	53.9 g	53.9 g	53.9 g
Vitamin A, D ₃ , E, K ₁ premix	47.40 g	47.40 g	47.40 g	47.40 g	47.40 g
Citric acid	29.77 g	29.77 g	29.77 g	29.77 g	29.77 g
Probiotic	1.0	0.95	0.90	1.05	1.10
Mixed carotenoid premix	26.40 g	26.40 g	26.40 g	26.40 g	26.40 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	3.31 g	3.31 g	3.31 g	3.31 g	3.31 g
Tricalcium phosphate	15.65 g	15.65 g	15.65 g	15.65 g	15.65 g
Potassium phosphate monobasic	13.67 g	13.67 g	13.67 g	13.67 g	13.67 g
Riboflavin	2.42 g	2.42 g	2.42 g	2.42 g	2.42 g
Potassium hydroxide	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 11-15

[0144] Examples 11-15 illustrate concentrated liquid emulsions of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 11	Ex. 12	Ex. 13	Ex. 14	Ex. 15
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	166.6	166.6	166.6	166.6	166.6
Lactose	106.1	106.1	106.1	106.1	106.1
High oleic safflower oil	27.16	27.16	27.16	27.16	27.16
Soybean oil	20.42	20.42	20.42	20.42	20.42
Coconut oil	19.48	19.48	19.48	19.48	19.48
2' fucosyllactose (2'FL)	0.1896	0.1188	0.0853	0.2414	0.2560
Galactooligosaccharides (GOS)	16.71	16.71	16.71	16.71	16.71
Whey protein concentrate	12.20	12.20	12.20	12.20	12.20
Potassium citrate	894.5 g	894.5 g	894.5 g	894.5 g	894.5 g
Calcium carbonate	1.072	1.072	1.072	1.072	1.072
Monoglycerides	690.0 g	690.0 g	690.0 g	690.0 g	690.0 g
Soy lecithin	690.0 g	690.0 g	690.0 g	690.0 g	690.0 g
ARA oil	684.2 g	684.2 g	684.2 g	684.2 g	684.2 g
Nucleotide/chloride premix	568.9 g	568.9 g	568.9 g	568.9 g	568.9 g
Potassium chloride	480.8 g	480.8 g	480.8 g	480.8 g	480.8 g
Ascorbic acid	958.6 g	958.6 g	958.6 g	958.6 g	958.6 g
Vitamin mineral premix	276.9 g	276.9 g	276.9 g	276.9 g	276.9 g
DHA oil	256.1 g	256.1 g	256.1 g	256.1 g	256.1 g
Carrageenan	200.0 g	200.0 g	200.0 g	200.0 g	200.0 g
Magnesium chloride	174.7 g	174.7 g	174.7 g	174.7 g	174.7 g
Ferrous sulfate	112.7 g	112.7 g	112.7 g	112.7 g	112.7 g
Choline chloride	104.8 g	104.8 g	104.8 g	104.8 g	104.8 g
Vitamin A, D ₃ , E, K ₁ premix	86.90 g	86.90 g	86.90 g	86.90 g	86.90 g
Citric acid	64.55 g	64.55 g	64.55 g	64.55 g	64.55 g
Mixed carotenoid premix	45.63 g	45.63 g	45.63 g	45.63 g	45.63 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	6.371 g	6.371 g	6.371 g	6.371 g	6.371 g
Riboflavin	2.921 g	2.921 g	2.921 g	2.921 g	2.921 g
Vitamin A Palmitate	1.504 g	1.504 g	1.504 g	1.504 g	1.504 g
Potassium hydroxide	659.8 g	659.8 g	659.8 g	659.8 g	659.8 g
Tricalcium phosphate	AN	AN	AN	AN	AN
Potassium phosphate monobasic	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 16-20

[0145] Examples 16-20 illustrate spray dried nutritional powders of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 16	Ex. 17	Ex. 18	Ex. 19	Ex. 20
Nonfat dry milk	456.9	456.9	456.9	456.9	456.9
Lactose	259.0	259.0	259.0	259.0	259.0
High oleic sunflower oil	93.9	93.9	93.9	93.9	93.9
Soy oil	70.4	70.4	70.4	70.4	70.4
Coconut oil	67.1	67.1	67.1	67.1	67.1
2' fucosyllactose (2'FL)	0.7584	0.7204	0.6824	0.7964	0.8344
Galactooligosaccharide (GOS)	53.5	53.5	53.5	53.5	53.5

Probiotic	1.0	0.95	0.90	1.05	1.10
Flavoring agent	6.2	6.2	6.2	6.2	6.2
Calcium carbonate	4.8	4.8	4.8	4.8	4.8
Potassium citrate	4.7	4.7	4.7	4.7	4.7
Oligofructose (FOS)	2.9	2.9	2.9	2.9	2.9
Ascorbic acid	2.0	2.0	2.0	2.0	2.0
Nucleotide/Choline Premix	1.8	1.8	1.8	1.8	1.8
ARA oil	1.8	1.8	1.8	1.8	1.8
Vitamin/Trace Mineral Premix	1.5	1.5	1.5	1.5	1.5
Sodium chloride	1.3	1.3	1.3	1.3	1.3
Lecithin	1.2	1.2	1.2	1.2	1.2
Sodium citrate	982.2 g	982.2 g	982.2 g	982.2 g	982.2 g
DHA oil	882.1 g	882.1 g	882.1 g	882.1 g	882.1 g
Magnesium chloride	477.4 g	477.4 g	477.4 g	477.4 g	477.4 g
Vitamin A, D3, E, K1 Premix	314.7 g	314.7 g	314.7 g	314.7 g	314.7 g
Ascorbyl Palmitate	278.8 g	278.8 g	278.8 g	278.8 g	278.8 g
Antioxidant	137.3 g	137.3 g	137.3 g	137.3 g	137.3 g
Tocopheryl acetate	32.0 g	32.0 g	32.0 g	32.0 g	32.0 g
Beta-carotene 30%	11.0 g	11.0 g	11.0 g	11.0 g	11.0 g
Potassium iodide	2.5 g	2.5 g	2.5 g	2.5 g	2.5 g
Riboflavin	2.0 g	2.0 g	2.0 g	2.0 g	2.0 g
Magnesium sulfate	499.5 mg	499.5 mg	499.5 mg	499.5 mg	499.5 mg
Potassium phosphate dibasic	AN	AN	AN	AN	AN
Potassium chloride	AN	AN	AN	AN	AN
Tricalcium phosphate	AN	AN	AN	AN	AN
Potassium hydroxide	AN	AN	AN	AN	AN
Calcium hydroxide	AN	AN	AN	AN	AN
Sodium hydroxide	AN	AN	AN	AN	AN
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.

AN = as needed

EXAMPLES 21-25

[0146] Examples 21-25 illustrate spray dried nutritional powders of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 21	Ex. 22	Ex. 23	Ex. 24	Ex. 25
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Corn syrup	308.9	308.9	308.9	308.9	308.9
Maltodextrin	297.1	297.1	297.1	297.1	297.1
Sucrose	112.4	112.4	112.4	112.4	112.4
High Oleic sunflower oil	84.9	84.9	84.9	84.9	84.9
Sodium caseinate	73.0	73.0	73.0	73.0	73.0
Calcium caseinate	50.2	50.2	50.2	50.2	50.2
2' fucosyllactose (2'FL)	0.7584	0.7204	0.6824	0.7964	0.8344
Inulin, oligofructose	47.0	47.0	47.0	47.0	47.0
Soy oil	38.3	38.3	38.3	38.3	38.3
Isolated soy protein	35.9	35.9	35.9	35.9	35.9
Milk protein isolate	16.3	16.3	16.3	16.3	16.3
Canola oil	13.7	13.7	13.7	13.7	13.7
Sodium citrate	9.8	9.8	9.8	9.8	9.8

Potassium citrate	9.7	9.7	9.7	9.7	9.7
Tricalcium phosphate	9.0	9.0	9.0	9.0	9.0
Flavoring agent	7.3	7.3	7.3	7.3	7.3
Magnesium chloride	6.2	6.2	6.2	6.2	6.2
Potassium chloride	5.5	5.5	5.5	5.5	5.5
Choline chloride	1.7	1.7	1.7	1.7	1.7
Vitamin premix	950.0 g	950.0 g	950.0 g	950.0 g	950.0 g
Ascorbic acid	755.0 g	755.0 g	755.0 g	755.0 g	755.0 g
Vitamin/trace mineral premix	465.0 g	465.0 g	465.0 g	465.0 g	465.0 g
Potassium hydroxide	215.9 g	215.9 g	215.9 g	215.9 g	215.9 g
Potassium phosphate dibasic	185.8 g	185.8 g	185.8 g	185.8 g	185.8 g
Ascorbyl palmitate	164.7 g	164.7 g	164.7 g	164.7 g	164.7 g
Antioxidant	82.3 g	82.3 g	82.3 g	82.3 g	82.3 g
Vitamin A, D3, E, K1 premix	82.3 g	82.3 g	82.3 g	82.3 g	82.3 g
Vitamin A palmitate	16.5 g	16.5 g	16.5 g	16.5 g	16.5 g
Ferrous sulfate	12.0 g	12.0 g	12.0 g	12.0 g	12.0 g
Beta carotene 30%	5.5 g	5.5 g	5.5 g	5.5 g	5.5 g
Vitamin D3 oil	1.0 g	1.0 g	1.0 g	1.0 g	1.0 g
Potassium iodide	800.0 mg	800.0 mg	800.0 mg	800.0 mg	800.0 mg
Citric acid	AN	AN	AN	AN	AN
Potassium hydroxide 40%	AN	AN	AN	AN	AN
Maltodextrin	AN	AN	AN	AN	AN
Magnesium sulfate	AN	AN	AN	AN	AN
Sodium chloride	AN	AN	AN	AN	AN
Calcium carbonate	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 26-30

[0147] Examples 26-30 illustrate ready-to-feed nutritional emulsions of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 26	Ex. 27	Ex. 28	Ex. 29	Ex. 30
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	86.64	86.64	86.64	86.64	86.64
Lactose	54.80	54.80	54.80	54.80	54.80
High oleic safflower oil	14.10	14.10	14.10	14.10	14.10
Soybean oil	10.6	10.6	10.6	10.6	10.6
Coconut oil	10.1	10.1	10.1	10.1	10.1
2' fucosyllactose (2'FL)	0.0948	0.09005	0.0853	0.0995	0.1043
6'-sialyllactose (6'SL)	0.0948	0.09005	0.0853	0.0995	0.1043
Galactooligosaccharides (GOS)	8.630	8.630	8.630	8.630	8.630
Whey protein concentrate	6.40	6.40	6.40	6.40	6.40
Potassium citrate	478.9 g	478.9 g	478.9 g	478.9 g	478.9 g
Calcium carbonate	448.28 g	448.28 g	448.28 g	448.28 g	448.28 g
Soy lecithin	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
Stabilizer	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
ARA oil	368.01 g	368.01 g	368.01 g	368.01 g	368.01 g
Nucleotide/chloride premix	293.26 g	293.26 g	293.26 g	293.26 g	293.26 g
Potassium chloride	226.45 g	226.45 g	226.45 g	226.45 g	226.45 g
Ascorbic acid	445.94 g	445.94 g	445.94 g	445.94 g	445.94 g

Vitamin mineral premix	142.88 g	142.88 g	142.88 g	142.88 g	142.88 g
DHA oil	137.8 g	137.8 g	137.8 g	137.8 g	137.8 g
Carrageenan	180.0 g	180.0 g	180.0 g	180.0 g	180.0 g
Magnesium chloride	55.0 g	55.0 g	55.0 g	55.0 g	55.0 g
Ferrous sulfate	58.0 g	58.0 g	58.0 g	58.0 g	58.0 g
Choline chloride	53.9 g	53.9 g	53.9 g	53.9 g	53.9 g
Vitamin A, D ₃ , E, K ₁ premix	47.40 g	47.40 g	47.40 g	47.40 g	47.40 g
Citric acid	29.77 g	29.77 g	29.77 g	29.77 g	29.77 g
Probiotic	1.0	0.95	0.90	1.05	1.10
Mixed carotenoid premix	26.40 g	26.40 g	26.40 g	26.40 g	26.40 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	3.31 g	3.31 g	3.31 g	3.31 g	3.31 g
Tricalcium phosphate	15.65 g	15.65 g	15.65 g	15.65 g	15.65 g
Potassium phosphate monobasic	13.67 g	13.67 g	13.67 g	13.67 g	13.67 g
Riboflavin	2.42 g	2.42 g	2.42 g	2.42 g	2.42 g
Potassium hydroxide	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 31-34

[0148] Examples 31-34 illustrate concentrated liquid human milk fortifiers of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient (Per 1000 Kg)	Ex. 31	Ex. 32	Ex. 33	Ex. 34
Water	Q.S.	Q.S.	Q.S.	Q.S.
Casein Hydrolysate	108	108	125	150
Maltodextrin	104	104	104	104
MCT Oil	17.3	17.3	17.3	17.3
Tricalcium Phosphate	16.0	16.0	16.0	16.0
Soy Oil	10.4	10.4	10.4	10.4
6' sialyllactose (6'SL)	0.0948	0.09005	0.0853	0.0995
Lacto-N-neotetraose (LNnT)	0.0948	0.09005	0.0853	0.0995
Galactooligosaccharides (GOS)	6.7704	6.7704	6.7704	6.7704
Gum Arabic	12.0	10.0	15.0	2.031
Starch	12.0	10.0	35.0	6.0
Coconut Oil	6.3	6.3	6.3	6.3
Potassium Citrate	6.9	6.9	6.9	6.9
Ascorbic Acid	2.9	2.9	2.9	2.9
Magnesium Chloride	4.0	4.0	4.0	4.0
ARA oil	2.6	2.6	2.6	2.6
Leucine	1.8	1.8	1.8	1.8
DHA oil	2.1	2.1	2.1	2.1
Potassium Chloride	1.1	1.1	1.1	1.1
Tyrosine	1.4	1.4	1.4	1.4
Monoglycerides	800 g	800 g	800 g	800 g
Mixed Carotenoid Premix	551 g	551 g	551 g	551 g
M-Inositol	529 g	529 g	529 g	529 g

Sodium Chloride	861 g	861 g	861 g	861 g
L-Carnitine	221 g	221 g	221 g	221 g
Tryptophan	331 g	331 g	331 g	331 g
Zinc Sulfate	309 g	309 g	309 g	309 g
Niacinamide	320 g	320 g	320 g	320 g
Tocopheryl Acetate	364 g	364 g	364 g	364 g
Gellan Gum	200 g	300 g	400 g	600 g
Ferrous Sulfate	106 g	106 g	106 g	106 g
Choline Chloride	353 g	353 g	353 g	353 g
Calcium Pantothenate	132 g	132 g	132 g	132 g
Vitamin A Palmitate	77 g	77 g	77 g	77 g
Riboflavin	33 g	33 g	33 g	33 g
Vitamin D3	13 g	13 g	13 g	13 g
Copper Sulfate	18 g	18 g	18 g	18 g
Pyridoxine Hydrochloride	20 g	20 g	20 g	20 g
Thiamin Hydrochloride	24 g	24 g	24 g	24 g
Folic Acid	3.3 g	3.3 g	3.3 g	3.3 g
Biotin	2.5 g	2.5 g	2.5 g	2.5 g
Manganese Sulfate	1.8 g	1.8 g	1.8 g	1.8 g
Phylloquinone	880 mg	880 mg	880 mg	880 mg
Sodium Selenate	90 mg	90 mg	90 mg	90 mg
Cyanocobalamin	88 mg	88 mg	88 mg	88 mg
Potassium Hydroxide	Q.S.	Q.S.	Q.S.	Q.S.

EXAMPLE 35

[0149] In this Example, the effect of 2'-fucosyllactose (2'FL) OR 3'-fucosyllactose (3'FL) on stimulating enteric nerve cells in the gastrointestinal tract of rodents is analyzed.

[0150] Specifically, a peristalsis model using luminally perfused mouse colon is used to test the stimulation effect of 2'FL or 3'FL on enteric nerve cells. Colon muscle is perfused with 2'FL or 3'FL, at concentrations of 1 mg/mL, 0.5 mg/mL, and 0.1 mg/mL, for 15 minutes. The frequency and amplitude of contractions of the muscle is analyzed. The results are shown in FIG. 1.

[0151] As shown in the results, there is a direct stimulation of nerve cells by 2'FL or 3'FL without involving gut microbiota and/or their metabolites. Specifically, the frequency and amplitude of contraction are reduced consistently and in a dose response fashion.

EXAMPLE 36

[0152] In this Example, the fermentation rates of various non-digestible carbohydrates are measured.

[0153] Inclusion/exclusion criteria for choosing eight infant participants include: the infant was full term at birth with a gestational age of 38 to 42 weeks; the infant was at or above the fifth percentile for weight at birth; the infant has no maternal medical history of diabetes, tuberculosis, or perinatal infection with proven adverse effects on the fetus; was a vaginal birth; was at least 2 months of age at study entry, but not older than 4 months of age; has no known cardiac, respiratory, gastrointestinal, or other systemic disease such as urinary tract infection or otitis media; is free of history of blood group incompatibility serious enough to result in hematological problems; and is not receiving any medications (except for supplemental vitamins) and has never received antibiotics. The eight infants are allowed to consume their normal diet of breast milk or infant formula. Four infants are exclusively breast fed and four infants are exclusively formula fed one of four commercially available infant formulas.

[0154] On the day of the *in vitro* experiments, a fecal sample is collected in the diaper and prepped within 15 minutes of defecation. For prepping, the sample is placed in a container with tepid water and analyzed. Fecal samples are diluted 1:10 (wt/vol) in anaerobic dilution solution by blending for 15 seconds in a Waring blender under a stream of CO₂. Blended, diluted feces are filtered through four layers of cheesecloth and sealed in 125-mL serum bottles under CO₂. Inoculum is stored at 37°C until inoculation of *in vitro* tubes.

[0155] Oligosaccharide substrates suitable for growing the bacterium include galactooligosaccharides (GOS) 95 (GOS; Inalco Pharmaceuticals, San Luis, California), α -(2-6')-N-Acetylneuraminyllactose sodium salt (6'SL; Inalco Pharmaceuticals, San Luis, California); 2'- α -L-Fucopyranosyl-D-Lactose (2'FL; Inalco Pharmaceuticals, San Luis, California); LNnT; Orafti[®] HP inulin (HP inulin) (BENEIO-Orafti, Belgium); and gum Arabic (Fisher Scientific, Pittsburgh, Pennsylvania).

In vitro fermentation model

[0156] Approximately 80 mg of each substrate is weighed in triplicate for each pull time into 16-mL Balch tubes that are used in a model that simulates large bowel fermentation. An aliquot (7.2 mL) of medium (Table 1; FIG. 2) is aseptically transferred into the Balch tubes, capped with butyl rubber stoppers, and sealed with aluminum caps. Tubes containing HP inulin and gum arabic are stored at 4°C for approximately 12 h to enable hydration of the substrates before initiating fermentation. These tubes are placed in a 37°C water bath approximately 30 min before inoculation. Due to the cost of the substrates and difficulty in obtaining samples from infants, tubes containing GOS, 6'SL, 2'FL, and LNnT are hydrated upon obtaining a fecal sample and placed in a 37°C water bath until inoculation.

[0157] Sample and blank tubes are aseptically inoculated with 0.8 ml of diluted feces. Tubes are incubated at 37°C with periodic mixing every 2 h for up to 12 h. At 0, 3, 6, and 12 h after inoculation, tubes are removed from the 37°C incubator and processed immediately for analyses. The pH of tube contents is measured with a standard pH meter. A 3-ml subsample of fluid is collected and used for short-chain fatty acid analysis.

Short-chain fatty acid (SCFA) analysis

[0158] The 3-mL aliquot of fluid removed from the sample tubes for SCFA analysis is immediately added to 0.75 mL of 25% metaphosphoric acid. Concentrations of acetate, propionate, and butyrate are determined using a Hewlett-Packard 5890A series II gas chromatograph and a glass column (180 cm x 4 mm i.d.) packed with 10% SP-1200/1% H₃PO₄ on 80/100+ mesh Chromosorb WAW (Supelco Inc., Bellefonte, PA). Oven temperature, detector temperature, and injector temperature are 125, 175, and 180°C, respectively. SCFA concentration values are corrected for blank tube production of SCFA and 0 h concentrations for each substrate. Total SCFA are calculated as the total amount of acetate, propionate, and butyrate.

Results and Discussion

[0159] The pH change from baseline decreases ($P < 0.0001$) over time for all substrates except gum arabic (FIG. 3). At 3, 6, and 12 h after inoculation, pH change from baseline is smallest ($P < 0.0001$) with the gum arabic substrate, and greatest in the LNnT,

2'FL, and GOS substrates. A decrease in pH is an indicator of fermentation, and these data are reflective of SCFA production.

[0160] Total SCFA production differs among substrates (FIG. 4) at 3, 6, and 12 h of fermentation ($P<0.0001$). Gum arabic produces the least amount of SCFA and does not change over time. After 3 and 6 h of fermentation, total SCFA production is lower ($P<0.05$) with HP inulin compared to all other substrates and is lower ($P<0.05$) with 6'SL compared to GOS. By 12 h of fermentation, total SCFA production remains lower ($P<0.05$) with HP inulin relative to 2'FL, 6'SL, GOS, and LNnT substrates. Also, after 12 h of fermentation, total SCFA production is greater ($P<0.05$) for the 6'SL and GOS substrates compared to 2'FL.

EXAMPLE 37

[0161] In this Example, probiotic fermentation parameters are determined for purified HMOs, HMO precursors, and other prebiotic oligosaccharides.

Bacterial Cultures

[0162] All bifidobacteria strains are initially inoculated from frozen stocks, grown in deMan Rogosa Sharpe (MRS) broth (Difco, Detroit, MI) supplemented with 0.5 g/L L-cysteine and incubated at 37°C for 24 h in an anaerobic chamber (90% N₂, 5% CO₂ and 5% H₂). Subsequently, the cultures are passed twice on a semi-synthetic MRS medium (sMRS) + 0.5 g/L L-cysteine which is supplemented with 1% (w/v) filter-sterilized glucose as the sole carbohydrate source. After the 2nd pass, cultures are prepared to use as inoculums for growth assays described below. For bifidobacteria strains, the same procedure is followed except all media are supplemented with 0.5 g/l L-cysteine/HCl. All bacterial strains for use in this Example are listed in the table below.

Table: Microorganisms

#	Culture Collection Number	Genus	Species	Strain
1	MJM29	<i>Bifidobacterium</i>	<i>adolescentis</i>	ATCC 15703
2	MJM30	<i>Bifidobacterium</i>	<i>infantis</i>	S12; ATCC 15697
3	MJM32	<i>Bifidobacterium</i>	<i>animalis</i> subsp. <i>lactis</i>	DSM 10140
4	MJM33	<i>Bifidobacterium</i>	<i>animalis</i> subsp. <i>animalis</i>	ATCC 25527
5	MJM34	<i>Bifidobacterium</i>	<i>bifidum</i>	ATCC 29521

6	MJM35	<i>Bifidobacterium</i>	<i>breve</i>	ATCC 15700
7	MJM37	<i>Bifidobacterium</i>	<i>bifidum</i>	ATCC 11617
8	MJM88	<i>Bifidobacterium</i>	<i>lactis</i>	Bf-6 (Cargill)
9	MJM92	<i>Bifidobacterium</i>	<i>longum</i>	BB536 (Morinaga)
10	MJM93	<i>Bifidobacterium</i>	<i>infantis</i>	M-63 (Morinaga)
11	MJM94	<i>Bifidobacterium</i>	<i>breve</i>	M-16V (Morinaga)
12	MJM95	<i>Bifidobacterium</i>	<i>lactis</i>	Bb12; (Chr. Hansen)

Bacterial Growth Assays

[0163] After the 2nd pass in sMRS + glucose + cysteine, the cultures are washed once with 10 mL of sterile sMRS + cysteine (no carbohydrate), resuspended in 10 ml of sterile sMRS + cysteine (no carbohydrate) and then used as a 1% inoculum. Carbohydrates for use in this Example are shown in the table below. The carbohydrates are sterilized with a 0.22 micron filter and used at a 1% final concentration. Cell growth is performed in 250 μ L of sMRS + cysteine covered with 50 μ L of mineral oil in a Bioscreen 100-well Honeycomb plate. Cell growth is monitored by measuring optical density at 600 nm (OD600) using a Bioscreen C Automated Microbiology Growth Curve Analysis System. The plate reader is operated in discontinuous mode, with absorbance readings performed in 30-minute intervals, and preceded by 30-second shaking intervals at maximum speed. Controls consist of inoculated medium lacking carbohydrate. Due to space limitations on the microtitre plate, the carbohydrates are divided into three separate groups: plate A (HMO precursors: glucose, galactose, lactose, NAG, fucose, fructose and sialic acid), plate B (Prebiotics: glucose, Purimune™ GOS, purified Purimune™ GOS, Vivinal® GOS, purified Vivinal® GOS, scFOS and PDX), and plate C (HMOs: glucose, 6'-SL, 3'-SL, 2'-FL, 3'-FL and LNnT). All three plates include a positive control (glucose) and negative control (no carbohydrate).

Table: Carbohydrates

Carbohydrate	Source
Dextrose (D-Glucose)	Fisher Scientific
D(+)-Galactose	ACROS-ORGANICS
α -Lactose	Fisher Scientific
L-(-) Fucose	SIGMA
D-Fructose	ALDRICH
Sialic acid (N-acetylneuraminic acid)	CALBIOCHEM
NAG (N-acetyl-D-glucosamine)	SIGMA
GOS (Purimune™ Galactooligosaccharide)	GTC Nutrition
Purified GOS (Purimune™ Galactooligosaccharide)	GTC Nutrition

Vivinal® GOS (Galactooligosaccharide)	Friesland Foods
Purified Vivinal® GOS (Galactooligosaccharide)	Friesland Foods
scFOS (Short-Chain Fructooligosaccharide)	Nutraflora® P-95 (GTC Nutrition)
PDX (Litesse® Polydextrose)	DANISCO
6'SL (6'-sialyllactose)	V-labs; SL 306 Lot#HGDX 21-163-1
3'SL (3'-sialyllactose)	V-labs; SL 302 Lot#HGDX 76-161-1
2'FL (2'-fucosyllactose)	V-labs; Lot# DX103
3'FL (3'-fucosyllactose)	V-labs; Lot# DX807
LNnT (Lacto-N-Neotetraose)	Abbott Nutrition

Bacterial growth curves

[0164] The OD600 data for each carbohydrate is corrected by subtracting the OD600 of the basal media (sMRS) from the sample plate for each probiotic. Maximum OD is determined by inspection of the corrected growth data. OD is determined by subtracting the initial corrected OD (time point 0) from the maximum corrected OD. Samples are grown in biologically independent triplicates and the resulting growth kinetic data are expressed as the mean of these replicates.

[0165] For the growth curve plots, OD600 vs. time is first plotted for the bacteria grown on medium lacking carbohydrate (sMRS). For all other carbohydrates, the OD600 data is corrected by subtracting the OD600 of sMRS.

Purification of GOS

[0166] Purified GOS is obtained by purification of Purimune™ GOS (GTC Nutrition) and Vivinal® GOS (Friesland Foods Domo). Stock solutions of 1.5 g/100 mL are applied to a XK column (XK 50/100 column, 5.0 x 100 cm, GE healthcare) packed with Sephadex G25 medium (Sigma). The column is eluted with pure distilled water at a rate of 8 ml/min and is collected in 12-mL fractions by a Gilson FC 203B fraction collector.

[0167] Detection of carbohydrate in every 2-3 fractions is performed using the phenol–sulfuric acid assay. Briefly, 50 µL of sample (2 µL of fraction and 48 µL of distilled water in a well) is added to 150 µL of concentrated sulfuric acid rapidly in a 96-well microtitre plate. Immediately thereafter, 30 µL of 5% phenol is added and the plate is kept in a static water bath for 30 minutes at 80°C. After cooling to room temperature for 5 minutes, it is wiped dry and absorbance at 490 nm is measured by a SpectraMax Plus384 Spectrophotometer. Based on carbohydrate analysis, fractions containing minimal di- and

monosaccharides are pooled and freeze dried (Freeze dry system/Freezezone 4.5/LABCONCO) for bacterial fermentation experiments. In addition, freeze dried GOS is pooled from multiple runs in order to generate enough purified GOS for growth experiments (5 runs with Purimune™ GOS and 3 runs with Vivinal® GOS).

RESULTS & DISCUSSION:

GOS Purification

[0168] GOS is produced by the transgalactosylation of lactose and has been used as a prebiotic supplement in pediatric nutrition. Due to issues with GOS synthesis, commercial GOS products are a mixture of many different carbohydrates which may include mono- and disaccharides. In order to test the fermentation parameters of GOS and not the mono- and disaccharides which would not normally reach the colon, a purified GOS fraction, essentially free of mono- and disaccharides is obtained. Glucose (monosaccharide), lactose (disaccharide) and raffinose (trisaccharide) are used as standards. Consistent with information from the suppliers, Purimune™ GOS has less mono- and disaccharides than Vivinal® GOS. For example, the Purimune™ GOS peaks before the raffinose peak suggesting that Purimune™ GOS consists primarily of trisaccharides or larger. For Vivinal® GOS, the peak is observed at a similar fraction number as lactose. Since lactose begins to appear in fraction 55, fractions 30 through 55 are used as the purified GOS from both suppliers.

HMO Precursor Fermentation

[0169] All bifidobacteria tested grow very little in the basal media (sMRS + cysteine) (FIG. 5A), whereas they all grow well in glucose (FIG. 5B). In general, the bifidobacteria, which is not able to ferment galactose (FIG. 5C), also has reduced growth on lactose (FIG. 5D). None of the bifidobacteria are able to ferment L-fucose (FIG. 5E) or sialic acid (FIG. 5F), two key constituents of HMOs and mucin. Only *B. breve* ATCC 15700 is able to ferment NAG (FIG. 5G), a key component of HMOs and mucin. Lastly, the majority of bifidobacteria is able to ferment fructose (FIG. 5H).

Prebiotic Fermentation

[0170] Removal of mono- and disaccharides from Purimune™ GOS results in a decrease in growth for all bifidobacteria (FIG. 6A). In fact, *B. lactis* DSM 10140, *B. animalis* ATCC 25527, *B. bifidum* ATCC 29521, *B. lactis* Bf-6 and *B. longum* are not able

to ferment the purified Purimune™ GOS (FIG. 6D). A similar pattern is seen with purified Vivinal® GOS (FIG. 6F), except more growth is seen with Vivinal® GOS (FIG. 6E) than Purimune™ GOS (FIG. 6C). In order to mimic the colonic situation, the free mono- and disaccharides present in these products need to be removed. Also, it is clear that Purimune™ GOS has a higher relative concentration of oligosaccharides. Both *B. infantis* strains are among the best growers on purified GOS as determined by Δ OD, confirming that GOS is a reasonable prebiotic to add to infant formula if the goal is to increase *B. infantis*. All bifidobacteria tested, except for *B. animalis* ATCC 25527, are able to ferment scFOS (FIG. 6G), whereas no bifidobacteria are able to ferment polydextrose (PDX) (FIG. 6H).

HMO Fermentation

[0171] Only *B. infantis* ATCC 15697 and *B. infantis* M-63 are able to ferment 6'-SL, 3'-SL, 2'-FL and 3'-FL (FIG. 7). In all cases, *B. infantis* M-63 grows better than *B. infantis* ATCC 15697. On the more complex LNnT, *B. breve* ATCC 15700 and the two *B. infantis* strains grow well but not *B. breve* M-16V. In addition, the ability of the two *B. infantis* strains to ferment HMOs correlates with the abundance of *B. infantis* found in breast fed infants. Curiously, both *B. infantis* strains are not able to ferment fucose or sialic acid.

CONCLUSIONS:

[0172] There are significant differences amongst the tested bifidobacteria strains regarding their abilities to ferment HMO precursors, prebiotics and HMOs. Of the 12 bifidobacteria strains tested, none are able to ferment sialic acid. Regarding prebiotics, most of the bifidobacteria are able to ferment GOS and scFOS, but they are not able to ferment PDX. Amongst the bifidobacteria strains tested, only *B. infantis* ATCC 15697 and *B. infantis* M-63 are able to ferment 6'-SL, 3'-SL, 2'-FL and 3'-FL. *B. breve* ATCC 15700, *B. infantis* ATCC 15697 and *B. infantis* M-63 are able to ferment LNnT.

WHAT IS CLAIMED IS:

1. A synthetic pediatric formula for promoting intestinal barrier integrity, the synthetic formula comprising a probiotic, a first oligosaccharide in a concentration of from about 1 mg/mL to about 4 mg/mL and selected from the group consisting of a galactooligosaccharide, a fructooligosaccharide, and combinations thereof; and a second
5 oligosaccharide in a concentration of from about 0.05 mg/mL to about 0.5 mg/mL and selected from the group consisting of 2'-fucosyllactose, 3'-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, lacto-N-neotetraose, and combinations thereof.
2. The synthetic pediatric formula of claim 1, wherein the probiotic is of human infant origin.
3. The synthetic pediatric formula of claim 2, wherein the probiotic is a *Bifidobacterium*.
4. The synthetic pediatric formula of claim 3, wherein the probiotic is *Bifidobacterium infantis*.
5. The synthetic pediatric formula of claim 4, wherein the probiotic is selected from the group consisting of *Bifidobacterium infantis* M-63, *Bifidobacterium infantis* ATCC 15697, *Bifidobacterium infantis* 35624, *Bifidobacterium infantis* CHCC2228, *Bifidobacterium infantis* BB-02, *Bifidobacterium infantis* DSM20088, *Bifidobacterium infantis* R-0033, and combinations thereof.
6. The synthetic pediatric formula of claim 1, comprising probiotics in a concentration of from about 10^3 CFU/g to about 10^{12} CFU/g.
7. A method of promoting the growth of beneficial microbiota in the gastrointestinal tract of an infant or toddler, the method comprising administering to the infant or toddler a synthetic pediatric formula comprising a probiotic, a first
5 oligosaccharide in a concentration of from about 1 mg/mL to about 4 mg/mL and selected from the group consisting of galactooligosaccharide, fructooligosaccharide, and combinations thereof; and a second oligosaccharide in a concentration of from about 0.05 mg/mL to about 0.5 mg/mL and selected from the group consisting of 2'-fucosyllactose,

3'-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, lacto-N-neotetraose, and combinations thereof.

8. The method of claim 7, wherein the gastrointestinal diseases and/or disorders are selected from the group consisting of irritable bowel syndrome, colitis, food intolerance, and food allergies.

9. The method of claim 8, wherein the gastrointestinal disease is necrotizing enterocolitis.

10. The method of claim 7, wherein the probiotic is of human infant origin.

11. The method of claim 10, wherein the probiotic is a *Bifidobacterium*.

12. The method of claim 11, wherein the probiotic is *Bifidobacterium infantis*.


13. The method of claim 12, wherein the probiotic is selected from the group consisting of *Bifidobacterium infantis* M-63, *Bifidobacterium infantis* ATCC 15697, *Bifidobacterium infantis* 35624, *Bifidobacterium infantis* CHCC2228, *Bifidobacterium infantis* BB-02, *Bifidobacterium infantis* DSM20088, *Bifidobacterium infantis* R-0033, and combinations thereof.

14. A method of stimulating enteric nerve cells in the gastrointestinal tract of an infant or toddler, the method comprising administering to the infant or toddler a synthetic pediatric formula comprising a probiotic, a first oligosaccharide in a concentration of from about 1 mg/mL to about 4 mg/mL and selected from the group consisting of
5 galactooligosaccharide, fructooligosaccharide, and combinations thereof; and a second oligosaccharide in a concentration of from about 0.05 mg/mL to about 0.05 mg/mL and selected from the group consisting of 2'-fucosyllactose, 3'-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, lacto-N-neotetraose, and combinations thereof.

15. The synthetic pediatric formula of claim 14, wherein the probiotic is of human infant origin.

1/15

FIG. 1



2/15

FIG. 2

Table 1. Composition of microbiological medium used in the *in vitro* experiment.

Component	Concentration in medium
	<i>mL/L</i>
Solution A ¹	330.0
Solution B ²	330.0
Trace mineral solution ³	10.0
Water-soluble vitamin solution ⁴	20.0
Folate:biotin solution ⁵	5.0
Riboflavin solution ⁶	5.0
Hemin solution ⁷	2.5
Resazurin ⁸	1.0
Distilled H ₂ O	296.1
	<i>g/L</i>
Na ₂ CO ₃	4.0
Cysteine HCl-H ₂ O	0.5
Trypticase	0.5
Yeast extract	0.5

¹Composition (g/L): NaCl, 5.4; KH₂PO₄, 2.7; CaCl₂-H₂O, 0.16; MgCl₂-6H₂O, 0.12; MnCl₂-4H₂O, 0.06; CoCl₂-6H₂O, 0.06; (NH₄)₂SO₄, 5.4.

²Composition (g/L): K₂HPO₄, 2.7.

³Composition (mg/L): ethylenediaminetetraacetic acid (disodium salt), 500; FeSO₄-7H₂O, 200; ZnSO₄-7H₂O, 10; MnCl₂-4H₂O, 3; H₃PO₄, 30; CoCl₂-6H₂O, 20; CuCl₂-2H₂O, 1; NiCl₂-6H₂O, 2; Na₂MoO₄-2H₂O, 3.

⁴Composition (mg/L): thiamin-HCl, 100; d-pantothenic acid, 100; niacin, 100; pyridoxine, 100; p-aminobenzoic acid, 5; vitamin B₁₂, 0.25.

⁵Composition (mg/L): folic acid, 10; d-biotin, 2; NH₄HCO₃, 100.


⁶Composition: riboflavin, 10 mg/mL in 5 mmol/L of Hepes.

⁷Composition: hemin, 500 mg/mL in 10 mmol/L of NaOH.

⁸Composition: resazurin, 1 g/L in distilled H₂O.


3/15

FIG. 3



^{a,b} Points not sharing a common superscript letter within each time differ ($P < 0.05$)

FIG. 4



^{a,b} Points not sharing a common superscript letter within each time differ ($P < 0.05$)

4/15

FIG. 5A





FIG. 5B



5/15

FIG. 5C





FIG. 5D



6/15

FIG. 5E





FIG. 5F



7/15

FIG. 5G





FIG. 5H



8/15

FIG. 6A





FIG. 6B



10/15

FIG. 6E





FIG. 6F



11/15

FIG. 6G





FIG. 6H



12/15

FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 7E




FIG. 7F





FIG. 7G



INTERNATIONAL SEARCH REPORT

International application No PCT/US2011/067012

A. CLASSIFICATION OF SUBJECT MATTER INV. A23L1/29 A23L1/30 A61K31/702 A61K35/74 A61P1/00 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols) A23L A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, FSTA, BIOSIS, MEDLINE				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A	WO 2004/112509 A2 (NESTEC SA [CH]; GARCIA-RODENAS CLARA LUCIA [CH]; BERGONZELLI GABRIELA) 29 December 2004 (2004-12-29) pages 3-11; example 1, -----	1-15		
A	EP 2 060 257 A1 (NESTEC SA [CH]) 20 May 2009 (2009-05-20) paragraphs [0016] - [0042]; claims 1-14; example 1 -----	1-15		
A	WO 2010/023178 A1 (CHR HANSEN AS [DK]; KILDSGAARD JENS [DK]; JANZEN THOMAS [DK]; FLAMBARD) 4 March 2010 (2010-03-04) page 1 - page 2, line 30; claims 1-14 ----- -/--	1-15		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.				
* Special categories of cited documents : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; border: none; vertical-align: top;"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
11 May 2012	24/05/2012			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Krajewski, Doris			

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2011/067012

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2007/046699 A2 (NUTRICIA NV [NL]; SCHMITT JOACHIM [DE]; BOEHM GUNTHER [DE]; BEERMANN C) 26 April 2007 (2007-04-26) pages 2-11; claims 1-12 -----	1-15
Y	WO 01/60346 A2 (AMERICAN HOME PROD [US]) 23 August 2001 (2001-08-23) page 3, line 22 - page 6, line 5; claims 1-14; examples 1-3 -----	7-13
Y	ROBERT E. WARD ET AL: "In vitro fermentability of human milk oligosaccharides by several strains of bifidobacteria", MOLECULAR NUTRITION & FOOD RESEARCH, vol. 51, no. 11, 1 November 2007 (2007-11-01), pages 1398-1405, XP55026719, ISSN: 1613-4125, DOI: 10.1002/mnfr.200700150 the whole document -----	7-13
A	US 6 083 934 A (PRIETO PEDRO A [US] ET AL) 4 July 2000 (2000-07-04) column 2, line 13 - column 4, line 45; claim 1; examples 1,2 -----	1-15
A	WO 2009/102193 A1 (NUTRICIA NV [NL]; SCHOLTENS PETRONELLA ANNA MARI [NL]; ALLES MARTINE S) 20 August 2009 (2009-08-20) page 1, line 26 - page 9, line 22 claims 1-10; examples 1-3 -----	1-15
A	RINNE M M ET AL: "Similar bifidogenic effects of prebiotic-supplemented partially hydrolyzed infant formula and breastfeeding on infant gut microbiota", FEMS IMMUNOLOGY AND MEDICAL MICROBIOLOGY, ELSEVIER SCIENCE B.V., AMSTERDAM, NL, vol. 43, no. 1, 1 January 2005 (2005-01-01), pages 59-65, XP025316645, ISSN: 0928-8244, DOI: 10.1016/J.FEMSIM.2004.07.005 [retrieved on 2005-01-01] the whole document -----	1-15
A	EP 2 127 661 A1 (NESTEC SA [CH]) 2 December 2009 (2009-12-02) paragraphs [0011] - [0012], [0021] - [0055]; examples 1-18 -----	1-15
	-/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/067012

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2007/101675 A1 (NESTEC SA [CH]; SPRENGER NORBERT [CH]; MORGAN FRANCOIS [FR]; BERROCAL) 13 September 2007 (2007-09-13) page 4, line 20 - page 7, line 12 page 8, line 15 - page 14, line 23; claims 1-19; examples 1-6 -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2011/067012

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-6, 14, 15

synthetic pediatric formula for promoting intestinal barrier integrity comprising a probiotic, a first oligosaccharide in a defined concentration and a second selected human milk oligosaccharide in a defined amount
method of stimulating enteric nerve cells using said pediatric formula

2. claims: 7-13

method of using said synthetic pediatric formula for promoting the growth of beneficial microbiota

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2011/067012

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004112509	A2	29-12-2004	AR 044888 A1 05-10-2005
			CA 2530437 A1 29-12-2004
			CN 1863463 A 15-11-2006
			EP 1638416 A2 29-03-2006
			TW 1342779 B 01-06-2011
			US 2007104700 A1 10-05-2007
			WO 2004112509 A2 29-12-2004

EP 2060257	A1	20-05-2009	AU 2008324205 A1 14-05-2009
			CA 2704977 A1 14-05-2009
			CN 101909615 A 08-12-2010
			EP 2060257 A1 20-05-2009
			EP 2217230 A1 18-08-2010
			RU 2010123174 A 20-12-2011
			TW 200936147 A 01-09-2009
			US 2010233129 A1 16-09-2010
			WO 2009059996 A1 14-05-2009

WO 2010023178	A1	04-03-2010	NONE

WO 2007046699	A2	26-04-2007	AR 056145 A1 19-09-2007
			AR 058132 A1 23-01-2008
			AR 058133 A1 23-01-2008
			AR 058136 A1 23-01-2008
			AT 432620 T 15-06-2009
			AT 440508 T 15-09-2009
			AT 442786 T 15-10-2009
			AT 476877 T 15-08-2010
			AU 2006303477 A1 26-04-2007
			AU 2006304992 A1 26-04-2007
			AU 2006304993 A1 26-04-2007
			BR PI0617507 A2 26-07-2011
			BR PI0617645 A2 02-08-2011
			BR PI0617653 A2 02-08-2011
			CA 2620821 A1 26-04-2007
			CA 2626615 A1 26-04-2007
			CA 2626636 A1 26-04-2007
			CN 101272701 A 24-09-2008
			CN 101330837 A 24-12-2008
			CN 101330838 A 24-12-2008
			CN 101360429 A 04-02-2009
			DK 1940246 T3 01-11-2010
			DK 1940247 T3 13-07-2009
			DK 1940250 T3 11-01-2010
			EP 1776877 A1 25-04-2007
			EP 1940245 A1 09-07-2008
			EP 1940246 A1 09-07-2008
			EP 1940247 A2 09-07-2008
			EP 1940250 A1 09-07-2008
			EP 2105055 A1 30-09-2009
			EP 2140771 A1 06-01-2010
			EP 2407036 A1 18-01-2012
			ES 2327781 T3 03-11-2009
ES 2332064 T3 25-01-2010			
ES 2350321 T3 21-01-2011			
JP 2009512686 A 26-03-2009			
MY 145311 A 13-01-2012			
PT 1940245 E 04-01-2010			

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2011/067012

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		PT 1940246 E	12-11-2010
		PT 1940247 E	09-09-2009
		PT 1940250 E	15-10-2009
		SG 176521 A1	29-12-2011
		US 2009162323 A1	25-06-2009
		US 2009220639 A1	03-09-2009
		US 2009221486 A1	03-09-2009
		US 2009263537 A1	22-10-2009
		WO 2007045502 A1	26-04-2007
		WO 2007046697 A1	26-04-2007
		WO 2007046698 A1	26-04-2007
		WO 2007046699 A2	26-04-2007

WO 0160346	A2	23-08-2001	AR 027450 A1
			AT 357854 T
			AU 3828501 A
			BR 0108478 A
			CA 2400737 A1
			CN 1406111 A
			EP 1255449 A2
			JP 2003522784 A
			MX PA02008016 A
			PT 1255449 E
			TW 1265008 B
			WO 0160346 A2

US 6083934	A	04-07-2000	AT 271788 T
			CA 2285066 A1
			DE 69825260 D1
			DE 69825260 T2
			DK 0973413 T3
			EP 0973413 A1
			ES 2229486 T3
			JP 4148535 B2
			JP 2001517950 A
			NO 994722 A
			PT 973413 E
			US 5906982 A
			US 6083934 A
			WO 9843495 A1

WO 2009102193	A1	20-08-2009	CN 101945658 A
			EP 2240186 A1
			US 2010322904 A1
			WO 2009102193 A1
			WO 2009102199 A1

EP 2127661	A1	02-12-2009	NONE

WO 2007101675	A1	13-09-2007	AU 2007222598 A1
			BR PI0708689 A2
			CA 2644968 A1
			CN 101432007 A
			EP 1993576 A1
			US 2009041736 A1
			WO 2007101675 A1
			ZA 200808503 A
