



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/12/01
(87) Date publication PCT/PCT Publication Date: 2018/09/07
(85) Entrée phase nationale/National Entry: 2020/08/19
(86) N° demande PCT/PCT Application No.: US 2017/064361
(87) N° publication PCT/PCT Publication No.: 2018/160245
(30) Priorités/Priorities: 2017/03/01 (US62/465,797);
2017/03/01 (US62/465,798); 2017/03/01 (US62/465,819);
2017/03/02 (US62/466,253); 2017/03/02 (US62/466,256);
2017/03/02 (US62/465,834); 2017/03/06 (US62/467,755);
2017/03/06 (US62/467,740); 2017/03/06 (US62/467,742);
2017/09/08 (US62/556,288)

(71) Demandeur/Applicant:
INDIGO AG, INC., US
(72) Inventeur/Inventor:
RILEY, RAYMOND, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : COMPOSITIONS D'ENDOPHYTE ET PROCEDES D'AMELIORATION DE CARACTERISTIQUES DE PLANTE
(54) Title: ENDOPHYTE COMPOSITIONS AND METHODS FOR IMPROVEMENT OF PLANT TRAITS

(57) **Abrégé/Abstract:**

This invention relates to methods and materials for modulating the characteristics of a plant, said plant having been heterologously disposed to an endophyte or a plurality of endophytes, or derived from a plant reproductive element heterologously disposed to an endophyte or a plurality of endophytes.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

07 September 2018 (07.09.2018)



(10) International Publication Number

WO 2018/160245 A1

(51) International Patent Classification:

A01N 63/04 (2006.01) C05F 11/08 (2006.01)

C12N 1/14 (2006.01) A01H 17/00 (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2017/064361

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

(22) International Filing Date:

01 December 2017 (01.12.2017)

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/465,819	01 March 2017 (01.03.2017)	US
62/465,797	01 March 2017 (01.03.2017)	US
62/465,798	01 March 2017 (01.03.2017)	US
62/466,253	02 March 2017 (02.03.2017)	US
62/465,834	02 March 2017 (02.03.2017)	US
62/466,256	02 March 2017 (02.03.2017)	US
62/467,742	06 March 2017 (06.03.2017)	US
62/467,740	06 March 2017 (06.03.2017)	US
62/467,755	06 March 2017 (06.03.2017)	US
62/556,288	08 September 2017 (08.09.2017)	US

(71) Applicant: INDIGO AG, INC. [US/US]; 500 Rutherford Avenue, North Building, Boston, MA 02129 (US).

(72) Inventor: RILEY, Raymond; c/o Indigo Ag, Inc., 500 Rutherford Avenue, North Building, Boston, MA 02129 (US).

(74) Agent: HUBL, Susan, T. et al.; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Mountain View, CA 94041 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: ENDOPHYTE COMPOSITIONS AND METHODS FOR IMPROVEMENT OF PLANT TRAITS

(57) Abstract: This invention relates to methods and materials for modulating the characteristics of a plant, said plant having been heterologously disposed to an endophyte or a plurality of endophytes, or derived from a plant reproductive element heterologously disposed to an endophyte or a plurality of endophytes.

 WO 2018/160245 A1

ENDOPHYTE COMPOSITIONS AND METHODS FOR IMPROVEMENT OF PLANT TRAITS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to Provisional Application No. 62/556,288, filed September 8, 2017; Provisional Application No. 62/467,740, filed March 6, 2017; Provisional Application No. 62/467,742, filed March 6, 2017; Provisional Application No. 62/467,755, filed March 6, 2017; Provisional Application No. 62/466,253, filed March 2, 2017; Provisional Application No. 62/465,834, filed March 2, 2017; Provisional Application No. 62/466,256, filed March 2, 2017; Provisional Application No. 62/465,797, filed March 1, 2017; Provisional Application No. 62/465,819, filed March 1, 2017; and Provisional Application No. 62/465,798, filed March 1, 2017, the disclosures of which are incorporated by reference in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing with 71 sequences which has been submitted via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on November 29, 2017, is named 39061_10101_Sequence_Listing.txt, and is 54,920 bytes in size.

FIELD OF THE INVENTION

[0003] This invention relates to compositions and methods for improving the traits of plants, particularly plants important for human or animal consumption, for example rice (*Oryza sativa* and related varieties), soy (*Glycine max* and related varieties), wheat (*Triticum aestivum* and related varieties), and corn (*Zea mays* and related varieties). For example, this invention describes microbes that are capable of living within or heterologously disposed to a plant, and which can be used to impart improved traits to plants with which they are or have been heterologously disposed. The disclosed invention also describes methods of improving plant element characteristics by introducing microbes to parental plants. Further, this invention also provides methods of treating plant elements with microbes that are capable of living within a plant, particularly rice, soy, wheat, and corn, to impart improved yield, and other agronomic characteristics to that plant.

BACKGROUND

[0004] According the United Nations Food and Agricultural Organization, the world's population will exceed 9.6 billion people by the year 2050, which will require significant

improvements in agricultural to meet growing food demands. There is a need for improved agricultural plants that will enable the nearly doubled food production demands with fewer resources and more environmentally sustainable inputs, for plants with improved responses to various abiotic stresses.

[0005] Today, crop performance is optimized primarily via technologies directed towards the interplay between crop genotype (e.g., plant breeding, genetically-modified (GM) crops) and its surrounding environment (e.g., fertilizer, synthetic herbicides, pesticides). While these paradigms have assisted in doubling global food production in the past fifty years, yield growth rates have stalled in many major crops, and shifts in the climate have been linked to production instability as well as changing pest and disease pressures, driving an urgent need for novel solutions to crop improvement. In addition to their long development and regulatory timelines, public fears of GM-crops and synthetic chemicals have challenged their use in many key crops and countries, resulting in a lack of acceptance for many GM traits and the exclusion of GM crops and many synthetic chemistries from some global markets. Thus, there is a significant need for innovative, effective, environmentally-sustainable, and publicly-acceptable approaches to improving the yield and other agronomically important characteristics of plants.

[0006] Provided herein are methods and compositions for improving agronomically important characteristics of plants by associating those plants with the disclosed endophytes.

SUMMARY OF INVENTION

[0007] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the corn plant element is the variety Stine 9734 or a closely related variety thereof.

[0008] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide

sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

[0009] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62 and wherein the soybean plant is a variety selected from the group consisting of Pfister 38R25, Dairyland DSR1808R2Y, Stine 3920, and closely related varieties thereof.

[0010] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

[0011] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, and wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0012] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

[0013] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62 and wherein the peanut plant is a variety selected from the group consisting of AT-9899, FloRun 107, Tamnut OL06, Georgia-06G, and closely related varieties thereof.

[0014] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and root dry weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

[0015] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and root dry weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, and wherein the soybean is variety Stine 33E22 or a closely related variety thereof.

[0016] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

[0017] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, and wherein the peanut plant is a variety selected from the group consisting of FloRun 107, Georgia-06G, and closely related varieties thereof.

[0018] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase root length in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

[0019] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[0020] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the canola plant is the variety Brett Young 552 or a closely related variety thereof.

[0021] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element,

wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66.

[0022] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, and wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.

[0023] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66.

[0024] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0025] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at

least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45.

[0026] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45, wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.

[0027] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45.

[0028] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0029] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the corn plant element is the variety Stine 9734 or a closely related variety.

[0030] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67.

[0031] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.

[0032] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67.

[0033] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0034] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the

endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69.

[0035] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69, wherein the canola plant is variety NCC1015 or a closely related variety thereof.

[0036] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

[0037] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the trait of agronomic importance is yield and the soybean plant is variety Pfister 38R25 or a closely related variety thereof.

[0038] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of decreasing grain moisture and increasing yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at

least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

[0039] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of decreasing grain moisture and increasing yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0040] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the peanut plant is a variety Georgia-06G or a closely related variety.

[0041] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase dry shoot biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

[0042] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing a first endophyte and a second endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of plant height, fresh root weight, and fresh shoot weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a

polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[0043] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing a first endophyte and a second endophyte to a rice plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[0044] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68.

[0045] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68, wherein the soybean is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.

[0046] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68.

[0047] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0048] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68 and the peanut plant is the variety AT-9899 or a closely related variety thereof.

[0049] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing a first endophyte and a second endophyte to a soybean plant element in an amount effective to increase dry root biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, the second endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, and soybean plant is variety Stine 33E22 or a closely related variety thereof.

[0050] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[0051] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[0052] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the trait of agronomic importance is yield and the soybean plant is the variety Pfister 38R25 or a closely related variety thereof.

[0053] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root length and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[0054] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root length and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ

ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.

[0055] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the corn plant element is the variety Stine 9734 or a closely related variety thereof, wherein the plant element is a seed, optionally a modified seed.

[0056] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the plant element is a seed, optionally a modified seed.

[0057] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the plant element is a seed, optionally a modified seed.

[0058] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the plant element is a seed, optionally a modified seed.

[0059] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and root dry weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the plant element is a seed, optionally a modified seed.

[0060] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the plant element is a seed, optionally a modified seed.

[0061] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase root length in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the plant element is a seed, optionally a modified seed.

[0062] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the plant element is a seed, optionally a modified seed.

[0063] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the canola plant is the variety Brett Young 552 or a closely related variety thereof, wherein the plant element is a seed, optionally a modified seed.

[0064] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, wherein the plant element is a seed, optionally a modified seed.

[0065] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, wherein the plant element is a seed, optionally a modified seed.

[0066] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45, wherein the plant element is a seed, optionally a modified seed.

[0067] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45, wherein the plant element is a seed, optionally a modified seed.

[0068] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the corn plant element is the variety Stine 9734 or a closely related variety, wherein the plant element is a seed, optionally a modified seed.

[0069] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the plant element is a seed, optionally a modified seed.

[0070] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the plant element is a seed, optionally a modified seed.

[0071] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated

plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69, wherein the plant element is a seed, optionally a modified seed.

[0072] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the plant element is a seed, optionally a modified seed.

[0073] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of decreasing grain moisture and increasing yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the plant element is a seed, optionally a modified seed.

[0074] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the peanut plant is a variety Georgia-06G or a closely related variety, wherein the plant element is a seed, optionally a modified seed.

[0075] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase dry shoot biomass in the plant derived from the

treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the plant element is a seed, optionally a modified seed.

[0076] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing a first endophyte and a second endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of plant height, fresh root weight, and fresh shoot weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the plant element is a seed, optionally a modified seed.

[0077] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing a first endophyte and a second endophyte to a rice plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the plant element is a seed, optionally a modified seed.

[0078] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide

sequence at least 97% identical to SEQ ID NO: 68, wherein the plant element is a seed, optionally a modified seed.

[0079] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68, wherein the plant element is a seed, optionally a modified seed.

[0080] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68 and the peanut plant is the variety AT-9899 or a closely related variety thereof, wherein the plant element is a seed, optionally a modified seed.

[0081] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing a first endophyte and a second endophyte to a soybean plant element in an amount effective to increase dry root biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, the second endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, and soybean plant is variety Stine 33E22 or a closely related variety thereof, wherein the plant element is a seed, optionally a modified seed.

[0082] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at

least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the plant element is a seed, optionally a modified seed.

[0083] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the plant element is a seed, optionally a modified seed.

[0084] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root length and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the plant element is a seed, optionally a modified seed.

[0085] In some embodiments, any of the endophyte is heterologously disposed to the plant element in a formulation, said formulation further comprising one or more of the following: a stabilizer, a preservative, a carrier, a surfactant, a fungicide, a nematocide, a bactericide, an insecticide, or herbicide, or any combination thereof.

[0086] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the corn plant element is the variety Stine 9734 or a closely related variety thereof, wherein the percent identify is

determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0087] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0088] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0089] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0090] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and root dry weight in the plant derived from the treated plant element relative to a plant derived from a reference plant

element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0091] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0092] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase root length in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0093] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0094] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola

plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 70, wherein the canola plant is the variety Brett Young 552 or a closely related variety thereof, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0095] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0096] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0097] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44,

and 45, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0098] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[0099] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 67, wherein the corn plant element is the variety Stine 9734 or a closely related variety, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00100] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 67, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00101] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide

sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 67, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00102] In one aspect, the invention provides a method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 69, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00103] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00104] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of decreasing grain moisture and increasing yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00105] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated

plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the peanut plant is a variety Georgia-06G or a closely related variety, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00106] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase dry shoot biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00107] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing a first endophyte and a second endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of plant height, fresh root weight, and fresh shoot weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00108] In one aspect, the invention provides a method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing a first endophyte and a second endophyte to a rice plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant

element, wherein the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00109] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 68, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00110] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 68, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00111] In one aspect, the invention provides a method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 68 and the peanut plant is the variety AT-9899 or a closely related variety thereof, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00112] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing a first endophyte and a second endophyte to a soybean plant element in an amount effective to increase dry root biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the first endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, the second endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, and soybean plant is variety Stine 33E22 or a closely related variety thereof, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00113] In one aspect, the invention provides a method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00114] In one aspect, the invention provides a method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00115] In one aspect, the invention provides a method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat

plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root length and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00116] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

[00117] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

[00118] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62 further comprising a second endophyte, wherein the second endophyte is of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[00119] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[00120] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70.

[00121] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66.

[00122] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68.

[00123] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69.

[00124] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

[00125] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68 and wherein the microbial active ingredient further comprises a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide

sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[00126] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[00127] In some embodiments, the formulation oil comprises rapeseed, NEEM, or erucic acid, or comprises herbicidal or insecticidal properties. In some embodiments, the formulation surfactant is a non-ionic detergent, Tween 20, or Triton X-100. In some embodiments, the formulation polymer is Flo Rite®, DISCOTM, or Kannar® Universal Wonder. In some embodiments, the formulation microbial active ingredient comprises a spore suspension, spray dried spores, or whole cell broth. In some embodiments, the formulation further comprises one or more of the following: fungicide, nematocide, bactericide, insecticide, herbicide, stabilizer, preservative, carrier, anticomplex agent, or any combination thereof. In some embodiments, the endophyte of the formulation is shelf-stable

[00128] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00129] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00130] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus

Exserohilum and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62 further comprising a second endophyte, wherein the second endophyte is of the genus Chaetomium and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00131] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus Coniochaeta and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00132] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus Epicoccum and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 70, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00133] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus Curvularia and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00134] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus Cladosporium and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group

consisting of SEQ ID NOs: 67 and 68, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00135] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 69, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00136] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00137] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68 and wherein the microbial active ingredient further comprises a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00138] In one aspect, the invention provides an agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the percent identity is

determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00139] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00140] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a corn plant element and the improved trait of agronomic importance is yield.

[00141] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element.

[00142] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety selected from the group consisting of: Pfister 38R25, Dairyland DSR1808R2Y, Stine 3920, and closely related varieties thereof, and the improved trait of agronomic importance is yield.

[00143] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the

genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the improved trait of agronomic importance is yield.

[00144] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a peanut plant element and the improved trait of agronomic importance is yield.

[00145] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a peanut variety selected from the group consisting of: AT9899, FloRun 107, Georgia-06G, Tamnut OL06, and closely related varieties thereof.

[00146] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00147] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte,

wherein the plant element is a soybean plant element and the improved trait of agronomic importance is selected from the group consisting of root area, root length, and root dry weight.

[00148] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety Stine 33E22 or a closely related variety thereof.

[00149] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a peanut plant element and the improved trait of agronomic importance is yield.

[00150] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a peanut plant element is selected from the group consisting of the variety FloRun 107, Georgia-06G, and closely related varieties thereof and the improved trait of agronomic importance is yield.

[00151] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte,

and further comprising a second endophyte, wherein the second endophyte is of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00152] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is the soybean variety Stine 33E22 and the trait of agronomic importance is dry root biomass.

[00153] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00154] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.

[00155] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the synthetic composition is capable of providing an improved

trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00156] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is canola variety Brett Young 5525 or a closely related variety

[00157] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00158] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

[00159] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof, and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

[00160] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.

[00161] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof, and the trait of agronomic importance is yield.

[00162] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00163] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

[00164] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%

identical to SEQ ID NO: 67, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof, and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

[00165] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00166] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a corn plant element and the trait of agronomic importance is yield.

[00167] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is corn variety Stine 9734 and the trait of agronomic importance is yield.

[00168] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of

agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

[00169] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof and the trait of agronomic importance is yield.

[00170] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.

[00171] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof and the trait of agronomic importance is yield.

[00172] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of

agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the endophyte comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68, the plant element is a peanut of variety AT9899, and the trait of agronomic importance is yield.

[00173] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00174] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is canola variety NCC1015 and the trait of agronomic importance is yield.

[00175] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00176] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element and the trait of agronomic importance is dry shoot biomass.

[00177] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the

genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is grain moisture or yield.

[00178] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is wheat variety SDSU Select and the trait of agronomic importance is yield.

[00179] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is peanut variety Georgia-06G and the trait of agronomic importance is yield.

[00180] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a rice plant element and the trait of agronomic importance is dry shoot biomass.

[00181] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48,

49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, and further comprising a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00182] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, and further comprising a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is selected from the group consisting of plant height, fresh root biomass, and fresh shoot weight.

[00183] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, and further comprising a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further

comprising the endophyte, wherein the plant element is a rice plant element and the trait of agronomic importance is yield.

[00184] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

[00185] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, dry shoot biomass, and yield.

[00186] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof, and the trait of agronomic importance is yield.

[00187] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further

comprising the endophyte, wherein the plant element is a wheat plant element and the trait of agronomic importance is root length or yield.

[00188] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof and the trait of agronomic importance is yield.

[00189] In some embodiments, any of the synthetic compositions described herein further comprise a plant element is a seed, optionally wherein the seed is modified. In some embodiments, wherein the improved trait of agronomic importance is conferred under normal watering conditions. In some embodiments, the plant element is placed into a substrate that promotes plant growth, optionally soil. In some embodiments, a plurality of said plant elements are placed in the soil in rows, with substantially equal spacing between each seed within each row.

[00190] In some embodiments, any of the synthetic compositions described herein further comprise a formulation that comprising one or more of the following: stabilizer, preservative, carrier, surfactant, anticomplex agent, or any combination thereof and/or one or more of the following: fungicide, nematocide, bactericide, insecticide, or herbicide. In some embodiments, any of the synthetic compositions described herein are confined within an object selected from the group consisting of: bottle, jar, ampule, package, vessel, bag, box, bin, envelope, carton, container, silo, shipping container, truck bed, or case. In some embodiments, any of the synthetic compositions described herein are shelf-stable.

[00191] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00192] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00193] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00194] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 70, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00195] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identity is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00196] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 67, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00197] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00198] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to SEQ ID NO: 69, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00199] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

[00200] In one aspect, the invention provides a synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97%, at least 98%, at least 99%, or 100% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte, wherein the percent identify is determined over a region of alignment of at least 100, 200, 300, 400, or at least 500 nucleotides.

BRIEF DESCRIPTION OF THE DRAWINGS

[00201] Figure 1A shows exemplary rice plants grown as described in Example 29. The pot on the left contains untreated control rice plants. The pot on the right shows rice plants treated with MIC-68178.

[00202] Figure 2 shows the washed roots of exemplary rice plants grown as described in Example 29. The roots of untreated control rice plants are on the left. The roots of rice plants treated with MIC-68178 are on the right.

[00203] Figure 3 shows exemplary soybean roots of soybean plants grown as described in Example 29. The roots of untreated control rice plants are on the left. The roots of soybean plants treated with a combination of MIC-68178 and MIC-33414 are on the right.

[00204] Figure 4 shows the pedigree of the wheat variety SDSU Focus.

[00205] Figure 5 shows the pedigree of the wheat variety SDSU Select.

DETAILED DESCRIPTION

[00206] As demonstrated herein, agricultural plants are heterologously disposed to symbiotic microorganisms, termed endophytes, particularly bacteria and fungi, which contribute to plant survival, performance, and characteristics.

[00207] Described herein are endophytes that are capable of living within or otherwise heterologously disposed to plants to improve plant characteristics. Described herein are methods of using endophytes that are heterologously disposed to plants to impart novel characteristics to a host plant, as well as to distinct plant elements of the host plant. In some embodiments, endophyte compositions are isolated and purified from plant or fungal sources, and heterologously disposed with a plant element to impart improved agronomic potential and/or improved agronomic traits to the host plant. In some embodiments, endophytes that are capable of living within plants are isolated and purified from their native source(s) and

heterologously disposed, e.g., manually, mechanically, or artificially combined, with a plant element, to impart improved agronomic potential and/or improved agronomic traits to the host plant or the host plant's elements. Such endophytes that are capable of living within plants may be further manipulated or combined with additional elements prior to combining with the plant element(s).

[00208] As described herein, endophytes can be robustly derived from heterologous, homologous, or engineered sources, optionally cultured, manually, mechanically or artificially applied heterologously to plant elements, e.g., heterologously disposed, and, as a result of the manual, mechanical or artificial application, confer multiple beneficial properties. This is surprising given the variability observed in the art in endophytic microbe isolation and the previous observations of inefficient plant element pathogen colonization of plant host's tissues.

[00209] In part, the present invention provides preparations of endophytes that are capable of living within plants, and the creation of synthetic compositions of plant elements and/or seedlings with heterologously disposed endophytes, and formulations comprising the synthetic compositions, as well as the recognition that such synthetic compositions display a diversity of beneficial and unexpected properties present in the agricultural plants and/or the heterologous endophyte populations. Beneficial properties include, but are not limited to metabolism, transcript expression, proteome alterations, morphology, resilience to a variety of environmental stresses, and any combination of such properties. The present invention also provides methods of using endophytes described herein to benefit the host plant with which they are heterologously disposed.

Definitions

[00210] Terms used in the claims and specification are defined as set forth below unless otherwise specified.

[00211] It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise.

[00212] An "endophyte" is an organism capable of living on a plant element (e.g., rhizoplane or phyllosphere) or within a plant element, or on a surface in close physical proximity with a plant element, e.g., the rhizosphere, or e.g., on a seed. A "beneficial" endophytes does not cause disease or harm the host plant otherwise. Endophytes can occupy the intracellular or extracellular spaces of plant tissue, including the leaves, stems, flowers, fruits, seeds, or roots. An endophyte can be, for example, a bacterial or fungal organism, and can confer a beneficial

property to the host plant such as an increase in yield, biomass, resistance, or fitness. An endophyte can be a fungus or a bacterium. As used herein, the term “microbe” is sometimes used to describe an endophyte.

[00213] A “population” of endophytes, or an “endophyte population”, refers to one or more endophytes that share a common genetic derivation, e.g., one or more propagules of a single endophyte, i.e., endophytes grown from a single picked colony. In some embodiments, a population refers to endophytes of identical taxonomy. In some cases, a population of endophytes refers to one or more endophytes of the same genus. In some cases, a population of endophytes refers to one or more endophytes of the same OTU.

[00214] A “plurality of endophytes” means two or more types of endophyte entities, e.g., of bacteria or fungi, or combinations thereof. In some embodiments, the two or more types of endophyte entities are two or more individual endophytic organisms, regardless of genetic derivation or taxonomic relationship. In some embodiments, the two or more types of endophyte entities are two or more populations of endophytes. In other embodiments, the two or more types of endophyte entities are two or more species of endophytes. In yet other embodiments, the two or more types of endophyte entities are two or more genera of endophytes. In yet other embodiments, the two or more types of endophyte entities are two or more families of endophytes. In yet other embodiments, the two or more types of endophyte entities are two or more orders of endophytes. In yet other embodiments, the two or more types of endophyte entities are two or more classes of endophytes. In yet other embodiments, the two or more types of endophyte entities are two or more phyla of endophytes. In some embodiments, a plurality refers to three or more endophytes, either distinct individual organisms or distinct members of different genetic derivation or taxa. In some embodiments, a plurality refers to four or more either distinct individual endophytic organisms or distinct members of different genetic derivation or taxa. In some embodiments, a plurality refers to five or more, ten or more, or an even greater number of either distinct individual endophytic organisms or distinct members of different genetic derivation or taxa. In some embodiments, the term “consortium” or “consortia” may be used as a collective noun synonymous with “plurality”, when describing more than one population, species, genus, family, order, class, or phylum of endophytes.

[00215] As used herein, the term “microbe” or “microorganism” refers to any species or taxon of microorganism, including, but not limited to, archaea, bacteria, microalgae, fungi (including mold and yeast species), mycoplasmas, microspores, nanobacteria, oomycetes, and protozoa. In some embodiments, a microbe or microorganism is an endophyte, for example a

bacterial or fungal endophyte, which is capable of living within a plant. In some embodiments, a microbe or microorganism encompasses individual cells (e.g., unicellular microorganisms) or more than one cell (e.g., multi-cellular microorganism). A “population of microorganisms” may thus refer to multiple cells of a single microorganism, in which the cells share common genetic derivation.

[00216] As used herein, the term “bacterium” or “bacteria” refers in general to any prokaryotic organism, and may reference an organism from either Kingdom Eubacteria (Bacteria), Kingdom Archaeobacteria (Archae), or both. In some cases, bacterial genera have been reassigned due to various reasons (such as, but not limited to, the evolving field of whole genome sequencing), and it is understood that such nomenclature reassignments are within the scope of any claimed genus. For example, certain species of the genus *Erwinia* have been described in the literature as belonging to genus *Pantoea* (Zhang, Y. & Qiu, S. Antonie van Leeuwenhoek (2015) 108: 1037).

[00217] The term 16S refers to the DNA sequence of the 16S ribosomal RNA (rRNA) sequence of a bacterium. 16S rRNA gene sequencing is a well-established method for studying phylogeny and taxonomy of bacteria.

[00218] As used herein, the term “fungus” or “fungi” refers in general to any organism from Kingdom Fungi. Historical taxonomic classification of fungi has been according to morphological presentation. Beginning in the mid-1800’s, it was recognized that some fungi have a pleomorphic life cycle, and that different nomenclature designations were being used for different forms of the same fungus. In 1981, the Sydney Congress of the International Mycological Association laid out rules for the naming of fungi according to their status as anamorph, teleomorph, or holomorph (Taylor JW. One Fungus = One Name: DNA and fungal nomenclature twenty years after PCR. IMA Fungus 2(2):113-120. 2011.). With the development of genomic sequencing, it became evident that taxonomic classification based on molecular phylogenetics did not align with morphological-based nomenclature (Shenoy BD, Jeewon R, Hyde KD. Impact of DNA sequence-data on the taxonomy of anamorphic fungi. Fungal Diversity 26(10) 1-54. 2007). As a result, in 2011 the International Botanical Congress adopted a resolution approving the International Code of Nomenclature for Algae, Fungi, and Plants (Melbourne Code) (International Code of Nomenclature for algae, fungi, and plants (Melbourne Code), adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011), with the stated outcome of designating “One Fungus = One Name” (Hawksworth DL. Managing and coping with names of pleomorphic fungi in a period of transition. IMA Fungus 3(1):15-24. 2012.). However, systematics experts have not aligned

on common nomenclature for all fungi, nor are all existing databases and information resources inclusive of updated taxonomies. As such, many fungi referenced herein may be described by their anamorph form but it is understood that based on identical genomic sequencing, any pleomorphic state of that fungus may be considered to be the same organism. For example, the genus *Alternaria* is the anamorph form of the teleomorph genus *Lewia* (Kwasna H and Kosiak B. *Lewia avenicola* sp. nov. and its *Alternaria* anamorph from oat grain, with a key to the species of *Lewia*. Mycol Res 2003; 107(Pt 3):371-6.), ergo both would be understood to be the same organism with the same DNA sequence. For example, it is understood that the genus *Acremonium* is also reported in the literature as genus *Sarocladium* as well as genus *Tilachilidium* (Summerbell R. C., C. Gueidan, H-J. Schroers³, G.S. de Hoog, M. Starink, Y. Arocha Rosete, J. Guarro and J.A. Scott. *Acremonium* phylogenetic overview and revision of *Gliomastix*, *Sarocladium*, and *Trichothecium*. Studies in Mycology 68: 139–162. 2011.). For example, the genus *Cladosporium* is an anamorph of the teleomorph genus *Davidiella* (Bensch K, Braun U, Groenewald JZ, Crous PW. The genus *Cladosporium*. Stud Mycol. 2012 Jun 15; 72(1): 1–401.), and is understood to describe the same organism. In some cases, fungal genera have been reassigned due to various reasons, and it is understood that such nomenclature reassignments are within the scope of any claimed genus.

[00219] “Internal Transcribed Spacer” (ITS) refers to the spacer DNA (non-coding DNA) situated between the small-subunit ribosomal RNA (rRNA) and large-subunit (LSU) rRNA genes in the chromosome or the corresponding transcribed region in the polycistronic rRNA precursor transcript. ITS gene sequencing is a well-established method for studying phylogeny and taxonomy of fungi. In some cases, the “Large SubUnit” (LSU) sequence is used to identify fungi. LSU gene sequencing is a well-established method for studying phylogeny and taxonomy of fungi. Some fungal endophytes may be described by an ITS sequence and some may be described by an LSU sequence. Both are understood to be equally descriptive and accurate for determining taxonomy.

[00220] As used herein with respect to fungi and bacteria, the term “marker gene” refers to a conserved gene comprising sequence variation among related organisms, e.g. an organism’s 16S (for bacteria) or ITS (for fungi) polynucleotide sequence, *fusA* gene, or unique genomic regions, by which a microbe may be specifically identified and assigned taxonomic nomenclature. In some embodiments, marker genes include, but are not limited to, actin, elongation factor G (*fusA*), tubulin, largest subunit of RNA polymerase II (RPB1), long

subunit rRNA gene (LSU), second largest subunit of RNA polymerase II (RPB2), small subunit rRNA gene (SSU), phosphoglycerate kinase, beta-tubulin, and combinations thereof.

[00221] The terms “pathogen” and “pathogenic” in reference to a bacterium or fungus includes any such organism that is capable of causing or affecting a disease, disorder or condition of a host comprising the organism.

[00222] A “spore” or a population of “spores” refers to bacteria or fungi that are generally viable, more resistant to environmental influences such as heat and bactericidal or fungicidal agents than other forms of the same bacteria or fungi, and typically capable of germination and out-growth. Bacteria and fungi that are “capable of forming spores” are those bacteria and fungi comprising the genes and other necessary abilities to produce spores under suitable environmental conditions.

[00223] “Biomass” means the total mass or weight (fresh or dry), at a given time, of a plant tissue, plant tissues, an entire plant, or population of plants. Biomass is usually given as weight per unit area. The term may also refer to all the plants or species in the community (community biomass).

[00224] The term “isolated” is intended to specifically reference an organism, cell, tissue, polynucleotide, or polypeptide that is removed from its original source.

[00225] As used herein, an isolated endophyte or microbe is an endophyte or microbe that has been removed from its natural milieu. “Pure cultures” or “isolated cultures” are cultures in which the organisms present are only of one particular genus and species. This is in contrast to “mixed cultures,” which are cultures in which more than one genus and/or species of microorganism are present. As such, the term “isolated” does not necessarily reflect the extent to which the microbe has been purified. A “substantially pure culture” of the microbe refers to a culture which contains substantially no other endophytes or microbes than the desired endophyte or microbe. In other words, a substantially pure endophyte or microbe culture is substantially free of other contaminants, which can include microbial contaminants. Further, as used herein, “biologically pure” is intended to mean the endophyte or microbe separated from materials with which it is normally found in nature. A microbe or endophyte heterologously disposed to other microbes or endophytes, or with compounds or materials that it is not normally found with in nature, is still defined as “biologically pure.” A monoculture is, of course, “biologically pure.” As used herein, the term “enriched culture” of an isolated microbe or endophyte refers to a culture that contains more than 50%, 60%, 70%, 80%, 90%, or 95% of the isolated endophyte or microbe.

[00226] A “host plant” includes any plant, particularly a plant of agronomic importance, within which or onto which a microbe, such as an endophyte, is heterologously disposed. As used herein, a microbe is said to colonize a plant, plant element, or seed, when it can exist as an endophyte in relationship with a plant or plant element during at least part of either the plant’s or the microbe’s life cycle. In some embodiments, an endophyte is said to “colonize” a plant or plant element when it can be stably detected within the plant or plant element over a period time, such as one or more days, weeks, months or years. Some of the compositions and methods described herein involve a plurality of microbes in an amount effective to colonize a plant.

[00227] A “non-host target” means an organism or chemical compound that is altered in some way after contacting a host plant that comprises an endophyte, as a result of a property conferred to the host plant by the endophyte.

[00228] As used herein, a nucleic acid has “homology” or is “homologous” to a second nucleic acid if the nucleic acid sequence has a similar sequence to the second nucleic acid sequence. The terms “identity”, “percent identity”, “percent sequence identity” or “identical” in the context of nucleic acid sequences refer to the nucleotides in the two sequences that are the same when aligned for maximum correspondence. There are different algorithms known in the art that can be used to measure nucleotide sequence identity. Nucleotide sequence identity can be measured by a local or global alignment, preferably implementing an optimal local or optimal global alignment algorithm. For example, a global alignment may be generated using an implementation of the Needleman-Wunsch algorithm (Needleman, S.B. & Wunsch, C.D. (1970) *Journal of Molecular Biology*. 48(3):443-53). For example, a local alignment may be generated using an implementation of the Smith-Waterman algorithm (Smith T.F & Waterman, M.S. (1981) *Journal of Molecular Biology*. 147(1):195-197). Optimal global alignments using the Needleman-Wunsch algorithm and optimal local alignments using the Smith-Waterman algorithm are implemented in USEARCH, for example USEARCH version v8.1.1756_i86osx32.

[00229] A gap is a region of an alignment wherein a sequence does not align to a position in the other sequence of the alignment. In global alignments, terminal gaps are discarded before identity is calculated. For both local and global alignments, internal gaps are counted as differences. A terminal gap is a region beginning at the end of a sequence in an alignment wherein the nucleotide in the terminal position of that sequence does not correspond to a nucleotide position in the other sequence of the alignment and extending for all contiguous positions in that sequence wherein the nucleotides of that sequence do not correspond to a

nucleotide position in the other sequence of the alignment. An internal gap is a gap in an alignment which is flanked on the 3' and 5' end by positions wherein the aligned sequences are identical.

[00230] In some embodiments, the nucleic acid sequence to be aligned is a complete gene. In some embodiments, the nucleic acid sequence to be aligned is a gene fragment. In some embodiments, the nucleic acid sequence to be aligned is an intergenic sequence. In a preferred embodiment, inference of homology from a sequence alignment is made where the region of alignment is at least 85% of the length of the query sequence.

[00231] The term “substantial homology” or “substantial similarity,” when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 76%, 80%, 85%, or at least about 90%, or at least about 95%, 96%, at least 97%, 98%, 99% or 100% of the positions of the alignment, wherein the region of alignment is at least about 50%, 60%, 70%, 75%, 85%, or at least about 90%, or at least about 95%, 96%, 97%, 98%, 99% or 100% of the length of the query sequence. In a preferred embodiment, the region of alignment contains at least 100 positions inclusive of any internal gaps. In some embodiments, the region of alignment comprises at least 100 nucleotides of the query sequence. In some embodiments, the region of alignment comprises at least 200 nucleotides of the query sequence. In some embodiments, the region of alignment comprises at least 300 nucleotides of the query sequence. In some embodiments, the region of alignment comprises at least 400 nucleotides of the query sequence. In some embodiments, the region of alignment comprises at least 500 nucleotides of the query sequence. In some embodiments, the query sequence is selected from the group consisting of SEQ ID NOs 38-59.

[00232] As used herein, the terms “operational taxonomic unit,” “OTU,” “taxon,” “hierarchical cluster,” and “cluster” are used interchangeably. An operational taxonomic unit (OTU) refers to a group of one or more organisms that comprises a node in a clustering tree. The level of a cluster is determined by its hierarchical order. In some embodiments, an OTU is a group tentatively assumed to be a valid taxon for purposes of phylogenetic analysis. In another embodiment, an OTU is any of the extant taxonomic units under study. In yet another embodiment, an OTU is given a name and a rank. For example, an OTU can represent a domain, a sub-domain, a kingdom, a sub-kingdom, a phylum, a sub-phylum, a class, a subclass, an order, a sub-order, a family, a subfamily, a genus, a subgenus, or a species. In some embodiments, OTUs can represent one or more organisms from the kingdoms Eubacteria,

Bacteria, Protista, or Fungi, at any level of a hierarchical order. In some embodiments, an OTU represents a prokaryotic or fungal order.

[00233] In some embodiments, the present invention contemplates the synthetic compositions comprising the combination of a plant element, seedling, or whole plants and an endophyte population, in which the endophyte population is “heterologously disposed”. In some embodiments, one or more endophytes of the synthetic composition are heterologously disposed when they are mechanically or manually applied, artificially inoculated or disposed onto or into a plant element, seedling, plant or onto or into a plant growth medium or onto or into a treatment formulation so that the endophyte exists on or in said plant element, seedling, plant, plant growth medium, or treatment formulation in a manner not found in nature prior to the application of the one or more endophytes, e.g., said combination which is not found in nature. In some embodiments, such a manner is contemplated to be selected from the group consisting of: the presence of the endophyte; presence of the endophyte in a different number of cells, concentration, or amount; the presence of the endophyte in a different plant element, tissue, cell type, or other physical location in or on the plant; the presence of the endophyte at different time period, e.g. developmental phase of the plant or plant element, time of day, time of season, and combinations thereof. In some embodiments, one or more endophytes of a synthetic composition are heterologously disposed when the one or more endophytes are artificially inoculated, e.g., is manually or mechanically inoculated, or artificially applied, e.g. manually or mechanically applied, to a different plant element or at a different developmental stage than that with which the one or more endophytes are naturally found or at a greater concentration, number, or amount than that which is naturally found in or on said plant element, seedling, or plant. In some embodiments, “heterologously disposed” refers to the relationship between the endophyte and the inoculated host plant as compared to the type of host plant with which said endophyte is normally associated. In one example, endophytes used in a synthetic composition can be obtained from a different individual plant of the same variety as that of the host inoculated plant to which it becomes heterologously disposed, a plant of a different variety but the same genus and species, a plant of a different cultivar, or a plant of a different genus. In an embodiment, the endophyte is an endophytic microbe isolated from a different plant than the inoculated plant. For example, in an embodiment, the endophyte is an endophyte isolated from a different plant of the same species as the inoculated plant. In some cases, the endophyte is isolated from a species related to the inoculated plant. In another example, endophytes used in a synthetic composition can be obtained from different individual plants of the same variety, each of which has been

subjected to different growth conditions. For example, an endophyte derived from a drought-affected plant of one variety can be isolated and coated onto the plant element that was derived from a plant of the same variety not subjected to drought. In such cases, the endophyte is considered to be heterologously disposed to the plant element onto which it is manually, mechanically, or artificially applied. In some embodiments, “heterologously disposed” means that the endophyte applied to a different tissue or cell type of the plant element than that in which the microbe is naturally found. In some embodiments, an endophyte is heterologously disposed on a seedling if that endophyte is normally found at the flowering stage of a plant and not at a seedling stage. In some embodiments, an endophyte is heterologously disposed the endophyte is normally found in the root tissue of a plant element but not in the leaf tissue, and the endophyte is applied to the leaf. In yet another non-limiting example, if an endophyte is naturally found in the mesophyll layer of leaf tissue but is being applied to the epithelial layer, the endophyte would be considered to be heterologously disposed. In some embodiments, “heterologously disposed” means that the native plant element, seedling, or plant does not contain detectable levels of the microbe in that same plant element, seedling, or plant. For example, if said plant element or seedling or plant does not naturally have the endophyte heterologously disposed to it and the endophyte is applied, the endophyte would be considered to be heterologously disposed. In some embodiments, “heterologously disposed” means that the endophyte being applied is at a greater concentration, number, or amount to the plant element, seedling, or plant, than that which is naturally found in said plant element, seedling, or plant. For example, an endophyte is heterologously disposed when present at a concentration that is at least 1.5 times greater, between 1.5 and 2 times greater, 2 times greater, between 2 and 3 times greater, 3 times greater, between 3 and 5 times greater, 5 times greater, between 5 and 7 times greater, 7 times greater, between 7 and 10 times greater, 10 times greater, or even greater than 10 times higher number, amount, or concentration than the concentration that was present prior to the disposition of said endophyte. In some embodiments, “heterologously disposed” means that the endophyte is applied to a developmental stage of the plant element, seedling, or plant in which said endophyte is not naturally found, but may be associated at other stages. In some embodiments, “heterologously disposed” means that the endophyte was isolated from plants or plant elements under an environmental condition different than that which is normally found (for example but not limited to: different soil pH, different mean air temperature, different soil temperature, different rainfall conditions, different soil nutrient composition, or different environmental salinity). In one example, if an endophyte is normally found at the

flowering stage of a plant and no other stage, an endophyte applied at the seedling stage may be considered to be heterologously disposed. In another example, an endophyte that is normally heterologously disposed to leaf tissue of a plant is considered heterologous to a leaf tissue of another plant that naturally lacks said endophyte. In another example, an endophyte that is normally found at low levels in a plant is considered heterologous to that plant if a higher concentration of that endophyte is introduced into the plant. In yet another example, an endophyte that is heterologously disposed to a tropical grass species would be considered heterologous to a different grass species that naturally lacks said endophyte.

[00234] An “inoculated” plant or plant element has been artificially introduced to a heterologous endophyte at some point during the plant’s or plant element’s growth or development (including vegetative or reproductive phases). In some embodiments, the heterologous endophyte is transiently or permanently incorporated into the plant or plant element, and is detectable using methods known in the art or described herein. In some embodiments, a seed is inoculated with an endophyte by manually or mechanically contacting the seed with a formulation comprising said endophyte, which is detectable in or on the seed. In some embodiments, a plant is said to be inoculated with an endophyte if it is grown from a reproductive element (*e.g.* a seed) that was itself manually or mechanically contacted with a formulation comprising said endophyte, which is subsequently detectable in or on the plant. In some embodiments, a plant is said to be inoculated with an endophyte if any one or more of its plant elements (*e.g.*, leaf, stem, or root) is manually or mechanically contacted with a formulation comprising said endophyte, which is subsequently detectable either in the same plant element that was originally contacted with said formulation or in a different plant element of that plant. The term “inoculation” may also refer to the manual or mechanical contact of an endophyte population to any substance, that is detectable in or on said substance subsequent to endophyte contact. In one example, said substance is soil or other plant growth medium. In another example, said substance is a storage medium such as glycerol. In some cases, “inoculation” may refer to the contact of an endophyte population to a non-plant living organism, for example, but not limited to, an insect or a fungus.

[00235] The term “isoline” is a comparative term, and references organisms that are genetically identical, but may differ in treatment. In one example, two genetically identical maize plant embryos may be separated into two different groups, one receiving a treatment (such as transformation with a heterologous polynucleotide, to create a genetically modified plant) and one control, *e.g.*, reference, that does not receive such treatment. Any phenotypic differences between the two groups may thus be attributed solely to the treatment and not to

any inherency of the plant's genetic makeup. In another example, two genetically identical soybean seeds may be treated with a formulation, one that introduces an endophyte composition and one that does not. Any phenotypic differences between the plants derived from (e.g., grown from or obtained from) those seeds may be attributed to the endophyte treatment, thus forming an isoline comparison.

[00236] Similarly, by the term “reference agricultural plant,” it is meant an agricultural plant of the same species, variety, or cultivar to which a treatment, formulation, composition or endophyte preparation as described herein is not administered/contacted. A reference agricultural plant, therefore, is identical to the treated plant with the exception of the presence of the endophyte and can serve as a control for detecting the effects of the endophyte that is conferred to the plant. In some embodiments, the phrase “reference isoline plant” is used herein to describe a reference plant that is genetically identical and subject to the same conditions, i.e., a control plant, to the treated plant.

[00237] A “reference environment” refers to the environment, treatment or condition of the plant in which a measurement is made. For example, production of a compound in a plant heterologously disposed to an endophyte can be measured in a reference environment of drought stress, and compared with the levels of the compound in a reference agricultural plant under the same conditions of drought stress. Alternatively, the levels of a compound in plant heterologously disposed to an endophyte and reference agricultural plant can be measured under identical conditions of no stress.

[00238] A “plant element” is intended to generically reference either a whole plant or a plant component, including but not limited to plant tissues, parts, and cell types. A plant element is preferably one of the following: whole plant, seedling, meristematic tissue, ground tissue, vascular tissue, dermal tissue, seed, leaf, root, shoot, stem, flower, fruit, stolon, bulb, tuber, corm, keikis, shoot, bud. As used herein, a “plant element” is synonymous to a “portion” of a plant, and refers to any part of the plant, and can include distinct tissues and/or organs, and may be used interchangeably with the term “tissue” throughout.

[00239] Similarly, a “plant reproductive element” is intended to generically reference any part of a plant that is able to initiate other plants via either sexual or asexual reproduction of that plant, for example but not limited to: seed, seedling, root, shoot, cutting, scion, graft, stolon, bulb, tuber, corm, keikis, or bud.

[00240] A “progeny seed”, as used herein, refers to the seed produced by a host plant that has been inoculated with, or heterologously disposed to, an endophyte. For example, in the present invention, a seed, plant element, or whole plant may become heterologously disposed

to an endophyte, and the plant that is grown from said seed, or plant that is grown in heterologous association with said endophyte, may itself produce progeny seeds that comprise altered nutritional composition compared to seeds obtained from plants that were not grown from a plant element heterologously disposed to an endophyte or obtained from a parental (host) plant that had become heterologously disposed to an endophyte at some point in its life cycle. In the general sense, the phrase “progeny seed” may be construed to represent any plant propagative unit produced by the host plant that is capable of becoming another individual of that same plant species.

[00241] A “population” of plants refers to more than one plant, that are of the same taxonomic category, typically be of the same species, and will also typically share a common genetic derivation.

[00242] As used herein, an “agricultural seed” is a seed used to grow a plant typically used in agriculture (an “agricultural plant”). The seed may be of a monocot plant, and may be planted for the production of an agricultural product, for example feed, food, fiber, fuel, industrial uses, etc. As used herein, an agricultural seed is a seed that is prepared for planting, for example, in farms for growing.

[00243] “Agricultural plants” or “plants of agronomic importance include plants that are cultivated by humans for food, feed, fiber, fuel, and/or industrial purposes. In some embodiments, plants (including seeds and other plant elements) treated in accordance with the present invention are monocots. In some embodiments, plants (including seeds or other plant elements) treated in accordance with the present invention are dicots, excluding cotton and sorghum. In some embodiments, plants treated in accordance with the present invention include, but are not limited to: agricultural row, agricultural grass plants or other field crops: wheat, rice, barley, buckwheat, beans (soybean, snap, dry), corn (grain, seed, sweet corn, silage, popcorn, high oil), canola, peas (dry, succulent), peanuts, safflower, sunflower, alfalfa hay, forage crops (alfalfa, clover, vetch, and trefoil), berries and small fruits (blackberries, blueberries, currants, elderberries, gooseberries, huckleberries, loganberries, raspberries, strawberries, bananas and grapes), bulb crops (garlic, leeks, onions, shallots, and ornamental bulbs), citrus fruits (citrus hybrids, grapefruit, kumquat, lines, oranges, and pummelos), cucurbit vegetables (cucumbers, melons, gourds, pumpkins, and squash), flowers, bedding plants, ornamentals, fruiting vegetables (eggplant, sweet and hot peppers, tomatillos, and tomatoes), herbs, spices, mints, hydroponic crops (cucumbers, tomatoes, lettuce, herbs, and spices), leafy vegetables and cole crops (arugula, celery, chervil, endive, fennel, lettuce (head and leaf), parsley, radicchio, rhubarb, spinach, Swiss chard, broccoli, Brussels sprouts,

cabbage, cauliflower, collards, kale, kohlrabi, and mustard greens), asparagus, legume vegetable and field crops (snap and dry beans, lentils, succulent and dry peas, and peanuts), pome fruit (pears and quince), root crops (beets, sugarbeets, red beets, carrots, celeriac, chicory, horseradish, parsnip, radish rutabaga, salsify, and turnips), deciduous trees (maple and oak), pine, rye, wheat, millet, stone fruits (apricots, cherries, nectarines, peaches, plums, and prunes), tree nuts (almonds, beech nuts, Brazil nuts, butternuts, cashews, chestnuts, filberts, hickory nuts, macadamia nuts, pecans, pistachios, and walnuts), and tuber crops (potatoes, sweet potatoes, yams, artichoke, cassava, and ginger). In a particular embodiment, the agricultural plant is selected from the group consisting of rice (*Oryza sativa* and related varieties), soy (*Glycine max* and related varieties), wheat (*Triticum aestivum* and related varieties), corn (*Zea mays* and related varieties), peanuts (*Arachis hypogaea* and related varieties), canola (*Brassica napus*, *Brassica rapa* and related varieties), coffee (*Coffea* spp.), cocoa (*Theobroma cacao*), melons, and tomatoes (*Solanum lycopersicum* and related varieties).

[00244] A “closely related variety” comprises a common genetic derivation with a plant variety. In some embodiments, a closely related variety has at least one grandparental line in common with the plant variety. In some embodiments, a closely related variety has at least one parental line in common with the plant variety. In some embodiments, a closely related variety has at least 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.7%, 99.9%, 99.99% of the same SNPs detected in the plant variety. In some embodiments, a closely related variety has at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or 20 or more of the same SNPs detected in the plant variety. In some embodiments, a closely related variety has at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, at least 200, at least 300, at least 400, at least 500, at least 600, at least 700, at least 800, at least 900, or 100 or more of the same SNPs detected in the plant variety. In some embodiments, a closely related variety has at least 1000, at least 2000, at least 3000, at least 4000, at least 5000, at least 6000, at least 7000, at least 8000, at least 9000, or 10000 or more of the same SNPs detected in the plant variety. In some embodiments, a closely related variety has at least 10000, at least 20000, at least 30000, at least 40000, at least 50000, at least 60000, at least 70000, at least 80000, at least 90000, or 100000 or more of the same SNPs detected in the plant variety.

[00245] A “synthetic composition” comprises one or more endophytes combined by human endeavor with a heterologously disposed plant element or a treatment formulation, said combination which is not found in nature. In some embodiments, the term “synthetic composition” means one or more plant elements or formulation components combined by human endeavor with an isolated, purified endophyte composition. In some embodiments, said purified endophyte composition is mechanically or manually applied, artificially inoculated or disposed on a plant element in a manner that is not found on or in the plant element before application of the purified endophyte composition, e.g., said combination or association which is not found in nature.

[00246] In some embodiments, “synthetic composition” is used to refer to a treatment formulation comprising an isolated, purified population of endophytes heterologously disposed to a plant element. In some embodiments, “synthetic composition” refers to a purified population of endophytes in a treatment formulation comprising additional compositions with which said endophytes are not found in nature.

[00247] A “treatment formulation” refers to a mixture of chemicals that facilitate the stability, storage, and/or application of the endophyte composition(s). Treatment formulations may comprise any one or more agents such as: surfactant, a buffer, a tackifier, a microbial stabilizer, a fungicide, an anticomplex agent, an herbicide, a nematocide, an insecticide, a plant growth regulator, a rodenticide, a desiccant, a nutrient, an excipient, a wetting agent, a salt.

[00248] In some embodiments, an “agriculturally compatible carrier” can be used to formulate an agricultural formulation or other composition that includes a purified endophyte preparation. As used herein an “agriculturally compatible carrier” refers to any material, other than water, that can be added to a plant element without causing or having an adverse effect on the plant element (e.g., reducing seed germination) or the plant that grows from the plant element, or the like.

[00249] “Plant health” is demonstrated by the presence or improvement of a trait of agronomic importance found in a plant or plant element as compared to a reference plant or plant element. The compositions and methods herein may provide for an improved “agronomic trait” or “trait of agronomic importance” to a host plant, which include, but are not limited to disease resistance, drought tolerance, heat tolerance, cold tolerance, salinity tolerance, metal tolerance, herbicide tolerance, improved water use efficiency, improved nitrogen utilization, improved nitrogen fixation, pest resistance, herbivore resistance, pathogen resistance, yield improvement, health enhancement, vigor improvement, growth

improvement, photosynthetic capability improvement, nutrition enhancement, altered protein content, altered oil content, increased biomass, increased shoot length, increased root length, increased root area, improved root architecture, modulation of a metabolite, modulation of the proteome, increased seed weight, altered seed carbohydrate composition, altered seed oil composition, altered seed protein composition, altered seed nutrient composition, and combinations thereof, as compared to reference plant derived from a seed without said seed treatment formulation.

[00250] In some embodiments, a treatment is heterologously disposed on a plant element in an amount effective to improve a trait of agronomic importance. In some embodiments, treatments capable of improving plant health are applied in an amount effective to improve a trait of agronomic importance or tolerance by at least 0.1%, at least 0.5%, at least 1%, at least 2%, at least 3%, between 3% and 5%, at least 5%, between 5% and 10%, least 10%, between 10% and 15%, for example at least 15%, between 15% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 75%, at least 75%, between 75% and 100%, at least 100%, between 100% and 150%, at least 150%, between 150% and 200%, at least 200%, between 200% and 300%, at least 300% or more, as compared to a reference plant element not further comprising said endophyte.

[00251] In some embodiments, an improvement in a trait of agronomic importance is measured by the “win rate”. The win rate is the proportion of replicates where the treatment shows an improvement in a trait of agronomic importance relative to reference replicates. In some embodiments, replicates are individual plants. In some embodiments, replicates are plots, e.g. replicated plots within a randomized complete block design field trial. In some embodiments, replicates are field trials conducted at diverse geographies.

[00252] In some embodiments, the endophyte is capable of improving a trait of agronomic importance at concentrations detected on or in the treated plant element of at least 10^2 CFU or spores per plant element, between 10^2 and 10^3 CFU or spores per plant element, about 10^3 CFU or spores per plant element, between 10^3 and 10^4 CFU or spores per plant element, about 10^4 CFU or spores per plant element, or between 10^4 , of about 10^5 CFU or spores per plant element, at least 10^5 CFU or spores per plant element, between 10^5 and 10^6 CFU or spores per plant element, about 10^6 CFU or spores per plant element, between 10^6 and 10^7 CFU or spores per plant element, about 10^7 CFU or spores per plant element, between 10^7 and 10^8 CFU or spores per plant element, about 10^8 CFU or spores

per plant element, or even greater than 10^8 CFU or spores per plant element. In some embodiments, the plant element is a seed.

[00253] The phrase “nutritional quality trait” includes any measurable parameter of a seed that either directly or indirectly influences the value (nutritional or economic) of said seed, for example, but not limited to: protein, fat, carbohydrate, ash, moisture, fiber, and calories. In some cases, “nutritional quality trait” is synonymous with “nutritional quality trait” or “seed nutritional quality trait”, and can refer to any composition of the associated plant element, most particularly compositions providing benefit to other organisms that consume or utilize said plant element. As used herein, “oil” and “fat” are used interchangeably.

[00254] An increased “seed yield” can refer to any increase in seed or fruit weight, size, or abundance per a unit of measure, for example, per plant, per number of plants, per mass of plants, per acre planted, per acre harvested. In some embodiments, seed yield is reported as pounds or bushels of seed produced per acre harvested. The terms seed and grain are used interchangeably herein. Yield may also refer to the recovery of a particular component of a plant tissue upon processing, for example, the amount of oil which can be extracted per unit of seed. Typically, the particular characteristic is designated when referring to increased yield, e.g., increased seed yield or increased oil yield. Where the characteristic is not specified it may be assumed yield refers to seed yield and the terms may be used interchangeably.

[00255] As used herein, the terms “water-limited condition” and “drought condition,” or “water-limited” and “drought,” may be used interchangeably. For example, a method or composition for improving a plant’s ability to grow under drought conditions means the same as the ability to grow under water-limited conditions. In such cases, the plant can be further said to display improved tolerance to drought stress.

[00256] As used herein, the terms “normal watering” and “well-watered” are used interchangeably, to describe a plant grown under typical growth conditions with no water restriction.

[00257] Additionally, “altered metabolic function” or “altered enzymatic function” may include, but not be limited to, the following: altered production of an auxin, altered nitrogen fixation, altered production of an antimicrobial compound, altered production of a siderophore, altered mineral phosphate solubilization, altered production of a cellulase, altered production of a chitinase, altered production of a xylanase, altered production of acetoin, altered utilization of a carbon source.

[00258] “Nutrient” or “seed nutrient” refers to any composition of the associated plant element, most particularly compositions providing benefit to other organisms that consume or utilize said plant element.

[00259] “Agronomic trait potential” is intended to mean a capability of a plant element for exhibiting a phenotype, preferably an improved agronomic trait, at some point during its life cycle, or conveying said phenotype to another plant element with which it is associated in the same plant. For example, a plant element may comprise an endophyte that will provide benefit to leaf tissue of a plant from which the plant element is grown; in such case, the plant element comprising such endophyte has the agronomic trait potential for a particular phenotype (for example, increased biomass in the plant) even if the plant element itself does not display said phenotype.

[00260] In some cases, the present invention contemplates the use of compositions that are “compatible” with agricultural chemicals, including but not limited to, a fungicide, an anti-complex compound, a bactericide, a virucide, an herbicide, a nematocide, a parasiticide, a pesticide, or any other agent widely used in agriculture which has the effect of killing or otherwise interfering with optimal growth of another organism. As used herein, a composition is “compatible” with an agricultural chemical when the organism is modified, such as by genetic modification, e.g., contains a transgene that confers resistance to an herbicide, or is adapted to grow in, or otherwise survive, the concentration of the agricultural chemical used in agriculture. For example, an endophyte disposed on the surface of a plant element is compatible with the fungicide metalaxyl if it is able to survive the concentrations that are applied on the plant element surface.

[00261] As used herein, a “colony-forming unit” (“CFU”) is used as a measure of viable microorganisms in a sample. A CFU is an individual viable cell capable of forming on a solid medium a visible colony whose individual cells are derived by cell division from one parental cell. In some embodiments, a cell is a fungal spore.

[00262] The terms “decreased,” “fewer,” “slower” and “increased” “faster” “enhanced” “greater” as used herein refers to a decrease or increase in a characteristic of the endophyte treated plant element or resulting plant compared to an untreated plant element or resulting plant. For example, a decrease in a characteristic may be at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, between 5% and 10%, at least 10%, between 10% and 20%, at least 15%, at least 20%, between 20% and 30%, at least 25%, at least 30%, between 30% and 40%, at least 35%, at least 40%, between 40% and 50%, at least 45%, at least 50%, between 50% and 60%, at least about 60%, between 60% and 70%, between 70% and 80%, at least

75%, at least about 80%, between 80% and 90%, at least about 90%, between 90% and 100%, at least 100%, between 100% and 200%, at least 200%, at least about 300%, at least about 400% or more lower than the untreated control and an increase may be at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, between 5% and 10%, at least 10%, between 10% and 20%, at least 15%, at least 20%, between 20% and 30%, at least 25%, at least 30%, between 30% and 40%, at least 35%, at least 40%, between 40% and 50%, at least 45%, at least 50%, between 50% and 60%, at least about 60%, between 60% and 70%, between 70% and 80%, at least 75%, at least about 80%, between 80% and 90%, at least about 90%, between 90% and 100%, at least 100%, between 100% and 200%, at least 200%, at least about 300%, at least about 400% or more higher than the untreated control.

[00263] As used herein, a microbe or plant or plant element is “modified” when it comprises an artificially introduced genetic or epigenetic “modification”. In some embodiments, the modification is introduced by a genome engineering technology. In some embodiments, the modification is introduced by a targeted nuclease. In some embodiments, targeted nucleases include, but are not limited to, transcription activator-like effector nuclease (TALEN), zinc finger nuclease (ZNF), clustered regulatory interspaced short palindromic repeats (CRISPR), CRISPR/Cas9, CRISPR/CPF1, and combinations thereof. In some embodiments, the modification is an epigenetic modification. In some embodiments, the modification is introduced by treatment with a DNA methyltransferase inhibitor such as 5-azacytidine, or a histone deacetylase inhibitor such as 2-amino-7-methoxy-3H-phenoxazin-3-one. In some embodiments, the modification is introduced via tissue culture. In some embodiments, a modified microbe or plant or plant element comprises a transgene.

Endophyte Compositions

[00264] The endophytes described herein provide several unexpected and significant advantages to agricultural plants over other plant-associated microbes, as demonstrated in the Examples.

[00265] Novel endophyte compositions are described herein. In some embodiments, the endophyte is selected from Table 4. In some embodiments, the endophyte is selected from Table 6.

[00266] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 67. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5%

identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 67.

[00267] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 68. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 68.

[00268] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 69. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 69.

[00269] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 70. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 70.

[00270] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 65 and 66. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 65 and 66. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 65 and 66. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 65 and 66.

[00271] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID

NOs: 63, 64, and 71. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 63, 64, and 71. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 63, 64, and 71. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 63, 64, and 71. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

[00272] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 60, 61, and 62. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 60, 61, and 62. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 60, 61, and 62. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

[00273] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are at least 97% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

[00274] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least two

polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are at least 97% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45.

[00275] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group

consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are at least 97% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least five polynucleotide sequences that are at least 97% identical to at least five sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least five polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least five sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least six polynucleotide sequences that are at least 97% identical to at least six sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51. In some embodiments, the endophyte comprises at least six polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least six sequences selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

[00276] In some embodiments, the endophyte comprises a polynucleotide sequence that is at least 97% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises a polynucleotide sequence that is between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, at least 99.5% identical, or 100% identical to at least one sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are at least 97% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least two polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least two sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are at least 97% identical to

at least three sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least three polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least three sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are at least 97% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least four polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least four sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least five polynucleotide sequences that are at least 97% identical to at least five sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least five polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least five sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least six polynucleotide sequences that are at least 97% identical to at least six sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least six polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least six sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least seven polynucleotide sequences that are at least 97% identical to at least seven sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59. In some embodiments, the endophyte comprises at least seven polynucleotide sequences that are between 97% and 98% identical, at least 98% identical, between 98.0% identical and 99.5% identical, or at least 99.5% identical, or 100% identical to at least seven sequences selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

[00277] In some embodiments, the endophyte is a *Exserohilum rostrata*. In some embodiments, the endophyte is a *Curvularia spicifera*. In some embodiments, the endophyte

is a *Curvularia protuberata*. In some embodiments, the endophyte is an *Acremonium alternatum*. In some embodiments, the endophyte is a *Cladosporium oxysporum*. In some embodiments, the endophyte is a *Chaetomium globosum*. In some embodiments, the endophyte is a *Epicoccum nigrum*. In some embodiments, the endophyte is a *Paecilomyces inflatus*. In some embodiments, the endophyte is of the taxonomy *Coniochaeta prunicola*.

[00278] In some cases, the endophyte, or one or more components thereof, is of monoclonal origin, providing high genetic uniformity of the endophyte population in an agricultural formulation or within a plant element or synthetic combination with the endophyte.

[00279] In some embodiments, the endophyte can be cultured on a culture medium or can be adapted to culture on a culture medium.

[00280] The synthetic compositions provided herein are preferably stable. The endophyte may be shelf-stable, where at least 0.01%, of the CFUs are viable after storage in desiccated form (i.e., moisture content of 30% or less) for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or greater than 10 weeks at 4°C or at room temperature. Optionally, a shelf-stable formulation is in a dry formulation, a powder formulation, or a lyophilized formulation. In some embodiments, the formulation is formulated to provide stability for the population of endophytes. In an embodiment, the formulation is substantially stable at temperatures between about -20°C and about 50°C for at least about 1, 2, 3, 4, 5, or 6 days, or 1, 2, 3 or 4 weeks, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months, or one or more years. In another embodiment, the formulation is substantially stable at temperatures between about 4°C and about 37°C for at least about 5, 10, 15, 20, 25, 30 or greater than 30 days.

Endophytes and Synthetic Compositions with Plants and Plant Elements

[00281] It is contemplated that the methods and synthetic compositions may be used to improve a characteristic of agronomic importance to a plant.

[00282] The methods described herein can also be used with transgenic plants comprising one or more exogenous transgenes, for example, to yield additional trait benefits conferred by the newly introduced endophytic microbes.

[00283] For example, the endophyte may provide an improved benefit or tolerance to a plant that is of at least 3%, between 3% and 5%, at least 5%, between 5% and 10%, least 10%, between 10% and 15%, for example at least 15%, between 15% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 75%, at least 75%, between 75% and 100%, at least 100%, between 100% and 150%, at least 150%,

between 150% and 200%, at least 200%, between 200% and 300%, at least 300% or more, when compared with uninoculated plants grown under the same conditions.

[00284] In one embodiment, it is contemplated that the plant of the present invention is rice (*Oryza spp.*), in particular *O. sativa* and *O. glaberrima*, and members of the major *O. sativa* subspecies japonica, javanica, and indica. In one embodiment, it is contemplated that the plant of the present invention is the rice variety Rex, and closely related varieties thereof. In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a rice plant element or rice plant to which it is heterologously disposed.

[00285] In one embodiment, it is contemplated that the plant of the present invention is corn (*Zea spp.*), in particular *Zea mays ssp.* such as *Zea mays indenata*, *Zea mays indurata*, *Zea mays amylacea*, *Zea mays saccharata*, and *Zea mays everta*. In one embodiment, it is contemplated that the plant of the present invention is the corn variety Stine 9734, and closely related varieties thereof. In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a corn plant element or corn plant to which it is heterologously disposed.

[00286] In one embodiment, it is contemplated that the plant of the present invention is wheat (*Triticum spp.*) including species *T. aestivum* and *T. durum*. In one embodiment, it is contemplated that the plant of the present invention is hard red winter (HRW), hard red spring (HRS), hard white (HW), durum, soft white (SW), or soft red winter (SRW). In one embodiment, it is contemplated that the plant of the present invention is the wheat variety SDSU Focus, SDSU Select, and closely related varieties thereof. In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a wheat plant element or wheat plant to which it is heterologously disposed.

[00287] In one embodiment, it is contemplated that the plant of the present invention is soy (*Glycine max*). In one embodiment, it is contemplated that the plant of the present invention is the soy variety Dairyland DSR1808R2Y, Pfister 38R25, Stine 3920, Stine 33E22, or closely related varieties thereof.

[00288] In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a soy plant element or soy plant to which it is heterologously disposed.

[00289] In one embodiment, it is contemplated that the plant of the present invention is peanut (*Arachis hypogaea*). In one embodiment, it is contemplated that the plant of the present invention is the peanut variety AT9899, FloRun 107, Georgia-06G, Tamnut OL06, or

closely related varieties thereof. In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a peanut plant element or peanut plant to which it is heterologously disposed.

[00290] In one embodiment, it is contemplated that the plant of the present invention is a member of the genus *Brassica*. In one embodiment, it is contemplated that the plant of the present invention is *Brassica napus*. In one embodiment, it is contemplated that the plant of the present invention is a low erucic acid and low glucosinolate cultivar of *Brassica napus*. In one embodiment, it is contemplated that the plant of the present invention is canola. In one embodiment, it is contemplated that the plant of the present invention is the canola variety Brett Young 5525, NCC1015, or closely related varieties thereof. In some embodiments, the present invention contemplates the use of endophytes that can confer a beneficial agronomic trait upon a canola plant element or canola plant to which it is heterologously disposed.

[00291] In some cases, the endophytes described herein are capable of moving from one tissue type to another. For example, the present invention's detection and isolation of endophytes within the mature tissues of plants after coating on the exterior of a plant element demonstrates their ability to move from the plant element into the vegetative tissues of a maturing plant. Therefore, in some embodiments, the population of endophytes is capable of moving from the plant element exterior into the vegetative tissues of a plant. In some embodiments, the endophyte that is coated onto the plant element of a plant is capable, upon germination of the plant element into a vegetative state, of localizing to a different tissue of the plant. For example, endophytes can be capable of localizing to any one of the tissues in the plant, including: the root, adventitious root, seminal root, root hair, shoot, leaf, flower, bud, tassel, meristem, pollen, pistil, ovaries, stamen, fruit, stolon, rhizome, nodule, tuber, trichome, guard cells, hydathode, petal, sepal, glume, rachis, vascular cambium, phloem, and xylem. In an embodiment, the endophyte is capable of localizing to the root and/or the root hair of the plant. In another embodiment, the endophyte is capable of localizing to the photosynthetic tissues, for example, leaves and shoots of the plant. In other cases, the endophyte is localized to the vascular tissues of the plant, for example, in the xylem and phloem. In still another embodiment, the endophyte is capable of localizing to the reproductive tissues (flower, pollen, pistil, ovaries, stamen, fruit) of the plant. In another embodiment, the endophyte is capable of localizing to the root, shoots, leaves and reproductive tissues of the plant. In still another embodiment, the endophyte colonizes a fruit or plant element tissue of the plant. In still another embodiment, the endophyte is able to colonize the plant such that it is present in the surface of the plant (i.e., its presence is

detectably present on the plant exterior, or the episphere of the plant). In still other embodiments, the endophyte is capable of localizing to substantially all, or all, tissues of the plant. In certain embodiments, the endophyte is not localized to the root of a plant. In other cases, the endophyte is not localized to the photosynthetic tissues of the plant.

[00292] In some cases, endophytes are capable of replicating within the host plant and colonizing the plant.

[00293] The endophyte populations described herein are capable of colonizing a host plant. Successful colonization can be confirmed by detecting the presence of the endophyte population within the plant. For example, after applying the endophyte to the plant elements, high titers of the endophyte can be detected in the roots and shoots of the plants that germinate from the plant elements. Detecting the presence of the endophyte inside the plant can be accomplished by measuring the viability of the endophyte after surface sterilization of the plant element or the plant: endophyte colonization results in an internal localization of the endophyte, rendering it resistant to conditions of surface sterilization. The presence and quantity of endophyte can also be established using other means known in the art, for example, immunofluorescence microscopy using microbe-specific antibodies, or fluorescence in situ hybridization. Alternatively, specific nucleic acid probes recognizing conserved sequences from an endophyte can be employed to amplify a region, for example by quantitative PCR, and correlated to CFUs by means of a standard curve.

[00294] In another embodiment, the endophyte is heterologously disposed, for example, on the surface of a reproductive element of an agricultural plant, in an amount effective to be detectable in the mature agricultural plant. In some embodiments, the endophyte is heterologously disposed in an amount effective to be detectable in an amount of at least about 100 CFU between 100 and 200 CFU, at least about 200 CFU, between 200 and 300 CFU, at least about 300 CFU, between 300 and 400 CFU, at least about 500 CFU, between 500 and 1,000 CFU, at least about 1,000 CFU, between 1,000 and 3,000 CFU, at least about 3,000 CFU, between 3,000 and 10,000 CFU, at least about 10,000 CFU, between 10,000 and 30,000 CFU, at least about 30,000 CFU, between 30,000 and 100,000 CFU, at least about 100,000 CFU or more in the mature agricultural plant.

[00295] In some cases, the endophyte is capable of colonizing particular plant elements or tissue types of the plant. In an embodiment, the endophyte is heterologously disposed on the plant element or seedling in an amount effective to be detectable within a target tissue of the mature agricultural plant selected from a fruit, a seed, a leaf, or a root, or portion thereof. For example, the endophyte can be detected in an amount of at least about 100 CFU, at least

about 200 CFU, at least about 300 CFU, at least about 500 CFU, at least about 1,000 CFU, at least about 3,000 CFU, at least about 10,000 CFU, at least about 30,000 CFU, at least about 100,000 CFU or more, in the target tissue of the mature agricultural plant.

Endophytes compatible with agrichemicals

[00296] In certain embodiments, the endophyte is selected on the basis of its compatibility with commonly used agrichemicals. As described herein, plants, particularly agricultural plants, can be treated with a vast array of agrichemicals, including fungicides, biocides (anti-complex agents), herbicides, insecticides, nematicides, rodenticides, bactericides, virucides, fertilizers, and other agents.

[00297] In some embodiments, the endophytes display tolerance to an agrichemical selected from the group consisting of: Aeris®, Avicta® DuoCot 202, Cruiser®, Syntenta CCB ® (A), Clariva®, Albaugh, Dynasty®, Apron®, Maxim®, Gaucho®, Provoke® ST, Syngenta CCB®, Trilex®, WG Purple, WG Silver, Azoxystrobin, Carboxin, Difenconazole, Fludioxonil, fluxapyroxad, Ipconazole, Mefenoxam, Metalaxyl, Myclobutanil, Penflufen, pyraclostrobin, Sedaxane, TCMTB, Tebuconazole, Thiram, Triadimenol (Baytan®), Trifloxystrobin, Triticonazole, Tolclofos-methyl, PCNB, Abamectin, Chlorpyrifos, Clothianidin, Imidacloprid, Thiamethoxam, Thiodicarb.

[00298] In some cases, it can be important for the endophyte to be compatible with agrichemicals, particularly those with anticomplex properties, in order to persist in the plant although, as described herein, there are many such anticomplex agents that do not penetrate the plant, at least at a concentration sufficient to interfere with the endophyte. Therefore, where a systemic anticomplex agent is used in the plant, compatibility of the endophyte to be inoculated with such agents will be an important criterion.

[00299] In an embodiment, natural isolates of endophytes that are compatible with agrichemicals can be used to inoculate the plants according to the methods described herein. For example, endophytes that are compatible with agriculturally employed anticomplex agents can be isolated by plating a culture of endophytes on a petri dish comprising an effective concentration of the anticomplex agent, and isolating colonies of endophytes that are compatible with the anticomplex agent. In another embodiment, an endophyte that is compatible with an anticomplex agent is used for the methods described herein.

[00300] Bactericide-compatible endophyte can also be isolated by selection on liquid medium. The culture of endophytes can be plated on petri dishes without any forms of mutagenesis; alternatively, endophytes can be mutagenized using any means known in the art. For example, endophyte cultures can be exposed to UV light, gamma-irradiation, or

chemical mutagens such as ethylmethanesulfonate (EMS), ethidium bromide (EtBr) dichlorvos (DDVP, methyl methane sulphonale (MMS), triethylphosphate (TEP), trimethylphosphate (TMP), nitrous acid, or DNA base analogs, prior to selection on fungicide comprising media. Finally, where the mechanism of action of a particular bactericide is known, the target gene can be specifically mutated (either by gene deletion, gene replacement, site-directed mutagenesis, etc.) to generate an endophyte that is resilient against that particular chemical. It is noted that the above-described methods can be used to isolate endophytes that are compatible with both bacteriostatic and bactericidal compounds.

[00301] It will also be appreciated by one skilled in the art that a plant may be exposed to multiple types of anticomplex compounds, either simultaneously or in succession, for example at different stages of plant growth. Where the target plant is likely to be exposed to multiple anticomplex agents, an endophyte that is compatible with many or all of these agrichemicals can be used to inoculate the plant. An endophyte that is compatible with several agents can be isolated, for example, by serial selection. An endophyte that is compatible with the first agent can be isolated as described above (with or without prior mutagenesis). A culture of the resulting endophyte can then be selected for the ability to grow on liquid or solid media comprising the second agent (again, with or without prior mutagenesis). Colonies isolated from the second selection are then tested to confirm its compatibility to both agents.

[00302] Likewise, endophytes that are compatible to biocides (including herbicides such as glyphosate or anticomplex compounds, whether bacteriostatic or bactericidal) that are agriculturally employed can be isolated using methods similar to those described for isolating compatible endophytes. In some embodiments, mutagenesis of the endophyte population can be performed prior to selection with an anticomplex agent. In another embodiment, selection is performed on the endophyte population without prior mutagenesis. In still another embodiment, serial selection is performed on an endophyte: the endophyte is first selected for compatibility to a first anticomplex agent. The isolated compatible endophyte is then cultured and selected for compatibility to the second anticomplex agent. Any colony thus isolated is tested for compatibility to each, or both anticomplex agents to confirm compatibility with these two agents.

[00303] Compatibility with an antimicrobial agent can be determined by a number of means known in the art, including the comparison of the minimal inhibitory concentration of the unmodified and modified endophytes. In some embodiments, the present invention discloses an isolated modified endophyte, wherein the endophyte is modified such that it exhibits at

least 3 fold greater, for example, at least 5 fold greater, between 5 and 10 fold greater, at least 10 fold greater, between 10 and 20 fold greater, at least 20 fold greater, between 20 and 30 fold greater, at least 30 fold greater or more minimal inhibitory concentration to an antimicrobial agent when compared with the unmodified endophyte.

[00304] In a particular embodiment, disclosed herein are endophytes with enhanced compatibility to the herbicide glyphosate. In some embodiments, the endophyte has a doubling time in growth medium comprising at least 1 mM glyphosate, for example, between 1 mM and 2mM glyphosate, at least 2 mM glyphosate, between 2 mM and 5 mM glyphosate, at least 5mM glyphosate, between 5 mM and 10 mM glyphosate, at least 10mM glyphosate, between 10 mM and 15 mM glyphosate, at least 15mM glyphosate or more, that is no more than 250%, between 250% and 100%, for example, no more than 200%, between 200% and 175%, no more than 175%, between 175% and 150%, no more than 150%, between 150% and 125%, or no more than 125%, of the doubling time of the endophyte in the same growth medium comprising no glyphosate. In one particular embodiment, the endophyte has a doubling time in growth medium comprising 5mM glyphosate that is no more than 150% the doubling time of the endophyte in the same growth medium comprising no glyphosate.

[00305] In another embodiment, the endophyte has a doubling time in a plant tissue comprising at least 10 ppm glyphosate,, between 10 and 15 ppm, for example, at least 15 ppm glyphosate, between 15 and 10 ppm, at least 20 ppm glyphosate, between 20 and 30 ppm, at least 30 ppm glyphosate, between 30 and 40 ppm, at least 40 ppm glyphosate or more, that is no more than 250%, between 250% and 200%, for example, no more than 200%, between 200% and 175%, no more than 175%, between 175% and 150%, no more than 150%, between 150% and 125%, or no more than 125%, of the doubling time of the endophyte in a reference plant tissue comprising no glyphosate. In one particular embodiment, the endophyte has a doubling time in a plant tissue comprising 40 ppm glyphosate that is no more than 150% the doubling time of the endophyte in a reference plant tissue comprising no glyphosate.

[00306] The selection process described above can be repeated to identify isolates of endophytes that are compatible with a multitude of agents.

[00307] Candidate isolates can be tested to ensure that the selection for agrichemical compatibility did not result in loss of a desired bioactivity. Isolates of endophytes that are compatible with commonly employed agents can be selected as described above. The resulting compatible endophyte can be compared with the parental endophyte on plants in its ability to promote germination.

[00308] The agrichemical compatible endophytes generated as described above can be detected in samples. For example, where a transgene was introduced to render the endophyte compatible with the agrichemical(s), the transgene can be used as a target gene for amplification and detection by PCR. In addition, where point mutations or deletions to a portion of a specific gene or a number of genes results in compatibility with the agrichemical(s), the unique point mutations can likewise be detected by PCR or other means known in the art. Such methods allow the detection of the endophyte even if it is no longer viable. Thus, commodity plant products produced using the agrichemical compatible endophytes described herein can readily be identified by employing these and related methods of nucleic acid detection.

Beneficial Attributes of Synthetic Compositions of Plant Elements and Endophytes

[00309] The present invention contemplates the establishment of a relationship between a symbiont and a plant element. In some embodiments, endophyte association results in a detectable change to the plant element, or the whole plant. The detectable change can be an improvement in a number of agronomic traits (e.g., improved general health, increased response to biotic or abiotic stresses, or enhanced properties of the plant or a plant element, including fruits and grains). Alternatively, the detectable change can be a physiological or biological change that can be measured by methods known in the art. The detectable changes are described in more detail in the sections below. As used herein, an endophyte is considered to have conferred an improved agricultural trait whether or not the improved trait arose from the plant, the endophyte, or the concerted action between the plant and endophyte. Therefore, for example, whether a beneficial hormone or chemical is produced by the plant or the endophyte, for purposes, the endophyte will be considered to have conferred an improved agronomic trait upon the host plant, as compared to an isoline plant that has not been heterologously disposed to said endophyte.

[00310] In some embodiments, provided herein, are methods for producing a plant element of a plant with a heritably altered trait. The trait of the plant can be altered without known genetic modification of the plant genome, and comprises the following steps. First, a preparation of an isolated endophyte that is heterologously disposed to the plant element of the plant is provided, and optionally processed to produce an endophyte formulation. The endophyte formulation is then contacted with the plant. The plants are then allowed to go to seed, and the seeds are collected.

Improved general health

[00311] Also described herein are plants, and fields of plants, that are heterologously disposed to beneficial endophytes, such that the overall fitness, productivity or health of the plant or a portion thereof, is maintained, increased and/or improved over a period of time. Improvement in overall plant health can be assessed using numerous physiological parameters including, but not limited to, height, overall biomass, root and/or shoot biomass, emergence, seed germination, seedling survival, photosynthetic efficiency, transpiration rate, seed/fruit number or mass, plant grain yield, leaf chlorophyll content, photosynthetic rate, root length, or any combination thereof.

Drought tolerance

[00312] In some cases, a plant resulting from seeds or other plant elements treated with an endophyte can exhibit a physiological change, such as a compensation of the stress-induced reduction in photosynthetic activity. Fv/Fm tests whether or not plant stress affects photosystem II in a dark adapted state. Fv/Fm is one of the most commonly used chlorophyll fluorescence measuring parameter. The Fv/Fm test is designed to allow the maximum amount of the light energy to take the fluorescence pathway. It compares the dark-adapted leaf pre-photosynthetic fluorescent state, called minimum fluorescence, or Fo, to maximum fluorescence called Fm. In maximum fluorescence, the maximum number of reaction centers have been reduced or closed by a saturating light source. In general, the greater the plant stress, the fewer open reaction centers available, and the Fv/Fm ratio is lowered. Fv/Fm is a measuring protocol that works for many types of plant stress. For example, there would be a difference in the Fv/Fm after exposure of an endophyte treated plant that had been subjected to heat shock or drought conditions, as compared to a corresponding control, a genetically identical plant that does not contain the endophytes grown in the same conditions. In some cases, the inoculated plant as disclosed herein can exhibit an increased change in photosynthetic activity $\Delta Fv(\Delta Fv/Fm)$ after heat-shock or drought stress treatment, for example 1, 2, 3, 4, 5, 6, 7 days or more after the heat-shock or drought stress treatment, or until photosynthesis ceases, as compared with corresponding control plant of similar developmental stage but not comprising endophytes. For example, a plant having an endophyte able to confer heat and/or drought-tolerance can exhibit a $\Delta Fv/Fm$ of from about 0.1 to about 0.8 after exposure to heat-shock or drought stress or a $\Delta Fv/Fm$ range of from about 0.03 to about 0.8 under one day, or 1, 2, 3, 4, 5, 6, 7, or over 7 days post heat-shock or drought stress treatment, or until photosynthesis ceases. In some embodiments, stress-induced reductions in photosynthetic activity can be compensated by at least about 0.25% (for example, at least about 0.5%, between 0.5% and 1%, at least about 1%, between 1% and 2%,

at least about 2%, between 2% and 3%, at least about 3%, between 3% and 5%, at least about 5%, between 5% and 10%, at least about 8%, at least about 10%, between 10% and 15%, at least about 15%, between 15% and 20%, at least about 20%, between 20% and 25%, at least about 25%, between 25% and 30%, at least about 30%, between 30% and 40%, at least about 40%, between 40% and 50%, at least about 50%, between 50% and 60%, at least about 60%, between 60% and 75%, at least about 75%, between 75% and 80%, at least about 80%, between 80% and 85%, at least about 85%, between 85% and 90%, at least about 90%, between 90% and 95%, at least about 95%, between 95% and 99%, at least about 99% or at least 100%) as compared to the photosynthetic activity decrease in a corresponding reference agricultural plant following heat shock conditions. Significance of the difference between inoculated and reference agricultural plants can be established upon demonstrating statistical significance, for example at $p < 0.05$ with an appropriate parametric or non-parametric statistic, e.g., Chi-square test, Student's t-test, Mann-Whitney test, or F-test based on the assumption or known facts that the inoculated plant and reference agricultural plant have identical or near identical genomes (isoline comparison).

[00313] In some embodiments, the plants comprise endophytes able to increase heat and/or drought-tolerance in sufficient quantity, such that increased growth or improved recovery from wilting under conditions of heat or drought stress is observed. For example, an endophyte population described herein can be present in sufficient quantity in a plant, resulting in increased growth as compared to a plant that does not contain endophytes, when grown under drought conditions or heat shock conditions, or following such conditions. Increased heat and/or drought tolerance can be assessed with physiological parameters including, but not limited to, increased height, overall biomass, root and/or shoot biomass, seed germination, seedling survival, photosynthetic efficiency, transpiration rate, seed/fruit number or mass, plant grain or fruit yield, leaf chlorophyll content, photosynthetic rate, root length, wilt recovery, turgor pressure, or any combination thereof, as compared to a reference agricultural plant grown under similar conditions. For example, the endophyte may provide an improved benefit or tolerance to a plant that is of at least 3%, between 3% and 5%, at least 5%, between 5% and 10%, at least 10%, between 10% and 15%, for example at least 15%, between 15% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 75%, at least 75%, between 75% and 100%, at least 100%, between 100% and 150%, at least 150%, between 150% and 200%, at least 200%, between 200% and

300%, at least 300% or more, when compared with uninoculated plants grown under the same conditions.

[00314] In various embodiments, endophytes heterologously disposed to the plant can confer various benefits to the plant, including but not limited to: thermal tolerance, herbicide tolerance, drought resistance, insect resistance, fungus resistance, virus resistance, bacteria resistance, male sterility, cold tolerance, salt tolerance, increased yield, enhanced nutrient use efficiency, increased nitrogen use efficiency, increased protein content, increased fermentable carbohydrate content, reduced lignin content, increased antioxidant content, enhanced water use efficiency, increased vigor, increased germination efficiency, earlier or increased flowering, increased biomass, altered root-to-shoot biomass ratio, enhanced soil water retention, or a combination thereof. A difference between the inoculated plant (e.g., a plant to which one or more endophytes have been heterologously disposed) and a reference agricultural plant can also be measured using other methods known in the art.

Formulations for Agricultural Use

[00315] The endophyte populations described herein are intended to be useful in the improvement of agricultural plants, and as such, may be formulated with other compositions as part of an agriculturally compatible carrier. It is contemplated that such carriers can include, but not be limited to: seed treatment, root wash, seedling soak, foliar application, soil inocula, in-furrow application, sidedress application, soil pre-treatment, wound inoculation, drip tape irrigation, vector-mediation via a pollinator, injection, osmopriming, hydroponics, aquaponics, aeroponics. The carrier composition with the endophyte populations, may be prepared for agricultural application as a liquid, a solid, or a gas formulation. Application to the plant may be achieved, for example, as a powder for surface deposition onto plant leaves, as a spray to the whole plant or selected plant element, as part of a drip to the soil or the roots, or as a coating onto the plant element prior to planting. Such examples are meant to be illustrative and not limiting to the scope of the invention.

[00316] The formulation useful for these embodiments generally and typically include at least one member selected from the group consisting of a buffer, a tackifier, a microbial stabilizer, a fungicide, an anticomplex agent, an herbicide, a nematicide, an insecticide, a bactericide, a virucide, a plant growth regulator, a rodenticide, a desiccant, and a nutrient.

[00317] The carrier can be a solid carrier or liquid carrier, and in various forms including microspheres, powders, emulsions and the like. The carrier may be any one or more of a number of carriers that confer a variety of properties, such as increased stability, wettability, or dispersability. Wetting agents such as natural or synthetic surfactants, which can be

nonionic or ionic surfactants, or a combination thereof can be included in a composition of the invention. Water-in-oil emulsions can also be used to formulate a composition that includes the purified population (see, for example, U.S. Patent No. 7,485,451). Suitable formulations that may be prepared include wettable powders, granules, gels, agar strips or pellets, thickeners, biopolymers, and the like, microencapsulated particles, and the like, liquids such as aqueous flowables, aqueous suspensions, water-in-oil emulsions, etc. The formulation may include grain or legume products, for example, ground grain or beans, broth or flour derived from grain or beans, starch, sugar, or oil.

[00318] In some embodiments, the agricultural carrier may be soil or a plant growth medium. Other agricultural carriers that may be used include water, fertilizers, plant-based oils, humectants, or combinations thereof. Alternatively, the agricultural carrier may be a solid, such as diatomaceous earth, loam, silica, alginate, clay, bentonite, vermiculite, seed cases, other plant and animal products, or combinations, including granules, pellets, or suspensions. Mixtures of any of the aforementioned ingredients are also contemplated as carriers, such as but not limited to, pesta (flour and kaolin clay), agar or flour-based pellets in loam, sand, or clay, etc. Formulations may include food sources for the cultured organisms, such as barley, rice, wheat or other biological materials such as seed, plant elements, sugar cane bagasse, hulls or stalks from grain processing, ground plant material or wood from building site refuse, sawdust or small fibers from recycling of paper, fabric, or wood. Other suitable formulations will be known to those skilled in the art.

[00319] In an embodiment, the formulation can include a tackifier or adherent. Such agents are useful for combining the complex population of the invention with carriers that can contain other compounds (e.g., control agents that are not biologic), to yield a coating composition. Such compositions help create coatings around the plant or plant element to maintain contact between the endophyte and other agents with the plant or plant element. In some embodiments, adherents are selected from the group consisting of: alginate, gums, starches, lecithins, formononetin, polyvinyl alcohol, alkali formononetinate, hesperetin, polyvinyl acetate, cephalins, Gum Arabic, Xanthan Gum, carragennan, PGA, other biopolymers, Mineral Oil, Polyethylene Glycol (PEG), Polyvinyl pyrrolidone (PVP), Arabino-galactan, Methyl Cellulose, PEG 400, Chitosan, Polyacrylamide, Polyacrylate, Polyacrylonitrile, Glycerol, Triethylene glycol, Vinyl Acetate, Gellan Gum, Polystyrene, Polyvinyl, Carboxymethyl cellulose, Gum Ghatti, and polyoxyethylene-polyoxybutylene block copolymers. Other examples of adherent compositions that can be used in the synthetic

preparation include those described in EP 0818135, CA 1229497, WO 2013090628, EP 0192342, WO 2008103422 and CA 1041788.

[00320] It is also contemplated that the formulation may further comprise an anti-caking agent.

[00321] The formulation can also contain a surfactant, wetting agent, emulsifier, stabilizer, or anti-foaming agent. Non-limiting examples of surfactants include nitrogen-surfactant blends such as Prefer 28 (Cenex), Surf-N(US), Inhance (Brandt), P-28 (Wilfarm) and Patrol (Helena); esterified seed oils include Sun-It II (AmCy), MSO (UAP), Scoil (Agasco), Hasten (Wilfarm) and Mes-100 (Drexel); and organo-silicone surfactants include Silwet L77 (UAP), Silikin (Terra), Dyne-Amic (Helena), Kinetic (Helena), Sylgard 309 (Wilbur-Ellis) and Century (Precision), polysorbate 20, polysorbate 80, Tween 20, Tween 80, Scattics, Alktest TW20, Canarcel, Peogabsorb 80, Triton X-100, Conco NI, Dowfax 9N, Igebapl CO, Makon, Neutronyx 600, Nonipol NO, Plytergent B, Renex 600, Solar NO, Sterox, Serfonic N, T-DET-N, Tergitol NP, Triton N, IGEPAL CA-630, Nonident P-40, Pluronic. In some embodiments, the surfactant is present at a concentration of between 0.01% v/v to 10% v/v. In another embodiment, the surfactant is present at a concentration of between 0.1% v/v to 1% v/v. An example of an anti-foaming agent would be Antifoam-C.

[00322] In certain cases, the formulation includes a microbial stabilizer. Such an agent can include a desiccant. As used herein, a “desiccant” can include any compound or mixture of compounds that can be classified as a desiccant regardless of whether the compound or compounds are used in such concentrations that they in fact have a desiccating effect on the liquid inoculant. Such desiccants are ideally compatible with the population used, and should promote the ability of the endophyte population to survive application on the seeds and to survive desiccation. Examples of suitable desiccants include one or more of trehalose, sucrose, glycerol, and methylene glycol. Other suitable desiccants include, but are not limited to, non-reducing sugars and sugar alcohols (e.g., mannitol or sorbitol). The amount of desiccant introduced into the formulation can range from about 5% to about 50% by weight/volume, for example, between about 10% to about 40%, between about 15% and about 35%, or between about 20% and about 30%. In some embodiments, components of a sugar-based microbial stabilizer include, but are not limited to, glucose, sucrose, polyvinylpyrrolidone K 30 (PVP30K), mineral oil, soy lecithin, peptone, monopotassium phosphate (KH₂PO₄) and dipotassium phosphate (K₂HPO₄). In an alternate embodiment, components of a non-sugar based microbial stabilizer include, but are not limited to, polyvinylpyrrolidone K 30 (PVP30K), polyvinylpyrrolidone/vinyl acetate (PVP-VA), soy

lecithin, peptone, mineral oil, hydroxypropyl-guar (HP-Guar), monopotassium phosphate (KH_2PO_4) and dipotassium phosphate (K_2HPO_4). Components of exemplary microbial stabilizers for use with the invention described herein are depicted in Table 1 and Table 2.

Table 1. Exemplary Sugar Based Microbial Stabilizer

Component	Percentage (%), by weight
Glucose	11.4
Sucrose	11.4
PVP30K	2.8
Mineral oil	5.7
Soy lecithin	0.3
Peptone	11.4
KH_2PO_4	0.78
K_2HPO_4	0.99
Non-chlorinated water	55

Table 2: Exemplary Non-sugar Based Microbial Stabilizer

Component	Percentage (%), by weight
PVP30K	3.8
PVP-VA	3.8
Soy lecithin	0.4
Peptone	15.4
Mineral oil	6.0
HP-Guar	0.2
KH_2PO_4	0.96
K_2HPO_4	1.23
Non-chlorinated water	68

[00323] In some cases, it is advantageous for the formulation to contain agents such as a fungicide, an anticomplex agent, an herbicide, a nematocide, an insecticide, a plant growth regulator, a rodenticide, a bactericide, a virucide, or a nutrient. Such agents are ideally compatible with the agricultural plant element or seedling onto which the formulation is applied (e.g., it should not be deleterious to the growth or health of the plant). Furthermore, the agent is ideally one which does not cause safety concerns for human, animal or industrial use (e.g., no safety issues, or the compound is sufficiently labile that the commodity plant product derived from the plant contains negligible amounts of the compound).

[00324] In the liquid form, for example, solutions or suspensions, endophyte populations can be mixed or suspended in water or in aqueous solutions. Suitable liquid diluents or carriers include water, aqueous solutions, petroleum distillates, or other liquid carriers.

[00325] Solid compositions can be prepared by dispersing the endophyte populations of the invention in and on an appropriately divided solid carrier, such as peat, wheat, bran, vermiculite, clay, talc, bentonite, diatomaceous earth, fuller's earth, pasteurized soil, and the like. When such formulations are used as wettable powders, biologically compatible dispersing agents such as non-ionic, anionic, amphoteric, or cationic dispersing and emulsifying agents can be used.

[00326] In some cases, a flowability polymer, also referred to as a plantability polymer such as Flo Rite[®] e.g., Flo-Rite[®] 1706 (BASF, Ludwigshafen, Germany). In some embodiments, a flowability or plantability polymer is DISCO[™] AG (Incotec, Enkhuizen, the Netherlands). In some embodiments, a flowability or plantability polymer is Kannar[®] Universal Wonder (Kannar Earth Science, Ltd., Buford, GA).

[00327] The solid carriers used upon formulation include, for example, mineral carriers such as kaolin clay, pyrophyllite, bentonite, montmorillonite, diatomaceous earth, acid white soil, vermiculite, and perlite, and inorganic salts such as ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride, and calcium carbonate. Also, organic fine powders such as wheat flour, wheat bran, and rice bran may be used. The liquid carriers include vegetable oils such as soybean oil, neem oil, cottonseed oil, and other compositions such as glycerol, ethylene glycol, polyethylene glycol, propylene glycol, polypropylene glycol, etc.

[00328] In an embodiment, the formulation is ideally suited for coating of a population of endophytes onto plant elements. The endophytes populations described in the present invention are capable of conferring many fitness benefits to the host plants. The ability to confer such benefits by coating the populations on the surface of plant elements has many potential advantages, particularly when used in a commercial (agricultural) scale.

[00329] The endophyte populations herein can be combined with one or more of the agents described above to yield a formulation suitable for combining with an agricultural plant element, seedling, or other plant element. Endophyte populations can be obtained from growth in culture, for example, using a synthetic growth medium. In addition, endophytes can be cultured on solid media, for example on petri dishes, scraped off and suspended into the preparation. Endophytes at different growth phases can be used. For example, endophytes at lag phase, early-log phase, mid-log phase, late-log phase, stationary phase, early death phase,

or death phase can be used. Endophytic spores may be used for the present invention, for example but not limited to: arthospores, sporangiospores, conidia, chlamadospores, pycnidiospores, endospores, zoospores.

[00330] The formulations comprising endophyte populations typically contains between about 0.1 to 95% by weight, for example, between about 1% and 90%, between about 3% and 75%, between about 5% and 60%, between about 10% and 50% in wet weight of the population. It is preferred that the formulation contains at least about 10^3 CFU per ml of formulation, for example, at least about 10^4 , at least about 10^5 , at least about 10^6 , at least about 10^7 CFU, at least about 10^8 CFU per ml of formulation. It is preferred that the formulation be applied to the plant element at about 10^2 CFU/seed, between 10^2 and 10^3 CFU, at least about 10^3 CFU, between 10^3 and 10^4 CFU, at least about 10^4 CFU, between 10^4 and 10^5 CFU, at least about 10^5 CFU, between 10^5 and 10^6 CFU, at least about 10^6 CFU, between 10^6 and 10^7 CFU, at least about 10^7 CFU, between 10^7 and 10^8 CFU, or even greater than 10^8 CFU per seed.

Populations of Plant Elements

[00331] In another embodiment, the invention provides for a substantially uniform population of synthetic compositions comprising plant elements (PEs), comprising two or more PEs comprising the endophytic population, as described herein above. Substantial uniformity can be determined in many ways. In some cases, at least 10%, between 10% and 20%, for example, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 70%, at least 70%, between 70% and 75%, at least 75%, between 75% and 80%, at least 80%, between 80% and 90%, at least 90%, between 90% and 95%, at least 95% or more of the PEs in the population, comprises the endophytic population in an amount effective to colonize a plant, or plants, derived from said PEs when disposed on the surface of the PEs. In other cases, at least 10%, between 10% and 20%, for example, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 70%, at least 70%, between 70% and 75%, at least 75%, between 75% and 80%, at least 80%, between 80% and 90%, at least 90%, between 90% and 95%, at least 95% or more of the plant elements in the population, contains at least 1, between 10 and 10, 10, between 10 and 100, or 100 CFU on the plant element surface or per gram of plant element, for example, between 100 and 200 CFU, at least 200 CFU, between 200 and 300 CFU, at least 300 CFU, between 300 and 1,000 CFU, at least 1,000 CFU, between 1,000 and 3,000 CFU, at least 3,000 CFU,

between 3,000 and 10,000 CFU, at least 10,000 CFU, between 10,000 and 30,000 CFU, at least 30,000 CFU, between 30,000 and 100,000 CFU, at least 100,000 CFU, between 100,000 and 300,000 CFU, at least 300,000 CFU, between 300,000 and 1,000,000 CFU, or at least 1,000,000 CFU per plant element or more.

[00332] In a particular embodiment, the population of plant elements is packaged in a bag or container suitable for commercial sale. Such a bag contains a unit weight or count of the plant elements comprising the endophytic population as described herein, and further comprises a label. In an embodiment, the bag or container contains at least 100 plant elements, between 100 and 1,000 plant elements, 1,000 plant elements, between 1,000 and 5,000 plant elements, for example, at least 5,000 plant elements, between 5,000 and 10,000 plant elements, at least 10,000 plant elements, between 10,000 and 20,000 plant elements, at least 20,000 plant elements, between 20,000 and 30,000 plant elements, at least 30,000 plant elements, between 30,000 and 50,000 plant elements, at least 50,000 plant elements, between 50,000 and 70,000 plant elements, at least 70,000 plant elements, between 70,000 and 80,000 plant elements, at least 80,000 plant elements, between 80,000 and 90,000, at least 90,000 plant elements or more. In another embodiment, the bag or container can comprise a discrete weight of plant elements, for example, at least 1 lb, between 1 and 2 lbs, at least 2 lbs, between 2 and 5 lbs, at least 5 lbs, between 5 and 10 lbs, at least 10 lbs, between 10 and 30 lbs, at least 30 lbs, between 30 and 50 lbs, at least 50 lbs, between 50 and 70 lbs, at least 70 lbs or more. The bag or container comprises a label describing the plant elements and/or said endophytic population. The label can contain additional information, for example, the information selected from the group consisting of: net weight, lot number, geographic origin of the plant elements, test date, germination rate, inert matter content, and the amount of noxious weeds, if any. Suitable containers or packages include those traditionally used in plant seed commercialization. The invention also contemplates other containers with more sophisticated storage capabilities (e.g., with microbiologically tight wrappings or with gas-or water-proof containments).

[00333] In some cases, a sub-population of seeds comprising the endophytic population is further selected on the basis of increased uniformity, for example, on the basis of uniformity of microbial population. For example, individual plant elements of pools collected from individual cobs, individual plants, individual plots (representing plants inoculated on the same day) or individual fields can be tested for uniformity of microbial density, and only those pools meeting specifications (e.g., at least 80% of tested plant elements have minimum

density, as determined by quantitative methods described elsewhere) are combined to provide the agricultural seed sub-population.

[00334] The methods described herein can also comprise a validating step. The validating step can entail, for example, growing some plant elements collected from the inoculated plants into mature agricultural plants, and testing those individual plants for uniformity. Such validating step can be performed on individual seeds collected from cobs, individual plants, individual plots (representing plants inoculated on the same day) or individual fields, and tested as described above to identify pools meeting the required specifications.

[00335] In some embodiments, methods described herein include planting a synthetic composition described herein. Suitable planters include an air seeder and/or fertilizer apparatus used in agricultural operations to apply particulate materials including one or more of the following, seed, fertilizer and/or inoculants, into soil during the planting operation. Seeder/fertilizer devices can include a tool bar having ground-engaging openers thereon, behind which is towed a wheeled cart that includes one or more containment tanks or bins and associated metering means to respectively contain and meter therefrom particulate materials.

[00336] In certain embodiments, a composition described herein may be in the form of a liquid, a slurry, a solid, or a powder (wetttable powder or dry powder). In another embodiment, a composition may be in the form of a seed coating. Compositions in liquid, slurry, or powder (e.g., wetttable powder) form may be suitable for coating plant elements. When used to coat plant elements, the composition may be applied to the plant elements and allowed to dry. In embodiments wherein the composition is a powder (e.g., a wetttable powder), a liquid, such as water, may need to be added to the powder before application to a seed.

[00337] In still another embodiment, the methods can include introducing into the soil an inoculum of one or more of the endophyte populations described herein. Such methods can include introducing into the soil one or more of the compositions described herein. The inoculum(s) or compositions may be introduced into the soil according to methods known to those skilled in the art. Non-limiting examples include in-furrow introduction, spraying, coating seeds, foliar introduction, etc. In a particular embodiment, the introducing step comprises in-furrow introduction of the inoculum or compositions described herein.

[00338] In an embodiment, plant elements may be treated with composition(s) described herein in several ways but preferably via spraying or dripping. Spray and drip treatment may be conducted by formulating compositions described herein and spraying or dripping the

composition(s) onto a seed(s) via a continuous treating system (which is calibrated to apply treatment at a predefined rate in proportion to the continuous flow of seed), such as a drum-type of treater. Batch systems, in which a predetermined batch size of seed and composition(s) as described herein are delivered into a mixer, may also be employed.

[00339] In another embodiment, the treatment entails coating plant elements. One such process involves coating the inside wall of a round container with the composition(s) described herein, adding plant elements, then rotating the container to cause the plant elements to contact the wall and the composition(s), a process known in the art as “container coating.” Plant elements can be coated by combinations of coating methods. Soaking typically entails using liquid forms of the compositions described. For example, plant elements can be soaked for about 1 minute to about 24 hours (e.g., for at least 1 min, between 1 and 5 min, 5 min, between 5 and 10 min, 10 min, between 10 and 20 min, 20 min, between 20 and 40 min, 40 min, between 40 and 80 min, 80 min, between 80 min and 3 hrs, 3 hrs, between 3 hrs and 6 hrs, 6 hr, between 6 hrs and 12 hrs, 12 hr, between 12 hrs and 24 hrs, 24 hrs).

Population of Plants/Agricultural Fields

[00340] A major focus of crop improvement efforts has been to select varieties with traits that give, in addition to the highest return, the greatest homogeneity and uniformity. While inbreeding can yield plants with substantial genetic identity, heterogeneity with respect to plant height, flowering time, and time to seed, remain impediments to obtaining a homogeneous field of plants. The inevitable plant-to-plant variability is caused by a multitude of factors, including uneven environmental conditions and management practices. Another possible source of variability can, in some cases, be due to the heterogeneity of the endophyte population inhabiting the plants. By providing endophyte populations onto plant reproductive elements, the resulting plants generated by germinating the plant reproductive elements have a more consistent endophyte collection, and thus are expected to yield a more uniform population of plants.

[00341] Therefore, in another embodiment, the invention provides a substantially uniform population of plants. The population can include at least 5 plants, between 5 and 10 plants at least 10 plants, between 10 and 100 plants, for example, at least 100 plants, between 100 and 300 plants, at least 300 plants, between 300 and 1,000 plants, at least 1,000 plants, between 1,000 and 3,000 plants, at least 3,000 plants, between 3,000 and 10,000 plants, at least 10,000 plants, between 10,000 and 30,000 plants, at least 30,000 plants, between 30,000 and 100,000 plants, at least 100,000 plants or more. The plants may be derived from plant reproductive

elements comprising endophyte populations as described herein. The plants are cultivated in substantially uniform groups, for example in rows, groves, blocks, circles, or other planting layout.

[00342] The uniformity of the plants can be measured in a number of different ways. In some embodiments, there is an increased uniformity with respect to endophytes within the plant population. For example, In some embodiments, a substantial portion of the population of plants, for example at least 10%, between 10% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 70%, at least 70%, between 70% and 75%, at least 75%, between 75% and 80%, at least 80%, between 80% and 90%, at least 90%, between 90% and 95%, at least 95% or more of the plant elements or plants in a population, contains a threshold number of an endophyte population. The threshold number can be at least 10 CFU, between 10 and 100 CFU, at least 100 CFU, between 100 and 300 CFU, for example at least 300 CFU, between 300 and 1,000 CFU, at least 1,000 CFU, between 1,000 and 3,000 CFU, at least 3,000 CFU, between 3,000 and 10,000 CFU, at least 10,000 CFU, between 10,000 and 30,000 CFU, at least 30,000 CFU, between 30,000 and 100,000 CFU, at least 100,000 CFU or more, in the plant or a part of the plant. Alternatively, in a substantial portion of the population of plants, for example, in at least 1%, between 1% and 10%, at least 10%, between 10% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 70%, at least 70%, between 70% and 75%, at least 75%, between 75% and 80%, at least 80%, between 80% and 90%, at least 90%, between 90% and 95%, at least 95% or more of the plants in the population, the endophyte population that is provided to the seed or seedling represents at least 0.1%, between 0.1% and 1% at least 1%, between 1% and 5%, at least 5%, between 5% and 10%, at least 10%, between 10% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60%, between 60% and 70%, at least 70%, between 70% and 80%, at least 80%, between 80% and 90%, at least 90%, between 90% and 95%, at least 95%, between 95% and 99%, at least 99%, between 99% and 100%, or 100% of the total endophyte population in the plant/seed.

[00343] In an embodiment, there is increased genetic uniformity of a substantial proportion or all detectable endophytes within the taxa, genus, or species of a component relative to an uninoculated control. This increased uniformity can be a result of the endophyte being of

monoclonal origin or otherwise deriving from a population comprising a more uniform genome sequence and plasmid repertoire than would be present in the endophyte population a plant that derives its endophyte community largely via assimilation of diverse soil symbionts.

[00344] In another embodiment, there is an increased uniformity with respect to a physiological parameter of the plants within the population. In some cases, there can be an increased uniformity in the height of the plants when compared with a population of reference agricultural plants grown under the same conditions. For example, there can be a reduction in the standard deviation in the height of the plants in the population of at least 5%, between 5% and 10%, for example, at least 10%, between 10% and 15%, at least 15%, between 15% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60% or more, when compared with a population of reference agricultural plants grown under the same conditions. In other cases, there can be a reduction in the standard deviation in the flowering time of the plants in the population of at least 5%, between 5% and 10%, for example, at least 10%, between 10% and 15%, at least 15%, between 15% and 20%, at least 20%, between 20% and 30%, at least 30%, between 30% and 40%, at least 40%, between 40% and 50%, at least 50%, between 50% and 60%, at least 60% or more, when compared with a population of reference agricultural plants grown under the same conditions.

Commodity Plant Products

[00345] The present invention provides a commodity plant product, as well as methods for producing a commodity plant product, that is derived from a plant. As used herein, a “commodity plant product” refers to any composition or product that is comprised of material derived from a plant, seed, plant cell, or plant element of the present invention. Commodity plant products may be sold to consumers and can be viable or nonviable. Nonviable commodity products include but are not limited to nonviable plant elements and grains; processed seeds, seed parts, and plant elements; dehydrated plant tissue, frozen plant tissue, and processed plant tissue; seeds and plant elements processed for animal feed for terrestrial and/or aquatic animal consumption, oil, meal, flour, flakes, bran, fiber, paper, tea, coffee, silage, crushed or whole grain, and any other food for human or animal consumption such as the fruit or other edible portion of the plant; and biomasses and fuel products; and raw material in industry.

[00346] Industrial uses of oils derived from the agricultural plants described herein include ingredients for paints, plastics, fibers, detergents, cosmetics, lubricants, and biodiesel fuel. Plant oils may be split, inter-esterified, sulfurized, epoxidized, polymerized, ethoxylated, or

cleaved. Designing and producing plant oil derivatives with improved functionality and improved oleochemistry is a rapidly growing field. For example, a mixture of triglycerides is usually split and separated into pure fatty acids, which are then combined with petroleum-derived alcohols or acids, nitrogen, sulfonates, chlorine, or with fatty alcohols derived from fats and oils to produce the desired type of oil or fat. Commodity plant products also include industrial compounds, such as a wide variety of resins used in the formulation of adhesives, films, plastics, paints, coatings and foams.

Methods of Using Endophytes and Synthetic Compositions Comprising Endophytes

[00347] As described herein, purified endophyte populations and synthetic compositions comprising the same (e.g., formulations) can be used to confer beneficial traits to the host plant.

[00348] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims. Each patent application, journal article, citation, and other references are herein incorporated by reference in their entirety, as if each has been incorporated by reference individually.

EXAMPLES**Example 1. Isolation and Identification of Endophytes**

[00349] Isolation and cultivation of endophytic microbes from agricultural plants was performed using methods well known in the art. DNA was extracted from the ground tissues using the DNeasy DNA extraction kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The endophytes were characterized by the sequences of genomic regions, these sequences are listed in Table 4. Primers that amplify genomic regions of the endophytes of the present invention are listed in Table 3.

Table 3. Primer sequences useful in identifying microbes of the present invention

Primers	Genomic locus
27f (5' - AGAGTTTGATYMTGGCTCAG - 3') (SEQ ID NO: 1) 1492r (5' - GGTTACCTTGTTACGACTT - 3') (SEQ ID NO: 2)	16S
515f (5' - GTGYCAGCMGCCGCGGTAA - 3') (SEQ ID NO: 3) 806r (5' - GGACTACNVGGGTWTCTAAT - 3') (SEQ ID NO: 4)	16S
ITS_1 (5' - CTTGGTCATTTAGAGGAAGTAA - 3') (SEQ ID NO: 5) LR5 (5' - TCCTGAGGGAAACTTCG - 3') (SEQ ID NO: 8)	ITS
ITS_2 (5' - GCTGCGTTCTTCATCGATGC - 3') (SEQ ID NO: 6) ITS_3 (5' - GCATCGATGAAGAACGCAGC - 3') (SEQ ID NO: 7)	ITS
MIC-19994, unique genomic region, primer - amplicon F (5' - TGCTGGTAGTGCGAATGAAA - 3') (SEQ ID NO: 9), MIC-19994, unique genomic region, primer - amplicon R (5' - CTTTCGGGTTCCATCAGGT - 3') (SEQ ID NO: 10)	unique genomic region
MIC-31593, unique genomic region, primer - amplicon F (5' - CTACCGCAAGAGCAACTGTG - 3') (SEQ ID NO: 11) MIC-31593, unique genomic region, primer - amplicon R (5' - ACTTCCTCCTCCTCCTCCTC - 3') (SEQ ID NO: 12)	unique genomic region
MIC-96038, unique genomic region, primer - amplicon F (5' - GTCCTCGCCTAATCAGGAGTC - 3') (SEQ ID NO: 13) MIC-96038, unique genomic region, primer - amplicon R (5' - TCCTATTCCCTGACGTGCTAC - 3') (SEQ ID NO: 14)	unique genomic region
MIC-33414, unique genomic region, primer - amplicon F (5' - GAGGAGGAGGAGGAGAGGTT - 3') (SEQ ID NO: 15) MIC-33414, unique genomic region, primer - amplicon R (5' - CGTCCGTCTCCCAGACTATT - 3') (SEQ ID NO: 16)	unique genomic region
MIC-68178, primer - amplicon F (5' - CTCCTCCTCCTCCTCCTGAT - 3') (SEQ ID NO: 36) MIC-68178, primer - amplicon R (5' - TCACAGAGCTACGCGACTTG - 3') (SEQ ID NO: 37)	unique genomic region
MIC-68390 (5' - CTTCCAGGCATAGTAATGTGGA - 3') (SEQ ID NO: 34) MIC-68390 (5' - ACTTCCACTACCATGAGCAATTC - 3') (SEQ ID NO: 35)	unique genomic region
PGK (5' - GTYGAYTTCAAYGTYCC - 3') (SEQ ID NO: 32) PGK (5' - ACACCDGGDGGRRCCGTTCCA - 3') (SEQ ID NO: 33)	Phosphoglycerate kinase

Primers	Genomic locus
ACT512f, Actin, primer - amplicon F (5' - ATGTGCAAGGCCGGTTTCG - 3') (SEQ ID NO: 17) ACT783r, Actin, primer - amplicon R (5' - TACGAGTCCTTCTGGCCCAT - 3') (SEQ ID NO: 18)	Actin
fusA-f2, elongation factor G, primer - amplicon F (5' - TCGCGTTCGTTAACAAAATGGACCGTAT - 3') (SEQ ID NO: 19) fusA-R2, elongation factor G, primer - amplicon R (5' - TCGCCAGACGGCCCAGAGCCAGACCCAT - 3') (SEQ ID NO: 20)	elongation factor G
RPB1-Af, largest subunit of RNA polymerase II, primer - amplicon F (5' - GARTGYCCDGGDCAYTTYGG - 3') (SEQ ID NO: 21) RPB1-Cr, largest subunit of RNA polymerase II, primer - amplicon R (5' - CCNGCDATNTCRRTRTCCATRTA - 3') (SEQ ID NO: 22)	largest subunit of RNA polymerase II
LR0R, long subunit rRNA gene, primer - amplicon F (5' - ACCCGCTGAACTTAAGC - 3') (SEQ ID NO: 23) LR5, long subunit rRNA gene, primer - amplicon R (5' - TCCTGAGGGAACTTCG - 3') (SEQ ID NO: 24)	long subunit rRNA gene
bRPB2-7.1R, second largest subunit of RNA polymerase II, primer - amplicon R (5' - CCCATRGCYTGYTTMCCCATDGC - 3') (SEQ ID NO: 25) fRPB2-5F, second largest subunit of RNA polymerase II, primer - amplicon F (5' - GAYGAYMGWGATCAYTTYGG - 3') (SEQ ID NO: 26)	second largest subunit of RNA polymerase II
NS1 (5' - GTAGTCATATGCTTGTCTC - 3') (SEQ ID NO: 27) NS4 (5' - CTTCCGTCAATTCCTTTAAG - 3') (SEQ ID NO: 28)	SSU, small subunit rRNA gene
SR1R (5' - TACCTGGTTGATQCTGCCAGT - 3') (SEQ ID NO: 29) NS4 (5' - CTTCCGTCAATTCCTTTAAG - 3') (SEQ ID NO: 28)	SSU, small subunit rRNA gene
Btub2Fd, beta-tubulin, primer - amplicon F (5' - GTBCACCTYCARACCGGYCARTG - 3') (SEQ ID NO: 30) Btub4Rd, beta-tubulin, primer - amplicon R (5' - CCRGAYTGRCCRAARACRAAGTTGTC - 3') (SEQ ID NO: 31)	Beta-tubulin

Table 4. Exemplary sequences of endophytes of the present invention

SEQ ID	MIC ID	Gene/Locus	Sequence
38	MIC-19994	ITS	TCTTTGGTCATTTAGAGGAAGTAAAAAGTCGTAACAAAGGTCTCCGTTGGTGAACACGCGGAGGGGATCATTTACAAGAAAG CCGAAAGGCTACTTTCAAACCATCGTGAACCTATCCAAGTTGCTTCGGCGGCGGGCTCCCTCTCGCGGGGTGCGGCA GCCCGCCCCCTCGGGGGTGGTGGGCGCCCGCGGAGGTATTAACACTCTCCCGTATTTATAGTGGTATTTCTGAAGTA AAAACAAATAAGTTAAAACTTTCAACAAACGGATCTCTTTGGTTCTGGCATCGATGAAGAAACGACGAGAAATGCGGATA AGTAATGTGAATTCAGAAATTCAGTGAATCATCGAATCTTTTGAACGCACATTCGCCCCGCTAGTATTTCTAGCGGGC ATGCCTGTTTCGAGCGTCATTTCAACCCCTCAAGCCCTGCTTGGTGTTTGGGGCCCTACGGCTGCCGTAGGCCCTGA AGAAGTGGCGGGCTCGTGCAACTCCGAGCGTAGTAATTCATTTATCTCGTAGGGAGGCGCGGGTGTCTCTCTGCC GTTAAAGACCATCTTTAAACCAAAGTTGACCTCGGATCAGGTAGGAATACCCGCTGAACTTTAAGCATATCAATAAG CGGAGGAAAAAGAAACCAACAGGGATTGCCCTAGTAACCGCGAGTGAACGGGCAACAGCTCAAAATTTGAAATCTGGC TTTCCGCCCGAGTTGTAATTTGTAGAGGATGCTTTTGGTGAGGTGCTTTCTGAGTTTCTTGGAAACGGGACGCCAGAG AGGGTGAGAGCCCCGTATAGTCGGCCACCGATCTCTGTAAAGCTCTTTTGACGAGTCGAGTAGTTTGGGAAATGCT GCTCAAAATGGGAGGTATATCTCTTTCTAAAGCTAAATATAGGCCAGAGACCGGATAGCGCAAAAGTAGAGTGATCGA AAGATGAAAAAGCACTTTGAAAAAGAGGGTTAAATAGCACGTGAAATTTGTTGAAAGGGAAGCGCTTGTGACCAAGACTT GCGCCGGGCTGATCATCCAGTGTTCTCACTGGTGCACTCGACCCGGCTCAGGCCAGCGTCGGTTCTCGCAGGGGGA TAAAAAGCTTCGGGAACGTGGCACCTTCGGGTGTGTTATAGCCCCGTCTTAATAACCCCGTGGGACCGAGGTTTCG CGCTCTGCAAGGACGCTGGCATAATGGTCATCAGCGACCCGCTCTTGAACACAGGAGTCGAGGTTTTCG CGAGTGTTTGGGTGTAAAAACCCGACACGCGTAATGAAAAGTGAACGTAGGTGAGAGCTTCGGCGCATCATCGACCGCAT CCTGATGTATTCGGATGGATTTGAGTAAGAGCGTATAGCCTCGGACCCGAAAGATGGTGAACATATGCTGAATAGG GGGAAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGCA
39	MIC-19994	LSU	TCTTGAGGGAAACTTCGGCGGTAAACCAAGCTACTAGATGGTTTCGATAGTCTTTTCGCCCCCATGCCCAAATTTGACG ATCGATTTGCACGTGAGAACCGCTCGAGCCTCCACAGAGTTTCTCTGGCTTCACCTATATTCAGGCATAGTTCA CCATCTTTCCGGTCCGAGGCTATACGCTCTTACTCAAAATCCATCCGAATACATCAGGATCGGTTCGATGATGCGCCG AAGCTCTCACTACGTTCACTTTCAATTACGCGTGGGGTTTACACCAAAACACTCGCGCAAAACCTCGACTCCTTT GGTCCGTGTTTCAAGACGGGTGCTGATGACCATATGCCAGCGTCTTTGCAGAGCGCGAAACCTCGGTCCCCACAG GGGTATTAAGCAGCGGGCTATAACACACCCGAAAGTGCCAGTTTCCCGAAGCTTTTATATCCCTTCGAGAACCGGAC GCTGGCCTGAGCCGGGTGAGTGCACCGAGTGAGAACACTGGATGATCAGCCCGGCGCAAGTGTGGTCACAAAGCGCT TCCCTTTCAACAAATTTACGTGCTATTTAACCTCTTTTCAAAAGTGCTTTTTCATCTTTTCGATCACTTACTTTGTGC GCTATCGGTCTCTGGCCTATATTTAGCTTTAGAAAGAGATATACCTCCCATTTTGGAGCGCATTTCCAAACTACTCG ACTCGTGAAGGAGCTTTACAGAGGATCGGTGGCCGACTATACGGGGCTCTCACCTCTCTGGCGTCCCGTTCCAG GGAACCTCAGAAAGGCACCTCACCAAAAAGCATCCTCTACAAATTAACAACCTCGGGCCGAAAGCCAGATTTCAAAATTTGAG CTGTTGCCGCTTCACTCGCCGTTACTAGGGCAATCCCTGTGGTTCTTTTCTCCGCTTATTGATATGCTTAAGT

SEQ ID	MIC ID	Gene/Locus	Sequence
40	MIC-19994	SSU	CTTCCGTCAATTTCTTTAAAGTTTCAGCCTTGGGACCATACTCCCCCAGAACCCAGAAAACTTTACTTTTCGTGTAAG GTGCCGAACGAGTCAAAATATAACATCGTCCGATCCCTAGTCGGCATAGTTTATGGTTAAGACTACGACGGTATCT GATCGTCTTCGATCCCCTAACCTTCGTTCTGATTAATGAAAAACATCCTTGGCAAATGCTTTCGAGTAGTTAGTC TTCAATAAATCCAAGAAATTCACCTCTGACAAATGAATACTGATGCCCCGACTGTCCCTATTAAATCATTACGGCG GTCTAGAAAAACAACAAATAGAACACACACGTCCTATTCTATTATTCATGCTAAATGATTCGAGCATAGGCCTTC TTTAAAGCGATCTAATTTGTTCAAAGTAAAAGTCCTGGTTCCCCGACACACCCAGTGAAGGGCATGCGGTTCCCCAG AGGGAAGGCCCGCGGACAGTGCACGCGGTGAGCGGACCGGCCAGGCCCAAGGTTTCAACTACGAGCTT TTTAACTGCAACAACATTTAATATACGCTATTGGAGCTGGAATTACCGCGGCTGCTGGCACGAGACTTGCCCTCCAA TTGTTTCTTCGTTAAGGGATTTAAATTTGTACTCATTCCAATTACAAGACCCAAAAGAGCCCTGTATCAGTATTTATT GTCACTACCTCCCCGAATCGGGATTGGGTAAATTTGCGCGCCTGCTGCCCTTCCTTGGATGTAGTAGCCGTTTCTCAG GCTCCTTCTCCGGGTGAGCCCTAACCCCTCCGTTACCCGTTGTCAACATGGCTGGCCAAAGACCCAGCCGTCGAAA GTTGATAGGGCAGAAATTTGAATGAACCATCGCCGGCGCAAGCCGTGCGATTGAGAAATTATTATGAATCACCA GAGCCCCGAAGGCATTGGTTTTTAATCTAATAAATACATCCCTCCGAAGTCGGGATTTTGTAGCATGTATTAG CTCTAGAATTACCACGGTTATCCAAGTAGTAAGTACTATCAAAATAAACGATAACTGATTTAATGAGCCATTGCA GTTTCGCGGTATAATTGCTTATACTTAGACATGCATGGCTTAATCTTTGAGACAAGCATATGACTACTGGCAGAAT CAACCAGGTAA
41	MIC-19994	Unique genomic region	TGCTGGTAGTCGGAATGAAAATGGCTGGTTCCAGGATATAACGGGTAATCGACTGCACTTTAAACAAGGCTATGCGA GTACTTTTGCAGACCATGGTCTTGCAGAAGCAGATCCGCCGACGAAAGAGCACCGGTTTCGGAGTCTGGAGGGTACAGTG TGCACGGATGTGTGCACTCCTGGATGGTAAACGTCCTCAACCCAGACAGGAGATGCGGAGATGGCACGTCCTGGCTTT GAGGTGTGTGGCTAGCCATGTGCCAAGCACGGAGGAGGTGAGTATTTGGCGGGTACAGCGCGCCTCCTTCTGCAC GCAGACCAATGCTTGAATTTGATGGAAGAGGGTCAGGAGGAGGAAGGCAATGGATGGGTATTCATATAATCTAGGAG ATCTCTACAAAGCCCCAAGGGCGGTTCAAGGAAGCAGAAGCCATGTACGAGCGGGCGCTTCGAGGCAAGGAGAAGGC ATGGGACCCAGACACAGTCGACACTCGACACAGTCAACAATCTGGGTCTCGTCGCCGACACAACAGCCAGCCAC ACCAACATCAAGTTCATTTCTGTTCCCCGTCTTTGTGCTGGCAGACAAAAACCTGATGGAACCCGAAAG

SEQ ID	MIC ID	Gene/Locus	Sequence
42	MIC-31593	ITS	AGGTGAACCTGCGGAGGGATCATTACACAATAAAATACGAAGGCCGTTGCGGGCTGGACTATTTATTACCCCTTGTC TTTTTGGCAGCTTGTGTTTCTTGGCGGGTTCGTGCGCCACGAGGACCACAATATAAACCTTTTTTATGCAGTTGC AATCAGCGTCAGTATAACAAATGTAATCATTTACAACTTTCAACAACGGATCTCTTGGTCTGGCATCGATGAAG AACGCAGCGAAATCGATACGTAGTGAATTCAGAAATTCAGTGAATCATCGAATCTTTGAACGCACATTCGCC CTTTGGTATTCAAAAGGGCATGCTGTTTCGAGCGTCATTTGTACCTCAAGCTTTGCTTGGTGTGGCGTTTTTG TCCTTTGGCCCGCCAAAGACTCGCCTTAAATGATTTGGCAGCCGGCTACTGTTTTCGACGCGCAGCACATTTTTC GCTTGCAATCAGCAAAAGAGGACGGCAATCCATCAAGACTCTCTTCTACGTTTGACCTCGGATCAGGTAGGGATAC CCGTGAACCTTAAGCATATCAATAAGCGGAGGAAAAAGAAACCAACAGGGATTGCCCTAGTAACGGCGAGTGAAGCG GCAACAGCTCAAAATTTGAAATCTGGCTCTTTTCAGAGTCCGAGTTGTAATTTGCAGAGGGCGCTTTTGGCTTTGGCAG CGGTCCAAGTTCTTTGGAACAGGACGTACACAGAGGTGAGAAATCCCGTACGTGGTTCGTAGCTATTCGCCGTGTAAA GCCCTTCGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGGAGGTAATTTCTTCTAAAGCTAAATATTGG CCAGAGACCGATAGCGCACAAGTAGAGTGATCGAAAGATGAAAAGCACTTTGGAAGAGAGTCAAACAGCACGTGA AATTGTTGAAAGGAAAGCGCTTCAGCCAGACTTGCTTGCAGTTGCTCATCCGGCTTTTCCCGGTGCACTCTTC TGCAGGACGGCCAGCATCAGTTTGGCGGTGGGATAAAGTCTCTGTACGTACCTTCTTCCGGTTGGCCTTATA GGGAGACGCCATACCAAGCTTGACTGAGTCCGCGCATCTGTAGGATCTGGCGTAATGGCTGTAAGCGGC CCGTCTTGAAACACGACCAAGGAGTCTAACATCTATCGGAGTGTGGGTGTCAAGCCCGAGCGGTAATGAAAG TGAACGGAGGTGGAAACCCGCAAGGTGCACCATCGACCGATCCTGAAGTTTACGGAAGGATTTGAGTAAGAGCAT GGCTGTTGGGACCCGAAAGATGGTGAACATATGCTTGAATAGGGTGAAGCCAGAGGAAACTCTGGTGGAGGCTCGCA GCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGCATAGGGCGCAAGACTAATCGAA
43	MIC-31593	RPB2	CCCATAGCTTTGCTTACCATAGCAGATTGGTATGTGTTACGGGCGACTGGTTGTGATCTGGGAAGGGAATGATAC TGGCGCAAAATACCAAGATCATAGCTGGGTGAATCTCACAAATGGGTGTAGGCGTGGATCGAGGATCCGGTAGTGG CTTGAGACGGCGGAGTCGATCCTTGCCCTTCAGTAGATCGCTCAGCTGCAGGCAAGCCCATCTTCATTTCTCGCCAT TCTTCCAAGTCTCGGGAGAGAAATGTTATCATTCAGTTTCTTCTTCCGCGCATCGAGGTATTCACACGACACCGT CTTGAATGAGACCTCTCCAGCCGTATGAGCCTGCTCGACTTCTTCTTGAATCTCTTGAATCTCTGTCTGCTGCTC TTGCTGTTTCCAGCTTGGCTGATTTCTTGGTAAAGATGAGGTGGTCCGGTTTGGCTTTCGAATATCG TTTTCTACAACGAACAAAGGCTCATGACACGACCCGCTCATGTGAATATCTTGAATCTCTGTCTGCGGAATATCAC GAATCAAACTCATCTCGTAAGACAGAGTACCATTCGGCGGAAGCTCCTGCACGACTGTGACAAAGCTGCTGAGCAT TGAATGAACACCAACCCAGACACCGTTAACGAAGACCTTGGTCGCATCCGGGTTCTGGTTCGTACTCCTCG AGAAAGTTGCATGTTTACGTTGTGTCATGAAGTCGATAATGGCGGATGCATCGCTACCAACACTGACATAACACATAA GAGACAAAGTTCTTGACAGACCCGAAGCCTGTCTTTCGGGCGTCTCAGCAGGGCAGACAAGACCCCAATGAGAGTT GTGAAGCTGTTCGGGCTTTTGCACCTTACCATCACGTCCAACGGGGTGTTCGTTTCGACGAGATGGGATAGTGTG GAGGCATAGGTGTATCGGTTCAACACCTGCGAAACACAGCCTTGGCAGATGCAGCCTTCTTCTGATCACCCCAAT TGCTGTAGCCAGAGAGTACTTCAGGCCGTTTGTGATGATGCTGGCTTTCACAGCCATTTGAACATGAAGTCTTG GTTGTTTTCCACGACCGCTGGAGGTACTTTGAGACATCCTTGGTGAAGTTCAGGAACAAGATTCGGAACAAGTTG GCAATCAGAGGTCAGCCAGATCTAGTCGCTTCTTTCCAAAGTGATCACGATCGTCC

SEQ ID	MIC ID	Gene/Locus	Sequence
44	MIC-31593	Beta-tubulin	GTTACACCTCCAGACCGGTCAATGCGTAAGTCTCGGCGCGCCTGAAAAACACCACGGGAACGACTGCTAACAGCCGCTACAGGGTAACCAAAATTTGGTCCGCCCTTCTGGCAGACCATTTCCGGCGAGCATGGCCTCGACGGCTCTGGTGTCTACACGGCACCTCTGACCTTCAGCTCGAGCGCATGAACGTCTACTTCAACGAAGTACGTCCCTCGTGAAGCTCCAACAGACAAAAGACCAATACTGATGTGCAGCAGGCTTCCAACAACAAGTTCTGTCGCCCTCGCTCGTCTGATCTCGAGCTCGGCTGATGACGCTGTCCGGCTGGTCCCTTTGGTCAGCTCTTCCGCCCCGACAACTTCGTCTTCGGCCA
45	MIC-31593	Unique genomic region	GTCGGGTCTACCGCAAGAGCAACTGTGCAAGTCCAGCTTTCAGTCCCTTTCGATCCCCCTCGCCCCAGGAAGAGATTCACCCCACGAGCGCAACCCAGTCCGAGGCGGTCCAGCGTCCACGTCTTATGAGCGTTTGACGCAACTCGCCAGCAGTTTCAGCACACCAAAAGAGCCTCCATCTACGTCTCCGACGCTTCCATCATCTAGCGCAACGCTCACCCATGGCTTCTCCGGTTTCCACACAGACTCCATGTCTCCCCCATCCATGAATCGTCTGATGCCGTGACCACTATGCTATCCTGGAGATCACCCCTAGAGCAACTACCGATGAGGTCAAGGTGCCCTACCGCCGACTACGGGTCTGCTACTTCTCAAAGTGACGCGAAGAAGTACCGGACACTGCAGCGGCCCTTCGACGCTTTCGATGGACCCGCAATCCCGCGAAGCTTACGACGCAACCTATCAACAACTGCCGACACCAAGTATCGTCGCTAGCATTTGGTAGATCCTGGACTCGGGAAAGCTATGGCGACAGGACACGCCCCACGGAGACCCAGTAATCCAGAAGAGGAAGAGGAGGAGGAAAGT
46	MIC-96038	ITS	TCTTTGGTCATTTAGAGGAAGTAAAGTCTGTAACAAGTCTCCGTTGGTGAACCCAGCGGAGGGATCATTTACCGAGTGTAAAACTCCCCAAACCATTTGAACTTACCACTGTTGCTTCGGCGGCTCGCCCCGGCGCTTCGCGGGCCCCGACCCAGGCGTCCGCGGAGGCTCCAAACTCTTGTCTTTAGTGATTTCTGAGTGGCATAAGCAAAATAAATCAAACTTTTCAGCAACGGATCTCTTTGGTTCTGGCATCGATGAAGAACGACGCAAAATCGATAAGTAATGTGAATTCAGAAATTCAGTGAATCATCGAATCTTTGAACGCACATTCGCCCGCCAGTATTTGGCGGGCATGCCTGTCTGAGCGTCATTTTCAACCTCAGGACCCGTTTCGCGGGACCTGGCGTTTGGGATCAGCCTGCCCTGGCGGGCTGGCCCTGAAATCCAGTGGCGGTTTCCCTCGCAACTCTCCGTGCAGTAATTAACCTCTCGCGCAGGATAGCGGTTGAACACACGCCGTTAAACCCCCACTTCTCAAGGTTGACCTCAGATCAGGTAGGAATACCCGTGAACTTAAGCATATCAATAAGCGGAGGAAAACCAACAGGGATTGCCCTAGTAACGGCGAGTGAAGCGGCAACAGCTCAAAATTTGAAATCTGGCCTCACGGTCCGAATTTGTAATTTGTAGAGGATGTTTCTGGCGACGTGTCTTCGAGTTCCCTGGAACGGGACGCCATAGAGGTGAGAGCCCCGTCCGGTCGTACACCTAGCCTCTGTGAAACTCCTTCGACGAGTCGAGTAGTTTGGGAATGCTCTAAATGGAGGTATACGTCTTTAAAGCTAAATACCGGCCAGAGACCAGTACGCAAGTAGAGTGATCGAAGATGAAAAGCACTTTGAAAAGAGGTTAAATAGTACGTGAAATTCGTAAGGGAAGCGCTTATGACCAAGCTTGGGCTCGGTGAATCATCCGGCGTTCTCGCGGTGCATTTGCCGTCCAGGCCAGCATCAGTTCGCGCCGGGGATAAAGGTTTCGGGAATGTAGCTCCTTCGGGAGTGTTATAGCCCGTTTCGTAATACCTGGCGTGGACTGAGGTCGCGCTCTGCAAGGATGCTGGCGTAATGGTCATCAGTGACCCGTTCTTGAACACGGACCAAGGAGTCGTCTTCGTATGCAGTGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGTAGAGAGAGCTTCGGCGCATCTCCGACCGATCTGTATGTTCTCGGATGGATTTGAGTAAGAGCATACGGGGCCGGACCCGAAAAGAGGTGAACATATGCCTGTATAGGGTGAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAAATATGGGCATGGGGCGGAAAGCTAATC
47	MIC-96038	Actin	ATGTGCAAGGCGGTTTCGCCGGTGACGATGCTCCCCGAGCTGTTTTCCGTAAGTCAACCCCACTTTCGTTCCCAAGCTCTTAATCGCCACACCTGGCGATATGGGCTTTGGGGCCTGTAAGCAGCCGACACAAGACTAACCGGATGCGCCAGCTTCCATTGTCCGTCCGCCCGTCACCATGGGTAAGTACGCGCGCAATGACACCTGTGAGCCCCCTCGACAGCGCACAGGCTTGACCATTCGATAGTATCATGATTGGTATGGGACAGAAGGACTCGTAC

SEQ ID	MIC ID	Gene/Locus	Sequence
48	MIC-96038	Beta-tubulin	CCGGACTGGCCGAAGAGCGAAGTTGTCTGGGACGGGAAGAGCTGACCGAAAGGACCGGCGCGGACGGCATCCATGGTAC CGGGCTCGAGATCGACGAGGACAGCGCGAGGAACGTACTTGTGGCAGAGGCTACAGAGGTCAGCTTGGCCACA GACTCGGGGATACTCCAAATTGCTCACCTCGTTGAAGTAGACGCTCATCGCTCGAGCTGAGCTCAGAGGTGCCG TTGTAGACACCGTTGCTGTCGAGGCCATGCTCGCCAGAGATGGTCTGCCAAGGACGACCAATCTGTTTACCCCT GCTCGGAGGTTAGACATGGTAGCGGATATCACATATGGCGAAGTACTTACGCACTGACCGGTCTGGAGGTGAACC GATGATCGTGATCACTTCGGGAAAAAGCGCCTTGACCTGGCTGGGCCCTCTTAGCTAAATTTGTTCCGCAACATTA TTTCGAGGATCAACAACGAGCTGTGCACCTACCTCAGGCGATGTGTGAGGGCGGCGAGGAACCTTCAACCTCGCTGT CGGCATCAAGCCTGGCACACTGTGGAACGGGTTGAAGTACTCTTTGGCAACAGGCAACTGGGGAGACCAAAAGAAAG GCAATGAGCTCGGTTGCTGGAGTGTCCAGGTTCTCAACCGCTACACATTTGCGTCAACCTTGTCTCATTTGAGGC GCACCAACACCCCATTTGGCCGTGATGGAAGCTGGCGAAGCCTCGGACGCTACACAACACGCAATTTGGGTTCTTGT GTGTCCCGCCGAGACCCCGAGGGTCAGGCTTGTGGTTTGGTGAAGAACCTGTCACTGATGTGTACGTTTCCGTT GGCACACTAGCGAACCTCTCTACGGATACTTCAACAACCTGGCATGGAAGTGCTCGAAGAGTACGAGCCCCCAGC GGTTCCTCAACGCCAACAGGTGTTCAACAACGCTGCTGGTGGTGTGCACACAAGCCCCGAAAGATCTCGTGGGA TAGCATCATGATCTCGGGCGCTATGGTGACCTGAACCATGAAGCTTCCGTCACTCCGCGACATTCGGGATCGAGAG TTCAGGGTCGTACGGATGCTGGTCGTGTTATGCGCCCATGATTCACCGTGACGCAAGAACAGCTAGACGGGC CCGAGAAAGGGTCGTTGTGCATGACCAAGGAGCATCTTGCCGGTTTGGATGACTGGCATCTGGTCAACGAGGAGAG GGAAGAGATGGCCACGGGCTGGGAGTACCTCGTGAAGAGTGGTGTATTTAGTACTTTGGACGCCGGAAGAAAGAGAG ACGGCAATGATTTGCATGACACCAAGAGACTTGGAGTCTTACCGCAAGGAGAAAGTACCTCGATCAGAAAACCCGAGG AGCACAACTGGAAGCCGAGCCCCAACAAAGCGACTCAAGACGAAGACCAACCCGACGACACACATGTACACCCACTG CGAGATTCACTCCAGTATGATCTCGGTATCTGCGCCAGCATCATCCCCCTTCCCGGATCATAAACGAGCATGTCTC TACGCCACCAAGACCTCGAGATTACTTAATAATTGCATCTAGTCGCCCCCGTAATACTTACCAATCTGCCATGGGC AAGCAGGCCATGGGC
49	MIC-96038	RPB2	
50	MIC-96038	RPB1	GAATGCCCCGGTCATTTTGGTCACATCGAGCTGGCAAGCCCGTTTACCACCCCGGCTTCATCAAGAAAAGTCAAGA AGATTTTGGAGATTGTCTGCCACAACTGCAGCAAGGTCTTGGCCGATGAAGTTGGTCTCACCTGATCCATGTCTTG TTCCTTAGATGCTAACATGGACCTCTCAGAGCGACCCCGAGTTTGTCACTCGATCCGTACCGCGCACCCGAAAGT CCGCTTCCAGCGAGTCTGGGCTGTGTCAAGAAGAAGCGGAAGTGTGAGAACGAGGATCGCCAAGACAAGAAGGAA GAGGAGTTCGCGCCCGCATGAAGCCGAGACGCAACAACCGCGGCTGTGGAACGAGATGCCCGGGTTCGTGTC AAGCTGGTTTTCGTCTCAACGCGCAGTTCGAGATCAAGGAAGAGGGCGGAGCTAAGCGCAAGGATACTCAAGTTAT CCTGCCCGACCAAGCTCACACAATCCTGCGGCGGATATCGGAACAGGACCTCCGACACATGGGCCTCAACTCAGAG TATGCCCGCCCGAGAGTGGATGGTTCTTACCGTCTTCCGGTCCCCCGCTCCCGTTTGTCCAAGTATTTCCATGG ACGGCACCTGGCACGGGAATGCGGAACGAGGATGATTTGACTTACAAGCTTGGTGATATCATCCGAGCCAACGGAAA CGTCAAGCAGGCTATCCGCGAAGGCTCGCCGGCCCAATTTGCTCGCGATTTTGAAGAAGCTGCTCCAAGTACCATGTA GCCACCTACATGGATAAATGATATTGCTGGA

104

SEQ ID	MIC ID	Gene/Locus	Sequence
54	MIC-33414	LSU	TCCTGAGGGAAACTTCGGGGTAACACGCTACTAGATGGTTCGATTAGTCTTTGCGCCCCCATGCTCAGATTTGACG ATCGATTTGCACGTCAAAACCGCTGCGAGCCTCCACAGAGTTTCTCTGGCTTACCCCTATCCAAGCATAGTTCA CCATCTTTGCGGTCCAAGGTTAACGCTCTACTCAAATCCATCCGAAACATCAGGATCGGTGATGCGCCG AAGCTCTCACCTACGTTCACTTTCACTACGCGTGGGGTTTACACCCAAACACTCGCGCAAAACCTTGACTCCTT GGTCCGTGTTTCAAGACGGGTGCTGATGACCATTAACGCGAGCATCTTTCAGATGCGGGAACCTCGGTCCCCGCG AGGGCATTAACGCCCGGGCTATAACACTCCCGGAGGAGTACGTTCCAGGACCTTTATCCCCCGCGAGAACCGGA TGCTGGCCTGAGCCGGGGAGTGACCGGTGAGAACACCGGATGATCCGCCGGCGCAAGTCTGGTCACAAGCGC TTCCCTTTCAACAATTTACAGTGCTATTTAAACCTCTTTTCAAAGTGCTTTTCACTCTTTTCGATCACTCTACTTGTG CGCTATCGGTCTCTGGCCGGTATTTAGCTTTTAAAGAAAATTTACCTCCCATTTTGAAGAGCATTTCCAAACTACTC GACTCGTCAAGGAGCTTTACACAGGCTAGGCATCCAACATAACGGGCTCTCACCTCTATGGCGCCCCGTCCA GGGACTCAGAAGGTGCCGCGCTAAAGCTTCTCTGCAAAATTAACAACCTCGGGCCGAAGCCAGATTTCAAATTTGA GCTGTTGCCGCTTCACTCGCCGTTACTAGGGCAATCCCTGTTGGTTTCTTTTCTCCGCTTATTGATATGCTTAAG TTCAGCGGGTC
55	MIC-33414	RPB1	GAGTGCCAGGTCACTTTGGCCACATTTAGCTATCCAGACCCGTTTCCACCCCGGGTTTCATCAGGCGTGTCAAAA AGTTGCTCGAGATGGTCTGCCACAACCTGCAGCAAGGTGTTGGCTGATCGTGTAGTGACCTTGCCTGACCGAGTG ATGATTTGTTTGGCATGCTAACTCTTACCAGGAGGACGAGCAATATGCTGCTGCCATGCGGATTCGGGACCCCA AAGTACGCTTCAAGCGAGTTTGGGATATTTGCAAGAGTAAGAACGCTGCGAAAACGAAAGTGCACAAGGGGAAAGA TGGCGAGTTCAAAACCGACAGCGAAAACCAAGCCGAGAGGTTGGCCATGGAGGATGTGGCAACACGCGAGCCAGTC ATTGCCAGCAGGCTCTCACCTGTGGGGCAGCGTTCGAGACCAAGGACGAGGATGGTGTGAAGACCAAGGAGAAGA AGGTCAATCAACCCAGAAATGGCCCTGAACATCTTCCGTGCGATGTGGAACGACGAGATGATTGACATTTGGCCTCAA TATTTCCCAAGCTCGTCCGGAATGGATGATCATCACGGTTCTTCTGTCCCGCTCCTCCGGTGCGCCCAAGTAT TCCATGGACGGAACCTGGAACAGGCTTGCGGAATGAGGACGATCTGACGTATAAACTCGGCGATATCATCCGCGCCA ATGGCAACGTCCGCCAGGCTATTGCCGAGGGCTCTCCTCAGCATATCATCACCGACTTTGAGAACCTACTCCAGTA CCACGTCGCTACGTACATGGATAATGACATCGCCGGT

106

SEQ ID	MIC ID	Gene/Locus	Sequence
59	MIC-33414	Unique genomic region	GAGGAGGAGGAGGAGGTTGGAGAGGGAGGCGCTTGGCTGCCAGAGGCGCTTGTGAGGTGAGGCGGGTTATGGCGGCTGCTGGAGGATACGCTGCTTGGGACGGGCGGAGTGGTGGGCGGTGGTGGCGGTGATGGTGGCGAGTGAGGGTCCGAGAAAAGGGCCGACGTTGGCGGATATCGAGGCCGTGGGTGCTTCACTGGATGATTGGCATTCTTGGTCCGTGTTCAACGCCGGGTATGTAGCGCCGAGCGGTTTCCGCGGTGTATCGTGGTGGCGCGGTTTCAGGCGGCGTTGGGCGGGAAGGCCGGGTGGTGGTGAAGGCGATGAGCGGGAGGAGGCGCGGTAGTGTGAAGGGGCAGAGAGAAGGGGTAGGATATTTTGAAGAAGGAGGGGAGGTGGACGCCGCGGACCCGATCGTCAAGGTGTACGGAATTGAGAGAAAAGGGAGCAGGGTCGAGGTGTGGCCGACGGACTCCGGGGCTGGGCATCGGGATCAGGGCTGCCTGGTGA
60	MIC-68390	ITS	GCGTCCGACGCCGAGGAAATAGTCTGGGAGACGGACGTCTTTGGTGGTCAATTTAGAGGAAAGTAAAGTCTGTAACAAAGTCTCCGTAGGTGAACCTGCGGAGGGATCATTTACACAACAAAAATGAGGGTGTGGTCTTGGCAACAGCGTCCGCCCAAGTATTTTCACCCATGTCTTTTGCGCACATTTTGTCTTCTGGCGAGTTCGTCGCCAGGTTTCGTCGCCACAGGACCCAAACCATAAACCTTTTATGCAAGTTCGAATCAGCGTCAGTATAATAATCAATTTATAAACTTTCAACAACGGATCTCTTGGTCTTGGCATCGATGAAGAACGACGCGAAAATGCGATACGTAGTGTGAATTCAGAAATTCAGTGAATCATCGAATCTTTGAACGCACATTTGCGCCCTTTGGTATTTCCAAAGGCATGCCCTGTCGAGCGTCATTTGTACCTCAAGCTTTGCTTGGTGGTGGCGCTCTTTTGTCTCTCCCTTGTGGGGAGACTCGCCTTAAACGATTGGCAGCCGACCTACTGTTTTTCGGAGCGCAGCACAAATTTGCGCCTTCCAAATCCACGGGGCGCATCCAGCAAGCCTTTGTTTTCTATAACAAATCCACATTTTGACCTCGGATCAGGTAGGGATACCCGTGAACCTTAAGCATATCAATAAGCGGAGGAAAGAAACCAACAGGGATTGCCCTAGTAACGGCGAGTGAAGCGCAACAGCTCAAAATTTGAAATCTGGCTCTTTTCAGAGTCCGAGTTGTAATTTGCAGAGGGCGCTTTGGCTTTGGCAGCGGTCCAAGTTCCCTTGGAACAGGACGTACAGAGGGTGAGAAATCCCGTACGTGGTCGTAGCTATTGCCGTGTAAA
			GCCCCCTTCGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGAGGTAAATTTCTTCTAAAGCTAAATATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAAGCACTTTGGAAAAGAGAGTCAAAACAGCACGTGA
			AAATGTTGAAAGGGAAGCGCTTGACGCCAGACTTGCTTGCAAGTTGCTCATCCGGGCTTTTGCCCCGTGCACTCTTC
			TGCAGGCAGGCCAGCATCAGTTTGGCGGTGGGATAAAGTCTCTGTCAATGTACCTCTCTTCGGGGAGGCCCTTATA
			GGGAGGCGGACATACCAACGCTAGACTGAGGTCCGCGCATCTGTAGGATGCTGGCGTAATGGCTGTAAGCGGC
			CCGTCTTGAAACACGGACCAAGGAGTCTAACATCTATCGAGTGTGGTGTCAAGCCCGAGCGCGTAATGAAAG
			TGAACGGAGGTGGAAACCCGCAAGGTGCACCATCGACCGATCCTGAAGTTTACGGAAGGATTTGAGTAAGAGCAT
			GGCTGTTGGGACCCGAAAGATGGTGAACATGCTTGAATAGGTGAAGCCAGAGGAAACTCTGGTGGAGGCTCGCA
			GCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGCATAGGGGCGGAAAGACTAATCGA

SEQ ID	MIC ID	Gene/Locus	Sequence
61	MIC-68390	second largest subunit of RNA polymerase II	CCCATGGCTTGCTTGCCCATAGCAGATTGGTAGGTGTTACGAGGCGACTGGTTGTGATCTGGGAAAGGAATGATAC TGGCGCAAAATACCCAGAATCATGGCCGGATGGATTTACAGTGAGTGAGGCATGGATACGAGGATCTGGTAGGGG CTTGAGACGGCGAAGACGGTCTTACCTCTGTGGAGCGCTCCGCTGAGGTAGGCCCATCTTCATCTCTCGCCAT TCTTCTAAGTCTCTGGGAGAGAATGTAATCATGGCAGTCTCTTCTCTCAGCATCGAGGTATTCGACAACACCAT CTTGGATAAGACCTCTCCAGCCATAAGTAGCCTGCTCAACCTCCTCCTGACTCCATCCTTGCCTTGTGCTAGTCTC TTGTTGTTTTCAGCCTTGAGCTTGTTACTAATCTCCTTGGTGAAGATGAGGTGGTTTTCGGTTTTCGAAATATCG TTCCTACAAACAAATAGAGGTCTCATGACACGACGACCATCTGTGAAGATCTTGAACCTCTCGGTTCACGAATGTCA GAATCAAACTCATCTCGTAAGATAGAGTTCCGTTTTCGTGAAAGCTCCTGCACGACCGTAACAAAGTTGTTGAGCAT GGAATGAACCCCAACCCAAACACCATTAACGAAAAACCTTGGTTCGCATCGGGGTTTTCGGTTCGTATTCCTCG AGCAGCTGCATGTTACGTTGCGTCATGAAGTCGATAATGGGCGACGCATCGCTACCGACACTGACGTAACACATGA GAGACAAAGTTCTTGACCCAGACCCGAAGCCTGTCTTCAGGCGTCTCAGCAGGGCAGACAAGACCCAGTGGGAGTT GTGCAGTTGACGGGGCTTGGCCAAATTTACCATCACGTCCAACAGGAGTATTTGTTTCGACGAAGATGGGACAAATGTG GAGCGTAAGTGTATCGGTTCAACACCTGCGAGACACCGCCTTGGCAGATGCTGCCTTCTTCTGGTCACCCAGT TGCCTGTAGCCAGAGAGTACTTCAATCCGTTTCGTGATGATGCTGGCTTTTACAGCCATCTGAACGTTAAAGTCTTG GTTGTTCTCAACGCACCGCTGGAGGTACTTGTAGACGTCTTGGTAAGCTTCAAGAAGAGGATACGGAAACAAGTTG GCGATTAAAGGTCCAGCAAGATCCAGGCGCTTCTTTCCGAAATGATCACGGTCATCC
62	MIC-68390	unique genomic region	CTTCCAGGCATAGTAATGTGGATATTAGGTGAGAGCGAAATATAAGTGTCCCTAGAAAGTGATAGTGAGAAAGGCTAT GGTGAGGTTGAAGAAGGTAGATGGCATATTGGTAATTATGAACATCATATAATCTAATGAGTCGAAATCATTAAT TTTTTTTTTAAACTAATTACCATTTACTCTGTTTCATTTCTAATCCTTTTTTGTGTTTCATTCATGCTAGGCCTAGAGA TAGAATTGTGACTAGATAAAGGCTATAATTATTATAGTAGAGGTTTTTAATTTGTTTGAATTTGCTCATGGTAGTGGA AGT

SEQ ID	MIC ID	Gene/Locus	Sequence
63	MIC-68178	ITS	TCTTGGTCATTTAGAGGAAGTAAAGTCGTAAACAAGGTTTCCGTAGGTGAACCTGCGGAAGGATCATTACCTAGAG TTTTGTGGAACCTTCGGTCTGCTACCTCTTACCCATGTCTTTTGGTACCTTCTGTTTCTCGCGGGTCCGCCCGCCGG TTGGACAACATTCAAACCCCTTTCAGTTGCAATCAGCGTCTGAAAAAATTAAATAGTTACAACCTTCAACAACGGA TCTCTTGGTTCTGGCATCGATGAAGAACGACGAGAAATCGATAAGTAGTGTGAATTCAGAAATTCAGTGAATCAT CGAATCTTTGAACGCACATTCGCCCCCTTGGTATTCATGGGCATGCCCTGTTGAGCGTCATTTGTACCTTCAAG CTCTGCTTGGTGTGGTGTCTCGCCCTCGCGCGCAGACTCGCCCTTAAAAACAATTTGGCAGCCCGCGTATTG ATTTCCGGAGCGCATACATCTCGCGCTTTTGCACTCATAAACGACGACGTCCAAAAAGTACATTTTACACTCTTGACC TCGGATCAGGTAGGGATACCCGCTGAACTTAAGCATATCAATAAGCGGAGAAAAAGAAACCAACAGGGATTGCCCT AGTAACGGCGAGTGAAGCGGCAACAGCTCAAAATTTGAAATCTGGCGTCTTTGGCGTCCGAGTTGTAATTTGCAGAG GGCGCTTTGGCATTTGGCAGCGGTCCAAGTTCTTTGGAACAGGACGTCAAGAGGTTGAGAAATCCCCTACGTGGTCTG CTAGCCCTTACCGTGTAAAGCCCCCTTCGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGGAGGTAAATTTTC TTCTAAAGCTAAATACTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAAGCACTTTGGAAAG AGAGTTAAAAAGCACGTGAAATTTGTTGAAAGGAAAGCGCTTGACCCAGACTTGCCTGTAGTTGCTCATCCGGGTT TCTACCCGGTGCACCTCTTACGGGCAGGCCAGCATCAGTTTGGCGGTTGGATAAAGGTCCTGTCTCATGTACCTC CCTTCGGGGAGATCTTATAGGGGAGACGACATGCAACCGACCAAGGAGTCTAACATCTATGCGAGTGTTCGGGTGTCAAG CGTAATGGCTGTAAGCGGCCCGTCTTGAAACACGGACCAAGGAGTCTAACATCTATGCGAGTGTTCGGGTGTCAAG CCCGAGCGCGTAATGAAAGTGAACGGAGGTGGGAACCTTTTCGGGGTGCACCATCGACCGATCCTGATGTCTTCGGA TGGATTTGAGTAAGAGCATAGCTGTTGGGACCCGAAAGATGTTGAACTATGCTTGAATAGGTTGAAGCCAGAGGAA ACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGCATAGGGCGAAAAGAC CCAGACTGGCCGAAGACGAAAGTTATCGGGACGGAAGAGCTGGCCGAAGGGCCGGCGGACAGCGTCCATTGTAC CGGGCTCCAAGTCGACGAGGACGGCACGGGGAACGAACTTGTTCAGAGAGGCTGCGGGAGGTCAAGCACTCGCAGT CCGTCTCAGGAAAGCGTGTGCTTCTAGTACCTCGTTGAAGTAGACGTTTCATGCGCTCGAGCTGGAGGTCCGAGGT GCCGTTGTAGACACCGGAGCCGTGAGGCCATGCTCGCCGGAGATGGTCTGCCAGAGGACGACCGATTTGGTTA CCCTGTCCCTTGTGAGTGGCTGCCGTCCATGAGAGAACATGCAAGTGGTGTACTTACGCACTGACCCGCTCTGGAGGTGA ACC
64	MIC-68178	Tubulin	

SEQ ID	MIC ID	Gene/Locus	Sequence
65	MIC-07010	ITS	TCTTGGTCATTTAGAGGAAGTAAAGTCGTAACAAGGTCTCCGTAGGTGAACCTCGGAGGGATCATTACACAATA ACATATGAAGGCTGTACGCCGTGCGCCCCCGGGCCAGTTGGCTGAGGCTGGATTATTTATACCTTGTCTTTTG CGCACTTGTGTCTTCTGGCGGGTTTCTGGCGGCTCCAGGACCACCATAAACCTTTTATGCAGTTGCAATCA GCGTCAGTACAACAAATGTAATCATTTACAACCTTTCAACAACGGATCTCTTGGTCTGGCATCGATGAAGAACGC AGCGAAATCGGATACGTAGTGTGAATTCAGAAATTCAGTGAATCATCGAATCTTTGAACGCACATTCGCGCCCTTG GTATTCAAAAGGGCATGCCCTGTTTCGAGCGTCATTTGTACCCCTCAAGCTTGGTGGTGGCGGTTTTTGTCTTT GGTTTGCCAAAGACTCGCCTTAAACAGATTGGCAGCCGGCCTCTGTGTACGCAGCGCACATTTTTCGCGCTTG CAATCAGCAAGAGGGCGGCACTCCATCAAGACTCTTCTCACGTTTGACCTCGGATCAGGTAGGGATACCCGCTGA ACTTAAGCATATCAATTAAGCGGAGGAAAGAAACAAACAGGGATTGCCCTAGTAACGGCGAGTGAAGCGGCAACAG CTCAAAATTTGAAATCTGGCTCTTTTAGGGTCCGAGTTGTAATTTGCAGAGGGCGCTTTGGCTTTGGCAGCGGTCCA AGTTCTCTTGAACAGGACGTACAGAGGGTGAGAAATCCCGTACGTGGTCCGTAGCTATTTGCCGTGTAAGAGCCCTT CGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGGAGGTAAATTTCTTCTAAAGCTAAATATTTGGCCAGAGA CCGATAGCGCACAAGTAGATGATCGAAAGATGAAAAGCACTTTGAAAAGAGAGTCAAACAGCACCTGAAAATTTGTT GAAAGGAAAGCGCTTCAGCCAGACTTGCTTGCAGTTGCTCATCCGGGCTTTTCCCGGTGCACCTCTTCTGTAGGC AGGCCAGCATCAGTTTGGCGGTGGGATAAAGTCTCTGACACGTTCTTCTTCCGCTTGGGTTGGCCATATAGGGGAGA CGTCATACCAACGCTGACTGAGGTCGCGCATCTGCTAGGATGCTGGCGTAATGGCTGTAAGCGGCCCTCTT GAAACACGGACCAAGGAGTCTAACATCTATGCGAGTGTGGGTGTCAAGCCCCGAGCGCGTAATGAAAGTGAACGG AGGTGGGAACCCGCAAGGGCGCACCATCGACCGATCTGAAAGTTTACGGAAGGATTTGAGTAAGAGCATGGCTGTT GGGACCCGAAAAGATGGTGAACATATGCTTGAATAGGTTGAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCT TGACGTGCAAAATCGATCGTCAAAATTTGGGCATAGGGGCGAAAGACTAATCGA
66	MIC-07010	phosphogly cerate kinase	ACACGGCAGGGCCGTTCCAGAGGATGGTCTGCGCCTCATCGATGGCCTCCTTGTAAAGCTTGATCGACTTCTCTC CACAGTCGAGACCCATCCAGCCATCTGGGATACCATCCTTGTCTCGGCATAACCGACGTTGGCGTCTTGTGAA CTTGTGGCGGTGATGTAGTCAACAGGCAGACAATCTTGACATTTGTTCTTCTCGCCTTCTCCACGAGGTCTTG ACGGTCTTGCTACCACTCATCGAACAAAGCTTTCACCAATCTTGACGCCCTCGAGAGTCTTCTTGAAGGTGAAGG ACATGCCCTCCGCAATGATCAGGCTGTTGACCTTGCCAAAGCAGGTTGTGATCAATTTGAATCTTGTGAGAGACCTT GGCACCAACCAAGGATGGCGAGGAAAGTCTGCTTGGGTTCTCAAGCGCTTGTGCAAGTAATCAAGCTCCTTCTTG ACAAAGGAAGCCAGAGGCCCTTTTGTGGAGGTGACACCGACCATGGAGCTGTGCGCGCGGTGAGCAGTACCAAAAG CGTCGTCTGTAAACCGTCAGCCTCGTGTCTTTCGCCCATGAATTCATAGTTACTTACTAATGTAGACGTGCGCCAGA GCAGTCAGTCCCTTCTTAAACTCATCGACCTTGTCTTGTGACCTTCTGCTTCTTGGCCGCATCATCCTTGTAGC TACCCCTCCTCCTCAGCGTGAAGCGCAGGTTCTCGAGGAGGATGACCTGACCAACCGCTAGCGTTGTTGACGGTATC CTCTACCGACTTGCCGACGAGTCGTGGTGAAGGTAAACGCTCTTTCGCCGAGGAGCTTCTCGAGTTTCGGGAACAAC GGCTTGAGGCTGTACTTCGCAATTTGGGCTTACCGTCTGGCCGGCCAAGGTGGGACATGAGAAATGACGGCCTTGGCGC CATTTGTCGACGGCGTACTTGATTGTGGGAAGTGGCCCAACAAATGCGCTGGTGTGTTGGTGATCTTCTTGTGCGCGCTC GAGAGGACGTTGAAGTCGACC

SEQ ID	MIC ID	Gene/Locus	Sequence
67	MIC-48747	ITS	TCTTGGTCATTTAGAGGAAGTAAAAAGTCGTAAACAAGGTCTCCGTAGGTGAACCTCGGAGGGATCATTTACAAGTGA CCCCGGTCTAACCAACCGGGATGTTATAAACCCCTTTGTTGTCGACTCTGTTGCCTCCGGGGCGACCCCTGCCTTCGG GCGGGGCTCCGGGTGGACACTTCAAACCTCTTGCCTAACTTTGCAGTCTGAGTAAACTTAATTAATAAATAAACAAC TTTTTAACAACGGATCTCTTGGTCTTGGCATCGATGAAGAACGACGAGAAATCGATAAGTAATGTGAATTCAGAA TTTCAGTGAATCATCGAATCTTTGAACGCACATTCGCCCCCTGGTATTCGGGGGGCATGCCTGTTTCGAGCGTCAT TTCACCACTCAAGCCTCGCTTGGTATTTGGGCATCGCGGTCCGCCGCTGCTCAAAATCGACCGGCTGGGTCTTCTG TCCCCTAAGCGTTGTGAAACATATTCGCTAAAGGGTGTTCGGGAGGCTACGCCGTAAAAACAACCCCATTTCTAAGG TTGACCTTCGGATCAGGTAGGGATACCCGCTGAACCTTAAGCATATCAATAAGCGGAGGAAAAAGAAAAACAACAGGGAT TGCTCTAGTTAACGGCGAGTGAAGCAGCAATAAGCTCAAAATTTGAAATCTGGCGTCTTCGACGTCCGAGTTGTAATTT GTAGAGGATGCTTCTGAGTAAACCAACCGACTTAAGTTCTTGGAACAGGACGTATAGAGGTTGAGAAATCCCCTATG CGGTCCGAAAGGTGCTCTATACGTAGCTCCTTCGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGGAGGTA AATTTCTTTCTAAAGCTAAATATTTGGCCAGAGACCGATAGCCACAAGTAGAGTGATCGAAAGATGAAAAGCACTTT GGAAGAGAGTTAAAAAGCACGTGAAATTTGTTAAAGGGAAGGATGCAACACAGACTTGCTCGCGGTGTTCCGCC GGTCTTCTGACCGGTCTACTCGCCGCTTGCAGGCCAGCATCGTCTGGTCCGCTGGATAAGACTTGAGGAATGTA GCTCCTTCGGGAGTTATAGCCTCTTGTGATGCAGCAGCGCCGGCGAGGTCCGCGCTTCGGCTAGGATGCTGG CGTAATGGTCGTAATCCGCCCGTCTTTGAAACACGGACCAAGGAGTCTAACATCTATGCGAGTGTTCGGGTGTCAAA CCCCACGCGTAATGAAAGTGAACGGAGGTGAGAACCCGCAAGGTGCATCATCGACCGATCCTGATGTCTTCGGATG GATTTGAGTAAGAGCATAGCTGTTGGGACCCGAAAGATGTTGAACTATGCCTGAATAGGTTGAAGCCAGAGGAAAC TCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGTATAGGGCGAAAAGACTAATCG
68	MIC-50414	ITS	TCTTGGTCATTTAGAGGAAGTAAAAAGTCGTAAACAAGGTCTCCGTAGGTGAACCTCGGAGGGATCATTTACAAGTGA CCCCGGTCTAACCAACCGGGATGTTATAAACCCCTTTGTTGTCGACTCTGTTGCCTCCGGGGCGACCCCTGCCTTCGG GCGGGGCTCCGGGTGGACACTTCAAACCTCTTGCCTAACTTTGCAGTCTGAGTAAACTTAATTAATAAATAAACAAC TTTTTAACAACGGATCTCTTGGTCTTGGCATCGATGAAGAACGACGAGAAATCGATAAGTAATGTGAATTCAGAA TTTCAGTGAATCATCGAATCTTTGAACGCACATTCGCCCCCTGGTATTCGGGGGGCATGCCTGTTTCGAGCGTCAT TTCACCACTCAAGCCTCGCTTGGTATTTGGCAACGCGGTCCGCCGCTGCTCAAAATCGACCGGCTGGGTCTTCTG TCCCCTAAGCGTTGTGAAACATATTCGCTAAAGGTGCTCGGAGGCTACGCCGTAAAAACAACCCCATTTCTAAGG TTGACCTCGGATCAGGTAGGGATACCCGCTGAACCTTAAGCATATCAATAAGCGGAGGAAAAAGAAAAACAACAGGGAT TGCTCTAGTTAACGGCGAGTGAAGCAGCAATAAGCTCAAAATTTGAAATCTGGCGTCTTCGACGTCCGAGTTGTAATTT GTAGAGGATGCTTCTGAGTAACCAACCGACTTAAGTTCTTGGAACAGGACGTATAGAGGTTGAGAAATCCCCTATG CGGTCCGAAAGGTGCTCTATACGTAGCTCCTTCGACGAGTCGAGTTGTTTGGGAATGCAGCTCTAAATGGGAGGTA AATTTCTTTCTAAAGCTAAATATTTGGCCAGAGACCGATAGCCACAAGTAGAGTGATCGAAAGATGAAAAGCACTTT GGAAGAGAGTTAAAAAGCACGTGAAATTTGTTAAAGGGAAGGATGCAACACAGACTTGCTCGCGGTGTTCCGCC GGTCTTCTGACCGGTCTACTCGCCGCTTGCAGGCCAGCATCGTCTGGTCCGCTGGATAAGACTTGAGGAATGTA GCTCCCTCGGAGTTATAGCCTCTTGTGATGCAGCAGCGCCGGCGAGGTCCGCCGCTTCGGCTAGGATGCTGG CGTAATGGTCGTAATCCGCCCGTCTTTGAAACACGGACCAAGGAGTCTAACATCTATGCGAGTGTTCGGGTGTCAAA CCCCACGCGTAATGAAAGTGAACGGAGGTGAGAACCCGCAAGGTGCATCATCGACCGATCCTGATGTCTTCGGATG GATTTGAGTAAGAGCATAGCTGTTGGGACCCGAAAGATGTTGAACTATGCCTGAATAGGTTGAAGCCAGAGGAAAC TCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAAATTTGGGTATAGGGCGGAAAG

SEQ ID	MIC ID	Gene/Locus	Sequence
69	MIC-50989	ITS	TCTTGGTCATTTAGAGGAAGTAAAGTCGTAAACAAGGTCTCCGTTGGTGAACACGCGGAGGGATCATTTACAGAGTT TAACGACTCCCAACCACTGTGAACATACCCGTACCGTTGCCTCGCGGGCGGCCCCAGGGCGGGCCGACGCTC CCCAGCGAGGCGCCCGCAGGTGCAAAAACATACTATATTAGTGGCATCTCTGAGTAACTTCCAAACAAT CAAACTTTCAACAACGGATCTCTTGGTTCTGGCATCGATGAAGAACGACGAAATCGGATAAGTAATGTGAATT GCAGAAATTCAGTGAATCATCGAATCTTTGAACGCACATTCGCCCCCGCCAGCATTTCTGGCGGCATGCCTGTCCGAG CGTCATTTCAACCCCTCAAGCCCTGCTTGGTGTGGGGCACTACGCGGAGCGTAGGCCCTCAAAATCAGTGGCGGA CCCCGTGGAGGTCGGGGCTAGTAACACATCTGCCCCGAGGTCCCAGCGTCCCCTGCCGTTAAACCCCAAAAT TACAGAAAGGTTGACCTCGGATCAGGTAGGAATACCCGCTGAACCTTAAGCATATCAATAAGCGGAGGAAAAAGAAACC AACAGGGATTGCCCTAGTAACGGCGAGTGAAGCGGCAACAGCTCGAATTTGAAATCTGGCCTCGGCCCGAGTTGTA ATCTGTAGAGGATGCTTTTGGCGCGGTGCCCTTCGAGTGCCTTGGAAACGGACGCCACAGAGGTGAGAGCCCCGT ATGGTCGGACACCAAGCCTGTGTAAAGCTCCTTCGACGAGTCGGGTAGCTTGGAAATGCTGCTCTAAGTGGAGGT AAACTTTCTTCTAAAGCTAAATACTGGCCAGAGACCGCATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAAGCACTT TGAAAAGAGGGTCAAATAGTACGTGAAATTTGTTGAAAGGGAAGCGCTCATGACCAGACTTGCGCCGGCTGATCAT CCAGTGGTCTCCACTGGTGCACTCTGCCCCGGCTCAGGCCAGCGTCGGCTGTACGGGGGACAAAAAGCACTGGGAA AGTAGCTCTCTCCGGGGAGTTATAGCCCTATGCAGAAATACCCCGCGGGCCGAGGTCCGCGCTCTGCAAGGA CGCTGGCGTAATGGTCATCAGCGACCCGTCCTTGAAACACGGACCAAGGAGTCGAGGTTTTGCGCGAGTGTTCGGGT GCAAAAGCCCCAGCGCGTAATTAAGTGAACGTAGGTGAGAGCTTCGGCGCATCATCGACCGATCCTGATGTATTCTG GATGGATTTGAGTAGGAGCGTAAAGCCTCGGACCCGAAAGATGGTGAACATATGCCTGTATAGGGTGAAGCCAGAGG AAACTCTGGTGGAGGCTCGCAGCGTTCTGACGTGCAAAATCGATCGTCAAAATATGGCATGGGGCGAAAAAG
70	MIC-85555	ITS	TCTTGGTCATTTAGAGGAAGTAAAGTCGTAAACAAGGTTTTCCGTAGGTGAACCTCGGGAAGGATCATTTACCTAGAG TTTTGTGGACTTCGGTCTGTACCTCTTACCCATGTCTTTTGTAGTACCTTCGTTCCTCGCGGGTCCGCCCGCCGG TTGGACAAACATTCAAACCCCTTGCAGTTGCAATCAGCGTCTGAAAAAATTAATAGTTACAACCTTCAACAACGGA TCTCTTGGTTCTGGCATCGATGAAGAACGACGCGAAATGCGATAAGTAGTGTGAATTCGAGAATTCAGTGAATCAT CGAATCTTTGAACGCACATTCGCCCTTGGTATTCATGGGCATGCCTGTCGAGCGTCATTTGTACCTTCAAG CTCTGCTTGGTGGTGTCTCGCTCCGCGCGCAGACTCGCTTAAAAACAATGGCAGCCCGCGTATTG ATTTCCGAGCGCAGTACATCTCGCGCTTTCACCTATAACGACGACGTCCAAAAGTACATTTTACACTCTTGACC TCGGATCAGGTAGGGATACCCGCTGAACCTTAAGCATATCAATAAGCGGAGGAAAAAGAACCAACAGGGATTGCCCT AGTAACGGCGAGTGAAGCGGCAACAGCTCAAATTTGAAATCTGGCGTCTTTGGCGTCCGAGTTGTAATTTGCAGAG GGCGCTTTGGCATTTGGCAGCGGTCCAAGTTCTTGGAACAGGACGTCACAGAGGTTGAGAATCCCCTACGTGGTCG CTAGCCTTTACCGTGTAAAGCCCTTCGACGAGTCGAGTTGTTTGGAAATGCAGCTCTAAATGGGAGGTAAATTTTC TTCTAAAGCTAAATACTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAAGATGAAAAGCACTTTGGAAG AGAGTTAAAAGCACGTGAAATTTGTTGAAAGGGAAGCGCTTCGAGCCAGACTTGCCTGTAGTTGCTCATCCGGGTT TCTACCCGGTGCACTTTCTACGGGCAGGCCAGCATCAGTTTGGCGGTTGGATAAAGTCTCTGTCTACCTC CCTTCGGGGAGATCTTATAGGGGAGACGACATGCAACAGCCTGGACTGAGTCCGCGCATCTGCTAGGATGCTGG CGTAATGGCTGAAGCGGCCCTTTGAAACACGGACCAAGGATCTAACATCTATCGAGTGTTCGGGTGTCAAG CCCGAGCGGTAATGAAAGTGAACGGAGGTGGAAACCTTTCCGGGTGCACCATCGACCGATCCTGATGTCTTCGGA TGGATTTGAGTAAGAGCATAGCTGTTGGACCCGAAAGATGGTGAACATATGCTTGAATAGGTGAAGCCAGAGGAA ACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCAAAATCGATCGTCAAATTTGGGCATAGGGGCGAAAGACTAAT

SEQ ID	MIC ID	Gene/Locus	Sequence
71	MIC-68178	Unique genomic region	CTCCTCCTCCTCCTGATCGAACTCGCCCCCCCACCAACTCCACGAGCCCCAGCCGCCCCAAAAACCTCCCCA TCCCCGCTAGCCGCCCCCATGGCCCGGACAAGCAGCANNNNNNNNNNNGTCCGGGATGGCCTTAGGTTGCTCT CGAGCTGCTCCAGCCCGACAGGACATGCAGCAGCTCCCTGCGCAGCGAGTCCGGCCCGGCATCGGCCCCCGCAG CGGCTGGTCGGCGATGTATGTCTTGAGCGGGAGCGCGCGCAGGATGAGGTAGGGCGCCGCTGCCCTGGGCGAGT TTGATGTGCGAGTGCAGGGCGAGGGCGAGGTACTGGCGGGGCGTGGGCCACGAGGGCGAAGAGGGAGGCCGTGC AGGTGTAGCTCATCTTTGTGCGCAAGGTGGCGGGGAGGACGGCCGCTTGCCCGGTGCGGACGGAGAGGTTGGC GAGGGCGGAGGTGGTGATGGGAGGGAGTTCTCCTTGTACGGGAGTGTGGATGAGGGAGGTGGTGAAGAGGTTGCGG GCGTATGTCTGCGGAGGGTGTGCGGGAGGGAGGGGAGTTTAGAGGGCCGCTGATCAAGTTCGCGTAGCTCTGTGA

Example 2: Isolation and identification of endophytes using marker gene sequences.

[00350] The fungal endophytes of the present invention can be identified by the sequence of one or more of the following loci: long subunit rRNA (LSU), small subunit rRNA (SSU), largest subunit of RNA polymerase II (RPB1), second largest subunit of RNA polymerase II (RPB2), beta-tubulin, actin, phosphoglycerate kinase (PGK). PCR amplification of the largest subunit of RNA polymerase I (RPB1) using primer sequences RPB1-Af (SEQ ID NO: 21) and RPB1-Cr (SEQ ID NO: 22) is described in Cendejas-Bueno E, Kolečka A, Alastruey-Izquierdo A, et al. Reclassification of the *Candida haemulonii* Complex as *Candida haemulonii* (*C. haemulonii* Group I), *C. duobushaemulonii* sp. nov. (*C. haemulonii* Group II), and *C. haemulonii* var. *vulnera* var. nov.: Three Multiresistant Human Pathogenic Yeasts. *Journal of Clinical Microbiology*. 2012;50(11):3641-3651. PCR amplification of second largest subunit of RNA polymerase II (RPB2) using primer sequences fRPB2-5F (SEQ ID NO: 26) and fRPB2-7.1R (SEQ ID NO: 25) is described in Riess K, Oberwinkler F, Bauer R, Garnica S. High genetic diversity at the regional scale and possible speciation in *Sebacina epigaea* and *S. incrustans*. *BMC Evolutionary Biology*. 2013;13:102. doi:10.1186/1471-2148-13-102. PCR amplification of beta-tubulin using primer sequences Btub2Fd (SEQ ID NO: 30) and Btub4Rd (SEQ ID NO: 31) is described in Aveskamp et al. (2009) DNA phylogeny reveals polyphyly of *Phoma* section *Peyronellaea* and multiple taxonomic novelties *Mycologia*, 101(3):363–382. PCR amplification of the LSU using primer sequences LR0R (SEQ ID NO: 23) and LR5 (SEQ ID NO: 24) is described in Stielow JB, Lévesque CA, Seifert KA, et al. One fungus, which genes? Development and assessment of universal primers for potential secondary fungal DNA barcodes. *Persoonia : Molecular Phylogeny and Evolution of Fungi*. 2015;35:242-263. doi:10.3767/003158515X689135. PCR amplification of the SSU using primer sequences SR1R (SEQ ID NO: 29) and NS4 (SEQ ID NO: 28) is described in Zhu et al. (2016) *Helminthosporium velutinum* and *H. aquaticum* sp. nov. from aquatic habitats in Yunnan Province, China. *Phytotaxa* 253 (3): 179–190. PCR amplification of Actin using primer sequences ACT512f (SEQ ID NO: 17) and ACT783r (SEQ ID NO: 18) is described in Carbone, I. & Kohn, L.M. (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia*, 91(3):552-556. PCR amplification of largest subunit of RNA polymerase II (RPB1) using primer sequences RPB1-Af (SEQ ID NO: 21) and RPB1-Cr (SEQ ID NO: 22) is described in Urbina H, Blackwell M (2012) Multilocus Phylogenetic Study of the *Scheffersomyces* Yeast Clade and Characterization of the N-Terminal Region of Xylose Reductase Gene. *PLoS ONE* 7(6): e39128.

[00351] MIC-68390 can be identified by one or more of the following exemplary sequences: RPB2 sequence (SEQ ID NO: 61). MIC-68178 can be identified by one or more of the following exemplary sequences: beta-tubulin sequence (SEQ ID NO: 64). MIC-07010 can be identified by one or more of the following exemplary sequences: phosphoglycerate kinase sequence (SEQ ID NO: 66). MIC-96038 can be identified by one or more of the following exemplary sequences: actin sequence (SEQ ID NO: 47), beta-tubulin sequence (SEQ ID NO: 48), RPB2 sequence (SEQ ID NO: 49), RPB1 sequence (SEQ ID NO: 50). MIC-33414 can be identified by one or more of the following exemplary sequences: its Actin sequence (SEQ ID NO: 53), RPB1 sequence (SEQ ID NO: 55), beta-tubulin sequence (SEQ ID NO: 58), LSU sequence (SEQ ID NO: 54), SSU sequence (SEQ ID NO: 56), SSU sequence (SEQ ID NO: 57). MIC-31593 can be identified by one or more of the following exemplary sequences: its RPB2 sequence (SEQ ID NO: 43), beta-tubulin sequence (SEQ ID NO: 44). Exemplary LSU and SSU sequences of MIC-19994 are SEQ ID NO: 39 and SEQ ID NO: 40, respectively.

Example 3: Isolation and Identification of bacterial and fungal endophytes using ITS sequence

[00352] Classification of the fungal strain using ITS sequences was done by the following methodology.

[00353] Total genomic DNA was extracted from individual fungal isolates, using the DNeasy Plant Mini Kit (Qiagen, Germantown, MD). Polymerase Chain Reaction (PCR) was used to amplify a genomic region including the nuclear ribosomal internal transcribed spacers (ITS) using a primer pair ITS_1 (5'-CTTGGTCATTTAGAGGAAGTAA-3') (SEQ ID NO: 5) and LR5 (5'-TCCTGAGGGAACTTCG-3') (SEQ ID NO: 8). Each 25 microliter-reaction mixture included 22.5 microliters of Invitrogen Platinum Taq supermix, 0.5 microliter of each primer (10 uM), and 1.5 microliters of DNA template (~2-4ng). Cycling reactions were run with MJ Research PTC thermocyclers and consisted of 94°C for 5 min, 35 cycles of 94°C for 30 s, 54°C for 30 s, and 72°C for 1 min, and 72°C for 10 min. Sanger sequencing of was performed at Genewiz (South Plainfield, NJ) using primers: ITS_1 (5'-CTTGGTCATTTAGAGGAAGTAA-3') (SEQ ID NO: 5), ITS_2 (5'-GCTGCGTTCTTCATCGATGC-3') (SEQ ID NO: 6), ITS_3 (5'-GCATCGATGAAGAACGCAGC-3') (SEQ ID NO: 7), and LR5 (5'-TCCTGAGGGAACTTCG-3') (SEQ ID NO: 8). Sequencing primers were chosen so that overlapping regions were sequenced. Raw chromatograms were converted to sequences, and corresponding quality scores were assigned using TraceTuner v3.0.6beta (US 6,681,186).

These sequences were quality filtered, aligned and a consensus sequence generated using Geneious v 8.1.8 (Biomatters Limited, Auckland NZ).

[00354] Taxonomic classifications were assigned to the sequences using the highest probability of assignment based on the results of industry standard taxonomic classification tools: LCA (runs USEARCH (Edgar, R. C. (2010) *Bioinformatics*. 26(19):2460-2461) with option search_global, then for all best match hits, returns lowest taxonomic rank shared by all best hits for a query), SPINGO (Allard et al. (2015) *BMC Bioinformatics*. 16: 324), and UTAX (Edgar, R.C., 2016), using the WARCUP Fungal ITS trainset 1 (Deshpande et al. (2016) *Mycologia* 108(1):1-5) and UNITE (Koljalg et al. (2013) *Molecular Ecology*, 22: 5271–5277). The classifier and database combinations listed in Table 5 were used to assign taxonomy to fungal sequences.

Table 5: The classifier and database combinations used to classify ITS sequences

<i>Classifier</i>	<i>Database</i>
LCA	UNITE, Fungal ITS trainset 07/04/2014
RDP	UNITE, Fungal ITS trainset 07/04/2014
	WARCUP, Fungal ITS trainset 1
SPINGO	UNITE, Fungal ITS trainset 07/04/2014
UTAX	UNITE, Fungal ITS trainset 07/04/2014
	WARCUP, Fungal ITS trainset 1

Table 6. Taxonomic classification of endophytes of the present invention

MIC ID	Phylum	Class	Order	Family	Genus	Species
MIC-68390	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Pleosporales</i>	<i>Pleosporaceae</i>	<i>Exserohilum</i>	<i>rostrata</i>
MIC-68178	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Pleosporales</i>	<i>Pleosporaceae</i>	<i>Epicoccum</i>	<i>nigrum</i>
MIC-07010	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Pleosporales</i>	<i>Pleosporaceae</i>	<i>Curvularia</i>	<i>protuberata</i>
MIC-31593	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Pleosporales</i>	<i>Pleosporaceae</i>	<i>Curvularia</i>	<i>spicifera</i>
MIC-48747	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Capnodiales</i>	<i>Cladosporiaceae</i>	<i>Cladosporium</i>	<i>cladosporioides</i>
MIC-96038	<i>Ascomycota</i>	<i>Sordariomycetes</i>	<i>Hypocreales</i>	<i>Hypocreaceae</i>	<i>Acremonium</i>	<i>alternatum</i>
MIC-50414	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Capnodiales</i>	<i>Cladosporiaceae</i>	<i>Cladosporium</i>	<i>oxysporum</i>
MIC-33414	<i>Ascomycota</i>	<i>Sordariomycetes</i>	<i>Sordariales</i>	<i>Chaetomiaceae</i>	<i>Chaetomium</i>	<i>globosum</i>
MIC-85555	<i>Ascomycota</i>	<i>Dothidiomycetes</i>	<i>Pleosporales</i>	<i>Pleosporaceae</i>	<i>Epicoccum</i>	<i>nigrum</i>
MIC-50989	<i>Ascomycota</i>	<i>Sordariomycetes</i>	<i>Sordariales</i>	<i>Cephalothecaceae</i>	<i>Paecilomyces</i>	<i>inflatus</i>
MIC-19994	<i>Ascomycota</i>	<i>Sordariomycetes</i>	<i>Coniochaetales</i>	<i>Coniochaetaceae</i>	<i>Coniochaeta</i>	<i>prunicola</i>

[00355] MIC-68390 can be identified by the sequence of its ITS sequence (SEQ ID NO: 60). MIC-68178 can be identified by the sequence of its ITS sequence (SEQ ID NO: 63). MIC-07010 can be identified by the sequence of its ITS sequence (SEQ ID NO: 65). MIC-31593 can be identified by the sequence of its ITS sequence (SEQ ID NO: 42). MIC-48747 can be

identified by the sequence of its ITS sequence (SEQ ID NO: 67). MIC-96038 can be identified by the sequence of its ITS sequence (SEQ ID NO: 46). MIC-50414 can be identified by the sequence of its ITS sequence (SEQ ID NO: 68). MIC-33414 can be identified by the sequence of its ITS sequence (SEQ ID NO: 52). MIC-85555 can be identified by the sequence of its ITS sequence (SEQ ID NO: 70). MIC-50989 can be identified by the sequence of its ITS sequence (SEQ ID NO: 69).

Example 4. Assessment of Improved Plant Characteristics, Seedling Vigor

Assay of soy seedling vigor

[00356] *Seed preparation:* The lot quality of soybean seeds was first assessed by testing germination of 100 seeds. Seeds were placed, 8 seeds per petri dish, on filter paper in petri dishes, 12 mL of water was added to each plate and plates are incubated for 3 days at 24°C. The percent germination was greater than 95%. One thousand soybean seeds were then surface sterilized by co-incubation with chlorine gas in a 20 x 30 cm container placed in a chemical fume hood for 16 hours. Percent germination of 50 seeds, per sterilization batch, was tested as above and confirmed to be greater than 95%.

[00357] *Preparation and heterologous disposition of endophytes:* Spore solutions were made by rinsing and scraping spores from agar slants which have been growing for about 1 month. Rinsing was done with 0.05% Silwet. Solutions were passed through Miracloth to filter out mycelia. Spores per ml were counted under a microscope using a hemocytometer. The stock suspension was then diluted into 10^6 spores/ml utilizing water. 3 μ l of spore suspension was used per soy seed ($\sim 10^3$ CFUs/seed is obtained). Control treatments were prepared by adding equivalent volumes of sterile water to seeds.

[00358] *Assay of seedling vigor:* Two rolled pieces of germination paper were placed in a sterile glass jar with 50 mL sterile water, then removed when completely saturated. Then the papers were separated and inoculated seeds were placed at approximately 1 cm intervals along the length of one sheet of moistened germination paper, at least 2.5 cm from the top of the paper and 3.8 cm from the edge of the paper. The second sheet of was placed on top of the soy seeds and the layered papers and seeds were loosely rolled into a tube. Each tube was secured with a rubber band around the middle and placed in a single sterile glass jar and covered loosely with a lid. For each treatment, three jars with 15 seeds per jar were prepared. The position of jars with the growth chamber was randomized. Jars were incubated at 60% relative humidity, and 22°C day, 18°C night with 12 hours light and 12 hours dark for 4 days and then the lids were removed and the jars incubated for an additional 7 days. Then the

germinated soy seedlings were weighed and photographed and root length and root surface area scored as follows.

[00359] Dirt, excess water, seed coats and other debris was removed from seedlings to allow accurate scanning of the roots. Individual seedlings were laid out on clear plastic trays and trays are arranged on an Epson Expression 11000XL scanner (Epson America, Inc., Long Beach CA). Roots were manually arranged to reduce the amount of overlap. For root measurements, shoots were removed if the shape of the shoot causes it to overlap the roots.

[00360] The WinRHIZO software version Arabidopsis Pro2016a (Regents Instruments, Quebec Canada) was used with the following acquisition settings: greyscale 4000 dpi image, speed priority, overlapping (1 object), Root Morphology: Precision (standard), Crossing Detection (normal). The scanning area was set to the maximum scanner area. When the scan was completed, the root area was selected and root length and root surface area were measured.

[00361] Statistical analysis was performed using R (R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. R-project.org/). Results are summarized in Tables 7 and 8.

Table 7. Root traits of endophyte treated and untreated soybean seedlings.

MIC ID	Average Root Length	Standard Deviation, Root Length	Average Root Area	Standard Deviation, Root Area	Root Length % difference NT	Root Area % difference NT
Untreated control (NT)	79.4	32.5	8.9	4.1	0.0	0.0
MIC-68178	81.2	34.6	9.8	3.4	2.2	9.6
MIC-07010	94.1	29.5	10.5	3.2	18.5	17.7
MIC-31593	85.6	29.7	10.1	3.4	7.7	12.9
MIC-48747	96.7	26.0	11.7	3.0	21.7	31.4
MIC-96038	88.5	33.1	10.7	4.0	11.4	19.8

Table 8. Percent increase in soybean seedling root length of endophyte treatment relative to untreated controls.

	% Gain over Control Vigor
Treatment	Soy
MIC-33414	27.8

Assay of corn seedling vigor

[00362] *Seed preparation:* The lot quality of corn seeds is first evaluated for germination by transfer of 100 seeds and with 3.5 mL of water to a filter paper lined petri dish. Seeds are

incubated for 3 days at 24°C, and to ensure that percent germination is greater than 95%. One thousand corn seeds are then surface sterilized by co-incubation with chlorine gas in a 20 x 30 cm container in a chemical fume hood for 12 hours. Percent germination of 50 seeds, per sterilization batch, is tested as above and confirmed to be greater than 95%.

[00363] *Optional reagent preparation:* 7.5% PEG 6000 (Calbiochem, San Diego, CA) is prepared by adding 75 g of PEG to 1000 mL of water, then stirred on a warm hot plate until the PEG is fully dissolved. The solution is then autoclaved.

[00364] *Preparation and heterologous disposition of endophytes:* Spore solutions are made by rinsing and scraping spores from agar slants which have been growing for about 1 month. Rinsing is done with 0.05% Silwet. Solutions are passed through Miracloth to filter out mycelia. Spores per ml are counted under a microscope using a hemocytometer. The stock suspension is then diluted into 10^6 spores/ml utilizing water. 3 μ l of spore suspension is used per corn seed ($\sim 10^3$ CFUs/seed is obtained). Control treatments are prepared by adding equivalent volumes of sterile water to seeds.

[00365] *Assay of seedling vigor:* Either 25 ml of sterile water (or optionally, 25 ml of PEG solution as prepared above) is added to each CygTM germination pouch (Mega International, Newport, MN) and place into pouch rack (Mega International, Newport, MN). Sterile forceps are used to place corn seeds prepared as above into every other perforation in the germination pouch. Seeds are fitted snugly into each perforation to ensure they did not shift when moving the pouches. Before and in between treatments forceps are sterilized using ethanol and flame and workspace wiped down with 70% ethanol. For each treatment, three pouches with 15 seeds per pouch are prepared. The germination racks with germination pouches are placed into plastic tubs, and covered with perforated plastic wrap to prevent drying. Tubs are incubated at 60% relative humidity, and 22°C day, 18°C night with 12 hours light and 12 hours dark for 6 days to allow for germination and root length growth. Placement of pouches within racks and racks/tubs within the growth chamber is randomized to minimize positional effect. At the end of 6 days the corn seeds are scored manually for germination, root and shoot length.

[00366] Statistical analysis is performed using R (R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. R-project.org/).

Assay of wheat seedling vigor

[00367] *Seed preparation:* The lot of wheat seeds was first evaluated for germination by transfer of 100 seeds and with 8 mL of water to a filter paper lined petri dish. Seeds were

incubated for 3 days at 24°C, and percent germination was greater than 95%. Wheat seeds were then surface sterilized by co-incubation with chlorine gas in a 20 x 30 cm container in a chemical fume hood for 12 hours. Percent germination of 50 seeds, per sterilization batch, was tested as above and confirmed to be greater than 95%.

[00368] *Reagent preparation:* 7.5% polyethylene glycol (PEG) was prepared by adding 75 g of PEG to 1000 mL of water, then stirring on a warm hot plate until the PEG is fully dissolved. The solution was then autoclaved.

[00369] *Preparation and heterologous disposition of endophytes:* Spore solutions were made by rinsing and scraping spores from agar slants which had been growing for about 1 month. Rinsing was done with 0.05% Silwet. Solutions were passed through Miracloth to filter out mycelia. Spores per ml were counted under a microscope using a hemocytometer. The stock suspension was then diluted into 10^6 spores/ml utilizing water. 3 μ l of spore suspension was used per wheat seed ($\sim 10^3$ CFUs/seed was obtained). Seeds and spores were combined a 50 mL falcon tube and gently shaken for 5-10 seconds until thoroughly coated. Control treatments were prepared by adding equivalent volumes of sterile water to seeds.

[00370] *Assay of seedling vigor:* Petri dishes were prepared by adding four sheets of sterile heavy weight seed germination paper, then adding 50 mL of PEG solution as prepared above to each plate then allowing the liquid to thoroughly soak into all sheets. The sheets were positioned and then creased so that the back of the plate and one side wall were covered, two sheets were then removed and placed on a sterile surface. Along the edge of the plate across from the covered side wall 15 inoculated wheat seeds were placed evenly at least one inch from the top of the plate and half an inch from the sides. Seeds were placed smooth side up and with the pointed end of the seed pointing toward the side wall of the plate covered by germination paper. The seeds were then covered by the two reserved sheets, and the moist paper layers smoothed together to remove air bubbles and secure the seeds, and then the lid was replaced. For each treatment, at least three plates with 15 seeds per plate were prepared. The plates were then randomly distributed into stacks of 8-12 plates and a plate without seeds was placed on the top. The stacks were incubated at 60% relative humidity, and 22°C day, 18°C night with 12 hours light and 12 hours dark for 24 hours, then each plate was turned to a semi-vertical position with the side wall covered by paper at the bottom. The plates were incubated for an additional 5 days, then wheat seeds scored manually for germination, root and shoot length.

[00371] Statistical analysis was performed using R (R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. R-project.org/). Results are summarized in Table 9.

Table 9. Percent increase in wheat seedling root length of endophyte treatment relative to untreated controls.

	% Gain over Control Vigor
SYM	Wheat
MIC-33414	6.5

Example 5: Culture preparations and heterologous disposition of endophytes for greenhouse experiments

[00372] Strains may be cultured by the methods described herein and methods well known in the art.

Preparation of fungal biomass

Method 1

[00373] Biomass for MIC-19994 and MIC-96038 was produced by growing for two weeks in liquid medium (PDB). The resulting biomass was homogenized by sonication (50% amplitude for 30 seconds) or in a FastPrep-24 (MP Biomedicals, Santa Ana, CA, USA) set to 4.5m/s for 30 seconds.

Method 2

Spore solutions were made by rinsing and scraping spores from agar slants which have been growing for about 1 month. Rinsing was done with 0.05% Silwet. Solutions were passed through Miracloth to filter out mycelia. Spores per ml were counted under a microscope using a hemocytometer. The stock suspension was then diluted into 10^6 spores/ml utilizing water. 3 μ l of spore suspension was used per seed ($\sim 10^3$ CFUs/seed is obtained). Control treatments were prepared by adding equivalent volumes of sterile water to seeds.

Method 3

[00374] Preparation: Molasses broth was prepared by dissolving 30 g molasses and 5 g yeast extract per liter deionized water in an autoclavable container and autoclaving (15 psi, 121°C) for 45min. Potato dextrose agar (PDA) plates were prepared by dissolving 39.0g PDA powder per liter deionized water in an autoclavable container and autoclaving (15 psi, 121°C) for 45 min. The agar was allowed to cool to 50-60°C, before pouring into sterile petri plates (30mL per 90mm plate).

[00375] Liquid biomass: All equipment and consumables were thoroughly sterilized and procedures performed in a biosafety cabinet. The inoculant is prepared by placing 1 plug

from a cryopreserved stock on a fresh PDA plate, sealing the plate with Parafilm® and incubating at room temperature in the dark for 5-10 days. Then ~5x5 mm plugs were cut from the PDA plates and 10-12 plugs were transferred into flasks containing the sterile molasses broth, covered, secured in a shaker and incubated for at least 10 days with shaking at ~130 rpm. Then the culture was placed in a blender for 5 seconds and 1 mL of the blended was centrifuged and the supernatant was discarded and the pellet resuspended in 0.5 mL 1x Phosphate Buffered Saline (PBS) to generate inoculum.

[00376] Dry biomass: All equipment and consumables were thoroughly sterilized and procedures performed in a biosafety cabinet. The inoculant is prepared by placing 1 plug from a cryopreserved stock on a fresh PDA plate, sealing the plate with Parafilm® and incubating at room temperature in the dark for 5-10 days. Then ~5x5 mm plugs were cut from the PDA plates and 10-12 plugs were transferred into flasks containing the sterile molasses broth, covered, secured in a shaker and incubated for at least 10 days with shaking at ~130 rpm. In sterile conditions, the liquid culture was carefully decanted using 150 mm sterile filter paper on a sterilized Buchner funnel over a sterile flask. Once all liquid had passed through the funnel, the pellet was rinsed with sterile water until the filtrate ran clear. When dry, the pellet was transferred to a drying cabinet and dried until brittle. The pellet was then ground into a fine powder, and sample used to generate CFU counts.

Seed inoculation

[00377] Unless otherwise specified, inoculum was added to seeds to reach a targeted dose of 10^4 CFU. Where low, medium and high doses are indicated, and in not specified otherwise, a high dose is 10^5 CFU/seed, a medium dose is 10^4 CFU/seed, and a low dose 10^3 CFU/seed. The seeds were agitated to disperse the inoculum evenly on the seeds.

Formulation control treatments were prepared using equivalent volumes of PBS. The seeds were allowed to dry for approximately 2 mins, then 2 fluid ounces per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds.

Example 6. Assessment of Improved Plant Characteristics: Greenhouse

Rice

[00378] Rice seeds of variety Rex were treated with commercial fungicidal and insecticidal treatment CruiserMaxx® Rice (Syngenta, Basel, Switzerland). Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 28, untreated seeds (lacking formulation and endophyte) were also planted. The loaded dose of each microbe is shown in Table 42. The endophyte treatment

MIC-68178/MIC-33414 was co-inoculated with MIC-68178 and MIC-33414. In the co-inoculation treatment, MIC-68178 made up 17.45% of the loaded dose, and MIC-33414 made up 82.55% of the loaded dose.

Table 10. Loaded doses (CFU per seed) of soybean seeds treated with endophytes.

	Loaded doses, CFU per seed
MIC-68178	1.64E+03
MIC-33414	2.39E+03
MIC-68178/MIC-33414	1.44E+03

[00379] Each pot was filled with Cahaba/Wickham type, fine sandy loam soil, and two seeds were sown evenly spaced in each pot. Ten pots were planted per treatment/control. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Plants were watered twice daily. Upon emergence of true leaves, plants were fertilized weekly at 250 PPM N using Peter's Peat-Lite 20-1020 water-soluble fertilizer.

[00380] At day 7, pots were thinned to 1 seedling/pot. Root tissue was harvested from the experiment six weeks post-planting, and the soil was washed from the roots. The tissues from each plant was placed in an unlined paper bag. The tissue was dried in an oven set to 85°C for 3 days. Once completely dried, the root biomass of individual plants were weighed and recorded.

[00381] MIC-68178 demonstrated an increase of 51.5% in dry root weight compared to the untreated control, with over 80% confidence by Bayesian analysis.

Table 11. Percent difference in root dry weight in rice variety Rex in greenhouse conditions

	Rice variety Rex
	% difference Untreated control, root dry weight
Untreated control	0
Formulation control	52.0
MIC-68178	51.5

Soy

[00382] Sandy loam was mixed in a ratio of 60% loam and 40% mortar sand. Prior to mixing, both planting media were sifted through a 3/8" square steel mesh screen to remove larger particles and debris. Soy seeds of variety Stine 33E22 were treated with commercial fungicidal and insecticidal treatment CruiserMaxx® Soy (Syngenta, Basel, Switzerland). Seeds were heterologously disposed with the endophyte formulations and formulation control

(lacking any endophyte) as described in Example 28, untreated seeds (lacking formulation and endophyte) were also planted.

[00383] Each pot was filled with 1000 mL of soil, watered with 225 mL of water and one seed sown per pot. Ten pots were planted per treatment/control. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Environmental conditions were set at 12h photoperiod, at 23/20°C temperature for day/night period and light intensity was set at 550 $\mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds were watered to maintain approximately 75% soil capacity.

[00384] Root tissue was harvested from the experiment three weeks post-planting, and the soil was washed from the roots. The tissues from each plant was placed in an unlined paper bag. The tissue was dried in an oven set to 85°C for 3 days. Once completely dried, the root biomass of individual plants were weighed and recorded.

[00385] Roots treated with the endophyte MIC-68178 demonstrated an increase of 16.9% in dry root weight compared to the untreated control, but with less than 80% confidence by Bayesian analysis. Roots treated with the endophyte MIC-33414 demonstrated a decrease of 23.5% in dry root weight compared to the untreated control, but with less than 80% confidence by Bayesian analysis. However, the endophyte formulation comprising both MIC-68178 and MIC-33414 demonstrated an increase in dry root weight of 36.8% compared to the untreated control, with over 80% confidence by Bayesian analysis. A photo taken prior to drying of exemplary untreated control and MIC-68178 and MIC-33414 treated soybean roots is shown in Figure 3.

Table 12. Percent difference in root dry weight in soybean variety Stine 33E22 in greenhouse conditions.

	Soybean variety Stine 33E22
	% difference Untreated control, root dry weight
Untreated control	0
Formulation control	7.4
MIC-68178	16.9
MIC-68178/MIC-33414	36.8
MIC-33414	-23.5

Example 7: Cultivation of Endophyte-Treated Plants in Greenhouse Experiment 1

Wheat

[00386] A sandy loam and a commercial potting soil (Farfard®, Agawam, MA) were used in this experiment. Sandy loam was mixed in a ratio of 60% loam and 40% mortar sand (Northeast Nursery, Peabody, MA). Prior to mixing, both planting media were sifted through

a 3/8" square steel mesh screen to remove larger particles and debris. Wheat seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 5, untreated seeds (lacking formulation and endophyte) were also planted. Endophyte treatments were applied to seed in three target doses: high (10^5 CFU/seed), medium (10^4 CFU/seed), low (10^3 CFU/seed). Each pot was filled with 600 mL of its respective soil, watered with 200 mL of water and then, nine seeds were sown evenly spaced in each pot (in a 3x3 pattern). Soil was then overlaid atop the seeds (estimated average planting depth of 1 inch) and an additional 110 mL of water was added to moisten the overlaying soil substrate. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Environmental conditions were set at 12h photoperiod, at 22/18°C temperature for day/night period and light intensity was set at $650 \mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds were watered to maintain approximately 80% soil capacity.

[00387] Wheat seedlings emergence was recorded on days 4, 5, and 7 after planting, with days 4 and 5 representing early emergence percentage and day 7 representing final emergence percentage. At day 7, all pots were thinned to 3 seedlings/pot. Above ground tissue was harvested from the experiment three weeks post-planting. The tissues from individual replicate treatments (pots) were pooled and placed in an unlined paper bag. All tissues were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot biomass of individual treatment replicates (pots) was weighed and recorded

[00388] MIC-96038, MIC-96038/MIC-19994 co-cultured, and MIC-19994 treatments all increased dry plant shoot biomass at the medium dose. The MIC-19994 treatment resulted in a 6% ($p < 0.05$) increase in dry shoot biomass at 10^4 CFU/seed.

Table 13. Dry shoot biomass of endophyte treated wheat

	Avg dry shoot biomass (g)	Standard deviation	Standard error of the mean	% change over untreated control	% change over formulation control
Untreated control	0.949	0.0852	0.011	0	0.21
Formulation control	0.947	0.072	0.0093	-0.2	0.00
MIC-19994, 10^3	0.97	0.0684	0.0153	2.2	2.43
MIC-19994, 10^4	1.009	0.0539	0.0121	6.32	6.55

MIC-19994 , 10 ⁵	0.941	0.0865	0.0193	-0.79	-0.63
MIC-96038 pure, 10 ³	0.962	0.0631	0.0141	1.41	1.58
MIC-96038 pure, 10 ⁴	0.992	0.0831	0.0186	4.57	4.75
MIC-96038 pure, 10 ⁵	0.934	0.0891	0.0199	-1.55	-1.37
MIC-96038/ MIC-19994 , 10 ³	0.943	0.0644	0.0144	-0.65	-0.42
MIC-96038/ MIC-19994 , 10 ⁴	0.989	0.0775	0.0173	4.27	4.44
MIC-96038/ MIC-19994 , 10 ⁵	0.898	0.089	0.0199	-5.38	-5.17

Soy

[00389] A sandy loam growth substrate was mixed in the greenhouse and consisting of 60% loam and 40% mortar sand (Northeast Nursery, Peabody, MA). Prior to mixing, loam was sifted through a 3/8" square steel mesh screen to remove larger particles and debris. Soybean seeds were treated with commercial fungicidal and insecticidal treatment CruiserMaxx® Vibrance (Syngenta, Basel, Switzerland) according to the manufacturer's instructions. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 11, untreated seeds (lacking formulation and endophyte) were also planted. Endophyte treatments were applied to seed in three target doses: high (10⁵ CFU/seed), medium (10⁴ CFU/seed), low (10³ CFU/seed). Each pot was filled with 600 mL of its respective soil, watered with 200 mL of water and then, nine seeds were sown evenly spaced in each pot (in a 3x3 pattern). Soil was then overlaid atop the seeds (estimated average planting depth of 1 inch) and an additional 110 mL of water was added to moisten the overlaying soil substrate. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Environmental conditions were set at 12h photoperiod, at 22/18C temperature for day/night period and light intensity was set at 650 $\mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds grown in normal watering conditions were watered to maintain approximately 80% soil capacity and the seeds grown in drought conditions were watered to maintain approximately 40% soil capacity. Above ground tissue was harvested from the experiment three weeks post-planting. The tissues from individual replicate treatments (pots) were pooled and placed in an unlined paper bag. All tissues were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot biomass of individual treatment replicates (pots) was weighed and recorded.

Table 14. Endophyte treatments associated with increases in dry shoot biomass under normal watering conditions relative to formulation for endophyte treated soybeans

	Normal watering		Limited watering	
	Dry weight (g)	% change over formulation control	Dry weight (g)	% change over formulation control
MIC-96038	1.987	7.3%	1.145	-0.72%
MIC-96038/MIC-19994	1.945	5.07%	1.155	0.17%

	Normal watering		Limited watering	
	Dry weight (g)	% change over formulation control	Dry weight (g)	% change over formulation control
MIC-19994	1.937	4.64%	1.106	-4.11%

Rice

[00390] A sandy loam growth substrate was mixed in the greenhouse and consisting of 60% sifted loam and 40% mortar sand (Northeast Nursery, Peabody, MA). Prior to mixing, loam was sifted through a 3/8 inch. square steel mesh screen to remove larger particles and debris.

[00391] Endophyte treated seeds and control (no endophyte) seeds prepared in Example 26 were planted in a growth room experiment. For each treatment or control 20 replicates were prepared as follows. Each pot was filled with 600 mL of soil, watered with 200 mL of water and then, six seeds were sown evenly spaced in two rows (in a 2x3 pattern). Soil was then overlaid atop the seeds (estimated average planting depth of 0.5 inches) and an additional 100 mL of water was added to moisten the overlaying substrate. The experimental design called for a completely randomized pattern of each treatment within each block/replicate.

Environmental conditions were set at 12h photoperiod, at 22/18C temperature for day/night period and light intensity was set at 650 $\mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds were watered with 200 mL of water every other day. Pots were thinned down to 3 seedlings at approximately 9 days post-planting.

[00392] Emergence was recorded on days 4, 5, 7, 8 and 9 days after planting, with days 4 and 5 representing early emergence percentage and day 9 representing final emergence percentage. Above ground tissue was harvested from the experiment three weeks post-planting. The tissues from individual replicate treatments (pots) were pooled and placed in an unlined paper bag. All tissues were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot biomass of individual treatment replicates (pots) was weighed and recorded.

Table 15. Shown are mean dry biomass (g) of seeds treated with Flo Rite®

Treatment	Dose	Mean dry biomass (g)	Std Dev	Std Err of the mean	% vs untreated control	% vs formulation control
Untreated control		0.107	0.0323	0.00417	0	6.82
Formulation control		0.1002	0.0294	0.0038	-6.38	0
MIC-68178	Low	0.1053	0.0542	0.01211	-1.61	5.09
	Medium	0.1195	0.0263	0.00588	11.61	19.21
	High	0.0967	0.0229	0.00513	-9.67	-3.51
MIC-96038	Low	0.1302	0.0204	0.00456	21.65	29.94
	Medium	0.1244	0.0244	0.00545	16.19	24.1
	High	0.1298	0.0241	0.00538	21.26	29.53

Table 16. Shown are mean dry biomass (g) of seeds not treated with Flo Rite®

Treatment	Dose	Mean dry biomass (g)	Standard deviation	Standard error of the mean	% over untreated control	% over formulation control
Untreated control		0.107	0.0323	0.00417	0.0	6.8
Formulation control		0.1002	0.02945	0.0038	-6.4	0.0
MIC-68178	Low	0.0891	0.04152	0.00928	-16.7	-11.1
	Medium	0.0653	0.0156	0.00349	-39.0	-34.9
	High	0.1294	0.01668	0.00373	20.9	29.1
MIC-96038	Low	0.144	0.03537	0.00791	34.5	43.7
	Medium	0.1203	0.02521	0.00564	12.4	20.1
	High	0.1187	0.02709	0.00606	10.9	18.4

Example 8: Oil-based Formulation of Endophyte Seed Treatments

[00393] Endophyte compositions were generated comprising an oil high in erucic acid, a non-ionic surfactant, and a plantability polymer. MIC-31593 and MIC-33414 were prepared in three different formulations (A_2, B_2, C_2) and MIC-96038 was prepared in formulation A_1, described in Table 15.

Table 17. Components of endophyte compositions, volumes per 50 g of seed

Formulation	Oil	Oil volume (mL)	0.5% Triton X-100 volume (mL)	Flo Rite® (mL)
A_2	Rapeseed	0.2545	0.12727	0.049
B_2	Rapeseed	0.1273	0.2545	0.049
C_2	None	0	0.38182	0.049
A_1	Rapeseed	0.1273	0.2545	0.049
A_4	None	0	0	0.049

Example 9. Vigor Assessment in Greenhouse Experiment 2*Wheat*

[00394] Endophyte inocula as prepared in Example 5 was applied to wheat seeds which had not previously been treated with chemical insecticide or fungicide. Endophyte treated seeds were prepared at 5 target doses: 10^1 , 10^2 , 10^3 , 10^4 , 10^5 CFU/seed. Ten biological replicates each were planted for each treatment and control condition (no endophyte) in individual containers containing commercial potting media.

[00395] Emergence was recorded at 7 days post planting, and plant height (cm) scored at 7, 14, 21 and 28 days post planting. Above and below ground tissue was harvested from the experiment four weeks post-planting. The fresh weight of roots and shoots were recorded, then the samples were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot and root biomass was weighed and recorded.

Table 18. Germination percentage, plant height and weight of endophyte treated wheat formulation controls. DAP = Days after planting

		Formulation control	MIC-96038/MIC-19994, 10 ¹ CFU/seed	MIC-96038/MIC-19994, 10 ² CFU/seed	MIC-96038/MIC-19994, 10 ³ CFU/seed	MIC-96038/MIC-19994, 10 ⁴ CFU/seed	MIC-96038/MIC-19994, 10 ⁵ CFU/seed
% Germ	7 DAP	100	100	100	100	90	90
Plant height (cm)	7 DAP	10.4	11.1	10.8	11.2	11	11.6
Plant height (cm)	14 DAP	15.3	16.2	17.1	17.2	16.9	17.7
Plant height (cm)	21 DAP	16.1	17.2	17.3	17.8	17.2	18.5
Plant height (cm)	28 DAP	16.29	17.21	17.66	18	17.767	18.74
Fresh Root Weight (g)	Wt (g)	0.367	0.419	0.41	0.437	0.413	0.429
Fresh Top Weight (g)	Wt (g)	0.21	0.2406	0.253	0.265	0.276	0.267
Fresh Total Weight (g)	Wt (g)	0.577	0.6596	0.663	0.702	0.689	0.696
Dry Root Weight (g)	Wt (g)	0.0627	0.0729	0.0713	0.0756	0.07	0.074
Dry Top Weight (g)	Wt (g)	0.0582	0.0688	0.0695	0.0733	0.071	0.074
Dry Total Weight (g)	Wt (g)	0.1209	0.1417	0.1408	0.1489	0.141	0.149

Table 19. Percent change over formulation control of germination percentage, plant height and weight in endophyte treated wheat. DAP = Days after planting

		MIC-96038/MIC-19994, 10 ¹ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ² CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ³ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁴ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁵ CFU/seed, % change over formulation control
% Germination	7 DAP	0.0	0.0	0.0	-10.0	-10.0
Plant height (cm)	7 DAP	6.7	3.8	7.7	5.8	11.5
Plant height (cm)	14 DAP	5.9	11.8	12.4	10.5	15.7
Plant height (cm)	21 DAP	6.8	7.5	10.6	6.8	14.9
Plant height (cm)	28 DAP	5.6	8.4	10.5	9.1	15.1
Fresh Root Weight (g)	Wt (g)	14.2	11.7	19.1	12.6	16.9
Fresh Top Weight (g)	Wt (g)	14.6	20.5	26.2	31.2	27.0
Fresh Total Weight (g)	Wt (g)	14.3	14.9	21.7	19.4	20.5
Dry Root Weight (g)	Wt (g)	16.3	13.7	20.6	11.1	18.6
Dry Top Weight (g)	Wt (g)	18.2	19.4	25.9	22.0	27.9
Dry Total Weight (g)	Wt (g)	17.2	16.5	23.2	16.3	23.1

Rice

[00396] Endophyte inocula as prepared in Example 4 was applied to rice seeds which had not previously been treated with chemical insecticide or fungicide. Endophyte treated seeds were prepared at 5 target doses: 10¹, 10², 10³, 10⁴, 10⁵ CFU/seed. Ten biological replicates each were planted for each treatment and control condition (no endophyte) in individual containers containing commercial potting media.

[00397] Emergence was recorded at 7 days post planting, and plant height (cm) scored at 7, 14, 21 and 28 days post planting. Above and below ground tissue was harvested from the experiment four weeks post-planting. The fresh weight of roots and shoots were recorded, then the samples were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot and root biomass was weighed and recorded.

Table 20. Germination percentage, plant height and weight of endophyte treated rice formulation controls. DAP = Days after planting

		Formulation control	MIC-96038/MIC-19994, 10 ³ CFU/seed	MIC-96038/MIC-19994, 10 ⁴ CFU/seed	MIC-96038/MIC-19994, 10 ⁵ CFU/seed	MIC-96038/MIC-19994, 10 ⁶ CFU/seed	MIC-96038/MIC-19994, 10 ⁷ CFU/seed
% Germ	7 DAP	90	100	100	100	80	90
Plant height (cm)	7 DAP	7.1	7.1	7.7	7.7	7.3	7.3
Plant height (cm)	14 DAP	11.7	11.5	12.9	13.1	14	12.6
Plant height (cm)	21 DAP	14.1	13.5	14.9	15.2	15.7	14.4
Plant height (cm)	28 DAP	14.57	13.99	15.35	15.67	15.8875	15.325
Fresh Root Weight (g)	Wt (g)	0.189	0.196	0.218	0.2056	0.245	0.222125
Fresh Top Weight (g)	Wt (g)	0.0879	0.0862	0.0993	0.1057	0.1135	0.108125
Fresh Total Weight (g)	Wt (g)	0.277	0.2828	0.3173	0.3113	0.3585	0.33025
Dry Root Weight (g)	Wt (g)	0.031	0.0291	0.0321	0.0335	0.036625	0.035875
Dry Top Weight (g)	Wt (g)	0.022	0.0211	0.0233	0.025	0.0275	0.0265
Dry Total Weight (g)	Wt (g)	0.053	0.0502	0.0554	0.0585	0.064125	0.062375

Table 21. Percent change over formulation control of germination percentage, plant height and weight in endophyte treated rice. DAP = Days after planting

		MIC-96038/MIC-19994, 10 ³ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁴ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁵ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁶ CFU/seed, % change over formulation control	MIC-96038/MIC-19994, 10 ⁷ CFU/seed, % change over formulation control
% Germination	7 DAP	11%	11%	11%	-11%	0%
Plant height (cm)	7 DAP	0%	8%	8%	3%	3%
Plant height (cm)	14 DAP	-2%	10%	12%	20%	8%
Plant height (cm)	21 DAP	-4%	6%	8%	11%	2%
Plant height (cm)	28 DAP	-4%	5%	8%	9%	5%
Fresh Root Weight (g)	Wt (g)	4%	15%	9%	30%	18%
Fresh Top Weight (g)	Wt (g)	-2%	13%	20%	29%	23%
Fresh Total Weight (g)	Wt (g)	2%	15%	12%	30%	19%
Dry Root Weight (g)	Wt (g)	-6%	3%	8%	18%	15%
Dry Top Weight (g)	Wt (g)	-5%	5%	13%	24%	20%
Dry Total Weight (g)	Wt (g)	-6%	4%	10%	20%	17%

Example 10: Cultivation and of Endophyte-Treated Plants in Greenhouse Experiment 2

[00398] A sandy loam growth substrate was mixed in the greenhouse and consisting of 60% loam and 40% mortar sand (Northeast Nursery, Peabody, MA). Prior to mixing, loam was sifted through a 3/8" square steel mesh screen to remove larger particles and debris. Soybean seeds were treated with commercial fungicidal and insecticidal treatment CruiserMaxx® Vibrance (Syngenta, Basel, Switzerland) according to the manufacturer's instructions. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 14, untreated seeds (lacking formulation and endophyte) were also planted. Each pot was filled with 600 mL of its respective soil, watered with 200 mL of water and then, nine seeds were sown evenly spaced in each pot (in a 3x3 pattern). Soil was then overlaid atop the seeds (estimated average planting depth of 1 inch) and an additional 110 mL of water was added to moisten the overlaying soil substrate. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Environmental conditions were set at 12h photoperiod, at 22/18C temperature for day/night period and light intensity was set at 650 $\mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds were watered to maintain approximately 80% soil capacity. Above ground tissue was harvested from the experiment three weeks post-planting. The tissues from individual replicate treatments (pots) were pooled and placed in an unlined paper bag. All tissues were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot biomass of individual treatment replicates (pots) was weighed and recorded

Results

[00399] All Formulations tested in soybean resulted in a less than 5% change in dry shoot biomass. Formulation C2 was neutral (-0.09%). Formulations A2 and A4 had a slight negative effect on dry shoot biomass (-3.57% and -1.8%, respectively). The difference between formulation that had slight beneficial (B_2) and slight negative effect (A_2) was in concentrators of oil and Triton X-100. MIC-31593 was the best overall performer with the formulations tested in this experiment; the greatest effect was observed with "B_2 low dose", "A2 medium dose" and "C_2 low dose", in that order, with an increase of dry shoot biomass by 6.9%, 6.87% and 5.26%; respectively, compared to the control. MIC-33414 demonstrated the largest positive effect when paired with C_2 Formulation at "medium dose", followed by "A_2 medium dose", "A_2 high dose" and "A_2 low dose", in that order of magnitude, with an increase of dry shoot biomass by 6.52%, 5.27% and 3.96% and 3.23; respectively,

compared to the non-treated control. MIC-96038 paired with Formulation A_f1, exhibited a slight increase in biomass at doses High and Low (2.05% and 4.25%, respectively).

Table 22. Dry shoot biomass of endophyte treated soybeans grown in sandy loam soil, using three different formulations

MIC ID	Formulation	Dose	Average dry biomass (g)	% change over untreated	Standard deviation	Standard error of the mean
	Untreated control		1.19	0	0.1027	0.0236
	Formulation control – A_2		1.15	-3.57	0.1485	0.0332
	Formulation control – B_2		1.22	2.88	0.1047	0.024
	Formulation control – C_2		1.19	-0.09	0.0957	0.022
	Formulation control – A_1		1.2	0.76	0.1039	0.0245
	Formulation control – A_4		1.17	-1.8	0.0899	0.0206
MIC-31593	A_2	low	1.23	3.45	0.1082	0.0248
		medium	1.27	6.87	0.1069	0.0245
		high	1.24	3.9	0.1218	0.0279
	B_2	low	1.27	6.9	0.0959	0.022
		medium	1.23	3.12	0.1038	0.0238
		high	1.2	0.47	0.1162	0.0267
	C_2	low	1.25	5.26	0.1279	0.0301
		medium	1.22	2.78	0.0872	0.0206
		high	1.21	1.68	0.1079	0.0247
MIC-96038	A_1	low	1.24	4.25	0.1273	0.0292
		medium	1.2	0.88	0.0776	0.0178
		high	1.21	2.05	0.1117	0.0256
MIC-33414	A_2	low	1.23	3.23	0.1099	0.0252
		medium	1.25	5.27	0.0985	0.0226
		high	1.24	3.96	0.0903	0.0207
	B_2	low	1.21	1.38	0.0716	0.0169
		medium	1.22	2.2	0.1161	0.0266
		high	1.21	1.58	0.084	0.0193
	C_2	low	1.17	-1.45	0.1148	0.0263
		medium	1.27	6.52	0.1044	0.0246
		high	1.18	-0.6	0.1076	0.0247

Example 11: Cultivation and of Endophyte-Treated Plants in Greenhouse Experiment 3

[00400] A sandy loam growth substrate was mixed in the greenhouse and consisting of 60% loam and 40% mortar sand (Northeast Nursery, Peabody, MA). Prior to mixing, loam was sifted through a 3/8" square steel mesh screen to remove larger particles and debris. Soybean seeds were treated with commercial fungicidal and insecticidal treatment CruiserMaxx® Vibrance (Syngenta, Basel, Switzerland) according to the manufacturer's instructions. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 11, untreated seeds (lacking formulation

and endophyte) were also planted. Endophyte treatments were applied to seed in two target doses: high (10^5 CFU/seed), medium (10^4 CFU/seed). Each pot was filled with 600 mL of its respective soil, watered with 200 mL of water and then, nine seeds were sown evenly spaced in each pot (in a 3x3 pattern). Soil was then overlaid atop the seeds (estimated average planting depth of 1 inch) and an additional 110 mL of water was added to moisten the overlaying soil substrate. The experimental design called for a completely randomized pattern of each treatment within each block/replicate. Environmental conditions were set at 12h photoperiod, at 22/18°C temperature for day/night period and light intensity was set at $650 \mu\text{Mol m}^{-2} \text{s}^{-1}$. Post-planting, the seeds grown in normal watering conditions were watered to maintain approximately 80% soil capacity and the seeds grown in drought conditions were watered to maintain approximately 40% soil capacity. Above ground tissue was harvested from the experiment three weeks post-planting. The tissues from individual replicate treatments (pots) were pooled and placed in an unlined paper bag. All tissues were dried in an oven set to 85°C for 3 days. Once completely dried, the shoot biomass of individual treatment replicates (pots) was weighed and recorded.

Table 23. Dry shoot biomass of endophyte treated wheat seedlings grown in sandy loam soil under normal conditions

Treatment	Average dry biomass (g)	% change over untreated	% change over formulation
Formulation control	1.852	-3.46	0
Untreated control	1.918	0	3.59
MIC-96038 10^4	1.987	3.58	7.3
MIC-96038 10^5	1.843	-3.91	-0.46
MIC-96038/MIC-19994 10^4	1.945	1.43	5.07
MIC-96038/MIC-19994 10^5	1.885	-1.72	1.8
MIC-19994 10^4	1.937	1.01	4.64
MIC-19994 10^5	1.915	-0.14	3.44

Example 12. Methods of Preparing Fungal Biomass and Treating Seeds for Field Experiments

Preparation of fungal endophytes

[00401] Preparation of molasses broth and potato dextrose agar: Molasses broth was prepared by dissolving 30 g molasses and 5 g yeast extract per liter deionized water in an autoclavable container and autoclaving (15 psi, 121°C) for 45 min. Potato dextrose agar (PDA) plates were prepared by dissolving 39.0 g PDA powder per liter deionized water in an autoclavable container and autoclaving (15 psi, 121°C) for 45 min. The agar was allowed to cool to 50-60°C, before pouring into sterile petri plates (30 mL per 90 mm plate).

[00402] Liquid biomass: All equipment and consumables were thoroughly sterilized and procedures performed in a biosafety cabinet. The inoculant is prepared by placing 1 plug from a cryopreserved stock on a fresh PDA plate, sealing the plate with Parafilm® and incubating at room temperature in the dark for 5-10 days. Then ~5x5 mm plugs were cut from the PDA plates and 10-12 plugs were transferred into flasks containing the sterile molasses broth, covered, secured in a shaker and incubated for at least 10 days with shaking at ~130 rpm. Then the culture was placed in a blender for 5 seconds and 1 mL of the blended was centrifuged and the supernatant was discarded and the pellet resuspended in 0.5 mL 1x Phosphate Buffered Saline (PBS) to generate inoculum.

[00403] Dry biomass: All equipment and consumables were thoroughly sterilized and procedures performed in a biosafety cabinet. The inoculant is prepared by placing 1 plug from a cryopreserved stock on a fresh PDA plate, sealing the plate with Parafilm® and incubating at room temperature in the dark for 5-10 days. Then ~5x5 mm plugs were cut from the PDA plates and 10-12 plugs were transferred into flasks containing the sterile molasses broth, covered, secured in a shaker and incubated for at least 10 days with shaking at ~130 rpm. In sterile conditions, the liquid culture was carefully decanted using 150 mm sterile filter paper on a sterilized Buchner funnel over a sterile flask. Once all liquid had passed through the funnel, the pellet was rinsed with sterile water until the filtrate ran clear. When dry, the pellet was transferred to a drying cabinet and dried until brittle. The pellet was then ground into a fine powder, and sample used to generate CFU counts.

Preparation of sodium alginate and talc for seed treatments

[00404] A 2% weight/volume solution of sodium alginate for the seed coatings is prepared by the following method. An Erlenmeyer flask is filled with the appropriate volume of deionized water and warmed to 50 degrees Celsius on a heat plate with agitation using a stir bar. The appropriate mass of sodium alginate powder for the desired final concentration solution is slowly added until dissolved. The solution is autoclaved at 121 degrees Celsius at 15 PSI for 30 minutes to sterilize.

[00405] Talcum powder for the powdered seed coatings is prepared by the following method. Talcum powder is aliquoted into Ziploc bags or 50 mL Falcon tubes, and autoclaved in dry cycle (121 degrees Celsius at 15 PSI for 30 minutes) to sterilize.

[00406] Seeds were heterologously disposed to each endophyte according to the following seed treatment protocol.

[00407] Liquid formulation: The 2% sodium alginate solution prepared above added to the seeds at a rate of 15 ml per kg of seeds. Liquid fungal culture as prepared in Example 10 was

added to the seeds at a rate of 8.3 ml per kg of seeds. Control treatments were prepared using equivalent volumes of sterile broth. The seeds were then agitated to disperse the solution evenly on the seeds.

[00408] Then 12.5 g of talc powder per kg of seed was added and the seeds were agitated to disperse the powder evenly on the seeds. Then 17 ml per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds. The final concentration of endophyte was targeted to be at least 10^4 CFU. Treated seeds were allowed to dry overnight in a well-ventilated space before planting.

[00409] Dry formulation: The 2% sodium alginate solution prepared above was added to the seeds at a rate of 20 ml per kg of seeds. Equal parts of the fungal powder prepared and the talc prepared in above were mixed. The solution is applied to the prepared seeds so that an equivalent of 12.5 g of fungal powder was applied per kg of seeds. Control treatments were prepared using equivalent volumes of talc. The seeds were then agitated to disperse the solution evenly on the seeds.

[00410] Then 17 ml per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds. The final concentration of endophyte was targeted to be at least 10^4 CFU. Treated seeds were allowed to dry overnight in a well-ventilated space before planting.

Heterologous disposition of endophytes on soy or peanut seeds

[00411] Seeds were heterologously disposed to each endophyte according to the following seed treatment protocol.

[00412] Liquid formulation: The 2% sodium alginate solution prepared in Example 10 was added to the seeds at a rate of 8.3 ml per kg of seeds. Liquid fungal culture as prepared in Example 9 was added to the seeds at a rate of 8.3 ml per kg of seeds. Control treatments were prepared using equivalent volumes of sterile broth. The seeds were then agitated to disperse the solution evenly on the seeds.

[00413] Then 15 g per kg of seed of the talc powder prepared in Example 10 was added and the seeds were agitated to disperse the powder evenly on the seeds. Then 13.3 ml per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds. The final concentration of endophyte was targeted to be at least 10^4 CFU. Treated seeds were allowed to dry overnight in a well-ventilated space before planting.

[00414] Dry formulation: The 2% sodium alginate solution prepared in Example 10 was added to the seeds at a rate of 16.6 ml per kg of seeds. Equal parts of the fungal powder prepared in Example 9 and the talc prepared in Example 10 were mixed. The solution was applied so that an equivalent of 10 g of fungal powder was applied per kg of seeds. Control treatments were prepared using equivalent volumes of talc. The seeds were then agitated to disperse the solution evenly on the seeds.

[00415] Then 13.3 ml per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds. The final concentration of endophyte was targeted to be at least 10^4 CFU. Treated seeds were allowed to dry overnight in a well-ventilated space before planting.

Heterologous disposition of endophytes on corn seeds

[00416] Seeds were heterologously disposed to each endophyte according to the following seed treatment protocol.

[00417] Dry formulation: The 2% sodium alginate solution prepared as above was added to the seeds at a rate of 23 ml per kg of seeds. Equal parts of the fungal powder and the talc as prepared above were mixed. The solution was applied so that an equivalent of 10 g of fungal powder was applied per kg of seeds. Control treatments were prepared using equivalent volumes of talc. The seeds were then agitated to disperse the solution evenly on the seeds.

[00418] Then 16.6 ml per kg of seed of Flo-Rite® 1706 (BASF, Ludwigshafen, Germany) was added and the seeds were agitated to disperse the powder evenly on the seeds. The final concentration of endophyte was targeted to be at least 10^4 CFU. Treated seeds were allowed to dry overnight in a well-ventilated space before planting.

Heterologous disposition of endophytes on rice

[00419] Seeds were treated with the commercial fungicidal and insecticidal product Sativa® IMF MAX (Nufarm Americas, Alsip, IL) per manufacturer's instructions (3.4oz/cwt). Chemically treated rice seeds were heterologously disposed to each endophyte according to one of two different formulation protocols (Formulation Protocol A, Formulation Protocol B). A corresponding seed formulation control, lacking any endophyte, was also prepared included for each type of formulation. Further seeds lacking any formulation and endophyte were planted as a non-treated baseline control. Formulation A included only diluent 0.05% silwet and microbial preparations. Formulation B included the same diluent and the seed plantability polymer Flo Rite® 1706 applied at 2.0oz/cwt seed per the manufacturer. Microbe and polymer were applied sequentially.

[00420] For endophytes formulated by Formulation Protocol A, microbial preparations were applied to the seeds at a rate of 1 μ L/seed and the seeds were agitated for at least 20 seconds to disperse the microbe.

[00421] For endophytes formulated by Formulation Protocol B, microbial preparations were applied to the seeds at a rate of 1 μ L/seed. Then Flo Rite[®] 1706 plantability polymer was applied to seeds per the manufacturer's suggestion (2.0 oz/cwt of seed) and agitated for 20 seconds to disperse the polymer.

Example 13: Cultivation of Endophyte-Treated Plants in Field Experiment 1

Assay of seed yield under field conditions, soy

[00422] Field trials were conducted under non-irrigated (dryland) conditions at multiple locations, preferably in diverse geographic regions. Seeds were prepared with the endophyte formulations as described in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer. Only the middle two rows of the 4 row plots were harvested to present border effects. Seeds were sown in regularly spaced rows in soil at 40,000 seeds/acre seeding density. At each location, at least 3 replicate plots were planted per endophyte or control treatment in a randomized complete block design. Each plot consisted of four 15.24 m (40 ft.) rows, each separated by 76.2 cm (30 in).

[00423] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer. Only the middle two rows of the 4 row plots were harvested to prevent border effects.

Table 24. Yield of treated endophyte treated soybeans and untreated and formulation controls, Field Experiment 1

	Location 1 yield (bu/ac)	Location 2 yield (bu/ac)	Location 3 yield (bu/ac)	Location 4 yield (bu/ac)	Overall yield (bu/ac)	Overall % Δ untreated control	Overall % Δ formulation control
Condition	Severe water stress	Water stress	Good moisture, some weed pressure	Weed pressure			
Days between treatment and planting	9	13	28	28			
Untreated control	27.7	68.4	61	41.6	49.7		-1.4
Formulation control	27.1	68.9	64.8	42.5	50.4	1.4	0

MIC-31593	27.9	70.2	62.1	48	52.1	4.8	3.4
MIC-96038	27	69.5	63.4	43.4	50.8	2.2	0.8
MIC-33414	27.9	72	67.2	43.6	52.1	4.8	3.4
Trial Average	25.7	68.1	62.5	45.1	50.4		

Assay of seed yield under field conditions, wheat

[00424] Field trials were conducted under non-irrigated (dryland) conditions at multiple locations, preferably in diverse geographic regions. Wheat seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were heterologously disposed with the endophyte formulations described above in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. Seeds were sown in regularly spaced rows in soil at 1.2 million seeds/acre seeding density. At each location, at least 3 replicate plots were planted for each endophyte or control treatment in a randomized complete block design. Each plot consisted of seven, 15.24 m (40 ft.) rows.

[00425] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

Table 25. Average yield of endophyte treated spring wheat and untreated and formulation controls by location, Field Experiment 1

	Location 1 yield (bu/ac)	Location 2 yield (bu/ac)	Location 3 yield (bu/ac)	Location 4 yield (bu/ac)	Overall yield (bu/ac)	Overall % Δ untreated control	Overall % Δ formulation control
Untreated control	25.73	39.50	38.95	47.02	37.80		-2.0
Formulation control	29.72	41.76	35.15	48.17	38.57	2.0	
MIC-96038	28.00	40.37	37.63	50.95	39.24	3.8	1.7
Trial Average	28.75	41.60	38.10	47.97	39.10		

Table 26. Average yield of endophyte treated spring wheat and untreated and formulation controls by plant variety, Field Experiment 1

	Variety 1, yield (bu/ac)	Variety 2, yield (bu/ac)	Variety 3, yield (bu/ac)	Variety 4, yield (bu/ac)	Overall yield (bu/ac)	Overall % Δ untreated control	Overall % Δ formulation control
Untreated	24.83	26.63	40.96	42.69	37.80		-2.0
Formulation control	31.22	28.22	38.83	44.23	38.57	2.0	
MIC-96038	32.54	23.46	42.60	43.37	39.24	3.8	1.7
Variety Average	30.62	28.44	40.77	43.95			

Assay of seed yield under field conditions, Rice

[00426] Seeds were prepared with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 12. Seeds were sown with a research sized

grain drill at a planting rate of 45 pounds per acre. Four replicate, 30x175 ft plots were planted per endophyte or control treatment in a randomized complete block design. Plots were flooded, weeds and insects were controlled with local standard practices.

[00427] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

Table 27. Yield of treated endophyte rice and untreated controls, Field Experiment 1

	% emergence	Grain moisture	Weight (lb/bushel)	Yield (lb/plot)	Yield (bushel/acre)	% change over control
Control treatment	65.3	19.33	43.95	16.405	119.6	0
MIC-96038 + MIC-19994	67.8	18.88	41.33	17.165	125.9	5.27

Example 14: Cultivation of Endophyte-Treated Plants in Field Experiment 2

Assay of seed yield under field conditions, wheat

[00428] Field trials were conducted under non-irrigated (dryland) conditions. Two varieties of spring wheat seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 12, untreated seeds (lacking formulation and endophyte) were also planted. Seeds were sown in regularly spaced rows in soil at 1.2 million seeds/acre seeding density. At each location replicate plots were planted for each endophyte or control treatment in a randomized complete block design). Each plot consisted of seven, 15.24 m (40 ft.) rows.

[00429] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield and grain percent moisture were calculated by the on-board computer.

Table 28. Percent emergence of endophyte treated spring wheat and untreated and formulation controls by location, Field Experiment 2

	Variety 1, % emergence	Variety 2, % emergence	Overall, % emergence	Overall % Δ untreated control	Overall % Δ formulation control
Untreated control	98.3	96.7	97.5		-0.8
Formulation control	98.3	98.3	98.3	0.8	
MIC-96038	98.3	98.3	98.3	0.8	0
MIC-19994	98.3	93.3	95.8	-1.7	-2.5

Table 29. Grain moisture of endophyte treated spring wheat and untreated and formulation controls by location, Field Experiment 2

	Variety 1, grain moisture (%)	Variety 2, grain moisture (%)	Overall, grain moisture (%)	Overall % Δ untreated control	Overall % Δ formulation control
Untreated control	19.049	16.657	17.85		1.0
Formulation control	19.397	15.970	17.68	-0.9	
MIC-96038	13.573	16.567	15.07	-15.6	-14.8
MIC-19994	18.657	16.930	17.79	-0.3	0.6

Table 30. Percent protein of endophyte treated spring wheat and untreated and formulation controls by location, Field Experiment 2

	Variety 1, % protein	Variety 2, % protein	Overall, % protein	Overall % Δ untreated control	Overall % Δ formulation control
Untreated control	14.55087	14.76347	14.657		0.2
Formulation control	14.68427	14.56693	14.626	-0.2	
MIC-96038	14.90133	14.71067	14.806	1.0	1.2
MIC-19994	14.30587	14.21200	14.259	-2.7	-2.5

Table 31. Yield of endophyte treated spring wheat and untreated and formulation controls by location, Field Experiment 2

	Variety 1, yield (bu/acre)	Variety 2, yield (bu/acre)	Overall, yield (bu/acre)	Overall % Δ untreated control	Overall % Δ formulation control
Untreated control	24.832	26.631	25.731		-13.3
Formulation control	31.172	28.217	29.694	15.4	
MIC-96038	32.539	23.457	27.998	8.8	-5.7
MIC-19994	31.172	26.892	29.032	12.8	-2.2

Example 15. Cultivation of Endophyte-Treated Plants in Field Experiment 3

[00430] Field trials were conducted in 2016, under non-irrigated (dryland) conditions. Wheat seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were heterologously disposed with the endophyte formulations and formulation control (lacking any endophyte) as described in Example 12, untreated seeds (lacking formulation and endophyte) were also planted. Seeds were sown in regularly spaced rows in soil at 1.2 million seeds/acre seeding density. At each location replicate plots were planted for each endophyte or control treatment in a randomized complete block design). Each plot consisted of seven, 15.24 m (40 ft.) rows.

[00431] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

Table 32. Yield of endophyte treated wheat and untreated and formulation controls by location, field experiment 3

	Variety 1, yield (bu/ac)	Variety 2, yield (bu/ac)	Variety 3, yield (bu/ac)	Variety 4, yield (bu/ac)	Overall, yield (bu/ac)	Overall % Δ	Overall % Δ
--	-----------------------------	-----------------------------	-----------------------------	-----------------------------	------------------------------	-----------------------	-----------------------

						untreated control	formulation control
Untreated control	27.9	27.4	39.8	43.2	34.6		-3.0
Formulation control	31.2	28.2	39.0	44.2	35.7	3.1	
MIC-96038	32.5	23.5	42.6	43.4	35.5	2.7	-0.4
MIC-19994	31.2	26.9	40.4	43.9	35.6	3.0	-0.1

Example 16. Assessment of Improved Plant Characteristics: Field Conditions

Assay of seed yield under field conditions, wheat

[00432] Field trials were conducted under non-irrigated (dryland) conditions at multiple locations, preferably in diverse geographic regions. Wheat seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were heterologously disposed with the dry endophyte formulations as described in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. Seeds were sown in regularly spaced rows in soil at 1.2 million seeds/acre seeding density. At each location, at least 3 replicate plots were planted for each endophyte or control treatment in a randomized complete block design. Each plot consisted of seven, 15.24 m (40 ft.) rows.

[00433] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

[00434] The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-68178, MIC-07010, MIC-31593, MIC-48747, MIC-96038, MIC-50414 or MIC-33414, resulted in average increases in yield of 7-16% in the wheat variety SDSU Focus. The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-68178, MIC-07010, MIC-31593, MIC-48747, MIC-96038, MIC-50414 or MIC-33414, resulted in average increases in yield of 14-22% in the wheat variety SDSU Select.

Table 33: Average yield of wheat treated with endophytes in field trials

	SDSU Focus, Variety 3		SDSU Select, Variety 4	
	Average yield (BU/acre)	% difference Untreated	Average yield (BU/acre)	% difference Untreated
Untreated control	36.9	0	37.7	0%
MIC-68390	39.3	7%	45.0	19%
MIC-68178	40.9	11%	46.1	22%
MIC-07010	41.1	11%	43.9	16%
MIC-31593	42.1	14%	44.3	18%
MIC-48747	42.8	16%	44.1	17%
MIC-96038	42.6	15%	43.4	15%

MIC-50414	40.0	8%	45.7	21%
MIC-33414	41.6	13%	42.9	14%

Assay of seed yield under field conditions, corn

[00435] Field trials were conducted at multiple locations, preferably in diverse geographic regions. Plots were non-irrigated (dryland) or maintained with suboptimal irrigation at a rate to target approximately 25% reduction in yield. Seeds were prepared with the endophyte formulations (dry) and formulation control (dry, lacking any endophyte) as described in Example 12, untreated seeds (lacking formulation and endophyte) were also planted. Seeds were sown in regularly spaced rows in soil at planting densities typical for each region. At each location 3 replicate plots were planted per endophyte or control treatment in a randomized complete block design. Each plot consisted of four 15.24 m (40 ft.) rows, each separated by 76.2 cm (30 in).

[00436] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer. Only the middle two rows of the 4 row plots were harvested to prevent border effects.

[00437] The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-48747, or MIC-33414, resulted in average increases in yield of 0.9-1.5% relative to formulation control and average increases in yield of 1.0-1.6% relative to the untreated control, in the corn variety Stine 9734.

Table 34: Average yield of corn variety Stine 9734 treated with endophytes in field trials

	Stine 9734, Variety 2		
	Average yield (BU/acre)	% difference Formulation control	% difference Untreated control
Untreated	185.5		0.0%
Formulation control (dry)	185.7	0.0%	
MIC-68390	187.4	0.9%	1.0%
MIC-48747	187.7	1.1%	1.2%
MIC-33414	188.5	1.5%	1.6%

Assay of seed yield under field conditions, soy

[00438] Field trials were conducted under non-irrigated (dryland) conditions at multiple locations, preferably in diverse geographic regions. Seeds were prepared with the endophyte

formulations as described in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. MIC-68178 and MIC-33414 were formulated with the dry formulation; MIC-68390, MIC-07010, MIC-31593, MIC-48747, MIC-96038, and MIC-50414 were formulated with the liquid formulation.

[00439] Seeds were sown in regularly spaced rows in soil at 40,000 seeds/acre seeding density. At each location, at least 3 replicate plots were planted per endophyte or control treatment in a randomized complete block design. Each plot consisted of four 15.24 m (40 ft.) rows, each separated by 76.2 cm (30 in).

[00440] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer. Only the middle two rows of the 4 row plots were harvested to prevent border effects.

[00441] The endophyte treatments, each comprising one of the following microbes: MIC-68390 MIC-31593, or MIC-33414, resulted in average increases in yield of 1.5-6.2% in the soy variety Dairyland DSR1808R2Y. The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-31593, or MIC-33414, resulted in average increases in yield of 1.5-6.2% in the soy variety Dairyland DSR1808R2Y. The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-68178, MIC-07010, MIC-31593, MIC-48747, MIC-96038, MIC-50414, or MIC-33414, resulted in average increases in yield of 2.5-15% in the soy variety Pfister 38R25. The endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-68178, MIC-07010, MIC-31593, MIC-48747, MIC-50414, or MIC-33414, resulted in average increases in yield of 1.1-6% in the soy variety Stine 3920.

Table 35: Average yield of soy variety Dairyland DSR1808R2Y treated with endophytes in field trials

	Dairyland DSR1808R2Y, Variety 1	
	Average yield (pounds/acre)	% difference Untreated control
Untreated control	33.9	0.0%
MIC-68390	35.1	3.5%
MIC-31593	34.4	1.5%
MIC-33414	36.0	6.2%

Table 36. Average yield of soy variety Pfister 38R25 treated with endophytes in field trials

	Pfister 38R25, Variety 2	
Row Labels	Average yield (pounds/acre)	% difference Untreated control
Untreated control	56.8	0%
MIC-68390	58.8	3.5%
MIC-68178	60.0	5.6%
MIC-07010	60.0	5.6%
MIC-31593	65.3	15.0%
MIC-48747	58.2	2.5%
MIC-96038	61.2	7.7%
MIC-50414	62.6	10.2%
MIC-33414	63.9	12.5%

Table 37. Average yield of soy variety Stine 3920 treated with endophytes in field trials

	Stine 3920, Variety 4	
Row Labels	Average yield (pounds/acre)	% difference Untreated control
Untreated control	56.9	0%
MIC-68390	58.4	2.6%
MIC-68178	59.7	4.9%
MIC-07010	58.9	3.5%
MIC-31593	60.3	6.0%
MIC-48747	57.5	1.1%
MIC-50414	57.5	1.1%
MIC-33414	60.2	5.8%

Assay of seed yield under field conditions, canola

[00442] Field trials were conducted at multiple locations, preferably in diverse geographic regions. Plots were non-irrigated (dryland) or maintained with suboptimal irrigation at a rate to target approximately 25% reduction in yield. Canola seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were prepared with the liquid endophyte formulations and liquid formulation control (lacking any endophyte) as described in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. At each location, at least 3 replicate plots were planted for each endophyte or control treatment in a randomized complete block design.

[00443] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

[00444] The endophyte treatments comprising MIC-85555 resulted in an average increase in yield of 0.3% relative to formulation control and an average increase in yield of 0.2% relative to the untreated control, in the canola variety Brett Young 5525. The endophyte treatments comprising MIC-50989 resulted in an average increase in yield of 8.2% relative to formulation control and an average increase in yield of 10.3% relative to the untreated control, in the canola variety NCC1015.

Table 38. Average yield of canola variety Brett Young 5525 treated with endophytes in field trials

	Brett Young 5525, Variety 1		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated	1111.0		0.0%
Formulation control (liquid)	1109.5	0.0%	
MIC-85555	1112.7	0.3%	0.2%

Table 39. Average yield of canola variety NCC1015 treated with endophytes in field trials

	NCC1015, Variety 2		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated	1120.3		0.0%
Formulation control (liquid)	1142.8	0.0%	
MIC-50989	1236.1	8.2%	10.3%

Assay of seed yield under field conditions, Peanut

[00445] Field trials were conducted at multiple locations, preferably in diverse geographic regions. Plots were non-irrigated (dryland) or maintained with suboptimal irrigation at a rate to target approximately 25% reduction in yield. Peanut seeds were treated with commercial fungicidal and insecticidal treatment. Seeds were prepared with either the endophyte formulations and formulation control (lacking any endophyte) as described in Example 12 and untreated seeds (lacking formulation and endophyte) were also planted. MIC-68390 was formulated with both the dry and liquid formulation; MIC-50414, MIC-68178, and MIC-96038 were formulated with the liquid formulation. At each location, at least 3 replicate plots were planted for each endophyte or control treatment in a randomized complete block design.

[00446] At the end of the field trial employing endophyte treatment and control treatment plants, plots were machine harvested with a 5-ft research combine and yield calculated by the on-board computer.

[00447] In dry formulation, the endophyte treatments comprising MIC-68390 resulted in an average increase in yield of 9.1% relative to formulation control and an average increase in yield of 0.7% relative to the untreated control, in the peanut variety AT-9899. In liquid formulation, the endophyte treatments comprising MIC-50414 resulted in an average increase in yield of 3.8% relative to formulation control and an average increase in yield of 0.7% relative to the untreated control, in the peanut variety AT-9899. In liquid formulation, the endophyte treatments, each comprising one of the following microbes: MIC-68390 or MIC-68178, resulted in an average increase in yield of 4.1-4.5% relative to formulation control and an average increase in yield of 10.0-10.5% relative to the untreated control, in the peanut variety FloRun 107. However, in dry formulation, endophyte treatments comprising MIC-68390 resulted in an average decrease in yield of 6.3% in the peanut variety FloRun 107. In liquid formulation, the endophyte treatments, each comprising one of the following microbes: MIC-68390, MIC-68178, or MIC-96038, resulted in an average increase in yield of 0.2-3.6% relative to the formulation control and an average increase in yield of 6.4-10.0% relative to untreated control, in the peanut variety Georgia-06G. However, in dry formulation, endophyte treatments comprising MIC-68390 resulted in an average decrease in yield of 4.9% in the peanut variety Georgia-06G. In liquid formulation, the endophyte treatments comprising MIC-68390 resulted in an average increase in yield of 11.2% relative to formulation control and an average increase in yield of 4.9% relative to the untreated control, in the peanut variety Tamnut OL06. However, in dry formulation, endophyte treatments comprising MIC-68390 resulted in an average decrease in yield of 0.4% in the peanut variety Tamnut OL06.

Table 40. Average yield of peanut variety AT-9899 treated with endophytes in dry formulation in field trials

	AT-9899, Variety 2		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated control	3567.8		0.0%
Formulation control (dry)	3294.0	0.0%	
MIC-68390	3592.7	9.1%	0.7%

Table 41. Average yield of peanut variety AT-9899 treated with endophytes in liquid formulation in field trials

	AT-9899, Variety 2		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control

Untreated control	3567.8		0.0%
Formulation control (liquid)	3459.9	0.0%	
MIC-50414	3592.7	3.8%	0.7%

Table 42. Average yield of peanut variety FloRun 107 treated with endophytes in liquid formulation in field trials

	FloRun 107, Variety 3		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated control	4140.7		0.0%
Formulation control (liquid)	4375.1	0.0%	
MIC-68390	4555.7	4.1%	10.0%
MIC-68178	4574.0	4.5%	10.5%

Table 43. Average yield of peanut variety Georgia-06G treated with endophytes in liquid formulation in field trials

	Georgia-06G, Variety 4		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated control	4592.3		0.0%
Formulation control (liquid)	4876.3	0.0%	
MIC-68390	5053.2	3.6%	10.0%
MIC-68178	4884.1	0.2%	6.4%
MIC-96038	4966.5	1.9%	8.2%

Table 44. Average yield of peanut variety Tamnut OL06 treated with endophytes in liquid formulation in field trials

	Tamnut OL06, Variety 5		
	Average yield (pounds/acre)	% difference Formulation control	% difference Untreated control
Untreated control	4098.8		0.0%
Formulation control	3866.5	0.0%	
MIC-68390	4297.9	11.2%	4.9%

Example 17. Identification of sequence variants across core genes

[00448] Phylogenomic analysis of whole genome sequences of endophytes can be used to identify distinguishing sequence variants. Sets of genes suitable for phylogenomic analysis as well as methods for identifying the same are well known in the art, for example Floutas et al. (2012) The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31

fungus genomes. *Science*, 336(6089):1715-9. doi: 10.1126/science.1221748 and James TY, Pelin A, Bonen L, Ahrendt S, Sain D, Corradi N, Stajich JE. Shared signatures of parasitism and phylogenomics unite Cryptomycota and microsporidia. *Curr Biol*. 2013;23(16):1548-53. doi: 10.1016/j.cub.2013.06.057. Orthologous genes to the reference set are identified in protein data bases derived from the genome of each species. Orthologous genes can be identified in the genomes using methods well known including reciprocal best hits (Ward N, Moreno-Hagelsieb G. Quickly Finding Orthologs as Reciprocal Best Hits with BLAT, LAST, and UBLAST: How Much Do We Miss? de Crécy-Lagard V, ed. *PLoS ONE*. 2014;9(7):e101850. doi:10.1371/journal.pone.0101850) and Hidden Markov Models (HMMs). The best hits are extracted and a multiple sequence alignment generated for each set of orthologous genes. The alignments are used to build phylogenetic trees using methods well known in the art including Bayesian inference and maximum likelihood methods, for example using software tools MrBayes (Huelsenbeck, J.P. & Ronquist (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17(8):754-755) and RAxML (Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30 (9): 1312-1313. doi: 10.1093/bioinformatics/btu033). Sequence variants which distinguish between closely related species are identified.

Example 18. Identification of unique genes in an endophyte of interest

[00449] Whole genome analysis of endophytes can be used to identify genes whose presence, absence or over or under representation (“differential abundance”) are associated with desirable phenotypes. To identify genes with differential abundance in the genome of an endophyte of interest, protein sequences predicted from the genomes of the endophyte and closely related species compared in an all-vs-all pairwise comparison (for example, using BLAST) followed by clustering of the protein sequences based on alignment scores (for example, using MCL: Enright A.J., Van Dongen S., Ouzounis C.A. An efficient algorithm for large-scale detection of protein families. *Nucleic Acids Research* 30(7):1575-1584 (2002)). Additional software tools useful for this analysis are well known in the art and include OMA, OrthoMCL and TribeMCL (Roth AC, Gonnet GH, Dessimoz C. Algorithm of OMA for large-scale orthology inference. *BMC Bioinformatics*. 2008;9:518. doi: 10.1186/1471-2105-9-518, Enright AJ, Kunin V, Ouzounis CA. Protein families and TRIBES in genome sequence space. *Nucleic Acids Res*. 2003;31(15):4632-8.; Chen F, Mackey AJ, Vermunt JK, Roos DS. Assessing performance of orthology detection strategies applied to eukaryotic genomes. *PLoS One*. 2007;2(4):e383.). The protein clusters are queried to identify clusters

with differential abundance of proteins derived from endophytes having desirable phenotypes. Proteins of these clusters define the unique properties of these endophytes, and the abundance of genes encoding these proteins may be used to identify endophytes of the present invention.

[00450] Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other embodiments, advantages, and modifications are within the scope of the following claims.

WE CLAIM:

1. A method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the corn plant element is the variety Stine 9734 or a closely related variety thereof.
2. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.
3. The method of Claim 2, wherein the soybean plant is a variety selected from the group consisting of Pfister 38R25, Dairyland DSR1808R2Y, Stine 3920, and closely related varieties thereof.
4. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.
5. The method of Claim 4, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
6. A method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

7. The method of Claim 6, wherein the peanut plant is a variety selected from the group consisting of AT-9899, FloRun 107, Tamnut OL06, Georgia-06G, and closely related varieties thereof.
8. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and root dry weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.
9. The method of Claim 8, wherein the soybean is variety Stine 33E22 or a closely related variety thereof.
10. A method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.
11. The method of Claim 10, wherein the peanut plant is a variety selected from the group consisting of FloRun 107, Georgia-06G, and closely related varieties thereof.
12. A method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase root length in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.
13. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

14. A method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the canola plant is the variety Brett Young 552 or a closely related variety thereof.
15. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66.
16. The method of Claim 15, wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
17. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 65 and 66.
18. The method of Claim 17, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
19. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45.

20. The method of Claim 19, wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
21. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, and 45.
22. The method of Claim 21, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
23. A method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the corn plant element is the variety Stine 9734 or a closely related variety.
24. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67.
25. The method of Claim 24, wherein the trait of agronomic importance is yield and the soybean plant is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
26. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67.

27. The method of Claim 26, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
28. A method of improving a trait of agronomic importance in a canola plant, comprising heterologously disposing an endophyte to a canola plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69.
29. The method of Claim 28, wherein the canola plant is variety NCC1015 or a closely related variety thereof.
30. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.
31. The method of Claim 30, wherein the trait of agronomic importance is yield and the soybean plant is variety Pfister 38R25 or a closely related variety thereof.
32. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of decreasing grain moisture and increasing yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.
33. The method of Claim 32, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
34. A method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein:

- a. the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and
 - b. the peanut plant is a variety Georgia-06G or a closely related variety.
35. A method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing an endophyte to a rice plant element in an amount effective to increase dry shoot biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.
36. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing a first endophyte and a second endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of plant height, fresh root weight, and fresh shoot weight in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein:
- a. the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and
 - b. the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.
37. A method of improving a trait of agronomic importance in a rice plant, comprising heterologously disposing a first endophyte and a second endophyte to a rice plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein:
- a. the first endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and
 - b. the second endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

38. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68.
39. The method of Claim 38, wherein the soybean is a variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
40. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68.
41. The method of Claim 40, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
42. A method of improving a trait of agronomic importance in a peanut plant, comprising heterologously disposing an endophyte to a peanut plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68 and the peanut plant is the variety AT-9899 or a closely related variety thereof.
43. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing a first endophyte and a second endophyte to a soybean plant element in an amount effective to increase dry root biomass in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein:
- the first endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71,
 - the second endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59,
 - and soybean plant is variety Stine 33E22 or a closely related variety thereof.

44. A method of improving a trait of agronomic importance in a corn plant, comprising heterologously disposing an endophyte to a corn plant element in an amount effective to increase yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.
45. A method of improving a trait of agronomic importance in a soybean plant, comprising heterologously disposing an endophyte to a soybean plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root area, root length, dry shoot biomass, and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.
46. The method of Claim 45, wherein the trait of agronomic importance is yield and the soybean plant is the variety Pfister 38R25 or a closely related variety thereof.
47. A method of improving a trait of agronomic importance in a wheat plant, comprising heterologously disposing an endophyte to a wheat plant element in an amount effective to improve a trait of agronomic importance selected from the group consisting of root length and yield in the plant derived from the treated plant element relative to a plant derived from a reference plant element, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.
48. The method of Claim 47, wherein the wheat plant is a variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
49. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the plant element is a seed.
50. The method of Claim 49, wherein the seed is modified.
51. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the endophyte is heterologously disposed to the plant element in a formulation, said formulation further

comprising one or more of the following: a stabilizer, a preservative, a carrier, a surfactant, a fungicide, a nematocide, a bactericide, an insecticide, or herbicide, or any combination thereof.

52. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is at least 98%.

53. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is at least 99%.

54. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is 100%.

55. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is determined over a region of alignment of at least 100 nucleotides.

56. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is determined over a region of alignment of at least 200 nucleotides.

57. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is determined over a region of alignment of at least 300 nucleotides.

58. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is determined over a region of alignment of at least 400 nucleotides.

59. The method of any one of Claims 1, 2, 4, 6, 8, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 26, 28, 30, 32, 34, 35, 36, 37, 38, 40, 42, 43, 44, 45, and 47, wherein the percent identity is determined over a region of alignment of at least 500 nucleotides.

60. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62.

61. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active

ingredient comprises a first endophyte of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71.

62. The agrochemically active microbial formulation of Claim 60, further comprising a second endophyte, wherein the second endophyte is of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

63. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

64. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70.

65. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66.

66. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68.

67. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69.

68. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Acremonium* and comprises at least one

polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51.

69. The agrochemically active microbial formulation of Claim 66, wherein the microbial active ingredient further comprises a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41.

70. An agrochemically active microbial formulation, comprising at least one oil, surfactant, polymer, and a microbial active ingredient, wherein the microbial active ingredient comprises a first endophyte of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59.

71. The formulation of any one of Claims 60-70, wherein the oil is rapeseed or NEEM.

72. The formulation of any one of Claims 60-70, wherein the oil comprises erucic acid.

73. The formulation of any one of Claims 60-70, wherein the oil comprises herbicidal or insecticidal properties.

74. The formulation of any one of Claims 60-70, wherein the surfactant is a non-ionic detergent.

75. The formulation of any one of Claims 60-70, wherein the surfactant is Tween 20 or Triton X-100.

76. The formulation of any one of Claims 60-70, wherein the polymer is Flo Rite®, DISCO™, or Kannar® Universal Wonder.

77. The formulation of any one of Claims 60-70, wherein the microbial active ingredient comprises a spore suspension.

78. The formulation of any one of Claims 60-70, wherein the microbial active ingredient comprises spray dried spores.

79. The formulation of any one of Claims 60-70, wherein the microbial active ingredient comprises whole cell broth.

80. The formulation of any one of Claims 60-70 that further comprises one or more of the following: fungicide, nematocide, bactericide, insecticide, or herbicide.

81. The formulation of any one of Claims 60-70 that further comprises one or more of the following: stabilizer, preservative, carrier, anticomplex agent, or any combination thereof.

82. The formulation of any one of Claims 60-70, wherein the endophyte is shelf-stable.

83. The formulation of any one of Claims 60-70, wherein the percent identity is at least 98%.
84. The formulation of any one of Claims 60-70, wherein the percent identity is at least 99%.
85. The formulation of any one of Claims 60-70, wherein the percent identity is 100%.
86. The formulation of any one of Claims 60-70, wherein the percent identity is determined over a region of alignment of at least 100 nucleotides.
87. The formulation of any one of Claims 60-70, wherein the percent identity is determined over a region of alignment of at least 200 nucleotides.
88. The formulation of any one of Claims 60-70, wherein the percent identity is determined over a region of alignment of at least 300 nucleotides.
89. The formulation of any one of Claims 60-70, wherein the percent identity is determined over a region of alignment of at least 400 nucleotides.
90. The formulation of any one of Claims 60-70, wherein the percent identity is determined over a region of alignment of at least 500 nucleotides.
91. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Exserohilum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 61, and 62, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
92. The synthetic composition of Claim 91, wherein the plant element is a corn plant element and the improved trait of agronomic importance is yield.
93. The synthetic composition of Claim 91, wherein the plant element is a soybean plant element.
94. The synthetic composition of Claim 93, wherein the plant element is a soybean variety selected from the group consisting of: Pfister 38R25, Dairyland DSR1808R2Y, Stine 3920, and closely related varieties thereof, and the improved trait of agronomic importance is yield.
95. The synthetic composition of Claim 91, wherein the plant element is a wheat plant element and the improved trait of agronomic importance is yield.
96. The synthetic composition of Claim 91, wherein the plant element is a peanut plant element and the improved trait of agronomic importance is yield.

97. The synthetic composition of Claim 91, wherein the plant element is a peanut variety selected from the group consisting of: AT9899, FloRun 107, Georgia-06G, Tamnut OL06, and closely related varieties thereof.
98. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 63, 64, and 71, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
99. The synthetic composition of Claim 98, wherein the plant element is a soybean plant element and the improved trait of agronomic importance is selected from the group consisting of root area, root length, and root dry weight.
100. The synthetic composition of Claim 98, wherein the plant element is a soybean variety Stine 33E22 or a closely related variety thereof.
101. The synthetic composition of Claim 98, wherein the plant element is a peanut plant element and the improved trait of agronomic importance is yield.
102. The synthetic composition of Claim 101, wherein the peanut plant element is selected from the group consisting of the variety FloRun 107, Georgia-06G, and closely related varieties thereof.
103. The synthetic composition of Claim 98 further comprising a second endophyte, wherein the second endophyte is of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
104. The synthetic composition of Claim 99, wherein the plant element is the soybean variety Stine 33E22 and the trait of agronomic importance is dry root biomass.
105. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

106. The synthetic composition of Claim 105, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.
107. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Epicoccum* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 70, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
108. The synthetic composition of Claim 107, wherein the plant element is canola variety Brett Young 5525 or a closely related variety.
109. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Curvularia* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 42, 43, 44, 45, 65, and 66, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
110. The synthetic composition of Claim 109, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.
111. The synthetic composition of Claim 110, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
112. The synthetic composition of Claim 109, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.
113. The synthetic composition of Claim 112, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
114. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 67, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
115. The synthetic composition of Claim 114, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.

116. The synthetic composition of Claim 115, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof.
117. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Cladosporium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 67 and 68, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
118. The synthetic composition of Claim 117, wherein the plant element is a corn plant element and the trait of agronomic importance is yield.
119. The synthetic composition of Claim 118, wherein the plant element is corn variety Stine 9734.
120. The synthetic composition of Claim 117, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, and yield.
121. The synthetic composition of Claim 120, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof and the trait of agronomic importance is yield.
122. The synthetic composition of Claim 117, wherein the plant element is a wheat plant element and the trait of agronomic importance is yield.
123. The synthetic composition of Claim 122, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof.
124. The synthetic composition of Claim 117, wherein the endophyte comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 68, the plant element is a peanut of variety AT9899, and the trait of agronomic importance is yield.
125. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Paecilomyces* and comprises at least one polynucleotide sequence at least 97% identical to SEQ ID NO: 69, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.
126. The synthetic composition of Claim 125, wherein the plant element is canola variety NCC1015 and the trait of agronomic importance is yield.

127. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Acremonium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 46, 47, 48, 49, 50, and 51, and wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

128. The synthetic composition of Claim 127, wherein the plant element is a soybean plant element and the trait of agronomic importance is dry shoot biomass.

129. The synthetic composition of Claim 127, wherein the plant element is a wheat plant element and the trait of agronomic importance is grain moisture or yield.

130. The synthetic composition of Claim 129, wherein the plant element is wheat variety SDSU Select and the trait of agronomic importance is yield.

131. The synthetic composition of Claim 127, wherein the plant element is peanut variety Georgia-06G and the trait of agronomic importance is yield.

132. The synthetic composition of Claim 127, wherein the plant element is a rice plant element and the trait of agronomic importance is dry shoot biomass.

133. The synthetic composition of Claim 127 further comprising a second endophyte, wherein the second endophyte is of the genus *Coniochaeta* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 38, 39, 40, and 41, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

134. The synthetic composition of Claim 133, wherein the plant element is a wheat plant element and the trait of agronomic importance is selected from the group consisting of plant height, fresh root biomass, and fresh shoot weight.

135. The synthetic composition of Claim 133, wherein the plant element is a rice plant element and the trait of agronomic importance is yield.

136. A synthetic composition comprising a plant element and a heterologously disposed endophyte, wherein the endophyte is a member of the genus *Chaetomium* and comprises at least one polynucleotide sequence at least 97% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 52, 53, 54, 55, 56, 57, 58, and 59, wherein the synthetic composition is capable of providing an improved trait of agronomic importance as compared to a reference plant element not further comprising the endophyte.

137. The synthetic composition of Claim 136, wherein the plant element is a soybean plant element and the trait of agronomic importance is selected from the group consisting of root area, root length, dry shoot biomass, and yield.
138. The synthetic composition of Claim 137, wherein the plant element is a soybean variety selected from the group consisting of Pfister 38R25, Stine 3920, and closely related varieties thereof, and the trait of agronomic importance is yield.
139. The synthetic composition of Claim 136, wherein the plant element is a wheat plant element and the trait of agronomic importance is root length or yield.
140. The synthetic composition of Claim 139, wherein the plant element is a wheat variety selected from the group consisting of SDSU Focus, SDSU Select, and closely related varieties thereof and the trait of agronomic importance is yield.
141. The synthetic composition of any one of Claims 91-140, wherein the improved trait of agronomic importance is conferred under normal watering conditions.
142. The synthetic composition of any one of Claims 91-140, wherein the plant element is a seed.
143. The synthetic composition of Claim 142, wherein the seed is modified.
144. The synthetic composition of any one of Claims 91-140, wherein the plant element is placed into a substrate that promotes plant growth.
145. The synthetic composition of Claim 144, wherein the substrate that promotes plant growth is soil.
146. A plurality of synthetic compositions of Claim 145, wherein a plurality of said plant elements are placed in the soil in rows, with substantially equal spacing between each seed within each row.
147. The synthetic composition of any one of Claims 91-140 further comprising a formulation that comprises one or more of the following: stabilizer, preservative, carrier, surfactant, anticomplex agent, or any combination thereof.
148. The synthetic composition of any one of Claims 91-140 further comprising a formulation that comprises one or more of the following: fungicide, nematocide, bactericide, insecticide, or herbicide.
149. The synthetic composition of any one of Claims 91-140, wherein said compositions are confined within an object selected from the group consisting of: bottle, jar, ampule, package, vessel, bag, box, bin, envelope, carton, container, silo, shipping container, truck bed, or case.

150. The synthetic composition of any one of Claims 91-140, wherein the synthetic combinations are shelf-stable.
151. The synthetic composition of any one of Claims 91-140, wherein the percent identity is at least 98%.
152. The synthetic composition of any one of Claims 91-140, wherein the percent identity is at least 98%.
153. The synthetic composition of any one of Claims 91-140, wherein the percent identity is 100%.
154. The synthetic composition of any one of Claims 91-140, wherein the percent identity is determined over a region of alignment of at least 100 nucleotides.
155. The synthetic composition of any one of Claims 91-140, wherein the percent identity is determined over a region of alignment of at least 200 nucleotides.
156. The synthetic composition of any one of Claims 91-140, wherein the percent identity is determined over a region of alignment of at least 300 nucleotides.
157. The synthetic composition of any one of Claims 91-140, wherein the percent identity is determined over a region of alignment of at least 400 nucleotides.
158. The synthetic composition of any one of Claims 91-140, wherein the percent identity is determined over a region of alignment of at least 500 nucleotides.


Figure 1A



Control


MIC-68178

Figure 2



Control

MIC-68178

Figure 3

Control

MIC-68178/MIC-33414

Figure 4




Figure 5

