



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 47/00	A2	(11) International Publication Number: WO 98/01037 (43) International Publication Date: 15 January 1998 (15.01.98)
(21) International Application Number: PCT/US97/12018 (22) International Filing Date: 10 July 1997 (10.07.97) (30) Priority Data: 08/676,617 10 July 1996 (10.07.96) US (71) Applicant: THE DOW CHEMICAL COMPANY [US/US]; 2030 Dow Center, Midland, MI 48674 (US). (72) Inventors: SHANKAR, Ravi, B.; 4300 Congress Drive, Midland, MI 48642 (US). ROMER, Duane, R.; 4025 Lowell Court, Midland, MI 48642 (US). PEWS, R., Garth; 4403 Andre Street, Midland, MI 48642 (US). (74) Agent: JONES, S., Preston; Patent Dept., P.O. Box 1967, Midland, MI 48641-1967 (US).		(81) Designated States: CA, JP, NO, SG, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: THIOCYANOMETHYLOXO(AND THIOCYANOTHIOMETHYL)-1,2,3-BENZATRIAZIN-4(3H)ONES, COMPOSITIONS CONTAINING THEM AND THEIR USE AS ANTIMICROBIAL AND MARINE ANTIFOULING AGENTS (Ia) (Ib) (57) Abstract Disclosed are thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)-one compounds corresponding to one of formulae (Ia) or (Ib) wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C ₁ -C ₇ straight or branched chain alkyl, C ₁ -C ₇ straight or branched chain alkoxy, -CF ₃ , -OCF ₃ , -CN, -COOR wherein R is C ₁ -C ₇ straight or branched chain alkyl and n is an integer of from 0-4. These compounds have been found to exhibit antimicrobial and marine antifouling activity in industrial and commercial applications and compositions containing these compounds are so employed.		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THIOCYANOMETHYLOXO(AND THIOCYANOTHIOMETHYL)-1,2,3-
-BENZATRIAZIN-4(3H)ONES, COMPOSITIONS CONTAINING THEM AND THEIR
USE AS ANTIMICROBIAL AND MARINE ANTIFOULING AGENTS

The present invention is directed to thiocyanomethyloxo(and
thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compounds, compositions
containing them and their use as antimicrobial and marine antifouling agents.

The present invention is directed to thiocyanomethyloxo(and
thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compounds corresponding to one of
the formulae:



wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C_1-C_7 straight or
branched chain alkyl, C_1-C_7 straight or branched chain alkoxy, $-CF_3$, $-OCF_3$, -CN,
-COOR wherein R is -H or C_1-C_7 straight or branched chain alkyl and n is an integer of
from 0-4.


The present invention is also directed to antimicrobial compositions
comprising an inert diluent in admixture with an antimicrobially-effective amount of a
thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compound.

The present invention is further directed to a method for inhibiting
microorganisms in a microbial habitat comprising contacting said microbial habitat with
a composition containing an antimicrobially effective amount of a
thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compound.

The antimicrobial compositions of the present invention can also be employed to treat surfaces exposed to a marine environment in which marine organisms grow to prevent the growth of said marine organisms on said surfaces.

The preferred compounds of the present invention which are used in the antimicrobial compositions of the present invention include those wherein Z is oxygen and n is 0.

The 3-thiocyanomethyloxo(and thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compounds of the present invention can be prepared by reacting, with agitation and in a polar solvent, a 3-halomethyloxo(or 3-halomethylthiomethyl)-1,2,3-benzotriazin-4(3H)one corresponding to the formula



wherein halo represents bromo, chloro, fluoro or iodo with an alkali metal thiocyanate.

In the present specification and claims, the term "alkali metal" is employed to designate cesium, lithium, potassium or sodium.

Representative polar solvents which can be employed in the present process include dimethylformamide, dimethylsulfoxide, dimethylacetamide, N-methylpyrrolidinone and acetone.

The reactions are typically carried out at a temperature of from room temperature to 100°C. under a nitrogen atmosphere. The reactants may be added to the reaction mixture in any order of addition; conventionally they are added as a solution in the solvent used for the reaction. The reaction is allowed to continue over a period of from 10 minutes to 15 hours. The reaction consumes the reactants in the ratio of one mole equivalent of the halide reactant per mole of the thione reactant. To insure the completion of the reaction, an excess of the alkali metal thiocyanate reactant is normally employed.

After the reaction is complete, the product normally precipitates out and is recovered by filtration. To insure product precipitation, an equal volume amount of water and/or methanol can be added to the reaction mixture. The recovered product is washed with water or methanol and dried.

The following examples illustrate the present invention and the manner by which it can be practiced but, as such, should not be construed as limitations upon the overall scope of the same. In the following examples, certain specific alkali metals, halo groups, solvents and the like are set forth. These specific representations are only presented for convenience and are not to be considered as an indication that these specific representations are the only groups or materials which can be employed.

Since the hereinabove and hereinafter set forth compound preparation procedures employ only standard chemistry practices and it is known that slightly different reactants can require slightly different reaction parameters from those for other reactants, it is to be understood that minor modifications to the reaction parameters set forth such as the use of an excess of one reactant, the use of a catalyst, the use of temperatures slightly higher than room temperature and/or high speed mixing and other such conventional changes are within the scope of the present invention.

In the present invention, it is to be noted that all substituent groups are sterically compatible with each other. The term "sterically compatible" is employed to designate substituent groups which are not affected by steric hindrance as this term is defined in "The Condensed Chemical Dictionary", 7th edition, Reinhold Publishing Co., N.Y. page 893 (1966) which definition is as follows:


"steric hindrance. A characteristic of molecular structure in which the molecules have a spatial arrangement of their atoms such that a given reaction with another molecule is prevented or retarded in rate."

Sterically compatible may be further defined as reacting compounds having substituents whose physical bulk does not require confinement within volumes insufficient for the exercise of their normal behavior as discussed in *Organic Chemistry* by D. J. Cram and G. Hammond, 2nd edition, McGraw-Hill Book Company, N.Y., page 215 (1964).

The structure identity of all compounds is confirmed by proton nuclear magnetic resonance spectroscopy (^1H NMR), recorded at 300 MHz; carbon nuclear magnetic resonance spectroscopy (^{13}C NMR) recorded at 75 MHz; infrared spectroscopy (IR) and mass spectrometry (MS). All of the reactions are conducted under a positive pressure of nitrogen.

EXAMPLE I:

3-Thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one



5 A mixture of 1.28 grams(g) (0.006 mol) of 3-chloromethyloxo-1,2,3-benzotriazin-4(H)one and 0.65 g (1.1 equivalents) of potassium thiocyanate in 30 mL of dimethylformamide was heated at 85°C. overnight. The reaction mixture was poured into water and the product was collected by filtration. The title compound was recovered as a tan solid which melted at 120-121°C. in a yield of 43 percent of theoretical. ¹H NMR (DMSO, d₆) δ 6.16 (s, 2H), 7.99 (t, J=7.5 Hz, 1H), 8.16 (t, J=7.7 Hz, 1H), 8.32 (t, J=7.7 Hz, 1H); ¹³C NMR (DMSO, d₆) 80.69, 112.12, 122.12, 125.00, 128.43, 132.96, 135.66, 143.36, 150.26; MS (EI) m/e 234 (M⁺), 148, 134, 90.

Preparation of Starting Materials:

15 The 3-halomethyloxo-1,2,3-benzotriazin-4(3H)one employed as a starting material can be prepared by reacting 3-hydro-1,2,3-benzotriazin-4(3H)one with a dihalomethane in a halogenated solvent and in the presence of a halide acceptor such as trimethyl amine, pyridine or the like. The thus formed halo product can be separated employing conventional separatory procedures. Other halomethoxy products can be, if desired, prepared by conventional halogen exchange using procedures known in the art.

20 Example II:

3-Chloromethyloxo-1,2,3-benzotriazin-4(H)one

25 To a slurry of 1.63 g (0.01 mol) of 3-hydro-1,2,3-benzotriazin-4(3H)one slurry in 7 mL of bromochloromethane was added 1.02 g of triethylamine. The mixture was stirred at 25°C. overnight, diluted with methylene chloride and washed with water. The organic layer was separated, dried and the solvent evaporated off to give 1.65 g of the titled compound as a substantially white solid.

Antimicrobial Activity

The compounds of this invention are useful as antimicrobial additives, and they can be added to industrial products such as paints, inks, adhesives, soaps,

cutting oils, textiles, and paper and pigment slurries and to styrene-butadiene latexes used for paper coatings.

The compounds are also useful as antimicrobial additives in such personal care products as hand creams, lotions, shampoos, and hand soaps. A further advantage of this invention is its cost-effectiveness for applications which need to have an antimicrobial continuously replenished, such as in cooling towers and pulp and paper mills.

As appreciated in the art, not all of the compounds disclosed herein are active at the same concentrations or against the same microbial species. That is, there is some compound-to-compound variation in antimicrobial potency and spectrum of antimicrobial activity.

The present invention is also directed to a method for inhibiting microorganisms which comprises contacting said microorganisms or habitat thereof with a composition containing an antimicrobially effective amount of at least one of the compounds of this invention set forth hereinabove of Formula 1b or 1c.

The antimicrobial compounds of this invention may be added directly to aqueous formulations susceptible to microbial growth, either undiluted or dissolved in inert diluents, such as organic solvents, such as glycols, alcohols, or acetone. They may also be added alone or in combination with other preservatives.

As used herein, the term "microorganism" is meant to refer to bacteria, fungi, viruses, algae, subviral agents and protozoa.

As used herein, the term "antimicrobially-effective amount" refers to that amount of one or a mixture of two or more of the compounds, or of a composition comprising such compound or compounds, of this invention needed to exhibit inhibition of selected microorganisms. Typically, this amount varies from providing 1 part per million (ppm) to 5,000 ppm by weight of the compound to a microbial habitat being contacted with the compound. Such amounts vary depending upon the particular compound tested and microorganism treated. Also, the exact concentration of the compounds to be added in the treatment of industrial and consumer formulations may vary within a product type depending upon the components of the formulation. A preferred effective amount of the compound is from 1 ppm to 500 ppm, more preferably from 1 ppm to 50 ppm by weight, of a microbial habitat.

The term "microbial habitat" refers to a place or type of site where a microorganism naturally or normally lives or grows. Typically, such a microbial habitat

will be an area that comprises a moisture, nutrient, and/or an oxygen source such as, for example, a cooling water tower or an air washing system.

The terms "inhibition", "inhibit" or "inhibiting" refer to the suppression, stasis, kill, or any other interference with the normal life processes of microorganisms that is adverse to such microorganisms, so as to destroy or irreversibly inactivate existing microorganisms and/or prevent or control their future growth and reproduction.

The antimicrobial activity of the compounds of the present invention is demonstrated by the following techniques.

The antimicrobial activity of the compounds of the present invention is set forth as the minimum inhibitory concentration (MIC) for the active compounds and is determined for nine (9) bacteria, using nutrient agar, and seven (7) yeast and fungi, using malt yeast agar. This determination is conducted using a one percent solution of the test compound prepared in a mixture of acetone and water.

Nutrient agar is prepared at pH 6.8, representing a neutral medium, and at pH 8.2, representing an alkaline medium. The nutrient agars are prepared by adding 23 g of nutrient agar to one liter of deionized water. In addition, the alkaline medium is prepared by adjusting a 0.04 M solution of N-[tris-(hydroxymethyl)methyl]-glycine buffered deionized water with concentrated sodium hydroxide to a pH of 8.5.

Malt yeast agar is prepared by adding 3 g yeast extract and 45 g malt agar per liter of deionized water. The specific agar is dispensed in 30 mL aliquots into 25 x 200 mm test tubes, capped and autoclaved for 15 minutes at 115°C.

The test tubes containing the agar are cooled in a water bath until the temperature of the agar is 48°C. Then, an appropriate amount of the one percent solution of the test compound is added (except in the controls where no compound is added) to the respective test tubes so that the final concentrations are 500, 250, 100, 50, 25, 10, 5, 2.5, 1.0 and zero parts per million of the test compound in the agar, thus having a known concentration of test compound dispersed therein. The contents of the test tubes are then transferred to respective petri plates. After drying for 24 hours, the petri plates containing nutrient agar are inoculated with bacteria and those containing malt yeast agar are inoculated with yeast and fungi.

The inoculation with bacteria is accomplished by using the following procedure. Twenty-four hour-cultures of each of the bacteria are prepared by incubating the respective bacteria in tubes containing nutrient broth for 24 hours at

30°C in a shaker. Dilutions of each of the 24 hour-cultures are made so that nine separate suspensions (one for each of the nine test bacteria) are made, each containing 10^8 colony forming units (CFU) per mL of suspension of a particular bacteria. Aliquots of 0.3 mL of each of the bacterial suspensions are used to fill the individual wells of Steer's Replicator. For each microbial suspension, 0.3 mL was used to fill three wells (that is, three wells of 0.3 mL each) so that for the nine different bacteria, 27 wells are filled. The Steer's Replicator is then used to inoculate both the neutral and alkaline pH nutrient agar petri plates.

The inoculated petri plates are incubated at 30°C for 48 hours and then read to determine if the test compound which is incorporated into the agar prevented growth of the respective bacteria.

The inoculation with the yeast and fungi is accomplished as follows. Cultures of yeast and fungi are incubated for seven days on malt yeast agar at 30°C. These cultures are used to prepare suspensions by the following procedure. A suspension of each organism is prepared by adding 10 mL of sterile saline and 10 microliters of octylphenoxy polyethoxy ethanol to the agar slant of yeast or fungi. The sterile saline/octylphenoxy polyethoxy ethanol solution is then agitated with a sterile swab to suspend the microorganism grown on the slant. Each resulting suspension is diluted into sterile saline (1 part suspension to 9 parts sterile saline). Aliquots of these dilutions are placed in individual wells of Steer's Replicator and petri plates inoculated as previously described. The petri plates are incubated at 30°C and read after 48 hours for yeast and 72 hours for fungi.

Table I lists the bacteria, yeast and fungi used in the MIC test described above along with their respective American Type Culture Collection (ATCC) identification numbers.

TABLE I
Organisms used in the Minimum
Inhibitory Concentration Test

Organism	ATCC No.
Bacteria	
Bacillus subtilis (Bs)	8473
Enterobacter aerogenes (Ea)	13048
Escherichia coli (Ec)	11229
Klebsiella pneumoniae (Kp)	8308
Proteus vulgaris (Pv)	881
Pseudomonas aeruginosa (Pa)	10145
Pseudomonas aeruginosa (PRD-10)	15442
Salmonella choleraesuis (Sc)	10708
Staphylococcus aureus (Sa)	6538
Yeast/Fungi	
Aspergillus niger (An)	16404
Candida albicans (Ca)	10231
Penicillium chrysogenum (Pc)	9480
Saccharomyces cerevisiae (Sc)	4105
Trichoderma viride (Tv)	8678
Aureobasidium pullulan (Ap)	16622
Fusarium oxysporum (Fo)	48112

In Tables II and III, the MIC values of the compounds of the present invention as compared to the MIC of a standard commercial preservative (with 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride as the active agent, and referred to in Tables II and III as "STANDARD") are set forth for the bacteria organisms and yeast/fungi organisms which are listed in Table I.

TABLE II
Minimum Inhibitory Concentrations for Test Compounds
in Bacteria Species (in ppm)

Compound (Example No.)	ORGANISMS									
	Bs	Ea	Ec	Kp	Pv	PRD	Pa	Sc	Sa	
STANDARD										
pH 6.8	<10	100	50	25	50	>500	>500	50	25	
pH 8.2	250	500	>500	500	500	>500	>500	>500	500	
(I)	<10	250	50	50	25	250	250	25	25	
pH 8.2	250	500	500	500	500	>500	500	500	500	

TABLE III
Minimum Inhibitory Concentrations for Test Compounds
in Yeast/Fungi Species (in ppm)

Compound (Example No.)	ORGANISMS						
	An	Ca	Pc	Sc	Tv	Ap	Fo
STANDARD	>500	>500	>500	500	>500	>500	>500
I	<1	25	2.5	10	50	2.5	5

Marine Antifouling Activity

The present invention is also directed to a method for inhibiting marine organisms. The term "marine organisms" is meant to include marine animals, such as barnacles, serpulid, bryozoa, oysters and hydroids, and marine plants, such as green algae and brown algae. The method for inhibiting marine organisms comprises contacting a surface exposed to a marine environment in which marine organisms grow with a marine antifouling effective amount of the compound of this invention.

As appreciated by those skilled in the art, not all of the compounds disclosed herein are active at the same concentrations or against the same marine organism species. That is, there may be some compound-to-compound variation in marine antifouling potency and spectrum of marine antifouling activity. Furthermore, the level of a specific compound's marine antifouling activity may be dependent on various factors including the specific materials with which the compound is formulated.

As used herein, the term "marine antifouling effective amount" refers to that amount of one or a mixture of two or more of the compounds of this invention needed to exhibit inhibition of selected marine organisms. Typically, this amount varies from providing 1 weight percent to 30 weight percent of the compound to a marine antifouling composition which is used to treat a surface exposed to a marine environment in which marine organisms live or grow. Such amounts vary depending upon the particular compound tested and marine organism to be treated. Also, the exact concentration of the compounds to be added in the preparation of industrial and consumer formulations may vary within a product type depending upon the components of the formulation.

A composition comprising a marine antifouling effective amount of the compound will also comprise an inert diluent which may be, for example, in the form of a paint. Particularly preferred are those paints having a vinyl resin binder such as, for example, a plasticized polyvinyl chloride or a polyvinyl chloride-polyvinyl acetate type. Preferably, the binders are formulated as latexes or emulsions. In a paint composition,

the compound of the present invention is preferably used in an amount from 1 to 30 weight percent and, most preferably, from 10 to 25 weight percent. In addition to vinyl resin binder paints, epoxy and polyurethane binder paints containing the compound may also be useful. Coatings and films prepared from paints comprising the compound of the present invention typically remain substantially free from build-up of marine organisms for periods ranging from 3 to 12 months, depending upon the concentration of the compound and the thickness of the applied coating or film.

The term "a surface exposed to a marine environment" refers to a surface where a marine organism naturally or normally lives or grows. Typically, such a surface will be an area that is in continual or periodic contact with a marine environment such as an ocean or other body of water. Typical surfaces include, for example, a ship hull.

The marine antifouling activity of the compounds of the present invention is demonstrated by the following techniques.

Test panels are prepared from clear, rigid polyvinyl chloride film that is 0.381×10^{-3} m thick and has one textured surface. The test panels are 0.1524 m by 0.1524 m squares that have 0.00635 m holes punched at corners on 0.127 m centers. A 0.102 square template, with a 0.067 m diameter hole at the center, is attached to the center of the textured surface of the test panels.

The candidate marine antifoulant compound (1.0 g) is stirred into a resinous latex binder (9.0 g). A portion of the compound/binder mixture (1.5 g) is added to the center of the test panel and uniformly spread over the circular area inside the template.

Water is added dropwise as needed to properly spread the compound/binder mixture. The template prevents the compound/binder mixture from spreading beyond the uncovered area. The test panel is allowed to sit for between 10 to 30 minutes until the edge of the spread compound/binder mixture has dried. The template is then removed. The test panel is then allowed to dry for 8 to 12 hours at room temperature.

Two test panels are prepared for each candidate marine antifoulant compound. Two control test panels are also prepared by only treating with the resinous latex binder. One test panel of each candidate marine antifoulant compound is attached over a white background to the topside of an exposure support apparatus. The second test panel is attached over a black background to the underside of the exposure support apparatus. The exposure support apparatus is placed horizontally

0.0254 m under a marine surface with the white background topside facing up. The exposure support apparatus is exposed to the marine environment for both 6 and 10 weeks during which time the control test panels become substantially covered with mature marine organism growth on both the topside and underside exposures.


After being removed from the exposure support apparatus, each test panel is inspected and rated for marine organism growth on both the treated and untreated areas of the test panel. The marine organisms present on the treated and untreated areas are noted. The presence of algae spores and bacterial slime are noted but not included in rating each test panel. The test panels are rated on a scale from 10 (representing completely free of marine organism growth) to 0 (representing completely covered with marine organism growth).

In Table IV, the marine antifouling rating values for 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one is set forth, as well as the ratings for control panels (with no marine antifouling compound and referred to as "Control").

TABLE IV
Marine Antifouling Rating for Test Compounds

Test Compound	Marine Antifouling Ratings					
	Top Panel at indicated time in weeks			Bottom Panel at indicated time in weeks		
	6	10	16	6	10	16
3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one	9	7	8	9	8	7
Control	6	3	-	0	0	-


1. A thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)one compounds corresponding to one of the formulae:



wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C₁-C₇ straight or branched chain alkyl, C₁-C₇ straight or branched chain alkoxy, -CF₃, -OCF₃, -CN, -COOR wherein R is C₁-C₇ straight or branched chain alkyl and n is an integer of from 0-4.

2. The compound of Claim 1 which is 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one.

3. An antimicrobial composition comprising an inert diluent and an antimicrobially-effective amount of a thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)-one compound corresponding to one of the formulae:




wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C₁-C₇ straight or branched chain alkyl, C₁-C₇ straight or branched chain alkoxy, -CF₃, -OCF₃, -CN,

-COOR wherein R is C₁-C₇, straight or branched chain alkyl and n is an integer of from 0-4.

4. The composition of Claim 3 wherein the compound is 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one.

5. A method for inhibiting microorganisms in a microbial habitat comprising contacting said microbial habitat with a composition containing an inert diluent and an antimicrobially-effective amount of an active compound which is a thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)-one compound corresponding to one of the formulae:




wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C₁-C₇, straight or branched chain alkyl, C₁-C₇, straight or branched chain alkoxy, -CF₃, -OCF₃, -CN, -COOR wherein R is C₁-C₇, straight or branched chain alkyl and n is an integer of from 0-4.

6. The method of Claim 5 wherein the active compound is 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one.

7. The method of Claim 5 wherein the active compound is present in the composition in an amount to provide from 1 part per million to 5,000 parts per million by weight of the compound to the microbial habitat.


8. A composition useful in preventing the growth of marine organisms on a surface exposed to a marine environment in which marine organisms grow comprising an inert diluent and a marine antifouling effective amount of a thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)-one compound corresponding to one of the formulae:



wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C₁-C₇ straight or branched chain alkyl, C₁-C₇ straight or branched chain alkoxy, -CF₃, -OCF₃, -CN, -COOR wherein R is C₁-C₇ straight or branched chain alkyl and n is an integer of from 0-4.

9. The composition of Claim 8 wherein the compound is 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one.

10. A method for preventing the growth of marine organisms on a surface exposed to a marine environment in which marine organisms grow comprising contacting said surface with a composition containing an inert diluent and a marine antifouling effective amount of an active material which is a thiocyanomethyloxo(or thiocyanothiomethyl)-1,2,3-benzotriazin-4(3H)-one compound corresponding to one of the formulae:



wherein Z represents oxygen or sulfur; X represents -Cl, -F, -I, C₁-C₇ straight or branched chain alkyl, C₁-C₇ straight or branched chain alkoxy, -CF₃, -OCF₃, -CN, -COOR wherein R is C₁-C₇ straight or branched chain alkyl and n is an integer of from 0-4.

5 11. The method of Claim 10 wherein the compound is 3-thiocyanomethyloxo-1,2,3-benzotriazin-4(3H)one.

 12. The method of Claim 10 wherein the active compound comprises from 1 to 30 weight percent of the composition.