



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/09/10
(87) **Date publication PCT/PCT Publication Date:** 2016/03/17
(85) **Entrée phase nationale/National Entry:** 2017/02/17
(86) **N° demande PCT/PCT Application No.:** US 2015/049461
(87) **N° publication PCT/PCT Publication No.:** 2016/040657
(30) **Priorité/Priority:** 2014/09/10 (US62/048,770)

(51) **Cl.Int./Int.Cl.** **A61K 39/395** (2006.01),
A61K 45/06 (2006.01), **A61P 35/00** (2006.01)
(71) **Demandeur/Applicant:**
THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
US
(72) **Inventeurs/Inventors:**
MCCORMICK, FRANK, US;
WANG, MAN-TZU, US
(74) **Agent:** FETHERSTONHAUGH & CO.

(54) **Titre : CIBLAGE DE VOIES DE SIGNALISATION MEDIEES PAR K-RAS ET DE MALIGNITE PAR DES ANTICORPS ANTI-HLIF**
(54) **Title: TARGETING K-RAS-MEDIATED SIGNALING PATHWAYS AND MALIGNANCY BY ANTI-HLIF ANTIBODIES**




FIG. 5F

(57) **Abrégé/Abstract:**

The present invention provides method of treating a K-Ras-expressing cancer in a subject comprising administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF). Compositions and kits for treating a K-Ras-expressing cancer in a subject are also provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2016/040657 A1(43) International Publication Date
17 March 2016 (17.03.2016)

(51) International Patent Classification:

A61K 39/395 (2006.01) *A61P 35/00* (2006.01)
A61K 45/06 (2006.01)

(21) International Application Number:

PCT/US2015/049461

(22) International Filing Date:

10 September 2015 (10.09.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/048,770 10 September 2014 (10.09.2014) US

(71) Applicant: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, California 94607-5200 (US).(72) Inventors: **MCCORMICK, Frank**; 5 Bernice Street, San Francisco, California 94103 (US). **WANG, Man-Tzu**; 550 Gene Friend Way, Apartment 614, San Francisco, California 94158 (US).(74) Agents: **FLANAGAN, Lisa** et al.; Kilpatrick Townsend & Stockton LLP, Two Embarcadero Center, 8th Floor, San Francisco, California 94111 (US).(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: TARGETING K-RAS-MEDIATED SIGNALING PATHWAYS AND MALIGNANCY BY ANTI-hLIF ANTIBODIES


(57) **Abstract:** The present invention provides method of treating a K-Ras-expressing cancer in a subject comprising administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF). Compositions and kits for treating a K-Ras-expressing cancer in a subject are also provided.

FIG. 5F

WO 2016/040657 A1

WO 2016/040657 A1

Published:

— *with international search report (Art. 21(3))*

TARGETING K-RAS-MEDIATED SIGNALING PATHWAYS AND MALIGNANCY BY ANTI-hLIF ANTIBODIES

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 62/048,770, filed September 10, 2014, the entire content of which is incorporated by reference herein for all purposes.

BACKGROUND OF THE INVENTION

[0002] Pancreatic cancer is a cancer that often has a poor prognosis, even when detected in its early stages. It is estimated that for all stages of pancreatic cancer combined, only 6% of patients survive five years after diagnosis. The most common form of pancreatic cancer, pancreatic ductal adenocarcinoma (PDAC), is known to have an extremely poor prognosis. Although survival time improves for patients who undergo a surgical resection, PDAC frequently is not diagnosed in time for surgical resection to be feasible.

[0003] The oncogene K-Ras is frequently mutated in cancers, such as pancreatic, lung, and colorectal cancers, with activating K-Ras mutations present in over 90% of PDACs. However, to date there have been no successes in developing small molecule inhibitors that directly block K-Ras function and show efficacy in pre-clinical model.

BRIEF SUMMARY OF THE INVENTION

[0004] In one aspect, methods of treating a cancer in a subject are provided. In some embodiments, the method comprises administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF).

[0005] In some embodiments, the cancer is a K-Ras-expressing cancer. In some embodiments, the K-Ras-expressing cancer is a cancer that expresses wild-type K-Ras. In some embodiments, the K-Ras-expressing cancer is a cancer that expresses a mutated K-Ras.

[0006] In some embodiments, the cancer is a pancreatic cancer, a colorectal cancer, or a lung cancer. In some embodiments, the cancer is pancreatic cancer (*e.g.*, pancreatic ductal adenocarcinoma).

[0007] In some embodiments, the agent that antagonizes LIF is an anti-LIF antibody. In some embodiments, the anti-LIF antibody is a monoclonal antibody. In some embodiments, the anti-LIF antibody is an antibody fragment selected from the group consisting of a Fab, a F(ab')₂, and a Fv.

[0008] In some embodiments, the agent that antagonizes LIF is administered orally, intravenously, or intraperitoneally.

[0009] In some embodiments, the agent that antagonizes LIF is administered in combination with a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is a nucleoside analog. In some embodiments, the chemotherapeutic agent is gemcitabine. In some embodiments, the agent that antagonizes LIF and the chemotherapeutic agent are administered concurrently. In some embodiments, the agent that antagonizes LIF is administered and the chemotherapeutic agent are administered sequentially.

[0010] In another aspect, compositions and kits for treating a cancer are provided. In some embodiments, the composition or kit comprises:

an agent that antagonizes leukemia inhibitory factor (LIF); and
a chemotherapeutic agent.

[0011] In some embodiments, the cancer is a K-Ras-expressing cancer. In some embodiments, the K-Ras-expressing cancer is a cancer that expresses wild-type K-Ras. In some embodiments, the K-Ras-expressing cancer is a cancer that expresses a mutated K-Ras. In some embodiments, the cancer is a pancreatic cancer, a colorectal cancer, or a lung cancer. In some embodiments, the cancer is pancreatic cancer (*e.g.*, pancreatic ductal adenocarcinoma).

[0012] In some embodiments, the chemotherapeutic agent is a nucleoside analog. In some embodiments, the chemotherapeutic agent is gemcitabine.

[0013] In another aspect, compositions comprising an agent that antagonizes LIF for use in treating a cancer are provided. In some embodiments, the cancer is a K-Ras-expressing cancer (*e.g.*, a cancer that expresses wild-type K-Ras or a cancer that expresses a mutated K-Ras). In some embodiments, the cancer is a pancreatic cancer, a colorectal cancer, or a lung

cancer. In some embodiments, the composition comprising an agent that antagonizes LIF is used in combination with a chemotherapeutic agent. In some embodiments, the composition comprising an agent that antagonizes LIF further comprises a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is gemcitabine.

5 [0014] In still another aspect, the use of a composition comprising an agent that antagonizes LIF for the manufacture of a medicament for the treatment of a cancer is provided. In some embodiments, the cancer is a K-Ras-expressing cancer (e.g., a cancer that expresses wild-type K-Ras or a cancer that expresses a mutated K-Ras). In some
10 embodiments, the cancer is a pancreatic cancer, a colorectal cancer, or a lung cancer. In some embodiments, the composition comprising an agent that antagonizes LIF further comprises a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is gemcitabine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] **Figure 1. LIF-pSTAT3 signaling is regulated by oncogenic K-Ras in**
15 **pancreatic cancer.** (A) Microarray analysis (affymetrix Gene ST1.0) revealed that LIF was significantly unregulated in K-Ras^{V12}-transformed NIH/3T3 cells when compared to vector control or H-Ras^{V12}-transformed one. (N=3). (B) qPCR (left panel) and Western blotting analysis (right panel) confirmed the elevated LIF expressions for mRNA and protein in K-Ras^{V12}-transformed NIH/3T3 cells. (Right panel) The increased LIF expression referred to
20 enhanced phosphorylated levels of STAT3 in K-Ras^{V12}-transformed. (N=3 in qPCR analysis; ** P <0.01; *** P<0.001). (C) qPCR analysis suggested that mutant K-Ras driven mouse pancreatic cancers had higher LIF expression when compared to mutant B-Raf induced mouse pancreatic tumors. (N=3; ** P<0.01). (D) LIF/LIFR expressions in pancreatic cancer tissues and other types of cancers according to K-Ras mutation status. (E) LIF expressions in
25 established pancreatic cancer cell lines are uncorrelated to specific K-Ras mutation isoforms. (F) Knock-down LIF by shRNA suppressed sphere forming ability in oncogenic K-Ras driven mPCACs (N=8, *** P <0.0001). (G) In two different lines of oncogenic K-Ras driven mPCACs, knock-down of LIF dramatically reduced the recurrent colony forming efficiency post-5FU treatments when compared to control cells. (H) Knock-down of LIF reduced the
30 formation of macro-metastatic spleen lesions and increased media survival time in oncogenic K-Ras driven mPDACs.

[0016] Figure 2. LIF-pSTAT3 signaling is regulated by oncogenic K-Ras in pancreatic**cancer.** (A-E) LIF expression levels are regulated by K-Ras in pancreatic cancer cell lines.

(A) PANC2.13 with knock-down expression of K-Ras showed reduced expressions of LIF and phospho-STAT3 from Western blot. (B) PANC1.0 and (C) PANC2.03 with knock-down

5 expression of K-Ras showed reduced expressions of LIF and phospho-STAT3 at mRNA levels. (N=3; * P <0.05; *** P<0.001). (D) CaPanI and (E) HcG25 with knock-down

expression of K-Ras showed reduced expressions of LIF and phospho-STAT3 at mRNA levels. (N=3; ** P <0.01; *** P<0.001; **** P<0.0001). (F) LIF ELISA revealed that human

pancreatic cancer cell lines with knock-down expression of K-Ras secreted significantly

10 decreased LIF in culture media in the comparison to control cells. (N=4; * P<0.05; ****

P<0.00001). (G) Western blot suggested that the pancreatic cancer cells, with knock-down expression of K-Ras and sequentially down-regulated expression of LIF, showed decreased

phospho-STAT3 levels. (H) phospho-STAT luciferase reporter assays suggested that the pancreatic cancer cells with knock-down expression of K-Ras had significantly decreased

15 STAT3 transcriptional activity. (N=3; ** P<0.01; *** P<0.001).

[0017] Figure 3. LIF plays important roles in human pancreatic cancer**growth/initiation.** (A) Western blot to confirm the knock-down efficiency of shRNA

targeting human LIF in human pancreatic cancer cell lines. (B) Tumor free survival curve of PANC2.03 in subcutaneous xenograft model suggested that the cancer cells with knock-down

20 expression of LIF possessed dramatically reduced tumor-initiating ability, when compared to control cells. (N=6). (C) The pancreatic tumors in subcutaneous xenograft with knock-down

expression of LIF grew in a significantly slow rate when compared to control tumors. (N=6; *

P<0.01). (D) Knock-down of LIF expression reduced tumor initiation rate in the PANC1-driven tumors in orthotopic model. (N=4).

[0018] Figure 4. LIF is required for resistance to gemcitabine treatments in pancreatic**cancers.** An MTS ((3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-

sulfophenyl)-2H-tetrazolium) assay, which is a colorimetric assay for assessing cell viability, suggested that knock-down of LIF sensitized PANC2.03 cells to gemcitabine treatment when

compared to control cells. (N=6) Vehicle treatment (DMSO) was used for normalization.

[0019] Figure 5. LIF neutralizing antibody prevents tumor initiation and improves**therapeutic efficacy of gemcitabine in pancreatic cancer cells.** (A) LIF ELISA with anti-

hLIF antibody competition assay confirmed the neutralizing abilities of our target antibodies.

(B) 50,000 cells of PANC2.03 were subcutaneously injected into nude mice for the tumor initiation assay. The anti-LIF AB (clone D25.1.4) was given before the tumor inoculation at 10 mg/kg. The treatment was given three times a week. As shown, LIF antibodies dramatically prevent tumor initiation. (C) In drug sensitization assay, 0.2×10^6 cells of PANC2.03 cells were first subcutaneously injected into nude mice. The tumors were formed within 14 days post-inoculation. The mice were randomly separated into 4 different groups: control, Ab only, gemcitabine only, and combination. The combination treatment of gemcitabine and LIF Ab caused the complete regressions in 8 out of 10 tumors, whereas control IgG, LIF antibody (D25.1.4) alone, or gemcitabine alone did not lead to the tumor regressions. (D) Tumor volume changing curve in the PANC2.03 subcutaneous tumors with the treatment model of (C). (E) The combination treatment of gemcitabine and LIF Ab dramatically reduced the tumor proliferation rate, whereas the tumors treated with gemcitabine alone still had a positive growth rate. Tumor proliferation rate = (Tumor vol on the later day – Tumor vol on the initial day)/Tumor vol on the initial day * 100. (F) Fold change on tumor volumes suggested that the combination treatment of gemcitabine and LIF Ab dramatically reduced the tumor proliferation rate, whereas the tumors treated with gemcitabine alone still had a positive growth rate.

[0020] Figure 6. LIF expression is enriched in multiple types of cancers. (A-I) The online software Oncomine™ (Invitrogen) was used to analyze different published datasets to determine LIF expression levels for multiple types of cancers as compared to normal tissues. (A-B) LIF expression in TCGA colorectal cancer dataset. (C) LIF expression in D'Errico gastric cancer dataset. (D) LIF expression in Wang gastric cancer dataset. (E) LIF expression in Bredel brain cancer dataset. (F) LIF expression in Barretina cell line dataset. LIF expression was found to be enriched in pancreatic cancer. (G) LIF expression in Garnett cell line dataset. LIF expression was found to be enriched in pancreatic cancer. (H) LIF expression in Pei pancreatic cancer dataset. (I) LIF expression in Garnett cell line dataset. LIF expression at mRNA was significantly enhanced in the established cancer cell lines with mutant K-Ras when compared to cancer cell lines with wild-type K-Ras expression.

[0021] Figure 7. LIF expression at mRNA is dramatically decreased in chemotherapy-sensitive cancers. (A-I) The online software Oncomine™ (Invitrogen) was used to analyze different published datasets with gene profiles of chemotherapy-sensitive and –resistant tumor specimens to determine LIF expression levels. (A) LIF expression in Garnett cell line dataset (cytarabine-resistant and cytarabine-sensitive brain and CNS cancer cell lines). (B)

LIF expression in Garnett cell line dataset (vorinostat-resistant and vorinostat-sensitive multi-cancer cell lines). (C) LIF expression in Garnett cell line dataset (AZD8055-resistant and AZD8055-sensitive brain and CNS cancer cell lines). (D) LIF expression in Garnett cell line dataset (tretinoin-resistant and tretinoin-sensitive brain and CNS cancer cell lines).

5

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

[0022] The present invention is based in part on the surprising discovery that leukemia inhibitor factor (LIF), a stem cell and STAT3 regulated chemokine that is highly expressed in human pancreatic cancer cell lines, is regulated by oncogenic K-Ras. Without being bound to a particular theory, it is believed that LIF acts as a downstream effector essential for K-Ras-driven pancreatic cancer by regulating the stemness of pancreatic cancer cells with activated K-Ras.

[0023] Accordingly, in one aspect the invention provides methods of treating a cancer, such as a cancer that expresses wild-type K-Ras or a cancer that expresses a mutated K-Ras, in a subject by administering a therapeutic amount of an agent that antagonizes LIF. In another aspect, the invention also provides compositions and kits for treating a cancer, such as a K-Ras-expressing cancer, comprising an agent that antagonizes LIF, optionally in combination with a chemotherapeutic agent.

II. Definitions

[0024] As used herein, the term "K-Ras" refers to "Kirsten rat sarcoma viral oncogene homolog." The protein encoded by the K-Ras gene is a small GTPase that functions in intracellular signal transduction. Human K-Ras gene and protein sequences are set forth in, *e.g.*, Genbank Accession Nos. M54968.1 and AAB414942.1. Some common K-Ras genes and proteins found in human cancers contain mutations at codon 12, codon, codon 61, codon 146, and/or other concurrent sites. Non-limiting examples of K-Ras mutations include mutations at codon 5 (*e.g.*, K5E), codon 9 (*e.g.*, V9I), codon 12 (*e.g.*, G12A, G12C, G12D, G12F, G12R, G12S, G12V, G12Y), codon 13 (*e.g.*, G13C, G13D, G13V), codon 14 (*e.g.*, V14I, V14L), codon 18 (*e.g.*, A18D), codon 19 (*e.g.*, L19F), codon 22 (*e.g.*, Q22K), codon 23 (*e.g.*, L23R), codon 24 (*e.g.*, I24N), codon 26 (*e.g.*, N26K), codon 33 (*e.g.*, D33E), codon 36 (*e.g.*, I36L, I36M), codon 57 (*e.g.*, D57N), codon 59 (*e.g.*, A59E, A59G, A59T), codon 61

(*e.g.*, Q61H, Q61K, Q61L, Q61R), codon 62 (*e.g.*, E62G, E62K), codon 63 (*e.g.*, E63K), codon 64 (*e.g.*, Y64D, Y64H, Y64N), codon 68 (*e.g.*, R68S), codon 74 (*e.g.*, T74P), codon 92 (*e.g.*, D92Y), codon 97 (*e.g.*, R97I), codon 110 (*e.g.*, P110H, P110S), codon 117 (*e.g.*, K117N), codon 118 (*e.g.*, C118S), codon 119 (*e.g.*, D119N), codon 135 (*e.g.*, R135T), codon 138 (*e.g.*, G138V), codon 140 (*e.g.*, P140H), codon 146 (*e.g.*, A146T, A146V), codon 147 (*e.g.*, K147N), codon 153 (*e.g.*, D153N), codon 156 (*e.g.*, F156L), codon 160 (*e.g.*, V160A), codon 164 (*e.g.*, R164Q), codon 171 (*e.g.*, I171M), codon 176 (*e.g.*, K176Q), codon 185 (*e.g.*, C185R, C185S), and codon 188 (*e.g.*, M188V).

[0025] A "K-Ras-expressing cancer" refers to a cancer that has a detectable level of expression of K-Ras (either wild-type or its mutant forms). In some embodiments, a cancer has a detectable level of expression when at least 0.1% of cells in the cancer tissue sample are positive for K-Ras activation (*e.g.*, wild-type K-Ras or a K-Ras activating mutation at codon 12, codon 13, codon 61, and/or other codons). In some embodiments, the cancer has a detectable level of expression of wild-type K-Ras. In some embodiments, the cancer has a detectable level of expression of a mutated K-Ras. In some embodiments, a K-Ras-expressing cancer has a level of expression of K-Ras (*e.g.*, wild-type K-Ras or mutated K-Ras) that is at least 5%, 10%, 20%, 30%, 40%, 50%, 75%, 100%, 150%, or 200% greater than the level of K-Ras expression in a control (*e.g.*, a non-diseased cell or tissue that does not express K-Ras, such as normal human peripheral lymphocytes).

[0026] The term "cancer" refers to a disease characterized by the uncontrolled growth of aberrant cells. The term includes all known cancers and neoplastic conditions, whether characterized as malignant, benign, soft tissue, or solid, and cancers of all stages and grades including pre- and post-metastatic cancers. Examples of different types of cancer include, but are not limited to, digestive and gastrointestinal cancers such as gastric cancer (*e.g.*, stomach cancer), colorectal cancer, gastrointestinal stromal tumors, gastrointestinal carcinoid tumors, colon cancer, rectal cancer, anal cancer, bile duct cancer, small intestine cancer, and esophageal cancer; breast cancer; lung cancer; gallbladder cancer; liver cancer; pancreatic cancer; appendix cancer; prostate cancer, ovarian cancer; renal cancer; cancer of the central nervous system; skin cancer (*e.g.*, melanoma); lymphomas; gliomas; choriocarcinomas; head and neck cancers; osteogenic sarcomas; and blood cancers. As used herein, a "tumor" comprises one or more cancerous cells. In some embodiments, the cancer is pancreatic cancer.

[0027] "Leukemia inhibitory factor (LIF)" refers to an interleukin class 6 cytokine that inhibits cell differentiation. Human LIF gene and protein sequences are set forth in, *e.g.*, Genbank Accession Nos. AK315310 and AAC05174.

[0028] An "agent that antagonizes leukemia inhibitory factor" or "agent that antagonizes LIF" is any agent that inhibits, inactivates, decreases, blocks, or downregulates the expression or activity of LIF. In some embodiments, an agent antagonizes LIF if it decreases the expression or activity of LIF in a biological sample (*e.g.*, cell or tissue) contacted with the agent by at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more relative to a control sample (*e.g.*, the biological sample prior to the contacting). In some
 5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995
 1000
 1005
 1010
 1015
 1020
 1025
 1030
 1035
 1040
 1045
 1050
 1055
 1060
 1065
 1070
 1075
 1080
 1085
 1090
 1095
 1100
 1105
 1110
 1115
 1120
 1125
 1130
 1135
 1140
 1145
 1150
 1155
 1160
 1165
 1170
 1175
 1180
 1185
 1190
 1195
 1200
 1205
 1210
 1215
 1220
 1225
 1230
 1235
 1240
 1245
 1250
 1255
 1260
 1265
 1270
 1275
 1280
 1285
 1290
 1295
 1300
 1305
 1310
 1315
 1320
 1325
 1330
 1335
 1340
 1345
 1350
 1355
 1360
 1365
 1370
 1375
 1380
 1385
 1390
 1395
 1400
 1405
 1410
 1415
 1420
 1425
 1430
 1435
 1440
 1445
 1450
 1455
 1460
 1465
 1470
 1475
 1480
 1485
 1490
 1495
 1500
 1505
 1510
 1515
 1520
 1525
 1530
 1535
 1540
 1545
 1550
 1555
 1560
 1565
 1570
 1575
 1580
 1585
 1590
 1595
 1600
 1605
 1610
 1615
 1620
 1625
 1630
 1635
 1640
 1645
 1650
 1655
 1660
 1665
 1670
 1675
 1680
 1685
 1690
 1695
 1700
 1705
 1710
 1715
 1720
 1725
 1730
 1735
 1740
 1745
 1750
 1755
 1760
 1765
 1770
 1775
 1780
 1785
 1790
 1795
 1800
 1805
 1810
 1815
 1820
 1825
 1830
 1835
 1840
 1845
 1850
 1855
 1860
 1865
 1870
 1875
 1880
 1885
 1890
 1895
 1900
 1905
 1910
 1915
 1920
 1925
 1930
 1935
 1940
 1945
 1950
 1955
 1960
 1965
 1970
 1975
 1980
 1985
 1990
 1995
 2000
 2005
 2010
 2015
 2020
 2025
 2030
 2035
 2040
 2045
 2050
 2055
 2060
 2065
 2070
 2075
 2080
 2085
 2090
 2095
 2100
 2105
 2110
 2115
 2120
 2125
 2130
 2135
 2140
 2145
 2150
 2155
 2160
 2165
 2170
 2175
 2180
 2185
 2190
 2195
 2200
 2205
 2210
 2215
 2220
 2225
 2230
 2235
 2240
 2245
 2250
 2255
 2260
 2265
 2270
 2275
 2280
 2285
 2290
 2295
 2300
 2305
 2310
 2315
 2320
 2325
 2330
 2335
 2340
 2345
 2350
 2355
 2360
 2365
 2370
 2375
 2380
 2385
 2390
 2395
 2400
 2405
 2410
 2415
 2420
 2425
 2430
 2435
 2440
 2445
 2450
 2455
 2460
 2465
 2470
 2475
 2480
 2485
 2490
 2495
 2500
 2505
 2510
 2515
 2520
 2525
 2530
 2535
 2540
 2545
 2550
 2555
 2560
 2565
 2570
 2575
 2580
 2585
 2590
 2595
 2600
 2605
 2610
 2615
 2620
 2625
 2630
 2635
 2640
 2645
 2650
 2655
 2660
 2665
 2670
 2675
 2680
 2685
 2690
 2695
 2700
 2705
 2710
 2715
 2720
 2725
 2730
 2735
 2740
 2745
 2750
 2755
 2760
 2765
 2770
 2775
 2780
 2785
 2790
 2795
 2800
 2805
 2810
 2815
 2820
 2825
 2830
 2835
 2840
 2845
 2850
 2855
 2860
 2865
 2870
 2875
 2880
 2885
 2890
 2895
 2900
 2905
 2910
 2915
 2920
 2925
 2930
 2935
 2940
 2945
 2950
 2955
 2960
 2965
 2970
 2975
 2980
 2985
 2990
 2995
 3000
 3005
 3010
 3015
 3020
 3025
 3030
 3035
 3040
 3045
 3050
 3055
 3060
 3065
 3070
 3075
 3080
 3085
 3090
 3095
 3100
 3105
 3110
 3115
 3120
 3125
 3130
 3135
 3140
 3145
 3150
 3155
 3160
 3165
 3170
 3175
 3180
 3185
 3190
 3195
 3200
 3205
 3210
 3215
 3220
 3225
 3230
 3235
 3240
 3245
 3250
 3255
 3260
 3265
 3270
 3275
 3280
 3285
 3290
 3295
 3300
 3305
 3310
 3315
 3320
 3325
 3330
 3335
 3340
 3345
 3350
 3355
 3360
 3365
 3370
 3375
 3380
 3385
 3390
 3395
 3400
 3405
 3410
 3415
 3420
 3425
 3430
 3435
 3440
 3445
 3450
 3455
 3460
 3465
 3470
 3475
 3480
 3485
 3490
 3495
 3500
 3505
 3510
 3515
 3520
 3525
 3530
 3535
 3540
 3545
 3550
 3555
 3560
 3565
 3570
 3575
 3580
 3585
 3590
 3595
 3600
 3605
 3610
 3615
 3620
 3625
 3630
 3635
 3640
 3645
 3650
 3655
 3660
 3665
 3670
 3675
 3680
 3685
 3690
 3695
 3700
 3705
 3710
 3715
 3720
 3725
 3730
 3735
 3740
 3745
 3750
 3755
 3760
 3765
 3770
 3775
 3780
 3785
 3790
 3795
 3800
 3805
 3810
 3815
 3820
 3825
 3830
 3835
 3840
 3845
 3850
 3855
 3860
 3865
 3870
 3875
 3880
 3885
 3890
 3895
 3900
 3905
 3910
 3915
 3920
 3925
 3930
 3935
 3940
 3945
 3950
 3955
 3960
 3965
 3970
 3975
 3980
 3985
 3990
 3995
 4000
 4005
 4010
 4015
 4020
 4025
 4030
 4035
 4040
 4045
 4050
 4055
 4060
 4065
 4070
 4075
 4080
 4085
 4090
 4095
 4100
 4105
 4110
 4115
 4120
 4125
 4130
 4135
 4140
 4145
 4150
 4155
 4160
 4165
 4170
 4175
 4180
 4185
 4190
 4195
 4200
 4205
 4210
 4215
 4220
 4225
 4230
 4235
 4240
 4245
 4250
 4255
 4260
 4265
 4270
 4275
 4280
 4285
 4290
 4295
 4300
 4305
 4310
 4315
 4320
 4325
 4330
 4335
 4340
 4345
 4350
 4355
 4360
 4365
 4370
 4375
 4380
 4385
 4390
 4395
 4400
 4405
 4410
 4415
 4420
 4425
 4430
 4435
 4440
 4445
 4450
 4455
 4460
 4465
 4470
 4475
 4480
 4485
 4490
 4495
 4500
 4505
 4510
 4515
 4520
 4525
 4530
 4535
 4540
 4545
 4550
 4555
 4560
 4565
 4570
 4575
 4580
 4585
 4590
 4595
 4600
 4605
 4610
 4615
 4620
 4625
 4630
 4635
 4640
 4645
 4650
 4655
 4660
 4665
 4670
 4675
 4680
 4685
 4690
 4695
 4700
 4705
 4710
 4715
 4720
 4725
 4730
 4735
 4740
 4745
 4750
 4755
 4760
 4765
 4770
 4775
 4780
 4785
 4790
 4795
 4800
 4805
 4810
 4815
 4820
 4825
 4830
 4835
 4840
 4845
 4850
 4855
 4860
 4865
 4870
 4875
 4880
 4885
 4890
 4895
 4900
 4905
 4910
 4915
 4920
 4925
 4930
 4935
 4940
 4945
 4950
 4955
 4960
 4965
 4970
 4975
 4980
 4985
 4990
 4995
 5000
 5005
 5010
 5015
 5020
 5025
 5030
 5035
 5040
 5045
 5050
 5055
 5060
 5065
 5070
 5075
 5080
 5085
 5090
 5095
 5100
 5105
 5110
 5115
 5120
 5125
 5130
 5135
 5140
 5145
 5150
 5155
 5160
 5165
 5170
 5175
 5180
 5185
 5190
 5195
 5200
 5205
 5210
 5215
 5220
 5225
 5230
 5235
 5240
 5245
 5250
 5255
 5260
 5265
 5270
 5275
 5280
 5285
 5290
 5295
 5300
 5305
 5310
 5315
 5320
 5325
 5330
 5335
 5340
 5345
 5350
 5355
 5360
 5365
 5370
 5375
 5380
 5385
 5390
 5395
 5400
 5405
 5410
 5415
 5420
 5425
 5430
 5435
 5440
 5445
 5450
 5455
 5460
 5465
 5470
 5475
 5480
 5485
 5490
 5495
 5500
 5505
 5510
 5515
 5520
 5525
 5530
 5535
 5540
 5545
 5550
 5555
 5560
 5565
 5570
 5575
 5580
 5585
 5590
 5595
 5600
 5605
 5610
 5615
 5620
 5625
 5630
 5635
 5640
 5645
 5650
 5655
 5660
 5665
 5670
 5675
 5680
 5685
 5690
 5695
 5700
 5705
 5710
 5715
 5720
 5725
 5730
 5735
 5740
 5745
 5750
 5755
 5760
 5765
 5770
 5775
 5780
 5785
 5790
 5795
 5800
 5805
 5810
 5815
 5820
 5825
 5830
 5835
 5840
 5845
 5850
 5855
 5860
 5865
 5870
 5875
 5880
 5885
 5890
 5895
 5900
 5905
 5910
 5915
 5920
 5925
 5930
 5935
 5940
 5945
 5950
 5955
 5960
 5965
 5970
 5975
 5980
 5985
 5990
 5995
 6000
 6005
 6010
 6015
 6020
 6025
 6030
 6035
 6040
 6045
 6050
 6055
 6060
 6065
 6070
 6075
 6080
 6085
 6090
 6095
 6100
 6105
 6110
 6115
 6120
 6125
 6130
 6135
 6140
 6145
 6150
 6155
 6160
 6165
 6170
 6175
 6180
 6185
 6190
 6195
 6200
 6205
 6210
 6215
 6220
 6225
 6230
 6235
 6240
 6245
 6250
 6255
 6260
 6265
 6270
 6275
 6280
 6285
 6290
 6295
 6300
 6305
 6310
 6315
 6320
 6325
 6330
 6335
 6340
 6345
 6350
 6355
 6360
 6365
 6370
 6375
 6380
 6385
 6390
 6395
 6400
 6405
 6410
 6415
 6420
 6425
 6430
 6435
 6440
 6445
 6450
 6455
 6460
 6465
 6470
 6475
 6480
 6485
 6490
 6495
 6500
 6505
 6510
 6515
 6520
 6525
 6530
 6535
 6540
 6545
 6550
 6555
 6560
 6565
 6570
 6575
 6580
 6585
 6590
 6595
 6600
 6605
 6610
 6615
 6620
 6625
 6630
 6635
 6640
 6645
 6650
 6655
 6660
 6665
 6670
 6675
 6680
 6685
 6690
 6695
 6700
 6705
 6710
 6715
 6720
 6725
 6730
 6735
 6740
 6745
 6750
 6755
 6

[0032] Examples of antibody functional fragments include, but are not limited to, complete antibody molecules, antibody fragments, such as Fv, single chain Fv (scFv), complementarity determining regions (CDRs), VL (light chain variable region), VH (heavy chain variable region), Fab, F(ab)₂' and any combination of those or any other functional portion of an immunoglobulin peptide capable of binding to target antigen (*see, e.g.,* FUNDAMENTAL IMMUNOLOGY (Paul ed., 4th ed. 2001). As appreciated by one of skill in the art, various antibody fragments can be obtained by a variety of methods, for example, digestion of an intact antibody with an enzyme, such as pepsin; or de novo synthesis. Antibody fragments are often synthesized de novo either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, includes antibody fragments either produced by the modification of whole antibodies, or those synthesized de novo using recombinant DNA methodologies (*e.g.,* single chain Fv) or those identified using phage display libraries (*see, e.g.,* McCafferty *et al.*, (1990) *Nature* 348:552). The term "antibody" also includes bivalent or bispecific molecules, diabodies, triabodies, and tetrabodies. Bivalent and bispecific molecules are described in, *e.g.,* Kostelny *et al.* (1992) *J. Immunol.* 148:1547, Pack and Pluckthun (1992) *Biochemistry* 31:1579, Hollinger *et al.* (1993), *PNAS. USA* 90:6444, Gruber *et al.* (1994) *J Immunol.* :5368, Zhu *et al.* (1997) *Protein Sci.* 6:781, Hu *et al.* (1996) *Cancer Res.* 56:3055, Adams *et al.* (1993) *Cancer Res.* 53:4026, and McCartney, *et al.* (1995) *Protein Eng.* 8:301.

[0033] A "humanized" antibody is an antibody that retains the reactivity of a non-human antibody while being less immunogenic in humans. This can be achieved, for instance, by retaining the non-human CDR regions and replacing the remaining parts of the antibody with their human counterparts. *See, e.g.,* Morrison *et al.*, *PNAS USA*, 81:6851-6855 (1984); Morrison and Oi, *Adv. Immunol.*, 44:65-92 (1988); Verhoeyen *et al.*, *Science*, 239:1534-1536 (1988); Padlan, *Molec. Immun.*, 28:489-498 (1991); Padlan, *Molec. Immun.*, 31(3):169-217 (1994).

[0034] "Single chain Fv (svFv)" or "single chain antibodies" refers to a protein wherein the V_H and the V_L regions of a scFv antibody comprise a single chain which is folded to create an antigen binding site similar to that found in two chain antibodies. Methods of making scFv antibodies have been described in *e.g.,* Ward *et al.*, *Exp Hematol.* (5):660-4 (1993); and Vaughan *et al.*, *Nat Biotechnol.* 14(3):309-14 (1996). Single chain Fv (scFv) antibodies optionally include a peptide linker of no more than 50 amino acids, generally no more than 40 amino acids, preferably no more than 30 amino acids, and more preferably no more than

20 amino acids in length. In some embodiments, the peptide linker is a concatamer of the sequence Gly-Gly-Gly-Gly-Ser, *e.g.*, 2, 3, 4, 5, or 6 such sequences. However, it is to be appreciated that some amino acid substitutions within the linker can be made. For example, a valine can be substituted for a glycine. Additional peptide linkers and their use are well-known in the art. See, *e.g.*, Huston *et al.*, *Proc. Nat'l Acad. Sci. USA* 8:5879 (1988); Bird *et al.*, *Science* 242:4236 (1988); Glockshuber *et al.*, *Biochemistry* 29:1362 (1990); U.S. Patent No. 4,946,778, U.S. Patent No. 5,132,405 and Stemmer *et al.*, *Biotechniques* 14:256-265 (1993).

[0035] The phrase "specifically (or selectively) binds to an antibody", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein (*e.g.*, LIF or a portion thereof) and do not bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised against LIF can be selected to obtain antibodies specifically immunoreactive with that protein and not with other proteins, except for polymorphic variants, *e.g.*, proteins at least 80%, 85%, 90%, 95% or 99% identical to a sequence of interest. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays, Western blots, or immunohistochemistry are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See, Harlow and Lane *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, NY (1988) for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity. Typically, a specific or selective reaction will be at least twice the background signal or noise and more typically more than 10 to 100 times background.

[0036] The terms "polypeptide," "peptide," and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers. As used herein, the terms encompass amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds.

[0037] The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. "Amino acid mimetics" refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

[0038] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0039] As used herein, the terms "nucleic acid" and "polynucleotide" are used interchangeably. Use of the term "polynucleotide" includes oligonucleotides (*i.e.*, short polynucleotides). This term also refers to deoxyribonucleotides, ribonucleotides, and naturally occurring variants, and can also refer to synthetic and/or non-naturally occurring nucleic acids (*i.e.*, comprising nucleic acid analogues or modified backbone residues or linkages), such as, for example and without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs), and the like. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (*see, e.g.*, Batzer *et al.*, *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260:2605-2608 (1985); and Cassol *et al.* (1992); Rossolini *et al.*, *Mol. Cell. Probes* 8:91-98 (1994)).

[0040] A "biological sample" includes blood and blood fractions or products (*e.g.*, serum, plasma, platelets, red blood cells, and the like); sputum or saliva; kidney, lung, liver, heart, brain, nervous tissue, thyroid, eye, skeletal muscle, cartilage, or bone tissue; cultured cells, *e.g.*, primary cultures, explants, and transformed cells, stem cells, stool, urine, *etc.* Such biological samples also include sections of tissues such as biopsy and autopsy samples, and frozen sections taken for histologic purposes. A biological sample is typically obtained from a "subject," *i.e.*, a eukaryotic organism, most preferably a mammal such as a primate, *e.g.*, chimpanzee or human; cow; dog; cat; a rodent, *e.g.*, guinea pig, rat, or mouse; rabbit; or a bird; reptile; or fish.

[0041] A "therapeutic amount" or "therapeutically effective amount" of an agent (*e.g.*, an agent that antagonizes LIF) is an amount of the agent which prevents, alleviates, abates, or reduces the severity of symptoms of a cancer (*e.g.*, a K-Ras-expressing cancer) in a subject.

[0042] The terms "administer," "administered," or "administering" refer to methods of delivering agents, compounds, or compositions to the desired site of biological action. These methods include, but are not limited to, topical delivery, parenteral delivery, intravenous delivery, intradermal delivery, intramuscular delivery, colonic delivery, rectal delivery, or intraperitoneal delivery. Administration techniques that are optionally employed with the agents and methods described herein, include *e.g.*, as discussed in Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, current ed.; Pergamon; and Remington's, *Pharmaceutical Sciences* (current edition), Mack Publishing Co., Easton, PA.

III. Methods of Treating Cancers

[0043] In one aspect, methods for treating or preventing a cancer in a subject are provided. In some embodiments, the method comprises administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF). In some embodiments, the subject is a human, *e.g.*, a human adult or a human child.

[0044] In some embodiments, the cancer is a K-Ras-expressing cancer, *e.g.*, a cancer that expresses or overexpresses wild-type K-Ras or a cancer that expresses a mutated form of K-Ras. In some embodiments, the K-Ras-expressing cancer is a pancreatic cancer, a colorectal cancer, or a lung cancer. In some embodiments, the K-Ras-expressing cancer is a pancreatic cancer, *e.g.*, pancreatic ductal adenocarcinoma. In some embodiments, the method further comprises measuring the level of K-Ras expression in a sample (*e.g.*, a tumor tissue sample)

from the subject. In some embodiments, the method further comprises determining a K-Ras genotype that is expressed in a sample (*e.g.*, a tumor tissue sample) from the subject.

[0045] In some embodiments, the method further comprises:

detecting the level of K-Ras expression in a sample from the subject (*e.g.*, a tumor cell or tumor tissue sample from the subject);

determining whether the level of K-Ras expression in the sample from the subject is greater than the level of K-Ras expression of a control (*e.g.*, a non-diseased cell or tissue that does not express K-Ras, such as normal human peripheral lymphocytes); and

administering an agent that antagonizes LIF to the subject when the level of K-Ras expression in the sample from the subject is greater than the level of K-Ras expression of a control.

[0046] In some embodiments, the cancer is not a K-Ras-expressing or -overexpressing cancer. As a non-limiting example, in some embodiments the cancer is a pancreatic cancer (*e.g.*, a pancreatic ductal adenocarcinoma) that does not express or overexpress K-Ras.

K-Ras-Expressing Cancers

[0047] In some embodiments, the cancer is a cancer that expresses K-Ras at a detectable level. In some embodiments, a cancer has a detectable level of K-Ras expression when at least 0.1% of cells in the cancer tissue sample are positive for K-Ras activation (*e.g.*, wild-type K-Ras or a K-Ras activating mutation at codon 12, codon 13, codon 61, and/or other codons). In some embodiments, the cancer has a detectable level of expression of wild-type K-Ras. In some embodiments, the cancer has a detectable level of expression of a mutated K-Ras. In some embodiments, the K-Ras mutation is an activating mutation at one or more of codon 5 (*e.g.*, K5E), codon 9 (*e.g.*, V9I), codon 12 (*e.g.*, G12A, G12C, G12D, G12F, G12R, G12S, G12V, G12Y), codon 13 (*e.g.*, G13C, G13D, G13V), codon 14 (*e.g.*, V14I, V14L), codon 18 (*e.g.*, A18D), codon 19 (*e.g.*, L19F), codon 22 (*e.g.*, Q22K), codon 23 (*e.g.*, L23R), codon 24 (*e.g.*, I24N), codon 26 (*e.g.*, N26K), codon 33 (*e.g.*, D33E), codon 36 (*e.g.*, I36L, I36M), codon 57 (*e.g.*, D57N), codon 59 (*e.g.*, A59E, A59G, A59T), codon 61 (*e.g.*, Q61H, Q61K, Q61L, Q61R), codon 62 (*e.g.*, E62G, E62K), codon 63 (*e.g.*, E63K), codon 64 (*e.g.*, Y64D, Y64H, Y64N), codon 68 (*e.g.*, R68S), codon 74 (*e.g.*, T74P), codon 92 (*e.g.*, D92Y), codon 97 (*e.g.*, R97I), codon 110 (*e.g.*, P110H, P110S), codon 117 (*e.g.*, K117N), codon 118 (*e.g.*, C118S), codon 119 (*e.g.*, D119N), codon 135 (*e.g.*, R135T), codon 138 (*e.g.*, G138V), codon 140 (*e.g.*, P140H), codon 146 (*e.g.*, A146T, A146V), codon 147 (*e.g.*, K147N), codon

153 (*e.g.*, D153N), codon 156 (*e.g.*, F156L), codon 160 (*e.g.*, V160A), codon 164 (*e.g.*, R164Q), codon 171 (*e.g.*, I171M), codon 176 (*e.g.*, K176Q), codon 185 (*e.g.*, C185R, C185S), and codon 188 (*e.g.*, M188V). In some embodiments, the K-Ras mutation is a mutation at amino acid residue G12 (*e.g.*, a G12C, G12V, G12D, G12A, G12S, G12R, or G12F substitution). In some embodiments, the K-Ras mutation is a mutation at amino acid residue G13 (*e.g.*, a G13C or G13D substitution). In some embodiments, the K-Ras mutation is a mutation at amino acid residue Q61 (*e.g.*, a Q61H or Q61K substitution). In some embodiments, the K-Ras mutation is a mutation at amino acid residue A146 (*e.g.*, an A146T or A146V substitution). In some embodiments, the cancer that expresses wild-type or mutated K-Ras at a detectable level is a pancreatic cancer, a lung cancer, or a colorectal cancer.

[0048] In some embodiments, the cancer is a cancer that overexpresses K-Ras. As used herein a cancer "overexpresses" K-Ras if the level of expression of K-Ras (*e.g.*, wild-type K-Ras or mutated K-Ras) is increased relative to a threshold value or a control sample (*e.g.*, a non-diseased cell or tissue that does not express K-Ras, such as normal human peripheral lymphocytes, or a cancer sample from a subject known to be negative for expression of K-Ras). In some embodiments, a cancer overexpresses K-Ras if the level of expression of K-Ras (*e.g.*, wild-type K-Ras or mutated K-Ras) is at least 10%, 20%, 30%, 40%, 50%, 75%, 100%, 150%, or 200% greater than a threshold value or the level of K-Ras expression in a control sample (*e.g.*, a cancer known to be negative for expression of K-Ras). In some embodiments, a cancer overexpresses K-Ras if the level of expression of K-Ras (*e.g.*, wild-type K-Ras or mutated K-Ras) is at least 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or more relative to a threshold value or to the level of K-Ras expression in a control sample (*e.g.*, a cancer known to be negative for expression of K-Ras). In some embodiments, the cancer that overexpresses wild-type or mutated K-Ras is a pancreatic cancer, a lung cancer, or a colorectal cancer.

[0049] The level of expression of K-Ras in a cancer can be measured according to methods known in the art. In some embodiments, the level of K-Ras gene expression in a cancer is measured. In some embodiments, the level of K-Ras protein expression in a cancer is measured. The level of K-Ras gene or protein expression, or the detection of a K-Ras genotype, can be measured in a biological sample from a subject. In some embodiments, the biological sample comprises a cancer cell (*e.g.*, a cell obtained or derived from a tumor). In some embodiments, the biological sample is a tumor tissue sample.

[0050] The level of K-Ras protein expression can be measured using any of a number of immunoassays known in the art. Immunoassay techniques and protocols are generally described in Price and Newman, "Principles and Practice of Immunoassay," 2nd Edition, Grove's Dictionaries, 1997; and Gosling, "Immunoassays: A Practical Approach," Oxford University Press, 2000. A variety of immunoassay techniques, including competitive and non-competitive immunoassays, can be used (*see, e.g., Self et al., Curr. Opin. Biotechnol.*, 7:60-65 (1996)). The term immunoassay encompasses techniques including, without limitation, enzyme immunoassays (EIA) such as enzyme multiplied immunoassay technique (EMIT), enzyme-linked immunosorbent assay (ELISA), IgM antibody capture ELISA (MAC ELISA), and microparticle enzyme immunoassay (MEIA); capillary electrophoresis immunoassays (CEIA); radioimmunoassays (RIA); immunoradiometric assays (IRMA); immunofluorescence (IF); fluorescence polarization immunoassays (FPIA); and chemiluminescence assays (CL). If desired, such immunoassays can be automated. Immunoassays can also be used in conjunction with laser induced fluorescence (*see, e.g., Schmalzing et al., Electrophoresis*, 18:2184-93 (1997); Bao, *J. Chromatogr. B. Biomed. Sci.*, 699:463-80 (1997)).

[0051] Specific immunological binding of an antibody to a protein (*e.g., K-Ras*) can be detected directly or indirectly. Direct labels include fluorescent or luminescent tags, metals, dyes, radionuclides, and the like, attached to the antibody. An antibody labeled with iodine-125 (^{125}I) can be used. A chemiluminescence assay using a chemiluminescent antibody specific for the protein marker is suitable for sensitive, non-radioactive detection of protein levels. An antibody labeled with fluorochrome is also suitable. Examples of fluorochromes include, without limitation, DAPI, fluorescein, Hoechst 33258, R-phycoerythrin, B-phycoerythrin, R-phycoerythrin, rhodamine, Texas red, and lissamine. Indirect labels include various enzymes well known in the art, such as horseradish peroxidase (HRP), alkaline phosphatase (AP), β -galactosidase, urease, and the like. A horseradish-peroxidase detection system can be used, for example, with the chromogenic substrate tetramethylbenzidine (TMB), which yields a soluble product in the presence of hydrogen peroxide that is detectable at 450 nm. An alkaline phosphatase detection system can be used with the chromogenic substrate p-nitrophenyl phosphate, for example, which yields a soluble product readily detectable at 405 nm. Similarly, a β -galactosidase detection system can be used with the chromogenic substrate o-nitrophenyl- β -D-galactopyranoside (ONPG), which yields a

soluble product detectable at 410 nm. A urease detection system can be used with a substrate such as urea-bromocresol purple (Sigma Immunochemicals; St. Louis, MO).

[0052] A signal from the direct or indirect label can be analyzed, for example, using a spectrophotometer to detect color from a chromogenic substrate; a radiation counter to detect radiation such as a gamma counter for detection of ^{125}I ; or a fluorometer to detect fluorescence in the presence of light of a certain wavelength. For detection of enzyme-linked antibodies, a quantitative analysis can be made using a spectrophotometer such as an EMAX Microplate Reader (Molecular Devices; Menlo Park, CA) in accordance with the manufacturer's instructions. If desired, the assays of the present invention can be automated or performed robotically, and the signal from multiple samples can be detected simultaneously. In some embodiments, the amount of signal can be quantified using an automated high-content imaging system. High-content imaging systems are commercially available (*e.g.*, ImageXpress, Molecular Devices Inc., Sunnyvale, CA).

[0053] Antibodies can be immobilized onto a variety of solid supports, such as magnetic or chromatographic matrix particles, the surface of an assay plate (*e.g.*, microtiter wells), pieces of a solid substrate material or membrane (*e.g.*, plastic, nylon, paper), and the like. An assay strip can be prepared by coating the antibody or a plurality of antibodies in an array on a solid support. This strip can then be dipped into the test sample and processed quickly through washes and detection steps to generate a measurable signal, such as a colored spot.

[0054] Analysis of K-Ras nucleic acid expression levels or K-Ras genotype can be achieved using routine techniques such as Southern analysis, reverse-transcriptase polymerase chain reaction (RT-PCR), or any other methods based on hybridization to a nucleic acid sequence that is complementary to a portion of the coding sequence of interest (*e.g.*, slot blot hybridization) are also within the scope of the present invention. Applicable PCR amplification techniques are described in, *e.g.*, Ausubel *et al.* and Innis *et al.*, *supra*. General nucleic acid hybridization methods are described in Anderson, "Nucleic Acid Hybridization," BIOS Scientific Publishers, 1999. Amplification or hybridization of a plurality of nucleic acid sequences (*e.g.*, genomic DNA, mRNA or cDNA) can also be performed from mRNA or cDNA sequences arranged in a microarray. Microarray methods are generally described in Hardiman, "Microarrays Methods and Applications: Nuts & Bolts," DNA Press, 2003; and Baldi *et al.*, "DNA Microarrays and Gene Expression: From Experiments to Data Analysis and Modeling," Cambridge University Press, 2002.

[0055] Analysis of nucleic acid expression levels or genotype can also be performed using techniques known in the art including, without limitation, microarrays, polymerase chain reaction (PCR)-based analysis, sequence analysis, and electrophoretic analysis. A non-limiting example of a PCR-based analysis includes a Taqman® allelic discrimination assay
5 available from Applied Biosystems. Non-limiting examples of sequence analysis include Maxam-Gilbert sequencing, Sanger sequencing, capillary array DNA sequencing, thermal cycle sequencing (Sears *et al.*, *Biotechniques*, 13:626-633 (1992)), solid-phase sequencing (Zimmerman *et al.*, *Methods Mol. Cell Biol.*, 3:39-42 (1992)), sequencing with mass spectrometry such as matrix-assisted laser desorption/ionization time-of-flight mass
10 spectrometry (MALDI-TOF/MS; Fu *et al.*, *Nat. Biotechnol.*, 16:381-384 (1998)), pyrosequencing (Ronaghi *et al.*, *Science*, 281:363-365 (1998)), and sequencing by hybridization. Chee *et al.*, *Science*, 274:610-614 (1996); Drmanac *et al.*, *Science*, 260:1649-1652 (1993); Drmanac *et al.*, *Nat. Biotechnol.*, 16:54-58 (1998). Non-limiting examples of electrophoretic analysis include slab gel electrophoresis such as agarose or polyacrylamide
15 gel electrophoresis, capillary electrophoresis, and denaturing gradient gel electrophoresis. In some embodiments, methods for detecting nucleic acid variants include, *e.g.*, the INVADER® assay from Third Wave Technologies, Inc., restriction fragment length polymorphism (RFLP) analysis, allele-specific oligonucleotide hybridization, a heteroduplex mobility assay, single strand conformational polymorphism (SSCP) analysis, single-
20 nucleotide primer extension (SNUPE), and pyrosequencing.

[0056] A detectable moiety can be used in the assays described herein. A wide variety of detectable moieties can be used, with the choice of label depending on the sensitivity required, ease of conjugation with the antibody, stability requirements, and available instrumentation and disposal provisions. Suitable detectable moieties include, but are not
25 limited to, radionuclides, fluorescent dyes (*e.g.*, fluorescein, fluorescein isothiocyanate (FITC), Oregon Green™, rhodamine, Texas red, tetra-rhodamine isothiocyanate (TRITC), Cy3, Cy5, *etc.*), fluorescent markers (*e.g.*, green fluorescent protein (GFP), phycoerythrin, *etc.*), autoquenched fluorescent compounds that are activated by tumor-associated proteases, enzymes (*e.g.*, luciferase, horseradish peroxidase, alkaline phosphatase, *etc.*), nanoparticles,
30 biotin, digoxigenin, and the like.

[0057] The analysis can be carried out in a variety of physical formats. For example, the use of microtiter plates or automation could be used to facilitate the processing of large numbers of test samples.

[0058] Alternatively, for detecting the level of protein or nucleic acid expression, antibody or nucleic acid probes can be applied to subject samples immobilized on microscope slides. The resulting antibody staining or *in situ* hybridization pattern can be visualized using any one of a variety of light or fluorescent microscopic methods known in the art.

5 [0059] Analysis of the protein or nucleic acid can also be achieved, for example, by high pressure liquid chromatography (HPLC), alone or in combination with mass spectrometry (*e.g.*, MALDI/MS, MALDI-TOF/MS, tandem MS, *etc.*).

[0060] Methods of determining K-Ras genotype are described in the art. *See, e.g.*, Kramer *et al.*, *Cell Oncol.* 31:161-167 (2009); Chen *et al.*, *J. Chromatogr. A* 1216:5147-5154 (2009);
10 Lamy *et al.*, *Modern Pathology* 24:1090-1100 (2011); Galbiati *et al.*, *PLoS ONE* 8(3):359939 (2013); and WO 2010/048691.

Agents That Antagonize LIF

[0061] In some embodiments, a therapeutic amount of an agent that antagonizes LIF is administered to a subject in need thereof (*e.g.*, a subject having a cancer, *e.g.*, a K-Ras-expressing or -overexpressing cancer). In some embodiments, the agent that antagonizes LIF
15 is a peptide, protein, oligopeptide, circular peptide, peptidomimetic, antibody, polysaccharide, lipid, fatty acid, inhibitory RNA (*e.g.*, siRNA, miRNA, or shRNA), polynucleotide, oligonucleotide, aptamer, small organic molecule, or drug compound. The agent can be either synthetic or naturally-occurring.

20 ***Anti-LIF antibodies***

[0062] In some embodiments, the agent is an anti-LIF antibody. In some embodiments, the anti-LIF antibody is a monoclonal antibody. In some embodiments, the anti-LIF antibody is an antibody fragment such as a Fab, a F(ab')₂, and a Fv.

[0063] In some embodiments, the anti-LIF antibody is a monoclonal antibody produced by
25 the hybridoma cell deposited under American Type Culture Collection Accession Number ATCC HB11074 (Clone D25.1.4), ATCC HB11076 (Clone D3.14.1.), ATCC HB11077 (Clone D4.16.9), or ATCC HB11075 (Clone D62.3.2), or a humanized version thereof. Anti-LIF antibodies and methods of making anti-LIF antibodies are described in US Patent No. 5,654,157 and in WO 2011/124566, each of which is incorporated by reference herein. In
30 some embodiments, the anti-LIF antibody is an antibody that competes with an antibody produced by the hybridoma cell deposited under American Type Culture Collection Accession Number ATCC HB11074 (Clone D25.1.4), ATCC HB11076 (Clone D3.14.1.),

ATCC HB11077 (Clone D4.16.9), or ATCC HB11075 (Clone D62.3.2) for binding to an epitope. In some embodiments, the anti-LIF antibody is an antibody that binds the same epitope as an antibody produced by the hybridoma cell deposited under American Type Culture Collection Accession Number ATCC HB11074 (Clone D25.1.4), ATCC HB11076 (Clone D3.14.1.), ATCC HB11077 (Clone D4.16.9), or ATCC HB11075 (Clone D62.3.2). In some embodiments, the anti-LIF antibody is an antibody that binds to an epitope within the region comprising amino acids 160 to 202 of human LIF.

[0064] For preparing an antibody that antagonizes LIF (*e.g.*, a recombinant or monoclonal antibody), many techniques known in the art can be used. *See, e.g.*, Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985); Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *Antibodies, A Laboratory Manual* (1988); and Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986)).

[0065] The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, *e.g.*, the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chains of monoclonal antibodies can also be made from hybridoma or plasma cells. Alternatively, phage or yeast display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (*see, e.g.*, McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992); Lou *et al.* (2010) *PEDS* 23:311). Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity (*see, e.g.*, Kubly, *Immunology* (3rd ed. 1997)). Techniques for the production of single chain antibodies or recombinant antibodies (U.S. Patent 4,946,778, U.S. Patent No. 4,816,567) can also be adapted to produce antibodies. Antibodies can also be made bispecific, *i.e.*, able to recognize two different antigens (*see, e.g.*, WO 93/08829, Traunecker *et al.*, *EMBO J.* 10:3655-3659 (1991); and Suresh *et al.*, *Methods in Enzymology* 121:210 (1986)). Antibodies can also be heteroconjugates, *e.g.*, two covalently joined antibodies, or immunotoxins (*see, e.g.*, U.S. Patent No. 4,676,980, WO 91/00360; and WO 92/200373).

[0066] Antibodies can be produced using any number of expression systems, including prokaryotic and eukaryotic expression systems. In some embodiments, the expression system is a mammalian cell expression, such as a hybridoma, or a CHO cell expression system.

Many such systems are widely available from commercial suppliers. In embodiments in which an antibody comprises both a V_H and V_L region, the V_H and V_L regions may be expressed using a single vector, *e.g.*, in a di-cistronic expression unit, or under the control of different promoters. In other embodiments, the V_H and V_L region may be expressed using
5 separate vectors. A V_H or V_L region as described herein may optionally comprise a methionine at the N-terminus.

[0067] Methods for humanizing or primatizing non-human antibodies are also known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred
10 to as import residues, which are typically taken from an import variable domain.

Humanization can be essentially performed following the method of Winter and co-workers (*see, e.g.*, Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-327 (1988); Verhoeyen *et al.*, *Science* 239:1534-1536 (1988) and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992)), by substituting rodent CDRs or CDR sequences for the corresponding
15 sequences of a human antibody. Such humanized antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent
20 antibodies. Transgenic mice, or other organisms such as other mammals, can be used to express humanized or human antibodies (*see, e.g.*, U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, Marks *et al.*, *Bio/Technology* 10:779-783 (1992); Lonberg *et al.*, *Nature* 368:856-859 (1994); Morrison, *Nature* 368:812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14:845-51 (1996); Neuberger, *Nature Biotechnology*
25 14:826 (1996); and Lonberg & Huszar, *Intern. Rev. Immunol.* 13:65-93 (1995)).

[0068] As an alternative to humanization, human antibodies can be generated. As a non-limiting example, transgenic animals (*e.g.*, mice) can be produced that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the
30 homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. *See, e.g.*,

Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in Immun.*, 7:33 (1993); and U.S. Patent Nos. 5,591,669, 5,589,369, and 5,545,807.

[0069] In some embodiments, antibody fragments (such as a Fab, a Fab', a F(ab')₂, a scFv, or a dAB) are generated. Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (*see, e.g.*, Morimoto *et al.*, *J. Biochem. Biophys. Meth.*, 24:107-117 (1992); and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly using recombinant host cells. For example, antibody fragments can be isolated from antibody phage libraries. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* cells and chemically coupled to form F(ab')₂ fragments (*see, e.g.*, Carter *et al.*, *BioTechnology*, 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to those skilled in the art. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). *See, e.g.*, PCT Publication No. WO 93/16185; and U.S. Patent Nos. 5,571,894 and 5,587,458. The antibody fragment may also be a linear antibody as described, *e.g.*, in U.S. Patent No. 5,641,870.

[0070] In some embodiments, the antibody or antibody fragment can be conjugated to another molecule, *e.g.*, polyethylene glycol (PEGylation) or serum albumin, to provide an extended half-life *in vivo*. Examples of PEGylation of antibody fragments are provided in Knight *et al. Platelets* 15:409, 2004 (for abciximab); Pedley *et al.*, *Br. J. Cancer* 70:1126, 1994 (for an anti-CEA antibody); Chapman *et al.*, *Nature Biotech.* 17:780, 1999; and Humphreys, *et al.*, *Protein Eng. Des.* 20: 227, 2007).

[0071] An antibody or antibody fragment can be assayed for the ability to neutralize the activity of LIF. Methods of assaying inhibition of LIF activity are known in the art. As a non-limiting example, an assay can be performed to determine if the antibody or antibody fragment neutralizes the activity of LIF in a cell proliferation assay using the murine leukemic cell line DA-1a. *See*, Moreau *et al.*, *Nature* 15:690-692 (1988). Neutralizing antibodies can also be evaluated for their ability to block the binding of mLIF to LIFR in mouse pancreatic cancer cells driven by oncogenic *K-Ras*, for their ability to reduce pancreatic tumor formation, and/or for their ability to improve therapeutic responses of pancreatic tumors to gemcitabine in immuno-competent syngenic animal models.

Other LIF antagonists

[0072] Additional antagonists of LIF can be readily identified according to methods well known to those of skill in the art. In some embodiments, antagonists of LIF can be identified by screening potential antagonists for the ability to compete with LIF for binding to the leukemia inhibitory factor receptor (LIFR). Competition assays are well known in the art. Typically, a competitive binding assay uses a labeled known ligand (*e.g.*, LIF) in order to screen libraries (*e.g.*, compound or peptide libraries) for candidates that bind to the known receptor (*e.g.*, LIFR) with at least as much affinity as the known ligand.

[0073] In some embodiments, antagonists of LIF can be identified by screening potential antagonists for the ability to inhibit LIF bioactivity (*e.g.*, in a cell proliferation assay). Methods of assaying for antagonists of LIF are described in the art, *e.g.*, Fairlie *et al.*, *Biochemistry* 42:13193-13201 (2003); Zhou *et al.*, *J. Endod* 7:819-824 (2011).

[0074] Screening assays can be carried out *in vitro*, such as by using cell-based assays, or *in vivo*, such as by using animal models. In some embodiments, the assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (*e.g.*, in microtiter formats on microtiter plates in robotic assays). The agents screened as potential antagonists of LIF can be small organic molecules, peptides, peptidomimetics, peptoids, proteins, polypeptides, glycoproteins, oligosaccharides, or polynucleotides such as inhibitory RNA (*e.g.*, siRNA, antisense RNA).

[0075] Essentially any chemical compound can be tested for its ability to antagonize LIF. In some embodiments, the agents have a molecular weight of less than 1,500 daltons, and in some cases less than 1,000, 800, 600, 500, or 400 daltons. The relatively small size of the agents can be desirable because smaller molecules have a higher likelihood of having physiochemical properties compatible with good pharmacokinetic characteristics, including oral absorption than agents with higher molecular weight.

[0076] In some embodiments, high throughput screening methods involve providing a combinatorial library containing a large number of potential therapeutic compounds (potential LIF-antagonizing compounds). Such "combinatorial chemical or peptide libraries" can be screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The

compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

[0077] A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (*i.e.*, the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

[0078] The preparation and screening of combinatorial chemical libraries is well known to those of skill in the art (*see, e.g.*, Beeler *et al.*, *Curr Opin Chem Biol.*, 9:277 (2005); and Shang *et al.*, *Curr Opin Chem Biol.*, 9:248 (2005)). Libraries of use in the present invention can be composed of amino acid compounds, nucleic acid compounds, carbohydrates, or small organic compounds. Carbohydrate libraries have been described in, for example, Liang *et al.*, *Science*, 274:1520-1522 (1996); and U.S. Patent No. 5,593,853.

[0079] Representative amino acid compound libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent Nos. 5,010,175; 6,828,422; and 6,844,161; Furka, *Int. J. Pept. Prot. Res.*, 37:487-493 (1991); Houghton *et al.*, *Nature*, 354:84-88 (1991); and Eichler, *Comb Chem High Throughput Screen.*, 8:135 (2005)), peptoids (PCT Publication No. WO 91/19735), encoded peptides (PCT Publication No. WO 93/20242), random bio-oligomers (PCT Publication No. WO 92/00091), vinylogous polypeptides (Hagihara *et al.*, *J. Amer. Chem. Soc.*, 114:6568 (1992)), nonpeptidal peptidomimetics with β -D-glucose scaffolding (Hirschmann *et al.*, *J. Amer. Chem. Soc.*, 114:9217-9218 (1992)), peptide nucleic acid libraries (*see, e.g.*, U.S. Patent No. 5,539,083), antibody libraries (*see, e.g.*, U.S. Patent Nos. 6,635,424 and 6,555,310; PCT Application No. PCT/US96/10287; and Vaughn *et al.*, *Nature Biotechnology*, 14:309-314 (1996)), and peptidyl phosphonates (Campbell *et al.*, *J. Org. Chem.*, 59:658 (1994)).

[0080] Representative nucleic acid compound libraries include, but are not limited to, genomic DNA, cDNA, mRNA, inhibitory RNA (*e.g.*, RNAi, siRNA), and antisense RNA libraries. *See, e.g.*, Ausubel, *Current Protocols in Molecular Biology*, eds. 1987-2005, Wiley Interscience; and Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, 2000, Cold Spring Harbor Laboratory Press. Nucleic acid libraries are described in, for example,

U.S. Patent Nos. 6,706,477; 6,582,914; and 6,573,098. cDNA libraries are described in, for example, U.S. Patent Nos. 6,846,655; 6,841,347; 6,828,098; 6,808,906; 6,623,965; and 6,509,175. RNA libraries, for example, ribozyme, RNA interference, or siRNA libraries, are described in, for example, Downward, *Cell*, 121:813 (2005) and Akashi *et al.*, *Nat. Rev. Mol. Cell Biol.*, 6:413 (2005). Antisense RNA libraries are described in, for example, U.S. Patent Nos. 6,586,180 and 6,518,017.

[0081] Representative small organic molecule libraries include, but are not limited to, diversomers such as hydantoins, benzodiazepines, and dipeptides (Hobbs *et al.*, *Proc. Nat. Acad. Sci. USA*, 90:6909-6913 (1993)); analogous organic syntheses of small compound libraries (Chen *et al.*, *J. Amer. Chem. Soc.*, 116:2661 (1994)); oligocarbamates (Cho *et al.*, *Science*, 261:1303 (1993)); benzodiazepines (*e.g.*, U.S. Patent No. 5,288,514; and Baum, *C&EN*, Jan 18, page 33 (1993)); isoprenoids (*e.g.*, U.S. Patent No. 5,569,588); thiazolidinones and metathiazanones (*e.g.*, U.S. Patent No. 5,549,974); pyrrolidines (*e.g.*, U.S. Patent Nos. 5,525,735 and 5,519,134); morpholino compounds (*e.g.*, U.S. Patent No. 5,506,337); tetracyclic benzimidazoles (*e.g.*, U.S. Patent No. 6,515,122); dihydrobenzpyrans (*e.g.*, U.S. Patent No. 6,790,965); amines (*e.g.*, U.S. Patent No. 6,750,344); phenyl compounds (*e.g.*, U.S. Patent No. 6,740,712); azoles (*e.g.*, U.S. Patent No. 6,683,191); pyridine carboxamides or sulfonamides (*e.g.*, U.S. Patent No. 6,677,452); 2-aminobenzoxazoles (*e.g.*, U.S. Patent No. 6,660,858); isoindoles, isooxyindoles, or isooxyquinolines (*e.g.*, U.S. Patent No. 6,667,406); oxazolidinones (*e.g.*, U.S. Patent No. 6,562,844); and hydroxylamines (*e.g.*, U.S. Patent No. 6,541,276).

[0082] Devices for the preparation of combinatorial libraries are commercially available. See, *e.g.*, 357 MPS and 390 MPS from Advanced Chem. Tech (Louisville, KY), Symphony from Rainin Instruments (Woburn, MA), 433A from Applied Biosystems (Foster City, CA), and 9050 Plus from Millipore (Bedford, MA).

[0083] Agents that are initially identified as antagonizing LIF activity can be further tested to validate the apparent activity. Preferably such studies are conducted with suitable cell-based or animal models. The basic format of such methods involves administering a lead compound identified during an initial screen to an animal that serves as a model and then determining if in fact the activity of LIF is antagonized. The animal models utilized in validation studies generally are mammals of any kind. Specific examples of suitable animals

include, but are not limited to, primates (*e.g.*, chimpanzees, monkeys, and the like) and rodents (*e.g.*, mice, rats, guinea pigs, rabbits, and the like).

Administration and Combination Therapy

[0084] The route of administration of a therapeutic agent (*e.g.*, an agent that antagonizes LIF) can be oral, intraperitoneal, transdermal, subcutaneous, by intravenous or intramuscular injection, by inhalation, topical, intralesional, infusion; liposome-mediated delivery; topical, intrathecal, gingival pocket, rectal, intrabronchial, nasal, transmucosal, intestinal, ocular or otic delivery, or any other methods known in the art. In some embodiments, the agent that antagonizes LIF is administered orally, intravenously, or intraperitoneally.

[0085] In some embodiments, agent that antagonizes LIF is administered at a therapeutically effective amount or dose. A daily dose range of about 0.01 mg/kg to about 500 mg/kg, or about 0.1 mg/kg to about 200 mg/kg, or about 1 mg/kg to about 100 mg/kg, or about 10 mg/kg to about 50 mg/kg, can be used. The dosages, however, may be varied according to several factors, including the chosen route of administration, the formulation of the composition, patient response, the severity of the condition, the subject's weight, and the judgment of the prescribing physician. The dosage can be increased or decreased over time, as required by an individual patient. In certain instances, a patient initially is given a low dose, which is then increased to an efficacious dosage tolerable to the patient. Determination of an effective amount is well within the capability of those skilled in the art.

[0086] In some embodiments, the agent that antagonizes LIF is administered in combination with a second therapeutic agent. In some embodiments, the second therapeutic agent is a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is an alkylating agent (*e.g.*, cyclophosphamide, ifosfamide, chlorambucil, busulfan, melphalan, mechlorethamine, uramustine, thiotepa, nitrosoureas, or temozolomide), an anthracycline (*e.g.*, doxorubicin, adriamycin, daunorubicin, epirubicin, or mitoxantrone), a cytoskeletal disruptor (*e.g.*, paclitaxel or docetaxel), a histone deacetylase inhibitor (*e.g.*, vorinostat or romidepsin), an inhibitor of topoisomerase (*e.g.*, irinotecan, topotecan, amsacrine, etoposide, or teniposide), a kinase inhibitor (*e.g.*, bortezomib, erlotinib, gefitinib, imatinib, vemurafenib, or vismodegib), a nucleoside analog or precursor analog (*e.g.*, azacitidine, azathioprine, capecitabine, cytarabine, fluorouracil, gemcitabine, hydroxyurea, mercaptopurine, methotrexate, or thioguanine), a peptide antibiotic (*e.g.*, actinomycin or bleomycin), a platinum-based agent (*e.g.*, cisplatin, oxaloplatin, or carboplatin), or a plant alkaloid (*e.g.*,

vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, paclitaxel, or docetaxel). In some embodiments, the chemotherapeutic agent is a nucleoside analog. In some embodiments, the chemotherapeutic agent is gemcitabine.

[0087] Co-administered therapeutic agents (*e.g.*, the agent that antagonizes LIF, and a second therapeutic agent as described herein) can be administered together or separately, simultaneously or at different times. When administered, the therapeutic agents independently can be administered once, twice, three, four times daily or more or less often, as needed. In some embodiments, the administered therapeutic agents are administered once daily. In some embodiments, the administered therapeutic agents are administered at the same time or times, for instance as an admixture. In some embodiments, one or more of the therapeutic agents is administered in a sustained-release formulation.

[0088] In some embodiments, the agent that antagonizes LIF and a second therapeutic agent are administered concurrently. In some embodiments, the agent that antagonizes LIF is administered first, for example for about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 days or more prior to administering the second therapeutic agent (*e.g.*, chemotherapeutic agent). In some embodiments, the second therapeutic agent (*e.g.*, chemotherapeutic agent) is administered first, for example for about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 days or more prior to administering the agent that antagonizes LIF.

[0089] In some embodiments, the agent that antagonizes LIF (and optionally a second therapeutic agent, *e.g.*, a chemotherapeutic agent as described herein) is administered to the subject over an extended period of time, *e.g.*, for at least 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350 day or longer.

IV. Compositions and Kits

[0090] In another aspect, compositions and kits for use in treating or preventing a cancer (*e.g.*, a K-Ras-expressing or -overexpressing cancer) in a subject are provided.

[0091] In some embodiments, pharmaceutical compositions comprising an agent that antagonizes LIF for use in administering to a subject having a cancer (*e.g.*, a cancer in which wild-type K-Ras or mutated K-Ras is expressed or overexpressed) are provided. In some embodiments, the agent that antagonizes LIF (*e.g.*, an anti-LIF antibody) is as described in Section III above. In some embodiments, a combination of an agent that antagonizes LIF and

a second therapeutic agent (*e.g.*, a chemotherapeutic agent as described herein) are formulated into pharmaceutical compositions, together or separately, by formulation with appropriate pharmaceutically acceptable carriers or diluents, and can be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, pills, powders, granules, dragees, gels, slurries, ointments, solutions, suppositories, injections, inhalants and aerosols.

[0092] Guidance for preparing formulations for use in the present invention is found in, for example, in *Remington: The Science and Practice of Pharmacy*, 21st Ed., 2006, *supra*; *Martindale: The Complete Drug Reference*, Sweetman, 2005, London: Pharmaceutical Press; Niazi, *Handbook of Pharmaceutical Manufacturing Formulations*, 2004, CRC Press; and Gibson, *Pharmaceutical Preformulation and Formulation: A Practical Guide from Candidate Drug Selection to Commercial Dosage Form*, 2001, Interpharm Press, which are hereby incorporated herein by reference. The pharmaceutical compositions described herein can be manufactured in a manner that is known to those of skill in the art, *i.e.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. The following methods and excipients are merely exemplary and are in no way limiting.

[0093] In some embodiments, an agent that antagonizes LIF (and optionally a second therapeutic agent, *e.g.*, a chemotherapeutic agent as described herein) is prepared for delivery in a sustained-release, controlled release, extended-release, timed-release or delayed-release formulation, for example, in semi-permeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Current extended-release formulations include film-coated tablets, multiparticulate or pellet systems, matrix technologies using hydrophilic or lipophilic materials and wax-based tablets with pore-forming excipients (*see*, for example, Huang, *et al. Drug Dev. Ind. Pharm.* 29:79 (2003); Pearnchob, *et al. Drug Dev. Ind. Pharm.* 29:925 (2003); Maggi, *et al. Eur. J. Pharm. Biopharm.* 55:99 (2003); Khanvilkar, *et al., Drug Dev. Ind. Pharm.* 228:601 (2002); and Schmidt, *et al., Int. J. Pharm.* 216:9 (2001)). Sustained-release delivery systems can, depending on their design, release the compounds over the course of hours or days, for instance, over 4, 6, 8, 10, 12, 16, 20, 24 hours or more. Usually, sustained release formulations can be prepared using naturally-occurring or synthetic polymers, for instance, polymeric vinyl pyrrolidones, such as polyvinyl pyrrolidone (PVP); carboxyvinyl hydrophilic

polymers; hydrophobic and/or hydrophilic hydrocolloids, such as methylcellulose, ethylcellulose, hydroxypropylcellulose, and hydroxypropylmethylcellulose; and carboxypolymethylene.

[0094] The sustained or extended-release formulations can also be prepared using natural ingredients, such as minerals, including titanium dioxide, silicon dioxide, zinc oxide, and clay (see, U.S. Patent 6,638,521, herein incorporated by reference). Exemplary extended release formulations include those described in U.S. Patent Nos. 6,635,680; 6,624,200; 6,613,361; 6,613,358, 6,596,308; 6,589,563; 6,562,375; 6,548,084; 6,541,020; 6,537,579; 6,528,080 and 6,524,621, each of which is hereby incorporated herein by reference. Exemplary controlled release formulations include those described in U.S. Patent Nos. 6,607,751; 6,599,529; 6,569,463; 6,565,883; 6,482,440; 6,403,597; 6,319,919; 6,150,354; 6,080,736; 5,672,356; 5,472,704; 5,445,829; 5,312,817 and 5,296,483, each of which is hereby incorporated herein by reference. Those skilled in the art will readily recognize other applicable sustained release formulations.

[0095] For oral administration, an agent that antagonizes LIF (and optionally a second therapeutic agent, *e.g.*, a chemotherapeutic agent as described herein) can be formulated readily by combining with pharmaceutically acceptable carriers that are well known in the art. Such carriers enable the compounds to be formulated as tablets, pills, dragees, capsules, emulsions, lipophilic and hydrophilic suspensions, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by mixing the compounds with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents can be added, such as a cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[0096] The agent that antagonizes LIF (and optionally a second therapeutic agent, *e.g.*, a chemotherapeutic agent as described herein) can be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. For injection, the compound or

compounds can be formulated into preparations by dissolving, suspending or emulsifying them in an aqueous or nonaqueous solvent, such as vegetable or other similar oils, synthetic aliphatic acid glycerides, esters of higher aliphatic acids or propylene glycol; and if desired, with conventional additives such as solubilizers, isotonic agents, suspending agents, emulsifying agents, stabilizers and preservatives. In some embodiments, compounds can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. Formulations for injection can be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0097] The agent that antagonizes LIF (and optionally a second therapeutic agent, *e.g.*, a chemotherapeutic agent as described herein) can be administered systemically by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. For topical administration, the agents are formulated into ointments, creams, salves, powders and gels. In one embodiment, the transdermal delivery agent can be DMSO. Transdermal delivery systems can include, *e.g.*, patches. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. Exemplary transdermal delivery formulations include those described in U.S. Patent Nos. 6,589,549; 6,544,548; 6,517,864; 6,512,010; 6,465,006; 6,379,696; 6,312,717 and 6,310,177, each of which are hereby incorporated herein by reference.

[0098] In some embodiments, a pharmaceutical composition comprises an acceptable carrier and/or excipients. A pharmaceutically acceptable carrier includes any solvents, dispersion media, or coatings that are physiologically compatible and that preferably does not interfere with or otherwise inhibit the activity of the therapeutic agent. In some embodiments, the carrier is suitable for intravenous, intramuscular, oral, intraperitoneal, transdermal, topical, or subcutaneous administration. Pharmaceutically acceptable carriers can contain one or more physiologically acceptable compound(s) that act, for example, to stabilize the composition or to increase or decrease the absorption of the active agent(s). Physiologically acceptable compounds can include, for example, carbohydrates, such as glucose, sucrose, or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins, compositions that reduce the clearance or hydrolysis

of the active agents, or excipients or other stabilizers and/or buffers. Other pharmaceutically acceptable carriers and their formulations are well-known and generally described in, for example, *Remington: The Science and Practice of Pharmacy*, 21st Edition, Philadelphia, PA. Lippincott Williams & Wilkins, 2005. Various pharmaceutically acceptable excipients are
 5 well-known in the art and can be found in, for example, Handbook of Pharmaceutical Excipients (5th ed., Ed. Rowe *et al.*, Pharmaceutical Press, Washington, D.C.).

[0099] In some embodiments, kits for use in administering to a subject having a cancer (*e.g.*, a cancer in which wild-type K-Ras or mutated K-Ras is expressed or overexpressed) are provided. In some embodiments, the kit comprises:

- 10 an agent that antagonizes leukemia inhibitory factor (LIF); and
 a second therapeutic agent.

[0100] In some embodiments, the agent that antagonizes LIF (*e.g.*, an anti-LIF antibody) is as described in Section III above. In some embodiments, the second therapeutic agent is a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is an alkylating
 15 agent, an anthracycline, a cytoskeletal disruptor, a histone deacetylase inhibitor, an inhibitor of topoisomerase, a kinase inhibitor, a nucleoside analog or precursor analog, a peptide antibiotic, a platinum-based agent, or a plant alkaloid. In some embodiments, the chemotherapeutic agent is a nucleoside analog. In some embodiments, the chemotherapeutic agent is gemcitabine.

20 [0101] In some embodiments, the kits can further comprise instructional materials containing directions (*i.e.*, protocols) for the practice of the methods of this invention (*e.g.*, instructions for using the kit for treating a cancer). While the instructional materials typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this
 25 invention. Such media include, but are not limited to electronic storage media (*e.g.*, magnetic discs, tapes, cartridges, chips), optical media (*e.g.*, CD ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

V. Examples

[0102] The following examples are offered to illustrate, but not to limit, the claimed
 30 invention.

Example 1: Targeting leukemia inhibitory factor (LIF) to eradicate pancreatic cells expressing oncogenic K-Ras

[0103] Activating mutations of K-Ras occur in over 90% of pancreatic cancers, but effective approaches to target oncogenic *K-Ras* have been difficult to develop. Thus, identification of essential factors for *K-Ras*-mediated malignancy may provide an alternative way to block this "undruggable" oncogene. With a high degree of sequence homology as well as common sets of downstream effectors and upstream affecters, the three isoforms of Ras, N-, H- and K-Ras, have long been assumed to be functionally redundant. However, *K-Ras*, not *N-* or *H-Ras*, deficiency in mice leads to embryonic lethality, suggesting *K-Ras* may be required for the functions of stem cells (Koera *et al.*, *Oncogene* 15:1151-1159 (1997)). Cancer stem cells (CSCs), sharing certain similar gene expressing signatures and biological functions with normal stem cells, have been identified in numbers of human malignancies, including pancreatic adenocarcinoma (PADC) (Sampieri and Fodde, *Semin Cancer Biol* 22:187-193 (2012)). Due to their self-renewal, tumor initiation, chemo-resistance, and metastatic properties, CSCs are postulated to underlie treatment failures. Despite the putative role of K-Ras activation in pancreatic carcinogenesis, the roles of oncogenic *K-Ras* in pancreatic CSCs have not been convincingly demonstrated. In a preliminary study, we found that oncogenic *K-Ras*, unlike *H-Ras*, causes CSC-like properties in transformed mouse fibroblast and pancreatic cancer cells (data not shown).

[0104] The signal transducer and activator of transcription 3 (STAT3) is a crucial regulator of stem cell self-renewal, cancer cell survival, and metastasis. The majority of PDAC show constitutive activation of STAT3. In the *Pdx1-Cre; LSL-KRAS^{G12D}* transgenic mice model, loss of STAT3 reduced pancreatic tumor formation and progression, suggesting its potential as a therapeutic target in oncogenic *K-Ras*-induced pancreatic cancers (Corcoran *et al.*, *Cancer Res* 71:5020-5029 (2011)). However, most of the STAT3 inhibitors reported to date have presented limited clinical efficacy. Therefore, alternative approaches are needed to inhibit STAT3 as potential anti-pancreatic cancer therapy.

[0105] Through genome wide expression analysis in *Ras^{V12}*-transformed NIH3T3 cells, we identified leukemia inhibitory factor (LIF), a stem cell regulatory chemokine and STAT3 activator, as a factor markedly unregulated by K-Ras, but not by H-Ras (Fig. 1A). Moreover, oncogenic *K-Ras* induced mouse PDACs showed enhanced LIF expression compared to those induced by oncogenic *B-Raf* (Fig. 1C). Further studies demonstrated LIF is required in *K-Ras*-mediated stemness, including sphere formation ability and drug-resistance, in

pancreatic cancer cells (Fig. 1F-1G). In a syngenic animal model, mice with orthotopic transplantation of K-Ras-driven mPDACs which LIF had been knock-down showed greater probability of survival, and impaired spleen metastasis (Fig. 1H).

[0106] Based on these preliminary data, we hypothesized that LIF plays an essential role in the stemness of pancreatic cancer cells with activated K-Ras. Therefore, LIF represents a novel therapeutic target to eradicate *K-Ras*-driven pancreatic cancers. To test this hypothesis, we validated LIF as a therapeutic target of *K-Ras* driven pancreatic cancer by knocking down LIF via small hairpin RNA in mouse pancreatic cancer cells driven by mutant *K-Ras* (FVB background; LSC-K-Ras^{G12D}; p53^{F/+}, pDX^{CRE}). Knock-down efficiency was confirmed by quantitative PCR and western blotting analysis. Knock-down of K-Ras by shRNA repressed LIF protein expression in multiple human pancreatic cancer lines (Fig. 2A-2E). Additionally, a LIF ELISA revealed that human pancreatic cancer cell lines with knock-down expression of K-Ras secreted significantly decreased LIF in culture media in the comparison to control cells (Fig. 2F).

[0107] Levels and phosphorylation of STAT3 were examined by Western blot and using a STAT3 responsive luciferase assay. Western blot analysis and phospho-STAT luciferase reporter assays indicated that the pancreatic cancer cells, with knock-down expression of K-Ras and sequentially down-regulated expression of LIF, had decreased phospho-STAT3 levels (Fig. 2G) and significantly decreased STAT3 transcriptional activity (Fig. 2H).

[0108] For further validation that LIF-LIFR signaling acts as a potential clinical therapeutic target, LIF was also knocked-down in human pancreatic cancer cell lines with active *K-Ras* mutation via small hairpin RNA (Fig. 3A). As shown in Figure 3B, a tumor-free survival curve of PANC2.03 in subcutaneous xenograft model suggested that the cancer cells with knock-down expression of LIF possessed dramatically reduced tumor-initiating ability, when compared to control cells. Additionally, the pancreatic tumors in subcutaneous xenograft with knock-down expression of LIF grew at a significantly slower rate when compared to control tumors (Fig. 3C). Additionally, knock-down of LIF expression reduced tumor initiation rate in the PANC1-driven tumors in an orthotopic model (Fig. 3D).

[0109] The response to gemcitabine was analyzed in pancreatic cancer cells. As shown in Figure 4, an MTS assay suggested that knock-down of LIF sensitized PANC2.03 cells to gemcitabine treatment when compared to control cells. The effects of LIF-neutralizing antibody were also tested in mouse models. As shown in Figure 5B, anti-LIF antibodies

dramatically prevented tumor initiation when injected into nude mice prior to tumor inoculation. In a drug sensitization assay, PANC2.03 cells were subcutaneously injected into nude mice to form tumors, then the mice were treated with anti-LIF antibody, gemcitabine, or a combination of anti-LIF antibody and gemcitabine. The combination treatment of
5 gemcitabine and anti-LIF antibody caused complete regression in 8 out of 10 tumors, whereas anti-LIF antibody alone or gemcitabine alone did not lead to the tumor regression (Fig. 5C). Additionally, the combination treatment of gemcitabine and anti-LIF antibody dramatically reduced the tumor proliferation rate, whereas the tumors treated with gemcitabine alone still had a positive growth rate (Fig. 5E-5F).

10 [0110] We used the online software, Oncomine™ (Invitrogen), to analyze different published datasets (as indicated in Figure 6A-I). Oncomine™ is an online database, collecting large published genome-wide microarray data. The collected data are examined and published by other research groups. Interestingly, we found that human LIF was overexpressed in different types of cancers, including colon and pancreatic cancers, across
15 multiple microarray platforms and published databases. Furthermore, as shown in Figure 6I, *LIF* expression at mRNA was significantly enhanced in the established cancer cell lines with mutant *K-Ras* when compared to the ones with wild type *K-Ras* expression. These results suggest the potential role of LIF as an oncogene in different types of human cancers, especially K-Ras driven tumors.

20 [0111] We also used Oncomine™ to analyze LIF expression at mRNA in chemotherapy-sensitive and chemotherapy-resistant cancers. Oncomine™ provides some datasets with gene profiles of chemotherapy-sensitive and chemotherapy-resistant tumor specimens. Interestingly, as shown in Figure 7A-D, LIF expression was much higher in the tumors that were resistant to different chemotherapy treatments when compared to chemotherapy-
25 sensitive tumors, suggesting that LIF may play an important role in the resistance of human cancers toward chemotherapies. Therefore, targeting LIF with a neutralizing antibody may re-sensitize tumor cells to conventional chemotherapy treatments.

[0112] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, one of skill in the art will
30 appreciate that certain changes and modifications may be practiced within the scope of the appended claims. In addition, each reference provided herein is incorporated by reference in its entirety to the same extent as if each reference was individually incorporated by reference.

WHAT IS CLAIMED IS:

1. A method of treating a K-Ras-expressing cancer in a subject, the method comprising administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF).
2. The method of claim 1, wherein the K-Ras-expressing cancer is a pancreatic cancer, a colorectal cancer, or a lung cancer.
3. The method of claim 1 or 2, wherein the K-Ras-expressing cancer is a pancreatic cancer.
4. The method of any of claims 1-3, wherein the pancreatic cancer is a pancreatic ductal adenocarcinoma.
5. The method of any of claims 1-4, wherein the agent that antagonizes LIF is an anti-LIF antibody.
6. The method of claim 5, wherein the anti-LIF antibody is a monoclonal antibody.
7. The method of claim 5, wherein the anti-LIF antibody is an antibody fragment selected from the group consisting of a Fab, a F(ab')₂, and a Fv.
8. The method of any of claims 1-7, wherein the agent that antagonizes LIF is administered orally, intravenously, or intraperitoneally.
9. The method of any of claims 1-8, wherein the agent that antagonizes LIF is administered in combination with a chemotherapeutic agent.
10. The method of claim 9, wherein the chemotherapeutic agent is gemcitabine.
11. The method of claim 9 or 10, wherein the agent that antagonizes LIF and the chemotherapeutic agent are administered concurrently.
12. The method of claim 9 or 10, wherein the agent that antagonizes LIF and the chemotherapeutic agent are administered sequentially.

13. A method of treating a pancreatic cancer in a subject, the method comprising administering to the subject a therapeutic amount of an agent that antagonizes leukemia inhibitory factor (LIF).

14. The method of claim 13, wherein the pancreatic cancer is a pancreatic ductal adenocarcinoma.

15. The method of claim 13 or 14, wherein the agent that antagonizes LIF is an anti-LIF antibody.

16. The method of claim 15, wherein the anti-LIF antibody is a monoclonal antibody.

17. The method of claim 15, wherein the anti-LIF antibody is an antibody fragment selected from the group consisting of a Fab, a F(ab')₂, and a Fv.

18. The method of any of claims 13-17, wherein the agent that antagonizes LIF is administered orally, intravenously, or intraperitoneally.

19. The method of any of claims 13-18, wherein the agent that antagonizes LIF is administered in combination with a chemotherapeutic agent.

20. The method of claim 19, wherein the chemotherapeutic agent is gemcitabine.

21. The method of claim 19 or 20, wherein the agent that antagonizes LIF and the chemotherapeutic agent are administered concurrently.

22. The method of claim 19 or 20, wherein the agent that antagonizes LIF and the chemotherapeutic agent are administered sequentially.

23. A kit for treating a K-Ras-expressing cancer, the kit comprising:
an agent that antagonizes leukemia inhibitory factor (LIF); and
a chemotherapeutic agent.

24. The kit of claim 23, wherein the agent that antagonizes LIF is an anti-LIF antibody.

25. The kit of claim 24, wherein the anti-LIF antibody is a monoclonal antibody.
26. The kit of any of claims 23-25, wherein the chemotherapeutic agent is gemcitabine.

Probeld	GeneSymbol	GeneDescription	logFC	AveExpr	P.Value	adj.P.Val	Mean.HRas	Mean.KRas	Mean.Control
10373918	Lif	leukemia inhibitory factor	-2.2975	7.45033	3.43E-08	5.45E-05	6.88805	9.18554	6.27739

FIG. 1A





FIG. 1B



2/39




FIG. 1C




FIG. 1D




FIG. 1E





FIG. 1F





FIG. 1G



	TR30014	F1346978
Sample Size	5	5
Media Survival	24	31
Significance	P=0.0415 *	
Harzard Ratio	2.9654	
95% CI	0.7116-	
	2.3574	

FIG. 1H




FIG. 2A









FIG. 2B

PANC1.0
(G12D)



PANC2.03
(G12D)

FIG. 2C



CaPani
(G12V)

FIG. 2D




FIG. 2E




FIG. 2F






FIG. 2G

15/39




FIG. 2H




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4







FIG. 5A




Recipients with Development of PANC2.03 Tumors (55 th Day)		
	5X10 ⁴ cells	
Control IgG	6/6	
D25.1.4	0/4	
P Value	<0.0001	

FIG. 5B



Recipients with Development of PANC2.03 Tumors (14 th Day)	Recipients with Development of PANC2.03 Tumors (14 th Day)
Before treatments	32 days of treatments
Control IgG 9/9	Control IgG 9/9
D25.1.4 6/7	D25.1.4 7/7
Gemcitabine 10/10	Gemcitabine 10/10
Combination 10/10	Combination 2/10

FIG. 5C



24/39




FIG. 5D




FIG. 5E







FIG. 5F



LIF Expression in TCGA Colorectal
Colon Mucinous Adenocarcinoma vs. Normal

Over-expression Gene Rank: 86 (in top 1%) P-value: 3.49E-14
Reporter: A_23_P155017 t-Test: 11.484
Fold Change: 3.788




Legend		TCGA Colorectal	
1. Colon (19)	2. Rectum (3)	Nature 2012/07/18	237 samples
3. Colon Mucinous Adenocarcinoma (20)		mRNA	20,423 measured genes
		Platform not pre-defined in Oncomine	

FIG. 6A

LIF Expression in TCGA Colorectal
Colon Adenocarcinoma vs. Normal

Over-expression Gene Rank: 130 (in top 1%)
Reporter: NM_002309_2_3647
P-value: 6.16E-27
t-Test: 14.401
Fold Change: 2.422



Legend

- 1. Colon (19)
- 2. Rectum (3)
- 3. Colon Adenocarcinoma (102)

TCGA Colorectal

Nature 2012/07/18
mRNA
Platform not pre-defined in OncoPrint

237 samples
20,423 measured genes

FIG. 6B

LIF Expression in DErrico Gastric
Gastric Intestinal Type Adenocarcinoma vs. Normal

Over-expression Gene Rank: 93 (in top 1%)
Reporter: 205266_at
P-value: 2.32E-13
t-Test: 9.492
Fold Change: 4.128







FIG. 6C



LIF Expression in Wang Gastric
Gastric Cancer vs. Normal

Over-expression Gene Rank: 157 (in top 1%)
Reporter: 205266_at
P-value: 3.41E-5
t-Test: 5.040
Fold Change: 2.525




Legend

Wang Gastric	
1. Gastric Mucosa (12)	27 samples
2. Gastric Tissue (3)	19,574 measured genes
3. Gastric Cancer (12)	
Human Genome U133 Plus 2.0 Array	

FIG. 6D

LIF Expression in Bredel Brain 2
Glioblastoma vs. Normal

Over-expression Gene Rank: 727 (in top 5%)
Reporter: IMAGE:153025
P-value: 5.75E-6
t-Test: 5.305
Fold Change: 3.076




Legend		Bredel Brain 2	
1. Brain (4)	2. Glioblastoma (27)	Cancer Res 2005/10/01	54 samples
		mRNA	14,836 measured genes
		Platform not pre-defined in OncoPrint	

FIG. 6E

32/39

LIF Expression in Barretina CellLine
Cancer Type: Pancreatic Cancer

Over-expression Gene Rank: 238 (in top 2%)
Reporter: 205266_at
P-value: 1.02E-11
t-Test: 8.512
Fold Change: 3.483



Legend

Barretina CellLine

1. Bladder Cancer (21) 2. Brain and CNS Cancer (64) 3. Breast Cancer (56) 4. Colorectal Cancer (56) 5. Esophageal Cancer (25) 6. Gastric Cancer (35) 7. Head and Neck Cancer (41) 8. Kidney Cancer (21) 9. Leukemia (83) 10. Liver Cancer (29) 11. Lung Cancer (166) 12. Lymphoma (61) 13. Melanoma (57) 14. Myeloma (26) 15. Ovarian Cancer (44) 16. Pancreatic Cancer (44) 17. Prostate Cancer (7) 18. Sarcoma (39)


Nature 2012/03/28
mRNA
Human Genome U133 Plus 2.0 Array

917 samples
19,574 measured genes

FIG. 6F

LIF Expression in Garnett CellLine
Cancer Type: Pancreatic Cancer

Over-expression Gene Rank: 205 (in top 2%)
Reporter: 205266_at
P-value: 2.30E-5
t-Test: 5.517
Fold Change: 4.337



Legend

1. Bladder Cancer (16) 2. Brain and CNS Cancer (94) 3. Breast Cancer (39) 4. Cervical Cancer (12) 5. Colorectal Cancer (40) 6. Esophageal Cancer (23) 7. Gastric Cancer (19) 8. Head and Neck Cancer (33) 9. Kidney Cancer (22) 10. Leukemia (66) 11. Liver Cancer (13) 12. Lung Cancer (130) 13. Lymphoma (44) 14. Melanoma (41) 15. Myeloma (9) 16. Ovarian Cancer (22) 17. Pancreatic Cancer (16) 18. Prostate Cancer (5) 19. Sarcoma (46)


Garnett CellLine

Nature 2012/03/28 732 samples
mRNA 12,624 measured genes
Human Genome U133A Array

FIG. 6G

LIF Expression in Pei Pancreas
Pancreatic Carcinoma vs. Normal

Over-expression Gene Rank: 843 (in top 5%)
Reporter: 205266_at
P-value: 6.38E-6
t-Test: 5.120
Fold Change: 3.089



Legend

Pei Pancreas

1. Pancreas (16) 2. Pancreatic Carcinoma (36)

Cancer Cell 2009/09/08

52 samples


mRNA

Human Genome U133 Plus 2.0 Array 19,574 measured genes

FIG. 6H

LIF Expression in Garnett CellLine
Multi-cancer Cell Line - KRAS Mutation

Over-expression Gene Rank: 61 (in top 1%)
Reporter: 205266_at
P-value: 3.02E-10
t-Test: 6.846
Fold Change: 2.770



1. KRAS Wild Type (522) 2. KRAS Mutation (76)

Nature 2012/03/28 732 samples

mRNA 12,624 measured genes

Human Genome U133A Array

FIG. 6I

LIF Expression in Garnett CellLine
Cytarabine Sensitive - Brain and CNS Cancer Cell
Line

Under-expression Gene Rank: 56 (in top 1%)
Reporter: 205266_at
P-value: 6.23E-7
t-Test: -6.008
Fold Change: -4.297




FIG. 7A

LIF Expression in Garnett CellLine
Vorinostat Sensitive - Multi-cancer Cell Line

Under-expression Gene Rank: 29 (in top 1%)
Reporter: 205266_at
P-value: 2.94E-15
t-Test: -8.428
Fold Change: -3.245







FIG. 7B





LIF Expression in Garnett CellLine
AZD8055 Sensitive - Brain and CNS Cancer Cell Line

Under-expression Gene Rank: 146 (in top 2%)
Reporter: 205266_at
P-value: 1.17E-5
t-Test: -5.926
Fold Change: -4.362



Legend	Garnett CellLine	
1. AZD8055 Resistant (43)	Nature 2012/03/28	732 samples
2. AZD8055 Sensitive (4)	mRNA	12,624 measured genes
	Human Genome U133A Array	

FIG. 7C



LIF Expression in Garnett CellLine
Tretinoin Sensitive - Brain and CNS Cancer Cell Line

Under-expression Gene Rank: 187 (in top 2%)
Reporter: 205266_at
P-value: 6.01E-6
t-Test: -6.163
Fold Change: -3.791




FIG. 7D




FIG. 5F