(19) IPOS INTELLECTUAL PROPERTY OFFICE OF SINGAPORE

(11) Publication number: **SG 187410 A1**
(43) Publication date: **28.02.2013**
(51) Int. Cl: ;

(12) Patent Application

(21) Application number: **2012096343**
(22) Date of filing: **23.12.2008**
(30) Priority: **US 61/017,418 28.12.2007**
US 61/017,881 31.12.2007

(71) Applicant: **BAXTER INTERNATIONAL INC. ONE BAXTER PARKWAY, DEERFIELD, IL 60015 IN US**
BAXTER HEALTHCARE SA THURGAUERSTRASSE 130, 8152 GLATTPARK (OPFIKON) CH

(72) Inventor: **MATTHIESSEN, PETER VORGARTENSTRASSE 129-143/1/5, A-1020 VIENNA AT**
TURECEK, PETER WEIDLING, HAUPTSTRASSE 59G, A-3400 KLOSTERNEUBURG AT
SCHWARZ, HANS-PETER WEIMARER STRASSE 76, A-1180 VIENNA AT
SCHNECKER, KURT PROMENADEWEG 6, A-1170 VIENNA AT

(54) **Title:**
RECOMBINANT VWF FORMULATIONS

(57) **Abstract:**
RECOMBINANT VWF FORMULATIONS Abstract The present invention provides long-term stable pharmaceutical formulations of recombinant von-Willebrand Factor (rVWT) and methods for making and administering said formulations. (No suitable figure)

This PDF First Page has been artificially created from the Singaporian Abstracts

RECOMBINANT VWF FORMULATIONS

Abstract

The present invention provides long-term stable pharmaceutical formulations of recombinant von-Willebrand Factor (rVWF) and methods for making and administering said formulations.

(No suitable figure)

RECOMBINANT VWF FORMULATIONS

This application claims priority of U.S. Provisional Application No. 61/017,418, filed December 28, 2007, and U.S. Provisional Application No. 61/017,881 filed December 31, 2007, each of which is incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

[0001] Generally, the invention relates to formulations of recombinant VWF and methods for making a composition comprising recombinant VWF.

BACKGROUND OF THE INVENTION

[0002] Von Willebrand factor (VWF) is a glycoprotein circulating in plasma as a series of multimers ranging in size from about 500 to 20,000 kD. Multimeric forms of VWF are composed of 250 kD polypeptide subunits linked together by disulfide bonds. VWF mediates initial platelet adhesion to the sub-endothelium of the damaged vessel wall. Only the larger multimers exhibit hemostatic activity. It is assumed that endothelial cells secrete large polymeric forms of VWF and those forms of VWF which have a low molecular weight (low molecular weight VWF) arise from proteolytic cleavage. The multimers having large molecular masses are stored in the Weibel-Pallade bodies of endothelial cells and liberated upon stimulation.

[0003] VWF is synthesized by endothelial cells and megakaryocytes as prepro-VWF that consists to a large extent of repeated domains. Upon cleavage of the signal peptide, pro-VWF dimerizes through disulfide linkages at its C-terminal region. The dimers serve as protomers for multimerization, which is governed by disulfide linkages between the free end termini. The assembly to multimers is followed by the proteolytic removal of the propeptide sequence (Leyte et al., Biochem. J. 274 (1991), 257-261).

[0004] The primary translation product predicted from the cloned cDNA of VWF is a 2813-residue precursor polypeptide (prepro-VWF). The prepro-VWF consists of a 22 amino acid signal peptide and a 741 amino acid propeptide, with the mature VWF comprising 2050 amino acids (Ruggeri Z.A., and Ware, J., FASEB J., 308-316 (1993)).

[0005] Defects in VWF are causal to Von Willebrand disease (VWD), which is characterized by a more or less pronounced bleeding phenotype. VWD type 3 is the most severe form in which VWF is completely missing, and VWD type 1 relates to a quantitative loss of VWF and its phenotype can be very mild. VWD type 2 relates to qualitative defects of VWF and can be as severe as VWD type 3. VWD type 2 has many sub forms, some being associated with the loss or the decrease of high molecular weight multimers. Von Willebrand syndrome type 2a (VWS-2A) is characterized by a loss of both intermediate and large multimers. VWS-2B is characterized by a loss of highest-molecular-weight multimers. Other diseases and disorders related to VWF are known in the art.

[0006] US. Patent Nos. 6,531,577, 7,166,709, and European Patent Application No. 04380188.5, describe plasma-derived VWF formulations. However, in addition to quantity and purity issues with plasma-derived VWF, there is also a risk of blood-born pathogens (e.g., viruses and Variant Creutzfeldt-Jakob disease (vCJD).

[0007] Thus there exists a need in the art to develop a stable pharmaceutical formulation comprising recombinant VWF.

SUMMARY OF THE INVENTION

[0008] The present invention provides formulations useful for compositions comprising recombinant VWF, resulting in a highly stable pharmaceutical composition. The stable pharmaceutical composition is useful as a therapeutic agent in the treatment of individuals suffering from disorders or conditions that can benefit from the administration of recombinant VWF.

[0009] In one embodiment, the invention provides a stable liquid pharmaceutical formulation of a recombinant von Willebrand Factor (rVWF) comprising: (a) a rVWF; (b) a buffering agent; (c) one or more salts; (d) optionally a stabilizing agent; and (e) optionally a surfactant; wherein the rVWF comprises a polypeptide selected from the group consisting of: a) the amino acid sequence set out in SEQ ID NO: 3; b) a biologically active analog, fragment or variant of a); c) a polypeptide encoded by the polynucleotide set out in SEQ ID NO: 1; d) a biologically active analog, fragment or variant of c); and e) a polypeptide encoded by a polynucleotide that hybridizes to the polynucleotide set out in SEQ ID NO: 1 under moderately stringent hybridization conditions; wherein the buffer is comprised of a pH buffering agent in a range of about 0.1 mM to about 500 mM and wherein the pH is in a range of about 2.0 to about 12.0; wherein the salt is at a concentration of about 1 to 500 mM;

wherein the stabilizing agent is at a concentration of about 0.1 to 1000 mM; and wherein the surfactant is at a concentration of about 0.01 g/L to 0.5 g/L.

[0010] In another embodiment, the aforementioned formulation is provided wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3. In another embodiment, an aforementioned formulation is provided wherein the buffering agent is selected from the group consisting of sodium citrate, glycine, histidine, Tris and combinations of these agents. In yet another embodiment, an aforementioned formulation is provided wherein the buffering agent is citrate. In still another embodiment of the invention, the aforementioned formulation is provided wherein pH is in the range of 6.0-8.0, or 6.5-7.3. In a related embodiment, the aforementioned formulation is provided wherein the pH is 7.0. In another embodiment, an aforementioned formulation is provided wherein the buffering agent is citrate and the pH is 7.0.

[0011] In still another embodiment, an aforementioned formulation is provided wherein the salt is selected from the group consisting of calcium chloride, sodium chloride and magnesium chloride. In another embodiment, the aforementioned formulation is provided wherein the salt is at a concentration range of 0.5 to 300 mM. In another embodiment, the aforementioned formulation is provided wherein the salt is calcium chloride at a concentration of 10 mM.

[0012] In another embodiment, an aforementioned formulation is provided wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agent is citrate and the pH is 7.0; and wherein the salt is calcium chloride at a concentration of 10 mM. In still another embodiment, an aforementioned formulation is provided wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agent is sodium citrate and the pH is 7.0; and wherein the salt is calcium chloride at a concentration of 10 mM and NaCl at a concentration of 100 mM.

[0013] Other formulations are also contemplated by the instant invention. For example, in one embodiment, an aforementioned formulation is provided wherein the one or more buffering agents is histidine and Tris at a concentration of 3.3 mM each. In another embodiment, the aforementioned formulation is provided wherein the pH is 7.0. In yet another embodiment, an aforementioned formulation is provided wherein the first salt is sodium chloride at a concentration of 30 mM and the second salt is calcium chloride at a concentration of 0.56 mM.

[0014] In still another embodiment of the invention, an aforementioned formulation is provided wherein the stabilizing agent is selected from the group consisting of mannitol, lactose, sorbitol, xylitol, sucrose, trehalose, mannose, maltose, lactose, glucose, raffinose, cellobiose,

gentiobiose, isomaltose, arabinose, glucosamine, fructose and combinations of these stabilizing agents. In another embodiment, the aforementioned formulation is provided wherein the stabilizing agents are trehalose at a concentration of 7.8 mM and mannitol at a concentration of 58.6 mM.

[0015] In another embodiment, an aforementioned formulation is provided wherein the surfactant is selected from the group consisting of digitonin, Triton X-100, Triton X-114, TWEEN-20, TWEEN-80 and combinations of these surfactants. In another embodiment, the aforementioned formulation is provided wherein the surfactant is TWEEN-80 at 0.03 g/L.

[0016] In one embodiment of the invention, an aforementioned formulation is provided wherein the rVWF comprises amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agents are histidine at a concentration of 3.3 mM and Tris at a concentration of 3.3 mM at pH 7.0; wherein the first salt is sodium chloride at a concentration of 30 mM and the second salt is calcium chloride at a concentration of 0.56 mM; wherein the stabilizing agents are trehalose at a concentration of 7.8 mM mannitol at a concentration of 58.6 mM; and wherein the surfactant is TWEEN-80 at 0.03 g/L.

BRIEF DESCRIPTION OF THE FIGURES

[0017] **Figure 1** shows that rVWF is not stable in Advate buffer after 26 weeks, due to the presence of glutathione.

[0018] **Figure 2** shows that rVWF is stable in Advate 1:3 buffer for up to 12 weeks at 4 °C.

[0019] **Figure 3** shows that the stability of a citrate-based formulation is better than Advate 1:3 buffer formulation containing 0.1M glutathione.

[0020] **Figure 4** shows shows that rVWF concentration is stable over 26 weeks in Advate buffer.

[0021] **Figure 5** shows that rVWF concentration is stable over time in Advate 1:3 buffer.

[0022] **Figure 6** shows that rVWF concentration is stable over time in citrate-based buffer.

[0023] **Figure 7** shows that most excipients increase the unfolding temperature of rVWF by about 1 or 2 °C.

[0024] **Figure 8** shows that 10 mM $CaCl_2$ increases unfolding temperature of rVWF by about 8 °C to about 67 °C.

[0025] **Figure 9** shows that the effect of $CaCl_2$ is similar at pH 7.3 and pH 6.5.

DETAILED DESCRIPTION OF THE INVENTION

Definition of terms

[0026] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton, et al., DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY (2d ed. 1994); THE CAMBRIDGE DICTIONARY OF SCIENCE AND TECHNOLOGY (Walker ed., 1988); THE GLOSSARY OF GENETICS, 5TH ED., R. Rieger, et al. (eds.), Springer Verlag (1991); and Hale and Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY (1991).

[0027] Each publication, patent application, patent, and other reference cited herein is incorporated by reference in its entirety to the extent that it is not inconsistent with the present disclosure.

[0028] It is noted here that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

[0029] As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0030] The term "comprising," with respect to a peptide compound, means that a compound may include additional amino acids at either or both amino and carboxy termini of the given sequence. Of course, these additional amino acids should not significantly interfere with the activity of the compound. With respect to a composition of the instant invention, the term "comprising" means that a composition may include additional components. These additional components should not significantly interfere with the activity of the composition.

[0031] The term "pharmacologically active" means that a substance so described is determined to have activity that affects a medical parameter (e.g., but not limited to blood pressure, blood cell count, cholesterol level) or disease state (e.g., but not limited to cancer, autoimmune disorders).

[0032] As used herein the terms "express," "expressing" and "expression" mean allowing or causing the information in a gene or DNA sequence to become manifest, for example, producing a

protein by activating the cellular functions involved in transcription and translation of a corresponding gene or DNA sequence. A DNA sequence is expressed in or by a cell to form an "expression product" such as a protein. The expression product itself, e.g. the resulting protein, may also be said to be "expressed." An expression product can be characterized as intracellular, extracellular or secreted. The term "intracellular" means inside a cell. The term "extracellular" means outside a cell, such as a transmembrane protein. A substance is "secreted" by a cell if it appears in significant measure outside the cell, from somewhere on or inside the cell.

[0033] As used herein a "polypeptide" refers to a polymer composed of amino acid residues, structural variants, related naturally-occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds. Synthetic polypeptides can be prepared, for example, using an automated polypeptide synthesizer. The term "protein" typically refers to large polypeptides. The term "peptide" typically refers to short polypeptides.

[0034] As used herein a "fragment" of a polypeptide is meant to refer to any portion of a polypeptide or protein smaller than the full-length polypeptide or protein expression product.

[0035] As used herein an "analog" refers to any of two or more polypeptides substantially similar in structure and having the same biological activity, but can have varying degrees of activity, to either the entire molecule, or to a fragment thereof. Analogs differ in the composition of their amino acid sequences based on one or more mutations involving substitution of one or more amino acids for other amino acids. Substitutions can be conservative or non-conservative based on the physico-chemical or functional relatedness of the amino acid that is being replaced and the amino acid replacing it.

[0036] As used herein a "variant" refers to a polypeptide, protein or analog thereof that is modified to comprise additional chemical moieties not normally a part of the molecule. Such moieties may modulate the molecule's solubility, absorption, biological half-life, etc. The moieties may alternatively decrease the toxicity of the molecule and eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed in Remington's Pharmaceutical Sciences (1980). Procedure for coupling such moieties to a molecule are well known in the art. For example, the variant may be a blood clotting factor having a chemical modification which confers a longer half-life in vivo to the protein. In various aspects, polypeptides are modified by glycosylation, pegylation, and/or polysialylation.

[0037] **Recombinant VWF**

[0038] The polynucleotide and amino acid sequences of prepro-VWF are set out in SEQ ID NO:1 and SEQ ID NO:2, respectively, and are available at GenBank Accession Nos. NM_000552 and NP_000543, respectively. The amino acid sequence corresponding to the mature VWF protein is set out in SEQ ID NO: 3 (corresponding to amino acids 764-2813 of the full length prepro-VWF amino acid sequence).

[0039] One form of useful rVWF has at least the property of in vivo-stabilizing, e.g. binding, of at least one Factor VIII (FVIII) molecule and having optionally a glycosylation pattern which is pharmacologically acceptable. Specific examples thereof include VWF without A2 domain thus resistant to proteolysis (Lankhof et al., Thromb. Haemost. 77: 1008-1013, 1997), and the VWF fragment from Val 449 to Asn 730 including the glycoprotein Ib-binding domain and binding sites for collagen and heparin (Pietu et al., Biochem. Biophys. Res. Commun. 164: 1339-1347, 1989). The determination of the ability of a VWF to stabilize at least one FVIII molecule can be carried out in VWF-deficient mammals according to methods known in the state in the art.

[0040] The rVWF of the present invention may be produced by any method known in the art. One specific example is disclosed in WO86/06096 published on Oct. 23, 1986 and U.S. Patent Application No. 07/559,509, filed on Jul. 23, 1990, which is incorporated herein by reference with respect to the methods of producing recombinant VWF. Thus, methods are known in the art for (i) the production of recombinant DNA by genetic engineering, e.g. via reverse transcription of RNA and/or amplification of DNA, (ii) introducing recombinant DNA into procaryotic or eucaryotic cells by transfection, e.g. via electroporation or microinjection, (iii) cultivating said transformed cells, e.g. in a continuous or batchwise manner, (iv) expressing VWF, e.g. constitutively or upon induction, and (v) isolating said VWF, e.g. from the culture medium or by harvesting the transformed cells, in order to (vi) obtain purified rVWF, e.g. via anion exchange chromatography or affinity chromatography. A recombinant VWF may be made in transformed host cells using recombinant DNA techniques well known in the art. For instance, sequences coding for the polypeptide could be excised from DNA using suitable restriction enzymes.

[0041] Alternatively, the DNA molecule could be synthesized using chemical synthesis techniques, such as the phosphoramidate method. Also, a combination of these techniques could be used.

[0042] The invention also provides vectors encoding polypeptides of the invention in an appropriate host. The vector comprises the polynucleotide that encodes the polypeptide operatively linked to appropriate expression control sequences. Methods of effecting this operative linking,

either before or after the polynucleotide is inserted into the vector, are well known. Expression control sequences include promoters, activators, enhancers, operators, ribosomal binding sites, start signals, stop signals, cap signals, polyadenylation signals, and other signals involved with the control of transcription or translation. The resulting vector having the polynucleotide therein is used to transform an appropriate host. This transformation may be performed using methods well known in the art.

[0043] Any of a large number of available and well-known host cells may be used in the practice of this invention. The selection of a particular host is dependent upon a number of factors recognized by the art, including, for example, compatibility with the chosen expression vector, toxicity of the peptides encoded by the DNA molecule, rate of transformation, ease of recovery of the peptides, expression characteristics, bio-safety and costs. A balance of these factors must be struck with the understanding that not all host cells are equally effective for the expression of a particular DNA sequence. Within these general guidelines, useful microbial host cells include bacteria, yeast and other fungi, insects, plants, mammalian (including human) cells in culture, or other hosts known in the art.

[0044] Next, the transformed host is cultured and purified. Host cells may be cultured under conventional fermentation conditions so that the desired compounds are expressed. Such fermentation conditions are well known in the art. Finally, the polypeptides are purified from culture by methods well known in the art.

[0045] Depending on the host cell utilized to express a compound of the invention, carbohydrate (oligosaccharide) groups may conveniently be attached to sites that are known to be glycosylation sites in proteins. Generally, O-linked oligosaccharides are attached to serine (Ser) or threonine (Thr) residues while N-linked oligosaccharides are attached to asparagine (Asn) residues when they are part of the sequence Asn-X-Ser/Thr, where X can be any amino acid except proline. X is preferably one of the 19 naturally occurring amino acids not counting proline. The structures of N-linked and O-linked oligosaccharides and the sugar residues found in each type are different. One type of sugar that is commonly found on both is N-acetylneuraminic acid (referred to as sialic acid). Sialic acid is usually the terminal residue of both N-linked and O-linked oligosaccharides and, by virtue of its negative charge, may confer acidic properties to the glycosylated compound. Such site(s) may be incorporated in the linker of the compounds of this invention and are preferably glycosylated by a cell during recombinant production of the polypeptide compounds (e.g., in mammalian cells such as CHO, BHK, COS). However, such sites may further be glycosylated by synthetic or semi-synthetic procedures known in the art.

[0046] Alternatively, the compounds may be made by synthetic methods. For example, solid phase synthesis techniques may be used. Suitable techniques are well known in the art, and include those described in Merrifield (1973), Chem. Polypeptides, pp. 335-61 (Katsoyannis and Panayotis eds.); Merrifield (1963), J. Am. Chem. Soc. 85: 2149; Davis et al. (1985), Biochem. Intl. 10: 394-414; Stewart and Young (1969), Solid Phase Peptide Synthesis; U.S. Pat. No. 3,941,763; Finn et al. (1976), The Proteins (3rd ed.) 2: 105-253; and Erickson et al. (1976), The Proteins (3rd ed.) 2: 257-527. Solid phase synthesis is the preferred technique of making individual peptides since it is the most cost-effective method of making small peptides.

[0047] **Fragments, variants and analogs of VWF**

[0048] Methods for preparing polypeptide fragments, variants or analogs are well-known in the art.

[0049] Fragments of a polypeptide are prepared using, without limitation, enzymatic cleavage (e.g., trypsin, chymotrypsin) and also using recombinant means to generate a polypeptide fragments having a specific amino acid sequence. Polypeptide fragments may be generated comprising a region of the protein having a particular activity, such as a multimerization domain or any other identifiable VWF domain known in the art.

[0050] Methods of making polypeptide analogs are also well-known. Amino acid sequence analogs of a polypeptide can be substitutional, insertional, addition or deletion analogs. Deletion analogs, including fragments of a polypeptide, lack one or more residues of the native protein which are not essential for function or immunogenic activity. Insertional analogs involve the addition of, e.g., amino acid(s) at a non-terminal point in the polypeptide. This analog may include insertion of an immunoreactive epitope or simply a single residue. Addition analogs, including fragments of a polypeptide, include the addition of one or more amino acids at either of both termini of a protein and include, for example, fusion proteins.

[0051] Substitutional analogs typically exchange one amino acid of the wild-type for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide without the loss of other functions or properties. In one aspect, substitutions are conservative substitutions. By "conservative amino acid substitution" is meant substitution of an amino acid with an amino acid having a side chain of a similar chemical character. Similar amino acids for making conservative substitutions include those having an acidic side chain (glutamic acid, aspartic acid); a basic side chain (arginine, lysine, histidine); a polar amide side chain (glutamine, asparagine); a hydrophobic, aliphatic side chain (leucine, isoleucine, valine, alanine, glycine); an

aromatic side chain (phenylalanine, tryptophan, tyrosine); a small side chain (glycine, alanine, serine, threonine, methionine); or an aliphatic hydroxyl side chain (serine, threonine).

[0052] Analogs may be substantially homologous or substantially identical to the recombinant VWF from which they are derived. Preferred analogs are those which retain at least some of the biological activity of the wild-type polypeptide, *e.g.* blood clotting activity.

[0053] Polypeptide variants contemplated include polypeptides chemically modified by such techniques as ubiquitination, glycosylation, including polysialation, conjugation to therapeutic or diagnostic agents, labeling, covalent polymer attachment such as pegylation (derivatization with polyethylene glycol), introduction of non-hydrolyzable bonds, and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally occur in human proteins. Variants retain the same or essentially the same binding properties of non-modified molecules of the invention. Such chemical modification may include direct or indirect (e.g., via a linker) attachment of an agent to the VWF polypeptide. In the case of indirect attachment, it is contemplated that the linker may be hydrolyzable or non-hydrolyzable.

[0054] Preparing pegylated polypeptide analogs will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the binding construct polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the percentage of poly-pegylated product. In some embodiments, the binding construct will have a single PEG moiety at the N-terminus. Polyethylene glycol (PEG) may be attached to the blood clotting factor to provide a longer half-life *in vivo*. The PEG group may be of any convenient molecular weight and may be linear or branched. The average molecular weight of the PEG ranges from about 2 kiloDalton ("kD") to about 100 kDa, from about 5 kDa to about 50 kDa, or from about 5 kDa to about 10 kDa. The PEG groups are attached to the blood clotting factor via acylation or reductive alkylation through a natural or engineered reactive group on the PEG moiety (e.g., an aldehyde, amino, thiol, or ester group) to a reactive group on the blood clotting factor (e.g., an aldehyde, amino, or ester group) or by any other technique known in the art.

[0055] Methods for preparing polysialylated polypeptide are described in United States Patent Publication 20060160948, Fernandes et Gregoriadis; Biochim. Biophys. Acta 1341: 26-34, 1997, and Saenko et al., Haemophilia 12:42-51, 2006. Briefly, a solution of colominic acid

containing 0.1 M $NaIO_4$ is stirred in the dark at room temperature to oxidize the CA. The activated CA solution is dialyzed against, e.g., 0.05 M sodium phosphate buffer, pH 7.2 in the dark and this solution was added to a rVWF solution and incubated for 18 h at room temperature in the dark under gentle shaking. Free reagents can then be separated from the rVWF-polysialic acid conjugate by ultrafiltration/diafiltration. Conjugation of rVWF with polysialic acid may also be achieved using glutaraldehyde as cross-linking reagent (Migneault et al., Biotechniques 37: 790-796, 2004).

[0056] It is further contemplated that a polypeptide of the invention may be a fusion protein with a second agent which is a polypeptide. In one embodiment, the second agent which is a polypeptide, without limitation, is an enzyme, a growth factor, an antibody, a cytokine, a chemokine, a cell-surface receptor, the extracellular domain of a cell surface receptor, a cell adhesion molecule, or fragment or active domain of a protein described above. In a related embodiment, the second agent is a blood clotting factor such as Factor VIII, Factor VII, Factor IX. The fusion protein contemplated is made by chemical or recombinant techniques well-known in the art.

[0057] It is also contemplated that prepro-VWF and pro-VWF polypeptides may provide a therapeutic benefit in the formulations of the present invention. For example, US Patent No. 7,005,502 describes a pharmaceutical preparation comprising substantial amounts of pro-VWF that induces thrombin gemeation *in vitro*. In addition to recombinant, biologically active fragments, variants, or analogs of the naturally-occuring mature VWF, the present invention contemplates the use of recombinant biologically active fragments, variants, or analogs of the prepro-VWF (set out in SEQ ID NO:2) or pro-VWF polypeptides (amino acid residues 23 to 764 of SEQ ID NO: 2) in the formulations described herein.

[0058] Polynucleotides encoding fragments, variants and analogs may be readily generated by a worker of skill to encode biologically active fragments, variants, or analogs of the naturally-occurring molecule that possess the same or similar biological activity to the naturally-occurring molecule. These polynucleotides can be prepared using PCR techniques, digestion/ligation of DNA encoding molecule, and the like. Thus, one of skill in the art will be able to generate single base changes in the DNA strand to result in an altered codon and a missense mutation, using any method known in the art, including, but not limited to site-specific mutagenesis. As used herein, the phrase "moderately stringent hybridization conditions" means, for example, hybridization at 42°C in 50% formamide and washing at 60°C in 0.1 x SSC, 0.1% SDS. It is understood by those of skill in the art that variation in these conditions occurs based on the length and GC nucleotide base content of the sequences to be hybridized. Formulas standard in the art are appropriate for determining exact

hybridization conditions. See Sambrook et al., 9.47-9.51 in Molecular Cloning, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989).

[0059] Formulations and excipients in general

[0060] Excipients are additives that are included in a formulation because they either impart or enhance the stability and delivery of a drug product. Regardless of the reason for their inclusion, excipients are an integral component of a drug product and therefore need to be safe and well tolerated by patients. For protein drugs, the choice of excipients is particularly important because they can affect both efficacy and immunogenicity of the drug. Hence, protein formulations need to be developed with appropriate selection of excipients that afford suitable stability, safety, and marketability.

[0061] The principal challenge in developing formulations for therapeutic proteins is stabilizing the product against the stresses of manufacturing, shipping and storage. The role of formulation excipients is to provide stabilization against these stresses. Excipients may also be employed to reduce viscosity of high concentration protein formulations in order to enable their delivery and enhance patient convenience. In general, excipients can be classified on the basis of the mechanisms by which they stabilize proteins against various chemical and physical stresses. Some excipients are used to alleviate the effects of a specific stress or to regulate a particular susceptibility of a specific protein. Other excipients have more general effects on the physical and covalent stabilities of proteins. The excipients described herein are organized either by their chemical type or their functional role in formulations. Brief descriptions of the modes of stabilization are provided when discussing each excipient type.

[0062] Given the teachings and guidance provided herein, those skilled in the art will know what amount or range of excipient can be included in any particular formulation to achieve a biopharmaceutical formulation of the invention that promotes retention in stability of the biopharmaceutical (e.g., a polypeptide). For example, the amount and type of a salt to be included in a biopharmaceutical formulation of the invention can be selected based on the desired osmolality (i.e., isotonic, hypotonic or hypertonic) of the final solution as well as the amounts and osmolality of other components to be included in the formulation. Similarly, by exemplification with reference to the type of polyol or sugar included in a formulation, the amount of such an excipient will depend on its osmolality.

[0063] By way of example, inclusion of about 5% sorbitol can achieve isotonicity while about 9% of a sucrose excipient is needed to achieve isotonicity. Selection of the amount or range of

concentrations of one or more excipients that can be included within a biopharmaceutical formulation of the invention has been exemplified above by reference to salts, polyols and sugars. However, those skilled in the art will understand that the considerations described herein and further exemplified by reference to specific excipients are equally applicable to all types and combinations of excipients including, for example, salts, amino acids, other tonicity agents, surfactants, stabilizers, bulking agents, cryoprotectants, lyoprotectants, anti-oxidants, metal ions, chelating agents and/or preservatives.

[0064] Further, where a particular excipient is reported in molar concentration, those skilled in the art will recognize that the equivalent percent (%) w/v (e.g., (grams of substance in a solution sample/mL of solution) X 100%) of solution is also contemplated.

[0065] Of course, a person having ordinary skill in the art would recognize that the concentrations of the excipients described herein share an interdependency within a particular formulation. By way of example, the concentration of a bulking agent may be lowered where, e.g., there is a high polypeptide concentration or where, e.g., there is a high stabilizing agent concentration. In addition, a person having ordinary skill in the art would recognize that, in order to maintain the isotonicity of a particular formulation in which there is no bulking agent, the concentration of a stabilizing agent would be adjusted accordingly (i.e., a "tonicifying" amount of stabilizer would be used). Common excipients are known in the art and can be found in Powell et al., Compendium of Excipients fir Parenteral Formulations (1998), PDA J. Pharm. Sci. Technology, 52:238-311.

[0066] Buffers and buffering agents

[0067] The stability of a pharmacologically active polypeptide formulation is usually observed to be maximal in a narrow pH range. This pH range of optimal stability needs to be identified early during pre-formulation studies. Several approaches, such as accelerated stability studies and calorimetric screening studies, have been demonstrated to be useful in this endeavor (Remmele R.L. Jr., et al., *Biochemistry*, 38(16): 5241-7 (1999)). Once a formulation is finalized, the drug product must be manufactured and maintained throughout its shelf-life. Hence, buffering agents are almost always employed to control pH in the formulation.

[0068] Organic acids, phosphates and Tris have been employed routinely as buffers in protein formulations. The buffer capacity of the buffering species is maximal at a pH equal to the pKa and decreases as pH increases or decreases away from this value. Ninety percent of the

buffering capacity exists within one pH unit of its pKa. Buffer capacity also increases proportionally with increasing buffer concentration.

[0069] Several factors need to be considered when choosing a buffer. First and foremost, the buffer species and its concentration need to be defined based on its pKa and the desired formulation pH. Equally important is to ensure that the buffer is compatible with the polypeptide and other formulation excipients, and does not catalyze any degradation reactions. A third important aspect to be considered is the sensation of stinging and irritation the buffer may induce upon administration. For example, citrate is known to cause stinging upon injection (Laursen T, et al., *Basic Clin Pharmacol Toxicol.*, 98(2): 218-21 (2006)). The potential for stinging and irritation is greater for drugs that are administered via the subcutaneous (SC) or intramuscular (IM) routes, where the drug solution remains at the site for a relatively longer period of time than when administered by the IV route where the formulation gets diluted rapidly into the blood upon administration. For formulations that are administered by direct IV infusion, the total amount of buffer (and any other formulation component) needs to be monitored. One has to be particularly careful about potassium ions administered in the form of the potassium phosphate buffer, which can induce cardiovascular effects in a patient (Hollander-Rodriguez JC, et al., *Am. Fam. Physician.*, 73(2): 283-90 (2006)).

[0070] The buffer system present in the compositions is selected to be physiologically compatible and to maintain a desired pH of the pharmaceutical formulation. In one embodiment, the pH of the solution is between pH 2.0 and pH 12.0. For example, the pH of the solution may be 2.0, 2.3, 2.5, 2.7, 3.0, 3.3, 3.5, 3.7, 4.0, 4.3, 4.5, 4.7, 5.0, 5.3, 5.5, 5.7, 6.0, 6.3, 6.5, 6.7, 7.0, 7.3, 7.5, 7.7, 8.0, 8.3, 8.5, 8.7, 9.0, 9.3, 9.5, 9.7, 10.0, 10.3, 10.5, 10.7, 11.0, 11.3, 11.5, 11.7, or 12.0.

[0071] The pH buffering compound may be present in any amount suitable to maintain the pH of the formulation at a predetermined level. In one embodiment, the pH buffering concentration is between 0.1 mM and 500 mM (1 M). For example, it is contemplated that the pH buffering agent is at least 0.1, 0.5, 0.7, 0.8 0.9, 1.0, 1.2, 1.5, 1.7, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500 mM.

[0072] Exemplary pH buffering agents used to buffer the formulation as set out herein include, but are not limited to glycine, histidine, glutamate, succinate, phosphate, acetate, citrate, Tris and amino acids or mixtures of amino acids, including, but not limited to aspartate, histidine, and glycine..

[0073] **Salts**

[0074] Salts are often added to increase the ionic strength of the formulation, which can be important for protein solubility, physical stability, and isotonicity. Salts can affect the physical stability of proteins in a variety of ways. Ions can stabilize the native state of proteins by binding to charged residues on the protein's surface. Alternatively, salts can stabilize the denatured state by binding to peptide groups along the protein backbone (-CONH-). Salts can also stabilize the protein native conformation by shielding repulsive electrostatic interactions between residues within a protein molecule. Salts in protein formulations can also shield attractive electrostatic interactions between protein molecules that can lead to protein aggregation and insolubility. In formulations provided, the salt concentration is between 0.1, 1, 10, 20, 30, 40, 50, 80, 100, 120, 150, 200, 300, and 500 mM.

[0075] **Stabilizers and bulking agents**

[0076] In the present pharmaceutical formulations, a stabilizer (or a combination of stabilizers) may be added to prevent or reduce storage-induced aggregation and chemical degradation. A hazy or turbid solution upon reconstitution indicates that the protein has precipitated or at least aggregated. The term "stabilizer" means an excipient capable of preventing aggregation or other physical degradation, as well as chemical degradation (for example, autolysis, deamidation, oxidation, etc.) in an aqueous state. Stabilizers that are conventionally employed in pharmaceutical compositions include, but are not limited to, sucrose, trehalose, mannose, maltose, lactose, glucose, raffinose, cellobiose, gentiobiose, isomaltose, arabinose, glucosamine, fructose, mannitol, sorbitol, glycine, arginine HCL, poly-hydroxy compounds, including polysaccharides such as dextran, starch, hydroxyethyl starch, cyclodextrins, N-methyl pyrollidene, cellulose and hyaluronic acid, sodium chloride, [Carpenter et al., Develop. Biol. Standard 74:225, (1991)]. In the present formulations, the stabilizer is incorporated in a concentration of about 0.1, 0.5, 0.7, 0.8 0.9, 1.0, 1.2, 1.5, 1.7, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 700, 900, or 1000 mM.

[0077] If desired, the formulations also include appropriate amounts of bulking and osmolarity regulating agents. Bulking agents include, for example, mannitol, glycine, sucrose, polymers such as dextran, polyvinylpyrolidone, carboxymethylcellulose, lactose, sorbitol, trehalose, or xylitol. In one embodiment, the bulking agent is mannitol. The bulking agent is incorporated in a concentration of about 0.1, 0.5, 0.7, 0.8 0.9, 1.0, 1.2, 1.5, 1.7, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 700, 900, or 1000 mM.

[0078] **Surfactants**

[0079] Protein molecules have a high propensity to interact with surfaces making them susceptible to adsorption and denaturation at air-liquid, vial-liquid, and liquid-liquid (silicone oil) interfaces. This degradation pathway has been observed to be inversely dependent on protein concentration and results in either the formation of soluble and insoluble protein aggregates or the loss of protein from solution via adsorption to surfaces. In addition to container surface adsorption, surface-induced degradation is exacerbated with physical agitation, as would be experienced during shipping and handling of the product.

[0080] Surfactants are commonly used in protein formulations to prevent surface-induced degradation. Surfactants are amphipathic molecules with the capability of out-competing proteins for interfacial positions. Hydrophobic portions of the surfactant molecules occupy interfacial positions (e.g., air/liquid), while hydrophilic portions of the molecules remain oriented towards the bulk solvent. At sufficient concentrations (typically around the detergent's critical micellar concentration), a surface layer of surfactant molecules serve to prevent protein molecules from adsorbing at the interface. Thereby, surface-induced degradation is minimized. The most commonly used surfactants are fatty acid esters of sorbitan polyethoxylates, i.e. polysorbate 20 and polysorbate 80. The two differ only in the length of the aliphatic chain that imparts hydrophobic character to the molecules, C-12 and C-18, respectively. Accordingly, polysorbate-80 is more surface-active and has a lower critical micellar concentration than polysorbate-20.

[0081] Detergents can also affect the thermodynamic conformational stability of proteins. Here again, the effects of a given detergent excipient will be protein specific. For example, polysorbates have been shown to reduce the stability of some proteins and increase the stability of others. Detergent destabilization of proteins can be rationalized in terms of the hydrophobic tails of the detergent molecules that can engage in specific binding with partially or wholly unfolded protein states. These types of interactions could cause a shift in the conformational equilibrium towards the more expanded protein states (i.e. increasing the exposure of hydrophobic portions of the protein molecule in complement to binding polysorbate). Alternatively, if the protein native state exhibits some hydrophobic surfaces, detergent binding to the native state may stabilize that conformation.

[0082] Another aspect of polysorbates is that they are inherently susceptible to oxidative degradation. Often, as raw materials, they contain sufficient quantities of peroxides to cause oxidation of protein residue side-chains, especially methionine. The potential for oxidative damage arising from the addition of stabilizer emphasizes the point that the lowest effective concentrations of excipients should be used in formulations. For surfactants, the effective concentration for a given protein will depend on the mechanism of stabilization. It has been postulated that if the mechanism

of surfactant stabilization is related to preventing surface-denaturation the effective concentration will be around the detergent's critical micellar concentration. Conversely, if the mechanism of stabilization is associated with specific protein-detergent interactions, the effective surfactant concentration will be related to the protein concentration and the stoichiometry of the interaction (Randolph T.W., et al., *Pharm Biotechnol.*, 13:159-75 (2002)).

[0083] Surfactants may also be added in appropriate amounts to prevent surface related aggregation phenomenon during freezing and drying [Chang, B, J. Pharm. Sci. 85:1325, (1996)]. Exemplary surfactants include anionic, cationic, nonionic, zwitterionic, and amphoteric surfactants including surfactants derived from naturally-occurring amino acids. Anionic surfactants include, but are not limited to, sodium lauryl sulfate, dioctyl sodium sulfosuccinate and dioctyl sodium sulfonate, chenodeoxycholic acid, N-lauroylsarcosine sodium salt, lithium dodecyl sulfate, 1-octanesulfonic acid sodium salt, sodium cholate hydrate, sodium deoxycholate, and glycodeoxycholic acid sodium salt. Cationic surfactants include, but are not limited to, benzalkonium chloride or benzethonium chloride, cetylpyridinium chloride monohydrate, and hexadecyltrimethylammonium bromide. Zwitterionic surfactants include, but are not limited to, CHAPS, CHAPSO, SB3-10, and SB3-12. Non-ionic surfactants include, but are not limited to, digitonin, Triton X-100, Triton X-114, TWEEN-20, and TWEEN-80. Surfactants also include, but are not limited to lauromacrogol 400, polyoxyl 40 stearate, polyoxyethylene hydrogenated castor oil 10, 40, 50 and 60, glycerol monostearate, polysorbate 40, 60, 65 and 80, soy lecithin and other phospholipids such as dioleyl phosphatidyl choline (DOPC), dimyristoylphosphatidyl glycerol (DMPG), dimyristoylphosphatidyl choline (DMPC), and (dioleyl phosphatidyl glycerol) DOPG; sucrose fatty acid ester, methyl cellulose and carboxymethyl cellulose. Compositions comprising these surfactants, either individually or as a mixture in different ratios, are therefore further provided. In the present formulations, the surfactant is incorporated in a concentration of about 0.01 to about 0.5 g/L.

[0084] **Other common excipient components**

[0085] Amino acids

[0086] Amino acids have found versatile use in protein formulations as buffers, bulking agents, stabilizers and antioxidants. Histidine and glutamic acid are employed to buffer protein formulations in the pH range of 5.5 – 6.5 and 4.0 – 5.5 respectively. The imidazole group of histidine has a pKa = 6.0 and the carboxyl group of glutamic acid side chain has a pKa of 4.3 which makes these amino acids suitable for buffering in their respective pH ranges. Glutamic acid is particularly useful in such cases (e.g., Stemgen®). Histidine is commonly found in marketed protein

formulations (e.g., Xolair®, Herceptin®, Recombinate®),and this amino acid provides an alternative to citrate, a buffer known to sting upon injection. Interestingly, histidine has also been reported to have a stabilizing effect, as observed in formulations with ABX-IL8 (an IgG2 antibody), with respect to aggregation when used at high concentrations in both liquid and lyophilized presentations (Chen B, et al., *Pharm Res.*, 20(12): 1952-60 (2003)). Histidine (up to 60 mM) was also observed to reduce the viscosity of a high concentration formulation of this antibody. However, in the same study, the authors observed increased aggregation and discoloration in histidine containing formulations during freeze-thaw studies of the antibody in stainless steel containers. The authors attributed this to an effect of iron ions leached from corrosion of steel containers. Another note of caution with histidine is that it undergoes photo-oxidation in the presence of metal ions (Tomita M, et al., *Biochemistry*, 8(12): 5149-60 (1969)). The use of methionine as an antioxidant in formulations appears promising; it has been observed to be effective against a number of oxidative stresses (Lam XM, et al., *J Pharm Sci.*, 86(11): 1250-5 (1997)).

[0087] The amino acids glycine, proline, serine and alanine have been shown to stabilize proteins by the mechanism of preferential exclusion. Glycine is also a commonly used bulking agent in lyophilized formulations (e.g., Neumega ®, Genotropin®, Humatrope®). Arginine has been shown to be an effective agent in inhibiting aggregation and has been used in both liquid and lyophilized formulations (e.g., Activase®, Avonex®, Enbrel® liquid). Furthermore, the enhanced efficiency of refolding of certain proteins in the presence of arginine has been attributed to its suppression of the competing aggregation reaction during refolding.

[0088] Antioxidants

[0089] Oxidation of protein residues arises from a number of different sources. Beyond the addition of specific antioxidants, the prevention of oxidative protein damage involves the careful control of a number of factors throughout the manufacturing process and storage of the product such as atmospheric oxygen, temperature, light exposure, and chemical contamination. The most commonly used pharmaceutical antioxidants are reducing agents, oxygen/free-radical scavengers, or chelating agents. Antioxidants in therapeutic protein formulations are water-soluble and remain active throughout the product shelf-life. Reducing agents and oxygen/free-radical scavengers work by ablating active oxygen species in solution. Chelating agents such as EDTA are effective by binding trace metal contaminants that promote free-radical formation. For example, EDTA was utilized in the liquid formulation of acidic fibroblast growth factor to inhibit the metal ion catalyzed oxidation of cysteine residues. EDTA has been used in marketed products like Kineret® and Ontak®.

[0090] In addition to the effectiveness of various excipients to prevent protein oxidation, the potential for the antioxidants themselves to induce other covalent or physical changes to the protein is of concern. For example, reducing agents can cause disruption of intramolecular disulfide linkages, which can lead to disulfide shuffling. In the presence of transition metal ions, ascorbic acid and EDTA have been shown to promote methionine oxidation in a number of proteins and peptides (Akers MJ, and Defelippis MR. Peptides and Proteins as Parenteral Solutions. In: Pharmaceutical Formulation Development of Peptides and Proteins. Sven Frokjaer, Lars Hovgaard, editors. Pharmaceutical Science. Taylor and Francis, UK (1999)); Fransson J.R., *J. Pharm. Sci.* 86(9): 4046-1050 (1997); Yin J, et al., *Pharm Res.*, 21(12): 2377-83 (2004)). Sodium thiosulfate has been reported to reduce the levels of light and temperature induced methionine-oxidation in rhuMab HER2; however, the formation of a thiosulfate-protein adduct was also reported in this study (Lam XM, Yang JY, et al., *J Pharm Sci.* 86(11): 1250-5 (1997)). Selection of an appropriate antioxidant is made according to the specific stresses and sensitivities of the protein.

[0091] Metal ions

[0092] In general, transition metal ions are undesired in protein formulations because they can catalyze physical and chemical degradation reactions in proteins. However, specific metal ions are included in formulations when they are co-factors to proteins and in suspension formulations of proteins where they form coordination complexes (e.g., zinc suspension of insulin). Recently, the use of magnesium ions (10 –120 mM) has been proposed to inhibit the isomerization of aspartic acid to isoaspartic acid (WO 2004039337).

[0093] Two examples where metal ions confer stability or increased activity in proteins are human deoxyribonuclease (rhDNase, Pulmozyme®), and Factor VIII. In the case of rhDNase, Ca^{+2} ions (up to 100 mM) increased the stability of the enzyme through a specific binding site (Chen B, et al., *J Pharm Sci.*, 88(4): 477-82 (1999)). In fact, removal of calcium ions from the solution with EGTA caused an increase in deamidation and aggregation. However, this effect was observed only with Ca^{+2} ions; other divalent cations Mg^{+2}, Mn^{+2} and Zn^{+2} were observed to destabilize rhDNase. Similar effects were observed in Factor VIII. Ca^{+2} and Sr^{+2} ions stabilized the protein while others like Mg^{+2}, Mn^{+2} and Zn^{+2}, Cu^{+2} and Fe^{+2} destabilized the enzyme (Fatouros, A., et al., *Int. J. Pharm.*, 155, 121–131 (1997). In a separate study with Factor VIII, a significant increase in aggregation rate was observed in the presence of Al^{+3} ions (Derrick TS, et al., *J. Pharm. Sci.*, 93(10): 2549-57 (2004)). The authors note that other excipients like buffer salts are often contaminated with Al^{+3} ions and illustrate the need to use excipients of appropriate quality in formulated products.

[0094] Preservatives

[0095] Preservatives are necessary when developing multi-use parenteral formulations that involve more than one extraction from the same container. Their primary function is to inhibit microbial growth and ensure product sterility throughout the shelf-life or term of use of the drug product. Commonly used preservatives include benzyl alcohol, phenol and m-cresol. Although preservatives have a long history of use, the development of protein formulations that includes preservatives can be challenging. Preservatives almost always have a destabilizing effect (aggregation) on proteins, and this has become a major factor in limiting their use in multi-dose protein formulations (Roy S, et al., *J Pharm Sci.*, 94(2): 382-96 (2005)).

[0096] To date, most protein drugs have been formulated for single-use only. However, when multi-dose formulations are possible, they have the added advantage of enabling patient convenience, and increased marketability. A good example is that of human growth hormone (hGH) where the development of preserved formulations has led to commercialization of more convenient, multi-use injection pen presentations. At least four such pen devices containing preserved formulations of hGH are currently available on the market. Norditropin® (liquid, Novo Nordisk), Nutropin AQ® (liquid, Genentech) & Genotropin (lyophilized – dual chamber cartridge, Pharmacia & Upjohn) contain phenol while Somatrope® (Eli Lilly) is formulated with m-cresol.

[0097] Several aspects need to be considered during the formulation development of preserved dosage forms. The effective preservative concentration in the drug product must be optimized. This requires testing a given preservative in the dosage form with concentration ranges that confer anti-microbial effectiveness without compromising protein stability. For example, three preservatives were successfully screened in the development of a liquid formulation for interleukin-1 receptor (Type I), using differential scanning calorimetry (DSC). The preservatives were rank ordered based on their impact on stability at concentrations commonly used in marketed products (Remmele RL Jr., et al., *Pharm Res.*, 15(2): 200-8 (1998)).

[0098] Some preservatives can cause injection site reactions, which is another factor that needs consideration when choosing a preservative. In clinical trials that focused on the evaluation of preservatives and buffers in Norditropin, pain perception was observed to be lower in formulations containing phenol and benzyl alcohol as compared to a formulation containing m-cresol (Kappelgaard A.M., *Horm Res.* 62 Suppl 3:98-103 (2004)). Interestingly, among the commonly used preservative, benzyl alcohol possesses anesthetic properties (Minogue SC, and Sun DA., *Anesth Analg.*, 100(3): 683-6 (2005)).

[0099] Lyophilization

[00100] It is also contemplated that the formulations comprising a VWF polypeptide of the invention may be lyophilized prior to administration. Lyophilization is carried out using techniques common in the art and should be optimized for the composition being developed [Tang et al., Pharm Res. 21:191-200, (2004) and Chang et al., Pharm Res. 13:243-9 (1996)].

[00101] A lyophilization cycle is, in one aspect, composed of three steps: freezing, primary drying, and secondary drying [A.P. Mackenzie, Phil Trans R Soc London, Ser B, Biol 278:167 (1977)]. In the freezing step, the solution is cooled to initiate ice formation. Furthermore, this step induces the crystallization of the bulking agent. The ice sublimes in the primary drying stage, which is conducted by reducing chamber pressure below the vapor pressure of the ice, using a vacuum and introducing heat to promote sublimation. Finally, adsorbed or bound water is removed at the secondary drying stage under reduced chamber pressure and at an elevated shelf temperature. The process produces a material known as a lyophilized cake. Thereafter the cake can be reconstituted with either sterile water or suitable diluent for injection.

[00102] The lyophilization cycle not only determines the final physical state of the excipients but also affects other parameters such as reconstitution time, appearance, stability and final moisture content. The composition structure in the frozen state proceeds through several transitions (e.g., glass transitions, wettings, and crystallizations) that occur at specific temperatures and can be used to understand and optimize the lyophilization process. The glass transition temperature (Tg and/or Tg') can provide information about the physical state of a solute and can be determined by differential scanning calorimetry (DSC). Tg and Tg' are an important parameter that must be taken into account when designing the lyophilization cycle. For example, Tg' is important for primary drying. Furthermore, in the dried state, the glass transition temperature provides information on the storage temperature of the final product.

[00103] Methods of Preparation

[00104] The present invention further contemplates methods for the preparation of pharmaceutical formulations. A variety of aqueous carriers, e.g., sterile water for injection, water with preservatives for multi dose use, or water with appropriate amounts of surfactants (for example, polysorbate-20), 0.4% saline, 0.3% glycine, or aqueous suspensions may contain the active compound in admixture with excipients suitable for the manufacture of aqueous suspensions. In various aspects, such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum

tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyl-eneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate.

[00105] **Administration**

[00106] To administer compositions to human or test animals, in one aspect, the compositions comprises one or more pharmaceutically acceptable carriers. The phrases "pharmaceutically" or "pharmacologically" acceptable refer to molecular entities and compositions that are stable, inhibit protein degradation such as aggregation and cleavage products, and in addition do not produce allergic, or other adverse reactions when administered using routes well-known in the art, as described below. "Pharmaceutically acceptable carriers" include any and all clinically useful solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like, including those agents disclosed above.

[00107] The pharmaceutical formulations may be administered orally, topically, transdermally, parenterally, by inhalation spray, vaginally, rectally, or by intracranial injection. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intracisternal injection, or infusion techniques. Administration by intravenous, intradermal, intramusclar, intramammary, intraperitoneal, intrathecal, retrobulbar, intrapulmonary injection and or surgical implantation at a particular site is contemplated as well. Generally, compositions are essentially free of pyrogens, as well as other impurities that could be harmful to the recipient.

[00108] Single or multiple administrations of the compositions can be carried out with the dose levels and pattern being selected by the treating physician. For the prevention or treatment of disease, the appropriate dosage will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether drug is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the drug, and the discretion of the attending physician.

[00109] **Kits**

[00110] As an additional aspect, the invention includes kits which comprise one or more pharmaceutical formulations packaged in a manner which facilitates their use for administration to subjects. In one embodiment, such a kit includes pharmaceutical formulation described herein (e.g., a composition comprising a therapeutic protein or peptide), packaged in a container such as a sealed bottle or vessel, with a label affixed to the container or included in the package that describes use of the compound or composition in practicing the method. In one embodiment, the pharmaceutical formulation is packaged in the container such that the amount of headspace in the container (e.g., the amount of air between the liquid formulation and the top of the container) is very small. Preferably, the amount of headspace is negligible (i.e., almost none). In one embodiment, the kit contains a first container having a therapeutic protein or peptide composition and a second container having a physiologically acceptable reconstitution solution for the composition. In one aspect, the pharmaceutical formulation is packaged in a unit dosage form. The kit may further include a device suitable for administering the pharmaceutical formulation according to a specific route of administration. Preferably, the kit contains a label that describes use of the pharmaceutical formulations.

[00111] **Dosages**

[00112] The dosage regimen involved in a method for treating a condition described herein will be determined by the attending physician, considering various factors which modify the action of drugs, e.g. the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors. By way of example, a typical dose of a recombinant VWF of the present invention is approximately 50 U/kg, equal to 500 μg/kg.

[00113] Formulations of the invention may be administered by an initial bolus followed by a continuous infusion to maintain therapeutic circulating levels of drug product. As another example, the inventive compound may be administered as a one-time dose. Those of ordinary skill in the art will readily optimize effective dosages and administration regimens as determined by good medical practice and the clinical condition of the individual patient. The frequency of dosing will depend on the pharmacokinetic parameters of the agents and the route of administration. The optimal pharmaceutical formulation will be determined by one skilled in the art depending upon the route of administration and desired dosage. See for example, Remington's Pharmaceutical Sciences, 18th Ed. (1990, Mack Publishing Co., Easton, PA 18042) pages 1435-1712, the disclosure of which is hereby incorporated by reference. Such formulations may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the administered agents. Depending on the route of administration, a suitable dose may be calculated according to body weight, body surface area or

organ size. Appropriate dosages may be ascertained through use of established assays for determining blood level dosages in conjunction with appropriate dose-response data. The final dosage regimen will be determined by the attending physician, considering various factors which modify the action of drugs, e.g. the drug's specific activity, the severity of the damage and the responsiveness of the patient, the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors. As studies are conducted, further information will emerge regarding the appropriate dosage levels and duration of treatment for various diseases and conditions.

[00114] The following examples are not intended to be limiting but only exemplary of specific embodiments of the invention.

Example 1

Shaking experiments

[00115] In order to determine the amount of precipitation of rVWF in various formulations, the percent recovery of rVWF following turbulent shaking was tested under a variety of conditions.

[00116] rVWF in Advate buffer (90 mM NaCl, 1.68 mM $CaCl_2$, 10 mM L-histidine, 10 mM tris, 0.26 mM glutathione, 23.4 mM trehalose, 175.7 mM mannitol, and 0.1g/L TWEEN-80, pH 7.0)or Advate 1:3 buffer (Advate buffer diluted 3-fold in water) was subjected to turbulent shaking on a shaker at room temperature (RT) for 0 min, 1 min, 2.5 hrs, or 4 days, and percent recovery of the rVWF was measured relative to the starting material prior to shaking. As shown in Table 1, losses of about 40-80% were observed in the Advate buffer while losses of about 20-30% were observed in the Advate 1:3 buffer. VWF antigen VWF:Ag corresponds to the amount of VWF which can be detected in an VWF-specific ELISA using polyclonal anti-VWF antibody, while VWF:RCo corresponds to the amount of VWF which causes agglutination of stabilized platelets in the presence of ristocetin.. In both cases human reference plasma calibrated against the actual WHO standard was used as standard (1 ml of reference plasma usually contains 1U VWF).

[00117] **Table 1. Influence of turbulent shaking time on rVWF recovery**

rVWF	Turbulent shaking at RT	VWF:Ag [U/ml]	Recovery [%]	VWF:RCo [U/ml]	Recovery [%]	RCo/VWF:Ag [U/U]
Advate	0 min	213	100%	104	100%	0.49
	1 min	120	56%			
	2.5 hr	139	65%			
	4 d	37	17%	7	7%	0.19
Advate 1:3	0 min	206	100%	134	100%	0.65
	1 min	152	74%			
	2.5 hr	170	82%			
	4 d	138	67%	131	98%	0.95

[00118] The effect of freeze/thawing and lyophilization was also tested in the shaking experiments. Freezing was performed at –20°C in an –20°C cold room or on dry ice, thawing in both cases at RT and both started from the liquid formulations. As for lyophilization, the formulated VWF samples described herein were frozen within a pilot scale lyophilizer at <=-40°C and were lyophilized using a standard lyo program. Shaking was performed directly with the liquid formulations (2ml in 5 ml vials). As shown in Table 2, percent recovery of rVWF was higher in Advate 1:3 buffer compared to Advate buffer.

[00119] Table 2.

RVWF		VWF:Ag [U/ml]	VWF:Ag recovery [%]	VWF:RCo [U/ml]	VWF:RCo recovery [%]	RCo:Ag [U/U]
Advate	**Frozen**	213	100%	104	100%	0.49
	Frozen – 3x at -20°C	229	107%	84	81%	0.37
	Frozen – 3x with dry ice	231	108%	72	69%	0.31
	Lyo	242	113%	61	59%	0.25
	Starting material	213	100%	104	100%	0.49
	Heavily shaken for 4 days at RT	37.0	17%	7.2	6.9%	0.19
Advate 1:3	**Frozen**	206	100%	134	100%	0.65
	Frozen – 3x at -20°C	184	89%	132	99%	0.72
	Frozen – 3x with dry ice	195	94%	128	96%	0.66
	Lyo	195	94%	107	80%	0.55
	Starting material	206	100%	134	100%	0.65
	Heavily shaken for 4 days at RT	138	67%	131	98%	0.95

[00120] Percent recovery was also measured in the shaking experiments with rVWF being stored in syringes with headspace and without headspace. Interestingly, when rVWF is stored in syringes without headspace and shaken as described above, no rVWF precipitation was observed. In contrast, when rVWF is stored in syringes with headspace, some precipitation was observed.

[00121] In summary, turbulent shaking resulted in at least 30% loss of rVWF in Advate buffer or Advate 1:3 buffer, with Advate buffer showing higher loss of recovery compared to Advate 1:3 buffer. Interestingly, the same precipitates observed in the turbulent shaking experiments were not observed when rVWF was stored and transported ~5000 km in an automobile (representing

the expected shaking during transport). Precipitation of rVWF could be eliminated by storage in syringes without headspace.

Example 2

Stability of recombinant VWF

[00122] The stability of rVWF was tested by assessing the activity level of rVWF present in a various formulations.

[00123] As shown in Figure 1, rVWF is not stable in Advate buffer after 26 weeks due to the presence of 0.3 mM glutathione. As shown in Figure 2, however, rVWF is more stable in Advate 1:3 buffer (e.g., for up 12 weeks at 4°C.)

[00124] As shown in Figure 3, the stability of a citrate-based formulation (15 mM sodium citrate, 10 mM $CaCl_2$, 100 mM NaCl, pH 7.0) is better than Advate 1:3 buffer formulation containing 0.1M glutathione.

[00125] Likewise, the concentration of rVWF was measured over time in various buffers. As shown in Figure 4, Figure 5 and Figure 6, rVWF concentration is stable over time in Advate buffer, Advate 1:3 buffer, and citrate-based buffer, respectively.

Example 4

Characterization of the liquid formulations

[00126] Differential scanning calorimetry (DSC) was used to assess the extent of protein (rVWF) unfolding in various buffers. As shown in Table 3, Advate buffer pH 7.0 is the optimum for stabilization.

[00127] DSC is a thermoanalytical technique in which the difference in the amount of heat required to increase the temperature of a sample and references are measured as a function of temperature. The result of a DSC experiment is a curve of heat flux versus temperature or versus time.

[00128] The Differential Scanning Calorimeter can scan through a range of temperatures while heating and cooling and it determines a phase transition, i.e. melting, crystallization, or glass transition, by measuring the amount of heat needed to reach a set temperature. The calorimeter was calibrated with a set of pure metals (zinc, indium, and tin) that have a known heat capacity, Cp and melting point, Tm. The respective reference buffer was placed into the reference capillary and the rVWF sample was placed into the sample capillary of the

instrument.

[00129] **Table 3. Unfolding temperature in various buffers**

Lot	Buffer	pH	T unfold [°C]
rVWF161A	Advate	7.0	66.0
rVWF161B	Immunate	6.8	64.5
rVWF161C	Citrate	6.8	61.2
rVWF161D	NovoSeven	6.8	64.9
rVWF158	Hepes	7.4	61.3

Buffer components and concentrations:

A) Advate: 5.26g/l NaCl
0.248g/l CaCl2
32g/l D-Mannitol
8g/l Trehalose
1.56g/l L-Histidine
1.2g/l Tris
0.08g/l Glutadione red. pH=7.0

B) Immunate: 5.25g/l Glycin
2.2g/l NaCl
5.25g/l NaCit3
5.25g/l Lysin-HCl
0.62g/l CaCl2 pH=6.8

C) Citrat: 3g/l Glycin
2.92g/l NaCl
2.5g/l NaCit3
30g/l D-Mannitol
10g/l Trehalose pH=6.8

D) NovoSeven: 0.75g/l Glycin
2.92g/l NaCl
1.47g/l CaCl2
30g/l D-Mannitol pH=6,8

rVWF158: 20 mM Hepes, 150 mM NaCl, 5g/L sucrose, pH 7.4

[00130] Further, as shown in Figure 7, most formulation excipients increase the unfolding temperature by about 1-2 °C. Figure 8 shows that 10 mM $CaCl_2$ increases the unfolding

temperature by ~8 °C to ~67 °C, an unfolding temperature which can also be reached by Advate buffer. This effect of $CaCl_2$ is similar at pH 7.3 and 6.5, as shown in Figure 9. Finally, the effect of trehalose and sucrose were analyzed on the unfolding temperature. Compared to citrate alone, neither trehalose nor sucrose increased the unfolding temperature of rVWF. A summary of the unfolding temperature (Tmax) data for rVWF in the presence of various excipients is set out in Table 4.

Table 4.

15 mM Sodium Citrate buffer	-	15 mM Tris	15 mM Glycine	50 mM NaCl
ΔH [kJ/mol]	128494.3	656259.7	157352.2	124985.8
Unfolding T [°C] - Peak 1	58.6		59.1	61
Peak 2	65.2	68.5	65.5	
Peak 3	80.4		80.1	81
Peak 4				
15mM Sodium Citrate buffer	15 mM Histidine	20.52 g/L Mannitol	10.26 g/L Trehalose	
ΔH [kJ/mol]	134044.5	1588590.1	612235.9	
Unfolding T [°C] Peak 1	59.2	58.5	58.5	
Peak 2	65.2	65.5	71.3	
Peak 3	79.3	78.2	81.5	
Peak 4	88.5		92.7	
15mM Sodium Citrate buffer	1 mM $CaCl_2$	10 mM $CaCl_2$	32 g/L Saccharose	0.25 mM Saccharose
ΔH [kJ/mol]	266008.2	308171.3	115082.4	246904.6
Unfolding T [°C] - Peak 1	64.5	67.2	59.2	60
Peak 2			66	67
Peak 3	81	83.1	81.1	81.7
Peak 4	91.8	93		
15mM Sodium Citrate buffer	0.1 g/L TWEEN-80	32 g/L Raffinose	Na_2HPO_4/$NaH PO_4$	7.8 mM Trehalose
ΔH [kJ/mol]	338792.7	127329.2	197967.5	135573.3
Unfolding T [°C] - Peak 1	58.7	60.1	61.4	58.4
Peak 2	64.4	65.8		65.4
Peak 3	81.6	80.3	80.4	80.4
Peak 4			89.2	

[00131] In addition to the various buffers, DSC was used to assess unfolding temperature of rVWF at various pH values in Advate buffer. The results are shown in Table 5, below. Advate buffer pH 7.0 is the optimum for stabilization (i.e., highest unfolding temperature; Peak 1) of rVWF.

[00132] **Table 5.**

pH	Peak 1	Peak 2
5.0	59.5	62.0
6.0	65.2	75.4
7.0	67.2	82.8
8.0	66.6	85.6
9.0	65.0	84.9

[00133] The fluorescence spectrum of rVWF in Advate buffer and Advate 1:3 buffer was assessed after storage at various temperatures for various lengths of time. No (or only slight) change in fluorescence spectrum was observed after storage at 40 °C from 0 to 28 days in either Advate or Advate 1:3 buffers. No difference was observed at other temperatures.

[00134] Likewise, degradation of rVWF was assessed using gelfiltration (Superose 6). While some degradation was observed after 26 weeks at 4 °C in Advate buffer, almost no degradation of rVWF in Advate 1:3 buffer was observed after 26 weeks at 4 °C. At 40 °C, glutathione increased the amount of degradation over time (albeit to a slower extent in Advate 1:3 buffer).

[00135] Based on the above Examples, Advate 1:3 buffer offers an advantage with respect to freeze/thawing and recovery after lyophilization as compared to the undiluted Advate buffer. Moreover, Advate 1:3 buffer can stabilize (e.g., maintain biological activity) rVWF activity during incubation at 40 °C better that Advate buffer. rVWF in Advate 1:3 buffer is stable for 4 weeks of incubation at 4 °C. Finally, DSC has demonstrated that pH 7.0 is optimum for preventing degradation of rVWF (i.e., showed the highest unfolding temperature).

[00136] Thus, in view of the data presented herein, a formulation was proposed for rVWF including 15 mM citrate (or glycine or hitidine), 10 mM $CaCl_2$, pH 6.5-7.3, adjusted to the

desired osmolarity by NaCl. For example, in one embodiment, the citrate-based formula is 15 mM sodium citrate, 10 mM $CaCl_2$, 100 mM NaCl, pH 7.0.

[00137] Alternatively, an Advate or Advate 1:3 buffer, without glutathione, is also contemplated: Advate: 90mM NaCl, 1.68mM $CaCl_2$, 10mM L-histidine, 10mM Tris, 0.26mM glutathione, 23.4mM trehalose, 175.7mM mannitol, and 0.1g/L TWEEN-80, pH 7.0; Advate 1:3: 30 mM NaCl, 0.56 mM $CaCl_2$, 3.3 mM L-histidine, 3.3 mM tris, 7.8 mM trehalose, 58.6 mM mannitol, and 0.03g/L TWEEN-80, ph 7.0.

What is claimed is:

1. A stable liquid pharmaceutical formulation of a recombinant von Willebrand Factor (rVWF) comprising: (a) a rVWF; (b) a buffering agent; (c) one or more salts; (d) optionally a stabilizing agent; and (e) optionally a surfactant;

wherein said rVWF comprises a polypeptide selected from the group consisting of:

a) the amino acid sequence set out in SEQ ID NO: 3;

b) a biologically active analog, fragment or variant of a);

c) a polypeptide encoded by the polynucleotide set out in SEQ ID NO: 1;

d) a biologically active analog, fragment or variant of c); and

e) a polypeptide encoded by a polynucleotide that hybridizes to the polynucleotide set out in SEQ ID NO: 1 under moderately stringent hybridization conditions;

wherein said buffer is comprised of a pH buffering agent in a range of about 0.1 mM to about 500 mM and wherein the pH is in a range of about 2.0 to about 12.0;

wherein said salt is at a concentration of about 1 to 500 mM;

wherein said stabilizing agent is at a concentration of about 0.1 to 1000 mM; and

wherein said surfactant is at a concentration of about 0.01 g/L to 0.5 g/L.

2. The formulation of claim 1 wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3.

3. The formulation of claim 1 wherein the buffering agent is selected from the group consisting of sodium citrate, glycine, histidine, Tris and combinations of these agents.

4. The formulation of claim 3 wherein the buffering agent is sodium citrate at a concentration of 15 mM..

5. The formulation of claim 1 wherein pH is in the range of 6.0-8.0.

6. The formulation of claim 5 wherein pH is in the range of 6.5-7.3.

7. The formulation of claim 4 wherein the pH is 7.0.

8. The formulation of claim 1 wherein the buffering agent is citrate and the pH is 7.0.

9. The formulation of claim 1 wherein the salt is selected from the group consisting of calcium chloride, sodium chloride and magnesium chloride.

10. The formulation of claim 9 wherein the salt is at a concentration range of 0.5 to 300 mM.

11. The formulation of claim 10 wherein the salt is calcium chloride at a concentration of 10 mM.

12. The formulation of claim 1 wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agent is citrate and the pH is 7.0; and wherein the salt is calcium chloride at a concentration of 10 mM.

13. The formulation of claim 1 wherein the rVWF comprises the amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agent is sodium citrate at a concentration of 15 mM and the pH is 7.0; and wherein the salt is calcium chloride at a concentration of 10 mM and NaCl at a concentration of 100 mM.

14. The formulation of claim 3 wherein the one or more buffering agents is histidine and Tris at a concentration of 3.3 mM each.

15. The formulation of claim 3 wherein the pH is 7.0.

16. The formulation of claim 9 wherein the one or more salts is sodium chloride at a concentration of 30 mM and calcium chloride at a concentration of 0.56 mM.

17. The formulation of claim 1 wherein the stabilizing agent is selected from the group consisting of mannitol, lactose, sorbitol, xylitol, sucrose, trehalose, mannose, maltose, lactose, glucose, raffinose, cellobiose, gentiobiose, isomaltose, arabinose, glucosamine, fructose and combinations of these stabilizing agents.

18. The formulation of claim 17 wherein the stabilizing agents are trehalose at a concentration of 7.8 mM and mannitol at a concentration of 58.6 mM.

19. The formulation of claim 1 wherein the surfactant is selected from the group consisting of digitonin, Triton X-100, Triton X-114, TWEEN-20, TWEEN-80 and combinations of these surfactants.

20. The formulation of claim 1 wherein the surfactant is TWEEN-80 at 0.03 g/L.

21. The formulation of claim 1 wherein the rVWF comprises amino acid sequence set out in SEQ ID NO: 3; wherein the buffering agents are histidine at a concentration of 3.3 mM and Tris at a concentration of 3.3 mM at pH 7.0; wherein the salts are sodium chloride at a concentration of 30 mM and calcium chloride at a concentration of 0.56 mM; wherein the stabilizing agents are trehalose at a concentration of 7.8 mM and mannitol at a concentration of 58.6 mM.; and wherein the surfactant is TWEEN-80 at 0.03 g/L.

SEQUENCE LISTING

<110> Matthiessen, et al.

<120> Recombinant VWF Formulations

<130> 31315/43550

<150> 61/017,881
<151> 2007-12-31

<160> 3

<170> PatentIn version 3.5

<210> 1
<211> 8833
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Polynucleotide

<220>
<221> misc_feature
<223> prepro-vWF

<400> 1

agctcacagc tattgtggtg ggaaagggag ggtggttggt ggatgtcaca gcttgggctt 60

tatctccccc agcagtgggg actccacagc ccctgggcta cataacagca agacagtccg 120

gagctgtagc agacctgatt gagcctttgc agcagctgag agcatggcct agggtgggcg 180

gcaccattgt ccagcagctg agtttcccag ggaccttgga gatagccgca gccctcattt 240

gcaggggaag atgattcctg ccagatttgc cggggtgctg cttgctctgg ccctcatttt 300

gccagggacc ctttgtgcag aaggaactcg cggcaggtca tccacggccc gatgcagcct 360

tttcggaagt gacttcgtca acacctttga tgggagcatg tacagctttg cgggatactg 420

cagttacctc ctggcagggg gctgccagaa acgctccttc tcgattattg gggacttcca 480

gaatggcaag agagtgagcc tctccgtgta tcttggggaa ttttttgaca tccatttgtt 540

tgtcaatggt accgtgacac agggggacca aagagtctcc atgccctatg cctccaaagg 600

gctgtatcta gaaactgagg ctgggtacta caagctgtcc ggtgaggcct atggctttgt 660

ggccaggatc gatggcagcg gcaactttca agtcctgctg tcagacagat acttcaacaa 720

gacctgcggg ctgtgtggca actttaacat ctttgctgaa gatgacttta tgacccaaga 780

agggaccttg acctcggacc cttatgactt tgccaactca tgggctctga gcagtggaga 840

acagtggtgt gaacgggcat ctcctcccag cagctcatgc aacatctcct ctggggaaat 900

gcagaagggc ctgtgggagc agtgccagct tctgaagagc acctcggtgt ttgcccgctg 960

ccaccctctg gtggaccccg agccttttgt ggccctgtgt gagaagactt tgtgtgagtg 1020

tgctgggggg ctggagtgcg cctgccctgc cctcctggag tacgcccgga cctgtgccca 1080

ggagggaatg gtgctgtacg gctggaccga ccacagcgcg tgcagcccag tgtgccctgc 1140

tggtatggag tataggcagt gtgtgtcccc ttgcgccagg acctgccaga gcctgcacat 1200

caatgaaatg tgtcaggagc gatgcgtgga tggctgcagc tgccctgagg gacagctcct 1260

ggatgaaggc ctctgcgtgg agagcaccga gtgtccctgc gtgcattccg gaaagcgcta 1320

ccctcccggc acctccctct ctcgagactg caacacctgc atttgccgaa acagccagtg 1380

gatctgcagc aatgaagaat gtccagggga gtgccttgtc acaggtcaat cacacttcaa 1440

gagctttgac aacagatact tcaccttcag tgggatctgc cagtacctgc tggcccggga 1500

ttgccaggac cactccttct ccattgtcat tgagactgtc cagtgtgctg atgaccgcga 1560

cgctgtgtgc acccgctccg tcaccgtccg gctgcctggc ctgcacaaca gccttgtgaa 1620

actgaagcat ggggcaggag ttgccatgga tggccaggac gtccagctcc ccctcctgaa 1680

aggtgacctc cgcatccagc atacagtgac ggcctccgtg cgcctcagct acggggagga 1740

cctgcagatg gactgggatg gccgcgggag gctgctggtg aagctgtccc ccgtctatgc 1800

cgggaagacc tgcggcctgt gtgggaatta caatggcaac cagggcgacg acttccttac 1860

cccctctggg ctggcggagc cccgggtgga ggacttcggg aacgcctgga agctgcacgg 1920

ggactgccag gacctgcaga agcagcacag cgatccctgc gccctcaacc cgcgcatgac 1980

caggttctcc gaggaggcgt gcgcggtcct gacgtccccc acattcgagg cctgccatcg 2040

tgccgtcagc ccgctgccct acctgcggaa ctgccgctac gacgtgtgct cctgctcgga 2100

cggccgcgag tgcctgtgcg gcgccctggc cagctatgcc gcggcctgcg cggggagagg 2160

cgtgcgcgtc gcgtggcgcg agccaggccg ctgtgagctg aactgcccga aaggccaggt 2220

gtacctgcag tgcgggaccc cctgcaacct gacctgccgc tctctctctt acccggatga 2280

ggaatgcaat gaggcctgcc tggagggctg cttctgcccc ccagggctct acatggatga 2340

gaggggggac tgcgtgccca aggcccagtg cccctgttac tatgacggtg agatcttcca 2400

gccagaagac atcttctcag accatcacac catgtgctac tgtgaggatg gcttcatgca 2460

ctgtaccatg agtggagtcc ccggaagctt gctgcctgac gctgtcctca gcagtcccct 2520

gtctcatcgc agcaaaagga gcctatcctg tcggcccccc atggtcaagc tggtgtgtcc 2580

cgctgacaac ctgcgggctg aagggctcga gtgtaccaaa acgtgccaga actatgacct 2640

ggagtgcatg agcatgggct gtgtctctgg ctgcctctgc cccccgggca tggtccggca 2700

tgagaacaga tgtgtggccc tggaaaggtg tccctgcttc catcagggca aggagtatgc 2760

ccctggagaa acagtgaaga ttggctgcaa cacttgtgtc tgtcgggacc ggaagtggaa 2820

ctgcacagac catgtgtgtg atgccacgtg ctccacgatc ggcatggccc actacctcac 2880

cttcgacggg ctcaaatacc tgttccccgg ggagtgccag tacgttctgg tgcaggatta 2940

ctgcggcagt aaccctggga cctttcggat cctagtgggg aataagggat gcagccaccc 3000

ctcagtgaaa tgcaagaaac gggtcaccat cctggtggag ggaggagaga ttgagctgtt 3060

tgacggggag gtgaatgtga agaggcccat gaaggatgag actcactttg aggtggtgga 3120

gtctggccgg tacatcattc tgctgctggg caaagccctc tccgtggtct gggaccgcca 3180

cctgagcatc tccgtggtcc tgaagcagac ataccaggag aaagtgtgtg gcctgtgtgg 3240

gaattttgat ggcatccaga acaatgacct caccagcagc aacctccaag tggaggaaga 3300

ccctgtggac tttgggaact cctggaaagt gagctcgcag tgtgctgaca ccagaaaagt 3360

gcctctggac tcatccctg ccacctgcca taacaacatc atgaagcaga cgatggtgga 3420

ttcctcctgt agaatcctta ccagtgacgt cttccaggac tgcaacaagc tggtggaccc 3480

cgagccatat ctggatgtct gcatttacga cacctgctcc tgtgagtcca ttggggactg 3540

cgcctgcttc tgcgacacca ttgctgccta tgcccacgtg tgtgcccagc atggcaaggt 3600

ggtgacctgg aggacggcca cattgtgccc ccagagctgc gaggagagga atctccggga 3660

gaacgggtat gagtgtgagt ggcgctataa cagctgtgca cctgcctgtc aagtcacgtg 3720

tcagcaccct gagccactgg cctgccctgt gcagtgtgtg gagggctgcc atgcccactg 3780

ccctccaggg aaaatcctgg atgagctttt gcagacctgc gttgaccctg aagactgtcc 3840

agtgtgtgag gtggctggcc ggcgttttgc ctcaggaaag aaagtcacct tgaatcccag 3900

tgaccctgag cactgccaga tttgccactg tgatgttgtc aacctcacct gtgaagcctg 3960

ccaggagccg ggaggcctgg tggtgcctcc cacagatgcc ccggtgagcc ccaccactct 4020

gtatgtggag gacatctcgg aaccgccgtt gcacgatttc tactgcagca ggctactgga 4080

cctggtcttc ctgctggatg gctcctccag gctgtccgag gctgagtttg aagtgctgaa 4140

ggcctttgtg gtggacatga tggagcggct gcgcatctcc cagaagtggg tccgcgtggc 4200

cgtggtggag taccacgacg gctcccacgc ctacatcggg ctcaaggacc ggaagcgacc 4260

gtcagagctg cggcgcattg ccagccaggt gaagtatgcg ggcagccagg tggcctccac 4320

cagcgaggtc ttgaaataca cactgttcca aatcttcagc aagatcgacc gccctgaagc 4380

ctcccgcatc accctgctcc tgatggccag ccaggagccc caacggatgt cccggaactt 4440

tgtccgctac gtccagggcc tgaagaagaa gaaggtcatt gtgatcccgg tgggcattgg 4500

gccccatgcc aacctcaagc agatccgcct catcgagaag caggcccctg agaacaaggc 4560

cttcgtgctg agcagtgtgg atgagctgga gcagcaaagg gacgagatcg ttagctacct 4620

ctgtgacctt gcccctgaag ccctcctctc tactctgccc cccgacatgg cacaagtcac 4680

tgtgggcccg gggctcttgg gggtttcgac cctggggccc aagaggaact ccatggttct 4740

ggatgtggcg ttcgtcctgg aaggatcgga caaaattggt gaagccgact tcaacaggag 4800

caaggagttc atggaggagg tgattcagcg gatggatgtg ggccaggaca gcatccacgt 4860

cacggtgctg cagtactcct acatggtgac tgtggagtac cccttcagcg aggcacagtc 4920

caaaggggac atcctgcagc gggtgcgaga gatccgctac cagggcggca acaggaccaa 4980

cactgggctg gccctgcggt acctctctga ccacagcttc ttggtcagcc agggtgaccg 5040

ggagcaggcg cccaacctgg tctacatggt caccggaaat cctgcctctg atgagatcaa 5100

gaggctgcct ggagacatcc aggtggtgcc cattggagtg ggccctaatg ccaacgtgca 5160

ggagctggag aggattggct ggcccaatgc ccctatcctc atccaggact ttgagacgct 5220

cccccgagag gctcctgacc tggtgctgca gaggtgctgc tccggagagg ggctgcagat 5280

ccccaccctc tcccctgcac ctgactgcag ccagcccctg gacgtgatcc ttctcctgga 5340

tggctcctcc agtttcccag cttcttattt tgatgaaatg aagagtttcg ccaaggcttt 5400

catttcaaaa gccaatatag ggcctcgtct cactcaggtg tcagtgctgc agtatggaag 5460

catcaccacc attgacgtgc catggaacgt ggtcccggag aaagcccatt tgctgagcct 5520

tgtggacgtc atgcagcggg agggaggccc cagccaaatc ggggatgcct tgggctttgc 5580

tgtgcgatac ttgacttcag aaatgcatgg tgccaggccg ggagcctcaa aggcggtggt 5640

catcctggtc acggacgtct ctgtggattc agtggatgca gcagctgatg ccgccaggtc 5700

caacagagtg acagtgttcc ctattggaat tggagatcgc tacgatgcag cccagctacg 5760

gatcttggca ggcccagcag gcgactccaa cgtggtgaag ctccagcgaa tcgaagacct 5820

ccctaccatg gtcaccttgg gcaattcctt cctccacaaa ctgtgctctg gatttgttag 5880

gatttgcatg gatgaggatg ggaatgagaa gaggcccggg gacgtctgga ccttgccaga 5940

ccagtgccac accgtgactt gccagccaga tggccagacc ttgctgaaga gtcatcgggt 6000

caactgtgac cgggggctga ggccttcgtg ccctaacagc cagtcccctg ttaaagtgga 6060

agagacctgt ggctgccgct ggacctgccc ctgcgtgtgc acaggcagct ccactcggca 6120

catcgtgacc tttgatgggc agaatttcaa gctgactggc agctgttctt atgtcctatt 6180

tcaaaacaag gagcaggacc tggaggtgat tctccataat ggtgcctgca gccctggagc 6240

aaggcagggc tgcatgaaat ccatcgaggt gaagcacagt gccctctccg tcgagctgca 6300

cagtgacatg gaggtgacgg tgaatgggag actggtctct gttccttacg tgggtgggaa 6360

catggaagtc aacgtttatg gtgccatcat gcatgaggtc agattcaatc accttggtca 6420

catcttcaca ttcactccac aaaacaatga gttccaactg cagctcagcc ccaagacttt 6480

tgcttcaaag acgtatggtc tgtgtgggat ctgtgatgag aacggagcca atgacttcat 6540

gctgagggat ggcacagtca ccacagactg gaaaacactt gttcaggaat ggactgtgca 6600

gcggccaggg cagacgtgcc agcccatcct ggaggagcag tgtcttgtcc ccgacagctc 6660

ccactgccag gtcctcctct taccactgtt tgctgaatgc cacaaggtcc tggctccagc 6720

cacattctat gccatctgcc agcaggacag ttgccaccag gagcaagtgt gtgaggtgat 6780

cgcctcttat gcccacctct gtcggaccaa cggggtctgc gttgactgga ggacacctga 6840

tttctgtgct atgtcatgcc caccatctct ggtctacaac cactgtgagc atggctgtcc 6900

ccggcactgt gatggcaacg tgagctcctg tggggaccat ccctccgaag gctgtttctg 6960

ccctccagat aaagtcatgt tggaaggcag ctgtgtccct gaagaggcct gcactcagtg 7020

cattggtgag gatggagtcc agcaccagtt cctggaagcc tgggtcccgg accaccagcc 7080

ctgtcagatc tgcacatgcc tcagcgggcg gaaggtcaac tgcacaacgc agccctgccc 7140

cacggccaaa gctcccacgt gtggcctgtg tgaagtagcc cgcctccgcc agaatgcaga 7200

ccagtgctgc cccgagtatg agtgtgtgtg tgacccagtg agctgtgacc tgcccccagt 7260

gcctcactgt gaacgtggcc tccagcccac actgaccaac cctggcgagt gcagacccaa 7320

cttcacctgc gcctgcagga aggaggagtg caaaagagtg tccccaccct cctgcccccc 7380

gcaccgtttg cccacccttc ggaagaccca gtgctgtgat gagtatgagt gtgcctgcaa 7440

ctgtgtcaac tccacagtga gctgtcccct tgggtacttg gcctcaactg ccaccaatga 7500

ctgtggctgt accacaacca cctgccttcc cgacaaggtg tgtgtccacc gaagcaccat 7560

ctaccctgtg ggccagttct gggaggaggg ctgcgatgtg tgcacctgca ccgacatgga 7620

ggatgccgtg atgggcctcc gcgtggccca gtgctcccag aagccctgtg aggacagctg 7680

tcggtcgggc ttcacttacg ttctgcatga aggcgagtgc tgtggaaggt gcctgccatc 7740

tgcctgtgag gtggtgactg gctcaccgcg gggggactcc cagtcttcct ggaagagtgt 7800

cggctcccag tgggcctccc cggagaaccc ctgcctcatc aatgagtgtg tccgagtgaa 7860

ggaggaggtc tttatacaac aaaggaacgt ctcctgcccc cagctggagg tccctgtctg 7920

cccctcgggc tttcagctga gctgtaagac ctcagcgtgc tgcccaagct gtcgctgtga 7980

gcgcatggag gcctgcatgc tcaatggcac tgtcattggg cccgggaaga ctgtgatgat 8040

cgatgtgtgc acgacctgcc gctgcatggt gcaggtgggg gtcatctctg gattcaagct 8100

ggagtgcagg aagaccacct gcaacccctg ccccctgggt tacaaggaag aaaataacac 8160

aggtgaatgt tgtgggagat gtttgcctac ggcttgcacc attcagctaa gaggaggaca 8220

gatcatgaca ctgaagcgtg atgagacgct ccaggatggc tgtgatactc acttctgcaa 8280

ggtcaatgag agaggagagt acttctggga gaagagggtc acaggctgcc cacccttga 8340

tgaacacaag tgtctggctg agggaggtaa aattatgaaa attccaggca cctgctgtga 8400

cacatgtgag gagcctgagt gcaacgacat cactgccagg ctgcagtatg tcaaggtggg 8460

aagctgtaag tctgaagtag aggtggatat ccactactgc cagggcaaat gtgccagcaa 8520

agccatgtac tccattgaca tcaacgatgt gcaggaccag tgctcctgct gctctccgac 8580

acggacggag cccatgcagg tggccctgca ctgcaccaat ggctctgttg tgtaccatga 8640

ggttctcaat gccatggagt gcaaatgctc ccccaggaag tgcagcaagt gaggctgctg 8700

cagctgcatg ggtgcctgct gctgcctgcc ttggcctgat ggccaggcca gagtgctgcc 8760

agtcctctgc atgttctgct cttgtgccct tctgagccca caataaaggc tgagctctta 8820

tcttgcaaaa ggc 8833

<210> 2
<211> 2783
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Polypeptide

<220>
<221> MISC_FEATURE
<223> prepro-vWF

<400> 2

```
Met Ile Pro Ala Arg Phe Ala Gly Val Leu Leu Leu Ile Leu Pro Gly
1               5                  10                  15

Thr Leu Cys Ala Glu Gly Thr Arg Gly Arg Ser Ser Thr Ala Arg Cys
            20                  25                  30

Ser Leu Phe Gly Ser Asp Phe Val Asn Thr Phe Asp Gly Ser Met Tyr
        35                  40                  45

Ser Phe Ala Gly Tyr Cys Ser Tyr Leu Leu Ala Gly Gly Cys Gln Lys
    50                  55                  60

Arg Ser Phe Ser Ile Ile Gly Asp Phe Gln Asn Gly Lys Arg Val Ser
65                  70                  75                  80

Leu Ser Val Tyr Leu Gly Glu Phe Phe Asp Ile His Leu Phe Val Asn
                85                  90                  95

Gly Thr Val Thr Gln Gly Asp Gln Arg Val Ser Met Pro Tyr Ala Ser
            100                 105                 110

Lys Leu Glu Thr Glu Ala Gly Tyr Tyr Lys Leu Ser Gly Glu Ala Tyr
        115                 120                 125

Gly Phe Val Ala Arg Ile Asp Gly Ser Gly Asn Phe Gln Val Leu Leu
    130                 135                 140

Ser Asp Arg Tyr Phe Asn Lys Thr Cys Gly Leu Cys Gly Asn Phe Asn
145                 150                 155                 160

Ile Phe Ala Glu Asp Asp Phe Met Thr Gln Glu Gly Thr Leu Thr Ser
                165                 170                 175

Asp Pro Tyr Asp Phe Ala Asn Ser Trp Ala Leu Ser Ser Gly Glu Gln
            180                 185                 190

Trp Cys Glu Arg Pro Ser Ser Ser Cys Asn Ile Ser Ser Gly Glu Met
        195                 200                 205

Gln Lys Gly Leu Trp Glu Gln Cys Gln Leu Leu Lys Ser Thr Ser Val
    210                 215                 220

Phe Ala Arg Cys His Pro Leu Val Asp Pro Glu Pro Phe Cys Glu Lys
225                 230                 235                 240

Thr Leu Cys Glu Cys Ala Gly Gly Leu Glu Cys Ala Cys Pro Ala Leu
                245                 250                 255

Leu Glu Tyr Ala Arg Thr Cys Ala Gln Glu Gly Met Val Leu Tyr Gly
            260                 265                 270
```

```
Trp Thr Asp His Ser Ala Cys Ser Pro Val Cys Pro Ala Gly Met Glu
        275                 280                 285

Tyr Arg Gln Cys Val Ser Pro Cys Ala Arg Thr Cys Gln Ser Leu His
    290                 295                 300

Ile Asn Glu Met Cys Gln Glu Arg Cys Val Asp Gly Cys Ser Cys Pro
305                 310                 315                 320

Glu Gly Gln Leu Leu Asp Glu Gly Leu Cys Val Glu Ser Thr Glu Cys
                325                 330                 335

Pro Cys Val His Ser Gly Lys Arg Tyr Pro Pro Gly Thr Ser Leu Ser
            340                 345                 350

Arg Asp Cys Asn Thr Cys Ile Cys Arg Asn Ser Gln Trp Ile Cys Ser
        355                 360                 365

Asn Glu Glu Cys Pro Gly Glu Cys Leu Val Thr Gly Gln Ser His Phe
    370                 375                 380

Lys Ser Phe Asp Asn Arg Tyr Phe Thr Phe Ser Gly Ile Cys Gln Tyr
385                 390                 395                 400

Leu Leu Ala Arg Asp Cys Gln Asp His Ser Phe Ser Ile Val Ile Glu
                405                 410                 415

Thr Val Gln Cys Ala Asp Asp Arg Asp Ala Val Cys Thr Arg Ser Val
            420                 425                 430

Thr Val Arg Leu Pro Gly Leu His Asn Ser Leu Val Lys Leu Lys His
        435                 440                 445

Gly Ala Gly Val Ala Met Asp Gly Gln Asp Val Gln Leu Pro Leu Leu
    450                 455                 460

Lys Gly Asp Leu Arg Ile Gln His Thr Val Thr Ala Ser Val Arg Leu
465                 470                 475                 480

Ser Tyr Gly Glu Asp Leu Gln Met Asp Trp Asp Gly Arg Gly Arg Leu
                485                 490                 495

Leu Val Lys Leu Ser Pro Val Tyr Ala Gly Lys Thr Cys Gly Leu Cys
            500                 505                 510

Gly Asn Tyr Asn Gly Asn Gln Gly Asp Asp Phe Leu Thr Pro Ser Gly
        515                 520                 525

Leu Ala Glu Pro Arg Val Glu Asp Phe Gly Asn Ala Trp Lys Leu His
    530                 535                 540
```

```
Gly Asp Cys Gln Asp Leu Gln Lys Gln His Ser Asp Pro Cys Ala Leu
545                 550                 555                 560

Asn Pro Arg Met Thr Arg Phe Ser Glu Glu Ala Cys Ala Val Leu Thr
                565                 570                 575

Ser Pro Thr Phe Glu Ala Cys His Arg Ala Val Ser Pro Leu Pro Tyr
            580                 585                 590

Leu Arg Asn Cys Arg Tyr Asp Val Cys Ser Cys Ser Asp Gly Arg Glu
        595                 600                 605

Cys Leu Cys Gly Ser Tyr Ala Ala Ala Cys Ala Gly Arg Gly Val Arg
    610                 615                 620

Val Ala Trp Arg Glu Pro Gly Arg Cys Glu Leu Asn Cys Pro Lys Gly
625                 630                 635                 640

Gln Val Tyr Leu Gln Cys Gly Thr Pro Cys Asn Leu Thr Cys Arg Ser
                645                 650                 655

Leu Ser Tyr Pro Asp Glu Glu Cys Asn Glu Ala Cys Leu Glu Gly Cys
            660                 665                 670

Phe Cys Pro Pro Met Asp Glu Arg Gly Asp Cys Val Pro Lys Ala Gln
        675                 680                 685

Cys Pro Cys Tyr Tyr Asp Gly Glu Ile Phe Gln Pro Glu Asp Ile Phe
    690                 695                 700

Ser Asp His His Thr Met Cys Tyr Cys Glu Asp Gly Phe Met His Cys
705                 710                 715                 720

Thr Met Ser Gly Val Pro Gly Ser Leu Leu Pro Asp Ala Val Leu Ser
                725                 730                 735

Ser Pro Leu Ser His Arg Ser Lys Arg Ser Leu Ser Cys Arg Pro Pro
            740                 745                 750

Met Val Lys Leu Val Cys Pro Ala Asp Asn Leu Arg Ala Glu Gly Leu
        755                 760                 765

Glu Cys Thr Lys Thr Cys Gln Asn Tyr Asp Leu Glu Cys Met Ser Met
    770                 775                 780

Gly Cys Val Ser Gly Cys Leu Cys Pro Pro Gly Met Val Arg His Glu
785                 790                 795                 800

Asn Arg Cys Glu Arg Cys Pro Cys Phe His Gln Gly Lys Glu Tyr Ala
                805                 810                 815
```

```
Pro Gly Glu Thr Val Lys Ile Gly Cys Asn Thr Cys Val Cys Arg Asp
            820                 825                 830

Arg Lys Trp Asn Cys Thr Asp His Val Cys Asp Ala Thr Cys Ser Thr
        835                 840                 845

Ile Gly Met Ala His Tyr Leu Thr Phe Asp Gly Leu Lys Tyr Leu Phe
    850                 855                 860

Pro Gly Glu Cys Gln Tyr Val Leu Val Gln Asp Tyr Cys Gly Ser Asn
865                 870                 875                 880

Pro Gly Thr Phe Arg Ile Leu Val Gly Asn Lys Gly Cys Ser His Pro
                885                 890                 895

Ser Val Lys Cys Lys Lys Arg Val Thr Ile Leu Val Glu Gly Gly Glu
            900                 905                 910

Ile Glu Leu Phe Asp Gly Glu Val Asn Val Lys Arg Pro Met Lys Asp
        915                 920                 925

Glu Thr His Phe Glu Val Val Glu Ser Gly Arg Tyr Ile Ile Leu Leu
    930                 935                 940

Leu Gly Lys Ala Leu Ser Val Val Trp Asp Arg His Leu Ser Ile Ser
945                 950                 955                 960

Val Val Leu Lys Gln Thr Tyr Gln Glu Lys Val Cys Gly Leu Cys Gly
                965                 970                 975

Asn Phe Asp Gly Ile Gln Asn Asn Asp Leu Thr Ser Ser Asn Leu Gln
            980                 985                 990

Val Glu Glu Asp Pro Val Asp Phe  Gly Asn Ser Trp Lys  Val Ser Ser
        995                 1000                1005

Gln Cys  Ala Asp Thr Arg Lys  Val Pro Leu Asp Ser  Ser Pro Ala
    1010                1015                1020

Thr Cys  His Asn Asn Ile Met  Lys Gln Thr Met Val  Asp Ser Ser
    1025                1030                1035

Cys Arg  Ile Leu Thr Ser Asp  Val Phe Gln Asp Cys  Asn Lys Leu
    1040                1045                1050

Val Asp  Pro Glu Pro Tyr Leu  Asp Val Cys Ile Tyr  Asp Thr Cys
    1055                1060                1065

Ser Cys  Glu Ser Ile Gly Asp  Cys Ala Cys Phe Cys  Asp Thr Ile
    1070                1075                1080
```

```
Ala Ala  Tyr Ala His Val Cys  Ala Gln His Gly Lys  Val Val Thr
    1085                 1090                 1095

Trp Arg  Thr Ala Thr Leu Cys  Pro Gln Ser Cys Glu  Glu Arg Asn
    1100                 1105                 1110

Leu Arg  Glu Asn Gly Tyr Glu  Cys Glu Trp Arg Tyr  Asn Ser Cys
    1115                 1120                 1125

Ala Pro  Ala Cys Gln Val Thr  Cys Gln His Pro Glu  Pro Leu Ala
    1130                 1135                 1140

Cys Pro  Val Gln Cys Val Glu  Gly Cys His Ala His  Cys Pro Pro
    1145                 1150                 1155

Gly Lys  Ile Leu Asp Glu Leu  Leu Gln Thr Cys Val  Asp Pro Glu
    1160                 1165                 1170

Asp Cys  Pro Val Cys Glu Val  Ala Gly Arg Arg Phe  Ala Ser Gly
    1175                 1180                 1185

Lys Lys  Val Thr Leu Asn Pro  Ser Asp Pro Glu His  Cys Gln Ile
    1190                 1195                 1200

Cys His  Cys Asp Val Val Asn  Leu Thr Cys Glu Ala  Cys Gln Glu
    1205                 1210                 1215

Pro Gly  Gly Leu Val Val Pro  Pro Thr Asp Ala Pro  Val Ser Pro
    1220                 1225                 1230

Thr Thr  Leu Tyr Val Glu Asp  Ile Ser Glu Pro Pro  Leu His Asp
    1235                 1240                 1245

Phe Tyr  Cys Ser Arg Leu Leu  Asp Leu Val Phe Leu  Leu Asp Gly
    1250                 1255                 1260

Ser Ser  Arg Leu Ser Glu Ala  Glu Phe Glu Val Leu  Lys Ala Phe
    1265                 1270                 1275

Val Val  Asp Met Met Glu Arg  Leu Arg Ile Ser Gln  Lys Trp Val
    1280                 1285                 1290

Arg Val  Ala Val Val Glu Tyr  His Asp Gly Ser His  Ala Tyr Ile
    1295                 1300                 1305

Gly Leu  Lys Asp Arg Lys Arg  Pro Ser Glu Leu Arg  Arg Ile Ala
    1310                 1315                 1320

Ser Gln  Val Lys Tyr Ala Gly  Ser Gln Val Ala Ser  Thr Ser Glu
    1325                 1330                 1335
```

Val Leu Lys Tyr Thr Leu Phe Gln Ile Phe Ser Lys Ile Asp Arg
1340 1345 1350

Pro Glu Ala Ser Arg Ile Thr Leu Leu Leu Met Ala Ser Gln Glu
1355 1360 1365

Pro Gln Arg Met Ser Arg Asn Phe Val Arg Tyr Val Gln Gly Leu
1370 1375 1380

Lys Lys Lys Lys Val Ile Val Ile Pro Val Gly Ile Gly Pro His
1385 1390 1395

Ala Asn Leu Lys Gln Ile Arg Leu Ile Glu Lys Gln Ala Pro Glu
1400 1405 1410

Asn Lys Ala Phe Val Leu Ser Ser Val Asp Glu Leu Glu Gln Gln
1415 1420 1425

Arg Asp Glu Ile Val Ser Tyr Leu Cys Asp Leu Ala Pro Glu Ala
1430 1435 1440

Pro Pro Pro Thr Leu Pro Pro Asp Met Ala Gln Val Thr Val Gly
1445 1450 1455

Pro Gly Leu Leu Gly Val Ser Thr Leu Gly Pro Lys Arg Asn Ser
1460 1465 1470

Met Val Leu Asp Val Ala Phe Val Leu Glu Gly Ser Asp Lys Ile
1475 1480 1485

Gly Glu Ala Asp Phe Asn Arg Ser Lys Glu Phe Met Glu Glu Val
1490 1495 1500

Ile Gln Arg Met Asp Val Gly Gln Asp Ser Ile His Val Thr Val
1505 1510 1515

Leu Gln Tyr Ser Tyr Met Val Thr Val Glu Tyr Pro Phe Ser Glu
1520 1525 1530

Ala Gln Ser Lys Gly Asp Ile Leu Gln Arg Val Arg Glu Ile Arg
1535 1540 1545

Tyr Gln Gly Gly Asn Arg Thr Asn Thr Gly Leu Ala Leu Arg Tyr
1550 1555 1560

Leu Ser Asp His Ser Phe Leu Val Ser Gln Gly Asp Arg Glu Gln
1565 1570 1575

Ala Pro Asn Leu Val Tyr Met Val Thr Gly Asn Pro Ala Ser Asp
1580 1585 1590

```
Glu Ile  Lys Arg Leu Pro Gly  Asp Ile Gln Val Val  Pro Ile Gly
    1595                 1600                 1605

Val Gly  Pro Asn Ala Asn Val  Gln Glu Leu Glu Arg  Ile Gly Trp
    1610                 1615                 1620

Pro Asn  Ala Pro Ile Leu Ile  Gln Asp Phe Glu Thr  Leu Pro Arg
    1625                 1630                 1635

Glu Ala  Pro Asp Leu Val Leu  Gln Arg Cys Cys Ser  Gly Glu Gly
    1640                 1645                 1650

Leu Gln  Ile Pro Thr Leu Ser  Pro Ala Pro Asp Cys  Ser Gln Pro
    1655                 1660                 1665

Leu Asp  Val Ile Leu Leu Leu  Asp Gly Ser Ser Ser  Phe Pro Ala
    1670                 1675                 1680

Ser Tyr  Phe Asp Glu Met Lys  Ser Phe Ala Lys Ala  Phe Ile Ser
    1685                 1690                 1695
```

```
Lys Ala  Asn Ile Gly Pro Arg  Leu Thr Gln Val Ser  Val Leu Gln
    1700                 1705                 1710

Tyr Gly  Ser Ile Thr Thr Ile  Asp Val Pro Trp Asn  Val Val Pro
    1715                 1720                 1725

Glu Lys  Ala His Leu Leu Ser  Leu Val Asp Val Met  Gln Arg Glu
    1730                 1735                 1740

Gly Gly  Pro Ser Gln Ile Gly  Asp Ala Leu Gly Phe  Ala Val Arg
    1745                 1750                 1755

Tyr Leu  Thr Ser Glu Met His  Gly Ala Arg Pro Gly  Ala Ser Lys
    1760                 1765                 1770

Ala Val  Val Ile Leu Val Thr  Asp Val Ser Val Asp  Ser Val Asp
    1775                 1780                 1785

Ala Ala  Ala Asp Ala Ala Arg  Ser Asn Arg Val Thr  Val Phe Pro
    1790                 1795                 1800

Ile Gly  Ile Gly Asp Arg Tyr  Asp Ala Ala Gln Leu  Arg Ile Leu
    1805                 1810                 1815

Ala Gly  Pro Ala Gly Asp Ser  Asn Val Val Lys Leu  Gln Arg Ile
    1820                 1825                 1830

Glu Asp  Leu Pro Thr Met Val  Thr Leu Gly Asn Ser  Phe Leu His
    1835                 1840                 1845
```

Lys Leu Cys Ser Gly Phe Val Arg Ile Cys Met Asp Glu Asp Gly
1850 1855 1860

Asn Glu Lys Arg Pro Gly Asp Val Trp Thr Leu Pro Asp Gln Cys
1865 1870 1875

His Thr Val Thr Cys Gln Pro Asp Gly Gln Thr Leu Leu Lys Ser
1880 1885 1890

His Arg Val Asn Cys Asp Arg Gly Leu Arg Pro Ser Cys Pro Asn
1895 1900 1905

Ser Gln Ser Pro Val Lys Val Glu Glu Thr Cys Gly Cys Arg Trp
1910 1915 1920

Thr Cys Pro Cys Val Cys Thr Gly Ser Ser Thr Arg His Ile Val
1925 1930 1935

Thr Phe Asp Gly Gln Asn Phe Lys Leu Thr Gly Ser Cys Ser Tyr
1940 1945 1950

Val Leu Phe Gln Asn Lys Glu Gln Asp Leu Glu Val Ile Leu His
1955 1960 1965

Asn Gly Ala Cys Ser Pro Gly Ala Arg Gln Gly Cys Met Lys Ser
1970 1975 1980

Ile Glu Val Lys His Ser Ala Leu Ser Val Glu Leu His Ser Asp
1985 1990 1995

Met Glu Val Thr Val Asn Gly Arg Leu Val Ser Val Pro Tyr Val
2000 2005 2010

Gly Gly Asn Met Glu Val Asn Val Tyr Gly Ala Ile Met His Glu
2015 2020 2025

Val Arg Phe Asn His Leu Gly His Ile Phe Thr Phe Thr Pro Gln
2030 2035 2040

Asn Asn Glu Phe Gln Leu Gln Leu Ser Pro Lys Thr Phe Ala Ser
2045 2050 2055

Lys Thr Tyr Gly Leu Cys Gly Ile Cys Asp Glu Asn Gly Ala Asn
2060 2065 2070

Asp Phe Met Leu Arg Asp Gly Thr Val Thr Thr Asp Trp Lys Thr
2075 2080 2085

Leu Val Gln Glu Trp Thr Val Gln Arg Pro Gly Gln Thr Cys Gln
2090 2095 2100

```
Pro Glu  Gln Cys Leu Val Pro  Asp Ser Ser His Cys  Gln Val Leu
    2105                 2110                 2115

Leu Leu  Pro Leu Phe Ala Glu  Cys His Lys Val Leu  Ala Pro Ala
    2120                 2125                 2130

Thr Phe  Tyr Ala Ile Cys Gln  Gln Asp Ser Cys His  Gln Glu Gln
    2135                 2140                 2145

Val Cys  Glu Val Ile Ala Ser  Tyr Ala His Leu Cys  Arg Thr Asn
    2150                 2155                 2160

Gly Val  Cys Val Asp Trp Arg  Thr Pro Asp Phe Cys  Ala Met Ser
    2165                 2170                 2175

Cys Pro  Pro Ser Leu Val Tyr  Asn His Cys Glu His  Gly Cys Pro
    2180                 2185                 2190

Arg His  Cys Asp Gly Asn Val  Ser Ser Cys Gly Asp  His Pro Ser
    2195                 2200                 2205

Glu Gly  Cys Phe Cys Pro Pro  Asp Lys Val Met Leu  Glu Gly Ser
    2210                 2215                 2220

Cys Val  Pro Glu Glu Ala Cys  Thr Gln Cys Ile Gly  Glu Asp Gly
    2225                 2230                 2235

Val Gln  His Gln Phe Leu Glu  Ala Trp Val Pro Asp  His Gln Pro
    2240                 2245                 2250

Cys Gln  Ile Cys Thr Cys Leu  Ser Gly Arg Lys Val  Asn Cys Thr
    2255                 2260                 2265

Thr Gln  Pro Cys Pro Thr Ala  Lys Ala Pro Thr Cys  Gly Leu Cys
    2270                 2275                 2280

Glu Val  Ala Arg Leu Arg Gln  Asn Ala Asp Gln Cys  Cys Pro Glu
    2285                 2290                 2295

Tyr Glu  Cys Val Cys Asp Pro  Val Ser Cys Asp Leu  Pro Pro Val
    2300                 2305                 2310

Pro His  Cys Glu Arg Gly Leu  Gln Pro Thr Leu Thr  Asn Pro Gly
    2315                 2320                 2325

Glu Cys  Arg Pro Asn Phe Thr  Cys Ala Cys Arg Lys  Glu Glu Cys
    2330                 2335                 2340

Lys Arg  Val Ser Pro Pro Ser  Cys Pro Pro His Arg  Leu Pro Thr
    2345                 2350                 2355
```

```
Leu Arg  Lys Thr Gln Cys Cys  Asp Glu Tyr Glu Cys  Ala Cys Asn
    2360                 2365                 2370

Cys Val  Asn Ser Thr Val Ser  Cys Pro Leu Gly Tyr  Leu Ala Ser
    2375                 2380                 2385

Thr Ala  Thr Asn Asp Cys Gly  Cys Thr Thr Thr Thr  Cys Leu Pro
    2390                 2395                 2400

Asp Lys  Val Cys Val His Arg  Ser Thr Ile Tyr Pro  Val Gly Gln
    2405                 2410                 2415

Phe Trp  Glu Glu Gly Cys Asp  Val Cys Thr Cys Thr  Asp Met Glu
    2420                 2425                 2430

Asp Ala  Val Met Gly Leu Arg  Val Ala Gln Cys Ser  Gln Lys Pro
    2435                 2440                 2445

Cys Glu  Asp Ser Cys Arg Ser  Gly Phe Thr Tyr Val  Leu His Glu
    2450                 2455                 2460

Gly Glu  Cys Cys Gly Arg Cys  Leu Pro Ser Ala Cys  Glu Val Val
    2465                 2470                 2475

Thr Gly  Ser Pro Arg Gly Asp  Ser Gln Ser Ser Trp  Lys Ser Val
    2480                 2485                 2490

Gly Ser  Gln Trp Glu Asn Pro  Cys Leu Ile Asn Glu  Cys Val Arg
    2495                 2500                 2505

Val Lys  Glu Glu Val Phe Ile  Gln Gln Arg Asn Val  Ser Cys Pro
    2510                 2515                 2520

Gln Leu  Glu Val Pro Val Cys  Pro Ser Gly Phe Gln  Leu Ser Cys
    2525                 2530                 2535

Lys Thr  Ser Ala Cys Cys Pro  Ser Cys Arg Cys Glu  Arg Met Glu
    2540                 2545                 2550

Ala Cys  Met Leu Asn Gly Thr  Val Ile Gly Pro Gly  Lys Thr Val
    2555                 2560                 2565

Met Ile  Asp Val Cys Thr Thr  Cys Arg Cys Met Val  Gln Val Gly
    2570                 2575                 2580

Val Ile  Ser Gly Phe Lys Leu  Glu Cys Arg Lys Thr  Thr Cys Asn
    2585                 2590                 2595

Pro Cys  Pro Leu Gly Tyr Lys  Glu Glu Asn Asn Thr  Gly Glu Cys
    2600                 2605                 2610
```

```
Cys Gly  Arg Cys Leu Pro Thr  Ala Cys Thr Ile Gln  Leu Arg Gly
    2615                 2620                 2625

Gly Gln  Ile Met Thr Leu Lys  Arg Asp Glu Thr Leu  Gln Asp Gly
    2630                 2635                 2640

Cys Asp  Thr His Phe Cys Lys  Val Asn Glu Arg Gly  Glu Tyr Phe
    2645                 2650                 2655

Trp Glu  Lys Arg Val Thr Gly  Cys Pro Pro Phe Asp  Glu His Lys
    2660                 2665                 2670

Cys Leu  Ala Glu Gly Gly Lys  Ile Met Lys Ile Pro  Gly Thr Cys
    2675                 2680                 2685

Cys Asp  Thr Cys Glu Glu Pro  Glu Cys Asn Asp Ile  Thr Ala Arg
    2690                 2695                 2700

Leu Gln  Tyr Val Lys Val Gly  Ser Cys Lys Ser Glu  Val Glu Val
    2705                 2710                 2715
```

```
Asp Ile  His Tyr Cys Gln Gly  Lys Cys Ala Ser Lys  Ala Met Tyr
    2720                 2725                 2730

Ser Ile  Asp Ile Asn Asp Val  Gln Asp Gln Cys Ser  Cys Cys Ser
    2735                 2740                 2745

Pro Thr  Arg Thr Glu Pro Met  Gln His Cys Thr Asn  Gly Ser Val
    2750                 2755                 2760

Val Tyr  His Glu Val Leu Asn  Ala Met Glu Cys Lys  Cys Ser Pro
    2765                 2770                 2775

Arg Lys  Cys Ser Lys
    2780
```

<210> 3
<211> 2050
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Polypeptide

<220>
<221> MISC_FEATURE
<223> mature vWF

<400> 3

```
Ser Leu Ser Cys Arg Pro Pro Met Val Lys Leu Val Cys Pro Ala Asp
1               5                  10                  15
```

```
Asn Leu Arg Ala Glu Gly Leu Glu Cys Thr Lys Thr Cys Gln Asn Tyr
            20                  25                  30

Asp Leu Glu Cys Met Ser Met Gly Cys Val Ser Gly Cys Leu Cys Pro
        35                  40                  45

Pro Gly Met Val Arg His Glu Asn Arg Cys Val Ala Leu Glu Arg Cys
    50                  55                  60

Pro Cys Phe His Gln Gly Lys Glu Tyr Ala Pro Gly Glu Thr Val Lys
65                  70                  75                  80

Ile Gly Cys Asn Thr Cys Val Cys Arg Asp Arg Lys Trp Asn Cys Thr
                85                  90                  95

Asp His Val Cys Asp Ala Thr Cys Ser Thr Ile Gly Met Ala His Tyr
            100                 105                 110

Leu Thr Phe Asp Gly Leu Lys Tyr Leu Phe Pro Gly Glu Cys Gln Tyr
        115                 120                 125

Val Leu Val Gln Asp Tyr Cys Gly Ser Asn Pro Gly Thr Phe Arg Ile
    130                 135                 140

Leu Val Gly Asn Lys Gly Cys Ser His Pro Ser Val Lys Cys Lys Lys
145                 150                 155                 160

Arg Val Thr Ile Leu Val Glu Gly Gly Glu Ile Glu Leu Phe Asp Gly
                165                 170                 175

Glu Val Asn Val Lys Arg Pro Met Lys Asp Glu Thr His Phe Glu Val
            180                 185                 190

Val Glu Ser Gly Arg Tyr Ile Ile Leu Leu Leu Gly Lys Ala Leu Ser
        195                 200                 205

Val Val Trp Asp Arg His Leu Ser Ile Ser Val Val Leu Lys Gln Thr
    210                 215                 220

Tyr Gln Glu Lys Val Cys Gly Leu Cys Gly Asn Phe Asp Gly Ile Gln
225                 230                 235                 240

Asn Asn Asp Leu Thr Ser Ser Asn Leu Gln Val Glu Glu Asp Pro Val
                245                 250                 255

Asp Phe Gly Asn Ser Trp Lys Val Ser Ser Gln Cys Ala Asp Thr Arg
            260                 265                 270

Lys Val Pro Leu Asp Ser Ser Pro Ala Thr Cys His Asn Asn Ile Met
        275                 280                 285
```

```
Lys Gln Thr Met Val Asp Ser Ser Cys Arg Ile Leu Thr Ser Asp Val
    290                 295                 300

Phe Gln Asp Cys Asn Lys Leu Val Asp Pro Glu Pro Tyr Leu Asp Val
305                 310                 315                 320

Cys Ile Tyr Asp Thr Cys Ser Cys Glu Ser Ile Gly Asp Cys Ala Cys
                325                 330                 335

Phe Cys Asp Thr Ile Ala Ala Tyr Ala His Val Cys Ala Gln His Gly
            340                 345                 350

Lys Val Val Thr Trp Arg Thr Ala Thr Leu Cys Pro Gln Ser Cys Glu
        355                 360                 365

Glu Arg Asn Leu Arg Glu Asn Gly Tyr Glu Cys Glu Trp Arg Tyr Asn
    370                 375                 380

Ser Cys Ala Pro Ala Cys Gln Val Thr Cys Gln His Pro Glu Pro Leu
385                 390                 395                 400

Ala Cys Pro Val Gln Cys Val Glu Gly Cys His Ala His Cys Pro Pro
                405                 410                 415

Gly Lys Ile Leu Asp Glu Leu Leu Gln Thr Cys Val Asp Pro Glu Asp
            420                 425                 430

Cys Pro Val Cys Glu Val Ala Gly Arg Arg Phe Ala Ser Gly Lys Lys
        435                 440                 445

Val Thr Leu Asn Pro Ser Asp Pro Glu His Cys Gln Ile Cys His Cys
    450                 455                 460

Asp Val Val Asn Leu Thr Cys Glu Ala Cys Gln Glu Pro Gly Gly Leu
465                 470                 475                 480

Val Val Pro Pro Thr Asp Ala Pro Val Ser Pro Thr Thr Leu Tyr Val
                485                 490                 495

Glu Asp Ile Ser Glu Pro Pro Leu His Asp Phe Tyr Cys Ser Arg Leu
            500                 505                 510

Leu Asp Leu Val Phe Leu Leu Asp Gly Ser Ser Arg Leu Ser Glu Ala
        515                 520                 525

Glu Phe Glu Val Leu Lys Ala Phe Val Val Asp Met Met Glu Arg Leu
    530                 535                 540

Arg Ile Ser Gln Lys Trp Val Arg Val Ala Val Val Glu Tyr His Asp
545                 550                 555                 560
```

Gly Ser His Ala Tyr Ile Gly Leu Lys Asp Arg Lys Arg Pro Ser Glu
565 570 575

Leu Arg Arg Ile Ala Ser Gln Val Lys Tyr Ala Gly Ser Gln Val Ala
580 585 590

Ser Thr Ser Glu Val Leu Lys Tyr Thr Leu Phe Gln Ile Phe Ser Lys
595 600 605

Ile Asp Arg Pro Glu Ala Ser Arg Ile Thr Leu Leu Leu Met Ala Ser
610 615 620

Gln Glu Pro Gln Arg Met Ser Arg Asn Phe Val Arg Tyr Val Gln Gly
625 630 635 640

Leu Lys Lys Lys Lys Val Ile Val Ile Pro Val Gly Ile Gly Pro His
645 650 655

Ala Asn Leu Lys Gln Ile Arg Leu Ile Glu Lys Gln Ala Pro Glu Asn
660 665 670

Lys Ala Phe Val Leu Ser Ser Val Asp Glu Leu Glu Gln Gln Arg Asp
675 680 685

Glu Ile Val Ser Tyr Leu Cys Asp Leu Ala Pro Glu Ala Pro Pro Pro
690 695 700

Thr Leu Pro Pro Asp Met Ala Gln Val Thr Val Gly Pro Gly Leu Leu
705 710 715 720

Gly Val Ser Thr Leu Gly Pro Lys Arg Asn Ser Met Val Leu Asp Val
725 730 735

Ala Phe Val Leu Glu Gly Ser Asp Lys Ile Gly Glu Ala Asp Phe Asn
740 745 750

Arg Ser Lys Glu Phe Met Glu Glu Val Ile Gln Arg Met Asp Val Gly
755 760 765

Gln Asp Ser Ile His Val Thr Val Leu Gln Tyr Ser Tyr Met Val Thr
770 775 780

Val Glu Tyr Pro Phe Ser Glu Ala Gln Ser Lys Gly Asp Ile Leu Gln
785 790 795 800

Arg Val Arg Glu Ile Arg Tyr Gln Gly Gly Asn Arg Thr Asn Thr Gly
805 810 815

Leu Ala Leu Arg Tyr Leu Ser Asp His Ser Phe Leu Val Ser Gln Gly
820 825 830

```
Asp Arg Glu Gln Ala Pro Asn Leu Val Tyr Met Val Thr Gly Asn Pro
        835                 840                 845

Ala Ser Asp Glu Ile Lys Arg Leu Pro Gly Asp Ile Gln Val Val Pro
    850                 855                 860

Ile Gly Val Gly Pro Asn Ala Asn Val Gln Glu Leu Glu Arg Ile Gly
865                 870                 875                 880

Trp Pro Asn Ala Pro Ile Leu Ile Gln Asp Phe Glu Thr Leu Pro Arg
                885                 890                 895

Glu Ala Pro Asp Leu Val Leu Gln Arg Cys Cys Ser Gly Glu Gly Leu
            900                 905                 910

Gln Ile Pro Thr Leu Ser Pro Ala Pro Asp Cys Ser Gln Pro Leu Asp
        915                 920                 925

Val Ile Leu Leu Leu Asp Gly Ser Ser Ser Phe Pro Ala Ser Tyr Phe
    930                 935                 940

Asp Glu Met Lys Ser Phe Ala Lys Ala Phe Ile Ser Lys Ala Asn Ile
945                 950                 955                 960

Gly Pro Arg Leu Thr Gln Val Ser Val Leu Gln Tyr Gly Ser Ile Thr
                965                 970                 975

Thr Ile Asp Val Pro Trp Asn Val Val Pro Glu Lys Ala His Leu Leu
            980                 985                 990

Ser Leu Val Asp Val Met Gln Arg  Glu Gly Gly Pro Ser  Gln Ile Gly
        995                 1000                1005

Asp Ala  Leu Gly Phe Ala Val  Arg Tyr Leu Thr Ser  Glu Met His
    1010                1015                1020

Gly Ala  Arg Pro Gly Ala Ser  Lys Ala Val Val Ile  Leu Val Thr
    1025                1030                1035

Asp Val  Ser Val Asp Ser Val  Asp Ala Ala Ala Asp  Ala Ala Arg
    1040                1045                1050

Ser Asn  Arg Val Thr Val Phe  Pro Ile Gly Ile Gly  Asp Arg Tyr
    1055                1060                1065

Asp Ala  Ala Gln Leu Arg Ile  Leu Ala Gly Pro Ala  Gly Asp Ser
    1070                1075                1080

Asn Val  Val Lys Leu Gln Arg  Ile Glu Asp Leu Pro  Thr Met Val
    1085                1090                1095
```

```
Thr Leu  Gly Asn Ser Phe Leu  His Lys Leu Cys Ser  Gly Phe Val
    1100                 1105                 1110

Arg Ile  Cys Met Asp Glu Asp  Gly Asn Glu Lys Arg  Pro Gly Asp
    1115                 1120                 1125

Val Trp  Thr Leu Pro Asp Gln  Cys His Thr Val Thr  Cys Gln Pro
    1130                 1135                 1140

Asp Gly  Gln Thr Leu Leu Lys  Ser His Arg Val Asn  Cys Asp Arg
    1145                 1150                 1155

Gly Leu  Arg Pro Ser Cys Pro  Asn Ser Gln Ser Pro  Val Lys Val
    1160                 1165                 1170

Glu Glu  Thr Cys Gly Cys Arg  Trp Thr Cys Pro Cys  Val Cys Thr
    1175                 1180                 1185

Gly Ser  Ser Thr Arg His Ile  Val Thr Phe Asp Gly  Gln Asn Phe
    1190                 1195                 1200

Lys Leu  Thr Gly Ser Cys Ser  Tyr Val Leu Phe Gln  Asn Lys Glu
    1205                 1210                 1215

Gln Asp  Leu Glu Val Ile Leu  His Asn Gly Ala Cys  Ser Pro Gly
    1220                 1225                 1230

Ala Arg  Gln Gly Cys Met Lys  Ser Ile Glu Val Lys  His Ser Ala
    1235                 1240                 1245

Leu Ser  Val Glu Leu His Ser  Asp Met Glu Val Thr  Val Asn Gly
    1250                 1255                 1260

Arg Leu  Val Ser Val Pro Tyr  Val Gly Gly Asn Met  Glu Val Asn
    1265                 1270                 1275

Val Tyr  Gly Ala Ile Met His  Glu Val Arg Phe Asn  His Leu Gly
    1280                 1285                 1290

His Ile  Phe Thr Phe Thr Pro  Gln Asn Asn Glu Phe  Gln Leu Gln
    1295                 1300                 1305

Leu Ser  Pro Lys Thr Phe Ala  Ser Lys Thr Tyr Gly  Leu Cys Gly
    1310                 1315                 1320

Ile Cys  Asp Glu Asn Gly Ala  Asn Asp Phe Met Leu  Arg Asp Gly
    1325                 1330                 1335

Thr Val  Thr Thr Asp Trp Lys  Thr Leu Val Gln Glu  Trp Thr Val
    1340                 1345                 1350
```

```
Gln Arg  Pro Gly Gln Thr Cys  Gln Pro Ile Leu Glu  Glu Gln Cys
    1355                 1360                 1365

Leu Val  Pro Asp Ser Ser His  Cys Gln Val Leu Leu  Leu Pro Leu
    1370                 1375                 1380

Phe Ala  Glu Cys His Lys Val  Leu Ala Pro Ala Thr  Phe Tyr Ala
    1385                 1390                 1395

Ile Cys  Gln Gln Asp Ser Cys  His Gln Glu Gln Val  Cys Glu Val
    1400                 1405                 1410

Ile Ala  Ser Tyr Ala His Leu  Cys Arg Thr Asn Gly  Val Cys Val
    1415                 1420                 1425

Asp Trp  Arg Thr Pro Asp Phe  Cys Ala Met Ser Cys  Pro Pro Ser
    1430                 1435                 1440

Leu Val  Tyr Asn His Cys Glu  His Gly Cys Pro Arg  His Cys Asp
    1445                 1450                 1455

Gly Asn  Val Ser Ser Cys Gly  Asp His Pro Ser Glu  Gly Cys Phe
    1460                 1465                 1470

Cys Pro  Pro Asp Lys Val Met  Leu Glu Gly Ser Cys  Val Pro Glu
    1475                 1480                 1485

Glu Ala  Cys Thr Gln Cys Ile  Gly Glu Asp Gly Val  Gln His Gln
    1490                 1495                 1500

Phe Leu  Glu Ala Trp Val Pro  Asp His Gln Pro Cys  Gln Ile Cys
    1505                 1510                 1515

Thr Cys  Leu Ser Gly Arg Lys  Val Asn Cys Thr Thr  Gln Pro Cys
    1520                 1525                 1530

Pro Thr  Ala Lys Ala Pro Thr  Cys Gly Leu Cys Glu  Val Ala Arg
    1535                 1540                 1545

Leu Arg  Gln Asn Ala Asp Gln  Cys Cys Pro Glu Tyr  Glu Cys Val
    1550                 1555                 1560

Cys Asp  Pro Val Ser Cys Asp  Leu Pro Pro Val Pro  His Cys Glu
    1565                 1570                 1575

Arg Gly  Leu Gln Pro Thr Leu  Thr Asn Pro Gly Glu  Cys Arg Pro
    1580                 1585                 1590

Asn Phe  Thr Cys Ala Cys Arg  Lys Glu Glu Cys Lys  Arg Val Ser
    1595                 1600                 1605
```

```
Pro Pro  Ser Cys Pro Pro His  Arg Leu Pro Thr Leu  Arg Lys Thr
    1610                 1615                 1620

Gln Cys  Cys Asp Glu Tyr Glu  Cys Ala Cys Asn Cys  Val Asn Ser
    1625                 1630                 1635

Thr Val  Ser Cys Pro Leu Gly  Tyr Leu Ala Ser Thr  Ala Thr Asn
    1640                 1645                 1650

Asp Cys  Gly Cys Thr Thr Thr  Thr Cys Leu Pro Asp  Lys Val Cys
    1655                 1660                 1665

Val His  Arg Ser Thr Ile Tyr  Pro Val Gly Gln Phe  Trp Glu Glu
    1670                 1675                 1680

Gly Cys  Asp Val Cys Thr Cys  Thr Asp Met Glu Asp  Ala Val Met
    1685                 1690                 1695

Gly Leu  Arg Val Ala Gln Cys  Ser Gln Lys Pro Cys  Glu Asp Ser
    1700                 1705                 1710

Cys Arg  Ser Gly Phe Thr Tyr  Val Leu His Glu Gly  Glu Cys Cys
    1715                 1720                 1725

Gly Arg  Cys Leu Pro Ser Ala  Cys Glu Val Val Thr  Gly Ser Pro
    1730                 1735                 1740

Arg Gly  Asp Ser Gln Ser Ser  Trp Lys Ser Val Gly  Ser Gln Trp
    1745                 1750                 1755

Ala Ser  Pro Glu Asn Pro Cys  Leu Ile Asn Glu Cys  Val Arg Val
    1760                 1765                 1770

Lys Glu  Glu Val Phe Ile Gln  Gln Arg Asn Val Ser  Cys Pro Gln
    1775                 1780                 1785

Leu Glu  Val Pro Val Cys Pro  Ser Gly Phe Gln Leu  Ser Cys Lys
    1790                 1795                 1800

Thr Ser  Ala Cys Cys Pro Ser  Cys Arg Cys Glu Arg  Met Glu Ala
    1805                 1810                 1815

Cys Met  Leu Asn Gly Thr Val  Ile Gly Pro Gly Lys  Thr Val Met
    1820                 1825                 1830

Ile Asp  Val Cys Thr Thr Cys  Arg Cys Met Val Gln  Val Gly Val
    1835                 1840                 1845

Ile Ser  Gly Phe Lys Leu Glu  Cys Arg Lys Thr Thr  Cys Asn Pro
    1850                 1855                 1860
```

```
Cys Pro  Leu Gly Tyr Lys Glu  Glu Asn Asn Thr Gly  Glu Cys Cys
    1865                1870                1875

Gly Arg  Cys Leu Pro Thr Ala  Cys Thr Ile Gln Leu  Arg Gly Gly
    1880                1885                1890

Gln Ile  Met Thr Leu Lys Arg  Asp Glu Thr Leu Gln  Asp Gly Cys
    1895                1900                1905

Asp Thr  His Phe Cys Lys Val  Asn Glu Arg Gly Glu  Tyr Phe Trp
    1910                1915                1920

Glu Lys  Arg Val Thr Gly Cys  Pro Pro Phe Asp Glu  His Lys Cys
    1925                1930                1935

Leu Ala  Glu Gly Gly Lys Ile  Met Lys Ile Pro Gly  Thr Cys Cys
    1940                1945                1950

Asp Thr  Cys Glu Glu Pro Glu  Cys Asn Asp Ile Thr  Ala Arg Leu
    1955                1960                1965

Gln Tyr  Val Lys Val Gly Ser  Cys Lys Ser Glu Val  Glu Val Asp
    1970                1975                1980

Ile His  Tyr Cys Gln Gly Lys  Cys Ala Ser Lys Ala  Met Tyr Ser
    1985                1990                1995

Ile Asp  Ile Asn Asp Val Gln  Asp Gln Cys Ser Cys  Cys Ser Pro
    2000                2005                2010

Thr Arg  Thr Glu Pro Met Gln  Val Ala Leu His Cys  Thr Asn Gly
    2015                2020                2025

Ser Val  Val Tyr His Glu Val  Leu Asn Ala Met Glu  Cys Lys Cys
    2030                2035                2040

Ser Pro  Arg Lys Cys Ser Lys
    2045                2050
```

FIG. 1


rVWF 164A 4°C (132U)
rVWF164A 4°C (30U)
rVWF 164A 40°C (132U)
rVWF164A 40°C (30U)
rVWF 164A 29°C (132U)
rVWF 164A 29°C (30U)
VWF:RCo [%]
storage time [weeks]
100
90
80
70
60
50
40
30
20
10
0
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

FIG. 2


VWF:RCo [%]
100
90
80
70
60
50
40
30
20
10
0
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26
storage time [weeks]
rVWF 164B 4°C (132U)
rVWF164B 4°C (30U)
rVWF 164B 40°C (132U)
rVWF164B 40°C (30U)
rVWF 164B 29°C (132U)
rVWF 164B 29°C (30U)

FIG. 3


rVWF209 4°C
rVWF209 30°C
rVWF209 40°C
rVWF209 -20°C
rvWF164B 4°C
rvWF164B 29°C
rvWF164B 40°C
VWF:RCo [U/ml]
0
20
40
60
80
100
120
140
160
storage time [week]
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

FIG. 4


VWF:Ag [%]
storage time [weeks]
rVWF 164A 4°C (132U)
rVWF164A 4°C (30U)
rVWF 164A 40°C (132U)
rVWF164A 40°C (30U)
rVWF 164A 29°C (132U)
rVWF 164A 29°C (30U)

FIG. 5


VWF:Ag [%]
160
140
120
100
80
60
40
20
0
storage time [weeks]
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26
rVWF 164B 4°C (132U)
rVWF164B 4°C (30U)
rVWF 164B 40°C (132U)
rVWF164B 40°C (30U)
rVWF 164B 29°C (132U)
rVWF 164B 29°C (30U)

FIG. 6


VWF:Ag [U/ml]
450
400
350
300
250
200
150
100
50
0
storage time [week]
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26
rVWF209 4°C
rVWF209 30°C
rVWF209 40°C
rVWF209 -20°C

FIG. 7


DSC103 rvWF183-1 (Citrat pH 6.5)
DSC105 rvWF183-3 (Citrat/Glycin pH 6.5)
DSC107 rvWF183-5 (Citrat/Histidin pH 6.5)
DSC112 rvWF183-10 (Citrat/32g/lSaccharose pH 6.5)
DSC114 rvWF183-12 (Citrat/Tween80 pH 6.5)
DSC115 rvWF183-13 (Citrat/Raffinose pH 6.5)
DSC117 rvWF183-15 (Citrat/7.8mMTrehalose pH 6.5)
Molar Heat Capacity [kJ/(K mol)]
Temperature [°C]
9700
8700
7700
6700
5700
4700
3700
2700
1700
700
-300
50
55
60
65
70
75
80
85
90
58.6°C
60.1°C
58.4°C
59.2°C
65.8°C
65.2°C
59.2°C
65.4°C
66.0°C
59.1°C
65.2°C
58.7°C
65.5°C
64.4°C
79.3°C
80.4°C
80.1°C
81.1°C
80.4°C
80.3°C
88.5°C
81.6°C

FIG. 8


DSC106 rvWF183-4 (Citrat/NaCl pH 6.5)
DSC110 rvWF183-8 (Citrat/1mMCaCl2 pH 6.5)
DSC111 rvWF183-9 (Citrat/10mMCaCl2 pH 6.5)
DSC113 rvWF183-11 (Citrat/0.25MSaccharose pH 6.5)
DSC116 rvWF183-14 (Citrat/Na2HPO4/NaH2PO4 pH 6.5)
Molar Heat Capacity [kJ/(K mol)]
Temperature [°C]
19700
17700
15700
13700
11700
9700
7700
5700
3700
1700
-300
40
50
60
70
80
90
93.0°C
81.7°C
83.1°C
81.0°C
80.4°C
91.8°C
81.0°C
89.2°C
67.2°C
64.5°C
67.0°C
61.4°C
61.0°C
60.0°C


Molar Heat Capacity [kJ/(K mol)]
17900
15900
13900
11900
9900
7900
5900
3900
1900
-100
50
55
60
65
70
75
80
85
90
95
100
Temperature [°C]
DCS124 rWF186-1 (Citral/1CaCl2 pH 7.3)
DCS118 rWF184-1 (Citral/1CaCl2 pH 6.5)
63.8°C
64.7°C
81.7°C
80.1°C
89.8°C
96.0°C

FIG. 9