



- (51) International Patent Classification:  
*A61K 38/22* (2006.01)
- (21) International Application Number:  
PCT/US2016/018602
- (22) International Filing Date:  
19 February 2016 (19.02.2016)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
62/118,431 19 February 2015 (19.02.2015) US
- (71) Applicant: THE REGENTS OF THE UNIVERSITY  
OF CALIFORNIA [US/US]; 1111 Franklin Street, 5th  
Floor, Oakland, CA 94607-5200 (US).
- (72) Inventor: BANSAL, Vishal; 200 West Arbor, San Diego,  
CA 92103 (US).
- (74) Agent: BAKER, Joseph, R., Jr.; Gavrilovich, Dodd &  
Lindsey LLP, 4660 La Jolla Village Drive, Suite 750, San  
Diego, CA 92122 (US).
- (81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,  
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,  
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,  
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,  
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,  
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,  
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,  
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: METHODS OF TREATING MILD BRAIN INJURY AND POST-TRAUMATIC STRESS DISORDER




FIG. 1

(57) Abstract: The present disclosure provides methods for treating Post-Traumatic Stress Disorder (PTSD) and other neurological disorders in a subject, comprising administering to the subject an effective amount of a composition comprising ghrelin. This disclosure provides for methods of treating mild brain injury (mBI) and PTSD in a subject having such an injury or disorder, comprising administering to the subject an effective amount of ghrelin or ghrelin agonist, thereby treating the mBI or disorder.

## METHODS OF TREATING MILD BRAIN INJURY AND POST-TRAUMATIC STRESS DISORDER

### CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. §119 from Provisional Application Serial No. 62/118,431, filed February 19, 2015, the disclosure of which is incorporated herein by reference in its entirety.

### FIELD OF THE INVENTION

[0002] This invention provides for methods for treating Post-Traumatic Stress Disorder (PTSD), mild brain injuries and other neurological disorders arising from such an injury in a subject by administering to the subject an effective amount of a composition comprising ghrelin.

### BACKGROUND

[0003] Ghrelin, a 28-amino acid peptide predominantly secreted by gastric mucosa, is a neuroendocrine hormone that acts as an endogenous ligand for growth hormone secretagogue receptor. Beyond the known effects on hunger regulation, ghrelin is known to have potent anti-inflammatory properties and has been shown to be protective in several models of severe neuronal injury.

[0004] Post traumatic brain injury (*i.e.*, TBI) edema and the complications associated with increased intracranial pressure (ICP) account for approximately 50% of death in hospitalized patients. Medically managing intracranial hypertension utilizes a strategy combining sedation and osmotic agents such as mannitol and hypertonic saline.

[0005] Post-Traumatic Stress Disorder (PTSD) is a disorder that can develop after exposure to one or more traumatic event or ordeal, such as one in which grave physical harm occurred or was threatened to oneself or others, sexual assault, warfare, serious injury, or threats of imminent death, that result in feelings of intense fear, horror, and/or powerlessness.

[0006] The cost of treating post-traumatic stress disorder (PTSD) for soldiers participating in Iraq and Afghanistan from 2003-2010 has been approximately \$1.4 billion. Approximately 21% of soldiers have been observed to develop PTSD after deployment to Iraq or Afghanistan.

**SUMMARY**

**[0007]** This disclosure provides for methods of treating mild brain injury (mBI) and PTSD in a subject having such an injury or disorder, comprising administering to the subject an effective amount of ghrelin or ghrelin agonist, thereby treating the mBI or disorder. In one embodiment, ghrelin comprises a polypeptide having at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO:1). In another embodiment, the serine at amino acid position 2 and/or serine at amino acid position 3 are/is acylated with an octanoyl group ( $-C(O)(CH_2)_6CH_3$ ). In yet further or alternative embodiments, amino acid(s) at position 1 to 5 of SEQ ID NO:1 of the ghrelin or ghrelin agonist can each independently be modified. In yet other embodiments, ghrelin increases uncoupling protein-2 (UCP-2) expression. In yet another embodiment, ghrelin increases UCP-2 expression in mitochondria. In yet another embodiment of any of the foregoing, the amount ghrelin (or modified for thereof) delivered according to the methods herein, can be for example, a dosage of 2  $\mu\text{g/kg}$  per dose and/or per day. In another embodiment, ghrelin is des-acyl-ghrelin or ghrelin with no modification of the primary amino acid sequence as provided in SEQ ID NO:1. In another embodiment, ghrelin binds to a receptor other than GHSR-1a or ghrelin receptor, and wherein binding to a receptor other than GHSR-1a or ghrelin receptor provides a therapeutic benefit following mBI, for example, neuroprotection following mBI, repeated mBI or CTE. The therapeutic benefit such as neuroprotection following mBI or repeated mBI may include reduced oxidative stress or reduced apoptosis.

**[0008]** The disclosure provides methods of treating Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0009]** The disclosure provides methods of preventing the onset of Post-Traumatic Stress Disorder (PTSD) or delaying PTSD resulting from one or more of mild brain injuries in a subject in need

thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby preventing or delaying the onset of the PTSD, or treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0010]** In some embodiments, the PTSD is associated with a single mild brain injury or concussion. In some embodiments, the PTSD is associated with repetitive or multiple mild brain injuries. In some embodiments, ghrelin is used as a sole active agent for treating the PTSD. In some embodiments, ghrelin is used in a combination with one or more therapeutic agents.

**[0011]** The disclosure provides compositions comprising a therapeutically effective amount of ghrelin for use in the treatment of Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject the composition, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0012]** The disclosure provides compositions comprising a therapeutically effective amount of ghrelin for use in the treatment of preventing or delaying the onset of Post-Traumatic Stress Disorder (PTSD) or delaying the PTSD resulted from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject composition, thereby preventing the onset of the PTSD or treating the PTSD or reduce the severity of the PTSD and/or the related symptoms.

**[0013]** In some embodiments, the PTSD is associated with a single mild brain injury or concussion. In some embodiments, the PTSD is associated with repetitive or multiple mild brain injuries. In some embodiments, ghrelin is used as a sole active agent for treating the PTSD. In some embodiments, ghrelin is used in a combination with one or more therapeutic agents. In some embodiments, when administered the ghrelin is the sole active agent being administered to the subject.

**[0014]** In some embodiments, the ghrelin has the sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO: 1). In some embodiments, the ghrelin is modified with one or more fatty acids. In some embodiments, the fatty acid is an octanoic acid. In some

embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine at amino acid position 3 of SEQ ID NO: 1. In some embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine amino acid position 3 with an octanoic acid. In some embodiments, modification of ghrelin is acylation of ghrelin at serine at amino acid position 2 and/or serine at amino acid position 3 with an octanoic acid. In some embodiments, the ghrelin so modified is an octanoyl-ghrelin with an acylation at serine at amino acid position 3 by the octanoic acid.

**[0015]** In some embodiments, the subject is a mammal. In some embodiments, the ghrelin is administered in an amount that results in ghrelin blood levels being at least two times greater than the endogenous ghrelin blood levels of the subject. In some embodiments, ghrelin is administered in a single dose. In some embodiments, ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per day. In some embodiments, ghrelin is administered at a dosage of 2 µg/kg per day.

**[0016]** This disclosure also provides methods of treating Post-Traumatic Stress Disorder (PTSD) in a subject, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition consisting essentially of isolated ghrelin.

**[0017]** In some embodiments, the mild brain injury comprises a concussion. In some embodiments, the subject is a mammal. In some embodiments, the subject is a human. In some embodiments, ghrelin is administered within not more than about 8, 24 or 72 hours after the mBI. In some embodiments, ghrelin is administered within not more than about 24 hours after the mBI. In some embodiments, ghrelin is administered within or at about 0.1, 0.3, 0.5, 0.7, 1, 2, 3, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 36, 48, or 72 (or any integer there between) hours after the mBI.

**[0018]** This disclosure provides methods of reducing the incidence or severity of mBI in a subject, comprising administering to the subject an effective amount of ghrelin, thereby reducing the incidence or severity of the mBI. In some embodiments, ghrelin is administered prior to an event or activity with a potential for occurrence of mBI.

**[0019]** This disclosure provides for methods of reducing the amount of time required for recovery from a mild brain injury, comprising administering to a patient suffering from a mild brain injury a therapeutically effective amount of ghrelin within 72 hours of the mild brain injury. In some embodiments, ghrelin is administered in a single dose. In some embodiments, ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per day.

**[0020]** Still further, ghrelin can be used in an assay to assess the ability of candidate compounds to effect increased uncoupling protein-2 (UCP-2) expression including increased UCP-2 expression in mitochondria. In such assays, ghrelin is used as a control to determine the relative efficacy of the candidate compound or compounds. Suitable assays include, by way of example only, competitive assays for binding of a candidate compound or compounds to growth hormone secretagogue receptor 1a (*i.e.*, GHSR) in the presence of ghrelin as well as frontal affinity chromatography.

**[0021]** In yet another embodiment, a patient suffering loss of cognitive or motor skills due to mBI and, in particular, repetitive mBI, can be monitored for therapy or progression of such skills by correlating the ghrelin level in the patient's brain over time. As the ghrelin levels decrease, there will be an increased need for intervention. This invention also provides for methods of measuring ghrelin levels before an activity, for example before the start of football, soccer, rugby or any other sport or activity season, and monitoring during the season to ascertain if the player is at a level not qualified to play or participate by utilizing a test or an assay for measuring ghrelin levels, such as a test or assay for determining levels in the blood.

**[0022]** In some embodiments, the disclosure provides for a method for treating a subject suffering from metabolic derangements associated with mBI or concussion, wherein such method comprises administration of an effective amount of ghrelin to the subject, thereby treating the subject suffering from metabolic derangements associated with mBI or concussion.

**[0023]** In some embodiments, the disclosure provides for a method for treating a subject suffering from increased levels of reactive oxidative species (ROS) in neurons associated with mBI or

concussion, wherein such method comprises administration of an effective amount of ghrelin to the subject so as to decrease levels of ROS in neurons associated with mBI or concussion, thereby treating the subject suffering from metabolic derangements associated with mBI or concussion.

**[0024]** In some embodiments, the disclosure provides for a method for preventing chronic traumatic encephalopathy (CTE) associated with repeated mBI or concussions in a subject, wherein such method comprises administration of one or more doses of ghrelin to the subject, thereby preventing chronic traumatic encephalopathy (CTE) associated with repeated mBI or concussions in a subject.

**[0025]** In some embodiments, the disclosure provides for a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) in a subject with one or more incidence of mBI or concussion.

**[0026]** In some embodiments, the disclosure provides for a method for treating injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI, including concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject, thereby treating injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI, including concussion. In one embodiment, the injured or disrupted neuronal or axonal connection is mechanical disruption of neuronal or axonal cytoskeleton. In another embodiment, the injured or disrupted neuronal or axonal connection is disruption or altered neuronal or axonal transport. In another embodiment, the injured or disrupted neuronal or axonal connection is proteolysis, die-back disconnection and reorganization. In one embodiment, treating injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI, including concussion,

restores neuronal or axonal connection in the brain of a subject following one or more incidence of mBI, including concussion. In another embodiment, treating injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI or concussion hastens restoration of neuronal or axonal connection in the brain of a subject following one or more incidence of mBI, including concussion.

**[0027]** In some embodiments, the disclosure provides for a method for preventing or diminishing occurrence of an injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI, including concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject, thereby reducing or preventing or diminishing occurrence of an injured or disrupted neuronal or axonal connection in the brain of a subject with one or more incidence of mBI, including concussion. In one embodiment, the injured or disrupted neuronal or axonal connection is mechanical disruption of neuronal or axonal cytoskeleton. In another embodiment, the injured or disrupted neuronal or axonal connection is disruption or altered neuronal or axonal transport. In another embodiment, the injured or disrupted neuronal or axonal connection is proteolysis, die-back disconnection and reorganization. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI, including concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI, including concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI, including concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0028]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI, including concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore normal neuronal depolarization, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI, including concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal

brain physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0029]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI, including concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain normal neuronal depolarization, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI, including concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI, including concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI, including concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI, including concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0030]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore normal release of excitatory neurotransmitters, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal brain physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0031]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain normal release of excitatory neurotransmitters, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI or concussion. In

another embodiment, ghrelin is administered immediately after the occurrence of an mBI or concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI or concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0032]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore normal cerebral blood flow, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal brain physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0033]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain normal cerebral blood flow, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI or concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI or concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI or concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0034]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore proper or normal axonal function, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal brain

physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0035]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain proper or normal axonal function, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI or concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI or concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI or concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0036]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore brain metabolic function, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal brain physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0037]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain brain metabolic function or balance, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI or concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI or concussion. In another embodiment, ghrelin is administered prior

to and following occurrence of one or more mBI or concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0038]** In some embodiments, the disclosure provides for a method for treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to restore glucose metabolism, thereby treating perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, treating perturbed brain physiology with ghrelin restores normal brain physiology. In another embodiment, treating perturbed brain physiology with ghrelin hastens restoration of normal brain physiology.

**[0039]** In some embodiments, the disclosure provides for a method for preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion, wherein such method comprises administration of one or more doses of ghrelin to the subject so as to maintain glucose metabolism, thereby reducing or preventing perturbed brain physiology in a subject with one or more incidence of mBI or concussion. In one embodiment, ghrelin is administered prior to the occurrence of one or more incidence of mBI or concussion. In another embodiment, ghrelin is administered immediately after the occurrence of an mBI or concussion. In another embodiment, ghrelin is administered prior to and following occurrence of one or more mBI or concussion. In another embodiment, multiple doses of ghrelin are administered.

**[0040]** In some embodiments, the disclosure provides for a method for preventing memory loss or headaches in a subject with mBI or concussion, wherein such method comprises administration of an effective amount of ghrelin to the subject in one or more doses, thereby preventing memory loss and/or headaches in a subject with mBI or concussion.

**[0041]** Some embodiments relate to formulations for administration to a subject, which formulations can include a pharmaceutically acceptable carrier and ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein said formulation is suitable for delivery of an effective amount of ghrelin to the brain of said patient so as to treat mild brain

injuries. For example, any of the methods described above and elsewhere herein can utilize a ghrelin molecule have a C14 content of less than 1 ppt.

**[0042]** Some embodiments relate to methods of monitoring a mild brain injury, the severity of an injury and/or the recovery from such an injury. The methods can include, for example, administering a purified ghrelin compound, including for example, ghrelin with a C14 content of less than 1 ppt, in a pharmaceutically acceptable composition to a subject that has suffered a mild brain injury.

**[0043]** Some embodiments relate to methods of treating mild brain injury, reducing the incidence or severity of mBI in a subject, and/or reducing the amount of time needed to recover from a mild brain injury. The methods include providing or administering to a subject in need (e.g., a subject that has suffered, is at risk of suffering, is prone to suffer, and/or is about to participate in an activity with a high risk for suffering, a mBI) an amount of ghrelin (including ghrelin with a C14 content of less than 1 ppt) sufficient to provide a therapeutically effective *in vivo* level of ghrelin to treat or reduce according to the method, wherein the level is greater than the endogenous level of ghrelin in the subject. For example, the amount of administered ghrelin can be an amount sufficient to provide a blood level of ghrelin that is greater than the usual or average endogenous blood level of ghrelin, such as 1.5, 2, 3, 5, 10, 20, 50, 100, 1,000 or up to 2,000 times the normal endogenous blood level (or any sub value or sub range there between). In some case, the amount administered can result in a blood or plasma concentration of at least 55 picograms per milliliter. In some embodiments the greater ghrelin levels can be achieved within hours of the injury (e.g., less than 8 hours after the injury). They also can be maintained above endogenous levels for some period of time that is sufficient to provide the desired therapeutic benefit, for example, for at least 30 minutes to 24 hours (or any sub value or sub range there between). Endogenous ghrelin levels are not sufficient for treating mBI or reducing the incidence, severity or the time needed to recover as readily evidenced by the long term damage done to the

brain by repetitive concussive injuries (mBI). The instant embodiments provide a benefit and result that do not occur naturally in the body with endogenous levels. Such a benefit was unknown prior to the instant embodiments associated with the invention. In some embodiments, the methods can further include selecting or identifying a subject that has suffered, is at risk of suffering, is prone to suffer, and/or is about to participate in an activity with a high risk for suffering, a mBI, prior to administration of the ghrelin.

**[0044]** Some embodiments relate to methods for determining the efficacy of a compound treating a patient suffering mild brain injury (mBI). The methods can include, for example, i) determining the expression level of uncoupling protein-2 (UCP-2) in a biological sample obtained from the patient treated with the compound; ii) comparing the expression level of UCP-2 to a biological sample obtained from a subject treated with ghrelin (as described herein including with the modifications described herein); and iii) determining the efficacy of the compound, wherein the compound is efficacious when the expression level of UCP-2 induced by the compound is equal to (within at least 10%) or greater than the expression level of UCP-2 induced by ghrelin.

**[0045]** Some embodiments relate to methods for treating a patient suffering loss of cognitive or motor skills due to mild brain injury (mBI). The methods can include, for example, i) determining the level of ghrelin in the patient's brain over a period of time; ii) administering ghrelin (including the modified and C14 versions described herein) to the patient; and iii) periodically repeating step ii) during treatment when the level of ghrelin falls below a normal range as a basis to determine the efficacy of the treatment, wherein an increase in the level of ghrelin in the brain demonstrates an improvement in the patient's cognitive or motor skill condition.

**[0046]** The details of one or more embodiments of the disclosure are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the disclosure will be apparent from the description and drawings, and from the claims.

**BRIEF DESCRIPTION OF THE DRAWINGS**

[0047] **FIG. 1** depicts principle of assaying oxidative burst within inflammatory cells. Dihydrorhodamine 123 (DHR 123) diffuses across the cell membrane. When it encounters reactive oxygen species (ROS), DHR 123 is oxidized and fluoresces green. The fluorescence is then measured and presented as arbitrary fluorescence unit (AFU), wherein higher intensity means greater oxidative burst and, therefore greater concentration of amount of ROS (See Chen, et al., Measurement of oxidative burst in neutrophils," Methods Mol Biol., 844: 115-124 (2012); which is incorporated herein by reference in its entirety).

[0048] **FIG. 2** depicts subcutaneous administration of ghrelin resulting in reduced oxidative burst in inflammatory cells following mBI. Rightward shift in peak reflects higher oxidative burst. Higher oxidative burst is observed in mBI alone, and lower oxidative burst is observed in ghrelin treated mBI. For the ghrelin treatment, the peak is shifted left in line with control meaning lower oxidative burst.

[0049] **FIG. 3** depicts oxidative burst of ipsilateral whole brain following mild BI. Mild BI plus ghrelin reduces oxidative burst as quantified by arbitrary fluorescence unit (AFU). Oxidative burst is increased in animals following mBI compared to sham control (7963 AFU  $\pm$  2900 (mild BI) vs. 4624 AFU  $\pm$  1858 (sham); n=5 animals in each group). Ghrelin treatment reduces the oxidative burst compared to mBI (7963 AFU  $\pm$  2900 (mild BI) vs. 3257 AFU  $\pm$  1031 (mild BI + ghrelin (2 doses)); p-value of 0.048 using Student's t-test). AFU equals mean fluorescence intensity.

[0050] The figures are exemplification; hence, the figures are presented for the purpose of illustrating one or more embodiments with the explicit understanding that they will not be used to limit the scope or the meaning of the claims that follow below.

**DETAILED DESCRIPTION**

[0051] Unless specifically noted otherwise herein, the definitions of the terms used are standard definitions used in the art of pharmaceutical sciences. As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise.

Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

**[0052]** Also, the use of "or" means "and/or" unless stated otherwise. Similarly, "comprise," "comprises," "comprising," "include," "includes," and "including" are interchangeable and not intended to be limiting.

**[0053]** It is to be further understood that where descriptions of various embodiments use the term "comprising," those skilled in the art would understand that in some specific instances, an embodiment can be alternatively described using language "consisting essentially of" or "consisting of."

**[0054]** Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this disclosure belongs. Although any methods and reagents similar or equivalent to those described herein can be used in the practice of the disclosed methods and compositions, the exemplary methods and materials are now described.

**[0055]** All publications mentioned herein are incorporated herein by reference in full for the purpose of describing and disclosing the methodologies presented therein.

**[0056]** It is to be understood that the present disclosure is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

**[0057]** As used herein, the term "about" when used before a numerical designation, e.g., temperature, time, amount, concentration, and such other, including a range, indicates approximations which may vary by (+) or (-) 10%, 5% or 1%.

**[0058]** As used herein, the term "administration" can be effected in one dose, continuously or intermittently or by several subdoses which in the aggregate provide for a single dose. Dosing can be conducted throughout the course of treatment. Methods of determining the most effective means and dosage of administration are known to those of skill in the art and will vary with the

composition used for therapy, the purpose of the therapy, the target cell or tissue being treated and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician. Suitable dosage formulations and methods of administering the agents are known in the art. Route of administration can also be determined and method of determining the most effective route of administration are known to those of skill in the art and will vary with the composition used for treatment, the purpose of the treatment, the health condition or disease stage of the subject being treated and target cell or tissue. Non-limiting examples of route of administration include oral administration, vaginal, nasal administration, injection, topical application, sublingual, pulmonary, and by suppository.

**[0059]** As used herein, the term "affinity" refers to the strength of binding between receptors and their ligands, for example, between an antibody and its antigen.

**[0060]** As used herein, the term "amino acid residue" refers to an amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. Unless otherwise specified, the amino acid encompasses L-amino acid, including both natural amino acid and synthetic amino acid or the like as long as the desired functional property is retained by the polypeptide.  $\text{NH}_2$  refers to the free amino group present at the amino terminus of a polypeptide.  $\text{COOH}$  refers to the free carboxy group present at the carboxy terminus of a polypeptide. Standard polypeptide abbreviations for amino acid residues are as follows: A (Ala or Alanine); C (Cys or Cysteine); D (Asp or Aspartic Acid); E (Glu or Glutamic Acid); F (Phe or Phenylalanine); G (Gly or Glycine); H (His or Histidine); I (Ile or Isoleucine); K (Lys or Lysine); L (Leu or Leucine); M (Met or Methionine); N (Asn or Asparagine); P (Pro or Proline); Q (Gln or Glutamine); R (Arg or Arginine); S (Ser or Serine); T (Thr or Threonine); V (Val or Valine); W (Trp or Tryptophan); X (Xaa or Unknown or Other); Y (Tyr or Tyrosine); and Z (Glx/Gln/Glu or Glutamic Acid/Glutamine). All amino acid residue sequences represented herein by formula have a left-to-right orientation in the conventional direction of amino terminus to

carboxy terminus. The phrase "amino acid residue" is broadly defined to include the naturally occurring and modified and non-naturally occurring amino acids. A dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues or a covalent bond to an amino-terminal group such as NH<sub>2</sub> or acetyl or to a carboxy-terminal group such as COOH.

**[0061]** As used herein, the term "ghrelin receptor" refers to any naturally occurring molecule to which ghrelin binds and induces a biological activity. Ghrelin is known to bind to growth hormone secretagogue receptor 1a (*i.e.*, GHSR), however, the present disclosure is not limited to a specific type of receptor.

**[0062]** As used herein, the term "individual" or "subject" is an animal or human susceptible to a condition, in particular mBI or concussion. In some embodiments, the individual is a mammal, including human and non-human mammals such as dogs, cats, pigs, cows, sheep, goats, horses, rats, and mice.

**[0063]** As used herein, the term "mild brain injury" (mBI) sometimes referred as a "mild traumatic brain injury" (mTBI) refers to a non-disease event commonly caused by an injury resulting in an insult to the brain. mBI may be caused, for example, by impact forces, in which the head strikes or is struck by something, or impulsive forces, in which the head moves without itself being subject to blunt trauma (for example, when the chest hits something and the head snaps forward; as a result of rapid acceleration or deceleration of the head; acceleration or deceleration of the brain of a subject producing a change in a qualitative or quantifiable clinical parameter related to functional well-being of the subject; or differential acceleration or deceleration of the brain of the subject relative to the skull of the subject wherein differential acceleration or deceleration produces a change in a qualitative or quantifiable clinical parameter related to functional well-being of the subject). mBI commonly results, for example, from a sports-related injury, a motor vehicle accident, an accidental fall, or an assault. Although the vast majority of such injuries improve through natural recovery the damage caused by such an injury or

repetitive injuries can cause long term deficits in cognitive and/or motor skill functions.

**[0064]** Mild BI is different from and has a distinct pathology as compared to diseases such as acute traumatic brain insults such as strokes (ischemic or hemorrhagic), AVM's, brain tumors, and the like.

**[0065]** Moreover, mBI is distinct from severe TBI. Mild brain injury and severe traumatic brain injury are dramatically different injuries. Mild BI and severe TBI have different characteristic and symptoms, different mortality rates, different treatment protocols, etc. Severe TBI is an injury state where there is an unacceptable level of mortality and permanent disability in humans within hours or days of the injury. According to the Centers for Disease Control, such injuries are a contributing factor to a third (30%) of all injury-related deaths in the United States (See, National Vital Statistics System (NVSS), 2006-2010). Severe TBIs often are caused, for example, by automobile accidents and combat gunshot and bomb wounds to the head. Severe TBI typically has direct and overt brain hemorrhage, brain cell destruction, severe brain edema and severe cellular apoptosis, all characteristic of anatomical changes or anatomical injury to the brain of the injured patient—all of which can be detected by non-invasive anatomical imaging methods. Due to the severity of such symptoms and risk of imminent death and disability, operative intervention and critical care are paramount. Severe TBI results in adverse anatomical changes in the patient's brain (observable at a gross anatomy level) resulting in immediate and permanent loss of brain function, if not death.

**[0066]** On the other hand, mild brain injury (mBI) causes changes in the brain function at a metabolic / molecular level in the affected neurons. As such, it does not typically cause mortality or permanent disability in the time frame provided above and patients often resume normal activities of daily living and functionality within hours or days of the mBI.

**[0067]** The body's response to severe TBI consists of a large number of differing responses including those initiated by hypoxia, blood loss and significant neuron death, to name just a few. The body's response to mBI is significantly different as hemorrhaging

and blood loss, tissue loss, hypoxia, significant neuron death etc. are not part of the manifestations of this condition.

[0068] Consequently, therapeutic treatments for each injury are significantly different. The same therapeutic methodology used for severe TBI, for example, surgery, deep sedation, mannitol therapy and operative decompression, would be contra-indicated for treatment of mBI, for example, concussion, which is a type of mild brain injury. The inverse, such as rest and mental relaxation with little intervention for recovering from mBI, is likewise true.

[0069] As used herein, the term "polypeptide" or "peptide" is intended to encompass a singular "polypeptide" as well as plural "polypeptides," and refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term "polypeptide" refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, "protein," "amino acid chain," or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of "polypeptide," and the term "polypeptide" may be used instead of, or interchangeably with any of these terms. The term "polypeptide" is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, acylation, acylation by fatty acid, fatty acid-modification, and derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non-naturally occurring amino acids. Modification may be fatty acid modification or triglyceride modification. Fatty acid modification may be a short to medium-chain fatty acid. The short fatty acid may be a two-carbon fatty acid or acetic acid. Medium chain fatty acid may be 14-carbon fatty acid or tetradecanoic acid. Modification with a fatty acid may be acylation of SEQ ID NO: 1 at serine amino acid position 2 and/or serine amino acid position 3. Modification may be catalyzed by ghrelin O-acyl transferase (GOAT) of fatty acid thioester and ghrelin as substrates. In one embodiment, post-translationally modified ghrelin may be bound and/or recognized by growth hormone secretagogue receptor type 1a

(GHSR-1a) or ghrelin receptor. In one embodiment, post-translationally modified ghrelin may be fatty acid-acylated ghrelin at serine amino acid position 2 and/or serine amino acid position 3 bound and/or recognized by GHSR-1a or ghrelin receptor. A polypeptide may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis. The term "Polypeptide" or "Peptide" also refers to a compound comprising a plurality of amino acids linked therein via peptide linkages. Here, the amino acid (also called an amino acid residue) includes naturally occurring amino acids represented by formula:  $\text{NH}_2\text{-CH(R')-COOH}$ , wherein R' is a naturally occurring substituent group, as well as its D, L-optical isomers etc. There is also a peptide, wherein a certain naturally occurring amino acid is replaced by a modified amino acid. The modified amino acid includes the amino acids of the above formula wherein the substituent group R' is further modified, its D, L-optical isomers thereof, and non-natural amino acids wherein e.g. various substituent groups are bound to the substituent group R' of the above formula via or not via an ester, ether, thioester, thioether, amide, carbamide or thiocarbamide linkage. The modified amino acid also includes non-natural amino acids whose amino groups are replaced by lower alkyl groups.

**[0070]** As used herein, the term "secretagogue" is a substance stimulating growth hormone release, such as ghrelin. In some embodiments, the ghrelin binds to the growth hormone secretagogue receptor GHS-R 1a (GHSR). The ghrelin compounds described herein are active at the receptor for growth hormone secretagogue (GHS), e.g., the receptor GHS-R 1a. The compounds can bind to GHS-R 1a, and stimulate receptor activity. In some embodiments, the compounds can bind other receptors and, optionally, stimulate their activity. In some embodiments, ghrelin increases uncoupling protein-2 (UCP-2) expression. In some embodiments, ghrelin increases UCP-2 expression in mitochondria. In some embodiments, ghrelin prevents the metabolic consequence of mBI and any associated chronic conditions.

[0071] As used herein, the term "ghrelin" refers to a polypeptide that may have a 28 amino acid sequence as set forth in SEQ ID NO:1, and can include the octanoyl acylation as described herein. Human ghrelin refers to a polypeptide having the amino acid sequence as set forth in GenBank® Accession No. NP-057446 or Swiss-Prot Identifier GHRL\_HUMAN (each of which is incorporated herein by reference). Human ghrelin preprotein has 117 amino acids. This preprotein may undergo a post-translational processing. For example, the signal peptide (amino acids 1-23) can be removed and the remaining 94 amino acids cleaved by a protease to provide a mature 28 amino acid ghrelin (amino acids 24-51) or a mature 27 amino acid ghrelin (amino acids 24-50) and a mature 23 amino acid obestatin (amino acids 76-98). In another embodiment, "ghrelin" comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Ser-Gln-Lys-Pro-Gln-Asn-Lys-Val-Lys-Ser-Ser-Arg-Ile (e.g., SEQ ID NO:4). In another embodiment, "ghrelin" comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Lys-Ala-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO:5). In another embodiment, "ghrelin" comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Lys-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Ala-Ala-Lys-Leu-Lys-Pro-Arg (SEQ ID NO:6). In another embodiment, "ghrelin" comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Ala-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO:7). In another embodiment, "ghrelin" comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Thr-Tyr-Lys-Asn-Ile-Gln-Gln-Gln-Lys-Asp-Thr-Arg-Lys-Pro-Thr-Ala-Arg-Leu-His (SEQ ID NO:8). In yet another embodiment, ghrelin comprises a polypeptide comprising at least one modification to the natural form of an amino acid sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Lys-Leu-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg

(SEQ ID NO:9). Modifications to ghrelin may include adding or removing amino acids so long as the ghrelin so modified retains its therapeutic value (e.g., by adding or removing any of 1, 2, 3, 4 or 5 amino acids from the molecule). Other modifications can include replacing an L-amino acid with a D-amino acids.

**[0072]** Ghrelin has been shown to prevent blood brain barrier (BBB) permeability, intestinal dysfunction and systemic inflammation following TBI. The mechanism of the above observations are unclear, but may, at least in part, be a result of decreasing transcription of inflammatory cytokines and end-cascade effects including decreasing apoptosis, and blood brain barrier leakage. Ghrelin treated TBI animals had a significant decrease in brain fluorescence correlating with decreased BBB vascular permeability. It is shown that ghrelin prevents post-TBI up-regulation of Aquaporin 4 (AQP-4). Decreased expression of AQP-4 is associated with prevention of BBB breakdown and brain edema.

**[0073]** Mild brain injuries (mBI), typically including concussions, having "your bell rung", and the like, describe an insult to the brain that, in turn, can cause long term injury to the brain. To date, there has been little to no credible treatment of such mild brain injuries (mBI). Thus, there is a significant unmet need for a therapy for treating mBI that does not impose undesirable costs and delays and are effective.

**[0074]** Posttraumatic stress disorder (PTSD) is a severe anxiety central nervous system disorder that may develop in response to exposure to an event resulting in psychological trauma. PTSD may be less frequent and more enduring than the more commonly seen posttraumatic stress. PTSD is believed to be triggered by a subject witnessing or experiencing any of a wide range of events that produce intense negative feelings of fear, helplessness, or horror. This experienced fear may trigger many split-second changes in the body to prepare to defend against or avoid the danger. The "fight-or-flight" response is a healthy reaction meant to protect a person from harm. But it is believed that with PTSD, this reaction is altered. People suffering from PTSD may feel stressed or frightened even when they are not in danger. PTSD symptoms may include reliving the traumatic event in the form of flashbacks, or

nightmares, for example. Further, symptoms of PTSD may include avoidance of places or things that are reminders of the experience; feeling emotionally numb; feeling anxious; and/or losing interest in formerly enjoyable activities. People suffering from PTSD may also experience hyperarousal symptoms such as being easily startled; feeling tense; and having difficulty sleeping for example.

**[0075]** PTSD is an anxiety disorder that can develop after exposure to a terrifying event or ordeal in which grave physical harm occurred or was threatened to oneself or others. A single or repetitive mBI may leave the patients at a high risk for longer-term neuropsychological and neurodegenerative disorders, such as PTSD, which is associated with a high rate of suicide. PTSD was described in veterans of the American Civil War, and was called "shell shock," "combat neurosis," and "operational fatigue." PTSD symptoms can be grouped into three categories: (1) re-experiencing symptoms; (2) avoidance symptoms; and (3) hyperarousal symptoms. Exemplary re-experience symptoms include flashbacks (e.g., reliving the trauma over and over, including physical symptoms like a racing heart or sweating), bad dreams, and frightening thoughts. Re-experiencing symptoms may cause problems in a person's everyday routine. They can start from the person's own thoughts and feelings. Words, objects, or situations that are reminders of the event can also trigger re-experiencing. Symptoms of avoidance include staying away from places, events, or objects that are reminders of the experience; feeling emotionally numb; feeling strong guilt, depression, or worry; losing interest in activities that were enjoyable in the past; and having trouble remembering the dangerous event. Things that remind a person of the traumatic event can trigger avoidance symptoms. These symptoms may cause a person to change his or her personal routine. For example, after a bad car accident, a person who usually drives may avoid driving or riding in a car. Hyperarousal symptoms include being easily startled, feeling tense or "on edge", having difficulty sleeping, and/or having angry outbursts. Hyperarousal symptoms are usually constant, instead of being triggered by things that remind one of the traumatic event. They can make the person feel stressed and angry.

These symptoms may make it hard to do daily tasks, such as sleeping, eating, or concentrating. Therefore, generally, PTSD symptoms can include nightmares, flashbacks, emotional detachment or numbing of feelings (emotional self-mortification or dissociation), insomnia, avoidance of reminders and extreme distress when exposed to the reminders ("triggers"), loss of appetite, irritability, hypervigilance, memory loss (may appear as difficulty paying attention), excessive startle response, clinical depression, stress, and anxiety. The symptoms may last for a month, for three months, or for longer periods of time. Some embodiments herein, relate to methods of treating, reducing the symptoms or severity of symptoms of PTSD, delaying the onset of, and/or minimizing PTSD or the severity of PTSD, including PTSD resulting from one or more mild brain injuries. Such methods can include, for example, providing or administering a ghrelin compound as described herein to a subject that has experienced a mild brain injury. In some cases, the method can apply to or can include the selection of a subject that is susceptible to PTSD, has experienced it previously, or whose injury is a type of injury or is an injury that occurred in connection with an activity with a higher incidence of PTSD (e.g., military combat, injury where there was death of another person as part of the same injury causing event, etc.). In some embodiments, using a ghrelin as a therapeutic for treating PTSD associated with a one or more mild brain injuries may reduce the severity of PTSD and/or the related symptoms, and/or delay the onset of PTSD and/or the related symptoms. In some embodiments, a ghrelin is used as a therapeutic for preventing the onset PTSD or delaying PTSD, resulting from a single or from multiple mild brain injuries. In some embodiments, using ghrelin as a therapeutic for treating PTSD linked to a single injury or to multiple mild brain injuries may reduce the severity of PTSD and/or the related symptoms, and/or delay the onset of PTSD and/or the related symptoms.

**[0076]** The disclosure provides methods and compositions useful to treat or reduce the symptoms of PTSD. In some embodiments, ghrelin is used as a therapeutic for treating the PTSD associated with a single mild brain injury or concussion in an amount effective to

reduce the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, ghrelin is used as a therapeutic for preventing or delaying the onset of the PTSD or delaying the PTSD resulted from one or more of mild brain injuries. In some embodiments, using ghrelin as a therapeutic for treating the PTSD linked to repetitive or multiple mild brain injuries may reduce or delay the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, using ghrelin as a therapeutic in a combination with one or more therapeutic agents disclosed herein for treating the PTSD associated with a single mild brain injury or concussion may reduce the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, using ghrelin as a therapeutic in a combination with one or more therapeutic agents disclosed herein for treating the PTSD linked to repetitive or multiple mild brain injuries may reduce the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, using ghrelin *per se* as a therapeutic or as a sole active agent for treating the PTSD associated with a single mild brain injury or concussion may reduce the severity of PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, ghrelin *per se* is used as a therapeutic or as a sole active agent for preventing the onset of the PTSD or delaying the PTSD resulted from one or more of mild brain injuries. In some embodiments, using ghrelin *per se* as a therapeutic or as a sole active agent for treating the PTSD linked to repetitive or multiple mild brain injuries may reduce the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, using ghrelin *per se* as a therapeutic or as a sole active agent in a combination with one or more therapeutic agents disclosed herein for treating the PTSD associated with a single mild brain injury or concussion may reduce the severity of PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms. In some embodiments, using ghrelin

*per se* as an agent or as a sole active agent in a combination with one or more therapeutic agents disclosed herein for treating the PTSD linked to repetitive or multiple mild brain injuries may reduce the severity of the PTSD and/or the related symptoms, and/or delay the onset of the PTSD and/or the related symptoms.

**[0077]** The disclosure provides methods of treating Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0078]** The disclosure provides methods of preventing the onset of Post-Traumatic Stress Disorder (PTSD) or delaying the PTSD resulting from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby preventing or delaying the onset of the PTSD, or treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0079]** The disclosure provides methods of treating Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to a subject suffering from the PTSD or a related symptom a therapeutically effective amount of ghrelin to treat the PTSD and/or the related symptoms.

**[0080]** The disclosure provides methods method of preventing the onset of Post-Traumatic Stress Disorder (PTSD) or delaying the PTSD resulting from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin to delay or prevent the onset of the PTSD or to treat or reduce the severity of the PTSD and/or the related symptoms.

**[0081]** The disclosure provides compositions comprising a therapeutically effective amount of ghrelin for use in the treatment of Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject the composition, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.

**[0082]** The disclosure provides compositions comprising a therapeutically effective amount of ghrelin for use in the

treatment of preventing or delaying the onset of Post-Traumatic Stress Disorder (PTSD) or delaying the PTSD resulted from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject composition, thereby preventing the onset of the PTSD or treating the PTSD or reduce the severity of the PTSD and/or the related symptoms.

**[0083]** In some embodiments, the PTSD is associated with a single mild brain injury or concussion. In some embodiments, the PTSD is associated with repetitive or multiple mild brain injuries. In some embodiments, ghrelin is used as a sole active agent for treating the PTSD. In some embodiments, ghrelin is used in a combination with one or more therapeutic agents.

**[0084]** In some embodiments of any of the foregoing, the methods and composition can utilize ghrelin molecules that have a carbon 14 (C14) content less than found in endogenously produced ghrelin molecules or in ghrelin that has a C14 content about the same as atmospheric C14 levels. For example, ghrelin molecules can have at least one carbon atom or carbon containing moiety that is from fossil derived reagents that have a C14 content less than found in endogenous molecules or less than atmospheric levels. In some embodiments, the ghrelin molecules can have all, substantially all or at least a some carbon having a C14 content less than found endogenously or less than atmospheric levels. For example, one or more of the amino acids of a sequence can include carbon and have a C14 content less than found in endogenous amino acids or less than atmospheric levels. In other cases an entire sequence can include carbon and have a C14 content less than found endogenously or less than atmospheric levels. Still, in other embodiments, a ghrelin molecule can be modified, for example to have an octanoyl or other like group, and that octanoyl group can have a C14 content less than endogenous ghrelin C14 levels or less than atmospheric levels. In yet other embodiments, the gherlin can comprise one or more D-amino acids (e.g., .1, 2, 3, 4 or 5 D-amino acids). Further examples and embodiments are described below and elsewhere herein.

**[0085]** In some embodiments, ghrelin molecule can have a C14 content of less than 0.9 ppt, 0.95 ppt, 1.05 ppt, 1.10 ppt, 1.15 ppt, 1.2 ppt or atmospheric content of C14. In some embodiments,

ghrelin molecule can have a C14 content that is from about 1% to 50% (or any value or sub range therein) less than the content of C14 in endogenous ghrelin or the content of atmospheric C14. For example, a molecule according to some embodiments can have about 5% to about 11% less C14 content. Ghrelin with C14 content of less than 0.9 ppt, 0.95 ppt, 1.0 ppt, 1.05 ppt, 1.10 ppt, 1.15 ppt, 1.2 ppt or atmospheric content of C14, or with a lesser percentage of C14 as discussed herein, may be obtained by peptide or chemical synthesis using reactants with carbons free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. Alternatively, ghrelin with C14 content of less than 0.9 ppt, 0.95 ppt, 1.0 ppt, 1.05 ppt, 1.10 ppt, 1.15 ppt, 1.2 ppt or atmospheric content of C14 may be produced *in vitro* by enzymatic methods using starting materials with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. Such enzymatic methods may include cell-free protein synthesis system or coupled *in vitro* transcription-translation system based on cellular extracts prepared from bacteria, yeast, wheat germ, insect and/or mammalian cells using aminoacyl-tRNAs charged with amino acids with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. In an alternative method, ghrelin with C14 content of less than 0.9 ppt, 0.95 ppt, 1.0 ppt, 1.05 ppt, 1.10 ppt, 1.15 ppt, 1.2 ppt or atmospheric content of C14 may be produced by recombinant methods in bacterial, yeast, insect and/or mammalian cells following introduction of an expression system with a cDNA comprising ghrelin-encoded sequences and culturing the cells in a medium with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. Alternatively, the medium may include glucose, galactose, sugars, glycerol, pyruvate, acetates, metabolites, fatty acids, and/or amino acids with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. Methods for changing stable isotopic content of proteins may be found in Becker et al., 2008 (G. W. Becker, "Stable isotopic labeling of proteins for

quantitative proteomic applications," Briefings in Functional Genomics Proteins 7 (5): 371-382, 2008; which is incorporated herein by reference in its entirety) Ghrelin may be co-expressed with or exposed to ghrelin O-acyl transferase (GOAT) to permit fatty acid modification of the primary sequence of ghrelin at serine at amino acid position 3 so as to produce a biologically active ghrelin capable of being bound and activating the ghrelin receptor (GHSR-1a or growth hormone secretagogue receptor type 1a). The modification may be an octanoic acid modification of ghrelin so as to produce octanoyl-ghrelin with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14.

**[0086]** In some embodiments, ghrelin with C14 content of less than 0.9 ppt, 0.95 ppt, 1.0 ppt, 1.05 ppt, 1.10 ppt, 1.15 ppt, 1.2 ppt or atmospheric content of C14 (or having a percentage as discussed herein) may be obtained following modification of the primary sequence of ghrelin (SEQ ID NO: 1) with a fatty acid with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. Such fatty acids may be chemically synthesized with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14 or produced in a cell cultured in a medium wherein carbon source used to synthesize the fatty acid or fatty acids is free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. In some embodiments, the fatty acid or fatty acids are conjugated to coenzyme A (CoA) and the fatty acid in the resulting fatty acid-CoA thioesters is transferred to serine at amino acid position 3 of ghrelin by ghrelin O-acyl transferase (GOAT), so as to produce a fatty acid-modified ghrelin with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14. In some embodiments, fatty acids are straight chain fatty acids with a carbon content of C2, C3, C4, C5, C6, C7, C8, C9, C10, C11, C12, C13, C14, C15, C16, C17, C18, C19 or C20 and having a general chemical formula of  $(\text{CH})_3-(\text{CH}_2)_{n-2}-\text{COOH}$ , wherein "n" is the number of carbons in the fatty acid. In a

preferred embodiment, the fatty acid is a C8 octanoic acid or C14 tetradecanoic acid. In a more preferred embodiment, the fatty acid is octanoic acid and the fatty acid-modified ghrelin is octanoyl-modified ghrelin at serine amino acid position 3.

**[0087]** In some embodiments, fatty acid or fatty acids may be conjugated to ghrelin at serine amino acid position 3. In some embodiments, fatty acid or fatty acids may be conjugated to ghrelin at a position other than serine amino acid position 3. In some embodiments, fatty acid or fatty acids may be conjugated to ghrelin at serine amino acid position 2. In some embodiments, fatty acid or fatty acids may be conjugated to ghrelin at serine amino acid position 2 and serine amino acid position 3. In some embodiments, fatty acid or fatty acids may be conjugated to ghrelin at one or more amino acids.

**[0088]** In some embodiments, fatty acid or fatty acids may be conjugated to immature ghrelin (such as preproghrelin or proghrelin) and then fatty acid- or fatty acids-modified ghrelin is processed to a mature ghrelin that can activate the ghrelin receptor (GHSR-1a). Processing of immature ghrelin may be *in vitro* or *in vivo* and may be carried out by proteolytic enzymes. In some embodiments, fatty acid or fatty acids may be conjugated to a mature ghrelin having the amino acid sequence as provided in SEQ ID NO: 1.

**[0089]** In some embodiments, ghrelin with one or more modifications is an isolated ghrelin with one or more modifications. In a preferred embodiment, ghrelin with one or more modifications is an isolated ghrelin with one or more fatty acid modifications. In a more preferred embodiment, ghrelin with one or more modifications is an isolated ghrelin acylated at serine 3 with octanoic acid, such as an isolated octanoyl-ghrelin.

**[0090]** C14-deficient starting material used in the synthesis of ghrelin with a carbon content free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14 relative to the atmospheric content of C14 may be obtained from carbon sources not participating in atmospheric carbon cycle or by fractionating naturally occurring carbon isotope to obtain carbons free of C14, substantially free of C14, less than 1 ppt C14 or deficient in C14

relative to the atmospheric content of C14. Such carbons will be enriched in carbon-12 (C12) and/or carbon-13 (C13) and depleted of C14. Methods for isotope fractionation, enrichment or depletion are known in the art and may be based on diffusion, centrifugation, electromagnetism, laser excitation, kinetic isotope effect, chemical methods, gravity, evaporation, and cryogenic distillation among many other methods of isotope fractionation.

**[0091]** The ghrelin as used in the various embodiments described herein can be any suitable ghrelin, including for example, ghrelin that is produced using recombinant methods and ghrelin that is chemically synthesized. Regardless of the source, the ghrelin can be used and administered as described herein. The ghrelin can be isolated ghrelin, for example, ghrelin obtained following fractionation or separation of ghrelin away from other constituents present in a sample in which ghrelin is produced or synthesized as well as modified. The isolated ghrelin need not be pure ghrelin. While it is desirable to have pure ghrelin or essentially pure ghrelin, isolated ghrelin may be used without achieving absolute purity so long as ghrelin has been fractionated or separated away from constituents whose presence is not desirable in a preparation of the isolated ghrelin. It is noted that following fractionation of ghrelin, the ghrelin may be introduced into solution or used as a mixture or admixture in the presence of other components such as pharmaceutical carrier(s), lubricant(s), preservative(s), buffer(s), salt(s), sugar(s), colorant(s), pharmaceutically acceptable excipient(s) and combination thereof.

**[0092]** The disclosure also provide methods for the use for ghrelin in mBI. Mild BI, including concussion, has a significantly different pathology than other traumas associated with brain disease and illness and severe traumatic brain injuries (severe TBI) such as those causing ischemia. Mild BI does not cause the massive tissue and cellular damage as observed in severe traumatic brain disease. Rather mBI causes subtle metabolic changes within the brain, specifically oxidative stress and overproduction of reactive oxygen species (ROS) which, in turn, can damage neuroconnectivity and lead to neuron damage and encephalopathic and psychological changes with recurrent injury.

**[0093]** The disclosure utilizes ghrelin in treating mBI. Unlike severe traumatic injury to the brain, mBI does not show acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema (*i.e.*, blood brain barrier breakdown). Rather, mBI is a clinical condition generally associated with a list of clinical symptoms. It sometimes is associated with loss of consciousness but does not necessarily show significant radiological changes. Symptoms that often are reported include, without limitation, headache, "pressure in head," neck pain, nausea or vomiting, dizziness, blurred vision, balance problems, sensitivity to light, sensitivity to noise, feeling slowed down, feeling like "in a fog," "don't feel right," difficulty concentrating, difficulty remembering, fatigue or low energy, confusion, drowsiness, trouble falling asleep, more emotional, irritability, sadness and being nervous or anxious. In some non-limiting embodiments the methods herein can include or relate to treating or reducing one or more of the above listed symptoms. Most importantly, repeated concussions, as observed in the military and in sports such as football and hockey, cause a series of metabolic disturbances and the formation of ROS. These series of events lead to chronic traumatic encephalopathy (CTE) as witnessed in the autopsies of several deceased professional football players.

**[0094]** In some embodiments, ghrelin has the ability to decrease ROS and the subsequent damage caused to neurons following mBI. This is especially important for chronic concussions where preventing the inflammatory, excitatory milieu of ROS would have significant clinical impact. Ghrelin treatments decrease ROS, and therefore prevent the metabolic consequence of concussions and the chronic conditions that are associated thereof.

**[0095]** In some embodiments, the therapeutic effect of ghrelin in mBI is compared to its therapeutic effect in severe TBI. The etiology, imaging, and assessment of mBI lack a cohesive explanation for the observed cognitive deficits of chronic headaches, memory loss and sleep impairment. Only the most advanced and cumbersome technology can even detect subtle changes associated with mBI. Therefore, unlike severe TBI, where a

definitive mass lesion is identified with anatomic and cellular changes, mBI is undetectable radiologically, nevertheless, specific metabolic derangements occur. The disclosure demonstrates that ghrelin can be potent neuro-conservative agents in mBI.

**[0096]** Without being limited by any theory, the disclosure also describes the biological function of ghrelin following mBI. After mBI, the metabolic needs of the cell, including glucose requirements, increase. Mild injured cells have significant metabolic derangements causing reactive oxygen species on neurons. This metabolic stress and increased metabolic needs is the fundamental concept underlying acute concussion management. Decreasing ROS and improving glucose uptake of the brain or neuronal cells and axons can restore intracellular function and remain viable. This prevents or reduces concussion-induced dysfunction and overall improves neurocognitive outcome, as well as prevents memory loss, and chronic mBI states of headaches and development of chronic traumatic encephalopathy.

**[0097]** The disclosure provides for a method of treating mild brain injury (mild BI or mBI) in a subject, comprising administering to the subject (e.g., a subject that has mBI) an effective amount of a compound comprising ghrelin, thereby treating the mBI. The ghrelin can be administered for the purpose of treating the mBI in a therapeutically effective amount for the mBI. In some embodiments, the methods can further include selecting or identifying a subject that has suffered, is at risk of suffering, is prone to suffer, and/or is about to participate in an activity with a high risk for suffering, a mBI, prior to administration of the ghrelin.

**[0098]** In some embodiments, the mild brain injury comprises a concussion. In some embodiments, the subject that undergoes the method of treatment is a mammal. In some embodiments, the subject is a human, monkey, cow, goat, sheep, mouse, rat, cat, dog, horse, hamster, pig, fish and chicken. In some embodiments, ghrelin is administered within about 72 hours of the mBI. In some embodiments, ghrelin is administered within about 24 hours of the mBI. In some embodiments, ghrelin is administered at about 0.1,

0.3, 0.5, 0.7, 1, 2, 3, 6, 12, 18, 24, 36, 48, or 72 hours after the mBI.

**[0099]** Thus, disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the mBI or PTSD, wherein the ghrelin is the sole active agent. In some embodiments, the ghrelin has the sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO: 1). In some embodiments, the ghrelin is modified with one or more fatty acids. In some embodiments, the fatty acid is an octanoic acid. In some embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine at amino acid position 3 of SEQ ID NO: 1. In some embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine amino acid position 3 with an octanoic acid. In some embodiments, modification of ghrelin is acylation of ghrelin at serine at amino acid position 2 and/or serine at amino acid position 3 with an octanoic acid. In some embodiments, the ghrelin so modified is an octanoyl-ghrelin with an acylation at serine at amino acid position 3 by the octanoic acid. In some embodiments, the ghrelin comprise one or more D-amino acids (e.g., from 1-5, 2-6, 3-7, or 4-8 D-amino acids. In some embodiments, the mBI comprises or the PTSD results from a concussion. In some embodiments, the subject is a mammal. In some embodiments, ghrelin is administered, at least for a first administration (but not limited to), within or in less than about 24 hours, or less than 48 hours, or less than 72 hours of the mBI or the onset of PTSD or suspected PTSD, or any sub value or subrange between immediately after the injury and those time periods. In some embodiments, ghrelin is administered within not more than about 8 hours of the mBI or a PTSD causing event. In some embodiments the ghrelin can be administered with about 6 hours of the mBI or a PTSD causing event. In some embodiments, the ghrelin can be administered a single dose, or in multiple doses, for example 2-3 doses. The multiple doses can have at least a first dose administered with the time period mentioned herein and one or more subsequent doses after the time period. In some embodiments,

the ghrelin is administered in an amount that results in ghrelin blood levels being at least 1, at least 1.5, at least two times (or more) greater than the endogenous ghrelin blood levels of the subject. In some embodiments, ghrelin is administered in a single dose. In some embodiments, ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per day or per dose. In some embodiments, the dosage of ghrelin can be between 1 microgram and 15 micrograms per kg, for example about 10 µg/kg. In some embodiments, ghrelin is administered at a dosage of 2 µg/kg per dose or in some case per day. In some embodiments, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt).

**[00100]** The disclosure also provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD. The ghrelin can be the sole active agent. In some embodiments, the ghrelin has the sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO: 1). In some embodiments, the ghrelin is modified with one or more fatty acids. In some embodiments, the fatty acid is an octanoic acid. In some embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine at amino acid position 3 of SEQ ID NO: 1. In some embodiments, the ghrelin is modified at serine at amino acid position 2 and/or serine at amino acid position 3 with an octanoic acid. In some embodiments, modification of ghrelin is acylation of ghrelin at serine at amino acid position 2 and/or serine at amino acid position 3 with an octanoic acid. In some embodiments, the ghrelin so modified is an octanoyl-ghrelin with an acylation at serine at amino acid position 3 by the octanoic acid. In some embodiments, the ghrelin comprises at least one D-amino acids (e.g., 1-5, 2-6, 3-7 or 4-8 D-amino acids). In some embodiments, the mBI comprises or PTSD results from a concussion. In some embodiments, ghrelin is administered in a single dose. In some embodiments, the amount administered provides a blood level of at least 1.0 to 100 times greater the amount found endogenously in

the subject. In some embodiments, the amount administered results in a blood level of at least 55 picograms/mL. In some embodiments, ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per dosage and/or per day. In some embodiments, the dosage of ghrelin can be between 1 microgram and 15 micrograms per kg, for example about 10 µg/kg. In some embodiments, ghrelin is administered at a dosage of 2 µg/kg per dose or in some case per day. In some embodiments, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In some embodiments, the ghrelin is administered in an amount that provides blood levels of ghrelin that are at least 1.0, 2.0 times, or more, greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI.

**[00101]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition consisting essentially of ghrelin. In some embodiment, ghrelin is the sole active agent. In some embodiments, the ghrelin is administered to the subject in an amount that provides blood levels of ghrelin that are at least 1.0, 1.5, 2.0 or more times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD. In some embodiments, the ghrelin is administered in an amount that provides blood levels of ghrelin that are at least 2.0 times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD.

**[00102]** In some embodiments, the subject is free of intracranial hypertension. In some embodiments, the subject is free of blood brain barrier vascular permeability. In some embodiments, the subject is free of necrosis of brain tissue. In some embodiment, the subject is free of massive brain tissue or cellular damage associated with brain injury. In some embodiment, the subject is free of severe neuronal inflammation or significant cellular or vasogenic edema in the subject's head. In some embodiments, the subject is free of injury-associated gross anatomical changes to the subject's brain.

**[00103]** In some embodiments, the subject is free of intracranial hypertension, blood brain barrier vascular permeability, or

necrosis of brain tissue. In some embodiments, the subject is free of massive brain tissue or cellular damage associated with brain injury, severe neuronal inflammation or significant cellular or vasogenic edema in the subject's head, or injury-associated gross anatomical changes to the subject's brain.

**[00104]** In some embodiments, the subject is free of intracranial hypertension, blood brain barrier vascular permeability, necrosis of brain tissue, massive brain tissue or cellular damage associated with brain injury, severe neuronal inflammation or significant cellular or vasogenic edema in the subject's head, or injury-associated gross anatomical changes to the subject's brain. In some embodiments, the subject is free of intracranial hypertension, blood brain barrier vascular permeability, necrosis of brain tissue, massive brain tissue or cellular damage associated with brain injury, severe neuronal inflammation or significant cellular or vasogenic edema in the subject's head, and injury-associated gross anatomical changes to the subject's brain or combinations thereof.

**[00105]** In some embodiments, mBI is associated with loss of cognitive or motor skills. In some embodiments, mBI is associated with metabolic derangements. In some embodiments, mBI is associated with chronic traumatic encephalopathy (CTE). In some embodiments, mBI is associated with neuroconnectivity damage. In some embodiments, mBI is associated with encephalopathic and/or psychological changes. In some embodiments, mBI is associated with loss of consciousness.

**[00106]** In some embodiments, mBI is associated with loss of cognitive or motor skills and metabolic derangements. In some embodiment, mBI is associated with chronic traumatic encephalopathy (CTE) and neuroconnectivity damage. In some embodiments, mBI is associated with psychological changes and loss of consciousness. In some embodiments, mBI is associated with loss of cognitive or motor skills, metabolic derangements, chronic traumatic encephalopathy (CTE), neuroconnectivity damage, psychological changes, or loss of consciousness. In some embodiments, mBI is associated with loss of cognitive or motor skills, metabolic derangements, chronic traumatic encephalopathy (CTE),

neuroconnectivity damage, psychological changes, and loss of consciousness or combination thereof.

**[00107]** The disclosure provides a method for reducing, inhibiting and/or minimizing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or the onset or causation of PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin so administered reduces oxidative stress and overproduction of ROS, thereby inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or the onset or causation of PTSD in a subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and ghrelin is the sole active agent.

**[00108]** The disclosure further provides a method for inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI), including concussion, or concussion or the onset or causation of PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin so administered reduces oxidative stress and overproduction of ROS and wherein the ghrelin is the sole active agent, thereby inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00109]** The disclosure additionally provides a method for inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or the onset or causative event for PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt),

wherein the ghrelin so administered reduces oxidative stress and overproduction of ROS, thereby inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00110]** The disclosure provides a method for inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or the onset or causative event for PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein the ghrelin so administered reduces oxidative stress and overproduction of ROS and wherein the ghrelin is the sole active agent, thereby inhibiting damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject

**[00111]** Additionally, the disclosure provides a method of restoring intracellular function after mild brain injury (mBI) in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin so as to decrease ROS and improve glucose uptake of brain or neuronal cells and axons, thereby restoring intracellular function after mBI in the subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00112]** The disclosure further provides a method of restoring intracellular function after mild brain injury (mBI) or after or during a PTSD event in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin so as to decrease ROS and improve glucose uptake of brain or neuronal cells and axons, wherein the ghrelin is the sole active agent, thereby restoring intracellular function after mBI or PTSD in the subject.

**[00113]** The disclosure further still provides a method of restoring intracellular function after mild brain injury (mBI) in a

subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt) so as to decrease ROS and improve glucose uptake of brain or neuronal cells and axons, thereby restoring intracellular function after mBI in the subject.

**[00114]** The disclosure also provides a method of restoring intracellular function after mild brain injury (mBI) or after a PTSD event of during a PTSD event in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt) so as to decrease ROS and improve glucose uptake of brain or neuronal cells and axons, wherein the ghrelin is the sole active agent, thereby restoring intracellular function after mBI or PTSD in the subject.

**[00115]** Further, the disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis, or massive tissue or cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the mBI or PTSD. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00116]** The disclosure additionally provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis, or massive tissue or cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD.

**[00117]** The disclosure also provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis, or massive tissue or cellular damage associated with brain injury, comprising administering to the

subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating the mBI or PTSD.

**[00118]** The disclosure also provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis, or massive tissue or cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD.

**[00119]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis or massive tissue and cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00120]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis or massive tissue and cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD.

**[00121]** The disclosure yet further provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular

permeability, necrosis or massive tissue and cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt) and in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD.

**[00122]** The disclosure also provides a method of treating mild brain injury (mBI) or PTSD in a subject free of brain edema, intracranial hypertension, blood brain barrier vascular permeability, necrosis or massive tissue and cellular damage associated with brain injury, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt) and in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD.

**[00123]** The disclosure further provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00124]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin is the sole active agent, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00125]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a

therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00126]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein the ghrelin is the sole active agent, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00127]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00128]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin is the sole active agent, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00129]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that

provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00130]** The disclosure provides a method of treating metabolic derangements associated with mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent, thereby treating metabolic derangements associated with mBI or concussion or PTSD in the subject.

**[00131]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00132]** The disclosure also provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS and wherein the ghrelin is the sole active agent, thereby preventing damage to neurons

associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00133]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein ghrelin treatment reduces oxidative stress and overproduction of ROS, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00134]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein ghrelin treatment reduces oxidative stress and overproduction of ROS and wherein the ghrelin is the sole active agent, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00135]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or

concussion or PTSD in the subject. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00136]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS and wherein the ghrelin is the sole active agent, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00137]** The disclosure provides a method for preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS and wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00138]** The disclosure provides a method for reducing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of mild brain injury (mBI) or concussion or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are

at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein ghrelin treatment reduces oxidative stress and overproduction of ROS and wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent, thereby preventing damage to neurons associated with oxidative stress and overproduction of reactive oxidative species (ROS) with one or more incidence of the mBI or concussion or PTSD in the subject.

**[00139]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00140]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00141]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00142]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00143]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00144]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin is the sole active agent, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00145]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00146]** The disclosure provides a method of treating mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject, free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema, a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent, thereby treating the mBI or PTSD in the subject free of acute neuronal histological changes, severe neuronal inflammation or significant cellular or vasogenic edema.

**[00147]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating or reducing symptoms of the mBI or PTSD. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00148]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin, wherein the ghrelin is the sole active agent, thereby treating or reducing symptoms of the mBI or PTSD.

**[00149]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), thereby treating or reducing symptoms of the mBI or PTSD.

**[00150]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin having a carbon 14 (C14) content of less than 1 part per trillion (ppt), wherein the ghrelin is the sole active agent, thereby treating or reducing symptoms of the mBI or PTSD.

**[00151]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, thereby treating or reducing symptoms of the mBI or PTSD. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt). In one embodiment, the ghrelin is the sole active agent. In one embodiment, the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent.

**[00152]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin is the sole active agent, thereby treating or reducing symptoms of the mBI or PTSD.

**[00153]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin has a carbon 14

(C14) content of less than 1 part per trillion (ppt), thereby treating or reducing symptoms of the mBI or PTSD.

**[00154]** The disclosure provides a method of treating or reducing symptoms of mild brain injury (mBI) or PTSD in a subject, comprising administering to the subject a therapeutically effective amount of ghrelin in an amount that provides blood levels of ghrelin that are at least 1.5 times greater than endogenous ghrelin blood levels of the subject, wherein the ghrelin has a carbon 14 (C14) content of less than 1 part per trillion (ppt) and is the sole active agent, thereby treating or reducing symptoms of the mBI or PTSD.

**[00155]** In some embodiments, the symptoms of mBI are any of headache, "pressure in head," neck pain, nausea or vomiting, dizziness, blurred vision, balance problems, sensitivity to light, sensitivity to noise, feeling slowed down, feeling like "in a fog," "don't feel right," difficulty concentrating, difficulty remembering, fatigue or low energy, confusion, drowsiness, loss of consciousness, trouble falling asleep, more emotional, irritability, sadness and being nervous or anxious.

**[00156]** As discussed herein, ghrelin can be used to treat and/or reduce the severity of mild brain injuries, such as for example, concussions. The ghrelin can be provided or administered as described herein. As described elsewhere herein, ghrelin can be the sole active agent administered to a subject. For example, ghrelin can be the only active ingredient for treating the mBI or reducing the severity of the mBI. In some embodiments, additional active agents are specifically excluded. For example, in some of the methods herein where ghrelin is administered, the methods can specifically exclude the use or administration of other compounds, such as, for example, selective androgen receptor modulators, steroid receptor modulators, ligand for a glucocorticoid receptor, or a modulator of the NF-kB signaling pathway.

**[00157]** In the embodiments described herein it should be understood that mBI is different from and is not severe TBI.

**[00158]** In some embodiments, mBI or concussion is diagnosed by non-invasive functional imaging following impact. In some embodiments, mBI or concussion is diagnosed by neuropsychological

assessment following impact. In some embodiment, mBI or concussion is diagnosed by neuroanatomical assessment of the brain following impact. In some embodiment, the neuroanatomical assessment of the brain following impact is at the cellular level or molecular level. In some embodiment, mBI or concussion is diagnosed by cognitive assessment following impact. In some embodiment, mBI or concussion is diagnosed by assessment of alertness following impact. In some embodiments, mBI and concussion can be diagnosed and/or assessed by evaluating one or more of consciousness, assessment of alertness, assessment of orientation, assessment of cognitive function, assessment of psychological state, neuropsychological assessment.

**[00159]** In some embodiments, mBI or concussion is diagnosed using a Glasgow Coma Scale (GCS) having a score of 13 to 15. In some embodiments, mBI or concussion is diagnosed using a Glasgow Coma Scale (GCS) having a score of 13 or greater.

**[00160]** In contrast, by way of example, severe TBI generally has a GCS score of 8 or below.

**[00161]** In some embodiments, an intravenous injection of ghrelin is employed. The administration route must ensure that the non-degraded, bioactive form of the peptide will be the dominating form in the circulation, which will reach and stimulate the ghrelin receptors in order to obtain the maximum effect of ghrelin treatment on mBI. In some embodiments, ghrelin is administered within about 30 minutes of the incident that results in mBI. In some embodiments, ghrelin is administered within about 30 minutes to about 2 hours of the incident that results in mBI. In some embodiments, ghrelin is administered within about 30 minutes to about 6 hours of the incident that results in mBI. In some embodiments, ghrelin is administered within about 30 minutes to about 12 hours of the incident that results in mBI.

**[00162]** A typical dosage is in a concentration equivalent to from 10 ng to 10 mg ghrelin per kg bodyweight. In some embodiments, ghrelin is administered in a concentration equivalent to from about 0.1 µg to about 1 mg ghrelin per kg bodyweight, such as from about 0.5 µg to about 0.5 mg ghrelin per kg bodyweight, such as from about 1.0 µg to about 0.1 mg ghrelin per kg

bodyweight, such as from about 1.0 µg to about 50 µg ghrelin per kg bodyweight, such as from about 1.0 µg to about 10 µg ghrelin per kg bodyweight. In some embodiments, about 10 µg ghrelin powder is reconstituted in about 100 µL of a sterile saline solution before administration. In some embodiments, the sterile saline solution is contained in an IV bag for ease of delivery.

**[00163]** In some embodiments, the ghrelin and ghrelin or ghrelin agonist is used in an assay to assess the ability of candidate compounds to effect increased uncoupling protein-2 (UCP-2) expression including increased UCP-2 expression in mitochondria. In such assays, ghrelin is used as a control to determine the relative efficacy of the candidate compound or compounds. Suitable assays include by way of example only competitive assays for binding of a candidate compound or compounds to growth hormone secretagogue receptor 1a (*i.e.*, GHSR-1a) in the presence of ghrelin as well as frontal affinity chromatography.

**[00164]** Any competitive binding assay known in the art is applicable for binding of a candidate compound or compounds to growth hormone secretagogue receptor in the presence of ghrelin, using either heterogeneous or homogeneous methods, with one or more reagents, and with labels and detection methods. By way of non-limiting example, detection methods may include radioactive methods; enzyme techniques using intact enzymes of many types including, for example,  $\beta$ -galactosidase, glucose 6-phosphate dehydrogenase, alkaline phosphatase, horseradish peroxidase, or glucose oxidase; techniques using enzyme fragments, such as  $\beta$ -galactosidase complementation assays; detection systems including chromogenic substrates; fluorescent methods detected by direct fluorescence, time-resolved fluorescence, fluorescence polarization, or fluorescence energy transfer; and chemical or bioluminescence detection systems.

**[00165]** In some embodiments, frontal affinity chromatography (FAC) can be used for screening of compound libraries. The basic premise of FAC is that continuous infusion of a compound will allow for equilibration of the ligand between the free and bound states, where the precise concentration of free ligand is known. The detection of compounds eluting from the column can be accomplished

using methods such as fluorescence, radioactivity, or electrospray mass spectrometry. The former two methods usually make use of either a labeled library, or use a labeled indicator compound, which competes against known unlabeled compounds, getting displaced earlier if a stronger binding ligand is present.

**[00166]** In some embodiments, a patient suffering loss of cognitive or motor skills due to mBI and, in particular, repetitive mBI, can be monitored for therapy or progression of such skills by correlating the ghrelin level in the patient's brain over time. As the ghrelin levels decrease, there will be an increased need for intervention.

**[00167]** This invention also provides for methods for measuring ghrelin levels before starting a sport or activity, for example prior to the beginning of football season (or any other sport or activity, including those listed elsewhere herein), and monitoring ghrelin levels during the season to ascertain if the player or participant is at a level not qualified to play or participate. The methods can include the use of any suitable measurement or assay technique for measuring ghrelin levels, such as from blood to determine if blood levels correlate to brain levels.

**[00168]** With the benefit of the instant embodiments, the skilled artisan can select any suitable technique for measuring ghrelin levels. A number of assays known in the art for measuring a protein or hormone level are applicable for measuring ghrelin levels. By way of non-limiting example, assays such as a blood sugar test by extracting a drop of blood and putting it into a device, can quantitatively assess the amount of ghrelin, or an assay involving measuring a range of substances whereby a specific reaction chemistry is followed photo-metrically with time, for example by utilizing an antibody specific to ghrelin that is coated onto latex particles and measuring the increased turbidity that is produced when ghrelin being measured promotes aggregation of the latex particles as the reaction between ghrelin and anti-ghrelin antibody proceeds. This measurement of increasing turbidity can be achieved using a conventional photometer and using the associated scientific principles of photometric measurements. Such

concentration dependent turbidity is then compared to that produced by standards which are established in the art.

**[00169]** Further compatible, but non-limiting, methodologies include carrying out a series of enzyme-linked reactions in solution, where ghrelin in the plasma fraction of a whole blood sample is altered by an enzyme-promoted reaction to ultimately derive a colored dye from colorless reaction constituents. The color is developed in a time dependent way and monitored photometrically. This measurement of color change can also be achieved using a conventional photometer using the associated scientific principles of photometric measurements. Such concentration dependent change in transmission is then compared to that produced by standards.

**[00170]** In whole blood samples, hematocrit or percentage of red blood cells by volume in the whole blood sample is a variable and can be taken into account when analyzing ghrelin levels that are present in the plasma component. As the hematocrit of a patient's blood rises, so does the volume of plasma in a fixed volume sample, which is introduced into the test device decreases and vice versa. Since it is the plasma component which exclusively carries the ghrelin levels being measured, then the lower the volume of plasma component added to the reaction mix, the lower the resulting concentration of the substance being measured in that reaction mix and the resulting assayed value and vice versa.

**[00171]** Any analysis that produces a concentration of a plasma substance in whole blood may be corrected for variations in hematocrit to give a true plasma concentration. It can be most useful in these situations to measure two substances, one of which is ghrelin under investigation and the other which is considered to be a marker by which to estimate or normalize the sample hematocrit. The hemoglobin concentration of whole blood, after red blood cells are lysed, is directly proportional to the red blood cell volume in the whole blood sample.

**[00172]** Ghrelin described herein can be formulated as a pharmaceutical composition, e.g., flash frozen or lyophilized for storage and/or transport. In some embodiments, the compound can be in a composition with sterile saline, for example. In some

embodiments, a ghrelin can be reconstituted in such saline or other acceptable diluent. In some embodiments, about 10 µg ghrelin powder is reconstituted in about 100 µL saline solution before administration. In addition, the composition can be administered alone or in combination with a carrier, such as a pharmaceutically acceptable carrier or a biocompatible scaffold. Compositions of the invention may be conventionally administered parenterally, by injection, for example, intravenously, subcutaneously, or intramuscularly. Additional formulations which are suitable for other modes of administration include oral formulations. Oral formulations include such normally employed excipients such as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10% to about 95% of active ingredient, about 25% to about 70%.

**[00173]** Typically, compositions are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective for the disease or condition by treated. The quantity to be administered depends on the subject to be treated. Precise amounts of the composition to be administered depend on the judgment of the practitioner. Suitable regimes for initial administration and boosters are also variable, but are typified by an initial administration followed by subsequent administrations.

**[00174]** In some embodiments, additional pharmaceutical compositions are administered to a subject to support or augment the compositions as described herein. Different aspects of the present invention involve administering an effective amount of the composition to a subject. Additionally, such compositions can be administered in combination with other agents. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

**[00175]** The phrases "pharmaceutically acceptable" or "pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic, or other

untoward reaction when administered to an animal, or human. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredients, its use in immunogenic and therapeutic compositions is contemplated.

**[00176]** Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid or lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene or water. Nasal aerosol or inhalation formulations may be prepared, for example, as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, employing fluorocarbons, and/or employing other solubilizing or dispersing agents.

**[00177]** An effective amount of therapeutic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed above in association with its administration, *i.e.*, the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the result and/or protection desired. Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is

therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

**[00178]** Ghrelin compositions can be produced using techniques well known in the art. For example, a polypeptide region of ghrelin can be chemically or biochemical synthesized and modified. Techniques for chemical synthesis of polypeptides are well known in the art (Lee V. H. L. in "Peptide and Protein Drug Delivery", New York, N.Y., M. Dekker, 1990). Examples of techniques for biochemical synthesis involving the introduction of a nucleic acid into a cell and expression of nucleic acids are provided in Ausubel F. M. et al., "Current Protocols in Molecular Biology", John Wiley, 1987-1998, and Sambrook J. et al., "Molecular Cloning, A Laboratory Manual", 2d Edition, Cold Spring Harbor Laboratory Press, 1989, each of which is incorporated herein by reference. Another exemplary technique described in U.S. Pat. No. 5,304,489, incorporated herein by reference, is the use of a transgenic mammal having mammary gland-targeted mutations, which result in the production and secretion of synthesized ghrelin in the milk of the transgenic mammal.

**[00179]** Ghrelin can also be produced recombinantly using routine expression methods known in the art. The polynucleotide encoding the desired ghrelin (e.g., primarily sequence of ghrelin as given in SEQ ID NO: 1) is operably linked to a promoter into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems are used in forming recombinant ghrelin. Ghrelin is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Isolated ghrelin may be modified or further modified at serine amino acid position 2 and/or serine amino acid position 3 by fatty acid acylation *in vivo* or *in vitro*, with the latter *in vitro* acylation reaction condition comprising fatty acid thioester, ghrelin, and microsomes comprising ghrelin O-acyl transferase (GOAT). In some embodiments, acyl ghrelin modified with fatty acid at serine amino acid position 2 and/or serine amino acid position 3 is isolated from cellular or reaction components.

**[00180]** Ghrelin compositions can include pharmaceutically acceptable salts of the compounds therein. These salts will be ones which are acceptable in their application to a pharmaceutical use, meaning that the salt will retain the biological activity of the parent compound and the salt will not have untoward or deleterious effects in its application and use in treating diseases. Pharmaceutically acceptable salts are prepared in a standard manner.

**[00181]** Ghrelin compositions may be administered in the form of an alkali metal or earth alkali metal salt thereof, concurrently, simultaneously, or together with a pharmaceutically acceptable carrier or diluent, especially and in the form of a pharmaceutical composition thereof, whether by various routes (e.g., oral, rectal, parenteral, subcutaneous) in an effective amount.

**[00182]** The disclosure provides for a method of reducing the incidence or severity of mBI or PTSD in a subject, comprising administering to the subject an effective amount of a compound comprising ghrelin, thereby reducing the incidence or severity of the mBI or PTSD. The disclosure also provides for methods of reducing the incidence or severity of mBI or PTSD in a subject, comprising administering to the subject an effective amount of ghrelin, thereby reducing the incidence or severity of the mBI or PTSD. This invention further provides for methods of reducing the amount of time needed to recover from a mild brain injury, comprising administering to a patient suffering from a mild brain injury a therapeutically effective amount of ghrelin within a certain period (e.g., 72 hours) of the mild brain injury.

**[00183]** In some embodiments, ghrelin is administered prior to an event or activity with a potential for occurrence of mBI or PTSD. In some embodiments, the event or activity is participation in a sporting event, physical training, or combat. In some embodiments, the event or activity is baseball, basketball, rugby, football, hockey, lacrosse, soccer, cycling, boxing, gymnastics, a martial art, a mixed martial art, a military exercise, automobile racing, snow skiing, snowboarding, ice skating, skateboarding, motocross, mountain biking, motorcycle and ATV riding, and the like. In some embodiments, the subject has not suffered mBI or PTSD. In some

embodiments, the subject has a history of mBI or PTSD or is susceptible to mBI or PTSD.

**[00184]** In some embodiments, an administration route for a ghrelin is selected from: buccal delivery, sublingual delivery, transdermal delivery, inhalation and needle-free injection, such as using the methods developed by PowderJet. For inhalation, ghrelin can be formulated using methods known to those skilled in the art, for example an aerosol, dry powder or solubilized such as in microdroplets, in a device intended for such delivery (such as commercially available devices and formulation technologies from Aradigm Corp. (Hayward, Calif.), Alkermes, Inc. (Cambridge, Mass.), Nektar Therapeutics (San Carlos, Calif.), Windgap Medical, Inc. (e.g., Abiliject™) or MannKind Corporation (Valencia, Calif.; e.g., Technosphere®, Dreamboat®, and Cricket™ technologies)).

**[00185]** In some embodiments, the ghrelin is administered via a powder or stable formulation, wherein ghrelin is formulated in a dosage form selected from the group consisting of: liquid, beverage, medicated sports drink, powder, capsule, chewable tablet, caplet, swallowable tablet, buccal tablet, troche, lozenge, soft chew, solution, suspension, spray, suppository, tincture, decoction, infusion, and a combination thereof.

**[00186]** In some embodiments, ghrelin is administered via inhalation, oral, intravenous, parenteral, buccal, subcutaneous, transdermal, patch, sublingual, intramuscular, or intranasal. In some embodiments, ghrelin is administered in a single dose. In some embodiments, ghrelin is administered in multi-doses. In some embodiments, ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per day (or any sub value or sub range there between, e.g., 0.1 µg/kg per day to 5 mg/kg per day). In some embodiments, a dosing regimen (2 µg/kg, for example delivered intravenously) is administered within 8 hours following injury. The dosing is a one-time dose with possible recurrent dosing based on patient symptoms.

**[00187]** Nasal delivery is a non-invasive route for therapeutics targeting the central nervous system because of relatively high permeability of nasal epithelium membrane, avoidance of hepatic first pass elimination. Nasal delivery is easy to administer and

allows for self-medication by an individual. Nasal mucociliary clearance is an important limiting factor to nasal drug delivery. Nasal mucociliary clearance severely limits the time allowed for drug absorption to occur and may effectively prevent sustained drug administration. However, it has been documented that nasal administration of certain hormones has resulted in a more complete administration. In some embodiments, the present disclosure utilizes nasal delivery of ghrelin.

**[00188]** In some embodiments, a composition comprising ghrelin that is suitable for nasal administration may include one or more bioadhesive polymers. Some polymers such as carbopol, can adhere onto the nasal mucosa for reasonably prolonged periods, preventing rapid nasal clearance. In some embodiments, a composition suitable for nasal administration, the percentage of bioadhesive polymer in a suitable solution of ghrelin is about 0.1%. In some embodiments, a composition suitable for nasal administration, the percentage of bioadhesive polymer in a suitable solution of ghrelin is about 0.5%. In some embodiments, a composition suitable for nasal administration, the percentage of bioadhesive polymer in a suitable solution of ghrelin is about 1%. In some embodiments, a composition suitable for nasal administration, the percentage of bioadhesive polymer in a suitable solution of ghrelin is about 5%.

**[00189]** In some embodiments, a composition comprising ghrelin that is suitable for nasal administration may include one or more surfactants. Surfactants that may be used in the compositions of the present invention include different polyethylene glycols (PEGs) and polyethylene glycol-derivatives. In some embodiments, a composition suitable for nasal administration, the percentage of surfactant in a suitable solution of ghrelin is about 1%. In some embodiments, a composition suitable for nasal administration, the percentage of surfactant in a suitable solution of ghrelin is about 2%. In some embodiments, a composition suitable for nasal administration, the percentage of surfactant in a suitable solution of ghrelin is about 5%. In some embodiments, a composition suitable for nasal administration, the percentage of surfactant in a suitable solution of ghrelin is about 10%.

**[00190]** In some embodiments, a composition comprising ghrelin that is suitable for nasal administration may include one or more buffering agents for controlling the pH of the composition. Buffering agents that may be used in the compositions of the present invention include citric acid and sodium citrate dihydrate. In some embodiments, a composition suitable for nasal administration, the percentage of buffering agent in a suitable solution of ghrelin is about 0.001%. In some embodiments, a composition suitable for nasal administration, the percentage of buffering agent in a suitable solution of ghrelin is about 0.005%. In some embodiments, a composition suitable for nasal administration, the percentage of buffering agent in a suitable solution of ghrelin is about 0.01%. In some embodiments, a composition suitable for nasal administration, the percentage of buffering agent in a suitable solution of ghrelin is about 0.1%.

**[00191]** In some embodiments, the osmolarity of the composition comprising ghrelin may be controlled by propylene glycol. When a composition comprising ghrelin is a gel, the composition may include a gelling agent such as hydroxylpropyl cellulose, carbopols, carboxymethylcellulose, and ethylcellulose. In some embodiments, the composition comprising ghrelin may include a preservative such as ethylenediaminetetraacetic acid (EDTA) and benzalkonium chloride. Non-limiting examples of suitable solvents for compositions of the present invention include water, vegetable oil and ethanol. In some embodiments, the use of a nasal inhalant reduces the concentration required to treat mBI and prevent unwanted side effects.

**[00192]** In some embodiments, nasal administration is a more practical means of delivery in a military or sport setting. In some embodiments, the present invention provides a method for improving the standard of care for preventing or treating mBI in military personnel or athletes through a prophylactic and post-acute intranasal therapeutic. In some embodiments, the active ingredient of the therapeutic is ghrelin. In some embodiments, ghrelin may be part of a formulation that is delivered intranasally to facilitate ease of access and use in the field and to minimize the dose required further limiting side effects. Using ghrelin as a

therapeutic may reduce poor outcomes following injury, especially neuropsychological and neurodegenerative disorders including Chronic Traumatic Encephalopathy (CTE) and Post-Traumatic Stress Disorder (PTSD) linked to repetitive brain injuries, an increasing concern for today's military personnel and athletes.

**[00193]** In some embodiments, the present invention provides compositions comprising ghrelin that are applied as nasal drops, eye drops and nasal sprays. For the nasal application, a solution or suspension may be used which is applied as spray, *i.e.*, in the form of a fine dispersion in air or by means of a conventional spray-squeeze bottle or pump. Suitable nontoxic pharmaceutically acceptable carriers for use in a drug delivery system for intranasal administration of ghrelin may include, but not limited to, carriers used for nasal pharmaceutical formulations for other steroids, such as estrogen.

**[00194]** In some embodiments, formulations of the present invention may contain a preservative and/or stabilizer. These include, but not limited to, ethylene diamine tetraacetic acid (EDTA) and its alkali salts (for example dialkali salts such as disodium salt, calcium salt, calcium-sodium salt), lower alkyl p-hydroxybenzoates, chlorhexidine (for example in the form of the acetate or gluconate) and phenyl mercury borate. Other suitable preservatives are: pharmaceutically useful quaternary ammonium compounds, for example cetylpyridinium chloride, tetradecyltrimethyl ammonium bromide, generally known as "cetrimide", N-Benzyl-N,N-dimethyl-2-{2-[4-(2,4,4-trimethylpentan-2-yl)phenoxy]ethoxy}ethanaminium chloride, generally known as "benzethonium chloride" and myristyl picolinium chloride. Each of these compounds may be used in a concentration of about 0.002 to 0.05%, for example about 0.02% (weight/volume in liquid formulations, otherwise weight/weight). In some embodiments, preservatives among the quaternary ammonium compounds are, but not limited to, alkylbenzyl dimethyl ammonium chloride and mixtures thereof, for example, benzalkonium chloride.

**[00195]** In some embodiments, the present invention provides for a treatment strategy for athletes who have suffered a mBI that may

not only reduce the time required for safe return to play but also provide protection from future mBI.

**[00196]** Intranasal (IN) administrations may have fewer side effects than intraperitoneal (IP) administrations due to a shift in pharmaceutical research to nasal sprays, drops and gels: the nasal route of drug administration continues to receive increasing attention from pharmaceutical scientists and clinicians because this route circumvents hepatic first pass elimination associated with oral delivery, is easily accessible and suitable for self-medication. Intranasal administration also particularly suits drugs targeting the brain because certain drug solutions can bypass the blood-brain barrier (BBB) and reach the central nervous system (CNS) directly from the nasal cavity—uptake of these drugs depends on their molecular weight and lipophilicity. The intranasal delivery increases brain levels of the drug while decreasing systemic concentrations and thus should have less harmful side effects.

**[00197]** In some embodiments, the present invention provides a method of prophylactically administering a composition comprising ghrelin to individuals who are involved in activities, such as contact sports or serving in the armed forces, where there is a possibility of the individuals suffering mBI. In some embodiments, the present invention provides a method for acutely treating individuals who have suffered mBI. For acute treatments, nasal administration of the composition comprising ghrelin may reduce the time for uptake and increase the concentration of ghrelin that reaches the blood or brain.

**[00198]** For oral administration, such excipients include, e.g., pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, gelatin, sucrose, magnesium carbonate, and the like. In powders, the carrier is a finely divided solid, which is a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. The powders and tablets contain from one to about seventy percent of the active compound. Suitable carriers are magnesium carbonate,

magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include a composition comprising an active compound disclosed herein with encapsulating material as carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be as solid forms suitable for oral administration.

**[00199]** Also included are solid form preparations, which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

**[00200]** Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, toothpaste, gel dentifrice, chewing gum, or solid form preparations which are intended to be converted shortly before use to liquid form preparations. Emulsions may be prepared in solutions in aqueous propylene glycol solutions or may contain emulsifying agents such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents. Solid form preparations include solutions, suspensions, and emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

**[00201]** The administration of ghrelin is based on suitable dosing regimens that take into account factors well-known in the art including, e.g., type of subject being dosed; age, weight, sex and medical condition of the subject; the route of administration; the renal and hepatic function of the subject; the desired effect; and the particular compound employed. Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

**[00202]** In some embodiments, ghrelin composition is administered subcutaneously. In some embodiments, ghrelin composition is administered as a bolus, wherein the administration form may be any suitable parenteral form.

**[00203]** Pharmaceutical compositions for parenteral administration include sterile aqueous and non-aqueous injectable solutions, dispersions, suspensions or emulsions, as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Other suitable administration forms include suppositories, sprays, ointments, creams, gels, inhalants, dermal patches, implants, pills, tablets, lozenges and capsules.

**[00204]** A typical non-limiting dosage is in a concentration equivalent to from 10 ng to 10 mg ghrelin per kg bodyweight. In some embodiments, ghrelin is administered in a concentration equivalent to from about 0.1 µg to about 1 mg ghrelin per kg bodyweight, such as from about 0.5 µg to about 0.5 mg ghrelin per kg bodyweight, such as from about 1.0 µg to about 0.1 mg ghrelin per kg bodyweight, such as from about 1.0 µg to about 50 µg ghrelin per kg bodyweight, such as from about 1.0 µg to about 10 µg ghrelin per kg bodyweight.

**[00205]** In some embodiments, an intravenous injection of ghrelin is employed. The administration route must ensure that the non-degraded, bioactive form of the peptide will be the dominating form in the circulation, which will reach and stimulate the ghrelin receptors in order to obtain the maximum effect of ghrelin treatment on mBI. In some embodiments, ghrelin is administered

within about 30 minutes of the incident that results in mBI. In some embodiments, the ghrelin is administered within about 30 minutes to about 2 hours of the incident that results in mBI. In some embodiments, ghrelin is administered within about 30 minutes to about 6 hours of the incident that results in mBI. In some embodiments, ghrelin is administered within about 30 minutes to about 12 hours of the incident that results in mBI.

**[00206]** Ghrelin compositions may be formulated for parenteral administration (e.g., by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for example solutions in aqueous polyethylene glycol. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution for constitution before use with a suitable vehicle, e.g., sterile, pyrogen-free water. Aqueous solutions should be suitably buffered if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

**[00207]** The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions comprising the active ingredient that are adapted for administration by encapsulation in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. Sterile injectable solutions are prepared by incorporating ghrelin or pharmaceutical acceptable salt thereof in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by, e.g., filter sterilization.

**[00208]** Ghrelin compounds can also be delivered topically. Regions for topical administration include the skin surface and also mucous membrane tissues of the rectum, nose, mouth, and

throat. Compositions for topical administration via the skin and mucous membranes should not give rise to signs of irritation, such as swelling or redness.

**[00209]** Ghrelin compounds may include a pharmaceutical acceptable carrier adapted for topical administration. Thus, the composition may take the form of, for example, a suspension, solution, ointment, lotion, cream, foam, aerosol, spray, suppository, implant, inhalant, tablet, capsule, dry powder, syrup, balm or lozenge. Methods for preparing such compositions are well known in the pharmaceutical industry.

**[00210]** Ghrelin compounds may be administered transdermally, which involves the delivery of a pharmaceutical agent for percutaneous passage of the drug into the systemic circulation of the patient. The skin sites include anatomic regions for transdermally administering the drug and include the forearm, abdomen, chest, back, buttock, and the like. Transdermal delivery is accomplished by exposing a source of the active compound to a patient's skin for an extended period of time. Transdermal patches can add advantage of providing controlled delivery of a compound complex to the body. Such dosage forms can be made by dissolving, dispersing, or otherwise incorporating ghrelin compound in a proper medium, such as an elastomeric matrix material. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate-controlling membrane or dispersing the compound in a polymer matrix or gel.

**[00211]** Ghrelin compositions may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses. The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. The compounds can be provided in a kit. Such a kit typically contains an active compound in dosage forms for administration. The kit comprises an amount of dosage units corresponding to the relevant dosage regimen. In some embodiment, the kit comprises a pharmaceutical composition comprising a ghrelin compound or a pharmaceutically acceptable salt thereof and pharmaceutically acceptable carrier, vehicles and/or

excipients, said kit having multiple dosage units. The dosage units comprise an amount of ghrelin or a salt thereof equivalent to from about 0.3 µg to about 600 mg ghrelin, such as from about 2.0 µg to about 200 mg ghrelin, such as from about 5.0 µg to about 100 mg ghrelin, such as from about 10 µg to about 50 mg ghrelin, such as from about 10 µg to about 5 mg ghrelin, such as from about 10 µg to about 1.0 mg ghrelin.

**[00212]** The kit contains instructions indicating the use of the dosage form to achieve a desirable affect and the amount of dosage form to be taken over a specified time period. Accordingly, in one embodiment the kit comprises instructions for administering the pharmaceutical composition. In particular said instructions may include instructions referring to administration of said pharmaceutical composition after mBI or concussion, or at the most about 12 hours after the incident causing mBI or concussion, such as at the most about 6 hours after the incident causing mBI or concussion, such as at the most about 3 hours after the incident causing mBI or concussion, such as at the most about 1 hours after the incident causing mBI or concussion, such as at the most about 30 minutes after the incident causing mBI or concussion, such as at the most about 10 minutes after the incident causing mBI or concussion, such as at the most about 5 minutes after the incident causing mBI or concussion.

#### **EXAMPLES**

**[00213]** The following Examples are intended to further illustrate certain embodiments of the disclosure and are not intended to limit its scope.

#### **EXAMPLE 1**

**[00214]** Ghrelin administration reduces oxidative burst in inflammatory cells following mBI. Since no well-accepted animal model exists for concussions, a very small cerebral lesion that closely mimics mBI is used as a model of mBI. C57/B6 mice anesthetized with 5% isoflurane in oxygen (1.7 L/min) are given 0.3 mg/kg buprenorphine subcutaneously for analgesia prior to infliction of the mild brain injury. Anesthesia is assessed by paw pinch reflex. After creating a burr hole through the dura with a dental drill, a lesion using a controlled cortical impactor (CCI)

is used to create injury 1 mm lateral and posterior to the bregma (5.0 mm/sec at a depth of 1.0 mm).

**[00215]** Animals are separated into three treatment groups: 1) Sham, 2) mBI, and 3) mBI plus ghrelin. A variety of doses can be tested depending upon the particular ghrelin. Ghrelin can be obtained for example from Phoenix Pharmaceuticals, Inc. (Burlingame, CA). For example, 1 to 50 µg at one or more time points. As one example, treatment with subcutaneous ghrelin: 1 dose of 10 µg following mBI and additional dose 10 µg 1 hour after. Brain tissue is harvested at 2-8 hours, preferably about 6 hours post-injury. Reactive oxygen species (ROS) is measured by neutrophil oxidative burst in leukocytes isolated from brain tissue as shown in FIG. 1, which depicts the principle of assaying oxidative burst within inflammatory cells. Dihydrorhodamine 123 (DHR 123) diffuses across the cell membrane. When it encounters reactive oxygen species (ROS), DHR 123 is oxidized and fluoresces green. The fluorescence is then measured and presented as arbitrary fluorescence unit (AFU), wherein higher intensity means greater oxidative burst and, therefore greater concentration or amount of ROS. Brain cells are isolated (for example by collagenase Dispase/DNase and a Percoll gradient with resuspension of the pellet using 100 µL of HBSS). Oxidative burst, which is a marker for ROS, can be measured by flow cytometry (e.g., FL1 channel with 488-nm laser).

**[00216]** Oxidative burst is increased in animals following mBI compared to control. The number of animals was three in each group (n=3 animals in each group). Ghrelin treatment reduced the oxidative burst compared to untreated animal with mBI (FIG. 2). FIG. 2 shows that subcutaneous ghrelin reduces oxidative burst in inflammatory cells following mBI. The impact on mBI when ghrelin is administered can be quantified, for example, by arbitrary fluorescence unit (AFU) (FIG. 3). FIG. 3 depicts quantitation of oxidative burst of ipsilateral whole brain following mild BI. Oxidative burst increased in animals following mBI compared to control. Ghrelin treatment reduced the oxidative burst compared to untreated mBI (FIG. 2 and FIG. 3).

[00217] As quantified in FIG. 3, oxidative burst was increased in animals following mBI compared to sham control (7963 AFU  $\pm$  2900 (mild BI) vs. 4624 AFU  $\pm$  1858 (sham); n=5 animals in each group). Ghrelin treatment reduced the oxidative burst compared to mBI (7963 AFU  $\pm$  2900 (mild BI) vs. 3257 AFU  $\pm$  1031 (mild BI + ghrelin (2 doses))); p-value of 0.048 using Student's t-test).

#### EXAMPLE 2

[00218] The binding ability of ghrelin to GHS-R can be determined by a binding assay. Chinese hamster ovary cell line cells, CHO-K1, are prepared to express the human recombinant GHS receptor.

[00219] The cells can be prepared by any suitable method. One such method can include: The cDNA for human growth hormone secretagogue receptor (hGHS-R1a, or ghrelin receptor) is cloned by Polymerase Chain Reaction (PCR) using human brain RNA as a template (Clontech, Palo Alto, Calif.), gene specific primers flanking the full-length coding sequence of hGHS-R, (S: 5'-ATGTGGAACGCGACGCCAGCGAAGAG-3' (SEQ ID NO: 2) and AS: 5'-TCATGTATTAATACTAGATTCTGTCCA-3') (SEQ ID NO: 3), and Advantage 2 PCR Kit (Clontech). The PCR product is cloned into the pCR2.1 vector using Original TA Cloning Kit (Invitrogen, Carlsbad, Calif.). The full length human GHS-R is subcloned into the mammalian expression vector pcDNA 3.1 (Invitrogen). The plasmid is transfected into the Chinese hamster ovary cell line, CHO-K1 (American Type Culture Collection, Rockville, Md.), by calcium phosphate method (Wigler, M et al., Cell 11, 223, 1977). Single cell clones stably expressing the hGHS-R are obtained by selecting transfected cells grown in cloning rings in RPMI 1640 media supplemented with 10% fetal bovine serum and 1 mM sodium pyruvate containing 0.8 mg/ml G418 (Gibco, Grand Island, N.Y.).

[00220] GHS-R binding assay is performed by homogenizing the CHO-K1 cells expressing the human recombinant GHS receptor in 20 ml of ice-cold 50 mM Tris-HCl with a Brinkman Polytron (Westbury, N.Y.). The homogenates are washed twice by centrifugation (39,000 g/10 min), and the final pellets are resuspended in 50 mM Tris-HCl, containing 2.5 mM MgCl<sub>2</sub>, and 0.1% BSA. For the GHS-R binding assay, aliquots (0.4 ml) are incubated with 0.05 nM (<sup>125</sup>I)ghrelin

(~2000 Ci/mmol, Perkin Elmer Life Sciences, Boston, Mass.), with and without 0.05 ml of unlabeled competing test peptides. After a 60 min incubation (4° C.), the bound (<sup>125</sup>I)ghrelin is separated from the free by rapid filtration through GF/C filters (Brandel, Gaithersburg, Md.), which have been previously soaked in 0.5% polyethyleneimine/0.1% BSA. The filters then are washed three times with 5ml aliquots of 50 mM Tris-HCl and 0.1% bovine serum albumin, and the bound radioactivity trapped on the filters is counted by gamma spectrometry (Wallac LKB, Gaithersburg, Md.). Specific binding is defined as the total (<sup>125</sup>I)ghrelin bound minus that bound in the presence of 1000 nM ghrelin.

### Example 3

**[00221]** The functional activity of ghrelin is examined using GHS-R functional activity assays *in vitro* and *in vivo*. Ghrelin binding to GSH receptor can mediate intracellular iCa<sup>2+</sup> mobilization *in vitro*. Ghrelin may also be tested for ability to stimulate or suppress release of growth hormone (GH) *in vivo*.

**[00222]** Cells expressing human GSH receptor can be used. For example, CHO-K1 cells expressing the human GSH receptor are harvested by incubating in a 0.3% EDTA/phosphate buffered saline solution (25° C), and are washed twice by centrifugation. The washed cells are resuspended in Hank's-buffered saline solution (HBSS) for loading of the fluorescent Ca<sup>2+</sup> indicator Fura-2AM. Cell suspensions of approximately 10<sup>6</sup> cells/ml are incubated with 2 μM Fura-2AM for 30 min at about 25° C. Unloaded Fura-2AM is removed by centrifugation twice in HBBS, and the final suspensions are transferred to a spectrofluorometer (Hitachi F-2000) equipped with a magnetic stirring mechanism and a temperature-regulated cuvette holder. After equilibration to 37° C., ghrelin is added for measurement of intracellular Ca<sup>2+</sup> mobilization. The excitation and emission wavelengths can be, for example, 340 and 510 nm, respectively.

**[00223]** Ghrelin may be tested for its ability to stimulate or suppress release of growth hormone (GH) *in vivo* (Deghenghi, R., et al., *Life Sciences*, 1994, 54, 1321-1328; International Application No. WO 02/08250; each of which is incorporated herein by reference in its entirety). Thus, for example, in order to ascertain

ghrelin's ability to stimulate GH release *in vivo* the compound may be injected subcutaneously in 10-day old rats at a dose of, *e.g.*, 300 mg/kg. The circulating GH may be determined at, *e.g.*, 15 minutes after injection and compared to GH levels in rats injected with a solvent control.

**[00224]** It is to be understood that while the present disclosure has been described in conjunction with the above embodiments, that the foregoing description and examples are intended to illustrate and not limit the scope of the present disclosure. Other aspects, advantages and modifications within the scope of the present disclosure will be apparent to those skilled in the art to which the present disclosure pertains.

**[00225]** The present disclosure is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the present disclosure, and any compositions or methods, which are functionally equivalent are within the scope of this disclosure. It will be apparent to those skilled in the art that various modifications and variations can be made in the methods and compositions of the present disclosure without departing from the spirit or scope of the disclosure. Thus, it is intended that the present disclosure cover the modifications and variations of this disclosure provided they come within the scope of the appended claims and their equivalents.

**CLAIMS**

What is claimed is:

1. A method of treating Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.
2. A method of preventing the onset of Post-Traumatic Stress Disorder (PTSD) or delaying PTSD resulting from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of ghrelin, thereby preventing or delaying the onset of the PTSD, or treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.
3. A method of treating Post-Traumatic Stress Disorder (PTSD) in a subject, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition consisting essentially of isolated ghrelin.
4. The method of any one of claims 1-3, wherein the PTSD is associated with a single mild brain injury or concussion.
5. The method of any one of claims 1-3, wherein the PTSD is associated with repetitive or multiple mild brain injuries.
6. The method of any one of claims 1-5, wherein ghrelin is used as a sole active agent for treating the PTSD.
7. The method of any one of claims 1-6, wherein ghrelin is used in a combination with one or more therapeutic agents.
8. The method of any one of claims 1-3, wherein the subject is a mammal.

9. The method of any one of claims 1-3, wherein the ghrelin is administered in an amount that results in ghrelin blood levels being at least two times greater than the endogenous ghrelin blood levels of the subject.
10. The method of any one of claims 1-3, wherein ghrelin is administered in a single dose.
11. The method of any one of claims 1-3, wherein ghrelin is administered at a dosage from 10 ng/kg per day to 10 mg/kg per day.
12. The method of any one of claims 1-3, wherein ghrelin is administered at a dosage of 2 µg/kg per day.
13. A composition comprising a therapeutically effective amount of ghrelin for use in the treatment of Post-Traumatic Stress Disorder (PTSD) in a subject in need thereof, comprising administering to the subject the composition, thereby treating the PTSD or reducing the severity of the PTSD and/or the related symptoms.
14. A composition comprising a therapeutically effective amount of ghrelin for use in the treatment of preventing or delaying the onset of Post-Traumatic Stress Disorder (PTSD) or delaying the PTSD resulted from one or more of mild brain injuries in a subject in need thereof, comprising administering to the subject composition, thereby preventing the onset of the PTSD or treating the PTSD or reduce the severity of the PTSD and/or the related symptoms.
15. The composition of claim 13 or 14, wherein the PTSD is associated with a single mild brain injury or concussion.
16. The composition of claim 13 or 14, wherein the PTSD is associated with repetitive or multiple mild brain injuries.
17. The composition of any one of claims 13-16, wherein ghrelin is used as a sole active agent for treating the PTSD.

18. The composition of any one of claims 13-17, wherein ghrelin is used in a combination with one or more therapeutic agents.
19. The composition of claim 13-17, wherein when administered the ghrelin is the sole active agent being administered to the subject.
20. The composition of claim 13-19, wherein the ghrelin has the sequence of Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (SEQ ID NO: 1).
21. The composition of claim 20, wherein the ghrelin is modified with one or more fatty acids.
22. The composition of claim 21, wherein the fatty acid is an octanoic acid.
23. The composition of claim 20, wherein the ghrelin is modified at serine at amino acid position 2 and/or serine at amino acid position 3 of SEQ ID NO: 1.
24. The composition of claim 20, wherein the ghrelin is modified at serine at amino acid position 2 and/or serine amino acid position 3 with an octanoic acid.
25. The composition of claim 20, wherein modification of ghrelin is acylation of ghrelin at serine at amino acid position 2 and/or serine at amino acid position 3 with an octanoic acid.
26. The composition of claim 20, wherein the ghrelin is an octanoyl-ghrelin with an acylation at the serine at amino acid position 3 by the octanoic acid.




FIG. 1




FIG. 2




FIG. 3

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/018602

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 38/22 (2016.01)

CPC - A61K 38/22 (2016.02)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A61K 38/00, 38/16, 38/17, 38/22 (2016.01)

CPC - A61K 38/00, 38/16, 38/17, 38/22 (2016.02)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

U.S. Classes - 514/1, 1.1, 4.8, 5.3 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Orbit, Google Patents, PubMed, Google.

Search terms used: treat Post-Traumatic Stress Disorder administer ghrelin brain injury concussion

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                 | Relevant to claim No. |
|-----------|----------------------------------------------------------------------------------------------------|-----------------------|
| A         | US 2012/0302526 A1 (BARTH et al) 29 November 2012 (29.11.2012) entire document                     | 1-5, 8-16             |
| A         | US 2014/0255517 A1 (KERRY J. RESSLER) 11 September 2014 (11.09.2014) entire document               | 1-5, 8-16             |
| A         | WO 2009/071283 A2 (CHARITE - UNIVERSITÄTSMEDIZIN BERLIN) 11 June 2009 (11.06.2009) entire document | 1-5, 8-16             |
| A         | US 8,791,149 B2 (MCKNIGHT et al) 29 July 2014 (29.07.2014) entire document                         | 1-5, 8-16             |
| A         | US 8,324,151 B2 (WANG) 04 December 2012 (04.12.2012) entire document                               | 1-5, 8-16             |



Further documents are listed in the continuation of Box C.



See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

11 April 2016

Date of mailing of the international search report

02 MAY 2016

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/018602

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 6, 7, 17-26  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.