



(86) Date de dépôt PCT/PCT Filing Date: 2015/09/03
(87) Date publication PCT/PCT Publication Date: 2016/03/10
(85) Entrée phase nationale/National Entry: 2017/02/23
(86) N° demande PCT/PCT Application No.: US 2015/048316
(87) N° publication PCT/PCT Publication No.: 2016/036937
(30) Priorité/Priority: 2014/09/05 (US62/046,682)

(51) Cl.Int./Int.Cl. *C07K 16/28* (2006.01),
A61K 39/395 (2006.01), *A61P 35/00* (2006.01),
C07K 16/30 (2006.01), *C07K 16/46* (2006.01),
C12N 15/13 (2006.01), *C12N 5/10* (2006.01)

(71) Demandeur/Applicant:
JANSSEN PHARMACEUTICA NV, BE


(72) Inventeurs/Inventors:
GAUDET, FRANCOIS, US;
ATTAR, RICARDO, US;
HARMAN, BENJAMIN C., US;
LI, YINGZHE, US;
LUO, JINQUAN, US;
...

(74) Agent: NORTON ROSE FULBRIGHT CANADA

(54) Titre : AGENTS DE LIAISON CD123 ET LEURS UTILISATIONS


(54) Title: CD123 BINDING AGENTS AND USES THEREOF

Figure 1.



(57) Abrégé/Abstract:

Provided herein are antibodies that immunospecifically bind to CD123. Also described are related polynucleotides capable of encoding the provided CD123-specific antibodies or antigen-binding fragments, cells expressing the provided antibodies or antigen-binding fragments, as well as associated vectors and detectably labeled antibodies or antigen-binding fragments. In addition, methods of using the provided antibodies are described. For example, the provided antibodies may be used to diagnose, treat, or monitor CD123-expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with CD123-expressing cancer and thus may be amenable to treatment with a CD123-specific anti-cancer therapeutic, such as the multispecific antibodies against CD123 and CD3 described herein.



(72) **Inventeurs(suite)/Inventors(continued)**: MCDAID, RONAN, US; POMERANTZ, STEVEN C., US; TAM, SUSAN H., US; TEPLYAKOV, ALEXEY, US; WHEELER, JOHN, US; WU, SHENG-JIUN, US; NEMETH, JENNIFER F., US

(74) **Agent(suite/continued)**: LLP/S.E.N.C.R.L., S.R.L.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2016/036937 A1(43) International Publication Date
10 March 2016 (10.03.2016)

(51) International Patent Classification:

C07K 16/28 (2006.01) *A61K 39/395* (2006.01)
C07K 16/30 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:

PCT/US2015/048316

(22) International Filing Date:

3 September 2015 (03.09.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/046,682 5 September 2014 (05.09.2014) US

(71) Applicant: JANSSEN PHARMACEUTICA NV
[BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors: GAUDET, Francois; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). ATTAR, Ricardo; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). HARMAN, Benjamin C.; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). LI, Yingzhe; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). LUO, Jinquan; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). MCDAID, Ronan; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). POMERANTZ, Steven C.; 1400 McKean Road,

Spring House, Pennsylvania 19477 (US). TAM, Susan H.; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). TEPLYAKOV, Alexey; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). WHEELER, John; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). WU, Sheng-Jiun; 1400 McKean Road, Spring House, Pennsylvania 19477 (US). NEMETH, Jennifer F.; 1400 McKean Road, Spring House, Pennsylvania 19477 (US).

(74) Agents: PLANTZ, Bernard F. et al.; JOHNSON & JOHNSON, One Johnson & Johnson Plaza, New Brunswick, New Jersey 08933 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: CD123 BINDING AGENTS AND USES THEREOF

Figure 1.



Comments for pDisplay™:
 5325 nucleotides

(57) Abstract: Provided herein are antibodies that immunospecifically bind to CD123. Also described are related polynucleotides capable of encoding the provided CD123-specific antibodies or antigen-binding fragments, cells expressing the provided antibodies or antigen-binding fragments, as well as associated vectors and detectably labeled antibodies or antigen-binding fragments. In addition, methods of using the provided antibodies are described. For example, the provided antibodies may be used to diagnose, treat, or monitor CD123-expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with CD123-expressing cancer and thus may be amenable to treatment with a CD123-specific anti-cancer therapeutic, such as the multispecific antibodies against CD123 and CD3 described herein.

WO 2016/036937 A1



- GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

 - as to the identity of the inventor (Rule 4.17(i))
 - as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

 - with international search report (Art. 21(3))
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
 - with sequence listing part of description (Rule 5.2(a))

CD123 BINDING AGENTS AND USES THEREOF

TECHNICAL FIELD

[0001] The disclosure provided herein relates to monoclonal antibodies that immunospecifically bind cluster determinant 123 (CD123; also known as IL-3R α), multispecific antibodies that immunospecifically bind CD123 and cluster determinant 3 (CD3), and methods of producing and using the described antibodies.

BACKGROUND

[0002] Approximately every three minutes, a new diagnosis of a blood cancer is made. The most common blood cancers are leukemia, lymphoma and myeloma, which will account for 156,420 new people to be diagnosed in the United States in 2014. Approximately every 10 minutes, someone in the United States dies from a blood cancer. Blood cancers are diseases that can affect the bone marrow, the blood cells, the lymph nodes and other parts of the lymphatic system. These cancers disproportionately target young people, with leukemia being the most common type of cancer in children and adolescents younger than 20.

[0003] One type of blood cancer cell expresses a cell marker known as CD123 (IL-3R α). Examples of blood cancer cells that express CD123 include blasts and leukemia stem cells. Diseases associated with the expression of CD123 include acute myeloid leukemia (AML), myelodysplastic syndrome (MDS; low and high risk), acute lymphocytic leukemia (ALL, all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), and blastic plasmacytoid dendritic cell neoplasm (DPDCN).

[0004] Currently, treatments for these diseases include over 50 individual drugs with others under study and in clinical trials. Radiation therapy (RT) is also commonly used to treat blood cancers and sometimes it is administered along with drug therapy. Immunotherapy, gene therapy and personalized medicine are also used. However, these therapies can have significant side effects and adverse reactions. Thus, there is a need for new and improved treatments for CD123 (IL-3R α)-expressing blood cancers.

SUMMARY

[0005] Provided herein are antibodies that immunospecifically bind to CD123 and antigen-binding fragments thereof. Also described are related polynucleotides capable of encoding the provided CD123-specific antibodies and antigen-binding fragments, cells expressing the provided antibodies and antigen-binding fragments, as well as associated vectors and detectably labeled antibodies and antigen-binding fragments. In addition, methods of using the provided antibodies and antigen-binding fragments are described. For example, the CD123-specific antibodies and antigen-binding fragments may be used to diagnose or monitor CD123-expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with CD123-expressing cancer and thus may be amenable to treatment with a CD123-specific anti-cancer therapeutic, such as the multispecific antibodies against CD123 and CD3 described herein.

[0006] Further provided herein are multispecific antibodies that immunospecifically bind to CD123 and CD3 and multispecific antigen-binding fragments thereof. Also described are related polynucleotides capable of encoding the provided CD123 x CD3-multispecific antibodies, cells expressing the provided antibodies, as well as associated vectors and detectably labeled multispecific antibodies. In addition, methods of using the provided multispecific antibodies are described. For example, the CD123 x CD3-multispecific antibodies may be used to diagnose or monitor CD123-expressing cancer progression, regression, or stability; to determine whether or not a patient should be treated for cancer; or to determine whether or not a subject is afflicted with CD123-expressing cancer and thus may be amenable to treatment with a CD123-specific anti-cancer therapeutic, such as the CD123 x CD3-multispecific antibodies described herein.

CD123-Specific Antibodies

[0007] Described herein are isolated antibodies and antigen-binding fragments specific for CD123. In some embodiments, the CD123-specific antibodies and antigen-binding fragments bind human CD123 SP1 (SEQ ID NO: 1). In some embodiments, the CD123-specific antibodies and antigen-binding fragments bind human CD123 SP2 (SEQ ID NO: 2). In some embodiments, the CD123-specific antibodies and antigen-binding fragments bind human CD123 SP1 and SP2. In some embodiments, the CD123-specific antibodies and antigen-binding fragments bind human CD123 SP1 and cynomolgus monkey CD123 (SEQ ID NO: 3). In some embodiments, the CD123-specific antibodies and antigen-binding fragments bind to an epitope including one or more residues from (i) the segment of CD123 SP2 extracellular domain (ECD)

comprising residues 195 - 202 (RARERVYE (SEQ ID NO: 234)) and/or the segment of CD123 SP2 ECD comprising residues 156-161 (RKFRYE (SEQ ID NO:232)) and/or the segment of CD123 SP2 ECD comprising residues 173 – 178 (TEQVRD (SEQ ID NO: 233)) or (ii) the segment of CD123 SP2 ECD comprising residues 164-175 (IQKRMQPVITEQ (SEQ ID NO: 228)). and/or the segment of CD123 SP2 ECD comprising residues 184-189 (LLNPGT (SEQ ID NO: 229)). This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less.

[0008] In some embodiments, the CD123-specific antibody or antigen-binding fragment competes for binding to CD123 with a CD123-specific antibody or antigen-binding fragment that binds to an epitope including one or more residues from (i) the segment of CD123 SP2 ECD comprising residues 195-202 (RARERVYE (SEQ ID NO: 234)) or (ii) the segment of CD123 SP2 ECD comprising residues 164-175 (IQKRMQPVITEQ (SEQ ID NO: 228)). Antibodies or fragments binding to at least one residue in these epitopes may also bind to additional residues in the CD123 ECD including one or more residues from (i) the segment of CD123 SP2 ECD comprising residues 156-161 (RKFRYE (SEQ ID NO:232)) and/or the segment of CD123 SP2 ECD comprising residues 173 – 178 (TEQVRD (SEQ ID NO: 233)) or ii) one or more residues form the segment of CD123 SP2 ECD comprising residues 184-189 (LLNPGT (SEQ ID NO: 229)). This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less.

[0009] In some embodiments, the CD123-specific antibodies and antigen-binding fragments, such as those discussed in the preceding two paragraphs, are neutralizing antibodies. A neutralizing CD123-specific antibody or antigen-binding fragment includes those that are capable of inhibiting the binding of IL-3 to CD123 as determined by measuring the decrease in STAT5 phosphorylation upon stimulation of TF-1 cells with rhIL-3.

[0010] In some embodiments, the CD123-specific antibodies and antigen-binding fragments can prevent IL-3 binding to the CD123(IL3Ra)/CD131(IL3Rb) receptor. In other embodiments, the CD123-specific antibodies and antigen-binding fragments can prevent the association of the α and β chains of the of the IL3R receptor, (CD123(IL3Ra)/CD131(IL3Rb)). An antibody or antigen binding fragment includes those that are capable of inhibiting the binding of IL3 and/or capable of inhibiting heteromerization of CD123/CD133 as determined by measuring the decrease in association between CD123 and CD131 and measuring the loss of

heteromerization with increasing antibody concentration. Table 1 provides a summary of examples of some CD123-specific antibodies described herein:

Table 1. CDR sequences of mAbs generated from phage panning against human CD123 (SEQ ID NO:)

ID	H-CDR1	H-CDR2	H-CDR3	L-CDR1	L-CDR2	L-CDR3
I3RB1	DYGMG (6)	YTRGGGSKYYADSVKG (7)	HGGSPRNELDY (8)	KSQQSVLYSSNNKNYLA (9)	WASTRES (10)	QQYYSTPLT (11)
I3RB2	GYWMH (12)	AIRSDGSSKYYADSVKG (13)	DGVIEDTFDY (14)	RASQSVSSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB3	SYAMS (18)	GKYGGGSKYYADSVKG (19)	KDMSYFDY (20)	KSSQSVLYSSNNKNYLA (9)	WASTRES (10)	QQYYSTPLT (11)
I3RB4	GYGMS (21)	AISGSGGSTYYADSVKG (22)	GNWYYGLGFDY (23)	RASQSVSSSYLA (24)	GASSRAT (25)	QQYGSSPLT (26)
I3RB5	GYWML (27)	GINYDGSTYYADSVKG (28)	DHFLAEFDY (29)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB6	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GLFNWSNVALDY (35)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB7	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GKRWLADAGDFDY (36)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB8	SYAIS (33)	GIPIFGTANYAQKFQG (34)	HGFAWNDYSLDY (37)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB9	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GAPWNPENLDY (38)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB10	SYGIS (39)	WISAIFGNTNYAQKFQG (40)	GGLLYYASYLDY (41)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB11	SYGIS (39)	GIPIFGTANYAQKFQG (34)	DLESWRYNFDY (42)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB12	SYAIS (33)	GIPIFGTANYAQKFQG (34)	ADRVWDYYLDY (43)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB13	SYGIS (39)	GIPIFGTANYAQKFQG (44)	GGHPYVRLDY (45)	RASQSVSSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB14	SYGIS (39)	WISAIFGTTNYAQKFQG (46)	GGPLRYNHFDY (47)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB15	SYAIS (33)	GIPIFGTANYAQKFQG (34)	DLESLRYSTFDY (48)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB16	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GAVWGDQWFDY (49)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB17	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GAISLWYSTFDY (50)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB18	SYWIS (51)	IIDPSDSDTRYSPSFQG (52)	GDGSTDLDY (53)	RASQSVSSSYLA (24)	GASSRAT (25)	QQDYGFPT (54)
I3RB19	NYAMS (55)	GIRNGESTYYADSVKG (56)	GGPIGAPFDYLDY (57)	RASQSISSYLN (58)	AASSLQS (59)	QQSYSTPLT (60)
I3RB20	SYAIS (33)	GIPIFGTANYAQKFQG (34)	DDQIWGSYHLDY (60)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB21	SYAIS (33)	GIPIFGTANYAQKFQG (34)	ELWVGCKTFDY (61)	RASQSVANFLA (62)	DASNRAT (63)	QQTHWPT (64)
I3RB22	SYAIS (33)	GIPIFGTANYAQKFQG (34)	NLFYWADSVYLDY (65)	RASQSVNKLWA (66)	YASNRAT (67)	QQGIDWPRT (68)
I3RB23	SYGIS (39)	GIPIFGTANYAQKFQG (34)	EGSWMKNDRYVFDY (69)	RASQSISSYLN (30)	AASSLQS (31)	QQYDFPLT (70)
I3RB24	SYAIS (33)	GIPIFGTANYAQKFQG (34)	HTDAWGYRLDY (71)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB25	SYGIS (39)	GIPIFGTANYAQKFQG (72)	RPKWVESYFDY (73)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB26	SYGIS (39)	GIPIFGTANYAQKFQG (34)	NGFAWSVSGNLDY (74)	RASQSVDNWLA (75)	GASNRAT (76)	QQSISAPYT (77)

ID	H-CDR1	H-CDR2	H-CDR3	L-CDR1	L-CDR2	L-CDR3
I3RB27	SYAIS (33)	GIIPFGTANYAQKFQG (34)	ACWWMLRYGLDY (78)	RASQSVAKSLA (79)	AASNRAT (63)	QQFICWPIT (80)
I3RB28	SYAIS (33)	GIIPFGTANYAQKFQG (34)	APFTWDYSRLDY (81)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB29	SYAIS (33)	GIIPFGTANYAQKFQG (34)	DEKRWSESLDY (82)	RASQSIGEWLN (83)	AASSLQS (31)	QQYHFEPLT (84)
I3RB30	SYAIS (33)	WIIPFGTANYAQKFQG (85)	LVYSSDFDY (86)	RASQSVANWLA (87)	YASNRAT (67)	QQYDGWPRT (88)
I3RB31	SYAIS (33)	GISAYFTNANYAQKFQG (89)	SYEGDAYFDY (90)	RASQSVDKDLA (91)	GASNRAT (76)	QQYDRAPIT (92)
I3RB32	SYGIS (39)	GIIPFGTANYAQKFQG (34)	GAWWAYDTYLDY (93)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB33	SYGIS (39)	GIIPFGTANYAQKFQG (34)	GYWNNYDYLDY (94)	RASQSVNDWLA (95)	GASNRAT (76)	QQYKRAPYT (96)
I3RB34	SYAIS (33)	GIIPFGTANYAQKFQG (34)	GWSYYRLDY (97)	RASQSVDKWLA (98)	YASNRAT (67)	QQFDRAPFT (99)
I3RB35	SYAIS (33)	GIIPFGTANYAQKFQG (34)	HLFWLAGPLDY (100)	RASQSISSYLN (30)	AASSLQS (31)	QQYISPEYT (101)
I3RB36	SYGIS (39)	GIIPFGTANYAQKFQG (34)	DLHVWAYSNEFDY (102)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB37	SYAIS (33)	GIIPFGTANYAQKFQG (34)	DKTDFPSELDY (103)	RASQSIATWLN (104)	AASSLQS (31)	QQYITFEPLT (105)
I3RB38	SYGIS (39)	GIIPFGTANYAQKFQG (34)	DLMIWRFENFDY (106)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB39	SYAIS (33)	GIIPFGTANYAQKFQG (34)	EYGLDY (107)	RASQTVADFLA (108)	GASNRAT (109)	QQYNWPWT (110)
I3RB40	SYAIS (33)	GIIPFGTANYAQKFQG (34)	GQWWADTWFDY (111)	RASQSVAKWLA (112)	GASNRAT (76)	QQYHTAPWT (113)
I3RB41	SYAMS (114)	AISGSGGSTYYADSVKG (22)	VAYWEEFVYESLDY (115)	RASQSVSSSYLA (14)	GASNRAT (25)	QQYGSPLT (26)
I3RB42	SYAMS (114)	AISGSGGSTYYADSVKG (22)	HDWAFWIVFLDY (116)	RASQSVSSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB43	SYMMH (117)	AISGSGSKYYADSVKG (13)	DGIWMTFDY (118)	RASQSVSSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB44	SYWIS (51)	IIDPSDSDTRYSPSFQG (52)	GDGSTDLDY (53)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB47	SYAIS (33)	GIIPFGTANYAQKFQG (34)	HLFSWRYSEFDY (42)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)

[0011] In some embodiments are provided a CD123-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments are provided a CD123-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments described herein, the CD123-specific antibody or antigen-binding fragment thereof competes for binding to CD123 with an antibody or antigen-binding comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1.

[0012] The IgG class is divided in four isotypes: IgG1, IgG2, IgG3 and IgG4 in humans. They share more than 95% homology in the amino acid sequences of the Fc regions but show major differences in the amino acid composition and structure of the hinge region. The Fc region mediates effector functions, such as antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). In ADCC, the Fc region of an antibody binds to Fc receptors (FcγRs) on the surface of immune effector cells such as natural killers and macrophages, leading to the phagocytosis or lysis of the targeted cells. In CDC, the antibodies kill the targeted cells by triggering the complement cascade at the cell surface. The antibodies described herein include antibodies with the described features of the variable domains in combination with any of the IgG isotypes, including modified versions in which the Fc sequence has been modified to effect different effector functions.

[0013] For many applications of therapeutic antibodies, Fc-mediated effector functions are not part of the mechanism of action. These Fc-mediated effector functions can be detrimental and potentially pose a safety risk by causing off-mechanism toxicity. Modifying effector functions can be achieved by engineering the Fc regions to reduce their binding to FcγRs or the complement factors. The binding of IgG to the activating (FcγRI, FcγRIIa, FcγRIIIa and FcγRIIIb) and inhibitory (FcγRIIb) FcγRs or the first component of complement (C1q) depends on residues located in the hinge region and the CH2 domain. Mutations have been introduced in IgG1, IgG2 and IgG4 to reduce or silence Fc functionalities. The antibodies described herein may include these modifications.

[0014] In one embodiment, the antibody comprises an Fc region with one or more of the following properties: (a) reduced effector function when compared to the parent Fc; (b) reduced affinity to FcγRI, FcγRIIa, FcγRIIb, FcγRIIIb and/or FcγRIIIa, (c) reduced affinity to FcγRI (d) reduced affinity to FcγRIIa (e) reduced affinity to FcγRIIb, (f) reduced affinity to FcγRIIIb or (g) reduced affinity to FcγRIIIa.

[0015] In some embodiments, the antibodies or antigen-binding fragments are IgG, or derivatives thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the antibody has an IgG1 isotype, the antibody contains L234A, L235A, and/or K409R substitution(s) in its Fc region. In some embodiments wherein the antibody has an IgG4 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region. The antibodies described herein may include these modifications.

[0016] In some embodiments the described antibodies are capable of binding to CD123 with a dissociation constant of 5 nM or less as measured by surface plasmon resonance (SPR). In

some embodiments, the antibodies comprise the CDRs of the antibodies presented in Table 1 above. Assays for measuring affinity by SPR include assays performed using a BIAcore 3000 or Biacore T200 machine, where the assay is performed at room temperature (*e.g.* at or near 25°C), wherein the antibody capable of binding to CD123 is captured on the BIAcore sensor chip by an anti-Fc antibody (*e.g.* goat anti-human IgG Fc specific antibody Jackson ImmunoResearch laboratories Prod # 109-005-098) to a level around 75RUs, followed by the collection of association and dissociation data at a flow rate of 40µl/min.

[0017] In addition to the described CD123-specific antibodies and antigen-binding fragments, also provided are polynucleotide sequences capable of encoding the described antibodies and antigen-binding fragments. Vectors comprising the described polynucleotides are also provided, as are cells expressing the CD123-specific antibodies or antigen-binding fragments provided herein. Also described are cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as 293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells (such as *E. coli*). The described antibodies may also be produced by hybridoma cells.

Methods of using CD123-Specific Antibodies

[0018] Methods of using the described CD123-specific antibodies or antigen-binding fragments are also disclosed. Particular antibodies for use in the methods discussed in this section include those with the set of CDRs described for antibodies in Table 1 above or antibodies that compete for binding to CD123 with one of the antibodies in Table 1. For example, these antibodies or antigen-binding fragments may be useful in treating cancer, by inhibiting a biological effect of IL-3 by preventing IL-3 from binding to IL-3R or where the antibody is conjugated to a toxin, so targeting the toxin to the CD123-expressing cancer. Further, these antibodies or antigen-binding fragments may be useful for detecting the presence of CD123 in a biological sample, such as blood or serum; for quantifying the amount of CD123 in a biological sample, such as blood or serum; for diagnosing CD123-expressing cancer; determining a method of treating a subject afflicted with cancer; or monitoring the progression of CD123-expressing cancer in a subject. In some embodiments, CD123-expressing cancer may be a hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN). The described methods may be carried out before the subject

receives treatment for CD123-expressing cancer, such as treatment with a multispecific antibody against CD123 and CD3. Furthermore, the described methods may be carried out after the subject receives treatment for CD123-expressing cancer, such as treatment with a multispecific antibody against CD123 and CD3 described herein.

[0019] The described methods of detecting CD123 in a biological sample include exposing the biological sample to one or more of the CD123-specific antibodies or antigen-binding fragments described herein.

[0020] The described methods of diagnosing CD123-expressing cancer in a subject also involve exposing the biological sample to one or more of the CD123-specific antibodies or antigen-binding fragments described herein; however, the methods also include quantifying the amount of CD123 present in the sample; comparing the amount of CD123 present in the sample to a known standard or reference sample; and determining whether the subject's CD123 levels fall within the levels of CD123 associated with cancer.

[0021] Also described herein are methods of monitoring CD123-expressing cancer in a subject. The described methods include exposing the biological sample to one or more of the CD123-specific antibodies or antigen-binding fragments described herein; quantifying the amount of CD123 present in the sample that is bound by the antibody, or antigen-binding fragment thereof; comparing the amount of CD123 present in the sample to either a known standard or reference sample or the amount of CD123 in a similar sample previously obtained from the subject; and determining whether the subject's CD123 levels are indicative of cancer progression, regression or stable disease based on the difference in the amount of CD123 in the compared samples.

[0022] The samples obtained, or derived from, subjects are biological samples such as urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated, tissues, surgically resected tumor tissue, biopsies, fine needle aspiration samples, or histological preparations.

[0023] The described CD123-specific antibodies or antigen-binding fragments may be labeled for use with the described methods, or other methods known to those skilled in the art. For example, the antibodies described herein, or antigen-binding fragments thereof, may be labeled with a radiolabel, a fluorescent label, an epitope tag, biotin, a chromophore label, an ECL label, an enzyme, ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, or poly-histidine or similar such labels known in the art.

CD123-Specific Antibody Kits

[0024] Described herein are kits including the disclosed CD123-specific antibodies or antigen-binding fragments thereof. The described kits may be used to carry out the methods of using the CD123-specific antibodies or antigen-binding fragments provided herein, or other methods known to those skilled in the art. In some embodiments the described kits may include the antibodies or antigen-binding fragments described herein and reagents for use in detecting the presence of CD123 in a biological sample. Accordingly, the described kits may include one or more of the antibodies, or an antigen-binding fragment(s) thereof, described herein and a vessel for containing the antibody or fragment when not in use, instructions for use of the antibody or fragment, the antibody or fragment affixed to a solid support, and/or detectably labeled forms of the antibody or fragment, as described herein.

CD123 x CD3-Multispecific Antibodies

[0025] Described herein are isolated multispecific antibodies that bind CD123 and CD3 (“CD123 x CD3 multispecific antibodies”) and multispecific antigen-binding fragments thereof. In some embodiments an isolated antibody, or an antigen-binding fragment thereof, that binds immunospecifically to CD123 SP2 (IL3-R α) and CD123 SP1 (IL3-R α) is provided.

[0026] In some embodiments, the CD123-specific arm of the multispecific antibody binds human CD123 and/or cynomolgus monkey CD123. In some embodiments, the CD123-specific arm of the CD123 x CD3-multispecific antibodies or antigen-binding fragments binds the SP1 and/or SP2 fragment of human CD123. In preferred embodiments, the CD123 x CD3 multispecific antibody or antigen-binding fragment is a bispecific antibody or antigen-binding fragment. In some embodiments, an isolated CD123 (IL3-R α) x CD3 bispecific antibody comprising: a) a first heavy chain (HC1); b) a second heavy chain (HC2); c) a first light chain (LC1); and d) a second light chain (LC2), wherein the HC1 and the LC1 pair to form a first antigen-binding site that immunospecifically binds CD123 (IL3-R α), and the HC2 and the LC2 pair to form a second antigen-binding site that immunospecifically binds CD3, or a CD123 (IL3-R α) x CD3-bispecific binding fragment thereof is provided. In another embodiment, an isolated cell expressing the antibody or bispecific binding fragment is provided. In some embodiments, the CD123-binding arm (or “CD123-specific arm”) of the CD123 x CD3 multispecific antibody is derived from a CD123 antibody described herein (for example, from an antibody having the CDR sequences listed in Table 1).

[0027] In some embodiments, the CD123-specific arm of the CD123 x CD3-multispecific antibodies or antigen-binding fragments are IgG, or derivatives thereof. In some embodiments the described CD123 x CD3-multispecific antibodies are capable of binding to CD123 with a dissociation constant of 5 nM or less as measured by surface plasmon resonance, or MSD-CAT.

[0028] In some embodiments, the CD3-binding arm (or “CD3-specific arm”) of the CD123 x CD3 multispecific antibody is derived from the mouse monoclonal antibody SP34, a mouse IgG3/lambda isotype. (Pessano, S., et al , 1995. EMBO J. 4, 337-344). In some embodiments, the CD3-binding arm of the CD123 x CD3 multispecific antibody comprises one VH domain and one VL domain selected from Table 2. Table 2 provides a summary of examples of some the heavy chains and light chains of the CD3-specific antibodies and antigen-binding fragments.

Table 2. Heavy chains and light chains of the CD3-specific antibodies and antigen-binding fragments.

VH	VL
CD3H141 (SEQ ID NO:184): IGHV3-72*01 with mouse CDRs+ Gly49Ala EVQLVESGGGLVQPGGSLRLSCAASGFTF NTYAMNWVRQAPGKGLEWVARIRSKYNNY ATYYAASVKGRFTISRDDSKNSLYLQMNS LKTEDTAVYYCARHGNFGNSYVSWFAYWG QGTLVTVSS	CD3L63 (SEQ ID NO:188): IGLV7-46*01 with mouse CDRs + F38V,A48G,Y51G,W59G QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTS NYANWVQQKPGQAPRGLIGGTNKRAPGTPARF SGSLLGKKAALTLSGAQPEDEAEYYCALWYSN LWVFGGGTKLTVL
CD3H142 (SEQ ID NO:185): IGHV3-23*01 with mouse CDRs+ Ser49Ala EVQLLES GGGLVQPGGSLRLSCAASGFTF NTYAMNWVRQAPGKGLEWVARIRSKYNNY ATYYADSVKGRFTISRDN SKNTLYLQMNS LRAEDTAVYYCAKHGNFGNSYVSWFAYWG QGTLVTVSS	CD3L64 (SEQ ID NO:189): IGLV1-51*01 with mouse CDRs + Y38V, L48G, Y51G QSVLTQPPSVSAAPGQKVTISCRSSTGAVTTS NYANWVQQLPGTAPKGLIGGTNKRAPGIPDRF SGSKSGTSATLGITGLQTGDEADYYCALWYSN LWVFGGGTKLTVL
CD3H143 (SEQ ID NO:186): IGHV3-23*01 with mouse CDRs+ Ser49Ala, Ala99Val EVQLLES GGGLVQPGGSLRLSCAASGFTF	CD3L66 (SEQ ID NO:190): IGLV7-43*01 with mouse CDRs + F38V,A48G,Y51G,W59G QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTS NYANWVQQKPGQAPRGLIGGTNKRAPGTPARF

NTYAMNWVRQAPGKGLEWVARIRSKYNNY ATYYADSVKGRFTISRDN SKNTLYLQMNS LRAEDTAVYYCVKHG NFGNSYVSWFAYWG QGTLVTVSS	SGSLLGGKAAL T LSGVQPEDEAEYYCALWYSN LWVFGGG TKLTVL
CD3H144(SEQ ID NO:187): IGHV3-73*01 with mouse CDRs + Asn57Gly EVQLVESGGGLVQP GGSLKLSCAA SGFTFNTYAMNWVRQASGKGLEWVGRI RS KYN GYATYYAASVKGRFTISRDDSKNTAY LQMNSLKTEDTAVYYCTR HGNFGNSYVSW FAYWGQGTLVTVSS	

[0029] In some embodiments, the CD3-specific antibodies and antigen-binding fragments comprise a heavy chain from Table 3 and a light chain from Table 3. Table 3 provides a summary of the matrix of the heavy chains and light chains of the CD3-specific antibodies and antigen-binding fragments.

Table 3. The antibodies created by combining the heavy and light chains.

Heavy chain	Light chain		
	CD3L63	CD3L64	CD3L66
CD3H141	CD3B143	CD3B144	CD3B146
CD3H142	CD3B147	CD3B148	CDB150
CD3H143	CD3B151	CD3B152	CD3B154
CD3H144	CD3B155	CD3B156	CD3B158

[0030] The IgG class is divided in four isotypes: IgG1, IgG2, IgG3 and IgG4 in humans. They share more than 95% homology in the amino acid sequences of the Fc regions but show major differences in the amino acid composition and structure of the hinge region. The Fc

region mediates effector functions, such as antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). In ADCC, the Fc region of an antibody binds to Fc receptors (FcγRs) on the surface of immune effector cells such as natural killers and macrophages, leading to the phagocytosis or lysis of the targeted cells. In CDC, the antibodies kill the targeted cells by triggering the complement cascade at the cell surface.

[0031] For many applications of therapeutic antibodies, Fc-mediated effector functions are not part of the mechanism of action. These Fc-mediated effector functions can be detrimental and potentially pose a safety risk by causing off-mechanism toxicity. Modifying effector functions can be achieved by engineering the Fc regions to reduce their binding to FcγRs or the complement factors. The binding of IgG to the activating (FcγRI, FcγRIIa, FcγRIIIa and FcγRIIIb) and inhibitory (FcγRIIb) FcγRs or the first component of complement (C1q) depends on residues located in the hinge region and the CH2 domain. Mutations have been introduced in IgG1, IgG2 and IgG4 to reduce or silence Fc functionalities. Silencing mutations can include, but are not limited to IgG1 AA (F234A, L235A), or IgG4 PAA (S228P, F234A, L235A), or IgG2 AA (V234A, G237A), or IgG1 FEA (L234F, L235E, D265A), or IgG1 FES (L234F/L235E/P331S).

[0032] In one embodiment, the antibody comprises an Fc region with one or more of the following properties: (a) reduced effector function when compared to the parent Fc; (b) reduced affinity to FcγRI, FcγRIIa, FcγRIIb, FcγRIIIb and/or FcγRIIIa, (c) reduced affinity to FcγRI (d) reduced affinity to FcγRIIa (e) reduced affinity to FcγRIIb, (f) reduced affinity to FcγRIIIb or (g) reduced affinity to FcγRIIIa.

[0033] In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG, or a derivative thereof. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG1, or a derivative thereof. In some embodiments, for example, the Fc region of the CD3-specific IgG1 antibody from which the CD3-binding arm is derived comprises L234A, L235A, and F405L substitutions in its Fc region. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG4, or a derivative thereof. In some embodiments, for example, the Fc region of the CD3-specific IgG4 antibody from which the CD3-binding arm is derived comprises S228P, L234A, L235A, F405L, and R409K substitutions in its Fc region. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is

derived is IgG-AA Fc. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived is IgG-AA Fc-L234A, L235A, and F405L (where L234A, L235A, and F405L are mutations). In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived binds CD3 ϵ on primary human T cells and/or primary cynomolgus T cells. In some embodiments, the CD3-specific antibody or antigen-binding fragment from which the CD3-specific arm of the multispecific antibody is derived activates primary human CD4⁺ T cells and/or primary cynomolgus CD4⁺ T cells. In some embodiments, the described CD123 x CD3 multispecific antibodies are capable of binding to CD3 on human or cynomolgous monkey T-cells with a dissociation constant of less than 500, or less than 100 or less than 20 nM as determined by competition binding with a labeled anti-CD3 antibody with known affinity

[0034] In addition to the described CD123 x CD3-multispecific antibodies, also provided are polynucleotide sequences capable of encoding the described CD123 x CD3-multispecific antibodies. In some embodiments, an isolated synthetic polynucleotide encoding the HC1, the HC2, the LC1 or the LC2 of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment is provided. Vectors comprising the described polynucleotides are also provided, as are cells expressing the CD123 x CD3-multispecific antibodies provided herein. Also described are cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as 293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells (such as E. coli). The described antibodies may also be produced by hybridoma cells. In some embodiments, methods for generating the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment by culturing cells is provided.

[0035] Further provided herein are pharmaceutical compositions comprising the CD123 (IL3-R α) x CD3 multispecific antibodies or antigen-binding fragments and a pharmaceutically acceptable carrier.

Methods of using CD123 x CD3-Multispecific Antibodies

[0036] Methods of using the described CD123 x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof are also disclosed. For example, the CD123 x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof may be useful in the treatment of a CD123-expressing cancer in a subject in need thereof. In some embodiments, the CD123-expressing cancer is a hematological cancer, such as acute myeloid

leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN).

[0037] The described methods of treating CD123-expressing cancer in a subject in need thereof include administering to the subject a therapeutically effective amount of a described CD123 x CD3-multispecific antibody or multispecific antigen-binding fragment thereof. In some embodiments, the subject is a mammal, preferably a human. In preferred embodiments are provided methods for treating a subject having cancer by administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific antigen-binding fragment to a patient in need thereof for a time sufficient to treat the cancer.

[0038] Further provided herein are methods for inhibiting growth or proliferation of cancer cells by administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment to inhibit the growth or proliferation of cancer cells.

[0039] Also provided herein are methods of redirecting a T cell to a CD123-expressing cancer cell by administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment to redirect a T cell to a cancer.

CD123 x CD3-Specific Antibody Kits

[0040] Described herein are kits including the disclosed CD123 x CD3-multispecific antibodies. The described kits may be used to carry out the methods of using the CD123 x CD3-multispecific antibodies provided herein, or other methods known to those skilled in the art. In some embodiments the described kits may include the antibodies described herein and reagents for use in treating a CD123-expressing cancer. Accordingly, the described kits may include one or more of the multispecific antibodies, or a multispecific antigen-binding fragment(s) thereof, described herein and a vessel for containing the antibody or fragment when not in use, and/or instructions for use of the antibody or fragment, the antibody or fragment affixed to a solid support, and/or detectably labeled forms of the antibody or fragment, as described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0041] Figure 1. Figure 1 shows the pDisplay vector used for cloning CD123 extracellular domains.

[0042] Figure 2. Figure 2 (Figure 2A, Figure 2B, and Figure 2C) shows a cell binding assay that demonstrated the binding potential of phage panel positive binders to CD123 expressing cells.

[0043] Figure 3. Figure 3 shows a competition ELISA between the antibody panel and the anti-CD123 antibody 7G3.

[0044] Figure 4. Figure 4 (Figure 4A and Figure 4B) shows a CD123 cell-based STAT5 functional assay. Figure 4C shows a dose-dependence CD123 cell-based STAT5 functional assay for I3RB18 and 7G3 antibodies.

[0045] Figure 5. Figure 5 (Figure 5A, Figure 5B, and Figure 5C) shows the binding of Mabs I3RB2, I3RB18, and 7G3 to endogenous CD123 expressed on AML cell line, OCI-AML5.

[0046] Figure 6. Figure 6 (Figure 6A and Figure 6B) shows a competitive binding assay between labeled I3RB2 and I3RB18 mAbs and other anti-CD123 Abs identified in the screen.

[0047] Figure 7. Figure 7 (Figure 7A (SEQ ID NO: 232) and Figure 7B (SEQ ID NO:232)) shows the results of epitope mapping studies by hydrogen/deuterium exchange-mass spectrometry (HDX-MS) showing differences in deuterium levels for CD123 SP2 in the presence or absence of Fab.

[0048] Figure 8. Figure 8 shows the Antibody residues involved in binding of CD123 sp2 observed in the cocrystal structure of the I3RB18 derived scFv and CD123 SP2 ECD. Numbering: CD123 sp2 in ovals; CDRs of I3RB18 in squares.

[0049] Figure 9. Figure 9A shows the co-crystal structure of CD123 sp2:I3RB18 (labeled B18) and Figure 9B shows the cocrystal structure of CD123 sp1:CSL362 Fab, a humanized form of mAb 7G3 from PDB entry 4JZJ.

[0050] Figure 10. Figure 10 shows the amino acid sequence of SP34 with sequential numbering. CDRs in AbM definition (K.R. Abhinandan and A. C. Martin, 2008. Mol. Immunol. 45, 3832-3839) are underlined. Ser230 is the last HC residue present in papain-cleaved Fab. Residues 231-455 are from IGHG3_MOUSE (mouse IgG3, isoform 2).

[0051] Figure 11. Figure 11 shows the variable domain of SP34 with key residues at VL/VH interface shown. Residues 38, 48, and 51 in VL (labeled) are in contact with CDR-H3.

[0052] Figure 12. Figure 12 shows the Human Framework Adaptation (“HFA”) variants for V_H (SEQ ID NOS 5 and 184-187, respectively, in order of appearance) and V_L (SEQ ID NOS 4 and 188-190, respectively, in order of appearance). The numbering is sequential;

CDRs in the AbM definition are underlined; residues that differ from SP34 are highlighted in bold; back mutations in HFA variants are bold and underlined.

[0053] Figure 13. Figure 13 shows binding of SP34 HFA variants to primary Human T cells.

[0054] Figure 14. Figure 14 shows binding of SP34 HFA variants to Cynomolgus primary T cells.

[0055] Figure 15. Figure 15 shows that SP34 HFA variants activate primary human T cells *in vitro*. Negative controls are shown in white and positive controls are shown in black.

[0056] Figure 16. Figure 16 shows that SP34 HFA variants activate primary cynomolgus T cells *in vitro*. Negative controls are shown in white and positive controls are shown in black.

[0057] Figure 17. Figure 17 shows the correlation of binding and activation by SP34 HFA variants. Average binding and CD69 Mean Fluorescence Intensity (“MFI”) values for human (Figure 17A) and cynomolgus (Figure 17B) were plotted against each other.

[0058] Figure 18. Figure 18 shows a T-cell mediated cytotoxicity assay for donor M6587 (Figure 18A) and donor M7020 (Figure 18B) with the MV4-11 cell line.

[0059] Figure 19. Figure 19 shows a T-cell mediated cytotoxicity assay for donor M6587 (Figure 19A) and donor M7020 (Figure 19B) with the OCI-M2 cell line.

[0060] Figure 20. Figure 20 shows a T-cell mediated cytotoxicity assay for donor M6587 (Figure 20A) and donor M7020 (Figure 20B) with the OCI-AML cell line.

[0061] Figure 21. Figure 21 shows the efficacy of I3RB186 in the KG-1 tumor xenograft model.

[0062] Figure 22. Figure 22 shows the efficacy of I3RB186 in the KG-1 tumor xenograft model by fluorescence-activated cell sorting (FACS) analysis of peripheral blood on day 30 at CD45+ (Figure 22A) and CD8+/CD4+ (Figure 22B).

[0063] Figure 23. Figure 23 shows the efficacy of I3RB186 in KG-1 tumor xenograft model by FACS analysis of peripheral blood on day 53 post tumor implantation at CD45+ (Figure 23A) and CD8+/CD4+ (Figure 23B).

[0064] Figure 24. Figure 24 shows the efficacy of I3RB186 in KG-1 tumor xenograft model by showing body weight change with treatment.

[0065] Figure 25. Figure 25 shows the efficacy of CD123 x CD3 bispecific Ab I3RB186 with control null arm bispecific Abs I3RB191 and I3RB192 in the KG-1 tumor xenograft model.

[0066] Figure 26. Figure 26 shows the efficacy of CD123 x CD3 bispecific Ab I3RB186 with control null arm bispecific Abs I3RB191 and I3RB192 in the KG-1 tumor xenograft model by FACS analysis on day 36 post tumor implantation at CD45+ (Figure 26A) and CD8+/CD4+ (Figure 26B).

[0067] Figure 27. Figure 27 shows the efficacy of CD123 x CD3 bispecific Ab I3RB186 with control null arm bispecific Abs I3RB191 and I3RB192 in the KG-1 tumor xenograft model by FACS analysis on day 63 post tumor implantation at CD45+ (Figure 27A) and CD8+/CD4+ (Figure 27B).

[0068] Figure 28. Figure 28 shows the efficacy of CD123 x CD3 bispecific Ab I3RB186 with control null arm bispecific Abs I3RB191 and I3RB192 in the KG-1 tumor xenograft model by showing body weight change with treatment.

[0069] Figure 29. Figure 29 shows saturation binding curves used to determine the cell binding affinity (Kd) for SP34-2 on primary human T cells (Figure 29A) and cynomolgus monkey T cells (Figure 29B).

[0070] Figure 30. Figure 30 shows competition binding experiments on primary human T cells (Figure 30A) and cynomolgus monkey T cells (Figure 30B) using labelled antibody. Alexa Fluor^R 488B146, and increasing concentrations of unlabeled CD123 x CD3 antibodies.

[0071] Figure 31. Figure 31 shows T-cell mediated cytotoxicity assay for donor M6948 (Figure 31A) and donor M6521 (Figure 31B) with the OCI-AML cell line.

[0072] Figure 32. Figure 32 shows T-cell mediated cytotoxicity assay for donor M6948 (Figure 32A) and donor M6521 (Figure 32B) with the KG-1 cell line.

[0073] Figure 33. Figure 33 shows T-cell mediated cytotoxicity assay for donor M6948 (Figure 33A) and donor M6521 (Figure 33B) with the JIM3 cell line.

[0074] Figure 34. Figure 34 A, B, C and D shows the effect of CD123 x CD3 antibodies on the IL-3 induced heteromerization of CD123 and CD131 for I3RB218 (Figure 34A), 8747 (Figure 34B), I3RB217 (Figure 34C) and 7959 (Figure 34D)

[0075] Figure 35. Figure 35 shows the efficacy of CD123 x CD3 Ab 7959, and Ab 9958 in the KG-1 tumor xenograft model by comparison of mean tumor volume.

[0076] Figure 36. Figure 36 shows the efficacy of CD123 x CD3 Ab 3978 in the KG-1 tumor xenograft model by comparison of mean tumor volume.

[0077] Figure 37. Figure 37 shows the efficacy of CD123 x CD3 Ab 8747 in the KG-1 tumor xenograft model by comparison of mean tumor volume.

[0078] Figure 38. Figure 38 shows the efficacy of CD123 x CD3 Ab 8876 in the KG-1 tumor xenograft model by comparison of mean tumor volume.

[0079] Figure 39. Figure 39 shows the efficacy of CD123 x CD3 Ab 7959 and Ab 9958 in the KG-1 tumor xenograft model by comparison of body weight change with treatment.

[0080] Figure 40. Figure 40 shows the efficacy of CD123 x CD3 Ab 3978 in the KG-1 tumor xenograft model by comparison of body weight change with treatment.

[0081] Figure 41. Figure 41 shows the efficacy of CD123 x CD3 Ab 8747 in the KG-1 tumor xenograft model by comparison of body weight change with treatment.

[0082] Figure 42. Figure 42 shows the efficacy of CD123 x CD3 Ab 8876 in the KG-1 tumor xenograft model by comparison of body weight change with treatment.

[0083] Figure 43. Figure 43 shows the in-vivo mouse PK of CD123 x CD3 bispecific antibodies 3978, 7955, 7959, 9958

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Definitions

[0084] Various terms relating to aspects of the description are used throughout the specification and claims. Such terms are to be given their ordinary meaning in the art unless otherwise indicated. Other specifically defined terms are to be construed in a manner consistent with the definitions provided herein.

[0085] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a cell” includes a combination of two or more cells, and the like.

[0086] The term “about” as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of up to $\pm 10\%$ from the specified value, as such variations are appropriate to perform the disclosed methods. Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the following specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

[0087] Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0088] “Isolated” means a biological component (such as a nucleic acid, peptide or protein) has been substantially separated, produced apart from, or purified away from other biological components of the organism in which the component naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids, peptides and proteins that have been “isolated” thus include nucleic acids and proteins purified by standard purification methods. “Isolated” nucleic acids, peptides and proteins can be part of a composition and still be isolated if such composition is not part of the native environment of the nucleic acid, peptide, or protein. The term also embraces nucleic acids, peptides and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids. An “isolated” antibody or antigen-binding fragment, as used herein, is intended to refer to an antibody or antigen-binding fragment which is substantially free of other antibodies or antigen-binding fragments having different antigenic specificities (for instance, an isolated antibody that specifically binds to CD123 is substantially free of antibodies that specifically bind antigens other than CD123). An isolated antibody that specifically binds to an epitope, isoform or variant of CD123 may, however, have cross-reactivity to other related antigens, for instance from other species (such as CD123 species homologs).

[0089] “Polynucleotide,” synonymously referred to as “nucleic acid molecule,” “nucleotides” or “nucleic acids,” refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, “polynucleotide” embraces

chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

[0090] The meaning of “substantially the same” can differ depending on the context in which the term is used. Because of the natural sequence variation likely to exist among heavy and light chains and the genes encoding them, one would expect to find some level of variation within the amino acid sequences or the genes encoding the antibodies or antigen-binding fragments described herein, with little or no impact on their unique binding properties (e.g., specificity and affinity). Such an expectation is due in part to the degeneracy of the genetic code, as well as to the evolutionary success of conservative amino acid sequence variations, which do not appreciably alter the nature of the encoded protein. Accordingly, in the context of nucleic acid sequences, “substantially the same” means at least 65% identity between two or more sequences. Preferably, the term refers to at least 70% identity between two or more sequences, more preferably at least 75% identity, more preferably at least 80% identity, more preferably at least 85% identity, more preferably at least 90% identity, more preferably at least 91% identity, more preferably at least 92% identity, more preferably at least 93% identity, more preferably at least 94% identity, more preferably at least 95% identity, more preferably at least 96% identity, more preferably at least 97% identity, more preferably at least 98% identity, and more preferably at least 99% or greater identity. The percent identity between two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The percent identity between two nucleotide or amino acid sequences may e.g. be determined using the algorithm of E. Meyers and W. Miller, *Comput. Appl. Biosci* 4, 11-17 (1988) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences may be determined using the Needleman and Wunsch, *J. Mol. Biol.* 48, 444-453 (1970) algorithm.

[0091] The degree of variation that may occur within the amino acid sequence of a protein without having a substantial effect on protein function is much lower than that of a nucleic acid sequence, since the same degeneracy principles do not apply to amino acid sequences. Accordingly, in the context of an antibody or antigen-binding fragment,

“substantially the same” means antibodies or antigen-binding fragments having 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to the antibodies or antigen-binding fragments described. Other embodiments include CD123 specific antibodies, or antigen-binding fragments, that have framework, scaffold, or other non-binding regions that do not share significant identity with the antibodies and antigen-binding fragments described herein, but do incorporate one or more CDRs or other sequences needed to confer binding that are 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to such sequences described herein. A “vector” is a replicon, such as plasmid, phage, cosmid, or virus in which another nucleic acid segment may be operably inserted so as to bring about the replication or expression of the segment.

[0092] A “clone” is a population of cells derived from a single cell or common ancestor by mitosis. A “cell line” is a clone of a primary cell that is capable of stable growth in vitro for many generations. In some examples provided herein, cells are transformed by transfecting the cells with DNA.

[0093] The terms “express” and “produce” are used synonymously herein, and refer to the biosynthesis of a gene product. These terms encompass the transcription of a gene into RNA. These terms also encompass translation of RNA into one or more polypeptides, and further encompass all naturally occurring post-transcriptional and post-translational modifications. The expression or production of an antibody or antigen-binding fragment thereof may be within the cytoplasm of the cell, or into the extracellular milieu such as the growth medium of a cell culture.

[0094] The terms “treating” or “treatment” refer to any success or indicia of success in the attenuation or amelioration of an injury, pathology or condition, including any objective or subjective parameter such as abatement, remission, diminishing of symptoms or making the condition more tolerable to the patient, slowing in the rate of degeneration or decline, making the final point of degeneration less debilitating, improving a subject’s physical or mental well-being, or prolonging the length of survival. The treatment may be assessed by objective or subjective parameters; including the results of a physical examination, neurological examination, or psychiatric evaluations.

[0095] An “effective amount” or “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result. A therapeutically effective amount of a CD123 x CD3 antibody may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody to elicit a desired response in the individual. A therapeutically effective amount is also one in which any

toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

[0096] “Antibody” refers to all isotypes of immunoglobulins (IgG, IgA, IgE, IgM, IgD, and IgY) including various monomeric, polymeric and chimeric forms, unless otherwise specified. Specifically encompassed by the term “antibody” are polyclonal antibodies, monoclonal antibodies (mAbs), and antibody-like polypeptides, such as chimeric antibodies and humanized antibodies.

[0097] Antigen-binding fragments are any proteinaceous structure that may exhibit binding affinity for a particular antigen. Antigen-binding fragments include those provided by any known technique, such as enzymatic cleavage, peptide synthesis, and recombinant techniques. Some antigen-binding fragments are composed of portions of intact antibodies that retain antigen-binding specificity of the parent antibody molecule. For example, antigen-binding fragments may comprise at least one variable region (either a heavy chain or light chain variable region) or one or more CDRs of an antibody known to bind a particular antigen. Examples of suitable antigen-binding fragments include, without limitation diabodies and single-chain molecules as well as Fab, F(ab')₂, Fc, Fabc, and Fv molecules, single chain (Sc) antibodies, individual antibody light chains, individual antibody heavy chains, chimeric fusions between antibody chains or CDRs and other proteins, protein scaffolds, heavy chain monomers or dimers, light chain monomers or dimers, dimers consisting of one heavy and one light chain, a monovalent fragment consisting of the VL, VH, CL and CH1 domains, or a monovalent antibody as described in WO2007059782, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region, a Fd fragment consisting essentially of the V.sub.H and C.sub.H1 domains; a Fv fragment consisting essentially of the VL and VH domains of a single arm of an antibody, a dAb fragment (Ward et al., Nature 341, 544-546 (1989)), which consists essentially of a VH domain and also called domain antibodies (Holt et al; Trends Biotechnol. 2003 Nov.; 21(11):484-90); camelid or nanobodies (Revets et al; Expert Opin Biol Ther. 2005 Jan.; 5(1):111-24); an isolated complementarity determining region (CDR), and the like. All antibody isotypes may be used to produce antigen-binding fragments. Additionally, antigen-binding fragments may include non-antibody proteinaceous frameworks that may successfully incorporate polypeptide segments in an orientation that confers affinity for a given antigen of interest, such as protein scaffolds. Antigen-binding fragments may be recombinantly produced or produced by enzymatic or chemical cleavage of intact antibodies. The phrase “an antibody or antigen-binding fragment thereof” may be used to denote that a given antigen-binding fragment

incorporates one or more amino acid segments of the antibody referred to in the phrase. When used herein in the context of two or more antibodies or antigen-binding fragments, the term "competes with" or "cross-competes with" indicates that the two or more antibodies or antigen-binding fragments compete for binding to CD123, e.g. compete for CD123 binding in the assay described in Example 9. For some pairs of antibodies or antigen-binding fragments, competition or blocking in the assay of the Examples is only observed when one antibody is coated on the plate and the other is used to compete, and not vice versa. Unless otherwise defined or negated by context, the terms "competes with" or "cross-competes with" when used herein is also intended to cover such pairs of antibodies or antigen-binding fragments.

[0098] The term "epitope" means a protein determinant capable of specific binding to an antibody. Epitopes usually consist of surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents. The epitope may comprise amino acid residues directly involved in the binding and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked or covered by the specifically antigen binding peptide (in other words, the amino acid residue is within the footprint of the specifically antigen binding peptide).

[0099] "Specific binding" or "immunospecific binding" or derivatives thereof when used in the context of antibodies, or antibody fragments, represents binding via domains encoded by immunoglobulin genes or fragments of immunoglobulin genes to one or more epitopes of a protein of interest, without preferentially binding other molecules in a sample containing a mixed population of molecules. Typically, an antibody binds to a cognate antigen with a K_d of less than about 1×10^{-8} M, as measured by a surface plasmon resonance assay or a cell binding assay. Phrases such as "[antigen]-specific" antibody (e.g., CD123-specific antibody) are meant to convey that the recited antibody specifically binds the recited antigen.

[00100] The term " k_d " (sec^{-1}), as used herein, refers to the dissociation rate constant of a particular antibody-antigen interaction. Said value is also referred to as the k_{off} value.

[00101] The term " k_a " ($\text{M}^{-1} \text{sec}^{-1}$), as used herein, refers to the association rate constant of a particular antibody-antigen interaction.

[00102] The term " K_D " (M), as used herein, refers to the dissociation equilibrium constant of a particular antibody-antigen interaction.

[00103] The term " K_A " (M^{-1}), as used herein, refers to the association equilibrium constant of a particular antibody-antigen interaction and is obtained by dividing the k_a by the k_d .

[00104] The term "subject" refers to human and non-human animals, including all vertebrates, e.g., mammals and non-mammals, such as non-human primates, mice, rabbits, sheep, dogs, cats, horses, cows, chickens, amphibians, and reptiles. In many embodiments of the described methods, the subject is a human.

[00105] The term "sample" as used herein refers to a collection of similar fluids, cells, or tissues (e.g., surgically resected tumor tissue, biopsies, including fine needle aspiration), isolated from a subject, as well as fluids, cells, or tissues present within a subject. In some embodiments the sample is a biological fluid. Biological fluids are typically liquids at physiological temperatures and may include naturally occurring fluids present in, withdrawn from, expressed or otherwise extracted from a subject or biological source. Certain biological fluids derive from particular tissues, organs or localized regions and certain other biological fluids may be more globally or systemically situated in a subject or biological source. Examples of biological fluids include blood, serum and serosal fluids, plasma, lymph, urine, saliva, cystic fluid, tear drops, feces, sputum, mucosal secretions of the secretory tissues and organs, vaginal secretions, ascites fluids such as those associated with non-solid tumors, fluids of the pleural, pericardial, peritoneal, abdominal and other body cavities, fluids collected by bronchial lavage and the like. Biological fluids may also include liquid solutions contacted with a subject or biological source, for example, cell and organ culture medium including cell or organ conditioned medium, lavage fluids and the like. The term "sample," as used herein, encompasses materials removed from a subject or materials present in a subject.

[00106] A "known standard" may be a solution having a known amount or concentration of CD123, where the solution may be a naturally occurring solution, such as a sample from a patient known to have early, moderate, late, progressive, or static cancer, or the solution may be a synthetic solution such as buffered water having a known amount of CD123 diluted therein. The known standards, described herein may include CD123 isolated from a subject, recombinant or purified CD123 protein, or a value of CD123 concentration associated with a disease condition.

[00107] The term "CD3" refers to the human CD3 protein multi-subunit complex. The CD3 protein multi-subunit complex is composed to 6 distinctive polypeptide chains. These include a CD3 γ chain (SwissProt P09693), a CD3 δ chain (SwissProt P04234), two CD3 ϵ chains (SwissProt P07766), and one CD3 ζ chain homodimer (SwissProt 20963), and which is

associated with the T cell receptor α and β chain. The term “CD3” includes any CD3 variant, isoform and species homolog which is naturally expressed by cells (including T cells) or can be expressed on cells transfected with genes or cDNA encoding those polypeptides, unless noted.

[00108] As used herein, the terms “alpha subunit of the IL-3 receptor,” “IL3R α ,” “CD123,” “IL3R α chain” and “IL3R α subunit” refer interchangeably to an antigenic determinant detectable on leukemia precursor cells, which immunobinds interleukin-3 (IL3). In a specific embodiment, the CD123 is the human CD123. In a specific embodiment, the CD123 is cynomolgus monkey CD123. In a specific embodiment, the CD123 is CD123 SP1. In a specific embodiment, the CD123 is CD123 SP2. The term “CD123” includes any CD123 variant, isoform and species homolog, unless noted.

[00109] A “CD123 x CD3 antibody” is a multispecific antibody, optionally a bispecific antibody, which comprises two different antigen-binding regions, one of which binds specifically to the antigen CD123 and one of which binds specifically to CD3. A multispecific antibody can be a bispecific antibody, diabody, or similar molecule (see for instance *PNAS USA* 90(14), 6444-8 (1993) for a description of diabodies). The bispecific antibodies, diabodies, and the like, provided herein may bind any suitable target in addition to a portion of CD123. The term “bispecific antibody” is to be understood as an antibody having two different antigen-binding regions defined by different antibody sequences. This can be understood as different target binding but includes as well binding to different epitopes in one target.

[00110] A “reference sample” is a sample that may be compared against another sample, such as a test sample, to allow for characterization of the compared sample. The reference sample will have some characterized property that serves as the basis for comparison with the test sample. For instance, a reference sample may be used as a benchmark for CD123 levels that are indicative of a subject having cancer. The reference sample does not necessarily have to be analyzed in parallel with the test sample, thus in some instances the reference sample may be a numerical value or range previously determined to characterize a given condition, such as CD123 levels that are indicative of cancer in a subject. The term also includes samples used for comparative purposes that are known to be associated with a physiologic state or disease condition, such as CD123-expressing cancer, but that have an unknown amount of CD123.

[00111] The term “progression,” as used in the context of progression of CD123-expressing cancer, includes the change of a cancer from a less severe to a more severe state. This may include an increase in the number or severity of tumors, the degree of metastasis, the speed with which the cancer is growing or spreading, and the like. For example, “the

progression of colon cancer” includes the progression of such a cancer from a less severe to a more severe state, such as the progression from stage I to stage II, from stage II to stage III, etc.

[00112] The term “regression,” as used in the context of regression of CD123-expressing cancer, includes the change of a cancer from a more severe to a less severe state. This could include a decrease in the number or severity of tumors, the degree of metastasis, the speed with which the cancer is growing or spreading, and the like. For example, “the regression of colon cancer” includes the regression of such a cancer from a more severe to a less severe state, such as the progression from stage III to stage II, from stage II to stage I, etc.

[00113] The term “stable” as used in the context of stable CD123-expressing cancer, is intended to describe a disease condition that is not, or has not, changed significantly enough over a clinically relevant period of time to be considered a progressing cancer or a regressing cancer.

[00114] The embodiments described herein are not limited to particular methods, reagents, compounds, compositions or biological systems, which can, of course, vary.

CD123-Specific Antibodies and Antigen-Binding Fragments

[00115] Described herein are isolated monoclonal antibodies or antigen-binding fragments that specifically bind CD123. The general structure of an antibody molecule comprises an antigen binding domain, which includes heavy and light chains, and the Fc domain, which serves a variety of functions, including complement fixation and binding antibody receptors.

[00116] The described CD123-specific antibodies or antigen-binding fragments include all isotypes, IgA, IgD, IgE, IgG and IgM, and synthetic multimers of the four-chain immunoglobulin structure. The described antibodies or antigen-binding fragments also include the IgY isotype generally found in hen or turkey serum and hen or turkey egg yolk.

[00117] The CD123-specific antibodies and antigen-binding fragments may be derived from any species by recombinant means. For example, the antibodies or antigen-binding fragments may be mouse, rat, goat, horse, swine, bovine, chicken, rabbit, camelid, donkey, human, or chimeric versions thereof. For use in administration to humans, non-human derived antibodies or antigen-binding fragments may be genetically or structurally altered to be less antigenic upon administration to a human patient.

[00118] In some embodiments, the antibodies or antigen-binding fragments are chimeric. As used herein, the term “chimeric” refers to an antibody, or antigen-binding fragment thereof, having at least some portion of at least one variable domain derived from the antibody

amino acid sequence of a non-human mammal, a rodent, or a reptile, while the remaining portions of the antibody, or antigen-binding fragment thereof, are derived from a human.

[00119] In some embodiments, the antibodies are humanized antibodies. Humanized antibodies may be chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin sequence. The humanized antibody may include at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

[00120] The antibodies or antigen-binding fragments described herein can occur in a variety of forms, but will include one or more of the antibody CDRs shown in Table 1.

[00121] Described herein are isolated antibodies and antigen-binding fragments that immunospecifically bind to CD123. In some embodiments, the CD123-specific antibodies or antigen-binding fragments are human IgG, or derivatives thereof. While the CD123-specific antibodies or antigen-binding fragments exemplified herein are human, the antibodies or antigen-binding fragments exemplified may be chimerized.

[00122] In some embodiments are provided a CD123-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1. In some embodiments are provided a CD123-specific antibody, or an antigen-binding fragment thereof, comprising a heavy chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1 and a light chain comprising a CDR1, a CDR2, and a CDR3 of any one of the antibodies described in Table 1.

[00123] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 006, a heavy chain CDR2 comprising SEQ ID NO: 007, and a heavy chain CDR3 comprising SEQ ID NO: 008. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy

chain CDR1 comprising SEQ ID NO: 006, a heavy chain CDR2 comprising SEQ ID NO: 007, a heavy chain CDR3 comprising SEQ ID NO: 008, a light chain CDR1 comprising SEQ ID NO: 009, a light chain CDR2 comprising SEQ ID NO: 010, and a light chain CDR3 comprising SEQ ID NO: 011. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 119. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 119 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 164. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00124] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 012, a heavy chain CDR2 comprising SEQ ID NO: 013, and a heavy chain CDR3 comprising SEQ ID NO: 014. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 012, a heavy chain CDR2 comprising SEQ ID NO: 013, a heavy chain CDR3 comprising SEQ ID NO: 014, a light chain CDR1 comprising SEQ ID NO: 015, a light chain CDR2 comprising SEQ ID NO: 016, and a light chain CDR3 comprising SEQ ID NO: 017. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 120. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 120 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 165. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00125] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 018, a heavy chain CDR2 comprising SEQ ID NO: 019, and a heavy chain CDR3 comprising SEQ ID NO: 020. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 018, a heavy chain CDR2 comprising SEQ ID NO: 019, a heavy chain CDR3 comprising SEQ ID NO: 020, a light chain CDR1 comprising SEQ ID NO: 009, a light chain CDR2 comprising SEQ ID NO: 010, and a light chain CDR3 comprising SEQ ID NO: 011. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 121. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 121 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 164. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00126] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 021, a heavy chain CDR2 comprising SEQ ID NO: 022, and a heavy chain CDR3 comprising SEQ ID NO: 023. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 021, a heavy chain CDR2 comprising SEQ ID NO: 022, a heavy chain CDR3 comprising SEQ ID NO: 023, a light chain CDR1 comprising SEQ ID NO: 024, a light chain CDR2 comprising SEQ ID NO: 025, and a light chain CDR3 comprising SEQ ID NO: 026. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 122. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 122 and a light chain variable domain substantially the same as, or

identical to, SEQ ID NO: 166. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00127] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 027, a heavy chain CDR2 comprising SEQ ID NO: 028, and a heavy chain CDR3 comprising SEQ ID NO: 029. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 027, a heavy chain CDR2 comprising SEQ ID NO: 028, a heavy chain CDR3 comprising SEQ ID NO: 029, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 123. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 123 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00128] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 035. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 035, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same

as, or identical to, SEQ ID NO: 124. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 124 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00129] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 036. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 036, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 125. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 125 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00130] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 037. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 037, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to

CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 126. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 126 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00131] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 038. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 038, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 127. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 127 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00132] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 040, and a heavy chain CDR3 comprising SEQ ID NO: 041. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 040, a heavy chain CDR3 comprising SEQ ID NO: 041, a light chain CDR1 comprising SEQ ID NO:

030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 128. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 128 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00133] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 042. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 042, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 129. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 129 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00134] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 043. In some

embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 034, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 043, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 130. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 130 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00135] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 044, and a heavy chain CDR3 comprising SEQ ID NO: 045. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 044, a heavy chain CDR3 comprising SEQ ID NO: 045, a light chain CDR1 comprising SEQ ID NO: 015, a light chain CDR2 comprising SEQ ID NO: 016, and a light chain CDR3 comprising SEQ ID NO: 017. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 131. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 131 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 165. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00136] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 046, and a heavy chain CDR3 comprising SEQ ID NO: 047. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 046, a heavy chain CDR3 comprising SEQ ID NO: 047, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 132. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 132 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00137] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 048. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 048, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 133. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 133 and a light chain variable domain substantially the same as, or

identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00138] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 049. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 049, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 134. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 134 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00139] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 050. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 050, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same

as, or identical to, SEQ ID NO: 135. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 135 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00140] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 051, a heavy chain CDR2 comprising SEQ ID NO: 052, and a heavy chain CDR3 comprising SEQ ID NO: 053. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 051, a heavy chain CDR2 comprising SEQ ID NO: 052, a heavy chain CDR3 comprising SEQ ID NO: 053, a light chain CDR1 comprising SEQ ID NO: 024, a light chain CDR2 comprising SEQ ID NO: 025, and a light chain CDR3 comprising SEQ ID NO: 054. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 136. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 136 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 168. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00141] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 055, a heavy chain CDR2 comprising SEQ ID NO: 056, and a heavy chain CDR3 comprising SEQ ID NO: 057. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 055, a heavy chain CDR2 comprising SEQ ID NO: 056, a heavy chain CDR3 comprising SEQ ID NO: 057, a light chain CDR1 comprising SEQ ID NO: 058, a light chain CDR2 comprising SEQ ID NO: 059, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to

CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 137. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 137 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 169. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00142] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 060. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 060, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 138. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 138 and a light chain variable domain substantially the same as, or identical to, gSEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00143] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 061. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 061, a light chain CDR1 comprising SEQ ID NO:

062, a light chain CDR2 comprising SEQ ID NO: 063, and a light chain CDR3 comprising SEQ ID NO: 064. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 139. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 139 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 170. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00144] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 065. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 065, a light chain CDR1 comprising SEQ ID NO: 066, a light chain CDR2 comprising SEQ ID NO: 067, and a light chain CDR3 comprising SEQ ID NO: 068. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 140. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 140 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 171. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00145] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 069. In some

embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 069, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 070. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 141. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 141 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 172. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00146] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 071. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 071, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 142. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 142 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00147] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 072, and a heavy chain CDR3 comprising SEQ ID NO: 073. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 072, a heavy chain CDR3 comprising SEQ ID NO: 073, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 143. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 143 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00148] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 074. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 074, a light chain CDR1 comprising SEQ ID NO: 075, a light chain CDR2 comprising SEQ ID NO: 076, and a light chain CDR3 comprising SEQ ID NO: 077. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 144. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 144 and a light chain variable domain substantially the same as, or

identical to, SEQ ID NO: 173. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00149] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 078. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 078, a light chain CDR1 comprising SEQ ID NO: 079, a light chain CDR2 comprising SEQ ID NO: 063, and a light chain CDR3 comprising SEQ ID NO: 080. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 145. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 145 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 174. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00150] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 081. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 081, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same

as, or identical to, SEQ ID NO: 146. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 146 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00151] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 082. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 082, a light chain CDR1 comprising SEQ ID NO: 083, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 084. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 147. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 147 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 175. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00152] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 085, and a heavy chain CDR3 comprising SEQ ID NO: 086. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 085, a heavy chain CDR3 comprising SEQ ID NO: 086, a light chain CDR1 comprising SEQ ID NO: 087, a light chain CDR2 comprising SEQ ID NO: 067, and a light chain CDR3 comprising SEQ ID NO: 088. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to

CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 148. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 148 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 176. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00153] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 089, and a heavy chain CDR3 comprising SEQ ID NO: 090. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 089, a heavy chain CDR3 comprising SEQ ID NO: 090, a light chain CDR1 comprising SEQ ID NO: 091, a light chain CDR2 comprising SEQ ID NO: 076, and a light chain CDR3 comprising SEQ ID NO: 092. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 149. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 149 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 177. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00154] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 093. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 093, a light chain CDR1 comprising SEQ ID NO:

030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 150. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 150 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00155] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 094. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 094, a light chain CDR1 comprising SEQ ID NO: 095, a light chain CDR2 comprising SEQ ID NO: 076, and a light chain CDR3 comprising SEQ ID NO: 096. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 151. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 151 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 178. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00156] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 097. In some

embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 097, a light chain CDR1 comprising SEQ ID NO: 098, a light chain CDR2 comprising SEQ ID NO: 067, and a light chain CDR3 comprising SEQ ID NO: 099. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 152. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 152 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 179. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00157] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 100. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 100, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 101. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 153. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 153 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 180. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00158] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 102. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 102, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 154. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 154 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00159] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 103. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 103, a light chain CDR1 comprising SEQ ID NO: 104, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 105. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 155. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 155 and a light chain variable domain substantially the same as, or

identical to, SEQ ID NO: 181. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00160] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 106. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 039, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 106, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 156. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 156 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00161] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 107. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 107, a light chain CDR1 comprising SEQ ID NO: 108, a light chain CDR2 comprising SEQ ID NO: 109, and a light chain CDR3 comprising SEQ ID NO: 110. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same

as, or identical to, SEQ ID NO: 157. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 157 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 182. . The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00162] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 111. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 111, a light chain CDR1 comprising SEQ ID NO: 112, a light chain CDR2 comprising SEQ ID NO: 076, and a light chain CDR3 comprising SEQ ID NO: 113. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 158. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 158 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 183. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00163] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 114, a heavy chain CDR2 comprising SEQ ID NO: 022, and a heavy chain CDR3 comprising SEQ ID NO: 115. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 114, a heavy chain CDR2 comprising SEQ ID NO: 022, a heavy chain CDR3 comprising SEQ ID NO: 115, a light chain CDR1 comprising SEQ ID NO: 024, a light chain CDR2 comprising SEQ ID NO: 025, and a light chain CDR3 comprising SEQ ID NO: 026. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to

CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 159. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 159 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 166. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00164] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 114, a heavy chain CDR2 comprising SEQ ID NO: 022, and a heavy chain CDR3 comprising SEQ ID NO: 116. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 114, a heavy chain CDR2 comprising SEQ ID NO: 022, a heavy chain CDR3 comprising SEQ ID NO: 116, a light chain CDR1 comprising SEQ ID NO: 015, a light chain CDR2 comprising SEQ ID NO: 016, and a light chain CDR3 comprising SEQ ID NO: 017. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 160. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 160 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 165. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00165] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 117, a heavy chain CDR2 comprising SEQ ID NO: 013, and a heavy chain CDR3 comprising SEQ ID NO: 118. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 117, a heavy chain CDR2 comprising SEQ ID NO: 013, a heavy chain CDR3 comprising SEQ ID NO: 118, a light chain CDR1 comprising SEQ ID NO:

015, a light chain CDR2 comprising SEQ ID NO: 016, and a light chain CDR3 comprising SEQ ID NO: 017. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 161. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 161 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 165. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00166] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 051, a heavy chain CDR2 comprising SEQ ID NO: 052, and a heavy chain CDR3 comprising SEQ ID NO: 053. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 051, a heavy chain CDR2 comprising SEQ ID NO: 052, a heavy chain CDR3 comprising SEQ ID NO: 053, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 162. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 162 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00167] In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, and a heavy chain CDR3 comprising SEQ ID NO: 042. In some

embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain CDR1 comprising SEQ ID NO: 033, a heavy chain CDR2 comprising SEQ ID NO: 034, a heavy chain CDR3 comprising SEQ ID NO: 042, a light chain CDR1 comprising SEQ ID NO: 030, a light chain CDR2 comprising SEQ ID NO: 031, and a light chain CDR3 comprising SEQ ID NO: 032. This CD123-specific antibody or antigen-binding fragment may comprise human framework sequences. This CD123-specific antibody or antigen-binding fragment may bind to CD123 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 163. In some embodiments, the CD123-specific antibodies and antigen-binding fragments comprise a heavy chain variable domain substantially the same as, or identical to, SEQ ID NO: 163 and a light chain variable domain substantially the same as, or identical to, SEQ ID NO: 167. The heavy chain variable domain and light chain variable domain of antibodies discussed in this paragraph are suitable for inclusion in bispecific constructs in which one arm is an anti-CD123 arm.

[00168] The anti-CD123 antibodies and antigen-binding fragments provided by the invention also include antibodies which compete for binding with the antibodies described above. Competition for binding can be determined using a competition binding ELISA, in line with the technique described below in Example 5. Competitive binding may be determined by detecting at least 20% inhibition of the binding of a first antibody by a second antibody, irrespective of the order in which the antibodies are bound to CD123 (*i.e.* if when antibody A is bound to CD123 before antibody B, only 10% inhibition is observed, but when antibody B is bound to CD123 before antibody A, 30% inhibition is observed, then because greater than 20% inhibition has been observed in one of the experiments, competitive binding may be concluded).

[00169] In some embodiments, the antibodies or antigen-binding fragments are IgG, or derivatives thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the antibody has an IgG1 isotype, the antibody contains L234A, L235A, and K409R substitution(s) in its Fc region. In some embodiments wherein the antibody has an IgG4 isotype, the antibody contains S228P, L234A, and L235A substitutions in its Fc region. The specific antibodies defined by CDR and/or variable domain sequence discussed in the above paragraphs may include these modifications.

[00170] Also disclosed are isolated polynucleotides that encode the antibodies or antigen-binding fragments that immunospecifically bind to CD123. The isolated polynucleotides

capable of encoding the variable domain segments provided herein may be included on the same, or different, vectors to produce antibodies or antigen-binding fragments.

[00171] Polynucleotides encoding recombinant antigen-binding proteins also are within the scope of the disclosure. In some embodiments, the polynucleotides described (and the peptides they encode) include a leader sequence. Any leader sequence known in the art may be employed. The leader sequence may include, but is not limited to, a restriction site or a translation start site.

[00172] The CD123-specific antibodies or antigen-binding fragments described herein include variants having single or multiple amino acid substitutions, deletions, or additions that retain the biological properties (e.g., binding affinity or immune effector activity) of the described CD123-specific antibodies or antigen-binding fragments. In the context of the present invention the following notations are, unless otherwise indicated, used to describe a mutation; i) substitution of an amino acid in a given position is written as e.g. K409R which means a substitution of a Lysine in position 409 with an Arginine; and ii) for specific variants the specific three or one letter codes are used, including the codes Xaa and X to indicate any amino acid residue. Thus, the substitution of Arginine for Lysine in position 409 is designated as: K409R, or the substitution of any amino acid residue for Lysine in position 409 is designated as K409X. In case of deletion of Lysine in position 409 it is indicated by K409*. The skilled person may produce variants having single or multiple amino acid substitutions, deletions, or additions.

[00173] These variants may include: (a) variants in which one or more amino acid residues are substituted with conservative or nonconservative amino acids, (b) variants in which one or more amino acids are added to or deleted from the polypeptide, (c) variants in which one or more amino acids include a substituent group, and (d) variants in which the polypeptide is fused with another peptide or polypeptide such as a fusion partner, a protein tag or other chemical moiety, that may confer useful properties to the polypeptide, such as, for example, an epitope for an antibody, a polyhistidine sequence, a biotin moiety and the like. Antibodies or antigen-binding fragments described herein may include variants in which amino acid residues from one species are substituted for the corresponding residue in another species, either at the conserved or nonconserved positions. In other embodiments, amino acid residues at nonconserved positions are substituted with conservative or nonconservative residues. The techniques for obtaining these variants, including genetic (deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art.

[00174] The CD123-specific antibodies or antigen-binding fragments described herein may embody several antibody isotypes, such as IgM, IgD, IgG, IgA and IgE. In some embodiments the antibody isotype is IgG1, IgG2, IgG3, or IgG4 isotype, preferably IgG1 or IgG4 isotype. Antibody or antigen-binding fragment thereof specificity is largely determined by the amino acid sequence, and arrangement, of the CDRs. Therefore, the CDRs of one isotype may be transferred to another isotype without altering antigen specificity. Alternatively, techniques have been established to cause hybridomas to switch from producing one antibody isotype to another (isotype switching) without altering antigen specificity. Accordingly, such antibody isotypes are within the scope of the described antibodies or antigen-binding fragments.

[00175] The CD123-specific antibodies or antigen-binding fragments described herein have binding affinities for CD123 SP1 that include a dissociation constant (K_D) of less than about 5×10^{-7} M, preferably less than about 5×10^{-8} M. In some embodiments, the CD123-specific antibodies or antigen-binding fragments described herein have binding affinities for CD123 SP2 that include a dissociation constant (K_D) of less than about 5×10^{-7} M, preferably less than about 5×10^{-8} M. The affinity of the described CD123-specific antibodies, or antigen-binding fragments, may be determined by a variety of methods known in the art, such as surface plasmon resonance or ELISA-based methods. Assays for measuring affinity by SPR include assays performed using a BIAcore 3000 machine, where the assay is performed at room temperature (*e.g.* at or near 25°C), wherein the antibody capable of binding to CD123 is captured on the BIAcore sensor chip by an anti-Fc antibody (*e.g.* goat anti-human IgG Fc specific antibody Jackson ImmunoResearch laboratories Prod # 109-005-098) to a level around 75RUs, followed by the collection of association and dissociation data at a flow rate of 40 μ l/min.

[00176] Also provided are vectors comprising the polynucleotides described herein. The vectors can be expression vectors. Recombinant expression vectors containing a sequence encoding a polypeptide of interest are thus contemplated as within the scope of this disclosure. The expression vector may contain one or more additional sequences such as but not limited to regulatory sequences (*e.g.*, promoter, enhancer), a selection marker, and a polyadenylation signal. Vectors for transforming a wide variety of host cells are well known and include, but are not limited to, plasmids, phagemids, cosmids, baculoviruses, bacmids, bacterial artificial chromosomes (BACs), yeast artificial chromosomes (YACs), as well as other bacterial, yeast and viral vectors.

[00177] Recombinant expression vectors within the scope of the description include synthetic, genomic, or cDNA-derived nucleic acid fragments that encode at least one

recombinant protein which may be operably linked to suitable regulatory elements. Such regulatory elements may include a transcriptional promoter, sequences encoding suitable mRNA ribosomal binding sites, and sequences that control the termination of transcription and translation. Expression vectors, especially mammalian expression vectors, may also include one or more nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, other 5' or 3' flanking nontranscribed sequences, 5' or 3' nontranslated sequences (such as necessary ribosome binding sites), a polyadenylation site, splice donor and acceptor sites, or transcriptional termination sequences. An origin of replication that confers the ability to replicate in a host may also be incorporated.

[00178] The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. Exemplary vectors may be constructed as described by Okayama and Berg, 3 *Mol. Cell. Biol.* 280 (1983).

[00179] In some embodiments, the antibody- or antigen-binding fragment-coding sequence is placed under control of a powerful constitutive promoter, such as the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin, human myosin, human hemoglobin, human muscle creatine, and others. In addition, many viral promoters function constitutively in eukaryotic cells and are suitable for use with the described embodiments. Such viral promoters include without limitation, Cytomegalovirus (CMV) immediate early promoter, the early and late promoters of SV40, the Mouse Mammary Tumor Virus (MMTV) promoter, the long terminal repeats (LTRs) of Maloney leukemia virus, Human Immunodeficiency Virus (HIV), Epstein Barr Virus (EBV), Rous Sarcoma Virus (RSV), and other retroviruses, and the thymidine kinase promoter of Herpes Simplex Virus. In one embodiment, the CD123-specific antibody or antigen-binding fragment thereof coding sequence is placed under control of an inducible promoter such as the metallothionein promoter, tetracycline-inducible promoter, doxycycline-inducible promoter, promoters that contain one or more interferon-stimulated response elements (ISRE) such as protein kinase R 2',5'-oligoadenylate synthetases, Mx genes, ADAR1, and the like.

[00180] Vectors described herein may contain one or more Internal Ribosome Entry Site(s) (IRES). Inclusion of an IRES sequence into fusion vectors may be beneficial for enhancing expression of some proteins. In some embodiments the vector system will include one or more polyadenylation sites (e.g., SV40), which may be upstream or downstream of any of the aforementioned nucleic acid sequences. Vector components may be contiguously linked, or arranged in a manner that provides optimal spacing for expressing the gene products (i.e., by the

introduction of “spacer” nucleotides between the ORFs), or positioned in another way. Regulatory elements, such as the IRES motif, may also be arranged to provide optimal spacing for expression.

[00181] The vectors may comprise selection markers, which are well known in the art. Selection markers include positive and negative selection markers, for example, antibiotic resistance genes (e.g., neomycin resistance gene, a hygromycin resistance gene, a kanamycin resistance gene, a tetracycline resistance gene, a penicillin resistance gene), glutamate synthase genes, HSV-TK, HSV-TK derivatives for ganciclovir selection, or bacterial purine nucleoside phosphorylase gene for 6-methylpurine selection (Gadi et al., 7 *Gene Ther.* 1738-1743 (2000)). A nucleic acid sequence encoding a selection marker or the cloning site may be upstream or downstream of a nucleic acid sequence encoding a polypeptide of interest or cloning site.

[00182] The vectors described herein may be used to transform various cells with the genes encoding the described antibodies or antigen-binding fragments. For example, the vectors may be used to generate CD123-specific antibody or antigen-binding fragment-producing cells. Thus, another aspect features host cells transformed with vectors comprising a nucleic acid sequence encoding an antibody or antigen-binding fragment thereof that specifically binds CD123, such as the antibodies or antigen-binding fragments described and exemplified herein.

[00183] Numerous techniques are known in the art for the introduction of foreign genes into cells and may be used to construct the recombinant cells for purposes of carrying out the described methods, in accordance with the various embodiments described and exemplified herein. The technique used should provide for the stable transfer of the heterologous gene sequence to the host cell, such that the heterologous gene sequence is heritable and expressible by the cell progeny, and so that the necessary development and physiological functions of the recipient cells are not disrupted. Techniques which may be used include but are not limited to chromosome transfer (e.g., cell fusion, chromosome mediated gene transfer, micro cell mediated gene transfer), physical methods (e.g., transfection, spheroplast fusion, microinjection, electroporation, liposome carrier), viral vector transfer (e.g., recombinant DNA viruses, recombinant RNA viruses) and the like (described in Cline, 29 *Pharmac. Ther.* 69-92 (1985)). Calcium phosphate precipitation and polyethylene glycol (PEG)-induced fusion of bacterial protoplasts with mammalian cells may also be used to transform cells.

[00184] Cells suitable for use in the expression of the CD123-specific antibodies or antigen-binding fragments described herein are preferably eukaryotic cells, more preferably cells of plant, rodent, or human origin, for example but not limited to NSO, CHO, CHOK1, perC.6,

Tk-ts13, BHK, HEK293 cells, COS-7, T98G, CV-1/EBNA, L cells, C127, 3T3, HeLa, NS1, Sp2/0 myeloma cells, and BHK cell lines, among others. In addition, expression of antibodies may be accomplished using hybridoma cells. Methods for producing hybridomas are well established in the art.

[00185] Cells transformed with expression vectors described herein may be selected or screened for recombinant expression of the antibodies or antigen-binding fragments described herein. Recombinant-positive cells are expanded and screened for subclones exhibiting a desired phenotype, such as high level expression, enhanced growth properties, or the ability to yield proteins with desired biochemical characteristics, for example, due to protein modification or altered post-translational modifications. These phenotypes may be due to inherent properties of a given subclone or to mutation. Mutations may be effected through the use of chemicals, UV-wavelength light, radiation, viruses, insertional mutagens, inhibition of DNA mismatch repair, or a combination of such methods.

Methods of using CD123-specific antibodies for treatment

[00186] Provided herein are CD123-specific antibodies or antigen-binding fragments thereof for use in therapy. In particular, these antibodies or antigen-binding fragments may be useful in treating cancer, such as CD123-expressing cancer. Accordingly, the invention provides a method of treating cancer comprising administering an antibody as described herein, such as CD123-specific antibodies or antigen-binding fragments. For example, the use may be by inhibiting a biological effect of IL-3 by preventing IL-3 from binding to IL-3R or where the antibody is conjugated to a toxin, so targeting the toxin to the CD123-expressing cancer. In some embodiments CD123-expressing cancer includes hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN). The antibodies for use in these methods include those described herein above, for example a CD123-specific antibody or antigen-binding fragment that binds to an epitope including one or more residues from the segment of CD123 SP2 ECD comprising residues 195 - 202 (RARERVYE (SEQ ID NO: 234)) and/or the segment of CD123 SP2 ECD comprising residues 156-161 (RKFRYE (SEQ ID NO:232)) and/or the segment of CD123 SP2 ECD comprising residues 173 – 178 (TEQVRD (SEQ ID NO: 233)). Also useful for use in these methods are antibodies with

the features set out in Table 1, for example the CDRs or variable domain sequences, and in the further discussion of these antibodies.

[00187] In some embodiments described herein, immune effector properties of the CD123-specific antibodies may be enhanced or silenced through Fc modifications by techniques known to those skilled in the art. For example, Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cell-mediated phagocytosis (ADCP), down regulation of cell surface receptors (e.g., B cell receptor; BCR), etc. may be provided and/or controlled by modifying residues in the Fc responsible for these activities.

[00188] "Antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a cell-mediated reaction in which non-specific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell.

[00189] The ability of monoclonal antibodies to induce ADCC can be enhanced by engineering their oligosaccharide component. Human IgG1 or IgG3 are N-glycosylated at Asn297 with the majority of the glycans in the well known biantennary G0, G0F, G1, G1F, G2 or G2F forms. Antibodies produced by non-engineered CHO cells typically have a glycan fucose content of about at least 85%. The removal of the core fucose from the biantennary complex-type oligosaccharides attached to the Fc regions enhances the ADCC of antibodies via improved Fc.gamma.RIIIa binding without altering antigen binding or CDC activity. Such mAbs can be achieved using different methods reported to lead to the successful expression of relatively high defucosylated antibodies bearing the biantennary complex-type of Fc oligosaccharides such as control of culture osmolality (Konno et al., *Cytotechnology* 64:249-65, 2012), application of a variant CHO line Lec13 as the host cell line (Shields et al., *J Biol Chem* 277:26733-26740, 2002), application of a variant CHO line EB66 as the host cell line (Olivier et al., *MAbs*; 2(4), 2010; Epub ahead of print; PMID:20562582), application of a rat hybridoma cell line YB2/0 as the host cell line (Shinkawa et al., *J Biol Chem* 278:3466-3473, 2003), introduction of small interfering RNA specifically against the .alpha. 1,6-fucosyltransferase (FUT8) gene (Mori et al., *Biotechnol Bioeng* 88:901-908, 2004), or coexpression of .beta.-1,4-N-acetylglucosaminyltransferase III and Golgi .alpha.-mannosidase II or a potent alpha-mannosidase I inhibitor, kifunensine (Ferrara et al., *J Biol Chem* 281:5032-5036, 2006, Ferrara et al., *Biotechnol Bioeng* 93:851-861, 2006; Xhou et al., *Biotechnol Bioeng* 99:652-65, 2008).

[00190] In some embodiments described herein, ADCC elicited by the CD123 antibodies may also be enhanced by certain substitutions in the antibody Fc. Exemplary substitutions are for example substitutions at amino acid positions 256, 290, 298, 312, 356, 330, 333, 334, 360, 378 or 430 (residue numbering according to the EU index) as described in U.S. Pat. No. 6,737,056.

Methods of detecting CD123

[00191] Provided herein are methods for detecting CD123 in a biological sample by contacting the sample with an antibody, or antigen-binding fragment thereof, described herein. As described herein, the sample may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like. In some embodiments the described methods include detecting CD123 in a biological sample by contacting the sample with any of the CD123-specific antibodies or antigen-binding fragments thereof described herein.

[00192] In some embodiments the sample may be contacted with more than one of the CD123-specific antibodies or antigen-binding fragments described herein. For example, a sample may be contacted with a first CD123-specific antibody, or antigen-binding fragment thereof, and then contacted with a second CD123-specific antibody, or antigen-binding fragment thereof, wherein the first antibody or antigen-binding fragment and the second antibody or antigen-binding fragment are not the same antibody or antigen-binding fragment. In some embodiments, the first antibody, or antigen-binding fragment thereof, may be affixed to a surface, such as a multiwell plate, chip, or similar substrate prior to contacting the sample. In other embodiments the first antibody, or antigen-binding fragment thereof, may not be affixed, or attached, to anything at all prior to contacting the sample.

[00193] The described CD123-specific antibodies and antigen-binding fragments may be detectably labeled. In some embodiments labeled antibodies and antigen-binding fragments may facilitate the detection CD123 via the methods described herein. Many such labels are readily known to those skilled in the art. For example, suitable labels include, but should not be considered limited to, radiolabels, fluorescent labels, epitope tags, biotin, chromophore labels, ECL labels, or enzymes. More specifically, the described labels include ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase

and beta-galactosidase, poly-histidine (HIS tag), acridine dyes, cyanine dyes, fluorone dyes, oxazin dyes, phenanthridine dyes, rhodamine dyes, Alexa Fluor® dyes, and the like.

[00194] The described CD123-specific antibodies and antigen-binding fragments may be used in a variety of assays to detect CD123 in a biological sample. Some suitable assays include, but should not be considered limited to, western blot analysis, radioimmunoassay, surface plasmon resonance, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

[00195] In some embodiments described herein detection of CD123-expressing cancer cells in a subject may be used to determine that the subject may be treated with a therapeutic agent directed against CD123.

[00196] CD123 is present at detectable levels in blood and serum samples. Thus, provided herein are methods for detecting CD123 in a sample derived from blood, such as a serum sample, by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds CD123. The blood sample, or a derivative thereof, may be diluted, fractionated, or otherwise processed to yield a sample upon which the described method may be performed. In some embodiments, CD123 may be detected in a blood sample, or a derivative thereof, by any number of assays known in the art, such as, but not limited to, western blot analysis, radioimmunoassay, surface plasmon resonance, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

Methods for Diagnosing Cancer

[00197] Provided herein are methods for diagnosing CD123-expressing cancer in a subject. In some embodiments CD123-expressing cancer includes hematological cancers, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (BPDCN). In some embodiments, as described above, detecting CD123 in a biological sample, such as a blood sample or a serum sample, provides the ability to diagnose cancer in the subject from whom the sample was obtained. Alternatively, in some embodiments other samples such as a histological sample, a fine needle aspirate sample, resected tumor tissue, circulating cells, circulating tumor

cells, and the like, may also be used to assess whether the subject from whom the sample was obtained has cancer. In some embodiments, it may already be known that the subject from whom the sample was obtained has cancer, but the type of cancer afflicting the subject may not yet have been diagnosed or a preliminary diagnosis may be unclear, thus detecting CD123 in a biological sample obtained from the subject can allow for, or clarify, diagnosis of the cancer. For example, a subject may be known to have cancer, but it may not be known, or may be unclear, whether the subject's cancer is CD123-expressing.

[00198] In some embodiments the described methods involve assessing whether a subject is afflicted with CD123-expressing cancer by determining the amount of CD123 that is present in a biological sample derived from the subject; and comparing the observed amount of CD123 with the amount of CD123 in a control, or reference, sample, wherein a difference between the amount of CD123 in the sample derived from the subject and the amount of CD123 in the control, or reference, sample is an indication that the subject is afflicted with a CD123-expressing cancer. In another embodiment the amount of CD123 observed in a biological sample obtained from a subject may be compared to levels of CD123 known to be associated with certain forms or stages of cancer, to determine the form or stage of the subject's cancer. In some embodiments the amount of CD123 in the sample derived from the subject is assessed by contacting the sample with an antibody, or an antigen-binding fragment thereof, that immunospecifically binds CD123, such as the CD123-specific antibodies described herein. The sample assessed for the presence of CD123 may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like. In some embodiments CD123-expressing cancer includes hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN). In some embodiments the subject is a human.

[00199] In some embodiments the method of diagnosing a CD123-expressing cancer will involve: contacting a biological sample of a subject with a CD123-specific antibody, or an antigen-binding fragment thereof (such as those derivable from the antibodies and fragments provided in Table 1), quantifying the amount of CD123 present in the sample that is bound by the antibody or antigen-binding fragment thereof, comparing the amount of CD123 present in the sample to a known standard or reference sample; and determining whether the subject's CD123

levels fall within the levels of CD123 associated with cancer. In an additional embodiment, the diagnostic method can be followed with an additional step of administering or prescribing a cancer-specific treatment. In another embodiment, the diagnostic method can be followed with an additional step of transmitting the results of the determination to facilitate treatment of the cancer. In some embodiments the cancer-specific treatment may be directed against CD123-expressing cancers, such as the CD123 x CD3 multispecific antibodies described herein.

[00200] In some embodiments the described methods involve assessing whether a subject is afflicted with CD123-expressing cancer by determining the amount of CD123 present in a blood or serum sample obtained from the subject; and comparing the observed amount of CD123 with the amount of CD123 in a control, or reference, sample, wherein a difference between the amount of CD123 in the sample derived from the subject and the amount of CD123 in the control, or reference, sample is an indication that the subject is afflicted with a CD123-expressing cancer.

[00201] In some embodiments the control, or reference, sample may be derived from a subject that is not afflicted with CD123-expressing cancer. In some embodiments the control, or reference, sample may be derived from a subject that is afflicted with CD123-expressing cancer. In some embodiments where the control, or reference, sample is derived from a subject that is not afflicted with CD123-expressing cancer, an observed increase in the amount of CD123 present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is afflicted with CD123-expressing cancer. In some embodiments where the control sample is derived from a subject that is not afflicted with CD123-expressing cancer, an observed decrease or similarity in the amount of CD123 present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is not afflicted with CD123-expressing cancer. In some embodiments where the control or reference sample is derived from a subject that is afflicted with CD123-expressing cancer, an observed similarity in the amount of CD123 present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is afflicted with CD123-expressing cancer. In some embodiments where the control or reference sample is derived from a subject that is afflicted with CD123-expressing cancer, an observed decrease in the amount of CD123 present in the test sample, relative to that observed for the control or reference sample, is an indication that the subject being assessed is not afflicted with CD123-expressing cancer.

[00202] In some embodiments the amount of CD123 in the sample derived from the subject is assessed by contacting the sample with an antibody, or an antigen-binding fragment thereof, that specifically binds CD123, such as the antibodies described herein. The sample assessed for the presence of CD123 may be derived from a blood sample, a serum sample, circulating cells, circulating tumor cells, cells that are not tissue associated (*i.e.*, free cells), tissues (*e.g.*, surgically resected tumor tissue, biopsies, including fine needle aspiration), histological preparations, and the like.

[00203] In various aspects, the amount of CD123 is determined by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds CD123. In some embodiments, the sample may be contacted by more than one type of antibody, or antigen-binding fragment thereof, that specifically binds CD123. In some embodiments, the sample may be contacted by a first antibody, or antigen-binding fragment thereof, that specifically binds CD123 and then contacted by a second antibody, or antigen-binding fragment thereof, that specifically binds CD123. CD123-specific antibodies or antigen-binding fragments such as those described herein may be used in this capacity.

[00204] Various combinations of the CD123-specific antibodies and antigen-binding fragments can be used to provide a “first” and “second” antibody or antigen-binding fragment to carry out the described diagnostic methods. In some embodiments CD123-expressing cancer includes a hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN).

[00205] In certain embodiments, the amount of CD123 is determined by western blot analysis, radioimmunoassay, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

[00206] In various embodiments of the described diagnostic methods a control or reference sample is used. This sample may be a positive or negative assay control that ensures the assay used is working properly; for example, an assay control of this nature might be commonly used for immunohistochemistry assays. Alternatively, the sample may be a standardized reference for the amount of CD123 in a biological sample from a healthy subject. In some embodiments, the observed CD123 levels of the tested subject may be compared with CD123 levels observed in samples from subjects known to have CD123-expressing cancer. In

some embodiments, the control subject may be afflicted with a particular cancer of interest. In some embodiments, the control subject is known to have early stage cancer, which may or may not be CD123-expressing cancer. In some embodiments, the control subject is known to have intermediate stage cancer, which may or may not be CD123-expressing cancer. In some embodiments, the control subject is known to have late stage, which may or may not be CD123-expressing cancer.

Methods for Monitoring Cancer

[00207] Provided herein are methods for monitoring CD123-expressing cancer in a subject. In some embodiments CD123-expressing cancer includes a hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (BPDCN). In some embodiments the described methods involve assessing whether CD123-expressing cancer is progressing, regressing, or remaining stable by determining the amount of CD123 that is present in a test sample derived from the subject; and comparing the observed amount of CD123 with the amount of CD123 in a biological sample obtained, in a similar manner, from the subject at an earlier point in time, wherein a difference between the amount of CD123 in the test sample and the earlier sample provides an indication of whether the cancer is progressing, regressing, or remaining stable. In this regard, a test sample with an increased amount of CD123, relative to the amount observed for the earlier sample, may indicate progression of a CD123-expressing cancer. Conversely, a test sample with a decreased amount of CD123, relative to the amount observed for the earlier sample, may indicate regression of a CD123-expressing cancer.

[00208] Accordingly, a test sample with an insignificant difference in the amount of CD123, relative to the amount observed for the earlier sample, may indicate a state of stable disease for a CD123-expressing cancer. In some embodiments the amount of CD123 in a biological sample derived from the subject is assessed by contacting the sample with an antibody, or an antibody fragment thereof, that specifically binds CD123, such as the antibodies described herein. The sample assessed for the presence of CD123 may be derived from urine, blood, serum, plasma, saliva, ascites, circulating cells, circulating tumor cells, cells that are not tissue associated (i.e., free cells), tissues (e.g., surgically resected tumor tissue, biopsies,

including fine needle aspiration), histological preparations, and the like. In some embodiments the subject is a human.

[00209] In some embodiments the methods of monitoring a CD123-expressing cancer will involve: contacting a biological sample of a subject with a CD123-specific antibody, or antigen-binding fragment thereof (such as those derivable from the antibodies and fragments provided in Table 1), quantifying the amount of CD123 present in the sample, comparing the amount of CD123 present in the sample to the amount of CD123 determined to be in a biological sample obtained, in a similar manner, from the same subject at an earlier point in time; and determining whether the subject's CD123 level has changed over time. A test sample with an increased amount of CD123, relative to the amount observed for the earlier sample, may indicate progression of cancer. Conversely, a test sample with a decreased amount of CD123, relative to the amount observed for the earlier sample, may indicate regression of a CD123-expressing cancer. Accordingly, a test sample with an insignificant difference in the amount of CD123, relative to the amount observed for the earlier sample, may indicate a state of stable disease for a CD123-expressing cancer. In some embodiments, the CD123 levels of the sample may be compared to a known standard or a reference sample, alone or in addition to the CD123 levels observed for a sample assessed at an earlier point in time. In an additional embodiment, the diagnostic method can be followed with an additional step of administering a cancer-specific treatment. In some embodiments the cancer-specific treatment may be directed against CD123-expressing cancers, such as the CD123 x CD3 multispecific antibodies described herein.

[00210] In various aspects, the amount of CD123 is determined by contacting the sample with an antibody, or antigen-binding fragment thereof, that specifically binds CD123. In some embodiments, the sample may be contacted by more than one type of antibody, or antigen-binding fragment thereof, that specifically binds CD123. In some embodiments, the sample may be contacted by a first antibody, or antigen-binding fragment thereof, that specifically binds CD123 and then contacted by a second antibody, or antigen-binding fragment thereof, that specifically binds CD123. Antibodies such as those described herein may be used in this capacity.

[00211] Various combinations of the antibodies and antigen-binding fragments described in Table 1 can be used to provide a "first" and "second" antibody or antigen-binding fragment to carry out the described monitoring methods. In some embodiments CD123-expressing cancer includes a hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL,

including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN).

[00212] In certain embodiments, the amount of CD123 is determined by western blot analysis, radioimmunoassay, immunofluorimetry, immunoprecipitation, equilibrium dialysis, immunodiffusion, electrochemiluminescence (ECL) immunoassay, immunohistochemistry, fluorescence-activated cell sorting (FACS) or ELISA assay.

Kits for Detecting CD123

[00213] Provided herein are kits for detecting CD123 in a biological sample. These kits include one or more of the CD123-specific antibodies described herein, or an antigen-binding fragment thereof, and instructions for use of the kit.

[00214] The provided CD123-specific antibody, or antigen-binding fragment, may be in solution; lyophilized; affixed to a substrate, carrier, or plate; or detectably labeled.

[00215] The described kits may also include additional components useful for performing the methods described herein. By way of example, the kits may comprise means for obtaining a sample from a subject, a control or reference sample, *e.g.*, a sample from a subject having slowly progressing cancer and/or a subject not having cancer, one or more sample compartments, and/or instructional material which describes performance of a method of the invention and tissue specific controls or standards.

[00216] The means for determining the level of CD123 can further include, for example, buffers or other reagents for use in an assay for determining the level of CD123. The instructions can be, for example, printed instructions for performing the assay and/or instructions for evaluating the level of expression of CD123.

[00217] The described kits may also include means for isolating a sample from a subject. These means can comprise one or more items of equipment or reagents that can be used to obtain a fluid or tissue from a subject. The means for obtaining a sample from a subject may also comprise means for isolating blood components, such as serum, from a blood sample. Preferably, the kit is designed for use with a human subject.

Multispecific Antibodies

[00218] The binding domains of the anti-CD123 antibodies described herein recognize cells expressing CD123 on their surface. As noted above, CD123 expression can be indicative of a cancerous cell. More specific targeting to particular subsets of cells can be achieved by making

bispecific molecules, such as antibodies or antibody fragments, which bind to CD123 and to another target. Examples of such further targets include CD3 and CD33. This is achieved by making a molecule which comprises a first region binding to CD123 and a second binding region binding to the further antigen. The antigen-binding regions can take any form that allows specific recognition of the target, for example the binding region may be or may include a heavy chain variable domain or an Fv (combination of a heavy chain variable domain and a light chain variable domain). Accordingly, bispecific molecules comprising two different antigen-binding regions which bind CD123 and another antigen, respectively, are provided.

[00219] Some of the multispecific antibodies described herein comprise two different antigen-binding regions which bind CD123 and CD3, respectively. In preferred embodiments, multispecific antibodies that bind CD123 and CD3 (CD123 x CD3-multispecific antibodies) and multispecific antigen-binding fragments thereof are provided. In some embodiments, the CD123 x CD3-multispecific antibody comprises a first heavy chain (HC1) and a first light chain (LC1) that pair to form a first antigen-binding site that immunospecifically binds CD123 and a second heavy chain (HC2) and a second light chain (LC2) that pair to form a second antigen-binding site that immunospecifically binds CD3. In preferred embodiments, the CD123 x CD3-multispecific antibody is a bispecific antibody comprising a CD123-specific arm comprising a first heavy chain (HC1) and a first light chain (LC1) that pair to form a first antigen-binding site that immunospecifically binds CD123 and a CD3-specific arm comprising second heavy chain (HC2) and a second light chain (LC2) that pair to form a second antigen-binding site that immunospecifically binds CD3. In some embodiments, the bispecific antibodies of the invention include antibodies having a full length antibody structure. "Full length antibody" as used herein refers to an antibody having two full length antibody heavy chains and two full length antibody light chains. A full length antibody heavy chain (HC) includes heavy chain variable and constant domains VH, CH1, CH2, and CH3. A full length antibody light chain (LC) includes light chain variable and constant domains VL and CL. The full length antibody may be lacking the C-terminal lysine (K) in either one or both heavy chains. The term "Fab-arm" or "half molecule" refers to one heavy chain-light chain pair that specifically binds an antigen.

[00220] The CD123-binding arm of the multispecific antibodies provided herein may be derived from any of the CD123-specific antibodies described above. In some embodiments, the CD123-binding arm binds to an epitope including one or more residues from (i) the segment of CD123 SP2 ECD comprising residues 195 - 202 (RARERVYE (SEQ ID NO: 234)) and/or the segment of CD123 SP2 ECD comprising residues 156-161 (RKFRYE (SEQ ID NO:232)) and/or

or or the segment of CD123 SP2 ECD comprising residues 173 – 178 (TEQVRDR (SEQ ID NO: 233)) or (ii) the segment of CD123 SP2 ECD comprising residues 164 - 175 (IQKRMQPVITEQ (SEQ ID NO: 228)) and/or the segment of CD123 SP2 ECD comprising residues 184-189 (LLNPGT (SEQ ID NO: 229)). In some embodiments, the CD123-binding arm competes for binding to CD123 with a CD123-specific antibody or antigen-binding fragment that binds to an epitope including one or more residues from (i) the segment of CD123 SP2 ECD comprising residues 195 - 202 (RARERVYE (SEQ ID NO: 234)) and/or the segment of CD123 SP2 ECD comprising residues 156-161 (RKFRYE (SEQ ID NO: 232)) and/or the segment of CD123 SP2 ECD comprising residues 173 – 178 (TEQVRDR (SEQ ID NO: 233)) or (ii) the segment of CD123 SP2 ECD comprising residues 164 - 175 (IQKRMQPVITEQ (SEQ ID NO: 228)) and/or the segment of CD123 SP2 ECD comprising residues 184-189 (LLNPGT (SEQ ID NO: 229)). CD123-binding arms binding to at least one residue in these epitopes may also bind to additional residues in the CD123 ECD. In some embodiments, the CD123-binding arm is neutralizing. A neutralizing CD123-binding arm includes those that are capable of inhibiting the binding of IL-3 to CD123 as determined by measuring the decrease in STAT5 phosphorylation upon stimulation of TF-1 cells with rhIL-3. In some embodiments of the bispecific antibodies, the CD123-binding arm binds human CD123 SP1, preferably the extracellular domain thereof.

[00221] In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises a heavy chain CDR1, CDR2, and CDR3 derived from an antibody clone as described in Table 1. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain CDR1, CDR2, and CDR3 and light chain CDR1, CDR2, and CDR3 derived from an antibody clone as described in Table 1. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain CDR1, CDR2, and CDR3 of clone I3RB1, I3RB2, I3RB5, I3RB6, I3RB7, I3RB8, I3RB9, I3RB11, I3RB12, I3RB16, I3RB17, I3RB18, I3RB19, I3RB20, I3RB21, I3RB22, I3RB24, I3RB28, I3RB29, I3RB30, I3RB32, I3RB33, I3RB34, I3RB35, I3RB36, I3RB37, I3RB38, I3RB40, or I3RB47. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain CDR1, CDR2, and CDR3 and light chain CDR1, CDR2, and CDR3 of clone I3RB1, I3RB2, I3RB5, I3RB6, I3RB7, I3RB8, I3RB9, I3RB11, I3RB12, I3RB16, I3RB17, I3RB18, I3RB19, I3RB20, I3RB21, I3RB22, I3RB24, I3RB28, I3RB29, I3RB30, I3RB32, I3RB33, I3RB34, I3RB35, I3RB36, I3RB37, I3RB38, I3RB40, or I3RB47. In some exemplary embodiments of such CD123 SP1-binding arms, the

first antigen-binding region which binds CD123 comprises a heavy chain variable domain derived from an antibody clone as described in Table 1. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain variable domain and light chain variable domain derived from an antibody clone as described in Table 1. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain variable domain of clone I3RB1, I3RB2, I3RB5, I3RB6, I3RB7, I3RB8, I3RB9, I3RB11, I3RB12, I3RB16, I3RB17, I3RB18, I3RB19, I3RB20, I3RB21, I3RB22, I3RB24, I3RB28, I3RB29, I3RB30, I3RB32, I3RB33, I3RB34, I3RB35, I3RB36, I3RB37, I3RB38, I3RB40, or I3RB47. In some exemplary embodiments of such CD123 SP1-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain variable domain and light chain variable domain of clone I3RB1, I3RB2, I3RB5, I3RB6, I3RB7, I3RB8, I3RB9, I3RB11, I3RB12, I3RB16, I3RB17, I3RB18, I3RB19, I3RB20, I3RB21, I3RB22, I3RB24, I3RB28, I3RB29, I3RB30, I3RB32, I3RB33, I3RB34, I3RB35, I3RB36, I3RB37, I3RB38, I3RB40, or I3RB47.

[00222] In some embodiments of the bispecific antibodies, the CD123-binding arm binds human CD123 SP2, preferably the extracellular domain thereof. In preferred embodiments of the bispecific antibodies, the CD123-binding arm binds human CD123 SP1 and human CD123 SP2, and more preferably the extracellular domains thereof. In some exemplary embodiments of such CD123 SP2-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain CDR1, CDR2, and CDR3 of clone I3RB1, I3RB2, I3RB5, I3RB18, I3RB19, or I3RB30. In some exemplary embodiments of such CD123 SP2-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain CDR1, CDR2, and CDR3 and light chain CDR1, CDR2, and CDR3 of clone I3RB1, I3RB2, I3RB5, I3RB18, I3RB19, or I3RB30. In some exemplary embodiments of such CD123 SP2-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain variable domain of clone I3RB1, I3RB2, I3RB5, I3RB18, I3RB19, or I3RB30. In some exemplary embodiments of such CD123 SP2-binding arms, the first antigen-binding region which binds CD123 comprises heavy chain variable domain and light chain variable domain of clone I3RB1, I3RB2, I3RB5, I3RB18, I3RB19, or I3RB30.

[00223] In some embodiments of the bispecific antibodies, the CD123-binding arm also binds cynomolgus CD123, preferably the extracellular domain thereof.

[00224] In some embodiments of the bispecific antibodies, the CD123-binding arm is derived from a CD123-specific antibody that competes for binding to CD123 with antibody

clone I3RB2, I3RB60, I3RB70, I3RB79, or I3RB118. In some embodiments of the bispecific antibodies, the CD123-binding arm is derived from a CD123-specific antibody that competes for binding to CD123 with antibody clone I3RB18, I3RB49, or I3RB55. Competition for binding can be determined using a competition binding ELISA, in line with the technique described below in Example 5. Competitive binding may be determined by detecting at least 20% inhibition of the binding of a first antibody by a second antibody, irrespective of the order in which the antibodies are bound to CD123 (*i.e.* if when antibody A is bound to CD123 before antibody B, only 10% inhibition is observed, but when antibody B is bound to CD123 before antibody A, 30% inhibition is observed, then because greater than 20% inhibition has been observed in one of the experiments, competitive binding may be concluded).

[00225] In some embodiments, the CD123-binding arm of the multispecific antibody is IgG, or a derivative thereof, e.g., IgG1, IgG2, IgG3, and IgG4 isotypes. In some embodiments wherein the CD123-binding arm has an IgG1 isotype, it contains L234A, L235A, and K409R substitution(s) in its Fc region. In some embodiments wherein the CD123-binding arm has an IgG4 isotype, it contains S228P, L234A, and L235A substitution(s) in its Fc region.

[00226] In some embodiments of the bispecific antibodies, the second antigen-binding arm binds human CD3. In some preferred embodiments, the CD3-specific arm of the CD123 x CD3 bispecific antibody is derived from a CD3-specific antibody that binds and activates human primary T cells and/or cynomolgus monkey primary T cells. In some embodiments, the CD3-binding arm binds to an epitope at the N-terminus of CD3 ϵ . In some embodiments, the CD3-binding arm contacts an epitope including the six N-terminal amino acids of CD3 ϵ . In some embodiments, the CD3-specific binding arm of the bispecific antibody is derived from the mouse monoclonal antibody SP34, a mouse IgG3/lambda isotype. In some embodiments, the CD3-binding arm comprises the CDRs of antibody SP34. Such CD3-binding arms may bind to CD3 with an affinity of 5×10^{-7} M or less, such as 1×10^{-7} M or less, 5×10^{-8} M or less, 1×10^{-8} M or less, 5×10^{-9} M or less, or 1×10^{-9} M or less. The CD3-specific binding arm may be a humanized version of an arm of mouse monoclonal antibody SP34. Human framework adaptation (HFA) may be used to humanize the anti-CD3 antibody from which the CD3-specific arm is derived. In some embodiments of the bispecific antibodies, the CD3-binding arm comprises a heavy chain and light chain pair selected from Table 2. In some embodiments, the CD3-binding arm of the CD123 x CD3 bispecific antibody is derived from Table 3.

[00227] In some embodiments, the CD3-binding arm is IgG, or a derivative thereof. In some embodiments, the CD3-binding arm is IgG1, IgG2, IgG3, or IgG4. In some embodiments

wherein the CD3-binding arm has an IgG1 isotype, it contains L234A, L235A, and F405L substitution(s) in its Fc region. In some embodiments wherein the CD3-binding arm has an IgG4 isotype, it contains S228P, L234A, L235A, F405L, and R409K substitution(s) in its Fc region. In some embodiments, the antibodies or antigen-binding fragments are IgG-AA Fc. In some embodiments, the antibodies or antigen-binding fragments are IgG-AA Fc-L234A, L235A, and F405L. In some embodiments, the antibodies or antigen-binding fragments bind CD3 ϵ on primary human T cells. In some embodiments, the antibodies or antigen-binding fragments bind CD3 ϵ on primary cynomolgus T cells. In some embodiments, the antibodies or antigen-binding fragments bind CD3 ϵ on primary human and cynomolgus T cells. In some embodiments, the antibodies or antigen-binding fragments activate primary human CD4⁺ T cells. In some embodiments, the antibodies or antigen-binding fragments activate primary cynomolgus CD4⁺ T cells.

[00228] In some embodiments are provided a CD123 x CD3 bispecific antibody having a CD123-binding arm comprising a heavy chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466. In some embodiments are provided a CD123 x CD3 bispecific antibody having a CD123-binding arm comprising a heavy chain and light chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466. In some embodiments are provided a CD123 x CD3 bispecific antibody having a CD3-binding arm comprising a heavy chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466. In some embodiments are provided a CD123 x CD3 bispecific antibody having a CD3-binding arm comprising a heavy chain and light chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466. In some embodiments are provided a CD123 x CD3 bispecific antibody having a CD123-binding arm comprising a heavy chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, mAb 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466 and a CD3-binding arm comprising a heavy chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, AbB 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466. In some

embodiments are provided a CD123 x CD3 bispecific antibody having a CD123-binding arm comprising a heavy chain and light chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466 and a CD3-binding arm comprising a heavy chain and light chain of antibody clone I3RB179, I3RB180, I3RB181, I3RB182, I3RB183, I3RB186, I3RB187, I3RB188, I3RB189, CD3B191, Ab 7959, Ab3978, Ab 7955, Ab 9958, Ab 8747, Ab 8876, Ab 4435, or Ab 5466.

[00229] Preferred CD123 x CD3 bispecific antibodies are provided in Tables 13 and 17.

[00230] Different formats of bispecific antibodies have been described and were recently reviewed by Chames and Baty (2009) *Curr Opin Drug Disc Dev* 12: 276.

[00231] In some embodiments, the bispecific antibody of the present invention is a diabody, a cross-body, or a bispecific antibody obtained via a controlled Fab arm exchange as those described in the present invention.

[00232] In some embodiments, the bispecific antibodies include IgG-like molecules with complementary CH3 domains to force heterodimerisation; recombinant IgG-like dual targeting molecules, wherein the two sides of the molecule each contain the Fab fragment or part of the Fab fragment of at least two different antibodies; IgG fusion molecules, wherein full length IgG antibodies are fused to an extra Fab fragment or parts of Fab fragment; Fc fusion molecules, wherein single chain Fv molecules or stabilized diabodies are fused to heavy-chain constant-domains, Fc-regions or parts thereof; Fab fusion molecules, wherein different Fab-fragments are fused together; ScFv- and diabody-based and heavy chain antibodies (e.g., domain antibodies, nanobodies) wherein different single chain Fv molecules or different diabodies or different heavy-chain antibodies (e.g. domain antibodies, nanobodies) are fused to each other or to another protein or carrier molecule.

[00233] In some embodiments, IgG-like molecules with complementary CH3 domains molecules include the Triomab/Quadroma (Trion Pharma/Fresenius Biotech), the Knobs-into-Holes (Genentech), CrossMAbs (Roche) and the electrostatically-matched (Amgen), the LUZ-Y (Genentech), the Strand Exchange Engineered Domain body (SEEDbody)(EMD Serono), the Biclonic (Merus) and the DuoBody (Genmab A/S).

[00234] In some embodiments, recombinant IgG-like dual targeting molecules include Dual Targeting (DT)-Ig (GSK/Domantis), Two-in-one Antibody (Genentech), Cross-linked Mabs (Karmanos Cancer Center), mAb2 (F-Star) and CovX-body (CovX/Pfizer).

[00235] In some embodiments, IgG fusion molecules include Dual Variable Domain (DVD)-Ig (Abbott), IgG-like Bispecific (InnClone/Eli Lilly), Ts2Ab (MedImmune/AZ) and BsAb (Zymogenetics), HERCULES (Biogen Idec) and TvAb (Roche).

[00236] In some embodiments, Fc fusion molecules include to ScFv/Fc Fusions (Academic Institution), SCORPION (Emergent BioSolutions/Trubion, Zymogenetics/BMS), Dual Affinity Retargeting Technology (Fc-DART) (MacroGenics) and Dual(ScFv).sub.2-Fab (National Research Center for Antibody Medicine--China).

[00237] In some embodiments, Fab fusion bispecific antibodies include F(ab)2 (Medarex/AMGEN), Dual-Action or Bis-Fab (Genentech), Dock-and-Lock (DNL) (ImmunoMedics), Bivalent Bispecific (Biotechnol) and Fab-Fv (UCB-Celltech). ScFv-, diabody-based and domain antibodies include but are not limited to Bispecific T Cell Engager (BITE) (Micromet), Tandem Diabody (Tandab) (Affimed), Dual Affinity Retargeting Technology (DART) (MacroGenics), Single-chain Diabody (Academic), TCR-like Antibodies (AIT, ReceptorLogics), Human Serum Albumin ScFv Fusion (Merrimack) and COMBODY (Epigen Biotech), dual targeting nanobodies (Ablynx), dual targeting heavy chain only domain antibodies.

[00238] Full length bispecific antibodies of the invention may be generated for example using Fab arm exchange (or half molecule exchange) between two mono specific bivalent antibodies by introducing substitutions at the heavy chain CH3 interface in each half molecule to favor heterodimer formation of two antibody half molecules having distinct specificity either in vitro in cell-free environment or using co-expression. The Fab arm exchange reaction is the result of a disulfide-bond isomerization reaction and dissociation-association of CH3 domains. The heavy-chain disulfide bonds in the hinge regions of the parent mono specific antibodies are reduced. The resulting free cysteines of one of the parent monospecific antibodies form an inter heavy-chain disulfide bond with cysteine residues of a second parent mono specific antibody molecule and simultaneously CH3 domains of the parent antibodies release and reform by dissociation-association. The CH3 domains of the Fab arms may be engineered to favor heterodimerization over homodimerization. The resulting product is a bispecific antibody having two Fab arms or half molecules which each bind a distinct epitope, i.e. an epitope on CD123 (IL3-R α) and an epitope on CD3.

[00239] "Homodimerization" as used herein refers to an interaction of two heavy chains having identical CH3 amino acid sequences. "Homodimer" as used herein refers to an antibody having two heavy chains with identical CH3 amino acid sequences.

[00240] "Heterodimerization" as used herein refers to an interaction of two heavy chains having non-identical CH3 amino acid sequences. "Heterodimer" as used herein refers to an antibody having two heavy chains with non-identical CH3 amino acid sequences.

[00241] The "knob-in-hole" strategy (see, e.g., PCT Int. Publ. No. WO 2006/028936) may be used to generate full length bispecific antibodies. Briefly, selected amino acids forming the interface of the CH3 domains in human IgG can be mutated at positions affecting CH3 domain interactions to promote heterodimer formation. An amino acid with a small side chain (hole) is introduced into a heavy chain of an antibody specifically binding a first antigen and an amino acid with a large side chain (knob) is introduced into a heavy chain of an antibody specifically binding a second antigen. After co-expression of the two antibodies, a heterodimer is formed as a result of the preferential interaction of the heavy chain with a "hole" with the heavy chain with a "knob". Exemplary CH3 substitution pairs forming a knob and a hole are (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain): T366Y/F405A, T366W/F405W, F405W/Y407A, T394W/Y407T, T394S/Y407A, T366W/T394S, F405W/T394S and T366W/T366S_L368A_Y407V.

[00242] Other strategies such as promoting heavy chain heterodimerization using electrostatic interactions by substituting positively charged residues at one CH3 surface and negatively charged residues at a second CH3 surface may be used, as described in US Pat. Publ. No. US2010/0015133; US Pat. Publ. No. US2009/0182127; US Pat. Publ. No. US2010/028637 or US Pat. Publ. No. US2011/0123532. In other strategies, heterodimerization may be promoted by the following substitutions (expressed as modified position in the first CH3 domain of the first heavy chain/modified position in the second CH3 domain of the second heavy chain): L351Y_F405AY407V/T394W, T366I_K392M_T394W/F405A_Y407V, T366L_K392M_T394W/F405A_Y407V, L351Y_Y407A/T366A_K409F, L351Y_Y407A/T366V_K409F_Y407A/T366A_K409F, or T350V_L351Y_F405A_Y407V/T350V_T366L_K392L_T394W as described in U.S. Pat. Publ. No. US2012/0149876 or U.S. Pat. Publ. No. US2013/0195849.

[00243] In addition to methods described above, bispecific antibodies of the invention may be generated in vitro in a cell-free environment by introducing asymmetrical mutations in the CH3 regions of two mono specific homodimeric antibodies and forming the bispecific heterodimeric antibody from two parent monospecific homodimeric antibodies in reducing conditions to allow disulfide bond isomerization according to methods described in Int. Pat.

Publ. No. W02011/131746. In the methods, the first monospecific bivalent antibody (e.g., anti-CD123 (IL3-R α) antibody) and the second monospecific bivalent antibody (e.g., anti-CD3 antibody) are engineered to have certain substitutions at the CH3 domain that promotes heterodimer stability; the antibodies are incubated together under reducing conditions sufficient to allow the cysteines in the hinge region to undergo disulfide bond isomerization; thereby generating the bispecific antibody by Fab arm exchange. The incubation conditions may optimally be restored to non-reducing conditions. Exemplary reducing agents that may be used are 2-mercaptoethylamine (2-MEA), dithiothreitol (DTT), dithioerythritol (DTE), glutathione, tris (2-carboxyethyl)phosphine (TCEP), L-cysteine and beta-mercaptoethanol, preferably a reducing agent selected from the group consisting of: 2-mercaptoethylamine, dithiothreitol and tris (2-carboxyethyl)phosphine. For example, incubation for at least 90 min at a temperature of at least 20° C in the presence of at least 25 mM 2-MEA or in the presence of at least 0.5 mM dithiothreitol at a pH from 5-8, for example at pH of 7.0 or at pH of 7.4 may be used.

[00244] In addition to the described CD123 x CD3-multispecific antibodies, also provided are polynucleotide sequences capable of encoding the described CD123 x CD3-multispecific antibodies. Vectors comprising the described polynucleotides are also provided, as are cells expressing the CD123 x CD3-multispecific antibodies provided herein. Also described are cells capable of expressing the disclosed vectors. These cells may be mammalian cells (such as 293F cells, CHO cells), insect cells (such as Sf7 cells), yeast cells, plant cells, or bacteria cells (such as *E. coli*). The described antibodies may also be produced by hybridoma cells.

Therapeutic composition and methods of treatment using multispecific antibodies and multispecific antigen-binding fragments thereof

[00245] The CD123 bispecific antibodies discussed above, for example the CD123 x CD3 bispecific antibodies discussed above, are useful in therapy. In particular, the CD123 bispecific antibodies are useful in treating cancer. Also provided herein are therapeutic compositions for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a multispecific antibody or multispecific antigen-binding fragment described herein and a pharmaceutically acceptable carrier. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. In one embodiment said pharmaceutical composition is for the treatment of a CD123-expressing

cancer, including (but not limited to) the following: CD123-expressing hematological cancers, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN); and other cancers yet to be determined in which CD123 is expressed. Particular bispecific antibodies that may be used to treat cancer, such as hematological cancer, including the specific cancers discussed above, include antibodies 7959, 3978, 7955, 9958, 8747, 4435, and 5466. One example of a useful bispecific antibody for treating cancer, such as hematological cancer, including these specific cancers is antibody 9958. Another example of a useful bispecific antibody for treating cancer, such as hematological cancer, including these specific cancers is antibody 3978. Another example of a useful bispecific antibody for treating cancer, such as hematological cancer, including these specific cancers is antibody 8747. Another example of a useful bispecific antibody for treating cancer, such as hematological cancer, including these specific cancers is antibody 7959.

[00246] The pharmaceutical compositions provided herein comprise: a) an effective amount of a multispecific antibody or antibody fragment of the present invention, and b) a pharmaceutically acceptable carrier, which may be inert or physiologically active. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. As used herein, the term "pharmaceutically acceptable carriers" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, and the like that are physiologically compatible. Examples of suitable carriers, diluents and/or excipients include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, and the like, as well as any combination thereof. In many cases, it will be preferable to include isotonic agents, such as sugars, polyalcohols, or sodium chloride in the composition. In particular, relevant examples of suitable carrier include: (1) Dulbecco's phosphate buffered saline, pH.about.7.4, containing or not containing about 1 mg/mL to 25 mg/mL human serum albumin, (2) 0.9% saline (0.9% w/v sodium chloride (NaCl)), and (3) 5% (w/v) dextrose; and may also contain an antioxidant such as tryptamine and a stabilizing agent such as Tween 20 ®.

[00247] The compositions herein may also contain a further therapeutic agent, as necessary for the particular disorder being treated. Preferably, the multispecific antibody or antibody fragment and the supplementary active compound will have complementary activities

that do not adversely affect each other. In a preferred embodiment, the further therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. In a preferred embodiment, the further therapeutic agent is a chemotherapeutic agent.

[00248] The compositions of the invention may be in a variety of forms. These include for example liquid, semi-solid, and solid dosage forms, but the preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions. The preferred mode of administration is parenteral (e.g. intravenous, intramuscular, intraperitoneal, subcutaneous). In a preferred embodiment, the compositions of the invention are administered intravenously as a bolus or by continuous infusion over a period of time. In another preferred embodiment, they are injected by intramuscular, subcutaneous, intra-articular, intrasynovial, intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects.

[00249] Sterile compositions for parenteral administration can be prepared by incorporating the antibody, antibody fragment or antibody conjugate of the present invention in the required amount in the appropriate solvent, followed by sterilization by microfiltration. As solvent or vehicle, there may be used water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, and the like, as well as combination thereof. In many cases, it will be preferable to include isotonic agents, such as sugars, polyalcohols, or sodium chloride in the composition. These compositions may also contain adjuvants, in particular wetting, isotonizing, emulsifying, dispersing and stabilizing agents. Sterile compositions for parenteral administration may also be prepared in the form of sterile solid compositions which may be dissolved at the time of use in sterile water or any other injectable sterile medium.

[00250] The multispecific antibody or antibody fragment may also be orally administered. As solid compositions for oral administration, tablets, pills, powders (gelatine capsules, sachets) or granules may be used. In these compositions, the active ingredient according to the invention is mixed with one or more inert diluents, such as starch, cellulose, sucrose, lactose or silica, under an argon stream. These compositions may also comprise substances other than diluents, for example one or more lubricants such as magnesium stearate or talc, a coloring, a coating (sugar-coated tablet) or a glaze.

[00251] As liquid compositions for oral administration, there may be used pharmaceutically acceptable solutions, suspensions, emulsions, syrups and elixirs containing inert diluents such as water, ethanol, glycerol, vegetable oils or paraffin oil. These compositions

may comprise substances other than diluents, for example wetting, sweetening, thickening, flavoring or stabilizing products.

[00252] The doses depend on the desired effect, the duration of the treatment and the route of administration used; they are generally between 5 mg and 1000 mg per day orally for an adult with unit doses ranging from 1 mg to 250 mg of active substance. In general, the doctor will determine the appropriate dosage depending on the age, weight and any other factors specific to the subject to be treated.

[00253] Also provided herein are methods for killing a CD123+ cell by administering to a patient in need thereof a multispecific antibody which binds said CD123 and is able to recruit T cells to kill said CD123+ cell (i.e., T cell redirection). Any of the multispecific antibodies or antibody fragments of the invention may be used therapeutically. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof.

[00254] In a preferred embodiment, multispecific antibodies or antibody fragments of the invention are used for the treatment of a hyperproliferative disorder in a mammal. In a more preferred embodiment, one of the pharmaceutical compositions disclosed above, and which contains a multispecific antibody or antibody fragment of the invention, is used for the treatment of a hyperproliferative disorder in a mammal. In one embodiment, the disorder is a cancer. In particular, the cancer is a CD123-expressing cancer, including (but not limited to) the following: CD123-expressing hematological cancers, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN); and other cancers yet to be determined in which CD123 is expressed. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof.

[00255] Accordingly, the pharmaceutical compositions of the invention are useful in the treatment or prevention of a variety of cancers, including (but not limited to) the following: a CD123-expressing cancer, including (but not limited to) the following: CD123-expressing hematological cancers, such as acute myeloid leukemia (AML), myelodysplastic syndrome

(MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (DPDCN); and other cancers yet to be determined in which CD123 is expressed.

[00256] Similarly, further provided herein is a method for inhibiting the growth of selected cell populations comprising contacting CD123-expressing target cells, or tissue containing such target cells, with an effective amount of a multispecific antibody or antibody fragment of the present invention, either alone or in combination with other cytotoxic or therapeutic agents, in the presence of a peripheral blood mononuclear cell (PBMC). In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. In a preferred embodiment, the further therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. In a preferred embodiment, the further therapeutic agent is a chemotherapeutic agent. The method for inhibiting the growth of selected cell populations can be practiced *in vitro*, *in vivo*, or *ex vivo*.

[00257] Examples of *in vitro* uses include treatments of autologous bone marrow prior to their transplant into the same patient in order to kill diseased or malignant cells; treatments of bone marrow prior to its transplantation in order to kill competent T cells and prevent graft-versus-host-disease (GVHD); treatments of cell cultures in order to kill all cells except for desired variants that do not express the target antigen; or to kill variants that express undesired antigen. The conditions of non-clinical *in vitro* use are readily determined by one of ordinary skill in the art.

[00258] Examples of clinical *ex vivo* use are to remove tumor cells from bone marrow prior to autologous transplantation in cancer treatment. Treatment can be carried out as follows. Bone marrow is harvested from the patient or other individual and then incubated in medium containing serum to which is added the cytotoxic agent of the invention. Concentrations range from about 10 μ M to 1 μ M, for about 30 min to about 48 hr at about 37 °C. The exact conditions of concentration and time of incubation, i.e., the dose, are readily determined by one of ordinary skill in the art. After incubation the bone marrow cells are washed with medium containing serum and returned to the patient by i.v. infusion according to known methods. In circumstances where the patient receives other treatment such as a course of ablative chemotherapy or total-

body irradiation between the time of harvest of the marrow and reinfusion of the treated cells, the treated marrow cells are stored frozen in liquid nitrogen using standard medical equipment.

[00259] For clinical *in vivo* use, a therapeutically effective amount of the multispecific antibody or antigen-binding fragment is administered to a subject in need thereof. For example, the CD123 x CD3-multispecific antibodies and multispecific antigen-binding fragments thereof may be useful in the treatment of a CD123-expressing cancer in a subject in need thereof. In some embodiments, the CD123-expressing cancer is a hematological cancer, such as acute myeloid leukemia (AML), myelodysplastic syndrome (MDS, low or high risk), acute lymphocytic leukemia (ALL, including all subtypes), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML), or blastic plasmacytoid dendritic cell neoplasm (BPDCN). In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. In some embodiments, the subject is a mammal, preferably a human. In some embodiments, the multispecific antibody or antigen-binding fragment will be administered as a solution that has been tested for sterility.

[00260] Dosage regimens in the above methods of treatment and uses are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage.

[00261] The efficient dosages and the dosage regimens for the multispecific antibodies and fragments depend on the disease or condition to be treated and may be determined by one skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of a compound of the present invention is about 0.001-10 mg/kg, such as about 0.001-5 mg/kg, for example about 0.001-2 mg/kg, such as about 0.001-1 mg/kg, for instance about 0.001, about 0.01, about 0.1, about 1 or about 10 mg/kg.

[00262] A physician or veterinarian having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the multispecific antibody or fragment employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is

achieved. In general, a suitable daily dose of a bispecific antibody of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Administration may e.g. be parenteral, such as intravenous, intramuscular or subcutaneous. In one embodiment, the multispecific antibody or fragment may be administered by infusion in a weekly dosage of calculated by mg/m^2 . Such dosages can, for example, be based on the mg/kg dosages provided above according to the following: $\text{dose (mg/kg)} \times 70$: 1.8. Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hr, such as of from 2 to 12 hr. In one embodiment, the multispecific antibody or fragment may be administered by slow continuous infusion over a long period, such as more than 24 hours, in order to reduce toxic side effects.

[00263] In one embodiment, the multispecific antibody or fragment may be administered in a weekly dosage of calculated as a fixed dose for up to eight times, such as from four to six times when given once a week. Such regimen may be repeated one or more times as necessary, for example, after six months or twelve months. Such fixed dosages can, for example, be based on the mg/kg dosages provided above, with a body weight estimate of 70 kg. The dosage may be determined or adjusted by measuring the amount of bispecific antibody of the present invention in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the CD123 antigen binding region of the multispecific antibodies of the present invention.

[00264] In one embodiment, the multispecific antibody or fragment may be administered by maintenance therapy, such as, e.g., once a week for a period of six months or more.

[00265] A multispecific antibody or fragment may also be administered prophylactically in order to reduce the risk of developing cancer, delay the onset of the occurrence of an event in cancer progression, and/or reduce the risk of recurrence when a cancer is in remission.

[00266] The multispecific antibodies and fragments thereof as described herein may also be administered in combination therapy, i.e., combined with other therapeutic agents relevant for the disease or condition to be treated. Accordingly, in one embodiment, the antibody-containing medicament is for combination with one or more further therapeutic agent, such as a chemotherapeutic agent. In some embodiments, the other therapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2. Such combined

administration may be simultaneous, separate or sequential, in any order. For simultaneous administration the agents may be administered as one composition or as separate compositions, as appropriate.

[00267] In one embodiment, a method for treating a disorder involving cells expressing CD123 in a subject, which method comprises administration of a therapeutically effective amount of a multispecific antibody or fragment, such as a CD123 x CD3 bispecific antibody described herein, and radiotherapy to a subject in need thereof is provided. In one embodiment is provided a method for treating or preventing cancer, which method comprises administration of a therapeutically effective amount of a multispecific antibody or fragment, such as a CD123 x CD3 antibody described herein, and radiotherapy to a subject in need thereof. Radiotherapy may comprise radiation or associated administration of radiopharmaceuticals to a patient is provided. The source of radiation may be either external or internal to the patient being treated (radiation treatment may, for example, be in the form of external beam radiation therapy (EBRT) or brachytherapy (BT)). Radioactive elements that may be used in practicing such methods include, e.g., radium, cesium-137, iridium-192, americium-241, gold-198, cobalt-57, copper-67, technetium-99, iodide-123, iodide-131, and indium-111.

Kits

[00268] Also provided herein are includes kits, e.g., comprising a described multispecific antibody or antigen-binding fragment thereof and instructions for the use of the antibody or fragment for killing of particular cell types. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. The instructions may include directions for using the multispecific antibody or antigen-binding fragment thereof in vitro, in vivo or ex vivo.

[00269] Typically, the kit will have a compartment containing the multispecific antibody or antigen-binding fragment thereof. The multispecific antibody or antigen-binding fragment thereof may be in a lyophilized form, liquid form, or other form amendable to being included in a kit. The kit may also contain additional elements needed to practice the method described on the instructions in the kit, such a sterilized solution for reconstituting a lyophilized powder, additional agents for combining with the multispecific antibody or antigen-binding

fragment thereof prior to administering to a patient, and tools that aid in administering the multispecific antibody or antigen-binding fragment thereof to a patient.

Diagnostic Uses

[00270] The multispecific antibodies and fragments described herein may also be used for diagnostic purposes. Thus, also provided are diagnostic compositions comprising a multispecific antibody or fragments as defined herein, and to its use. In preferred embodiments, the multispecific antibody is a CD123 x CD3-multispecific antibody as described herein, or a multispecific antigen-binding fragment thereof, and more preferably a CD123 x CD3-bispecific antibody as described herein, or a CD123 x CD3-bispecific antigen-binding fragment thereof. In one embodiment, the present invention provides a kit for diagnosis of cancer comprising a container comprising a bispecific CD123 x CD3 antibody, and one or more reagents for detecting binding of the antibody to CD123. Reagents may include, for example, fluorescent tags, enzymatic tags, or other detectable tags. The reagents may also include secondary or tertiary antibodies or reagents for enzymatic reactions, wherein the enzymatic reactions produce a product that may be visualized. For example, the multispecific antibodies described herein, or antigen-binding fragments thereof, may be labeled with a radiolabel, a fluorescent label, an epitope tag, biotin, a chromophore label, an ECL label, an enzyme, ruthenium, ¹¹¹In-DOTA, ¹¹¹In- diethylenetriaminepentaacetic acid (DTPA), horseradish peroxidase, alkaline phosphatase and beta-galactosidase, or poly-histidine or similar such labels known in the art.

[00271] The following examples are provided to supplement the prior disclosure and to provide a better understanding of the subject matter described herein. These examples should not be considered to limit the described subject matter. It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be apparent to persons skilled in the art and are to be included within, and can be made without departing from, the true scope of the invention.

Example 1: Materials

Generation of CD123 cell lines

[00272] A set of pDisplay™ vectors presenting human CD123 SP1 ECD (amino acids 20 - 305) (SEQ ID NO:1), human CD123 SP2 ECD (amino acids 19 - 227 of SEQ ID NO:2), and cyno CD123 ECD (amino acid 19 – 305 of SEQ ID NO:3) were generated for use as screening tools to assess the anti-CD123 leads. A mammalian expression vector that allows display of

proteins on the cell surface, pDisplay (Invitrogen) was used (Figure 1). Proteins expressed from pDisplay™ are fused at the N-terminus to the murine Ig κ-chain leader sequence, which directs the protein to the secretory pathway, and at the C-terminus to the platelet derived growth factor receptor (PDGFR) transmembrane domain, which anchors the protein to the plasma membrane, displaying it on the extracellular side. Recombinant proteins expressed from pDisplay™ contain the hemagglutinin A and myc epitopes for detection by western blot or immunofluorescence. The CMV promoter drives expression.

[00273] Vectors were transiently transfected into HEK293T cells using standard methods. Transfected 293F adherent cells were selected for stable plasmid integration, then single cell sorted and the CD123 surface receptor expression was quantified by FACS using the BangsLabs Quantum FITC-5kit (Catalog #855, Bangs Laboratories, Inc). A set of 10 single cell clones for each cell line were selected for screening, and quantified for CD123 ECD expression. The cell lines used for subsequent hit screening had surface expression of approximately 500,000 CD123 ECD copies per cell.

Generation of Soluble CD123 ECD Protein

[00274] Recombinant human CD123 SP1 ECD-His tag protein (Lot #LV081110A), corresponding to amino acid 20 to 305 of CD123 SP1 (SEQ ID NO:1) was obtained from R&D Systems (#301-R3/CF) for use in phage panning and hit screening. The protein was tested for endotoxin prior to use and was biotinylated for phage panning studies. This material was also used for binding and affinity measurements.

[00275] Recombinant human CD123 SP2 ECD protein corresponding to amino acids 18-225 of human CD123 SP2 (SEQ ID NO: 2) was purified for use in binding and affinity measurements. cDNA was prepared using gene synthesis techniques (U.S. Pat. No. 6,670,127; U.S. Pat. No. 6,521,427). Plasmids for expression of the synthetic soluble CD123 ECD SP2 were prepared using standard molecular biology techniques. The CD123 ECD SP2 gene fragment with an N-terminal gp67 signal sequence and a c-terminal 6-His tag was cloned into the Eco RI and Not I sites of pFastbac1 (Invitrogen) and expressed with the Bac to Bac system (Invitrogen) in High Five Cells (Invitrogen). The secreted protein (SEQ ID NO: 226) was purified through HisTrap (GE) and Superdex 75 (GE) columns. This material was used for binding and affinity measurements and epitope mapping.

[00276] The soluble CD123 ECD proteins were biotinylated using the SureLink Biotinylation Kit (KPL #86-00-01) as per the manufacturer's instructions. Proteins were run on SDS/PAGE to confirm monomeric state.

[00277]

Anti-CD3 antibody for x-ray crystallography

[00278] SP34 mAb, mouse IgG3/lambda isotype, was purchased from BD Biosciences Pharmingen (San Diego, CA), Cat. No. 556611 and comprising the Light and Heavy chains shown in SEQ ID NOs: 4 and 5, respectively.

Example 2: Identification of Anti-human CD123 mAbs

[00279] Solution panning of the de novo Human Fab-pIX libraries [Shi, L., et al J Mol Biol, 2010. 397(2): p. 385-396. WO 2009/085462], consisting of VH1-69, 3-23 and 5-51 heavy chain libraries paired with Vk1-39, 3-11, 3-20 and 4-1 light chain libraries, was performed using a biotinylated antigen-streptavidin magnetic bead capture method as described (Rothe et al., J. Mol. Biol. 376:1182-1200, 2008; Steidl et al., Mol. Immunol. 46: 135-144, 2008) in four subsequent rounds.

[00280] The pIX gene was excised from phagemid DNA following the fourth round of panning to generate soluble his-tagged Fab coding regions. Fabs were expressed in E. coli and screened for binding to recombinant human CD123 SP1 ECD-His tag protein in an ELISA. Briefly, 96-well Nunc Maxisorp plates (Nunc #437111) were coated with sheep anti-human Fd (The Binding Site #PC075) in PBS at 1 µg/mL overnight at 4°C. Bacterial colonies containing the Fab expression vector were grown in 450 µL of 2xYT (Carbenecillin) in deep-well culture plates until turbid (OD600 ≈ 0.6). Fab expression was induced by the addition of IPTG to a concentration of 1 mM. Cultures were grown overnight at 30°C and then clarified by centrifugation. Anti-Fd coated Maxisorp plates were washed once with TBS, 0.5% Tween-20 (Sigma #79039-10PAK) and blocked with 200 µL PBS-Tween (0.5%) + nonfat dried milk (3%) per well for one hr at room temperature. At this step and all subsequent steps plates are washed three times with TBS, 0.5% Tween-20 (Sigma #79039-10PAK). Each well received 50 µL of Fab supernatant followed by one hr incubation at room temperature. After washing, 50 µL of biotinylated CD123 was added and incubated for one hour at room temperature. After washing, 50 µL of Streptavidin:HRP (Pierce #21130) was added at a 1:5000 dilution and plates were incubated for one hour at room temperature. Plates were washed and 50 µL chemiluminescent

substrate, PoD (Roche # 121-5829500001), was added according to manufacturer’s instructions. Plates were then read for luminescence on an EnVision (Perkin Elmer) plate reader. Wells displaying signal >5-fold over background were considered hits.

[00281] Clones that demonstrated binding to recombinant human CD123 SP1 ECD-His tag protein were sequenced in the heavy (HC) and light chain (LC) variable regions. A total of 52 unique Fab sequences were identified via phage panning and 45 were ultimately converted to IgG1 isotype by in-fusion cloning. (Table 1) In-fusion cloning was performed by PCR-amplification using PCR SuperMix High Fidelity kit (Life Technologies # 10790-020), of the HC and LC variable regions and cloning into Esp3I sites in vDR149 for HC and vDR157 for LC using the In-Fusion® HD Cloning Plus kit (Clontech # 638909). VH and VL of the hits are shown below in Table 4.

Table 1. CDR sequences of mAbs generated from phage panning against recombinant human CD123 SP1 ECD-His tag protein (corresponding SEQ ID NOs are listed in parentheses)

ID	H-CDR1	H-CDR2	H-CDR3	L-CDR1	L-CDR2	L-CDR3
I3RB1	DYGMG (6)	YTRSGGSKYYADSVKG (7)	HGGSPRNELDY (8)	KSSQSVLYSSNNKNYLA (9)	WASTRES (10)	QYYSTPLT (11)
I3RB2	GYWMH (12)	AIRSDGSSKYYADSVKG (13)	DGVIEDTFDY (14)	RASQSVSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB3	SYWMG (18)	GPKYDGGSKYYADSVKG (19)	KMMSYFDY (20)	KSSQSVLYSSNNKNYLA (9)	WASTRES (10)	QYYSTPLT (11)
I3RB4	GYGMS (21)	AISGSGGSTYYADSVKG (22)	GNWYYGLGFDY (23)	RASQSVSSSYLA (24)	GASSRAT (25)	QYGSSPLT (26)
I3RB5	GYWMG (27)	GINYDGGSTYYADSVKG (28)	DHFLAEFDY (29)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB6	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GLFNWSNVALDY (35)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB7	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GKRWIADAGDFDY (36)	RASQSISSYLN (30)	AASSLQS (31)	QYGYSTPLT (32)
I3RB8	SYAIS (33)	GIPIFGTANYAQKFQG (34)	HGFAWNDYSLLDY (37)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB9	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GAPWENPPENLDY (38)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB10	SYGIS (39)	WISAIFGNTNYAQKFQG (40)	GGLLYYASYLDY (41)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB11	SYGIS (39)	GIPIFGTANYAQKFQG (34)	DLESWRYSNFDY (42)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB12	SYAIS (33)	GIPIFGTANYAQKFQG (34)	ADRVWDYYLDY (43)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB13	SYGIS (39)	GIPIFGNTNYAQKFQG (44)	DGGFYVRLDY (45)	RASQSVSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB14	SYGIS (39)	WISAIFGTTNYAQKFQG (46)	GGPLRYYNHFDY (47)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB15	SYAIS (33)	GIPIFGTANYAQKFQG (34)	DLESRLYSFLDY (48)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB16	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GAVWGDQWFDY (49)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)
I3RB17	SYAIS (33)	GIPIFGTANYAQKFQG (34)	GALSLWYSFLDY (50)	RASQSISSYLN (30)	AASSLQS (31)	QSYSTPLT (32)

ID	H-CDR1	H-CDR2	H-CDR3	L-CDR1	L-CDR2	L-CDR3
I3RB18	SYWIS (51)	IIDPSDSDTRYSPSFQG (52)	GDGSTDLDY (53)	RASQSVSSSYLA (24)	GASSRAT (25)	QQDYGFPWT (54)
I3RB19	NYAMS (55)	GIRGNSSSTYYADSVKG (56)	GHPITAAFPDYLDY (57)	RASQSIGDFLN (58)	YASSLQS (59)	QYGYTHPLT (62)
I3RB20	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	DDQIWGSYHLDY (60)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB21	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	ECWWSQCKFDY (61)	RASQSVANFLA (62)	AASNRAT (63)	QQYTHWPYT (64)
I3RB22	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	NLFYWADSVYLDY (65)	RASQSVNKWLA (66)	YASNRAT (67)	QQGIDWPRT (68)
I3RB23	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	ECSSWKNPRYVFDY (69)	RASQSISSYLN (30)	AASSLQS (31)	QQYDFEPLT (70)
I3RB24	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	HTDAWGYRLDY (71)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB25	SYGIS (39)	GISATFGNANYAQKFQG (72)	RPKWWSYFDY (73)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB26	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	NGFAWSVSGNLDY (74)	RASQSVDNWLA (75)	GASNRAT (76)	QQSISAPYT (77)
I3RB27	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	ACWWNLRYGLDY (78)	RASQSVAKSLA (79)	AASNRAT (83)	QQFICWPIT (80)
I3RB28	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	APFTWDYSRLDY (81)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB29	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	DRIWSEFLDY (82)	RASQSIGEWLN (83)	AASSLQS (31)	QQYHFEPLT (84)
I3RB30	SYAIS (33)	WIIPIFGTANYAQKFQG (85)	LVYSSDFDY (86)	RASQSVANWLA (87)	YASNRAT (67)	QQYDGWPRT (88)
I3RB31	SYAIS (33)	GISAYFENANYAQKFQG (89)	SYEGDAYFDY (90)	RASQSVDKDLA (91)	GASNRAT (76)	QQYDRAPIT (92)
I3RB32	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	GAWWAYDTYLDY (93)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB33	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	GYWAWNIDYLDY (94)	RASQSVNDWLA (95)	GASNRAT (76)	QQYKRAPYT (96)
I3RB34	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	GWSYYRLDY (97)	RASQSVDKWLA (98)	YASNRAT (67)	QQFDRAPFT (99)
I3RB35	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	HLFWDAGPLDY (100)	RASQSISSYLN (30)	AASSLQS (31)	QQYTSPPYT (101)
I3RB36	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	DLHVWAYSNEFDY (102)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB37	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	EKTEPPSRLDY (103)	RASQSIATWLN (104)	AASSLQS (31)	QQYITEPLT (105)
I3RB38	SYGIS (39)	GIIPIFGTANYAQKFQG (34)	DLMIWRFENFDY (106)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB39	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	EYGSLDY (107)	RASQSVADFLA (108)	AASNRAT (109)	QQYMGWPWT (110)
I3RB40	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	GQWWADTWFDY (111)	RASQSVAKWLA (112)	GASNRAT (76)	QQYHTAPWT (113)
I3RB41	SYAMS (114)	AISGSGGSTYYADSVKG (22)	VAYWEEFVYESLDY (115)	RASQSVSSSYLA (14)	GASSRAT (25)	QQYGSPLT (26)
I3RB42	SYAMS (114)	AISGSGGSTYYADSVKG (22)	HDWAFWIVFLDY (116)	RASQSVSSSYLA (15)	DASNRAT (16)	QQRSNWPLT (17)
I3RB43	SYWMB (117)	ATRGDGSSKYADSVKG (13)	DEIVMUTFDY (118)	RASQSVSSSYLA (15)	DASNRAT (16)	QYRSNWPLT (17)
I3RB44	SYWIS (51)	IIDPSDSDTRYSPSFQG (52)	GDGSTDLDY (53)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)
I3RB47	SYAIS (33)	GIIPIFGTANYAQKFQG (34)	ELFSWRYSNFDY (12)	RASQSISSYLN (30)	AASSLQS (31)	QQSYSTPLT (32)

Table 4: V_H and V_L sequences of mAbs generated from phage panning against CD123

mAb AA ID	VH Amino Acid Sequence	SEQ ID NO:	VL Amino Acid Sequence	SEQ ID NO
I3RB01	EVQLLESGGGLVQPGGSLRLSC AASGFTFSDYGMSWVRQAPGKG LEWVSVIRGGGSSKYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKHSGSFRFNELDY WGQGTLVTVSS	119	DIVMTQSPDSLAVSLGERAT INCKSSQSVLYSSNNKNYLA WYQQKPGQPPKLLIYWASTR ESGVPDRFSGSGSGTDFTLT ISSLQAEDVAVYYCQQYYST PLTFGQGGTKVEIK	164
I3RB02	EVQLLESGGGLVQPGGSLRLSC AASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWG QGTLVTVSS	120	EIVLTQSPATLSLSPGERAT LSCRASQSVSSYLA WYQQK GQAPRLLIYDASN RATGIPA RFSGSGSGTDFTLT ISSLEP EDFAVYYCQQR SNWPLTFGQ GTKVEIK	165
I3RB03	EVQLLESGGGLVQPGGSLRLSC AASGFTFSSYWMSWVRQAPGKG LEWVSGIKYDGGSKYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKKWMSYFDYWGG TLVTVSS	121	DIVMTQSPDSLAVSLGERAT INCKSSQSVLYSSNNKNYLA WYQQKPGQPPKLLIYWASTR ESGVPDRFSGSGSGTDFTLT ISSLQAEDVAVYYCQQYYST PLTFGQGGTKVEIK	164
I3RB04	EVQLLESGGGLVQPGGSLRLSC AASGFTFSGYGMSWVRQAPGKG LEWVSAISGSGGSTYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKGNWYYGLGFDYW GGTLVTVSS	122	EIVLTQSPGTLSLSPGERAT LSCRASQSVSSSYLA WYQQK PGQAPRLLIYGASSRATGIP DRFSGSGSGTDFTLTISRLE PEDFAVYYCQQYGSSPLTFG QGGTKVEIK	166
I3RB05	EVQLLESGGGLVQPGGSLRLSC AASGFTFSGYWMSWVRQAPGKG LEWVSGINYDGGSTYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDHFLAEFDYWGG GTLVTVSS	123	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB06	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGLFNWSNVALDY WGQGTLVTVSS	124	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB07	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGKRWLADAGDFD YWGQGTLVTVSS	125	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB08	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS	126	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP	167

	EDTAVYYCARHGFAWNDYSLLD YWGQGTTLVTVSS		EDFATYYCQQSYSTPLTFGQ GTKVEIK	
I3RB09	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGARWFNPPENLD YWGQGTTLVTVSS	127	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB10	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGWISAIFGNTNYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGGLLYASYLDY WGQGTTLVTVSS	128	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB11	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDLFSWRYSNFDY WGQGTTLVTVSS	129	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB12	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARADRVWDYYLDY WGQGTTLVTVSS	130	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB13	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGNTNYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARQSGFYVVRLDY WGQGTTLVTVSS	131	EIVLTQSPATLSLSPGERAT LSCRASQSVSSYLAWYQQKP GQAPRLLIYDASNRAIGIPA RFSGSGSGTDFTLTISLEP EDFAVYYCQQRNWLPLTFGQ GTKVEIK	165
I3RB14	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGWISAIFGTTNYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGGPLRYNHFY WGQGTTLVTVSS	132	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB15	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDLFSRLYSFLDY WGQGTTLVTVSS	133	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB16	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGAVWGDQWFDY	134	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ	167

	GQGTLVTVSS		GTKVEIK	
I3RB17	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGALSLWYSFLDY WGQGTLLVTVSS	135	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB18	EVQLVQSGAEVKKPGESLKISC KSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQG QVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQ GTLVTVSS	136	EIVLTQSPGTLSSLSPGERAT LSCRASQSVSSSYLAWYQQK PGQAPRLLIYGASSRATGIP DRFSGSGSGTDFTLTISRLE PEDFAVYYCQQDYGFPWTFG QGTKVEIK	168
I3RB19	EVQLLESGGGLVQPGGSLRLSC AASGFTFSNYAMSWVRQAPGKG LEWVSGIRGNGSSTYYADSVKG RFTISRDNKNTLYLQMNSLRA EDTAVYYCAKGGPIGARFPDYL DYWGQGTLLVTVSS	137	DIQMTQSPSSLSASVGDRV ITCRASQSIGDFLNWYQQKP GKAPKLLIYYASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	169
I3RB20	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDQIWGSYHLDY WGQGTLLVTVSS	138	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB21	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCAREGWGQKFDYW GQGTLLVTVSS	139	EIVLTQSPATLSLSPGERAT LSCRASQSVANFLAWYQQKP GQAPRLLIYAASNRA TGPAPRFSGSGSGTDFTLTISL LEPEDFAVYYCQQYFHW PYTFGQGTKVEIK	170
I3RB22	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARNLFYWADSVYLD YWGQGTLLVTVSS	140	EIVLTQSPATLSLSPGERAT LSCRASQSVNKLAWYQQKP GQAPRLLIYYASNRA TGPAPRFSGSGSGTDFTLTISL LEPEDFAVYYCQQGIDW PRTFGQGTKVEIK	171
I3RB23	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCAREGSSWKNPRYV FDYWGQGTLLVTVSS	141	EIVLTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQYFDFPLTFGQ GTKVEIK	172
I3RB24	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARHTDAWGYRLDYW	142	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ	167

	GQGTLLVTVSS		GTKVEIK	
I3RB25	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGISAIFGNANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARRFKWWESEYFDYW GQGTLLVTVSS	143	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB26	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARNGFAWSVSGNLD YWGQGTLLVTVSS	144	DIQMTQSPATLSLSPGERAT LSCRASQSVDNWLAWYQQKP GQAPRLLIYGASN RATGIPA RFSGSGSGTDFTLTISLEP EDFAVYYCQQSISAPYTFGQ GTKVEIK	173
I3RB27	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAIWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARAGWWNLRYGLDY WGQGTLLVTVSS	145	EIVLTQSPATLSLSPGERAT LSCRASQSVAKSLAWYQQKP GQAPRLLIYAASN RATGIPA RFSGSGSGTDFTLTISLEP EDFAVYYCQQFIGWPITFGQ GTKVEIK	174
I3RB28	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAIWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARAPFTWDYSRLDY WGQGTLLVTVSS	146	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB29	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAIWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARD SRIWSFSLDYW GQGTLLVTVSS	147	DIQMTQSPSSLSASVGDRV ITCRASQSIGEWNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQYYHFPLTFGQ GTKVEIK	175
I3RB30	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAIWVRQAPGQG LEWMGWIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARLVYSSDFDYWGQ GTLVTVSS	148	EIVLTQSPATLSLSPGERAT LSCRASQSVANWLAWYQQKP GQAPRLLIYYASN RATGIPA RFSGSGSGTDFTLTISLEP EDFAVYYCQQYDGWPRTFGQ GTKVEIK	176
I3RB31	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAIWVRQAPGQG LEWMGGISAYFGNANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARSYFGDAYFDYWG QGTLLVTVSS	149	EIVLTQSPATLSLSPGERAT LSCRASQSVDKDLAWYQQKP GQAPRLLIYGASN RATGIPA RFSGSGSGTDFTLTISLEP EDFAVYYCQQYDRAPITFGQ GTKVEIK	177
I3RB32	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGAWWAYDTYLDY	150	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISLQ EDFATYYCQQSYSTPLTFGQ	167

	WGQGTLLVTVSS		GTKVEIK	
I3RB33	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGYWHWNYDYLDY WGQGTLLVTVSS	151	EIVLTQSPATLSLSPGERAT LSCRASQSVNDWLAWYQQKP GQAPRLLIYGASNRATGIPA RFSGSGSGTDFTLTISSELP EDFAVYYCQQYKRAPYTFGQ GTKVEIK	178
I3RB34	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGWSYYRLDYWGQ GTLVTVSS	152	EIVLTQSPATLSLSPGERAT LSCRASQSVDKWLAWYQQKP GQAPRLLIYYASNATGIPA RFSGSGSGTDFTLTISSELP EDFAVYYCQQFDRAPFTFGQ GTKVEIK	179
I3RB35	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARHLFWDAGPLDYW GQGTLLVTVSS	153	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISSELP EDFATYYCQQYFSPPYTFGQ GTKVEIK	180
I3RB36	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDLHVWAYSNFY WGQGTLLVTVSS	154	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISSELP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB37	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDKTDFPSRLDYW GQGTLLVTVSS	155	DIQMTQSPSSLSASVGDRV ITCRASQSIATWLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISSELP EDFATYYCQQYITFPLTFGQ GTKVEIK	181
I3RB38	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYGISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDLMIWRFENFDY WGQGTLLVTVSS	156	DIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLTISSELP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB39	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCAREYGSLDYWGQGT LTVSS	157	EIVLTQSPATLSLSPGERAT LSCRASQSVADFLAWYQQKP GQAPRLLIYKASNRATGIPA RFSGSGSGTDFTLTISSELP EDFAVYYCQQYNGWPWTFGQ GTKVEIK	182
I3RB40	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGIIPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARGQWWADTWFDYW	158	EIVLTQSPATLSLSPGERAT LSCRASQSVAKWLAWYQQKP GQAPRLLIYGASNRATGIPA RFSGSGSGTDFTLTISSELP EDFAVYYCQQYHTAPWTFGQ	183

	GQGTLVTVSS		GTKVEIK	
I3RB41	EVQLLES GGGLVQPGGSLRLSC AASGFTFSSYAMSWVRQAPGKG LEWVSAISGSGGSTYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKVAYWEFFVYESL DYWGQGTLVTVSS	159	EIVLTQSPGTL SLSPGERAT LSCRASQSVSSSYLAWYQQK PGQAPRL LIYGASSRATGIP DRFSGSGSGTDFTLTISRLE PEDFAVYYCQQYGSSPLTFG QGTKVEIK	166
I3RB42	EVQLLES GGGLVQPGGSLRLSC AASGFTFSSYAMSWVRQAPGKG LEWVSAISGSGGSTYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKHDWAFWIVFLDY WGQGTLVTVSS	160	EIVLTQSPATL SLSPGERAT LSCRASQSVSSSYLAWYQQKP GQAPRL LIYDASN RATGIPA RFSGSGSGTDFTLT ISSLEP EDFAVYYCQQR SNWPLTFGQ GTKVEIK	165
I3RB43	EVQLLES GGGLVQPGGSLRLSC AASGFTFSSYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKG RFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGI VMDTFDYWG QGTLVTVSS	161	EIVLTQSPATL SLSPGERAT LSCRASQSVSSSYLAWYQQKP GQAPRL LIYDASN RATGIPA RFSGSGSGTDFTLT ISSLEP EDFAVYYCQQR SNWPLTFGQ GTKVEIK	165
I3RB44	EVQLLES GA EVKKPGESLKISC KSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQG QVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDL DYWGQ GTLVTVSS	162	DIQMTQSPSSL SASVGDRV T ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167
I3RB47	QVQLVQSGAEVKKPGSSVKVSC KASGGTFSSYAISWVRQAPGQG LEWMGGI IPIFGTANYAQKFQG RVTITADESTSTAYMELSSLRS EDTAVYYCARDLFSWRYSNFDY WGQGTLVTVSS	163	DIQMTQSPSSL SASVGDRV T ITCRASQSISSYLNWYQQKP GKAPKLLIYAASSLQSGVPS RFSGSGSGTDFTLT ISSLQP EDFATYYCQQSYSTPLTFGQ GTKVEIK	167

Example 3: MSD Cell Binding to hCD123 SP1, hCD123 SP2, and cynoCD123 SP1

[00282] Binding of CD123 antibodies to engineered pDisplay cells was assessed using a MSD (Mesoscale) cell binding assay. The object of the screening assay was to identify antibodies that bound to cells expressing hCD123 SP1 and SP2 as well as cross reactivity with cells expressing cynoCD123 SP1.

[00283] Cells were immobilized and phages were assayed in triplicate. Briefly, expression supernatants or purified CD123 antibodies were normalized to 10 µg/mL. 5000 cells per well were plated into a 384 well plate (MA6000, cat. L21XB, MSD) and allowed to adhere for 2 hr. Cells were then blocked with 20% FBS in PBS (Gibco) for 15 mins. Antibody

supernatants were then added and left at RT for 1 hr. Cells were washed 3 times with PBS and a ruthenium labeled secondary antibody (Jackson Immuno Research) was then added at 1 $\mu\text{g/mL}$ and incubated for 1 hr at room temperature. A further washing step was then applied and 35 μL per well of MSD Read buffer T (surfactant free) was then added and incubated for 30 min for detection. Plates were then read using Sector Imager 2400 (MSD). Data was normalized to controls and graphed using GraphPad Prism Version 5. A positive binder was determined to be a hit with a signal 3x greater than background (Figure 2A, B and C). The assay was repeated for data consistency and top binders were selected for further development. The following hits were positive for binding to all three cell lines: I3RB2, I3RB5, I3RB8, I3RB18, I3RB20, I3RB21, and I3RB35.

Example 4: Affinity measurements by SPR.

ProteOn Affinity Measurements

[00284] The affinities of 29 anti-CD123 candidates to recombinant human CD123 SP1 ECD and CD123 SP2 ECD were measured by Surface Plasmon Resonance (SPR) using a ProteOn XPR36 protein interaction array system (BioRad).

[00285] The rates of CD123 SP1 ECD or CD123 SP2 ECD association and dissociation were measured for each variant. The biosensor surface was prepared by covalently coupling Goat anti-Human IgG (Fc) to the surface of a GLC chip (BioRad) using the manufacturer instructions for amine-coupling chemistry. Approximately 8800 RU (response units) of Goat anti-Human IgG (Fc) antibody (Jackson ImmunoResearch laboratories Prod # 109-005-098) were immobilized. The RU immobilized also included a goat anti-mouse Fc antibody that was added to capture other antibodies not included in the ones reported here. Since the mixture was 1:1 about 50% of these RU immobilized are expected to be goat anti-human Fc. The kinetic experiments were performed at 25 °C in running buffer (PBS pH 7.4, 0.005% P20, 3 mM EDTA). 4-fold (1:3) serial dilutions of human CD123 SP1 ECD and CD123 SP2 ECD, starting at 400 nM were prepared in running buffer. An average of 300 RU of mAb (174-600) were captured on each channel of the sensor chip. The reference spots (Goat anti-Human IgG (Fc)-modified surface) containing no candidate captured were used as a reference surface. Capture of mAb was followed by 3 min injection (association phase) of antigen at 40 $\mu\text{L/min}$, followed by 10 min of buffer flow (dissociation phase). The chip surface was regenerated by injection of 0.85% phosphoric acid at 100 $\mu\text{L/min}$. Data was processed on the instrument software. Double

reference subtraction of the data was performed by subtracting the curves generated by buffer injection from the reference-subtracted curves for analyte injections. Kinetic analysis of the data was performed using 1:1 Langmuir binding model with group fit. The result for each mAb was reported in the format of K_a (kon or on-rate), K_d (koff or off-rate), K_D (Equilibrium dissociation constant) (Table 5).

[00286] The results indicated that all 29 mAbs bound to CD123 SP1 ECD, but only six of those showed binding to CD123 SP2 ECD. In order to assess data reproducibility, four of the antibodies were run at least in duplicate. In general, the results indicated good reproducibility between replicates, except for I3RB1 which has slow on-rates.

Table 5. Affinity assessment for phage panel 1 hits by SPR

CD123 SP1	CD123 SP1			CD123 SP2		
Sample Name	kon (1/Ms)	koff (1/s)	K_D (nM)	kon (1/Ms)	koff (1/s)	K_D (nM)
I3RB1	2.37E+04	5.69E-04	24.00	1.48E+04	4.57E-04	30.8
I3RB1	6.22E+03	1.88E-04	30.30	3.52E+03	3.70E-04	105
I3RB1	5.97E+04	7.82E-05	1.31	2.67E+04	$\leq 5e-5$	≤ 1.87
I3RB1	6.06E+04	2.45E-04	4.05	1.57E+04	1.50E-04	9.59
I3RB2	1.06E+06	4.77E-03	4.50	1.81E+06	3.35E-03	1.85
I3RB5	8.91E+05	1.14E-02	12.80	1.32E+06	6.43E-03	4.88
I3RB5	8.61E+05	1.11E-02	12.90	1.52E+06	6.23E-03	4.09
I3RB6	5.14E+05	5.93E-03	11.50	NBO		
I3RB7	9.54E+05	1.47E-02	15.40	NBO		
I3RB8	5.68E+05	1.95E-03	3.43	NBO		
I3RB9	6.80E+05	8.43E-03	12.40	NBO		
I3RB11	8.74E+05	2.53E-03	2.89	NBO		
I3RB12	8.12E+05	7.80E-03	9.61	NBO		
I3RB16	4.24E+05	2.12E-03	5.00	NBO		
I3RB16	3.87E+05	2.23E-03	5.77	NBO		
I3RB17	5.85E+05	2.01E-03	3.44	NBO		
I3RB18	1.44E+06	8.20E-04	0.57	2.69E+06	9.78E-04	0.363
I3RB19	2.11E+05	2.51E-02	119.00	3.34E+05	1.61E-02	48.3
I3RB20	6.31E+05	1.06E-03	1.68	NBO		

I3RB21	5.21E+05	1.14E-03	2.19	NBO		
I3RB22	2.57E+05	1.06E-03	4.12	NBO		
I3RB24	1.13E+06	2.26E-01	201.00	NBO		
I3RB28	5.28E+05	2.11E-03	3.99	NBO		
I3RB29	2.24E+05	1.32E-03	5.90	NBO		
I3RB30	7.25E+05	3.02E-03	4.17	1.45E+05	4.80E-02	330
I3RB32	8.68E+05	9.42E-04	1.09	NBO		
I3RB33	4.17E+05	1.77E-03	4.23	NBO		
I3RB34	4.97E+05	2.83E-02	56.80	NBO		
I3RB35	1.04E+06	2.93E-03	2.83	NBO		
I3RB36	6.75E+05	1.66E-03	2.47	NBO		
I3RB37	1.07E+06	6.69E-03	6.27	NBO		
I3RB37	1.21E+06	6.21E-03	5.15	NBO		
I3RB38	8.88E+05	4.34E-04	0.49	NBO ¹		
I3RB40	5.74E+05	3.46E-03	6.02	NBO		
I3RB47	1.59E+05	2.12E-03	13.40	NBO		

¹NBO = no binding observed

Biacore Affinity Measurements.

[00287] Affinity of several antibodies for the CD123 SP1 ECD and CD123 SP2 ECD was also measured by surface plasmon resonance (SPR) in both mAb and Fab format using a Biacore instrument. Kinetic studies were performed at 25° C using a Biacore 3000 (BIAcore, Inc., now part of GE Healthcare). Goat anti-Human IgG (Fc) specific antibody (Jackson ImmunoResearch laboratories Prod # 109-005-098) was covalently attached to two flow cells (normally 1 and 2) of the carboxymethyl dextran coated gold surfaces (CM-5 Chip, Biacore). Sheep anti-Human Fd specific antibody (The binding site Prod # PC075) was covalently attached to two flow cells (normally 3 and 4) of the carboxymethyl dextran coated gold surfaces (CM-5 Chip, Biacore). The carboxymethyl groups of dextran were activated with N-Ethyl-N'-(3-Dimethylaminopropyl)carbodiimide (EDC) and N-hydroxysuccinimide (NHS). The antibodies were coupled at pH 4.5 in 10 mM sodium acetate. Any remaining reactive sites on the surface were blocked by reaction with ethanolamine. For kinetic binding measurements, anti-CD123

antibodies were captured onto the anti-human Fc γ specific antibody, while the Fabs were captured onto the anti-Fd specific antibody by injecting the anti-CD123 molecules at a flow rate of 5 or 6 μ L/min. About 75 RU of antibody and about 50 RU of Fab were captured, respectively. Ab and Fab capture was followed by injection of human CD123 SP1 or human CD123 SP2 at concentrations between 1.6 nM and 400 nM at 40 μ L/min. Association data was collected for 2 min followed by 10 min of dissociation. The surface was regenerated with 30 μ L of 100 mM H₃PO₄ 100 μ L/min. All samples were prepared in D-PBS containing 3 mM EDTA and 0.005% surfactant P20. Data reported is the difference in SPR signal between the flow cell containing the captured antibody or Fab and a reference cell without captured antibody or Fab. Additional instrumental contributions to the signal were removed by subtraction of the data from the blank injection from the reference-subtracted signal. Data were performed in triplicate and analyzed by fitting association and dissociation phases at all concentrations (global fit) with a 1:1 binding model using the BIAevaluation software (BIAcore, Inc.). Duplicate experiments were performed and were in good agreement. Data presented is an average.

[00288] The results showed that the affinity of CD123 SP1 ECD and CD123 SP2 ECD binding to mAbs (I3RB2, I3RB18, I3RB35, I3RB37) are in agreement with their corresponding Fabs (I3RB120, I3RB119, I3RB121, I3RB122) (Table 6.). The results for all the anti-CD123 analyzed also showed that the affinity range for the Fab binding to CD123 SP1 ECD and CD123 SP2 ECD is 1.8-46.9 nM and 0.4-12.5 nM, respectively; while the affinity range for the mAb binding is 1.2-52 nM and 0.3-11.7 nM, respectively.

Table 6. Affinity and on-/off-rate values for anti-CD123 Phage 1 hits obtained by SPR (Biacore).

Construct	Class	rhCD123 SP1 K _D (nM)	rhCD123 SP2 K _D (nM)	rhCD123 SP1 k on Ave(M ⁻¹ s ⁻¹)	rhCD123 SP1 k off ave (s ⁻¹)
I3RB2	Mab	7.7	1.4	4.81E+05	3.72E-03
I3RB120	Fab	8.5	1.4	3.57E+05	3.04E-03
I3RB18	Mab	1.2	0.3	6.88E+05	8.08E-04
I3RB119	Fab	1.8	0.4	4.93E+05	8.91E-04

Construct	Class	rhCD123 SP1 K _D (nM)	rhCD123 SP2 K _D (nM)	rhCD123 SP1 k on Ave(M ⁻¹ s ⁻¹)	rhCD123 SP1 k off ave (s ⁻¹)
I3RB35	Mab	4.8	ND	5.40E+05	2.58E-03
I3RB121	Fab	6.3	1.2**	3.87E+05	2.45E-03
I3RB37	Mab	9.7	ND	5.45E+05	5.30E-03
I3RB122	Fab	11.5	ND	3.93E+05	4.50E-03

**Assay response is lower than expected

ND: apparent binding, but signal outside of acceptance criteria; (< 5 RU and bad data quality or irregular sensogram)

Example 5: Competition with 7G3

CD123 Competitive assay by ELISA

[00289] The CD123 antibody panel was screened in a 7G3 binding competition ELISA. 7G3 is a neutralizing monoclonal antibody, the epitope for which has been localized to within the first 50 amino acids of the CD123 SP1 antigen (US6177078B1). 7G3 mAb was purchased from BD Biosciences Pharmingen (San Diego, CA, Cat. No. 554526) and labeled with MSD Sulfo-Tag™ NHS-ester according to manufacturer's instructions (Meso Scale Discovery).

[00290] For CD123 competitive ELISA, 96-well clear maxisorb plates were treated with 100 µL/well of 2 µg/mL anti-6x histidine (R&D Systems Cat #: MAB050) made in bicarbonate buffer, pH 9.4 (Pierce #: 28382) and incubated at 4 °C overnight. The plates were then washed three times with ELISA wash buffer, (PBS, 0.01% Tween-20) and then blocked with 300 µL/well of StartingBlock containing Tween-20, PBST, (Thermo Scientific #: 37539). All wells were treated with 1 ng of recombinant huCD123 ECD SP1 and the plates were incubated at room temperature for 1 hr. Unbound huCD123 ECD SP1 was washed with ELISA wash buffer. 7G3 or mouse IgG2A (mIgG2A), was prepared in expression media (FreeStyle™ Expression media, Gibco #: 12338-018) at 20 µg/mL and added in duplicates to the plate at 50 µl/well to their respective wells whereas the test anti-CD123 mAbs were added at 50 µl/well of 2 µg/mL or neat to the remaining wells and the plates were incubated for 1 hr at room temperature with moderate shaking. Biotinylated 7G3 was then added to a final concentration of 100 ng/mL to all of the wells and the plates were incubated for an additional 1 hr. The plates were then

washed three times with ELISA wash buffer and bound biotinylated 7G3 was detected using SA-HRP conjugate at an optical density of 450 nm.

[00291] Anti-CD123 mAbs that inhibited 7G3:CD123 binding were defined at 20% inhibition of activity. That is, an antibody was considered to be an inhibitor if it was able to inhibit the binding of the biotinylated 7G3 to the human CD123 ECD by at least 20%. Based on this selection criterion, three inhibitors were identified: 13RB18, 13RB34, and 13RB44 (Figure 3).

Example 6: Functional pSTAT5 Assay

[00292] To assess agonist or antagonist activity of the antibodies, the panel was screened in a cell-based assay of IL-3-induced STAT5 phosphorylation using TF-1 cells (where purchased). The presence of anti-CD123 mAb inhibitor causes a decrease in STAT5 phosphorylation upon stimulation with rhIL-3. A 20% inhibition criterion was used in the STAT5 functional assay (20% inhibition of rhIL-3 activity).

[00293] Approximately 50,000 TF-1 (human erythroleukaemia) cells were plated in each well of a 96-well plate in 60 μ L of RPMI containing 10%FBS and incubated at 37 °C with 5% CO₂ incubator overnight. All samples were prepared in expression media (FreeStyleTM Expression media, Gibco #: 12338-018). The control samples received 70 μ L/well of either 20 μ g/mL 7G3, or mIgG2A isotype control. To the remaining wells, 70 μ L/well of 2 μ g/mL or neat anti-humanCD123 mAb samples were added. All samples were incubated for 1 hr at 37 °C with 5% CO₂ incubator. The cells were then treated with recombinant human IL-3, rhIL-3, (PeproTech catalog#: 200-03) at a final concentration of 10 ng/mL in RPMI containing 10% FBS with the exception of zero-, 7G3-, or isotype-only treated cells. The samples were then incubated for additional 15 min at 37 °C with 5% CO₂ incubator. Cells were lysed with 46.7 μ L ice-cold complete lysis buffer per well and the samples were incubated on ice for 30 min. Lysates were mixed by pipetting up and down 10 times. Phosphorylated STAT5 (pSTAT5a,b) was then determined using Phospho(Tyr694)/Total STAT5a,b kit from Meso Scale Discovery (MSD #: K15163D-2) and following the manufacturer's instructions.

[00294] Anti-CD123 mAbs that inhibited STAT5 phosphorylation by rhIL-3 were defined at 20% inhibition of activity. That is, an antibody was considered to be an inhibitor if it was able to inhibit the phosphorylation of STAT5 by rhIL-3 by at least 20%.

[00295] Five mAbs demonstrated ability to block IL-3 stimulation of STAT5 (Figure 4A). These five included 13RB18 as well as 13RB19, 13RB30, 13RB34, and 13RB44. However, when tested at 1 μ g/mL, only one antibody, 13RB18, blocked the IL-3 stimulation of

STAT5 phosphorylation in TF-1 cells (Figure 4B). Furthermore, 13RB18 (B18) showed dose dependence in this assay (Figure 4C). From these data, it was concluded that I3RB18 is the only antagonistic antibody.

Example 7: Confirmation of monovalent affinity on hCD123

[00296] The Fab binding of the two anti-CD123 hits (I3RB120 (I3RB2 Fab), I3RB119 (I3RB18 Fab) to cell-surface expressed human or cyno CD123 SP1 was analyzed in duplicate by MSD-Cell Affinity Technology to obtain a measure of the monovalent binding to cell-surface CD123.

[00297] Monovalent affinities of the selected anti-CD123 leads for cell-surface expressed hCD123 or cynoCD123 were performed using MSD-cell affinity technique (MSD-CAT) method. The MSD-CAT was developed in-house as a label-free method to determine affinity using intact cells in a high throughput format. These experiments were performed to assess the binding affinity and specificity of anti-CD123 candidates to cell-surface human or cynomolgus (cyno) CD123 SP1. This analysis allowed comparing the affinities of the anti-CD123 candidates to the human and cyno antigen in the absence of recombinant soluble cyno CD123. Cell lines used were human pDisplay CD123SP1 and cyno pDisplay CD123SP1. In order to measure the affinity of these interactions using the MSD-CAT method, a series of mixtures with a fixed concentration of anti-CD123 (1000, 200, 40 and/or 8 pM) and varying concentrations of cells (1.5×10^7 - 0.762×10^7 cells/mL) were prepared and allowed to reach equilibrium by rotating the plates for 24 hr at 4 °C. These samples were prepared in DMEM Glutamax medium containing 0.05% Azide, 1% BSA, 3 mM EDTA. The receptor numbers of $(3.15-4.18) \times 10^6$ hCD123/cell and $(4.78-9.24) \times 10^6$ cyCD123/cell were converted to M receptor concentration in the mixture on the basis of the volume of reaction, the cell density (cells/L) and the Avogadro's number. This resulted in a concentration range of 104 nM to 5.3 pM for human CD123 and 12 nM to 0.6 pM for cyno CD123. After equilibration the plate was centrifuged for 5 min ~1000 rpm and free anti-CD3 detected on the supernatant. The free anti-CD123 in the mixture was detected by electro chemiluminescence (ECL) using Mesoscale Discovery (MSD) reader instrument. For detection of free anti-CD123 in the equilibrated mixture by Electrochemiluminescence Immunoassays (ECL) detection plates were prepared. To prepare detection plates (plate bound antigen on SA-MSD plates) MSD Streptavidin Standard plates were blocked with 50 µL/well of assay buffer (PBS, (Life Sciences GIBCO 14190-136), 0.05%

Tween 20, 0.2% BSA) for 5 min. The assay buffer was removed without washing and 50 μL /well of 0.7 $\mu\text{g}/\text{mL}$ of biotinylated antigen in assay buffer were added to MSD plates and incubated overnight (~ 16 hr at 4°C). After overnight incubation, the plates were blocked by adding 150 μL /well of assay buffer without removing coating antigen, incubated for ~ 1 hr at ambient temperature and washed 5 times with wash buffer (assay buffer without BSA). 50 μL /well of the supernatants from samples plate were transferred to antigen-coated plates, incubated for 60 min, and then washed three times with wash Buffer. After this 50 μL per well of ruthenium labeled detection antibody(anti-human H+L) were added and incubated for 1 hr. After 1 hr the plates were washed and 150 μL of MSD Read Buffer (prepared by diluting 1:4 of stock into d. H_2O) were added per well. The plates were read immediately on the MSD Sector Imager 6000 Reader for luminescence levels. ECL signal detected by MSD was expressed in term of % free antibody in the mixture and the data was analyzed to determine affinity using a user defined equation (derived from the law of mass action) introduced in Prism software. The data show that I3RB18 and its Fab (I3RB119) are the tightest binders to cell-surface CD123 SP1 with pM affinity (or apparent affinity for the mAb) but binds >10 -fold weaker to cyno CD123 SP1. For I3RB18 and its Fab (I3RB119) it was not possible to get an affinity value for either the mAb or Fab against cynoSP1 expressing cells. All that can be said is that the affinity is > 12 nM. However, while I3RB120 binds with nM affinity to both antigens its binds with equal or < 5 -fold affinities to human and cyno CD123 SP1. The affinities obtained via SPR for hCD123 SP1 are weaker than observed on cells. This difference is most likely due to the presentation of the antigen on the cell surface and the location of the antibody's epitope. Results are shown in Table 7.

Table 7. Affinity values of Fabs to CD123 cells obtained by MSD-CAT

	hCD123 cells	hCD123 cells	cynoCD123 cells	cynoCD123 cells
	K_D (assay-1)	K_D (assay-2)	K_D (assay-1)	K_D (assay-2)
Fab I3RB119	293 pM	367 pM	>15 nM ^a	>11.9 nM ^a
Fab I3RB120	~ 3.37 pM 3.44 nM ^b	~ 3.84 nM 3.81 nM ^b	2.4 nM	>11.9 nM ^a
mAb I3RB18	55 ^c pM	343 ^c pM	832 ^c pM	$>11.9^c$ nM ^a
mAb 7G3	-	154 pM	-	57 pM

^a This K_D is greater than the value listed, but an actual value could not be determined.

^b In this fit a parameter called Bo was constrained to obtain an exact number instead of an approximation. The fitting algorithm sometimes gives an approximation when there is variability in the curve

^c This is apparent K_D because it could be affected by avidity due to bivalent binding.

[00298] The affinity measured for the I3RB2 Fab is consistent with the mAb data obtained via Proteon. Additionally, there is good cynoCD123 cell binding with this Fab, giving a clear indication that I3RB2 is a cross-reactive hit. The assessment of the I3RB18 mAb, and its corresponding Fab (I3RB119) indicate that the affinities obtained via Proteon for recombinant CD123 SP1 are weaker than observed on cells; 1 nM for recombinant protein vs 55-300 pM for cells. This difference is most likely due to the presentation of the antigen on the cell surface and the location of the antibody's epitope. It was not possible to get an affinity value for either the mAb or Fab (affinity > 12 nM). This would suggest that the antibody is not cross-reactive in a monovalent format. The previous cell binding data indicated cross-reactivity, which was most likely facilitated by the bivalent binding to the cell surface.

Example 8: Endogenous Cell Binding

[00299] Confirmation of binding of I3RB2 and I3RB18 to endogenous CD123 on AML cells was measured. OCI-AML5 cells (DSMZ), which express approximately 75,000 copies of CD123 on the cell surface, were used in a dose dependent MSD cell binding assay. Binding of CD123 antibodies to AML cells was assessed using a MSD (Mesoscale) cell binding assay. Briefly, expression supernatants or purified CD123 antibodies were used at a dose range of 40 µg/mL to 0.039 µg/mL. 50,000 cells per well were plated into a 96 well plate (Mesoscale high bind plate) and allowed to adhere for 2 hr. Cells were then blocked with 20% FBS in PBS plus Fc blocker (Fc blocker is the purified Fc portion of a papain-cleaved antibody (SEQ ID NO 209) for 15 min. Antibody supernatants were then added and left at RT for 1 hr. Cells were washed 3 times with PBS and a ruthenium labeled secondary antibody (Jackson Immuno Research) was then added at 1 µg/mL and incubated for 1 hr at room temperature. A further washing step was then applied and 150 µL of MSD Read buffer T (surfactant free) was then added and incubated for 30 mins for detection. Plates were then read using Sector Imager 2400 (MSD). Data was normalized to controls and graphed using GraphPad Prism Version 5.

[00300] The results showed that I3RB2 and I3RB18 bind to the endogenous CD123 expressed on OCI-AML5 cells in a dose dependent manner (Figures 5 A and B). The positive control, mAb 7G3, was also included in this assay as a comparator (Figure 5C).

Example 9: Competition binding analysis of CD123mABs with I3RB2 and I3RB18

[00301] A competition study was conducted for I3RB2 and I3RB18 against other cross-reactive CD123 SP1/SP2 hits and the 7G3 control to determine the anti-CD123 antibody competition groups or “epitope bins”.

[00302] For competitive ELISA, 5 μ L (20 μ g/mL) of purified human CD123 ECD protein generated as described in Example 1 was coated on MSD HighBind plate (Meso Scale Discovery, Gaithersburg, MD) per well for 2 hr at room temperature. A 150 μ L-aliquot of 5% MSD Blocker A buffer (Meso Scale Discovery) was added to each well and incubated for 2 hr at room temperature. Plates were washed three times with 0.1 M HEPES buffer, pH 7.4, followed by the addition of the mixture of labeled anti-CD123 mAb with different competitor anti-CD123 mAbs. Labeled antibodies (20 nM) were incubated 2 μ M of unlabeled anti-CD123 competitor antibodies, and then added to the designated wells in a volume of 25 μ L mixture. After a 2-hr incubation with gentle shaking at room temperature, plates were washed 3 times with 0.1 M HEPES buffer (pH 7.4). MSD Read Buffer T was diluted with distilled water (4-fold) and dispensed at a volume of 150 μ L/well and analyzed with a SECTOR Imager 6000. Antibodies were labeled with MSD Sulfo-TagTM NHS-ester according to manufacturer’s instructions (Meso Scale Discovery).

[00303] The competition ELISA results indicate that I3RB2 competes with I3RB60, I3RB70, I3RB79 and I3RB118 but does not compete with other antibodies including I3RB18 (Figure 6A). It should be noted, that when I3RB2 was labeled, competition was observed with I3RB60; however, when I3RB60 was labeled, competition was not observed. One possible reason for this is some non-specific binding interactions. When I3RB18 was assessed, it was found to compete with I3RB49 and I3RB55, but not with I3RB2 (Figure 6B).

[00304] The competition binning analysis defined two competition groups for the cross-reactive CD123 SP1/SP2 antibodies (Table 8). Monoclonal antibody I3RB2 does not compete with I3RB18 and they belong to different epitope groups. Group 1 (Dark Grey) includes mAbs I3RB2, I3RB60, I3RB70, I3RB79 and I3R118. Group 2 (Light Grey) consists of mAbs I3RB18, I3RB49 and I3RB55. The commercial mAb 7G3 does not compete with any in-house anti-CD123 antibodies.

Table 8. Results of Competition binding of Ru-labeled I3RB2 and I3RB18 to anti-CD123 Abs

Competitor	Ru-labeled antibody						
	I3RB2	I3RB70	I3RB79	I3RB18	I3RB55	I3RB60	7G3
I3RB2	+	+	+	—	—	—	—
I3RB60	+	+	+	—	±	+	—
I3RB70	+	+	+	—	±	—	—
I3RB79	+	+	+	—	±	—	—
I3RB118 (B102)	+	+	+	—	—	—	—
I3RB18	—	—	—	+	±	—	—
I3RB49	—	—	—	+	+	—	—
I3RB55	—	—	—	+	+	—	—
7G3	—	—	—	—	—	—	+

Example 10: Epitope Mapping of I3RB2 and I3RB18

H/D Exchange studies.

[00305] To identify the epitopes for I3RB2 and I3RB18 on human CD123, solution hydrogen/deuterium exchange-mass spectrometry (HDX-MS) was performed using the corresponding Fabs. For H/D exchange, the procedures used to analyze the Fab perturbation were similar to that described previously (Hamuro *et al.*, J. Biomol. Techniques 14:171–182, 2003; Horn *et al.*, Biochemistry 45:8488-8498, 2006) with some modifications. The CD123 SP2 ECD antigen was used for these studies since the antigen is less complex than the SP1 molecule due to a reduced number of glycosylation sites. Recombinant CD123 SP2 ECD (SEQ ID NO:226) was incubated in a deuterated water solution for predetermined times resulting in deuterium incorporation at exchangeable hydrogen atoms. The deuterated CD123 SP2 ECD was in complex with either I3RB119 (Fab of I3RB18) or I3RB120 (Fab of I3RB2) in 43 μL deuterium oxide (D₂O) at 4 °C for 30 sec, 2 min, 10 min and 60 min. The exchange reaction was quenched by low pH and the proteins were digested with pepsin. The deuterium levels at the identified peptides were monitored from the mass shift on LC-MS. As a reference control, CD123 SP2 ECD sample was processed similarly except that it was not in complex with the Fab molecules. Regions bound to the Fab were inferred to be those sites relatively protected from exchange and thus contain a higher fraction of deuterium than the reference CD123 SP2 ECD sample. About 94% of the protein could be mapped to specific peptides.

[00306] The solution HDX-MS perturbation maps of CD123 ECD SP2 with I3RB119 and I3RB120 are shown in Figure 7A and 7B, respectively. One segment, residues 176-184 (RARERVYEF (SEQ ID NO: 227)), corresponding to amino acid residues 195 – 202 of CD123 sp2, is strongly protected by I3RB119. Two different regions, residues 145-156 (IQKRMQPVITEQ (SEQ ID NO: 228)) and residues 165-170 (LLNPGT (SEQ ID NO: 229)), corresponding to residues 164 – 175 and residues 184 – 189 of CD123 sp2 respectively, were recognized by I3RB120. These HDX-MS results suggest the peptide level epitopes for I3RB119 and I3RB120. There were no overlapped epitope regions for these two antibodies. These results are in agreement with the previous competition binding data that I3RB2 and I3RB18 do not compete with each other.

Example 11: Epitope mapping of anti-CD123 antibody I3RB18 by Crystal Structure

[00307] The binding epitope of antibody I3RB18 was determined by X-ray crystallography.

[00308] The single-chain Fv fragment of anti-CD123 mAb I3RB18 was produced in the form: VL-(Gly4Ser)4-VH-Gly-His6 (SEQ ID NO:230). It was expressed in HEK293 Expi cells and purified by affinity (HisTrap) and ion exchange (Source 15S and Mono S) chromatography.

[00309] The sp2 isoform of human CD123 ECD (SEQ ID NO:231) with a C-terminal 8xHis tag was expressed in baculovirus-infected insect cells and purified by affinity (HisTrap) and size-exclusion (Superdex 75) chromatography.

[00310] The CD123:I3RB18 scFv complex was prepared by mixing 1.8 mg CD123 (1.1 mg/mL) with 2.4 mg scFv (1.6 mg/mL) at an approximate molar ratio of 1:1.2 (excess of scFv) and incubated overnight at 4°C. A small-scale (150 µg) SEC indicated complex formation. The protein was concentrated to 18 mg/mL in 20 mM HEPES, pH 7.5, 100 mM NaCl.

[00311] Crystallization was carried out by the vapor diffusion method at 20°C in a sitting drop format in MRC 2-well crystallization plates (Swissci). The crystals of the complex suitable for X-ray experiment were obtained under conditions: 2.0 M (NH₄)₂SO₄, 0.1 M MES buffer, pH 6.5. Crystal data are given in Table 9. One crystal was transferred to the mother liquor supplemented with 24% glycerol, frozen in liquid nitrogen, and used for X-ray diffraction data collection. The structure was determined at 3.5 Å resolution.

Table 9. Crystal data, X-ray data, and refinement statistics.

Crystal data

Space group	P4 ₁ 2 ₁ 2
Unit cell axes (Å)	111.32, 111.32, 192.19
Molecules/asym.unit	2
V _m (Å ³ /Da)	2.86
Solvent content (%)	57

X-ray data

Resolution (Å)	50–3.56	(3.70–3.56)
No.measured reflections	136,381	(5,853)
No.unique reflections	13,977	(929)
Completeness (%)	93.4	(64.2)
Redundancy	9.8	(6.3)
R-merge	0.195	(0.490)
<I/σ>	10.8	(2.3)
B-factor (Wilson) (Å ²)	66.1	

Refinement

Resolution (Å)	20–3.56
No. refls used in refinement	13,128
Completeness (%)	92.1
Number of all atoms	6568
Number of water molecules	0
R-factor (%)	23.1
R-free (%) (5% data)	32.3
RMSD bond lengths (Å)	0.005
RMSD bond angles (°)	1.1
Mean B-factor (Å ²)	120.3

Values for the highest-resolution shell are in parentheses.

[00312] I3RB18 binds CD123 sp2 at the C-terminal (proximal to cell surface) domain of the ECD. The epitope is conformational and includes three segments of the CD123 sp2 chain, residues 156 – 161 (RKFRYE, (SEQ ID NO:232)), 173 – 178 (TEQVRD, (SEQ ID NO: 233)) and 195 – 202 (RARERVYE (SEQ ID NO: 234)) corresponding to residues 234 – 239, 251 – 256 and 273 – 280 of CD123 sp1.. The antibody-antigen interactions are predominantly electrostatic. The epitope on CD123 sp2 contains a large number of basic residues, whereas the CDRs of I3RB18 are populated with acidic residues. The antibody residues involved in binding of CD123 include 7 residues from the light chain and 9 residues from the heavy chain (Fig. 8). All CDRs except LCDR2 are involved in binding.

[00313] The binding of I3RB18 to CD123 sp2 (Figure 9A) differentiates it from another anti-CD123 antibody, 7G3, which binds the N-terminal domain 1 of the CD123 sp1 ECD

as shown in the crystal structure of the humanized 7G3 Fab, CSL362, in complex with CD123 sp1 (Figure 9B) (pdb:4JZJBroughton et al. Cell Rep. 2014; 8:410-419).

Example 11: Crystal Structure of an anti-CD3 Fab

[00314] The crystal structure of the SP34 Fab was determined at 2.1 Å resolution. It revealed the complete amino acid sequence and identified the possible mouse germ lines from which the SP34 mAb was derived.

Materials

[00315] SP34 mAb, mouse IgG3/lambda isotype, was purchased from BD Biosciences Pharmingen (San Diego, CA), Cat. No. 556611. According to the technical data sheet, it was purified from tissue culture supernatant by affinity chromatography and stored at 4 °C. The Fab fragment was produced by papain digestion of mAb (Pierce, Cat # 44985, Thermofisher) and was separated from Fc using Nab Protein A Plus Spin column (Pierce, Cat # 44985, Thermofisher) according to manufacturer's protocol. The Fab was further purified on a MonoS column (GE Healthcare) equilibrated with 20 mM MES, pH 6.5 (buffer A). Elution was performed with buffer A in 13-28 % gradient of 1 M NaCl in 50 column volumes. Fractions corresponding to the main peak were pooled, concentrated to 9.2 mg/mL and used for crystallization.

Crystallization

[00316] Crystallization was carried out by the vapor diffusion method at 20 °C in a sitting drop format in 96-well Corning 3550 plates. The Fab crystal used for X-ray analysis was obtained from 12% PEG 3350, 0.2 M K/Na tartrate (pH 7.4), 3% isopropanol and 3% dioxane. Crystal data are given in Table 10.

Table 10 Crystal Data, X-ray data and refinement statistics

Crystal data

Space group	P21
Unit cell axes (Å)	55.14, 141.23, 61.29
Unit cell angles (°)	90, 99.02, 90
Molecules/asym.unit	2
V _m (Å ³ /Da)	2.48

Solvent content (%) 50

X-ray data

Resolution (Å)	30–2.1	(2.15–2.10)*
No.measured reflections	179,420	(11,506)
No.unique reflections	53,483	(3,667)
Completeness (%)	98.9	(92.5)
Redundancy	3.4	(3.1)
R-merge	0.038	(0.393)
<I/σ>	18.7	(3.8)
B-factor (Wilson) (Å ²)	45.4	

Refinement

Resolution (Å)	15–2.1
No. refls used in refinement	52,212
Completeness (%)	96.8
No. all atoms	6,886
No water molecules	219
R-factor (%)	20.5
R-free (%)	26.2
RMSD bond lengths (Å)	0.008
RMSD bond angles (°)	1.2
RMSD B-factor main-chain (Å ²)	2.7
Mean B-factor (Å ²)	53.7

* Numbers in parentheses are for the highest resolution shell.

X-ray data collection and structure determination

[00317] For X-ray data collection, one crystal was soaked for a few seconds in the mother liquor supplemented with 20% glycerol and flash frozen in liquid nitrogen. Diffraction data were collected at the Advanced Photon Source (Argonne, IL) IMCA beamline using a Pilatus CCD detector. X-ray data statistics are given in Table 10.

[00318] The structure was solved by molecular replacement using a Fab model constructed from mouse anti-Thomsen-Friedenreich Antigen antibody Jaa-F11 (PDB 3gnm),

which is a IgG3/kappa isotype. All crystallographic calculations were performed with the CCP4 suite of programs [CCP4. 1994, Acta Crystallogr. D50:760-763.]. Model adjustments were carried out using the program COOT [Emsley P, and Cowtan K. 2004. Acta Crystallogr. D60:2126-2132.]. The refinement statistics are given in Table 10.

[00319] The sequence of SP34 is shown in Figure 10, with residues 1 – 215 of the light chain and residues 1-230 of the heavy chain derived directly from the electron density map, and with residues 231 – 455 derived from IGHG3_MOUSE (mouse IgG3, isoform 2).

Example 12: Human Framework Adaptation of anti-CD3 antibody SP34

[00320] Anti-CD3 murine antibody SP34 was humanized by the Human Framework Adaptation method (Fransson, et al, JMB, 2010 398(2):214-31). Four different heavy chains were combined with three different light chains to produce 12 humanized variants.

SP34 Humanization and Affinity Maturation

Selection of human germlines

[00321] A matrix of four human heavy and three light v region sequences were selected for testing. Selection of human germlines were based solely on the overall sequence similarity to SP34 in the framework region (FR). Neither the CDR sequences, nor their length or canonical structures, were considered in this selection.

[00322] The closest matches for the heavy chain are human GLs IGHV3-72 and IGHV3-73. Another GL, IGHV3-23 was selected because of its high frequency of occurrence in the human B-cell repertoire.

[00323] The closest matches for the light chain are human lambda GLs IGLV7-43 (aka 7a), IGLV7-46 (aka 7b) and IGLV1-51 (aka 1b). IGLV7-46 is virtually identical to IGLV7-43, but has an advantage of Ala at position 2, i.e. as in SP34.

[00324] Selected J-regions are the following: IGHJ1 for the heavy chain; IGLJ3 for the lambda light chain

Back mutations

[00325] To preserve the conformation of CDR-H3, residues in several framework positions in VL, most notably positions Val38, Gly48 and Gly51 (Figure11) must be retained. These ‘back mutations’ were added into the humanization plan.

[00326] The Asn at position 57 of the heavy chain does not have good side chain density in the structure. It also sits in the middle of CDR-H2 and points away from the typical

binding site. Based upon this analysis, it may not contribute to binding significantly. In addition, the backbone geometry sits in a region most favorable for a Gly residue in the Ramachadran plot. Thus it was truncated to Gly in the maturation plan to allow necessary flexibility and potentially improve stability (by reducing non-glycine related locat structural strain) while not impacting binding.

[00327] There were several other considerations made in the humanization design. First, human GLs IGLV7-46 and IGLV7-43 introduce a Trp at position 59 with an unwanted oxidation potential. Two other GLs have Gly at this position, which corresponds to the mouse sequence. Therefore, Gly59 was preserved in both IGLV7-46 and IGLV7-43 variants. Finally, Ala at position 49 of VH may be essential. Also, the residue at position 99 (Val in SP34) may impact antigen binding. To test these positions, back mutations were introduced in some variants (Figure 12)

HFA matrix

[00328] The HFA matrix (Table 11) is composed of four variants of VH and three variants of VL (Figure 12). For the purpose of HFA, AbM CDR definition (K.R. Abhinandan and A. C. Martin, 2008. Mol. Immunol. 45, 3832-3839) is used.

The variants for VH:

CD3H141 (SEQ ID NO:184): IGHV3-72*01 with mouse CDRs+ Gly49Ala

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVARIRSKYNNYATYYAA
SVKGRFTISRDDSKNSLYLQMNSLKTEDTAVYYCARHGNFGNSYVSWFAYWGQGTLVTVSS

CD3H142 (SEQ ID NO:185): IGHV3-23*01 with mouse CDRs+ Ser49Ala

EVQLLES GGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVARIRSKYNNYATYYAD
SVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKHGNFGNSYVSWFAYWGQGTLVTVSS

CD3H143 (SEQ ID NO:186): IGHV3-23*01 with mouse CDRs+ Ser49Ala, Ala99Val

EVQLLES GGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVARIRSKYNNYATYYAD
SVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCVKHGNFGNSYVSWFAYWGQGTLVTVSS

CD3H144(SEQ ID NO:187): IGHV3-73*01 with mouse CDRs + Asn57Gly

EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKGLEWVGRIRSKYNGYATYYAA
SVKGRFTISRDDSKNTAYLQMNSLKTEDTAVYYCTR HGNFGNSYVSWFAYWGQGTLVTVSS

The variants for VL:

CD3L63 (SEQ ID NO:188): IGLV7-46*01 with mouse CDRs + F38V,A48G,Y51G,W59G
 QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGTPARF
 SGSLLGGAALTLGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

CD3L64 (SEQ ID NO:189): IGLV1-51*01 with mouse CDRs + Y38V, L48G, Y51G
 QSVLTQPPSVSAAPGQKVTISCRSSTGAVTTSNYANWVQQLPGTAPKGLIGGTNKRAPGIPDRF
 SGSKSGTSATLGITGLQTGDEADYYCALWYSNLWVFGGGTKLTVL

CD3L66 (SEQ ID NO:190): IGLV7-43*01 with mouse CDRs + F38V,A48G,Y51G,W59G
 QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGTPARF
 SGSLLGGAALTLGSGVQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Table 11 Matrix of CD3 Heavy and Light chains

(All were prepared with IgG1-AA Fc containing L234A, L235A, and F405L)

	CD3L63 (LV7-46/W59G) SEQ ID NO:188	CD3L64 (LV1-51) SEQ ID NO:189	CD3L66 (LV7-43/W59G) SEQ ID NO: 190
CD3H141 (HV3-72 + G49A) SEQ ID NO: 184	CD3B143	CD3B144	CD3B146
CD3H142 (HV3-23 + S49A) SEQ ID NO:185	CD3B147	CD3B148	CD3B150
CD3H143 (HV3-23 +S49A,A99V) SEQ ID NO: 186	CD3B151	CD3B152	CD3B154
CD3H144 (VH3-73 with G49) SEQ ID NO:187	CD3B155	CD3B156	CD3B158

[00329] Amino acid sequences were back-translated to DNA and cDNA was prepared using gene synthesis techniques (U.S. Pat. No. 6,670,127; U.S. Pat. No. 6,521,427). Heavy chain (HC) v regions were subcloned onto human IgG1-AA Fc containing L234A, L235A, and F405L mutations using an in-house expression vector with the CMV promoter using standard molecular biology techniques. Light chain (LC) variable regions were subcloned onto a human Lambda (λ) constant regions using an in-house expression vector with the CMV promoter using standard molecular biology techniques. Resulting plasmids were transfected into Expi293F cells (Invitrogen) and mAbs were expressed. Purification was by standard methods using a Protein A

column (hiTrap MAbSelect SuRe column). After elution, the pools were dialyzed into D-PBS, pH 7.2. The VH and VL sequence of the antibodies are shown in Table 12.

Table 12. The VH and VL sequences of anti-CD3 antibodies

mAb	HC	VH Amino Acid sequence	SEQ ID NO:	LC	VL Amino Acid sequence	SEQ ID NO:
CD3B1 43	CD3H 141	EVQLVESGGGLVQPG GSLRLSCAASGFTN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYAASVKGRFTISR DSKNSLYLQMNSLKT EDTAVYYCARHGNFG NSYVSWFAYWGQGT LTVSS	184	CD3L 63	QAVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGAQPED EAEYYCALWYSNLW VFGGGTKLTVL	188
CD3B1 44	CD3H 141	EVQLVESGGGLVQPG GSLRLSCAASGFTN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYAASVKGRFTISR DSKNSLYLQMNSLKT EDTAVYYCARHGNFG NSYVSWFAYWGQGT LTVSS	184	CD3L 64	QSVLTQPPSVSAAP GQKVTISCRSSTGA VTTSNYANWVQQLP GTAPKGLIGGTNKR APGIPDRFSGSKSG TSATLGITGLQTGD EADYYCALWYSNLW VFGGGTKLTVL	189
CD3B1 46	CD3H 141	EVQLVESGGGLVQPG GSLRLSCAASGFTN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYAASVKGRFTISR DSKNSLYLQMNSLKT EDTAVYYCARHGNFG NSYVSWFAYWGQGT LTVSS	184	CD3L 66	QTVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGVQPED EAEYYCALWYSNLW VFGGGTKLTVL	190
CD3B1 47	CD3H 142	EVQLLESGGGLVQPG GSLRLSCAASGFTN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYADSVKGRFTISR NSKNTLYLQMNSLRA EDTAVYYCAKHGNFG NSYVSWFAYWGQGT LTVSS	185	CD3L 63	QAVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGAQPED EAEYYCALWYSNLW VFGGGTKLTVL	188
CD3B1 48	CD3H 142	EVQLLESGGGLVQPG GSLRLSCAASGFTN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT	185	CD3L 64	QSVLTQPPSVSAAP GQKVTISCRSSTGA VTTSNYANWVQQLP GTAPKGLIGGTNKR	189

		YYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKHGNFG NSYVSWFAYWGQGT LTVSS			APGIPDRFSGSKSG TSATLGITGLQTGD EADYYCALWYSNLW VFGGGTKLTVL	
CD3B1 50	CD3H 142	EVQLLES GGGLVQPG GSLRLSCAASGFTFN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKHGNFG NSYVSWFAYWGQGT LTVSS	185	CD3L 66	QTVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGVQPED EAEYYCALWYSNLW VFGGGTKLTVL	190
CD3B1 51	CD3H 143	EVQLLES GGGLVQPG GSLRLSCAASGFTFN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCVKHGNFG NSYVSWFAYWGQGT LTVSS	186	CD3L 63	QAVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGAQPED EAEYYCALWYSNLW VFGGGTKLTVL	188
CD3B1 52	CD3H 143	EVQLLES GGGLVQPG GSLRLSCAASGFTFN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCVKHGNFG NSYVSWFAYWGQGT LTVSS	186	CD3L 64	QSVLTQPPSVSAAP GQKVTISCRSSTGA VTTSNYANWVQQLP GTAPKGLIGGTNKR APGIPDRFSGSKSG TSATLGITGLQTGD EADYYCALWYSNLW VFGGGTKLTVL	189
CD3B1 54	CD3H 143	EVQLLES GGGLVQPG GSLRLSCAASGFTFN TYAMNWVRQAPGKGL EWVARIRSKYNNYAT YYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCVKHGNFG NSYVSWFAYWGQGT LTVSS	186	CD3L 66	QTVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGVQPED EAEYYCALWYSNLW VFGGGTKLTVL	190
CD3B1 55	CD3H 144	EVQLVES GGGLVQPG GSLKLSAASGFTFN TYAMNWVRQASGKGL EWSGRIRSKYNGYAT YYAASVKGRFTISRDN SKNTAYLQMNSLKT EDTAVYYCTRHGNFG NSYVSWFAYWGQGT LTVSS	187	CD3L 63	QAVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGAQPED EAEYYCALWYSNLW VFGGGTKLTVL	188

CD3B1 56	CD3H 144	EVQLVESGGGLVQPG GSLKLSCAASGFTFN TYAMNWVRQASGKGL EWVGRIRSKYNGYAT YYAASVKGRFTISR DSKNTAYLQMNSLKT EDTAVYYCTRHGNE NSYVSWFAYWGQGT LTVSS	187	CD3L 64	QSVLTQPPSVSAAP GQKVTISCRSSTGA VTTSNYANWVQQLP GTAPKGLIGGTNKR APGIPDRFSGSKSG TSATLGITGLQTGD EADYYCALWYSNLW VFGGGTKLTVL	189
CD3B1 58	CD3H 144	EVQLVESGGGLVQPG GSLKLSCAASGFTFN TYAMNWVRQASGKGL EWVGRIRSKYNGYAT YYAASVKGRFTISR DSKNTAYLQMNSLKT EDTAVYYCTRHGNE NSYVSWFAYWGQGT LTVSS	187	CD3L 66	QTVVTQEPSLTVSP GGTVTLTCRSSTGA VTTSNYANWVQQKP GQAPRGLIGGTNKR APGTPARFSGSLLG GKAALTLSGVQPED EAEYYCALWYSNLW VFGGGTKLTVL	190

[00330] A monospecific anti-CD3 antibody CD3B143 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 184 and the VL of SEQ ID NO: 188 and an IgG1 constant region with L234A, L235A, F405L substitution. A monospecific anti-CD3 antibody CD3B144 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 184 and the VL of SEQ ID NO: 189 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B146 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 184 and the VL of SEQ ID NO: 190) and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B147 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 185 and the VL of SEQ ID NO: 188) and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B148 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 185 and the VL of SEQ ID NO: 189 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B150 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 185 and the VL of SEQ ID NO: 190 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B151 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 186 and the VL of SEQ ID NO: 188 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B152 was generated comprising the VH

and VL regions having the VH of SEQ ID NO: 186 and the VL of SEQ ID NO: 189 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B154 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 186 and the VL of SEQ ID NO: 190 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B155 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 187 and the VL of SEQ ID NO: 188 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B156 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 187 and the VL of SEQ ID NO: 189 and an IgG1 constant region with L234A, L235A, and F405L substitutions. A monospecific anti-CD3 antibody CD3B158 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 187 and the VL of SEQ ID NO: 190 and an IgG1 constant region with L234A, L235A, and F405L substitutions.

Example 13: Endogenous cell binding of the humanized anti-CD3 hits to primary T cells

[00331] The resulting panel of anti-CD3 antibodies was tested for binding against cell-surface CD3 ϵ on primary human T cells. To do this, binding of antibodies from expression supernatants was visualized using a polyclonal anti-human secondary antibody and analyzed by flow cytometry. Briefly, binding of anti-CD3 antibodies to cell-surface CD3 ϵ was assessed by flow cytometry using primary Human T lymphocytes purified by negative selection (Biological Specialty, Colmar, USA). Expression supernatants or purified antibodies were normalized to 10 μ g/ml in media or FACS buffer (BD BioSciences), respectively. 2x10⁵ cells were aliquoted into wells of a 96 well round-bottomed plate (CoStar) for labeling. Antibodies in expression supernatant were added to cells and incubated for 45 min at 4 °C. Following centrifugation at 1300rpm for 3 min and removal of supernatant, 50 μ L of anti-human IgG (H+L) Alexa Fluor 647 secondary antibody (Life technologies Inc.) was incubated with the cells at a final concentration of 10 μ g/mL for 30 min at 4 °C away from direct light. Following washing and resuspension in 30 μ L FACS buffer (BD BioSciences). Sample collection was performed on an Intellicyt HTFC system using ForeCyt software. Viable single cells were gated prior to analysis of binding using the green or red fixable live/dead dyes (Life Technologies Inc.) and forward/side scatter area and height parameters, respectively. Graphs were generated in GraphPad Prism version 5 using mean fluorescence intensity values.

[00332] Although a titration series was run, an intermediate concentration is presented in Figure 13 for clarity. Two in-house phage-derived antibodies with the same Fc region as the

therapeutic antibodies were used as controls: G11 (HC SEQ ID NO:222, LC SEQ ID NO:223), a non-cyno cross-reactive, agonistic antibody was used as a positive control and CD3B94 (HC-SEQ ID NO:224, LC – SEQ ID NO:225) a non-binder / non-agonistic antibody was used to assess non-specific binding. The commercial SP34 antibody was not used as a comparator in this assay since it is a mouse antibody and the use of a different secondary detection reagent would have prohibited direct comparison with the variants tested.

[00333] The data demonstrates an array of binding potential within the panel of humanized anti-CD3 hits, with two antibodies (CD3B144, CD3B152) showing complete loss of binding to human T cells. The remaining antibodies showed a range of binding potential that could be broadly split into strong and weak binders using G11 binding as an arbitrary threshold. Using these parameters, seven strong binders and seven weak binders were identified from the panel of variants (Figure 13).

[00334] Binding analysis of the anti-CD3 hits to primary cynomolgus CD4⁺ T cells was then tested in order to assess the retention of cross-reactivity. Purified CD4⁺ T cells from the peripheral blood of cynomolgus monkeys (Zen Bio, Triangle Research Park, USA) were used. Assay protocols were similar to those described above. Since G11 does not cross-react with cynomolgus CD3 ϵ , CD3B124, an in-house chimeric SP34-derived antibody having the VH and VL of SP34 with murine framework and a human IgG1 Fc was used as a positive control in this assay (Figure 14). Interestingly, several variants showed decreased binding potential compared to that seen with human cells. This included the strong binders CD3B150, CD3B151 and CD3B154, in which binding was reduced, and several weak binders where binding could no longer be detected over background. This loss of binding was not related to a specific immunoglobulin chain, suggesting that the combination of heavy and light chains played a role in the loss of cross-reactivity. Together, these assays allowed the identification of variants that retained species cross-reactivity between human and cynomolgus CD3 ϵ .

Example 14: Functional analysis of the humanized anti-CD3 hits in primary T cells

[00335] Binding analysis demonstrated that the panel of humanized anti-CD3 hits showed a range of binding potential to human and cynomolgus T-cells. To investigate the capacity of each variant to induce activation in via CD3 ϵ crosslinking, primary T-cells were cultured overnight in the presence of bead-conjugated antibody. The following day, cells were harvested and labeled with an anti-CD69 antibody to measure activation (Figure 15). Humanized anti-CD3 antibodies were bound to protein A coated magnetic beads (SpheroTech, Lake forest,

USA) by overnight incubation with antibody at 10 µg/mL. The following day, 2×10^5 primary human T cells were plated in round-bottomed cell culture plates in triplicate and 2×10^5 coated beads were added. Following overnight culture at 37 °C, cells were harvested and labeled with anti-CD69 Alexa Fluor® 488 antibody (clone FN50; Biolegend) to assess the up-regulation of this activation marker. Sample collection and analysis were performed as described above for binding. Several negative controls were run, including T-cells alone, T-cells with non-coated beads, and T-cells with isotype control (CD3B94)-coated beads. All of these showed similar mean fluorescence intensity values comparable to unstained T-cells indicating that background was low in this assay. Several positive controls were run for comparison, including OKT3 (US5929212) and commercially available SP34-2 antibody.

[00336] The humanized anti-CD3 hits were then tested for their capacity to activate primary cynomolgus CD4+ T cells (Zen Bio, Triangle Research Park, USA) in the same assay (Figure 16). The FN50 anti-CD69 antibody has been described as being cross-reactive with non-human protein and could therefore be used to test activation of these cells.

[00337] The human and cynomolgus activation data correlated with the binding data in that the panel of hits displayed a range of activation potentials. A number of the strong binders showed the capacity to activate human T-cells to an equivalent or greater extent when compared to commercially available SP34-2. Several variants showed activation potential that was lower compared SP34-2, whereas some binders did not show evidence of CD69 stimulation. The inability to activate was only seen in the variants that showed no or weak binding and all strong binders showed some level of activation, suggesting a correlation between binding and activation potentials for both human (Figure 17A) and cynomolgus(Figure 17B).

Example 15: Preparation of the Antibodies in a Bispecific Format in IgG1 L234A, L235A

[00338] Several monospecific CD123 antibodies were expressed as IgG1, having Fc substitutions L234A, L235A , and K409R (on anti-CD123) (numbering according to the EU index) in their Fc regions. The monospecific antibodies were expressed in HEK cell lines. The monospecific CD3 antibodies were IgG1 with Fc substitutions L234A, L235A, and F405L.

[00339] A monospecific anti-CD123 antibody I3RB135-K409R was generated comprising the VH and VL regions of an anti-CD123 antibody I3RB2 having the VH of SEQ ID NO: 120 and the VL of SEQ ID NO: 165 and an IgG1 constant region with L234A, L235A, and K409R substitution.

[00340] A monospecific anti-CD123 antibody I3RB125-K409R was generated comprising the VH and VL regions of an anti-CD123 antibody I3RB18 having the VH of SEQ ID NO: 136 and the VL of SEQ ID NO: 168 and an IgG1 constant region with L234A, L235A, and K409R substitution.

[00341] As a control, a monospecific anti-RSV antibody, B21M, was generated comprising the VH and VL regions having the VH of SEQ ID NO: 191 and the VL of SEQ ID NO: 192 and an IgG1 constant region with L234A, L235A, and either K409R or F405L to partner as the null arm with either the CD3 or CD123 arm of a bispecific antibody.

[00342] The monospecific antibodies were purified using standard methods using a Protein A column (HiTrap MabSelect SuRe column). After elution, the pools were dialyzed into D-PBS, pH 7.2.

[00343] The monospecific anti-CD123 antibodies were combined in matrix in *in-vitro* Fab arm exchange to generate bispecific antibodies that were subsequently characterized further (Table 13).

Table 13. Matrix of CD123 x CD3 mAbs to form bispecific antibodies

		CD123 ARMS		Control
		I3RB135 (I3RB2)	I3RB125 (I3RB18)	B21M, 409R
CD3 mAb	CD3B146	I3RB179	I3RB186	I3RB192
	CD3B147	I3RB180	I3RB187	I3RB193
	CD3B151	I3RB181	I3RB188	I3RB194
	CD3B154	I3RB182	I3RB189	I3RB195
	CD3B155	I3RB183	CD3B191	I3RB196
Control	B21M, F405L	I3RB185	I3RB191	I3RB198

mAb				
-----	--	--	--	--

[00344] Bispecific CD123 x CD3 antibodies were generated by combining a monospecific CD3 mAb and a monospecific CD123 mAb in *in-vitro* Fab arm exchange (as described in WO2011/131746). Briefly, at about 1-20 mg/mL at a molar ratio of 1.08:1 of anti-CD123/anti-CD3 antibody in PBS, pH 7-7.4 and 75 mM 2-mercaptoethanolamine (2-MEA) was mixed together and incubated at 25-37 °C. for 2-6 hr, followed by removal of the 2-MEA via dialysis, diafiltration, tangential flow filtration and/or spun cell filtration using standard methods. Control bispecific antibodies with an anti-RSV-(B21M) arm were generated similarly.

[00345] The generated monospecific anti-CD3 and CD123 antibodies were mixed for in vitro Fab arm exchange in matrix and characterized in various assays. The bispecific antibody I3RB179-Ab comprises the CD3 binding arm of mAb CD3B146-F405L and the CD123 binding arm of mAb I3RB135-K409R. The bispecific antibody I3RB186-Ab comprises the CD3 binding arm of mAb CD3B146-F405L and the CD123 binding arm of mAb I3RB125-K409R. The bispecific antibody I3RB180-Ab comprises the CD3 binding arm of mAb CD3B147-F405L and the CD123 binding arm of mAb I3RB135-K409R. The bispecific antibody I3RB187-Ab comprises the CD3 binding arm of mAb CD3B147-F405L and the CD123 binding arm of mAb I3RB125-K409R. The bispecific antibody I3RB181-Ab comprises the CD3 binding arm of mAb CD3B151-F405L and the CD123 binding arm of mAb I3RB135-K409R. The bispecific antibody I3RB188-Ab comprises the CD3 binding arm of mAb CD3B155-F405L and the CD123 binding arm of mAb I3RB125-K409R. The bispecific antibody I3RB182-Ab comprises the CD3 binding arm of mAb CD3B154-F405L and the CD123 binding arm of mAb I3RB135-K409R. The bispecific antibody I3RB189-Ab comprises the CD3 binding arm of mAb CD3B154-F405L and the CD123 binding arm of mAb I3RB125-K409R. The bispecific antibody I3RB183-Ab comprises the CD3 binding arm of mAb CD3B155-F405L and the CD123 binding arm of mAb I3RB135-K409R. The bispecific antibody CD3B191-Ab comprises the CD3 binding arm of mAb CD3B155-F405L and the CD123 binding arm of mAb I3RB125-K409R.

[00346] For control bispecific antibodies, anti-RSV antibody, B21M (HC SEQ ID NO: 207 – shown with F405L mutation, LC SEQ ID NO:208), was combined with either the CD3 arm or CD123 arms as follows. The bispecific antibody I3RB185-Ab comprises the anti-RSV binding arm of mAb B21M-F405L and the CD123 binding arm of mAb I3RB135-K409R. The

bispecific antibody I3RB191-Ab comprises the anti-RSV binding arm of mAb B21M-F405L and the CD123 binding arm of mAb I3RB125-K409R. The bispecific antibody I3RB192-Ab comprises the anti-RSV binding arm of mAb B21M-K409R and the CD3 binding arm of mAb CD3B146-F405L. The bispecific antibody I3RB193-Ab comprises the RSV binding arm of mAb B2M-F409R and the CD3 binding arm of mAb CD3B147-F405L. The bispecific antibody I3RB194-Ab comprises the anti-RSV binding arm of mAb B2M-F409R and the CD3 binding arm of mAb CD3B151-F405L. The bispecific antibody I3RB195-Ab comprises the anti-RSV binding arm of mAb B21M-K409R and the CD3 binding arm of mAb CD3B154-F405L. The bispecific antibody I3RB196-Ab comprises the RSV binding arm of mAb B21M-K409R and the CD3 binding arm of mAb CD3B155-F405L.

[00347] Heavy and Light chains for the CD123 x CD3 bispecific Abs are shown below in Table 14.

Table 14. Heavy and Light Chain Sequences for bispecific IgG1 antibodies

Ab		Amino Acid Sequence
I3RB179	Heavy chain 1 CD3B146 (SEQ ID NO:193)	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRIIRSKYNGYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTRHGNFGNSYVSWFAYWGQGLTVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYITLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFLLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B146	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK

	(SEQ ID NO:194)	ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB135 (I3RB2) (SEQ ID NO:203)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSRLTVDKSRWQQGNV FSCSVMHEALHNHYTQ KSLSLSPGK
	Light Chain 2 I3RB135 (I3RB2) (SEQ ID NO:204)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB180	Heavy chain 1 CD3B147 (SEQ ID NO:195)	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCAKHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFLLYSKLTVDKSRWQQGNV FSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ

	1 CD3B147 (SEQ ID NO: 196)	APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB135 (I3RB2) (SEQ ID NO:203)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSEFFLYSRLTVDKSRWQQGNVVFSCSVMHEALHNHYTQ KSLSLSPGK
	Light Chain 2 I3RB135 (I3RB2) (SEQ ID NO:204)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFI FPPSDEQLKSGTASV VCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGEC
I3RB181	Heavy chain 1 CD3B151 (SEQ ID NO:197)	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTT PPVLDSDGSEFLLYSKLTVDKSRWQQGNVVFSCSVMHEAL

		HNHYTQKSLSLSPGK
	Light Chain 1 CD3B151 (SEQ ID NO: 198)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB135 (I3RB2) (SEQ ID NO:203)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSRLTVDKSRWQQGNV FSCSVMHEALHNHYTQ KSLSLSPGK
	Light Chain 2 I3RB135 (I3RB2) (SEQ ID NO:204)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB182	Heavy chain 1 CD3B154 (SEQ ID NO:199)	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR

		EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFLLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B154 (SEQ ID NO:200)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB135 (I3RB2) (SEQ ID NO:203)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSRLTVDKSRWQQGNVFSCSVMHEALHNHYTQ KSLSLSPGK
	Light Chain 2 I3RB135 (I3RB2) (SEQ ID NO:204)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB183	Heavy chain 1 CD3B155 (SEQ ID NO: 201)	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRI RSKYNGYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTR HGNFGNSYVSWFAYWGQGT LVT VSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT

		PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFLLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B155 (SEQ ID NO: 202)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB135 (I3RB2) (SEQ ID NO:203)	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSRLTVDKSRWQQGNVFSCSVMHEALHNHYTQ KSLSLSPGK
	Light Chain 2 I3RB135 (I3RB2) (SEQ ID NO:204)	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLT ISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFI FPPSDEQLKSGTASV VCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB186	Heavy chain 1 CD3B146 (SEQ ID	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRI RSKYNGYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTR HGNFGNSYVSWFAYWGQGT LVT VSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH

	NO:193)	TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYITLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFLLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B146 (SEQ ID NO:194)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB125 (I3RB18) (SEQ ID NO: 205)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSDDL DYWGQGTLVTVSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYITL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP PVLDSDGSFFLYSRLTVDKSRWQQGNVFSCSVMHEALHNHYTQK SLSLSPGK
	Light Chain 2 I3RB125 (I3RB18) (SEQ ID NO: 206)	EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB187	Heavy chain	EVQLLES GGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG

	1 CD3B147 (SEQ ID NO:195)	LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCAKHG NFGNSYVSWFAYWGQGLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVF LFPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYITLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDS DGSFLLYSKLTVDKSRWQQGNV FSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B147 (SEQ ID NO: 196)	QAVVTQEPSLTVSPGGTVTLTCSRSTGAVTT SNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALT LSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTL FPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVET TTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain I3RB125 (I3RB18) (SEQ ID NO: 205)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDL DYWGQGLVTVSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPEAAGGPSVF LFPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYITL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PVLDS DGSFFLYSRLTVDKSRWQQGNV FSCSVMHEALHNHYTQK LSLSLSPGK
	Light Chain I3RB125 (I3RB18) (SEQ ID NO: 206)	EIVLTQSPGTL SLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

I3RB188	Heavy chain 1 CD3B151 (SEQ ID NO:197)	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHG NFGNSYVSWFAYWGQGLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDS DGSFLLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B151 (SEQ ID NO: 198)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain I3RB125 (I3RB18) (SEQ ID NO: 205)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGLVTVSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP PVLDS DGSFFLYSRLTVDKSRWQQGNVFSCSVMHEALHNHYTQK SLSLSPGK
	Light Chain I3RB125 (I3RB18) (SEQ ID	EIVLTQSPGTL SLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL

	NO: 206)	SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
I3RB189	Heavy chain 1 CD3B154 (SEQ ID NO:199)	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHG NFGNSYVSWFAYWGQGLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYITLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFLLYSKLTVDKSRWQQGNV FSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B154 (SEQ ID NO:200)	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain I3RB125 (I3RB18) (SEQ ID NO: 205)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGLVTVSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYITL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP PVLDSDGSFFLYSRLTVDKSRWQQGNV FSCSVMHEALHNHYTQK SLSLSPGK
	Light Chain I3RB125 (I3RB18)	EIVLTQSPGTL SLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS

	(SEQ ID NO: 206)	VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
CD3B19 1	Heavy chain 1 CD3B155 (SEQ ID NO: 201)	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRIIRSKYNGYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTRHGNFGNSYVSWFAYWGQGLTVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFLLYSKLTVDKSRWQQGNVVFSCSVMHEAL HNHYTQKSLSLSPGK
	Light Chain 1 CD3B155 (SEQ ID NO: 202)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKGDSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain I3RB125 (I3RB18) (SEQ ID NO: 205)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGLTVTVSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP PVLDSDGSFFLYSRLTVDKSRWQQGNVVFSCSVMHEALHNHYTQK SLSLSPGK
	Light Chain I3RB125	EIVLTQSPGTLISLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY

	(I3RB18) (SEQ ID NO: 206)	CQQDYGFPTWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
--	--	--

Example 16: Evaluation of Bispecific Antibodies in Functional Cell Killing Assay

[00348] T-cell mediated cytotoxicity assay is a functional assay to evaluate the CD123 x CD3 bispecific antibodies for cell lysis using T-cells from healthy donors.

[00349] The protocol of Laszlo, et al was followed (Laszlo, G., et al 2014 BLOOD 123:4, 554-561). Briefly, effector cells were harvested, counted, washed, and resuspended to 1×10^6 cells/ml in RPMI (10% FBS) cell media. Target cells were labeled with CFSE (Invitrogen #C34554) and resuspended to 2×10^5 cells/mL in RPMI (Invitrogen #61870-036) with 10% FBS (Invitrogen #10082-147). Effectors and CFSE-labeled target cells were mixed at E:T=5:1 in sterile 96-well round bottom plates. A 5 μ L aliquot of each bispecific antibody was added to each well containing various concentrations. Cultures were incubated for 48 hrs at 37 °C under 5% CO₂. After 48hr, The LIVE/DEAD® Fixable Near-IR Dead Cell Stain buffer (life technologies Cat# L10119) was added to samples, and cultures were incubated for 20 min in the dark at RT, washed, and resuspended in 170 μ L FACs buffer. The drug-induced cytotoxicity was determined using CANTO II flow cytometer (BD Biosciences) and analyzed with FlowJo Software or Dive software (BD Biosciences). The population of interest is the double positive CFSE+/ live/dead+ cells.

[00350] The results of the T-cell mediated cell lysis of AML cell lines MV4-11 (Figure 18 A and B), OCI-AML5 (Figure 19 A and B), and OCI-M2 (Figure 20 A and B) after 48 hr incubation at 37 °C, 5% CO₂ are shown. The MV4-11 and OCI-AML5 are CD123 expression cell lines, and the OCI-M2 has significant low CD123 expression. The Effector/Target ratio for this study was 5:1. A 2 mg/mL aliquot of Fc blocker was added to block Fc function.

[00351] Both I3RB2 and I3RB18 antibodies, when combined with an anti-CD3 antibody into a bispecific format, are efficacious at specifically killing CD123+ cells. Additionally, the data allow for a clear ranking between the I3RB135 (I3RB2-based) and I3RB125 (I3RB18-based) bispecific antibodies with the I3RB125 x CD3 bispecific antibodies being more potent than I3RB135 x CD3 bispecific antibodies. Within each family, the CD3B146- and CD3B155- based bispecific antibodies (higher affinity mAbs) were more potent than the CD3B151- and CD3B154- based bispecific antibodies. Low levels of dose-dependent background cytotoxicity are seen with low CD123 expression cell line OCI-M2.

Example 17: Evaluation of Bispecific Antibody, I3RB186 in a Tumor Model of Disease**Materials and Methods**

[00352] **Cell line.** In order to determine the efficacy of the bispecific antibody, I3RB186 in vivo, commercially available tumor cell lines with high CD123 expression were chosen for efficacy studies. The KG-1 (DSMZ, catalog number ACC 14) human acute myelogenous leukemia (AML) tumor cells were maintained in vitro in RPMI medium supplemented with heat inactivated fetal bovine serum (10% v/v) at 37 °C in an atmosphere of 5% CO₂ in air. The cells were routinely subcultured two to three times weekly. The cells growing in an exponential growth phase were harvested and counted for tumor cell inoculation.

[00353] **Preparation of Human PBMCs for engrafting.** Human, Mononuclear Enriched Cells (Catalog 213-15-04), obtained from Biological Specialty Corporation (Colmar, PA), were used for hIgG1-AA molecule testing. PBMCs were isolated via Ficoll density gradient separation (Ficoll-Paque TM Plus, GE Healthcare Bio-Sciences AB, Catalog 17-1440-03), and aliquoted at 50x10⁶ cells per vial in freezing media (Recovery Cell Culture Freezing Medium, Gibco, Catalog 12648-010). Vials were stored at -80 °C for approximately 24 hours, and then transferred to liquid nitrogen for long term storage. Frozen isolated peripheral blood mononuclear cell vials (100x10⁶ cells per vial, Catalog PB009-3) obtained from HemaCare (Van Nuys, CA) were used for IgG4 molecule testing. To thaw PBMCs, frozen vials were placed in a water bath at 37 °C. Cells were transferred to a conical tube containing cold thawing media. The conical tube was centrifuged, and cells were resuspended in sterile PBS. Cell viability was assessed using trypan blue exclusion method. Cells were resuspended to a cell concentration of 50x10⁶ cells per mL in sterile PBS for injection.

[00354] **Peripheral blood collection for FACS analysis.** For this, 50 µL of blood was collected from each animal via retro-orbital sinus into lithium heparin coated tubes. A 25 µL aliquot of blood from each sample was placed into 175 µL media (RPMI with 10% FBS) in each of two 96-well plates. The plates were centrifuged and red blood cells lysed using three treatments with ACK lysing buffer. Remaining cells were consolidated for each sample and stained for CD45, CD3, CD8, and CD4 to quantify circulating human T lymphocytes (see Mouse Peripheral Blood Harvesting/Staining: Protocol for Leukocyte Isolation and FACS analysis).

[00355] **Protocol for Leukocyte FACS analysis. Protocol for Leukocyte FACS analysis.** Peripheral blood was collected up to two times during the study for Fluorescence-activated Cell Sorting (FACS) analysis of circulating human PBMCs. Whole blood (25µL) was

diluted in 175 μ L of RPMI media in 96 well plates. Plates were centrifuged at 1400 rpm for 4 min and supernatant was decanted. Cells were resuspended in 200 μ L of ACK lysing buffer and incubated on ice for 5 min. After centrifugation at 1300 rpm for 5 min, supernatant was aspirated. Cells were retreated with ACK lysing buffer two more times and were washed once in 200 μ L PBS and recentrifuged at 1500 rpm for 5 min. Cell pellets were resuspended in 50 μ L/well of antibody cocktail in PBS containing Live/Dead stain (Invitrogen, cat# L10119, 0.25 μ L/well of stock. Stock is 1 vial diluted in 150 μ L DMSO) and incubated at room temperature in the dark for 30 min. The following antibodies were used to label cells: CD4 (Becton Dickinson Cat. 557922, 0.5 μ L/well), CD8 (Invitrogen, Q010055, 0.5 μ L of a 1:10 dilution in PBS/well), CD3 (Becton Dickinson, cat. 558117, 0.5 μ L/well), CD45 (BioLegend cat. 304006, 0.5 μ L/well). Cells were washed 3X with FACS buffer (200 μ L/well) and resuspended in 170 μ L FACS Buffer. Sample collection was performed on a BD LSR Fortessa Flow Cytometry Analyzer. Viable single cells were gated prior to analysis using Near-IR live/dead dye (Life Technologies Inc.) and forward/side scatter area and height parameters, respectively. Data was analyzed using BD FACS Diva software version 7.

[00356] In vivo design. Female NSG (NOD.Cg-*Prkdc*^{scid} *Il2rg*^{tm1Wjl}/SzJ) mice were subcutaneously inoculated with KG-1 cells (5×10^6 cells in phosphate buffered saline in a volume of 200 μ L) on the dorsal flank of each animal. The day of tumor cell inoculation was denoted as day 0. Tumor measurements were monitored twice weekly beginning seven days post-implantation, until tumor volumes ranged between 100-150 mm³ (fourteen days post-implantation), at which point mice were randomized by tumor volume into treatment groups. Mice were then intravenously (lateral tail vein) engrafted with human peripheral blood mononuclear cells (PBMCs) (10×10^6 cells in phosphate buffered saline in a volume of 200 μ L). Immediately following PBMC engraftment, mice received intravenous therapy bispecific Ab I3RB186 (bispecific diluted in PBS and dosed at a volume of 100 μ L). Treatment occurred approximately every other day for a total of five doses (see Table 15 for exact dosing days). Tumor measurements and body weights were recorded twice weekly.

[00357] The endpoints of the studies were tumor growth inhibition, maximal tumor burden (group mean greater than 1500 mm³), and body weight loss greater than 20% treatment initiation body weight. Tumor size was measured twice weekly in two dimensions using a caliper and the volume was expressed in mm³ using the formula: $V = 0.5 \times a \times b^2$ where a and b are the long and short diameters of the tumor, respectively. Complete tumor regression (CR) is defined as tumors that are reduced to below the limit of palpation (50 mm³). Partial tumor regression

(PR) is defined as tumors that are reduced from initial tumor volume. A minimum duration of CR or PR in three or more successive tumor measurements is required for a CR or PR to be considered durable.

[00358] The engraftment of human PBMCs leads to eventual graft-versus-host disease (GVHD) in the mice, where the engrafted donor T cells become activated and infiltrate the host tissues, leading to organ failure, extreme body weight loss, and inevitably, death. To monitor the onset and severity of GVHD in this model, body weight was recorded twice weekly and expressed in grams (g). Percent body weight change was calculated using the formula: Body weight change = $[(C-I)/I]*100$ where C is the current body weight and I is the body weight at the initiation of treatment.

[00359] Summary statistics, including mean and the standard error of the mean (SEM), are provided for the tumor volume of difference in tumor volume among each group at each time-point are shown in corresponding study tables. Statistical analysis of difference in tumor volume among the groups were evaluated using a two-way ANOVA repeated measures test, followed by Bonferroni post-test, using GraphPad Prism version 5.01. $p<0.05$ was considered to be statistically significant.

Efficacy of CD123xCD3 IgG1, F234A, L235A Bispecific Abs

[00360] NSG mice were subcutaneously inoculated with KG-1 cells, and then intravenously engrafted with human PBMCs described previously and dosed with the CD123 x CD3 bispecific Ab, I3RB186 at doses of 0.01, 0.1, 1, and 10 μg per animal, when tumors were established (mean tumor volume = $102 \pm 5.9 \text{ mm}^3$), as described previously. A subset of tumor-bearing mice were not engrafted with PBMCs but were dosed, as controls for the mechanism of the bispecific in the absence of control bispecific Abs. Also, a subset of non-tumor-bearing mice were engrafted with PBMCs and dosed, as controls for peripheral blood FACS analysis (see **Table 15** for study design).

Table 15. Dosing Schedule for in-vivo efficacy of I3RB186

Group	N	Tumor	PBMC	Treatment	Dose ($\mu\text{g}/\text{animal}$)	Dosing Route	Dosing Schedule (Days Post-tumor Implantation)	Blood Sampling (Days Post-tumor Implantation)
-------	---	-------	------	-----------	---	-----------------	--	---

1	10	+	-	PBS	0	i.v.	14, 17, 20, 22, 24	30
2	10	+	-	I3RB186	10	i.v.	14, 17, 20, 22, 24	30
3	10	+	-	I3RB186	1	i.v.	14, 17, 20, 22, 24	30
4	10	+	-	I3RB186	0.1	i.v.	14, 17, 20, 22, 24	30
5	10	+	-	I3RB186	0.01	i.v.	14, 17, 20, 22, 24	30
6	10	+	+	PBS	0	i.v.	14, 17, 20, 22, 24	30
7	10	+	+	I3RB186	10	i.v.	14, 17, 20, 22, 24	30
8	10	+	+	I3RB186	1	i.v.	14, 17, 20, 22, 24	30, 53
9	10	+	+	I3RB186	0.1	i.v.	14, 17, 20, 22, 24	30, 53
10	10	+	+	I3RB186	0.01	i.v.	14, 17, 20, 22, 24	30
11	5	-	+	PBS	0	i.v.	14, 17, 20, 22, 24	30, 53
12	5	-	+	I3RB186	10	i.v.	14, 17, 20, 22, 24	30, 53
13	5	-	+	I3RB186	1	i.v.	14, 17, 20, 22, 24	30, 53
14	5	-	+	I3RB186	0.1	i.v.	14, 17, 20, 22, 24	30, 53
15	5	-	+	I3RB186	0.01	i.v.	14, 17, 20, 22, 24	30, 53

Results of in-vivo efficacy study

[00361] Figure 21 shows the efficacy of CD123xCD3 IgG1-AA bispecific, I3RB186 – IgG1, F234A, L235A, in KG-1 human AML xenografts when human PBMCs are present, at two doses, 0.1 and 1 µg per animal ($p < 0.001$). Bispecific at 1 µg per animal (gray closed square) showed more immediate anti-tumor efficacy than at 0.1 µg, with complete regressions occurring in 3/8 animals, and partial regressions occurring in 3/8 animals. However, tumor regrowth was seen in 6/8 mice beginning at day 55 post-tumor implantation. Bispecific at 0.1 µg per animal (gray closed diamond) showed delayed but better efficacy with complete and partial regressions occurring in all animals. The data demonstrate the necessity of the presence of effector T lymphocytes for target cell killing with bispecific antibodies.

[00362] Figure 22 shows the FACS analysis of peripheral blood collected from mice on day 30 post-tumor implantation. An increase in CD45+ cells, driven by an increase in CD8+ T lymphocytes, was apparent in tumor-bearing animals treated with 0.1 and 1 µg bispecific antibodies. This expansion of CD8+ T lymphocytes only occurred when target cells (KG-1) were present, in groups where anti-tumor efficacy was observed. Alternately, 10 µg bispecific appeared to clear CD45+ PBMCs from peripheral blood. This clearance of effector cells may account for the lack of efficacy seen at this dose.

[00363] Figure 23 shows the FACS analysis of peripheral blood collected from mice on day 53 post-tumor implantation. CD45+, CD8+, and CD4+ cells were at similar levels in tumor-bearing mice treated with 0.1 and 1 µg bispecific, as in non-tumor bearing mice treated with PBS and 0.01 and 0.1 µg bispecific. Non-tumor bearing mice treated with 1 and 10 µg bispecific had very low levels of CD45+, CD8+, and CD4+ cells; the cause of this is currently unknown.

[00364] Figure 24 shows the mean body weight change of treatment groups over time. As described previously, body weight loss is correlated with onset and severity of GVHD, which is caused by activated T cells. In both tumor-bearing and non-tumor bearing mice, body weight loss was most severe with treatment with 0.1 µg bispecific antibody. Tumor-bearing mice treated with 1 µg bispecific did not experience severe body weight loss. T lymphocytes were present at day 53 post-tumor implantation (by FACS analysis, Figure 23), however the lack of body weight loss and GVHD onset indicates a loss of activated T cells, which may account for the tumor regrowth seen in this group beginning on day 55 post-tumor implantation (Figure 21).

Example 18. Evaluation of I3RB186 and control bispecific Abs (I3RB191 and I3RB192) *in-vivo*

[00365] In the second in -vivo experiment, bispecific Ab controls were added, I3RB191, a CD3 null arm and I3RB192, a CD123 null arm Ab. The protocol was the same as for Example 16. KG-1 human AML tumor xenografts were subcutaneously implanted into female NSG mice. Fourteen days after implant, mice were randomized by tumor volume to treatment groups. Human PBMCs were intravenously implanted, followed by intravenous treatment with I3RB186, and I3RB191 and I3RB192 control bispecific Abs at 1 µg per animal (see dosing schedule on Table 16). Treatment occurred on days 14, 16, 18, 21, and 23 days after tumor implant. Arrows in the figure show the bispecific Ab administration days.

Table 16 Dosing Schedule for 2nd in-vivo experiment

Group	N	Tumor	PBMC	Treatment	Dose (µg/animal)	Dosing Route	Dosing Schedule (Days Post-tumor Implantation)	Blood Sampling (Days Post-tumor Implantation)
1	8	+	+	PBS	0	i.v.	14, 16, 18, 21, 23	36
2	8	+	+	I3RB192	1	i.v.	14, 16, 18, 21, 23	36
3	8	+	+	I3RB191	1	i.v.	14, 16, 18, 21, 23	36
4	8	+	+	I3RB186	1	i.v.	14, 16, 18, 21, 23	36, 63
5	8	+	-	PBS	0	i.v.	14, 16, 18, 21, 23	N/A
6	8	+	-	I3RB192	1	i.v.	14, 16, 18, 21, 23	N/A
7	8	+	-	I3RB191	1	i.v.	14, 16, 18, 21, 23	N/A
8	8	+	-	I3RB186	1	i.v.	14, 16, 18, 21, 23	N/A
9	4	-	+	PBS	0	i.v.	14, 16, 18, 21, 23	36, 63
10	4	-	+	I3RB186	1	i.v.	14, 16, 18, 21, 23	36, 63

[00366] The anti-tumor activity of the bispecific Abs is shown as change in tumor size (mm³) over time (Figure 25). Treatment with I3RB186 at 1 µg significantly inhibited tumor growth (p<0.001) compared to that of PBS and control bispecific Ab-treated animals.

[00367] On day 36 post-tumor implantation, peripheral blood was collected for FACS analysis of circulating human PBMCs. Unlike the first study, there was no difference in the frequency of human CD45+ PBMCs (a) or CD8+ and CD4+ T lymphocyte frequencies (b) in animals treated with I3RB186 compared with PBS and I3RB191 (Figure 26). CD45+, CD8+, and CD4+ cells were at lower frequencies in tumor-bearing and non-tumor bearing animals treated with I3RB192, the CD123 null arm control bispecific Ab.

[00368] On day 63 post-tumor implantation, peripheral blood was collected for FACS analysis of circulating human PBMCs. Of the tumor-bearing animals, only animals treated with I3RB186 at 1 µg remained (Figure 27). There was an elevation in frequency of CD45+ human PBMCs (a) and CD8+ T lymphocytes (b) in tumor-bearing animals treated with 1 µg I3RB186, compared with non-tumor bearing animals treated with PBS or 1 µg I3RB186 (Figure 25). CD4+ T lymphocytes were at similar frequencies across all remaining groups. Non-tumor bearing mice treated with PBS and 1 µg I3RB186 had very low frequencies of CD45+, CD8+, and CD4+ cells.

[00369] Figure 28 shows the mean body weight change of treatment groups over time. As described previously, body weight loss is correlated with onset and severity of GVHD, which is caused by activated T cells. In tumor-bearing mice, there was a greater loss in body weight with treatment with 1 µg bispecific antibody, compared to all other groups. This is contradictory to the first study, where tumor-bearing mice treated with 1 µg bispecific did not experience severe body weight loss. T lymphocytes were present at day 63 post-tumor implantation (Figure 27), however the efficacy at the 1 µg dose was not as pronounced as in the first study (Figures 21, 25).

Example 19. Preparation of the Antibodies in a Bispecific Format in IgG4 S228P, F234A, L235A

[00370] Several of the monospecific CD3 and CD123 antibodies were expressed as IgG4, having Fc substitutions S228P, F234A, and L235Ax (CD123 arm) or S228P, F234A, L235A, F405L, and R409K(CD3 arm) (numbering according to EU index) in their Fc regions. The monospecific antibodies were expressed in CHO cell lines under CMV promoters.

[00371] A monospecific anti-CD3 antibody CD3B219 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 184 and the VL of SEQ ID NO: 190 and an IgG4 constant region with S228P, F234A, L235A, F405L, and R409K substitutions. A monospecific anti-CD3 antibody CD3B217 was generated comprising the VH and VL regions

having the VH of SEQ ID NO: 186 and the VL of SEQ ID NO: 188 and an IgG4 constant region with S228P, F234A, L235A, F405L, and R409K substitution. A monospecific anti-CD3 antibody CD3B218 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 186 and the VL of SEQ ID NO: 190 and IgG4 constant region with S228P, F234A, L235A, F405L, and R409K substitutions. A monospecific anti-CD3 antibody CD3B220 was generated comprising the VH and VL regions having the VH of SEQ ID NO: 187 and the VL of SEQ ID NO: 188 and IgG4 constant region with S228P, F234A, L235A, F405L, and R409K substitutions.

[00372] A monospecific anti-CD123 antibody I3RB218 was generated comprising the VH and VL regions of an anti-CD123 antibody I3RB2 having the VH of SEQ ID NO: 120 and the VL of SEQ ID NO: 165 and an IgG4 constant region with S228P, F234A, and L235A substitutions. A monospecific anti-CD123 antibody I3RB217 was generated comprising the VH and VL regions of an anti-CD123 antibody I3RB18 having the VH of SEQ ID NO: 136 and the VL of SEQ ID NO: 168 and an IgG4 constant region with S228P, F234A, and L235A substitutions.

[00373] As a control, a monospecific anti-RSV antibody, derived from B21M, was generated comprising the VH and VL regions having the VH of SEQ ID NO: 191 and the VL of SEQ ID NO: 192 and an IgG4 constant region with S228P, F234A, L235A, or F234A, L235A, R409K, F405L to partner as the null arm with either the CD3 or CD123 arm of a bispecific antibody.

[00374] The monospecific antibodies were purified, and the generated monospecific anti-CD3 and CD123 antibodies were mixed for in vitro Fab arm exchange in matrix (Table 12) as previously described in Example 15 and characterized in various assays. The bispecific antibody -Ab 7959 comprises the CD3 binding arm of mAb CD3B219 -F405L, R409K and the CD123 binding arm of mAb I3RB217 -R409. The bispecific antibody Ab 3978 comprises the CD3 binding arm of mAb CD3B217 -F405L, R409K and the CD123 binding arm of mAb I3RB217 -R409. The bispecific antibody Ab 7955 comprises the CD3 binding arm of mAb CD3B218 -F405L, R409K and the CD123 binding arm of mAb I3RB217 -R409. The bispecific antibody 9958 Ab comprises the CD3 binding arm of mAb CD3B220 -F405L, R409K and the CD123 binding arm of mAb I3RB217 -R409. The bispecific antibody Ab 8747 comprises the CD3 binding arm of mAb CD3B219 -F405L, R409K and the CD123 binding arm of mAb I3RB218 -R409. The bispecific antibody Ab 8876 comprises the CD3 binding arm of mAb CD3B217 -F405L, R409K and the CD123 binding arm of mAb I3RB218 -R409. The bispecific

antibody Ab 4435 comprises the CD3 binding arm of mAb CD3B218 -F405L, R409K and the CD123 binding arm of mAb I3RB218 -R409. The bispecific antibody Ab 5466 comprises the CD3 binding arm of mAb CD3B220 -F405L, R409K and the CD123 binding arm of mAb I3RB218 -R409.

[00375] For control bispecific antibodies, B2M1 in the IgG4 PAA format was generated, purified and, combined with either the CD3 arm or CD123 arms following the matrix in the table 17 below.

Table 17 Matrix of IgG4 bispecific antibodies

	CD3B219 (I3RB146) SEQ ID NO:210, 211	CD3B217 (I3RB151) SEQ ID NO:212, 213	CD3B218 (I3RB154) SEQ ID NO:214, 215	CD3B220 (I3RB155) SEQ ID NO:216, 217	B21M IgG4, F045L CD3 null
I3RB217 (I3RB18) SEQ ID NO:218, 219	7959	3978	7955	9958	CD3 null 1 (4309)
I3RB218 (I3RB2) SEQ ID NO:220, 221	8747	8876	4435	5466	CD3 null 2 (6601)
B21M IgG4, K409R CD123 null	CD123 null 1	CD123 null 2	CD123 null 3	CD123 null 4	CD123 CD3 null (3244)

[00376] Heavy and Light chains for CD123 x CD3 bispecific antibodies are shown in Table 18.

Table 18. Heavy and Light Chain Sequences for bispecific Abs IgG4-PAA

Ab	Chain	Amino Acid Sequence
7959	Heavy chain 1 CD3B219 (I3RB146) SEQ ID NO:210	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYAASVKGRFTISRDDSKNSLYLQMNSL KTEDTAVYYCARHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTKTYTCNVDHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTI SKAKGQPREPQ

		VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 1 CD3B219 (I3RB146) SEQ ID NO:211	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB217 (I3RB18) SEQ ID NO:218,	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGTLVTVSSASTKGPSVFPLAPC SRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGKTKYTCNVDHKPSNTKVKRVESKY GPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPS QEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVL DSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 2 I3RB217 (I3RB18) SEQ ID NO:219	EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGFPTWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
3978	Heavy chain 1 CD3B217 (I3RB151) SEQ ID NO:212	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNGFGNSYVSWFAYWGQGT LVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGKTKYTCNVDHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY

		KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 1 CD3B217 (I3RB151) SEQ ID NO:213	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB217 (I3RB18) SEQ ID NO:218,	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDSDTRYSPSFQGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGTLVTVSSASTKGPSVFPLAPC SRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGKTKTYTCNVDPHKPSNTKVDKRVESKY GPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPS QEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVL DSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 2 I3RB217 (I3RB18) SEQ ID NO:219	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGFPTWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
7955	Heavy chain 1 CD3B218 (I3RB154) SEQ ID NO:214	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNGFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGKTKTYTCNVDPHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK

	Light Chain 1 CD3B218 (I3RB154) SEQ ID NO:215	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB217 (I3RB18) SEQ ID NO:218,	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDS DTRYSPSFQGGVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDL DYWGQGTLVTVSSASTKGPSVFPLAPC SRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLG TKTYTCNV DHKPSNTKVDKRVESKY GPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPS QEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV L DSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLS LSLGK
	Light Chain 2 I3RB217 (I3RB18) SEQ ID NO:219	EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
9958	Heavy chain 1 CD3B220 (I3RB155) SEQ ID NO:216	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRI RSKYNAYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTR HGNFGNSYVSWFAYWGQGT LVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLG TKTYTCNV DHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	CD3B220 (I3RB155)	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK

	SEQ ID NO:217	ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB217 (I3RB18) SEQ ID NO:218,	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKG LEWMGIIDPSDS DTRYSPSFQGGQVTISADKSISTAYLQWSSLKA SDTAMYYCARGDGSTDLDYWGQGTLVTVSSASTKGPSVFPLAPC SRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVVTVPSSSLGTKTYTCNVDHKPSNTKVDKRVESKY GPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPS QEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLS LSLGK
	Light Chain 2 I3RB217 (I3RB18) SEQ ID NO:219	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQA PRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQDYGF PWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTAS VVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
8747	Heavy chain 1 CD3B219 (I3RB146) SEQ ID NO:210	EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYAASVKGRFTISRDDSKNSLYLQMNSL KTEDTAVYYCARHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTKTYTCNVDHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 1 CD3B219 (I3RB146) SEQ ID NO:211	QTVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS

	Heavy chain 2 I3RB218 (I3RB2) SEQ ID NO:220	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVI EDTFDYWGQGTLVTVSSASTKGPSVFPLAP CSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLG TKTYTCNV DHKPSNTKVDKRVESK YGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVD VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL SLSLGK
	Light Chain 2 I3RB218 (I3RB2) SEQ ID NO:221	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASN RATGIPARFSGSGSGTDFTLTISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYSL STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
8876I	Heavy chain 1 CD3B217 (I3RB151) SEQ ID NO:212	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNFGNSYVSWFAYWGQGTLVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLG TKTYTCNV DHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 1 CD3B217 (I3RB151) SEQ ID NO:213	QAVVTQEPSLTVSPGGTVTLTCSRSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB218 (I3RB2) SEQ	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVI EDTFDYWGQGTLVTVSSASTKGPSVFPLAP CSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL

	ID NO:220	QSSGLYSLSSVVTVPSSSLGKTYTCNVDPHKPSNTKVDKRVESK YGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVD VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL SLSLGK
	Light Chain 2 I3RB218 (I3RB2) SEQ ID NO:221	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASNRAITGIPARFSGSGSGTDFTLTISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPRFAKVKQWVDNALQSGNSQESVTEQDSKDSTYSLS STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
4435	Heavy chain 1 CD3B218 (I3RB154) SEQ ID NO:214	EVQLLESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKG LEWVARIRSKYNNYATYYADSVKGRFTISRDN SKNTLYLQMNSL RAEDTAVYYCVKHGNGNSYVSWFAYWGQGT LVT VSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVT VSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGKTYTCNVDPHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
	Light Chain 1 CD3B218 (I3RB154) SEQ ID NO:215	QTVVTTQEPSTLVSPGGTVTLTCSRSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYA ASSYLSLTPEQWKSHRYSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB218 (I3RB2) SEQ ID	EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGT LVT VSSASTKGPSVFPLAP CSRSTSESTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGKTYTCNVDPHKPSNTKVDKRVESK YGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVD VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVL

5466	NO:220	HQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL SLSLGK
	Light Chain 2 I3RB218 (I3RB2) SEQ ID NO:221	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASNRATGIPARFSGSGSGTDFTLTISSLEPEDFAVYYC QQRSNWPLTFGQGKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYSL STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
	Heavy chain 1 CD3B220 (I3RB155) SEQ ID NO:216	EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQASGKG LEWVGRIRSKYNAYATYYAASVKGRFTISRDDSKNTAYLQMNSL KTEDTAVYYCTRHGNGFNSYVSWFAYWGQGLTVTVSSASTKGPS VFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTCTYTCNVDHKPSNTKVD KRVESKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQ VYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTTPPVLDSDGSFLLYSKLTVDKSRWQEGNVFSCSVMHEALHNH YTQKSLSLSLGK
5466	CD3B220 (I3RB155) SEQ ID NO:217	QAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQQKPGQ APRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYA ASSYLSLTPEQWKSHRYSYSCQVTHEGSTVEKTVAPTECS
	Heavy chain 2 I3RB218 (I3RB2) SEQ ID NO:220	EVQLLES GGGLVQPGGSLRLSCAASGFTFSGYWMHWVRQAPGKG LEWVSAIRSDGSSKYYADSVKGRFTISRDN SKNTLYLQMNSLRA EDTAVYYCAKDGVIEDTFDYWGQGLTVTVSSASTKGPSVFPLAP CSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVVTVPSSSLGTCTYTCNVDHKPSNTKVDKRVESK YGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVD VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL

		SLSLGK
	Light Chain 2 I3RB218 (I3RB2) SEQ ID NO:221	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAP RLLIYDASNRAITGIPARFSGSGSGTDFTLTISSLEPEDFAVYYC QQRSNWPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDESTYSL STLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

[00377] Example 20. CD123 monovalent affinity of bispecific antibodies in IgG4-PAA format using recombinant antigen

[00378] Surface plasmon resonance (SPR) experiments were performed to determine the kinetics and affinity for the binding of CD3XCD123 bispecific antibodies to human CD123 SP1 ECD and CD123 SP2 ECD.

[00379] The affinities of anti-CD123 xCD3 bispecific Abs 3978, 7955, 7959, 9958 8876, 8747, 5466 for recombinant human CD123 SP1 and recombinant human CD123 SP2 ECD were measured by surface plasmon resonance (SPR) using a Biacore instrument. Kinetic studies were performed at 25° C using a Biacore T200 (Biacore, Inc., now part of GE Healthcare). Goat anti-Human IgG (Fc) specific antibody (Jackson ImmunoResearch laboratories Prod # 109-005-098) was covalently attached to the carboxymethyl dextran coated gold surfaces of a CM-5 sensor chip (GE Healthcare). The carboxymethyl groups of dextran were activated with N-Ethyl-N'-(3-Dimethylaminopropyl)carbodiimide (EDC) and N-hydroxysuccinimide (NHS). The anti-Fc antibody was coupled at pH 4.5 in 10 mM sodium acetate. Any remaining reactive sites on the surface were blocked by reaction with ethanolamine. For kinetic binding measurements, anti-CD123 antibodies were captured onto the anti-human Fcγ specific antibody,. 40-70 RU of antibody were captured. Ab capture was followed by injection of human CD123 SP1 or human CD123 SP2 at concentrations between 0.4nM and 400 nM at 40 μL/min. Association data was collected for 2 min followed by 10 min of dissociation. The surface was regenerated with 30 μL of 100 mM H3PO4 100 μL/min, followed by 50 mM NaOH. The samples for kinetic analysis were prepared in PBS-based buffer (D-PBS containing 3 mM EDTA and 0.005% surfactant

P20). Data reported is the difference in SPR signal between the flow cell containing the captured antibody and a reference cell without captured antibody. Additional instrumental contributions to the signal were removed by subtraction of the data from the blank injection from the reference-subtracted signal. Data were analyzed by fitting association and dissociation phases at all concentrations (global fit) with a 1:1 binding model using the BIAevaluation software (BIAcore, Inc.). Table 20 and 21 summarize the kinetic and affinity results obtained by Biacore. Both tables show the data obtained during three or more independent experiments.

[00380] Biacore data show that within the same family I3RB18-derived bispecific Abs and I3RB2-derived bispecific Abs bind with similar affinities to CD123 SP1 (Table 19) and with similar affinities to CD123 SP2 (Table 20) I3RB18-derived bispecific Abs bind to recombinant CD123 SP1 > 10-fold tighter than I3RB2-derived bispecific Abs with affinities ~1 nM and 14 nM, respectively. When binding to recombinant CD123 SP2, I3RB18 derived bispecific Abs bind > 5-fold tighter than I3RB2 derived bispecific Abs with affinities ~ 0.3 nM and 1.7 nM, respectively. Standard deviations in Tables 19 and 20 indicate that the data were very reproducible.

Table 19 Biacore kinetic and affinity data for the binding of bispecific antibodies to recombinant human CD123 SP1.

Sample ID	Common name	k_{on} Ave ($M^{-1} s^{-1}$)	k_{on} STDEV ($M^{-1} s^{-1}$)	k_{off} Ave (s^{-1})	k_{off} STDEV (s^{-1})	K_D Ave (nM)	K_D STDEV (nM)
3978	CD123(B18) x CD3(B151)	5.64E+05	3.82E+04	8.30E-04	4.70E-05	1.47	0.129
7955	CD123(B18) x CD3(B154)	5.62E+05	4.53E+04	8.40E-04	5.30E-05	1.49	0.153
7959	CD123(B18) x CD3(B146)	5.79E+05	3.55E+04	8.80E-04	5.40E-05	1.53	0.132
9958	CD123(B18) x CD3(B155m)	5.87E+05	4.57E+04	7.90E-04	5.00E-05	1.34	0.135

8876	CD123(B2) x CD3(B151)	3.43E+05	1.10E+04	4.90E-03	1.20E-04	14.4	0.583
4435	CD123(B2) x CD3(B154)	3.37E+05	1.25E+04	4.80E-03	1.60E-04	14.3	0.713
8747	CD123(B2) x CD3(B146)	3.37E+05	1.25E+04	4.80E-03	2.10E-04	14.3	0.821
5466	CD123(B2) x CD3(B155m)	3.71E+05	6.43E+03	5.10E-03	6.70E-05	13.7	0.298
3244	B21M x B21M	NB	NB	NB	NB	NB	NB
I3RB18	Mab for Fab I3RB119	7.73E+05	5.68E+04	7.20E-04	3.60E-05	0.935	0.083

NB – no bniding

Table 20. Biacore kinetic and affinity data for the binding of anti-CD123 bispecific antibodies to recombinant human CD123 SP2

Sample ID	Common name	k _{on} Ave (M ⁻¹ s ⁻¹)	k _{on} STDEV (M ⁻¹ s ⁻¹)	k _{off} Ave (s ⁻¹)	k _{off} STDEV (s ⁻¹)	K _D Ave (nM)	K _D STDEV (nM)
3978	CD123(B18) x CD3(B151)	3.12E+06	6.34E+05	1.10E-03	5.30E-05	0.356	0.074
7955	CD123(B18) x CD3(B154)	3.33E+06	9.37E+05	1.10E-03	2.90E-05	0.344	0.097
7959	CD123(B18) x CD3(B146)	3.78E+06	5.43E+05	1.30E-03	1.30E-04	0.335	0.06
9958	CD123(B18) x CD3(B155m)	3.57E+06	9.82E+05	1.10E-03	6.90E-05	0.311	0.088

8876	CD123(B2) x CD3(B151)	2.83E+06	4.07E+05	5.00E-03	1.30E-04	1.75	0.255
4435	CD123(B2) x CD3(B154)	2.88E+06	5.51E+05	5.00E-03	3.20E-04	1.74	0.349
8747	CD123(B2) x CD3(B146)	3.19E+06	1.05E+06	5.20E-03	4.50E-04	1.63	0.558
5466	CD123(B2) x CD3(B155m)	2.88E+06	2.86E+05	4.90E-03	2.80E-04	1.69	0.193
3244	B21M x B21M	NB	NB	NB	NB	NB	NB

NB – no binding

[00381] Example 21. CD123 monovalent affinity of bispecific antibodies in IgG4-PAA format to cell-surface expressed antigen by MSD-CAT

[00382] Monovalent affinities of the selected anti-CD123 bispecific antibodies for cell-surface expressed hCD123 SP1 and SP2 were performed using MSD-cell affinity technique (MSD-CAT) method. The MSD-CAT was developed in-house as a label-free method to determine affinity using intact cells in a high throughput format. These experiments were performed to assess the binding affinity and specificity of anti-CD123 candidates to cell-surface human CD123 SP1 and CD123 SP2. Cell lines used were human pDisplay CD123SP1 and pDisplay CD123SP2. A negative control antibody was used to test if the bispecific Abs scaffold bound nonspecifically to the cells and differentiate nonspecific versus specific binding to CD123. In order to measure the affinity of these interactions using the MSD-CAT method, a series of mixtures with a fixed concentration of anti-CD123 (800, 160, 32 and 6 pM) and varying concentrations of cells (20 Million to 1016 cells/mL) were prepared and allowed to reach equilibrium by rotating the plates for 24 hours at 4°C. These samples were prepared in DMEM Glutamax medium containing 0.05% Azide, 1% BSA, 3 mM EDTA. The receptor numbers of $(0.29-1.08) \times 10^6$ hCD123 SP1/cell and $(0.57-1.5) \times 10^6$ hCD123 SP2/ were converted to M receptor concentration in the mixture on the basis of the volume of reaction, the cell density (cells/L) and the Avogadro's number. This resulted in a concentrations ranging from of 35 nM to 0.5 M for human CD123 SP1 ; and 49nM to 0.97 pM for human CD123 SP2. After equilibration the plate was centrifuged for 5 minutes ~1000 rpm and free anti-CD3 detected on

the supernatant. The free anti-CD123 in the mixture was detected by electro chemiluminescence (ECL) using mesoscale discovery (MSD) reader instrument. For detection of free anti-CD123 in the equilibrated mixture by Electrochemiluminescence Immunoassays (ECL) detection plates were prepared. To prepare detection plates (plate bound antigen on SA-MSD plates) MSD Streptavidin Standard plates were blocked with 50 uL/well of assay buffer (PBS, (Life Sciences GIBCO 14190-136), 0.05% Tween 20, 0.2% BSA) for 5 minutes. The assay buffer was removed without washing and 50 uL/well of 0.7 ug/mL of biotinylated antigen in assay buffer were added to MSD plates and incubated overnight (~16 hours at 4°C). After overnight incubation, the plates were blocked by adding 150 uL/well of assay buffer without removing coating antigen, incubated for ~1 hour at ambient temperature and washed 5 times with wash buffer (assay buffer without BSA). 50 uL/well of the supernatants from samples plate were transferred to antigen-coated plates, incubated for 60 minutes, and then washed 3 times with wash buffer. After this 50 uL per well of ruthenium labeled detection antibody(anti-human H+L) were added and incubated for 1 hour. After 1 hour the plates were washed with wash buffer and 150 uL of MSD Read Buffer (Read Buffer T 4X, R92TD-2, MSD) were added per well. The plates were read immediately on the MSD Sector Imager 6000a Reader for luminescence levels. ECL signal detected by MSD was expressed in terms of % free antibody in the mixture and the data was analyzed to determine affinity using a user defined equation (derived from the law of mass action) introduced in Prism software. Results for MSD-CAT experiments are shown in Table 21.

Table 21. MSD-CAT affinity data show the binding of anti-CD123 molecules to cell-surface human CD123 SP1 and human CD123 SP2. The data were fit using non-linear least square analysis with a 1:1 binding model.

Sample ID	K_d [pM] human CD123 SP1 cells	K_d [pM] human CD123 SP2 cells
3978	153 ± 124	528 ± 296
7955	136 ± 105	436 ± 255
7959	149 ± 98	461 ± 290
9958	121 ± 80	538 ± 430

8876	1291 ± 556	2450 ± 2104
4435	1531 ± 1093	3701 ± 1898
8747	1761 ± 1337	2211 ± 1003
5466	2431 ± 1222	1722 ± 1638
3244	No binding	No binding
I3RB18 mAb	*47 ± 14	*49 ± 36
I3RB2 mAb	NA	*36 ± 20

NA = not applicable; assay was not performed

[00383] MSD-CAT affinities of Bispecific Abs for cell-surface CD123 SP1 are >6-fold tighter than SPR data for recombinant CD123 SP1; However, the affinities for cell-surface CD123 SP2 are similar to recombinant CD123 SP2 (< 2-fold different). The difference in SPR versus MSD-CAT affinities for CD123 SP1 is most likely due to the presentation of the antigen on the cell surface in comparison to the recombinant antigen. MSD-CAT showed that I3RB18-derived bispecific Abs (3978, 7955, 7959, 9958) are the tightest binders to cell-surface human CD123 SP1 and human CD123 SP2 with pM affinities. I3RB18-derived affinities are about 10-fold and about 5-fold tighter than I3RB2-derived bispecific Abs to cell-surface CD123 SP1 and CD123 SP2, respectively. The affinities were similar for bispecific Abs within the same family.

[00384] Overall, molecular interaction analyses using Biacore and MSD-CAT are in agreement showing that I3RB18-derived bispecific Abs bind tighter to recombinant and cell-surface human CD123 (SP1 and SP2) than for I3RB2-derived bispecific Abs.

[00385] **Example 22. CD123 monovalent affinity of bispecific antibodies in IgG4-PAA format to cell-surface expressed antigen by flow cytometry**

[00386] Flow cytometry was used to measure affinity values of several CD123xCD3 bispecific Abs for CD3 on human T cells (Biological Specialty, Colmar, USA) and cynomolgus monkey T cells (Zen Bio, Triangle Research Park, USA). The format involved competition binding using a fixed concentration of labeled anti-CD3 mAb of known affinity and increasing concentrations of unlabeled test Abs (Ashkenazi A et al. PNAS: 88:10535, 1991.). The anti-CD3

mAb used was CD3B146 hu IgG1-AlaAla F405L antibody with an affinity value similar to SP34-2. The K_d for SP34-2 was determined using saturation binding and examples of human and cynomolgous monkey T-cell binding curves are shown in Figure 29. Figure 30 shows the competition binding with labeled B146 and various concentrations of unlabeled CD123 x CD3 bispecific antibodies obtained for human (Figure 30 A) and cynomolgous (Figure 30 B) T-cells. Comparable values were obtained for human and cynomolgous monkey T cells. There appear to be three CD3 affinity groups among the samples analyzed: high (9-15 nM), medium (25-50 nM) and low (110-270 nM) which are summarized in Table 22.

TABLE 22. Affinity values (K_d) for CD123 x CD3 bispecific antibodies to human or cynomolgus T cells – competition binding using labeled B146 and increasing concentrations of unlabeled antibodies

	Human T-cells	Cyno T-cells
bispecific Abs	K _d (nM)	K _d (nM)
3978	241.2 +/- 57.3	215.0 +/- 17.1
8876	169.2 +/- 27.9	109.6 +/- 4.8
CD123 null 2	266 +/- 78.0	217 +/- 18.0
7955	209.6 +/- 31.8	169.1 +/- 8.8
4435	173.6 +/- 48.6	138.9 +/- 2.8
CD123 null 3	200.5 +/- 67.3	236.7 +/- 16.4
7959	11.0 +/- 4.3	11.2 +/- 0.3
8747	9.6 +/- 1.5	9.5 +/- 0.1
CD123 null 1	13.2 +/- 3.0	13.4 +/- 0.3
9958	43.0 +/- 10.6	29.1 +/- 1.2
5466	27.9 +/- 9.3	25.3 +/- 1.1
CD123 null 4	48.6 +/- 14.8	36.8 +/- 0.5
CD3B146	3.2 +/- 1.2	1.1 +/- 0.1

Example 23 Evaluation of IgG4-PAA CD123 x CD3 Bispecific Abs in Functional Cell Killing Assay

[00387] T-cell mediated cytotoxicity assay as described in Example 16 was used to evaluate the CD123 x CD3 bispecific Abs for cell lysis using T-cells from two healthy donors. For these experiments, OCI-AML5, KG-1 and JIM3 cells were used. JIM3 is a myeloma tumor line and has no CD123 expression and was used as a control. Cells were treated for 48 hours with bispecific Abs. The E:T ratio for this study was 5:1, and 2mg/mL Fc blocker was added to block Fc function.

[00388] The results of the T-cell mediated cell lysis of AML cell lines OCI-AML (Figure 31), KG-1 (Figure 32), and JIM3 (Figure 33) after 48 hr incubation at 37 °C, 5% CO₂ are shown. The MV4-11 and OCI-AML5 are CD123 expression cell lines, and the JIM3 has very little or no CD123 expression. The Effector/Target ratio for this study was 5:1. A 2 mg/mL aliquot of Fc blocker was added to block Fc function.

[00389] Results are similar to the previous cell-killing experiments with CD123 x CD3 bispecific Abs in the IgG1-AA format. Both I3RB217 (I3RB18) and I3RB218 (I3RB2) antibodies, when combined with an anti-CD3 antibody into a bispecific format, are efficacious at specifically killing CD123+ cells. Cell-killing is specific to CD123-containing cells, as demonstrated by the lack of effect on JIM3 cells. Additionally, the data allow for a clear ranking between the I3RB218 (I3RB2-based) and I3RB217 (I3RB18-based) bispecific antibodies with the I3RB217 x CD3 bispecific Abs being more potent than I3RB218 x CD3 Bispecific Abs, in agreement with previous cell killing data.

[00390] Example 24. Evaluation of Bispecific Antibodies in Receptor heterodimerization assay

[00391] The DiscoverX Receptor Dimerization assay for IL3RA/CD131 (DiscoverX 93-0969-C1) was used to evaluate the ability of the CD123 antibodies to prevent the IL3-induced heteromerization of IL3R α (CD123)/IL3R β (CD131). The CD123 and CD131 are tagged with ProLink™ (PK) or Enzyme Acceptor (EA). Upon IL3-induced activation, the proteins dimerize to form the IL3 receptor, forcing the two β -gal components to complement and create an active enzyme. Active β -gal generates a chemiluminescent signal in the presence of substrate. Anti-CD123 antibodies or bispecific antibodies that show decreasing signal with increasing antibody concentration are positive for preventing heterodimerization.

[00392] The cells were tested for increases in enzyme activity in the presence of the IL-3 ligand using PathHunter® Detection Reagents (DiscoverX) according to the manufacturer's protocol. HEK293 IL3RA-PK / CSF2RB-EA cell lines were plated in 20 uL assay media in quadruplicate on 384-well plates with 5,000 cells / well. Antibody stocks were serially diluted in 0.1% BSA / PBS such that the high concentration of compound was 10 ug / mL. The high dose was serially diluted 1:3 with 11 doses tested. 5 µl of diluted antibody was added to the wells. Cells were incubated for 1 hour at 37C. A recombinant human IL-3 stock solution at 100 µg / mL was diluted such that 5 µl of a 60 ng / mL dilution of IL-3 was added to each well. The final concentration of IL-3 used was 10 ng / mL. Cells were incubated an additional 6 hours at 37C. PathHunter Flash Detection Reagent containing lysis buffer and enzyme substrate was added to the cells, incubated 30 minutes at room temperature and read on the Envision luminometer. Data was analyzed using GraphPad Prism 6. Curves are fit using a sigmoidal dose response with variable slope (four parameter) with no constraints; fit method= least squares (normal fit).

[00393] IgG4 PAA bispecific antibodies 8747 and 7959, as well as the parental antibodies I3RB218 and I3RB217 were run in the assay. The assay was run two independent times in the presence of 10 ng/ml of IL-3 and the positive control CD123 antibody 7G3 was used as a comparator in the assay. Antibodies that contained the anti-CD123 arm I3RB18 sequence, I3RB217 and 7959, (Figure 34 C and D) were able to prevent formation of a functional IL-3 receptor in the presence of IL-3 ligand. Antibodies that contained the anti-CD123 arm I3RB2, I3RB218 and 8747 (Figure 34 A and B) did not prevent formation of functional IL-3 receptor in this assay. This correlates with previous data that showed I3RB18 could inhibit downstream signaling associated with a functional IL-3 receptor

Example 24. Evaluation of Several Bispecific Antibodies in the KG-1 tumor model

[00394] Several of the CD123 x CD3 bispecific Abs were evaluated for efficacy in the KG-1 AML murine model as previously described. The protocol was the same for this study as in Examples 16 and 17, except that frozen isolated peripheral blood mononuclear cell vials (100×10^6 cells per vial, Catalog PB009-3) obtained from HemaCare (Van Nuys, CA) were used for testing the IgG4 bispecific antibodies. NSG mice were subcutaneously inoculated with KG-1 cells, and then intravenously engrafted with human PBMCs when tumors were established (mean tumor volume = 135.7 ± 4.7 mm³). Mice were then dosed with IgG4 PAA CD123 x CD3 bispecific Abs with various affinities and corresponding control bispecific Abs at a range of doses, as described in Table 23.

Table 23. Dosing Schedule for 3rd in vivo study

Group	N	Tumor	PBMC	Treatment	Dose (μ g/animal)	Dosing Route	Dosing Schedule (Days Post-tumor Implantation)	Blood Sampling (Days Post- tumor Implantation)
1	10	+	+	PBS	0	i.v.	14, 16, 18, 21, 23	31
2	10	+	+	7959	0.1	i.v.	14, 16, 18, 21, 23	31
3	10	+	+	7959	1	i.v.	14, 16, 18, 21, 23	31
4	10	+	+	9958	0.1	i.v.	14, 16, 18, 21, 23	31
5	10	+	+	9958	1	i.v.	14, 16, 18, 21, 23	31
6	10	+	+	8747	0.1	i.v.	14, 16, 18, 21, 23	31
7	10	+	+	8747	1	i.v.	14, 16, 18, 21, 23	31
8	10	+	+	8747	10	i.v.	14, 16, 18, 21, 23	31
9	10	+	+	3978	0.1	i.v.	14, 16, 18, 21, 23	31
10	10	+	+	3978	1	i.v.	14, 16, 18, 21, 23	31
11	10	+	+	3978	10	i.v.	14, 16, 18, 21, 23	31
12	10	+	+	8876	0.1	i.v.	14, 16, 18, 21, 23	31
13	10	+	+	8876	1	i.v.	14, 16, 18, 21, 23	31
14	10	+	+	8876	10	i.v.	14, 16, 18, 21, 23	31
15	10	+	+	CD3 null 1	0.1	i.v.	14, 16, 18, 21, 23	31
16	10	+	+	CD3 null 1	1	i.v.	14, 16, 18, 21, 23	31
17	10	+	+	CD3 null 1	10	i.v.	14, 16, 18, 21, 23	31
18	10	+	+	CD3 null 2	0.1	i.v.	14, 16, 18, 21, 23	31
19	10	+	+	CD3 null 2	1	i.v.	14, 16, 18, 21, 23	31
20	10	+	+	CD3 null 2	10	i.v.	14, 16, 18, 21, 23	31

21	10	+	+	CD123 null 1	0.1	i.v.	14, 16, 18, 21, 23	31
22	10	+	+	CD123 null 1	1	i.v.	14, 16, 18, 21, 23	31
23	10	+	+	CD123 null 1	10	i.v.	14, 16, 18, 21, 23	31
24	10	+	+	CD123 null 2	0.1	i.v.	14, 16, 18, 21, 23	31
25	10	+	+	CD123 null 2	1	i.v.	14, 16, 18, 21, 23	31
26	10	+	+	CD123 null 2	10	i.v.	14, 16, 18, 21, 23	31

[00395] Results of in vivo efficacy studies with multiple CD123 x CD3 bispecific Abs are shown in Figures 35 - 42. Figures 35 - 38 show the efficacy of CD123 x CD3 IgG4-PAA bispecific Abs at various affinities and doses in KG-1 human AML xenografts. In Figure 35, bispecific Abs with high affinity CD123 and CD3 arms had significant efficacy compared to PBS and control bispecific Abs from days 25 through 36 post-tumor implantation ($p < 0.001$). Bispecific Ab 9958 at the 1 μ g dose had significant efficacy compared to 0.1 μ g, and both doses of bispecific Ab 7959 by day 36 post-tumor implantation ($p < 0.01$). This indicates high affinity CD123 and CD3 arms are necessary for pronounced efficacy in this model.

[00396] In Figure 36, bispecific Ab 3978 at the 10 μ g dose had significant efficacy compared to PBS and control bispecific Abs from day 28 ($p < 0.05$) through day 36 ($p < 0.001$) post-tumor implantation, the 1 μ g dose from day 32 ($p < 0.05$) through day 36 ($p < 0.01$) post-tumor implantation, and the 0.1 μ g dose from day 32 ($p < 0.01$) through day 36 ($p < 0.001$) post-tumor implantation. There is a dose-dependent response with this bispecific Ab, indicating a high affinity CD123 arm at a high dose can result in efficacy in this model.

[00397] In Figure 37, bispecific Ab 8747 at the 0.1 μ g dose had significant efficacy compared to PBS and control bispecific Abs from days 32 through 36 post-tumor implantation ($p < 0.001$), and compared to the 1 and 10 μ g doses by day 36 post-tumor implantation ($p < 0.001$). This indicates a high affinity CD3 arm at a low dose can result in efficacy in this model.

[00398] In Figure 38, bispecific Ab 8876 did not have significant efficacy compared to PBS and control bispecific Abs at any dose.

[00399] Figures 39 - 42 show the mean body weight change of treatment groups over time. As described previously, body weight loss is correlated with onset and severity of GVHD, which is caused by activated T cells.

[00400] Animals treated with bispecific Ab 7959 at 0.1 μ g and bispecific Ab 9958 at 1 μ g had more severe and earlier onset body weight loss than those treated with PBS, control bispecific Abs, and the other doses of bispecific Ab 7959 and bispecific Ab 9958 (Figure 39). This correlates with the significant anti-tumor efficacy seen at 1 μ g bispecific Ab 9958 (Figure 35).

[00401] Animals treated with bispecific Ab 3978 at the 10 μ g dose had more severe and earlier onset body weight loss compared with those treated with PBS and control bispecific Abs (Figure 40). The mice treated with the 1 μ g and 0.1 μ g doses followed in body weight loss in a dose-dependent manner. The dose-dependent weight loss correlates with the dose dependent anti-tumor efficacy seen in Figure 36).

[00402] Animals treated with bispecific Ab 8747 at the 0.1 μ g dose had similar body weight loss to that of the PBS-treated group, however, mice regained body weight beginning day 39 post-tumor implantation (Figure 41). There was no body weight loss with the 1 or 10 μ g doses. The weight loss seen at the 0.1 μ g dose correlates with anti-tumor efficacy seen at this dose (Figure 37).

[00403] Animals treated with bispecific Ab 8876 did not show weight loss different from that of PBS or control bispecific treated mice (Figure 42), corresponding to the lack of anti-tumor efficacy seen with this bispecific antibody (Figure 38).

[00404] In summary, the CD123 x CD3 bispecific Abs shows consistent efficacy in a CD123 expressing human AML cell line, KG-1, only in the presence of effector cells (T lymphocytes). T cell expansion was seen shortly after the dosing period, only in the presence of disease (KG-1 xenografts). Additionally, bispecific efficacy is correlated with GVHD onset as measured by body weight loss, indicating activated T lymphocytes are present. Together, these data indicate that the CD123 x CD3 bispecific has anti-tumor efficacy through the proposed mechanism of target and effector cell engagement, and T cell killing.

Example 24. *In vivo* Mouse PK studies

[00405] Test Ab articles were formulated in phosphate-buffered saline at 0.2 mg/mL. Concentrations were confirmed using the Nanodrop spectrophotometer, and then sterile-filtered with 0.2 micron syringe filters.

[00406] Transgenic animals used in these studies are derived from C57BL/6 mice. Tg32 licensed from the Jackson Laboratory (Bar Harbor) have their endogenous mouse FcRn α

gene knocked out and are transgenic with the human FcRn α gene under the control of the native human gene promoter. Tg32 hemi refer to mice hemizygous for the FcRn transgene, the latter derived by mating homozygous transgenic mice with FcRn α knockout mice. A significant correlation was observed between the PK of human antibodies and the PK in primates with the Tg32 hemi mouse model, and therefore it was used in the following PK studies to evaluate Ab half-life. All mouse breeding was done at SAGE Research Labs Boyertown, PA Facility.

[00407] For the study, 6 week old mice were used with 48 female Tg32 hemi mice injected IV with hIgG4-PAA bispecific Abs using 5 mice per group. Retro-orbital bleeds were taken at the same time points.

[00408] After sample collection, a serum analysis was conducted. Concentrations of human IgG in the serum samples were determined by an electrochemiluminescent immunoassay with the MESO Scale Discovery (MSD) format. Streptavidin MSD plates were coated with 50 μ L/well of 2 μ g/mL biotinylated F(ab')₂ goat anti hu IgG (H+L, Jackson lot 109-066-08) in Starting Block T20 (Thermo) overnight, 4 °C. Plates were washed with PBS buffer, and samples diluted in 10% mouse serum (Bioreclamations, NY) in Starting Block T20. Included on each plate was a standard curve of each test article, starting at 0.1 mg/mL with serial 2-fold dilutions. Plates were incubated for 2-3 h, RT on a shaker, washed and then incubated with 2 μ g/mL MSD-TAG (ruthenium-labeled anti-human IgG mAb, R10Z8E9, MSD) for 1 hr, RT on a shaker. Plates were washed and 200 μ L MSD Read Buffer (MSD) was added and read on the MSD Sector Imager 6000.

[00409] To determine whether the PK serum samples had notable immune titers that could affect the PK of test samples, an ELISA was performed on Maxisorb plates (Nunc) coated with the respective test article at 10 μ g/mL and incubated overnight at 4 °C. Serum samples were diluted in 1% BSA-PBS and incubated on the plates for 2-3 h with shaking at RT. Horseradish peroxidase-conjugated donkey anti-mouse IgG (Jackson ImmunoResearch) was used to detect captured antibody; followed by 3,3',5,5'-tetramethylbenzidine addition (Fitzgerald) for substrate development. Plates were read and spectrophotometer readings that were three times greater than buffer or control sera values were considered positive. Immune titers were expressed as 1/serum dilution. No immune titers were observed (data not shown).

[00410] Finally, the pharmacokinetics for the molecules was determined. Terminal half-life ($t_{1/2}$) calculations of the elimination phase for PK studies were determined using the 1-

phase exponential decay model fitted by linear regression of natural log concentration vs. time using Prism version 5.01 software (GraphPad Software, Inc.). Two phase models were ruled out because for each test article, the best-fit model was a 1-phase exponential decay model as determined by nonsignificance of the extra sum of squares F test ($p > 0.05$) for the majority of animals. The least squares nonlinear decay model was weighted by $1/\text{fitted concentration}$. Half-life calculations of the elimination phase were determined using the formula $t_{1/2} = \ln 2 / \beta$ where β is the $-\text{slope}$ of the line fitted by the least square regression analysis starting after first-dose.

[00411] In the PK study described here, the terminal half-life value for an antibody was determined by taking the average of the $t_{1/2}$ values calculated for each animal within the test group. Outliers in the studies were identified as animals either showing a mouse anti-human IgG titer greater than a 1 to 1000 about 7d after dose or an initial serum value that was more than 2-fold lower than values for other mice in the group, perhaps due to not being fully dosed.

[00412] The human PK predictions from the mouse data were based on observed half-life differences in huFcRn-transgenic mice vs humans for a panel of eight human IgG antibodies whose clearance was believed to not be significantly impacted by target binding in either mice or humans. Based on those analyses, it was estimated that the terminal half-life in humans for the CD123 x CD3 bispecific Abs would be 2-4-fold longer than what was observed in the huFcRn-transgenic mice, an extrapolation that assumes the influence of target binding on clearance is comparable in mice and humans. Table 24 summarizes the observed mouse half-life values for the various Bispecific antibody variants and the corresponding predicted human values which reflect that assumption. Because the well-known human PK prediction method based on allometric scaling across species has not been validated using the mouse PK data, allometric scaling was not used for the predictions. The PK results are shown in Figure 43 with the serum concentration vs. time. PK profiles display a linear decline of serum concentration over the course of 28 days. The estimated mouse half-life values for all the CD123 x CD3 Bispecific antibody Abs were similar, between 5.2-6.6 days. Minimal immune titers ($<1:40$) were observed in all groups. The mouse PK data (with mean $\pm$ standard deviation) along with predicted human clearance and human half-life values are summarized in Tables 24. The human half-life prediction assumes that target binding in humans is not greater than in mice.

[00413] The IgG4-PAA bispecific antibody Abs showed similar values between the I3RB2 and I3RB18 groups in mice. Mouse half-life calculations of the elimination phase were determined using the 1-phase exponential decay model fitted by linear regression of natural log concentrations vs time as described. The half-life values calculated for the eight Bispecific

antibodies Abs in Tg32 hemi mice were: 3978, 6.6 +/- 0.7 days; 7955, 5.2 +/- 0.4 days; 7959, 6.6 +/- 0.6 days; 9958, 6.4 +/- 0.7 days; 8876, 4.1 +/- 0.7 days; 4435, 5.4 +/- 1.0 days; 8747, 6.4 +/- 0.4 days; 5466, 5.6 +/- 0.1 days. The human PK predictions from the mouse data were based on observed half-life differences in huFcRn-transgenic mice vs humans. Based on those analyses, the estimated terminal half-life in humans for the CD123xCD3 bispecific antibodies would be 2 to 4-fold longer than what was observed in the huFcRn-transgenic mice, assuming the influence of target binding on clearance is comparable in mice and humans. Table 24 summarizes the observed mouse half-life values for the Bispecific antibody variants and the corresponding predicted human values which reflect that assumption.

Table 24. Summary of PK of CD123 x CD3 IgG4-PAA Bispecific Abs

Bispecific Ab	Animal No.	T1/2 (day)	Mean calc. T1/2 (day)	Predicted hT1/2 (day)
3978	2	6.71	6.59±0.66	13.2 – 26.4
	5	7.08		
	7	5.44		
	22	7.02		
	28	6.71		
7955	1	4.85	5.24±0.43	10.5 – 21.0
	6	4.80		
	8	5.31		
	9	5.39		
	10	5.85		
7959	3	6.31	6.63±0.57	13.2 – 26.4
	12	7.53		
	13	6.28		
	16	6.86		
	33	6.16		
9958	4	7.15	6.36±0.68	12.7 - 25.4
	15	5.60		
	18	6.88		
	19	5.77		
	20	6.40		
8876	21	4.65	5.19±0.70	10.4 – 20.8
	22	6.05		
	23	4.40		
	24	5.10		
	35	5.73		
4435	17	5.39	5.42±0.95	10.8 – 21.6
	27	3.96		
	29	5.66		
	30	5.47		
	36	6.60		

8747	14	6.38	6.37±0.37	12.7 - 25.4
	25	6.49		
	31	5.78		
	32	6.80		
	34	6.41		
5466	26	4.81	5.64±0.96	11.3 – 22.6
	37	5.68		
	38	4.53		
	39	6.61		
	40	6.54		

[00414] Results of mouse PK studies with CD123XCD3 bispecific antibodies show that the observed t1/2 values in Tg32 hemi mice compare favorably to 8 clinical antibodies profiled in the same manner.(Tam, et al, MAbs (2013) 5(3):3987-405).

What is Claimed:

1. An isolated antibody, or antigen-binding fragment thereof, that binds immunospecifically to CD123 SP2 (IL-3Ra) and CD123 SP1 (IL3Ra)
2. The isolated antibody, or an antigen-binding fragment of claim 1, comprising:
 - a. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 006, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 007, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 008,
 - b. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 012, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 013, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 014,
 - c. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 027, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 028, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 029,
 - d. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 051, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 052, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 053,
 - e. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 055, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 056, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 057, or
 - f. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 085, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 086.
3. The isolated antibody, or an antigen-binding fragment of claim 2, wherein
 - a. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 006, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 007, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 008 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 009, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 010, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 011;

b. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 012, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 013, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 014 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 015, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 016, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 017;

c. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 027, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 028, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 029 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

d. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 051, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 052, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 053 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 024, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 025, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 054;

e. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 055, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 056, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 057 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 058, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 059, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032; or

f. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 085, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 086 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 087, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 067, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 088.

4. The antibody or antigen-binding fragment of any one of claims 1 to 3, wherein the heavy chain of the antibody of (a) comprises the amino acid sequence of SEQ ID NO: 119; the heavy chain of the antibody of (b) comprises the amino acid sequence of SEQ ID NO: 120; the heavy chain of the antibody of (c) comprises the amino acid sequence of SEQ ID NO: 123; the heavy chain of the antibody of (d) comprises the amino acid sequence of SEQ ID NO: 136; the heavy chain of the antibody of (e) comprises the amino acid sequence of SEQ ID NO: 137; or the heavy chain of the antibody of (f) comprises the amino acid sequence of SEQ ID NO: 148.
5. The antibody or antigen-binding fragment of any one of claims 1 to 4, wherein the light chain of the antibody of (a) comprises the amino acid sequence of SEQ ID NO: 164; the light chain of the antibody of (b) comprises the amino acid sequence of SEQ ID NO: 165 the light chain of the antibody of (c) comprises the amino acid sequence of SEQ ID NO: 167; the light chain of the antibody of (d) comprises the amino acid sequence of SEQ ID NO: 168; the light chain of the antibody of (e) comprises the amino acid sequence of SEQ ID NO: 169; or the light chain of the antibody of (f) comprises the amino acid sequence of SEQ ID NO: 176.
6. The antibody or antigen-binding fragment of claim 1, wherein the heavy chain of the antibody comprises the amino acid sequence of SEQ ID NO: 120, and the light chain of the antibody comprises the amino acid sequence of SEQ ID NO: 165.
7. The antibody or antigen-binding fragment of claim 1, wherein the heavy chain of the antibody comprises the amino acid sequence of SEQ ID NO: 136 and light chain of the antibody comprises the amino acid sequence of SEQ ID NO: 168.
8. The antibody or antigen-binding fragment of any one of claims 1 to 7 wherein the antibody or antigen-binding fragment thereof binds to a polypeptide having the amino acid sequence of SEQ ID NO: 002.
9. The antibody or antigen-binding fragment of any one of claims 1 to 8 wherein the antibody or antigen-binding fragment is a human antibody or antigen-binding fragment.
10. The antibody or antigen-binding fragment of any one of claims 1 to 9 wherein the antibody or antigen-binding fragment is recombinant.
11. The antigen binding fragment of any one of claims 1 to 10 wherein the antigen binding fragment is a Fab fragment, a Fab2 fragment, or a single chain antibody.

12. The antibody or antigen-binding fragment of any one of claims 1 to 11 wherein the antibody or antigen-binding fragment thereof immunospecifically binds CD123 with an affinity of at least 5 nM as measured by surface plasmon resonance.
13. The antibody or antigen-binding fragment of any one of claims 1 to 12 wherein the antibody or antigen-binding fragment thereof immunospecifically binds CD123 SP1 or CD123 SP2 with an affinity of at least 5 nM as measured by surface plasmon resonance.
14. The antibody or antigen-binding fragment of any one of claims 1 to 13 wherein the antibody or antigen-binding fragment thereof immunospecifically binds CD123 SP1 and CD123 SP2 with an affinity of at least 5 nM as measured by surface plasmon resonance.
15. The antibody or antigen-binding fragment of any one of claims 1 to 14 wherein the antibody or antigen-binding fragment thereof are of IgG1, IgG2, IgG3, or IgG4 isotype.
16. The antibody or antigen-binding fragment of any of claims 1 to 15 is IgG1 or IgG4 isotype.
17. The antibody or antigen-binding fragment of any of claims 1 to 16 is IgG1 isotype.
18. The antibody of claim 17 wherein the IgG1 has a K409R substitution in its Fc region.
19. The antibody of claim 18 further comprising a L234A substitution and/or a L235A substitution in its Fc region.
20. The antibody or antigen-binding fragment of any one of claims 1 to 16 is IgG4 isotype.
21. The antibody or antigen-binding fragment of any one of claims 1 to 20 wherein the antibody or antigen-binding fragment thereof immunospecifically binds human CD123 and cynomolgus monkey CD123.
22. The antibody or antigen-binding fragment of claim 21 wherein the antibody or antigen-binding fragment immunospecifically binds cynomolgus monkey CD123 with a binding affinity within 10 times the binding affinity of said antibody or fragment for human SP1.
23. The antibody or antigen-binding fragment of claim 21 wherein the antibody or antigen-binding fragment immunospecifically binds cynomolgus monkey CD123 with a binding affinity within 5 times the binding affinity of said antibody or fragment for human SP1.

24. The isolated antibody, or an antigen-binding fragment of claim 1, wherein said antibody competes for binding to CD123 SP2 (IL3-R α) with an antibody comprising:

a. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 006, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 007, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 008,

b. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 012, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 013, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 014,

c. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 026, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 027, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 028,

d. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 051, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 052, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 053,

e. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 055, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 056, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 057, or

f. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 085, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 086.

25. The isolated antibody, or an antigen-binding fragment of claim 24, wherein the CD123 SP2 (IL3-R α) has the amino acid sequence of SEQ ID NO: 002.

26. An isolated antibody, or antigen-binding fragment thereof, that immunospecifically binds human CD123 SP1, said antibody comprising:

a. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 018, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 019, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 020;

- b. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 021, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 023;
- c. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 035;
- d. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 036;
- e. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 037;
- f. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 038;
- g. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 040, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 041;
- h. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 042;
- i. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 043;
- j. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 044, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 045;
- k. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 046, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 047;

- l. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 048;
- m. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 049;
- n. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 050;
- o. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 060;
- p. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 061;
- q. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 065;
- r. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 069;
- s. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 071;
- t. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 072, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 073;
- u. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 074;

- v. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 078;
- w. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 081;
- x. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 082;
- y. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 089, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 090;
- z. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 093;
- aa. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 094;
- bb. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 097;
- cc. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 100;
- dd. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 102;
- ee. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 103;

- ff. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 106;
- gg. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 107;
- hh. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 111;
- ii. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 114, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 115;
- jj. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 114, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 116;
- kk. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 117, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 013, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 118;
- ll. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 051, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 052, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 053; or
- mm. a heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, a heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and a heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 042.
27. The antibody, or antigen-binding fragment thereof, of claim 26, wherein
- a. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 018, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 019, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 020 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 009, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 010, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 011;

b. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 021, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 023 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 024, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 025, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 026;

c. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 035 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

d. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 036 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

e. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 037 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

f. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 038 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

g. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 040, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 041 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

h. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 042 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

i. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 043 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

j. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 044, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 045 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 015, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 016, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 017;

k. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 046, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 047 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

l. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 048 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

m. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 049 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

n. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 050 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

o. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 060 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

p. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 061 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 062, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 063, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 064;

q. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 065 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 066, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 067, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 068;

r. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 069 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 070;

s. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 071 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

t. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 072, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 073 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

u. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 074 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 075, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 076, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 077;

v. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 078 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 079, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 063, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 080;

w. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 081 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

x. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 082 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 083, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 084;

y. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 089, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 090 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 091, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 076, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 092;

z. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 093 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

aa. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 094 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 095, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 076, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 096;

bb. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 097 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 098, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 067, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 099;

cc. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033, said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 100 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 101;

dd. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 102 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

ee. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 103 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 104, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 105;

ff. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 039 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 106 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 30, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032;

gg. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 107 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 108, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 109, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 110;

hh. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 111 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 112, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 076, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 113;

ii. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 114 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 115 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 024, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 025, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 026;

jj. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 114 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 022, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 116 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 015, a light chain

CDR2 having the amino acid sequence of SEQ ID NO: 016, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 017;

kk. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 117 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 013, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 118 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 015 light chain CDR2 having the amino acid sequence of SEQ ID NO: 016 and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 017;

ll. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 051, heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 052 and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 053 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030 light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032; or

mm. said antibody comprising said heavy chain CDR1 having the amino acid sequence of SEQ ID NO: 033 said heavy chain CDR2 having the amino acid sequence of SEQ ID NO: 034, and said heavy chain CDR3 having the amino acid sequence of SEQ ID NO: 042 further comprises a light chain CDR1 having the amino acid sequence of SEQ ID NO: 030, a light chain CDR2 having the amino acid sequence of SEQ ID NO: 031, and a light chain CDR3 having the amino acid sequence of SEQ ID NO: 032.

28. An isolated cell expressing the antibody or antigen-binding fragment of any one of claims 1 to 27.

29. The cell of claim 28 wherein the cell is a hybridoma.

30. The cell of claim 28 wherein the antibody is recombinantly produced.

31. An isolated CD123 (IL3-R α) x CD3 bispecific antibody comprising:

- a) a first heavy chain (HC1);
- b) a second heavy chain (HC2);
- c) a first light chain (LC1); and
- d) a second light chain (LC2),

wherein the HC1 and the LC1 pair to form a first antigen-binding site that immunospecifically binds CD123 (IL3-R α), and the HC2 and the LC2 pair to form a second antigen-binding site that immunospecifically binds CD3, or a CD123 (IL3-R α) x CD3 -bispecific binding fragment thereof.

32. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 31 wherein the CD123 (IL3-R α) is CD123 (IL3-R α) SP1.

33. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 31 wherein the CD123 (IL3-R α) is CD123 (IL3-R α) SP2.

34. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 31 wherein the CD123 (IL3-R α) is CD123 (IL3-R α) SP1 and CD123 (IL3-R α) SP2.

35. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any of claims 31 to 34 wherein the antibody or bispecific binding fragment is IgG1, IgG2, IgG3, or IgG4 isotype.

36. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any of claims 31 to 35 wherein the antibody or bispecific binding fragment is IgG1 or IgG4 isotype.

37. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any of claims 31 to 36 wherein the antibody or bispecific binding fragment is IgG1 isotype.

38. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any of claims 31 to 36 wherein the antibody or bispecific binding fragment is IgG4 isotype.

39. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 37 wherein HC1 comprises SEQ ID NO: 203 and LC1 comprises SEQ ID NO: 204.

40. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39 wherein HC2 comprises SEQ ID NO: 193 and LC2 comprises SEQ ID NO: 194.

41. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39 wherein HC2 comprises SEQ ID NO: 195 and LC2 comprises SEQ ID NO: 196.

42. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39 wherein HC2 comprises SEQ ID NO: 197 and LC2 comprises SEQ ID NO: 198.

43. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39 wherein HC2 comprises SEQ ID NO: 199 and LC2 comprises SEQ ID NO: 200.
44. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39 wherein HC2 comprises SEQ ID NO: 201 and LC2 comprises SEQ ID NO: 202.
45. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31, to 37 wherein HC1 comprises SEQ ID NO: 205 and LC1 comprises SEQ ID NO: 206.
46. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 193 and LC2 comprises SEQ ID NO: 194.
47. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 195 and LC2 comprises SEQ ID NO: 196.
48. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 197 and LC2 comprises SEQ ID NO: 198.
49. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 199 and LC2 comprises SEQ ID NO: 200.
50. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 201 and LC2 comprises SEQ ID NO: 202.
51. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 31 to 50 wherein the antibody or bispecific binding fragment immunospecifically binds CD123 with an affinity of at least 5 nM as measured by surface plasmon resonance.
52. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39, 40, 45, 46, and 51 wherein the antibody or bispecific binding fragment immunospecifically binds CD3 with an affinity of at least 15 nM as measured by surface plasmon resonance.
53. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 39, 41 to 44, 47 to 51 wherein the antibody or bispecific binding fragment immunospecifically binds CD3 with an affinity of at least 75 nM as measured by surface plasmon resonance.
54. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 36, and 38 wherein HC1 comprises SEQ ID NO: 220 and LC1 comprises SEQ ID NO: 221.

55. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54 wherein HC2 comprises SEQ ID NO: 210 and LC2 comprises SEQ ID NO: 211.
56. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54 wherein HC2 comprises SEQ ID NO: 212 and LC2 comprises SEQ ID NO: 213.
57. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54 wherein HC2 comprises SEQ ID NO: 214 and LC2 comprises SEQ ID NO: 215.
58. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54 wherein HC2 comprises SEQ ID NO: 216 and LC2 comprises SEQ ID NO: 217.
59. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 36, and 38 wherein HC1 comprises SEQ ID NO: 218 and LC1 comprises SEQ ID NO: 219.
60. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 59 wherein HC2 comprises SEQ ID NO: 210 and LC2 comprises SEQ ID NO: 211.
61. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 59 wherein HC2 comprises SEQ ID NO: 212 and LC2 comprises SEQ ID NO: 213.
62. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 59 wherein HC2 comprises SEQ ID NO: 214 and LC2 comprises SEQ ID NO: 215.
63. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 45 wherein HC2 comprises SEQ ID NO: 216 and LC2 comprises SEQ ID NO: 217.
64. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54 to 63 wherein the antibody or bispecific binding fragment immunospecifically binds CD123 with an affinity of at least 5 nM as measured by surface plasmon resonance.
65. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 54, 55, 59, 60, and 64 wherein the antibody or bispecific binding fragment immunospecifically binds CD3 with an affinity of at least 15 nM as measured by surface plasmon resonance.
66. The CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of claim 59, 56 to 58, 61 to 64 wherein the antibody or bispecific binding fragment immunospecifically binds CD3 with an affinity of at least 75 nM as measured by surface plasmon resonance.

67. An isolated cell expressing the antibody or bispecific binding fragment of any one of claims 31 to 53.
68. The cell of claim 67 wherein the cell is a hybridoma.
69. The cell of claim 67 wherein the antibody or bispecific binding fragment is recombinantly produced.
70. A method for treating a subject having cancer, said method comprising:
administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 51 to a patient in need thereof for a time sufficient to treat the cancer.
71. A method for inhibiting growth or proliferation of cancer cells, said method comprising:
administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 51 to inhibit the growth or proliferation of cancer cells.
72. A method of redirecting a T cell to a CD123-expressing cancer cell, said method comprising:
administering a therapeutically effective amount of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 51 to redirect a T cell to a cancer.
73. The method of claim 70, 71, or 72 wherein the cancer is a hematological cancer.
74. The method of claim 73 wherein the hematological cancer is a CD123 (IL3-R α)-expressing hematological cancer.
75. The method of claim 74 wherein the CD123 (IL3-R α)-expressing hematological cancer, is acute myeloid leukemia (AML), myelodysplastic syndrome (MDS), acute lymphocytic leukemia (ALL), diffuse large B-cell lymphoma (DLBCL), chronic myeloid leukemia (CML) or blastic plasmacytoid dendritic cell neoplasm (DPDCN).
76. The method of claim 75 wherein the CD123 (IL3-R α)-expressing hematological cancer is AML.
77. The method of claim 70 comprising administering a second therapeutic agent.

78. The method of claim 77 wherein the second therapeutic agent is a chemotherapeutic agent or a targeted anti-cancer therapy.
79. The method of claim 78 wherein the chemotherapeutic agent is cytarabine, an anthracycline, histamine dihydrochloride, or interleukin 2.
80. The method of claim 77 wherein the second therapeutic agent is administered to said subject simultaneously with, sequentially, or separately from the bispecific antibody.
81. A pharmaceutical composition comprising the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 66 and a pharmaceutically acceptable carrier.
82. A method for generating the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 66 by culturing the cell of any one of claims 54 to 56.
83. An isolated synthetic polynucleotide encoding the HC1, the HC2, the LC1 or the LC2 of the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 66.
84. A kit comprising the CD123 (IL3-R α) x CD3 bispecific antibody or bispecific binding fragment of any one of claims 31 to 66 and packaging for the same.
85. An isolated antibody, or an antigen-binding fragment thereof, that binds immunospecifically to a polypeptide comprising the amino acid sequence of SEQ ID NO: 226.
86. An isolated antibody, or an antigen-binding fragment thereof, that binds immunospecifically to a polypeptide comprising the amino acid sequences of SEQ ID NO: 234 and SEQ ID NO: 228.

Figure 1.




Figure 2A)

a)




Figure 2 B)




Figure 2 C)




Figure 3.






Figure 4.


A)(Neat)



B) (1 ug/ml)



C) Dose dependence



	B18 + IL-3	7G3 + IL-3
IC ₅₀ , ng/ml	75.39	160.3

Figure 5.







Figure 6.**A) Ru-labeled I3RB2****B) Ru-labeled I3RB18**


Figure 7.

A)



Numbering begins at amino acid 19 of SP2

B)



Numbering begins at amino acid 19 of SP2

Figure 8.

Numbering: ovals refer to residues of CD123 SP2, rectangles refer to residues of SEQ ID NO:230




Figure 9.





Figure 10.**Light Chain (SEQ ID NO:4)**

```
      10      20      30      40      50      60
      . |      . |      . |      . |      . |      . |
QAVVTQESALTTSPGETVTLTCRSSTGAVTTSNYANWVQEKPDHLFTGLIGGTNKRAPGV
      70      80      90     100     110     120
      . |      . |      . |      . |      . |      . |
PARFSGSLIGDKAALTITGAQTEDEAIYFCALWYSNLWVFGGGTKLTVLGQPKSSPSVTL
      130     140     150     160     170     180
      . |      . |      . |      . |      . |      . |
FPPSSEETNKATLVCTITDFYPGVVTVDWKVDGTPVTQGMETTQPSKQSNNKYMASSY
      190     200     210
      . |      . |      . |
LTLTARAWERHSSYSCQVTHEGHTVEKSLSRADCS
```

Heavy Chain (SEQ ID NO:5)

```
      10      20      30      40      50      60
      . |      . |      . |      . |      . |      . |
EVKLLESGGGLVQPKGSLKLSCAASGFTFNTYAMNWVRQAPGKGLEWVARIRSKYNNYAT
      70      80      90     100     110     120
      . |      . |      . |      . |      . |      . |
YYADSVKDRFTISRDDSQSILYLQMNNLKTEDTAMYYCVRHGNFGNSYVSWFAYWGQGTL
      130     140     150     160     170     180
      . |      . |      . |      . |      . |      . |
VTVSAATTTAPSVYPLVPGCSDTSGSSVTLGCLVKGYFPEPVTVKWNYGALSSGVRTVSS
      190     200     210     220     230     240
      . |      . |      . |      . |      . |      . |
VLQSAFYSLSSLVTVPSSTWPSQTVICNVAHPASKTELKRIEPRIPKPPSTPPGSSCPPG
      250     260     270     280     290     300
      . |      . |      . |      . |      . |      . |
NILGGPSVFIFPPKPKDALMISLTPKVTCVVVDVSEDDPDVHVSFVFDNKEVHTAWTQPR
      310     320     330     340     350     360
      . |      . |      . |      . |      . |      . |
EAQYNSTFRVVSALPIQHQDWMRGKEFKCKVNNKALPAPIERTISKPKGRAQTPQVYTIP
      370     380     390     400     410     420
      . |      . |      . |      . |      . |      . |
PPREQMSKKKVSLTCLVTNFFSEAISVEWERNGELEQDYKNTPPILDSGTYFLYSKLTV
      430     440     450
      . |      . |      . |
DTDSWLQGEIFTCSVVHEALHNHHTQKNLSRSPGK
```

Fig. 11.



VH

	10	20	30	40	50	60
sp34	EVKLL	ESGGGLVQ	PKGSLKL	SCAASG	FTFNTY	AMNWVRQAPGKGLEWVARIRSKYNNYATY
CD3H141	EV QLV ESGGGLVQ	PG SL RL SCAASG	FTFNTY	AMNWVRQAPGKGLEWV A IRIRSKYNNYATY		
CD3H142	EV QL LESGGGLVQ	PG SL RL SCAASG	FTFNTY	AMNWVRQAPGKGLEWV A IRIRSKYNNYATY		
CD3H143	EV QL LESGGGLVQ	PG SL RL SCAASG	FTFNTY	AMNWVRQAPGKGLEWV A IRIRSKYNNYATY		
CD3H144	EV QLV ESGGGLVQ	PG SLKLSCAASG	FTFNTY	AMNWVRQ AS GKGLEWV G IRIRSKYNGYATY		

	70	80	90	100	110	120
sp34	YADSVKDRFTISRDDS	QSILYLQMN	NLKTEDT	AMYYCVRHGN	FGNSYVSWFAYWGQGT	LVTVSA
CD3H141	Y AAS VK G RFTISRDDS	SKNS LYLQMN S LKTEDT	AVYYC ARHGN	FGNSYVSWFAYWGQGT	LVTVSS	
CD3H142	YADSVK G RFTISRDN SKNT	LYLQMN SL RAEDT	AVYYC AKHGN	FGNSYVSWFAYWGQGT	LVTVSS	
CD3H143	YADSVK G RFTISRDN SKNT	LYLQMN SL RAEDT	AVYYC VKHGN	FGNSYVSWFAYWGQGT	LVTVSS	
CD3H144	Y AAS VK G RFTISRDDS	SKNTA YLQMN S LKTEDT	AVYYC TRHGN	FGNSYVSWFAYWGQGT	LVTVSS	

VL

	10	20	30	40	50	60
sp34	QAVVTQES-ALTTSPGETVT	LT CR SSTGAVTTS	SNYANWVQEKPDHLFTGLIG	GT NKRAPGV		
CD3L63	QAVVTQEP- SLT VSPGGT	VTLT CR SSTGAVTTS	SNYANW VQ QKP GQ APRGLI	G GTNKRAP GT		
CD3L64	Q SV LTQ PP - SVS AAPG Q KVT I S	CR SSTGAVTTS	SNYANW VQ QL P GTAPKGLI	G GTNKRAP G I		
CD3L66	Q T VVTQEP- SLT VSPGGT	VTLT CR SSTGAVTTS	SNYANW VQ QKP GQ APRGLI	G GTNKRAP GT		

	70	80	90	100	110
sp34	PARFSGSLIGDKAALTITGAQTEDEAIYFCALWYSNLWVF	GGG	TKLTVL		
CD3L63	PARFSGSL LGG KAA L TL S G	AQ P EDEA EYY CALWYSNLWVF	GGG	TKLTVL...	
CD3L64	PDRFSGS KSG TS A T L GIT G LQT G DEAD Y CALWYSNLWVF	GGG	TKLTVL...		
CD3L66	PARFSGSL LGG KAA L TL S GV Q P	EDEA EYY CALWYSNLWVF	GGG	TKLTVL...	

Figure 13.




Figure 14.




Fig. 15.




Fig. 16.





Figure 17

A) Human



B) Cynomolgous monkey





Figure 18.

A) MV4-11, donor M6587



B) MV4-11, donor M7020





Figure 19.

A) OCI-M2, Donor M6587



B) OCI-M2, Donor M7020






Figure 20.**A) OCI-AML cell line Donor M6587****B) OCI-AML cell line Donor 7020**

Figure 21.





Figure 22.

A) CD45+



B) CD8+/CD4+





Figure 23.

A)CD45+



B)CD8+/CD4+




Figure 24.




Figure 25.





Figure 26.

A. CD45 +



B. CD8+/CD4+





Figure 27.

A) CD45 +



B) CD8+/CD4+




Figure 28.





Figure 29.

A)



B)





Figure 30.

A)



B)





Figure 31.

A)



B)






Figure 32.


A)



B)



IgG4-PAA
JIM3 +Fc Blocker (48hrs)
Donor ID:M6948



IgG4-PAA
JIM3 +Fc Blocker (48hrs)
Donor ID:M6521




Figure 34.






Figure 35.








Figure 38.




Figure 39.




Figure 40.




Figure 41.




Figure 42.




Figure 43.





Figure 1.



Comments for pDisplay™:
5325 nucleotides