(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
23 November 2006 (23.11.2006)

PCT

(10) International Publication Number
WO 2006/124710 A1

(51) International Patent Classification:
C07C 261/00 (2006.01)

(21) International Application Number:
PCT/US2006/018613

(22) International Filing Date: 15 May 2006 (15.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/682,364 18 May 2005 (18.05.2005) US

(71) Applicant *(for all designated States except US)*: **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **ALI, Amjad** [GB/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **GREENLEE, Mark, L.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **MCVEAN, Carol, A.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **MEISSNER, Robert, S.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **TAYLOR, Gayle, E.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2006/124710 A1

(54) Title: D-HOMOANDROSTA-17-YL-CARBAMATE DERIVATIVES AS SELECTIVE GLUCOCORTICOID RECEPTOR LIGANDS

(57) Abstract: The present invention is directed to D-homoandrosta-17-yl-carbamate derivatives as selective glucocorticoid receptor ligands useful for treating a variety of autoimmune and inflammatory diseases or conditions. Pharmaceutical compositions and methods of use are also included.

TITLE OF THE INVENTION

D-HOMOANDROSTA-17-YL-CARBAMATE DERIVATIVES AS SELECTIVE GLUCOCORTICOID RECEPTOR LIGANDS

BACKGROUND OF THE INVENTION


Intracellular receptors (IR's) are a class of structurally related proteins involved in the regulation of gene expression. The steroid hormone receptors are a subset of this superfamily whose natural ligands are typically comprised of endogenous steroids such as estradiol, progesterone, and cortisol. Man-made ligands to these receptors play an important role in human health and, of these receptors, the glucocorticoid receptor has an essential role in regulating human physiology and immune response. Steroids that interact with the glucocorticoid receptor have been shown to be potent anti-inflammatory agents. The present invention is directed to a novel class of compounds that are selective glucocorticoid receptor modulators that have potent anti-inflammatory and immunosuppressive activity and possess advantages over steroidal glucocorticoid ligands with respect to side effects, efficacy, toxicity and/or metabolism.

SUMMARY OF THE INVENTION

The present invention is directed to D-homoandrosta-17-yl-carbamate derivatives as selective glucocorticoid receptor ligands useful for treating a variety of autoimmune and inflammatory diseases or conditions. Pharmaceutical compositions and methods of use are also included.

DETAILED DESCRIPTION OF THE INVENTION

The invention encompasses a genus of compounds of Formula I



or a pharmaceutically acceptable salt thereof, wherein

R^1 and R^2 are independently selected from the group consisting of: R^a, -C(O)-R^a, and –C(O)-N(R^a)R^a;

R^3 and R^4 are independently selected from the group consisting of: hydrogen, C_{1-10}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, C_{3-10}cycloalkyl, aryl, hetreoaryl, heterocyclyl, and C_{1-10}fluoroalkyl; and

each Ra is independently selected from the group consisting of: hydrogen, C_{1-10}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, C_{3-6}cycloalkyl, aryl, hetreoaryl, heterocyclyl, and C_{1-10}fluoroalkyl.

Within the genus, the invention encompasses a sub-genus of compounds of Formula I wherein R^2 is selected from hydrogen and CH_3-C(O)-.

Also within the genus, the invention encompasses a sub-genus of compounds of Formula I wherein R^1 is selected from hydrogen and CF_3-(CO)-.

Also within the genus, the invention encompasses a sub-genus of compounds of Formula I wherein R^3 is hydrogen. Within this sub-genus, the invention encompasses a class of compounds of Formula I wherein R^4 is selected from C_{1-10}alkyl and C_{3-10}cycloalkyl.

Also within the genus, the invention encompasses a sub-genus of compounds of Formula I wherein: R^1 is selected from hydrogen or CF_3-(CO)-; R^2 is selected from hydrogen and CH_3-C(O)-; R^3 is hydrogen; and R^4 is selected from C_{1-10}alkyl and C_{3-10}cycloalkyl.

Within this sub-genus, the invention encompasses a class of compounds of Formula I wherein R^4 is C_{1-6}alkyl.

Also within this sub-genus, the invention encompasses a class of compounds of Formula I wherein R^4 is selected from: cyclopentyl, cyclohexyl and adamantanyl.

Also within this sub-genus, the invention encompasses a class of compounds of Formula I wherein R^1 and R^2 are hydrogen.

Illustrating the invention are the following compounds:

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-pentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-methyl-pentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(3-methyl-2-propyl)carbamate,

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclopentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclohexyl)carbamate

or a pharmaceutically acceptable salt of any of the above.

Another embodiment of the invention encompasses a pharmaceutical composition comprising a compound of Formula I in combination with a pharmaceutically acceptable carrier.

Another embodiment of the invention encompasses a method for treating a glucocorticoid receptor mediated disease or condition in a mammalian patient in need of such treatment comprising administering the patient a compoud of Formula I in an amount that is effective for treating the glucocorticoid receptor mediated disease or condition.

Within this embodiment is encompassed the above method wherein the glucocorticoid receptor mediated disease or condition is selected from the group consisting of: tissue rejection, leukemias, lymphomas, Cushing's syndrome, acute adrenal insufficiency, congenital adrenal hyperplasia, rheumatic fever, polyarteritis nodosa, granulomatous polyarteritis, inhibition of myeloid cell lines, immune proliferation/apoptosis, HPA axis suppression and regulation, hypercortisolemia, stroke and spinal cord injury, hypercalcemia, hypergylcemia, acute adrenal insufficiency, chronic primary adrenal insufficiency, secondary adrenal insufficiency, congenital adrenal hyperplasia, cerebral edema, thrombocytopenia, Little's syndrome, obesity, metabolic syndrome, inflammatory bowel disease, systemic lupus erythematosus, polyartitis nodosa, Wegener's granulomatosis, giant cell arteritis,

rheumatoid arthritis, juvenile rheumatoid arthritis, uveitis, hay fever, allergic rhinitis, urticaria, angioneurotic edema, chronic obstructive pulmonary disease, asthma, tendonitis, bursitis, Crohn's disease, ulcerative colitis, autoimmune chronic active hepatitis, organ transplantation, hepatitis, cirrhosis, inflammatory scalp alopecia, panniculitis, psoriasis, discoid lupus erythematosus, inflamed cysts, atopic dermatitis, pyoderma gangrenosum, pemphigus vulgaris, buflous pernphigoid, systemic lupus erythematosus, dermatomyositis, herpes gestationis, eosinophilic fasciitis, relapsing polychondritis, inflammatory vasculitis, sarcoidosis, Sweet's disease, type I reactive leprosy, capillary hemangiomas, contact dermatitis, atopic dermatitis, lichen planus, exfoliative dermatitus, erythema nodosum, acne, hirsutism, toxic epidermal necrolysis, erythema multiform, cutaneous T-cell lymphoma, Human Immunodeficiency Virus (HIV), cell apoptosis, cancer, Kaposi's sarcoma, retinitis pigmentosa, cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, sleep disorders, and anxiety.

Another embodiment of the invention encompasses a method of selectively modulating the activation, repression, agonism and antagonism effects of the glucocorticoid receptor in a mammal comprising administering to the mammal a compound of Formula I in an amount that is effective to modulate the glucocorticoid receptor.

Exemplifying the invention are the compounds of the Examples disclosed hereunder.

Definitions

The invention is described using the following definitions unless otherwise indicated.

"Alkyl", as well as other groups having the prefix "alk", such as alkoxy, alkanoyl, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

"Fluoroalkyl" means alkyl as defined above wherein one or more of the hydrogen atoms are replaced with a fluoro atom, up to the maximum number of substitutable positions.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

"Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

"Cycloalkyl" means mono-, bi- or tri-cyclic saturated carbocyclic rings having the indicated number of carbon atoms. The term also includes monocyclic rings fused to an aryl group in

which the point of attachment is on the non-aromatic portion. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, decahydronaphthyl, indanyl, adamantanyl and the like.

"Aryl" means mono- or bicyclic aromatic rings containing only carbon atoms. The term also includes aryl group fused to a monocyclic cycloalkyl or monocyclic heterocyclyl group in which the point of attachment is on the aromatic portion. Examples of aryl include phenyl, naphthyl, indanyl, indenyl, tetrahydronaphthyl, 2,3-dihydrobenzofuranyl, dihydrobenzopyranyl, 1,4-benzodioxanyl, and the like.

"Heteroaryl" means a mono- or bicyclic aromatic ring containing at least one heteroatom selected from N, O and S, with each ring containing 5 to 6 atoms. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, furo(2,3-b)pyridyl, quinolyl, indolyl, isoquinolyl, and the like.

"Heterocyclyl" means mono- or bicyclic saturated rings containing at least one heteroatom selected from N, S and O, each of said ring having from 3 to 10 atoms in which the point of attachment may be carbon or nitrogen. The term also includes monocyclic heterocycle fused to an aryl or heteroaryl group in which the point of attachment is on the non-aromatic portion. Examples of "heterocyclyl" include pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, 2,3-dihydrofuro(2,3-b)pyridyl, benzoxazinyl, tetrahydrohydroquinolinyl, tetrahydroisoquinolinyl, dihydroindolyl, and the like. The term also includes partially unsaturated monocyclic rings that are not aromatic, such as 2- or 4-pyridones attached through the nitrogen or N-substituted-(1H,3H)-pyrimidine-2,4-diones (N-substituted uracils).

Abbreviations

The following abbreviations have the indicated meanings:

AIBN	=	2.2'-azobisisobutyronitrile
B.P.	=	benzoyl peroxide
Bn	=	benzyl
CCl_4	=	carbon tetrachloride
D	=	$-O(CH_2)_3O-$
DAST	=	diethylamine sulfur trifluoride
DCC	=	dicyclohexyl carbodiimide

	DCI carbodiimide	=	1-(3-dimethylaminopropyl)-3-ethyl
	DEAD	=	diethyl azodicarboxylate
	DIBAL	=	diisobutyl aluminum hydride
	DME	=	ethylene glycol dimethylether
	DMAP	=	4-(dimethylamino)pyridine
	DMF	=	N,N-dimethylformamide
	DMSO	=	dimethyl sulfoxide
	Et_3N	=	triethylamine
	LDA	=	lithium diisopropylamide
	m-CPBA	=	metachloroperbenzoic acid
	NBS	=	N-bromosuccinimide
	NSAID	=	non-steroidal anti-inflammatory drug
	PCC	=	pyridinium chlorochromate
	PDC	=	pyridinium dichromate
	Ph	=	phenyl
	1,2-Ph	=	1,2-benzenediyl
	Pyr	=	pyridinediyl
	Qn	=	7-chloroquinolin-2-yl
	R^S	=	$-CH_2SCH_2CH_2Ph$
	r.t.	=	room temperature
	rac.	=	racemic
	THF	=	tetrahydrofuran
	THP	=	tetrahydropyran-2-yl

Alkyl group abbreviations

Me	=	methyl
Et	=	ethyl
n-Pr	=	normal propyl
i-Pr	=	isopropyl
n-Bu	=	normal butyl
i-Bu	=	isobutyl
s-Bu	=	secondary butyl
t-Bu	=	tertiary butyl

c-Pr	=	cyclopropyl
c-Bu	=	cyclobutyl
c-Pen	=	cyclopentyl
c-Hex	=	cyclohexyl

Optical Isomers - Diastereomers - Geometric Isomers - Tautomers

Compounds of Formula I contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

Some of the compounds described herein contain olefinic double bonds, and unless specified otherwise, are meant to include both E and Z geometric isomers.

Some of the compounds described herein may exist with different points of attachment of hydrogen, referred to as tautomers. Such an example may be a ketone and its enol form known as keto-enol tautomers. The individual tautomers as well as mixture thereof are encompassed with compounds of Formula I.

Compounds of the Formula I may be separated into diastereoisomeric pairs of enantiomers by, for example, fractional crystallization from a suitable solvent, for example MeOH or EtOAc or a mixture thereof. The pair of enantiomers thus obtained may be separated into individual stereoisomers by conventional means, for example by the use of an optically active amine as a resolving agent or on a chiral HPLC column.

Alternatively, any enantiomer of a compound of the general Formula I or Ia may be obtained by stereospecific synthesis using optically pure starting materials or reagents of known configuration.

Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenedi-

amine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropyl-amine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

Dose Ranges

It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

The magnitude of prophylactic or therapeutic dose of a compound of Formula I will, of course, vary with the nature and the severity of the condition to be treated and with the particular compound of Formula I and its route of administration. It will also vary according to a variety of factors including the age, weight, general health, sex, diet, time of administration, rate of excretion, drug combination and response of the individual patient. In general, the daily dose from about 0.001 mg to about 100 mg per kg body weight of a mammal, preferably 0.01 mg to about 10 mg per kg. On the other hand, it may be necessary to use dosages outside these limits in some cases.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for oral administration to humans may contain from about 0.5 mg to about 5 g of active agent compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Dosage unit forms will generally contain from about 1 mg to about 2 g of an active ingredient, typically 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 800 mg, or 1000 mg.

Pharmaceutical Compositions

For the treatment of glucocorticoid receptor mediated diseases the compound of Formula I may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and

vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, etc., the compound of the invention is effective in the treatment of humans.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, solutions, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Patent 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredients is mixed with water-miscible solvents such as propylene glycol, PEGs and ethanol, or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or

condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. Cosolvents such as ethanol, propylene glycol or polyethylene glycols may also be used. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including

synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ambient temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, gels, solutions or suspensions, etc., containing a compound of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.) Topical formulations may generally be comprised of a pharmaceutical carrier, cosolvent, emulsifier, penetration enhancer, preservative system, and emollient.

Utilities

The ability of the compounds of Formula I to selectively modulate glucocorticoid receptors makes them useful for treating, preventing or reversing the progression of a variety of inflammatory and autoimmune diseases and conditions. Thus, the compounds of the present invention are useful to treat, prevent or ameliorate the following diseases or conditions: inflammation, tissue rejection, auto-immunity, various malianancies, such as leukemias and lymphomas, Cushing's syndrome, acute adrenal insufficiency, congenital adrenal hyperplasia, rheumatic fever, polyarteritis nodosa, granulomatous polyarteritis, inhibition of myeloid cell lines, immune proliferation/apoptosis, HPA axis suppression and regulation, hypercortisolemia, stroke and spinal cord injury, hypercalcemia, hypergylcemia, acute adrenal insufficiency, chronic primary adrenal insufficiency, secondary adrenal insufficiency, congenital adrenal hyperplasia, cerebral edema, thrombocytopenia, Little's syndrome, obesity and metabolic syndrome.

The compounds of the present invention are also useful for treating, preventing or reversing the progression of disease states involving systemic inflammation such as inflammatory bowel disease, systemic lupus erythematosus, polyartitis nodosa, Wegener's granulomatosis, giant cell arteritis, rheumatoid arthritis, juvenile rheumatoid arthritis, uveitis, hay fever, allergic rhinitis, urticaria, angioneurotic edema, chronic obstructive pulmonary disease, asthma, tendonitis, bursitis, Crohn's disease, ulcerative colitis, autoimmune chronic active hepatitis, organ transplantation, hepatitis, and cirrhosis. The compounds of the present invention are useful for treating, preventing or reversing the progression of a variety of topical diseases such as inflammatory scalp alopecia, panniculitis, psoriasis, discoid lupus erythematosus, inflamed cysts, atopic dermatitis, pyoderma gangrenosum, pemphigus vulgaris, buflous pernphigoid, systemic lupus erythematosus, dermatomyositis, herpes

gestationis, eosinophilic fasciitis, relapsing polychondritis, inflammatory vasculitis, sarcoidosis, Sweet's disease, type I reactive leprosy, capillary hemangiomas, contact dermatitis, atopic dermatitis, lichen planus, exfoliative dermatitus, erythema nodosum, acne, hirsutism, toxic epidermal necrolysis, erythema multiform, cutaneous T-cell lymphoma.

The compounds of the present invention are also useful in treating, preventing or reversing the progression of disease states associated with Human Immunodeficiency Virus (HIV), cell apoptosis, and cancer including, but not limited to, Kaposi's sarcoma, immune system activation and modulation, desensitization of inflammatory responses, IIL- I expression, natural killer cell development, lymphocytic leukemia, and treatment of retinitis pigmentosa. Cogitive and behavioral processes are also susceptible to glucocorticoid therapy where antagonists would potentially be useful in the treatment of processes such as cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, stroke, sleep disorders, and anxiety.

Preferably, the compounds of the invention are useful for treating the diseases or conditions set for the below.

1. Allergic States

Control of severe or incapacitating allergic conditions not responsive to adequate trials of conventional treatment; seasonal or perennial allergic rhinitis; bronchial asthma; contact dermatitis; atopic dermatitis; serum sickness; and drug hypersensitivity reactions.

2. Rheumatic Disorders

As adjunctive therapy for short-term administration during an acute episode or exacerbation of: psoriatic arthritis; rheumatoid arthritis including juvenile rheumatoid arthritis (selected cases may require low-dosemaintenance therapy); ankylosing spondylitis; acute and subacute bursitis; acute nonspecific tenosynovitis; acute gouty arthritis; post-traumatic osteoarthritis; synovitis of osteoarthritis; and epicondylitis

3. Dermatologic Diseases

Ppemphigus; bullous dermatitis herpetiformis; severe erythema multiforme (Stevens-Johnson syndrome); exfoliative dermatitis; mycosis fungoïdes; severe psoriasis; and severe seborrheic dermatitis.

4. Ophthalmic Diseases

Severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: allergic conjunctivitis; keratitis; allergic corneal marginal ulcers; herpes zoster ophthalmicus; iritis and iridocyclitis; chorioretinitis; anterior segment inflammation; diffuse posterior uveitis and choroiditis; optic neuritis; and sympathetic ophthalmia.

5. Endocrine Disorders

Primary or secondary adrenocortical insufficiency; congenital adrenal hyperplasia; nonsuppurative thyroiditis; and hypercalcemia associated with cancer.

6. Respiratory Diseases

Symptomatic sarcoidosis; Löffler's syndrome not manageable by other means; berylliosis; fulminating or disseminated pulmonary tuberculosis when concurrently accompanied by appropriate antituberculous chemotherapy; and aspiration pneumonitis.

7. Hematologic Disorders

Idiopathic thrombocytopenic purpura in adults; secondary thrombocytopenia in adults; acquired (autoimmune) hemolytic anemia; erythroblastopenia (RBC anemia); and congenital (erythroid) hypoplastic anemia.

8. Neoplastic Diseases

For palliative management of: leukemias and lymphomas in adults; and acute leukemia of childhood.

9. Edematous States

To induce a diuresis or remission of proteinuria in the nephrotic syndrome without uremia, of the idiopathic type or that due to lupus erythematosus. Compounds of Formula I may be used to treat patients with cerebral edema from various causes. It may be used also in the preoperative preparation of patients with increased intracranial pressure secondary to brain tumors, and also for palliation of patients with inoperable or recurrent brain neoplasms, and in the management of cerebral edema associated with neurosurgery. Some patients with cerebral edema due to head injury or pseudotumor cerebri also may benefit fromtherapy with compounds of Formula I.

10. Gastrointestinal Diseases

During a critical period of the disease in: ulcerative colitis and regional enteritis.

11. Miscellaneous

Tuberculous meningitis with subarachnoid block or impending block when concurrently accompanied by appropriate antituberculous chemotherapy; Trichinosis with neurologic or myocardial involvement; During an exacerbation or as maintenance therapy in selected cases of: Systemic lupus erythematosus and acute rheumatic carditis; in combination with ondansetron for the management of nausea and vomiting associated with cisplatin and non-cisplatin emetogenic chemotherapy.

Combination Therapy

The invention also encompasses a method for treating a glucocorticoid receptor mediated disease comprising concomitantly administering to a patient in need of such treatment a compound of Formula I and one or additional more agents. For treating or preventing asthma or chronic obstructive

pulmonary disease, the compounds of Formula I may be combined with one or more agents selected from the group consisting of: 9-agonists (e.g., salmeterol), theophylline, anticholinergics (e.g., atropine and ipratropium bromide), cromolyn, nedocromil and leukotriene modifiers (e.g., montelukast). For treating or preventing inflammation, the compounds of Formula I may be combined with one or the following: a salicylate, including acetylsalicylic acid, a non-steroidal antiinflammatory drug, including indomethacin, sulindac, mefenamic, meclofenamic, tolfenamic, tolmetin, ketorolac, dicofenac, ibuprofen, naproxen, fenoprofen, ketoprofen, flurbiprofin and oxaprozin, a TNF inhibitor, including etanercept and infliximab, an IL-1 receptor antagonist, a cytotoxic or immunosuppressive drug, including methotrexate, leflunomide, azathioprine and cyclosporine, a gold compound, hydroxychloroquine or sulfasalazine, penicillamine, darbufelone, and a p38 kinase inhibitor. The compound of Formula I may also be used in combination with bisphonates such as alendronate to treat a glucocorticoid mediated disease and simultaneously inhibit osteoclast-mediated bone resorption.

Methods of Synthesis and Examples

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

(i) all operations were carried out at room or ambient temperature, that is, at a temperature in the range 18-25°C,

(ii) evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 pascals: 4.5-30 mm. Hg) with a bath temperature of up to 60°C.,

(iii) the course of reactions was followed by thin layer chromatography (TLC) and reaction times are given for illustration only;

(iv) melting points are uncorrected and `d' indicates decomposition; the melting points given are those obtained for the materials prepared as described; polymorphism may result in isolation of materials with different melting points in some preparations;

(v) the structure and purity of all final products were assured by at least one of the following techniques: TLC, mass spectrometry, nuclear magnetic resonance (NMR) spectrometry or microanalytical data;


(vi) yields are given for illustration only;


(vii) when given, NMR data is in the form of delta (δ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, determined at 500 MHz or 600 MHz using the indicated solvent; conventional abbreviations used for signal shape are: s. singlet; d. doublet; t. triplet; m. multiplet; br. broad; etc.: in addition "Ar" signifies an aromatic signal;

(viii) chemical symbols have their usual meanings; the following abbreviations have also been used v (volume), w (weight), b.p. (boiling point), m.p. (melting point), L (litre(s)), mL (millilitres), g (gram(s)), mg (milligrams(s)), mol (moles), mmol (millimoles), eq (equivalent(s)).

EXAMPLE 1

SYNTHESIS OF (11β,16β,17α)-9-FLUORO-11-HYDROXY-16-METHYL-3,17A-DIOXO-17-(HYDROXYMETHYL)-D-HOMOANDROSTA-1,4-DIEN-17-YL-(1-ADAMANTYL)CARBAMATE





(11β,16β)-9-fluoro-17,21-dihydroxy-16-methyl-3,20-dioxo-11-trifluoroacetoxy-pregna-1,4-diene (1-3)

Betamethasone (1-1, 96 g, 245 mmol) and DMAP (6.0 g, 49 mmol) were dissolved in 2.8 L of dry THF with stirring. The solution was placed in an ice bath and cooled to 10-15°C. To the cooled solution was added trifluoroacetic anhydride (129 g, 614 mmol) via an addition funnel over a 20 minute period. The ice bath was allowed to expire and the solution was allowed to stir at room temperature for 15 hours. To the solution was added 675 mL of water, and the resulting mixture was allowed to stir for 15 hours. The mixture was reduced to one-quarter the original volume in vacuo and the residue was diluted with 2 L of water. The precipitate was isolated by filtration and dried for 15 hrs at reduced pressure to produce 122.5 g of 1-3 as a pale white solid.

MS (ESI): $m/z = 489.2$ (MH^+).

(11β,16β)-9-fluoro-17-hydroxy-16-methyl-3,20-dioxo-11-trifluoroacetoxy-21-acetoxy-pregna-1,4-diene (1-4)

To a solution of 1-3 (122.5 g) in THF (1 L) at RT was added DMAP (6.0 g, 49 mmol) and acetic anhydride (50.2 g, 491 mmol) and the mixture allowed to stir for 15 hours. The solvent was evaporated, and the residue diluted with 1 L of water. The precipitate was isolated by filtration and dried for 15 hrs in a warm (35°C) vacuum oven to produce 123.4 g (94.7%) 1-4 as an off-white solid.
MS (ESI): m/z = 531.2 (MH^+).

(11β,16β,17α)-9-fluoro-17-hydroxy-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-diene (1-5)

A suspension of 1-4 (42.0 g, 79.168 mmol) in 630mL anhydrous acetonitrile was cooled to -5°C and $SnCl_4$ (41.24 g, 158.335 mmol) was added dropwise. The resulting homogenous solution was warmed to ambient temperature stirred for 72h during which time a white solid precipitated. The solid was filtered and dried overnight at 35°C to give 34.10 g of 1-5. The filtrate was diluted with ethyl acetate (2 L) and diluted with a mixture of sodium bicarbonate (6.651 g, 79.168 mmol) in 300 mL water. The combined organic layers were washed with water, saturated brine solution, dried over sodium sulfate, filtered and concentrated in vacuo to give an additional 3 g of 1-5 as beige foamy solid.
MS (ESI): m/z = 531.3 (MH+).

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate (1-6)

A solution of 1-5 (0.815 g, 1.536 mmol) in 2.2 mL anhydrous N,N-dimethylformamide was purged with a stream of nitrogen for 10 minutes, then 1-adamantyl isocyanate (0.545 g, 3.072 mmol) and CuCl (0.152 g, 1.536 mmol) were added. The resulting green solution was stirred at ambient temperature for 1h. The reaction mixture was poured into 15mL water and extracted with ethyl acetate (2x 50mL). The combined organic layers were washed with 10 mL water, 10 mL saturated brine solution, dried over magnesium sulfate, filtered, and concentrated in vacuo to give 1.18 g of an orange viscous oil. The oil was dissolved in with toluene (15mL) and concentrated in vacuo to give a pale green solid (1.00g). The residue was subjected to flash chromatography through a column of 40g of silica gel, eluting with a gradient of 0% B/100% A to 20% B/80% A over 55 min (A = 50% dichloromethane/50% hexane, B = ethyl acetate) to provide 0.646g (59%) of 1-6 as a beige solid.
MS (ESI): m/z = 708.3 (MH+).

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate (1-7)

Solid 1-6 (610 g, 0.862 mmol) was dissolved in 30 mL of a premixed solution of methanol/chloroform/ 6N HCl (10:2:1) and the resulting mixture stirred at ambient temperature for 15 hours. An additional 1 mL of 6N HCl was added and the mixture stirred for a further 20 hours. The mixture was concentrated in vacuo to give 0.544 g of 1-7 as a beige solid.
MS (ESI): *m/z* = 666.3 (MH+).

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate (1-8)


To a solution of 1-7 (0.492 g, 0.739 mmol) in 36mL methanol was added triethylamine (0.206 mL, 1.478 mmol) and the resulting colorless solution was stirred at ambient temperature for 30 min then concentrated in vacuo to give a beige foamy solid. The solid was dissolved 7 mL hot dichloromethane, loaded onto a 40 g silica gel cartridge, and eluted with 0%B/100%A for 10 min, then with a gradient of 0%B/100%A to 50% B/50%A over 50 min to yield 0.330 g 1-8 as a white solid (78%).
MS (ESI):*m/z* = 570.3 (MH+).

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate (1-9)

To a solution of 1-6 (0.100 g, 0.141 mmol) in 7 mL methanol was added triethylamine (0.039 mL, 0.283 mmol) and the resulting colorless solution was stirred at ambient temperature for 30 min then concentrated in vacuo to give 0.076 g of 1-9 as a beige solid.
MS (ESI):*m/z* = 612.3 (MH+).

EXAMPLES 2-9

The following compounds were prepared using methods analogous to those described in the preceding example:



2	R^a = H R^b = H R^c = 2-pentyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-pentyl)carbamate
Mass Spectrum (ESI): *m/z* = 506.7 (MH^+)		

3	R^a = H R^b = H R^c = 2-methyl-pentyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-methyl-pentyl)carbamate
Mass Spectrum (ESI): *m/z* = 520.7 (MH^+)		

4	R^a = H R^b = H R^c = 3-methyl-2-propyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(3-methyl-2-propyl)carbamate
Mass Spectrum (ESI): *m/z* = 506.7 (MH^+)		

5	R^a = $COCF_3$ R^b = $COCH_3$ R^c = 1-adamantyl	(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate
Mass Spectrum (ESI): *m/z* = 708.3 (MH^+)		

6	R^a = H R^b = $COCH_3$ R^c = 1-adamantyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate
Mass Spectrum (ESI): *m/z* = 612.3 (MH^+)		

7	R^a = $COCF_3$ R^b = H R^c = 1-adamantyl	(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate

Mass Spectrum (ESI): *m/z* = 666.3 (MH^+)		

8	R^a = H R^b = H R^c = cyclopentyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclopentyl)carbamate
Mass Spectrum (ESI): *m/z* = 504.3 (MH^+)		

9	R^a = H R^b = H R^c = cyclohexyl	(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclohexyl)carbamate
Mass Spectrum (ESI): *m/z* = 518.3 (MH^+)		

Biological Evaluation

The compounds exemplified in the present application exhibited activity in one or more of the following assays.

Ligand binding assay

Materials:

Binding Buffer: TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM beta-mecaptoethanol, 10 mM Sodium Molybdate, pH 7.2)

50% HAP Slurry: Calbiochem Hydroxylapatite, Fast Flow, in 10 mM Tris, pH 8.0 and 1 mM EDTA.

Wash Buffer: 40 mM Tris, pH7.5, 100 mM KCl, 1 mM EDTA and 1 mM EGTA.

95% EtOH

Dexmethasone-methyl-^{3}H, (DEX*); (Amersham cat# TRK645)

Dexamethasone(DEX) (Sigma, cat# D1756):

Hydroxylapatite Fast Flow; Calbiochem Cat#391947

Molybdate = Molybdic Acid (Sigma, M1651)

MDA-MB-453 cell culture media:

RPMI 1640 (Gibco 11835-055) w/23.8

mM $NaHCO_3$, 2 mM L-glutamine

in 500 mL of complete media Final conc.

10 mL (1M Hepes)	20 mM
5 mL (200 mM L-glu)	4 mM
0.5 mL (10 mg/mL human insulin) in 0.01 N HCl Calbiochem#407694-S)	10 µg/mL
50 mL FBS (Sigma F2442)	10%
1 mL (10 mg/mL Gentamicin Gibco#15710-072)	20 µg /mL

Cell Passaging

Cells (Hall R. E., et al., European Journal of Cancer, 30A: 484-490 (1994)) MDA-MB-453 (ATCC) cultured in RPMI 1640 (Gibco 11835-055) containing 20 mM Hepes, 4 mM L-glu, 10 ug/ml of human insulin (Sigma, I-0259), 10% FBS and 20 ug/ml of Gentamicin (Gibco#15710-072) are rinsed twice in PBS. Phenol red-free Trypsin-EDTA is diluted in the same PBS 1:10. The cell layers are rinsed with 1X Trypsin, extra Trypsin is poured out, and the cell layers are incubated at 37°C for ~ 2 min. The flask is tapped and checked for signs of cell detachment. Once the cells begin to slide off the flask, the complete media is added. The cells are counted at this point, then diluted to the appropriate concentration and split into flasks or dishes for further culturing (Usually 1:3 to 1:6 dilution).

Preparation of MDA-MB-453 Cell Lysate

When the cells are 70 to 85% confluent, they are detached as described above, and collected by centrifuging at 1000 g for 10 minutes at 4°C. The cell pellet is washed twice with TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM beta-mercaptoethanol, 10 mM Sodium Molybdate, pH 7.2). After the final wash, the cells are resuspended in TEGM at a concentration of 10^7 cells/mL. The cell suspension is snap frozen in liquid nitrogen or ethanol/dry ice bath and transferred to –80°C freezer on dry ice. Before setting up the binding assay, the frozen samples are left on ice-water to just thaw (~1 hr). Then the samples are centrifuged at 12,500 g to 20,000 g for 30 min at 4°C. The supernatant is used to set-up assay right away. If using 50 µL of supernatant, the test compound can be prepared in 50 µL of the TEGM buffer.

Procedure for Multiple Compound Screening

1x TEGM buffer is prepared, and the isotope-containing assay mixture is prepared in the following order: EtOH (2% final concentration in reaction), ^{3}H-DEX (Amersham Biosciences) and 1x TEGM. [e.g. For 100 samples, 200 µL (100 x 2) of EtOH + 4.25 µL of 1:10 ^{3}H--Dex stock + 2300 µL (100 x 23) 1x TEGM]. The compound is serially diluted, e.g., if starting final conc. is 1 µM, and the

compound is in 25 μL of solution, for duplicate samples, 75 μL of 4x1 μM solution is made and 3 μL of 100 μM is added to 72 μL of buffer, and 1:5 serial dilution.

25μL of ^{3}H-DEX (6 nM) trace and 25 μL compound solution are first mixed together, followed by addition of 50 μL receptor solution. The reaction is gently mixed, spun briefly at about 200 rpm and incubated at 4°C overnight. 100 μL of 50% HAP slurry is prepared and added to the incubated reaction which is then vortexed and incubated on ice for 5 to 10 minutes. The reaction mixture is vortexed twice more to resuspend HAP while incubating reaction. The samples in 96-well format are then washed in wash buffer using The FilterMate™ Universal Harvester plate washer (Packard). The washing process transfers HAP pellet containing ligand-bound expressed receptor to Unifilter-96 GF/B filter plate (Packard). The HAP pellet on the filter plate is incubated with 50 μL of MICROSCINT (Packard) scintillint for 30 minutes before being counted on the TopCount microscintillation counter (Packard). IC_{50}s are calculated using DEX as a reference.

Examples 1 to 9 were tested in the ligand binding assay and demonstrated IC50s less than 1000 nM.

Trans-Activation Modulation of Glucocorticoid Receptor (GRAMMER)

This assay assesses the ability of test compounds to control transcription from the MMTV-LUC reporter gene in lung adenocarcinoma A549 cells or MDA-MB-453 cells, a human breast cancer cell line that naturally expresses the human GR. The assay measures induction of a modified MMTV LTR/promoter linked to the LUC reporter gene.

The routine transient assay consists of plating 7,000-25,000 cells/well of a white, clear-bottom 96-well plate. Alternatively, 384-well plates can be used at a cell concentration of 10,000 /well. The media that the cells are plated in is "exponential growth medium" which consists of phenol red-free RPMI1640 containing 10%FBS, 4mM L-glutamine, 20mM HEPES, 10ug/mL human insulin, and 20ug/mL gentamicin. Incubator conditions are 37°C and 5% CO_2. The transfection is done in batch mode. The cells are trypsinized and counted to the right cell number in the proper amount of fresh media. It is then gently mixed with the FuGene6/DNA mix and plated onto the 96 or 384-well plate, all the wells receive 100 uL or 40uL, respectively, of medium + lipid/DNA complex then incubated 37°C overnight. The transfection cocktail consists of serum-free OptiMEM, FuGene6 reagent and DNA. The manufacturer's (Roche Biochemical) protocol for cocktail setup is as follows: The lipid to DNA ratio is approximately 2.5:1 and the incubation time is 20 min at room temperature. Sixteen to 24 hours after transfection, the cells are treated with dexamethasone to a final concentration of 10nM as well as the compound of interest, such that final DMSO (vehicle) concentration is equal to or less than 1%. Each plate also contains samples that are treated with 10nM dexamethasone alone, which is used as the 100%

activity control. The cells are exposed to the compounds for 24 hours. After 24 hours, the cells are lysed by a Promega cell culture lysis buffer for approximately 30 min and then the luciferase activity in the extracts is assayed in the 96-well format luminometer. In 384-well format, Steady-Glo (Promega) or Steady-Lite (PerkinElmer) can be used by adding an equal volume of reagent to the media present in each well. Activity induced by 10nM dexamethasone alone is set at 100% activity. Antagonist activity is calculated by determining the decrease in dexamethasone-induced activity in response to compound treatment relative to samples that were treated with dexamethasone alone. Results are expressed as % inhibition of 10nM dexamethasone activity or as fold of 10nM dexamethasone activity. This transactivation assay can be performed in an agonist and antagonist mode to identify these different activities.

Activity of test compounds is calculated as the E_{max} relative to the activity obtained with 300 nM dexamethasone. Activity of test compounds is calculated as the E_{max} relative to the activity obtained with 300 nM DEX. The exemplified tissue selective glucocorticoid receptor modulators of the present invention display partial agonist activity in this assay of greater than 5%.

The action of compounds is also tested in an antagonist mode (Anti-GRAMMER) in which the cells are treated with medium containing an agonist such as 10 nM DEX and the ability to agents to inhibit the activation by an agonist is measured.

Transrepression assay

This assay assesses the ability of test compounds to control transcription from the TNFα-β-lactamase reporter gene in U937 cells, a human myelomonocytic leukemia cell line that naturally expresses the human GR. The assay measures compound dependent-repression of the TNFa promoter linked to a reporter gene.

The human U937cells that had been stablely transfected with the TNF-α promoter driving β-lactamase are used for this assay. U937 cells contain an endogenous glucocorticoid receptor (GR). Cells are maintained in RPMI 1640 Growth medium *(Gibco Cat#11875-093)* containing 25mM HEPES, 10% FBS, 2mM L-Glutamine, 1mM Sodium pyruvate, 25μg/ml Gentamicin *(Gibco Cat#15710-064)*, 1:1000 2-Mercaptoethanol *(Gibco Cat#21985-023) and* 0.8 mg/ml G418 *(Gibco Cat#10131-027)*. The density of the cells in the flask needs to be about $1X10^6 - 3X10^6$/ml at the time of harvest. Usually, the cells are split to $1.2\sim1.4x10^5$ /ml (1:10) 3 days prior to the assay. 50,000 cells/well are plated in 96 well black-walled plates the day of assay. Test compounds are added 10 μL/well, and cells are incubated at 37°C for 30~45 min. For assaying compounds, first dilute 1:10 in DMSO to make 1 mM, then further dilute 1:100 in medium to make 10X stock prior to adding to the cells. Add 50ng/ml PMA (Sigma, cat#

P8139) 10 μL/well to a final concentration 5ng/ml, and 1 μg/ml LPS (Sigma, cat# L4130) 10 μL/well to a final concentration 100ng/ml. Incubate cells at 37^{0}C overnight for ~18hr. PMA is stored frozen as 100 μg/ml stock in DMSO. Dilute 1:10 in DMSO for a working stock of 10 μg/ml and store at –20C. For assaying, dilute the 10 μg/ml working stock 1:200 in medium to make a 10X solution (50 ng/ml). Store frozen LPS at 1 mg/ml in PBS, dilute 1:1000 in medium to make 10X (1μg/ml) for the assay. Add 6X loading buffer (CCF2-AM) 20 μL/well, and incubate at room temperature for 70~90 min. Read plates on CytoFluor II Plate Reader according to manufacture suggested protocols. The activity repressed by 100nM dexamethasone alone is set as 100% activity.

Examples 1 to 9 were tested in the transrepression assay and demonstrated maximal activity greater than 5%.

Microarray analysis

All cell culture reagents were purchased from Invitrogen Life Tech, Carlsbad CA. A549 cells were grown in phenol red-free DMEM/F12 medium supplemented with 10% FBS. Cells were grown at 37^{o}C with 5% CO_2. Using the RNeasy Kit (Qiagen Corp, Valencia CA.), total RNA was extracted and purified from A549 cells treated with different GC compounds for 24 hours, at a fully active dose. These cells express large amount of the GR and are very responsive to GC treatment. All samples were compared against cells treated with vehicle. Expression levels of 23000 genes were measured using oligonucleotide microarrays purchased from Agilent Technologies, Inc. Each comparison was done on a pair of microarrays with reversed fluorophores. Raw image intensity data were processed according to the method described in Patent 6,351,712. The method was used to remove dye bias and to derive a Rosetta probability (p) and fold change value for each gene and each sample pair. Furthermore, for each gene an ANOVA model was constructed across all treatments to derive error estimates. P values for evaluating expression differences were computed using a Bayesian adjusted t-test that was developed by Lönnstedt and Speed (2002) and extended by Smyth (2003). A gene was declared differentially expressed in any particular comparison if it satisfied two critera:

1. The Rosetta p value had to be less than 0.1 and the Rosetta fold change value had to be greater than 1.4 in at least one of the treatments.

2. The ANOVA p value had to be less than 0.01 and the fold change greater than 2 in the comparison under consideration.


In Vivo inflammation Assay

Intact adult (6 month old) female Sprague-Dawley rats are used in the oxazolone (OX) contactdermatitis model. Rats were sensitized on the ventral abdomen with OX on Day 0. On Days 7 and 9, a randomly-selected ear was challenged (same ear each time) with OX; the other was treated with vehicle. Daily treatment begun on Day 7 and continued for 7d with test compounds at different doses and 1.3 mpk 6-methlyprednisolone or 0.1mpk DEX as positive controls. The thickness of both ears are measured on Days 11 and 14. Necropsy occurred on Day 14. The rat is first weighed, then anesthetized in a CO_2 chamber until near death. Approximately 5ml whole blood is obtained by cardiac puncture. The rat is then examined for certain signs of death and completeness. Tissues are dissected in a highly stylized fashion. The the following endpoints were evaluated: a) inhibiting ear inflammation induced by oxazalone, b) raising serum insulin, c) reducing serum ACTH, d) reducing spleen weight, e) reducing skin thickness, f) reducing body weight, g) increasing expression of bone-related genes with potential relationship to negative glucocorticoid effects on bone; e) changes in molecular markers that correlate with skin inflammation, skin thinning, muscle atrophy and glucose metabolism in liver. All blood samples were collected between 1330-1530 hours, ~4-5 hrs after the last compound treatment.

Primary data for this assay are left and right ear thickness. Inter-ear thickness difference (etd) is used for the estimating the level of inflammation and effectiveness of the compounds is determined by their ability to reduce the increase the thickness of the inflamed ear. Back of the rat skin thickness, spleen weight, serum insulin as well as the effects of gcs on the expression of molecular markers in skin inflammation, skin atrophy, muscle atrophy and glucose metabolism in liver are measured. Data are analyzed by anova plus fisher plsd post-hoc test to identify intergroup differences.

WHAT IS CLAIMED IS:

1. A compound of Formula I



or a pharmaceutically acceptable salt thereof, wherein

R^1 and R^2 are independently selected from the group consisting of: R^a, -C(O)-R^a, and –C(O)-N(R^a)R^a;

R^3 and R^4 are independently selected from the group consisting of: hydrogen, C_{1-10}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, C_{3-10}cycloalkyl, aryl, hetreoaryl, heterocyclyl, and C_{1-10}fluoroalkyl; and

each Ra is independently selected from the group consisting of: hydrogen, C_{1-10}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, C_{3-6}cycloalkyl, aryl, hetreoaryl, heterocyclyl, and C_{1-10}fluoroalkyl.

2. The compound according to Claim 1 wherein R^2 is selected from hydrogen and CH_3-C(O)-.

3. The compound according to Claim 1 wherein R^1 is selected from hydrogen and CF_3-(CO)-.

4. The compound according to Claim 1 wherein R^3 is hydrogen.

5. The compound according to Claim 4 wherein R^4 is selected from C_{1-10}alkyl and C_{3-10}cycloalkyl.

6. The compound according to Claim 1 wherein:

R^1 is selected from hydrogen or CF_3-(CO)-;

R^2 is selected from hydrogen and CH_3-C(O)-;

R^3 is hydrogen; and

R^4 is selected from C_{1-10}alkyl and C_{3-10}cycloalkyl.

7. The compound according to Claim 6 wherein R^4 is C_{1-6}alkyl.

8. The compound according to Claim 6 wherein R^4 is selected from: cyclopentyl, cyclohexyl and adamantanyl.

9. The compound according to Claim 6 wherein R^1 and R^2 are hydrogen.

10. The compound according to Claim 1 selected from the following group: (11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-pentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(2-methyl-pentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(3-methyl-2-propyl)carbamate,

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-[(acetyloxy)methyl]-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-trifluoroacetoxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(1-adamantyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclopentyl)carbamate,

(11β,16β,17α)-9-fluoro-11-hydroxy-16-methyl-3,17a-dioxo-17-(hydroxymethyl)-D-homoandrosta-1,4-dien-17-yl-(cyclohexyl)carbamate

or a pharmaceutically acceptable salt of any of the above.

11. A pharmaceutical composition comprising a compound according to Claim 1 in combination with a pharmaceutically acceptable carrier.

12. A method for treating a glucocorticoid receptor mediated disease or condition in a mammalian patient in need of such treatment comprising administering the patient a compound according to Claim 1 in an amount that is effective for treating the glucocorticoid receptor mediated disease or condition.

13. The method according to Claim 12 wherein the glucocorticoid receptor mediated disease or condition is selected from the group consisting of: tissue rejection, leukemias, lymphomas, Cushing's syndrome, acute adrenal insufficiency, congenital adrenal hyperplasia, rheumatic fever, polyarteritis nodosa, granulomatous polyarteritis, inhibition of myeloid cell lines, immune proliferation/apoptosis, HPA axis suppression and regulation, hypercortisolemia, stroke and spinal cord injury, hypercalcemia, hypergylcemia, acute adrenal insufficiency, chronic primary adrenal insufficiency, secondary adrenal insufficiency, congenital adrenal hyperplasia, cerebral edema, thrombocytopenia, Little's syndrome, obesity, metabolic syndrome, inflammatory bowel disease, systemic lupus erythematosus, polyartitis nodosa, Wegener's granulomatosis, giant cell arteritis, rheumatoid arthritis, juvenile rheumatoid arthritis, uveitis, hay fever, allergic rhinitis, urticaria, angioneurotic edema, chronic obstructive pulmonary disease, asthma, tendonitis, bursitis, Crohn's disease, ulcerative colitis, autoimmune chronic active hepatitis, organ transplantation, hepatitis, cirrhosis, inflammatory scalp alopecia, panniculitis, psoriasis, discoid lupus erythematosus, inflamed cysts, atopic dermatitis, pyoderma gangrenosum, pemphigus vulgaris, buflous pernphigoid, systemic lupus erythematosus, dermatomyositis, herpes gestationis, eosinophilic fasciitis, relapsing polychondritis, inflammatory

vasculitis, sarcoidosis, Sweet's disease, type I reactive leprosy, capillary hemangiomas, contact dermatitis, atopic dermatitis, lichen planus, exfoliative dermatitus, erythema nodosum, acne, hirsutism, toxic epidermal necrolysis, erythema multiform, cutaneous T-cell lymphoma, Human Immunodeficiency Virus (HIV), cell apoptosis, cancer, Kaposi's sarcoma, retinitis pigmentosa, cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, sleep disorders, and anxiety.

14. The method according to Claim 12 wherein the glucocorticoid receptor mediated disease or condition is selected from the group consisting of: tissue rejection, Cushing's syndrome, inflammatory bowel disease, systemic lupus erythematosus, rheumatoid arthritis, juvenile rheumatoid arthritis, hay fever, allergic rhinitis, asthma, organ transplantation, inflammatory scalp alopecia, psoriasis, discoid lupus erythematosus, and depression.

15. A method of selectively modulating the activation, repression, agonism and antagonism effects of the glucocorticoid receptor in a mammal comprising administering to the mammal a compound according to Claim 1 in an amount that is effective to modulate the glucocorticoid receptor.

16. A method of partially or fully antagonizing, repressing agonizing or modulating the glucocorticoid receptor in a mammal comprising administering to the mammal an effective amount of compound according to Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US06/18613

A. CLASSIFICATION OF SUBJECT MATTER
IPC(8): **C07C 261/00**(2006.01),**261/00**(2006.01)

USPC: 560/24,27,115,162,163,164
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 560/24,27,115,162,163,164

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EAST, WEST AND CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 47011068 B (TANABE SEIYAKU CO., LTD.) 04 April 1972, see entire document.	1-16
A	US 3,598,845 A (JOHNSON) 10 August 1971, see entire document.	1-16

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
05 August 2006 (05.08.2006)

Date of mailing of the international search report
29 AUG 2006

Name and mailing address of the ISA/US
Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450
Facsimile No. (571) 273-3201

Authorized officer
Jafar Parsa

Telephone No. (571)272-2717

Form PCT/ISA/210 (second sheet) (April 2005)