



(86) Date de dépôt PCT/PCT Filing Date: 2001/02/01

(87) Date publication PCT/PCT Publication Date: 2001/08/09

(85) Entrée phase nationale/National Entry: 2002/08/05

(86) N° demande PCT/PCT Application No.: US 2001/003293

(87) N° publication PCT/PCT Publication No.: 2001/057070

(30) Priorité/Priority: 2000/02/03 (60/179,967) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 7/08, A61K 38/00, C07K 7/52

(71) Demandeur/Applicant:
MERCK & CO., INC., US

(72) Inventeur/Inventor:
BEDNAREK, MARIA, US

(74) Agent: OGILVY RENAULT

(54) Titre : ANALOGUES DE L'HORMONE CONCENTRATRICE DE LA MELANINE

(54) Title: MELANIN-CONCENTRATING HORMONE ANALOGS

(57) **Abrégé/Abstract:**

The present invention features truncated MCH analogs active at the MCH receptor. The truncated MCH analogs are optionally modified peptide derivatives of mammalian MCH. The analogs can bind to the MCH receptor and, preferably, bring about signal transduction. MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57070 A1

(51) International Patent Classification⁷: **C07K 7/08**,
7/52, A61K 38/00

[US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number: PCT/US01/03293

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 1 February 2001 (01.02.2001)

(25) Filing Language: English

(81) Designated States (*national*): CA, JP, US.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(30) Priority Data:
60/179,967 3 February 2000 (03.02.2000) US

(71) Applicant (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

Published:
— *with international search report*

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **BEDNAREK, Maria**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/57070 A1

(54) Title: MELANIN-CONCENTRATING HORMONE ANALOGS

(57) Abstract: The present invention features truncated MCH analogs active at the MCH receptor. The truncated MCH analogs are optionally modified peptide derivatives of mammalian MCH. The analogs can bind to the MCH receptor and, preferably, bring about signal transduction. MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically.

TITLE OF THE INVENTION

MELANIN-CONCENTRATING HORMONE ANALOGS

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 The present application claims priority to provisional application U.S. Serial No. 60/179,967, filed February 3, 2000, which is hereby incorporated by reference herein.

BACKGROUND OF THE INVENTION

- 10 Neuropeptides present in the hypothalamus play a major role in mediating the control of body weight. (Flier, *et al.*, 1998. *Cell*, 92, 437-440.) Melanin-concentrating hormone (MCH) produced in mammals is a cyclic 19-amino acid neuropeptide synthesized as part of a larger pre-prohormone precursor in the hypothalamus which also encodes neuropeptides NEI and NGE. (Nahon, *et al.*, 1990. *Mol. Endocrinol.* 4, 632-637; Vaughan, *et al.*, U.S. Patent No. 5,049,655; and Vaughan, *et al.*, 1989. *Endocrinology* 125, 1660-1665.) MCH was first identified in salmon pituitary, and in fish MCH affects melanin aggregation thus affecting skin pigmentation. In trout and eels MCH has also been shown to be involved in stress induced or CRF-stimulated ACTH release. (Kawauchi, *et al.*, 1983. *Nature* 305, 20 321-323.)

- In humans two genes encoding MCH have been identified that are expressed in the brain. (Breton, *et al.*, 1993. *Mol. Brain Res.* 18, 297-310.) In mammals MCH has been localized primarily to neuronal cell bodies of the hypothalamus which are implicated in the control of food intake, including perikarya of the lateral hypothalamus and zona inertia. (Knigge, *et al.*, 1996. *Peptides* 17, 25 1063-1073.)

- Pharmacological and genetic evidence suggest that the primary mode of MCH action is to promote feeding (orexigenic). MCH mRNA is up regulated in fasted mice and rats, in the *ob/ob* mouse and in mice with targeted disruption in the gene for neuropeptide Y (NPY). (Qu, *et al.*, 1996. *Nature* 380, 243-247 and Erickson, *et al.*, 1996. *Nature* 381, 415-418.) Injection of MCH centrally (ICV) stimulates food intake and MCH antagonizes the hypophagic effects seen with α melanocyte stimulating hormone (α MSH). (Qu, *et al.*, 1996. *Nature* 380, 243-247.) MCH deficient mice are lean, hypophagic and have increased metabolic rate. 30 (Shimada, *et al.*, 1998. *Nature* 396, 670-673.) The administration of MCH has been


indicated to useful for promoting eating, appetite or the gain or maintenance of weight. (Maratos-Flier, U.S. Patent No. 5,849,708.)

MCH action is not limited to modulation of food intake as effects on the hypothalamic-pituitary-axis have been reported. (Nahon, 1994. *Critical Rev. in Neurobiol.* 8, 221-262.) MCH may be involved in the body response to stress as MCH can modulate the stress-induced release of CRF from the hypothalamus and ACTH from the pituitary. In addition, MCH neuronal systems may be involved in reproductive or maternal function.

10 SUMMARY OF THE INVENTION

The present invention features truncated MCH analogs active at the MCH receptor. The truncated MCH analogs are optionally modified peptide derivatives of mammalian MCH. The analogs can bind to the MCH receptor and, preferably, bring about signal transduction. MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically.

Thus, a first aspect of the present invention describes a truncated MCH analog. The truncated MCH analog is an optionally modified peptide having the structure:



wherein X¹ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X² is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X³ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

X⁴ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof;

5 X⁵ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

10 X⁶ is an optionally present amino acid that, if present is either arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof,

X⁷ is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

15 X⁸ is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof;

X⁹ is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof;

20 X¹⁰ is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, asparagine, serine, sarcosine, isobutyric, or gamma-aminobutyric acid, or a derivative thereof;

X¹¹ is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof;

X¹² is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof;

25 X¹³ is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO₂, NH₂, F, I, and Br; or a derivative thereof;

30 X¹⁴ is either arginine, lysine, histidine, norarginine, or 5-aminopentanoic acid or a derivative thereof;

X¹⁵ is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof;

X¹⁶ is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

X¹⁷ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

Z¹ is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z² is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;
or a labeled derivative of said peptide;
or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

Unless otherwise stated, those amino acids with a chiral center are provided in the L-enantiomer. Reference to "a derivative thereof" refers to the corresponding D-amino acid, N-alkyl-amino acid and β -amino acid.

Another aspect of the present invention describes a method of screening for a compound able to bind a MCH receptor. The method comprises the step of measuring the ability of the compound to effect binding of a truncated MCH analog to either the MCH receptor, a fragment of the receptor comprising a MCH binding site, a polypeptide comprising such a fragment, or a derivative of the polypeptide.

Another aspect of the present invention describes a method for increasing weight in a subject. The method comprises the step of administering to the subject an effective amount of a truncated MCH analog to produce a weight increase.

Another aspect of the present invention describes a method for increasing appetite in a subject. The method comprises the step of administering to the subject an effective amount of a truncated MCH analog to produce an appetite increase.

Another aspect of the present invention describes a method for measuring the ability of a compound to decrease weight or appetite in a subject. The method comprising the steps of:

a) administering to the subject an effective amount of a truncated MCH analog to produce a weight increase or appetite increase,

- b) administering the compound to the subject, and
- c) measuring the change in weight or appetite of the subject.

Other features and advantages of the present invention are apparent from the additional descriptions provided herein including the different examples.

- 5 The provided examples illustrate different components and methodology useful in practicing the present invention. The examples do not limit the claimed invention. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present invention.

10 BRIEF DESCRIPTION OF THE DRAWINGS

- Figure 1 illustrates the results of an alanine scan where different amino acid residues of human MCH were replaced with alanine. The binding assay was performed by measuring inhibition of (¹²⁵I-tyrosine, phenylalanine¹³)-MCH binding to cloned human MCH receptor (CHO clone). Cyclization sites (S-S) are indicated by
15 “*”.

DETAILED DESCRIPTION OF THE INVENTION

- Truncated MCH analogs contain about 10 to about 17 groups that are amino acids or amino acid derivatives. Using the present application as a guide
20 truncated MCH analogs can be produced having significant MCH receptor activity, and in some cases having activity equal to or better than naturally occurring mammalian MCH. The smaller size of truncated MCH analogs offers advantages over longer-length MCH such as ease of synthesis and/or increased solubility in physiological buffers.

- 25 The MCH receptor is a G-protein coupled receptor that appears to be able to couple to Gi and Gq. Several references describe a receptor that is indicated to be a MCH receptor. (Chambers, *et al.*, 1999. *Nature* 400, 261-265; Saito, *et al.*, 1999. *Nature* 400, 265-269; Bächner, *et al.*, 1999. *FEBS Letters* 457:522-524; and Shimomura, *et al.*, 1999. *Biochemical and Biophysical Research Communications*
30 261, 622-626. These references are not admitted to be prior art to the claimed invention.)

The nucleic acid encoding for different variants of a MCH receptor is provided for by SEQ. ID. NOS. 1-3. The encoded amino acid sequences of the variants are provided by SEQ. ID. NOS. 4-6. The variants differ from each other by

the presence of additional amino acids at the N-terminal. One or more of these variants may be a physiological MCH receptor.

Significant MCH activity is preferably at least about 50%, at least about 75%, at least about 90%, or at least about 95%, the activity of mammalian MCH as determined by a binding assay or MCH receptor activity assay. Examples of such assays are provided below.

MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically. Research tool applications generally involve the use of a truncated MCH analog and the presence of a MCH receptor or fragment thereof. The MCH receptor can be present in different environments such as a mammalian subject, a whole cell, and membrane fragments. Examples of research tool applications of truncated MCH analogs include screening for compounds active at the MCH receptor, determining the presence of the MCH receptor in a sample or preparation, examining the role or effect of MCH, and examining the role or effect of MCH antagonists.

Truncated MCH analogs can be used to screen for both MCH agonists and MCH antagonists. Screening for MCH agonists can be performed, for example, by using a truncated MCH analog in a competition experiment with test compounds. Screening for MCH antagonists can be performed, for example, by using a truncated MCH analog to produce MCH receptor activity and then measuring the ability of a compound to alter MCH receptor activity.

Truncated MCH analogs can be administered to a subject. A "subject" refers to a mammal including, for example, a human, a rat, a mouse, or a farm animal. Reference to subject does not necessarily indicate the presence of a disease or disorder. The term subject includes, for example, mammals being dosed with a truncated MCH analog as part of an experiment, mammals being treated to help alleviate a disease or disorder, and mammals being treated prophylactically to retard or prevent the onset of a disease or disorder.

MCH agonists can be used to achieve a beneficial effect in a subject. For example, a MCH agonist can be used to facilitate a weight gain, maintenance of weight and/or an appetite increase. Such effects are particularly useful for a patient having a disease or disorder, or under going a treatment, accompanied by weight loss. Examples of diseases or disorders accompanied by weight loss include anorexia, AIDS, wasting, cachexia, and frail elderly. Examples of treatments accompanied by weight loss include chemotherapy, radiation therapy, and dialysis.


MCH antagonists can also be used to achieve a beneficial effect in a patient. For example, a MCH antagonist can be used to facilitate weight loss, appetite decrease, weight maintenance, cancer (*e.g.*, colon or breast) treatment, pain reduction, stress reduction and/or treatment of sexual dysfunction.

5

Truncated MCH Analogs

A truncated MCH analog is an optionally modified peptide having the structure:

10



wherein X¹ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof; preferably, X¹ if present is aspartic acid or glutamic acid; more preferably, X¹ if present is aspartic acid; and more preferably, X¹ is not present;

X² is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof; preferably, X² if present is phenylalanine or tyrosine; more preferably, X² if present is phenylalanine; and more preferably, X² is not present;

X³ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X³ if present is aspartic acid or glutamic acid; more preferably, X³ if present is aspartic acid; and more preferably, X³ is not present;

X⁴ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof; preferably, X⁴ if

present is methionine, leucine, isoleucine, valine, alanine or norleucine; more preferably, X⁴ if present is methionine; and more preferably, X⁴ is not present;

X⁵ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X⁵ if present is leucine, methionine, isoleucine, valine or alanine; more preferably, X⁵ if present is leucine; and more preferably, X⁵ is not present;

X⁶ is an optionally present amino acid that, if present is either arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof; preferably X⁶ is not present or is either arginine, D-arginine, D-norleucine, D-proline, D-serine, or D-asparagine; more preferably X⁶ is arginine or D-arginine;

X⁷ is either cysteine, homocysteine, or penicillamine, or a derivative thereof; preferably, X⁷ is cysteine;

X⁸ is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof; preferably, X⁸ is methionine, norleucine, or N-methyl norleucine;

X⁹ is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X⁹ is leucine;

X¹⁰ is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, gamma-aminobutyric acid, asparagine, serine, sarcosine, or isobutyric or a derivative thereof; preferably, X¹⁰ is either glycine, alanine, leucine, norleucine, asparagine, serine, D-norleucine, D-proline, gamma-aminobutyric acid, or sarcosine; more preferably X¹⁰, is either glycine, leucine, norleucine, asparagine, or serine;

X¹¹ is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof; preferably, X¹¹ is arginine;

X¹² is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof; preferably, X¹² is valine;

X¹³ is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO₂, NH₂, F, I, and Br; or a derivative

thereof; preferably, X¹³ is phenylalanine, (2')naphthylalanine, p-fluoro-phenylalanine, tyrosine, or cyclohexylalanine;

X¹⁴ is either arginine, lysine, histidine or norarginine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X¹⁴ is arginine;

5 X¹⁵ is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X¹⁵ is proline or sarcosine;

X¹⁶ is either cysteine, homocysteine, or penicillamine, or a derivative thereof; preferably, X¹⁶ is cysteine or D-cysteine;

10 X¹⁷ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X¹⁷ if present is tyrosine or tryptophan; more preferably X¹⁷ is not present;

15 Z¹ is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z² is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;

or a labeled derivative of said peptide;

20 or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

The present invention is meant to comprehend diastereomers as well as their racemic and resolved enantiomerically pure forms. Truncated MCH analogs can contain D-amino acids, L-amino acids or a combination thereof. Preferably, amino
25 acids present in a truncated MCH analog are the L-enantiomer.

In different embodiments, MCH analogs contain a preferred (or more preferred) group at one or more different locations. More preferred embodiments contain preferred (or more preferred) groups in each of the different locations.

30 A protecting group covalently joined to the N-terminal amino group reduces the reactivity of the amino terminus under *in vivo* conditions. Amino protecting groups include optionally substituted -C₁₋₁₀ alkyl, optionally substituted -C₂₋₁₀ alkenyl, optionally substituted aryl, -C₁₋₆ alkyl optionally substituted aryl, -C(O)-(CH₂)₁₋₆-COOH, -C(O)-C₁₋₆ alkyl, -C(O)-optionally substituted aryl,

-C(O)-O-C₁₋₆ alkyl, or -C(O)-O-optionally substituted aryl. Preferably, the amino terminus protecting group is acetyl, propyl, succinyl, benzyl, benzyloxycarbonyl or *t*-butyloxycarbonyl.

5 A protecting group covalently joined to the C-terminal carboxy group reduces the reactivity of the carboxy terminus under *in vivo* conditions. The carboxy terminus protecting group is preferably attached to the α -carbonyl group of the last amino acid. Carboxy terminus protecting groups include amide, methylamide, and ethylamide.

"Alkyl" refers to carbon atoms joined by carbon-carbon single bonds.
10 The alkyl hydrocarbon group may be straight-chain or contain one or more branches or cyclic groups. Preferably, the alkyl group is 1 to 4 carbons in length. Examples of alkyl include methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, and *t*-butyl. Alkyl substituents are selected from the group consisting of halogen (preferably -F or -Cl) -OH, -CN, -SH, -NH₂, -NO₂, -C₁₋₂ alkyl substituted with 1 to 6 halogens (preferably
15 -F or -Cl, more preferably -F), -CF₃, -OCH₃, or -OCF₃.

"Alkenyl" refers to a hydrocarbon group containing one or more carbon-carbon double bonds. The alkenyl hydrocarbon group may be straight-chain or contain one or more branches or cyclic groups. Preferably, the alkenyl group is 2 to 4 carbons in length. Alkenyl substituents are selected from the group consisting of
20 halogen (preferably -F or -Cl), -OH, -CN, -SH, -NH₂, -NO₂, -C₁₋₂ alkyl substituted with 1 to 5 halogens (preferably -F or -Cl, more preferably -F), -CF₃, -OCH₃, or -OCF₃.

"Aryl" refers to an optionally substituted aromatic group with at least one ring having a conjugated pi- electron system, containing up to two
25 conjugated or fused ring systems. Aryl includes carbocyclic aryl, heterocyclic aryl and biaryl groups. Preferably, the aryl is a 5 or 6 membered ring, more preferably benzyl. Aryl substituents are selected from the group consisting of -C₁₋₄ alkyl, -C₁₋₄ alkoxy, halogen (preferably -F or -Cl), -OH, -CN, -SH, -NH₂, -NO₂, -C₁₋₂ alkyl substituted with 1 to 5 halogens (preferably -F or -Cl, more preferably -F), -CF₃, or
30 -OCF₃.

A labeled derivative indicates the alteration of a substituent with a detectable label. Examples of detectable labels include luminescent, enzymatic, and radioactive labels. A preferred radiolabel is ¹²⁵I. Both the type of label and the position of the label can effect MCH activity. Labels should be selected so as not to
35 substantially alter the activity of the truncated MCH analog at the MCH receptor. The

In naturally occurring full length MCH, alteration of the tyrosine at position 13 by labeling with ^{125}I substantially effects MCH activity. (Drozd, *et al.*, 1995. *FEBS letters* 359, 199-202.) ^{125}I labeled analogs of full length mammalian MCH having substantial activity can be produced, for example, by replacing the tyrosine at position 13 with a different group, then replacing valine at position 19 with tyrosine, and labeling the tyrosine. Examples of such analogs include [^{125}I][Phe¹³, Try¹⁹]-MCH and (D-(*p*-benzoylphenylalanine)¹³, tyrosine¹⁹)-MCH. (Drozd, *et al.*, *FEBS letters* 359, 199-202, 1995; and Drozd, *et al.*, *J. Peptide Sci.* 5, 234-242, 1999.)

15 $\overbrace{z^1-x^6-x^7-x^8-x^9-x^{10}-x^{11}-x^{12}-x^{13}-x^{14}-x^{15}-x^{16}-x^{17}-z^2}$

20 In different embodiments the truncated MCH analog is a peptide of SEQ. ID. NOS. 7, 8, 9, or 10, a labeled derivative of said peptide or a pharmaceutically acceptable salt of said peptide or of said labeled derivative. SEQ. ID. NOS. 7-12 are made up of L-amino acids and have the following sequences (“*” indicates cyclization (S-S)):

35 SEQ. ID. NO. 10:

* *

Asp-Phe-Asp-Met-Leu-Arg-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-amide;

* * *

SEQ. ID. NO. 12: Ac-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-Trp-Gln-Val;

SEQ. ID. NO. 13:

* * *

Asp-Phe-Asp-Nle-Leu-Arg-Cys-Nle-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-Trp-Gln-Val;

SEQ. ID. NO. 14:

* * *

Asp-Phe-Ala-Met-Leu-Arg-Cys-Met-Leu-Gly-Arg-Val-Phe-Arg-Pro-Cys-Trp-Gln-Tyr.

In additional embodiments the peptide has a sequence selected from the group consisting of SEQ. ID. NOs. 7, 8, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 57, 61, 62, 63, 65-67, 69-72, and 77, is a labeled derivative of said peptide or a pharmaceutically acceptable salt of said peptide or of said labeled derivative. Preferred sequences are those with an IC₅₀ less than 0.3 nM, preferably less than 0.1 nM; and/or those having a % activation greater than about 90%, preferably greater than 100%. Examples of preferred sequences are provided in Example 4, Tables 1-7.

Truncated MCH analogs can be produced using techniques well known in the art. For example, a polypeptide region of a truncated MCH analog can be chemically or biochemically synthesized and, if desired modified to produce a blocked N-terminus and/or blocked C-terminus. Techniques for chemical synthesis of polypeptides are well known in the art. (See *e.g.*, Vincent, in *Peptide and Protein Drug Delivery*, New York, N.Y., Dekker, 1990.) Examples of techniques for biochemical synthesis involving the introduction of a nucleic acid into a cell and expression of nucleic acids are provided in Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, and Sambrook, *et al.*, in *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, 1989.

MCH Receptor Binding Assay

Assays measuring the ability of a compound to bind a MCH receptor employ a MCH receptor, a fragment of the receptor comprising a MCH binding site, a

polypeptide comprising such a fragment, or a derivative of the polypeptide. Preferably, the assay uses the MCH receptor or a fragment thereof.

A polypeptide comprising a MCH receptor fragment that binds MCH can also contain one or more polypeptide regions not found in a MCH receptor. A
5 derivative of such a polypeptide comprises a MCH receptor fragment that binds MCH along with one or more non-peptide components.

The MCH receptor amino acid sequence involved in MCH binding can be readily identified using labeled MCH or truncated MCH analogs and different receptor fragments. Different strategies can be employed to select fragments to be
10 tested to narrow down the binding region. Examples of such strategies include testing consecutive fragments about 15 amino acids in length starting at the N-terminus, and testing longer length fragments. If longer length fragments are tested, a fragment binding MCH can be subdivided to further locate the MCH binding region. Fragments used for binding studies can be generated using recombinant nucleic acid
15 techniques.

Binding assays can be performed using individual compounds or preparations containing different numbers of compounds. A preparation containing different numbers of compounds having the ability to bind to the MCH receptor can be divided into smaller groups of compounds that can be tested to identify the
20 compound(s) binding to the MCH receptor. In an embodiment of the present invention a test preparation containing at least 10 compounds is used in a binding assay.

Binding assays can be performed using recombinantly produced MCH receptor polypeptides present in different environments. Such environments include,
25 for example, cell extracts and purified cell extracts containing the MCH receptor polypeptide expressed from recombinant nucleic acid or naturally occurring nucleic acid; and also include, for example, the use of a purified MCH receptor polypeptide produced by recombinant means or from naturally occurring nucleic acid which is introduced into a different environment.

30

Screening for MCH Receptor Active Compounds

Screening for MCH active compounds is facilitated using a recombinantly expressed MCH receptor. Using recombinantly expressed MCH receptor polypeptides offers several advantages such as the ability to express the
35 receptor in a defined cell system so that response to MCH receptor active compounds

can more readily be differentiated from responses to other receptors. For example, the MCH receptor can be expressed in a cell line such as HEK 293, COS 7, and CHO not normally expressing the receptor by an expression vector, wherein the same cell line without the expression vector can act as a control.

5 Screening for MCH receptor active compounds is facilitated through the use of a truncated MCH analog in the assay. The use of a truncated MCH analog in a screening assay provides for MCH receptor activity. The effect of test compounds on such activity can be measured to identify, for example, allosteric modulators and antagonists. Additionally, such assays can be used to identify
10 agonists.

 MCH receptor activity can be measured using different techniques such as detecting a change in the intracellular conformation of the MCH receptor, Gi or Gq activity, and/or intracellular messengers. Gi activity can be measured using techniques well known in the art such as a melonaphore assay, assays measuring
15 cAMP production, inhibition of cAMP accumulation, and binding of ^{35}S -GTP. cAMP can be measured using different techniques such as radioimmunoassay and indirectly by cAMP responsive gene reporter proteins.

 Gq activity can be measured using techniques such as those measuring intracellular Ca^{2+} . Examples of techniques well known in the art that can be
20 employed to measure Ca^{2+} include the use of dyes such as Fura-2 and the use of Ca^{2+} -bioluminescent sensitive reporter proteins such as aequorin. An example of a cell line employing aequorin to measure G-protein activity is HEK293/aeq17. (Button, *et al.*, 1993. *Cell Calcium* 14, 663-671, and Feighner, *et al.*, 1999. *Science* 284, 2184-2188, both of which are hereby incorporated by reference herein.)

25 Chimeric receptors containing a MCH binding region functionally coupled to a G protein can also be used to measure MCH receptor activity. A chimeric MCH receptor contains an N-terminal extracellular domain; a transmembrane domain made up of transmembrane regions, extracellular loop regions, and intracellular loop regions; and an intracellular carboxy terminus.
30 Techniques for producing chimeric receptors and measuring G protein coupled responses are provided for in, for example, International Application Number WO 97/05252, and U.S. Patent Number 5,264,565, both of which are hereby incorporated by reference herein.

Weight or Appetite Alteration

Truncated MCH analogs can be used in methods to increase or maintain weight and/or appetite in a subject. Such methods can be used, for example, as part of an experimental protocol examining the effects of MCH antagonists, to
5 achieve a beneficial effect in a subject and/or to further examine the physiological effects of MCH.

Experimental protocols examining the effects of MCH antagonists can be performed, for example, by using a sufficient amount of a truncated MCH analog to produce a weight or appetite increase in a subject and then examining the effect of
10 a test compound. Changes in weight and appetite can be measured using techniques well known in the art.

Increasing weight or appetite can be useful for maintaining weight or producing a weight or appetite gain in an under weight subject, or in a patient having a disease or undergoing treatment that effects weight or appetite. In addition, for
15 example, farm animals such as pigs, cows and chickens can be treated to gain weight.

Under weight subjects include those having a body weight about 10% or less, 20% or less, or 30% or less, than the lower end of a "normal" weight range or Body Mass Index ("BMI"). "Normal" weight ranges are well known in the art and take into account factors such as a patient age, height, and body type.

20 BMI measures your height/weight ratio. It is determined by calculating weight in kilograms divided by the square of height in meters. The BMI "normal" range is 19-22.

Administration

25 Truncated MCH analogs can be formulated and administered to a subject using the guidance provided herein along with techniques well known in the art. The preferred route of administration ensures that an effective amount of compound reaches the target. Guidelines for pharmaceutical administration in general are provided in, for example, *Remington's Pharmaceutical Sciences 18th Edition*, Ed.
30 Gennaro, Mack Publishing, 1990, and *Modern Pharmaceutics 2nd Edition*, Eds. Banker and Rhodes, Marcel Dekker, Inc., 1990, both of which are hereby incorporated by reference herein.

Truncated MCH analogs can be prepared as acidic or basic salts. Pharmaceutically acceptable salts (in the form of water- or oil-soluble or dispersible
35 products) include conventional non-toxic salts or the quaternary ammonium salts that

are formed, *e.g.*, from inorganic or organic acids or bases. Examples of such salts include acid addition salts such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate; and base salts such as ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine and lysine.

Truncated MCH analogs can be administered using different routes including oral, nasal, by injection, transdermal, and transmucosally. Active ingredients to be administered orally as a suspension can be prepared according to techniques well known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants.

Truncated MCH analogs may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form. When administered by injection, the injectable solution or suspension may be formulated using suitable non-toxic, parenterally-acceptable diluents or solvents, such as Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Suitable dosing regimens are preferably determined taking into factors well known in the art including type of subject being dosed; age, weight, sex and medical condition of the subject; the route of administration; the renal and hepatic function of the subject; the desired effect; and the particular compound employed.

Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug. The daily dose for a subject is expected to be
5 between 0.01 and 1,000 mg per subject per day.

Truncated MCH analogs can be provided in kit. Such a kit typically contains an active compound in dosage forms for administration. A dosage form contains a sufficient amount of active compound such that a weight or appetite increase can be obtained when administered to a subject during regular intervals, such
10 as 1 to 6 times a day, during the course of 1 or more days. Preferably, a kit contains instructions indicating the use of the dosage form for weight or appetite increase and the amount of dosage form to be taken over a specified time period.

EXAMPLES

15 Examples are provided below to further illustrate different features of the present invention. The examples also illustrate useful methodology for practicing the invention. These examples do not limit the claimed invention.

Example 1: Synthesis of MCH Analogs

20 MCH analogs were produced using the procedures described below and varying the stepwise addition of amino acid groups. Other procedures for producing and modifying peptides are well known in the art.

Elongation of peptidyl chains on 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)-phenoxy resin and the acetylation of the N-terminal amino groups of
25 the peptides was performed on a 431A ABI peptide synthesizer. Manufacture-supplied protocols were applied for coupling of the hydroxybenzotriazole esters of amino acids in N-methylpyrrolidone (NMP). The fluorenylmethyloxycarbonyl (Fmoc) group was used as a semipermanent alpha-amino protecting group, whereas the side chains protecting groups were: *tert*-butyl for aspartic acid and tyrosine,
30 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf) for arginine, and trityl for cysteine.

Peptides were cleaved from the resin with TFA containing 5 % of anisole. After 2 hours at room temperature the resin was filtered, washed with TFA and the combined filtrates were evaporated to dryness in vacuo. The residue was

trituated with ether, the precipitate which formed was filtered off, washed with ether, and dried.

Crude peptides were dissolved in 5 % acetic acid in water, and the pH of the solutions were adjusted to ca. 8.2 with diluted ammonium hydroxide. The reaction mixtures were stirred vigorously while 0.05 % solution of potassium ferricyanide ($K_3Fe(CN)_6$) in water was added dropwise till the reaction mixture remained yellow for about 5 minutes. After an additional 20 minutes oxidation was terminated with ca. 1 ml of acetic acid and the reaction mixtures were lyophilized.

Crude lyophilized peptides were analyzed by analytical reverse-phase high-pressure liquid chromatography (RP HPLC) on a C18 Vydac column attached to a Waters 600E system with automatic Wisp 712 injector and 991 Photodiode Array detector. A standard gradient system of 0-100% buffer B in 30 minutes was used for analysis: buffer A was 0.1% trifluoroacetic acid in water and buffer B was 0.1% trifluoroacetic acid in acetonitrile. HPLC profiles were recorded at 210 nm and 280 nm. Preparative separations were performed on a Waters Delta Prep 4000 system with a semipreparative C18 RP Waters column. The above-described solvent system of water and acetonitrile, in a gradient of 20-80 % buffer B in 60 minutes, was used for separation. The chromatographically homogenous compounds were analyzed by electrospray mass spectrometry.

20

Example 2: Aequorin Bioluminescence Functional Assay

The aequorin bioluminescence assay is a reliable test for measuring the activity of G protein-coupled receptors that couple through the $G\alpha$ protein subunit family consisting of G_q and G_{11} and leads to the activation of phospholipase C, mobilization of intracellular calcium and activation of protein kinase C.

Measurement of MCH receptor activity in the aequorin-expressing stable reporter cell line 293-AEQ17 (Button *et al.*, *Cell Calcium* 14:663-671, 1993) was performed using a Luminoskan RT luminometer (Labsystems Inc., Gaithersburg, MD). 293-AEQ17 cells (8×10^5 cells plated 18 hours before transfection in a T75 flask) were transfected with 22 μ g of human MCH receptor plasmid using 264 μ g lipofectamine. The open reading frame cDNA (SEQ. ID. NO. 1) encoding the human MCH receptor inserted in the mammalian expression vector pcDNA-3 (Invitrogen, Carlsbad, CA) was used for expression studies. Following approximately 40 hours of expression the apo-aequorin in the cells was charged for 4 hours with coelenterazine (10 μ M) under reducing conditions (300 μ M reduced glutathione) in ECB buffer (140

35

mM NaCl, 20 mM KCl, 20 mM HEPES-NaOH [pH=7.4], 5 mM glucose, 1 mM MgCl₂, 1 mM CaCl₂, 0.1 mg/ml bovine serum albumin).

The cells were harvested, washed once in ECB medium and resuspended to 500,000 cells/ml. 100 µl of cell suspension (corresponding to 5x10⁴ cells) was then injected into the test plate containing MCH or MCH analogs, and the integrated light emission was recorded over 30 seconds, in 0.5 second units. 20 µL of lysis buffer (0.1% final Triton X-100 concentration) was then injected and the integrated light emission recorded over 10 seconds, in 0.5 second units. The “fractional response” values for each well were calculated by taking the ratio of the integrated response to the initial challenge to the total integrated luminescence including the Triton X-100 lysis response.

Example 3: Radiolabeled MCH-R Binding Assay

Activity of truncated MCH analogs was assayed by measuring the ability of the analog to inhibit binding of [¹²⁵I]-human MCH (Phe¹³, Tyr¹⁹ substituted) to membranes prepared from cells stably expressing the human MCH receptor. Human MCH (Phe¹³, Tyr¹⁹ substituted) used in the assay was radiolabeled with ¹²⁵I at ¹⁹Tyr to a specific activity of ~2000 Ci/mmol (NEN Life Science Products, Boston, MA).

Cell membranes were prepared on ice. Each T-75 flask was rinsed twice with 10 ml of Enzyme-free Cell Dissociation Buffer (Specialty Media, Lavallete, NJ), and the cell monolayer was detached in an additional 10 ml of Enzyme-free Cell Dissociation Buffer by incubation at room temperature for 10 minutes. Dissociated cells were centrifuged (500 x g for 10 minutes at 4°C), resuspended in 5 ml homogenization buffer (10 mM Tris-HCl, pH 7.4, 0.01 mM Pefabloc, 10 µM phosphoramidon, 40 µg/ml bacitracin) and then homogenized using a glass homogenizer (10-15 strokes). The homogenate was centrifuged for 10 minutes (1,000 x g at 4°C). The resulting supernatant was then centrifuged at 38,700 x g for 15 minutes at 4°C. Pelleted membranes were resuspended (passed through 25 gauge needle 5 times), snap-frozen on liquid nitrogen, and stored at -80°C until use.

Binding was performed in a 96-well filter assay or Scintillation Proximity Assay (SPA)-based format using cell membranes from a stable CHO or HEK-293 cell line expressing the MCH receptor. For the filter assay, reactions were performed at 20°C for 1 hour in a total volume of 0.2 ml containing: 0.05 ml of membrane suspension (~3 µg protein), 0.02 ml of [¹²⁵I]-human MCH (Phe¹³, Tyr¹⁹

substituted; 30 pM), 0.01 ml of competitor and 0.12 ml of binding buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 2 mM EDTA, 200 µg/ml bacitracin, 1 µM phosphoramidon).

Bound radioligand was separated by rapid vacuum filtration (Packard
5 Filtermate 96-well cell harvester) through GF/C filters pretreated for 1 hour with 1 % polyethylenimine. After application of the membrane suspension to the filter, the filters were washed 3 times with 3 ml each of ice-cold 50 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 2 mM EDTA, 0.04 % Tween 20 and the bound radioactivity on the filters was quantitated by scintillation counting (TopCount device). Specific binding
10 (>80 % of total) is defined as the difference between total binding and non-specific binding conducted in the presence of 100 nM unlabeled human MCH.

For the SPA-based assay, WGA-PVT beads (NEN Life Sciences Products) were resuspended in Dulbecco's PBS with calcium and magnesium (500 mg beads in 4 ml PBS). For each 96-well assay plate, 0.18 ml of beads was pre-
15 coated with MCH receptor by mixing with 0.2 ml MCH receptor CHO cell membranes (~ 0.2-4 mg protein) and 1.5 ml SPA assay buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 2 mM EDTA, 0.1 % BSA, 12 % glycerol). The suspension was mixed gently for 20 minutes, 12.3 ml of assay buffer and protease inhibitors were added (final concentration given): 2 µg/ml leupeptin, 10 µM phosphoramidon, 40
20 µg/ml bacitracin, 5 µg/ml aprotinin, 0.1 mM Pefabloc.

Coated beads were kept on ice until use. For each well, 0.145 ml of beads were added to Optiplate assay plates (Packard 6005190), followed by 0.002-0.004 ml of competitor and 0.05 ml of [¹²⁵I]-human MCH (Phe¹³, Tyr¹⁹ substituted; 30 pM). Binding reactions were allowed to proceed at room temperature for 3 hours.
25 Quantitation was performed by scintillation counting (TopCount device).

Example 4: MCH Activity

The activity of different MCH analogs was measured using the procedures described in Examples 2 and 3 above. Tables 1-7 illustrate the activity of
30 different truncated MCH analogs and mammalian MCH (SEQ. ID. NO. 11). Figure 1 illustrates the results of replacing different amino acids of mammalian MCH with alanine. Based on the guidance provided herein, additional MCH analogs active at the MCH receptor can be obtained.

TABLE 1

SEQ. ID. NO.	Binding Assay IC ₅₀ (nM)	EC ₅₀ (nM)	% Activation at 10 μM
11	0.3	36	100
7	0.12	18	123
8	0.16	36	123
9	1.6	300	74
10	0.3		99
12	6.4	492	3
13	1.5		65.9
14	0.5		62.2

IC₅₀ was determined using a SPA based assay.
EC₅₀ (nM) and % Activation at 10 μM were determined using aequorin functional assays.

5

Table 2 illustrates the affect of different D-amino acids.

TABLE 2

Ac-Arg ⁶ -Cys ⁷ -Met ⁸ -Leu ⁹ -Gly ¹⁰ -Arg ¹¹ -Val ¹² -Tyr ¹³ -Arg ¹⁴ -Pro ¹⁵ -Cys ¹⁶ -NH ₂				
SEQ. ID. NO.	Compound	Activity		
		IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		0.5	20	99
15	D-Arg ⁶	0.46	45	86
16	D-Cys ⁷	7.78	909	34
17	D-Met ⁸	>1000	Inactive	
18	D-Leu ⁹	1520	Inactive	
19	D-Arg ¹¹	>1000	Inactive	
20	D-Val ¹²	381	Inactive	
21	D-Tyr ¹³	>1000	Inactive	
22	D-Arg ¹⁴	368	Inactive	
23	D-Pro ¹⁵	584	Inactive	
24	D-Cys ¹⁶	0.8	133	76

10

Table 3 illustrates the effect of different N-methyl-amino acids.

TABLE 3

$\text{Ac-Arg}^6\text{-Cys}^7\text{-Met}^8\text{-Leu}^9\text{-Gly}^{10}\text{-Arg}^{11}\text{-Val}^{12}\text{-Tyr}^{13}\text{-Arg}^{14}\text{-Pro}^{15}\text{-Cys}^{16}\text{-NH}_2$				
SEQ. ID. NO.	Compound	Activity		
		IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
25	N-Me-Nle ⁸	0.16	20	110
26	N-Me-Leu ⁹	10% @ 1	>10000	3
27	Sar ¹⁰	2.3	140	95
28	N-Me-Arg ¹¹	43	10	110
29	N-Me-Arg ¹⁴	643	>1000	
30	Sar ¹⁵	0.36	25	113

Table 4 illustrates the affect of different alterations to position 6 of the
 5 SEQ. ID. NO. 7 MCH analog.

TABLE 4

$\text{X}^6\text{-Cys}^7\text{-Met}^8\text{-Leu}^9\text{-Gly}^{10}\text{-Arg}^{11}\text{-Val}^{12}\text{-Tyr}^{13}\text{-Arg}^{14}\text{-Pro}^{15}\text{-Cys}^{16}\text{-NH}_2$				
SEQ. ID. NO.	Position 6 modification	Activity		
		IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
31	Ac-Ala	27	114	135
32	Ac-Nle	40	117	107
33	Ac-Pro	3.4	59	133
34	Ac-Asn	2.6	150	96
35	Ac-Ser	4.5	207	120
36	Ac-Glu	19	935	113
37	H	12	809	120
38	Ac	1.6	144	82
39	Arg	0.13	14	106
40	Δ NH ₂ -Arg	0.48	38.5	49
41	Ac-D-Arg	0.46	45	86
42	Ac-D-Nle	1.2	110	97
43	Ac-D-Pro	0.82	60	96
44	Ac-D-Asn	3	340	94
45	Ac-D-Ser	2.3	170	93
46	Ac-D-Glu	8	820	85

Table 5 illustrates the affect of different alterations to position 10 of the SEQ. ID. NO. 7 MCH analog.

TABLE 5

5

$\text{Ac-Arg}^6\text{-Cys}^7\text{-Met}^8\text{-Leu}^9\text{-X}^{10}\text{-Arg}^{11}\text{-Val}^{12}\text{-Tyr}^{13}\text{-Arg}^{14}\text{-Pro}^{15}\text{-Cys}^{16}\text{-NH}_2$				
SEQ. ID. NO.	Position 10 modification	Activity		
		IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		0.5	20	99
47	Ala	0.59	31	104
48	Leu	0.06	23	106
49	Nle	0.04	15	106
50	Pro	700	519	4
51	Asn	0.23	23	106
52	Ser	0.32	65	104
53	Lys	110	4500	25
54	Glu	190	> 10000	12
55	D-Leu	16	750	23
56	D-Nle	2.4	215	33
57	D-Pro	1.2	190	90
58	D-Glu	40% @ 1	>10000	
59	D-Lys	>1000	>10000	
60	β-Ala	390	>1000	3.2
61	γ-Abu	2.1	30.6	101

Table 6 illustrates the affect of different alterations to position 13 of the SEQ. ID. NO. 7 MCH analog.

TABLE 6

Ac-Arg ⁶ -Cys ⁷ -Met ⁸ -Leu ⁹ -Gly ¹⁰ -Arg ¹¹ -Val ¹² -Tyr ¹³ -Arg ¹⁴ -Pro ¹⁵ -Cys ¹⁶ -NH ₂				
		Activity		
SEQ. ID. NO.	Position 13 modification	IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
62	Phe	1	46	96
63	Trp	3.8	890	83
64	His	13.1	3400	66
65	(2')Nal	0.15	54	105
66	Phe(pF)	0.6	108	98
67	Phe(pNH ₂)	3.2	610	88
68	Phe(pCOOH)	>1000	>10000	
69	Cha	0.09	122	93

Table 7 illustrates the affect of some alteration combinations and some
5 alterations to position 8 of the SEQ. ID. NO. 7 MCH analog.

TABLE 7

Ac-Arg ⁶ -Cys ⁷ -Met ⁸ -Leu ⁹ -Gly ¹⁰ -Arg ¹¹ -Val ¹² -Tyr ¹³ -Arg ¹⁴ -Pro ¹⁵ -Cys ¹⁶ -NH ₂				
		Activity		
SEQ. ID. NO.	Compound	IC ₅₀ (nM)	EC ₅₀ (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
70	Ava ^{9,10}	3.7	587	82
71	D-Arg ⁶ ,Ava ^{9,10}	3.7	1080	72
72	Ava ^{14,15}	6.2	406	75
73	D-Arg ⁶ ,Ava ^{14,15}	19.5	1300	28
74	D-Pro ¹⁰ ,Ava ^{14,15}	700	1530	3
75	ΔArg ⁶ ,Ava ^{14,15}	250	>10000	3
76	Ava ^{9,10} ,Ava ^{14,15}	50	> 10000	3
77	Nle ⁸	0.5	44	105
78	ΔArg ⁶ ,D-Nle ¹⁰	25	72	4

Other embodiments are within the following claims. While several embodiments have been shown and described, various modifications may be made without departing from the spirit and scope of the present invention.

SEQUENCE LISTING

<110> Merck & Co., Inc.

<120> MELANIN-CONCENTRATING HORMONE ANALOGS

<130> 20590Y PCT

<150> 60/179,967

<151> 2000-02-03

<160> 78

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1062

<212> DNA

<213> Human

<400> 1

atggacctgg	aagcctcgct	gctgcccact	ggtecccaacg	ccagcaacac	ctctgatggc	60
cccgataacc	tcacttcggc	aggatcacct	cctcgcacgg	ggagcatctc	ctacatcaac	120
atcatcatgc	cttcggtgtt	cggcaccatc	tgcttcctgg	gcatcatcgg	gaactccacg	180
gtcatcttcg	cggtcgtgaa	gaagtccaag	ctgcactggg	gcaacaacgt	ccccgacatc	240
ttcatcatca	acctctcggt	agtagatctc	ctctttctct	tgggcatgcc	cttcatgata	300
caccagctca	tgggcaatgg	ggtgtggcac	tttggggaga	ccatgtgcac	cctcatcacg	360
gccatggatg	ccaatagtca	gttcaccagc	acctacatcc	tgaccgccat	ggccattgac	420
cgctacctgg	ccactgtcca	ccccatctct	tccacgaagt	tccggaagcc	ctctgtggcc	480
acctgggtga	tctgcctcct	gtggggccctc	tccttcatca	gcatcacccc	tgtgtggctg	540
tatgccagac	tcateccctt	cccaggaggt	gcagtgggct	gcggcatacg	cctgcccac	600
ccagacactg	acctctactg	gttcaccctg	taccagtttt	tcttggcctt	tgccctgcct	660
tttgtgggtca	tcacagccgc	atacgtgagg	atcctgcagc	gcatgacgtc	ctcagtggcc	720
cccgccctccc	agcgcagcat	ccggctgcgg	acaaagaggg	tgacccgcac	agccatcgcc	780
atctgtcttg	tcttctttgt	gtgctgggca	ccctactatg	tgctacagct	gacccagttg	840
tccatcagcc	gcccgaacct	cacctttgtc	tacttataca	atgcggccat	cagcttgggc	900
tatgccaaca	gctgcctcaa	cccctttgtg	tacatcgtgc	tctgtgagac	gttccgcaaa	960
cgcttggtcc	tgtcggtgaa	gcctgcagcc	caggggcagc	ttcgcgctgt	cagcaacgct	1020
cagacggctg	acgaggagag	gacagaaagc	aaaggcacct	ga		1062

<210> 2

<211> 1254

<212> DNA

<213> Human

<400> 2

atgaagaagg	gagtggggag	ggcagttggg	cttggaggcg	gcagcggctg	ccaggctacg	60
gaggaagacc	cccttcccaa	ctgcggggct	tgcgctccgg	gacaagggtg	caggcgctgg	120
aggctgccgc	agcctgcgtg	ggtggagggg	agctcagctc	ggttgtggga	gcaggcgacc	180
ggcactggct	ggatggacct	ggaagcctcg	ctgctgccc	ctgggtccca	cgccagcaac	240
acctctgatg	gccccgataa	cctcacttcg	gcaggatcac	ctcctcgcac	ggggagcatc	300
tcctacatca	acatcatcat	gccttcgggtg	ttcggcacca	tctgcctcct	gggcatcatc	360
gggaactcca	cggtcattct	cgcggtcgtg	aagaagtcca	agctgcactg	gtgcaacaac	420
gtccccgaca	tcttcatcat	caacctctcg	gtagtagatc	tcctctttct	cctgggcatg	480
cccttcatga	tccaccagct	catgggcaat	ggggtgtggc	actttgggga	gaccatgtgc	540
acctcatca	cggccatgga	tgccaatagt	cagttcacca	gcacctacat	cctgaccgcc	600
atggccattg	accgctacct	ggccactgtc	caccccatct	cttccacgaa	gttccggaag	660
ccctctgtgg	ccaccctggg	gatctgcctc	ctgtggggcc	tctccttcat	cagcatcacc	720
cctgtgtggc	tgtatgccag	actcatcccc	ttcccaggag	gtgcagtggtg	ctgcggcata	780
cgcctgccc	acccagacac	tgacctctac	tggttcaccc	tgtaccagtt	tttccctggc	840

WO 01/57070

PCT/US01/03293

tttgccctgc	cttttgtggt	catcacagcc	gcatacgtga	ggatcctgca	gcgcatgacg	900
tcctcagtgg	ccccgcctc	ccagcgcagc	atccggctgc	ggacaaagag	ggtgacccgc	960
acagccatcg	ccatctgtct	ggtcttcttt	gtgtgctggg	caccctacta	tgtgctacag	1020
ctgacccagt	tgtccatcag	ccgcccgcacc	ctcacctttg	tctacttata	caatgcggcc	1080
atcagcttgg	gctatgccaa	cagctgcctc	aacccttttg	tgtacatcgt	gctctgtgag	1140
acgttccgca	aacgcttggg	cctgtcgggtg	aagcctgcag	cccaggggca	gcttcgcgct	1200
gtcagcaacg	ctcagacggc	tgacgaggag	aggacagaaa	gcaaaggcac	ctga	1254

<210> 3

<211> 1269

<212> DNA

<213> Human

<400> 3

atgtcagtgg	gagccatgaa	gaagggagtg	gggagggcag	ttgggcttgg	aggcggcagc	60
ggctgccagg	ctacggagga	agaccccctt	cccaactgcg	gggcttgccg	tccgggacaa	120
ggtggcaggc	gctggaggct	gccgcagcct	gcgtgggttg	aggggagctc	agctcggttg	180
tgggagcagg	cgaccggcac	tggctggatg	gacctggaag	cctcgtgctg	gccactggt	240
cccaacgcc	gcaacacctc	tgatggcccc	gataacctca	cttcggcagg	atcacctcct	300
cgcacgggga	gcattctccta	catcaacatc	atcatgcctt	cggtgttcgg	caccatctgc	360
ctcctgggca	tcacgaggaa	ctccacggtc	atcttcgcgg	tcgtgaagaa	gtccaagctg	420
cactggtgca	acaacgtccc	cgacatcttc	atcatcaacc	tctcggtagt	agatctcctc	480
tttctcctgg	gcattgccctt	catgatccac	cagctcatgg	gcaatggggg	gtggcacttt	540
ggggagacca	tgtgcaccct	catcacggcc	atggatgcc	atagtcagtt	caccagcacc	600
tacatcctga	ccgccatggc	cattgaccgc	tacctggcca	ctgtccaccc	catctcttcc	660
acgaagtcc	ggaagccctc	tgtggccacc	ctggtgatct	gcctcctgtg	ggccctctcc	720
ttcatcagca	tcacccctgt	gtggctgtat	gccagactca	tccccttccc	aggaggtgca	780
gtgggctgcg	gcatacgcct	gcccaccca	gacactgacc	tctactgggt	caccctgtac	840
cagtttttcc	tggcctttgc	cctgcctttt	gtggctcatc	cagccgcata	cgtgaggatc	900
ctgcagcgca	tgacgtcctc	agtggccccc	gcctcccagc	gcagcatccg	gctgcggaca	960
aagaggggtga	ccgcacacag	catcgccatc	tgtctgggtc	tctttgtgtg	ctgggcaccc	1020
tactatgtgc	tacagctgac	ccagttgtcc	atcagccgcc	cgaccctcac	ctttgtctac	1080
ttatacaatg	cggccatcag	cttgggctat	gccaacagct	gcctcaaccc	ctttgtgtac	1140
atcgtgctct	gtgagacgtt	ccgcaaacgc	ttggctcctg	cggtgaagcc	tgcagcccag	1200
gggcagcttc	gcgctgtcag	caacgctcag	acggctgacg	aggagaggac	agaaagcaaa	1260
ggcacctga						1269

<210> 4

<211> 353

<212> PRT

<213> Human

<400> 4

Met	Asp	Leu	Glu	Ala	Ser	Leu	Leu	Pro	Thr	Gly	Pro	Asn	Ala	Ser	Asn
1				5					10					15	
Thr	Ser	Asp	Gly	Pro	Asp	Asn	Leu	Thr	Ser	Ala	Gly	Ser	Pro	Pro	Arg
			20					25					30		
Thr	Gly	Ser	Ile	Ser	Tyr	Ile	Asn	Ile	Ile	Met	Pro	Ser	Val	Phe	Gly
			35				40					45			
Thr	Ile	Cys	Leu	Leu	Gly	Ile	Ile	Gly	Asn	Ser	Thr	Val	Ile	Phe	Ala
			50			55					60				
Val	Val	Lys	Lys	Ser	Lys	Leu	His	Trp	Cys	Asn	Asn	Val	Pro	Asp	Ile
65					70				75					80	
Phe	Ile	Ile	Asn	Leu	Ser	Val	Val	Asp	Leu	Phe	Leu	Leu	Gly	Met	
			85					90					95		
Pro	Phe	Met	Ile	His	Gln	Leu	Met	Gly	Asn	Gly	Val	Trp	His	Phe	Gly
			100					105					110		
Glu	Thr	Met	Cys	Thr	Leu	Ile	Thr	Ala	Met	Asp	Ala	Asn	Ser	Gln	Phe
			115				120					125			
Thr	Ser	Thr	Tyr	Ile	Leu	Thr	Ala	Met	Ala	Ile	Asp	Arg	Tyr	Leu	Ala
			130				135				140				

Thr	Val	His	Pro	Ile	Ser	Ser	Thr	Lys	Phe	Arg	Lys	Pro	Ser	Val	Ala
145					150					155					160
Thr	Leu	Val	Ile	Cys	Leu	Leu	Trp	Ala	Leu	Ser	Phe	Ile	Ser	Ile	Thr
				165					170						175
Pro	Val	Trp	Leu	Tyr	Ala	Arg	Leu	Ile	Pro	Phe	Pro	Gly	Gly	Ala	Val
			180					185					190		
Gly	Cys	Gly	Ile	Arg	Leu	Pro	Asn	Pro	Asp	Thr	Asp	Leu	Tyr	Trp	Phe
		195					200					205			
Thr	Leu	Tyr	Gln	Phe	Phe	Leu	Ala	Phe	Ala	Leu	Pro	Phe	Val	Val	Ile
	210					215					220				
Thr	Ala	Ala	Tyr	Val	Arg	Ile	Leu	Gln	Arg	Met	Thr	Ser	Ser	Val	Ala
225					230					235					240
Pro	Ala	Ser	Gln	Arg	Ser	Ile	Arg	Leu	Arg	Thr	Lys	Arg	Val	Thr	Arg
				245					250					255	
Thr	Ala	Ile	Ala	Ile	Cys	Leu	Val	Phe	Phe	Val	Cys	Trp	Ala	Pro	Tyr
			260					265					270		
Tyr	Val	Leu	Gln	Leu	Thr	Gln	Leu	Ser	Ile	Ser	Arg	Pro	Thr	Leu	Thr
		275					280					285			
Phe	Val	Tyr	Leu	Tyr	Asn	Ala	Ala	Ile	Ser	Leu	Gly	Tyr	Ala	Asn	Ser
	290					295					300				
Cys	Leu	Asn	Pro	Phe	Val	Tyr	Ile	Val	Leu	Cys	Glu	Thr	Phe	Arg	Lys
305					310					315					320
Arg	Leu	Val	Leu	Ser	Val	Lys	Pro	Ala	Ala	Gln	Gly	Gln	Leu	Arg	Ala
				325					330					335	
Val	Ser	Asn	Ala	Gln	Thr	Ala	Asp	Glu	Glu	Arg	Thr	Glu	Ser	Lys	Gly
			340					345					350		
Thr															

<210> 5
 <211> 417
 <212> PRT
 <213> Human

<400> 5

Met	Lys	Lys	Gly	Val	Gly	Arg	Ala	Val	Gly	Leu	Gly	Gly	Gly	Ser	Gly
1				5					10					15	
Cys	Gln	Ala	Thr	Glu	Glu	Asp	Pro	Leu	Pro	Asn	Cys	Gly	Ala	Cys	Ala
			20					25					30		
Pro	Gly	Gln	Gly	Gly	Arg	Arg	Trp	Arg	Leu	Pro	Gln	Pro	Ala	Trp	Val
		35					40					45			
Glu	Gly	Ser	Ser	Ala	Arg	Leu	Trp	Glu	Gln	Ala	Thr	Gly	Thr	Gly	Trp
	50					55					60				
Met	Asp	Leu	Glu	Ala	Ser	Leu	Leu	Pro	Thr	Gly	Pro	Asn	Ala	Ser	Asn
65					70					75					80
Thr	Ser	Asp	Gly	Pro	Asp	Asn	Leu	Thr	Ser	Ala	Gly	Ser	Pro	Pro	Arg
				85					90					95	
Thr	Gly	Ser	Ile	Ser	Tyr	Ile	Asn	Ile	Ile	Met	Pro	Ser	Val	Phe	Gly
			100					105					110		
Thr	Ile	Cys	Leu	Leu	Gly	Ile	Ile	Gly	Asn	Ser	Thr	Val	Ile	Phe	Ala
		115					120					125			
Val	Val	Lys	Lys	Ser	Lys	Leu	His	Trp	Cys	Asn	Asn	Val	Pro	Asp	Ile
	130					135					140				
Phe	Ile	Ile	Asn	Leu	Ser	Val	Val	Asp	Leu	Leu	Phe	Leu	Leu	Gly	Met
145					150					155					160
Pro	Phe	Met	Ile	His	Gln	Leu	Met	Gly	Asn	Gly	Val	Trp	His	Phe	Gly
				165					170					175	
Glu	Thr	Met	Cys	Thr	Leu	Ile	Thr	Ala	Met	Asp	Ala	Asn	Ser	Gln	Phe
			180					185					190		
Thr	Ser	Thr	Tyr	Ile	Leu	Thr	Ala	Met	Ala	Ile	Asp	Arg	Tyr	Leu	Ala
		195					200					205			

WO 01/57070

PCT/US01/03293

```

Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Val Ala
 210                215                220
Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr
225                230                235                240
Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val
                245                250                255
Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe
                260                265                270
Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile
                275                280                285
Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala
                290                295                300
Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg
305                310                315                320
Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr
                325                330                335
Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr
                340                345                350
Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser
                355                360                365
Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys
370                375                380
Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Ala
385                390                395                400
Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly
                405                410                415
Thr

```

```

<210> 6
<211> 422
<212> PRT
<213> Human

```

```

<400> 6
Met Ser Val Gly Ala Met Lys Lys Gly Val Gly Arg Ala Val Gly Leu
 1                5                10                15
Gly Gly Gly Ser Gly Cys Gln Ala Thr Glu Glu Asp Pro Leu Pro Asn
                20                25                30
Cys Gly Ala Cys Ala Pro Gly Gln Gly Gly Arg Arg Trp Arg Leu Pro
35                40                45
Gln Pro Ala Trp Val Glu Gly Ser Ser Ala Arg Leu Trp Glu Gln Ala
50                55                60
Thr Gly Thr Gly Trp Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly
65                70                75                80
Pro Asn Ala Ser Asn Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala
                85                90                95
Gly Ser Pro Pro Arg Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met
100                105                110
Pro Ser Val Phe Gly Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser
115                120                125
Thr Val Ile Phe Ala Val Val Lys Lys Ser Lys Leu His Trp Cys Asn
130                135                140
Asn Val Pro Asp Ile Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu
145                150                155                160
Phe Leu Leu Gly Met Pro Phe Met Ile His Gln Leu Met Gly Asn Gly
165                170                175
Val Trp His Phe Gly Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp
180                185                190
Ala Asn Ser Gln Phe Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile
195                200                205

```


WO 01/57070

PCT/US01/03293

```

Asp Arg Tyr Leu Ala Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg
 210                215                220
Lys Pro Ser Val Ala Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser
225                230                235                240
Phe Ile Ser Ile Thr Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe
                245                250                255
Pro Gly Gly Ala Val Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr
                260                265                270
Asp Leu Tyr Trp Phe Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu
                275                280                285
Pro Phe Val Val Ile Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met
                290                295                300
Thr Ser Ser Val Ala Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr
305                310                315                320
Lys Arg Val Thr Arg Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val
                325                330                335
Cys Trp Ala Pro Tyr Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser
                340                345                350
Arg Pro Thr Leu Thr Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu
                355                360                365
Gly Tyr Ala Asn Ser Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys
                370                375                380
Glu Thr Phe Arg Lys Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln
385                390                395                400
Gly Gln Leu Arg Ala Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg
                405                410                415
Thr Glu Ser Lys Gly Thr
                420

```

```

<210> 7
<211> 11
<212> PRT
<213> Artificial Sequence

```

```

<220>
<221> ACETYLATION
<222> (1)...(1)

```

```

<221> AMIDATION
<222> (11)...(11)

```

```

<221> DISULFID
<222> (2)...(11)

```

```

<223> MCH Analog

```

```

<400> 7
Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1                5                10

```

```

<210> 8
<211> 12
<212> PRT
<213> Artificial Sequence

```

```

<220>
<221> ACETYLATION
<222> (1)...(1)

```

```

<221> AMIDATION
<222> (12)...(12)

```


WO 01/57070

PCT/US01/03293

<221> DISULFID
 <222> (2)...(11)

<223> MCH Analog

<400> 8
 Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys Tyr
 1 5 10

<210> 9
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> ACETYLTATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (10)...(10)

<221> DISULFID
 <222> (1)...(10)

<223> MCH Analog

<400> 9
 Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 10
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> AMIDATION
 <222> (16)...(16)

<221> DISULFID
 <222> (7)...(16)

<223> MCH Analog

<400> 10
 Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10 15

<210> 11
 <211> 19
 <212> PRT
 <213> Human

<220>
 <221> DISULFID
 <222> (7)...(16)

<400> 11
 Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10 15
 Trp Gln Val

WO 01/57070

PCT/US01/03293

<210> 12
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (1)...(10)

<221> ACETYLATION
 <222> (1)...(1)

<400> 12
 Cys Met Leu Gly Arg Val Tyr Arg Pro Cys Trp Gln Val
 1 5 10

<210> 13
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (7)...(16)

<221> MOD_RES
 <222> (4)...(4)
 <223> Xaa = Norleucine

<221> MOD_RES
 <222> (8)...(8)
 <223> Xaa = Norleucine

<400> 13
 Asp Phe Asp Xaa Leu Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10 15
 Trp Gln Val

<210> 14
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (7)...(16)

<400> 14
 Asp Phe Ala Met Leu Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys
 1 5 10 15
 Trp Gln Tyr

<210> 15

WO 01/57070

PCT/US01/03293

<211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analogs

<221> DISULFID
 <222> (2)...(11)

<221> ACETYLTATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = D-Arginine

<221> AMIDATION
 <222> (11)...(11)

<400> 15
 Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 16
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (2)...(11)

<221> MOD_RES
 <222> (2)...(2)
 <223> Xaa = D-Cysteine

<221> ACETYLTATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<400> 16
 Arg Xaa Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 17
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (2)...(11)

<221> MOD_RES
 <222> (3)...(3)

WO 01/57070

PCT/US01/03293

<223> Xaa = D-Methionine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 17

Arg	Cys	Xaa	Leu	Gly	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 18

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> DISULFID

<222> (2)...(11)

<221> MOD_RES

<222> (4)...(4)

<223> Xaa = D-Leucine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 18

Arg	Cys	Met	Xaa	Gly	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 19

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> DISULFID

<222> (2)...(11)

<221> MOD_RES

<222> (6)...(6)

<223> Xaa = D-Arginine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 19

Arg	Cys	Met	Leu	Gly	Xaa	Val	Tyr	Arg	Pro	Cys
1				5					10	

WO 01/57070

PCT/US01/03293

<210> 20
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (2)...(11)

<221> MOD_RES
 <222> (7)...(7)
 <223> Xaa = D-Valine

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<400> 20
 Arg Cys Met Leu Gly Arg Xaa Tyr Arg Pro Cys
 1 5 10

<210> 21
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> DISULFID
 <222> (2)...(11)

<221> MOD_RES
 <222> (8)...(8)
 <223> Xaa = D-Tyrosine

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<400> 21
 Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys
 1 5 10

<210> 22
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES

WO 01/57070

PCT/US01/03293

<222> (9)...(9)
 <223> Xaa = D-Arginine

 <221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 22
 Arg Cys Met Leu Gly Arg Val Tyr Xaa Pro Cys
 1 5 10

 <210> 23
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> MOD_RES
 <222> (10)...(10)
 <223> Xaa = D-Proline

 <221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 23
 Arg Cys Met Leu Gly Arg Val Tyr Arg Xaa Cys
 1 5 10

 <210> 24
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> MOD_RES
 <222> (11)...(11)
 <223> Xaa = D-Cysteine

 <221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 24
 Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Xaa
 1 5 10

<210> 25
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (3)...(3)
 <223> Xaa = N-Me-Norleucine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 25
 Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 26
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (4)...(4)
 <223> Xaa = N-Me-Leucine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 26
 Arg Cys Met Xaa Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 27
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
<222> (5)...(5)
<223> Xaa = MeGly

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 27
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
1 5 10

<210> 28
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLTATION
<222> (1)...(1)

<221> MOD_RES
<222> (6)...(6)
<223> Xaa = N-Me-Arginine

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 28
Arg Cys Met Leu Gly Xaa Val Tyr Arg Pro Cys
1 5 10

<210> 29
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLTATION
<222> (1)...(1)

<221> MOD_RES
<222> (9)...(9)
<223> Xaa = N-Me-Arginine

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 29

Arg Cys Met Leu Gly Arg Val Tyr Xaa Pro Cys
 1 5 10

<210> 30
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (10)...(10)
 <223> Xaa = MeGly

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 30
 Arg Cys Met Leu Gly Arg Val Tyr Arg Xaa Cys
 1 5 10

<210> 31
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 31
 Ala Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 32
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = Norleucine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<221> ACETYLATION

<222> (1)...(1)

<400> 32

Xaa	Cys	Met	Leu	Gly	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 33

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 33

Pro	Cys	Met	Leu	Gly	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 34

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 34

Asn	Cys	Met	Leu	Gly	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 35

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

WO 01/57070

PCT/US01/03293

<221> ACETYLATION
 <222> (1)...(1)

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 35
 Ser Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 36
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 36
 Glu Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 37
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> AMIDATION
 <222> (10)...(10)

<221> DISULFID
 <222> (1)...(10)

<400> 37
 Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 38
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

WO 01/57070

PCT/US01/03293

<221> DISULFID
<222> (1)...(10)

<221> AMIDATION
<222> (10)...(10)

<400> 38
Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
1 5 10

<210> 39
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> AMIDATION
<222> (11)...(11)

<221> DISULFID
<222> (2)...(11)

<400> 39
Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
1 5 10

<210> 40
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> MOD_RES
<222> (1)...(1)
<223> Xaa = Des-Amino-Arginine

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 40
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
1 5 10

<210> 41
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> MOD_RES
<222> (1)...(1)
<223> Xaa = D-Arginine

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<221> ACETYLATION
<222> (1)...(1)

<400> 41
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
1 5 10

<210> 42
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> MOD_RES
<222> (1)...(1)
<223> Xaa = D-Norleucine

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<221> ACETYLATION
<222> (1)...(1)

<400> 42
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
1 5 10

<210> 43
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> MOD_RES
<222> (1)...(1)
<223> Xaa = D-Proline

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<221> ACETYLATION
<222> (1)...(1)

<400> 43

WO 01/57070

PCT/US01/03293

Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 44
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = D-Asparagine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<221> ACETYLATION
 <222> (1)...(1)

<400> 44
 Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 45
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = D-Serine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<221> ACETYLATION
 <222> (1)...(1)

<400> 45
 Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 46
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> MOD_RES

WO 01/57070

PCT/US01/03293

<222> (1)...(1)
 <223> Xaa = D-Glutamic Acid

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<221> ACETYLATION
 <222> (1)...(1)

<400> 46
 Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 47
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> AMIDATION
 <222> (11)...(11)

<221> DISULFID
 <222> (2)...(11)

<221> ACETYLATION
 <222> (1)...(1)

<400> 47
 Arg Cys Met Leu Ala Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 48
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<221> DISULFID
 <222> (2)...(11)

<400> 48
 Arg Cys Met Leu Leu Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 49
 <211> 11
 <212> PRT
 <213> Artificial Sequence

WO 01/57070

PCT/US01/03293

<220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> MOD_RES
 <222> (5)...(5)
 <223> Xaa = Norleucine

 <221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 49
 Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
 1 5 10

 <210> 50
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> AMIDATION
 <222> (11)...(11)

 <221> DISULFID
 <222> (2)...(11)

 <400> 50
 Arg Cys Met Leu Pro Arg Val Tyr Arg Pro Cys
 1 5 10

 <210> 51
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> AMIDATION
 <222> (11)...(11)

 <221> DISULFID
 <222> (2)...(11)

 <400> 51
 Arg Cys Met Leu Asn Arg Val Tyr Arg Pro Cys
 1 5 10

WO 01/57070

PCT/US01/03293

<210> 52
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<221> DISULFID
 <222> (2)...(11)

<400> 52
 Arg Cys Met Leu Ser Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 53
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<221> DISULFID
 <222> (2)...(11)

<400> 53
 Arg Cys Met Leu Lys Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 54
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> AMIDATION
 <222> (11)...(11)

<221> DISULFID
 <222> (2)...(11)

<400> 54

WO 01/57070

PCT/US01/03293

Arg Cys Met Leu Glu Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 55
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (5)...(5)
 <223> Xaa = D-Leucine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 55
 Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 56
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (5)...(5)
 <223> Xaa = D-Norleucine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 56
 Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 57
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION

WO 01/57070

PCT/US01/03293

<222> (1)...(1)

<221> MOD_RES

<222> (5)...(5)

<223> Xaa = D-Proline

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 57

Arg	Cys	Met	Leu	Xaa	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 58

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> MOD_RES

<222> (5)...(5)

<223> Xaa = D-Glutamic Acid

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 58

Arg	Cys	Met	Leu	Xaa	Arg	Val	Tyr	Arg	Pro	Cys
1				5					10	

<210> 59

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> MOD_RES

<222> (5)...(5)

<223> Xaa = D-Lysine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

WO 01/57070

PCT/US01/03293

<400> 59
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
1 5 10

<210> 60
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> MOD_RES
<222> (5)...(5)
<223> Xaa = bAla

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 60
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
1 5 10

<210> 61
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> MOD_RES
<222> (5)...(5)
<223> Xaa = Gamma-Aminobutyric Acid

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 61
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
1 5 10

<210> 62
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> AMIDATION
<222> (11)...(11)

<221> DISULFID
<222> (2)...(11)

<400> 62
Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys
1 5 10

<210> 63
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> AMIDATION
<222> (11)...(11)

<221> DISULFID
<222> (2)...(11)

<400> 63
Arg Cys Met Leu Gly Arg Val Trp Arg Pro Cys
1 5 10

<210> 64
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> AMIDATION
<222> (11)...(11)

<221> DISULFID
<222> (2)...(11)

<400> 64
Arg Cys Met Leu Gly Arg Val His Arg Pro Cys
1 5 10

<210> 65
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

WO 01/57070

PCT/US01/03293

<221> ACETYLATION

<222> (1)...(1)

<221> MOD_RES

<222> (8)...(8)

<223> Xaa = (2')Naphthylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 65

Arg	Cys	Met	Leu	Gly	Arg	Val	Xaa	Arg	Pro	Cys
1				5					10	

<210> 66

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> MOD_RES

<222> (8)...(8)

<223> Xaa = P-Fluoro-Phenylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 66

Arg	Cys	Met	Leu	Gly	Arg	Val	Xaa	Arg	Pro	Cys
1				5					10	

<210> 67

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)

<221> MOD_RES

<222> (8)...(8)

<223> Xaa = P-Amino-Phenylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

WO 01/57070

PCT/US01/03293

<222> (11)...(11)

<400> 67

Arg	Cys	Met	Leu	Gly	Arg	Val	Xaa	Arg	Pro	Cys
1				5					10	

<210> 68

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLTATION

<222> (1)...(1)

<221> MOD_RES

<222> (8)...(8)

<223> Xaa = P-Carboxy-Phenylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 68

Arg	Cys	Met	Leu	Gly	Arg	Val	Xaa	Arg	Pro	Cys
1				5					10	

<210> 69

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLTATION

<222> (1)...(1)

<221> MOD_RES

<222> (8)...(8)

<223> Xaa - Cyclohexylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 69

Arg	Cys	Met	Leu	Gly	Arg	Val	Xaa	Arg	Pro	Cys
1				5					10	

<210> 70

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

WO 01/57070

PCT/US01/03293

<223> MCH Analog
 <221> ACETYLATION
 <222> (1)...(1)
 <221> MOD_RES
 <222> (4)...(5)
 <223> Xaa = 5-Aminopentanoic Acid
 <221> DISULFID
 <222> (2)...(11)
 <221> AMIDATION
 <222> (11)...(11)
 <400> 70
 Arg Cys Met Xaa Xaa Arg Val Tyr Arg Pro Cys
 1 5 10
 <210> 71
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> MCH Analog
 <221> ACETYLATION
 <222> (1)...(1)
 <221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = D-Arginine
 <221> MOD_RES
 <222> (4)...(5)
 <223> Xaa = 5-Aminopentanoic Acid
 <221> DISULFID
 <222> (2)...(11)
 <221> AMIDATION
 <222> (11)...(11)
 <400> 71
 Xaa Cys Met Xaa Xaa Arg Val Tyr Arg Pro Cys
 1 5 10
 <210> 72
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> MCH Analog
 <221> ACETYLATION
 <222> (1)...(1)
 <221> MOD_RES
 <222> (9)...(10)
 <223> Xaa = 5-Aminopentanoic Acid

WO 01/57070

PCT/US01/03293

<221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 72
 Arg Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys
 1 5 10

 <210> 73
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = D-Arginine

 <221> MOD_RES
 <222> (9)...(10)
 <223> Xaa = 5-Aminopentanoic Acid

 <221> DISULFID
 <222> (2)...(11)

 <221> AMIDATION
 <222> (11)...(11)

 <400> 73
 Xaa Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys
 1 5 10

 <210> 74
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> MCH Analog

 <221> ACETYLATION
 <222> (1)...(1)

 <221> MOD_RES
 <222> (5)...(5)
 <223> Xaa = D-Proline

 <221> MOD_RES
 <222> (9)...(10)
 <223> Xaa = 5-Aminopentanoic Acid

 <221> DISULFID
 <222> (2)...(11)

WO 01/57070

PCT/US01/03293

<221> AMIDATION
<222> (11)...(11)

<400> 74
Arg Cys Met Leu Xaa Arg Val Tyr Xaa Xaa Cys
1 5 10

<210> 75
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> MOD_RES
<222> (1)...(1)
<223> Xaa = Des-Amino-Arginine

<221> MOD_RES
<222> (9)...(10)
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 75
Xaa Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys
1 5 10

<210> 76
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> MCH Analog

<221> ACETYLATION
<222> (1)...(1)

<221> MOD_RES
<222> (4)...(5)
<223> Xaa = 5-Aminopentanoic Acid

<221> MOD_RES
<222> (9)...(10)
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID
<222> (2)...(11)

<221> AMIDATION
<222> (11)...(11)

<400> 76

WO 01/57070

PCT/US01/03293

Arg Cys Met Xaa Xaa Arg Val Tyr Xaa Xaa Cys
 1 5 10

<210> 77
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (3)...(3)
 <223> Xaa = Norleucine

<221> DISULFID
 <222> (2)...(11)

<221> AMIDATION
 <222> (11)...(11)

<400> 77
 Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys
 1 5 10

<210> 78
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCH Analog

<221> ACETYLATION
 <222> (1)...(1)

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = Des-Amino-Arginine

<221> MOD_RES
 <222> (5)...(5)
 <223> Xaa = D-Norleucine

<221> DISULFID
 <222> (2)...(11)


<221> AMIDATION
 <222> (11)...(11)

<400> 78
 Xaa Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys
 1 5 10

WHAT IS CLAIMED IS:

1. An optionally substituted peptide having the structure:

5



herein X¹ is an optionally present amino acid that, if present, is either
 10 alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X² is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine,
 15 serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X³ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine,
 20 aspartic acid or glutamic acid, or a derivative thereof;

X⁴ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof;

25 X⁵ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

X⁶ is an optionally present amino acid that, if present is either
 30 arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof,

X⁷ is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

35 X⁸ is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof;

X⁹ is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof;

X¹⁰ is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, asparagine, serine, sarcosine, isobutyric, or gamma-aminobutyric acid, or a derivative thereof;

X¹¹ is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof;

X¹² is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof;

X¹³ is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO₂, NH₂, F, I, and Br; or a derivative thereof;

X¹⁴ is either arginine, lysine, histidine, norarginine, or 5-aminopentanoic acid or a derivative thereof;

X¹⁵ is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof;

X¹⁶ is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

X¹⁷ is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

Z¹ is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z² is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;

or a labeled derivative of said peptide;

or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

2. The peptide of claim 1, wherein X¹, X², X³, X⁴, and X⁵ are not present; X⁶ is arginine; and X¹⁷ is tyrosine or tryptophan.

3. The peptide of claim 1, wherein X¹, X², X³, X⁴, X⁵, and X¹⁷ are not present; and X⁶ is arginine.
- 5 4. The peptide of claim 3, wherein Z¹ is -C(O)CH₃ and Z² is -NH₂.
5. The peptide of claim 1, wherein said peptide is either SEQ. ID. NO. 7, SEQ. ID. NO. 8, SEQ. ID. NO. 9, SEQ. ID. NO. 10, or
10 a pharmaceutically acceptable salt thereof.
6. The peptide of claim 1, wherein said peptide is either SEQ. ID. NO. 7, SEQ. ID. NO. 8, or a pharmaceutically acceptable salt thereof.
- 15 7. The peptide of claim 1, wherein X¹, X², X³, X⁴, X⁵ and X¹⁷ are not present;
X⁶ is either arginine, D-arginine, D-norleucine, D-proline, D-serine, or D-asparagine;
X⁷ is cysteine;
X⁸ is either methionine, norleucine, or N-methyl norleucine;
20 X⁹ is leucine;
X¹⁰ is either glycine, alanine, leucine, norleucine, asparagine, serine, D-norleucine, D-proline, gamma-aminobutyric acid, or sarcosine;
X¹¹ is arginine;
X¹² is valine;
25 X¹³ is phenylalanine, (2')naphthylalanine, p-fluoro-phenylalanine, tyrosine, or cyclohexylalanine;
X¹⁴ is arginine;
X¹⁵ is either proline or sarcosine; and
X¹⁶ is either cysteine or D-cysteine.
30
8. The peptide of claim 1, wherein said peptide consists of a sequence selected from the group consisting of: 7, 8, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 57, 61, 62, 63, 65-67, 69-72, and 77.

9. A method of screening for a compound able to bind a MCH receptor comprising the step of measuring the ability of said compound to effect binding of the peptide of any one of claims 1-8 to either said receptor, a fragment of said receptor comprising a MCH binding site, a polypeptide comprising said
5 fragment, or a derivative of said polypeptide.

10. The method of claim 9, wherein said method measures the ability of said peptide to bind to said receptor or said fragment thereof.

10 11. The method of claim 10, wherein said peptide is radiolabeled.

12. The method of claim 9, wherein said peptide is a radiolabeled derivative of SEQ. ID. NO. 8 or a pharmaceutically acceptable salt thereof.

15 13. A method for increasing weight in a subject comprising the step of administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce a weight increase.

20 14. A method for increasing appetite in a subject comprising the step of administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce an appetite increase.

15. A method for measuring the ability of a compound to decrease weight or appetite in a subject comprising the steps of:
25 a) administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce a weight increase or appetite increase,
b) administering said compound to said subject, and
c) measuring the change in weight or appetite of said subject.

30 16. The method of claim 15, wherein said subject is either a rat or a mouse.

ALANINE SCAN OF HUMAN MELANIN-CONCENTRATING HORMONE (hMCH)

1 3 5 * 9 11 13 15 * 17 19
Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys Trp Gln Val ACID

BINDING IC₅₀ (nM) 0.8 0.2 0.49 0.49 0.33 4.5 - 71 0.9 - 169 0.9 253 1.4 1.9 - 0.2 1.1 0.3

FUNCTION EC₅₀ (nM) ND ND ND ND ND ND - ND ND ND ND ND ND ND - ND ND ND

% ACTIVATION (100 nM) 97.7 96.3 97.7 95.7 90.8 11.4 - 0 86.5 - 3 82.9 0 90.4 76.3 - 100 78.3 94.4

hMCH BINDING ASSAY⁺, IC₅₀ = 0.3nM
 AEQUORIN FUNCTIONAL ASSAY⁺⁺, EC₅₀ = 36nM, 100% ACTIVATION AT 10μM

+ SPA ASSAY: INHIBITION OF (¹²⁵ I-F13, Y19-hMCH BINDING TO THE CLONED HUMAN MCH RECEPTOR (COS-7 CELLS)
++ FOR AEQUORIN FUNCTIONAL ASSAY, 100% ACTIVATION IS THE BIOLUMINESCENCE VALUE OBTAINED WITH 100 nM MCH.
* SITES OF CYCLIZATION (S-S)
ND-NOT DONE

FIG.1