

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 February 2006 (16.02.2006)

PCT

(10) International Publication Number
WO 2006/015445 A1

(51) International Patent Classification⁷: **A61K 48/00**,
A61P 1/04, C12N 1/21

(21) International Application Number:
PCT/AU2005/001211

(22) International Filing Date: 12 August 2005 (12.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2004904564 13 August 2004 (13.08.2004) AU
60/602,859 20 August 2004 (20.08.2004) US

(71) Applicant and

(72) Inventor: MARSHALL, Barry, J. [AU/AU]; 105 Hay
Street, SUBIACO, WA 6008 (AU).

(74) Agent: GRIFFITH HACK; Level 6, 256 Adelaide Ter-
race, PERTH, WA 6000 (AU).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ,
OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL,
SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: BACTERIAL DELIVERY SYSTEM



(57) Abstract: Helicobacter based preparations comprising a pharmacologically active molecule of interest are disclosed, as well as methods of preparing and using said preparations. In particular, Helicobacter pylori vectors, vector plasmids and recombinant cells that include a sequence encoding a pharmacologically active molecule of interest useful in therapeutic treatments and/or vaccination against disease are provided. Delivery of the pharmacologically active molecules is provided at the mucosal surface, such as the gastric mucosa or nasal membranes, to provide effective and continuous delivery of a pharmacologically active agent. Vectors and shuttle vector constructs are also provided.

WO 2006/015445 A1

BACTERIAL DELIVERY SYSTEMFIELD OF THE INVENTION

5 The present invention relates to the delivery of
biologically active agents *in vivo*. In particular, the
present invention relates to treating, palliating or
preventing diseases using a bacterial delivery system for
delivering biologically active agents directly to an
10 anatomical site *in vivo*. In one embodiment, the bacterial
delivery system comprises *Helicobacter* or bacterium
exhibiting *Helicobacter* features for delivering
heterologous nucleic acid into an animal or animal cell,
wherein the heterologous nucleic acid is expressed. In a
15 further embodiment the *Helicobacter* has been engineered to
contain a DNA vector that encodes heterologous nucleic
acid, wherein upon infection the nucleic acid vector is
expressed such that the biologically active agent is
delivered to the animal body, especially at the mucosa.

20

BACKGROUND TO THE INVENTION

There is a continuing need for long-term delivery of
pharmacologic and immunologic agents to individuals with
25 both congenital and acquired diseases. In most cases, this
involves repeated administration of therapeutic compounds
several times per day, daily or at intervals. However, it
is appreciated that any long-term therapeutic or preventive
regime has problems such as compliance, side effects and
30 drug resistance. Consequently, there is a constant need to
identify treatments, which will be usually 100% effective,
free from side effects and cheap.

One form of delivery that has been investigated in recent
35 times for various therapeutic and prophylactic agents has
been the use of microorganisms. Genes of interest from
various organisms including bacteria, viruses, parasites as

- 2 -

well as mammals have, for example, been cloned into a variety of bacteria, viruses and mycobacteria for the purpose of directing these micro-organisms to express foreign protein or impart certain desired properties. These
5 microorganisms have been used in vaccination programs, gene-replacement therapies and therapeutic composition delivery.

Microorganisms have also been used to transform animal
10 (host) cells *in vivo*. Host cell transformation can be accomplished using gene-delivery vectors comprising replication incompetent viruses (see for example U.S. Pat. No. 5,824,544), naked DNA, (see for example U.S. Pat. No. 6,261,834), liposomes containing recombinant expression
15 cassettes (see for example U.S. Pat. No. 6,271,207). Other molecular-based therapeutic composition delivery approaches include using replication incompetent recombinant viruses designed to express heterologous surface proteins (see, for example, U.S. Pat. No. 6,376,236).

20 In recent times, pharmaceutical researchers have also attempted to develop methods for *in vivo* therapeutic composition expression using recombinant organism-based vectors, inanimate vectors and naked DNA. Examples of
25 recombinant organism-based vectors include recombinant bacteria (see for example U.S. Pat. No. 5,547,664) and viruses such as alphaviruses (see for example U.S. Pat. No. 6,391,632), vaccinia viruses (see for example U.S. Pat. No. 6,267,965), adenoviruses (see for example U.S. Pat. No.
30 5,698,202) and adenovirus associated virus (AAV) (see for example U.S. Pat. No. 6,171,597). Inanimate vectors include lipidic gene delivery vector constructs such as DNA/cationic liposome complexes, DNA encapsulated in neutral or anionic liposomes, and liposome-entrapped,
35 polycation-condensed DNA (LPDI and LPDII). (see Ropert, 1999, *Braz J Med Biol Res*, 32(2):163-9).

- 3 -

Examples of other genes that have been delivered by various bacteria include cloning the invasion genes of *Shigella* into the normally non-invasive *E. coli* rendering the *E. coli* invasive and therefore more suitable for use as a vaccine strain, or cloning of *Plasmodium falciparum* malaria genes into *Salmonella typhimurium* which subsequently express these malaria proteins and, following oral administration of the bacteria, induce specific cytotoxic T cell immunity and protection in mice against malaria challenge (see, for example, Hone et al., 1991, *Vaccine*, 9:810-816; Tacket et al., 1992, *Infect. Immun.*, 60:536-541; Hone et al., 1992, *J. Clin. Invest.*, 90:412-420; Chatfield et al., 1992, *Vaccine*, 10:8-11; Tacket et al., 1992, *Vaccine*, 10:443-446; and Mims et al., 1993, *In: Medical Microbiology*, Eds., Mosby-Year Book Europe Ltd., London; Sadoff et al., 1988, *Science*, 240:336-338; Aggrawal et al., 1990, *J. Exp. Med.*, 172:1083-1090).

Attenuated or less virulent *Shigella* (see, for example, Noriega et al., 1994, *Infect. Immun.*, 62:5168-5172; US Pat. Appl. No. 20020176848), *Salmonella* (see, for example, US Pat. No. 6,531,313; US Pat. Appl. No. 20030170211), *Listeria* (see, for example, Schafer et al., 1992, *J. Immunol.*, 149:53-59; US Pat. Appl. No. 20030008839), and other bacteria have been given orally to immunise against subsequent infection with more virulent forms of these bacteria. Likewise, attenuated bacterial and mycobacterial organisms such as *Bacille Calmette-Guerin* (BCG) (Lagranderie et al., 1993, *Vaccine*, 11:1283-1290; Flynn, 1994, *Cell. Molec. Biol.*, 40(Suppl. 1):31-36) have been administered parenterally to protect against related organisms such as *M. tuberculosis*.

Some of the other bacterial species that have been used as vectors include *Yersinia enterocolitica* (van Damme et al., 1992, *Gastroenterol.*, 103:520-531) and *Vibrio cholerae* (Levine et al., 1994, *In: Vibrio cholerae*, Molecular to

- 4 -

Global Perspective's, Wachsmuth et al., Eds, ASM Press, Washington, D.C., pages 395-414).

Despite all of the research all of the above bacterial
5 delivery systems have technical difficulties, which need to
be overcome before these can be used *in vivo*. Indeed, most
of the bacterial systems require the bacteria themselves to
produce functional molecules and are dependent on a
bacterium which is sufficiently attenuated to be safe for
10 use in humans, but still able to produce biologically
active agents. However, all of the attenuated strains of
bacteria used previously are not capable of surviving
lengthy periods *in vivo*, without causing side effects. More
importantly, many of these bacterial species are not
15 capable of delivering therapeutic or prophylactic agents to
the mucosal epithelial cells of an animal. However, many
of the most important therapeutic agents are taken up via
the mucosa; therefore there is a need for a bacterial
delivery system capable of delivering biologically active
20 agents directly to the mucosa.

U.S. Pat. No. 5,877,159 to Powell et al., describes live
bacteria that can invade mucosal cells without establishing
a productive infection or causing disease to thereby
25 introduce a eukaryotic expression cassette encoding an
antigen capable of being expressed by the animal cells.
While this method allows delivery of the DNA vaccine to
mucosal surfaces, including easy administration, a concern
for vaccine delivery in developing countries, it does not
30 have the advantage of providing amplifiable mRNA encoding
the gene of interest. Moreover, the bacterium disclosed in
Powell et al. is not capable of sustained delivery.

Non-pathogenic, non-colonising, non-invasive food-grade
35 bacterium *Lactococcus lactis* has been used previously in
the past to deliver agents to the mucosa (see, for example,
UK patent GB-2278358B). However, while *Lactococcus lactis*

- 5 -

is non-invasive it is not capable of establishing a chronic infection, which is capable of delivering continuous therapeutic agents directly into mucosal epithelial cells.

5 Consequently, there is a continuing need for a delivery system, which is capable of establishing a non-invasive, chronic infection in close proximity to the mucosal epithelial cells of an animal such that biologically active agents can be delivered thereto. Moreover, there is a
10 continuing need for improved delivery mechanisms for pharmacologically active molecules at the mucosal surface sufficient to elicit a useful and beneficial immunogenic response. Such would provide an effective *in vivo* delivery system for pharmacological active agents, as well as an
15 effective method for immunization, i.e., antigen exposure at mucosal surface sufficient to elicit a general humoral and mucosal immune response.

SUMMARY OF THE INVENTION

20

The present invention is directed to overcoming the above-mentioned challenges and is exemplified in a number of implementations and applications, some of which are summarized below.

25

The inventors have now surprisingly found that *Helicobacter* and in particular *Helicobacter pylori*, which is capable of forming a chronic infection, can be used to produce continuous drug delivery. Moreover, *Helicobacter* of the
30 present invention can be activated and inactivated at various times and because of its chronicity, the *Helicobacter* can be used to deliver drugs through the gastric mucosa.

35 *Helicobacter pylori* is a gram-negative spiral shaped bacterium found almost exclusively in the human gastric mucosa. The acidity of the human stomach is an effective

- 6 -

barrier to colonization by essentially all bacteria, with the exception of *Helicobacter* species.

5 *H. pylori* has the unique ability to colonize and persist for decades within the human gastric mucosa, despite development of a mucosal inflammatory and immune response. This characteristic renders *H. pylori* an interesting candidate for the delivery of selected agents through the mucosa.

10

In accordance with some aspects, compositions, methods and systems are provided for preparing and using a *Helicobacter*-based construct comprising a *Helicobacter* sequence having a promoter region and a non-*Helicobacter* sequence encoding a non-*Helicobacter* pharmacologically active molecule. This construct in some embodiments is described as a vector or a plasmid vector, wherein the promoter sequence is capable of controlling the expression of the non-*Helicobacter* pharmacologically active molecule of interest.

20

Accordingly, in a first aspect the present invention provides a method of delivering one or more biologically active agents to a subject comprising the step of administering to a subject an effective amount, preferably a therapeutically or prophylactically effective amount of, *Helicobacter* cells or bacterial cells having the features of *Helicobacter*, which cells express said one or more biologically active agents.

30

In a second aspect, the present invention provides a non-invasive or non-pathogenic *Helicobacter* cell expressing one or more heterologous biologically active agents.

35 Preferably, the *Helicobacter* is of the species *H. pylori*.

The biologically active agent can be either homologous to

- 7 -

the *Helicobacter* genus or heterologous to cells of the genus or species of the *Helicobacter* cells used for delivery. Heterologous agents may be derived from either eukaryotic or prokaryotic sources.

5

In some aspects, the *Helicobacter*-based vector and vector plasmid constructs comprise biologically active agents such as an antigen, organic or inorganic molecule or substance, a pharmacological agent eg a therapeutic agent or
10 prophylactic agent, such as a gene product or gene sequence (isolated nucleic acid). By way of example, such agents may include an immunoregulatory agent, hormone, ligand, an enzyme or an anti-sense RNA, a catalytic RNA, a protein, a peptide or any other molecule which can be present on or
15 released from the bacterial cells so that it may be delivered to an animal or to an animal cell.

In some embodiments, the biologically active agent is encoded by a nucleic acid molecule, which is preferably
20 obtained in isolated form, and subsequently inserted into the bacterial delivery vehicle of this invention. It will be appreciated by those skilled in the art that the isolated nucleic acid molecule of the present invention may be cDNA, genomic DNA, RNA, or a hybrid molecule thereof.
25 Preferably, the nucleic acid is cDNA.

By way of example, a protein and/or peptide of interest may comprise ghrelin, amylin, insulin, motilin, β -glucosidase, a chemical chaperone, or other molecule useful in the
30 treatment of Gauchers disease, cell wasting, human immunodeficiency disease (AIDS), appetite suppression, preparations useful in the treatment of diabetes, etc.

The isolated nucleic acid is preferably incorporated into
35 an expression vector, which can be transformed into and maintained in the *Helicobacter* cells.

- 8 -

Accordingly, in a third aspect, the present invention provides a recombinant vector for delivering biologically active agents directly to an anatomical site in need thereof. The *Helicobacter* cell of the present invention has
5 preferably been adapted to secrete and/or express on its surface the biologically active agents.

In a fourth aspect, the present invention provides a vector or construct for delivering *in vivo* an effective amount of
10 a biologically active agent, comprising:

- a) a nucleotide sequence encoding a biologically active agent;
- b) operatively linked thereto, a control or regulatory sequence capable of controlling the expression
15 of the nucleotide sequence of (a) such that the biologically active agent is produced *in vivo* when the delivery vehicle comprising the vector is delivered to a subject.

20 The vector maybe modified chemically by means of chemical or enzymatic treatment, or *in vivo* by means of recombinant DNA technology to produce a modified or variant vector.

Such a construct may differ from those disclosed, for
25 example, by virtue of one or more nucleotide substitutions, deletions or insertions, but substantially retain a desired biological activity of the construct or nucleic acid molecule, or its encoded product, in accordance with this invention.

30 In another embodiment of the present invention, the *Helicobacter* vector or construct is provided with a reporter gene(s) expressed from a constitutive promoter cloned into the expression vector and used as a screening
35 tool. Non-limiting examples of reporter genes suitable for use herein include green fluorescent protein (GFP), β -galactosidase, amylase, and chloramphenicol acetyl

transferase (CAT).

In another embodiment of the present invention the *Helicobacter* cell is used to deliver a heterologous gene of interest to a subject in need thereof. The gene of interest may encode a therapeutic product (a transgene product), including, but not limited to a peptide hormone (such as, but not limited to α -melanocyte-stimulating hormone (α -MSH), insulin, growth hormone, and parathyroid hormone), a cytokine including, but not limited to an interferon, interleukin (IL)-2, IL-4, IL-10, IL-12, granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF) and erythropoietin (EPO).

In still another embodiment, the present invention provides methods for treating, palliating or preventing a disease in a subject. These methods are facilitated with the use of a therapeutic or prophylactic composition. Accordingly, in a fifth aspect, the present invention provides a pharmaceutical composition comprising (a) live non-pathogenic *Helicobacter* cells expressing and/or secreting a biologically active agent, together with (b) a therapeutically effective carrier.

In a sixth aspect the present invention provides a method of treating, preventing or palliating a disease comprising administering to a subject in need thereof an effective amount of a *Helicobacter* cells of the present invention that express the biologically active agent.

Non-limiting examples of subjects in which the present invention may be used, include mammals such as primates, equines, bovines, porcines, ovines and rodents. Also intended are fish and birds.

In one embodiment of the present invention the disease being treated, prevented or palliated is cancer, a disease

- 10 -

or condition of the immune/hematopoietic system, a disease or condition of the reproductive system, a disease or condition of the musculoskeletal system, a disease or condition of the cardiovascular system, a disease or a
5 condition described as mixed fetal, a disease or a condition of the excretory system, a disease or a condition of the neural/sensory system, a disease or a condition of the endocrine system, a disease or condition of the respiratory system, a disease or condition of the digestive
10 system and a disease or condition associated with connective/epithelial tissue or disease or conditions caused by bacterial, viral or parasitic infection.

The present invention provides a variety of
15 pharmaceutically acceptable preparations formulated for delivery to a patient, such as, gastrically, orally, or intranasally. In particular embodiments, the composition are suitable for delivery at a mucosal surface. In particular embodiments, the composition is suitable for
20 delivery to the mucosal surface of the gut.

By way of example, the mucosa may be that of the gastric, vaginal, nasal, oral, or ocular surface, or any other of the body characterized by the presence of a penetrable
25 mucosal surface or lining. In some embodiments, the mucosal surface is the gastric mucosal surface.

The various delivery forms of the compositions are readily prepared for use in the practice of the present invention
30 given the specific types and ratios of specific *Helicobacter*, *Helicobacter* constructs and other delivery vehicles described herein, and those formulation techniques known to those in the formulary arts, such as are described in Remington's Pharmaceutical Sciences, 20th edition, Mack
35 Publishing Company, which text is specifically incorporated herein by reference.

- 11 -

It is envisioned that the delivery system may be employed in animals, particularly primates, including humans, equines, bovines, ovines, and rodents, fish and birds. It is also anticipated that the preparations may be used on
5 both infants and adults, as well as parentally or for administration to pregnant or lactating animals. The preparations and methods may be further described as suitable for both male and female animals.

10 In yet another aspect, a method is provided for vaccinating an animal. In some embodiments, the method comprises administering a composition comprising a vaccine comprising cells transformed with the *Helicobacter*- based plasmid vector and/or plasmid vectors as described herein. In
15 other embodiments, the method provides for the delivery of an effective amount of the pharmacologically active molecule of interest sufficient to eliminate or inhibit a disease or physiological condition in the animal, or sufficient to elicit an immune response specific for the
20 pharmacologically active molecule of interest.

By way of example, the non-*Helicobacter* pharmacologically active molecule of interest useful in the vaccine may comprise a mammalian protein, peptide, enzyme, hormone, or
25 any combination of these. In particular embodiments, the pharmacologically active molecule of interest is further defined as a human pharmacologically active molecule of interest. In some embodiments the pharmacologically active molecule of interest is a human pathogen molecule/antigen,
30 human protein antigen, such as amylin or an analog or derivative thereof, or ghrelin, or an analog or derivative thereof.

In particular embodiments, the vaccine conveys immunity
35 against the human pathogen, Ebola virus, HIV virus, Marburg virus, influenza virus, and the like. Replication competent vaccines based on attenuated recombinant vesicular

- 12 -

stomatitis virus vectors have been described by Jones *et al.* (2005, *Nat. Med.* 11: 786-90) that include Ebola glycoprotein and Marburg glycoprotein. Hence, it is envisioned that constructs using the *Helicobacter*-based
5 vector systems and plasmid vector systems with these and other glycoprotein's associated with human pathogens may also be provided according to the present invention together with the disclosure provided herein.

10 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a schematic diagram of the plasmid construct pHPA1 (2.8kb).

15 Figure 2 is a schematic diagram of the plasmid construct pHP3 (3.4kb).

Figure 3 is a schematic diagram of the plasmid vector pTMI03-8.

20

Figure 4 shows the structure of sulfasalazine (SSN).

Figure 5 is a schematic diagram showing the use of ion exchange resin (Amberlite XE-96) conjugated with a dye
25 (Azure-A).

BRIEF DESCRIPTION OF THE SEQUENCES

The following nucleic acid and amino acid sequences are
30 referenced throughout the description of the present invention:

SEQ ID NO: 1 - Nucleotide sequence of plasmid pHP1 (2796 nucleotides);

35

SEQ ID NO: 2 - Nucleotide sequence of pHP1 (2796 nucleotides);

- 13 -

SEQ ID NO: 3 - Nucleotide sequence of plasmid pHP3 (3444 nucleotides);

- 5 SEQ ID NO: 4 - Hepatitis C virus core antigen (HCV) nucleotide Sequence;

- SEQ ID NO: 5 - Nucleotide sequence 135 bp (45 amino acids) immunogenic coding sequence from the Hepatitis C virus
10 (HCV) core antigen;

- SEQ ID NO: 6 - Nucleotide sequence of the surface exposed loop of the *HopE* gene (at nt504, aa position 168) of *H. pylori*;
15

SEQ ID NO: 7 - Upstream primer (29 nucleotides);

SEQ ID NO: 8 - Downstream Primer (28 nucleotides);

- 20 SEQ ID NO: 9 - Oligonucleotide Primer (15 nucleotides).

DEFINITION OF TERMS

- Prior to setting forth the invention, it may be helpful to
25 an understanding thereof to set forth definitions of certain terms that will be used hereinafter.

- An "antibiotic resistance gene" as defined herein includes heterologous nucleic acid sequences purposely provided to a
30 vector and used as a selection system. The term "antibiotic resistance gene" does not include other mechanisms or genes that impart antibiotic resistance to naturally occurring micro-flora organisms.

- 35 The term "attenuated" as used herein for example to describe a bacterial strain, particularly an *E. coli* or a *Helicobacter* strain such as *Helicobacter pylori*, is defined

- 14 -

as a strain that is less virulent and/or toxic (invasive) that a native, wild type bacterial strain.

The term "biologically active" as used herein refers to ability to perform a biological function and with reference to a polypeptide implies that the polypeptide adopts a stable conformation ("folded form") which is the same or closely analogous to its native conformation. When folded correctly or substantially correctly, for example with formation of proper folded units α -helices, β -sheets, domains, disulphide bridges etc., a polypeptide should have the ability to perform its natural function. Generally, the unit of function in a polypeptide is a domain.

Mere ability to be bound by an antibody or other receptor, either with or without elicitation of an immune response, is passive or does not constitute "biological activity". Any antigen has the ability to be bound by an antibody but is not necessarily biologically active.

"Clinical grade vector" as used herein means a plasmid or other expression vector that is capable of being expressed in *Helicobacter* or a non-pathogenic bacterium engineered to have features of *Helicobacter*. The clinical grade vectors of the present invention do not use antibiotic resistance markers for selection and/or have been modified to prevent replication outside the host e.g. such as a suicide vector.

An example of suicide system in *H. pylori* has been described by Panthel et al. 2003 (*Infection & Immunity*, 71: 109-116). This system introduces a plasmid into *H. pylori* which contains the PhiX174 lysis gene E. To eradicate the strain, incubation at 42°C for 5 hours was used. *In vivo* this would mean that the animal would consume a drink at 45-50°C to raise the temperature of the gastric environment above 42°C.

- 15 -

A second example is the *L-Dap* selection system, commonly used to allow survival of bacterial mutants on supplemented plates (see, for example, Kirata *et al.* 1997 (*Infection & Immunity*, 65: 4158-4164). In this system the animal subject
5 must supplement their diet with a missing substrate i.e. diamino-pimelic-acid (DAP), in order for the *DapE* deficient *H. pylori* mutant to survive. In order to eradicate the then DAP consumption is ceased.

10 A third possible system relates to metronidazole sensitivity of *H. pylori* because of its *rdxA* gene. Excessive replication of the *rdxA* gene is harmful to mammalian cells and *E. coli*. However, duplication may be tolerated by the bacterium. Therefore a *Helicobacter*
15 species of the present invention can be engineered to contain two copies of *rdxA* which prevent the normal mutation-dependant *rdxA* loss. The introduction of at least two functional *rdxA* genes into the *Helicobacter* genome should result in a *Helicobacter* strain, which is
20 permanently sensitive to metronidazole. Jeong *et al.* 2000 (*J. Bacteriol.*, 182: 5082-5090) showed that the nitroreductase produced by a functional *rdxA* gene converts metronidazole from a prodrug to a bactericidal compound. The mode of action of the active compound is to cause DNA
25 breaks of the *Helicobacter* genome.

"Detectable immune response" as used herein is either an antibody (humoral) or cytotoxic (cellular) response formed in an animal in response to an antigen that can be measured
30 using routine laboratory methods including, but not limited to enzyme-linked immunosorbant assays (ELISA), radio-immune assays (RIA), Enzyme-linked ImmunoSPOT (ELISPOT), immunofluorescent assays (IFA), complement fixation assays (CF), Western Blot (WB) or an equivalent thereto.

35

"Gene of interest" as used herein refers to any nucleic acid sequence encoding for a polypeptide or protein whose

- 16 -

expression is desired. The gene of interest may or may not include the promoter or other regulatory components. The gene of interest also includes constructs capable of producing anti-sense RNA.

5

"Gene therapy" as used herein is defined as the delivery of a gene of interest to an animal in need thereof using a recombinant vector. The gene of interest can be a transgene encoding for a therapeutic or prophylactic protein or polypeptide including, but not limited to cytokines, anti-inflammatory, anti-proliferatives, antibiotics, metabolic inhibitors/activators and immunologically active antigens and fragments thereof. Furthermore, "gene therapy" as used herein also includes gene replacement technologies directed at both inherited and non-inherited disorders.

The term *Helicobacter* includes all bacteria of the genus *Helicobacter* including *H. pylori* and *H. mustelae*. The term also includes bacteria that have similar biology to *H. pylori* in that they are capable of residing on the gastric mucosa of primates and/or capable of establishing a chronic, but isolated infection of the mucosa. The term also encompasses bacteria that have been modified so that the bacterium has *H. pylori* features such as the ability to reside on the gastric mucosa.

A "heterologous" polypeptide is one not native to *Helicobacter*, i.e., not expressed by *Helicobacter* in nature or prior to introduction into *Helicobacter*, or an ancestor thereof, of encoding nucleic acid for the biologically active agent.

"Host" as used herein defines the intended recipient of a therapeutic composition of the present invention. Host includes all animals. Specifically, hosts include, but are not limited to, primates (including man), bovine, equine, canine, feline, porcine, ovine, rabbits, rodents, birds and

- 17 -

fish.

"Immunologically inert" as used herein shall mean any substance, including microorganisms such as microflora that does not provoke a significant immune response in its host. Examples of immunologically inert materials as used herein include stainless steel, biocompatible polymers such as poly-L-lactide, medical grade plastics and the microflora organisms of the present invention. A "significant immune" response is any immune response that would limit or restrict the *in vivo* utility of a material or organism used in accordance with the teachings of the present invention. A detectable immune response is not necessarily a "significant immune response."

An "isolated nucleic acid" is a nucleic acid the structure of which is not identical to that of any naturally occurring nucleic acid or to that of any fragment of a naturally occurring genomic nucleic acid spanning more than three separate genes. The term therefore covers, for example, (a) a DNA molecule which has the sequence of part of a naturally occurring genomic DNA molecule but is not flanked by both of the coding sequences that flank that part of the molecule in the genome of the organism in which it naturally occurs; (b) a nucleic acid incorporated into a vector or into the genomic DNA of a prokaryote or eukaryote in a manner such that the resulting molecule is not identical to any naturally occurring vector or genomic DNA; (c) a separate molecule such as a cDNA, a genomic fragment, a fragment produced by polymerase chain reaction (PCR), or a restriction fragment; and (d) a recombinant nucleotide sequence that is part of a hybrid gene, i.e., a gene encoding a fusion protein. Specifically excluded from this definition are nucleic acids present in mixtures of (i) DNA molecules, (ii) transfected cells, and (iii) cell clones, e.g., as these occur in a DNA library such as a cDNA or genomic DNA library.

"Percent identity (homology)" of two amino acid sequences or of two nucleic acids is determined using the algorithm of Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA*, 87:2264-2268, 1990, modified as in Karlin and Altschul (*Proc. Natl. Acad. Sci. USA* 90:5873-5877, 1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (*J. Mol. Biol.* 215:403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, score=100, wordlength=12, to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches are performed with the XBLAST program, score=50, wordlength=3, to obtain amino acid sequences homologous to a reference polypeptide (eg., SEQ ID NO: 2). To obtain gapped alignments for comparison purposes, Gapped BLAST is utilised as described in Altschul et al. (*Nucleic Acids Res.* 25:3389-3402, 1997). When utilising BLAST and Gapped BLAST programs, the default parameters of the respective programs (eg., XBLAST and NBLAST) are used. These may be found on the World Wide Web at the URL "ncbi.nlm.nih.gov."

The term "reporter gene" as used herein is a nucleic acid sequence incorporated into (or adjacent to) the heterologous nucleic acid encoding for the gene of interest that provides the transformed vector expressing the gene of interest an identifiable phenotype. Non-limiting examples of reporter genes include GFP, β -galactosidase, amylase, and CAT.

"Screening marker" as used herein refers to an identifying characteristic (phenotype) provided to a transformed vector made in accordance with the teachings of the present invention. In one embodiment of the present invention the screening marker is a reporter gene.

"Selectable marker," "selectable gene," "reporter gene" and

- "reporter marker" (referred to hereinafter as a "selectable marker") as used herein refer to nucleic acid sequences encoding for phenotypic traits that permit the rapid identification and isolation of a transformed bacterial
- 5 vector. Generally, bacterial vectors deemed "clinical grade" and made in accordance with the teachings of the present invention are those vectors having selectable markers that do not encode for antibiotic resistance.
- 10 "Transgene" as used herein refers to a gene that is inserted, using cDNA technology, into a cell in a manner that ensures its function, replication and transmission as a normal gene.
- 15 "Transforming nucleic acid sequence" as used herein means a plasmid, or other expression cassette containing a nucleic acid sequence encoding a gene of interest. In some embodiments of the present invention the nucleic acid sequence can encode for one or more therapeutic agents.
- 20 "Transforming nucleic acid sequence" can also be used to mean a "transgene" in accordance with certain embodiments of the present invention. In another embodiment of the present invention the transforming nucleic acid sequence includes nucleic acid sequence encoding for a promoter
- 25 and/or other regulatory elements.
- The term "cancer" as used herein refers to neoplastic diseases (e.g., leukemia, cancers and "hyperproliferative disorders"). The neoplasm may be located in a tissue
- 30 selected from the group consisting of: colon, abdomen, bone, breast, digestive system, liver, pancreas, prostate, peritoneum, lung, blood (e.g., leukemia), endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), uterus, eye, head and neck, nervous (central and
- 35 peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

- 20 -

In one embodiment the term "cancer" also encompasses pre-neoplastic conditions selected from the group consisting of hyperplasia (e.g., endometrial hyperplasia), metaplasia (e.g., connective tissue metaplasia and/or dysplasia (e.g.,
5 cervical dysplasia, and bronchopulmonary dysplasia).

In another embodiment, the term "cancer" also encompasses benign dysproliferative disorder selected from the group consisting of: benign tumors, fibrocystic conditions, and
10 tissue hypertrophy.

The term "a disease or condition of the immune/haematopoietic system" as used herein refers to a disease or condition selected from the group consisting of:
15 anaemia, pancytopenia, leukopenia, thrombocytopenia, leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic anaemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, asthma, AIDS, autoimmune disease, rheumatoid arthritis, granulomatous disease,
20 immune deficiency, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, systemic lupus erythematosus, haemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, Celiac
25 disease and allergies.

The term "a disease or condition of the reproductive system" as used herein refers to a disease or condition selected from the group consisting of: cryptorchism,
30 prostatitis, inguinal hernia, varicocele, leydig cell tumours, verrucous carcinoma, prostatitis, malacoplakia, Peyronie's disease, penile carcinoma, squamous cell hyperplasia, dysmenorrhea, ovarian adenocarcinoma, Turner's syndrome, mucopurulent cervicitis, Sertoli-leydig tumours,
35 ovarian cancer, uterine cancer, pelvic inflammatory disease, testicular cancer, prostate cancer, Klinefelter's syndrome, Young's syndrome, premature ejaculation, diabetes

mellitus, cystic fibrosis, Kartagener's syndrome,
testicular atrophy, testicular feminization, anorchia,
ectopic testis, epididymitis, orchitis, gonorrhoea,
syphilis, testicular torsion, vasitis nodosa, germ cell
5 tumours, stromal tumours, dysmenorrhea, retroverted uterus,
endometriosis, fibroids, adenomyosis, anovulatory bleeding,
amenorrhoea, Cushing's syndrome, hydatidiform moles,
Asherman's syndrome, premature menopause, precocious
puberty, uterine polyps, dysfunctional uterine bleeding,
10 cervicitis, chronic cervicitis, mucopurulent cervicitis,
cervical dysplasia, cervical polyps, Nabothian cysts,
cervical erosion, cervical incompetence, cervical
neoplasms, pseudohermaphroditism, and premenstrual
syndrome.

15

The term "a disease or condition of the musculoskeletal
system" as used herein refers to a disease or condition
selected from the group consisting of: bone cancers (e.g.,
osteochondromas, benign chondromas, chondroblastoma,
20 chondromyxoid fibromas, osteoid osteomas, giant cell
tumours, multiple myeloma, osteosarcomas), Paget's Disease,
rheumatoid arthritis, systemic lupus erythematosus,
osteomyelitis, Lyme Disease, gout, bursitis, tendonitis,
osteoporosis, osteoarthritis, muscular dystrophy,
25 mitochondrial myopathy, cachexia, and multiple sclerosis.

The term "a disease or condition of the cardiovascular
system" as used herein refers to a disease or condition
selected from the group consisting of: myxomas, fibromas,
30 rhabdomyomas, cardiovascular abnormalities (e.g.,
congenital heart defects, cerebral arteriovenous
malformations, septal defects), heart disease (e.g., heart
failure, congestive heart disease, arrhythmia, tachycardia,
fibrillation, pericardial Disease, endocarditis), cardiac
35 arrest, heart valve disease (e.g., stenosis, regurgitation,
prolapse), vascular disease (e.g., hypertension, coronary
artery disease, angina, aneurism, arteriosclerosis,

peripheral vascular disease), hyponatremia, hypematremia, hypokalemia, and hyperkalemia.

The term "a disease or condition described as mixed fetal" as used herein refers to a disease or condition selected from the group consisting of: spina bifida, hydranencephaly, neurofibromatosis, fetal alcohol syndrome, diabetes mellitus, PKU, Down's syndrome, Patau syndrome, Edwards syndrome, Turner syndrome, Apert syndrome, Carpenter syndrome, Conradi syndrome, Crouzon syndrome, cutis laxa, Cornelia de Lange syndrome, Ellis-van Creveld syndrome, Holt-Oram syndrome, Kartagener syndrome, Meckel-Gruber syndrome, Noonan syndrome, Pallister-Hall syndrome, Rubinstein-Taybi syndrome, Scimitar syndrome, Smith-Lemli-Opitz syndrome, thrombocytopenia-absent radius (TAR) syndrome, Treacher Collins syndrome, Williams syndrome, Hirschsprung's disease, Meckel's diverticulum, polycystic kidney disease, Turner's syndrome, and gonadal dysgenesis, Klippel-Feil syndrome, Osteogenesis imperfecta, muscular dystrophy, Tay-Sachs disease, Wilm's tumour, neuroblastoma, and retinoblastoma.

The term "a disease or condition of the excretory system" as used herein refers to a disease or condition selected from the group consisting of: bladder cancer, prostate cancer, benign prostatic hyperplasia, bladder disorders (e.g., urinary incontinence, urinary retention, urinary obstruction, urinary tract infections, interstitial cystitis, prostatitis, neurogenic bladder, hematuria), renal disorders (e.g., hydronephrosis, proteinuria, renal failure, pyelonephritis, urolithiasis, reflux nephropathy, and unilateral obstructive uropathy).

The term "a disease or condition of the neural/sensory system" as used herein refers to a disease or condition selected from the group consisting of: brain cancer (e.g., brain stem glioma, brain tumours, central nervous system

- 23 -

(Primary) lymphoma, central nervous system lymphoma, cerebellar astrocytoma, and cerebral astrocytoma, neurodegenerative disorders (e.g., Alzheimer's Disease, Creutzfeldt-Jakob Disease, Parkinson's Disease, and
5 Idiopathic Presenile Dementia), encephalomyelitis, cerebral malaria, meningitis, metabolic brain diseases (e.g., phenylketonuria and pyruvate carboxylase deficiency), cerebellar ataxia, ataxia telangiectasia, and AIDS Dementia Complex, schizophrenia, attention deficit disorder,
10 hyperactive attention deficit disorder, autism, and obsessive compulsive disorders.

The term "a disease or condition of the respiratory system" as used herein refers to a disease or disorder selected
15 from the group consisting of: cancers of the respiratory system such as larynx cancer, pharynx cancer, trachea cancer, epiglottis cancer, lung cancer, squamous cell carcinomas, small cell (oat cell) carcinomas, large cell carcinomas, and adenocarcinomas. Allergic reactions, cystic
20 fibrosis, sarcoidosis, histiocytosis X, infiltrative lung diseases (e.g., pulmonary fibrosis and lymphoid interstitial pneumonia), obstructive airway diseases (e.g., asthma, emphysema, chronic or acute bronchitis), occupational lung diseases (e.g., silicosis and
25 asbestosis), pneumonia, and pleurisy.

The term "a disease or condition of the endocrine system" as used herein refers to a disease or condition selected from the group consisting of: cancers of endocrine tissues
30 and organs (e.g., cancers of the hypothalamus, pituitary gland, thyroid gland, parathyroid glands, pancreas, adrenal glands, ovaries, and testes), diabetes (e.g., diabetes insipidus, type I and type II diabetes mellitus), obesity, disorders related to pituitary glands (e.g.,
35 hyperpituitarism, hypopituitarism, and pituitary dwarfism), hypothyroidism, hyperthyroidism, goiter, reproductive disorders (e.g., male and female infertility), disorders

- related to adrenal glands (e.g., Addison's Disease, corticosteroid deficiency, and Cushing's Syndrome), kidney cancer (e.g., hypemephroma, transitional cell cancer, and Wilm's tumour), diabetic nephropathy, interstitial
- 5 nephritis, polycystic kidney disease, glomerulonephritis (e.g., IgM mesangial proliferative glomerulonephritis and glomerulonephritis caused by autoimmune disorders; such as Goodpasture's syndrome), and nephrocalcinosis.
- 10 The term "a disease or condition of the digestive system" as used herein refers to a disease or condition selected from the group consisting of: ulcerative colitis, appendicitis, Crohn's disease, hepatitis, hepatic encephalopathy, portal hypertension, cholelithiasis, cancer
- 15 of the digestive system (e.g., biliary tract cancer, stomach cancer, colon cancer, gastric cancer, pancreatic cancer, cancer of the bile duct, tumours of the colon (e.g., polyps or cancers), and cirrhosis), pancreatitis, ulcerative disease, pyloric stenosis, gastroenteritis,
- 20 gastritis, gastric atropy, benign tumours of the duodenum, distension, irritable bowel syndrome, malabsorption, congenital disorders of the small intestine, bacterial and parasitic infection, megacolon, Hirschsprung's disease, aganglionic megacolon, acquired megacolon, colitis,
- 25 anorectal disorders (e.g., anal fistulas, haemorrhoids), congenital disorders of the liver (e.g., Wilson's disease, hemochromatosis, cystic fibrosis, biliary atresia, and alpha 1-antitrypsin deficiency), portal hypertension, cholelithiasis, and jaundice.
- 30
- The term "a disease or condition of the connective / epithelial" as used herein refers to a disease or condition selected from the group consisting of: connective tissue metaplasia, mixed connective tissue disease, focal
- 35 epithelial hyperplasia, epithelial metaplasia, mucoepithelial dysplasia, graft v. host disease, polymyositis, cystic hyperplasia, cerebral dysplasia,

- 25 -

tissue hypertrophy, Alzheimer's disease, lymphoproliferative disorder, Waldenstrom's macroglobulinemia, Crohn's disease, pernicious anaemia, idiopathic Addison's disease, glomerulonephritis, bullous pemphigoid, Sjogren's syndrome, diabetes mellitus, cystic fibrosis, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, osteoporosis, osteoarthritis, periodontal disease, wound healing, relapsing polychondritis, vasculitis, polyarteritis nodosa, Wegener's granulomatosis, cellulitis, rheumatoid arthritis, psoriatic arthritis, discoid lupus erythematosus, systemic lupus erythematosus, scleroderma, CREST syndrome, Sjogren's syndrome, polymyositis, dermatomyositis, mixed connective tissue disease, relapsing polychondritis, vasculitis, Henoch-Schonlein syndrome, erythema nodosum, polyarteritis nodosa, temporal (giant cell) arteritis, Takayasu's arteritis, Wegener's granulomatosis, Reiter's syndrome, Behcet's syndrome, ankylosing spondylitis, cellulitis, keloids, Ehler Danlos syndrome, Marfan syndrome, pseudoxantoma elasticum, osteogenesis imperfecta, chondrodysplasias, epidermolysis bullosa, Alport syndrome, and cutis laxa.

The term "a" and "the" as used in the present descriptive is intended to include both one (the singular) and more than one (plural).

A "therapeutically effective amount" of an active agent or combination of agents as described herein is understood to comprise an amount effective to elicit the desired response but insufficient to cause a toxic reaction. A desired response, for example, may constitute the formation of a sufficient and/or acceptable detectable antibody titer level in a blood sample. The dosage and duration of treatment of the preparation to be administered to a subject will be determined by the health professional attending the subject in need of treatment, and will consider the age, sex, weight, extent of existing diseased

- 26 -

state and/or tissue damage of the subject, and specific formulation of *Helicobacter* and the gene of interest product being used as the treatment for the subject.

- 5 The phrase, "effective level" refers to the level of the desired activity of the molecules and not necessarily limited to the number of molecules. For example, the effective level of amylin may be decreased to stimulate ghrelin secretion by using amylin antagonists, without a
10 necessary concomitant decrease in the amount of free amylin present in a subject.

The phrase "ghrelin-associated diseases and disorders" refers to any condition that can be treated prevented or
15 ameliorated through the modulation of ghrelin activity. These include conditions that are enhanced, exacerbated or stimulated by ghrelin, for example, growth hormone release or drive to eat. The physiological actions of ghrelin are considered to include, by way of example, the stimulation
20 of growth hormone release, the stimulation of hormone secretion from lactotrophs and corticotropes, orexigenic and cardiovascular actions, anti-proliferative effects on thyroid and breast tumors and regulation of gastric motility and acid secretion through vagal mediation. (See
25 WO 2005021026).

Throughout the specification, unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply
30 the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

Where the definition of terms departs from the commonly
35 used meaning of the term, applicant intends to utilize the definitions provided herein, unless specifically indicated.

DETAILED DESCRIPTION OF THE INVENTION

Before describing the present invention in detail, it is to
5 be understood that this invention is not limited to
particularly exemplified methods and may, of course, vary.
It is also to be understood that the terminology used
herein is for the purpose of describing particular
embodiments of the invention only, and is not intended to
10 be limiting which will be limited only by the appended
claims.

All publications, patents and patent applications cited
herein, whether supra or infra, are hereby incorporated by
15 reference in their entirety. However, publications
mentioned herein are cited for the purpose of describing
and disclosing the protocols, reagents and vectors which
are reported in the publications and which might be used in
connection with the invention. Nothing herein is to be
20 construed as an admission that the invention is not
entitled to antedate such disclosure by virtue of prior
invention.

Furthermore, the practice of the present invention employs,
25 unless otherwise indicated, conventional immunological and
molecular biological techniques and pharmacology within the
skill of the art. Such techniques are well known to the
skilled worker, and are explained fully in the literature.
See, eg., Coligan et al. "Current protocols in Protein
30 Science" (1999) Volume I and II (John Wiley & Sons Inc.);
Sambrook et al., (Molecular Cloning: A Laboratory Manual,
2nd & 3rd Editions, Cold Spring Harbor Laboratory press
(1989) (2001); and Bailey, J.E. and Ollis, D.F.,
Biochemical Engineering Fundamentals, McGraw-Hill Book
35 Company, NY, 1986.

It must be noted that as used herein and in the appended

claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a nucleic acid" includes a plurality of such nucleic acids, and a
5 reference to "an isolated peptide" is a reference to one or more peptides, and so forth. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any
10 materials and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred materials and methods are now described.

15 Delivery of therapeutic compositions and nucleic acids to specific target sites within the animal body is an ongoing challenge faced by the drug development industry. The present inventor has developed a *Helicobacter*-based bacterial delivery system capable of carrying vectors
20 encoding biologically active agents, wherein these agents are expressed on the surface of the bacterium, or secreted therefrom. In one embodiment, the bacterium is a species of *Helicobacter*. More preferably, *H. pylori*. In some embodiments the strain of *H. pylori* can be any strain known
25 in the field. In some embodiments, the *H. pylori* strain is a non-pathogenic strain such as genomic strain 26695.

In another embodiment, a bacterium, other than *Helicobacter*, is utilised wherein the bacterium has been
30 genetically altered such that it has *Helicobacter* or *H. pylori* features including the ability to chronically colonise the gastric mucosa or other areas of gastrointestinal tract, urinary tract, bronchial epithelium or other mucosal organ, without significant toxicity to the
35 host.

In one embodiment, the *H. pylori* has been manipulated so

that some of the pathogenic features have been removed and/or attenuated. For example, the vacuolating cytotoxin and the *cag* pathogenicity island genes can be removed so that the *H. pylori* is less pathogenic. In one embodiment, 5 the *H. pylori* has been manipulated so that some of the pathogenic features have been removed and/or attenuated. For example, the vacuolating cytotoxin and the *cag* pathogenicity island genes can be removed so that the *H. pylori* are less pathogenic. Attenuating mutations can be 10 introduced into *Helicobacter* using non-specific mutagenesis either chemically, using N-methyl-N-nitro-N-nitrosoguanidine, or using recombinant DNA technologies.

The skilled person will appreciate that the methods of the 15 present invention could be used to deliver a range of biologically active agents. Examples of suitable agents include ones which are capable of functioning locally or systemically, eg an agent capable of exerting endocrine activities affecting local or whole-body metabolism and/or 20 an agent which is capable of regulating the activities of cells belonging to the immuno/haemopoietic system and/or an agent which is capable of affecting the viability, growth and differentiation of a variety of normal or neoplastic cells in the body or affecting the immune regulation or 25 induction of acute phase inflammatory responses to injury and infection and/or an agent which is capable of enhancing or inducing resistance to infection of cells and tissues mediated by chemokines acting on their target cell receptors, or the proliferation of epithelial cells or the 30 promotion of wound healing and/or an agent which modulates the expression or production of substances by cells in the body.

Specific examples of such biologically active agents 35 include insulin, growth hormone, prolactin, calcitonin, luteinising hormone, parathyroid hormone, somatostatin, thyroid stimulating hormone, vasoactive intestinal

- 30 -

polypeptide, a structural group 1 cytokine adopting an antiparallel 4 α helical bundle structure such as IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, IL-11, IL-12, IL-13, GM-CSF, M-CSF, SCF, IFN- γ , EPO, G-CSF, LIF, OSM, CNTF, GH, PRL or IFN α/β , a structural group 2 cytokine which are often cell-surface associated, form symmetric homotrimers and the subunits take up the conformation of β -jelly roll described for certain viral coat proteins such as the tumor necrosis factor (TNF) family of cytokines, eg TNF α , TNF β , CD40, CD27 or FAS ligands, the IL-1 family of cytokines, the fibroblast growth factor family, the platelet derived growth factors, transforming growth factor β and nerve growth factors, a structural group 3 cytokine comprising short chain α/β molecules, which are produced as large transmembrane pre-cursor molecules which each contain at least one epidermal growth factor (EGF) domain in the extracellular region, e.g., the EGF family of cytokines, the chemokines characterised by their possession of amino acid sequences grouped around conserved cysteine residues (the C--C or C--X--C chemokine subgroups) or the insulin related cytokines, a structural group 4 cytokine which exhibit mosaic structures such as the heregulins or neuregulins composed of different domains, e.g., EGF, immunoglobulin-like and kringle domains.

Alternatively, the biologically active agent can be a receptor or antagonist for biologically active agent as defined above.

In some embodiments, the *H. pylori*-based vector and/or vector plasmid construct is employed to create transformed cells (such as an *E. coli* or *Helicobacter* cell) that permits the expression and/or secretion of a biologically active agent from an isolated nucleic acid molecule contained within it at the mucosa surface of a host to which the transformed cell preparation is administered. The isolated nucleic acid contained within the transformed cell

- 31 -

(or vector) may comprise one or more nucleic acid constructs in which nucleic acid encoding the biologically active agent is under control of appropriate regulatory sequences for expression in the *H. pylori*.

5

Suitable vectors and shuttle vector sequences comprising nucleic acid for introduction into *H. pylori* can be chosen or constructed, to contain appropriate regulatory sequences, including promoter sequences, terminator
10 fragments, enhancer sequences, marker genes and other sequences as appropriate. Vectors may be plasmids, viral eg. phage or phagemid, as appropriate. For further details see, for example, Sambrook *et al.*, *supra*. Many known techniques and protocols for manipulation of nucleic acid,
15 for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in Short Protocols in Molecular Biology, Second Edition, Ausubel *et al.* eds., John Wiley & Sons, 1992. The
20 disclosures of Sambrook *et al.* *supra* and Ausubel *et al.* are incorporated herein by reference.

In some embodiments, the coding sequence for the biologically active agent is contained in an operon, i.e.,
25 a nucleic acid construct for multi-cistronic expression. In an operon, transcription from the promoter results in a mRNA which comprises more than one coding sequence, each with its own suitably positioned ribosome binding site upstream. Thus, more than one agent can be translated from
30 a single mRNA. Use of an operon enables expression of the biologically active agent to be coordinated.

A nucleic acid construct or vector comprising a coding sequence for a biologically active agent of the present
35 invention is preferably under the control of a promoter for expression in *H. pylori*.

In one embodiment, the promoter employed in accordance with the present invention is expressed constitutively in the *H. pylori*. Use of a constitutive promoter avoids the need to supply an inducer or other regulatory signal for expression to take place. Preferably, the promoter directs expression at a level at which the *H. pylori* host cell remains viable, i.e., retains some metabolic activity, even if growth is not reduced. Advantageously then, such expression may be at a low level. For example, where the expression product accumulates intracellularly, the level of expression may lead to accumulation of the expression product at less than about 10% of cellular protein, preferably about or less than about 5%, for example about 1-3%.

The promoter may be homologous to the *H. pylori* strain employed, i.e., one found in that strain of *H. pylori* in nature. In some embodiments the promoter is arabinose inducible promoter. Other promoters include FlaB sigma 54 promoter (Josenhans et al., 1998, *FEMS Microbiol Lett*, 161(2): 263-73), T7 promoter, and nir B promoter of salmonella (Chatfield et al., 1992, *Biotechnology*, 10(8): 888-92).

In another embodiment the promoter is inducible. Inducible promoters that may be used with the clinical grade vectors include, but are not limited to, a pH inducible promoter as described in U.S. Pat. No. 6,242,194 issued to Kullen et al., a lactose inducible promoter such as that used in *E. coli* plasmids (e.g., pBluescript™ from Stratagene) or the endogenous lactose promoter in *Lactobacillus*; promoters induced during anaerobic growth such as the promoter for alcohol dehydrogenase (*adhE*), as described in Aristarkhov et al., 1999, *J. Bacteriology*, Vol. 178(14), 4327-4332.

In one embodiment, the constructs of the present invention also include a toxic gene. These toxic genes are preferably under the control of inducible promoters so

- 33 -

that, on completion of treatment, the *Helicobacter* of the present invention can be readily eliminated by inducing the expression of the toxic gene. Non-limiting examples of toxic genes include bacterial autolysins under the control
5 of an inducible promoter. The autolysing gene may then be triggered at the appropriate time and place in the gastrointestinal tract through the use of one or more of the inducible promoters described immediately above.

10 In some embodiment, the engineered *Helicobacter* vector and plasmid vector constructs are sensitive to oxygen. This oxygen sensitivity is another method for limiting dissemination of the clinical grade vectors of the present invention. The environment of the human gut is very low in
15 oxygen, suitable for growth of anaerobic and microaerophilic microorganisms, including *Helicobacter*. Thus, an efficient means of eliminating *Helicobacter* delivery vehicles once they have exited the human body upon discharge of intestinal waste into the oxygen-rich outside
20 environment, is to engineer genes into the transformed micro-organisms that confer oxygen sensitivity.

The nucleic acid construct or constructs of the present invention may comprise a secretory signal sequence. Thus,
25 in some embodiments the nucleic acid encoding the biologically active agent eg non-*Helicobacter* polypeptide, may provide for secretion of the agent at a cell membrane by appropriately coupling a nucleic acid sequence encoding a secretory signal sequence to the nucleic acid sequence
30 encoding the molecule (polypeptide). The ability of *Helicobacter* harbouring the nucleic acid to secrete the polypeptide may be tested *in vitro* in culture conditions, which maintain viability of the *Helicobacter*.

35 Suitable secretory signal sequences include any of those with activity in Gram negative organisms such as *Escherichia*, *Klebsiella* and *Salmonella*. Secretory signal

- 34 -

sequences may include the secretion leader of the Staphylokinase enzyme secreted by some strains of Staphylococcus, which is known to function in both Gram-positive and Gram-negative hosts (see "Gene Expression
5 Using *Bacillus*", Rapoport (1990) *Curr Opin Biotech* 1:21-27).

Other secretory signal sequences that can be used include, for example, the β -lactamase gene (Talmadge et al., 1980,
10 *Proc. Natl. Acad. Sci. USA* 77:3369-3373) or the enteroinvasive *E. coli* hemolysin A (hlyA) (Su et al., 1992, *Microbial Pathogen*, 13:465-476). An illustrative list of secretory signal sequences is presented in Pugsley, 1988, Protein secretion across the outer membrane of gram-
15 negative bacteria. In: Protein Transfer and organelle Biogenesis, R. C. Dand and P. W. Robbins (eds), Academic Press, Inc., San Diego, pp 607-652.

Selectable markers provide researchers and technicians a
20 convenient means for distinguishing transformed microorganisms from non-transformed ones in a mixed population. One means of identifying transformed organism is to incorporate a selectable marker nucleic acid sequence into the plasmid containing the gene of interest. The
25 selectable marker sequence is generally inserted downstream of the gene of interest and is driven off the same promoter. As a result, cells successfully transformed with the gene of interest will also be transformed with selectable marker nucleic acid sequence. When antibiotic
30 resistance is used as the selectable marker, only transformed cells will survive and/or grow in media containing the antibiotic.

Thus, antibiotic resistance is a convenient and much used
35 phenotype when developing transformants. However, vectors having antibiotic resistant genes as selective markers are capable of horizontal gene transfer that can endow other

- 35 -

organisms with antibiotic-resistant phenotypes. This risk is especially acute when *Helicobacter* is used as part of a therapeutic vector.

5 In order to use *Helicobacter* as a gene delivery system to animals, the present disclosure presents, in some embodiments, a clinical grade vector system that does not use an antibiotic selection marker. One of the alternatives to using antibiotic resistance genes provided by the
10 present delivery systems includes clinical grade vectors having chromosomal deletions or lethal mutations in an essential "house-keeping" gene. Next, a functional analogous house-keeping gene is inserted into a plasmid encoding for a gene of interest. Consequently, the "house-
15 keeping" gene becomes the selectable marker allowing for the rapid isolation and identification of transformants.

Examples of essential "house-keeping" genes include genes that encode for any number of metabolic regulators and/or
20 enzymes including, but not limited to kinases, proteases, synthetases, dehydrogenases and others. Another alternative to antibiotic resistance genes provided by the present invention includes clinical grade vectors having reporter genes incorporated into the plasmid containing the gene of
25 interest. Other non-limiting examples of reporter genes used in accordance with the teachings of the present invention include green fluorescent Protein (GFP), β -galactosidase and amylase.

30 In one embodiment, the biologically active polypeptide preferably has cytokine activity. Cytokines are discussed in "The Cytokine Facts Book", Callard and Gearing (1994), Academic Press. Preferred polypeptides with cytokine activity are interleukins, including Interleukin-2 (IL-2)
35 and Interleukin 6 (IL-6).

In some embodiments, the *Helicobacter* delivery system,

- 36 -

vector or vector plasmid system comprises a nucleic acid construct as described above is introduced into a *Helicobacter* or other suitable host cell, to provide transformed cells. Thus, a further aspect of the present invention provides a method comprising introducing nucleic acid as disclosed into a non-pathogenic *Helicobacter*. The transformation of a culture of host cells, such as *Helicobacter*, may employ any available technique. For *H. pylori* cells, suitable techniques may include calcium chloride transformation, electroporation and transfection using bacteriophage.

The introduction of the vector plasmid into a *Helicobacter* cell may be followed by causing or allowing expression from the nucleic acid, e.g., by culturing *H. pylori* under conditions for expression of the gene. Growing the *Helicobacter* in culture under conditions for expression of the biologically active polypeptide may be employed to verify that the *Helicobacter* contain the encoding nucleic acid and is able to produce the encoded material.

In a further aspect, the present invention provides a method of delivering a therapeutic or prophylactic dose of a biologically active agent *in vivo*, the method comprising administering to a subject an effective amount of the non-pathogenic preparation of the *H. pylori* compositions and vaccines of the present invention.

It will be appreciated that the methods of the present invention and the use of a non-invasive or non-pathogenic *Helicobacter* as described herein provide a wide range of therapeutic methods which would enable the skilled person to manipulate, for instance, the immune response of a subject. Thus, in one aspect the present invention provides a method of regulating the survival, growth, differentiation, effector functions or susceptibility to infection of cells or tissues which comprises administering

to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

5 In another aspect, a method of boosting an immune response against tumour cells or an infection colonising a mucosal surface or adjacent or distant tissue is provided which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

10 In yet another aspect, a method of modulating the type of immune response (antibody versus cell-mediated) against a pathogenic infectious agent is provided which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

15 In another aspect, a method of modulating the infiltration of normal tissues with inflammatory or tumour cells is provided which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

20 In some aspects a method of controlling the rate of growth, rate of invasion or survival of tumour cells is provided which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

25 In yet another aspect a method of inducing apoptosis in tumour cells is provided which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* as defined herein.

30 Other aspects provided for a method of down-regulating an immune response is provided which comprises administering to a subject a non-invasive or non-pathogenic bacterium which expresses a biologically active agent.

35 In another aspect a method of treating an allergic autoimmune or other immune dysregulative disease state is

- 38 -

provided, which comprises administering to a subject a non-invasive or non-pathogenic *Helicobacter* which expresses a biologically active agent.

- 5 The subject can be any primate, equine, bovine, porcine, ovine, rodent, fish, or bird. In one embodiment, the subject is human. Administration may conveniently be nasal or oral.
- 10 In a therapeutic context, i.e., where the biological effect of delivery of the biologically active agent to a subject is beneficial to that subject, administration is preferably in a "therapeutically effective amount", this being sufficient to show benefit to a subject. Such benefit may
- 15 be at least amelioration or reduce the severity or occurrence of at least one symptom. In a prophylactic context, the amount may be sufficient to reduce the deleterious effect on the subject of a subsequent pathogenic challenge, for instance by enhancing the immune
- 20 response. The actual amount administered, and rate and time-course of administration will depend on the aim of the administration, e.g. the biological effect sought in view of the nature and severity of the challenge, and is the subject of routine optimisation. Prescription of treatment,
- 25 including prophylactic vaccination, for example decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors.

A composition comprising *Helicobacter* may be administered

30 in accordance with the present invention alone or in combination with other treatments, either simultaneously or sequentially.

The present invention also provides a pharmaceutical

35 composition comprising a *Helicobacter* as disclosed. Such a pharmaceutical composition is in one embodiment preferably suitable for application to a mucosal membrane.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may comprise, in addition to the *Helicobacter*, a
5 pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may
10 depend on the route of administration. For oral administration a parenterally acceptable aqueous solution may be employed which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions.

15 Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required. As discussed, a pharmaceutical comprising a *Helicobacter* for administration in accordance with the present invention may comprise one or more nutrient substances, e.g., an energy
20 source such as glucose, amino acids and so on.

In another aspect, the present invention provides a method of manufacture of pharmaceutical formulations comprising formulating *Helicobacter* as disclosed with a suitable
25 carrier medium suitable for administration to an individual. In one embodiment, the pharmaceutical is suitable for application to a mucosal membrane of an individual.

30 In another aspect, the present invention provides a non-pathogenic *Helicobacter* expressing a heterologous biologically active polypeptide for pharmaceutical use, e.g., for use in a method of treatment of the human or animal body by surgery or therapy, including prophylaxis
35 ("vaccination").

In one embodiment the methods of the present invention can

- 40 -

be used to treat, prevent or palliate a disease such as cancer. The methods and deliver system can also be used to treat or prevent a disease or condition of the immune/haematopoietic system, a disease or condition of the reproductive system, a disease or condition of the musculoskeletal system, a disease or condition of the cardiovascular system, a disease or condition described as mixed fetal, a disease or condition of the excretory system, a disease or condition of the neural/sensory system, a disease or condition of the endocrine system, a disease or condition of the respiratory system, a disease or condition of the digestive system and a disease or condition associated with connective/epithelial tissue or a disease or condition caused by bacterial, viral or parasitic infection.

In another embodiment, the *Helicobacter* delivery system described herein is capable of concomitant or sequential delivery of a number of different nucleic acid molecules, which encode products capable of treating a number of conditions or diseases as described herein. Moreover, preferred delivery systems would also deliver compositions capable of producing additional desirable physiological effects such as appetite suppression or enhancement.

An example of suicide system in *H. pylori* has been described by Panthel et al. 2003 (*Infection & Immunity*, 71: 109-116). This system introduces a plasmid into *H. pylori* which contains the PhiX174 lysis gene E. To eradicate the strain, incubation at 42°C for 5 hours was used. *In vivo* this would mean that the animal would consume a drink at 45-50°C to raise the temperature of the gastric environment above 42°C.

A second example is the L-Dap selection system, commonly used to allow survival of bacterial mutants on supplemented plates (see, for example, Kirata et al. 1997 (*Infection &*

Immunity, 65: 4158-4164). In this system the animal subject must supplement their diet with a missing substrate i.e., diamino-pimelic-acid (DAP), in order for the DapE deficient *H. pylori* mutant to survive. In order to eradicate the mutants, DAP consumption is ceased.

A third possible system relates to metronidazole sensitivity of *H. pylori* because of its *rdxA* gene. Excessive replication of the *rdxA* gene is harmful to mammalian cells and *E. coli*. However, duplication may be tolerated by the bacterium. Therefore a *Helicobacter* species of the present invention can be engineered to contain two copies of *rdxA* which prevent the normal mutation-dependant *rdxA* loss. The introduction of at least two functional *rdxA* genes into the *Helicobacter* genome will result in a *Helicobacter* strain that is permanently sensitive to metronidazole. Jeong et al. 2000 (*J. Bacteriol.*, 182: 5082-5090) showed that the nitroreductase produced by a functional *rdxA* gene converts metronidazole from a prodrug to a bactericidal compound. The mode of action of the active compound is to cause DNA breaks of the *Helicobacter* genome.

The invention will now be further described by reference only to the following non-limiting examples. It should be understood, however, that the examples following are illustrative, and should not be taken in any way as a restriction on the generality of the invention described herein. In particular, while the invention is described in detail in relation to the use of a specific *H. pylori* strain, it will be clearly understood that the findings herein are not limited to this strain.

EXAMPLE 1 VECTORS AND TRANSGENIC *H. pylori* ORGANISMS
FOR STABLE EXPRESSION OF FOREIGN PROTEINS

The genetic manipulation of *H. pylori* is uncommon. The
5 present example demonstrates the utility of the invention
for providing a genetically transformed *Helicobacter*,
particularly transformed *H. pylori*. The transformed
bacterium are prepared using plasmids and plasmid vectors
derived from *Helicobacter*, which have had been subject to
10 prior manipulation in a non-*Helicobacter* organism, such as
E. coli.

Several *H. pylori* plasmids described in the literature can
be successfully converted to *H. pylori*/*E. coli* shuttle
15 vectors. Many strains of *E. coli* have been reported to be
naturally competent for DNA uptake. Resistance markers for
streptomycin, rifampin and metronidazole have also been
successfully transformed into most strains of *H. pylori*.
However, while plasmid DNA from *E. coli* and other organisms
20 can be introduced into *H. pylori* these plasmids cannot be
stably maintained. Moreover, *H. pylori* plasmids cannot be
transformed into *E. coli* or *Helicobacter* species.
Accordingly, *H. pylori* shuttle vectors must be constructed.

25 Two plasmids from *H. pylori* are illustrated in the
schematics shown in Figures 1 and 2. Vectors pHPA1 (2.8kb)
(Fig. 1) and pHP3 (3.4kb) (Fig. 2) have been sequenced and
it has been revealed that pHPA1 replicated via the theta
mode of plasmid replication. In contrast to rolling-circle
30 replicating plasmids, theta plasmids do not generate
single-stranded DNA intermediates during replication and
are thus more stable vector candidates because they are
less prone to undergo illegitimate recombination.
Furthermore, the pHPA1 origin of replication (*ori*) contains
35 a series of direct repeat sequences (termed "iterons") that
are involved in replication control and maintaining stable
copy number. Vector pHP623 shares many of these features.

The nucleotide sequences for these two vectors are shown below.

(1) Plasmid pHP1 shown in double stranded form (top strand is SEQ ID NO:1; bottom strand is SEQ ID NO:2)

5	GTCATGCGCGTTGTTTTTAATTACATTTTAAACAACCTTGTTGTTGTTTTTACATGTTTTACTCGC CAGTACGCGCAACAAAATTAATGTAAATTTGTTGAACAACAACAAAATGTACAAAATGAGCG	65
10	ATGCGCGCGCGTGAGGGATTGGGGGTTGCAACCCCTAAATAACGAAGCTGTAGGGTTTCTCATT TACGCGCGCGCACTCCCTAACCCCAACGTTGGGGGATTATTTGCTTCGACATCCCAAAGAGTAA	130
	TTTGTGGTGAAAATGAATAAAACAGAACTTCTTGCCAACACTAACAGAACTTCTTGCCAACACTA AAACACCACTTTTACTTATTTTGTCTTGAAGAACGGTTGTGATTGTCTTGAAGAACGGTTGTGAT	195
15	ACAGAACTTCTTGCCAACACTAACAGAACTTCTTGCCAACACTAACAGAACTTCTTTATTTTAAA TGCTTGAAGAACGGTTGTGATTGTCTTGAAGAACGGTTGTGATTGTCTTGAAGAAATAAAATTT	260
	GTTATGATTATTAACAATTTTTAGACATAATAACAGCGTGTGAAGATACTTTTGTAGCGGTATTT CAATACTAATAATTGTTAAAAATCTGTATTATTGTGCGCACACTTCTATGAAAACATCGCCATAAA	325
20	CCTATGTGCGGCAAAATTTGGAGCAATTAGCTTGACTTGTTGAGTTAGTGGGTTGGAGGATAGA GGATACACGCCGTTTTAAACCTCGTTAATCGAACTGAACCAACTCAATCACCCAACCTCCTATCT	390
25	GAGGCGACACCTCGTTAGGAGGTATCAATGTGAAAGTATTTGTCGTATTAGTTCTAGTATTAGT CTCCCGCTGTGGAGCAATCCTCCATAGTTACACTTTCATAAACAGCATAATCAAGATCATAATCA	455
	AATTCTCGCACAAATGCTATATTAGGCTTATTCGTGGTCTAACCCCTGTTTATGGGGGTTGGCT TTAAGAGCGTGTTAACGATATAATCCGAATAAGCACCAGATTGGGGAACAAATACCCCAACCGA	520
30	CGTTATAAGCATACTGATACGATCACACTTATTATACACCAAAAGATAAGGAGTATAGAGTGGA GCAATATTCTGATGACTATGCTAGTGTGAATAATATGTGTTTTCTATTCTCATATCTCACCTT	585
	TTTGATCAATCAGATTTACAAAAGCGTTGAAAATATTAGATACACTCCACAAACCCACAAAT AAACTAGTTAGTCTAAATGTTTTTCGCAACTTTTATAATCTATGTGAGGGTGTTTGGGGTGTTTA	650
35	TGAGCTACAAAACAAGAAATACAAAACCGCATCAACAAAATAACAGAGACAATCATTAAAGAAT ACTCGATGTTTTTGTCTTTATGTTTTGGCGTAGTTGTTTTATTGTCTCTGTTAGTAATTTCTTA	715
40	TACTATCAAAGCATGAAATCAAGAAAGAAGAACTAGAACCCACTCTAACCCCAAAACCCACACCA ATGATAGTTTTCTACTTTAGTTCTTTCTTCTTGATCTTGGGTGAGATTGGGGTTTTGGGTGTGGT	780
	CTCAAAGAGCCACAAACCACCCCAACACCATGCAAAGATTTAGTGGTTAGCACCCCTAAAGATAA GAGTTTCTCGGTGTTTGGTGGGGTTGTGGTACGTTTCTAAATCACCAATCGTGGGGATTTCTATT	845
45	AACCTAATATCACCTACCACAATAACGCTAATAAGGTCAATCTAGGGAAATTGAGCGAAAGGGAA TTGGATTATAGTGGATGGTATTGCGGATTATCCAGTTAGATCCCTTTAACTCGCTTTCCCTT	910
	GCCAATCTTTTATTCGCTATTTTTCAAAAACCTCAAAGCCCAAGGGAATACCCTCATTCGTTTTGA CGTTAGAAAATAAGCGATAAAAAGTTTTTGAGTTTCGGGTTCCCTTATGGGAGTAAGCAAACT	975
50	ACCGCAAGATTTGAAACGCATGCTAAACATAGATATTTCTAATGAGCGCTTATCAGAAGTCGTTA TGGCGTTCTAAACTTTGCGTACGATTTGTATCTATAAAGATTACTCGCGAATAGTCTTCAGCAAT	1040
55	TTAAGCTGTGGGATAGCATTAAAACCGCTGATTTTTTGAAAATTAGCGAAACCGAAACTTCAATC AATTCGACACCTATCGTAATTTTGGCGACTAAAAACCTTTTAATCGCTTTGGCTTTGAAGTTAG	1105
	ATTCAAGAAAATTACATGCTTTTTAGTCGGTGTAATAATTGAATTGAACAAACCGAGTAAAGATTT TAAGTTCTTTAATGTACGAAAATCAGCCACATTTAACTTAAGTTGTTTGGCTCATTTCTAAA	1170
60	GAAGTATTTAGAAATCCAACCTCAACGATAACTATCAAGACTTACTCAACAATCTGGGCATGGGTC CTTCATAAATCTTTAGGTTGAGTTGCTATTGATAGTTCTGAATGAGTTGTTAGACCCGTACCCAG	1235
	AATACACTTCTTTCAATCTGTTAGAAATTTCAAAGAGTGAGGGGTAAATACGCTAAAACGCTCTAT TTATGTGAAGAAAGTTAGACAATCTTAAAGTTTCTCACTCCCATTTATGCGATTTTGGCGAGATA	1300
65	CGCTTGCTCAAGCAATACAAAAGCACAGGGATTTTGAGCGTGGAATGGACTCAATTCAGGGAGCT GCGAACGAGTTGTTTATGTTTTCTGTGCCCTAAAACCTCGCACCTTACCTGAGTTAAGTCCCTCGA	1365
	TTTAGACATTCCAAAAGACTACAAAATGGAAAACATCGATCAAAAAGTCTTAACCCCTCTCTCA	1430

	AAATCTGTAAGGTTTTCTGATGTTTTACCTTTTGTAGCTAGTTTTTCAGAATTGGGGGAGAGAGT	
	AAGAACTCAGAAAAATCTACCCTTTTGAACACTTGAGCTATAAAAAAGAACGCAAAAGCCATTAC	1495
5	TTCTTGAGTCTTTTTTAGATGGGAAAACCTGTGAACTCGATATTTTTCTTGCGTTTTTCGGTAATG	
	AAGCGCAAAGTAACCCACATTGATTTTTATTTTGAGCAATTTCCCTTAAGGCGAAAATAAGAAACA	1560
	TTGCGTTTTTCATTGGGTGTAACATAAAATAAACTCGTTAAAGGAATTCGCTTTTATTCTTTGT	
10	AAACAAAGCCGACAAGCAACGCGCTCAAAGGGACATCAAGCTTGTAGCATGGGATATTCAACACC	1625
	TTTGTTCGGCTGTTGCTTGCGCGAGTTTCCCTGTAGTTCGAACATCGTACCCTATAAGTGTGG	
	AAATCGCTAAAAGAAACGCAAAAGCCACTATGGAAGCTAGGTTTCTTGAATTGAAAACCTTGATC	1690
	TTTAGCGATTTTCTTGCGTTTTTCGGTGATACCTTCGATCCAAAGAACTTAACCTTTTGAACTAG	
15	GGCTATCAGTTCAGGAACAATGACAGTAGGAACAAATTAAGATTGACAACACCCTTTTGAAAG	1755
	CCGATAGTCAAGTCCTTGTTACTGTCATCCTTGTTTAATTTCTAACTGTTGTGGTGAAAACCTTC	
	AATCAAATGTATTTACATGTATCTTAACCTTAAAAATAAGCATAACCCCCAAAAATTCCTTGAT	1820
20	TTAGTTTACATAAATGTACATAGAATTGGGATTTTTATTTCGTATTGGGGGTTTTTAAGGAACATA	
	CCAACAAGACATTCGCATTGGAACACTATATATCAATAGATACAGCCTAAAAAAAAGACAACCTT	1885
	GGTTGTTCTGTAAGCGTAACCTTGATGATATATAGTTATCTATGTCGGATTTTTTTTCTGTTGAA	
25	GCTAGAAGAATTTAACCCCCAAAAATCCACCTATCACCAACGAACCTATCAAGGAATTTGCAGA	1950
	CGATCTTCTTAAATTGGGGGGTTTTAGGTGGGATAGTGGTTGCTTGGATAGTTCCTTAAACGTCT	
	ATACATCGGCAAAACGATTAACATCACCAACTTCAATGTGGATCAATGCCATGAGGGAATCAGCA	2015
	TATGTAGCCGTTTTGCTAATTGTAGTGGTTGAAGTTACACCTAGTTACGGTACTCCCTTAGTCGT	
30	ACTACCTGACAATCACTAGGATCGTGAACCTGGACGTAATCGGATCTGTATTTGGTCCAGATGTGG	2080
	TGATGGACTGTTAGTGATCCTAGCACTTGACCTGCATTAGCCTAGACATAAACAGGTCTACACC	
	ATAAGCCTGGGACTTCTCAAGCCTTTCATTGCTAAAGTGAGAAAATTTGGGGATTGGTTCAAGAA	2145
35	TATTCGGACCTGAAGAGTTCGGAAAGTAACGATTTCACTCTTTTAAACCCCTAACCAAGTTCTT	
	CACTACAGGTGAAAAGACAGATGCATGCTGACTAAACTCATAGAAAACTGAATCACGAAAGAAA	2210
	GTGATGTCCACTTTTCTGTCTACGTACGACTGATTTGAGTATCTTTTGACTTAGTGCTTCTTT	
40	GAATGCAAGCAGAAAACAAACACCTAAAAGAACAAGGACTAGAAAAAATCTACACTCAAAAAGAC	2275
	CTTACGTTTCGCTTTTGTGTTGTGGATTTTCTGTTCTGATCTTTTTTAGATGTGAGTTTTTCTG	
	TACGAGCAGTTAAAAGAACAGCATTTGAAAGAAATTGAAGCACTCAAAAAAGAAATCCAAAAAAC	2340
	ATGCTCGTCAATTTTCTTGTCGTAAACTTCTTTAACTTCGTGAGTTTTTTCTTTAGGTTTTTTG	
45	CAAGCAAGAAACATACACGCAACCAAAAGAATGTAGCCATTTAGCGCATTCTTTTAGCCCTAATT	2405
	GTTTCGTTCTTTGTATGTGCGTTGGTTTTCTTACATCGGTAAATCGCGTAAGAAAATCGGGATTAA	
	CATTCTTTCAATCAAAATCCGACTAATTCATCGGCTAAACGCTAAAAATCGCTTAAAACGAAAAA	2470
50	GTAAGAAAGTTAGTTTTAGGCTGATTAAGTAGCCGATTTGCGATTTTATAGCAATTTTGCTTTTT	
	TACAAAGCAAAAACTTCATTTCCCTTTTAGTCGTTAACCATTTAGCCAATCTAACTAGTTTAGC	2535
	ATGTTTCGTTTTTTGAAGTAAGGGGAAAATCAGCAATTGGTAAATCGGTTAGATTGATCAAATCG	
55	ATCTAAAGGCGAATCTATCTTGTGTTAGACATCCAACCTTACCAAAACCGCAGAGCGAGCTTAAG	2600
	TAGATTTCCGCTTAGATAGAACACAATCTGTAGGTTGGAATGGTTTTGGCGTCTCGCTCGAATTC	
	AGAGATTCAAGCGGTTTTGCACGATTGTTTGTGCTGCAAGAAAACCAACAGCGAAGTAAGGCGCA	2665
	TCTCTAAGTTCGCCAAAACGTGCTAACAAACGACGGTCTTTTGGTTGTTTCGCTTCATTCCGCGT	
60	TAGACAAAAGCGCATCGCAGTTTGAAAGCGTAGGCGTCAGAAAGTGGTTGCGTTAGAATCAAACA	2730
	ATCTGTTTTTCGCGTAGCGTCAAACCTTCGCATCCGCGTCTTACCACAAACGAATCTTAGTTTGT	
	AGATAGCGCAAAACCTGGCGTTAGGCTAAAAAACCCCTAAAAACTAAAAACCCCAAAATATGTAGTGC	2796
	TCTATCGCGTTTGGACCGCAATCCGATTTTTTGGGGATTTTTGATTTTGGGGTTTTATACATCAGC	
65	(	
	2) Plasmid pHP3 shown in single stranded form (SEQ ID	
	NO:3):	
70	TCTACACAATTAACAATCTTTAGCTACAATAACAGCGTGTGAAGATGCTTTCACAGCGGT	60
	ATTTCCCTATGTGCGGCAAAATTTGGAGCAATTAACCTTGACTTGGTTGGGTTAGTGGGTTG	120

	GAGGATAGAGAGGGCGACACCTCGTTAGGAGGTATCAATGTGAAAGTATTTGTCGTATTA	180
	GTTCTAGTATTAGTAATTTCTCGCACAAATGCTATATTAGGCTTATTTGTGGTCTAACCCC	240
	TTGTTTATGGGGGTTAGATCCTTATAAGCATACTGATACGATCACACTTATTATACACCA	300
	AAAGATAAGGAGTATAGAGTGAATTTGATCAATTAGAATCACAAAGATCAGACTTACAA	360
5	AAAGTGTTAAAAGAATTAGATACACTCCCCAAAACCCACAAATTGAGCTACAAAACAA	420
	GAAATACAAAACCGCATCAACAAAATAACAGACACAATCATTAAGAATTACTATCAAAA	480
	CATGAAATCAAAAAGAAGAACTAGAACCCACTCTAACCCCAAAACCCACACCAACAAA	540
	GAGCCACAAACCACCCCCACACCATGCAAAAATTTAGTGGTTAGCACCCCTAAAGATAAA	600
	ACCTATATCACCTACCACAATAACGCTAATAAGGTCAATCTAGGGAAATTGAGCGAAAGG	660
10	GAAGCCAATCTTTTATTCGCTATTTTTCAAAGGCTTAAAGATCAAGGGAATACCTCATT	720
	CGTTTTGAACGCAAGATTTAAAACGCATGATCATGGTCAAATCCAACCTTAACCAACAGG	780
	CAATTATTGCAAGTCTTAAAAAATTTGCTTGACAACATTAGCGGTGCTAATTTTTGGATC	840
	AATTAGAGAGCATGTTGAAAATGGCGAAATCTATGAAGATCACACTAGCTACATGCTTTT	900
	CAACAATTTGAAATCCGCATCCATAAGCCAACACAACTATAGAATACTTAGATGTCCA	960
15	ACTCAATGATAGCTATCAATACTTGCTCAACAATCTAGGAATGGGCGGTCAATACACTTC	1020
	TTTCAATCTCTTAGAATTTCAAAGGGTGAGGGGCAAATAGTGAGAGCGTTAAATTTCCCC	1080
	CCCCTATTCCCCTTAAAAAGGACCCCTTATCCCAGGGAATTTTTGGCCCCAATACAATTAG	1140
	GGCCAAAACCCGGTCCCTTCCATGGCTTAACCAACCCAATTGGGGGATTCCAATTTCCC	1200
	CTGGATGGGAATAACCCAAGGCTTTTTTTGAAAATTCACCTACCATTGGTCCAAAATT	1260
20	GGATGGACAATTCCAAATTCCAAATCTCTTTTCCAAGAATGGGGGCCAACCCCTTGACAA	1320
	ACTCCTTAAACCTTTTCATTTCGGCTAAAAGGTTGAAAAACATTTGGAAGATTTGGTTAA	1380
	GGAAATATTTATCGGGTGAAAAGACCAGATGCATGGCTAACTTAACTCCATAGAAAAAC	1440
	TGAATCACGAAAGAAAGATGCTATCAAAAATGGCATTACCACCTGATCCAAATCAAAT	1500
	TTTCTTACAACCTCCAATCGCATTGAAGGAAGCGGTTGACTTATGAACAAACCGCTCATA	1560
25	TTTTTGACAAATCCGTTCTCATAACTGAAAAAACACCAATATCAAACCTTGATGATATTT	1620
	TTGAAACTATCAATCATTTTGAATGCGTGAATTACTTGCTTGAAAGCTATAAAGAACCTT	1680
	TGAGTTTAGAATACTTTAAGAATTTACACAAAATCTTAAAAAGAATTGTTCTGATGAAG	1740
	TTATTGGTGATTTTAAAAACGCCCTAATTTTGTAGGCAATAGCGCCACAACAAGACCCA	1800
	AATTAGTTGAAAGCGAATTGACAAATCTGTGAAAAATTATCAACGCAACCTTGAAGTGA	1860
30	GTTTGAAAAACAATATCATGCCTTTTCATCATAGAAAACGAACACAAAGCCTTTTACTACA	1920
	GGGGCATCAAGAATATGACAACACAAAAGGCTACTTGAAAGACACCATTTTGCAAAGTC	1980
	AAGACAATTTCAATGAAATGGTTAGCTATTTCTTTTCTTGAGTGAAACCGCTTATTTTTG	2040
	CTTGTGTGCTTTTGTTTTTTGCGTTTTTAGTTGTAGGTGGTAAGAAATATCGGTTTTTTG	2100
	CTTTTCGTTGGTTGTAGGCGATTTTAGATAGCAAAAAACAGCTAAAAAATCCAAGCAAC	2160
35	TAATTGATTTCAAACCAACTTCATTTCCCTTTTAGTCGTTAGCCATTTAGCCAATCTAAC	2220
	TAGTTTAGCATCTAAAAGCGCATATAACTTGAGTTAGCAATCCAACCAATACTAAAACCG	2280
	CCTAGCGAAGCGTTAGCGAGCAAAATAAGCGGTTTTAGACCGATTGTTTGCTGACAAGCA	2340
	AACACCAATAAGCGAGCGTTAGCGAGCATGGACAAAAGCGCATCGCAGTTTGAAAGCGTA	2400
	GGCGTTAGCCGAAGCTGTTTTCGCTAAGCAAATCAAACAAGATAGCGCAAGCCGAGGTGC	2460
40	AGCCCCAAGAATTTGAATTATCCATGCGGTGTTTAGGGCGTTTTAGCGTGATCGCTTTAT	2520
	TACATGTTTTAAACAGCATGCTGTTTTTACATGTTTTACTCGCATGCGCGCGCTAGG	2580
	TATTGGTGGTTGGAATAGCCTAAATAACGCAGCTGTATGGTTTCTCATTTTTTCGGTGACA	2640
	ATGAATAAGGGGTAGTTCTTGCGAGTCATAAGTGTAGTTCTTGCGAGTCATAAGTGTAGT	2700
	TCTTGCGAGTCATAAGTGTAGTTCTTGCGAGTCATAAGTGTAGTTCTTTCACAATATCT	2760
45	ACACAATTCACAATCTCTAGCTACAATAACAGCGGTGGAAGATGCTTTACAGCGGTATT	2820
	TCCTATGTGCGGCAAAATTTGGAGCAATTAGCTTTAAAAGCTAGTGGGTTGGGAGTTTGT	2880
	AGCGGGTATGCACTCCGTTAGGAGGCACACCATGAAAGCATTTTTGATAGTAGTGATTTT	2940
	AGTGGTAATCTTGACACAGCCACTATATTAACCTTAGCGTTTTAATAACCCCTATAAG	3000
	TCCGCCAAGACTTCTTAAGGGTTTCACTCCTGTTATTATATCGTCTTTTGAAAAATAAGC	3060
50	ATTAAAAGGCGCTTAAATGCCCATGAATACGAATTTGAACAGCTTAGAAAACAGAATT	3120
	GGAATTACGAAAATTATTAGAAGAATTAGAAACGCTCCCACAAACCCACAAATTAAACT	3180
	GCAAAAACAAAAATACAACTTACATAGACAAGATAACACCAAGTATTTTGAGCGGTTT	3240
	TGATCAAAAATTCAAAGAAATTATAGAAAATCTATCAAATGAATTTGAAAAAGAAAAATC	3300
	CACACCACTCAAAGAGCCACAAACCCACACCATGCAAAGATTTAGTGGTTAGCAC	3360
55	CCCTAAAGATAACACCTATACCACCTACCACAATAACGCTAATAAGGTCAATCTAGGGAA	3420
	ATTGAGCGAAAGGGAAGCCAATCT	3444

An additional nucleotide sequence that was cloned is provided at SEQ ID NO: 4, which includes a 135 bp segment

- 46 -

that encodes a peptide of 45 amino acids (SEQ ID NO: 5). This smaller 45 amino acid peptide is an immunogenic polypeptide of the Hepatitis C virus (HCV) core antigen. The nucleic acid sequence encoding the 45 amino acid peptide is shown below with the indicated 135 nucleotides underscored. (SEQ ID NO: 5)

10 CATGAGCAGC AATCCTAAAC CTCAAAGAAA AACCAAACGT AACACCAACC GTCGCCCACA
 GGACGTCAAG TTCCCGGGTG GCGGTCAGAT CGTTGGTGGA GTTTACTTGT TGCCCGCGCAG
 GGGCCCTAGA TTGGGTGTGC GCGCGACGAG GAAGACTTCC GAGCGGTCCG AACCTCGAGG
 TAGACGTCAG CCTATCCCCA AGGCACGTCG GCCCCGAGGC AGGACCTGGG CTCAGCCCGG
 GTACCCTTGG CCCCCTCTATG GCAATGAGGG TTGCGGGTGG GCGGGATGGC TCCTGTCTCC
 CCGTGGCTCT CGGCCTAGCT GGGGCCCCAC AGACCCCCGG CGTAGGTCGC GCAATTTGGG
 15 TAAGGTCATC GATACCCTTA CGTGCGGCTT CGCCGACCTC ATGGGGTACA TACCGCTCGT
 CGGCGCCCCT CTTGGAGGCG CTGCCAGGGC CCTGGCGCAT GGCGTCCGGG TTCTGGAAGA
 CGGCGTGAAC TATGCAACAG GGAACCTTCC TGGTTGCTCT TTCTCTATCT TCCTTCTGGC
 CCTGCTCTCT TGCCTGACTG TGCCCCGCTT AGCCTACCAA
SEQ ID NO: 4

20 AATCCTAAAC CTCAAAGAAA AACCAAACGT AACACCAACC GTCGCCCACA GGACGTCAAG
TTCCCGGGTG GCGGTCAGAT CGTTGGTGGA GTTTACTTGT TGCCCGCGCAG GGGCCCTAGA
TTGGGTGTGC GCGCG
SEQ ID NO: 5

25 The nucleic acid of SEQ ID NO: 4 was cloned into the *hopE* gene (SEQ ID NO: 6, shown below), of *H. pylori* 26695 at nt504 of SEQ ID NO: 4 (noted in bold/underscore; corresponding to amino acid residue 168 of the protein product) so that the expressed product would be located as part of the surface exposed loop of the HopE gene product. This construct, designated as vector pTMI03-8 (Figure 6) was expressed on the surface of *E. coli*.

35 ATGCCATAGC ATTTTTATCC ATAAGATTAG CGGATCCTAC CTGACGCTTT
 TTATCGCAAC TCTCTACTGT TTCTCCATAC CCGTTTTTTT GGCTAACAGG
 AGGAATTAAC **C**

40 1 ATGGAATTTA TGAAAAAGTT TGTAGCTTTA GGGCTTCTAT CCGCAGTTTT
 51 AAGCTCTTCG TTGTTAGCCG AAGGTGATGG TGTATTATATA GGGACTAATT
 101 ATCAGCTTGG ACAAGCCCGT TTGAATAGTA ATATTTATAA TACAGGGGAT
 151 TGCACAGGGA GTGTTGTAGG TTGCCCCCA GGTCTTACCG CTAATAAGCA
 201 TAATCCAGGA GGCACCAATA TCAATTGGCA TGCTAAATAC GCTAATGGGG
 251 CTTTGAATGG TCTTGGGTTG AATGTGGGTT ATAAGAAGTT CTTCCAGTTC
 45 301 AAGTCTTTTG ATATGACAAG CAAGTGGTTT GGTTTTAGAG TGTATGGGCT
 351 TTTTGATTAT GGGCATGCCA CTTTAGGCAA GCAAGTTTAT GCACCTAATA
 401 AAATCCAGTT GGATATGGTC TCTTGGGGTG TGGGGAGCGA TTTGTTAGCT
 451 GATATTATTG ATAACGATAA CGCTTCTTTT GGTATTTTGG GTGGGGTCGC
 501 TATCGGCGGT AACACTTGGA AAAGCTCAGC GGCAAACTAT TGGAAAGAGC
 50 AAATCATTGA AGCTAAGGGT CCTGATGTTT GTACCCCTAC TTATTGTAAC
 601 CCTAACGCTC CTTATAGCAC CAAAACCTCA ACCGTCGCTT TTCAGGTATG
 651 GTTGAATTTT GGGGTGAGAG CCAATATTTA CAAGCATAAT GGCGTAGAGT
 701 TTGGCGTGAG AGTGCCGCTA CTCATCAACA AGTTTTTGAG TGCGGGTCTT
 751 AACGCTACTA ATCTTTATTA CCATTTGAAA CGGGATTATT CGCTTTATTT
 55 801 AGGGTATAAC TACACTTTTT

- 47 -

CTCGAGATCT GCAGCTGGTA CGATATGGGA ATTCGAAGCT TTCTAGAACA

5

AAAACATCATC TCAGAAGAGG ATCTGAATAG CGCCGTCGAC CATCATCATC
ATCATTGAGT TTAACGGTCT CCAGCTTGGC TGTTTTGGCG GATGAGAGAA

GATTTTCAGC CTGATACAGA TTAAATC

(SEQ ID NO:6)

Briefly, one method of accomplishing the isolation of *hopE* gene is amplification from *H. pylori* 22695 by using *Taq* DNA polymerase. The upstream primer
10 5'AAGGATCCGATAGGAATGTAAAGGAATGG-3' (SEQ ID NO:7) containing a *Bam*HI site and the downstream primer
5'CCGAATTCTAAAGGCATGAACGCTTGCA-3' (SEQ ID NO:8) containing a *Eco*RI site can be constructed by using a DNA synthesizer,
15 such as the Perkin-Elmer Applied Biosystems, Inc. model 332 (ABI; Mississauga, Ontario, Canada). The resulting PCR fragment can be blunt-end cloned into the *Eco*RV site in pBluescript II KS(+) in the same orientation as the *lac* promoter.

20

PCR primers can then be designed to insert two unique restriction enzyme sites into the *hopE* gene for insertion of the 135bp immunogenic coding sequence from the Hepatitis C virus (HCV) core antigen. The PCR amplification using *Taq*
25 DNA polymerase can be performed using a touchdown amplification procedure as follows. The PCR thermocycler is programmed for an initial denaturation step of 96°C for 4 min, followed by 18 cycles at an initial annealing temperature of 65°C (for 90 s), which is decreased by 0.5°C
30 for each successive cycle, an extension step at 72°C for 6 min, and denaturation at 96°C for 1 min. Subsequent to completion of the first 18 cycles, an additional 14 amplification cycles can be performed by using 72°C extension and 96°C denaturation steps with a constant 55°C
35 annealing temperature. The resulting amplicon is then purified by column, precipitated with ethanol, and made blunt by digestion with the Klenow fragment of DNA polymerase. The PCR products can be digested with restriction enzyme to remove the template DNA, and

- 48 -

religated into an appropriate vector such as pTM103.8 under the control of the arabinose inducible promoter, and transformed into *E. coli* JM105.

- 5 Recombinant clones can be identified by using oligonucleotide primer 5'-AGATCTAAGGACGTC-3' (SEQ ID NO:9) plus the reverse sequencing primer in PCR amplification reactions. Identified clones can be sequenced to verify that the inserted restriction endonuclease sites are in
10 frame and that no errors had been introduced into the *hopE* gene. The 135bp immunogenic coding sequence from the Hepatitis C virus (HCV) core antigen can then be inserted using standard techniques.
- 15 Once vector pTMI03-8 had been created it was transformed into *E. coli*, which was grown at 37°C. Cells were then harvested and the expression of the HCV insert was confirmed by Western Blot.
- 20 Briefly, a procedure for this is as follows. Cells are harvested from about 20 plates and resuspended in 20% sucrose with 50mg of DNase I (Boehringer Mannheim) in 10mM Tris-HCl (pH8.0). The cells are then disrupted with a French pressure cell at 15,000 lb/in². Broken cells are
25 overlaid on a sucrose step gradient of 1ml of 70% and 6ml of 70% sucrose in 10mM Tris-HCl (pH8.0). The outer membrane fraction is collected and pelleted at 150,000 × *g*, and the pellet is resuspended in 100µl of distilled water.
- 30 Alternatively, outer membranes from 500 ml of log-phase culture can be solubilized in 10mM Tris-HCl (pH 8.0)-3% *n*-octyl-polyoxyethylene incubated at 23°C for 1h and centrifuged for 30min at 173,000 × *g*. The pellet is resuspended in 10mM Tris-HCl-3% *n*-octyl-polyoxyethylene-5mM
35 EDTA (pH8.0), incubated at 23°C for 1h, and centrifuged for 30min at 173,000 × *g*, and the supernatant collected. A Western immunoblot indicated the presence of HCV/*hopE* in

- 49 -

the supernatant of the second solubilization step. The supernatant containing HCV/*hopE* is mixed with an equal volume of 0.125 M Tris-HCl (pH 6.8), 4% (wt/vol) sodium dodecyl sulfate (SDS), and 20% (vol/vol) glycerol and
5 subjected to SDS-12% polyacrylamide gel electrophoresis (PAGE). If required the HCV/*hopE* band can be excised from an unstained portion of the gel and eluted overnight at 4°C into 10mM Tris-HCl (pH8.0), 1mM EDTA (pH8.0), and 100mM NaCl. The elution supernatant can then be run on an SDS-
10 PAGE gel to check for purity, and a Western immunoblot using standard techniques undertaken. For example, isolated outer membranes can be loaded at a concentration of 15µg/lane. Electrophoresis is then carried out by SDS-PAGE on a discontinuous 12% polyacrylamide gel. Proteins are
15 then stained with Coomassie brilliant blue. For Western immunoblotting, unstained gels can be electroblotted onto Immobilon-P membranes (Millipore, Bedford, Mass.). After blocking for 2h at 23°C with 3% bovine serum albumin (BSA; Boehringer Mannheim)-0.1% Tween 20 (Sigma) in phosphate-
20 buffered saline (PBS), the membranes are then incubated with a 1/10,000 dilution of anti-HCV rabbit antiserum in 1% BSA-0.05% Tween 20 in PBS for 1 h at 37°C. The membranes are then washed with PBS and incubated with a 1/5,000 dilution of an alkaline phosphatase-conjugated secondary
25 antibody (Bio-Rad, Richmond, Calif.) for 1h at 37°C. The bound antibodies are detected with 5-bromo-4-chloro-3-indolylphosphate (BCIP, Calbiochem, La Jolla, Calif.) and nitroblue tetrazolium (NBT, Sigma).

30 EXAMPLE 2 EXPRESSION VECTORS AND SELECTION OF ANTIGENS
 FOR STABLE EXPRESSION

Because *HopE* is a native protein of *H. pylori* and is tolerated by this organism and can thus form a construct
35 useful for expressing foreign antigens or other heterologous gene products in *H. pylori*. Other *H. pylori*/*E. coli* shuttle vectors that can be readily

- 50 -

developed as described above include, for example, vectors comprising two plasmid *ori* sites and markers which are suitable for each host. Markers might include genes for chloramphenicol/kanamycin resistance as well as promoters
5 that can be recognised by both the *E. coli* and *H. pylori* transcriptional systems.

The requirements for replication in *E. coli* can be achieved by using any number of known *E. coli* plasmids (e.g.
10 pBR322).

Constructed shuttle vectors can be tested for replication in both *E. coli* and *H. pylori* *in vitro* and compared to existing shuttle plasmids described in the literature.
15

The choice of antigen or other heterologous gene product to be expressed would ideally be one that is not toxic to *H. pylori* and *E. coli* and that is highly immunogenic (or possesses another desirable property) when delivered to a
20 selected site in a mammal. In the case of an expressed antigen for immunization of the mammal, such a site might be a mucosal site.

As described in Example 1, *hopE*/HCV core antigen fusion
25 protein can be expressed at the *E. coli* surface. The product of pTMI03-8 (Figure 3) is preferably targeted to the *H. pylori* outer membrane, so that it would display the HCV core antigen in a mucosal environment.

30 Tetanus toxoid (TT) has been studied extensively as an antigen in humans, and immune responses to it are well characterized. TT elicits good mucosal immune responses when administered orally or intranasally when displayed on the surface of bacterial spores (Duc & Cutting, 2003,
35 *Expert Opinion Biol Ther*, 3(8), 1263-70). The tetanus toxin C fragment can be fused to the *hopE* gene product as described above or engineered to contain a membrane anchor

- 51 -

and cell surface target sequence. The advantage of this system is the existence of well-characterized murine models for assessment of the effectiveness of the vaccination procedure, whereas; in the case of HCV where the known
5 murine models typically rely on immune-deficient mice. The gB protein of cytomegalovirus (CMV) has been shown to immunize mice against a lethal dose of an engineered *Vaccinia* virus expressing the gB antigen. Accordingly, shuttle vectors such as those described herein, comprising
10 gB antigen in place of an HCV antigen can be constructed using the protocol described in Example 1.

A number of promoters can be used in the shuttle vectors. Ideally the promoter used is inducible either by a natural
15 *in vivo* colonization mechanism of *Helicobacter* or by induction with an innocuous foodstuff or chemical that can be consumed. For example, the promoter from the *H. pylori* histidine kinase HP165 that is reported to be induced by acidic pH and may be a virulence factor related to gastric
20 mucosa colonization is one such promoter. The benefit of this promoter is that a construct can be made *in vitro* with the foreign antigen only becoming expressed when exposed to the acidic environment of the gastric mucosa.

25 Other promoters include the arabinose inducible promoter used in pTMI03.8 and FlaB sigma 54 promoter (Josenhans et al., 1998, *FEMS Microbial Lett.*, 161(2), 263-73), the extensively studied T7 promoter used in many constitutive and inducible *E. coli* systems and the *nir B* promoter of
30 *Salmonella* which is induced in anaerobic environments (Chatfield et al., 1992, *Biotechnology*, 10(8): 888-92). The ability of any of these promoters to function in *H. pylori* can be tested using the system developed by Angelini et al. (2004) (*Plasmid*, 51:101-107) that uses CAT and GFP
35 reporters as a readout of promoter activity in a *H. pylori* plasmid vector.

- 52 -

The target sites of expression will depend on the antigen or other gene product used. Initial studies focused on the *HopE* proteins and fusion polypeptides which target the expressed polypeptide (e.g., antigen) to the cell surface of *H. pylori*.

Plasmid stability is also very important and, while the use of antibiotic resistance genes as a selective determinant for plasmid maintenance is useful *in vitro*, is less practical *in vivo*. An alternative is a balanced-lethal system, for example, the *asd* gene that is used inactivated in *Salmonella*. The *asd* gene, which exists natively in *H. pylori*, encodes aspartate- β -semialdehyde dehydrogenase (an enzyme in biosynthetic pathway for diaminopimelic acid (DAP), an essential component of the cell wall peptidoglycan of gram-negative bacteria). In the absence of DAP, *asd* mutants undergo lysis. Since DAP is not present in mammalian tissues, this balanced-lethal system imposes a requirement for all living *H. pylori* to carry the recombinant *asd* gene-containing plasmid.

In order to use the *asd* gene system the genomic copy of the *asd* gene is inactivated using standard gene knockout protocols. This strain of *H. pylori* will then only grow with the supply of DAP or with a plasmid that contains the *asd* gene.

Other systems that can be used for similar purposes include *E. coli* enterotoxin or Cholera toxin (CT) as mucosal adjuvants. Adjuvants can also be used to boost the mucosal immune response. Two such adjuvants are CT and *E. coli* enterotoxin (LT) wherein expressed antigens are fused to the LTB and CTB mutants that maintain their strong mucosal adjuvant properties but have dramatically reduced toxicity.

EXAMPLE 3 VIRULENCE, LD₅₀

As described in Examples 1 and 2, *Helicobacter*-based vectors such as pHP3 and pHP1 are capable of providing
5 protection against infection in a mammal, such as a mouse or human. In the present example, a murine model is used to demonstrate the utility of using the *Helicobacter*-based systems to provide delivery of a pharmacologically active molecule of interest to a mammal, including a human. The
10 murine model is employed to demonstrate the activity of a transgenic strain of *H. pylori* to elicit a serological response to an expressed surface antigen *in vivo*.

Mice are infected with wild-type *H. pylori*, while other
15 mice are inoculated by gavage with temperature-sensitive *H. pylori* as described in Example 2. Sera from both control and test animals are assayed for antibody and gastric histology are performed on sacrificed animals in accordance with the schedule shown in Table 1. A mouse urea breath
20 test can also be used.

As shown in Table 2, a 50% decrease in virulence (from 75% to 40%) was observed. Specific antibody titre increased 4 fold above baseline, indicating a serological response.
25 Serum samples were taken at baseline, 12, 24, 48 weeks. At these times 10, 10 and 20 animals were sacrificed and gastric histology performed.

- 54 -

TABLE 1IN VIVO STUDY

Week	Mice* Remaining	Mice Used	Serology	Histology	Adjistment
0	50	0	40	0	0
12	50	10	40	10	109.2
24	40	10	30	10	218.4
48	30	30	20	20	1310.4
49	0				
Totals		50	130	40	

5 *C57BL/6J females, 40 test and 10 control

TABLE 2POWER CALCULATION VIRULENCE STUDY

At $\alpha=0.05$ and Power=80% and various Infection Rates of *H. pylori*

Normal=75% vs TSHP=50% n=58 mice in each group

Normal=75% vs TSHP=40% n=30 mice in each group

Sample size of 20 will only give you 37% Power: to detect a difference (75% vs 50%) and 62% Power: to detect a difference (75% vs 40%).

In order to use a sample size of 20 you would need the infection rate in TSHP to be at least 32% or less (with 80% Power and $\alpha=0.05$)

EXAMPLE 4 COMPARISON OF VIRULENCE AND ANTIGENICITY OF
SEVERAL TEMPERATURE SENSITIVE *H. pylori*
STRAINS

- 5 In order detect a major change in virulence related to
expression/modification of an outer membrane protein, mice
are inoculated with temperature-sensitive *H. pylori* as
described in Example 3. An equal sized control group of
mice were infected with a wild type *H. pylori* strain. Non-
10 invasive means were used to determine presence or absence
of *H. pylori*. Mice were bled at 3 and 6 months for antibody
estimation. At sacrifice, histology was performed to assay
gastritis and confirm colonization. Table 3 shows the
serology and histology of the mice used.

- 57 -

TABLE 3VIRULENCE AND SEROLOGY COMPARISON STUDY

Week	Mice Remaining	Mice Used	Serology	Histology	Adjistment
0	180		180		
24	180		180		
48	180	180	180	180	7862.4
49	0				
Totals		180	540	180	

- 58 -

EXAMPLE 5 LD₅₀ STUDY TO EVALUATE TEMPERATURE SENSITIVE
H. pylori VACCINE EFFICACY FOR A
PNEUMOCOCCAL ANTIGEN

5 In order to demonstrate the *Helicobacter*-based vaccine
protection effect from a standard pathogen (pneumococcus),
mice were inoculated with temperature sensitive *H. pylori*
by gavage. An equal sized control group was infected with
the wild type *H. pylori* strain. Non-invasive means are
10 used to determine presence or absence of *H. pylori* as
described in Example 4. At 6 months post infection, all
mice were given intraperitoneal challenge with 10 times the
LD₅₀ of live virulent pneumococci type 4 (~20 CFU/mouse),
as per the method of Aaberge et al. (1995, *Microb. Pathog.*
15 18:141-152).

As shown in Table 4, allowing for 75% lethality in the
controls, the study has a power of 0.8 to detect a 50%
decrease in mortality (75% vs 50%).

- 59 -

TABLE 4LD₅₀ TO PNEUMOCOCCUS FOR IMMUNIZED MICE

Week	Mice Remaining	Mice Used	Serology	Histology	Adjistment
0	60	0	0		
24	60		0		
25	60	60	0	0	1365
26	0				
Totals		60	0	0	

- 60 -

EXAMPLE 6 DETERMINATION OF *H. pylori* STATUS OF MICE:
BREATH TEST METHOD

5 In the present Example, the urea breath test used in humans
was adapted use in mice.

10 Ten mice were fed a diet devoid of urease (uncooked soy).
Mice were then administered 3.7 kBq ¹⁴C urea in 200µl
flavoured citrate by gavage and placed in air-filled 2L
plastic ziplock bags for 20 minutes. Mice were then
removed without exchanging the air within the bag. Hyamine,
0.1 mmol in ethanol, was then introduced and scintillant
was added to the hyamine solution and counted for 10 min or
up to a count of 1,000 dpm.

15

EXAMPLE 7 HUMAN STUDIES

20 To confirm virulence and antibody response in humans, a
strain of *H. pylori* like the "Baylor Strain" will be
employed and the following criteria will be adopted:

- 25 1. The infected individuals have no symptoms, no more
than mild histologic damage, and no evidence of infection
with hepatitis viruses or HIV.
- 30 2. The isolate is a single strain, *cagA* negative, and
sensitive to metronidazole, clarithromycin, tetracycline,
and amoxicillin.
- 35 3. Volunteers to receive a challenge are healthy with
normal gastric histology, no history of peptic ulcer, no
young children at home, no regular contact with young
children, and no allergies to the antibiotics that might be
required to treat the infection.

35

Challenge will consist of 40mg famotidine at bedtime
followed by administration of *H. pylori* in beef broth

orally in the morning. Subjects are contacted daily for 14 days. A ^{13}C -UBT is performed after 7 and 14 days and endoscopy with quantitative culture and histology after 2 weeks and 3 months. Antibiotics are used to eradicate the infection.

EXAMPLE 8 DEVELOPMENT OF EXTERNAL CHEMICAL MARKER FOR
THE DETECTION OF WILD TYPE AND/OR TSHP IN
VIVO

10

An example of a chemical marker that may be used for the detection of wild type or TSHP *in vivo* is sulfasalazine (SSN), the structure of which is shown in Figure 4. Studies in germ free mice and conventional rats have shown that intestinal bacteria are solely responsible for the diazo-bond reduction, resulting in the reductive catabolism of SSN and the release of sulfapyridine and 5-aminosalicylate. The enzyme(s) which catalyses this reaction is referred to as diazoreductase(s) (synonym azoreductase(s)).

Conventional rats given SSN excrete 5-aminosalicylate and sulfapyridine (and their respective conjugates) in urine and faeces, whereas germ-free rats show no evidence of SSN degradation.

Several bacterial species have been shown to have diazoreductases (AZR's). Preliminary bioinformatic studies have indicated that *H. pylori* may not contain the AZR gene. The presence of similar analogous sequences has also produced a negative result. Under these circumstances a transgenic strain of *H. pylori* (TSHP) which has a viable and functional azoreductase (azr + TSHP) can be used to assess the use of these markers.

Plasmid pTMI03-02 is digested by *EcoR* I and *Hind*III, and ligated with the Azoreductase (AZR) gene from *Bacillus subtilis* treated with *EcoR* I and *Hind*III, to generate a vector containing both *HopE* 168aa and AZR named pTMI03-azr.

- 62 -

This plasmid is transformed into *E. coli* to assess whether expression of *HopE* and the *B. subtilis* AZR occurs. pTMI02 when similarly treated with full-length hepatitis C core antigen (HCCA) demonstrated transport of *HopE*::HCCA to *E. coli* outer membrane employing western blots and anti-*HopE* Abs.

Mice (n =30) are infected with the azr + TSHP by gavage and once AZR expression *in vivo* to produce 5-aminosalicylate and sulfapyridine (and their respective conjugates) in urine and faeces is established human trials can begin.

EXAMPLE 9USE OF DIAGNEX BLUE AS A MARKER

The diagnostic agent "Diagnex Blue" consists of an ion exchange resin (Amberlite XE-96) conjugated with a dye (Azure-A). This test relies on the fact that the resin-dye combination disassociates at pH less than 2.5 after which the dye is absorbed and appears in the urine. Persons without dye in the urine are achlorhydric. This principle is shown in Figure 5.

The same principle can be used to test for *H. pylori*. For example, a dye-resin combination which disassociates at pH > 7.0 could detect urease if the resin was given with urea. This would produce a pH > 7.0 in the mucus layer where *H. pylori* resides thus releasing the dye.

Mice (n =30) are inoculated with a wild type *H. pylori* strain while germ-free mice (n =30) are used as controls (Pilot study). After an optimal period allowing for the *H. pylori* to establish an active infection, the test group and the controls are introduced with a predetermined quantity of the resin-dye complex by gavage. This will be followed by a urea solution. (Range 0.01M to 0.5M). The mice are kept in metabolic cages and the excretion of the azure dye are monitored and quantified. Different ratios of the

- 63 -

resin and urea concentrations are tested to verify the optimal combinations to be used.

EXAMPLE 10DELIVERY FORMULATIONS

5

For administration by aerosol, the present invention can be delivered in the form of aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. The formulation would be prepared as a powder for administration by inhalation. Administration by inhalation can also be carried out by atomizing solutions or suspensions which contain the compositions according to the invention.

The compositions according to the invention may also be formulated in a liquid for oral digestion for administration to a subject as an intravenous preparation.

20

All of the various preparations of the invention may be prepared by procedures familiar to those skilled in the art, if appropriate using further suitable pharmaceutical auxiliaries. Compositions according to the invention advantageously contain the species of *Helicobacter*, alone or in combination with other desired ingredients.

25

Any of the above individual or combination of *Helicobacter* formulations may be included in a pharmaceutical composition comprising the pharmaceutically acceptable composition according to the present invention. The pharmaceutical compositions as described herein may be in solid (e.g. powder, particles, granules, sachets, tablets, capsules etc.), semi-solid (gels, pastes etc.) or liquid (solutions, dispersions, suspensions, emulsions, mixtures etc) form and adapted for administration via e.g. the gastrointestinal tract and gastric mucosa. The

30

35

- 64 -

pharmaceutical compositions may thus be in powder or particulate form adapted to be dispersed in an aqueous medium before use.

- 5 A pharmaceutical composition in liquid form may be in the form of a dispersion comprising the *Helicobacter* composition and an electrolyte solution such as, e.g. a composition that is adapted to physiological conditions e.g. a physiologically acceptable solution.

10

A pharmaceutical composition according to the invention may further comprise another therapeutically, prophylactically and/or diagnostically active substance.

- 15 In another aspect, the invention relates to a pharmaceutical kit comprising a first and a second container, the first container comprising a recombinant *Helicobacter* composition comprising the plasmid and/or plasmid vector according to the invention and the second
20 container comprising a dispersion medium for the *Helicobacter* composition, accompanied by instructions for administering and/or dosing the *Helicobacter* composition in the dispersion medium before use.

- 25 The *Helicobacter* composition according to the present invention contained in the kit may be in powder or particulate form.

- A pharmaceutical kit according to the present invention may
30 include instructions with recommendations for the time period during which the *Helicobacter* composition should be administered after dispersion in the dispersion medium.

35

- 65 -

EXAMPLE 11IMMUNE MODULATION WITH *Helicobacter* -
VACCINE PREPARATION

The TH1 response (T-helper cell type 1) is a cell mediated
5 response. Over activity of this is a presumed cause of
diseases such as rheumatoid arthritis (RA) and lupus. In
contrast, TH-2 is an antibody type serological response
characteristic of vaccines. The present example provides a
technique to obtain a TH-2 type response in an animal when
10 treated with a *Helicobacter*-based vaccine treatment
preparation according to the present invention.

Use of the *Helicobacter* vectors and vector plasmid systems
as described herein may be used to invoke antibody response
15 in an animal. By way of example, a system employing a gene
expression cassette in a construct that provided for the
transformation of the bacterium, *Clostridium*, and the
subsequent secretion of a protein (S-layer protein) from
the surface of the transformed *Clostridium*, this resulting
20 in initiation of mucosal vaccination, is described in
WO0194599, which disclosure is hereby incorporated herein
in its entirety. These constructs may also include a
secretory leader sequence selected from ORF1, ORF3, ORF5-
7., ORF7 or ORF11.

25 In accordance with some embodiments of the vaccine, the
Helicobacter-based vectors and vector plasmids may comprise
a sequence encoding a bacterial surface layer protein. A
surface layer protein is defined herein as any molecule of
30 proteinaceous nature, including e.g., protein, glycol-
protein or lipoprotein occurring in the outer membrane of a
bacterium and capable of being exposed on the surface of
the bacterium. S-layer proteins may be continuously and
spontaneously produced in larger amounts than any other
35 class of protein in the cell.

- 66 -

A process for preparation of a recombinant cell preparation comprising a gram negative host cell, *Clostridium*, having the S-layer protein, is also provided in WO-97/28263. The process may be modified and followed in accord with the
5 procedures described herein to incorporate an S-protein as part of the *Helicobacter* constructs of the present invention.

Accordingly, in some of the vector and vector plasmid
10 constructs, a fusion protein is provided that comprises a *Helicobacter* sequence and a non-*Helicobacter* pharmacologically active molecule of interest. In order to enhance the immunogenicity of a vaccine employing the *Helicobacter* constructs of the present invention, the
15 *Helicobacter* sequence of the fusion protein may comprise a sequence encoding an S-layer protein. *Bacillus* constructs that include the S-layer protein as part of a fusion protein have been reported to express the S-layer protein at the *Bacillus* surface. (See WO-95/19371, describing
20 *Bacillus sphaericus*), thus enhancing the immunogenicity of the preparation.

Mucosal immunization is already provided against some diseases, including an oral polio vaccine and an oral
25 (drinkable) vaccine against cholera and diarrhea due to *E. coli* (an inactivated vaccine). In some embodiments, it is contemplated that the vaccines of the invention may thus comprise an inactivated vaccine.

30 The present invention contemplates a live vaccine, as such will provide a single-dose, long lasting vaccination, because the carrier organism, *Helicobacter*, will continue to produce the antigen, i.e., non-*Helicobacter* pharmacologically active molecule of interest, and boost
35 immunity *in vivo*. In addition, the vaccines will be administered in combination with an adjuvant. These adjuvants' comprise molecules such as aluminium hydroxide

- 67 -

or lipid vesicles that increase the exposure time for the vaccine by slowing its removal from site of administration. Adjuvants' also act by evoking production of immunomodulatory peptides called cytokines and chemokines (Brewer et al. 1997, *J. Cytokines Cell Mol. Ther.*, 4:223-246). Thus, the present vaccines may comprise cytokine adjuvant to enhance immune response. The transformed *Helicobacter* or *E. coli* bacterium, when administered orally to a mammal such as a human or animal, will provide for the gastro intestinal colonization, production and presentation of the desired polypeptide through the gastric wall, which is the natural site of colonization. The gastro intestinal tract is surrounded by an immense immune apparatus specialized in mounting immune response of various types. Gastro intestinal colonization by recombinant *Helicobacter* vaccine or peptide producer strain thus enables a much longer immune stimulus than traditional vaccination. Additionally, antigen can be presented preferentially to the gut wall or the lumen.

20

EXAMPLE 12 *Helicobacter* AND USES THEREOF AS AN APPETITE SUPPRESSANT

The present example is provided to demonstrate the utility of the present invention as a method for employing *Helicobacter* in preparations and treatment regimens that provide for appetite suppression. In particular, delivery to the gut mucosa of a construct that comprises attenuated *Helicobacter* together with a non-*Helicobacter* pharmacologically active molecule of interest that regulates the level of ghrelin (a hormone) or an agonist of ghrelin, is expected to provide an effective means for providing suppression of the gut-brain axis that regulates appetite and sanity.

35

Studies have suggested that ghrelin is an appetite stimulant, i.e., ghrelin increases food intake in mice

- 68 -

(Asakawa *et al.* 2003, *Gut*, 52(7):947-52). Ghrelin has also been reported to reduce fat utilization in adipose tissue in rodents (Tschop *et al.*, 2000, *Nature*, 407: 908-13), as well as to be involved in rat adipogenesis (Choi *et al.* 5 (2003), *Endocrinology*, 144 (3):751-9). Ghrelin has also been reported to be a hunger signal, prompting the subject to eat when nutrition availability is low.

Ghrelin, an endogenous ligand for the growth hormone 10 secretagogue receptor (GHS-R), stimulates growth hormone (GH) release from cultured pituitary cells in a dose-dependent manner, and is produced and secreted from the A-like cells found mainly in the oxyntic glands of the gastric fundus. Ghrelin is now known to play a role in not 15 only GH release, but also in controlling the appetite and body weight.

Both parenterally and intracerebro-ventricularly administered ghrelin have been shown to stimulate food 20 intake and increase the body weight of mice and rats with free access to food, even those animals with GH deficiency. The control of appetite and body weight may be independent of GH release.

25 Ghrelin, a 28-amino-acid peptide, is activated when its third serine residue is acylated by n-octanoic acid, and GHS-R is responsive to the first four or five residues including the octanoylated serine residue of the whole ghrelin peptide. GHS-R has been shown to be present in the 30 pituitary, hypothalamus, adrenal glands, thyroid, pancreas, myocardium, spleen and testes. Ghrelin stimulates the expression of both NPY and AGRP mRNA in the hypothalamus. The central orexigenic effect of ghrelin is mediated by the NPY/AGRP-expressing neurons in the hypothalamus. Ghrelin 35 has also been reported to suppress vagal afferent activity. The peripheral orexigenic effect of ghrelin may be mediated, at least in part, by its suppressive effect on

- 69 -

the vagal afferent activity. IL-1 β is a pro-inflammatory cytokine that mediates the cachectic process by stimulating the expression and release of leptin, and/or by mimicking the effect on the hypothalamus of excessive negative-
5 feedback signalling from leptin.

It is proposed that antagonists to ghrelin if provided to the animal at the gut mucosa will reduce food intake by an animal and reduce body weight gain.

10

EXAMPLE 13 CELL WASTING ATTENDANT CANCER AND AIDS

The present example demonstrates the utility of the present invention for use as a preparation that will prevent or
15 inhibit cell wasting, particularly cell wasting associated with the diseased states of AIDS and cancer.

Cachexia is a condition characterized by wasting, emaciation, feebleness and inanition. It was recently
20 reported that the levels of both ghrelin peptide and ghrelin mRNA in the stomach were up-regulated in a mouse model of cancer cachexia. In cachectic mice with increased plasma levels of IL-1 β , the plasma concentrations of ghrelin also increased with the progression of cachexia.
25 This result suggests that a close relationship might exist between the ghrelin dynamics and the cachectic process mediated by IL-1. IL-1 β is an anorexigenic substance, just like CCK, leptin, gastrin-related protein and bombesin, and antagonizes the actions of ghrelin.

30

Asakawa *et al.* reported that parenterally administered IL-1 β decreased NPY mRNA expression in the hypothalamus and pre-proghrelin mRNA expression in the stomach, and that intraperitoneally administered ghrelin inhibited the
35 severity of IL-1 β -induced anorexia. *H. pylori* infection is known to be a major pathogenetic factor in the development of gastritis, peptic ulcer disease and gastric malignancy.

- 70 -

Attachment of *H. pylori* to the gastric mucosa induces inflammation, which is associated with the release of various cytokines, including IL-1 β .

5 It has been observed clinically that *H. pylori* eradication is often followed by improvement of some nutritional parameters, such as the body weight and the serum levels of total cholesterol, total protein and albumin. *H. pylori* infection has been reported to be capable of modifying the
10 plasma and gastric ghrelin dynamics in Mongolian gerbils. In humans, however, *H. pylori* infection has been reported not to be associated with any changes of the plasma ghrelin levels, although eradication of *H. pylori* has been shown by some to be associated with increases of the plasma ghrelin
15 levels.

It is proposed that *H. pylori* may be used as a carrier to provide amylin to a patient in need thereof, by, for example, acting to provide secretion of ghrelin to the
20 gastric mucosa. In some embodiments, the *Helicobacter* vector will be constructed that include a sequence encoding amylin, amylin analogs, and/or derivatives thereof, and amylin agonists (including calcitonins, calcitonin gene-related peptides, and analogs therefore), so as to decrease
25 ghrelin levels.

Amylin antagonists can increase ghrelin levels. Modulation of the effective levels of amylin, with amylin, amylin agonists, or other like compounds that decrease the
30 effective level of amylin, may inhibit, or stimulate in the case of antagonists and antibodies, ghrelin secretion. Hence, some embodiments of the method are directed to modulating endogenous levels of ghrelin by increasing the effective level of amylin or amylin agonists in the body,
35 by direct or indirect means, or by decreasing the effective level of amylin using amylin antagonists or inhibiting amylin production.

EXAMPLE 14 TREATMENT OF GAUCHERS DISEASE

The present example demonstrates the utility of the
5 invention for use as a treatment for a disease resulting
from an enzyme deficiency, such as Gaucher's disease.
Gaucher's disease is the most common lysosomal storage
disorder in humans, and results from a deficiency of the
enzyme, glucocerebrosidase (GC). (Nolta et al., (1992), *J.*
10 *Clin. Invest.* 90 (2):342-348).

Enzyme replacement therapy is provided with a *Helicobacter*
vaccine construct that comprises a sequence encoding
chemical chaperones. (Sawker et al., (2002), *PNAS USA*
15 99(24): 15428-15433) or glucocerebrosidase. An enhanced
level of functional β -glycosidase (β -Glu,
glucocerebrosidase) may thus be obtained. In particular, a
sequence encoding the chemical chaperone deoxynojirimycin
(NN-DNJ) may be used in the *H. pylori* construct and
20 administered to the patient orally or intragastrically.

As part of yet another embodiment, a *Helicobacter*-based
construct is provided comprising a vector having a non-
Helicobacter pharmacologically active molecule of interest,
25 in this case, encoding glucocerebrosidase (GC). Retroviral
mediated transfer of glucocerebrosidase cultured Gaucher
bone marrow is described as one approach for treating
Gauchers disease in Nolta et al. (1992). However, this
approach is extremely invasive. Alternative enzyme
30 replacement therapy employing the *Helicobacter*-based
constructs of the present invention that include a sequence
encoding for the deficient enzyme, glucocerebrosidase,
provides a much more attractive and less expensive
alternative to such a therapy.

EXAMPLE 16

The present example is presented to demonstrate the utility of the present invention to provide a useful preparation
5 that is suitable for treating and or inhibiting a bacterial induced malignancy, such as lymphoma, particularly MALT lymphoma, using a vaccination preparation comprising the *Helicobacter* vector and/or plasmid vectors as described herein.

10

Sutton et al. (2004) (*Vaccine*, 22 (20): 2541-6) report protection against a bacteria-induced malignancy, specifically primary gastric MALT lymphoma, as a result of vaccination/immunization of an animal against *Helicobacter*
15 *felis*. Therefore, the *H. pylori* constructs of the present disclosure that include a vector and/or plasmid vector suitable for providing an immunizing preparation against other than *H. felis* may also be used to provide vaccination protection against a bacterial-induced malignancy, and in
20 particular, against primary gastric MALT lymphoma.

All documents, patents, journal articles and other materials cited in the present application are hereby incorporated by reference.

25

Although the present invention has been fully described in conjunction with several embodiments thereof with reference to the accompanying drawings, it is to be understood that various changes and modifications may be apparent to those
30 skilled in the art. Such changes and modifications are to be understood as included within the scope of the present invention as defined by the appended claims, unless they depart therefrom.

BIBLIOGRAPHY

The following references are specifically incorporated herein by reference.

- 5
1. U.S. 6,570,004- Blaser et al (2003).
 2. U.S. 6,680,179 Collins et al.
 3. U.S. 6,383,496 - Curtiss et al (2002).
 4. U.S. 6,150,170 - Powell et al. (2000).
 - 10 5. U.S. 6,410,012 - Seizmore et al. (2002).
 6. U.S. 6,550,419 - Hone et al. (2002).
 7. U.S. 6,531,313 - Goudsmit et al. (2003).
 8. U.S. 6,682,729 - Powell et al (2004).
 9. U.S. Patent Publication 2005/0075298A1 - Chen et al.
 - 15 (2005).
 10. U.S. Patent Publication 2002/0176848A1 - Seizmore et al (2002).
 11. U.S. Patent Publication 2005/0096288 A1 - Guevara et al. (2005).
 - 20 12. U.S. Patent Publication 2004/0236072 A1 - Olmsted et al. (2004).
 13. U.S. Patent Publication 2004/0203039 A1 - Hensel et al. (2004).
 14. U.S. Patent Publication 2004/0005325 A1 - Kusters et al. (2004).
 - 25 15. U.S. Patent Publication 2002/0032152 A1 - Torossian (2002).
 16. U.S. Patent Publication 2003/0170264 A1 - Turner et al. (2003).
 - 30 17. U.S. Patent Publication 2003 0204068- Blassec et al (2003).
 18. U.S. Patent Publication 2002/0161192 A1- Meyer et al (2002).
 19. WO 96/33274 - Covacci et al. (1996).
 - 35 20. WO 99/21959 - Ellis et al. (1999).
 21. WO 01/94599 - Burman et al. (2001).
 22. WO 2005 021026 - Baron, et al. (2005).

23. Graham et al (2002), *Gastroenterology*, 123:1637-1648.
24. Liu et al (2005), *World Journal Gastroenterology*, 11(14): 2154-2156.
25. Conway, BR (2005), *Curr. Pharm. Res.*, 11(6) 775-90.
- 5 26. Sawker et al (2002), *PNAS USA*, 99(24): 15428-15433.
27. Sutton, P et al (2004), *Vaccine*, 22(20): 2541-6.
28. Kang et al (2005), *World Journal Gastroenterology*, 11(3): 454-456.
29. Moschos et al (2004), *Immunology and Cell Biology*, 82(6): 628-637.
- 10 30. Reddy et al (2004), *International Journal Antimicrob. Agents*, 24(6): 536-47.
31. Bai et al. (2003), *Sheng Wugong Cheng Xu Bao*, 19(4):433-8.
- 15 32. Nolta et al. (1992), *Journal of Clin. Invest*, 90(2): 342-348.
33. Shi et al. (2005), *Helicobacter*, 10(1): 71-9.
34. Deml et al. (2005), *Infection Immunity*, 73(8): 4732-42.
- 20 35. Cosima et al. (2005), *Trends in Immunology*, 26(4): 199-207.
36. Velin et al. (2005), *Gastroenterology*, 129(1): 142-155.
37. Kong et al. (2000), *Nucleic Acids Research*, 28(17) 3216-3223.
- 25 38. Mao et al. (2003), *World Journal of Gastroenterology*, 9(7): 1529-1536.
39. Chu et al. (2005), *World Journal of Gastroenterology*, 11(23): 3518-22.
- 30 40. Kathy Parton. (2000), *Institute of Veterinary, Animal and Biomedical Sciences*, "*Helicobacter mustelae* as vector for biological control in stoats" *Wildlife Health and Conservation Research Program; Landcare Research (Funding Body)*.
- 35 41. Forrester N.T., Parton, K. (2000), *New Zealand Veterinary Journal*, 48: 65-69. Title, "*Isolation of Helicobacter mustelae* from ferrets in New Zealand".

- 75 -

42. Spranger et al. (2005), Br. Nutr., 93 (6):765-71.
43. Garbom et al. (2004), Infect Immun., 72(3): 1333-1340.
44. Tschop et al. (2000) Nature, 407:908-13.
45. Choi et al (2003), Edocrinology, 144 (3).
- 5 46. Remington's Pharmaceutical Sciences, 20th edition, Mack Publishing Company.
47. Jones et al (2005) Nat. Med. 11 (7): 786-90.

- 76 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A composition comprising a *Helicobacter* molecular construct having a *Helicobacter* sequence and a non-
5 *Helicobacter* sequence encoding a non-*Helicobacter* pharmacologically active molecule of interest.
2. A composition according to claim 1, wherein the
10 *Helicobacter* is *Helicobacter pylori*.
3. A composition according to claim 1 or claim 2, wherein the pharmacologically active molecule of interest is heterologous to the *Helicobacter pylori* species.
- 15 4. A composition according to any one of claims 1 to 3, further comprising a promoter sequence.
5. A composition according to any one of claims 1 to 4, wherein the pharmacologically active molecule of
20 interest is ghrelin, amylin or an analog thereof.
6. A composition according to any one of claims 2 to 5, wherein the *Helicobacter pylori* is attenuated
25 *Helicobacter pylori*.
7. A composition according to any one of claims 2 to 6, wherein the pharmacologically active molecule of
30 interest comprises a protein, peptide or nucleic acid molecule.
8. A vaccine comprising a composition according to claim 4 and a pharmacologically acceptable carrier solution.
- 35 9. A vector plasmid comprising a composition according to claim 4.

- 77 -

10. A composition comprising a vaccine, wherein said vaccine comprises a *Helicobacter* vector plasmid comprising:

- (a) a *Helicobacter* nucleotide sequence comprising a promoter sequence capable of controlling expression of a sequence encoding a non-*Helicobacter* pharmacologically active molecule of interest; and
- (b) a sequence encoding a non-*Helicobacter* pharmacologically active molecule of interest.

11. A composition according to claim 10, further comprising an adjuvant.

12. A composition according to claim 10 or claim 11, further described as suitable for delivery to the mucosa.

13. A composition according to any one of claims 10 to 12, wherein the non-*Helicobacter* pharmacologically active molecule of interest is a protein, peptide or nucleic acid molecule.

14. A composition according to any one of claims 10 to 12, wherein the non-*Helicobacter* pharmacologically active molecule of interest is ghrelin, amylin or an antagonist or analog thereof.

15. A method of preparing an immunogenic composition comprising a non-*Helicobacter* immunogen comprising:

- (a) providing a culture comprising *Helicobacter pylori* cells transformable with a plasmid vector;
- (b) introducing into said culture a plasmid vector comprising a promoter sequence and a sequence encoding a non-*Helicobacter* immunogen under suitable conditions to provide transformed *Helicobacter pylori* cells, and
- (c) selecting transformed cells that express said non-*Helicobacter* immunogen to provide an immunogenic composition,

- 78 -

wherein said promoter sequence is capable of controlling the expression of the sequence encoding the non-*Helicobacter* immunogen.

5 16. A method according to claim 15, wherein the promoter sequence comprises an inducible promoter.

17. A method according to claim 15 or claim 16, further comprising a non-antibiotic resistance gene marker.

10

18. A method according to claim 15, wherein the promoter is constitutively expressed in *Helicobacter pylori*.

15 19. A method according to claim 15, the promoter is an arabinose inducible promoter.

20. A method according to any one of claims 15 to 19, wherein the *Helicobacter pylori* is strain 26695.

20

21. A method according to any one of claims 15 to 20, wherein the plasmid vector is further defined as comprising an operon construct.

25 22. A method of treating an animal comprising the steps of:

(a) providing a composition according to any one of claims 10 to 14; and

(b) administering to the animal an effective
30 amount of said composition.

23. A method according to claim 22, wherein the animal is a human.

35 24. A method according to claim 22, wherein the treated animal has an enhanced immune response.

- 79 -

25. A method according to any one of claims 22 to 24, wherein the composition is administered at a mucosal surface of the animal.

5 26. A method according to any one of claims 22 to 25, wherein one or more effective amounts of the composition are administered to the animal.

27. A composition comprising a recombinant cell
10 comprising a sequence encoding a non-*Helicobacter pylori* pharmacologically active molecule of interest comprising:
(a) at least one non-*Helicobacter* sequence encoding a pharmacologically active molecule of interest; and
15 (b) a *Helicobacter* sequence having a promoter sequence capable of controlling expression of the non-*Helicobacter* sequence,
wherein the recombinant cell is capable of expressing said nucleic acid or causing the expression of the nucleic acid
20 molecule in a target cell.

28. A composition according to claim 27, wherein the nucleotide sequence encoding the recombinant cell further comprises a secretory signal polypeptide.

25 29. A composition according to claim 27, wherein said recombinant cell is a recombinant *Helicobacter pylori*.

30. A composition according to claim 29, wherein the
30 recombinant *Helicobacter pylori* is pTM103-8.

31. A composition comprising a recombinant *Helicobacter pylori* comprising:
(a) a sequence encoding a fusion protein
35 comprising a non-*Helicobacter* pharmacologically active molecule of interest and a *Helicobacter* sequence; and

- 80 -

(b) an inducible *Helicobacter pylori* promoter sequence.

32. A plasmid vector comprising a composition
5 according to claim 31.

33. A pharmaceutical composition comprising as an active ingredient the plasmid vector of claim 32, together with a pharmaceutically acceptable dilute, carrier,
10 adjuvant or combination thereof.


34. A method for preparing a recombinant *Helicobacter pylori* comprising:

(a) providing a culture of *Helicobacter pylori*;
15 and

(b) providing a plasmid vector according to claim 32 to the culture to provide recombinant *Helicobacter pylori*,
wherein said recombinant *Helicobacter pylori* are capable of
20 expressing or causing expression of said *non-Helicobacter* pharmaceutically active molecule in a target cell.

35. A composition according to claim 31, wherein said recombinant *Helicobacter* further comprises a second nucleic acid molecule encoding an immunomodulatory polypeptide,
25 wherein said recombinant *Helicobacter* is capable of expressing said second nucleic acid molecule in a target cell.

1/5

**Fig.1**

2/5




Fig.2

3/5





Fig.3

4/5



5/5




Fig. 5

SEQUENCE LISTING

<110> Marshall, Barry

<120> BACTERIAL DELIVERY SYSTEM

<130> FP22313

<140> AU 2004904564

<141> 2005-08-12

<150> AU 2004904564

<151> 2004-08-13

<150> US 60/602,859

<151> 2004-08-20

<160> 9

<170> PatentIn version 3.3

<210> 1

<211> 2796

<212> DNA

<213> Helicobacter pylori

<400> 1

```

gtcatgcgcg ttgtttttaa ttacatttta aacaacttgt tgttgttttt acatgtttta      60
ctcgcgcgcg cgcgcggtgag ggattggggg ttgcaacccc ctaaataacg aagctgtagg      120
gtttctcatt tttgtggtga aaatgaataa aacagaactt cttgccaaca ctaacagaac      180
ttcttgccaa cactaacaga acttcttgcc aacactaaca gaacttcttg ccaacactaa      240
cagaacttct ttatttttaa gttatgatta ttaacaattt ttagacataa taacagcgtg      300
tgaagatact tttgtagcgg tatttcctat gtgcggcaaa atttgagca attagcttga      360
cttggttgag ttagtgggtt ggaggataga gagggcgaca cctcgtagg aggtatcaat      420
gtgaaagtat ttgtcgtatt agttctagta ttagtaattc tcgcacaatt gctatattag      480
gcttattcgt ggtctaacc cttgtttatg ggggttggct cgttataagc atactgatac      540
gatcacactt attatacacc aaaagataag gagtatagag tggaatttga tcaatcagat      600
ttacaaaaag cgttgaaaat attagataca ctcccacaaa cccacaaaat tgagctacaa      660
aaacaagaaa taaaaaacg catcaacaaa ataacagaga caatcattaa agaattacta      720
tcaaagcatg aaatcaagaa agaagaacta gaaccactc taaccccaaa acccacacca      780
ctcaaagagc cacaacccac cccaacacca tgcaaagatt tagtggtag caccctaaa      840
gataaaacct aatatcacct accacaataa cgctaataag gtcaatctag ggaaattgag      900
cgaaagggaa gccaatcttt tattcgctat ttttcaaaa ctcaaagccc aagggaatac      960
cctcattcgt tttgaaccgc aagatttgaa acgcatgcta aacatagata tttctaata      1020
gcgcttatca gaagtcgtta ttaagctgtg ggatagcatt aaaaccgctg atttttggaa      1080
aattagcgaa accgaaactt caatcattca agaaaattac atgcttttta gtcggtgtaa      1140
aattgaattg aacaaaccga gtaaagattt gaagtattta gaaatccaac tcaacgataa      1200
ctatcaagac ttactcaaca atctgggcat gggtaatac acttctttca atctgttaga      1260

```

```

atttcaaaga gtgaggggta aatacgctaa aacgctctat cgcttgctca agcaatacaa 1320
aagcacaggg attttgagcg tggaatggac tcaattcagg gagcttttag acattccaaa 1380
agactacaaa atggaaaaca tcgatcaaaa agtcttaacc ccctctctca aagaactcag 1440
aaaaatctac ctttttgaac acttgagcta taaaaaagaa cgcaaaagcc attacaagcg 1500
caaagtaacc cacattgatt tttattttga gcaatttcct taaggcgaaa ataagaaaca 1560
aaacaaagcc gacaagcaac gcgctcaaag ggacatcaag cttgtagcat gggatattca 1620
caaccaaadc gctaaaagaa acgcaaaagc cactatggaa gctagggttc ttgaattgaa 1680
aactttgatc ggctatcagt tcaggaacaa tgacagtagg aacaaattaa agattgacaa 1740
caccactttt gaaagaatca aatgtattta catgtatctt aaccctaaaa ataagcataa 1800
cccccaaaaa ttccttgat ccaacaagac attcgcattg gaactactat atatcaatag 1860
atacagccta aaaaaaagac aacttgctag aagaatttaa ccccccaaaa tccaccctat 1920
caccaacgaa cctatcaagg aatttgcgaga atacatcggc aaaacgatta acatcaccaa 1980
cttcaatgtg gatcaatgcc atgaggggaat cagcaactac ctgacaatca ctaggatcgt 2040
gaactggacg taatcggatc tgtattttgg ccagatgtgg ataagcctgg gacttctcaa 2100
gcctttcatt gctaaagtga gaaaatttgg ggattgggtc aagaacacta cagggtgaaaa 2160
gacagatgca tgctgactaa actcatagaa aaactgaatc acgaaagaaa gaatgcaagc 2220
agaaaacaaa cacctaaaag aacaaggact agaaaaaatc tacactcaaa aagactacga 2280
gcagttaaaa gaacagcatt tgaaagaaat tgaagcactc aaaaaagaaa tccaaaaaac 2340
caagcaagaa acatacacgc aaccaaaaaga atgtagccat ttagcgcat ctttttagccc 2400
taattcattc tttcaatcaa aatccgacta attcatcggc taaacgctaa aaatcgctta 2460
aaacgaaaaa tacaaagcaa aaaacttcat tccctttta gtcgttaacc atttagccaa 2520
tctaactagt ttagcatcta aaggcgaatc tatcttgtgt tagacatcca accttaccaa 2580
aaccgcagag cgagcttaag agagattcaa gcggttttgc acgattgttt gctgccaaga 2640
aaaccaacaa gcgaagtaag gcgcatagac aaaagcgcat cgagtttga aagcgtaggc 2700
gtcagaagtg gtttgcgta gaatcaaca agatagcgca aacctggcgt taggctaaaa 2760
aaccctaaa aactaaaacc ccaaaatatg tagtgc 2796

```

```

<210> 2
<211> 2796
<212> DNA
<213> Helicobacter pylori

```

```

<400> 2
cagtacgcgc aacaaaaatt aatgtaaaat ttgttgaaca acaacaaaaa tgtacaaaat 60
gagcgtacgc gcgcgcactc cctaaccccc aacgttgggg gatttattgc ttcgacatcc 120
caaagagtaa aaacaccact ttactttatt ttgtcttgaa gaacggttgt gattgtcttg 180
aagaacgggt gtgattgtct tgaagaacgg ttgtgattgt cttgaagaac ggttgtgatt 240

```

gtcttgaaga aataaaat	ttt caatacta	aat aattgt	taaa aatctgt	tatt attgtc	gcac 300	
acttctatga aaacatc	gcc ataaagg	ata cacgcc	gttt taaacct	cgtaaatcga	act 360	
gaaccaactc aatcaccca	aa cctcctat	ct ctcccgt	gt ggagcaat	cc tccatag	tta 420	
cactttcata aacagcata	aa tcaagat	cat aatcatta	ag agcgtgt	taa cgatata	aatc 480	
cgaataagca ccagattgg	ggaacaa	atac cccaacc	ga gcaatatt	cg tatgact	atg 540	
ctagtgtgaa taatatgt	gg ttttctat	tc ctcata	tct accttaa	act agttag	tcta 600	
aatgtttttc gcaacttt	taaatctat	gt gaggg	gtttt ggggtgt	tta actcgat	gtt 660	
tttgttcttt atgtttt	ggc gtagttg	ttt tattgt	tctt gttag	taatt tctta	atgat 720	
agtttcgtac tttagtt	ctt tctttg	at cttggg	tgg attggg	gttt tgggtg	tggt 780	
gagtttctcg gtgtttg	gtg gggttg	tgtt acgttt	ctaa atcacca	atc gtgggg	atgt 840	
ctattttgga ttatagt	gga tgggtgt	tatt gcgatt	attc cagttag	atc ccttta	actc 900	
gctttccctt cggttag	aaa ataagc	gata aaaag	tttt gagttt	cggg ttccc	ttatg 960	
ggagtaagca aaactt	ggcg ttctaa	actt tgcgtac	gat ttgtat	ctat aaag	attact 1020	
cgcgaaatagt cttcag	caat aattcga	cac cctatc	gtaa ttttgg	cgc taaaa	acctt 1080	
ttaatcgctt tggcttt	gaa gttagta	agt tctttt	aatg tacgaaa	aat cagcca	catt 1140	
ttacttaac ttgtttg	gct catttt	ctaaa cttcata	aat ctttag	gtt agttg	ctatt 1200	
gatagtctcg aatgag	tgt tagaccc	gta cccagt	tatg tgaagaa	agt tagaca	atct 1260	
taaagtttct cactcccc	at ttatg	cgaatt ttg	cgaata gcgaac	gagt tcg	ttatg 1320	
ttcgtgtccc taaaact	cgc accttac	ctg agtta	agtc ctcgaaa	atc tgta	agg 1380	
tctgatgttt taccttt	gt agctag	ttt tcaga	aattgg gggag	agagt ttctt	gagtc 1440	
tttttagatg ggaaa	acttg tgaact	cgaat attttt	tctt gcgttt	tcgg taatg	ttcgc 1500	
gtttcattgg gtgta	actaa aaataa	aaact cgttaa	agga attccg	cttt tattc	tttgt 1560	
tttgtttcgg ctgttc	gttg cgcgag	ttt cctgt	agttc gaacat	cgtta ccct	ataag 1620	
gttggttttag cgattt	tctt tgctgt	tttc gtgata	cctt cgatcca	aaag aact	taact 1680	
ttgaaactag ccgat	agtca agtc	cctgtt actgt	catcc ttgttt	aat tcta	actgt 1740	
gtggtgaaaa cttt	cttagt ttacata	aat gtacata	gaa ttggg	atttt tattc	gtatt 1800	
gggggttttt aagga	acata ggttg	tcttg taagc	gtaac cttgat	gata tatag	ttatc 1860	
tatgtcggat tttttt	tctg ttgaac	gatc ttctta	aat ggggg	gtttt aggtg	gggata 1920	
gtggttgctt ggatag	tcc ttaa	acgtct tatgt	agccg ttttg	cta	at tgtag	tggt 1980
gaagttacac ctagt	tacgg tactcc	ctta gtcgtt	gatg gactgt	tagt gatc	cctag	ca 2040
cttgacctgc attag	cctag acataa	acca ggtct	acacc tattc	ggacc ctga	agag	tt 2100
cggaaagtaa cgattt	cact ctttta	aaacc ccta	accaag ttctt	gtgat	gtcc	acttt 2160
ctgtctacgt acgact	gatt tgagt	atctt tttg	acttag tgctt	tctt	cttac	gttcg 2220
tcttttgttt gtggat	ttt cctga	tcttttt	tag atgtg	agttt	ttctg	atgct 2280

cgtcaat ttt cttgtcgtaa actttcttta acttcgtgag ttttttcttt aggttttttg 2340
 gttcgttctt tgtatgtgcg ttggttttct tacatcggt aatcgcgtaa gaaaatcggg 2400
 attaagtaag aaagttagtt ttaggctgat taagtagccg atttgcgatt tttagcgaat 2460
 ttgtcttttt atgtttcgtt ttttgaagta aggggaaaat cagcaattgg taaatcggtt 2520
 agattgatca aatcgtagat ttccgcttag atagaacaca atctgtaggt tggaatgggt 2580
 ttggcgtctc gctcgaattc tctctaagtt cgccaaaacg tgctaacaaa cgacggttct 2640
 ttgtgttggt cgcttcattc cgcgtatctg ttttcgcgta gcgtcaaaact ttcgcatccg 2700
 cagtcttcac caaacgcaat cttagtttgt tctatcgctg ttggaccgca atccgatttt 2760
 ttggggattt ttgattttgg ggttttatac atcacg 2796

<210> 3
 <211> 3444
 <212> DNA
 <213> *Helicobacter pylori*

<400> 3
 tctacacaat taacaatctt tagctacaat aacagcgtgt gaagatgctt tcacagcggg 60
 atttcctatg tgcggcaaaa ttgggagcaa ttaacttgac ttgggtgggt tagtgggttg 120
 gaggatagag agggcgacac ctcgttagga ggtatcaatg tgaaagtatt tgctgtatta 180
 gttctagtat tagtaattct cgcacaattg ctatattagg cttatttggt gtctaacccc 240
 ttgtttatgg gggtagatc cttataagca tactgatacg atcacactta ttataacca 300
 aaagataagg agtatagagt ggaatttgat caattagaat cacaagatc agacttaca 360
 aaagtgttaa aagaattaga tacactccca aaaacccac aaattgagct acaaaaaaca 420
 gaaatacaaa accgcatcaa caaaataaca gacacaatca ttaaagaatt actatcaaaa 480
 catgaaatca aaaaagaaga actagaaccc actctaacc caaaaccac accaacaana 540
 gagccacaaa ccacccccac accatgcaaa aatttagtgg ttagcaccac taaagataaa 600
 acctatatca cctaccacaa taacgcta ataggatcaat tagggaaatt gagcgaaagg 660
 gaagccaatc ttttattcgc tttttttcaa aggcctaaag atcaaggga taccctcatt 720
 cgttttgaac cgcaagattt aaaacgcgtg atcatgggtc aatccaactt aaccaacagg 780
 caattattgc aagtcttaaa aaatttgctt gacaacatta gcggtgctaa tttttggatc 840
 aattagagag catgttgaaa atggcgaaat ctatgaagat cacactagct acatgctttt 900
 caacaattt gaaatccgca tccataagcc aacacaaact atagaatact tagatgtcca 960
 actcaatgat agctatcaat acttgctcaa caatctagga atgggcgggtc aatacacttc 1020
 tttcaatctc ttagaatttc aaagggtgag gggcaaatag tgagagcggt aaatttcccc 1080
 cccctattcc ccttaaaaag gacccttatt ccagggaatt tttggcccca atacaattag 1140
 ggccaaaaac ccggtccctt ccatggctta accaaccaa ttggggggatt ccaatttccc 1200
 ctggatggga ataaccacaag gctttttttg aaaattccac ctaccatttg gtccaaaatt 1260
 ggatggacaa ttccaaattc caaatcttct tttccaagaa tgggggcca cccttgacaa 1320

actccttaaa	ccttttcatt	cggctaaaag	gttgaaaaac	atttgggaaga	tttggtttta	1380
ggaaatattt	atcgggtgaa	aagaccagat	gcatggctaa	cttaaaactcc	atagaaaaac	1440
tgaatcacga	aagaaagaat	gctatcaaaa	atggcattta	ccacttgatc	caaatcaa	1500
tttcttaca	ctccaatcgc	attgaaggaa	gcggtttgac	ttatgaaca	accgctcata	1560
tttttgaca	atccgttctc	ataactgaaa	aaaacaccaa	tatcaaactt	gatgatattt	1620
ttgaaactat	caatcatttt	gaatgcgtga	attacttgct	tgaaagctat	aaagaacctt	1680
tgagtttaga	atactttaag	aatttacaca	aaatcttgaa	aaagaattgt	tctgatgaag	1740
ttattggtga	ttttaaaaa	cgccctaatt	ttgtaggcaa	tagcgccaca	acaagacca	1800
aattagttga	aagcgaattg	acaaatcttg	tgaaaaatta	tcaacgcaac	cttgaagtga	1860
gtttgaaaa	caatatcatg	cctttcatca	tagaaaacga	acacaaagcc	ttttactaca	1920
ggggcatcaa	agaatatgac	aacacaaaag	gctacttgaa	agacaccatt	ttgcaaagtc	1980
aagacaattt	caatgaaatg	gttagctatt	tcttttcttg	agtgaaccg	cttatttttg	2040
cttggtgct	tttgttttt	gcgtttttag	ttgtagggtg	taagaaatat	cggttttttg	2100
cttttcgttg	gttgtaggcg	attttagata	gcaaaaaaca	gctaaaaaat	ccaagcaacc	2160
taattgattt	caaaccaact	tcatttcctt	tttagtcgtt	agccatttag	ccaatcta	2220
tagtttagca	tctaaaagcg	catataactt	gagttagcaa	tccaaccaat	actaaaaccg	2280
cctagcgaag	cgtttagcgag	caaaataagc	ggtttttagac	cgattgtttg	ctgacaagca	2340
aacaccaata	agcgagcggt	agcgagcatg	gacaaaagcg	catcgagtt	tgaaagcgta	2400
ggcgtagcc	gaagctgttt	tgcgtaagca	aatcaaaaa	gatagcgcaa	gccgaggtgc	2460
agcccaagaa	tttgaattaa	tccatgcggt	gttttagggcg	ttttagcgtg	atcgctttat	2520
tacatgtttt	aaacagcatg	ctgtttttta	catgttttac	tcgcatgcgc	gcgcgctagg	2580
tattggtggt	tggaatagcc	taaataacgc	agctgtatgg	tttctcattt	ttcggtgaca	2640
atgaataagg	ggtagttctt	gcgagtcata	agtgtagtct	ttgcgagtca	taagtgtagt	2700
tcttgcgagt	cataagtgtg	gttcttgcca	gtcataagtg	tagttctctt	cacaatatct	2760
acacaattca	caatctctag	ctacaataac	agcgtgtgaa	gatgctttca	cagcggattt	2820
tcctatgtgc	ggcaaaattt	ggagcaatta	gctttaaaag	ctagtgggtt	gggagtttgt	2880
agcgggtatg	cactccgtta	ggaggcacac	catgaaagca	tttttgatag	tagtgatttt	2940
agtggtaatc	ttgacacagc	cactatatta	aaaccttagc	gttttaataa	cccttataag	3000
tccgccaaga	cttcttaagg	gtttcactcc	tgttattata	tcgtcttttg	aaaaataagc	3060
attaaaaggc	gcttaaatgc	ccatgaatac	gaattttgaa	cagcttagaa	aacaagaatt	3120
ggaattacga	aaattattag	aagaattaga	aacgctccca	caaaccac	aaattaaact	3180
gcaaaaaaca	aaaatacaaa	cttacataga	caagataaca	ccaagtattt	tgagcggttt	3240
tgatcaaaaa	ttcaaagaaa	ttatagaaaa	tctatcaaat	gaatttgaaa	aagaaaaatc	3300
cacaccactc	aaagagccac	aaaccacccc	cacaccatgc	aaagatttag	tggttagcac	3360

ccctaaagat aacacctata ccacctacca caataacgct aataagggtca atctagggaa 3420
attgagcgaa agggaagcca atct 3444

<210> 4
<211> 580
<212> DNA
<213> Hepatitis C virus

<400> 4
catgagcacg aatcctaaac ctcaaagaaa aaccaaactg aacaccaacc gtcgcccaca 60
ggacgtcaag ttcccgggtg gcggtcagat cgttggtgga gtttacttgt tgccgcgcag 120
gggccctaga ttgggtgtgc gcgcgacgag gaagacttcc gagcggtcgc aacctcgagg 180
tagacgtcag cctatcccca aggcacgtcg gcccgagggc aggacctggg ctcagcccgg 240
gtacccttg g cccctctatg gcaatgaggg ttgcgggtgg gcgggatggc tcctgtctcc 300
ccgtggctct cggcctagct ggggccccac agacccccgg cgtaggtcgc gcaatttggg 360
taagggtcatc gataccctta cgtgcggctt cgccgacctc atgggggtaca taccgctcgt 420
cggcgccccct cttggaggcg ctgccagggc cctggcgcat ggcggtccggg ttctggaaga 480
cggcggtgaac tatgcaacag ggaaccttcc tggttgctct ttctctatct tccttctggc 540
cctgctctct tgcctgactg tgcccgttc agcctaccaa 580

<210> 5
<211> 135
<212> DNA
<213> Hepatitis C virus

<400> 5
aatcctaaac ctcaaagaaa aaccaaactg aacaccaacc gtcgcccaca ggacgtcaag 60
ttcccgggtg gcggtcagat cgttggtgga gtttacttgt tgccgcgcag gggccctaga 120
ttgggtgtgc gcgcg 135

<210> 6
<211> 1108
<212> DNA
<213> Helicobacter pylori

<400> 6
atgcatagc atttttatcc ataagattag cggatcctac ctgacgcttt ttatcgcaac 60
tctctactgt ttctccatac ccgttttttg ggctaacagg aggaattaac catggaattt 120
atgaaaaagt ttgtagcttt agggcttcta tccgcagttt taagctcttc gttgttagcc 180
gaagggtgatg gtgtttatat agggactaat tatcagcttg gacaagcccc tttgaatagt 240
aatatttata atacagggga ttgcacaggg agtggttagt gttgcccccc aggtcttacc 300
gctaataagc ataatccagg aggcaccaat atcaattggc atgctaaata cgctaattgg 360
gctttgaatg gtcttggggt gaatgtgggt tataagaagt tcttccagtt caagtctttt 420
gatatgacaa gcaagtgggt tggttttaga gtgtatgggc tttttgatta tgggcatgcc 480

```

actttaggca agcaagttta tgcacctaataaaaatccagttggatatggctccttggggt 540
gtggggagcg atttgttagc tgatattatt gataacgata acgcttcttt tggatatttt 600
ggtggggtcg ctatcggcgg taacacttgg aaaagctcag cggcaaacta ttggaaagag 660
caaatcattg aagctaaggg tcctgatgtt tgtacccta cttattgtaa ccctaacgct 720
ccttatagca ccaaaacttc aaccgtcgct tttcaggtat ggttgaattt tggggtgaga 780
gccaatattt acaagcataa tggcgtagag tttggcgtga gaggccgct actcatcaac 840
aagtttttga gtgcgggtcc taacgtact aatctttatt accatttgaa acgggattat 900
tcgctttatt tagggataa ctacactttt tctcgagatc tgcagctggt acgatatggg 960
aattcgaagc tttctagaac aaaaactcat ctcaagag gatctgaata gcgccgtcga 1020
ccatcatcat catcattgag tttaacggtc tccagcttgg ctgttttggc ggatgagaga 1080
agattttcag cctgatacag attaaatc 1108

```

```

<210> 7
<211> 29
<212> DNA
<213> Helicobacter pylori

```

```

<400> 7
aaggatccga taggaatgta aaggaatgg 29

```

```

<210> 8
<211> 28
<212> DNA
<213> Helicobacter pylori

```

```

<400> 8
ccgaattcta aaggcatgaa cgcttgca 28

```

```

<210> 9
<211> 15
<212> DNA
<213> Helicobacter pylori

```

```

<400> 9
agatctaagg acgtc 15

```

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU2005/001211

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl. ⁷: A61K 48/00, A61P 1/04, C12N 1/21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

AU* ; Inventor/Applicant search

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Derwent World Patent Index, Medline: helicobacter/pylori & vector/recombinant/nucleotide/control sequence & heterologous/homologous & expression

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2002 040516 A2 (Demi, Ludwig et al) 23 May 2002.	
X	WO 1998 027819 A2 (Vanderbilt University) 2 July 1998.	1 - 35
X	WO 2002 060933 A2 (Queen Elizabeth College, Dublin) 8 August 2002.	1 - 35
A	Londoño-Arcila, P. et al, Infection and Immunity , September 2002, Vol 70, No 9 pp 5096 - 5106	

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
12 September 2005

Date of mailing of the international search report
23 SEP 2005

Name and mailing address of the ISA/AU
AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaustalia.gov.au
Facsimile No. (02) 6285 3929

Authorized officer

K.G. ENGLAND

Telephone No : (02) 6283 2292

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU2005/001211

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Corthésy – Theulaz, I et al, Infection and Immunity, February 1998 Vol 66 No 2 pp 581 - 586	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU2005/001211

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	9827819	AU	59576/98	US	6570004	US	2003204068
WO	02060933	EP	1362098	IE	20020049	US	2004146526
Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001.							
END OF ANNEX							