



(51) International Patent Classification:

A61P 37/00 (2006.01) C07K 16/28 (2006.01)  
A61K 39/395 (2006.01)

(21) International Application Number:

PCT/EP2013/077898

(22) International Filing Date:

23 December 2013 (23.12.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/748,201 2 January 2013 (02.01.2013) US

(71) Applicant: **GLENMARK PHARMACEUTICALS S.A.**  
[CH/CH]; Chemin de la Combeta 5, CH-2300 La Chaux-  
de-Fonds (CH).

(72) Inventors: **ATTINGER, Antoine**; c/o Glenmark Pharma-  
ceuticals S.A., Chemin de la Combeta 5, CH-2300 La  
Chaux-de-Fonds (CH). **BACK, Jonathan Albert**; c/o  
Glenmark Pharmaceuticals S.A., Chemin de la Combeta 5,  
CH-2300 La Chaux-de-Fonds (CH). **BLEIN, Stanislas**;  
c/o Glenmark Pharmaceuticals S.A., Chemin de la  
Combeta 5, CH-2300 La Chaux-de-Fonds (CH). **LISSIL-  
AA, Rami**; c/o Glenmark Pharmaceuticals S.A., Chemin de  
la Combeta 5, CH-2300 La Chaux-de-Fonds (CH).  
**SKEGRO, Darko**; c/o Glenmark Pharmaceuticals S.A.,  
Chemin de la Combeta 5, CH-2300 La Chaux-de-Fonds  
(CH).

(74) Agent: **SCHOELLHORN, Andreas**; Latscha Schöllhorn  
Partner AG, Aeschenvorstadt 55, CH-4051 Basel (CH).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,  
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,  
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,  
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a  
patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))  
— with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTIBODIES THAT BIND TO TL1A AND THEIR USES

(57) Abstract: The present invention relates to antibodies or fragments thereof that bind to TL1A. More specifically, the present invention relates to an antibody or fragment thereof that binds to TL1A comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 1, and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, 5 and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and/or comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.



WO 2014/106602 A1

## Antibodies that bind to TL1A and their uses

### **Field of the Invention**

The present invention relates to antibodies or fragments thereof that bind to TL1A. More specifically, the present invention relates to an antibody or fragment thereof that binds to TL1A comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and/or comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.

### **Background of the Invention**

TNF-like ligand 1A (TL1A) is a member of the tumor necrosis factor (ligand) superfamily, member 15. TL1A is also known as TNFSF15 and VEGI and was identified in 1999 as an angiogenesis inhibitor that suppresses the growth of colon carcinomas *in vivo* (Zhai Y *et al* (1999) FASEB J, 13(1): 181-9). The protein is abundantly expressed in endothelial cells and activated cells of the hematopoietic lineage, including monocytes, macrophages, lymphocytes, lamina propria mononuclear cells, dendritic cells and plasma cells but is not expressed in either B or T cells (Tan KB *et al* (1997) Gene, 204: 35-46; Prehn JL *et al* (2007) J Immunol, 178: 4033-4038). It is also expressed in kidney, lung, prostate and thymus (Tan KB *et al* (1997), *supra*). It is a ligand for TNFRSF25/DR3 and decoy receptor TR6/DcR3 and its expression is inducible by TNF and IL-1 $\alpha$ . TNFRSF25/DR3 is a death domain-containing receptor that is upregulated during T cell activation. TL1A induces NF-kappaB activation and apoptosis in TNFRSF25/DR3-expressing cell lines, and in T cells, TL1A can act as a costimulator that increases IL-2 responsiveness and secretion of proinflammatory cytokines both *in vitro* and *in vivo*. The interaction of TL1A with DR3 can promote T cell expansion during an immune response (Migone TS *et al* (2002) Immunity, 16(3): 479-92). The secreted decoy receptor (DcR3), a soluble protein of the tumor necrosis factor receptor (TNFR) superfamily, blocks the action of TL1A. (Kim S & Zhang L (2005) J Immunol Methods, 298: 1-8). TL1A has been implicated as a potential therapeutic target in a number of diseases and disorders.

A major cause of lung inflammation in allergy and asthma is Th2 polarization of CD4 T cells with elevated IgE levels and production of IL-13 by NKT cells. TL1A plays a major role in allergic lung inflammation by co-stimulating IL-4 and IL-13 production in NKT cells. Blocking TL1A and DR3 interaction by TL1A antibody or dominant negative TL1A mutant abolishes lung inflammation (Fang L *et al.*, (2008) *J Exp Med*, 205(5): 1037-48). DcR3, the decoy receptor for TL1A is expressed in several lung and colon carcinomas and in some normal tissues, therefore suggesting a role for TL1A in lung and colon carcinomas. In addition, TL1A has also been reported to be angiostatic and to induce metalloproteinase and IL-8 gene expression (Su WB *et al.*, (2006) *Exp Cell Res*, 312: 266-277; Kang YJ *et al.*, (2005) *Cytokine*, 29: 229-235). TL1A and DR3 may also be involved in the pathogenesis of atherosclerosis by increasing the production of proinflammatory cytokines and chemokines and decreasing plaque stability by inducing extracellular matrix-degrading enzymes (Kang YJ *et al.*, (2005) *supra*). There is also evidence to suggest that TL1A/DR3 is involved in the etiology of rheumatoid arthritis (Bossen C *et al.*, (2006) *J Biol Chem*, 281(20): 13964-13971).

An association between the expression of TL1A and inflammatory bowel disease has been identified by researchers (Prehn JL *et al.*, (2004) *Clin Immunol*, 112: 66-77; Bamias G *et al.*, (2003) *J Immunol*, 171: 4868-4874). Crohn's disease, which is a severe inflammatory bowel disorder, is thought to originate from predisposing genetic and environmental factors that cause an imbalance of effector (proinflammatory) and regulatory T cell responses, resulting in inflammation of the gastrointestinal mucosa and disease. The TL1A/DR3 pathway has been shown to play an important role in intestinal diseases, such as Crohn's disease (Papadakis KA *et al.*, (2005) *J. Immunol*, 174: 4985-4990; Bamias G *et al.*, (2003) *supra*) and therefore, blockade of the TL1A/DR3 pathway may offer therapeutic opportunities in this disease.

Death receptors and their ligands play a key role in the maintenance of tissue homeostasis and the physiological regulation of programmed cell death. Binding of a death ligand induces oligomerization of the receptor, recruitment of an adapter protein via a conserved cytoplasmic signalling element termed the death domain, activation of caspases and induction of apoptosis (Young HA *et al.*, (2006) *Proc Natl Acad Sci USA*, 103(22): 8303-8304). Although death receptors such as Fas/Apo-1/CD95, TNF-R1, TRAIL-R1, TRAIL-R2, or DR3 were initially characterized as inducers of apoptosis, there is growing evidence that these receptors also have non-apoptotic functions, including regulation of the adaptive immune response. Bamias *et al.*,

reported that TL1A is expressed by lamina propria dendritic cells and that it functions by increasing the proliferation of memory cells, but not naïve CD4<sup>+</sup> T cells, and synergizes with IL-12 and/or low-dose stimulation of the T cell receptor to strongly enhance IFN- $\gamma$  gene expression (Bamias G *et al.*, (2006) Proc. Natl. Acad. Sci. USA, 103: 8441-8446). IFN- $\gamma$  expression in the gut has been considered a marker of inflammation and many strategies for treating Crohn's disease rely on broad attempts to suppress the immune-activated state. However, such approaches (steroid treatment and immunosuppressive drugs) do not focus on the gut specifically and therefore have their own complications. Targeted therapies based on the use of antagonists of TNF- $\alpha$  were introduced with success in the 1990s and the results suggest that therapy directed specifically against TL1A or its receptor may provide an alternative targeted therapy for this debilitating disorder.

Current treatments for Crohn's disease include the anti- TNF- $\alpha$  monoclonal antibodies Infliximab (Remicade<sup>®</sup>; Centocor) and Adalimumab (Humira<sup>®</sup>; Abbott), as well as anti-inflammatories (e.g., sulfasalazine), cortisone or steroids (e.g., prednisone), immune system suppressors (e.g., 6-mercaptopurine) and antibiotics. However, Infliximab is the only treatment option having a high degree of specificity compared to the other available treatments (Young HA *et al.*, (2006) *supra*). Although Infliximab is generally well tolerated it can cause a recurrence of tuberculosis infection, worsening of heart failure, demyelinating disease and an increased incidence of lymphoma.

Therefore there remains a need in the art for compositions that can be used in the treatment and diagnosis of diverse inflammatory and immune diseases and disorders.

## Summary of the Invention

The present disclosure relates generally to antibodies or fragments thereof that bind to TL1A, methods for their preparation and use, including methods for treating TL1A mediated disorders. The antibodies or fragments thereof of the present invention that bind to TL1A exhibit numerous desirable properties and may be useful for the treatment of various diseases that include but are not limited to inflammatory diseases and/or auto immune diseases, including *inter alia* inflammatory bowel diseases (e.g., ulcerative colitis and Crohn's disease), rheumatoid arthritis, multiple sclerosis (MS), atherosclerosis, transplant rejection, central nervous system injury, psoriasis, leukaemia or lymphoma (e.g., chronic lymphocytic leukaemia

(CLL)), atherosclerosis, and lung and colon carcinomas. The antibodies or fragments thereof of the present invention that bind to TL1A exhibit numerous desirable properties and may be useful for the treatment of various diseases that include but are not limited to inflammatory diseases and/or auto immune diseases, including *inter alia* inflammatory bowel diseases (e.g.,  
5 ulcerative colitis and Crohn's disease), rheumatoid arthritis, multiple sclerosis (MS), atherosclerosis, transplant rejection, central nervous system injury, psoriasis, leukaemia or lymphoma (e.g., chronic lymphocytic leukaemia (CLL)), atherosclerosis, and lung and colon carcinomas, chronic obstructive pulmonary disease COPD, optic neuritis, age related macular degeneration, systemic lupus erythematosus (SLE), sjogen's syndrome, scleroderma, systemic  
10 sclerosis, chronic Kidney disease, liver fibrosis, tuberculosis, idiopathic pulmonary fibrosis, tuberculosis induced lung fibrosis, retroperitoneal Fibrosis, pulmonary fibrosis , cystic fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis (bone marrow), retroperitoneal fibrosis, progressive massive fibrosis, pephrogenic systemic fibrosis, arthrofibrosis.

15

In one aspect, the present disclosure provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and/or  
20 comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.

25

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 1. In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable framework region that is the product of or derived from a human gene selected from the group consisting of: IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4), IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-  
30 2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID NO: 7).

30

In a further aspect the present invention provides an antibody or fragment thereof comprising a heavy chain variable framework region that is the product of or derived from human gene IGHV1-2\*02 (SEQ ID NO: 3) and wherein the heavy chain variable framework region

comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody.

5 In a further aspect the present invention provides an antibody or fragment thereof comprising a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 13 and wherein the heavy chain variable framework region comprises at least one amino acid modification from the corresponding heavy chain variable framework region of the corresponding murine antibody.

10 In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 2. In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a light chain variable framework region that is the product of or derived from a human gene selected from the group consisting of: IGKV1-33\*01 (SEQ ID NO: 8), IGKV1D-33\*01 (SEQ ID NO: 9), IGKV1D-12\*02 (SEQ ID NO: 10),  
15 IGKV1D-12\*01 (SEQ ID NO: 11) and IGKV1-12\*02 (SEQ ID NO: 12).

In a further aspect the present invention provides an antibody or fragment thereof comprising a light chain variable framework region that is the product of or derived from human gene IGKV1-33\*01 (SEQ ID NO: 8) and wherein the light chain variable framework region  
20 comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody.

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain sequence selected from the group consisting of SEQ ID NOS:  
25 16, 21, 22, 23 and 24. In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a light chain sequence selected from the group consisting of SEQ ID NOS: 17 and 25.

30 In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising:

- (a) a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 22 or 24; and
- (b) a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17.

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOS: 13, 26, 27, 28 and 29. In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a  
5 light chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOS: 14 and 30.

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising:

- 10 (a) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 27 or 29; and  
(b) a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14.

In a further aspect the present invention provides an antibody or fragment thereof that binds to  
15 TL1A, wherein the antibody comprises a human IgG4 Fc region, wherein the antibody has no Fc-mediated cytotoxicity activity. In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A, wherein the antibody comprises a human IGHG1 Fc region, wherein the antibody is competent for cytotoxicity mechanisms such as antibody dependent cellular cytotoxicity (ADCC). In a preferred aspect, the antibody or  
20 fragment thereof that binds to TL1A has a non fucosylated IGHG1 Fc region and exhibits enhanced Fc-mediated cytotoxicity mechanisms such as ADCC.

In another aspect, the present invention provides a cross-reactive antibody or fragment thereof which binds to human TL1A and which also binds to murine, rat and cynomologous TL1A. By  
25 "cross-reactive antibody" is meant an antibody that binds to an antigen from one species, e.g. human, and which also binds to the corresponding antigen in a different species, e.g. rat.

In another aspect, the disclosure of the present invention also describes humanized antibodies or fragments thereof that bind with a similar affinity to TL1A as the corresponding chimeric  
30 antibody e.g. retain at least 85% of the TL1A binding affinity ( $K_D$ ) of the corresponding chimeric antibody or have at least equivalent or higher TL1A binding affinity ( $K_D$ ) when compared to the corresponding chimeric antibody. In a preferred aspect the humanised

antibody or fragment thereof has approximately a three-fold higher TL1A binding affinity when compared to the corresponding chimeric antibody.

5 In a further aspect, the present invention also describes humanized antibodies or fragments thereof that bind to hTL1A and inhibits the interaction of hTL1A with both DR3 and DcR3.

The disclosure of the present invention also provides isolated nucleic acids encoding antibodies and fragments thereof that bind to TL1A, vectors and host cells comprising the nucleic acid or the vector. Compositions comprising the anti-TL1A antibody or fragment thereof and a  
10 pharmaceutically acceptable carrier and immunoconjugates comprising the antibody or fragment thereof linked to a therapeutic agent are also provided.

The present disclosure also provides methods for treating TL1A mediated disorders. In one aspect, in an *in vitro* model of TL1A-induced IFN $\gamma$  secretion by primed CD4 T cells, an anti-  
15 TL1A antibody or fragment thereof efficiently suppressed the production of IFN $\gamma$  induced by immune complex-stimulated monocytes. In another aspect, in an *in vivo* model of allergic asthma, an anti-TL1A antibody reduced the number of eosinophils in bronchoalveolar lavage fluid of asthmatic mice by approximately 4-fold. In a further aspect, in an *in vivo* model of acute colitis induced in mice with dextran sulphate sodium (DSS) and in rats by  
20 trinitrobenzenesulfonic acid (TNBS), an anti-TL1A antibody was effective in reducing the symptoms of disease.

The present disclosure also provides pharmaceutical compositions comprising an anti-TL1A antibody or fragments thereof and a carrier, such as a diluent or excipient.  
25

The present disclosure also provides kits and articles of manufacture comprising the antibody or fragments thereof, a composition or an immunoconjugate for the treatment of a TL1A mediated disorder.

### 30 **Brief Description of the Figures**

**Figure 1:** This figure shows the binding of hybridoma antibodies to human TL1A-his (Fig 1A) or the irrelevant protein-his (Fig 1B), detected using an HRP-labelled anti-mouse IgG secondary antibody and TMB substrate. Fig 1A displays the absorbance at 450nm of an ELISA



against coated human TL1A-his and Fig 1B displays the absorbance at 450nm of an ELISA against an irrelevant protein-his.

**Figure 2:** This figure shows the effect of purified hybridoma anti-TL1A antibodies at three  
5 different concentrations in a blocking ELISA, where the binding of human TNFRSF25 to TL1A was evaluated in the presence of the antibodies shown in Fig 2. Absorbance was read at 450nm. mIgG: mouse IgG isotype control.

**Figure 3:** Parental 5G6 candidate blocks the effect of soluble and membrane bound TL1A  
10 produced by activated monocytes: Fig 3A: PBMCs from healthy donors were stimulated with immune complexes then stained with fluorescent antibodies. The monocytes were gated based on large forward light scatter and high side scatter parameters. The histogram plot displays the PE fluorescence of monocytes gated population. The grey shaded histogram represents the isotype control and the blank histogram represents the staining with anti TL1A. Fig 3B: The  
15 supernatants of purified human monocytes from healthy donor PBMCs stimulated with immune complex were harvested and tested by ELISA for the presence of sTL1A proteins. The graph shows the interpolated TL1A concentration measured in the supernatants of indicated conditions. 'IC stim' means immune complex stimulated. 'NS' means not stimulated. Fig 3C: Naïve CD4 T cells purified from healthy donor PBMCs were incubated with IL-12, IL-  
20 18 and IC stimulated autologous monocytes. The parental chimeric 5G6 antibody was added at the concentrations indicated in the table at the same time as the monocytes. NA indicates that no IL-12 and IL-18 was added. The supernatants of cultures were quantified by ELISA. The graph shows the interpolated IFN- $\gamma$  concentration for each indicated condition.

**Figure 4:** Parental 5G6 candidate binds to mouse, rat, cynomologus monkey and human  
25 TL1A: The binding of 5G6 on the extracellular part of TL1A protein corresponding to human (*homo sapiens*), rat (*ratus norvegicus*), mouse (*mus musculus*) and cynomologus monkey (*macaca fascicularis*) sequences was determined by immunofluorescence. The graph shows the absorbance at 450nm according to the log of the concentration of 5G6 used.

**Figure 5:** Humanized 5G6 antibodies block TL1A-induced IFN- $\gamma$  secretion by primed CD4 T  
30 cells: Naïve CD4 T cells were incubated with IL-12, IL-18 and recombinant soluble human TL1A and the humanized 5G6 candidates (VH3/VL1, VH4/VL1, VH5/VL1 and VH2/VL2)

were added at the concentrations indicated in the table. NA indicates that no IL-12 and IL-18 was added. The culture supernatants were quantified by ELISA for concentration of IFN- $\gamma$ . Graphs 5A-5D shows the IFN- $\gamma$  concentration for each culture condition. Each graph displays the result of one humanized 5G6 candidate.

5

**Figure 6:** Humanized 5G6 antibody reduced the number of cells in bronchoalveolar lavage (BAL) fluid in a murine model of allergic asthma. Mice were treated with the humanized 5G6 candidate (VH5/VL1; format IgG4 hinge stabilised) at 50mg/kg or an equivalent amount of control human IgG, or dexamethasone at 5mg/kg (positive control), on day 28, 30 and 33 following the induction of an immunological response induced by ovalbumin challenge. The graph shows the number of eosinophils in BAL fluid for each mouse and the average number of eosinophils for each group was calculated. Standard deviation was calculated using a one way ANOVA \* indicates  $p < 0.05$ , and \*\* indicates  $p < 0.01$ .

10

15

**Figure 7:** Treatment by humanized 5G6 antibody ameliorates shortening of colon in a DSS-induced model of acute colitis. Mice were treated 3 x week with 50mg/kg of a humanized 5G6 antibody (VH5/VL1; format IgG4 hinge stabilised) or an equivalent amount of isotype control, or cyclosporine at 5mg/kg (positive control). The graph shows the entire colon length for each mouse and the average length per group. Standard deviation was calculated using a one way ANOVA \* indicates  $p < 0.05$ , \*\* indicates  $p < 0.01$  and \*\*\* indicates  $p < 0.001$ .

20

**Figure 8:** Treatment by humanized 5G6 antibody ameliorates disease severity in a TNBS-induced model of acute colitis. Rats were treated i.p with a single dose of humanized 5G6 antibody (50mg/kg) or an equal amount of isotype control, two hours after TNBS administration. Prednisolone was administered as a positive control. Disease severity was assessed using a colonic score for adhesions, strictures, ulcers and wall thickness and the average score is shown in the histograms. Standard deviation was calculated using a Student's t-test and \* indicates  $p < 0.05$ .

25

30

**Figure 9:** Binding of 5G6 to hTL1A is blocked by both hDcR3-Fc and hDR3-Fc. Histidine-tagged human TL1A was coated at 2  $\mu\text{g/ml}$  on an ELISA plate and incubated with 20  $\mu\text{g/ml}$  5G6 in the presence of 10  $\mu\text{g/ml}$  Fc fusions of the ectodomains of either human DcR3 (white

bar), DR3 (black bar) or an irrelevant receptor (Ctrl-Fc, hatched bar) followed by detection with peroxidase-conjugated anti-human IgG (Fab specific).

### Detailed description of the invention

5 The present disclosure relates to antibodies and fragments thereof that bind to TL1A.

The term “TL1A” as used herein includes variants, isoforms, and species homologs of TL1A. Accordingly, antibodies of this disclosure may bind to human TL1A and may cross-react with TL1A from species other than human, for example, mouse, rat or cynomolgous monkey. In  
10 certain embodiments, the antibodies may be completely specific for one or more human TL1A proteins and may not exhibit species or other types of non-human cross-reactivity. The complete amino acid sequence of an exemplary human TL1A has Swiss-Prot accession number O95150 (TNFSF15\_HUMAN; SEQ ID NO:38). TL1A is also known as TNFSF15; TNF-like protein 1A; VEGI; TNF $\gamma$  $\beta$ . Human TL1A is designated GeneID: 9966 by Entrez Gene, and  
15 HGNC: 11931 by HGNC. TL1A can be encoded by the gene designated TNFSF15 /TL1A. The complete amino acid sequence of an exemplary murine TL1A has Swiss-Prot accession number Q5UBV8 (TNFSF15\_MOUSE; SEQ ID NO: 39). Murine TL1A is designated GeneID: 326623 by Entrez Gene. The complete amino acid sequence of an exemplary rat  
20 TL1A has Swiss-Prot accession number Q8K3Y7 (TNFSF15\_RAT; SEQ ID NO: 40). Rat TL1A is designated GeneID: 252878 by Entrez Gene. The complete amino acid sequence of an exemplary cyno TL1A (*macaca fascicularis*) has SEQ ID NO: 41.

The use of “TL1A” herein encompasses all known or as yet undiscovered alleles and polymorphic forms of TL1A, preferably human TL1A.

25 The term “antibody or fragment thereof that binds to TL1A” as used herein includes antibodies or a fragment thereof that binds to TL1A e.g. human TL1A in isolated form, with an affinity ( $K_D$ ) of 850 pM or less, preferably 700nM or less, more preferably 300 nM or less, more preferably 260 nM or less, even more preferably 250 nM or less.

30 The term “antibody or fragment thereof that binds to TL1A” includes antibodies or antigenic binding fragments thereof.

The term “antibody” as referred to herein includes whole antibodies and any antigen binding fragments or single chains thereof. An “antibody” refers to a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding fragment thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR) with are hypervariable in sequence and/or involved in antigen recognition and/or usually form structurally defined loops, interspersed with regions that are more conserved, termed framework regions (FR or FW). Each VH and VL is composed of three CDRs and four FWs, arranged from amino- terminus to carboxy-terminus in the following order: FW1, CDR1, FW2, CDR2, FW3, CDR3, FW4. The amino acid sequences of FW1, FW2, FW3, and FW4 all together constitute the “non-CDR region” or “non-extended CDR region” of VH or VL as referred to herein.

The term “heavy chain variable framework region” as referred herein may comprise one or more (e.g., one, two, three and/or four) heavy chain framework region sequences (e.g., framework 1 (FW1), framework 2 (FW2), framework 3 (FW3) and/or framework 4 (FW4)). Preferably the heavy chain variable region framework comprises FW1, FW2 and/or FW3, more preferably FW1, FW2 and FW3. The term “light chain variable framework region” as referred herein may comprise one or more (e.g., one, two, three and/or four) light chain framework region sequences (e.g., framework 1 (FW1), framework 2 (FW2), framework 3 (FW3) and/or framework 4 (FW4)). Preferably the light chain variable region framework comprises FW1, FW2 and/or FW3, more preferably FW1, FW2 and FW3.

The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

Antibodies are grouped into classes, also referred to as isotypes, as determined genetically by

the constant region. Human constant light chains are classified as kappa (CK) and lambda (C $\lambda$ ) light chains. Heavy chains are classified as mu ( $\mu$ ), delta ( $\delta$ ), gamma ( $\gamma$ ), alpha ( $\alpha$ ), or epsilon ( $\epsilon$ ), and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. Thus, "isotype" as used herein is meant any of the classes and/or subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. The known human immunoglobulin isotypes are IgG1 (IGHG1), IgG2 (IGHG2), IgG3 (IGHG3), IgG4 (IGHG4), IgA1 (IGHA1), IgA2 (IGHA2), IgM (IGHM), IgD (IGHD), and IgE (IGHE). The so-called human immunoglobulin pseudo-gamma IGHGP gene represents an additional human immunoglobulin heavy constant region gene which has been sequenced but does not encode a protein due to an altered switch region (Bensmana M *et al.*, (1988) *Nucleic Acids Res.* 16(7): 3108). In spite of having an altered switch region, the human immunoglobulin pseudo-gamma IGHGP gene has open reading frames for all heavy constant domains (CH1-CH3) and hinge. All open reading frames for its heavy constant domains encode protein domains which align well with all human immunoglobulin constant domains with the predicted structural features. This additional pseudo-gamma isotype is referred herein as IgGP or IGHGP. Other pseudo immunoglobulin genes have been reported such as the human immunoglobulin heavy constant domain epsilon P1 and P2 pseudo-genes (IGHEP1 and IGHEP2). The IgG class is the most commonly used for therapeutic purposes. In humans this class comprises subclasses IgG1, IgG2, IgG3 and IgG4. In mice this class comprises subclasses IgG1, IgG2a, IgG2b, IgG2c and IgG3.

The term "murine antibody" as used herein includes antibodies in which the variable region sequences and the constant region sequences are derived from a mouse.

The term "chimeric antibody" as used herein includes antibodies in which the variable region sequences are derived from one species and the constant region sequences are derived from another species, such as an antibody in which the variable region sequences are derived from a mouse antibody and the constant region sequences are derived from a human antibody.

The term "humanized antibody" or "humanized anti-TL1A antibody" as used herein includes antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. Additional framework

region modifications may be made within the human framework sequences as well as within the CDR sequences derived from the germline of another mammalian species.

5 The term “neutralising antibody” includes an antibody that is capable of inhibiting and/or neutralising the biological activity of TL1A, for example by blocking binding or substantially reducing binding of TL1A to its receptor TNFRSF25/DR3 or the decoy receptor TNFRSF21/DR6 and thus inhibiting or reducing the signalisation pathway triggered by TL1A and/or inhibiting or reducing a TL1A-mediated cell response like e.g. lymphocyte proliferation, cytokine expression, or lymphocyte survival.

10

The terms “antagonistic antibody” or “antagonist antibody” are used herein equivalently and include an antibody that is capable of inhibiting and/or neutralising the biological signalling activity of TL1A, as described for a neutralising antibody *supra*.

15

The terms “agonistic antibody” or “agonist antibody” are used herein equivalently and include an antibody that is capable of activating and/or enhancing the biological signalling activity of TL1A, for example by increasing binding of TL1A to its receptor TNFRSF25/DR3 or the decoy receptor TNFRSF21/DR6 and thus activating or enhancing the signalisation pathway triggered by TL1A and/or activating or enhancing a TL1A-mediated cell response like e.g.

20

lymphocyte proliferation, cytokine expression, or lymphocyte survival.

The term “Fab” or “Fab region” as used herein includes the polypeptides that comprise the VH, CH1, VL, and CL immunoglobulin domains. Fab may refer to this region in isolation, or this region in the context of a full length antibody or antibody fragment.

25

The term “Fc” or “Fc region”, as used herein includes the polypeptide comprising the constant region of an antibody excluding the first constant region immunoglobulin domain. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, and the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-

30

terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, Fc comprises immunoglobulin domains C gamma 2 and C gamma 3 (C $\gamma$ 2 and C $\gamma$ 3) and the hinge between C gamma 1 (C $\gamma$ 1) and C gamma 2 (C $\gamma$ 2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to comprise residues C226

or P230 to its carboxyl-terminus, wherein the numbering is according to the EU numbering system. For human IgG1 the Fc region is herein defined to comprise residue P232 to its carboxyl-terminus, wherein the numbering is according to the EU numbering system (Edelman GM *et al.*, (1969) Proc Natl Acad Sci USA, 63(1): 78-85). Fc may refer to this region in  
5 isolation or this region in the context of an Fc polypeptide, for example an antibody.

The term “hinge” or “hinge region” or “antibody hinge region” herein includes the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. The “hinge region” as referred to herein is a sequence region of 6-62 amino acids in  
10 length, only present in IgA, IgD and IgG, which encompasses the cysteine residues that bridge the two heavy chains. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 231 (A231 in IgG1), wherein the numbering is according to the EU numbering system (Edelman GM *et al.*, *supra*).

15 The term “parent antibody” or “parent immunoglobulin” as used herein includes an unmodified antibody that is subsequently modified to generate a variant. Said parent antibody may be a naturally occurring antibody, or a variant or engineered version of a naturally occurring antibody. Parent antibody may refer to the antibody itself, compositions that comprise the  
20 parent antibody, or the amino acid sequence that encodes it. By “parent anti-TL1A antibody” as used herein is meant an antibody or immunoglobulin that binds TL1A and is modified to generate a variant. By “corresponding murine antibody” as used herein is meant a murine antibody or immunoglobulin that binds to TL1A and that can be modified to generate a variant, specifically the murine antibody 5G6 as disclosed herein. By “corresponding chimeric  
25 antibody” as used herein is meant a chimeric antibody or immunoglobulin that binds to TL1A and that can be modified to generate a variant.

The term “variant antibody” or “antibody variant” as used herein includes an antibody sequence that differs from that of a parent antibody sequence by virtue of at least one amino  
30 acid modification compared to the parent. The variant antibody sequence herein will preferably possess at least about 80%, most preferably at least about 90%, more preferably at least about 95% amino acid sequence identity with a parent antibody sequence. Antibody variant may refer

to the antibody itself, compositions comprising the antibody variant, or the amino acid sequence that encodes it.

5 The term “identity” or “substantial identity” or “substantially identical,” when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 80%, and more preferably at least about 90%, 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known algorithm of sequence identity, such as FASTA, BLAST or GAP, as discussed below.

10

As applied to polypeptides, the term “substantial similarity” or “substantially similar” means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80% sequence identity, even more preferably at least 90%, 95%, 98% or 99% sequence identity. Preferably, residue positions 15 which are not identical differ by conservative amino acid substitutions.

15

The term “amino acid modification” herein includes an amino acid substitution, insertion, and/or deletion in a polypeptide sequence. By “amino acid substitution” or “substitution” herein is meant the replacement of an amino acid at a particular position in a parent 20 polypeptide sequence with another amino acid. For example, the substitution R94K refers to a variant polypeptide, in this case a heavy chain variable framework region variant, in which the arginine at position 94 is replaced with a lysine. For the preceding example, 94K indicates the substitution of position 94 with a lysine. For the purposes herein, multiple substitutions are typically separated by a slash. For example, R94K/L78V refers to a double variant comprising 25 the substitutions R94K and L78V. By “amino acid insertion” or “insertion” as used herein is meant the addition of an amino acid at a particular position in a parent polypeptide sequence. For example, insert -94 designates an insertion at position 94. By “amino acid deletion” or “deletion” as used herein is meant the removal of an amino acid at a particular position in a parent polypeptide sequence. For example, R94- designates the deletion of arginine at position 30 94.

25

30

As used herein, the term “conservative modifications” or “conservative sequence modifications” is intended to refer to amino acid modifications that do not significantly affect



or alter the binding characteristics of the antibody containing the amino acid sequence. Such conservative modifications include amino acid substitutions, insertions and deletions.

Modifications can be introduced into an antibody of the invention by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis.

5 Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine,  
10 tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues within the CDR regions or within the framework regions of an antibody of the invention can be replaced with other amino acid residues from the same side  
15 chain family and the altered antibody (variant antibody) can be tested for retained function.

The term “epitope” refers to a region of an antigen that is bound by an antibody. An epitope may be defined as structural or functional. Functional epitopes are generally a subset of structural epitopes and have those residues that directly contribute to the affinity of the  
20 interaction. Epitopes may also be conformational, that is, composed of non-linear amino acids. In certain embodiments, epitopes may include determinants that are chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl groups, or sulfonyl groups, and, in certain embodiments, may have specific three-dimensional structural characteristics, and/or specific charge characteristics.

25 For all human immunoglobulin heavy chain constant domains numbering is according to the “EU numbering system” (Edelman GM *et al.*, (1969) Proc Natl Acad Sci USA, 63(1): 78-85). For the human kappa immunoglobulin light chain constant domain (IGKC), numbering is according to the “EU numbering system” (Edelman GM *et al.*, *supra*).

30 For the human lambda immunoglobulin light chain constant domains (IGLC1, IGLC2, IGLC3, IGLC6, and IGLC7), numbering is according to the “Kabat numbering system” (Kabat EA *et al.*, (1991) Sequences of proteins of immunological interest. 5th Edition - US Department of

Health and Human Services, NIH publication no 91-3242) as described by Dariavach P *et al.*, (1987) Proc Natl Acad Sci USA, 84(24): 9074-8 and Frangione B *et al.*, (1985) Proc Natl Acad Sci USA, 82(10): 3415-9.

5 The term “variable domain” refers to the domains that mediates antigen-binding and defines specificity of a particular antibody for a particular antigen. In naturally occurring antibodies, the antigen-binding site consists of two variable domains that define specificity: one located in the heavy chain (VH) and the other located in the light chain (VL). In some cases, specificity may exclusively reside in only one of the two domains as in single-domain antibodies from  
10 heavy-chain antibodies found in camelids. The V regions are usually about 110 amino acids long, and consist of relatively invariant stretches of amino acid sequence called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called “hypervariable regions” that are 9-12 amino acids long. The variable domains of native heavy and light chains comprise four FRs, largely adopting a beta-sheet configuration, connected by  
15 three hypervariable regions, which form loops. The hypervariable regions in each chain are held together in close proximity by FRs, and with the hypervariable regions from the other chain, contribute to the formation of the antigen binding site of antibodies (see Kabat EA *et al.*, *supra*). The term “hypervariable region” as used herein refers to the amino acid residues of an antibody which are responsible for antigen binding. The hypervariable region generally  
20 comprises amino acid residues from a “complementary determining region” or “CDR”, the latter being of highest sequence variability and/or involved in antigen recognition. For all variable domains numbering is according to Kabat (Kabat EA *et al.*, *supra*).

A number of CDR definitions are in use and are encompassed herein. The Kabat definition is  
25 based on sequence variability and is the most commonly used (Kabat EA *et al.*, *supra*). Chothia refers instead to the location of the structural loops (Chothia C & Lesk AM (1987) J. Mol. Biol. 196: 901-917). The AbM definition is a compromise between the Kabat and the Chothia definitions and is used by Oxford Molecular's AbM antibody modelling software (Martin ACR *et al.*, (1989) Proc. Natl Acad. Sci. USA, 86: 9268-72; Martin ACR *et al.*, (1991) Methods  
30 Enzymol. 203: 121-153; Pedersen JT *et al.*, (1992) Immunomethods, 1: 126-136; Rees AR *et al.*, (1996) In Sternberg M.J.E. (ed.), Protein Structure Prediction. Oxford University Press, Oxford, 141-172). The contact definition has been recently introduced (MacCallum RM *et al.*, (1996) J. Mol. Biol. 262: 732-745) and is based on an analysis of the available complex

structures available in the Protein Databank. The definition of the CDR by IMGT<sup>®</sup>, the international ImMunoGeneTics information system<sup>®</sup> (<http://www.imgt.org>) is based on the IMGT numbering for all immunoglobulin and T cell receptor V-REGIONS of all species (IMGT<sup>®</sup>, the international ImMunoGeneTics information system<sup>®</sup>; Lefranc MP *et al.*, (1991) Nucleic Acids Res. 27(1): 209-12; Ruiz M *et al.*, (2000) Nucleic Acids Res. 28(1): 219-21; Lefranc MP (2001) Nucleic Acids Res. 29(1): 207-9; Lefranc MP (2003) Nucleic Acids Res. 31(1): 307-10; Lefranc MP *et al.*, (2005) Dev. Comp. Immunol. 29(3): 185-203; Kaas Q *et al.*, (2007) Briefings in Functional Genomics & Proteomics, 6(4): 253-64).

10 All Complementarity Determining Regions (CDRs) discussed in the present invention, are defined preferably according to IMGT<sup>®</sup>. The variable domain residues for each of these CDRs are as follows (numbering according to Kabat EA, *et al.*, *supra*): LCDR1: 27-32, LCDR2: 50-52, LCDR3: 89-97, HCDR1: 26-35, HCDR2: 51-57 and HCDR3: 93-102. The “non-CDR region” of the VL region as used herein comprise the amino acid sequences: 1-26 (FR1), 33-15 49 (FR2), 53-88 (FR3), and 98- approximately 107 (FR4). The “non-CDR region” of the VH region as used herein comprise the amino acid sequences: 1-25 (FR1), 36-50 (FR2), 58-92 (FR3), and 103- approximately 113 (FR4).

The CDRs of the present invention may comprise “extended CDRs” which are based on the 20 aforementioned definitions and have variable domain residues as follows: LCDR1: 24-36, LCDR2: 46-56, LCDR3:89-97, HCDR1: 26-36, HCDR2:47-65, HCDR3: 93-102. These extended CDRs are numbered as well according to Kabat *et al.*, *supra*. The “non-extended CDR region” of the VL region as used herein comprise the amino acid sequences: 1-23 (FR1), 37-45 (FR2), 57-88 (FR3), and 98- approximately 107 (FR4). The “non-extended CDR 25 region” of the VH region as used herein comprise the amino acid sequences: 1-25 (FR1), 37-46 (FR2), 66-92 (FR3), and 103- approximately 113 (FR4).

The term “full length antibody” as used herein includes the structure that constitutes the natural biological form of an antibody, including variable and constant regions. For example, in most 30 mammals, including humans and mice, the full length antibody of the IgG class is a tetramer and consists of two identical pairs of two immunoglobulin chains, each pair having one light and one heavy chain, each light chain comprising immunoglobulin domains VL and CL, and each heavy chain comprising immunoglobulin domains VH, CH1 (C $\gamma$ 1), CH2 (C $\gamma$ 2), and CH3

(C $\gamma$ 3). In some mammals, for example in camels and llamas, IgG antibodies may consist of only two heavy chains, each heavy chain comprising a variable domain attached to the Fc region.

Antibody fragments include, but are not limited to, (i) the Fab fragment consisting of VL, VH, CL and CH1 domains, including Fab' and Fab'-SH, (ii) the Fd fragment consisting of the VH and CH1 domains, (iii) the Fv fragment consisting of the VL and VH domains of a single antibody; (iv) the dAb fragment (Ward ES *et al.*, (1989) *Nature*, 341: 544-546) which consists of a single variable, (v) F(ab')<sub>2</sub> fragments, a bivalent fragment comprising two linked Fab fragments (vi) single chain Fv molecules (scFv), wherein a VH domain and a VL domain are linked by a peptide linker which allows the two domains to associate to form an antigen binding site (Bird RE *et al.*, (1988) *Science* 242: 423-426; Huston JS *et al.*, (1988) *Proc. Natl. Acad. Sci. USA*, 85: 5879-83), (vii) bispecific single chain Fv dimers (PCT/US92/09965), (viii) "diabodies" or "triabodies", multivalent or multispecific fragments constructed by gene fusion (Tomlinson I & Hollinger P (2000) *Methods Enzymol.* 326: 461-79; WO94/13804; Holliger P *et al.*, (1993) *Proc. Natl. Acad. Sci. USA*, 90: 6444-48) and (ix) scFv genetically fused to the same or a different antibody (Coloma MJ & Morrison SL (1997) *Nature Biotechnology*, 15(2): 159-163).

The term "effector function" as used herein includes a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or ligand. Effector functions include Fc $\gamma$ R-mediated effector functions such as ADCC (antibody dependent cell-mediated cytotoxicity) and ADCP (antibody dependent cell-mediated phagocytosis), and complement-mediated effector functions such as CDC (complement dependent cytotoxicity). An effector function of an antibody may be altered by altering, i.e. enhancing or reducing, preferably enhancing, the affinity of the antibody for an effector molecule such as an Fc receptor or a complement component. Effector function may be determined using one or more cell based or *in vivo* assays. Such assays often involve monitoring the response of cells to the antibody, for example cell survival, cell death, change in cellular morphology or transcriptional activation such as cellular expression of a natural gene or reporter gene. For example, such assays may measure the ability of an antibody to elicit ADCC, ADCP, or CDC. For some assays additional cells or components, that is in addition to the target cells, may need to be added, for example serum complement or effector cells such as peripheral blood monocytes (PBMCs), NK cells, macrophages, and the like. Enhanced effector function can be determined by comparing the

effector function of an altered antibody with a control antibody and detecting, for example, an increase in ADCC, ADCP or CDC measured by one of more of the aforementioned assays.

5 Binding affinity will generally be varied by modifying the effector molecule binding site and in this case it is appropriate to locate the site of interest and modify at least part of the site in a suitable way. It is also envisaged that an alteration in the binding site on the antibody for the effector molecule need not alter significantly the overall binding affinity but may alter the geometry of the interaction rendering the effector mechanism ineffective as in non-productive binding. It is further envisaged that an effector function may also be altered by modifying a site  
10 not directly involved in effector molecule binding, but otherwise involved in performance of the effector function. By altering an effector function of an antibody it may be possible to control various aspects of the immune response, e.g. enhancing or suppressing various reactions of the immune system, with possible beneficial effects in diagnosis and therapy.

15 As used herein, the term "TL1A-mediated disorder" includes conditions such as inflammatory diseases and/or auto immune diseases, including *inter alia* inflammatory bowel diseases (e.g., ulcerative colitis and Crohn's disease), rheumatoid arthritis, multiple sclerosis (MS), atherosclerosis, transplant rejection, central nervous system injury, psoriasis, leukemia or lymphoma (e.g., chronic lymphocytic leukemia (CLL)), atherosclerosis, and lung and colon  
20 carcinomas, chronic obstructive pulmonary disease COPD, optic neuritis, age related macular degeneration, systemic lupus erythematosus (SLE), sjogen's syndrome, scleroderma, systemic sclerosis, chronic Kidney disease, liver fibrosis, tuberculosis, idiopathic pulmonary fibrosis, tuberculosis induced lung fibrosis, retroperitoneal Fibrosis, pulmonary fibrosis , cystic fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis (bone marrow),  
25 retroperitoneal fibrosis, progressive massive fibrosis, pephrogenic systemic fibrosis, arthrofibrosis.

As used herein, the term "subject" includes any human or nonhuman animal. The term "nonhuman animal" includes all vertebrates, e.g., mammals and non-mammals, such as  
30 nonhuman primates, sheep, dogs, cats, horses, cows, chickens, amphibians, reptiles, etc. Preferably the subject is human.

### **Anti-TL1A antibodies**

In a first aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and/or  
5 comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.

In some embodiments the antibody or fragment thereof that binds to TL1A comprises an  
10 extended heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 57, and/or an extended heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 58, and/or an extended heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 59; and/or comprises an extended light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 60, and/or an extended light chain CDR2 comprising the amino acid sequence of  
15 SEQ ID NO: 61 and/or an extended light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 62.

Preferably the antibody or fragment thereof that binds to TL1A and comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, a heavy chain CDR2  
20 comprising the amino acid sequence of SEQ ID NO: 52, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53 and/or a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56. More preferably the antibody or fragment thereof that binds to TL1A comprises a heavy  
25 chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53 and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55 and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.  
30 Preferably, the antibody or fragment thereof binds to human TL1A and is cross reactive with murine, rat and cyno TL1A.

It is well known in the art that the CDR3 domain, independently from the CDR1 and/or CDR2

domain(s), alone can determine the binding specificity of an antibody for a cognate antigen and that multiple antibodies can predictably be generated having the same binding specificity based on a common CDR3 sequence. See, for example, Klimka A *et al.*, (2000) Br. J. Cancer, 83(2): 252-260 (describing the production of a humanized anti-CD30 antibody using only the heavy chain variable domain CDR3 of murine anti-CD30 antibody Ki-4); Beiboer SH *et al.*, (2000) J. Mol. Biol. 296: 833-849 (describing recombinant epithelial glycoprotein-2 (EGP-2) antibodies using only the heavy chain CDR3 sequence of the parental murine MOC-31 anti-EGP-2 antibody); Rader C *et al.*, (1998) Proc. Natl. Acad. Sci USA, 95: 8910-8915 (describing a panel of humanized anti-integrin  $\alpha v \beta 3$  antibodies using a heavy and light chain variable CDR3 domain of a murine anti-integrin  $\alpha v \beta 3$  antibody LM609 wherein each member antibody comprises a distinct sequence outside the CDR3 domain and capable of binding the same epitope as the parental murine antibody with affinities as high or higher than the parental murine antibody); Barbas C *et al.*, (1994) J. Am. Chem. Soc. 116: 2161-62 (disclosing that the CDR3 domain provides the most significant contribution to antigen binding).

Accordingly, the present invention provides antibodies and fragments thereof that bind to TL1A comprising one or more heavy and/or light chain CDR3 domains, in particular comprising heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53 and/or light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56, wherein the antibody is capable of binding to TL1A. Within some embodiments, such inventive antibodies comprising one or more heavy and/or light chain CDR3 domain from a non-human antibody (a) are capable of competing for binding with; (b) retain the functional characteristics; (c) bind to the same epitope; and/or (d) have a similar binding affinity as the corresponding parental non-human e.g. murine antibody.

In further aspect the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 1. In further aspect the antibody or fragment thereof comprises a non-CDR region of a heavy chain variable region sequence which is at least 80 % identical to the non-CDR region of the heavy chain variable region sequence of SEQ ID NO: 1.

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable region sequence comprising the amino acid sequence selected from SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28 and SEQ ID NO: 29.

In a further aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 29. In another aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 14. In some embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 29 and a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 14. Preferably, the antibody or fragment thereof binds to human TL1A and is cross reactive with murine, rat and cyno TL1A.

10

In another aspect the present invention provides variants of an antibody or fragment thereof that binds to TL1A. Thus the present invention provides antibodies or fragments thereof that have an amino acid sequence of the non-CDR regions of the heavy and/or light chain variable region sequence which is at least 80% identical (having at least 80% amino acid sequence identity) to the amino acid sequence of the non-CDR regions of the heavy and/or light chain variable region sequence of the parent antibody of either the heavy or the light chain e.g. of either the heavy and light variable region sequences as in SEQ ID NO: 13, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29 or SEQ ID NO: 14, respectively. As well antibodies or fragments thereof that have an amino acid sequence of the non-extended CDR regions of the heavy and/or light chain variable region sequence which is at least 80% identical to the amino acid sequence of the non-extended CDR regions of the heavy and/or light chain variable region sequence of the parent antibody of either the heavy or the light chain are provided by the present invention. Preferably the amino acid sequence identity of the non-CDR regions or of the non-extended CDR regions of the heavy and/or light chain variable region sequence is at least 85%, more preferably at least 90%, and most preferably at least 95%, in particular 96%, more particular 97%, even more particular 98%, most particular 99%, including for example, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and 100%. Identity or homology with respect to an amino acid sequence is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the antibody or fragment thereof that binds to TL1A, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Thus sequence identity can be determined by standard methods that are commonly used to compare the similarity in position of the amino acids of two

30



polypeptides. Using a computer program such as BLAST or FASTA, two polypeptides are aligned for optimal matching of their respective amino acids (either along the full length of one or both sequences or along a pre-determined portion of one or both sequences). The programs provide a default opening penalty and a default gap penalty, and a scoring matrix such as PAM250 (a standard scoring matrix; see Dayhoff MO *et al.*, (1978) in Atlas of Protein Sequence and Structure, vol 5, supp. 3) can be used in conjunction with the computer program. For example, the percent identity can be calculated as: the total number of identical matches multiplied by 100 and then divided by the sum of the length of the longer sequence within the matched span and the number of gaps introduced into the longer sequences in order to align the two sequences.

In some embodiments the present disclosure thus provides an antibody or fragment thereof that binds to TL1A, wherein the antibody or fragment thereof comprises a heavy chain variable framework region sequence which is at least 65 % identical to the framework region sequence of SEQ ID NOS: 3, 4, 5, 6 or 7 and/or a light chain variable framework region sequence which is at least 75 % identical to the framework region sequence of SEQ ID NOS: 8, 9, 10, 11 and 12. In some embodiments the present disclosure provides an antibody or fragment thereof that binds to TL1A, wherein the antibody or fragment thereof comprises a heavy chain variable framework region sequence which is at least 69% identical to the framework region sequence of SEQ ID NO: 3 and/or a light chain variable framework region sequence which is at least 80 % identical to the framework region sequence of SEQ ID NO: 8.

In another aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising the heavy and or light chain CDRs as described *supra* and further comprising a heavy chain variable framework region that is the product of or derived from a human gene selected from the group consisting of IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4), IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID NO: 7), preferably a heavy chain variable framework region that is the product of or derived from human gene IGHV1-2\*01 (SEQ ID NO: 3). The heavy chain variable framework region may comprise one or more (e.g., one, two, three and/or four) heavy chain framework region sequences (e.g., framework 1 (FW1), framework 2 (FW2), framework 3 (FW3) and/or framework 4 (FW4)) present in the product of or derived from those human genes. Preferably the heavy chain variable region framework comprises FW1, FW2 and/or

FW3, more preferably FW1, FW2 and FW3 present in the product of or derived from a human gene selected from the group consisting of IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4), IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID NO: 7). Heavy chain framework region sequences as used herein include FW1  
5 (position 1 to position 25), FW2 (position 36 to position 49), FW3 (position 66 to position 94) and FW 4 (position 103 to position 113), wherein the amino acid position is indicated utilizing the numbering system set forth in Kabat.

10 In some embodiments the present disclosure provides an antibody or fragment thereof, wherein the antibody or fragment thereof comprises a heavy chain variable framework region that is the product of or derived from human gene IGHV1-2\*01 (SEQ ID NO: 3) and wherein the heavy chain variable framework region comprises at least one amino acid modification from the corresponding heavy chain variable framework region of the corresponding murine antibody.

15 In some embodiments the present disclosure provides an antibody or fragment thereof comprising a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 16 and wherein the heavy chain variable framework region comprises at least one amino acid modification from the corresponding heavy chain variable framework region of the corresponding murine antibody.

20 Preferably the amino acid modification comprises an amino acid substitution at amino acid position selected from the group consisting of 37, 48, 50, 67, 69, 71 and 75, more preferably at amino acid positions selected from the group consisting of 37, 48, 50, 67 and 71, most preferred at amino acid position 37, wherein the amino acid position of each group member is  
25 indicated according to the Kabat numbering. Specifically the amino acid modification comprises an amino acid substitution selected from the group consisting of 37A, 48I, 50E, 67A, 69L, 71V and 75S, preferably an amino acid substitution selected from the group consisting of V37A, M48I, W50E, V67A, M69L, R71V and I75S, whereas V37A is the most preferred amino acid substitution wherein the amino acid position of each group member is  
30 indicated according to the Kabat numbering.

In another aspect the present invention provides an antibody or fragment thereof that binds to TL1A comprising a light chain variable framework region that is the product of or derived

from a human gene selected from the group consisting of IGKV1-33\*01 (SEQ ID NO: 8), IGKV1D-33\*01 (SEQ ID NO: 9), IGKV1D-12\*02 (SEQ ID NO: 10), IGKV1D-12\*01 (SEQ ID NO: 11), and IGKV1-12\*02 (SEQ ID NO: 12), preferably a light chain variable framework region that is the product of or derived from human gene IGKV1-33\*01 (SEQ ID NO: 8). The light chain variable region framework region may comprise one or more (e.g., one, two, three and/or four) light chain framework region sequences (e.g., framework 1 (FW1), framework 2 (FW2), framework 3 (FW3) and/or framework 4 (FW4)) present in the product of or derived from those human genes. Preferably the light chain variable region framework comprises FW1, FW2 and/or FW3, more preferably FW1, FW2 and FW3 present in the product of or derived from a human gene selected from the group consisting of IGKV1-33\*01 (SEQ ID NO: 8), IGKV1D-33\*01 (SEQ ID NO: 9), IGKV1D-12\*02 (SEQ ID NO: 10), IGKV1D-12\*01 (SEQ ID NO: 11), and IGKV1-12\*02 (SEQ ID NO: 12). Light chain framework region sequences as used herein include FW1 (position 1 to position 23), FW2 (position 35 to position 49), FW3 (position 57 to position 88) and FW 4 (position 98 to position 108), wherein the amino acid position is indicated utilizing the numbering system set forth in Kabat.

In some embodiments the present disclosure provides an antibody or fragment thereof comprising a light chain variable framework region that is the product of or derived from human gene IGKV1-33\*01 (SEQ ID NO: 8) and wherein the light chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody.

In further aspect the antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 2. In further aspect the antibody or fragment thereof comprises a non-CDR region of a light chain variable region sequence which is at least 80 % identical to the non-CDR region of the heavy chain variable region sequence of SEQ ID NO: 2.

In some embodiments the present disclosure provides an antibody or fragment thereof comprising a light chain sequence comprising the amino acid sequence selected from SEQ ID NO: 17 and SEQ ID NO: 25.

In some embodiments the present disclosure provides an antibody or fragment thereof comprising a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17. Alternatively, the light chain variable framework region of the light chain sequence comprises

at least one amino acid modification from the corresponding light chain variable framework region of the corresponding murine antibody.

The amino acid modification may comprise an amino acid substitution at an amino acid  
5 position selected from the group consisting of 5 and 34, wherein the amino acid position of  
each group member is indicated according to the Kabat numbering. Specifically the amino acid  
modification comprises an amino acid substitution selected from the group consisting of 5N,  
and 34S, preferably T5N and N34S, wherein the amino acid position of each group member is  
indicated according to the Kabat numbering. Particularly preferred is a light chain sequence  
10 comprising the amino acid sequence of SEQ ID NO: 17, without any amino acid modifications.

In some embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy  
chain variable framework region that is the product of or derived from a human gene selected  
from the group consisting of IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4),  
15 IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID  
NO: 7) and a light chain variable framework region that is the product of or derived from a  
human gene selected from the group consisting of IGKV1-33\*01 (SEQ ID NO: 8), IGKV1D-  
33\*01 (SEQ ID NO: 9), IGKV1D-12\*02 (SEQ ID NO: 10), IGKV1D-12\*01 (SEQ ID NO:  
11), and IGKV1-12\*02 (SEQ ID NO: 12), preferably a heavy chain variable framework region  
20 that is the product of or derived from human gene IGHV1-2\*02 (SEQ ID NO: 3), and a light  
chain variable framework region that is the product of or derived from human gene IGKV1-  
33\*01 (SEQ ID NO: 8). As well combinations of heavy chain variable framework regions  
which are present in the product of or derived from different human genes mentioned *supra*  
and/or of light chain variable region framework regions which are present in the product of or  
25 derived from different human genes mentioned *supra* are encompassed by the present  
invention.

Germline DNA sequences for human heavy and light chain variable region genes can be found  
in the "VBase" human germline sequence database (available on the Internet at  
30 [www.mrcpe.cam.ac.uk/vbase](http://www.mrcpe.cam.ac.uk/vbase)), as well as in Kabat EA *et al.*, *supra*; Tomlinson IM *et al.*,  
(1992) J. Mol. Biol. 227: 776-798 and Cox JPL *et al.*, (1994) Eur. J. Immunol. 24: 827-836.  
As another example, the germline DNA sequences for human heavy and light chain variable  
region genes can be found in the Genbank database.

In another aspect, the present disclosure also provides an antibody or fragment thereof that binds to TL1A, wherein at least one of the heavy chain CDRs and/or at least one of the light chain CDRs comprises at least one amino acid modification. Site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the modification(s) and the effect on antibody binding, or other functional property of interest, can be evaluated in *in vitro* or *in vivo* assays. Preferably conservative modifications are introduced. The modification(s) may be amino acid substitutions, additions or deletions, but are preferably substitutions. Typically, no more than five, preferably no more than four, more preferably no more than three, even more preferably no more than two, most preferably no more than one amino acid modifications are performed within a CDR region.

In certain embodiments, framework sequences can be used to engineer variable regions to produce variant antibodies. Variant antibodies of the invention include those in which modifications have been made to framework residues within VH and/or VK, e.g. to improve the properties of the antibody. Typically such framework modifications are made to decrease the immunogenicity of the antibody. For example, one approach is to “backmutate” one or more framework residues to the corresponding murine sequence or to “backmutate” one or more framework residues to a corresponding germline sequence.

Thus in a further aspect the present disclosure provides an antibody or fragment thereof that binds to TL1A, wherein at least one of the framework region sequences of the heavy chain variable region of the antibody or fragment thereof comprises at least one amino acid modification from the corresponding framework region of the heavy chain variable region of the corresponding murine antibody. Preferably the amino acid modification is an amino acid substitution. Typically, no more than seven, preferably no more than six, preferably no more than five, preferably no more than four, more preferably no more than three, even more preferably no more than two, most preferably no more than one amino acid modifications are performed within a framework region. In some embodiments the present disclosure provides an antibody or fragment thereof that binds to TL1A, wherein the amino acid modification of the framework regions of the heavy chain variable region comprise an amino acid substitution at amino acid position selected from the group consisting of 37, 48, 50, 67, 69, 71, and 75 and wherein the amino acid position of each group member is indicated according to the Kabat

numbering. Preferred amino acid substitution of the framework regions of the heavy chain variable region are at amino acid positions selected from the group consisting of 37, 48, 50, 67 and 71. More preferred amino acid substitutions of the framework regions of the heavy chain variable region are selected from the group consisting of V37A, M48I, W50E, V67A, M69L, R71V and I75S, whereas V37A is the most preferred amino acid substitution of the framework regions of the heavy chain variable region.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A, wherein at least one of the framework region sequences of the light chain variable region of the antibody or fragment thereof may comprise at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody. Preferably the amino acid modification is an amino acid substitution. Typically, no more than two, more preferably no more than one and most preferably, no amino acid modifications are performed within a framework region. In some embodiments the present disclosure provides an antibody or fragment thereof, wherein the amino acid modification of the framework regions of the light chain variable region sequence comprises an amino acid substitution at amino acid position selected from the group consisting of 5 and 34. The amino acid modifications of the framework regions of the light chain variable region sequence comprise a substitution selected from the group consisting of a 5N and 34S, preferably T5N and N34S and wherein the amino acid position of each group member is indicated according to the Kabat numbering. In some embodiments the antibody or fragment thereof of the present invention may comprise amino acid modifications of the framework regions of the heavy chain variable region sequence as set out above and amino acid modifications of the framework regions of the light chain variable region sequence as set out above.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A that comprises a heavy chain variable region selected from the group consisting of SEQ ID NOS: 13, 26, 27, 28 and 29, preferably selected from the group consisting of SEQ ID NOS: 26, 27, 28 and 29, more preferably from the group consisting of SEQ ID NOS: 27, 28 and 29 and even more preferably from the group consisting of SEQ ID NOS: 27 and 29. The present disclosure also provides an antibody or fragment thereof that binds to TL1A that comprises a light chain variable region selected from the group consisting of SEQ ID NOS: 14 and 30, more preferably SEQ ID NO: 14. In some embodiments the antibody or fragment

thereof that binds to TL1A comprises a heavy chain variable region selected from the group consisting of SEQ ID NOS: 26, 27, 28 and 29, and a light chain variable region selected from the group consisting of SEQ ID NOS: 14 and 30. Given that each of these heavy and light chain variable region sequences can bind to TL1A, the heavy and light chain variable region sequences can be "mixed and matched" to create anti-TL1A binding molecules of the invention. TL1A binding of such "mixed and matched" antibodies can be tested using the binding assays described e.g. in the Examples.

In some embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain variable region selected from the group consisting of SEQ ID NOS: 27 and 29, and a light chain variable region selected from the group consisting of SEQ ID NOS: 14 and 30. In more preferred embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 27 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14 or a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 29 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14. Most preferred is an antibody or fragment thereof that binds to TL1A comprising a heavy chain variable region selected from the group consisting of SEQ ID NOS: 27 and 29, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A that comprises a heavy chain sequence selected from the group consisting of SEQ ID NOS: 16, 21, 22, 23 and 24, preferably selected from the group consisting of SEQ ID NOS: 22, 23 and 24 and more preferably from the group consisting of SEQ ID NOS: 22 and 24. The present disclosure also provides an antibody or fragment thereof that binds to TL1A that comprises a light chain sequence selected from the group consisting of SEQ ID NOS: 17 and 25, more preferably SEQ ID NO: 17. In some embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain sequence selected from the group consisting of SEQ ID NOS: 21, 22, 23 and 24, and a light chain sequence selected from the group consisting of SEQ ID NOS: 17 and 25. Given that each of these heavy and light chain variable region sequences can bind to TL1A, the heavy and light chain variable region sequences can be "mixed and matched" to create anti-TL1A binding molecules of the invention. TL1A binding of such

"mixed and matched" antibodies can be tested using the binding assays described e.g. in the Examples.

5 In some embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain sequence selected from the group consisting of SEQ ID NOS: 22 and 24, and a light chain sequence selected from the group consisting of SEQ ID NOS: 17 and 25. In more preferred embodiments the antibody or fragment thereof that binds to TL1A comprises a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 22 and a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17 or a heavy chain sequence  
10 comprising the amino acid sequence of SEQ ID NO: 24 and a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17. Most preferred is an antibody or fragment thereof that binds to TL1A comprising a heavy chain sequence selected from the group consisting of SEQ ID NOS: 22 and 24, and a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17.

15 In one embodiment of the present disclosure, the antibody or fragment thereof is a humanized antibody. Preferably, the antibody or fragment thereof is a humanized monoclonal antibody.

The present disclosure also provides a monovalent antibody or fragment thereof that binds to  
20 TL1A, i.e. an antibody which consists of a single antigen binding arm. The present disclosure also provides a fragment of a antibody that binds to TL1A selected from the group consisting of Fab, Fab', Fab'-SH, Fd, Fv, dAb , F(ab')<sub>2</sub>, scFv, bispecific single chain Fv dimers, diabodies, triabodies and scFv genetically fused to the same or a different antibody. Preferred fragments are scFv, bispecific single chain Fv dimers and diabodies. The present disclosure also provides  
25 a full length antibody that binds to TL1A.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A which further comprises a heavy and/or light constant region in particular a human heavy and/or a human light constant region. Human heavy constant regions may be selected from the group of  
30 human immunoglobulins consisting of IgG1 (IGHG1), IgG2 (IGHG2), IgG3 (IGHG3), IgG4 (IGHG4), IgA1 (IGHA1), IgA2 (IGHA2), IgM (IGHM), IgD (IGHD), or IgE (IGHE), whereas the human heavy constant region IgG, in particular IgG1 (IGHG1) is preferred. Human light constant region may be selected from the group of human immunoglobulins



consisting of kappa or lambda constant regions, whereas human kappa constant region is preferred. In a preferred embodiment the antibody or fragment thereof that binds to TL1A comprises a human IgG1 (IGHG1) heavy constant domain and a human light kappa constant domain.

5

In addition or alternative to modifications made within the framework regions or CDR regions, antibodies of the invention may be engineered to include modifications within the Fc region, typically to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding, and/or antigen-dependent cellular cytotoxicity.

10

Furthermore, an antibody of the invention may be chemically modified (e.g., one or more chemical moieties can be attached to the antibody) or be modified to alter its glycosylation.

15

Each of these embodiments is described in further detail below. Modifications within the Fc region as outlined below are according to the EU numbering of residues in the Fc region. In one embodiment, the hinge region of CH1 is modified such that the number of cysteine

20

residues in the hinge region is altered, e.g., increased or decreased. This approach is described further in U.S. Patent No. 5,677,425 by Bodmer *et al.* The number of cysteine residues in the hinge region of CH1 is altered to, for example, facilitate assembly of the light and heavy chains or to increase or decrease the stability of the antibody. In another embodiment, the Fc hinge

25

region of an antibody is mutated to decrease the biological half life of the antibody. More specifically, one or more amino acid mutations are introduced into the CH2-CH3 domain interface region of the Fc-hinge fragment such that the antibody has impaired Staphylococcal protein A (SpA) binding relative to native Fc-hinge domain SpA binding. This approach is

30

described in further detail in U.S. Patent No. 6,165,745 by Ward *et al.* In another embodiment, the antibody is modified to increase its biological half life. Various approaches are possible. For example, one or more of the following mutations can be introduced: T252L, T254S, T256F, as described in U.S. Patent No. 6,277,375 to Ward. Alternatively, to increase the biological half

35

life, the antibody can be altered within the CH1 or CL region to contain a salvage receptor binding epitope taken from two loops of a CH2 domain of an Fc region of an IgG, as described in U.S. Patent Nos. 5,869,046 and 6,121,022 by Presta *et al.* In a further embodiment Fc region is altered by replacing at least one amino acid residue with a different amino acid

40


residue to alter the effector function(s) of the antibody. For example, one or more amino acids selected from amino acid residues 234, 235, 236, 237, 297, 318, 320 and 322 can be replaced with a different amino acid residue such that the antibody has an altered affinity for an effector

ligand but retains the antigen-binding ability of the parent antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Patent Nos. 5,624,821 and 5,648,260, both by Winter *et al.* In another example, one or more amino acids selected from amino acid residues 329, 331 and 322 can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Patent Nos. 6,194,551 by Idusogie *et al.* In another example, one or more amino acid residues within amino acid positions 231 to 238 in the N-terminal region of the CH2 domain are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in PCT Publication WO94/29351 by Bodmer *et al.* In yet another example, the Fc region is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or to increase the affinity of the antibody for an Fcγ receptor by modifying one or more amino acids at the following positions: 238, 239, 248, 249, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 301, 303, 305, 307, 309, 312, 315, 320, 322, 324, 326, 327, 329, 330, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 378, 382, 388, 389, 398, 414, 416, 419, 430, 434, 435, 437, 438 or 439. This approach is described further in PCT Publication WO00/42072 by Presta.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A comprising human heavy and/or light constant regions, wherein the human heavy constant region comprises an isotypic variant comprising the CH1 region, the hinge region, the CH2 region and CH3 region from human IgG4 (IGHG4) and wherein the hinge region comprises a substitution of serine at position 228 to proline. Preferably the antibody comprising the isotypic variant is a full length antibody. A particular preferred antibody or fragment thereof that binds to TL1A comprising an isotypic variant comprising the CH1 from human IgG4 (IGHG4), the hinge from human IgG4 (IGHG4), having S228P substitution and the CH2 and CH3 from human IgG4 (IGHG4). It has been found that the isotypic variant exhibits no Fc-mediated cytotoxicity mechanisms such as ADCC compared to an antibody or fragment thereof that binds to TL1A which comprises a human heavy constant region from human IgG1 (IGHG1) (which is usually a native human IgG1), i.e. as compared to an antibody or fragment thereof

that binds to TL1A that only differs from the isotypic variant with regard to the modified heavy constant region.

The present disclosure also provides an antibody or fragment thereof that binds to TL1A which  
 5 comprises a human IgG Fc region, wherein the mature core carbohydrate structure attached to the human IgG Fc region lacks fucose (referred herein alternatively as “non fucosylated”). The term “mature core carbohydrate structure” as used herein includes a processed core carbohydrate structure attached to an Fc region which generally consists of the carbohydrate structure GlcNAc (Fucose)-GlcNAc-Man-(Man-GlcNAc)<sub>2</sub> typical of biantennary  
 10 oligosaccharides represented schematically below:



This term specifically includes G-1 forms of the core mature carbohydrate structure lacking a  
 β1,2 GlcNAc residue. Preferably, however, the core carbohydrate structure includes both β1,2  
 20 GlcNAc residues. The mature core carbohydrate structure herein generally is not hypermannosylated. The mature core carbohydrate structure is attached to the Fc region of the glycoprotein, generally via N-linkage to Asn297 of a CH2 domain of the Fc region.

Preferably the antibody comprises a human IgG1 (IGHG1) Fc region, wherein the mature core  
 25 carbohydrate structure attached to the human IgG1 (IGHG1) Fc region lacks fucose. More preferred is a full-length antibody comprising a human IgG1 (IGHG1) Fc region, wherein the mature core carbohydrate structure attached to the human IgG1 (IGHG1) Fc region lacks fucose. It is known from WO03/035835 that lack of fucose in the mature core carbohydrate structure attached to the human IgG Fc region may enhance ADCC. Thus in a further  
 30 embodiment the antibody or fragment thereof of the present disclosure comprises a human IgG1 (IGHG1) Fc region, wherein the mature core carbohydrate structure attached to the human IgG1 (IGHG1) Fc region lacks fucose, whereas the antibody lacking fucose exhibits enhanced ADCC compared to the parent antibody or fragment thereof not lacking fucose.

Methods to generate antibodies which lack fucose are, for example (a) use of an engineered or mutant host cell that is deficient in fucose metabolism such that it has a reduced ability (or is unable to) fucosylate proteins expressed therein; (b) culturing cells under conditions which prevent or reduce fucosylation; (c) post-translational removal of fucose (e. g. with a fucosidase enzyme); (d) post-translational addition of the desired carbohydrate, e. g. after recombinant expression of a non-glycosylated glycoprotein; or (e) purification of the glycoprotein so as to select for product which is not fucosylated. Preferably used are methods described in Example 14 of WO10/095031 e.g. methods described in Longmore *et al.*, (1982) Carbohydr. Res. 365-92 or in Imai-Nishiya *et al.*, (2007), BMC Biotechnol. 7: 84.

Also provided by the present invention is an antibody or fragment thereof that binds to TL1A and which binds to the same epitope as the antibody comprising the heavy chain variable sequence comprising the amino acid sequence of SEQ ID NO. 27 or 29 and the light chain variable sequence comprising the amino acid sequence of SEQ ID NO. 14. Also provided by the present invention is a specific region or epitope of TL1A, which is bound by an antibody provided by the present invention, in particular by an antibody comprising the heavy chain variable sequence comprising the amino acid sequence of SEQ ID NO. 27 or SEQ ID NO 29 and the light chain variable sequence comprising the amino acid sequence of SEQ ID NO. 14. This specific region or epitope of the TL1A polypeptide can be identified by any suitable epitope mapping method known in the art in combination with any one of the antibodies provided by the present invention. Examples of such methods include screening peptides of varying lengths derived from TL1A for binding to the antibody of the present invention with the smallest fragment that can specifically bind to the antibody containing the sequence of the epitope recognised by the antibody. The TL1A peptides may be produced synthetically or by proteolytic digestion of the TL1A polypeptide. Peptides that bind the antibody can be identified by, for example, mass spectrometric analysis. In another example, NMR spectroscopy or X-ray crystallography can be used to identify the epitope bound by an antibody of the present invention. Once identified, the epitopic fragment which binds an antibody of the present invention can be used, if required, as an immunogen to obtain additional antibodies which bind the same epitope.

#### **Anti-TL1A antibody properties**

Standard assays to evaluate the binding ability of the antibodies toward e.g. TL1A are known

in the art, including for example, ELISAs, BIAcore<sup>®</sup>, Western blots, RIAs, and flow cytometry analysis. Suitable assays are described in detail in the Examples. The binding kinetics (e.g., binding affinity like  $K_D$ ) of the antibodies also can be assessed by standard assays known in the art, such as by Scatchard or BIAcore<sup>®</sup> system analysis. The relative binding affinity  $K_i$  can be assessed by standard competition assays known in the art.

5

In a further aspect the present invention provides antibodies or fragment thereof that bind to human, mouse, rat and cynomologus monkey TL1A as visualized by ELISA or BIAcore<sup>®</sup> methods. Binding ELISA can be carried out and measured according to Example 3.

10

In a further aspect the present invention provides antibodies or fragments thereof that bind to recombinant or naturally produced human TL1A and prevent activation and cytokine secretion by CD4 T lymphocytes. For example, the antibodies or fragments thereof of the invention may suppress the production of  $INF\gamma$  induced by immune complex stimulated monocytes. An assay to determine such TL1A-mediated cytokine secretion by CD4 T lymphocytes can be carried out and measured according to Examples 3 and 6.

15

In a further aspect the present invention provides antibodies or fragment thereof that bind to TL1A, in particular TL1A in isolated form, with an affinity ( $K_D$ ) of 850 pM or less, preferably 700nM or less, more preferably 300 nM or less, more preferably 260 nM or less, even more preferably 250 nM or less, e.g. measured by Surface Plasmon Resonance (SPR) on a BIAcore<sup>®</sup> instrument (GE Healthcare Europe GmbH, Glattbrugg, Switzerland) by capturing the antibody on a protein-A coupled CM5 research grade sensor chip (GE Healthcare Europe GmbH, Glattbrugg, Switzerland; BR-1000-14) with a human soluble TL1A polypeptide (encoded by SEQ ID NO: 116) used as analyte as detailed in Example 5. In a preferred aspect, the present invention provides a humanized antibody or fragment thereof that retains at least 85% of the TL1A binding affinity ( $K_D$ ) of the corresponding chimeric antibody. Preferably the humanized antibody or fragment thereof retains at least 90% of the TL1A binding affinity ( $K_D$ ) of the corresponding chimeric antibody, more preferably at least 95% of the binding affinity ( $K_D$ ) of the corresponding chimeric antibody. Preferably, the humanized antibody or fragment thereof binds human TL1A with equivalent affinity to the corresponding chimeric antibody. By “equivalent affinity” is meant an affinity value that is within a range of  $\pm 10\%$  of the TL1A binding affinity of the corresponding chimeric antibody. More preferably, the present invention

20

25

30

provides a humanized antibody or fragment thereof that binds human TL1A with a higher affinity than the corresponding chimeric antibody. Preferably the humanized antibody or fragment thereof binds human TL1A with two-fold higher affinity than the corresponding chimeric antibody, more preferably with three-fold higher affinity than the corresponding chimeric antibody. In a preferred aspect of the present invention, humanized antibodies or fragment thereof that bind to human TL1A are provided that have a binding affinity ( $K_D$ ) of 900 pM or less, 700 pM or less, preferably 500 pM or less, more preferably 300 nM or less, more preferably 260 pM or less, even more preferably 250 pM or less e.g. measured by Surface Plasmon Resonance (SPR) on a BIAcore<sup>®</sup> instrument (GE Healthcare Europe GmbH, Glattbrugg, Switzerland) by capturing the antibody on a protein-A coupled CM5 research grade sensor chip (GE Healthcare Europe GmbH, Glattbrugg, Switzerland; BR-1000-14) with a human soluble TL1A polypeptide (encoded by SEQ ID NO: 116) used as analyte as detailed in Example 5.

A further aspect of the present invention provides antibodies or fragments thereof that bind to TL1A and which have good thermal stability. In a preferred embodiment, an antibody or fragment thereof that binds to TL1A has a FAB fragment thermostability temperature greater than 70°C, preferably greater than 75°C and even more preferably greater than 80°C. For analysis of FAB fragment thermostability differential scanning calorimetry measurements are used, whereas a mid-point melting temperature of the FAB fragment in context of a full-length IgG is identified. These kind of calorimetric measurements are known to the skilled person and can be carried out according to e.g. Garber E & Demarest SJ (2007) *Biochem Biophys Res Commun*, 355: 751-7, as further described in Example 5.

25

### **Nucleic acids, Vectors and Host Cells**

The present disclosure also provides isolated nucleic acids encoding the antibodies and fragments thereof that bind to TL1A, vectors and host cells comprising the nucleic acid or the vector. The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is "isolated" or "rendered substantially pure" when  
5 purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art, see e.g. F. Ausubel, *et al.*, ed. (1987) *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York. A nucleic acid of the invention can be, for example, DNA or  
10 RNA and may or may not contain intron sequences. In a preferred embodiment, the nucleic acid is a cDNA molecule.

Nucleic acids of the invention can be obtained using standard molecular biology techniques e.g. cDNAs encoding the light and heavy chains of the antibody or encoding VH and VL segments  
15 can be obtained by standard PCR amplification or cDNA cloning techniques. For antibodies obtained from an immunoglobulin gene library (e.g., using phage display techniques), one or more nucleic acids encoding the antibody can be recovered from the library. The methods of introducing exogenous nucleic acid into host cells are well known in the art, and will vary with the host cell used. Techniques include but are not limited to dextran- mediated transfection,  
20 calcium phosphate precipitation, calcium chloride treatment, polyethylenimine mediated transfection, polybrene mediated transfection, protoplast fusion, electroporation, viral or phage infection, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. In the case of mammalian cells, transfection may be either transient or stable.

25 Preferred nucleic acids molecules of the invention are those encoding the heavy chain sequence selected from the group consisting of SEQ ID NOS: 42, 43, 44, 45 and 46 and/or the light chain sequence selected from the group consisting of SEQ ID NOS: 47 and 48. Preferred nucleic acids molecules of the invention are those encoding the heavy chain variable region selected from the group consisting of SEQ ID NOS: 31, 32, 33, 34 and 35 and/or the light  
30 chain variable region selected from the group consisting of SEQ ID NOS: 36 and 37.

Preferred nucleic acids molecules of the invention are those encoding the heavy chain variable region of SEQ ID NO: 1 and/or the light chain variable region of SEQ ID NO: 2, e.g. DNA

encoding the heavy chain variable region comprising the nucleic acid sequence of SEQ ID NO: 63 and/or DNA encoding the light chain variable region comprising the nucleic acid sequence of SEQ ID NO: 64. More preferred nucleic acid molecules of the invention are those encoding the heavy chain variable region of SEQ ID NOS: 27 or 29 and/or the light chain variable region of SEQ ID NO: 14, e.g. DNA encoding the heavy chain variable region comprising the nucleic acid sequence of SEQ ID NOS: 33 or 35 and/or DNA encoding the light chain variable region comprising the nucleic acid sequence of SEQ ID NO: 36, which are most preferred.

Once DNA fragments encoding VH and VL segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, or to fragments genes corresponding to the fragments described *supra* like Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker.

The term “operatively linked”, as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame. The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat EA *et al.*, *supra*) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1 (IGHG1), IgG2 (IGHG2), IgG3 (IGHG3), IgG4 (IGHG4), IgA1 (IGHA1), IgA2 (IGHA2), IgM (IGHM), IgD (IGHD), or IgE (IGHE) constant region, but most preferably is an IgG1 (IGHG1) constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region. The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., Kabat EA *et al.*, *supra.*) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. In preferred embodiments, the light chain constant region can be a kappa or lambda constant region, preferably a kappa constant region. To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another



fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly<sub>4</sub>-Ser)<sub>3</sub>, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird RE *et al.*, (1988) *Science*, 242: 423-426; Huston JS *et al.*, (1988) *Proc. Natl. Acad. Sci. USA*, 85: 5879-83; McCafferty J *et al.*, (1990) *Nature*, 348: 552-554). Various techniques have been developed for the production of antibody fragments of antibodies. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto K *et al.*, (1992) *J. Biochem. & Biophysical Methods*, 24: 107-117 and Brennan M *et al.*, (1985) *Science*, 229: 81-3). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from E. coli and chemically coupled to form F(ab')<sub>2</sub> fragments (Carter P *et al.*, (1992) *Bio/ Technology*, 10: 163-167). According to another approach, F(ab')<sub>2</sub> fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single-chain Fv fragment (scFv), see e.g. WO 1993/16185; U.S. Patent. No. 5,571,894 and U.S. Patent No. 5,587,458. The antibody fragment may also be a "linear antibody", e.g., as described in U.S. Patent No. 5,641,870, for example.

The nucleic acids that encode the antibodies of the present invention may be incorporated into a vector, preferably an expression vector in order to express the protein. A variety of expression vectors may be utilized for protein expression. Expression vectors may comprise self-replicating extra- chromosomal vectors or vectors which integrate into a host genome. Expression vectors are constructed to be compatible with the host cell type. Thus vectors, preferably expression vectors, which find use in the present invention include but are not limited to those which enable protein expression in mammalian cells, bacteria, insect cells, yeast, and in *in vitro* systems. As is known in the art, a variety of expression vectors are available, commercially or otherwise, that may find use in the present invention for expressing antibodies.

Expression vectors typically comprise a protein operably linked with control or regulatory sequences, selectable markers, any fusion partners, and/or additional elements. By "operably linked" herein is meant that the nucleic acid is placed into a functional relationship with another

nucleic acid sequence. The term “regulatory sequence” is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel (Gene Expression Technology, Methods in Enzymology  
5 185, Academic Press, San Diego, CA (1990)). Generally, these expression vectors include transcriptional and translational regulatory nucleic acid operably linked to the nucleic acid encoding the antibody, and are typically appropriate to the host cell used to express the protein. In general, the transcriptional and translational regulatory sequences may include promoter sequences, ribosomal binding sites, transcriptional start and stop sequences,  
10 translational start and stop sequences, and enhancer or activator sequences. As is also known in the art, expression vectors typically contain a selection gene or marker to allow the selection of transformed host cells containing the expression vector. Selection genes are well known in the art and will vary with the host cell used. For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into  
15 which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr- host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

Suitable host cells for cloning or expressing the DNA in the vectors herein are prokaryote,  
20 yeast, or higher eukaryote cells. Suitable prokaryotes for this purpose include eubacteria, including gram-negative or gram-positive organisms, for example, *Enterobacteriaceae* such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis*, *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. Suitable  
25 *E. coli* cloning hosts include *E. coli* 294 (ATCC 31,446), *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325). In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly  
30 available and useful, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts including *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. WaltH* (AJCC 56,500), *K. drosopmarum* (ATCC 36,906), *K. thermotolerans*, or *K. marxianusyarrowia* (EP402226); *Pichia pastoris* (EP183070); *Candida*; *Trichoderma*

*reesia* (EP244234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi including *Neurospora*, *Penicillium*, *Tolyocladium*, or *Aspergillus* hosts such as *A. nidulans* or *A. niger*.

- 5 Suitable host cells for the expression of the antibodies of the invention are derived from multicellular organisms. Examples of invertebrate cells include plaril and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes augypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly) and *Bombyx mori* have been identified. A
- 10 variety of viral strains for transfection are publicly available, for example, the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.
- 15 Host cells for expressing the recombinant antibodies of the invention are preferably mammalian host cells which include Chinese Hamster Ovary (CHO cells) (including dhfr<sup>-</sup> CHO cells, described in Urlaub G & Chasin LA (1980) Proc. Natl. Acad. Sci, USA, 77: 4216-4220, used with a DHFR selectable marker, e.g., as described in Kaufman RJ & Sharp PA (1982) J. Mol. Biol, 159: 601-621), NSO myeloma cells, COS cells and SP2 cells. In particular, for use with
- 20 NSO myeloma cells, another preferred expression system is the GS gene expression system disclosed in WO 87/04462 (to Wilson), WO 89/01036 (to Bebbington) and EP338841 (to Bebbington). When recombinant antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, for secretion of the antibody
- 25 into the culture medium in which the host cells are grown. Host cells useful for producing antibodies that bind to TL1A may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland), Minimal Essential Medium (MEM; Sigma-Aldrich Chemie GmbH), RPMI-1640 (Sigma-Aldrich Chemie GmbH, Basel, Switzerland), and Dulbecco's Modified Eagle's Medium ((DMEM; Sigma-
- 30 Aldrich Chemie GmbH) are suitable for culturing the host cells. Antibodies can be recovered from the culture medium using standard protein purification methods.

Antibodies may be operably linked to a fusion partner to enable targeting of the expressed protein, purification, screening, display, and the like. Fusion partners may be linked to the antibody sequence via a linker sequences. The linker sequence will generally comprise a small number of amino acids, typically less than ten, although longer linkers may also be used.

5 Typically, linker sequences are selected to be flexible and resistant to degradation. As will be appreciated by those skilled in the art, any of a wide variety of sequences may be used as linkers. For example, a common linker sequence comprises the amino acid sequence GGGGS. A fusion partner may be a targeting or signal sequence that directs antibody and any associated fusion partners to a desired cellular location or to the extracellular media. As is known in the art, certain signalling sequences may target a protein to be either secreted into the growth  
10 media, or into the periplasmic space, located between the inner and outer membrane of the cell. A fusion partner may also be a sequence that encodes a peptide or protein that enables purification and/or screening. Such fusion partners include but are not limited to polyhistidine tags (His-tags) (for example H6 and H10 or other tags for use with Immobilized Metal Affinity  
15 Chromatography (IMAC) systems (e.g. Ni<sup>2+</sup> affinity columns)), GST fusions, MBP fusions, Strep-tag, the BSP biotinylation target sequence of the bacterial enzyme BirA, and epitope tags which are targeted by antibodies (for example c-myc tags, flag-tags, and the like). As will be appreciated by those skilled in the art, such tags may be useful for purification, for screening, or both.

20

### **Construction and Production of Antibodies**

Antibodies generated against the TL1A polypeptide may be obtained by immunisation of an animal i.e. by administering the polypeptides to an animal, preferably a non-human animal, using well-known and routine protocols, see for example Handbook of Experimental  
25 Immunology (Weir DM (ed.), Vol 4, Blackwell Scientific Publishers, Oxford, England, 1986). Many warm-blooded animals, such as rabbits, mice, rats, sheep, cows, camels or pigs may be immunized. However, mice, rabbits, pigs and rats in particular mice are generally most suitable. Antibodies can be produced as well by recombinant DNA techniques known to the skilled person. In additional antibodies can be produced by enzymatic or chemical cleavage of  
30 naturally occurring antibodies. Humanized antibodies of the present invention may be constructed by transferring one or more CDRs or portions thereof from VH and/or VL regions from a non-human animal (e.g., mouse) to one or more framework regions from human VH and/or VL regions. Optionally, human framework residues thus present in the VH and/or VL

regions may be replaced by corresponding non-human (e.g., mouse) residues when needed or desired for decreasing immunogenicity of the antibody and/or maintaining binding affinity.

Optionally, non-human amino acid residues present in the CDRs may be replaced with human residues. Chimeric or humanized antibodies of the present invention can be prepared based on  
5 the sequence of a non-human monoclonal antibody prepared as described above. DNA encoding the heavy and light chain immunoglobulins can be obtained from the non- human hybridoma of interest and engineered to contain non-murine (e.g., human) immunoglobulin sequences using standard molecular biology techniques. For example, to create a chimeric antibody, murine variable regions can be linked to human constant regions using methods  
10 known in the art (see e.g., U.S. Patent No. 4,816,567 to Cabilly *et al*). To create a humanized antibody, murine CDR regions can be inserted into a human framework using methods known in the art (see e.g., U.S. Patent No. 5,225,539 to Winter, and U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen *et al*).

15 Humanized antibodies of the present invention may be constructed wherein the human acceptor molecule for the heavy chain variable region is selected based on homology considerations between potential acceptor molecule variable regions and the heavy chain variable region of the murine antibody. Germline candidate human acceptor molecules are preferred to reduce potential immunogenicity. Germline databases are made up of antibody  
20 sequences that read through the end of the heavy chain FW3 region and partially into the CDR3 sequence. For selection of a FW4 region, databases of mature antibody sequences which have been derived from the selected germline molecule can be searched or antibody sequences which have been derived from the selected germline molecule from a human donor can be used. Human acceptor molecules are preferably selected from the same heavy chain  
25 class as the murine donor molecule, and of the same canonical structural class of the variable region of the murine donor molecule. Secondary considerations for selection of the human acceptor molecule for the heavy chain variable region elude homology in CDR length between the murine donor molecule and the human acceptor molecule. Human acceptor antibody molecules are preferably selected by homology search to the V-BASE database, although other  
30 databases such as the Kabat and the public NCBI databases may be used as well.

Humanized antibodies of the present invention may be constructed wherein the human acceptor molecule for the light chain variable region is selected based on homology

considerations between potential acceptor molecule variable regions and with the light chain variable region of the murine antibody. Germline candidate human acceptor molecules are preferred to reduce potential immunogenicity. Germline databases are made up of antibody sequences that read through the end of the heavy chain FW3 region and partially into the CDR3 sequence. For selection of a FW4 region, databases of mature antibody sequences which have been derived from the selected germline molecule can be searched or antibody sequences which have been derived from the selected germline molecule from a human donor can be used. Human acceptor molecules are preferably selected from the same light chain class as the murine donor molecule, and of the same canonical structural class of the variable region of the murine donor molecule. Secondary considerations for selection of the human acceptor molecule for the light chain variable region include homology in CDR length between the murine donor molecule and the human acceptor molecule. Human acceptor antibody molecules are preferably selected by homology searches to the V-BASE database, and other databases such as the Kabat and the public NCBI databases may be used as well. Methods for humanizing a non-human antibody are described herein, including in Example 5, below.

The present invention provides a method of producing an antibody or fragment thereof that binds to TL1A comprising culturing a host cell comprising an isolated nucleic acid encoding the antibody or fragment thereof that binds to TL1A or a vector comprising an isolated nucleic acid encoding the antibody or fragment thereof that binds to TL1A so that the nucleic acid is expressed and the antibody produced. Preferably the antibody is isolated. For host cells, nucleic acids and vectors, the ones described above can be used. Expression of the nucleic acids can be obtained by, e.g. a combination of recombinant DNA techniques and gene transfection methods as is well known in the art (e.g., Morrison S (1985) Science 229: 1202) and as further outlined above. For example, to express the antibodies, or antibody fragments thereof, DNAs encoding partial or full-length light and heavy chains, can be obtained by standard molecular biology techniques (e.g., PCR amplification or cDNA cloning using a hybridoma that expresses the antibody of interest) and the DNAs can be inserted into vectors such as expression vectors. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or, more typically, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the

antibody gene fragment and vector, or blunt end ligation if no restriction sites are present). The light and heavy chain variable regions of the antibodies described herein can be used to create full-length antibody genes of any antibody isotype by inserting them into expression vectors already encoding heavy chain constant and light chain constant regions of the desired isotype such that the VH segment is operatively linked to the CH1 segment(s) within the vector and the VK segment is operatively linked to the CK segment within the vector.

### **Characterization and Purification of Anti-TL1A antibodies**

Screening for antibodies can be performed using assays to measure binding to TL1A and/or assays to measure the ability to block the binding of TL1A to its receptor TNFRSF25. An example of a binding assay is an ELISA, in particular, using a fusion protein of TL1A and human Fc, which is immobilized on plates, and employing a conjugated secondary antibody to detect anti-TL1A antibody bound to the fusion protein. An example of a blocking assay is a flow cytometry based assay measuring the blocking of TL1A fusion protein binding to TNFRSF25 on human CD4 cells. A fluorescently labelled secondary antibody is used to detect the amount of TL1A fusion protein binding to the cell. This assay is looking for a reduction in signal as the antibody in the supernatant blocks the binding of ligand fusion protein to TNFRSF25. A further example of a blocking assay is an assay where the blocking of costimulation of naive human T cells mediated by TL1A fusion protein coated to a plate is measured by measuring thymidine incorporation. As an assay for evaluating the functional activity of anti-TL1A antibodies e.g. the reduction of cytokine secretion by CD4 T lymphocytes as described in Examples 3 and 6 can be used.

Antibodies of the present invention may be isolated or purified in a variety of ways known to those skilled in the art. Standard purification methods include chromatographic techniques, including ion exchange, hydrophobic interaction, affinity, sizing or gel filtration, and reversed-phase, carried out at atmospheric pressure or at high pressure using systems such as FPLC and HPLC. Purification methods also include electrophoretic, immunological, precipitation, dialysis, and chromatofocusing techniques. Ultrafiltration and diafiltration techniques, in conjunction with protein concentration, are also useful. To purify TL1A antibodies, selected host cells can be grown in e.g. spinner-flasks for monoclonal antibody purification. Supernatants can be filtered and concentrated before affinity chromatography with protein A-sepharose (Pharmacia, Piscataway, NJ). Eluted antibodies can be checked by gel

electrophoresis and high performance liquid chromatography to ensure purity. A preferred antibody of the present invention is thus an isolated and/or purified antibody that binds to TL1A.

## 5 **Immunoconjugates**

In another aspect, the present invention provides a TL1A antibody or a fragment thereof that binds to TL1A, linked to a therapeutic agent, such as a cytotoxin, a drug (e.g., an immunosuppressant) or a radiotoxin. Such conjugates are referred to herein as "immunoconjugates". Immunoconjugates that include one or more cytotoxins are referred to as "immunotoxins." A cytotoxin or cytotoxic agent includes any agent that is detrimental to (e.g., kills) cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents also include, for example, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). Other examples of therapeutic cytotoxins that can be linked to an antibody of the invention include duocarmycins, calicheamicins, maytansines and auristatins, and derivatives thereof. An example of a calicheamicin antibody conjugate is commercially available (Mylotarg(R); American Home Products). Cytotoxins can be linked to antibodies of the invention using linker technology available in the art. Examples of linker types that have been used to conjugate a cytotoxin to an antibody include, but are not limited to, hydrazones, thioethers, esters, disulfides and peptide-containing linkers. A linker can be chosen that is, for example, susceptible to cleavage by low pH within the lysosomal compartment or susceptible to cleavage by proteases, such as proteases preferentially expressed in tumor tissue such as cathepsins (e.g., cathepsins B, C, D). For further discussion of types of cytotoxins, linkers and methods for conjugating therapeutic agents to antibodies, see also Saito G *et al.*, (2003) *Adv. Drug Deliv. Rev.* 55: 199-215; Trail



PA *et al.*, (2003) *Cancer Immunol. Immunother.* 52: 328-337; Payne G (2003) *Cancer Cell*, 3: 207-212; Allen TM (2002) *Nat. Rev. Cancer*, 2: 750-763; Pastan I & Kreitman RJ (2002) *Curr. Opin. Investig. Drugs*, 3: 1089-1091; Senter PD & Springer CJ, (2001) *Adv. Drug Deliv. Rev.* 53: 247-264. Antibodies of the present invention also can be linked to a radioactive isotope to generate cytotoxic radiopharmaceuticals, also referred to as radioimmunoconjugates. Examples of radioactive isotopes that can be conjugated to antibodies for use diagnostically or therapeutically include, but are not limited to, iodine<sup>131</sup>, indium<sup>111</sup>, yttrium<sup>90</sup> and lutetium<sup>177</sup>. Methods for preparing radio-immunconjugates are established in the art. Examples of radioimmunoconjugates are commercially available, including Zevalin<sup>®</sup> (EDEC Pharmaceuticals) and Bexxar<sup>®</sup> (Corixa Pharmaceuticals) and similar methods can be used to prepare radioimmunoconjugates using the antibodies of the invention. The antibody immunoconjugates of the invention can be used to modify a given biological response, and the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an enzymatically active toxin, or active fragment thereof, such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor or interferon- $\gamma$ ; or, biological response modifiers such as, for example, lymphokines, interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), or other growth factors.

Techniques for linking such therapeutic agents to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.*, (eds.), pp. 243- 56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.*, (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.*, (eds.), pp. 475-506 (1985); "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy", in *Monoclonal Antibodies for Cancer Detection and Therapy*, Baldwin *et al.*, (eds.), pp. 303-16 (Academic Press 1985), and Thorpe PE & Ross WC (1982) *Immunol. Rev.* 62: 119-58.

In another aspect, the present invention provides a TL1A antibody or a fragment thereof that binds to TL1A, administered together with a therapeutic agent, such as a cytotoxin, a drug (e.g., an immunosuppressant) or a radiotoxin.

5

### Pharmaceutical Compositions

In another aspect, the present invention provides a composition, e.g., a pharmaceutical composition, comprising the antibody or fragment thereof, of the present invention, and a pharmaceutically acceptable carrier. Such compositions may include one or a combination of (e.g., two or more different) antibodies, and/or immunoconjugates of the invention and/or a therapeutic agent, such as a cytotoxin, a drug (e.g., an immunosuppressant) or a radiotoxin as described *supra*. For example, a pharmaceutical composition of the invention can comprise a combination of antibodies (or immunoconjugates) that bind to different epitopes on the target antigen or that have complementary activities. Pharmaceutical compositions of the invention also can be administered in combination therapy, i.e., combined with other agents. For example, the combination therapy can include a TL1A antibody of the present invention combined with at least one other anti-inflammatory or immunosuppressant agent.

As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound, i.e., antibody or immunoconjugate, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the invention is contemplated.

Supplementary active compounds can also be incorporated into the compositions.

In another aspect, the present invention provides a composition comprising an

immunoconjugate comprising the antibody or fragment thereof that binds to TL1A linked to a therapeutic agent and a pharmaceutically acceptable carrier. Immunoconjugates and therapeutic agents which can be used are as described *supra*.

5 In another aspect, the present invention provides a composition comprising the antibody or fragment thereof of the present invention which further comprises another pharmaceutically active agent. Preferably the another pharmaceutically active agent is one or more of: a) another antagonist to TL1A, b) an anti-inflammatory agent, c) an immune suppressive agent e.g. TNF $\alpha$  antagonist, cortisone or steroids etc) and/or d) an anti-allergy agent.

10 A pharmaceutical composition of the invention may also include a pharmaceutically acceptable antioxidant. Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil- soluble antioxidants, such as ascorbyl  
15 palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic-acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like. Examples of suitable aqueous and nonaqueous carriers that may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol,  
20 propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying  
25 agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, *supra*, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be  
30 brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

### **Therapeutic and other uses**

The antibodies of the present invention have numerous *in vitro* and *in vivo* diagnostic and therapeutic utilities involving the diagnosis and treatment of TL1A mediated disorders. For example, these molecules can be administered to cells in culture, *in vitro* or *ex vivo*, or to human subjects, e.g., *in vivo*, to treat, prevent and to diagnose a variety of TL1A-mediated disorders. Preferred subjects are human and include patients having disorders mediated by TL1A activity (TL1A mediated disorders). The neutralizing antibodies of the present invention can be effective in treating patients independent of their TL1A costimulatory status. More preferred subjects are human and include patients expressing a high level of TL1A.

A “patient” for the purposes of the present invention includes both humans and other animals, preferably mammals and most preferably humans. Thus the antibodies of the present invention have both human therapy and veterinary applications. The term “treatment” or “treating” in the present invention is meant to include therapeutic treatment, as well as prophylactic, or suppressive measures for a disease or disorder. Thus, for example, successful administration of an antibody prior to onset of the disease results in treatment of the disease. As another example, successful administration of an antibody after clinical manifestation of the disease to combat the symptoms of the disease comprises treatment of the disease. “Treatment” and “treating” also encompasses administration of an antibody after the appearance of the disease in order to eradicate the disease. Successful administration of an antibody after onset and after clinical symptoms have developed, with possible abatement of clinical symptoms and perhaps amelioration of the disease, comprises treatment of the disease. Those “in need of treatment” include mammals already having the disease or disorder, as well as those prone to having the disease or disorder, including those in which the disease or disorder is to be prevented.

In a particular embodiment, the antibodies are used *in vivo* to treat, prevent or diagnose a variety of TL1A-mediated disorders. Thus the invention provides a method for treating a TL1A mediated disorder in a subject, the method comprising administering to the subject a therapeutically effective amount of the antibody or fragment thereof. Exemplary TL1A mediated disorders include, but are not limited to, inflammatory diseases and/or autoimmune diseases, for example, inflammatory bowel disease (IBD) including ulcerative colitis and Crohn’s disease, rheumatoid arthritis, MS, type 1 and type 2 diabetes, psoriasis, psoriatic arthritis, ankylosing spondylitis, atopic dermatitis; allergic reactions or conditions, including for example, asthma and allergic lung inflammation; cancers, atherosclerosis, infections,

neurodegenerative diseases, graft rejection, graft versus host diseases (GVHD) and cardiovascular disorders/ diseases. The invention also provides a method for treating a TL1A mediated disorder in a subject, the method comprising administering to the subject a therapeutically effective amount of the antibody or fragment thereof. Exemplary TL1A mediated disorders include, but are not limited to, inflammatory diseases and/or autoimmune diseases, for example, inflammatory bowel disease (IBD) including ulcerative colitis and Crohn's disease, rheumatoid arthritis, MS, type 1 and type 2 diabetes, psoriasis, psoriatic arthritis, ankylosing spondylitis, atopic dermatitis; allergic reactions or conditions, including for example, asthma and allergic lung inflammation; cancers, atherosclerosis, infections, neurodegenerative diseases, graft rejection, graft versus host diseases (GVHD) and cardiovascular disorders/ diseases, chronic obstructive pulmonary disease COPD, optic neuritis, age related macular degeneration, systemic lupus erythematosus (SLE), sjogén's syndrome, scleroderma, systemic sclerosis, chronic Kidney disease, liver fibrosis, tuberculosis, idiopathic pulmonary fibrosis, tuberculosis induced lung fibrosis, retroperitoneal Fibrosis, pulmonary fibrosis , cystic fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis (bone marrow), retroperitoneal fibrosis, progressive massive fibrosis, pephrogenic systemic fibrosis, arthrofibrosis. Preferably, the TL1A mediated disorders include inflammatory diseases and/or auto immune diseases, including *inter alia* inflammatory bowel diseases (e.g., ulcerative colitis and Crohn's disease), rheumatoid arthritis, MS and atherosclerosis.

Preferred TL1A mediated disorders to be treated with the antibody of the invention are selected from the group consisting of inflammatory bowel disease, multiple sclerosis, rheumatoid arthritis and asthma. A particular preferred TL1A mediated disorder to be treated with the antibody of the invention is inflammatory bowel disease.

Animal model for evaluating the functional activity of anti-TL1A antibodies in TL1A-mediated disorders are described in Example 7 for asthma and in Examples 8 and 9 for IBD.

In one embodiment, the antibodies of the invention can be used to detect levels of TL1A, or levels of cells which contain TL1A on their membrane surface, which levels can then be linked to certain disease symptoms. Alternatively, the antibodies can be used to inhibit or block TL1A function which, in turn, can be linked to the prevention or amelioration of certain disease

symptoms, thereby implicating TL1A as a mediator of the disease. This can be achieved by contacting a sample and a control sample with the TL1A antibody under conditions that allow for the formation of a complex between the antibody and TL1A. Any complexes formed between the antibody and TL1A are detected and compared in the sample and the control. In light of the specific binding of the antibodies of the invention for TL1A, the antibodies of the invention can be used to specifically detect TL1A expression on the surface of cells e.g. can be used to detect a patient having low or high expression levels of TL1A. The antibodies of the invention can also be used to purify TL1A via immunoaffinity purification.

10 In another embodiment, the antibodies of the invention can be initially tested for binding activity associated with therapeutic or diagnostic use *in vitro*. For example, compositions of the invention can be tested using flow cytometric assays.

The present disclosure further provides the use of an antibody or fragment thereof as a medicament and the use of an antibody or fragment thereof in the preparation of a medicament for the treatment of a TL1A mediated disorder. In a further embodiment the present disclosure provides the antibody or fragment thereof for use as a medicament. Also provided by the present disclosure is the antibody or fragment thereof for use in a method for treating a TL1A mediated disorder. TL1A mediated disorders are the ones as described *supra*. The antibody or fragment thereof of the present invention may be particularly useful for treating TL1A mediated disorders independent of the DR3 costimulatory status of a patient. In a preferred embodiment, the antibody or fragment thereof can be used for treating a TL1A mediated disorder wherein a patient expresses a high level of TL1A.

25 As previously described, anti-TL1A antibodies of the invention can be co-administered with one or other more therapeutic agents, e.g., a cytotoxic agent, a radiotoxic agent or an immunosuppressive agent. The antibody can be linked to the agent (as an immunoconjugate as described *supra*) or can be administered separate from the agent. In the latter case (separate administration), the antibody can be administered before, after or concurrently with the agent or can be co-administered with other known therapies, e.g., an anti-cancer therapy, e.g., radiation.

For administration of the antibody, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 10 mg/kg, of the host body weight. An exemplary treatment regime entails administration once per week, once every two weeks, once every three weeks, once every four weeks, once a month, once every three months or once every three to six months.

5 The antibody is usually administered on multiple occasions. Intervals between single dosages can be, for example, weekly, monthly, every three months or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to the target antigen in the patient. In some methods, dosage is adjusted to achieve a plasma antibody concentration of about 1-1000  $\mu\text{g/ml}$  and in some methods about 25-300  $\mu\text{g/ml}$ . Alternatively the antibody can be  
10 administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients  
15 continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated.

Actual dosage levels of the active ingredients, i.e. the antibody in the pharmaceutical  
20 compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the  
25 time of administration, the rate of excretion of the particular antibody being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

30 A "therapeutically effective amount" of a TL1A antibody of the invention preferably results in a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, and/or a prevention of impairment or disability due to the disease affliction. The ability of a compound for the treatment of a TL1A mediated disorder can be

evaluated in an animal model system predictive of efficacy in human. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit cell growth, such inhibition can be measured *in vitro* by assays known to the skilled practitioner. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

The antibody or the composition of the present invention can be administered via one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Preferred routes of administration include intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other parenteral routes of administration, for example by injection or infusion. More preferred routes of administration are intravenous or subcutaneous. The phrase "parenteral administration" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion. Alternatively, an antibody of the invention can be administered via a non- parenteral route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically.

#### **Article of manufacture and kit**

In another embodiment of the disclosure, an article of manufacture comprising the antibody or fragment thereof, the composition or the immunoconjugate of the invention for the treatment of a TL1A mediated disorder is provided. The article of manufacture may comprise a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials or syringes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that may be effective for treating the condition and may have a sterile access port (e.g., the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition may be the antibody described herein. The label or package insert may indicate that the composition may be used for treating the



condition of choice, such as cancer. In one embodiment, the label or package insert may indicate that the composition comprising the antibody may be used to treat a TL1A -mediated disorder.

5 Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises the antibody herein, and (b) a second container with a composition contained therein, wherein the composition comprises a therapeutic agent other than the antibody. The article of manufacture in this embodiment of the disclosure may further comprise a package insert indicating that the first and second  
10 compositions can be used in combination to treat a TL1A mediated disease or disorder. Such therapeutic agent may be any of the adjunct therapies described in the preceding section (e.g., a thrombolytic agent, an anti-platelet agent, a chemotherapeutic agent, an anti-angiogenic agent, an anti-hormonal compound, a cardioprotectant, and/or a regulator of immune function in a mammal, including a cytokine). Alternatively, or additionally, the article of manufacture may  
15 further comprise a second (or third) container comprising a pharmaceutically acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

20 Also within the scope of the present invention are kits comprising the antibody, the compositions or the immunoconjugates of the invention and instructions for use. The kit can further contain one more additional reagents, such as an immunosuppressive reagent, a cytotoxic agent or a radiotoxic agent, or one or more additional antibodies of the invention (e.g., an antibody having a complementary activity which binds to an epitope in the TL1A  
25 antigen distinct from the first antibody).

Without further description, it is believed that one of ordinary skill in the art may, using the preceding description and the following illustrative examples, make and utilize the agents of the present disclosure and practice the claimed methods. The following working examples are  
30 provided to facilitate the practice of the present disclosure, and are not to be construed as limiting in any way the remainder of the disclosure.

## Examples

### **Example 1:**

#### **Generation and screening of mouse anti-human TL1A antibodies**

5 To produce the recombinant human TL1A-Fc protein, a cDNA for the human TL1A gene was purchased from Source BioScience (Nottingham, UK; clone number: IRATp970G02115D). This cDNA was used as a template to amplify the coding region of the processed secreted version of human TL1A (SEQ ID NO: 116) using PCR. The PCR was performed using primers GlnPr994 and GlnPr995 (SEQ ID NOs: 119 and 120, respectively). Primer GlnPr994  
10 adds a BamHI restriction site 5' of the extracellular region and cleaves the native signal peptide. Primer GlnPr995 adds a HindIII restriction site 3' of the extracellular region. The amplicon was cut using the flanking restriction sites BamHI and HindIII and cloned into a modified mammalian expression vector based on the pcDNA3.1(-) plasmid from Invitrogen (Invitrogen AG, Basel, Switzerland), expressing an Fc-fusion construct. The expression vector  
15 contains the human CMV promoter with the Ig donor acceptor fragment (first intron) described in U.S. Patent 5,924,939, the OriP sequence (Koons MD *et al.*, (2001) J Virol. 75(22): 10582-92), the SV40 enhancer, and the SV40 polyA fused to the gastrin terminator as described by Kim D, *et al.*, (2003) Biotechnol. Prog. 19(5): 1620-2. The expression cassette contains a kozak region upstream of the open reading frame of the Fc fusion protein, followed  
20 by a signal peptide, terminated by a BamHI restriction site for convenient cloning. The Fc region of the fusion protein is started by a HindIII restriction site for convenient cloning, followed by a small glycine-serine linker. In order to release the previous construct upstream of the Fc region, the vector was cut using BamHI (NEB, Ipswich, MA, USA) and HindIII (NEB, Ipswich, MA, USA), treated using CIP (NEB, Ipswich, MA, USA) and gel purified. The insert  
25 coding for the extracellular region of human TL1A was ligated in the backbone and transformed in *E.coli* Top 10 cells (Life Technologies, Carlsbad, CA, USA) leading to the Fc fusion expression construct (SEQ ID NO: 117). The first 20 amino acids of SEQ ID NO: 117 correspond to the signal sequence, which was not present in the final Fc fusion expression construct.

30

This recombinant plasmid allowed for expression of the human TL1A-Fc fusion protein in mammalian cells with secretion into the cell culture medium driven by the signal peptide. For recombinant protein production, the aforementioned recombinant vector was transfected into

suspension-adapted HEK 293 EBNA cells (Life Technologies, Carlsbad, CA, USA) using jetPEI<sup>TM</sup> transfection reagent (Polyplus-transfection S.A., Strasbourg, France; Distributor: Brunschwig, Basel, Switzerland). The cell culture supernatant was collected after five days and further purified using a Protein A affinity purification column (HiTrap Protein A sepharose  
5 column; GE Healthcare Europe GmbH, Glattbrugg, Switzerland) operated on an ÄKTA FPLC system (GE Healthcare Europe GmbH, Glattbrugg, Switzerland).

To produce the secreted recombinant human TL1A protein with his-tag, the DNA sequence coding for the processed version of secreted human TL1A was amplified by PCR using primers  
10 GlnPr1542 and GlnPr1543 (SEQ ID NOs: 121 and 122, respectively) adding an N-terminal 6xHis linker. A second round of PCR using primers GlnPr1544 (SEQ ID NO: 123) and GlnPR1543 adds a signal peptide and convenient flanking restriction sites for cloning (5': NheI; 3': XhoI). The PCR product was cut using NheI and XhoI (NEB, Ipswich, MA, USA) and subsequently cloned in the modified pcDNA3.1(-) plasmid described above, cut using the  
15 same enzymes and CIPed. This restriction digest releases the open reading frame of the entire Fc-fusion protein previously present in the expression vector. After ligation and transformation in Top 10 *E. coli* cells, the final plasmid was chosen based on restriction digest and sequencing of the expression construct for secreted TL1A with an N-terminal his-tag (SEQ ID No: 118). The first 20 amino acids of SEQ ID NO: 118 correspond to the signal sequence, which was not  
20 present in the final N-terminal his-tag expression construct.

This recombinant plasmid allowed for the expression of the human TL1A-his protein in mammalian cells with secretion into the cell culture media. For protein production, the recombinant vector was transfected into suspension-adapted HEK 293 EBNA cells (Life  
25 Technologies, Carlsbad, CA, USA) using jetPEI<sup>TM</sup> transfection reagent (Polyplus-transfection S.A., Strasbourg, France; Distributor: Brunschwig, Basel, Switzerland). The cell culture supernatant was collected five days after transfection and purified using a Ni<sup>2+</sup>-NTA affinity purification column (HiTrap Ni<sup>2+</sup>-NTA sepharose column; GE Healthcare Europe GmbH, Glattbrugg, Switzerland) operated on an ÄKTA FPLC system (GE Healthcare Europe GmbH,  
30 Glattbrugg, Switzerland). The buffer of the recombinant human TL1A-Fc and TL1A-his proteins was changed into phosphate buffer saline (PBS). Recombinant human TL1A-Fc protein dissolved in PBS was mixed with an equal volume of Stimune adjuvant (Prionics, Switzerland) and an emulsion was prepared. The emulsion was transferred to 0.5 mL insulin

syringes (BD Pharmingen, Allschwil, Switzerland) and BALB/c animals (Harlan, Netherlands) were immunized sub-cutaneously in the back footpads, the base of the tail and the neck with 50 µg of the emulsified protein. The immunization was repeated two weeks later with the same amount of antigen and the same route of injection.

5

The presence of circulating anti-human TL1A antibodies in the immunized mouse sera was evaluated by direct ELISA using plates coated with the recombinant human TL1A-his protein. A serial dilution (from 1:10<sup>0</sup> to 1:10<sup>9</sup>) of the different mouse sera was added to the plates and the bound antibodies were detected using a goat anti-mouse H+L whole molecule-HRP  
10 (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland). A final sub-cutaneous boost with 50 µg of antigen without adjuvant was performed in animals displaying the best anti-human TL1A IgG serum titer three days before sacrifice.

15

Animals were euthanized and the inguinal, axillary, brachial, popliteal and sciatic lymph nodes were collected to prepare a single cell suspension by disturbing the lymph node architecture with two 25G needles in a DNase (Roche Diagnostics (Schweiz) AG, Rotkreuz, Switzerland) and collagenase (Roche Diagnostics (Schweiz) AG, Rotkreuz, Switzerland) solution. Single cell suspensions were fused to a myeloma cell line X63AG8.653 (mouse BALB/c myeloma cell line; ATCC accession number: CRL 1580; Kearney JF *et al.*, (1979) J. Immunol. 123(4):  
20 1548-1550) at a ratio of 7:1 (fusion partner-to-harvested lymph node cells) with polyethylene glycol 1500 (Roche Diagnostics (Schweiz) AG, Rotkreuz, Switzerland). The fused cells were plated into 96 well flat bottom plates containing mouse macrophages in DMEM-10 medium (Invitrogen AG, Basel, Switzerland) supplemented with 10% fetal bovine serum (FBS, PAA Laboratories, Pasching, Austria), 2mM L-glutamine, 100U/ml (Biochrom AG, Germany)  
25 penicillin, 100 µg/ml streptomycin (Biochrom AG, Germany), 10mM HEPES (Invitrogen AG, Basel, Switzerland), 50 µM β-mercaptoethanol (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland), HAT (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) and 1% Growth factor (Hybridokine, Interchim/Uptima, Montluçon, France).

30

Approximately 800 wells from the fusions were screened by ELISA for the presence of mouse IgG that recognized human TL1A and blocked the binding of human TL1A to its receptor. Positive wells were expanded and subjected to two rounds of sub cloning. Cells were collected and the heavy and light chains were cloned and sequenced.

**Example 2:****Cloning and sequencing of the VH and VL chains of the anti-TL1A antibodies from hybridoma cells**

For each positively selected hybridoma, total RNA was prepared, reverse-transcribed into cDNA and VH and VL genes were respectively amplified by PCR. These PCR products were ligated into a rescue-vector (pDrive vector; QIAGEN AG, Hombrechtikon, Switzerland), allowing for the DNA sequencing of individual PCR products and the determination of mono- or poly-clonality of the selected hybridomas. This vector allowed for blue/white selection on LB-agar plates containing IPTG and X-gal (colonies with no insert were blue because of the degradation of X-gal by the LacZ  $\alpha$ -peptide). Recombinant plasmids from positive (white) bacterial clones were prepared and sequenced using standard DNA sequencing primers specific for the vector backbone (M13rev, M13fwd, T7 or SP6). DNA sequences were finally subcloned into an expression vector for recombinant expression of the antibody of interest in mammalian cells.

15

*RNA isolation*

Total RNA was isolated from  $2-10 \times 10^6$  cells using the RNeasy Mini Kit from QIAGEN (QIAGEN AG, Hombrechtikon, Switzerland) according to the manufacturer's protocol; samples were quantified using a NanoDrop ND-1000 spectrophotometer (WITEC AG, Littau, Switzerland).

20

*One step RT-PCR*

The total RNA preparations described above were further reverse-transcribed into cDNA, and the VH and VL fragments were amplified by PCR using two different mixtures of degenerated primers, each one allowing the recovery of all the different subfamilies of mouse immunoglobulin heavy chain variable fragments and variable heavy chain junction regions or the recovery of all mouse immunoglobulin light chain kappa variable fragments and variable light chain kappa junction regions. The primers used for reverse transcription and amplification were synthesized by Microsynth (Balgach, Switzerland), and were HPLC purified (Tables 1-4). Both reverse-transcription and PCR amplification were performed simultaneously using the QIAGEN one step RT-PCR kit (QIAGEN AG, Hombrechtikon, Switzerland). Since the technique used specific primers, each mRNA sample was then treated in duplicate allowing for the individual reverse-transcription and amplification of either the VH or the VL fragments.

30

2µg of total RNA dissolved into RNase-free water to a final volume of 30µl were mixed with: 10µl of a 5x stock solution of QIAGEN OneStep RT-PCR Buffer, 2µl of a dNTPs mix at a concentration of 10mM, 3µl of primer mix at a concentration of 10µM and 2µl of QIAGEN OneStep RT-PCR Enzyme Mix. The final mixture was then placed in a PCR tube, and cycled  
5 in a PCR-thermocycler (BioRad iCycler version 4.006, Bio-Rad Laboratories AG, Reinach, Switzerland) using the following settings:

30 min at 50 °C  
15 min at 95 °C  
10  
40 cycles: 30 sec at 94 °C  
30 sec at 55 °C  
1 min at 72 °C  
15  
10 min at 72 °C  
Hold at 4 °C

#### pDrive cloning

PCR products were run onto 2% agarose gels. Following DNA electrophoresis, the fragments  
20 of interest (~450bp) were excised from the agarose gels, and further extracted using the Macherey-Nagel NucleoSpin Extract II kit 250 (Macherey-Nagel, Oensingen, Switzerland). For DNA sequencing, the extracted PCR products were cloned into the rescue-vector described above (pDrive vector, QIAGEN AG, Hombrechtikon, Switzerland) and transformed into the *E. coli* TOP10 strain (Invitrogen AG, Basel, Switzerland).

25

#### Miniprep extraction

Positive colonies were cultured overnight at 37 °C (shaking 250 RPM) in 1.5ml of Luria Bertani (LB) medium supplemented with 100µg/ml ampicillin seeded in Macherey-Nagel Square-well Block plates (Macherey-Nagel, Oensingen, Switzerland). The next day DNA  
30 miniprep extractions were performed using the NucleoSpin Multi-8 Plasmid kit (Macherey-Nagel, Oensingen, Switzerland).

Sequencing and Sequence analysis

Samples were sent for DNA sequencing to the DNA sequencing service company FASTERIS (Plan-les-Ouates, Switzerland). The standard primers: M13rev, M13fwd, T7, SP6 were used (Table 5). To analyse the DNA sequences, the Clone Manager 9 Professional Edition (Scientific & Educational Software, NC, USA) and the BioEdit Sequence Alignment Editor (Hall TA (1999) Nucl Acids Symp Ser 41: 95-98) were used.

Cloning of expression vector for recombinant chimeric antibody expression

For recombinant expression in mammalian cells, the isolated murine VH and VL fragments were formatted as chimeric immunoglobulins using assembly-based PCR methods. These chimeric antibodies consist of a heavy chain where the murine heavy chain variable domain is fused to the human IgG1 heavy chain constant domains ( $\gamma$ 1, hinge,  $\gamma$ 2, and  $\gamma$ 3 regions) and a light chain where the murine light chain variable domain is fused to a human kappa constant domain (C $\kappa$ ). PCR- assembled murine variable and human constant parts were subsequently cloned into a modified mammalian expression vector based on the modified pcDNA3.1(-) vector from Invitrogen mentioned in Example 1 with the difference that a human immunoglobulin light chain kappa leader peptide was employed to drive protein secretion. For protein production of the immunoglobulin candidates, equal quantities of heavy and light chain vector DNA were co-transfected into suspension-adapted HEK-293 (ATCC number: CRL-1573). The cell culture supernatant was collected after five days and purified using a Protein A affinity purification column (HiTrap Protein A sepharose column) operated on an ÄKTA FPLC system (both from GE Healthcare Europe GmbH, Glattbrugg, Switzerland).

**Table 1: primer Mix VH – back**

1	GTG ATC GCC ATG GCG TCG ACC GAK GTR MAG CTT CAG GAG TC	SEQ ID NO: 65
2	GTG ATC GCC ATG GCG TCG ACC GAG GTB CAG CTB CAG CAG TC	SEQ ID NO: 66
3	GTG ATC GCC ATG GCG TCG ACC CAG GTG CAG CTG AAG SAR TC	SEQ ID NO: 67
4	GTG ATC GCC ATG GCG TCG ACC GAG GTC CAR CTG CAA CAR TC	SEQ ID NO: 68
5	GTG ATC GCC ATG GCG TCG ACC CAG GTY CAG CTB CAG CAR TC	SEQ ID NO: 69
6	GTG ATC GCC ATG GCG TCG ACC CAG GTY CAR CTG CAG CAR TC	SEQ ID NO: 70
7	GTG ATC GCC ATG GCG TCG ACC CAG GTC CAC GTG AAG CAR TC	SEQ ID NO: 71
8	GTG ATC GCC ATG GCG TCG ACC GAG GTG AAS STG GTG GAR TC	SEQ ID NO: 72
9	GTG ATC GCC ATG GCG TCG ACC GAV GTG AWG STG GTG GAG TC	SEQ ID NO: 73

10	GTG ATC GCC ATG GCG TCG ACC GAG GTG CAG STG GTG GAR TC	SEQ ID NO: 74
11	GTG ATC GCC ATG GCG TCG ACC GAK GTG CAM CTG GTG GAR TC	SEQ ID NO: 75
12	GTG ATC GCC ATG GCG TCG ACC GAG GTG AAG CTG ATG GAR TC	SEQ ID NO: 76
13	GTG ATC GCC ATG GCG TCG ACC GAG GTG CAR CTT GTT GAR TC	SEQ ID NO: 77
14	GTG ATC GCC ATG GCG TCG ACC GAR GTR AAG CTT CTC GAR TC	SEQ ID NO: 78
15	GTG ATC GCC ATG GCG TCG ACC GAA GTG AAR STT GAG GAR TC	SEQ ID NO: 79
16	GTG ATC GCC ATG GCG TCG ACC CAG GTT ACT CTR AAA SAR TC	SEQ ID NO: 80
17	GTG ATC GCC ATG GCG TCG ACC CAG GTC CAA CTV CAG CAR CC	SEQ ID NO: 81
18	GTG ATC GCC ATG GCG TCG ACC GAT GTG AAC TTG GAA SAR TC	SEQ ID NO: 82
19	GTG ATC GCC ATG GCG TCG ACC GAG GTG AAG GTC ATC GAR TC	SEQ ID NO: 83

**Table 2: primer Mix VH - forward**

1	CCTCCACCACTCGAGCC CGA GGA AAC GGT GAC CGT GGT	SEQ ID NO: 84
2	CCTCCACCACTCGAGCC CGA GGA GAC TGT GAG AGT GGT	SEQ ID NO: 85
3	CCTCCACCACTCGAGCC CGC AGA GAC AGT GAC CAG AGT	SEQ ID NO: 86
4	CCTCCACCACTCGAGCC CGA GGA GAC GGT GAC TGA GGT	SEQ ID NO: 87

**Table 3: primer Mix VL - back**

1	GGCGGTGGC GCT AGC GAY ATC CAG CTG ACT CAG CC	SEQ ID NO: 88
2	GGCGGTGGC GCT AGC CAA ATT GTT CTC ACC CAG TC	SEQ ID NO: 89
3	GGCGGTGGCGCT AGC GAY ATT GTG MTM ACT CAG TC	SEQ ID NO: 90
4	GGCGGTGGC GCT AGC GAY ATT GTG YTR ACA CAG TC	SEQ ID NO: 91
5	GGCGGTGGC GCT AGC GAY ATT GTR ATG ACM CAG TC	SEQ ID NO: 92
6	GGCGGTGGC GCT AGC GAY ATT MAG ATR AMC CAG TC	SEQ ID NO: 93
7	GGCGGTGGC GCT AGC GAY ATT CAG ATG AYD CAG TC	SEQ ID NO: 94
8	GGCGGTGGCGCT AGC GAY ATY CAG ATG ACA CAG AC	SEQ ID NO: 95
9	GGCGGTGGC GCT AGC GAY ATT GTT CTC AWC CAG TC	SEQ ID NO: 96
10	GGCGGTGGCGCT AGC GAY ATT GWG CTS ACC CAA TC	SEQ ID NO: 97
11	GGCGGTGGC GCT AGC GAY ATT STR ATG ACC CAR TC	SEQ ID NO: 98
12	GGCGGTGGC GCT AGC GAY RTT KTG ATG ACC CAR AC	SEQ ID NO: 99
13	GGCGGTGGCGCT AGC GAY ATT GTG ATG ACB CAG KC	SEQ ID NO: 100
14	GGCGGTGGC GCT AGC GAY ATT GTG ATA ACY CAG GA	SEQ ID NO: 101
15	GGCGGTGGC GCT AGC GAY ATT GTG ATG ACC CAG WT	SEQ ID NO: 102
16	GGCGGTGGC GCT AGC GAY ATT GTG ATG ACA CAA CC	SEQ ID NO: 103



17	GGCGGTGGCGCT AGC GAY ATT TTG CTG ACT CAG TC	SEQ ID NO: 104
18	GGCGGTGGC GCT AGC GAA ACA ACT GTG ACC CAG TC	SEQ ID NO: 105
19	GGCGGTGGCGCT AGC GAA AAT GTK CTS ACC CAG TC	SEQ ID NO: 106
20	GGCGGTGGCGCT AGC CAG GCT GTT GTG ACT CAG GAA TC	SEQ ID NO: 107

**Table 4: primer Mix VL - forward**

1	ATGCTGAC GC GGC CGC ACG TTT KAT TTC CAG CTT GG	SEQ ID NO: 108
2	ATGCTGAC GC GGC CGC ACG TTT TAT TTC CAA CTT TG	SEQ ID NO: 109
3	ATGCTGAC GC GGC CGC ACG TTT CAG CTC CAG CTT GG	SEQ ID NO: 110
4	ATGCTGAC GC GGC CGC ACC TAG GAC AGT CAG TTT GG	SEQ ID NO: 111

**Table 5: sequencing primers**

M13-Fwd	GTAAAACGACGGCCAGT	SEQ ID NO: 112
M13-Rev	AACAGCTATGACCATG	SEQ ID NO: 113
T7	TAATACGACTCACTATAGG	SEQ ID NO: 114
SP6	GATTTAGGTGACACTATAG	SEQ ID NO: 115

5

**Example 3:****Biological characterization of anti-human TL1A antibodies***TL1A-specific Antibody Detection ELISA*

Antibody titers, specificity and production by hybridomas and recombinant antibody candidates were determined by a direct ELISA. Briefly, 96 well-microtiter plates (Costar USA, distributor VWR AG, Nyon, Switzerland) were coated with 100 µl of recombinant human TL1A-his at 2 µg/ml in PBS (see Example 1 for the generation of the TL1A-his protein). Plates were incubated overnight at 4°C and were then blocked with PBS 2% BSA (Bovine Serum Albumin, PAA Laboratories, Pasching, Austria) at room temperature (RT) for one hour. The blocking solution was removed and the hybridoma supernatants or purified antibodies were added. The plates were incubated at RT for 30 minutes, then washed nine times with PBS 0.01% Tween-20 (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) and a Horseradish Peroxidase (HRP) labelled-Goat anti-mouse H+L-detection antibody (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) was added at a dilution of 1:1000. To detect recombinant chimeric antibodies (see Example 2) that possess a human Fc, a HRP-labelled rabbit anti

15

20

human IgG antibody (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) at a dilution of 1:1000 was used as the detection antibody. Plates were incubated for 30 minutes at room temperature (RT), washed nine times with PBS 0.01% Tween-20 and the TMB substrate (Bio-rad Laboratories AG, Reinach, Switzerland) was added to the plates and the reaction stopped after two to six minutes by adding H<sub>2</sub>SO<sub>4</sub>. Absorbance was then read at 450 nm by a microplate reader (Biotek, USA; distributor: WITTEC AG, Littau, Switzerland). Figure 1 shows that the parental hybridoma supernatants of various clones recognize the human TL1A-his coated protein and not irrelevant his-tagged protein.

#### 10 *TNFRSF25 blocking ELISA*

The recombinant human TNFRSF25 receptor protein (TNFRSF25) was generated as follows: the cDNA for human TNFRSF25 (clone name: IRCMp5012F0812D) was purchased from Source Biosystems (Nottingham, UK) and the extracellular portion (amino acids 25-199) of human TNFRSF25 (numbering according to the Uniprot Q93038 sequence) was amplified with flanking restriction sites. The resulting PCR product encompassing an N-terminal 8-His tag sequence was subsequently cloned into a modified version of the pcDNA3.1 vector from Invitrogen (Invitrogen AG, Basel, Switzerland) carrying a CMV promoter, a Bovine Growth Hormone poly-adenylation, and the murine VJ2C leader peptide to drive the secretion of the recombinant protein. For recombinant protein production, the recombinant vector was transfected into suspension-adapted HEK 293 cells (ATCC number CRL 1573) using jetPEI<sup>TM</sup> transfection reagent (Polyplus-transfection S.A., Strasbourg, France; Distributor: Brunschwig, Basel, Switzerland). Cell culture supernatant was collected after five days and purified using a Protein A affinity purification column (HiTrap Protein A sepharose column; GE Healthcare Europe GmbH, Glattbrugg, Switzerland) operated on an ÄKTA FPLC system (GE Healthcare Europe GmbH, Glattbrugg, Switzerland).

In order to determine if the generated anti-TL1A antibodies can block the binding of TL1A to the TNFRSF25 receptor, a blocking ELISA was developed. Ninety-six well-microtiter plates (Costar, USA; distributor VWR AG, Nyon, Switzerland) were coated with 100 µl of recombinant human TL1A-Fc (see Example 1) at 2 µg/ml in PBS or recombinant untagged TL1A (R&D Systems, Minneapolis, USA). Plates were incubated overnight at 4°C and were then blocked with PBS 2% BSA at RT for one hour. The blocking solution was removed and the hybridoma supernatants or purified antibodies were added to the plate. Five minutes later,

50  $\mu$ l of recombinant human TNFRSF25-Fc-his (R&D Systems) at 4  $\mu$ g/ml was added to each well. Plates were incubated at RT for 60 minutes, then washed nine times with PBS 0.01% Tween-20 and mouse anti-poly histidine-HRP (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) was added at a dilution of 1:2000. Plates were incubated for 30 minutes at RT, washed 9 times with PBS 0.01% Tween-20 and the TMB substrate (Bio-rad Laboratories AG, Reinach, Switzerland) was added to the plates and the reaction stopped after 6 minutes by adding H<sub>2</sub>SO<sub>4</sub>. Absorbance was then read at 450 nm by a microplate reader (Biotek, USA; distributor: WITTEC AG, Littau, Switzerland). Figure 2 shows that the purified antibodies are able to block the interaction between TL1A and TNFRSF25 in a dose dependent manner.

10

*Inhibition of TL1A-induced IFN- $\gamma$  secretion by primed CD4 T cells*

CD4 T cells primed by IL-12 and IL-18 cytokines polarize toward a TH1 phenotype and secrete IFN- $\gamma$ . TL1A has been shown to enhance IFN- $\gamma$  production by primed CD4 T cells. We therefore tested if the chimeric 5G6 antibody could block this TL1A-dependent increase in IFN- $\gamma$  production.

15

To purify human CD4 T cells from peripheral blood mononuclear cells (PMBC), filters containing human leukocytes were collected from the Blood Collection Centre from La Chaux-de-Fonds, Switzerland (Centre de Transfusion Sanguine et Laboratoire de Sérologie, rue Sophie-Mairet 29, CH-2300). Cells were removed from the filters by back flushing with 60 mL of PBS containing 10U/mL of liquemin (Drossapharm AG, Lucern, Switzerland). PMBCs were then purified with 50mL Blood-Sep-Filter Tubes (distributor: Brunschwig, Basel, Switzerland) following manufacturer's instructions. Cells were washed three times with phosphate buffered saline (PBS) and then used for CD4 purification using naïve CD4 T cell purification kit from Miltenyi (Gladbach, Germany) according to the manufacturer's instructions.

25

The 5G6 antibody was tested to determine whether it could inhibit the effect of naturally produced TL1A. IL-12 and IL-18 primed CD4 T cells were incubated with monocytes that had been pre-activated by immune complexes (IC), and the production of IFN- $\gamma$  was measured. Monocytes were isolated from PMBCs (see above) using a monocyte isolation kit II from Miltenyi (Gladbach, Germany) according to the manufacturer's instructions. Monocyte IC stimulation was performed as follows: chrompure human IgG (Jackson ImmunoResearch

30

Europe Ltd, Newmarket, UK) was coated on a 12-well cell-culture plate (TPP, Trasadingen, Switzerland) at 50µg/mL for 2 hrs at room temperature in PBS. The plate was then washed with PBS and incubated with mouse anti-human IgG (Jackson ImmunoResearch) for 1 hr at room temperature. The coated plate was washed once with PBS before plating of the purified monocytes. IC-stimulated monocytes were harvested after 48-72 hrs incubation at 37°C in a 5% CO<sub>2</sub> incubator.

For flow cytometry analysis, IC-stimulated monocytes were stained with an anti-TL1A-PE antibody (GeneTex, distributed by Lucerna Chem. AG, Lucerna, Switzerland) or a rabbit isotype control (BD Pharmingen, Allschwil, Switzerland) at 10µg/mL in a V-bottom 96-well microtiter plate (TPP, Trasadingen, Switzerland) at 4°C for 30min. The dilution buffer (FACS buffer) was PBS supplemented with 2% fetal calf serum (FCS, Amimed distributed by Bioconcept, Allschwil, Switzerland) and 10% Versene (Gibco Life Technologies). After incubation, 100µL of FACS buffer was added to each well and the plate was centrifuged at 300g for 3min. Supernatant was discarded and samples were resuspended in 100µL of a PE-anti rabbit secondary antibody solution at 0.2µg/mL in FACS buffer. Samples were incubated at 4°C for 20min. The plate was washed as described above and samples were resuspended in 300µL of FACS buffer and immediately acquired on a FACSCyan flow cytometer (Beckman Coulter International S.A., Nyon, Switzerland). For soluble TL1A quantification, supernatant from IC-stimulated monocyte culture were harvested at different time points (20, 48 and 72 hours) and sTL1A was quantified using human TL1A development ELISA kit (Peprotech), according to the manufacturer's recommendations.

For the monocyte-T cell co-culture, IC stimulated monocytes (10<sup>4</sup>) and CD4 purified T cells (10<sup>5</sup>) were seeded in flat bottom-96 well plates (TPP) with IL-12 (Peprotech) at 8ng/mL and IL-18 (MBL International, distributed by LabForce AG, Nunningen, Switzerland) at 200ng/mL in a CO<sub>2</sub> incubator at 37°C. The supernatants of culture were harvested after 72 hrs and IFN-γ was quantified as described above.

IC stimulated monocytes expressed TL1A on their membrane (mTL1A) (Figure 3A) but more substantially as a soluble factor (sTL1A) (Figure 3B) and induced a strong production of IFN-γ by co-cultured CD4 T cells (Figure 3C). The 5G6 antibody suppressed completely the

production of IFN- $\gamma$  induced by the IC-stimulated monocytes showing a potent blocking of both mTL1A and sTL1A-mediated effect.

**Example 4:**

**Parental 5G6 candidate binds to mouse, rat, cynomologus monkey and human TL1A**

5 The reactivity of the parental 5G6 antibody (produced as a chimeric antibody with Human Fc) on the extracellular part of TL1A from diverse species was tested by ELISA. The extracellular part of TL1A protein corresponding to human (*homo sapiens*), rat (*ratus norvegicus*), mouse (*mus musculus*) and cynomologus monkey (*macaca fascicularis*) sequences (SEQ ID NOS: 38, 39, 40 and 41, respectively) was immobilized on high binding 96-well plates (Costar, USA; distributor VWR AG, Nyon, Switzerland) overnight at a concentration of 2 $\mu$ g/mL at 4°C in 10 PBS. Plates were blocked with 2% BSA albumin (BSA, Sigma-Aldrich Chemie, Buchs, Switzerland) at RT for one hour. The blocking solution was removed and a dose dilution of 5G6 antibody was applied to the plates. Plates were incubated at RT for 60 minutes, washed six times with PBS 0.01% Tween-20 and the TMB substrate (Bio-rad Laboratories AG, 15 Reinach, Switzerland) was added to the plates. The reaction was stopped after 6 minutes by adding H<sub>2</sub>SO<sub>4</sub> and the binding of 5G6 on the different TL1A proteins was revealed using an HRP-labelled anti human IgG secondary antibody (Sigma-Aldrich Chemie, Buchs, Switzerland), added at a dilution of 1:1000. The plates were read for absorbance at 450nm by a microplate reader (Biotek, USA; distributor: WITTEC AG, Littau, Switzerland). Figure 4 20 shows that 5G6 recognized, in a dose dependent manner, the TL1A protein from all tested species.

**Example 5:**

**Humanization of mouse monoclonal antibody 5G6**

25 Humanizing the anti-human TL1A mouse antibody 5G6 including selection of human acceptor frameworks, back mutations, and mutations that substantially retain and/or improve the binding properties of human CDR-grafted acceptor frameworks is described herein.

Design of the reshaped variable regions

30 Homology matching was used to choose human acceptor frameworks to graft 5G6 CDRs. Databases e.g. a database of germline variable genes from the immunoglobulin loci of human and mouse (the IMGT database, *supra*) or the VBASE2 (Retter I *et al.*, (2005) Nucleic Acids Res. 33, Database issue D671-D674) or the Kabat database (Johnson G *et al.*, (2000) Nucleic

Acids Res. 28: 214-218) or publications (e.g., Kabat EA *et al.*, *supra*) may be used to identify the human subfamilies to which the murine heavy and light chain V regions (SEQ ID NO: 1 and 2, respectively) belong and determine the best-fit human germline framework to use as the acceptor molecule. Selection of heavy and light chain variable sequences (VH and VL) within these subfamilies to be used as acceptor may be based upon sequence homology and/or a match of structure of the CDR1 and CDR2 regions to help preserve the appropriate relative presentation of the six CDRs after grafting.

For example, use of the IMGT database indicates good homology between the 5G6 heavy chain variable domain framework and the members of the human heavy chain variable domain subfamily 1. Highest homologies and identities of both CDRs and framework sequences were observed for germline sequences: IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4), IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID NO: 7), all of which had sequence identity above 67% for the whole sequence up to CDR3. IGHV1-2\*02 and IGHV1-2\*04 showed 69.4% sequence identity while IGHV1-2\*01 and IGHV1-46\*01 showed a sequence identity of 68.4 and 67.3%, respectively.

Using the same approach, 5G6 light chain variable domain sequence showed good homology to the members of the human light chain variable domain kappa subfamily 1. Highest homologies and identities of both CDRs and framework sequences were observed for germline sequences: IGKV1-33\*01 (SEQ ID NO: 8) and IGKV1D-33\*01 (SEQ ID NO: 9) exhibited the highest identity (both having 80.0% identity), closely followed by another group consisting of IGKV1D-12\*02 (SEQ ID NO: 10), IGKV1D-12\*01 (SEQ ID NO: 11), and IGKV1-12\*02 (SEQ ID NO: 12) all exhibiting the same degree of sequence identity (75.8%).

As starting point to the humanization process, human IGHV1-2\*01 (SEQ ID NO: 3), and IGKV1-33\*01 (SEQ ID NO: 8) variable domains were selected as acceptors to the 5G6 CDRs. A first humanized antibody of human gamma one isotype was prepared (see below). The antibody encompassed a human-mouse hybrid heavy chain variable domain and a human-mouse hybrid light chain variable domain. The hybrid heavy chain variable domain was based on the human heavy chain variable domain IGHV1-2\*01 wherein germline CDR1 and 2 were respectively replaced for 5G6 heavy chain CDR1 and 2. Best matching JH segment sequence to the human acceptor framework was identified from the IMGT searches mentioned above.

The resulting human-mouse hybrid heavy chain variable sequence having human IGHV1-2\*01 framework regions, 5G6 mouse CDRs, and best matching JH to human acceptor is referred herein as heavy chain variable domain VH1 with SEQ ID NO: 13. Similarly, the human-mouse hybrid light chain variable domain used for this first humanized antibody candidate had human IGKV1-33\*01 framework regions, 5G6 mouse CDRs, and best matching JK to human acceptor, and is referred herein as light chain variable domain VL1 with SEQ ID NO: 14. The first humanized antibody encompassing VH1 and VL1 is abbreviated herein VH1/VL1 antibody.

10 *Production of the first humanized antibody prototype*

Coding DNA sequences (cDNAs) for VH1 and VL1 were synthesized in a scFv format by GENEART AG (Regensburg, Germany) thereby allowing for a single cDNA sequence to encompass both variable domains (SEQ ID NO: 15). Individual variable domain cDNAs were retrieved from this scFv construct by PCR, and further assembled upstream of their respective constant domain cDNA sequence(s) using PCR assembly techniques. Finally, the complete heavy and light chain cDNAs were ligated in independent vectors that are based on a modified pcDNA3.1 vector (Invitrogen, CA, USA) carrying the CMV promoter and a Bovine Growth Hormone poly-adenylation signal. The light chain specific vector allowed expression of human kappa isotype light chains by ligation of the light chain variable domain cDNA of interest in front of the kappa light chain constant domain cDNA using BamHI and BsiWI restriction enzyme sites; while the heavy chain specific vector was engineered to allow ligation of the heavy chain variable domain cDNA of interest in front of the cDNA sequence encoding the human IGHG1 CH1, IGHG1 hinge region, IGHG1 CH2, and IGHG1 CH3 constant domains using BamHI and Sall restriction enzyme sites. In both heavy and light chain expression vectors, secretion was driven by the mouse VJ2C leader peptide containing the BamHI site. The BsiWI restriction enzyme site is located in the kappa constant domain; whereas the Sall restriction enzyme site is found in the IGHG1 CH1 domain.

The VH1/VL1 antibody (having heavy chain SEQ ID NO: 16 and light chain SEQ ID NO: 17) was transiently produced by co-transfecting equal quantities of heavy and light chains vectors into suspension-adapted HEK293-EBNA1 cells (ATCC<sup>®</sup> catalogue number: CRL-10852) using polyethylenimine (PEI, Sigma, Buchs, Switzerland). Typically, 100 ml of cells in suspension at a density of 0.8-1.2 million cells per ml is transfected with a DNA-PEI mixture

containing 50 µg of expression vector encoding the heavy chain and 50 µg of expression vector encoding the light chain. When recombinant expression vectors encoding antibody genes are introduced into the host cells, antibodies are produced by further culturing the cells for a period of 4 to 5 days to allow for secretion into the culture medium (EX-CELL 293, HEK293-serum-free medium; Sigma, Buchs, Switzerland), supplemented with 0.1% pluronic acid, 4 mM glutamine, and 0.25 µg/ml geneticin).

The VH1/VL1 antibody was purified from cell-free supernatant using recombinant protein-A streamline media (GE Healthcare Europe GmbH, Glattbrugg, Switzerland), and buffered exchanged into phosphate buffer saline prior to assays.

*Kinetic binding affinity constants by surface plasmon resonance (SPR)*

Kinetic binding affinity constants (KD) were measured on protein-A captured antibody using recombinant histidine tagged TL1A as analyte. Measurements were conducted on a BIAcore 2000 (GE Healthcare - BIAcore, GE Healthcare Europe GmbH, Glattbrugg, Switzerland) at room temperature, and analyzed with the BiaEvaluation software (BIAcore; v4.1, GE Healthcare Europe GmbH).

A CM5 research grade sensor chip (GE Healthcare Europe GmbH; ref. BR-1000-14) was activated by injecting 35 µl of a 1:1 N-hydroxysulfosuccinimide (NHS)/ 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide Hydrochloride (EDC) solution (v/v; 5 µl/min flow-rate; on flow paths 1 and 2). Protein-A (ref. P7837; Sigma-Aldrich Chemie GmbH, Buchs, Switzerland) was diluted to a final concentration of 50 µg/ml in acetate buffer pH 4.5 (GE Healthcare Europe GmbH, BR-1003-50; one pH unit below pI) and subsequently immobilized on the previously activated CM5 sensor chip by injecting 35 µl on both flow path 1 and 2 (5 µl/min); this corresponded to approximately 1500 response units (RUs). The protein-A-CM5 sensor chip was then deactivated by injecting 35 µl of ethanolamine solution (5µl/min). Finally, two injections of 10 µl of glycine solution (GE Healthcare Europe GmbH, ref. BR-1003-54; 10 mM; pH 1.5) were performed to release non-crosslinked protein-A molecules. For affinity measurements, the chimeric and humanized antibody stored in 1 x PBS buffer were diluted in HBS-EP buffer (GE Healthcare Europe GmbH, ref. BR-1001-88; 0.01 M HEPES, 0.15 M NaCl, EDTA 3 mM, 0.005% Surfactant P20, pH 7.4) and subsequently injected on the flow-path 2 of the protein-A CM5 chip (30 µl/min) to reach about 180 RUs. Following this



capture step, the recombinant histidine tagged human TL1A was injected at different concentrations (1.25 to 125 nM) on the flow-path 1 and 2 (flow-path 1 being used as reference) at a 30  $\mu$ l/min flow rate. After each binding event, surface was regenerated with glycine buffer pH 1.5 injected for 20 sec (30  $\mu$ l/min).

5

Measurements (sensorgram: fc2-fc1) were best fitted with a 1:1 analyte model without mass transfer. To account for the experimental variations in protein-A captured antibody at the beginning of each measurement, the Rmax value was set to local in all fits. Dissociation times were of at least 300-600 seconds. Measurements were performed in duplicate and included zero-concentration samples for referencing. The Chi2 value represents the sum of squared differences between the experimental data and reference data at each point; while the plots of residuals indicate the difference between the experimental and reference data for each point in the fit. Both Chi2 and residual values were used to evaluate the quality of a fit between the experimental data and individual binding models.

15

#### Back mutations of grafted human frameworks

Since straight grafting of CDRs from 5G6 mouse antibody led to a candidate having no binding to human TL1A (Table 6), mutagenesis wherein human residues are substituted for mouse residues was initiated. This process is called back-mutation and is the most unpredictable procedure in the humanization of monoclonal antibodies. It necessitates the identification and the selection of critical framework residues from the mouse antibody that need to be retained in order to preserve affinity while at the same time minimizing potential immunogenicity in the humanized antibody.

20

To identify residues that may impact the most CDR conformation and/or inter-variable domain packing, a 3D model for the VH1-VL1 pair of variable domains was calculated using the structure homology-modelling server SWISS-MODEL (Arnold K *et al.*, (2006) Bioinformatics, 22(2): 195-201; <http://swissmodel.expasy.org>) set in automated mode. Model analysis allowed the selection of a subset of positions based on their putative influence on CDR regions and/or heavy chain-light chain variable domain packing. This subset of positions consisted of variable heavy chain positions: 37, 48, 50, 67, 69, 71 and 75 as well as variable light chain positions: 5 and 34 (Kabat numbering).

25

30

Further humanized candidates having back mutations at the selected positions mentioned above were prepared using gene synthesis and standard mutagenesis methods. A single cDNA sequence encompassing both VH2 and VL2 variable domains (SEQ ID NO: 18) was synthesised and used as a starting point for further mutagenesis. Antibody expression and purification followed the methods described above. Humanized antibody candidates were assayed for their binding affinity by SPR as previously described.

Binding properties (KD) of selected humanized antibodies based on these single or combination of substitutions are shown in Table 6. Amongst humanized variants, VH3/VL1 antibody had the highest affinity for TL1A antigen, exhibiting a lower KD than the 5G6 chimeric antibody.

*Thermostability of selected humanized anti-TL1A antibodies by differential scanning calorimetry*

The thermal stabilities of the humanized antibodies were measured using differential scanning calorimetry (DSC). Monoclonal antibodies melting profiles are characteristic of their isotypes (Garber E & Demarest SJ (2007) Biochem. Biophys. Res. Commun. 355: 751-7), however the mid-point melting temperature of the FAB fragment can be easily identified even in the context of a full-length IgG. Such mid-point melting of FAB portion was used to monitor monoclonal stability of the humanized candidates.

Calorimetric measurements were carried out on a VP-DSC differential scanning microcalorimeter (GE Healthcare Europe GmbH). The cell volume was 0.128 ml, the heating rate was 200°C/h, and the excess pressure was kept at 65 p.s.i. All antibodies were used at a concentration of 1 mg/ml in PBS (pH 7.4). The molar heat capacity of antibody was estimated by comparison with duplicate samples containing identical buffer from which the antibody had been omitted. The partial molar heat capacities and melting curves were analyzed using standard procedures. Thermograms were baseline corrected and concentration normalized before being further analyzed using a Non-Two State model in the software Origin v7.0.

Humanized variant VH5/VL1 FAB fragment displayed a single transition at 83.8°C with a shape and amplitude consistent with a cooperative unfolding which is generally observed for a

compactly folded FAB fragments indicating that the engineering process was successful at retaining FAB stability. Overall the humanized variant showed a good thermal stability.

**Table 6: humanized anti human TL1A antibodies**

Antibody variant (IGHG1)	SEQ ID NOs	Back-mutations VH/VL	K <sub>D</sub> (pM)
Chimera	19, 20	N.A./N.A.	728
VH1/VL1	16, 17	N.A./N.A.	no binding
VH2/VL2	21, 25	<u>VH</u> : V37A-M48I-W50E-V67A-M69L-R71V-I75S <u>VL</u> : N34S-T5N	857
VH3/VL1	22, 17	<u>VH</u> : V37A-M48I-W50E-V67A-R71V	249
VH4/VL1	23, 17	<u>VH</u> : V37A-M48I-W50E-R71V	681
VH5/VL1	24, 17	<u>VH</u> : V37A-W50E-R71V	259

5

**Example 6:**

**5G6 humanized candidates can block TL1A-induced IFN- $\gamma$  secretion by primed CD4 T cells**

Humanized candidates of the 5G6 antibody were tested to determine whether they could inhibit the TL1A-dependent increase in IFN- $\gamma$  production.

10

Human CD4 T cells were purified from peripheral blood mononuclear cells (PMBC) as described in Example 3 above. All cell cultures were performed in Roswell Park Memorial Institute medium (RPMI-1640, PAA Laboratories, Pasching, Austria) supplemented with 10% heat inactivated foetal calf serum (FCS, Amimed distributed by Bioconcept, Allschwil, Switzerland), non essential amino acids (PAA, distributed by Chemie Brunschwig AG, Basel, Switzerland), ultraglutamine (Lonza, Basel, Switzerland), sodium pyruvate (PAA) and penicillin/streptomycin mix (Gibco Life technologies). CD4 purified T cells ( $10^5$  cells/well) were incubated with IL-12 (Peprotech, Hamburg Germany) at 8ng/mL, IL-18 (MBL International, distributed by LabForce AG, Nunningen, Switzerland) at 200ng/mL (priming factors) and human soluble TL1A with an N-terminal his-tag (encoded by SEQ ID NO: 118) for 72hrs in presence of the blocking 5G6 humanized antibody candidates (VH3/VL1– SEQ ID

15

20

NOS: 22 and 17, VH4/VL1– SEQ ID NOS: 23 and 17, VH5/VL1– SEQ ID NOS: 24 and 17 and VH2/VL2– SEQ ID NO: 21 and 25), added at the concentrations 100, 10, 1 and 0.1µg/mL (see tables in Figure 5), at the same time. The isotype control was added at 100µg/mL in a flat bottom 96-well cell culture plate (TPP AG, Trasadingen, Switzerland). The supernatants were harvested after 72 hrs and IFN-γ was quantified by ELISA using OptEIA kit (BD Pharmingen, Allschwil, Switzerland) according to the manufacturer's instructions. Figure 5 shows that humanized anti-TL1A antibodies were able to inhibit substantially the production of IFN-γ.

#### 10 **Example 7:**

##### **5G6 humanized antibody is effective in a murine model of allergic asthma**

Asthma does not spontaneously develop in mice therefore to investigate this disease in mice an asthmatic-like reaction needs to be induced in the airways. A variety of acute allergen challenge models have been developed and in this example BALB/c mice were used as they develop a good T helper cell 2 (Th2)-biased immunological response (Boyce JA & Austin KF (2005) J Exp Med, 201: 1869-73). Ovalbumin, derived from chicken egg is an allergen that induces a robust, allergic pulmonary inflammation in mice and therefore is frequently used in murine models of allergic asthma (Kumar RK *et al.*, (2008) Curr Drug Targets, 9: 485-94).

20 In this example, the following immunization protocol was used to induce allergic asthma: BALB/c mice were sensitized by i.p. injection of 100µg of ovalbumin (Albumin from chicken egg white, Grade V, Sigma Aldrich, Switzerland) adsorbed on 1mg of a suspension of aluminium hydroxide and magnesium hydroxide (Imject Alum, Thermo Scientific, Switzerland) on day 0 and day 14. On day 28, 30 and 33, mice were treated i.p. with 50mg/kg of humanized 5G6 antibody (VH5/VL1; format IgG4 hinge stabilised; SEQ ID NO: 124 and 17) or an equivalent amount of the control human IgG. As a positive control, dexamethasone (Sigma, Switzerland) was used at 5mg/kg. Four hours following treatment, mice were anesthetized with 30mg/kg of xylazine and 150mg/kg of Ketamin (Xylazol and Ketazol from Graeub Veterinary products, Switzerland) and injected intranasally with 10µg of ovalbumin. Three days later, mice were sacrificed and after cannulation of their trachea, bronchoalveolar lavage (BAL) was performed by injecting 2ml of cold PBS into the lung. Cells in BAL fluids were counted and eosinophils detected by flow cytometry using CD11c and Siglec F cell surface markers.

The results are shown in Figure 6 where it can be observed that treatment with the humanized antibody 5G6 resulted in approximately a 4-fold reduction in the number of eosinophils in BAL fluid of asthmatic mice.

5

**Example 8:****5G6 humanized antibody is effective in a treating DSS-induced acute colitis in mice**

Many different animal models of inflammatory bowel disease (IBD) have been developed and these are valuable tools for investigating the involvement of various factors into the pathogenesis of IBD and to evaluate therapeutic options. The dextran sulphate sodium (DSS) induced colitis model is a widely used model of inflammatory bowel disease because of its simplicity and it has many similarities to human IBD, particularly ulcerative colitis (Perše M & Cerar A (2012) J Biomed Biotechnol, 2012; 718617; Wirtz S *et al.*, (2007) Nat. Protoc, 2: 541-6).

15

To evaluate the potential effect of humanized 5G6 antibody (VH5/VL1; format IgG4 hinge stabilised; SEQ ID NO: 124 and 17) in IBD, a condition of acute colitis was induced in C57Bl/6 mice by administering 2% of DSS (MW 36-50kDa; MP Biomedicals) in the drinking water of the test group for 5 days. The control group were given untreated tap water. Following DSS exposure, tap water was given to the test group of mice for 7 days. Mice were treated i.p 3x/week with 50mg/kg of humanized 5G6 antibody or an equal amount of isotype control. As a positive control, cyclosporine (Sandimmune, Novartis Pharma, Switzerland) was used at 5mg/kg. Mice were monitored daily for weight loss and stool consistency. At day 12, all mice were sacrificed and their entire colon lengths measured. As shown in Figure 7, it can be observed that treatment of mice with humanized antibody 5G6 resulted in reduction of colon shortening induced by DSS.

25

**Example 9:****5G6 humanized antibody is effective in a treating TNBS colitis in rats**

Intestinal inflammation in rats can be induced by intrarectal administration of trinitrobenzenesulfonic acid (TNBS). The resulting localised ulceration and inflammation inflammation is believed to involve a T-cell mediated response against hapten-modified

30

autologous proteins or luminal antigens (Wirtz S *et al.*, *supra*). Symptoms include diarrhoea, occult blood and weight loss.

To evaluate the potential effect of a humanized 5G6 antibody (VH5/VL1; format IgG4 hinge stabilised; SEQ ID NO: 124 and 17) in IBD, a condition of colitis was induced in Sprague-Dawley rats by intrarectal administration of TNBS Solution (50% TNBS: 50% 200 proof ethanol; 16 mg/ml TNBS (Sigma, Cat# 92822) at 64 mg/kg (4 ml/kg) into the colon of anaesthetised rats in the treatment groups. The control group received no TNBS Solution. Two hours after TNBS administration, rats were treated i.p with a single dose of humanized 5G6 antibody (50mg/kg) or an equal amount of isotype control. As a positive control, prednisolone (Sigma) was administered orally at a dose of 10mg/kg two hours after TNBS administration and daily for the following 5 days. Rats were sacrificed on day 7 and disease severity assessed as a colonic score using the following scoring system:

- 1) Adhesions: none = 0, minimal = 1, involving several bowel loops = 2
- 2) Strictures: none = 0, mild = 2, severe, proximal dilatation = 3
- 3) Ulcers: none = 0, linear ulceration < 1 cm = 1, two linear ulcers < 1 cm = 2, more sites of ulceration or one large ulcer = 3
- 4) Wall thickness: less than 1 mm = 0, 1-3 mm = 1, > 3 mm = 2

As shown in Figure 8, it can be observed that treatment of rats with humanized antibody 5G6 resulted in reduction of disease parameters induced by TNBS.

#### **Example 10:**

#### **Binding of 5G6 humanized antibody to hTL1A is blocked by both hDcR3-Fc and hDR3-Fc**

As discussed above TL1A is a ligand for TNFRSF25/DR3 and the decoy receptor DcR3. DR3 is a death domain-containing receptor that is upregulated during T cell activation and the interaction of TL1A with DR3 can promote T cell expansion during an immune response (Migone TS *et al.*, *supra*). The secreted decoy receptor (DcR3), a soluble protein of the tumor necrosis factor receptor (TNFR) superfamily, blocks the action of TL1A. (Kim S & Zhang L *supra*).

To evaluate whether the 5G6 humanized antibody could interfere with the interaction of hTL1A to the receptor DR3 and/or to the decoy receptor DcR3, a his-tagged human TL1A

was coated at 2  $\mu\text{g/ml}$  on an ELISA plate and incubated with 20  $\mu\text{g/ml}$  of 5G6 humanized antibody (VH5/VL1; format IgG4 hinge stabilised; SEQ ID NO: 124 and 17) in the presence of 10  $\mu\text{g/ml}$  of Fc fusions of the ectodomains of either human DcR3, DR3 (both R&D systems) or an irrelevant receptor (Ctrl-Fc) followed by detection with peroxidase-  
5 conjugated anti-human IgG (Fab specific). As can be observed in Figure 9, binding of the 5G6 humanized antibody to hTL1A is blocked by both hDcR3-Fc and hDR3-Fc. This confirms that 5G6 humanized antibody binding to hTL1A inhibits the interaction of hTL1A with both DR3 and DcR3.

**Claims**

1. An antibody or fragment thereof that binds to TL1A comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 51, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 52, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and/or comprises a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 54, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 55, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 56.  
5
2. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof is a murine antibody, chimeric antibody or a humanized antibody.  
10
3. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof is a humanized antibody.
4. The antibody or fragment thereof of claims 1 or 2, wherein the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 1  
15
5. The antibody or fragment thereof of claims 1 or 2, wherein the antibody or fragment thereof comprises a non-CDR region of a heavy chain variable region sequence which is at least 80 % identical to the non-CDR region of the heavy chain variable region sequence of SEQ ID NO: 1.  
20
6. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence selected from the group consisting of SEQ ID NOS: 26, 27, 28 and 29.
7. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a non-CDR region of a heavy chain variable region sequence which is at least 80% identical to the non-CDR region of the heavy chain variable region sequence selected from the group consisting of SEQ ID NOS: 26, 27, 28 and 29.  
25



8. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 29.
- 5 9. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a non-CDR region of a heavy chain variable region sequence which is at least 80 % identical to the non-CDR region of the heavy chain variable region sequence of SEQ ID NO: 29.
- 10 10. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a heavy chain sequence comprising the amino acid sequence selected from the group consisting of SEQ ID NOS: 21, 22, 23 and 24.
11. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a heavy chain sequence comprising a non-CDR region which is at least 80% identical to the non-CDR region of the heavy chain variable region sequence of the heavy chain sequence selected from the group consisting of SEQ ID NOS: 21, 22, 23 and 24.
- 15 12. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a heavy chain variable framework region that is the product of or derived from a human gene selected from the group consisting of IGHV1-2\*02 (SEQ ID NO: 3), IGHV1-2\*04 (SEQ ID NO: 4), IGHV1-2\*05 (SEQ ID NO: 5), IGHV1-2\*01 (SEQ ID NO: 6), and IGHV1-46\*01 (SEQ ID NO: 7).
- 20 13. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof comprises a heavy chain variable framework region that is the product of or derived from human gene IGHV1-2\*01 (SEQ ID NO: 3) and wherein the heavy chain variable framework region comprises at least one amino acid modification from the corresponding heavy chain variable framework region of a corresponding murine antibody.
- 25 14. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof comprises a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 16 and wherein the heavy chain variable framework region comprises at least one amino acid

modification from the corresponding heavy chain variable framework region of a corresponding murine antibody.

15. The antibody or fragment thereof of claim 13 or 14, wherein the amino acid modification comprises an amino acid substitution at amino acid position selected from the group consisting of 37, 48, 50, 67, 69, 71 and 75, wherein the amino acid position of each group member is indicated according to the Kabat numbering.

16. The antibody or fragment thereof of claim 13 or 14, wherein the amino acid modification comprises an amino acid substitution selected from the group consisting of 37A, 48I, 50E, 67A, 69L, 71V and 75S, wherein the amino acid position of each group member is indicated according to the Kabat numbering.

17. The antibody or fragment thereof of claims 1 or 2, wherein the antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 2.

18. The antibody or fragment thereof of claims 1 or 2, wherein the antibody or fragment thereof comprises a non-CDR region of a light chain variable region sequence which is at least 80 % identical to the non-CDR region of the light chain variable region sequence of SEQ ID NO: 8.

19. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence selected from the group consisting of SEQ ID NOS: 14 and 30.

20. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a non-CDR region of a light chain variable region sequence which is at least 80 % identical to the non-CDR region of the light chain variable region sequence selected from the group consisting of SEQ ID NO: 14 and 30.

21. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a light chain sequence comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 17 and 25.
22. The antibody or fragment thereof of claim 1, wherein the light chain sequence comprises  
5 a non-CDR region which is at least 80% identical to the non-CDR region of the light chain variable region sequence of the light chain sequence selected from the group consisting of SEQ ID NO: 17 and 25.
23. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises a light chain variable framework region that is the product of or derived from a  
10 human gene selected from the group consisting of IGKV1-33\*01 (SEQ ID NO: 8), IGKV1D-33\*01 (SEQ ID NO: 9), IGKV1D-12\*02 (SEQ ID NO: 10), IGKV1D-12\*01 (SEQ ID NO: 11) and IGKV1-12\*02 (SEQ ID NO: 12).
24. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof comprises a light chain variable framework region that is the product of or derived from human  
15 gene IGKV1-33\*01 (SEQ ID NO: 8) and wherein the light chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of a corresponding murine antibody.
25. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof comprises a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17 and  
20 wherein the light chain variable framework region comprises at least one amino acid modification from the corresponding light chain variable framework region of the corresponding murine antibody.
26. The antibody or fragment thereof of claim 24 or 25, wherein the amino acid modification  
25 comprises an amino acid substitution at amino acid position selected from the group consisting of 5 and 34, wherein the amino acid position of each group member is indicated according to the Kabat numbering.

27. The antibody or fragment thereof of claim 24 or 25, wherein the amino acid modification comprises an amino acid substitution selected from the group consisting of 5N and 34S, wherein the amino acid position of each group member is indicated according to the Kabat numbering.
- 5 28. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises:  
(a) a heavy chain sequence comprising the amino acid sequence of SEQ ID NO: 22 or SEQ ID NO: 24; and  
(b) a light chain sequence comprising the amino acid sequence of SEQ ID NO: 17.
- 10 29. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof comprises:  
(a) a heavy chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 27 or SEQ ID NO: 29; and  
(b) a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO:  
15 14.
30. The antibody or fragment thereof of any one of claims 1 to 29, wherein at least one of the heavy chain CDRs and/or at least one of the light chain CDRs comprises at least one amino acid modification.
- 20 31. The antibody or fragment thereof of any one of claims 1 to 30, further comprising heavy and/or light constant regions.
32. The antibody or fragment thereof of claim 31, wherein the human heavy constant region is selected from the group of human immunoglobulins consisting of IGHG1, non fucosylated IGHG1 and IGHG4.
- 25 33. The antibody or fragment thereof of any one of claims 1 to 31, wherein the antibody has a non fucosylated IGHG1 Fc region.

34. The antibody or fragment thereof of any one of claims 1 to 31, wherein the antibody comprises an isotypic variant comprising the CH1 from human IgG4 (IGHG4), the hinge from human IgG4 (IGHG4), having S228P substitution and the CH2 and CH3 from human IgG4 (IGHG4).

5

35. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody or fragment thereof binds to human TL1A.

36. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody or fragment thereof binds to human TL1A and is cross reactive with murine, rat and cynomologus TL1A.

10

37. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody or fragment thereof that bind to hTL1A and inhibits the interaction of hTL1A with both DR3 and DcR3.

15

38. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody or fragment thereof is an antagonist antibody.

39. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody or fragment thereof is a neutralizing antibody.

20

40. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody is a monovalent antibody.

41. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody is a full length antibody.

25

42. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody is an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH, Fd, Fv, dAb, F(ab')<sub>2</sub>, scFv, bispecific single chain Fv dimers, diabodies, triabodies and scFv genetically fused to the same or a different antibody.

30

43. The antibody or fragment thereof of any one of claims 1 to 34, wherein the antibody comprises a variant Fc region which comprises at least one amino acid modification relative to the Fc region of the parent antibody, whereas the antibody comprising the variant Fc region exhibits altered effector function compared to the parent antibody.

5

44. The antibody or fragment thereof of any one of claims 1 to 43, wherein the antibody or fragment thereof binds to human TL1A with an affinity ( $K_D$ ) of 700 pM or less.

45. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof retains at least 85% of the TL1A binding affinity ( $K_D$ ) of the corresponding chimeric antibody.

10

46. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof has equivalent or higher TL1A binding affinity ( $K_D$ ) when compared to the corresponding chimeric antibody.

47. The antibody or fragment thereof of claim 3, wherein the antibody or fragment thereof binds human TL1A with three-fold higher affinity than the corresponding chimeric antibody

15

48. The antibody or fragment thereof of any one of claims 1 to 43, wherein the antibody has a FAB fragment thermostability temperature greater than 80°C.

49. An antibody or fragment thereof that binds to human TL1A and which binds to the same epitope as the antibody of any one of claims 1 to 43.

20

50. An epitope on soluble human TL1A which is bound by the antibody of any one of claims 1 to 43.

25

51. An isolated nucleic acid encoding the antibody or fragment thereof of any one of claims 1 to 49.

52. The isolated nucleic acid of claim 51 comprising DNA encoding the heavy chain variable region comprising the nucleotide acid sequence of SEQ ID NO: 33 or 35; and/or DNA

encoding the light chain variable region comprising the nucleotide acid sequence of SEQ ID NO: 36.

53. A vector comprising the isolated nucleic acid of claim 51 or 52.

5

54. A host cell comprising the isolated nucleic acid of claim 51 or 52 or the vector of claim 53.

10

55. A method of producing an antibody or fragment thereof that binds to human TL1A comprising culturing the host cell of claim 54 so that the nucleic acid is expressed and the antibody produced.

15

56. An antibody or fragment thereof that binds to human TL1A encoded by the isolated nucleic acid of claim 51 or 52.

57. A composition comprising the antibody or fragment thereof of any one of claims 1 to 49 and a pharmaceutically acceptable carrier.

20

58. An immunoconjugate comprising the antibody or fragment thereof of any one of claims 1 to 49 linked to a therapeutic agent.

59. A composition comprising the immunoconjugate of claim 58 and a pharmaceutically acceptable carrier.

25

60. The composition of claim 57 or 59 further comprising another pharmaceutically active agent.

30

61. A method for treating a TL1A mediated disorder in a subject, the method comprising administering to the subject a therapeutically effective amount of the antibody or fragment thereof of any one of claims 1 to 49.

62. The method of claim 61, wherein the TL1A mediated disorder is selected from the group consisting of inflammatory bowel disease (IBD) including ulcerative colitis and Crohn's

disease, rheumatoid arthritis, MS, type 1 and type 2 diabetes, psoriasis, psoriatic arthritis, ankylosing spondylitis, atopic dermatitis; allergic reactions or conditions, including for example, asthma and allergic lung inflammation; cancers, atherosclerosis, infections, neurodegenerative diseases, graft rejection, graft versus host diseases (GVHD) and cardiovascular disorders/diseases.

63. The method of claim 61 or 62, wherein the antibody has a non fucosylated IGHG1 Fc region and exhibits enhanced cytotoxicity as compared to the antibody having human heavy chain constant region IGHG1.

64. Use of an antibody or fragment thereof of any one of claims 1 to 49 as a medicament.

65. Use of an antibody or fragment thereof of any one of claims 1 to 49 in the preparation of a medicament for the treatment of a TL1A mediated disorder.

66. The antibody or fragment thereof of any one of claims 1 to 49 for use as a medicament.


67. The antibody or fragment thereof of any one of claims 1 to 49 for use in a method for treating a TL1A mediated disorder.

68. An article of manufacture comprising the antibody or fragment thereof of any one of claims 1 to 49, the composition of claim 57 or 59 or the immunoconjugate of claim 58 for the treatment of a TL1A mediated disorder.

69. A kit comprising the antibody or fragment thereof of any one of claims 1 to 49, the composition of claim 57 or 59 or the immunoconjugate of claim 58 for the treatment of a TL1A mediated disorder.



FIG. 1






FIG. 2

FIG. 3





FIG. 3 cont.

**C**



IL-12 + IL18	-	+	+	+	+	+	+
IC monocytes	-	-	+	+	+	+	+
5G6 (µg/ml)	-	-	-	10	1	0.1	-
Isotype control	-	-	-	-	-	-	+

FIG. 4




FIG. 5







FIG. 6




FIG. 7






FIG. 8




FIG. 9



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2013/077898

## Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
  - a. (means)
    - on paper
    - in electronic form
  - b. (time)
    - in the international application as filed
    - together with the international application in electronic form
    - subsequently to this Authority for the purpose of search
2.  In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

# INTERNATIONAL SEARCH REPORT

International application No <b>PCT/EP2013/077898</b>
--

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> INV. A61P37/00      A61K39/395      C07K16/28 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b>				
Minimum documentation searched (classification system followed by classification symbols) C07K A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, BIOSIS, EMBASE				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	WO 2012/064682 A1 (REGENERON PHARMA [US]; CLASSON BRENDAN J [US]; SKOKOS DIMITRIS [US]) 18 May 2012 (2012-05-18) examples 1-7 -----	1-69		
X	WO 2012/161856 A1 (US GOVERNMENT [US]; SIEGEL RICHARD M [US]; MEYLAN FRANCOISE [US]; SONG) 29 November 2012 (2012-11-29) paragraphs [0031], [0362]; figures 22,26-28 -----	1-69		
X	WO 2009/064854 A2 (COGENESYS INC [US]; SMITH RODGER [US]; KANAKARAJ PALANISAMY [US]; ROSC) 22 May 2009 (2009-05-22) examples 1-6 ----- -/--	1-69		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.				
* Special categories of cited documents : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;">                     "A" document defining the general state of the art which is not considered to be of particular relevance                      "E" earlier application or patent but published on or after the international filing date                      "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)                      "O" document referring to an oral disclosure, use, exhibition or other means                      "P" document published prior to the international filing date but later than the priority date claimed                 </td> <td style="width: 50%; border: none; vertical-align: top;">                     "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention                      "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone                      "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art                      "&amp;" document member of the same patent family                 </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
12 February 2014	24/02/2014			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Cilensek, Zoran			

INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2013/077898

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>HIDETOSHI TAKEDATSU ET AL: "TL1A (TNFSF15) regulates the development of chronic colitis by modulating both T-helper 1 and T-helper 17 activation", GASTROENTEROLOGY, ELSEVIER, PHILADELPHIA, PA, vol. 135, no. 2, 1 August 2008 (2008-08-01), pages 552-567, XP002668871, ISSN: 0016-5085, DOI: 10.1053/J.GASTRO.2008.04.037 [retrieved on 2008-05-07] page 555, left-hand column, lines 16-19; figures 4-6</p>	1-69
X	<p>-----                      WO 2008/106451 A2 (CEDARS SINAI MEDICAL CENTER [US]; ROTTER JEROME I [US]; TAYLOR KENT D) 4 September 2008 (2008-09-04) example 25</p>	1-69
X	<p>-----                      WO 2007/027751 A2 (UNIV MIAMI [US]; PODACK ECKHARD [US]; DEYEV VADIM [US]; LEVY ROBERT [U] 8 March 2007 (2007-03-08) examples 15,18</p> <p>-----</p>	1-69

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2013/077898
---

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012064682 A1	18-05-2012	AR 083785 A1	20-03-2013
		AU 2011326097 A1	09-05-2013
		CA 2816799 A1	18-05-2012
		CN 103261228 A	21-08-2013
		EA 201370105 A1	30-12-2013
		EP 2638069 A1	18-09-2013
		KR 20130132842 A	05-12-2013
		SG 190176 A1	28-06-2013
		TW 201233796 A	16-08-2012
		US 2012114654 A1	10-05-2012
		UY 33715 A	31-05-2012
		WO 2012064682 A1	18-05-2012
		-----	
WO 2012161856 A1	29-11-2012	AU 2012259312 A1	12-12-2013
		CA 2836898 A1	29-11-2012
		US 2012263718 A1	18-10-2012
		WO 2012161856 A1	29-11-2012
-----			
WO 2009064854 A2	22-05-2009	AU 2008320987 A1	22-05-2009
		CA 2705292 A1	22-05-2009
		CN 101903402 A	01-12-2010
		EA 201070596 A1	30-12-2010
		EP 2220122 A2	25-08-2010
		EP 2500362 A2	19-09-2012
		JP 2011502522 A	27-01-2011
		KR 20100097683 A	03-09-2010
		NZ 585559 A	30-11-2012
		US 2009280116 A1	12-11-2009
		US 2012308480 A1	06-12-2012
		WO 2009064854 A2	22-05-2009
		-----	
WO 2008106451 A2	04-09-2008	US 2010190162 A1	29-07-2010
		WO 2008106451 A2	04-09-2008
-----			
WO 2007027751 A2	08-03-2007	AU 2006284922 A1	08-03-2007
		BR PI0617057 A2	12-07-2011
		CA 2621083 A1	08-03-2007
		CN 101253199 A	27-08-2008
		EP 1919954 A2	14-05-2008
		JP 5164167 B2	13-03-2013
		JP 2009505678 A	12-02-2009
		US 2007128184 A1	07-06-2007
		US 2011243951 A1	06-10-2011
		US 2012321645 A1	20-12-2012
		WO 2007027751 A2	08-03-2007
		-----	