



(51) International Patent Classification:

A61K 31/4545 (2006.01) A61P 25/06 (2006.01)

A61K 9/00 (2006.01)

(21) International Application Number:

PCT/US2021/043791

(22) International Filing Date:

29 July 2021 (29.07.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/103,353	29 July 2020 (29.07.2020)	US
63/070,449	26 August 2020 (26.08.2020)	US
63/087,175	02 October 2020 (02.10.2020)	US
63/092,211	15 October 2020 (15.10.2020)	US
63/129,362	22 December 2020 (22.12.2020)	US
63/201,254	20 April 2021 (20.04.2021)	US

(71) Applicant: ALLERGAN PHARMACEUTICALS INTERNATIONAL LIMITED [IE/IE]; Clonsaugh Business & Technology Park, Dublin, D17 E400 (IE).

(72) Inventors; and

(71) Applicants: TRUGMAN, Joel [US/US]; 1025 Maxwell Lane, Unit 703, Hoboken, New Jersey 07030 (US). FIN-NEGAN, Michelle [US/US]; 358 78th Street, Brooklyn, New York 11209 (US).

(74) Agent: JOHANNES, Laura et al.; c/o AbbVie Inc., AP34-2 V0377, 1 North Waukegan Road, North Chicago, Illinois 60064 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

(54) Title: TREATMENT OF MIGRAINE




FIGURE 1

(57) Abstract: The present disclosure provides methods for the treatment of migraine by the administration of atogepant or a pharmaceutically acceptable salt thereof.

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

TREATMENT OF MIGRAINE

FIELD

[0001] The present disclosure is related to medicaments and methods for treating migraine.

CROSS-REFERENCE TO RELATED APPLICATIONS

5 [0002] The present application claims the benefit of Provisional Application No. 63/103,353, filed July 29, 2020; Provisional Application No. 63/070,449, filed August 26, 2020; Provisional Application 63/087,175, filed October 2, 2020; Provisional Application No. 63/092,211, filed October 15, 2020; Provisional Application No. 63/129,362, filed December 22, 2020; and Provisional Application No. 63/201,254, filed April 20, 2021, the disclosures of which are
10 incorporated herein by reference.

BACKGROUND

[0003] Migraine is a highly prevalent, severe, and disabling neurological condition with a significant unmet need for effective treatments. (Holland, P. R. & Goadsby, P. J. Neurotherapeutics (2018). Migraine affects over 1 billion people worldwide, and it was reported
15 as the second leading cause of disability in the 2016 Global Burden of Disease study. *See* GBD 2019 Diseases and Injuries Collaborators. Global Burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systemic analysis for the Global Burden of Disease Study 2019, *Lancet* 2020;396:1204-22.

[0004] When attacks are frequent or disabling, prevention becomes a focus of migraine
20 treatment. Current preventive treatments for migraine include oral medications, such as valproic acid, flunarizine, topiramate, and propranolol, as well as injectable treatments, such as monoclonal antibodies targeting calcitonin gene-related peptide (CGRP).

[0005] Currently available CGRP-targeted preventive treatments are limited to monoclonal antibodies, administered by injection. Oral CGRP treatments are approved for acute but not
25 preventive treatment, and yet patients and physicians overwhelmingly prescribe oral migraine preventives, suggesting a treatment gap. There remains a need for optimized and targeted methodologies and dosing regimens to use oral CGRP treatments to prophylactically treat migraines.

SUMMARY

[0006] In embodiments, the present disclosure provides methods of prophylactically treating migraine, involving administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, to a patient in need of treatment.

5 [0007] In embodiments, the present disclosure provides methods of prophylactically treating migraine, involving administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, wherein treatment with atogepant results in a reduction in mean monthly migraine days of at least 3.6 days, or at least 3.8 days, or at least 4.2 days, relative to no treatment.

10 [0008] In embodiments, the present disclosure provides a method of prophylactically treating migraine, involving administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, wherein the prophylactic treatment results in a reduction in mean monthly headache days of at least 3.9 days, or at least 4 days, or at least 4.2 days relative to no treatment.

15 [0009] In embodiments, the present disclosure provides a method of prophylactically treating migraine, involving administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, wherein the prophylactic treatment results in a reduction in mean monthly acute medication use days of at least 3.6 days, or at least 3.8 days relative to no treatment.

20 [0004] In embodiments, the present disclosure provides methods of prophylactically treating migraine, involving administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, wherein the prophylactic treatment results in at least 50% reduction in monthly migraine days relative to no treatment.

BRIEF DESCRIPTION OF THE FIGURES

25 [0010] Figures 1 and 2 show the least square mean (+/- standard error of the least squares (SE)) of change from baseline in monthly migraine days (MMRM – mixed effects model for repeated measures) during the double-blind treatment period for the mITT population in Study A. There was a significant reduction in mean monthly migraine days across all three atogepant doses.

[0011] Figure 3 shows the distribution of change from baseline in monthly migraine days across 30 12 weeks by treatment group. A treatment benefit over placebo for all doses of atogepant is seen across a range of changes from baseline in mean monthly migraine days.

[0012] Figure 4 shows the LS mean change from baseline in monthly migraine days across 12 weeks by treatment group.

[0013] Figure 5A shows the mean change from baseline in weekly migraine days during the first month (weeks 1-4) by treatment group. Figure 5B shows the proportion of participants with a migraine each day during the first week of treatment (mITT population). Atogepant provided an early and sustained reduction in migraine days including statistically significant reductions in each of the three 4-week intervals, each week during the first 4-week interval, and as early as the first full day after study drug initiation.

[0014] Figure 6 shows the response profile for percent reduction from baseline in 3-month average of monthly migraine days for the mITT population.

[0015] Figure 7 shows the percent of participants with a $\geq 50\%$ reduction in 3-month average monthly migraine days (MMDs) for placebo, atogepant 10 mg, atogepant 30 mg, and atogepant 60 mg. With respect to the proportion of patients who achieved a $\geq 50\%$ reduction in their 3-month average of monthly migraine days, all three atogepant doses had a significantly greater 50% responder rate versus placebo, with nearly double the proportion of responders achieving at least a 50% reduction in monthly migraine days across all doses when compared to placebo.

[0016] Figure 8A shows the proportion of participants achieving $\geq 25\%$, $\geq 50\%$, $\geq 75\%$, and 100% responder rates in the 12-week average of monthly migraine days. Figure 8B shows a cumulative distribution function graph of the percent reduction from baseline in 12-week average MMDs.

Figure 8C shows the proportion of participants with $\geq 25\%$ reduction in mean monthly migraine days, and Figure 8D shows the proportion of participants with $\geq 50\%$, $\geq 75\%$, and 100% reduction in monthly migraine days for month 1 (weeks 1-4), month 2 (weeks 5-8), and month 3 (weeks 9-12) across the 12-week treatment period.

[0017] Figure 9A shows the mean change from baseline in moderate/severe headache days per month (mITT population). Figure 9B shows the mean change from baseline in monthly headache days per month.

[0018] Figure 10A shows the LS mean change (SE) from baseline in mean monthly acute medication use days across the 12-week treatment period for atogepant 10 mg, atogepant 30 mg, atogepant 60 mg, and placebo. Figure 10B shows the LS mean change from baseline in acute medication use days during the first treatment period (weeks 1-4), second treatment period (weeks 5-8) and third treatment period (weeks 9-12) for atogepant 10 mg, atogepant 30 mg,

atogepant 60 mg, and placebo. Figure 10C shows the LS mean reduction from baseline in mean cumulative headache hours during the first treatment period (weeks 1-4), second treatment period (weeks 5-8) and third treatment period (weeks 9-12) for atogepant 10 mg, atogepant 30 mg, atogepant 60 mg, and placebo.

5 [0019] Figure 11 shows the migraine-specific quality of life questionnaire role function-restrictive domain score for weeks 4, 8, and 12 for all three atogepant doses and placebo. Significant differences vs. placebo were observed at the earliest time point assessed (week 4) and seen throughout the treatment period. All atogepant groups achieved within-group minimally important difference among the MSQ v 2.1 domains at weeks 4, 8 and 12.

10 [0020] Figure 12 provides a schematic for the design of the study evaluating the safety and tolerability of long-term atogepant treatment (Study B).

[0021] Figure 13 shows the LS mean change from baseline in number of migraine days over time (mITT population) in the 52-week long term safety study of atogepant. Administration of 60 mg atogepant led to a rapid reduction in monthly migraine days in month 1, with additional
15 gradual improvement over time through week 52.

[0022] Figure 14 shows the LS mean change from baseline in monthly acute medication use days at each monthly period (mITT population) in the 52-week long-term safety study of atogepant. Administration of 60 mg atogepant led to a rapid reduction in monthly acute medication use days in month 1, with additional gradual improvement over time through week
20 52.

[0023] Figure 15A shows the results for $\geq 50\%$ improvement (reduction) in monthly migraine days at each monthly period (mITT population).

[0024] Figure 15B shows the proportion of responders with $\geq 50\%$, $\geq 75\%$, and $\geq 100\%$ reduction in monthly migraine days (MMDs) (mITT population, observed cases) in the 52-week,
25 multicenter, open-label trial (Study B) of atogepant in adult patients with episodic migraine with or without aura.

[0025] Figure 16A shows the change from baseline in AIM-D monthly PDA domain score: reduced impairment in performance of daily activities (mITT population). Figure 16B shows the change from baseline in AIM-D monthly PI domain score: reduced physical impairment (mITT
30 population) over the 52-week, multicenter, open-label, 52-week, long-term safety trial (Study B) of atogepant in adult patients with episodic migraine with or without aura.

[0026] Figure 17 shows the LS mean change from baseline in MSQ v 2.1 – RFR Domain Score (mITT population, MMRM analysis) over the 52-week, multicenter, open-label trial (Study B) of atogepant in adult patients with episodic migraine with or without aura. Findings indicated rapid improvement over the first 12 weeks, and sustained improvement from 12 to 52 weeks. Non-zero confidence intervals throughout indicate significant improvement at each time point assessed.

[0027] Figure 18 shows the change from baseline in HIT-6 Total Score (mITT population).


[0028] Figure 19 shows CDF plots for percentage change from baseline in body weight (kg) at the end of the double-blind treatment period for Study A (safety population). Statistically significant weight loss was observed in the atogepant 30 mg once daily and 60 mg once daily groups compared with placebo.

[0029] Figure 20 shows CDF plots for percentage change from baseline in body weight (kg) at the end of the open-label treatment period for Study B (safety population). As shown in Figure 20, treatment with atogepant over the 52-week treatment period was associated with weight loss.

[0030] Figures 21-24 show CDF plots of weight loss for the safety population in the Study B long-term safety study. In particular, Figure 21 is a CDF plot of weight loss for the safety population in the long-term safety study at 3 months; Figure 22 is a CDF plot of weight loss for the safety population in the long-term safety study at 6 months; Figure 23 is a CDF plot of weight loss for the safety population in the long-term safety study at 9 months; and Figure 24 is a CDF plot of weight loss for the safety population in the long-term safety study at 12 months.

DETAILED DESCRIPTION

[0031] The present disclosure provides methods for treating migraine in a patient in need thereof. In embodiments, the present disclosure provides methods for the prophylactic treatment of patients suffering from migraine. In embodiments, the present disclosure provides methods for the treatment of migraine comprising administering a prophylactically effective amount of atogepant or a pharmaceutically acceptable salt thereof. The structure of atogepant is shown below:



[0032] Atogepant is a small molecule CGRP receptor antagonist which may be administered orally, reaching maximum plasma concentrations by 2 hours, with a half-life of approximately 11 hours.

- 5 [0033] Prophylactic treatment can reduce the frequency and intensity of migraine attacks. In embodiments, the prophylactic methods can result in freedom from symptoms associated with migraine attacks, including headaches. In embodiments, the administration of atogepant may provide for fewer symptoms or symptoms of reduced intensity. In embodiments, the non-headache symptoms of migraine may be reduced or eliminated. In embodiments, the
- 10 prophylactic methods of the present disclosure are directed to the entire range of symptoms experienced by a patient during a migraine attack, and not solely at the prevention of headaches associated with a migraine attack.

[0034] The indications for preventative treatment of migraine have been published by the American Academy of Neurology. *See Update: Pharmacologic Treatment for Episodic Migraine Prevention in Adults*, American Academy of Neurology, 2012. Prophylactic treatment is

15 generally proposed for patients who suffer from two or more migraine attacks per month. Prophylactic treatment can also be used for patients who experience less frequent migraine attacks that are more potent or even disabling.

[0035] In embodiments, a patient in need of treatment may suffer from one or more symptoms of

20 migraine including, for example, sinusitis, nausea, nasopharyngitis, photophobia, appetite changes, cognition and concentration difficulties, cold extremities, diarrhea or other bowel changes, excitement or irritability, fatigue, frequent urination, memory changes, weakness, yawning, stretching, seeing bright spots or flashes of light, vision loss, seeing dark spots, tingling

sensations, speech problems, aphasia, tinnitus, gastric stasis, pulsating or throbbing pain on one or both sides of the head, extreme sensitivity to light, sounds, or smells, worsening pain during physical activity, and vomiting, abdominal pain or heartburn, loss of appetite, lightheadedness, blurred vision, and fainting. In embodiments, the administration of a prophylactically effective amount of atogepant results in the improvement, reduced frequency, or reduced intensity of symptoms.

[0036] In embodiments, the present disclosure provides a method for the prophylactic treatment of migraine comprising regularly administering to a patient in need thereof a therapeutically effective amount of atogepant, or a pharmaceutically acceptable salt thereof. In embodiments, atogepant is administered orally at a once-daily dose of 10 mg. In embodiments, atogepant is administered orally at a once-daily dose of 30 mg. In embodiments, atogepant is administered orally at a once-daily dose of 60 mg. In embodiments, atogepant is administered orally at a once-daily dose of 10 mg for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks.

[0037] In embodiments, atogepant is administered orally at a once-daily dose of 30 mg. In embodiments, atogepant is administered orally at a once-daily dose of 30 mg for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks.

[0038] In embodiments, atogepant is administered orally at a once-daily dose of 60 mg. In embodiments, atogepant is administered orally at a once-daily dose of 60 mg for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks.

[0039] In embodiments, atogepant, or a pharmaceutically acceptable salt thereof, is administered to a patient identified as susceptible to treatment with atogepant or a pharmaceutically acceptable salt thereof. A patient that suffers from episodic migraine may be considered susceptible to treatment with atogepant (for example 10 mg QD, 30 mg QD, or 60 mg QD) if after a treatment period of one month, two months, or three months, the patient achieves at least 70% reduction in

migraine or probable migraine days, or at least 75% reduction, at least 80% reduction, at least 85% reduction, at least 90% reduction, at least 95% reduction or 100% reduction in migraine or probable migraine days.

[0040] In embodiments, atogepant or a pharmaceutically acceptable salt thereof, is administered to a patient suffering from episodic migraine for at least one month, or at least two months, or at least three months, and results in at least three fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 mean monthly migraine days, or at least 4 fewer mean monthly migraine days, or at least 4.2 fewer monthly migraine days relative to no treatment.

[0041] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 mean monthly migraine days, or at least 4 fewer mean monthly migraine days, or at least 4.2 fewer monthly migraine days.

[0042] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg QD, 30 mg QD, or 60 mg QD to a population of human patients, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 mean monthly migraine days, or at least 4 fewer mean monthly migraine days, or at least 4.2 fewer monthly migraine days. In embodiments, at least about 70% of the patients in the population of human patients have taken at least one prior preventive migraine therapy. In embodiments, the atogepant is administered for at least one month, or at least two months, or at least three months.

[0043] For example, in embodiments, the present disclosure provides a method for prophylactically treating migraine, comprising administering atogepant in an amount of 10 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.65 fewer mean monthly migraine days, or at least 3.69 fewer mean monthly migraine days.

[0044] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 10 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, administering atogepant results in at least 3.4 fewer monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.65 fewer mean monthly migraine days, or at least 3.69 fewer mean monthly migraine days.

[0045] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 10 mg QD to a population of human patients, resulting in a reduction in mean monthly migraine days. In embodiments, at least about 70% of the patients in the population have taken at least one prior preventive migraine therapy. In embodiments, administering atogepant results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.65 fewer mean monthly migraine days, or at least 3.69 fewer mean monthly migraine days.

[0046] In embodiments, the present disclosure provides a method for prophylactically treating migraine, comprising administering atogepant in an amount of 30 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.86 fewer mean monthly migraine days.

[0047] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 30 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.86 fewer mean monthly migraine days.

[0048] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 30 mg QD to a population of human patients, resulting in a reduction in mean monthly migraine days. In embodiments, at least about 70% of the patients in the population have taken at least one prior migraine therapy. In embodiments, treatment results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.86 fewer mean monthly migraine days.

[0049] In embodiments, the present disclosure provides a method for prophylactically treating migraine, comprising administering atogepant in an amount of 60 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, treatment results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.9 fewer mean monthly migraine days, or at least 4.0 fewer mean monthly migraine days, or at least 4.1 fewer mean monthly migraine days, or at least 4.2 fewer mean monthly migraine days.

[0050] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month),

wherein the method comprises administering atogepant in an amount of 60 mg QD, resulting in a reduction in mean monthly migraine days. In embodiments, administering atogepant results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.9 fewer mean monthly migraine days, or at least 4.0 fewer mean monthly migraine days, or at least 4.1 fewer mean monthly migraine days, or at least 4.2 fewer mean monthly migraine days.

[0051] Treatment with atogepant can result in an early and sustained reduction in monthly migraine days. For example, treatment with atogepant can result in a treatment effect as early as the first full day after starting treatment with atogepant. In embodiments, the present disclosure provides a method of preventing migraine, such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering atogepant in an amount of 10 mg or 30 mg or 60 mg, wherein patients treated with atogepant are less likely to have a migraine on the day following atogepant administration than patients who received placebo.

[0052] In embodiments, the present disclosure provides a method for preventing migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), wherein the method comprises administering atogepant in an amount of 60 mg QD to a population of human patients, resulting in a reduction in mean monthly migraine days. In embodiments, at least about 70% of the patients in the population have taken at least one prior migraine therapy. In embodiments, administering atogepant results in at least 3.4 fewer mean monthly migraine days, or at least 3.5 fewer mean monthly migraine days, or at least 3.6 fewer mean monthly migraine days, or at least 3.7 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.8 fewer mean monthly migraine days, or at least 3.9 fewer mean monthly migraine days, or at least 4.0 fewer mean monthly migraine days, or at least 4.1 fewer mean monthly migraine days, or at least 4.2 fewer mean monthly migraine days.

[0053] In embodiments, the present disclosure provides a method for the prophylactic treatment of migraine comprising administering 10 mg atogepant or 30 mg atogepant or 60 mg atogepant

to a patient in need thereof, wherein the patient achieves at least 50% reduction in 3-month average of monthly migraine days.

[0054] In embodiments, the present disclosure provides a method for preventing migraine, such as a method of preventing migraine in a patient having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant or 30 mg atogepant or 60 mg atogepant, wherein the patient achieves at least 50% reduction in 3-month average of monthly migraine days.

[0055] In embodiments, the present disclosure provides a method for the prophylactic treatment / prevention of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant or 30 mg atogepant or 60 mg atogepant to a population of patients, wherein at least about 50% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 55% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 58% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 60% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days. In embodiments, at least about 70% of the patients in the population of patients have taken at least one prior migraine therapy.

[0056] In embodiments, the present disclosure provides a method for the prevention of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant to a population of patients, wherein at least about 50% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 55% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days.

[0057] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least four weeks to a population of patients, wherein at least about 40%, or at least about 45%, or at least about 49% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the

preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least four weeks to a population of patients, wherein at least about 20% of patients, or at least about 25% of patients, or at least about 27% of patients, achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least four weeks to a population of patients, wherein at least about 7%, or at least about 10%, or at least about 12%, or at least about 14% of patients achieve 100% reduction in monthly migraine days.

[0058] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 50%, or at least about 55%, or at least about 60% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 30%, or at least about 35%, or at least about 39% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 15%, or about 17%, or about 21% of patients achieve a 100% reduction in monthly migraine days.

[0059] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 12 weeks to a population of patients, wherein at least

about 55%, or 60%, or 64% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 35%, or 40%, or 43% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 15%, or about 17%, or about 21% of patients achieve a 100% reduction in monthly migraine days.

[0060] In embodiments, the present disclosure provides a method for the prevention of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant to a population of patients, wherein at least about 50% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 55% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 58% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days.

[0061] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 4 weeks to a population of patients, wherein at least about 40%, or about 45%, or about 48%, or about 49% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 4 weeks to a population of patients, wherein at least about 20%, or about 25%, or about 26%, or about 27% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine

days per month), the method comprising administering 30 mg atogepant for at least four weeks to a population of patients, wherein at least about 7%, or at least about 10%, or at least about 11% of patients achieve 100% reduction in monthly migraine days.

[0062] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 50%, or at least about 55%, or at least about 60%, or at least about 61% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 30%, or at least about 35%, or at least about 36% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 15%, or about 17%, or about 18% of patients achieve a 100% reduction in monthly migraine days.

[0063] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 50%, or 55%, or 60% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 35%, or 40%, or 43% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4

and 14 migraine days per month), the method comprising administering 30 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 15%, or about 20%, or about 25%, or about 27% of patients achieve a 100% reduction in monthly migraine days.

[0064] In embodiments, the present disclosure provides a method for the prevention of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant to a population of patients, wherein at least about 50% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 55% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 58% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days, or at least about 60% of patients achieve $\geq 50\%$ reduction in a 3-month average of monthly migraine days.

[0065] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 4 weeks to a population of patients, wherein at least about 50%, or about 55%, or about 60%, or about 61% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 4 weeks to a population of patients, wherein at least about 25%, or about 30%, or about 35%, or about 39% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least four weeks to a population of patients, wherein at least about 10%, or at least about 15%, or at least about 19% of patients achieve 100% reduction in monthly migraine days.

[0066] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 8 weeks to a population of patients, wherein at least

about 50%, or at least about 55%, or at least about 60%, or at least about 65%, or at least about 66% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 30%, or at least about 35%, or at least about 40% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 8 weeks to a population of patients, wherein at least about 15%, or about 20%, or about 22%, or about 24% of patients achieve a 100% reduction in monthly migraine days.

[0067] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 50%, or at least about 60%, or at least about 65%, or at least about 70%, or at least about 71% of patients achieve $\geq 50\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 40%, or at least about 45%, or at least about 47%, or at least about 49% of patients achieve $\geq 75\%$ reduction in monthly migraine days. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant for at least 12 weeks to a population of patients, wherein at least about 15%, or at least about 20%, or at least about 25%, or at least about 27% of patients achieve a 100% reduction in monthly migraine days.

[0068] In embodiments, the present disclosure provides a method for prophylactically treating migraine in a patient in need thereof comprising administering atogepant in a therapeutically

effective amount, resulting in fewer headache days per month. In embodiments, treatment results in a change from baseline of at least 3.8 fewer mean monthly headache days, or at least 3.9 fewer monthly mean monthly headache days, or at least 4 fewer monthly mean monthly headache days, or at least 4.1 fewer monthly mean monthly headache days, or at least 4.2 fewer mean monthly
5 headache days.

[0069] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg QD, 30 mg QD, or 60 mg QD to a population of human patients, resulting in fewer headache days per month. In embodiments,
10 at least about 70% of patients in the population have taken at least one prior migraine therapy. In embodiments, treatment results in a change from baseline of at least 3.8 fewer mean monthly headache days, or at least 3.9 fewer monthly mean monthly headache days, or at least 4 fewer monthly mean monthly headache days, or at least 4.1 fewer monthly mean monthly headache days, or at least 4.2 fewer mean monthly headache days.

[0070] For example, in embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg QD to a population of human patients, resulting in fewer headache days per month. In embodiments, at
15 least about 70% of patients in the population have taken at least one prior migraine therapy. In embodiments, administering atogepant results in a change from baseline of at least 3.8 fewer mean monthly headache days, or at least 3.9 fewer monthly mean monthly headache days.

[0071] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen headache days per month), wherein atogepant is administered in an amount of 30 mg QD to a population of human
25 patients, resulting in fewer headache days per month. In embodiments, at least about 70% of patients in the population have taken at least one prior migraine therapy. In embodiments, treatment results in a change from baseline of at least 3.8 fewer mean monthly headache days, or at least 3.9 fewer monthly mean monthly headache days, or at least 4 fewer monthly mean monthly headache days.

[0072] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per
30

month), wherein atogepant is administered in an amount of 30 mg QD to a population of human patients, resulting in fewer headache days per month. In embodiments, at least about 70% of patients in the population have taken at least one prior migraine therapy. In embodiments, treatment results in a change from baseline of at least 3.8 fewer mean monthly headache days, or at least 3.9 fewer monthly mean monthly headache days, or at least 4 fewer monthly mean monthly headache days, or at least 4.1 fewer monthly mean monthly headache days, or at least 4.2 fewer mean monthly headache days.

[0073] In embodiments, the present disclosure provides a method for prophylactically treating migraine in a patient in need thereof, the method comprising administering atogepant in an amount of 10 mg, 30 mg, or 60 mg, resulting in fewer acute medication use days per month. In embodiments, treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 3.5 days, or at least 3.6 days, or at least 3.66 days, or at least 3.68 days, or at least 3.7 days, or at least 3.8 days, or at least 3.85 days.

[0074] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg QD, 30 mg QD, or 60 mg QD to a population of human patients, resulting in fewer acute medication use days per month. In embodiments, at least about 70% of the patients in the population of human patients have taken at least one prior migraine therapy. In embodiments, treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 3.5 days, or at least 3.6 days, or at least 3.66 days, or at least 3.68 days, or at least 3.7 days, or at least 3.8 days, or at least 3.85 days.

[0075] For example, in embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg QD. In embodiments, at least about 70% of the patients in the population of human patients have taken at least one prior migraine therapy. In embodiments, treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 3.5 days, or at least 3.6 days, or at least 3.66 days.

[0076] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per

month), wherein atogepant is administered in an amount of 30 mg QD. In embodiments, at least about 70% of the patients in the population of human patients have taken at least one prior migraine therapy. In embodiments, treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 3.5 days, or at least 3.6 days, or at least 3.66 days, or at least 3.68 days.

[0077] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 60 mg QD. In embodiments, at least about 70% of the patients in the population of human patients have taken at least one prior migraine therapy. In embodiments, treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 3.5 days, or at least 3.6 days, or at least 3.66 days, or at least 3.68 days, or at least 3.7 days, or at least 3.8 days, or at least 3.85 days.

[0078] In embodiments, the present disclosure provides methods for preventing or prophylactically treating migraine to result in improved patient function. In embodiments, the present disclosure provides methods for preventing migraine (such as a method of preventing migraine in patients having fewer than fifteen migraine days per month) by administering atogepant, resulting in an improvement in the physical impairment or quality-of-life impact scores reported by patients as compared to a pre-treatment baseline and/or a patient not receiving atogepant. Migraines can impact patient quality of life, impact productivity, and prevent patients from engaging in leisure and everyday activities. "Improved patient function" can be defined as an improvement measured by factors such as a reduced pain, reduced time spent in bed, increased ambulation, healthier attitude, more varied lifestyle and/or healing permitted by normal muscle tone. Improved patient function may be measured with an improved quality of life (QOL) or Health-Related Quality of Life (HRQL). These effects can be assessed, for example, using questionnaires or surveys, such as the Migraine-Specific Quality of Life Questionnaire (MSQ), the Headache Impact Test ("Headache Impact Test-6" or "HIT-6"), or the Activity Impairment in Migraine Diary (AIM-D). Scores obtained can be compared to published values available for various general and patient populations.

[0079] The Migraine-Specific Quality of Life Questionnaire Version 2.1 is one of the most frequently utilized disease-specific tools assessing the impact of migraine on HRQL. The MSQ is a validated tool that assess the impact of migraine on function (i.e., daily social and work-

related activities) over the past 4 weeks across three dimensions: Role Function-Restrictive (RR, or “RFR”, relating to how migraines limit the patient’s daily social and work related activities), Role Function-Preventive (RP, e.g., how migraines prevent the patient’s daily social and work related activities), and Emotional Function (EF, e.g., the emotions associated with a patient’s migraines). Patients respond to items using a 6-point scale: “none of the time,” “a little bit of the time,” “some of the time”, “a good bit of the time,” “most of the time” and “all of the time,” which are assigned scores of 1 to 6, respectively. Raw dimension scores are computed as a sum of item responses and rescaled from a 0 to 100 scale such that higher scores indicate better quality of life. See Bagley et al., *Validating Migraine-Specific Quality of Life Questionnaire v2.1 in Episodic and Chronic Migraine*, Headache, 2012 Mar; 52(3):409-21.

[0080] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg, 30 mg, or 60 mg once daily, and wherein administration of atogepant results in improvements in the ability to perform daily activities. In embodiments, the improvement in the ability to perform daily activities is measured by the MSQ v2.1. In embodiments, treatment with atogepant results in a change from baseline in the MSQ v2.1 RFR Domain of greater than about 21.0 points, such as greater than about 25.0 points, or greater than about 30.0 points, or greater than about 30.1 points, or greater than about 30.2 points, or greater than about 30.3 points, or greater than about 30.4 points, or greater than about 30.5 points, or greater than about 31.0 points, or greater than about 31.3 points relative to baseline.

[0081] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of about 10 mg QD, wherein the administration of atogepant results in an improvement in the ability to perform daily activities as measured by the MSQ v2.1. In embodiments, treatment with atogepant results in a change from baseline in the MSQ v2.1 RFR Domain of greater than about 25.0 points, or greater than about 30.0 points, or greater than about 30.1 points, or greater than about 30.2 points, or greater than about 30.3 points, or greater than about 30.4 points relative to baseline. For example, in embodiments, atogepant is administered in

an amount of about 10 mg QD, and the ability to perform daily activities as measured by the MSQ v2.1 RFR Domain improves by about 30.4 points relative to baseline.

[0082] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of about 30 mg QD, wherein administration of atogepant results in an improvement in the ability to perform daily activities as measured by the MSQ v2.1. In embodiments, treatment with atogepant results in a change from baseline in the MSQ v2.1 RFR Domain of greater than about 25.0 points, or greater than about 30.0 points, or greater than about 30.1 points, or greater than about 30.2 points, or greater than about 30.3 points, or greater than about 30.4 points, or greater than about 30.5 points relative to baseline. For example, in embodiments, atogepant is administered in an amount of about 30 mg QD, and the ability to perform daily activities as measured in the MSQ v2.1 RFR Domain improves by about 30.5 points relative to baseline.

[0083] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of about 60 mg QD, wherein administration of atogepant results in an improvement in the ability to perform daily activities as measured by the MSQ v2.1. In embodiments, treatment with atogepant results in a change from baseline in the MSQ v2.1 RFR Domain of greater than about 25.0 points, or greater than about 30.0 points, or greater than about 30.1 points, or greater than about 30.2 points, or greater than about 30.3 points, or greater than about 30.4 points, or greater than about 30.5 points, or greater than about 31.0 points, or greater than about 31.3 points relative to baseline. For example, in embodiments, atogepant is administered in an amount of about 60 mg QD, and the ability to perform daily activities as measured in the MSQ v2.1 RFR Domain improves by about 31.3 points relative to baseline.

[0084] The Activity in Migraine Diary (AIM-D) evaluates the impact of migraine on the performance of daily activities and physical impairment using an electronic daily diary. In particular, the AIM-D is comprised of two domains that evaluate Performance of Daily Activities (PDA) and Physical Impairment (PI). Assessed items for the AIM-D PDA domain include household chores, errands, leisure activities at home, social or leisure activities outside the home,

strenuous physical activities, concentration, and thinking clearly. Assessed items for the AIM-D PI Domain include walking, moving one's body, bending forward, and moving one's head.

Response options for each item range from "not difficult at all" to "I could not do it at all" on a 6-point rating scale. The AIM-D domain scores are scaled from 0-100, with higher scores

5 indicating greater impact of migraine, and reductions from baseline in scores indicate improvement. *See* Cala et al., The Activity Impairment in Migraine – Diary (AIM-D): A Novel Migraine-Specific Patient-Reported Outcome Measure to Assess Functioning Based on Activity Impairment in Episodic and Chronic Migraine Patients, MTIS2018-005 (September 5, 2018).

[0085] In embodiments, the present disclosure provides a method for preventing migraine (such
10 as a method for preventing migraine in a patient having fewer than fifteen migraine days per month), wherein atogepant is administered in an amount of 10 mg, 30 mg, or 60 mg once daily, wherein administration of atogepant results in improvements in performance of daily activities and physical impairment. In embodiments, the improvement is assessed using a daily diary. In embodiments, the daily diary is the Activity Impairment in Migraine Diary (AIM-D). In

15 embodiments, the daily activities include household chores, errands, leisure activities at home, social or leisure activities outside the home, strenuous physical activities, concentration, and thinking clearly. In embodiments, administration of atogepant results in less physical impairment in activities such as walking, moving the body, bending forward, and moving one's head.

[0086] In embodiments, the present disclosure provides a method for preventing migraine in a

20 patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, and reduces the patient's AIM-D

Performance of Daily Activities score relative to baseline by more than about 6 points, or more than about 7 points, or more than about 7.5 points or more than about 8 points, or more than

25 about 8.3 points, or more than about 8.6 points, or more than about 9 points, more than about 9.4

points. In embodiments, the present disclosure provides a method for preventing migraine in a

patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in

an amount of 10 mg QD, 30 mg QD, or 60 mg QD, and reduces the patient's AIM-D Physical

30 Impairment score by more than about 4 points, or more than about 5 points, or more than about 5.5 points, or more than about 6 points, or more than about 6.5 points relative to baseline.

[0087] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of about 30 mg QD, resulting in a reduction in the AIM-D Performance of Daily Activities score of more than about 7.5 points relative to baseline, such as more than about 7.7 points, or more than about 8 points, or more than about 8.2 points, or more than about 8.4 points, or more than about 8.6 points relative to baseline. In embodiments, the reduction in the AIM-D PDA Score is about 8.6 points relative to baseline (i.e., -8.6 relative to baseline).

[0088] In embodiments, the present disclosure provides a method for preventing migraine in a patient or a group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of about 60 mg QD, resulting in a reduction in the AIM-D Performance of Daily Activities Score of more than about 7.5 points relative to baseline, such as more than about 7.7 points, or more than about 8 points, or more than about 8.2 points, or more than about 8.4 points, or more than about 8.6 points, or more than about 8.8 points, or more than about 9 points, or more than about 9.2 points, or more than about 9.4 points relative to baseline. In embodiments, the reduction in the AIM-D PDA score is about 9.4 points relative to baseline (i.e., -9.4 relative to baseline).

[0089] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of about 30 mg QD, resulting in a reduction in the AIM-D Physical Impairment Score of more than about 5.2 points relative to baseline, such as more than about 5.5 points, or more than about 5.7 points, or more than about 6.0 points relative to baseline. In embodiments, the reduction in the AIM-D PI score is about 6.0 points relative to baseline (i.e., -6.0 relative to baseline).

[0090] In embodiments, the present disclosure provides a method for preventing migraine in a patient or group of patients (such as a method for preventing migraine in a patient or group of patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of about 60 mg QD, resulting in a reduction in the AIM-D Physical Impairment Score of more than about 5.2 points relative to baseline, such as more than about 5.5 points, or more

than about 5.7 points, or more than about 6.0 points, or more than about 6.3 points, or more than about 6.5 points relative to baseline. In embodiments, the reduction in the AIM-D PI Score is about 6.5 points relative to baseline (i.e., -6.5 relative to baseline).

[0091] In embodiments, the present disclosure provides a method for preventing migraine (such as a method for preventing migraine in patients having fewer than fifteen migraine days per month) wherein atogepant is administered in an amount of 10 mg QD, 30 mg QD, or 60 mg QD, wherein administration of atogepant results in a reduction in a 6-item Headache Impact Test (HIT-6) disability score relative to baseline. The Headache Impact Test (HIT-6) is a well-known tool for assessing migraine intensity, and uses six questions to measure the adverse impact of headache on, for example, social functioning, role functioning, vitality, cognitive functioning, and psychological distress. Responses are based on frequency using a 5-point verbal response scale ranging from “never” to “always”. The HIT-6 total score is the sum of the responses, and ranges from 36 to 78. *See Yang et al., Validation of the Headache Impact Test (HIT-6TM) Across Episodic and Chronic Migraine, Cephalalgia 2011 Feb; 31(3): 357-367.* In embodiments, treatment with atogepant results in a greater than five (≥ 5) point decrease in HIT-6 disability score.

[0092] Long-Term Treatment with Atogepant

[0093] In embodiments, atogepant may be administered daily for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks. Based on the clinical data, atogepant may be administered safely for at least up to 52 weeks.

[0094] For example, in embodiments, atogepant may be administered at a dose of 10 mg QD, or 30 mg QD, or 60 mg QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks.

[0095] For example, in an embodiment, atogepant may be administered in an amount of 60 mg once daily for at least 52 weeks. In embodiments, treatment with atogepant results in a reduction in mean monthly migraine days of at least 3.8 days, or at least 4.1 days, or at least 4.3 days, or at

least 4.6 days, or at least 4.7 days, or at least 4.8 days, or at least 4.9 days, or at least 5 days, or at least 5.1 days, or at least 5.19 days.

[0096] In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering atogepant in an amount of 10 mg, 30 mg, or 60 mg for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks. For example, in embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having between 4-14 migraine days per month, the method comprising administering atogepant 60 mg QD for at least 52 weeks, wherein the treatment results in a reduction in mean monthly migraine days of at least 5 days, or at least 5.1 days, or at least 5.19 days.

[0097] In embodiments, the present disclosure provides a method for the preventive treatment of migraine comprising administering 10 mg, 30 mg, or 60 mg atogepant for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, wherein treatment results in a $\geq 50\%$ improvement (reduction) in monthly migraine days at each monthly period. In embodiments, at least 60% of patients achieve $\geq 50\%$ improvement (reduction) in monthly migraine days at each monthly period, or at least 70%, or at least 80%. In embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month, the method comprising administering atogepant 60 mg QD for at least 52 weeks, wherein treatment results in at least 50% improvement (reduction) in monthly migraine days at each monthly period.

[0098] In embodiments, the present disclosure provides methods for the prevention of migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg, 30 mg, or 60 mg QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52

weeks, resulting in fewer headache days per month. For example, in embodiments, the present disclosure provides a method for the preventive treatment of migraine in patients having fewer than 15 migraine days per month, the method comprising administering atogepant 60 mg QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20
5 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, wherein treatment with atogepant results in a reduction from baseline in monthly headache days of at least 4 days, or at least 5 days, or at least 5.3 days, or at least 5.6 days, or at least 5.9 days, or at least 5.99 days. In embodiments, the present disclosure provides a method for the preventive treatment of
10 migraine, the method comprising administering atogepant 60 mg QD for at least 52 weeks, resulting in a reduction from baseline in monthly headache days of at least 5.9 days.

[0099] In embodiments, the present disclosure provides methods for the prevention of migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 10 mg, 30
15 mg, or 60 mg QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, resulting in a reduction in mean monthly acute medication use days. For example, in
20 embodiments, the present disclosure provides methods for the preventive treatment of migraine in patients having fewer than 15 migraine days per month, the method comprising administering 60 mg atogepant QD for at least 4 weeks, or 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks,
25 wherein treatment with atogepant results in a reduction in mean monthly acute medication use days of at least 4 days, or at least 4.4 days, or at least 4.5 days, or at least 4.6 days, or at least 4.7 days, or at least 4.9 days, or at least 4.93 days. In embodiments, the present disclosure provides methods for the preventive treatment of migraine, the method comprising administering atogepant 60 mg QD for at least 52 weeks, resulting in a change from baseline in acute monthly medication use days of about 4.93 days.

[0100] Atogepant and Weight Loss

[0101] Migraine and obesity are co-morbid conditions. Obesity has previously been linked both to an increased prevalence of migraine and also to increased migraine attack frequency leading to progression from episodic to chronic migraine. See Kristoffersen et al., *Migraine, Obesity, and Body Fat Distribution – a Population-Based Study*, J. Headache & Pain 21: 97 (2020); Bigal ME, Liberman JN, Lipton RB (2006) Obesity and migraine: a population study. Neurology. 66(4):545–550; Bigal ME, Tsang A, Loder E, Serrano D, Reed ML, Lipton RB (2007) Body mass index and episodic headaches: a population-based study. Arch Intern Med 167(18):1964–1970; Peterlin BL, Rosso AL, Rapoport AM, Scher AI (2010) Obesity and migraine: the effect of age, gender and adipose tissue distribution. Headache. 50(1):52–62; Scher AI, Stewart WF, Ricci JA, Lipton RB (2003) Factors associated with the onset and remission of chronic daily headache in a population-based study. Pain. 106(1–2):81–89; Gelaye B, Sacco S, Brown WJ, Nitchie HL, Ornello R, Peterlin BL (2017) Body composition status and the risk of migraine: a meta-analysis. Neurology. 88(19):1795–1804; Ornello R, Ripa P, Pistoia F, Degan D, Tiseo C, Carolei A et al (2015) Migraine and body mass index categories: a systematic review and meta-analysis of observational studies. J Headache Pain. 16:27; Winter AC, Berger K, Buring JE, Kurth T (2009) Body mass index, migraine, migraine frequency and migraine features in women. Cephalalgia. 29(2):269–278. Each of these references are incorporated by reference herein in their entireties.

[0102] 40% of patients with migraine disease are obese ($\text{BMI} \geq 30 \text{ kg / m}^2$). Further, migraine patients with $\text{BMI} \geq 27 \text{ kg / m}^2$ with obesity-related comorbidities (type 2 diabetes, hypertension, dyslipidemia, sleep apnea, cardiovascular disease). Treatment of obesity has been associated with both improved general health and improved migraine disease.

[0103] It has been discovered that certain dosages of atogepant, when administered over a sufficient period of time, are associated with weight loss.

[0104] Accordingly, in embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering atogepant to a patient in an amount effective to both prevent migraine and reduce the patient's body weight. For example, in embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering atogepant in an amount effective to reduce both the patient's monthly migraine days and the patient's body weight. In embodiments, the patient's body weight is reduced as compared to the patient's body weight before taking atogepant.

[0105] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering atogepant daily to a patient for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, wherein administration of atogepant results in a reduction in body weight by at least 0.5 kg, or at least 0.6 kg, or at least 0.7 kg, or at least 0.8 kg, or at least 0.9 kg, or at least 1 kg, or at least 1.1 kg, or at least 1.2 kg, or at least 1.3 kg, or at least 1.4 kg, or at least 1.42 kg. In embodiments, atogepant is administered daily in an amount of 10 mg, 30 mg, or 60 mg.

[0106] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg or 60 mg atogepant QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, resulting in a reduction in body weight by at least 0.5 kg, or at least 0.6 kg, or at least 0.7 kg, or at least 0.8 kg, or at least 0.9 kg, or at least 1 kg, or at least 1.1 kg, or at least 1.2 kg, or at least 1.3 kg, or at least 1.4 kg, or at least 1.42 kg. Based on the clinical data, atogepant may be administered safely for at least up to 52 weeks.

[0107] For example, in embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, resulting in a reduction in body weight by at least 0.5 kg, or at least 0.6 kg.

[0108] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg atogepant QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, wherein the mean difference for percentage change from baseline in body weight is at least about 0.7%, or at least about 0.8%, or at least about 0.9%, or at least about 0.98%. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 30 mg atogepant QD for at least about 12 weeks, wherein the mean difference for percentage change from baseline in body weight is about 0.98%. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 30 mg once daily for at least about 12 weeks, wherein treatment with atogepant results in at least about 3.2% of patients achieving a weight decrease of greater than about 7% from baseline.

[0109] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, resulting in a reduction in body weight by at least about 0.5 kg, or at least about 0.6 kg, or at least about 0.7 kg, or at least about 0.8 kg, or at least about 0.9 kg, or at least about 1 kg, or at least about 1.1 kg, or at least about 1.2 kg, or at least about 1.27 kg. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg atogepant QD for at least 4 weeks, resulting in a reduction in body weight from baseline of at least about 0.6 kg, or at least about 0.7 kg, or at least about 0.8 kg, or at least about 0.81 kg. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method

comprising administering 60 mg atogepant QD for at least about 52 weeks, resulting in a reduction in body weight from baseline of at least about 1.1 kg, or at least about 1.2 kg, or at least about 1.3 kg, or at least about 1.4 kg, or at least about 1.42 kg.

[0110] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant QD for at least 4 weeks, or at least 8 weeks, or at least 12 weeks, or at least 16 weeks, or at least 20 weeks, or at least 24 weeks, or at least 28 weeks, or at least 32 weeks, or at least 36 weeks, or at least 40 weeks, or at least 44 weeks, or at least 48 weeks, or at least 52 weeks, wherein the mean difference for percentage change from baseline in body weight is at least about 1.0%, or at least about 1.3%, or at least about 1.5%, or at least about 1.6%. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg atogepant QD for at least about 12 weeks, wherein the mean difference for percentage change from baseline in body weight is about 1.64%.

[0111] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg QD atogepant for at least about 12 weeks, or at least about 16 weeks, or at least about 20 weeks, or at least about 24 weeks, or at least about 28 weeks, or at least about 32 weeks, or at least about 36 weeks, or at least about 40 weeks, or at least about 44 weeks, or at least about 48 weeks, or at least about 52 weeks to a population of patients, wherein at least about 10% of patients, or at least about 12% of patients, or at least about 14% of patients, achieve a weight decrease of greater than about 7% from baseline. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, such as a method for preventing migraine in patients having fewer than 15 migraine days per month (such as ≤ 14 migraine days per month, or between 4 and 14 migraine days per month), the method comprising administering 60 mg atogepant once daily for at least about 12 weeks, wherein treatment with atogepant results in at least about 4.9% of patients achieving a weight decrease of greater than about 7% from baseline. In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising

administering 60 mg QD atogepant for at least about 6 months to a population of patients, wherein at least about 15% of patients, or at least about 18% of patients, or at least about 10% of patients, or at least about 22% of patients, achieve a weight decrease of greater than about 5% from baseline.

5 [0112] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg QD atogepant for at least about six months to a population of patients, wherein at least about 8 % of patients, or at least about 10% of patients, or at least about 12% of patients, achieve a weight decrease of greater than about 7% from baseline.

10 [0113] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg QD atogepant for at least about 9 months to a population of patients, wherein at least about 15% of patients, or at least about 18% of patients, or at least about 10% of patients, or at least about 22% of patients, or at least about 25% of patients, achieve a weight decrease of greater than about 5% from baseline.

15 [0114] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg QD atogepant for at least about nine months to a population of patients, wherein at least about 8% of patients, or at least about 10% of patients, or at least about 12% of patients, or at least about 14% of patients, achieve a weight decrease of greater than about 7% from baseline.

20 [0115] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg QD atogepant for at least about 12 months to a population of patients, wherein at least about 18% of patients, or at least about 20% of patients, or at least about 22% of patients, or at least about 25% of patients, achieve a weight decrease of greater than about 5% relative to baseline. In embodiments, the present disclosure
25 provides a method for the preventive treatment of migraine, the method comprising administering 60 mg atogepant QD for at least about 12 months, wherein at least about 22.4% of patients achieve a weight decrease of greater than about 5% relative to baseline body weight.

[0116] In embodiments, the present disclosure provides a method for the preventive treatment of migraine, the method comprising administering 60 mg QD atogepant for at least about 12
30 months to a population of patients, wherein at least about 10% of patients, or at least about 12%

of patients, or at least about 15% of patients, achieve a weight decrease of greater than about 7% from baseline.

[0117] Definitions

[0118] "Administration" or "to administer" means the step of giving (i.e., administering) a pharmaceutical composition to a subject, or alternatively a subject receiving a pharmaceutical composition.

[0119] Calcitonin-Gen-Related-Peptide, or CGRP, encompasses any member of the calcitonin family, including any calcitonin gene related peptide and analogs, calcitonin, amylin, adrenomedullin and their analogs.

[0120] "CGRP antagonist" refers to any molecule that exhibits any one or more of the following characteristics: (a) bind to CGRP or CGRP-R and the binding results in a reduction or inhibition of CGRP activity; (b) block CGRP from binding to its receptor(s); (c) block or decrease CGRP receptor activation; (d) inhibit CGRP biological activity or downstream pathways mediated by CGRP signaling function; (e) increase clearance of CGRP; and (f) inhibit or reduce CGRP synthesis, production or release. CGRP antagonists include but are not limited to antibodies to CGRP, antibodies to the CGRP-R, small molecules that antagonize CGRP, and small molecules that antagonize CGRP-R.

[0121] The term "prophylactic" or "preventative" refers to the prevention of the onset, recurrence or spread of a disease or disorder, or of one or more symptoms associated with the disease or disorder. In one embodiment, such symptoms are those known to a person of skill in the art to be associated with the disease or disorder being prevented. In certain embodiments, the terms refer to the treatment with or administration of a compound provided herein prior to the onset of symptoms, particularly to patients at risk of disease or disorders provided herein. The terms encompass the inhibition or reduction of a symptom of the particular disease. For frequent migraines, prophylactic or preventative treatments are employed to reduce the frequency of migraines and also to reduce the severity and duration of migraines and their associated symptoms when they occur.

[0122] A "prophylactically effective amount" of a compound is an amount sufficient to prevent a disease or disorder, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of therapeutic agent that can provide a prophylactic or preventative benefit in the prevention of the disease. The term "prophylactically effective amount" can encompass an

amount that improves overall prophylaxis or prevention or enhances the prophylactic or preventative efficacy of another prophylactic agent.

[0123] "Effective amount" as applied to the biologically active ingredient means that amount of the ingredient which is generally sufficient to effect a desired change in the subject. For example, where the desired effect is a reduction in an autoimmune disorder symptom, an effective amount of the ingredient is that amount which causes at least a substantial reduction of the autoimmune disorder symptom, and without resulting in significant toxicity.

[0124] "Headache day" refers to any calendar day on which headache pain lasting two hours or longer occurs unless an acute headache medication (e.g., ibuprofen, triptan) was used after the start of the headache, in which case no minimum duration is specified.

[0125] "Migraine day" refers to any calendar day on which a headache occurs which meets criteria A, B, and C, or meets criteria D and E, as provided herein:

(A) Headache has at least two of the following four characteristics:

- (i) unilateral location,
- (ii) pulsating quality,
- (iii) moderate or severe pain intensity,
- (iv) aggravated by or causing avoidance of routine physical activity (e.g., walking or climbing stairs)

(B) At least one of the following:

- (i) Nausea and/or vomiting;
- (ii) Photophobia and phonophobia;
- (iii) Typical aura (i.e., visual, sensory, or speech/language) accompanying or

within sixty minutes before headache begins

(C) Duration of headache lasting two hours or longer on a calendar day unless an acute, migraine-specific medication (i.e., triptan or ergot derivative) was used after the start of the headache, in which case no minimum duration is specified

OR

(D) Any headache which fulfills one criterion from (1) and at least one criterion from (2)
OR fulfills at least two criteria from (1) and no criteria from (2):

(1) Headache characteristics:

- (i) Unilateral location;

- (ii) Pulsating quality;
- (iii) Moderate or severe pain intensity;
- (iv) Aggravated by or causing avoidance of routine physical activity (e.g.,

walking or climbing stairs)

5 (2) Symptoms

- (i) Nausea and/or vomiting;
 - (ii) Photophobia and phonophobia;
 - (iii) Typical aura (i.e., visual, sensory, or speech/language) accompanying
- or within sixty minutes before headache begins.

10 (E) Duration of headache lasting two hours or longer on a calendar day unless an acute, migraine-specific medication (i.e., triptan or ergot derivative) was used after the start of the headache, in which case no minimum duration is specified.

[0126] "Patient" means a human or non-human subject receiving or in need of medical or veterinary care. Accordingly, the compositions as disclosed herein can be used in treating any
15 animal, such as, for example, mammals, or the like.

[0127] "Pharmaceutical composition" means a composition comprising an active pharmaceutical ingredient, such as, for example, a CGRP antagonist, and at least one additional ingredient, such as, for example, a stabilizer or excipient or the like. A pharmaceutical composition is therefore a formulation which is suitable for diagnostic or therapeutic administration to a subject, such as a
20 human patient. The pharmaceutical composition can be, for example, in a lyophilized or vacuum dried condition, a solution formed after reconstitution of the lyophilized or vacuum dried pharmaceutical composition, or as a solution or solid which does not require reconstitution.

[0128] "Treating" means to alleviate (or to eliminate) at least one symptom of a condition or disorder, such as, for example, sinusitis, nausea, nasopharyngitis, photophobia, appetite changes, cognition and concentration difficulties, cold extremities, diarrhea or other bowel changes, excitement or irritability, fatigue, frequent urination, memory changes, weakness, yawning, stretching, seeing bright spots or flashes of light, vision loss, seeing dark spots, tingling
25 sensations, speech problems, aphasia, tinnitus, gastric stasis, pulsating or throbbing pain on one or both sides of the head, extreme sensitivity to light, sounds, or smells, worsening pain during physical activity, and vomiting, abdominal pain or heartburn, loss of appetite, lightheadedness,
30 blurred vision or fainting or the like, either temporarily or permanently.

[0129] The invention of the present disclosure will be more clearly understood by reference to the following examples, which are included for purposes of illustration only and are not intended to be limiting.

[0130] EXAMPLE 1

5 [0131] Study A: A phase 3, multicenter, randomized, double-blind, placebo-controlled, parallel-group study was conducted to evaluate the efficacy, safety, and tolerability of oral atogepant for the prevention of migraine in participants with episodic migraine (EM).

[0132] This study comprised a 4-week screening and baseline period, a 12-week double-blind treatment period, and a 4-week follow-up period. The total study duration was 20 weeks.

10 [0133] Adults 18 to 80 years of age with 4 to 14 migraine days per month in the 3 months prior to visit 1, and 4 to 14 migraine days during the 28-day baseline period per electronic Diary, were allowed to enroll. Inclusion criteria included at least a 1-year history of migraine with or without aura consistent with a diagnosis, diagnosed as specified in the International Classification of Headache Disorders, 3rd edition, version (ICHD-3), and the age of the participant at the time of
15 migraine onset was <50 years. Exclusion criteria included a history of migraine accompanied by diplopia or decreased level of consciousness or retinal migraine; a current diagnosis of chronic migraine, new persistent daily headache, trigeminal autonomic cephalgia (e.g., cluster headache), or painful cranial neuropathy as defined by ICHD-3, or if they averaged 15 or more migraine days per month across the 3 months prior to Visit 1 or during the 28-day baseline period; a
20 history of an inadequate response to >4 medications (2 of which have different mechanisms of action) prescribed for the treatment of migraine; and participants with clinically significant hematologic, endocrine, cardiovascular, pulmonary, renal, hepatic, gastrointestinal, or neurologic disease. Use of opioids or barbiturate on more than 2 days per month, triptans or ergots on 10 or more days per month, or simple analgesics (e.g., aspirin, non-steroidal anti-inflammatory drugs, acetaminophen) on 15 or more days per month in the 3 months prior to Visit 1 or during the 28-
25 day baseline period were also excluded. Use of barbiturates was also excluded 30 days prior to screening and throughout the duration of the trial.

[0134] Randomization was stratified based on prior exposure (yes/no) to a migraine prevention medication with proven efficacy, such as antiepileptics, tricyclic antidepressants, beta-blockers,
30 calcium channel blocker, angiotensin receptor blocker or converting enzyme inhibitor, or serotonin-norepinephrine reuptake inhibitors. Approximately 70% of randomized participants

took at least one prior migraine prevention medication with proven efficacy. Participants were not on concurrent medication during the study.

[0135] The trial included 910 randomized participants (223 to placebo, 222 to atogepant 10 mg, 230 to atogepant 30 mg, and 235 to atogepant 60 mg). The safety population included 902 participants and the modified-intent-to-treat population (efficacy analysis population) included 873 participants; >87% of participants (805/910, 88.5%) completed the double-blind treatment period across all treatment groups. Baseline demographics and clinical characteristics were similar across treatment groups in the safety population. Participants were between 18 and 73 years of age, with a mean age of 41.6 years. 89% of participants were female, and 83% white, with a baseline BMI of 30.6 kg/m². At screening, 99.3% of participants reported use of medication for the acute treatment of migraine and 70.3% reported having previously used a preventive treatment for migraine, with or without proven efficacy. The baseline demographics of the safety population (i.e., all participants who received at least one dose of study intervention) are summarized in Table 1.

TABLE 1: Baseline Demographics (Safety Population*)

Baseline Demographics		Placebo (N=222)	Atogepant 10 mg QD (N=221)	Atogepant 30 mg QD (N=228)	Atogepant 60 mg QD (N=231)	Total (N=902)
Age	Mean (SD) (years)	40.3 (12.81)	41.4 (12.05)	42.1 (11.68)	42.5 (12.41)	41.6 (12.25)
	Min, max	18, 69	18, 73	19, 70	18, 72	18, 73
Sex	Male	24 (10.8)	21 (9.5)	24 (10.5)	32 (13.9)	101 (11.2)
	Female	198 (89.2)	200 (90.5)	204 (89.5)	199 (86.1)	801 (88.8)
Race, n (%) **	White	194 (87.4)	191 (81.9)	195 (81.1)	192 (83.1)	752 (83.4)
	Black or African American	24 (10.8)	34 (15.4)	38 (16.7)	28 (12.1)	124 (13.7)
	Asian	2 (0.9)	2 (0.9)	1 (0.4)	7 (3.0)	12 (1.3)
	American Indian or Alaska Native	0 (0.00)	1 (0.5)	1 (0.4)	1 (0.4)	3 (0.3)
	Multiple***	2 (0.9)	3 (1.4)	3 (1.3)	2 (0.9)	10 (1.1)
Ethnicity	Non-Hispanic	89.6%	90.5%	91.7%	93.9%	91.5%
BMI	Mean (SD) (kg/m ²)	30.83 (8.713)	30.35 (7.597)	31.15 (7.631)	29.91 (7.318)	30.56 (7.828)
	Min, max	16.9, 82.0	16.4, 60.3	17.1, 66.2	16.3, 57.3	16.3, 82.0
Prior Exposure to a Migraine Prevention Medication with Proven Efficacy	Yes	71.2%	67.0%	70.6%	68.0%	69.2%
Number of headache days per month in the last 3 months	Mean (SD)	9.5 (2.8)	9.3 (2.7)	9.2 (2.7)	9.1 (2.7)	9.3 (2.7)
	Min, Max	4, 14	4, 14	4, 14	4, 14	4, 14
Number of migraine days per month in the last 3 months	Mean (SD)	7.7 (2.6)	7.2 (2.5)	7.3 (2.4)	7.3 (2.4)	7.4 (2.5)
	Min, Max	4, 14	4, 14	4, 14	4, 13	4, 14

* The safety population included all randomized participants who took at least 1 dose of trial treatment.

** Excludes one participant with missing data randomized to atogepant 60 mg.

5 *** Only includes participants who reported multiple races.

[0136] A total of 910 patients were randomized 1:1:1:1 to one of the four treatment groups (10 mg QD, 30 mg QD, 60 mg QD, or placebo). Efficacy analyses were based on the Modified Intent-to-Treat Population (mITT) of 873 patients. The Modified Intent-to-Treat (mITT)

5 Population includes all randomized participants who received at least one dose of study intervention, had an evaluable baseline period of eDiary data, and had at least one evaluable post-baseline 4-week period (Weeks 1 to 4, 5 to 8, and 9 to 12) of eDiary data during the double-blind treatment period.

[0137] All participants were instructed to take trial treatment orally at approximately the same
10 time once daily for 12 weeks. To maintain the blinding, all participants took 3 tablets of trial treatment provided in identical blister cards: placebo (placebo 10 mg / placebo 30 mg / placebo 60 mg), atogepant 10 mg (atogepant 10 mg / placebo 30 mg / placebo 60 mg), atogepant 30 mg (placebo 10 mg, atogepant 30 mg, placebo 60 mg), and atogepant 60 mg (placebo 10 mg / placebo 30 mg / atogepant 60 mg). Participants, site personnel, and trial sponsor personnel were
15 blinded to the treatment assignment. Tablet formulations for CGRP compounds are described, for example, in U.S. Patent No. 10,117,936.

[0138] Participants were allowed to take acute treatments for migraine which included triptans, ergot derivatives, opioids, acetaminophen, non-steroidal anti-inflammatory drugs, and antiemetic agents. Participants were not allowed to take any preventive treatments for migraine 30 days
20 prior to visit 1 and throughout the trial.

[0139] Efficacy assessments were recorded by the participant in an electronic diary at home or via eTablet at the trial site during clinic visits. Headache duration, headache clinical features (headache pain severity, unilateral location, aggravated by or causing avoidance of routine physical activity), non-headache associated symptoms (nausea and/or vomiting; photophobia, phonophobia, and aura), and acute medication use were recorded. Additional health outcomes
25 measures were collected.

[0140] The primary efficacy endpoint was change from baseline in mean monthly migraine days across the 12-week treatment period. Secondary efficacy endpoints, tested in hierarchical order, were change from baseline in mean monthly headache days across the 12-week treatment period, change from baseline in mean monthly acute medication use days across the 12 week treatment
30 period, $\geq 50\%$ reduction in 3-month average of monthly migraine days, change from baseline in

MSQ v 2.1 Role Function-Restrictive domain score at week 12, change from baseline in mean monthly Performance of Daily Activities domain score of the AIM-D across the 12-week treatment period, and change from baseline in mean monthly Physical Impairment domain score of the AIM-D across the 12-week treatment period. An exploratory analysis of the time course of efficacy for atogepant by 4-week intervals, based on the least square mean change from baseline for the number of monthly migraine days over the 12-week treatment period was also conducted. [0141] The primary efficacy variable was the change from baseline in mean monthly migraine days across the 12-week treatment period. The results are shown in Table 2. Figures 1 and 2 show the least square mean (+/- standard error of the least squares (SE)) of change from baseline in monthly migraine days (MMRM – mixed effects model for repeated measures) during the double-blind treatment period for the mITT population. Figure 3 shows the distribution of change from baseline in monthly migraine days across 12 weeks by treatment group.

TABLE 2: Primary Endpoint – Change from Baseline in Mean Monthly Migraine Days Across the 12-Week Treatment Period (mITT Population)

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Baseline number of monthly migraine days, mean (SD)	7.51 (2.39)	7.45 (2.46)	7.86 (2.32)	7.75 (2.31)
LS Mean Change (SE)	-2.48 (0.21)	-3.69 (0.21)	-3.86 (0.21)	-4.20 (0.21)
LSMD vs. Placebo (95% CI)	-	-1.21 (-1.78, -0.64)	-1.38 (-1.94, -0.82)	-1.72 (-2.28, -1.15)
Nominal p-value	-	<.0001	<.0001	<.0001
Adjusted p-value	-	<.0001	<.0001	<.0001

SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI = Confidence Interval; LSMD = least squares mean difference.

[0142] As shown in Table 2 and Figures 1, 2, and 3, all atogepant dose groups met the primary endpoint and demonstrated statistically significantly greater decreases in monthly migraine days compared to placebo. Patients treated in the 10 mg, 30 mg, and 60 mg atogepant arms experienced a decrease of 3.69, 3.86, and 4.20 days, respectively, all compared to patients in the placebo arm, who experienced a decrease of 2.48 days (all dose groups vs. placebo, $p < .0001$).

[0143] The LS mean change from baseline in monthly migraine days across 12 weeks by treatment group is shown in Figure 4. Figure 5A shows the mean change from baseline in weekly migraine days during the first month (weeks 1-4) by treatment group. During weeks 1-4 of treatment, mean change from baseline in mean monthly migraine days was -3.1 for atogepant 10 mg, -3.4 for atogepant 30 mg, -3.9 for atogepant 60 mg, and -1.6 for placebo. This significantly greater decrease in monthly migraine days was maintained during the subsequent two 4-week intervals of double-blind treatment for all atogepant doses (weeks 5-8: -3.7 for atogepant 10 mg, -3.9 for atogepant 30 mg, -4.2 for atogepant 60 mg, and -2.9 for placebo, $P \leq 0.012$ for all atogepant groups; weeks 9-12: -4.2 for atogepant 10 mg, -4.3 for atogepant 30 mg, -4.4 for atogepant 60 mg, and -3.0 for placebo, $P \leq 0.0002$ for all atogepant groups).

[0144] During the 28-day baseline period, the mean weekly migraine days were 1.9 in the overall mITT population. Mean change from baseline in weekly migraine days during the first week of treatment was -0.77 atogepant 10 mg, -0.94 atogepant 30 mg, -1.03 atogepant 60 mg, and -0.29 placebo ($P < 0.0001$ all dose groups). The greater reduction in weekly migraine days compared with placebo was consistent throughout each week of the first 4-week interval of treatment ($P \leq 0.0397$ all dose groups). On the first full day after study drug administration, the proportion of participants who reported a migraine day was 14.1% for atogepant 10 mg, 10.8% for atogepant 30 mg, and 12.3% for atogepant 60 mg versus 25.2% in the placebo group ($P \leq 0.0071$ all dose groups). The proportions of participants with a migraine each day during the first week of treatment (mITT population) are shown in Figure 5B. On post dose days 2-6, the proportions of participants reporting a migraine was consistently lower across the treatment groups compared with placebo, with the majority of days reaching significance vs. placebo ($P < 0.05$) in the atogepant 30 mg and 60 mg dose groups. The odds ratio vs placebo for reporting a migraine on post dose day 1 was 0.49 with atogepant 10 mg, 0.33 with atogepant 30 mg, and 0.39 with atogepant 60 mg. Atogepant provided an early and sustained reduction in migraine days including statistically significant reductions in each of the three 4-week intervals, each week during the first 4-week interval, and as early as the first full day after study drug initiation.

[0145] A secondary endpoint measured the proportion of patients that achieved a 50% reduction in mean monthly migraine days across the 12-week treatment period. The results are shown in Table 3. The response profile for percent reduction from baseline in 3-month average of monthly migraine days (mITT population) is set forth in Figures 6 and 7. In particular, Figure 6 shows the

response profile for percent reduction from baseline in 3-month average of monthly migraine days for the mITT population. Figure 7 shows the percent of participants with a $\geq 50\%$ reduction in 3-month average monthly migraine days.

TABLE 3: $\geq 50\%$ Reduction in 3-Month Average of Monthly Migraine Days (mITT Population)

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Responder, n (%)	62 (29.0)	119 (55.6)	131 (58.7)	135 (60.8)
Odds ratio vs. Placebo (95% CI)	-	3.06 (2.05, 4.56)	3.53 (2.37, 5.26)	3.82 (2.56, 5.71)
Nominal p-value	-	<.0001	<.0001	<.0001
Adjusted p-value	-	<.0001	<.0001	<.0001

CI = confidence interval

[0146] As shown in Table 3, 55.6%, 58.7%, and 60.8% of patients in the 10 mg, 30 mg, and 60 mg atogepant arms, respectively, achieved at least a 50% reduction in mean monthly migraine days across the 12-week treatment period, compared to 29% of patients in the placebo arm (all dose groups vs. placebo, $p < .0001$).

[0147] Prespecified additional endpoints included proportions of participants achieving $\geq 25\%$, $\geq 75\%$, and 100% reductions in the 12-week average of MMDs. A 100% reduction in MMDs represented individuals reporting no migraine days from the day the participant received the first dose of study treatment (day 1) through the end of week 12. The results are shown in Tables 4-6.

Figure 8A illustrates the proportion of participants achieving various responder rates by treatment group across the 12-week treatment period.

TABLE 4: $\geq 25\%$ Reduction in 3-Month Average of Monthly Migraine Days

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Responder, n (%)	126 (58.9)	157 (73.4)	172 (77.1)	180 (81.1)

$P < .0018$ for all comparisons

TABLE 5: ≥ 75 Reduction in 3-Month Average of Monthly Migraine Days

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Responder, n (%)	23 (10.7)	65 (30.4)	66 (29.6)	84 (37.8)

$P < .0001$ for all comparisons

TABLE 6: 100% Reduction in 3-Month Average of Monthly Migraine Days

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Responder, n (%)	2 (0.9)	17 (7.9)	11 (4.0)	17 (7.7)

5 $P < .0207$ for all comparisons

[0148] Atogepant-treated participants were significantly more likely to experience a $\geq 50\%$ reduction in mean MMDs across 12 weeks (range 56%-61% vs 29% with placebo, $P < 0.0001$).

Atogepant treated patients were also significantly more likely than placebo-treated participants to experience $\geq 25\%$ (range 73%-81% vs 59% for placebo, $P < 0.01$), $\geq 75\%$ (range 30%-38% vs 11% for placebo, $P < 0.0001$), and 100% (range 5%-8%, vs 1% for placebo, $P < 0.05$) reductions in MMDs across 12 weeks.

[0149] Figure 8B provides a cumulative distribution function graph of the percent reduction from baseline in 12-week average MMDs. As shown in Figure 8B, response rates were better for each of the three atogepant groups vs. placebo. A consistently higher proportion of participants in each of the three atogepant groups showed improvement in 12-week average MMDs compared with placebo across all levels of improvement that were evaluated.

[0150] Additional prespecified exploratory endpoints included $\geq 25\%$, $\geq 50\%$, $\geq 75\%$, and 100% reductions in MMDs by 4-week intervals (weeks 1-4, 5-8, 9-12). Figure 8C shows the proportion of participants with $\geq 25\%$ reduction in mean monthly migraine days, and 8D shows the proportion of participants with $\geq 50\%$, $\geq 75\%$, and 100% reduction in monthly migraine days across the 12-week treatment period, for month 1 (weeks 1-4), month 2 (weeks 5-8) and month 3 (weeks 9-12). Response to atogepant treatment was evident as early as the first four weeks of treatment and increased over time. All differences for $\geq 50\%$, $\geq 75\%$, and 100% responder rates

by 4-week intervals were statistically significant in favor of atogepant. For $\geq 25\%$ responder rates, all differences were statistically significant in favor of atogepant vs placebo, except for atogepant 10 mg for weeks 5-8.

[0151] Relative to placebo, atogepant treated participants were ≥ 3 times more likely to achieve $\geq 50\%$ reduction in the 12-week average of MMDs. At all doses, atogepant was effective in the first 4 weeks for $\geq 25\%$, $\geq 50\%$, $\geq 75\%$, and 100% responders, suggesting robust treatment effects in the first month. The proportion of participants experiencing a $\geq 25\%$, $\geq 50\%$, $\geq 75\%$, and 100% reduction in mean MMDs significantly increased with duration of treatment for all atogepant doses and efficacy was sustained for three months, noting that placebo response rates also increased throughout the study period. The responder rates were higher with increasing doses among the atogepant treatment groups.

[0152] A secondary endpoint measured the change from baseline in mean monthly headache days across the 12-week treatment period. The results are shown in Table 7.

TABLE 7: Change from Baseline in Mean Monthly Headache Days across the 12-Week Treatment Period (mITT Population)

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Baseline number of mean monthly headache days, mean (SD)	8.43 (2.55)	8.41 (2.75)	8.78 (2.62)	9.00 (2.56)
LS Mean Change (SE)	-2.52 (0.23)	-3.94 (0.22)	-4.04 (0.22)	-4.23 (0.22)
LSMD vs. Placebo (95% CI)	-	-1.42 (-2.03, -0.81)	-1.53 (-2.13, -0.92)	-1.71 (-2.32, -1.10)
Nominal p-value	-	<.0001	<.0001	<.0001
Adjusted p-value	-	<.0001	<.0001	<.0001

SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI = Confidence Interval; LSMD = least squares mean difference.

[0153] As shown in Table 7, patients in the 10 mg, 30mg, and 60 mg atogepant arms experienced a decrease of 3.94, 4.04, and 4.23 days, respectively, all compared to patients in the placebo arm, who experienced a decrease of 2.52 days (all dose groups vs. placebo, $p < .0001$).

[0154] The mean change from baseline in moderate / severe headache days per month (mITT population) is shown in Figure 9A, and the mean change from baseline in monthly headache days per month (mITT population) is shown in Figure 9B. The LS mean change from baseline in moderate/severe headache days in the first treatment period (weeks 1-4) was -3.0 for atogepant 10mg, -3.2 for atogepant 30 mg, -3.8 for atogepant 60 mg, and -1.7 for placebo ($P < 0.0001$ for all atogepant groups). The LS mean change from baseline in mean headache days in the first treatment period was -3.2 for atogepant 10 mg, -3.4 for atogepant 30 mg, -3.8 for atogepant 60 mg, and -1.4 for placebo ($P \leq 0.0001$ for all atogepant groups).

[0155] A secondary endpoint measured change from baseline in mean monthly acute medication use days across the 12-week treatment period. The results are shown in Table 8 and Figure 10A.

TABLE 8: Change from Baseline in Mean Monthly Acute Medication Use Days Across the 12-Week Treatment Period (mITT Population)

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Baseline number of monthly acute medication use days, mean (SD)	6.48 (3.15)	6.57 (2.99)	6.69 (3.02)	6.89 (3.17)
LS Mean Change (SE)	-2.35 (0.18)	-3.66 (0.18)	-3.68 (0.18)	-3.85 (0.18)
LSMD vs. Placebo (95% CI)	-	-1.31 (-1.81, -0.82)	-1.33 (-1.82, -0.83)	-1.50 (-2.00, -1.01)
Nominal p-value	-	<.0001	<.0001	<.0001
Adjusted p-value	-	<.0001	<.0001	<.0001

SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI = Confidence Interval; LSMD = least squares mean difference.

[0156] As shown in Table 8 and Figure 10A, patients in the 10 mg, 30 mg, and 60 mg atogepant arms experienced a decrease of 3.66, 3.68, and 3.85 days, respectively, across the 12-week treatment period compared to patients in the placebo arm, who experienced a decrease of 2.35 days (all dose groups vs. placebo, $p < .0001$).

[0157] The LS mean change from baseline in acute medication use days during the first treatment period (weeks 1-4) was -3.3 for atogepant 10 mg, -3.4 for atogepant 30 mg, -3.7 for

atogepant 60 mg, and -1.7 for placebo ($P<0.0001$ for all atogepant groups). The LS mean change in acute medication use days showed a nominally significant difference from placebo starting in the first treatment period and persisting into the second and third treatment periods, as shown in Figure 10B. Baseline mean cumulative headache hours ranged from 47.4 – 51.1 in the mITT population. The LS mean reduction from baseline in mean cumulative headache hours during the first treatment period was -23.3 for atogepant 10 mg, -23.6 for atogepant 30 mg, -25.1 for atogepant 60 mg, and -9.5 for placebo ($P<0.0001$ for all atogepant groups). LS mean reduction from baseline in mean cumulative headache hours is illustrated in Figure 10C.

Table 9: Baseline Parameters on Efficacy Measures (mITT Population)

Baseline Parameters, mean (SD)	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Monthly Cumulative Headache Hours	51.1 (34.5)	47.4 (27.3)	49.5 (26.7)	50.4 (27.4)
Monthly Moderate / Severe Headache days	6.5 (2.6)	6.4 (2.6)	6.9 (2.5)	6.9 (2.6)
Weekly migraine days*	1.9 (0.6)	1.9 (0.6)	2.0 (0.6)	1.9 (0.6)

* For weekly data, baseline was defined as monthly migraine days divided by 4, and change from baseline in weekly migraine days was calculated for consecutive 7-day periods beginning with day 1.

[0158] A secondary endpoint measured the change from baseline in MSQ v2.1 Role Function Restrictive Domain Score at Week 12. The Migraine-Specific Quality of Life Questionnaire Version 2.1 is one of the most frequently utilized disease-specific tools assessing the impact of migraine on HRQL. The MSQ measures the impact of migraine on the patient's HRQL over the past 4 weeks across three dimensions: Role Function-Restrictive (RR), Role Function-Preventive (RP), and Emotional Function (EF). The results from this secondary endpoint analysis are shown in Table 10.

TABLE 10: Change from Baseline in MSQ v2.1 Role Function Restrictive Domain* Score at Week 12 (mITT Population)

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Baseline MSQ v2.1 role function restrictive domain score, mean (SD)	46.8 (19.67)	44.9 (21.37)	44.0 (19.61)	46.8 (20.36)
Week 12, mean (SD)	66.3 (23.9)	75.6 (23.5)	75.4 (24.2)	78.0 (25.0)
Change from baseline, mean (SD)	19.5 (24.3)	30.8 (27.6)	31.1 (24.9)	30.5 (25.1)
MMRM**, LS Mean Change (SE)	20.45 (1.62)	30.35 (1.64)	30.53 (1.59)	31.25 (1.59)
LSMD vs. Placebo (95% CI)	-	9.90 (5.45, 14.36)	10.08 (5.71, 14.46)	10.80 (6.42, 15.18)
Nominal p-value	-	<.0001	<.0001	<.0001
Adjusted p-value	-	<.0001	<.0001	<.0001

SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI =

5 Confidence Interval; LSMD = least squares mean difference.

*Role Function-Restrictive domain assesses how migraines limit one's daily social and work-related activities. Participants respond to items using a 6-point scale ranging from "none of the time" to "all of the time." Raw domain scores are rescaled to a 0 to 100 scale, where higher scores indicate better quality of life. Items included leisure time activities; work or daily

10 activities; getting done as much at work or home; concentrate on work or daily activities; left you too tired; dealt with family, friends, and others; felt energetic.

** The MMRM model includes baseline as a covariate, prior exposure to migraine prevention medications (y/n), treatment group, and visit (month) as fixed factors, and treatment group by visit and baseline-by-visit as interaction terms, with an unstructured covariance matrix. P values

15 are from the test between the atogepant dose group and placebo.

[0159] As shown in Table 10, at week 12, statistically significant and clinically meaningful improvements for all dose groups vs placebo were observed in the RFR domain (10 mg, 9.9 [5.5 – 14.4]; 30 mg, 10.1 [5.7-14.5]; 60 mg, 10.8 [6.4 – 15.2]), in the RFP domain (10 mg, 5.8 [1.9 –

5.6]; 30 mg, 6.9 [3.1 – 10.7], 60 mg, 7.1 [3.3 – 10.9]), and in the EF domain (10 mg, 8.3 [3.4 – 13.1]; 30 mg, 9.7 [4.9-14.4]; 60 mg, 10.5 [5.8 – 15.3]). Figure 11 illustrates the Migraine-Specific Quality of Life Questionnaire Role Function-Restrictive Domain Score at Weeks 4, 8, and 12. As shown in Figure 11, for the RFR domain, significant differences vs. placebo were

5 observed at the earliest time point assessed (week 4) and seen throughout the treatment period. [0160] In people with 4-14 migraine days per month, oral atogepant produced significant and clinically meaningful between- and within-group reductions in emotional impact, improvements in functioning in daily social and work-related activities, the ability to undertake daily activities that had previously been prevented by migraine, and reduction in the emotional effect of

10 migraine, as measured by the MSQ v 2.1. These results indicate that preventive treatment with atogepant is associated with statistically significant and clinically meaningful improvements in all domains of migraine-specific quality of life.

[0161] A secondary endpoint evaluated the change from baseline in mean monthly performance of daily activities domain score of the AIM-D across the 12-week treatment period. Another

15 secondary endpoint evaluated the change from baseline in mean monthly physical impairment domain score of the AIM-D across the 12-week treatment period. The Activity Impairment in Migraine – Diary (AIM-D) is an 11-item daily diary that is comprised of two domains that evaluate Performance of Daily Activities (PDA; 7 items) and Physical Impairment (PI; 4 items). *See* Cala et al., “The Activity Impairment in Migraine Diary (AIM-D): A novel migraine-specific

20 patient-reported outcome measure to assess functioning based on activity impairment in episodic and chronic migraine patients”, *Cephalalgia* 2018; 38:1-115, which is incorporated by reference herein in its entirety. In particular, the AIM-D assesses the impact of migraine on the performance of daily activities and physical impairment using a 6-point rating scale ranging from “Not difficult at all,” “A little difficult,” “Somewhat difficult,” “Very Difficult,” “Extremely

25 Difficult”, and “I could not do it at all.” Items assessed for the AIM-D PDA domain included household chores, errands, leisure activities at home, social or leisure activities outside the home, strenuous physical activities, concentration, and thinking clearly. Items assessed for the AIM-D PI Domain included walking, moving body, bending forward, and moving head. Raw domain scores are recalled to a 0 to 100 scale, where higher scores indicate greater impact of migraine,

30 and reductions from baseline in scores indicate improvement. The AIM-D was collected daily via an electronic diary with the same set of questions administered in headache (HA) and non-

headache (NHA) versions. Monthly domain scores are calculated by summing the non-missing daily domain scores (HA and NHA days combined) and dividing the number of non-missing daily scores. The secondary endpoints for AIM-D PDA and PI domain scores were evaluated as change from baseline in mean monthly PDA or PI domain scores of the AIM-D across the 12-week treatment period. The results with respect to daily activity are shown in Table 11. The results with respect to change from baseline in mean monthly physical impairment domain score are shown in Table 12.

TABLE 11: Change from Baseline in Mean Monthly Performance of Daily Activities
Domain Score of the AIM-D across the 12-week treatment period**

mITT Population	Placebo (n=214)	Atogepant 10 mg QD (n=214)	Atogepant 30 mg QD (n=223)	Atogepant 60 mg QD (n=222)
Change from baseline to week 12	n=178	n=182	n=181	n=178
Baseline monthly performance of daily activities domain score, mean (SD)	15.2 (8.25)	15.5 (8.85)	16.9 (8.02)	15.9 (8.34)
Month 1-3 ^a , mean (SD)	9.6 (7.2)	8.6 (9.5)	8.0 (8.1)	6.6 (7.1)
Change from baseline, mean (SD)	-5.5 (7.6)	-7.0 (8.1)	-8.9 (7.5)	-9.2 (7.4)
MMRM ^b , LS mean change (SE)	-6.1 (0.5)	-7.3 (0.5)	-8.6 (0.5)	-9.4 (0.5)
MMRM ^b , LS Mean Change (SE)	-6.09 (0.50)	-7.28 (0.50)	-8.63 (0.50)	-9.41 (0.50)
LSMD vs. Placebo (95% CI)	-	-1.19 (-2.56, 0.17)	-2.54 (-3.91, -1.18)	-3.32 (-4.68, -1.96)
Nominal p-value	-	0.0856	0.0003	<.0001
Adjusted p-value	-	0.0856	0.0005	<.0001

AIM-D, Activity Impairment in Migraine-Diary; SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI = Confidence Interval; LSMD = least squares mean difference; mITT, modified intent to treat; MMRM, mixed-effects model for repeated measures for change from baseline; QD, once daily.

^aMonth 1-3 = average of monthly Performance of Daily Activity domain scores across the 12-week treatment period

^bThe MMRM model includes baseline as a covariate, prior exposure to migraine prevention medications (y/n), treatment group, and visit (month) as fixed factors, and treatment group by visit and baseline-by-visit as interaction terms, with an unstructured covariance matrix. P values are from the test between the atogepant dose group and placebo.

** Items included household chores, errands, leisure activities at home, social or leisure activities outside the home, strenuous physical activities, concentration, and thinking clearly.

TABLE 12: Change from Baseline in Mean Monthly Physical Impairment Domain*
Score of the AIM-D Across the 12-Week Treatment Period (mITT population)**

	Placebo (N=214)	Atogepant 10 mg QD (N=214)	Atogepant 30 mg QD (N=223)	Atogepant 60 mg QD (N=222)
Baseline monthly physical impairment domain score, mean (SD)	11.2 (8.11)	11.7 (8.46)	13.0 (8.00)	11.6 (7.85)
LS Mean Change (SE)	-4.03 (0.44)	-5.11 (0.44)	-6.02 (0.44)	-6.49 (0.44)
LSMD vs. Placebo (95% CI)	-	-1.08 (-2.27, 0.11)	-1.99 (-3.18, -0.80)	-2.46 (-3.65, -1.28)
Nominal p-value	-	0.0743	0.0011	<.0001
Adjusted p-value	-	0.0856	0.0021	0.0002

SD = Standard Deviation; LS = Least Squares; SE = Standard error of the least squares; CI = Confidence Interval; LSMD = least squares mean difference.

*** Items included walking, moving body, bending forward, and moving head.

[0162] As shown in Tables 11 and 12, compared with placebo, all atogepant groups demonstrated improvement of function in PDA and PI domain scores across the 12-week treatment period. For both AIM-D domains, differences were statistically significant for participants in the atogepant 60 mg and 30 mg groups (least squares mean difference [LSMD] vs placebo: PDA, -3.32 for 60 mg, -2.54 for 30 mg; PI, -2.46 for 60 mg, -1.99 for 30 mg). The improvement in PDA and PI domain scores for the atogepant 10 mg group did not reach statistical significance vs. placebo (LSMD: -1.19 and -1.08, respectively).

[0163] As shown above, with respect to the secondary endpoints, treatment with 30mg and 60 mg atogepant doses resulted in statistically significant improvements in all secondary endpoints, and treatment with 10 mg atogepant resulted in statistically significant improvements in four out of the six secondary endpoints. Atogepant demonstrated significant improvements in patients' functioning in daily activities, and reductions in physical impairment and the impact of headaches, confirming its role as a promising treatment among people with migraine.

[0164] The Headache Impact Test (HIT-6, completed monthly) represented an additional exploratory measure. The HIT-6 test is a well-known tool for assessing migraine intensity that uses six questions to capture the impact of headache and its treatment on an individual's

functional health and well-being. HIT-6 was completed monthly and was evaluated as the change from baseline in HIT-6 total score at weeks 4, 8, and 12. HIT-6 results are shown in Table 13.

Table 13: HIT-6: Change from Baseline to Week 12

mITT Population	Placebo (n=214)	Atogepant 10 mg QD (n=214)	Atogepant 30 mg QD (n=223)	Atogepant 60 mg QD (n=222)
Change from baseline to week 12	n=178	n=182	n=181	n=178
Baseline, mean (SD)	64.5 (4.5)	64.1 (5.4)	64.3 (4.9)	63.6 (5.6)
Week 12, mean (SD)	59.2 (6.5)	55.8 (8.3)	56.1 (8.6)	54.5 (9.0)
Change from baseline, mean (SD)	-5.3 (6.8)	-8.3 (7.8)	-8.1 (7.9)	-9.9 (8.7)
MMRM ^b , LS mean change (SE)	-5.2 (0.5)	-8.4 (0.5)	-8.1 (0.5)	-9.2 (0.5)
LSMD vs placebo (95% CI)		-3.2 (-4.7, -1.7)	-2.9 (-4.3, -1.4)	-4.0 (-5.4, -2.5)
P value		<0.0001	0.0001	<0.0001
Responders^c, n/N (%)				
Week 4	127/206 (38.3)	101/208 (48.6)	102/219 (46.8)	122/217 (56.2)
Odds ratio vs. Placebo (95% CI)		1.53 (1.03, 2.27)	1.43 (0.97, 2.12)	2.18 (1.47, 3.25)
P value		0.0386	0.738	0.0001
Week 8	96/203 (47.3)	113/187 (60.4)	130 / 209 (62.2)	134 / 207 (64.7)
Odds ratio vs. Placebo (95% CI)		1.79 (1.19, 2.68)	1.95 (1.31, 2.90)	2.25 (1.50, 3.37)
P value		0.0053	0.0011	<0.0001
Week 12	103/198 (52.0)	120 / 186 (64.5)	128 / 204 (62.7)	131 / 201 (65.2)
Odds ratio vs. Placebo (95% CI)		1.74 (1.15, 2.62)	1.54 (1.03, 2.30)	1.77 (1.18, 2.66)
P value		0.0086	0.0334	0.0055

HIT-6, 6-item headache impact test; LSMD, least squares mean difference; mITT, modified intent to treat; MMRM, mixed-effects model for repeated measures change from baseline; QD, once daily; SD, standard deviation; SE, standard error of the least squares.

- 5 ^bThe MMRM model includes baseline as a covariate, prior exposure to migraine prevention medications (y/n), treatment group, and visit (month) as fixed factors, and treatment group by

visit and baseline-by-visit as interaction terms, with an unstructured covariance matrix. P values are from the test between the atogepant dose group and placebo.

^cResponders defined as patients experiencing a ≥ 5 point reduction in the HIT-6 score.

[0165] All atogepant groups demonstrated significant improvement in HIT-6 scores vs. placebo at weeks 4, 8, and 12. Significantly greater proportions of atogepant vs. placebo-treated participants were HIT-6 responders (≥ 5 point decrease) with all atogepant doses (except 30mg at week 4).

[0166] The AIM-D and HIT-6 results demonstrate that atogepant significantly improved patient functioning in daily activities, and reduced physical impairment and impact of headaches.

[0167] Additional prespecified responder analyses included the Patient Global Impression of Change (PGI-C), defined as “much better” or “very much better” at week 12, and satisfaction with study medication, defined as “satisfied” or “extremely satisfied” at weeks 4, 8, and 12. The results are shown in Table 14.

Table 14: PGI-C and Treatment Satisfaction After 12 Weeks of Treatment (mITT Population)

Responder Criteria, Week 12	Placebo (n=214)	Atogepant 10 mg (n=214)	Atogepant 30 mg (n=223)	Atogepant 60 mg (n=222)
PGI-C, ^a n/N	95/206	145/201	153/209	162/213
Percentage	46.1%	72.1%	73.2%	76.1%
OR (95% CI)	---	3.05 (2.01, 4.61)	3.33 (2.20, 5.05)	3.83 (2.52, 5.84)
P value		<0.0001	<0.0001	<0.0001
Satisfaction with Study Medication ^b	109/199	146/188	162/203	166/201
Percentage	54.8%	77.7%	80.3%	82.6%
OR (95% CI)	---	2.77 (1.81, 4.24)	3.44 (2.23, 5.30)	3.58 (2.31, 5.54)
P value		<0.0001	<0.0001	<0.0001

^aResponse defined as “much better” or “very much better”

^bResponse defined as “satisfied” or “extremely satisfied.”

OR, odds ratio; PGI-C, Patient Global Impression of Change.

[0168] Significantly higher proportions of patients receiving atogepant were responders at week 12 for PGI-C (72%-76%) and were satisfied with treatment (78% - 83%) vs placebo (46% and 55%, $P < 0.0001$), as shown in Table 14. The odds of achieving PGI-C response in each atogepant group was ≥ 3 times that of placebo. A greater proportion of participants in each of the three atogepant arms met treatment satisfaction responder criteria at weeks 4, 8, and 12, compared with placebo, as shown in Table 14. The odds for being treatment satisfaction responders at week 12 for atogepant were approximately 2-4 times greater than placebo.

[0169] Participants treated with atogepant were more satisfied with treatment and reported greater improvement in their migraine attacks compared with placebo. Even if a participant did not achieve the thresholds for being considered a treatment responder, a high proportion of participants (70-80%) reported feeling “much better” or “very much better”, demonstrating that some individuals experience a treatment benefit without exceeding specified responder definitions.

[0170] The study further showed that atogepant was safe and well-tolerated. Adverse events were reported by participants throughout the trial, and at a 4-week follow-up visit. Adverse event information was collected and documented during each clinic visit. Participants could also report adverse events via telephone call between visits. Causality of each adverse event was determined by the investigator who was blinded to the treatment. In addition, clinical laboratory tests, vital signs, electrocardiograms (ECG), and Columbia-Suicide Severity Rating Scale were evaluated. Based on the potential hepatotoxicity of prior gepants, treatment-emergent elevations in ALT or $AST \geq 3$ times the upper limit of normal, as well as potential Hy’s law cases, were evaluated as prespecified adverse events of special interest and were reviewed by an independent panel of liver experts blinded to treatment (see, e.g., Negro A et al., CGRP Receptor Antagonists: an expanding drug class for acute migraine? *Expert Opin Investig Drugs* 2012; 21:807-18; Messina R et al., CGRP – a target for acute therapy in migraine: Clinical Data. *Cephalalgia* 2019;39:420-7).

[0171] Rates of adverse events (AEs) were similar across all treatment groups. Treatment-emergent adverse events were reported in 53.9% of participants (486 of 902 participants); the frequency of events was similar between placebo and atogepant treatment groups and no dose relationship was observed. Serious adverse events occurred in 0.9% of patients treated in the

atogepant 10 mg arm compared to 0.9% of patients in the placebo arm (2 participants in both the placebo and atogepant 10 mg groups). In particular, serious adverse events were reported in 2 participants treated with placebo (gastric ulcer hemorrhage; post-surgical laryngospasm with hypoxic brain injury) and 2 participants treated with atogepant 10 mg (asthma attack; optic neuritis). The asthma attack was considered unrelated to trial treatment and clinical evidence did not support the diagnosis of optic neuritis.

[0172] No patients in the atogepant 30 mg or 60 mg treatment arms experienced a serious adverse event. The most common adverse events reported with a frequency of $\geq 5\%$ in at least one atogepant treatment arm, and greater than placebo, were constipation (6.9-7.7% across all doses vs. 0.5% for placebo), nausea (4.4-6.1% across all doses vs. 1.8% for placebo), and upper respiratory tract infection (3.9-5.7% across all doses vs. 4.5% for placebo). The majority of cases of constipation, nausea and upper respiratory tract infection were mild or moderate in severity and did not lead to discontinuation. Cases of constipation were primarily mild (71.4%) or moderate (26.5%) in severity. One case of constipation was considered severe in the atogepant 10 mg group. This was reported as a worsening of pre-existing constipation and the participant was treated with over-the-counter medication, completed the trial, and entered an open-label extension study. All reported cases of nausea were mild (77.1%) or moderate (22.9%) in severity.

[0173] Rates of discontinuation due to AEs were low across all treatment groups and not dose-dependent. Rates of discontinuation for the highest doses were the same or lower than placebo. The rates of adverse events are summarized in Table 15:

Table 15: Adverse Events

	Placebo (N=222) n(%)	Atogepant 10mg QD (N=221) n(%)	Atogepant 30mg QD (N=228) n(%)	Atogepant 60mg QD (N=231) n(%)
Adverse Event	126 (56.8)	117 (52.9)	119 (52.2)	124 (53.7)
Serious AE	2 (0.9)	2 (0.9)	0	0
AE leading to Discontinuation	6 (2.7)	9 (4.1)	4 (1.8)	6 (2.6)
Treatment Emergent Adverse Event	126 (56.8)	117 (52.9)	119 (52.2)	124 (53.7)
TEAEs reported in >= 2% of participants in any treatment group				
Constipation	1 (0.5)	17 (7.7)	16 (7.0)	16 (6.9)
Upper Respiratory Tract Infection	10 (4.5)	9 (4.1)	13 (5.7)	9 (3.9)
Nausea	4 (1.8)	11 (5.0)	10 (4.4)	14 (6.1)
Urinary Tract Infection	8 (3.6)	3 (1.4)	9 (3.9)	9 (3.9)
Nasopharyngitis	8 (3.6)	4 (1.8)	8 (3.5)	8 (3.5)
Fatigue	4 (1.8)	3 (1.4)	7 (3.1)	9 (3.9)
Somnolence	2 (0.9)	7 (3.2)	4 (1.8)	4 (1.7)
Blood creatine phosphokinase increased	2 (0.9)	5 (2.3)	2 (0.9)	7 (3.0)
Sinusitis	3 (1.4)	4 (1.8)	3 (1.3)	5 (2.2)
Gastroenteritis	4 (1.8)	2 (0.9)	5 (2.2)	3 (1.3)

Alanine aminotransferase increased	6 (2.7)	3 (1.4)	2 (0.9)	2 (0.9)
Influenza	2 (0.9)	3 (1.4)	2 (0.9)	5 (2.2)
Sinus Congestion	5 (2.3)	1 (0.5)	2 (0.9)	4 (1.7)
Aspartate aminotransferase increased	6 (2.7)	2 (0.9)	2 (0.9)	1 (0.4)
Anxiety	2 (0.9)	2 (0.9)	1 (0.4)	5 (2.2)
Treatment-related TEAE	20 (9.0)	51 (23.1)	34 (14.9)	45 (19.5)
Treatment-emergent serious adverse event	2 (0.9)	2 (0.9)	0 (0.0)	0 (0.0)
Treatment related TESAE	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)
Deaths	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
TEAE leading to treatment discontinuation	6 (2.7)	9 (4.1)	4 (1.8)	6 (2.6)

Safety population included all randomized participants who took at least 1 dose of trial treatment. Participants are only counted once within each category. Percentages are calculated at $100 \times (n/N)$, where n is the number of participants within a specific category and N is the number of participants in the safety population for the treatment group. Adverse events are listed in decreasing order based on frequency reported in the overall population.

[0174] There were no hepatic safety issues identified in this trial. ALT and/or AST increases were low across all atogepant doses and occurred at a lower rate than placebo. Cases of ALT or AST elevations that were ≥ 3 x upper limit of normal (ULN) occurred at a lower rate compared to placebo (where 4 cases were reported), and did not increase with an increase in dose. No cases of liver disease were reported, and no case met the criteria for potential Hy's Law following daily dosing for 12 weeks. Tables 16 and 17 summarize postbaseline hepatic laboratory parameter values.

Table 16: Postbaseline Hepatic Laboratory Parameter Values of Clinical Interest – Safety Population

Parameter Criterion	Placebo (N=222) n/N1	Atogepant 10 mg QD (N=221) n/N1	Atogepant 30 mg QD (N=228) n/N1	Atogepant 60 mg QD (N=231) n/N1
ALT or AST (U/L)				
$\geq 1 * \text{ULN}$	35/220 (15.9)	19/220 (8.6)	18/225 (8.0)	28/228 (12.3)
$\geq 1.5 * \text{ULN}$	14/220 (6.4)	3/220 (1.4)	7/225 (3.1)	7/228 (3.1)
$\geq 2 * \text{ULN}$	6/220 (2.7)	2/220 (0.9)	2/225 (0.9)	4/228 (1.8)
$\geq 3 * \text{ULN}$	4/220 (1.8)	2/220 (0.9)	2/225 (0.9)	1/228 (0.4)
$\geq 5 * \text{ULN}$	1/220 (0.5)	1/220 (0.5)	1/225 (0.4)	1/228 (0.4)
$\geq 10 * \text{ULN}$	0	1/220 (0.5)	0	0
$\geq 20 * \text{ULN}$	0	0	0	0
Potential Hy's Law				
ALT or AST $\geq 3 * \text{ULN}$ <u>AND</u> Bilirubin Total $\geq 2 * \text{ULN}$ <u>AND</u> ALP $< 2 * \text{ULN}$	0	0	0	0

N1 = Number of participants with at least one non-missing postbaseline value.

5 n = Number of participants within a specific category.

ULN = Upper limit of normal value

Table 17: Hepatic Safety Summary

	Placebo (N=222)	Atogepant 10 mg QD (N=221)	Atogepant 30 mg QD (N=228)	Atogepant 60 mg QD (N=231)
ALT or AST (U/L), n/N1 (%) ≥3x upper limit of normal (ULN)	4/220 (1.8)	2/220 (0.9)	2/225 (0.9)	1/228 (0.4)
Potential Hy's Law ALT or AST ≥3x ULN AND Bilirubin Total ≥2x ULN AND ALP <2x ULN	0	0	0	0

N1 = number of participants with at least one non-missing postbaseline value

n = number of participants within a specific category.

[0175] In summary, as shown above, atogepant 10 mg, 30 mg, and 60 mg groups demonstrated statistically significant and clinically meaningful improvements over the placebo group for the primary efficacy endpoint (reduction in mean monthly migraine days). There was a clinically relevant dose response relationship. Significant improvements were also observed for all secondary endpoints, with atogepant 30 mg and 60 mg meeting all six secondary endpoints, and atogepant 10 mg meeting four secondary endpoints. Atogepant 10 mg, 30 mg, and 60 mg groups demonstrated statistically significant and clinically meaningful improvements over the placebo group in the proportion of patients with 50% reduction in mean monthly migraine days (56-61% reduction in monthly migraine days vs. 29% for placebo).

[0176] Atogepant was safe and well-tolerated. The most common TEAEs were constipation (~7%) and nausea (~5%). No hepatic safety issues were identified.

[0177] Example 2

[0178] Study B: A phase 3, multicenter, randomized, open label study was conducted to evaluate the long-term safety and tolerability of oral atogepant for the prevention of migraine in participants with episodic migraine (Study B). The study objective was to evaluate the safety and tolerability of daily treatment with atogepant 60 mg QD when administered over 52 weeks for the prevention of migraine in participants with episodic migraine.

[0179] Figure 12 provides a schematic for the long-term safety study design. The study comprised a 4-week baseline/screening period, followed by a 52-week open-label treatment period, and a 4-week safety follow-up period.

[0180] A total of 744 participants from 106 sites were randomized in a ratio of 5:2 to the following treatment groups: atogepant 60 mg QD or oral standard of care (SOC) migraine prevention medication. The latter arm was included to contextualize safety data from the atogepant-treated participants, and efficacy measures were collected from the atogepant arm only. Efficacy measures were evaluated using the modified intent-to-treat (mITT) population and a mixed-effects model for repeated measures and included changes from baseline in monthly migraine days (MDDs), moderate/severe headache days, and acute medication use days, as well as the proportion of responders based on reductions in MMDs.

[0181] A total of 739 patients (n=546 atogepant) were included in the safety population. Participants included: (1) eligible participants who completed study NCT02848326 (visit 8) without significant protocol deviations (e.g., non-compliance with protocol-required procedures) and who did not experience an AE that, in the investigator's opinion, may indicate an unacceptable safety risk; and (2) de novo participants – adults (18-80 years) with a history (≥ 1 year) of migraine and 4-14 migraine days per month.

[0182] For participants in the SOC group, participants who did not tolerate the initially prescribed migraine preventive medication, or for whom the medication was not sufficiently effective (per investigator judgment), the investigator was allowed to prescribe an alternative medication or not prescribe any migraine preventive medication. Regardless of which of these options was chosen, the participant was allowed to continue in the study. In addition, the investigator had the option to select alternative preventive medication, or discontinue migraine prevention medication, for a participant in the SOC migraine preventive medication arm as often as needed throughout the study.

[0183] Table 18 provides a summary of the subject population.

Table 18: Subject Population

Population	Standard of Care	Atogepant 60 mg QD	Overall
Screened, N	--	--	1727
Intent-to-Treat, N	198	546	744
De Novo, N	170	467	637
CGP-MD-01 Completers, N	28	79	107
Safety, N	196	543	739
Modified Intent-to- Treat, N	--	521	521

[0184] All Screened Participants include those screened participants that signed informed consent forms. The Intent-to-Treat Population (ITT) includes all of the randomized participants.

- 5 The safety population includes all participants who received at least one dose of study intervention (atogepant or SOC medication). The modified intent-to-treat population (mITT) includes all randomized participants who received at least one dose of atogepant, had an evaluable baseline period of eDiary data, and had at least one evaluable post-baseline 4-week period of eDiary data.

- 10 [0185] Table 19 provides a summary of patient disposition during the open-label treatment period for all randomized participants.

Table 19: Participant Disposition During Open-Label Treatment Period – All Randomized Participants

Disposition	Standard of Care (N=198) n (%)	Atogepant 60 mg QD (N=546) n (%)	Overall (N=744) n (%)
Number of Participants Entered	198 (100.0)	546 (100.0)	744 (100.0)
Number of Participants Completed	136 (68.7)	373 (68.3)	509 (68.4)
Number of Participants Discontinued	62 (31.3)	173 (31.7)	235 (31.5)
Reason for Discontinuation			
Adverse events	5 (2.5)	31 (5.7)	36 (4.8)
Lack of Efficacy	2 (1.0)	5 (0.9)	7 (0.9)
Withdrawal by Subject	29 (14.6)	75 (13.7)	104 (14.0)
Lost to Follow-Up	16 (8.1)	23 (4.2)	39 (5.2)
Pregnancy	2 (1.0)	4 (0.7)	6 (0.8)
Protocol Deviation	7 (3.5)	31 (5.7)	38 (5.1)
Non-compliance with study drug	1 (0.5)	3 (0.5)	4 (0.5)
Other	0 (0.0)	1 (0.2)	1 (0.1)

Table 20 provides a summary of the baseline demographics of the safety population. Table 21 provides the migraine history of the safety population.

5 **Table 20: Baseline Demographics (Safety Population)**

Baseline Demographics		Standard of Care (N=196)	Atogepant 60 mg QD (N=543)	Total (N=739)
Age	Mean (SD) years	41.1 (12.09)	42.5 (12.03)	42.2 (12.05)
Sex	Female	87.8%	88.2%	88.1%
Race	White	74.0%	76.6%	75.9%
Ethnicity	Hispanic	14.8%	15.3%	15.2%
BMI	Mean (SD) (kg/m ²)	30.60 (8.028)	30.54 (7.440)	30.55 (7.595)
Prior Exposure to a Migraine Prevention Medication with Proven Efficacy	Yes	23.5%	27.6%	26.5%

Table 21: Migraine History (Safety Population)

Parameter		Standard of Care (N=196)	Atogepant 60 mg QD (N=543)	Total (N=739)
Migraine Diagnosis	With aura /without aura / both	24.0% / 42.3% / 33.7%	21.4% / 41.1% / 37.6%	21.1% / 41.4 % / 36.5 %
Migraine Disorder Duration in Years	Mean (SD)	19.6 (12.41)	20.9 (12.38)	20.6 (12.39)
Migraine Prevention Medication in the Past	Yes	28.1%	33.0%	31.7%
Average number of migraine days per month in the last 3 months	Mean (SD)	7.2 (2.56)	7.3 (2.59)	7.3 (2.58)
Average number of headache days per month in the last 3 months	Mean (SD)	9.4 (2.72)	9.2 (2.74)	9.3 (2.74)
Acute Medication Treatment	Yes	97.4%	99.3%	98.8%
Advice on Lifestyle Alterations	Yes	49.5%	51.2%	50.7%

[0186] Treatment duration in summarized in Table 22.

Table 22: Treatment Duration (Study Population)

Treatment Duration (days)	Standard of care (N=196)	Atogepant 60 mg QD (N=543)
Algorithm	Treatment Duration = Last treatment date – first treatment date + 1	Treatment duration = last study treatment date – first study treatment date + 1
Mean (SD)	278.9 (143.08)	291.6 (123.40)
≥ 30 days	180 (91.8)	508 (93.6)
≥ 90 days	160 (81.6)	459 (84.5)
≥ 180 days	140 (71.4)	428 (78.8)
≥ 270 days	123 (62.8)	397 (73.1)
≥ 360 days	115 (58.7)	362 (66.7)

[0187] Adverse events were reported by 67.0% of participants treated with atogepant; 18.0% of participants reported AEs that were considered related to atogepant by the investigator. Most commonly reported AEs ($\geq 5\%$ of participants) were upper respiratory tract infection (10.3%), constipation (7.2%), nausea (6.3%), and urinary tract infection (5.2%) following treatment with atogepant. Serious AEs were reported by 4.4% of participants treated with atogepant, which included a broad variety of common medical conditions; no event was seen in more than one participant and no event was considered related to atogepant. Two deaths were reported in participants treated with atogepant (victim of homicide and a group A beta-hemolytic streptococcal sepsis [toxic shock syndrome]); both were considered not related to atogepant. Discontinuation due to AEs was 5.7% following treatment with atogepant. Table 23 summarizes the adverse events in the safety population.

Table 23: Summary of Adverse Events (Safety Population)

	Standard of care (N=196) n (%)	Atogepant 60 mg QD (N=543) n (%)
Treatment-emergent adverse event (TEAE)	154 (78.6)	364 (67.0)
Treatment-related TEAE	71 (36.2)	98 (18.0)
Death	0 (0.0)	2 (0.4)
Treatment-emergent serious adverse events (TESAE)	7 (3.6)	24 (4.4)
TEAE leading to study discontinuation	5 (2.6)	31 (5.7)

[0188] Cases of alanine aminotransferase / aspartate aminotransferase (ALT/AST) levels ≥ 3 times the upper limit of normal were reported for 2.4% of participants (n=13/531) treated with atogepant and 3.2% for standard-of-care (n=6/190). No cases of potential Hy's Law were reported. Table 24 provides postbaseline hepatic laboratory parameter values of clinical interest for the safety population.

Table 24: Postbaseline Hepatic Laboratory Parameter Values of Clinical Interest (Safety Population)

Parameter (unit) Criteria	Standard of care (N=196) n / N1	Atogepant 60 mg QD (N=543) n / N1
ALT or AST (U/L)		
≥ 1 x ULN	55 / 190 (28.9)	112 / 531 (21.1)
≥ 1.5 x ULN	16 / 190 (8.4)	46 / 531 (8.7)
≥ 2 x ULN	11 / 190 (5.8)	26 / 531 (4.9)
≥ 3 x ULN	6 / 190 (3.2)	13 / 531 (2.4)
≥ 5 x ULN	1 / 190 (0.5)	7 / 531 (1.3)
≥ 10 x ULN	0 / 190 (0.0)	3 / 531 (0.6)
≥ 20 x ULN	0 / 190 (0.0)	0 / 531 (0.0)
Potential Hy's Law (ALT or AST ≥ 3 x ULN <u>and</u> Bilirubin Total ≥ 2 x ULN <u>and</u> ALP < 2 x ULN)	0 / 190 (0.0)	0 / 531 (0.0)

N1 = number of participants with at least one non-missing baseline value.

n = number of participants within a specific category

- 5 [0189] Efficacy endpoints for this study included change from baseline in monthly migraine days at each monthly period; change from baseline in monthly headache days at each monthly period; change from baseline in monthly acute medication use days at each monthly period; and ≥ 50% improvement (reduction) in monthly migraine days at each monthly period.

- 10 [0190] LS mean change from baseline in monthly migraine days over time (mITT population) is shown in Figure 13. As shown in Figure 13, atogepant 60 mg led to a reduction from baseline in monthly migraine days of approximately 4 days in month 1, followed by additional improvement over time for the remaining months. That is, a rapid reduction in monthly migraine days was observed in month 1, with additional gradual improvement over time through week 52. The change from baseline in monthly migraine days over time (mITT population) is shown in Table 25.
- 15

Table 25: Change from Baseline in Monthly Migraine Days Over Time (mITT Population)

Derived Visit	N	Atogepant 60 mg QD (N=521)	
		Observed Mean (SD)	Change from Baseline LS means (95% CI) - MMRM
Baseline	521	7.31 (2.61)	-
Weeks 1-4	513	3.47 (3.35)	-3.84 (-4.10, -3.57)
Weeks 5-8	490	3.13 (3.47)	-4.15 (-4.44, -3.87)
Weeks 9-12	466	2.85 (3.25)	-4.38 (-4.65, -4.11)
Weeks 13-16	440	2.52 (3.14)	-4.69 (-4.95, -4.43)
Weeks 17-20	419	2.33 (3.01)	-4.76 (-5.02, -4.50)
Weeks 21-24	417	2.37 (3.20)	-4.72 (-5.00, -4.43)
Weeks 25-28	398	2.20 (3.21)	-4.85 (-5.14, -4.56)
Weeks 29-32	390	2.12 (3.06)	-4.82 (-5.11, -4.53)
Weeks 33-36	379	2.06 (3.03)	-4.95 (-5.23, -4.67)
Weeks 37-40	375	2.08 (3.08)	-4.97 (-5.26, -4.68)
Weeks 41-44	363	2.09 (3.21)	-4.95 (-5.25, -4.65)
Weeks 45-48	352	1.91 (3.22)	-5.13 (-5.44, -4.83)
Weeks 49 - 52	335	1.84 (3.28)	-5.19 (-5.50, -4.87)

MMRM = Mixed-effects model for repeated measures

[0191] Table 26 provides the change from baseline in monthly headache days over time (mITT population).

5 **Table 26: Change from Baseline in Monthly Headache Days Over Time (mITT Population)**

Derived Visit	N	Atogepant 60 mg QD (N=521)	
		Observed Mean (SD)	Change from Baseline LS means (95% CI) - MMRM
Baseline	521	8.42 (2.95)	-
Weeks 1-4	513	4.40 (3.75)	-4.00 (-4.30, -3.71)
Weeks 5-8	490	3.97 (3.75)	-4.40 (-4.71, -4.10)
Weeks 9-12	466	3.51 (3.52)	-4.80 (-5.09, -4.51)
Weeks 13-16	440	3.19 (3.40)	-5.12 (-5.40, -4.84)
Weeks 17-20	419	2.84 (3.32)	-5.35 (-5.64, -5.06)
Weeks 21-24	417	2.87 (3.47)	-5.30 (-5.61, -4.99)
Weeks 25-28	398	2/74 (3.59)	-5.42 (-5.74, -5.09)
Weeks 29-32	390	2.52 (3.41)	-5.41 (-5.72, -5.09)
Weeks 33-36	379	2.47 (3.31)	-5.62 (-5.92, -5.32)
Weeks 37-40	375	2.45 (3.28)	-5.69 (-6.00, -5.38)
Weeks 41-44	363	2.45 (3.45)	-5.66 (-5.99, -5.34)
Weeks 45-48	352	2.23 (3.43)	-5.90 (-6.22, -5.58)
Weeks 49 - 52	335	2.13 (3.54)	-5.99 (-6.33, -5.66)

MMRM = Mixed-effects model for repeated measures

[0192] Figure 14 shows the LS mean change from baseline in monthly acute medication use days at each monthly period (mITT population), which is also summarized in Table 27.

Table 27: Change from Baseline in Monthly Acute Medication Use Days At Each Monthly Period (mITT Population)

Derived Visit	N	Atogepant 60 mg QD (N=521)	
		Observed Value Mean (SD)	Change from Baseline LS means (95% CI) – MMRM
Baseline	521	6.59 (3.27)	-
Weeks 1-4	513	2.60 (3.00)	-4.04 (-4.28, -3.81)
Weeks 5-8	490	2.49 (3.06)	-4.11 (-4.37, -3.86)
Weeks 9-12	466	2.39 (2.90)	-4.18 (-4.42, -3.94)
Weeks 13-16	440	2.09 (2.58)	-4.46 (-4.68, -4.23)
Weeks 17-20	419	1.92 (2.68)	-4.56 (-4.80, -4.33)
Weeks 21-24	417	2.01 (2.65)	-4.49 (-4.72, -4.25)
Weeks 25-28	398	1.90 (2.78)	-4.59 (-4.84, -4.34)
Weeks 29-32	390	1.84 (2.64)	-4.54 (-4.79, -4.28)
Weeks 33-36	379	1.75 (2.54)	-4.66 (-4.90, -4.41)
Weeks 37-40	375	1.75 (2.52)	-4.68 (-4.93, -4.44)
Weeks 41-44	363	1.69 (2.62)	-4.71 (-4.97, -4.46)
Weeks 45-48	352	1.51 (2.46)	-4.90 (-5.15, -4.66)
Weeks 49 - 52	335	1.50 (2.73)	-4.93 (-5.20, -4.66)

5

[0193] Figure 15A shows the results for $\geq 50\%$ improvement (reduction) in monthly migraine days at each monthly period (mITT population) which is also summarized in Table 28.

Table 28: $\geq 50\%$ Improvement (reduction) in monthly migraine days at each monthly period (mITT population)

Derived Visit	Atogepant 60 mg QD (N=521) Responder n/N1 (%)
Weeks 1-4	310 / 513 (60.4)
Weeks 5-8	319 / 490 (65.1)
Weeks 9-12	327 / 466 (70.2)
Weeks 13-16	326 / 440 (74.1)
Weeks 17-20	308 / 419 (73.5)
Weeks 21-24	314 / 417 (75.3)
Weeks 25-28	299 / 398 (75.1)
Weeks 29-32	299 / 390 (76.7)
Weeks 33-36	305 / 379 (80.5)
Weeks 37-40	293 / 375 (78.1)
Weeks 41-44	285 / 363 (78.5)
Weeks 45-48	290 / 352 (82.4)
Weeks 49 - 52	282 / 335 (84.2)

[0194] The proportion of responders with $\geq 50\%$, $\geq 75\%$, and 100% reduction in MMDs (mITT population, observed cases), are shown in Figure 15B. Of participants who remained in the trial at weeks 49-52, 84.2% (282/335) experienced $\geq 50\%$ reduction in MMDs, 69.9% (234/335) experienced $\geq 75\%$ reduction in MMDs, and 48.4% (162/335) experienced a 100% reduction in MMDs. Proportions of responders in each category increased over the course of the trial.

[0195] As shown above, results demonstrate that atogepant is well-tolerated in this 52-week open label study. ICH E1 exposure requirements for 6-month exposure (≥ 300 patients) and one-year exposure (≥ 100 patients) are met. The efficacy of atogepant treatment in terms of reduction in monthly migraine days, reduction in monthly headache days, reduction in acute medication use days, and $\geq 50\%$ responder rate are well-maintained for the one-year atogepant 60 mg QD treatment. A slight improvement over time was observed for the four endpoints.

[0196] The change from baseline in AIM-D PDA and PI domains for the first and last timepoints assessed are shown in Table 29. The negative change scores with confidence intervals that exclude 0 indicate a significant reduction in impairment due to migraine in PDA and PI.

Table 29: LS mean changes from baseline in AIM-D Performance of Daily Activities (PDA) and Physical Impairment (PI) Domain scores (mITT population, MMRM analysis)

		AIM-D PDA Domain		AIM-D PI Domain	
		Baseline LS mean (SE) = 15.0 (8.80); n = 445		Baseline LS Mean (SE) = 11.4 (8.85); n=445	
Weeks	N	LS Mean (SE) change from baseline	95% CI	LS Mean (SE) Change from baseline	95% CI
1-4	397	-7.61 (0.329)	(-8.86, -6.96)	-5.56 (0.290)	(-6.13, -4.99)
49-52	247	-10.17 (0.370)	(-10.90, -9.44)	-7.20 (0.357)	(-7.91, -6.50)

[0197] The change from baseline in AIM-D monthly PDA domain score (mITT population) is shown in Figure 16A and Table 30. The change from baseline in AIM-D monthly PI domain score (mITT population) is shown in Figure 16B and Table 31. The LS mean change from baseline in the AIM-D PDA and PI scores demonstrated an early and consistent reduction in impairment due to migraine over the 52 week trial. Change scores with confidence intervals excluding zero were present at each time point assessed, indicating significant reduction in impairment due to migraine during the 52-week trial.

Table 30: Change from Baseline in AIM-D Monthly PDA Domain Score: Reduced Impairment in Performance of Daily Activities (mITT Population)

Weeks	n	LS Mean Δ from baseline	95% CI
1-4	397	-7.61	-8.26, -6.96
5-8	360	-8.16	-8.89, -7.42
9-12	346	-8.53	-9.22, -7.84
13-16	334	-8.66	-9.37, -7.94
17-20	323	-9.02	-9.72, -8.32
21-24	313	-9.11	-9.80, -8.43
25-28	301	-9.16	-9.88, -8.44
29-32	288	-9.22	-9.93, -8.51
33-36	288	-9.61	-10.28, -8.94
37-40	283	-9.61	-10.27, -8.95
41-44	273	-9.77	-10.48, -9.06
45-48	262	-9.79	-10.48, -9.10
49-52	247	-10.17	-10.90, -9.44

Table 31: Change from Baseline in AIM-D Monthly PI Domain Score: Reduced Physical Impairment (mITT Population)

Weeks	n	LS Mean Δ from baseline	95% CI
1-4	397	-5.56	-6.13, -4.99
5-8	360	-5.82	-6.50, -5.13
9-12	346	-6.13	-6.75, -5.51
13-16	334	-6.22	-6.87, -5.56
17-20	323	-6.38	-7.04, -5.73
21-24	313	-6.43	-7.08, -5.77
25-28	301	-6.40	-7.09, -5.72
29-32	288	-6.51	-7.18, -5.84
33-36	288	-6.94	-7.57, -6.31
37-40	283	-6.93	-7.56, -6.30
41-44	273	-7.01	-7.68, -6.34
45-48	262	-7.02	-7.68, -6.36
49-52	247	-7.20	-7.91, -6.50

[0198] The Migraine-Specific Quality of Life Questionnaire version 2.1 (MSQv2.1) Role

- 5 Function-Restrictive (RFR) domain least squares (LS) mean (standard error [SE]) at baseline was 48.1 (20.26) and LS mean changes from baseline throughout the trial are shown in Figure 17 and in Table 32. Findings indicated rapid improvement over the first 12 weeks and sustained improvement from 12 to 52 weeks, and non-zero confidence intervals throughout indicate significant improvement at each timepoint assessed.

Table 32: Change from Baseline in MSQ v2.1 Domain Score (mITT Population)

Weeks	n	LS Mean Δ from baseline	95% CI
12	495	30.02	28.16, 31.87
24	436	30.79	28.83, 32.76
36	404	32.34	30.35, 34.34
48	377	34.32	32.38, 36.25
52	363	34.70	32.74, 36.66

[0199] The mean (SE) HIT-6 total score at baseline was 64.07 (4.89) and the LS mean change (95% confidence interval) at Week 4 was -7.63 (-8.31, -6.95) and at Week 52 was -12.10 (-12.98, -11.22), demonstrating reductions in headache impact. The proportion of HIT-6 responders (≥ 5 points from baseline) were 59.92% and 80.77% of participants in the first (week 4) and last (week 52) timepoints assessed, respectively. Change from baseline in HIT-6 Total Score (mITT Population) is shown in Figure 18 and Table 33. HIT-6 total score responder rates (mITT) population are shown in Table 33. A responder on the HIT-6 was defined as a participant with a ≥ 5 point improvement from baseline.

Table 33: Change from Baseline in HIT-6 Total Score (mITT Population)

Week	n	LS Mean Δ from baseline	95% CI
4	514	-7.63	-8.31, -6.95
8	491	-9.14	-9.87, -8.42
12	471	-6.60	-10.33, -8.87
16	448	-10.15	-10.93, -9.37
20	442	-10.56	-11.35, -9.76
24	423	-10.55	-11.39, -9.71
28	418	-10.99	-11.83, -10.15
32	407	-11.15	-11.98, -10.32
36	402	-11.29	-12.14, -10.43
40	394	-11.76	-12.60, -10.91
44	384	-11.46	-12.32, -10.60
48	377	-11.86	-12.72, -10.99
52	364	-12.10	-12.98, -11.22

Table 34: HIT-6 Total Score Responder Rates

Week	Responders	Nonresponders	Total (N)	% Responder
4	308	206	514	59.92
8	327	164	491	66.60
12	326	145	471	69.21
16	321	127	448	71.65
20	327	115	442	73.98
24	313	110	423	74.00
28	320	98	418	76.56
32	317	90	407	77.89
36	304	98	402	75.62
40	318	76	394	80.71
44	301	83	384	78.39
48	302	75	377	80.11
52	294	70	364	80.77

Total (N) = Number of participants with non-missing values at the post-baseline analysis visit.

[0200] Long-term daily use of atogepant 60 mg for the preventive treatment of migraine was associated with reductions in the impact of migraine on the AIM-D in performance of daily

5 activities and physical impairment, improvements in migraine-specific quality of life on the RFR domain, and reductions in the impact of headaches as assessed by changes from baseline, which were significant as indicated by non-zero confidence intervals. Improvements were observed at the earliest time point assessed and increased over the 52-week trial.

[0201] Example 3

10 [0202] As discussed in Example 1, a phase 3 trial (Study A) demonstrated that atogepant dosed once daily results in a clinically meaningful reduction in mean monthly migraine days. An open-label extension study for trial completers evaluated the long-term safety and tolerability of oral atogepant 60 mg daily for the prevention of migraine in participants with episodic migraine.

15 [0203] Participants in this trial (Study C) rolled over from the lead in trial (Study A) and were treated with atogepant 60 mg once daily for 40-weeks, with a 4-week safety follow-up period. Only safety data were collected.

[0204] Of 695 participants screened, a total of 685 participants took at least one dose of study medication and were included in the safety population. Overall, the mean age was 41.8 years. Female participants accounted for 88.2% of the Safety Population. White and Black or African American patients accounted for 84.4% and 12.6%, respectively. Mean BMI was 30.58 kg/m².

5 Demographics data for the Safety Population are summarized in Table 35.

Table 35: Demographics and Baseline Characteristics (Safety Population)

Parameter	Atogepant 60 mg QD N=685
Age (years)	
Mean (SD)	41.8 years (12.30)
Sex, n (%)	
Male	81 (11.8%)
Female	604 (88.2)
Race Group, n (%)	
White	578 (84.4)
Black or African American	86 (12.6)
Asian	9 (1.3)
American Indian or Alaska Native	3 (0.4)
Native Hawaiian or Other Pacific Islander	0
Multiple [1]	8 (1.2)
Missing	1 (0.1)
Race Group, n (%)	
White	578 (84.4)
All other races	106 (15.5)
Missing	1 (0.1)
Ethnicity, n (%)	
Hispanic or Latino	54 (7.9)
Not Hispanic or Latino	631 (92.1)
Weight (kg)	
Mean (SD)	84.21 (23.267)

N	685
BMI (kg / m ²)	
Mean (SD)	30.58 (7.820)
n	685

[1] Participants who reported multiple races are only included in the “Multiple” category.

N = number of patients in the safety population

n=Number of participants within a specific category

Percentages are calculated as $100 \times (n / N)$

- 5 [0205] The majority of participants (74.6%) completed the open label treatment period. Overall, 62.5% of participants experienced a treatment-emergent adverse event (TEAE), with 8.8% considered treatment-related by the investigator; serious adverse events occurred in 3.4% of participants, none of which were treatment related. Table 36 reports the most frequent AEs leading to discontinuation; Table 37 reports the most frequent TEAEs observed. No deaths and
- 10 no hepatic safety issues were observed.

Table 36: Overall Summary of Adverse Events – Safety Population

Category	Total (N=685) n/N (%)
Treatment-emergent adverse event (TEAE)	428 (62.5)
Deaths	0 (0.0)
Treatment-emergent serious adverse events (SAE)	23 (3.4)
Treatment-related TEAE	60 (8.8)
TEAE Leading to Study Discontinuation [1, 2]	22 (3.2)
Nausea	3 (0.4)
Abdominal pain	2 (0.3)
Vomiting	2 (0.3)
Weight Decreased	2 (0.3)
Dizziness	2 (0.3)
Migraine	2 (0.3)

Participants are counted only once within each category.

N = number of participants in the Safety Population

n=number of participants within a specific category

Percentages are calculated as $100 \times (n/N)$

[1] The inset AEs represent the most frequent AEs leading to discontinuation in the extension study.

[2] Three patients with AE onset date in the lead-in trial rolled over to the extension study and discontinued in the extension study.

Table 37: Most Frequent ($\geq 2\%$) Treatment Emergent Adverse Events (TEAEs) by Preferred Term – Safety Population

TEAEs by Preferred Term	Atogepant 60 mg QD (N=685) n (%)
Upper respiratory tract infection	38 (5.5)
Urinary tract infection	36 (5.3)
Nasopharyngitis	33 (4.8)
Sinusitis	25 (3.6)
Constipation	23 (3.4)
Nausea	23 (3.4)
Weight decreased	18 (2.6)
Back pain	17 (2.5)
Dizziness	17 (2.5)
Influenza	17 (2.5)
Arthralgia	15 (2.2)
Corona virus infection	15 (2.2)
Gastroenteritis	15 (2.2)
Anxiety	14 (2.0)
Bronchitis	14 (2.0)
Vomiting	14 (2.0)

Participants are counted only once within each preferred term.

N = number of participants in the Safety Population

n = number of participants within a specific category

Percentages are calculated as $100 \times (n/N)$.

[0206] The study supported that atogepant 60 mg once daily is safe and well tolerated. The study results contributed 538 participants exposed to atogepant for 6 months and 509 participants exposed to atogepant for 9 months.

[0207] Example 4

[0208] The populations of Study A (described in Example 1) and Study B (described in Example 2) were analyzed for weight-related changes over the course of treatment.

[0209] Data was also evaluated from Study C (described in Example 3) and Study D, a Phase 2b/3 clinical trial evaluating the efficacy, safety, and tolerability of orally administered atogepant. In Study D, after a 28 day baseline period, a total of 652 patients were randomized 1:2:2:2 to receive either atogepant 10 mg (n=94), atogepant 30 mg (n = 185), atogepant 60 mg (n = 187) or placebo (n=186).

[0210] In the atogepant Study A population, 44.8% of patients had a BMI of $\geq 30 \text{ kg} / \text{m}^2$, and 58.6% of patients had a BMI $\geq 27 \text{ kg} / \text{m}^2$ with obesity-related comorbidities (e.g., Type 2 diabetes, hypertension, dyslipidemia, sleep apnea, cardiovascular disease). In the atogepant Study B population, 45.3% of patients had BMI $\geq 30 \text{ kg} / \text{m}^2$, and 54.9% of patients had BMI $\geq 27 \text{ kg} / \text{m}^2$ and obesity-related comorbidities.

[0211] Statistically significant weight loss was observed in the atogepant 30 mg once daily and 60 mg once daily groups compared with placebo in Study A. The LS mean difference for percentage change from baseline in body weight at the end of the treatment period was 0.98% (p=0.0005) in the atogepant 30 mg once daily group and 1.64% (p<0.0001) in the atogepant 60 mg once daily group. CDF plots for percentage change from baseline in body weight (kg) at the end of the double-blind treatment period (week 12) for Study A (safety population) are shown in Figure 19.

[0212] Treatment with atogepant over the 52-week treatment period was associated with modest weight loss in Study B. Statistically significant weight loss was observed in the atogepant 60 mg once daily group compared with SOC. The LS mean difference for percentage change from baseline in body weight at the end of the treatment period versus SOC was -1.76% (p=0.0003).

In addition, treatment with atogepant was associated with weight loss as measured by the percentage of participants who lost at least 5% of baseline body weight at the end of the

treatment period. In Study B, the percentage of participants in the atogepant 60 mg once daily group who met this weight loss threshold was 22.4% compared with 14.2% in the SOC group. CDF plots for percentage change from baseline in body weight (kg) at the end of the open-label treatment period (week 52) for Study B (safety population) are shown in Figure 20.

- 5 [0213] When evaluated with both study A and study CGP-MD-01 (NCT02848326), patients had a mean body weight of 83.6 kg and a mean BMI of 30.34 kg / m². In these studies, a dose-dependent mean change in body weight was observed at the end of the double-blind treatment period as follows: +0.32 kg for placebo, +0.07 kg for atogepant 10 mg, -0.40 kg for atogepant 30 mg, and -0.81 kg for atogepant 60 mg. The proportion of patients with weight decrease $\geq 7\%$ at
- 10 any point was 2.8% for placebo, 3.8% for atogepant 10 mg, 3.2% for atogepant 30 mg, and 4.9% for atogepant 60 mg. No patients in either study discontinued atogepant due to decreased weight.

Table 38: Weight Loss (Study A and Study D)

	Placebo N=408	Atogepant 10 mg N=314	Atogepant 30 mg N=411	Atogepant 60 mg (N=417)
Mean Change in weight by end of study (kg)	+0.40	+0.07	-0.40	-0.81
PCS Weight Loss > 7% baseline	11 / 399 (2.8%)	12 / 312 (3.8%)	13 / 404 (3.2%)	20 / 409 (4.9%)

Table 39: Weight Loss Adverse Events (CGP-MD-01 and Study A)

TEAE	Placebo N=408	Atogepant 10 mg N=314	Atogepant 30 mg N=411	Atogepant 60 mg (N=417)	All Atogepant (N=1142)
Weight decreased	3 (0.7%)	1 (0.3%)	0	4 (1.0%)	5 (0.4%)

[0214] In the 52-week open label long-term safety Study B, patients had a mean weight of 83.9 kg and a mean BMI of 30.55 kg / m². In this study, a mean change in body weight was also observed at the end of treatment period as follows: +0.20 kg for oral preventive standard of care group and -1.42 kg for the atogepant 60 mg group. The proportion of patients with weight decrease $\geq$ 7% at any point was 14.7% for the oral migraine preventive standard of care group and 24.1% for the atogepant 60 mg group. One patient (0.1%) in the long-term safety study discontinued atogepant 60 mg due to decreased weight.

Table 40: Weight Loss Adverse Events (Study B)

TEAE	SOC N=196	Atogepant 60 mg (N=1198)
Weight decreased	3 (1.5%)	25 (2.1%)

[0215] Table 41 provides a subset analysis for percentage change from baseline in body weight (kg) at the end of the double-blind treatment period (week 12 (LOCF) ANCOVA) for Study A (safety population). Table 42 provides a subset analysis for 5% responders in body weight (kg) at the end of the double-blind treatment period [week 12 (LOCF) ANCOVA] for Study A (that is, Table 42 sets out a subset analysis for participants who lost at least 5% of baseline body weight at the end of the treatment period). The “Guidance Population” refers to patients with BMIs greater than or equal to 30 kg/m² or BMI greater than or equal to 27 kg/m² in the presence of obesity-related comorbidities (e.g., type 2 diabetes, hypertension, dyslipidemia, sleep apnea, cardiovascular disease).

Table 41: Subset Analysis for Percentage Change from Baseline in Body Weight (kg) at the end of the double-blind treatment period [week 12 (LOCF) ANCOVA] Study A, Safety Population

Subgroup	Statistics	Placebo	Atogepant 10 mg	Atogepant 30 mg	Atogepant 60 mg
Overall N=902	N	222	221	228	231
	Baseline weight	84.61	83.6	85.69	82.52
	LS Mean (SE)	0.37 (0.20)	0.14 (0.20)	-0.61 (0.20)	-1.27 (0.20)
	LSMD vs. Placebo (95% CI)		-0.23 (-0.79, 0.32)	-0.98 (-1.53, -0.43)	-1.54 (-2.19, -1.09)
	p value		0.4138	0.0005	<.0001
BMI $\geq$ 30 (S.E) N=404	N	97	98	119	90
	Baseline weight	104.07	102.06	101.56	102.68
	LS Mean (SE)	0.52 (0.29)	0.14 (0.29)	-0.36 (0.26)	-1.29 (0.30)
	LSMD vs. Placebo (95% CI)		-0.39 (-1.17, 0.40)	-0.88 (-1.63, -0.13)	-1.81 (-2.62, -1.01)
	p value		0.3345	0.021	<.0001
Guidance Population (S.E) N=529	N	126	127	148	128
	Baseline weight	98.15	96.83	97.02	95.18
	LS Mean (SE)	0.36 (0.27)	0.13 (0.27)	-0.63 (0.25)	-1.40 (0.27)
	LSMD vs. Placebo (95% CI)		-0.23 (-0.96, 0.49)	-1.00 (-1.69, -0.30)	-1.76 (-2.49, -1.04)

Table 42: Subset Analysis for 5% Responders in Body Weight (kg) at the end of the double-blind treatment period [Week 12 (LOCF) ANCOVA] – Study A

Subgroup and Statistics	Placebo	Atogepant 10 mg	Atogepant 30 mg	Atogepant 60 mg
Overall (N)	222	221	228	231
Responder, n/N1	9/220 (5.0)	11/220 (5.0)	9/225 (4.0)	32/228 (14.0)
Odds Ratio vs. Placebo (95% CI)		1.22 (0.50, 3.02)	1.00 (0.39, 2.56)	3.78 (1.76, 8.13)
Nominal p-value		0.6613	0.9937	0.0007
BMI $\geq$ 30 (N)	97	98	119	90
S.E				
Responder, n/N1	5/96 (5.2)	3/97 (3.1)	1/117 (0.9)	8.88 (9.1)
Odds Ratio vs. Placebo (95% CI)		0.57 (0.13, 2.47)	0.15 (0.02, 1.34)	1.82 (0.57, 5.82)
Nominal p-value		0.4545	0.0898	0.312
Guidance Population (N)	126	127	148	128
S.E				
Responder, n/N1	7/125 (5.6)	5/126 (4.0)	4/146 (2.7)	17/126 (13.5)
Odds Ratio vs. Placebo (95% CI)		0.69 (0.21, 2.26)	0.47 (0.13, 1.66)	2.55 (1.01, 6.44)
Nominal p-value		0.542	0.2429	0.0479

[0216] Table 43 provides a subset analysis for percentage change from baseline in body weight (kg) at the end of the open label treatment period [week 52 (LOCF) Logistic Regression Approach] for Study B. Table 44 presents data regarding participants who lost at least 5% of

baseline body weight (kg) at the end of the open-label treatment period [Week 52 (LOCF) ANCOVA] for Study B (safety population).

Table 43: Subset Analysis for Percentage Change from Baseline in Body Weight (kg) at the end of the Open-Label Treatment Period [Week 52 (LOCF) Logistic Regression Approach]

5 – Study B Safety Population

Subgroup	Statistics	SOC	Atogepant 60 mg QD
<u>Overall</u> N=739	N	196	543
	Baseline weight	83.94	84.2
	LS Mean (SE)	0.19 (0.423)	-1.57 (0.249)
	LSMD vs. Placebo (95% CI)		-1.76 (-2.72, -0.80)
	p value		0.0003
<u>BMI >=30</u> N=335	N	85	250
	Baseline weight	102.99	101.1
	LS Mean (SE)	0.38 (0.60)	-2.27 (0.35)
	LSMD vs. Placebo (95% CI)		-2.65 (-4.02, -1.27)
	p value		0.0002
<u>Guidance Population</u> (S.E) N=406	N	99	307
	Baseline weight	99.77	97.42
	LS Mean (SE)	0.40 (0.55)	-2.09 (0.31)
	LSMD vs. Placebo (95% CI)		-2.49 (-3.73, -1.26)
	p value		<.0001

Table 44: Participants who lost at least 5% of baseline body weight (kg) at the end of the open label treatment period [week 52 (LOCF) ANCOVA] – Study B Safety Population

Statistics	SOC	Atogepant 60 mg QD
<u>Overall</u>		
Responder, n/N1 (%)	27/190 (14.2)	119/531 (22.4)
Odds Ratio vs. Placebo (95% CI)		1.75 (1.11, 2.75)
Nominal p-value	-	0.0165
<u>BMI >=30</u>		
Responder, n/N1 (%)	9/84 (10.7)	58/247 (23.5)
Odds Ratio vs. Placebo (95% CI)		2.56 (1.20, 5.42)
Nominal p-value	--	0.0145
<u>Guidance Population</u>		
Responder, n/N1	9/98 (9.2)	70/302 (23.2)
Odds Ratio vs. Placebo (95% CI)		2.98 (1.43, 6.23)
Nominal p-value	--	0.0036

[0217] As shown above, a clear dose-dependent weight reduction was observed for percentage change from baseline in body weight for Study A based on ANCOVA analysis. For Study A, the responder rate ($\geq 5\%$ weight loss) at week 12 was higher in the atogepant 60 mg group compared to atogepant 10 mg and 30 mg. For Study B, a clear difference in the percentages of patients who lost at least 5% of baseline body weight was observed in the atogepant 60 mg daily group (22.4%) compared to the SOC group (14.2%); it should be noted that patients in the SOC group were heterogeneous, and may have taken medications associated with weight loss, weight gain, or neither.

[0218] In Study A, baseline BMI did not affect the % weight change; the placebo-adjusted difference was consistent across BMI categories. In Study B, a slightly larger weight reduction was observed in the special populations (BMI ≥ 30) based on ANCOVA analysis.

[0219] The time course of weight loss associated with atogepant use was also evaluated in the Study B population. Figures 21-24 show CDF plots at 3 months (Figure 21), 6 months (Figure 22), 9 months (Figure 23), and 12 months (Figure 24) in the safety population based on observed data. The change from baseline in weight over time for the Study B safety population is summarized in Table 45. The percentage change from baseline in weight over time for Study B (Safety Population) is shown in Table 46.

Table 45: Change from baseline in weight over time – Study B Safety Population

Week	Atogepant 60 mg		SOC	
	n	Mean (SD)	n	Mean (SD)
4	529	-0.39 (1.79)	189	-0.01 (1.61)
8	498	-0.72 (2.29)	180	-0.06 (2.14)
12	480	-1.04 (2.76)	175	0.07 (2.71)
16	456	-1.27 (3.24)	172	0.09 (3.33)
20	446	-1.62 (3.81)	166	0.12 (3.93)
24	243	-1.82 (4.10)	157	-0.05 (4.23)
28	426	-1.79 (4.41)	150	0.13 (4.78)
32	412	-2.00 (4.79)	147	0.11 (5.00)
36	406	-2.01 (4.82)	142	0.12 (5.48)
40	398	-2.11 (5.62)	140	0.03 (5.54)
44	388	-1.84 (5.22)	141	0.12 (5.83)
48	382	-1.70 (5.28)	139	0.43 (5.99)
52	366	-1.54 (5.33)	131	0.24 (6.04)
End of the OL Period	531	-1.42 (4.87)	190	0.20 (5.33)

Table 46: Percentage change from baseline in weight over time – Study B, Safety Population

Week	Atogepant 60 mg		SOC	
	n	Mean (SD)	n	Mean (SD)
4	529	-0.42 (2.11)	189	-0.04 (1.95)
8	498	-0.82 (2.65)	180	-0.13 (2.57)
12	480	-1.18 (3.16)	175	0.08 (3.22)
16	456	-1.48 (3.75)	172	0.13 (3.83)
20	446	-1.86 (4.36)	166	0.15 (4.49)
24	243	-2.08 (4.78)	157	-0.07 (4.73)
28	426	-2.02 (5.14)	150	0.10 (5.14)
32	412	-2.24 (5.45)	147	0.10 (5.49)
36	406	-2.26 (5.60)	142	0.14 (6.05)
40	398	-2.38 (6.70)	140	0.00 (6.02)
44	388	-2.04 (6.14)	141	0.13 (6.28)
48	382	-1.85 (6.23)	139	0.53 (6.49)
52	366	-1.71 (6.39)	131	0.33 (6.61)
End of the OL Period	531	-1.57 (5.75)	190	0.19 (5.83)

[0220] As discussed above, the SOC group was heterogeneous, and encompassed multiple potential options for standard of care medications. A comparison of atogepant 60 mg to those patients who initially took topiramate is shown in Table 47.

Table 47: Percentage Change from Baseline in Weight Over Time, Study B, Safety Population

Week	Atogepant 60 mg		Topiramate Initial Use	
	n	Mean (SD)	n	Mean (SD)
4	529	-0.42 (2.11)	57	-0.57 (1.74)
8	498	-0.82 (2.65)	53	-0.93 (3.06)
12	480	-1.18 (3.16)	48	-1.17 (3.76)
16	456	-1.48 (3.75)	47	-0.57 (5.32)
20	446	-1.86 (4.36)	45	-0.94 (6.55)
24	243	-2.08 (4.78)	43	-1.53 (7.06)
28	426	-2.02 (5.14)	40	-1.35 (8.17)
32	412	-2.24 (5.45)	38	-1.31 (8.79)
36	406	-2.26 (5.60)	38	-1.41 (9.46)
40	398	-2.38 (6.70)	36	-1.57 (9.69)
44	388	-2.04 (6.14)	35	-1.80 (9.67)
48	382	-1.85 (6.23)	35	-1.51 (9.58)
52	366	-1.71 (6.39)	30	-1.90 (10.03)
End of the OL Period	531	-1.57 (5.75)	58	-1.14 (7.76)

[0221] Table 48 provides a summary of final total daily dose among initial treatment in Study B.

5 The most common daily dose for topiramate in Study B was 25 mg.

Table 48: Summary of final total daily dose among initial treatment

	10 mg	12.5 mg	20 mg	25 mg	50 mg	100 mg	Total
Amitriptyline	19	1	1	21	1	1	43
Topiramate	-	2	-	33	29	6	70

[0222] As shown above, in the Study B long term safety study, atogepant led to a gradual weight reduction from week 4 to week 40 (2.11 kg reduction at week 40); the weight reduction then slightly decreased to -1.54 kg at week 52, and -1.42 at the end of the OL period. At the end of the

10

OL period, the percentage reduction of weight was -1.57 for atogepant 60 mg, and -1.14 for topiramate.

CLAIMS

1. A method of prophylactically treating migraine, the method comprising administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD.
- 5 2. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean monthly migraine days of at least 3.6 days.
3. The method of claim 2, wherein the prophylactic treatment results in a reduction in mean monthly migraine days of at least 3.8 days.
4. The method of claim 3, wherein the prophylactic treatment results in a reduction in mean
10 monthly migraine days of at least 4.2 days.
5. The method of claim 1, wherein the prophylactic treatment results in at least 50% reduction in monthly migraine days.
6. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean monthly headache days of at least 3.9 days.
- 15 7. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean monthly headache days of at least 4 days.
8. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean monthly headache days of at least 4.2 days.
9. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean
20 monthly acute medication use days of at least 3.6 days.
10. The method of claim 1, wherein the prophylactic treatment results in a reduction in mean monthly acute medication use days of at least 3.8 days.
11. A method of preventing migraine in a patient having fewer than fifteen migraine days per month, the method comprising administering atogepant in an amount of 10 mg QD, 30 mg QD,
25 or 60 mg QD, wherein administering atogepant results in a reduction in mean monthly migraine days of at least about 3.6 days.
12. The method according to claim 11, wherein administering atogepant results in a reduction in mean monthly migraine days of at least about 3.8 days.
13. The method according to claim 11, wherein administering atogepant results in a reduction
30 in mean monthly migraine days of at least 4.2 days.

14. The method according to claim 11, wherein administering atogepant results in a reduction in mean monthly headache days of at least about 4.2 days.

15. The method according to claim 11, wherein administering atogepant results in a reduction in mean monthly acute medication use days of at least 3.8 days.

5 16. A method of preventing migraine, the method comprising administering atogepant in an amount of about 10 mg QD, about 30 mg QD, or about 60 mg QD, wherein administration of atogepant results in an improvement in ability to perform daily activities as compared to a pre-treatment baseline or a patient not receiving atogepant.

10 17. The method according to claim 16, wherein the improvement in ability to perform daily activities is assessed by the MSQ v2.1.

18. The method according to claim 17, wherein administration of atogepant results in a change from baseline in the MSQ v2.1 RFR domain of greater than about 25 points.

15 19. The method according to claim 18, wherein atogepant is administered in an amount of about 10 mg QD, and wherein administration of atogepant results in a change in the MSQ v2.1 RFR domain of about 30.4 points from baseline.

20. The method according to claim 18, wherein atogepant is administered in an amount of about 30 mg QD, and wherein administration of atogepant results in a change in the MSQ v2.1 RFR domain of about 30.5 points relative to baseline.

20 21. The method according to claim 18, wherein atogepant is administered in an amount of about 60 mg QD, and wherein administration of atogepant results in a change in the MSQ v2.1 RFR domain of about 31.3 points relative to baseline.

22. The method according to claim 16, wherein the improvement in ability to perform daily activities is assessed using the Activity Impairment in Migraine Diary (AIM-D).

25 23. The method according to claim 22, wherein the daily activity is selected from the group consisting of household chores, errands, leisure activities at home, social or leisure activities outside the home, strenuous physical activities, concentration, and thinking clearly.

24. The method according to claim 22, wherein the daily activity is selected from the group consisting of walking, moving body, bending forward, or moving head.

30 25. The method according to claim 22, wherein administration of atogepant results in an improvement from baseline in Performance of Daily Activities score of greater than about 6 points.

26. The method according to claim 25, wherein atogepant is administered at about 30 mg QD, and wherein the improvement from baseline in Performance of Daily Activities score is about 8.6 points relative to baseline.

27. The method according to claim 25, wherein atogepant is administered at about 60 mg QD, and wherein the improvement from baseline in the AIM-D Performance of Daily Activities score is about 9.4 points.

28. The method according to claim 22, wherein administration of atogepant results in an improvement from baseline in the AIM-D Physical Impairment Score of more than about 5.2 points.

29. The method according to claim 28, wherein atogepant is administered at about 30 mg QD, and wherein the improvement from baseline in the AIM-D Physical Impairment Score is about 6.0 points.

30. The method according to claim 28, wherein atogepant is administered at about 60 mg QD, and wherein the improvement from baseline in the AIM-D Physical Impairment score is about 6.5 points.

31. The method of claim 11, wherein atogepant is administered for at least about 4 weeks, and wherein administration of atogepant results in a reduction in body weight from baseline of at least about 0.5 kg.

32. A method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month, the method comprising administering atogepant or a pharmaceutically acceptable salt thereof in an amount of 10 mg QD, 30 mg QD, or 60 mg QD for at least about six months.

32. The method of claim 31, wherein atogepant is administered daily for at least about nine months.

33. The method of claim 31, wherein atogepant is administered daily for at least about 12 months.

34. The method of claim 33, wherein administration of atogepant results in a reduction in body weight from baseline of at least about 1.42 kg.

35. The method of claim 33, wherein at least about 22% of patients achieve a weight decrease of greater than about 5% relative to baseline.

36. A method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month, the method comprising administering atogepant in an amount of about 30 mg once daily for at least about 12 weeks, wherein administration of atogepant results in a reduction in body weight of at least about 0.4 kg.

5 37. The method of claim 36, wherein atogepant is administered daily for at least about 52 weeks.

38. The method of claim 36, wherein at least about 3.2% of patients achieve a weight loss of greater than 7% relative to baseline body weight.

10 39. A method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month, the method comprising administering atogepant in an amount of about 60 mg once daily for at least about 12 weeks, wherein administration of atogepant results in a reduction in body weight of at least about 0.8 kg.

40. The method of claim 39, wherein at least about 4.9% of patients achieve a weight loss of greater than 7% relative to baseline body weight.

15 41. A method for the preventive treatment of migraine in patients having fewer than fifteen migraine days per month, the method comprising administering atogepant in an amount of about 60 mg once daily for at least about 52 weeks, wherein administration of atogepant results in a reduction in body weight of at least about 1.4 kg.

20 42. The method of claim 41, wherein at least about 22.4% of patients achieve a weight loss of at least 5% relative to baseline body weight.

1/33




FIGURE 1




FIGURE 2






FIGURE 3



* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ versus placebo


FIGURE 4



*P<0.05; **P<0.01; ***P<0.001.
LS, least-squares; mITT, modified intent-to-treat; SE, standard error.
For weekly data, baseline was defined as monthly migraine days/4, and change from baseline in weekly migraine days was to be calculated for consecutive 7-day periods beginning with day 1.

FIGURE 5A

6/33



* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

mITT, modified intent-to-treat.

^a Day of initial dose excluded, as migraine attacks occurring prior to study drug administration were included.

FIGURE 5B

7/33




FIGURE 6





FIGURE 7

9/33

Proportions of Participants Achieving Various Responder Rates by Treatment Group: 12-Week Average Responder Rates (mITT Population)



* $P < 0.05$ vs placebo; $^†P < 0.01$ vs placebo; $^{††}P < 0.0001$ vs placebo.
mITT, modified intent-to-treat.

FIGURE 8A

Cumulative Distribution Function: Proportions of Participants Achieving Reduction in Mean MMDs Across 12 Weeks (mITT Population)





FIGURE 8B

11/33

Reductions of $\geq 25\%$ in Mean MMDs by Treatment Group and by 4-Week intervals (mITT population)

$\geq 25\%$ Reduction in Mean MMDs



* $P < 0.05$ vs placebo; [†] $P < 0.01$ vs placebo; [‡] $P < 0.0001$ vs placebo.
MMD, monthly migraine day.




FIGURE 8C

Proportion of participants with $\geq 50\%$, $\geq 75\%$, and 100% reduction in migraine days





FIGURE 8D

Mean Change from Baseline in moderate/severe monthly headache days per month (mITT population)

13/33




* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

LS, least square; mITT, modified intent-to-treat; SE, standard error.

FIGURE 9A

14/33

Mean Change from Baseline in Monthly Headache Days Per Month (mITT Population)



* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

LS, least square; mITT, modified intent-to-treat; SE, standard error.

FIGURE 9B





FIGURE 10A

16/33

Mean change from baseline in acute medication use days per month (mITT population)




* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

LS, least square; mITT, modified intent-to-treat; SE, standard error.

FIGURE 10B

17/33

Mean Change from Baseline in Cumulative Headache Hours Per Month (mITT Population)




* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

LS, least square; mITT, modified intent-to-treat; SE, standard error.

Figure 10C

18/33

Migraine-Specific Quality of Life Questionnaire Role Function-Restrictive Domain^a Score: Weeks 4, 8, and 12



BL, baseline mean; LSM, least squares mean; LSMD, least squares mean difference; MSQ, Migraine-Specific Quality of Life Questionnaire; QD, once daily; RFR, Role Function-Restrictive Domain; SE, standard error

^a Role Function-Restrictive domain assesses how migraines limit one's daily social and work-related activities. Participants respond to items using a 6-point scale ranging from "none of the time" to "all of the time." Raw domain scores are rescaled to a 0 to 100 scale, where higher scores indicate better quality of life.

[†]LSMD (95% CI): Week 4: 8.6 (4.4, 12.9); $P<0.0001$; Week 8: 11.3 (7.1, 15.5); $P<0.0001$; Week 12: 9.9 (5.5, 14.4); $P<0.0001$;

[‡]LSMD (95% CI): Week 4: 11.2 (7.0, 15.4); $P<0.0001$; Week 8: 9.9 (5.8, 14.0); $P<0.0001$; Week 12: 10.1 (5.7, 14.5); $P<0.0001$;

[§]LSMD (95% CI): Week 4: 12.3 (8.1, 16.5); $P<0.0001$; Week 8: 10.8 (6.7, 14.9); $P<0.0001$; Week 12: 10.8 (6.4, 15.2); $P<0.0001$.

FIGURE 11





FIGURE 12






FIGURE 14




FIGURE 15A

23/33





FIGURE 15B


Change from Baseline in AIM-D Monthly PDA Domain Score: Reduced Impairment in Performance of Daily Activities (mITT Population)



AIM-D, Activity Impairment in Migraine-Diary; LS, least-squares; mITT, modified intent-to-treat; PDA, Performance of Daily Activities.
n=number of participants with evaluable data at both baseline and a specific time point in the mITT population.


FIGURE 16A

Change from Baseline in AIM-D Monthly PI Domain Score: Reduced Physical Impairment (mITT Population)



AIM-D, Activity Impairment in Migraine-Diary; LS, least-squares; mITT, modified intent-to-treat; PI, Physical Impairment.
n=number of participants with evaluable data at both baseline and a specific time point in the mITT population.


FIGURE 16B



LS mean baseline (SE)=48.1 (20.26); n=516.
LS=least squares, mITT=modified intent-to-treat, MMRM=mixed-effects model for repeated measures, SE=standard error.
n=number of participants with evaluable value at both baseline and a specific time point in the mITT population.

FIGURE 17
Change from Baseline in HIT-6 Total Score (mITT Population)

27/33



HIT-6, Headache Impact Test-6; LS, least-squares; mITT, modified intent-to-treat; SE, standard error.

n=number of participants with evaluable data at both baseline and a specific time point in the mITT population.

FIGURE 18




FIGURE 19




FIGURE 20




FIGURE 21

MONTH 6




FIGURE 22

MONTH 9




FIGURE 23

MONTH 12, COMPLETER'S POPULATION




FIGURE 24

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2021/043791

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/4545; A61K 9/00; A61P 25/06 (2021.01)

CPC - A61K 31/4545; A61K 9/0053; A61P 25/06 (2021.08)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

see Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

see Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

see Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	GOADSBY et al., Orally Administered Atogepant Was Efficacious, Safe, and Tolerable for the Prevention of Migraine: Results From a Phase 2b/3 Study (S17.001), Neurology, Vol. 92, S15, 06 May 2019 [retrieved on 14 October 2021]. Retrieved from the Internet: <URL: https://n.neurology.org/content/92/15_Supplement/S17.001.abstract >. Abstract	1-16, 31, 32a, 32b, 33-42 ----- 17-30
Y	WO 2020/051137 A1 (ELI LILLY AND COMPANY) 12 March 2020 (12.03.2020) entire document	17-21
Y	CALA et al., The Activity Impairment in Migraine-Diary (AIM-D): A Novel Migraine-Specific Patient-Reported Outcome Measure to Assess Functioning based on Activity Impairment in Episodic and Chronic Migraine Patients, MTIS 2018 Abstracts, Vol. 38, Issue 1S, MTIS2018-005, 2018 [retrieved on 14 October 2021], Retrieved from the Internet: <URL: https://journals.sagepub.com/doi/10.1177/0333102418789865 >. Pgs. 4-5	22-30
A	US 2019/0374518 A1 (ALLERGAN PHARMACEUTICALS INTERNATIONAL LIMITED) 12 December 2019 (12.12.2019) entire document	1-31, 32a, 32b, 33-42

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance

“D” document cited by the applicant in the international application

“E” earlier application or patent but published on or after the international filing date

“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

“O” document referring to an oral disclosure, use, exhibition or other means

“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

14 October 2021

Date of mailing of the international search report

NOV 18 2021

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Harry Kim

Telephone No. PCT Helpdesk: 571-272-4300