

**(12) PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 200117017 B2  
(10) Patent No. 757538**

(54) Title  
**Stable, high glycerol liquids comprising sulfosuccinic acid monoesters**

(51)<sup>7</sup> International Patent Classification(s)  
**A61K 007/48 C11D 001/32  
A61K 007/50**

(21) Application No: **200117017** (22) Application Date: **2000.11.13**

(87) WIPO No: **WO01/41729**

(30) Priority Data

(31) Number **9928820** (32) Date **1999.12.06** (33) Country **GB**

(43) Publication Date : **2001.06.18**

(43) Publication Journal Date : **2001.08.30**

(44) Accepted Journal Date : **2003.02.27**

(71) Applicant(s)  
**Unilever PLC**

(72) Inventor(s)  
**Masaaki Arai; Mitsunobu Kobayashi; Yukie Mishina; Naohiko Takada**

(74) Agent/Attorney  
**UNILEVER AUSTRALIA LIMITED, B F Jones, Private Mailbag 2, EPPING NSW 2121**

(56) Related Art  
**US 5130056  
US 5132037  
WO 99/29409**

# AU 200117017

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 June 2001 (14.06.2001)

PCT

(10) International Publication Number  
**WO 01/41729 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 7/48, 7/50, C11D 1/32**

Innovation Center, 38, Hagadai, Haga-machi, Haga-gun, Tochigi, Tochigi 321-3325 (JP). MISHINA, Yukie; 1-22, Kita 6-chome, Nishi 14-jo, Obihiro-shi, Hokkaido 080-0044 (JP). TAKADA, Naohiko; Nippon Lever B.V., Innovation Center, 38, Hagadai, Haga-machi, Haga-gun, Tochigi, Tochigi 321-3325 (JP).

(21) International Application Number: **PCT/EP00/11359**

(74) Agent: ROTHS, Maria, Johanna, Francisca; Unilever PLC, Patent Dept., Colworth House, Sharnbrook, Bedford, Bedfordshire MK44 1LQ (GB).

(22) International Filing Date: **13 November 2000 (13.11.2000)**

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(25) Filing Language: **English**

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(26) Publication Language: **English**

## Published:

(30) Priority Data: **9928820.1 6 December 1999 (06.12.1999) GB**

- *With international search report.*
- *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

(71) Applicant (for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW only): **UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (for all designated States except AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, IN, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW): **UNILEVER NV [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).**

(71) Applicant (for IN only): **HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).**

(72) Inventors: **ARAI, Masaaki; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). KOBAYASHI, Mitsunobu; Nippon Lever B.V.,**

**WO 01/41729 A1**

(54) Title: **STABLE, HIGH GLYCEROL LIQUIDS COMPRISING SULFOSUCCINIC ACID MONOESTERS**

(57) Abstract: The invention discloses high glycerol content (greater than or equal to 10% by wt.) liquid compositions comprising sulfosuccinic acid monoesters wherein the compositions are stable. Stability is provided by a stabilization system comprising a minimum level of a defined salt and a defined thickener. A method of enhancing stability of high content glycerin compositions is also disclosed.

- 1 -

STABLE, HIGH GLYCEROL LIQUIDS COMPRISING SULFOSUCCINIC ACID MONOESTERS

Field of the Invention

5

The present invention relates to high glycerin content liquid compositions comprising one or more defined sulfosuccinic acid monoesters. More specifically, while high glycerin content has been found to destabilize such 10 compositions, applicants have found that a combination of a minimum level of fatty acid salt (e.g., alkali metal salt or alkanolamine salt), and a thickener selected from acrylate copolymers, hydroxy alkyl cellulose, cationic guar gums (e.g., Jaguar C13S) stabilize the compositions. The 15 compositions of the invention may be used in skin cleansing, shower gel, and hair compositions.

Compositions containing sulfosuccinic acid monoesters are disclosed in U.S. Patent No. 4,749,515 to Miyamoto et al..

20

However, because of its moisturizing properties, it would also be greatly beneficial to add glycerin to such compositions to relieve skin dryness. In such compositions, however, it has been found that large amounts of glycerin 25 (e.g., 10% by wt. and above) lead to instability and phase separation. While not wishing to be bound by theory, it is thought that this may be due to the high specific gravity of glycerin.

- 2 -

Unexpectedly, applicants have now found that this instability may be overcome by using a stabilization system comprising:

5 (a) at least 4% by wt. of an alkali metal soap of C<sub>12</sub>-C<sub>24</sub> fatty acid and/or alkanolamine soap; and

(b) 0.01 to 5% by wt. of a thickening copolymer selected from acrylate copolymers, hydroxy alkyl cellulose (e.g., hydroxy ethyl or propyl cellulose), cationic 10 guar gum, and isostearic acid. Acrylate copolymers are particularly preferred.


Brief Summary of the Invention

15



The present invention relates to stable, high glycerin liquid detergent compositions. More particularly, the compositions comprise:

20 (1) 1-25%, preferably 5 to 20% of one or more sulfosuccinic acid monoesters represented by Formula (I), (II), (III) or (IV);


- 3 -



10



or



- 4 -

wherein each of  $M_1$  to  $M_4$  is H,  $NH_4$ , an alkali metal or a hydroxyalkyl substituted ammonium.  $R_1$  and  $R_2$  are each an alkyl or hydroxyalkyl group having about 8 to 20 carbon atoms,  $R_3$  is H or  $CH_3$ ,  $AO$  is an oxyalkylene group having 2 or 3 carbon atoms, and  $n$  represents an integer from 0 to 20;

- (2) 10% and above, preferably 10 to 50% by wt., preferably 15 to 40% by wt. of glycerin;
- 10 (3) at least 4% by wt., preferably at least 5% to 15% by wt. of an alkali metal soap of  $C_{12}$ - $C_{24}$  fatty acid and/or alkanolamine soap;
- 15 (4) 0.01 to 5% by wt., preferably 0.1 to 3% by wt. (levels may vary depending on which thickener is used) of a thickener selected from acrylate copolymer (e.g., alkyl acrylate copolymer), hydroxy alkyl cellulose, cationic guar gum and isostearic acid; and
- 20 (5) balance water.

In a second embodiment, the invention relates to a method of enhancing the stability of high glycerin compositions comprising defined sulfosuccinic acid monoesters by adding a stabilization system comprising components (3) and (4) above.

The pH of the compositions is about 6 to 8.5, preferably 6.5 to 8.5 and viscosity is about 100 to 5000 centistokes

- 5 -

preferably 100 to 200 measured using a Brookfield Viscometer, Spindle No.2 or 3, 12 rpm, for about 30 seconds at 25°C.

5 The stability of the composition is defined by the absence of separation after 4 weeks storage at room temperature.


Detailed Description of the Invention

10 The present invention relates to liquid compositions comprising one or more defined sulfosuccinic acid monoesters and relatively high amounts of glycerin while retaining stability. More specifically, by using a combination of fatty acid salts and specific thickeners (preferably 15 acrylate copolymers and isostearic acid), it has been found that stable, high content glycerin (greater than or equal to 10% by wt. to about 50% by wt. glycerin), sulfosuccinic monoester compositions may be formed.


20 Compositions of the present invention are set forth in greater detail below.

The compositions of the present invention comprise 1-25%, preferably 5-20% by wt. of one or more sulfosuccinic acid 25 monoesters represented by Formula I, II, III or IV and mixtures thereof as follows:


- 6 -




5



10



or



(IV)

- 7 -

In these formulae, each of  $M_1$  to  $M_4$  may be selected from H,  $NH_4$ , an alkali metal and hydroxyalkyl-substituted ammonium groups.  $M_1$  and  $M_2$ , and  $M_3$  and  $M_4$  may be the same or 5 different. Alkali metals may include lithium, sodium, potassium and the like. The hydroxyalkyl substituted ammonium groups, which may preferably have 1 to 3 carbon atoms in the hydroxyalkyl group, may include monoethanol ammonium, diethanol ammonium, triethanol ammonium, methyl 10 diethanol ammonium and the like. Preferably,  $M_1$  to  $M_4$  are hydrogen, sodium or triethanol ammonium.

$R_1$  and  $R_2$  are each straight or branched alkyl or hydroxyalkyl groups having about 8 to 20 carbon atoms, for 15 example, hexyl, decyl, hydroxydecyl, dodecyl, hydroxytetradecyl, tetradecyl, nonadecyl, or the like. Any alkyl group having less than 8 or more than 20 carbon atoms may optionally be contained in the molecule, provided that the total number of carbon atoms in the sulfosuccinic acid 20 monoester contained in the detergent composition of the present invention is in the range from 8 to 20. This number of carbon atoms provides good foaming properties, whereas when the number of carbon atoms is either less than 8 or more than 20 less foaming is observed.

25

In the formulae (II) and (IV),  $R_3$  is either H or  $CH_3$ .

In the formulae,  $AO$  represents an oxyalkylene group having 2 or 3 carbon atoms, that is, oxyethylene or oxypropylene

- 8 -

group. Both oxyethylene and oxypropylene groups may be present in the molecule.

The letter n represents an integer from 0 to 20, preferably 5 1 to 10. If n is more than 20, the foaming properties of the resulting liquid detergent compositions may be poor.

The compounds represented by the formulae (I), (II), (III) and (IV) may be prepared by any known methods.

10

For example, the sulfosuccinic acid monoesters represented by the general formulae (I) or (II) may be prepared by reacting an alkylene oxide adduct of a higher fatty acid with maleic anhydride to produce an ester of maleic 15 anhydride and further reacting the ester of maleic anhydride with a sulfite.

Alternatively, the sulfosuccinic acid monoesters represented by the general formulae (II) or (IV) may be prepared by 20 reacting a lower alcohol ester of a higher fatty acid with an alkanolamine, adding an alkylene oxide to the reaction product, further reacting the resulting addition product with maleic anhydride to produce an ester of maleic anhydride, and reacting the ester of maleic anhydride with a 25 sulfite.

As part of the required stability system of the invention, compositions of the invention also require at least 4%, preferably at least 5% to about 15% by wt. of an alkali 30 metal soap of a C<sub>12</sub>-C<sub>14</sub> fatty and/or alkanolamine salt.

- 9 -

Suitable fatty acid soaps include, but are not limited to, sodium or potassium salts of lauric or myristic acid. Suitable alkanolamine salts include but are not limited to ethanolamine, triethanolamine etc.

5

A second component of the stability system is a thickener. The thickener is selected from acrylate copolymers, hydroxy alkyl celluloses, cationic guar gums, and isostearic acid.

10 Among the acrylates which may be used are included alkyl acrylate copolymer e.g. Leoarl MS200 ex Lion and Acuryn 33 ex Rohm and Haas; Acryl copolymer e.g. Auryn 22 ex Rohm and Haas. Hydroxy alkyl celluloses include hydroxy ethyl, hydroxy propyl or hydroxy methyl cellulose. Cationic guar  
15 gums include Jaguar gums.

The thickener may be used in amounts between 0.01 to 5% by wt. although the amounts may depend on which thickener is used. For example, acrylate copolymer may comprise 0.1-3%  
20 by wt. while isostearic acid may comprise from 1-5% by wt of the composition.

The balance of the composition (generally about 20-70%, preferably 25-65% by wt.) is water.

- 10 -

Optional Components

Compositions of the invention may contain 0.1 to 10% by wt. of a surfactant selected from anionic, nonionic, amphoteric 5 and cationic surfactants and mixtures thereof.

Suitable anionic detergents include, but are not limited, to alkyl and alkylene carboxylates having 10 to 20 carbon atoms, alkyl ether carboxylates, fatty alcohol sulfates, 10 fatty alcohol ether sulfates, alkylol amido-sulfates and sulfonates, fatty acid alkylol amido-polyglycol ether sulfates, alkane sulfonates and hydroxyalkane sulfonates, olefin sulfonates, acyl esters of isethionates,  $\alpha$ -sulfo fatty acid esters, alkyl-benzene sulfonates, alkylphenol 15 glycol ether sulfonates, sulfo-succinates, sulfosuccinic acid half and di-esters, fatty alcohol ether phosphates, protein fatty acid condensation products, alkyl monoglyceride sulfates and sulfonates, alkyl glyceride ether sulfonates, fatty acid methyl taurides, fatty acid 20 sarcosinates, sulfo-ricinoleates. These compounds and mixtures thereof are used in the form of water-soluble or water-dispersible salts, for example sodium, potassium, magnesium, ammonium, mono-, di- and triethanol-ammonium salts and analogous alkylol ammonium salts.

25

A particularly preferred anionic surfactant is polyoxyalkylene alkylether sulfate which may be used in an amount of, for example of 0.01 to 5% by wt.

30 Suitable nonionic surface-active agents include, for example fatty alcohol ethoxylates (alkyl-polyethylene glycols),

- 11 -

alkylphenol-polyethylene glycols, alkylation-polyethylene glycols, fatty amine ethoxylates (alkylamino-polyethylene glycols), fatty acid ethoxylates (acyl polyethylene glycols), polypropylene glycol ethoxylates (Pluronic 5 (Registered Trademark)), fatty acid alkyl amides (fatty acid amino polyethylene glycols), saccharose esters, sorbitol esters, and polyglycol ether.

Suitable amphoteric surface-active agents to be added to the 10 shampoos include N-alkyl-β-amino dipropionates having 12 to 18 alkyl carbon atoms as alkali metal salts and mono-, di- and trialkylol-ammonium salts, N<sup>+</sup>acylamido-alkyl-N, N-dimethylacetobetaine, preferably N-acyl-amidopropyl-N,N-dimethylacetobetaine having 8 to 18 acyl carbon atoms, 15 alkyl-dimethylsulfopropyl-betaine having 12 to 18 alkyl carbon atoms, amphoteric surfactants of the imidazoline type (Trademarks: Miranol Steinapon), preferably the sodium salt of 1-(β-carboxy-methyloxethyl)-1-(carboxymethyl)-2-lauryl-imidazolinium; amine oxides, for example alkyl-dimethyl-20 amine oxide having 12 to 18 alkyl carbon atoms, fatty acid amidoalkyl-dimethylamine oxide.

A particularly preferred amphoteric surfactant is cocoamide propyl betaine.

25 Among suitable cationics are, for example, quaternary ammonium salts, such as dialkyl-dimethyl-ammonium, chloride or bromide having 10 to 24, preferably 12 to 18 carbon atoms, in the alkyl portion, alkyl-dimethyl-ethylammonium 30 chloride or bromide having 10 to 24 alkyl carbon atoms, alkyl-trimethyl-ammonium chloride or bromide having 10 to 24

- 12 -

alkyl carbon atoms, preferably cetyl-trimethyl ammonium chloride or bromide, alkyl-trimethyl ammonium chloride or bromide having 10 to 22 alkyl carbon atoms, or alkyl-dimethyl-benzyl ammonium chloride or bromide having 10 to 5 24, preferably 12 to 18, carbon atoms in the alkyl portion.

Other optional components are as follows:

The liquid personal cleansing compositions of the present 10 invention may optionally also include water-dispersible, gel-forming polymers. This polymer is preferably an anionic, nonionic, cationic or hydrophobically modified polymer, selected from cationic polysaccharides of the cationic guar gum class with molecular weights of 1,000 to 15 3,000,000, anionic, cationic and nonionic homopolymers derived from acrylic and/or methacrylic acid, anionic, cationic and nonionic cellulose resins; cationic copolymers of dimethyldialkyl ammonium chloride and acrylic acid; cationic homopolymers of dimethyldialkyl ammonium chloride; 20 cationic polyalkylene and ethoxypolyalkylene imines polyethylene glycol having molecular weights from 10,000 to 4,000,000; and mixtures thereof. Preferably, the polymer is selected from Sodium Polyacrylate, Hydroxy Ethyl Cellulose, Cetyl Hydroxy Ethyl Cellulose, and Polyquaternium 10.

25

The polymer is preferably included in the compositions of the present invention at a level of from about 0.1 parts to 1 part, more preferably 0.1 parts to 0.5 parts of the composition. The polymers improve the sensory feel of the 30 lipid on skin in addition to providing product stabilization. The improved sensory feel results from

- 13 -

reduced tackiness and greasiness and improved smoothness. In an especially preferred embodiment a mixture of polymers is used, for example those polymers preferred for product stabilization, and/or those preferred for improved sensory 5 feel. Preferred polymers for improved sensory feel are selected from polyethylene glycol, hydroxypropyl guar, guar hydroxypropyl trimonium chloride, polyquaternary 3, 5, 6, 7, 10, 11 and 24 and mixtures thereof.

10 A variety of additional ingredients may be incorporated into the compositions of the present invention. These materials include, but not limited to, liquid appearance aids, salts and their hydrates and other "filler materials" as listed in U.S. Patent No. 5,340,492 to Kacher et al., and U.S. Patent 15 No, 4,919,934 to Deckner et al. ,both of which are herein incorporated by reference.

Other non-limiting examples of these additional ingredients include vitamins and derivatives thereof (e.g., ascorbic 20 acid, vitamin E, tocopherol acetate, and the like); sunscreens; thickening agents (e.g., polyol alkoxy ester, available as Crothix from Croda at levels up to 2% and xanthan gum at levels up to 2%); preservatives for maintaining the anti-microbial integrity of the 25 compositions; anti-acne medicaments (resorcinol, salicylic acid, and the like); antioxidants; skin soothing and healing agents such as aloe vera extract, allantoin and the like; chelators (e.g., EDTA and hydroxy ethan diphosphoric acid) and sequestrants; and agents suitable for aesthetic 30 purposes such as fragrances, essential oils, skin sensates, pigments, pearlescent agents (e.g., mica and titanium

- 14 -

dioxide), additives to impart a draggy rinse feel (e.g., fumed silica), additives to enhance deposition (e.g., maleated soybean oil at levels up to 3%), lakes, colorings, and the like (e.g., clove oil, menthol, camphor, eucalyptus 5 oil and eugenol).

The compositions of the present invention have a pH of about 6 to 8.5, preferably 6.5 to 8.5.

10 Compositions of the present invention have a viscosity of 100-5000 centistokes measured using a Brookfield Viscometer, Spindle 2 and 3 at 12 rpm for 30 seconds at 25°C.

15 The compositions of the present invention exhibit no precipitation or phase separation after 4 weeks of storage at room temperature (i.e., about 25°C).

**EXAMPLES**

20 Examples 1, 2 and 3

The following examples were prepared as follows:

- 15 -

	Example1	Example2	Example3
Disodium Laurylether sulfsuccinate(DSLES)	4.25%	5%	7%
Disodium Lauryl sulfsuccinate(DSLS)	4.25%	5%	7%
Potassium Laurate		3%	3%
Potassium Myristate		2%	2%
Lauric acid	4.25%		
Isostearic acid			4%
Potassium hydroxyde	0.6%		
Cocamido propyl betaine(CAPB)	2.6%	3%	0.6%
Sodium Laurylether sulfate(SLES)	1.7%	2%	0.4%
Glycerine	20%	20%	20%
Dipropylene glycol		3%	
Amphoteric polymer	0.3%		0.3%
Cationic polymer		0.6%	
Alkyl acrylate polymer	0.5%	0.5%	0.5%
Ethylene glycol distearate(EGDS)	2%	2%	2%
Dibutylhydroxy toluene(BHT)	0.05%	0.05%	0.05%
EDTA	0.05%	0.05%	0.05%
Hydroxy ethane diphosphoric acid		0.2%	0.2%
Preservative	0.3%	0.3%	0.3%
Perfume	0.7%	0.7%	0.7%
Water	To 100%	To 100%	To 100%

Processing: Example 1

5

Example 1 above was processed as follows:

10 1) A mixture of glycerin, water, potassium hydroxide and SLES was heated to 75-80°C until becoming quite fluid(mixture-1);

2) Lauric acid, preservative and BHT were added to mixture-1 with agitation(mixture-2);

- 16 -

- 3) DSLES, DSLS and CAPB were added to mixture-2 with agitation(mixture-3);
- 4) A mixture of EDTA and water was mixed at 75-80°C (mixture-4);
- 5) A mixture of amphoteric polymer and water was added into mixture-4 with agitation (mixture-5);
- 6) EGDS and alkyl acrylate polymer were separately added to mixture-5 with agitation(mixture-6);
- 7) Mixture-6 was cooled to 45°C;
- 10 8) Perfume was added to mixture-9 and cooled to 35°C.

- 17 -

Examples 5, 6 & Comparative Example A

	Example 5	Comparative A	Example 6	Comparative A
Disodium Laurylether sulfsuccinate(DSLES)	7%	7%	4.25%	4.25%
Disodium Lauryl sulfsuccinate(DSLS)	7%	7%	4.25%	4.25%
Potassium Laurate	3%	3%		
Potassium Myristate	2%	2%		
Lauric acid			4.25%	4.25%
Isostearic acid	4%			
Potassium hydroxyde			0.6%	0.6%
Cocamido propyl betaine(CAPB)	0.6%	0.6%	2.6%	2.6%
Sodium Laurylether sulfate(SLES)	0.4%	0.4%	1.7%	1.7%
Glycerine	20%	20%	20%	20%
Dipropylene glycol				
Amphoteric polymer	0.3%	0.3%	0.3%	0.3%
Cationic polymer				
Alkyl acrylate polymer	0.5%	0.5%	0.5%	
Ethylene glycol distearate(EGDS)	2%	2%	2%	2%
Dibutylhydroxy toluene(BHT)	0.05%	0.05%	0.05%	0.05%
EDTA	0.05%	0.05%	0.05%	0.05%
Hydroxy ethane diphosphoric acid	0.2%	0.2%		
Preservative	0.3%	0.3%	0.3%	0.3%
Perfume	0.7%	0.7%	0.7%	0.7%
Water	To 100%	To 100%	To 100%	To 100%
RT stability: Separation	good	no good	good	no good

5 From the data obtained for Example 5 and Comparative A, it can be seen that 4% isostearate provided good stability (no phase separation) while an absence of isostearic acid (Comparative A) resulted in instability. Similarly, the use of alkyl acrylate (Example 6) resulted in stability while 10 the absence (Comparative B) resulted in instability.


- 18 -

CLAIMS


5 1. A stable, high glycerol liquid detergent composition comprising:

(a) 1-25%, by wt. of one or more sulfosuccinic acid monoesters represented by Formulae (I), (II), (III) or (IV).

10




15




20

- 19 -



or



15

wherein each of  $M_1$  to  $M_4$  is H,  $NH_4$ , an alkali metal or a hydroxyalkyl substituted ammonium;  $R_1$  and  $R_2$  are each an alkyl or hydroxyalkyl group having about 8 to 20 carbon atoms;  $R_3$  is H or  $CH_3$ ;  $AO$  is an oxyalkylene group having 2 or 3 carbon atoms, and  $n$  represents an integer of 0 to 20;

(b) 10% to 50% by wt. glycerin;

(c) at least 4% by wt. of an alkali metal soap of  $C_{12-}$

25 C<sub>24</sub> fatty acid and/or alkanol soap; and

(d) 0.01-5%, by wt. of a thickener selected from acrylate copolymer, hydroxy alkyl cellulose, cationic guar gums and isostearic acid; and

- 20 -

(e) balance water.

2. A composition according to claim 1, wherein the composition additionally comprises 0.01-10% of a surfactant selected from anionic, nonionic and amphoteric and cationic surfactants and mixtures thereof.
- 10 3. A composition according to 2, wherein the amphoteric surfactant is a betaine.
4. A composition according to any one of claims 1 to 3, wherein the pH of the composition is about 6-8.5.
- 15 5. A composition according to any of the preceding claims, wherein the composition has a viscosity of 100 to 5000 centistokes.
- 20 6. A composition according to any of the preceding claims, wherein the composition has no precipitation or phase separation after 4 weeks at room temperature.