



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2857575 A1 2013/06/06

(21) **2 857 575**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2012/11/30
(87) Date publication PCT/PCT Publication Date: 2013/06/06
(85) Entrée phase nationale/National Entry: 2014/05/30
(86) N° demande PCT/PCT Application No.: GB 2012/052973
(87) N° publication PCT/PCT Publication No.: 2013/079967
(30) Priorité/Priority: 2011/12/02 (GB1120772.7)

(51) Cl.Int./Int.Cl. *A23G 1/48* (2006.01),
A23G 1/32 (2006.01), *A23G 1/42* (2006.01),
A23L 1/303 (2006.01)
(71) Demandeur/Applicant:
IP SCIENCE LIMITED, GB
(72) Inventeur/Inventor:
PETYAEV, IVAN, GB
(74) Agent: MARKS & CLERK

(54) Titre : PRODUITS ALIMENTAIRES A BASE DE CACAO
(54) Title: COCOA-BASED FOOD PRODUCTS

(57) **Abrégé/Abstract:**

The invention is concerned with food products comprising one or more cocoa bean products and a carotenoid compound, particularly with food products which are, or comprise, chocolate. The products of the invention may be used in reducing elevated total cholesterol, triglycerides and inflammatory damage, as well as improving tissue microcirculation and tissue oxygenation.



(51) **International Patent Classification:**
A23G 1/48 (2006.01) A23L 1/303 (2006.01)
A23G 1/42 (2006.01) A23G 1/32 (2006.01)

(21) **International Application Number:**
PCT/GB2012/052973

(22) **International Filing Date:**
30 November 2012 (30.11.2012)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
1120772.7 2 December 2011 (02.12.2011) GB

(71) **Applicant: IP SCIENCE LIMITED** [GB/GB]; 2nd Floor,
The Platinum Building, St John's Innovation Park, Cowley
Road, Cambridge Cambridgeshire CB4 0DS (GB).

(72) **Inventor: PETYAEV, Ivan**; 1 Cambridge Road, Coton,
Cambridge CB23 7PJ (GB).

(74) **Agent: J A KEMP**; 14 South Square, Gray's Inn, London,
Greater London WC1R 5JJ (GB).

(81) **Designated States** (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

(54) **Title:** COCOA-BASED FOOD PRODUCTS

(57) **Abstract:** The invention is concerned with food products comprising one or more cocoa bean products and a carotenoid compound, particularly with food products which are, or comprise, chocolate. The products of the invention may be used in reducing elevated total cholesterol, triglycerides and inflammatory damage, as well as improving tissue microcirculation and tissue oxygenation.

WO 2013/079967 A1

Cocoa-Based Food Products

Field of the Invention

5

This invention relates to cocoa-based food products, such as chocolate, which have beneficial effects on parameters of metabolism in individuals including levels of triglycerides, cholesterol and other lipids, molecular oxygen transport and its
10 metabolism, oxygen tissue saturation and microcirculation, control of hypoxia/ischaemia, as well as markers of inflammation and inflammatory oxidative damage.

Background to the Invention

15

Food products based on cocoa beans, such as chocolate and other products containing cocoa solids, cocoa butter, cocoa liquor, and/or their derivatives, are widespread in the Western society. Although these cocoa-based products often contain flavonols and
20 flavonoids, which have been associated with certain health benefits, they also contain high levels of cocoa butter or other high-fat ingredients. This high fat content means that cocoa-based food products are one of the dietary factors responsible for the growth of Metabolic Syndrome, Diabetes II and Obesity in the
25 modern society.

Cocoa-based food products, such as chocolate, have been associated with health benefits, such as improvements in endothelial vascular function, including positive effects on blood pressure, and with
30 antioxidant and anti-inflammatory properties [Keen *et al Am J Clin. Nutr.* 2005, 81(suppl):298S-303S].

However, the effects of chocolate on blood lipid concentrations are either inconclusive or negative. Consumption of 105 g of dark
35 chocolate has been reported to result in a mild reduction of total blood lipid concentrations of 11% [Cesar *et al Clinical & Developmental Immunology*, 2005; 12(1): 11-17], but this trial was

done on an exceptional group of young elite sportsmen and positive blood lipid changes may be attributed to the overall improvement of their physical performance.

5 Meta-analysis of eight clinical trials on dark chocolate or cocoa powder showed an observed reduction in the cholesterol levels of up to 5.82 mg/dL, which was statistically insignificant [Lei *et al* *Am J Clin Nutr* 2010; 92 : 218-25]. Furthermore, the amount of the daily administered doses of cocoa phenols in these trials was
10 equivalent of consumption of 100g or more of the dark chocolate [Cheng *et al* *Molecules* 2009, 14: 200-209]. In other studies, when dark chocolate was consumed from 30 to 100 g daily, no changes were observed in the blood lipids [Taubert *et al* *JAMA*. 2007; 4, 298(1):49-60, Grassi *et al* *Hypertension*. 2005; 46(2):398-405].

15

The level of triglycerides in all the above studies above either had insignificant trends or did not change at all.

There are no reports of any health benefit, including on blood
20 lipids, arising from consumption of the most common forms of chocolate, including milk chocolate and white chocolate.

Lycopene is known to be as a potent antioxidant. Its mild cholesterol-lowering effect has been reported from 13 clinical
25 studies and was 7.55+6.15 mg/dL [Ried *et al* *Maturitas*. 2011, 68(4):299-310]. However this effect was only observed for daily doses 25 mg of lycopene or above. There were no reports on the effect of lycopene on elevated triglycerides in human.

30 Although 25 mg of lycopene and above is considered to be safe for certain periods of administration, it is far above the daily level which could be consumed with a diet rich with tomato or tomato processed products (about 6-10 mg). Daily consumption of 6-10 mg lycopene has been reported to have no effect on cholesterol or
35 other blood lipids [Bose *et al* *Singapore Med J* 2007; 48 (5): 415-420; Upritchard *et al* *Diabetes Care*, 2000, 23 6: 733-735].

Summary of the Invention

The present invention is based on the unexpected finding that the combination of a carotenoid and a cocoa bean product can be used
5 to reduced elevated cholesterol, reduce elevated triglyceride levels, reduce inflammatory oxidative damage and improve tissue microcirculation and oxygen transport. Given the high fat and sugar content of chocolate, such a finding was highly unexpected and indeed counterintuitive.

10

Accordingly, the invention provides a food product comprising one or more cocoa bean products and a carotenoid compound.

15

The invention also provides such a food and/or beverage and/or nutraceutical product for use in:

- (a) reducing levels of elevated cholesterol, LDL and/or triglyceride in an individual, preferably where the individual has elevated levels of cholesterol, LDL and/or triglyceride;
- 20 (b) reducing subclinical or clinical inflammation; reducing anti-inflammatory oxidative damage; increasing plasma molecular oxygen transport, microcirculation and tissue oxygen saturation, reducing already developed liver (micro-) damage and liver steatosis, liver and other organs, including
25 peripheral, tissue hypoxia or ischaemia, increasing antioxidant activity and/or reducing or delaying symptoms of ageing in an individual;
- (c) reducing postprandial cholesterol- and triglyceride-aemias, reducing size of chylomicrons and increasing rate of their
30 clearance, reducing postprandial inflammatory and oxidative stress, reducing postprandial or other liver (micro-) damage and liver steatosis, liver and other organs, including peripheral tissue hypoxia or ischaemia, or delaying of above mentioned symptoms of fat, or excessive, or imbalance food
35 intake in an individual;
- (d) increasing oxygen transport in a subject, preferably where the subject has a respiratory disorder and/or lung damage;

and/or strenuous physical or mental performance; and/or
muscle wasting conditions; and/or

(e) slimming, weight reduction or dieting.

5 The invention additionally provides a method of:

(a) improving the appearance and performance of an individual comprising administering a nutracosmetic formulation or food or beverage product of the invention to the individual; and/or

10 (b) reducing or delaying signs of aging in an individual, preferably visible signs of aging, comprising administering a food product of the invention to the individual.

The invention also provides a method of:

15 (a) reducing levels of elevated cholesterol, LDL and/or triglyceride in the blood of an individual comprising administering a food product of the invention to an individual in need thereof;

(b) reducing subclinical or clinical inflammation; reducing anti-inflammatory oxidative damage; increasing plasma molecular oxygen transport, microcirculation and tissue oxygen saturation, reducing already developed liver (micro-) damage and liver steatosis, liver and other organs, including peripheral, tissue hypoxia or ischaemia; increasing
20 antioxidant activity and/or reducing or delaying symptoms of ageing in an individual; comprising administering a food product of the invention to an individual in need thereof;

(c) reducing postprandial cholesterol- and triglyceride- aemias, reducing size of chylomicrons and increasing rate of their clearance, reducing postprandial inflammatory and oxidative stress, reducing postprandial or other liver (micro-) damage and liver steatosis, liver and other organs, including peripheral tissue hypoxia or ischaemia, or delaying of above mentioned symptoms of fat, or excessive, or imbalance food
25 intake in an individual; comprising administering a food product of the invention to an individual in need thereof;

- (d) providing nutrition to an individual comprising administering a food product of the invention to an individual in need thereof; and/or strenuous physical or mental performance; and/or muscle wasting conditions; and/or
- 5 (e) slimming, weight reduction or dieting comprising administering a food product of the invention.

The invention also provides a chocolate bar and/or chocolate beverage comprising a carotenoid, preferably where the carotenoid
10 is a lycopene compound.

Detailed Description of the Invention

- This invention relates to the unexpected finding that
15 incorporating carotenoid compounds into food products which contain cocoa-bean based products, such as cocoa solids, cocoa butter, cocoa liquor, and/or their derivatives, causes these food products to exert a positive effect on levels of triglycerides, cholesterol, LDL, and other metabolic parameters in individuals,
20 despite being rich in saturated and unsaturated fats. Given the high fat content of food products such as chocolate, the finding that the combination of a carotenoid and chocolate is able to reduce those parameters was unexpected and counter-intuitive.
- 25 An aspect of the invention provides a food product which comprises one or more cocoa-bean products and a carotenoid compound. Typically, the carotenoid compound is an isolated carotenoid compound.
- 30 The food product may comprise a homogenous matrix which contains the cocoa-bean products and the carotenoid compound. For example, the cocoa-bean products and carotenoid compound may be blended together in a chocolate or cocoa-butter matrix.
- 35 A cocoa bean product is an extract, fraction or isolate from cocoa beans (i.e. beans of the cacao tree (*Theobroma cacao*)). Suitable cocoa bean products are well-known in the art and include cocoa

solid, cocoa liquor and/or cocoa butter. For example, a food product may comprise one or more of cocoa solid, cocoa liquor and/or cocoa butter.

- 5 In some instances cocoa nibs or fragments thereof, chocolate liquor, partially and fully-defatted cocoa solids (e.g. cocoa powder), cocoa extract or a fraction thereof may be employed.

Cocoa solid (also known as cocoa powder) is a low-fat extract of
10 cocoa beans, which contains flavanols, flavanoids, caffeine and theobromine. Cocoa solid may be produced by removing the fat component (cocoa butter) from the cocoa bean and grinding the remaining material, excluding the shell, to a powder using techniques which are well-known in the art, such as Broma
15 processing. In some embodiments, cocoa powder may be treated with an alkaline substance such as potassium carbonate to reduce acidity and darken the colour (Dutch processing).

Cocoa butter is a high-fat extract of cocoa beans which is high in
20 stearic acid, palmitic acid and other saturated fats. Cocoa butter may be produced from whole or ground cocoa beans using techniques which are well-known in the art.

Cocoa liquor is a cocoa bean extract which contains both cocoa
25 solid and cocoa butter. Cocoa liquor may be produced by grinding and melting the cocoa bean nib (centre) to a smooth liquid state in accordance with techniques which are well-known in the art. Chocolate liquor does not contain non-cocoa vegetable fat and may also be referred to as "chocolate", "unsweetened chocolate",
30 "baking chocolate", or "bitter chocolate".

In other embodiments, cocoa bean products may include derivatives or fermentation products of cocoa bean extracts, isolates or fractions.

35

Preferably, the food product comprises cocoa butter; cocoa solid; or both cocoa butter and cocoa solid.

For example the food product may contain at least 1% by weight, at least 10% by weight, at least 15% by weight, at least 20% by weight, at least 25% by weight or at least 30%, or at least 40% by weight cocoa butter. The food product may contain an amount of cocoa butter in a range comprising any of the above two values as endpoints.

In some embodiments, a food product may further comprise non-cocoa fats, such as vegetable or animal fats in addition to cocoa butter.

In some embodiments, a food product may be devoid of cocoa butter. For example, a food product may contain animal or non-cocoa vegetable fat instead of cocoa butter. Non-cocoa vegetable fats may include vegetable oils. Suitable vegetable oils, such as palm oil, soybean oil rapeseed oil and olive oil, are well known in the art.

The total fat content of a food product described herein may be at least 10% by dry weight, at least 15% by dry weight, at least 20% by dry weight, at least 25% by dry weight, at least 30% by dry weight or at least 35% by dry weight or at least 40% by dry weight. The fat content may be, for instance, in a range comprising any two such values as endpoints.

Additionally or alternatively, the food product may contain at least 5% by weight, at least 15% by weight, at least 20% by weight, at least 25% by weight, at least 30% by dry weight or at least 35% by weight, or at least 40% by weight dry cocoa solid. In some instances, the amount of cocoa solid may be at least 50% by weight, at least 60% by weight, at least 75% by weight, at least 80% by weight, at least 85% by weight, at least 90% by weight or even at least 95% by weight dry cocoa solid, particularly when the food stuff is a dark chocolate. The amount of weight of dry cocoa solid may be, for instance, in the range comprising any two of those values as endpoints.

In some embodiments, a food product may be devoid of cocoa solid.

For the avoidance of doubt, aspects of the invention provide food
5 products which comprise all combinations of the above parameters
of cocoa solid, cocoa butter and total fat.

In some embodiments, the cocoa bean products may form a chocolate
matrix. The carotenoid compound may be incorporated into the
10 chocolate matrix by blending or admixing.

Any cocoa-based food product may be supplemented with a carotenoid
compound as described herein. For example, the food product may be
a foodstuff, a beverage or a dietary supplement or nutraceutical
15 product.

Foodstuff products include bread, flour, cereal, biscuit, pastry,
dairy products, such as cheese spread, cheese, cream and yoghurt,
fillings, pastes, sauces and mousses. Other suitable foodstuffs
20 are well known in the art.

In some preferred embodiments, foodstuff products may include
confectionary products, such as chocolate. Especially preferred
embodiments of the invention provide chocolate comprising a
25 carotenoid compound, as described herein.

Chocolate may include dark chocolate, milk chocolate, or white
chocolate.

30 In one preferred instance, the foodstuff of the invention may be a
chocolate bar, for instance a dark, plain or milk chocolate bar
comprising a carotenoid, such as any of those discussed herein.
The amount of carotenoid in the bar may be, for instance, any of
the amounts of carotenoid specified herein.

35 Dark chocolate, milk chocolate and white chocolate are subject to
defined identity standards (for example, by the Food and Drug

Administration (USA), EU and Food Standards Agency (UK); see for example EU directive 2000/36/EC; FDA 21 CFR Part 163 Federal Register: 2002 67 193 62171-62178). In one instance, a composition of the invention may be a standard of identity (SOI) chocolate, in
5 others it is a non-SOI chocolate.

The ingredients of dark chocolate, milk chocolate, white chocolate or other forms of chocolate are well-known in the art. For example, dark chocolate typically comprises sugar, cocoa butter
10 (e.g. at least 12% by weight), cocoa solids (e.g. at least 35% by weight), and optionally vanilla. Fat content may vary but averages between 30%-35%. Dark chocolate is sometimes referred to as sweet or semi-sweet chocolate. Milk chocolate may comprise sugar, cocoa butter, cocoa solids, vanilla or other flavourings, and milk, milk
15 powder or cream. Milk chocolate typically contains at least 20% cocoa solid and at least 12% milk solids by weight. White chocolate may comprise sugar, cocoa butter, milk or milk powder, and vanilla and lacks cocoa solids. White chocolate typically contains at least 20% cocoa butter, 14% total milk solids, and
20 less than 55% sugar.

In one instance, the food product of the invention may be about 100g, 150g, 200g, 250g, 300g, 400g or 500g in weight or may have a weight in a range with any two of those values as endpoints. In a
25 preferred instance, the foodstuff may be a chocolate bar of such weight.

The foodstuff may be a candy bar, for instance a chocolate coated candy bar. The foodstuff may take the form of individual
30 chocolates, bagged chocolates or a box of chocolates. The chocolate may be in a formed shape. In one instance the foodstuff is an Easter egg. The invention may be provided in the form of chocolate icing or a cake comprising a carotenoid and chocolate. The invention also provides fruit or nuts coated with a chocolate
35 of the invention. The invention also provides sweets or candy coated with a chocolate of the invention.

The invention also provides a chocolate of the invention provided in the form of a single serving dose, for instance in 10 to 30g amounts, as well as a packet of such single serving doses. The invention also provides a chocolate bar of the invention
5 segmented, for instance segmented so that it can be broken into single serving dosages.

The foodstuff of the invention may be, in other instances, a cake, cheesecake, baked snack, brownie, cookie or biscuit, a meal
10 replacement bar, a rice cake, ice cream or other pudding or dessert. In some instances, the invention provides such products coated in, or comprising, a chocolate of the invention. The products may for instance comprise the chocolate in the form of chips or in a central region.

15 Beverages may include any drink which comprises cocoa-bean products and may include cocoa, drinking chocolate, milk shakes, and other dairy and non-dairy drinks. Beverages may be non-alcoholic or alcoholic. The formulation of suitable beverages is well-known in the art. In one preferred instance, the beverage is
20 a chocolate milkshake. In another instance, a powder, gel or cube for making up as a beverage is also provided. The invention also provides a hot chocolate, chocolate or cocoa drink, as well as a chocolate/cocoa shot drink comprising a product of the invention.

25 Dietary supplements or nutraceutical products may be in any form suitable for oral administration (e.g., by ingestion) and may be presented as discrete units such as capsules, cachets or tablets; as a powder or granules; as a solution or suspension in an aqueous
30 or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion; as a bolus; as an electuary; or as a paste.

The invention also provides a food-stuff intended for dieters
35 which is, or comprises, a foodstuff of the invention. The invention provides for the use of the products described herein for slimming, dieting or weight reduction. The invention also

provides for products for diabetics comprising, or consisting of, a foodstuff of the invention. In one instance, the invention provides a diabetic chocolate, where the chocolate is a chocolate of the invention.

5

In one preferred instance, a foodstuff of the invention may be provided with packaging and/or wrapping. Such packaging/wrapping may indicate the benefits of the invention and/or suggest consumption at, or near, mealtimes for maximal benefit. The
10 packaging/wrapping may indicate the benefits of the product in slimming, decreasing cholesterol, and/or triglyceride levels. In another instance, the packaging may refer to the ability of the product to improve oxygen transport. The packaging may refer to treating or ameliorating any of the conditions mentioned herein.
15 The packaging may be a sachet, for instance where the product is to be made up as a beverage.

The invention also provides food products targeted at sports people. For instance, the products may be used to reduce weight in
20 such subjects or bring about any of the other benefits highlighted herein for the products of the invention. The products may be packaged or wrapped and include an indication of their ability to increase oxygen transport. The products of the invention may be used to reduce recovery time. The products may be used by
25 climbers, particularly those climbing at altitude

In addition to cocoa-bean products, food products described herein comprise a carotenoid compound.

30 The carotenoid compound may be isolated in the products of the invention. An isolated carotenoid compound is outside the physical milieu or environment in which it occurs in nature. For example, an isolated carotenoid compound may be free or substantially free from its natural environment e.g. it is not
35 contained in the natural plant material with which it is naturally associated. Isolated carotenoid compounds include compounds which have been isolated, concentrated, purified or partially purified

from natural sources, such as plants, and compounds which have been produced synthetically.

5 The food product will typically provide an effective amount of carotenoid, such as lycopene. The food product may comprise 0.0001% to 1%; 0.001% to 1%; or 0.01% to 0.1% by weight of carotenoid compound. For example, the food product may comprise 0.001 to 10mg of carotenoid compound per one gram of food product, for example, 0.01 to 10 mg per gram or 0.1 to 1 mg per one gram of
10 food product. The product of the invention will typically provide an effective amount of the carotenoid, for instance an amount effective to alter one of the parameters referred to herein.

The food product may be in a unit dose form which allows a
15 controlled daily dose of carotenoid, preferably lycopene, to be consumed. For example, the food product may be formulated to provide a daily dose of 0.1mg to 100mg of lycopene, preferably 0.5 to 50 mg of lycopene. In some instances, a product of the invention may provide about 0.1mg, 0.2mg, 0.5mg, 1mg, 1.5mg, 2mg,
20 2.5mg or more of carotenoid, such as about 3, 4, 5, 6, 7, 8, 9, or 10mg of carotenoid. In some instances, the amount of carotenoid may be about 10, 15, 20 or 25mg, or up to those levels. Preferably the carotenoid is lycopene. The product may comprise an amount of carotenoid which is in a range with any two of the values
25 mentioned herein as endpoints.

In one instance, the foodstuff or product provides from 0.1 to 1.0mg of carotenoid per gram of food product for example at least 0.2, 0.3, 0.4 or 0.5mg carotenoid per gram of food product, with
30 in some instances, up to 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0mg of carotenoid per gram of product. In one preferred instance, those values are employed where the product is chocolate and/or carotenoid is lycopene, preferably both. In another instance, the level of foodstuff administered is enough to reduce
35 any of the makers discussed herein, preferably to near, or at, or below baseline levels for a healthy control or below baseline prior to administration of the product.

In some instances, the ratio of carotenoid to triglyceride or other fat molecules in the products of the invention may be from 1:1000 to 1:100,000, for instance from 1:2000 to 1:50,000, or from
5 1:5000 to 1:25,000.

The carotenoid and cocoa bean product may be present in a synergistic amount. For instance, they may be present where the combination produces a greater effect on any of the parameters
10 mentioned herein than either individually when provided in the same amount. The invention therefore also provides a synergistic combination of a carotenoid and cocoa bean product. The invention also provides for the use of a carotenoid, such as any of those referred to herein, to treat any of the conditions mentioned
15 herein, where the subject is also being administered chocolate and also the use of chocolate to treat any of the conditions mentioned herein where the subject is also being administered a carotenoid. Typically the carotenoid and chocolate will be administered together, for instance eaten together, or within 5, 10, 15, 30, 45
20 or 60 minutes of each other. The two may be given simultaneously.

Carotenoid compounds are tetraterpenoids which contain long polyene chains. Carotenoid compounds include xanthophylls such as lutein, capsanthin and zeaxanthin, and carotenes, such as beta-
25 carotene, alpha-carotene, zeto-carotene, and lycopene compounds.

Lycopene compounds may include lycopene, 1-HO-3', 4'-didehydrolycopene, 3, 1'-(HO)2-gamma-carotene, 1,1'-(HO)2-3, 4, 3', 4'-tetrahydrolycopene, 1, 1'-(HO)2-3, 4-didehydrolycopene.
30

In preferred embodiments, the carotenoid compound is lycopene.

Lycopene is an open-chain unsaturated C₄₀ carotenoid of structure I (Chemical Abstracts Service Registry Number 502-65-8).
35

Structure I.



Lycopene

Molecular Weight = 536.89

Exact Mass = 536

Molecular Formula = C₄₀H₅₆

Molecular Composition = C 89.49% H 10.51%

5 Lycopene occurs naturally in plants such as tomatoes, guava, rosehip, watermelon and pink grapefruit and any such sources of lycopene may be, for instance, employed.

10 Lycopene for use as described herein may comprise one or more different isomers. For example, lycopene may include *cis*-lycopene isomers, *trans*-lycopene isomers and mixtures of the *cis*- and *trans*-isomers. Lycopene may comprise at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70% , at least 80% , at least 90% , or at least 95% (Z)-isomers,
 15 (all-E)-isomers, or *cis*-isomers, such as 5-*cis*- or 9-*cis*- or 13-*cis*-isomers, which have improved bioavailability relative to *trans* isomers. *Trans* isomers may isomerise into *cis* forms *in vivo*, or during storage and processing.

20 Carotenoid compounds, such as lycopene, for use as described herein may be natural i.e. obtained from a natural source, for example, extracted from a plant, such as a tomato or melon. In one instance, oleoresin, particularly tomato oleoresin, may be employed in the invention. A range of methods for extracting,
 25 concentrating and/or purifying carotenoids from plants are known in the art. For example, solvent extraction using ethanol, DMSO, ethyl acetate, hexane, acetone, soya or other vegetable oil, or non-vegetable oils may be employed.

30 Carotenoid compounds, such as lycopene, for use as described herein may be synthetic i.e. produced by artificial means, for example, by chemical synthesis. A range of methods for chemical synthesis of lycopene and other carotenoids are known in the art.

For example, a three-stage chemical synthesis based on the standard Wittig olefination reaction scheme for carotenoid synthesis may be employed, in which an organic solution of C₁₅ phosphonium methanesulfonate in dichloromethane (DCM) and an
5 organic solution of C₁₀ dialdehyde in toluene are produced, and the two organic solutions are gradually combined with sodium methoxide solution and undergo a condensation reaction to form crude lycopene. The crude lycopene may then be purified using routine techniques, for example by adding glacial acetic acid and
10 deionized water to the mixture, stirring vigorously, allowing the aqueous and organic phases to separate, and extracting the organic phase containing DCM and crude lycopene with water. Methanol is added to the organic phase and the DCM removed via distillation under reduced pressure. The crude methanolic lycopene solution is
15 then be heated and cooled to crystalline slurry that is filtered and washed with methanol. The lycopene crystals may then be recrystallized and dried under heated nitrogen. Synthetic carotenoids, such as lycopene, are also available from commercial suppliers (e.g. BASF Corp, NJ USA).

20

Synthetic carotenoid compounds, such as lycopene, may comprise an increased proportion of cis isomers relative to natural carotenoid compounds. For example, synthetic lycopene may be up to 25% 5-cis, 1% 9-cis, 1% 13-cis, and 3% other cis isomers, whilst lycopene
25 produced by tomatoes may be 3-5% 5-cis, 0-1% 9-cis, 1% 13-cis, and <1% other cis isomers. Since cis-lycopene has increased bioavailability relative to trans-lycopene, synthetic lycopene is preferred in some embodiments.

30 Derivatives of carotenoids as described above may be produced by chemical synthesis analogous to the synthesis described above or by chemical modification of natural carotenoids extracted from plant material.

35 A food product as described herein may contain a single carotenoid compound (e.g. lycopene) or more than one carotenoid compound

(e.g. lycopene and beta-carotene). Typically, each carotenoid compound will be present in a range of different isomeric forms.

The food product may be produced by admixing or blending the
5 cocoa-bean products, such as cocoa butter and cocoa solids, and optionally one or more other ingredients, and the carotenoid compound under conditions which allow the carotenoid compound to incorporate into the matrix of the food product.

10 Other ingredients may include sugar, vanilla, milk, milk powder, emulsifying agents, such as soy lecithin or polyglycerol polyricinoleate (PGPR; E476), whey or potato peptides and/or proteins, soy products, such as soy proteins, soy extracts and/or soy isoflavones, vegetable oils or animal fats, nut-based
15 products, such as nut powders and nut extract, starch and polysaccharides.

The cocoa-bean products may be in a dry, liquid, aerosol, frozen or melted form for admixing or blending with the carotenoid
20 compound. For example, chocolate for blending may be in liquid form (i.e. melted chocolate).

In some preferred embodiments, the cocoa-bean products and the carotenoid compound in mixable forms and have the same or similar
25 viscosities.

Suitable methods of mixing and blending, including mechanical blending, are well-known in the art.

30 In one instance, a carotenoid is added whilst the chocolate is being made or chocolate is melted and the carotenoid added. The chocolate may be added to a mould to give products of a particular shape and/or size.

35 The invention also provides for a method of producing a food product, such as a food product of the invention, which comprises adding a carotenoid during production of the food product. For

instance, the carotenoid may be added during the preparation of chocolate.

Products of the invention may also contain other ingredients such as flavourings, emulsifiers, colourings and/or preservatives. In some cases the products may comprise nuts, particularly where the product is a chocolate, such as walnuts, hazelnuts, almonds or brazil nuts.

Food products as described herein are shown to have an unexpected effect on levels of blood cholesterol, low density lipoprotein, triglycerides and/or other lipids or lipid particles, such as LDL particles, in an individual. Given chocolate is perceived as a high fat food, that was unexpected.

Aspects of the invention provide a food product as described above for use in reducing blood levels of cholesterol, low density lipoprotein, triglycerides and/or other lipids or lipid particles, such as LDL particles, in an individual and a method of reducing blood levels of cholesterol, low density lipoprotein, triglycerides and/or other lipids or lipid particles, such as LDL particles, in an individual comprising administering a food product described above to the individual.

Another aspect of the invention provides the use of a carotenoid compound and one or more cocoa bean products, as described above, in the manufacture of a food product for use in reducing blood levels of cholesterol, low density lipoprotein, triglycerides and/or other lipids or lipid particles, such as LDL particles, in an individual.

This may be useful in the treatment or prevention of cardio- and cerebro-vascular disorders, or Metabolic Syndrome, high blood pressure, pre-diabetes and type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30) and hypercholesterolaemia. The invention may be employed, for instance, with any of those subjects. The products of the invention may be used in dieting.

The invention also provides a method of dieting comprising consuming a product of the invention as part of the diet.

An individual is preferably a human, though use in animals is also possible. The individual may have normal blood levels of cholesterol, LDL and/or triglycerides or elevated blood levels of cholesterol, LDL and/or triglycerides. In some instances, the subject may have a total serum cholesterol of more than 200mg/dL, for instance more than 210mg/dL. In some cases a subject may additionally, or alternatively have, triglyceride levels above 150mg/dL. In some cases, the subject may be apparently healthy, but be identified as having such elevated levels of cholesterol and/or triglycerides, in other instances the subject may have a history of heart disease and/or atherosclerosis. The subject may be overweight and may be obese. The subject may be one taking statins, aspirin and/or blood pressure reducing medication. The subject may be one on a diet.

Methods of measuring levels of cholesterol, LDL, triglycerides and other lipids in an individual are well-known in the art.

In some embodiments, the individual may be at suffering from, or at risk of suffering from, a cardio- or cerebro-vascular disorders, such as coronary heart disease, metabolic syndrome, high blood pressure, pre-diabetes and type II diabetes, being overweight (e.g. BMI>25) or obesity (e.g. BMI>30). The subject may have had a heart attack. The subject may have had a stroke.

Food products as described herein are also shown to reduce levels of markers of inflammatory oxidative damage in an individual. In some cases the subject may have elevated levels of inflammatory oxidative damage. For instance, they may have ≥ 20 -39 μ M MDA and/or at least 0.25 to 0.30u/ml of P_x-1gG. Such levels may be in addition to, or alternative to, the above specified levels of total cholesterol and/or triglycerides.

The food products may therefore also be useful in reducing inflammation; reducing anti-inflammatory oxidative damage; increasing antioxidant activity and/or reducing or delaying symptoms of aging in an individual. The invention may be used to
5 reduce the visible signs of aging.

Examples of possible daily doses of 0.1mg to 100mg of carotenoid compound, such as lycopene, preferably 0.5 to 50 mg, may be administered to the individual. Any of the amounts referred to
10 herein may be administered.

In some embodiments, a suitable individual may be a mature or elderly individual, for example at least 50, 60, 65, 70, 75 or more years old or be of an age in the range defined by any of
15 those two values.

Food products as described herein may also be useful in providing nutrition to an individual.

20 For example, food products may be useful as sports nutrition products or in providing nutrition to mature or elderly individuals (e.g. >50 years old) or individuals undergoing body mass control or reduction, i.e. for "slimming" purposes.

25 In other examples, food products may be useful in providing nutrition to individual having or recovering from a clinical condition. For example, food products described herein may be useful in the nutrition of an individual recovering from injury, operation, or trauma; an individual having or recovering from
30 chemo- or radio-therapy; or an individual having or at risk of Metabolic Syndrome, obesity, diabetes II, atherosclerosis and their clinical complications.

The invention may be used to help treat ischemia or hypoxia. The
35 invention may be, in some instances, administered after blood flow has been cut off to a particular tissue or organ. In one instance,

the invention may be administered to subjects who have had a stroke.

Food products as described herein may also be useful in the
5 treatment or prevention of cardio- and cerebro-vascular disorders, hypertension, metabolic syndrome, high blood pressure, pre-diabetes and type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30) or other medical conditions such as anaemia, rheumatism, rheumatoid arthritis, non-rheumatoid arthritis,
10 prostate or testes malfunctions, erectile dysfunctions, loss of libido, cellulite, sarcopenia and cachexia.

In some instances, the subject the invention is applied to may have an auto-immune disease; an allergic condition; hypertension;
15 atherosclerosis; cardio pathologies, such as Coronary Heart Disease; vascular pathologies, such as endocarditis, myocarditis, heart failure, heart valve disease, arrhythmias, atherosclerosis, hypertension, vasculitis, endarteritis, varicose veins, endophlebitis, endothelial damage; cerebral pathologies; obesity;
20 diabetes type 2; cancer, sarcopenia; metabolic dysfunction; Metabolic Syndrome; cellulite and aging tissue degradation; gastritis; stomach or duodenum ulcers; or arthritis; or dermatitis, psoriasis, acne, chronic skin ulcerations, or other age-related or not skin conditions, including skin and other
25 tissues burns and wounds; sport, trauma, operation and other injuries; cachexia, side-effects of chemotherapies and radiation treatment, or radiation exposure; the subject may be at risk of such a condition.

30 Due to the ability of the invention to increase oxygen transport, the invention may also be used to treat conditions where such increased oxygen transport may be beneficial. For instance, a subject with a respiratory disorder such as emphysema, COPD, cystic fibrosis, asthma, or ARDS. The subject may have reduced
35 lung function, for instance due to lung damage or lung cancer. In one instance, the subject may be a smoker.

The invention may also be used to treat impairment of tissue oxygenation, for instance due to reduction of blood supply due to circulatory dysfunction or circulatory disease. The subject may have had an injury, disease or disorder causing reduced blood
5 flow, for instance one that results from blood flow to an organ and/or tissue being reduced or cut-off.

The invention may be used to increase tissue oxygenation and treat circulatory disease. In one instance the circulatory disorder may
10 be due to traumatic, compressive, occlusive, tumors/malformations and/or vasospastic reduction in oxygenation. The subject may have atherosclerosis resulting in reduced tissue oxygenation or DVT. The subject may be one with angina, such as angina pectoris, acute coronary syndrome, or had a myocardial infraction. The invention
15 may also be used to treat individuals with reduced tissue inflammation due to ongoing inflammatory conditions or processes in the tissue, such as any of those referred to herein.

Given the ability of the invention to reduce inflammatory markers,
20 the invention may also be employed to help treat inflammatory or autoimmune disorders, for instance arthritis, inflammatory bowel disease and atherosclerosis.

Another aspect of the invention provides a nutracosmetic
25 formulation comprising one or more cocoa bean products and a carotenoid compound.

Suitable cocoa bean products and carotenoid compounds are described in more detail above.

30

A nutracosmetic formulation which comprises one or more cocoa bean products and a carotenoid compound as defined above, may further comprise one or more cosmetically or nutritionally acceptable carriers, adjuvants, excipients, sweeteners, diluents, fillers,
35 buffers, stabilisers, preservatives, colourings, lubricants, or other materials well known to those skilled in the art.

The term "nutraceutically acceptable" as used herein pertains to compounds, materials, compositions, and/or dosage forms which are in common or widespread usage in food and dietary products and are generally considered non-toxic, for example, compounds may have
5 the US FDA designation "GRAS" (Generally Recognised as Safe), or equivalent food additive status in other jurisdictions.

Nutracosmetic formulations are generally intended for oral administration and may be formulated accordingly.

10

Nutracosmetic formulations may be useful in improving the appearance of an individual or in reducing, delaying or masking visual signs of aging in an individual.

15 The invention may be administered to treat, ameliorate, prevent, or reduce the severity of symptoms in any of the conditions referred to herein. In one instance, the invention is administered prophylactically to help prevent the onset of any of the conditions mentioned herein. The invention may result in reduction of any of
20 the parameters discussed herein, it may, for instance, reduce cholesterol, triglyceride, inflammatory damage, weight or body fat.

Various further aspects and embodiments of the present invention
25 will be apparent to those skilled in the art in view of the present disclosure.

All documents mentioned in this specification are incorporated herein by reference in their entirety.

30

The term "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. For example "A and/or B" is to be taken as specific disclosure of each of (i) A, (ii) B and (iii) A
35 and B, just as if each is set out individually herein.

In instances herein where the terms "*comprises*" or "*comprising*" are used, the invention may also provide what is described when it "*consists essentially of*" or "*consisting of*" the specified constituents.

5

Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the invention and apply equally to all aspects and embodiments which are described.

10

The following is a list of some further numbered embodiments of the invention:

(1) A food product comprising one or more cocoa bean products and an isolated carotenoid compound.

(2) A food product according to (1) which comprises a homogenous matrix containing the cocoa-bean products and the carotenoid compound.

(3) A food product according to (1) or (2) wherein the cocoa bean products comprise one or more of cocoa solids, cocoa powder, cocoa liquor and/or cocoa butter.

(4) A food product according to any of (1) to (3) where the one or more cocoa bean products are in the form of a chocolate or cocoa butter matrix, said matrix incorporating the carotenoid compound.

(5) A food product according to any one (1) to (4) which comprises 0.001 to 10mg of carotenoid compound per gram of food product.

(6) A food product according to any one of (1) to (5) the wherein the carotenoid compound is a lycopene compound.

(7) A food product according to any one of (1) to (6) wherein the carotenoid compound is comprised in a carotenoid-rich product, such as tomato or other fruit, vegetable or plant paste, sauce, concentrate, oleoresin, fraction or extract.

(8) A food product according to any one of (1) to (6) wherein the carotenoid compound is comprised in a carotenoid rich fruit, vegetable or other plant, or fungus, algae or bacterium.

(9) A food product according to any one of (1) to (8) wherein the lycopene compound is lycopene.

(10) A food product according to any one of (1) to (8) wherein the food product is produced by admixing or blending together the
5 cocoa bean products, the carotenoid compound and optionally one or more additional ingredients.

(11) A food product according to (10) wherein the cocoa bean products are admixed or blended together with the carotenoid compound in a dry, liquid, aerosol, frozen or melted form.

10 (12) A food product according to any one of (1) to (11) wherein the food product is a foodstuff.

(13) A food product according to (12) wherein the foodstuff is bread, flour, cereal, biscuit, pastry, spread, filling, paste, sauce, mousse, cream, or yogurt.

15 (14) A food product according to (12) wherein the foodstuff is a confectionary product.

(15) A food product according to (14) wherein the foodstuff is chocolate.

20 (16) A food product according to (15) wherein the chocolate is dark, milk, or white chocolate.

(17) A food product according to any one of (1) to (11) wherein the food product is a beverage.

(18) A food product according to any one of (1) to (11) wherein the food product is a dietary supplement, nutracosmetic or
25 nutraceutical product.

(19) A food product according to any one of (1) to (18)_ for use in reducing levels of cholesterol, LDL and/or triglyceride in an individual.

30 (20) A food product according to any one of (1) to (19) for use in reducing inflammation; reducing anti-inflammatory oxidative damage; increasing antioxidant activity and/or reducing or delaying symptoms of aging in an individual.

(21) A food product according to any one of (1) to (20) for use in the nutrition of an individual.

35 (22) A food product for use according to (21) wherein the individual is mature or elderly.

(23) A food product for use according to any one of (19) to (22) wherein the individual is undergoing body mass control or body mass reduction.

(24) A food product for use according to any one of (19) to (22) wherein the individual is suffering from; at risk of suffering from; or recovering from a clinical condition.

(25) A food product for use according to (24) wherein the individual is recovering from injury, operation, or trauma or undergoing or recovering from chemo- or radio-therapy; or having or being at risk of having Metabolic Syndrome, obesity, diabetes II, atherosclerosis and clinical complications thereof.

(26) A food product for use according to any one of (1) to (20) for the treatment of a clinical condition.

(27) A food product for use according to (26) wherein the clinical condition is a cerebro-vascular disorder, cardio-vascular disorder, hypertension, metabolic syndrome, high blood pressure, pre-diabetes, type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30), anaemia, rheumatism, rheumatoid arthritis, non-rheumatoid arthritis, prostate or testes malfunction, erectile dysfunction, loss of libido, cellulite, sarcopenia and cachexia.

(28) A method of improving the appearance of an individual comprising administering a nutracosmetic formulation according to any one of (1) to (18) to the individual.

(29) A method of reducing or delaying visible signs of aging in an individual comprising administering a nutracosmetic formulation according to any one of (1) to (18) to the individual.

(30) A method of reducing levels of cholesterol, LDL and/or triglyceride in the blood of an individual comprising administering a food product according to any one of (1) to (18) to an individual in need thereof.

(31) A method of reducing inflammation; reducing anti-inflammatory oxidative damage; increasing antioxidant activity and/or reducing or delaying symptoms of aging in an individual; comprising administering a food product according to any one of (1) to (18) to an individual in need thereof.

(32) A method of providing nutrition to an individual comprising administering a food product according to any one of (1) to (18) to an individual in need thereof.

(33) A method according to (32) wherein the individual is mature
5 or elderly.

(34) A method according to any one of (30) to (33) wherein the individual is undergoing body mass control or body mass reduction.

(35) A method according to any one of (30) to (33) wherein the individual is suffering from; at risk of suffering from; or
10 recovering from a clinical condition.

(36) A method according to (35) wherein the individual is recovering from injury, operation, or trauma or undergoing or recovering from chemo- or radio-therapy; or having or being at risk of having Metabolic Syndrome, obesity, type II diabetes, atherosclerosis and clinical complications thereof.
15

(37) A method of treatment of a clinical condition comprising administering a food product according to any one of (1) to (18) to an individual in need thereof.

(38) A method of treatment according to (37) wherein the clinical
20 condition is cerebro-vascular disorder, cardio-vascular disorder, hypertension, metabolic syndrome, high blood pressure, pre-diabetes, type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30), anaemia, rheumatism, rheumatoid arthritis, non-rheumatoid arthritis, prostate or testes malfunction, erectile
25 dysfunction, loss of libido, cellulite, sarcopenia or cachexia.

Certain aspects and embodiments of the invention will now be illustrated by way of example and with reference to the tables described below.

30

Table 1 shows the effect of lycopene on lipid parameters, and markers of IOD and inflammation in CHD patients.

Table 1 shows the effect of 30g dark chocolate on lipid
35 parameters, and markers of IOD and inflammation in CHD patients.

Table 3 shows the effect of 30 g of L-chocolate on lipid parameters, and markers of IOD and inflammation in CHD patients.

The results presented in the other Tables are discussed in the individual Examples below.

EXAMPLES

Experiments

10

Lipid-lowering chocolate (L-chocolate)

Commercially available dark chocolate (Green & Black's Dark Chocolate; 85% cocoa) was melted at 70°C. The melted chocolate was mixed with tomato oleoresin, containing 15% of lycopene (Lyc-O-Mato), in the ratio of 1.57 mg of oleoresin to 1 g of the chocolate. The mixture was blended for 10 minutes and then divided into daily 10 g or 30 g portions and cooled down to the room temperature.

Each 10 g or 30 g chocolate portion contained 47.1 mg of tomato oleoresin or about 7 mg of lycopene.

Control samples of chocolate

The melting and mixing procedures were performed as described above using the same commercially available dark chocolate, but instead of tomato oleoresin, sunflower oil (Floratm) was used.

Lycopene

47.1 mg of tomato oleoresin was pre-dissolved in ethanol and mixed with Whey Protein as described in Richelle *et al* (2002) *J Nutr* 132 404-408, WO01/091588 and US2002/01072992. Then the mixture was placed into gelatine capsules.

All products were kept in cool dry, protected from light conditions.

Validation in Clinical trials*CHD patients*

18 male CHD patients, age 47-69, were recruited for this study.

5 Main inclusion criteria were:

- elevated total serum cholesterol above 200 mg/dL and/or triglycerides above 150 mg/dL,
- all patients were naive for any lipid-lowering medications,
- stable clinical conditions and regimen of medications was t

10 the last 3 months.

Secondary inclusion criteria were:

- positive blood markers on inflammatory oxidative damage, IOD, > 20-30 μ M MDA
- 15 - positive blood on an antibody inflammatory marker, Px-IgG > 0.250-0.300 U/ml

All patients were randomised and divided into three equal groups of 6 patients each. Two groups receiving chocolate were blinded; 20 the group receiving lycopene preparation along was open labelled. The period of the trial was 4 weeks.

Results

The results of the ongoing trial are presented in the tables 1 to 25 3 below.

It was observed that after two weeks of administration of 7 mg of lycopene, there were no changes in any patients on their levels of elevated cholesterol, triglycerides and markers of oxidative 30 damage or inflammation (table 1).

Similar results were observed in the group where patients were taking 30 g of the control bar of dark chocolate (table 2).

35 However, in the group of patients taking 30 g of L-chocolate, a reduction in concentration of both total cholesterol and its LDL fraction was observed in every patient in the group even after the

first seven days of the trial (table 3). The positive trend in triglyceride levels was also detected in 5 out of 6 patients.

Furthermore, the elevated level of transferases in two patients in this group also started to decline, indicating a positive effect of L-chocolate on their liver damage status. In addition, for the majority of the patients administration of the L-chocolate was accompanied by reduction of inflammatory oxidative damage markers, indicating that this product has not only lipid-lowering properties but anti-inflammatory as well.

Baseline		Lycopene								
ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		$\mu\text{M MDA}$	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
13	48	101	0.765	225	161	39	153	6.5	44	25
14	69	162	0.698	231	150	42	159	5.6	45	36
15	54	79	0.811	204	134	41	135	3.8	34	24
16	49	95	0.803	219	126	44	161	4.4	27	35
17	66	83	0.751	243	165	37	186	5.9	49	29
18	53	49	0.743	210	157	40	147	6.1	25	26
56.5		95	0.762	222	149	40.5	157	5.4	37.3	29.2
Week 1										
ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		$\mu\text{M MDA}$	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
13	48	99	0.823	224	160	39	153	6.4	47	31
14	69	158	0.746	231	152	42	160	5.7	46	33
15	54	85	0.809	205	137	40	134	4.9	36	29
16	49	94	0.867	217	130	43	160	3.6	31	34
17	66	81	0.851	241	164	38	185	5.1	44	33
18	53	57	0.839	209	159	40	149	6.1	34	29
		96	0.823	221	150	40.3	157	5.3	39.7	31.5
Week 2										
ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		$\mu\text{M MDA}$	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
13	48	95	0.812	223	159	39	152	6.6	43	29
14	69	139	0.809	230	151	41	158	5.2	42	31
15	54	84	0.815	208	138	40	133	5.4	31	28
16	49	91	0.844	216	132	42	159	4.1	38	32
17	66	75	0.830	239	162	39	183	3.9	41	31
18	53	68	0.799	208	158	41	150	6	30	34
		92	0.818	221	150	40.3	156	5.2	37.5	30.8

Week 3

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
13	48	94	0.834	221	155	39	151	6.2	42	32
14	69	136	0.781	227	149	42	157	4.8	39	29
15	54	85	0.84	210	135	40	132	5.1	35	25
16	49	93	0.795	214	142	41	155	5	36	31
17	66	76	0.809	232	161	38	179	4.4	42	28
18	53	81	0.774	211	154	41	152	5.8	33	31
		94	0.806	219	149	40.2	154	5.22	37.8	29.3

Table 1

Baseline

Chocolate

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
7	49	132	0.902	209	165	41	157	5.6	35	25
8	55	96	0.933	232	183	40	153	4.2	42	36
9	51	145	0.998	198	182	45	124	6.6	28	24
10	62	53	0.756	227	144	39	179	5.9	41	35
11	50	61	0.854	217	136	41	166	3.8	33	29
12	53	110	0.941	221	179	38	148	5.8	37	26
53.3		99.5	0.897	217	165	40.7	155	5.3	36	29.2

Week 1

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
7	49	123	0.912	208	168	41	156	5.8	37	27
8	55	105	0.875	229	181	40	154	4.9	41	43
9	51	132	0.914	201	178	44	128	6.4	35	28
10	62	96	0.665	225	149	40	177	5.9	40	38
11	50	83	0.806	216	147	42	165	4.7	36	31
12	53	105	0.915	219	180	38	149	5.6	39	30
		107	0.848	216	167	40.8	155	5.5	38	32.3

Week 2

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		µM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
7	49	119	0.945	207	169	40	157	5.9	32	29
8	55	99	0.927	230	179	40	154	5.6	39	44
9	51	141	0.983	197	185	45	126	6.4	29	32
10	62	67	0.844	223	155	40	176	5.8	42	37
11	50	78	0.915	218	146	41	165	3.7	35	33
12	53	114	0.926	215	182	39	147	5.9	35	35
		103	0.923	215	169	40.8	154	5.55	35.3	35

Week 3

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		µM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
7	49	118	0.999	201	165	40	156	5.2	32	28
8	55	99	0.876	233	182	41	157	4.9	39	43
9	51	133	0.858	195	164	44	134	6.7	29	36
10	62	69	0.761	219	153	41	175	5.1	42	47
11	50	83	0.944	226	132	42	161	4.2	35	31
12	53	105	0.832	209	167	38	149	5	35	37
		101	0.878	214	160	41	155	5.2	35.3	37

Table 2

Baseline				Chocolate + Lycopene						
ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		µM MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
1	52	73	0.904	217	121	40	132	4.5	22	40
2	55	46	0.842	211	200	37	169	4.7	30	33
3	63	88	0.871	249	199	42	174	5.3	30	27
4	59	150	0.901	136	170	37	167	6.2	48	110
5	47	112	0.660	228	168	40	150	4.6	40	45
6	49	123	0.789	227	113	42	130	5.5	120	154
54		98.7	0.827	211	162	39.7	154	5.1	48.7	68.2

Week 1

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μ M MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
1	52	76	0.943	195	120	40	130	4.2	22	40
2	55	38	0.912	183	200	37	167	4.8	30	32
3	63	69	0.838	233	179	42	170	5	30	26
4	59	143	0.522	132	162	37	162	6.1	49	73
5	47	97	0.720	208	161	40	149	4.4	38	42
6	49	101	0.324	193	99	42	127	5	74	137
		87.3	.710	191	153	39.7	151	4.9	40.5	58.3

Week 2

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μ M MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
1	52	44	0.452	193	119	40	129	4.3	22	40
2	55	19	0.81	183	202	37	167	4.7	27	36
3	63	58	0.448	223	178	42	170	5.1	30	28
4	59	63	0.522	130	161	37	160	6.2	45	70
5	47	88	0.23	200	159	40	147	4.5	38	41
6	49	89	0.214	191	89	42	125	5.8	70	132
		60.2	0.446	187	151	39.7	150	5.1	38.7	57.8

Week 3

ID	Age	IOD	Px-IgG	TC	TG	HDL	LDL	GL	AST	ALT
		μ M MDA	U/ml	mg/dL	mg/dL	mg/dL	mg/dL	mmol/L	U/L	U/L
1	52	53	0.129	193	119	40	129	4.3	22	40
2	55	21	0.742	182	188	38	160	4.5	27	35
3	63	41	0.081	225	171	42	170	5.2	30	27
4	59	50	0.096	130	158	38	159	6.1	44	70
5	47	61	0	189	154	40	137	5.0	38	39
6	49	54	0.120	190	84	43	122	5.5	72	130
		46.7	0.195	185	146	40.1	146	5.1	38.1	56.8

Table 3

Clinically healthy volunteers with hypercholesterolaemia*Dose effect*

5 52 clinically healthy volunteers 26 males and 26 females, age 35-61 years old, were recruited for this study.

Main inclusion criteria were:

elevated total serum cholesterol above 200 mg/dL and/or triglycerides above 150 mg/dL,

10 all patients were naive, for at least 3 months prior to the study, for any lipid-lowering medications, dietary supplements or special lipid or weight management diets, willing to participate in the study.

15 All volunteers were randomised and divided into six groups. 10 participants were included in the control group and group which received chocolate containing 0.36 mg of lycopene per 1 gram of the product. Other four groups were formed from 8 volunteers each. Every participant received the one week supply of the same size of
20 10 g of the chocolate bar with different concentration of lycopene, or without it at all.

All chocolate samples were blinded so participants did not know what exactly composition of chocolate they were ingesting. All
25 collected blood samples were also blinded so the analytical laboratory was not aware from which volunteers and from which group the samples were analysed. Every week participants were invited to the clinic when compliance of the ingesting chocolate was verified, blood from these persons was collected, and new
30 batch of one week supply of chocolate was given.

The duration of the trial was 4 weeks. Effects of the following concentrations of lycopene, "L- tug", in the chocolate was studied: 0.0, 0.1, 0.2, 0.3, 0.35, 0.7 mg of lycopene per 1 g of
35 chocolate.

Results

Cholesterol

The effects of chocolate with different concentration of lycopene
5 on the elevated level of the total cholesterol, and other
biochemical parameters of the participants are presented in table
4a and table 5.

These results showed that the chocolate with 0.1 mg of lycopene
10 per 1 g of the product was already able to reduce elevated total
cholesterol although the changes were not statistically
significant. However, chocolate with 0.2 mg of lycopene and
above, per 1 g of the product, were consistently producing
cholesterol lowering effect in the serum of the participants. The
15 significant effect was already registered from the second week of
the trial and reached it maximum by the fourth week.

Biochemistry

The effect of this lycopene / L-tug chocolate on other biochemical
20 parameters was insignificant for the doses studied (table 5).
Presumably this was because in most groups these parameters were
within their physiological norms and there were no much room for
their normalisation / "improvement".

25 *Inflammation and oxidation*

However, the majority of the participants were positive on markers
of Inflammatory Oxidative Damage, IOD, or in some cases on
presence of such inflammatory markers as LDL-Px and Chl.-IgG. This
was probably due that the majority of the participants were
30 between 50 and 60 years old, and these markers can frequently be
detected on a subclinical level even in apparently healthy people
of this age and above.

It all groups taking L-tug chocolate the reduction of the IOD was
35 significant. However, reduction of two other inflammatory markers
was observed only in some groups and was not apparently dose

dependent. This inconsistency could be a result of a small number of participants in the tested groups.

Plasma Oxygen transport

5 The other interest observation was that ingestion of the L-tug chocolate resulted in the dose-dependent increase of the plasma oxygen transport. This useful property could be used to increase and / or restore supply of the molecular O₂ depression or reduction of which may occur not only in many clinical conditions but also
10 during strenuous exercises, or with ageing.

Timing of the ingestion of chocolate in relation to food intake.

To evaluate a possible hypothesis that incorporation of lycopene
15 into chylomicrons and lipoproteins, during their re-assembly at the time of digesting food fat, we undertake the following study. We recruited a group of clinically healthy volunteers of similar age and similar level of hypercholesterolaemia.

The design of the study was the same as the study above but
20 instead of taking L-tug chocolate with main food we asked participants to ingest the chocolate between meals - at least two hours after their breakfast, or lunch, and at least two hours before their next meal - lunch or dinner.

25 The product used was the same format of 10 g containing 0.7 mg of lycopene per 1 g of chocolate.

Results

The results of this study are presented in the table 4b.
30

It was observed that this regiment was also able to reduce elevated total cholesterol but the significant reduction was only observed on the week 3 of the trial. The maximum of the reduction was on the last 4th week.

35

It was interesting to note, that although the cholesterol-lowering effect was prominent it was still significantly lower than when the same chocolate was taken during the main meal.

If the week 4 is taken as a reference point, the 0.7 mg dose of the L-tug chocolate ingested on the "empty stomach" was more effective than the dose 0.1 but less effective than dose 0.2 mg when they were taken with food.

Table 4a Dose dependency - chocolate ingestion with main food

L-tug, per 1 g chocolate	Total serum cholesterol, mg/dL				
	0 w	1 w	2 w	3 w	4 w
0.0 mg	217 + 4.2	215 + 5.6 $\Delta = -2, p^*$	215 + 5.1 $\Delta = -2, p^*$	214 + 4.9 $\Delta = -3, p^*$	213 + 5.3 $\Delta = -4, p^*$
0.1 mg	229 + 3.0	228 + 6.5 $\Delta = -1, p^*$	224 + 3.8 $\Delta = -5, p^*$	224 + 2.8 $\Delta = -5, p^*$	218 + 1.5 $\Delta = -11, p = 0.01$
0.2 mg	228 + 4.1	216 + 8.3 $\Delta = -12, p^*$	199 + 2.0 $\Delta = -29, p < 0.001$	191 + 5.0 $\Delta = -37, p < 0.001$	191 + 5.3 $\Delta = -37, p < 0.001$
0.3 mg	226 + 6.0	202 + 9.3 $\Delta = -24, p^*$	198 + 6.8 $\Delta = -28, p < 0.05$	196 + 7.3 $\Delta = -30, p < 0.05$	192 + 4.3 $\Delta = -34, p = 0.002$
0.35 mg	221 + 2.8	215 + 3.0 $\Delta = -6, p^*$	209 + 2.6 $\Delta = -12, p < 0.05$	208 + 2.5 $\Delta = -13, p < 0.05$	195 + 8.5 $\Delta = -26, p = 0.02$
0.7 mg	241 + 14.0	231 + 14.8 $\Delta = -10, p^*$	190 + 7.0 $\Delta = -51, < 0.05$	188 + 6.3 $\Delta = -53, p = 0.01$	186 + 5.3 $\Delta = -55, p < 0.01$

Table 4b. The chocolate ingestion between meals two hours after and two hours before any food intake

0.7 mg	243 + 13.7	230 + 10.1 $\Delta = -13, p^*$ $\Delta_{F[+/-]} = -1$ $P_{F[+/-]} < 0.05$	218 + 4.3 $\Delta = -25, p < 0.05$ $\Delta_{F[+/-]} = +28$ $P_{F[+/-]} > 0.01$	210 + 5.1 $\Delta = -33, p < 0.05$ $\Delta_{F[+/-]} = +22$ $P_{F[+/-]} > 0.05$	208 + 5.8 $\Delta = -35, p < 0.05$ $\Delta_{F[+/-]} = +22$ $P_{F[+/-]} > 0.01$
--------	------------	--	---	---	---

$p_{F[+/-]}$ - difference between the same time points in groups which ingested L-tug chocolate with or without food

Table 5. Effect of different doses of L-tug Chocolate on biochemical parameters and markers of inflammation and oxidation

L-tug, per 1g chocolate	TG mg/dL	HDL mg/dL	LDL mg/dL	glucose	AST	ALT	CRP	LDL in μ M	LDL-Px EUSA $\times 10^3$	Cys-p-IgG EUSA $\times 10^3$	Plasma-O ₂
0	165 $\pm$ 12.8 162 $\pm$ 11.5 $\Delta=3$, p > 0.05	40 $\pm$ 0.9 40 $\pm$ 0.8 p > 0.05	155 $\pm$ 9.9 154 $\pm$ 8.7 p > 0.05	5.9 $\pm$ 0.7 5.8 $\pm$ 0.6 p > 0.05	32 $\pm$ 4.1 34 $\pm$ 3.9 p > 0.05	41 $\pm$ 5.6 40 $\pm$ 5.1 p > 0.05	5.5 $\pm$ 1.4 6.1 $\pm$ 1.2 p > 0.05	133 $\pm$ 11.4 134 $\pm$ 12.7 p > 0.05	354 $\pm$ 41 401 $\pm$ 38 p > 0.05	675 $\pm$ 55 722 $\pm$ 64 p > 0.05	0.812 $\pm$ 75 0.825 $\pm$ 66 $\Delta=11$, p > 0.05
0.1 mg	99 $\pm$ 3.8 88 $\pm$ 3.3 $\Delta=11$, p > 0.05	43 $\pm$ 1.8 44 $\pm$ 1.3 p > 0.05	131 $\pm$ 3.8 127 $\pm$ 4.0 $\Delta=4$, p > 0.05	5.4 $\pm$ 0.6 5.4 $\pm$ 0.3 p > 0.05	40 $\pm$ 3.8 36 $\pm$ 2.8 p > 0.05	34 $\pm$ 3.8 30 $\pm$ 2.7 p > 0.05	6.0 $\pm$ 0.9 5.4 $\pm$ 0.5 p > 0.05	115 $\pm$ 10.8 9 $\pm$ 4.6 p < 0.001	175 $\pm$ 33 212 $\pm$ 59 p > 0.05	577 $\pm$ 98 521 $\pm$ 67 p > 0.05	1.155 $\pm$ 83 1.263 $\pm$ 19 $\Delta=0.110$, p > 0.05
0.2 mg	146 $\pm$ 10.3 130 $\pm$ 8.5 $\Delta=26$, p < 0.05	44 $\pm$ 0.6 45 $\pm$ 0.5 p > 0.05	145 $\pm$ 9.0 135 $\pm$ 8.5 $\Delta=10$, p > 0.05	5.1 $\pm$ 0.5 6.1 $\pm$ 0.4 p > 0.05	29 $\pm$ 2.1 38 $\pm$ 1.6 p > 0.05	25 $\pm$ 3.0 25 $\pm$ 2.3 p > 0.05	5.1 $\pm$ 1.3 4.9 $\pm$ 1.2 p > 0.05	35 $\pm$ 7.3 19 $\pm$ 6.5 p < 0.001	142 $\pm$ 18 9 $\pm$ 1.8 p < 0.001	444 $\pm$ 22 260 $\pm$ 18 p < 0.001	0.811 $\pm$ 51 1.383 $\pm$ 37 $\Delta=0.572$, p < 0.001
0.3 mg	162 $\pm$ 10.7 143 $\pm$ 9.1 $\Delta=20$, p > 0.05	40 $\pm$ 1.0 41 $\pm$ 0.8 p > 0.05	154 $\pm$ 3.6 142 $\pm$ 1.8 $\Delta=10$, p < 0.05	4.9 $\pm$ 0.4 4.7 $\pm$ 0.2 p > 0.05	49 $\pm$ 8.6 36 $\pm$ 4.7 p > 0.05	60 $\pm$ 9.1 53 $\pm$ 10 p > 0.05	6.7 $\pm$ 1.5 5.5 $\pm$ 1.1 p > 0.05	159 $\pm$ 10.7 38 $\pm$ 4.1 p < 0.05	511 $\pm$ 97 0 $\pm$ 38 p < 0.05	828 $\pm$ 99 288 $\pm$ 49 p < 0.05	0.781 $\pm$ 37 0.963 $\pm$ 24 $\Delta=0.223$, p < 0.01
0.35 mg	136 $\pm$ 28.1 106 $\pm$ 16.5 $\Delta=30$, p > 0.05	43 $\pm$ 1.0 44 $\pm$ 1.5 p > 0.05	140 $\pm$ 10.0 132 $\pm$ 9.3 $\Delta=8$, p > 0.05	5.3 $\pm$ 0.4 5.5 $\pm$ 0.3 p > 0.05	41 $\pm$ 2.4 39 $\pm$ 2.3 p > 0.05	49 $\pm$ 4.0 45 $\pm$ 4.1 p > 0.05	5.4 $\pm$ 1.7 5.8 $\pm$ 1.6 p > 0.05	130 $\pm$ 30.8 53 $\pm$ 8.8 p < 0.05	160 $\pm$ 36 206 $\pm$ 47 p > 0.05	552 $\pm$ 41 506 $\pm$ 66 p > 0.05	0.988 $\pm$ 34 1.283 $\pm$ 19 $\Delta=0.275$, p < 0.001
0.7 mg	128 $\pm$ 10.8 108 $\pm$ 6.6 $\Delta=20$, p > 0.05	42 $\pm$ 0.6 43 $\pm$ 0.7 p > 0.05	153 $\pm$ 3.8 141 $\pm$ 3.0 $\Delta=8$, p > 0.05	5.4 $\pm$ 0.2 6.0 $\pm$ 0.4 p > 0.05	29 $\pm$ 2.5 39 $\pm$ 2.0 p > 0.05	29 $\pm$ 3.5 27 $\pm$ 1.8 p > 0.05	6.8 $\pm$ 0.8 6.6 $\pm$ 0.5 p > 0.05	35 $\pm$ 9.9 23 $\pm$ 7.8 p < 0.05	66 $\pm$ 12 3 $\pm$ 8.5 p < 0.05	negative negative	1.099 $\pm$ 87 1.438 $\pm$ 85 $\Delta=0.331$, p < 0.01

5

Postprandial study

8 clinically healthy volunteers 4 males and 4 females, age 35-60 years old, were recruited for this study.

10 Main inclusion criteria were:

- all patients were naive, for at least 3 months prior to the study, for any lipid-lowering medications, dietary supplements or special lipid or weight management diets,
- willing participate in the study.

15

All volunteers received standardise fat rich meal comprising of 50 g of butter as a part of a sandwich with 2 slices of white bread. Then without any break volunteers were asked to ingestion 10 g chocolate bar without lycopene. During intake of this test meal
20 volunteers was given 200 ml of warm decaffeinated tea with skimmed milk containing no more than 1% of dairy fat.

The blood was collected just before the intake of the meal and every hour for 4 hours after that.

25

After one week of break the same volunteers were ask to take exactly the same meal of 50 g of the same butter with two slices of the same white bread. Then each of them was asked to take 10 g of the chocolate bar with 7 mg of lycopene blended in at the protocol described above. The same type and the volume of tea was allowed, and the blood was collected at the same protocol as in the previous week.

The results of this cross-over study are presented in the table 6. They show that ingestion of lycopene-contained chocolate was able to reduce elevation of postprandial triglycerides about two fold and cholesterol between 2 and 3 times. It was also interesting to note that the postprandial glycaemia was also reduced by this L-tug chocolate.

The most remarkable changes were observed when this chocolate with the embedded lycopene was not only able to prevent increase of the inflammatory oxidative markers in the postprandial blood but even cause their reduction below the baseline level (table 6).

Postprandial time	Trial A (control) 50 g butter + 10 g chocolate (n = 8)					Trial B 50 g butter + 10 g L-tug chocolate (n = 8)				
	TC in mg/dL	TG in mg/dL	LDL in mg/dL	Glucose in mmol/l	ROS, pM MDA	TC in mg/dL	TG in mg/dL	LDL in mg/dL	Glucose in mmol/l	ROS, pM MDA
baseline	164	108	133	5.9	87	168	169	133	5.9	85
1 h	208 $\Delta = 24 \pm 5.7$	115 $\Delta = 11 \pm 3.9$	136 $\Delta = 3 \pm 0.2$	6.9 $\Delta = 1.0 \pm 0.3$	94 $\Delta = 7 \pm 0.7$	206 $\Delta = 18 \pm 4.5$ $p(A-B) < 0.05$	114 $\Delta = 15 \pm 2.9$ $p(A-B) < 0.05$	135 $\Delta = 3 \pm 0.2$ $p(A-B) < 0.05$	5.9 $\Delta = 0.9 \pm 0.2$ $p(A-B) < 0.05$	87 $\Delta = 1 \pm 0.5$ $p(A-B) < 0.05$
2 h	215 $\Delta = 33 \pm 5.2$	129 $\Delta = 21 \pm 3.3$	136 $\Delta = 3 \pm 0.2$	6.1 $\Delta = 1.3 \pm 0.1$	97 $\Delta = 10 \pm 0.6$	201 $\Delta = 14 \pm 4.7$ $p(A-B) < 0.01$	126 $\Delta = 11 \pm 2.8$ $p(A-B) < 0.05$	135 $\Delta = 2 \pm 0.1$ $p(A-B) < 0.05$	5.7 $\Delta = 0.2 \pm 0.1$ $p(A-B) < 0.05$	88 $\Delta = 1 \pm 0.6$ $p(A-B) < 0.05$
3 h	204 $\Delta = 29 \pm 5.3$	120 $\Delta = 12 \pm 3.1$	135 $\Delta = 2 \pm 0.1$	5.6 $\Delta = 0.6 \pm 0.2$	97 $\Delta = 10 \pm 0.6$	184 $\Delta = 6 \pm 4.3$ $p(A-B) < 0.05$	114 $\Delta = 5 \pm 2.4$ $p(A-B) < 0.05$	133 $\Delta = 0 \pm 0.1$ $p(A-B) < 0.05$	5.4 $\Delta = 0.4 \pm 0.1$ $p(A-B) < 0.05$	72 $\Delta = -14 \pm 1.3$ $p(A-B) < 0.01$
4 h	193 $\Delta = 9 \pm 4.4$	110 $\Delta = 2 \pm 3.4$	134 $\Delta = 1 \pm 0.2$	5.3 $\Delta = 0.5 \pm 0.2$	99 $\Delta = 12 \pm 1.3$	195 $\Delta = 7 \pm 4.2$ $p(A-B) < 0.05$	110 $\Delta = 1 \pm 2.7$ $p(A-B) < 0.05$	133 $\Delta = 0 \pm 0.2$ $p(A-B) < 0.05$	5.1 $\Delta = 0.1 \pm 0.2$ $p(A-B) < 0.05$	70 $\Delta = -16 \pm 1.1$ $p(A-B) < 0.01$

TC - total cholesterol, TG - triglycerides,
 Δ - changes in mean concentrations with the baseline,
 $p(A - B)$ - statistical differences in same parameters at the same time points between trial A and trial B.

8 clinically healthy volunteers, 4 men and 4 women, average age - 35 - 60 years old

Statistically significant between
control and L-tug trials

Table 6. Effect of L-tug chocolate on Postprandial Biochemistry
Profile and Markers of Inflammatory Oxidation
5 in Serum of Healthy Volunteers - Cross-over Study

**New opportunities to control lipid metabolism, inflammation and
tissue oxygenation**

10 The results presented here open a possible new mechanisms and new
ways not only to control already developed changes in the lipid
metabolism but also to prevent these changes.

These results also provide for the development of new ways to
15 control subclinical and other forms of inflammation and / or
boosting transport of the plasma molecular oxygen, which could be
useful to restore tissue oxygen saturation which could be
important in many clinical conditions and to delay ageing.

20 The results described herein are unexpected because the benefit of
adding any ingredient with additional health value to a chocolate
product would be expected to be outweighed by the potential
harmful consequences of consuming increased amounts of this high-
fat food product.

25 Reducing fat content in food products is the standard way to
minimise their fat load to the body. However this approach is not
generally useful for cocoa-based products, such as chocolate,
because fat reduction negatively affects the melting, feeling and
30 taste properties.

The results described herein show the unusual and unexpected
outcome of the blending of carotenoids such as lycopene with
cocoa-based products. Not only is cocoa butter prevented from

contributing to the rise of blood lipids, but the blend actively reduces lipids which are already at an elevated level.

In other words, the invention described herein not only makes
5 cocoa products, such as chocolate, "safer" from the health impact point of view, but may also make it useful as a proactive interventional product for slimming, lipid-lowering purposes and anti-aging purposes, and for prevention and help in management of metabolic, pre-diabetes, cardio-vascular and other conditions.

CLAIMS

1. A food product comprising one or more cocoa bean products and a carotenoid compound.

5

2. A food product according to claim 1, where:

(a) the food product comprises the carotenoid compound provided as an isolated carotenoid compound or a carotenoid rich product;

10 (b) the food product comprises a homogenous matrix containing the cocoa-bean products and the carotenoid compound;

(c) the cocoa bean products comprise one or more of cocoa solids, cocoa powder, cocoa liquor and/or cocoa butter;

15 (d) the one or more cocoa bean products are in the form of a chocolate or cocoa butter matrix, said matrix incorporating the carotenoid compound; and/or

(e) the food-product comprises 0.001 to 10mg of carotenoid compound per gram of food product.

20 3. A food product according to any one of the preceding claims wherein the carotenoid compound is, or comprises, a lycopene compound, preferably lycopene.

25 4. A food product according to any one of the preceding claims where:

(a) the food product is a foodstuff;

30 (b) the carotenoid compound is provided as a carotenoid rich product selected from a tomato paste, sauce, concentrate, oleoresin, fractions, extracts, or synthetic or purified carotenoid molecules;

(c) the carotenoid compound is provided as a carotenoid rich product from suitable fruit or vegetable or other plants, or fungi, yeast, algae, or bacteria, or as synthetic or purified carotenoid molecules

35 (d) the food product is produced by admixing or blending together the cocoa bean products, the carotenoid compound and optionally one or more additional ingredients; and/or

- (e) the cocoa bean products are admixed, or blended, or sonicated, or sprayed-dried, or embedded by other means together with the carotenoid compound in a dry, liquid, aerosol, frozen or melted form.

5

5. A food product according to any one the preceding claims, wherein:

- (a) the foodstuff is bread, flour, cereal, biscuit, pastry, spread, filling, paste, sauce, mousse, cream, ice cream, cake or yogurt;
- (b) the foodstuff is a confectionary product;
- (c) the foodstuff is, or comprises, chocolate, preferably where the chocolate is dark, milk, or white chocolate; or
- (d) the food product is a beverage, preferably a chocolate, or cacao, or milkshake or liquid supplement.

10

15

6. A food product according to any one of the preceding claims wherein the food product is a dietary supplement, nutraceutical or nutraceutical product.

20

7. A food product according to any one of the preceding claims for use in:

- (a) reducing elevated levels of cholesterol, LDL and/or triglyceride in an individual, preferably where the individual has elevated levels of cholesterol, LDL and/or triglyceride;
- (b) reducing subclinical or clinical inflammation; reducing anti-inflammatory oxidative damage; increasing plasma molecular oxygen transport, microcirculation and tissue oxygen saturation, reducing already developed liver (micro-) damage and liver steatosis, liver and other organs, including peripheral, tissue hypoxia or ischaemia, increasing antioxidant activity and/or reducing or delaying symptoms of ageing in an individual;
- (c) reducing postprandial cholesterol- and triglyceride-aemias, reducing size of chylomicrons and increasing rate of their clearance, reducing postprandial inflammatory and oxidative

25

30

35

- stress, reducing postprandial or other liver (micro-) damage and liver steatosis, liver and other organs, including peripheral tissue hypoxia or ischaemia, or delaying of above mentioned symptoms of fat, or excessive, or imbalance food intake in an individual;
- 5 (d) increasing oxygen transport in a subject, preferably where the subject has a respiratory disorder and/or lung damage; and/or strenuous physical or mental performance; and/or muscle wasting conditions; and/or
- 10 (e) slimming, weight reduction or dieting.
8. A food product according to any one claims 1 to 6 for use in the nutrition of an individual, preferably for use in the nutrition of the individual to treat one of the conditions
- 15 referred to in claim 7.
9. A food product for use according to claim 7 or 8 where:
- (a) the individual is mature or elderly;
- (b) the individual is undergoing body mass control or body mass
- 20 reduction;
- (c) the individual is recovering from injury, operation, or trauma or undergoing or recovering from chemo- or radio-therapy; or having or being at risk of having Metabolic Syndrome, obesity, diabetes II, atherosclerosis and clinical
- 25 complications thereof;
- (d) the food product is for using in treating a clinical condition, preferably where the clinical condition is a cerebro-vascular disorder, cardio-vascular disorder, hypertension, metabolic syndrome, high blood pressure, pre-
- 30 diabetes, type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30), anaemia, rheumatism, rheumatoid arthritis, non-rheumatoid arthritis, prostate or testes malfunction, erectile dysfunction, loss of libido, cellulite, age-related functions of liver, skin and other organs,
- 35 sarcopenia and cachexia;and/or
- (e) the food product is consumed during a mealtime.

10. A method of:

(a) improving the appearance and performance of an individual comprising administering a nutracosmetic formulation according to any one of claims 1 to 6 to the individual;
5 and/or

(b) reducing or delaying signs of aging in an individual, preferably visible signs of aging, comprising administering a food product according to any one of claims 1 to 6 to the individual.

10

11. A method of:

(a) reducing elevated levels of cholesterol, LDL and/or triglyceride in the blood of an individual comprising administering a food product according to any one of claims 1
15 to 6 to an individual in need thereof;

(b) reducing subclinical or clinical inflammation; reducing anti-inflammatory oxidative damage; increasing plasma molecular oxygen transport, microcirculation and tissue oxygen saturation, reducing already developed liver (micro-) damage
20 and liver steatosis, liver and other organs, including peripheral, tissue hypoxia or ischaemia; increasing antioxidant activity and/or reducing or delaying symptoms of ageing in an individual; comprising administering a food product according to any one of claims 1 to 6 to an
25 individual in need thereof;

(c) reducing postprandial cholesterol- and triglyceride- aemias, reducing size of chylomicrons and increasing rate of their clearance, reducing postprandial inflammatory and oxidative stress, reducing postprandial or other liver (micro-) damage
30 and liver steatosis, liver and other organs, including peripheral tissue hypoxia or ischaemia, or delaying of above mentioned symptoms of fat, or excessive, or imbalance food intake in an individual; comprising administering a food product according to any one of claims 1 to 6 to an
35 individual in need thereof;

(d) providing nutrition to an individual comprising administering a food product according to any one of claims 1

to 6 to an individual in need thereof; and/or strenuous physical or mental performance; and/or muscle wasting conditions; and/or

- (e) slimming, weight reduction or dieting comprising administering a food product according to any one of claims 1 to 6.

12. The method of claim 11 wherein:

- (a) the individual is mature or elderly;
- 10 (b) the individual is undergoing body mass control or body mass reduction;
- (c) the individual is recovering from injury, operation, or trauma or undergoing or recovering from chemo- or radio-therapy; or having or being at risk of having Metabolic Syndrome, obesity, diabetes II, atherosclerosis and clinical complications thereof;
- 15 (d) the food product is for using in treating a clinical condition, preferably where the clinical condition is a cerebro-vascular disorder, cardio-vascular disorder, hypertension, metabolic syndrome, high blood pressure, pre-diabetes, type II diabetes, being overweight (e.g. BMI>25), obesity (e.g. BMI>30), anaemia, rheumatism, rheumatoid arthritis, non-rheumatoid arthritis, prostate or testes malfunction, erectile dysfunction, loss of libido, cellulite, age-related functions of liver, skin and other organs, sarcopenia and cachexia; and/or
- 20 (e) the food product is consumed during a mealtime.

13. A method of producing a food product which comprises adding a carotenoid compound during production of the food product, so producing a food product as defined in any one of claims 1 to 6.

14. A chocolate bar comprising a carotenoid, preferably where the carotenoid is a lycopene compound.

35

15 A chocolate bar according to claim 14 which comprises at least 0.2 mg of the carotenoid.