

(19) **DANMARK**

(10) **DK/EP 2816111 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 15/10 (2006.01)** **C 12 Q 1/68 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-06-06**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-04-13**
- (86) Europæisk ansøgning nr.: **14181813.8**
- (86) Europæisk indleveringsdag: **2010-08-13**
- (87) Den europæiske ansøgnings publiceringsdag: **2014-12-24**
- (30) Prioritet: **2009-08-14 US 234044 P**
- (62) Stamansøgningsnr: **10808806.3**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Epicentre Technologies Corporation, 726 Post Road, Madison, WI 53713, USA**
- (72) Opfinder: **SOOKNANAN, Roy R., 322 James-Shaw, Beaconsfield, Québec H9W 6G5, Canada**
- (74) Fuldmægtig i Danmark: **Hegner & Partners A/S, Banemarksvej 50, 2605 Brøndby, Danmark**
- (54) Benævnelse: **Fremgangsmåder, sammensætninger og kits til generering af rRNA-depleterede prøver eller isolering af rRNA fra prøver**
- (56) Fremdragne publikationer:
WO-A2-2007/019444
**Darshini Mehta: "RiboMinus Transcriptome Isolation Kit (Human/Mouse)", Invitrogen User Manual , 12 September 2005 (2005-09-12), XP002688465, Retrieved from the Internet:
URL: http://tools.invitrogen.com/content/sf/s/manuals/ribominus_human_mouse_man.pdf [retrieved on 2012-12-03]**
SU CHUNLEI ET AL: "A simple method to enrich mRNA from total prokaryotic RNA", MOLECULAR BIOTECHNOLOGY, vol. 10, no. 1, August 1998 (1998-08), pages 83-85, XP008158531, ISSN: 1073-6085

DESCRIPTION

[0001] The present application claims priority to U.S. Provisional Application Serial Number 61/234,044, filed August 14, 2009.

FIELD OF THE INVENTION

[0002] The present invention relates to methods, compositions, and kits for generating rRNA-depleted samples and for isolating rRNA from samples. In particular, the present invention provides compositions comprising affinity-tagged antisense rRNA molecules that exhibit sequences complementary to substantially all of at least one full-length rRNA molecule encoded by a rRNA gene (e.g., compositions comprising affinity-tagged antisense rRNA molecules that exhibit sequences which, either alone or in combination, are complementary to substantially all or the complete sequence of at least one rRNA molecule selected from among 28S, 26S, 25S, 18S, 5.8S and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S and 5S prokaryotic rRNA molecules) and methods for using such compositions to generate rRNA-depleted samples or to isolate rRNA from samples.

BACKGROUND OF THE INVENTION

[0003] Since rRNA comprises about 95% to about 98% of the RNA in a cell, its presence can complicate various types of analyses of other RNA molecules of interest in a sample (e.g., gene expression analyses by arrays or microarrays, next-generation sequencing of tagged cDNA molecules made from one or more types of RNA molecules in samples (e.g., using the massively parallel digital sequencing methods referred to as "RNA-seq"), etc.). The problems caused by rRNA are especially difficult for analyses of RNA molecules of interest that are fragmented. For example, a considerable and continuing problem in the art is to find better methods for removing degraded rRNA from formalin-fixed paraffin-embedded (FFPE) tissue sections. If better methods were available to remove degraded rRNA from samples (e.g., FFPE-derived samples), it is believed that the enormous quantities of clinical specimens, for which medical outcomes of various diseases and various treatments are recorded in the medical records, would provide extremely valuable information related to identifying RNAs involved in the cause, maintenance, response, diagnosis, or prognosis of many diseases, such as cancer. Still further, better methods for removing rRNA, including degraded rRNA, from non-rRNA RNA molecules of interest would greatly improve the applicability and success of methods that comprise deliberately degrading the RNA as part of the particular method (such as the method of Ingolia et al., Science 324: 218-23, 2009).

[0004] WO 2007/019444 A2 discloses a method of removing rRNA using biotinylated oligonucleotides. The biotinylated oligonucleotides used to remove the rRNA are shown to be about 20 nucleobases in length. They comprise only one affinity-tag.

[0005] The manual for the RiboMinus™ Kit of Invitrogen: "RiboMinus Transcriptome Isolation Kit (Human/Mouse), retrieved from the internet under [URL: http://tools.invitrogen.com/content/sfs/manuals/ribominus_human_mouse_man.pdf](http://tools.invitrogen.com/content/sfs/manuals/ribominus_human_mouse_man.pdf) on March 12, 2012 discloses a method of removing rRNA using oligonucleotide probes, which are biotinylated. The probes comprise one affinity-tag.

[0006] Su and Sordillo in Molecular Biotechnology, vol. 10, no. 1, August 1998, on pages 83-85, disclose a simple method to enrich mRNA from total prokaryotic RNA disclose. The method uses biotin-labeled antisense rRNA to subtract rRNA.

SUMMARY OF THE INVENTION

[0007] The present invention provides methods, compositions, and kits as defined in the claims for generating rRNA-depleted samples or for isolating rRNA from samples. In particular, the present invention provides methods for generating compositions comprising affinity-tagged antisense rRNA molecules, kits comprising such compositions, and methods for using such compositions to generate rRNA-depleted samples or to isolate rRNA from a sample (e.g., for further analysis and use).

[0008] In some embodiments, the present invention provides methods for generating a composition comprising affinity-tagged antisense rRNA molecules that exhibit sequences which, either alone or in combination, are complementary to the complete sequence of at least one rRNA molecule selected from among 28S, 26S, 25S, 18S, 5.8S and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S and 5S prokaryotic rRNA molecules, wherein the composition is for use in a method for generating a rRNA-depleted sample or for isolating rRNA from a sample. In some

embodiments, the composition comprises affinity-tagged antisense rRNA molecules that exhibit sequences which, either alone or in combination, are complementary to multiple different rRNA molecule selected from among 28S, 26S, 25S, 18S, 5.8S and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S and 5S prokaryotic rRNA molecules from one or multiple cells or organisms. In some embodiments, the present invention provides methods for using the composition comprising the affinity-tagged antisense rRNA molecules for generating a rRNA-depleted sample and/or for isolating rRNA from a sample.

[0009] In some embodiments, the present invention provides a composition comprising affinity-tagged antisense rRNA molecules that, either alone or in combination, exhibit one or more sequences that are complementary to substantially all of the sequence exhibited by at least one full-length rRNA molecule selected from among 28S, 26S, 25S, 18S, 5.8S and/or 5S eukaryotic cytoplasmic rRNA molecules, and 12S and 16S eukaryotic mitochondrial rRNA molecules. In some embodiments, the present invention provides a composition comprising affinity-tagged antisense rRNA molecules that, either alone or in combination, exhibit one or more sequences that are complementary to substantially all of the sequence exhibited by the at least one full-length rRNA molecule selected from among 23S, 16S and 5S prokaryotic rRNA molecules. In particular embodiments, the at least one full-length cytoplasmic rRNA molecule includes both the 28S rRNA and the 18S rRNA (or both the 25S rRNA and the 18S rRNA, or both the 26S rRNA and the 18S rRNA) from one or multiple eukaryotic cells, tissues, organs, or organisms. In some embodiments, the at least one full-length rRNA molecule includes both the 23S rRNA and the 16S rRNA from one or multiple prokaryotic organisms. In some embodiments wherein the at least one full-length rRNA molecule includes both the 28S rRNA and the 18S rRNA (or both the 25S rRNA and the 18S rRNA, or both the 26S rRNA and the 18S rRNA) from one or multiple eukaryotic cells, tissues, organs, or organisms, the affinity-tagged antisense rRNA molecules also correspond to the 5.8S rRNA molecule and the 5S rRNA molecule from the one or multiple eukaryotic cells, tissues, organs, or organisms. In some embodiments, the affinity-tagged antisense rRNA molecules also correspond to at least one full-length rRNA molecule that includes both the 12S and 16S eukaryotic mitochondrial rRNA molecules from the one or multiple eukaryotic cells, tissues, organs, or organisms. In some embodiments, the affinity-tagged antisense rRNA molecules also correspond to at least one full-length rRNA molecule that includes both the 23S rRNA and the 16S rRNA from one or multiple prokaryotic organisms, and in some embodiments, the affinity-tagged antisense rRNA molecules also correspond to at least one full-length rRNA molecule that includes the 5S rRNA molecules from the one or multiple prokaryotic organisms. In further embodiments of any of the compositions comprising affinity-tagged antisense rRNA molecules, the compositions are substantially free of non-rRNA RNA molecules comprising the affinity tags. In further embodiments of any of the compositions comprising affinity-tagged antisense rRNA molecules, the compositions further comprise a binding matrix comprising affinity-tag-binding molecules.

[0010] In some embodiments wherein the composition comprising affinity-tagged antisense rRNA molecules corresponds to all of at least one rRNA molecule selected from among 28S, 26S, 25S, and 18S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and prokaryotic 23S and 16S rRNA molecules, said affinity-tagged antisense rRNA molecules are fragmented (e.g., by controlled nuclease fragmentation or by controlled fragmentation using a divalent metal cation, such as Mg^{2+} , and heat). In some embodiments, the composition comprising fragmented affinity-tagged antisense rRNA molecules comprises fragments ranging in size from about 3,500 nucleotides to about 240 nucleotides.

[0011] In some embodiments, the present invention provides a method for generating a composition comprising affinity-tagged antisense rRNA molecules comprising: a) generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and b) contacting the double-stranded DNA molecules comprising the RNA polymerase promoter with an RNA polymerase and ribonucleoside-5'-triphosphates complementary to all of the nucleobases, including at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase, one of which pair comprises an affinity tag and the other of which pair does not comprise an affinity tag, and incubating under conditions wherein affinity-tagged antisense rRNA molecules are generated corresponding to substantially all of the sequence of the at least one rRNA molecule.

[0012] With respect to such methods, work conduct during the development of embodiments of the present invention determined that the concentration of the ribonucleoside-5'-triphosphate molecules comprising the affinity tag relative to the concentration of the ribonucleoside-5'-triphosphate molecules that lacked the affinity tag in the at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase was important for generating a composition comprising affinity-tagged antisense rRNA molecules that could be used to remove >95% of the at least one rRNA molecules from a sample, and that the relative concentration of the ribonucleoside-5'-triphosphate molecules comprising the affinity tag needed to be higher than had been previously used in the art. In some embodiments of the present method for generating a composition comprising antisense rRNA molecules, at least about 35% of the ribonucleoside-5'-triphosphate molecules comprising the at least one pair of ribonucleoside-5'-triphosphates comprise an affinity tag and about 65% of the ribonucleoside-5'-triphosphate molecules comprising the pair do not have an affinity tag. In some embodiments of this method, at least about 40% of the ribonucleoside-5'-

triphosphate molecules comprising the at least one pair of ribonucleoside-5'- triphosphates have an affinity tag and about 60% of the ribonucleoside-5'- triphosphate molecules comprising the pair do not have an affinity tag. In some embodiments of this method, at least about 50% of the ribonucleoside-5'- triphosphate molecules comprising the at least one pair of ribonucleoside-5'- triphosphates have an affinity tag and about 50% of the ribonucleoside-5'-triphosphate molecules comprising the pair do not have an affinity tag. In some embodiments of this method, at least about 60% of the ribonucleoside-5'-triphosphate molecules comprising the at least one pair of ribonucleoside-5'-triphosphates have an affinity tag and about 40% of the ribonucleoside-5'-triphosphate molecules comprising the pair do not have an affinity tag. In some embodiments of this method, wherein the at least one rRNA molecule is selected from 5.8S and 5S eukaryotic cytoplasmic rRNA, and prokaryotic 5S rRNA, at least about 75% of the ribonucleoside-5'- triphosphate molecules comprising the at least one pair of ribonucleoside-5'- triphosphates have an affinity tag and 25% of the ribonucleoside-5'- triphosphate molecules comprising the pair do not have an affinity tag.

[0013] In some other embodiments, the present invention provides a method for generating a composition comprising affinity-tagged antisense rRNA molecules comprising: a) generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; b) contacting the double-stranded DNA molecules comprising the RNA polymerase promoter with an RNA polymerase and ribonucleoside-5'-triphosphates complementary to all of the nucleobases, including at least one ribonucleoside-5'-triphosphate that comprises an affinity-tag-reactive moiety (e.g., an allylamino group on the 5 position of uridine), and incubating under conditions wherein antisense rRNA molecules comprising the affinity-tag-reactive moiety are generated, which antisense rRNA molecules correspond to substantially all of the sequence of the at least one rRNA molecules; and c) contacting the antisense rRNA molecules comprising the affinity-tag-reactive moiety with a quantity of an affinity tag reagent (e.g., Biotin-X-X-NHS, EPICENTRE Biotechnologies, Madison, WI) under conditions wherein the affinity tag reagent reacts with at least a portion of the affinity-tag-reactive moieties in the antisense rRNA molecules comprising the affinity-tag-reactive moiety and affinity-tagged antisense rRNA molecules are generated.

[0014] Work conducted during the development of embodiments of the present invention found that the relative number of the affinity tags per given number of nucleobases in the affinity-tagged antisense rRNA molecules is important for generating a composition comprising affinity-tagged antisense rRNA molecules that can be used to remove >95% of the at least one rRNA molecules from a sample, and that the number of the affinity tags per given number of nucleobases in the affinity-tagged antisense rRNA molecules generated using the method varies based on the amount of the affinity-tag-reactive moieties present in the antisense rRNA molecules generated in step b), the properties and concentration of the affinity tag reagent, the reaction conditions, and the reaction time. Thus, in some embodiments of this method, 100% of the ribonucleoside-5'-triphosphate molecules comprising one particular nucleobase have an affinity-tag-reactive moiety. In some other embodiments of this method, the ribonucleoside-5'-triphosphates includes at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase, one of which pair comprises an affinity-tag-reactive moiety (e.g., UTP having an allylamino reactive group on the 5 position of uridine or "AA-UTP") and the other of which pair does not comprise an affinity tag. In some embodiments of this method, at least about 50% of the ribonucleoside-5'- triphosphate molecules comprising the at least one pair of ribonucleoside-5'- triphosphates have an affinity-tag-reactive moiety and about 50% of the ribonucleoside-5'- triphosphate molecules comprising the pair do not have an affinity-tag-reactive moiety. In some embodiments of this method, at least about 75% of the ribonucleoside-5'- triphosphate molecules comprising the at least one pair of ribonucleoside-5'-triphosphates have an affinity-tag-reactive moiety and 25% of the ribonucleoside-5'- triphosphate molecules comprising the pair do not have an affinity-tag-reactive moiety. In some embodiments of this method, the affinity tag reagent is a biotinylation reagent (e.g., Biotin-X-X-NHS) and the affinity tag comprises biotin. In some embodiments, the biotin is joined via a spacer arm to the nucleobase (e.g., a spacer arm joined to the allylamino group on the 5 position of uridine, e.g., from incorporation of AA-UTP).

[0015] In further embodiments, the present invention provides a method for generating a composition comprising antisense rRNA molecules comprising: a) generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and b) contacting the double-stranded DNA molecules comprising the RNA polymerase promoter with an RNA polymerase and ribonucleoside-5'-triphosphates complementary to all of the nucleobases, including at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase, one of which pair comprises an affinity tag and the other of which pair does not comprise an affinity tag, and incubating under conditions wherein affinity-tagged antisense rRNA molecules are generated corresponding to substantially all of the sequence of the at least one rRNA molecule, wherein the affinity tags are present at a ratio of at least two affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on fluorescence quantification of 200 ng of affinity-tagged antisense rRNA according to the instructions supplied with the Fluorescence Biotin Quantitation Kit, Pierce Biotechnology, Rockford, IL; Cat. #: Thermo 46610) after digestion of the affinity-

tagged antisense rRNA with RNase 1 at 36°C for 45 min and heat inactivation of the enzyme according to the RNase 1 literature provided by the manufacturer, EPICENTRE Biotechnologies, Madison, WI). In some other embodiments of this method, the affinity tags are present at a ratio of at least about three to five affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some other embodiments of this method, the affinity tags are present at a ratio of at least about four to six affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some other embodiments of this method, the affinity tags are present at a ratio of at least about six to eight affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some embodiments of this method wherein the at least one rRNA molecule is selected from among 28S, 26S, 25S, and 18S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S and 16S prokaryotic rRNA molecules, at least about 35% to about 50% of the ribonucleoside-5'-triphosphates comprising the at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase comprise the affinity tag (e.g., an affinity tag comprising biotin; e.g., biotin-16-UTP) and the affinity tags are present in the affinity-tagged antisense rRNA molecules generated at a ratio of at least about two to eight affinity tags per hundred nucleobases (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some embodiments of this method wherein the at least one rRNA molecule is selected from among 5.8S and 5S eukaryotic rRNA molecules and 5S prokaryotic rRNA molecules, at least about 60% to about 75% of the ribonucleoside-5'-triphosphates comprising the at least one pair of ribonucleoside-5'-triphosphates complementary to the same nucleobase comprise the affinity tag (e.g., an affinity tag comprising biotin; e.g., biotin-16-UTP) and the affinity tags are present in the affinity-tagged antisense rRNA molecules generated at a ratio of at least about two to eight affinity tags per hundred nucleobases (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit).

[0016] In further embodiments, the present invention provides a method for generating a composition comprising antisense rRNA molecules comprising: a)

generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; b) contacting the double-stranded DNA molecules comprising the RNA polymerase promoter with an RNA polymerase and ribonucleoside-5'-triphosphates complementary to all of the nucleobases, including at least one ribonucleoside-5'-triphosphate that comprises an affinity-tag-reactive moiety (e.g., an allylamino group on the 5 position of uridine), and incubating under conditions wherein antisense rRNA molecules comprising the affinity-tag-reactive moiety are generated, which antisense rRNA molecules correspond to substantially all of the sequence of the at least one rRNA molecules; and c) contacting the antisense rRNA molecules comprising the affinity-tag-reactive moiety with a quantity of an affinity tag reagent (e.g., Biotin-X-X-NHS, EPICENTRE Biotechnologies, Madison, WI) under conditions wherein the affinity tag reagent reacts with the affinity-tag-reactive moieties in the antisense rRNA molecules comprising the affinity-tag-reactive moiety and affinity-tagged antisense rRNA molecules are generated, wherein the affinity tags are present at a ratio of at least about two affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on fluorescence quantification of RNase 1-digested affinity-tagged antisense rRNA molecules (200 nanograms) using the Fluorescence Biotin Quantitation Kit from Pierce Biotechnology, Rockford, IL). In some other embodiments of this method, the affinity tags are present at a ratio of at least about three to five affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some other embodiments of this method, the affinity tags are present at a ratio of at least about four to six affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit). In some other embodiments of this method, the affinity tags are present at a ratio of at least about six to eight affinity tags per hundred nucleobases of the affinity-tagged antisense rRNA molecules (e.g., based on the fluorescence data obtained using the Pierce Fluorescence Biotin Quantitation Kit).

[0017] In some embodiments of any of the methods herein for generating affinity-tagged antisense rRNA molecules, the affinity tag comprises biotin. In some embodiments, the biotin is joined via a spacer arm to the 5 position of uridine. In some embodiments of any of these methods, the RNA polymerase is selected from among T7 RNA polymerase, T3 RNA polymerase, and SP6 RNA polymerase. In some embodiments of any of these methods, wherein the at least one rRNA molecule is selected from among eukaryotic cytoplasmic 25S, 26S, and 28S rRNA molecules, the RNA polymerase is SP6 RNA polymerase. In some embodiments of any of these methods, wherein the at least one rRNA molecule is a prokaryotic 23S rRNA, the RNA polymerase is SP6 RNA polymerase.

[0018] In certain embodiments of any of the above methods for generating affinity-tagged antisense rRNA molecules, the methods further comprise, after generating the antisense rRNA molecules, contacting the double-stranded DNA molecules comprising the RNA polymerase promoter with a DNase enzyme under conditions wherein the double-stranded DNA molecules

comprising the RNA polymerase promoter are digested.

[0019] In some embodiments of any of the methods for generating affinity-tagged antisense rRNA molecules, step a) of the method (comprising "generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules") comprises either: i) incubating the at least one rRNA molecule with a DNA polymerase and at least one primer pair under conditions wherein double-stranded DNA molecules that comprise an RNA polymerase promoter that is capable of directing RNA synthesis of antisense RNA corresponding to all of the at least one rRNA molecule are generated, wherein each said at least one primer pair comprises a forward primer and a reverse primer, wherein the reverse primer anneals to the at least one rRNA molecule and has a 5' portion that exhibits the sequence of one strand of an RNA polymerase promoter and the forward primer anneals to the DNA generated from the reverse primer; or: ii) incubating the at least one rRNA molecule with a DNA polymerase and at least one primer pair under conditions wherein double-stranded DNA molecules are generated, wherein each said at least one primer pair comprises a forward primer and a reverse primer, wherein the reverse primer primes DNA synthesis after annealing to the at least one rRNA molecule and the forward primer primes DNA synthesis after annealing to the DNA generated from DNA polymerase extension of the reverse primer, and then ligating said double-stranded DNA molecules generated from each at least one primer pair into a DNA vector that comprises an RNA polymerase promoter, wherein RNA polymerase promoter is capable of directing synthesis of antisense RNA that is complementary to the at least one rRNA molecule using said double-stranded DNA that is ligated into said DNA vector as a template.

[0020] In some embodiments of any of the methods for generating affinity-tagged antisense rRNA molecules, step a) of the method (comprising "generating double-stranded DNA molecules comprising an RNA polymerase promoter that directs RNA synthesis of antisense RNA corresponding to all of at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules") comprises obtaining a genomic DNA fragment that encodes the at least one rRNA molecule and either: i) contacting said genomic DNA fragment with a DNA polymerase and at least one primer pair, wherein the first primer of each of said at least one primer pair has a 5' portion that exhibits the sequence of one strand of an RNA polymerase promoter and a 3' portion that is complementary to a portion of a first strand of the genomic DNA fragment and the second primer of each of said at least one primer pair is complementary to a portion of the second strand of the genomic DNA fragment, and incubating under conditions wherein RNA polymerase promoter-containing double-stranded DNA copies of at least a portion of the genomic DNA fragment are generated, wherein the RNA polymerase promoter is capable of directing RNA synthesis of antisense RNA corresponding to all of the at least one rRNA molecule; or: ii) ligating said genomic DNA fragment or a PCR amplification product thereof into a DNA vector that comprises an RNA polymerase promoter to obtain a genomic clone, wherein the RNA polymerase promoter in said genomic clone is capable of directing synthesis of antisense RNA that is complementary to the at least one rRNA molecule using said double-stranded DNA that is ligated into said DNA vector as a template.

[0021] In some embodiments of the methods for generating affinity-tagged antisense rRNA molecules and of the compositions generated using the method, the affinity-tagged antisense rRNA molecules do not exhibit any rRNA internal transcribed spacer sequences. In some other embodiments, the affinity-tagged antisense rRNA molecules exhibit internal transcribed spacer sequences selected from among: i) a sequence exhibited by the ITS1 rRNA spacer region, which is located between a eukaryotic 18S rRNA gene and a eukaryotic 5.8S rRNA gene; ii) a sequence exhibited by the ITS2 rRNA spacer region, which is located between a eukaryotic 5.8S rRNA gene and a eukaryotic 28S rRNA gene; iii) a sequence exhibited by a prokaryotic 16S-23S ITS; and iv) a sequence exhibited by a prokaryotic 23S-5S rRNA ITS.

[0022] In some embodiments, the present invention provides a composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix comprising affinity-tag-binding molecules, wherein the affinity-tagged antisense rRNA molecules, alone or in combination, exhibit sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules.

[0023] A composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix comprising affinity-tag-binding molecules is generated by incubating the composition comprising affinity-tagged antisense rRNA molecules with a binding matrix comprising affinity-tag-binding molecules under binding conditions, wherein the affinity tag binds to the affinity-tag-binding molecules that are attached to the matrix or solid support (e.g., microparticles) to form a specific binding pair. In some embodiments, this method further comprises the step of washing the composition under conditions wherein affinity-tagged antisense rRNA molecules that are not specifically bound to the binding matrix are removed, thereby generating a purified composition comprising affinity-tagged antisense rRNA molecules that are bound to the binding matrix.

[0024] In particular embodiments of this aspect of the invention, the composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix can be any composition of affinity-tagged antisense rRNA molecules described herein and can be generated using any method for generating a composition comprising antisense rRNA molecules described herein.

[0025] In one particular embodiment of this composition, the affinity-tagged antisense rRNA molecules comprise biotin as the affinity tag, wherein the biotin is joined to at least about two nucleobases per hundred nucleobases of the antisense rRNA molecules, and the binding matrix comprises a microparticle to which an affinity-tag-binding molecule comprising streptavidin or avidin is attached. In some other embodiments of this composition, the affinity-tagged antisense rRNA molecules comprise biotin as the affinity tag, wherein the biotin is joined to at least about two to four nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least three to five nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least four to six nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least six to eight nucleobases per hundred nucleobases of the antisense rRNA molecules.

[0026] In some embodiments of the invention, any of the compositions comprising affinity-tagged antisense rRNA molecules, including any of the compositions comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix, is used in a method of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule.

[0027] In some embodiments, the present invention provides methods for generating a rRNA-depleted sample from an initial sample comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and at least one non-rRNA RNA molecule of interest; ii) a composition comprising affinity-tagged antisense rRNA molecules that, alone or in combination, exhibit sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix and are removed from the treated sample, thereby generating a rRNA-depleted sample, wherein the rRNA-depleted sample is substantially free of rRNA sequences exhibited by the at least one rRNA molecule and comprises substantially all of the at least one non-rRNA RNA molecule of interest present in the initial sample (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the at least one non-rRNA RNA molecule of interest present in the initial sample).

[0028] In certain embodiments, one or more steps are repeated and/or additional methods are performed to generate the rRNA-depleted sample. In particular embodiments, the methods are performed with repeated rounds of subtraction in order to generate the rRNA-depleted sample. In particular embodiments, at least a portion of the RNA molecules in the initial sample are highly fragmented, and wherein the rRNA-depleted sample is at least 50% ... 60% ... 70% ... 80% ... 90.0% ... 95% ... 98% ... 99% ... or 100% free of rRNA sequences exhibited by the at least one rRNA molecule.

[0029] In certain embodiments, the present invention provides methods for generating a rRNA-depleted sample from an initial sample comprising: a) providing: i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and at least one non-rRNA RNA molecule of interest; ii) a composition comprising affinity-tagged antisense rRNA molecules that, alone or in combination, are complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix and are removed from the treated sample, thereby generating an rRNA-depleted sample, wherein the rRNA-depleted sample comprises substantially all (e.g., >90%..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9%) of the at least one non-rRNA RNA molecule of interest present in the initial sample and is substantially free (e.g., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of rRNA sequences exhibited by the at least one rRNA molecule present in the initial sample.

[0030] In additional embodiments, the present invention provides methods for generating a rRNA-depleted sample from an initial sample comprising: a) providing: i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA

molecules and at least one non-rRNA RNA molecules of interest; ii) a composition comprising affinity-tagged antisense rRNA molecules complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the affinity tags are present on the antisense rRNA molecules at a ratio of at least about two to at least four affinity-tags per hundred nucleobases of the antisense rRNA molecules; and iii) a binding matrix comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions wherein at least a portion of the double-stranded rRNA hybrids bind to the binding matrix and are removed from the treated sample, thereby generating a rRNA-depleted sample, wherein the rRNA-depleted sample comprises substantially all (e.g., >90%..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9%) of the at least one non-rRNA RNA molecule of interest present in the initial sample and is substantially free (e.g., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of molecules that, either alone or in combination, exhibit the sequences within the at least one rRNA molecule present in the initial sample.

[0031] In certain embodiments, the present invention provides methods for generating a rRNA-depleted sample from an initial sample comprising: a) providing; i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and at least one non-rRNA RNA molecule of interest; ii) a composition comprising affinity-tagged antisense rRNA molecules complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix and are removed from the treated sample, thereby generating a rRNA-depleted sample, wherein the rRNA-depleted sample comprises substantially all (e.g., >90%..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9%) of the at least one non-rRNA RNA molecule of interest present in the initial sample) and wherein, the rRNA-depleted sample is either: i) $\geq 99.0\%$ free (e.g., >99.0%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9%, 99.95%, or 99.99% free) of RNA molecules that, either alone or in combination, exhibit a sequence within the at least one rRNA molecule present in the initial sample, or ii) contains undetectable levels of the at least one rRNA molecules from the initial sample as measured using agarose gel electrophoresis and ethidium bromide staining.

[0032] In some embodiments, the present invention provides methods for generating a rRNA-depleted sample from an initial sample comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and at least one non-rRNA RNA molecule of interest; and ii) a composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix (e.g., microparticles comprising affinity-tag-binding molecules), wherein the affinity-tagged antisense rRNA molecules, alone or in combination, exhibit sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules in the composition and at least some of the rRNA molecules form double-stranded rRNA hybrids that are bound to the binding matrix, thereby generating a treated sample; and c) removing the binding matrix to which the double-strand rRNA hybrids are bound, thereby generating a rRNA-depleted sample, wherein the rRNA-depleted sample is: i) substantially free of rRNA sequences exhibited by the at least one rRNA molecule and comprises substantially all of the at least one non-rRNA RNA molecule of interest present in the initial sample (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the at least one non-rRNA RNA molecule of interest present in the initial sample), and ii) either, substantially free (e.g., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of rRNA sequences exhibited by the at least one rRNA molecule present in the initial sample, or, contains undetectable levels of the at least one rRNA molecules from the initial sample as measured using agarose gel electrophoresis and ethidium bromide staining. In particular embodiments of this method of the invention, the composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix is any composition of affinity-tagged antisense rRNA molecules described herein and/or is generated using any method for generating a composition comprising antisense rRNA molecules described herein.

[0033] In particular embodiments of any of the methods for generating an rRNA-depleted sample, the rRNA-depleted sample is 98.0% free of rRNA molecules that exhibit sequences within the at least one rRNA molecule (e.g., 98.0% free, 98.5% free, 99.0% free, 99.5% free, 99.7% free, 99.9% free, 99.99% free, or 100% free). In other embodiments, the at least one rRNA molecule includes both the 28S rRNA and the 18S rRNA, and wherein the rRNA-depleted sample is 99.0% free (e.g., 99.5% free) of rRNA molecules that exhibit sequences within the 28S rRNA gene and the 18S rRNA gene. In other embodiments, the rRNA-depleted

sample contains undetectable levels of rRNA molecules that exhibit sequences within the at least one rRNA molecule (e.g., undetectable as measured using agarose gel electrophoresis and ethidium bromide staining). In some embodiments of the method, the composition comprising affinity-tagged antisense rRNA molecules does not exhibit any rRNA internal transcribed spacer sequences. In some other embodiments of the method, the composition comprising the antisense rRNA molecules exhibits internal transcribed spacer sequences selected from among: i) a sequence exhibited by the ITS1 rRNA spacer region, which is located between a eukaryotic 18S rRNA gene and a eukaryotic 5.8S rRNA gene; ii) a sequence exhibited by the ITS2 rRNA spacer region, which is located between a eukaryotic 5.8S rRNA gene and a eukaryotic 28S rRNA gene; iii) a sequence exhibited by a prokaryotic 16S-23S ITS; and iv) a sequence exhibited by a prokaryotic 23S-5S rRNA ITS. In particular embodiments of the methods for generating rRNA-depleted samples or for isolating rRNA from a sample, step b) is performed using a composition comprising antisense rRNA molecules that do not exhibit any rRNA internal transcribed spacer sequences, whereas, in other embodiments, step b) is performed using a composition comprising affinity-tagged antisense rRNA molecules that exhibit one or more of the internal transcribed spacer sequences.

[0034] In some embodiments of any of the methods for generating a rRNA-depleted sample from an initial sample wherein the initial sample comprises RNA molecules that comprise rRNA molecules and at least one non-rRNA RNA molecule of interest, the RNA molecules in the initial sample are fragmented (e.g., from an FFPE or other sample comprising degraded RNA molecules, or from a sample wherein the RNA molecules are deliberately degraded (e.g. by incubation in the presence of heat and Mg²⁺) prior to being provided in the initial sample). In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample, the at least one non-rRNA RNA molecule of interest comprises a multiplicity of non-rRNA molecules of interest. In some embodiments, the at least one non-rRNA RNA molecule of interest comprises substantially all of the non-rRNA RNA molecules present in the initial sample (e.g., including both the intact and fragmented non-rRNA RNA molecules present in the initial sample). In some embodiments, the at least one non-rRNA RNA molecule of interest comprises substantially all of the eukaryotic mRNA molecules present in the initial sample. In some embodiments, the at least one non-rRNA RNA molecule of interest comprises substantially all of the non-rRNA RNA molecules comprising the transcriptome (minus the at least one rRNA molecule) from one or multiple eukaryotic cells, tissues, organs, or organisms (e.g., for use in making sequencing templates or labeled target nucleic acid, e.g., for analysis of the expression or relative expression of said at least one non-rRNA RNA molecule of interest by digital expression analysis or microarray analysis, respectively). In some embodiments, the at least one non-rRNA RNA molecule of interest comprises the non-rRNA RNA molecules comprising a transcriptome (minus the at least one rRNA molecule) of one or multiple eukaryotic cells, tissues, organs, or organisms (e.g., for use in making sequencing templates or labeled target nucleic acid, e.g., for analysis of the expression or relative expression of all of said non-rRNA RNA molecules by digital expression analysis or microarray analysis, respectively, e.g., for medical or agricultural analysis). In some embodiments, the at least one non-rRNA RNA molecule of interest comprises a subfraction of the non-rRNA RNA molecules comprising the transcriptome (minus the at least one rRNA molecules) of one or multiple cells, tissues, organs, or organisms (e.g., a subfraction selected from among mRNA molecules, miRNA molecules, ncRNA molecules, piwiRNA, snRNA, etc.) (e.g., for use in making sequencing templates or labeled target nucleic acid, e.g., for analysis of the expression or relative expression of said subfraction of non-rRNA RNA molecules, e.g., by digital expression analysis or microarray analysis, respectively, e.g., for medical or agricultural analysis). In some embodiments, the at least one non-rRNA RNA molecule of interest comprises substantially all of the non-rRNA RNA molecules comprising a transcriptome (minus the at least one rRNA molecule) of one or multiple prokaryotic cells or of one or multiple cells comprising both eukaryotic and prokaryotic cells. In some embodiments, the at least one non-rRNA RNA molecule of interest comprises or consists of one or multiple RNA molecules that exhibit specific nucleic acid sequences (e.g., wherein the presence or absence or the quantity of said one or multiple RNA molecules that exhibit the specific nucleic acid sequences is used to detect a pathogen or a medical condition, e.g., for screening (e.g., for screening for the presence of a pathogen in water, on a surface, such as a hospital surface, etc.), or e.g., for diagnostic or theranostic assay (e.g., for diagnosing or monitoring the quantity of a pathogen, or the status of a disease or medical condition for deciding on a therapy or treatment)).

[0035] In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample, the method further comprises using the rRNA-depleted sample or the at least one non-rRNA RNA molecule of interest contained therein for further analysis or use. In some embodiments, the method further comprises using the at least one non-rRNA RNA molecule of interest as part of a method for generating templates for next-generation DNA sequencing (e.g., for digital expression analysis or RNA-Seq, miRNA profiling, etc.). In some embodiments of the method for generating a rRNA-depleted sample, the method further comprises using the at least one non-rRNA RNA molecule of interest as part of a method for generating labeled target nucleic acid molecules for hybridization to probes of an array or microarray on a porous or non-porous surface (e.g., to probes on an array or microarray, a dot blot, etc.). In some embodiments of the method for generating a rRNA-depleted sample, the method further comprises using the rRNA-depleted sample for performing a diagnostic or theranostic assay to detect for the presence of the at least one non-rRNA RNA molecule of interest (e.g., wherein the presence or quantity of said at least one non-rRNA RNA molecule of interest is indicative of the presence or status of a health or disease state). In some embodiments of the method for generating a rRNA-depleted sample, the method further comprises amplifying the at least one non-rRNA RNA molecule of interest for further analysis or use. In some embodiments of the method for generating a rRNA-depleted sample, the method further

comprises using the at least one non-rRNA RNA molecule of interest or an amplification product thereof for transfection of a eukaryotic cell (e.g., an antigen-presenting cell (APC), such as a dendritic cell, a macrophage, or an artificial APC for immunotherapeutic use). In some embodiments, the at least one non-rRNA RNA molecule of interest or an amplification product thereof is used for transfection of a human or animal cell from the same individual from whom the at least one non-rRNA RNA molecule of interest was obtained (e.g., to make a vaccine comprising the APC-transfected cell for immunotherapeutic use to treat a disease, e.g., cancer, in a human or animal individual). In some embodiments, the at least one non-rRNA RNA molecule of interest or an amplification product thereof is used for transfection of a cell from a different human or animal than the cell from whom the at least one non-rRNA RNA molecule of interest was obtained (e.g., to make a vaccine comprising the APC-transfected cell for immunotherapeutic use to treat a disease, e.g., cancer, in a human or animal individual). In certain embodiments, the RNA vaccine is made using the at least one non-rRNA RNA of interest from a first individual and the RNA vaccine is used to vaccinate a second individual. In some embodiments of the method for generating a rRNA-depleted sample, the method further comprises using the at least one non-rRNA RNA molecule of interest or an amplification product thereof to manufacture an RNA vaccine for therapeutic use. In some embodiments of the method, the at least one non-rRNA RNA molecule of interest or a sense RNA amplification product thereof is used to manufacture an RNA vaccine that is used to directly inoculate a patient for therapeutic use. In some embodiments, the RNA vaccine is made using the at least one non-rRNA RNA of interest from a cell, tissue, or organ from first individual and the RNA vaccine is administered to said first individual as an immunotherapeutic treatment. In some embodiments, the RNA vaccine is made using the at least one non-rRNA RNA of interest from a first individual and the RNA vaccine is used to vaccinate a second individual.

[0036] In still other embodiments, the invention provides methods for using the composition comprising affinity-tagged antisense rRNA molecules for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules.

[0037] Thus, in some embodiments, the present invention provides methods for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule, the method comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and non-rRNA RNA molecules; ii) a composition comprising affinity-tagged antisense rRNA molecules that exhibits sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix; d) removing the binding matrix to which are bound the double-stranded rRNA hybrids comprising the affinity-tagged antisense rRNA molecules and the at least some of the rRNA molecules from the treated sample; and e) incubating the binding matrix in a solution under conditions wherein the at least some rRNA molecules from the treated sample are released into the solution, thereby isolating substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule present in the initial sample (e.g., at least >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the at least one full-length rRNA molecules present in the initial sample).

[0038] In some embodiments, the present invention provides methods for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule, the method comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and non-rRNA RNA molecules; ii) a composition comprising affinity-tagged antisense rRNA molecules that exhibits sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix; d) removing the binding matrix to which are bound the double-stranded rRNA hybrids comprising the affinity-tagged antisense rRNA molecules and the at least some of the rRNA molecules from the treated sample; and e) incubating the binding matrix in a solution under conditions wherein the at least some rRNA molecules from the treated sample are released into the solution, thereby generating an isolated rRNA sample comprising substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule present in the initial sample (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the at least one full-length rRNA molecule present in the initial sample) and wherein the

isolated rRNA sample is substantially free (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of the non-rRNA RNA molecules present in the initial sample.

[0039] In some embodiments, the present invention provides methods for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule, the method comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and non-rRNA RNA molecules; ii) a composition comprising affinity-tagged antisense rRNA molecules that exhibits sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; and iii) a binding matrix (e.g., microparticles) comprising affinity-tag-binding molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules and at least some of the rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; c) contacting the treated sample with the binding matrix under conditions such that at least a portion of the double-stranded rRNA hybrids bind to the binding matrix; d) removing the binding matrix to which are bound the double-stranded rRNA hybrids comprising the affinity-tagged antisense rRNA molecules and the at least some of the rRNA molecules from the treated sample; and e) incubating the binding matrix in a solution under conditions wherein the at least some rRNA molecules from the treated sample are released into the solution, thereby generating an isolated rRNA sample comprising substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule present in the initial sample (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the at least one full-length rRNA molecule present in the initial sample) and wherein, either i) the isolated rRNA sample is substantially free (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of the non-rRNA RNA molecules present in the initial sample, or ii) contains undetectable levels of non-rRNA RNA from the initial sample as measured using agarose gel electrophoresis and ethidium bromide staining.

[0040] In some embodiments of the methods for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule, the isolated rRNA sample is from an initial sample (e.g., comprising a biological specimen, including a medical specimen, such as saliva, sputum, feces, urine, or a cell, tissue, or organ sample, an environmental sample, a metagenomic sample, or any other type of sample which may contain the RNA of interest), whether the sample is unprepared or prepared (e.g., by fixation using a solution, e.g., formalin or ethanol; mounting on a slide; or using other procedures known in the art), and the isolated rRNA is analyzed to determine the genera, species or strains of origin, thereby indicating what particular genera, species or strains were present in the initial sample, and therefore, in the particular specimen or environment from which the sample was collected. For example, in some embodiments, the method is used to identify and study the organisms present in a human or animal or plant "microbiome", e.g., for research, medical, or agricultural applications. In some embodiments, the isolate rRNA sample is from an initial sample comprising a sample from a human, animal, or plant specimen that is provided for medical diagnostic or theranostic testing or analysis (e.g., to detect a pathogenic microorganism, such as a fungal or bacterial pathogen, that is or may be related to or causative of a disease condition, e.g., a pathogenic bacterium that can be detected or diagnosed based on analysis of at least one rRNA molecule, e.g., as described by Chakravorty, S et al., J. Microbiol. Methods 69: 330-339, 2007 and by Millar, BC et al. in Current Issues Mol. Biol. 9: 21-40, 2007) In some embodiments, the isolated rRNA sample is analyzed using any of the many next-generation sequencing methods known in the art (e.g., after tagging the 3' end or the 3' and 5' ends of the isolated rRNA molecules and reverse transcribing them to make tagged or di-tagged linear ssDNA templates or circular ssDNA templates for next-generation sequencing (e.g., using a Roche 454™, Illumina Solexa™, Life Technologies SOLID™, Pacific Biosciences, Intelligent Biosystems, Helicos, Qiagen, or another next-generation sequencing platform. In still other embodiments, the isolated rRNA sample is analyzed by real-time PCR.

[0041] In some embodiments, the present invention provides methods for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule, the method comprising: a) providing i) an initial sample comprising RNA molecules, wherein the RNA molecules comprise rRNA molecules and non-rRNA RNA molecules; and ii) a composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix (e.g., microparticles comprising affinity-tag-binding molecules), wherein the affinity-tagged antisense rRNA molecules, alone or in combination, exhibit sequences corresponding to substantially all of at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules; b) contacting the initial sample with the composition under conditions such that at least some of the affinity-tagged antisense rRNA molecules in the composition and at least some of the rRNA molecules form double-stranded rRNA hybrids that are bound to the binding matrix, thereby generating a treated sample; and c) removing the binding matrix to which the double-strand rRNA hybrids are bound, thereby isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule. In some embodiments of this method, the method further comprises the step of washing the binding matrix to which the double-stranded rRNA hybrids are bound in order to remove non-specifically-bound non-rRNA RNA molecules from the sample. In some embodiments, the treated

sample is stored on the binding matrix for future analysis or use. In some embodiments, the method further comprises the step of incubating the binding matrix in a solution under conditions wherein the at least some rRNA molecules from the treated sample are released into the solution, thereby generating a solution of the isolated rRNA sample comprising substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule present in the initial sample (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule present in the initial sample) and wherein, either i) the isolated rRNA sample is substantially free (e.g., at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% free) of the non-rRNA RNA molecules present in the initial sample, or ii) contains undetectable levels of non-rRNA RNA from the initial sample as measured using agarose gel electrophoresis and ethidium bromide staining. In some embodiments, the isolated rRNA sample is from an environmental or metagenomic sample and the isolated rRNA is analyzed to determine the genera, species or strains that were present in the sample, and therefore, the particular environment from which the sample was collected. In some embodiments, the isolated rRNA sample is analyzed (e.g., by next-generation sequencing, real-time reverse transcription PCR, or another method, such as a method described elsewhere herein).

[0042] In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the method uses any of the compositions comprising affinity-tagged antisense rRNA molecules obtained using any of the methods described herein for generating affinity-tagged antisense rRNA molecules.

[0043] In some embodiments of any of the methods for generating a rRNA-depleted sample or for isolating substantially all of the rRNA that, either alone or in combination, exhibits a sequence within at least one full-length rRNA molecule from an initial sample, the composition comprising affinity-tagged antisense rRNA molecules that are bound to a binding matrix comprises biotin as the affinity tag, wherein the biotin is joined to at least about two to eight nucleobases per hundred nucleobases of the antisense rRNA molecules, and the binding matrix comprises microparticles to which streptavidin or avidin is attached as the affinity-tag-binding molecule. In some other embodiments of this method, the affinity-tagged antisense rRNA molecules comprise biotin as the affinity tag, wherein the biotin is joined to at least about two to four nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least about three to five nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least about four to six nucleobases per hundred nucleobases of the antisense rRNA molecules, or to at least about six to eight nucleobases per hundred nucleobases of the antisense rRNA molecules.

[0044] In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the rRNA that, either alone or in combination, exhibits a sequence within at least one full-length rRNA molecule from an initial sample, the initial sample contains degraded RNA and the method is used for generating a rRNA-depleted sample for further analysis and use. For example, in some embodiments, the initial sample is an FFPE sample that contains degraded RNA, and the method is used for generating a rRNA-depleted sample that contains at least one non-rRNA RNA molecule of interest, wherein the rRNA-depleted sample is used for analysis of expression or relative expression of one or more RNA molecules, selected from among mRNA, miRNA, ncRNA, piwiRNA, and RNA comprising the whole transcriptome (minus the rRNA) from one or multiple cells, tissues, organs, or organisms, wherein the method of analysis is selected from among microarray analysis (e.g., Affymetrix, NimbleGen Systems, Agilent Systems), next-generation sequencing (or so-called "digital" analysis), including, among others, next-gen RNA sequencing methods and techniques referred to by terms such as "RNA-Seq," "digital mRNA profiling," "transcriptome profiling," and "ribosome profiling" (e.g., Ingolia et al., Science 324: 218-223, 2009), screening analysis, and analysis using a diagnostic or theranostic assay, including a diagnostic or theranostic assay comprising reverse-transcription qPCR, and a diagnostic or theranostic assay comprising RNA amplification and/or detection of a sequencing using a labeled probe.

[0045] In certain embodiments of any of the methods of the invention for using a composition comprising affinity-tagged antisense rRNA molecules for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the composition of affinity-tagged antisense rRNA molecules are generated from a first species or organism, and used for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule from a second species or organism different from the first species or organism (e.g., the first species or organism is mouse or rat and the second species or organism is human). In some embodiments, the affinity-tagged antisense rRNA molecules are generated from any at least one rRNA molecule from the first species of organism wherein said affinity-tagged antisense rRNA molecules hybridize with sequences exhibited within all of the at least one rRNA molecules present in the initial sample from the second species or organism under the conditions used in the method for generating a rRNA-depleted sample or for isolating at least one rRNA molecule from the initial sample. In further embodiments, the first species or organism is a non-human mammal (e.g., cat, dog, sheep, mouse, rat, monkey, etc.) and the second species or organism is *homo sapiens*. In

particular embodiments, the first species or organism comprises non-*E. coli* bacteria, and the second species or organism is *E. coli*. In some embodiments, the method for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule is performed using a metagenomic or environmental sample containing multiple species or organisms. Thus, in some embodiments wherein the composition comprising affinity-tagged antisense rRNA molecules corresponds to at least one full-length prokaryotic rRNA molecule (e.g., generated from at least one full-length rRNA molecule from *E. coli*), the composition is used for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit a sequence within at least one full-length rRNA molecule comprising multiple rRNA molecules from multiple species or organisms.

[0046] In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, instead of using the compositions comprising affinity-tagged antisense rRNA molecules, the method uses compositions comprising affinity-tagged single-stranded DNA molecules that, alone or in combination, are complementary to substantially all of the sequence exhibited by the at least one full-length rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the affinity tags are present at a ratio of at least about two to at least four affinity tags per hundred nucleobases of the affinity-tagged single-stranded DNA molecules.

[0047] Word conducted during developments of the present invention found that the method can be performed using samples wherein the initial sample comprises generally any amount of total RNA molecules. In one embodiment, the amount of total RNA present in the initial sample is between about 10 picograms and 50 nanograms. In another embodiment, the amount of total RNA present in the initial sample is between about 50 nanograms and one microgram. In still another embodiment, the amount of total RNA present in the initial sample is between about one microgram and five micrograms. In some embodiments of these methods, at least a portion of the RNA molecules in the initial sample are highly fragmented (e.g., previously digested with an RNase enzyme or from older RNA samples). In some embodiments of these methods, the RNA molecules in the initial sample are from a paraffin-embedded sample (e.g., paraffin-embedded formalin-fixed sample). In further embodiments of these methods, the affinity-tagged antisense RNA molecules corresponds to the at least one rRNA molecules from the same species or organism (e.g., the affinity-tagged antisense RNA molecules correspond to the at least one rRNA molecule from human cells or the affinity-tagged antisense RNA molecules correspond to the at least one rRNA molecule from *E. coli* cells). In certain embodiments of these methods wherein the affinity-tagged antisense RNA molecules corresponds to the at least one rRNA molecule from human cells, the at least one rRNA molecule is human 28S rRNA.

[0048] In some embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the ratio of the affinity-tag-binding molecules to the affinity tags during the contacting in step b) is at least 2 to 1 (e.g., 2:1, 3:1, 4:1, 5:1, ... or 10:1).

[0049] In certain embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the initial sample comprising RNA molecules represents total RNA isolated from a cell or tissue sample or from an environmental sample (e.g., from a cell line, a tissue biopsy sample, a bodily fluid sample, a swab from a hospital surface, a water sample, etc.). In some embodiments of these methods, the method further comprises providing a control sample of RNA (e.g., comprising total RNA) from a cell or tissue sample, and also performing the method using the control sample. In certain embodiments, the control sample comprises total RNA from HeLa cells and/or from *E. coli* cells.

[0050] In additional embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the initial sample is substantially free of salts and/or organic liquids. In other of these embodiments, the initial sample comprises RNase-free water and/or TE buffer. In particular embodiments, the method further comprises treating the rRNA-depleted sample or the isolated rRNA sample under conditions such that the RNA molecules present in the sample are further purified (e.g., by further purification using ethanol, isopropanol or ammonium acetate precipitation). In particular embodiments, the rRNA-depleted sample or the isolated rRNA sample is used in methods for further analysis, such as for analysis by next-gen sequencing, for preparing labeled target for microarray analysis, for screening or diagnostic or theranostic analysis (e.g., using a plant, human, or animal, screening, diagnostic or theranostic kit). In some embodiments, the nucleic acids in the rRNA-depleted sample or the isolate rRNA sample is amplified prior to further analysis. In other embodiments of these methods, the contacting in step b) is conducted in the presence of an RNase inhibitor. In some embodiments, the compositions used in the method further comprise RNase free water.

[0051] In further embodiments of any of the methods of the invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence of at least one full-length rRNA molecule, the condition in step b) comprise incubating at a temperature of about 60-75 °C for a first time period (e.g., 5-15 minutes) and incubating at about room temperature for a second time period (e.g., 10-25 minutes). In some embodiments of these methods, the conditions in step b) include the presence of hybridization buffer. In particular embodiments of these methods, the conditions in step c) include at least occasional mixing at room temperature and/or at 35-60°C, or 25-70°C, or about 50°C. In further embodiments of these methods, the binding matrix comprises a plurality of individual particles (e.g., nanoparticle, magnetic particles, macroporous beads, etc.) In further embodiments of these methods, the binding matrix comprises a binding column or a membrane. In some embodiments of these methods, the affinity tags are selected from biotin, avidin or streptavidin, digoxigenin, antibodies (or antibody fragments), and other useful binding molecules. In certain embodiments of these methods, the affinity-tag-binding molecules are selected from biotin, avidin or streptavidin, digoxigenin, antibodies (or antibody fragments), and other useful binding molecules.

[0052] In some embodiments of the invention, one or more of the sequences in the at least one rRNA molecules, or in one or more of the non-rRNA RNA molecules, or in one or more other nucleic acid molecules present in the initial sample is amplified (e.g., using any appropriate method known in the art for RNA and/or DNA amplification, e.g., real-time PCR or reverse transcription-PCR, transcription-mediated amplification, RNA amplification using a RiboMultiplier™ Kit, Epicentre Biotechnologies, Madison, WI, strand-displacement amplification, LAMP, ICAN™, UCAN™ (TAKARA), rolling circle amplification, etc.) either, prior to, or after generating the rRNA-depleted sample or isolating the at least one rRNA molecule, respectively, using one of the methods of the present invention for generating a rRNA-depleted sample or for isolating substantially all of the RNA molecules that, either alone or in combination, exhibit the sequence within at least one full-length rRNA molecule. In some of these embodiments, the sample comprising the respective one or more amplified sequences of non-rRNA RNA molecules, or rRNA molecules, and/or other nucleic acid molecules is used for further analysis.

[0053] In other embodiments, the present invention provides a composition comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, wherein the composition is substantially free of non-rRNA RNA molecules comprising the affinity tags. In certain embodiments, the composition further comprises a binding matrix, wherein the binding matrix comprises affinity-tag-binding molecules (e.g., microparticles). In particular embodiments, the antisense rRNA molecules are bound to the binding matrix via the affinity-tag - affinity tag binding molecule interaction.

[0054] In some embodiments, the present invention provides compositions comprising: a) a composition comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity tags, and b) non-rRNA RNA molecules that are affinity-tag free. In certain embodiments, the compositions further comprise a binding matrix comprising affinity-tag-binding molecules.

[0055] In particular embodiments, the present invention provides compositions comprising an rRNA-depleted sample comprising non-rRNA RNA molecules, wherein the composition is substantially free of rRNA sequences exhibited by at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules.

[0056] In further embodiments, the present invention provides compositions comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, and wherein the composition is substantially free of non-rRNA RNA molecules comprising the affinity tags.

[0057] In other embodiments, the present invention provides compositions comprising: a) compositions comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, and b) non-rRNA RNA molecules that are affinity-tag free. In further embodiments, the compositions further comprise a binding matrix comprising affinity-tag-binding molecules.

[0058] In some embodiments, the present invention provides compositions comprising: an rRNA-depleted sample comprising non-rRNA RNA molecules, wherein the composition is substantially free of rRNA sequences exhibited by at least one rRNA molecule selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules. In some particular embodiments, the compositions are substantially free of rRNA sequences exhibited by two, three, or four rRNA molecules selected from among 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules. In some particular embodiments, the compositions are substantially free of rRNA sequences exhibited by both 12S and 16S eukaryotic mitochondrial rRNA molecules.

[0059] In certain embodiments, the present invention provides compositions comprising: a) compositions comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, and wherein the antisense rRNA molecules are from a first species or organism, and b) non-rRNA RNA molecules that are affinity-tag free, wherein the non-rRNA RNA molecules are from a second species or organism different from the first species or organism. In other embodiments, the compositions further comprise a binding matrix comprising affinity-tag-binding molecules.

[0060] In some embodiments, the present invention provides compositions comprising: a) affinity-tagged antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the affinity tags are present on the antisense rRNA molecules at a ratio of at least about two to eight affinity tags per hundred nucleobases of the antisense rRNA molecules, and b) non-rRNA RNA molecules that are affinity-tag free. In some embodiments, composition comprises affinity-tagged antisense rRNA molecules from a first species or organism and non-rRNA RNA molecules that are affinity-tag free from a second species or organism. In certain embodiments, the compositions further comprise a binding matrix comprising affinity-tag-binding molecules.

[0061] In additional embodiments, the present invention provides compositions comprising: an rRNA-depleted sample comprising non-rRNA RNA molecules, wherein the composition is at least about 99.0% free of rRNA sequences exhibited by at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules.

[0062] The present invention provides kits or systems for performing any of the methods of the invention, including any of the steps of said methods.

[0063] For example, in some further embodiments, the present invention provides kits and systems comprising: a) a first component comprising a composition comprising antisense rRNA molecules complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, and wherein the composition is substantially free of non-rRNA RNA molecules comprising the affinity tags; and b) at least one second component selected from the group consisting of: i) a binding matrix comprising affinity-tag-binding molecules; ii) a control sample comprising total RNA from a cell or tissue sample; iii) a solution comprising an RNase inhibitor; iv) a binding matrix wash solution that is RNase-free; v) a volume of RNase-free water; vi) a hybridization buffer; vii) a total RNA purification reagent; and viii) a binding matrix resuspension solution, wherein said solution is RNase-free. In certain embodiments, the second component is the binding matrix. In further embodiments, the second component is the control sample. In some embodiments, the second component is the solution comprising an RNase inhibitor.

[0064] In some embodiments, the present invention provides kits and systems comprising: a) a first component comprising a composition comprising antisense rRNA molecules complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity-tags, and wherein the composition is substantially free of non-rRNA RNA molecules comprising the affinity tags; and b) at least one second component selected from the group consisting of: i) a binding matrix comprising affinity-tag-binding molecules; ii) a control sample comprising total RNA from a cell or tissue sample; iii) a solution comprising an RNase inhibitor; iv) a binding matrix wash solution that is RNase-free; v) a volume of RNase-free water; vi) a hybridization buffer; and vii) a total RNA purification reagent.

[0065] In additional embodiments, the present invention provides kits and systems comprising: a) a first component comprising a composition comprising affinity-tagged antisense rRNA molecules complementary to substantially all of the sequence exhibited by the at least one rRNA molecule selected from: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules, 12S

and 16S eukaryotic mitochondrial rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein the antisense rRNA molecules comprise affinity tags, wherein the affinity tags are present on the antisense rRNA molecules at a ratio of at least about two to eight affinity tags per hundred nucleobases of the antisense rRNA molecules, and wherein the composition is substantially free of non-rRNA RNA molecules comprising the affinity tags; and b) at least one second component selected from the group consisting of: i) a binding matrix comprising affinity-tag-binding molecules; ii) a control sample comprising total RNA from a cell or tissue sample; iii) a solution comprising an RNase inhibitor; iv) a binding matrix wash solution that is RNase-free; v) a volume of RNase-free water; vi) a hybridization buffer; and vii) a total RNA purification reagent.

[0066] In some embodiments of a kit or system, the affinity-tags are present in the composition of affinity-tagged antisense rRNA molecules at a ratio of at least about two to about eight affinity tags per every 100 nucleobases of the antisense rRNA molecules (e.g., at least 2:100, 3:100, 4:100, 5:100, 6:100, 7:100, or 8:100). In certain embodiments, the affinity tag is associated with only one nucleobase of the antisense rRNA molecules. In some of these embodiments, the affinity-tags are associated with only one type of nucleobase selected from: adenine (A), cytosine (C), guanine (G) and uracil (U).

[0067] In other embodiments of a kit or system, the compositions comprising affinity-tagged antisense rRNA molecules correspond to at least 95.0% of all of the at least one rRNA sequence (e.g., 95.0% ... 95.5% ... 96.0% ... 96.5% ... 97.0% ... 97.5% ... 98% ... 98.5% ... 99.0% ... 99.5% ... 99.9 ... or 100% of the at least one rRNA molecule (e.g., wherein the compositions comprise antisense rRNA molecules that exhibit, alone or in combination, sequences corresponding to all of the full-length rRNA sequence for a particular rRNA molecule).

BRIEF DESCRIPTION OF THE DRAWINGS

[0068] The foregoing summary and description is better understood when read in conjunction with the accompanying drawings which are included by way of example and not by way of limitation.

Figure 1A shows an ethidium bromide stained agarose gel containing the following PCR amplicons: Lane M - DNA molecular weight ladder; Lane 1 - 300 ng of PCR amplicon for human 18S rRNA; Lane 2 - 300 ng of PCR amplicon for human 5.8S rRNA; Lane 3 - 300 ng of PCR amplicon for human 5S rRNA; Lane 4 - 300 ng of PCR amplicon for *E. coli* 23S rRNA 5' segment; Lane 5 - 300 ng of PCR amplicon for *E. coli* 23S rRNA 3' segment; Lane 6 - 300 ng of PCR amplicon for *E. coli* 16S rRNA; and Lane 7 - 300 ng of PCR amplicon for *E. coli* 5S rRNA

Figure 1B shows an ethidium bromide stained agarose gel containing the following antisense sequences: Lane M - DNA molecular weight ladder; Lane 1 - 300 ng of human antisense 18S rRNA; Lane 2 - 300 ng of human antisense 5.8S rRNA; Lane 3 - 300 ng of human antisense 5S rRNA; Lane 4 - 300 ng of *E. coli* antisense 23S rRNA 5' segment; Lane 5 - 300 ng of *E. coli* antisense 23S rRNA 3' segment; Lane 6 - 300 ng of *E. coli* antisense 16S rRNA; and Lane 7 - 300 ng of *E. coli* antisense 5S rRNA.

Figure 1C shows an ethidium bromide stained agarose gel containing the following lanes: Lane M - DNA molecular weight ladder; Lane 1 - 300 ng of T7-based 28S biotinylated antisense rRNA; and Lane 2 - 300 ng of SP6-based 28S biotinylated antisense rRNA.

Figure 2A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Biotinylated antisense rRNA following purification with microspheres directly on column; Lane 2 - Biotinylated antisense rRNA following purification from supernatant after removal of microspheres; and Lane 3 - Biotinylated antisense rRNA control.

Figure 2B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Biotinylated antisense rRNA control; Lane 2 - Biotinylated antisense rRNA following purification after 20 μ l of microspheres; Lane 3 - Biotinylated antisense rRNA following purification after 2x 20 μ l of microspheres; and Lane 4 - Biotinylated antisense rRNA following purification after 40 μ l of microspheres

Figure 3A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - 10% biotinylated antisense rRNA minus microspheres; Lane 2 - 10% biotinylated antisense rRNA plus 20 μ l microspheres; Lane 3 - 10% biotinylated antisense rRNA plus 40 μ l microspheres; Lane 4 - 20% biotinylated antisense rRNA minus microspheres; Lane 5 - 20% biotinylated antisense rRNA plus 20 μ l microspheres; and Lane 6 - 20% biotinylated antisense rRNA plus 40 μ l microspheres.

Figure 3B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - 35% biotinylated antisense rRNA minus streptavidin microspheres; Lane 2 - 35% biotinylated antisense rRNA plus streptavidin microspheres; Lane 3 - 50% biotinylated antisense rRNA minus streptavidin microspheres; Lane 4 - 50% biotinylated antisense rRNA plus streptavidin microspheres; Lane 5 - 60% biotinylated antisense rRNA minus streptavidin microspheres; and Lane 6 - 60% biotinylated antisense rRNA plus

streptavidin microspheres.

Figure 3C shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR of 35% biotinylated antisense rRNA minus streptavidin microspheres; Lane 2 - PCR of 35% biotinylated antisense rRNA plus streptavidin microspheres; Lane 3 - PCR of 50% biotinylated antisense rRNA minus streptavidin microspheres; Lane 4 - PCR of 50% biotinylated antisense rRNA plus streptavidin microspheres; Lane 5 - PCR of 60% biotinylated antisense rRNA minus streptavidin microspheres; and Lane 6 - PCR of 60% biotinylated antisense rRNA plus streptavidin microspheres. Panel 1 shows 5'-23S rRNA primers. Panel 2 shows 3'-23S rRNA primers. Panel 3 shows 5'-16S rRNA primers. Panel 4 shows 3'-16S rRNA primers.

Figure 4 shows the following: Lane M - DNA molecular weight ladder; Lane 1 - *L. plantarum* total RNA plus *E. coli* biotinylated antisense rRNA mixture subtraction; and Lane 2 - *L. plantarum* total RNA minus *E. coli* biotinylated antisense rRNA mixture subtraction.

Figure 5, panels A-D, showing the following: Lane M - DNA molecular weight ladder; Lanes 1, 2 - PCR result for 50 ng input *E. coli* total RNA plus subtraction; Lanes 3, 4 - PCR result for 100 ng input *E. coli* total RNA plus subtraction; Lanes 5, 6 - PCR result for 500 ng input *E. coli* total RNA plus subtraction; Lanes 7, 8 - PCR result for 1000 ng input *E. coli* total RNA plus subtraction; Lanes 9, 10 - PCR result for 2500 ng input *E. coli* total RNA plus subtraction; Lanes 11, 12 - PCR result for 5000 ng input *E. coli* total RNA plus subtraction; Lanes 13, 14 - PCR result for 500 ng input *E. coli* total RNA minus subtraction; and Lane 15 - PCR result for no template control reaction. Panel 5A shows 5'-23S rRNA RT-PCR. Panel 5B shows 5'-16S rRNA RT-PCR. Panel 5C shows 5S rRNA RT-PCR. Panel 5D shows *ompA* mRNA RT-PCR.

Figure 6A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - 300 ng intact *E. coli* total RNA; and Lane 2 - 300 ng fragmented *E. coli* total RNA.

Figure 6B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact *E. coli* total RNA minus subtraction; Lane 2 - Intact *E. coli* total RNA plus subtraction; Lane 3 - Fragmented *E. coli* total RNA minus subtraction; and Lane 4 - Fragmented *E. coli* total RNA plus subtraction. Panel 1 shows an ethidium bromide stained agarose gel. Panel 2 shows *ompA* mRNA Northern blot. Panel 3 shows 23S rRNA Northern blot. Panel 4 shows 16S rRNA Northern blot. Panel 5 shows 5S rRNA Northern blot.

Figure 6C shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact *E. coli* total RNA minus subtraction; Lane 2 - Intact *E. coli* total RNA plus subtraction; Lane 3 - Fragmented *E. coli* total RNA minus subtraction; and Lane 4 - Fragmented *E. coli* total RNA plus subtraction. Panel 1 shows *ompA* RT-PCR. Panel 2 shows 23S rRNA 5' RT-PCR. Panel 3 shows 16S rRNA 5' RT-PCR. Panel 4 shows 5S rRNA RT-PCR.

Figure 7A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - biotinylated antisense 28S rRNA; Lane 2 - biotinylated antisense 18S rRNA; Lane 3 - Human total RNA; Lane 4 - Human total RNA plus biotinylated antisense 28S rRNA; Lane 5 - Human total RNA plus biotinylated antisense 18S rRNA; Lane 6 - Mouse total RNA; Lane 7 - Mouse total RNA plus biotinylated antisense 28S rRNA; Lane 8 - Mouse total RNA plus biotinylated antisense 18S rRNA; Lane 9 - Rat total RNA; Lane 10 - Rat total RNA plus biotinylated antisense 28S rRNA; and Lane 11 - Rat total RNA plus biotinylated antisense 18S rRNA.

Figure 7B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Human total RNA (HeLa) minus subtraction; Lane 2 - Human total RNA (HeLa) plus subtraction; Lane 3 - Mouse total RNA (3T3) minus subtraction; Lane 4 - Mouse total RNA (3T3) minus subtraction; Lane 5 - Rat total RNA (NRK) minus subtraction; and Lane 6 - Rat total RNA (NRK) plus subtraction.

Figure 8 shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR result for 100 ng input human total RNA plus subtraction; Lane 2 - PCR result for 500 ng input human total RNA plus subtraction; Lane 3 - PCR result for 5.0 µg input human total RNA plus subtraction; Lane 4 - PCR result for 500 ng input human total RNA minus subtraction; and Lane 5 - PCR result for no template control reaction. Panel A shows 5' 28S rRNA RT-PCR. Panel B shows 3' 28S rRNA RT-PCR. Panel C shows 5' 18S rRNA RT-PCR. Panel D shows 3' 18S rRNA RT-PCR. Panel E shows 5.8S rRNA RT-PCR. Panel F shows 5S rRNA RT-PCR. Panel G shows 5' GAPDH mRNA RT-PCR.

Figure 9A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact HeLa total RNA minus subtraction; Lane 2 - Intact HeLa total RNA plus subtraction; Lane 3 - Fragmented (1 minute) HeLa total RNA minus subtraction; Lane 4 - Fragmented (1 minute) HeLa total RNA plus subtraction; Lane 5 - Fragmented (2 minute) HeLa total RNA minus subtraction; Lane 6 - Fragmented (2 minute) HeLa total RNA plus subtraction; Lane 7 - Fragmented (3 minute) HeLa total RNA minus subtraction; and Lane 8 - Fragmented (3 minute) HeLa total RNA plus subtraction.

Figure 9B shows the following: Lane 1 - PCR result for intact HeLa total RNA minus subtraction; Lane 2 - PCR result for intact HeLa total RNA plus subtraction; Lane 3 - PCR result for fragmented (1 minute) HeLa total RNA minus subtraction; Lane 4 - PCR result for fragmented (1 minute) HeLa total RNA plus subtraction; Lane 5 - PCR result for fragmented (2 minute) HeLa total RNA minus subtraction; Lane 6 - PCR result for fragmented (2 minute) HeLa total RNA plus subtraction; Lane 7 - PCR result for

fragmented (3 minute) HeLa total RNA minus subtraction; Lane 8 - PCR result for fragmented (3 minute) HeLa total RNA plus subtraction; and Lane 9 - PCR result for no template control reaction. Panel 1 shows 5' 28S rRNA. Panel 2 shows 5' 18S rRNA. Panel 3 shows 5.8S rRNA. Panel 4 shows 5S rRNA. Panel 5 shows 5' GAPDH mRNA

Figure 10A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact *E. coli* total RNA minus subtraction; Lane 2 - Intact *E. coli* total RNA plus subtraction; Lane 3 - Fragmented *E. coli* total RNA minus subtraction; and Lane 4 - Fragmented *E. coli* total RNA plus subtraction. Panel 1 shows ompA mRNA. Panel 2 shows 23S rRNA. Panel 3 shows 16S rRNA. Panel 4 shows 5S rRNA.

Figure 10B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR result for intact *E. coli* total RNA minus subtraction; Lane 2 - PCR result for intact *E. coli* total RNA plus subtraction; Lane 3 - PCR result for fragmented *E. coli* total RNA minus subtraction; and Lane 4 - PCR result for fragmented *E. coli* total RNA plus subtraction. Panel 1 shows ompA mRNA. Panel 2 shows 23S rRNA. Panel 3 shows 16S rRNA. Panel 4 shows 5S rRNA.

Figure 11A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact HeLa total RNA minus subtraction; Lane 2 - Intact HeLa total RNA plus subtraction; Lane 3 - Fragmented HeLa total RNA minus subtraction; and Lane 4 - Fragmented HeLa total RNA plus subtraction. Panel 1 shows results of an exemplary method of the present invention. Panel 2 shows the results of the OLIGOTEX method.

Figure 11B shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR results for intact HeLa total RNA minus subtraction; Lane 2 - PCR results for intact HeLa total RNA plus subtraction; Lane 3 - PCR results for fragmented HeLa total RNA minus subtraction; and Lane 4 - PCR results for fragmented HeLa total RNA plus subtraction. Panel 1 shows 5'- 28S rRNA. Panel 2 shows 3'- 28S rRNA. Panel 3 shows 5'- 18S rRNA. Panel 4 shows 3'- 18S rRNA. Panel 5 shows 5.8S rRNA. Panel 6 shows 5S rRNA.

Figure 11C shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR results for intact HeLa total RNA minus subtraction; Lane 2 - PCR results for intact HeLa total RNA plus subtraction; Lane 3 - PCR results for fragmented HeLa total RNA minus subtraction; and Lane 4 - PCR results for fragmented HeLa total RNA plus subtraction. Panel 1 shows 5'- GAPDH. Panel 2 shows 3'- GAPDH. Panel 3 shows 5'- PGK1. Panel 4 shows 3'- PGK1.

Figure 11D shows the following: Lane M - DNA molecular weight ladder; Lane 1 - PCR results for intact HeLa total RNA minus subtraction; Lane 2 - PCR results for intact HeLa total RNA plus subtraction; Lane 3 - PCR results for fragmented HeLa total RNA minus subtraction; and Lane 4 - PCR results for fragmented HeLa total RNA plus subtraction. Panel 1 shows Poly A- RNA #1. Panel 2 shows Poly A- RNA #3. Panel 3 shows Poly A- RNA #15. Panel 4 shows Bimorphic RNA #5.

Figure 12A shows the following: Lane M - DNA molecular weight ladder; Lane 1 - Intact HeLa total RNA minus subtraction; Lane 2 - Intact HeLa total RNA plus subtraction; Lane 3 - Fragmented HeLa total RNA minus subtraction; and Lane 4 - Fragmented HeLa total RNA plus subtraction.

Figure 12B shows Lane M - DNA molecular weight ladder; Lane 1 - PCR results for intact HeLa total RNA minus subtraction; Lane 2 - PCR results for intact HeLa total RNA plus subtraction; Lane 3 - PCR results for fragmented HeLa total RNA minus subtraction; and Lane 4 - PCR results for fragmented HeLa total RNA plus subtraction; Panel 1 - 5'- 28S rRNA; Panel 2 - 3'- 28S rRNA; Panel 3 - 5'- 18S rRNA; Panel 4 - 3'- 18S rRNA; Panel 5 - 5.8S rRNA; and Panel 6 - 5S rRNA.

Figure 12C shows Lane M - DNA molecular weight ladder; Lane 1 - PCR results for intact HeLa total RNA minus subtraction; Lane 2 - PCR results for intact HeLa total RNA plus subtraction; Lane 3 - PCR results for fragmented HeLa total RNA minus subtraction; Lane 4 - PCR results for fragmented HeLa total RNA plus subtraction; Panel 1- 5'- GAPDH; and Panel 2 - 3'- GAPDH.

DEFINITIONS

[0069] It is to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. Further, unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In describing and claiming the present invention, the following terminology and grammatical variants will be used in accordance with the definitions set forth below.

[0070] When the terms "for example", "e.g.", "such as", "include", "including" or variations thereof are used herein, these terms will not be deemed to be terms of limitation, and will be interpreted to mean "but not limited to" or "without limitation."

[0071] As used herein, a "composition comprising affinity-tagged antisense rRNA molecules" means "a composition comprising RNA molecules that, either alone or in combination, exhibit one or more sequences that are complementary to substantially all of the sequence exhibited by the at least one full-length rRNA molecule, wherein at least a portion of the nucleotides in said RNA molecules are joined to an affinity tag."

[0072] As used herein, the phrase "a composition comprising antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule" or "a composition comprising antisense rRNA molecules corresponding to substantially all of" (or "all of the sequence exhibited by") "at least one rRNA molecule" or "a composition comprising antisense rRNA molecules that, alone or in combination, are complementary to substantially all" (or "all of the sequence") "of at least one full-length rRNA molecule" refers to a composition comprising affinity-tagged antisense rRNA molecules that exhibit, that will specifically hybridize with or anneal with or complex with, at least 95% of the RNA molecules or fragments of RNA molecules that exhibit a sequence of a particular full-length rRNA molecule. Preferably, the antisense rRNA molecules in the composition will specifically hybridize with at least 95% of the molecules of a particular rRNA molecule or fragments thereof in a sample, such that, when affinity-tagged, said antisense rRNA molecules can be used with a binding matrix to remove at least about 95% of the RNA molecules or fragments of RNA molecules that exhibit a sequence of said particular rRNA molecule from the sample. It is not necessary that the antisense nucleic acid sequences exhibited in the composition share 95% sequence identity with the complement of a particular rRNA molecule, but instead, it is only necessary that the antisense nucleic acid sequences are able to specifically hybridize, anneal or complex with at least 95% of the RNA molecules or fragments of RNA molecules that exhibit a sequence of said particular rRNA molecule. The antisense nucleic acid sequences in the composition do not need to be present in a single nucleic acid strand (although they may be), but instead, the composition comprising antisense rRNA molecule can consist of antisense rRNA fragments that will collectively hybridize with at least 95% of the RNA molecules or fragments of RNA molecules that exhibit a sequence of a particular rRNA molecule. In certain embodiments, the antisense nucleic acid sequences will specifically hybridize with at least 95% ... 96% ... 97% ... 98% ... 98.5% ... 99.0% ... 99.3% ... 99.5% ... 99.8%, 99.0% ... 99.5% ... 99.9 ... or 100% of the RNA molecules or fragments of RNA molecules that exhibit the sequence of a particular rRNA molecule.

[0073] As used herein, the phrase "an rRNA-depleted sample that comprises substantially all of said at least one non-rRNA RNA molecule of interest present in said initial sample," refers to a purified sample that originally contained rRNA molecules and a first amount of the at least one non-rRNA RNA molecule of interest and that has been purified by removing a certain amount of rRNA molecules, while retaining at least 90% of said first amount of said at least one non-rRNA RNA molecule of interest. In certain embodiments, at least 91% ... 93% ... 96% ... 97% ... 98% ... 98.5% ... 99.0% ... 99.3% ... 99.5% ... 99.8% ..., or 99.9% of said first amount of said at least one non-rRNA RNA molecule of interest is present in the rRNA-depleted sample. Those with knowledge in the art will know or easily find methods for assaying for the first amount of the at least one non-rRNA RNA molecule of interest in the initial sample and assaying for the amount of the at least one non-rRNA RNA molecule of interest remaining in the rRNA-depleted sample, and using said information to calculate the percentage of said first amount of said at least one non-rRNA RNA molecule of interest that is present in the rRNA-depleted sample. For example, in one embodiment, the percentage of the at least one non-rRNA RNA molecule of interest remaining in the rRNA-depleted sample is determined based on the amounts of the at least one non-rRNA RNA molecule of interest in the initial sample and in the rRNA-depleted sample as determined using reverse transcription real-time PCR (also called "RT-qPCR").

[0074] As used herein, the phrase "an isolated rRNA sample that is substantially free of the non-rRNA RNA molecules present in the initial sample" refers to a purified sample that initially contained non-rRNA RNA molecules and a first amount of the at least one rRNA molecule and that has been purified by removing a certain amount of the non-rRNA RNA molecules, while retaining at least 90% of said first amount of said at least one rRNA molecule. In certain embodiments, at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of said first amount of said at least one rRNA molecule is present in the isolated rRNA sample. In certain embodiments, at least >90% ..., >95% ..., >98% ..., >99% ..., >99.8% ..., or >99.9% of the non-rRNA RNA molecules that were present in the initial sample are not present in the isolated rRNA sample (e.g., wherein the amounts of the at least one rRNA molecule and/or of the non-rRNA molecules are determined using RT-qPCR as described above).

[0075] As used herein, the phrase "substantially free of RNA molecules that exhibit sequences of the at least one rRNA molecule" refers to a purified sample wherein 5% or less of all the molecules present in the initial sample that exhibited a nucleic acid sequence from said at least one rRNA molecule are still present in the rRNA-depleted sample. In certain embodiments, 5% ... 4% ... 3% ... 2% ... 1.5% ... 1.0% ... 0.5% ... or 0.1% or less of all the molecules present in the initial sample that exhibited a nucleic acid sequence from a particular rRNA molecule are still present in the purified sample. Those with knowledge in the art will know or easily find methods for assaying for the amounts of RNA molecules that exhibit sequences of the at least one rRNA molecule that are present in the initial sample and in the rRNA-depleted sample. For example, in one embodiment, the percentage of RNA molecules that exhibit sequences of the at least one rRNA molecules is determined based on the amounts of the RNA molecules

that exhibit sequences of the at least one rRNA molecule in the initial sample and in the rRNA-depleted sample as determined using reverse transcription real-time PCR (also called "RT-qPCR").

[0076] As used herein, the phrase "substantially free of non-rRNA RNA molecules comprising affinity tags," refers to a composition wherein 2% or less of all the affinity- tagged nucleic acid sequences present are non-rRNA RNA molecules having affinity tags. In certain embodiments, 2% ... 1.5% ... 1.0% ... 0.5% ... 0.1% or less of all the affinity-tagged nucleic acid molecules present are non-rRNA RNA molecules having affinity tags.

[0077] As used herein, the terms "an rRNA-depleted sample" or "a sample that is substantially free of rRNA molecules" is sometimes referred to herein as "a rRNA subtracted sample" or "a subtracted sample," and the "methods for generating rRNA-depleted samples" or the "methods for isolating rRNA from samples" are sometimes referred to herein as "methods for rRNA subtraction" or, more simply, as "rRNA subtraction." The term "rRNA subtraction" herein shall mean and refer to "a method for generating a rRNA-depleted sample or a method for isolating rRNA from a sample." Similarly, when a form of the verb "to subtract" is used herein, it shall mean "a method or the process of performing a method for generating a rRNA-depleted samples or for isolating rRNA from a sample" and the term "subtracted" herein shall mean or refer to the rRNA-depleted state of the sample after performing the method or process.

[0078] As used herein, the phrase "binding matrix" refers to any type of substrate, whether porous or non-porous, that will bind affinity-tagged nucleic acid molecules such that affinity-tagged nucleic acid molecules, and the sequences they are hybridized to, can be preferentially removed from a sample. Affinity-tag-binding molecules are associated with the binding matrix to allow the binding matrix to bind affinity-tagged nucleic acid molecules. Examples of such binding matrices include, but are not limited to, nylon membranes or particles, silica membranes or particles, cellulose acetate membranes or particles, membranes or particles composed of silica and Fe₂O₃, and other similar membranes, fibers, coated plates, solid supports, and particles.

[0079] As used herein, a "specific binding pair" refers to two molecules that have affinity for and "bind" to each other under certain conditions, referred to as "binding conditions." Biotin and streptavidin or avidin are examples of a "specific binding pair" or "affinity binding molecules," but the invention is not limited to use of this particular specific binding pair. In many embodiments of the present invention, one member of a particular specific binding pair is referred to as the "affinity tag molecule" or the "affinity tag" and the other as the "affinity-tag-binding molecule" or the "affinity tag binding molecule." For example, but without limitation, in some embodiments, biotin is referred to as the affinity tag or affinity tag molecule, and a streptavidin or avidin molecule, whether it is free, attached to a surface, attached to another molecule, or labeled with a detectable molecule such as a dye, is referred to as the affinity-tag-binding molecule. In other embodiments, streptavidin is the affinity tag and biotin is the affinity-tag-binding molecule, since streptavidin and biotin function together as a specific binding pair or as affinity binding molecules. A wide variety of other specific binding pairs or affinity binding molecules, including both affinity tag molecules and affinity-tag-binding molecules, are known in the art (e.g., see U.S. Pat. No. 6,562,575), which can be used in the present invention. For example, an antigen (which itself may be an antibody) and an antibody, including a monoclonal antibody, that binds the antigen is a specific binding pair. Also, an antibody and an antibody binding protein, such as *Staphylococcus aureus* Protein A, can be employed as a specific binding pair. Other examples of specific binding pairs include, but are not limited to, a carbohydrate moiety which is bound specifically by a lectin and the lectin; a hormone and a receptor for the hormone; and an enzyme and an inhibitor of the enzyme. Usually, molecules that comprise a specific binding pair interact with each other through non-covalent bonds such as hydrogen-bonding, hydrophobic interactions (including stacking of aromatic molecules), van der Waals forces, and salt bridges. Without being bound by theory, it is believed in the art that these kinds of non-covalent bonds result in binding, in part due to complementary shapes or structures of the molecules involved in the binding pair. The term "binding" according to the invention refers to the interaction between an affinity binding molecules or specific binding pairs (e.g., between biotin as an affinity tag molecule and streptavidin as an affinity-tag-binding molecule) as a result of non-covalent bonds, such as, but not limited to, hydrogen bonds, hydrophobic interactions, van der Waals bonds, and ionic bonds. Based on the definition for "binding," and the wide variety of affinity binding molecules or specific binding pairs, it is clear that "binding conditions" vary for different specific binding pairs. Those skilled in the art can easily determine conditions whereby, in a sample, binding occurs between the affinity binding molecules. In particular, those skilled in the art can easily determine conditions whereby binding between affinity binding molecules that would be considered in the art to be "specific binding" can be made to occur. As understood in the art, such specificity is usually due to the higher affinity between the affinity binding molecules than for other substances and components (e.g., vessel walls, solid supports) in a sample. In certain cases, the specificity might also involve, or might be due to, a significantly more rapid association of affinity binding molecules than with other substances and components in a sample.

[0080] In some embodiments of the invention, an "affinity tag reagent" or and "affinity tag having a reactive moiety" is used, by which we mean herein, a molecule that comprises both an affinity tag and a reactive chemical group or moiety that is capable of reacting with one or more atoms or groups of the molecule with which it reacts to form one or more covalent chemical bonds

between the molecule comprising the affinity tag and the molecule with which it reacts. By way of example, but without limitation, in some embodiments, the affinity tag reagent is an acylating reagent (e.g., an N-hydroxysuccinimidyl ester), wherein the affinity tag is chemically joined to an atom in the molecule with which it reacts via an acyl linkage. In other embodiments, the affinity tag reagent is an alkylating reagent, group, wherein the affinity tag is chemically joined to an atom in the molecule with which it reacts via an alkyl linkage. In other embodiments, the affinity tag reagent reacts via an electrocyclic type of chemical reaction, such as a 1,3-dipolar cycloaddition (e.g., cycloaddition of an alkyne with an azide). Thus, the term "reactive" moiety with respect to, for example, an "affinity tag reagent" or "affinity tag having a reactive moiety" is used to refer to a moiety or group that is involved in or responsible for the chemical reaction whereby a molecule comprising the affinity tag reacts chemically to form a covalent chemical bond with one or more atoms in the molecule with which it reacts, rather than to the binding that results between affinity binding molecules due to non-covalent forces and bonds.

[0081] When we refer to attaching the "affinity-tag-binding molecule" or the "affinity tag binding molecule," such as streptavidin or avidin, directly to the surface or solid support, it usually, but not always means that the affinity-tag-binding molecule is covalently attached to the surface by means of a chemical linker that is joined to the surface and to the affinity-tag-binding molecule. When we refer to attaching the "affinity-tag-binding molecule" or the "affinity tag binding molecule," such as streptavidin or avidin, indirectly to the surface, it is meant that the affinity-tag-binding molecule is bound to another molecule with which it has affinity (e.g., an anti-streptavidin antibody) that is in turn bound to the surface. In some embodiments, the affinity-tag-binding molecule, such as streptavidin or avidin, is not attached to a surface, but is bound by another molecule, such as an antibody or Protein A, and the biotinylated modified-nucleotide-capped RNA is recovered by precipitation or by binding to a second antibody or other molecule using methods and compositions known in the art.

[0082] As used herein, the term "about" means encompassing plus or minus 25%. For example, "about 200 nucleotides" refers to a range encompassing between 150 and 250 nucleotides.

[0083] As used herein, the term "hybridization" or "hybridize" is used in reference to the pairing of complementary nucleic acids. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between the nucleic acids) is influenced by such factors as the degree of complementarity between the nucleic acids, stringency of the conditions involved, the melting temperature (T_m) of the formed hybrid, and the G:C ratio within the nucleic acids. A single molecule that contains pairing of complementary nucleic acids within its structure is said to be "self-hybridized." An extensive guide to nucleic acid hybridization may be found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes*, part I, chapter 2, "Overview of principles of hybridization and the strategy of nucleic acid probe assays," Elsevier (1993).

DETAILED DESCRIPTION

[0084] The present invention provides methods, compositions, and kits as defined in the claims for generating rRNA-depleted samples and for isolating rRNA from samples. In particular, the present invention provides compositions comprising affinity-tagged antisense rRNA molecules corresponding to substantially all of the sequence of at least one rRNA molecule (e.g., a eukaryotic 28S, 26S, 25S, 18S, 5.8S, and/or 5S rRNA and/or a prokaryotic 23S, 16S, and/or 5S rRNA) and methods of using such compositions to generate rRNA-depleted samples.

[0085] Ribosomal RNA constitutes a majority mass of the RNA content of a cell and thus, presents a tremendous background contamination burden when studying the less abundant and, often, more relevant transcripts. Methods to reduce rRNA are available but are severely limiting since, for example, they require only "high quality intact RNA" samples for good rRNA removal. Fragments of rRNA in the sample without the complementary consensus sequences will not be removed using the methods in the art and, thus, still contribute to significant rRNA contamination of the sample, which limits the efficacy of downstream transcript analysis and significantly increases cost burden from unnecessary use of consumables and manpower. Furthermore, even in the best case wherein the RNA in the sample comprises so-called "intact total RNA," the sample can still comprise a population of fragmented rRNA, which still gives rise to a rRNA background using the methods in the art. Still further, the methods in the art do not remove sufficient quantities of all of the sequence exhibited by the rRNA molecules, so the rRNA background is high when the sample is used for modern analysis methods, such as for generating tagged sequencing templates from RNA for use in next-generation sequencing (e.g., for so-called "digital" gene expression methods). For example, it has been reported that more than half of the sequence reads can be for rRNA sequences, even after one or more rounds of rRNA removal using commercial methods known in the art (e.g., using RiboMinus™, Life Technologies, Carlsbad, CA). Another requirement of these existing methods is the need for a relatively large sample size, which is frequently prohibitive. Often, total RNA extracted from especially long-term stored cell/tissue samples is invariably very fragmented and limited in quantity. Thus, a method to simultaneously remove both fragmented and intact rRNA and to do so from a small sample size is needed.

[0086] The present invention addresses this need as it provides, for example, methods for the efficient removal of either completely fragmented or variably fragmented rRNA molecules, leaving the non-rRNA RNA transcripts generally unperturbed. Furthermore, the method operates independently of any "unique feature" of the non-ribosomal RNA transcripts such as, a poly(A) tail as in the case of eukaryotic mRNA and thus, presents no selection bias in the recovery of all non-ribosomal RNA transcripts. In certain embodiments to achieve these objectives, one or more affinity-tagged antisense rRNA representing substantially all of the full-length sequence of each rRNA in whole or in part (e.g., eukaryotic 28S, 26S, 25S, 18S, 5.8S and/or 5Ss and/or prokaryotic 23S, 16S and 5S) are synthesized and hybridized to their respective complementary rRNA sequences in the test sample, and the hybrids as well as any residual unhybridized affinity-tagged antisense rRNA molecules are then physically removed using affinity-tag-binding molecules (e.g., linked to a binding matrix), leaving the non-rRNA RNA transcripts unperturbed. Since substantially the entire (or the entire) sequence of at least one, and preferably each rRNA, is represented in the composition comprising affinity-tagged antisense rRNA molecules, any size fragments of the native rRNA contained in the samples are also able to form hybrids under the same hybridization conditions and are efficiently removed along with any intact rRNA sequences that may also be present.

[0087] In particular embodiments, in order to achieve efficient rRNA removal, a molar excess of the affinity-tagged antisense rRNA molecules is used in the hybridization reaction, which drives the hybrid formation process to completion. Excess affinity-tagged antisense rRNA molecules may also be removed since these molecules themselves may contribute to various types of background in different downstream methods or analyses. To accomplish this, the affinity-tagged antisense rRNA molecules preferably contain an optimal amount of affinity-tag molecules, and an appropriate amount of the binding matrix comprising the affinity-tag-binding molecules is used for removal of both rRNA-hybridized and unhybridized affinity-tagged antisense rRNA molecules. As described in the Examples, conditions were developed that allow the methods to be performed using samples comprising a broader dynamic range of total RNA than existing methods, including using samples comprising sub-microgram quantities of total RNA.

Exemplary rRNA Sequences and Primers

[0088] The present invention includes compositions comprising affinity-tagged antisense rRNA molecules corresponding to substantially all of at least one rRNA molecule. The present invention is not limited to compositions comprising antisense rRNA molecules that exhibit sequences that are exactly complementary to particular rRNA molecules in the sample, but instead includes compositions comprising antisense rRNA molecules that hybridize with or anneal to or complex with any rRNA molecules in the sample, whether or not said rRNA molecules are from the same organism.

Exemplary Embodiment

[0089] Described below is an exemplary embodiment that may be used to generate rRNA-depleted samples using affinity-tag-binding molecule linked microspheres (herein after "binding microspheres" or "microspheres"). This embodiment is not meant to limit the present invention and instead is simply an exemplary embodiment.

[0090] Wash the binding microspheres and suspend in solution (20 mM Tris-HCl pH 7.5, 1 M NaCl, 1 mM EDTA and 0.0005% RNase-Free Triton-X100). Allow the binding microspheres to reach room temperature. Vigorously vortex the microspheres for 20 seconds to produce a homogeneous suspension. For each reaction, pipette 25 μ l of the resuspended microspheres into a microsphere wash tube. Centrifuge the dispensed microspheres at 10,000 rpm in a bench top microcentrifuge for 3 minutes. Carefully pipette off and discard the supernatant, without disturbing the microsphere pellet. Wash the microspheres by adding 1 volume of wash solution (20 mM Tris-HCl pH 7.5, 1 M NaCl and 1 mM EDTA) equal to the original volume of microspheres (e.g., add 25 μ l of wash solution for every 25 μ l of microspheres) to the tube. Resuspend the microspheres by vigorous vortex mixing. Centrifuge the tube at 10,000 rpm for 3 minutes in a bench top microcentrifuge. Carefully pipette off and discard the supernatant, without disturbing the microsphere pellet. Repeat the wash, resuspension, centrifuge, and pipetting steps.

[0091] Resuspend the microspheres in 1 volume resuspension solution. For example, add 25 μ l of resuspension solution for every 25 μ l of microspheres that were originally used. Resuspend the microspheres by vigorous vortex mixing to produce a homogeneous suspension. Add 0.25 μ l of ScriptGuard™ RNase Inhibitor (EPICENTRE, Madison, WI) to the tube for every 25 μ l of resuspended microspheres and store them at room temperature.

[0092] For a sample containing, for example, 50 ng - 1 μ g to total RNA (e.g., human, mouse, or rat), in a 0.2-ml thin-walled

microcentrifuge tube, combine: 10X reaction buffer (0.5M Tris-HCl pH7.5 and 1 M NaCl), a total RNA sample, a rRNA removal solution (1.2 pmoles each of affinity-tagged antisense rRNA molecules corresponding to 28S, 18S, 5.8S and 5S human rRNA molecules), and RNase-free water. Gently mix the reaction(s) and incubate at 68°C for 10 minutes. Remove the reaction tube(s) to room temperature and incubate for at least 15 minutes.

[0093] Resuspend the washed microspheres from above by pipetting up and down. For each sample, pipette 25 µl of the microspheres to a new RNase-free 1.5-ml microcentrifuge tube. Add the room temperature total RNA sample to the appropriate tube containing the microspheres. Mix each thoroughly by pipetting the tube contents up and down. Incubate each tube at room temperature for 15 minutes with occasional (every 3 - 4 minutes) mixing by *gentle* vortex (low setting) for 5 seconds. Place the tube at 37°C for 5 minutes. Immediately centrifuge the tube at 14,000 rpm in a microcentrifuge for 5 minutes at room temperature. Carefully pipette off each supernatant, which contains the rRNA-depleted sample, to an RNase-free 1.5-ml microcentrifuge tube. If there are microspheres still visible in the sample, and ethanol precipitation is desired to be used, repeat the above procedure. The rRNA-depleted sample can, for example, be purified by ethanol precipitation or column method.

[0094] For ethanol precipitation, adjust the volume of each sample to 180 µl using RNase-free water. Add 18 µl of 3M Sodium Acetate to each tube. Add 2 µl of Glycogen (10 mg/ml) to each tube and mix by gentle vortex. Add 3 volumes (600 µl) of ice cold 100% ethanol to each tube and mix thoroughly by gentle vortex. Place the tubes at -20°C for at least 1 hour. Centrifuge the tubes at >10,000 x g in a microcentrifuge for 30 minutes. Carefully remove and discard the supernatant. Wash the pellet with ice cold 70% ethanol and centrifuge at >10,000 x g for 5 minutes. Carefully remove and discard the supernatant. Repeat the ethanol and centrifuge step. Centrifuge briefly to collect any residual supernatant. Carefully remove and discard the supernatant and allow the pellet to air dry at room temperature for 5 minutes. Dissolve the pellet in the desired volume of RNase-free water or buffer.

[0095] For column purification, the RNA Clean & Concentrator™-5 Column (Zymo Research; Catalog Numbers R1015, R1016) can be used to purify the rRNA-depleted RNA sample. If using the RNA Clean & Concentrator-5 Column, follow the manufacturer's procedure entitled "*To recover total RNA including small RNAs*". The eluted RNA can be used immediately or stored at -70°C to -80°C.

Sequencing Technologies

[0096] Purified RNA samples generated in accordance with the present invention can be sequenced using any type of suitable sequencing technology. The present invention is not limited by the type of sequencing method employed. Exemplary sequencing methods are described below.

[0097] Illustrative non-limiting examples of nucleic acid sequencing techniques include, but are not limited to, chain terminator (Sanger) sequencing and dye terminator sequencing. Chain terminator sequencing uses sequence-specific termination of a DNA synthesis reaction using modified nucleotide substrates. Extension is initiated at a specific site on the template DNA by using a short radioactive, or other labeled, oligonucleotide primer complementary to the template at that region. The oligonucleotide primer is extended using a DNA polymerase, standard four deoxynucleotide bases, and a low concentration of one chain terminating nucleotide, most commonly a di-deoxynucleotide. This reaction is repeated in four separate tubes with each of the bases taking turns as the di-deoxynucleotide. Limited incorporation of the chain terminating nucleotide by the DNA polymerase results in a series of related DNA fragments that are terminated only at positions where that particular di-deoxynucleotide is used. For each reaction tube, the fragments are size-separated by electrophoresis in a slab polyacrylamide gel or a capillary tube filled with a viscous polymer. The sequence is determined by reading which lane produces a visualized mark from the labeled primer as you scan from the top of the gel to the bottom.

[0098] Dye terminator sequencing alternatively labels the terminators. Complete sequencing can be performed in a single reaction by labeling each of the di-deoxynucleotide chain-terminators with a separate fluorescent dye, which fluoresces at a different wavelength.

[0099] A set of methods referred to as "next-generation sequencing" techniques have emerged as alternatives to Sanger and dye-terminator sequencing methods (Voelkerding et al., Clinical Chem., 55: 641-658, 2009; MacLean et al., Nature Rev. Microbiol., 7: 287-296). Most current methods describe the use of next-generation sequencing technology for de novo sequencing of whole genomes to determine the primary nucleic acid sequence of an organism. In addition, targeted re-sequencing (deep sequencing) allows for sensitive mutation detection within a population of wild-type sequence. Some examples include recent work describing the identification of HIV drug-resistant variants as well as EGFR mutations for determining response to anti-TK therapeutic drugs. Recent publications describing the use of bar code primer sequences permit the

simultaneous sequencing of multiple samples during a typical sequencing run including, for example: Margulies, M. et al. "Genome Sequencing in Microfabricated High-Density Picolitre Reactors", *Nature*, 437, 376-80 (2005); Mikkelsen, T. et al. "Genome-Wide Maps of Chromatin State in Pluripotent and Lineage-Committed Cells", *Nature*, 448, 553-60 (2007); McLaughlin, S. et al. "Whole-Genome Resequencing with Short Reads: Accurate Mutation Discovery with Mate Pairs and Quality Values", ASHG Annual Meeting (2007); Shendure J. et al. "Accurate Multiplex Polony Sequencing of an Evolved Bacterial Genome", *Science*, 309, 1728-32 (2005); Harris, T. et al. "Single-Molecule DNA Sequencing of a Viral Genome", *Science*, 320, 106-9 (2008); Simen, B. et al. "Prevalence of Low Abundance Drug Resistant Variants by Ultra Deep Sequencing in Chronically HIV-infected Antiretroviral (ARV) Naive Patients and the Impact on Virologic Outcomes", 16th International HIV Drug Resistance Workshop, Barbados (2007); Thomas, R. et al. "Sensitive Mutation Detection in Heterogeneous Cancer Specimens by Massively Parallel Picoliter Reactor Sequencing", *Nature Med.*, 12, 852-855 (2006); Mitsuya, Y. et al. "Minority Human Immunodeficiency Virus Type 1 Variants in Antiretroviral-Naive Persons with Reverse Transcriptase Codon 215 Revertant Mutations", *J. Vir.*, 82, 10747-10755 (2008); Binladen, J. et al. "The Use of Coded PCR Primers Enables High-Throughput Sequencing of Multiple Homolog Amplification Products by 454 Parallel Sequencing", *PLoS ONE*, 2, e197 (2007); and Hoffmann, C. et al. "DNA Bar Coding and Pyrosequencing to Identify Rare HIV Drug Resistance Mutations", *Nuc. Acids Res.*, 35, e91 (2007).

[0100] Compared to traditional Sanger sequencing, next-gen sequencing technology produces large amounts of sequencing data points. A typical run can easily generate tens to hundreds of megabases per run, with a potential daily output reaching into the gigabase range. This translates to several orders of magnitude greater than a standard 96-well plate, which can generate several hundred data points in a typical multiplex run. Target amplicons that differ by as little as one nucleotide can easily be distinguished, even when multiple targets from related species or organisms are present. This greatly enhances the ability to do accurate genotyping. Next-gen sequence alignment software programs used to produce consensus sequences can easily identify novel point mutations, which could result in new strains with associated drug resistance. The use of primer bar coding also allows multiplexing of different patient samples within a single sequencing run.

[0101] Next-generation sequencing (NGS) methods share the common feature of massively parallel, high-throughput strategies, with the goal of lower costs in comparison to older sequencing methods. NGS methods can be broadly divided into those that require template amplification and those that do not. Amplification-requiring methods include pyrosequencing commercialized by Roche as the 454 technology platforms (e.g., GS 20 and GS FLX), the Solexa platform commercialized by Illumina, and the Supported Oligonucleotide Ligation and Detection (SOLiD) platform commercialized by Applied Biosystems. Non-amplification approaches, also known as single-molecule sequencing, are exemplified by the HeliScope platform commercialized by Helicos BioSciences, and emerging platforms commercialized by VisiGen and Pacific Biosciences, respectively.

[0102] In pyrosequencing (Voelkerding et al., *Clinical Chem.*, 55: 641-658, 2009; MacLean et al., *Nature Rev. Microbiol.*, 7: 287-296; U.S. Pat. No. 6,210,891; U.S. Pat. No. 6,258,568), template DNA is fragmented, end-repaired, ligated to adaptors, and clonally amplified in-situ by capturing single template molecules with beads bearing oligonucleotides complementary to the adaptors. Each bead bearing a single template type is compartmentalized into a water-in-oil microvesicle, and the template is clonally amplified using a technique referred to as emulsion PCR. The emulsion is disrupted after amplification and beads are deposited into individual wells of a picotitre plate functioning as a flow cell during the sequencing reactions. Ordered, iterative introduction of each of the four dNTP reagents occurs in the flow cell in the presence of sequencing enzymes and luminescent reporter such as luciferase. In the event that an appropriate dNTP is added to the 3' end of the sequencing primer, the resulting production of ATP causes a burst of luminescence within the well, which is recorded using a CCD camera. It is possible to achieve read lengths greater than or equal to 400 bases, and 1×10^6 sequence reads can be achieved, resulting in up to 500 million base pairs (Mb) of sequence.

[0103] In the Solexa/Illumina platform (Voelkerding et al., *Clinical Chem.*, 55: 641-658, 2009; MacLean et al., *Nature Rev. Microbiol.*, 7: 287-296; U.S. Pat. No. 6,833,246; U.S. Pat. No. 7,115,400; U.S. Pat. No. 6,969,488), sequencing data are produced in the form of shorter-length reads. In this method, single-stranded fragmented DNA is end-repaired to generate 5'-phosphorylated blunt ends, followed by Klenow-mediated addition of a single A base to the 3' end of the fragments. A-addition facilitates addition of T-overhang adaptor oligonucleotides, which are subsequently used to capture the template-adaptor molecules on the surface of a flow cell that is studded with oligonucleotide anchors. The anchor is used as a PCR primer, but because of the length of the template and its proximity to other nearby anchor oligonucleotides, extension by PCR results in the "arching over" of the molecule to hybridize with an adjacent anchor oligonucleotide to form a bridge structure on the surface of the flow cell. These loops of DNA are denatured and cleaved. Forward strands are then sequenced with reversible dye terminators. The sequence of incorporated nucleotides is determined by detection of post-incorporation fluorescence, with each fluor and block removed prior to the next cycle of dNTP addition. Sequence read length ranges from 36 nucleotides to over 50 nucleotides, with overall output exceeding 1 billion nucleotide pairs per analytical run.

[0104] Sequencing nucleic acid molecules using SOLiD technology (Voelkerding et al., Clinical Chem., 55: 641-658, 2009; MacLean et al., Nature Rev. Microbiol., 7: 287-296; U.S. Pat. No. 5,912,148; U.S. Pat. No. 6,130,073) also involves fragmentation of the template, ligation to oligonucleotide adaptors, attachment to beads, and clonal amplification by emulsion PCR. Following this, beads bearing template are immobilized on a derivatized surface of a glass flow-cell, and a primer complementary to the adaptor oligonucleotide is annealed. However, rather than utilizing this primer for 3' extension, it is instead used to provide a 5' phosphate group for ligation to interrogation probes containing two probe-specific bases followed by 6 degenerate bases and one of four fluorescent labels. In the SOLiD system, interrogation probes have 16 possible combinations of the two bases at the 3' end of each probe, and one of four fluors at the 5' end. Fluor color and thus identity of each probe corresponds to specified color-space coding schemes. Multiple rounds (usually 7) of probe annealing, ligation, and fluor detection are followed by denaturation, and then a second round of sequencing using a primer that is offset by one base relative to the initial primer. In this manner, the template sequence can be computationally reconstructed, and template bases are interrogated twice, resulting in increased accuracy. Sequence read length averages 35 nucleotides, and overall output exceeds 4 billion bases per sequencing run.

[0105] In certain embodiments, nanopore sequencing is employed (see, e.g., Astier et al., J Am Chem Soc. 2006 Feb 8;128(5):1705-10). The theory behind nanopore sequencing has to do with what occurs when the nanopore is immersed in a conducting fluid and a potential (voltage) is applied across it: under these conditions a slight electric current due to conduction of ions through the nanopore can be observed, and the amount of current is exceedingly sensitive to the size of the nanopore. If DNA molecules pass (or part of the DNA molecule passes) through the nanopore, this can create a change in the magnitude of the current through the nanopore, thereby allowing the sequences of the DNA molecule to be determined.

[0106] HeliScope by Helicos BioSciences (Voelkerding et al., Clinical Chem., 55: 641-658, 2009; MacLean et al., Nature Rev. Microbiol., 7: 287-296; U.S. Pat. No. 7,169,560; U.S. Pat. No. 7,282,337; U.S. Pat. No. 7,482,120; U.S. Pat. No. 7,501,245; U.S. Pat. No. 6,818,395; U.S. Pat. No. 6,911,345; U.S. Pat. No. 7,501,245) is the first commercialized single-molecule sequencing platform. This method does not require clonal amplification. Template DNA is fragmented and polyadenylated at the 3' end, with the final adenosine bearing a fluorescent label. Denatured polyadenylated template fragments are ligated to poly(dT) oligonucleotides on the surface of a flow cell. Initial physical locations of captured template molecules are recorded by a CCD camera, and then label is cleaved and washed away. Sequencing is achieved by addition of polymerase and serial addition of fluorescently-labeled dNTP reagents. Incorporation events result in fluor signal corresponding to the dNTP, and signal is captured by a CCD camera before each round of dNTP addition. Sequence read length ranges from 25-50 nucleotides, with overall output exceeding 1 billion nucleotide pairs per analytical run. Other emerging single molecule sequencing methods real-time sequencing by synthesis using a VisiGen platform (Voelkerding et al., Clinical Chem., 55: 641-658, 2009; U.S. Pat. No. 7,329,492; U.S. Pat. App. Ser. No. 11/671956; U.S. Pat. App. Ser. No. 11/781166) in which immobilized, primed DNA template is subjected to strand extension using a fluorescently-modified polymerase and fluorescent acceptor molecules, resulting in detectable fluorescence resonance energy transfer (FRET) upon nucleotide addition. Another real-time single molecule sequencing system developed by Pacific Biosciences (Voelkerding et al., Clinical Chem., 55: 641-658, 2009; MacLean et al., Nature Rev. Microbiol., 7: 287-296; U.S. Pat. No. 7,170,050; U.S. Pat. No. 7,302,146; U.S. Pat. No. 7,313,308; U.S. Pat. No. 7,476,503) utilizes reaction wells 50-100 nm in diameter and encompassing a reaction volume of approximately 20 zeptoliters (10×10^{-21} L). Sequencing reactions are performed using immobilized template, modified phi29 DNA polymerase, and high local concentrations of fluorescently labeled dNTPs. High local concentrations and continuous reaction conditions allow incorporation events to be captured in real time by fluor signal detection using laser excitation, an optical waveguide, and a CCD camera.

binding matrix solid supports

[0107] The binding matrix employed in the present invention may be any surface or solid support to which the affinity-tag-binding molecule can be attached and which does not exhibit substantial non-specific binding of the affinity-tagged antisense rRNA molecules. The present invention is not limited to any one solid support. In some embodiments, polystyrene plates containing either 96 or 384 wells are employed. In some embodiments, streptavidin (SA) coated 96-well or 384-well microtiter plates (Boehringer Mannheim Biochemicals, Indianapolis, IN) are used as solid supports. In some embodiments, particles or beads are employed. The particles can be made of any suitable material, including, but not limited to, latex. In some embodiments, minicolumns are employed. The columns may contain affinity-tag-binding molecules. In other embodiments, a BEADARRAY (Illumina, San Diego, CA) is employed. A variety of other solid supports are contemplated including, but not limited to, glass microscope slides, glass wafers, gold, silicon, microchips, and other plastic, metal, ceramic, or biological surfaces.

EXAMPLES

[0108] It is understood that the examples and embodiments described herein are for illustrative purposes only and are not intended to limit the scope of the claimed invention.

Example 1

Synthesis of biotinylated antisense rRNA molecules representing the full-length sequences of ribosomal RNA (rRNA) molecules of eukaryotes and prokaryotes

[0109] In order to synthesize antisense rRNA corresponding to the complete sequence of each rRNA molecule, PCR templates were made containing a phage promoter sequence in all but one case as follows.

Human 18S, 5.8S and 5S rRNA PCR templates for *in vitro* synthesis of antisense rRNA

[0110] PCR amplicons representing the full-length 18S (Accession# NR_003286), 5.8S (Accession# U13369 REGION: 6623-6779) and 5S (Accession # NR_023363) rRNA sequences were synthesized. Each 100- μ l PCR reaction comprised 20 pmoles of a forward primer, 20 pmoles of a reverse primer containing T7 RNA polymerase phage promoter sequence (shown in italics) and 5 ng random primed first-strand cDNA made from Universal Human Reference total RNA (Stratagene) under FailSafe PCR reaction conditions optimized as described by the manufacturer (EPICENTRE Biotechnologies). The PCR amplification reactions were performed for 25 cycles to 30 cycles depending on the primer pair and the PCR amplicons corresponding to the full-length 18S, 5.8S and 5S rRNA sequences and PCR products were purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). The following is a list of the respective forward and reverse primers used:

18S rRNA

forward primer (SEQ ID NO: 1: 5'-CCTACCTACCTGGTTGATCC) and

reverse primer (SEQ ID NO: 2: 5'-AATTCTAATACGACTCACTATAGGGAGAGATC CTTCCGCAGGTTACCTAC).

5.8S rRNA

forward primer (SEQ ID NO: 3: 5'-CGACTCTTAGCGGTGGATCACTC) and

reverse primer (SEQ ID NO: 4: 5'-AATTCTAATACGACTCACTATAGGGAGAGATC CTTCCGCAGGTTACCTAC)

5S rRNA

forward primer (SEQ ID NO: 5: 5'-GTCTACGGCCATACCACCCTGAA) and

reverse primer (SEQ ID NO: 6: 5'-AATTCTAATACGACTCACTATAGGGAGAAAG CCTACAGCACCCGGTATTC)

Escherichia coli 23S, 16S and 5S rRNA PCR templates for *in vitro* synthesis of antisense rRNA

[0111] PCR amplicons representing the full-length 23S (Accession# EG30077), 16S (Accession# EG30084) and 5S (Accession # EG30070) rRNA sequences were synthesized. Each 100 μ l PCR reaction comprised 20 pmoles of a forward primer, 20 pmoles of a reverse primer containing T7 RNA polymerase phage promoter sequence (shown in italics) and 5 ng random primed first-strand cDNA made from *E. coli* K-12 total RNA under FailSafe PCR reaction conditions optimized as described by the manufacturer (EPICENTRE Biotechnologies). The PCR amplification reactions were performed for 25 cycles to 30 cycles depending on the primer pair and the PCR amplicons corresponding to the full-length 23S, 16S and 5S rRNA sequences and PCR products were purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). The following is a list of the respective forward and reverse primers used:

23S rRNA

[0112] In order to efficiently amplify the complete 23S rRNA sequence, two pairs of primers were used whereby, the full-length 23S rRNA sequence was divided into approximately two equal segments. The 5' 23S rRNA segment comprised forward primer (SEQ ID NO: 7: 5'-AAGCGACTAAGCGTACACGGTGA) and reverse primer (SEQ ID NO: 8: 5'-AATTCTAATACGACTCACTATAGGGAGATTCTGGAAGCAGG GCATITGTTG) and the 3' 23S rRNA segment comprised forward primer (SEQ ID NO: 9: 5'-CAACAAATGCCCTGCTTCCAGGAA) and reverse primer (SEQ ID NO: 10: 5'-AATTCTAATACGACTCACTATAGGGAGACACGGTTCATTAGTACCGGTTAGCT).

16S rRNA

forward primer (SEQ ID NO: 11: 5'-AGAGTTTGATCCTGGCTCAG) and reverse primer (SEQ ID NO: 12: 5'-AATTCTAATACGACTCACTATAGGGAGAGGAG GTGATCCAACCGCAGGT).

5S rRNA

forward primer (SEQ ID NO: 13: 5'-TGCCTGGCGGCAGTAGCGCGGT) and

reverse primer (SEQ ID NO: 14: 5'-AATTCTAATACGACTCACTATAGGGAGATGC CTGGCAGTTCCTACTCTC).

[0113] Each PCR amplicon (300 ng) representing the respective human or *E. coli* rRNA sequence was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 1A.

Human 28S ribosomal RNA clone

[0114] PCR amplification of the complete 28S rRNA (Accession# NR_003287) sequence was found not to be very efficient, especially for the approximately 1200-nt 5'-end sequence, even after dividing the full-length sequence into segments ranging from 1 Kb to 2 Kb. Thus, in order to generate the required template for *in vitro* synthesis of the 28S antisense rRNA, the 28S rRNA sequence was cloned into both a T7 phage promoter-containing plasmid (pSP73; Promega Corporation) and a SP6 phage promoter-containing plasmid (pSP64; Promega Corporation) using standard cloning methods and strategies (Maniatis et. al. (1982) Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York and information described by Maden et. al. (1987) Biochem. J. 246: 519-527.

In vitro synthesis of biotinylated antisense rRNA molecules from the respective PCR templates for human 18S, 5.8S and 5S and *E. coli* 23S, 16S and 5S rRNA sequences

[0115] Either an AmpliScribe™ T7 High Yield Transcription Kit or an AmpliScribe™ T7-FLASH Transcription Kit (EPICENTRE, Madison, WI) was used for *in vitro* transcription (IVT) reactions comprising 500 ng to 1000 ng of the respective templates (human 18S, 5.8S and 5S and *E. coli* 23S, 16S and 5S) were performed as described by the manufacturer (EPICENTRE Biotechnologies) with the following modification - the uridine triphosphate (UTP) was replaced with mixtures of biotin-16-UTP and UTP comprising 10%, 20%, 35%, 50% or 60% biotin-16-UTP. The IVT reactions were incubated at 37°C for 4 to 6 hours and the DNA template used in each reaction was then digested with DNase I as recommended by the manufacturer. Each biotinylated antisense rRNA was subsequently purified using illustra RNAspin Mini columns as recommended by the manufacturer (GE Healthcare) and eluted in RNase-free water. Each purified antisense rRNA was treated with Baseline-Zero™ DNase I as recommended by the manufacturer (EPICENTRE Biotechnologies) to remove all traces of the DNA template used for transcription. The biotinylated antisense rRNA was again purified using illustra RNAspin Mini columns, eluted in RNase-free water and the RNA concentration determined by measuring the absorbance at 260 nm with a spectrophotometer. (Note: subsequent experiments showed that best results for generating rRNA-depleted samples with respect to 5.8S and 5S eukaryotic rRNA or 5S prokaryotic rRNA were obtained using at least about 75% biotin-16-UTP in the *in vitro* transcription reaction.)

[0116] Each purified antisense RNA (300 ng), representing the entire human or *E. coli* rRNA sequence, as analyzed by ethidium bromide stained agarose gel electrophoresis, is shown in Figure 1B.

In vitro synthesis of biotinylated antisense rRNA from the plasmid templates for human 28S rRNA sequence

[0117] Initially, 1 µg of the linearized T7-28S rRNA plasmid clone was used in an AmpliScribe™ T7 High Yield Transcription Kit for *in vitro* transcription (IVT) as described by the manufacturer (EPICENTRE Biotechnologies) with the following modification - a 1:1 mixture of bio-UTP:ITP instead of 100% UTP. Incubation was performed at 37°C for 4 hours and the DNA template then removed by digesting with DNase I as recommended by the manufacturer (EPICENTRE Biotechnologies). An aliquot (300 ng) of the purified transcription reaction product was analyzed by gel electrophoresis. It was evident from the results (Figure 1C, Lane 1) that very little, if any of the expected full-length antisense 28S rRNA was produced in the IVT reaction. Instead, RNA molecules ranging in size from low to high molecular weights were observed. Thus, it appeared that the T7 RNA polymerase was unable to efficiently transcribe the 28S DNA template.

[0118] Next, 1 µg of the linearized SP6-28S rRNA plasmid clone was tested in a standard AmpliScribe™ SP6 High Yield Transcription Kit for *in vitro* transcription (IVT) as described by the manufacturer (EPICENTRE Biotechnologies) using a similar 1:1 mixture of bio-UTP and UTP. Incubation was performed at 37°C for 4 hours and the DNA template then removed by digesting with DNase I as recommended by the manufacturer (EPICENTRE Biotechnologies). The biotinylated antisense rRNA produced was subsequently purified using illustra RNAspin Mini columns as recommended by the manufacturer (GE Healthcare) and eluted in RNase-free water. Thereafter, the purified antisense 28S rRNA was treated with Baseline-Zero™ DNase I as recommended by the manufacturer (EPICENTRE Biotechnologies) to remove all traces of the DNA template used for transcription. The biotinylated antisense rRNA was again purified using illustra-RNAspin Mini columns, eluted in RNase-free water and the RNA concentration determined by measuring the absorbance at 260 nm with a spectrophotometer. An aliquot (300 ng) of the transcription reaction was analyzed by gel electrophoresis and it was evident from the results that the expected antisense 28S rRNA was produced in the SP6-IVT reaction (Figure 1C, Lane 2).

Example 2

Removal of biotinylated antisense rRNA and resulting ds-rRNA hybrids formed using *E. coli* total RNA using streptavidin-coated microspheres

[0119] *E. coli* rRNA was used as the model to test the amount of ProActive® Streptavidin Coated Microspheres (Bangs Laboratories) needed to remove a fixed quantity biotinylated antisense rRNA from solution. The streptavidin-coated microspheres were washed and resuspended in the bind/wash buffer as recommended by the manufacturer (Bangs Laboratories). *E. coli* biotinylated antisense rRNA materials synthesized using 35% biotin-UTP as described in Example 1 were used to prepare a mixture comprising one microgram each of the 23S rRNA 5' segment, the 23S rRNA 3' segment and the full-length 16S rRNA in a 4-µl volume of RNase-free water. Each 4-µl biotinylated antisense rRNA mixture would represent at least a >2-fold molar excess when mixed with one microgram of native *E. coli* total RNA, which concentration was selected to drive the hybridization of the sense and antisense rRNA to completion or near completion when mixed together.

[0120] The amount of streptavidin-coated microspheres required to efficiently remove the added biotinylated antisense rRNA added to a hybridization reaction was determined. Three 4-µl aliquots of the *E. coli* biotinylated antisense rRNA mixture were prepared in 0.2 ml microcentrifuge tubes to a final reaction volume of 20 µl comprising 1x hybridization buffer (50 mM Tris-HCl, pH7.5 and 100 mM NaCl). The reactions were incubated at 68°C for 5 minutes and then at room temperature for 15 minutes. Reactions #1 and #2 were each then added to a fresh 1.5-ml microcentrifuge tube containing 20 µl of the washed and resuspended streptavidin-coated microspheres. Control reaction #3 was added to a fresh 1.5-ml microcentrifuge tube with only 20 µl bind/wash buffer. All reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex. The biotinylated antisense rRNA remaining in solution following capture by the microspheres was then purified using RNA Clean-up Kit™ -5 columns as recommended by the manufacturer (ZYMO Research). Reactions #1 and #3 were added directly in the ZYMO RNA Clean-up procedure whereas, reaction #2 was first spun at 12,000 rpm for 5 minutes using a benchtop centrifuge to pellet the microspheres and the supernatant removed and then added to the ZYMO RNA Clean-up procedure. Each RNA sample was eluted in 10 µl RNase-free water and a 5-µl aliquot analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 2A.

[0121] Figure 2A, lanes 1 and 2 show the residual biotinylated antisense rRNA following binding to the streptavidin coated microspheres compared to the control (Lane 3). The amount of antisense RNA remaining was more pronounced in Lane 2 where the microspheres were removed prior to the ZYMO RNA clean-up procedure. While unclear, it is possible that adding the hybridization reaction mixture with the microspheres directly to the RNA purification column reduced the binding capacity of the column resulting in a further loss of the biotinylated antisense RNA. Nevertheless, it was evident from this experiment that 20 µl of the streptavidin coated microspheres was insufficient to remove the entire amount of biotinylated antisense rRNA used.

[0122] Thus, an additional 20 μ l of washed microspheres was tested to determine if it would be sufficient to remove the 4 μ l of biotinylated antisense rRNA. Four reactions, each comprising 4 μ l of the *E. coli* biotinylated antisense rRNA were prepared in 1x hybridization buffer to a final volume of 20 μ l. The reactions were incubated at 68°C for 5 minutes and then at room temperature for 15 minutes. Each of reactions #2 and #3 was added to a fresh 1.5-ml microcentrifuge tube containing 20 μ l of the washed and resuspended streptavidin-coated microspheres. Reaction #4 was added to a fresh 1.5-ml microcentrifuge tube containing 40 μ l of the washed and resuspended streptavidin-coated microspheres. Control reaction #1 was added to a fresh 1.5-ml microcentrifuge tube with only bind/wash buffer. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex. The reactions were spun at 12,000 rpm for 5 minutes using a benchtop centrifuge to pellet the microspheres and each supernatant transferred to a fresh 1.5-ml microcentrifuge tubes. To the supernatant of reaction #3, an additional 20 μ l of the washed streptavidin-coated microspheres was added and again incubated at room temperature with occasional gentle mixing (3-4 minutes). Reaction #3 was centrifuged at 12,000 rpm for 5 minutes once more and the supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The biotinylated antisense RNA contained in each of the four reactions was purified using the ZYMO RNA Clean-up procedure and eluted in 10 μ l RNase-free water. The entire eluate for each was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 2B.

[0123] Figure 2B, lane 2 again shows that biotinylated antisense rRNA is not completely removed when only 20 μ l of microspheres is used. However, when either 20 μ l of microspheres followed by a second 20 μ l of microspheres or 40 μ l of microspheres was used, there appears to be complete removal of the added biotinylated antisense rRNA (Lane 3 and Lane 4, respectively). Thus, the one treatment with 40 μ l of microspheres was generally preferred since it required fewer steps.

[0124] Next, the amount of microspheres used as described in Figure 2B was tested to determine if it was also sufficient to efficiently remove the double-stranded ribosomal RNA (ds-rRNA) hybrids formed after annealing the biotinylated antisense rRNA and the native rRNA contained in total RNA. Firstly, in order to demonstrate the formation of the ds-rRNA hybrid, 500 ng of *E. coli* total RNA was mixed with 4 μ l of the *E. coli* biotinylated antisense rRNA in a reaction comprising 1x hybridization buffer in a final volume of 20 μ l. The reaction was incubated at 68°C for 5 minutes and then at room temperature for 15 minutes, purified using the ZYMO RNA Clean-up procedure and eluted in 10 μ l RNase-free water. A 5- μ l aliquot was analyzed by ethidium bromide stained agarose gel electrophoresis along with equivalent amounts of the individual *E. coli* total RNA and biotinylated antisense rRNA as shown in Figure 2C. Figure 2C, lane 3 shows the efficient formation of the ds-rRNA hybrids compared to the individual sense and antisense RNA samples (Lanes 1 and 2, respectively).

[0125] Next, the removal of the ds-rRNA hybrids was tested using the quantities of washed microspheres described in Figure 2B. Three reactions (#1, #2 and #3), each comprising 500 ng of *E. coli* total RNA and 4 μ l of the *E. coli* biotinylated antisense rRNA in 1x hybridization buffer in a final volume of 20 μ l were prepared, along with two control reactions - one containing only the *E. coli* biotinylated antisense rRNA (#4) and the other, only the *E. coli* total RNA (#5). The reactions were incubated at 68°C for 5 minutes and then at room temperature for 15 minutes. Reactions #1 and #3 were each added to a fresh 1.5-ml microcentrifuge tube containing 20 μ l of the washed and resuspended streptavidin-coated microspheres. Reactions #2 and #4 were each added to a fresh 1.5-ml microcentrifuge tube containing 40 μ l of the washed and resuspended streptavidin-coated microspheres. Control reaction #5 was added to a fresh 1.5-ml microcentrifuge tube containing only bind/wash buffer. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex. The reactions were spun at 12,000 rpm for 5 minutes using a benchtop centrifuge to pellet the microspheres and each supernatant transferred to a fresh 1.5-ml microcentrifuge tubes. To the supernatant of reaction #3, an additional 20 μ l of the washed streptavidin-coated microspheres was added and incubated at room temperature with occasional gentle mixing (3-4 minutes). Reaction #3 was again centrifuged at 12,000 rpm for 5 minutes and the supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The RNA contained in each of the five reactions was purified using the ZYMO RNA Clean-up procedure and eluted in 10 μ l RNase-free water. A 5- μ l aliquot of each was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 2D. In addition, 250 ng of untreated *E. coli* total RNA was run as a control.

[0126] Figure 2D, Lane 1 shows the residual ds-rRNA when only 20 μ l of microspheres was used. However, with either 2x 20 μ l (Lane 2) or 1x 40 μ l of microspheres (Lane 3), the ds-rRNA hybrids were no longer visible indicating that the biotinylated antisense rRNA was capable of annealing to its complementary sense rRNA sequences and facilitate their removal by the streptavidin microspheres. The quantity of *E. coli* total RNA did not appear to be affected by the addition of 40 μ l of microspheres (Lane 5) when compared to a similar amount of untreated *E. coli* total RNA (Lane 6). Thus, it was evident that 40 μ l of microspheres was sufficient to remove either the added biotinylated antisense rRNA or the resulting ds-rRNA hybrids formed after hybridization of the biotinylated antisense rRNA to the rRNA.

Example 3

Effect of using different levels of biotin-UTP for synthesis of biotinylated antisense rRNA and its removal with the Streptavidin Coated Microspheres

[0127] The experiments described in Example 2 above used biotinylated antisense rRNA synthesized using a UTP:biotin-UTP mixture comprising 35% biotin-UTP. For the amount of antisense rRNA used in the hybridization reaction, 40 μ l of the streptavidin microspheres was needed for its efficient removal. Thus, it was thought that antisense rRNA made with lesser amounts of biotin-UTP should require less streptavidin microspheres for its quantitative removal, which would result in an overall reduction in the cost associated with the amount of both the biotin-UTP and microspheres required. Biotinylated antisense 16S and 23S rRNA were prepared using either 10% or 20% biotin-UTP solutions as described above in Example 1 and standard 4- μ l mixtures of the respective biotinylated antisense rRNA were then made. The percentage of biotin-UTP used in the *in vitro* transcription reaction for its synthesis is used to refer to the biotinylated antisense rRNA product; for example, if the percentage of biotin-UTP used in the *in vitro* transcription reaction is 10%, with the remaining 90% being UTP, the product is referred to herein as 10% biotinylated antisense rRNA, even though the percentage of biotin-UTP nucleotides incorporated into the product may be less than 10% compared to the UTP nucleotides incorporated into the product.

[0128] Two sets of three reactions each comprising 4 μ l of either the 10% (reaction #1, #2 and #3) or 20% (reactions #4, #5 and #6) biotinylated antisense rRNA mixture were prepared in a final volume of 20 μ l in 1x hybridization buffer. The reactions were incubated at 68°C for 5 minutes and then at room temperature for 15 minutes. Reactions # 2 and #5 were each transferred to a fresh 1.5-ml microcentrifuge tube containing 20 μ l of washed streptavidin microspheres and reactions # 3 and #6 were each transferred to a fresh 1.5-ml microcentrifuge tube containing 40 μ l of washed streptavidin microspheres. Reactions #1 and #4 were each transferred to a fresh 1.5-ml microcentrifuge tubes containing 40 μ l of bind/wash buffer. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex. The reactions were spun at 12,000 rpm for 5 minutes on a benchtop centrifuge and the supernatant transferred to fresh 1.5-ml microcentrifuge tubes. Any biotinylated antisense rRNA remaining in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 10 μ l RNase-free water. The entire amount of each purified sample was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 3A.

[0129] Figure 3A, Lanes 1 and 4 show the 10% and 20% biotinylated antisense rRNA not treated with microspheres, respectively. Lanes 2 and 3 show the 10% biotinylated antisense rRNA treated with 20 μ l and 40 μ l of microspheres, respectively. Lanes 5 and 6 show the 20% biotinylated antisense rRNA treated with 20 μ l and 40 μ l of microspheres, respectively. Clearly, there was a significant amount of both the 10% and 20% biotinylated antisense rRNA remaining independent of either 20 μ l or 40 μ l of microspheres used (Lanes 2, 3, 5 and 6). Nevertheless, the 20% condition (Lanes 5 and 6) was clearly better than the 10% condition (Lanes 2 and 3). Thus, it appeared that using these lower levels of biotin-UTP were used to synthesize the biotinylated rRNA, at least some of the antisense rRNA product synthesized may not contain sufficient biotin molecules to facilitate its removal by the streptavidin microspheres.

[0130] Thus, the question arose as to whether the 35% biotin-UTP used in Example 2 was the optimal concentration to ensure that all antisense rRNA synthesized contained sufficient biotin molecules for its subsequent removal by the streptavidin beads. To answer this question, the following experiment tested biotinylated antisense rRNAs made as described in Example 1 using either 50% or 60% biotin-UTP solutions. The standard 4- μ l mixture of the respective biotinylated antisense 16S and 23S rRNA synthesized using the 50% and 60% levels of biotin-UTP was then made. The standard 4 μ l-mixture of the 35% biotinylated antisense rRNA included as a control.

[0131] Three sets of two reactions, each comprising 4 μ l of the 35% (reactions #1 and #2) or 50% (reactions #3 and #4) or 60% (reactions #5 and #6) biotinylated antisense rRNA mixture were prepared in a final volume of 20 μ l in 1x hybridization buffer. The reactions were incubated at 68°C for 5 minutes and then at room temperature for 15 minutes. Reactions # 2, #4 and #6 were each transferred to a fresh 1.5-ml microcentrifuge tube containing 40 μ l of washed streptavidin microspheres and reactions # 1, #3 and #5 were each transferred to a fresh 1.5-ml microcentrifuge tube containing 40 μ l of bind/wash buffer. All reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex. The reactions were spun at 12,000 rpm for 5 minutes on a benchtop centrifuge and the supernatant transferred to fresh 1.5-ml microcentrifuge tubes. Any biotinylated antisense rRNA remaining in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 μ l RNase-free water. Fifty percent of each purified sample was analyzed by ethidium bromide-stained agarose gel electrophoresis as shown in Figure 3B.

[0132] Figure 3B, Lanes 1, 3 and 5 show the biotinylated antisense rRNA remaining in the absence of any microspheres for the 35%, 50% and 60% biotin-UTP samples, respectively. With the addition of 40 µl of streptavidin microspheres, there was no biotinylated antisense rRNA visibly remaining in any of the corresponding reactions (Lanes 2, 4 and 6). It was not possible to observe any differences in the performance of the 35%, 50% and 60% biotinylated antisense rRNA samples following treatment with 40 µl of the streptavidin microspheres by agarose gel analysis.

[0133] In order to further determine if there was a difference between the use of the 35%, 50% and 60% biotinylated antisense rRNA, the remaining half of each reaction was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5-µl aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' and 3' regions of 23S (SEQ ID NO: 15: 5'-GACGTGCTAATCTGCGATAAGC, SEQ ID NO: 16: 5'-ATGGATTCAAGTTAATGATAGTGTGTCG, SEQ ID NO: 17: 5'-CTGAAAGCATCTAAGCACGAACTTGC and SEQ ID NO: 18: 5'-CCTATCAACGTCGTCGTCTTCAAC) and 16S (SEQ ID NO: 19: 5'-GCCTAACACATGCAAGTCGAAC, SEQ ID NO: 20: 5'-AGCTACCGTTTCCAGTAGTTATCC, SEQ ID NO: 21: 5'-CGGAATCGCTAGTAATCGTGGAT and SEQ ID NO: 22: 5'-TCCCGAAGGTTAAGCTACCTACTT) rRNAs for 20 PCR cycles. PCR amplification is notably more sensitive than ethidium bromide agarose gel analysis and should therefore better demonstrate any differences between the removal efficiencies of the 35%, 50% and 60% biotinylated antisense rRNA by the microspheres. The results are shown in Figure 3C.

[0134] Figure 3C, Lanes 1, 3 and 5 show the expected PCR amplicons for the 5' 23S rRNA (Panel 1), 3' 23S rRNA (Panel 2), 5' 16S rRNA (Panel 3) and 3' 16S rRNA (Panel 4) with the 35%, 50% and 60% biotinylated antisense rRNA mixtures, respectively. Lanes 2, 4 and 6 show the corresponding PCR reaction products following the use of the streptavidin microspheres. As seen in Lane 2 (Panels 2 and 4), residual carryover of 35% biotinylated antisense rRNA resulted in still some specific PCR-amplified products that were not observed with the 50% and 60% biotinylated antisense rRNA materials. Both the 50% and 60% biotinylated antisense rRNA materials behaved similarly and thus, the 50% condition was selected.

Example 4

Removal of the 23S and 16S rRNA from another prokaryotic species (*Lactobacillus plantarum*) using the *E. coli* biotinylated antisense 23S and 16S rRNA mixture

[0135] In order to test if the *E. coli* biotinylated antisense rRNA mixture would be capable of removing the 23S and 16S rRNA sequences exhibited by a relatively diverse prokaryotic species, total RNA from *Lactobacillus plantarum*, which shares approximately 80% homology with *E. coli* 16S and 23S rRNA was used. In one reaction, 500 ng of *L. plantarum* total RNA was mixed with 4 µl of the *E. coli* biotinylated antisense rRNA mixture described in Example 2 above in 1x hybridization buffer in a final volume of 20 µl. A second reaction to serve as a control contained everything as the first reaction minus the *E. coli* biotinylated antisense rRNA mixture. The reactions were incubated at 68°C for 10 minutes and then at room temperature for 15 minutes. Each reaction was then added to a fresh 1.5-ml microcentrifuge tube containing 50 µl of washed and resuspended streptavidin-coated microspheres. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow for formation of the biotin-streptavidin complex. The reactions were then spun at 12,000 rpm for 5 minutes using a benchtop centrifuge to pellet the microspheres, the supernatant removed and the RNA purified using the Zymo RNA Clean-up procedure. Each RNA sample was eluted in 13.5 µl RNase-free water and 50% of each analyzed by ethidium bromide stained agarose gel electrophoresis as shown Figure 4.

[0136] Figure 4, Lane 1 compared to Lane 2 shows that both the 16S and 23S rRNA bands for *L. plantarum* were no longer visible following subtraction with the *E. coli* biotinylated antisense rRNA mixture indicating that they were efficiently subtracted as well even though the rRNA sequences were only 80% homologous.

Example 5

Removal of 23S, 16S and 5S rRNA from a range of inputs of *E. coli* total RNA using the *E. coli* biotinylated antisense rRNA mixture

[0137] An *E. coli* biotinylated antisense rRNA mixture was prepared, now containing 23S, 16S and 5S antisense rRNA, and the following reactions for generating rRNA-depleted samples and control reactions were set up as shown in Table 1, and were performed in duplicate in a final volume of 20 μ l in 1x hybridization buffer:

Table 1:

Reaction #	<i>E. coli</i> total RNA Inputs	<i>E. coli</i> biotinylated antisense rRNA mixture (23S, 16S, 5S)	Streptavidin coated microspheres
1,2	50 ng	2 μ l	25 μ l
3,4	100 ng	2 μ l	25 μ l
5,6	500 ng	4 μ l	50 μ l
7,8	1000 ng	4 μ l	50 μ l
9,10	2500 ng	8 μ l	100 μ l
11,12	5000 ng	10 μ l	125 μ l
13,14	500 ng	-	50 μ l

The reactions were incubated at 68°C for 10 minutes and then at room temperature for 15 minutes. Each reaction was then added to a fresh 1.5-ml microcentrifuge tube containing the appropriate quantity of washed and resuspended streptavidin-coated microspheres. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow for formation of the biotin-streptavidin complex. The reactions were then spun at 12,000 rpm for 5 minutes using a benchtop centrifuge to pellet the microspheres and the supernatant removed and the RNA purified using the ZYMO RNA Clean-up procedure. Each RNA sample was eluted in 13.5 μ l RNase-free water. The entire RNA sample for each was then converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5- μ l aliquot of each first-strand cDNA sample was used in separate PCR amplification reactions containing primers specific to the terminal 5' regions of 23S (SEQ ID NO: 15 and SEQ ID NO: 16) and 16S rRNA (SEQ ID NO: 19 and SEQ ID NO: 20), the complete 5S rRNA (SEQ ID NO: 13 and SEQ ID NO: 23: 5'-TGCCTGGCAGTTCCTACTCTC) and the *ompA* mRNA (SEQ ID NO: 24: 5'-ACCAGGTTAACCCGTATGTTGGCT and SEQ ID NO: 25: 5'-ACCGATGTTGTTGGTCCACTGGTA) for 20 cycles. A control reaction minus template was also performed for each primer pair. A 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 5.

[0138] Figure 5, Lanes 1-12 show excellent reduction in the amount of 23S (Panel A), 16S (Panel B) and 5S (Panel C) rRNA sequences contained in the different inputs of *E. coli* total RNA in duplicate. Whereas, the *ompA* mRNA (Panel D) transcripts was still clearly detected in all samples and appeared to be unperturbed compared to the corresponding non-subtracted samples (Lanes 7 and 8 versus Lane 13 and 14). Clearly, this method for rRNA subtraction appears to be applicable across a broad range of total RNA inputs with at least a 100-fold dynamic range in this example.

Example 6

Subtraction of fragmented *E. coli* 23S, 16S and 5S rRNA sequences using the *E. coli* biotinylated antisense rRNA mixture

[0139] Ten micrograms of *E. coli* total RNA was fragmented in 1x fragmentation buffer (Ambion) at 65°C for 3 minutes. The fragmented RNA was purified using the ZYMO RNA Clean-up procedure, eluted in 20 μ l RNase-free water and the concentration determined by measuring the absorbance at 260 nm with a spectrophotometer. A 300 ng aliquot of the fragmented RNA compared to the intact RNA was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 6A.

[0140] Figure 6A, Lane 1 shows the intact *E. coli* total RNA and Lane 2 shows the corresponding fragmented total RNA. The intact 23S and 16S rRNA bands were no longer visible in the fragmented sample contained in Lane 2.

[0141] Next, ribosomal RNA subtraction was applied to a 2.5 μ g sample of the fragmented *E. coli* total RNA sample comparing it to a similar amount of intact *E. coli* total RNA. The following four reactions were assembled - reactions #1 and #2 contained 2.5 μ g of intact *E. coli* total RNA each and reactions #3 and #4 contained 2.5 μ g of the fragmented *E. coli* total RNA each. Next, to each of reactions #2 and #4, 8 μ l of the *E. coli* biotinylated antisense rRNA mixture was added and the final reaction volume adjusted to 40 μ l in 1x hybridization buffer. The reactions were incubated at 68°C for 10 minutes and then at room temperature for 15 minutes. Each reaction was transferred to a fresh 1.5-ml microcentrifuge tube containing 50 μ l of washed streptavidin

microspheres. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow for formation of the biotin-streptavidin complex. The reactions were spun at 12,000 rpm for 5 minutes on a benchtop centrifuge and each supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The RNA contained in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 µl RNase-free water. Fifty percent of each purified sample was analyzed by ethidium bromide stained agarose gel electrophoresis and Northern blot analysis as shown in Figure 6B.

[0142] Figure 6B, Panel 1 (Lanes 2 and 4) show the RNA remaining following rRNA subtraction. In the case of the intact total RNA, it is clearly evident that the 23S and 16S rRNA bands are removed (Lane 2 *versus* Lane 1). Whereas, for the fragmented total RNA, there is an overall reduction in the amount of RNA following subtraction as would be expected if the fragments of rRNA were also subtracted (Lanes 4 *versus* Lane 3). The Northern blot analysis for *ompA* mRNA (Panel 2) and 23S (Panel 3), 16S (Panel 4) and 5S (Panel 5) rRNA clearly show similar and excellent subtraction of the different rRNA sequences in both the intact total RNA (Panel 3, 4, and 5 - Lane 2) and fragmented total RNA (Panel 3, 4, and 5 - Lane 4) compared to the respective non-subtracted samples (Panel 3, 4, and 5 - Lane 1 and 3). At the same time, the amount of *ompA* mRNA sequence appears to be unperturbed by the subtraction process for both the intact total RNA (Panel 2 - Lane 2 *versus* Lane 1) and the fragmented total RNA (Panel 2 - Lane 4 *versus* Lane 3).

[0143] In addition, the remaining 50% of each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5-µl aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' regions of 23S (SEQ ID NO: 15 and SEQ ID NO: 16) and 16S rRNA (SEQ ID NO: 19 and SEQ ID NO: 20), the complete 5S rRNA (SEQ ID NO: 13 and SEQ ID NO: 23) and the *ompA* mRNA (SEQ ID NO: 24 and SEQ ID NO: 25) for 20 cycles. A 5-µl aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 6C.

[0144] Figure 6C shows that by RT-PCR analysis, there was excellent subtraction of 23S (Panel 2), 16S (Panel 3) and 5S rRNA (Panel 4) for both the intact (Lane 2 *versus* Lane 1) and the fragmented (Lane 4 *versus* Lane 3) *E. coli* total RNA. Whereas, the *ompA* mRNA was equally preserved in both cases before and after subtraction (Panel 1).

Example 7

Specific hybridization of human, mouse and rat 28S and 18S rRNA to human biotinylated antisense 28S and 18S rRNA, and their removal with streptavidin microspheres

[0145] Intact total RNA (500 ng) samples for human (reactions #4 and #5), mouse (reactions #7 and #8) and rat (reactions #10 and #11) were mixed with either 2.1 µg of biotinylated antisense 28S rRNA or 1.0 µg of biotinylated antisense 18S rRNA in 1x hybridization buffer in a final volume of 20 µl. Corresponding intact total RNA samples (reactions #3, #6 and #9) and biotinylated antisense rRNA alone (#1 and #2) were included as controls. All reactions were incubated at 68°C for 10 minutes followed by room temperature for 15 minutes. The RNA contained in each sample was then purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 µl RNase-free water. The entire sample for each was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 7A.

[0146] Figure 7A, Lanes 1 and 2 show the individual biotinylated antisense 23S and 18S rRNA, respectively. Lanes 3, 6 and 9 show the individual intact total RNA corresponding to human, mouse and rat, respectively. Lanes 4, 7 and 10 show the human, mouse and rat total RNA samples, respectively following hybridization to the biotinylated antisense 28S rRNA. Clearly, the 28S rRNA band in the different total RNA samples have hybridized to its complementary antisense rRNA since it now migrates with a high molecular weight higher whereas, the 18S rRNA remained unperturbed. When the different total RNA samples were then hybridized to the biotinylated antisense 18S rRNA, clearly the 18S rRNA band now migrated with a higher molecular weight (Lane 5, 8 and 11) and now, the 28S band was unperturbed. It's evident from these results that each biotinylated antisense rRNA was hybridizing specifically to its complementary target and equally well for human, mouse and rat that share at least 99% sequence homology.

[0147] Next, a biotinylated antisense rRNA mixture for human comprising approximately 1.2 pmoles each of 28S and 18S antisense rRNA in a final volume of 4 µl of RNase-free water was prepared using the *in vitro* synthesized RNA made as described in Example 1. Intact total RNA (2.5 µg) from human (HeLa), mouse (3T3) and rat (NRK) were each mixed with 8 µl of the

biotinylated antisense rRNA solution in 1x hybridization buffer in a final volume of 40 μ l. A control reaction containing only 2.5 μ g of the corresponding intact total RNA was also included. The reactions were incubated at 68°C for 10 minutes followed by room temperature for 15 minutes. Each reaction was transferred to a fresh 1.5-ml microcentrifuge tube containing 100 μ l of washed streptavidin microspheres, respectively. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex and then 37°C for 5 minutes. The reactions were spun at 14,000 rpm for 5 minutes on a benchtop centrifuge and each supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The RNA contained in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 μ l RNase-free water. Fifty percent of each purified RNA sample was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 7B.

[0148] Figure 7B, Lanes 1, 3 and 5 show the individual total RNA without any 28S and 18S rRNA subtraction, whereas, Lanes 2, 4, and 6 show the corresponding RNA remaining following 28S and 18S rRNA subtraction. It is clear from these results that a majority of the 28S and 18S rRNA bands were removed following the subtraction procedure as described.

Example 8

Subtraction of 28S, 18S, 5.8S and 5S rRNA from varying amounts of intact human total RNA using a human biotinylated antisense rRNA mixture

[0149] A biotinylated antisense rRNA mixture for human comprising approximately 1.2 pmoles each of 28S, 18S, 5.8S and 5S antisense rRNA in a final volume of 4 μ l of RNase-free water was prepared using the *in vitro* synthesized RNA made as described in Example 1. Ribosomal RNA subtraction reactions comprising 100 ng (reaction #1), 500 ng (reaction #2) and 5.0 μ g (reaction #3) of intact HeLa total RNA with 2 μ l, 4 μ l and 8 μ l of the biotinylated human antisense rRNA mixture, respectively were prepared in 1x hybridization buffer in a final volume of 20 μ l. A control reaction (reaction #4) with only 500 ng of the intact HeLa total RNA was also included. The reactions were incubated at 68°C for 10 minutes followed by room temperature for 15 minutes. Each reaction was transferred to a fresh 1.5-ml microcentrifuge tube containing 25 μ l, 50 μ l, 125 μ l and 50 μ l of washed streptavidin microspheres, respectively. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex and then 37°C for 5 minutes. The reactions were spun at 14,000 rpm for 5 minutes on a benchtop centrifuge and each supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The RNA contained in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 μ l RNase-free water. Next, each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5- μ l aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' and 3' regions of 28S (SEQ ID NO: 26: 5'-CTCAGTAACGGCGAGTGAAC, SEQ ID NO: 27: 5'-GCCTCGATCAGAAGGACTTG, SEQ ID NO: 28: 5'-TACCACAGGGA TAACTGGCT and SEQ ID NO: 29: 5'-TAGGAAGAGCCGACATCGAA) and 18S rRNA (SEQ ID NO: 30: 5'-CCTACCTGGTTGATCCTGCC, SEQ ID NO: 31: 5'-CCAAGTAGGAGAGGAGCGAG, SEQ. ID. No. 32: 5'-CCCAGTAAGTGC GGGTCATA and SEQ ID NO: 33: 5'-TCACTAAACCATCCAATCGGTAGTA) the complete 5.8S rRNA (SEQ ID NO: 3 and SEQ ID NO: 34: 5'-GATCCTTCGCGAGGT TCACCTAC), the complete 5S rRNA (SEQ ID NO: 5 and SEQ ID NO: 35: 5'-AAGCCTACAGCACCCGGTATTC) and 5' GAPDH mRNA (SEQ ID NO: 36: 5'-TCGACAGTCAGCCGCATCTTCTTT and SEQ ID NO: 37: 5'-ACCAAATCCGTTG ACTCCGACCTT) for 20 cycles. A control reaction for each primer pair minus template was included. A 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis shown in Figure 8.

[0150] Figure 8, Lanes 1,2 and 3 show the RT-PCR results for 28S rRNA (Panel A), 18S rRNA (Panel B), 5.8S rRNA (Panel C), 5S rRNA (Panel D) and GAPDH mRNA (Panel E) for the 100 ng, 500 ng and 5.0 μ g inputs of total RNA subtracted with the biotinylated human antisense rRNA mixture, respectively. Lane 4 shows the PCR for the 500 ng non-subtracted total RNA input. It is evident from the results that there is excellent subtraction over a broad range of total RNA inputs for the different ribosomal RNA sequences using the biotinylated antisense rRNA mixture (Lanes 1, 2 and 3 *versus* Lane 4).

Example 9

Subtraction of human rRNA sequences exhibited by various levels of fragmented human total RNA,

[0151] HeLa intact total RNA was fragmented in 1x fragmentation buffer (Ambion) for 1, 2 and 3 minutes. The RNA samples were purified using the ZYMO RNA Clean-up procedure, eluted RNase-free water and the concentration determined at an absorbance of 260 nm using a spectrophotometer.

[0152] Duplicate reactions comprising either 2.5 µg of the intact (reactions # and #2) or fragmented (reactions #3, #4, #5, #6, #7 and #8) total RNA samples were prepared in 1x hybridization buffer. To one of each duplicate reaction (#2, #4, #6 and #8), 8 µl of the biotinylated human antisense rRNA mixture was added and the final volume for each reaction adjusted to 40 µl. The reactions were incubated at 68°C for 10 minutes followed by room temperature for 15 minutes. Each reaction was transferred to a fresh 1.5-ml microcentrifuge tube containing 100 µl of washed streptavidin microspheres. The reactions were further incubated at room temperature for 15 minutes with occasional gentle mixing (3-4 minutes) in order to allow formation of the biotin-streptavidin complex and then 37°C for 5 minutes. The reactions were spun at 14,000 rpm for 5 minutes on a benchtop centrifuge and each supernatant transferred to a fresh 1.5-ml microcentrifuge tube. The RNA contained in the supernatants were purified using the ZYMO RNA Clean-up procedure and eluted in 13.5 µl RNase-free water. Fifty percent of each purified RNA sample was analyzed by ethidium bromide stained gel electrophoresis as shown in Figure 9A.

[0153] Figure 9A, Lanes 4, 6 and 8, containing the fragmented total RNA and biotinylated antisense rRNA show a significant reduction in the amount of RNA as a result of subtraction compared to the respective controls (Lanes 3, 5 and 7). In Lane 2 compared to Lane 1 with the intact total RNA, the full-length rRNA bands were no longer visible.

[0154] Next, the remaining 50% of each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5-µl aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' regions of 28S (SEQ ID NO: 26 and SEQ ID NO: 27) and 18S rRNA (SEQ ID NO: 30 and SEQ ID NO: 31), the complete 5.8S rRNA (SEQ ID NO: 3 and SEQ ID NO: 34), the complete 5S rRNA (SEQ ID NO: 5 and SEQ ID NO: 35) and 5' GAPDH mRNA (SEQ ID NO: 36 and SEQ ID NO: 37) for 20 cycles. A control reaction minus template was also performed for each primer pair. A 5-µl aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis shown in Figure 9B.

[0155] Figure 9B, Lanes 2, 4, 6 and 8 show the PCR results for 28S rRNA (Panel 1), 18S rRNA (Panel 2), 5.8S rRNA (Panel 3), 5S rRNA (Panel 4) and GAPDH mRNA (Panel 5) for the intact, and fragmented (1, 2 and 3 minutes) total RNA, respectively following subtraction. The corresponding non-subtracted control reactions are shown in Lanes 1, 3, 5 and 7. It is evident from the results that there is excellent subtraction over a broad range of fragmented total RNA inputs similar to the intact total RNA for the different ribosomal RNA sequences using the biotinylated antisense rRNA mixture.

Example 10

Comparison of *E. coli* rRNA subtraction using the method of the present invention and the MICROBExpress™ method

[0156] A kit that uses conserved rRNA (23S and 16S) oligonucleotide sequences to subtract the corresponding rRNA from inputs of 2 µg - 10 µg from prokaryotic total RNA was purchased from Ambion for comparison to the exemplary methods in these Examples. An 15 µg aliquot of the intact control *E. coli* total RNA supplied in the MICROBExpress™ kit was first fragmented in 1x fragmentation buffer (Ambion) at 65°C for 2 minutes, purified using the ZYMO RNA Clean-up procedure, eluted RNase-free water and the concentration determined at an absorbance of 260 nm using a spectrophotometer. An equal aliquot (2.5 µg) of either the intact or fragmented *E. coli* total RNA was then used in subtraction reactions following either the exemplary method described in Example 9 ("Exemplary Method") for the appropriate RNA samples or the MICROBExpress™ method exactly as described by the manufacturer (Ambion). Control reactions representing no rRNA subtraction were included for both methods. Each RNA sample was then purified by EtOH precipitation as described for the MICROBExpress™ procedure and resuspended in 12 µl RNase-free water. Fifty percent of each purified RNA sample was analyzed by ethidium bromide stained agarose gel electrophoresis and Northern blot as shown in Figure 10A.

[0157] Figure 10A, Lanes 1, 2 and 3, 4 show the relative rRNA subtraction from intact and fragmented total RNA using the Exemplary Method and the MICROBExpress™ method for 23S rRNA (Panel 2), 16S rRNA (Panel 3) and 5S rRNA (Panel 4). Clearly, for both the intact and fragmented total RNA samples, the different rRNA sequences were subtracted significantly better

using the Exemplary Method invention (Exemplary Method - Lanes 2 and 4) compared to the MICROBExpress method (MICROBExpress™ Panel - Lanes 2 and 4). It appears that even the control intact RNA supplied with the MICROBExpress™ contain a level of fragmented rRNA sequences that were clearly not removed by the MicroBExpress method (MICROBExpress™ Panel - Lane 2) but were clearly removed using the Exemplary Method described herein (Exemplary Method, Panel - Lane 2). In addition, Panel 1 shows that the *ompA* mRNA sequence remain unperturbed before and after subtraction for both methods.

[0158] Next, the remaining 50% of each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5- μ l aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' region of 23S (SEQ ID NO: 15 and SEQ ID NO: 16) and 16S rRNA (SEQ ID NO: 19 and SEQ ID NO: 20) the complete 5S rRNA (SEQ ID NO: 13 and SEQ ID NO: 23) and *ompA* mRNA (SEQ ID NO: 24 and SEQ ID NO: 25) for 20 cycles. A 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis shown in Figure 10B.

[0159] Figure 10B, Lanes 2 and 4 show the PCR results for 23S rRNA (Panel 2), 16S rRNA (Panel 3), 5S rRNA (Panel 4) and *ompA* mRNA (Panel 1) for the intact and fragmented *E. coli* total RNA, respectively following subtraction using the Exemplary Method and the MICROBExpress™ method. The corresponding non-subtracted control reactions are shown in Lanes 1 and 3. It is evident from the results that there is excellent subtraction of the different rRNA sequences exhibited by both the intact and fragmented total RNA samples using the Exemplary Method (Exemplary Method Panel) compared to the MICROBExpress™ method (MicroExpress Panel). In both cases though, the *ompA* mRNA was similarly maintained.

Example 11

Comparison of Human rRNA subtraction using an Exemplary Method vs. the OLIGOTEX poly A+ mRNA purification method

[0160] A kit that uses oligonucleotide dT to isolate mRNA from eukaryotic total RNA was purchased from Qiagen for comparison to the Exemplary Method described in Example 9 above. When fragmented RNA was used, the RNA was first fragmented in 1x fragmentation buffer (Ambion) at 65°C for 2 minutes. The intact or fragmented RNA was purified using the Zymo RNA Clean-up procedure, eluted RNase-free water and the concentration determined at an absorbance of 260 nm using a spectrophotometer. An equal aliquot (2.5 μ g) of either the intact or fragmented *E. coli* total RNA was then used in rRNA subtraction reactions using the Exemplary Method for the appropriate RNA samples or the Oligotex poly A+ mRNA purification method exactly as described by the manufacturer (Qiagen). Control reactions representing no rRNA subtraction were included for both methods. Each RNA sample was then purified by ethanol precipitation and resuspended in 12 μ l RNase-free water. Fifty percent of each purified RNA sample was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 11A.

[0161] Figure 11A, Lanes 1, 2 and 3, 4 show the relative rRNA removal for intact and fragmented total RNA using the Exemplary Method and the OLIGOTEX poly A+ mRNA purification method. Clearly, for both the intact and fragmented total RNA samples, the different rRNA sequences were visibly removed using the Exemplary Method (Panel 1 - Lanes 2 and 4) and the OLIGOTEX method (Panel 2 - Lanes 2 and 4) compared to the respective controls (Panel 1 - Lanes 1 and 3 and Panel 2 - Lanes 1 and 3). In addition, it appears that there was very little mRNA visibly remaining following the OLIGOTEX method and all of the types of small RNA molecules (tRNA, miRNA etc.) appear to have been eliminated as well (Panel 2 - Lanes 2 and 4).

[0162] Next, the remaining 50% of each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5- μ l aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' and 3' regions of 28S (SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28 and SEQ ID NO: 29) and 18S rRNA (SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32 and SEQ ID NO: 33), the complete 5.8S rRNA (SEQ ID NO: 3 and SEQ ID NO: 34) and the complete 5S rRNA (SEQ ID NO: 5 and SEQ ID NO: 35) for 20 cycles. A 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis shown in Figure 11B.

[0163] Figure 11B, Lanes 2 and 4 show the PCR results for 28S rRNA (Panels 1 and 2), 18S rRNA (Panel 3 and 4), 5.8S rRNA (Panel 5) and 5S rRNA (Panel 6) for the intact and fragmented HeLa total RNA, respectively following rRNA subtraction using the Exemplary Method and the OLIGOTEX method. The corresponding non-subtracted control reactions are shown in Lanes 1 and 3. It is evident from the results that there is overall excellent subtraction of the different rRNA sequences exhibited by both the intact and fragmented total RNA samples using using the Exemplary Method (Exemplary Method Panel) and to the OLIGOTEX method

(OLIGOTEX Panel). In fact, the Exemplary Method appears to be somewhat better than the OLIGOTEX method for both the 28S and 18S rRNA sequences (Panels 1-4), whereas, for the 5.8S and 5S rRNA sequences, the OLIGOTEX method appears to be slightly better (Panel E and F, respectively).

[0164] In addition, PCR primers specific for the terminal 5' and 3' regions of GAPDH (SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38: 5'-CACAGAGGAAGAGAGAGA CCCTCA and SEQ ID NO: 39: 5'-TTGATGGTACATGACAAGGTGCGG) and PGK1 (SEQ ID NO: 40: 5'-GAATCACCGACCTCTCTCCC, SEQ. ID. No. 41: 5'-CGACTCTCATAACGACCCGC, SEQ ID NO: 42: 5'-CCAGAGGTGACCACTTTCAA and SEQ ID NO: 43: 5'-ATGTGGAACAGAGCCTTCCTC) mRNA were used in PCR (GAPDH: 22 cycles and PGK1: 26 cycles) with the random primed cDNA templates and a 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 11C. It is clearly evident from the PCR results that there is a reduction in the amount of mRNA for both GAPDH (Panels 1 and 2) and PGK 1 (Panels 3 and 4) in the OLIGOTEX method for both intact and fragmented total RNA (OLIGOTEX Panel, Lanes 2 and 4, respectively) compared to the non-rRNA- subtracted samples (OLIGOTEX Panel, Lanes 1 and 3, respectively), using the Exemplary Method (Exemplary Method Panel, Lanes 2 and 4). Furthermore, the Exemplary Method did not show any significant loss in the same mRNA sequences for both intact and fragmented total RNA (Exemplary Method Panel, Lanes 2 and 4, respectively) compared to the non-rRNA subtracted samples (Exemplary Method Panel, Lanes 1 and 3, respectively). In addition, the 5' regions of both GAPDH (Panel 1, Lane 4) and PGK1 (Panel 3, Lane 4) for the fragmented total RNA sample were lost with the OLIGOTEX method (OLIGOTEX Panel) since the poly A tail was no longer present but not so for the Exemplary Method (Exemplary Method Panel, Lane 4 (Panel 1 and 3), respectively). Overall, these results demonstrate a clear benefit of the Exemplary Method compared to the OLIGOTEX method.

[0165] Next, PCR primers specific for Poly A- and bimorphic transcripts (SEQ ID NO:

44: 5'-CACGTTTTCTCAGCTGCTTG and SEQ ID NO: 45: 5'-TTCACCTTTTCATC CAAGGC for transcript #1, SEQ ID NO: 46: 5'-GTGTGGTGGTGTGTGCCTAT and SEQ ID NO: 47: 5'-GAGACATGGTCTTGCTCCGT for transcript #3, SEQ ID NO: 48: 5'-TAGCTCAGTGGTAGAGCGCA and SEQ ID NO: 49: 5'-GATTGCTCAGCA GCACGTA for transcript #15 and SEQ ID NO: 50: 5'-CACTTGGGGACACTTTCCAG and SEQ ID NO: 51: 5'-TCAGGGAAAATGAGCCAATC for bimorphic transcript #5 (Poly A-Transcripts Expressed in HeLa Cells Qingfa Wu et. al. (July 2008). PLoS One (www followed by "plosone.org/article/info:doi/10.1371/journal.pone.0002803")) were used in PCR with the random primed cDNA templates (36, 22, 28 and 38 cycles, respectively) and a 5- μ l aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 11D. It is clearly evident from the PCR results that the poly A- and biomorphic transcripts were no longer present following the OLIGOTEX mRNA purification method for both intact and fragmented total RNA (OLIGOTEX Panel, Lanes 2 and 4, respectively) whereas, these sequences were maintained following rRNA subtraction using the Exemplary Method for both intact and fragmented total RNA (Exemplary Method PANEL, Lane 2 and 4, respectively). These results point to another clear benefit of the rRNA subtraction methods described in this application for maintaining a broader spectrum of transcripts allowing for better representation of the cellular content.

Example 12

Comparison of human rRNA subtraction using an Exemplary Method of the present invention vs. the RiboMinus™ Eukaryote Kit for RNA-Seq

[0166] A kit (RiboMinus™ Eukaryote Kit for RNA-Seq) that uses two conserved oligonucleotide sequences for each of 28S, 18S, 5.8S and 5S rRNA to subtract the corresponding rRNA from inputs of 2 μ g - 10 μ g from eukaryotic total RNA was purchased from InVitrogen (Burlington, ON) for comparison to an exemplary method of the present invention. When fragmented total RNA was used, the initial total RNA was first fragmented in 1x fragmentation buffer (Ambion, Austin, TX) at 65°C for 2 minutes. The intact or fragmented total RNA was purified using the ZYMO RNA Clean-up procedure, eluted RNase-free water and the concentration determined at an absorbance of 260 nm using a spectrophotometer. An equal aliquot (2.5 μ g) of either the intact or fragmented HeLa total RNA was then used in rRNA subtraction reactions using the Exemplary Method for the appropriate RNA samples or the RiboMinus™ Eukaryote Kit for RNA-Seq method exactly as described by the manufacturer (InVitrogen). Control reactions representing no rRNA subtraction were included for both methods. Each RNA sample was then purified by ethanol precipitation and resuspended in 12 μ l RNase-free water. Fifty percent of each purified RNA sample was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 12A.

[0167] Figure 12A, Lanes 1,2 and 3, 4 show the relative rRNA removal for intact and fragmented total RNA using the Exemplary

Method and the RiboMinus™ Eukaryote Kit for RNA-Seq method. Clearly, for both the intact and fragmented total RNA samples, the different rRNA sequences were visibly removed using the Exemplary Method (Exemplary method - Lanes 2 and 4) whereas for the RiboMinus™ Eukaryote Kit for RNA-Seq method, only the intact total RNA showed reduction of the rRNA sequences (RiboMinus™ method - Lanes 2) compared to the respective non-rRNA-subtracted samples (Exemplary method - Lanes 1 and 3 and RiboMinus™ method - Lane 1). There appeared to be only minimal reduction of the rRNA-subtracted fragmented total RNA sample using the RiboMinus™ Eukaryote Kit for RNA-Seq as judged by the ethidium bromide stained intensity (RiboMinus™ method - Lane 4) compared to the non-rRNA- subtracted sample (RiboMinus™ method - Lane 3) likely due to the fact that those ribosomal fragments that are not represented by the consensus oligonucleotide sequences in the kit would therefore not be removed.

[0168] Next, the remaining 50% of each purified RNA sample was converted to first-strand cDNA in a standard reverse transcriptase reaction containing random hexamers and purified using the Qiaquick PCR purification kit as recommended by the manufacturer (Qiagen). A 5-µl aliquot of each was then used in separate PCR amplification reactions containing primers specific to the terminal 5' and 3' regions of 28S (SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28 and SEQ ID NO: 29) and 18S rRNA (SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32 and SEQ ID NO: 33), the complete 5.8S rRNA (SEQ ID NO: 3 and SEQ ID NO: 34) and the complete 5S rRNA (SEQ ID NO: 5 and SEQ ID NO: 35) for 20 cycles. A 5-µl aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis shown in Figure 12B.

[0169] Figure 12B, Lanes 2 and 4 show the PCR results for 28S rRNA (Panels 1 and 2), 18S rRNA (Panel 3 and 4), 5.8S rRNA (Panel 5) and 5S rRNA (Panel 6) for the intact and fragmented HeLa total RNA, respectively following rRNA subtraction using the Exemplary Method and the RiboMinus™ Eukaryote Kit for RNA-Seq method. The corresponding non-rRNA-subtracted control reactions are shown in Lanes 1 and 3. It is evident from the results that there is overall excellent subtraction of the 28S rRNA (Panels 1 and 2) and 18S rRNA (Panel 3 and 4) sequences contained in both the intact and fragmented total RNA samples using the Exemplary Method (Exemplary Method Panel) but not the RiboMinus™ Eukaryote Kit for RNA-Seq method (RiboMinus™ Panel). However, both methods showed similar removal of the 5.8S (Panel 5) and 5S rRNA sequences (Panel 6).

[0170] In addition, PCR primers specific for the terminal 5' and 3' regions of GAPDH (SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38: 5'-CACAAAGAGGAAGAGAGAGAG CCCTCA and SEQ ID NO: 39: 5'-TTGATGGTACATGACAAGGTGCGG) mRNA was used in PCR (GAPDH: 24 cycles) with the random primed cDNA templates and a 5-µl aliquot of each PCR reaction was analyzed by ethidium bromide stained agarose gel electrophoresis as shown in Figure 12C. It is clearly evident from the PCR results that there is no obvious reduction in the amount of mRNA for GAPDH (Panels 1 and 2) for both methods with either the intact or fragmented HeLa total RNA (Lanes 2 and 4) compared to the non-rRNA- subtracted samples (Lanes 1 and 3), respectively. Overall, these results demonstrate a clear benefit of the Exemplary Method in the removal of 18S and 28S rRNA sequences independent of the state (intact or fragmented) of the total RNA sample compared to the RiboMinus™ Eukaryote Kit for RNA-Seq method. In fact, even for the intact total RNA sample, the Exemplary method appeared to be more efficient at removal of 18S and 28S rRNA sequences likely due, at least in part, to the fact that there is invariably some degree of fragmented rRNA in even a most intact total RNA sample that would not be removed by the RiboMinus™ method. In fact, the RiboMinus™ Eukaryote Kit for RNA-Seq method clearly states that this method requires use of only high-quality total RNA samples.

[0171] Next, the random primed cDNA samples were analyzed by QPCR as follows: The cDNA samples were diluted to anywhere from 2-fold to 10-fold depending on the starting amount. Then, 1 µl of each dilution was added to a 25 µl qPCR reaction comprising 1x FS PreMix E (GREEN), 12.5 pmole of forward and reverse PCR primers and 1 unit of FS Enzyme Mix. The cycling conditions were 98°C for 2 minutes, followed by 40-45 cycles of 98°C for 5 seconds, 60°C for 15 seconds and 72°C for 30 seconds using the Bio-Rad iGyler (Bio-Rad, Hercules, CA). The following QPCR primer pairs were used for 18S (SEQ. ID. No. 30: SEQ. ID. No. 31, SEQ. ID. No. 52: 5'-CTTAGAGGGACAAGTGGCG, SEQ. ID. No. 53: 5'-GTAGGGTAGGCACACGCTGA, SEQ. ID. No. 54: 5'-GAAACTTAAAGGAATTGACGGAAG, SEQ. ID. No. 55: 5'-GAATCGAGAAAGAGCTATCAATC, SEQ. ID. No. 56: 5'-CGATTGGATGGTTAGTGAGG and SEQ. ID. No. 57: 5'-CCTTGTTACGACTTTTACTTCCTCTAG), 28S (SEQ. ID. No. 58: 5'-GCCGAAACGATCTCAACCTA, SEQ. ID. No. 59: 5'-CGCCAGTTCTGCTTACCAAA, SEQ. ID. No. 60: 5'-CGGACCAAGGAGTCTAACA, SEQ. ID. No. 61: 5'-CAGGCATAGTTACCATCTTTTCG, SEQ. ID. No. 62: 5'-GGAGAGGGTGTAATCTCGC, SEQ. ID. No. 63: 5'-GCCGACTTCCCTTACCTACA, SEQ. ID. No. 64: 5'-GTGTCAGAAAAGTTACCACAGG, SEQ. ID. No. 65: 5'-GGCGAATTCTGCTTACAATGATAG, SEQ. ID. No. 66: 5'-GGGAGTAAGTATGACTCTCTTAAGGT, SEQ. ID. No. 67: 5'-TTGGCTGTGGTTTCGCTGGAT, SEQ. ID. No. 68: 5'-GTGAACAGCAGTTGAACATGG and SEQ. ID. No. 69: 5'-CTTCACAAAGAAAGAGAACTCTCCC), 5.8S (SEQ. ID. No. 70: 5'-CGACTCTTAGCGGTGGATCA and SEQ. ID. No. 71: 5'-AAGCGACGCTCAGACAG) and 5S (SEQ. ID. NO. 5 and SEQ. ID. No. 72: 5'-AAAGCCTACAGCACCCGGTATTC) rRNA sequences.

[0172] The QPCR results are shown in Table 2 below:

TABLE 2

QPCR Primer Sets	Percentage Ribosomal RNA Reduction			
	Exemplary Method		RiboMinus™ Method	
	Intact Hela total RNA (2.5 µg)	Fragmented Hela total RNA (2.5 µg)	Intact Hela total RNA (2.5 µg)	Fragmented Hela total RNA (2.5 µg)
1BS.5' (nt 100 - nt 247)	> 99.9%	> 99.9%	93.30%	50%
18S.3' (nt 1544 - nt 1663)	> 99.9%	> 99.9%	97.10%	81.10%
18S.F3/R3 (nt 1288 - nt 1417)	> 99.9%	> 99.9%	96.40%	34%
18S.F4/R4 (nt 1818 - nt 1937)	> 99.9%	> 99.9%	97.40%	88.30%
28S.3.5K (nt 1748 - nt 1867)	> 99.9%	> 99.9%	85.60%	0%
28S.#2 (nt 1324 - nt 1530)	> 99.9%	99.60%	92.80%	90.50%
28S.F5/R5 (nt 4341 - nt 4456)	> 99.9%	> 99.9%	93.80%	78.20%
28S.5#2 (nt 2740 - nt 2843)	> 99.9%	> 99.9%	92.80%	29.30%
28S.F3/R3 (nt 2401 - nt 2630)	> 99.9%	> 99.9%	83.50%	0%
28S.F41R4 (nt 3732 - nt 3851)	> 99.9%	> 99.9%	82.30%	0%
5.8S (nt 1 - nt 157)	> 99.9%	> 99.9%	99.40%	99.60%
5S (nt 1- nt 121)	96.80%	97.80%	91.80%	88.90%

[0173] Clearly, from the results in Table 2, the exemplary method was significantly more efficient at removal of all ribosomal RNA sequences independent of the primer sets used for both 28S and 18S rRNA sequences with either intact or fragmented total RNA. Whereas, for the RiboMinus™ method with the fragmented RNA, there was little or no ribosomal subtraction depending on the location of the different primer sets for both 28S and 18S rRNA sequences.

Example 13

Significant reduction of rRNA background and improvement in uniquely mappable reads using rRNA removal method disclosed herein compared to the RiboMinus™ method

[0174] Intact and partially fragmented Universal Human Reference RNA (UHRR) (2 x 2.5 µg each) were treated with either the method as described in Example 9 above or the RiboMinus™ Eukaryote Kit for RNA-Seq rRNA removal kit. The respective rRNA-depleted samples were pooled and, for each, Illumina RNA-Seq libraries were prepared in triplicate using rRNA-depleted RNA from the equivalent of 1 µg total RNA. Replicates of the respective RNA-Seq libraries were pooled and sequencing was performed using Illumina® GAIIx next generation sequencer with 36-nt reads. The data were analyzed using Illumina's Pipeline Eland_rna Module and CASAVA Software as well as the TopHat Software for mapping splice junctions (see <http://tophat.cbcb.umd.edu/index.html>). The mapping results showed that rRNA background was significantly reduced by the methods described in Example 9 compared to the RiboMinus™ method (see Table 3 below). In addition, the uniquely mappable sequences not including rRNA sequences were significantly increased (Table 3) in the sample treated with the methods as described in Example 9 compared to the RiboMinus™. Furthermore, for the fragmented samples, the Example 9 methods considerably outperformed the competitive kit, both in terms of reducing rRNA background and improving the uniquely mappable sequences (Table 3).

TABLE 3

Total RNA Sample	rRNA Removal Method	% rRNA Background	% Uniquely Mappable Sequences
Intact UHRR	Example 9	1.4%	58.1%
Intact UHRR	RiboMinus™	18.4%	51.4%
Fragmented UHRR	Example 9	2.1%	59.6%
Fragmented UHRR	RiboMinus™	63.3%	24.6%

Table 3 show that the rRNA removal methods described in Example 9 significantly reduces rRNA background and improves RNA-Seq results. Intact and partially fragmented Universal Human Reference RNA (UHRR) was treated with either the method from Example 9 or the commercial kit (RiboMinus™ method). The rRNA-depleted RNAs were then used to prepare RNA-Seq libraries that were sequenced on an Illumina® GAllx sequencer.

Example 14

Prophetic Example for Plant rRNA Removal

[0175] Plants generally comprises rRNA sequences corresponding to chloroplast, mitochondrial and of nuclear origins. For chloroplast origin, the rRNA comprise 23S, 16S, 5S and 4.5S sequences (e.g. *Arabidopsis thaliana*; Accession # AP000423.1); for mitochondrial origin, the rRNA comprise 18S and 5S sequences (e.g., *Arabidopsis thaliana*; Accession # Y08501.2) and for nuclear origin, the rRNA comprise 25/26S, 17/18S and 5.8S sequences (e.g. *Arabidopsis thaliana*; AC006837.16). PCR templates corresponding to each of the plant rRNA sequence could be synthesized (in full or in part), as well as the respective biotinylated rRNA sequences as described in Example 1 for *E. coli*.

[0176] The respective biotinylated antisense could then be mixed either in a single ratio or several different ratios in order to efficiently remove all rRNA sequences from the different plant tissues (e.g. leaf, root, stem etc.) where it is known that the representation of the different rRNA are present in varying amounts, especially dependent on the chloroplast content. It is contemplated that the various plant rRNA sequences will be effectively removed similar to that described for human and *E. coli* rRNA sequences as in Examples 5-12 herein.

[0177] Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

Preferred embodiments are outlined in the following paragraphs

EMBODIMENTS

[0178]

1. 1. A method of generating a rRNA-depleted sample from an initial sample comprising:
 1. a) providing;
 1. i) an initial sample comprising RNA molecules derived from at least one eukaryotic organism or species, wherein said RNA molecules comprise rRNA molecules and non-rRNA RNA molecules;
 2. ii) a composition comprising antisense rRNA molecules complementary to substantially all of the sequence of at least one rRNA molecule selected from the group consisting of: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules and 12S and 16S eukaryotic mitochondrial rRNA molecules, wherein said antisense rRNA molecules comprise affinity-tags at a ratio of at least one affinity-tag per every 10 nucleobases of said antisense rRNA molecules; and
 3. iii) a binding matrix comprising affinity-tag-binding molecules;
 2. b) contacting said initial sample with said composition comprising antisense rRNA molecules under conditions such that at least some of said affinity-tagged antisense rRNA molecules and at least some of said rRNA molecules form double-stranded rRNA hybrids, thereby generating a treated sample;
 3. c) contacting said treated sample with said binding matrix under conditions wherein the ratio of said affinity-tag-binding molecules to said affinity tags present in the antisense rRNA molecules used in step b) is at least 8 to 1, such

that at least a portion of said double-stranded rRNA hybrids bind to said binding matrix and are removed from said treated sample, thereby generating an rRNA-depleted sample, wherein said rRNA-depleted sample comprises at least a portion of said non-rRNA RNA molecules present in said initial sample and is substantially depleted or free of rRNA molecules that exhibit sequences exhibited by said at least one rRNA molecule.

2. 2. The method of Paragraph 1, wherein said at least one eukaryotic organism or species is selected from the group consisting of a human, an animal, a plant, and a fungal organism or species.
3. 3. The method of Paragraph 2, wherein said RNA molecules in said initial sample and said at least one rRNA molecule are derived from the same eukaryotic organism or species.
4. 4. The method of Paragraph 1, wherein said RNA molecules are from a first eukaryotic organism or species, and wherein said at least one rRNA molecule is from a second eukaryotic organism for species.
5. 5. The method of Paragraph 4, wherein one of said first or second organisms or species is *Homo sapiens*, and wherein the other of said first or second organisms or species is a non-human mammal.
6. 6. The method of any of Paragraphs 1-5, wherein said affinity-tags are associated with at least one nucleobase selected from the group consisting of: adenine (A), cytosine (C), guanine (G) and uracil (U); and wherein said affinity-tags are present on said antisense rRNA molecules at a ratio of at least one affinity-tag per every 8 nucleobases of said antisense rRNA molecules; and/or wherein the ratio of said affinity-tag-binding molecules to said affinity tags during said contacting in step b) is at least 10 to 1.
7. 7. The method of any of Paragraphs 1-6, wherein the number of molecules that exhibit the sequence of said at least one rRNA molecule in said rRNA-depleted sample is depleted by 99.0% relative to the number of molecules that exhibit the sequence of said at least one rRNA molecule in said initial sample.
8. 8. The method of any of Paragraphs 1-7, wherein said at least one rRNA molecule includes either both said 28S rRNA molecules and said 18S rRNA molecules, or both said 25S or 26S rRNA molecules and said 16S rRNA molecules.
9. 9. The method of any of Paragraphs 1-8, wherein said at least one rRNA molecule additionally includes both said 5.8 S rRNA and said 5S rRNA molecules, and/or wherein said at least one rRNA molecule additionally includes both said 12S and said 16S eukaryotic mitochondrial rRNA molecules, and/or wherein said at least one rRNA molecule is additionally selected from the group consisting of 16S, 23S, 4.5S and 5S eukaryotic chloroplast rRNA molecules.
10. 10. The method of any of Paragraphs 1-9, wherein said initial sample comprising RNA molecules derived from a eukaryotic organism or species additionally comprises RNA molecules derived from at least one prokaryotic species, wherein said prokaryotic RNA molecules comprise rRNA molecules and non-rRNA RNA molecules, and wherein said method comprises:

providing a composition that additionally comprises affinity-tagged antisense rRNA molecules complementary to substantially all of the sequence of at least one rRNA molecule selected from the group consisting of 23S, 16S, and 5S prokaryotic rRNA molecules; and

contacting said initial sample with said composition that additionally comprises prokaryotic affinity-tagged antisense rRNA molecules under conditions such that at least some of said affinity-tagged antisense rRNA molecules and at least some of said prokaryotic rRNA molecules form double-stranded rRNA hybrids thereby generating a treated sample; and

contacting said treated sample with said binding matrix in step c) under conditions wherein said rRNA-depleted sample also comprises at least a portion of said non-rRNA prokaryotic RNA molecules present in said initial sample and is substantially depleted or free of prokaryotic rRNA molecules that exhibit sequences exhibited by said at least one prokaryotic rRNA molecule.
11. 11. The method of any of Paragraphs 1-10, wherein at least a portion of said RNA molecules in said initial sample are fragmented.
12. 12. The method of any of Paragraphs 1-11, wherein said RNA molecules in said initial sample comprise RNA extracted, isolated, or purified from a source selected from the group consisting of: a tissue sample, a cell sample, a paraffin-embedded sample, a paraffin-embedded formalin-fixed (FFPE) sample, and an environmental sample consisting of soil, water, growth medium, or a biological fluid or specimen.
13. 13. The method of any of Paragraphs 1-12, wherein said conditions in step b) comprise incubating in the presence of a hybridization buffer at a temperate of about 60-75 °C for a first time period and incubating at about room temperature for a second time period, and wherein said binding matrix in step c) comprises a plurality of individual particles or a binding column, and wherein said conditions in step c) include at least occasional mixing at room temperature and/or at 35-60 °C.
14. 14. The method of any of Paragraphs 1-13, wherein the affinity-tag comprises biotin.
15. 15. A composition comprising antisense rRNA molecules complementary to substantially all of the sequence exhibited by at least one rRNA molecule selected from the group consisting of 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic

rRNA molecules, 12S and 16S eukaryotic mitochondrial rRNA molecules, 16S, 23S, 4.5S and 5S eukaryotic chloroplast rRNA molecules, and 23S, 16S, and 5S prokaryotic rRNA molecules, wherein said antisense rRNA molecules comprise affinity-tags at a ratio of at least one affinity-tag per every 10 nucleobases of said antisense rRNA molecules, or wherein said antisense rRNA molecules comprise affinity-tags at a ratio of at least one affinity-tag per every 8 nucleobases of said antisense rRNA molecules.

16. 16. The composition of Paragraph 15, wherein said antisense rRNA molecules comprise multiple different anti-sense rRNA molecules complementary to rRNA molecules selected from the groups consisting of: at least four rRNA molecules selected from 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules; two rRNA molecules consisting of 12S and 16S eukaryotic mitochondrial rRNA molecules; four rRNA molecules consisting of 16S, 23S, 4.5S and 5S eukaryotic chloroplast rRNA molecules; and three rRNA molecules consisting of 23S, 16S, and 5S prokaryotic rRNA molecules.
17. 17. The composition of any of Paragraphs 15 or 16, additionally comprising RNA molecules derived from at least one organism or species, wherein said RNA molecules comprise rRNA molecules and non-rRNA RNA molecules, wherein at least some of said affinity-tagged antisense rRNA molecules in said composition are present as double-stranded hybrids with at least some of said rRNA molecules.
18. 18. The composition of Paragraph 17, wherein the said RNA molecules comprise RNA molecules derived from multiple organisms or species.
19. 19. The composition of Paragraph 17 or 18, further comprising a binding matrix comprising affinity-tag-binding molecules, wherein the ratio of said affinity-tag-binding molecules to said affinity tags present in the antisense rRNA molecules used in step b) is at least 8 to 1, or wherein the ratio of said affinity-tag-binding molecules to said affinity tags present in the antisense rRNA molecules used in step b) is at least 10 to 1.
20. 20. A composition comprising an rRNA-depleted sample comprising non-rRNA RNA molecules, wherein, compared to a sample that is not rRNA-depleted, said composition is substantially free of rRNA molecules that exhibit the sequence of at least one rRNA molecule selected from the group consisting of: 25S, 26S, 28S, 18S, 5.8S, and 5S eukaryotic cytoplasmic rRNA molecules; 12S and 16S eukaryotic mitochondrial rRNA molecules; 16S, 23S, 4.5S and 5S eukaryotic chloroplast rRNA molecules; and 23S, 16S, and 5S prokaryotic rRNA molecules.

SEQUENCE LISTING

[0179]

<110> Epicentre Technologies Corporation Sooknanan, Roy R.

<120> Methods, Compositions, and Kits for Generating rRNA-Depleted Samples or Isolating rRNA from Samples

<130> EPICE-30968/WO-1/ORD

<150> US 61/234,044

<151> 2009-08-14

<160> 72

<170> Patent In version 3.5

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 1

cctacctacc tgggtgatcc 20

<210> 2

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 2

aattctaata cgactcacta tagggagaga tcctccgca gggtcaccta c 51

<210> 3

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 3

cgactcttag cggtaggatca ctc 23

<210> 4

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 4

aattctaata cgactcacta tagggagaga tcctccgca gggtcaccta c 51

<210> 5

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 5

gtctacggcc ataccaccct gaa 23

<210> 6

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 6

aattctaata cgactcacta tagggagaaa gcctacagca cccggtattc 50

<210> 7

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 7

aagcgactaa gcgtacacgg tgga 24

<210> 8

<211> 52

<212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 8
 aattctaata cgactcacta tagggagatt cctggaagca gggcattgt tg 52

 <210> 9
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 9
 caacaaatgc cctgctcca ggaa 24

 <210> 10
 <211> 53
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 10
 aattctaata cgactcacta tagggagaca cgggtcatta gtaccggta gct 53

 <210> 11
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 11
 agagtttgat cctggctcag 20

 <210> 12
 <211> 50
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 12
 aattctaata cgactcacta tagggagagg aggtgatcca accgcaggtt 50

 <210> 13
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 13
 tgctggcgg cagtagcgcg gt 22

<210> 14
 <211> 50
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 14
 aattctaata cgactcacta tagggagatg cctggcagtt ccctactctc 50

 <210> 15
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 15
 gacgtgctaa tctgcgataa gc 22

 <210> 16
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 16
 atggattcag ttaatgatag tgtgtcg 27

 <210> 17
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 17
 ctgaaagcat ctaagcacga aacttgc 27

 <210> 18
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 18
 cctatcaacg tcgtgtctt caac 24

 <210> 19
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

<400> 19
 gcctaacaca tgcaagtcga ac 22

 <210> 20
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 20
 agctaccgtt tccagtagtt atcc 24

 <210> 21
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 21
 cggaatcgct agtaatcgtg gat 23

 <210> 22
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 22
 tccgaaggt taagctacct actt 24

 <210> 23
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 23
 tgctggcag ttcctactc tc 22

 <210> 24
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 24
 accaggttaa cccgtatggt ggct 24

 <210> 25
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

 <400> 25
 accgatgttg ttgtccact ggta 24

 <210> 26
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 26
 ctcaagtaacg gcgagtgaac 20

 <210> 27
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <900> 27
 gcctcgatca gaaggacttg 20

 <210> 28
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 28
 taccacaggg ataactggct 20

 <210> 29
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 29
 taggaagagd cgacatcgaa 20

 <210> 30
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 30
 cctacctggt tgatcctgcc 20

 <210> 31
 <211> 20

<212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 31
 ccaagtagga gaggagcgag 20

 <210> 32
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 32
 ccagtaagt gcgggtcata 20

 <210> 33
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 33
 tcactaaacc atccaatcgg tagta 25

 <210> 34
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 34
 gatccttcg caggttcacc tac 23

 <210> 35
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 35
 aagcctacag caccggtat tc 22

 <210> 36
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 36
 tcgacagtca gccgcatctt cttt 24

<210> 37
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 37
 accaaatccg ttgactccga cctt 24

<210> 38
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 38
 cacaagagga agagagagac cctca 25

<210> 39
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 39
 ttgatgtac atgacaaggt gcgg 24

<210> 40
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 40
 gaatcacga cctctctccc 20

<210> 41
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 41
 cgactctcat aacgacccgc 20

<210> 42
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 42
 ccagaggtga ccacttcaa 20

 <210> 43
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 43
 atgtggaaca gagccttcct c 21

 <210> 44
 <211> 20
 <212> DNA
 <213> Artificial Sequences

 <220>
 <223> Synthetic

 <400> 44
 cacgttttct cagctgcttg 20

 <210> 45
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 45
 ttcacctttt catccaaggc 20

 <210> 46
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 46
 gtgtggtggt gtgtgcctat 20

 <210> 47
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 47
 gagacatggt ctgctccgt 20

 <210> 48
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

 <400> 48
 tagctcagtg gtagagcgca 20

 <210> 49
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 49
 gatttgctca gcagcacgta 20

 <210> 50
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 50
 cacttgggga cactttccag 20

 <210> 51
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 51
 tcagggaata tgagccaatc 20

 <210> 52
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 52
 ctagaggga caagtggcg 19

 <210> 53
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 53
 gtaggtagg cacacgctga 20

 <210> 54
 <211> 24

<212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 54
 gaaacttaa ggaattgacg gag 24

 <210> 55
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 55
 gaatcgagaa agagctatca atc 23

 <210> 56
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 56
 cgattgcatg gtttagtgag g 21

 <210> 57
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 57
 cctgttacg actttactt cctctag 27

 <210> 58
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 58
 gccgaaacga tctcaaccta 20

 <210> 59
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 59
 cgccagtct gcttacaaa 20

<210> 60
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 60
 cggaccaagg agtctaaca 19

<210> 61
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 61
 caggcatagt tcacatctt tcg 23

<210> 62
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 62
 ggagagggtg taaatctgc 20

<210> 63
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 63
 gccgactcc cttacctaca 20

<210> 64
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 64
 gtgtcagaaa agttaccaca gg 22

<210> 65
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 65
 ggcgattct gctcacaat gatag 25

 <210> 66
 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 66
 gggagtaact atgactctt taagg 26

 <210> 67
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 67
 ttggctgtgg ttgcgtgga t 21

 <210> 68
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 68
 gtgaacagca gttgaacatg g 21

 <210> 69
 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 69
 ctcacaaag aaaagagaac tctcc 26

 <210> 70
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic

 <400> 70
 cgactcttag cgggtgatca 20

 <210> 71
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Synthetic

<400> 71

aagcgacgct cagacag 17

<210> 72

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 72

aaagcctaca gcacccggta ttc 23

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US6123404A \[0001\]](#)
- [WQ2007019444A2 \[0004\]](#)
- [US6562575B \[0079\]](#)
- [US6210891B \[0102\]](#)
- [US6258568B \[0102\]](#)
- [US6833246B \[0103\]](#)
- [US7115400B \[0103\]](#)
- [US6969498B \[0103\]](#)
- [US5912148A \[0104\]](#)
- [US6130073A \[0104\]](#)
- [US7169560B \[0106\]](#)
- [US7282337B \[0106\]](#)
- [US7482126B \[0106\]](#)
- [US7501245B \[0106\] \[0106\]](#)
- [US6818395B \[0106\]](#)
- [US6911345B \[0106\]](#)
- [US7329492B \[0106\]](#)
- [US671956A \[0106\]](#)
- [US781166A \[0106\]](#)
- [US7170050B \[0106\]](#)
- [US7302146B \[0106\]](#)
- [US7313308B \[0106\]](#)
- [US7476503B \[0106\]](#)
- [US6123404B \[0179\]](#)

Non-patent literature cited in the description

- **INGOLIA et al.** Science, 2009, vol. 324, 218-23 [0003]
- **SUSORDILLO** Molecular Biotechnology, 1998, vol. 10, 183-85 [0006]
- **CHAKRAVORTY, S et al.** J. Microbiol. Methods, 2007, vol. 69, 330-339 [0040]
- **MILLAR, BC et al.** Current Issues Mol. Biol., 2007, vol. 9, 21-40 [0040]
- **INGOLIA et al.** Science, 2009, vol. 324, 218-223 [0044]
- Overview of principles of hybridization and the strategy of nucleic acid probe assays **TJJSSEN** Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes Elsevier 19930000 [0083]
- **VOELKERDING et al.** Clinical Chem., 2009, vol. 55, 641-658 [0099] [0102] [0103] [0104] [0106] [0106] [0106]
- **MACLEAN et al.** Nature Rev. Microbiol., vol. 7, 287-296 [0099] [0102] [0103] [0104] [0106] [0106]
- **MARGULIES, M. et al.** Genome Sequencing in Microfabricated High-Density Picolitre Reactors Nature, 2005, vol. 437, 376-80 [0099]
- **MIKKELSEN, T. et al.** Genome-Wide Maps of Chromatin State in Pluripotent and Lineage-Committed Cells Nature, 2007, vol. 448, 553-60 [0099]
- **MCLAUGHLIN, S. et al.** Whole-Genome Resequencing with Short Reads: Accurate Mutation Discovery with Mate Pairs and Quality Values ASHG Annual Meeting, 2007, [0099]
- **SHENDURE J. et al.** Accurate Multiplex Polony Sequencing of an Evolved Bacterial Genome Science, 2005, vol. 309, 1728-32 [0099]
- **HARRIS, T. et al.** Single-Molecule DNA Sequencing of a Viral Genome Science, 2008, vol. 320, 106-9 [0099]
- **SIMEN, B. et al.** Prevalence of Low Abundance Drug Resistant Variants by Ultra Deep Sequencing in Chronically HIV-infected Antiretroviral (ARV) Naive Patients and the Impact on Virologic Outcomes 16th International HIV Drug Resistance Workshop, 2007, [0099]
- **THOMAS, R. et al.** Sensitive Mutation Detection in Heterogeneous Cancer Specimens by Massively Parallel Picoliter Reactor Sequencing Nature Med., 2006, vol. 12, 852-855 [0099]
- **MITSUYA, Y. et al.** Minority Human Immunodeficiency Virus Type 1 Variants in Antiretroviral-Naive Persons with Reverse Transcriptase Codon 215 Revertant Mutations J. Vir., 2008, vol. 82, 10747-10755 [0099]
- **BINLADEN, J. et al.** The Use of Coded PCR Primers Enables High-Throughput Sequencing of Multiple Homolog Amplification Products by 454 Parallel Sequencing PLoS ONE, 2007, vol. 2, e197- [0099]
- **HOFFMANN, C. et al.** DNA Bar Coding and Pyrosequencing to Identify Rare HIV Drug Resistance Mutations Nuc. Acids Res., 2007, vol. 35, e91- [0099]
- **ASTIER et al.** J Am Chem Soc., 2006, vol. 128, 51705-10 [0105]
- **MANIATIS** Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory 19820000 [0114]
- **MADEN** Biochem. J., 1987, vol. 246, 519-527 [0114]
- **WU** PLoS One, 2008, [0165]

PATENTKRAV

1. Sammensætning, der omfatter antisense-rRNA-molekyler, hvor
5 antisense-rRNA-molekylerne omfatter multiple forskellige antisense-rRNA-
molekyler, som er komplementære til i det væsentlige alle de sekvenser, der
udvises af rRNA-molekyler, som er valgt fra grupperne bestående af: mindst fire
rRNA-molekyler, som er valgt blandt 25S, 26S, 28S, 18S, 5.8S og 5S eukaryote
cytoplasmatiske rRNA-molekyler; to rRNA-molekyler bestående af 12S og 16S
10 eukaryote mitokondriske rRNA-molekyler; fire rRNA-molekyler bestående af 16S,
23S, 4.5S og 5S eukaryote chloroplast-rRNA-molekyler; samt tre rRNA-molekyler
bestående af 23S, 16S og 5S prokaryote rRNA-molekyler; hvor antisense-rRNA-
molekylerne omfatter affinitetsmærker i et forhold på mindst ét affinitetsmærke
for hver 10 nukleobaser af antisense-rRNA-molekylerne, eller hvor antisense-
15 rRNA-molekylerne omfatter affinitetsmærker i et forhold på mindst ét
affinitetsmærke for hver 8 nukleobaser af antisense-rRNA-molekylerne.
2. Sammensætningen ifølge krav 1, hvor affinitetsmærket omfatter biotin.
- 20 3. Sammensætningen ifølge et hvilket som helst af kravene 1 eller 2,
hvilken sammensætning yderligere omfatter RNA-molekyler afledt af mindst én
organisme eller species, hvor RNA-molekylerne omfatter rRNA-molekyler og
ikke-rRNA-RNA-molekyler, hvor mindst nogle af de affinitetsmærkede antisense-
rRNA-molekyler i sammensætningen forekommer som dobbeltstrengede
25 hybrider med mindst nogle af rRNA-molekylerne.
4. Sammensætningen ifølge krav 3, hvilken sammensætning yderligere
omfatter en bindingsmatrix, der omfatter affinitetsmærkebindende molekyler, hvor
forholdet mellem de affinitetsmærkebindende molekyler og de i antisense-rRNA-
30 molekylerne forekommende affinitetsmærker er mindst 8 til 1, eller hvor forholdet
mellem de affinitetsmærkebindende molekyler og de i antisense-rRNA-
molekylerne forekommende affinitetsmærker er mindst 10 til 1.

5. Fremgangsmåde til generering af en rRNA-depleteret prøve fra en initial prøve, hvilken fremgangsmåde omfatter, at:

a) der tilvejebringes;

5

i) en initial prøve, der omfatter RNA-molekyler afledt af mindst én eukaryot organisme eller species,

10 hvor RNA-molekylerne omfatter rRNA-molekyler og ikke-rRNA-RNA-molekyler;

ii) sammensætningen ifølge et hvilket som helst af kravene 1-3;

15 iii) en bindingsmatrix, der omfatter affinitetsmærkebindende molekyler;

15

b) den initiale prøve bringes i kontakt med sammensætningen, der omfatter antisense-rRNA-molekyler, under sådanne betingelser, at mindst nogle af de affinitetsmærkede antisense-rRNA-molekyler og mindst nogle af rRNA-molekylerne danner dobbeltstrengede rRNA-hybrider, hvorved der genereres en behandlet prøve;

20

c) den behandlede prøve bringes i kontakt med bindingsmatrixen under betingelser, hvor forholdet mellem de affinitetsmærkebindende molekyler og de affinitetsmærker, som forekommer i de i trin b) anvendte antisense-rRNA-molekyler, er mindst 8 til 1, således at mindst en del af de dobbeltstrengede rRNA-hybrider binder til bindingsmatrixen og fjernes fra den behandlede prøve, hvorved der genereres en rRNA-depleteret prøve, hvor den rRNA-depleterede prøve omfatter mindst en del af de i den initiale prøve forekommende ikke-rRNA-RNA-molekyler og i det væsentlige er depleteret eller fri for rRNA-molekyler, der udviser sekvenser, som udvises af det mindst ene rRNA-molekyle.

30

6. Fremgangsmåde ifølge krav 5, hvor mindst den ene eukaryote organisme eller species er valgt fra gruppen bestående af et menneske, et dyr, en plante og en svampeorganisme eller species.

7. Fremgangsmåden ifølge krav 6, hvor RNA-molekylerne i den initiale prøve og det mindst ene rRNA-molekyle er afledt af den samme eukaryote organisme eller species, eller hvor RNA-molekylerne er fra en første eukaryot organisme eller species, og hvor det mindst ene rRNA-molekyle er fra en anden eukaryot organisme eller species.

8. Fremgangsmåden ifølge et hvilket som helst af kravene 5-7, hvor affinitetsmærkerne er associeret med mindst én nukleobase, der er valgt fra gruppen bestående af: adenin (A), cytosin (C), guanin (G) og uracil (U); og

10 hvor affinitetsmærkerne forekommer på antisense-rRNA-molekylerne i et forhold på mindst ét affinitetsmærke for hver 8 nukleobaser af antisense-rRNA-molekylerne; og/eller

15 hvor forholdet mellem de affinitetsmærkebindende molekyler og affinitetsmærkerne under kontakten i trin b) er mindst 10 til 1.

9. Fremgangsmåden ifølge et hvilket som helst af kravene 5-8, hvor det antal molekyler, der udviser sekvensen for det mindst ene rRNA-molekyle i den rRNA-depleterede prøve, depleteres med 99,0 % i forhold til det antal molekyler, der udviser sekvensen for det mindst ene rRNA-molekyle i den initiale prøve.

10. Fremgangsmåden ifølge et hvilket som helst af kravene 5-9, hvor RNA-molekylerne i den initiale prøve omfatter RNA, der er ekstraheret, isoleret eller oprenset fra en kilde, der er valgt fra gruppen bestående af: en vævsprøve, en celleprøve, en i paraffin indlejret prøve, en i paraffin indlejret formalinfikseret (FFPE) prøve og en miljøprøve bestående af jord, vand, vækstmedium eller en biologisk væske eller prøve.

11. Fremgangsmåden ifølge et hvilket som helst af kravene 5-10, hvor betingelserne i trin b) omfatter inkubering i nærværelse af en hybridiseringsbuffer ved en temperatur på ca. 60-75 °C i et første tidsrum og inkubering ved omtrent stuetemperatur i et andet tidsrum, og hvor bindingsmatrixen i trin c) omfatter en

flerhed af individuelle partikler eller en bindingssøjle, og hvor betingelserne i trin c) indbefatter mindst lejlighedsvis blanding ved stuetemperatur og/eller ved 35-60 °C.

12. Kit, der omfatter

5

a) sammensætningen ifølge et hvilket som helst af kravene 1-3; og

b) en bindingsmatrix, der omfatter affinitetsmærkebindende molekyler, hvor forholdet mellem de affinitetsmærkebindende molekyler og de i antisense-rRNA-

10 molekylerne forekommende affinitetsmærker er mindst 8 til 1, eller hvor forholdet mellem de affinitetsmærkebindende molekyler og de i antisense-rRNA-molekylerne forekommende affinitetsmærker er mindst 10 til 1, og hvor bindingsmatrixen omfatter en flerhed af individuelle partikler, der er valgt blandt nanopartikler, magnetiske partikler og en flerhed af småkugler, såsom makroporøse småkugler;

15

hvor kittet eventuelt yderligere omfatter:

c) en hybridiseringsbuffer;

20 d) en vaskeopløsning;

e) glycogen;

f) natriumacetat;

25

g) RNase-frit vand;

h) RNase-inhibitor;

30 i) en bindingsmatrixresuspenderingsopløsning;

j) en kontrolprøve, der omfatter total-RNA fra en celle- eller vævsprøve og/eller

k) et total-RNA-oprensningsreagens.

35

DRAWINGS

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 2A




FIGURE 2B




FIGURE 3A




FIGURE 3B




FIGURE 3C




FIGURE 4




FIGURE 5




FIGURE 6A




FIGURE 6B




FIGURE 6C




FIGURE 7A




FIGURE 7B




FIGURE 8




FIGURE 9A




FIGURE 9B




FIGURE 10A




FIGURE 10B




FIGURE 11A




FIGURE 11B




FIGURE 11C




FIGURE 11D




FIGURE 12A




FIGURE 12B




FIGURE 12C

