



(22) Date de dépôt/Filing Date: 1990/10/02

(41) Mise à la disp. pub./Open to Public Insp.: 1991/04/04

(62) Demande originale/Original Application: 2 026 703

(30) Priorités/Priorities: 1989/10/03 (8902452) NL;
1990/07/27 (9001711) NL

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 1/00, C12N 15/79, C12N 15/11,
C12N 5/10, C12N 15/09


(71) Demandeurs/Applicants:
AVEVE N.V., BE;
CLOVIS MATTON N.V., BE

(72) Inventeurs/Inventors:
VAN HAUTE, EDDIE, BE;
AMELOOT, PAUL, BE;
DE LAFONTEYNE, JEAN, BE;
FIERS, WALTER, BE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : MANIPULATIONS GENETIQUES AVEC DE L'ADN RECOMBINANT COMPRENANT DES SEQUE CES DE
VIRUS A ARN

(54) Title: GENETIC MANIPULATIONS WITH RECOMBINANT DNA COMPRISING SEQUENCES DERIVED FROM RNA
VIRUS



(57) Abrégé/Abstract:

The invention relates to genetic manipulations of eukaryotic organisms, with recombinant DNA comprising RNA virus derived sequences for protecting such organisms against RNA viruses or enabling inducible or tissue-specific production of foreign proteins/peptides or RNAs. One embodiment of the recombinant DNA according to the invention comprises recombinant DNA,



(57) Abrégé(suite)/Abstract(continued):

comprising two, 12-1250 base pair long, inverted repeat nucleotide sequences with therebetween at least one nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, said RNA virus derived sequence comprising at least cis elements for replication but no gene that codes for viral RNA/RNA polymerase and no gene that codes for viral coat protein. The invention also relates to eukaryotic or prokaryotic cells or organisms which incorporate the recombinant DNA according to the invention. Further the invention relates to a method of protecting such cells or organisms by genetically incorporating recombinant DNA according to the invention.

~~63~~

Title: Genetic manipulations with recombinant DNA comprising sequences derived from RNA virus.

ABSTRACT

The invention relates to genetic manipulations of eukaryotic organisms, with recombinant DNA comprising RNA virus derived sequences for protecting such organisms against RNA viruses or enabling inducible or tissue-specific production of foreign proteins/peptides or RNAs. One embodiment of the recombinant DNA according to the invention comprises recombinant DNA, comprising two, 12-1250 base pair long, inverted repeat nucleotide sequences with therebetween at least one nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, said RNA virus derived sequence comprising at least cis elements for replication but no gene that codes for viral RNA/RNA polymerase and no gene that codes for viral coat protein. The invention also relates to eukaryotic or prokaryotic cells or organisms which incorporate the recombinant DNA according to the invention. Further the invention relates to a method of protecting such cells or organisms by genetically incorporating recombinant DNA according to the invention.

\184-302D

1

Title: Genetic manipulations with recombinant DNA comprising
sequences derived from RNA virus

This application is a divisional application of
Canadian patent application 2,206,703 entitled "Genetic
5 manipulation with recombinant DNA comprising sequences derived
from RNA virus" filed on October 2, 1990.

The invention relates to the field of genetic
engineering by means of DNA recombinant techniques, more
particularly to the field of the genetic engineering of
10 eukaryotic organisms, such as yeasts, fungi and particularly
plants.

The invention is particularly directed to genetic
manipulations which lead to resistance of the host organism
against one or more RNA viruses, and to genetic manipulations
15 which render the host organism capable of an inducible or
tissue-specific production of foreign proteins/peptides or
RNAs.

For such genetic manipulations recombinant DNA is
used comprising one or more sequences derived from RNA virus,
20 or, more particularly, at least one nucleotide sequence which
is derived from RNA virus which for its replication is
dependent on a viral RNA/RNA polymerase (i.e. an RNA-dependent
RNA polymerase, which is sometimes referred to as "replicase").
These RNA viruses may belong to the group of double-stranded
25 RNA viruses (i.e. the genome of the virus consists of double-
stranded RNA), to the group of positive-strand RNA viruses
(i.e. the genome of the virus consists of "sense" or messenger
single-stranded RNA), or to the group of negative-strand RNA
viruses (i.e. the genome of the virus consists of "antisense"
30 single-stranded RNA).

Not all RNA viruses, however, are dependent on a viral RNA/RNA polymerase for their replication. This holds in particular for the viruses belonging to the family of Retroviridae, which for their multiplication are dependent on DNA replication after the genomic RNA is transcribed into DNA by means of reverse transcriptase.

Examples of RNA viruses which for their replication are dependent on a viral RNA/RNA polymerase, are double-stranded viruses from the families of Reoviridae and Birnaviridae, negative-strand viruses from the families of Arenaviridae, Bunyaviridae, Orthomyxoviridae, Paramyxoviridae and Rhabdoviridae, and positive-strand viruses from the families of Togaviridae, Flaviviridae, Coronaviridae, Nodaviridae, Picornaviridae and Caliciviridae. Concrete examples of positive-strand viruses with a plant host are Tobacco Mosaic Virus, Tobacco Necrosis Virus, Brome Mosaic Virus, Cucumber Mosaic Virus, Tobacco Streak Virus, Tobacco Rattle Virus, Cowpea Mosaic Virus, Tomato Black Ring Virus, Potato Y Virus, Turnip Yellow Mosaic Virus, Tomato Bushy Stunt Virus, Southern Bean Mosaic Virus, Barley Yellow Dwarf Virus, Potatovirus X, Sugar Beet Yellow Virus, Carnation Latent Virus, Carnation Ringspot Virus, Barley Stripe Mosaic Virus, Alfalfa Mosaic Virus, Pea Enation Mosaic Virus and Tomato Spotted Wilt Virus.

Subject to the above-mentioned limitation to RNA viruses which for their replication are dependent on a viral RNA/RNA polymerase, the term "RNA virus" as used herein includes viruses and virusoids which for their replication are

dependent on the help of another (helper) virus. As is well known, RNA virus infections may be accompanied by coinfections of, for instance, satellite viruses with a coat protein of their own, satellite RNA which is packed in mixed particles, and virusoids (a small circular RNA genome, packed in mixed particles). The term "RNA virus" as used herein also includes viroids, i.e. autonomous small bare RNA molecules. The further explanation of the invention in the experimental section to follow will be given with reference to an RNA satellite virus, viz. the Satellite Tobacco Necrosis Virus (STNV), a small plant virus (1.85×10^3 kD), which for its replication is entirely dependent on the presence of the helper Tobacco Necrosis Virus (TNV). The RNA genome of STNV contains 1239 nucleotides and codes for a coat protein of 195 amino acids.

The invention comprises incorporating genetic information into the genome of eukaryotes by means of genetic engineering, which information does not as such, nor through transcription products derived therefrom, constitute a burden to the host, and yet accomplishes a very effective protection of the host against viral infections, or enables an inducible or tissue-specific, very efficient production of foreign proteins (or peptides) or RNAs. The genetic information to be incorporated according to the invention comprises an expression-cassette for the host to be transformed containing two, 12-250 base pair long, inverted repeat nucleotide sequences with therebetween at least one nucleotide sequence which is derived from RNA virus which for its replication is dependent on a

viral RNA/RNA polymerase, said RNA virus derived sequence comprising at least cis elements for replication, but no gene that codes for viral RNA/RNA polymerase and no gene that codes for viral coat protein.

5 The genome of RNA viruses which for their replication are dependent on a viral RNA/RNA polymerase comprises various cis elements, i.e. elements or functions which function only for the nucleic acid by which they are encoded, such as structure elements of the nucleic acid. In addition to cis elements for
10 replication (including in any case the binding site for an RNA/RNA polymerase) the genome of RNA viruses mostly also includes cis elements for transport (the binding site of transport proteins), cis elements for packing the nucleic acid in phage envelopes to form virus particles, and cis elements
15 for translation of messenger RNA into protein, in particular coat protein. Examples of trans elements of the genome of RNA viruses are the genes which code for coat protein, transport protein and RNA/RNA polymerase.

 An essential element of the invention is that the
20 expression cassette incorporated into the genome of the host leads to transcription of the sequence derived from RNA virus to form a messenger RNA molecule with a panhandle structure. No strict requirements are set to the elements of the expression cassette which regulate this transcription, such as
25 in particular the transcription promoter. The promoter may be, and in many cases will even preferably be, a relatively weak promoter so that the host will be virtually unburdened by this

transcription and the transcription products formed in the process. Suitable promoters are known for many organisms. Naturally the expression cassette should also comprise a suitable polyadenylation site, while the expression cassette
5 is flanked at both ends by so-called integration sites enabling integration into the genome of the intended host.

Various experiments have demonstrated that a successful expression in the host of the genetic information incorporated into the genome requires the presence of two, 12-250 base pair
10 long, inverted repeat nucleotide sequences flanking the DNA in-between. These inverted repeat nucleotide sequences may for instance consist of dG-dC base pairs or dC-dG base pairs. The fact that the presence of inverted repeat nucleotide sequences leads to both replication and expression in infected cells of
15 the host, is ascribed to the formation of RNA molecules with a stabilizing panhandle structure (see Van Emmelo et al., Virology 157, 1987, 480-487). For this purpose it is necessary that the inverted repeat nucleotide sequences have a length of at least 12 base pairs. Preferably, however, the inverted
20 repeat nucleotide sequences have a length of at least 15 base pairs. Inverted repeat nucleotide sequences of a length of more than 250 base pairs are not very practical. Preferably, they are not longer than about 50 base pairs.

A further essential feature is the absence of RNA/RNA
25 polymerase (or replicase), or the gene coding therefor, so that the amount of RNA virus-specific messenger RNA present in the cells of the host due to transcription is not further

increased. When the RNA virus derived sequence to be incorporated is derived from a satellite virus, such as STNV, this requirement is automatically satisfied because the STNV genome does not contain an RNA/RNA polymerase gene (which
5 explains why the satellite virus depends for its replication on the helper virus, which provides the required RNA/RNA polymerase).

According to the invention it is further of great importance that no viral coat protein is produced and that the
10 nucleotide sequence derived from RNA virus does not contain a gene that codes for viral coat protein.

Finally it is also essential according to the invention that the messenger RNA derived from RNA virus contains at least those elements which in the presence of RNA/RNA
15 polymerase enable replication. In other words, at least cis elements for replication should be present. Depending on the object contemplated, it may be desirable that other elements of the viral genome are present as well. Thus, in particular in the case of protection of the host against virus
20 infections, it is preferable that the genetic information incorporated into the genome of the host also comprises cis elements for transport.

On the other hand, it is not necessary according to the invention that the sequence derived from RNA virus is
25 incorporated into the genome in the sense orientation. The same holds for the orientation of other sequences located between the inverted repeat nucleotide sequences, such as a

sequence coding for a ribozyme and a sequence coding for a type-foreign protein/peptide. Owing to the fact that according to the invention both a sense orientation and an anti-sense orientation can be chosen, it is possible to accomplish both a maximum protection under normal, infection-free conditions and an adequate reaction in the case of infection.

Accordingly, the invention primarily provides a recombinant DNA, comprising two, 12-250 base pair long, inverted repeat nucleotide sequences with therebetween at least one nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, said RNA virus derived sequence comprising at least cis elements for replication, but no gene that codes for RNA/RNA polymerase, and no gene that codes for viral coat protein.

To be employed, the recombinant DNA according to the invention must be incorporated in an expression cassette for a host to be transformed enabling transcription to take place in the host to form an RNA molecule with a panhandle structure.

Preferably, in a recombinant DNA according to the invention the RNA virus derived sequence contains both cis elements for replication and cis elements for transport.

The invention further includes such a recombinant DNA in which the RNA virus derived sequence also comprises cis elements for packing in coat protein and such a recombinant DNA in which the RNA virus derived sequence also comprises cis elements for translation. In the case of uses where these

functions play no role, however, they are preferably absent, because they may have an adverse effect on the replication efficiency of the RNA formed by transcription. This holds in particular for the use of the invention for virus protection.

5 This is one of the reasons why no gene that codes for viral coat protein may be present. Similarly, preferably no other unnecessary trans elements are present, such as a gene that codes for viral transport protein.

Accordingly, in the most preferable embodiments of the
10 invention the RNA virus derived sequence, present in the recombinant DNA, corresponds to a stripped viral replicon which without outside help (such as infection with a helper virus) is incapable of multiplying and actively moving to other cells of the host, and which does not code for any viral
15 protein. Such a substantially stripped viral replicon, however, exhibits in the presence of RNA/RNA polymerase a much more efficient replication than the complete viral genome. This is what makes the recombinant DNA according to the invention useful for the protection of the host against virus
20 infections. Thus, when an infection of the genetically modified host by an RNA virus with the relevant RNA/RNA polymerase occurs, the number of RNA virus derived RNA molecules which are already present in the cells of the host will very rapidly increase and accordingly will increasingly
25 appropriate the RNA/RNA polymerase at the expense of the multiplication of the infecting virus. Therefore the substantially stripped replicon has an important advantage

over the complete viral genome in the battle for the available RNA/RNA polymerase. This advantage can be further increased when the RNA virus derived sequence incorporated in the genome of the host also contains cis elements for transport, in virtue of the fact that then also transport protein of the infecting virus, responsible for expansion of the infection to other cells of the host, is appropriated by the rapidly increasing RNA virus derived RNA molecules which are derived from the DNA incorporated.

10 A preferred embodiment of the invention, which concerns such use for protection of the host against viral infections, consists of recombinant DNA, in which between the two inverted repeat nucleotide sequences, in addition to the RNA virus derived sequence, also at least one nucleotide sequence is located which codes for a ribozyme which can cut viral RNA or mRNA.

When in the case of a viral infection by a virus which supplies the RNA/RNA polymerase in question, to be referred to hereinafter as a compatible virus, the RNA virus derived RNA present in the cells of the host under attack is rapidly multiplied, in this preferred embodiment the present amount of ribozyme with specificity for the infecting virus is also increased at the same time. This ribozyme takes care of an active destruction of the RNA of the infecting virus, or messenger RNA derived therefrom.

For further information on the structure and action of ribozymes, reference is made to Nature 334, page 197, 1988.

For further information on the formation and properties of substantially stripped replicons, reference is made to Perrault, Current Topics in Microbiol. and Immunol. 93, 1981, 152-209; Lazzarini et al., Cell 26, 1981, 145-154; Nayak, 5 Ann.Rev. Microbiol. 34, 1980, 619-644; Barrett and Dimmock, Current Topics in Microbiol. and Immunol. 128, 1986, 55-84; "RNA genetics", 1989, E. Domingo et al., Eds., CRC Press, Boca Raton, Florida; Strauss and Strauss, Current Topics in Microbiol. and Immunol. 105, 1983, 1-89. There they are 10 designated as "defective interfering viruses/particles/RNAs". A highly defective replicon of an RNA virus can for instance be obtained by infecting a suitable host with bare viral RNA, isolating the replicated viral RNA and using it for a subsequent infection. Thus the need for coat protein is 15 artificially eliminated, as a result of which deletion mutants are formed that do not contain a coat protein gene anymore. By further repeats of this procedure each time in the presence of a small amount of added RNA of this coat protein deletion mutant, also RNA can be isolated which does not contain an 20 RNA/RNA polymerase gene anymore (due to the fact that the added RNA provides the necessary RNA/RNA polymerase, also RNA/RNA polymerase-deletion mutants undergo replication). Other methods of preparation, however, are also possible, such as a method in which cells or tissues with a very high dose of 25 virus particles (100 or more per cell) are infected and RNA deletion mutants are isolated; or a method in which cells or tissues are infected with virus stocks, which contain

184-302D

11

satellite viruses or virusoids, an RNA deletion mutants are isolated; or a method in which a subgenome is isolated from an RNA virus with a bipartite or tripartite (or, more generally, polypartite) genome; or a method in which cells or tissues
5 which produce a suitable RNA/RNA polymerase are infected with viral RNAs, optionally already partly deleted, and further deleted RNAs are isolated; or a method in which targeted genetic manipulations in the cDNA of a viral, satellite or virusoid RNA are performed.

10 One aspect of the invention provides an eukaryotic organism comprising recombinant DNA incorporated into its genome, said recombinant DNA comprising an expression cassette comprising two inverted repeat nucleotide sequences of 12 to 1250 basepairs with there between at least one nucleotide
15 sequence which is a stripped viral replicon of an RNA virus whose replication is dependent upon a viral RNA/RNA polymerase, wherein said stripped viral replicon comprises cis elements for replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is
20 unable to replicate independently, and wherein the replicon, in the presence of viral RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA.

 Another aspect of the invention provides a method of protecting eukaryotic cells or plants against an RNA-based
25 virus which for its replication is dependent upon a viral RNA/RNA polymerase, comprising incorporating into the genome of the eukaryotic cells or plants an expression cassette functional in eukaryotes, said cassette comprising: a stripped viral replicon of said RNA-based virus, wherein the stripped
30 viral replicon comprises cis elements for replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is unable to replicate independently wherein the replicon, in the presence of viral

20184-302

11a

RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA, and wherein the replicon is between two inverted repeat nucleotide sequences of 12-250 bp; wherein greater efficiency of replication of the replicon in the presence of RNA/RNA polymerase compared to wild-type viral RNA protects the eukaryotic cells or plants against the RNA-based virus.

Another aspect of the invention provides use of an expression cassette functional in animals comprising a stripped viral replicon of an RNA-based animal virus whose replication is dependent upon a viral RNA/RNA polymerase, for protecting an animal against said RNA-based virus, said replicon comprising *cis* elements for replication but being devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, said replicon being unable to replicate independently, said replicon, in the presence of viral RNA/RNA polymerase, being able to replicate more efficiently than wild-type viral RNA, and said replicon being present between two inverted repeat nucleotide sequences of 12-250 bp.

Another aspect of the invention provides a system for protecting an animal against an RNA-based animal virus whose replication is dependent upon a viral RNA/RNA polymerase, comprising: (a) an expression cassette functional in animals comprising a stripped viral replicon of the RNA-based virus, said replicon comprising *cis* elements for replication but being devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, said replicon being unable to replicate independently, said replicon, in the presence of viral RNA/RNA polymerase, being able to replicate more efficiently than wild-type viral RNA, and said replicon being present between two inverted repeat nucleotide sequences of 12-250 bp; (b) means for incorporating the expression cassette into the genome of the animal; and wherein greater efficiency of replication of

20184-302

11b

the replicon in the presence of RNA/RNA polymerase compared to wild-type viral RNA protects the animal against the RNA-based animal virus.

Another aspect of the invention provides a method of inducing or amplifying production in eukaryotic cells or plants, of at least one non-viral product selected from the group consisting of protein, polypeptide, peptide and RNA, comprising: (a) incorporating into the genome of the eukaryotic cells or plants an expression cassette functional in eukaryotes, said cassette comprising, between two inverted repeat nucleotide sequences of 12-250 bp, (i) a stripped viral replicon of a RNA-based virus whose replication is dependent upon a viral RNA/RNA polymerase, wherein said replicon comprises *cis* elements for replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is unable to replicate independently, and wherein the replicon, in the presence of viral RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA; and (ii) a nucleotide sequence encoding at least one non-viral product, the non-viral nucleotide sequence being in sense or anti-sense orientation; (b) growing the eukaryotic cells or plants produced in step (a) under conditions where RNA/RNA polymerase is present in the eukaryotic cells or plants; wherein the RNA/RNA polymerase induces or amplifies expression of the non-viral product in the eukaryotic cells or plants; and (c) isolating the non-viral product produced.

Another aspect of the invention provides use of an expression cassette functional in animals, in conjunction with an RNA/RNA polymerase, for inducing or amplifying expression in an animal or animal cell of a non-viral product selected from the group consisting of protein, polypeptide, peptide and RNA, said cassette comprising, between two inverted repeat

20184-302

11c

nucleotide sequences of 12-250 bp, (i) a stripped viral replicon of a RNA-based virus whose replication is dependent upon a viral RNA/RNA polymerase, wherein said replicon comprises *cis* elements for replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is unable to replicate independently, and wherein the replicon, in the presence of viral RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA; and (ii) a nucleotide sequence encoding at least one non-viral product, the non-viral nucleotide sequence being in sense or anti-sense orientation.

Another aspect of the invention provides a system for inducing or amplifying production in eukaryotic cells or plants of a non-viral product selected from the group consisting of protein, polypeptide, peptide and RNA, comprising: (a) an expression cassette functional in eukaryotes, comprising, between two inverted repeat nucleotide sequences of 12-250 bp, (i) a stripped viral replicon of a RNA-based virus whose replication is dependent upon a viral RNA/RNA polymerase, wherein said replicon comprises *cis* elements for replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is unable to replicate independently, and wherein the replicon, in the presence of viral RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA; and (ii) a nucleotide sequence encoding at least one non-viral product, the non-viral nucleotide sequence being in sense or anti-sense orientation; and (b) means for incorporating the expression cassette into the genome of the eukaryotic cells or plants; and (c) means for providing the eukaryotic cells or plants, into whose genome the expression cassette is incorporated, with RNA/RNA polymerase; wherein the RNA/RNA polymerase induces or amplifies expression of the non-viral product in the eukaryotic cells or plants.

20184-302

11d

The invention has important advantages over the previously proposed methods of protecting eukaryotic host organisms against virus infections, such as the formation of antisense viral RNA to block viral messenger RNA (see for instance Cuozzo et al., Biotechnology 6, 1988, 549-557), the formation of viral coat protein (see the cited article of Cuozzo et al., and Hoekema et al., Biotechnology 7, 1989, 273-279), or the formation of viral satellite RNA (see EP-A-0242016). The invention is subject to fewer limitations than the known virus protection methods. For instance, the invention is not limited to viruses of which a satellite virus is known, and is not limited to the incorporation of "sense" or "antisense" DNA (the RNA virus derived RNA formed by transcription may appropriate both in the sense and in the antisense orientation the trans functions for replication and possibly transport of the infecting virus). The invention also offers more possibilities for achieving greater effectiveness, such as the possibility of incorporating one or more ribozyme

sequences. The invention combines a smallest possible burdening of the host and a highest possible protection for the host under infection-free conditions with an extremely fast and effective reaction, as soon as an infection by a compatible virus occurs, which reaction gives the host the required protection against the virus, exclusively at locations where that is necessary. The protection the invention offers is of a permanent nature because it is integrated into the genome of the host and, due to its being a minor burden on the host, exerts virtually no selection pressure. A simultaneous protection of the host against several different viruses is a real possibility of the invention owing to the fact that it constitutes virtually no extra burden on the host, i.e. the host need hardly use any energy (the transcription of the DNA inserted requires only very little energy, while in the preferred embodiment no translation into protein is involved). According to the invention the inverted repeat sequences may contain several different RNA virus-specific sequences (i.e. defective replicons of different RNA viruses), and/or several different ribozyme sequences (a ribozyme sequence which is specific for a first virus, a ribozyme sequence which is specific for a second virus, etc.). Naturally, however, these several different RNA virus-specific sequences and several different ribozyme sequences may also be incorporated each within two inverted repeat sequences of their own. In the known methods of virus protection such simultaneous protection against

different viruses is not very well possible because the large amounts of entities providing protection, which must be produced continuously throughout the host to ensure effective protection constitute, due to their being a burden, a
5 selective drawback of the host as compared with others of its type and/or are detrimental to its growth and development.

The above-mentioned properties of the recombinant DNA according to the invention are also responsible for its usability as regards methods of producing foreign
10 proteins/peptides and RNAs using genetically modified, prokaryotic and particularly eukaryotic organisms. This concerns recombinant DNA according to the invention in which between the two inverted repeat nucleotide sequences, in addition to the RNA virus derived sequence, a non-viral
15 nucleotide sequence is located which codes for one or more RNAs, and more particularly recombinant DNA according to the invention in which the RNA virus derived sequence comprises at least cis elements for replication and cis elements for translation and between the two inverted repeat nucleotide
20 sequences, in addition to the RNA virus derived sequence, a non-viral nucleotide sequence is located which codes for one or more proteins/peptides.

In virtue of the above-mentioned properties of the recombinant DNA according to the invention the host produces
25 at most a negligible amount of the foreign RNAs or proteins/peptides in the absence of RNA/RNA polymerase, while in the presence of RNA/RNA polymerase a sharp increase of the

messenger RNA occurs, which results in a considerable production of the desired RNAs or proteins/peptides. This very strong production can moreover be induced at any desired moment, or be limited to specific tissues of the host by
5 regulating the presence of the RNA/RNA polymerase accordingly.

An inducible production may for instance be accomplished by infecting the genetically modified host (or host cells in a cell culture, for instance) with a compatible virus at the desired moment. Normally, however, this method will not be
10 preferable because it is laborious, the efficiency of the infection may be very variable, the infection mostly takes place at the expense of the host (a possible consequence being a breakdown of the desired product) and/or the use of the virus involves risks. Another, more attractive option, which
15 may also be chosen to accomplish a tissue-specific production consists in a double transformation of the host, wherein the genome of the host is provided not only with the above described recombinant DNA according to the invention but also with recombinant DNA comprising genetic information for a
20 viral RNA/RNA polymerase or an RNA/RNA polymerase construct in an inducible or tissue-specific expression cassette. For this purpose use can be made of known per se inducible (for instance by heat, UV irradiation, or chemicals such as salicylic acid) or tissue-specific (for instance patatin for
25 expression in the tubers of potatoes) promoters.

The phrase "genetic information for an RNA/RNA polymerase construct" refers to gene constructions which may or may not

effect coding for another RNA/RNA polymerase. A possible gene construction consists, for instance, of a mutation through which an internal facultative stop codon which may occur in genes coding for RNA/RNA polymerases, is replaced by a
5 sequence not recognizable as a stop codon anymore. Another possibility is a gene construction which consists of a fusion of different RNA/RNA polymerase genes, naturally with the same above-mentioned option of using a mutation of an internal stop codon.

10 The invention is further embodied in eukaryotic or prokaryotic cells or organisms which through genetic engineering are provided with recombinant DNA according to the invention and optionally, through genetic engineering, are also provided with recombinant DNA comprising genetic
15 information for a viral RNA/RNA polymerase or an RNA/RNA polymerase construct in an inducible or tissue-specific expression cassette.

The invention is further embodied in a method of protecting eukaryotic organisms, in particular plants, yeasts
20 and fungi, against an RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase by performing a genetic manipulation of the organism to be protected into the genome of this organism recombinant DNA which in an active expression cassette comprises two, 12-250 base pair long,
25 inverted repeat nucleotide sequences with therebetween a nucleotide sequence derived from RNA virus, this RNA virus derived sequence comprising at least cis elements for

replication, but no gene that codes for viral RNA/RNA polymerase and no gene that codes for viral coat protein.

Preferably, in such a method the RNA virus derived sequence comprises both cis elements for replication and cis
5 elements for transport. The RNA virus derived sequence may further also comprise cis elements for packing in coat protein and cis elements for translation.

Most preferably, in such a method, between the two inverted repeat nucleotide sequences, in addition to the RNA
10 virus derived sequence, at least one nucleotide sequence is located which codes for a ribozyme which can cut viral RNA or mRNA.

The invention is further embodied in a method of inducibly producing one or more proteins/peptides by culturing
15 a prokaryotic or eukaryotic organism, or cells of a prokaryotic or eukaryotic organism, which organism, through genetic engineering, is provided with recombinant DNA which in an active expression cassette comprises two, 12-250 base pair long, inverted repeat nucleotide sequences with therebetween
20 one nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, this RNA virus derived sequence comprising at least cis elements for replication and cis elements for translation, but no gene that codes for viral RNA/RNA
25 polymerase, and no gene that codes for viral coat protein, and there being located between the two inverted repeat nucleotide sequences, in addition to the RNA virus derived sequence, a

non-viral nucleotide sequence which codes for one or more proteins/peptides, and infecting the organisms or cells thereof with the virus.

Preferable, however, is a method of producing in an
5 inducible manner or in a tissue-specific manner one or more proteins/peptides by culturing a eukaryotic organism, whose genome, through genetic manipulation, incorporates recombinant DNA which in an active expression cassette comprises two, 12-250 base pair long, inverted repeat nucleotide sequences with
10 therebetween a nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, with this RNA virus derived sequence comprising at least cis elements for replication and cis elements for translation, but no gene that codes for viral
15 RNA/RNA polymerase, and no gene that codes for viral coat protein, and there being located between the two inverted repeat nucleotide sequences, in addition to the RNA virus derived sequence, a non-viral nucleotide sequence which codes for one or more proteins/peptides, as well as recombinant DNA
20 which comprises genetic information for a viral RNA/RNA polymerase or an RNA/RNA polymerase construct in an inducible or tissue-specific expression cassette.

The invention is further embodied in a method of inducibly producing one or more RNAs, such as ribozymes,
25 antisense RNAs and double-stranded RNAs, by culturing a prokaryotic or eukaryotic organism, or cells of a prokaryotic or eukaryotic organism, which organism through genetic

engineering is provided with recombinant DNA which in an active expression cassette comprises two, 12-250 base pair long, inverted repeat nucleotide sequences with therebetween a nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, this RNA virus derived sequence comprising at least cis elements for replication but no gene which codes for viral RNA/RNA polymerase, and no gene which codes for viral coat protein, and there being located between the two inverted repeat nucleotide sequences, in addition to the RNA virus derived sequence, a non-viral nucleotide sequence which codes for one or more RNAs, and infecting said organism or cells thereof with the virus.

More preferable, however, is a method of producing in an inducible manner or in a tissue-specific manner one or more RNAs, such as ribozymes, antisense RNAs and double-stranded RNAs, by culturing a eukaryotic organism, whose genome, through genetic engineering, incorporates recombinant DNA which in an active expression cassette comprises two, 12-250 base pair long, inverted repeat nucleotide sequences with therebetween a nucleotide sequence which is derived from RNA virus which for its replication is dependent upon a viral RNA/RNA polymerase, this RNA virus derived sequence comprising at least cis elements for replication, but no gene which codes for viral RNA/RNA polymerase, and no gene which codes for viral coat protein, and there being located between the two inverted repeat nucleotide sequences, in addition to the RNA

virus derived sequence, a non-viral nucleotide sequence which codes for one or more RNAs, as well as recombinant DNA which comprises genetic information for a viral RNA/RNA polymerase or an RNA/RNA polymerase construct in an inducible or
5 tissue-specific expression cassette.

Thus the invention can be used broadly for the production in eukaryotes or cells of eukaryotes of one or more products such as proteins, oligo and polynucleotides, oligo and polypeptides, enzymes, antibodies, antigenic substances,
10 antiviral compounds, anticancer substances, hormones, vitamins, medicines and pharmaceuticals, primary and secondary metabolites.

A further aspect of the invention is recombinant DNA comprising a nucleotide sequence which codes for a viral
15 RNA/RNA polymerase or an RNA/RNA polymerase construct.

More particularly, according to a preferred embodiment, the invention provides such a recombinant DNA comprising the part of the nucleotide sequence shown in Fig. 4 that codes for a viral RNA/RNA polymerase, or constructs derived therefrom,
20 such as a substitution mutant which has sequence TAT instead of the sequence TAG at the positions 656-658 according to the numbering used in Fig. 4, and substitution mutants in which a part of the sequence shown in Fig. 4 is replaced by a corresponding part of another gene which codes for a viral
25 RNA/RNA polymerase.

Such a recombinant DNA in which the nucleotide sequence which codes for a viral RNA/RNA polymerase or an RNA/RNA

polymerase construct is located in an inducible or tissue-specific expression cassette constitutes yet another preferred embodiment of the invention.

The invention will now be explained in and by the following Examples.

Example I illustrates how tobacco plants can be protected against infection by TNV through transformation with a replicon derived from STNV. The replicon derived from STNV is a highly defective replicon which does not code for a protein and under infection-free conditions is produced only in a very limited amount, i.e. only a weak transcription of the DNA incorporated into the genome occurs. However, when the plant is infected with TNV, which codes for an RNA/RNA polymerase which is capable of multiplying the STNV derived replicon, massive reproduction of the STNV derived replicon occurs, irrespective of whether the STNV information is incorporated into the genome in sense or anti-sense orientation. As a result, the plant is protected against the infecting virus. This protection is much more effective when the DNA incorporated into the genome also comprises the information for a ribozyme which is directed against mRNA for TNV coat protein.

Example II illustrates how the invention can be used to accomplish an inducible production of a type-foreign protein. As a model for this purpose the beta-glucuronidase gene of E. coli was selected, which was fused with the initiation codon of the STNV coat protein. In the example given the expression

was induced by infecting the transformed tobacco plants with TNV.

Example III describes the isolation of the replicase gene of TNV and the construction of a plasmid pSPTNV rep-1, which
5 contains this replicase gene.

Example IV describes expression experiments in which amplification of the messenger RNA by contact with the TNV replicase was effected. The viral replicon contained as foreign DNA a fusion of a part of the gene which codes for the
10 viral coat protein of STNV, and the chloramphenicol acetyl transferase (CAT) gene of E.coli.

Example I

(a) Recombinant plasmids

Starting from the plasmid pSTNV-413 described by Van
15 Emmelo et al. in Virology 157, 480-487 (1987), a number of new insertion mutants were constructed. The starting plasmid was linearized with RsaI, which has 9 cutting sites in the STNV genome, by incubating at 28°C with 0.1 µg enzyme per µg DNA for 15 min. Ligation with the same 14-mer linker as described
20 by Van Emmelo et al: 5' TCCATGGGAATTCT 3' led, among other things, to the insertion mutant pBR STNV N162, which contains the linker with the NcoI site at the 5' end at position 162 of the STNV genome.

From this insertion mutant and the insertion mutants pBR
25 STNV N198, N322 and E531, already described by Van Emmelo et al., all of which contain the reading frame interfering insertion of the above-mentioned 14-mer linker in the gene

coding for coat protein, mutants with a recovered reading frame were constructed. To that end the 14-mer insertion was converted into an 18-mer insertion by cutting with NcoI, filling in the single-stranded ends with Klenow enzyme in the presence of the 4 dNTP's, and religation of the plasmid. Thus the 14-mer insertion is converted into the following 18-mer insertion: 5' TCCATGCATGGGAATTCT 3', in which the NcoI site (CCATGG) is replaced by an NsiI site (ATGCAT). Thus the insertion mutants pBR STNV N164, N200, N324 and E533 were obtained which theoretically code for a coat protein which is increased by 6 amino acids.

Further the double insertion mutant pBR STNV N164N843 was constructed by isolating from pBR STNV N843 the EcoRV fragment of bases 198 to 962 in the STNV genome and substituting it for the corresponding EcoRV fragment of pBR STNV N164.

From the 18-mer insertion mutants pBR STNV N164 and N200 the mutants pBR STNV S164 and S200 were constructed by inserting an additional linker of 18 bases as NsiI fragment in the NsiI site of the 18-mer insertion:

in S164: 5' TCCATGCAATCGAGGGTAGGCATGCATGGGAATTCT 3'

in S200: 5' TCCATGCATGCCTACCCTCGATTGCATGGGAATTCT 3'

These mutants have an insertion of 36 bases and code theoretically for a shorter coat protein as a result of a reading frame mutation in the case of S164 and for a coat protein with 12 extra amino acids in the case of S200.

By inserting the same 18-mer in the unique NsiI site at the end of the coat protein gene (base 613), the mutant pBR STNV S613 was obtained.

At the location of the RsaI site at position 162 a linker of 30 bases was inserted in a similar way as described hereinabove for the 14-mer linker. In the reading frame that is formed thus this linker codes for the neuropeptide bradykinin: 5' GCGGCCGCCCGGGTTTAGTCCTTTTAGGTT 3'

ArgProProGlyPheSerProPheArg

10 This insertion mutant was designated pBR STNV Brad.

From pBR STNV N164 the deletion mutant pBR STNV D1 was made by a deletion of the NsiI fragment of 167 to 631. The genome of this mutant has a length of 793 bases as compared with 1239 for wild type STNV.

15 For expression studies in E.coli the various STNV constructions were transcloned as PstI fragment into pPLC 2820. The restriction enzyme PstI cuts precisely at the end of the poly GC regions which flank the 5' and 3' ends of STNV. These constructions in pPLC2820 are designated as pPLC, 20 followed by the name of the STNV mutant, further followed by .1 when the 5' end of the STNV genome adjoins the pL promoter of pPLC 2820, or by .2 when the 3' end adjoins the promoter. Thus the plasmid pPLC STNV N164.1 is the plasmid with the STNV mutant N164 cloned into pPLC 2820, with the STNV mutant being 25 inserted into the plasmid such that upon transcription by pL the equivalent of the positive-strand genomic RNA of STNV is made as mRNA.

The STNV mutants cloned as PstI fragment into pPLC 2820 are preceded at the 5' end by a unique BamHI site and at the 3' end followed by a SalI site. Since neither STNV, nor the mutants derived therefrom, contain a BamHI or a SalI site, the different constructions can be transcloned as BamHI/SalI fragment into pPCV 520 (Fig. 1) behind the plant promoter pTR1'. The plasmids thus obtained are designated as pPCV, followed by the name of the STNV mutant, further followed by .1 when the 5' end, and by .2 when the 3' end of STNV is turned towards the promoter pTR1'. The pPCV derivatives were used for transformations of tobacco plants as part of a binary vector system, as described by Hoekema et al. in EMBO Journal 3, 2485 (1984) and by De Framond et al. in Mol. Gen. Genet. 202, 125 (1986). The plasmids derived from pPCV 520 functioned as the T-plasmid (i.e. the plasmid that contains the T-DNA which is integrated into the genome of the plant through the transformation). The most important properties of pPCV 520 (see Fig. 1) are:

- ColE1 replication in E.coli, which makes it suitable for use in all current E.coli strains and which makes it a multicopy plasmid (efficient DNA preparation and clone analysis)
- bacterial resistancies against the antibiotics ampicillin and chloramphenicol
- the P-type replication origin of pRK2 is active in E.coli and in Agrobacterium, provided that the RK2-gene trfa is

expressed (in Agrobacterium, moreover, selection is necessary for stability)

--- the P-type transfer origin of pRK2 permits an efficient transfer of the plasmid from strains in which the pRK2-genes
5 tra1, tra2 and tra3 are expressed (for E.coli this is the case for the strain SM10 and for Agrobacterium for strains which contain the plasmid pMP90RK)

--- the border sequences of the T-DNA flank all elements of the vector which are important in plants

10 --- the NPT-II kanamycin resistance gene under the control of the plant promoter pTR2' permits selection of transformed plants

--- the octopine synthase gene in the T-DNA makes it possible to confirm the transformation of kanamycin-resistant plants
15 using octopine synthesis

--- the constitutive plant promoters pTR1' and pnos, are followed by the unique restriction sites for SalI and BamHI and for BglII and BclI, respectively and by the polyadenylation sequences of the T-DNA genes 7 and 4,
20 respectively

--- back cloning of T-DNA sequences from transformed plants is simplified by the pBR322 oriV and ampicillin resistance in the T-DNA.

The T-plasmid pPCV 520, therefore, can replicate both in
25 E.coli and in Agrobacterium. The cloning steps for the introduction of the STNV mutants into this vector can be performed in E.coli. Then, by conjugation, the vector is

transferred from E.coli SM10 into the Agrobacterium strain
GV3100 (pMP90RK), which is used for the plant transformations.
The plasmid pMP90RK (Fig. 2; see Koncz and Schell, Mol. Gen.
Genet. 204, 1986, 383-396) functions as virulence plasmid. It
5 is derived from the nopaline Ti-plasmid pGV3100 by (1) a
combined deletion and insertion mutagenesis, leading to a
complete deletion of the T-DNA and to the incorporation of a
gentamycin resistance gene (the construction thus obtained is
designated as pMP90), and (2) the insertion of a fragment of
10 the P-type plasmid pRK2013 with thereon the genes for
kanamycin resistance, the genes acting in trans for transfer
and replication of P-type plasmids and the P-type origin of
transfer (Figurski and Helinski, PNAS USA 76, 1648 (1979)).
The most important properties of this plasmid pMP90RK (see
15 Fig. 2) are:
--- a stable replication only in Agrobacterium
--- compatibility with P-type plasmids
--- it codes for all Ti-virulence functions which are
necessary and sufficient for the integration of the T-DNA into
20 the genome of plants during transformation
--- it contains gentamycin and kanamycin resistance genes as
genetic markers which are easy to select
--- it complements transfer and replication of P-type mutants
by the expression of the pRK2-genes tra1, tra2, and tra3 and
25 trfa
--- it conjugates very efficiently owing to the P-type oriT.

(b) Techniques

The transformation of plants (tobacco SR1) was according to the leaf segment method, described by De Block et al. in EMBO Journal 3, 1681 (1984) and by Horsch et al. in Science 5 223, 496 (1984). The procedure was as follows:

--- by incisions with a scalpel tobacco leave are wounded and the freshly wounded leave are incubated for 48 h with a 1/50 dilution of a fresh matured culture of the transforming Agrobacteria. This is conducted in liquid MS-plant medium 10 (Murashige and Skoog medium, Gibco)

--- then the leaf fragments are washed twice for 12 h in liquid MS-medium to which claforan (500 µg/ml) is added (claforan is an antibiotic which acts only against bacteria)

--- then the leaf fragments are incubated on solid MS-medium 15 (+ 0.8% agar) to which cytokinin (6-benzyl-aminopurine or BAP, 1 mg/l) and auxin (α-naphthalene acetic acid or NAA, 0.1 mg/l) are added in a ratio which promotes shoot formation at the location of the wounds, kanamycin (100 µg/ml) for the selection of transformed shoots and claforan for the further 20 selection against the transforming Agrobacteria

--- after 4 to 10 weeks transformed shoots grow from this medium which, when they are sufficiently developed, can be transferred to hormone-free medium where they can develop to normal plants with roots.

25 Callus induction was performed on sterile leaf and stem fragments. For that purpose the fragments were incubated on solid MS-medium (+ 0.8 % agar) to which the cytokinin BAP

(0.5 mg/l) and the auxin NAA (1.0 mg/l) were added in a concentration and ratio which promote callus growth.

For octopine tests small particles of plant tissue (about 50 mg) were incubated overnight in a solution of 100 mM L-arginine and 50 mM pyruvate dissolved in liquid MS-medium. After incubation the liquid was removed and the tissue was washed in MS-medium. Then the plant fragment was crushed and centrifuged, and 3 to 5 μ l of the extract was separated by paper electrophoresis. This was conducted on Wattman 3MM paper, in a solution of 15 % HAc, 5 % HCOOH and 80 % H₂O. Octopine spots on the paper were made visible under UV lighting after a staining reaction with a fresh mixture of 1 part 0.02 % phenanthrenequinone in 95 % EtOH and 1 part 10 % NaOH in 60 % EtOH.

Plasmid infections of cowpea were performed to investigate the infectivity of the STNV mutants. To that end cowpea was inoculated with a mixture of TNV helper virus and chimeric plasmids according to the method described by Van Emmelo.

Northern analyses were performed essentially in the manner described by Van Emmelo. For the analyses of the transformed plants single- and double-stranded RNA were not separated by LiCl precipitation for gel electrophoresis. Besides denaturing glyoxal agarose gels, denaturing formamide agarose gels were used, as described by Lechrach et al in Biochemistry 16, 4743 (1977) and by Maniatis et al in Molecular Cloning: a laboratory Manual, Cold Spring Harbor

Laboratory, Cold Spring Harbor, N.Y., USA. In this method the RNA samples are first denatured at 55°C for 15 min in a mixture of 50% formamide, 17.5% formaldehyde, 20 mM morpholino propane sulfonic acid (MOPS) (pH 7.0), 5 mM NaAc and 1 mM EDTA (pH 8.0). Gel electrophoresis was conducted in a 1.2% agarose gel in 20 mM MOPS (pH 7.0), 5 mM NaAc and 1 mM EDTA (pH 8.0) as a transporting buffer. The further treatment of the samples for blotting and hybridization was carried out in the manner described by Van Emmelo.

Western analyses for detecting TNV and STNV coat protein were performed as follows. Infected leaves were picked after two days and using a potter were ground at 4°C in 1 volume part PBS (PBS= 0.13 M NaCl, 10 mM sodium phosphate buffer pH 7.4) which contained 100 µg/ml PMSF. Cell residues were removed by centrifugation and the proteins were precipitated with trichloroacetic acid (TCA, final concentration 10%). The precipitate was washed in ethanol, ethanol ether (1:1) and ether, dried and resuspended in 1x Laemmli-buffer (Nature 227, 680-685, 1970). Fractions thereof were separated on a 10% PAG by electrophoresis, and via electrotransfer for 90 min at 0.5 A transferred to Pall-biodyne filter in a transfer buffer (1 mM sodium acetate, 5 mM MOPS pH 7.5 in 20% ethanol). The filter was blocked overnight in PBS which contained 10% low-fat milk powder and was incubated with anti-STNV coat protein serum of rabbits, diluted 1:200 in PBS, 2% low-fat milk powder at room temperature for 1 h. The filter was washed four times with PBS, 0.1% Triton-X-100* and incubated with anti-

rabbit alkaline phosphatase labeled antibodies from goats, diluted 1:250 in PBS, 2 % low-fat milk powder. After washing out non-bound antibodies, a staining reaction was carried out using Nitro Blue Tetrazolium (NBT) and 5-bromo-4-chloro-3-indolyl phosphate (BCIP), under the conditions as indicated by the producer (Promega Biotec).

(c) Experiments and results

10 (c1) expression studies in E. coli NF1

A number of pPLC STNV plasmids with the modified STNV genome in the sense orientation (the .1 constructions) and the plasmids pPLC STNV N164.2 and Brad.2 (with the antisense orientation) were transformed in E. coli NF1. This strain has a ts (temperature-sensitive) mutation of the c1 repressor gene of pL, by which the pL promoter is repressed when the bacteria are cultured at 28°C. At a temperature of 42°C the repressor is unstable and the promoter yields pL transcription.

Using Western blotting the synthesis of coat protein was investigated, i.e. the presence or absence of coat protein and the difference in size of the proteins encoded by the different STNV mutants. The results are summarized in Table 1. It reveals that

--- the STNV mutants with a reading frame mutation (+14 or +36 bases) yield a shortened coat protein whose length is greater according as the insertion is located further towards the 3' end

--- the insertion mutants with a recovered reading frame (+18, +30 or +36 bases) yield a larger coat protein than the wild type, while the mutual mobility (+18 > +30 > +36) corresponds to the higher molecular weight, which is expected on the basis of the number of added amino acids (+6, +10 and +12)

--- only the plasmids in which the modified STNV is inserted in the .1 orientation relative to the promoter, yield coat protein.

(c2) plasmid infections

Unless indicated otherwise, for the plasmid infections pBR STNV plasmids were used. The plasmid pSTNV 413 was used as a positive control in the infections to allow correlation of the various experiments. As a negative control pBR STNV N843 was used, which had been previously established as replication-deficient. When the signals after the primary infection were very weak or absent, secondary infections with RNA extracts were performed on cowpea to obtain clearer RNA and protein signals.

In the analysis of the inoculated leaf material the synthesis of coat protein was assayed by Western blotting and that of RNA by Northern blotting. For the RNA single- and double-stranded RNA were separated and their presence was determined separately. The results are also summarized in Table 1. It reveals that

--- dsRNA (double-stranded RNA) is present in all mutants with an insertion in the coat protein gene and is absent only in the mutants with an insertion at position 843

--- coat protein and normal amounts of ssRNA (single-stranded RNA) are only found in the mutants N164, N200 and Brad

--- in the insertion mutants in which no coat protein is found, the plants also contain much less ssRNA

5 --- the coat proteins which are produced for the mutants N164, N200 and Brad in the plants and in E. coli NF1 are of equal length.

By hybridization with a strand-specific probe (with an SP6 RNA transcript of STNV or with the suitable single-
10 stranded linker for the different mutants) it was established that the ssRNA isolated in the different infections is positive-strand and accordingly corresponds to the genomic RNA (also when no coat protein, no particles and only small amounts of ssRNA are present).

15 Besides the various pBR STNV plasmids, for the mutants N164.1, N164.2, Brad.1 and Brad.2 also the pPLC and pPCV plasmids were used for plasmid infections, with the same results.

20 (c3) expression of STNV mutants in transformed plants

Transformations of tobacco SR1 were performed with the following T-plasmids:

pPCV 520

pPCV STNV N164.1 and .2

25 pPCV STNV N164N843.1 and .2

pPCV STNV Brad.1 and .2

Transformation with pPCV 520 was performed as a check on the transformation experiment itself. Using the infective STNV mutant N164 with a linker insertion in the coat protein gene enables both the coat protein (+6 amino acids) of the mutant and the RNA (by hybridization with the specific linker) of the wild type to be distinguished in the analysis of the results. The mutant N164N843 was selected as replication-deficient genome. In this mutant the insertion N164 is used to identify the coat protein and to distinguish it from the wild type, while the two inserted linkers enable identification of the RNA. The mutant STNV Brad was used for transformation to establish whether the expression of the fusion protein between the STNV coat protein and the bradykinin is amplified by an infection of the transformants with the helper virus TNV. The transformed plants are designated by SR1, followed by the identification of the STNV mutant and further followed by a number used to identify the independent transformants with the same mutant.

The kanamycin resistant plants which after transfer to hormone-free medium with kanamycin developed roots in a normal manner, were increased for further investigation. Of these plants also callus cultures were produced to enable confirmation of the transformation of the kanamycin resistant plants by octopine tests. The point is that the octopine synthase gene in the T-DNA of pPCV 520 is under the control of a tissue-specific promoter of T-DNA gene 5, which is expressed only in stem fragments (very weakly) and in callus tissue, so

that the octopine tests are preferably performed on callus tissue. More than 90 % of the kanamycine resistant plants proved indeed to be transformed.

Transformed plants were obtained under sterile growth conditions. For TNV infection small top shoots were first allowed to grow roots in vitro and then transferred to potting soil to be further cultured under greenhouse conditions. The plants were allowed to develop further and infected with TNV when they were sufficiently large. The mechanical infection was performed with a fresh TNV-inoculum, isolated from tobacco SR1. After 72 h the infected leaves were picked, frozen in liquid nitrogen and preserved at -70°C. Besides the TNV infected leaves, non-infected leaves of the same plant were picked and analysed.

The presence of STNV RNA and coat protein in the various samples were determined by Northern and Western blotting, respectively, Table 2 summarizing the results. The Table reveals that

--- TNV infection of plants transformed with STNV under the control of a constitutive plant promoter and flanked by inverted poly-GC-regions leads to the replication and expression of the STNV genome and to the formation of STNV particles

--- the insertion mutant N164N843 characterized as replication-deficient replicates normally in transformed plants

--- the STNV construction with a fusion between the coat protein and the bradykinin, after infection with TNV, yields replication of the mRNA and amplified synthesis of the fusion protein

5 --- the detected STNV RNA according to the hybridizations with the various linkers corresponds to the mutant genome which the plants had been transformed with

--- the plants with the STNV genome in the .2 orientation, i.e. they produce the anti-sense mRNA, after infection with
10 TNV also yield STNV replication and expression, without observable differences with the .1 plants

--- the differences in the intensity of the RNA and protein signals do not correlate with the type or the orientation of the mutant STNV genomes, but are typical of the individual
15 character of independent transformants with the same genome.

Owing to the fact that for the assessments only 10 mg plant material was used, neither RNA nor coat protein could be detected in the transformed plants without TNV infection.

Repeat experiments all produced qualitatively and
20 quantitatively similar results. Significant variations were observed only for the efficiency of the TNV infection, depending on various factors such as age and the differentiation level of the infected plants, the quality of the inoculum, the temperature, the humidity, etc. When the TNV
25 infection was insufficiently strong, only weak RNA signals and no protein signals were obtained. With a proper TNV infection, on the other hand, much stronger STNV signals were obtained

than in the case of the plasmid infections. On the basis of the intensity of the signals obtained in Western analysis, the amount of coat protein could be estimated at about 100 ng in a sample prepared from 10 mg leaf material. Taking into account
5 the limitation of the synthesis of the STNV coat protein to the region of the necrotic lesions where TNV is also present, which lesions constitute only about 5 % of the total leaf, it can be calculated from this that the STNV signal constitutes about 0.02 % of the total, wet leaf weight or about 0.5 % of
10 the dry weight. This relatively large amount of protein is formed in only 3 days, after expression by TNV infection is induced.

The results with the STNV N164 and Brad mutants reveal that the 18-mer insertion and the fusion with bradykinin do
15 not influence the replicability and the expression of the STNV-RNA. This proves that the coat protein can be replaced in whole or in part without adverse effects on replicability and expression of the STNV-RNA. See also Example 2, to follow hereinafter.

20 Quite unexpectedly, it turned out that the mutant N164N843, unlike in the case of the plasmid infections, replicated normally in transformed plants (by linker hybridization it was proved that the 14-mer insertion at position N843 was indeed present in the detected and
25 replicated RNA). Apparently, the absence of ssRNA and dsRNA in plasmid infections with STNV N843 is not the result of a deficient replication. Further it appeared that replication of

the STNV-RNA is not dependent upon the presence or absence of STNV coat protein or STNV particles. Since in these mutants the spread of STNV infection from cell to cell occurs in a normal manner, only the viral RNA (ds or ss) can be responsible for this. However, the different behaviour of the mutant N843 in plasmid infections and in transformed plants can be explained by assuming that this mutation prevents the normal cell to cell spread of the STNV infection by RNA. In plasmid infections this mutant is not infective. In transformed plants, however, the STNV-mRNA occurs in all cells and spreading is not necessary to obtain normal replication. In addition to structural elements for replication, therefore, the STNV-RNA also contains structural elements which are necessary for cell to cell transport. Probably a TNV-encoded protein plays an active role here.

In the transformants where STNV is inserted in the .2 orientation relative to the plant promoter, the anti-sense or negative-strand RNA is formed as messenger RNA. Previous attempts to initiate an STNV infection with the negative-strand RNA using SP6-transcripts had failed. In transformed plants this appears to be possible after induction, and as efficiently as with the STNV .1-transformants. It is therefore possible to repress completely the expression of a gene during the entire growth phase of the plant due to the fact that only the anti-sense messenger RNA of a protein is produced, and to limit expression to the period when the helper functions for replication are present, while a strong expression can be

obtained in a short time by replication and amplification of the anti-sense to encoding sense RNA. Especially for the synthesis of toxic products it may be of great importance to repress entirely their expression during the growth period so
5 as to prevent a negative influence on the development of the plants.

(c4) In vivo deletions

In plasmid infections with STNV, wild type and mutants,
10 often secondary infections were performed to obtain clearer signals for RNA and protein in the analysis of the mutants. These infections were done with total RNA (ds and ss) from these plants, while both the STNV and the TNV infection were transferred. By thus continuing the same infection three to
15 four times consecutively, deletion mutants of the STNV could be isolated. Northern blotting made the RNA of these mutants visible as clear, faster migrating discrete bands. In different infection lines different mutants were formed independently of each other, the length of the replicating RNA
20 varying from about 600 to 900 bases. It appeared that when in an infection line such deletion mutants of STNV formed, the original wild type genome disappeared after 1 or 2 further infections. Also, in existing and fairly large deletion
mutants, additional deletions could lead to the formation of
25 smaller derivatives. Upon continuation of such an infection line, the larger mutant disappeared. Generally it was established that sufficient repetition of these infections led

to the formation of STNV deletion mutants with a genome of about 600 bases. Western blotting demonstrated that the STNV deletion mutants no longer code for the coat protein. By hybridizations with different fragments of the STNV genome it was determined that the deletion is located at the 5' end of the genome.

Elements which seem to play a role in the formation of these deletion mutants are:

--- the lack of a coat protein due to which the mutants are no longer encapsulated and all the positive-strand RNA in the infected cells remains available for replication and possibly for the spread of the infection

--- since larger deletions (without coat protein gene) evolve further to even smaller genomes, it can be assumed that the mutants replicate faster and undergo a more efficient infection according as they are smaller (invariably more STNV-RNA is found in the deletion mutant plants than in the wild type plants, and the amount increases according as the mutants become smaller).

The most important observation in the performance of these deletion mutants, however, is their strong interference with the TNV helper virus. When in an RNA infection line STNV deletion mutants are formed, the amount of TNV decreases very sharply. This decrease is so strong after 1 or 2 further infections that without addition of fresh TNV or TNV-RNA to the RNA inoculum no new infections can be carried out any more. Accordingly, the STNV mutants which are formed here by

the unnatural RNA inoculations appear to yield a very efficient repression of the TNV infection.

For comparison the in vitro constructed deletion mutant STNV D1 was infected on cowpea by plasmid infection. When this mutant was propagated by RNA inoculation in accordance with the in vivo deletions, its replication proved to be more efficient than that of the original N164 mutant. Like the mutants that form in vivo, STNV D1 represses the TNV replication. This is evidenced by a minor infectivity of an RNA inoculum from these plants.

The deletion mutant STNV D1 constructed in vitro was also transformed to tobacco SR1 to investigate whether the influence on TNV replication would lead to protection of the transformed plants against TNV infection. To determine the effect of the expression of the RNA of the STNV deletion mutant on the development and the course of the infection with TNV, for comparison the following plants were grown under identical conditions as much as possible, infected and further observed:

- SR1
- SR1 STNV N164N843.1/1 and /2, with strong and weak expression of STNV, respectively
- SR1 STNV D1.1/1

In each of these plants 4 leaves were infected which had a development and differentiation of a similar nature. For the infection a fresh inoculum, prepared from TNV-infected cowpea, and a frozen inoculum from infected tobacco were used, both

diluted 1/1 (designated inoculum 1 and 3, respectively) and 1/10 (inocula 2 and 4). Two half leaves of the various plants were infected with each of the 4 inocula at similar locations; these leaves were designated a, b, c, and d from the bottom to the top. Thus the following combinations were made: a1 and a2, b3 and b4, c2 and c4, and d1 and d3.

After 72, 96 and 170 h the development of the infection was observed and photographically recorded. Also, after 96 h particles were taken from each leaf and photographed under UV lighting to investigate the hypersensitivity response of the plants. After 96 h the half leaf of each plant with the best developed lesions (b3) was picked. A part of it was frozen and retained for further analysis of RNA and protein composition. The other part was used for the preparation of an inoculum to infect cowpea leaves. For each of the tobacco plants 4 cowpea leaves were infected:

- with an extract from one single lesion (1 leaf)
- with an extract from 200 mg leaf tissue (2 leaves)
- with an 1/5 dilution thereof (1 leaf)

The photographs (not shown) revealed that a great difference occurred in the development of the TNV infection in the various tobacco plants. The most important differences were:

- the number of lesions on the various leaves differed according to the inoculum used ($3 > 4 > 1 > 2$), with a minor difference between the plants (SR1 > SR1 N164N843.1/1 and /2 > SR1 D1.1/1)

--- the lesions in SR1 D1.1/1 are clearly smaller than in the other plants; tobacco SR1 and N164N843.1/1 differ most, while N164N843.1/2 is in-between. The differences in size of the lesions on one and the same plant depending on the leaf (a > b > c > d) are normal and are associated with the stage of development of the leaves

--- particularly after 170 h it is clear that fairly large portions of the infected leaves of SR1 and SR1 N164N843.1/1 necrotize completely, while the corresponding leaves of D1/1, it is true, show lesions, but otherwise remain normally green --- UV lighting makes visible fluorescent rings and stains which are a result of the hypersensitivity reaction of the plant. Due to this reaction infected parts of the plant are isolated by a ring of tissue which leads to these parts dying off. For SR1 D1.1/1 these parts are much smaller than for the other plants. This points to a less extensive infection of the leaf.

After 96 h of each plant a part of the half leaf b3 was used to prepare an inoculum for the cowpea leaves to be subsequently infected with. After 72 h very clear differences in the extent of infection of the cowpea leaves were to be observed. The plant which had been infected with the inocula prepared from the deletion plant yielded lesions but each of the leaves was still green. After 72 h the necrotization of the other plants had progressed much further and several of the leaves were so infected that they were already withered completely and almost fell off.

To compare the amount of TNV-RNA in the various plants, it was attempted to demonstrate the presence of TNV ds and ssRNA in EtBr-stained agarose gels. The amount of TNV-coat protein was determined by Western blotting. It was found that the amount of TNV-RNA in SR1 D1.1/1 was smaller than in the other plants. The amount of coat protein in SR1 D1.1/1 was about 20 times lower than that in tobacco SR1, while the amount in SR1 N164N843.1/1 and/2 was in between the two. The secondarily infected cowpea leaves were also examined for the presence of TNV-RNA and this yielded the same picture as in the tobacco plants.

In the case of grand-scale infection of the plant or portions thereof, the presence of RNA of STNV deletion mutants leads to no, less, or delayed complete necrotization and loss of the infected portions. In infected but protected plants much less TNV is produced, which has an inhibitory effect on the spread of the infection to other plants. The total protective effect is accomplished with the synthesis of a negligible amount of RNA. Only when the infection actually occurs do the virally encoded replication functions produce high and protection-yielding concentrations of deletion-RNA. Moreover this occurs only in the infected cells, so that the energy burden on the plant is negligible.

25 Example II

In the experiments described herein use was made of the glucuronidase gene of E. coli, because the enzyme

glucuronidase retains its activity as fusion protein, no plant glucuronidase activity is known and the enzyme activity can be detected in a quantitative and very sensitive manner. The glucuronidase (GUS) gene (*uidA*) used in the various constructions was isolated as PstI-EcoRI fragment from the plasmid pRAJ255, described by Jefferson et al in PNAS 83, 8447-8451 (1986). To enable the GUS gene to be inserted in the desired position and in the proper reading frame in the STNV-cDNA, the following adjustments were carried out:

--- by substitution of the 1549 bp ScaI-NdeI fragment of the plasmid pPLC 321 by the corresponding fragment of plasmid pPLC 2820 (which leads to the removal of the PstI site in the ampicillin gene) the plasmid pPLC 322 was constructed

--- by insertion in the unique NcoI site of pPLC 322 of a synthetic linker of the structure:

CATGGCTAGCAGCTGCAGGAATTCATGCATC

CGATCGTCGACGTCCTTAAGTACGTAGGTAC

the plasmid pPLC 323 was obtained

--- by cloning the PstI-EcoRI fragment containing the GUS gene from pRAJ255 into pPLC 323, the plasmid pPLC GUS1 was obtained, in which the GUS gene is flanked by two NcoI cutting sites

--- in pPLC GUS1 a unique PstI cutting site is located right before the coding portion of the GUS gene; by cutting the plasmid with PstI and removing the 4-base long 3'-overhanging ends by means of T4-polymerase, the plasmid pPLC GUS dPst1.1 was obtained, which still contains the two NcoI sites.

The insertions of the GUS gene were performed in the following STNV plasmids:

--- pPLC STNV N160.1 and .2 (the sense and the anti-sense orientation relative to the pL promoter), with a complete STNV genome, provided with a 14 bp linker insertion at position 160. The inserted linker contains a unique NcoI cutting site --- by linker mutagenesis at position 28 of the STNV genome the base A was replaced by the base C, which leads to the formation of an NcoI cutting site which overlaps the initiation codon of the coat protein gene. The introduction of this mutation into STNV N160 yielded the plasmids pPLC STNV N160Nco1.1 and .2. After cutting with NcoI and ligation the 134-bp long NcoI fragment of these plasmids can be deleted, which leads to the plasmids pPLC STNV dNco1.1 and .2.

The insertions of the GUS gene in STNV were accomplished as follows:

--- the GUS gene was cloned as NcoI fragment from pPLC GUS1 into pPLC STNV N160.1 and .2. In the proper orientation this yielded the plasmids pPLC STNV GUS.1 and .2, the GUS gene being fused in the proper reading frame with the 5'-portion of the STNV coat protein. Translation of the fusion protein stops at the location of the stop codon of the GUS gene, the 3'-portion of the coat protein is not translated. In fact these constructions represent the pure insertion of the GUS gene in STNV

--- the GUS gene was cloned as NcoI fragment from pPLC GUS dPst1.1 into the NcoI cutting site of pPLC STNV dNco1.1 and

.2, which yielded the plasmids pPLC STNV GUS dNco1.1 and .2. The insertion was such that the AUG initiation codon of STNV was retained, followed by 7 codons from the synthetic linker and adjoining the initiation codon of the GUS gene in the
5 proper reading frame. Translation stops likewise at the location of the stop codon of the GUS gene and thus provides the glucuronidase with 8 additional amino acids at the amino-terminal end

--- the plasmids pPLC STNV GUS.1 and .2 and pPLC STNV GUS
10 dNCO1.1 and .2 were cut with NsiI (this restriction enzyme cuts right behind the GUS gene and right at the end of the coat protein gene) and ligated. Thus an NsiI fragment of 468 bp was deleted and the plasmids pPLC STNV GUS dNsi1.1 and .2, and pPLC STGUS.1 and .2 were obtained, respectively. These
15 last two plasmids pPLC STGUS.1 and .2 contain the constructions in which the coat protein gene is entirely substituted by the GUS gene.

Starting from the pPLC plasmids the constructions STNV GUS and STGUS were transcloned as BamHI-SalI fragment in the
20 plant vector pPCV 520, behind the pTR1' promoter. In the process the orientation relative to the promoter is reversed so that, for example, the plasmid pPCV STGUS.2 is formed from pPLC STGUS.1.

Expression in plants was tested by plasmid infections of
25 cowpea with the various STNV-GUS constructions. For that purpose the plasmids pPLC STNV GUS.1 and .2, and pPCV STNV GUS.1 en .2 were used. Cowpea leaves were infected with

plasmid and helper virus and picked 3 days after inoculation. A freshly picked leaf was incubated overnight in a 2 mM solution of X-Gluc (5-bromo-4-chloro-3-indolyl-beta-D-glucuronic acid) in 40 mM phosphate buffer, pH 7.4, at 37°C.

5 Then the leaves were fixated in glutaraldehyde (6 h) and decolourized with an ethanol series (from 30 to 95 %). Conversion of the X-Gluc by the glucuronidase leads to the formation of an insoluble blue precipitate.

With the four above-mentioned plasmids four independent
10 infection series were performed on cowpea, in two cases of which a localized blue reaction was observed in the incubated leaves. With pPLC STNV GUS.1 a blue edge was observed at the location of a lesion. Microscopic examination of the fixated material revealed that the stain reaction was localized in the
15 plant cells which were located right at the edge of the lesion. With pPLC STNV GUS.2 the reaction was visible in the form of a 0.3 mm stain. A microscopic examination could not establish whether the reaction was intracellular, because the plant cells had been damaged too severely by infection through
20 bacterial contamination during the incubation with the substrate. However, because the blue colour was preserved during fixation, decolourizing, embedding and cutting, it can be assumed with fairly great certainty that the reaction was plant-specific.

25 Further, also a transformation of tobacco was performed, using the T-plasmids pPCV STNV GUS.1 and .2. For both constructions transformants were obtained. GUS tests before

and after TNV infections of the grown transformants proved to meet the expectations.

Example III

5 Isolation of the gene for replicase of Tobacco Necrosis Virus

Used as starting material was Tobacco Necrosis Virus (TNV) Kassanis strain A and serotype A, which is capable of multiplying Satellite Tobacco Necrosis Virus (STNV) SV-1 (SV-A
10 serotype). This strain was obtained from Dr. Wieringa Brants, Phytopathologic Laboratory, Baarn, the Netherlands. TNV, free of STNV, was obtained as described by Van Emmelo et al., Virology 157, 480-487 (1987).

TNV virus was increased in Phaseolus leaves (Vigna
15 unguiculata, cowpea), by infection with virus and carborundum. Genomic RNA was prepared from purified TNV particles and converted into cDNA with AMV reverse transcriptase (Boehringer) using random primers. The second strand synthesis was performed by E. coli DNA polymerase I (Klenow fragment)
20 and RNAaseH (Boehringer) according to the method of Gubler and Hoffman (Gene 25, 263-269, 1983). Then T4 DNA polymerase was added to blunt the ends. After cloning into plasmid pSP64 (Promega), cut with SmaI with simultaneous treatment of the ends with phosphatase, the clones were transferred to E. coli
25 MC1061 by transformation and the colonies were tested by colony-hybridization with fragmented ³²P ds TNV RNA.

Subgenomic ssRNA prepared from Phaseolus leaves infected with TNV was used for Northern hybridization with the selected TNV clones. The lengths of these subgenomic RNAs were 1.25 kb, 1.5 kb and 3.8 kb, respectively. After agarose gel electrophoresis, the subgenomic RNA was immobilized on a nylon membrane filter (Pall Biodyne) and hybridized with ³²P labeled clone DNA. Selected was a clone (pSPTNV127), which hybridized with the two 3' subgenomic TNV RNAs and contained a fragment located between positions 1900 and 2300 of the TNV RNA.

Restriction analysis demonstrated that this fragment contained an internal HpaII fragment which could be used as a primer for the cDNA synthesis of the replicase gene which is located 5' terminally on the TNV RNA. For the preparatory cDNA cloning, however, use was made of a synthetic oligonucleotide, derived from the sequence of pSPTNV127: 5' GCTTGTGAGTATCA 3'. This sequence is complementary to the genomic RNA.

The synthesis of the cDNA was performed as described hereinabove but in the presence of methyl-mercury-hydroxide to make it easier for the RNA to be copied (Lenstra et al., Gen. Anal. Techn. 5, 57-61, 1988). Thus a 2.2 kb strand was obtained which corresponds to the region of the replicase gene of TNV. After cloning into pSP64, from 1500 clones three clones were selected which contain a 2.2 kb fragment in the sense orientation relative to the SP6 promoter.

Using SP6 polymerase and starting from the clone pSPTNV94, RNA was prepared which, in a wheat germ extract, led

inter alia to synthesis of a 73 kD protein, which is the expected molecular weight of the TNV replicase.

In this experiment ^{35}S methionine was used during the protein synthesis and the 73 kD protein was detected after SDS
5 PAGE and autoradiography.

The sequence of pSPTNV94 was determined by the dideoxy method of Sanger et al. (Proc. Natl. Acad. Sci. USA 74, 5463-5467, 1977) after cloning of fragments into pUC18 plasmid. The strategy for determining the nucleotide sequence is presented
10 in Fig. 3 and the sequence in Fig. 4.

The fragment in pSPTNV127 appeared to contain the 3' end of the replicase gene. To obtain a complete clone of the replicase gene of TNV, use was made of a unique XmaI site in pSPTNV94 and pSPTNV127 DNA and a unique SacI site in the
15 polylinker of pSP64. By a ligation of the small XmaI-SacI fragment of pSPTNV127 in the large pSPTNV94 fragment with XmaI and SacI ends, pSPTNV rep-1 was obtained. The clone is 2236 bases long, the longest open reading frame being found from base 50 to base 2221 (absolute reading frame relative to the
20 first base: 2) with an internal stop codon at position 656. This open reading frame, after translation and use of a suppressor tRNA at position 656-658, codes for a replicase of 724 amino acids. For further uses, the internal stop codon TAG, by means of specific mutagenesis, was changed into a
25 codon TAT, coding for Tyr.

An E. coli strain MC1061 [pSPTNV rep-1], which contains the plasmid pSPTNV rep-1, was registered on July 26, 1990 with

the Centraal Bureau voor Schimmelculturen (CBS), Baarn, the Netherlands, under number CBS 336.90.

Example IV

5 Expression, with amplification of the messenger RNA by TNV replicase, of a chloramphenicol-acetyl-transferase gene, fused with a part of the STNV coat protein gene

1. Construction of a DNA sequence in which a part of the
10 STNV coat protein gene is fused with the chloramphenicol-acetyl-transferase gene

The plasmid pSTNV N198 was described by Van Emmelo et al. (Virology 157, 480-487, 1987). It contains a 14 bp NcoI linker
15 in the EcoRV site of the coat protein gene of STNV. By cutting with NcoI, filling in the ends with E. coli polymerase and back ligation the reading frame was recovered again:
Asp(56)/Ser-Met-His-Gly-Asn-Ser/Ile(57).

The plasmid obtained gives rise to the formation of STNV
20 infective particles in coinfections with TNV of Phaseolus leaves (*Vigna unguiculata*, cowpea). This plasmid was called pSTNV N202. The PstI fragment, containing the STNV cDNA N202, was transcloned into pSP65 (Promega). The plasmid obtained was called pSPSTNV N202 (Fig. 5). From pBR325 the chloramphenicol-
25 acetyl-transferase gene (CAT gene) was taken as a TagI fragment, which was then blunt-ended with the Klenow fragment of E. coli DNA polymerase. pSPSTNV N202 was cut with NsiI and

blunt-ended. Then the TagI fragment was cloned into the NsiI deleted pSPSTNV N202 plasmid.

The sense orientation relative to the SP6 promoter was called pSPSTNV CAT-I and the non-sense orientation pSPSTNV
5 CAT-2 (see Fig. 5).

2. Expression of the chimeric CAT gene using TNV virus in Phaseolus (Vigna unguiculata) leaves

Leaves of cowpea were infected with either TNV and STNV
10 plasmid, or TNV and pSPSTNV CAT-1. Forty hours after inoculation the leaves were extracted with PBS buffer with 0.1% Triton X100 and the extract was treated with anti-STNV coat protein serum. Immunoprecipitation with protein A-sepharose was followed by frequent washing.

15 The chloramphenicol-acetyl-transpherase activity was determined on the complexes with 1-¹⁴C-acetyl-coenzyme A and chloramphenicol as described by Gorman et al. (Molec. Cell. Biol. 2, 1044-1081, 1982) and by De Block et al. (EMBO J. 3, 1681-1689, 1984).

20 CAT activity could only be demonstrated in the leaves which were infected with TNV and pSPSTNV CAT-1 and the extract of which was treated with STNV antiserum (the results are not shown here).

25 3. Multiplication of the STNV-CAT RNA

Two days after inoculation of cowpea leaves with TNV and pSPSTNV CAT-1 DNA, the leaves were picked and the nucleic acid fraction was isolated. DNA was removed with RNAase-free DNAase I and the RNA was analyzed by Northern hybridization with CAT specific, ^{32}P labeled DNA (TagI fragment from pBR325). Both in the ss RNA and the ds RNA fraction a band of about 1750 nucleotides was detected (the results are not shown here). This clearly demonstrates that the pSPSTNV CAT-1 plasmid DNA is converted by TNV into ss and ds STNV CAT-1 RNA.

10

Description of the Figures

Fig. 1 shows the map of the plasmid pPCV 520. The T-DNA of this plasmid is indicated by a thick line and is bounded by left-hand (LB) and right-hand (RB) border sequences; the unique restriction sites for SalI and BamHI can be used for cloning fragments between the plant promoter pTR1' and the polyadenylation signal pAg7, and the restriction sites BglII and BclI for cloning between pnos and pAg4. The kanamycin resistance gene NPT-II is expressed by pTR2' and pAocs. The octopine gene is expressed with the tissue-specific promoter pg5 and pAocs. The pBR322-sequences with the replication-origin (oriV, colE1) and the ampicillin resistance (Ap) are located in the T-DNA. The pRK2 origin of replication (ori V) and transfer (ori T), the chloramphenicol resistance gene (Cm) and the cohesive ends of lambda (cos) are located outside the T-DNA.

25

Fig. 2 shows the map of the plasmid pMP90RK. At the location where the KpnI fragments 9 and 12 of pGV 3100 adjoin one another, the T-DNA inclusive of the border sequences was deleted. The virulence region is entirely intact. By exchange
5 recombination the gene for gentamycin resistance (Gm) and a fragment of pRK 2013, which carries the genes for transfer (tra1, tra2 and tra3), replication (trfa) and kanamycin resistance (Km), were inserted.

Fig. 3 schematically shows the restriction map of the
10 region of the TNV replicase gene and the fragments used for determining the sequence. After cloning into pUC18 the fragments were analyzed in both directions or twice according to the dideoxy method of Sanger to determine the nucleotide sequence.

15 Fig. 4 shows the complete sequence of the replicase gene of TNV. The amino acid sequence exhibits homology with the replicase genes of CarMV (Carnation Mottle Virus) and TuCV (Turnip Crinkle Virus). The amber stop codon at position 656 is present in the natural genome. For further uses this stop
20 codon was replaced by TAT which codes for the amino acid Tyr. The initiation and stop codons are indicated by ▲ and ▼, respectively. The amino acid sequence starts at nucleotide position 50 and stops at position 2221.

Fig. 5 schematically shows the construction of the
25 plasmids pSPSTNVCAT1 and pSPSTNVCAT2. Details about this construction are described in Example IV.

Table 1

	STNV mutant	<u>E. coli</u> NF1 coat protein	cowpea coat protein	ssRNA	dsRNA
5	-----	-----	-----	-----	-----
	wild type	+	+	++	++
	N162	+ (short)	-	+	++
	N164	+ (+6 a.a.)	+ (+6 a.a.)	++	++
	S164	+ (short)	-	+	++
10	Brad	+ (+10 a.a.)	+ (+10 a.a.)	++	++
	N198	+ (short)	-	+	++
	N200	+ (+6 a.a.)	+ (+6 a.a.)	++	++
	S200	+ (+12 a.a.)	-	+	++
	N320	+ (short)	-	+	++
15	N322	+ (+6 a.a.)	-	+	++
	N531	+ (short)	-	+	++
	N533	+ (+6 a.a.)	-	+	++
	S613	+ (+6 a.a.)	-	+	++
	N843	+	-	-	-
20	N164N843	+ (+6 a.a.)	-	-	-

"a.a." stands for amino acids; "short" indicates a shortened coat protein. The presence and the size of the coat protein were determined by Western blotting. RNA was hybridized with an STNV-DNA probe, which was labeled with ^{32}P .

Table 2

	STNV and control plants	STNV coat protein		STNV RNA (ds and ss)				
		- TNV	+ TNV	- TNV	+ TNV		14	18
				with STNV DNA				30
	N164.1	-	+ to +++	-	+ to +++	-	+	-
	N164.2	-	+ to +++	-	+ to +++	-	+	-
35	N164N843.1	-	++ to +++	-	++ to +++	+	+	-
	N164N843.2	-	+ to +++	-	+ to +++	+	+	-
	Brad.1	-	+ to +++	-	+ to +++	-	-	+
	Brad.2	-	++	-	++	-	-	+
	tobacco SR1	-	-	-	-			
40	520	-	-	-	-			

Of the transformed plants, per STNV mutant several independently isolated plants were tested. STNV coat protein (+6 and +10 amino acids) was determined by Western blotting, RNA by Northern hybridization with ^{32}P -labeled STNV-DNA. Independent transformants with the same mutant genome yielded signals of different intensity, which is indicated by the

number of + signs. For the Western and Northern blottings per sample 10 mg leaf material was used. By hybridization with the various linkers (14-, 18- and 30-mer) the identity of the STNV genomes was confirmed.

184-302D

57

CLAIMS:

1. An eukaryotic organism comprising recombinant DNA incorporated into its genome, said recombinant DNA comprising an expression cassette comprising two inverted
5 repeat nucleotide sequences of 12 to 1250 basepairs with there between at least one nucleotide sequence which is a stripped viral replicon of an RNA virus whose replication is dependent upon a viral RNA/RNA polymerase, wherein said stripped viral replicon comprises *cis* elements for
10 replication but is devoid of genes coding for viral RNA/RNA polymerase and viral coat protein, wherein the replicon is unable to replicate independently, and wherein the replicon, in the presence of viral RNA/RNA polymerase, is able to replicate more efficiently than wild-type viral RNA.
- 15 2. The eukaryotic organism according to claim 1, wherein said stripped viral replicon further comprises *cis* elements for transport.
3. The eukaryotic organism according to claim 1 or 2, wherein said stripped viral replicon also comprises *cis*
20 elements for packing in coat protein.
4. The eukaryotic organism according to any one of claims 1-3, wherein said stripped viral replicon also comprises *cis* elements for translation.
5. The eukaryotic organism according to any one of
25 claims 1-4, wherein the expression cassette further comprises, between the two inverted repeat nucleotide sequences, at least one non-viral nucleotide sequence coding for one or more proteins or peptides.
6. The eukaryotic organism according to any one of
30 claims 1-5, further comprising a second recombinant DNA

184-302D

58

incorporated into its genome, which second recombinant DNA comprises a polynucleotide whose expression is inducible or tissue-specific and which codes for a viral RNA/RNA polymerase or an RNA/RNA polymerase construct.

5 7. The eukaryotic organism according to any one of claims 1-6, which is a plant.

8. The eukaryotic organism according to any one of claims 1-6, which is an animal.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS

1/8




FIG.1




FIG.2

2/8




FIG.3

3/8

10 20 30 40 50 60
| | | | |
CCAAGAATACCAATAGGTGCAAGGCCCTTACTCAGCTAAAGAGTCTAAAATGGAGCTACC
GlnGluTyrGlnIleGlyAlaArgProTyrSerAlaLysGluSerLysMETGluLeuPro
▲

70 80 90 100 110 120
| | | | |
AAACCAACACAAGCAATCAGCCGCCGAGGGTITTTGTATCTTTCCTAAACTGGCTATGCAA
AsnGlnHisLysGlnSerAlaAlaGluGlyPheValSerPheLeuAsnTrpLeuCysAsn

130 140 150 160 170 180
| | | | |
CCCATGGAGACGACAGCGAACAGTCAACGCTGCAGTTGTGTTCCAAAAGATCTTCTAGC
ProTrpArgArgGlnArgThrValAsnAlaAlaValValPheGlnLysAspLeuLeuAla

190 200 210 220 230 240
| | | | |
CATTGAGGATTCGAGCATTGGATGATATCAATGAGTGTTCGAAGAATCTGCTGGGGC
IleGluAspSerGluHisLeuAspAspIleAsnGluCysPheGluGluSerAlaGlyAla

250 260 270 280 290 300
| | | | |
ACAATCCCAGCGAACTAAGGTGTGCGCCGACGGAGCATATGCCCCGCAAAATCCAATAG
GlnSerGlnArgThrLysValValAlaAspGlyAlaTyrAlaProAlaLysSerAsnArg

310 320 330 340 350 360
| | | | |
GACCCGCCGAGTTCGTAAAGCAGAAAAAGCACAAGTTTGTCAAATATCTTGTCAACGAAGC
ThrArgArgValArgLysGlnLysLysHisLysPheValLysTyrLeuValAsnGluAla

370 380 390 400 410 420
| | | | |
TCGTGCCGAGTTTGGATTGCCTAAACCAACTGAGGCAAACAGACTCATGGTCCAACATTT
ArgAlaGluPheGlyLeuProLysProThrGluAlaAsnArgLeuMETValGlnHisPhe

FIG. 4

4/8

430 440 450 460 470 480
CTTGCTCAGGGTGTGTAAAGGATTGGGGCGTTGTTACAGCCCACGTACACGGCAACGTTGC
LeuLeuArgValCysLysAspTrpGlyValValThrAlaHisValHisGlyAsnValAla

490 500 510 520 530 540
ACTAGCTTTGCCACTGGTGTTCATTCCAACGGAAGATGATCTGCTATCACGAGCATTGAT
LeuAlaLeuProLeuValPheIleProThrGluAspAspLeuLeuSerArgAlaLeuMET

550 560 570 580 590 600
GAACACACATGCTACTAGAGCTGCTGTACGAGGCATGGACAAATGTCCAAGGCCAAGGGTG
AsnThrHisAlaThrArgAlaAlaValArgGlyMETAspAsnValGlnGlyGlnGlyTrp

610 620 630 640 650 660
GTGGAACAATAGGTTGGGGATTGGGGGCCAGATCGGACTGGCCTTCCGGTCCAAATAGGG
TrpAsnAsnArgLeuGlyIleGlyGlyGlnIleGlyLeuAlaPheArgSerLys---Gly
▼

670 680 690 700 710 720
GTGCCTTGAAAGGAGGCCAGGATCTCCACGTCCGTTTCGCGTGGGGAACATCCTGATCT
CysLeuGluArgArgProGlyPheSerThrSerValSerArgGlyGluHisProAspLeu

730 740 750 760 770 780
GGTGGTCATACCATCAGGGCGCCCTGAGAAACAGCGTCAGTTGTTACGCTACAGTGGTAT
ValValIleProSerGlyArgProGluLysGlnArgGlnLeuLeuArgTyrSerGlyIle

790 800 810 820 830 840
AGGCGGCCATTTATTAATCGGCATCCACAACAACTCTCTTTCCAACCTTGCGTAGGGGCTT
GlyGlyHisLeuLeuIleGlyIleHisAsnAsnSerLeuSerAsnLeuArgArgGlyLeu

850 860 870 880 890 900
GATGGAGAGAGTATTCTACGTCGAGGGGCCCAATGGGCTCCAAGACGCCCTAAGCCCGT
METGluArgValPheTyrValGluGlyProAsnGlyLeuGlnAspAlaProLysProVal

5/8

910 920 930 940 950 960
 CAAGGGAGCTTTCCGGACCCTTGATAAGTTTCGTGATCTCTATACTAAAAATAGTTGGCG
 LysGlyAlaPheArgThrLeuAspLysPheArgAspLeuTyrThrLysAsnSerTrpArg

970 980 990 1000 1010 1020
 TCATACCCCTGTAAGTAGTGAACAATTCCTAATGAATTACACGGGCAGGAAACTGACTAT
 HisThrProValThrSerGluGlnPheLeuMEIAsnTyrThrGlyArgLysLeuThrIle

1030 1040 1050 1060 1070 1080
 TTACAGAGAGGCGGTTGATAGTTTGTGCGCATCAACCCCTTAGCTCACGAGATGCGAAGCT
 TyrArgGluAlaValAspSerLeuSerHisGlnProLeuSerSerArgAspAlaLysLeu

1090 1100 1110 1120 1130 1140
 AAAGACATTCGTGAAGGCCGAAAAATTAAACCTTTCTAAGAAGCCTGACCCTGCTCCCAG
 LysThrPheValLysAlaGluLysLeuAsnLeuSerLysLysProAspProAlaProArg

1150 1160 1170 1180 1190 1200
 GGTCATACAACCTAGATCGCCTCGGTATAACGTTTGTGTTGGGCAGGTACCTCCGACATTA
 ValIleGlnProArgSerProArgTyrAsnValCysLeuGlyArgTyrLeuArgHisTyr

1210 1220 1230 1240 1250 1260
 TGAACATCACGCGTTTAAAACCATTGCCAAGTGCTTTGGGGAAATCACGGTCTTCAAAGG
 GluHisHisAlaPheLysThrIleAlaLysCysPheGlyGluIleThrValPheLysGly

1270 1280 1290 1300 1310 1320
 GTTTACTCTGGAGCAACAAGGGGAAATCATGCGCTCGAAGTGGAATAAATATGTTAATCC
 PheThrLeuGluGlnGlnGlyGluIleMETArgSerLysTrpAsnLysTyrValAsnPro

1330 1340 1350 1360 1370 1380
 CGTTGCGGTCGGACTTGACGCCAGTCGTTTCGACCAACACGTGTCTGTTGAAGCACTCGA
 ValAlaValGlyLeuAspAlaSerArgPheAspGlnHisValSerValGluAlaLeuGlu

1390 1400 1410 1420 1430 1440
 GTATGAGCATGAATTTTATCTCAGAGATTACCCAATGATAAACAGCTAAAATGGCTGCT
 TyrGluHisGluPheTyrLeuArgAspTyrProAsnAspLysGlnLeuLysTrpLeuLeu

6/8

1450 1460 1470 1480 1490 1500
AAAGCAGCAATTGTGCAACGTGGGAACGGCATTGCGCCAGTGACGGTGTTATAAAATACAA
LysGlnGlnLeuCysAsnValGlyThrAlaPheAlaSerAspGlyValIleLysTyrLys

1510 1520 1530 1540 1550 1560
GAAGAAGGGTTGTAGAATGAGCGGAGACATGAACACGAGTTTGGGCAACTGCATTITTAAT
LysLysGlyCysArgMETSerGlyAspMETAsnThrSerLeuGlyAsnCysIleLeuMET

1570 1580 1590 1600 1610 1620
GTGCGCCATGGTCTACGGGTGAAAGAACACTTAAACATCAATTTGTCCCTTGCAAATAA
CysAlaMETValTyrGlyLeuLysGluHisLeuAsnIleAsnLeuSerLeuAlaAsnAsn

1630 1640 1650 1660 1670 1680
TGGGGATGACTGCGTCATTGTCTGTGAGAAAGCGGATTTAAAGAAATTGACGAGCAGCAT
GlyAspAspCysValIleValCysGluLysAlaAspLeuLysLysLeuThrSerSerIle

1690 1700 1710 1720 1730 1740
CGAGCCATATTTCAAGCAGTTTGGATTCAAGATGGAAGTGGA AAAACCCGTGGATATCTT
GluProTyrPheLysGlnPheGlyPheLysMETGluValGluLysProValAspIlePhe

1750 1760 1770 1780 1790 1800
TGAGCGTATAGAATTTTGCCAAACCCAGCCTGTGTTTGATGGATCCCAATATATCATGGT
GluArgIleGluPheCysGlnThrGlnProValPheAspGlySerGlnTyrIleMETVal

1810 1820 1830 1840 1850 1860
ACGCAAACCTTCAGTTGTAACATCCAAAGACGTCACCAGCCTCATCCCATGTCAAACGAA
ArgLysProSerValValThrSerLysAspValThrSerLeuIleProCysGlnThrLys

1870 1880 1890 1900 1910 1920
AGCACAATACGCAGAATGGCTGCAAGCTGTGGGTGAGTGTGGCATGAGCATCAATGGTGG
AlaGlnTyrAlaGluTrpLeuGlnAlaValGlyGluCysGlyMETSerIleAsnGlyGly

7/8

1930 1940 1950 1960 1970 1980
GATTCCTGTTATGCAGAATTTCTACCAGATGCTCCAAACTGGCATCCGCCGCACAAAATT
IleProValMETGlnAsnPheTyrGlnMETLeuGlnThrGlyIleArgArgThrLysPhe

1990 2000 2010 2020 2030 2040
CACCAAGACCGGCGAGTTCCAGACGAACGGATTGGGGTATCACTCTAGATTTATGCATAG
ThrLysThrGlyGluPheGlnThrAsnGlyLeuGlyTyrHisSerArgPheMETHisArg

2050 2060 2070 2080 2090 2100
AGTGGCCCGGGTCCCTTCGCCTGAAACCCGTTTATCCTTCTATCTAGCTTTCGGTATCAC
ValAlaArgValProSerProGluThrArgLeuSerPheTyrLeuAlaPheGlyIleThr

2110 2120 2130 2140 2150 2160
ACCAGACCTCCAAGAAGCAATGGAGATCTTCTATGATACTCACAAGCTTGATTGGATGA
ProAspLeuGlnGluAlaMETGluIlePheTyrAspThrHisLysLeuAspLeuAspAsp

2170 2180 2190 2200 2210 2220
TGTTATCCCGACTGATACCTACCAAGTGTCAGGAGAGCATTGATCAATGGATTACCAA
ValIleProThrAspThrTyrGlnValSerGlyGluHisLeuIleAsnGlyLeuProAsn

2230
CTGATGTAACGGAGGA

---CysAsnGlyGly



8/8




FIG.5

