

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Processes for producing 3-substituted 2-chloro-5-fluoropyridine or salt thereof

(51)⁶ International Patent Classification(s)
C07D 213/80 20060101AFI2005100
(2006.01) 8BMEP
C07D 213/80 PCT/JP2004/006267

(21) Application No: 2004236085 (22) Application Date: 2004 .04 .30

(87) WIPO No: W004/099149

(30) Priority Data

(31) Number	(32) Date	(33) Country
2003-132137	2003 .05 .09	JP

(43) Publication Date : 2004 .11 .18

(71) Applicant(s)
Asahi Glass Company, Limited

(72) Inventor(s)
Yamada, Yasuhiro; Matsumura, Yasushi; Mori, Nobuaki

(74) Agent/Attorney
Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
Setliff et al. Journal of Chemical and Engineering Data 1972, 17(4), 515-516
WO 1998/050362
US 5204478
EP 333020
Winn et al. J. Med. Chem. 1993, 36(18), 2676-2688
US 5250548
EP 634413
EP1229034

(19) 世界知的所有権機関
国際事務局(43) 国際公開日
2004年11月18日 (18.11.2004)


PCT

(10) 国際公開番号
WO 2004/099149 A1


- (51) 国際特許分類: C07D 213/803, 213/85 (74) 代理人: 泉名 謙治, 外 (SENMYO, Kenji et al.); 〒1010042 東京都千代田区神田東松下町3番地 島本鋼業ビル Tokyo (JP).
- (21) 国際出願番号: PCT/JP2004/006267
- (22) 国際出願日: 2004年4月30日 (30.04.2004)
- (25) 国際出願の言語: 日本語
- (26) 国際公開の言語: 日本語
- (30) 優先権データ:
特願2003-132137 2003年5月9日 (09.05.2003) JP
- (71) 出願人 (米国を除く全ての指定国について): 旭硝子株式会社 (ASAHI GLASS COMPANY, LIMITED) [JP/JP]; 〒1008405 東京都千代田区有楽町一丁目1番1号 Tokyo (JP).
- (72) 発明者: および
- (75) 発明者/出願人 (米国についてのみ): 松村 靖 (MATSUMURA, Yasushi) [JP/JP]; 〒2908566 千葉県市原市五井海岸1番地 旭硝子株式会社内 Chiba (JP). 森信明 (MORI, Nobuaki) [JP/JP]; 〒2908566 千葉県市原市五井海岸1番地 旭硝子株式会社内 Chiba (JP). 山田 康弘 (YAMADA, Yasuhiro) [JP/JP]; 〒2908566 千葉県市原市五井海岸1番地 旭硝子株式会社内 Chiba (JP).
- (81) 指定国 (表示のない限り、全ての種類の国内保護が可能): AF, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) 指定国 (表示のない限り、全ての種類の広域保護が可能): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), ユーラシア (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), ヨーロッパ (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- 添付公開書類:
国際調査報告書
- 2文字コード及び他の略語については、定期発行される各PCTガゼットの巻頭に掲載されている「コードと略語のガイダンスノート」を参照。

(54) Title: PROCESSES FOR PRODUCING 3-SUBSTITUTED 2-CHLORO-5-FLUOROPYRIDINE OR SALT THEREOF


(54) 発明の名称: 2-クロロ-5-フルオロ-3-置換ピリジンまたはその塩の製造方法



(1)



(2)



(4)

(57) Abstract: Processes for producing in high yield a 3-substituted 2-chloro-5-fluoropyridine or salt thereof from an easily available material through a short step under mild reaction conditions using a reagent which can be reacted by a simple procedure and is easy to handle. One of the processes is a process for producing the compound (2) shown below or a salt thereof which comprises reducing the compound (1) shown below or a salt thereof selectively at the 6-position chlorine atom. The other is a process for producing the compound (4) shown below or a salt thereof which comprises replacing the group Z¹ of the compound (2) or salt obtained by the reduction reaction with the group Z². In the following formulae, Z¹ represents -CO₂R¹, etc., wherein R¹ represents hydrogen, alkyl, alkenyl, aryl, aralkyl, or cycloalkyl; and Z² is a group different from Z¹ and represents -CO₂R², etc., wherein R² represents alkyl, alkenyl, aryl, aralkyl, or cycloalkyl.


[続葉有]

WO 2004/099149 A1




(57) 要約:


入手容易な原料から、反応操作が簡便で取扱いの容易な試薬を用いて、短い工程で、かつ温和な反応条件で、2-クロロ-5-フルオロ-3-置換-ピリジンまたはその塩を高収率で製造する方法を提供する。すなわち、下記化合物(1)またはその塩の6位の塩素原子を選択的に還元する下記化合物(2)またはその塩の製造方法を提供する。また、該還元反応によって得た下記化合物(2)またはその塩の Z^1 基を Z^2 基に置換することによる、下記化合物(4)またはその塩の製造方法を提供する。ただし、 Z^1 は、 $-\text{CO}_2\text{R}^1$ 等であり、 R^1 は、水素原子、アルキル基、アルケニル基、アリール基、アルアルキル基またはシクロアルキル基を示す。 Z^2 は Z^1 と異なる基であり、 $-\text{CO}_2\text{R}^5$ 等であり、 R^5 はアルキル基、アルケニル基、アリール基、アルアルキル基またはシクロアルキル基を示す。



(1)



(2)



(4)

1
DESCRIPTION

METHOD FOR PRODUCING 3-SUBSTITUTED 2-CHLORO-5-FLUORO-
PYRIDINE OR ITS SALT

5

TECHNICAL FIELD

The present invention relates to a novel method for producing a 3-substituted 2-chloro-5-fluoro-pyridine or its salt, such as 2-chloro-5-fluoronicotinic acid or its
10 salt. The 3-substituted 2-chloro-5-fluoro-pyridine or its salt is a compound useful as an intermediate for pharmaceuticals or for agricultural chemicals and further as an intermediate or the like for production of various functional materials.

15

BACKGROUND ART

With respect to a method for producing a 3-substituted 2-chloro-5-fluoro-pyridine or its salt, the following methods have heretofore been proposed.

20

(1) A method wherein 2,6-dichloro-5-fluoronicotinic acid is used as the starting material, and it is converted to an ethyl ester of 2,6-dichloro-5-fluoronicotinic acid and then reacted with sodium thiomethoxide to obtain ethyl 2-chloro-5-fluoro-6-
25 methylthionicotinate, and further, the methylthio group at the 6-position is reduced in the presence of a Raney nickel catalyst to obtain ethyl 2-chloro-5-fluoro-

nicotinate (U.S. Patent 5,250,548 (Examples 180A to C)).

(2) A method wherein 2-hydroxy nicotinic acid is used as the starting material, and the 5-position is nitrated, then the 2-position is chlorinated, then the
5 nitro group at the 5-position is reduced in hydrochloric acid by means of tin(II) chloride to obtain 5-amino-2-chloronicotinic acid hydrochloride, which is then diazotized by means of tetrafluoroboric acid and sodium nitrite to convert the 5-position to diazonium
10 tetrafluoroborate, followed by thermal decomposition in 1,2-dichlorobenzene to obtain 2-chloro-5-fluoronicotinic acid (European Patent No. 634413A1 (pages 12 and 13)).

(3) A method wherein 2-chloro-3-methyl-5-nitropyridine is used as the starting material, and the
15 nitro group is reduced and then diazotized in HPF_6 to convert the 5-position to diazonium hexafluorophosphate, followed by thermal decomposition to obtain 2-chloro-5-fluoro-3-methylpyridine, which is further oxidized with potassium permanganate to obtain 2-chloro-5-
20 fluoronicotinic acid (Frank L. Setliff, Gary O. Rankin, "J. Chemical and Engineering Data", (U.S.A.), 1972, Vol. 17, No. 4, p. 515).

However, the above-methods have the following drawbacks.

25 In the method (1), the reaction for introducing a methylthio group is a reaction accompanying a bad odor. Further, in that method, after substituting a hydrogen

2004236085 23 Feb 2010

3

atom by a methylthio group, the methylthio group is again substituted to a hydrogen atom by a reduction reaction, and as such, the method is inefficient. Further, there is an additional drawback that the yield in such a reduction
5 reaction is low at a level of 30%. The method (2) has a drawback that the reaction process is long. Further, after synthesizing an unstable diazonium salt, the diazonium salt is thermally decomposed at a high temperature, and accordingly, the method has a drawback that the overall
10 yield is low. The method (3) also has a drawback that the reaction process is long. Further, it has a drawback that the yield in the oxidation reaction is so low that the method is poor in practical applicability.


15 DISCLOSURE OF THE INVENTION

In one or more aspects the present invention may advantageously provide a method for producing in good yield a 3-substituted 2-chloro-5-fluoro-pyridine or its salt which is useful as e.g. an intermediate for production of
20 pharmaceuticals, agricultural chemicals and various functional materials, from a starting material which is industrially readily available, through a short process under mild reaction conditions using a reagent which is easy in handling and simple in the reaction operation.


25 Namely, the present invention provides the

following.


(1) A method for producing a compound represented by the following formula (2) or its salt, which comprises selectively reducing a chlorine atom at the 6-position of
 5 a compound represented by the following formula (1) or its salt, provided that in the following formulae, Z^1 is a group represented by $-\text{CO}_2\text{R}^1$, $-\text{CONR}^2\text{R}^3$ or $-\text{CN}$, wherein each of R^1 , R^2 and R^3 which are independent of one another, is a hydrogen atom, an alkyl group, an alkenyl
 10 group, an aryl group, an aralkyl group or a cycloalkyl group:



(2) A method for producing a compound represented by the following formula (2), which comprises selectively
 15 reducing a chlorine atom at the 6-position of a compound represented by the following formula (1), provided that in the following formulae, Z^1 is a group represented by $-\text{CO}_2\text{R}^1$, $-\text{CONR}^2\text{R}^3$ or $-\text{CN}$, wherein each of R^1 , R^2 and R^3 which are independent of one another, is a hydrogen atom,
 20 an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group:



(1)




(2)

(3) The method according to (1) or (2), wherein Z^1 is $-CO_2R^1$ (wherein R^1 is as defined above).

(4) The method according to any one of (1) to (3),
 5 wherein the reduction is carried out by letting a protic solvent act in the presence of a metal or a metal salt.

(5) The method according to any one of (1) to (4), wherein the reduction is carried out by letting a protic solvent act in the presence of zinc.

10 (6) A method for producing a compound represented by the following formula (4) or its salt, which comprises carrying out a substitution reaction of a compound represented by the formula (2) or its salt obtained by the method as defined in any one of (1) to (5), to
 15 substitute the Z^1 group at the 3-position by a Z^2 group, thereby to obtain a compound represented by the following formula (4) or its salt, provided that Z^2 in the following formula is a group different from Z^1 and is a group represented by $-CO_2R^5$, $-CONR^6R^7$ or $-CN$, wherein R^5
 20 is an alkyl group, an alkenyl group, an aryl group, an aralkyl group, or a cycloalkyl group, and each of R^6 and R^7 which are independent of each other, is a hydrogen atom, an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group:



(4)

(7) The method according to (6), wherein from the compound of the formula (2) or its salt wherein the Z^1 group is a group represented by $-\text{COOH}$ or $-\text{COOR}^{10}$ (wherein R^{10} is an alkyl group), the compound (4) wherein Z^2 is a $-\text{COOR}^5$ group (wherein R^5 is as defined above), is obtained by an esterification reaction in a case where Z^1 is a $-\text{COOH}$ group, or by an ester exchange reaction in a case where Z^1 is a $-\text{COOR}^{10}$ group (wherein R^{10} is as defined above).

BEST MODE FOR CARRYING OUT THE INVENTION

In this specification, the compound represented by the formula (1) may be referred to also as the compound (1). The same applies to other compounds.

In this specification, "an alkyl group" may be linear or branched. Unless otherwise specified, the alkyl group is preferably a C_{1-6} lower alkyl group. As an example of the alkyl group, a methyl group, an ethyl group, a n-propyl group, an isopropyl group, a butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group or a hexyl group may, for example, be mentioned.

"An alkenyl group" may be linear or branched. Unless

otherwise specified, the alkenyl group is preferably a C₂₋₆ alkenyl group. As an example of the alkenyl group, an allyl group, an isopropenyl group or a 3-butenyl group may, for example, be mentioned.

5 "An aryl group" means a monovalent aromatic hydrocarbon group, and a phenyl group is preferred.

"An aralkyl group" means an alkyl group substituted by an aryl group. As the aryl group moiety, a phenyl group is preferred. Further, as the alkyl group moiety
10 in the aralkyl group, a C₁₋₄ alkyl group is preferred. As an example of the aralkyl group, a benzyl group, a benzhydryl group, a trityl group or a phenylethyl group may, for example, be mentioned.

"A cycloalkyl group" means an at least 3-membered
15 cyclic alkyl group, and a 3- to 8-membered cycloalkyl group is preferred. As an example of the cycloalkyl group, a cyclopentyl group, a cyclohexyl group or a cycloheptyl group may, for example, be mentioned.

In the compound (1) or its salt, Z¹ is a group
20 represented by -CO₂R¹, -CONR²R³ or -CN, preferably -CO₂R¹ or -CN from the viewpoint of availability, particularly preferably -CO₂R¹ from the viewpoint of the yield and the efficiency in the post treatment after the reaction.
Here, each of R¹, R² and R³ which are independent of one
25 another, is a hydrogen atom, an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group. R¹ is preferably a hydrogen atom or a lower alkyl

group, more preferably a hydrogen atom, a methyl group or an ethyl group, particularly preferably a hydrogen atom. Each of R^2 and R^3 which are independent of each other, is preferably a hydrogen atom, a methyl group or an ethyl group. It is particularly preferred that both R^2 and R^3 are hydrogen atoms, methyl groups or ethyl groups.

The compound (1) as the starting material in the present invention is commercially available or can easily be obtained by a usual preparation method. For example, it can be produced by the method disclosed in e.g. European Patent 333020 or U.S. Patent 5,204,478. As the compound (1), one having a purity usually available may be used as it is without carrying out purification or the like. A salt of the compound (1) is commercially available or can easily be obtained by a usual preparation method. As such a salt, one having a purity usually available may be used as it is without carrying out purification or the like.

The following compounds may be mentioned as specific examples of the compound (1).

2,6-Dichloro-5-fluoronicotinic acid, methyl 2,6-dichloro-5-fluoronicotinate, ethyl 2,6-dichloro-5-fluoronicotinate, isopropyl 2,6-dichloro-5-fluoronicotinate, butyl 2,6-dichloro-5-fluoronicotinate, 2,6-dichloro-5-fluoronicotinonitrile, 2,6-dichloro-5-fluoronicotinic acid amide, 2,6-dichloro-5-fluoronicotinic acid dimethylamide, 2,6-dichloro-5-

fluoronicotinic acid diethylamide, and 2,6-dichloro-5-fluoro-3-cyanopyridine. As the compound (1), 2,6-dichloro-5-fluoronicotinic acid is preferred since it is readily available and easy to handle.

5 The salt of the compound (1) may, for example, be a hydrochloride, a nitrate, a sulfate, an acetate, a formate, a trifluoroacetate or a phosphate, of the above compound.

10 In the present invention, the chlorine atom at the 6-position of the compound (1) or its salt is selectively reduced. For such a reduction reaction, conditions for a common reduction reaction may be employed. For example, the method for reducing the carbon-halogen bond as disclosed in "Shin Jikken Kagaku Kouza (New Experimental
15 Chemistry) 14-Yukikagoubutsu no Gousei to Hannou (Syntheses and reactions of organic compounds) (I), published by Maruzen" may, for example, be employed. The reduction reaction of the present invention can be carried out under usual reduction reaction conditions,
20 but only the 6-position is selectively reduced among chlorine atoms at the 2- and 6-positions. As a specific example for such a reduction reaction, any one of the following methods 1, 2 and 3 is preferred.

Method 1: A method wherein catalytic hydrogenolysis
25 is carried out by using hydrogen in the presence of a metal reduction catalyst such as 5% Pd/calcium carbonate (one poisoned with lead, so-called Lindlar catalyst), Pd-

barium sulfate, Pd-C, or Raney nickel.

In this method 1, the amount of the metal reduction catalyst is preferably from 0.1 to 100 mass%, particularly preferably from 1 to 30 mass%, based on the
5 compound (1) or its salt. Further, the amount of hydrogen is preferably at least 1 time by mol, more preferably from 1 to 100 times by mol, particularly preferably from 1 to 10 times by mol, relative to the compound (1).

10 Method 2: A method wherein reduction is carried out by letting a protic solvent act in the presence of an alkali metal such as sodium or lithium; a metal such as magnesium, aluminum, tin, copper, zinc, iron or nickel, or an alloy thereof; or a metal salt such as lithium
15 iodide or tin(II) chloride.

In the method 2, it is preferred to employ a metal, particularly a metal in a powder form. Further, as the metal, tin, copper, zinc or iron is, for example, preferred, and zinc is particularly preferred.

20 In the method 2, the amount of the metal or metal salt is preferably from 0.1 to 1,000 mass%, particularly preferably from 0.2 to 300 mass%, based on the compound (1) or its salt. Further, the amount of the protic solvent is preferably an amount whereby the compound (1)
25 or its salt can be dissolved, dispersed or suspended, and it is preferably from 0.5 to 100 times by mass, particularly preferably from 1 to 10 times by mass,

relative to the compound (1) or its salt.

The protic solvent in the method 2 is a compound which will be a hydrogen source in the reduction reaction and is a compound capable of being a solvent in the
5 reaction of the present invention. As such a protic solvent, various solvents may be employed which are useful for reduction reactions. For example, water, an acid (hereinafter an acid as a protic solvent will be referred to as an acid (A)), an alkaline aqueous
10 solution, an alcohol solvent, an ester solvent, a hydrocarbon solvent, an ether solvent, or a solvent mixture thereof, may, for example, be employed. As a specific example of such a solvent, an acid (A) such as acetic acid, formic acid, propionic acid or hydrochloric
15 acid; an alcohol solvent such as methanol, ethanol, 2-propanol or t-butanol; or an aqueous alkaline solution such as an aqueous sodium hydroxide solution or an aqueous potassium hydroxide solution, may be mentioned. The protic solvent is preferably an acid selected from
20 acetic acid, formic acid and hydrochloric acid, or an aqueous solution of such an acid; an alcohol solvent such as methanol, ethanol, 2-propanol or t-butanol; a solvent mixture having an acid (A) such as acetic acid, formic acid, propionic acid or hydrochloric acid, added to such
25 an alcohol solvent, particularly preferably methanol, or a solvent mixture of methanol with acetic acid.

In a case where a solvent mixture having two or more

solvents mixed, is used as the protic solvent, such solvents may preliminarily be mixed, or the respective solvents may separately be added to the reaction system and mixed in the reaction system to form a solvent

5 mixture. In a case where the solvent mixture is a solvent mixture of an alcohol solvent with an acid (A), the acid (A) is preferably added lastly to the reaction system to form a solvent mixture, whereby the operation efficiency will be good.

10 The amount of the protic solvent is preferably from 0.5 to 50 times by mass, particularly preferably from 2 to 10 times by mass, relative to the compound (1) or its salt. In a case where a solvent mixture of an alcohol solvent with an acid (A) is used as the solvent mixture, 15 the amount of the acid (A) is preferably from 0.1 to 10 times by mol, more preferably from 0.5 to 2 times by mol, particularly preferably from 1.0 to 1.3 times by mol, relative to the compound (1) or its salt.

The method 2 is preferably carried out by either one 20 of the following operations methods.

Method 2-1: A method wherein the compound (1) or its salt is dissolved, dispersed or suspended in a protic solvent, and then, a metal or a metal salt is added and reacted. In this method, the metal or the metal salt may 25 be added all at once at the initial stage of the reaction or may be added dividedly in a plurality of times.

Method 2-2: A method wherein a metal or a metal salt

is dissolved, dispersed or suspended in a protic solvent, and then, the compound (1) or its salt is dropwise added and reacted. In this method, it is preferred that the compound (1) or its salt is dropwise added in the form of
5 a solution or suspension diluted with a protic solvent, whereby the reaction heat can be suppressed.

The method 2 is preferably carried out by the method 2-2, whereby the operation is good, and control of the reaction will be easy.

10 Method 3: A method for a reaction with a hydride reactant, such as a metal hydride such as a trialkyltin hydride or a trialkylsilane, or a metal/hydrogen complex compound such as sodium borohydride or lithium aluminum hydride.

15 In the method 3, the amount of the hydride reactant is preferably from 0.6 to 10 times by mol, particularly preferably from 0.8 to 5 times by mol, relative to the compound (1) or its salt.


The reduction method of the present invention is
20 carried out preferably by the method 2 among them. According to the reduction reaction of the present invention, the chlorine atom at the 6-position of the compound (1) or its salt, is selectively reduced to form the compound (2) or its salt. However, depending upon
25 the structure of the compound (1) or its salt as the starting material, and reaction conditions, the following compound (3) having both chlorine atoms at the 2- and 6-

positions reduced, may sometimes be formed as a by-product (provided that Z^1 in the formula (2) and the formula (3) is the same group as Z^1 in the formula (1)).

However, in a case where the reduction reaction is

5 carried out by the method 2, it is possible to suppress the production of the compound (3) as a by-product.


Among the method 2, particularly preferred is the method 2 wherein a metal is employed.



10 The reaction temperature for the reduction reaction of the compound (1) or its salt is preferably -20 to +100°C, particularly preferably from -5 to +50°C. The reaction time is not particularly limited. The termination of the reaction may suitably be changed,
15 usually, by analyzing the progress of the reaction by means of high performance liquid chromatography (hereinafter referred to as HPLC) or the like, and the reaction time is preferably from 0.5 to 72 hours, particularly preferably from about 1 to 25 hours.

20 The crude reaction product obtained as a result of the reduction reaction may be subjected to purification treatment, as the case requires. As a method for such purification treatment, filtration, evaporation of solvent, extraction, washing, high performance liquid
25 chromatography, recrystallization and distillation may, for example, be mentioned.

In the method of the present invention, by utilizing the reactivity of the Z^1 group at the 3-position of the compound (2) or its salt obtained by the above reduction reaction, the Z^1 group may be converted to another group
 5 (Z^2 group) thereby to produce the following compound (4) or its salt.




In the formula (4), Z^2 is a group different from Z^1 and is a group represented by $-CO_2R^5$, $-CONR^6R^7$ or $-CN$.
 10 From the usefulness of the compound (4) or its salt, Z^2 is preferably $-CO_2R^5$. Here, R^5 is an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group, and each of R^6 and R^7 which are independent of each other, is a hydrogen atom, an alkyl
 15 group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group. R^5 is preferably a lower alkyl group, particularly preferably an ethyl group. Each of R^6 and R^7 which are independent of each other, is preferably a hydrogen atom, a methyl group or an ethyl
 20 group, and it is particularly preferred that both R^6 and R^7 are hydrogen atoms, methyl groups or ethyl groups.

As the method for converting the compound (2) or its salt to the compound (4) or its salt, a known or well known method may be employed and may suitably be selected

for use depending upon the types of Z^1 and Z^2 .


For example, the following compound (4a) i.e. the compound (4) wherein Z^1 is $-\text{COOR}^5$, or its salt, can be produced by an esterification reaction of the following compound (2a-1) i.e. the compound (2) wherein Z^1 is $-\text{COOH}$, or its salt, or by an ester exchange reaction of the following compound (2a-2) i.e. the compound (2) wherein Z^1 is $-\text{COOR}^{10}$ (wherein R^{10} is an alkyl group), or its salt.



The esterification reaction of the compound (2a-1) or its salt can be carried out by the following method 4-1 or 4-2, preferably the method 4-1, whereby the reactivity is good.

15 Method 4-1: A method wherein the compound (2a-1) or its salt is reacted with a chlorination agent to obtain the following compound (5) or its salt, and the compound (5) or its salt, and a compound represented by the formula $R^5\text{OH}$ (wherein R^5 is as defined above) are
20 subjected to an esterification reaction.

Method 4-2: A method wherein the compound (2a-1) or its salt, and a compound represented by the formula $R^5\text{OH}$ (wherein R^5 is as defined above) are subjected to an esterification reaction.



(5)

The chlorination agent in the method 4-1 is preferably thionyl chloride or oxalyl chloride, particularly preferably thionyl chloride. The amount of the chlorination agent is preferably from 1 to 10 times by mol relative to the compound (2a-1) or its salt, particularly preferably from 1 to 2 times by mol, since the post treatment is thereby simple.

The chlorination reaction may be carried out in the presence or absence of the solvent, preferably in the presence of a solvent. The solvent may suitably be selected from solvents inert to the chlorination reaction, and an aromatic hydrocarbon solvent such as toluene or xylene, or a halogenated hydrocarbon solvent such as methylene chloride or HFC-225, may, for example, be mentioned. Among these solvents, an aromatic hydrocarbon solvent such as xylene or toluene is preferred.

Further, in order to let the chlorination reaction proceed smoothly, it is preferred to add N,N-dimethylformamide. The amount of the N,N-dimethylformamide is preferably from 0.001 to 1 time by mol, particularly preferably from 0.001 to 0.5 time by mol, relative to the compound (2a-1) or its salt.

The temperature for the chlorination reaction is

preferably from +20 to +100°C, particularly preferably from +50 to +90°C.

The time for the chlorination reaction may suitably be changed depending upon the progress of the reaction, and it is preferably from 1 to 24 hours, particularly preferably from 5 to 15 hours, in an industrial production.

In the esterification reaction of the compound (5) or its salt, and the compound represented by the formula R^5OH (wherein R^5 is as defined above), the amount of the compound represented by the formula R^5OH is preferably from 1 to 100 times by mol, more preferably from 1 to 20 times by mol, particularly preferably from 1 to 10 times by mol, relative to the compound (5) or its salt.

The esterification reaction may be carried out in the presence or absence of a solvent, preferably in the presence of a solvent. The solvent may be suitably selected from solvents inert to the esterification reaction, and an aromatic hydrocarbon solvent such as toluene or xylene; an aliphatic hydrocarbon solvent such as hexane or heptane; a halogenated hydrocarbon solvent such as HCFC-225 or methylene chloride; or an ether solvent such as diisopropyl ether or tert-butyl methyl ether, may, for example, be mentioned, and an aromatic hydrocarbon solvent such as toluene or xylene is preferred.

The temperature for the esterification reaction is

preferably from +50 to +120°C, particularly preferably from +70 to +90°C.

For the esterification reaction, the compound (5) or its salt isolated from the crude reaction solution
5 obtained by the above chlorination reaction may be employed, or the crude reaction solution may be employed as it is. In the latter case, the reaction may be carried out by using the same reactor as the reactor used for the chlorination reaction. As an example of the
10 latter method, a method may be mentioned wherein the compound (2a-1) or its salt is dissolved or suspended in a solvent, and a chlorination agent is added, followed by heating to obtain the compound (5) or its salt, and then the compound represented by the formula R^5OH is dropwise
15 added thereto.

The reaction time for the esterification reaction may be suitably changed depending upon the progress of the reaction, and in an industrial production, the reaction time is preferably from 3 to 20 hours,
20 particularly preferably from 4 to 15 hours.

The esterification reaction in the method 4-2 may be carried out in accordance with the esterification reaction in the method 4-1, and in the method 4-2, it is preferred to use an acid catalyst. The acid catalyst is
25 preferably a Lewis acid or a protic acid, and from the viewpoint of the economical efficiency, a protic acid is preferred. The protic acid may, for example, be

concentrated sulfuric acid, hydrochloric acid, p-toluene sulfonic acid or trifluoromethanesulfonic acid, and concentrated sulfuric acid is preferred. The amount of the acid catalyst is preferably from 0.001 to 1 time by 5 mol, particularly preferably from 0.01 to 0.6 time by mol, relative to the compound (5) or its salt.

In order to let the esterification reaction proceed smoothly, it is preferably carried out while water formed as a by-product is distilled off from the reaction 10 system.

The time for the esterification reaction may suitably be changed depending upon the progress of the reaction, and in an industrial production, it is preferably from 1 to 24 hours, particularly preferably 15 from 5 to 15 hours.

The ester exchange reaction of the compound (2a-2) or its salt can be carried out by reacting the compound (2a-2) or its salt with the compound represented by the formula R^5OH (wherein R^5 is as defined above), and can be 20 carried out in the same manner as for the esterification reaction disclosed in the method 4-2. In the ester exchange reaction, in order to let the reaction proceed smoothly, it is preferably carried out while the compound represented by the formula R^1OH , formed as a by-product, 25 is distilled off from the reaction system.

The reaction time for the ester exchange reaction may suitably be changed depending upon the progress of

the reaction, and in an industrial production, it is preferably from 1 to 24 hours, particularly preferably from 5 to 15 hours.

In the reduction reaction of the present invention,
5 when the compound (1) is used as the starting material, the product may be the compound (2) or a salt of the compound (2), and when a salt of the compound (1) is used as the starting material, the product may be the compound (2) or a salt of the compound (2). Further, depending
10 upon the reaction conditions for the reduction reaction, the product may be the compound (2) only, a salt of the compound (2) only, or a mixture of the compound (2) and a salt of the compound (2).

The same applies to the substitution reaction of the
15 present invention. Namely, when the compound (2) is used as the starting material, the product may be the compound (4) or a salt of the compound (4), and when a salt of the compound (2) is used as the starting material, the product may be the compound (4) or a salt of the compound
20 (4). Further, depending upon the reaction conditions of the substitution reaction, the product may be the compound (4) only, a salt of the compound (4) only, or a mixture of the compound (4) and a salt of the compound (4). The compounds (1) to (4) and salts corresponding
25 thereto, are equivalent compounds showing similar reactivities in the respective reactions in the methods of the present invention.

In a case where the product in the method of the present invention is a compound forming no salt (hereinafter referred to also as a free compound) or its salt, or a mixture thereof, it can be converted to the
5 respective corresponding salt or free compound by a conventional method. For example, as a method for converting a free compound to its salt, a method may be mentioned wherein an acid (hereinafter referred to as an acid (B)) is reacted to the compound (2) or (4) to
10 convert it to a salt of the corresponding compound (2) or (4) (hereinafter referred to also as an acid treatment step). As the acid (B), hydrochloric acid, sulfuric acid, nitric acid, formic acid, acetic acid, trifluoroacetic acid or citric acid may, for example, be
15 mentioned, and hydrochloric acid is preferred. The amount of the acid (B) is preferably from 1 to 10 times by mol, particularly preferably from 1 to 2 times by mol, relative to the compound (2).

The step of reacting the compound (2) with the acid
20 (B) is preferably carried out in the presence of a solvent, whereby stirring and control of heat generation may be facilitated. Such a solvent is a solvent not involved in the reaction and is preferably water or a water-soluble organic solvent. The water-soluble organic
25 solvent may, for example, be an alcohol solvent such as methanol, ethanol or isopropyl alcohol; or a nitrile solvent such as acetonitrile or propionitrile. These

solvents may be used alone or in combination as a solvent mixture of two or more of them. As the solvent for the acid treatment step, water, an alcohol solvent or a solvent mixture of water and an alcohol solvent, is preferred, and water, methanol or a solvent mixture of water and methanol, is particularly preferred.

The acid treatment step is preferably carried out by dropwise adding the acid (B) to a solution having the compound (2) dissolved in the solvent. If the compound (2) is precipitated at the time of the dropwise addition of the acid (B), stirring may become difficult, or the purity of the desired compound (2) is likely to be low.

The compound (2) to be used for the acid treatment step may be the compound (2) isolated from the crude reaction solution obtained by the above reduction reaction, or the compound (2) contained in such a crude reaction solution. It is preferably the latter, since the operation can be thereby simplified. In the latter, in a case where the solvent to be used for the acid treatment step is different from the protic solvent used in the reduction step, it is preferred that with respect to the crude reaction solution after completion of the reduction step, the solvent substitution operation is carried out, or the crude reaction solution is concentrated to such a degree that it will not be solidified, and the solvent for the acid treatment step is added, and then, the acid is dropwise added. Whereas,

in a case where the solvent to be used for the acid treatment step is the same as the protic solvent used in the reduction step, the acid (B) may be dropwise added after concentrating the crude reaction solution to such a
5 degree that it will not be solidified, or the acid (B) may be dropwise added to the crude reaction solution from the reduction step.

In the acid treatment step, the temperature at the time of the dropwise addition of the acid (B) is
10 preferably from +30 to +100°C, particularly preferably from +40 to +80°C.

The salt of the compound (2) obtained in the acid treatment step is preferably purified, as the case requires. The purification method may suitably be
15 selected from methods such as distillation, sublimation and crystallization, depending upon the physical properties of the salt of the compound (2).

For example, as a method for producing a salt of the compound (2a-1) from the compound (2a-1), the compound
20 (2a-1) is dissolved in a solvent mixture of water and methanol under heating at a level of from +50 to +60°C, and hydrochloric acid is dropwise added to the solution. Then, the content is cooled to a temperature of from +25 to +35°C, whereby a salt of the compound (2a-1) will
25 precipitate. Further, the solution is maintained at that temperature for from 1 to 24 hours to sufficiently precipitate the salt. Then, the precipitate is separated

from the mother liquor by a method such as filtration or centrifugal separation, then washed and dried to isolate the salt of the compound (2a-1).


The step of reacting the compound (4) with the acid
5 (B) can be carried out in the same manner as the step of reacting the compound (2) with the acid (B). For example, the method of converting the compound (2a-1) to a salt of the compound (2a-1) can be employed also as a method of converting the compound (4) to a salt of the
10 compound (4).

As a method for converting a salt of the compound to a free compound, a method may be mentioned wherein a salt of the compound (2) or (4) is reacted with a basic compound to convert it to the compound (2) or (4),
15 respectively. The basic compound may, for example, be an inorganic base, such as an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide, or an alkali metal carbonate such as sodium carbonate or potassium carbonate, or an organic base such
20 as triethylamine or pyridine. From the viewpoint of the handling efficiency and the economical efficiency, an inorganic base is preferred, and an alkali metal carbonate such as sodium carbonate or potassium carbonate, is particularly preferred. Such an inorganic
25 base is preferably used in the form of an aqueous solution.

The compound (2) or its salt, and the compound (4)

or its salt, obtained by the methods of the present invention, are compounds useful as intermediates for pharmaceuticals, agricultural chemicals, etc. or as intermediates for production of various functional materials.

As a preferred embodiment of the present invention, a method for producing the following compound (4a-1) may be mentioned. Namely, it is a method wherein a compound (1a-1) is reduced by reacting it with a protic solvent in the presence of zinc to obtain a compound (2a-1), then the compound (2a-1) is reacted with hydrochloric acid to obtain a salt of the compound (2a-1), and then the salt of the compound (2-1) is reacted with thionyl chloride and then with ethanol to obtain the compound (4a-1).



EXAMPLES

Now, the present invention will be described in further detail with reference to Examples, but the present invention is by no means restricted to these Examples. In the following, "%" in the analytical results by HPLC represents area% of each peak in the chromatogram. As the detector for HPLC, an ultraviolet absorption detector was employed. The detection

wavelength was 254 nm in Examples 1 to 4 and Reference Example. Further, in Examples 5 and 6, the detection wavelength was 276 nm, and the analytical values were corrected based on the differences in the absorption coefficient depending upon the compounds.

EXAMPLE 1: Preparation of 2-chloro-5-fluoronicotinic acid (First Example)

Into a 50 mL round bottomed flask, 2,6-dichloro-5-fluoronicotinic acid (1 g), acetic acid (5 mL) and water (0.5 mL) were charged, and under cooling with ice, zinc powder (200 mg) was added, followed by stirring at room temperature for 3 hours. Then, zinc powder (200 mg) was added, followed by stirring for one hour, and then zinc powder (400 mg) was further added, followed by stirring for 3 hours. Further, zinc powder (200 mg) was added, followed by stirring for one hour and then by filtration with celite and washing with ethyl acetate and ethanol. The solvent was distilled off under reduced pressure, and ethyl acetate (20 mL) and a saturated sodium bicarbonate aqueous solution (10 mL) were added for liquid separation. The aqueous layer was extracted three times with ethyl acetate (20 mL). The extract was dried over magnesium sulfate and then concentrated under reduced pressure to obtain a reddish brown oil (400 mg). As a result of the HPLC analysis, the formed reddish brown oil was found to be a mixture comprising 71% of the title compound, 27% of the starting material and 2% of 5-

fluoronicotinic acid. Yield: 48%.

¹HNMR (CD₃OD): δ (ppm) 7.86 (dd, J=7.8, 3.0 Hz, 1H),
8.29 (d, J=2.7 Hz, 1H)

EXAMPLE 2: Preparation of 2-chloro-5-fluoronicotinic

5 acid (Second Example)

Into a 200 mL round bottomed flask, 2,6-dichloro-5-fluoronicotinic acid (1 g), methanol (10 mL), triethylamine (0.96 g) and 5% Pd/calcium carbonate (one poisoned by lead) (107 mg) were charged under cooling
10 with ice and stirred at room temperature for one hour in a hydrogen atmosphere. The catalyst was filtered off, and the filtrate was concentrated to obtain a crude product (1.3 g). The concentrated filtrate was analyzed by HPLC and found to be a mixture comprising 26% of the
15 title compound, 57% of 2,6-dichloro-5-fluoronicotinic acid and 17% of 5-fluoronicotinic acid.

EXAMPLE 3: Preparation of ethyl 2-chloro-5-fluoronicotinate (First Example)

Into a 500 mL four-necked flask, ethyl 2,6-dichloro-5-fluoronicotinate (20 g) was put and dissolved by adding
20 acetic acid (190 mL) and water (10 mL). Under cooling with ice, zinc powder (4 g) was added, followed by stirring at room temperature for 0.5 hour. Then, zinc powder (4 g) was added, followed by stirring for 0.5
25 hour, and zinc powder (4 g) was further added, followed by stirring for 0.5 hour. The solvent was distilled off under reduced pressure, and a 5% sodium bicarbonate

aqueous solution (100 mL) and ethyl acetate (100 mL) were added for liquid separation. The aqueous layer was extracted twice with ethyl acetate (100 mL). The organic layer was washed with a 5% sodium bicarbonate aqueous solution (100 mL) and washed with a saturated sodium chloride aqueous solution (100 mL). The solvent was distilled off to obtain a reddish brown oil (15.6 g). It was purified by distillation under reduced pressure to obtain the title compound (10.6 g, yield: 62%). As a result of the HPLC analysis, the purity of the title compound was 99%.

¹H NMR (CDCl₃): δ (ppm) 1.43 (t, J=7.2 Hz, 3H), 4.43 (q, J=7.2 Hz, 2H), 7.91 (dd, J=7.8, 2.7 Hz, 1H), 8.39 (d, J=3.0 Hz, 1H)

¹³C NMR (CDCl₃): δ (ppm) -129.2 (d, J=7.6 Hz)

EXAMPLE 4: Preparation of 2-chloro-5-fluoronicotinonitrile

Into a 50 mL round bottomed flask, 2,6-dichloro-5-fluoronicotinonitrile (1 g), acetic acid (5 mL) and water (0.5 mL) were charged, and reacted and post-treated in the same manner as in Example 1 and then purified by silica gel column chromatography (hexane/ethyl acetate=6/1) to obtain the title compound (420 mg, yield: 52%). As a result of the HPLC analysis, the purity was 100%.

¹H NMR (CDCl₃): δ (ppm) 7.76 (dd, J=7.2, 3.0 Hz, 1H), 8.49 (d, J=3.3 Hz, 1H)

¹⁹F NMR (CDCl₃): δ (ppm) -126.9 (d, J=7.6Hz)

EXAMPLE 5: Preparation of ethyl 2-chloro-5-fluoronicotinate (Second Example)

EXAMPLE 5-1: Preparation of a salt of 2-chloro-5-

5 fluoronicotinic acid

Into a 2 L round bottomed flask, zinc powder (847 g) and methanol (4,080 mL) were charged and dispersed with stirring. A methanol (2,720 mL) solution of 2,6-dichloro-5-fluoronicotinic acid (1,700 g) was slowly
10 added dropwise thereto, and then, acetic acid (535 g) was further added dropwise thereto. The internal temperature was raised to 35°C, followed by stirring at that temperature for 5 hours. After confirming that the starting material became at most 3% by the HPLC analysis,
15 the residue of zinc powder was removed by filtration. Methanol was distilled off under reduced pressure, and then, water (3,000 mL) was added, whereby crystals precipitated. Then, the crude reaction product containing the crystals was transferred to a 20 L reactor
20 by means of water (3,750 mL). The internal temperature was heated to at least 50°C, and then, 6 mol/L hydrochloric acid (1,960 g) was dropwise added thereto. After stirring for 3 hours as it was, the mixture was gradually cooled to room temperature, and a precipitated
25 solid was collected by filtration and washed with water. The obtained crystals were dried under reduced pressure (60°C, 30.7 Pa, 12 hours) to obtain a salt of 2-chloro-5-

fluoronicotinic acid (1,000 g).

NMR spectrum of the salt of 2-dichloro-5-fluoronicotinic acid:

¹H NMR (CD₃COCD₃): δ (ppm) 8.05 (dd, J=2.9, 8.2Hz, 1H), 8.47 (d, J=2.9Hz, 1H)

¹⁹F NMR (CDCl₃): δ (ppm) -130.4 (d, J=9.2Hz)

EXAMPLE 5-2: Esterification reaction of the salt of 2-chloro-5-fluoronicotinic acid

Into a 5 L four-necked flask, the salt (1,000 g) of 2-chloro-5-fluoronicotinic acid obtained in Example 5-1 was put and dissolved by adding toluene (3,000 mL), thionyl chloride (617 g) and N,N-dimethylformamide (0.95 g). The flask was heated by dipping it in an oil bath set at 80°C. Six hours later, after confirming by the HPLC analysis that 2-chloro-5-fluoronicotinic acid became at most 5%, ethanol (2,173 g) was added. The temperature of the oil bath was raised to 95°C, and the reaction was carried out for 14 hours. Then, concentration under reduced pressure was carried out, and to the obtained crude liquid, ethyl acetate (4,000 mL) and a 7.5% sodium carbonate aqueous solution (4,000 mL) were added, followed by stirring. After confirming that the pH of the aqueous layer was at least 8, filtration and separation were carried out, and an organic layer was recovered. An aqueous layer was extracted with ethyl acetate (2,000 mL), and the extract was put together with the previous organic layer, followed by washing with

water (2,000 mL) and then with a 5% sodium chloride aqueous solution (2,000 mL). The organic solvent was distilled off under reduced pressure to obtain a reddish brown oil (1,234.7 g). It was purified by distillation under reduced pressure to obtain ethyl 2-chloro-5-fluoronicotinate (624.8 g). The yield from 2,6-dichloro-5-fluoronicotinic acid was 66%. As a result of the HPLC analysis, the purity of ethyl 2-chloro-5-fluoronicotinate was 99%.

10 EXAMPLE 6: Preparation of a salt of 2-chloro-5-fluoronicotinic acid (Second Example)

The reaction and post-treatment were carried out in the same manner as in Example 5-1 except that the internal temperature at the time of dropwise addition of 6 mol/L hydrochloric acid was changed to at least 60°C, and after completion of the dropwise addition, the internal temperature was further heated to 80°C, and stirring was continued at that temperature for 30 minutes, whereby a salt of 2-chloro-5-fluoronicotinic acid (1,000 g) was obtained.

20 REFERENCE EXAMPLE: Preparation of a 5-fluoronicotinic acid

Into a 200 mL eggplant type flask, 2,6-dichloro-5-fluoronicotinic acid (1 g), methanol (10 mL), triethylamine (0.96 g) and 5% Pd-C (107 mg) were charged under cooling with ice and stirred at room temperature for 4 hours in a hydrogen atmosphere. After completion

of the stirring, Pd-C was filtered off, and the filtrate was concentrated. The obtained crude product was analyzed by HPLC and as a result, found to be a mixture containing 98.6% of the above-identified compound and
5 1.4% of 2-chloro-5-fluoronicotinic acid.

INDUSTRIAL APPLICABILITY

The present invention provides a method for selectively producing a 3-substituted 2-chloro-5-fluoropyridine or its salt by a single step reduction reaction
10 from a readily available 3-substituted 2,6-dichloro-5-fluoropyridine or its salt. The reduction reaction can be carried out without using a special reagent and without employing a special reaction condition, and the
15 yield is high. Thus, the method is very advantageous for industrial operation. Further, the 3-substituted 2-chloro-5-fluoropyridine or its salt obtained by the above reduction reaction may be subjected to conversion of the substituent by utilizing the reactivity of the
20 substituent at the 3-position. Thus, the present invention may provide efficient methods for producing various 3-substituted 2-chloro-5-fluoropyridines or their salts.

2004236085 23 Feb 2010

33a

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a
5 stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any
10 matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.


15

2004236085 23 Feb 2010


34

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for producing a compound represented by the following formula (2) or its salt, which comprises
 5 selectively reducing a chlorine atom at the 6-position of a compound represented by the following formula (1) or its salt, provided that in the following formulae, Z^1 is a group represented by $-\text{CO}_2\text{R}^1$, $-\text{CONR}^2\text{R}^3$ or $-\text{CN}$, wherein each of R^1 , R^2 and R^3 which are independent of one another, is a hydrogen
 10 atom, an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group:




(1)




(2)

2. A method for producing a compound represented by the
 15 following formula (2), which comprises selectively reducing a chlorine atom at the 6-position of a compound represented by the following formula (1), provided that in the following formulae, Z^1 is a group represented by CO_2R^1 , $-\text{CONR}^2\text{R}^3$ or $-\text{CN}$, wherein each of R^1 , R^2 and R^3 which are independent of one
 20 another, is a hydrogen atom, an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group:




(1)



(2)

3. The method according to Claim 1 or 2, wherein Z^1 is $-\text{CO}_2\text{R}^1$ (wherein R^1 is as defined above).
4. The method according to any one of Claims 1 to 3, wherein the reduction is carried out by letting a protic solvent act in the presence of a metal or a metal salt.
5. The method according to any one of Claims 1 to 4, wherein the reduction is carried out by letting a protic solvent act in the presence of zinc.
6. A method for producing a compound represented by the following formula (4) or its salt, which comprises carrying out a substitution reaction of a compound represented by the formula (2) or its salt obtained by the method as defined in any one of Claims 1 to 5, to substitute the Z^1 group at the 3-position by a Z^2 group, thereby to obtain a compound represented by the following formula (4) or its salt, provided that Z^2 in the following formula is a group different from Z^1 and is a group represented by $-\text{CO}_2\text{R}^5$, $-\text{CONR}^6\text{R}^7$ or $-\text{CN}$, wherein R^5 is an alkyl group, an alkenyl group, an aryl group, an aralkyl group, or a cycloalkyl group, and each of R^6 and R^7 which are independent of each other, is a hydrogen atom, an alkyl group, an alkenyl group, an aryl group, an aralkyl group or a cycloalkyl group:



(4)

2004236085 23 Feb 2010

36

7. The method according to Claim 6, wherein from the compound of the formula (2) or its salt wherein the Z^1 group is a group represented by $-\text{COOH}$ or $-\text{COOR}^{10}$ (wherein R^{10} is an alkyl group), the compound (4) wherein Z^2 is a $-\text{COOR}^5$ group (wherein R^5 is as defined above), is obtained by an esterification reaction in a case where Z^1 is a $-\text{COOH}$ group, or by an ester exchange reaction in a case where Z^1 is a $-\text{COOR}^{10}$ group (wherein R^{10} is as defined above).

10

8. The method according to Claim 3, wherein R^1 is a hydrogen atom.

9. The method according to Claim 1 or 2, wherein Z^1 is $-\text{CONR}^2\text{R}^3$ wherein R^2 and R^3 , independently of each other, are selected from the group consisting of a hydrogen atom, a methyl group and an ethyl group.

10. The method according to Claim 1 or 2, wherein the compound of formula (I) is selected from the group consisting of 2,6-dichloro-5-fluoronicotinic acid, methyl 2,6-dichloro-5-fluoronicotinate, ethyl 2,6-dichloro-5-fluoronicotinate, isopropyl 2,6-dichloro-5-fluoronicotinate, butyl 2,6-dichloro-5-fluoronicotinate, 2,6-dichloro-5-fluoronicotinonitrile, 2,6-dichloro-5-fluoronicotinic acid amide, 2,6-dichloro-5-fluoronicotinic acid dimethylamide, 2,6-dichloro-5-fluoronicotinic acid diethylamide, and 2,6-dichloro-5-fluoro-3-cyanopyridine.

11. The method according to Claim 1 or 2, wherein the compound of formula (I) is 2,6-dichloro-5-fluoronicotinic acid.

2004236085 23 Feb 2010

37

12. The method according to Claim 1 or 2, wherein reducing includes carrying out a catalytic hydrogenolysis with hydrogen in the presence of a metal reduction catalyst.
- 5 13. The method according to Claim 1 or 2, wherein the reducing is carried out in the presence of a protic solvent and at least one of sodium and lithium.
- 10 14. The method according to Claim 1 or 2, wherein the reducing is carried out by reacting the compound of formula (I) with a hydride reactant.
- 15 15. The method according to Claim 14, wherein the hydride reactant is at least one selected from the group consisting of a trialkyl tin hydride, a trialkyl silane, sodium borohydride, and lithium aluminum hydride.
- 20 16. The method according to Claim 1 or 2, wherein the reducing is carried out by directly replacing the chlorine atom with a hydrogen atom.
- 25 17. The method according to Claim 1 or 2, wherein the reducing is carried out in one step.
18. The method according to any one of claims 1, 2 or 6 substantially as hereinbefore described with reference to any one of the Examples.
- 30 19. A compound produced by the method of any one of claims 1-17.