



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/09/22	(51) Cl.Int./Int.Cl. <i>C12N 5/0783</i> (2010.01), <i>A61K 35/17</i> (2015.01), <i>C07K 14/195</i> (2006.01), <i>C07K 14/71</i> (2006.01), <i>C07K 14/725</i> (2006.01), <i>C12N 9/22</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2018/03/29	(71) Demandeur/Applicant: FRED HUTCHINSON CANCER RESEARCH CENTER, US
(85) Entrée phase nationale/National Entry: 2019/02/25	(72) Inventeurs/Inventors: BLEAKLEY, MARIE, US; DOSSA, ROBSON, US; SOMMERMEYER, DANIEL, DE
(86) N° demande PCT/PCT Application No.: US 2017/053112	(74) Agent: GOWLING WLG (CANADA) LLP
(87) N° publication PCT/PCT Publication No.: 2018/058002	
(30) Priorité/Priority: 2016/09/23 (US62/399,291)	

(54) Titre : TCR SPECIFIQUES DE L'ANTIGENE HA-1 D'HISTOCOMPATIBILITE (H) MINEURE ET SES UTILISATIONS
(54) Title: TCRS SPECIFIC FOR MINOR HISTOCOMPATIBILITY (H) ANTIGEN HA-1 AND USES THEREOF




FIG. 30

(57) **Abrégé/Abstract:**
The present disclosure provides compositions and methods for targeting a minor histocompatibility (H) antigen (HA-1H?) to, for example, prevent or manage relapse of a hematological malignancy after allogeneic hematopoietic stem cell transplantation (HCT). Also provided are transgene constructs encoding engineered binding proteins, such as a T cell receptor or a chimeric antigen receptor, optionally encoding additional components such as a co-receptor and/or safety switch. Such transgene constructs can be transduced into an immune cell, such as a T cell, and used as an immunotherapy in a subject having a hematological malignancy or at risk for recurrence of the hematological malignancy (e.g., leukemia, lymphoma, myeloma).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
29 March 2018 (29.03.2018)



(10) International Publication Number
WO 2018/058002 A1

(51) International Patent Classification:

C12N 5/0783 (2010.01) C07K 14/725 (2006.01)
C12N 9/22 (2006.01) C07K 14/71 (2006.01)
C07K 14/195 (2006.01) A61K 35/17 (2014.01)

(21) International Application Number:

PCT/US2017/053112

(22) International Filing Date:

22 September 2017 (22.09.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/399,291 23 September 2016 (23.09.2016) US

(71) Applicant: **FRED HUTCHINSON CANCER RESEARCH CENTER** [US/US]; 1100 Fairview Avenue North, Seattle, Washington 98109 (US).

(72) Inventors: **BLEAKLEY, Marie**; 8202 Ashworth Avenue North, Seattle, Washington 98103 (US). **DOSSA, Robson**; 730 North 198th Street, Shoreline, Washington 98133

(US). **SOMMERMEYER, Daniel**; Noerdliche Auffahrt-sallee 43, 80638 Munich (DE).

(74) Agent: **MORGAN, John, A.** et al.; Seed Intellectual Property Law Group LLP, Suite 5400, 701 Fifth Avenue, Seattle, Washington 98104-7064 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: TCRS SPECIFIC FOR MINOR HISTOCOMPATIBILITY (H) ANTIGEN HA-1 AND USES THEREOF




FIG. 30

(57) Abstract: The present disclosure provides compositions and methods for targeting a minor histocompatibility (H) antigen (HA-1H?) to, for example, prevent or manage relapse of a hematological malignancy after allogeneic hematopoietic stem cell transplantation (HCT). Also provided are transgene constructs encoding engineered binding proteins, such as a T cell receptor or a chimeric antigen receptor, optionally encoding additional components such as a co-receptor and/or safety switch. Such transgene constructs can be transduced into an immune cell, such as a T cell, and used as an immunotherapy in a subject having a hematological malignancy or at risk for recurrence of the hematological malignancy (e.g., leukemia, lymphoma, myeloma).

WO 2018/058002 A1

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

TCRS SPECIFIC FOR MINOR HISTOCOMPATIBILITY (H) ANTIGEN HA-1 AND USES THEREOF

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format
5 in lieu of a paper copy, and is hereby incorporated by reference into the specification.
The name of the text file containing the Sequence Listing is
360056_455WO_SEQUENCE_LISTING.txt. The text file is 200 KB, was created on
September 22, 2017, and is being submitted electronically via EFS-Web.

STATEMENT OF GOVERNMENT INTEREST

10 This invention was made with government support under CA154532 awarded
by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND

Patients with hematologic malignancies can be treated with allogeneic
hematopoietic stem cell transplantation (HCT). In the United States, for example,
15 allogeneic HCT transplants have risen steadily over the past 35 years, with
approximately 8,000 transplants per year since 2013 (*see* CIBMTR 2016 Summary).
However, relapse of the hematologic malignancy can occur thereafter. Currently, a
significant number of patients who receive HCT for the treatment of acute leukemia
relapse (approximately 2,000 patients relapse post-HCT each year in the U.S. alone, or
20 about 25 to 50%; *see* Sucheston-Campbell *et al.*, *Curr. Hematol. Malig. Rep.* 10:45-58,
2015). Relapse rates are especially high in patients who are not able to achieve deep
complete remissions and/or are unable to tolerate intensive conditioning regimens prior
to HCT. The prognosis for patients with post-HCT relapse is abysmal: two year
survival rates for patients relapsing at < 100, 100-200 and >200 days after HCT are 3%,
25 9% and 19%, respectively. Patients who receive a second HCT may have better
outcomes, but to be eligible for a second HCT, the patient must first achieve remission,
which typically only occurs in about 30% of patients.

Acute leukemia relapses can, in some cases, be treated with donor lymphocyte infusions from the original stem cell donor. This graft-versus-leukemia (GVL) effect of donor lymphocyte infusion, however, is often accompanied by graft-versus-host disease (GVHD), causing serious mortality and morbidity and is not always effective. If GVL could be selectively increased without enhancing immune responses against normal tissues (graft-versus-host disease, GVHD), post-HCT relapses might be prevented.

Certain minor H antigens are expressed on leukemic stem cells and blasts (*see, e.g., Bleakley and Riddell, Nat. Rev. Cancer* 4:371-380, 2004; Bleakley *et al., Blood* 115:4923-4933, 2010; Bleakley and Riddell, *Immunol. Cell. Biol.* 89:396-407, 2011; van der Harst *et al., Blood* 83:1060-1066, 1994; Bonnet *et al., Proc. Natl. Acad. Sci. USA* 96:8639-8644, 1999; Hambach *et al., Leukemia* 20:371-374, 2006), and have been targeted using cancer-specific T cells. In a small clinical trial of minor H antigen-targeted T cell immunotherapy in patients with post-HCT relapse, clinical responses were observed in some patients (Warren *et al., Blood* 115:3869-3878, 2010). Technical advances in genetic modification of T cells and growing knowledge of T cell biology means that therapeutic doses of antigen-specific T cells can now be prepared efficiently, given to patients, and persist and exert potent anti-tumor effects *in vivo* (Heemskerk *et al., J. Exp. Med.* 199:885-894, 2004; Morgan *et al., Science* 314:126-129, 2006; Griffioen *et al., Haematologica* 93:1535-1543, 2008; Ochi *et al., J. Biomed. Biotechnol.* 2010:5212248, 2010; Schmitt *et al., Hum. Gene Ther.* 20:1240-1248, 2009; Stromnes *et al., Immunol Rev.* 257:145-164, 2014). However, there is a need for cell-based therapies that target leukemia-associated antigens. Presently disclosed embodiments address these needs and provide other related advantages.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts a process used to isolate and characterize HA-1^H-specific T cell clones of the present disclosure. (Top left of Figure 1) To isolate HA-1^H-specific T cells, CD8⁺ cells were primed with dendritic cells (DCs) pulsed with HA1^H peptide (VLHDDLLEA; SEQ ID NO:1) and expanded in microcultures. (Middle left) Following 12 days of incubation in CTL media containing IL-12 and IL-15, aliquots of the T cells were evaluated in a split well microcytotoxicity assay, where T2 cells were

pulsed with the VLHDDLLEA peptide or not. (Bottom left) Cells that specifically reacted against the T2⁺ VLHDDLLEA were cloned by limiting dilution, reassessed for cytotoxicity, and rapidly expended for further evaluation (8 clones). (Top right) Flow cytometry data showing HA-1^H specificity of the indicated seven (7) clones. Staining
 5 for HLA-A2-HA-1^H dextramer staining (y-axis) and CD8 (x-axis). (Bottom right) Data from a chromium release assay (CRA) experiment wherein the indicated CTL clones were tested for specific lysis of ⁵¹CR-labeled T2 cells pulsed with the cognate peptide at the indicated titrations.

Figures 2A-2C show characterization of isolated cytotoxic T cell clones that
 10 were obtained using the method shown in Figure 1. (A) HLA-A2/HA-1^H multimer and CD8⁺ monoclonal antibody staining of seven (7) representative HA-1^H specific clones (1, 2, 10, 13, 4, 16, and 5) and a control clone specific for another tumor antigen. (B) Data from a chromium release assay in which the seven HA-1^H-specific clones were tested for killing HA-1 peptide-pulsed target cells. (C) Data from cytotoxicity assays in
 15 which the HA-1^H-specific CTL clones were incubated with peptide-pulsed T2 cells, HA-1^{H+} AML cell line THP-1, HA-1^{H+} primary AML, or HA-1⁻ AML.

Figure 3 depicts a representative HA-1^H TCR-encoding lentiviral construct of the present disclosure.

Figure 4 shows a procedure for evaluating T cells transduced to express HA-1
 20 TCRs.

Figures 5A-5C show expression and activity of HA-1^H-transduced CD8⁺ T cells. (A) Flow cytometry data showing HA-1^H dextramer binding and CD8 expression of T cells transduced with TCR2 or TCR16. (B) Killing activity of CD8⁺ T cells transduced with TCR2 or TCR16, as well as of the corresponding 'parental clones', *i.e.*,
 25 the T cell clones from which TCR2 and TCR16 were isolated and of a control clone specific for a control a different antigen (SMCY). Left, lysis of HA-1^H-pulsed T2 cells. Right, T2 cells pulsed with irrelevant SMCY peptide. (C) Specific lysis of T2 cells pulsed with the indicated amount of HA-1^H peptide (x-axis) by HA-1^H TCR2 (dashed line with circles), HA-1^H TCR16 (dashed line with squares), parental TCR2 clone (solid line with circles), parental TCR16 clone (solid line with squares), and a heterologous or parent clone specific for SMCY peptide (lower two lines on graph).
 30

Figures 6A-6E show transduction and activity of the HA-1^H TCRs into CD8⁺ T cells using a lentiviral vector. (A) Flow cytometry showing HLA-A2/HA-1^H multimer staining of CD8⁺ T cells transduced to contain a polynucleotide encoding a HA-1^H specific TCR (specifically, TCR clones 1, 2, 10, 16, and 5), which were delivered with a lentiviral vector (LV), or with a TCR specific for a different minor H antigen (control). (B-E) A chromium release assay (CRA) was used to evaluate specific lytic activity. (B) Lysis of T2 target cells pulsed with HA-1 peptide antigen at various concentrations by TCR-transduced CD8⁺ T cells (solid lines and symbols-TCR2 circles, TCR16 squares), HA-1^H-specific T cell clones (dashed lines, open symbols-clone 2=circles, clone 16=squares), or T cell clone control (diamonds). (C) Lysis of HLA-A2⁺HA-1^H⁺ LCL by CD8⁺ T cells transduced with HA-1^H-specific TCR2 (circles) or HA-1^H-specific TCR16 (squares) at the indicated effector: target (E:T) ratios. (D) Lysis of HLA-A2⁺/HA-1⁺ homozygous (H/H) (circles n=7), HLA-A2⁺/HA1^H⁺ heterozygous (H/R) (square n=22), HLA-A2⁺/HA-1^H⁻ (R/R) (triangles n=17) or HLA-A2 negative (inverted triangles n=41) hematopoietic cell (LCL) targets by HA-1^H TCR2 transduced CD8⁺ T cells. (E) Lysis of LCL with common HLA alleles by HA-1^H TCR2 transduced CD8⁺ T cells. An E:T ratio of 20:1 was used unless otherwise specified. Data comparable to that shown in (D) and (E) were also obtained with HA-1^H TCR16 (not shown).

Figure 7 provides data from cytotoxicity experiments in which target cells that endogenously express HA-1 (HA-1^H⁺ LCL (H/H or H/R) and HA-1^H⁻LCL (R/R) by were incubated with TCR2 transduced cells or with 'parental' clone 2 cells. Also shown is a control clone specific for a Y chromosome-associated minor H antigen (FIDSYICQV).

Figures 8A-8E show specific killing of HA-1⁺ leukemia cells by HA-1^H-TCR2-transduced CD8⁺ T cells. (A) HA-1^H-specific expression of CD107a on HA-1^H-TCR2 transduced CD8⁺ T cells showing degranulation after 5h co-culture (1:1) with a panel of primary AML samples. (B-E) CRA showing lysis of leukemia and lymphoma targets by HA-1 TCR-transduced CD8⁺ T cells; (B) Lysis of primary HA-1^H⁺ AML or HA-1^H⁻ AML by HA-1^HTCR-transduced CD8⁺ T cells (dark grey bars) and HA-1^H-specific T cell clone 2 (light grey bars); (C) HLA-A2⁺/ HA-1^H⁺ primary AML (AML1) at various

E:T ratios; (D) B-ALL lines (1) BALL-1, (2) RS4;11, T-ALL lines (1) MOLT4, (2) CEM (3) RPMI-8402 (4) HSB-2 and AML line NB-4; (E) T cell lymphoma (SUP-M2 HLA-A2⁺, HA-1⁺; SU-DHL-1 HLA-A2⁺ HA-1⁻) cell lines. In (D) HLA-A2⁻ and/or HA-1^{H-} (WT) cell lines were transduced (TD) with LV encoding *HLA-A2 or **HLA-A2 and HA-1^H minigene if the WT was HLA-A2⁻ or had a HA-1^{H-} genotype. An E:T ratio of 20:1 was used, unless otherwise specified.

Figures 9A and 9B show cytotoxicity of T cells transduced with TCR2 or TCR16 (as well as of the corresponding parental clones) against HA-1^{H+} or HA-1^{H-} primary leukemia cells. 9A: specific lysis (4h CRA) of the indicated cell lines. 9B: specific lysis of target cells at the indicated effector:target ratios.

Figure 10 shows data from a cytotoxicity assay in which TCR2- and TCR16-transduced cells (and parental clones) were incubated with HA-1^H genotypically positive dermal fibroblasts, with or without exposure of the fibroblasts to interferon gamma (IFN γ).

Figures 11A-11C show characterization of CD4⁺ T cells transduced with the HA-1^H TCR2 and CD8 co-receptor variants. (A) (i, ii) Mean fluorescence intensity (MFI) of HA-1^H/HLA-A2 multimer staining of CD4⁺ T cells transduced with CD8 α and/or β M1-M5 chains as indicated.. (iii) MFI of the various CD8 co-receptor constructs is summarized in the graph. (B) CRA showing lysis of T2 pulsed with HA-1^H peptide at various concentrations by HA-1^H-specific CD8⁺T cells (solid circles), CD4⁺ T cells transduced with the CD8 α and β chains (squares, diamonds, downward triangles), CD8 α chains alone (upward triangles), or HA-1^H TCR only (open circle). (C) Proliferation assay showing dilution of the carboxyfluorescein (CFSE) dye with cell division in CD4⁺ T cells transduced with (top to bottom) the HA-1^H TCR alone, with CD8 α chain, CD8 α and β M1 chain, or CD8 α and β M4 chain, in response to stimulation with HLA-A2⁺ HA-1^{H+} LCL, HA-1^{H-} LCL, or media only.

Figures 12A and 12B show further functional characterization of CD4⁺ T cells transduced with the HA-1^H-TCR2 and a CD8 co-receptor. (A) Intracellular cytokine assay showing IL-2 and IFN- γ production by CD8⁺ T cells (left) and CD4⁺ T cells (right) transduced with HA-1^HTCR2 LV (upper panels) or HA-1^HTCR2-CD8 co-receptor LV (lower panels) in response to HLA-A2⁺/ HA-1^{H+} AML or HLA-A2⁺/ HA-

1^H- AML; (B) CFSE assay showing proliferation of CD8⁺ T cells (left) and CD4⁺ T cells (right) transduced with HA-1^H-specific TCR2 LV (upper panels) or HA-1^HTCR2-CD8 co-receptor LV (lower panels) in response to HLA-A2⁺/ HA-1^H primary AML , HLA-A2⁺/ HA-1^H- AML or media control.

5 **Figure 13** provides representative diagrams of safety switch gene constructs of the present disclosure (bottom) and schema using the safety switch gene constructs to kill T cells (top).

Figure 14 provides data from a cytotoxicity assay in which primary T cells were transduced with transgene constructs expressing safety switch genes and the HA-1^H specific TCR2 (or were transduced with TCR2 alone) and incubated with T2 cells pulsed with HA-1^H peptide.

Figure 15 shows percent survival of the transduced TCRs in the presence or absence of a "suicide drug" that activates the encoded safety switch.

Figure 16 shows survival of CD8⁺ T cells transduced with HA-1^H TCR2 plus safety genes after exposure to the cognate safety-switch activating drug at the indicated concentrations. Survival of iCasp9-TCR2, tEGFR-TCR2, RQR8-TCR2 and Myc-TCR2 CD8⁺-transduced T cells was measured after 24 hours of incubation with the indicated concentrations of the respective safety switch activating drug: AP1903; anti-EGFR mAb (Cetuximab) + complement; anti-CD20Mab (Rituximab) + complement; anti-myc mAb + complement. Residual HA-1 TCR2 transduced T cells were quantified by flow cytometry. The arrows indicate the drug concentrations that can be achieved and tolerated in humans *in vivo*.

Figure 17 shows five (5) different types of constructs for the evaluation of iCasp 9- HA-1^H TCR-CD8⁺ expression constructs: (i) iCasp 9 and TCR2; (ii) TCR 2 and CD8 co-receptor; (iii) iCasp 9, TCR 2 and CD8 co-receptor; (iv) iCasp 9, TCR 2 and CD8 co-receptor with RQR tag on the CD8 co-receptor; and (v) iCasp 9, TCR 2 and CD8 co-receptor with Q (CD34) tag on the alpha chain of the TCR.

Figure 18 depicts a flow chart for evaluating TCR-transduced cells of the present disclosure.

Figure 19 provides flow cytometry data showing expression of the engineered TCR in each of the indicated transgene constructs before (top row) and after (bottom row) enrichment with an HA-1 dextramer.

Figures 20A and 20B show specific lysis of peptide-pulsed target cells (20A) and LCL lines (20B) by T cells transduced with transgene constructs: iCasp-9-TCR (- -; first bar); TCR and CD8 co-receptor (-▲-; second bar); iCasp-9-TCR-CD8 co-receptor (-◆-; third bar); iCasp9-TCR-RQR-CD8 co-receptor (-■-; fourth bar); and iCas9-CD34tag-TCR-CD8 co-receptor (-●-; fifth bar).

Figure 21 shows cytokine elaboration by T cells transduced with the indicated transgene constructs following stimulation with HA-1^{H+} (top) or HA-1^{H-} (bottom) cell lines.

Figure 22 shows cytolytic activity (bottom) of T cells transduced with the transgene constructs (top) against indicated target cell lines.

Figure 23 shows the absence of cytolytic activity of T cells transduced with the indicated transgene constructs against indicated non-hematopoietic cells in the presence or absence of interferon-gamma. HA-1^{H+} hematopoietic control cells were killed by the T cells.

Figure 24 shows cytokine elaboration by cells transduced with the indicated transgene constructs when exposed to primary leukemia cells.

Figure 25 provides flow cytometry histograms showing proliferation of transduced T cells when stimulated with HA-1^{H+} primary leukemia cells. Top: scheme for measuring proliferation by staining F1 cells with CFSE (left), and representative proliferation data from a T cell transduced with a transgene constructs of the present disclosure. Bottom: proliferation of T cells transduced with the indicated transgene constructs.

Figure 26 shows survival (bottom) of T cells transduced with transgene constructs as shown (top) following introduction of the cognate suicide drug at the indicated concentrations.

Figure 27 shows an enrichment scheme for engineered T cells of the present disclosure. T cells are transduced with the indicated transgene constructs (top) and examined for expression (middle). Cells expressing a selectable transduction marker

(CD34 epitope; two right-most scatter plots) are selected using magnetic beads with anti-CD34 antibody.

Figures 28A and 28B provide flow cytometry data (28A) showing frequency of transduced T cells before (top row) and after (bottom row) magnetic selection. Four
 5 scatter plots at left: staining for CD34 selection marker and CD8. Four scatter plots at right: staining for HA-1^H dextramer and CD8. Also shown (28B) are cell counts of cells expressing the CD34 selection marker (left) or being specific for HA-1^H (right) before and after selection.

Figure 29 provides a schematic showing various functionalities of a TCR-safety
 10 gene construct of the present disclosure.

Figure 30 provides a diagram of a lentiviral delivery vector encoding an exemplary iC9-HA-1^H-TCR-RQR-CD8 construct of the present disclosure.

Figure 31 shows the functional characterization of CD8⁺ and CD4⁺ cells transduced with a TCR2-CD8 transgene construct of the present disclosure. Left panel:
 15 flow cytometry data showing cytokine release (IL-2; IFN- γ) in response to HA-1^H peptide antigen. Middle panel: quantification of the flow cytometry data. Right panels: proliferation of transduced cells in response to HA-1^H.

Figures 32A-32E show characterization of CD4⁺ and CD8⁺ T cells transduced with a (HA-1^H-specific TCR)-(RQR)-(CD8) and expanded to clinical scale. (A)
 20 Growth of transduced T cells. (B) HA-1^H TCR multimer binding and CD34 expression on transduced T cells by flow cytometry with HA-1/HLA-A2 multimer staining. (C) Expression of co-stimulatory and homing molecules on T cells at the time of the apheresis, after CD45RA depletion and following transduction and expansion (N=5). (D, E) Expression of 'exhaustion' markers on HA-1^H TCR CD8⁺ and CD4⁺ in the final
 25 cell product (N=3) (D) and a representative example (E).

Figures 33A-E show data from functional recognition assays in which the clinical-scale HA-1^H-TCR2-RQR-CD8-transduced T cells were incubated with target cells. (A) Lysis of target T2 cells pulsed with a range of VLH (solid lines, dark grey) and VLR (solid line, light grey) peptide concentrations by CD8⁺ (solid lines) and CD4⁺
 30 T cells (dashed lines) in CRA at ET ratio 20:1. (B) Lysis of HA-1^H A2⁺ LCL, HA-1^H-

A2⁺ LCL and AML HA-1^H + A2⁺ cell line (THP-1) by CD8⁺ (solid lines) in CRA (C) IL-2, IFN- γ , and TNF α production by T cells in response to stimulation by T2 cells pulsed with 10ng/ml of HA-1 peptide. (D) Pie charts displaying the number of cytokine types secreted by T cells. (E) Concentration of cytokines and granzyme B in media 24 hours after stimulation of T cells by T2 cells pulsed with VLH or VLR peptides, as measured by multiplex immunoassay.

Figure 34A shows (A) CD34 (left graph) and HA-1^H TCR (right graph) expression on T cells in the final product before and after enrichment by CD34 immunomagnetic beads (N=3). **Figure 34B** shows survival of T cells in the cell product after 24 hours of incubation with 5ng/ml AP1903 or media control only.

Figure 35 shows cytolytic activity of T cells transduced with another HA-1^H TCR of the present disclosure (circles) and of control cells (triangles) against T2 cells pulsed with the HA-1^H antigen at the indicated concentrations.

DETAILED DESCRIPTION

In some aspects, the present disclosure provides compositions and methods for treating hyperproliferative diseases characterized by expression of minor histocompatibility antigen HA-1^H. By way of background, human leukocyte antigen (HLA) testing is typically used to match organ, cell, and tissue transplant recipients with compatible donors. HLA testing identifies the major HLA genes a person has inherited and the corresponding antigens, or proteins, which are present on the surface of their cells. These antigens help the body's immune system distinguish which cells are "self", and which are "foreign" or "non-self." Any cells that are recognized as "non-self" can trigger an immune response, such as T cell-mediated cytotoxicity or the production of antibodies. Of note, tests for major HLA genes do not identify the minor HLA genes, which give rise to further antigens. Accordingly, even "HLA-matched" donor cells may attack healthy recipient cells expressing a perceived "foreign" minor HLA protein or peptide. Minor H antigens that are expressed on epithelial tissues are targets of alloreactive T cells, leading to graft-versus-host disease. However, some

minor H antigens are associated with genes that are expressed predominantly or exclusively in the hematopoietic system, including hematopoietic cells that can be affected by hematological malignancies. Thus, minor H antigens with restricted expression are potential targets for therapies that seek to augment the graft-versus-leukemia effect and thereby prevent relapse.

The minor histocompatibility antigen HA-1^H is encoded by the polymeric *HMHA1* gene (also called Rho GTPase-activating protein 45) and is highly expressed in leukemia cells and normal hematopoietic cells (*see, e.g., Griffioen et al., Front. Immunol.* 7:100, 2016; Spierings *et al. Biol. Blood Marrow Transpl.* 19:1244-1253, 2013, the HA-1 expression disclosure of which is incorporated herein by reference), but not in normal non-hematopoietic cells. *HMHA1* variants (rs1801284 A/A or A/G) present in 52% of individuals give rise to an immunogenic peptide containing a histidine residue in place of an arginine (VLHDDLLEA; SEQ ID NO:66) (R139H polymorphism) and HLA presentation of this peptide occurs in individuals with the common HLA-A*0201 (A2) allele (den Haan *et al., Science* 279:1054-1057, 1998). T cell therapies targeting HA-1^H are therefore applicable to approximately 25% of subjects transplanted for hematological malignancies and require a T cell donor who is either HLA-A2 negative or HA-1^H negative ("HA-1^R"; rs1801284 G/G - VLRDDLLEA; SEQ ID NO: 65).

In some aspects, the present disclosure provides engineered immune cells expressing binding proteins, such as TCRs and CARs, specific for HA-1^H. Such engineered immune cells can be used as a standalone therapy to treat a hematologic malignancy or to prevent a relapse or recurrence thereof, or such cells can be used as part of a therapeutic regimen comprising additional therapies or agents (*e.g., following, or in combination with, allogeneic HCT*).

Prior to setting forth this disclosure in more detail, it may be helpful to an understanding thereof to provide definitions of certain terms to be used herein. Additional definitions are set forth throughout this disclosure.

In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one

hundredth of an integer), unless otherwise indicated. Also, any number range recited herein relating to any physical feature, such as polymer subunits, size or thickness, are to be understood to include any integer within the recited range, unless otherwise indicated. As used herein, the term "about" means $\pm 20\%$ of the indicated range, value, or structure, unless otherwise indicated. It should be understood that the terms "a" and "an" as used herein refer to "one or more" of the enumerated components. The use of the alternative (*e.g.*, "or") should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the terms "include", "have", and "comprise" are used synonymously, which terms and variants thereof are intended to be construed as non-limiting.

In addition, it should be understood that the individual compounds, or groups of compounds, derived from the various combinations of the structures and substituents described herein, are disclosed by the present application to the same extent as if each compound or group of compounds was set forth individually. Thus, selection of particular structures or particular substituents is within the scope of the present disclosure.

The term "consisting essentially of" is not equivalent to "comprising" and refers to the specified materials or steps of a claim, or to those that do not materially affect the basic characteristics of a claimed subject matter. For example, a protein domain, region, or module (*e.g.*, a binding domain, hinge region, linker module) or a protein (which may have one or more domains, regions, or modules) "consists essentially of" a particular amino acid sequence when the amino acid sequence of a domain, region, module, or protein includes extensions, deletions, mutations, or a combination thereof (*e.g.*, amino acids at the amino- or carboxy-terminus or between domains) that, in combination, contribute to at most 20% (*e.g.*, at most 15%, 10%, 8%, 6%, 5%, 4%, 3%, 2% or 1%) of the length of a domain, region, module, or protein and do not substantially affect (*i.e.*, do not reduce the activity by more than 50%, such as no more than 40%, 30%, 25%, 20%, 15%, 10%, 5%, or 1%) the activity of the domain(s), region(s), module(s), or protein (*e.g.*, the target binding affinity of a binding protein).

As used herein, an "immune system cell" means any cell of the immune system that originates from a hematopoietic stem cell in the bone marrow, which gives rise to

two major lineages, a myeloid progenitor cell (which give rise to myeloid cells such as monocytes, macrophages, dendritic cells, meagakaryocytes and granulocytes) and a lymphoid progenitor cell (which give rise to lymphoid cells such as T cells, B cells and natural killer (NK) cells). Exemplary immune system cells include a CD4⁺ T cell, a
 5 CD8⁺ T cell, a CD4 - CD8 - double negative T cell, a $\gamma\delta$ T cell, a regulatory T cell, a natural killer cell, and a dendritic cell. Macrophages and dendritic cells can be referred to as "antigen presenting cells" or "APCs," which are specialized cells that can activate T cells when a major histocompatibility complex (MHC) receptor on the surface of the APC complexed with a peptide interacts with a TCR on the surface of a T cell.

10 A "T cell" or "T lymphocyte" is an immune system cell that matures in the thymus and produces T cell receptors (TCRs). T cells can be naïve ("T_N"; not exposed to antigen; increased expression of CD62L, CCR7, CD28, CD3, CD127, and CD45RA, and decreased or no expression of CD45RO as compared to T_{CM} (described herein)), memory T cells (T_M) (antigen experienced and long-lived), and effector cells (antigen-
 15 experienced, cytotoxic). T_M can be further divided into subsets of central memory T cells (T_{CM}, expresses CD62L, CCR7, CD28, CD45RO) and effector memory T cells (T_{EM}, express CD45RO, decreased expression of CD62L, CCR7, and CD28). Effector T cells (T_E) refers to antigen-experienced CD8⁺ cytotoxic T lymphocytes that express CD45RA, have decreased expression of CD62L, CCR7, and CD28 as compared to
 20 T_{CM}, and are positive for granzyme and perforin. Helper T cells (T_H) are CD4⁺ cells that influence the activity of other immune cells by releasing cytokines. CD4⁺ T cells can activate and suppress an adaptive immune response, and which of those two functions is induced will depend on presence of other cells and signals. T cells can be collected using known techniques, and the various subpopulations or combinations
 25 thereof can be enriched or depleted by known techniques, such as by affinity binding to antibodies, flow cytometry, or immunomagnetic selection. Other exemplary T cells include regulatory T cells, such as CD4⁺ CD25⁺ (Foxp3⁺) regulatory T cells and Treg17 cells, as well as Tr1, Th3, CD8⁺CD28⁻, and Qa-1 restricted T cells.

"T cell receptor" (TCR) refers to an immunoglobulin superfamily member
 30 (having a variable binding domain, a constant domain, a transmembrane region, and a short cytoplasmic tail; see, *e. g.*, Janeway *et al.*, Immunobiology: The Immune System

in Health and Disease, 3rd Ed., Current Biology Publications, p. 433, 1997) capable of specifically binding to an antigen peptide bound to a MHC receptor. A TCR can be found on the surface of a cell or in soluble form and generally is comprised of a heterodimer having α and β chains (also known as TCR α and TCR β , respectively), or γ and δ chains (also known as TCR γ and TCR δ , respectively).

Like other immunoglobulins (*e.g.*, antibodies), the extracellular portion of TCR chains (*e.g.*, α -chain, β -chain) contain two immunoglobulin domains, a variable domain (*e.g.*, α -chain variable domain or V $_{\alpha}$ chain variable domain or V $_{\beta}$; typically amino acids 1 to 116 based on Kabat numbering (Kabat *et al.*, "Sequences of Proteins of Immunological Interest, US Dept. Health and Human Services, Public Health Service National Institutes of Health, 1991, 5th ed.) at the N-terminus, and one constant domain (*e.g.*, α -chain constant domain or C $_{\alpha}$, typically 5 amino acids 117 to 259 based on Kabat, β -chain constant domain or C $_{\beta}$, typically amino acids 117 to 295 based on Kabat) adjacent the cell membrane. Also, like immunoglobulins, the variable domains contain complementary determining regions (CDRs) separated by framework regions (FRs) (see, *e.g.*, Jores *et al.*, *Proc. Nat'l Acad. Sci. USA* 87:9138, 1990; Chothia *et al.*, *EMBO J.* 7:3745, 1988; see also Lefranc *et al.*, *Dev. Comp. Immunol.* 27:55, 2003).

"Antigen" or "Ag" as used herein refers to an immunogenic molecule that provokes an immune response. This immune response may involve antibody production, activation of specific immunologically-competent cells (*e.g.*, T cells), or both. An antigen (immunogenic molecule) may be, for example, a peptide, glycopeptide, polypeptide, glycopolypeptide, polynucleotide, polysaccharide, lipid or the like. It is readily apparent that an antigen can be synthesized, produced recombinantly, or derived from a biological sample. Exemplary biological samples that can contain one or more antigens include tissue samples, tumor samples, cells, biological fluids, or combinations thereof. Antigens can be produced by cells that have been modified or genetically engineered to express an antigen, or that endogenously (*e.g.*, without modification or genetic engineering by human intervention) express a mutation or polymorphism that is immunogenic.

"Major histocompatibility complex" (MHC) refers to glycoproteins that deliver peptide antigens to a cell surface of all nucleated cells. MHC class I molecules are

heterodimers having a membrane spanning α chain (with three α domains) and a non-covalently associated β_2 microglobulin. MHC class II molecules are composed of two transmembrane glycoproteins, α and β , both of which span the membrane. Each chain has two domains. MHC class I molecules deliver peptides originating in the cytosol to the cell surface, where a peptide:MHC complex is recognized by CD8⁺ T cells. MHC class II molecules deliver peptides originating in the vesicular system to the cell surface, where they are recognized by CD4⁺ T cells. Human MHC is referred to as human leukocyte antigen (HLA).

The term "epitope" or "antigenic epitope" includes any molecule, structure, amino acid sequence or protein determinant that is recognized and specifically bound by a cognate binding molecule, such as an immunoglobulin, T cell receptor (TCR), chimeric antigen receptor, or other binding molecule, domain or protein. Epitopic determinants generally contain chemically active surface groupings of molecules, such as amino acids or sugar side chains, and can have specific three dimensional structural characteristics, as well as specific charge characteristics.

As used herein "specifically binds" or "specific for" refers to an association or union of a binding protein (*e.g.*, TCR receptor) or a binding domain (or fusion protein thereof) to a target molecule with an affinity or K_a (*i.e.*, an equilibrium association constant of a particular binding interaction with units of 1/M) equal to or greater than 10^5 M^{-1} (which equals the ratio of the on-rate [k_{on}] to the off-rate [k_{off}] for this association reaction), while not significantly associating or uniting with any other molecules or components in a sample. Binding proteins or binding domains (or fusion proteins thereof) may be classified as "high affinity" binding proteins or binding domains (or fusion proteins thereof) or as "low affinity" binding proteins or binding domains (or fusion proteins thereof). "High affinity" binding proteins or binding domains refer to those binding proteins or binding domains having a K_a of at least 10^7 M^{-1} , at least 10^8 M^{-1} , at least 10^9 M^{-1} , at least 10^{10} M^{-1} , at least 10^{11} M^{-1} , at least 10^{12} M^{-1} , or at least 10^{13} M^{-1} . "Low affinity" binding proteins or binding domains refer to those binding proteins or binding domains having a K_a of up to 10^7 M^{-1} , up to 10^6 M^{-1} , up to 10^5 M^{-1} . Alternatively, affinity can be defined as an equilibrium

dissociation constant (K_d) of a particular binding interaction with units of M (*e.g.*, 10^{-5} M to 10^{-13} M).

In certain embodiments, a receptor or binding domain may have "enhanced affinity," which refers to selected or engineered receptors or binding domains with stronger binding to a target antigen than a wild type (or parent) binding domain. For example, enhanced affinity may be due to a K_a (equilibrium association constant) for the target antigen that is higher than the wild type binding domain, due to a K_d (dissociation constant) for the target antigen that is less than that of the wild type binding domain, due to an off-rate (k_{off}) for the target antigen that is less than that of the wild type binding domain, or a combination thereof. In certain embodiments, enhanced affinity TCRs can be codon optimized to enhance expression in a particular host cell, such as a cell of the immune system, a hematopoietic stem cell, a T cell, a primary T cell, a T cell line, a NK cell, or a natural killer T cell (Scholten *et al.*, *Clin. Immunol.* 119:135, 2006). The T cell can be a CD4+ or a CD8+ T cell.

A variety of assays are known for identifying binding domains of the present disclosure that specifically bind a particular target, as well as determining binding domain or fusion protein affinities, such as multimer/tetramer staining, Western blot, ELISA, analytical ultracentrifugation, spectroscopy and surface plasmon resonance (Biacore®) analysis (*see, e.g.*, Dolton *et al.*, *Immunology* 146:11-22, 2015, Scatchard *et al.*, *Ann. NY Acad. Sci.* 51:660, 1949; Wilson, *Science* 20295:2103, 2002; Wolff *et al.*, *Cancer Res.* 53:2560, 1993; and U.S. Patent Nos. 5,283,173, 5,468,614, or the equivalent; all incorporated herein by reference).

The source of a TCR as used in the present disclosure can be from various animal species, such as a human, mouse, rat, rabbit or other mammal.

As used herein, the term "CD8 co-receptor" or "CD8" means the cell surface glycoprotein CD8, either as an alpha-alpha homodimer or an alpha-beta heterodimer. The CD8 co-receptor assists in the function of cytotoxic T cells ($CD8^+$) and functions through signaling via its cytoplasmic tyrosine phosphorylation pathway (Gao and Jakobsen, *Immunol. Today* 21:630-636, 2000; Cole and Gao, *Cell. Mol. Immunol.* 1:81-88, 2004). There are five (5) different CD8 beta chains (*see* UniProtKB identifier P10966) and a single CD8 alpha chain (*see* UniProtKB identifier P01732)

"CD4" is an immunoglobulin co-receptor glycoprotein that assists the TCR in communicating with antigen-presenting cells (*see*, Campbell & Reece, Biology 909 (Benjamin Cummings, Sixth Ed., 2002)). CD4 is found on the surface of immune cells such as T helper cells, monocytes, macrophages, and dendritic cells, and includes four immunoglobulin domains (D1 to D4) that are expressed at the cell surface. During antigen presentation, CD4 is recruited, along with the TCR complex, to bind to different regions of the MHCII molecule (CD4 binds MHCII $\beta 2$, while the TCR complex binds MHCII $\alpha 1/\beta 1$). Without wishing to be bound by theory, it is believed that close proximity to the TCR complex allows CD4-associated kinase molecules to phosphorylate the immunoreceptor tyrosine activation motifs (ITAMs) present on the cytoplasmic domains of CD3. This activity is thought to amplify the signal generated by the activated TCR in order to produce various types of T helper cells.

In certain embodiments, a TCR is found on the surface of T cells (or T lymphocytes) and associates with a CD3 complex. "CD3" is a multi-protein complex of six chains (*see*, Abbas and Lichtman, 2003; Janeway *et al.*, p. 172 and 178, 1999) that is associated with antigen signaling in T cells. In mammals, the complex comprises a CD3 γ chain, a CD3 δ chain, two CD3 ϵ chains, and a homodimer of CD3 ζ chains. The CD3 γ , CD3 β , and CD3 ϵ chains are highly related cell surface proteins of the immunoglobulin superfamily containing a single immunoglobulin domain. The transmembrane regions of the CD3 γ , CD3 β , and CD3 ϵ chains are negatively charged, which is a characteristic that allows these chains to associate with the positively charged T cell receptor chains. The intracellular tails of the CD3 γ , CD3 β , and CD3 ϵ chains each contain a single conserved motif known as an immunoreceptor tyrosine based activation motif or ITAM, whereas each CD3 ζ chain has three. Without wishing to be bound by theory, it is believed that the ITAMs are important for the signaling capacity of a TCR complex. CD3 as used in the present disclosure may be from various animal species, including human, mouse, rat, or other mammals.

As used herein, "TCR complex" refers to a complex formed by the association of CD3 with TCR. For example, a TCR complex can be composed of a CD3 γ chain, a CD3 β chain, two CD3 ϵ chains, a homodimer of CD3 ζ chains, a TCR α chain, and a TCR β chain. Alternatively, a TCR complex can be composed of a CD3 γ chain, a CD3 β

chain, two CD3 ϵ chains, a homodimer of CD3 ζ chains, a TCR γ chain, and a TCR β chain.

A "component of a TCR complex", as used herein, refers to a TCR chain (*i.e.*, TCR α , TCR β , TCR γ or TCR δ), a CD3 chain (*i.e.*, CD3 γ , CD3 δ , CD3 ϵ or CD3 ζ), or a
 5 complex formed by two or more TCR chains or CD3 chains (*e.g.*, a complex of TCR α and TCR β , a complex of TCR γ and TCR δ , a complex of CD3 ϵ and CD3 δ , a complex of CD3 γ and CD3 ϵ , or a sub-TCR complex of TCR α , TCR β , CD3 γ , CD3 δ , and two CD3 ϵ chains).

As used herein, the term "HA-1^H antigen" or "HA-1^H peptide antigen" or
 10 "HA-1^H-containing peptide antigen" (or "minor HA-1^H antigen" or "minor HA-1^H peptide antigen" or "minor HA-1^H-containing peptide antigen" or "minor Histocompatibility HA-1^H antigen peptide") refers to a naturally or synthetically produced peptide portion of a HMHA1 protein ranging in length from about 7 amino acids, about 8 amino acids, about 9 amino acids, about 10 amino acids, up to about
 15 20 amino acids, and comprising the R139H substitution polymorphism), which can form a complex with a MHC (*e.g.*, HLA) molecule, and a binding protein of this disclosure specific for a HA-1^H peptide:MHC (*e.g.*, HLA) complex can specifically bind to such as complex. An exemplary HA-1^H HA-1 peptide antigen comprises a peptide having the amino acid VL**H**D~~D~~LLEA (SEQ ID NO: 66), wherein the bolded
 20 histidine in the sequence represents the R139H polymorphism.

The term "HA-1^H-specific binding protein," as used herein, refers to a protein or polypeptide, such as a TCR or CAR, that specifically binds to an HA-1^H peptide antigen (or to an HA-1^H peptide antigen:HLA complex, *e.g.*, on a cell surface), and does not bind an HMHA peptide that does not contain the HA-1^H polymorphism (*e.g.*, a
 25 peptide comprising the amino acid sequence shown in SEQ ID NO:65) and does not bind to an HLA complex containing such an HMHA peptide.

In certain embodiments, a HA-1^H-specific binding protein specifically binds to an HA-1-containing peptide (or an HA-1^H peptide:HLA complex) with a K_d of less than about 10^{-8} M, less than about 10^{-9} M, less than about 10^{-10} M, less than about 10^{-11} M,
 30 less than about 10^{-12} M, or less than about 10^{-13} M, or with an affinity that is about the same as, at least about the same as, or is greater than at or about the affinity exhibited

by an exemplary HA-1^H-specific binding protein provided herein, such as any of the HA-1^H-specific TCRs provided herein, for example, as measured by the same assay. In certain embodiments, a HA-1-specific binding protein comprises a HA-1-specific immunoglobulin superfamily binding protein or binding portion thereof.

5 Principles of antigen processing by antigen presenting cells (APC) (such as dendritic cells, macrophages, lymphocytes or other cell types), and of antigen presentation by APC to T cells, including major histocompatibility complex (MHC)-restricted presentation between immunocompatible (*e.g.*, sharing at least one allelic form of an MHC gene that is relevant for antigen presentation) APC and T cells, are
 10 well established (*see, e.g.*, Murphy, Janeway's Immunobiology (8th Ed.) 2011 Garland Science, NY; chapters 6, 9 and 16). For example, processed antigen peptides originating in the cytosol (*e.g.*, tumor antigen, intracellular pathogen) are generally from about 7 amino acids to about 11 amino acids in length and will associate with class I MHC (HLA) molecules, whereas peptides processed in the vesicular system
 15 (*e.g.*, bacterial, viral) will vary in length from about 10 amino acids to about 25 amino acids and associate with class II MHC (HLA) molecules.

An "altered domain" or "altered protein" refers to a motif, region, domain, peptide, polypeptide, or protein with a non-identical sequence identity to a wild type motif, region, domain, peptide, polypeptide, or protein (*e.g.*, a wild type TCR α chain,
 20 TCR β chain, TCR α constant domain, TCR β constant domain) of at least 85% (*e.g.*, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9%).

Altered domains or altered proteins or derivatives can include those based on all possible codon choices for the same amino acid and codon choices based on
 25 conservative amino acid substitutions. For example, the following six group's each contain amino acids that are conservative substitutions for one another: 1) alanine (ala; A), serine (ser; S), threonine (thr; T); 2) aspartic acid (asp; D), glutamic acid (glu; E); 3) asparagine (asn; N), glutamine (gln; Q); 4) arginine (arg; R), lysine (lys; K); 5) Isoleucine (ile; I), leucine (L), methionine (met; M), valine (val; V); and
 30 6) phenylalanine (phe; F), tyrosine (tyr; Y), tryptophan (trp; W). (See also WO97/09433 at page 10, Lehninger, Biochemistry, 2nd Edition, Worth Publishers, Inc.,

NY, NY, pp. 71-77, 1975; Lewin Genes IV, Oxford University Press, NY and Cell Press, Cambridge, MA, p.8, 1990; Creighton, Proteins, W.H. Freeman and Company 1984). In addition, individual substitutions, deletions or additions that alter, add or delete, a single amino acid or a small percentage of amino acids in an encoded sequence
5 are also "conservative substitutions."

As used herein, "nucleic acid" or "nucleic acid molecule" refers to any of deoxyribonucleic acid (DNA), ribonucleic acid (RNA), oligonucleotides, polynucleotides, fragments thereof generated, for example, by the polymerase chain reaction (PCR) or by *in vitro* translation, and also to fragments generated by any of
10 ligation, scission, endonuclease action, or exonuclease action. In certain embodiments, the nucleic acids of the present disclosure are produced by PCR. Nucleic acids can be composed of monomers that are naturally occurring nucleotides (such as deoxyribonucleotides and ribonucleotides), analogs of naturally occurring nucleotides (*e.g.*, α -enantiomeric forms of naturally occurring nucleotides), or a combination of
15 both. Modified nucleotides can have modifications in or replacement of sugar moieties, or pyrimidine or purine base moieties. Nucleic acid monomers can be linked by phosphodiester bonds or analogs of such linkages. Analogous of phosphodiester linkages include phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoranilidate, phosphoramidate, and
20 the like. Nucleic acid molecules can be either single stranded or double stranded.

The term "isolated" means that the material is removed from its original environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally occurring nucleic acid or polypeptide present in a living animal is not isolated, but the same nucleic acid or polypeptide, separated from some or all of the co-existing
25 materials in the natural system, is isolated. Such a nucleic acid could be part of a vector and/or such nucleic acid or polypeptide could be part of a composition (*e.g.*, a cell lysate), and still be isolated in that such vector or composition is not part of the natural environment for the nucleic acid or polypeptide. The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and
30 following the coding region ("leader and trailer") as well as intervening sequences (introns) between individual coding segments (exons).

As used herein, the terms "recombinant" and "engineered" refer to a cell, microorganism, nucleic acid molecule, polypeptide, protein, plasmid, or vector that has been modified by introduction of an exogenous nucleic acid molecule, or refers to a cell or microorganism that has been genetically engineered by human intervention—that is, modified by introduction of of a heterologous nucleic acid molecule, or refers to a cell or microorganism that has been altered such that expression of an endogenous nucleic acid molecule or gene is controlled, deregulated or constitutive, where such alterations or modifications can be introduced by genetic engineering. Human-generated genetic alterations can include, for example, modifications introducing nucleic acid molecules (which may include an expression control element, such as a promoter) encoding one or more proteins or enzymes, or other nucleic acid molecule additions, deletions, substitutions, or other functional disruption of or addition to a cell's genetic material. Exemplary modifications include those in coding regions or functional fragments thereof of heterologous or homologous polypeptides from a reference or parent molecule.

As used herein, "mutation" refers to a change in the sequence of a nucleic acid molecule or polypeptide molecule as compared to a reference or wild-type nucleic acid molecule or polypeptide molecule, respectively. A mutation can result in several different types of change in sequence, including substitution, insertion or deletion of nucleotide(s) or amino acid(s). In certain embodiments, a mutation is a substitution of one or three codons or amino acids, a deletion of one to about 5 codons or amino acids, or a combination thereof.

A "conservative substitution" is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are well known in the art (*see, e.g.*, WO 97/09433 at page 10; Lehninger, Biochemistry, 2nd Edition; Worth Publishers, Inc. NY, NY, pp.71-77, 1975; Lewin, Genes IV, Oxford University Press, NY and Cell Press, Cambridge, MA, p. 8, 1990).

The term "construct" refers to any polynucleotide that contains a recombinant nucleic acid molecule. A "transgene" or "transgene construct" refers to a construct that contains two or more genes operably linked in an arrangement that is not found in nature. The term "operably-linked" (or "operably linked" herein) refers to the

association of two or more nucleic acid molecules on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably-linked with a coding sequence when it can affect the expression of that coding sequence (*i.e.*, the coding sequence is under the transcriptional control of the promoter).

- 5 "Unlinked" means that the associated genetic elements are not closely associated with one another and the function of one does not affect the other. In some embodiments, the genes present in a transgene are operably linked to an expression control sequence (*e.g.*, a promoter).

A construct (*e.g.*, a transgene) can be present in a vector (*e.g.*, a bacterial vector, 10 a viral vector) or can be integrated into a genome. A "vector" is a nucleic acid molecule that is capable of transporting another nucleic acid molecule. Vectors can be, for example, plasmids, cosmids, viruses, a RNA vector or a linear or circular DNA or RNA molecule that can include chromosomal, non-chromosomal, semi-synthetic or synthetic nucleic acid molecules. Exemplary vectors are those capable of autonomous replication 15 (episomal vector) or expression of nucleic acid molecules to which they are linked (expression vectors). Vectors useful in the compositions and methods of this disclosure are described further herein.

The term "expression", as used herein, refers to the process by which a polypeptide is produced based on the encoding sequence of a nucleic acid molecule, 20 such as a gene. The process can include transcription, post-transcriptional control, post-transcriptional modification, translation, post-translational control, post translational modification, or any combination thereof.

The term "introduced" in the context of inserting a nucleic acid molecule into a cell, means "transfection", or "transformation", or "transduction" and includes reference 25 to the incorporation of a nucleic acid molecule into a eukaryotic or prokaryotic cell wherein the nucleic acid molecule can be incorporated into the genome of a cell (*e.g.*, a chromosome, a plasmid, a plastid, or a mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (*e.g.*, transfected mRNA).

As used herein, "heterologous" or "exogenous" nucleic acid molecule, construct 30 or sequence refers to a nucleic acid molecule or portion of a nucleic acid molecule that is not native to a host cell, but can be homologous to a nucleic acid molecule or portion

of a nucleic acid molecule from the host cell. The source of the heterologous or exogenous nucleic acid molecule, construct or sequence can be from a different genus or species. In certain embodiments, a heterologous or exogenous nucleic acid molecule is added (*i.e.*, not endogenous or native) to a host cell or host genome by, for example, 5 conjugation, transformation, transfection, transduction, electroporation, or the like, wherein the added molecule can integrate into the host genome or exist as extra-chromosomal genetic material (e.g., as a plasmid or other form of self-replicating vector), and can be present in multiple copies. In addition, "heterologous" refers to a non-native enzyme, protein or other activity encoded by an exogenous nucleic acid 10 molecule introduced into the host cell, even if the host cell encodes a homologous protein or activity.

As described herein, more than one heterologous or exogenous nucleic acid molecule can be introduced into a host cell as separate nucleic acid molecules, as a plurality of individually controlled genes, as a polycistronic nucleic acid molecule, as a 15 single nucleic acid molecule encoding a fusion protein, or any combination thereof. For example, as disclosed herein, a host cell can be modified to express two or more heterologous or exogenous nucleic acid molecules encoding the desired TCR specific for a minor histocompatibility (H) antigen HA-1^H peptide (*e.g.*, TCR α and TCR β). When two or more exogenous nucleic acid molecules are introduced into a host cell, it 20 is understood that the two or more exogenous nucleic acid molecules can be introduced as a single nucleic acid molecule (*e.g.*, on a single vector), on separate vectors, integrated into the host chromosome at a single site or multiple sites, or any combination thereof. The number of referenced heterologous nucleic acid molecules or protein activities refers to the number of encoding nucleic acid molecules or the number 25 of protein activities, not the number of separate nucleic acid molecules introduced into a host cell.

As used herein, the term "endogenous" or "native" refers to a gene, protein, or activity that is normally present in a host cell. Moreover, a gene, protein or activity that is mutated, overexpressed, shuffled, duplicated or otherwise altered as compared to a 30 parent gene, protein or activity is still considered to be endogenous or native to that particular host cell. For example, an endogenous control sequence from a first gene

(*e.g.*, a promoter, translational attenuation sequences) can be used to alter or regulate expression of a second native gene or nucleic acid molecule, wherein the expression or regulation of the second native gene or nucleic acid molecule differs from normal expression or regulation in a parent cell.

5 The term "homologous" or "homolog" refers to a molecule or activity found in or derived from a host cell, species or strain. For example, a heterologous or exogenous nucleic acid molecule can be homologous to a native host cell gene, and can optionally have an altered expression level, a different sequence, an altered activity, or any combination thereof.

10 "Sequence identity," as used herein, refers to the percentage of amino acid residues in one sequence that are identical with the amino acid residues in another reference polypeptide sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. The percentage sequence
15 identity values can be generated using the NCBI BLAST 2.0 software as defined by Altschul *et al.* (1997), *Nucl. Acids Res.* 25:3389-3402, with the parameters set to default values.

HA-1^H-Specific Binding Proteins, Accessory Proteins, and Engineered Host Cells

20 In certain aspects, the present disclosure provides engineered immune cells comprising a heterologous polynucleotide that encodes a binding protein that specifically binds to an HA-1^H antigen. In certain embodiments, the encoded binding protein is an HA-1^H antigen-specific T cell receptor (TCR) or an HA-1^H antigen-specific chimeric antigen receptor (CAR). In further embodiments, a binding protein is expressed as part of a transgene construct that encodes additional accessory proteins,
25 such as a safety switch protein, a tag, a selection marker, a CD8 co-receptor β -chain, α -chain or both, or any combination thereof.

30 In any of the embodiments described herein, an encoded polypeptide of this disclosure (*e.g.*, iCasp9, TCR β -chain, TCR α -chain, CD8 β -chain, CD8 α -chain) can comprise a "signal peptide" (also known as a leader sequence, leader peptide, or transit peptide). Signal peptides target newly synthesized polypeptides to their appropriate location inside or outside the cell. A signal peptide may be removed from the

polypeptide during or once localization or secretion is completed. Polypeptides that have a signal peptide are referred to herein as a "pre-protein" and polypeptides having their signal peptide removed are referred to herein as "mature" proteins or polypeptides. Representative signal peptides include the amino acids from position 1 to position 21 of
 5 any one of SEQ ID NOS:1-3, 5-9, and 70-75, or the amino acids from position 1 to position 19 of any one of SEQ ID NOS:4, 10, and 12.

Binding proteins of this disclosure, such as TCRs and CARs, will contain a binding domain specific for a target (in this case, HA-1^H). A "binding domain" (also referred to as a "binding region" or "binding moiety"), as used herein, refers to a
 10 molecule or portion thereof (*e.g.*, peptide, oligopeptide, polypeptide, protein) that possesses the ability to specifically and non-covalently associate, unite, or combine with a target (*e.g.*, HA-1^H peptide or HA-1^H peptide:MHC complex). A binding domain includes any naturally occurring, synthetic, semi-synthetic, or recombinantly produced binding partner for a biological molecule, a molecular complex (*i.e.* complex
 15 comprising two or more biological molecules), or other target of interest. Exemplary binding domains include single chain immunoglobulin variable regions (*e.g.*, single chain TCR (scTCR), single chain Fv (scFv)), receptor ectodomains, ligands (*e.g.*, cytokines, chemokines), or synthetic polypeptides selected for their specific ability to bind to a biological molecule, a molecular complex or other target of interest.

20 In certain embodiments, an HA-1^H-specific binding domain alone (*i.e.*, without any other portion of a HA-1-specific binding protein) can be soluble and can bind to HA-1^H with a K_d of less than about 10^{-8} M, less than about 10^{-9} M, less than about 10^{-10} M, less than about 10^{-11} M, less than about 10^{-12} M, or less than about 10^{-13} M. In particular embodiments, an HA-1^H-specific binding domain includes an HA-1^H-specific
 25 scTCR (*e.g.*, single chain $\alpha\beta$ TCR proteins such as $V\alpha$ -L- $V\beta$, $V\beta$ -L- $V\alpha$, $V\alpha$ -C α -L- $V\alpha$, or $V\alpha$ -L- $V\beta$ -C β , wherein $V\alpha$ and $V\beta$ are TCR α and β variable domains respectively, C α and C β are TCR α and β constant domains, respectively, and L is a linker).

The term "variable region" or "variable domain" refers to the domain of an immunoglobulin superfamily binding protein (*e.g.*, a TCR α -chain or β -chain (or γ
 30 chain and δ chain for $\gamma\delta$ TCRs)) that is involved in binding of the immunoglobulin superfamily binding protein (*e.g.*, TCR) to antigen. The variable domains of the

α -chain and β -chain (V_α and V_β , respectively) of a native TCR generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three CDRs. The V_α domain is encoded by two separate DNA segments, the variable gene segment and the joining gene segment (V-J); the V_β domain is encoded by three
 5 separate DNA segments, the variable gene segment, the diversity gene segment, and the joining gene segment (V-D-J). A single V_α or V_β domain may be sufficient to confer antigen-binding specificity. Furthermore, TCRs that bind a particular antigen may be isolated using a V_α or V_β domain from a TCR that binds the antigen to screen a library of complementary V_α or V_β domains, respectively.

10 The terms "complementarity determining region," and "CDR," are synonymous with "hypervariable region" or "HVR," and are known in the art to refer to non-contiguous sequences of amino acids within TCR variable regions, which confer antigen specificity and/or binding affinity. In general, there are three CDRs in each α -chain variable region (α CDR1, α CDR2, α CDR3) and three CDRs in each β -chain
 15 variable region (β CDR1, β CDR2, β CDR3). CDR3 is thought to be the main CDR responsible for recognizing processed antigen. CDR1 and CDR2 mainly interact with the MHC.

In certain embodiments, an encoded binding protein comprises: (a) a T cell receptor (TCR) α chain variable (V_α) domain having an amino acid sequence encoded
 20 by a TRAV17 gene, a TRAV21 gene, or a TRAV10 gene, and a TCR β -chain variable (V_β) domain comprising a CDR3 amino acid sequence as shown in any one of SEQ ID NOS:13-17 and 86; (b) a TCR V_α domain comprising a CDR3 amino acid sequence as shown in any one of SEQ ID NOS:87-92, and a TCR V_β domain having an amino acid sequence encoded by a TRBV7-9 gene; or (c) a TCR V_α domain comprising a CDR3
 25 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V_β domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86, wherein the encoded binding protein is capable of specifically binding to a peptide containing an HA-1^H minor antigen and does not bind to a peptide that does not contain an HA-1^H minor antigen.

30 In further embodiments, an encoded binding protein comprises a TCR V_α domain and a TCR V_β domain, wherein: (a) the encoded V_β CDR3 comprises the amino

acid sequence of SEQ ID NO:13, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:87; (b) the encoded V_{β} CDR3 comprises the amino acid sequence shown in SEQ ID NO:14, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:88; (c) the encoded V_{β} CDR3 comprises the amino acid sequence shown in SEQ ID NO:15, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:89; (d) the encoded V_{β} CDR3 comprises the amino acid sequence shown in SEQ ID NO:16, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:90; (e) the encoded V_{β} CDR3 comprises the amino acid sequence shown in SEQ ID NO:17, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:91; or (f) the encoded V_{β} CDR3 comprises the amino acid sequence shown in SEQ ID NO:86, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:92.

In further embodiments, an encoded binding protein comprises a V_{α} domain, wherein the encoded V_{α} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:2, 4, 6, 8, 10, and 12. In additional embodiments, an encoded binding protein comprises a V_{β} domain, wherein the encoded V_{β} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:1, 3, 5, 7, 9, and 11.

In some embodiments, the encoded V_{α} domain comprises no change in amino acid sequence of CDR1, the encoded V_{β} domain comprises no change in amino acid sequence of CDR1, or the CDR1 of the encoded V_{α} domain and the CDR1 of the encoded V_{β} domain comprise no change in amino acid sequence. In further embodiments, the encoded V_{α} domain comprises no change in amino acid sequence of CDR2, the encoded V_{β} domain comprises no change in amino acid sequence of CDR2, or the CDR2 of the encoded V_{α} domain and the CDR2 of the encoded V_{β} domain comprise no change in amino acid sequence.

In particular embodiments, an encoded binding protein comprises a TCR V_{α} domain and a TCR V_{β} domain, wherein: (a) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:1, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:2; (b) the encoded V_{β}

domain comprises or consists of the amino acid sequence of SEQ ID NO:3, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:4; (c) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:5, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:6; (d) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:7, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:8; (e) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:9, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:10; or (f) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:11, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:12.

Exemplary binding proteins of this disclosure expressed by a cell may include a signal peptide (*e.g.*, binding pre-proteins), and the cell may remove the signal peptide to generate a mature binding protein. In certain embodiments, a binding protein comprises two components, such as an α -chain and a β -chain, which can associate on the cell surface to form a functional binding protein. The two associated components may comprise mature proteins. In certain embodiments, a binding protein of this disclosure comprises a mature V_{β} domain, wherein the mature V_{β} domain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:96, 98, 100, 102, 104, or 106. In further embodiments, a binding protein of this disclosure comprises a mature V_{α} domain, wherein the mature V_{α} domain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:97, 99, 101, 103, 105, or 107. In still further embodiments, a binding protein of this disclosure comprises a mature V_{β} domain and a mature V_{α} domain, wherein the mature V_{β} domain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:96, 98, 100, 102, 104, or 106, and the mature V_{α} domain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:97, 99, 101, 103, 105, or 107. In certain embodiments, a binding protein of this disclosure comprises a mature TCR β -chain, wherein the mature TCR β -chain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:108, 110, 112, 114, 116, or 118. In further embodiments, a binding protein of this disclosure

comprises a mature TCR α -chain, wherein the mature TCR α -chain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:109, 111, 113, 115, 117, or 119. In yet further embodiments, a binding protein of this disclosure comprises a mature TCR β -chain and a mature TCR α -chain, wherein the mature TCR β -chain

5 comprises or consists of the amino acid sequence of any one of SEQ ID NOS:108, 110, 112, 114, 116, or 118, and the mature TCR α -chain comprises or consists of the amino acid sequence of any one of SEQ ID NOS:109, 111, 113, 115, 117, or 119. In certain embodiments, a binding protein of this disclosure is expressed with a CD8 β -chain and the CD8 β -chain comprises a mature CD8 β -chain, wherein the mature CD8 β -chain

10 comprises or consists of the amino acid sequence shown in any one of SEQ ID NOS:121-125. In further embodiments, a binding protein of this disclosure is expressed with a CD8 α -chain and the CD8 α -chain comprises a mature CD8 α -chain, wherein the mature CD8 α -chain comprises or consists of the amino acid sequence of SEQ ID NO:120. In more embodiments, a binding protein of this disclosure is expressed with a

15 CD8 β -chain and a CD8 α -chain, wherein the CD8 β -chain and α -chain comprises a mature CD8 β -chain and α -chain, wherein the mature CD8 β -chain comprises or consists of the amino acid sequence shown in any one of SEQ ID NOS:121-125, and the mature CD8 α -chain comprises or consists of the amino acid sequence of SEQ ID NO:120.

20 In further embodiments, an encoded binding protein comprises a mature TCR V_α domain and a mature TCR V_β domain, wherein: (a) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:96, and the V_α domain comprising or consisting of the amino acid sequence of SEQ ID NO:97; (b) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:98, and the

25 V_α domain comprising or consisting of the amino acid sequence of SEQ ID NO:99; (c) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:100, and the V_α domain comprising or consisting of the amino acid sequence of SEQ ID NO:101; (d) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:102, and the V_α domain comprising or consisting of the amino acid

30 sequence of SEQ ID NO:103; (e) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:104, and the V_α domain comprising or consisting of the

amino acid sequence of SEQ ID NO:105; or (f) the V_β domain comprising or consisting of the amino acid sequence of SEQ ID NO:106, and the V_α domain comprising or consisting of the amino acid sequence of SEQ ID NO:107.

5 An encoded binding protein contained in an engineered immune cell of the present disclosure may, in some embodiments, comprise a TCR constant domain. In certain embodiments, a TCR constant domain is modified to enhance pairing of desired TCR chains. For example, enhanced pairing between a heterologous TCR α -chain and a heterologous TCR β -chain due to a modification results in the preferential assembly of a TCR comprising two heterologous chains over an undesired mispairing of a
10 heterologous TCR chain with an endogenous TCR chain (*see, e.g., Govers et al., Trends Mol. Med. 16(2):77 (2010), the TCR modifications of which are herein incorporated by reference*). Exemplary modifications to enhance pairing of heterologous TCR chains include the introduction of complementary cysteine residues in each of the heterologous TCR α -chain and β -chain. In some embodiments, a
15 polynucleotide encoding a heterologous TCR α -chain encodes a cysteine at amino acid position 48 (corresponding to the full-length, mature human TCR α -chain sequence) and a polynucleotide encoding a heterologous TCR β -chain encodes a cysteine at amino acid position 57 (corresponding to the full-length mature human TCR β -chain sequence).

20 In certain embodiments, the encoded binding protein comprises a TCR α -chain constant (C_α) domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:19, 22, 24 and 26. In further embodiments, the encoded binding protein comprises a TCR C_α domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:19, 22, 24 and
25 26, provided that the TCR C_α domain retains the introduced cysteine residue at position 48. In still further embodiments, the encoded binding protein comprises a TCR C_α domain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:19, 22, 24 and 26.

30 In certain embodiments, the encoded binding protein comprises a TCR β -chain constant (C_β) domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:18, 23 and 25. In further embodiments, the

encoded binding protein comprises a TCR C β domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:18, 23 and 25, provided that the TCR C β domain retains the introduced cysteine residue at position 57. In still further embodiments, the encoded binding protein comprises a TCR C β domain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:18, 23 and 25.

In certain embodiments, an encoded binding protein comprises a TCR α -chain (*e.g.*, a TCR V α domain operatively associated with a TCR C α domain) having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36 and 38, optionally wherein the TCR C α domain retains the cysteine at position 47 (as counted from the beginning of the C α domain). In further embodiments, an encoded binding protein comprises a TCR α -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36 and 38. In other embodiments, an encoded binding protein comprises a TCR β -chain (*e.g.*, a TCR V β domain operatively associated with a TCR C β domain) having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37, optionally wherein the TCR C β domain retains the cysteine at position 57 (as counted from the beginning of the C β domain). In still further embodiments, the encoded binding protein comprises a TCR β -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37.

A binding protein encoded by an engineered immune cell of this disclosure may comprise any of the presently disclosed TCR α -chains in association with any of the disclosed TCR β -chains. For example, in certain embodiments, an encoded binding protein comprises: (a) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:27, and a TCR α -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:28; (b) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:29, and a TCR α -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:30; (c) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:31, and the TCR α -chain comprising or consisting of the amino acid sequence

shown in SEQ ID NO:32; (d) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:33, and a TCR α -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:34; (e) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:35, and a TCR- α chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:36; or (f) a TCR β -chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:37, and a TCR- α chain comprising or consisting of the amino acid sequence shown in SEQ ID NO:38.

An engineered immune cell of the present disclosure may comprise a single polynucleotide that encodes a binding protein as described herein, or the binding protein may be encoded by more than one polynucleotide. In other words, components or portions of a binding protein may be encoded by two or more polynucleotides, which may be contained on a single nucleic acid molecule or may be contained on two or more nucleic acid molecules.

In certain embodiments, a polynucleotide encoding two or more components or portions of a binding protein of the present disclosure comprises the two or more coding sequences operatively associated in a single open reading frame. Such an arrangement can advantageously allow coordinated expression of desired gene products, such as, for example, contemporaneous expression of alpha- and beta-chains of a TCR, such that they are produced in about a 1:1 ratio. In certain embodiments, two or more substituent gene products of a binding protein of this disclosure, such as a TCR (*e.g.*, alpha- and beta-chains) or CAR, are expressed as separate molecules and associate post-translationally. In further embodiments, two or more substituent gene products of a binding protein of this disclosure are expressed as a single peptide with the parts separated by a cleavable or removable segment. For instance, self-cleaving peptides useful for expression of separable polypeptides encoded by a single polynucleotide or vector are known in the art and include, for example, a Porcine teschovirus-1 2A (P2A) peptide, such as a peptide encoded by a polynucleotide having the nucleotide sequence shown in any one of SEQ ID NOS:76-81, a Thoseaasigna virus 2A (T2A) peptide, such as a peptide encoded by a polynucleotide having the nucleotide sequence shown in SEQ ID NO:82, an Equine rhinitis A virus (ERAV) 2A (E2A) peptide, such as a peptide

encoded by a polynucleotide having the nucleotide sequence shown in SEQ ID NO:83, and a Foot-and-Mouth disease virus 2A (F2A) peptide, such as a peptide encoded by a polynucleotide having the nucleotide sequence shown in SEQ ID NO:84.

In certain embodiments, a binding protein of the present disclosure comprises
5 one or more junction amino acids. "Junction amino acids" or "junction amino acid residues" refer to one or more (*e.g.*, 2 to about 10) amino acid residues between two adjacent motifs, regions or domains of a polypeptide, such as between a binding domain and an adjacent constant domain or between a TCR chain and an adjacent self-cleaving peptide. Junction amino acids can result from the design of a construct that encodes a
10 fusion protein (*e.g.*, amino acid residues resulting from the use of a restriction enzyme site during the construction of a nucleic acid molecule encoding a fusion protein), or from cleavage of, for example, a self-cleaving peptide adjacent one or more domains of an encoded binding protein of this disclosure (*e.g.*, a P2A peptide disposed between a TCR α -chain and a TCR β -chain, the self-cleavage of which can leave one or more
15 junction amino acids in the α -chain, the TCR β -chain, or both).

Binding proteins contained in engineered immune cells of this disclosure can, in certain embodiments, specifically bind to an HA-1^H peptide:HLA complex. For example, in specific embodiments, a binding protein of this disclosure is capable of specifically binding to an HA-1^H peptide:HLA complex, wherein the HLA can
20 comprise HLA-A*0201. In particular embodiments, the HA-1^H peptide comprises the amino acid sequence VLHDDLLEA (SEQ ID NO:66).

In any of the aforementioned embodiments, an encoded binding protein contained in an engineered immune cell can comprise a TCR, an antigen-binding fragment of a TCR (*e.g.*, a single chain TCR ("scTCR")), or a chimeric antigen receptor
25 ("CAR").

In certain embodiments, an antigen-binding fragment of a TCR comprises a single chain TCR (scTCR), which comprises both the TCR V α and TCR V β domains, but only a single TCR constant domain (C α or C β). In further embodiments, an antigen-binding fragment of a TCR or a chimeric antigen receptor is chimeric (*e.g.*, comprises
30 amino acid residues or motifs from more than one donor or species), humanized (*e.g.*,

comprises residues from a non-human organism that are altered or substituted so as to reduce the risk of immunogenicity in a human), or human.

"Chimeric antigen receptor" (CAR) refers to a fusion protein that is engineered to contain two or more naturally-occurring amino acid sequences linked together in a way that does not occur naturally or does not occur naturally in a host cell, which fusion protein can function as a receptor when present on a surface of a cell. CARs of the present disclosure include an extracellular portion comprising an antigen binding domain (i.e., obtained or derived from an immunoglobulin or immunoglobulin-like molecule, such as an scFv derived from an antibody or TCR specific for a cancer antigen, or an antigen binding domain derived or obtained from a killer immunoreceptor from an NK cell) linked to a transmembrane domain and one or more intracellular signaling domains (optionally containing co-stimulatory domain(s)) (*see, e.g., Sadelain et al., Cancer Discov., 3(4):388 (2013); see also Harris and Kranz, Trends Pharmacol. Sci., 37(3):220 (2016), and Stone et al., Cancer Immunol. Immunother., 63(11):1163 (2014).*)

Methods for producing engineered TCRs are described in, for example, Bowerman *et al., Mol. Immunol., 46(15):3000 (2009)*, the techniques of which are herein incorporated by reference. Methods for making CARs are well known in the art and are described, for example, in U.S. Patent No. 6,410,319; U.S. Patent No. 7,446,191; U.S. Patent Publication No. 2010/065818; U.S. Patent No. 8,822,647; PCT Publication No. WO 2014/031687; U.S. Patent No. 7,514,537; and Brentjens *et al., 2007, Clin. Cancer Res. 13:5426*, the techniques of which are herein incorporated by reference.

Engineered immune cells of this disclosure can be administered as therapies for, *e.g., cancer*. In some circumstances, it may be desirable to reduce or stop the activity associated with a cellular immunotherapy. Thus, in certain embodiments, an engineered immune cell of the present disclosure comprises a heterologous polynucleotide encoding a binding protein and an accessory protein, such as a safety switch protein, which can be targeted using a cognate drug or other compound to selectively modulate the activity (*e.g., lessen or ablate*) of such cells when desirable. Safety switch proteins used in this regard include, for example, a truncated EGF

receptor polypeptide (huEGFRt) that is devoid of extracellular N-terminal ligand binding domains and intracellular receptor tyrosine kinase activity but retains the native amino acid sequence, type I transmembrane cell surface localization, and a conformationally intact binding epitope for pharmaceutical-grade anti-EGFR

5 monoclonal antibody, cetuximab (Erbitux) tEGF receptor (tEGFr; Wang *et al.*, *Blood* 118:1255-1263, 2011), a caspase polypeptide (*e.g.*, iCasp9; Straathof *et al.*, *Blood* 105:4247-4254, 2005; Di Stasi *et al.*, *N. Engl. J. Med.* 365:1673-1683, 2011; Zhou and Brenner, *Exp. Hematol.* pii:S0301-472X(16)30513-6. doi:10.1016/j.exphem.2016.07.011), RQR8 (Philip *et al.*, *Blood* 124:1277-1287, 2014),

10 a 10 amino acid tag of the human c-myc protein (Myc) (Kieback *et al.*, *Proc. Natl. Acad. Sci. USA* 105:623-628, 2008), as discussed herein, and a marker/safety switch polypeptide, such as RQR (CD20 + CD34; Philip *et al.*, 2014).

Other accessory components useful for therapeutic cells comprise a tag or selection marker that allows the cells to be identified, sorted, isolated, enriched, or

15 tracked. For example, marked immune cells having desired characteristics (*e.g.*, an antigen-specific TCR and a safety switch protein) can be sorted away from unmarked cells in a sample and more efficiently activated and expanded for inclusion in a therapeutic product of desired purity.

As used herein, the term "selection marker" comprises a nucleic acid construct

20 that confers an identifiable change to a cell permitting detection and positive selection of immune cells transduced with a polynucleotide comprising a selection marker. RQR is a selection marker that comprises a major extracellular loop of CD20 and two minimal CD34 binding sites. In some embodiments, an RQR-encoding polynucleotide comprises a polynucleotide that encodes the 16 amino acid CD34 minimal epitope. In

25 some embodiments, such as certain embodiments provided in the examples herein, the CD34 minimal epitope is incorporated at the amino terminal position of the CD8 stalk domain (Q8). In further embodiments, the CD34 minimal binding site sequence can be combined with a target epitope for CD20 to form a compact marker/suicide gene for T cells (RQR8) (Philip *et al.*, 2014, incorporated by reference herein). This construct

30 allows for the selection of immune cells expressing the construct, with for example, CD34 specific antibody bound to magnetic beads (Miltenyi) and that utilizes clinically

accepted pharmaceutical antibody, rituximab, that allows for the selective deletion of a transgene expressing engineered T cell (Philip *et al.*, 2014).

Further exemplary selection markers also include several truncated type I transmembrane proteins normally not expressed on T cells: the truncated low-affinity nerve growth factor, truncated CD19, and truncated CD34 (see for example, Di Stasi *et al.*, *N. Engl. J. Med.* 365:1673-1683, 2011; Mavilio *et al.*, *Blood* 83:1988-1997, 1994; Fehse *et al.*, *Mol. Ther.* 1:448-456, 2000; each incorporated herein in their entirety). A particularly attractive feature of CD19 and CD34 is the availability of the off-the-shelf Miltenyi CliniMACs™ selection system that can target these markers for clinical-grade sorting. However, CD19 and CD34 are relatively large surface proteins that may tax the vector packaging capacity and transcriptional efficiency of an integrating vector. Surface markers containing the extracellular, non-signaling domains or various proteins (e.g., CD19, CD34, LNGFR) also can be employed. Any selection marker may be employed and should be acceptable for Good Manufacturing Practices. In certain embodiments, selection markers are expressed with a polynucleotide that encodes a gene product of interest (e.g., a binding protein of the present disclosure, such as a TCR or CAR). Further examples of selection markers include, for example, reporters such as GFP, EGFP, β -gal or chloramphenicol acetyltransferase (CAT). In certain embodiments, a selection marker, such as, for example, CD34 is expressed by a cell and the CD34 can be used to select enrich for, or isolate (e.g., by immunomagnetic selection) the transduced cells of interest for use in the methods described herein. As used herein, a CD34 marker is distinguished from an anti-CD34 antibody, or, for example, a scFv, TCR, or other antigen recognition moiety that binds to CD34.

In certain embodiments, a selection marker comprises an RQR polypeptide, a truncated low-affinity nerve growth factor (tNGFR), a truncated CD19 (tCD19), a truncated CD34 (tCD34), or any combination thereof.

By way of background, inclusion of CD4⁺ T cells in an immunotherapy cell product can provide antigen-induced IL-2 secretion and augment persistence and function of transferred cytotoxic CD8⁺ T cells (see, e.g., Kennedy *et al.*, *Immunol. Rev.* 222:129 (2008); Nakanishi *et al.*, *Nature* 462(7272):510 (2009)). In certain circumstances, a class I restricted TCR in CD4⁺ T cells may require the transfer of a

CD8 co-receptor to enhance sensitivity of the TCR to class I HLA peptide complexes. CD4 co-receptors differ in structure to CD8 and cannot effectively substitute for CD8 co-receptors (*see, e.g., Stone & Kranz, Front. Immunol. 4:244 (2013); see also Cole et al., Immunology 137(2):139 (2012).* Thus, another accessory protein for use in the compositions and methods of this disclosure comprises a CD8 co-receptor or component thereof.

Engineered immune cells comprising a heterologous polynucleotide encoding a binding protein of the present disclosure may, in certain embodiments, further comprise a heterologous polynucleotide encoding a CD8 co-receptor protein, or a beta-chain or alpha-chain component thereof. An encoded CD8 co-receptor includes, in some embodiments, a β -chain comprising the amino acid sequence of any one of SEQ ID NOS:71-75. In further embodiments, the encoded CD8 co-receptor is a recombinant CD8 co-receptor further comprising a RQR polypeptide having the amino acid sequence of SEQ ID NO:69. Without wishing to be bound by theory, it is believed that distance from the host cell surface is important for RQR polypeptides to function as selection markers/safety switches (Philip *et al.*, 2010 (*supra*)). In some embodiments, the encoded RQR polypeptide is contained in a β -chain, an α -chain, or both, of the encoded CD8 co-receptor. In specific embodiments, an engineered immune cell comprises a heterologous polynucleotide encoding iCasp9 and a heterologous polynucleotide encoding a recombinant CD8 co-receptor protein that comprises a β -chain containing a RQR polypeptide and further comprises a CD8 α -chain. In particular embodiments, the encoded CD8 α -chain comprises the amino acid sequence shown in SEQ ID NO:70.

In further embodiments, an engineered immune cell comprises a heterologous polynucleotide encoding iCasp9 and a heterologous polynucleotide encoding a recombinant CD8 co-receptor protein that comprises an α -chain containing a RQR polypeptide and further comprises a CD8 β -chain. In some embodiments, both of the encoded CD8 α -chain and the encoded CD8 β -chain contain a RQR polypeptide.

An engineered immune cell may be efficiently transduced to contain, and may efficiently express, a single polynucleotide that encodes the binding protein, safety switch protein, selection marker, and CD8 co-receptor protein. For example, in some

embodiments, an engineered immune cell of the present disclosure comprises a heterologous polynucleotide that encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a porcine teschovirus 2A (P2A) peptide]-[a TCR β -chain]-[a P2A peptide]-[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]). In specific embodiments, the TCR β -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:41, and the TCR α -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:42.

In particular embodiments, an engineered immune cell contains a heterologous polynucleotide that comprises or consists of the nucleotide sequence of SEQ ID NO:85.

Any suitable immune cell may be engineered to include a heterologous polynucleotide encoding a binding protein of this disclosure, including, for example, a T cell, a NK cell, or a NK-T cell. In some embodiments, an engineered immune cell comprises a CD4⁺T cell, a CD8⁺ T cell, or both. Methods for transfecting/transducing T cells with desired nucleic acids have been described (*e.g.*, U.S. Patent Application Pub. No. US 2004/0087025) as have adoptive transfer procedures using T cells of desired target-specificity (*e.g.*, Schmitt *et al.*, *Hum. Gen.* 20:1240, 2009; Dossett *et al.*, *Mol. Ther.* 17:742, 2009; Till *et al.*, *Blood* 112:2261, 2008; Wang *et al.*, *Hum. Gene Ther.* 18:712, 2007; Kuball *et al.*, *Blood* 109:2331, 2007; US 2011/0243972; US 2011/0189141; Leen *et al.*, *Ann. Rev. Immunol.* 25:243, 2007), such that adaptation of these methodologies to the presently disclosed embodiments is contemplated, based on the teachings herein.

Any appropriate method can be used to transfect or transduce the cells, for example, the T cells, or to administer the polynucleotides or compositions of the present methods. Known methods for delivering polynucleotides to host cells include, for example, use of cationic polymers, lipid-like molecules, and certain commercial products such as, for example, IN-VIVO-JET PEI. Other methods include *ex vivo* transduction, injection, electroporation, DEAE-dextran, sonication loading, liposome-mediated transfection, receptor-mediated transduction, microprojectile bombardment, transposon-mediated transfer, and the like. Still further methods of transfecting or transducing host cells employ vectors, described in further detail herein.

In any of the foregoing embodiments, an engineered immune cell may be a "universal donor" cell that is modified to reduce or eliminate expression of one or more endogenous genes that encode a polypeptide involved in immune signaling or other related activities. Exemplary gene knockouts include those that encode PD-1, LAG-3, CTLA4, TIM3, an HLA molecule, a TCR molecule, or the like. Without wishing to be bound by theory, certain endogenously expressed immune cell proteins may be recognized as foreign by an allogeneic host receiving the engineered immune cells, which may result in elimination of the engineered immune cells (*e.g.*, an HLA allele), or may downregulate the immune activity of the engineered immune cells (*e.g.*, PD-1, LAG-3, CTLA4), or may interfere with the binding activity of a heterologously expressed binding protein of the present disclosure (*e.g.*, an endogenous TCR that binds a non-HA-1^H antigen and thereby interferes with the engineered immune cell binding a cell that expresses HA-1^H antigen). Accordingly, decreasing or eliminating expression or activity of such endogenous genes or proteins can improve the activity, tolerance, and persistence of the engineered immune cells within an allogeneic host, and allows for universal, "off-the-shelf" cells for administration (*e.g.*, to any recipient regardless of HLA type).

In certain embodiments, an engineered immune cell of this disclosure comprises a chromosomal gene knockout of one or more of a gene that encodes PD-1, LAG-3, CTLA4, TIM3, an HLA component (*e.g.*, a gene that encodes an α 1 macroglobulin, an α 2 macroglobulin, an α 3 macroglobulin, a β 1 microglobulin, or a β 2 microglobulin), or a TCR component (*e.g.*, a gene that encodes a TCR variable region or a TCR constant region) (*see, e.g.*, Torikai *et al.*, *Nature Sci. Rep.* 6:21757 (2016); Torikai *et al.*, *Blood* 119(24):5697 (2012); and Torikai *et al.*, *Blood* 122(8):1341 (2013), the gene editing techniques and compositions of which are herein incorporated by reference in their entirety). As used herein, the term "chromosomal gene knockout" refers to a genetic alteration in an engineered immune cell that prevents production, by the engineered immune cell, of a functionally active endogenous polypeptide product. Alterations resulting in a chromosomal gene knockout can include, for example, introduced nonsense mutations (including the formation of premature stop codons), missense mutations, gene deletion, and strand breaks, as well as the heterologous expression of

inhibitory nucleic acid molecules that inhibit endogenous gene expression in the engineered immune cell.

A chromosomal gene knockout may be introduced by chromosomal editing of the immune cell. In certain embodiments, the chromosomal gene knockout is made by chromosomal editing of the immune cell. Chromosomal editing can be performed using, for example, endonucleases. As used herein "endonuclease" refers to an enzyme capable of catalyzing cleavage of a phosphodiester bond within a polynucleotide chain. In certain embodiments, an endonuclease is capable of cleaving a targeted gene thereby inactivating or "knocking out" the targeted gene. An endonuclease may be a naturally occurring, recombinant, genetically modified, or fusion endonuclease. The nucleic acid strand breaks caused by the endonuclease are commonly repaired through the distinct mechanisms of homologous recombination or non-homologous end joining (NHEJ). During homologous recombination, a donor nucleic acid molecule may be used for gene "knock-in" to inactivate a target gene. NHEJ is an error-prone repair process that often results in changes to the DNA sequence at the site of the cleavage, *e.g.*, a substitution, deletion, or addition of at least one nucleotide. NHEJ may be used to "knock-out" a target gene. Methods of disrupting or knocking out genes or gene expression in immune cells using endonucleases are known in the art and described, for example, in PCT Publication Nos. WO 2015/066262; WO 2013/074916; and WO 2014/059173; methods from each of which is incorporated by reference. Examples of endonucleases include zinc finger nucleases, TALE-nucleases, CRISPR-Cas nucleases, and meganucleases.

As used herein, a "zinc finger nuclease" (ZFN) refers to a fusion protein comprising a zinc finger DNA-binding domain fused to a non-specific DNA cleavage domain, such as a FokI endonuclease. Each zinc finger motif of about 30 amino acids binds to about 3 base pairs of DNA, and amino acids at certain residues can be changed to alter triplet sequence specificity (*see, e.g., Desjarlais et al., Proc. Natl. Acad. Sci. 90:2256-2260, 1993; Wolfe et al., J. Mol. Biol. 285:1917-1934, 1999*). Multiple zinc finger motifs can be linked in tandem to create binding specificity to desired DNA sequences, such as regions having a length ranging from about 9 to about 18 base pairs. By way of background, ZFNs mediate genome editing by catalyzing the formation of a

site-specific DNA double strand break (DSB) in the genome, and targeted integration of a transgene comprising flanking sequences homologous to the genome at the site of DSB is facilitated by homology directed repair. Alternatively, a DSB generated by a ZFN can result in knock out of target gene via repair by non-homologous end joining (NHEJ), which is an error-prone cellular repair pathway that results in the insertion or deletion of nucleotides at the cleavage site. In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, made using a ZFN molecule.

As used herein, a "transcription activator-like effector nuclease" (TALEN) refers to a fusion protein comprising a TALE DNA-binding domain and a DNA cleavage domain, such as a FokI endonuclease. A "TALE DNA binding domain" or "TALE" is composed of one or more TALE repeat domains/units, each generally having a highly conserved 33-35 amino acid sequence with divergent 12th and 13th amino acids. The TALE repeat domains are involved in binding of the TALE to a target DNA sequence. The divergent amino acid residues, referred to as the Repeat Variable Diresidue (RVD), correlate with specific nucleotide recognition. The natural (canonical) code for DNA recognition of these TALEs has been determined such that an HD sequence at positions 12 and 13 leads to a binding to cytosine (C), NG binds to T, NI to A, NN binds to G or A, and NG binds to T and non-canonical (atypical) RVDs are also known (*see, e.g.*, U.S. Patent Publication No. US 2011/0301073, which atypical RVDs are incorporated by reference herein in its entirety). TALENs can be used to direct site-specific double-strand breaks (DSB) in the genome of T cells. Non-homologous end joining (NHEJ) ligates DNA from both sides of a double-strand break in which there is little or no sequence overlap for annealing, thereby introducing errors that knock out gene expression. Alternatively, homology directed repair can introduce a transgene at the site of DSB providing homologous flanking sequences are present in the transgene. In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, and made using a TALEN molecule.

As used herein, a "clustered regularly interspaced short palindromic repeats/Cas" (CRISPR/Cas) nuclease system refers to a system that employs a CRISPR RNA (crRNA)-guided Cas nuclease to recognize target sites within a genome (known

as protospacers) via base-pairing complementarity and then to cleave the DNA if a short, conserved protospacer associated motif (PAM) immediately follows 3' of the complementary target sequence. CRISPR/Cas systems are classified into three types (*i.e.*, type I, type II, and type III) based on the sequence and structure of the Cas nucleases. The crRNA-guided surveillance complexes in types I and III need multiple Cas subunits. Type II system, the most studied, comprises at least three components: an RNA-guided Cas9 nuclease, a crRNA, and a *trans*-acting crRNA (tracrRNA). The tracrRNA comprises a duplex forming region. A crRNA and a tracrRNA form a duplex that is capable of interacting with a Cas9 nuclease and guiding the Cas9/crRNA:tracrRNA complex to a specific site on the target DNA via Watson-Crick base-pairing between the spacer on the crRNA and the protospacer on the target DNA upstream from a PAM. Cas9 nuclease cleaves a double-stranded break within a region defined by the crRNA spacer. Repair by NHEJ results in insertions and/or deletions which disrupt expression of the targeted locus. Alternatively, a transgene with homologous flanking sequences can be introduced at the site of DSB via homology directed repair. The crRNA and tracrRNA can be engineered into a single guide RNA (sgRNA or gRNA) (*see, e.g.*, Jinek *et al.*, *Science* 337:816-21, 2012). Further, the region of the guide RNA complementary to the target site can be altered or programmed to target a desired sequence (Xie *et al.*, *PLOS One* 9:e100448, 2014; U.S. Pat. Appl. Pub. No. US 2014/0068797, U.S. Pat. Appl. Pub. No. US 2014/0186843; U.S. Pat. No. 8,697,359, and PCT Publication No. WO 2015/071474; the techniques and compositions of each of which are incorporated by reference). In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, and made using a CRISPR/Cas nuclease system.

As used herein, a "meganuclease," also referred to as a "homing endonuclease," refers to an endodeoxyribonuclease characterized by a large recognition site (double stranded DNA sequences of about 12 to about 40 base pairs). Meganucleases can be divided into five families based on sequence and structure motifs: LAGLIDADG, GIY-YIG, HNH, His-Cys box and PD-(D/E)XK. Exemplary meganucleases include I-SceI, I-CeuI, PI-PspI, PI-Sce, I-SceIV, I-CsmI, I-PanI, I-SceII, I-PpoI, I-SceIII, I-CreI, I-TevI, I-TevII and I-TevIII, whose recognition sequences are known (*see, e.g.*, U.S.

Patent Nos. 5,420,032 and 6,833,252; Belfort *et al.*, *Nucleic Acids Res.* 25:3379-3388, 1997; Dujon *et al.*, *Gene* 82:115-118, 1989; Perler *et al.*, *Nucleic Acids Res.* 22:1125-1127, 1994; Jasin, *Trends Genet.* 12:224-228, 1996; Gimble *et al.*, *J. Mol. Biol.* 263:163-180, 1996; Argast *et al.*, *J. Mol. Biol.* 280:345-353, 1998).

5 In certain embodiments, naturally-occurring meganucleases may be used to promote site-specific genome modification of a target selected from PD-1, LAG3, TIM3, CTLA4, an HLA-encoding gene, or a TCR component-encoding gene. In other embodiments, an engineered meganuclease having a novel binding specificity for a target gene is used for site-specific genome modification (*see, e.g.*, Porteus *et al.*, *Nat.*
10 *Biotechnol.* 23:967-73, 2005; Sussman *et al.*, *J. Mol. Biol.* 342:31-41, 2004; Epinat *et al.*, *Nucleic Acids Res.* 31:2952-62, 2003; Chevalier *et al.*, *Molec. Cell* 10:895-905, 2002; Ashworth *et al.*, *Nature* 441:656-659, 2006; Paques *et al.*, *Curr. Gene Ther.* 7:49-66, 2007; U.S. Patent Publication Nos. US 2007/0117128; US 2006/0206949; US 2006/0153826; US 2006/0078552; and US 2004/0002092).

15 In certain embodiments, a chromosomal gene knockout comprises an inhibitory nucleic acid molecule that is introduced into an engineered immune cell comprising a heterologous polynucleotide encoding an antigen-specific receptor that specifically binds to a tumor associated antigen, wherein the inhibitory nucleic acid molecule encodes a target-specific inhibitor and wherein the encoded target-specific inhibitor
20 inhibits endogenous gene expression (*i.e.*, of PD-1, TIM3, LAG3, CTLA4, an HLA component, a TCR component, or any combination thereof) in the engineered immune cell.

A chromosomal gene knockout can be confirmed directly by DNA sequencing of the engineered immune cell following use of the knockout procedure or agent.
25 Chromosomal gene knockouts can also be inferred from the absence of gene expression (*e.g.*, the absence of an mRNA or polypeptide product encoded by the gene) following the knockout.

In another aspect, compositions are provided herein that comprise an engineered immune cell of the present disclosure and a pharmaceutically acceptable carrier,
30 diluent, or excipient. Also provided herein are unit doses that comprise an effective amount of an engineered immune cell or of a composition comprising the engineered

immune cell. In certain embodiments, a unit dose comprises (i) a composition comprising at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% engineered CD4⁺ T cells, combined with (ii) a composition

5 comprising at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells (*i.e.*, has less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about

10 10%, less than about 5%, or less than about 1% the population of naïve T cells present in a unit dose as compared to a patient sample having a comparable number of PBMCs).

In some embodiments, a unit dose comprises (i) a composition comprising at least about 50% engineered CD4⁺ T cells, combined with (ii) a composition comprising

15 at least about 50% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In further embodiments, a unit dose comprises (i) a composition comprising at least about 60% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 60% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or

20 substantially no naïve T cells. In still further embodiments, a unit dose comprises (i) a composition comprising at least about 70% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 70% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells.

In some embodiments, a unit dose comprises (i) a composition comprising at least about

25 80% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 80% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In some embodiments, a unit dose comprises (i) a composition comprising at least about 85% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 85% engineered CD8⁺ T

30 cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In some embodiments, a unit dose comprises (i) a

composition comprising at least about 90% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 90% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells.

In any of the embodiments described herein, a unit dose comprises equal, or
 5 approximately equal numbers of engineered CD45RA⁻ CD3⁺ CD8⁺ and engineered CD45RA⁻ CD3⁺ CD4⁺ T_M cells.

Polynucleotides, Transgenes and Vectors

In further aspects, the present disclosure provides an isolated polynucleotide that encodes a binding protein as described herein (*e.g.*, an HA-1^H-specific TCR, scTCR, or
 10 CAR that comprises TCR V_α and V_β domains as described herein (and optionally further comprises constant domains or other components as described herein)), and may additionally encoded a safety switch protein, a selection marker, a CD8 co-receptor β-chain, or a CD8 co-receptor α-chain, or any combination thereof, provided that at least a portion of the isolated polynucleotide is codon-optimized for expression in a host cell
 15 (*e.g.*, an engineered immune cell as disclosed herein).

In particular, any of the aforementioned heterologous polynucleotides comprised in the engineered immune cells (*e.g.*, encoding any of the binding proteins of the present disclosure) may also or alternatively be provided in an isolated form, wherein the polynucleotide is codon-optimized for expression in a host cell. For
 20 example, in certain embodiments, an isolated polynucleotide encodes a TCR β-chain of an HA-1^H-specific binding protein and comprises or consists of the nucleotide sequence of any one of SEQ ID NOS:39, 41, 43, 45, 47, 49, or 51. In further embodiments, an isolated polynucleotide encodes a TCR α-chain of an HA-1^H-specific binding protein and comprises or consists of the nucleotide sequence of any one of SEQ ID NOS:40,
 25 42, 46, 48, 50, or 52.

In certain embodiments, a heterologous polynucleotide encoding a TCR α-chain and a heterologous polynucleotide encoding a TCR β-chain are contained in a single open reading frame comprised in the engineered immune cell, wherein the single open reading frame further comprises a polynucleotide encoding a self-cleaving peptide
 30 disposed between the α-chain-encoding polynucleotide and the β-chain-encoding polynucleotide. In some embodiments, the polynucleotide encoding the self-cleaving

peptide comprises or consists of the nucleotide sequence of any one of SEQ ID NOS:76-84.

In further embodiments, the single open reading frame comprises a nucleotide sequence that is at least about 80% identical to the nucleotide sequence of any one of SEQ ID NOS:59-63. In specific embodiments, the single open reading frame comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 59-63. In still further embodiments, the encoded ([TCR β -chain]-[self-cleaving-peptide]-[TCR α -chain]) comprises or consists of the amino acid sequence of any one of SEQ ID NOS: 53-57, which exists before the cell removes the signal peptide, and before the β -chain and α -chains are separated by the self cleaving peptide.

An isolated polynucleotide of this disclosure may further comprise a polynucleotide encoding a safety switch protein, a selection marker, a CD8 co-receptor beta chain (*e.g.*, SEQ ID NOS:71-75), or a CD8 co-receptor alpha chain (*e.g.*, SEQ ID NO:70) as disclosed herein, or may comprise a polynucleotide encoding any combination thereof. In specific embodiments, an isolated comprising a heterologous polynucleotide encoding iCasp9 and a heterologous polynucleotide encoding a recombinant CD8 co-receptor protein that comprises a β -chain or α -chain that contains a RQR polypeptide.

In some embodiments, an isolated polynucleotide comprises a single open reading frame containing, from 5' to 3', ([a polynucleotide encoding a safety switch protein]-[a polynucleotide encoding a self-cleaving peptide]-[the polynucleotide encoding a TCR β -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a TCR α -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a CD8 α -chain]).

In further embodiments, an isolated polynucleotide comprises a single open reading frame that encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a porcine teschovirus 2A (P2A) peptide]-[a TCR β chain]-[a P2A peptide]-[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]). In certain embodiments, the TCR β -chain-encoding polynucleotide

comprises or consists of the nucleotide sequence of SEQ ID NO:41, and wherein the TCR α -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:42.

In specific embodiments, an isolated polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:85.

In any of the embodiments described herein, an isolated polynucleotide is codon-optimized for expression in an immune cell, such as a T cell.

In another aspect, transgene constructs are provided herein, wherein a transgene construct comprises an expression control sequence (*e.g.*, a promoter sequence) operatively linked to a single open reading frame comprising (a) a polynucleotide encoding a safety switch protein; (b) a polynucleotide encoding a TCR β -chain; (c) a polynucleotide encoding a TCR α -chain; (b) a polynucleotide encoding a selection marker; (c) a polynucleotide encoding a CD8 co-receptor β -chain; and (d) a polynucleotide encoding a CD8 co-receptor α -chain.

Construction of an transgene construct for genetically engineering and producing a polypeptide of interest can be accomplished by using any suitable molecular biology engineering technique known in the art. To obtain efficient transcription and translation, a polynucleotide in each transgene construct of the present disclosure includes, in certain embodiments, at least one appropriate expression control sequence (also called a regulatory sequence), such as a leader sequence and particularly a promoter operably (*i.e.*, operatively) linked to the nucleotide sequence encoding the polypeptide of interest. In certain embodiments, a transgene construct comprises a polynucleotide that encodes a safety switch protein, wherein the encoded safety switch protein comprises: (i) a truncated EGF receptor (tEGFR); (ii) iCasp9; (iii) a RQR polypeptide; (iv) a myc epitope; or (v) any combination thereof.

In further embodiments, the encoded selection marker comprises: (i) a RQR polypeptide; (ii) a truncated low-affinity nerve growth factor (tNGFR); (iii) a truncated CD19 (tCD19); (iv) a truncated CD34 (tCD34); or (v) any combination thereof.

In some embodiments, the encoded CD8 co-receptor is a recombinant CD8 co-receptor comprising a RQR polypeptide having the amino acid sequence shown in SEQ ID NO:69. In particular embodiments, a transgene construct includes a polynucleotide

that encodes an RQR polypeptide that is contained in an encoded CD8 β -chain. In further embodiments, a transgene construct includes a polynucleotide that encodes an RQR polypeptide that is contained in an encoded CD8 α -chain.

For example, a transgene construct of the present disclosure comprises, in certain embodiments, an open reading frame containing (a) a polynucleotide encoding a safety switch protein; (b) a polynucleotide encoding a TCR β -chain; (c) a polynucleotide encoding a TCR α -chain; (d) a polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide; and (e) a polynucleotide encoding a CD8 α -chain. Any arrangement of the component polynucleotides is contemplated herein, including, for example, a single open reading frame that comprises, from 5' to 3', ([the polynucleotide encoding a safety switch protein]-[a polynucleotide encoding a self-cleaving peptide]-[the polynucleotide encoding a TCR β -chain]-[the polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a TCR α -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide]-[a polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a CD8 α -chain]).

In specific embodiments, a transgene construct of the instant disclosure comprises a single open reading frame that encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a P2A peptide]-[a TCR β -chain]-[a P2A peptide]-[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]).

In further embodiments, a transgene construct can comprise an expression control sequence operatively linked to a polynucleotide as described herein. For example, a transgene construct can comprise an expression control sequence operatively linked to a polynucleotide that encodes a binding protein of the present disclosure, wherein the binding protein includes (a) a T cell receptor (TCR) α chain variable (V_α) domain having an amino acid sequence encoded by a TRAV17 gene, a TRAV21 gene, or a TRAV10 gene, and a TCR β -chain variable (V_β) domain comprising a CDR3 amino acid sequence as shown in any one of SEQ ID NOS:13-17 and 86; (b) a TCR V_α domain comprising a CDR3 amino acid sequence as shown in any one of SEQ ID NOS:87-92, and a TCR V_β domain having an amino acid sequence

encoded by a TRBV7-9 gene; or (c) a TCR V α domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V β domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86, wherein the encoded binding protein is capable of specifically binding to a peptide containing an HA-1^H antigen and does not bind to a peptide that does not contain an HA-1^H antigen.

Also provided herein are vectors that comprise a transgene construct of the instant disclosure. Some examples of vectors include plasmids, viral vectors, cosmids, and others. Some vectors may be capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors), whereas other vectors may be integrated into the genome of a host cell or promote integration of the polynucleotide insert upon introduction into the host cell and thereby replicate along with the host genome (*e.g.*, lentiviral vector, retroviral vector). Additionally, some vectors are capable of directing the expression of genes to which they are operatively linked (these vectors may be referred to as "expression vectors"). According to related embodiments, it is further understood that, if one or more agents (*e.g.*, polynucleotides encoding binding proteins as described herein) are co administered to a subject, that each agent may reside in separate or the same vectors, and multiple vectors (each containing a different agent or the same agent) may be introduced to a cell or cell population or administered to a subject.

In certain embodiments, polynucleotides of the present disclosure may be operatively linked to certain elements of a vector. For example, polynucleotide sequences that are needed to effect the expression and processing of coding sequences to which they are ligated may be operatively linked. Expression control sequences may include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (*i.e.*, Kozak consensus sequences); sequences that enhance protein stability; and possibly sequences that enhance protein secretion. Expression control sequences may be operatively linked if they are contiguous with the gene of interest and

expression control sequences that act in *trans* or at a distance to control the gene of interest.

In certain embodiments, the vector comprises a plasmid vector or a viral vector (*e.g.*, a vector selected from lentiviral vector or a γ -retroviral vector). Viral vectors
5 include retrovirus, adenovirus, parvovirus (*e.g.*, adeno-associated viruses), coronavirus, negative strand RNA viruses such as ortho-myxovirus (*e.g.*, influenza virus), rhabdovirus (*e.g.*, rabies and vesicular stomatitis virus), paramyxovirus (*e.g.*, measles and Sendai), positive strand RNA viruses such as picornavirus and alphavirus, and double-stranded DNA viruses including adenovirus, herpesvirus (*e.g.*, Herpes Simplex
10 virus types 1 and 2, Epstein-Barr virus, cytomegalovirus), and poxvirus (*e.g.*, vaccinia, fowlpox and canarypox). Other viruses include Norwalk virus, togavirus, flavivirus, reoviruses, papovavirus, hepadnavirus, and hepatitis virus, for example. Examples of retroviruses include avian leukosis-sarcoma, mammalian C-type, B-type viruses, D type viruses, HTLV-BLV group, lentivirus, and spumavirus (Coffin, J. M., *Retroviridae: The*
15 *viruses and their replication*, In *Fundamental Virology*, Third Edition, B. N. Fields et al., Eds., Lippincott-Raven Publishers, Philadelphia, 1996).

"Retroviruses" are viruses having an RNA genome, which is reverse-transcribed into DNA using a reverse transcriptase enzyme, the reverse-transcribed DNA is then incorporated into the host cell genome. "Gammaretrovirus" refers to a genus of the
20 retroviridae family. Examples of gammaretroviruses include mouse stem cell virus, murine leukemia virus, feline leukemia virus, feline sarcoma virus, and avian reticuloendotheliosis viruses. "Lentiviral vector," as used herein, means HIV-based lentiviral vectors for gene delivery, which can be integrative or non-integrative, have relatively large packaging capacity, and can transduce a range of different cell types.
25 Lentiviral vectors are usually generated following transient transfection of three (packaging, envelope and transfer) or more plasmids into producer cells. Like HIV, lentiviral vectors enter the target cell through the interaction of viral surface glycoproteins with receptors on the cell surface. On entry, the viral RNA undergoes reverse transcription, which is mediated by the viral reverse transcriptase complex. The
30 product of reverse transcription is a double-stranded linear viral DNA, which is the substrate for viral integration into the DNA of infected cells.

In certain embodiments, the viral vector can be a gammaretrovirus, e.g., Moloney murine leukemia virus (MLV)-derived vectors. In other embodiments, the viral vector can be a more complex retrovirus-derived vector, e.g., a lentivirus-derived vector. HIV-1-derived vectors belong to this category. Other examples include

5 lentivirus vectors derived from HIV-2, FIV, equine infectious anemia virus, SIV, and Maedi-Visna virus (ovine lentivirus). Methods of using retroviral and lentiviral viral vectors and packaging cells for transducing mammalian host cells with viral particles containing TCR or CAR transgenes are known in the art and have been previously

10 described, for example, in: U.S. Patent 8,119,772; Walchli *et al.*, *PLoS One* 6:327930, 2011; Zhao *et al.*, *J. Immunol.* 174:4415, 2005; Engels *et al.*, *Hum. Gene Ther.* 14:1155, 2003; Frecha *et al.*, *Mol. Ther.* 18:1748, 2010; and Verhoeven *et al.*, *Methods Mol. Biol.* 506:97, 2009. Retroviral and lentiviral vector constructs and expression systems are also commercially available. Other viral vectors also can be used for polynucleotide delivery including DNA viral vectors, including, for example adenovirus-based vectors

15 and adeno-associated virus (AAV)-based vectors; vectors derived from herpes simplex viruses (HSVs), including amplicon vectors, replication-defective HSV and attenuated HSV (Krisky *et al.*, *Gene Ther.* 5:1517, 1998).

Other vectors recently developed for gene therapy uses can also be used with the compositions and methods of this disclosure. Such vectors include those derived from

20 baculoviruses and α -viruses. (Jolly, D J. 1999. Emerging Viral Vectors. pp 209-40 in Friedmann T. ed. *The Development of Human Gene Therapy*. New York: Cold Spring Harbor Lab), or plasmid vectors (such as Sleeping Beauty or other transposon vectors).

When a viral vector genome comprises a plurality of polynucleotides to be expressed in a host cell as separate transcripts, the viral vector may also comprise

25 additional sequences between the two (or more) transcripts allowing for bicistronic or multicistronic expression. Examples of such sequences used in viral vectors include internal ribosome entry sites (IRES), furin cleavage sites, viral 2A peptide, or any combination thereof.

In certain embodiments, a vector is capable of delivering the transgene construct

30 to a host cell (e.g., a hematopoietic progenitor cell or a human immune system cell). In specific embodiments, a vector is capable of delivering a transgene construct to human

immune system cell, such as, for example, a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁺ CD8⁺ double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a dendritic cell, or any combination thereof. In further embodiments, a vector is capable of delivering a transgene construct to a naïve T cell, a central memory T cell, an effector memory T
5 cell, or any combination thereof. In some embodiments, a vector that encodes a polynucleotide or transgene construct of the present disclosure may further comprise a polynucleotide that encodes a nuclease that can be used to perform a chromosomal knockout in a host cell (*e.g.*, a CRISPR-Cas endonuclease or another endonuclease as disclosed herein) or that can be used to deliver a therapeutic transgene or portion
10 thereof to a host cell in a gene therapy replacement or gene repair therapy. Alternatively, a nuclease used for a chromosomal knockout or a gene replacement or gene repair therapy can be delivered to a host cell independent of a vector that encodes a polynucleotide or transgene construct of this disclosure.

Uses

15 In still other aspects, the present disclosure provides methods for treating or for preventing a relapse of a hyperproliferative disorder characterized by expression of an HA-1 antigen in a subject, the method comprising administering to the subject a unit dose comprising an engineered immune cell of this disclosure (or a composition comprising an engineered immune cell), thereby treating the hyperproliferative
20 disorder.

"Treat" or "treatment" or "ameliorate" refers to medical management of a disease, disorder, or condition of a subject (*e.g.*, a human or non-human mammal, such as a primate, horse, cat, dog, goat, mouse, or rat). In general, an appropriate dose or treatment regimen comprising an engineered immune cell of the present disclosure, and
25 optionally an adjuvant, is administered in an amount sufficient to elicit a therapeutic or prophylactic benefit. Therapeutic or prophylactic/preventive benefit includes improved clinical outcome; lessening or alleviation of symptoms associated with a disease; decreased occurrence of symptoms; improved quality of life; longer disease-free status; diminishment of extent of disease, stabilization of disease state; delay of disease
30 progression; remission; survival; prolonged survival; or any combination thereof.

A "therapeutically effective amount" or "effective amount", as used herein, refers to an amount of engineered immune cells sufficient to result in a therapeutic effect, including improved clinical outcome; lessening or alleviation of symptoms associated with a disease; decreased occurrence of symptoms; improved quality of life; longer disease-free status; diminishment of extent of disease, stabilization of disease state; delay of disease progression; remission; survival; or prolonged survival in a statistically significant manner. When referring to an individual active ingredient or a cell expressing a single active ingredient, administered alone, a therapeutically effective amount refers to the effects of that ingredient or cell expressing that ingredient alone.

When referring to a combination, a therapeutically effective amount refers to the combined amounts of active ingredients or combined adjunctive active ingredient with a cell expressing an active ingredient that results in a therapeutic effect, whether administered serially or simultaneously. A combination may also be a cell expressing more than one active ingredient.

The term "pharmaceutically acceptable excipient or carrier" or "physiologically acceptable excipient or carrier" refer to biologically compatible vehicles, e.g., physiological saline, which are described in greater detail herein, that are suitable for administration to a human or other non-human mammalian subject and generally recognized as safe or not causing a serious adverse event.

As used herein, "statistically significant" refers to a p value of 0.050 or less when calculated using the Students t-test and indicates that it is unlikely that a particular event or result being measured has arisen by chance.

The presently disclosed methods may be useful to, for example, treat or prevent a relapse of a hyperproliferative disorder characterized by expression of HA-1 antigen in a subject, wherein the HA-1^H antigen is present in an HLA complex expressed by hyperproliferating cells in the subject.

Examples of hyperproliferative disorders characterized by HA-1^H:HLA complexes include hematological malignancies. In certain embodiments, the hematological malignancy comprises a leukemia (e.g., an acute leukemia or a chronic leukemia). In specific embodiments, the leukemia comprises acute myeloid leukemia (AML), acute lymphocytic leukemia (ALL), mixed phenotype acute leukemia (MPAL),

chronic myeloid leukemia (CML), B cell prolymphocytic leukemia, hairy cell leukemia, or chronic lymphocytic leukemia (CLL). In certain embodiments, the hematological malignancy comprises a lymphoma. In certain embodiments, the lymphoma comprises Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), a central nervous

5 system lymphoma, small lymphocytic lymphoma (SLL), CD37+ dendritic cell lymphoma, lymphoplasmacytic lymphoma, splenic marginal zone lymphoma, plasma cell myeloma, extraosseous plasmacytoma, extra-nodal marginal zone B-cell lymphoma of mucosa-associated (MALT) lymphoid tissue, nodal marginal zone B-cell lymphoma, follicular lymphoma, mantle cell lymphoma, diffuse large B-cell lymphoma,

10 mediastinal (thymic) large B-cell lymphoma, precursor B-lymphoblastic lymphoma, immunoblastic large cell lymphoma, intravascular large B-cell lymphoma, primary effusion lymphoma, Burkitt's lymphoma/leukemia, B-cell proliferations of uncertain malignant potential, lymphomatoid granulomatosis, and post-transplant lymphoproliferative disorder. In certain embodiments, the hematological malignancy

15 comprises a myelodysplastic disorder, such as, for example, refractory cytopenia with unilineage dysplasia (refractory anemia, refractory neutropenia, and refractory thrombocytopenia), refractory anemia with ring sideroblasts (RARS), refractory anemia with ring sideroblasts – thrombocytosis (RARS-t), refractory cytopenia with multinieage dysplasia (RCMD), refractory cytopenia with multinieage dysplasia and

20 ring sideroblasts (RCMD-RS), refractory anemia with excess blasts (RAEB), myelodysplasia unclassifiable, and refractory cytopenia of childhood. In further embodiments, the hematological malignancy comprises a myeloma. Subjects that can be treated by the present invention are, in general, human and other primate subjects, such as monkeys and apes for veterinary medicine purposes. In any of the aforementioned

25 embodiments, the subject may be a human subject. The subjects can be male or female and can be any suitable age, including infant, juvenile, adolescent, adult, and geriatric subjects. Cells according to the present disclosure may be administered in a manner appropriate to the disease, condition, or disorder to be treated as determined by persons skilled in the medical art. In any of the above embodiments, an engineered immune cell

30 or unit dose as described herein is administered intravenously, intraperitoneally, intratumorally, into the bone marrow, into a lymph node, or into the cerebrospinal fluid

so as to encounter target cells (*e.g.*, leukemia cells). An appropriate dose, suitable duration, and frequency of administration of the compositions will be determined by such factors as a condition of the patient; size, type, and severity of the disease, condition, or disorder; the particular form of the active ingredient; and the method of
5 administration.

The amount of cells in a composition or unit dose is at least one cell (for example, one engineered CD8⁺ T cell subpopulation; one engineered CD4⁺ T cell subpopulation) or is more typically greater than 10² cells, for example, up to 10⁶, up to 10⁷, up to 10⁸ cells, up to 10⁹ cells, or more than 10¹⁰ cells. In certain embodiments, the
10 cells are administered in a range from about 10⁶ to about 10¹⁰ cells/m², preferably in a range of about 10⁵ to about 10⁹ cells/m². The number of cells will depend upon the ultimate use for which the composition is intended as well the type of cells included therein. For example, cells modified to contain a fusion protein specific for a particular antigen will comprise a cell population containing at least 30%, 35%, 40%, 45%, 50%,
15 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or more of such cells. For uses provided herein, cells are generally in a volume of a liter or less, 500 mls or less, 250 mls or less, or 100 mls or less. In embodiments, the density of the desired cells is typically greater than 10⁴ cells/ml and generally is greater than 10⁷ cells/ml, generally 10⁸ cells/ml or greater. The cells may be administered as a single infusion or in multiple
20 infusions over a range of time. A clinically relevant number of immune cells can be apportioned into multiple infusions that cumulatively equal or exceed 10⁶, 10⁷, 10⁸, 10⁹, 10¹⁰, or 10¹¹ cells. In certain embodiments, a unit dose of the engineered immune cells can be co-administered with (*e.g.*, simultaneously or contemporaneously) hematopoietic stem cells from an allogeneic donor (*e.g.*, a donor that is HLA^H-negative, HLA-A2-
25 negative, or both).

Also contemplated are pharmaceutical compositions (*i.e.*, compositions) that engineered immune cells as disclosed herein and a pharmaceutically acceptable carrier, diluents, or excipient. Suitable excipients include water, saline, dextrose, glycerol, or the like and combinations thereof. In embodiments, compositions comprising fusion
30 proteins or host cells as disclosed herein further comprise a suitable infusion media. Suitable infusion media can be any isotonic medium formulation, typically normal

saline, Normosol R (Abbott) or Plasma-Lyte A (Baxter), 5% dextrose in water, Ringer's lactate can be utilized. An infusion medium can be supplemented with human serum albumin or other human serum components.

Pharmaceutical compositions may be administered in a manner appropriate to the disease or condition to be treated (or prevented) as determined by persons skilled in the medical art. An appropriate dose and a suitable duration and frequency of administration of the compositions will be determined by such factors as the health condition of the patient, size of the patient (i.e., weight, mass, or body area), the type and severity of the patient's condition, the particular form of the active ingredient, and the method of administration. In general, an appropriate dose and treatment regimen provide the composition(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit (such as described herein, including an improved clinical outcome, such as more frequent complete or partial remissions, or longer disease-free and/or overall survival, or a lessening of symptom severity).

An effective amount of a pharmaceutical composition refers to an amount sufficient, at dosages and for periods of time needed, to achieve the desired clinical results or beneficial treatment, as described herein. An effective amount may be delivered in one or more administrations. If the administration is to a subject already known or confirmed to have a disease or disease-state, the term "therapeutic amount" may be used in reference to treatment, whereas "prophylactically effective amount" may be used to describe administering an effective amount to a subject that is susceptible or at risk of developing a disease or disease-state (e.g., recurrence) as a preventative course.

The pharmaceutical compositions described herein may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers may be frozen to preserve the stability of the formulation until infusion into the patient. In certain embodiments, a unit dose comprises an engineered immune cell as described herein at a dose of about 10^7 cells/ m^2 to about 10^{11} cells/ m^2 . The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including e.g., parenteral or intravenous administration or formulation.

If the subject composition is administered parenterally, the composition may also include sterile aqueous or oleaginous solution or suspension. Suitable non-toxic parenterally acceptable diluents or solvents include water, Ringer's solution, isotonic salt solution, 1,3-butanediol, ethanol, propylene glycol or polythethylene glycols in mixtures with water. Aqueous solutions or suspensions may further comprise one or more buffering agents, such as sodium acetate, sodium citrate, sodium borate or sodium tartrate. Of course, any material used in preparing any dosage unit formulation should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations. Dosage unit form, as used herein, refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit may contain a predetermined quantity of engineered immune cells or active compound calculated to produce the desired effect in association with an appropriate pharmaceutical carrier.

In general, an appropriate dosage and treatment regimen provides the active molecules or cells in an amount sufficient to provide a benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated subjects as compared to non-treated subjects. Increases in preexisting immune responses to a tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which are routine.

For prophylactic use, a dose should be sufficient to prevent, delay the onset of, or diminish the severity of a disease associated with disease or disorder. Prophylactic benefit of the immunogenic compositions administered according to the methods described herein can be determined by performing pre-clinical (including in vitro and in vivo animal studies) and clinical studies and analyzing data obtained therefrom by appropriate statistical, biological, and clinical methods and techniques, all of which can readily be practiced by a person skilled in the art.

As used herein, administration of a composition refers to delivering the same to a subject, regardless of the route or mode of delivery. Administration may be effected continuously or intermittently, and parenterally. Administration may be for treating a

subject already confirmed as having a recognized condition, disease or disease state, or for treating a subject susceptible to or at risk of developing such a condition, disease or disease state. Co-administration with an adjunctive therapy may include simultaneous and/or sequential delivery of multiple agents in any order and on any dosing schedule
5 (e.g., engineered immune cells with one or more cytokines; immunosuppressive therapy such as calcineurin inhibitors, corticosteroids, microtubule inhibitors, low dose of a mycophenolic acid prodrug, or any combination thereof).

In certain embodiments, a plurality of doses of an engineered immune cell described herein is administered to the subject, which may be administered at intervals
10 between administrations of about two to about four weeks.

Treatment or prevention methods of this disclosure may be administered to a subject as part of a treatment course or regimen, which may comprise additional treatments prior to, or after, administration of the instantly disclosed unit doses, cells, or compositions. For example, in certain embodiments, a subject receiving a unit dose of
15 the engineered immune cell is receiving or had previously received a hematopoietic cell transplant (HCT; including myeloablative and non-myeloablative HCT). In specific embodiments, the HCT comprises donor cells that are HA-1⁺, HLA-A2⁺, or both, and the subject receiving the HCT donor cells is HA-1⁺/HLA-A2⁺. In any of the foregoing embodiments, a hematopoietic cell used in an HCT may be a "universal donor" cell that
20 is modified to reduce or eliminate expression of one or more endogenous genes that encode a polypeptide product selected from an HLA molecule or a TCR molecule (e.g., by a chromosomal gene knockout according to the methods described herein).

Techniques and regimens for performing HCT are known in the art and can comprise transplantation of any suitable donor cell, such as a cell derived from umbilical cord
25 blood, bone marrow, or peripheral blood, a hematopoietic stem cell, a mobilized stem cell, or a cell from amniotic fluid. Accordingly, in certain embodiments, an engineered immune cell of the present disclosure can be administered with or shortly after hematopoietic stem cells in a modified HCT therapy.

In further embodiments, the subject had previously received lymphodepleting
30 chemotherapy prior to receiving the engineered immune cells or HCT. In certain embodiments, a lymphodepleting chemotherapy comprises a conditioning regimen

comprising cyclophosphamide, fludarabine, anti-thymocyte globulin, or a combination thereof.

Methods according to this disclosure may further include administering one or more additional agents to treat the disease or disorder in a combination therapy. For example, in certain embodiments, a combination therapy comprises administering an engineered immune cell with (concurrently, simultaneously, or sequentially) an immune checkpoint inhibitor. In some embodiments, a combination therapy comprises administering an engineered immune cell with an agonist of a stimulatory immune checkpoint agent. In further embodiments, a combination therapy comprises administering an engineered immune cell with a secondary therapy, such as chemotherapeutic agent, a radiation therapy, a surgery, an antibody, or any combination thereof.

As used herein, the term "immune suppression agent" or "immunosuppression agent" refers to one or more cells, proteins, molecules, compounds or complexes providing inhibitory signals to assist in controlling or suppressing an immune response. For example, immune suppression agents include those molecules that partially or totally block immune stimulation; decrease, prevent or delay immune activation; or increase, activate, or up regulate immune suppression. Exemplary immunosuppression agents to target (*e.g.*, with an immune checkpoint inhibitor) include PD-1, PD-L1, PD-L2, LAG3, CTLA4, B7-H3, B7-H4, CD244/2B4, HVEM, BTLA, CD160, TIM3, GAL9, KIR, PVR1G (CD112R), PVRL2, adenosine, A2aR, immunosuppressive cytokines (*e.g.*, IL-10, IL-4, IL-1RA, IL-35), IDO, arginase, VISTA, TIGIT, LAIR1, CEACAM-1, CEACAM-3, CEACAM-5, Treg cells, or any combination thereof.

An immune suppression agent inhibitor (also referred to as an immune checkpoint inhibitor) may be a compound, an antibody, an antibody fragment or fusion polypeptide (*e.g.*, Fc fusion, such as CTLA4-Fc or LAG3-Fc), an antisense molecule, a ribozyme or RNAi molecule, or a low molecular weight organic molecule. In any of the embodiments disclosed herein, a method may comprise an engineered immune cell with one or more inhibitor of any one of the following immune suppression components, singly or in any combination.

In certain embodiments, an engineered immune cell is used in combination with a PD-1 inhibitor, for example a PD-1-specific antibody or binding fragment thereof, such as pidilizumab, nivolumab, pembrolizumab, MEDI0680 (formerly AMP-514), AMP-224, BMS-936558 or any combination thereof. In further embodiments, an
5 engineered immune cell of the present disclosure (or an engineered host cell expressing the same) is used in combination with a PD-L1 specific antibody or binding fragment thereof, such as BMS-936559, durvalumab (MEDI4736), atezolizumab (RG7446), avelumab (MSB0010718C), MPDL3280A, or any combination thereof.

In certain embodiments, an engineered immune cell of the present disclosure is
10 used in combination with a LAG3 inhibitor, such as LAG525, IMP321, IMP701, 9H12, BMS-986016, or any combination thereof.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of CTLA4. In particular embodiments, an engineered immune cell is used in combination with a CTLA4 specific antibody or binding fragment thereof, such as
15 ipilimumab, tremelimumab, CTLA4-Ig fusion proteins (e.g., abatacept, belatacept), or any combination thereof.

In certain embodiments, an engineered immune cell is used in combination with a B7-H3 specific antibody or binding fragment thereof, such as enoblituzumab (MGA271), 376.96, or both. A B7-H4 antibody binding fragment may be a scFv or
20 fusion protein thereof, as described in, for example, *Dangaj et al., Cancer Res. 73:4820*, 2013, as well as those described in U.S. Patent No. 9,574,000 and PCT Patent Publication Nos. WO /201640724A1 and WO 2013/025779A1.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of CD244.

25 In certain embodiments, an engineered immune cell is used in combination with an inhibitor of BLTA, HVEM, CD160, or any combination thereof. Anti CD-160 antibodies are described in, for example, PCT Publication No. WO 2010/084158.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of TIM3.

30 In certain embodiments, an engineered immune cell is used in combination with an inhibitor of Gal9.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of adenosine signaling, such as a decoy adenosine receptor.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of A2aR.

5 In certain embodiments, an engineered immune cell is used in combination with an inhibitor of KIR, such as lirilumab (BMS-986015).

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of an inhibitory cytokine (typically, a cytokine other than TGF β) or Treg development or activity.

10 In certain embodiments, an engineered immune cell is used in combination with an IDO inhibitor, such as levo-1-methyl tryptophan, epacadostat (INCB024360; Liu *et al.*, *Blood* 115:3520-30, 2010), ebselen (Terentis *et al.*, *Biochem.* 49:591-600, 2010), indoximod, NLG919 (Mautino *et al.*, American Association for Cancer Research 104th Annual Meeting 2013; Apr 6-10, 2013), 1-methyl-tryptophan (1-MT)-tira-pazamine, or
15 any combination thereof.

In certain embodiments, an engineered immune cell is used in combination with an arginase inhibitor, such as N(omega)-Nitro-L-arginine methyl ester (L-NAME), N-omega-hydroxy-nor-l-arginine (nor-NOHA), L-NOHA, 2(S)-amino-6-boronoheptanoic acid (ABH), S-(2-boronoethyl)-L-cysteine (BEC), or any combination thereof.

20 In certain embodiments, an engineered immune cell is used in combination with an inhibitor of VISTA, such as CA-170 (Curis, Lexington, Mass.).

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of TIGIT such as, for example, COM902 (Compugen, Toronto, Ontario Canada), an inhibitor of CD155, such as, for example, COM701 (Compugen), or both.

25 In certain embodiments, an engineered immune cell is used in combination with an inhibitor of PVRIG, PVRL2, or both. Anti-PVRIG antibodies are described in, for example, PCT Publication No. WO 2016/134333. Anti-PVRL2 antibodies are described in, for example, PCT Publication No. WO 2017/021526.

30 In certain embodiments, an engineered immune cell is used in combination with a LAIR1 inhibitor.

In certain embodiments, an engineered immune cell is used in combination with an inhibitor of CEACAM-1, CEACAM-3, CEACAM-5, or any combination thereof.

In certain embodiments, an engineered immune cell is used in combination with an agent that increases the activity (*i.e.*, is an agonist) of a stimulatory immune
5 checkpoint molecule. For example an engineered immune cell can be used in combination with a CD137 (4-1BB) agonist (such as, for example, urelumab), a CD134 (OX-40) agonist (such as, for example, MEDI6469, MEDI6383, or MEDI0562), lenalidomide, pomalidomide, a CD27 agonist (such as, for example, CDX-1127), a CD28 agonist (such as, for example, TGN1412, CD80, or CD86), a CD40 agonist (such
10 as, for example, CP-870,893, rhuCD40L, or SGN-40), a CD122 agonist (such as, for example, IL-2) an agonist of GITR (such as, for example, humanized monoclonal antibodies described in PCT Patent Publication No. WO 2016/054638), an agonist of ICOS (CD278) (such as, for example, GSK3359609, mAb 88.2, JTX-2011, Icos 145-1, Icos 314-8, or any combination thereof). In any of the embodiments disclosed herein, a
15 method may comprise administering an engineered immune cell with one or more agonist of a stimulatory immune checkpoint molecule, including any of the foregoing, singly or in any combination.

In certain embodiments, a combination therapy comprises an engineered immune cell and a secondary therapy comprising one or more of: an antibody or antigen
20 binding-fragment thereof that is specific for a cancer antigen expressed by the non-inflamed solid tumor, a radiation treatment, a surgery, a chemotherapeutic agent, a cytokine, RNAi, or any combination thereof.

In certain embodiments, a combination therapy method comprises administering a fusion protein and further administering a radiation treatment or a surgery. Radiation
25 therapy is well-known in the art and includes X-ray therapies, such as gamma-irradiation, and radiopharmaceutical therapies. Surgeries and surgical techniques appropriate to treating a given cancer in a subject are well-known to those of ordinary skill in the art.

In certain embodiments, a combination therapy method comprises administering
30 an engineered immune cell and further administering a chemotherapeutic agent. A chemotherapeutic agent includes, but is not limited to, an inhibitor of chromatin

function, a topoisomerase inhibitor, a microtubule inhibiting drug, a DNA damaging agent, an antimetabolite (such as folate antagonists, pyrimidine analogs, purine analogs, and sugar-modified analogs), a DNA synthesis inhibitor, a DNA interactive agent (such as an intercalating agent), and a DNA repair inhibitor. Illustrative chemotherapeutic agents include, without limitation, the following groups: anti-metabolites/anti-cancer agents, such as pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and cytarabine) and purine analogs, folate antagonists and related inhibitors (mercaptopurine, thioguanine, pentostatin and 2- chlorodeoxyadenosine (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones and navelbine, epidipodophyllotoxins (etoposide, teniposide), DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil, cisplatin, cyclophosphamide, Cytosan, dactinomycin, daunorubicin, doxorubicin, epirubicin, hexamethylmelamineoxaliplatin, iphosphamide, melphalan, merchloroethamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, temozolamide, teniposide, triethylenethiophosphoramide and etoposide (VP 16)); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin (adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin) and mitomycin; enzymes (L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkyl sulfonates -busulfan, nitrosoureas (carmustine (BCNU) and analogs, streptozocin), trazenes— dacarbazine (DTIC); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin

salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine,

5 mycophenolate mofetil); anti-angiogenic compounds (TNP470, genistein) and growth factor inhibitors (vascular endothelial growth factor (VEGF) inhibitors, fibroblast growth factor (FGF) inhibitors); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab, rituximab); chimeric antigen receptors; cell cycle inhibitors and differentiation inducers (tretinoin); mTOR inhibitors,

10 topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin, irinotecan (CPT-11) and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); growth factor signal transduction kinase inhibitors; mitochondrial dysfunction inducers,

15 toxins such as Cholera toxin, ricin, Pseudomonas exotoxin, Bordetella pertussis adenylate cyclase toxin, or diphtheria toxin, and caspase activators; and chromatin disruptors.

Cytokines are increasingly used to manipulate host immune response towards anticancer activity. *See, e.g., Floros & Tarhini, Semin. Oncol. 42(4):539-548, 2015.*

20 Cytokines useful for promoting immune anticancer or antitumor response include, for example, IFN- α , IL-2, IL-3, IL-4, IL-10, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18, IL-21, IL-24, and GM-CSF, singly or in any combination with an engineered immune cell of this disclosure.

Also provided herein are methods for modulating an adoptive immunotherapy,

25 wherein the methods comprise administering, to a subject who has previously received an engineered immune cell of the present disclosure that comprises a heterologous polynucleotide encoding a safety switch protein, a cognate compound of the safety switch protein in an amount effective to ablate in the subject the previously administered engineered immune cell.

30 As used herein, the term "adoptive immune therapy" or "adoptive immunotherapy" refers to administration of naturally occurring or genetically

engineered, disease- or antigen-specific immune cells (*e.g.*, T cells). Adoptive cellular immunotherapy may be autologous (immune cells are from the recipient), allogeneic (immune cells are from a donor of the same species) or syngeneic (immune cells are from a donor genetically identical to the recipient).

5 In certain embodiments, the safety switch protein comprises tEGFR and the cognate compound is cetuximab, or the safety switch protein comprises iCasp9 and the cognate compound is AP1903 (*e.g.*, dimerized AP1903), or the safety switch protein comprises a RQR polypeptide and the cognate compound is rituximab, or the safety switch protein comprises a myc binding domain and the cognate compound is an
10 antibody specific for the myc binding domain.

 In still further aspects, methods are provided for manufacturing a composition, or unit dose of the present disclosure. In certain embodiments, the methods comprise combining (i) an aliquot of a host cell transduced with a vector of the present disclosure with (ii) a pharmaceutically acceptable carrier. In certain embodiments, vectors of the
15 present disclosure are used to transfect/transduce a host cell (*e.g.*, a T cell) for use in adoptive transfer therapy (*e.g.*, targeting a cancer antigen).

 In some embodiments, the methods further comprise, prior to the aliquotting, culturing the transduced host cell and selecting the transduced cell as having incorporated (*i.e.*, expressing) the vector. In further embodiments, the methods
20 comprise, following the culturing and selection and prior to the aliquotting, expanding the transduced host cell. In any of the embodiments of the instant methods, the manufactured composition or unit dose may be frozen for later use. Any appropriate host cell can be used for manufacturing a composition or unit dose according to the instant methods, including, for example, a hematopoietic stem cell, a T cell, a primary T
25 cell, a T cell line, a NK cell, or a NK-T cell. In specific embodiments, the methods comprise a host cell which is a CD4⁺ T cell or a CD8⁺ T cell.

EXAMPLES

EXAMPLE 1

ISOLATION AND CLONING OF HA-1^H-SPECIFIC TCRs

HA-1^H-specific CD8⁺ T cell clones were isolated using *in vitro* methods previously described (Bleakley *et al.*, *Blood* 115:4923-4933, 2010 (Figure 1). Specifically, CD8⁺ T cells were isolated from HLA-A2⁺ donor (2 donors) peripheral blood mononuclear cells (PBMC), using a CD8⁺ T cell isolation kit and anti-CD45RO immunomagnetic beads (Miltenyi Biotec). Autologous dendritic cells (DCs) were pulsed with 1μg/mL HA-1^H peptide (VLHDDLLEA) for 3-6 hours at 37°C. Purified CD8⁺ T_N were combined in complete T lymphocyte (CTL) medium with peptide-pulsed DCs at a T_N to DC ratio of 30:1, and co-cultured in 96-well plates at 6x10⁴ T cells/well, supplemented with 10 ng/mL IL-12 from initiation and 10ng/mL IL-15 from day 7. On day 11-13, cells were evaluated for HA-1^H-specific cytotoxicity in split-well microchromium release assays (CRA; μCRA). T cell lines that lysed T2 cells pulsed with 1μg/mL HA-1^H peptide (>20% lysis and >5 fold more lysis of peptide-pulsed versus unpulsed targets) were subsequently cloned by limiting dilution using anti-CD3 monoclonal antibody (mAb), interleukin-2 (IL-2) and feeder cells.

Clones were screened by μCRA on day 11-13. T cell clones from wells showing specific cytotoxicity, using the above criteria, were expanded using anti-CD3 mAb, IL-2 and feeder cells, by the Rapid Expansion Protocol (REP). The specificity of expanded clones was evaluated by CRA, HA-1/HLA-A2 multimer staining, and intracellular cytokine staining (ICC). The HA-1^H specificity of the CTL clones was verified with HA-1^H/HLA A2 multimers. In particular, HLA-A2/ HA-1^H multimer and CD8⁺ monoclonal antibody (mAb) were used to stain HA-1 specific clones (clones 1, 2, 10, 13, 14, 16, and 5) and a control clone specific for another tumor antigen (Figure 2A).

In addition, a chromium release assay (CRA) was used to test 7 of the HA-1-specific CTL clones (1, 2, 10, 13, 14, 16 and 5) for killing of HA-1^H peptide-pulsed targets. The T cell clones recognized HA-1^H peptide (VLHDDLLEA)-pulsed target

cells at very low peptide concentrations and half-maximal lysis is seen at a peptide concentration of 10 pM. (Figure 2B) (ICC data not shown). Cytotoxicity of the the isolated clones against cells with or without endogenous HA-1^H expression was also examined. The assays show 7 HA-1^H-specific CTL clones (TCR1, TCR2, TCR10, TCR13, TCR14, TCR16 and TCR5) lysing HA-1^{H+} acute myeloid leukemia (AML) cell line (THP-1) and HA-1^{H+} primary AML but not HA-1^{H-} AML (Figure 2C).

Next, the isolated HA-1^H TCRs were cloned. The TCR V_β and V_α genes of the 8 CTL were sequenced and 6 distinct TCRs were identified (TCR1 and TCR13 were found to be identical, as were TCR2 and TCR14). The genes (and specific alleles) encoding the 6 TCRs, as well as the V_β CDR3 amino acid sequences of the TCRs, are shown in Table 1:

Table 1. Genes and βCDR3 sequences of isolated TCRs

	Alpha		Beta			
	V gene	J gene	V gene	D gene	J gene	CDR3 (aa)
TCR 1	TRAV17*01 F	TRAJ28*01 F	TRVB7-9*03	TRBD1*01	TRBJ21*01	CASSSTGGHNEQFF
TCR 13	TRAV17*01 F	TRAJ28*01 F	TRVB7-9*03	TRBD1*01	TRBJ21*01	CASSSTGGHNEQFF
TCR 2	TRAV21*02 F	TRAJ40*01 F	TRVB7-9*03	TRBD1*01	TRBJ1-4*01	CASSLVKGEKLFF
TCR 14	TRAV21*02 F	TRAJ40*01 F	TRVB7-9*03	TRBD1*01	TRBJ1-4*01	CASSLVKGEKLFF
TCR 10	TRAV10*01F	TRAG45*01 F	TRVB7-9*03	TRBD2*01	TRBJ27*01	CASSMLTNYEQYF
TCR 16	TRAV21*01 or *02	TRAF20*01 F	TRVB7-9*03	TRBD1*01 F	TRBJ21*01	CASSLVVGNEQFF
TCR 5	TRAV17	TRAJ29	TRVB7-9*03	TRBD1/2	TRBJ2-7	CASSLTTLDEQY
TCR 24	TRAV8-3	TRAJ27	TRBV15	TRBD1*01	TRBJ2-5	ATSKTRIAQETQYF

A seventh TCR, "TCR29" was also identified and sequenced. After sequencing, the genes encoding the HA-1^H TCRs (except for TCR24, which had poor function in transduced CD8⁺ T cells) were codon optimized to maximize expression and cysteine modifications were introduced and to reduce the risk of mispairing with endogenous TCR chains, as described below. . The nucleotide sequences of the CTL clone TCRβ

and TCR α genes after codon optimization and cysteine modification are provided in SEQ ID NOs 39-48 and 51-52.

RNA was extracted from each HA-1^H-specific T cell clone. 5'-first-strand cDNA amplification and Rapid Amplification of cDNA Ends (RACE) PCR were performed to identify full-length TCR regions, using a SMARTer RACE cDNA Amplification Kit (Clontech Laboratories). cDNA was synthesized from RNA using 5' CDS Primer A, SMARTer IIA oligo and SMARTScribe Reverse Transcriptase. Subsequently, the cDNA was used to perform a RACE PCR reaction, using Phusion® High Fidelity DNA polymerase, and Gene-Specific Primers for the TCR alpha (α)- (5'-GGTGAATAGGCAGACAGACTT -3') (SEQ ID NO:93) or TCR beta (β)- chain (GTGGCCAGGCACACCAGTGT) (SEQ ID NO:94). The RACE PCR product was purified and sequenced to identify the TCR α - and β -chains. IMGT/V-QUEST was used to define the TCR variable (V), diversity (D) and joining (J) regions.

Complementary cysteine residues at positions 48 (Thr to Cys) and 57 (Ser to Cys) were incorporated into the constant domains of the TCR α and β genes to increase exogenous TCR pairing and decrease mispairing with the endogenous TCR. To ensure coordinated gene expression, the TCR chains were separated by 2A elements from the porcine teschovirus (P2A). The transgenes were codon-optimized to enhance expression and synthesized by GeneArt (Life Technologies), and were cloned into the pRRLSIN.cPPT.MSCV.WPRE LV vector by restriction digestion and ligation. The amino acid sequences of the encoded TCR constructs are provided in SEQ ID NOS: 53-57.

EXAMPLE 2

HETEROLOGOUS EXPRESSION AND ACTIVITY OF HA-1^H-SPECIFIC TCRs

Next, codon-optimized and cysteine modified TCRs were tested for expression and activity.

Lentiviral (LV) vectors were used to transduce primary T cells to deliver a polynucleotide encoding the engineered HA-1^Hspecific TCR constructs. The transduced T cells were sorted, expanded, and tested for HA-1^Hspecificity (Figure 4).

Five of six HA-1^H TCR LVs efficiently transduced primary CD8⁺ and CD4⁺ T cells (Figures 5A and 6A), and conferred specific recognition of HLA-A2⁺ cells pulsed with low amounts of HA-1^H peptide. TCR2 and TCR16 showed the strongest cytotoxic activity and were selected for further experiments. T cells transduced with these TCRs killed HA-1^H-pulsed cells (Figures 5B, 5C, 6B, and 35), cell lines with endogenous HA-1^H expression (Figures 6C-6E and Figure 7) and primary leukemia cells with endogenous HA-1^H (Figures 8A-8E, Figure 9), but these engineered cells were not activated by, and did not kill, HA-1^H-negative cells (Figure 7), primary leukemia cells (Figures 8A, 8B and 9) or fibroblasts (Figure 10). In addition, CD8⁺ and CD4⁺ HA-1^H TCR T cells secreted IFN γ and IL-2 in response to HA-1^H peptide stimulation (data not shown). HA-1^H TCR CD4⁺ cells killed target cells pulsed with high peptide concentrations, although LV transduction with the HA-1^H TCR alone did not make CD4⁺ T cells responsive to primary leukemia cells with native levels of antigen (data not shown).

In sum, these data demonstrate that T cells effectively express transduced HA-1^H TCRs and have antigen-specific killing activity against lymphoid cells.

EXAMPLE 3

CD8 CO-RECEPTOR FUNCTION IN CD4⁺ HA-1^H TCR CELLS

Inclusion of CD4⁺ T cells in an immunotherapy cell product can provide antigen-induced IL-2 secretion and augment persistence and function of transferred cytotoxic CD8⁺ T cells (*see, e.g., Kennedy et al., Immunol. Rev.* 222:129 (2008); Nakanishi *et al., Nature* 462(7272):510 (2009)). However, optimal function of many class I restricted TCR in CD4⁺ T cells requires the transfer of a CD8 co-receptor to enhance sensitivity of the TCR to class I HLA peptide complexes. CD4 co-receptors differ in structure to CD8 and cannot effectively substitute for CD8 co-receptors (*see, e.g., Stone & Kranz, Front. Immunol.* 4:244 (2013); *see also Cole et al., Immunology* 137(2):139 (2012). Relatively high HA-1^H peptide concentrations were required to induce cytolytic activity in CD4⁺ T cells transduced with an HA-1^H TCR alone, and

HA-1 TCR CD4⁺ T cells did not recognize cell lines or leukemia, implying CD8 co-receptor dependency of the TCR (data not shown).

Various options for including a CD8 co-receptor in the transgene construct were explored. CD8 co-receptors exist on the surface of human conventional $\alpha\beta$ TCR T cells, typically as dimers of CD8 α - and β -chains, and there are five β -chain variants with different intracytoplasmic tail sequences (β M1-5) (*see, e.g., Thakral et al., J. Immunol. 180(11):7431 (2008); see also Thakral et al., PLoS One 8(3):e59374 (2013)*). The amino acid sequences of the CD8 co-receptor chains are provided in SEQ ID NOs: 48-53.

HA-1^H TCR2 constructs were generated that included one or both of the CD8 α -chain and the CD8 β -chain as full-length or truncated variants. When used to transduce primary CD4⁺ T cells, the α - and β -chains were expressed on the cell surface; $\alpha\beta$ dimers and α monomers increased HA-1^H/HLA-A2 multimer binding by TCR-transduced T cells to a greater extent than did β monomers (Figure 11A(i-ii)). CD8 α - and β -chains with truncations of the intracellular chain components did not increase multimer binding above the TCR alone (Figure 11A(iii)). Transduction with the CD8 α and β M1 or β M4 variants improved HA-1^H TCR function in CD4⁺ T cells more than did the CD8 α and β M2 or β M5 chains (Figure 11B). In functional assays, incorporation of the β M1 or β M4 chain improved the CD4⁺ T cell function to a greater extent than CD8 α monomers (Figures 11B and 11C). β M1 provided greater specificity of response than did β M4. The β M1 variant was therefore selected and used in a multi-cistronic LV that included CD8 α and β M1 sequences as well as the HA-1^H TCR. CD4⁺ T cells transduced with the vector secreted IL-2 and interferon gamma (IFN γ) (Figure 12A) and proliferated when co-cultured with HA-1^{H+} AML cells (Figure 12B).

EXAMPLE 4

INTRODUCTION OF A SAFETY SWITCH INTO THE HA-1^H TCR CONSTRUCTS

To ensure that HA-1^H TCR-transduced T cells can be rapidly depleted in case of any unexpected toxicity, four codon-optimized "safety switch" HA-1^H TCR constructs were generated: (1) The inducible caspase 9 (iCasp9) is based on the fusion of human

caspase 9 to a modified human FK-binding protein, allowing conditional dimerization; when exposed to a synthetic dimerizing drug, iCasp9 becomes activated and initiates rapid death of cells expressing this construct (*see, e.g., Straathof et al., Blood* 105(11):4247 (2005)); (2) The truncated human EGFR ("tEGFR") is a polypeptide devoid of extracellular N-terminal ligand binding domains and intracellular receptor tyrosine kinase activity, but retains type I transmembrane cell surface localization and a binding epitope for pharmaceutical-grade anti-EGFR mAb, cetuximab (*see Wang et al., Blood* 118(5):4255 (2011)); (3) RQR8 is a compact combined marker and safety switch for T cells, combining target epitopes from both CD34 (a marker recognized by antibody QBEnd10) and CD20 antigens (extracellular loop mimotopes) presented on a truncated CD8 co-receptor stalk; RQR8 is bound by the pharmaceutical-grade anti-CD20 mAb, rituximab (*see Philip et al., Blood* 124(8):1255 (2011)); (4) Myc-tagged TCR incorporate a 10-amino acid tag of the human c-Myc protein that is bound by a tag-specific mAb (*see Kieback et al., PNAS* 105(2):623 (2008)).

Binding of the respective mAbs to tEGFR, RQR8 or myc-tag provides a target for complement-dependent or antibody-dependent cellular cytotoxicity and elimination of transduced cells. The safety switch molecules were cloned into TCR2 LV constructs upstream of, and operatively associated with, the the TCR β and TCR α coding sequences. The TCR β and TCR α sequences were separated by P2A elements from the porcine teschovirus to ensure coordinated gene expression. T cells transduced with iCasp9-HA-1^H TCR, tEGFR-HA-1^H TCR, RQR8-HA-1^H TCR or Myc-tagged HA-1^H -TCR all demonstrated HA-1^H TCR expression and HA-1^{H+} target cell recognition similar to recognition by T cells transduced with the HA-1^H TCR alone (Figure 14).

To test the ability of the safety switches to eliminate T cells, transduced T cells were incubated for 24 hours with the optimal concentration (Figure 16) of the respective cognate drug (the dimerizer AP1903 for iCasp9/HA-1^H TCR; complement plus appropriate mAb (anti-EGFR mAb for tEGFR-HA-1^H TCR; anti-CD20 mAb plus for RQR8-HA-1^H TCR; and anti-Myc tag mAb for Myc-tagged HA-1^H-TCR in all other constructs). All of the safety switch transduced T cells were susceptible to their respective trigger (Figure 16, *see also* Figure 15). However, iCasp9 with AP1903

consistently provided the most rapid and complete elimination of transduced T cells and was selected for further evaluation.

EXAMPLE 5

DESIGN AND SELECTION OF A TRANSGENE CONSTRUCT

5 Next, the HA-1^H TCR transgenes incorporating both iCasp9 and the CD8 $\alpha\beta$ M1 co-receptor were analyzed for functionality. To assist in the selection and tracking of transduced T cells, two marker configurations were designed. In one, the minimal CD34 epitope ("Q") of the RQR polypeptide was nested into the α -chain of the HA-1^H TCR. In the other, the RQR polypeptide was incorporated into the β -chain of the full-
10 length functional CD8 $\alpha\beta$ M1 co-receptor. Five LV transgene constructs were then created, used to transduce CD8⁺ T cells, and compared: (1) iCasp9- HA-1^H TCR2; (2) HA-1^H -TCR2-CD8 co-receptor; (3) iCasp9- HA-1^H TCR2-CD8; (4) iCasp9- HA-1^H TCR2-RQR-CD8; and (5) iCasp9-CD34- HA-1^H TCR2-CD8 (see Figures 17 and 18).

All of the constructs produced T cells that specifically secreted cytokines and
15 killed HA-1^{H+}, but not HA-1^{H-} AML cells or fibroblasts, and had similar function (see Figures 19-26), except the iCasp9-CD34-HA-1^H TCR2-CD8 construct (with the CD34 epitope embedded in the TCR α -chain), which performed poorly. The iCasp9-HA-1^H TCR2-RQR-CD8 transgene, which contains all of the desired elements (including the capacity for immunomagnetic selection; see Figures 27-29) and functioned as well as
20 the less complex constructs, was selected for further studies. A schematic diagram of this construct is shown in Figure 30, and the nucleotide sequence of the construct is provided in SEQ ID NO:85.

EXAMPLE 6

CLINICAL-SCALE PRODUCTION AND TESTING OF HA-1^H TCR T CELLS

25 The cellular composition of T cell immunotherapy products can have important downstream effects on the persistence and function of antigen-specific T cells after adoptive T cell transfer (see, e.g., Sommermeyer *et al.*, *Leukemia* 30(2):492 (2016); see also Wang *et al.*, *Blood* 117(6):1888 (2011) and Hinrichs *et al.*, *PNAS* 106*41):17469

(2009)). In general, infusion of antigen-specific T cells derived from "younger" T cell subsets, including T_N, T memory stem cells (T_{SCM}) and central memory T cells (T_{CM}), appears advantageous. In the context of post-HCT T cell immunotherapy, it is also important to consider the potential for GVHD mediated by the native TCR of donor T cells. T_N cause severe GVHD in murine models and depletion of CD45RA⁺ T_N from PBSC grafts reduces the risk of severe and/or chronic GVHD in humans (*see, e.g.,* Bleakley *et al., J. Clin. Invest.* 125(7):2677 (2015)). It is also desirable to include both CD4⁺ and CD8⁺ cells specific for the same antigen, as CD4⁺ T helper cells can enhance anti-tumor CTL responses by enhancing clonal expansion at the tumor site and preventing activation-induced cell death (*see, e.g.,* Giuntoli *et al., Clin. Cancer Res.* 8(3):922 (2002); *see also* Kennedy and Celis, *J. Immunol.* 177(5):2862 (2006)). Therefore, the T cell product (1-2 x 10⁹ PBSC/PBMC) was first depleted of CD45RA⁺ T_N cells to minimize the risk of serious GVHD, and depleted of CD14⁺ monocytes to optimize LV transduction efficiency, prior to separating CD8⁺ and CD4⁺ enriched fractions to ensure a consistent CD4:CD8 composition (approx. 3x10⁶ cells of each cell type), and stimulating (CD3/CD28 microbeads beads) and transducing the T cells with the iCasp9-HA1 TCR2-RQR-CD8 LV.

The transduced cells were flow-sorted using HA-1^H/HLA-A2 multimers and CD34 mAb 4-5 days later, and cultured in G-Rex flasks using REP (Rapid Expansion Protocol) comprising OKT3 cells, PBMC, HA-1⁺ LCL, and IL-2. The CD4⁺ and CD8⁺ HA-1^HTCR memory T cells expanded efficiently, with an average 2000-fold expansion (Figure 32A; *see also* Figure 31, left-most panels). The CD4⁺ and CD8⁺-transduced T cells were harvested and combined (total 3-6x10⁹ cells at days 16-20), then enriched via selection for CD34 to produce an enriched population of ≥1.5x 10⁹ cells. Release assays were then performed to test for purity (>75%), viability, function, specificity, presence or absence of virus, and sterility. The final T cell product retained expression of the HA-1^H TCR (Figure 32B), had a predominantly CD45RO⁺CD28⁺ phenotype with variable expression of CD62L, CCR7 and CD27 (Figure 32C), and included cells that did not express exhaustion markers such as PD-1 (Figures 32D, 32E).

The expanded CD8⁺ and CD4⁺ HA-1^H TCR T cells retained their ability to specifically kill and secrete cytokines in response to stimulation with HA-1^H-pulsed

cells (Figure 33A) or HA-1^H leukemia cell lines (Figure 33B), and many HA-1^H TCR CD8⁺ and CD4⁺ cells secreted multiple cytokines (Figure 33C; *see also* Figure 31 middle and right-hand panels). Further, the cells could be enriched using anti-CD34 immunomagnetic beads (Figure 34A) and were efficiently eliminated by exposure to the AP1903 dimerizer drug (Figure 34B). Finally, the native TCR in HA-1^H TCR CD4⁺ and CD8⁺ T cells in the cell product was evaluated using TCR immunosequencing (Adaptive Biotechnologies) and a diverse polyclonal population was observed (data not shown). Moreover, the product contained numerous very-low-frequency TCR, a finding that has recently been associated with the potential for persistence and expansion after adoptive T cell transfer (Chapuis *et al.*, *Sci. Immunol.* 2(8) (2017).

EXAMPLE 7

CLINICAL STUDY USING THE HA-1^H T CELL THERAPY PRODUCT

A feasibility and safety study is conducted using the HA-1^H cell therapy product described in Example 5 (CD8⁺ and CD4⁺ memory T cells transduced with pRRLSIN iC9- HA-1^H-TCR2-RQR-CD8; abbreviated hereafter as "HA-1^H TCR LV"). The patient sample comprises children, adolescents and adults with recurrent leukemia (AML, ALL, another acute leukemia, or CML). Specifically, patients aged 0-70 are enrolled into two age groups of approximately 12 subjects each: one group aged ≥ 16 years, and one group aged < 16 years. group. All patients express HLA-A*0201 and have the HA-1(H) genotype (RS_1801284: A/G, A/A). The patients also have an adult donor for HCT who is adequately HLA-matched by institutional standards, and are currently undergoing or have previously undergone allogeneic HCT for AML, ALL, another type of acute leukemia, or chronic myeloid leukemia.

Bone marrow samples are taken subsequent to a suspected relapse. Patients then generally receive lymphodepleting chemotherapy (fludarabine) prior to infusion with the T cell product. Thereafter, patients are administered a single dose of HA-1^H TCR LV-T cells (approx. 1:1 CD4⁺:CD8⁺ T_M) when three criteria are satisfied: (1) there is evidence of recurrent or refractory disease after HCT; (2) HA-1^H TCR T cells have been generated; and (3) lymphodepleting chemotherapy has been administered (if

indicated). HA-1^H TCR LV-T cells are administered by infusion (as rapidly as tolerated through a central venous catheter via gravity or a syringe pump) according to the dosage schedule provided in Table 2.

Table 2. Clinical Dosing Schedule

Dose level	Dose (HA-1 ^H TCR T cells)	
-1	Up to $3 \times 10^5/\text{kg}$	
0	Up to $1 \times 10^6/\text{kg}$	
1	Up to $3 \times 10^6/\text{kg}$	Starting dose
2	Up to $10 \times 10^6/\text{kg}$	
3	Up to $30 \times 10^6/\text{kg}$	

5 At least 1-2 subjects ≥ 16 y.o. are treated prior to treatment of a subject in the younger cohort. In each age group (≥ 16 and < 16 years old), patients are treated in cohorts of three or more patients at one of five dose levels of HA-1^H TCR T cells, starting at dose level 1 (3×10^6 HA-1^H TCR T cells/kg). A 28-day period between administration of the investigational agent to consecutive subjects within each age group is observed. Bone marrow samples are taken at days 4, 18, and 32 following infusion of the T cell product. Other aspects of the study include monitoring: the *in vivo* persistence of transferred HA-1^H TCR T cells in peripheral blood; the ability of HA-1^H TCR T cells to migrate to bone marrow; the function of HA-1^H TCR T cells before and, if possible, after adoptive T cell transfer; whether infusion of HA-1^H TCR T cells is followed by a reduction of leukemia burden; whether infusion of HA-1^H TCR T cells is followed by a reduction of recipient hematopoietic chimerism; and whether infusion of HA-1^H TCR T cells is followed by the appearance or recurrence of signs or symptoms of graft-versus-host disease (GVHD).

20 U.S. Provisional Patent Application No. 62/399,291, filed September 23, 2016, to which the present application claims priority, is hereby incorporated herein by reference in its entirety.

The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet are
5 incorporated herein by reference, in their entirety. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, applications and publications to provide yet further embodiments.

These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be
10 construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

CLAIMS

What is claimed is:

1. An engineered immune cell, comprising a heterologous polynucleotide encoding a binding protein that includes:

(a) a T cell receptor (TCR) α -chain variable (V_α) domain having an amino acid sequence encoded by a TRAV17 gene, a TRAV21 gene, or a TRAV10 gene, and a TCR β -chain variable (V_β) domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86;

(b) a TCR V_α domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V_β domain having an amino acid sequence encoded by a TRBV7-9 gene; or

(c) a TCR V_α domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V_β domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86,

wherein the encoded binding protein is capable of specifically binding to a peptide containing an HA-1^H antigen and does not bind to a peptide that does not contain an HA-1^H antigen.

2. The engineered immune cell of claim 1, wherein:

(a) the encoded V_β CDR3 comprises the amino acid sequence of SEQ ID NO:14, and the encoded V_α CDR3 comprises the amino acid sequence of SEQ ID NO:88;

(b) the encoded V_β CDR3 comprises the amino acid sequence of SEQ ID NO:13, and the encoded V_α CDR3 comprises the amino acid sequence of SEQ ID NO:87;

(c) the encoded V_β CDR3 comprises the amino acid sequence of SEQ ID NO:15, and the encoded V_α CDR3 comprises the amino acid sequence of SEQ ID NO:89;

(d) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:16, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:90;

(e) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:17, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:91; or

(f) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:86, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:92.

3. The engineered immune cell of claim 1 or 2, wherein the encoded V_{α} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:2, 4, 6, 8, 10, and 12.

4. The engineered immune cell of any one of claims 1-3, wherein the encoded V_{β} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:1, 3, 5, 7, 9, and 11.

5. The engineered immune cell of claim 3 or 4, wherein the encoded V_{α} domain comprises no change in amino acid sequence of CDR1, the encoded V_{β} domain comprises no change in amino acid sequence of CDR1, or the CDR1 of the encoded V_{α} domain and the CDR1 of the encoded V_{β} domain comprise no change in amino acid sequence.

6. The engineered immune cell of any one of claims 3-5, wherein the encoded V_{α} domain comprises no change in amino acid sequence of CDR2, the encoded V_{β} domain comprises no change in amino acid sequence of CDR2, or the CDR2 of the encoded V_{α} domain and the CDR2 of the encoded V_{β} domain comprise no change in amino acid sequence.

7. The engineered immune cell of any one of claims 1-6, wherein:
- (a) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:3, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:4;
 - (b) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:1, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:2;
 - (c) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:5, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:6;
 - (d) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:7, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:8;
 - (e) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:9, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:10; or
 - (f) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:11, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:12.
8. The engineered immune cell of any one of claims 1-7, wherein the encoded binding protein comprises a TCR α -chain constant domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:19-22, 24, and 26.
9. The engineered immune cell of any one of claims 1-8, wherein the encoded binding protein comprises a TCR β -chain constant domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:18, 23, and 25.

10. The engineered immune cell of any one of claims 1-9, wherein the encoded binding protein comprises a TCR α -chain having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36, and 38.

11. The engineered immune cell of any one of claims 1-10, wherein the encoded binding protein comprises a TCR α -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36, and 38.

12. The engineered immune cell of any one of claims 1-11, wherein the encoded binding protein comprises a TCR β -chain having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37.

13. The engineered immune cell of any one of claims 1-12, wherein the encoded binding protein comprises a TCR β -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37.

14. The engineered immune cell of any one of claims 8-13, wherein the encoded binding protein comprises:

(a) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:27, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:28;

(b) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:29, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:30;

(c) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:31, and the TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:32;

(d) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:33, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:34;

(e) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:35, and a TCR- α chain comprising or consisting of the amino acid sequence of SEQ ID NO:36; or

(f) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:37, and a TCR- α chain comprising or consisting of the amino acid sequence of SEQ ID NO:38.

15. The engineered immune cell of any one of claims 10-14, wherein the (i) heterologous polynucleotide encoding the TCR α -chain and the (ii) heterologous polynucleotide encoding the TCR β -chain are contained in a single open reading frame, wherein the single open reading frame further comprises a polynucleotide encoding a self-cleaving peptide disposed between (i) and (ii).

16. The engineered immune cell of claim 15, wherein the polynucleotide encoding the self-cleaving peptide comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 76-84.

17. The engineered immune cell of claim 15 or 16, wherein the single open reading frame comprises a nucleotide sequence that is at least about 80% identical to the nucleotide sequence of any one of SEQ ID NOS: 59-63.

18. The engineered immune cell of claim 17, wherein the single open reading frame comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 59-63.

19. The engineered immune cell of any one of claims 15-18, wherein the encoded single open reading frame comprises or consists of the amino acid sequence of any one of SEQ ID NOS: 53-57.

20. The engineered immune cell of any one of claims 1-19, wherein the encoded binding protein is capable of specifically binding to a HA-1^H peptide:HLA complex.

21. The engineered immune cell of claim 20, wherein the HLA comprises HLA-A*0201.

22. The engineered immune cell of any one of claims 1-21, wherein the HA-1^H peptide comprises the amino acid sequence VLHDDLLEA (SEQ ID NO:66).

23. The engineered immune cell of any one of claims 1-22, wherein the encoded binding protein comprises a TCR, an antigen-binding fragment of a TCR, or a chimeric antigen receptor (CAR).

24. The engineered immune cell of claim 23, wherein the antigen-binding fragment of the TCR comprises a single chain TCR (scTCR).

25. The engineered immune cell of claim 23, wherein the encoded binding protein comprises a CAR.

26. The engineered immune cell of claim 23, wherein the encoded binding protein comprises a TCR.

27. The engineered immune cell of any one of claims 1-26, further comprising a heterologous polynucleotide encoding:

- (a) a safety switch protein;
- (b) a selection marker;
- (c) a CD8 co-receptor β -chain;
- (d) a CD8 co-receptor α -chain; or
- (e) any combination thereof.

28. The engineered immune cell of claim 27, wherein the encoded safety switch protein comprises:

- (i) a truncated EGF receptor (tEGFR);
- (ii) iCasp9;
- (iii) a RQR polypeptide;
- (iv) a myc epitope; or
- (v) any combination thereof.

29. The engineered immune cell of claim 27 or 28, wherein the encoded selection marker comprises:

- (i) a RQR polypeptide;
- (ii) a truncated low-affinity nerve growth factor (tNGFR);
- (iii) a truncated CD19 (tCD19);
- (iv) a truncated CD34 (tCD34); or
- (v) any combination thereof.

30. The engineered immune cell of any one of claims 27-29, wherein the encoded CD8 co-receptor comprises a β -chain comprising the amino acid sequence of any one of SEQ ID NOS:71-75.

31. The engineered immune cell of any one of claims 27-30, wherein the encoded CD8 co-receptor is a recombinant CD8 co-receptor comprising a RQR polypeptide having the amino acid sequence of SEQ ID NO:69.

32. The engineered immune cell of claim 31, wherein the encoded RQR polypeptide is contained in a β -chain of the encoded CD8 co-receptor.

33. The engineered immune cell of claim 31 or 32, wherein the encoded RQR polypeptide is contained in an α -chain of the encoded CD8 co-receptor.

34. The engineered immune cell of claim 32, comprising a heterologous polynucleotide encoding iCasp9 and a heterologous polynucleotide encoding a recombinant CD8 co-receptor protein that comprises a β -chain containing a RQR polypeptide.

35. The engineered immune cell of claim 32 or 34, comprising a heterologous polynucleotide that encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a porcine teschovirus 2A (P2A) peptide]-[a TCR β -chain]-[a P2A peptide]-[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]).

36. The engineered immune cell of claim any one of claims 27-35, wherein the TCR β -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:41, and wherein the TCR α -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:42.

37. An engineered immune cell, comprising a heterologous transgene polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO:85.

38. The engineered immune cell of any one of claims 1-37, wherein the immune cell is a T cell, a NK cell, or a NK-T cell.

39. The engineered immune cell of claim 38, wherein the immune cell is a CD4⁺ T cell or a CD8⁺ T cell.

40. The engineered immune cell of claim 39, wherein the immune cell comprises a chromosomal gene knockout of a PD-1 gene; a LAG3 gene; a TIM3 gene; a CTLA4 gene; an HLA component gene; a TCR component gene, or any combination thereof.

41. The engineered immune cell of claim 40, wherein the chromosomal gene knockout comprises a knockout of an HLA component gene selected from an $\alpha 1$ macroglobulin gene, an $\alpha 2$ macroglobulin gene, an $\alpha 3$ macroglobulin gene, a $\beta 1$ microglobulin gene, or $\beta 2$ microglobulin gene.

42. The engineered immune cell of claim 41, wherein the chromosomal gene knockout comprises a knockout of a TCR component gene selected from a TCR α variable region gene, a TCR β variable region gene, a TCR constant region gene, or a combination thereof.

43. A composition, comprising an engineered immune cell of any one of claims 1-42 and a pharmaceutically acceptable carrier, diluent, or excipient.

44. A unit dose, comprising an effective amount of (i) the engineered immune cell of any one of claims 1-42 or (ii) a composition according to claim 43.

45. The unit dose of claim 44, comprising (i) a composition comprising at least about 30% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 30% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains substantially no naïve T cells.

46. A method for treating or for preventing a relapse of a hyperproliferative disorder characterized by expression of an HA-1^H antigen in a subject, the method comprising administering to the subject a unit dose of claim 44 or 45, thereby treating the hyperproliferative disorder.

47. The method of claim 46, wherein the HA-1^H antigen is present in an HLA complex expressed by hyperproliferating cells in the subject.

48. The method of claim 46 or 47, wherein the hyperproliferative disorder comprises a hematological malignancy.

49. The method of claim 48, wherein the hematological malignancy comprises a leukemia, a lymphoma, a myelodysplastic disorder, or a myeloma.

50. The method of claim 49, wherein the hematological malignancy comprises a leukemia.

51. The method of claim 50, wherein the leukemia is selected from acute myeloid leukemia (AML), acute lymphocytic leukemia (ALL), mixed phenotype acute leukemia (MPAL), chronic myeloid leukemia (CML), B cell prolymphocytic leukemia, hairy cell leukemia, or chronic lymphocytic leukemia (CLL).

52. The method of claim 51, wherein the hematological disorder comprises a lymphoma.

53. The method of claim 52, wherein the lymphoma is selected from Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), a central nervous system lymphoma, small lymphocytic lymphoma (SLL), CD37+ dendritic cell lymphoma, lymphoplasmacytic lymphoma, splenic marginal zone lymphoma, extra-nodal marginal zone B-cell lymphoma of mucosa-associated (MALT) lymphoid tissue, nodal marginal zone B-cell lymphoma, follicular lymphoma, mantle cell lymphoma, diffuse large B-cell lymphoma, mediastinal (thymic) large B-cell lymphoma, precursor B-lymphoblastic lymphoma, immunoblastic large cell lymphoma, intravascular large B-cell lymphoma, primary effusion lymphoma, or Burkitt's lymphoma.

54. The method of claim 49, wherein the hematological malignancy comprises a myelodysplastic disorder,

55. The method of claim 54, wherein the myelodysplastic disorder is selected from refractory cytopenia with unilineage dysplasia (refractory anemia, refractory neutropenia, and refractory thrombocytopenia), refractory anemia with ring sideroblasts (RARS), refractory anemia with ring sideroblasts – thrombocytosis

(RARS-t), refractory cytopenia with multilineage dysplasia (RCMD), refractory cytopenia with multilineage dysplasia and ring sideroblasts (RCMD-RS), refractory anemia with excess blasts (RAEB), myelodysplasia unclassifiable, or refractory cytopenia of childhood.

56. The method of any one of claims 46-55, wherein the subject is receiving or previously received a hematopoietic cell transplant (HCT).

57. The method of claim 56, wherein the HCT comprises a donor hematopoietic cell comprising a chromosomal knockout of a gene that encodes an HLA component, a chromosomal knockout of a gene that encodes a TCR component, or both.

58. The method of any one of claims 46-57, wherein the subject had previously received lymphodepleting chemotherapy.

59. The method of claim 58, wherein the lymphodepleting chemotherapy comprised cyclophosphamide, fludarabine, anti-thymocyte globulin, or a combination thereof.

60. The method of any one of claims 46-59, wherein one or more of the engineered cells comprised in the unit dose is allogeneic to the subject.

61. A method for modulating an adoptive immunotherapy, the method comprising administering, to a subject who has previously received an engineered immune cell according to any one of claims 27-42 and comprising a heterologous polynucleotide encoding a safety switch protein, a cognate compound of the safety switch protein in an amount effective to ablate in the subject the previously administered engineered immune cell.

62. The method of claim 61, wherein:

- (a) the safety switch protein comprises tEGFR and the cognate compound is cetuximab;
- (b) the safety switch protein comprises iCasp9 and the cognate compound is AP1903;
- (c) the safety switch protein comprises a RQR polypeptide and the cognate compound is rituximab;
- (d) the safety switch protein comprises a myc binding domain and the cognate compound is an antibody specific for the myc binding domain; or
- (e) any combination of (a)-(d).

63. The method according to claim 62, wherein the safety switch comprises iCasp9 and the AP1903 bound to the iCasp9.

64. An isolated polynucleotide that encodes a binding protein, wherein the encoded binding protein comprises:

- (a) a T cell receptor (TCR) α -chain variable (V_α) domain having an amino acid sequence encoded by a TRAV17 gene, a TRAV21 gene, or a TRAV10 gene, and a TCR β -chain variable (V_β) domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86;
- (b) a TCR V_α domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V_β domain having an amino acid sequence encoded by a TRBV7-9 gene; or
- (c) a TCR V_α domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:87-92, and a TCR V_β domain comprising a CDR3 amino acid sequence of any one of SEQ ID NOS:13-17 and 86,

wherein the encoded binding protein is capable of specifically binding to a peptide containing an HA-1^H antigen and does not bind to a peptide that does not contain an HA-1^H antigen, and wherein the polynucleotide is codon-optimized for expression in a host cell.

65. The isolated polynucleotide of claim 64, wherein:
- (a) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:14, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:88;
 - (b) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:13, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:87;
 - (c) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:15, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:89;
 - (d) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:16, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:90; or
 - (e) the encoded V_{β} CDR3 comprises the amino acid sequence of SEQ ID NO:17, and the encoded V_{α} CDR3 comprises the amino acid sequence of SEQ ID NO:91.

66. The isolated polynucleotide of claim 64 or 65, wherein the encoded V_{α} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:2, 4, 6, 8, 10, and 12.

67. The isolated polynucleotide of any one of claims 64-66, wherein the encoded V_{β} domain comprises an amino acid sequence that has at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:1, 3, 5, 7, 9, and 11.

68. The isolated polynucleotide of claim 66 or 67, wherein the encoded V_{α} domain comprises no change in amino acid sequence of CDR1, the encoded V_{β} domain comprises no change in amino acid sequence of CDR1, or the CDR1 of the encoded V_{α} domain and the CDR2 of the encoded V_{β} domain comprise no change in amino acid sequence.

69. The isolated polynucleotide of any one of claims 66-68, wherein the encoded V_{α} domain comprises no change in amino acid sequence of CDR2, the encoded V_{β} domain comprises no change in amino acid sequence of CDR2, or the CDR2 of the encoded V_{α} domain and the CDR2 of the encoded V_{β} domain comprise no change in amino acid sequence.

70. The isolated polynucleotide of any one of claims 64-69, wherein:

(a) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:3, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:4;

(b) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:1, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:2;

(c) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:5, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:6;

(d) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:7, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:8;

(e) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:9, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:10;

(f) the encoded V_{β} domain comprises or consists of the amino acid sequence of SEQ ID NO:11, and the encoded V_{α} domain comprises or consists of the amino acid sequence of SEQ ID NO:12.

71. The isolated polynucleotide of any one of claims 64-70, wherein the encoded binding protein comprises a TCR α -chain constant domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:19-22, 24 and 26.

72. The isolated polynucleotide of any one of claims 64-71, wherein the encoded binding protein comprises a TCR β -chain constant domain having at least about 90% sequence identity to the amino acid sequence of any one of SEQ ID NOS:18, 23, and 25.

73. The isolated polynucleotide of any one of claims 64-72, wherein the encoded binding protein comprises a TCR α -chain having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36 and 38.

74. The isolated polynucleotide of any one of claims 64-73, wherein the encoded binding protein comprises a TCR α -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:28, 30, 32, 34, 36 and 38.

75. The isolated polynucleotide of claim 74, wherein the TCR α -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of any one of SEQ ID NOS:40, 42, 46, 48, 50, or 52.

76. The isolated polynucleotide of any one of claims 64-75, wherein the encoded binding protein comprises a TCR β -chain having an amino acid sequence that is at least about 90% identical to the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37.

77. The isolated polynucleotide of any one of claims 64-76, wherein the encoded binding protein comprises a TCR β -chain comprising or consisting of the amino acid sequence of any one of SEQ ID NOS:27, 29, 31, 33, 35, and 37.

78. The isolated polynucleotide of claim 77, wherein the TCR β -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 39, 41, 43, 45, 47, 49, or 51.

79. The isolated polynucleotide of any one of claims 64-78, wherein the encoded binding protein comprises:

(a) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:27, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:28;

(b) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:29, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:30;

(c) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:31, and the TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:32;

(d) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:33, and a TCR α -chain comprising or consisting of the amino acid sequence of SEQ ID NO:34;

(e) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:35, and a TCR- α chain comprising or consisting of the amino acid sequence of SEQ ID NO:36; or

(f) a TCR β -chain comprising or consisting of the amino acid sequence of SEQ ID NO:37, and a TCR- α chain comprising or consisting of the amino acid sequence of SEQ ID NO:38.

80. The isolated polynucleotide of any one of claims 64-79, comprising a polynucleotide encoding a self-cleaving peptide disposed between the polynucleotide encoding the TCR β -chain and the polynucleotide encoding the TCR α -chain.

81. The isolated polynucleotide of claim 80, wherein the encoded self-cleaving peptide comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 76-84.

82. The isolated polynucleotide of claim 80 or 81, wherein the single open reading frame comprises a nucleotide sequence that is at least about 80% identical to the nucleotide sequence of any one of SEQ ID NOS: 59-63.

83. The isolated polynucleotide of claim 82, wherein the single open reading frame comprises or consists of the nucleotide sequence of any one of SEQ ID NOS: 59-63.

84. The isolated polynucleotide of any one of claims 80-83, wherein the encoded ([TCR β -chain, self-cleaving-peptide, TCR α -chain]) comprises or consists of the amino acid sequence of any one of SEQ ID NOS: 53-57.

85. The isolated polynucleotide of any one of claims 64-84, wherein the encoded binding protein is capable of specifically binding to a HA-1^H peptide:HLA complex.

86. The isolated polynucleotide claim 85, wherein the HLA comprises HLA-A*0201.

87. The isolated polynucleotide of any one of claims 64-86, wherein the HA-1^H peptide comprises the amino acid sequence VLHDDLLEA (SEQ ID NO:66).

88. The isolated polynucleotide of any one of claims 64-87, wherein the encoded binding protein comprises a TCR, an antigen binding fragment of a TCR, or a chimeric antigen receptor (CAR).

89. The isolated polynucleotide of claim 88, wherein the antigen binding fragment of the TCR comprises a single chain TCR (scTCR).

90. The isolated polynucleotide of claim 88, wherein the encoded binding protein comprises a CAR.

91. The isolated polynucleotide of claim 88, wherein the encoded binding protein comprises a TCR.

92. The isolated polynucleotide of any one of claims 64-91, further comprising a polynucleotide encoding:

- (a) a safety switch protein;
- (b) a selection marker;
- (c) a CD8 co-receptor β -chain;
- (d) a CD8 co-receptor α -chain; or
- (e) any combination thereof.

93. The isolated polynucleotide of claim 92, wherein the encoded safety switch protein comprises:

- (i) a truncated EGF receptor (tEGFR);
- (ii) iCasp9;
- (iii) a RQR polypeptide;
- (iv) a myc epitope; or
- (v) any combination thereof.

94. The isolated polynucleotide of claim 92 or 93, wherein the encoded selection marker comprises:

- (i) a RQR polypeptide;
- (ii) a truncated low-affinity nerve growth factor (tNGFR);
- (iii) a truncated CD19 (tCD19);
- (iv) a truncated CD34 (tCD34); or
- (v) any combination thereof.

95. The isolated polynucleotide of any one of claims 92-94, wherein the encoded CD8 co-receptor comprises a β -chain comprising the amino acid sequence of any one of SEQ ID NOS:71-75.

96. The isolated polynucleotide of any one of claims 92-95, wherein the encoded CD8 co-receptor is a recombinant CD8 co-receptor comprising a RQR polypeptide having the amino acid sequence of SEQ ID NO:69.

97. The isolated polynucleotide of claim 96, wherein the polynucleotide encoding the RQR polypeptide is contained in a polynucleotide encoding a β -chain of the encoded CD8 co-receptor.

98. The isolated polynucleotide of claim 96 or 97, wherein the polynucleotide encoding the RQR polypeptide is contained in a polynucleotide encoding an α -chain of the encoded CD8-co-receptor.

99. The isolated polynucleotide of claim 98, comprising a polynucleotide encoding iCasp9 and a polynucleotide encoding a recombinant CD8 co-receptor protein that comprises a β -chain containing a RQR polypeptide.

100. The isolated polynucleotide of 99, comprising a single open reading frame containing, from 5' to 3', ([a polynucleotide encoding a safety switch protein]-[a polynucleotide encoding a self-cleaving peptide]-[the polynucleotide encoding a TCR β -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a TCR α -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide]-[a polynucleotide encoding a self-cleaving polypeptide]-[a polynucleotide encoding a CD8 α -chain]).

101. The isolated polynucleotide of claim 100, wherein the single open reading frame encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a porcine teschovirus 2A (P2A) peptide]-[a TCR β -chain]-[a P2A peptide]-[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]).

102. The isolated polynucleotide any one of claims 92-101, wherein the TCR β -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:41, and wherein the TCR α -chain-encoding polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:42.

103. An isolated polynucleotide, wherein the polynucleotide comprises or consists of the nucleotide sequence of SEQ ID NO:85.

104. The isolated polynucleotide of any one of claims 64-103, wherein the isolated polynucleotide is codon-optimized for expression in a T cell.

105. A transgene construct, comprising an expression control sequence operatively linked to the polynucleotide of any one of claims 64-104.

106. A transgene construct, comprising an expression control sequence operatively linked to a single open reading frame comprising:

- (a) a polynucleotide encoding a safety switch protein;
- (b) a polynucleotide encoding a TCR β -chain;
- (c) a polynucleotide encoding a TCR α -chain;
- (b) a polynucleotide encoding a selection marker;
- (c) a polynucleotide encoding a CD8 co-receptor β -chain; and
- (d) a polynucleotide encoding a CD8 co-receptor α -chain.

107. The transgene construct of claim 106, wherein the encoded safety switch protein comprises:

- (i) a truncated EGF receptor (tEGFR);
- (ii) iCasp9;
- (iii) a RQR polypeptide;
- (iv) a myc epitope; or
- (v) any combination thereof.

108. The transgene construct of claim 106 or 107, wherein the encoded selection marker comprises:

- (i) a RQR polypeptide;
- (ii) a truncated low-affinity nerve growth factor (tNGFR);
- (iii) a truncated CD19 (tCD19);
- (iv) a truncated CD34 (tCD34); or
- (v) any combination thereof.

109. The transgene construct of any one of claims 106-108, wherein the encoded CD8 co-receptor is a recombinant CD8 co-receptor comprising a RQR polypeptide having the amino acid sequence of SEQ ID NO:69.

110. The transgene construct of claim 109, wherein the encoded RQR polypeptide is contained in the encoded CD8 β -chain or in the encoded CD8 α -chain.

111. The transgene construct of claim 110, wherein the single open reading frame comprises:

- (a) a polynucleotide encoding a safety switch protein;
- (b) a polynucleotide encoding a TCR β -chain;
- (c) a polynucleotide encoding a TCR α -chain;
- (d) a polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide; and
- (e) a polynucleotide encoding a CD α -chain.

112. The transgene construct of claim 111, wherein the single open reading frame encodes, from 5' to 3', ([the polynucleotide encoding a safety switch protein]-[a polynucleotide encoding a self-cleaving peptide]-[the polynucleotide encoding a TCR β -chain]-[the polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a TCR α -chain]-[a polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a CD8 β -chain that contains an RQR polypeptide]-[a

polynucleotide encoding a self-cleaving polypeptide]-[the polynucleotide encoding a CD8 α -chain]).

113. The transgene construct of claim 112, wherein the single open reading frame encodes, from 5' to 3', ([an iCasp9 polypeptide]-[a P2A peptide]-[a TCR β -chain]-[a P2A peptide] -[a TCR α -chain]-[a P2A peptide]-[a CD8 β -chain comprising an RQR polypeptide]-[a P2A peptide]-[a CD8 α -chain]).

114. A vector, comprising the transgene construct of any one of claims 106-113.

115. The vector of claim 114, wherein the vector is capable of delivering the transgene construct to a host cell.

116. The vector of claim 115, wherein the host cell is a hematopoietic progenitor cell or a human immune system cell.

117. The vector of claim 116, wherein the human immune system cell is a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁻ CD8⁻ double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a dendritic cell, or any combination thereof.

118. The vector of claim 117, wherein the immune system cell is a CD4⁺ cell or a CD8⁺ T cell.

119. The vector of claim 117, wherein the T cell is a naïve T cell, a central memory T cell, an effector memory T cell, or any combination thereof.

120. The vector of any one of claims 114-119, wherein the vector is a viral vector.

121. The expression vector according to claim 120, wherein the viral vector is a lentiviral vector or a γ -retroviral vector.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C





FIG. 3

4/43

*FIG. 4*




FIG. 5A




FIG. 5B

6/43




FIG. 5C




FIG. 6A




FIG. 6B

FIG. 6C




FIG. 6E




FIG. 6D




FIG. 7




FIG. 8A




FIG. 8B




FIG. 8C

11/43




FIG. 8D

FIG. 8E




FIG. 9A




FIG. 9B

13/43




FIG. 10

14/43




FIG. 11A

15/43




FIG. 11B




FIG. 11C




FIG. 12A




FIG. 12B




FIG. 13




FIG. 14

20/43




FIG. 15




FIG. 16







FIG. 17

23/43

*FIG. 18*



TCR expression before sorting



TCR expression after sort and REP






FIG. 19







FIG. 21




FIG. 22





FIG. 23





FIG. 24




FIG. 25





FIG. 26




FIG. 27




FIG. 28A




FIG. 28B




FIG. 29




FIG. 30




FIG. 31




FIG. 32B




FIG. 32A




FIG. 32C




FIG. 32E

FIG. 32D




FIG. 33A




FIG. 33B




FIG. 33C




FIG. 33D




FIG. 33E




FIG. 34A




FIG. 34B




FIG. 35




FIG. 30