

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Angiogenesis inhibitors

(51)⁶ International Patent Classification(s)

C07D 405/04	20060101ALI2005100	
(2006.01)	8BMEP	A61P
A61K 31/416	1/04	
(2006.01)	20060101ALI2005122	
A61K 31/4162	0BMJP	A61P
(2006.01)	3/06	
A61K 31/4745	20060101ALI2005122	
(2006.01)	0BMJP	A61P
A61K 31/519	3/10	
(2006.01)	20060101ALI2005122	
A61P 1/04 (2006.01)	0BMJP	A61P
A61P 3/06 (2006.01)	9/00	
A61P 3/10 (2006.01)	20060101ALI2005122	
A61P 9/00 (2006.01)	0BMJP	A61P
A61P 9/10 (2006.01)	9/10	
A61P 15/00 (2006.01)	20060101ALI2005122	
A61P 17/06 (2006.01)	0BMJP	A61P
A61P 19/02 (2006.01)	20060101ALI2005122	
A61P 27/02 (2006.01)	0BMJP	A61P
A61P 29/00 (2006.01)	17/06	
A61P 35/00 (2006.01)	20060101ALI2005122	
C07D 405/04	0BMJP	A61P
20060101AFI2005122	19/02	
0BMJP	A61K	20060101ALI2005122
31/416	0BMJP	A61P
20060101ALI2005100	27/02	
8BMEP	A61K	20060101ALI2005122
31/4162	0BMJP	A61P
20060101ALI2008053	29/00	
1BMEP	A61K	20060101ALI2005122
31/4745	0BMJP	A61P
20060101ALI2005100	35/00	
8BMEP	A61K	20060101ALI2005122
31/519	0BMJP	
	PCT/US03/09968	

(21) Application No: 2003222148

(22) Application Date: 2003.03.31

(87) WIPO No: WO03/082274

(30) Priority Data

(31) Number	(32) Date	(33) Country
10/147,445	2002.05.16	US
60/368,892	2002.03.29	US

(43) Publication Date: 2003.10.13

(43) Publication Journal Date: 2003.11.20

(71) Applicant(s)

Carlsbad Technology, Inc.

(72) Inventor(s)

Teng, Che-Ming, Pan, Shiow-Lin, Guh, Jih-Hwa, Kuo, Shen-Chu, Lee, Fang, Yu

(74) Agent/Attorney

Shelston IP, 60 Margaret Street, Sydney, NSW, 2000

(56) Related Art

WO2001/053268

EP0667345
W02001/054680
JP2002/080367
W02001/002369
W02002/010137
EP1166785

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number
WO 03/082274 A1

(51) International Patent Classification⁷: A61K 31/416,
31/00

Sec. 1, Chung Shan Road, Tchia Taichung (TW). PAN,
Shlow-Lin [—/—]; 3F-1, No. 28m, Lane 265, Sin-Yi
Road Sec. 4, Taipei (TW). GUII, Jih-Iwa [—/—]; B1,
No. 60, Lane 394, Wu-Hsing Road, Taipei (TW).

(21) International Application Number: PCT/US03/09968

(22) International Filing Date: 31 March 2003 (31.03.2003) (74) Agent: TSAO, Y., Rocky, Fish & Richardson P.C., 225
Franklin Street, Boston, MA 02110-2804 (US).

(25) Filing Language: English

(81) Designated States (national): AE, AG, AL, AM, AT, AU,

(26) Publication Language: English

AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, NL, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(30) Priority Data:

60/368,892 29 March 2002 (29.03.2002) US
10/147,445 16 May 2002 (16.05.2002) US

(84) Designated States (regional): ARIPO patent (GII, GM,
KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, MI, MR, NF, SN, TD, TG).

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier applications:

US 60/368,892 (CON)
Filed on 29 March 2002 (29.03.2002)
US 10/147,445 (CON)
Filed on 16 May 2002 (16.05.2002)


Published:
— with international search report

(71) Applicant (for all designated States except US): CARLSBAD TECHNOLOGY, INC. [US/US]; 5923 Balfour Court, Carlsbad, CA 92008 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANGIOGENESIS INHIBITORS

WO 03/082274 A1



(57) Abstract: A method for treating an angiogenesis-related disorder. The method includes administrating to a subject in need thereof an effective amount of a compound of the formula (I). Each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thiienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl; each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NR'R', (CH₂)_nOR, (CH₂)_nSR, (CH₂)_nNR'R', (CH₂)_nCN, (CH₂)_nC(O)OR, (CH₂)_nCHO, (CH₂)_nCH=NR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(C₂H₄)_nO, in which each of R and R', independently, is H or C₁—C₆ alkyl; and n is 0, 1, 2, 3, 4, 5, or 6, and n is 0, 1, 2, or 3.

- 1 -

Angiogenesis Inhibitors

BACKGROUND

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common 5 general knowledge in the field.

Angiogenesis, formation of new blood vessels, occurs in the healthy body for healing wounds and restoring blood flow to tissues after injury. The angiogenic process is tightly controlled by various positive and negative regulatory factors. In many disease states, the body loses control over angiogenesis.

10 Excessive blood vessel growth may be triggered by certain pathological conditions such as cancer, age-related macular degeneration, rheumatoid arthritis, and psoriasis. As a result of excessive angiogenesis, new blood vessels feed diseased tissues and destroy normal tissues. In cancer, the new vessels allow tumor cells to escape into the circulation and lodge in other organs.


15 Angiogenesis occurs via a series of sequential steps, including division and migration of endothelial cells that form the walls of blood vessels. About 15 proteins are known to activate endothelial cell growth and movement. Therefore, angiogenesis can be suppressed by inhibitors of these activating proteins, e.g., angiogenin, epidermal growth factor, estrogen, fibroblast growth factor, interleukin 8, prostaglandins E1 and 20 E2, tumor necrosis factor, vascular endothelial growth factor, or granulocyte colony-stimulating factor.

Excessive angiogenesis-related disorders include cancer (both solid and hematologic tumors), cardiovascular diseases (e.g., atherosclerosis), chronic inflammation (e.g., rheumatoid arthritis or Crohn's disease), diabetes (e.g., diabetic 25 retinopathy), psoriasis, endometriosis, and adiposity. See, e.g., Pharmacological Reviews 52: 237-268, 2001. Compounds that effectively inhibit angiogenesis are drug candidates for treating or preventing these disorders.

SUMMARY

According to a first aspect, the present invention provides a method for treating 30 an angiogenesis-related disorder, comprising administrating to a subject in need thereof an effective amount of a compound of the formula:

- 1a -




wherein

each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

5 each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and

10 n is 0, 1, 2, or 3.

According to a second aspect, the present invention provides a method for treating cancer, comprising administrating to a subject in need thereof an effective amount of a compound of the formula:



15 wherein


each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and

- 1b -

n is 0, 1, 2, or 3.

According to a third aspect, the present invention provides use of a compound of the formula




5 wherein

each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

10 each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R' independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and

15 n is 0, 1, 2, or 3, for the manufacture of a medicament for the treatment of an angiogenesis-related disorder.

15 According to a fourth aspect, the present invention provides use of a compound of the formula



wherein

20 each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN,

- 1c -


$(CH_2)_nC(O)OR$, $(CH_2)_nCHO$, $(CH_2)_nCH=NOR$, or R_1 and R_2 together, R_3 and R_4 together, or R_5 and R_6 together are $O(CH_2)_mO$, in which each of R and R' , independently, is H or $C_1\sim C_6$ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and n is 0, 1, 2, or 3, for the manufacture of a medicament for the treatment of


5 cancer.

Unless the context clearly requires otherwise, throughout the description and the claims, the words "comprise", "comprising", and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".

10 This invention relates to methods of inhibiting angiogenesis with fused pyrazolyl compounds.

In one embodiment, this invention features a method for treating an angiogenesis-related disorder (e.g., cardiovascular disease, chronic inflammation, diabetes, psoriasis, endometriosis, or adiposity). The method includes administrating to
15 a subject in need thereof an effective amount of a compound of the formula:





Each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thiienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl; each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and n is 0, 1, 2, or 3. (CH₂)_m can be branched or linear. Note that the left atom shown in any substituted group described above is closest to the fused pyrazolyl ring. Also note that when there are one or more R or (CH₂)_m moieties in a fused pyrazolyl compound, the R or the (CH₂)_m moieties can be the same or different.

A subset of the above-described compounds are those in which each of Ar₁, Ar₂, and Ar₃ is phenyl or furyl. Further, each of R₁, R₂, R₅, and R₆ is H, and n is 1, e.g., 1-benzyl-3-(5'-hydroxymethyl-2'-furyl)indazole (Compound 1).

The term "Ar," as used herein, refers to both aryl and heteroaryl groups. Aryl, e.g., phenyl, is a hydrocarbon ring system having at least one aromatic ring. Heteroaryl is a hydrocarbon ring system having at least one aromatic ring which contains at least one heteroatom such as O, N, or S. Examples of heteroaryl include, but are not limited to, thiienyl, furyl, pyrrolyl, pyridinyl, and pyrimidinyl. An "Ar" may contain one, two, three, or more substituents on its ring. In addition to those assigned to R₁, R₂, R₃, R₄, R₅, and R₆ (see above), the substituents can also be nitro, C₂~C₆ alkenyl, C₂~C₆ alkynyl, aryl, heteroaryl, cyclyl, or heterocyclyl. Alkyl, alkenyl, alkynyl, alkoxy, aryl, heteroaryl, cyclyl, and heterocyclyl, as used herein, are optionally substituted with C₁~C₆ alkyl, halogen, amino, hydroxyl, mercapto, cyano, or nitro. Note that the term "alkyl" refers to both linear alkyl and branched alkyl.

The fused pyrazolyl compounds described above include the compounds themselves, as well as their salts and their prodrugs, if applicable. Such salts, for

example, can be formed by interaction between a negatively charged substituent (e.g., carboxylate) on a fused pyrazolyl compound and a cation. Suitable cations include, but are not limited to, sodium ion, potassium ion, magnesium ion, calcium ion, and an ammonium cation such as teteramethylammonium ion. Likewise, a positively charged 5 substituent (e.g., amino) can form a salt with a negatively charged counterion. Suitable counterions include, but are not limited to, chloride, bromide, iodide, sulfate, nitrate, phosphate, or acetate. Examples of prodrugs include esters and other pharmaceutically acceptable derivatives, which, upon administration to a subject, are capable of providing the fused pyrazolyl compounds described above.

10 The above-described compounds can also be used to treat cancer (e.g., lung cancer). More specifically, one or more of the compounds are administered an effective amount to a subject suffering from cancer.

As used herein, "cancer" refers to cellular tumor. Cancer cells having the capacity for autonomous growth, i.e., an abnormal state or condition characterized by 15 rapidly proliferating cell growth. The term is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type, or stage of invasiveness. Examples of cancers include, but are not limited to, carcinoma and sarcoma such as 20 leukemia, sarcomas, osteosarcoma, lymphomas, melanoma, ovarian cancer, skin cancer, testicular cancer, gastric cancer, pancreatic cancer, renal cancer, breast cancer, prostate colorectal cancer, cancer of head and neck, brain cancer, esophageal cancer, bladder cancer, adrenal cortical cancer, lung cancer, bronchus cancer, endometrial cancer, nasopharyngeal cancer, cervical or hepatic cancer, or cancer of unknown primary site.

Also within the scope of this invention are a composition containing one or 25 more of the fused pyrazolyl compounds described above for use in treating the aforementioned diseases, and the use of such a composition for the manufacture of a medicament for the just-described treatment.

Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF DRAWINGS

FIG. 1 shows the effect of Compound 1 on nude mice administered with a Matrigel plug containing 150 ng/mL vascular endothelial growth factor (VEGF) or basic fibroblast growth factor (bFGF).

5 FIG. 2 shows the effect of Compound 1 on nude mice implanted with A549 lung tumor cells.

DETAILED DESCRIPTION

A fused pyrazolyl compound used to practice the method of this invention can be prepared by procedures well known to a skilled person in the art (see, e.g., U.S. 10 Patent No. 5,574,168). They include the following synthetic route: An aryl aryl ketone is first prepared by coupling an arylcarbonyl chloride with another aryl compound. Either aryl compound is optionally mono- or multi-substituted. The ketone then reacts with an arylalkylhydrazine, the aryl group of which is also optionally mono- or multi-substituted, to form a hydrazone containing three aryl groups. The hydrazone group is 15 transformed into a fused pyrazolyl core via an alkylene linker, another aryl group is fused at 4-C and 5-C of the pyrazolyl core, and the third aryl group is directly connected to 3-C of the pyrazolyl core. Derivatives of the fused pyrazolyl compound may be obtained by modifying the substituents on any of the aryl groups.

The chemicals used in the above-described synthetic route may include, for 20 example, solvents, reagents, catalysts, protecting group and deprotecting group reagents. The methods described above may also additionally include steps, either before or after the steps described specifically herein, to add or remove suitable protecting groups in order to ultimately allow synthesis of the fused pyrazolyl compound. In addition, various synthetic steps may be performed in an alternate 25 sequence or order to give the desired compounds. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing applicable fused pyrazolyl compounds are known in the art and include, for example, those described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic 30 Synthesis*, 2d. Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette,

ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.

A fused pyrazolyl compound thus synthesized can be further purified by a method such as column chromatography, high pressure liquid chromatography, or 5 recrystallization.

This invention features a method for treating an angiogenesis-related disorder (e.g., cancer or an ocular disease). The method includes administering to a subject in need thereof an effective amount of one or more fused pyrazolyl compounds and a pharmaceutically acceptable carrier. The term "treating" is defined as the application 10 or administration of a composition including the fused pyrazolyl compound to a subject, who has a angiogenesis-related disease, a symptom of such a disease, or a predisposition toward such a disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptoms of the disease, or the predisposition toward the disease. "An effective amount" is defined as the 15 amount of a fused pyrazolyl compound which, upon administration to a subject in need thereof, is required to confer therapeutic effect on the subject. An effective amount of a fused pyrazolyl compound may range from about 1 mg/Kg to about 100 mg/Kg. Effective doses also vary, as recognized by those skilled in the art, depending on route 20 of administration, excipient usage, and the possibility of co-usage with other agents for treating an angiogenesis-related disorder.

To practice the method of the present invention, a fused pyrazolyl compound can be administered orally, parenterally, by inhalation spray, or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, 25 intrathecal, intralesional and intracranial injection or infusion techniques.

A composition for oral administration can be any orally acceptable dosage form including, but not limited to, tablets, capsules, emulsions and aqueous suspensions, dispersions and solutions. Commonly used carriers for tablets include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added to 30 tablets. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions or emulsions are administered orally, the active ingredient can be suspended or dissolved in an oily phase combined with

emulsifying or suspending agents. If desired, certain sweetening, flavoring, or coloring agents can be added.

A sterile injectable composition (e.g., aqueous or oleaginous suspension) can be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium (e.g., synthetic mono- or diglycerides). Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions can also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents.

An inhalation composition can be prepared according to techniques well-known in the art of pharmaceutical formulation and can be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

A carrier in a pharmaceutical composition must be "acceptable" in the sense of being compatible with the active ingredient of the formulation (and preferably, capable of stabilizing it) and not deleterious to the subject to be treated. For example, solubilizing agents, such as cyclodextrins (which form specific, more soluble complexes with fused pyrazolyl compounds), can be utilized as pharmaceutical excipients for delivery of fused pyrazolyl compounds. Examples of other carriers include colloidal silicon dioxide, magnesium stearate, cellulose, sodium lauryl sulfate, and D&C Yellow # 10.

A suitable *in vitro* assay can be used to preliminarily evaluate the efficacy of a fused pyrazolyl compound in inhibiting the activities of fibroblast growth factor (FGF) or vascular endothelial growth factor (VEGF). *In vivo* assays can also be performed by following procedures well known in the art to screen for efficacious fused pyrazolyl compounds. See the specific examples below.

Without further elaboration, it is believed that the above description has adequately enabled the present invention. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

5

Synthesis of 1-benzyl-3-(5'-hydroxymethyl-2'-furyl)indazole (Compound 1)

Calcium borohydride was first prepared by stirring anhydrous calcium chloride (88.8 mg, 0.8 mmole) with sodium borohydride (60 mg, 1.6 mmole) in anhydrous THF (20 mL) for 4 hrs. Then a 30 mL THF solution containing 88.0 mg 1-benzyl-3-(5'-methoxycarbonyl-2'-furyl)indazole (0.27 mmole) was added dropwise to the calcium borohydride solution at 30±2 °C. The mixture was heated under reflux for 6 hrs, cooled, quenched into crushed ice, placed at a reduced pressure to remove THF, and filtered to obtain a solid product. The solid was extracted with dichloromethane. The extract was concentrated to 50 mL and a solid precipitated after petroleum ether was added. The precipitate was collected and purified by column chromatography (silica gel-benzene) to obtain 70.0 mg 1-benzyl-3-(5'-hydroxymethyl-2'-furyl)indazole at a yield of 87%. mp: 108-109°C.

MS (%), m/z: 304 (M⁺).

IR (KBr) ν _{max}: 3350 cm⁻¹ (-OH).

20 ¹H-NMR (DMSO-d₆, 200 MHz) δ : 4.51 (2H, d, J=5.5 Hz, -CH₂O-), 5.31 (1H, t, J=5.5 Hz, -OH), 5.70 (2H, s, =NCH₂-), 6.48 (1H, d, J=3.4 Hz, H-4'), 6.97 (1H, d, J=3.4 Hz, H-3'), 7.21-7.31 (6H, m, H-5, phenyl), 7.45 (1H, t, J=8.2 Hz, H-6), 7.75 (1H, dd, J=8.2, 1.8 Hz, H-7), 8.12 (1H, dd, J=8.2, 1.0 Hz, C4-H).

25 **Inhibition of DNA synthesis**

Human umbilical vein endothelial cells (HUVECs) were incubated in the absence of Compound 1 (basal and control) or presence of Compound 1 (with a concentration of 0.1 μ M, 0.03 μ M, 0.1 μ M, 0.3 μ M, or 1 μ M). Vascular endothelial growth factor (VEGF) or basic fibroblast growth factor (bFGF) was added (except for basal) to induce DNA synthesis, which was detected based on [³H]thymidine incorporation. The results show that Compound 1 inhibited VEGF- and bFGF- induced

cell proliferation of HUVECs in a concentration-dependent manner. Unexpectedly, Compound 1 has IC₅₀ values of 9.0×10^{-8} M and 1.4×10^{-7} M, for VEGF and bFGF, respectively.

Additional 23 fused pyrazolyl compounds were also tested. All of them 5 inhibited VEGF-induced cell proliferation of HUVECs, some as potent as Compound 1.

Inhibition of tube formation

HUVECs were cultured onto chamberslide, which was pre-coated with Matrigel 10 (10 mg/mL). Cells were treated without Compound 1 (control) or with Compound 1 (10 μ M). VEGF (10 ng/mL) or bFGF (10 ng/mL) was added to induce tube formation. All photos were taken at 100X magnification. The results show that Compound 1 inhibited VEGF- and bFGF-induced formation of networks of elongated endothelial 15 cells.

15

Inhibition of angiogenic effect

Nude mice were subcutaneously injected with a Matrigel plug containing 150 ng/mL VEGF or bFGF. Vehicle or Compound 1 was administrated to the mice orally (1 mg/kg/day, 3 mg/kg/day, 10 mg/kg/day, 30 mg/kg/day, or 100 mg/kg/day) for seven 20 days. The angiogenic response was monitored visually through the transparent skin. Matrigel itself did not elicit an angiogenic response. After seven days the mice were sacrificed and the Matrigel plugs were observed in situ to quantify the ingrowth of blood vessels. The plugs were removed, fixed in 4% formaldehyde, embedded in paraffin, sectioned at 5- μ m thick for histological analysis, and blood vessel growth 25 quantitated by hematoxylin-eosin staining. All photos were taken at 40X magnification. The results show that oral administration of Compound 1 for seven days effectively inhibited VEGF or bFGF-induced angiogenic effect in a dose-dependent manner.

In a quantitative analysis of angiogenic effect, nude mice were treated as 30 described above, and the plugs were removed and dissolved. Hemoglobin concentrations were measured using a hemoglobin detection kit (Sigma Chem. Co.) as indices of angiogenesis. Means \pm S.E. (n=3) were presented (see FIG.1). Symbol

“***” represents $P<0.001$ that are compared with the control. The results illustrates that Compound 1 effectively inhibited VEGF or bFGF-induced angiogenic effect.

Anti-tumor activity

5 10^6 A549 lung tumor cells were introduced into the pleural space of nude mice. Compound 1 was administrated to the mice orally (10 mg/kg/day). The survival rates of Compound 1-treated mice and control mice were compared (FIG. 2). The life span (i.e., the medium survival time) of Compound 1-treated mice was about 1.8 times that of control mice, as analyzed by a %T/C value [(median survival time of
10 treatment/median survival time of control) x 100].


OTHER EMBODIMENTS

All of the features disclosed in this specification may be combined in any combination. Each feature disclosed in this specification may be replaced by an
15 alternative feature serving the same, equivalent, or similar purpose. Thus, unless expressly stated otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit
20 and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. For example, a compound structurally analogous to a fused pyrazolyl compound can also be used to practice the present invention. Thus, other embodiments are also within the claims.

WHAT IS CLAIMED IS:

1. A method for treating an angiogenesis-related disorder, comprising administrating to a subject in need thereof an effective amount of a compound of the formula:



5 wherein

each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN,

10 (CH₂)_mC(O)OR, (CH₂)_nCHO, (CH₂)_nCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and

n is 0, 1, 2, or 3.

15 2. The method of claim 1, wherein Ar₁ is phenyl.

3. The method of claim 2, wherein Ar₂ is furyl.

4. The method of claim 3, wherein Ar₂ is 5'-furyl.

20

5. The method of claim 4, wherein Ar₃ is phenyl.


6. The method of claim 5, wherein each of R₁, R₂, R₅, and R₆ is H.

25 7. The method of claim 6, wherein n is 1.

- 11 -

8. The method of claim 7, wherein one of R₃ and R₄ is substituted at position 2 of furyl.
9. The method of claim 8, wherein one of R₃ and R₄ is H, and the other is CH₂OH.
10. The method of claim 1, wherein Ar₂ is furyl.
- 5 11. The method of claim 10, wherein Ar₂ is 5'-furyl.
12. The method of claim 11, wherein one of R₃ and R₄ is substituted at position 2 of furyl.
13. The method of claim 1, wherein Ar₃ is phenyl.
14. The method of claim 13, wherein n is 1.
- 10 15. The method of any one of claims 1 to 14, wherein the angiogenesis-related disorder is a cardiovascular disease, chronic inflammation, diabetes, psoriasis, endometriosis, or adiposity.
16. A method for treating cancer, comprising administrating to a subject in need thereof an effective amount of a compound of the formula:

15



wherein

20 each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thiienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', 5 independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and n is 0, 1, 2, or 3.

17. The method of claim 16, wherein Ar₁ is phenyl.

10 18. The method of claim 17, wherein Ar₂ is furyl.

19. The method of claim 18, wherein Ar₂ is 5'-furyl.

20. The method of claim 19, wherein Ar₃ is phenyl.

15 21. The method of claim 20, wherein each of R₁, R₂, R₅, and R₆ is H.

22. The method of claim 21, wherein n is 1.

20 23. The method of claim 22, wherein one of R₃ and R₄ is substituted at position 2 of furyl.

24. The method of claim 23, wherein one of R₃ and R₄ is H, and the other is CH₂OH.

25 25. The method of claim 16, wherein Ar₂ is furyl.

26. The method of claim 25, wherein Ar₂ is 5'-furyl.


27. The method of claim 26, wherein one of R₃ and R₄ is substituted at position 2 of 30 furyl.

28. The method of claim 16, wherein Ar₃ is phenyl.

- 13 -

29. The method of claim 28, wherein n is 1.
30. The method of claim 16, wherein the cancer is lung cancer.
31. The method of claim 30, wherein Ar₁ is phenyl.
32. The method of claim 31, wherein Ar₂ is furyl.
- 5 33. The method of claim 32, wherein Ar₂ is 5'-furyl.
34. The method of claim 33, wherein Ar₃ is phenyl.
35. The method of claim 34, wherein each of R₁, R₂, R₅, and R₆ is H.
- 10 36. The method of claim 35, wherein n is 1.
37. The method of claim 36, wherein one of R₃ and R₄ is substituted at position 2 of furyl.
38. The method of claim 37, wherein one of R₃ and R₄ is H, and the other is CH₂OH.
39. The method of claim 30, wherein Ar₂ is furyl.
40. The method of claim 39, wherein Ar₂ is 5'-furyl.
- 15 41. The method of claim 40, wherein one of R₃ and R₄ is substituted at position 2 of furyl.
42. The method of claim 30, wherein Ar₃ is phenyl.
43. The method of claim 42, wherein n is 1.
44. Use of a compound of the formula

- 14 -




wherein

each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

5 each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and

10 n is 0, 1, 2, or 3, for the manufacture of a medicament for the treatment of an angiogenesis-related disorder.

45. Use of a compound of the formula



wherein

15 each of Ar₁, Ar₂, and Ar₃, independently, is phenyl, thienyl, furyl, pyrrolyl, pyridinyl, or pyrimidinyl;

each of R₁, R₂, R₃, R₄, R₅, and R₆, independently, is R, nitro, halogen, C(O)OR, C(O)SR, C(O)NRR', (CH₂)_mOR, (CH₂)_mSR, (CH₂)_mNRR', (CH₂)_mCN, (CH₂)_mC(O)OR, (CH₂)_mCHO, (CH₂)_mCH=NOR, or R₁ and R₂ together, R₃ and

- 15 -

R₄ together, or R₅ and R₆ together are O(CH₂)_mO, in which each of R and R', independently, is H or C₁~C₆ alkyl; and m is 0, 1, 2, 3, 4, 5, or 6, and n is 0, 1, 2, or 3, for the manufacture of a medicament for the treatment of cancer.

5 46. A method according to claim 1 or claim 16; or use according to claim 44 or claim 45, and substantially as herein described with reference to any one of the embodiments of the invention illustrated in the figures and/or examples.

