



(86) Date de dépôt PCT/PCT Filing Date: 2001/08/06
(87) Date publication PCT/PCT Publication Date: 2002/02/14
(45) Date de délivrance/Issue Date: 2017/07/11
(85) Entrée phase nationale/National Entry: 2003/02/05
(86) N° demande PCT/PCT Application No.: US 2001/024335
(87) N° publication PCT/PCT Publication No.: 2002/012478
(30) Priorité/Priority: 2000/08/07 (US60/223,483)

(51) Cl.Int./Int.Cl. C12N 15/00 (2006.01),
C12N 15/52 (2006.01), C12N 15/82 (2006.01)
(72) Inventeurs/Inventors:
BORONAT, ALBERT, ES;
CAMPOS, NARCISO, ES;
RODRIGUEZ-CONCEPCION, MANUEL, ES;
ROHMER, MICHEL, FR;
SEEMANN, MYRIAM, FR;
VALENTIN, HENRY E., US;
VENKATESH, TYAMAGONDLU V., US;
VENKATRAMESH, MYLAVARAPU, US
(73) Propriétaire/Owner:
MONSANTO TECHNOLOGY, LLC, US
(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : GENES DE LA VOIE METHYL-D-ERYTHRITOL PHOSPHATE
(54) Title: METHYL-D-ERYTHRITOL PHOSPHATE PATHWAY GENES

↓

Ecgcpe : MHNQAPIORRKSTRIYVGNVPIGDG : 25
Atgcpe : MATGVLPAVSGIKIPDSKVGFCKSMNLVRICDVRSLRSARRRVSVIRNSNQGSDLAELQPASEGSPLLVPRQKYCESLHKTVRKRTRTVMVGNVALGSE : 100

Ecgcpe : APIAVUSMINRTITDVEATVNOAKATERVAGADIVRVGVPTMDAAEAF.....KLEKQOVNVLVADITHFDYRTALKVAEYGVCLRTNPGNIGN..... : 114
Atgcpe : HPIRLEQIMTTSDTKDITGTVDEVMREADKAGADIVRITVQKKEADACFEIKDKLVQLNYNIPLVADIIHFAPTVALRVAEC.FUKIRVNPNGNFADRRRAQFE : 199

Ecgcpe : ERIRMVVLCAKDKNIPIRIGVNAGSLEKDIQEKYCEPTFOALLIESAMRHVDHLDRLNFDQEKVSVKASDVFLAVESVRLLAQO : 198
Atgcpe : TIDYTEDEYQKBLQHIEQVFTPEVEKCKKYGRAMRIGTNEGSI.SDRIMSYGDI.SPRGMVESAFEFARICRKLIDYHNEVFSMKASNPVIMVQAYRLLVAE : 298

Ecgcpe : I.....DQPLHLGITEAGGARSQAVKSAIGIGLILSEGIGDTLRVSLAADFVEETK..... : 249
Atgcpe : MYVHGWDYPLHLGVTEAGEGEDCRMKSAIGIGITLQDGLGDTLRVSLTEPPEEEDPCRRLANLGTAAKLQOGAPFEKRRHYFDFQRRRTGDLFVQKEG : 398

Ecgcpe : : -
Atgcpe : EEVDYRNVLHRDGSVLMSISLDQLKAPELLYRSLATKLVGMPFKDLATVDSILLRELPEVDDQVARLALKRLIDVSMGVIAPLSEQLTKPLPNAMVLVN : 498

Ecgcpe : : -
Atgcpe : LKELSGGAYKLLPEGTRLVVSRLRGDEPYEELEILKNIDATMILHDVPFTEDKVS RVHAAARRLFEBLSENSVNFVPIHHINFPTGIHRDELVIHAGTYAGG : 598

Ecgcpe : VGFDILKSLIRSRGINETACEHCSSRQBFDFVIGTVNATEQRLIEDITPMDVSTIGCVNGPGGEALVSTLGVTCGNK : 325
Atgcpe : LLVDGLGDGVMLEAPDQDFDLRNTSENHQQGCMRNTKTEYVSCSGRTLFDLQELSAELREKTSHPG.VSITATMGCVNGPGGEMADAFGVVGGSP : 697

Ecgcpe : .KSCLYEDG.VRKDRIDNNDMITOLEARTRAKASQLDEAPRTIDVQOVEK : 372
Atgcpe : GKIDLYVGKTVVKRGAMTATATATILIKKEHG.....RWMDPPVADE : 740

(57) Abrégé/Abstract:
A nucleic acid molecule comprising as operably linked components: (A) a promoter; and (B) a heterologous nucleic acid molecule that encodes an amino acid sequence of SEQ ID NO: 50, wherein said nucleic acid molecule is operably linked to a sequence encoding a chloroplast transit peptide, and wherein said promoter is a seed-specific promoter. Also disclosed are transformed cells and transgenic plant cells comprising the nucleic acid molecule.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 February 2002 (14.02.2002)

PCT

(10) International Publication Number
WO 02/012478 A2(51) International Patent Classification⁷: C12N 15/00

(21) International Application Number: PCT/US01/24335

(22) International Filing Date: 6 August 2001 (06.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/223,483 7 August 2000 (07.08.2000) US

(71) Applicant (for all designated States except US): MON-SANTO TECHNOLOGY, LLC [US/US]; 800 N. Lindbergh Boulevard, St. Louis, MO 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BORONAT, Albert [ES/ES]; Ronda Montmany, 43, E-08780 Pallesja (Barcelona) (ES). CAMPOS, Narciso [ES/ES]; C/ Riera Blanca 182, 4-3, E-08902 L'Hospitalet de Llobregat (Barcelona) (ES). RODRIGUEZ-CONCEPCION, Manuel [ES/ES]; Pl. Navas 7, 5-3, E-08004 Barcelona (ES). ROHMER, Michel [FR/FR]; 19, rue de l'Abbé

Hanauer, F-67100 Strasbourg (FR). SEEMAN, Myriam [FR/FR]; 10, rue des Ormes, F-68170 Rixheim (FR). VALENTIN, Henry, E. [DE/US]; 873 M Foxsprings Drive, Chesterfield, MO 63017 (US). VENKATESH, Tyamagondlu, V. [IN/US]; 12374 Summerhouse Drive, Apt. #12, St. Louis, MO 63146 (US). VENKATRAMESH, Mylavaram [US/US]; 901 Julianna Drive, Ballwin, MO 63011 (US).

(74) Agents: MARSH, David, R. et al.; Arnold & Porter, 555 12th Street, N.W., Washington, DC 20004 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

[Continued on next page]

(54) Title: METHYL-D-ERYTHRITOL PHOSPHATE PATHWAY GENES

Ecgcpe : MHNQAPIORRKSTRIYVGNVPTCDG : 25
Atgcpe : MATGVLPAPVSGIKIPDSKVGFGKSMNLVRICDVRSLRSARRRVSVIRNSNQSDLAELQPASEGSPLLVPRQKYESLHKTVERKTRTMVGNVALGSE : 100

Ecgcpe : APTAVQSMTNBRTTDMETVNOIKALERVGADIVRVSVETMDARDF.....KIKQDVNVPLVADIFHDYRTAALVASVGVCLRNPCNIGN..... : 114
Atgcpe : HPTRCMTTSDIKDITGTVDVEMRRADKAGADIVRTVQGGKEADACFEIKDKLVQLVNIPLVADIFHAPTVALRVASC.FEKERVNPGNFADRRAQFE : 199

Ecgcpe : BERIRMVDCARDKNIPTRIGVNAESLEKDKQEKVCEPTDOLISSAMRHVDHLDRLNPDQKVSVKASDVFLAVESYRLIAQ : 198
Atgcpe : TIDYTEDEYQKELQHIHOVFTPLVERCKKYGRAMRIGTNHGSLSDRFMSYVCE.SPRGMVESAFEPARICRKLIDYHNFVFSKASNPVIMVOYRLIVAE : 298

Ecgcpe : I.....DPLHLGTEAGGARSQAVKSAICGCLISECHGDTLRVSI.AADPVEETK..... : 249
Atgcpe : MYVHGWDYPLHLGTEAGEGEDGRMKAICGCLLOCGHGDTRLVSLTEPPEEBIDPCRRLANLGTAKAKLQGGAPFEKRRHYFDFQRTGDLPLVQKEG : 398

Ecgcpe : : -
Atgcpe : EEVDYRNVLRHDSVLSISLDQLKAPELLYRSLATKLTVGMPFKDLATVDSILRELPPVDDQVARLAKRLIDVSMGVIAPLSEQLTKPLPNAMVLVN : 498

Ecgcpe : : -
Atgcpe : LKELSGGAYKLLPBGTRLVVSLRGDEPYEELEILKNIDATMILHDVPFTEDKVS RVHAAARLFELSENSVNFPIHHINFPTGIHRDELVIHAGTYAGG : 598

Ecgcpe : VGFDIKSLRIRSRGINETACPTCSRQEPFVIGTVNAHEQRLDDITFMDVSIICCVNCPGEBALVSTLGVTCGNK : 325
Atgcpe : LLVDGLDGVMLEAPDQDFDRLNTSNLQGCMMRNTKTENVSCPSGRTLPDQEI SAETREKTSHPG.VSFAIMGCI/VNGPGEMADADRCYVGGSP : 697

Ecgcpe : .KSGLYEDG.VRDRDNDNMIDOLEARIRAKASQLEARRRIVQOVAK : 372
Atgcpe : GKIDLVGKTVVVRGAMTATDALISLKEHG.....RNVPPVADE : 740

(57) Abstract: The present invention provides and includes nucleic acids, proteins and antibodies associated with novel genes in the MEP pathway. The invention further encompasses methods utilizing such molecules, for example in gene isolation, gene analysis and the production of transgenic plants. The present invention also includes transgenic plants modified to express proteins associated with the MEP pathway.

WO 02/012478 A2

WO 02/012478 A2



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(15) Information about Correction:

see PCT Gazette No. 27/2002 of 4 July 2002, Section II

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(48) Date of publication of this corrected version:

4 July 2002

METHYL-D-ERYTHRITOL PHOSPHATE PATHWAY GENES

5 The present invention is in the field of plant genetics and biochemistry. More specifically, the invention relates to genes associated with the methyl-D-erythritol phosphate (MEP) pathway. The present invention provides and includes nucleic acid molecules, proteins, and antibodies associated with the genes of the MEP pathway and also provides methods utilizing such agents, for example in gene isolation, gene analysis and the
10 production of transgenic plants. Moreover, the present invention includes transgenic plants modified to express proteins associated with the MEP pathway and methods for the production of products from the MEP pathway.

 Tocopherols are an important component of mammalian diets. Epidemiological evidence indicates that tocopherol supplementation can result in decreased risk for
15 cardiovascular disease and cancer, can aid in immune function, and is associated with prevention or retardation of a number of degenerative disease processes in humans. Tocopherols function, in part, by stabilizing the lipid bilayer of biological membranes, reducing polyunsaturated fatty acid (PUFA) free radicals generated by lipid oxidation, and scavenging oxygen free radicals, lipid peroxy radicals and singlet oxygen species.

20 α -Tocopherol, often referred to as vitamin E, belongs to a class of lipid-soluble antioxidants that includes α , β , γ , and δ -tocopherols and α , β , γ , and δ -tocotrienols. Although α , β , γ , and δ -tocopherols and α , β , γ , and δ -tocotrienols are sometimes referred to collectively as "vitamin E", vitamin E is more appropriately defined chemically as α -tocopherol. α -Tocopherol is significant for human health, in part because it is readily
25 absorbed and retained by the body, and therefore has a higher degree of bioactivity than other tocopherol species. However, other tocopherols such as β , γ , and δ -tocopherols, also have significant health and nutritional benefits.

 Tocopherols are primarily synthesized only by plants and certain other photosynthetic organisms, including cyanobacteria. As a result, mammalian dietary
30 tocopherols are obtained almost exclusively from these sources. Plant tissues vary considerably in total tocopherol content and tocopherol composition, with α -tocopherol the predominant tocopherol species found in green, photosynthetic plant tissues. Leaf tissue can contain from 10-50 μ g of total tocopherols per gram fresh weight, but most of the world's major staple crops (*e.g.*, rice, corn, wheat, potato) produce low to extremely low levels of

total tocopherols, of which only a small percentage is α -tocopherol. Oil seed crops generally contain much higher levels of total tocopherols, but α -tocopherol is present only as a minor component in most oilseeds.

The recommended human daily dietary intake of 15-30 mg of vitamin E is quite
5 difficult to achieve from the average American diet. For example, it would take over 750 grams of spinach leaves in which α -tocopherol comprises 60% of total tocopherols, or 200-400 grams of soybean oil to satisfy this recommended daily vitamin E intake. While it is possible to augment the diet with supplements, most of these supplements contain primarily synthetic vitamin E, having eight stereoisomers, whereas natural vitamin E is predominantly
10 composed of only a single isomer. Furthermore, supplements tend to be relatively expensive, and the general population is disinclined to take vitamin supplements on a regular basis.

In addition to the health benefits of tocopherols, increased α -tocopherol levels in crops have been associated with enhanced stability and extended shelf life of fresh and processed plant products. Further, tocopherol supplementation of swine, beef, and poultry
15 feeds has been shown to significantly increase meat quality and extend the shelf life of post-processed meat products by retarding post-processing lipid oxidation, which contributes to undesirable flavor components.

Tocopherols are a member of the class of compounds referred to as the isoprenoids. Other isoprenoids include carotenoids, gibberellins, terpenes, chlorophyll and abscisic acid.
20 The chloroplasts of higher plants exhibit interconnected biochemical pathways leading to secondary metabolites including tocopherols. One tocopherol biosynthetic pathway in higher plants involves condensation of homogentisic acid and phytylpyrophosphate to form 2-methyl-6 phytylplastoquinol.

This plant tocopherol pathway can be divided into four parts: 1) synthesis of
25 homogentisic acid, which contributes to the aromatic ring of tocopherol; 2) synthesis of phytylpyrophosphate, which contributes to the side chain of tocopherol; 3) joining of HGA and phytylpyrophosphate via a prenyltransferase followed by a subsequent cyclization; 4) and S-adenosyl methionine-dependent methylation of an aromatic ring, which affects the relative abundance of each of the tocopherol species.

30 Homogentisic acid (HGA) is the common precursor to both tocopherols and plastoquinones. In at least some bacteria the synthesis of HGA is reported to occur via the conversion of chorismate to prephenate and then to p-hydroxyphenylpyruvate via a bifunctional prephenate dehydrogenase. Examples of bifunctional bacterial prephenate dehydrogenase enzymes include the proteins encoded by the *tyrA* genes of *Erwinia herbicola*
35 and *Escherichia coli*. The *tyrA* gene product catalyzes the production of prephenate from

chorismate, as well as the subsequent dehydrogenation of prephenate to form p-hydroxyphenylpyruvate (p-HPP), the immediate precursor to HGA. p-HPP is then converted to HGA by hydroxyphenylpyruvate dioxygenase (HPPD). In contrast, plants are believed to lack prephenate dehydrogenase activity, and it is generally believed that the synthesis in
5 plants of HGA from chorismate occurs via the synthesis and conversion of the intermediate arogenate. Because pathways involved in HGA synthesis are also responsible for tyrosine formation, any alterations in these pathways can also result in the alteration in tyrosine synthesis and the synthesis of other aromatic amino acids.

HGA is then combined with either phytyl-pyrophosphate or solanyl-pyrophosphate
10 by phytyl/prenyl transferase to form methyl-plastoquinols, which are precursors to plastoquinones and tocopherols. The major structural difference between each of the tocopherol species is the position of the methyl groups around the phenyl ring. This methylation process is S-adenosyl methionine-dependent. Methyl Transferase 1 (MT1) catalyzes the formation of plastoquinol-9 and γ -tocopherol by methylation of the 7 position.
15 Subsequent methylation at the 5 position of γ -tocopherol by γ -tocopherol methyl-transferase generates the biologically active α -tocopherol.

Phytylpyrophosphate, which is the central constituent of the tocopherol side chain, is formed from geranylgeranyldiphosphate (GGDP). GGDP is itself produced via a biosynthetic pathway in which isopentenyl diphosphate (IPP) plays a major role. IPP is a
20 central intermediate in the production of isoprenoids. Two pathways that generate IPP have been reported: a cytoplasmic-based pathway referred to as the mevalonate pathway; and a plastid-based pathway referred to as the MEP pathway. The cytoplasmic-based pathway involves the enzymes acetoacetyl CoA thiolase, HMGC_oA synthase, HMGC_oA reductase, mevalonate kinase, phosphomevalonate kinase, and mevalonate pyrophosphate
25 decarboxylase.

Evidence for the existence of an alternative, plastid-based, isoprenoid biosynthetic pathway recently emerged from studies in the research groups of Rohmer and Arigoni, who found that the isotope labeling patterns observed in studies on certain eubacterial and plant terpenoids could not be explained in terms of the mevalonate pathway. Eisenreich *et al.*,
30 *Chem. Bio.* 5:R221-233 (1998); Rohmer, *Prog. Drug. Res.* 50:135-154 (1998); Rohmer, 2 *Comprehensive Natural Products Chemistry* 45-68, Barton and Nakanishi (eds.), Pergamon Press, Oxford, England (1999). Arigoni and coworkers subsequently showed that 1-deoxyxylulose, or a derivative thereof, serves as an intermediate of the novel pathway, now referred to as the MEP pathway. Rohmer *et al.*, *Biochem. J.* 295:517-524 (1993); Schwarz,
35 Ph.D. thesis, Eidgenössische Technische Hochschule, Zurich, Switzerland (1994).

In the first step of the MEP pathway, DXP synthase, an enzyme encoded by the *dxs* gene, catalyzes the formation of 1-deoxy-D-xylulose-5-phosphate (DXP) from one molecule each of D-glyceraldehyde-3-phosphate and pyruvate. DXP is then converted into 2-C-methyl-D-erythritol-4-phosphate (MEP) by DXP reductoisomerase, which is encoded by the *dxr* gene. The conversion of MEP into 4-diphosphocytidyl-2-C-methyl-D-erythritol (CDP-ME) is catalyzed by CDP-ME synthase, which is encoded by the *ygbP* gene. CDP-ME kinase, which is encoded by the *ygbB* gene, catalyzes the conversion of CDP-ME into 4-diphosphocytidyl-2-C-methyl-D-erythritol 2-phosphate (CDP-MEP). CDP-MEP is then converted into 2-C-methyl-D-erythritol-2,4-cyclodiphosphate by ME-CDP synthase, which is encoded by the *ygbB* gene. The *ygbP* and *ygbB* genes are tightly linked on the *E. coli* genome. Herz *et al.*, *PNAS* 97(6):2485-2490 (2000).

Identification of further genes included in the MEP pathway will provide new approaches to increasing tocopherol levels in plants, which is a topic of the present application.

SUMMARY OF THE INVENTION

The present invention provides a novel gene essential to the MEP pathway: *gcpE*. *gcpE* is tightly linked to *ygbP* and *ygbB*. Expression of GCPE (protein) in organisms such as plants can increase the levels of tocopherol substrates such as isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) biosynthesis. The present invention also provides transgenic organisms expressing a GCPE protein, which can nutritionally enhance food and feed sources.

In particular, the present invention includes and provides a substantially purified nucleic acid molecule that encodes a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4 and 48 through 50. The present invention also includes and provides a substantially purified nucleic acid molecule that encodes a protein comprising an amino acid sequence of SEQ ID NO: 4. Further provided by the present invention is a substantially purified nucleic acid molecule that encodes a protein comprising an amino acid sequence of SEQ ID NO: 48.

The present invention includes and provides a substantially purified nucleic acid molecule that encodes a protein comprising an amino acid sequence of SEQ ID NO: 49. The present invention also includes and provides a substantially purified nucleic acid molecule that encodes a protein comprising an amino acid sequence of SEQ ID NO: 50. Further provided by the present invention is a substantially purified nucleic acid molecule that encodes a GCPE protein, where the nucleic acid molecule comprises a nucleic acid sequence

selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof.

The present invention includes and provides a recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter; and (B) a heterologous nucleic acid molecule that encodes an amino sequence selected from the group consisting of SEQ ID NOs: 4 and 48 through 50. The present invention also includes and provides transformed cells comprising such nucleic acid molecules.

Further provided by the present invention is a transgenic plant comprising a recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter; and (B) a heterologous nucleic acid molecule that encodes an amino sequence selected from the group consisting of SEQ ID NOs: 4 and 48 through 50.

The present invention includes and provides such a transgenic plant that exhibits an increased tocopherol level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. Also provided are seeds derived from such transgenic plants, and oil derived from such seeds. The present invention includes and provides such a transgenic plant that exhibits an increased monoterpene level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a transgenic plant that exhibits an increased carotenoid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a transgenic plant that exhibits an increased tocotrienol level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

The present invention includes and provides such a transgenic plant that produces a seed with an increased tocopherol level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a transgenic plant that produces a seed with an increased monoterpene level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a transgenic plant that produces a seed with an increased carotenoid level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a transgenic plant which produces a seed with an increased tocotrienol level relative to a plant with a similar genetic background but lacking the recombinant nucleic acid molecule.

The present invention includes and provides a recombinant nucleic acid molecule comprising as operably linked components: (A) an exogenous promoter; and (B) a nucleic

acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof. The present invention also includes and provides transformed cells comprising such nucleic acid molecules.

Further provided by the present invention is a transgenic plant comprising a
5 recombinant nucleic acid molecule comprising as operably linked components: (A) an exogenous promoter; and (B) a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof. The present invention includes and provides such a transgenic plant which is selected from the group consisting of *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed,
10 maize, mustard, oil palm, peanut, rapeseed, rice, safflower, sesame, soybean, sunflower, and wheat. The present invention includes and provides such a transgenic plant which is selected from the group consisting of coconut, crambe, maize, oil palm, peanut, rapeseed, safflower, sesame, soybean, and sunflower.

The present invention further includes and provides a seed derived from such a
15 transgenic plant. Also provided are oil and meal derived from such seeds. The present invention includes and provides such a seed which exhibits an increased tocopherol level relative to seed from a plant having a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a seed which exhibits an increased α -tocopherol level relative to seed from a plant having a similar genetic
20 background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a seed which exhibits an increased monoterpene level relative to seed from a plant having a similar genetic background but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a seed which exhibits an increased carotenoid level relative to seed from a plant having a similar genetic background
25 but lacking the recombinant nucleic acid molecule. The present invention includes and provides such a seed which exhibits an increased tocotrienol level relative to seed from a plant having a similar genetic background but lacking the recombinant nucleic acid molecule.

The present invention includes and provides a recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant cell to
30 cause production of an mRNA molecule; and (B) a nucleic acid sequence that hybridizes under moderate stringency conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof.

The present invention includes and provides a recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant cell to
35 cause production of an mRNA molecule; and (B) a nucleic acid sequence that has greater

than 85% identity to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof.

The present invention includes and provides a substantially purified protein comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 48, and 49. The present invention also includes and provides an antibody capable of specifically binding a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 48 and 49.

The present invention includes and provides a transgenic plant comprising a nucleic acid molecule that encodes a GCPE protein, where the nucleic acid molecule comprises a promoter operably linked to a heterologous nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof. The present invention includes and provides such a transgenic plant where the promoter is a seed-specific promoter. The present invention includes and provides such a transgenic plant where the seed-specific promoter is selected from the group consisting of napin, phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean α' subunit of b-conglycinin (soy 7s), and oleosin promoters.

The present invention includes and provides such a transgenic plant, where the plant exhibits an increased isoprenoid compound level relative to a plant with a similar genetic background but lacking the heterologous nucleic acid sequence. The present invention includes and provides such a transgenic plant, where the isoprenoid compound is selected from the group consisting of tocotrienols, tocopherols, terpenes, gibberellins, carotenoids, and xanthophylls. The present invention includes and provides such a transgenic plant, where the isoprenoid compound is a monoterpene. The present invention includes and provides such a transgenic plant, where the isoprenoid compound is selected from the group consisting of IPP and DMAPP. The present invention includes and provides such a transgenic plant, where the plant exhibits an increased tocopherol level relative to a plant with a similar genetic background but lacking the heterologous nucleic acid sequence. Also included and provided are feedstock, plant parts, and seeds derived from such plants. Further provided are containers of such seeds.

The present invention includes and provides a method of producing a transgenic plant with an increased isoprenoid compound level comprising: (A) transforming the plant with a nucleic acid molecule to produce a transgenic plant, where the nucleic acid molecule comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof; and (B) growing the transgenic plant.

The present invention includes and provides a method of producing a transgenic plant having seed with an increased isoprenoid compound level comprising: (A) transforming the plant with a nucleic acid molecule to produce a transgenic plant, where the nucleic acid molecule encodes a protein with an amino acid sequence selected from the group consisting of SEQ ID NOs: 4 and 48-50; and (B) growing the transgenic plant.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 sets forth chemical compounds that were determined as non-GCPE reaction products.

Figure 2 sets forth the diacetate of 2-methylbut-2-ene-1,4-diol.

Figure 3 sets forth (E)-1-(4-hydroxy-3-methylbut-2-enyl) diphosphate.

Figure 4 sets forth an alignment between proteins encoded by the *gcpE* gene from *E. coli* (SEQ ID NO: 78) and clone 135H1 from *A. thaliana* (SEQ ID NO: 79).

Figure 5 sets forth cloning of a truncated *Arabidopsis* cDNA to create pQE-AGH.

DESCRIPTION OF THE NUCLEIC AND AMINO ACID SEQUENCES

SEQ ID NO: 1 is an *Arabidopsis thaliana* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 2 is a rice nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 3 is an *E. coli* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 4 is an amino acid sequence derived from a rice *gcpE* gene.

SEQ ID NO: 5 is a partial *A. thaliana* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 6 is a partial soybean nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 7 is a partial tomato nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 8 is a partial *Mesembryanthemum crystallinum* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 9 is a partial rice nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 10 is a partial maize nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 11 is a partial Loblolly pine nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 12 is a partial *Physcomitrella patens* nucleotide sequence of a *gcpE* gene.

SEQ ID NOs: 13 through 20 are partial *A. thaliana* nucleotide sequences of a *gcpE* gene.

SEQ ID NOs: 21 through 32 are partial maize nucleotide sequences of a *gcpE* gene.

SEQ ID NOs: 33 through 46 are partial soybean nucleotide sequences of a *gcpE* gene.

SEQ ID NO: 47 is a partial *Brassica napus* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 48 is an amino acid sequence derived from an *A. thaliana gcpE* gene.

5 SEQ ID NO: 49 is an amino acid sequence derived from a rice *gcpE* gene.

SEQ ID NO: 50 is an amino acid sequence derived from an *E. coli gcpE* gene.

SEQ ID NOs: 51 through 77 are primer nucleotide sequences.

SEQ ID NO: 78 is an *E. coli* amino acid sequence derived from the *gcpE* gene.

SEQ ID NO: 79 is an *A. thaliana* amino acid sequence derived from clone 135H1.

10 SEQ ID NO: 80 is a partial *A. thaliana* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 81 is an amino acid sequence derived from an *A. thaliana gcpE* gene.

SEQ ID NO: 82 is a partial *A. thaliana* nucleotide sequence of a *gcpE* gene.

SEQ ID NO: 83 is an amino acid sequence derived from an *A. thaliana gcpE* gene.

SEQ ID NO: 84 is a partial *A. thaliana* nucleotide sequence of a *gcpE* gene.

15 SEQ ID NO: 85 is an amino acid sequence derived from an *A. thaliana gcpE* gene.

DEFINITIONS

The following definitions are provided as an aid to understanding the detailed description of the present invention.

The abbreviation "EP" refers to patent applications and patents published by the
20 European Patent Office, and the term "WO" refers to patent applications published by the World Intellectual Property Organization. "PNAS" refers to *Proc. Natl. Acad. Sci. (U.S.A.)*.

"Amino acid" and "amino acids" refer to all naturally occurring L-amino acids. This definition is meant to include norleucine, norvaline, ornithine, homocysteine, and homoserine.

25 "Chromosome walking" means a process of extending a genetic map by successive hybridization steps.

The phrases "coding sequence," "structural sequence," and "structural nucleic acid sequence" refer to a physical structure comprising an orderly arrangement of nucleic acids.

The coding sequence, structural sequence, and structural nucleic acid sequence may be
30 contained within a larger nucleic acid molecule, vector, or the like. In addition, the orderly arrangement of nucleic acids in these sequences may be depicted in the form of a sequence listing, figure, table, electronic medium, or the like.

A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity, *i.e.*, every nucleotide of one of the

molecules is complementary to a nucleotide of the other. Two molecules are “minimally complementary” if they can hybridize to one another with sufficient stability to remain annealed to one another under at least conventional “low-stringency” conditions. Similarly, the molecules are “complementary” if they can hybridize to one another with sufficient stability to remain annealed to one another under conventional “high-stringency” conditions. Conventional stringency conditions are described by Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Haymes *et al.*, *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, DC (1985).

10 The phrases “DNA sequence,” “nucleic acid sequence,” and “nucleic acid molecule” refer to a physical structure comprising an orderly arrangement of nucleic acids. The DNA sequence or nucleic acid sequence may be contained within a larger nucleic acid molecule, vector, or the like. In addition, the orderly arrangement of nucleic acids in these sequences may be depicted in the form of a sequence listing, figure, table, electronic medium, or the like. “Nucleic acid” refers to deoxyribonucleic acid (DNA) and ribonucleic acid (RNA).

15 An “elite soybean line” is any soybean line that has resulted from breeding and selection for superior agronomic performance. Elite soybean lines are commercially available to farmers or soybean breeders, *e.g.*, HARTZ™ variety H4452 Roundup Ready™ (HARTZ SEED, Stuttgart, Arkansas, USA); QP4544 (Asgrow Seeds, Des Moines, Iowa, USA); DeKalb variety CX445 (DeKalb, Illinois).

 “Exogenous genetic material” is any genetic material, whether naturally occurring or otherwise, from any source that is capable of being inserted into any organism.

 The term “expression” refers to the transcription of a gene to produce the corresponding mRNA and translation of this mRNA to produce the corresponding gene product (*i.e.*, a peptide, polypeptide, or protein). The term “expression of antisense RNA” refers to the transcription of a DNA to produce a first RNA molecule capable of hybridizing to a second RNA molecule. Formation of the RNA-RNA hybrid inhibits translation of the second RNA molecule to produce a gene product.

 “Fungi” as used herein includes the phyla *Ascomycota*, *Basidiomycota*, *Chytridiomycota* and *Zygomycota*, as well as the *Oomycota* and all mitosporic fungi, and “filamentous fungi” include all filamentous forms of the subdivision *Eumycota* and *Oomycota*. These terms are defined in Hawksworth *et al.*, in: Ainsworth and Bisby’s *Dictionary of The Fungi*, 8th edition, CAB International, University Press, Cambridge, UK (1995).

“Homology” refers to the level of similarity between two or more nucleic acid or amino acid sequences in terms of percent of positional identity (*i.e.*, sequence similarity or identity). Homology also refers to the concept of similar functional properties among different nucleic acids or proteins.

5 As used herein, a “homolog protein” molecule or fragment thereof is a counterpart protein molecule or fragment thereof in a second species (*e.g.*, maize GCPE is a homolog of *Arabidopsis* GCPE). A homolog can also be generated by molecular evolution or DNA shuffling techniques, so that the molecule retains at least one functional or structure characteristic of the original protein (*see, e.g.*, U.S. Patent No. 5,811,238).

10 The phrase “heterologous” refers to the relationship between two or more nucleic acid or protein sequences that are derived from different sources. For example, a promoter is heterologous with respect to a coding sequence if such a combination is not normally found in nature. In addition, a particular sequence may be “heterologous” with respect to a cell or organism into which it is inserted (*i.e.* does not naturally occur in that particular cell or
15 organism).

“Hybridization” refers to the ability of a strand of nucleic acid to join with a complementary strand via base pairing. Hybridization occurs when complementary nucleic acid sequences in the two nucleic acid strands contact one another under appropriate conditions.

20 The “MEP pathway” is the pathway associated with the biosynthesis of isopentenyl diphosphate or dimethylallyldiphosphate where deoxy-D-xylulose-5-phosphate or a derivative thereof serves as an intermediate.

 The phrase “operably linked” refers to the functional spatial arrangement of two or more nucleic acid regions or nucleic acid sequences. For example, a promoter region may be
25 positioned relative to a nucleic acid sequence such that transcription of a nucleic acid sequence is directed by the promoter region. Thus, a promoter region is “operably linked” to the nucleic acid sequence.

 “Phenotype” refers to traits exhibited by an organism resulting from the interaction of genotype and environment, such as disease resistance, pest tolerance, environmental
30 tolerance such as tolerance to abiotic stress, male sterility, quality improvement or yield *etc.*

 “Polyadenylation signal” or “polyA signal” refers to a nucleic acid sequence located 3’ to a coding region that promotes the addition of adenylate nucleotides to the 3’ end of the mRNA transcribed from the coding region.

 The term “promoter” or “promoter region” refers to a nucleic acid sequence, usually
35 found upstream (5’) to a coding sequence, which is capable of directing transcription of a

nucleic acid sequence into mRNA. The promoter or promoter region typically provide a recognition site for RNA polymerase and the other factors necessary for proper initiation of transcription. As contemplated herein, a promoter or promoter region includes variations of promoters derived by inserting or deleting regulatory regions, subjecting the promoter to
5 random or site-directed mutagenesis, *etc.* The activity or strength of a promoter may be measured in terms of the amounts of RNA it produces, or the amount of protein accumulation in a cell or tissue, relative to a promoter whose transcriptional activity has been previously assessed.

The term “protein” or “peptide molecule” includes any molecule that comprises five
10 or more amino acids. It is well known in the art that proteins may undergo modification, including post-translational modifications, such as, but not limited to, disulfide bond formation, glycosylation, phosphorylation, or oligomerization. Thus, as used herein, the term “protein” or “peptide molecule” includes any protein that is modified by any biological or non-biological process.

15 A “protein fragment” is a peptide or polypeptide molecule whose amino acid sequence comprises a subset of the amino acid sequence of that protein. A protein or fragment thereof that comprises one or more additional peptide regions not derived from that protein is a “fusion” protein.

“Recombinant vector” refers to any agent such as a plasmid, cosmid, virus,
20 autonomously replicating sequence, phage, or linear single-stranded, circular single-stranded, linear double-stranded, or circular double-stranded DNA or RNA nucleotide sequence. The recombinant vector may be derived from any source and is capable of genomic integration or autonomous replication.

“Regeneration” refers to the process of growing a plant from a plant cell or plant
25 tissue (e.g., plant protoplast or explant).

“Regulatory sequence” refers to a nucleotide sequence located upstream (5’), within, or downstream (3’) to a coding sequence. Transcription and expression of the coding sequence is typically impacted by the presence or absence of the regulatory sequence.

An antibody or peptide is said to “specifically bind” to a protein or peptide molecule
30 of the invention if such binding is not competitively inhibited by the presence of non-related molecules.

“Substantially homologous” refers to two sequences which are at least 90% identical in sequence, as measured by the BestFit program described herein (Version 10; Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI), using
35 default parameters.

“Substantially purified” refers to a molecule separated from substantially all other molecules normally associated with it in its native state. More preferably a substantially purified molecule is the predominant species present in a preparation. A substantially purified molecule may be greater than 60% free, preferably 75% free, more preferably 90% free, and most preferably 95% free from the other molecules (exclusive of solvent) present in the natural mixture. The term “substantially purified” is not intended to encompass molecules present in their native state.

“Transcription” refers to the process of producing an RNA copy from a DNA template. “Transformation” refers to the introduction of nucleic acid into a recipient host. The term “host” refers to bacteria cells, fungi, animals or animal cells, plants or seeds, or any plant parts or tissues including plant cells, protoplasts, calli, roots, tubers, seeds, stems, leaves, seedlings, embryos, and pollen.

“Transgenic” refers to organisms into which exogenous nucleic acid sequences are integrated. “Transgenic plant” refers to a plant where an introduced nucleic acid is stably introduced into a genome of the plant, for example, the nuclear or plastid genomes.

“Vector” refers to a plasmid, cosmid, bacteriophage, or virus that carries exogenous DNA into a host organism.

“Yeast” as used herein includes *Ascosporogenous* yeast (*Endomycetales*), *Basidiosporogenous* yeast and yeast belonging to the *Fungi Imperfecti* (*Blastomycetes*).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

One skilled in the art may refer to general reference texts for detailed descriptions of known techniques discussed herein or equivalent techniques. These texts include Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc. (1995); Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2d ed.), Cold Spring Harbor Press, Cold Spring Harbor, New York (1989); Birren *et al.*, *Genome Analysis: A Laboratory Manual*, volumes 1 through 4, Cold Spring Harbor Press, Cold Spring Harbor, New York (1997-1999); *Plant Molecular Biology: A Laboratory Manual*, Clark (ed.), Springer, New York (1997); Richards *et al.*, *Plant Breeding Systems* (2d ed.), Chapman & Hall, The University Press, Cambridge (1997); and Maliga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, New York (1995). These texts can, of course, also be referred to in making or using an aspect of the invention.

Utilizing a methodology for the isolation and characterization of essential MEP pathway genes, an essential and novel gene, termed *gcpE*, was isolated. *gcpE* is tightly

linked to *ygbP* and *ygbB*, which are other MEP pathway genes. As an essential MEP pathway component, enhanced expression or overexpression of GCPE in a variety of organisms such as plants can result in higher levels of tocopherol precursors such as IPP and DMAPP and ultimately in enhanced levels of tocopherols in such organisms. Moreover, the present invention provides a number of agents, for example, nucleic acid molecules encoding a GCPE protein, and provides uses of such agents.

The agents of the invention will preferably be "biologically active" with respect to either a structural attribute, such as the capacity of a nucleic acid to hybridize to another nucleic acid molecule, or the ability of a protein to be bound by an antibody (or to compete with another molecule for such binding). Alternatively, such an attribute may be catalytic and thus involve the capacity of the agent to mediate a chemical reaction or response. The agents will preferably be substantially purified. The agents of the invention may also be recombinant.

It is understood that any of the agents of the invention can be substantially purified and/or be biologically active and/or recombinant. It is also understood that the agents of the invention may be labeled with reagents that facilitate detection of the agent, *e.g.*, fluorescent labels, chemical labels, modified bases, and the like.

A. Nucleic Acid Molecules

Agents of the invention include nucleic acid molecules. In a preferred aspect of the present invention the nucleic acid molecule comprises a nucleic acid sequence which encodes a GCPE protein. In a preferred embodiment, the GCPE protein is derived from an organism having a MEP pathway. Examples of GCPE proteins are those proteins having an amino acid sequence selected from the group consisting of SEQ ID NO: 4, 48, 49, or 50.

In another preferred aspect of the present invention the nucleic acid molecule comprises a nucleic acid sequence that is selected from: (1) any of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof, or fragments of these sequences; (2) the group consisting of SEQ ID NOs: 1, 2, complements thereof, and fragments of these sequences; (3) the group consisting of SEQ ID NOs: 1, 2, 3, complements thereof and fragments of these sequences; (4) the group consisting of SEQ ID NOs: 1, 2, 13 through 47, complements thereof and fragments of these sequences; (5) the group consisting of SEQ ID NOs: 5 through 12, complements thereof and fragments of these sequences; or (6) the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof and fragments of these sequences.

In a further aspect of the present invention the nucleic acid molecule comprises a nucleic acid sequence encoding an amino acid sequence selected from: (1) any of SEQ ID

NOs: 4, 48, 49 or 50; (2) the group consisting of SEQ ID NO: 4, 48, and 49 and fragments of these sequences; or (3) the group consisting of SEQ ID NO: 4, 48, 49, 50 and fragments of these sequences.

It is understood that in a further aspect of the nucleic acid sequences of the present invention can encode a protein which differs from any of the proteins in that amino acid have been deleted, substituted or added without altering the function. For example, it is understood that codons capable of coding for such conservative amino acid substitutions are known in the art.

The present invention provides nucleic acid molecules that hybridize to the above-described nucleic acid molecules. Nucleic acid hybridization is a technique well known to those of skill in the art of DNA manipulation. The hybridization properties of a given pair of nucleic acids is an indication of their similarity or identity.

The nucleic acid molecules preferably hybridize, under low, moderate, or high stringency conditions, with a nucleic acid sequence selected from: (1) any of SEQ ID NOs: 1 through 3, 5 through 47, or complements thereof; (2) the group consisting of SEQ ID NOs: 1, 2, and complements thereof; (3) the group consisting of SEQ ID NOs: 1, 2, 3, and complements thereof; (4) the group consisting of SEQ ID NOs: 1, 2, 13 through 47, and complements thereof; (5) the group consisting of SEQ ID NOs: 5 through 12, and complements thereof; or (6) the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof. Fragments of these sequences are also contemplated.

The hybridization conditions typically involve nucleic acid hybridization in about 0.1X to about 10X SSC (diluted from a 20X SSC stock solution containing 3 M sodium chloride and 0.3 M sodium citrate, pH 7.0 in distilled water), about 2.5X to about 5X Denhardt's solution (diluted from a 50X stock solution containing 1% (w/v) bovine serum albumin, 1% (w/v) ficoll, and 1% (w/v) polyvinylpyrrolidone in distilled water), about 10 mg/mL to about 100 mg/mL fish sperm DNA, and about 0.02% (w/v) to about 0.1% (w/v) SDS, with an incubation at about 20°C to about 70°C for several hours to overnight. The stringency conditions are preferably provided by 6X SSC, 5X Denhardt's solution, 100 mg/mL fish sperm DNA, and 0.1% (w/v) SDS, with an incubation at 55°C for several hours.

The hybridization is generally followed by several wash steps. The wash compositions generally comprise 0.1X to about 10X SSC, and 0.01% (w/v) to about 0.5% (w/v) SDS with a 15 minute incubation at about 20°C to about 70°C. Preferably, the nucleic acid segments remain hybridized after washing at least one time in 0.1X SSC at 65°C. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 X SSC at 50°C to a high stringency of about 0.2 X SSC at 65°C. In addition, the

temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed.

5 Low stringency conditions may be used to select nucleic acid sequences with lower sequence identities to a target nucleic acid sequence. One may wish to employ conditions such as about 6.0 X SSC to about 10 X SSC, at temperatures ranging from about 20°C to about 55°C, and preferably a nucleic acid molecule will hybridize to one or more of the above-described nucleic acid molecules under low stringency conditions of about 6.0 X SSC
10 and about 45°C. In a preferred embodiment, a nucleic acid molecule will hybridize to one or more of the above-described nucleic acid molecules under moderately stringent conditions, for example at about 2.0 X SSC and about 65°C. In a particularly preferred embodiment, a nucleic acid molecule of the present invention will hybridize to one or more of the above-described nucleic acid molecules under high stringency conditions such as 0.2 X SSC and
15 about 65°C.

 In an alternative embodiment, the nucleic acid molecule comprises a nucleic acid sequence that is greater than 85% identical, and more preferably greater than 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through 3 and 5 through 47, complements thereof, and
20 fragments of any of these sequences.

 The percent identity is preferably determined using the "Best Fit" or "Gap" program of the Sequence Analysis Software Package™ (Version 10; Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI). "Gap" utilizes the algorithm of Needleman and Wunsch to find the alignment of two sequences that maximizes the
25 number of matches and minimizes the number of gaps. "BestFit" performs an optimal alignment of the best segment of similarity between two sequences and inserts gaps to maximize the number of matches using the local homology algorithm of Smith and Waterman. The percent identity calculations may also be performed using the Megalign program of the LASERGENE bioinformatics computing suite (default parameters,
30 DNASTAR Inc., Madison, Wisconsin). The percent identity is most preferably determined using the "Best Fit" program using default parameters.

 The present invention also provides nucleic acid molecule fragments that hybridize to the above-described nucleic acid molecules and complements thereof, fragments of nucleic acid molecules that exhibit greater than 80%, 85%, 90%, 95% or 99% sequence

identity with the above-described nucleic acid molecules and complements thereof, or fragments of any of these molecules.

Fragment nucleic acid molecules may consist of significant portion(s) of, or indeed most of, the nucleic acid molecules of the invention. In an embodiment, the fragments are
5 between about 3000 and about 1000 consecutive nucleotides, about 1800 and about 150 consecutive nucleotides, about 1500 and about 500 consecutive nucleotides, about 1300 and about 250 consecutive nucleotides, about 1000 and about 200 consecutive nucleotides, about 800 and about 150 consecutive nucleotides, about 500 and about 100 consecutive nucleotides, about 300 and about 75 consecutive nucleotides, about 100 and about 50
10 consecutive nucleotides, about 50 and about 25 consecutive nucleotides, or about 20 and about 10 consecutive nucleotides long of a nucleic molecule of the present invention.

In another embodiment, the fragment comprises at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 500, or 750 consecutive nucleotides of a nucleic acid sequence of the present invention.

15 Exemplary Uses

Nucleic acid molecules of the invention and fragments thereof may be employed to obtain other nucleic acid molecules from the same species (*e.g.*, nucleic acid molecules from maize may be utilized to obtain other nucleic acid molecules from maize). Exemplary nucleic acid molecules that may be obtained include, but are not limited to, nucleic acid
20 molecules that encode the complete coding sequence of a protein and promoters and flanking sequences of such molecules, and nucleic acid molecules that encode for other isozymes or gene family members.

Nucleic acid molecules of the invention and fragments thereof may also be employed to obtain nucleic acid homologs. Such homologs include the nucleic acid molecules of other
25 plants or other organisms, including the nucleic acid molecules that encode, in whole or in part, protein homologs of other plant species or other organisms, or sequences of genetic elements, such as promoters and transcriptional regulatory elements.

Promoters that may be isolated include, but are not limited to promoters of cell enhanced, cell specific, tissue enhanced, tissue specific, developmentally or environmentally
30 regulated expression profiles. Promoters obtained utilizing the nucleic acid molecules of the invention could also be modified to affect their control characteristics. Examples of such modifications would include but are not limited to enhancer sequences. Such genetic elements could be used to enhance gene expression of new and existing traits for crop improvement.

The above-described molecules can be readily obtained by using the above-described nucleic acid molecules or fragments thereof to screen cDNA or genomic libraries obtained from such plant species. These methods are known to those of skill in the art, as are methods for forming such libraries. In one embodiment, such sequences are obtained by incubating
5 nucleic acid molecules of the present invention with members of genomic libraries and recovering clones that hybridize to such nucleic acid molecules thereof. In a second embodiment, methods of chromosome walking or inverse PCR may be used to obtain such sequences.

Any of a variety of methods may be used to obtain one or more of the above-
10 described nucleic acid molecules. Automated nucleic acid synthesizers may be employed for this purpose. In lieu of such synthesis, the disclosed nucleic acid molecules may be used to define a pair of primers that can be used with the polymerase chain reaction to amplify and obtain any desired nucleic acid molecule or fragment.

In a preferred embodiment, nucleic acid molecules having SEQ ID NOs: 1 through 3
15 and 5 through 47, and complements thereof, and fragments of any of these sequences can be utilized to obtain such homologs. Such homolog molecules may differ in their nucleotide sequences from those found in one or more of SEQ ID NOs: 1 through 3, and 5 through 47 or complements thereof because complete complementarity is not needed for stable hybridization. The nucleic acid molecules of the invention therefore also include molecules
20 that, although capable of specifically hybridizing with the nucleic acid molecules may lack "complete complementarity."

In a preferred embodiment, the molecules are obtained from alfalfa, apple, *Arabidopsis*, banana, barley, *Brassica*, *Brassica campestris*, *Brassica napus*, broccoli, cabbage, canola, castor bean, chrysanthemum, citrus, coconut, coffee, cotton, crambe,
25 cranberry, cucumber, *Cuphea*, dendrobium, dioscorea, eucalyptus, fescue, fir, garlic, gladiolus, grape, hordeum, lentils, lettuce, liliacea, linseed, maize, millet, muskmelon, mustard, oat, oil palm, oilseed rape, onion, an ornamental plant, papaya, pea, peanut, pepper, perennial ryegrass, *Phaseolus*, pine, poplar, potato, rapeseed (including Canola and High Erucic Acid varieties), rice, rye, safflower, sesame, sorghum, soybean, strawberry, sugarbeet,
30 sugarcane, sunflower, tea, tomato, triticale, turf grasses, and wheat.

In a more preferred embodiment, the molecules are obtained from *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed, maize, mustard, oil palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), rice, safflower, sesame, soybean, sunflower, and wheat, and in a particularly preferred

embodiment from coconut, crambe, maize, oil palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), safflower, sesame, soybean, and sunflower.

The Sequence Analysis Software Package™ (Version 10; Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI) contains a number of
5 other useful sequence analysis tools for identifying homologs of the presently disclosed nucleotide and amino acid sequences. For example, programs such as “BLAST”, “FastA”, “TfastA”, “FastX”, and “TfastX” can be used to search for sequences similar to a query sequence. *See, e.g., Altschul et al., Journal of Molecular Biology* 215: 403-410 (1990); Lipman and Pearson, *Science* 227:1435-1441 (1985); Pearson and Lipman, 85:2444-2448
10 (1988); Pearson, “Rapid and Sensitive Sequence Comparison with FASTP and FASTA” in *Methods in Enzymology*, (R. Doolittle, ed.), 183:63-98, Academic Press, San Diego, California, USA (1990).

Short nucleic acid sequences having the ability to specifically hybridize to complementary nucleic acid sequences may be produced and utilized in the present
15 invention, *e.g.*, as probes to identify the presence of a complementary nucleic acid sequence in a given sample. Alternatively, the short nucleic acid sequences may be used as oligonucleotide primers to amplify or mutate a complementary nucleic acid sequence using PCR technology. These primers may also facilitate the amplification of related complementary nucleic acid sequences (*e.g.*, related sequences from other species).

20 Use of these probes or primers may greatly facilitate the identification of transgenic plants which contain the presently disclosed promoters and structural nucleic acid sequences. Such probes or primers may also be used to screen cDNA or genomic libraries for additional nucleic acid sequences related to or sharing homology with the presently disclosed promoters and structural nucleic acid sequences. The probes may also be PCR probes, which are
25 nucleic acid molecules capable of initiating a polymerase activity while in a double-stranded structure with another nucleic acid.

A primer or probe is generally complementary to a portion of a nucleic acid sequence that is to be identified, amplified, or mutated and of sufficient length to form a stable and sequence-specific duplex molecule with its complement. The primer or probe preferably is
30 about 10 to about 200 nucleotides long, more preferably is about 10 to about 100 nucleotides long, even more preferably is about 10 to about 50 nucleotides long, and most preferably is about 14 to about 30 nucleotides long.

The primer or probe may, for example without limitation, be prepared by direct chemical synthesis, by PCR (U.S. Patent Nos. 4,683,195 and 4,683,202), or by excising the
35 nucleic acid specific fragment from a larger nucleic acid molecule. Various methods for

determining the structure of PCR probes and PCR techniques exist in the art. Computer-generated searches using programs such as Primer 3, STSPipeline,

or GeneUp (Pesole *et al.*, *BioTechniques* 25:112-123, 1998), for example, can be used to identify potential PCR primers.

B. Protein and Peptide Molecules

Agents of the invention include proteins, peptide molecules, and fragments thereof encoded by nucleic acid agents of the invention. Preferred classes of protein and peptide molecules include: (1) GCPE proteins and peptide molecules; (2) GCPE proteins and peptide molecules derived from an organism having a MEP pathway; (3) GCPE proteins and peptide molecules derived from plants; and (4) GCPE proteins and peptide molecules derived from oilseed plants, including, but not limited to *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed, maize, mustard, oil palm, peanut, rapeseed, rice, safflower, sesame, soybean, sunflower, and wheat.

Other preferred proteins are those proteins having an amino acid sequence: (1) selected from the group consisting of SEQ ID NOs: 4, 48, 49, and 50; (2) selected from the group consisting of SEQ ID NOs: 4, 48 and 49; (3) selected from the group consisting of SEQ ID NOs: 4 and 49; (4) of SEQ ID NO: 4; (5) of SEQ ID NO: 48; (6) of SEQ ID NO: 49; and (7) of SEQ ID NO: 50.

In another preferred aspect of the present invention the protein or peptide molecule is encoded by a nucleic acid agent of the invention, including, but not limited to a nucleic acid sequence that is selected from: (1) any of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof, or fragments of these sequences; (2) the group consisting of SEQ ID NOs: 1, 2, complements thereof, and fragments of these sequences; (3) the group consisting of SEQ ID NOs: 1, 2, 3, complements thereof and fragments of these sequences; (4) the group consisting of SEQ ID NOs: 1, 2, 13 through 47, complements thereof and fragments of these sequences; (5) the group consisting of SEQ ID NOs: 5 through 12, complements thereof and fragments of these sequences; or (6) the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof and fragments of these sequences.

Any of the nucleic acid agents of the invention may be linked with additional nucleic acid sequences to encode fusion proteins. The additional nucleic acid sequence preferably encodes at least one amino acid, peptide, or protein. Many possible fusion combinations exist. For instance, the fusion protein may provide a "tagged" epitope to facilitate detection of the fusion protein, such as GST, GFP, FLAG, or polyHIS. Such fusions preferably encode

between 1 and about 50 amino acids, more preferably between about 5 and about 30 additional amino acids, and even more preferably between about 5 and about 20 amino acids.

Alternatively, the fusion may provide regulatory, enzymatic, cell signaling, or intercellular transport functions. For example, a sequence encoding a plastid transit peptide
5 may be added to direct a fusion protein to the chloroplasts within seeds. Such fusion partners preferably encode between 1 and about 1000 additional amino acids, more preferably between about 5 and about 500 additional amino acids, and even more preferably between about 10 and about 250 amino acids.

The above-described protein or peptide molecules may be produced via chemical
10 synthesis, or more preferably, by expression in a suitable bacterial or eukaryotic host. Suitable methods for expression are described by Sambrook *et al.*, *supra*, or similar texts. Fusion protein or peptide molecules of the invention are preferably produced via recombinant means. These proteins and peptide molecules may be derivatized to contain carbohydrate or other moieties (such as keyhole limpet hemocyanin, *etc.*).

Also contemplated are protein and peptide agents, including fragments and fusions
15 thereof, in which conservative, non-essential or non-relevant amino acid residues have been added, replaced or deleted. A further particularly preferred class of protein is a GCPE protein, in which conservative, non-essential or non-relevant amino acid residues have been added, replaced or deleted. Computerized means for designing modifications in protein
20 structure are known in the art. *See, e.g.*, Dahiyat and Mayo, *Science* 278:82-87 (1997).

A protein of the invention can also be a homolog protein. In a preferred embodiment, the nucleic acid molecules of the present invention, complements thereof, and fragments of these sequences can be utilized to obtain such homologs. In another preferred embodiment, the homolog is selected from the group consisting of alfalfa, apple,
25 *Arabidopsis*, banana, barley, *Brassica*, *Brassica campestris*, *Brassica napus*, broccoli, cabbage, canola, castor bean, chrysanthemum, citrus, coconut, coffee, cotton, crambe, cranberry, cucumber, Cuphea, dendrobium, dioscorea, eucalyptus, fescue, fir, garlic, gladiolus, grape, hordeum, lentils, lettuce, liliacea, linseed, maize, millet, muskmelon, mustard, oat, oil palm, oilseed rape, onion, an ornamental plant, papaya, pea, peanut, pepper,
30 perennial ryegrass, *Phaseolus*, pine, poplar, potato, rapeseed (including Canola and High Erucic Acid varieties), rice, rye, safflower, sesame, sorghum, soybean, strawberry, sugarbeet, sugarcane, sunflower, tea, tomato, triticale, turf grasses, and wheat.

In a more preferred embodiment, the homolog is selected from *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed, maize, mustard, oil
35 palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), rice, safflower,

sesame, soybean, sunflower, and wheat, and in a particularly preferred embodiment from coconut, crambe, maize, oil palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), safflower, sesame, soybean, and sunflower.

Agents of the invention include proteins comprising at least about a contiguous 10 amino acid region preferably comprising at least about a contiguous 20 amino acid region, even more preferably comprising at least about a contiguous 25, 35, 50, 75 or 100 amino acid region of a protein of the present invention. In another preferred embodiment, the proteins of the present invention include between about 10 and about 25 contiguous amino acid region, more preferably between about 20 and about 50 contiguous amino acid region, and even more preferably between about 40 and about 80 contiguous amino acid region.

Due to the degeneracy of the genetic code, different nucleotide codons may be used to code for a particular amino acid. A host cell often displays a preferred pattern of codon usage. Nucleic acid sequences are preferably constructed to utilize the codon usage pattern of the particular host cell. This generally enhances the expression of the nucleic acid sequence in a transformed host cell. Any of the above described nucleic acid and amino acid sequences may be modified to reflect the preferred codon usage of a host cell or organism in which they are contained. Modification of a nucleic acid sequence for optimal codon usage in plants is described in U.S. Patent No. 5,689,052. Additional variations in the nucleic acid sequences may encode proteins having equivalent or superior characteristics when compared to the proteins from which they are engineered.

It is understood that certain amino acids may be substituted for other amino acids in a protein or peptide structure (and the nucleic acid sequence that codes for it) without appreciable change or loss of its biological utility or activity. For example, amino acid substitutions may be made without appreciable loss of interactive binding capacity in the antigen-binding regions of antibodies, or binding sites on substrate molecules. The modifications may result in either conservative or non-conservative changes in the amino acid sequence. The amino acid changes may be achieved by changing the codons of the nucleic acid sequence, according to the codons given in Table 1.

Table 1: Codon degeneracy of amino acids

Amino acid	One letter	Three letter	Codons
Alanine	A	Ala	GCA GCC GCG GCT
Cysteine	C	Cys	TGC TGT
Aspartic acid	D	Asp	GAC GAT
Glutamic acid	E	Glu	GAA GAG
Phenylalanine	F	Phe	TTC TTT
Glycine	G	Gly	GGA GGC GGG GGT

Amino acid	One letter	Three letter	Codons
Histidine	H	His	CAC CAT
Isoleucine	I	Ile	ATA ATC ATT
Lysine	K	Lys	AAA AAG
Leucine	L	Leu	TTA TTG CTA CTC CTG CTT
Methionine	M	Met	ATG
Asparagine	N	Asn	AAC AAT
Proline	P	Pro	CCA CCC CCG CCT
Glutamine	Q	Gln	CAA CAG
Arginine	R	Arg	AGA AGG CGA CGC CGG CGT
Serine	S	Ser	AGC AGT TCA TCC TCG TCT
Threonine	T	Thr	ACA ACC ACG ACT
Valine	V	Val	GTA GTC GTG GTT
Tryptophan	W	Trp	TGG
Tyrosine	Y	Tyr	TAC TAT

It is well known in the art that one or more amino acids in a native sequence can be substituted with other amino acid(s), the charge and polarity of which are similar to that of the native amino acid, *i.e.*, a conservative amino acid substitution, resulting in a silent change. Conservative substitutes for an amino acid within the native polypeptide sequence can be selected from other members of the class to which the amino acid belongs. Amino acids can be divided into the following four groups: (1) acidic (negatively charged) amino acids, such as aspartic acid and glutamic acid; (2) basic (positively charged) amino acids, such as arginine, histidine, and lysine; (3) neutral polar amino acids, such as glycine, serine, threonine, cysteine, cystine, tyrosine, asparagine, and glutamine; and (4) neutral nonpolar (hydrophobic) amino acids such as alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine.

In a further aspect of the present invention, nucleic acid molecules of the present invention can comprise sequences that differ from those encoding a protein or fragment thereof selected from the group consisting of SEQ ID NOs: 4 and 48 through 50 due to the fact that the different nucleic acid sequence encodes a protein having one or more conservative amino acid changes.

In a preferred aspect, biologically functional equivalents of the proteins or fragments thereof of the present invention can have about 10 or fewer conservative amino acid changes, more preferably about 7 or fewer conservative amino acid changes, and most preferably about 5 or fewer conservative amino acid changes. In a preferred embodiment, the protein has between about 5 and about 500 conservative changes, more preferably between about 10 and about 300 conservative changes, even more preferably between about 25 and about 150 conservative changes, and most preferably between about 5 and about 25 conservative changes or between 1 and about 5 conservative changes.

Non-conservative changes include additions, deletions, and substitutions that result in an altered amino acid sequence. In a preferred embodiment, the protein has between about 5 and about 500 non-conservative amino acid changes, more preferably between about 10 and about 300 non-conservative amino acid changes, even more preferably between about 25 and about 150 non-conservative amino acid changes, and most preferably between about 5 and about 25 non-conservative amino acid changes or between 1 and about 5 non-conservative changes.

In making such changes, the role of the hydropathic index of amino acids in conferring interactive biological function on a protein may be considered. *See* Kyte and Doolittle, *J. Mol. Biol.* 157:105-132 (1982). It is accepted that the relative hydropathic character of amino acids contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, *e.g.*, enzymes, substrates, receptors, DNA, antibodies, antigens, *etc.* It is also understood in the art that the substitution of like amino acids may be made effectively on the basis of hydrophilicity, as the greatest local average hydrophilicity of a protein is known to correlate with a biological property of the protein. U.S. Patent No. 4,554,101.

Each amino acid has been assigned a hydropathic index and a hydrophilic value, as shown in Table 2.

Table 2: Amino Acid Hydropathic Indices and Hydrophilic Values

Amino acid	Hydropathic Index	Hydrophilic Value
Alanine	+1.8	-0.5
Cysteine	+2.5	-1.0
Aspartic acid	-3.5	+3.0 $\pm$ 1
Glutamic acid	-3.5	+3.0 $\pm$ 1
Phenylalanine	+2.8	-2.5
Glycine	-0.4	0
Histidine	-3.2	-0.5
Isoleucine	+4.5	-1.8
Lysine	-3.9	+3.0
Leucine	+3.8	-1.8
Methionine	+1.9	-1.3
Asparagine	-3.5	+0.2
Proline	-1.6	-0.5 $\pm$ 1
Glutamine	-3.5	+0.2
Arginine	-4.5	+3.0
Serine	-0.8	+0.3
Threonine	-0.7	-0.4
Valine	+4.2	-1.5
Tryptophan	-0.9	-3.4
Tyrosine	-1.3	-2.3

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic or hydrophilic index, score or value, and still result in a protein with similar biological activity, *i.e.*, still obtain a biologically functional protein. In making such changes, the substitution of amino acids whose hydropathic indices or
5 hydrophilic values are within ± 2 is preferred, those within ± 1 are more preferred, and those within ± 0.5 are most preferred.

As outlined above, amino acid substitutions are therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of the
10 foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine, and isoleucine.

These amino acid changes may be effected by mutating the nucleic acid sequence coding for the protein or peptide. Mutations to a nucleic acid sequence may be introduced in
15 either a specific or random manner, both of which are well known to those of skill in the art of molecular biology. Mutations may include deletions, insertions, truncations, substitutions, fusions, shuffling of motif sequences, and the like. A myriad of site-directed mutagenesis techniques exist, typically using oligonucleotides to introduce mutations at specific locations in a structural nucleic acid sequence. Examples include single strand rescue, unique site
20 elimination, nick protection, and PCR. Random or non-specific mutations may be generated by chemical agents (for a general review, *see* Singer and Kusmierek, *Ann. Rev. Biochem.* 52:655-693, 1982) such as nitrosoguanidine and 2-aminopurine; or by biological methods such as passage through mutator strains (Greener *et al.*, *Mol. Biotechnol.* 7:189-195, 1997).

C. Recombinant Vectors and Constructs

25 Exogenous and/or heterologous genetic material may be transferred into a host cell by use of a vector or construct designed for such a purpose. Any of the nucleic acid sequences described above may be provided in a recombinant vector. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the
30 genome of the host. Means for preparing recombinant vectors are well known in the art. Methods for making recombinant vectors particularly suited to plant transformation are described in U.S. Patent Nos.: 4,971,908, 4,940,835, 4,769,061 and 4,757,011.

Typical vectors useful for expression of nucleic acids in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of

Agrobacterium tumefaciens. Other vector systems suitable for introducing transforming DNA into a host plant cell include, but are not limited to the pCaMV CN transfer control vector, binary artificial chromosome (BIBAC) vectors (Hamilton *et al.*, *Gene* 200:107-116, 1997), and transfection with RNA viral vectors (Della-Cioppa *et al.*, *Ann. N.Y. Acad. Sci.* 792: 57-61, 1996). Additional vector systems also include plant selectable YAC vectors such as those described in Mullen *et al.*, *Molecular Breeding* 4:449-457 (1988).

A construct or vector may include a promoter, *e.g.*, a recombinant vector typically comprises, in a 5' to 3' orientation: a promoter to direct the transcription of a nucleic acid sequence of interest and a nucleic acid sequence of interest. Suitable promoters include, but are not limited to, those described herein. The recombinant vector may further comprise a 3' transcriptional terminator, a 3' polyadenylation signal, other untranslated nucleic acid sequences, transit and targeting nucleic acid sequences, selectable markers, enhancers, and operators, as desired.

The vector may be an autonomously replicating vector, *i.e.*, a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Alternatively, the vector may be one that, when introduced into the cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. This integration may be the result of homologous or non-homologous recombination.

Integration of a vector or nucleic acid into the genome by homologous recombination, regardless of the host being considered, relies on the nucleic acid sequence of the vector. Typically, the vector contains nucleic acid sequences for directing integration by homologous recombination into the genome of the host. These nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location or locations in one or more chromosomes. To increase the likelihood of integration at a precise location, there should be preferably two nucleic acid sequences that individually contain a sufficient number of nucleic acids, preferably about 400 bp to about 1500 bp, more preferably about 800 bp to about 1000 bp, which are highly homologous with the corresponding host cell target sequence. These nucleic acid sequences may be any sequence that is homologous with a host cell target sequence and, furthermore, may or may not encode proteins.

Vectors suitable for replication in mammalian cells may include viral replicons, or sequences that ensure integration of the appropriate sequences encoding HCV epitopes into

the host genome. For example, another vector used to express foreign DNA is vaccinia virus. Such heterologous DNA is generally inserted into a gene that is non-essential to the virus, for example, the thymidine kinase gene (tk), which also provides a selectable marker.

5 Expression of the HCV polypeptide then occurs in cells or animals that are infected with the live recombinant vaccinia virus.

In general, plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell are used in connection with bacterial hosts. The vector ordinarily carries a replication site, as well as marking sequences that are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, which contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, or other microbial plasmid or phage, also generally contains, or is modified to contain, promoters that can be used by the microbial organism for expression of the selectable marker genes.

15 Promoters

Promoters used in the context of the present invention are selected on the basis of the cell type into which the vector will be inserted. Promoters that function in bacteria, yeast, and plants are all taught in the art. The promoters may also be selected on the basis of their regulatory features, e.g., enhancement of transcriptional activity, inducibility, tissue specificity, and developmental stage-specificity. Additional promoters that may be utilized are described, for example, in U.S. Patent Nos. 5,378,619; 5,391,725; 5,428,147; 5,447,858; 5,608,144; 5,614,399; 5,633,441; 5,633,435; and 4,633,436.

Particularly preferred promoters in the recombinant vector include the nopaline synthase (*nos*) promoter; mannopine synthase (*mas*) promoter; octopine synthase (*ocs*) promoter; the cauliflower mosaic virus (CaMV) 19S and 35S promoters; the enhanced CaMV 35S promoter (eCaMV); the Figwort Mosaic Virus (FMV) 35S promoter; the light-inducible promoter from the small subunit of ribulose-1,5-bisphosphate carboxylase (ssRUBISCO); the EIF-4A promoter from tobacco; corn sucrose synthetase 1; corn alcohol dehydrogenase 1; corn light harvesting complex; corn heat shock protein; the chitinase promoter from *Arabidopsis*; the LTP (Lipid Transfer Protein) promoters from broccoli; petunia chalcone isomerase; bean glycine rich protein 1; potato patatin; the ubiquitin promoter from maize; the Adh promoter; the R gene complex promoter; and the actin promoter from rice.

The promoter is most preferably the *nos*, *ocs*, *mas*, CaMV19S, CaMV35S, eCaMV, ssRUBISCO, FMV, CaMV derived AS4, tobacco RB7, wheat POX1, tobacco EIF-4, lectin protein (Le1), or rice RC2 promoter. The promoter is preferably seed selective, tissue selective, constitutive, or inducible.

5 Often-used constitutive promoters include the CaMV 35S promoter, the eCaMV 35S promoter, the FMV promoter, the *mas* promoter, the *nos* promoter, and the *ocs* promoter, which is carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*.

Useful inducible promoters include promoters induced by salicylic acid or polyacrylic acids (PR-1), induced by application of safeners (substituted benzenesulfonamide herbicides), heat-shock promoters, a nitrate-inducible promoter derived from the spinach
10 nitrite reductase structural nucleic acid sequence, hormone-inducible promoters, and light-inducible promoters associated with the small subunit of RuBP carboxylase and LHCP families.

For the purposes of expression in specific tissues of the plant, such as the leaf, seed,
15 root or stem, it is preferred that the promoters utilized have relatively high expression in these specific tissues or organs. Examples reported in the literature include the chloroplast glutamine synthetase GS2 promoter from pea, the chloroplast fructose-1,6-biphosphatase (FBPase) promoter from wheat, the nuclear photosynthetic ST-LS1 promoter from potato, the serine/threonine kinase (PAL) promoter and the glucoamylase (CHS) promoter from *A.*
20 *thaliana*.

Also reported to be active in photosynthetically active tissues are the ribulose-1,5-bisphosphate carboxylase (RbcS) promoter from eastern larch (*Larix laricina*), the promoters for the *cab* genes of pine, wheat, spinach, and rice, the pyruvate orthophosphate dikinase (PPDK) promoter from maize, the promoter for the tobacco Lhcb1*2 gene, the *A. thaliana*
25 SUC2 sucrose-H⁺ symporter promoter and the promoter for the thylakoid membrane proteins from spinach (*psaD*, *psaF*, *psaE*, *PC*, *FNR*, *atpC*, *atpD*, *cab*, *rbcS*). Other promoters for the chlorophyll a/b-binding proteins may also be utilized in the invention, such as the promoters for *LhcB* gene and *PsbP* gene from white mustard.

For the purpose of expression in sink tissues of the plant, such as the tuber of the
30 potato plant, the fruit of tomato, or the seed of maize, wheat, rice and barley, it is preferred that the promoters utilized in the invention have relatively high expression in these specific tissues. A number of promoters for genes with tuber-specific or tuber-enhanced expression are known, including the class I patatin promoter, the promoter for the potato tuber ADPGPP genes, both the large and small subunits, the sucrose synthase promoter, the promoter for the
35 major tuber proteins including the 22 kd protein complexes and protease inhibitors, the

promoter for the granule-bound starch synthase gene (GBSS) and other class I and II patatins promoters.

Plant functional promoters useful for preferential expression in seeds include those from plant storage proteins and from proteins involved in fatty acid biosynthesis in oilseeds.

5 Examples of such promoters include the 5' regulatory regions from such genes as napin, phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean α' subunit of β -conglycinin (soy 7s), and oleosin. Further examples include the promoter for β -conglycinin and the lectin promoter from soybean. Seed-specific regulation is further discussed in EP 255 378.

10 Also included are promoters for the zeins, which are a group of storage proteins found in maize endosperm. Genomic clones for zein genes have been isolated and the promoters from these clones, including the 15 kD, 16 kD, 19 kD, 22 kD, 27 kD and genes, can also be used. Other promoters known to function, for example, in maize include the promoters for the following genes: *waxy*, *Brittle*, *Shrunken 2*, Branching enzymes I and II,
15 starch synthases, debranching enzymes, oleosins, glutelins and sucrose synthases. A particularly preferred promoter for maize endosperm expression is the promoter for the glutelin gene from rice, more particularly the Osgt-1 promoter.

Examples of promoters suitable for expression in wheat include those promoters for the ADPGlucose pyrosynthase (ADPGPP) subunits, the granule bound and other starch
20 synthase, the branching and debranching enzymes, the embryogenesis-abundant proteins, the gliadins and the glutenins. Preferred promoters in rice include promoters for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases and the glutelins, and particularly preferred is the promoter for rice glutelin, Osgt-1. Preferred promoters for barley include those promoters
25 for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases, the hordeins, the embryo globulins and the aleurone specific proteins.

Root specific promoters can also be used. An example of such a promoter is the promoter for the acid chitinase gene. Expression in root tissue can also be accomplished by
30 utilizing the root specific subdomains of the CaMV35S promoter that have been identified. Other root cell specific promoters include those reported by Conkling *et al.* *Plant Physiol.* 93:1203-1211 (1990).

Examples of suitable promoters for use with filamentous fungi are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase,
35 *Aspergillus niger* neutral alpha-amylase, *A. niger* acid stable alpha-amylase, *A. niger* or *A.*

awamori glucoamylase (glaA), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase and hybrids thereof. In a yeast host, preferred promoters include the *Saccharomyces cerevisiae* enolase (eno-1), the TAKA amylase, NA2-tpi (a hybrid of the promoters from the genes encoding *A. niger* neutral alpha-amylase and *A. oryzae* triose phosphate isomerase), glaA, *S. cerevisiae* GAL1 (galactokinase) and *S. cerevisiae* GPD (glyceraldehyde-3-phosphate dehydrogenase) promoters.

Suitable promoters for mammalian cells are also known in the art and include viral promoters, such as those from Simian Virus 40 (SV40), Rous sarcoma virus (RSV), adenovirus (ADV), cytomegalovirus (CMV), and bovine papilloma virus (BPV), as well as mammalian cell-derived promoters. Other preferred promoters include the hematopoietic stem cell-specific, *e.g.*, CD34, glucose-6-phosphatase, interleukin-1 alpha, CD11c integrin gene, GM-CSF, interleukin-5R alpha, interleukin-2, c-fos, h-ras, and DMD gene promoters.

Inducible promoters suitable for use with bacteria hosts include the β -lactamase and lactose promoter systems, the arabinose promoter system, alkaline phosphatase, a tryptophan (trp) promoter system and hybrid promoters such as the tac promoter. However, other known bacterial inducible promoters are suitable. Promoters for use in bacterial systems also generally contain a Shine-Dalgarno sequence operably linked to the DNA encoding the polypeptide of interest.

Examples of suitable promoters for an algal host are light harvesting protein promoters obtained from photosynthetic organisms, *Chlorella* virus methyltransferase promoters, CaMV 35 S promoter, PL promoter from bacteriophage λ , nopaline synthase promoter from the Ti plasmid of *A. tumefaciens*, and bacterial trp promoter.

Vectors for use with insect cells or insects may utilize a baculovirus transcriptional promoter including, *e.g.*, but not limited to the viral DNAs of *Autographa californica* MNPV, *Bombyx mori* NPV, *Trichoplusia ni* MNPV, *Rachiplusia ou* MNPV or *Galleria mellonella* MNPV, wherein the baculovirus transcriptional promoter is a baculovirus immediate-early gene IEl or IEN promoter; an immediate-early gene in combination with a baculovirus delayed-early gene promoter region selected from the group consisting of 39K and a *HindIII*-*k* fragment delayed-early gene; or a baculovirus late gene promoter.

Additional Nucleic Acid Sequences of Interest

The recombinant vector may also contain one or more additional nucleic acid sequences of interest. These additional nucleic acid sequences may generally be any sequences suitable for use in a recombinant vector. Such nucleic acid sequences include,

without limitation, any of the nucleic acid sequences, and modified forms thereof, described above. The additional nucleic acid sequences may also be operably linked to any of the above described promoters. The one or more additional nucleic acid sequences may each be operably linked to separate promoters. Alternatively, the additional nucleic acid sequences
5 may be operably linked to a single promoter (*i.e.* a single operon).

The additional nucleic acid sequences include, without limitation, those encoding seed storage proteins, fatty acid pathway enzymes, tocopherol biosynthetic enzymes, amino acid biosynthetic enzymes, and starch branching enzymes. Preferred seed storage proteins include zeins, 7S proteins, brazil nut protein, phenylalanine-free proteins, albumin, β -
10 conglycinin, 11S proteins, alpha-hordothionin, arcelin seed storage proteins, lectins, and glutenin. Preferred fatty acid pathway enzymes include thioesterases and desaturases.

Preferred tocopherol biosynthetic enzymes include *tyrA*, *slr1736*, *ATPT2*, *dxs*, *dxr*, *GGPPS*, *HPPD*, *GMT*, *MT1*, *AANT1*, *slr1737*, and an antisense construct for homogentisic acid dioxygenase. Preferred additional nucleic acid sequences encode MEP pathway proteins
15 including *ygbB*, *ygbP*, *ychB*, *yfgA*, *yfgB*, *dxs* and *dxr*. More preferred nucleic acid sequences include *yfgA* and *yfgB*, and still other preferred nucleic acid sequences include *ygbB*, *ychB* and *ygbP*. Preferred amino acid biosynthetic enzymes include anthranilate synthase, tryptophan decarboxylase, threonine decarboxylase, threonine deaminase, and aspartate kinase. Preferred starch branching enzymes include those set forth in U.S. Patent Nos. 6,232,122 and
20 6,147,279, and WO 97/22703.

Alternatively, the additional nucleic acid sequence may be designed to down-regulate a specific nucleic acid sequence. This is typically accomplished by operably linking the additional nucleic acid sequence, in an antisense orientation, with a promoter. One of ordinary skill in the art is familiar with such antisense technology. Any nucleic acid sequence may be
25 negatively regulated in this manner. Preferable target nucleic acid sequences contain a low content of essential amino acids, yet are expressed at relatively high levels in particular tissues. For example, β -conglycinin and glycinin are expressed abundantly in seeds, but are nutritionally deficient with respect to essential amino acids. This antisense approach may also be used to effectively remove other undesirable proteins, such as antifeedants (*e.g.*, lectins),
30 albumin, and allergens, from plant-derived foodstuffs.

Selectable and Screenable Markers

A vector or construct may also include a selectable marker. Selectable markers can also be used to select for plants or plant cells that contain the exogenous genetic material. Examples of such include, but are not limited to: a *neo* gene, which codes for kanamycin

resistance and can be selected for using kanamycin, RptII, G418, hpt *etc.*; a bar gene, which codes for bialaphos resistance; a mutant EPSP synthase gene, *aadA*, which encodes glyphosate resistance; a nitrilase gene, which confers resistance to bromoxynil; a mutant acetolactate synthase gene (ALS), which confers imidazolinone or sulphonylurea resistance, ALS, and a methotrexate resistant DHFR gene. The selectable marker is preferably GUS, green fluorescent protein (GFP), neomycin phosphotransferase II (*nptII*), luciferase (LUX), an antibiotic resistance coding sequence, or an herbicide (*e.g.*, glyphosate) resistance coding sequence. The selectable marker is most preferably a kanamycin, hygromycin, or herbicide resistance marker.

10 A vector or construct can also include a screenable marker. Screenable markers are useful to monitor expression. Exemplary screenable markers include: a β -glucuronidase or *uidA* gene (GUS), which encodes an enzyme for which various chromogenic substrates are known; an R-locus gene, which encodes a product that regulates the production of anthocyanin pigments (red color) in plant tissues; a β -lactamase gene, which encodes an enzyme for which various chromogenic substrates are known (*e.g.*, PADAC, a chromogenic cephalosporin); a luciferase gene; a *xyIE* gene, which encodes a catechol dioxygenase that can convert chromogenic catechols; an α -amylase gene; a tyrosinase gene, which encodes an enzyme capable of oxidizing tyrosine to DOPA and dopaquinone which in turn condenses to melanin; an α -galactosidase, which will turn a chromogenic α -galactose substrate.

20 Included within the terms "selectable or screenable marker genes" are also genes that encode a secretable marker whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers that encode a secretable antigen that can be identified by antibody interaction, or even secretable enzymes that can be detected catalytically. Secretable proteins fall into a number of classes, including small, diffusible proteins that are detectable, (*e.g.*, by ELISA), small active enzymes that are detectable in extracellular solution (*e.g.*, α -amylase, β -lactamase, phosphinothricin transferase), or proteins that are inserted or trapped in the cell wall (such as proteins which include a leader sequence such as that found in the expression unit of extension or tobacco PR-S). Other possible selectable and/or screenable marker genes will be apparent to those of skill in the art.

Other Elements in the Recombinant Vector

Various cis-acting untranslated 5' and 3' regulatory sequences may be included in the recombinant nucleic acid vector to produce desirable regulatory features. A vector or

construct may also include regulatory elements. Examples of such include the Adh intron 1, the sucrose synthase intron and the TMV omega element. These and other regulatory elements may be included when appropriate, and may be provided by the DNA sequence encoding the gene of interest or a convenient transcription termination region derived from a
5 different gene source.

A 3' non-translated region typically provides a transcriptional termination signal, and a polyadenylation signal that functions in plants to cause the addition of adenylate nucleotides to the 3' end of the mRNA. Such 3' non-translated regions can be obtained from the 3' regions of the nopaline synthase (*nos*) coding sequence, a soybean 7S α ' storage protein
10 coding sequence, the arcelin-5 coding sequence, the albumin coding sequence, and the pea ssRUBISCO E9 coding sequence. Particularly preferred 3' nucleic acid sequences include Arcelin-5 3', *nos* 3', E9 3', *adr12* 3', 7S α ' 3', 11S 3', USP 3', and albumin 3'.

Translational enhancers may also be incorporated as part of the recombinant vector, such as one or more 5' non-translated leader sequences that serve to enhance expression of
15 the nucleic acid sequence. Such enhancer sequences may be desirable to increase or alter the translational efficiency of the resultant mRNA. Preferred 5' nucleic acid sequences include dSSU 5', PetHSP70 5', and GmHSP17.9 5'. Such sequences can be derived from the promoter selected to express the gene or can be specifically modified to increase translation of the mRNA. Such regions can also be obtained from viral RNAs, from suitable eukaryotic
20 genes, or from a synthetic gene sequence. For a review of optimizing expression of transgenes, see Koziel *et al.*, *Plant Mol. Biol.* 32:393-405 (1996).

The recombinant vector can further comprise a nucleic acid sequence encoding a transit peptide. This peptide may be useful for directing a protein to the extracellular space, a plastid, or to some other compartment inside or outside of the cell. (see, *e.g.*, EP 0218571;
25 U.S. Patent Nos.: 4,940,835, 5,610,041, 5,618,988, and 6,107,060). The nucleic acid sequence in the recombinant vector may comprise introns. The introns may be heterologous with respect to the structural nucleic acid sequence. Preferred introns include the rice actin intron and the corn HSP70 intron.

A protein or fragment thereof encoding nucleic acid molecule of the invention may
30 also be operably linked to a suitable leader sequence. A leader sequence is a nontranslated region of a mRNA that is important for translation by the host. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the protein or fragment thereof. A polyadenylation sequence may also be operably linked to the 3' terminus of the nucleic acid sequence of the invention. The polyadenylation sequence is a sequence

that when transcribed is recognized by the host to add polyadenosine residues to transcribed mRNA.

A protein or fragment thereof encoding nucleic acid molecule of the invention may also be linked to a propeptide coding region. A propeptide is an amino acid sequence found at the amino terminus of a proprotein or proenzyme. Cleavage of the propeptide from the proprotein yields a mature biochemically active protein. The resulting polypeptide is known as a propolypeptide or proenzyme (or a zymogen in some cases). Propolypeptides are generally inactive and can be converted to mature active polypeptides by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide or proenzyme.

The recombinant vectors can further comprise one or more sequences that encode one or more factors that are advantageous in the expression of the protein or peptide, for example, an activator (*e.g.*, a trans-acting factor), a chaperone and a processing protease. An activator is a protein that activates transcription of a nucleic acid sequence encoding a polypeptide, a chaperone is a protein that assists another protein in folding properly, and a processing protease is a protease that cleaves a propeptide to generate a mature biochemically active polypeptide. The nucleic acids encoding one or more of these factors are preferably not operably linked to the nucleic acid encoding the protein or fragment thereof.

D. Transgenic Organisms, and Methods for Producing Same

One or more of the nucleic acid molecules or recombinant vectors of the invention may be used in plant transformation or transfection. For example, exogenous genetic material may be transferred into a plant cell and the plant cell regenerated into a whole, fertile or sterile plant. In a preferred embodiment, the exogenous genetic material includes a nucleic acid molecule of the present invention, preferably a nucleic acid molecule encoding a GCPE protein. In another preferred embodiment, the nucleic acid molecule has a sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof and fragments of these sequences. Other preferred exogenous genetic material are nucleic acid molecules that encode a protein or fragment thereof having an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, and 48 through 50 or fragments thereof.

The invention is also directed to transgenic plants and transformed host cells that comprise, in a 5' to 3' orientation, a promoter operably linked to a heterologous nucleic acid sequence of interest. Additional nucleic acid sequences may be introduced into the plant or host cell, such as 3' transcriptional terminators, 3' polyadenylation signals, other untranslated

nucleic acid sequences, transit or targeting sequences, selectable markers, enhancers, and operators. Preferred nucleic acid sequences of the present invention, including recombinant vectors, structural nucleic acid sequences, promoters, and other regulatory elements, are described above in parts A through C of the Detailed Description. Another embodiment of the invention is directed to a method of producing such transgenic plants which generally comprises the steps of selecting a suitable plant, transforming the plant with a recombinant vector, and obtaining the transformed host cell.

A transformed host cell may generally be any cell which is compatible with the present invention. A transformed host plant or cell can be or derived from a plant, or from a cell or organism such as a mammalian cell, mammal, fish cell, fish, bird cell, bird, algae cell, algae, fungal cell, fungus, or bacterial cell. Preferred host and transformants include: fungal cells such as *Aspergillus*, yeasts, mammals, particularly bovine and porcine, insects, bacteria, and algae. Methods to transform such cells or organisms are known in the art. See, e.g., EP 238023; Becker and Guarente, in: Abelson and Simon (eds.), *Guide to Yeast Genetics and Molecular Biology, Methods Enzymol.* 194: 182-187, Academic Press, Inc., New York; Bennett and LaSure (eds.), *More Gene Manipulations in Fungi*, Academic Press, CA, 1991; Hinnen *et al.*, *PNAS* 75:1920, 1978; Ito *et al.*, *J. Bacteriology* 153:163, 1983; Malardier *et al.*, *Gene* 78:147-156, 1989; Yelton *et al.*, *PNAS* 81:1470-1474, 1984.

Transfer of a nucleic acid that encodes a protein can result in expression or overexpression of that protein in a transformed cell, transgenic organism or transgenic plant. One or more of the proteins or fragments thereof encoded by nucleic acid molecules of the invention may be overexpressed in a transformed cell, transgenic organism or transgenic plant. Such expression or overexpression may be the result of transient or stable transfer of the exogenous genetic material.

In a preferred embodiment, expression or overexpression of a GCPE protein in a host provides in that host, relative to an untransformed host with a similar genetic background, an increased level of: (1) tocotrienols; (2) tocopherols; (3) α -tocopherols; (4) γ -tocopherols; (5) isopentenyl diphosphate (IPP); (6) DMAPP; (7) a GCPE protein in a plastid; (8) isoprenoids; (9) carotenoids; (10) an isoprenoid-related compound selected from the group consisting of IPP, DMAPP, and a GCPE protein; or (11) an isoprenoid compound selected from the group consisting of tocotrienols, tocopherols, terpenes, gibberellins, carotenoids, xanthophylls, α -tocopherols, γ -tocopherols, IPP, DMAPP, and a GCPE protein.

The expressed protein may be detected using methods known in the art that are specific for the particular protein or fragment. These detection methods may include the use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme

substrate. For example, if the protein has enzymatic activity, an enzyme assay may be used. Alternatively, if polyclonal or monoclonal antibodies specific to the protein are available, immunoassays may be employed using the antibodies to the protein. The techniques of enzyme assay and immunoassay are well known to those skilled in the art.

5 The resulting protein may be recovered by methods known in the arts. For example, the protein may be recovered from the nutrient medium by procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation. The recovered protein may then be further purified by a variety of chromatographic procedures, *e.g.*, ion exchange chromatography, gel filtration chromatography, affinity
10 chromatography, or the like. Reverse-phase high performance liquid chromatography (RP-HPLC), optionally employing hydrophobic RP-HPLC media, *e.g.*, silica gel, further purify the protein. Combinations of methods and means can also be employed to provide a substantially purified recombinant polypeptide or protein.

 In another preferred embodiment, overexpression of the GCPE protein in a
15 transgenic plant may provide tolerance to a variety of stresses, *e.g.*, oxidative stress tolerance such as to oxygen or ozone, UV tolerance, heat tolerance, drought tolerance, cold tolerance, or fungal/microbial pathogen tolerance.

 As used herein in a preferred aspect, a tolerance or resistance to stress is determined by the ability of a plant, when challenged by a stress such as cold, to produce a plant having a
20 higher yield than one without such tolerance or resistance to stress. In a particularly preferred aspect of the present invention, the tolerance or resistance to stress is measured relative to a plant with a similar genetic background to the tolerant or resistance plant except that the plant expresses or overexpresses a GCPE protein.

Host Cells and Organisms

25 Preferred host plants and cells can be or be derived from alfalfa, apple, *Arabidopsis*, banana, barley, *Brassica*, *Brassica campestris*, *Brassica napus*, broccoli, cabbage, canola, castor bean, chrysanthemum, citrus, coconut, coffee, cotton, crambe, cranberry, cucumber, Cuphea, dendrobium, dioscorea, eucalyptus, fescue, fir, garlic, gladiolus, grape, hordeum, lentils, lettuce, liliacea, linseed, maize, millet, muskmelon, mustard, oat, oil palm, oilseed
30 rape, onion, an ornamental plant, papaya, pea, peanut, pepper, perennial ryegrass, *Phaseolus*, pine, poplar, potato, rapeseed (including Canola and High Erucic Acid varieties), rice, rye, safflower, sesame, sorghum, soybean, strawberry, sugarbeet, sugarcane, sunflower, tea, tomato, triticale, turf grasses, and wheat.

In a more preferred embodiment, the host plants and cells are, or are derived from, *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed, maize, mustard, oil palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), rice, safflower, sesame, soybean, sunflower, and wheat, and in a particularly preferred embodiment from coconut, crambe, maize, oil palm, peanut, rapeseed (including Canola and High Erucic Acid varieties), safflower, sesame, soybean, and sunflower.

In another preferred embodiment, the plant or cell is or derived from canola. In another preferred embodiment, the plant or cell is or derived from *Brassica napus*. In a particularly preferred embodiment, the plant or cell is or derived from soybean. The soybean cell or plant is preferably a cell or plant of an elite soybean line.

Other preferred plants and plant host cells for use in the methods of the present invention include, but are not limited to Acacia, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beet, blackberry, blueberry, broccoli, brussel sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, cherry, chicory, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, garlic, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, nectarine, oat, oil palm, oilseed rape, okra, onion, orange, an ornamental plant, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, tritcale, turf, turnip, a vine, watermelon, wheat, yams, and zucchini.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines.

The fungal host cell may, for example, be a yeast cell, a fungi, or a filamentous fungal cell. In one embodiment, the fungal host cell is a yeast cell, and in a preferred embodiment, the yeast host cell is a cell of the species of *Candida*, *Kluyveromyces*, *Saccharomyces*, *Schizosaccharomyces*, *Pichia* and *Yarrowia*. In another embodiment, the fungal host cell is a filamentous fungal cell, and in a preferred embodiment, the filamentous fungal host cell is a cell of the species of *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Myceliophthora*, *Mucor*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium* and *Trichoderma*.

Suitable host bacteria include archaeobacteria and eubacteria, especially eubacteria and most preferably *Enterobacteriaceae*. Examples of useful bacteria include *Escherichia*, *Enterobacter*, *Azotobacter*, *Erwinia*, *Bacillus*, *Pseudomonas*, *Klebsiella*, *Proteus*, *Salmonella*, *Serratia*, *Shigella*, *Rhizobia*, *Vitreoscilla* and *Paracoccus*. Suitable *E. coli* hosts include *E. coli* W3110 (ATCC 27325), *E. coli* 294 (ATCC 31446), *E. coli* B and *E. coli* X1776 (ATCC 31537) (American Type Culture Collection, Manassas, Virginia). Mutant cells of any of the above-mentioned bacteria may also be employed. These hosts may be used with bacterial expression vectors such as *E. coli* cloning and expression vector BluescriptTM (Stratagene, La Jolla, CA); pIN vectors (Van Heeke and Schuster 1989), and pGEX vectors (Promega, Madison Wis.), which may be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST).

Preferred insect host cells are derived from *Lepidopteran* insects such as *Spodoptera frugiperda* or *Trichoplusia ni*. The preferred *Spodoptera frugiperda* cell line is the cell line Sf9 (ATCC CRL 1711). Other insect cell systems, such as the silkworm *B. mori* can also be used. These host cells are preferably used in combination with Baculovirus expression vectors (BEVs), which are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter in place of the viral gene, *e.g.*, polyhedrin (U.S. Patent No. 4,745,051).

Methods for Introducing Nucleic Acid Molecules into Organisms

Technology for introduction of nucleic acids into cells is well known to those of skill in the art. Common methods include chemical methods, microinjection, electroporation (U.S. Patent No. 5,384,253), particle acceleration, viral vectors, and receptor-mediated mechanisms. Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts and regeneration of the cell wall. The various techniques for transforming mammalian cells are also well known.

Algal cells may be transformed by a variety of known techniques, including but not limit to, microprojectile bombardment, protoplast fusion, electroporation, microinjection, and vigorous agitation in the presence of glass beads. Suitable procedures for transformation of green algal host cells are described in EP 108580. A suitable method of transforming cells of diatom *Phaeodactylum tricornutum* species is described in WO 97/39106. Chlorophyll C-containing algae may be transformed using the procedures described in U.S. Patent No. 5,661,017.

Methods for introducing nucleic acids into plants are also well known. Suitable methods include bacterial infection (*e.g.*, *Agrobacterium*), binary bacterial artificial

chromosome vectors, direct delivery of nucleic acids (*e.g.*, via PEG-mediated transformation), desiccation/inhibition-mediated nucleic acid uptake, electroporation, agitation with silicon carbide fibers, and acceleration of nucleic acid coated particles, *etc.* (reviewed in Potrykus *et al.*, *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42:205, 1991). For
5 example, electroporation has been used to transform maize protoplasts.

Alternatively, nucleic acids can be directly introduced into pollen by directly injecting a plant's reproductive organs. In another transformation technique, nucleic acids may also be injected into immature embryos. Plastids of higher plants can be stably transformed via particle gun delivery of DNA containing a selectable marker and targeting of
10 the DNA to the plastid genome through homologous recombination (U.S. Patent Nos. 5,451,513 and 5,545,818).

Methods for transforming dicots, primarily by use of *Agrobacterium tumefaciens* and obtaining transgenic plants, have been published for cotton, soybean, *Brassica*, peanut, papaya, pea and *Arabidopsis thaliana*. *E.g.*, U.S. Patent Nos. 5,004,863, 5,159,135,
15 5,416,011 5,463,174, 5,518,908, and 5,569,834. The latter method for transforming *Arabidopsis thaliana* is commonly called "dipping" or vacuum infiltration or germplasm transformation. Transformation of monocotyledons using electroporation, particle bombardment and *Agrobacterium* has also been reported. Transformation and plant regeneration have been achieved in asparagus, barley, maize, oat, orchard grass, rice, rye,
20 sugarcane, tall fescue, and wheat.

Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation and combinations of these treatments. Application of these systems to different plant strains depends upon the ability to regenerate that particular plant strain from protoplasts. Illustrative methods for the
25 regeneration of cereals from protoplasts are described in Abdullah *et al.*, *Biotechnology* 4:1087 (1986); Fujimura *et al.*, *Plant Tissue Culture Letters* 2:74 (1985); Toriyama *et al.*, *Theor Appl. Genet.* 205:34 (1986); and Yamada *et al.*, *Plant Cell Rep.* 4:85 (1986).

To transform plant strains that cannot be successfully regenerated from protoplasts, other ways to introduce DNA into intact cells or tissues can be utilized. For example, cereals
30 may be regenerated from immature embryos or explants. In addition, "particle gun" or high-velocity microprojectile technology can be utilized. Using the latter technology, DNA is carried through the cell wall and into the cytoplasm on the surface of small metal particles. The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants. A particular advantage of microprojectile
35 bombardment, in addition to it being an effective means of reproducibly transforming

monocots, is that neither the isolation of protoplasts (Christou *et al.*, *Plant Physiol.* 87:671-674, 1988), nor the susceptibility to *Agrobacterium* infection is required. *See also* Yang and Christou (eds.), *Particle Bombardment Technology for Gene Transfer*, Oxford Press, Oxford, England (1994).

5 An illustrative embodiment of a method for delivering DNA into maize cells by acceleration is a biolistics α -particle delivery system, which can be used to propel tungsten particles coated with DNA through a screen, such as a stainless steel or Nytex screen, onto a filter surface covered with corn cells cultured in suspension. Alternatively, immature embryos or other target cells may be arranged on solid culture medium. The screen disperses
10 the tungsten nucleic acid particles so that they are not delivered to the recipient cells in large aggregates. A particle delivery system suitable for use with the invention is the helium acceleration PDS-1000/He gun, which is available from Bio-Rad Laboratories (Bio-Rad, Hercules, California).

 Through the use of techniques set forth herein, one may obtain about 1000 or more
15 loci of cells transiently expressing a marker gene. The number of cells in a focus which express the exogenous gene product 48 hours post-bombardment often ranges from one to ten, and average one to three.

 In bombardment transformation, one may optimize the pre-bombardment culturing conditions and the bombardment parameters to yield the maximum numbers of stable
20 transformants. Important physical parameters to adjust include physical parameters such as gap distance, flight distance, tissue distance and helium pressure. In addition, biological factors, such as the nature of transforming DNA (e.g., linearized DNA or intact supercoiled plasmids) and the manipulation of cells before and immediately after bombardment, may affect transformation optimization. It is believed that pre-bombardment manipulations are
25 especially important for successful transformation of immature embryos. One may also minimize the trauma reduction factors by modifying conditions that influence the physiological state of the recipient cells and which may therefore influence transformation and integration efficiencies. For example, the osmotic state, tissue hydration and the subculture stage or cell cycle of the recipient cells may be adjusted for optimum
30 transformation.

Agrobacterium-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. Further, the integration of the Ti-DNA is a relatively precise process resulting in few rearrangements.

The region of DNA to be transferred is defined by the border sequences and intervening DNA is usually inserted into the plant genome as described.

Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations. Moreover, technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. Available vectors have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes. In addition, *Agrobacterium* containing both armed and disarmed Ti genes can be used for the transformations. In those plant strains where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer.

A transgenic plant formed using *Agrobacterium* transformation methods typically contains a single gene on one chromosome. Such transgenic plants can be referred to as being heterozygous for the added gene. More preferred is a transgenic plant that is homozygous for the added structural gene; *i.e.*, a transgenic plant that contains two added genes, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) an independent segregant, transgenic plant that contains a single added gene, germinating some of the seed produced and analyzing the resulting plants produced for the gene of interest.

Transgenic Plants

Regeneration, development, and cultivation of plants from single plant protoplast transformants or various transformed explants is taught in the art, *e.g.*, by Weissbach and Weissbach (eds.), *Methods for Plant Molecular Biology*, Academic Press, Inc., San Diego, CA (1988); and Horsch *et al.*, *Science* 227:1229-1231 (1985). There are a variety of methods for the regeneration of plants from plant tissue. The particular method of regeneration will depend on the starting plant tissue and the particular plant species to be regenerated.

Transformants are generally cultured in the presence of a selective media that selects for the successfully transformed cells and induces the regeneration of plant shoots. Such shoots are typically obtained within two to four months. Shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Many of the shoots will develop roots, which are then transplanted to soil

or other media to allow the continued development of roots. The method, as outlined, will generally vary depending on the particular plant employed.

Preferably, the regenerated transgenic plants are self-pollinated to provide homozygous transgenic plants. Alternatively, pollen obtained from the regenerated
5 transgenic plants may be crossed with seed-grown or non-transgenic plants, preferably plants of agronomically important lines. Conversely, pollen from seed-grown or non-transgenic plants may be used to pollinate the regenerated transgenic plants. A transgenic plant of the invention containing a desired polypeptide is cultivated using methods well-known to one skilled in the art.

10 A transgenic plant may pass along the nucleic acid sequence encoding the enhanced gene expression to its progeny. The transgenic plant is preferably homozygous for the nucleic acid encoding the enhanced gene expression and transmits that sequence to all of its offspring upon as a result of sexual reproduction. Progeny may be grown from seeds produced by the transgenic plant. These additional plants may then be self-pollinated to
15 generate a true breeding line of plants.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes that encode a polypeptide of interest. Back-crossing to a parental plant and out-
20 crossing with a non-transgenic plant are also contemplated, as is vegetative propagation.

The progeny from these plants are evaluated, among other things, for gene expression. The gene expression may be detected by several common methods such as western blotting, northern blotting, immunoprecipitation, and ELISA. Assays for gene expression based on the transient expression of cloned nucleic acid constructs have been
25 developed by introducing the nucleic acid molecules into plant cells by polyethylene glycol treatment, electroporation, or particle bombardment. Transient expression systems may be used to functionally dissect gene constructs (*see generally*, Maliga *et al.*, *Methods in Plant Molecular Biology, A Laboratory Course Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York, 1995).

30 Any of the nucleic acid molecules of the invention may be introduced into a plant cell in a permanent or transient manner in combination with other genetic elements such as vectors, promoters, enhancers, *etc.* Further, any of the nucleic acid molecules of the invention may be introduced into a plant cell in a manner that allows for expression or overexpression of the protein or fragment thereof encoded by the nucleic acid molecule, for
35 cosuppression of an endogenous protein, or for postranscriptional gene silencing of an

endogenous transcript. In addition, the activity of a protein in a plant cell may be reduced or depressed by growing a transgenic plant cell containing a nucleic acid molecule whose non-transcribed strand encodes a protein or fragment thereof.

5 Cosuppression is the reduction in expression levels, usually at the level of RNA, of a particular endogenous gene or gene family by the expression of a homologous sense construct that is capable of transcribing mRNA of the same strandedness as the transcript of the endogenous gene. Cosuppression may result from stable transformation with a single copy nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell or with multiple copies of a nucleic acid molecule that is homologous to a nucleic acid
10 sequence found with the cell. Genes, even though different, linked to homologous promoters may result in the cosuppression of the linked genes.

Posttranscriptional gene silencing (PTGS) can result in virus immunity or gene silencing in plants. PTGS is induced by dsRNA and is mediated by an RNA-dependent RNA polymerase, present in the cytoplasm, that requires a dsRNA template. The dsRNA is formed
15 by hybridization of complementary transgene mRNAs or complementary regions of the same transcript. Duplex formation can be accomplished by using transcripts from one sense gene and one antisense gene colocated in the plant genome, a single transcript that has self-complementarity, or sense and antisense transcripts from genes brought together by crossing. The dsRNA-dependent RNA polymerase makes a complementary strand from the transgene
20 mRNA and RNase molecules attach to this complementary strand (cRNA). These cRNA-RNase molecules hybridize to the endogene mRNA and cleave the single-stranded RNA adjacent to the hybrid. The cleaved single-stranded RNAs are further degraded by other host RNases because one will lack a capped 5' end and the other will lack a poly(A) tail. See Waterhouse *et al.*, *PNAS* 95: 13959-13964 (1998).

25 Antisense approaches are a way of preventing or reducing gene function by targeting the genetic material. The objective of the antisense approach is to use a sequence complementary to the target gene to block its expression and create a mutant cell line or organism in which the level of a single chosen protein is selectively reduced or abolished. Antisense techniques have several advantages over other 'reverse genetic' approaches. The
30 site of inactivation and its developmental effect can be manipulated by the choice of promoter for antisense genes or by the timing of external application or microinjection. Antisense can manipulate its specificity by selecting either unique regions of the target gene or regions where it shares homology to other related genes.

Under one embodiment, the process involves the introduction and expression of an
35 antisense gene sequence. Such a sequence is one in which part or all of the normal gene

sequences are placed under a promoter in inverted orientation so that the 'wrong' or complementary strand is transcribed into a noncoding antisense RNA that hybridizes with the target mRNA and interferes with its expression. An antisense vector can be constructed by standard procedures and introduced into cells by transformation, transfection, electroporation, microinjection, infection, *etc.* The type of transformation and choice of vector will determine whether expression is transient or stable. The promoter used for the antisense gene may influence the level, timing, tissue, specificity, or inducibility of the antisense inhibition.

Feed, Meal, Protein and Oil Preparations

Plants or agents of the present invention can be utilized in methods, for example without limitation, to obtain a seed that expresses a *gcpE* nucleic acid molecule in that seed, to obtain a seed enhanced in a product of a *gcpE* gene, to obtain meal enhanced in a product of a *gcpE* gene, to obtain feedstock enhanced in a product of a *gcpE* gene, and to obtain oil enhanced in a product of a *gcpE* gene.

The present invention also provides for parts of the plants, particularly reproductive or storage parts, of the present invention. Plant parts, without limitation, include seed, endosperm, mesocarp, ovule and pollen. In a particularly preferred embodiment of the present invention, the plant part is a seed. In one embodiment the seed is a constituent of animal feed. In another embodiment, the plant part is a fruit, more preferably a fruit with enhanced shelf life. In another preferred embodiment, the fruit has increased levels of a tocopherol.

Plants utilized in such methods may be processed. A plant or plant part may be separated or isolated from other plant parts. A preferred plant part for this purpose is a seed. It is understood that even after separation or isolation from other plant parts, the isolated or separated plant part may be contaminated with other plant parts. In a preferred aspect, the separated plant part is greater than about 50% (w/w) of the separated material, more preferably, greater than about 75% (w/w) of the separated material, and even more preferably greater than about 90% (w/w) of the separated material. Plants or plant parts of the present invention generated by such methods may be processed into products using known techniques.

Preferred products are meal, feedstock and oil. Methods to produce feed, meal, protein and oil preparations are known in the art. *See, e.g.,* U.S. Patents 4,957,748, 5,100,679, 5,219,596, 5,936,069, 6,005,076, 6,146,669, and 6,156,227. In a preferred embodiment, the protein preparation is a high protein preparation. Such a high protein

preparation preferably has a protein content of greater than about 5% w/v, more preferably about 10% w/v, and even more preferably about 15% w/v.

In a preferred embodiment, the oil preparation is a high oil preparation with an oil content derived from a plant or part thereof of the present invention of greater than about 5% w/v, more preferably greater than about 10% w/v, and even more preferably greater than about 15% w/v. In a preferred embodiment the oil preparation is a liquid and of a volume greater than about 1, 5, 10 or 50 liters. The present invention provides for oil produced from plants of the present invention or generated by a method of the present invention. Such oil may be a minor or major component of any resultant product. Moreover, such oil may be blended with other oils.

In a preferred embodiment, the oil produced from plants of the present invention or generated by a method of the present invention constitutes greater than about 0.5%, 1%, 5%, 10%, 25%, 50%, 75% or 90% by volume or weight of the oil component of any product. In another embodiment, the oil preparation may be blended and can constitute greater than about 10%, 25%, 35%, 50% or 75% of the blend by volume. Oil produced from a plant of the present invention can be admixed with one or more organic solvents or petroleum distillates.

Seed containers

Seeds of the plants may be placed in a container. As used herein, a container is any object capable of holding such seeds. A container preferably contains greater than about 500, 1,000, 5,000, or 25,000 seeds where at least about 10%, 25%, 50%, 75% or 100% of the seeds are derived from a plant of the present invention. The present invention also provides a container of over about 10,000, more preferably about 20,000, and even more preferably about 40,000 seeds where over about 10%, more preferably about 25%, more preferably 50% and even more preferably about 75% or 90% of the seeds are seeds derived from a plant of the present invention. The present invention also provides a container of over about 10 kg, more preferably about 25 kg, and even more preferably about 50 kg seeds where over about 10%, more preferably about 25%, more preferably about 50% and even more preferably about 75% or 90% of the seeds are seeds derived from a plant of the present invention.

30 E. Antibodies

One aspect of the invention concerns antibodies, single-chain antigen binding molecules, or other proteins that specifically bind to one or more of the protein or peptide molecules of the invention and their homologs, fusions or fragments. In a particularly

preferred embodiment, the antibody specifically binds to a protein having the amino acid sequence set forth in SEQ ID NOs: 4, 48, 49 and 50, or an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1 through 3 and 5 through 47. Such antibodies may be used to quantitatively or qualitatively detect the protein or peptide molecules of the invention.

Nucleic acid molecules that encode all or part of the protein of the invention can be expressed, via recombinant means, to yield protein or peptides that can in turn be used to elicit antibodies that are capable of binding the expressed protein or peptide. Such antibodies may be used in immunoassays for that protein. Such protein-encoding molecules, or their fragments may be a "fusion" molecule (*i.e.*, a part of a larger nucleic acid molecule) such that, upon expression, a fusion protein is produced. It is understood that any of the nucleic acid molecules of the invention may be expressed, via recombinant means, to yield proteins or peptides encoded by these nucleic acid molecules.

The antibodies that specifically bind proteins and protein fragments of the invention may be polyclonal or monoclonal and may comprise intact immunoglobulins, or antigen binding portions of immunoglobulins fragments (such as (F(ab'), F(ab')₂), or single-chain immunoglobulins producible, for example, via recombinant means. It is understood that practitioners are familiar with the standard resource materials that describe specific conditions and procedures for the construction, manipulation and isolation of antibodies (*see, e.g.*, Harlow and Lane, in: *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York, 1988).

As discussed below, such antibody molecules or their fragments may be used for diagnostic purposes. Where the antibodies are intended for diagnostic purposes, it may be desirable to derivatize them, for example with a ligand group (such as biotin) or a detectable marker group (such as a fluorescent group, a radioisotope or an enzyme).

The ability to produce antibodies that bind the protein or peptide molecules of the invention permits the identification of mimetic compounds derived from those molecules. These mimetic compounds may contain a fragment of the protein or peptide or merely a structurally similar region and nonetheless exhibits an ability to specifically bind to antibodies directed against that compound.

Antibodies have been expressed in plants. Cytoplasmic expression of a scFv (single-chain Fv antibody) has been reported to delay infection by artichoke mottled crinkle virus. Transgenic plants that express antibodies directed against endogenous proteins may exhibit a physiological effect. For example, expressed anti-abscisic antibodies have been reported to result in a general perturbation of seed development. See, *e.g.*, Hiatt *et al.*, *Nature* 342:76-78

(1989); Conrad and Fielder, *Plant Mol. Biol.* 26:1023-1030 (1994); Philips *et al.*, *EMBO J.* 16:4489-4496 (1997); Marion-Poll, *Trends in Plant Science* 2:447-448 (1997).

Antibodies that are catalytic may also be expressed in plants (abzymes). The principle behind abzymes is that because antibodies may be raised against many molecules, this recognition ability can be directed toward generating antibodies that bind transition states to force a chemical reaction forward. Persidas, *Nature Biotechnology* 15:1313-1315 (1997); Baca *et al.*, *Ann. Rev. Biophys. Biomol. Struct.* 26:461-493 (1997). The catalytic abilities of abzymes may be enhanced by site directed mutagenesis. Examples of abzymes are, for example, set forth in U.S. Patent Nos. 5,658,753; 5,632,990; 5,631,137; 5,602,015; 5,559,538; 5,576,174; 5,500,358; 5,318,897; 5,298,409; 5,258,289; and 5,194,585. It is understood that any of the antibodies of the invention may be expressed in plants and that such expression can result in a physiological effect. It is also understood that any of the expressed antibodies may be catalytic.

F. Markers

Another subset of the nucleic acid molecules of the invention includes nucleic acid molecules that are markers. The markers can be used in a number of ways in the field of molecular genetics. Such markers include nucleic acid molecules SEQ ID NOs: 1 through 3 and 5 through 47 or complements thereof or fragments of either that can act as markers and other nucleic acid molecules of the present invention that can act as markers.

Genetic markers of the invention include "dominant" or "codominant" markers. "Codominant markers" reveal the presence of two or more alleles (two per diploid individual) at a locus. "Dominant markers" reveal the presence of only a single allele per locus. The presence of the dominant marker phenotype (*e.g.*, a band of DNA) is an indication that one allele is in either the homozygous or heterozygous condition. The absence of the dominant marker phenotype (*e.g.*, absence of a DNA band) is merely evidence that "some other" undefined allele is present. In the case of populations where individuals are predominantly homozygous and loci are predominately dimorphic, dominant and codominant markers can be equally valuable. As populations become more heterozygous and multi-allelic, codominant markers often become more informative of the genotype than dominant markers. Marker molecules can be, for example, capable of detecting polymorphisms such as single nucleotide polymorphisms (SNPs).

The genomes of animals and plants naturally undergo spontaneous mutation in the course of their continuing evolution. A "polymorphism" is a variation or difference in the sequence of the gene or its flanking regions that arises in some of the members of a species.

The variant sequence and the "original" sequence co-exist in the species' population. In some instances, such co-existence is in stable or quasi-stable equilibrium.

A polymorphism is thus said to be "allelic," in that, due to the existence of the polymorphism, some members of a species may have the original sequence (i.e., the original "allele") whereas other members may have the variant sequence (i.e., the variant "allele"). In the simplest case, only one variant sequence may exist and the polymorphism is thus said to be di-allelic. In other cases, the species' population may contain multiple alleles and the polymorphism is termed tri-allelic, etc. A single gene may have multiple different unrelated polymorphisms. For example, it may have a di-allelic polymorphism at one site and a multi-allelic polymorphism at another site.

The variation that defines the polymorphism may range from a single nucleotide variation to the insertion or deletion of extended regions within a gene. In some cases, the DNA sequence variations are in regions of the genome that are characterized by short tandem repeats (STRs) that include tandem di- or tri-nucleotide repeated motifs of nucleotides. Polymorphisms characterized by such tandem repeats are referred to as "variable number tandem repeat" (VNTR) polymorphisms. VNTRs have been used in identity analysis (EP 370719; U.S. Patent Nos. 5,075,217 and 5,175,082; WO 91/14003).

The detection of polymorphic sites in a sample of DNA may be facilitated through the use of nucleic acid amplification methods. Such methods specifically increase the concentration of polynucleotides that span the polymorphic site, or include that site and sequences located either distal or proximal to it. Such amplified molecules can be readily detected by gel electrophoresis or other means.

In an alternative embodiment, such polymorphisms can be detected through the use of a marker nucleic acid molecule that is physically linked to such polymorphism(s). For this purpose, marker nucleic acid molecules comprising a nucleotide sequence of a polynucleotide located within 1 mb of the polymorphism(s) and more preferably within 100kb of the polymorphism(s) and most preferably within 10kb of the polymorphism(s) can be employed. Alternatively, marker nucleic acid molecules comprising a nucleotide sequence of a polynucleotide located within 25 cM of the polymorphism(s) and more preferably within 15 cM of the polymorphism(s) and most preferably within 5 cM of the polymorphism(s) can be employed.

The identification of a polymorphism can be determined in a variety of ways. By correlating the presence or absence of it in a plant with the presence or absence of a phenotype, it is possible to predict the phenotype of that plant. If a polymorphism creates or destroys a restriction endonuclease cleavage site, or if it results in the loss or insertion of

DNA (*e.g.*, a VNTR polymorphism), it will alter the size or profile of the DNA fragments that are generated by digestion with that restriction endonuclease. As such, organisms that possess a variant sequence can be distinguished from those having the original sequence by restriction fragment analysis. Polymorphisms that can be identified in this manner are
5 termed "restriction fragment length polymorphisms" (RFLPs) (UK Patent Application 2135774; WO 90/13668; WO 90/11369).

Polymorphisms can also be identified by Single Strand Conformation Polymorphism (SSCP) analysis, random amplified polymorphic DNA (RAPD), and cleaveable amplified polymorphic sequences (CAPS). *See, e.g.*, Lee *et al.*, *Anal. Biochem.* 205:289-293 (1992);
10 Sarkar *et al.*, *Genomics* 13:441-443 (1992); Williams *et al.*, *Nucl. Acids Res.* 18:6531-6535 (1990); and Lyamichev *et al.*, *Science* 260:778-783 (1993). It is understood that one or more of the nucleic acids of the invention, may be utilized as markers or probes to detect polymorphisms by SSCP, RAPD or CAPS analysis.

Polymorphisms may also be found using a DNA fingerprinting technique called
15 amplified fragment length polymorphism (AFLP), which is based on the selective PCR amplification of restriction fragments from a total digest of genomic DNA to profile that DNA. Vos *et al.*, *Nucleic Acids Res.* 23:4407-4414 (1995). This method allows for the specific co-amplification of high numbers of restriction fragments, which can be visualized by PCR without knowledge of the nucleic acid sequence. It is understood that one or more of
20 the nucleic acids of the invention may be utilized as markers or probes to detect polymorphisms by AFLP analysis or for fingerprinting RNA.

Single Nucleotide Polymorphisms (SNPs) generally occur at greater frequency than other polymorphic markers and are spaced with a greater uniformity throughout a genome than other reported forms of polymorphism. The greater frequency and uniformity of SNPs
25 means that there is greater probability that such a polymorphism will be found near or in a genetic locus of interest than would be the case for other polymorphisms. SNPs are located in protein-coding regions and noncoding regions of a genome. Some of these SNPs may result in defective or variant protein expression (*e.g.*, as a result of mutations or defective splicing). Analysis (genotyping) of characterized SNPs can require only a plus/minus assay
30 rather than a lengthy measurement, permitting easier automation.

SNPs can be characterized using any of a variety of methods. Such methods include the direct or indirect sequencing of the site, the use of restriction enzymes, enzymatic and chemical mismatch assays, allele-specific PCR, ligase chain reaction, single-strand conformation polymorphism analysis, single base primer extension (U.S. Patent Nos.
35 6,004,744 and 5,888,819), solid-phase ELISA-based oligonucleotide ligation assays, dideoxy

fingerprinting, oligonucleotide fluorescence-quenching assays, 5'-nuclease allele-specific hybridization TaqMan™ assay, template-directed dye-terminator incorporation (TDI) assay (Chen and Kwok, *Nucl. Acids Res.* 25:347-353, 1997), allele-specific molecular beacon assay (Tyagi *et al.*, *Nature Biotech.* 16: 49-53, 1998), PinPoint assay (Haff and Smirnov, *Genome Res.* 7: 378-388, 1997), dCAPS analysis (Neff *et al.*, *Plant J.* 14:387-392, 1998), pyrosequencing (Ronaghi *et al.*, *Analytical Biochemistry* 267:65-71, 1999; WO 98/13523; WO 98/28440), using mass spectrometry, *e.g.* the Masscode™ system (WO 99/05319; WO 98/26095; WO 98/12355; WO 97/33000; WO 97/27331; and U.S. Patent No. 5,965,363), invasive cleavage of oligonucleotide probes, and using high density oligonucleotide arrays (Hacia *et al.*, *Nature Genetics* 22:164-167).

Polymorphisms may also be detected using allele-specific oligonucleotides (ASO), which, can be for example, used in combination with hybridization based technology including Southern, northern, and dot blot hybridizations, reverse dot blot hybridizations and hybridizations performed on microarray and related technology.

The stringency of hybridization for polymorphism detection is highly dependent upon a variety of factors, including length of the allele-specific oligonucleotide, sequence composition, degree of complementarity (*i.e.* presence or absence of base mismatches), concentration of salts and other factors such as formamide, and temperature. These factors are important both during the hybridization itself and during subsequent washes performed to remove target polynucleotide that is not specifically hybridized. In practice, the conditions of the final, most stringent wash are most critical. In addition, the amount of target polynucleotide that is able to hybridize to the allele-specific oligonucleotide is also governed by such factors as the concentration of both the ASO and the target polynucleotide, the presence and concentration of factors that act to "tie up" water molecules, so as to effectively concentrate the reagents (*e.g.*, PEG, dextran, dextran sulfate, *etc.*), whether the nucleic acids are immobilized or in solution, and the duration of hybridization and washing steps.

Hybridizations are preferably performed below the melting temperature (T_m) of the ASO. The closer the hybridization and/or washing step is to the T_m , the higher the stringency. T_m for an oligonucleotide may be approximated, for example, according to the following formula: $T_m = 81.5 + 16.6 \times (\log_{10}[\text{Na}^+]) + 0.41 \times (\%G+C) - 675/n$; where $[\text{Na}^+]$ is the molar salt concentration of Na^+ or any other suitable cation and n = number of bases in the oligonucleotide. Other formulas for approximating T_m are available and are known to those of ordinary skill in the art.

Stringency is preferably adjusted so as to allow a given ASO to differentially hybridize to a target polynucleotide of the correct allele and a target polynucleotide of the incorrect allele. Preferably, there will be at least a two-fold differential between the signal produced by the ASO hybridizing to a target polynucleotide of the correct allele and the level of the signal produced by the ASO cross-hybridizing to a target polynucleotide of the incorrect allele (*e.g.*, an ASO specific for a mutant allele cross-hybridizing to a wild-type allele). In more preferred embodiments of the present invention, there is at least a five-fold signal differential. In highly preferred embodiments of the present invention, there is at least an order of magnitude signal differential between the ASO hybridizing to a target polynucleotide of the correct allele and the level of the signal produced by the ASO cross-hybridizing to a target polynucleotide of the incorrect allele. While certain methods for detecting polymorphisms are described herein, other detection methodologies may be utilized.

The present invention includes and provides a method for detecting a polymorphism in a plant whose presence is predictive of a mutation affecting a level or pattern of a protein comprising: (A) incubating under conditions permitting nucleic acid hybridization: (i) a marker nucleic acid molecule having a nucleic acid sequence that hybridizes to a sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof; and (ii) a complementary nucleic acid molecule obtained from a sample, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule permits the detection of a polymorphism; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention includes and provides a method of determining a degree of association between a polymorphism and a plant trait comprising: (A) hybridizing a nucleic acid molecule specific for the polymorphism to genetic material of a plant, wherein the nucleic acid molecule has a sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof, and fragments of these sequences; and (B) calculating the degree of association between the polymorphism and the plant trait.

The present invention includes and provides a method of isolating a nucleic acid that encodes a protein or fragment thereof comprising: (A) incubating under conditions permitting nucleic acid hybridization: (i) a first nucleic acid molecule comprising a sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, complements thereof, and fragments of these sequences; and (ii) a complementary second nucleic acid

molecule obtained from a plant cell or plant tissue; (B) permitting hybridization between the first nucleic acid molecule and the second nucleic acid molecule obtained from the plant cell or plant tissue; and (C) isolating the second nucleic acid molecule.

G. Plant Breeding

5 Plants of the present invention can be part of or generated from a breeding program. The choice of breeding method depends on the mode of plant reproduction, the heritability of the trait(s) being improved, and the type of cultivar used commercially (*e.g.*, F₁ hybrid cultivar, pureline cultivar, etc). Selected, non-limiting approaches, for breeding the plants of the present invention are set forth below. A breeding program can be enhanced using marker
10 assisted selection of the progeny of any cross. It is further understood that any commercial and non-commercial cultivars can be utilized in a breeding program. Factors such as, for example, emergence vigor, vegetative vigor, stress tolerance, disease resistance, branching, flowering, seed set, seed size, seed density, standability, and threshability etc. will generally dictate the choice.

15 For highly heritable traits, a choice of superior individual plants evaluated at a single location will be effective, whereas for traits with low heritability, selection should be based on mean values obtained from replicated evaluations of families of related plants. Popular selection methods commonly include pedigree selection, modified pedigree selection, mass selection, and recurrent selection. In a preferred embodiment a backcross or recurrent
20 breeding program is undertaken.

 The complexity of inheritance influences choice of the breeding method. Backcross breeding can be used to transfer one or a few favorable genes for a highly heritable trait into a desirable cultivar. This approach has been used extensively for breeding disease-resistant cultivars. Various recurrent selection techniques are used to improve quantitatively inherited
25 traits controlled by numerous genes. The use of recurrent selection in self-pollinating crops depends on the ease of pollination, the frequency of successful hybrids from each pollination, and the number of hybrid offspring from each successful cross.

 Breeding lines can be tested and compared to appropriate standards in environments representative of the commercial target area(s) for two or more generations. The best lines
30 are candidates for new commercial cultivars; those still deficient in traits may be used as parents to produce new populations for further selection.

 One method of identifying a superior plant is to observe its performance relative to other experimental plants and to a widely grown standard cultivar. If a single observation is inconclusive, replicated observations can provide a better estimate of its genetic worth. A

breeder can select and cross two or more parental lines, followed by repeated selfing and selection, producing many new genetic combinations.

The development of new cultivars requires the development and selection of varieties, the crossing of these varieties and the selection of superior hybrid crosses. The hybrid seed can be produced by manual crosses between selected male-fertile parents or by using male sterility systems. Hybrids are selected for certain single gene traits such as pod color, flower color, seed yield, pubescence color, or herbicide resistance, which indicate that the seed is truly a hybrid. Additional data on parental lines, as well as the phenotype of the hybrid, influence the breeder's decision whether to continue with the specific hybrid cross.

Pedigree breeding and recurrent selection breeding methods can be used to develop cultivars from breeding populations. Breeding programs combine desirable traits from two or more cultivars or various broad-based sources into breeding pools from which cultivars are developed by selfing and selection of desired phenotypes. New cultivars can be evaluated to determine which have commercial potential.

Pedigree breeding is used commonly for the improvement of self-pollinating crops. Two parents who possess favorable, complementary traits are crossed to produce an F_1 . An F_2 population is produced by selfing one or several F_1 's. Selection of the best individuals from the best families is carried out. Replicated testing of families can begin in the F_4 generation to improve the effectiveness of selection for traits with low heritability. At an advanced stage of inbreeding (*i.e.*, F_6 and F_7), the best lines or mixtures of phenotypically similar lines are tested for potential release as new cultivars.

Backcross breeding has been used to transfer genes for a simply inherited, highly heritable trait into a desirable homozygous cultivar or inbred line, which is the recurrent parent. The source of the trait to be transferred is called the donor parent. The resulting plant is expected to have the attributes of the recurrent parent (*e.g.*, cultivar) and the desirable trait transferred from the donor parent. After the initial cross, individuals possessing the phenotype of the donor parent are selected and repeatedly crossed (backcrossed) to the recurrent parent. The resulting parent is expected to have the attributes of the recurrent parent (*e.g.*, cultivar) and the desirable trait transferred from the donor parent.

The single-seed descent procedure in the strict sense refers to planting a segregating population, harvesting a sample of one seed per plant, and using the one-seed sample to plant the next generation. When the population has been advanced from the F_2 to the desired level of inbreeding, the plants from which lines are derived will each trace to different F_2 individuals. The number of plants in a population declines each generation due to failure of some seeds to germinate or some plants to produce at least one seed. As a result, not all of

the F₂ plants originally sampled in the population will be represented by a progeny when generation advance is completed.

5 In a multiple-seed procedure, breeders commonly harvest one or more pods from each plant in a population and thresh them together to form a bulk. Part of the bulk is used to plant the next generation and part is put in reserve. The procedure has been referred to as modified single-seed descent or the pod-bulk technique. The multiple-seed procedure has been used to save labor at harvest. It is considerably faster to thresh pods with a machine than to remove one seed from each by hand for the single-seed procedure. The multiple-seed procedure also makes it possible to plant the same number of seed of a population each
10 generation of inbreeding.

Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several reference books (*e.g.*, Fehr, *Principles of Cultivar Development*, Vol. 1 (1987)).

A transgenic plant of the present invention may also be reproduced using apomixis.
15 Apomixis is a genetically controlled method of reproduction in plants where the embryo is formed without union of an egg and a sperm. There are three basic types of apomictic reproduction: 1) apospory where the embryo develops from a chromosomally unreduced egg in an embryo sac derived from the nucleus, 2) diplospory where the embryo develops from an unreduced egg in an embryo sac derived from the megaspore mother cell, and
20 3) adventitious embryony where the embryo develops directly from a somatic cell. In most forms of apomixis, pseudogamy or fertilization of the polar nuclei to produce endosperm is necessary for seed viability. In apospory, a nurse cultivar can be used as a pollen source for endosperm formation in seeds. The nurse cultivar does not affect the genetics of the aposporous apomictic cultivar because the unreduced egg of the cultivar develops
25 parthenogenetically, but makes possible endosperm production. Apomixis is economically important, especially in transgenic plants, because it causes any genotype, no matter how heterozygous, to breed true. Thus, with apomictic reproduction, heterozygous transgenic plants can maintain their genetic fidelity throughout repeated life cycles. Methods for the production of apomictic plants are known in the art. *See, e.g.*, U.S. Patent No. 5,811,636.

30 Requirements for marker-assisted selection in a plant breeding program are: (1) the marker(s) should co-segregate or be closely linked with the desired trait; (2) an efficient means of screening large populations for the molecular marker(s) should be available; and (3) the screening technique should have high reproducibility across laboratories and preferably be economical to use and be user-friendly.

The genetic linkage of marker molecules can be established by a gene mapping model such as, without limitation, the flanking marker model reported by Lander and Botstein, *Genetics* 121:185-199 (1989), and the interval mapping model, based on maximum likelihood methods described by Lander and Botstein, and implemented in the software package MAPMAKER/QTL (Lincoln and Lander, *Mapping Genes Controlling Quantitative Traits Using MAPMAKER/QTL*, Whitehead Institute for Biomedical Research, Massachusetts, 1990). Additional software includes Qgene, Version 2.23 (1996), Department of Plant Breeding and Biometry, 266 Emerson Hall, Cornell University, Ithaca, NY). Use of Qgene software is a particularly preferred approach.

A maximum likelihood estimate (MLE) for the presence of a marker is calculated, together with an MLE assuming no QTL effect, to avoid false positives. A $\log_{10}$ of an odds ratio (LOD) is then calculated as: $\text{LOD} = \log_{10}(\text{MLE for the presence of a QTL} / \text{MLE given no linked QTL})$.

The LOD score essentially indicates how much more likely the data are to have arisen assuming the presence of a QTL than in its absence. The LOD threshold value for avoiding a false positive with a given confidence, say 95%, depends on the number of markers and the length of the genome. Graphs indicating LOD thresholds are set forth in Lander and Botstein, *supra*, and further described by Arús and Moreno-González, Plant Breeding, (Hayward *et al.*, eds.) Chapman & Hall, London, pp. 314-331 (1993).

In a preferred embodiment of the present invention the nucleic acid marker exhibits a LOD score of greater than about 2.0, more preferably about 2.5, even more preferably greater than about 3.0 or 4.0 with the trait or phenotype of interest. In a preferred embodiment, the trait of interest is altered tocopherol levels or compositions.

Additional models can be used. Many modifications and alternative approaches to interval mapping have been reported, including the use non-parametric methods. Kruglyak and Lander, *Genetics* 139:1421-1428 (1995). Multiple regression methods or models can be also be used, in which the trait is regressed on a large number of markers. Weber and Wricke, *Advances in Plant Breeding*, Blackwell, Berlin (1994). Procedures may combine interval mapping with regression analysis, whereby the phenotype is regressed onto a single putative QTL at a given marker interval and at the same time onto a number of markers that serve as 'cofactors.' Generally, the use of cofactors reduces the bias and sampling error of the estimated QTL positions, thereby improving the precision and efficiency of QTL mapping. Zeng, *Genetics* 136:1457-1468 (1994). These models can be extended to multi-environment experiments to analyze genotype-environment interactions. Jansen *et al.*, *Theo. Appl. Genet.* 91:33-37 (1995).

It is understood that one or more of the nucleic acid molecules of the invention may be used as molecular markers. It is also understood that one or more of the protein molecules of the invention may be used as molecular markers.

In a preferred embodiment, the polymorphism is present and screened for in a mapping population, *e.g.* a collection of plants capable of being used with markers such as polymorphic markers to map genetic position of traits. The choice of appropriate mapping population often depends on the type of marker systems employed. Consideration must be given to the source of parents (adapted vs. exotic) used in the mapping population. Chromosome pairing and recombination rates can be severely disturbed (suppressed) in wide crosses (adapted x exotic) and generally yield greatly reduced linkage distances. Wide crosses will usually provide segregating populations with a relatively large number of polymorphisms when compared to progeny in a narrow cross (adapted x adapted).

An F_2 population is the first generation of selfing (self-pollinating) after the hybrid seed is produced. Usually a single F_1 plant is selfed to generate a population segregating for all the genes in Mendelian (1:2:1) pattern. Maximum genetic information is obtained from a completely classified F_2 population using a codominant marker system (Mather, 1938). In the case of dominant markers, progeny tests (*e.g.*, F_3 , BCF_2) are required to identify the heterozygotes, in order to classify the population. However, this procedure is often prohibitive because of the cost and time involved in progeny testing. Progeny testing of F_2 individuals is often used in map construction where phenotypes do not consistently reflect genotype (*e.g.* disease resistance) or where trait expression is controlled by a QTL. Segregation data from progeny test populations *e.g.* F_3 or BCF_2) can be used in map construction. Marker-assisted selection can then be applied to cross progeny based on marker-trait map associations (F_2 , F_3), where linkage groups have not been completely disassociated by recombination events (*i.e.*, maximum disequilibrium).

Recombinant inbred lines (RIL) (genetically related lines; usually $>F_5$, developed from continuously selfing F_2 lines towards homozygosity) can be used as a mapping population. Information obtained from dominant markers can be maximized by using RIL because all loci are homozygous or nearly so. Under conditions of tight linkage (*i.e.*, about $<10\%$ recombination), dominant and co-dominant markers evaluated in RIL populations provide more information per individual than either marker type in backcross populations. However, as the distance between markers becomes larger (*i.e.*, loci become more independent), the information in RIL populations decreases dramatically when compared to codominant markers.

Backcross populations (*e.g.*, generated from a cross between a successful variety (recurrent parent) and another variety (donor parent) carrying a trait not present in the former) can be utilized as a mapping population. A series of backcrosses to the recurrent parent can be made to recover most of its desirable traits. Thus a population is created
5 consisting of individuals nearly like the recurrent parent but each individual carries varying amounts or mosaic of genomic regions from the donor parent. Backcross populations can be useful for mapping dominant markers if all loci in the recurrent parent are homozygous and the donor and recurrent parent have contrasting polymorphic marker alleles.

Information obtained from backcross populations using either codominant or
10 dominant markers is less than that obtained from F₂ populations because one, rather than two, recombinant gamete is sampled per plant. Backcross populations, however, are more informative (at low marker saturation) when compared to RILs as the distance between linked loci increases in RIL populations (*i.e.* about .15% recombination). Increased recombination can be beneficial for resolution of tight linkages, but may be undesirable in
15 the construction of maps with low marker saturation.

Near-isogenic lines (NIL) (created by many backcrosses to produce a collection of individuals that is nearly identical in genetic composition except for the trait or genomic region under interrogation) can be used as a mapping population. In mapping with NILs, only a portion of the polymorphic loci is expected to map to a selected region.

20 Bulk segregant analysis (BSA) is a method developed for the rapid identification of linkage between markers and traits of interest (Michelmore *et al.*, *PNAS* 88:9828-9832 (1991). In BSA, two bulked DNA samples are drawn from a segregating population originating from a single cross. These bulks contain individuals that are identical for a particular trait (resistant or susceptible to particular disease) or genomic region but arbitrary
25 at unlinked regions (*i.e.* heterozygous). Regions unlinked to the target region will not differ between the bulked samples of many individuals in BSA.

H. Determining the Level of Expression Response

In an aspect of the present invention, one or more of the nucleic molecules of the present invention are used to determine the level (*i.e.*, the concentration of mRNA in a
30 sample, *etc.*) or pattern (*i.e.*, the kinetics of expression, rate of decomposition, stability profile, *etc.*) of the expression of a protein encoded in part or whole by one or more of the nucleic acid molecule of the present invention (collectively, the "Expression Response" of a cell or tissue).

As used herein, the Expression Response manifested by a cell or tissue is said to be “altered” if it differs from the Expression Response of cells or tissues of plants not exhibiting the phenotype. To determine whether a Expression Response is altered, the Expression Response manifested by the cell or tissue of the plant exhibiting the phenotype is compared
5 with that of a similar cell or tissue sample of a plant not exhibiting the phenotype. As will be appreciated, it is not necessary to re-determine the Expression Response of the cell or tissue sample of plants not exhibiting the phenotype each time such a comparison is made; rather, the Expression Response of a particular plant may be compared with previously obtained values of normal plants.

10 A change in genotype or phenotype may be transient or permanent. Also as used herein, a tissue sample is any sample that comprises more than one cell. In a preferred aspect, a tissue sample comprises cells that share a common characteristic (e.g. derived from root, seed, flower, leaf, stem or pollen etc.).

In one aspect of the present invention, an evaluation can be conducted to determine
15 whether a particular mRNA molecule is present. One or more of the nucleic acid molecules of the present invention are utilized to detect the presence or quantity of the mRNA species. Such molecules are then incubated with cell or tissue extracts of a plant under conditions sufficient to permit nucleic acid hybridization. The detection of double-stranded probe-mRNA hybrid molecules is indicative of the presence of the mRNA; the amount of such
20 hybrid formed is proportional to the amount of mRNA. Thus, such probes may be used to ascertain the level and extent of the mRNA production in a plant’s cells or tissues. Such nucleic acid hybridization may be conducted under quantitative conditions (thereby providing a numerical value of the amount of the mRNA present). Alternatively, the assay may be conducted as a qualitative assay that indicates either that the mRNA is present, or
25 that its level exceeds a user set, predefined value.

A number of methods can be used to compare the expression response between two or more samples of cells or tissue. These methods include hybridization assays, such as northern, RNase protection assays, and *in situ* hybridization. Alternatively, the methods include PCR-type assays. In a preferred method, the expression response is compared by
30 hybridizing nucleic acids from the two or more samples to an array of nucleic acids. The array contains a plurality of suspected sequences known or suspected of being present in the cells or tissue of the samples.

An advantage of *in situ* hybridization over more other techniques for the detection of nucleic acids is that it allows an investigator to determine the precise spatial population. *In*
35 *situ* hybridization may be used to measure the steady-state level of RNA accumulation. A

number of protocols have been devised for *in situ* hybridization, each with tissue preparation, hybridization and washing conditions.

In situ hybridization also allows for the localization of proteins within a tissue or cell. It is understood that one or more of the molecules of the invention, preferably one or
5 more of the nucleic acid molecules or fragments thereof of the invention or one or more of the antibodies of the invention may be utilized to detect the level or pattern of a protein or mRNA thereof by *in situ* hybridization.

Fluorescent *in situ* hybridization allows the localization of a particular DNA sequence along a chromosome, which is useful, among other uses, for gene mapping,
10 following chromosomes in hybrid lines, or detecting chromosomes with translocations, transversions or deletions. *In situ* hybridization has been used to identify chromosomes in several plant species. It is understood that the nucleic acid molecules of the invention may be used as probes or markers to localize sequences along a chromosome.

Another method to localize the expression of a molecule is tissue printing. Tissue
15 printing provides a way to screen, at the same time on the same membrane many tissue sections from different plants or different developmental stages. *See, e.g., Barres et al., Neuron* 5:527-544 (1990); Cassab and Varner, *J. Cell. Biol.* 105:2581-2588 (1987); Harris and Chrispeels, *Plant Physiol.* 56:292-299 (1975); Reid and Pont-Lezica, *Tissue Printing: Tools for the Study of Anatomy, Histochemistry and Gene Expression*, Academic Press, New
20 York, New York (1992); Reid *et al.*, *Plant Physiol.* 93:160-165 (1990); Spruce *et al.*, *Phytochemistry* 26:2901-2903 (1987); Ye *et al.*, *Plant J.* 1:175-183 (1991); Yomo and Taylor, *Planta* 112:35-43 (1973).

A microarray-based method for high-throughput monitoring of gene expression may also be utilized to measure Expression Response. This 'chip'-based approach involves
25 microarrays of nucleic acid molecules as gene-specific hybridization targets to quantitatively measure expression of the corresponding mRNA. Hybridization to a microarray can be used to efficiently analyze the presence and/or amount of a number of nucleotide sequences simultaneously.

Several microarray methods have been described. One method compares the
30 sequences to be analyzed by hybridization to a set of oligonucleotides representing all possible subsequences. A second method hybridizes the sample to an array of oligonucleotide or cDNA molecules. An array consisting of oligonucleotides complementary to subsequences of a target sequence can be used to determine the identity of a target sequence, measure its amount, and detect single nucleotide differences between the target
35 and a reference sequence. Nucleic acid molecule microarrays may also be screened with

protein molecules or fragments thereof to determine nucleic acid molecules that specifically bind protein molecules or fragments thereof.

The microarray approach may be used with polypeptide targets (U.S. Patent Nos. 5,445,934; 5,143,854; 5,079,600; and 4,923,901). Essentially, polypeptides are synthesized
5 on a substrate (microarray) and these polypeptides can be screened with either protein molecules or fragments thereof or nucleic acid molecules in order to screen for either protein molecules or fragments thereof or nucleic acid molecules that specifically bind the target polypeptides.

In a preferred embodiment of the present invention microarrays may be prepared that
10 comprise nucleic acid molecules where preferably at least about 10%, preferably at least about 25%, more preferably at least about 50% and even more preferably at least about 75%, 80%, 85%, 90% or 95% of the nucleic acid molecules located on that array are selected from the group of nucleic acid molecules that hybridize under low, moderate or high stringency conditions to one or more nucleic acid molecules having a nucleic acid sequence selected
15 from the group of SEQ ID NO: 1 through 3, 5 through 47, and complements thereof.

In another preferred embodiment of the present invention microarrays may be prepared that comprise nucleic acid molecules where preferably at least about 10%, preferably at least about 25%, more preferably at least about 50% and even more preferably at least about 75%, 80%, 85%, 90% or 95% of the nucleic acid molecules located on that
20 array are selected from the group of nucleic acid molecules having a nucleic acid sequence selected from the group of SEQ ID NO: 1 through 3, 5 through 47, complements thereof, and fragments of these sequences.

In a preferred embodiment of the present invention microarrays may be prepared that comprise nucleic acid molecules where such nucleic acid molecules encode at least one,
25 preferably at least two, more preferably at least three, even more preferably at least four, five or six proteins or fragments thereof selected from the group consisting of *gcpE*, *ygbB*, *ygbP*, *yehB*, *dxs* and *dxr*.

The present invention includes and provides a method for determining a level or pattern of a protein in a plant cell or plant tissue comprising (A) incubating under conditions
30 permitting nucleic acid hybridization: (i) a marker nucleic acid molecule having a nucleic acid sequence that hybridizes to a sequence selected from the group consisting of SEQ ID NOs: 1 through 3, 5 through 47, and complements thereof; and (ii) a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule
35 permits the detection of an mRNA for the protein; (B) permitting hybridization between the

marker nucleic acid molecule; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the protein in the plant.

The present invention also includes and provides a method for determining a level or pattern of a protein in a plant cell or plant tissue comprising (A) assaying the concentration of the protein in a first sample obtained from the plant cell or plant tissue; (B) assaying the concentration of the protein in a second sample obtained from a reference plant cell or a reference plant tissue with a known level or pattern of the protein; and (C) comparing the assayed concentration of the protein in the first sample to the assayed concentration of the protein in the second sample.

I. Screening Uses

The present invention provides methods and agents that can be used to screen for and isolate genes associated with the MEP pathway. Because the MEP pathway is an essential pathway, disruption of any essential gene in the MEP pathway will result in the death of the cell or organism. While not being limited to any particular biological process, the present invention provides a method and the agents associated with such a method where mutations that result in loss of function of a MEP pathway gene do not result in cell or organism death by providing a second pathway capable of synthesizing IPP and DMAPP. The present invention provides cells and organisms having a second pathway capable of synthesizing IPP and DMAPP.

In a preferred aspect, a cell or organism comprising: (a) a first DNA sequence encoding an enzyme having catalytic activity of mevalonate kinase; (b) a second DNA sequence encoding an enzyme having catalytic activity of 5-phosphomevalonate kinase; (c) a third DNA sequence encoding an enzyme having catalytic activity of 5-diphosphomevalonate-decarboxylase; and (d) a fourth DNA sequence encoding an enzyme having catalytic activity of isopentenyl diphosphate isomerase; wherein at least two of said first, second, third, or fourth DNA sequences have a foreign DNA sequence.

In a preferred aspect, the second pathway capable of synthesizing IPP and DMAPP has at least one, more preferably at least two, even more preferably at least three or four enzymes selected from the group consisting of: mevalonate kinase, 5-phosphomevalonate kinase, 5-diphosphomevalonate decarboxylase and isopentenyl diphosphate isomerase. In a more preferred embodiment, at least two, even more preferably at least three or four of the enzymes selected from the group consisting of: mevalonate kinase, 5-phosphomevalonate kinase, 5-diphosphomevalonate decarboxylase and isopentenyl diphosphate isomerase are

encoded by a foreign DNA sequence. Any foreign DNA encoding such enzymes may be utilized such as human 5-phosphomevalonate kinase (Genbank Accession No. HO9914).

Any cell or organism that possesses the MEP pathway may be used in this aspect of the invention. By providing a second pathway capable of synthesizing IPP and DMAPP, such cells can be utilized in methods to examine the function of a gene, determine whether a gene is associated with the MEP pathway, and identify a gene associated with the MEP pathway.

The present invention includes and provides a cell comprising: (a) a first DNA sequence encoding an enzyme having catalytic activity of mevalonate kinase; (b) a second DNA sequence encoding an enzyme having catalytic activity of 5-phosphomevalonate kinase; (c) a third DNA sequence encoding an enzyme having catalytic activity of 5-diphosphomevalonate-decarboxylase and (d) a fourth DNA sequence encoding an enzyme having catalytic activity of isopentenyl diphosphate isomerase; wherein at least two of the first, second, third or fourth DNA sequence have a foreign DNA sequence.

The present invention includes and provides a method for examining the function of a gene associated with the MEP pathway, comprising: (a) rendering inoperative the gene in a first cell capable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; (b) rendering inoperative the gene in a second cell incapable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; and (c) determining the viability of the first cell and the second cell.

The present invention includes and provides a method for determining whether a gene is associated with the MEP pathway, comprising: (a) rendering inoperative the gene in a first cell capable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; (b) rendering inoperative the gene in a second cell incapable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; and (c) determining the viability of the first cell and the second cell.

The present invention includes and provides a method for identifying a gene associated with the MEP pathway, comprising: (a) rendering inoperative the gene in a first cell capable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; (b) rendering inoperative the gene in a second cell incapable of converting mevalonic acid to isopentenyl diphosphate and dimethylallyl diphosphate; and (c) determining the viability of the first cell and the second cell.

Application of the teachings of the present invention to a specific problem or environment is within the capabilities of one having ordinary skill in the art in light of the teachings contained herein. Examples of the products and processes of the present invention

appear in the following examples, which are provided by way of illustration, and are not intended to be limiting of the present invention.

EXAMPLE 1

ISOLATION AND MUTAGENESIS OF THE CODING SEQUENCES OF THE MVA⁺ TRANSCRIPTION UNIT

Yeast Diphosphomevalonate Decarboxylase (yPMD, ORF YNR043w, *ERG19*)

The coding sequence of yPMD is amplified by PCR using genomic DNA using *Saccharomyces cerevisiae* strain FY1679 as template. The reaction mixture of the PCR is prepared in a final volume of 25 µl containing 1 µg of template, 0.5 µM of primers CINCO (SEQ ID NO: 51) and SEIS (SEQ ID NO: 52), 100 µM of each deoxynucleoside triphosphate (dNTPs) and *Pfu* reaction buffer (20 mM of Tris-HCl adjusted to pH 8.8, 2 mM of MgSO₄, 10 mM of KCl, 10 mM of (NH₄)₂SO₄, 0.1 % of TritonTM X-100, 100 µg/ml of BSA). The sample is covered with mineral oil, incubated at 96° C for 3 minutes and cooled to 80° C. *Pfu* DNA polymerase (1 unit, Stratagene) is added and the reaction mixture is incubated for 30 cycles consisting of 1 minute at 94° C and 4 minutes 30 sec at 72° C, followed by a final step of 10 minutes at 72° C. The PCR product (1879 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+.

Nde I and *Eco* RI restriction sites are introduced, respectively, at the 5' and 3' end of the yPMD coding sequence by PCR, using plasmid DNA as template. The reaction mixture of the PCR is prepared in a final volume of 50 µl containing 200 ng of template, 1 µM of primers MPD-Nde5' (SEQ ID NO: 53) and MPD-Eco3' (SEQ ID NO: 54), 100 µM of dNTPs, *Pfu* reaction buffer and 1.25 units of *Pfu* DNA polymerase. The sample is denatured for 2 minutes at 94° C and incubated for 10 cycles consisting of 1 minute at 94° C, 1 minute at 61° C and 2 minutes 30 sec at 72° C. The PCR product (1207 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+. Sequencing is performed to ensure that no additional mutation had been introduced during amplification.

Human 5-Phosphomevalonate Kinase (hPMK)

A *Hpa* I restriction site is introduced at both ends of the coding sequence of the human 5-phosphomevalonate kinase by PCR, using the cDNA clone ym0505.rl from Soares infant brain 1NIB as template. The clone ym0505.rl (I.M.A.G.E. 46897; GenBank accession number H09914) is obtained from Research Genetics, Inc (Huntsville, Alabama). The reaction mixture of the PCR is prepared in a final volume of 50 µl containing 200 ng of

template, 1 μ M of primers hPMK1 (SEQ ID NO: 55) and hPMK4 (SEQ ID NO: 56), 100 μ M of dNTPs, *Pfu* reaction buffer and 1.25 units of *Pfu* DNA polymerase. The sample is denatured for 2 minutes at 94° C and incubated for 10 cycles consisting of 30 sec at 94° C, 40 sec at 65° C and 1 minute 45 sec at 72° C. The PCR product (601 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+ and sequenced.

Yeast Mevalonate Kinase (yMVK, ORF YMR208w, *ERG12*)

The coding sequence of yMVK is amplified by PCR using genomic DNA from *Saccharomyces cerevisiae* strain FY1679 as template. The reaction mixture of the PCR is prepared in a final volume of 25 μ l containing 1 μ g of template, 0.5 μ M of primers UNO (SEQ ID NO: 57) and DOS (SEQ ID NO: 58), 100 μ M of dNTPs and *Pfu* reaction buffer. The sample is covered with mineral oil, incubated at 96° C for 3 minutes and cooled to 80° C. One unit of *Pfu* DNA polymerase is added and the reaction mixture is incubated for 30 cycles consisting of 1 minute at 94° C and 4 minutes 30 sec at 72° C, followed by a final step of 10 minutes at 72° C. The PCR product (1744 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+.

A *Hpa* I restriction site is introduced at both ends of the yPMK coding sequence by PCR, using plasmid DNA as template. The reaction mixture of the PCR is prepared in a final volume of 50 μ l containing 200 ng of template, 1 μ M of primers MK-Hpa5' (SEQ ID NO: 59) and MK-Hpa3' (SEQ ID NO: 60), 100 μ M of dNTPs, *Pfu* reaction buffer and 1.25 units of *Pfu* DNA polymerase. The sample is denatured for 2 minutes at 94° C and incubated for 10 cycles consisting of 45 sec at 94° C, 45 sec at 57° C and 2 minutes 50 sec at 72° C. The PCR product (1351 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+ and sequenced.

Isopentenyl Diphosphate Isomerase from *Escherichia coli* (ecIDI)

The coding sequence of the isopentenyl diphosphate isomerase from *E. coli* is amplified by PCR, using genomic DNA from strain W3110 as template. In this PCR, a *Xho* I restriction site is introduced at both ends of the coding sequence. The reaction mixture of the PCR is prepared in a final volume of 50 μ l containing 200 ng of template, 0.5 μ M of primers idi5X (SEQ ID NO: 61) and idi3X (SEQ ID NO: 62), 100 μ M of dNTPs and *Pfu* reaction buffer. The sample is covered with mineral oil, incubated at 96° C for 3 minutes and cooled to 80° C. *Pfu* DNA polymerase (1.5 units) is added and the reaction mixture is incubated for 5 cycles consisting of 30 sec at 94° C, 40 sec at 55° C and 1 minute 45 sec at 72° C and 25

cycles consisting of 30 sec at 94° C and 2 minutes 15 sec at 72° C. The PCR product (569 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+.

EXAMPLE 2

ASSEMBLY OF THE MVA⁺ TRANSCRIPTION UNIT

5 The transcription unit is assembled in a derivative of the expression vector pBAD-GFPuv (Clontech, Palo Alto, California; GenBank accession number U62637). This is a high copy number plasmid that belongs to the pMB1/ColE1 incompatibility group. The final transcription unit is composed of four ORFs coding for yPMD, hPMK, yMVK and ecIDI. The coding sequences are preceded by ribosomal binding sites that consist of a Shine-
10 Dalgarno sequence followed by an AT-rich translation spacer of eight bases (optimal distance to the ATG start codon; Makrides, *Microbiol. Rev.* 60:512+ (1996)). The whole construct is under control of the *P*_{BAD} promoter, which can be induced in the presence of L-(+)-arabinose and repressed in the presence of D-(+)-glucose and absence of L-(+)-arabinose. Lobell and Schleif, *Science* 250:528-532 (1990); Guzman *et al.*, *J. Bacteriol.* 177:4121-4130
15 (1995).

As a preliminary step, the *Nde* I restriction site located between pBR322ori and the *araC* coding region of pBAD-GFPuv (position 4926-4931) is eliminated by site-directed mutagenesis as described (Kunkel *et al.*, *Meth. Enzymology* 154:367-382, 1987), using the oligonucleotide pBAD-mut1 (SEQ ID NO: 63) as mutagenic primer. The mutation is
20 confirmed by restriction analysis and sequencing. The plasmid obtained is named pAB-M0. The GFP coding sequence of pAB-M0 is substituted by the yPMD coding sequence. This sequence was cloned between *Nde* I and *Eco* RI restriction sites, taking advantage of the modifications introduced at the ends of the yPMD sequence. The yPMD sequence is the first of the transcription unit.

25 To clone the other coding sequences, a polylinker is first introduced between *Eco*RI and *Sal* I restriction sites. The polylinker is generated by annealing the oligonucleotides pBAD-Link1 (SEQ ID NO: 64) and pBAD-Link2 (SEQ ID NO: 65). It contains the restriction sites *Pme* I and *Sna* BI, flanked by cohesive ends of *Eco* RI and *Sal* I sites. Sites *Pme* I, *Sna* BI and *Sal* I are preceded by the Shine-Dalgarno consensus sequence
30 "TAAGGAGG". The modified inserts coding for hPMK and yMVK are digested with *Hpa* I and blunt ligated, respectively, into *Pme* I and *Sna* BI restriction sites. The modified insert coding for ecIDI is digested with *Xho* I and ligated into *Sal* I restriction site. Insert orientation is confirmed after every step by PCR and sequencing.

The plasmid containing yPMD, hPMK and yMVK is named pAB-M2. The plasmid containing, in addition, ecIDI is named pAB-m3.

EXAMPLE 3

STABLE INTEGRATION OF THE MVA⁺ TRANSCRIPTION

5 UNIT INTO THE *E. coli* CHROMOSOME

Transfer of the MVA⁺ transcription unit to the chromosome from *E. coli* is achieved with a genetic system based in two elements: the *E. coli* strain TE2680 (Elliott, *J. Bacteriol.* 174:245-253, 1992) and a pRS550-derived plasmid (Simons *et al.*, *Gene* 53:85-96, 1987). Strain TE2680 is a *recD* (tet^r) mutant host that allows efficient recombination of a linear
10 (restriction enzyme-cleaved) DNA with homologous sequences present in the chromosome. The new sequence is incorporated as a single copy and is perpetuated through cell division.

The sequence of interest, the MVA⁺ transcription unit in this case, can be cloned in pRS550 vector, between a functional kanamycin resistance (Kan^R) gene and a promoterless version of the *lac* operon. A similar cassette is present in the recipient host (strain TE2680),
15 interrupting the *trp* operon. This strain is auxotrophic for tryptophan. In this case, however, a non-functional kanamycin resistance (Kan^S) gene and the deleted version of the *lac* operon are flanking a functional chloramphenicol resistance (Cam^R) gene. A double crossover affecting the *Kan* gene and the deleted version of the *lac* operon substitutes the sequence of interest for the Cam^R gene in the chromosome. As a consequence of the crossover, the
20 recipient strain, originally Kan^S and Cam^R, becomes Kan^R and Cam^S.

The MVA⁺ transcription unit is amplified by PCR using the pAB-M3 plasmid as template and oligonucleotides pBAD-D2 (SEQ ID NO: 66) and pBAD-U3 (SEQ ID NO: 67) as primers. The reaction mixture of the PCR is prepared in a final volume of 50 µl containing 200 ng of template, 1 µM of primers, 200 µM of dNTPs, *Pfu* reaction buffer and
25 1.75 units of *Pfu* DNA polymerase. The sample is denatured for 2 minutes at 94° C and incubated for 10 cycles consisting of 40 sec at 94° C, 50 sec at 59° C and 8 minutes 15 sec at 72° C. The amplified sequence (4126 bp) contains the complete promoter, including the regulatory sequences that respond to arabinose and glucose, and the four ORFs that allow conversion of MVA to IPP and DMAPP, but lacks the transcription termination signals that
30 are originally present in the expression cassette.

A polylinker is introduced in the vector pRS550, to allow cloning of the PCR product containing the MVA⁺ transcription unit. The polylinker is generated by annealing the oligonucleotides pRS-L1 (SEQ ID NO: 68) and pRS-L2 (SEQ ID NO: 69). It contains the restriction sites *Pme* I, *Sma* I/*Srf* I and *Not* I, flanked by cohesive ends of *Bam* HI and *Eco* RI.

sites. Plasmid pRS2110 is generated by cloning the polylinker between *Bam* HI and *Eco* RI restriction sites of vector pRS550. The MVA⁺ transcription is cloned in the *Pme* I restriction site of vector pRS2110, with the same orientation than the promoterless *lac* operon, thus restoring transcription of the *lac* operon. The plasmid obtained is named pRS-MVA⁺.

- 5 Plasmid pRS-MVA⁺ are digested with *Sal* I and *Sca* I restriction enzymes. This digestion rendered a 3196 bp fragment containing the ampicillin resistance gene and a 13406 bp fragment containing the Kan gene, the MVA⁺ transcription unit and the deleted version of the *lac* operon. Strain EcAB3-1 is obtained by transformation of strain TE2680 with the linear plasmid DNA. The presence of the MVA⁺ transcription unit in the chromosome of this strain is confirmed by PCR. The activity of this transcription unit is confirmed by the appearance of blue colonies in plates containing 5-bromo-4-chloro-3-indolyl β -D-galactopyranoside (Xgal). Strain EcAB3-1 is resistant to kanamycin (25 μ g/ml) and tetracycline (6 μ g/ml) and sensitive to chloramphenicol (17 μ g/ml) and ampicillin (50 μ g/ml). The MVA⁺ transcription unit is transduced to *E. coli* strain MG1655 using phage P1.
- 10
- 15 The strain obtained is named EcAB4-1.

EXAMPLE 4

IDENTIFICATION AND FEATURES OF THE *gcpE* GENE FROM *E. coli* AND A PUTATIVE HOMOLOG FROM *Arabidopsis thaliana*

- To identify genes potentially involved in the MEP pathway, a bioinformatic approach is adopted. Because bacterial genes with related functions are often organized in operons, uncharacterized open reading frames (ORFs) that are beside known genes of the MEP pathway are examined. An ORF of 1195 bp with unknown function is found just upstream of a DXS coding sequence of *Streptomyces coelicolor* (cosmid 6A5, Accession Number AL049485). This ORF is homologous to an essential gene of *Escherichia coli* named *gcpE* (Baker *et al.*, *FEMS Microbiol. Lett.* 94:175-180, 1992 (accession number X64451)). An homolog of this gene, named *aarC*, is identified in *Providencia stuartii* and described as an essential gene involved in density-dependent regulation of the 2'-N-acetyltransferase (Rather *et al.*, *J. Bacteriol.* 179:2267-2273, 1997). However, no precise function was assigned to the *aarC* gene.
- 20
- 25

- 30 The *gcpE* gene is broadly distributed in evolution. The occurrence of this gene in completely sequenced genomes strictly correlates with the occurrence of the gene encoding 1-deoxy-D-xylulose 5-phosphate reductoisomerase (*dxr*), which catalyses the first committed step of the MEP pathway. Fourteen out of 26 sequenced genomes contain both *dxr* and *gcpE*. Twelve of these sequenced genomes do not contain *dxr* nor *gcpE*. The *gcpE* gene is also

highly conserved in plants. *GcpE* homologs are found as EST entries in *Arabidopsis thaliana* (gb T46582, SEQ ID NO: 5), *Glycine max* (gb AW152929, SEQ ID NO: 6), *Lycopersicon esculentum* (gb AW040413, SEQ ID NO: 7), *Mesembryanthemum crystallinum* (gb AI822799, SEQ ID NO: 8), *Oryza sativa* (gb AA753160, SEQ ID NO: 9), *Zea mays* (gb
 5 AW126434, SEQ ID NO: 10), *Pinus taeda* (gb AW042702, SEQ ID NO: 11) and *Physcomitrella patens* (gb AW497432, SEQ ID NO: 12).

A cDNA clone from *Arabidopsis* coding for a *gcpE* homolog (EST clone 135H1T7, accession number T46582) is obtained from the *Arabidopsis* Biological Resource Center (ABRC). This clone encodes a full length protein. The cDNA contains an ORF of 2223 bp
 10 that encodes a protein of 740 amino acid residues (SEQ ID NO: 1). The *Arabidopsis gcpE* gene corresponding to this cDNA is located in chromosome V (genomic P1 clone MUP24, accession number AB005246). This gene contains 20 exons that extend along 4 kb of genomic sequence.

Alignment of the *E. coli* and *Arabidopsis gcpE* proteins shows high similarity but
 15 also striking differences. The first 75 amino acid residues of the *Arabidopsis* sequence constitute a region that is not present in the bacterial counterpart. A transit peptide for plastids is predicted at this region with the ChloroP V1.0 program.

According to this program, the processing site of the transit peptide would be located between Arg38 and Ser39 (CS-score
 20 2.392). *In vivo* import experiments to chloroplasts demonstrated that the N-terminal region of the *Arabidopsis* protein is a functional transit peptide for plastids.

The putative mature *gcpE* protein from *Arabidopsis* is significantly larger than the *E. coli* counterpart (78 versus 41 kDa). Although the two proteins align and show high similarity at the N- and C-terminal regions, the *Arabidopsis* isoform possesses several
 25 additional amino acid sequences between these two regions, particularly a domain of 268 amino acid residues (30 kDa) which is only present in the *Arabidopsis* protein (SEQ ID NO: 1).

EXAMPLE 5

DELETION OF THE *gcpE* CODING SEQUENCE IN THE *E. coli* GENOME

30 To confirm whether *gcpE* from *E. coli* is indeed involved in the MEP pathway, *gcpE* is deleted in strain EcAB3-1. As mentioned above, mutants of the MEP pathway can be rescued in this strain, in the presence of MVA. Deletion of the *gcpE* gene is accomplished by homologous recombination using construct GC5CAT3 as the donor cassette. In this construct, the *CAT* gene is surrounded by the *gcpE* flanking regions. Substitution of the *CAT*

gene for the *gcpE* coding sequence in the genome can be selected by chloramphenicol resistance.

Four PCR reactions are necessary to prepare the GC5CAT3 construct. First, a genomic region of 3231 bp, encompassing the *gcpE* ORF (1116 bp), together with flanking regions, is amplified by PCR, using genomic DNA from strain MC4100 as template. The reaction mixture of the PCR is prepared in a final volume of 50 μ l containing 250 ng of template, 0.4 μ M of primers 1PE (SEQ ID NO: 70) and 4PE (SEQ ID NO: 73), 200 μ M of dNTPs, 1 mM of $MgSO_4$, *Pfx* reaction buffer and 1.25 units of PLATINUM *Pfx* DNA polymerase (Life Technologies Inc., Rockville, Maryland). The sample is denatured for 2 minutes at 94 C and incubated for 30 cycles consisting of 40 seconds at 94 C, 50 seconds at 67 C and 3 minutes 30 seconds at 68 C.

The regions flanking the *gcpE* coding sequence are amplified by PCR using the PCR product of primers 1PE and 4PE as template. Primers 1PE (SEQ ID NO: 70) and 22PE (SEQ ID NO: 71) are used to amplify the 5' flanking region. In this PCR, primer 22PE generates a *Sma* I restriction site. Primers 3PE (SEQ ID NO: 72) and 4PE (SEQ ID NO: 73) are used to amplify the 3' flanking region. In this PCR, primer 3 PE generates a *Pme* I restriction site. The reaction mixtures of these PCRs are prepared in final volumes of 50 μ l containing 150 ng of template, 4 μ M of primers, 200 μ M of dNTPs, *Pfx* reaction buffer and 1.25 units of PLATINUM *Pfx* DNA polymerase. The samples are denatured for 2 minutes at 94 C and incubated for 10 cycles consisting of 40 seconds at 94 C and 2 minutes at 68 C. The PCR product corresponding to the 3' flanking region (1061 bp) is cloned in the *Sma* I restriction site of plasmid pBluescript SK+. The plasmid obtained is named GC3. Subsequently, the PCR product corresponding to the 5' flanking region (1102 bp) is cloned in the *Pme* I restriction site of plasmid GC3. The relative orientation of the 3' and 5' flanking regions is the same than that in the *E. coli* genome. The plasmid with the two *gcpE* flanking regions is named GC53.

The *CAT* gene is amplified by PCR using the plasmid pCAT19 (Fuqua, 1992) as template and oligonucleotide CAT1 (SEQ ID NO: 74) and CAT4 (SEQ ID NO: 75) as primers. The reaction mixture of the PCR is prepared in a final volume of 50 μ l containing 100 ng of template, 1 μ M of primers, 100 μ M of dNTPs, *Pfx* reaction buffer and 1.25 units of PLATINUM *Pfx* DNA polymerase. The sample is denatured for 2 minutes at 94 C and incubated for 20 cycles consisting of 40 seconds at 94 C, 50 seconds at 53 C and 1 minute at 68 C. The PCR product (960 bp) is cloned in the *Sma* I restriction site of plasmid GC53. The construct obtained is named GC5CAT3. In this construct, the *CAT* gene has the same orientation than the *gcpE* gene previously deleted.

Plasmid containing GC5CAT3 construct is digested with *Hind*III, *Xba* I and *Xho* I restriction enzymes to release the recombination cassette. This cassette is amplified by PCR using oligonucleotides 1PE (SEQ ID NO: 70) and 4PE (SEQ ID NO: 73) as primers. The PCR product is used to transform electrocompetent cells of strain EcAB3-1. These cells are
 5 plated on 2xTY medium containing 1.5 % agar (w/v), 17 µg/ml chloramphenicol, 6µg/ml tetracycline, 25 µg/ml kanamycin, 0.2 % (w/v) L-(+)-arabinose and 1 mM MVA.

The presence of the *CAT* gene in place of the *gcpE* coding sequence in the genome of transformants is confirmed by PCR using oligonucleotides 0PE and 5PE as primers. The identity of the PCR product is verified by restriction analysis. Oligonucleotides 0PE (SEQ
 10 ID NO: 76) and 5PE (SEQ ID NO: 77) are complementary to genomic sequences located outside of the region included in the recombination construct. Analysis of transformants confirms both the absence of the original *gcpE* gene and the presence of the *CAT* gene. The novel strain is named EcAB3-3.

Strain EcAB3-3 can grow only in the presence of MVA. A control strain carrying a
 15 disruption of *dxs* gene (EcAB3-2) is also auxotrophic for MVA.

EXAMPLE 6

IDENTIFICATION OF GCPE FUNCTION

Example 5 describes the generation of *E. coli* strain with a deletion of the *gcpE* coding sequence (strain EcAB3-3). In addition to the *gcpE* deletion the strain also carries a
 20 MVA⁺ transcription unit as described in Examples 1, 2 and 3 which makes it auxotrophic for mevalonic acid or mevalonate (MVA). This strain is used to find out which intermediate accumulates due to the disruption of the *gcpE* gene. The *gcpE* deletion disrupts the MEP pathway blocking the formation of IPP and DMAPP, creating the need for exogenous MVA to synthesize IPP and DMAPP.

25 A culture of the *E. coli* strain with a disrupted *gcpE* gene is made in the presence of MVA. After growth, the cells are harvested by centrifugation, washed with culture medium containing no MVA and resuspended for 16 hours in a culture medium containing [³H]ME (Methylerythritol). Thin layer chromatography separation of the water/ethanol (30:70) extract of the cells affords a radioactive band co-eluting with methylerythritol
 30 cyclodiphosphate (isopropanol/water/ethyl acetate, 60:30:10, R_f = 0.56). Carrier material is obtained for the latter compound from *Corynebacterium ammoniagenes* treated with benzylviologen. Additional data is collected, suggesting that the radioactive compound might correspond to methylerythritol cyclodiphosphate. On HF hydrolysis, it releases free methylerythritol. Like methylerythritol cyclodiphosphate, it is not affected by alkaline

phosphatase, which normally cleaves acyclic diphosphates. This compound is not accumulated by the *mva⁺/dxr⁻* *E. coli* strain with an intact *gcpE* gene. In the latter experiment [³H]ME is incorporated into ubiquinone and menaquinone, which are not labeled in the *gcpE* disrupted strain.

- 5 Further conformation of function for *gcpE* will require cell-free assays using radiolabeled methylerythritol cyclodiphosphate as described below.

EXAMPLE 7

GCPE ENZYME ASSAYS

Enzymatic preparation of [¹⁴C]methylerythritol 2,4-cyclodiphosphate

- 10 The substrate methylerythritol cyclodiphosphate cannot be readily chemically synthesized. Attempts to accumulate the tritiated compound from [³H]ME by the *mva⁺/dxr⁻/gcpE⁻* mutant described above result in very low yields. Enzymatic synthesis of [¹⁴C]methylerythritol cyclodiphosphate is thus required. This can be achieved using all the known enzymes of the MEP pathway, viz., *dxs*, *dxr*, *ygbP*, *ychB*, and *ygbB*.

- 15 Enzymatic syntheses of [¹⁴C]-deoxy-D-xylulose-5-phosphate (DXP) and MEP from [¹⁴C]pyruvate isotopomers and D-glyceraldehyde-3-phosphate (GAP) are performed using *E. coli* strains overexpressing *dxs* and *dxr* genes. In order to prepare the subsequent [¹⁴C]methylerythritol cyclodiphosphate from the [¹⁴C]MEP the following scheme is used.

- 20 Three *E. coli* strains are generated with each one overexpressing one of the three remaining genes in the MEP pathway, viz., *ygbP* (pQE31-*ygbP*, pREP4), *ychB* (pQE30-*ychB*, pREP4) and *ygbB* (pQE30-*ygbB*, pREP4). Each strain is grown on LB medium containing ampicillin and kanamycin at 37°C overnight. Each culture (2ml) is used to inoculate the same medium (50 mL), which are then grown for 3 hours until a 0.5 OD (600 nm) is reached, then induced using IPTG (final concentration 0.1 mM) for 4.5 hours. After
25 centrifugation, the cells of each culture are resuspended in 100 mM Tris-HCl (3 mL, pH 8) and disrupted by sonication (3 x 30 s with 1 min cooling) at 0 °C. After centrifugation, the supernatant is stirred for 1 hour at 0°C in the presence of a 50% Ni-NTA slurry (1 mL, Qiagen Inc., Valencia, California).

- 30 The lysate-Ni-NTA mixture is loaded onto a column and the flow-through is collected. The column is washed twice with 100 mM Tris-HCl (4 mL, pH8) containing 50mM imidazole. The proteins are eluted with 100 mM Tris-HCl (2 mL, pH 8) containing 200 mM imidazole. Additional 100 mM Tris-HCl (1.5 mL, pH 8) is added to each protein, and the resulting solution is dialyzed against 100 mM Tris-HCl (pH 8) containing 20%

glycerol. On a 12% SDS-PAGE gel, the 6xHis-tagged MEP cytidylyl transferase (*ygbP*), CDP-ME kinase (*ychB*) and 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (*ygbB*) are separated from other cellular components.

Using these pure proteins, [^{14}C]2-C-methyl-D-erythritol 2,4-cyclodiphosphate is prepared in a one-pot procedure. In a typical incubation, [^{14}C]MEP (10 μL , 2.27×10^6 cpm, 15.8 Ci/mol) is incubated with the purified MEP cytidylyl transferase (100 μL , 0.4 mg/mL), 6xHis-tagged CDP-ME kinase (200 μL , 0.15 mg/mL) and 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (200 μL , 0.6 mg/mL) solutions in 100 mM Tris-HCl (1 mL, pH 8) containing 5 mM CTP, 1 mM ATP, 5 mM MnCl_2 and 5 mM MgCl_2 . The incubation is performed at 37°C for 10 hours.

An aliquot (3 μL) is analyzed on a silica gel plate eluted with isopropanol/water/ethyl acetate (6:3:1). Radioactivity is monitored with a PhosphorImager. A single radioactive compound is detected. It coelutes with unlabeled 2-C-methyl-D-erythritol 2,4-cyclodiphosphate. No radioactivity is found comigrating with ME-CDP. An aliquot is incubated in the presence of alkaline phosphatase and no [^{14}C]methyerythritol is detected, indicating that no [^{14}C]MEP remained in the incubation mixture.

GCPE Enzyme Test

When purified His-tagged GCPE is assayed with the [^{14}C] 2-C-methyl-D-erythritol 2,4-cyclodiphosphate as prepared above there is no reaction product detected. One reason for lack of activity could be that GCPE needs other proteins to form a complex with diverting 2-C-methyl-D-erythritol 2,4-cyclodiphosphate into the two branches of the MEP pathway. Because of the genetic link of *yfgB* and *yfgA* with *gcpE* (all three are on the same operon of the *E. coli* genome), it is possible that these proteins could be part of this hypothetical enzyme complex. Thus, an expression plasmid containing the genomic region covering *yfgB*, *yfgA* and *gcpE* is constructed and stably transformed into *E. coli* creating the strain BL21(DE3)pLys[PET-T7-gcpE-yfgA-yfgB]. This strain and the BL21(DE3)pLys[PET-T7] and BL21(DE3)pLys[PET-T7-yfgA-yfgB] or [MVA^+ ,gcpE-PQE30-AT-gcpE] strains are grown and induced with IPTG using standard conditions.

In a typical experiment, the *E. coli* strain BL21(DE3)pLys[PET-T7-gcpE-yfgA-yfgB] is grown at 30°C in LB medium (50 mL) containing chloramphenicol (34 $\mu\text{g/mL}$) and ampicillin (100 mg/mL) until reaching a 0.65 OD (600 nm). Induction is then performed with IPTG (0.5 mM) for 6 hours. The cells are harvested by centrifugation (7000g, 10 min) resuspended in buffer (4 mL, 50 mM Tris HCl pH = 8, 1 mM PMSF, 1 mM DTT, 5 mM

MgCl₂) and broken at 0°C by sonication (2 x 30 s, with 1 min cooling). The cell debris is removed by centrifugation (16000 g, 10 min).

The resulting crude cell-free material (130 L) is completed with buffer (20 L) and used for the enzyme assays at 37°C for 7 hours and 20 hours with the [¹⁴C]2-C-methyl-D-erythritol 2,4-cyclodiphosphate solution (50 L) obtained as described above. Controls consist in the same mixture, but the enzyme preparation is replaced by buffer. After incubation, an aliquot (9 l) of each assay is analyzed on a silica plate eluted with isopropanol/water/ethyl acetate (6:3:1). Radioactivity is monitored with a PhosphoImager.

For unknown reasons, only the assay with *E. coli* BL21(DE3)pLys[PET-T7-gcpE-yfgA-yfgB] extract is successful. In all assays performed with enzyme preparations from other strains, the entire radioactivity comigrated with unlabeled 2-C-methyl-D-erythritol 2,4-cyclodiphosphate, indicating that no reaction occurred. The TLC migration profile is the same as that observed for the control without enzyme.

In the case of all assays performed with the cell system prepared from the BL21(DE3)pLys[PET-T7-gcpE-yfgA-yfgB] strain, there is decrease of the substrate concentration and the accumulation of a new compound. According to its TLC behavior ($R_f = 0.85$, isopropanol/water/ethyl acetate, 60:30:10), this compound corresponds to a non-phosphorylated derivative. Such a dephosphorylation is most likely, as the test is performed with a crude cell-free system containing probably phosphatases, and as no phosphatase inhibitor was added to the incubation buffer. Dephosphorylation of the reaction product might favor displacement of the reaction, the full consumption of the substrate and finally accumulation of a single major product.

The same compound is obtained when only MgCl₂ was present in the assay, suggesting that the cofactors tested are not necessary. It is possible that the fact the product is dephosphorylated *in situ* helped to its accumulation. The dephosphorylated new compound ($R_f = 0.56$, CHCl₃/CH₃OH, 8:2) is characterized by a R_f between those of methylerythritol ($R_f = 0.22$) and isopentenol ($R_f = 0.56$). TLC comparison with unlabeled synthetic carriers indicates that compounds 1 to 9 (shown in Figure 1) do not correspond to the non-phosphorylated new compound.

To fully characterize the dephosphorylated product, a larger-scale incubation (10X) is performed and the residue is acetylated (pyridine/Ac₂O, 10 ml) overnight. After the removal of the reagents, the residue is resuspended in CHCl₃ (12 ml) and the resulting precipitate is removed by filtration. The filtrate is concentrated to dryness (836000 cpm, 1.1g) and purified on a silica column (8g) eluted with hexane/ethyl acetate (3:1) and fractions of 5 ml are collected. An aliquot (4 l) of each fraction is spotted on TLC plates (hexane/ethyl

acetate, 3:1) and the radioactivity monitored by PhosphoImager. The radioactive fractions of same R_f are pooled together.

Three radioactive products can be detected: Fraction A (200 mg) contains the acetate of the dephosphorylated new compound ($R_f = 0.4$), fraction B (20 mg) contains the 2-C-methyl-D-erythritol triacetate ($R_f = 0.2$), and fraction C (100 mg) contains another new compound ($R_f = 0.25$) which is not yet identified. Fraction A is further purified on a silica column (9g) eluted first with CH_2Cl_2 in order to remove almost all impurities and then with ethyl acetate in order to recover the radioactive product. As previously described, an aliquot (4 l) of each 2 ml fraction is checked for radioactivity and the radioactive fractions are pooled together, concentrated to dryness and almost pure acetate of the dephosphorylated new compound (1 mg) is obtained.

This compound is analyzed by ^1H -NMR and from the resulting spectrum it is concluded that the acetate of the putative dephosphorylated GCPE product could be diacetate of (*E*)-2-methylbut-2-ene-1,4-diol. The spectrum is compared with a reference synthetic diacetate of (*E*)-2-methylbut-2-ene-1,4-diol synthesized by LiAlH_4 reduction of methylfumaric acid as previously described for the reduction of 3-methylfuran-2(5H)-one or citraconic anhydride (Duvold *et al.*, *Tetrahedron Letters* 38: 6181-6184, 1997). All signals of the enzymatic product match the corresponding signals in the synthetic standard. Furthermore the coelution of the enzymatic radioactive product and the synthetic diacetate of (*E*)-2-methylbut-2-ene-1,4-diol is observed (CH_2Cl_2 , $R_f = 0.25$). Therefore, one product of the incubation is identified as diacetate of (*E*)-2-methylbut-2-ene-1,4-diol (Figure 2). This positive identification suggests that the product of GCPE reaction with 2-C-methyl-D-erythritol 2,4-cyclodiphosphate is (*E*)-1-(4-hydroxy-3-methylbut-2-enyl) diphosphate (Figure 3).

25

EXAMPLE 8

CHARACTERIZATION OF *ARABIDOPSIS* GCPE

Upon identification of the *Escherichia coli* *gcpE* gene as involved in the trunk line of the MEP pathway for isoprenoid biosynthesis, the available databases are searched for plant homologs. As described in Example 4, clone 135H1 (Genbank accession number T46582) is identified as containing an *Arabidopsis thaliana* cDNA encoding a protein with homology to the product of the bacterial *gcpE* gene. As shown in Figure 4, however, the putative *Arabidopsis* GCPE protein (SEQ ID NO: 79), contains several domains that are absent from the *E. coli* protein (SEQ ID NO: 78). Identical residues are in black boxes and conservative changes in grey boxes. Gaps are indicated with dots. The predicted cleavage site for the

plastidial targeting peptide (according to the ChloroP program)
is indicated with an arrow (see Figure 4).

To determine whether the *Arabidopsis* protein encoded by clone 135H1 is indeed a GCPE protein, a complementation assay is carried out using the *E. coli* strain EcAB3-3. In this strain, which is engineered to synthesize IPP and DMAPP from mevalonic acid (MVA), the chromosomal *gcpE* gene is disrupted by insertion of the *CAT* marker conferring chloramphenicol resistance. Because the disruption of *gcpE* is lethal, mutant EcAB3-3 cells require MVA for growth (see Example 5).

For the complementation assay, plasmid pQE-AGH is created by subcloning a *Bgl*III-*Sph*I fragment (coding sequence SEQ ID NO: 80 and deduced amino acid sequence SEQ ID NO: 81) from clone 135H1 into the *Bam*HI-*Sph*I sites of the pQE30 expression vector (coding sequence SEQ ID NO: 82 and deduced amino acid sequence SEQ ID NO: 83) (Qiagen) (Figure 5). The resulting construct encodes a His-tagged protein (coding sequence SEQ ID NO: 84 and deduced amino acid sequence SEQ ID NO: 85) lacking the N-terminal sequence predicted to be a plastidial targeting peptide with the ChloroP program (Figure 5). Expression from plasmid pQE-AGH is under the control of the IPTG-inducible *T5* promoter. Figure 5 depicts the coding sequences in uppercase, and the deduced amino acid sequences are shown below the respective coding sequences. The predicted cleavage site for the plastidial targeting peptide is indicated with an arrow.

EcAB3-3 cells are transformed with plasmid pQE-AGH and plated on LB plates containing 100 mg/l kanamycin (to select for the MVA operon), 34 mg/l chloramphenicol (to select for the *gcpE* gene disruption), 100 mg/l ampicillin (to select for transformants containing pQE-AGH), 0.04 % arabinose (to induce expression of the MVA operon genes), and 0.5 mM MVA (to be used for IPP and DMAPP biosynthesis). The resulting strain, EcAB3-3(pQE-AGH), is able to grow in absence of MVA at 30°C and 37°C, confirming that MVA auxotrophy can be overcome by the presence of plasmid pQE-AGH. These results demonstrate that the cloned *Arabidopsis* cDNA encodes a protein with the same activity as the *E. coli* GCPE protein.

In order to study whether the truncated *Arabidopsis* GCPE protein cloned in plasmid pQE-AGH is active in converting ME-cPP to the next intermediate of the MEP pathway, the protein is expressed at high levels in *E. coli*. Strains XL1Blue or M15 (Qiagen Inc., Valencia, California) are used for expression under several experimental conditions: growth at 23°C, 30°C, or 37°C and induction with 1 or 0.4 mM IPTG, with unsuccessful results. When strain EcAB3-3(pQE-AGH) is used, however, expression of the cloned protein is detected.

An overnight culture of EcAB3-3(pQE-AGH) cells grown in LB medium supplemented with kanamycin, chloramphenicol, ampicillin, arabinose and with or without MVA at the concentrations described above is diluted 1:50 in fresh medium and incubated at 37°C until reaching an OD₆₀₀ of ca. 0.3. Although cells grew better when MVA is added to the medium, the presence of plasmid pQE-AGH is sufficient to allow growth in the absence of any exogenous source for isoprenoid synthesis. Expression of the truncated *Arabidopsis* GCPE protein is induced by adding IPTG to a final concentration of 0.4 mM.

After incubation at 30°C for 4 hours, cells are collected by centrifugation and resuspended in a 1/50 volume of homogenization buffer (Tris-HCl 20 mM pH 8.0, 1 mM - mercatoethanol, 1mg/ml lysozyme, 80 mg/l PMSF, and 1 tablet/20 ml of Complete Mini, EDTA-free Protease Inhibitor Cocktail Tablets (Roche Molecular Biosystems, Indianapolis, Indiana)). Following incubation at room temperature for 20 minutes, cells are sonicated 5 times for 30 seconds at 30W. The insoluble fraction is pelleted by centrifugation at 5000xg for 30 minutes and the supernatant (soluble fraction) is collected. Electrophoresis on SDS-PAGE of an aliquot of this soluble fraction shows that a protein of the expected size (ca. 78 kD) is expressed in cells grown with or without MVA.

Purification of the His-tagged protein from the soluble extract is carried out using HiTrap™ columns (Pharmacia, Uppsala, Sweden). Flux through the column is kept constant at 2.5 ml/min during all the steps. After applying the sample to a column and washing unbound proteins with 20 ml of washing buffer (20 mM Tris-HCl pH 8.0, 10 mM imidazole, 500 mM NaCl), elution is performed with 50 ml of a gradient solution containing from 10 mM to 500 mM imidazole and 2.5 ml fractions are collected afterwards. The truncated *Arabidopsis* GCPE protein elutes at 100 mM imidazole and is virtually pure.

EXAMPLE 9

PREPARATION OF PLANT EXPRESSION VECTORS WITH GCPE

Rice, soybean and *E. coli gcpE* genes are chosen for plant expression. An *E. coli* gene (SEQ ID NO: 3) is cleaved by *NcoI* / *EcoRI* restriction digest, gel purified, and ligated into *NcoI* / *EcoRI*-digested and gel purified pMON26541 resulting in the formation of a shuttle vector. These ligations fuse the bacterial *gcpE* gene to CTP1, which is the chloroplast target peptide of the small subunit of the ribulose bisphosphate carboxylase from *Arabidopsis*, and place it under e35S promoter control.

To place the *gcpE* gene under napin promoter control, the shuttle vector is digested with *EcoRI*, ends are filled in using the Klenow fragment, and the gel purified vector is digested with *Bgl* II. The smaller fragment encoding the *gcpE* gene fused to CTP1 is gel

purified. pCGN3224 is digested with *Pst*I, ends are filled in with Klenow fragment and subsequently the vector is digested with *Bgl* II and gel purified. The purified vector and the purified CTP1::*gcpE* fusion are then ligated into digested and gel purified pGCN3223.

To transfer the *E. coli gcpE* gene into an *Arabidopsis* binary vector, pGCN3223 is
 5 digested with *Hind*III and *Sac* I and the gel purified fragment carrying the e35S promoter fused to CTP1 and *gcpE* is ligated into *Hind*III / *Sac*I-digested and gel purified pMON26543, resulting in a vector containing *gcpE* under e35S promoter control. The pNapin binary expression vector is obtained by ligating the gel purified *Not*I fragment harboring the pNapin::*CTP1::gcpE::napin* 3' expression cassette into *Not*I digested pMON36176.

10 Seed-specific expression vectors for a rice *gcpE* (SEQ ID NO: 2) and a soybean *gcpE* (SEQ ID NO: 6) sequence are constructed using a pBin19 (Bevan, *Nucleic Acids Research* 12: 8711-8720, 1984) derivative. The plasmid contains the *Vicia faba* seed-specific promoter from the Legumin B4 gene (Bäumlein *et al.*, *Nucleic Acids Research* 14: 2707-2719, 1996), the sequence encoding the transit peptide of the *Nicotiana tabacum* transketolase (TkTp) (R.
 15 Badur, Ph.D. thesis, Georg August University of Göttingen, Germany, 1998) and the transcriptional termination sequence from the octopin synthase gene (Gielen *et al.*, *EMBO J.* 3:835-846, 1984). A rice *gcpE* (SEQ ID NO: 2) sequence is cloned in sense orientation as a *Bam* HI fragment into the *Bam* HI site of the pBin-LePTkTp9 vector, resulting in a recombinant rice *gcpE* expression vector. A recombinant soybean *gcpE* (SEQ ID NO: 6)
 20 expression vector is similarly created.

EXAMPLE 10

TRANSFORMATION OF PLANTS

Agrobacterium transformed with the vectors of Example 9, and with pQE-AGH (which contains the *Arabidopsis gcpE* gene), are prepared as follows. 100μl of an overnight
 25 culture is spread on an agar LB plate with antibiotics. The plate is placed upside down in a 30°C chamber overnight. The plates are removed after colonies have grown (24-48 hours). A small scale culture is started by placing 10 ml of liquid LB media in a 50 ml tube. 10μl Kanamycin (50 μg/μL), 10μl Spectinomycin (75-100 μg/μL), and 10μl Chloramphenicol (25 μg/μL) are added. *Agrobacterium* is added from a plate, and the tube is shaken and placed in
 30 a 30°C shaker overnight.

Following overnight growth of the 10 ml culture, the culture is removed to a 500 ml flask. 200 ml of liquid LB is placed in a flask, 200μl Kanamycin (50 μg/μL), 200μl Spectinomycin (75-100 μg/μL), and 200μl of Chloramphenicol (25 μg/μL) are added, and the entire 10ml overnight culture is then added. The 500 ml flask is placed in a 30°C shaker

and grown overnight. The entire 200 ml culture is placed in a centrifuge tube and centrifuged for 25 minutes at 3,750 rpm and 19°C. After centrifugation, the liquid is poured off and the pellet is resuspended in 25 ml of 5% Sucrose (0.05% Silwet™) solution.

900µl of the sucrose solution and 100µl of the 25 ml bacterial culture are placed in a
5 cuvette, and the cuvette is shaken with a covering of parafilm. A blank OD reading is taken with 1 ml of sucrose solution, and then readings of all the bacterial solutions are taken. The OD (at a wavelength of 600) of each culture is recorded. The following calculations are then performed: $C_1V_1 = C_2V_2$; $C_1V_1 = (0.8)(200\text{ml})$; $C_1V_1 = 160$; $V_1 = 160 / C_1$; and $V_1 = X \text{ ml}/10$ to determine $\text{OD}_{600} = 0.8$ of an *Agrobacterium* culture.

10 Plants are soaked for at least 30 minutes in water prior to dipping. The bacterial solution is poured into a shallow plastic container, and above ground parts of the plant (bolts, rosettes) are dipped into the solution for 3-5 seconds with gentle agitation. Dipped plants are placed on their side in a diaper lined black tray, and covered by a dome overnight (16-24 hours) to maintain a high humidity. The cover is removed and normal plant growth
15 conditions are resumed for 4 weeks.

Following the transformation and high humidity treatment, plants are maintained at 22°C, 60% RH, and a 16 hour photoperiod for 4 weeks. 5-7 days after transformation, plants are coned. Fertilization with a weak 20-20-20 fertilizer is done weekly. After 4 weeks of growth, plants are placed in the greenhouse and all watering is stopped to encourage plant
20 dry down for seed harvest. Plants are ready for seed harvest after 1-1.5 weeks of dry down. Seeds are harvested by cutting the base of the plant below the cones, holding the plant over a seed sieve and a white piece of paper, running bolts through the cone hole, and collecting clean seeds through sieving.

Seeds are sterilized by connecting a vacuum desiccator hose to a vacuum in a fume
25 hood/flow bench. 100 ml of bleach is placed in a 250 ml beaker, and 3 ml of concentrated HCl is added to the bleach. The beaker is placed in the desiccator, and seeds in seed tubes in a tube holder are placed in the desiccator. A cover is placed on the desiccator, and the vacuum is operated. The desiccator is left overnight but no longer than 16 hours.

Once sterilized, seeds are plated on selection media (prepared by adding 10g (2g/L)
30 Phyta-Gel, 10.75 g (2.15 g/L) MS Basal Salts (M-5524 from Sigma), 50 g (10g/L) sucrose, and 6 ml (1.2 ml/L) Kanamycin solution (950mg/ml), 5ml (1ml/L) Cefotaxime Solution (250 mg/ml), and 5 ml (1 ml/L) Carbenecillin Solution (250 mg/ml) to a total volume of 5 liters at a pH of 5.7). Seed tubes are tapped lightly over a plate in order to distribute the seeds sparsely. The plates are wrapped in parafilm and placed in a 4°C refrigerator for 1-2 days of

cold treatment. After this cold treatment the plates are placed in a 28°C chamber for germination.

Selected plantlets are green and have secondary leaves developing. The selected plantlets are moved to soil after secondary leaves have developed. The plantlets are potted in
5 soil and covered with a dome for 5 days to maintain high humidity. The plantlets are moved to a greenhouse after the bottom siliques begin to turn yellow.

Seeds from the selected plantlets are grown in 2.5 inch pots with soil (½ Metro-200; ½ PGX Mix). The soil is mounded and the pot is covered with mesh screen. The screen is fastened to the pot with a rubber band. Seeds are sown and covered with a germination
10 dome. The seedlings are grown in a 12 hour photoperiod in 70% relative humidity at 22°C. Water is supplied every other day as needed and Peter's 20-20-20 fertilizer is applied from below, bi-weekly.

EXAMPLE 11

PRODUCTION OF SEEDS FROM TRANSGENIC PLANTS

15 Transgenic seed plants from Example 10 representing 20 independent transformation events are grown and seeds harvested to produce T₂ seeds. The T₂ seeds are grown and tested for tocopherol levels. Tocopherol levels are determined by adding 10 to 15 mg of *Arabidopsis* seed into a 2 mL microtube. A mass of 1 g of 0.5mm microbeads (Biospecifics Technologies Corp., Lynbrook, NY) and 500 µl 1% pyrogallol (Sigma Chem, St. Louis, MO)
20 in ethanol containing 5 µg/mL tocol, are added to the tube. The sample is shaken twice for 45 seconds in a FastPrep (Bio101/Savant) at a speed of 6.5. The extract is filtered (Gelman PTFE acrodisc 0.2 µm, 13 mm syringe filters, Pall Gelman Laboratory Inc, Ann Arbor, MI) into an autosampler tube. HPLC is performed on a Zorbax silica HPLC column, 4.6 mm x 250 mm (5 µm) with a fluorescent detection using a Hewlett Packard HPLC (Agilent
25 Technologies, Palo Alto CA). Sample excitation is performed at 290 nm, and emission is monitored at 336 nm. Tocopherols are separated with a hexane methyl-t-butyl ether gradient using an injection volume of 20 µl, a flow rate of 1.5 ml/min, and a run time of 12 min (40°C). Tocopherol concentration and composition is calculated based on standard curves for α, β, δ, and γ-tocopherol using Chemstation software (Agilent Technologies, Palo Alto
30 CA).

EXAMPLE 12

TRANSGENIC PLANTS WITH GCPE AND OTHER TOCOPHEROL BIOSYNTHESIS
GENES

Canola, *Brassica napus* and soybean plants are transformed with a variety of DNA
5 constructs using a particle bombardment approach
or using *Agrobacterium* mediated transformation. Two sets of DNA constructs are produced.

The first set of constructs are "single gene constructs" in which the *gcpE* gene is
inserted into a plant DNA construct under the control of an arcelin 5, 7S alpha or napin
promoter (Kridl *et al.*, *Seed Sci. Res.* 1:209-219, 1991). The products of the *gcpE* gene can
10 be targeted to the plastid by an encoded plastid target peptide such as CTP1 (Keegstra, *Cell*,
56(2):247-253, 1989; Nawrath, *et al.*, *PNAS* 91:12760-12764, 1994).

A second set of DNA constructs is generated and referred to as the "multiple gene
constructs". The multiple gene constructs contain multiple genes each under the control of a
napin promoter and the products of each of the genes are targeted to the plastid by an
15 encoded plastid target peptide, such as a natural plastid target peptide present in the trans
gene, or an encoded plastid target peptide such as CTP1.

The multiple gene construct contains the *gcpE* gene and one or more genes for other
MEP pathway proteins, including, but not limited to: a *ygbB* gene; a *ygbP* gene; a *ychB* gene;
a *yfgA* gene; a *yfgB* gene; a bifunctional prephenate dehydrogenase such as the *E. herbicola*
20 or *E. coli tyrA* gene (Xia *et al.*, *J. Gen. Microbiol.* 138:1309-1316, 1992), a
phytylprenyltransferase such as the *slr1736* gene (in Cyanobase)
or the *ATPT2* gene (Smith *et al.*, *Plant J.* 11: 83-92, 1997), a
deoxyxylulose synthase such as the *E. coli dxs* gene (Lois *et al.*, *PNAS* 95(5):2105-2110,
1998), a deoxyxylulose reductoisomerase such as the *dxr* gene (Takahashi *et al.* *PNAS*
25 95(17), 9879-9884, 1998), an *Arabidopsis thaliana* HPPD gene (Norris *et al.*, *Plant Physiol.*
117:1317-1323, 1998), an *Arabidopsis thaliana* GGPPS gene (Bartley and Scolnik, *Plant*
Physiol. 104:1469-1470, 1994), a transporter such as the *AANT1* gene (Saint Guily, *et al.*,
Plant Physiol. 100(2):1069-1071, 1992), a GMT gene (WO 00/32757, WO 00/10380), an
MT1 gene, a tocopherol cyclase such as the *slr1737* gene (in Cyanobase) or its *Arabidopsis*
30 ortholog, an isopentenyl diphosphate isomerase (IDI) gene, and an antisense construct for
homogentisic acid dioxygenase (Sato *et al.*, *J. DNA Res.* 7 (1):31-63, 2000).

Each construct is transformed into at least one canola, *Brassica napus* and soybean
plant. Plants expressing each of these genes are selected to participate in additional crosses.

The tocopherol composition and level in each plant is also analyzed using the method set forth in Example 11.

5 The tocopherol composition and level in each plant generated by the crosses (including all intermediate crosses) is also analyzed using the method set forth in Example 11. Progeny of the transformants from these constructs will be crossed with each other to stack the additional genes to reach the desired level of tocopherol.

10 Crosses are carried out for each species to generate transgenic plants having one or more of the following combination of introduced genes: *gcpE*, *ygbB*, *ygbP*, *ychB*; *yfgA*; *yfgB*; *tyrA*, *slr1736*, *ATPT2*, *dxs*, *dxr*, *GGPPS*, *HPPD*, *GMT*, *AANT1*, *slr1737*, *IDI* and an antisense construct for homogentisic acid dioxygenase.

15 The above description, sequences, drawings and examples are only illustrative of preferred embodiments that achieve the objects, features and advantages of the present invention. The scope of the claims should not be limited by the preferred embodiments set forth herein, but should be given the broadest interpretation consistent with the description as a whole.

1

gtt gat gag gtt atg aga ata gcg gat aaa gga gct gat att gta agg	558
Val Asp Glu Val Met Arg Ile Ala Asp Lys Gly Ala Asp Ile Val Arg	
120 125 130 135	
ata act gtt caa ggg aag aaa gag gcg gat gcg tgc ttt gaa ata aaa	606
Ile Thr Val Gln Gly Lys Lys Glu Ala Asp Ala Cys Phe Glu Ile Lys	
140 145 150	
gat aaa ctc gtt cag ctt aat tac aat ata ccg ctg gtt gca gat att	654
Asp Lys Leu Val Gln Leu Asn Tyr Asn Ile Pro Leu Val Ala Asp Ile	
155 160 165	
cat ttt gcc cct act gta gcc tta cga gtc gct gaa tgc ttt gac aag	702
His Phe Ala Pro Thr Val Ala Leu Arg Val Ala Glu Cys Phe Asp Lys	
170 175 180	
atc cgt gtc aac cca gga aat ttt gcg gac agg cgg gcc cag ttt gag	750
Ile Arg Val Asn Pro Gly Asn Phe Ala Asp Arg Arg Ala Gln Phe Glu	
185 190 195	
acg ata gat tat aca gaa gat gaa tat cag aaa gaa ctc cag cat atc	798
Thr Ile Asp Tyr Thr Glu Asp Glu Tyr Gln Lys Glu Leu Gln His Ile	
200 205 210 215	
gag cag gtc ttc act cct ttg gtt gag aaa tgc aaa aag tac ggg aga	846
Glu Gln Val Phe Thr Pro Leu Val Glu Lys Cys Lys Lys Tyr Gly Arg	
220 225 230	
gca atg cgt att ggg aca aat cat gga agt ctt tct gac cgt atc atg	894
Ala Met Arg Ile Gly Thr Asn His Gly Ser Leu Ser Asp Arg Ile Met	
235 240 245	
agc tat tac ggg gat tct ccc cga gga atg gtt gaa tct gcg ttt gag	942
Ser Tyr Tyr Gly Asp Ser Pro Arg Gly Met Val Glu Ser Ala Phe Glu	
250 255 260	
ttt gca aga ata tgt cgg aaa tta gac tat cac aac ttt gtt ttc tca	990
Phe Ala Arg Ile Cys Arg Lys Leu Asp Tyr His Asn Phe Val Phe Ser	
265 270 275	
atg aaa gcg agc aac cca gtg atc atg gtc cag gcg tac cgt tta ctt	1038
Met Lys Ala Ser Asn Pro Val Ile Met Val Gln Ala Tyr Arg Leu Leu	
280 285 290 295	
gtg gct gag atg tat gtt cat gga tgg gat tat cct ttg cat ttg gga	1086
Val Ala Glu Met Tyr Val His Gly Trp Asp Tyr Pro Leu His Leu Gly	
300 305 310	
gtt act gag gca gga gaa ggc gaa gat gga cgg atg aaa tct gcg att	1134
Val Thr Glu Ala Gly Glu Gly Glu Asp Gly Arg Met Lys Ser Ala Ile	
315 320 325	
gga att ggg acg ctt ctt cag gac ggg ctc ggt gac aca ata aga gtt	1182
Gly Ile Gly Thr Leu Leu Gln Asp Gly Leu Gly Asp Thr Ile Arg Val	
330 335 340	
tca ctg acg gag cca cca gaa gag gag ata gat ccc tgc agg cga ttg	1230
Ser Leu Thr Glu Pro Pro Glu Glu Glu Ile Asp Pro Cys Arg Arg Leu	
345 350 355	
gct aac ctc ggg aca aaa gct gcc aaa ctt caa caa ggc gca ccg ttt	1278
Ala Asn Leu Gly Thr Lys Ala Ala Lys Leu Gln Gln Gly Ala Pro Phe	
360 365 370 375	

gaa gaa aag cat agg cat tac ttt gat ttt cag cgt cgg acg ggt gat	1326
Glu Glu Lys His Arg His Tyr Phe Asp Phe Gln Arg Arg Thr Gly Asp	
380 385 390	
cta cct gta caa aaa gag gga gaa gag gtt gat tac aga aat gtc ctt	1374
Leu Pro Val Gln Lys Glu Gly Glu Glu Val Asp Tyr Arg Asn Val Leu	
395 400 405	
cac cgt gat ggt tct gtt ctg atg tcg att tct ctg gat caa cta aag	1422
His Arg Asp Gly Ser Val Leu Met Ser Ile Ser Leu Asp Gln Leu Lys	
410 415 420	
gca cct gaa ctc ctc tac aga tca ctc gct aca aag ctt gtc gtg ggt	1470
Ala Pro Glu Leu Leu Tyr Arg Ser Leu Ala Thr Lys Leu Val Val Gly	
425 430 435	
atg cca ttc aag gat ctg gca act gtt gat tca atc tta tta aga gag	1518
Met Pro Phe Lys Asp Leu Ala Thr Val Asp Ser Ile Leu Leu Arg Glu	
440 445 450 455	
cta ccg cct gta gat gat caa gtg gct cgt ttg gct cta aaa cgg ttg	1566
Leu Pro Pro Val Asp Asp Gln Val Ala Arg Leu Ala Leu Lys Arg Leu	
460 465 470	
att gat gtc agt atg gga gtt ata gca cct tta tca gag caa cta aca	1614
Ile Asp Val Ser Met Gly Val Ile Ala Pro Leu Ser Glu Gln Leu Thr	
475 480 485	
aag cca ttg ccc aat gcc atg gtt ctt gtc aac ctc aag gaa cta tct	1662
Lys Pro Leu Pro Asn Ala Met Val Leu Val Asn Leu Lys Glu Leu Ser	
490 495 500	
ggt ggc gct tac aag ctt ctc cct gaa ggt aca cgc ttg gtt gtc tct	1710
Gly Gly Ala Tyr Lys Leu Leu Pro Glu Gly Thr Arg Leu Val Val Ser	
505 510 515	
cta cga ggc gat gag cct tac gag gag ctt gaa ata ctc aaa aac att	1758
Leu Arg Gly Asp Glu Pro Tyr Glu Glu Leu Glu Ile Leu Lys Asn Ile	
520 525 530 535	
gat gct act atg att ctc cat gat gta cct ttc act gaa gac aaa gtt	1806
Asp Ala Thr Met Ile Leu His Asp Val Pro Phe Thr Glu Asp Lys Val	
540 545 550	
agc aga gta cat gca gct cgg agg cta ttc gag ttc tta tcc gag aat	1854
Ser Arg Val His Ala Ala Arg Arg Leu Phe Glu Phe Leu Ser Glu Asn	
555 560 565	
tca gtt aac ttt cct gtt att cat cac ata aac ttc cca acc gga atc	1902
Ser Val Asn Phe Pro Val Ile His His Ile Asn Phe Pro Thr Gly Ile	
570 575 580	
cac aga gac gaa ttg gtg att cat gca ggg aca tat gct gga ggc ctt	1950
His Arg Asp Glu Leu Val Ile His Ala Gly Thr Tyr Ala Gly Gly Leu	
585 590 595	
ctt gtg gat gga cta ggt gat ggc gta atg ctc gaa gca cct gac caa	1998
Leu Val Asp Gly Leu Gly Asp Gly Val Met Leu Glu Ala Pro Asp Gln	
600 605 610 615	
gat ttt gat ttt ctt agg aat act tcc ttc aac tta tta caa gga tgc	2046
Asp Phe Asp Phe Leu Arg Asn Thr Ser Phe Asn Leu Leu Gln Gly Cys	
620 625 630	

aga atg cgt aac act aag acg gaa tat gta tcg tgc ccg tct tgt gga 2094
 Arg Met Arg Asn Thr Lys Thr Glu Tyr Val Ser Cys Pro Ser Cys Gly
 635 640 645
 aga acg ctt ttc gac ttg caa gaa atc agc gcc gag atc cga gaa aag 2142
 Arg Thr Leu Phe Asp Leu Gln Glu Ile Ser Ala Glu Ile Arg Glu Lys
 650 655 660
 act tcc cat tta cct ggc gtt tcg atc gca atc atg gga tgc att gtg 2190
 Thr Ser His Leu Pro Gly Val Ser Ile Ala Ile Met Gly Cys Ile Val
 665 670 675
 aat gga cca gga gaa atg gca gat gct gat ttc gga tat gta ggt ggt 2238
 Asn Gly Pro Gly Glu Met Ala Asp Ala Asp Phe Gly Tyr Val Gly Gly
 680 685 690 695
 tct ccc gga aaa atc gac ctt tat gtc gga aag acg gtg gtg aag cgt 2286
 Ser Pro Gly Lys Ile Asp Leu Tyr Val Gly Lys Thr Val Val Lys Arg
 700 705 710
 ggg ata gct atg acg gag gca aca gat gct ctg atc ggt ctg atc aaa 2334
 Gly Ile Ala Met Thr Glu Ala Thr Asp Ala Leu Ile Gly Leu Ile Lys
 715 720 725
 gaa cat ggt cgt tgg gtc gac ccg ccc gtg gct gat gag tag 2376
 Glu His Gly Arg Trp Val Asp Pro Pro Val Ala Asp Glu
 730 735 740
 atttcaaaac ggagaaagat ggggtgggcca ttctttgaaa actgtgagag aagatatata 2436
 tatttgtgtg tgtatatcat ctgtttgttg tgtattgcat catcattttg aacaaatgtc 2496
 caaatctctt aagttgataa aagt 2520

<210> 2
 <211> 33675
 <212> DNA
 <213> Oryza sativa

<220>
 <221> CDS
 <222> (6924)..(7019), (7163)..(7269), (7344)..(7444), (7525)..(7634),
 <222> (7694)..(7813), (7923)..(8153), (8253)..(8369), (8515)..(8589),
 <222> (9012)..(9071), (9163)..(9225), (9328)..(9472), (9589)..(9730),
 <222> (9951)..(10028), (10134)..(10293), (10694)..(10798),
 <222> (11028)..(11129)

<220>
 <221> unsure
 <222> (1..33675)
 <223> unsure at all n locations

<400> 2
 ctttaaccctc gccgactgcc tggagattcg tgccgatcga tacacgtggc agcgcctaac 60
 gcgtaacccc tccctcactt ggagattcgt gcaagcaact cgattaatgc attaatgctg 120
 tcgcgtaggt ttccctacgg aagagctgag tttcgtaacg aaaaaaaccg gccacgtttc 180
 gcatcgagcc tacttttaatt agcgtgggaa aataattcaa agtagcgacc tgtaccctgt 240

ggcaacctag cgcgcgcggc catggctctt gttccgctcg tgacagtgct cctgttcgcc 300
 ggctcatgcc tcggatcagc gccgccgacg acatcgccgg cggcgtcggc ggcgtccacg 360
 gcgacacgta cggtagtagt cgacggcatt acggccatct acaaacctcg gcgactcgct 420
 gtccgacacc gcaacctcgc caggcaaggc gccaccggcg ggctgctccg gtacaccacg 480
 aggcttcctt acggcgtcac cgtcggccgc gccaccggcc ggtgctccga cggctacctc 540
 atcatcgact tcctcgggtga cgtcatcagt ttaattttct tctctcttcc gtctgaaaaa 600
 tggaagaaac aatattatat tacgttatat atatatgcgt ttttgtttcg gattaaattg 660
 tggatatgat cgatcgatgt gcagctagag atcttggcct ccctctgctc aacccgtaac 720
 tcgacgaggg cgcggacttc gccacggcg tcaacttcgc cgtcgcggcg gccaccgcgc 780
 tcaacacgac ggcgctcgcc gccaggcgga tcaccgtccc ccacaccaac agccccctcg 840
 acgtgcagct cagatTTTTT ttgttttaga gaagggtatt ttttaccgg cctctacatc 900
 caaccggata tatacggcta ttgaagtagg gaacttaacc ctgtaaacia tccatccata 960
 gaggatatga acctaagacc ttgaggtaact acttcaaccg gatataatcg tgcagctcag 1020
 atggttcaag gaattcatga actccacaac tagttctcct caagggtgaac gaacaaactg 1080
 aaacgcattt cagcttaatt tcgaccggtg cctgatcagt gccagtcagc aatgctgtat 1140
 ctcacaaata attaaagctaa tgtacagctt ttcagtgcta gaatgacttt catatagaga 1200
 aatcttgtgt tatatatata tacttttttc tgaaagaaaa aagttctttt gtgtgagcat 1260
 tgcattgcag agatccgtga aaagctgtcg aagtcactgg ttatgctggg agagatcgga 1320
 ggaaacgact acaactacgc ctctctccag acctggccga tggacgggtg atacagcctc 1380
 ggcaacgtca cacgcatgat cgaaagcggt gccacggccg tcgatcttgt accggaagtc 1440
 gtgcagtcca tagccagcgc agccaaggta cacaccattc ttttccatta atttttggga 1500
 ccttatTTTT aaaataataa tcctggctac aaagtaatta attaagaact aaattaattt 1560
 ttgtgggttt tgtgacacag gaggtgctcg acatgggcgc gacgcgggtg gtgatcccgg 1620
 gcaacctccc gctgggttgc gtgccgagct acatgagcgc ggtgaacgcg acggaccggg 1680
 cggcgtacga cgcgcgcgga tgctcgtcg cgctcaacct ctctgcggcg ctgcacaacg 1740
 cgtggctgcg ccgcgccgtc ggggagctgc ggcgcgcgta ccggggcgcc gcggtggtcg 1800
 cgtacgcgga ctactccgcc gcgtacgcgc cgacgctgga cggggcagcg gcgctcggct 1860
 tcgacgagcg gcgcgtgttc agggcgtgct gcggcaaggg cggcgggggc gcgtacgggt 1920
 tcgacgtgcg cgcgatgtgc ggcgcgccgg ggacggcggc gtgcgcggac ccggggaggt 1980
 acgtgagctg ggacggcgtc cacctgacgc agcgcgcgta cggcgtcatg gccgagctgc 2040
 tgttccgccc tggcctcgtg caccgcctc cgataaattt cacgaacagc gcgcgcgcgt 2100
 gaggcggtgt tgcattggctt gcgcgttttt tctgatcaaa actactcaag tttgagccgt 2160

```

tttgatttat aaataaaacc atatgcgatt ttgctaaacg tttgtcgcgt gatttctctt 2220
cggaagaaaa aatctcacc c gagt gatgca taggcgggtcc caaccatatg tgccctgacc 2280
tttctctgct tccttcgcgt cgtgcactga caacctcaca gtatgttttt ggtatgggcg 2340
cttgcgggccc aactcaatct gtaatacatt gggctgtcgt attgggtttg ttggacttca 2400
tagactggat cggagaaagt tgggtaattg actttttcat ttttgctata aaatgattaa 2460
ttaaacagtc taggataatt actgtagact ctaataatat tgtttggtta agtattatta 2520
tacattcctg tatttgacac tctaagagca tggccaagag ttgcctgaaa gtctcttcct 2580
aaatctgcct ttcattctct aatgagaatt taaggattaa aaatatactt attttcaata 2640
gacagcataa atttaattcc ctagaataaa aaaatgcccc cctaacaaca gaaattagat 2700
tcctctaccc gcacctcatc agatcgctcg atttaagatc acgccatctg acaccgccct 2760
cccgctcgct cttctctagt gtgggagtct cgcgctcaag agacggaaat cgggaacaag 2820
aatgattcct agcttagcga gaatgaaggg gaagacatat gtcataccta caccacata 2880
agtatgcct agcacaaggg atgaaaacgg atcgaaaacg gatggaaact agctttatca 2940
tattcgtttt catttttttt tcggaatcgg attcgaaatc gaaaactcgg atacggaaat 3000
aaaattgaat attatcgaat acagatacgg agcgaatata agatggaacg aatacagtag 3060
cgaatattta ccggtatata aaaaaccct caaattgagt ttcttgatta agaaagagat 3120
atcgcttatt attttagtta aatatctcca acatttatat cgtcaatttt atagacggtt 3180
ccacaatcgt atgtgaaaat cgattttcat ggttgttcct ctaagagatc catatgcaaa 3240
tatgattatc attttctatt ccaagacctt ttactagatg tataacttat ttaccattgc 3300
ataaattgga gatgttattt attttacttc acatcttcga aacttgtaat gtatgtatta 3360
tactttaaat gctttcaagt acaaagtta taaactacaa agtggttagat cccgttgagc 3420
tctacaactt tgatatggaa cacatctcca tcagatgtcg tttgaattgt agatctgaga 3480
ttttgtaaaa tttaatatgg tatattataa tgaatattta gacccttaaa tgaccttaaa 3540
taataaaata gtcaataata aagttgtaga tctcatcgag ctctataatg ttgatatgaa 3600
gtttgtcttc atctgattcc gtatgaaaaa gttatgtata tatacatgtt tttttataaa 3660
atltgctcaa tatctgcgga tatccgaaaa aaatttcgga tagtttttaa ccgtttttcg 3720
attccgatgg atagtatcct tactgtattc gttttcgttt ccgagaaaaa atatccaaat 3780
tcgtttccga atccgagaat ttttggataa ttccgacaga aactatccga atccgaaaaa 3840
tggttcggac ggacggaaac tatccaaacc agtttcatcc cggctagcac gcatttaaatt 3900
tcacatgagg ttgcacattt atctgaggta aaaagattgg aaacgggttac tggttcgtca 3960
agaattttcc gtatttatca gtataactat tcaatgacga catcaacata acagaaaatt 4020
aaaacaacat gagtcgattt tatatataac tagaaacgaa aacagtataa ctgttacgaa 4080

```

aacactagat tgatgggtcg aaaatttcca ccacgggtttt tatgcctacc tttcaacgct 4140
 cccaaagttc ccacgaccca aaacatgtgt gggagaactt ccgcccacat ggagacgggt 4200
 gtctcagga aacgtgccat ctgctttgct ccagggtcaac acatgtggtg tgactgaact 4260
 ggccatcgtc tcaatattgt catctacccg tcataccatg ccaccggacc agaaggatgat 4320
 tatgggtcttc ggcggccgtc gcgcgcggat gccttgctcc acaacaagtc agccgctcaa 4380
 accacactcc cctttggcat tgaacatgag gtttgacgac gatgtgtgtg tatgtttggg 4440
 caggtagctt tgtttcaagc tgcactagct aattaagatc gatctccttg tcaaagtcac 4500
 gatcaaacat cgaaagtaca tgcattggaag aaatgttgaa atgtaatgaa ctaaattgatg 4560
 tccttttctc cccttattaa acaacatcaa gtttctttta tttctaaaga atgttaatat 4620
 cctttttatt tcttcaataa atagtactgc actccctatg gtttttggtg ttttagcatct 4680
 tgactttcgg gcatacgttt tatgatattat cttattacaa aatataatta tcattttattt 4740
 tatcattaca aatactttta aaataacatt atcagctgat tttgaattaa aactaaaatt 4800
 acaccttaat tacaatatac ttcacatagc aattataata taactatata caacttacac 4860
 tataagttat gttcaaaaata tttttcctac aaaaactatc accagattct tagacagtcc 4920
 cattccacca cctcagctgc cgtgaaagaa ctttgggtct taaataagtc caaatttatc 4980
 tttttgtttt ctcaataaaa tattcgaatt atccaacaaa tcaaggaaaa aacatccttc 5040
 gatgacccat gaatattcgt gaagtttctc ctctagccag taacaatacg gaacaatcag 5100
 acaattttat ctggctcaag caccatctct cgcaccagat taaactattt ttttttcatg 5160
 gtacaataca atcccatgcc ggccacgaaa aacaaatggc agaaataata aacgaacaaa 5220
 acagcctctc tccatcgtga actaataaaa aataaaaata aaacaaaaca aaatgataat 5280
 ggaattacga agcgcattggg aaaacgacgg gcacgattaa atcatggcgg ggagagcccg 5340
 gaacccact tccacacctc caacccacg ccgtcagcct tcccctccca tgcacccgggt 5400
 ccaccaacac ctcatctctt ggacccca cgcagccact gccacggca acgcggtgct 5460
 cgtgcaccga gtccacacga cgcgcgcgc ggtgcggggg cgccggcctc tggggataaa 5520
 tgggctaate cggtagaaag cccaccactc gctcgccagt tcgtcgtcct cttcgccgag 5580
 ctgcgagct ctgcactct gtctccatcc ccgcatcgca tcgcctcgcc gctgctgatc 5640
 tcgtcgcggt cgccggaggg gagctacgag gttggggagc cttatctcta cttcctgaga 5700
 tttctagtag ctttgtgtat gtgtgtgtgt ttgtgtgttg gggggacgcc gatcgggtgg 5760
 atcctcctgt ggtggttggg tgggcgcaat tcgtgcttgg tttatttgct ggaattctag 5820
 cgggggagct ggcgttgctg gtgctaattg ctgcggggga gctgctggaa ttcgtgcttc 5880
 tgcttgggaa ttagaagggt tgggttttta tgattcagag ggctgtagag ctcttgagat 5940
 tggctgcgaa aattcgggat ttgatcaact tagagagcat tatctttgga ttaggaggga 6000

```

tttttcttaa tttttcttag ttttttttga gctatcaaga gttcatgcca tcttatttct 6060
ccctttgttc ttagccggaa ggatacacga atcagttttt tttttttaaa aaaaatattt 6120
atctcaattt tctgcaagca tgttcaattt ctaagtggaa atgctattta aaagaccagg 6180
cttattgatt ggtgctatac tttgattttt tttggaattg tagtagaagc atcagtttct 6240
tcatgctgtc ctaccaacct ctcttattat tagcaaagta aagttattaa atttgctaatt 6300
tgttgatatg tcagtatttt gtacgaattg tgaaatagtt aattttcaat aactacacac 6360
catggttgtc ctggtgttgg actggaagca ataagggaat attccatttc tgtccattaa 6420
aaccacaaaa gatgaccctg tgctcatctc taccattgcc atgcacctgt ttgtaggatt 6480
gcctaacca gaagttggtg cttcgagata gccatggcca ccggagtggc accagcgccg 6540
ctcccatatg tcagggtccg tgatgggtggc atcggcttca cgaggagcgt cgactttgct 6600
aagatcttgt cggttcctgc tactctaagg gtgggctcat caagaggcag ggtgcttgtg 6660
gccaagagct caagtaccgg ttctgatacc atggagctcg agccatcttc agaaggaagc 6720
ccacttttag gtataactcg ccggctgttg ttcaccttgc atgtatattc gtgttagttg 6780
ttcttagtgc ttttaactga atgaacattt tttctgtaaa gaatctgaca gcatgtcttt 6840
tgcccttttg ttattcttta gttcccaggc aaaagtattg tgaatctata tatgagacaa 6900
ggaggagaaa aaccgcact gtg atg gtt. ggg aat gtg cca ctt ggc agt gat 6953
                Met Val Gly Asn Val Pro Leu Gly Ser Asp
                  1                5                10

cat ccc att agg att cag act atg acc acc tcg gat acc aag gat gtt 7001
His Pro Ile Arg Ile Gln Thr Met Thr Thr Ser Asp Thr Lys Asp Val
                15                20                25

gct aaa acc gta gag gag gtacactcct atttgaagtt ctatgtttta 7049
Ala Lys Thr Val Glu Glu
                30

gtttttaatt ctatgcttga ataattgaat gctgggcatg cattaatcat gtgttctttt 7109
agatgttcta tgtttcatga ctagtgaaat aacgaagtat agcactggtc cag gtt 7165
                Val

atg agg ata gca gat aaa ggg gct gat ttt gtt aga ata aca gtc cag 7213
Met Arg Ile Ala Asp Lys Gly Ala Asp Phe Val Arg Ile Thr Val Gln
                35                40                45

ggt aga aag gaa gct gat gcc tgc ttt gag att aag aac act ctt gtt 7261
Gly Arg Lys Glu Ala Asp Ala Cys Phe Glu Ile Lys Asn Thr Leu Val
                50                55                60                65

cag aag aa gtaagagtca tcatttttcc agattcagtg agttttcatg 7309
Gln Lys Asn

aatgaattct catcttgctt ttgcatttca acag t tac aac atc ccc cta gtg 7362
                Tyr Asn Ile Pro Leu Val
                  70

```

gct gat att cat ttt gcc ccg aca gtt gct tta aga gtg gct gaa tgc 7410
 Ala Asp Ile His Phe Ala Pro Thr Val Ala Leu Arg Val Ala Glu Cys
 75 80 85 90

ttt gac aaa att cgt gtc aac cca ggg aat ttt g gtgagtgaaa 7454
 Phe Asp Lys Ile Arg Val Asn Pro Gly Asn Phe
 95 100

taatgatgtg tatcatttta gtgtcaatat cttatcaact ctgtgcatat gctgagaact 7514

ctacttgacag ct gat cgc cgt gcc caa ttt gag cag ctt gaa tat act 7562
 Ala Asp Arg Arg Ala Gln Phe Glu Gln Leu Glu Tyr Thr
 105 110

gaa gat gat tat caa aaa gag ctt gag cat atc gag aag gtt cca aat 7610
 Glu Asp Asp Tyr Gln Lys Glu Leu Glu His Ile Glu Lys Val Pro Asn
 115 120 125 130

atc tca ctc ttt agt gtt aat tta gtcagtaaga atgtgcagta tgtttcctta 7664
 Ile Ser Leu Phe Ser Val Asn Leu
 135

cttgcatagc cacttccata tcatttcag gtc ttc tcc ccg ttg gtt gag aaa 7717
 Val Phe Ser Pro Leu Val Glu Lys
 140 145

tgc aag cag tat gga aga gca atg cgt ata gga aca aat cat gga agt 7765
 Cys Lys Gln Tyr Gly Arg Ala Met Arg Ile Gly Thr Asn His Gly Ser
 150 155 160

ctg tct gac cgc ata atg agt tac tat ggt gat tct cca cgc gga atg 7813
 Leu Ser Asp Arg Ile Met Ser Tyr Tyr Gly Asp Ser Pro Arg Gly Met
 165 170 175

gtattatttc ctttctgggg atttcattca aataactttt cgtttcatgg atgtcttcaa 7873

ttaatgatcg ttttgataga tgaatgacat gttctacaaa taatttcag gtt gag tct 7931
 Val Glu Ser
 180

gct ttg gaa ttt gcc agg atc tgt cgg aag ctg gac ttc cat aac ttt 7979
 Ala Leu Glu Phe Ala Arg Ile Cys Arg Lys Leu Asp Phe His Asn Phe
 185 190 195

gtg ttt tca atg aaa gca agt aac cct gtt atc atg gtc caa gca tat 8027
 Val Phe Ser Met Lys Ala Ser Asn Pro Val Ile Met Val Gln Ala Tyr
 200 205 210

cgc ttg ctt gta gca gaa atg tat aac cta ggg tgg gat tat cct ttg 8075
 Arg Leu Leu Val Ala Glu Met Tyr Asn Leu Gly Trp Asp Tyr Pro Leu
 215 220 225

cac ttg gga gtt aca gaa gct gga gag ggt gaa gat ggg agg atg aag 8123
 His Leu Gly Val Thr Glu Ala Gly Glu Gly Glu Asp Gly Arg Met Lys
 230 235 240 245

tct gcc att ggc att gga aca ctt ctg atg gtaattgcat ttttactttg 8173
 Ser Ala Ile Gly Ile Gly Thr Leu Leu Met
 250 255

tgtattatat tgcatatatc atatctttcc atctgcaaag ggtaagcatg ccttatgtct 8233

tccttttggt gtcttacag gat ggc ttg ggc gat aca atc cgt gtc tcc ctc	8285
Asp Gly Leu Gly Asp Thr Ile Arg Val Ser Leu	
260 265	
acg gaa cca cct gaa gaa gag att gat cct tgc cgg aga ttg gca aat	8333
Thr Glu Pro Pro Glu Glu Glu Ile Asp Pro Cys Arg Arg Leu Ala Asn	
270 275 280	
ctt ggc aca cat gcc gca gac ctt caa ata gga gtg gtaacgattt	8379
Leu Gly Thr His Ala Ala Asp Leu Gln Ile Gly Val	
285 290	
attacctttc tctagtttta cacttttctc ttgttttagct gccaatgccca cacattaatt	8439
ttgactatatt ttagtagtgt tttgttctat ttgttctttt aagaatttct atttatatac	8499
attatatggt ctcag gct cct ttt gaa gaa aag cac agg cgc tat ttt gat	8550
Ala Pro Phe Glu Glu Lys His Arg Arg Tyr Phe Asp	
295 300 305	
ttc cag cgt aga agt ggt cag ttg cct tta caa aag gag gttagttcaa	8599
Phe Gln Arg Arg Ser Gly Gln Leu Pro Leu Gln Lys Glu	
310 315	
aataactcct atagtccata gttatcataa aaacaatagt gctagatttc ttattagttg	8659
cacttatgac aggggtgagga agtagactac agaggggtct tgcaccgtga tggctctggt	8719
ttgatgtcag tttccttgga tcagttgaag gtaactcaca tatttggttac ccttttggtgc	8779
aatgtgttga tcttgtgtaa ctttaccaaa atatatttca agacaatagt ctattttgta	8839
atatacaatt ctacaacatg atattttcag tagccatggt ccatgcattc tatgcatagt	8899
tcatagtaca tagtgagaat agcaatagca aaaagaaggc attgattttt ttctatctga	8959
atcaaatacaa ttgatgcatt ttgtaatgat ggaaggctct cttatttttc ag gct cct	9017
Ala Pro	
320	
gag ctc ctt tat agg tct ctt gct gca aag ctt gtg gtt ggc atg cct	9065
Glu Leu Leu Tyr Arg Ser Leu Ala Ala Lys Leu Val Val Gly Met Pro	
325 330 335	
ttc aag gtctgaccc tatagctgta cattctagca aacaactaaa ctttattggt	9121
Phe Lys	
acttcagtct aaactgatgt taatttttct atgaatatca g gat ctg gca act gta	9177
Asp Leu Ala Thr Val	
340	
gat tct att ctt ttg aag gag ctc cca cct gta gaa gat gct caa gct	9225
Asp Ser Ile Leu Leu Lys Glu Leu Pro Pro Val Glu Asp Ala Gln Ala	
345 350 355 360	
gtgagttcct tcaacattat ttgttctttt cacaatacac aagcttatat taacattcta	9285
ttccttttaaa atttttgtgt tgaaatctgt aaaatggtac ag agg ctt gca ctc	9339
Arg Leu Ala Leu	
aaa aga tta gtt gac atc agc atg ggt gtg ttg act ccc tta tca gag	9387
Lys Arg Leu Val Asp Ile Ser Met Gly Val Leu Thr Pro Leu Ser Glu	
365 370 375 380	

caa ctg aca aag cca ctc cca cat gca att gct ctt gtc aat gtg gat 9435
 Gln Leu Thr Lys Pro Leu Pro His Ala Ile Ala Leu Val Asn Val Asp
 385 390 395

gaa ctg tca agc ggt gca cac aaa ctt ttg cca gaa g gtagacattt 9482
 Glu Leu Ser Ser Gly Ala His Lys Leu Leu Pro Glu
 400 405

gaatttgata atgatctttg ttgttttgtg aattgtgttt atgtcatttt ctgtatttta 9542

acattttgct tagtctgttt tattgatgaa tctttttttt atgtag gc act aga 9596
 Gly Thr Arg
 410

ttg gct gtc acc ctt cgt gga gat gaa tca tat gaa cag cta gat ctt 9644
 Leu Ala Val Thr Leu Arg Gly Asp Glu Ser Tyr Glu Gln Leu Asp Leu
 415 420 425

ctt aag ggt gtt gat gat ata aca atg tta ctg cac agt gtt cct tat 9692
 Leu Lys Gly Val Asp Asp Ile Thr Met Leu Leu His Ser Val Pro Tyr
 430 435 440

ggt gaa gag aag act ggc aga gta cac gct gct agg ag gtaagtgaac 9740
 Gly Glu Glu Lys Thr Gly Arg Val His Ala Ala Arg Arg
 445 450 455

acagtaggcc agttaatacc actccctcca ttattaccat ttgttgggat gaaccgatag 9800

tcaattctaa gttacacatt aagcatgaaa aatgaaaatg gatttgactc tgcagaaaac 9860

tgacatacag accaatgttt ccacctgggt ttccattggt ctgtacttct ctttacctaa 9920

aattttatatt tttttaataa tgttttgcag g tta ttt gag tac tta gaa acc 9972
 Leu Phe Glu Tyr Leu Glu Thr
 460

aac ggt ttg aac ttc cct gta atc cat cac ata gaa ttc ccc aaa agc 10020
 Asn Gly Leu Asn Phe Pro Val Ile His His Ile Glu Phe Pro Lys Ser
 465 470 475

gtg aac ag gtactatgaa gtgcttatta agagatgcat tgaccgcca 10068
 Val Asn Arg
 480

tccttaccce ttgaaattac tgtaccttta ttctcttgtg cttatttgag ttaaattata 10128

tgcag a gat gac ctt gtt att ggt gct ggg gca aat gtt ggt gct ctt 10176
 Asp Asp Leu Val Ile Gly Ala Gly Ala Asn Val Gly Ala Leu
 485 490 495

cta gtt gat ggt ctt ggt gat ggt gta ctt ctt gaa gct gct gac cag 10224
 Leu Val Asp Gly Leu Gly Asp Gly Val Leu Leu Glu Ala Ala Asp Gln
 500 505 510

gaa ttt gag ttt ttg agg gac aca tcc ttc aac ttg tta cag ggc tgc 10272
 Glu Phe Glu Phe Leu Arg Asp Thr Ser Phe Asn Leu Leu Gln Gly Cys
 515 520 525

agg atg cgc aac aca aaa acg gtaagctgat gaattcttct ctgttagact 10323
 Arg Met Arg Asn Thr Lys Thr
 530 535

gtagatccca tgaacaacgt caacctttaa ctogtgagat atcatgaaga agtgcaaaat 10383

```

tgcactttta acagtaaag aaccttatag cctaccgaag aggataaata acttttaggca 10443
attctctctt gtgaagcaga acattctttt ggcgatttct gaccgttaat taatgctgca 10503
ggaatatgtc tcttgtcctt cttgtgggag gacactcttt gacctccaag aagtcagtgc 10563
tcagattaga gagaagacct ctcatctgcc aggcgtctct gtaaactctc ttacagacct 10623
tctgcctccc ttgttttcaa tcgcatatta gctagcctga tggctaataca tgtctacatt 10683
tgcctggcag att gct atc atg ggt tgc att gtc aat ggg cca ggg gag 10732
      Ile Ala Ile Met Gly Cys Ile Val Asn Gly Pro Gly Glu
              540                      545

atg gcc gat gct gat ttc gga tac gtt gga ggt gct cct ggg aag atc 10780
Met Ala Asp Ala Asp Phe Gly Tyr Val Gly Gly Ala Pro Gly Lys Ile
      550                      555                      560

gac ctt tat gtt ggc aag gtaacctttt cctatacttg tggaagttga 10828
Asp Leu Tyr Val Gly Lys
565                      570

atcatatcaa atggaataat ggaaatcacg gtatatcgtt gaacatagct gcaagtcaat 10888
atttgtacat gatcatgcaa acacaatcaa cagtagggat gttaactgca tggcatatat 10948
atgctctttg agctgaaaca aaaacttaga gctgccattt tccttccatt aacacaagtt 11008
ctacttgttt tgggtgcag acc gtc gtg caa cgg ggc att gca atg gag ggg 11060
      Thr Val Val Gln Arg Gly Ile Ala Met Glu Gly
              575                      580

gcc act gac gcc ttg att cag tta atc aag gac cat ggc cgt tgg gtg 11108
Ala Thr Asp Ala Leu Ile Gln Leu Ile Lys Asp His Gly Arg Trp Val
      585                      590                      595

gat cct cct gtt gag gag tag gccgtagcat gtagttcata tatgtactcc 11159
Asp Pro Pro Val Glu Glu
      600

tccataaaca atgttgtagc tgaggcacat tgtattgtat ccacggagta cataaataca 11219
cgttctgtac atcagtttag aaataaagta ggaatagggg tggctgcaac tttgtaacac 11279
cctcgtgaag catcggcaaa tccaaattag aagcgtcctg aaatcagtga aaaagaattg 11339
atactgctat tttttgtacc aattgaaaaa aaaaaggaat acatgatatg actaaatcat 11399
gggttacatc ttcgtcaaaa aatgtcacag cttacattat tttcactact tgcaaatacc 11459
agacgatcta ctgggtgcggg aacttgacgg gtgcaggaga cgcgaagccc ttgtggtaga 11519
gaagctcggc catgacgctg tacgcgcgct gagtcagggtg gacgccacgc catcccagct 11579
gatctgctcc atctcgaagt tgtacttccc gccgcagcac gccttgggtca gcgccacgcc 11639
gtcgaacccc gtgtcgcgcg cgcctccag catccgcacg tacgcgccgg agtagtcggc 11699
gtacgcgata gtggcctccg gttatgaccg cctcagctcc cggatcccct gctgcagcag 11759
cacgttgtgc atctgcgcga acaggttgag acccagagg caccggttcc cgtcgtacgc 11819
cgcgcgctcc gtctcgtcca ccgcgcgcag gtagctcggc gcgcaacca gcgggaagtt 11879

```

gcccgggatc accacccgcg tcgcgctcat ctcgagcacc tccctcgccg cgctcaccac 11939
 gcgaccgcac cacctctggt acgagcacca ccgactccac cagcccggtc atcatgcgcc 11999
 cgacgtccgc gcggcgctac gacctcctgt tcgcggcctg ttgcgggca tctctccac 12059
 catcaccagc aagctcggcg cagcttctct tgctcgagaa ttttcagaat atgccaccga 12119
 atatgcaccg ttttcaggat agaccactca attcgacta ctttcataat atggcatttg 12179
 gacgcgatat tttcttcgtt ccgtgacact ctcactcttc caccgtcagc gccagtaatt 12239
 ccgttcgcac accaacagct ctctctgagc gtccagctcc agtgggggag ttgttggtgc 12299
 gcggcgcggt aacaccgatc ctcgcgaggg ccgcccgcgc gagggcggtg gcgcccgtga 12359
 cggcgaaagt tgacaccgta ggagaagtcg gcgcctttgt cgatgtacgg gttgagcagc 12419
 ggcagcccta ggtcgttggc gaggtagtcg atcatgaggt acccgtcgtc ggagcactgc 12479
 cccgtggcgc tgccgatggc cgcgccgtac gtagggaggg gccacggtgt gctccatcaa 12539
 ggcgaggaag ttgccggtgt ccgagatgga gtccccgaag ttgtagatgt ccgtgatgcc 12599
 gtccaccacc gcccccttcg ccgccgatga caacgacgac gacgcggcct tccccggagc 12659
 cggccttgcc tggcaagtgc cgacgaggag cagcgccaag aacgcgacga ggattggatg 12719
 aaccggccta ctcgccatgg cgctcggtgc aagtgcaggt ggggtgcgacg cagcagttgt 12779
 tgtggcatgg cgcgcgcgcg gtgtggaatt cgattggaaa cgatttaagc tgagacatag 12839
 tccaactccg aaacccaaat taaccataca tacagtata caggtgaatc gacgagatga 12899
 tcatgcacta cttaaaaaaa accgtcaaaa cacatTTTTg taggcgggtca aatactctat 12959
 gtacttaaag gcctgcgaaa ataacgcccc aaaagtcgtt tcttagtagt gatgcatacg 13019
 caattgctgc aataacttaa aaagggatgat ttttattgca tcaacgtaac acgtacactg 13079
 cattagtcct cctacattga aagcaciaat taaaccagta tgggtgcaac ttgagacaca 13139
 caaaggatgat cgatcgagaa ggtagctat aaacagcacc ccaaattggca cgaattaata 13199
 atgtagttct ttctgcatgc tgacaaaaat ttcattttct ttttctctcc cctcgtcatt 13259
 aaaaaaagg tttaaagaca gaattacaag ctaattaatc atcagtggat cgagaattaa 13319
 ttaagggatc acaatggctg caccocgcta tttcggagta gctagctcca tgcactcact 13379
 catgcatgca ggcatgcata tacatgtccc ttgccatgtc ctatctaaca atttacacat 13439
 ttcgacaaaa tgctcacggt cgatttgat tgtgtcactg acattaattg gttcatgcat 13499
 ccacgcatgc gttactctca aggaaatatg aaagtatcat ccgtaatcag ggttccaaac 13559
 taaggataga tacctttcan nnnnnnnnnn nnnnnnnnnn nnnnnnnnag gcctgctgca 13619
 gcaagtgcac ttctcctgct catgcttcag agcctgcacg cagaaagacg acacaaaatt 13679
 caaaagttta tatcgcttct gttttggagc ctcggctaaa aaatgaaaat atgaacaacc 13739
 aaaaaaggca acacgtacga gttctaacca agtatataac cattataatg gcaaatgtga 13799

tctatacttt tgtagacgaa gacaattaat gatagtagca gtgaatatgc tagctatata 13859
cttttatcaa ctacttatcc gatcaatatg cttcagcatt acaaactagt tcttatatat 13919
atatttcttc tatcttattt catctctaaa atacaaagtt tatagtgtaa agagatcccc 13979
agggatgaat atatcttcta acacacctcg tagttaattt gttccaaaca atactagcat 14039
gcatataatt tgtagttatt tgtagcaaag cacggctatt tcgctaacaa atctaaatag 14099
aaaatatgtt atctctcagc cttgagaggt gtattaatta ccagcccata catcacttga 14159
gagggaaaag atttaaataa gacaaattga ttagaacaaa agggaaatgat agacaatgtc 14219
ggtttttttt egtttcttcc tttccttcgc ataggctcgt ctagctgggt gcgttatgta 14279
acaaaacctc ttttcttttt aatatattga tgggcgcgcc ttttgccgat tcacgaaaaa 14339
aatgtaaat gtgaattttc aatcttatcc cctacttgcg ggattagtcc ttgtgaagaa 14399
atcctcaaat atgcgtacct gcagctggct ctgcagaccc ttgatgtgct caactgcaag 14459
gtccaacatg tccgctgtgc ttgtttgctg ttgcaacacg aacataatta attactcaat 14519
tggttgcat attcatgcgc aaaaaatgtt accgctaatt aatattagct agaactagat 14579
gagagaacgt acgaccctt tcacttatat acaataatca tgaatttggt gagaaagcat 14639
gtttggtagt gtgttggagt tgtggctgct atgcacaaa gctctaattc cagtgcctat 14699
agaatttaac tacacaaaca tggatagct ttttctagaa attctattag gttatgattt 14759
tgcgcttgggt gtccatgaat ttgttgagca tgtgttaagg gacacttcac agtgcacact 14819
catgggtgaa tgcgtgtgca tttgccatgt ctattattaa ggcgagaaac atgaatctgt 14879
gtgctaattg cacaagaaat gtggaaagtt tttttttaa agaaaatact tagctaggga 14939
tgttcctttc ttcctcaaatt atcatgtaaa tataggtagt aacattatgc aaagttcaaa 14999
tcgtaatggc caccttgtcc atgttgggca ccagctcctg cagcttcctg agcttctcgc 15059
taattctcgt cctccgttcc tacggacgcg catcgatcac accgacgtac atgctcatgt 15119
gtcaagatct gaagagaaag caaaagcaaa tatagaggcg ttttgatcat gatattgcgt 15179
acgtaccctc tccgcgatgc tcctgggggtg cgtcgcgcag ccgcgcttgg cccgcacttt 15239
gaacggcacc tggatcatgt gcagctgcag gtacctgtcc atgccggcca tctccagcgc 15299
cgacgtgctc gccatgccgc cgaactgcc ccatttccaa cagcccaag aaatcagaac 15359
acatcgcgat atatatatat atatatatat atatatatat atatatatat atatatatat 15419
cacaacaca gcaaagctag ctactacttc ttcctctgtt ttacattaat tattataagt 15479
tgttttgagt tttgaataga ttcatacatg tataaatgta tgtgtttcat acatgtgtcc 15539
aaattcttat gaatgttagt aaatataaac aaggatgaa gagatcaaga agcctttagt 15599
tgtacaatga ttcaatgaag gtagccctag caatcaaatt tgccgagcaa tctttacctg 15659
ggactcgtac ccgccgaggg tggagatgat gtccctggac tcctcccacg gcccgacgat 15719

ggagaacccg ccgccgccgc tgctgccgcc ggcggagaag gtgcggggca cggaggcctc 15779
 ggcgccggcg cggtcgggga aggcgccgtc ctccgcgatg tgcgagaggt gcggcggccc 15839
 ggccgtgaag ctacagctggg acttcacatctt cctcccgccg ctgctgctgc cgctgccgct 15899
 gccggccatg gaaggggtggt ggtgggcttc ggctccgctg cctccccgc cttttgagcc 15959
 tggaaagcct gcttagttta ttgccaagta gcaagcacgg aaattaacta atgacgcgta 16019
 attagttaaa ttaactgtgt gtgtgagaga aagagctact gttacccaaa cgctagttga 16079
 aaactgccaa gtgtgacaag taaacaatag tttacggtat tagcataccg ttagagctag 16139
 ctctataggt acacgtgttg agcaataagt ttaacctaga tgtgatggga tgttcaaact 16199
 tgcttctcca aggttgaatg gagtagtgtg tatttgattc tacaatattt ttctgtagta 16259
 ggtgcacgta attaaggtta ggtttgattc tcatgttcaa atgtgtgttt aactgcaggt 16319
 gtaatgttat atatgcatag tggttctata aatattttca taattaaaca ctaccaaatt 16379
 tctatttgaa atccatgtac aaattaaact tgactaatca ccggttatta tagttaaaca 16439
 taacttaaac cacaacaatt accattcatc aacactatgc actactaact aattaaaaaa 16499
 aattacaagc tagcactacg aaattaaaag tggcccggcc gagttgcccc agcacaaaat 16559
 agcacgatag atacaggata tacttcctcc gtttctaaat attttacacc gttaactttt 16619
 tagcacatgt ttgaccattc atcttattca aaaatttttg tgaaatatat aaaactatat 16679
 gtatacataa aagtatatat aacaatgaat caaatgatag gaaaaaata atacttattt 16739
 aaaatttttg aataagacga acggtcaaac atgttttaaa aagtcaacgg catcgaatat 16799
 ttagaaacgg agggagtata tgagaggaat attctcgtga ctagaaccat atgttccaga 16859
 aagttgtact ccatccattt taaaatgtaa ggtctatttt gagtgggtcac aagtattaag 16919
 aatatgaaac ttacagaaag atgagttcaa acgaccacct taattagaaa gagtagtaga 16979
 tcgttagtga gacgaatatt atatatatga aagagacaaa aacaattaaa attagtgttt 17039
 gcatttgctg tcatctttac tagctattac tagttactta taagcacatc gtcaaacatg 17099
 tacttacgtg ttgcaactta atttctactc cctccaattc agtattgggtc gttttggatg 17159
 aaaataatat caaagttagc aatccggccg taaccatttt ttcaaacctt gtatgccccaa 17219
 tagttacatc gctattcaaa tcaaaggttt caaattttgg attactattg ggtcccaata 17279
 gaagcccaaa aagtatttga attttttaac ttaggccccg tttagttccc taaatttttt 17339
 ttcaaaaaac atcacatcga atttgtgaac acatgcatga agcattaaat atagataaga 17399
 gataatccct catatgccac taaaaattga tctgatccct tatatgccac taaaaattgg 17459
 ctctccctt atatgccatt ggtctaaatt tgcgtaccct ctcatgtcac taccgtcagt 17519
 tgaccgtgtg ttgaccgtta actctcaagt aaaaaagaca tattgccctc tctgagttgt 17579
 taggcatgcc ctatactcag aagggtaaat acgtcttttt tccttaagaa ttaacgggtca 17639

acacatgtca actgacggga gtggcgtgag agagtgtgca aatttggatc aatggcatat 17699
 aaggaagaa ctaattgtca atggcatata agggattaga ccaactttcg gtggtatata 17759
 agggattctc tctataaata aatgaaaaat ctaattgcac agttagggag gaaatcgcg 17819
 gacgaatctt ttgagcctaa ttaatccatg attagccata agtgctacag taaccacat 17879
 gtgctaata ga cggattaatt aggcttaaaa gattcgtctc gcagtttcca tgcaagttat 17939
 gaaattatct ttttcattcg tatctgaaaa acccttccga catccgggtca aacatccgat 17999
 atgacacca aatgtttct tttcgcaaac taaacaggcc cttagcaaaa tggttggtta 18059
 tcaactttta aaatatgttg acagtgtctg tgacgacttc atgacgggtcc tctttaaagg 18119
 tgcttatata gtgatagggt gtgctgtgat gttcagagcg ttgagtatgc atgtgtatat 18179
 atgcatgttt gtgtctgtac tgtgttaaaa aagaaaatcc caagatctag cctaaaattt 18239
 tcattaaaaa cattgaaatt ttggcccccac gattttttta ttccacaatg taaatttcta 18299
 gtcaaattgc tgcaatgac gcgaaaatta ttttctgacc agtgaactga catgcacaca 18359
 ttacactata tttattttat atttattttg aacgtaccta cgactacttc caggggatcg 18419
 atcttattct cctcaaatta ataagaacaa gtactctctc catttcaaaa tacaacaacc 18479
 taagaatatg gataattttc ttcatagaat cggatgggtt cttcgggttt tttgtactac 18539
 gatgcgaaca gatggtatat tgaagcctac cggacacgct agcacgtgca tgccgcgtgc 18599
 cggcccgtgc atatgagcaa gcctcgacg ctgacataga cgcagccaag agagaaagca 18659
 aacgccaat caagaagccg agcaatcacg catgccatct caacgcaccg taggtcacta 18719
 tcttttagcga ggcaagaccg tgacgtcacc gtcaggccat cagcagagga gctgaacctg 18779
 gacaaaccgg ggggcccacc ccgcaggcca agttgcggcg acacacacgt ggtccccgcc 18839
 ttacattaag gcaagtggcg ccctaattaa tccattgatc aaaaattaat taatccacaa 18899
 attaatcaaa tgccctcatc tttttctttt tgccctgggt agggttcgag gcactaagat 18959
 ccactggtta ttttaattgtg cttgctgtct tgataactaat taattgatca tatatgcgca 19019
 agttggtcta tctagagcag aatctagagt gcaactgggt gccgcattga aagaaatgct 19079
 gctacatggg ctccactgaa agacatttga ctctttttaa ctttactcga ggctattcct 19139
 acctcgatca aagtataatt actaaattta gtactgggtg agtacttata tgtggatttc 19199
 gacatttcta ctggtactat ttttatcctt accaattgtt gtatacaggt tgctcgggtca 19259
 aaaggccatt ttagatgttg gtatatatgt agtgtgaaaa ttaattataa cataactcta 19319
 tgttcatatt gatctgcatt tcaaaaagat attgacacac ttattcctaa tttttgaata 19379
 aatgatattt tgaagttttc attaaagggt tattatctct gtatgctcta aaacgttgaa 19439
 tatttgtgac gcagaattaa ttaatactc atggaataaa taatgatggg gcataatttt 19499
 gcaatgattt tcatcaaag aggtgcatat aggtatcctt tatatgaaat gagaatactt 19559

gccaaaaaca tttttaaaaa gagcttggtt tagctagcta gggtggtgaa tgggtgatact 19619
 aattaatcaa atgtacatat ttgtgcaa at cctggaagat gaatgcatgg ttttctagtc 19679
 ttattatgaa caaattaaat tagaaaaaaa aacatctatc tctttgctct ctccactata 19739
 gcttcaaatt gtttttttttc cccatgtcta ctattgtagt gaagaatgga ttgtcatgcg 19799
 caatgacttt gcaactgaaa ataatggatc aaatgagaga gagggacacc aggtgcaagt 19859
 ggcaaaaaaa ctaagccatt tatagcaagt tgcaatagaa aataagacaa tctagagaca 19919
 ctcgattata aaaagcgtac gtaaaaagaa taaaagcgggt gtattcaaaa ccctagaccc 19979
 cacatttcac tatcgatgat accctacttg agaaaacccg ctcctgtgtg agcccatagt 20039
 tttccatcgt ccttcttaca cgccgagcca aatttgtgca ctctcgttaa taacatatgc 20099
 cttaaaaact tgaactcata ttacattatc acgaaaacaa ttaagccgca taatctcatg 20159
 gatataacat ctcattggtg atccttaatt aacagcttat atatatatat atatatatat 20219
 atatatatat atatatatat atatatatat atatatatat tgaccctaac tgtggcaaac 20279
 atgcattatt atcacacaaa agttactaac cacatatagg agcctatggc taatggctct 20339
 gagtagaaaa atgggcacag aggatctcca tgatactatt tatggcaact cacgtagcaa 20399
 aaagccgcag actaacacat ccatggatat ccacaacgca tactgatagt agtctgatat 20459
 acacactagc tcctcccatg acggccttag cgaaaaccac tttttaaccc aaaaaaaaaa 20519
 ccagttagga ccggtgaaaa gtcgcacgcg atgatcgatt cacgcgcgcg ccgcagaagc 20579
 aacttgcaaa agggatcgag cttagctaga tagcgcgagc tcatcagcat ttcgtcgtcg 20639
 ccgagcgcg tagtggcttt ggcagttagt agtgatggga gttgcataga agttaagaac 20699
 caggtagaca gagatcgatc gattgatcaa acccgtttgg tttcggataa gtatgggaag 20759
 aatctgaaac agtgtggagg aaacactgag agagaaagaa caccattaac aataatatcg 20819
 atggaattcg ttttttttgg tggttgttgc tagaagccta gaacagcaat tcatgtgatc 20879
 gatcgatact tcgatcgtgt gcgtgtgtga cgagaaagag atggggcatg tgaaggcaaa 20939
 gacgaggttg acatttgac agctagccgt tctctcctga cagaattaag ctagaaattg 20999
 aagatccgtg actctgagta gtcctaacca attagctata cgcctataca cgatgggcta 21059
 gctatgcacg cacgcgacgc caaattgaac acggatgaac aaataaaatc gaacaatggg 21119
 ttggctagcg caatcgatcg atcgatctta ccgttgctgg ccatgaggtt ggagaagaat 21179
 ccggccggcg agctgctgtg ccgcgccagc aagtcacgc tcccgctctg cagctgatgc 21239
 ccatgccctc cgccccctcc gccgccgcgc ccgcgcgcgc cgtgcggggc cagcgagatg 21299
 tccccccac cgaaccgcag cccgcgcgc tccgcctccc ttggctgcgg cgtcgtcgac 21359
 gacgacgacg gctccacccc tctcctcct cctccccca ccggcagaaa cctcctcatc 21419
 ataccctcc ttggatcgat cgatcaactc cccccccgc gaccgagacg cggcctctcg 21479

tcgatcgatc tgcagctcgc gcaggcgcag gtaggcaggc ggcgcggtgt gtgggtggaa 21539
 atttcggcgt gaaaattaac aaaacgacgg gggcggccta tactatagct agtagaggag 21599
 agaagagggg aaggggaagg aagggggagg tgaggtggtg gaggtggtgg ggctaggcgc 21659
 aagtgggagg agaggtggtt gggattttta agggaagcga ggccccgtga ttggttctcg 21719
 gggcgtgtgg cgccgtgggg accagcggac cggccggggc cgggcaagtg gatgtctcgc 21779
 gcggagtga gtgggtctct gcactgcgca gcagcagcag tagcaagccg taggtggcgt 21839
 cgcgcgcccc gccccggaac cggcaggcat ctctctcggc ttttcgctgc atctttggtg 21899
 ctagattttt gtgttgata tatgatgctg atcgaggaaa gggaaggaag aagaaaaaaa 21959
 aaaggatttt ttggtgtgg cttagatttt tggatgcttt ctttcctctg ctgcggactg 22019
 cggggactag aggatgaact cgataatcaa tgggtggtggc ggcaaagtgt tatacttcct 22079
 cagtctttta tatttacctt ttgtgatatg gaggaacaa gctggtttgt ggtgttgtgc 22139
 actcgtagga ggggaggtac gtagttaacg gcaaagatcg atcatgcaag ttggttgggt 22199
 caatttgggt gtcgagctga cctatgttcg cccatcctct cgatactttt ctcatctaga 22259
 ctttttctac gacgctaaca gactgattat cacagtcatt ggatagatcg acatggtcac 22319
 ttgaaattgt tcgatataac tggtttaagt tcaaacaat ccaagctaaa ttttattttg 22379
 cggaaaaaat gtttgaattt cacttgtttt caaccgttat tgctgttagc gaccttgccg 22439
 ttagggagcg gttttttaac ctccggcacat ccgtaaactc tattgcaggg gagtcagtgt 22499
 tatgtctaac agtagtataa ttttatcaca atgatttgtc tctttacgag ttgtattata 22559
 aactcacggt gttccccgca aaaaataaaa aataaactca cggtatgtgt aaatggaatt 22619
 aggtcaaaat ttaggaatga aatgaataat caattgggtg tgaatgggtc aatgcactaa 22679
 accatatgtt ttgctcacta gatatgacaa ggaaaaaccg aaccatcaat aacactggaa 22739
 accatgtttt tgtggtgacg cttagttaac tcatacatca attataatct tttctctatc 22799
 caattccact ttggtctatt ttgtctattt gaaatcatgt ttcagctatc ttctaagtaa 22859
 agcaaacttg aaaacctagt acatctaaac ctagctccac tagtgtggtc caaaagcagt 22919
 ggagtaatat cataagagga agacaacaaa aaataggata gagatagtct tagcttgtgc 22979
 cgcagtaatt cattcgatag attattagta attcattcga taagatatga taatgatgaa 23039
 atgattcggc cgacgggtgg ttgtgaagtc ggagccatga tatgtggcat cgaaagcatt 23099
 agtcaaacgg acttcggttt tggtcaggta aagttgtgtt ccttggttct taattcttat 23159
 caaaattgga gtcgcctgat catgtgtgcg gtggtgtgat gacgaatgac ggcgagtttt 23219
 taccaatgta cagtgaactg cgttttgttt taaggctgtt agttttgttg tcgtgggttat 23279
 gctttgctag ctagttaggg tgattcctat tttttgtcag gtcttatgaa agttaaaaat 23339
 attttttaaa atatgagttt ttttattgtg aataatgagg aacaaatgaa gttttgggag 23399

gatacatggc tagaaaacat ggctttttaa gataaatatc catctttata ttatatagtt 23459
 cgaaggaaaa atttatctat tgctaatagcc atgggatctg ttccgcttaa tgtttctttt 23519
 agaagagttt tagttggtca gaatcttgta tattggcatg aattgcgtgc ttctattgta 23579
 catattcagt tgaatcaatc tactgactat tttagatgga attatcatca aaatgggttta 23639
 ttttctgtaa ggtcaatgta tctagcctta agccttaatt aataatgggtt acattgagag 23699
 aaataagatt atttagaaac ttaagatgcc gcttaaaaatt aagattttta tgtgggtactt 23759
 gcttaaaggg gttatgttaa caaagacaat ttggcaaaac ggaattggaa tggcagctta 23819
 agatgttggt tatgtatgaa aaatgagact attcaatatc tttttataga ttgtcatttt 23879
 gcaaaatttg tttggggagc gtttcagtac tcttttggtt tataccttcc tactttcata 23939
 cattgtatgt ttgatgggtg gctttggggg tgaacaagaa aaggagcaaa ctaattcttg 23999
 tagaagcttg tactatatgt taggctctgt gattgagtag gaatgatatg atttttgaca 24059
 aatcactatc tatttcattt atgcaggcat tcttcagagc aacatatggg ctccggtttt 24119
 gggcacaacc gtaaaagtgt gatgaagatg gagagctttt gaaagttata tgtcgtaagc 24179
 ttgagacgac ggttatgcaa ctttttgcca actatggatg gagattcaca aatagactta 24239
 aataattgtg tgctccttat attggtctgg tcattttttt tatgtttagg tgtgtgttta 24299
 attctatttg aactacactc tttgttaagt gctagattgt aataattggc tgtagctctg 24359
 ttgagcaaag gccgagatgt tatctattcc attattaaaa aaaagctagt tagtgtattt 24419
 attgttgat ggcggtttta gcccgattgt tctaaatcaa ctgaatatta atttgctctt 24479
 ttttagagaa acaccagag gtcttccggc tgggttagat gaccttggtc cttatccctt 24539
 ctaattattt gatattaggt acttcactaa tattcgatc ttttttaa atatttgctc 24599
 tcttttttaa ctattcatct tttctttaat atagcactaa attaaccgtg atctttcaaa 24659
 agaaaaggcg aaagggtgtga atatgcatga aagatcgagt ggacaccccc caaaaaaaaa 24719
 aaaccctagt tgttggtcacg tgactctcaa agtccatttg aggacttact aactgtttga 24779
 aattaatgga taaggctcca gctaagtagg cgggaaaaga tcaaacgtgt tcagtggatt 24839
 tataccaaat gtggtccgtg cgacatgttg gtccataaaa gggcatatga aagtttcctt 24899
 tcagctaatt aaagccagtg tcgaatactt atacagtata gttttcgaaa taagttttac 24959
 ttctacaatg taatccattc acggatgaaa aagctgtgcg cccaacagct atagctatac 25019
 aactatatct atttgttaat taagaggttc atatcttggg cacacaaagg ttctgtttga 25079
 atcttctgaa gataaatatg aagatcaaat gttttacgta aaacgagggtg gtaataactt 25139
 ttgattaatt agattttaat tattacaaac ttaaaaaaaaa gattaatctg atattttata 25199
 acaactttca tatagaaaat tttcacacga aacgcaccgt ttaacagttt gaaaagcgtg 25259
 ccacgaaaat ctagaactta atctgccctt tgttgggttc tcgaacagga ccaaacttca 25319

tgtccatact ccgtactgta cataccaact atactaaata tcgctaaaac gttttaaaaa 25379
 tattatacat atactttcaa tactattata cgtatgcgta aagttttatc ctcaaattca 25439
 ttataatttca tacttaaaaa aaatttctaag agctttatga atataagtct tagatttttt 25499
 tctccatata tatatatgat aaattttaaag atgggacttc acgcgtatat ataaatacta 25559
 tttaaagtac atgtacattt ttctaaaaaa ataatatgtg ttagtttgta tacattgtgt 25619
 gtatacgtga aggctcacgt agacattttg cactctcaat tatttatact agactaataa 25679
 ccacctaaat attgttctta gcggttttga cttgagctta cctacgagat gccaacgtgt 25739
 cagtccagtc agcaaaaaag ttttaaaaaa actccgtggg ccacttgtc atacttctcc 25799
 ctcaatctaa cgcctccccg gtcaccctac tctctcttcc tcgtgcgcac gctcgtgcgg 25859
 ccaacggcgg agtgggtgcgg tgttggtgtt ggtgggtgcgt ccgcgtcggg cacgacggcg 25919
 gtgttgccgg agagatggag ctacgcaaga ggcgggcgtc gacaagtatg gaccccagga 25979
 ggtggaggct acgcgcgagc gcgtcggccg cgccttttgc cttcaacggc gacgacagca 26039
 ttggccgctc gttctcggcc tcgcctctct gaccgcaggt cggcctcagc ctagcagtag 26099
 tcggccacgc acgttggcct cattttcgtc accgtgttct tggggctggc atgcaggcgg 26159
 gagaaggagg gatagcggca tggactgcgt gtgcgcctgt gcagtgacct ggggtggatac 26219
 tatgtcaagt tggagctctg caagtcggcg ctctgcgacg gcgacggcaa cagggacaca 26279
 tcgtcgttgt caccgtgctg cgcggcacga gaaagatggg ggcactgacg gtgaggcttg 26339
 cgatgacaaa tgtgatgggt acaagggccc aacgtcgagg tcgttgccct ggaatagaat 26399
 ccgatcggca gctctagcgg tggcaacggc tcagtgctgg cgtagagta cgagcgcggc 26459
 cgggtggccac agggcgggtac cgcggcgcgc gggagggtgt ccaggcggcg cagtgtccc 26519
 cccgtgagaa cgagcgggtt agcatcatgg ccatcgtcgc gcaccgtcac cgcctgggc 26579
 ttctctagcc gaccacgctc cacctcaagc aaccggggga gtagtctttg ctgccaccgc 26639
 ggccttctcc tcactgctcg gttcaaggat gagagagaga tcgagggtgga aggaggtaga 26699
 agagatgagg tgagaatata tggatcactg acaagtgggc ctgttatattt ttgccgcgtt 26759
 agaaatgcca agtcagctaa cctagcctaa aaccgtccaa aatagtgcc cggtatcgt 26819
 ctggttttta gagttttgag gtattgaata caatatatgt tattatagtt tagagggtaa 26879
 attgtactac cgtaccataa tagttcgggg gtaaattgta cttcctctgt actcataatg 26939
 gaagtcgttt aggacaatat ttaagtcaaa cattgggaat ataaatcatg aataactctc 26999
 aagttgttga gtttgaaaat gtaaaaatta tatgaataga tttttcttga aaaatatattt 27059
 cataaaagta tacatatatc actttttaat atatatattt atagaaacaa gaagtcaaaa 27119
 ttatgttttg gagaccgtgt cgctgtccaa aacgagtacg gaggaatac tttttactcg 27179
 tagtttacaa tatcgatctg ttaactgttt ataagagtat ttggatccat gcagtattgt 27239

agtagtagta gcagtacatt tgagaatatt agagtacgaa ttaggtggtg tttggataca 27299
 gagacttaac tttagtcttt gtatttagac actaathtag aatattaaat atagactact 27359
 tacaaaacta attatataaa tgaaagctaa tttgcgagac aaatTTTTTTT agcctaatta 27419
 atctataatt agagaatggt tattgtagca tcatataggc taattatgga ttaattaggc 27479
 tcaaaagatt tgtctcgtga attagtctaa gattatgaat gagTTTTtatt aatagtctac 27539
 gtttaatatatt tataattaat tttcaaacad ctgatgtaat agggacttaa aagactTTTta 27599
 actaccatTTT aaacagggtc actccaatgg taggtgaaat tcaacagctg ggaaatgcac 27659
 tagtgcggtg tgtcagtaaa tttcgtacta gtaccacgag acagctagac agacacgtca 27719
 ggtcacgacg cagcactgca gcagggtgt agcctgtacg ggaggcgtag gcgcaacatc 27779
 tcgaaaatTTT tgttccgtag ctaaagcccc cccaaagcca gccgcggttt tcatggattg 27839
 cacaggcgtt cctctccgcc ggattccgga aagaaaaaag aaaaaacaag atgtccgttc 27899
 cctgggtggt gcatccgttt tctgacaggt gcatgcacct ctcgctcgct accgcggtag 27959
 cgccacacg aaccacgttg gctttcggcc aacttgcccc attctTTaat cccctcacga 28019
 cgtacgtcgc tgtccaataa aaagTTTTta caccaactat agtaaccagc ttaattTTta 28079
 taaaaccaa gaaattctt aattacttag agcatctcca acagggtcct caaacaagt 28139
 ccctaaataa gTTTTgagag ttgatgcaaa aaaatatagg tccagcagat tccctactag 28199
 agccccaat ctagggaggc ccctagatca ctctccaag ccccagtcg ggggggctca 28259
 accccacagc ccccatcctc tttttttttg gcgggggaaa tttctgagcg cgcgccatcg 28319
 tcatcctccc tcccgcgtca tcgccatcct cccacaacct cccagcgagc aagccgccag 28379
 gtcgttttgc tccggtcggc gaccaccga catccctcgg gaagaaactt cggcgagatc 28439
 ccatggtgtg ccgcctcaag ccatcacaat catcggtttt actccgtcac tcaactgtcg 28499
 cgacctcctt gcatcttcgg ccaacgcacg tcgtcaccga ttgatcatgc ccagggtgagt 28559
 ctcgacgtcc tccccgaact agctgccac ccgctgtcca caaggcctag cgccgaccat 28619
 cgacagctat ctatcaccgg ccgcacacat gctgcagtaa aaaaattcag agatgatggt 28679
 gattaaaaca aatcacaata gtaaagttca gtttcgtatg tctgagtctc cttgtttgat 28739
 tttgatcttt atgggcttac taggcgtcta ggccatcta aatcatcgc acagcaaaac 28799
 gtacattgtc atcattcatg ttttatatgc agtgtcttgt tctatgtcag agagctaatc 28859
 ttgcagagca tatataatat ttaagaaata aatttgtgtt tgcactgagt ccttagtact 28919
 gcgcaaccaa tatatatgct aaataaatac atattgcaaa cagtataacc tgatgtacat 28979
 tgcaatcact tgttgatggt tctgagatag attggaaagg ttgtcaattt atatatttat 29039
 tgcagtgact agattgcaat gacaagtgga ggtgattcct ttgtgcgcat gatgtccgag 29099
 gacactgatg tcgaagtgtt aatgccaaat gaagaccttc gtacttcaac aaatgggtgca 29159

aaaggaagtg ccaaaagatc aagcaactat actcataagg aggacattca attgtgcatt 29219
 tcatggcaga gcattagctc agatcctatt attggcaatg agcaaccagg gaaggcatat 29279
 tggcagagga tcgcagagca ctaccatgct aaccgtgatt ttgagtctga taggaatgca 29339
 aactctcttg agcaccattg gggtaacatt cagaaggaag taagcaagtt tcaaggttgc 29399
 tacaatcaaa ttgagcgtcg tcatccaagt ggcataccac atcaagagct tgtaagttaa 29459
 attgtttatt tattattatt aataacaatc ttgtatgtat gtgaattaaa acttaaatta 29519
 tgttgcaggt tcttgaagct gaggcattat actcgtccac tgcaccaaag aatagggcat 29579
 ttcagtttaa tcattgttgg ctcaagttga ggaattctcc aaagtttcaa acactagaat 29639
 cccacaagag gccacggtct aggaagtctt cgaccccaat tgagagagct ggtgaagaag 29699
 atgaaggaga tgatgctagc aagagtacag ctcttgattt atctcagccg agtgctaaaa 29759
 agagaccaat aggtaggaag caagcaaagg aaaagttgaa gaatggagga caagatggac 29819
 catacaaaga ggcgatgaaa gatttgcttg acgctaaaga gaaagaagcg aaattgaaag 29879
 aagagagatg gaaggaaact aaggagattc aagagcgcaa gctcttattt gctgagcgta 29939
 agttagtggt ggatcaagaa cagaagatta tgttttgtga tgtttccacc ttggaaccgg 29999
 atgtgagaac gtatgtgttg gctatgaggg cacagattgc agcttcaaag gtggctgccc 30059
 tcaatggtgg atttgatggg agtagtggct ttggaggtga gtttggtggc ggtaatggag 30119
 aagtttgagc acttcgatgg aataagttgg attctattgg atgatccatg tgcctttac 30179
 tagtaggata tgccattatc acgattgggc tttggagtc ttttttgtta attatttcca 30239
 caataatttt agtgtcactt gctagtagga catatattac tttcagattt gttatttata 30299
 atcgaatcat tcatggttgt aggatgtatt atttttaaat tatataatgc atcattgggt 30359
 tcacatagtg tattttttat gagcaatttt cattttcatt ggtgaattac gaatcttggt 30419
 tgcactcttg tgcgtatat ggcactgtac ccataccata tttacatgtt taaaaatttt 30479
 aattttgtat tcgaattgta gtgtttgaaa ttgtgaattt aagtatgggt aaattatgtg 30539
 agttagaaat aattgtgttc gaatttttgt ggtgttaaac atactgtata tggattgtat 30599
 tttaaaatac aagataaaca tgagtaggga ctaagaaata ggggctactg ctggagttgg 30659
 aggcattttt tagtccttga gaaatggggg cagccctcat ttaactttta gacgcttcaa 30719
 aataaggtct attgctggag atgctcttag gtcccatcg tttccttcaa tcagcattag 30779
 ccgctaccaa aatttgaaat tttaaagttt ttcacgaag tttattttcc agcattggta 30839
 ttttaagtcgc taaaaacaca tatatgaaag tcttatctgt aaattattat tattttgcta 30899
 atacgccgaa tggcgtatta tatgtatttg gccaaaggat gggggcctta aaccttagcc 30959
 ttagtcgtgc cctacaaaag acacacgcct cgtcagggca agggctactg agcgtggagg 31019
 catggttcgc aagccatggg cggcgaggcc atgctctagc aatgcggtgc agtccacctc 31079

ctctccgagc gcggagctcc aacgggtgat ggccaatgaa agaaggagac cgacttgccg 31139
 ttggtttag tagcatgtaaatt tcttgcaactt tcttaataaa ttctcgctag tgttcgctag 31199
 ctcgacccaaa aaaaagagag gctaattgatg gggttaggaa gtgaaaacaa gcgcagtggg 31259
 ggagaagaag atcgagaggc ctatattgtat gatgctttgt cgatgtagat ttagtcccat 31319
 gctcatctca tccctcagcc acaacaatcc catcattgta gagctcatca gcttgctcta 31379
 ccatctctcc ttgtttatgg gccactccca acctgctacc catcgctga tctatgaatc 31439
 tagctgtcaa tgacctcatt ggccctagtc ttgagatcac ccagtggatc ctttggcaaa 31499
 gtggatccgc ctttgttttg ctttggagaa agaaaacgat gacttagcta aagatctcgt 31559
 cgggtcaaaaa gagagatgcc ttatgatat gctgaaaaat agaggagagg cagtgtcagc 31619
 tggagagctc tttatccaca cccgtgggga tcgagcttat tggcgtagag ggagagacat 31679
 tgagggagag agagtgcagg gggatTTTTT tgtaatttct agatttggtg gtgttttagtg 31739
 caatactttg aacttatttg taaattaagt aaaacatgat tgtaatagaa aatatcataa 31799
 actgacatag aaaaacaaag ataacaattg aagccactag cgctatggag aaaatgtgtg 31859
 acctcgggtc acatataacg gctatgtgtt attaccatgt cacttctaaa actaccatat 31919
 aaccatatac gtttttctcc tacttatcaa aaatataatt aacaaatttt tttaccgggt 31979
 tagtttaca gaaaaaagt tgactgcatt gttgataccc taccatcctt gtacgaaggc 32039
 aggcgctaca caacaccgct gccgctgccg tcgcccgcgt aagctaaggc tgtcacgccg 32099
 gcgaccggcc acggccgacg tggaaagcga cctaattctgt aaagtgtaaa cccaccctat 32159
 agaaaaaccc ggttggtggg acgagaatca ccgaatcagc gtcgacgacg acggccgacg 32219
 actccagcag cgggggtcac gagactcgga gccgagagag agaaagagga ccacgcgcgc 32279
 attcactcaa ctgcataaaa aaacccccgc gcggcggctg cgcagtcacg tctacgctcg 32339
 cgggatcgct cgatgaaatc aaccaaattc ttaaacaaac cgaaccaacc aaccaaccgt 32399
 cgcgcgtgtg cgcgcgaggc gctcgattag cggagacgca aacccatgta acaccgtgcg 32459
 gaaaaactta aagaaatccg cgtcgctcgc gccgtcgcgc gcgcgggggg cgcgtagtac 32519
 ctccacacac gattctgcac ttgtactacc acgcgaacct gatgcgggtt accggtcac 32579
 gattggctgc gaggcttgct gttactggtg gtggtagact ggtagtacgt tgcttgtagt 32639
 acctcactca tgtctggaga ttactacact tcgatctttt cctctgtttt gttaattgag 32699
 atttggaggt gttactgttc gctgtgtggt taagtatat ggtgtataac tacaagttgg 32759
 tactctcaa gggaaaaaaa ggtactgcaa attggctaatt ctatgattct attctgcaca 32819
 tgcatataga taagcactat aataaggac tgaggatcgt gaaaagtggc attaattata 32879
 acaggacat gtacgactat accactggca gggatttcac ggaatcaact ataggagtag 32939
 gttagttggc acttggcaag gttgattgat tcactaacgt ggggaaaaga acacacgaga 32999

```

tcaaaggctg tcgtgggctt aaaataaaaag ggcccatctg ggatcagctc ttttaagccc 33059
acatcactag ccaggaggct aggagtccag tattgcctcg tactgggccc tcctctgaaa 33119
tttggaggcc ctgtctaaaa ttctaataca gccttaaact taagtgacaa aataaaaaga 33179
ggtagactat ataacagcat accattacaa cggaatagct gtcgttagca cgatactcta 33239
tatgcatcag atatgggtacc aggtactata ccgacgttag catgatccga taggtatagg 33299
atctgggtgta cctagatatt atgctaacat aatcatgaca tcagctattc cattggaatg 33359
atataccggg ggtatcttcg gtaaattgtg agcatgctag gaatttaagt aaagggcctt 33419
aggggttaaaa tcacacgttc ttagtcactg cactatcaag tgcatttcaa ccctaatagcc 33479
cttttatgat ctatatctgc cctcctagcc tattttggac gaggctccct cgtcctagaa 33539
gtaaatcatc gtatccataa tccaaccgat tagtagagaa aaaacatact tttcgaacgc 33599
aacagttctt gtcattctgt gctctcaaat gttcattttc cccttactta aaggacatgg 33659
aaaacagaac agaccc 33675

```

```

<210>      3
<211>     1119
<212>      DNA
<213>     Escherichia coli

```

```

<220>
<221>      CDS
<222>      (1)..(1119)

```

```

<400>      3

```

```

atg cat aac cag gct cca att caa cgt aga aaa tca aca cgt att tac      48
Met His Asn Gln Ala Pro Ile Gln Arg Arg Lys Ser Thr Arg Ile Tyr
1              5              10              15

gtt ggg aat gtg ccg att ggc gat ggt gct ccc atc gcc gta cag tcc      96
Val Gly Asn Val Pro Ile Gly Asp Gly Ala Pro Ile Ala Val Gln Ser
              20              25              30

atg acc aat acg cgt acg aca gac gtc gaa gca acg gtc aat caa atc      144
Met Thr Asn Thr Arg Thr Thr Asp Val Glu Ala Thr Val Asn Gln Ile
              35              40              45

aag gcg ctg gaa cgc gtt ggc gct gat atc gtc cgt gta tcc gta ccg      192
Lys Ala Leu Glu Arg Val Gly Ala Asp Ile Val Arg Val Ser Val Pro
              50              55              60

acg atg gac gcg gca gaa gcg ttc aaa ctc atc aaa cag cag gtt aac      240
Thr Met Asp Ala Ala Glu Ala Phe Lys Leu Ile Lys Gln Gln Val Asn
65              70              75              80

gtg ccg ctg gtg gct gac atc cac ttc gac tat cgc att gcg ctg aaa      288
Val Pro Leu Val Ala Asp Ile His Phe Asp Tyr Arg Ile Ala Leu Lys
              85              90              95

gta gcg gaa tac ggc gtc gat tgt ctg cgt att aac cct ggc aat atc      336
Val Ala Glu Tyr Gly Val Asp Cys Leu Arg Ile Asn Pro Gly Asn Ile
              100              105              110

```

ggt aat gaa gag cgt att cgc atg gtg gtt gac tgt gcg cgc gat aaa Gly Asn Glu Glu Arg Ile Arg Met Val Val Asp Cys Ala Arg Asp Lys 115 120 125	384
aac att ccg atc cgt att ggc gtt aac gcc gga tcg ctg gaa aaa gat Asn Ile Pro Ile Arg Ile Gly Val Asn Ala Gly Ser Leu Glu Lys Asp 130 135 140	432
ctg caa gaa aag tat ggc gaa ccg acg ccg cag gcg ttg ctg gaa tct Leu Gln Glu Lys Tyr Gly Glu Pro Thr Pro Gln Ala Leu Leu Glu Ser 145 150 155 160	480
gcc atg cgt cat gtt gat cat ctc gat cgc ctg aac ttc gat cag ttc Ala Met Arg His Val Asp His Leu Asp Arg Leu Asn Phe Asp Gln Phe 165 170 175	528
aaa gtc agc gtg aaa gcg tct gac gtc ttc ctc gct gtt gag tct tat Lys Val Ser Val Lys Ala Ser Asp Val Phe Leu Ala Val Glu Ser Tyr 180 185 190	576
cgt ttg ctg gca aaa cag atc gat cag ccg ttg cat ctg ggg atc acc Arg Leu Leu Ala Lys Gln Ile Asp Gln Pro Leu His Leu Gly Ile Thr 195 200 205	624
gaa gcc ggt ggt gcg cgc agc ggg gca gta aaa tcc gcc att ggt tta Glu Ala Gly Gly Ala Arg Ser Gly Ala Val Lys Ser Ala Ile Gly Leu 210 215 220	672
ggt ctg ctg ctg tct gaa ggc atc ggc gac acg ctg cgc gta tcg ctg Gly Leu Leu Leu Ser Glu Gly Ile Gly Asp Thr Leu Arg Val Ser Leu 225 230 235 240	720
gcg gcc gat ccg gtc gaa gag atc aaa gtc ggt ttc gat att ttg aaa Ala Ala Asp Pro Val Glu Glu Ile Lys Val Gly Phe Asp Ile Leu Lys 245 250 255	768
tcg ctg cgt atc cgt tcg cga ggg atc aac ttc atc gcc tgc ccg acc Ser Leu Arg Ile Arg Ser Arg Gly Ile Asn Phe Ile Ala Cys Pro Thr 260 265 270	816
tgt tcg cgt cag gaa ttt gat gtt atc ggt acg gtt aac gcg ctg gag Cys Ser Arg Gln Glu Phe Asp Val Ile Gly Thr Val Asn Ala Leu Glu 275 280 285	864
caa cgc ctg gaa gat atc atc act ccg atg gac gtt tcg att atc ggc Gln Arg Leu Glu Asp Ile Ile Thr Pro Met Asp Val Ser Ile Ile Gly 290 295 300	912
tgc gtg gtg aat ggc cca ggt gag gcg ctg gtt tct aca ctc ggc gtc Cys Val Val Asn Gly Pro Gly Glu Ala Leu Val Ser Thr Leu Gly Val 305 310 315 320	960
acc ggc ggc aac aag aaa agc ggc ctc tat gaa gat ggc gtg cgc aaa Thr Gly Gly Asn Lys Lys Ser Gly Leu Tyr Glu Asp Gly Val Arg Lys 325 330 335	1008
gac cgt ctg gac aac aac gat atg atc gac cag ctg gaa gca cgc att Asp Arg Leu Asp Asn Asn Asp Met Ile Asp Gln Leu Glu Ala Arg Ile 340 345 350	1056
cgt gcg aaa gcc agt cag ctg gac gaa gcg cgt cga att gac gtt cag Arg Ala Lys Ala Ser Gln Leu Asp Glu Ala Arg Arg Ile Asp Val Gln 355 360 365	1104

WO 02/12478

PCT/US01/24335

cag gtt gaa aaa taa
 Gln Val Glu Lys
 370

1119

<210> 4
 <211> 686
 <212> PRT
 <213> Oryza sativa
 <400> 4

Met Ala Thr Gly Val Ala Pro Ala Pro Leu Pro His Val Arg Val Arg
 1 5 10 15
 Asp Gly Gly Ile Gly Phe Thr Arg Ser Val Asp Phe Ala Lys Ile Leu
 20 25 30
 Ser Val Pro Ala Thr Leu Arg Val Gly Ser Ser Arg Gly Arg Val Leu
 35 40 45
 Val Ala Lys Ser Ser Ser Thr Gly Ser Asp Thr Met Glu Leu Glu Pro
 50 55 60
 Ser Ser Glu Gly Ser Pro Leu Leu Gly Ile Thr Arg Arg Leu Leu Phe
 65 70 75 80
 Thr Leu His Met Val Gly Asn Val Pro Leu Gly Ser Asp His Pro Ile
 85 90 95
 Arg Ile Gln Thr Met Thr Thr Ser Asp Thr Lys Asp Val Ala Lys Thr
 100 105 110
 Val Glu Glu Val Met Arg Ile Ala Asp Lys Gly Ala Asp Phe Val Arg
 115 120 125
 Ile Thr Val Gln Gly Arg Lys Glu Ala Asp Ala Cys Phe Glu Ile Lys
 130 135 140
 Asn Thr Leu Val Gln Lys Asn Tyr Asn Ile Pro Leu Val Ala Asp Ile
 145 150 155 160
 His Phe Ala Pro Thr Val Ala Leu Arg Val Ala Glu Cys Phe Asp Lys
 165 170 175
 Ile Arg Val Asn Pro Gly Asn Phe Ala Asp Arg Arg Ala Gln Phe Glu
 180 185 190
 Gln Leu Glu Tyr Thr Glu Asp Asp Tyr Gln Lys Glu Leu Glu His Ile
 195 200 205
 Glu Lys Val Pro Asn Ile Ser Leu Phe Ser Val Asn Leu Val Phe Ser
 210 215 220
 Pro Leu Val Glu Lys Cys Lys Gln Tyr Gly Arg Ala Met Arg Ile Gly
 225 230 235 240
 Thr Asn His Gly Ser Leu Ser Asp Arg Ile Met Ser Tyr Tyr Gly Asp
 245 250 255
 Ser Pro Arg Gly Met Val Glu Ser Ala Leu Glu Phe Ala Arg Ile Cys
 260 265 270

Arg	Lys	Leu	Asp	Phe	His	Asn	Phe	Val	Phe	Ser	Met	Lys	Ala	Ser	Asn	275	280	285
Pro	Val	Ile	Met	Val	Gln	Ala	Tyr	Arg	Leu	Leu	Val	Ala	Glu	Met	Tyr	290	295	300
Asn	Leu	Gly	Trp	Asp	Tyr	Pro	Leu	His	Leu	Gly	Val	Thr	Glu	Ala	Gly	305	310	315
Glu	Gly	Glu	Asp	Gly	Arg	Met	Lys	Ser	Ala	Ile	Gly	Ile	Gly	Thr	Leu	325	330	335
Leu	Met	Asp	Gly	Leu	Gly	Asp	Thr	Ile	Arg	Val	Ser	Leu	Thr	Glu	Pro	340	345	350
Pro	Glu	Glu	Glu	Ile	Asp	Pro	Cys	Arg	Arg	Leu	Ala	Asn	Leu	Gly	Thr	355	360	365
His	Ala	Ala	Asp	Leu	Gln	Ile	Gly	Val	Ala	Pro	Phe	Glu	Glu	Lys	His	370	375	380
Arg	Arg	Tyr	Phe	Asp	Phe	Gln	Arg	Arg	Ser	Gly	Gln	Leu	Pro	Leu	Gln	385	390	395
Lys	Glu	Ala	Pro	Glu	Leu	Leu	Tyr	Arg	Ser	Leu	Ala	Ala	Lys	Leu	Val	405	410	415
Val	Gly	Met	Pro	Phe	Lys	Asp	Leu	Ala	Thr	Val	Asp	Ser	Ile	Leu	Leu	420	425	430
Lys	Glu	Leu	Pro	Pro	Val	Glu	Asp	Ala	Gln	Ala	Arg	Leu	Ala	Leu	Lys	435	440	445
Arg	Leu	Val	Asp	Ile	Ser	Met	Gly	Val	Leu	Thr	Pro	Leu	Ser	Glu	Gln	450	455	460
Leu	Thr	Lys	Pro	Leu	Pro	His	Ala	Ile	Ala	Leu	Val	Asn	Val	Asp	Glu	465	470	475
Leu	Ser	Ser	Gly	Ala	His	Lys	Leu	Leu	Pro	Glu	Gly	Thr	Arg	Leu	Ala	485	490	495
Val	Thr	Leu	Arg	Gly	Asp	Glu	Ser	Tyr	Glu	Gln	Leu	Asp	Leu	Leu	Lys	500	505	510
Gly	Val	Asp	Asp	Ile	Thr	Met	Leu	Leu	His	Ser	Val	Pro	Tyr	Gly	Glu	515	520	525
Glu	Lys	Thr	Gly	Arg	Val	His	Ala	Ala	Arg	Arg	Leu	Phe	Glu	Tyr	Leu	530	535	540
Glu	Thr	Asn	Gly	Leu	Asn	Phe	Pro	Val	Ile	His	His	Ile	Glu	Phe	Pro	545	550	555
Lys	Ser	Val	Asn	Arg	Asp	Asp	Leu	Val	Ile	Gly	Ala	Gly	Ala	Asn	Val	565	570	575
Gly	Ala	Leu	Leu	Val	Asp	Gly	Leu	Gly	Asp	Gly	Val	Leu	Leu	Glu	Ala	580	585	590
Ala	Asp	Gln	Glu	Phe	Glu	Phe	Leu	Arg	Asp	Thr	Ser	Phe	Asn	Leu	Leu	595	600	605

WO 02/12478

PCT/US01/24335

Gln Gly Cys Arg Met Arg Asn Thr Lys Thr Ile Ala Ile Met Gly Cys
610 615 620

Ile Val Asn Gly Pro Gly Glu Met Ala Asp Ala Asp Phe Gly Tyr Val
625 630 635 640

Gly Gly Ala Pro Gly Lys Ile Asp Leu Tyr Val Gly Lys Thr Val Val
645 650 655

Gln Arg Gly Ile Ala Met Glu Gly Ala Thr Asp Ala Leu Ile Gln Leu
660 665 670

Ile Lys Asp His Gly Arg Trp Val Asp Pro Pro Val Glu Glu
675 680 685

<210> 5
<211> 594
<212> DNA
<213> Arabidopsis thaliana

<220>
<221> unsure
<222> (1..594)
<223> unsure at all n locations

<400> 5

aaaatcgtca atccctctca aactcttctc accactaatt tcttcctctg gaacattctc 60
ttctctatta ttttgattcc cttggcctca acactgggtt ctcaattgca tgatcttggc 120
tcgtcttcag ttactttgat tcaactgagaa aaatggcgac tggagtattg ccagctccgg 180
tttctgggat caagataccg gattcgaaag tcgggttttg taaaagcatg aatcttgtga 240
gaatttgtna tgtaggagt ctaagatctg ctaggagaag agtttcgggt atccggaatt 300
caaaccaagg ctctgattta gctgagcttc aaccctgcat ccgaaggaaa gccctcttc 360
ttagtgccaa ggcaggaaat attgtgaatc attgcataa gcggttagga ggaagnctcg 420
gacctgtaat ggttgaaatg tcgncccttn gaagnnaca ccggtanggg tcaaacggtg 480
ccttcttngg gtacaaaang tnttccttgg ancctnttng tgggggtttt gggattgcgg 540
aaaaaggggc tgnnttttnaa ggnnacctnn caaggnagna agggnggggc tttt 594

<210> 6
<211> 615
<212> DNA
<213> Glycine max

<220>
<221> unsure
<222> (1..615)
<223> unsure at all n locations

<400> 6

accagaagtg atgagcctta tgaagaactg gacattctta aggggtgttg tgctactatg 60
cttttccatg accttcctta tacagaagac agaattagca gaggatgcgc aaccagacgg 120

ttatttgagt acctatctga caattctcta aacttccttg ttattcacca tattcagttc 180
 ccaaattggga ttcacagggg tgacttggtg attgggtgctg gttctgatgc tggagccctt 240
 ctgggttgatg ggcttggaga tggactactt ttggaagccc cggacaagga ttttgaattt 300
 attagaaaca cttctttcaa tttgttgcaa ggctgcagaa tgagaaatac aaagacagag 360
 tatgtctcat gtccatcctg tggcagaaca ttgtttgatc ttcaagaagt aagtgcacaa 420
 attcgggaga agacatcaca cctncctggg gtttcgattg caatcatggg atgcattggt 480
 aatggaccag gggagatggc tgatgcagac tttgggtatg tgggaagcac tccccggaag 540
 attgacctct atgttgggaa gactgggtgtg aagcgtggga attcaatgga gcatgccaac 600
 catggcttga tccga 615

<210> 7
 <211> 589
 <212> DNA
 <213> Lycopersicon esculentum
 <400> 7

tggcgatgaa tcacatgatg agttggaaat cctgaagagc tctgatgtta caatgattct 60
 tcataatctg ccatatacag aggaaaaaat tggcaggggt caagcagcca ggaggctttt 120
 tgagtatctt tccgagaatt ccttgaactt tccagtgatt catcacatac aatttcccag 180
 caacacccac agagatgact tagtgattgg tgccgggaca aatgcgggag ccctcttggt 240
 agatgggctt ggtgatggac ttctcttgga agctccagac aaggattttg attttctcag 300
 aaatacatct ttcaatttgc ttcaagggtg cagaatgcgg aacacaaaaa cggaatatgt 360
 atcatgcca tcctgtggca gaactttatt cgatcttcaa gagataagcg ctcaaattag 420
 agagaagacg tcacacttgc ctggtgtttc aattgccatc atgggttgca ttgtgaatgg 480
 acctggggag atggctgatg ctgacttttg atatgttggg ggtgctcctg gaaagattga 540
 cctttacgtc ggcaagacag tggtgaaacg ccctattgaa atggagcat 589

<210> 8
 <211> 617
 <212> DNA
 <213> Mesembryanthemum crystallinum
 <400> 8

gaaaagcata gacattattt tgactttcaa cgtagaactg gtcaattacc gattcagaaa 60
 gagggatgaag atgtggacta tagaggtgtc ctacaccgtg atggttctgt cctcatgact 120
 gtttccttgg acatgttgaa gacacctgaa ctcttttaca agtcattagc agcaaagctt 180
 gttgttggca tgccatttaa ggatctggct actgtagact ctatttttct gagagagctt 240
 tcaccagtag atgactctga tgctcggcta gctctgaaga ggttaataga tataagtatg 300

ggtgtcatag ctcttttttc tgagcaactg acaaagccct tgccaaatgc aattgtattg 360
 gtgaacctta aagagttgtc aaccggtgca tacaagcttt taccagtagg aaccgcttg 420
 gcagtatctg tgcgaggtga tgaaccatat tgagacattg gagatcctta aagatattga 480
 tgcttcaatg gctttttatg aactgtcttt taccgagagg atattcacac agtgcattgct 540
 ggaccaaagc ttttgaggtc ctatcagata agcttggacc tcccgtaatt aacatattct 600
 atcccttcgg attaagg 617

<210> 9
 <211> 416
 <212> DNA
 <213> Oryza sativa

<220>
 <221> unsure
 <222> (1..416)
 <223> unsure at all n locations

<400> 9

ggattcggca cgagtctaatt tgatggtctt ggtgatggtg tacttcttga aagctgctga 60
 ccaagaaatt tgagtttttg agggacacat cctccaactt gttacagggc tgcaggatgc 120
 gcaacacaaa aacggaatat ttccctgggc ctcttgggtg ggggacacnc ttnaccncc 180
 aaaaattcan tgctcaaatt aaanaaaaaa ccnctcatct gccaggcntc tctattgcta 240
 tcatgggtng cattgtcaat gggccagggg aaatggccaa tcctaattnc ggatacttng 300
 gaggtgccct ggagaaaatc nacctntatn ttggttnttt ttttttnnaac ggggcatngc 360
 aanagaaggg ggcccnacc ccnanatnch ttcnccgggn ccngggccgn ggggtt 416

<210> 10
 <211> 621
 <212> DNA
 <213> Zea mays

<400> 10

gaattcggca ccagaagcca ctcccacatg caattgtact tgtcaacctc gacgaattgt 60
 caagtgggtg acacaaactt ttgccagaag gcactagact agctgtcact ctctgtgggtg 120
 atgaatcata cgagcagcta gatattctta aggatgttga tgatataaca atgttggttac 180
 ataatgttcc atatggtgag gagaagacag gcagggtgca tgctgctagg aggttatttg 240
 agtacttaca ggccaatggc ttgaacttcc ctgtaattca tcacataaat ttccctgaaa 300
 ccattgacag agatggtctt gtcattgggtg ctggggccaa cgttgggtgct ctcttagtgc 360
 atggtcttgg tgatggtgta ttccctgaag ctgctgacca ggaatttgag tttctgaggg 420
 acacatcttt caacttgctc caagggtgca ggatgctgca cacaaaaact gaatatgtgt 480

cttgtccttc ctgcgccga acactctttg accttcagga aatcagcgct gagattagag 540
 aaaagacctc tcatctgccg ggtgtctcga tcgctatcat gggctgtatt gcaatggacc 600
 aggagagatg gctgatgccg a 621

<210> 11
 <211> 601
 <212> DNA
 <213> Pinus taeda

<220>
 <221> unsure
 <222> (1..601)
 <223> unsure at all n locations

<400> 11

aatgcaagaa gtacggaagg gcaatgcaa ttggcacaaa ccatggaagt ctttccgac 60
 gtactatgag ttattatggt gattctccca ggggtatggt ggaatcagca tttgaatttg 120
 cacgcatttg ccggaagttg ggttttcata attttgtgtt ttcaatgaaa gcgagcgac 180
 ctgtagtcat ggttcaggca taccgtttac ttgttgccga gatgtatgtg caaggatggg 240
 attatccatt gcatttagga gttactgaag ctggtgaagg tgaagatgga cgcataaagt 300
 ctgcaattgg cattggaaca cttttgcagg atggtttggg tgatactatt cgagtttccc 360
 ttacagaacc tccagaagag gagatcaatc cctgtagaag acttgcaaat cttgggatgc 420
 aagctgcaaa gctanggaaa ggagtggctc cttttgagga gaacatcgtc attactttac 480
 tttccaacgc angactggcn agctccagta cagaaggagg gtgatgaggt ggatacagag 540
 gagtccgcat cgtgatggtc tgttctaatt tcagtgtcct tgacagntga agacacanaa 600
 a 601

<210> 12
 <211> 443
 <212> DNA
 <213> Physcomitrella patens

<400> 12

gcacgtatct gccgcaaaca tgactatatt aatttcttgt tttctatgaa agcaagcaat 60
 ccggtcgtaa tggttcaagc atatcggctt ttagtatctg agatgtatgt gaacaactgg 120
 gactacccat tacatcttgg tgttactgag gctggagagg gagaggatgg tcgcatgaag 180
 tcagctatcg gcattggtgc tttacttcag gatggtctcg gtgacaccat acgtgtttca 240
 ttgacggaag ctctgaaga agaaattgat ccttgcaaaa agcttgcaaa ccttggcatg 300
 aagatttctg cagaacagaa gggggtggct gaattcgaag agaagcaccg gcgatacttt 360
 gacttccaac gaaggaccgg ccaacttcca ctgcagaggg agggagagtt ggtggactac 420
 agaaacgttc tgcaccgtga tgg 443

```

<210>      13
<211>      938
<212>      DNA
<213>      Arabidopsis thaliana

<220>
<221>      unsure
<222>      (1..938)
<223>      unsure at all n locations

<400>      13

atgatactgc cagctannnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn   60
nnnnnnnnnn nnnnnnnnnn nnnccacgcg tccgaaaacg ttttatcctg agtttctttc   120
accatccagc ttcatttgtg aaaaatcgtc aatccctctc aaactcttct caccactaat   180
ttcttcctct ggaacattct cttctctatt attttgattc ccttggcctc aacactgggt   240
tctcaattgc atgatcttgg ctcgctttca gttactttga ttcactgaga aaaatggcga   300
ctggagtatt gccagctccg gtttctggga tcaagatacc ggattcgaaa gtcggggtttg   360
gtaaaagcat gaatcttgtg agaatttgtg atgttaggag tctaagatct gctaggagaa   420
gagtttcggt tatccggaat tcaaaccaag gctctgattt agctgagctt caacctgcat   480
ccgaaggaag ccctctctta gtgccaagac agaaatattg tgaatcattg cataagacgg   540
tgagaaggaa gactcgtact gttatggttg gaaatgtcgc ccttggaagc gaacatccga   600
taaggattca aacgatgact acttcggata caaaagatat tactggaact gttgatgagg   660
ttatgagaat agcggataaa ggagctgata ttgtaaggat aactgtccaa gggaagaaag   720
aggcggatgc gtgctttgaa ataaaagata aactcgttca gcttaattac aatataccgc   780
tggttgcaga tattcattgt gccctactg tagccttacg agtcgctgaa tgctttgaca   840
agatccgtgt caaccagga aattttgcgg acaggcgggc ccagtttgag acgattgatt   900
atacagaaga tgaatatcag aaagaactcc agcatatc                               938

```

```

<210>      14
<211>      432
<212>      DNA
<213>      Arabidopsis thaliana

<400>      14

agcataacaa ggctctgatt tagctgagct tcaacctgca tccgaaggaa gccctctctt   60
agtgccaaaga cagaaatatt gtgaatcatt gcataagacg gtgagaagga agactcgtac   120
tgttatgggt ggaaatgtcg cccttggaag cgaacatccg ataaggattc aaacgatgac   180
tacttcggat acaaaagata ttactggaac tgttgatgag gttatgagaa tagcggataa   240
aggagctgat attgtaagga taactgttca agggaagaaa gaggcggatg cgtgctttga   300

```

aataaaagat aaactcggtc agcttaatta caatataccg ctgggtgcag atattcattt 360
 tgcccctact gtagccttac gagtcgctga atgctttgac aagatccgtg tcaaccaag 420
 aaattttgcg ga 432

<210> 15
 <211> 528
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <221> unsure
 <222> (1..528)
 <223> unsure at all n locations

<400> 15

tgatacgcca gctctatacg actcactatt aggggaagctg gtacgcctgc aggtacccgg 60
 tccgggaatt ccnngggtcg acccacgcgt ccgaaagaac tccagcatat cgagcaggtc 120
 ttcactcctt tgggttgagaa atgcaaaaag tacgggagag caatgcgtat tgggacaaat 180
 catggaagtc tttctgaccg tatcatgagc tattacgggg attctccccg aggaatgggt 240
 gaatctgcgt ttgagtttgc aagaatatgt cggaaattag actatcacia ctttgttttc 300
 tcaatgaaag cgagcaaccc agtgatcatg gtccaggcgt accgtttact tgtggctgag 360
 atgtatgttc atggatggga ttatcctttg catttgaggag ttactgaggc aggagaaggc 420
 gaagatggac ggatgaaatc tgcgattgga attgggacgc ttcttcagga cgggctcggg 480
 gacacaataa gagtttcact gacggagcca ccagaagagg agatagat 528

<210> 16
 <211> 379
 <212> DNA
 <213> Arabidopsis thaliana

<400> 16

gcgtattggg acaaatcatg gaagtctttc tgaccgtatc atgagctatt acggggattc 60
 tccccgagga atgggtgaat ctgcgtttga gtttgcaaga atatgtcgga aattagacta 120
 tcacaacttt gttttctcaa tgaaagcgag caaccagtg atcatgggtcc aggcgtaccg 180
 ttacttgtg gctgagatgt atgttcatgg atgggattat cttttgcatt tgggagttac 240
 tgaggcagga gaaggcgaag atggacggat gaaatctgcg attggaattg ggacgcttct 300
 tcaggacggg ctcggtgaca caataagagt ttactgacg gagccaccag aagaggagat 360
 agatccctgc aagcgattg 379

WO 02/12478

PCT/US01/24335

<210> 17
 <211> 395
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 17
 aaagaactcc agcatatcga gcaggtcttc actccttttg ttgagaaatg caaaaagtac 60
 gggagagcaa tgcgtattgg gacaaatcat ggaagtcttt ctgaccgtat catgagctat 120
 tacggggatt ctccccgagg aatgggttgaa tctgcgtttg agtttgcaag aatatgtcgg 180
 aaattagact atcacaaactt tgttttctca atgaaagcga gcaaccagc gatcatgggc 240
 caggcgtacc gtttacttgt ggctgagatg tatgttcatg gatgggatta tcctttgcat 300
 ttgggagtta ctgaggcagg agaaggcgaa gatggacgga tgaaatctgc gattggaatt 360
 ggggacactt cttcaggacg ggctcgggtga cacaa 395

<210> 18
 <211> 395
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 18
 aaagaactcc agcatatcga gcaggtcttc actccttttg ttgagaaatg caaaaagtac 60
 gggagagcaa tgcgtattgg gacaaatcat ggaagtcttt ctgaccgtat catgagctat 120
 tacggggatt ctccccgagg aatgggttgaa tctgcgtttg agtttgcaag aatatgtcgg 180
 gaattagact atcacaaactt tgttttctca atgaaagcga gcaaccagc gatcatgggc 240
 caggcgtacc gtttacttgt ggctgagatg tatgttcatg gatgggatta tcctttgcat 300
 ttgggagtta ctgatgcagg agaaggcgaa gatggacgga tgaaatctgc gattggaatt 360
 gggacgcttc ttcaggacgg gctcgggtgac acaat 395

<210> 19
 <211> 412
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 19
 atgctggagg ccttcttggt gatggactag gtgatggcgt aatgctcgaa gcacctgacc 60
 aagatthtga ttttcttagg aatacttcct tcaacttatt acaaggatgc agaatgcgta 120
 acactaagac ggaatatgta tcgtgcccggt cttgtggaag aacgcttttc gacttgcaag 180
 aaatcagcgc cgagatccga gaaaagactt cccatttacc tggcgtttcg atcgcaatca 240
 tgggatgcat tgtgaatgga ccaggagaaa tggcagatgc tgatttcgga tatgtagggtg 300
 gttctcccgg aaaaatcgac ctttatgtcg gaaagacggg ggtgaagcgt gggatagcta 360
 tgacggaggc aacagatgct ctgatcgggc tgatcaaaga acatgggtcgt tg 412

<210> 20
 <211> 1172
 <212> DNA
 <213> Arabidopsis thaliana

 <220>
 <221> unsure
 <222> (1..1172)
 <223> unsure at all n locations

 <400> 20

 gggatatgcca ttcaaggatc tggcaactgt tgattcaatc ttattaaaga gagctaccgc 60
 ctgtagatga tcaagtggct cgtttggctc taaaacgggt gattgatgtc agtatgggag 120
 ttatagcacc tttatcagag caactaaca agccattgcc caatgccatg gttcttgtca 180
 acctcaagga actatctggt ggcgcttaca agcttctccc tgaagggtaca cgcttggttg 240
 tctctctacg aggcgatgag ccttacgagg agcttgaaat actcaacaac attgatgcta 300
 cgatgattct ccatgatgta cctttcactg aagacaaagt tagcagagta catgcagctc 360
 ggaggctatt cgagttctta tccgagaatt cagttaactt tcctggttatt catcacataa 420
 acttcccaac cggaatccac agagacgaat tgggtgattca tgcagggaca tatgctggag 480
 gccttcttgt ggatggacta cgtgatggcg taatgctcga agcacctgac caagattttg 540
 attttcttag gaatacttcc ttcaacttat tacaaggatg cagaatgcgt aacactaaga 600
 cggaatatgt atcgtgcccc tcttgtggaa gaacgctttt cgacttgcaa gaaatcagcg 660
 ccgagatccg agaaaagact tcccatctac ctggcgtttc gatcgcaatc atgggatgca 720
 ttgtgaatgg accaggagaa atggcagatg ctgatttcgg atatgtaggt ggttctcccc 780
 gaaaaatcga cctttatgtc ggaaagacgg tggatgaagcg tgggatagct atgacggagg 840
 caacagatgc tctgatcggc ctgatcaaag aacatggctc ttgggtcgac ccgcccgtgg 900
 ccgatgagta gatttcaaaa cggagaaaga tgggtgggccc attctttgaa aactgtgaga 960
 ggagatatat atatttgtgt gtgtatatca tctgtttgtt gtgtattgca tcattcattt 1020
 tggacaaatg tccaaattct ctttaagttga taaaagttct taggccaaat taaatttaat 1080
 ataaaaaaaa aaaaaaaaaa gcnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 1140
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nn 1172

<210> 21
 <211> 584
 <212> DNA
 <213> Zea mays

 <400> 21

 caggttaatt aattcctgta cgccgtcggc ttcgggtact cgtttaattt cttcccgacc 60

WO 02/12478

PCT/US01/24335

acggttgatg gcaatgtaac cggcttgttt acccacatag ccatagtcgg catcggccat 120
 ttccccgggg ccattgacaa tacagcccat gacggcgatg tctaaacccg ttagatgttt 180
 agtggcttct cggacttcat gtaacacgtc ttccaagttg aacaacgtgc ggccacagga 240
 aggacaggcc acatattcca ccatggtttt cgcgcaaccc agcgccctgga gaatgctgta 300
 gcaaacggga atttcttttt cgggggcttc ggtgagggat acccggatag tatcgccaat 360
 gccatcagct aaaagggtgg caatgccagc ggtggattta atgcggccat attccccatc 420
 cccggcttcg gtaacccta gatggagggg ataatccatg cccaactcgt tcatacgttt 480
 caccatgagg cgataggcgg ccaacattac cggtaaccgg gacgctttca tggaaacgac 540
 taggttgagg aaatctaaag actcacaat tttgatgaat tcca 584

<210> 22
 <211> 670
 <212> DNA
 <213> Zea mays

<400> 22

caggtcgact ctagaggatc ggcgttaacc atggttctct ctccgaaaga atgcttttac 60
 ctacttttta cccccgaggg catggtgcaa tcggccctgg aattcatcaa aatttgtgag 120
 tccttagatt tccgcaacct agtcgtttcc atgaaagcgt cccgggtacc ggtaatgttg 180
 gccgcctatc gcctcatggt gaaacgtatg gacgagttgg gcatggatta tccccccat 240
 ctaggggtta ccgaagccgg ggatggggaa tatggccgca ttaaattccac cgctggcatt 300
 gccacccttt tagctgatgg cattggcgat actatccggg tatccctcac cgaagcccc 360
 gaaaaagaaa ttcccgtttg ctacagcatt ctccaggcgc tgggtttgcg gaaaaccatg 420
 gtggaatatg tggcctgtcc ttctgtggc cgcacgttgt tcaacttgga agacgtgtta 480
 catgaagtcc gagatgccac taaacatcta acgggttttag actttcgccg tcatgggctg 540
 tattgtcaat ggccccgggg caatggccga tgccgactat ggctatgtgg gtaaacaagc 600
 cggttacatt gccatcaacc gtggtcggga agaaattaaa cgagtaccgg aaaccgacgg 660
 cgtacaggaa 670

<210> 23
 <211> 596
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (1..596)
 <223> unsure at all n locations

<400> 23

caggtcgact ctagaggatc ggcgttaacc atggttctct ctccgaaaga atgcttttac 60

ctacttttta cccccgaggg catggtgcaa tcggccctgg aattcatcaa aatttgtgag 120
 tccttagatt tccgcaacct agtcgtttcc atgaaagcgt cccgggtaac ggtaatgttg 180
 gccgcctatc gcctcatggt gaaacgtatg gacgagttgg gcatggatta tcccctccat 240
 ctaggggtta ccgaagccgg ggatggggaa tatggccgca ttaaataccac cgctggcatt 300
 gccacccttt tagctgatgg cattggcgat actatccggg tatccctcac cgaagcccc 360
 gaaaaagaaa ttcccgtttg ctacagcatt ctccaggcgc tgggtttgcg gaaaaccatg 420
 gtggaatatg tggcctgtcc ttccctgtggc cgcacgttgt tcaacttgga agacgtgtta 480
 catgaagtcc gagatgccac taaacatcta acgtgttttag actttcgncg tcatgtgctg 540
 tattgtcaat ggccccggtg caatggccga tgccgactat ggctatgtgg gtaaac 596

<210> 24
 <211> 403
 <212> DNA
 <213> Zea mays

<400> 24

cagacaagga ggaggaaaac tcgaactgtg atggtgggga atgtgccact tgggagtgat 60
 caccataa ggattcaaac catgacgact tcagatacca aggatgttgc gaaaacagta 120
 gaggaggtga tgaggatagc agataaagga gctgatcttg ttagaataac agtccagggt 180
 aggaaggaag ctgatgcctg ctttgagatc aagaacactc tggttcagaa gaattacaac 240
 attccactag tggccgatat tcattttgct cctacggtag ctctaaagggt ggcagaatgt 300
 tttgacaaaa ttctgttgaa cccaggaaat tttgctgac gtcgtgctca atttgaaaag 360
 ctggaatata ctgacgacga ctacaaaaa gagctagagc ata 403

<210> 25
 <211> 293
 <212> DNA
 <213> Zea mays

<400> 25

cagacaaggc ggaggaaaac tcgaactgtg atggtgggga atgtgccact tggcagtgat 60
 caccataa ggattcaaac catgacgact tcagatacca aggatgttgc gaaaacagta 120
 gaggaggtga tgaggatagc agataaagga gctgatcttg ttagaataac agtccagggt 180
 aggaaggaag ctgatgcctg ctttgagatc aagaacactc tggttcagaa gaattacaac 240
 attccactag tggccgatat tcattttgct cctacggtag ctctaagggt ggc 293

WO 02/12478

PCT/US01/24335

<210> 26
 <211> 456
 <212> DNA
 <213> Zea mays

<400> 26

cagacaaggc ggaggaaaac tcgaactgtg atggtgggga atgtgccact tggcagtgat 60
 caccataa ggattcaaac catgacgact tcagatacca aggatgttgc gaaaacagta 120
 gaggaggtga tgaggattgc agataaagga gctgatcttg ttagaataac agtccagggt 180
 aggaaggaag ctgatgcctg ctttgagatc aagaacaact ctggttcaga agaattacaa 240
 ccttccacta gtggacctga tattcatttt gtccttcag tagcttttaa ggtggcagaa 300
 tgtttggaca aattaattga aacacacaat ttcttggtga tagtgtacct taattagaaa 360
 agctggaatt taccggctac gacttccata aagcgcttgg gcttgtttaa caattggttt 420
 ttaccttaat cgaatatattc acagaaattt gaattt 456

<210> 27
 <211> 619
 <212> DNA
 <213> Zea mays

<400> 27

caccgaaggc ttctaattta tttctcagat ctcaataaat gtacaaaatg tgtagggatg 60
 atgtacattg tatgctcagt tcctgcattg cgtgtttcgc tttacagaat atataaacta 120
 cagacttggc tacagcctac agccctactc ctccggcagga ggatccaccc atcggccatg 180
 gtccttgatc agctggatca aggcgtcagt tgcaccttc atggcgatgg cgcgctgcac 240
 aacggtcttg ccaacataaa ggtcgatctt tccgggagcg cctccaacgt atccgaaatc 300
 ggcatcagcc atctctcctg gtccattgac aatacaaccc atgatagcga tcgaaacacc 360
 tggcagatga gaggtctttt ctctaatttc agcgctgatt tcctgaaggc caaagagtgt 420
 tcggccgcag gaaggacaag acacatatcc agtttttgtg ttgcgcatcc tgcaaccttg 480
 gagcaagttg aaagatgtgt ccctcaggaa ctcaaattcc tggtcagcag cttcaaggaa 540
 tacaccatca ccaagaccat cgactaagag agcaccaacg ttggccccag caccaatgac 600
 aagaccatct ctgtcaatg 619

<210> 28
 <211> 422
 <212> DNA
 <213> Zea mays

<400> 28

tcgcttgcac ttgggtgtta cagaagctgg agagggtgaa gatggaagga tgaaatctgc 60
 tattggcatt gggacactgc taatggatgg tttgggtgat acaatccgtg tctccctcac 120

agaaccacca gaagaagaga ttgatccttg ccaaagggttg gcaaactcttg ggacgcaggc 180
 cgcaaaccctt caaattgggg tggccccatt tgaagaaaag cacaggcgct attttgattt 240
 ccagcgtagg agtgggtcaat tgcctttgca gaaggaggga ggcgatagtt gactacagaa 300
 atgtcctgca tcgtgatggg atctgactga tggcagtttc cctggatcag ttgaaggctc 360
 ctgatctcct ttataggtat attgcagcaa agcttgcgga tggcatgcct ttcaaggatc 420
 tg 422

<210> 29
 <211> 430
 <212> DNA
 <213> Zea mays

<400> 29

tcgcttgac ttgggtgtta cagaagctgg agaggggtgaa gatggaagga tgaaatctgc 60
 tattggcatt gggacactgc taatggatgg tttgggtgat acaatccgtg tctccctcac 120
 agaaccacca gaagaagaga ttgatccttg ccaaagggttg gcaaactcttg ggacgcaggc 180
 tgcaaaccctt caaattgggg tggccccatt tgaagaaaag cacaggcgctt attttgattt 240
 ccagcgtagg agtgggtcaat tgcctttgca gaaggagggt gaggaagttg actacagaaa 300
 tgtcctgcat cgtgatggta tctgtactga tggcagtttc cctggatcag ttgaaggctc 360
 ctgatctcct ttataggtct cttgcagcaa agcttgcggt tggcatgcct ttcaaggatc 420
 tggctactgt 430

<210> 30
 <211> 528
 <212> DNA
 <213> Zea mays

<400> 30

gacaggcagg gtgcatgctg ctaggagggtt atttgagtac ttacaggcca atggcttgaa 60
 cttccctgta attcatcaca taaatttccc tgaaaccatt gacagagatg gtcttgtcat 120
 tggggctggg gccaacgttg gtgctctctt agtcgatggg cttgggtgatg gtgtattcct 180
 tgaggcggct gaccaggaat ttgagttcct gagggacaca tctttcaact tgctccaagg 240
 ttgcaggatg cgcaacacaa aaactgaata tgtgtcttgt ccttcctgcg gccgaacact 300
 ctttgacctt caggaaatca gcgctgagat tagcgaaaag acctctcatc tgccacgtgt 360
 ttcgatcgct atcatgggtt gtattgtcaa tggaccagga gcgctggctg atgccgattt 420
 cggatacgct ggccggcgctc ccggaaagat cgacctttat attggcacga ccgttatgca 480
 gcgcgccatc gccatggacg gtgcaactga cgccttgatc cagctgat 528

WO 02/12478

PCT/US01/24335

<210> 31
 <211> 303
 <212> DNA
 <213> Zea mays

<400> 31

ggggccaacg ttggtgctct cttagtcgat ggtcttggtg atggtgtatt ccttgaggcg 60
 gctgaccagg aatttgagtt cctgagggac acatctttca acttgctcca aggttgcagg 120
 atgcgcaaca caaaaactga atatgtgtct tgtccttcct gcggccgaac actctttgac 180
 cttcaggaaa tcagcgctga gattagagaa aagacctctc atctgccacg tgtttcgatac 240
 gctatcatgg gttgtattgt caatggacca ggagagatgg ctgatgccga tttcggatac 300
 gtt 303

<210> 32
 <211> 613
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (1..613)
 <223> unsure at all n locations

<400> 32

cgagatggcg ttccatgccn ggcccttcct cctcttcctc ttcttctgcc cccccgctgg 60
 cttggaaaag ggagagaaac tcgcgcactc gggtatcgaa gggaggagcg cgggagagg 120
 tgaggtttcg cccacacgga gctgcgaggt gtttgttagga tctcctaggt gagccccctgc 180
 tgcttggaga cagccatggc caccggcggtg gctccagctc ctctcccaca tgtcagagt 240
 cgtcatgggg gcgtcgggtt caccaggagc gtcgattttg cgaaggctct gtctgctccc 300
 ggtgccggca cgatgagagc aagctcctct agaggcaggg cgctcgtggc gaagagctct 360
 agtactggct cggagaccat ggagctcgag ccatcttcag aaggaagccc acttttagta 420
 cccaggcaga agtactgtga atcaacacac cagacaagga ggaggaaaac tcgaactgtg 480
 atggtgggga atgtgccact tggcagtgat catcccataa ggattcaaac catgacgact 540
 tcagatacca aggatgttgc aaaaacagta gaggaggtga tgaggatagc agataaagga 600
 gctgatcttg tta 613

<210> 33
 <211> 464
 <212> DNA
 <213> Glycine max

<400> 33

agagcatgaa atcttctgcg aggaaaaggg tgtcaattat cacgaactca aatcctggcc 60

aagatattgc tgaacttcaa cctgcatccc caggaagccc tcttttgggt cctaggcaaa 120
 agtattgtga atcattgcac aaacccatca ggagaaaaac aagcacagta atgggttggt 180
 acgtggctat tggtagcgag catcctataa gaattcagac catgactaca actgacacta 240
 aggatgttgc tgggacagtt gaacaggtga tgagaatagc agataaagga gctgatattg 300
 tacggataac agttcaaggg aagaaagaag ctgatgcttg ttttgagatt aaaaacaccc 360
 ttgtgcagaa aaattacaac ataccctggg tggctgatat tcattttgct ccctctgttg 420
 ctttgcggt agctgaatgc tttgataaga ttcgtgtaaa ccct 464

<210> 34
 <211> 705
 <212> DNA
 <213> Glycine max

<400> 34

gtagctgaat gctttgataa gattcgtgta aaccctggaa attttgctga tagacgggct 60
 caatttgaaa cattagagta cacagaagaa gactatcaga aagaacttga gcatattgaa 120
 aaggttttca caccattggg tgagaaatgt aagaaatatg ggagagcaat gcgcattggg 180
 acaaaccatg gaagtctttc tgatcgtata atgagctact atggagactc gcctagggga 240
 atggtagaat ctgcttttga atttgcaagg atatgccgaa agttagacta tcacaatttt 300
 gttttttcta tgaaagcaag caaccagtt atcatgggtc aggcataccg cttacttgtg 360
 gctgaaatgt atgtccaagg ctgggattat ccattacact tgggtgttac tgaagctgga 420
 gaaggtgagg atgggaggat gaagtctgca ataggcattg gaactcttct tcaggatgga 480
 ttgggtgata caattagggg ttctctcaca gaaccaccag aggaggagat agacccttgc 540
 agaaggttgg caaatcttgg aatgatagct tctgaactcc agaagggggg ggaacctttt 600
 gaagaaaagc acagacatta ttttcgactt tcagcgccga tctggtcaat tgccagtgca 660
 aaaagagggt gaggaggtgg attacagagg tgtactgcac cgtga 705

<210> 35
 <211> 564
 <212> DNA
 <213> Glycine max

<220>
 <221> unsure
 <222> (1..564)
 <223> unsure at all n locations

<400> 35

aagcncggaa ttcggctcga gaggaactca aatcctggcc aagatattgc tgaacttcaa 60
 cctgtatccc caggaagccc tcttttgggt cctaggcaaa agtattgtga atgattacac 120
 aaaactgtca ggagaaaaac aaacacagtg atgggttggt acgtggctat tggtagcgag 180

catcctataa gaattcagac catgactacg actgacacta aggatggtgc tgggacagtt 240
 gaacaggtga tgagaatagc agataaagga gctgatattg tacggataac agttcaaggg 300
 aagaaagaag ctgatgcttg ttttgagatt aaaaacaccc ttgttcagaa aaattacaac 360
 atactcgtgg tggctgatat tcatttttgct ccctctggtg ctttgcggtg agctgaatgc 420
 tttgataaga ttcgtgtaaa ccctggaaat tttgctgata gacgggctca atttgaaaca 480
 ttagagtaca cagatgatga ctatcagaaa gaacttgagc atattgaaaa ggttttcaca 540
 ccattggttg agaaatgtaa gaaa 564

<210> 36
 <211> 511
 <212> DNA
 <213> Glycine max

<400> 36

aaaccatgga agtctttctg atcgtataat gagctactat ggagactcgc ctaggggaat 60
 ggtagaatct gcttttgaat ttgcaaggat atgccgaaag ttagactatc acaattttgt 120
 tttttctatg aaagcaagca acccagttat catgggttcag gcataaccgt tacttgtggc 180
 tgaaatgtat gtccaaggct gggattatcc attacacttg ggtgttactg aagctggaga 240
 aggtgaggat gggaggatga agtctgcaat aggcattgga actcttcttc aggatggatt 300
 gggtgataca attagggttt ctctcacaga accaccagag gaggagatag acccttgcag 360
 aaggttggca aatcttgga tgatagcttc tgaactccag aagggggtgg aaccttttga 420
 agaaaagcac agacattatt ttgactttca gcgccgatct ggtcaattgc cagtgcataa 480
 agaggggtgag gaggtggatt acagaggtgt a 511

<210> 37
 <211> 498
 <212> DNA
 <213> Glycine max

<220>
 <221> unsure
 <222> (1..498)
 <223> unsure at all n locations

<400> 37

cggaggtggc gtgaatgctt tgataagatt cgtgtaaacc ctggaaattt tgctgataga 60
 cgggctcaat ttgaaacatg agagtggaca naataagact atgagaaaga acttgagcat 120
 attgaaaagg ttttcacacc attggttgag aaatgtaaga aatatgggag agcaatgcgc 180
 attgggacaa accatggaag tctttctgat cgtataatga gctactatgg agactcgcct 240
 aggggaatgg tagaatctgc ttttgaattt gcaaggatat gccgaaagtt agactatcac 300

WO 02/12478

PCT/US01/24335

aattttgttt tttctatgaa agcaagcaac ccagttatca tgggttcaggc ataccgctta 360
 cttgtggctg aaatgtatgt ccaaggctgg gattatccat tacacttggg tgttactgaa 420
 gctggagaag gtgaggatgg gaggatgaag tctgcaatag gcattggaac tcttcttcag 480
 gatggattgg gtgataca 498

<210> 38
 <211> 440
 <212> DNA
 <213> Glycine max

<400> 38

gtagctgaat gctttgataa gattcgtgta aaccctggaa attttgttga tagacgggct 60
 caatttgaaa cattagagta cacagaagaa gactatcata aagaacttga gcatattgaa 120
 aaggttttca caccattggg tgagaaatgt aagaaatatg ggagagcaat gcgcattggg 180
 acaaaccatg gaagtctttc tgatcgtata atgagctact atggagactc gcctagggga 240
 atggtagaat ctgcttttga atttgcaagg atatgccgaa agtttagacta tcacaatttt 300
 gttttttcta tgaaagcaag caaccagtt atcatgggtc aggcataccg cttacttgtg 360
 gctgaaatgt atgttcaagg ctgggattat ccattacact tgggtgttac tgaagctgga 420
 aaaagtgagg atgggaggat 440

<210> 39
 <211> 353
 <212> DNA
 <213> Glycine max

<400> 39

aattcggctc gagaggaact caaatcctgg ccaagatatt gctgaacttc aacctgcac 60
 cccaggaagc cctcttttgg ttcctaggca aaagtattgt gaatcattac acaaaactgt 120
 caggagaaaa acaaacacag tgatggttgg taacgtggct attggtagcg agcatcctat 180
 aagaattcag accatgacta cgactgacac taaggatgtt gctgggacag ttgaacaggt 240
 gatgagaata gcagataaag gagctgatat tgtacggata acagttcaag ggaagaaaga 300
 agctgatgct tgttttgaga ttaaaaacac ccttgttcaa aaaaattaca aca 353

<210> 40
 <211> 577
 <212> DNA
 <213> Glycine max

<400> 40

gatgtttttg tcgtgtattc tttcctatt gcattcagct cactgatttc aattacaaag 60
 tcaattttgt aaatcagagg cagagagagt tgtaaagagc ctctgaattt tgatcacacc 120

acacccttct tctcatctcc accagaaatg gctaccggag ctgctgtgcc aactacgttt 180
 tctaccctca agacatggga ttccagtttg gggtttgcaa aaaacataga ttttgtgaga 240
 gtttccgata tgaagagcat gaaatcttct gcgaggaaaa ggggtgtcaat tatcaggaac 300
 tcaaatcctg gccaagatat tgctgaactt caacctgcat cccaggaag ccctcttttg 360
 gttcctaggc aaaagtattg tgaatcattg cacaaacca tcaggagaaa aacaagcaca 420
 gtaatggttg gtaacgtggc tattggtagc gagcatccta taagaattca gaccatgact 480
 acaactgaca ctaaggatgt tgctgggaca gttgaaccgg tgatgagaat agcagataaa 540
 ggagctgata ttgtacggat aacagttcaa gggaaga 577

<210> 41
 <211> 551
 <212> DNA
 <213> Glycine max
 <400> 41

tggtgctggc tctgatgctg gagcccttct ggtggatggg cttggagatg gacttctttt 60
 ggaagcgcca gacaaggatt ttgaatttat tagaaacact tctttcaatt tgttgcaagg 120
 ctgcagaatg agaaatacaa agacagagta tgtctcatgt ccattcctgtg gcagaacatt 180
 gtttgatctt caagaagtaa gtgcacaaat tcgggagaag acatcacacc tccccggtgt 240
 ttcgattgca atcatgggat gcattgtaaa tggaccaggg gagatggctg atgcagactt 300
 tgggtatgtg ggaggcactc ccgggaagat tgacctctat gttgggaaga ctgtggtgaa 360
 gcgtggaatt gcaatggagc atgcaaccaa tgccttgatc gatctaataa aagaacatgg 420
 acgatgggtg gaccctcctg ccgaggagta aaagcaagag ctttaattttg agattggcat 480
 tcaaggccat agtaagatga gcattgtcat atccaattat tggacacatg taatataagc 540
 atacactcaa t 551

<210> 42
 <211> 869
 <212> DNA
 <213> Glycine max
 <400> 42

gaagcatagt agcatcaatg ccttccttat acagaagact aaaattagca gagtgcattg 60
 ggccaggcgg ttatttgagt acctatccga caattctcta aacttccctg ttattcacca 120
 tattcagttc ccaaattggga ttcacagaga tgacttggtg attggtgctg gttctgatgc 180
 tggagccctt ctggtggatg ggcttggaga tggacttctt ttggaagcgc cagacaagga 240
 ttttgaattt attagaaaca cttctttcaa tttgttgcaa ggctgcagaa tgagaaatac 300
 aaagacagag tatgtctcat gtccatcctg tggcagaaca ttgtttgatc ttcaagaagt 360

aagtgcacaa attcgggaga agacatcaca cctccctggt gtttcgattg caatcatggg 420
atgcattgta aatggaccag gggagatggc tgatgcagac tttgggtatg tgggaggcac 480
tcccgggaag attgacctct atgttgggaa gactgtggtg aagcgtggaa ttgcaatgga 540
gcatgcaacc aatgccttga tcgatctaataaaaagaacat ggacgatggg tggaccctcc 600
tgccgaggag taaaagcaag agcttaattt tgagattggc attcaaggcc atagtaagat 660
gagcattgtc atatccaatt attgtacaca tgtaataataa gataacactc aatgcttaag 720
tttgagccta gttttaagtt ctttttgaga aagatcccaa ttaaagcttg ttgtgaggaa 780
atcgacagct agaacatgta tacagataac agtgtattgc tttgccccat cagccatcaa 840
taataatgag aatctcttag aatagtgcc 869

<210> 43
<211> 291
<212> DNA
<213> Glycine max

<220>
<221> unsure
<222> (1..291)
<223> unsure at all n locations

<400> 43

gangnactca aatcctgggc caagatattg ctgaacttca nccctgcatc cccaggnggc 60
cctcttttgg ttctaggca aaagtattgt gaatcattnc cacaaaactg nccagganaa 120
aaacaaacac agtgatggtt ggtaacgtgg ctattggttag cgagcatcct ataagaattc 180
agaccatgac tacgacngac actaaggatg ttgctgggac agtngaacng gtgatgagaa 240
tagcagataa aggagctgat attgtacgga taacagttca agggaagaaa g 291

<210> 44
<211> 388
<212> DNA
<213> Glycine max

<400> 44

cccggatatat gggttcaggca taccgtttac ttgtggctga aatgtatgtc caaggctggg 60
attatccatt acacttgggt gttactgaag ctggagaagg tgaggatggg aggatgaagt 120
ctgcaattgg cattggaact cttcttcagg atggattggg tgatacaatt agggtttctc 180
tcacagaacc accagaagag gagatagatc cttgcagaag gttggcaaatt cttggaatga 240
gagcttctga actccagaag ggggtggaac cttttgaaga aaagcacaga cattattttg 300
acttccagcg ccgatctggt caattgccag tgcaaaaaga gggtgaggag gtggattaca 360
gaggtgcact gcaccgtgac gggtctgt 388

WO 02/12478

PCT/US01/24335

<210> 45
 <211> 211
 <212> DNA
 <213> Glycine max

<400> 45

cccggttatc atggcgcagg cataccgctt acttgtggct gaaatgtatg tccaaggctg 60
 ggattatcca ttacacttgg gtgttactga agctggagga ggtgaggatg acaggatgaa 120
 gtctgcaatt ggcatgggaa ctcttcttca ggatggattg ggtgatacaa ttaggggtgtc 180
 tcgcacagaa ccaccagaag aggagataga t 211

<210> 46
 <211> 276
 <212> DNA
 <213> Glycine max

<400> 46

tgggcttgga gatggactac ttttgggaagc cccggacaag gatttttgaat ttattagaaa 60
 cacttctttc aatttggttc aaggctgcag aatgagaaat acaaagacag agtatgtctc 120
 atgtccatcc tgtggcagaa cattgtttga tcttcaagaa gtaagtgcac aaattcggga 180
 gaagacatca cacctccctg gtgtttcgat tgcaatcatg ggatgcattg taaatggacc 240
 aggggagatg gctgatgcag actttgggta tgtggg 276

<210> 47
 <211> 399
 <212> DNA
 <213> Brassica napus

<400> 47

cccacgcgtc cgcagggatt cacagggacg agttggtgat ccacgcaggg acatacgtg 60
 gggcacttct agtggatgga cttggagatg gtgtaatgct agaagcacct gatcaagact 120
 tcgagtttct taggaacact tctttcaact tgttacaagg ctgcaggatg cgtaacacca 180
 agacggaata cgtatcgtgc ccgtcttggtg gaagaactct gttcgacttg caagaaatca 240
 gcgctgagat cagagaaaag acttcgcatt tgcctggcgt ttcgattgca ataatggggt 300
 gcattgtgaa tggacctggc gaaatggctg atgctgattt cggttatgta ggcggttctc 360
 ccgggaaaat cgacctttac gttggaaaga cggtggtca 399

WO 02/12478

PCT/US01/24335

<210> 48
 <211> 740
 <212> PRT
 <213> Arabidopsis thaliana

<400> 48

```

Met Ala Thr Gly Val Leu Pro Ala Pro Val Ser Gly Ile Lys Ile Pro
1          5          10          15

Asp Ser Lys Val Gly Phe Gly Lys Ser Met Asn Leu Val Arg Ile Cys
          20          25          30

Asp Val Arg Ser Leu Arg Ser Ala Arg Arg Arg Val Ser Val Ile Arg
          35          40          45

Asn Ser Asn Gln Gly Ser Asp Leu Ala Glu Leu Gln Pro Ala Ser Glu
          50          55          60

Gly Ser Pro Leu Leu Val Pro Arg Gln Lys Tyr Cys Glu Ser Leu His
65          70          75          80

Lys Thr Val Arg Arg Lys Thr Arg Thr Val Met Val Gly Asn Val Ala
          85          90          95

Leu Gly Ser Glu His Pro Ile Arg Ile Gln Thr Met Thr Thr Ser Asp
          100          105          110

Thr Lys Asp Ile Thr Gly Thr Val Asp Glu Val Met Arg Ile Ala Asp
          115          120          125

Lys Gly Ala Asp Ile Val Arg Ile Thr Val Gln Gly Lys Lys Glu Ala
          130          135          140

Asp Ala Cys Phe Glu Ile Lys Asp Lys Leu Val Gln Leu Asn Tyr Asn
145          150          155          160

Ile Pro Leu Val Ala Asp Ile His Phe Ala Pro Thr Val Ala Leu Arg
          165          170          175

Val Ala Glu Cys Phe Asp Lys Ile Arg Val Asn Pro Gly Asn Phe Ala
          180          185          190

Asp Arg Arg Ala Gln Phe Glu Thr Ile Asp Tyr Thr Glu Asp Glu Tyr
          195          200          205

Gln Lys Glu Leu Gln His Ile Glu Gln Val Phe Thr Pro Leu Val Glu
          210          215          220

Lys Cys Lys Lys Tyr Gly Arg Ala Met Arg Ile Gly Thr Asn His Gly
225          230          235          240

Ser Leu Ser Asp Arg Ile Met Ser Tyr Tyr Gly Asp Ser Pro Arg Gly
          245          250          255

Met Val Glu Ser Ala Phe Glu Phe Ala Arg Ile Cys Arg Lys Leu Asp
          260          265          270

Tyr His Asn Phe Val Phe Ser Met Lys Ala Ser Asn Pro Val Ile Met
          275          280          285

Val Gln Ala Tyr Arg Leu Leu Val Ala Glu Met Tyr Val His Gly Trp
          290          295          300

```

Asp Tyr Pro Leu His Leu Gly Val Thr Glu Ala Gly Glu Gly Glu Asp
 305 310 315 320
 Gly Arg Met Lys Ser Ala Ile Gly Ile Gly Thr Leu Leu Gln Asp Gly
 325 330 335
 Leu Gly Asp Thr Ile Arg Val Ser Leu Thr Glu Pro Pro Glu Glu Glu
 340 345 350
 Ile Asp Pro Cys Arg Arg Leu Ala Asn Leu Gly Thr Lys Ala Ala Lys
 355 360 365
 Leu Gln Gln Gly Ala Pro Phe Glu Glu Lys His Arg His Tyr Phe Asp
 370 375 380
 Phe Gln Arg Arg Thr Gly Asp Leu Pro Val Gln Lys Glu Gly Glu Glu
 385 390 395 400
 Val Asp Tyr Arg Asn Val Leu His Arg Asp Gly Ser Val Leu Met Ser
 405 410 415
 Ile Ser Leu Asp Gln Leu Lys Ala Pro Glu Leu Leu Tyr Arg Ser Leu
 420 425 430
 Ala Thr Lys Leu Val Val Gly Met Pro Phe Lys Asp Leu Ala Thr Val
 435 440 445
 Asp Ser Ile Leu Leu Arg Glu Leu Pro Pro Val Asp Asp Gln Val Ala
 450 455 460
 Arg Leu Ala Leu Lys Arg Leu Ile Asp Val Ser Met Gly Val Ile Ala
 465 470 475 480
 Pro Leu Ser Glu Gln Leu Thr Lys Pro Leu Pro Asn Ala Met Val Leu
 485 490 495
 Val Asn Leu Lys Glu Leu Ser Gly Gly Ala Tyr Lys Leu Leu Pro Glu
 500 505 510
 Gly Thr Arg Leu Val Val Ser Leu Arg Gly Asp Glu Pro Tyr Glu Glu
 515 520 525
 Leu Glu Ile Leu Lys Asn Ile Asp Ala Thr Met Ile Leu His Asp Val
 530 535 540
 Pro Phe Thr Glu Asp Lys Val Ser Arg Val His Ala Ala Arg Arg Leu
 545 550 555 560
 Phe Glu Phe Leu Ser Glu Asn Ser Val Asn Phe Pro Val Ile His His
 565 570 575
 Ile Asn Phe Pro Thr Gly Ile His Arg Asp Glu Leu Val Ile His Ala
 580 585 590
 Gly Thr Tyr Ala Gly Gly Leu Leu Val Asp Gly Leu Gly Asp Gly Val
 595 600 605
 Met Leu Glu Ala Pro Asp Gln Asp Phe Asp Phe Leu Arg Asn Thr Ser
 610 615 620
 Phe Asn Leu Leu Gln Gly Cys Arg Met Arg Asn Thr Lys Thr Glu Tyr
 625 630 635 640

Val Ser Cys Pro Ser Cys Gly Arg Thr Leu Phe Asp Leu Gln Glu Ile
645 650 655

Ser Ala Glu Ile Arg Glu Lys Thr Ser His Leu Pro Gly Val Ser Ile
660 665 670

Ala Ile Met Gly Cys Ile Val Asn Gly Pro Gly Glu Met Ala Asp Ala
675 680 685

Asp Phe Gly Tyr Val Gly Gly Ser Pro Gly Lys Ile Asp Leu Tyr Val
690 695 700

Gly Lys Thr Val Val Lys Arg Gly Ile Ala Met Thr Glu Ala Thr Asp
705 710 715 720

Ala Leu Ile Gly Leu Ile Lys Glu His Gly Arg Trp Val Asp Pro Pro
725 730 735

Val Ala Asp Glu
740

<210> 49
<211> 603
<212> PRT
<213> Oryza sativa
<400> 49

Met Val Gly Asn Val Pro Leu Gly Ser Asp His Pro Ile Arg Ile Gln
1 5 10 15

Thr Met Thr Thr Ser Asp Thr Lys Asp Val Ala Lys Thr Val Glu Glu
20 25 30

Val Met Arg Ile Ala Asp Lys Gly Ala Asp Phe Val Arg Ile Thr Val
35 40 45

Gln Gly Arg Lys Glu Ala Asp Ala Cys Phe Glu Ile Lys Asn Thr Leu
50 55 60

Val Gln Lys Asn Tyr Asn Ile Pro Leu Val Ala Asp Ile His Phe Ala
65 70 75 80

Pro Thr Val Ala Leu Arg Val Ala Glu Cys Phe Asp Lys Ile Arg Val
85 90 95

Asn Pro Gly Asn Phe Ala Asp Arg Arg Ala Gln Phe Glu Gln Leu Glu
100 105 110

Tyr Thr Glu Asp Asp Tyr Gln Lys Glu Leu Glu His Ile Glu Lys Val
115 120 125

Pro Asn Ile Ser Leu Phe Ser Val Asn Leu Val Phe Ser Pro Leu Val
130 135 140

Glu Lys Cys Lys Gln Tyr Gly Arg Ala Met Arg Ile Gly Thr Asn His
145 150 155 160

Gly Ser Leu Ser Asp Arg Ile Met Ser Tyr Tyr Gly Asp Ser Pro Arg
165 170 175

Gly Met Val Glu Ser Ala Leu Glu Phe Ala Arg Ile Cys Arg Lys Leu
180 185 190

Asp	Phe	His	Asn	Phe	Val	Phe	Ser	Met	Lys	Ala	Ser	Asn	Pro	Val	Ile	195	200	205	
Met	Val	Gln	Ala	Tyr	Arg	Leu	Leu	Val	Ala	Glu	Met	Tyr	Asn	Leu	Gly	210	215	220	
Trp	Asp	Tyr	Pro	Leu	His	Leu	Gly	Val	Thr	Glu	Ala	Gly	Glu	Gly	Glu	225	230	235	240
Asp	Gly	Arg	Met	Lys	Ser	Ala	Ile	Gly	Ile	Gly	Thr	Leu	Leu	Met	Asp	245	250	255	
Gly	Leu	Gly	Asp	Thr	Ile	Arg	Val	Ser	Leu	Thr	Glu	Pro	Pro	Glu	Glu	260	265	270	
Glu	Ile	Asp	Pro	Cys	Arg	Arg	Leu	Ala	Asn	Leu	Gly	Thr	His	Ala	Ala	275	280	285	
Asp	Leu	Gln	Ile	Gly	Val	Ala	Pro	Phe	Glu	Glu	Lys	His	Arg	Arg	Tyr	290	295	300	
Phe	Asp	Phe	Gln	Arg	Arg	Ser	Gly	Gln	Leu	Pro	Leu	Gln	Lys	Glu	Ala	305	310	315	320
Pro	Glu	Leu	Leu	Tyr	Arg	Ser	Leu	Ala	Ala	Lys	Leu	Val	Val	Gly	Met	325	330	335	
Pro	Phe	Lys	Asp	Leu	Ala	Thr	Val	Asp	Ser	Ile	Leu	Leu	Lys	Glu	Leu	340	345	350	
Pro	Pro	Val	Glu	Asp	Ala	Gln	Ala	Arg	Leu	Ala	Leu	Lys	Arg	Leu	Val	355	360	365	
Asp	Ile	Ser	Met	Gly	Val	Leu	Thr	Pro	Leu	Ser	Glu	Gln	Leu	Thr	Lys	370	375	380	
Pro	Leu	Pro	His	Ala	Ile	Ala	Leu	Val	Asn	Val	Asp	Glu	Leu	Ser	Ser	385	390	395	400
Gly	Ala	His	Lys	Leu	Leu	Pro	Glu	Gly	Thr	Arg	Leu	Ala	Val	Thr	Leu	405	410	415	
Arg	Gly	Asp	Glu	Ser	Tyr	Glu	Gln	Leu	Asp	Leu	Leu	Lys	Gly	Val	Asp	420	425	430	
Asp	Ile	Thr	Met	Leu	Leu	His	Ser	Val	Pro	Tyr	Gly	Glu	Glu	Lys	Thr	435	440	445	
Gly	Arg	Val	His	Ala	Ala	Arg	Arg	Leu	Phe	Glu	Tyr	Leu	Glu	Thr	Asn	450	455	460	
Gly	Leu	Asn	Phe	Pro	Val	Ile	His	His	Ile	Glu	Phe	Pro	Lys	Ser	Val	465	470	475	480
Asn	Arg	Asp	Asp	Leu	Val	Ile	Gly	Ala	Gly	Ala	Asn	Val	Gly	Ala	Leu	485	490	495	
Leu	Val	Asp	Gly	Leu	Gly	Asp	Gly	Val	Leu	Leu	Glu	Ala	Ala	Asp	Gln	500	505	510	
Glu	Phe	Glu	Phe	Leu	Arg	Asp	Thr	Ser	Phe	Asn	Leu	Leu	Gln	Gly	Cys	515	520	525	

Arg Met Arg Asn Thr Lys Thr Ile Ala Ile Met Gly Cys Ile Val Asn
 530 535 540
 Gly Pro Gly Glu Met Ala Asp Ala Asp Phe Gly Tyr Val Gly Gly Ala
 545 550 555 560
 Pro Gly Lys Ile Asp Leu Tyr Val Gly Lys Thr Val Val Gln Arg Gly
 565 570 575
 Ile Ala Met Glu Gly Ala Thr Asp Ala Leu Ile Gln Leu Ile Lys Asp
 580 585 590
 His Gly Arg Trp Val Asp Pro Pro Val Glu Glu
 595 600

<210> 50
 <211> 372
 <212> PRT
 <213> Escherichia coli
 <400> 50

Met His Asn Gln Ala Pro Ile Gln Arg Arg Lys Ser Thr Arg Ile Tyr
 1 5 10 15
 Val Gly Asn Val Pro Ile Gly Asp Gly Ala Pro Ile Ala Val Gln Ser
 20 25 30
 Met Thr Asn Thr Arg Thr Thr Asp Val Glu Ala Thr Val Asn Gln Ile
 35 40 45
 Lys Ala Leu Glu Arg Val Gly Ala Asp Ile Val Arg Val Ser Val Pro
 50 55 60
 Thr Met Asp Ala Ala Glu Ala Phe Lys Leu Ile Lys Gln Gln Val Asn
 65 70 75 80
 Val Pro Leu Val Ala Asp Ile His Phe Asp Tyr Arg Ile Ala Leu Lys
 85 90 95
 Val Ala Glu Tyr Gly Val Asp Cys Leu Arg Ile Asn Pro Gly Asn Ile
 100 105 110
 Gly Asn Glu Glu Arg Ile Arg Met Val Val Asp Cys Ala Arg Asp Lys
 115 120 125
 Asn Ile Pro Ile Arg Ile Gly Val Asn Ala Gly Ser Leu Glu Lys Asp
 130 135 140
 Leu Gln Glu Lys Tyr Gly Glu Pro Thr Pro Gln Ala Leu Leu Glu Ser
 145 150 155 160
 Ala Met Arg His Val Asp His Leu Asp Arg Leu Asn Phe Asp Gln Phe
 165 170 175
 Lys Val Ser Val Lys Ala Ser Asp Val Phe Leu Ala Val Glu Ser Tyr
 180 185 190
 Arg Leu Leu Ala Lys Gln Ile Asp Gln Pro Leu His Leu Gly Ile Thr
 195 200 205
 Glu Ala Gly Gly Ala Arg Ser Gly Ala Val Lys Ser Ala Ile Gly Leu
 210 215 220

Gly Leu Leu Leu Ser Glu Gly Ile Gly Asp Thr Leu Arg Val Ser Leu
 225 230 235 240
 Ala Ala Asp Pro Val Glu Glu Ile Lys Val Gly Phe Asp Ile Leu Lys
 245 250 255
 Ser Leu Arg Ile Arg Ser Arg Gly Ile Asn Phe Ile Ala Cys Pro Thr
 260 265 270
 Cys Ser Arg Gln Glu Phe Asp Val Ile Gly Thr Val Asn Ala Leu Glu
 275 280 285
 Gln Arg Leu Glu Asp Ile Ile Thr Pro Met Asp Val Ser Ile Ile Gly
 290 295 300
 Cys Val Val Asn Gly Pro Gly Glu Ala Leu Val Ser Thr Leu Gly Val
 305 310 315 320
 Thr Gly Gly Asn Lys Lys Ser Gly Leu Tyr Glu Asp Gly Val Arg Lys
 325 330 335
 Asp Arg Leu Asp Asn Asn Asp Met Ile Asp Gln Leu Glu Ala Arg Ile
 340 345 350
 Arg Ala Lys Ala Ser Gln Leu Asp Glu Ala Arg Arg Ile Asp Val Gln
 355 360 365
 Gln Val Glu Lys
 370

<210> 51
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named CINCO

<400> 51

cgctgcccag aatggacctc cctag

25

<210> 52
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named SEIS

<400> 52

cagccgcgtt ttgacttgaa acgtgc

26

<210> 53
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named MPD-Nde5'

<400> 53

gccatatgac cgtttacaca gcatccg

27

<210> 54
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named MPD-Eco3'

<400> 54

tcgaattctc attattcctt tggtagacca gtctt

35

<210> 55
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named hPMK1

<400> 55

tggttaacat atggccccgc tgggaggcgc

30

<210> 56
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named hPMK4

<400> 56

aggttaactc aattaaagtc tggagcggat aaattctatc

40

<210> 57
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named UNO

<400> 57

cgggcctcgt ttggctgtcg cactg

25

<210> 58
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named DOS

<400> 58

cgcggggtgga aggaccttgt ggagg

25

<210> 59
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named MK-Hpa5'

<400> 59

aagttaacat atgtcattac cgttcttaac ttc

33

<210> 60
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named MK-Hpa3'

<400> 60

cggttaactc attatgaagt ccatggtaaa ttcg

34

<210> 61
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named idi5X

<400> 61

cccctcgaga ttatgcaaac ggaacacgtc

30

<210> 62
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named idi3X

<400> 62

ggctcgagtt atttaagctg ggtaaata gca g

31

<210> 63
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pBAD-mut1

<400> 63

ctgagagtgc accatctgcg gtgtgaaata cc 32

<210> 64
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pBAD-Link1

<400> 64

aattctaagg aggtttaaac taaggaggta cgtaaggagg 40

<210> 65
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pBAD-Link2

<400> 65

tcgacctcct tacgtacctc cttagtttaa acctccttag 40

<210> 66
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pBAD-D2

<400> 66

tcatactccc gccattcaga g 21

<210> 67
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pBAD-U3

<400> 67

ccgccaaaac agccaagctt g 21

<210> 68
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pRS-L1

<400> 68

gatccgttta aacgcccggg cggccgcg

28

<210> 69
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named pRS-L2

<400> 69

aattcgcggc cgcccggggt tttaaacg

28

<210> 70
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named 1PE

<400> 70

cgcggtgtgg gtgagcatga tg

22

<210> 71
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named 22PE

<400> 71

aaatctcccg ggttaccgt ctgttactgc

30

<210> 72
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named 3PE

<400> 72

gcgtttaaac tggacgaagc gcgtcgaatt gac

33

<210> 73
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named 4PE

<400> 73

tgcacgaccg cccagttggt cc 22

<210> 74
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named CAT1

<400> 74

gagtccgaat aaatacctgt g 21

<210> 75
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named CAT4

<400> 75

ccgaatttct gccattcatc c 21

<210> 76
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named OPE

<400> 76

tgggctttgt cacgagcaca c 21

<210> 77
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed primer named 5PE

<400> 77

ggcccatagc aaaaccgaca g 21

WO 02/12478

PCT/US01/24335

<210> 78

<211> 372

<212> PRT

<213> Escherichia coli

<400> 78

```

Met His Asn Gln Ala Pro Ile Gln Arg Arg Lys Ser Thr Arg Ile Tyr
 1              5              10              15

Val Gly Asn Val Pro Ile Gly Asp Gly Ala Pro Ile Ala Val Gln Ser
      20              25              30

Met Thr Asn Thr Arg Thr Thr Asp Val Glu Ala Thr Val Asn Gln Ile
      35              40              45

Lys Ala Leu Glu Arg Val Gly Ala Asp Ile Val Arg Val Ser Val Pro
      50              55              60

Thr Met Asp Ala Ala Glu Ala Phe Lys Leu Ile Lys Gln Gln Val Asn
 65              70              75              80

Val Pro Leu Val Ala Asp Ile His Phe Asp Tyr Arg Ile Ala Leu Lys
      85              90              95

Val Ala Glu Tyr Gly Val Asp Cys Leu Arg Ile Asn Pro Gly Asn Ile
      100              105              110

Gly Asn Glu Glu Arg Ile Arg Met Val Val Asp Cys Ala Arg Asp Lys
      115              120              125

Asn Ile Pro Ile Arg Ile Gly Val Asn Ala Gly Ser Leu Glu Lys Asp
      130              135              140

Leu Gln Glu Lys Tyr Gly Glu Pro Thr Pro Gln Ala Leu Leu Glu Ser
      145              150              155              160

Ala Met Arg His Val Asp His Leu Asp Arg Leu Asn Phe Asp Gln Phe
      165              170              175

Lys Val Ser Val Lys Ala Ser Asp Val Phe Leu Ala Val Glu Ser Tyr
      180              185              190

Arg Leu Leu Ala Lys Gln Ile Asp Gln Pro Leu His Leu Gly Ile Thr
      195              200              205

Glu Ala Gly Gly Ala Arg Ser Gly Ala Val Lys Ser Ala Ile Gly Leu
      210              215              220

Gly Leu Leu Leu Ser Glu Gly Ile Gly Asp Thr Leu Arg Val Ser Leu
      225              230              235              240

Ala Ala Asp Pro Val Glu Glu Ile Lys Val Gly Phe Asp Ile Leu Lys
      245              250              255

Ser Leu Arg Ile Arg Ser Arg Gly Ile Asn Phe Ile Ala Cys Pro Thr
      260              265              270

Cys Ser Arg Gln Glu Phe Asp Val Ile Gly Thr Val Asn Ala Leu Glu
      275              280              285

Gln Arg Leu Glu Asp Ile Ile Thr Pro Met Asp Val Ser Ile Ile Gly
      290              295              300

```

Cys Val Val Asn Gly Pro Gly Glu Ala Leu Val Ser Thr Leu Gly Val
305 310 315 320

Thr Gly Gly Asn Lys Lys Ser Gly Leu Tyr Glu Asp Gly Val Arg Lys
325 330 335

Asp Arg Leu Asp Asn Asn Asp Met Ile Asp Gln Leu Glu Ala Arg Ile
340 345 350

Arg Ala Lys Ala Ser Gln Leu Asp Glu Ala Arg Arg Ile Asp Val Gln
355 360 365

Gln Val Glu Lys
370

<210> 79

<211> 740

<212> PRT

<213> Arabidopsis thaliana

<400> 79

Met Ala Thr Gly Val Leu Pro Ala Pro Val Ser Gly Ile Lys Ile Pro
1 5 10 15

Asp Ser Lys Val Gly Phe Gly Lys Ser Met Asn Leu Val Arg Ile Cys
20 25 30

Asp Val Arg Ser Leu Arg Ser Ala Arg Arg Arg Val Ser Val Ile Arg
35 40 45

Asn Ser Asn Gln Gly Ser Asp Leu Ala Glu Leu Gln Pro Ala Ser Glu
50 55 60

Gly Ser Pro Leu Leu Val Pro Arg Gln Lys Tyr Cys Glu Ser Leu His
65 70 75 80

Lys Thr Val Arg Arg Lys Thr Arg Thr Val Met Val Gly Asn Val Ala
85 90 95

Leu Gly Ser Glu His Pro Ile Arg Ile Gln Thr Met Thr Thr Ser Asp
100 105 110

Thr Lys Asp Ile Thr Gly Thr Val Asp Glu Val Met Arg Ile Ala Asp
115 120 125

Lys Gly Ala Asp Ile Val Arg Ile Thr Val Gln Gly Lys Lys Glu Ala
130 135 140

Asp Ala Cys Phe Glu Ile Lys Asp Lys Leu Val Gln Leu Asn Tyr Asn
145 150 155 160

Ile Pro Leu Val Ala Asp Ile His Phe Ala Pro Thr Val Ala Leu Arg
165 170 175

Val Ala Glu Cys Phe Asp Lys Ile Arg Val Asn Pro Gly Asn Phe Ala
180 185 190

Asp Arg Arg Ala Gln Phe Glu Thr Ile Asp Tyr Thr Glu Asp Glu Tyr
195 200 205

Gln Lys Glu Leu Gln His Ile Glu Gln Val Phe Thr Pro Leu Val Glu
210 215 220

Lys Cys Lys Lys Tyr Gly Arg Ala Met Arg Ile Gly Thr Asn His Gly
 225 230 235 240
 Ser Leu Ser Asp Arg Ile Met Ser Tyr Tyr Gly Asp Ser Pro Arg Gly
 245 250 255
 Met Val Glu Ser Ala Phe Glu Phe Ala Arg Ile Cys Arg Lys Leu Asp
 260 265 270
 Tyr His Asn Phe Val Phe Ser Met Lys Ala Ser Asn Pro Val Ile Met
 275 280 285
 Val Gln Ala Tyr Arg Leu Leu Val Ala Glu Met Tyr Val His Gly Trp
 290 295 300
 Asp Tyr Pro Leu His Leu Gly Val Thr Glu Ala Gly Glu Gly Glu Asp
 305 310 315 320
 Gly Arg Met Lys Ser Ala Ile Gly Ile Gly Thr Leu Leu Gln Asp Gly
 325 330 335
 Leu Gly Asp Thr Ile Arg Val Ser Leu Thr Glu Pro Pro Glu Glu Glu
 340 345 350
 Ile Asp Pro Cys Arg Arg Leu Ala Asn Leu Gly Thr Lys Ala Ala Lys
 355 360 365
 Leu Gln Gln Gly Ala Pro Phe Glu Glu Lys His Arg His Tyr Phe Asp
 370 375 380
 Phe Gln Arg Arg Thr Gly Asp Leu Pro Val Gln Lys Glu Gly Glu Glu
 385 390 395 400
 Val Asp Tyr Arg Asn Val Leu His Arg Asp Gly Ser Val Leu Met Ser
 405 410 415
 Ile Ser Leu Asp Gln Leu Lys Ala Pro Glu Leu Leu Tyr Arg Ser Leu
 420 425 430
 Ala Thr Lys Leu Val Val Gly Met Pro Phe Lys Asp Leu Ala Thr Val
 435 440 445
 Asp Ser Ile Leu Leu Arg Glu Leu Pro Pro Val Asp Asp Gln Val Ala
 450 455 460
 Arg Leu Ala Leu Lys Arg Leu Ile Asp Val Ser Met Gly Val Ile Ala
 465 470 475 480
 Pro Leu Ser Glu Gln Leu Thr Lys Pro Leu Pro Asn Ala Met Val Leu
 485 490 495
 Val Asn Leu Lys Glu Leu Ser Gly Gly Ala Tyr Lys Leu Leu Pro Glu
 500 505 510
 Gly Thr Arg Leu Val Val Ser Leu Arg Gly Asp Glu Pro Tyr Glu Glu
 515 520 525
 Leu Glu Ile Leu Lys Asn Ile Asp Ala Thr Met Ile Leu His Asp Val
 530 535 540
 Pro Phe Thr Glu Asp Lys Val Ser Arg Val His Ala Ala Arg Arg Leu
 545 550 555 560

WO 02/12478

PCT/US01/24335

Phe Glu Phe Leu Ser Glu Asn Ser Val Asn Phe Pro Val Ile His His
 565 570 575
 Ile Asn Phe Pro Thr Gly Ile His Arg Asp Glu Leu Val Ile His Ala
 580 585 590
 Gly Thr Tyr Ala Gly Gly Leu Leu Val Asp Gly Leu Gly Asp Gly Val
 595 600 605
 Met Leu Glu Ala Pro Asp Gln Asp Phe Asp Phe Leu Arg Asn Thr Ser
 610 615 620
 Phe Asn Leu Leu Gln Gly Cys Arg Met Arg Asn Thr Lys Thr Glu Tyr
 625 630 635 640
 Val Ser Cys Pro Ser Cys Gly Arg Thr Leu Phe Asp Leu Gln Glu Ile
 645 650 655
 Ser Ala Glu Ile Arg Glu Lys Thr Ser His Leu Pro Gly Val Ser Ile
 660 665 670
 Ala Ile Met Gly Cys Ile Val Asn Gly Pro Gly Glu Met Ala Asp Ala
 675 680 685
 Asp Phe Gly Tyr Val Gly Gly Ser Pro Gly Lys Ile Asp Leu Tyr Val
 690 695 700
 Gly Lys Thr Val Val Lys Arg Gly Ile Ala Met Thr Glu Ala Thr Asp
 705 710 715 720
 Ala Leu Ile Gly Leu Ile Lys Glu His Gly Arg Trp Val Asp Pro Pro
 725 730 735
 Val Ala Asp Glu
 740

<210> 80
 <211> 155
 <212> LLA
 <213> Arabidopsis thaliana

<400> 80

aaaaatcgga aaaatggcga ctggagtatt gccagctccg gtttctggga tcaagatacc 60
 ggattcgaaa gtcgggtttg gtaaaagcat gaatcttggtg agaatttggtg atgtaggag 120
 tctaagatct gctgatgagt agatttcata aaagt 155

<210> 81
 <211> 42
 <212> PRT
 <213> Arabidopsis thaliana

<400> 81

Met Ala Thr Gly Val Leu Pro Ala Pro Val Ser Gly Ile Lys Ile Pro
 1 5 10 15
 Asp Ser Lys Val Gly Phe Gly Lys Ser Met Asn Leu Val Arg Ile Cys
 20 25 30
 Asp Val Arg Ser Leu Arg Ser Ala Asp Glu
 35 40

WO 02/12478

PCT/US01/24335

<210> 82
<211> 45
<212> DNA
<213> Arabidopsis thaliana

<400> 82

atgagaggat cg caycayca ycaycaycay cayggatccg catgc 45

<210> 83
<211> 12
<212> PRT
<213> Arabidopsis thaliana

<400> 83

Met Arg Gly Ser His His His His His His Gly Ser
1 5 10

<210> 84
<211> 59
<212> DNA
<213> Arabidopsis thaliana

<400> 84
atgagaggat cg caycayca ycaycaycay ggatctgctg atgagtagat ttcgcatgc 59

<210> 85
<211> 15
<212> PRT
<213> Arabidopsis thaliana

<400> 85

Met Arg Gly Ser His His His His His His Gly Ser Ala Asp Glu
1 5 10 15

The embodiments of the present invention for which an exclusive property or privilege is claimed are defined as follows:

1. A nucleic acid molecule comprising as operably linked components:
 - (A) a promoter; and
 - (B) a heterologous nucleic acid molecule that
 - (i) encodes an amino acid sequence of SEQ ID NO:50, or
 - (ii) has greater than 85% identity to a nucleic acid sequence of SEQ ID NO:3 and encodes a protein having the activity of SEQ ID NO:50, or
 - (iii) comprises a nucleic acid that hybridizes, under conditions of about 0.2X to about 2.0X SSC at about 55°C-70°C, to the complement of SEQ ID NO:3 and encodes a protein having the activity of SEQ ID NO:50,


wherein said nucleic acid molecule is operably linked to a sequence encoding a chloroplast transit peptide, and wherein said promoter is a seed-specific promoter.

2. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule comprises the nucleic acid sequence of SEQ ID NO:3.
3. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule is substantially purified.
4. The nucleic acid molecule of claim 1, wherein said heterologous nucleic acid molecule hybridizes under moderate stringency conditions of 2.0 x SSC at 65°C, to the complement of a nucleic acid sequence of SEQ ID NO:3 and encodes a protein having the activity of SEQ ID NO:50.

5. A substantially purified nucleic acid molecule that encodes a GCPE protein, wherein said nucleic acid molecule comprises a nucleic acid sequence of SEQ ID NO:3 or a nucleic acid that hybridizes, under conditions of about 0.2X to about 2.0X SSC at about 55°C-70°C to the complement of SEQ ID NO:3 and encodes a protein having the activity of SEQ ID NO:50, and a seed-specific promoter.
6. The nucleic acid molecule according to claim 1, wherein said nucleic acid molecule is a recombinant nucleic acid molecule.
7. The recombinant nucleic acid molecule of claim 6, wherein the seed-specific promoter is a napin promoter.
8. A recombinant nucleic acid molecule comprising as operably linked components: (A) an exogenous promoter; and (B) a nucleic acid sequence of SEQ ID NO:3 or a nucleic acid that hybridizes, under conditions of about 0.2X to about 2.0X SSC at 55°C-70°C to the complement of SEQ ID NO:3, wherein the nucleic acid encodes a GCPE protein having the same biological activity as the GCPE protein corresponding to SEQ ID NO:50, and wherein the promoter is a seed-specific promoter.
9. The recombinant nucleic acid molecule of claim 8, wherein the seed-specific promoter is a napin promoter.
10. The recombinant nucleic acid molecule of claim 8, wherein said nucleic acid molecule further comprises a second nucleic acid sequence that encodes at least one methyl-D-erythritol phosphate (MEP) pathway protein.
11. The recombinant nucleic acid molecule of claim 10, wherein said at least one MEP pathway protein comprises a yfgA protein or a yfgB protein.

12. A recombinant nucleic acid molecule comprising as operably linked components: (A) a promoter that functions in a plant cell to cause production of an mRNA molecule; and (B) a nucleic acid sequence that has greater than 85% identity to a nucleic acid sequence of SEQ ID NO:3 or a nucleic acid that hybridizes, under conditions of about 0.2X to about 2.0X SSC at about 55°C-70°C to the complement of SEQ ID NO:3 and encodes a protein having the same biological activity of SEQ ID NO:50, and wherein said promoter is a seed-specific promoter.
13. The recombinant nucleic acid molecule of claim 12, wherein the conditions are moderate stringency conditions of 2.0 x SSC at about 65°C.
14. A transformed cell comprising the recombinant nucleic acid molecule of claim 6.
15. A transformed cell comprising the recombinant nucleic acid molecule of claim 8.
16. The transformed cell of claim 15, wherein the cell is selected from the group consisting of a bacterial cell, a mammalian cell, an insect cell, a plant cell, and a fungal cell.
17. The transformed cell of claim 16, wherein the bacterial cell is *Agrobacterium tumefaciens*.
18. The transformed cell of claim 14, wherein the transformed cell is a transgenic plant cell.
19. The transformed cell of claim 15, wherein the transformed cell is a transgenic plant cell.
20. The transgenic plant cell of claim 19, wherein said transgenic plant cell is selected from the group consisting of *Brassica campestris*, *Brassica napus*, canola, castor bean, coconut, cotton, crambe, linseed, maize, mustard, oil palm, peanut, rapeseed, rice, safflower, sesame, soybean, sunflower, and wheat plant cells.
21. The transgenic plant cell of claim 19, wherein said transgenic plant cell is selected from the group consisting of coconut, crambe, maize, oil palm, peanut, rapeseed, safflower, sesame, soybean, and sunflower plant cells.

22. A transgenic plant cell comprising a nucleic acid molecule that encodes a GCPE protein, wherein said nucleic acid molecule comprises a promoter operably linked to a heterologous nucleic acid sequence of SEQ ID NO:3 or a nucleic acid that hybridizes, under conditions of about 0.2X to about 2.0X SSC at 55°C-70°C to the complement of SEQ ID NO:3, wherein said promoter is a seed-specific promoter.
23. The transgenic plant cell of claim 22, wherein the seed-specific promoter is selected from the group consisting of napin, phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean a' subunit of b-conglycinin (soy 7s), and oleosin promoters.
24. Use of a transgenic plant having the plant cell of claim 19 to produce an oil.
25. Use of a seed of a transgenic plant having the plant cell of claim 19 to produce a meal.
26. Use of a seed of a transgenic plant having the plant cell of claim 18 to produce an oil.
27. Use of a transgenic plant having the plant cell of claim 22 to produce a feedstock.

**Figure 1**




FIGURE 2




FIGURE 3




Figure 4

pZL1 (135H1):

... [SalI]aaaaatcg(...) gaaaaATGGCGACTGGAGTATTGCCAGCTCCGGTTTCTGGGATCAAG

M A T G V L P A P V S G I K

ATACCGGATTGCGAAAGTCGGGTTTGGTAAAGCATGAATCTTGTGAGAATTTGTGATGTTAGGAGTCTA

I P D S K V G F G K S M N L V R I C D V R S L

BglII

AGATCTGCT(...) GATGAGTAGatttc(...) ataaaagt [NotI] [XbaI] [BamHI] [HindIII] [SphI]...

R S A (...) D E *

pQE30: ...[prom. T5][2x operator lac][RBS]ATGAGAGGATCG[6xHis]GGATCCGCATGC...

pQE-AGH: BamHI/BglII SphI
 ... [RBS]ATGAGAGGATCG[6xHis]GGATCTGCT(...)GATGAGTAGatttc(...)GCATGC..
 M R G S HHHHHH G S A (...) D E *

Figure 5



Ecgcpe : MHNQAPIQRRKSTRIYVGNVPIGDG : 25
Atgcpe : MATGVLPAPVSGIKIPDSKVGF GKSMNLVRI CDVRSLSARRRVSVIRNSNQGSDLAELQPASEGSPLLVP RQKYCESLHKT VRRKTRTVMVGNVALGSE : 100

Ecgcpe : APIAVQSMINIRTTDVEATVNQIKALERVGADIVRVSVPTMDAAEAF.....KLTKQOVNVPLVADIHF DYRIALKVAEYGVDCLRINPGNIGN..... : 114
Atgcpe : HPIRIQIMTTSDTKDITGTVDEVMRLADKGADIVRITVQ GKKEADACFEIKDKLVQLNYNIPLVADIHFAPTVALRVAEC.FDKIRVNP GNFADRRAQFE : 199

Ecgcpe : EERIRMVVD CARDKNIPIRIGVNA GSLEKDEQEKYCEPTTOALL ESAMRHVDHLDRLNFDQFKVSVKASDVFLAVESYRLIAKQ : 198
Atgcpe : TIDYTEDEYQKELQHIEQVFTPEVVKCKKYGRAMRIGTNEGSI SDRIMSYGD.SPRGMVESAFEEFARICRKIDYHNFVFSMKASNPVIMVQAYRIJVAE : 298

Ecgcpe : I.....DQPLHLGITEAGGARS GAVKSAIGLGLLLSEGIGDTLRVSLAADPVEEIK..... : 249
Atgcpe : MYVHGWDYPLHLGVTEAGEGEDGRMKSAIGIGILLQDGLGDTIRVSLTEPPEEKIDPCRRLANLGTKAAKLQQGAPFEEKHRHYFDFQRRRTGDLFPVQKEG : 398

Ecgcpe : : -
Atgcpe : EEVDYRNVLHRDGSVLMSISLDQLKAPELLYRSLATKLVVGMPFKDLATVDSILLRELPPVDDQVARLALKRLIDVSMGVIAPLSEQLT KPLPNAMVLVN : 498

Ecgcpe : : -
Atgcpe : LKELSGGAYKLLPEGTRLVVS LRGDEPYEELEILKNIDATMILHDVPFTEDKVS RVHAARRLFEFLSENSVNF PVIHHINFPTGIHRDELVIHAGTYAGG : 598

Ecgcpe : VGFDILKSLRIRSRGINETACPTCSRQEFDVIGTVNATEQRLEDITTPMDVSTIGCVVNGPGEALVSTLGVTGGNK : 325
Atgcpe : LLVDGLGDGVMLEAPDQDFDFLRNTSEMLLQGC RMRNTKTEYVSCPSCGRTLFDLQEI SAETREKTSHPG.VSTAIMGCIVNGPGEMADADFGYVGGSP : 697

Ecgcpe : .KSGLYEDG.VRKDRIDNNMIDOLEARIRAKASOLDEARRIDVQOVEK : 372
Atgcpe : GRIDLIVGKTVVKRGIAMTEATDALIGLIKEHG.....RWVDPPVADE : 740