



(86) Date de dépôt PCT/PCT Filing Date: 1995/07/25

(87) Date publication PCT/PCT Publication Date: 1996/02/08

(45) Date de délivrance/Issue Date: 2004/04/20

(85) Entrée phase nationale/National Entry: 1996/03/20

(86) N° demande PCT/PCT Application No.: EP 1995/002938

(87) N° publication PCT/PCT Publication No.: 1996/003119

(30) Priorité/Priority: 1994/07/26 (940100369) GR

(51) Cl.Int.⁶/Int.Cl.⁶ A61K 9/70, A61K 31/565, A61K 47/12

(72) Inventeur/Inventor:
FOTINOS, SPIROS, GR


(73) Propriétaire/Owner:
LAVIPHARM S.A., GR

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DISPOSITIF D'ADMINISTRATION TRANSDERMIQUE CONTENANT UN OESTROGENE

(54) Title: TRANSDERMAL DELIVERY DEVICE CONTAINING AN ESTROGEN

MODIFIED FRANZ DIFFUSION CELL



(57) Abrégé/Abstract:


A composition for use in a transdermal drug delivery system is described. The composition comprises an adhesive, an estrogen, and a flux enhancer.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/70, 31/565, 47/12		A1	(11) International Publication Number: WO 96/03119
			(43) International Publication Date: 8 February 1996 (08.02.96)
(21) International Application Number: PCT/EP95/02938 (22) International Filing Date: 25 July 1995 (25.07.95) (30) Priority Data: 940100369 26 July 1994 (26.07.94) GR (71) Applicant (for all designated States except US): LAVIPHARM S.A. [GR/GR]; Agias Marina Street, P.O. Box 59, GR-190 02 Peania Attica (GR). (72) Inventor; and (75) Inventor/Applicant (for US only): FOTINOS, Spiros [GR/GR]; 18a I. Statha Street, GR-106 02 Kolonaki (GR). (74) Agents: CRISP, David, Norman et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: TRANSDERMAL DELIVERY DEVICE CONTAINING AN ESTROGEN

MODIFIED FRANZ DIFFUSION CELL

(57) Abstract

A composition for use in a transdermal drug delivery system is described. The composition comprises an adhesive, an estrogen, and a flux enhancer.

Transdermal delivery device containing an estrogen

5 The present invention relates to a device for the administration of estradiol alone or in combination with progestin(s), encompassing a specific enhancer that achieves elevated transdermal fluxes and optionally an anti-oxidant that achieves better product stability and to a method for manufacturing such device.

10 17-estradiol is the main estrogen produced by the ovaries in pre-menopausal women (J.A. Balfur and R.C. Heel, Drugs 40, 4, 561 - 582, 1990).

17-estradiol, the naturally occurred estrogen, has mainly been used in two areas, such as fertility control and estrogen replacement therapy.

15 Oral administration of estradiol results in an almost complete degradation of this hormone in the digestive tract, due to the phenomenon of first-pass hepatic metabolism. Since a large amount of the administered estradiol, approximately 90%, is destroyed, a large excess should be administered in
20 order to achieve an effective therapeutic result.

It is well-known, that oral administration of estradiol is associated with a number of major side effects such as thrombophlebitis and thrombosis, pulmonary embolism, coronary thrombosis, myocardial infarction, cerebral
25 thrombosis, cerebral hemorrhage and hypertension.

Estrogen replacement therapy is a special need for females on menopause or oophorectomy (loss of one or both ovaries by surgery) and/or pituitary failure. It can also contribute to osteoporosis (loss of bone mass) and atherosclerosis.

Administration of estradiol to post-menopausal women has been found to make post-menopausal symptoms (hot flushes, sweating, nervousness, sleep disturbance) less intense.

5 Co-administration of progestin has been shown to be advantageous for eliminating the side-effects caused by the administration of estradiol itself. Thus, in both fertility control and estrogen replacement therapy, the available therapeutic dosage schemes contain an effective amount of progestin. [Y.W. Chien, T-Y. Chien and Y-C. Huang, U.S patent #4906169 (transdermal
10 Estrogen/Progestin Dosage Unit, System and Process), March 6, 1990].

It has been of great importance to develop a delivery system which will provide certain advantages such as minimization of side effects, prolonged and controlled rate of administration of the hormones, rapid termination of the
15 treatment, and improvement of patient compliance.

The introduction of transdermal systems was found to satisfy the above requirements, which, thus, permits the use of the natural estrogen, 17 α -estradiol, and the use of lower daily doses with the same efficacy because of
20 reduced first-pass hepatic metabolism and continuous drug input.

A number of transdermal delivery systems of various designs exist. They are in general well tolerated, with only 2.5-7% of patients overall having been reported to discontinue the use of transdermal delivery due to severe irritation
25 problems.

The most frequent problem relates to insufficient adhesion of the transdermal "patch" to the skin during wear, resulting in patch loss. Typically, estradiol delivery systems are designed to be worn and to deliver the drug for 3 - 4
30 days.

Several sizes of transdermal estradiol patches already exist on the market, for example, 5, 10, 16 and 20 cm² containing 2, 4, 3.2 and 8 mg of estradiol respectively; the drug is delivered at a rate of 0.21 µg/cm²/hr corresponding to delivery rates of 0.025, 0.05, 0.08 and 0.1 mg per 24 hours (for up to 4 days).

A number of patented systems exist for the delivery of estradiol through the skin, wherein, some selectively presented, (e.g. - Chien, Yie., W. and Chien, Te-Yen, WO 87/07138, Transdermal Absorption Dosage Unit For Estradiol and other Estrogenic Steroids and Process for Administration - Transdermal drug delivery device using a polymer-filled microporous membrane to achieve delayed onset, by Venkatraman S., Cygnus Therapeutic Systems WO 93/03693 - Solid matrix system for transdermal drug delivery, by Chia-Ming Chiang et al, Cygnus Therapeutic Systems, US Patent # 5,252,334 (12.10.1993) - Estradiol transdermal delivery system, by Kim B. et al., Paco Pharmaceutical Services, U.S. Patent #4,906,475/6.3.1990 - Transdermal estrogen/progestin dosage unit, system and process, by Chien Y. et al., Rutgers, the State University of New Jersey, WO 90/06736).

There are basically two types of transdermal drug delivery systems:

a) *Liquid reservoir*

Drug impermeable covering 1.

Drug formulation reservoir 2.

Rate-controlling membrane 3.

Adhesion layer 4.

Release liner 5.

(as illustrated in Figure 1)

2172265

-4-

In this case, estradiol is formulated in a liquid medium and the drug is released from the system via a polymeric membrane.

b) Matrix

- 5 Occlusive cover sheet 6.
Adhesive matrix drug reservoir 7.
Release liner 8.

(as illustrated in Figure 2)

10

In this type of system the drug is formulated directly in the adhesive used which is placed directly in contact with the skin.

15

In general, matrix transdermal delivery system enjoy better patient acceptance, because they are considered to be more convenient than liquid reservoir systems. Furthermore, liquid patches developed for estradiol usually exhibit interactions between the adhesive substance and the semi-solid components of the liquid-reservoir, leading to reduced adhesiveness and altered skin flux rates.

20

All the available enhancers could influence the physicochemical properties of a final transdermal delivery system, such as adhesion, plasticity etc. In general, the selection of all the ingredients for a transdermal delivery system is guided by the desired physicochemical properties.

25

In the present invention, a percentage concentration range of the enhancer is proposed, while an optimum concentration, satisfying our needs for the desirable "patch" with the appropriate skin flux adhesion and removal properties, has been identified.

30

In this invention, linoleic acid is used as skin permeation enhancer. This unsaturated aliphatic acid alone or in combination has been for years in many topical formulations such as cosmetics, toiletries, etc.

5 It is an object of this invention to provide an improved transdermal estradiol delivery system suitable for use in women on fertility control and estrogen replacement therapy which overcomes the problems of the prior art systems.

10 According to a first aspect of the present invention there is provided a composition for use in a transdermal drug delivery system comprising:-

- (a) an adhesive;
- (b) an estrogen; and
- (c) a flux enhancer.

15 Preferably, the adhesive is at least one acrylic adhesive.

Preferably, the adhesive is a mixture of two acrylic adhesives.

20 Preferably, the two acrylic adhesives are in a dry weight ratio of about 80% to about 20% respectively.

Preferably, the amount of estrogen is from about 0.1 to about 8 % w/w.

25 Preferably, the amount of estrogen is 2% w/w.

Preferably, the estrogen is 17- β estradiol, ethimyl-estradiol or combinations thereof.

30 Preferably, the preceding claims wherein the amount of flux enhancer is from about 0.1 to about 20%.

2172265

-6-

Preferably, the amount of flux enhancer is about 5%.

Preferably, the flux enhancer is linoleic acid.

5 Preferably, the estrogen is combined with a progestin.

Preferably, the amount of progestin is from about 0.1 to about 10% w/w.

Preferably, the amount of progestin is about 4% w/w.

10

Preferably, the progestin is norethidrone acetate, levonorgestrel or medroxyprogesterone.

Preferably, the preceding claims further comprising an antioxidant.

15

Preferably, the antioxidant is butylhydroxytoluene (BHT).

Preferably, there are a plurality of compositions.

20 Preferably, each of the compositions contains different concentrations of adhesive, estrogen and flux enhancer.

According to a second aspect of the present invention, there is provided a transdermal delivery system comprising an adhesive, an estrogen, and a flux
25 enhancer.

25

Preferably, a part of the adhesive is in contact with a polymeric release liner which seals and protects the adhesive during storage, and wherein the liner
30 is capable of being peeled off and discarded prior to use of the transdermal delivery system.

30

-7-

Preferably, the release liner is of either polymeric or paper origin.

Preferably, at least a part of the adhesive is in contact with a backing layer which is substantially impermeable to the components of the composition the backing layer being occlusive or breathable.

Preferably, the controlled delivery of estrogen over a period of 1 to 11 days.

According to a third aspect of the present invention there is provided a method of administering estrogen through the skin by use of a transdermal delivery system, the delivery system comprising an adhesive, an an estrogen; and a flux enhancer.

Embodiments of the present invention will now be described by way of example with reference to the drawings in which:-

Figure 1 is a transdermal delivery system having a liquid reservoir.

Figure 2 is a transdermal delivery system having a matrix.

Figure 3 is a transdermal delivery system according to the present invention.

Figure 4 is a modified Franz diffusion cell.

Figure 5 is a graph showing the flux of estradiol through human stratum corneum in vitro.

Figure 6 is a graph showing the permeation of estradiol per day from a 10 sq. cm patch.

Figure 7 is a graph showing the flux of estradiol through human stratum corneum in vitro (after storage for 8 weeks at room temperature).

2172265

-8-

Figure 8 is a graph showing estradiol TDS drug release.

Figure 9 illustrates a shear test of the present invention.

5 It is an advantage of this invention to provide a transdermal delivery system which sufficiently delivers in vivo estrogen alone, or in combination with progestin, through intact skin at a controlled rate for 1-11 days.

10 It is another advantage of this invention to provide a transdermal estrogen and estrogen/progestin combination through a patch that is of smaller size for the same dose than other patches developed todate.

15 It is another advantage of this invention to provide a transdermal delivery system for administration of estrogen or in combination with progestin with better stability characteristics than other developed patches.

20 The present invention is also advantageous in that it relates to the use of linoleic acid together with an anti-oxidant, which enhances the rate of estradiol permeation and maintains superior product stability, that is innovative and unique as well.

25 This invention offers a simple and effective means of delivering estrogen, preferably estradiol and estrogen/progestin combinations through the skin. The hormones are formulated with a mixture of dermatologically acceptable acrylic adhesives, a flux enhancer preferably linoleic acid and optionally an anti-oxidant compound.

30 The transdermal estradiol or estradiol/progestin delivery systems (illustrated in Figure 3) of this invention comprise:

- a) a backing layer 9, which is substantially impermeable to the estradiol and progestin(s) to be delivered transdermally;

-9-

- b) an adhesive matrix 10, which is in contact with said backing layer and has distributed therein an effective amount of estradiol or estradiol/progestin combinations with an enhancing agent such as linoleic acid and an anti-oxidant compound, such as butylhydroxytoluene (BHT) and which secures the dosage unit in contact with the skin of the subject being treated for a period of 1-11 days to permit the hormones to be absorbed transdermally at a controlled rate; and
- c) a release liner 11, which is in contact with said adhesive matrix and which is stripped from the unit prior to the use.

The backing layer can be made of any suitable material which is impermeable to the estradiol alone or in combination with progestin(s) which are dissolved or otherwise distributed in the adhesive matrix. Preferably the materials for making the backing layer are commercially available films of Saranex™ sold by Dow Chemicals or of polyethylene sold by 3M, providing thus the desirable properties in the adhesive matrix.

The adhesive layer must be made of materials biologically acceptable and compatible with said hormones and the skin permeation enhancer. In this invention, preferably a pressure-sensitive material is used such as the vinyl acetate-acrylate multipolymer of Gelva™ 737 and/or Gelva™ 788 resin solution sold by Monsanto.

The adhesive layer is then covered with release liner which is made of materials impermeable to the hormones, the skin permeation enhancer and any other components of the dosage unit, and which is easily stripped or released prior the use. Preferably the release liner is made of siliconized polyester substance sold by release by Release Technologies.

Skin permeation enhancers, which can be employed for carrying out this invention, can widely vary. Specific skin permeation enhancers are distributed within the adhesive matrix and result in a good absorption of the hormones.

5 The above formulation description can be presented in the following tables which demonstrate the composition of a transdermal delivery unit consisting of (a) a single adhesive substance and (b) a mixture of adhesives.

Table 1
Single Adhesive Formulation

10	COMPONENT	QUALITY	QUANTITY % w/w (on a dry basis)
	17 α - estradiol	Pharmaceutically pure	2.0
	Gelva™ 737	Multipolymer Solution	92.9
	Pamolyn™ 200	77% min linoleic acid	5.0
	Butyl Hydroxy Toluene (BHT)		0.1

15 Table 2
Mixed Adhesive Formulation

20	COMPONENT	QUALITY	QUANTITY %w/w (on a dry basis)
	17 α - estradiol	Pharmaceutically pure	2.0
	Gelva™ 737/788 (80/20 ratio by weight in a dried coating)	Multipolymer Solution	92.9
	Linoleic acid	>96%	5.0
	Butyl Hydroxy Toluene (BHT)		0.1

25 This invention is illustrated further by the following examples. Examples are not to be seen as limiting the scope of the invention.

Example no. 1.*Preparation of adhesive mixture and transdermal delivery device*

5 The following ingredients are used in preparing the hormone containing adhesive matrix: estradiol, Gelva, linoleic acid and butylhydroxytoluene (BHT)

10 To 75g of adhesive (Gelva 737 with a total solids content of about 33.8%) solution in toluene ethylacetate and ethanol, estradiol (0.5g) and linoleic acid (1.25g) are added. 200il of BHT solution in isopropanol (0.6g BHT per 100ml of isopropanol) are added and the mixture is stirred at room temperature until all estradiol is dissolved.

When a mixture of adhesives is used, it consists of 60.5 gr Gelva 737 and 11.0 gr Gelva 788.

15 The adhesive mixture is formulated in a transdermal system as follows:

Using an appropriate coating device such as 15 mil casting knives, a layer of the adhesive mixture is coated onto the silicone treated side of polyethylene sheet. The coating is dried in an oven at 70° C for 15 minutes and it is then laminated with a polyester layer on the corona-treated (roughened) side.

20 The process ends with cutting the multi-layer laminate to shapes of the desired geometry and size.

Example no. 2.*Permeability of the transdermal delivery device*

25 The dosage units obtained, as described in the Example 1, are evaluated. The transdermal absorption (flux) of estradiol from the adhesive matrix of this invention is determined in vitro by using human cadaver skin, according to the procedure described by Franz T., In Percutaneous absorption on the relevance of in vitro data, J. Invest. Derm. 64, 190-195, 1975.

30

5 A typical experimental setup for flux measurements (modified Franz diffusion cell), illustrated in Figure 4, comprises a) Glass lid 12, b) Backing sheet 9, c) Drug in the adhesive matrix 10, d) Skin 13, e) Receiving medium 14, f) Sampling arm 15, g) Stirrer 16, h) Receiver compartment 17 and i) Clamp (not shown).

10 For in vitro flux studies, the stratum corneum of human cadaver skin was used. Using fresh, post-mortem skin samples, the stratum corneum was separated from the skin by the technique described by Kligman, A.M. and Christophers, E., in Preparation of isolated sheets of human stratum corneum. Arch. Derm., 88, 702, 1963.

15 The stratum corneum and the transdermal system of the Example 1 were cut into approximately 2-3 cm² squares. A sample of stratum corneum was placed onto the flat surface of the flux cell. After removing the release liner from the transdermal system, the adhesive matrix was laminated onto the stratum corneum, the whole was covered by a glass lid, and secured by a clamp.

20 The receiver medium was a 0.02 wt.% aqueous solution of sodium azide. The assembled cells were placed in a circulating water bath calibrated to maintain the skin surface temperature at 32 ± 1°C.

25 At predetermined time intervals of 6, 12, 24, 36, 48, and 72 hours, the entire content of the receiver was collected for the quantitative determination of estradiol, employing an HPLC method adapted from T. Loftsson and N. Bodor, Acta Pharm. Nord., 1, 4, 1989.

30 The receiver was then again filled with fresh receiving medium. Special care was taken to avoid accumulation of air bubbles at the skin/solution interface.

The cumulative amount of estradiol ($Q_t = \text{ig} \cdot \text{cm}^{-2}$) permeated per unit of area at any time (t) is calculated by using the following formula.

$$Q_t = \sum_{n=0}^t (C_n \cdot V) / A$$

where C_n is the concentration ($\mu\text{g/ml}$) of the drug in the receiving medium for each sampling time, V is the volume (6.3ml) of receiver solution, and A is the cell's diffusion area (0.636cm^2).

5 Typical results are shown in the graph of Figure 5.

The flux of estradiol released from a transdermal delivery system of a given size can be calculated using the above data. The results obtained are given in the graph of Figure 6.

10 A comparative study of skin flux determination between estradiol transdermal system described in the Example 1 and SYSTEN™ 50 Lot 3A01AZ (a matrix patch introduced by Jhonson and Jhonson) is presented in the following Table:

Table 3

15 Competitive skin flux studies between Estradiol transdermal system and SYSTEN™ 50

Transdermal system	Cumulative fluxes ($\mu\text{g/cm}^2$)			“incremental” fluxes ($\mu\text{g/cm}^2$) ratio (sample to reference)			Average
	(0-24h)	(0-48h)	(0-72h)	(0-24h)	(24-48h)	(48-72h)	
Estradiol	2.11	5.04	7.83	2.11 1.10	2.93 1.87	2.79 2.05	2.61 1.67
SYSTEN™ 50	1.92	3.49	4.85	1.92 1.00	1.57 1.00	1.36 1.00	1.62 1.00

20 The results from the Table 3 indicate that the estradiol transdermal delivery system of the present invention exhibits elevated skin fluxes comparing to that of SYSTEN™ 50. As skin flux rate are primary parameter of performance of a transdermal delivery system, the rates of the above table

25 clearly indicate that the transdermal delivery system described in this invention, can be of a smaller size than the commercially available. A smaller size “patch” signifies: 1) a smaller occluded area, therefore less risk of irritation, 2) better aesthetics and therefore, convenience, acceptability and compliance by patients under

2172265

-14-

treatment, and 3) higher production output.

Example no. 3.

Stability of the transdermal delivery device

5

The dosage units described in the Example 1 are evaluated.

10

The stability of estradiol in the adhesive matrix is determined by measuring fluxes of estradiol from the formulation at various intervals during storage of the transdermal systems. The graph of Figure 7 showing the results obtained after 8 weeks at room temperature, indicates that little change in the product's performance occurred during this period.

15

The stability of the transdermal system can also be evaluated by measuring the release rate of estradiol from the matrix employing a standard drug release test (U.S. Pharmacopeia XXII, p.1581).

The results of such measurements are given in the graph of Figure 8.

20

Samples of the transdermal system were kept at room temperature and at 37° C and 45° C for up to 12 weeks, (see also graph of Figure 8). The results again indicate that the system's performance remains satisfactory under these conditions.

25

Example no. 4.

The effect of enhancer on the flux of estradiol through human cadaver skin.

30

The transdermal units described in the Example 1 are evaluated. Various combinations of estradiol and linoleic acid amounts were tested to establish an optimum ratio of the drug and enhancer. The following examples, presented in the table below, illustrate the effect of various concentrations of the two components on the flux of estradiol through human cadaver skin.

Table 4

Effect of linoleic acid on estradiol flux

5	Transdermal delivery device	Flux in $\mu\text{g}/\text{cm}^2/\text{h}$	
		24	48
	A. 4% estradiol & 2% linoleic acid	0.073	0.072
10	4% estradiol & 5% linoleic acid	0.076	0.080
	B. 3% estradiol & 5% linoleic acid	0.133	0.138
15	Estraderm [®] Control	0.067	0.079
	C. 1.5% estradiol & 5% linoleic acid	0.090	0.100
20	D. 2% estradiol & 5% linoleic acid	0.120	0.130
	Estraderm [®] Control	0.060	0.110

As it is indicated above, various estradiol/linoleic acid combinations result in various flux rates.

Differences occurred in experimental results are mostly based upon the skin donor(s).

Example no. 5.

Skin irritation and sensitisation studies using estradiol transdermal system.

The transdermal units described in the Example 1 are tested.

A skin irritation study was conducted in six healthy rabbits of the New Zealand white variety.

The back and flanks of each animal are clipped free of fur with an electric clipper 12 hours before testing. Just prior to test article application, each rabbit

-16-

receives four parallel epidermal abrasions with a sterile needle at one test site (left flank), while the skin at the opposite site remains intact.

5 A 5cm² patch of the test article and a 5 cm² patch of the control article are then applied to each site (four sites per rabbit) to an area of skin approximately 2.7 x 3.7cm. The patches are covered with a non-reactive tape. The trunk of each animal is wrapped with a binder.

10 After 96 hour exposure, the binders, tape and patches are removed. The test sites are gently sponged with deionized water to remove any residue of test article. After at least 30 minutes, the sites are inspected and scored. Another evaluation is also conducted at 144 hours after application.

15 Each test site is examined for dermal reactions using the Draize scoring criteria. The primary irritation index of the test article is calculated following test completion. A substance with an empirical score of less than 5 would not be considered a primary irritant to the skin.

20 The primary irritation index of the test article (estradiol transdermal system of this invention) and of the control article (Estraderm™ TTS 25 µg/24h. Lot # 176700, expiration date 05/94) are calculated to be 1.70 and 1.75 respectively.

Individual experimental results are given in the following Table,. The results for the test article of this invention are given first, followed (after /) by the results for control article.

25 The table that follows presents the data for each rabbit.

Table 5
Skin irritation test

5	Rabbit No.	Reaction	96 hours		144 hours	
			Intact	Abraded	Intact	Abraded
10	No 1	Erythema:	2/0	2/0	1/0	0/1
		Oedema:	1/0	1/1	1/0	0/1
10	No 2	Erythema:	0/0	0/0	0/0	0/0
		Oedema:	0/0	0/0	0/0	0/0
15	No 3	Erythema:	1/2	1/1	1/1	1/1
		Oedema:	0/1	1/1	1/1	2/1
15	No 4	Erythema:	0/0	0/0	1/1	0/1
		Oedema:	0/0	0/0	1/0	0/1
15	No 5	Erythema:	1/1	2/2	0/1	1/2
		Oedema:	0/1	2/2	0/1	2/2
15	No 6	Erythema:	3/1	2/2	1/3	2/2
		Oedema:	2/1	2/2	1/1	2/2

20 Under the conditions of this test, the transdermal system tested could not be considered a primary skin irritant, since the primary irritation index was less than 5 and the difference between the test and control scores was not significant.

25 A skin sensitisation study was conducted in 35 healthy albino guinea pig of the Hartley strain to evaluate and compare the dermal sensitisation potential of estradiol transdermal system and Estraderm TTS patch, according to the method described by Buehler in Archives of Derm. 91, 171, 1965.

30 The estradiol transdermal system was occlusively patched to ten guinea pigs and the control patch was occlusively patched to ten guinea pigs, three times a week until nine induction applications were conducted. Similarly, five positive control guinea pigs were occlusively patched twice a week, with a 0.1% solution of 1-chloro-2,4-dinitrobenzene (DNCB) in propylene glycol, until six
35 inductions had been completed. Following a recovery period of three weeks, the original ten tests and ten controls, ten previously untreated negative control animals and the five positive control animals received a challenge patch. All

sites were scored at 24, 48, and 72 hours after challenge patch removal.

The results obtained are shown in the following Tables 6 and 7. Specifically the results referred to the induction of dermal reactions and challenge phase for the test article of this invention are given first, followed (after /) by the results for the control article.

Table 6
Skin sensitisation test: Dermal reactions - induction phase

Animal Number	OBS	Dermal Reaction The Day After Patch Removal								
		1	2	3	4	5	6	7	8	9
1/11	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0
2/12	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/1 0/1
3/13	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0*/0 0/0
4/14	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0
5/15	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0*/0 0/0
6/16	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0*/0 0/0
7/17	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0
8/18	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0
9/19	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0
10/20	ER OED	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0	0/0 0/0

ER: ERYTHEMA OED: OEDEMA * : BALDNESS

Table 7
Skin sensitisation test: Challenge phase

5 Animal Number	OBS	Hours After Patch Application		
		24	48	72
10 1/11	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
2/12	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
3/13	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
15 4/14	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
5/15	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
6/16	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
20 7/17	ER	0/1	0/0	0/0
	OED	0/1	0/0	0/0
8/18	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
25 9/19	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0
10/20	ER	0/0	0/0	0/0
	OED	0/0	0/0	0/0

ER: ERYTHEMA OED: OEDEMA

30 The following Table presents the results obtained during challenge phase in the negative control animals.

Table 8

Skin sensitisation test: Challenge phase (Negative control)

Animal Number	OBS	A	B	A	B	A	B
26	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
27	ER OED	1 0	1 0	1 0	1 0	0 0	0 0
28	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
29	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
30	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
31	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
32	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
33	ER OED	0 0	0 0	0 0	0 0	0 0	0 0
34	ER OED	1 1	1 1	1 0	1 0	0 0	0 0
35	ER OED	0 0	0 0	0 0	0 0	0 0	0 0

ER: ERYTHEMA OED: OEDEMA

A= Estraderm TTS Patch (Left Flank)

B= Estradiol Transdermal Patch (Right Flank)

The results of the positive control animals obtained from induction and challenge phase are presented below.

Table 9
Skin Sensitisation Test: Induction Phase (Positive Control)

5	Animal Number		Dermal Reactions The Day After Patch Removal					
		OBS	1	2	3	4	5	6
10	21	ER	0	0	0	0	0	0
		OED	0	0	0	0	0	0
	22	ER	0	0	0	0	0	0
		OED	0	0	0	0	0	0
	23	ER	0	0	0	0	0	0
		OED	0	0	0	0	0	0
15	24	ER	0	0	0	0	0	0
		OED	0	0	0	0	0	0
	25	ER	0	0	0	0	0	0
		OED	0	0	0	0	0	0

2172265

Table 10
Skin Sensitisation Test: Challenge Phase (Positive Control)

5	Animal Number		Hours After Patch Application		
		OBS	24	48	72
10	21	ER	2	1	1
		OED	1	1	0
10	22	ER	1	1	0
		OED	1	0	0
15	23	ER	0	0	0
		OED	0	0	0
15	24	ER	1	1	0
		OED	1	0	0
15	25	ER	0	0	0
		OED	0	0	0

ER: ERYTHEMA OED: OEDEMA

-23-

Under the conditions of this test, the estradiol transdermal system would not be considered a sensitizer in the guinea pig. There is no comparative difference of sensitization between the tested estradiol transdermal system and Estraderm control patch.

5

Skin irritation and sensitisation studies for each of the components (i.e. BHT, linoleic acid) of the transdermal delivery system described in this invention, did not show any degree of the irritation and sensitisation.

10

Example no. 7.

Shear test on estradiol transdermal delivery system.

The dosage units derived from the Example 1 are evaluated.

15

The adhesiveness of the system introduced by the present invention was evaluated by measuring the cohesive strength of the formulation matrix. A shear test, as illustrated in Figure 9, is performed according to a standard Test Procedure: A-0150 introduced by Monsanto Chemical Co.

20

In this test, the patch is attached to a standard steel plate 18 using a controlled pressure, and force 19 is applied to effect "sliding" of the patch from the test plate (adapted from British Pharmacopeia, Vol. II, p. A217, 1993). The results of the test are expressed as the time it takes for the patch to come off the plate, when a given force is applied.

25

The results of this test are summarized in the following table:

2172265

-24-

Table 11

Adhesion properties of the estradiol transdermal unit

5	POSITION	TEST SAMPLE	TIME (sec)	COMMENTS
	7	E1293A	398 346	Partial transfer of adhesive from the system to the steel plate occurred in all cases
	8		261 392	
	Average 342/std.dev. = 57			
10	9	E1293B	276 281	Partial transfer of adhesive from the system to the steel plate occurred in all cases
	10		279 334	
15	Average 293/std.dev. = 28			

- 25 -

I claim:

1. A composition for use in a transdermal drug delivery system comprising:
 - (a) an estrogen;
 - (b) a flux enhancer, wherein the flux enhancer is linoleic acid; and
 - (c) a dermatologically acceptable adhesive having an adhesive strength in the presence of the estrogen and the flux enhancer sufficient to maintain contact between the composition and the skin for a period of 1 - 11 days.
2. A composition according to claim 1 wherein the adhesive is at least one acrylic adhesive.
3. A composition according to claim 2 wherein the adhesive is a mixture of two acrylic adhesives.
4. A composition according to claim 3 wherein the two acrylic adhesives are in a dry weight ratio of 80% to 20% respectively.
5. A composition according to any one of claims 1 to 4 wherein the amount of flux enhancer is from 0.1 to 20%.
6. A composition according to claim 5 wherein the amount of flux enhancer is 5%.
7. A composition according to any one of claims 1 to 6 wherein the estrogen is combined with a progestin.
8. A composition according to claim 7 wherein the amount of progestin is from 0.1 to 10% w/w.
9. A composition according to claim 8 wherein the amount of progestin is 4% w/w.

- 26 -

10. A composition according to any one of claims 7 to 9 wherein the progestin is norethidrone acetate, levonorgestrel or medroxyprogesterone.
11. A composition according to any one of claims 1 to 10 wherein the amount of estrogen is from 0.1 to 8% w/w.
12. A composition according to claim 11 wherein the amount of estrogen is 2% w/w.
13. A composition according to any one of claims 1 to 12 wherein the estrogen is 17- β estradiol or ethinyl estradiol or combinations thereof.
14. A composition according to any one of claims 1 to 13 further comprising an antioxidant.
15. A composition according to claim 14 wherein the antioxidant is butylhydroxytoluene (BHT).
16. A composition according to any one of claims 1 to 15 consisting essentially of 17 α estradiol in amount of 2.0% w/w, GelvaTM 737 in an amount of 92.9% w/w, PamolynTM 200 in an amount of 5.0% w/w and butyl hydroxy toluene in an amount of 0.1% w/w.
17. A composition according to any one of claims 1 to 15 consisting essentially of 17 α estradiol in amount of 2.0% w/w, a mixture of GelvaTM 788 at a ratio of 80:20 in an amount of 92.9% w/w, PamolynTM 200 in an amount of 5.0% w/w and butyl hydroxy toluene in an amount of 0.1% w/w.
18. A transdermal delivery system comprising a composition as claimed in any one of claims 1 to 17.

- 27 -

19. A transdermal delivery system containing two or more compositions as claimed in any one of claims 1 to 17.
20. A transdermal delivery system according to claim 19 wherein each of the compositions contains different concentrations of adhesive, estrogen and flux enhancer.
21. A transdermal delivery system according to claim 20 wherein a part of the adhesive is in contact with a polymeric release liner which seals and protects the adhesive during storage, and wherein the liner is capable of being peeled off and discarded prior to use of the transdermal delivery system.
22. A transdermal delivery system according to claim 21 wherein the release liner is of either polymeric or paper origin.
23. A transdermal delivery system according to any one of the claims 18 to 22 for the controlled delivery of estrogen over a period of 1 to 11 days.
24. A transdermal drug delivery system according to any one of claims 18 to 23 further comprising a backing layer and wherein at least a part of the composition is in contact with the backing layer which is substantially impermeable to the components of the composition, said backing layer being occlusive or breathable.
25. Use of a transdermal delivery system for fertility control and estrogen replacement therapy, wherein the delivery system comprises a composition as claimed in any one of claims 1 to 17.
26. A composition for use in a transdermal delivery system, comprising:

- 28 -

- (a) an adhesive matrix having an adhesive strength sufficient to maintain an association between the composition and the skin for at least 3 days in the presence of a flux enhancer, wherein the enhancer is linoleic acid; and
 - (b) an effective dose of an estrogen, the flux enhancer and the estrogen being contained within the adhesive matrix.
27. A composition as defined in claim 26, wherein the adhesive matrix is formed from at least one acrylic adhesive.
28. A composition as defined in claim 27, wherein the adhesive matrix is formed from two acrylic adhesives.
29. A composition as defined in claim 28, wherein the acrylic adhesives are in a dry weight ratio of 80% to 20%.
30. A composition as defined in claim 29, wherein the estrogen is present in a concentration of 0.1-8% w/w.
31. A composition as defined in claim 30, wherein the estrogen is present at about 2% w/w.
32. A composition as defined in claim 26, wherein the estrogen is selected from the group consisting of 17 β estradiol, ethinyl estradiol and combinations thereof.
33. A composition as defined in claim 26, wherein the flux enhancer is present in a concentration range of 0.1-20% w/w.
34. A composition as defined in claim 33, wherein the concentration of the flux enhancer is 5% w/w.
35. A composition as defined in claim 26, further comprising a progestin.

- 29 -

36. A composition as defined in claim 35, wherein the concentration of progestin is in the range of 0.1 to 10% w/w.
37. A composition as defined in claim 36, wherein the concentration of progestin is 4% w/w.
38. A composition as defined in claim 35, wherein the progestin is selected from the group consisting of norethindrone acetate, levonorgestrel and medroxyprogesterone.
39. A composition as defined in claim 26, wherein the adhesive matrix additionally includes an antioxidant.
40. A composition as defined in claim 39, wherein the antioxidant is butylhydroxytoluene (BHT).
41. A composition as defined in claim 26, wherein the adhesive is comprised of two acrylic adhesives at a dry weight ratio of 80% to 20%, the adhesive matrix further comprising the hormone, estrogen, the estrogen having a concentration of 0.1-8% w/w and, optionally, further comprising progestin, at a concentration of 0.1-10% w/w, the flux enhancer comprising linoleic acid at a concentration of 0.1-20%, the matrix further containing an antioxidant.
42. A composition as defined in claim 41, wherein the surface area of the adhesive matrix is less than 5 sq. cm.
43. A transdermal delivery system comprising a composition as claimed in claim 41, wherein a first surface of the adhesive matrix is in contact with a polymeric release liner for sealing and protecting the adhesive during storage, wherein the liner is capable of being removed and discarded prior to use of the transdermal delivery system.

- 30 -

44. A transdermal delivery system as defined in claim 43, wherein the release liner is selected from the group consisting of a polymeric liner and a paper liner.
45. A transdermal delivery system as defined in claim 43, wherein a second surface of the adhesive matrix is in contact with a backing layer, the backing layer being substantially impermeable to the components of the composition, the backing layer being occlusive and breathable.
46. A transdermal delivery system as defined in claim 45, wherein estrogen is delivered in a controlled manner for up to 11 days.
47. A transdermal delivery system as defined in claim 46, having a surface area of less than 5 sq. cm.
48. Use of a transdermal delivery system according to claim 45, for applying the system to the skin, and administering estrogen through the skin.
49. A method of achieving fertility control in a female subject, comprising:
 - (a) applying a transdermal delivery system containing a composition according to claim 26; and
 - (b) permitting controlled release of the estrogen into the bloodstream of the patient.
50. A method as defined in claim 49, wherein the composition in step (a) comprises an adhesive matrix containing estrogen having a concentration of 0.1-8% w/w and optionally further comprising progestin at a concentration of 0.1-10% w/w, the flux enhancer comprising linoleic acid at a concentration of 0.1-20% w/w, and an antioxidant.
51. Use of a transdermal delivery system containing a composition according to claim 26 and use of controlled release of the estrogen into the bloodstream of the patient to achieve estrogen replacement in patients deficient in estrogen.

- 31 -

52. A use as defined in claim 51, wherein the composition comprises an adhesive matrix containing estrogen having a concentration of 0.1-8% w/w and optionally further comprising progestin at a concentration of 0.1-10% w/w, the flux enhancer comprising linoleic acid at a concentration of 0.1-20% w/w, and an antioxidant.
53. A composition for use in a three layer transdermal delivery system, comprising:
- (a) an adhesive matrix in the delivery system having an adhesive strength sufficient to maintain an association between the composition and the skin for at least 3 days in the presence of a flux enhancer and an antioxidant, wherein the enhancer is linoleic acid; and
 - (b) an effective dose of one or more steroid hormones with female sex related activity, the flux enhancer and the hormones being contained within the adhesive matrix, wherein the delivery system is capable of maintaining the delivery of a hormone through intact skin at a controlled rate for at least 3 days at a concentration of flux enhancer of no more than 55% w/w.
54. A composition for use in a three layer transdermal drug delivery system, comprising:
- (a) an adhesive matrix formed from an acrylate polymer having an adhesive strength sufficient to maintain an association between the composition and skin for at least 3 days in the presence of a linoleic acid flux enhancer and an antioxidant; and
 - (b) a steroid hormone with female sex related activity, the linoleic acid flux enhancer and the hormone being contained in the adhesive matrix.
55. A composition as defined in claim 54, wherein the flux enhancer has a concentration of no more than 5% w/w.

FIGURE 1




FIGURE 2




FIGURE 3




MODIFIED FRANZ DIFFUSION CELL

FIGURE 4




5/9

FIGURE 5

**FLUX OF ESTRADIOL THROUGH HUMAN
STRATUM CORNEUM IN VITRO**

6/9

FIGURE 6



7/9

FIGURE 7




FIGURE 8





FIGURE 9



MODIFIED FRANZ DIFFUSION CELL

