

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization

International Bureau



(10) International Publication Number

WO 2018/111556 A1

(43) International Publication Date

21 June 2018 (21.06.2018)

(51) International Patent Classification:

A23K 20/163 (2016.01) A61K 31/716 (2006.01)  
A23K 20/10 (2016.01) A61K 31/353 (2006.01)  
A23K 50/40 (2016.01)

(21) International Application Number:

PCT/US2017/063820

(22) International Filing Date:

30 November 2017 (30.11.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

15/381,783 16 December 2016 (16.12.2016) US

(71) Applicant: HILL'S PET NUTRITION, INC. [US/US];  
400 Southwest 8th Avenue, Topeka, Kansas 66603 (US).

(72) Inventors: JACKSON, Matthew; PO Box 4486, Topeka,  
Kansas 66604 (US). JEWELL, Dennis; 1752 E. 980 Road,  
Lawrence, Kansas 66049 (US).

(74) Agent: HEBLE, Nikhil A. et al.; 909 River Road, Piscataway,  
New Jersey 08854 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

**Published:**

— with international search report (Art. 21(3))

(54) Title: PET FOOD COMPOSITIONS

(57) Abstract: Described herein are pet food compositions comprising a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan); along with methods of making and using same.

WO 2018/111556 A1

## PET FOOD COMPOSITIONS

### BACKGROUND

[0001] Pets require a healthy diet and proper digestion for continued growth and ordinary wellbeing. Oral consumption of yeast beta glucan (BG) is practiced for its benefits to the immune system. However, it is not generally recognized as a prebiotic fiber; and its use in place of more typical prebiotic fibers (e.g. hemicellulose and pectin rich beet and citrus pulps) has traditionally been associated with sub-optimal colonic fermentation and acidification by commensal bacteria. Acidification of the colonic lumen by commensal bacteria is a critical to establishing a healthy microbiome and maintaining intestinal health.

[0002] Reversing the depressive effect of BG (relative to e.g. citrus and beet pulps) on colonic fermentation and acidification has been a challenge. Embodiments of the present invention are designed to meet these needs.

### BRIEF SUMMARY

[0003] In some embodiments, the present invention provides a pet food composition comprising a blood glucose management component comprising soy isoflavone (SI) and BG.

[0004] In some embodiments, the present invention provides methods of controlling post-prandial glucose levels in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising SI and yeast BG.

[0005] In some embodiments, the present invention provides methods of inducing satiety in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising SI and BG.

### DETAILED DESCRIPTION

[0006] As used herein, “satiety” refers to satisfaction of the need for nutrition and the extinguishment of the sensation of hunger, which is often described as “feeling full”. The satiety response refers to behavioral characteristics observed to be consistent with having consumed a sufficient amount of food, such as an abrupt or a tapered down cessation of eating. However, the

biological mechanisms which lead to the satiety response are often triggered in a gradual or delayed manner, such that they are usually out of phase with the amount of food taken in by the animal prior to cessation, which results in the animal consuming more nutritional content than is appropriate for the animal. Satiety inducing agents produce an accelerated onset of the satiety response, i.e., pet food compositions containing satiety inducing agents will trigger the satiety response at an earlier point in time than would a similar pet food composition without the satiety inducing agent.

[0007] In the context of the present disclosure, the terms “yeast beta glucan”, “ $\beta$ -glucan” and “BG” may be used interchangeably.

[0008] In the context of the present disclosure, the terms “soy isoflavone”, “soy isolate” and “SI” may also be used interchangeably.

[0009] In some embodiments, the present invention provides a pet food composition comprising a blood glucose management component comprising SI and BG.

[0010] In some embodiments, the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content greater than about 20% of the total SCFA produced. In other embodiments, the present invention provides a pet food composition wherein the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of greater than about 25% of the total SCFA produced. Still further embodiments provide a pet food composition wherein the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of about 30% of the total SCFA produced.

[0011] Further embodiments provide a pet food composition wherein the ratio of BG to SI (BG : SI) is from about 23:1 to about 7:1.

[0012] Still further embodiments provide a pet food composition wherein the blood glucose management component is present in an amount effective to produce, after about 6 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of about 20% of the total SCFA produced.

[0013] Other embodiments provide a pet food composition wherein the blood glucose management component is present in an amount effective to produce, after about 6 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of greater than about 20% of the total SCFA produced.

[0014] In some embodiments, the propionate content is greater after about 24 hours post-ingestion than the propionate content after about 6 hours post-ingestion.

[0015] Yet other embodiments provide a method of controlling post-prandial glucose levels in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising SI and BG. Some embodiments provide a method of controlling post-prandial glucose levels in a companion animal comprising the step of feeding the companion animal a pet food composition consisting essentially of an effective amount of a blood glucose management component comprising SI and BG.

[0016] The pet food compositions set forth herein may be formed by extrusion to form a kibble-type pet food composition. In some embodiments, the milled raw ingredients of the composition are extruded and then a surface coating comprising a palatant and/or a nutritional oil is applied. In some embodiments, the kibble is spray coated in a tumbling mixer with a composition comprising a palatant and/or a nutritional oil. In other embodiments, the kibble is coated using a vacuum enrobing technique, wherein the kibble is subjected to vacuum and then exposed to coating materials after which the release of the vacuum drives the coating materials inside the kibble.

[0017] The pet food compositions described herein can include any additional ingredients which provide adequate nutrition for the animal. For example, a typical canine or feline diet for use in the present invention may contain from about 17 to about 50% w/v crude protein (and preferably about 20 to about 40% w/v), from about 8 to about 40% w/v fat (and preferably about 12 to about 22% w/v), and from about 0.5 to about 20% w/v total dietary fiber (and preferably about 2 to about 10% w/v), along with the multiple starch source, all percentages by weight. However, no specific ratios or percentages of these nutrients are required.

[0018] In some embodiments, the compositions described herein comprise from about 17% to about 50% w/v crude protein. The crude protein material may comprise vegetable proteins such as soybean meal, soy protein concentrate, corn gluten meal, wheat gluten, cottonseed, and peanut

meal, or animal proteins such as casein, albumin, and meat protein. Examples of meat protein useful herein include beef, pork, lamb, equine, poultry, fish, and mixtures thereof.

[0019] In some embodiments, the compositions described herein comprise from about 8% to about 40% w/v fat. Examples of suitable fats include animal fats and vegetable fats. Preferably the fat source is an animal fat source such as tallow or grease. Vegetable oils such as corn oil, sunflower oil, safflower oil, rape seed oil, soy bean oil, olive oil and other oils rich in monounsaturated and polyunsaturated fatty acids, may also be used.

[0020] In some embodiments, the compositions described herein comprise from about 10% to about 50% w/v carbohydrate. Examples of suitable carbohydrates include grains or cereals such as rice, corn, millet, sorghum, alfalfa, barley, soybeans, canola, oats, wheat, rye, triticale and mixtures thereof. The compositions may also optionally comprise other materials such as dried whey and other dairy by-products.

[0021] In some embodiments, the compositions described herein comprise a fermentable fiber. In some embodiments, the fermentable fiber may be any fiber source which intestinal bacteria present in the animal can ferment to produce significant quantities of SCFAs. In some embodiments, the fermentable fiber is selected from beet pulp, gum arabic (including gum talha), psyllium, rice bran, carob bean gum, citrus pulp, pectin, fructooligosaccharides and inulin, mannanoligosaccharides and mixtures of these fibers.

[0022] The moisture content for the pet food compositions described herein can vary depending on the nature of the food composition. In some embodiments, the pet food compositions may be dry compositions (e.g., kibble), semi-moist compositions, wet compositions, or any mixture thereof. In some embodiments, the pet food composition is a complete and nutritionally balanced pet food. In some embodiments, the pet food may be a “wet food”, “dry food”, or food of “intermediate moisture” content.

[0023] As used herein, “wet food” describes a pet food that is typically sold in cans or foil bags and has a moisture content typically in the range of about 70% to about 90% w/v.

[0024] As used herein, “dry food” describes a pet food that is of a similar composition to wet food but contains a limited moisture content typically in the range of about 5% to about 15% or 20% w/v (typically in the form of small biscuit-like kibbles). In one embodiment, the compositions have moisture content from about 5% to about 20% w/v. Dry food products include a variety of foods of various moisture contents, such that they are relatively shelf-stable

and resistant to microbial or fungal deterioration or contamination. In some embodiments, dry food compositions are extruded food products such as pet foods or snack foods for companion animals.

**[0025]** In some embodiments, the pet food compositions described herein may also comprise one or more fiber sources. As used herein, the term “fiber” includes all sources of “bulk” in the pet food composition whether digestible or indigestible, soluble or insoluble, fermentable or nonfermentable. In some embodiments, the fiber comprises a fiber from a plant source such as marine plants, but microbial sources of fiber may also be used. A variety of soluble or insoluble fibers may be utilized, as will be known to those of ordinary skill in the art. The fiber source can be beet pulp (from sugar beet), gum arabic, gum talha, psyllium, rice bran, carob bean gum, citrus pulp, pectin, fructooligosaccharide, short chain oligofructose, mannanoligosaccharide, soy fiber, arabinogalactan, galactooligosaccharide, arabinoxylan, or mixtures thereof.

**[0026]** Alternatively, the fiber source can be a fermentable fiber. Fermentable fiber has previously been described to provide a benefit to the immune system of a companion animal. Fermentable fiber or other compositions known to skilled artisans that provide a prebiotic to enhance the growth of probiotics within the intestine may also be incorporated into the composition to aid in the enhancement of the benefit provided by the invention to the immune system of an animal.

**[0027]** In some embodiments, the ash content of the pet food composition ranges from less than 1% to about 15% w/v, preferably from about 5% to about 10% w/v.

**[0028]** In some embodiments, the pet food composition comprises from about 17% to about 50% w/v protein, from about 8% to about 40% w/v fat, from about 5% to about 10% w/v ash content, and has a moisture content of about 5% to about 20% w/v. In other embodiments, the pet food composition further comprises probiotics or prebiotics as described herein.

**[0029]** In some embodiments, the compositions described herein can be used as a dietary supplement and be co-administered with another pet food composition. The dietary supplement can have any suitable form such as a gravy, drinking water, beverage, yogurt, powder, granule, paste, suspension, chew, morsel, treat, snack, pellet, pill, capsule, tablet, sachet, or any other suitable delivery form. The dietary supplement can comprise the dietary formulations and optional compounds such as vitamins, preservatives, probiotics, prebiotics, and antioxidants. This permits the supplement to be administered to the animal in small amounts, or in the

alternative, can be diluted before administration. In some embodiments, the dietary supplement may be admixed with a pet food composition or with water or other diluent prior to administration to the animal. When administered in a dietary supplement, the dietary formulations comprise from about 0.1 to about 90% w/v of the supplement, preferably from about 3 to about 70% w/v, more preferably from about 5 to about 60% w/v.

**[0030]** In some embodiments, the compositions described herein are administered to an animal in the form of a nutraceutical composition. The nutraceutical composition comprises anyone of the pet food compositions described herein and one or more nutraceutically acceptable carriers, diluents, or excipients. Generally, nutraceutical compositions are prepared by admixing a compound or composition with excipients, buffers, binders, plasticizers, colorants, diluents, compressing agents, lubricants, flavorants, moistening agents, and the like, including other ingredients known to skilled artisans to be useful for producing nutraceuticals and formulating compositions that are suitable for administration to an animal as a nutraceutical. When administered in a nutraceutical composition, the dietary formulations comprise from about 0.1 to about 90% w/v of the composition, preferably from about 3 to about 70% w/v, more preferably from about 5 to about 60% w/v.

**[0031]** The compositions described herein can be administered to the animal on an as-needed, on an as-desired basis, or on a regular basis. A goal of administration on a regular basis is to provide the animal with a regular and consistent amount of SI and BG or the direct or indirect metabolites that result from such ingestion.

**[0032]** According to the methods of the invention, administration of the dietary formulations, including administration as part of a dietary regimen, can span a period ranging from parturition through the adult life of the animal. In various embodiments, the animal is a companion animal such as a dog or cat. In certain embodiments, the animal is a young or growing animal. In some embodiments, the animal is an aging animal. In other embodiments administration begins, for example, on a regular or extended regular basis, when the animal has reached more than about 30%, 40%, or 50% of its projected or anticipated lifespan. In some embodiments, the animal has attained 40, 45, or 50% of its anticipated lifespan. In yet other embodiments, the animal is older having reached 60, 66, 70, 75, or 80% of its likely lifespan. A determination of lifespan may be based on actuarial tables, calculations, estimates, or the like, and may consider past, present, and future influences or factors that are known to positively or negatively affect lifespan.

Consideration of species, gender, size, genetic factors, environmental factors and stressors, present and past health status, past and present nutritional status, stressors, and the like may also influence or be taken into consideration when determining lifespan.

[0033] The compositions described herein are administered to an animal for a time required to accomplish one or more objectives of the invention, e.g., managing or controlling blood glucose; preventing or treating insulin resistance; improving insulin sensitivity; extending the prime; improving the quality of life; and promoting the health and wellness in an animal. In some embodiments, the compositions described herein are administered to an animal on a regular basis.

[0034] In various embodiments, the compositions comprising the dietary formulations contain at least one of (1) one or more probiotics; (2) one or more inactivated probiotics; (3) one or more components of inactivated probiotics that promote health benefits similar to or the same as the probiotics, e.g., proteins, lipids, glycoproteins, and the like; (4) one or more prebiotics; and (5) combinations thereof. The probiotics or their components can be integrated into the compositions comprising the dietary formulations (e.g., uniformly or non-uniformly distributed in the compositions) or applied to the compositions comprising the dietary formulations (e.g., topically applied with or without a carrier). Such methods are known to skilled artisans, e.g., U.S. Pat. No. 5,968,569 and related patents.

[0035] Typical probiotics include, but are not limited to, probiotic strains selected from *Lactobacilli*, *Bifidobacteria*, or *Enterococci*, e.g., *Lactobacillus reuteri*, *Lactobacillus acidophilus*, *Lactobacillus animalis*, *Lactobacillus ruminis*, *Lactobacillus johnsonii*, *Lactobacillus casei*, *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum*, and *Bifidobacterium* sp., *Enterococcus faecium* and *Enterococcus* sp. In some embodiments, the probiotic strain is selected from the group consisting of *Lactobacillus reuteri* (NCC2581; CNCM 1-2448), *Lactobacillus reuteri* (NCC2592; CNCM 1-2450), *Lactobacillus rhamnosus* (NCC2583; CNCM 1-2449), *Lactobacillus reuteri* (NCC2603; CNCM 1-2451), *Lactobacillus reuteri* (NCC2613; CNCM 1-2452), *Lactobacillus acidophilus* (NCC2628; CNCM 1-2453), *Bifidobacterium adolescentis* (e.g., NCC2627), *Bifidobacterium* sp. NCC2657 or *Enterococcus faecium* SF68 (NCIMB 10415).

[0036] In some embodiments, the compositions described herein may contain one or more prebiotics, e.g., fructo-oligosaccharides, gluco-oligosaccharides, galacto-oligosaccharides,

isomalto-oligosaccharides, xylo-oligosaccharides, soybean oligosaccharides, lactosucrose, lactulose, and isomaltulose. In one embodiment, the prebiotic is chicory root, chicory root extract, insulin, or combinations thereof. Generally, prebiotics are administered in amounts sufficient to positively stimulate the healthy microflora in the gut and cause these “good” bacteria to reproduce.

**[0037]** The probiotics and prebiotics can be made part of the composition by any suitable means. Generally, the agents are mixed with the composition or applied to the surface of the composition, e.g., by sprinkling or spraying. In some embodiments, the pet food composition contains from about 0.1 to about 10% w/v prebiotic. The prebiotics can be integrated into the compositions using methods known to skilled artisans.

**[0038]** In some embodiments, the prebiotic fiber component comprises beet pulp, citrus pulp, a cellulosic material or a mixture thereof. In other embodiments, the prebiotic fiber component comprises a mixture of beet pulp and citrus pulp.

**[0039]** In some embodiments, the total fiber content of the pet food composition is about 0.01% w/v. In some embodiments, the SI is present at concentration of from about 0.005% to about 0.10% w/v. In some embodiments, the SI is present at concentration of from about 0.01% to about 0.05% w/v. In some embodiments, the SI is present at concentration of from about 0.02% to about 0.04% w/v. In some embodiments, the SI is present at about 0.03% w/v. In some embodiments, the SI is of the type available from Naturex of South Hackensack, New Jersey.

**[0040]** In some embodiments, the BG is present at concentration of from about 0.01% to about 1% w/v. In some embodiments, the BG is present at concentration of from about 0.05% to about 0.75% w/v. In some embodiments, the BG is present at concentration of from about 0.25% to about 0.6% w/v. In some embodiments, the BG is present at about 0.40% w/v. In some embodiments, the BG is of the type available from Sensient Flavors, LLC of Indianapolis, Indiana.

**[0041]** In some embodiments, the pet food compositions further comprise from about 17% to about 50% protein. In other embodiments, the pet food compositions of the present invention comprise a source of hydrolyzed animal or plant protein. In some embodiments, the source of hydrolyzed animal or plant protein comprises chicken liver. In further embodiments, the source of hydrolyzed animal or plant protein is present in an active content of from about 25 to about 45% w/v.

[0042] Further embodiments of the present invention provide methods of inducing satiety in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising SI and BG. While other embodiments of the present invention provide methods of inducing satiety in a companion animal comprising the step of feeding the companion animal a pet food composition consisting essentially of an effective amount of a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).

[0043] The invention will now be described in conjunction with the following, non-limiting examples.

## EXAMPLES

### Example 1

[0044] An exemplary pet food composition (Example 1) is prepared as set forth in Table 1 below. All amounts are provided in weight percent, based upon total weight of the pet food composition. The composition is formulated according to the nutrition standards set forth by the American Associated of Feed Control Officials (AAFCO) and the National Research Council (NRC). The composition may be produced by extrusion, dried, and then optionally coated with palatants.

Table 1

Ingredient	Preferred Active Content Range (% w/v)
SI	0.01-0.05
BG	0.1-1
Com, starch, common canning	40-50
Hydrolyzed Chicken Liver and Heart	30-35
Soybean oil, crude, degummed	1-5
Cellulose, Pelleted	1-5
Chicken, liver, digest, optimizor LDPE H	1-3
Lactic acid, food grade	1-2
Calcium carbonate	1-2
Dicalcium phosphate	1-2
Choice White Grease/Phos Acid	0.1-2
Flax Gen#1 + CWG	0.1-1
Glyceryl monostearate	0.1-1
Potassium chloride	0.1-1
Natural flavor, Pork, Liver, Digest, D'T	0.1-1
Sodium chloride, iodized	0.1-1
Choline chloride, liquid, 70%	0.1-1

## Example 2

[0045] Five (5) canines were randomly selected from a general population of beagles and mixed breed dogs. The canine subjects consisted of both neutered males and spayed females, who consumed varied, but typical, canine maintenance foods. The canines were selected to provide a varied compilation of feces that accurately represents the companion animal canine population at large; and would allow an objective assessment of the impact that a specific ingredient, or combination thereof, may have on a particular endpoint.

## Example 3

[0046] Feces were collected from dogs fasted overnight into plastic bags containing oxygen absorbing packs to reduce oxygen tension and maintain viability of anaerobic microbiota. Pooled feces were homogenized in bacterial minimal media and separated of the largest particulates by centrifugation. After adding glycerol as a cryoprotectant the aliquot of viable bacteria were frozen and stored at (-) 80 °C. Black tea and prebiotic fiber blends were sterilized via pasteurization prior to reconstitution in sterile water. Bacteria were incubated with prebiotic fiber as a carbon source as well as a published minimal media. During (6 hours) and after (24 hours) of anaerobic incubation, production of SCFA, including acetate, propionate and butyrate were measured. The results of this experiment are described in Table 2 (below).

Table 2

Ingredient	Ex. 1	Ex. 2	Ex. 3	Comp. Ex. 1	Comp. Ex. 2	Comp. Ex. 3	Comp. Ex. 4	Comp. Ex. 5
% w/w								
Beet Pulp	0.00286	0.00429	0.00571	0.00286	0.00429	0.00571	0.00714	0.00714
Citrus Pulp	0.005	0.0075	0.01	0.005	0.0075	0.01	0.0125	0.0125
Soy Isoflavone	0.00025	0.00025	0.00025	--	--	--	--	0.00025
Yeast $\beta$ -Glucan	0.00857	0.00571	0.00286	0.00857	0.00571	0.00286	--	--
Total Fiber	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Propionate: Total SCFA (%)								
6 hours ( $t_6$ )	22.39	22.11	19.88	16.29	16.43	15.07	14.38	17.18
24 hours ( $t_{24}$ )	29.66	25.26	20.78	13.89	11.54	10.66	10.18	18.31

[0047] As illustrated by the data reported in Table 2 (above), increasing BG and reducing typical prebiotics (e.g. beet pulp and citrus pulp) decreased SCFA production. However, the addition of soy isoflavone unexpectedly increased production of SCFA after both 6 hour and 24 hour incubation periods. It follows, therefore, that the combination of soy isoflavone and BG shifted the type of SCFA produced to favor propionate at the expense of other SCFA after both 6 hour

and 24 hour incubation periods. Propionate is gluconeogenic precursor and enterohepatically recirculated, microbially derived propionate can serve to normalize post-prandial glucose shifts and increase satiety. As such, these results demonstrate that exemplary compositions of the present invention are effective in normalizing post-prandial glucose shifts and increasing satiety, which correlates with an overall improvement in blood glucose management.

[0048] Without being bound by theory, it is believed that SE is not serving as a fermentable carbon source, as its inclusion rate is low. Rather, it appears that SE unexpectedly shifts the microbiome composition or substrate preference.

[0049] Although several embodiments of the invention have been disclosed in the foregoing specification, it is understood by those skilled in the art that many modifications and other embodiments of the invention will come to mind to which the invention pertains, having the benefit of the teaching presented in the foregoing description and associated drawings. It is thus understood that the invention is not limited to the specific embodiments disclosed hereinabove, and that many modifications and other embodiments are intended to be included within the scope of the appended claims. Moreover, although specific terms are employed herein, as well as in the claims which follow, they are used only in a generic and descriptive sense, and not for the purposes of limiting the described invention, nor the claims which follow.

## CLAIMS

### WHAT IS CLAIMED IS:

1. A pet food composition comprising a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).
2. The pet food composition according to claim 1, wherein the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content greater than about 20% of the total SCFA produced.
3. The pet food composition according to claim 1 or claim 2, wherein the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of greater than about 25% of the total SCFA produced.
4. The pet food composition according to any foregoing claim, wherein the blood glucose management component is present in an amount effective to produce, after about 24 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of about 30% of the total SCFA produced.
5. The pet food composition according to any foregoing claim, wherein the ratio of yeast beta glucan to soy isoflavone (yeast beta glucan : soy isoflavone) is from about 23:1 to about 7:1.
6. The pet food composition according to any foregoing claim, wherein the blood glucose management component is present in an amount effective to produce, after about 6 hours post-ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of about 20% of the total SCFA produced.
7. The pet food composition according to claim 3 or claim 4, wherein the blood glucose management component is present in an amount effective to produce, after about 6 hours post-

ingestion by a mammal, a short chain fatty acid (SCFA) complex having a propionate content of greater than about 20% of the total SCFA produced.

8. The pet food composition according to any foregoing claim, wherein the propionate content is greater after about 24 hours post-ingestion than the propionate content after about 6 hours post-ingestion.

9. The pet food composition according to any foregoing claim, further comprising a prebiotic fiber component.

10. The pet food composition according to claim 9, wherein the prebiotic fiber component comprises beet pulp, citrus pulp, a cellulosic material or a mixture thereof.

11. The pet food composition according to claim 9 or claim 10, wherein the prebiotic fiber component comprises a mixture of beet pulp and citrus pulp.

12. The pet food composition according to any foregoing claim, wherein the total fiber concentration is about 0.01% w/v.

13. The pet food composition according to any foregoing claim, wherein the soy isoflavone is present at about 0.03% w/v.

14. The pet food composition according to any foregoing claim, wherein the  $\beta$ -glucan is present at about 0.40% w/v.

15. The pet food composition according to any foregoing claim, further comprising from about 17% to about 50% w/v protein.

16. The pet food composition according to any foregoing claim, further comprising a source of hydrolyzed animal or plant protein.

17. The pet food composition of claim 16, wherein the source of hydrolyzed animal or plant protein comprises chicken liver.
18. The pet food composition according to claim 16 or claim 17, wherein the source of hydrolyzed animal or plant protein is present in an active content of from about 25 to about 45 w/v %.
19. The pet food composition according to any foregoing claim, further comprising from about 8% to about 40% w/v fat.
20. The pet food composition according to any foregoing claim having an ash content of from about 5% to about 10% w/v.
21. The pet food composition according to any foregoing claim having a moisture content of from about 5% to about 20% w/v.
22. The pet food composition according to any foregoing claim, further comprising high docosahexaenoate fish oil.
23. A method of controlling post-prandial glucose levels in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).
24. A method of controlling post-prandial glucose levels in a companion animal comprising the step of feeding the companion animal a pet food composition consisting essentially of an effective amount of a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).

25. A method of inducing satiety in a companion animal comprising the step of feeding the companion animal a pet food composition comprising an effective amount of a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).
26. A method of inducing satiety in a companion animal comprising the step of feeding the companion animal a pet food composition consisting essentially of an effective amount of a blood glucose management component comprising soy isoflavone and yeast beta-glucan ( $\beta$ -glucan).

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/063820

**A. CLASSIFICATION OF SUBJECT MATTER**  
 INV. A23K20/163 A23K20/10 A23K50/40 A61K31/716 A61K31/353  
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
**A23K A61K**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

**EPO-Internal, BIOSIS, FSTA, WPI Data**

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KR 100 439 147 B1 (KANG SANG MO [KR]) 5 July 2004 (2004-07-05) claims 1-4 examples 1-4 ----- Y WO 01/78534 A2 (PROCTER & GAMBLE [US]) 25 October 2001 (2001-10-25) claims 1,5,8 page 5, lines 1-22 page 6, lines 1-2 ----- Y WO 90/04334 A1 (ALPHA BETA TECHNOLOGY [US]) 3 May 1990 (1990-05-03) claims 7-13,16,23 page 5, lines 4-11 ----- -/-	1-4,6-8  1-26  1-26  -/-

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier application or patent but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
13 February 2018	09/03/2018
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  <b>Heirbaut, Marc</b>

## INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/063820

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BOEVER DE P ET AL: "FERMENTATION BY GUT MICROBIOTA CULTURED IN A SIMULATOR OF THE HUMAN INTESTINAL MICROBIAL ECOSYSTEM IS IMPROVED BY SUPPLEMENTING A SOYGERM POWDER", THE JOURNAL OF NUTRITION, AMERICAN SOCIETY FOR NUTRITION, US, vol. 130, no. 10, 1 October 2000 (2000-10-01), pages 2599-2606, XP001001515, ISSN: 0022-3166 abstract -----	1-26
A	WO 01/74345 A2 (INGRAM JONATHAN [US]) 11 October 2001 (2001-10-11) the whole document -----	1-26
A	WO 2005/089567 A1 (NESTEC SA [CH]; PAN YUANLONG [CH]) 29 September 2005 (2005-09-29) the whole document -----	1-26

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/063820

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
KR 100439147	B1	05-07-2004	NONE		
-----					
WO 0178534	A2	25-10-2001	AU 5339201 A		30-10-2001
			AU 2001253392 B2		02-02-2006
			CA 2405803 A1		25-10-2001
			EP 1272060 A2		08-01-2003
			US 2002012733 A1		31-01-2002
			US 2004086547 A1		06-05-2004
			WO 0178534 A2		25-10-2001
-----					
WO 9004334	A1	03-05-1990	AT 106193 T		15-06-1994
			AU 643670 B2		25-11-1993
			CA 2001754 A1		28-04-1990
			DE 68915773 D1		07-07-1994
			DE 68915773 T2		03-11-1994
			EP 0440725 A1		14-08-1991
			IE 64707 B1		23-08-1995
			IL 92098 A		12-04-1994
			JP H04505997 A		22-10-1992
			MX 168000 B		27-04-1993
			NZ 231176 A		26-05-1997
			US 4962094 A		09-10-1990
			WO 9004334 A1		03-05-1990
			ZA 8908174 B		30-10-1991
-----					
WO 0174345	A2	11-10-2001	AU 5307001 A		15-10-2001
			US 2002010141 A1		24-01-2002
			WO 0174345 A2		11-10-2001
-----					
WO 2005089567	A1	29-09-2005	AU 2005224017 A1		29-09-2005
			BR PI0508936 A		25-09-2007
			CA 2559974 A1		29-09-2005
			CN 1933739 A		21-03-2007
			EP 1750523 A1		14-02-2007
			EP 2208422 A1		21-07-2010
			ES 2346234 T3		13-10-2010
			ES 2559777 T3		15-02-2016
			JP 5283902 B2		04-09-2013
			JP 2007529211 A		25-10-2007
			JP 2012165746 A		06-09-2012
			RU 2366274 C2		10-09-2009
			US 2005222050 A1		06-10-2005
			US 2012270814 A1		25-10-2012
			WO 2005089567 A1		29-09-2005
			ZA 200608626 B		30-07-2008
-----					