



US00RE42704E

(19) **United States**  
(12) **Reissued Patent**  
**Prunkard et al.**

(10) **Patent Number:** **US RE42,704 E**  
(45) **Date of Reissued Patent:** **Sep. 13, 2011**

(54) **PRODUCTION OF FIBRINOGEN IN TRANSGENIC ANIMALS**

(75) Inventors: **Donna E. Prunkard**, Seattle, WA (US);  
**Donald C. Foster**, Lake Forest Park, WA (US)

(73) Assignee: **Pharming Intellectual Property B.V.**,  
Al Leiden (NL)

(21) Appl. No.: **09/232,488**

(22) Filed: **Jan. 15, 1999**

**Related U.S. Patent Documents**

Reissue of:

(64) Patent No.: **5,639,940**  
Issued: **Jun. 17, 1997**  
Appl. No.: **08/206,176**  
Filed: **Mar. 3, 1994**

(51) **Int. Cl.**  
**C12P 21/00** (2006.01)  
**A01K 67/00** (2006.01)  
**C12N 15/00** (2006.01)

(52) **U.S. Cl.** ..... **800/7; 800/14; 800/15; 800/16;**  
**800/17; 800/18; 800/25; 435/320.1**

(58) **Field of Classification Search** ..... **800/7, 14-18,**  
**800/25; 435/320.1**

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

4,873,316 A 10/1989 Meade et al.  
5,304,489 A 4/1994 Rosen  
5,366,894 A 11/1994 Clark et al.

**FOREIGN PATENT DOCUMENTS**

WO WO88/00239 1/1988  
WO WO 90/05188 5/1990  
WO WO91/03216 6/1991  
WO WO91/08216 6/1991  
WO WO92/11358 7/1992  
WO WO92/11757 7/1992

**OTHER PUBLICATIONS**

Garner's Preliminary Motion No. 1, Aug. 16, 2001.  
Declaration of Alan Colman, Aug. 16, 2001.  
Declaration of David Toman, Aug. 16, 2001.  
Declaration of Lothar Hennighausen, Aug. 16, 2001.  
Declaration of Jeffrey Rosen, Aug. 16, 2001.  
Yarus et al., *Genetic Engineering*, 18: 57-81 (1996).  
Reddy et al., *Animal Biotechnology* 2: 15-29 (1991).  
Tojo et al., *J. Reprod. Develop.* 39: 145-155 (1993).  
Farrell et al., *Biochemistry*, 30: 9414-9420 (1991).  
Velander's Preliminary Motion No. 1, Aug. 16, 2001.  
Garner's Opposition to Velander's Preliminary Motion No. 1, Aug. 16, 2001.  
Prunkard et al., *Nature Biotechnology*, 14:867-871, 870 (1996).  
Hurtley et al., *Annu. Rev. Cell Biol.* 5: 277-307 (1989).  
Prunkard et al., (Abstract) *J. Cell Biochem., Suppl. 17C* (1993).  
Drohan et al., *Transgenic Research*, 3: 355-364 (1994).  
Velander et al., *PNAS (USA)*, 89: 12003-12007 (1992).  
Pittius et al., *PNAS USA*, 85: 5874 (1988).  
Simons et al., *Biotechnology*, 6:179 (1988).  
Clark et al., *Biotechnology*, 7:487 (1989).

Garner's Motion to Correct Inventorship, Aug. 16, 2001.  
Garner's Preliminary Motion No. 2, Aug. 16, 2001.  
Hennighausen, *Protein Expression and Publication* 1: 3-8 (1990).  
Garner Exhibit No. 28—Typed Notes, Aug. 16, 2001.  
Masoud et al., *Ann. Zootech.* 45: -9 (1996).  
Garner's Reply in Support of Garner's Preliminary Motion 1, Aug. 16, 2001.  
Second Declaration of Alan Colman, Aug. 16, 2001.  
Declaration of Richard F. Lathe, Aug. 16, 2001.  
Behringer et al., "Synthesis Of Functional Human Hemoglobin In Transgenic Mice," *Science*, 245:971-973 (1989).  
Binnie et al., "Characterization Of Purified Recombinant Fibrinogen: Partial Phosphorylation Of Fibrinopeptide A," *Biochemistry*, 32:107-113 (1993).  
Bolyard, M.G., et al., "Expression In *Escherichia Coli* Of The Human Fibrinogen B $\beta$  Chains And Its Cleavage By Thrombin," *Blood*, 73:1202-1206, 1989.  
Burdon et al., "Expression Of A Whey Acidic Protein Transgene During Mammary Development," *The Journal of Biological Chemistry*, 266(11):6909-6914 (1991).  
Chung, Dominic W., et al., "Characterization Of A Complementary Deoxyribonucleic Acid and Genomic Deoxyribonucleic Acid For The  $\beta$  Chain Of Human Fibrinogen," *Biochemistry*, 22:3244-3250.  
Chung et al., "Characterization Of A Complementary Deoxyribonucleic Acid Coding For The  $\gamma$  Chain Of Human Fibrinogen," *Biochemistry*, 22(13):3250-3256 (1983).  
Chung et al., "Nucleotide Sequences Of The Three Genes Coding For Human Fibrinogen," *Adv. Exp. Med.*, 201:39-48 (1990).  
Clark et al., "Pharmaceuticals From Transgenic Livestock," *Tibtech*, 5:20-24 (1987).  
Clark et al., "Expression of human anti-hemophilic factor IX in the milk of transgenic sheep," *Bio/Technology*, 7:487-492 (1989).  
Clark et al., "Rescuing Transgene Expression By Co-Integration," *Bio/Technology*, 10:1450-1454.  
Danishefsky, K., et al., "Intracellular Fate Of Fibrinogen B $\beta$  Chain Expressed in COS Cells," *Biochem. Biophys. Acta*, 1048:202-208 (1990).  
James Darnell, *Molecular Cell Biology* abstract "The Organization Of DNA Differs Greatly In Prokaryotic and Eukaryotic Cells," p. 137 (Scientific American Books, 1986).  
DiTullio et al., "Production Of Cystic Fibrosis Transmembrane Conductance Regulator In The Milk of Transgenic Mice," *Bio/Technology*, 10:74-77 (1992).  
Dobrovolsky et al., "Human  $\gamma$ -Interferon Expression In The Mammary Gland Of Transgenic Mice," *FEBS Letters*, 319:181-184 (1993).

(Continued)

*Primary Examiner* — Deborah Crouch  
(74) *Attorney, Agent, or Firm* — Alston and Bird LLP

(57) **ABSTRACT**

Materials and methods for producing fibrinogen in transgenic non-human mammals are disclosed. DNA segments encoding A $\alpha$ , B $\beta$  and  $\gamma$  chains of fibrinogen are introduced into the germ line of a non-human mammal, and the mammal or its female progeny produces milk containing fibrinogen expressed from the introduced DNA segments. Non-human mammalian embryos and transgenic non-human mammals carrying DNA segments encoding heterologous fibrinogen polypeptide chains are also disclosed.

**17 Claims, 5 Drawing Sheets**

## OTHER PUBLICATIONS

- Drohan et al., "Inefficient Processing Of Human Protein C In The Mouse Mammary Gland," *Transgenic Research*, 3:355-364 (1994).
- Farrell, D.H. et al., "Binding Of Recombinant Fibrinogen Mutants To Platelets," *J. Biol. Chem.*, 269:226-231 (1994).
- Farrell et al., "Recombinant Human Fibrinogen And Sulfation Of The 65' Chain," *Biochemistry*, 30(39):9414-9420 (1991).
- Greenberg et al., "Expression Of Biologically Active Heterodimeric Bovine Follicle-Stimulating Hormone In Milk Of Transgenic Mice," *Biochemistry*, 88:8327-8331 (1991).
- Grosschedl, R., et al., "Introduction Of A  $\mu$  Immunoglobulin Gene Into The Mouse Germ Line; Specific Expression In Lymphoid Cells And Synthesis Of Functional Antibody," *Cell*, 38:647-658 (1984).
- Hartwig, R. et al., "Studies On The Assembly And Secretion Of Fibrinogen," *J. Biol. Chem.*, 266:6578-6585 (1991).
- Hennighausen, Lothar, "The Mammary Gland As A Bioreactor: Production Of Foreign Proteins In Milk," *Protein Expression And Purification*, 1:3-8 (1990).
- Hurtley et al., "Protein Obligomerization In The Endoplasmic Reticulum," *Annu. Rev. Cell Biol.*, 5: 277-307 (1989).
- Lee, S.H., et al., "Production Of Biomedical Proteins In The Milk Of Transgenic Dairy Cows; The State Of The Art," *J. Controlled Release*, 29:213-221 (1994).
- Lord, Susan T., "Expression Of A Cloned Human Fibrinogen cDNA In *Escherichia Coli*: Synthesis Of An A Alpha Polypeptide," *DNA*, 4:33-38 (1980).
- Lord, Susan T., et al., "Expression Of A Fibrinogen Fusion Peptide In *Escherichia Coli*: A Model Thrombin Substrate For Structure/Function Analysis," *Blood*, 73:166-171 (1989).
- Massoud et al., "The Deleterious Effects Of Human Erythropoietin Gene Driven By The Rabbit Whey Acidic Protein Gene Promoter In Transgenic Rabbits," *Ann Zootech.*, 45:1-9 (1996).
- McClenaghan et al., "Secretory Proteins Compete For Production In The Mammary Gland Of Transgenic Mice," *Biochemical Journal*, 310:637-641 (1995).
- Mishina et al., "Molecular Distinction Between Fetal And Adult Forms Of Muscle Acetylcholine Receptor," *Nature*, 321:406-411 (1986).
- Morocöl et al., "The Porcine Mammary Gland As A Bioreactor For Complex Proteins," *Recombinant DNA Technology II*, 721:21-233 (1994).
- Nicoll et al., "Estimates Of Parenchymal, Stromal, And Lymph Node Deoxyribonucleic Acid In Mammary Glands Of C311/Crgl/2 MICE," *Life Sciences*, 4(9):993-1001 (1965).
- Palmiter, Richard D., et al., "Transgenic Mice," *Cell*, 41:343-345 (1985).
- Pittius et al., "A Milk Protein Gene Promoter Directs The Expression Of Human Tissue Plasminogen Activator cDNA To The Mammary Gland In Transgenic Mice," *Biochemistry*, 85:5874-5878 (1988).
- Prunkard et al. UCLA Symposia On Molecular And Cellular Biology (Feb. 8-14, 1993, Taos, New Mexico), "High Level Secretion Of Recombinant Human Fibrinogen In BHK Cells Is Limited By A Post-Transcriptional Process," *Journal of Cellular Biochemistry*, vol. 17C, Abstract H123 (1993).
- Prunkard et al., "High-Level Expression Of Recombinant Human Fibrinogen In The Milk Of Transgenic Mice," *Nature Biotechnology*, 14:867-871 (1996).
- Reddy et al., "Expression Of Human Growth Hormone In The Milk Of Transgenic Mice," *Animal Biotechnology*, 2(1):15-29. (1991).
- Redman, C., et al., "Recombinant Production, Secretion, And Clotting Behavior Of Fibrinogen And Cell Line Used Therein," *Chemical Abstracts*, 115: 526 (1991).
- Rixon, M.W., et al., "Nucleotide Sequence Of The Gene For The  $\gamma$  Chain Of Human Fibrinogen," *Biochemistry*, 24:2077-2086 (1985).
- Rixon et al., "Characterization Of A Complementary Deoxyribonucleic Acid Coding For The  $\alpha$  Chain Of Human Fibrinogen," *Biochemistry*, 22(13):3237-3244 (1983).
- Roy et al., "Assembly And Secretion Of Recombinant Human Fibrinogen," *The Journal of Biological Chemistry*, 266(8):4758-4763 (1991).
- Shani et al., "Expression Of Human Serum Albumin In The Milk Of Transgenic Mice," *Transgenic Research*, 1(5):195-208 (1992).
- Simons et al., "Gene Transfer Into Sheep," *Bio/Technology*, 6:179-183 (1988).
- Sinai Yarus et al., "Engineering Transgenes For Use In The Mammary Gland," *Genetic Engineering*, 18:57-81 (1996).
- Storb et al., "Transgenic Mice With  $\mu$  And  $\kappa$  Genes Encoding Antiphosphorylcholine Antibodies," *The Journal of Experimental Medicine*, 164:627-641 (1986).
- Tojo et al., "Production And Characterization Of Transgenic Mice Expressing A hGH Fusion Gene Driven By The Promoter Of Mouse Whey Acidic Protein (mWAP) Putatively Specific To Mammary Gland," *Journal of Reproduction and Development*, 39(2):145-155 (1993).
- Turner et al., "Proteins Are Secreted By Both Constitutive And Regulated Secretory Pathways In Lactating Mouse Mammary Epithelial Cells," *The Journal of Cell Biology*, 117:269-278 (1992).
- Velander et al., "High-Level Expression Of A Heterologous Protein In The Milk Of Transgenic Swine Using The cDNA Encoding Human Protein C," *Applied Biological Sciences*, 89:12003-12007 (1992).
- Whitelaw et al., "Position-Independent Expression Of The Ovine  $\beta$ -Lactoglobulin Gene In Transgenic Mice," *Biochemical Journal*, 286:31-39 (1992).
- Wilkins et al., "Isolation Of Recombinant Proteins From Milk," *Journal of Cellular Biochemistry*, 49:333-338 (1992).
- Yu et al., "Fibrinogen Precursors," *The Journal of Biological Chemistry*, 259(16):10574-10581 (1984).
- Yu et al., "Functional Human CD4 Protein Produced In Milk Of Transgenic Mice," *Mol. Biol. Med.*, 6:255-261 (1989).
- Bolyard et al., *Gene* 66: 183-192, 1988.
- Wall, 1996, *Themogeneology* 45:57-68.
- Houdebine, 1994, *Journal of Biotechnology* 34:269-287.
- Genes, Lewin, ed., John Wiley and Sons, N.Y. pp. 87-96.
- D.F. Hoshier, "Disorders of Blood Coagulation" in *Cecil Textbook of Medicine*, 18<sup>th</sup> Ed., Wyngaarden et al., eds., W.B. Saunders Co., Philadelphia, 1988. pp. 1060-1065.

FIGURE 1




FIGURE 2




FIGURE 3






FIGURE 4



**FIGURE 5**



**PRODUCTION OF FIBRINOGEN IN  
TRANSGENIC ANIMALS**

**Matter enclosed in heavy brackets [ ] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.**

BACKGROUND OF THE INVENTION

The final step in the blood coagulation cascade is the thrombin-catalyzed conversion of the soluble plasma protein fibrinogen to insoluble fibrin. Thrombin cleaves a small peptide (fibrinopeptide A) from one of the three component chains (the  $\alpha$ -chain) of fibrinogen. Fibrin monomers subsequently polymerize and are cross-linked by activated factor XIII to form a stable clot.

Fibrinogen is a key component of biological tissue glues (see, e.g., U.S. Pat. Nos. 4,377,572 and 4,442,655), which mimic the formation of natural blood clots to promote hemostasis and repair damaged tissue. Tissue glues provide an adjunct or alternative to sutures, staples and other mechanical means for wound closure. However, the principal ingredients of these products (fibrinogen, factor XIII and thrombin) are prepared from pooled human plasma by cryoprecipitation (e.g. U.S. Pat. Nos. 4,377,572; 4,362,567; 4,909,251) or ethanol precipitation (e.g. U.S. Pat. No. 4,442,655) or from single donor plasma (e.g. U.S. Pat. No. 4,627,879; Spotnitz et al., *Am. Surg.* 55: 166-168, 1989). The resultant fibrinogen/factor XIII preparation is mixed with bovine thrombin immediately before use to convert the fibrinogen to fibrin and activate the factor XIII, thus initiating coagulation of the adhesive.

Commercially available adhesives are of pooled plasma origin. Because blood-derived products have been associated with the transmission of human immunodeficiency virus (HIV), hepatitis virus and other etiologic agents, the acceptance and availability of such adhesives is limited. At present they are not approved for use in the United States.

While the use of autologous plasma reduces the risk of disease transmission, autologous adhesives can only be used in elective surgery when the patient is able to donate the necessary blood in advance.

As noted above, fibrinogen consists of three polypeptide chains, each of which is present in two copies in the assembled molecule. These chains, designated the  $\alpha$ ,  $\beta$  and  $\gamma$ -chains, are coordinately expressed, assembled and secreted by the liver. While it might be expected that recombinant DNA technology could provide an alternative to the isolation of fibrinogen from plasma, this goal has proven to be elusive. The three fibrinogen chains have been individually expressed in *E. coli* (Lord, *DNA* 4: 33-38, 1985; Bolyard and Lord, *Gene* 66: 183-192, 1988; Bolyard and Lord, *Blood* 73: 1202-1206), but functional fibrinogen has not been produced in a prokaryotic system. Expression of biologically competent fibrinogen in yeast has not been reported. Cultured transfectant mammalian cells have been used to express biologically active fibrinogen (Farrell et al., *Blood* 74: 55a, 1989; Hartwig and Danishefsky, *J. Biol. Chem.* 266: 6578-6585, 1991; Farrell et al., *Biochemistry* 30: 9414-9420, 1991), but expression levels have been so low that production of recombinant fibrinogen in commercial quantities is not feasible. Experimental evidence suggests that lower transcription rates in cultured cells as compared to liver may be a factor in the low expression rates achieved to date, but increasing the amount of fibrinogen chain mRNA in transfected BHK cells did not produce corresponding increases in fibrinogen protein

secretion (Prunkard and Foster, XIV Congress of the International Society on Thrombosis and Haemostasis, 1993). These latter results suggest that proper assembly and processing of fibrinogen involves tissue-specific mechanisms not present in common laboratory cell lines.

There remains a need in the art for methods of producing large quantities of high quality fibrinogen for use in tissue adhesives and other applications. There is a further need for fibrinogen that is free of blood-borne pathogens. The present invention fulfills these needs and provides other, related advantages.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide commercially useful quantities of recombinant fibrinogen, particularly recombinant human fibrinogen. It is a further object of the invention to provide materials and methods for expressing fibrinogen in the mammary tissue of transgenic animals, particularly livestock animals such as cattle, sheep, pigs and goats.

Within one aspect, the present invention provides a method for producing fibrinogen comprising (a) providing a first DNA segment encoding a secretion signal operably linked to a fibrinogen  $\alpha$  chain, a second DNA segment encoding a secretion signal operably linked to a fibrinogen  $\beta$  chain, and a third DNA segment encoding a secretion signal operably linked to a fibrinogen  $\gamma$  chain, wherein each of the first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal; (b) introducing the DNA segments into a fertilized egg of a non-human mammalian species; (c) inserting the egg into an oviduct or uterus of a female of the species to obtain offspring carrying the DNA constructs; (d) breeding the offspring to produce female progeny that express the first, second and third DNA segments and produce milk containing biocompetent fibrinogen encoded by the segments; (e) collecting milk from the female progeny; and (f) recovering the fibrinogen from the milk. Within one embodiment, the egg containing the introduced segments is cultured for a period of time prior to insertion.

Within another aspect, the invention provides a method of producing fibrinogen comprising the steps of (a) incorporating a first DNA segment encoding a secretion signal operably linked to an  $\alpha$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a first gene fusion; (b) incorporating a second DNA segment encoding a secretion signal operably linked to a  $\beta$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a second gene fusion; (c) incorporating a third DNA segment encoding a secretion signal operably linked to a  $\gamma$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a third gene fusion; (d) introducing the first, second and third gene fusions into the germ line of a non-human mammal so that the DNA segments are expressed in a mammary gland of the mammal or its female progeny and biocompetent fibrinogen is secreted into milk of the mammal or its female progeny; (e) obtaining milk from the mammal or its female progeny; and (f) recovering the fibrinogen from the milk. Within preferred embodiments, the mammal is a sheep, pig, goat or bovine.

Within another aspect, the invention provides a method for producing fibrinogen comprising the steps of (a) providing a transgenic female non-human mammal carrying in its germline heterologous DNA segments encoding  $\alpha$ ,  $\beta$  and  $\gamma$  chains of fibrinogen, wherein the DNA segments are expressed in a mammary gland of the mammal and fibrinogen encoded by the DNA segments is secreted into milk of the

mammal; (b) collecting milk from the mammal; and (c) recovering the fibrinogen from the milk.

Within another aspect, the invention provides a non-human mammalian embryo containing in its nucleus heterologous DNA segments encoding  $\text{A}\alpha$ ,  $\text{B}\beta$  and  $\gamma$  chains of fibrinogen. Within a related aspect, the invention provides a transgenic non-human female mammal that produces recoverable amounts of human fibrinogen in its milk.

Within another aspect, the invention provides a method for producing a transgenic offspring of a mammal comprising the steps of (a) providing a first DNA segment encoding a fibrinogen  $\text{A}\alpha$  chain, a second DNA segment encoding a fibrinogen  $\text{B}\beta$  chain, and a third DNA segment encoding a fibrinogen  $\gamma$  chain, wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in a mammary gland of a host female mammal and secretion into milk of the host female mammal; (b) introducing the DNA segments into a fertilized egg of a mammal of a non-human species; (c) inserting the egg into an oviduct or uterus of a female of the non-human species to obtain an offspring carrying the first, second and third DNA segments. In a related aspect, the invention provides non-human mammals produced according to this process.

Within an additional aspect, the invention provides a non-human mammal carrying its germline DNA segments encoding heterologous  $\text{A}\alpha$ ,  $\text{B}\beta$  and  $\gamma$  chains of fibrinogen, wherein female progeny of the mammal express the DNA segments in a mammary gland to produce biocompetent fibrinogen.

These and other aspects of the invention will become evident to the skilled practitioner upon reference to the following detailed description and the attached drawings.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates the subcloning of a human fibrinogen  $\text{A}\alpha$  chain DNA sequence.

FIG. 2 is a partial restriction map of the vector Zem228. Symbols used are MT-1p, mouse metallothionein promoter; SV40t, SV40 terminator; and SV40p, SV40 promoter.

FIG. 3 illustrates the subcloning of a human fibrinogen  $\text{B}\beta$  chain DNA sequence.

FIG. 4 illustrates the subcloning of a human fibrinogen  $\gamma$  chain DNA sequence.

FIG. 5 is a partial restriction map of the vector Zem219b. Symbols used are MT-1p, mouse metallothionein promoter; hGHt, human growth hormone terminator; SV40p, SV40 promoter; DHFR, dihydrofolate reductase gene; and SV40t, SV40 terminator.

#### DETAILED DESCRIPTION OF THE INVENTION

Prior to setting forth the invention in detail, it will be helpful to define certain terms used herein:

As used herein, the term "biocompetent fibrinogen" is used to denote fibrinogen that polymerizes when treated with thrombin to form insoluble fibrin.

The term "egg" is used to denote an unfertilized ovum, a fertilized ovum prior to fusion of the pronuclei or an early stage embryo (fertilized ovum with fused pronuclei).

A "female mammal that produces milk containing biocompetent fibrinogen" is one that, following pregnancy and delivery, produces, during the lactation period, milk containing recoverable amounts of biocompetent fibrinogen. Those skilled in the art will recognize that such animals will produce milk and therefore the fibrinogen, discontinuously.

The term "progeny" is used in its usual sense to include children and descendants.

The term "heterologous" is used to denote genetic material originating from a different species than that into which it has been introduced, or a protein produced from such genetic material.

Within the present invention, transgenic animal technology is employed to produce fibrinogen within the mammary glands of a host female mammal. Expression in the mammary gland and subsequent secretion of the protein of interest into the milk overcomes many difficulties encountered in isolating proteins from other sources. Milk is readily collected, available in large quantities, and well characterized biochemically. Furthermore, the major milk proteins are present in milk at high concentrations (from about 1 to 15 g/l).

From a commercial point of view, it is clearly preferable to use as the host a species that has a large milk yield. While smaller animals such as mice and rats can be used (and are preferred at the proof-of-concept stage), within the present invention it is preferred to use livestock mammals including, but not limited to, pigs, goats, sheep and cattle. Sheep are particularly preferred due to such factors as the previous history of transgenesis in this species, milk yield, cost and the ready availability of equipment for collecting sheep milk. See WO 88/00239 for a comparison of factors influencing the choice of host species. It is generally desirable to select a breed of host animal that has been bred for dairy use, such as East Friesland sheep, or to introduce dairy stock by breeding of the transgenic line at a later date. In any event, animals of known, good health status should be used.

Fibrinogen produced according to the present invention may be human fibrinogen or fibrinogen of a non-human animal. For medical uses, it is preferred to employ proteins native to the patient. The present invention thus provides fibrinogen for use in both human and veterinary medicine. Cloned DNA molecules encoding the component chains of human fibrinogen are disclosed by Rixon et al. (Biochem. 22: 3237, 1983), Chung et al. (Biochem. 22: 3244, 1983), Chung et al. (Biochem. 22: 3250, 1983), Chung et al. (Adv. Exp. Med. Biol. 281: 39-48, 1990) and Chung et al. (Ann. NY Acad. Sci. 408: 449-456, 1983). Bovine fibrinogen clones are disclosed by Brown et al. (Nuc. Acids Res. 17: 6397, 1989) and Chung et al. (Proc. Natl. Acad. Sci. USA 78: 1466-1470, 1981). Other mammalian fibrinogen clones are disclosed by Murakawa et al. (Thromb. Haemost. 69: 351-360, 1993). Representative sequences of human  $\text{A}\alpha$ ,  $\text{B}\beta$  and  $\gamma$  chain genes are shown in SEQ ID NOS: 1, 3 and 5, respectively. Those skilled in the art will recognize that allelic variants of these sequences will exist; that additional variants can be generated by amino acid substitution, deletion, or insertion; and that such variants are useful within the present invention. In general, it is preferred that any engineered variants comprise only a limited number of amino acid substitutions, deletions, or insertions, and that any substitutions are conservative. Thus, it is preferred to produce fibrinogen chain polypeptides that are at least 90%, preferably at least 95, and more preferably 99% or more identical in sequence to the corresponding native chains. The term " $\gamma$  chain" is meant to include the alternatively spliced  $\gamma'$  chain of fibrinogen (Chung et al., Biochem. 23: 4232-4236, 1984). A human  $\gamma'$  chain amino acid sequence is shown in SEQ ID NO: 6. The shorter  $\gamma$  chain is produced by alternative splicing at nucleotides 9511 and 10054 of SEQ ID NO: 5, resulting in translation terminating after nucleotide 10065 of SEQ ID NO: 5.

To obtain expression in the mammary gland, a transcription promoter from a milk protein gene is used. Milk protein genes include those genes encoding caseins, beta-lactoglobulin (BLG),  $\alpha$ -lactalbumin, and whey acidic protein. The beta-lactoglobulin promoter is preferred. In the case of the ovine

beta-lactoglobulin gene, a region of at least the proximal 406 bp of 5' flanking sequence of the ovine BLG gene (contained within nucleotides 3844 to 4257 of SEQ ID NO:7) will generally be used. Larger portions at the 5' flanking sequence, up to about 5 kbp, are preferred. A larger DNA segment encompassing the 5' flanking promoter region and the region encoding the 5' non-coding portion of the beta-lactoglobulin gene (contained within nucleotides 1 to 4257 of SEQ ID NO:7) is particularly preferred. See Whitelaw et al., *Biochem J.* 28: 31-39, 1992. Similar fragments of promoter DNA from other species are also suitable.

Other regions of the beta-lactoglobulin gene may also be incorporated in constructs, as may genomic regions of the gene to be expressed. It is generally accepted in the art that constructs lacking introns, for example, express poorly in comparison with those that contain such DNA sequences (see Brinster et al., *Proc. Natl. Acad. Sci. USA* 85: 836-840, 1988; Palmiter et al., *Proc. Natl. Acad. Sci. USA* 88: 478-482, 1991; Whitelaw et al., *Transgenic Res.* 1: 3-13, 1991; WO 89/01343; WO 91/02318). In this regard, it is generally preferred, where possible, to use genomic sequences containing all or some of the native introns of a gene encoding the protein or polypeptide of interest. Within certain embodiments of the invention, the further inclusion of at least some introns from the beta-lactoglobulin gene is preferred. One such region is a DNA segment which provides for intron splicing and RNA polyadenylation from the 3' non-coding region of the ovine beta-lactoglobulin gene. When substituted for the natural 3' non-coding sequences of a gene, this ovine beta-lactoglobulin segment can both enhance and stabilize expression levels of the protein or polypeptide of interest. Within other embodiments, the region surrounding the initiation ATG of one or more of the fibrinogen sequences is replaced with corresponding sequences from a milk specific protein gene. Such replacement provides a putative tissue-specific initiation environment to enhance expression. It is convenient to replace the entire fibrinogen chain pre-pro and 5' non-coding sequences with those of, for example, the BLG gene, although smaller regions may be replaced.

For expression of fibrinogen, DNA segments encoding each of the three component polypeptide chains of fibrinogen are operably linked to additional DNA segments required for their expression to produce expression units. Such additional segments include the above-mentioned milk protein gene promoter, as well as sequences which provide for termination of transcription and polyadenylation of mRNA. The expression units will further include a DNA segment encoding a secretion signal operably linked to the segment encoding the fibrinogen polypeptide chain. The secretion signal may be a native fibrinogen secretion signal or may be that of another protein, such as a milk protein. The term "secretion signal" is used herein to denote that portion of a protein that directs it through the secretory pathway of a cell to the outside. Secretion signals are most commonly found at the amino-termini of proteins. See, for example, von Heinje, *Nuc. Acids Res.* 14: 4683-4690, 1986; and Meade et al., U.S. Pat. No. 4,873,316, which are incorporated herein by reference.

Construction of expression units is conveniently carried out by inserting a fibrinogen chain sequence into a plasmid or phage vector containing the additional DNA segments, although the expression unit may be constructed by essentially any sequence of ligations. It is particularly convenient to provide a vector containing a DNA segment encoding a milk protein and to replace the coding sequence for the milk protein with that of a fibrinogen chain (including a secretion signal), thereby creating a gene fusion that includes the expression control sequences of the milk protein gene. In any

event, cloning of the expression units in plasmids or other vectors facilitates the amplification of the fibrinogen sequences. Amplification is conveniently carried out in bacterial (e.g. *E. coli*) host cells, thus the vectors will typically include an origin of replication and a selectable marker functional in bacterial host cells.

In view of the size of the fibrinogen chain genes it is most practical to prepare three separate expression units, mix them, and introduce the mixture into the host. However, those skilled in the art will recognize that other protocols may be followed. For example, expression units for the three chains can be introduced individually into different embryos to be combined later by breeding. In a third approach, the three expression units can be linked in a single suitable vector, such as a yeast artificial chromosome or phage P1 clone. Coding sequences for two or three chains can be combined in polycistronic expression units (see, e.g., Levinson et al., U.S. Pat. No. 4,713,339).

The expression unit(s) is(are) then introduced into fertilized eggs (including early-stage embryos) of the chosen host species. Introduction of heterologous DNA can be accomplished by one of several routes, including microinjection (e.g. U.S. Pat. No. 4,873,191), retroviral infection (Jaenisch, *Science* 240: 1468-1474, 1988) or site-directed integration using embryonic stem (ES) cells (reviewed by Bradley et al., *Bio/Technology* 10: 534-539, 1992). The eggs are then implanted into the oviducts or uteri of pseudopregnant females and allowed to develop to term. Offspring carrying the introduced DNA in their germ line can pass the DNA on to their progeny in the normal, Mendelian fashion, allowing the development of transgenic herds. General procedures for producing transgenic animals are known in the art. See, for example, Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1986; Simons et al., *Bio/Technology* 6: 179-183, 1988; Wall et al., *Biol. Reprod.* 32: 645-651, 1985; Buhler et al., *Bio/Technology*: 140-143, 1990; Ebert et al., *Bio/Technology*: 835-838, 1991; Krimpenfort et al., *Bio/Technology* 9: 844-847, 1991; Wall et al. *J. Cell. Biochem.* 49: 113-120, 1992; and WIPO publications WO 88/00239, WO 90/05188, WO 92/11757; and GB 87/00458, which are incorporated herein by reference. Techniques for introducing foreign DNA sequences into mammals and their germ cells were originally developed in the mouse. See, e.g., Gordon et al., *Proc. Natl. Acad. Sci. USA* 77: 7380-7384, 1980; Gordon and Ruddle, *Science* 214: 1244-1246, 1981; Palmiter and Brinster, *Cell* 41: 343-345, 1985; Brinster et al., *Proc. Natl. Acad. Sci. USA* 82: 4438-1442, 1985; and Hogan et al. (*ibid.*). These techniques were subsequently adapted for use with larger animals, including livestock species (see e.g., WIPO publications WO 88/00239, WO 90/05188, and WO 92/11757; and Simons et al., *Bio/Technology* 6: 179-183, 1988). To summarize, in the most efficient route used to date in the generation of transgenic mice or livestock, several hundred linear molecules of the DNA of interest are injected into one of the pro-nuclei of a fertilized egg. Injection of DNA into the cytoplasm of a zygote can also be employed.

It is preferred to obtain a balanced expression of each fibrinogen chain to allow for efficient formation of the mature protein. Ideally, the three expression units should be on the same DNA molecule for introduction into eggs. This approach, however, may generate technical problems at, for example, the injection and manipulation stages. For example, the size of fibrinogen expression units may necessitate the use of yeast artificial chromosomes (YACs) or phage P1 to amplify and manipulate the DNA prior to injection. If this approach is followed, segments of DNA to be injected, con-

taining all three expression units, would be very large, thus requiring modification of the injection procedure using, for example, larger bore needles. In a more simple approach, a mixture of each individual expression unit is used. It is preferred to combine equimolar amounts of the three expression units, although those skilled in the art will recognize that this ratio may be varied to compensate for the characteristics of a given expression unit. Some expression, generally a reduced level, will be obtained when lesser molar amounts of one or two chains are used, and expression efficiencies can generally be expected to decline in approximate proportion to the divergence from the preferred equimolar ratio. In any event, it is preferred to use a mixture having a ratio of  $\alpha:\beta:\gamma$  expression units in the range of 0.5-1:0.5-1:0.5-1. When the ratio is varied from equimolar, it is preferred to employ relatively more of the  $\beta$  expression unit. Alternatively, one or a mixture of two of the expression units is introduced into individual eggs. However, animals derived by this approach will express only one or two fibrinogen chains. To generate an intact fibrinogen molecule by this approach requires a subsequent breeding program designed to combine all three expression units in individuals of a group of animals.

In general, female animals are superovulated by treatment with follicle stimulating hormone, then mated. Fertilized eggs are collected, and the heterologous DNA is injected into the eggs using known methods. See, for example, U.S. Pat No. 4,873,191; Gordon et al, Proc. Natl. Acad. Sci. USA 77: 7380-7384, 1980; Gordon and Ruddle, Science 214: 1244-1246, 1981; Palmiter and Brinster, Cell. 41: 343-345, 1985; Brinster et al., Proc. Natl. Acad. Sci. USA 82: 4438-4442, 1985; Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, 1986; Simons et al. Bio/Technology 6:179-183, 1988; Wall et al., Biol. Reprod. 32: 645-651, 1985; Buhler et al., Bio/Technology 8: 140-143, 1990; Ebert et al., Bio/Technology 9: 835-838, 1991; Krimpenfort et al., Bio/Technology 9: 844-847, 1991; Wall et al., J. Cell. Biochem. 49: 113-120, 1992; WIPO publications WO 88/00239, WO 90/05118, and WO 92/11757; and GB 87/00458, which are incorporated herein by reference.

For injection into fertilized eggs, the expression units are removed from their respective vectors by digestion with appropriate restriction enzymes. For convenience, it is preferred to design the vectors so that the expression units are removed by cleavage with enzymes that do not cut either within the expression units or elsewhere in the vectors. The expression units are recovered by conventional methods, such as electro-elution followed by phenol extraction and ethanol precipitation, sucrose density gradient centrifugation, or combinations of these approaches.

DNA is injected into eggs essentially as described in Hogan et al., *ibid.* In a typical injection, eggs in a dish of an embryo culture medium are located using a stereo room microscope ( $\times 50$  or  $\times 63$  magnification preferred). Suitable media include Hepes (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) or bicarbonate buffered media such as M2 or M16 (available from Sigma Chemical Co., St Louis, USA) or synthetic oviduct medium (disclosed below). The eggs are secured and transferred to the center of a glass slide on an injection rig using, for example, a drummond pipette complete with capillary tube. Viewing at lower (e.g.  $\times 4$ ) magnification is used at this stage. Using the holding pipette of the injection rig, the eggs are positioned centrally on the slide. Individual eggs are sequentially secured to the holding pipette for injection. For each injection process, the holding pipette/egg is positioned in the center of the viewing field. The injection needle is then positioned directly below the egg.

Preferably using  $\times 40$  Nomarski objectives, both manipulator heights are adjusted to focus both the egg and the needle. The pronuclei are located by rotating the egg and adjusting the holding pipette assembly as necessary. Once the pronucleus has been located, the height of the manipulator is altered to focus the pronuclear membrane. The injection needle is positioned below the egg such that the needle tip is in a position below the center of the pronucleus. The position of the needle is then altered using the injection manipulator assembly to bring the needle and the pronucleus into the same focal plane. The needle is moved, via the joy stick on the injection manipulator assembly, to a position to the right of the egg. With a short, continuous jabbing movement, the pronuclear membrane is pierced to leave the needle tip inside the pronucleus. Pressure is applied to the injection needle via the glass syringe until the pronucleus swells to approximately twice its volume. At this point, the needle is slowly removed. Reverting to lower (e.g.  $\times 4$ ) magnification, the injected egg is moved to a different area of the slide, and the process is repeated with another egg.

After the DNA is injected, the eggs may be cultured to allow the pronuclei to fuse, producing one-cell or later stage embryos. In general, the eggs are cultured at approximately the body temperature of the species used in a buffered medium containing balanced salts and serum. Surviving embryos are then transferred to pseudopregnant recipient females, typically by inserting them into the oviduct or uterus, and allowed to develop to term. During embryogenesis, the injected DNA integrates in a random fashion in the genomes of a small number of the developing embryos.

Potential transgenic offspring are screened via blood samples and/or tissue biopsies. DNA is prepared from these samples and examined for the presence of the injected construct by techniques such as polymerase chain reaction (PCR; see Mullis, U.S. Pat. No. 4,683,202) and Southern blotting (Southern, J. Mol. Biol. 98:503, 1975; Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, 1982). Founder transgenic animals, or G0s, may be wholly transgenic, having transgenes in all of their cells, or mosaic, having transgenes in only a subset of cells (see, for example, Wilkie et al., Develop. Biol. 118: 9-18, 1986). In the latter case, groups of germ cells may be wholly or partially transgenic. In the latter case, the number of transgenic progeny from a founder animal will be less than the expected 50% predicted from Mendelian principles. Founder G0 animals are grown to sexual maturity and mated to obtain offspring, or G1s. The G1s are also examined for the presence of the transgene to demonstrate transmission from founder G0 animals. In the case of male G0s, these may be mated with several non-transgenic females to generate many offspring. This increases the chances of observing transgene transmission. Female G0 founders may be mated naturally, artificially inseminated or superovulated to obtain many eggs which are transferred to surrogate mothers. The latter course gives the best chance of observing transmission in animals having a limited number of young. The above-described breeding procedures are used to obtain animals that can pass the DNA on to subsequent generations of offspring in the normal, Mendelian fashion, allowing the development of, for example, colonies (mice), flocks (sheep), or herds (pigs, goats and cattle) of transgenic animals.

The milk from lactating G0 and G1 females is examined for the expression of the heterologous protein using immunological techniques such as ELISA (see Harlow and Lane, Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, 1988) and Western blotting (Towbin et al., Proc. Natl. Acad. Sci. USA 76: 4350-4354, 1979). For a variety of rea-

sons known in the art, expression levels of the heterologous protein will be expected to differ between individuals.

A satisfactory family of animals should satisfy three criteria: they should be derived from the same founder G0 animal; they should exhibit stable transmission of the transgene; and they should exhibit stable expression levels from generation to generation and from lactation to lactation of individual animals. These principles have been demonstrated and discussed (Carver et al., *Bio/Technology* 11: 1263-1270, 1993). Animals from such a suitable family are referred to as a "line." Initially, male animals, G0 or G1, are used to derive a flock or herd of producer animals by natural or artificial insemination. In this way, many female animals containing the same transgene integration event can be quickly generated from which a supply of milk can be obtained.

The fibrinogen is recovered from milk using standard practices such as skimming, precipitation, filtration and protein chromatography techniques.

Fibrinogen produced according to the present invention is useful within human and veterinary medicine, such as in the formulation of surgical adhesives. Adhesives of this type are known in the art. See, for example, U.S. Pat. Nos. 4,377,572; 4,442,655; 4,462,567; and 4,627,879, which are incorporated herein by reference. In general, fibrinogen and factor XIII are combined to form a first component that is mixed just prior to use with a second component containing thrombin. The thrombin converts the fibrinogen to fibrin, causing the mixture to gel, and activates the factor XIII. The activated factor XIII cross links the fibrin to strengthen and stabilize the adhesive matrix. Such adhesives typically contain from about 30 mg/ml to about 100 mg/ml fibrinogen and from about 50 µg/ml to about 500 µg/ml factor XIII. They may also contain additional ingredients, such as aprotinin, albumin, fibrinectin, bulking agents, and solubilizers. Methods for producing factor XIII are known in the art. See, for example, U.S. Pat. No. 5,204,447. The fibrinogen is also useful for coating surfaces of polymeric articles, e.g. synthetic vascular grafts, as disclosed in U.S. Pat. No. 5,272,074 (incorporated herein by reference).

The invention is further illustrated by the following non-limiting examples.

## EXAMPLES

### Example I

The multiple cloning site of the vector pUC18 (Yanisch-Perron et al., *Gene* 33:103-119, 1985) was removed and replaced with a synthetic double stranded oligonucleotide (the strands of which are shown in SEQ ID NO: 8 and SEQ ID NO: 27) containing the restriction sites Pvu I/Mlu I/Eco RV/Xba I/Pvu I/Mlu I, and flanked by 5' overhangs compatible with the restriction sites Eco RI and Hind III. pUC18 was cleaved with both Eco RI and Hind III, the 5' terminal phosphate groups were removed with calf intestinal phosphatase, and the oligonucleotide was ligated into the vector backbone. The DNA sequence across the junction was confirmed by sequencing, and the new plasmid was called pUCPM.

The β-lactoglobulin (BLG) gene sequences from pSS1tgXS (disclosed in WIPO publication WO 88/00239) were excised as a Sal I-Xba I fragment and recloned into the vector pUCPM that had been cut with Sal I and Xba I to construct vector pUCXS. pUCXS is thus a pUC18 derivative containing the entire BLG gene from the Sal I site to the Xba I site of phage SS1 (Ali and Clark, *J. Mol. Biol.* 199: 415-426, 1988).

The plasmid pSS1tgSE (disclosed in WIPO publication WO 88/00239) contains a 1290 bp BLG fragment flanked by Sph I and EcoR I restriction sites, a region spanning a unique Not I site and a single Pvu II site which lies in the 5' untranslated leader of the BLG mRNA. Into this Pvu II site was ligated a double stranded, 8 bp DNA linker (5'-GGATATCC-3') encoding the recognition site for the enzyme Eco RV. This plasmid was called pSS1tgSE/RV. DNA sequences bounded by Sph I and Not I restriction sites in pSS1tgSE/RV were excised by enzymatic digestion and used to replace the equivalent fragment in pUCXS. The resulting plasmid was called pUCXSRV. The sequence of the BLG insert in pUCXSRV is shown in SEQ ID NO: 7, with the unique Eco RV site at nucleotide 4245 in the 5' untranslated leader region of the BLG gene. This site allows insertion of any additional DNA sequences under the control of the BLG promoter 3' to the transcription initiation site.

Using the primers BLGAMP3 (5'-TGG ATC CCC TGC CGG TGC CTC TGG-3'; SEQ ID NO: 9) and BLGAMP4 (5'-AAC GCG TCA TCC TCT GTG AGC CAG-3'; SEQ ID NO: 10) a PCR fragment of approximately 650 bp was produced from sequences immediately 3' to the stop codon of the BLG gene in pUCXSRV. The PCR fragment was engineered to have a BamH I site at its 5' end and an Mlu I site at its 3' end and was cloned as such into BamH I and Mlu I cut pGEM7zf (+) (Promega) to give pDAM200(+).

pUCXSRV was digested with Kpn I, and the largest, vector containing band was gel purified. This band contained the entire pUC plasmid sequences and some 3' non-coding sequences from the BLG gene. Into this backbone was ligated the small Kpn I fragment from pDAM200(+) which, in the correct orientation, effectively engineered a BamH I site at the extreme 5' end of the 2.6 Kbp of the BLG 3' flanking region. This plasmid was called pBLAC200. A 2.6 Kbp Cla I-Xba I fragment from pBLAC200 was ligated into Cla I-Xba I cut pSP72 vector (Promega), thus placing an EcoR V site immediately upstream of the BLG sequences. This plasmid was called pBLAC210.

The 2.6 Kbp Eco RV-Xba I fragment from pBLAC210 was ligated into Eco RV-Xba I cut pUCXSRV to form pMAD6. This, in effect, excised all coding and intron sequences from pUCXSRV, forming a BLG minigene consisting of 4.3 Kbp of 5' promoter and 2.6 Kbp of 3' downstream sequences flanking a unique EcoR V site. An oligonucleotide linker (ZC6839: ACTACGTAGT; SEQ ID NO: 11) was inserted into the Eco RV site of pMAD6. This modification destroyed the Eco RV site and created a Sna BI site to be used for cloning purposes. The vector was designated pMAD6-Sna. Messenger RNA initiates upstream of the Sna BI site and terminates downstream of the Sna BI site. The precursor transcript will encode a single BLG-derived intron, intron 6, which is entirely within the 3' untranslated region of the gene.

### Example II

Clones encoding the individual fibrinogen chains were obtained from the laboratory of Dr. Earl W. Davie, University of Washington, Seattle. A genomic fibrinogen Aα-chain clone (Chung et al., 1990, *ibid.*) was obtained from the plasmid BS4. This plasmid contains the Aα clone inserted into the Sal I and Bam HI sites of the vector pUC18, but lacks the coding sequence for the first four amino acids of the Aα chain. A genomic Bβ-chain DNA (Chung et al., *ibid.*) was isolated from a lambda Charon 4A phage clone (designated βλ4) as two EcoRI fragments of ca. 5.6 Kbp each. The two fragments were cloned separately into pUC19 that had been digested with Eco RI and treated with calf intestinal phosphatase. The

resulting clones were screened by digestion with the restriction enzyme Pvu II to distinguish plasmids with the 5' and 3' Bp inserts (designated Beta5'RI/puc and Beta3'RI/puc, respectively). Genomic  $\gamma$ -chain clones were isolated as described by Rixon et al. (Biochemistry 24: 2077-2086, 1985). Clone py12A9 comprises 5' non-coding sequences and approximately 4535 bp of  $\gamma$ -chain coding sequence. Clone py12F3 comprises the remaining coding sequence and 3' non-coding nucleotides. Both are pBR322-based plasmids with the fibrinogen sequences inserted at the EcoRI site. These plasmids were used as templates for the respective PCR reactions.

The fibrinogen chain coding sequences were tailored for insertion into expression vectors using the polymerase chain reaction (PCR) as generally described by Mullis (U.S. Pat. No. 4,683,202). This procedure removed native 5' and 3' untranslated sequences, added a 9 base sequence (CCT GCA GCC) upstream of the first ATG of each coding sequence, supplied the first four codons for the  $\alpha$ -chain sequence, removed an internal Mlu I site in the  $\alpha$ -chain sequence and added restriction sites to facilitate subsequent cloning steps.

Referring to FIG. 1, the 5' end of the  $\alpha$ -chain coding sequence was tailored in a PCR reaction containing 20 pmole for each of primers ZC6632 (SEQ ID NO: 12) and ZC6627 (SEQ ID NO: 13), approximately 10 ng of plasmid BS4 template DNA, 10  $\mu$ l of a mix containing 2.5 mM each dNTP, 7.5  $\mu$ l 10 $\times$  Pyrococcus furiosus (Pfu) DNA polymerase buffer #1 (200 mM Tris-HCl, pH 8.2, 100 mM KCl, 60 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM MgCl<sub>2</sub>, 1% Triton X-100, 100  $\mu$ g/ml nuclease free bovine serum albumin (Stratagene, La Jolla, Calif.), and water to 75  $\mu$ l. The mixture was heated to 94° C. in a DNA thermal cycler (Perkin-Elmer Corp., Norwalk, Conn.). To the heated mixture was added 25  $\mu$ l of a mixture containing 2.5  $\mu$ l 10 $\times$ Pfu buffer #1, 22  $\mu$ l H<sub>2</sub>O and 1  $\mu$ l 2.5 units/ $\mu$ l Pfu DNA polymerase (Stratagene). The reactions were run in a DNA thermal cycler (Perkin-Elmer) for five cycles of 94°, 45 seconds; 40°, 90 seconds; 72°, 120 seconds; 20 cycles of 94°, 45 seconds; 45°, 90 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. The 5' PCR-generated fragment was digested with Bam HI and Hind III, and the Bam HI-Hind III fragment was then ligated to an internal 2.91 Kbp Hind III-Xba I fragment and Bam HI, Xba I-digested pUC18. PCR-generated exon sequences were sequenced.

Referring again to FIG. 1, the 3' end of the  $\alpha$ -chain coding sequence was tailored in a series of steps in which the Mlu I site 563 bases upstream from the stop codon of the  $\alpha$ -chain sequence was mutated using an overlap extension PCR reaction (Ho et al., Gene 77: 51-59, 1989). In the first reaction 40 pmole of each of primers ZC6521 (SEQ ID NO: 14) and ZC6520 (SEQ ID NO: 15) were combined with approximately 10 ng of plasmid BS4 template DNA in a reaction mixture as described above. The reaction was run for 5 cycles of 94°, 45 seconds; 40°, 60 seconds; 72°, 120 seconds; 15 cycles of 94°, 45 seconds; 45°, 60 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. A second reaction was carried out in the same manner using 40 pmole of each of primers ZC6519 (SEQ ID NO: 16) and ZC6518 (SEQ ID NO: 17) and BS4 as template. The PCR-generated DNA fragments from the first and second reactions were isolated by gel electrophoresis and elution from the gel. Approximately 1/10 of each recovered reaction product was combined with 40 pmole of each of primers ZC6521 (SEQ ID NO: 14) and ZC6518 (SEQ ID NO: 17) in a PCR reaction in which the complementary 3' ends of each fragment (containing the single base change) annealed and served as a primer for the 3' extension of the complementary strand. PCR was carried out using the same reaction conditions as in the first and second 3' PCR steps. The reaction product was then digested with Xba I and

Bam HI, and the Xba I-Bam HI fragment was cloned into Xba I, Bam HI-digested pUC18. PCR-generated exons were sequenced.

As shown in FIG. 1, the 5' Bam HI-Xba I fragment (3.9 Kbp) and the 3' Xba I-Bam HI fragment (1.3 Kbp) were inserted into the Bam HI site of the vector Zem228. Zem228 is a pUC18 derivative comprising a Bam HI cloning site between a mouse MT-1 promoter and SV40 terminator, and a neomycin resistance marker flanked by SV40 promoter and terminator sequences. See European Patent Office Publication EP 319,944 and FIG. 2. The entire  $\alpha$ -chain coding sequence was isolated from the Zem228 vector as an Sna BI fragment, which was inserted into the Sna BI site of the plasmid pMAD6-Sna.

Referring to FIG. 3, the 5' end of the  $\beta$ -chain was tailored by PCR using the oligonucleotides ZC6629 (SEQ ID NO: 18), ZC6630 (SEQ ID NO: 19) and ZC6625 (SEQ ID NO: 20). These primers were used in pairwise combinations (ZC6629+ZC6625 or ZC6630+ZC6625) to generate  $\beta$ -chain coding sequences beginning at the first ATG codon (position 470 in SEQ ID NO: 3)(designated N1-Beta) or the third ATG codon (position 512 in SEQ ID NO: 3)(designated N3-Beta). Approximately 5 ng of Beta5'RI/puc template DNA was combined with 20 pmole of each of the primers (N1-Beta: ZC6629, SEQ ID NO: 18+ZC6625, SEQ ID NO: 20; or N3-Beta: ZC6630, SEQ ID NO: 19+ZC6625, SEQ ID NO: 20) in a reaction mixture as described above. The mixtures were incubated for 5 cycles of 94°, 45 seconds; 40°, 120 seconds; (N1-Beta) or 90 seconds (N3-Beta); 72°, 120 seconds; 20 cycles of 94°, 45 seconds; 45°, 120 seconds; (N1-Beta) or 90 seconds (N3-Beta); 72°, 120 seconds; then incubated at 72° for 7 minutes. The two reaction products N1, 555 bp or N3, 510 bp) were each digested with Eco RI and Bgl II, and the fragments were ligated to the internal Bgl II-Xba I fragment and Eco RI+Xba I-digested pUC19. The 3' end of the  $\beta$ -chain coding sequence was tailored in a reaction mixture as described above using the oligonucleotide primers ZC6626 (SEQ ID NO: 21) and ZC6624 (SEQ ID NO: 22) and approximately 5 ng of Beta3'RI/puc template. The mixtures were incubated for 5 cycles of 94°, 45 seconds; 40°, 90 seconds; 72°, 120 seconds; 15 cycles of 94°, 45 seconds; 45°, 90 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. A 990 bp Bgl II-Eco RI fragment was isolated. This 3' fragment was ligated to the adjacent coding fragment (340 bp, Sph I-Bgl II) and Sph I+Eco RI-digested pUC19. The 3' and 5' PCR-generated exons were sequenced. A third intermediate vector was constructed by combining two internal fragments (4285 bp Xba I-Eco RI and 383 kb Eco RI-Sph I) in Xba I+Sph I-digested pUC19. The entire  $\beta$ -chain coding sequence (two forms) was then assembled by ligating one of the 5' Eco RI-Xba I fragments, the internal Xba I-Sph I fragment, the 3' Sph I-Eco RI fragment and Eco RI-digested vector pUC19. The  $\beta$ -chain coding sequence was then isolated as a 7.6 Kbp Sna BI fragment and inserted into the Sna BI site of pMAD6-Sna.

Referring to FIG. 4, the 5' end of the gamma chain coding sequence was tailored by PCR using the oligonucleotide primers ZC6514 (SEQ ID NO: 23) and ZC6517 (SEQ ID NO: 24) and approximately 50 ng of py12A9 as template. The PCR reaction was run as described above using 40 pM of each primer. The reaction was run for 5 cycles of 94°, 45 seconds; 40°, 60 seconds; 72°, 120 seconds, followed by 15 cycles of 94°, 45 seconds; 45°, 60 seconds; 72°, 120 seconds. The resulting 213 bp fragment was digested with Bam HI and Spe I, and the resulting restriction fragment was ligated with the adjacent downstream 4.4 kb Spe I-Eco RI fragment and Bam HI+Eco RI digested pUC19. The 3' end of the gamma chain coding sequence was tailored using oligonucleotide primers ZC6516 (SEQ ID NO: 25) and ZC6515 (SEQ ID NO: 26) using 40 pM of each primer, approximately 50 ng of py12F3 template and the same thermal cycling schedule as used for the 5' fragment.

The resulting 500 bp fragment was digested with Spe I and Bam HI, and the resulting restriction fragment was ligated with the upstream 2.77 kb Eco RI-Spe I fragment and Eco RI+Bam HI-digested pUC19. All PCR-generated exons were sequenced. The entire  $\gamma'$ -chain coding sequence was then assembled by ligating a 4.5 Kbp Bam HI-Eco RI 5' fragment, a 1.1 Kbp Eco RI-Pst I internal fragment and a 2.14 Kbp Pst I-Xba I 3' fragment in Bam HI+Xba I-digested Zem219b. Zem219b is a pUC18-derived vector containing a mouse metallothionein promoter and a DHFR selectable marker operably linked to an SV40 promoter (FIG. 5). Plasmid Zem219b has been deposited with American Type Culture Collection as an E. coli XL1-blue transformant under Accession No. 68979. The entire  $\gamma$ -chain coding sequence was then isolated as a 7.8 Kbp Sna BI fragment and inserted into the Sna BI site of pMAD6-Sna.

#### Example III

Mice for initial breeding stocks (C57BL6J, CBACA) were obtained from Harlan Olac Ltd. (Bicester, UK). These were mated in pairs to produce F1 hybrid cross (B6CBAF1) for recipient female, superovulated females, stud males and vasectomized males. All animals were kept on a 14 hour light/10 hour dark cycle and fed water and food (Special Diet Services RM3, Edinburgh, Scotland) ad libitum.

Transgenic mice were generated essentially as described in Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1986, which is incorporated herein by reference in its entirety. Female B6CBAF1 animals were superovulated at 4-5 weeks of age by an i.p. injection of pregnant mares' serum gonadotrophin (FOLLIGON, Vet-Drug, Falkirk, Scotland) (5 iu) followed by an i.p. injection of human chorionic gonadotrophin (CHORULON, Vet-Drug, Falkirk, Scotland) (5 iu) 45 hours later. They were then mated with a stud male overnight. Such females were next examined for copulation plugs. Those that had mated were sacrificed, and their eggs were collected for microinjection.

DNA was injected into the fertilized eggs as described in Hogan et al. (ibid.) Briefly, each of the vectors containing the  $\alpha$ ,  $\beta$  and  $\gamma$  expression units was digested with Mlu I, and the expression units were isolated by sucrose gradient centrifugation. All chemicals used were reagent grade (Sigma Chemical Co., St. Louis, Mo., U.S.A.), and all solutions were sterile and nuclease-free. Solutions of 20% and 40% sucrose in 1M NaCl, 20 mM Tris pH 8.0, 5 mM EDTA were prepared using UHP water and filter sterilized. A 30% sucrose solution was prepared by mixing equal volumes of the 20% and 40% solutions. A gradient was prepared by layering 0.5 ml steps of the 40%, 30% and 20% sucrose solutions into a 2 ml polyallomer tube and allowed to stand for one hour. 100  $\mu$ l of DNA solution (max. 8  $\mu$ g DNA) was loaded onto the top of the gradient, and the gradient was centrifuged for 17-20 hours at 26,000 rpm, 15° C. in a Beckman TL100 ultracentrifuge using a TLS-55 rotor (Beckman Instruments, Fullerton, Calif., USA). Gradients were fractionated by puncturing the tube bottom with a 20 ga. needle and collecting drops in a 96 well microliter plate. 3  $\mu$ l aliquots were analyzed on a 1% agarose mini-gel. Fractions containing the desired DNA fragment were pooled and ethanol precipitated overnight at -20° C. in 0.3M sodium acetate. DNA pellets were resuspended in 50-100  $\mu$ l UHP water and quantitated by fluorimetry. The expression units were diluted in Dulbecco's phosphate buffered saline without calcium and magnesium (containing, per liter, 0.2 g KCl, 0.2 g KH<sub>2</sub>PO<sub>4</sub>, 8.0 g NaCl, 1.15 g Na<sub>2</sub>HPO<sub>4</sub>), mixed (using either the N1-Beta or N3-Beta expression unit) in a 1:1:1 molar ratio, concentration adjusted to about 6  $\mu$ g/ml, and injected into the eggs (~2 pl total DNA solution per egg).

Recipient females of 6-8 weeks of age are prepared by mating B6CBAF1 females in natural estrus with vasectomized males. Females possessing copulation plugs are then kept for transfer of microinjected eggs.

Following birth of potential transgenic animals, tail biopsies are taken, under anesthesia, at four weeks of age. Tissue samples are placed in 2 ml of tail buffer (0.3M Na acetate, 50 mM HCl, 1.5 mM MgCl<sub>2</sub>, 10 mM Tris-HCl, pH 8.5, 0.5% NP40, 0.5% Tween 20) containing 200  $\mu$ g/ml proteinase K (Boehringer Mannheim, Mannheim, Germany) and vortexed. The samples are shaken (250 rpm) at 55°-60° for 3 hours to overnight. DNA prepared from biopsy samples is examined for the presence of the injected constructs by PCR and Southern blotting. The digested tissue is vigorously vortexed, and 5  $\mu$ l aliquots are placed in 0.5 ml microcentrifuge tubes. Positive and negative tail samples are included as controls. Forty  $\mu$ l of silicone oil (BDH, Poole, UK) is added to each tube, and the tubes are briefly centrifuged. The tubes are incubated in the heating block of a thermal cycler (e.g. Omni-gene, Hybaid, Teddington, UK) to 95° C. for 10 minutes. Following this, each tube has a 45  $\mu$ l aliquot of PCR mix added such that the final composition of each reaction mix is: 50 mM KCl; 2 mM MgCl<sub>2</sub>; 10 mM Tris-HCl (pH 8.3); 0.01% gelatin; 0.1% NP40, 10% DMSO; 500 nM each primer, 200  $\mu$ M dNTPs; 0.02 U/ $\mu$ l Taq polymerase (Boehringer Mannheim, Mannheim, Germany). The tubes are then cycled through 30 repeated temperature changes as required by the particular primers used. The primers may be varied but in all cases must target the BLG promoter region. This is specific for the injected DNA fragments because the mouse does not have a BLG gene. Twelve  $\mu$ l of 5 $\times$  loading buffer containing Orange G marker dye (0.25% Orange G [Sigma] 15% Ficoll type 400 [Pharmacia Biosystems Ltd., Milton Keynes, UK]) is then added to each tube, and the reaction mixtures are electrophoresed on a 1.6% agarose gel containing ethidium bromide (Sigma) until the marker dye has migrated  $\frac{2}{3}$  of the length of the gel. The gel is visualized with a UV light source emitting a wavelength of 254 nm. Transgenic mice having one or more of the injected DNA fragments are identified by this approach.

Positive tail samples are processed to obtain pure DNA. The DNA samples are screened by Southern blotting using a BLG promoter probe (nucleotides 2523-4253 of SEQ ID NO: 7). Specific cleavages with appropriate restriction enzymes (e.g. Eco RI) allow the distinction of the three constructs containing the  $\alpha$ ,  $\beta$  and  $\gamma$  sequences.

Southern blot analysis of transgenic mice prepared essentially as described above demonstrated that more than 50% of progeny contained all three fibrinogen sequences. Examination of milk from positive animals by reducing SDS polyacrylamide gel electrophoresis demonstrated the presence of all three protein chains at concentrations up to 1 mg/ml. The amount of fully assembled fibrinogen was related to the ratios of individual subunits present in the milk. No apparent phenotype was associated with high concentrations of human fibrinogen in mouse milk.

#### Example IV

Donor ewes are treated with an intravaginal progesterone-impregnated sponge (CHRONOGEST Goat Sponge, Intervet, Cambridge, UK) on day 0. Sponges are left in situ for ten or twelve days.

Superovulation is induced by treatment of donor ewes with a total of one unit of ovine follicle stimulating hormone (OFSH) (OVAGEN, Horizon Animal Reproduction Technology Pty. Ltd., New Zealand) administered in eight intramuscular injections of 0.125 units per injection starting at 5:00 pm on day -4 and ending at 8:00 am on day 0. Donors are injected intramuscularly with 0.5 ml of a luteolytic agent (ESTRU-

15

MATE, Vet-Drug) on day -4 to cause regression of the corpus luteum, to allow return to estrus and ovulation. To synchronize ovulation, the donor animals are injected intramuscularly with 2 ml of a synthetic releasing hormone analog (RECEPTAL, Vet-Drug) at 5:00 pm on day 0.

Donors are starved of food and water for at least 12 hours before artificial insemination (A.I.). The animals are artificially inseminated by intrauterine laparoscopy under sedation and local anesthesia on day 1. Either xylazine (ROMPUN, Vet-Drug) at a dose rate of 0.05-0.1 ml per 10 kg bodyweight or ACP injection 10 mg/ml (Vet-Drug) at a dose rate of 0.1 ml per 10 kg bodyweight is injected intramuscularly approximately fifteen minutes before A.I. to provide sedation. A.I. is carried out using freshly collected semen from a Poll Dorset ram. Semen is diluted with equal parts of filtered phosphate buffered saline, and 0.2 ml of the diluted semen is injected per uterine horn. Immediately pre- or post-A.I., donors are given an intramuscular injection of AMOXYPEN (Vet-Drug).

Fertilized eggs are recovered on day 2 following starvation of donors of food and water from 5:00 pm on day 1. Recovery is carried out under general anesthesia induced by an intravenous injection of 5% thiopentone sodium (INTRAVAL SODIUM, Vet-Drug) at a dose rate of 3 ml per 10 kg bodyweight. Anesthesia is maintained by inhalation of 1-2% Halothane/O<sub>2</sub>/N<sub>2</sub>O after intubation. To recover the fertilized eggs, a laparotomy incision is made, and the uterus is exteriorized. The eggs are recovered by retrograde flushing of the oviducts with Ovum Culture Medium (Advanced Protein Products, Brierly Hill, West Midlands, UK) supplemented with bovine serum albumin of New Zealand origin. After flushing, the uterus is returned to the abdomen, and the incision is closed. Donors are allowed to recover post-operatively or are euthanized. Donors that are allowed to recover are given an intramuscular injection of Amoxypen L.A. at the manufacturer's recommended dose rate immediately pre- or post-operatively.

Plasmids containing the three fibrinogen chain expression units are digested with Mlu I, and the expression unit fragments are recovered and purified on sucrose density gradients. The fragment concentrations are determined by fluorimetry and diluted in Dulbecco's phosphate buffered saline without calcium and magnesium as described above. The concentration is adjusted to 6 µg/ml and approximately 2 µl of the mixture is microinjected into one pronucleus of each fertilized eggs with visible pronuclei.

All fertilized eggs surviving pronuclear microinjection are cultured in vitro at 38.5° C. in an atmosphere of 5% CO<sub>2</sub>:5% O<sub>2</sub>:90% N<sub>2</sub> and about ~100% humidity in a bicarbonate buffered synthetic oviduct medium (see Table) supplemented with 20% v/v vasectomized ram serum. The serum may be heat inactivated at 56° C. for 30 minutes and stored frozen at -20° C. prior to use. The fertilized eggs are cultured for a suitable period of time to allow early embryo mortality (caused by the manipulation techniques) to occur. These dead or arrested embryos are discarded. Embryos having developed to 5 or 6 cell divisions are transferred to synchronized recipient ewes.

TABLE

Synthetic Oviduct Medium	
Stock A (Lasts 3 Months)	
NaCl	6.29 g
KCl	0.534 g
KH <sub>2</sub> SO <sub>4</sub>	0.162 g
MgSO <sub>4</sub> ·7H <sub>2</sub> O	0.182 g
Penicillin	0.06 g
Sodium Lactate 60% syrup	0.6 mls
Super H <sub>2</sub> O	99.4 mls

16

TABLE-continued

Synthetic Oviduct Medium	
Stock B (Lasts 2 weeks)	
NaHCO <sub>3</sub>	0.21 g
Phenol red	0.001 g
Super H <sub>2</sub> O	10 mls
Stock C (Lasts 2 weeks)	
Sodium Pyruvate	0.051 g
Super H <sub>2</sub> O	10 mls
Stock D (Lasts 3 months)	
CaCl <sub>2</sub> ·2H <sub>2</sub> O	0.262 g
Super H <sub>2</sub> O	10 mls
Stock E (Lasts 3 months)	
Hepes	0.651 g
Phenol red	0.001 g
Super H <sub>2</sub> O	10 mls
To make up 10 mls of Bicarbonate Buffered Medium	
STOCK A	1 ml
STOCK B	1 ml
STOCK C	0.07 ml
STOCK D	0.1 ml
Super H <sub>2</sub> O	7.83 ml
Osmolarity should be 265-285 mOsm. Add 2.5 ml of heat inactivated sheep serum and filter sterilize.	
To make up 10 mls HEPES Buffered Medium	
STOCK A	1 ml
STOCK B	0.2 ml
STOCK C	0.07 ml
STOCK D	0.1 ml
STOCK E	0.8 ml
Super H <sub>2</sub> O	7.83 ml
Osmolarity should be 265-285 mOsm. Add 2.5 ml of heat inactivated sheep serum and filter sterilize.	

Recipient ewes are treated with an intravaginal progesterone-impregnated sponge (Chronogest Ewe Sponge or Chronogest Ewe-Lamb Sponge, Intervet) left in situ for 10 or 12 days. The ewes are injected intramuscularly with 1.5 ml (300 iu) of a follicle stimulating hormone substitute (P.M.S.G., Intervet) and with 0.5 ml of a luteolytic agent (Estrumate, Coopers Pitman-Moore) at sponge removal on day -1. The ewes are tested for estrus with a vasectomized ram between 8:00 am and 5:00 pm on days 0 and 1.

Embryos surviving in vitro culture are returned to recipients (starved from 5:00 pm on day 5 or 6) on day 6 or 7. Embryo transfer is carried out under general anesthesia as described above. The uterus is exteriorized via a laparotomy incision with or without laparoscopy. Embryos are returned to one or both uterine horns only in ewes with at least one suitable corpora lutea. After replacement of the uterus, the abdomen is closed, and the recipients are allowed to recover. The animals are given an intramuscular injection of Amoxypen L.A. at the manufacturer's recommended dose rate immediately pre- or post-operatively.

Lambs are identified by ear tags and left with their dams for rearing. Ewes and lambs are either housed and fed complete diet concentrates and other supplements and or ad lib. hay, or are let out to grass.

Within the first week of life (or as soon thereafter as possible without prejudicing health), each lamb is tested for the presence of the heterologous DNA by two sampling procedures. A 10 ml blood sample is taken from the jugular vein into an EDTA vacutainer. If fit enough, the lambs also have a second 10 ml blood sample taken within one week of the first.

Tissue samples are taken by tail biopsy as soon as possible after the tail has become desensitized after the application of a rubber elastrator ring to its proximal third (usually within 200 minutes after "tailing"). The tissue is placed immediately in a solution of tail buffer. Tail samples are kept at room temperature and analyzed on the day of collection. All lambs are given an intramuscular injection of Amoxypen L.A. at the manufacturer's recommended dose rate immediately post-biopsy, and the cut end of the tail is sprayed with an antibiotic spray.

DNA is extracted from sheep blood by first separating white blood cells. A 10 ml sample of blood is diluted in 20 ml of Hank's buffered saline (HBS; obtained from Sigma Chemical Co.). Ten ml of the diluted blood is layered over 5 ml of Histopaque (Sigma) in each of two 15 ml screw-capped tubes. The tubes are centrifuged at 3000 rpm (2000×g max.), low brake for 15 minutes at room temperature. White cell interfaces are removed to a clean 15 ml tube and diluted to 15 ml in HBS. The diluted cells are spun at 3000 rpm for 10 minutes at room temperature, and the cell pellet is recovered and resuspended in 2-5 ml of tail buffer.

To extract DNA from the white cells, 10% SDS is added to the resuspended cells to a final concentration of 1%, and the tube is inverted to mix the solution. One mg of fresh proteinase K solution is added, and the mixture is incubated overnight at 45° C. DNA is extracted using an equal volume of phenol/chloroform (×3) and chloroform/isoamyl alcohol (×1). The DNA is then precipitated by adding 0.1 volume of 3M NaOAc and 2 volumes of ethanol, and the tube is inverted to mix. The precipitated DNA is spooled out using a clean glass rod with a sealed end. The spool is washed in 70% ethanol, and the DNA is allowed to partially dry, then is redissolved in TE (10 mM Tris-HCl, 1 mM EDTA, pH 7.4).

DNA samples from blood and tail are analyzed by Southern blotting using probes for the BLG promoter region and the fibrinogen chain coding regions.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 27

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5943 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

(B) CLONE: Human Fibrinogen A-alpha chain

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
(B) LOCATION: join(31..84, 1154..1279, 1739..1922, 3055..3200, 3786..5210)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

GTCTAGGAGC CAGCCCCACC CTTAGAAAAG ATG TTT TCC ATG AGG ATC GTC TGC      54
                Met Phe Ser Met Arg Ile Val Cys
                1                5

CTA GTT CTA AGT GTG GTG GGC ACA GCA TGG GTATGGCCCT TTTCATTTTT      104
Leu Val Leu Ser Val Val Gly Thr Ala Trp
    10                15

TCTTCTTGCT TTCTCTCTGG TGTATTATCC ACAAAGAGCC TGGAGGTCAG AGTCTACCTG      164

CTCTATGTCC TGACACTC TTAGCTTTAT GACCCAGGC CTGGGAGGAA ATTTCTGGG      224

TGGGCTTGAC ACCTCAAGAA TACAGGGTAA TATGACACCA AGAGGAAGAT CTTAGATGGA      284

TGAGAGTGTA CAACTACAAG GGAAACTTTA GCATCTGTCA TTCAGTCTTA CCACATTTTG      344

TTTTGTTTTG TTTTAAAAG GGCAAGAATT ATTTGCCATC CTTGTACCTA TAAAGCCTTG      404

GTGCATTATA ATGCTAGTTA ATGGAATAAA ACATTTTATG GTAAGATTTG TTTTCTTTAG      464

TTATTAATTT CTTGCTACTT GTCCATAATA AGCAGAACTT TTAGTGTTAG TACAGTTTTG      524

CTGAAAGGTT ATTGTTGTGT TTGTCAAGAC AGAAGAAAAA GCAAACGAAT TATCTTTGGA      584

AATATCTTTG CAGTATCAGA AGAGATTAGT TAGTAAGGCA ATACGCTTTT CCGCAGTAAT      644

```

-continued

GGTATTCTTT TAAATTATGA ATCCATCTCT AAAGGTTACA TAGAACTTG AAGGAGAGAG	704
GAACATTCAG TTAAGATAGT CTAGGTTTTT CACTGAAGC AGCAATTACA GGAGAAAGAG	764
CTCTACAGTA GTTTTCAACT TTCTGTCTGC AGTCATTAGT AAAATGAAA AGGTAAAT	824
TAACTGATTT TATAGATTCA AATAATTTTC CTTTAGGAT GGATTCTTTA AAACCTCTAA	884
TATTTATCAA ATGCTTATTT AAGTGTCACA CACAGTTAAG AAATTTGTAC ACCTTGCTC	944
CTTTAATTCT CATAACAAC CCATAAAATG GGTCTAGGA TTTCCATTG AAGATAAGAA	1004
ACCTGAAGCT TGCCGAAGCC CTGTGTCTGC TCTCCTAAT CTCTGTGAGA GTGCCATCTC	1064
TTCTGGGGA CTGTAGGCA TGCCACTGTC TCCTCTTCTG GCTAACATTG CTGTTGCTCT	1124
CTTTGTGTA TGTGAATGAA TCTTTAAAG ACT GCA GAT AGT GGT GAA GGT GAC	1177
Thr Ala Asp Ser Gly Glu Gly Asp	
20 25	
TTT CTA GCT GAA GGA GGA GGC GTG CGT GGC CCA AGG GTT GTG GAA AGA	1225
Phe Leu Ala Glu Gly Gly Val Arg Gly Pro Arg Val Val Glu Arg	
30 35 40	
CAT CAA TCT GCC TGC AAA GAT TCA GAC TGG CCC TTC TGC TCT GAT GAA	1273
His Gln Ser Ala Cys Lys Asp Ser Asp Trp Pro Phe Cys Ser Asp Glu	
45 50 55	
GAC TGG GTAAGCAGTC AGCGGGGAA GCAGGAGATT CCTTCCCTCT GATGCTAGAG	1329
Asp Trp	
60	
GGGCTCACAG GCTGACCTGA TTGGTCCCAG AAACTTTTTT AAATAGAAAA TAATTGAATA	1389
GTTACCTACA TAGCAAATA AGAAAAGGAA CTTACTCCA AGAGCACTGT TTATTTACCT	1449
CCCCAATCT GGATCATTAG TGGGTGAACA GACAGGATTT CAGTTGCATG CTCAGGCAAA	1509
ACCAGGCTCC TGAGTATTGT GGCCTCAATT TCCTGGCACC TATTTATGGC TAAGTGGACC	1569
CTCATCCAG AGTTTCTCTG CGACCTCTAA CTAGTCTCT TACCTACTTT TAAGCCAAC	1629
TATCTGGAAG AGAAAAGGTA GGAAGAAATG GGGGCTGCAT GGAAACATGC AAAATTATTC	1689
TGAATCTGAG AGATAGATCC TTAAGTGAAT TTTCTCCCTT CACTTTCAG AAC TAC	1744
Asn Tyr	
AAA TGC CCT TCT GGC TGC AGG ATG AAA GGG TTG ATT GAT GAA GTC AAT	1792
Lys Cys Pro Ser Gly Cys Arg Met Lys Gly Leu Ile Asp Glu Val Asn	
65 70 75	
CAA GAT TTT ACA AAC AGA ATA AAT AAG CTC AAA AAT TCA CTA TTT GAA	1840
Gln Asp Phe Thr Asn Arg Ile Asn Lys Leu Lys Asn Ser Leu Phe Glu	
80 85 90	
TAT CAG AAG AAC AAT AAG GAT TCT CAT TCG TTG ACC ACT AAT ATA ATG	1888
Tyr Gln Lys Asn Asn Lys Asp Ser His Ser Leu Thr Thr Asn Ile Met	
95 100 105 110	
GAA ATT TTG AGA GGC GAT TTT TCC TCA GCC AAT A GTAAGTATTA	1932
Glu Ile Leu Arg Gly Asp Phe Ser Ser Ala Asn	
115 120	
CATATTTACT TCTTTGACTT TATAACAGAA ACAACAAAA TCCTAAATA ATATGATATC	1992
CGCTTATATC TATGACAATT TCATCCCAA GTACTTAGTG TAGAAACACA TACCTTCATA	2052
ATATCCCTGA AAATTTTAAG AGGGAGCTTT TGTTCGTT ATTTTTTCAA AGTAAAAGAT	2112
GTAACTGAG ATTTGTTAAG GTCACAAAAT AAGTCAGAAT TTTGGATTAA AACAAGAATT	2172
TAAATGTGTT CTTTTCAACA GTATATACTG AAAGTAGGAT GGGTCAGACT CTTTGAGTTG	2232
ATATTTTTGT TTCTGCTTG TAAAGGTGAA AACTGAGAGG TCAAGGAAC TGTCAAAGA	2292
CACAGAGCTG GGAATCAAC TCCCAGACT CACTGAGCTG ATTAGGTAGA TTTTAAAT	2352
TAAAATATAG GGTCAGCTA CGTCATTCTC ACAGTCTACT CATTAGGGTT AGGAAACATT	2412
GCATTCACTC TGGGCATGGA CAGCGAGTCT AGGGAGTCT CAGTTTCTCA AGTTTTGCTT	2472

-continued

TGCCTTTTCA	CACCTTCACA	AACACTTGAC	ATTTAAAATC	AGTGATGCCA	ACACTAGCTG	2532
GCAAGTGAGT	GATCCTGTTG	ACCCAAAACA	GCTTAGGAAC	CATTTCAAAT	CTATAGAGTT	2592
AAAAAGAAAA	GCTCATCAGT	AAGAAAATCC	AATATGTTCA	AGTCCCTTGA	TTAAGGATGT	2652
TATAAAATAA	TTGAAATGCA	ATCAAACCAA	CTATTTTAAAC	TCCAAATTAC	ACCTTTAAAA	2712
TTCCAAAGAA	AGTTCTTCTT	CTATATTTCT	TTGGGATTAC	TAATTGCTAT	TAGGACATCT	2772
TAACTGGCAT	TCATGGAAGG	CTGCAGGGCA	TAACATTATC	CAAAAGTCAA	ATGCCCCATA	2832
GGTTTTGAAC	TCACAGATTA	AACTGTAACC	AAAATAAAAT	TAGGCATATT	TACAAGCTAG	2892
TTTCTTTCTT	TCTTTTTTCT	CTTTCTTTCT	TTCTTTCTTT	CTTTCTTTCT	TTCTTTCTTT	2952
CTTTCTTTCT	TTCTCTTCC	TTCCTTTCTT	CCTTTCTTTT	TTGCTGGCAA	TTACAGACAA	3012
ATCACTCAGC	AGCTACTTCA	ATAACCATAT	TTTCGATTTC	AG	AC CGT GAT AAT	3065
					Asn Arg Asp Asn	125
ACC TAC AAC	CGA GTG TCA	GAG GAT CTG	AGA AGC AGA	ATT GAA GTC	CTG	3113
Thr Tyr Asn	Arg Val Ser	Glu Asp Leu	Arg Ser Arg	Ile Glu Val	Leu	
	130		135	140		
AAG CGC AAA	GTC ATA GAA	AAA GTA CAG	CAT ATC CAG	CTT CTG CAG	AAA	3161
Lys Arg Lys	Val Ile Glu	Lys Val Gln	His Ile Gln	Leu Leu Gln	Lys	
	145	150		155		
AAT GTT AGA	GCT CAG TTG	GTT GAT ATG	AAA CGA CTG	GAG GTAAGTATGT		3210
Asn Val Arg	Ala Gln Leu	Val Asp Met	Lys Arg Leu	Glu		
	160	165	170			
GGCTGTGGTC	CCGAGTGTCC	TTGTTTTTGA	GTAGAGGGAA	AAGGAAGGCG	ATAGTTATGC	3270
ACTGAGTGTC	TACTATATGC	AGAGAAAAGT	GTTATATCCA	TCATCTACCT	AAAAGTAGGT	3330
ATTATTTTCC	TCACTCCACA	GTTGAAGAAA	AAAAAATTCA	GAGATATTAA	GTAAATTTTC	3390
CAACGTACAT	AGATAGTAAT	TCAAAGCAAT	GTTCAGTCCC	TGTCTATTCC	AAGCCATTAC	3450
ATCACCCACAC	CTCTGAGCCC	TCAGCCTGAG	TTCACCAAGG	ATCATTTAAT	TAGCGTTTCC	3510
TTTGAGAGGG	AATAGCACCT	TACTCTTGAT	CCATTCTGAG	GCTAAGATGA	ATTAACAGC	3570
ATCCATTGCT	TATCCTGGCT	AGCCCTGCAA	TACCCAACAT	CTCTTCCACT	GAGGGTGCTC	3630
GATAGGCAGA	AAACAGAGAA	TATTAAGTGG	TAGGTCTCCG	AGTCAAAAAA	AATGAAACCA	3690
GTTTCCAGAA	GGAAAATTAA	CTACCAGGAA	CTCAATAGAC	GTAGTTTATG	TATTTGTATC	3750
TACATTTTCT	CTTTATTTTT	CTCCCCTCTC	TCTAG GTG	GAC ATT GAT	ATT AAG	3803
				Val Asp Ile	Asp Ile Lys	175
ATC CGA TCT	TGT CGA GGG	TCA TGC AGT	AGG GCT TTA	GCT CGT GAA	GTA	3851
Ile Arg Ser	Cys Arg Gly	Ser Cys Ser	Arg Ala Leu	Ala Arg Glu	Val	
	180	185		190		
GAT CTG AAG	GAC TAT GAA	GAT CAG CAG	AAG CAA CTT	GAA CAG GTC	ATT	3899
Asp Leu Lys	Asp Tyr Glu	Asp Gln Gln	Lys Gln Leu	Glu Gln Val	Ile	
	195	200	205			
GCC AAA GAC	TTA CTT CCC	TCT AGA GAT	AGG CAA CAC	TTA CCA CTG	ATA	3947
Ala Lys Asp	Leu Leu Pro	Ser Arg Asp	Arg Gln His	Leu Pro Leu	Ile	
	210	215	220			
AAA ATG AAA	CCA GTT CCA	GAC TTG GTT	CCC GGA AAT	TTT AAG AGC	CAG	3995
Lys Met Lys	Pro Val Pro	Asp Leu Val	Pro Gly Asn	Phe Lys Ser	Gln	
	225	230	235	240		
CTT CAG AAG	GTA CCC CCA	GAG TGG AAG	GCA TTA ACA	GAC ATG CCG	CAG	4043
Leu Gln Lys	Val Pro Pro	Glu Trp Lys	Ala Leu Thr	Asp Met Pro	Gln	
	245	250	255			
ATG AGA ATG	GAG TTA GAG	AGA CCT GGT	GGA AAT GAG	ATT ACT CGA	GGA	4091
Met Arg Met	Glu Leu Glu	Arg Pro Gly	Gly Asn Glu	Ile Thr Arg	Gly	
	260	265	270			

-continued

GGC Gly	TCC Ser	ACC Thr	TCT Ser	TAT Tyr	GGA Gly	ACC Thr	GGA Gly	TCA Ser	GAG Glu	ACG Thr	GAA Glu	AGC Ser	CCC Pro	AGG Arg	AAC Asn	4139
		275					280					285				
CCT Pro	AGC Ser	AGT Ser	GCT Ala	GGA Gly	AGC Ser	TGG Trp	AAC Asn	TCT Ser	GGG Gly	AGC Ser	TCT Gly	GGA Gly	CCT Pro	GGA Gly	AGT Ser	4187
		290				295					300					
ACT Thr	GGA Gly	AAC Asn	CGA Arg	AAC Asn	CCT Pro	GGG Gly	AGC Ser	TCT Ser	GGG Gly	ACT Thr	GGA Gly	GGG Gly	ACT Thr	GCA Ala	ACC Thr	4235
					310				315						320	
TGG Trp	AAA Lys	CCT Pro	GGG Gly	AGC Ser	TCT Ser	GGA Gly	CCT Pro	GGA Gly	AGT Ser	GCT Ala	GGA Gly	AGC Ser	TGG Trp	AAC Asn	TCT Ser	4283
				325					330					335		
GGG Gly	AGC Ser	TCT Ser	GGA Gly	ACT Thr	GGA Gly	AGT Ser	ACT Thr	GGA Gly	AAC Asn	CAA Gln	AAC Asn	CCT Pro	GGG Gly	AGC Ser	CCT Pro	4331
			340					345					350			
AGA Arg	CCT Pro	GGT Gly	AGT Ser	ACC Thr	GGA Gly	ACC Thr	TGG Trp	AAT Asn	CCT Pro	GGC Gly	AGC Ser	TCT Ser	GAA Glu	CGC Arg	GGA Gly	4379
		355					360						365			
AGT Ser	GCT Ala	GGG Gly	CAC His	TGG Trp	ACC Thr	TCT Ser	GAG Glu	AGC Ser	TCT Ser	GTA Val	TCT Ser	GGT Gly	AGT Ser	ACT Thr	GGA Gly	4427
		370				375				380						
CAA Gln	TGG Trp	CAC His	TCT Ser	GAA Glu	TCT Pro	GGA Gly	AGT Ser	TTT Phe	AGG Arg	CCA Pro	GAT Asp	AGC Ser	CCA Pro	GGC Gly	TCT Ser	4475
					390					395					400	
GGG Gly	AAC Asn	GCG Ala	AGG Arg	CCT Pro	AAC Asn	AAC Asn	CCA Pro	GAC Asp	TGG Trp	GGC Gly	ACA Thr	TTT Phe	GAA Glu	GAG Glu	GTG Val	4523
				405				410					415			
TCA Ser	GGA Gly	AAT Asn	GTA Val	AGT Ser	CCA Pro	GGG Gly	ACA Thr	AGG Arg	AGA Arg	GAG Glu	TAC Tyr	CAC His	ACA Thr	GAA Glu	AAA Lys	4571
			420				425						430			
CTG Leu	GTC Val	ACT Thr	TCT Ser	AAA Lys	GGA Gly	GAT Asp	AAA Lys	GAG Glu	CTC Leu	AGG Arg	ACT Thr	GGT Gly	AAA Lys	GAG Glu	AAG Lys	4619
		435				440						445				
GTC Val	ACC Thr	TCT Ser	GGT Gly	AGC Ser	ACA Thr	ACC Thr	ACC Thr	ACG Arg	CGT Arg	CGT Arg	TCA Ser	TGC Cys	TCT Ser	AAA Lys	ACC Thr	4667
		450				455					460					
GTT Val	ACT Thr	AAG Lys	ACT Thr	GTT Val	ATT Ile	GGT Gly	CCT Pro	GAT Asp	GGT Gly	CAC His	AAA Lys	GAA Glu	GTT Val	ACC Thr	AAA Lys	4715
		465			470				475						480	
GAA Glu	GTG Val	GTG Val	ACC Thr	TCC Ser	GAA Glu	GAT Asp	GGT Gly	TCT Ser	GAC Cys	TGT Pro	CCC Pro	GAG Glu	GCA Ala	ATG Met	GAT Asp	4763
				485					490					495		
TTA Leu	GGC Gly	ACA Thr	TTG Leu	TCT Ser	GGC Gly	ATA Ile	GGT Gly	ACT Thr	CTG Leu	GAT Asp	GGG Gly	TTC Phe	CGC Arg	CAT His	AGG Arg	4811
			500					505					510			
CAC His	CCT Pro	GAT Asp	GAA Glu	GCT Ala	GCC Ala	TTC Phe	TTC Phe	GAC Asp	ACT Thr	GCC Ala	TCA Ser	ACT Thr	GGA Gly	AAA Lys	ACA Thr	4859
		515					520						525			
TTC Phe	CCA Pro	GGT Gly	TTC Phe	TTC Phe	TCA Ser	CCT Pro	ATG Met	TTA Leu	GGA Gly	GAG Glu	TTT Phe	GTC Val	AGT Ser	GAG Glu	ACT Thr	4907
		530				535						540				
GAG Glu	TCT Ser	AGG Arg	GGC Gly	TCA Ser	GAA Glu	TCT Ser	GGC Gly	ATC Ile	TTC Phe	ACA Thr	AAT Asn	ACA Thr	AAG Lys	GAA Glu	TCC Ser	4955
		545			550					555					560	
AGT Ser	TCT Ser	CAT His	CAC His	CCT Pro	GGG Gly	ATA Ile	GCT Ala	GAA Glu	TTC Phe	CCT Pro	TCC Ser	CGT Arg	GGT Gly	AAA Lys	TCT Ser	5003
				565					570					575		
TCA Ser	AGT Ser	TAC Tyr	AGC Ser	AAA Lys	CAA Gln	TTT Phe	ACT Thr	AGT Ser	AGC Ser	ACG Thr	AGT Ser	TAC Tyr	AAC Asn	AGA Arg	GGA Gly	5051
				580				585						590		

-continued

GAC TCC ACA TTT GAA AGC AAG AGC TAT AAA ATG GCA GAT GAG GCC GGA	5099
Asp Ser Thr Phe Glu Ser Lys Ser Tyr Lys Met Ala Asp Glu Ala Gly	
595 600 605	
AGT GAA GCC GAT CAT GAA GGA ACA CAT AGC ACC AAG AGA GGC CAT GCT	5147
Ser Glu Ala Asp His Glu Gly Thr His Ser Thr Lys Arg Gly His Ala	
610 615 620	
AAA TCT CGC CCT GTC AGA GGT ATC CAC ACT TCT CCT TTG GGG AAG CCT	5195
Lys Ser Arg Pro Val Arg Gly Ile His Thr Ser Pro Leu Gly Lys Pro	
625 630 635 640	
TCC CTG TCC CCC TAGACTAAGT TAAATATTTT TGCACAGTGT TCCCATGGCC	5247
Ser Leu Ser Pro	
CCTTGCATTT CTTTCTTAAC TCTCTGTAC ACGTCATTGA AACTACACTT TTTTGGTCTG	5307
TTTTTGTGCT AGACTGTAAG TTCCTTGGGG GCAGGGCCTT TGTCTGTCTC ATCTCTGTAT	5367
TCCCAAATGC CTAACAGTAC AGAGCCATGA CTCATAAAT ACATGTTAAA TGGATGAATG	5427
AATTCCTCTG AAACCTTATT TGAGCTTATT TAGTCAAATT CTTTCACTAT TCAAAGTGTG	5487
TGCTATTAGA ATTGTACACC AACTGATTAA TCACATTTTT AGTATGTGTC TCAGTTGACA	5547
TTTAGGTCAG GCTAAATACA AGTTGTGTTA GTATTAAGTG AGCTTAGCTA CCTGTACTGG	5607
TTACTTGCTA TTAGTTTGTG CAAGTAAAT TCCAAATACA TTTGAGGAAA ATCCCCTTTG	5667
CAATTTGTAG GTATAAATAA CCGCTTATTT GCATAAGTTC TATCCCCTG TAAGTGCATC	5727
CTTCCCTAT GGAGGGAAGG AAAGGAGGAA GAAAGAAAGG AAGGAAAGA AACAGTATTT	5787
GCCTATTTA ATCTGAGCCG TGCCTATCTT TGTAAGTTA AATGAGAATA ACTTCTTCCA	5847
ACCAGCTTAA TTTTTTTTTT AGACTGTGAT GATGTCCTCC AAACACATCC TTCAGGTACC	5907
CAAAGTGGCA TTTTCAATAT CAAGCTATCC GGATCC	5943

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 644 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Phe Ser Met Arg Ile Val Cys Leu Val Leu Ser Val Val Gly Thr	
1 5 10 15	
Ala Trp Thr Ala Asp Ser Gly Glu Gly Asp Phe Leu Ala Glu Gly Gly	
20 25 30	
Gly Val Arg Gly Pro Arg Val Val Glu Arg His Gln Ser Ala Cys Lys	
35 40 45	
Asp Ser Asp Trp Pro Phe Cys Ser Asp Glu Asp Trp Asn Tyr Lys Cys	
50 55 60	
Pro Ser Gly Cys Arg Met Lys Gly Leu Ile Asp Glu Val Asn Gln Asp	
65 70 75 80	
Phe Thr Asn Arg Ile Asn Lys Leu Lys Asn Ser Leu Phe Glu Tyr Gln	
85 90 95	
Lys Asn Asn Lys Asp Ser His Ser Leu Thr Thr Asn Ile Met Glu Ile	
100 105 110	
Leu Arg Gly Asp Phe Ser Ser Ala Asn Asn Arg Asp Asn Thr Tyr Asn	
115 120 125	
Arg Val Ser Glu Asp Leu Arg Ser Arg Ile Glu Val Leu Lys Arg Lys	
130 135 140	
Val Ile Glu Lys Val Gln His Ile Gln Leu Leu Gln Lys Asn Val Arg	
145 150 155 160	

-continued

---

Ala Gln Leu Val Asp Met Lys Arg Leu Glu Val Asp Ile Asp Ile Lys  
165 170 175

Ile Arg Ser Cys Arg Gly Ser Cys Ser Arg Ala Leu Ala Arg Glu Val  
180 185 190

Asp Leu Lys Asp Tyr Glu Asp Gln Gln Lys Gln Leu Glu Gln Val Ile  
195 200 205

Ala Lys Asp Leu Leu Pro Ser Arg Asp Arg Gln His Leu Pro Leu Ile  
210 215 220

Lys Met Lys Pro Val Pro Asp Leu Val Pro Gly Asn Phe Lys Ser Gln  
225 230 235 240

Leu Gln Lys Val Pro Pro Glu Trp Lys Ala Leu Thr Asp Met Pro Gln  
245 250 255

Met Arg Met Glu Leu Glu Arg Pro Gly Gly Asn Glu Ile Thr Arg Gly  
260 265 270

Gly Ser Thr Ser Tyr Gly Thr Gly Ser Glu Thr Glu Ser Pro Arg Asn  
275 280 285

Pro Ser Ser Ala Gly Ser Trp Asn Ser Gly Ser Ser Gly Pro Gly Ser  
290 295 300

Thr Gly Asn Arg Asn Pro Gly Ser Ser Gly Thr Gly Gly Thr Ala Thr  
305 310 315 320

Trp Lys Pro Gly Ser Ser Gly Pro Gly Ser Ala Gly Ser Trp Asn Ser  
325 330 335

Gly Ser Ser Gly Thr Gly Ser Thr Gly Asn Gln Asn Pro Gly Ser Pro  
340 345 350

Arg Pro Gly Ser Thr Gly Thr Trp Asn Pro Gly Ser Ser Glu Arg Gly  
355 360 365

Ser Ala Gly His Trp Thr Ser Glu Ser Ser Val Ser Gly Ser Thr Gly  
370 375 380

Gln Trp His Ser Glu Ser Gly Ser Phe Arg Pro Asp Ser Pro Gly Ser  
385 390 395 400

Gly Asn Ala Arg Pro Asn Asn Pro Asp Trp Gly Thr Phe Glu Glu Val  
405 410 415

Ser Gly Asn Val Ser Pro Gly Thr Arg Arg Glu Tyr His Thr Glu Lys  
420 425 430

Leu Val Thr Ser Lys Gly Asp Lys Glu Leu Arg Thr Gly Lys Glu Lys  
435 440 445

Val Thr Ser Gly Ser Thr Thr Thr Thr Arg Arg Ser Cys Ser Lys Thr  
450 455 460

Val Thr Lys Thr Val Ile Gly Pro Asp Gly His Lys Glu Val Thr Lys  
465 470 475 480

Glu Val Val Thr Ser Glu Asp Gly Ser Asp Cys Pro Glu Ala Met Asp  
485 490 495

Leu Gly Thr Leu Ser Gly Ile Gly Thr Leu Asp Gly Phe Arg His Arg  
500 505 510

His Pro Asp Glu Ala Ala Phe Phe Asp Thr Ala Ser Thr Gly Lys Thr  
515 520 525

Phe Pro Gly Phe Phe Ser Pro Met Leu Gly Glu Phe Val Ser Glu Thr  
530 535 540

Glu Ser Arg Gly Ser Glu Ser Gly Ile Phe Thr Asn Thr Lys Glu Ser  
545 550 555 560

Ser Ser His His Pro Gly Ile Ala Glu Phe Pro Ser Arg Gly Lys Ser  
565 570 575

Ser Ser Tyr Ser Lys Gln Phe Thr Ser Ser Thr Ser Tyr Asn Arg Gly  
580 585 590



-continued

---

(ix) FEATURE:  
 (A) NAME/KEY: intron  
 (B) LOCATION: 7253..7870

(ix) FEATURE:  
 (A) NAME/KEY: exon  
 (B) LOCATION: 7871..8102

(ix) FEATURE:  
 (A) NAME/KEY: 3'UTR  
 (B) LOCATION: 8103..8537

(ix) FEATURE:  
 (A) NAME/KEY: misc\_RNA  
 (B) LOCATION: 8538..8878

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: join(470..583, 3258..3449, 3939..4122,  
 5831..5944, 6633..6758, 6967..7252, 7871..8102)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

```

GAATTCATGC CCCTTTTGAA ATAGACTTAT GTCATTGTCA GAAAACATAA GCATTTATGG      60
TATATCATTATGAGATCAGC ATTTTAGTGG TTGCCTTGTG AGTAGGTCAA ATTTACTAAG      120
CTTAGATTTG TTTTCTCACA TATTCTTTCG GAGCTTGTGT AGTTCCACA TTAATTTACC      180
AGAAACAAGA TACACACTCT CTTTGAGGAG TGCCCTAACT TCCCATCATT TTGTCCAATT      240
AAATGAATTG AAGAAATTTA ATGTTTCTAA ACTAGACCAA CAAAGAATAA TAGTTGTATG      300
ACAAGTAAAT AAGCTTTGCT GGGAAGATGT TGCTTAAATG ATAAAATGGT TCAGCCAACA      360
AGTGAACCAA AAATTAATAA TTAACCTAAG AAAGTAACC ATTTCTGAAG TCATTCCTAG      420
CAGAGGACTC AGATATATAT AGGATTGAAG ATCTCTCAGT TAAGTCTAC ATG AAA      475
                               Met Lys
                               1
AGG ATG GTT TCT TGG AGC TTC CAC AAA CTT AAA ACC ATG AAA CAT CTA      523
Arg Met Val Ser Trp Ser Phe His Lys Leu Lys Thr Met Lys His Leu
      5                10                15
TTA TTG CTA CTA TTG TGT GTT TTT CTA GTT AAG TCC CAA GGT GTC AAC      571
Leu Leu Leu Leu Leu Cys Val Phe Leu Val Lys Ser Gln Gly Val Asn
      20                25                30
GAC AAT GAG GAG GTGAATTTT TAAAGCATT TATATTATT AGTAGTATTA      623
Asp Asn Glu Glu
      35
TTAATATAAG ATGTAACATA ATCATATTAT GTGCTTATTT TAATGAAATT AGCATTGCTT      683
ATAGTTATGA AATGGAATTG TTAACCTCTG ACTTATTGTA TTTAAAGAAT GTTTCATAGT      743
ATTTCTTATA TAAAAACAAA GTAATTTCTT GTTTCTAGT TTATCACCTT TGTTTTCTTA      803
AGATGAGGAT GGCTTAGCTA ATGTAAGATG TGTTTTTCTC ACTTGCTATT CTGAGTACTG      863
TGATTTTCAT TTACTTCTAG CAATACAGGA TTACAATTAA GAGGACAAGA TCTGAAAATC      923
TCACAAACTA TAAAATAATA AAAGAGCAGA ATTTTAAGAT AAAAGAAACT GGTGGTAGGT      983
AGATTGTTCT TTGGTGAAGG AAGGTAATAT ATATTGTTAC TGAGATTACT ATTTATAAAA     1043
ATTATAACTA AGCCTAAAAG CAAAATACAT CAAGTGTAAT GATAGAAAAT GAAATATTGC     1103
TTTTTTCAGA TGAAAAGTTC AAATTAGAGT TAGTGTGTAT TGTATTATT AATAGTTATG     1163
AAACACGGTT CAGTCTAATT TATTTATTTG TAGAACAGTT TGTCCTCAAC TATTATTTTT     1223
GCTGACTTAT TGCTGTTAAT TTGCAGTTAC TAAAATACA GAAATGCATT TAGGACAATG     1283
GATATTTAAG AAATTTAAAT TTTATCATCA AACGTATCAT GGCCAAATTT CTTACATATA     1343
GCATAGTATC ATTAACTAG AAATAAGAAT ACACAATAAT ATTTAAATGA AGTGATTCAT     1403
TTCGGATCAT TATTGAGTTT CAAGGGAECT TGAGTGTGTG ACTTATCAGA CTCTACATGT     1463

```

-continued

AAGAACATAT	AGTTAATCTG	GTTGTGTGTG	TAAAAACATA	TGGTTAATCT	GGTTAAGTCT	1523
GGTTAATCAT	ATTAGGTAAG	AAAAATGTAA	AGAATGTGTA	AGACGAAAT	TTTGTAAAGT	1583
ACTCTGCAA	GCACTTTCAC	ATTTCTGCTT	ATCAACTAAA	CCTCACAGAG	ATAGTTTAAT	1643
AGTTTAGGCT	TTAAAATGGA	TTTTGATTAT	TCAACAAGTG	GCCTTCATAA	TTTCTTTAAG	1703
TGTTTTTCTT	TAAGTATATA	CTTCTTTTAA	ATATTTTTTA	AAATTCCTT	TTCTCTAGTA	1763
AAGCCAGACC	ATCCATGCTA	CCTCTCTAGT	GGCACTCTGA	AATAAAAAGA	AAATAGTTTT	1823
CTCTGTATA	ATGTATTTG	TAATAAGCAG	ATGAATCACA	TTTCTTAAAA	TTTGTTTTGT	1883
AGAGGGTAAG	CTCTGACTAG	GACCATGACT	TCAATGTGAA	ATATGTATAT	ATCCTCCGAA	1943
TCTTTACATA	TTAAGAAATG	ATATAGTCAA	CTGGTTAAAC	AGGAAATCT	GGAACAGCCT	2003
GGCTGGGTTT	TAATCTTAGC	ACCATCCTAC	TAAATGTTAA	ATAATATTAT	AATCTAATGA	2063
ATAAATGACA	ATGCAATTC	AAATAGAGTT	CATCTGATGA	CTTCTAGACT	CACAAAATTG	2123
CAAGAGAGCT	CAGTTGTTGC	TCAGTTGTTT	CAATCATGT	CGTTTGTAA	TTTGTAAATTA	2183
AGCTCCAAG	GATGTATAGC	TACTGACAAA	AAAAAAAATG	AGAATGTAGT	TAATCCAAT	2243
CAAACTTTC	CTATTGCAAT	GCGTATTTTC	TGCTTCATTA	TCCTTTAATA	TAATATTTTA	2303
AGTTAGCAAG	TAATTTTAA	TACAATGCAC	AAGCCTTGAG	AATTATTTTA	AATATAAGAA	2363
AATCATAATG	TTTGATAAAG	AAATCATGTA	AGAAATTTCA	AGATAATGGT	TTAACAAATA	2423
ATTTTGTGTA	TAGAAGATAA	GACTAAAAGT	GAAATTCGAA	GTGGAGAGGA	CACTTAAACT	2483
GTAGTACTTG	TTATGTGTGA	TTCCAGTAAA	AATAGTAATG	AGCACTTATT	ATTGCCAAGT	2543
ACTGTTCGA	GGGTACCATA	TGCAATAAGT	TATTTAATCC	TTACAATAAT	CTTGTAAAGC	2603
AGATTCAAAC	TATCATTACA	CTTATTTTAC	AGATGAGAAA	ACTGGGGCAC	AGATAAAGCA	2663
ACTTGCCCAA	GGTCTCATAG	CTGTAAGTCA	ACCCTACGGT	CAAGACCTAC	AAGTAGCCGA	2723
GCTCCAGAGT	ACATTATGAG	GGTCAAAGAT	TGTCTTATTA	CAAATAAATT	CCAAGTAGAA	2783
TCAACCTTTA	ATAAGTCTTT	AATGTCTCTT	AAATATGTTT	ATATAGGAGT	CTAATCACCA	2843
ATTCACAAAA	ATGAAAGTAG	GGAAATGATT	AACAATAATC	ATAGGAATCT	AACAATCCAA	2903
GTGGCTTGAG	AATATTCATT	CTTCTTGACA	GTATAGATTC	TTTACAATTT	CGTAAGTTCC	2963
AATGTATGTT	TTAGGAATAT	GAGGTCAATTA	CTATTCATAA	TCTGATACAG	CTTTATCCTA	3023
AGGCCTCTCT	TTAAAACATA	CACTGCATCA	TAGCTTTTTT	GTGCAGTTGG	TCTTCTACT	3083
GTTACTGAAC	AGTAAGCAAC	CTACAGATTC	ACTATCACCA	ACCAGCCAGT	TGATGGATCT	3143
TAAGCAAATT	ATCAAGCTTG	TGATAACCTA	AATTATAAAA	TGAGGGTGTT	GGAATAGTTA	3203
CATTCCAAAT	CTTCTATAAC	ACTCTGTATT	ATATTTCTGC	CTCATTCTTT	GTAG GGT Gly	3260
TTC TTC AGT GCC CGT GGT CAT CGA CCC CTT GAC AAG AAG AGA GAA GAG	3308					
Phe Phe Ser Ala Arg Gly His Arg Pro Leu Asp Lys Lys Arg Glu Glu						
40 45 50 55						
GCT CCC AGC CTG AGG CCT GCC CCA CCG CCC ATC AGT GGA GGT GGC TAT	3356					
Ala Pro Ser Leu Arg Pro Ala Pro Pro Pro Ile Ser Gly Gly Gly Tyr						
60 65 70						
CGG GCT CGT CCA GCC AAA GCA GCT GCC ACT CAA AAG AAA GTA GAA AGA	3404					
Arg Ala Arg Pro Ala Lys Ala Ala Ala Thr Gln Lys Lys Val Glu Arg						
75 80 85						
AAA GCC CCT GAT GCT GGA GGC TGT CTT CAC GCT GAC CCA GAC CTG	3449					
Lys Ala Pro Asp Ala Gly Gly Cys Leu His Ala Asp Pro Asp Leu						
90 95 100						
GTGGGTGCAC TGATGTTTCT TGCAGTGGTG GCTCTCTCAT GCAGAGAAAG CCTGTAGTCA	3509					
TGGCAGTCTG CTAATGTTTC ACTGACCCAC ATTACCATCA CTGTTATTTT GTTGTGTTTAT	3569					

-continued

TTTGAAATA AAATTCAAAA CATAAACATA TTGGGCCTTT GGTTTAGGCT TTCTTTCTTG	3629
TTTTCTTTGG TCTGGGCCCA AAATTTCAAA TTAGGATATG TGGGTGCCAC CTTCCATTT	3689
GTATTTTGCC ACTGCCTTTG TTTAGTTGGT AAAATTTTCA TAGCCCAATT ATATTTTTC	3749
TGGGGTAAGT AATATTTTAA ATCTCTATGA GAGTATGATG ATGACTTTCG AATTTCTGGT	3809
CTTACAGAAA ACCAAATAAT AAATTTTAT GTTGGCTAAT CGTATCGCTG AATTTTCCTA	3869
TGTGCTATTT TAACAAATGT CCATGACCCA AATCCTTCAT CTAATGCCTG CTATTTTCTT	3929
TGTTTTTAG GGG GTG TTG TGT CCT ACA GGA TGT CAG TTG CAA GAG GCT	3977
Gly Val Leu Cys Pro Thr Gly Cys Gln Leu Gln Glu Ala	
105 110 115	
TTG CTA CAA CAG GAA AGG CCA ATC AGA AAT AGT GTT GAT GAG TTA AAT	4025
Leu Leu Gln Gln Glu Arg Pro Ile Arg Asn Ser Val Asp Glu Leu Asn	
120 125 130	
AAC AAT GTG GAA GCT GTT TCC CAG ACC TCC TCT TCT TCC TTT CAG TAC	4073
Asn Asn Val Glu Ala Val Ser Gln Thr Ser Ser Ser Ser Phe Gln Tyr	
135 140 145	
ATG TAT TTG CTG AAA GAC CTG TGG CAA AAG AGG CAG AAG CAA GTA AAA G	4122
Met Tyr Leu Leu Lys Asp Leu Trp Gln Lys Arg Gln Lys Gln Val Lys	
150 155 160	
GTAGATATCC TTGTGCTTTC CATTGCTTTT TCAGCTATAA AATTGGAACC GTTAGACTGC	4182
CACGAGAATG CATGGTTGTG AGAAGATTAA CATTCTGGG TTAGTGAATA GCATTCATAC	4242
GCTTTTGGGC ACCTTCCCCT GCAACTTGCC AGATAAGCAC TATTCAGCTC TTATTCCCAG	4302
TCTGACATCA GCAAGTGTGA TTTTCTATGA AAAATTCTAC TATGACTCCT TATTTAAGT	4362
ATACAAGAAA CTGTGACTC AGAAGATAAT ATTTACAGAG TGGAAAAAAA CCCCTAGCAT	4422
TTATAGTTTT AACATTTGAG GTTTTGAATG AGAGAGTTAT CCATAATATA TTCAATTGTG	4482
TTGTGGATAA TGACACCTAA CCTGTGAATC TTGAGGTCAG AATGTTGAGT GCTGTTGACT	4542
TGGTGGTCAG GAAACAGCTA GTGCGTGAGC CTGGCACAGG CATCTCAGTG AGTAGCATA	4602
CCACAGTTGG AAATTTTCA AAGAAATCAA AGGAATCATG ACATCTTATA AATTTCAAGG	4662
TTCTGCTATA CTTATGTGAA ATGGATAAAT AAATCAAGCA TATCCACTCT GTAAGATTGA	4722
ACTTCTCAGA TGAAGACCC CAATACTGCT TTCTCCTCTT TTCCCTCACC AAAGAAATAA	4782
ACAACCTATT TCATTTATTA CTGGACACAA TCTTTAGCGT ATACCTATGG TAAATTACTA	4842
GTATGGTGGT TAGGATTTAT GTTAATTTGT ATATGTCATG CGCCAATCA TTTCCACTAA	4902
ATATGACTAT ATATCATAAC TGCTTGGTGA TAGCTCAGTG TTTAATAGTT TATCTCAGA	4962
AAATCAAAAT TGTATAGTTA AATACATTAG TTTTATGAGG CAAAAATGCT AACTATTTCT	5022
ACATAATTTT ATTTTCCAG AT AAT GAA AAT GTA GTC AAT GAG TAC TCC	5071
Asp Asn Glu Asn Val Val Asn Glu Tyr Ser	
165 170	
TCA GAA CTG GAA AAG CAC CAA TTA TAT ATA GAT GAG ACT GTG AAT AGC	5119
Ser Glu Leu Glu Lys His Gln Leu Tyr Ile Asp Glu Thr Val Asn Ser	
175 180 185	
AAT ATC CCA ACT AAC CTT CGT GTG CTT CGT TCA ATC CTG GAA AAC CTG	5167
Asn Ile Pro Thr Asn Leu Arg Val Leu Arg Ser Ile Leu Glu Asn Leu	
190 195 200 205	
AGA AGC AAA ATA CAA AAG TTA GAA TCT GAT GTC TCA GCT CAA ATG GAA	5215
Arg Ser Lys Ile Gln Lys Leu Glu Ser Asp Val Ser Ala Gln Met Glu	
210 215 220	
TAT TGT CGC ACC CCA TGC ACT GTC AGT TGC AAT ATT CCT GTG GTG TCT	5263
Tyr Cys Arg Thr Pro Cys Thr Val Ser Cys Asn Ile Pro Val Val Ser	
225 230 235	

-continued

GGC AAA G GTAACGTATT CATAAACATA TTTTGTAGAGA GTTCCAGAAG AACTCACACA	5320
Gly Lys	
CCAAAAATAA GAGAACAACA ACAACAACAA AAATGCTAAG TGGATTTTCC CAACAGATCA	5380
TAATGACATT ACAGTACATC ATAAAAATAT CCTTAGCCAG TTGTGTTTTG GACTGGCCTG	5440
GTGCATTTGC TGGTTTTGAT GAGCAGGATG GGGCACAGGT AGTCCCAGGG GTGGCTGATG	5500
TGTGCATCTG CGTACTGGCT TGAACAGATG GCAGAACCAC AGATAGATGT AGAAGTTTCT	5560
CCATTTTGTG TGTCTGGGA GCTCATGGAT ATTCCAGGAC ACAAAGGTG GAGAAGAGCT	5620
TTGTTTATCC TCTTAGCAGA TAAACGCTCT CAAAAGTGGG TTGGACTTAC TAAAGTAAAA	5680
TGAAAATCTA ATATTTGTTA TATTATTTTC AAAGTCTAT AATAACACAC TCCTTAGTAA	5740
CTTATGTAAT GTTATTTTAA AGAATTGGTG ACTAAATACA AAGTAATTAT GTCATAAACC	5800
CCTGAACATA ATGTTGTCTT ACATTGCGAG AA TGT GAG GAA ATT ATC AGG AAA	5853
Glu Cys Glu Glu Ile Ile Arg Lys	
240 245	
GGA GGT GAA ACA TCT GAA ATG TAT CTC ATT CAA CCT GAC AGT TCT GTC	5901
Gly Gly Glu Thr Ser Glu Met Tyr Leu Ile Gln Pro Asp Ser Ser Val	
250 255 260	
AAA CCG TAT AGA GTA TAC TGT GAC ATG AAT ACA GAA AAT GGA G	5944
Lys Pro Tyr Arg Val Tyr Cys Asp Met Asn Thr Glu Asn Gly	
265 270 275	
GTAAGCTTTC GACAGTTGTT GACCTGTTGA TCTGTAATTA TTTGGATACC GTAAAATGCC	6004
AGGAAACAAG GCCAGGTGTG GTGGCTCATA CCTGTAATTC CAGCACCTTG GGAGGCCAAA	6064
GTGGGCTGAT AGCTTGAGCC TAGGAGTTTG AACTAGCCT GGGCAACATA ATGAGACCCT	6124
AACTCTACAA AAAAAAAAAA AATACCAAAA AAAAAAAAAA AATCAGCTGT GTTGGTAGTA	6184
TGTGCCTGTA GTCCCAGCTA TCCAGGAGGC TGAGATGGGA GATCACCTGA GCCACAACC	6244
TGGAGTCTTG ATCATGCTAC TGAACGTAG CCTGGGCAAC AGAGGATAGT GAGATCCTGT	6304
CTCAAAAAAA AAAATTAATT AAAAAGCCAG GAAACAAGAC TTAGCTCTAA CATCTAACAT	6364
AGCTGACAAA GGAGTAATTT GATGTGGAAT TCAACCTGAT ATTTAAAAGT TATAAATAT	6424
CTATAATTCA CAATTTGGGG TAAGATAAAG CACTTGCAGT TTCCAAAGAT TTTACAAGTT	6484
TACCTCTCAT ATTTATTTCC TTATTGTGTC TATTTTAGAG CACCAATAT ATACTAAATG	6544
GAATGGACAG GGGATTGAGA TATTATTTTC AAAGTGACAT TATTGTCTGT TGGTTAATAT	6604
ATGCTCTTTT TGTCTGTGTC AACCAAAG GA TGG ACA GTG ATT CAG AAC CGT	6655
Gly Trp Thr Val Ile Gln Asn Arg	
280 285	
CAA GAC GGT AGT GTT GAC TTT GGC AGG AAA TGG GAT CCA TAT AAA CAG	6703
Gln Asp Gly Ser Val Asp Phe Gly Arg Lys Trp Asp Pro Tyr Lys Gln	
290 295 300	
GGA TTT GGA AAT GTT GCA ACC AAC ACA GAT GGG AAG AAT TAC TGT GGC	6751
Gly Phe Gly Asn Val Ala Thr Asn Thr Asp Gly Lys Asn Tyr Cys Gly	
305 310 315	
CTA CCA G GTAACGAACA GGCATGCAAA ATAAAATCAT TCTATTTGAA ATGGGATTTT	6808
Leu Pro	
TTTTAATTAA AAAACATTCA TTGTTGGAAG CCTGTTTTAG GCAGTTAAGA GGAGTTTCCT	6868
GACAAAAATG TGAAGCTAA AGATAAGGGA AGAAAGGCAG TTTTAGTTT CCAAAAATTT	6928
TATTTTGGT GAGAGATTTT ATTTTGTTTT TCTTTTAG GT GAA TAT TGG CTT	6980
Gly Glu Tyr Trp Leu	
320	
GGA AAT GAT AAA ATT AGC CAG CTT ACC AGG ATG GGA CCC ACA GAA CTT	7028
Gly Asn Asp Lys Ile Ser Gln Leu Thr Arg Met Gly Pro Thr Glu Leu	
325 330 335 340	

-continued

TTG ATA GAA ATG GAG GAC TGG AAA GGA GAC AAA GTA AAG GCT CAC TAT Leu Ile Glu Met Glu Asp Trp Lys Gly Asp Lys Val Lys Ala His Tyr 345 350 355	7076
GGA GGA TTC ACT GTA CAG AAT GAA GCC AAC AAA TAC CAG ATC TCA GTG Gly Gly Phe Thr Val Gln Asn Glu Ala Asn Lys Tyr Gln Ile Ser Val 360 365 370	7124
AAC AAA TAC AGA GGA ACA GCC GGT AAT GCC CTC ATG GAT GGA GCA TCT Asn Lys Tyr Arg Gly Thr Ala Gly Asn Ala Leu Met Asp Gly Ala Ser 375 380 385	7172
CAG CTG ATG GGA GAA AAC AGG ACC ATG ACC ATT CAC AAC GGC ATG TTC Gln Leu Met Gly Glu Asn Arg Thr Met Thr Ile His Asn Gly Met Phe 390 395 400	7220
TTC AGC ACG TAT GAC AGA GAC AAT GAC GGC TG GTATGTGTGG Phe Ser Thr Tyr Asp Arg Asp Asn Asp Gly Trp 405 410 415	7262
CACTCTTTGC TCCTGCTTTA AAAATCACAC TAATATCATT ACTCAGAATC ATTAACAATA	7322
TTTTTAATAG CTACCACTTC CTGGGCATT ACTGTCAGCC ACTGTCCTAA GCTCTTTATG	7382
CATCACTCGA AAGCATTTC ACTATAAGGT AGACATTCTT ATTCTCATTT TACAGATGAG	7442
ATTTAGAGAG ATTACGTGAT TTGTCCAATG TCACACAACCT ACCCAGAGAT AAAACTAGAA	7502
TTTGAGCACA GTTACTTTCT GAATAATGAG CATTAGATA AATACCTATA TCTCTATATT	7562
CTAAAGTGTG TGTGAAAACCT TTCATTTTCA TTTCCAGGGT TCTCTGATAC TAAGGGTTGT	7622
AAAAGCTATT ATTCCAGTAT AAAGTAACAA ACACAGTCCC TAGATGGATT GCCACAAAGG	7682
CCCAATTATC TCTCTTTCTT GCTATAGGGC ACAGGAGGTC TTTGGTGTAT TAGTGTGACT	7742
CTATGTATAG CACCCAAAGG AAAGACTACT GTGCACACGA GTGTAGCAGT CTTTATGGG	7802
TAATCTGCAA AACGTAACCT GACCACCGTA GTTCTGTTTC TAATAACGCC AAACACATTT	7862
TCTTTCAG G TTA ACA TCA GAT CCC AGA AAA CAG TGT TCT AAA GAA GAC Leu Thr Ser Asp Pro Arg Lys Gln Cys Ser Lys Glu Asp 420 425	7910
GGT GGT GGA TGG TGG TAT AAT AGA TGT CAT GCA GCC AAT CCA AAC GGC Gly Gly Gly Trp Trp Tyr Asn Arg Cys His Ala Ala Asn Pro Asn Gly 430 435 440	7958
AGA TAC TAC TGG GGT GGA CAG TAC ACC TGG GAC ATG GCA AAG CAT GGC Arg Tyr Tyr Trp Gly Gly Gln Tyr Thr Trp Asp Met Ala Lys His Gly 445 450 455 460	8006
ACA GAT GAT GGT GTA GTA TGG ATG AAT TGG AAG GGG TCA TGG TAC TCA Thr Asp Asp Gly Val Val Trp Met Asn Trp Lys Gly Ser Trp Tyr Ser 465 470 475	8054
ATG AGG AAG ATG AGT ATG AAG ATC AGG CCC TTC TTC CCA CAG CAA Met Arg Lys Met Ser Met Lys Ile Arg Pro Phe Phe Pro Gln Gln 480 485 490	8099
TAGTCCCAA TACGTAGATT TTTGCTCTTC TGTATGTGAC AACATTTTGG TACATTATGT	8159
TATTGGAATT TTCTTTCATA CATTATATTC CTCTAAAACCT CTCAAGCAGA CGTGAGTGTG	8219
ACTTTTTGAA AAAAGTATAG GATAAATTAC ATTAATAATAG CACATGATTT TCTTTTGTTT	8279
TCTTCATTTT TCTTGCTCAC CCAAGAAGTA ACAAAGTAT AGTTTTGACA GAGTTGGTGT	8339
TCATAATTC AGTTCAGTGT GATTGCGAGA ATTTTCAAAT AAGGAAGAGG GGTCTTTTAT	8399
CCTGTGCGTA GGAAAACCAT GACGGAAAGG AAAAAGTAT GTTTAAAAGT CCACCTTTAA	8459
AACTATATTT ATTTATGTAG GATCTGTCAA AGAAAACCTC CAAAAAGATT TATTAATTAA	8519
ACCAGACTCT GTTGCAATAA GTTAATGTTT TCTTGTTTTG TAATCCACAC ATTCAATGAG	8579
TTAGGCTTTG CACTTGTAAAG GAAGGAGAAG CGTTCACAAC CTCAAATAGC TAATAAACCG	8639
GTCTTGAATA TTTGAAGATT TAAAATCTGA CTCTAGGACG GGCACGGTGG CTCACGACTA	8699

-continued

---

TAATCCCAAC ACTTTGGGAG GCTGAGGCGG GCGGTCACAA GGTGAGGAGT TCAAGACCAG 8759  
 CCTGACCAAT ATGGTGAAAC CCCATCTCTA CTAAAAATAC AAAAAATTAGC CAGGCGTGGT 8819  
 GGCAGGTGCC TGTAGTCCC AGCTAGCCTG TGAGGTGGAG ATTGCATTGA GCCAAGATC 8878

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 491 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Lys Arg Met Val Ser Trp Ser Phe His Lys Leu Lys Thr Met Lys  
 1 5 10 15  
 His Leu Leu Leu Leu Leu Cys Val Phe Leu Val Lys Ser Gln Gly  
 20 25 30  
 Val Asn Asp Asn Glu Glu Gly Phe Phe Ser Ala Arg Gly His Arg Pro  
 35 40 45  
 Leu Asp Lys Lys Arg Glu Glu Ala Pro Ser Leu Arg Pro Ala Pro Pro  
 50 55 60  
 Pro Ile Ser Gly Gly Gly Tyr Arg Ala Arg Pro Ala Lys Ala Ala Ala  
 65 70 75 80  
 Thr Gln Lys Lys Val Glu Arg Lys Ala Pro Asp Ala Gly Gly Cys Leu  
 85 90 95  
 His Ala Asp Pro Asp Leu Gly Val Leu Cys Pro Thr Gly Cys Gln Leu  
 100 105 110  
 Gln Glu Ala Leu Leu Gln Gln Glu Arg Pro Ile Arg Asn Ser Val Asp  
 115 120 125  
 Glu Leu Asn Asn Asn Val Glu Ala Val Ser Gln Thr Ser Ser Ser Ser  
 130 135 140  
 Phe Gln Tyr Met Tyr Leu Leu Lys Asp Leu Trp Gln Lys Arg Gln Lys  
 145 150 155 160  
 Gln Val Lys Asp Asn Glu Asn Val Val Asn Glu Tyr Ser Ser Glu Leu  
 165 170 175  
 Glu Lys His Gln Leu Tyr Ile Asp Glu Thr Val Asn Ser Asn Ile Pro  
 180 185 190  
 Thr Asn Leu Arg Val Leu Arg Ser Ile Leu Glu Asn Leu Arg Ser Lys  
 195 200 205  
 Ile Gln Lys Leu Glu Ser Asp Val Ser Ala Gln Met Glu Tyr Cys Arg  
 210 215 220  
 Thr Pro Cys Thr Val Ser Cys Asn Ile Pro Val Val Ser Gly Lys Glu  
 225 230 235 240  
 Cys Glu Glu Ile Ile Arg Lys Gly Gly Glu Thr Ser Glu Met Tyr Leu  
 245 250 255  
 Ile Gln Pro Asp Ser Ser Val Lys Pro Tyr Arg Val Tyr Cys Asp Met  
 260 265 270  
 Asn Thr Glu Asn Gly Gly Trp Thr Val Ile Gln Asn Arg Gln Asp Gly  
 275 280 285  
 Ser Val Asp Phe Gly Arg Lys Trp Asp Pro Tyr Lys Gln Gly Phe Gly  
 290 295 300  
 Asn Val Ala Thr Asn Thr Asp Gly Lys Asn Tyr Cys Gly Leu Pro Gly  
 305 310 315 320  
 Glu Tyr Trp Leu Gly Asn Asp Lys Ile Ser Gln Leu Thr Arg Met Gly  
 325 330 335

-continued

---

Pro Thr Glu Leu Leu Ile Glu Met Glu Asp Trp Lys Gly Asp Lys Val  
 340 345 350

Lys Ala His Tyr Gly Gly Phe Thr Val Gln Asn Glu Ala Asn Lys Tyr  
 355 360 365

Gln Ile Ser Val Asn Lys Tyr Arg Gly Thr Ala Gly Asn Ala Leu Met  
 370 375 380

Asp Gly Ala Ser Gln Leu Met Gly Glu Asn Arg Thr Met Thr Ile His  
 385 390 395 400

Asn Gly Met Phe Phe Ser Thr Tyr Asp Arg Asp Asn Asp Gly Trp Leu  
 405 410 415

Thr Ser Asp Pro Arg Lys Gln Cys Ser Lys Glu Asp Gly Gly Gly Trp  
 420 425 430

Trp Tyr Asn Arg Cys His Ala Ala Asn Pro Asn Gly Arg Tyr Tyr Trp  
 435 440 445

Gly Gly Gln Tyr Thr Trp Asp Met Ala Lys His Gly Thr Asp Asp Gly  
 450 455 460

Val Val Trp Met Asn Trp Lys Gly Ser Trp Tyr Ser Met Arg Lys Met  
 465 470 475 480

Ser Met Lys Ile Arg Pro Phe Phe Pro Gln Gln  
 485 490

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10564 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vii) IMMEDIATE SOURCE:

- (B) CLONE: human fibrinogen gamma chain

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: join(1799..1876, 1973..2017, 2207..2390, 2510  
 ..2603, 4211..4341, 4645..4778, 5758..5942, 7426  
 ..7703, 9342..9571)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CTACACACTT CTTGAAGGCA AAGGCAATGC TGAAGTCACC TTTTCATGTTT AAATCATATT 60

AAAAAGTTAG CAAGATGTAA TTATCAGTGT ACTATGTAAA TCTTTGTGAA TGATCAATAA 120

TTACATATTT TCATTATATA TATTTTAGTA GATAATATTT ATATACATTC AACATTCTAA 180

ATATAGAAAG TTTACAGAGA AAAATAAAGC CTTTTTTTCC AATCCTGTCC TCCACCTCTG 240

CATCCCATTG TTCTTCACAG AGGCAACTGA TTCAAGTCAT TACATAGTTA TTGAGTGTTA 300

ACTACAACCTA TGTTAAGTAC AGCTATATAT GTTAGATGCC GTAGCCACAG AAATCAGTTT 360

ACAATCTAAT GCAGTGGATA CAGCATGTAT ACATATAATA TAAGGTTGCT ACAAATGCTA 420

TCTGAGGTAG AGCTGTTTGA AAGAATACTA ATACTTAAAT GTTTAATTCA ACTGACTTGA 480

TTGACAACCTG ATTAGCTGAG TGGAAAAGAT GGATGAGAAA GATTGTGAGA CTTAATTGGC 540

TGGTGGTATG GTGATATGAT TGACAATAAC TGCTAAGTCA GAGAGGGATA TATTAAGGAG 600

GAGAAGAAAA GCAACAAATC TGGTTTTGAT GTGTTCACTT TGTATAAATT ATTGATTATT 660

TACTGAATAT GAATATTTAT CTTTGTTTTT GAGTCAATAA ATATACCTTT GTAAAGACAG 720

AATTAAAGTA TTAGTATTTT TTTCAAACCTG GAGGCATTTT TCCCCTAAC ATATTTTCATC 780

AAAACCTTATA ATAAGCTTGG TTCCAGAGGA AGAAATGAGG GATAACCCAAA AATAGAGACA 840

-continued

TTAATAATAG TGTAACGCC AGTGATAAT CTCATAGGC AGTGATGACA GACATGTTTT	900
CCCAAACACA AGGATGCTGT AAGGGCCAAA CAGAAATGAT GGCCCTCCC CAGCACCTCA	960
TTTTGCCCTC TCCTTCAGCT ATGCCTCTAC TCTCCTTTAG ATACAAGGGA GGTGGATTTT	1020
TCTCTTCTCT GAGATAGCTT GATGGAACCA CAGGAACAAT GAAGTGGGCT CCTGGCTCTT	1080
TTCTCTGTGG CAGATGGGGT GCCATGCCCA CCTTCAGACA AAGGGAAGAT TGAGCTCAAA	1140
AGCTCCCTGA GAAGTGAAG CCTATGAACA TGGTTGACAC AGAGGGACAG GAATGTATTT	1200
CCAGGGTCAT TCATTCTGG GAATAGTGAA CTGGGACATG GGGGAAGTCA GTCTCCTCCT	1260
GCCACAGCCA CAGATTAAAA ATAATAATGT TAACTGATCC CTAGGCTAAA ATAATAGTGT	1320
TAACTGATCC CTAAGCTAAG AAGTTCTTT TGGTAATTCA GGTGATGGCA GCAGGACCCA	1380
TCTTAAGGAT AGACTAGGTT TGCTTAGTTC GAGGTCATAT CTGTTTGCTC TCAGCCATGT	1440
ACTGGAAGAA GTTGCACTAC ACAGCCTCCA GGACTGCCCT CCTCCTCACA GCAATGGATA	1500
ATGCTTCACT AGCCTTTGCA GATAATTTTG GATCAGAGAA AAAACCTTGA GCTGGGCCAA	1560
AAAGGAGGAG CTTCAACCTG TGTGCAAAAT CTGGGAACCT GACAGTATAG GTTGGGGGCC	1620
AGGATGAGGA AAAAGGAACG GGAAAGACCT GCCCACCTT CTGGTAAGGA GGCCCCGTGA	1680
TCAGCTCCAG CCATTTGCAG TCCTGGCTAT CCCAGGAGCT TACATAAAGG GACAATTGGA	1740
GCCTGAGAGG TGACAGTGT GACACTACAA GGCTCGGAGC TCCGGGCACT CAGACATC	1798
ATG AGT TGG TCC TTG CAC CCC CGG AAT TTA ATT CTC TAC TTC TAT GCT	1846
Met Ser Trp Ser Leu His Pro Arg Asn Leu Ile Leu Tyr Phe Tyr Ala	
1 5 10 15	
CTT TTA TTT CTC TCT TCA ACA TGT GTA GCA GTAAGTGTG TCTTCACAAA	1896
Leu Leu Phe Leu Ser Ser Thr Cys Val Ala	
20 25	
ACGTTGTTTA AAATGGAAAG CTGGAAAATA AAACAGATAA TAACTAGTG AAATTTTCGT	1956
ATTTTTTCTC TTTTAG TAT GTT GCT ACC AGA GAC AAC TGC TGC ATC TTA	2005
Tyr Val Ala Thr Arg Asp Asn Cys Cys Ile Leu	
30 35	
GAT GAA AGA TTC GTAAGTAGTT TTTATGTTTC TCCCTTGTG TGTGAAGTGG	2057
Asp Glu Arg Phe	
40	
AGAGGGGCAG AGGAATAGAA ATAATCCCT CATAAATATC ATCTGGCACT TGTAACTTTT	2117
TAAAAACATA GTCTAGGTTT TACCTATTTT TCTTAATAGA TTTTAAGAGT AGCATCTGTC	2177
TACATTTTTA ATCACTGTTA TATTTTCAG GGT AGT TAT TGT CCA ACT ACC TGT	2230
Gly Ser Tyr Cys Pro Thr Thr Cys	
45	
GGC ATT GCA GAT TTC CTG TCT ACT TAT CAA ACC AAA GTA GAC AAG GAT	2278
Gly Ile Ala Asp Phe Leu Ser Thr Tyr Gln Thr Lys Val Asp Lys Asp	
50 55 60 65	
CTA CAG TCT TTG GAA GAC ATC TTA CAT CAA GTT GAA AAC AAA ACA TCA	2326
Leu Gln Ser Leu Glu Asp Ile Leu His Gln Val Glu Asn Lys Thr Ser	
70 75 80	
GAA GTC AAA CAG CTG ATA AAA GCA ATC CAA CTC ACT TAT AAT CCT GAT	2374
Glu Val Lys Gln Leu Ile Lys Ala Ile Gln Leu Thr Tyr Asn Pro Asp	
85 90 95	
GAA TCA TCA AAA CCA A GTGAGAAAAT AAAGACTACT GACCAAAAAA	2420
Glu Ser Ser Lys Pro	
100	
TAATAATAAT AATCTGTGAA GTTCTTTTGC TGTTGTTTTA GTTGTCTAT TTGCTTAAGG	2480
ATTTTTATGT CTCTGATCCT ATATTACAG AT ATG ATA GAC GCT GCT ACT TTG	2532
Asn Met Ile Asp Ala Ala Thr Leu	
105 110	

-continued

AAG TCC AGG ATA ATG TTA GAA GAA ATT ATG AAA TAT GAA GCA TCG ATT	2580
Lys Ser Arg Ile Met Leu Glu Glu Ile Met Lys Tyr Glu Ala Ser Ile	
115 120 125	
TTA ACA CAT GAC TCA AGT ATT CG GTAAGGATTT TTGTTTTAAT TTGCTCTGCA	2633
Leu Thr His Asp Ser Ser Ile Arg	
130	
AGACTGATTT AGTTTTTATT TAATATTCTA TACTTGAGTG AAAGTAATTT TTAATGTGTT	2693
TTCCCCATTT ATAATATCCC AGTGACATTA TGCCTGATTA TGTGAGCAT AGTAGAGATA	2753
GAAGTTTTTA GTGCAATATA AATTACTCTG GGTATAAATT GCTTATTAAT AATCACATTG	2813
AAGAAAGATG TTCTAGATGT CTTCAAATGC TAGTTTGACC ATATTTATCA AAAATTTTTT	2873
CCCCATCCCC CATTATCTTT ACAACATAAA ATCAATCTCA TAGGAATTTG GGTGTTGAAA	2933
ATAAAATCCT CTTTATAAAA ATGCTGACAA ATTGGTGGTT AAAAAAATTA GCAAGCAGAG	2993
GCATAGTAAG GATTTTGGCT CCTAAAGTAA ATTATATTGA ATGTGGAGCA GGAAGAAACA	3053
TGTCTTGAGA GACTAAGTGT GGCAATATT GCAAAGCTCA TATTGATCAT TGCAGAATGA	3113
ACCTGCATAG TCTCTCCCTC TCATTGGAA GTGAATGTCT CTGTTAAAGC TTCTCAGGGA	3173
CTCATAAACT TTCTGAACAT AAGGTCTCAG ATACAGTTTT AATATTTTTT CCCAATTTTT	3233
TTTTCTGAAT TTTTCTCAA GCAGCTTGAG AAATTGAGAT AAATAGTAGC TAGGGAGAAG	3293
TGGCCAGGA AAGATTTCTC CTCTTTTTCG TATCAGAGGG CCCTTGTTAT TATTGTTATT	3353
ATTATTACTT GCATTATTAT TGTCCATCAT TGAAGTTGAA GGAGGTTATT GTACAGAAAT	3413
TGCCTAAGAC AAGGTAGAGG GAAAACGTGG ACAAATAGTT TGTCTACCCT TTTTACTTTC	3473
AAAGAAAGAA CGGTTTATGC ATTGTAGACA GTTTTCTATC ATTTTTGGAT ATTTGCAAGC	3533
CACCCTGTAA GTAACACAA AAGGAGGGTT TTTACTTCCC CCAGTCCATT CCCAAAGCTA	3593
TGTAACCAGA AGCATTAAAG AAGAAAGGGG AAGTATCTGT TGTTTTATTT TACATACAAT	3653
AACGTTCCAG ATCATGTCCC TGTGTAAGTT ATATTTTAGA TTGAAGCTTA TATGTATAGC	3713
CTCAGTAGAT CCACAAGTGA AAGGTATACT CCTTCAGCAC ATGTGAATTA CTGAAGTGA	3773
CTTTTCTGCT TTCTAAAGCA TCAGGGGGTG TTCCTATTAA CCAGTCTCGC CACTCTTGCA	3833
GGTGTCTATC TGCTGTCCCT TATGCATAAA GTAAAAAGCA AAATGTCAAT GACATTTGCT	3893
TATTGACAAG GACTTTGTTA TTTGTGTTGG GAGTTGAGAC AATATGCCCC ATTCTAAGTA	3953
AAAAGATTCA GGTCCACATT GTATTCTGT TTTAATTGAT TTTTGTATT GTTTTCTTT	4013
TTCAAAAAGT TTATAATTTT AATTCATGTT AATTTAGTAA TATAATTTTA CATTTTCTC	4073
AAGAATGGAA TAATTATCA GAAAGCACTT CTTAAGAAAA TACTTAGCAG TTTCCAAGA	4133
AAATATAAAA TTAATCTTCT GAAAGGAATA CTTATTTTGT TCTTCTTATT TTTGTTATCT	4193
TATGTTTCTG TTTGTAG A TAT TTG CAG GAA ATA TAT AAT TCA AAT AAT CAA	4244
Tyr Leu Gln Glu Ile Tyr Asn Ser Asn Asn Gln	
135 140 145	
AAG ATT GTT AAC CTG AAA GAG AAG GTA GCC CAG CTT GAA GCA CAG TGC	4292
Lys Ile Val Asn Leu Lys Glu Lys Val Ala Gln Leu Glu Ala Gln Cys	
150 155 160	
CAG GAA CCT TGC AAA GAC ACG GTG CAA ATC CAT GAT ATC ACT GGG AAA G	4341
Gln Glu Pro Cys Lys Asp Thr Val Gln Ile His Asp Ile Thr Gly Lys	
165 170 175	
GTAAGTATG AAGGTTATAT TGGGATTAGG TTCATCAAAG TAAGTAATGT AAAGGAGAAA	4401
GTATGTACTG GAAAGTATAG GAATAGTTTA GAAAGTGGCT ACCCATTAAG TCTAAGAATT	4461
TCAGTTGTCT AGACCTTTCT TGAATAGCTA AAAAAACAG TTTAAAAGGA ATGCTGATGT	4521

-continued

AAAAAGTAAG AAAATATTTC TTGGAAAATG AATAGTTTAC TACATGTTAA AAGCTATTTT	4581
TCAAGGCTGG CACAGTCTTA CCTGCATTTT AAACCACAGT AAAAGTCGAT TCTCCTTCTC	4641
TAG AT TGT CAA GAC ATT GCC AAT AAG GGA GCT AAA CAG AGC GGG CTT Asp Cys Gln Asp Ile Ala Asn Lys Gly Ala Lys Gln Ser Gly Leu 180 185 190	4688
TAC TTT ATT AAA CCT CTG AAA GCT AAC CAG CAA TTC TTA GTC TAC TGT Tyr Phe Ile Lys Pro Leu Lys Ala Asn Gln Gln Phe Leu Val Tyr Cys 195 200 205	4736
GAA ATC GAT GGG TCT GGA AAT GGA TGG ACT GTG TTT CAG AAG Glu Ile Asp Gly Ser Gly Asn Gly Trp Thr Val Phe Gln Lys 210 215 220	4778
GTAAATTTTT CCCACCATG TGTATTTAAT AAATTCCTAC ATTGTTTCTG CCATATGGCA	4838
GATACTTTTC TAAGCACCTT GTGAACCGTA GCTCATTTAA TCCTTGCAAT AGCCCTAAGA	4898
GGAAGGTACT TCTGTTACTC CTATTTACAG AAAAGGAAAC TGAGGCACAC AAGGTTAAAT	4958
AACTTGCCCA AGACCACATA ACTAATAAGC AACAGAGTCA GCATTTGAAC CTAGGCAGTA	5018
TAGTTTCAGA GTTTGTGACT TGA CTCTATA TTG TACTGGC ACTGACTTTG TAGATTCATG	5078
GTGGCACATA ATCATAGTAC CACAGTGACA AATAAAAAGA AGGAAACTCT TTTGTCAGGT	5138
AGGTCAAGAC CTGAGGTTTC CCATCACAAG ATGAGGAAGC CCAACACCAC CCCCACCAC	5198
CCCACCACCA TCACCACCCT TTCACACACC AGAGGATACA CTTGGGCTGC TCCAAGACAA	5258
GGAACCTGTG TTGCATCTGC CACTTGCTGA TACCCACTAG GAATCTTGGC TCCTTTACTT	5318
TCTGTTTACC TCCCACCCT GTTATAACTG TTTCTACAGG GGGCGCTCAG AGGGAATGAA	5378
TGGTGAAGC ATTAGTTGCC AGACACCGAT TGAGCAATGG GTTCCATCAT AAGTGAAGA	5438
ATCAGTAATA TCCAGCTAGA GTTCTGAAGT CGTCTAGGTG TCTTTTAAAT ATTACCACTC	5498
ATTTAGAATT TATGATGTGC CAGAAACCCT CTTAAGTATT TCTCTTATAT TCTCTCTCAT	5558
GATCCTTGCA GCAACCCTAA GAAGTAACCA TCATTTTTCC TATTTGATAC ATGAGGAAAC	5618
TGAGGTAGCT TGGCCAAGAT CACTTAGTTG GGAGTTGATA GAACCAAGTGC TCTGTATTTT	5678
TGACAAAATG TTGACAGCAT TCTCTTTACA TGCATTGATA GTCTATTTTC TCCTTTTGCT	5738
CTTGCAAATG TGTAATTAG AGA CTT GAT GGC AGT GTA GAT TTC AAG AAA AAC Arg Leu Asp Gly Ser Val Asp Phe Lys Lys Asn 225 230	5790
TGG ATT CAA TAT AAA GAA GGA TTT GGA CAT CTG TCT CCT ACT GGC ACA Trp Ile Gln Tyr Lys Glu Gly Phe Gly His Leu Ser Pro Thr Gly Thr 235 240 245	5838
ACA GAA TTT TGG CTG GGA AAT GAG AAG ATT CAT TTG ATA AGC ACA CAG Thr Glu Phe Trp Leu Gly Asn Glu Lys Ile His Leu Ile Ser Thr Gln 250 255 260 265	5886
TCT GCC ATC CCA TAT GCA TTA AGA GTG GAA CTG GAA GAC TGG AAT GGC Ser Ala Ile Pro Tyr Ala Leu Arg Val Glu Leu Glu Asp Trp Asn Gly 270 275 280	5934
AGA ACC AG GTACTGTTTT GAAATGACTT CCAACTTTTT ATTGTAAGA Arg Thr Ser	5982
TTGCCTGGAA TGTGCACTTT CCAACTATCA ATAGACAATG GCAAATGCAG CCTGACAAAT	6042
GCAAACAGCA CATCCAGCCA CCATTTTCTC CAGGAGTCTG TTTGGTTCTT GGGCAATCCA	6102
AAAAGGTAAA TTCTATTCAG GATGAATCTA AGTGTATTGG TACAATCTAA TTACCCTGGA	6162
ACCATTGAGA GTAATAGCTA ATTACTGAAC TTTTAATCAG TCCCAGGAAT TGAGCATAAA	6222
ATTATAATTT TATCTAGTCT AAATACTAT TTCATGAAGC AGGTATTATT ATTAATCCCA	6282
TTTTATAGAT TAACTTGCTC AAAGTCACAT TGCTGATAAG TGGTAGAGGT AGAATTCAGA	6342
CTCAAGTAGT TTAACCTTAG AGCCTGTCCT CTTAACAAC ATCCTGGTTG AAAAGCAAAT	6402

-continued

ACAGCCTCTT CAGACTTCTC AGTGCCTTGA TGGCCATTTA TTCTGTCAAA TCATGAGCTA	6462
CCCTAAAAGT AAACCAGCTA GCTCTTTTGA TGATCTAGAG GCTTCTTTTT GCTTGAGATA	6522
TTTGAAGGTT TTAAGCATTG TTACCTAATT AAAATGCAGA AAAATATCCA ACCCTCTTGT	6582
TATGTTAAG GAATAGTGAA ATATATTGTC TCAAAACACA TGGACTTTTT TTTATTGCTT	6642
GGTTGGTTTT TAATCCAGAA AGTGTATAG TCAGTAGACC TTCTTCTAGG AAAGGACCTT	6702
CCATTTCCCA GCCACTGGAG ATTAGAAAAT AAGCTAAATA TTTTCTGGAA ATTTCTGTTC	6762
ATTCATTAAG GCCCATCCTT TCCCCACTC TATAGAAGTG TTGTCCACTT GCACAATTTT	6822
TTCCAGGAAA GAATCTCTCT AACTCCTTCA GCTCACATGC TTTGGACCAC ACAGGGAAGA	6882
CTTTGATTGT GTAATGCCCT CAGAAGCTCT CCTTCTTGCC ACTACCACAC TGATTTGAGG	6942
AAGAAAATCC CTTTAGCACC TAACCCTTCA GGTGCTATGA GTGGCTAATG GAACTGTACC	7002
TCCTTCAAGT TTTGTGCAAT AATTAAGGGT CACTCACTGT CAGATACTTT CTGTGATCTA	7062
TGATAATGTG TGTGCAACAC ATAACATTTT AATAAAAGTA GAAAATATGA AATTAGAGTC	7122
ATCTACACAT CTGGATTGA TCTTAGAATG AAACAAGCAA AAAAGCATCC AAGTGAGTGC	7182
AATTATTAGT TTTCAGAGAT GCTTCAAAGG CTTCTAGGCC CATCCCGGA AGTGTTAATG	7242
AGCTGTGGAC TGGTTCACAT ATCTATTGCC TCTTGCCAGA TTTGCAAAA ACTTCACTCA	7302
ATGAGCAAAAT TTCAGCCTTA AGAAACAAAG TCAAAAATTC CAAGGAAGCA TCCTACGAAA	7362
GAGGGAACCT CTGAGATCCC TGAGGAGGGT CAGCATGTGA TGGTTGTATT TCCTTCTTCT	7422
CAG T ACT GCA GAC TAT GCC ATG TTC AAG GTG GGA CCT GAA GCT GAC	7468
Thr Ala Asp Tyr Ala Met Phe Lys Val Gly Pro Glu Ala Asp	
285 290 295	
AAG TAC CGC CTA ACA TAT GCC TAC TTC GCT GGT GGG GAT GCT GGA GAT	7516
Lys Tyr Arg Leu Thr Tyr Ala Tyr Phe Ala Gly Gly Asp Ala Gly Asp	
300 305 310	
GCC TTT GAT GGC TTT GAT TTT GGC GAT GAT CCT AGT GAC AAG TTT TTC	7564
Ala Phe Asp Gly Phe Asp Phe Gly Asp Asp Pro Ser Asp Lys Phe Phe	
315 320 325 330	
ACA TCC CAT AAT GGC ATG CAG TTC AGT ACC TGG GAC AAT GAC AAT GAT	7612
Thr Ser His Asn Gly Met Gln Phe Ser Thr Trp Asp Asn Asp Asn Asp	
335 340 345	
AAG TTT GAA GGC AAC TGT GCT GAA CAG GAT GGA TCT GGT TGG TGG ATG	7660
Lys Phe Glu Gly Asn Cys Ala Glu Gln Asp Gly Ser Gly Trp Trp Met	
350 355 360	
AAC AAG TGT CAC GCT GGC CAT CTC AAT GGA GTT TAT TAC CAA G	7703
Asn Lys Cys His Ala Gly His Leu Asn Gly Val Tyr Tyr Gln	
365 370 375	
GTATGTTTTT CTTTCTTAGA TTCCAAGTTA ATGTATAGTG TATACTATTT TCATAAAAA	7763
TAATAAATAG ATATGAAGAA ATGAAGAATA ATTTATAAAG ATAGTAGGGA TTTTATCATG	7823
TTCTTTATTT CAACTAAGTT CTTTGAAACT GGAAGTGGAT AATACCAAGT TCATGCCTAA	7883
AATTAGCCCT TCTAAAGAAA TCCACCTGCT GCAAAATATC CAGTAGTTTG GCATTATATG	7943
TGAAACTATC ACCATCATAG CTGGCACTGT GGGTTGTGGG ATCTCCTTTA GACATACAAC	8003
ATAAATGATC TGATGGATT AACATTACTA CATGGATGCT TGTTGACACA TTAACCTGGC	8063
TTCCCATGAG CTTTGTGTCA GATACACGCA GTGAACAGGT GTTTGGAGGA ACAGAATAAA	8123
GAGAAGGCAA GCACTGGTAA GGGCAGGGGT TTGTGAAAGC TTGAGAGAAG AGACCACTCT	8183
GAGGACAGTA GACACTTATT TTAGGATGGG GGTGGATGA GGAGGCTATA GTTTGCTATA	8243
AGCTTGAAT GGTTTGAAC ACTGGTTTCA CTCACCTACC CAGCAGTTAT GTGTGGGGAA	8303
GCCTTACCGA TGCTAAAGGA TCCATGTTAC AATAATGGCA TTATTTGGAA ATCCCAGTGG	8363

-continued

TATTCCATGA ATAAAACCAC TATGAAGATA ATCCCACTCA ACAGACTCTC CGTTGGAGAA	8423
GGACAGCAAC ACCACCCTGG GAAAGCCAAA CAGTCAGACC AGACCTGTTT AGCATCAGTA	8483
GGACTTCCCT ACCATATCTG CTGGGTAGAT GAGTGAAACC AGTGTTCCTAA ACCACTCCGG	8543
GCTTGTAGCA AACCATAGTC TCCTCATCTA CCAAGATGAG CAACCTTACC TCCTGATGTC	8603
CTAGCCAATC ACCAACTAGG AAACCTTGCA CAGTTTATTT AAAGTAACAG TTTGATTTTC	8663
ACAAATATTTT TAAATTGGAG AACATAACT TATCTTTGCA CTCACAAACC ACATAATGAG	8723
AAGAACTCT AAGGGAAAAT GCTTGATCTG TGTGACCCGG GGCGCCATGC CAGAGCTGTA	8783
GTTTCATGCCA GTGTTGTGCT CTGACAAGCC TTTTACAGAA TTACATGAGA TCTGCTTCCC	8843
TAGGACAAGG AGAAGCCAAA TCAACAGAGG CTGCACTTTA AAATGGAGAC ATAAAATAAC	8903
ATGCCAGAAC CATTTCCTAA AGCTCCTCAA TCAACCAACA AAATTGTGCT TTCAAATAAC	8963
CTGAGTTGAC CTCATCAGGA ATTTTGTGGC TCCTTCTCTT CTAACCTGCC TGAAGAAAGA	9023
TGGTCCACAG CAGCTGAGTC CGGGATGGAT AAGCTTAGGG ACAGAGGCCA ATTAGGGAAC	9083
TTTGGGTTTC TAGCCCTACT AGTAGTGAAT AAATTTAAAG TGTGGATGTG ACTATGAGTC	9143
ACAGCACAGA TGTTGTTTAA TAATATGTTT ATTTTATAAA TTGATATTTT AGGAATCTTT	9203
GGAGATATTT TCAGTTAGCA GATAATACTA TAAATTTTAT GTAACCTGGCA ATGCACTTCG	9263
TAATAGACAG CTCTTCATAG ACTTGCAGAG GTAAAAAGAT TCCAGAATAA TGATATGTAC	9323
ATCTACGACT TGTTTTAG GT GGC ACT TAC TCA AAA GCA TCT ACT CCT AAT	9373
Gly Gly Thr Tyr Ser Lys Ala Ser Thr Pro Asn	
380 385	
GGT TAT GAT AAT GGC ATT ATT TGG GCC ACT TGG AAA ACC CGG TGG TAT	9421
Gly Tyr Asp Asn Gly Ile Ile Trp Ala Thr Trp Lys Thr Arg Trp Tyr	
390 395 400	
TCC ATG AAG AAA ACC ACT ATG AAG ATA ATC CCA TTC AAC AGA CTC ACA	9469
Ser Met Lys Lys Thr Thr Met Lys Ile Ile Pro Phe Asn Arg Leu Thr	
405 410 415	
ATT GGA GAA GGA CAG CAA CAC CAC CTG GGG GGA GCC AAA CAG GTC AGA	9517
Ile Gly Glu Gly Gln Gln His His Leu Gly Gly Ala Lys Gln Val Arg	
420 425 430 435	
CCA GAG CAC CCT GCG GAA ACA GAA TAT GAC TCA CTT TAC CCT GAG GAT	9565
Pro Glu His Pro Ala Glu Thr Glu Tyr Asp Ser Leu Tyr Pro Glu Asp	
440 445 450	
GAT TTG TAGAAAATTA ACTGCTAACT TCTATTGACC CACAAAGTTT CAGAAAATCT	9621
Asp Leu	
CTGAAAGTTT CTTCCTTTTT TCTCTACTA TATTTATTGA TTTCAAGTCT TCTATTAAGG	9681
ACATTTAGCC TTCAATGGAA ATTAATACT ATTTAGGACT GTATTTCCAA ATTACTGATA	9741
TCAGAGTTAT TTAATAATG TTTATTTGAG GAGATAACAT TTCAACTTTG TTCCTAAATA	9801
TATAATAATA AAATGATTGA CTTTATTTGC ATTTTATGA CCACTTGTC TTTATTTTGT	9861
CTTCGTAAT TATTTTCATT ATATCAAATA TTTTAGTATG TACTTAATAA AATAGGAGAA	9921
CATTTTAGAG TTTCAAATTC CCAGGTATTT TCCTTGTTTA TTACCCCTAA ATCATTCCCTA	9981
TTTAATTCCT CTTTTTAAAT GGAGAAAATT ATGTCTTTTT AATATGGTTT TTGTTTGT	10041
ATATATTCAC AGGCTGGAGA CGTTTAAAG ACCGTTTCAA AAGAGATTTA CTTTTTTAAA	10101
GGACTTTATC TGAACAGAGA GATATAATAT TTTTCCTATT GGACAATGGA CTTGCAAAGC	10161
TTCACTTCAT TTTAAGAGCA AAAGACCCCA TGTTGAAAAC TCCATAACAG TTTTATGCTG	10221
ATGATAATTT ATCTACATGC ATTTCAATAA ACCTTTTGT TTCTAAGACT AGATACATGG	10281
TACCTTTATT GACCATTAAA AAACCACCAC TTTTGGCCAA TTTACCAATT ACAATTGGGC	10341
AACCATCAGT AGTAATTGAG TCCTCATTTT ATGCTAAATG TTATGCCTAA CTCTTTGGGA	10401

-continued

---

GTTACAAAGG AAATAGCAAT TATGGCTTTT GCCCTCTAGG AGATACAGGA CAAATACAGG 10461  
 AAAATACAGC AACCCAAACT GACAATACTC TATACAAGAA CATAATCACT AAGCAGGAGT 10521  
 CACAGCCACA CAACCAAGAT GCATAGTATC CAAAGTGCAG CTG 10564

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 453 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Ser Trp Ser Leu His Pro Arg Asn Leu Ile Leu Tyr Phe Tyr Ala  
 1 5 10 15  
 Leu Leu Phe Leu Ser Ser Thr Cys Val Ala Tyr Val Ala Thr Arg Asp  
 20 25 30  
 Asn Cys Cys Ile Leu Asp Glu Arg Phe Gly Ser Tyr Cys Pro Thr Thr  
 35 40 45  
 Cys Gly Ile Ala Asp Phe Leu Ser Thr Tyr Gln Thr Lys Val Asp Lys  
 50 55 60  
 Asp Leu Gln Ser Leu Glu Asp Ile Leu His Gln Val Glu Asn Lys Thr  
 65 70 75 80  
 Ser Glu Val Lys Gln Leu Ile Lys Ala Ile Gln Leu Thr Tyr Asn Pro  
 85 90 95  
 Asp Glu Ser Ser Lys Pro Asn Met Ile Asp Ala Ala Thr Leu Lys Ser  
 100 105 110  
 Arg Ile Met Leu Glu Glu Ile Met Lys Tyr Glu Ala Ser Ile Leu Thr  
 115 120 125  
 His Asp Ser Ser Ile Arg Tyr Leu Gln Glu Ile Tyr Asn Ser Asn Asn  
 130 135 140  
 Gln Lys Ile Val Asn Leu Lys Glu Lys Val Ala Gln Leu Glu Ala Gln  
 145 150 155 160  
 Cys Gln Glu Pro Cys Lys Asp Thr Val Gln Ile His Asp Ile Thr Gly  
 165 170 175  
 Lys Asp Cys Gln Asp Ile Ala Asn Lys Gly Ala Lys Gln Ser Gly Leu  
 180 185 190  
 Tyr Phe Ile Lys Pro Leu Lys Ala Asn Gln Gln Phe Leu Val Tyr Cys  
 195 200 205  
 Glu Ile Asp Gly Ser Gly Asn Gly Trp Thr Val Phe Gln Lys Arg Leu  
 210 215 220  
 Asp Gly Ser Val Asp Phe Lys Lys Asn Trp Ile Gln Tyr Lys Glu Gly  
 225 230 235 240  
 Phe Gly His Leu Ser Pro Thr Gly Thr Thr Glu Phe Trp Leu Gly Asn  
 245 250 255  
 Glu Lys Ile His Leu Ile Ser Thr Gln Ser Ala Ile Pro Tyr Ala Leu  
 260 265 270  
 Arg Val Glu Leu Glu Asp Trp Asn Gly Arg Thr Ser Thr Ala Asp Tyr  
 275 280 285  
 Ala Met Phe Lys Val Gly Pro Glu Ala Asp Lys Tyr Arg Leu Thr Tyr  
 290 295 300  
 Ala Tyr Phe Ala Gly Gly Asp Ala Gly Asp Ala Phe Asp Gly Phe Asp  
 305 310 315 320  
 Phe Gly Asp Asp Pro Ser Asp Lys Phe Phe Thr Ser His Asn Gly Met  
 325 330 335

-continued

---

Gln Phe Ser Thr Trp Asp Asn Asp Asn Asp Lys Phe Glu Gly Asn Cys  
 340 345 350

Ala Glu Gln Asp Gly Ser Gly Trp Trp Met Asn Lys Cys His Ala Gly  
 355 360 365

His Leu Asn Gly Val Tyr Tyr Gln Gly Gly Thr Tyr Ser Lys Ala Ser  
 370 375 380

Thr Pro Asn Gly Tyr Asp Asn Gly Ile Ile Trp Ala Thr Trp Lys Thr  
 385 390 395 400

Arg Trp Tyr Ser Met Lys Lys Thr Thr Met Lys Ile Ile Pro Phe Asn  
 405 410 415

Arg Leu Thr Ile Gly Glu Gly Gln Gln His His Leu Gly Gly Ala Lys  
 420 425 430

Gln Val Arg Pro Glu His Pro Ala Glu Thr Glu Tyr Asp Ser Leu Tyr  
 435 440 445

Pro Glu Asp Asp Leu  
 450

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10807 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (B) CLONE: ovine beta-lactoglobulin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

ACGCGTGTTCG ACCTGCAGGT CAACGGATCT CTGTGTCTGT TTTCATGTTA GTACCACACT 60

GTTTTGGTGG CTGTAGCTTT CAGCTACAGT CTGAAGTCAT AAAGCCTGGT ACCTCCAGCT 120

CTGTTCTCTC TCAAGATTGT GTTCTGCTGT TTGGGTCTTT AGTGTCTCCA CACAATTTTT 180

AGAATTGTTT GTTCTAGTTC TGTGAAAAAT GATGCTGGTA TTTTGATAAG GATTGCATTG 240

AATCTGTAAA GCTACAGATA TAGTCATTGG GTAGTACAGT CACTTTAACA ATATTAATC 300

TTACATCTG TGAGCATGAT ATATTTTCCC CCTCTATATC ATCTTCAATT CCTCCTATCA 360

GTTTCTTTCA TTGCAGTTTT CTGAGTACAG GTCTTACACC TCCTTGGTTA GAGTCATTCC 420

TCAGTATTTT ATTCCTTTGA TACAATTGTG AATGAGGTAA TTTTCTTAGT TTCTCTTTCT 480

GATAGCTCAT TGTTAGTGTA TATATAGAAA AGCAACAGAT TTCTATGTAT TAATTTTGTA 540

TCCTGCAACA GATTTCTATG TATTAATTTT GTATCCTGCT ACTTTACGGA ATTCATTAT 600

TAGCTTTTTG GTGACATCTT GAGGATTTTC TGAAGAAAAT GGCATGGTAT GGTAGGACAA 660

GGTGTATATG CATCTGCAA CAGTGGCAGT TTTCTTCTT CCCTTCCAAC CTGGATTTCT 720

TTGATTTCTT TCTGTCTGAG TACGACTAGG ATTCCAATA CTATACCGAA TAAAAGTGGC 780

AAGAGTGGAC ATCCTTGTCT TATTTTTCTG ACCTTAGAGG AAATGCTTTC AGTTTTTTCAC 840

CATTAATTAT AATGTTTACT GTGGGCTTGT CATATGTGGC CTTCAATATA TGGAGGTCTA 900

TTCCCTCTAT ACCCACCTTG TTGAGAGTTT TTATCATAAA AGTATGTGA ATTTTGTCAA 960

AAGTTTTTTC TGCATCTATT GAGATGATT TTAATCTTCA ATTCATTAAT GATTTTTATT 1020

CTTCATTTTG TTAATGATTT CCATTCTTCA ATTTGTTAAC GTGGTATATC ACATTGATTG 1080

ATTTGTGGAT ACCTTTGTAT CCCTGGGATA AACCTCACTT GATCATGAGC TTTCAATGTA 1140

TTTTTGAATT CACTTTGCTA ATATCTGTGT GGGTATTTTT GCATCTCTAT TCATCAATGA 1200

TATTGGCCTA AGAAAGGTTT TGTCTGGTTT TAGTATCAGG GTGATGCTGG CCTCATAGAG 1260

-continued

AGAGTTTAGA	AGCATTTCCCT	CCTCTTTGAT	TTTTCGGAAT	AGTTTGAGTA	GGATAGGTAT	1320
TAACTCTTCT	TAAATGTTT	GGGACTTCC	CTGGTGAGCC	GGTGGTTGAG	AATCCGCCTC	1380
AGGGATGTGG	GTTTGTATCC	TGGTCAGGA	ACCATTAATA	AGATCCCACA	TGCTGCAGGC	1440
AACAAGCCCC	CAAGCTGCAA	CCACTGAGCT	GCAACCGCTG	CAGTGCCAC	AGGCCACGAC	1500
CAGAGAAAGC	CCACATACAG	CAGGGAAGAC	CCAGCACAAAC	CGAAAAAGG	AGTTTGGTGG	1560
AATACAGCTG	TGAAGCCGTC	TGGTCCTGGA	CTCCTGCTTG	AGGGAATTTT	TTAAAAATTA	1620
TTGATTCAAT	TTCATTACTG	GTAACGTGTC	TGTTTCATATT	TTCTATTCT	TCCGGGTTC	1680
GTCTTGGGAG	ATTGTACATG	CCTAGGAATG	TGTCCGTTTC	TTCTAGGTTG	TCCATTTTAT	1740
TGGACATGCA	TGGGAGCACA	CAGCACCGAC	CAGCGAGACT	CATGCTGGCT	TCCTGGGGCC	1800
AGGCTGGGGC	CCCAAGCAGC	ATGGCATCCT	AGAGTGTGTG	AAAGCCCACT	GACCCTGCCC	1860
AGCCCCACAA	TTTCATTCTG	AGAAGTGATT	CCTTGCTTCT	GCACTTACAG	GCCCAGGATC	1920
TGACCTGCTT	CTGAGGAGCA	GGGGTTTTGG	CAGGACGGGG	AGATGCTGAG	AGCCGACGGG	1980
GGTCCAGGTC	CCCTCCAGG	CCCCCTGTC	TGGGGCAGCC	CTTGGGAAAG	ATTGCCCCAG	2040
TCTCCCTCCT	ACAGTGGTCA	GTCCCAGCTG	CCCCAGGCCA	GAGCTGCTTT	ATTCCGTCT	2100
CTCTCTCTGG	ATGGTATTCT	CTGGAAGCTG	AAGGTTCCCTG	AAGTTATGAA	TAGCTTTGCC	2160
CTGAAGGGCA	TGGTTTGTGG	TCACGGTTCA	CAGGAACTTG	GGAGACCTTG	CAGCTCAGAC	2220
GTCCCGAGAT	TGGTGGCACC	CAGATTCCT	AAGCTCGCTG	GGGAACAGGG	CGCTTGTTTC	2280
TCCCTGGCTG	ACCTCCCTCC	TCCCTGCATC	ACCCAGTTCT	GAAAGCAGAG	CGGTGCTGGG	2340
GTACAGCCT	CTCGCATCTA	ACGCCGGTGT	CCAAACCACC	CGTGTGGTG	TTCGGGGGGC	2400
TACCTATGGG	GAAGGGCTTC	TCACTGCAGT	GGTGCCCCC	GTCCCTCTG	AGATCAGAAG	2460
TCCAGTCCG	GACGTCAAAC	AGGCCGAGCT	CCCTCCAGAG	GCTCCAGGGA	GGGATCCTTG	2520
CCCCCCCGCT	GCTGCCTCCA	GCTCCTGGTG	CCGCACCCTT	GAGCCTGATC	TTGTAGACGC	2580
CTCAGTCTAG	TCTCTGCCTC	CGTGTTCACA	CGCCTTCTCC	CCATGTCCCC	TCCGTGTCCC	2640
CGTTTTCTCT	CACAAGGACA	CCGGACATTA	GATTAGCCCC	TGTTCCAGCC	TCACCTGAAC	2700
AGCTCACATC	TGTAAAGACC	TAGATTCCAA	ACAAGATTCC	AACCTGAAGT	TCCCGGTGGA	2760
TGTGAGTTCT	GGGGCGACAT	CCTTCAACCC	CATCACAGCT	TGCAGTTCAT	CGCAAAACAT	2820
GGAACCTGGG	GTTTATCGTA	AAACCAGGT	TCTTCATGAA	AACTGAGCT	TCGAGGCTTG	2880
TTGCAAGAAT	TAAAGTGCT	AATACAGATC	AGGGCAAGGA	CTGAAGCTGG	CTAAGCCTCC	2940
TCTTTCATC	ACAGGAAAGG	GGGGCCTGGG	GGCGGCTGGA	GGTCTGCTCC	CGTGAGTGAG	3000
CTCTTTCCTG	CTACAGTCAC	CAACAGTCTC	TCTGGGAAGG	AAACCAGAGG	CCAGAGAGCA	3060
AGCCGGAGCT	AGTTTAGGAG	ACCCCTGAAC	CTCCACCAA	GATGCTGACC	AGCCAGCGGG	3120
CCCCCTGGAA	AGACCCTACA	GTTTCAGGGG	GAAGAGGGGC	TGACCCGCCA	GGTCCCTGCT	3180
ATCAGGAGAC	ATCCCCGCTA	TCAGGAGATT	CCCCACCTT	GCTCCCGTTC	CCCTATCCCA	3240
ATACGCCAC	CCCACCCCTG	TGATGAGCAG	TTTAGTCACT	TAGAATGTCA	ACTGAAGGCT	3300
TTTGCAATCC	CTTTGCCAGA	GGCACAAGGC	ACCCACAGCC	TGCTGGGTAC	CGACGCCCAT	3360
GTGGATTCAG	CCAGGAGGCC	TGTCCTGCAC	CCTCCCTGCT	CGGGCCCCCT	CTGTGCTCAG	3420
CAACACACCC	AGCACCAGCA	TTCCCGCTGC	TCCTGAGGTC	TGCAGGCAGC	TCGCTGTAGC	3480
CTGAGCGGTG	TGGAGGGAAG	TGTCCTGGGA	GATTTAAAT	GTGAGAGGCG	GGAGGTGGGA	3540
GTTTGGGCC	TGTGGGCTG	CCCATCCAC	GTGCCTGCAT	TAGCCCAGT	GCTGCTCAGC	3600
CGTGCCCCG	CCGAGGGGT	CAGGTCACTT	TCCCGTCTG	GGTTATTAT	GACTCTTGTC	3660

-continued

---

ATTGCCATTG	CCATTTTTCG	TACCCTAACT	GGGCAGCAGG	TGCTTGACAG	GCCCTCGATA	3720
CCGACCAGGT	CCTCCCTCGG	AGCTCGACCT	GAACCCCATG	TCACCCCTGC	CCCAGCCTGC	3780
AGAGGGTGGG	TGACTGCAGA	GATCCCTTCA	CCCAAGGCCA	CGGTCACATG	GTTTGGAGGA	3840
GCTGGTGCCC	AAGGCAGAGG	CCACCTCCCA	GGACACACCT	GTCCCCAGTG	CTGGCTCTGA	3900
CCTGTCTTGG	TCTAAGAGGC	TGACCCCGGA	AGTGTTCCTG	GCACTGGCAG	CCAGCCTGGA	3960
CCCAGAGTCC	AGACACCCAC	CTGTGCCCCC	GCTTCTGGGG	TCTACCAGGA	ACCGTCTAGG	4020
CCCAGAGGGG	ACTTCTGTCT	TGGCCTTGGA	TGGAAGAAGG	CCTCCTATTG	TCCTCGTAGA	4080
GGAAGCCACC	CCGGGGCCTG	AGGATGAGCC	AAGTGGGATT	CCGGGAACCG	CGTGGCTGGG	4140
GGCCAGCCC	GGGCTGGCTG	GCCTGCATGC	CTCCTGTATA	AGGCCCAAG	CCTGCTGTCT	4200
CAGCCCTCCA	CTCCCTGCAG	AGCTCAGAAG	CACGACCCCA	GGGATATCCC	TGCAGCCATG	4260
AAGTGCCTCC	TGCTTGCCCT	GGGCCTGGCC	CTCGCCTGTG	GCGTCCAGGC	CATCATCGTC	4320
ACCCAGACCA	TGAAAGGCCT	GGACATCCAG	AAGGTTCCAG	GGTTGGCCGG	GTGGGTGAGT	4380
TGCAGGGCGG	GCAGGGGAGC	TGGGCCTCAG	AGAGCCAAGA	GAGGCTGTGA	CGTTGGGTTT	4440
CCATCAGTCA	GCTAGGGCCA	CCTGACAAAT	CCCCGCTGGG	GCAGCTTCAA	CCAGGCGTTC	4500
ACTGTCTTGC	ATTCTGGAGG	CTGGAAGCCC	AAGATCCAGG	TGTTGGCAGG	GCTGGCTTCT	4560
CCTGCGGCGG	CTCTCTGGGG	AGCAGACGGC	CGTCTTCTCC	AGTCTCTGTC	GCGCCCTGAT	4620
TTCTCTTCC	TGTGAGGCCA	CCAGGCCTGC	TGGAACACG	CCTGCCTGCG	CAGCTTCACA	4680
CGACCTTTGT	CATCTCTTTA	AAGGCCATGT	CTCCAGAGTC	ATGTGTTGAA	GTTCTGGGGG	4740
TTAGTGGGAC	ACAGTTCAGC	CCCTAAAAGA	GTCTCTCTGC	CCCTCAAATT	TTCCCCACCT	4800
CCAGCCATGT	CTCCCCAAGA	TCCAAATGTT	GCTACATGTG	GGGGGGCTCA	TCTGGGTCCC	4860
TCTTTGGGTT	CAGTGTGAGT	CTGGGAGAG	CATCCCCAG	GGTGCAGAGT	TGGGGGGAGT	4920
ATCTCAGGGC	TGCCCAGGCC	GGGGTGGGAC	AGAGAGCCCA	CTGTGGGGCT	GGGGGGCCCT	4980
TCCCACCCCC	AGAGTGCAAC	TCAAGGTCCC	TCTCCAGGTG	GCGGGGACTT	GGCACTCCTT	5040
GGCTATGGCG	GCCAGCGACA	TCTCCCTGCT	GGATGCCCCAG	AGTGCCCCCC	TGAGAGTGTA	5100
CGTGGAGGAG	CTGAAGCCCA	CCCCCGAGGG	CAACCTGGAG	ATCCTGTCTG	AGAAATGGTG	5160
GGCGTCTCTC	CCCAACATGG	AACCCCACT	CCCCAGGGCT	GTGGACCCCC	CGGGGGGTGG	5220
GGTGCAGGAG	GGACCAGGGC	CCCAGGGCTG	GGGAAGAGGG	CTCAGAGTTT	ACTGGTACCC	5280
GGCGCTCCAC	CCAAGCTGTC	CCACCCAGGG	CTTTTTTTTT	TTTTAAACTT	TTATTAATTT	5340
GATGCTTCAG	AACATCATCA	AACAAATGAA	CATAAAACAT	TCATTTTTGT	TTACTTGGA	5400
GGGGAGATAA	AATCCTCTGA	AGTGGAAATG	CATAGCAAAG	ATACATACAA	TGAGGCAGGT	5460
ATTCTGAATT	CCCTGTTAGT	CTGAGGATTA	CAAGTGTATT	TGAGCAACAG	AGAGACATTT	5520
TCATCATTTT	TAGTCTGAAC	ACCTCAGTAT	CTAAAATGAA	CAAGAAGTCC	TGGAAACGAA	5580
GCAGTGTGGG	GATAGCCCCG	TGTGAAGGCT	GCTGGGAGGC	AGCAGACCTG	GGTCTTCGGG	5640
CTCAAGCAGT	TCCCCTTACC	AGCCCTGTCC	ACCTCAGACG	GGGGTCAGGG	TGCAGGAGAG	5700
AGCTGGATGG	GTGTGGGGGC	AGAGATGGGG	ACCTGAACCC	CAGGGCTGCC	TTTTGGGGGT	5760
GCCTGTGGTC	AAGGCTCTCC	CTGACCTTTT	CTCTCTGGCT	TCATCTGACT	TCTCCTGGCC	5820
CATCCACCCG	GTCCCCTGTG	GCCTGAGGTG	ACAGTGAGTG	CGCCGAGGCT	AGTTGGCCAG	5880
CTGGCTCCTA	TGCCCATGCC	ACCCCTCTCC	AGCCCTCCTG	GGCCAGCTTC	TGCCCTGGC	5940
CCTCAGTTCA	TCCTGATGAA	AATGGTCCAT	GCCAATGGCT	CAGAAAGCAG	CTGTCTTTCA	6000
GGGAGAACGG	CGAGTGTGCT	CAGAAGAAGA	TTATTGCAGA	AAAAACCAAG	ATCCCTGCGG	6060

-continued

TGTTCAAGAT	CGATGGTGAG	TCCGGGTCCC	TGGGGGACAC	CCACCACCCC	CGCCCCGGG	6120
GACTGTGGAC	AGGTTACAGG	GGCTGGCGTC	GGGCCCTGGG	ATGCTAAGGG	ACTGGTGGTG	6180
ATGAAGACAC	TGCCTTGACA	CCTGCTTCAC	TTGCCTCCCC	TGCCACCTGC	CCGGGGCCTT	6240
GGGGCGGTGG	CCATGGGCAG	GTCCCGGCTG	GCGGGCTAAC	CCACCAGGGT	GACACCCGAG	6300
CTCTCTTTGC	TGGGGGGCGG	GCGGTGCTCT	GGGCCCTCAG	GCTGAGCTCA	GGAGGTACCT	6360
GTGCCCTCCC	AGGGGTAACC	GAGAGCCGTT	GCCCCTCCA	GGGGCCAGG	TGCCCCACGA	6420
CCCCAGCCCC	CTCCACAGCT	CCTTCATCTC	CTGGAGACAA	ACTCTGTCCG	CCCTCGCTCA	6480
TTCACTTGTT	CGTCCTAAAT	CCGAGATGAT	AAAGCTTCGA	GGGGGGTTG	GGGTTCCATC	6540
AGGGTGTCCC	TTCCGCCGGG	CAGCCTGGGC	CACATCTGCC	CTTGGCCCC	TCAGGACTCA	6600
CTCTGACTGG	AGGCCCTGCA	CTGACTGACG	CCAGGGTGCC	CAGCCAGGG	TCTCTGGCGC	6660
CATCCAGCTG	CACTGGGTTT	GGGTGCTGGT	CCTGCCCCCA	AGCTGCCCGG	ACACCACAGG	6720
CAGCCGGGGC	TGCCCCTGCG	CCTCGGTCAG	GGTGAGCCCC	AGCTGCCCCC	GCTCAGGGCT	6780
TGCCCCGACA	ATGACCCCAT	CCTCAGGACG	CACCCCCCTT	CCCTTGCTGG	GCAGTGTTCA	6840
GCCCCACCCG	AGATCGGGGG	AAGCCCTATT	TCTTGACAA	TCCAGTCCCT	GGGGGAGGGG	6900
GCCTCAGACT	GAGTGGTGAG	TGTTCCCAAG	TCCAGGAGGT	GGTGGAGGGT	CCTGGCGGAT	6960
CCAGAGTTGA	CAGTAGGGGC	TTCCTGGGCC	CCATGCGCCT	GGCAGTGCCA	GCAGGGAAGA	7020
GGAAGCACCA	TTTCAGGGGT	GGGGGATGCC	AGAGGCGCTC	CCCACCCCGT	CTTCGCCGGG	7080
TGGTGACCCC	GGGGGAGCCC	CGCTGGTCTG	GGAGGGTGCT	GGGGGCTGAC	TAGCAACCCC	7140
TCCCCCCCCG	TTGGAActCA	CTTTTCTCCC	GTCTTGACCG	CGTCCAGCCT	TGAATGAGAA	7200
CAAAGTCCTT	GTGCTGGACA	CCGACTACAA	AAAGTACCTG	CTCTTCTGCA	TGGAAAACAG	7260
TGCTGAGCCC	GAGCAAAGCC	TGGCCTGCCA	GTGCCTGGGT	GGGTGCCAAC	CCTGGCTGCC	7320
CAGGGAGACC	AGCTGCTGGG	TCCTTCTGTC	AACAGGGGGT	GGGGGGTGGG	AGCTTGATCC	7380
CCAGGAGGAG	GAGGGGTGGG	GGGTCCCTGA	GTCCCGCCAG	GAGAGAGTGG	TCGCATACCG	7440
GGAGCCAGTC	TGCTGTGGGC	CTGTGGGTGG	CTGGGGACGG	GGGCCAGACA	CACAGGCCGG	7500
GAGACGGGTG	GGCTGAGAAA	CTGTGACTGG	TGTGACCGTC	GCGATGGGGC	CGGTGGTCAC	7560
TGAATCTAAC	AGCCTTTGTT	ACCGGGGAGT	TTCAATTATT	TCCCAAATA	AGAACTCAGG	7620
TACAAAGCCA	TCTTTCAACT	ATCACATCCT	GAAACAAT	GGCAGGTGAC	ATTTTCTGTG	7680
CCGTAGCAGT	CCCACTGGGC	ATTTTCAGGG	CCCCTGTGCC	AGGGGGCGC	GGGCATCGGC	7740
GAGTGAGGGC	TCCTGGCTGT	GTCAGCCGGC	CCAGGGGAG	GAAGGGACCC	GGACAGCCAG	7800
AGGTGGGGGG	CAGGCTTTCC	CCCTGTGACC	TGCAGACCCA	CTGCACTGCC	CTGGGAGGAA	7860
GGGAGGGGAA	CTAGGCCAAG	GGGAAGGGC	AGGTGCTCTG	GAGGGCAAGG	GCAGACCTGC	7920
AGACCACCCCT	GGGGAGCAGG	GACTGACCCC	CGTCCCTGCC	CCATAGTCAG	GACCCCGGAG	7980
GTGGACAACG	AGGCCCTGGA	GAAATTCGAC	AAAGCCCTCA	AGGCCCTGCC	CATGCACATC	8040
CGGCTTGCCCT	TCAACCCGAC	CCAGCTGGAG	GGTGAGCACC	CAGGCCCGC	CCTTCCCAG	8100
GGCAGGAGCC	ACCCGGCCCC	GGGACGACCT	CCTCCCATGG	TGACCCCCAG	CTCCCCAGGC	8160
CTCCAGGAG	GAAGGGGTGG	GGTGACGAC	CCCGTGGGGG	CCCCCTCCCC	ACCCCTGCC	8220
AGGCCTCTCT	TCCCAGGTG	TCCAGTCCCA	TCCTGACCCC	CCCATGACTC	TCCCTCCCC	8280
ACAGGGCAGT	GCCACGTCTA	GGTGAGCCCC	TGCCGGTGCC	TCTGGGGTAA	GCTGCCTGCC	8340
CTGCCCCACG	TCCTGGGCAC	ACACATGGGG	TAGGGGTCT	TGGTGGGGCC	TGGGACCCCA	8400
CATCAGGCC	TGGGTCCCC	CCTGTGAGAA	TGGTGGAAG	CTGGGGTCCC	TCCTGGCGAC	8460

-continued

---

TGCAGAGCTG	GCTGGCCGCG	TGCCACTCTT	GTGGGTGACC	TGTGTCTTGG	CCTCACACAC	8520
TGACCTCCTC	CAGCTCCTTC	CAGCAGAGCT	AAGGCTAAGT	GAGCCAGAAT	GGTACCTAAG	8580
GGGAGGCTAG	CGGTCTTCT	CCCGAGGAGG	GGCTGTCTCTG	GAACCACCAG	CCATGGAGAG	8640
GCTGGCAAGG	GTCTGGCAGG	TGCCCCAGGA	ATCACAGGGG	GGCCCCATGT	CCATTTTCAGG	8700
GCCCCGGAGC	CTTGGACTCC	TCTGGGGACA	GACGACGTCA	CCACCGCCCC	CCCCCATCA	8760
GGGGGACTAG	AAGGGACCAG	GACTGCAGTC	ACCCTTCCTG	GGACCCAGGC	CCCTCCAGGC	8820
CCCTCCTGGG	GCTCCTGCTC	TGGGCAGCTT	CTCCTTCACC	AATAAAGGCA	TAAACCTGTG	8880
CTCTCCCTTC	TGAGTCTTTG	CTGGACGACG	GGCAGGGGGT	GGAGAAGTGG	TGGGGAGGGA	8940
GTCTGGCTCA	GAGGATGACA	GCGGGGCTGG	GATCCAGGGC	GTCTGCATCA	CAGTCTTGTG	9000
ACAACTGGGG	GCCCCACAC	ATCACTGCGG	CTCTTTGAAA	CTTTCAGGAA	CCAGGGAGGG	9060
ACTCGCAGA	GACATCTGCC	AGTTCACTTG	GAGTGTTCAG	TCAACACCCA	AACTCGACAA	9120
AGGACAGAAA	GTGGAATG	GCTGTCTCTT	AGTCTAATAA	ATATTGATAT	GAAACTCAAG	9180
TTGCTCATGG	ATCAATATGC	CTTTATGATC	CAGCCAGCCA	CTACTGTCGT	ATCAACTCAT	9240
GTACCCAAAC	GCACTGATCT	GTCTGGCTAA	TGATGAGAGA	TTCCAGTAG	AGAGCTGGCA	9300
AGAGTCAACA	GTGAGAACTG	TCTGCACACA	CAGCAGAGTC	CACCAGTCAT	CCTAAGGAGA	9360
TCAGTCCTGG	TGTTCAATTGG	AGGACTGATG	TTGAAGCTGA	AACTCCAATG	CTTTGGCCAC	9420
CTGATGTGAA	GAGCTGACTC	ATTTGAAAAG	ACCCTGATGC	TGGGAAAGAT	TGAGGGCAGG	9480
AGGAGAAGGG	GACGACAGAG	GATGAGATGG	TTGGATGGCA	TCACCAACAC	AATGGACATG	9540
GGTTTGGGTG	GACTCCAGGA	GTTGGTGATG	GACAGGGAGG	CCTGGCGTGC	TACGGAAGCG	9600
GTTTATGGGG	TCACAAAAGC	TGAGTGACTG	AACTGAGCTG	AACTGAATGG	AAATGAGGTA	9660
TACAGCAAAG	TGGGGATTTT	TTAGATAATA	AGAATATACA	CATAACATAG	TGTATACTCA	9720
TATTTTTATG	CATACCTGAA	TGCTCAGTCA	CTCAGTCGTA	TCTGACTCTG	TGACCTATGG	9780
ACCGTAGCCT	TCCAGGTTTC	TTCTGTCCAC	AGAATTCTCC	AAGGCAAGAA	TACTGGAGTG	9840
GGTAGCCATT	TCCTCTCCA	GGGATCCTC	CCGACCCAGG	GATTGAACCG	GCATCTCCTG	9900
TATTGGCAGG	TGGATTCTTT	ACCACTGTGC	CACCAGGAA	GCCCGTGTTA	CTCTCTATGT	9960
CCCCTTAAT	TACCAAAGCT	GCTCCAAGAA	AAAGCCCCTG	TGCCCTCTGA	GCTTCCCAGC	10020
CTGCAGAGGG	TGGTGGGGGT	AGACTGTGAC	CTGGGAACAC	CCTCCCGCTT	CAGGACTCCC	10080
GGGCCACGTG	ACCCACAGTC	CTGCAGACAG	CCGGGTAGCT	CTGCTCTTCA	AGGCTCATT	10140
TCTTTAAAA	AAACTGAGGT	CTATTTTGTG	ACTTCGCTGC	CGTAACTTCT	GAACATCCAG	10200
TGCGATGGAC	AGGACCTCCT	CCCCAGGCCT	CAGGGGCTTC	AGGGAGCCAG	CCTTCACCTA	10260
TGAGTCACCA	GACACTCGGG	GGTGGCCCCG	CCTTCAGGGT	GCTCACAGTC	TTCCCATCGT	10320
CCTGATCAAA	GAGCAAGACC	AATGACTTCT	TAGGAGCAAG	CAGACACCCA	CAGGACACTG	10380
AGGTTACCCA	GAGCTGAGCT	GTCTTTTGA	ACCTAAAGAC	ACACAGCTCT	CGAAGGTTTT	10440
CTCTTTAATC	TGGATTTAAG	GCCTACTTGC	CCCTCAAGAG	GGAAGACAGT	CCTGCATGTC	10500
CCCAGGACAG	CCACTCGGTG	GCATCCGAGG	CCACTTAGTA	TTATCTGACC	GCACCCCTGGA	10560
ATTAATCGGT	CCAACTGGA	CAAAAACCTT	GGTGGGAAGT	TTCATCCCAG	AGGCCTCAAC	10620
CATCCTGCTT	TGACCACCCT	GCATCTTTTT	TTCTTTTATG	TGTATGCATG	TATATATATA	10680
TATATATTTT	TTTTTTTTTC	ATTTTTTGGC	TGTGCTGGCT	GTTCTTGCA	GTTCTGGTGC	10740

-continued

---

CAGGCTTCTC TCTAGTTTCT CTCTAGTCTT CTCTATCAC AGAGCAGTCT CTAGACGATC 10800  
 GACGCGT 10807

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 47 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

AATTCGATC GACGCGTCGA CGATATACTC TAGACGATCG ACGCGTA 47

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 24 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: BLGAMP3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TGGATCCCCT GCCGGTGCCT CTGG 24

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 24 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: BLGAMP4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

AACGCGTCAT CCTCTGTGAG CCAG 24

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 10 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: ZC6839

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ACTACGTAGT 10

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 42 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: ZC6632

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

CGACGCGGAT CCTACGTACC TGCAGCCATG TTTCCATGA GG 42

-continued

## (2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6627

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

AGGGCTTCGG CAAGCTTCAG G

21

## (2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6521

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GCCAAAGACT TACTTCCCTC TAGA

24

## (2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6520

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GCATGAACGT CGCGTGGTGG TTGTGCTACC

30

## (2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6519

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

ACCACGCGAC GTTCATGCTC TAAAACCGTT

30

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6518

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

GCTGCGGGAT CCTACGTACT AGGGGACAG GGAAGG

36

-continued

## (2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 45 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6629

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

CGACGCGAAT TCTACGTACC TGCAGCCATG AAAAGGATGG TTTCT 45

## (2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 45 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6630

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

CGACGCGAAT TCTACGTACC TGCAGCCATG AAACATCTAT TATTG 45

## (2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6625

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GTGAGATTTT CAGATCTTGT C 21

## (2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6626

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

AAGAATTACT GTGCCTACC A 21

## (2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6624

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GCTGCGAAT TCTACGTACT ATTGCTGG GAA 33

-continued

## (2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 45 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6514

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

CGACGCGGAT CCTACGTACC TGCAGCCATG AGTTGGTCCT TGCAC 45

## (2) INFORMATION FOR SEQ ID NO: 24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: zc6517

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GTCTCTGGTA GCAACATACT A 21

## (2) INFORMATION FOR SEQ ID NO: 25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: zc6516

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

GGGTTTCTAG CCCTACTAGT AG 22

## (2) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: zc6515

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

GGGTTTCTAG CCCTACTAGT AG 22

## (2) INFORMATION FOR SEQ ID NO: 27:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 47 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

AAGCTACGCG TCGATCGTCT AGAGTATATC GTCGACGCGT CGATCGG 47

We claim:

1. A method for producing biocompetent fibrinogen comprising:

providing a first DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\text{A}\alpha$  chain, *the DNA segment comprising genomic DNA encoding the  $\text{A}\alpha$  chain*, a second DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\text{B}\beta$  chain, *the DNA segment comprising genomic DNA encoding the  $\text{B}\beta$  chain*, and a third DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\gamma$  chain, *the DNA segment comprising genomic DNA encoding the  $\gamma$  chain*, wherein each chain is from the same species, and wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal *and the first, second, third segments are linked in a single vector*;

introducing said DNA segments into a fertilized egg of a non-human mammalian species heterologous to the species of origin of said fibrinogen chains;

inserting said egg into an oviduct or uterus of a female of said mammalian species to obtain offspring carrying said DNA segments;

breeding said offspring to produce female progeny that express said first, second and third DNA segments and produce milk containing biocompetent fibrinogen encoded by said segments;

collecting milk from said female progeny; and  
and recovering the biocompetent fibrinogen from the milk.

2. A method according to claim 1 wherein said species into which said DNA segments are introduced is selected from the group consisting of sheep, pigs, goats, and cattle.

3. A method according to claim 1 wherein each of said first, second and third DNA segments comprises an intron.]

4. A method according to claim 1 wherein the molar ratio of said first, second and third DNA segments is within the range of 0.5-1:0.5-1:0.5-1.]

5. A method according to claim 1 wherein each of said first, second and third DNA segments is operably linked to a transcription promoter selected from the group consisting of casein,  $\beta$ -lactoglobulin,  $\alpha$ -lactalbumin and whey acidic protein gene promoters.

6. A method according to claim 1 wherein said first, second and third DNA segments are expressed under the control of a  $\beta$ -lactoglobulin promoter.

7. A method according to claim 1 wherein said introducing step comprises injecting said first, second and third DNA segments into a pronucleus of said fertilized egg.

8. A method according to claim 1 wherein said fibrinogen is human fibrinogen.

9. A method according to claim 1 wherein said second DNA segment comprises a sequence of nucleotides as shown in SEQ ID NO: 3 from nucleotide 470 to nucleotide 8100.

10. A method according to claim 1 wherein said second DNA segment comprises a sequence of nucleotides as shown in SEQ ID NO: 3 from nucleotide 512 to nucleotide 8100.

11. A method according to claim 1 wherein said species into which said DNA segments is introduced is sheep.

12. A method of producing biocompetent fibrinogen comprising:

incorporating *into operable linkage* a [first] DNA segment encoding a secretion signal [operably linked to], *a genomic DNA segment encoding an  $\text{A}\alpha$  chain of fibrinogen [into a  $\beta$ -lactoglobulin gene] and an additional segment required for expression of the  $\text{A}\alpha$  chain* in the mammary gland of a mammal to produce a first gene

fusion [comprising a  $\beta$ -lactoglobulin promoter operably linked to the first DNA segment];

incorporating *into operable linkage* a [second] DNA segment encoding a secretion signal [operably linked to], *a genomic DNA segment encoding a  $\text{B}\beta$  chain of fibrinogen [into a  $\beta$ -lactoglobulin gene] and an additional segment required for expression of the  $\text{B}\beta$  chain* to produce a second gene fusion [comprising a  $\beta$ -lactoglobulin promoter operably linked to the second DNA segment];

incorporating *into operable linkage* a [third] DNA segment encoding a secretion signal [operably linked to], *a genomic DNA segment encoding a  $\gamma$  chain of fibrinogen and an additional segment required for expression of the  $\gamma$  chain [into a  $\beta$ -lactoglobulin gene]* to produce a third gene fusion, [comprising a  $\beta$ -lactoglobulin promoter operably linked to the third DNA segment] wherein each of said first, second and third segments are of the same species;

*linking the first, second and third gene fusions in a single vector*; introducing said first, second and third gene fusions into the germ line of a non-human mammal so that said DNA segments are expressed in a mammary gland of said mammal or its female progeny and biocompetent fibrinogen is secreted into milk of said mammal or its female progeny;

obtaining milk from said mammal or its female progeny; and

recovering said fibrinogen from said milk.

13. A method according to claim 12 wherein said mammal is a sheep, pig, goat or cow.

14. A method according to claim 12 wherein each of said first, second and third gene fusions comprises an intron.]

15. A method according to claim 12 wherein the molar ratio of said first, second and third gene fusions introduced is within the range of 0.5-1:0.5-1:0.5-1.]

16. A method according to claim 12 wherein said introducing step comprises injecting said first, second and third gene fusions into a pronucleus of a fertilized egg and inserting said egg into an oviduct of a pseudopregnant female to produce female offspring carrying said gene fusions in the germ line, wherein said egg and said pseudopregnant female are of the same species.

17. A method according to claim 12 wherein said mammal is a sheep.

18. A method for producing biocompetent fibrinogen comprising:

providing a transgenic female non-human mammal carrying in its germline heterologous DNA segments encoding  $\text{A}\alpha$ ,  $\text{B}\beta$  and  $\gamma$  chains of fibrinogen, wherein said segments are expressed in a mammary gland of said mammal and biocompetent fibrinogen encoded by said segments is secreted into milk of said mammal;

collecting milk from said mammal; and  
recovering said biocompetent fibrinogen from said milk.]

19. A method according to claim 18 wherein said mammal is a sheep, pig, goat or cow.]

20. A method according to claim 18 wherein said mammal is a sheep.]

21. A transgenic non-human female mammal that produces recoverable amounts of biocompetent human fibrinogen in its milk, wherein said mammal comprises:

a first DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\text{A}\alpha$  chain,  
a second DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\text{B}\beta$  chain, and  
a third DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\gamma$  chain, and

further wherein each chain is derived from the same species and is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal.]

[22. A mammal according to claim 21 wherein said mammal is a sheep.]

23. A process for producing a transgenic offspring of a mammal comprising:

providing a first DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen A $\alpha$  chain, *the DNA segment comprising genomic DNA encoding the A $\alpha$  chain*; a second DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen B $\beta$  chain, *the DNA segment comprising genomic DNA encoding the B $\beta$  chain*; and a third DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\gamma$  chain, *the DNA segment comprising genomic DNA encoding the  $\gamma$  chain*; wherein each chain is derived from the same species, and wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal;

*linking the first, second and third segments in a single vector*;

introducing said DNA segments into a fertilized egg of a non-human mammalian species heterologous to the species of origin of said fibrinogen chains;

inserting said fertilized egg into an oviduct or uterus of a female of said mammalian species; and

allowing said fertilized egg to develop thereby producing transgenic offspring carrying said first, second and third DNA segments, wherein female progeny of said mammal express said DNA segments in a mammary gland to produce biocompetent fibrinogen.

24. A process according to claim 23 wherein said offspring is female.

25. A process according to claim 23 wherein said offspring is male.

[26. A non-human mammal produced according to the process of claim 23.]

[27. A non-human mammal according to claim 26 wherein said mammal is female.]

[28. A non-human female mammal according to claim 27 that produces milk containing biocompetent fibrinogen encoded by said DNA segments.]

[29. A non-human mammal according to claim 26 wherein said mammal is male.]

[30. A non-human mammal carrying in its germline DNA segments encoding human A $\alpha$ , B $\beta$  and  $\gamma$  chains of fibrinogen, wherein female progeny of said mammal express said DNA segments in a mammary gland to produce biocompetent human fibrinogen.]

[31. A mammal non-human according to claim 30 wherein said mammal is female.]

[32. A mammal non-human according to claim 30 wherein said mammal is male.]

[33. A mammal according to claim 30, wherein said mammal is a sheep.]

34. *A set of DNA sequences comprising:*

*a first DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen A $\alpha$  chain, the DNA segment comprising genomic DNA encoding the A $\alpha$  chain;*

*a second DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen B $\beta$  chain, the DNA segment comprising genomic DNA encoding the B $\beta$  chain; and*

*a third DNA segment encoding a secretion signal operably linked to a heterologous fibrinogen  $\gamma$  chain, the DNA segment comprising genomic DNA encoding the  $\gamma$  chain, wherein each chain is from the same species, and wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal;*

*and the first, second, third segments are linked in a single vector.*

\* \* \* \* \*