



(43) International Publication Date

16 April 2015 (16.04.2015)

(51) International Patent Classification:

A61K 31/045 (2006.01) *A61K 31/18* (2006.01)
A61K 31/10 (2006.01) *C12Q 1/26* (2006.01)

(21) International Application Number:

PCT/US2014/060174

(22) International Filing Date:

10 October 2014 (10.10.2014)

STAT, Michael [US/US]; 203 Glendale Road, Wallingford, PA 19806 (US). **TAYLOR-FISHWICK, David** [US/US]; 825 Fairfax Avenue, Suite 410, Norfolk, VA 23507 (US). **SIMEONOV, Anton** [US/US]; 7805 Green Twig Road, Bethesda, MD 20817 (US). **YASGAR, Adam** [US/US]; 2939 Van Ness Street, Apt. 703, Washington, DC 20008 (US).

(25) Filing Language:

English

Agents: **WISZ, Jamie, T.** et al.; Wilmer Cutler Pickering Hale And Dorr LLP, 1875 Pennsylvania Avenue, NW, Washington, DC 20006 (US).

(26) Publication Language:

English

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(30) Priority Data:

61/889,396 10 October 2013 (10.10.2013) US
61/987,129 1 May 2014 (01.05.2014) US

(81) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,


(71) Applicant: **EASTERN VIRGINIA MEDICAL SCHOOL** [US/US]; Office Of Technology Transfer, 721 Fairfax Avenue, 120 Andrews Hall, Norfolk, VA 23507 (US).

(84) Designated States (unless otherwise indicated, for every kind of national protection available): ARIGO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,

[Continued on next page]

(54) Title: 4-((2-HYDROXY-3-METHOXYBENZYL)AMINO) BENZENESULFONAMIDE DERIVATIVES AS 12-LIPOXY-GENASE INHIBITORS

Scheme 1. Synthesis of analogs 1-83.



Figures 1A – 1D

(57) Abstract: Human lipoxygenases (LOXs) are a family of iron-containing enzymes involved in catalyzing the oxidation of poly-unsaturated fatty acids to provide the corresponding bioactive hydroxyeicosatetraenoic acid (HETE) metabolites. These eicosanoid signaling molecules are involved in a number of physiologic responses such as platelet aggregation, inflammation, and cell proliferation. Platelet-type 12-(S)-LOX (12-LOX) is of particular interest because of its demonstrated role in skin diseases, diabetes, platelet hemostasis, thrombosis, and cancer. Disclosed herein is the identification and medicinal chemistry optimization of a 4-((2-hydroxy-3-methoxybenzyl)amino)benzenesulfonamide-based scaffold. The compounds display nM potency against 12-LOX and excellent selectivity over related lipoxygenases and cyclooxygenases. In addition to possessing favorable ADME properties, the compounds also inhibit PAR-4 induced aggregation and calcium mobilization in human platelets, and reduce 12-HETE in mouse/human beta cells. The compounds can also be used in methods for treating or preventing a 12-lipoxygenase mediated disease or disorder.



LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, **Published:**
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *with international search report (Art. 21(3))*
GW, KM, ML, MR, NE, SN, TD, TG).

4-((2-HYDROXY-3-METHOXYBENZYL)AMINO) BENZENESULFONAMIDE DERIVATIVES AS 12-LIPOXYGENASE INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This patent application claims the benefit of U.S. Provisional Application No. 61/889,396, filed on October 10, 2013, and U.S. Provisional Application No. 61/987,129, filed on May 1, 2014, which are both incorporated by reference.

STATEMENT CONCERNING GOVERNMENT RIGHTS IN FEDERALLY-SPONSORED RESEARCH

[0002] This invention was made with government support under the intramural research program of the National Center for Advancing Translational Sciences and the Molecular Libraries Initiative of the National Institutes of Health Roadmap for Medical Research grant no. U54MH084681, the National Institute of Health grant no. R01 GM56062, the Molecular Libraries Initiative of the National Institutes of Health Roadmap for Medical Research grant no. R03 MH081283, NIH grant no. S10-RR20939, the National Heart, Lung, and Blood Institute (NHLBI) grant no. HL114405, and the National Institute of General Medical Sciences (NIGMS) grant no. GM105671. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

[0003] Lipoxygenases are a class of non-heme iron-containing enzymes which regio- and stereospecifically oxidize polyunsaturated fatty acid substrates such as arachidonic acid (AA) and linoleic acid (LA). (Solomon, et al. *Chem. Biol.* **1997**, *4*, 795–808; Brash, *J. Biol. Chem.* **1999**, *274*, 23679–23682.) The position at which these *cis*, *cis*-1,4-pentadiene substrates are oxidized correspond to the requisite lipoxygenase, with the three major human lipoxygenases: 5-LOX, 12-LOX, and 15-LOX-1, oxidizing the C-5, C-12 and C-15 positions respectively. Lipoxygenases are involved in the first committed step in a cascade of metabolic pathways and the products of these enzymes (eicosanoids) are precursors of hormones such as leukotrienes and lipoxins, which mediate a wide array of cellular functions. (Serhan, et al. *Chem. Rev.* **2011**, *111*, 5922-5943.) Consequently, the lipoxygenase enzymes and their bioactive metabolites (e.g. hydroxyeicosatetraenoic acid (HETE) and leukotriene A₄) have been implicated in a variety of inflammatory diseases and cancers.

[0004] Specifically, 5-LOX has been implicated in cancer and inflammatory diseases, such as asthma and remains the only lipoxygenase enzyme for which there is an FDA-approved inhibitor (Zilueton) on the market.

[0005] Reticulocyte 15-LOX-1 has a role in atherogenesis, neurodegenerative diseases, and neuronal damage associated from an acute ischemic stroke event.

[0006] 12-LOX exists as three isozymes, platelet-type, leukocyte, and epidermal, but leukocyte 12-LOX is found in rat, mouse, pig and cow, but not in humans. (Yamamoto *Biochim. Biophys. Acta.* **1992**, *1128*, 117-131; Funk et al. *FEBS Lett.* **1997**, *402*, 162-166).

[0007] Skin Disease and Platelet Hemostasis: Further, 12-LOX has demonstrated a role in skin diseases and platelet hemostasis.

[0008] Transplants: 12-LOX inhibitors also have utility in transplantation/xenotransplantation scenarios, where, for example, islets can be treated *ex vivo* to improve survival prior to transplant.

[0009] Cancer: Platelet-type 12-(S)-LOX (12-LOX) has been found to be overexpressed in a variety of tumor tissues including prostate cancer, colorectal cancer, breast cancer and lung cancer. (Catalano et al. *Histol. Histopathol.* **2005**, *20*, 969-975; Nie, D et al. *Cancer Res.* **1998**, *58*, 4047-4051; Natarajan et al. *J. Clin. Endocrinol. Metab.* **1997**, *82*, 1790-1798; Kamitani et al. *Adv. Exp. Med. Biol.* **1999**, *469*, 593-598; Soriano et al. *Cancer Res.* **1999**, *59*, 6178-6184). Moreover, 12-HETE levels have been linked to increased cancer cell metastasis by facilitating tumor cell motility and angiogenesis. (Nappez et al. *Cancer Lett.* **1995**, *96*, 133-140; Timár et al. *Int. J. Cancer*, **1993**, *55*, 1003-1010; Honn et al. *Exp. Cell. Res.* **1994**, *214*, 120-130; Nie et al. *Blood* **2000**, *95*, 2304-2311).

[0010] Diabetes: Type 1 and Type 2 diabetes are serious disorders that can lead to major complications and reduced lifespan. There is an unmet medical need in identifying new ways to protect beta cells in these metabolic disorders. 12-LOX is expressed in human pancreatic islets, which is upregulated and activated by inflammatory cytokines leading to increased 12-LOX translocation. The resulting 12-HETE product leads to reduced insulin secretion, reduced metabolic activity and pancreatic β cell death through the amplification of the inflammatory response. (Chen et al. *Diabetologia* **2005**, *48*, 486-495). Both non-obese diabetic (NOD) 12-LOX and 12-LOX knock-out (“KO”) mice showed significant resistance to the development of diabetes compared to the controls, showing 12-LOX is a regulator in this disease.

[0011] Diabetic Kidney Disease: Further, studies show that activation of the 12-LOX pathway plays a role in the development of diabetic kidney disease (diabetic nephropathy) by multiple pathogenic mechanisms, including decreased expression of glomerular P-cadherin. (Guo, Q. et al. *Am. J. Physiol. Endocrinol. Metab.* **2011**, *300*, E708-E716).

[0012] Diabetic Nerve Disease: It has also been shown that increased aldose reductase, the first enzyme of the sorbitol pathway, activity plays a key role in diabetes-associated 12/15-LOX activation in the peripheral nerve and spinal cord. (Stavniichuk, R. et al. *Biochem Pharmacol.* 2012, 83, 932-940). Thus, inhibiting 12-LOX is attractive in treating diabetic nerve disease.

[0013] Diabetes and Cardiovascular Disease: A selective 12-LOX inhibitor would provide a new therapeutic approach to prevent and/or treat either form of diabetes (type I and type II). The development of 12-hLOX inhibitors provide a potent intracellular approach to decreasing the ability of platelets to form large clots in response to vessel injury or activation of the coagulation pathway. Thus, 12-hLOX inhibition has the ability to attenuate platelet-mediated clot formation caused by diabetes and/or cardiovascular disease and significantly decrease the occurrence of myocardial infarction, congestive heart failure, and stroke. Additionally, studies show that the gene *Alox15* that encodes the proteins 12-LOX and 15-LOX are up-regulated in heart failure. Thus, inhibition of 12-LOX could be a treatment for heart failure. (Kayama, Y. et al. *J.Exp.Med.* 2009, 206, 1565-1574).

[0014] Thrombosis: 12-LOX and its product 12-HETE have been implicated in the modulation of hemostasis and thrombosis via their role in regulating platelet function (reactivity, clot formation, calcium mobilization). (Brash, A. R. *Circulation* 1985, 72, 702-707). Additionally, Fc γ RIIa is the receptor on the human platelet responsible for heparin induced thrombocytopenia (HIT). It has been found that 12-LOX is essential for Fc γ RIIa-induced PLC γ 2 activity leading to activation of calcium mobilization, Rap 1 and PKC activation, and subsequent activation of the integrin α IIb β 3, which demonstrates the role of 12-LOX inhibitors in treating HIT. (Yeung, J. et al. *Blood* 2014, (DOI 10.1182/blood-2014-05-575878)). Further, 12-LOX has demonstrated a role in skin diseases and platelet hemostasis.

[0015] Alzheimer's disease: 12-LOX and 15-LOX are widely expressed in the central nervous system and have been reported to be involved in neurobiology of Alzheimer's disease because it modulates amyloid beta and APP processing. It has also been found that 12-LOX and 15-LOX modulate endogenous tau metabolism, making it an attractive therapeutic for treating Alzheimer's and related diseases. (Giannopoulos, P.F., et al. *Aging Cell* 2013, 12, 1082-1090).

[0016] Non-Alcoholic steatohepatitis: It has been shown that disruption of the gene encoding for 12-LOX, *Alox15*, protected mice against hepatic steatosis, insulin resistance, and inflammation in experimental liver disease of metabolic origin. (Martinez-Clemente, M. et al. *Hepatology* 2010, 52, 1980-1991; *see also* Tanaka, N. et al. *Hepatology* 2012, 56, 118-129).

[0017] One difficulty in being able to clearly define the role of 12-LOX in these systems has been the lack of potent and selective 12-LOX small molecule inhibitors.

[0018] A previously reported 12-LOX inhibitor, an 8-Hydroxyquinoline based compound (**ML127**), exhibited excellent selectivity, >50-100 fold selectivity over related lipoxygenases and cyclooxygenase. In contrast to many of the previously reported inhibitors, kinetic experiments revealed that **ML127** was a non-competitive and non-reductive inhibitor. Chiral HPLC separation of the probe molecule revealed a chiral preference for activity with the (−)-enantiomer being much more potent than the (+)-enantiomer (<0.5 μ M vs. >25 μ M, respectively). (Kenyon, V. et al. *J. Med. Chem.* **2011**, *54*, 5485-5497.) However, the chemical series was difficult to optimize further, given that subtle structural modifications led to diminished activity.

[0019] There exists a need for a potent, selective 12-LOX small molecule inhibitor that can be optimized without reducing activity to treat or prevent 12-LOX mediated diseases and disorders. The small molecule inhibitor should be soluble, have favorable ADME properties, and have good *in vivo* PK properties.

[0020] Platelets express three immunoreceptor tyrosine-based activation motif (ITAM) containing transmembrane receptors (glycoprotein VI (GPVI)/FcR γ complex, C-type lectin-like receptor 2 (CLEC-2), and Low affinity immunoglobulin gamma Fc region receptor II-a (Fc γ RIIa)). Ligation of ITAM containing receptors on the surface of platelets leads to a shared downstream signaling pathway culminating in platelet activation. These receptors engage in various degrees of hemostasis and thrombosis; however, they have non-redundant (patho) physiological functions. Fc γ RIIa, a broadly expressed immunoreceptor which is present on the surface of human but not mouse platelets is best known for its pathophysiological role in immune-mediated thrombocytopenia and thrombosis; a family of disorders including immune thrombocytopenia, thrombocytopenia associated with sepsis, and heparin-induced thrombocytopenia (HIT). Selectively inhibiting the Fc γ RIIa signaling pathway in platelets for prevention of immune-mediated thrombocytopenia and thrombosis has been a long sought approach for prevention of HIT (Reilly et al., *Blood* 2011, *117*, 2241-2246).

[0021] Heparin is widely used in the clinic to treat thrombosis. However, more than 3-5% of patients taking heparin will develop an immune response to heparin and be at high risk for heparin-induced thrombocytopenia (HIT), which can lead to a life-threatening thrombotic event mediated by the immune system. The current therapeutic approach to treatment of HIT is the removal of heparin treatment and replacement with direct thrombin inhibitors (DTIs) which have an inherently high risk for serious bleeding and must be monitored in the clinic. Even with this potentially fatal complication, heparin remains the standard anticoagulant for prevention and treatment of thrombosis.


[0022] Therefore, there is also a need for novel therapeutic approaches that directly target the pathogenesis of HIT.

SUMMARY OF THE INVENTION

[0023] The present invention provides compounds and methods comprising compounds useful as 12-LOX inhibitors.

[0024] The invention provides compounds that are potent and selective inhibitors of 12-LOX. In addition, the present invention provides compositions comprising these compounds and methods of using these compounds as therapeutic agents in the treatment or prevention of 12-lipoxygenase mediated diseases and disorders. 12-LOX mediated diseases and disorders include those in which 12-LOX is a direct mediator of the diseases and disorders, as well as those in which the inhibition of 12-LOX, although perhaps not a direct mediator, results in therapeutic value in the treatment or prevention of the diseases and disorders. These diseases or disorders include, but are not limited to, type 1 diabetes, type 2 diabetes, diabetic kidney disease, diabetic nerve disease, cardiovascular disease, congestive heart failure, myocardial infarction, stroke, Alzheimer's disease, Non-Alcoholic steatohepatitis, platelet hemostasis, skin diseases, heparin induced thrombocytopenia, thrombosis, prostate cancer, colorectal cancer, breast cancer and lung cancer.

[0025] The invention provides a compound of Formula (I):




wherein R₁ and R₂ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, F, Cl, Br, amine, nitrogen dioxide, indole, alkoxy, cycloalkyl, aryl, heterocycloalkyl, heteroaryl, each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, methoxy;

[0026] R₃ is selected from the group consisting of phenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, thiazole, benzothiazole, benzoxazole, imidazole, benzimidazole, thiophene, 1-naphthalene, 2-naphthalene, pyridine, quinoline, isoquinoline, 4N-boc-piperidine-3-phenyl, oxazole, benzothiophene, parathiazine, furan, pyran, chromene, benzofuran, pyrrole, pyrazole, pyrazine, pyrimidine, triazine, indole, purine, phthalazine,; each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, alkoxy, phenyl, cycloalkyl, aryl, piperazine, piperidine, pyridine, morpholine, pyrrolidine, pyrazolidine, imidazolidine, and thiomorpholine;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof;

with the proviso that the compound is not



[0027] The invention also provides pharmaceutical compositions comprising a compound, salt, enantiomers, a mixture of enantiomers, or diastereomers of the invention and a pharmaceutically acceptable carrier.

[0028] The invention further provides a method for treating or preventing a 12-lipoxygenase mediated disease or disorder, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of a compound, salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof of a compound of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] **Figures 1A-1D.** Show the reaction scheme for the synthesis of the analogs of the present invention.

[0030] **Figure 2.** PAR4-AP-induced calcium mobilization (A) and platelet aggregation (B) in human platelets.

[0031] **Figure 3.** Inhibition of 12-HETE in mouse beta cells (A) and human islets (B, C). Glucose-stimulated insulin secretion in human islets (D, E).

[0032] **Figure 4.** Steady-state kinetics data for the determination of K_i for 12-LOX with compound **35 (ML355)**. K_M/V_{max} (x-intercept, K_M/V_{max} units are $\mu\text{M}/\mu\text{mol}/\text{min}/\text{mg}$) versus [Inhibitor] (μM) is the secondary replot of the inhibition data, which yielded a K_i of $0.35 +/- 0.08 \mu\text{M}$.

[0033] **Figure 5.** Steady-state kinetics data for the determination of K_i for 12-LOX with compound **35 (ML355)**. $1/V_{max}$ (y-intercept, $1/V_{max}$ units are $1/\mu\text{mol}/\text{min}/\text{mg}$) versus [Inhibitor] (μM) is the secondary replot of the inhibition data, which yielded a K_i of $0.72 +/- 0.1 \mu\text{M}$.

[0034] **Figure 6.** Steady-state kinetics data for the determination of K_i for 12-LOX with compound **36**. K_M/V_{max} (x-intercept, K_M/V_{max} units are $\mu\text{M}/\mu\text{mol}/\text{min}/\text{mg}$) versus [Inhibitor] (μM) is the secondary replot of the inhibition data, which yielded a K_i of $0.53 +/- 0.2 \mu\text{M}$.

[0035] **Figure 7.** Steady-state kinetics data for the determination of K_i for 12-LOX with compound **36**. $1/V_{max}$ (y-intercept, $1/V_{max}$ units are $1/\mu\text{mol}/\text{min}/\text{mg}$) versus [Inhibitor] (μM) is the secondary replot of the inhibition data, which yielded a K_i of $0.63 +/- 0.1 \mu\text{M}$.

[0036] **Figure 8.** PAR4-AP-induced calcium mobilization (A) and platelet aggregation (B) in human platelets. For A: Washed human platelets (1×10^6 platelets/mL) were stimulated with $200 \mu\text{M}$ PAR4-AP in the absence or presence of increasing concentrations of compound **36**. Calcium mobilization was decreased as the concentration of compound **36** was increased. Calcium was measured in real time using a C6 Accuri flow cytometer. The experiments were done in triplicate. For B: Platelet aggregation of human platelets (3×10^8 platelets/mL) was measured in real-time using a Chronolog Lumi-Aggregometer (model 700D) following addition of PAR4-AP.

[0037] **Figure 9.** Inhibition of 12-HETE by compound **36** in mouse beta cells. Mouse beta cells (BTC3) were treated with arachidonic acid and calcium ionophore (AA/IONO) alone or in the presence of compound **36**. Graphed are the levels of 12-HETE expressed as a percentage of that detected in cells stimulated with AA/IONO alone. The data graphed in figure 3A is a representative experiment with each plotted data point being performed in triplicate. These plotted data are representative of four separate experimental determinations performed covering a lesser dose range. The graphed data are mean \pm SEM, $n=3$. Error bars for some points are masked by the symbol. The data was analyzed by non-linear regression for dose-response curve inhibition, using variable or restricted hill slope, $R^2 > 0.81$. This analysis was facilitated with Prism 5 software. In addition to compound, DMSO (stock solvent) was included with each condition. DMSO is also the solvent for the calcium ionophore (stimulant).

[0038] **Figure 10.** Stability of compound **35 (ML355)** measured as percent composition of probe molecule in aqueous solution (contains 20 % acetonitrile) at r.t. over the indicated time period in (A) pH 2 buffer (pH 7.4) (B) pH 10 buffer (C) PBS buffer (pH 7.4) (D) Lipoxygenase

UV-Vis assay buffer (1M HEPES buffer, pH 7.3) (E) in the presence of 5 mM glutathione (reduced form).

[0039] **Figures. 11A-11B** show that 12-LOX modulates Fc γ RIIa-mediated platelet aggregation. Washed human platelets were pretreated with DMSO (vehicle control) or ML355 (20 μ M) for 15 minutes and platelet aggregation was measured following Fc γ RIIa stimulation by **(Fig. 11A)** anti-CD9 or **(Fig. 11B)** crosslinking of IV.3, a Fc γ RIIa mouse monoclonal antibody, and goat anti-mouse Fab2 (IV.3 + GAM). Left panel depicts a representative aggregation tracing of anti-CD9 antibody stimulated platelets pre-treated with ML355 or DMSO. Right panel depicts final platelet aggregation of anti-CD9 stimulated platelets that had been treated with ML355 (n= 7) or DMSO (n=4). **P<.01.

[0040] **Figures. 12A-12C** show that murine platelets require 12-LOX for normal Fc γ RIIa-induced platelet aggregation. A dose response of anti-CD9-induced platelet aggregation was performed with washed platelets from hFcR/ALOX12^{+/+} or hFcR/ALOX12^{-/-} mice. Prior to aggregation fibrinogen (75 μ g/mL) and CaCl₂ (1 mM) were added to platelets. **(Fig. 12A)** Representative tracings of platelets stimulated with 0.25, 0.5 and 1 μ g/mL of anti-CD9. Inset: Western blots for 12-LOX, Fc γ RIIa, and GAPDH were performed with platelet lysate from hFcR/ALOX12^{+/+} or hFcR/ALOX12^{-/-} mice. The **(Fig. 12B)** lag time and **(Fig. 12C)** final aggregation was measured in hFcR/ALOX12^{+/+} or hFcR/ALOX12^{-/-} mice (n = 3-6 per group).

*P<.05

[0041] **Figures. 13A-13B** show that Fc γ RIIa-mediated Rap1 and integrin α IIb β 3 activation are potentiated by 12-LOX. Washed human platelets pre-treated with ML355 or DMSO were stimulated with IV.3 and GAM crosslinking and **(Fig. 13A)** α IIb β 3 integrin activation and **(Fig. 13B)** Rap1 activation were assessed. PAC1-FITC was used to measure α IIb β 3 activation by flow cytometry. A composite bar graph of PAC1-FITC fold changes relative to the unstimulated PAC1-FITC fluorescence. Activated Rap1 was pulled down using Ral-GDS and blotted with a Rap1 antibody. Active Rap1 was measured using LI-COR and then normalized to total Rap1 and unstimulated for fold change in Rap1 activity. ** P<.01

[0042] **Figures. 14A-14B** show that dense granule secretion mediated by Fc γ RIIa activation is regulated by 12-LOX. Washed human platelets pre-treated with DMSO or ML355 were stimulated by Fc γ RIIa crosslinking (IV.3 + GAM) and **(Fig. 14A)** alpha granule secretions was measured by using P-Selectin-PE conjugated antibody in a flow cytometer, while **(Fig. 14B)** ATP secretion was measured as a surrogate marker for dense granule secretion in an aggregometer. A composite of P-Selectin stained platelets treated with ML355 or DMSO (n=4). A bar graph of DMSO or ML355 treated platelets measured for ATP secretion following Fc γ RIIa crosslinking (n=4).

[0043] **Figures. 15A-15B** show the role of 12-LOX in regulating the Fc γ RIIa signaling complex. Washed human platelets were pre-treated with ML355 (20 μ M) or vehicle control prior to IV.3 and GAM stimulation. (**Fig. 15A**) Immunoprecipitation of Fc γ RIIa at 15, 30, and 60 seconds post crosslinking and measured for phosphorylation via Western blot. (**Fig. 15B**) Platelets were lysed following Fc γ RIIa cross-linking in the presence of ML355 or DMSO were immunoblotted for active (pY323) and total Syk.

[0044] **Figures. 16A-16C** show that 12-LOX modulates early signaling components of the Fc γ RIIa pathway in human platelets. Washed human platelets were pre-treated with ML355 (20 μ M) or vehicle control prior to IV.3 and GAM stimulation. (**Fig. 16A**) A time course of PLC γ 2 activation Western blot in which platelets stimulated with antibody crosslinking were stopped at 15, 30, 60, and 300 seconds. Samples of active PLC γ 2 activation were analyzed for Y759 phosphorylation by Western blotting. All samples were normalized to total PLC γ 2 and fold changes were quantified relative to the unstimulated condition. (**Fig. 16B**) Following crosslinking with IV.3 and GAM, calcium mobilization was measured by flow cytometry. Representative curves were quantitated in fold change of free calcium relative to the unstimulated condition over four minutes. Bar graphs represent the ratio of the calcium mobilization fold change. (**Fig. 16C**) Stimulated washed human platelets with or without 12-LOX inhibition were analyzed for PKC activity. A PKC substrate was blotted as a surrogate for PKC activation and Pleckstrin. Quantification of Pleckstrin phosphorylation is shown in the right panel. Data represents mean \pm S.E.M. *P<.05; **P<.01.

[0045] **Figure 17** is a schematic model of 12-LOX role in the regulation of Fc γ RIIa pathway. 12-LOX regulates early PLC γ 2 activation-mediated by Fc γ RIIa stimulation, which is essential for full calcium release in the platelets. Calcium flux is required for cPLA2 activity to generate free fatty acids, such as arachidonic acid (AA). Subsequently, Rap1 activation is also dependent on 12-LOX activity in order to activate integrin α IIb β 3 for platelet aggregation.

[0046] **Figure 18** shows 1 H NMR data for compound ML355 (**Fig. 18A**) and 13 C NMR for compound ML-355 (**Fig. 18B**).

[0047] **Figure 19** shows LC/MS characterization for compound ML355 at 220 nm (top) and 254 nm (bottom).

[0048] **Figure 20** shows 1 H NMR data for compound 36 (**Fig. 20A**) and 13 C NMR for compound 36 (**Fig. 20B**).

[0049] **Figure 21** shows LC/MS characterization for compound 36 at 220 nm (top) and 254 nm (bottom).

[0050] **Figure 22** shows 1 H NMR data for compound 37 (**Fig. 22A**) and 13 C NMR for compound 37 (**Fig. 22B**).

[0051] Figure 23 shows LC/MS characterization for compound 37 at 220 nm (top) and 254 nm (bottom).

DETAILED DESCRIPTION OF THE INVENTION

[0052] It is understood that the present invention is not limited to the particular methodologies, etc. described herein, as these may vary. It is also to be understood that the terminology used herein is used for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention.

[0053] In the description, the following terms are employed: LOX is lipoxygenase; 15-LOX-1 is human reticulocyte 15-lipoxygenase-1; 15-LOX-2 is human epithelial 15-lipoxygenase-2; 12-LOX is human platelet 12-lipoxygenase; COX is Cyclooxygenase; NDGA is nordihydroguaiaretic acid; AA is arachidonic acid; 12-HPETE is 12-(S)-hydroperoxyeicosatetraenoic acid; 12-HETE is 12-(S)-hydroxyeicosatetraenoic acid; LA is linoleic acid; V_{max} is maximal velocity (mmol/min); K_M is Henri-Michaelis-Menten Constant; [E] is total active enzyme concentration; IC_{50} is inhibitor constant at 50% inhibition; HTS is high-throughput screening; qHTS is quantitative high-throughput screening; PAR-4 is protease-activated receptor-4; EtOH is ethanol; MeOH is methanol; EtOAc is ethyl acetate; AcOH is acetic acid; MW is microwave; Xantphos is 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene; and SD is standard deviation.

[0054] For therapeutic use, “salts” of the compounds of the present invention will be physiologically acceptable, i.e., the salts will be derived from a physiologically acceptable acid or base. Salts of acids or bases, however, which are not physiologically acceptable may also find use in the preparation or purification of a physiologically acceptable compound. Thus, all salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the present invention.

[0055] “Diastereomer” refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another. Diastereomers have different physical properties, e.g., melting points, boiling points, spectral properties, and reactivities. Mixtures of diastereomers may separate under high resolution analytical procedures such as electrophoresis and chromatography.

[0056] “Enantiomers” refer to two stereoisomers of a compound which are non-superimposable mirror images of one another.

[0057] The term “pharmaceutically acceptable carrier,” as used herein, refers to any and all solvents, dispersion media, coatings, antibacterial and antifungal agent, isotonic and absorption

delaying agents for pharmaceutical active substances as are well known in the art. Except insofar as any conventional media or agent is incompatible with the compound, its use in the therapeutic compositions is contemplated. Supplementary compounds can also be incorporated into the compositions.

[0058] The expression “therapeutically effective amount” refers to an amount of a compound disclosed herein, that is effective for preventing, ameliorating, treating or delaying the onset of a disease or condition. The expression “prophylactically effective amount” refers to an amount of a compound disclosed herein, that is effective for inhibiting the onset or progression of a disorder.

[0059] “Alkyl” is C1-C6 hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms. Examples include, but are not limited to, methyl, ethyl, propyl, isopropyl, tert-butyl, and the like.

[0060] “Alkenyl” is C2-C6 hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, i.e. a carbon-carbon, sp² double bond. Examples include, but are not limited to, ethylene or vinyl (-CH=CH₂), allyl (-CH₂CH=CH₂), cyclopentenyl (-C₅H₇), and 5-hexenyl (-CH₂CH₂CH₂CH₂CH=CH₂), and the like.

[0061] “Alkynyl” is C2-C6 hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, i.e., a carbon-carbon, sp triple bond. Examples include, but are not limited to, acetylenic (-C=--CH) and propargyl (-CH₂C=CH), and the like.

[0062] “Cycloalkyl” is a saturated, unsaturated or aromatic ring system containing from, for example, 3 to 8 carbon atoms, preferably 3 to 7 carbon atoms, and more preferably 3 to 6 carbon atoms. Examples include, but are not limited to, cyclopentenyl and cyclohexenyl. The cyclic alkyl groups may be unsubstituted or further substituted.

[0063] “Heterocycloalkyl” means a saturated, unsaturated or aromatic ring system including at least one N, O, S, or P. Heterocycle thus includes heteroaryl groups. Heterocycle as used herein includes, but is not limited to heterocycles described in PAQUETTE, PRINCIPLES OF MODERN HETEROCYCLIC CHEMISTRY (W. A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; THE CHEMISTRY OF HETEROCYCLIC COMPOUNDS, A SERIES OF MONOGRAPHS (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; KATRITZKY ET AL., COMPREHENSIVE HETEROCYCLIC CHEMISTRY (Pergamon Press, 1996); and 82 J. AM. CHEM. SOC. 5566 (1960). Heterocycles include, but are not limited to pyridyl, dihydropyridyl, tetrahydropyridyl (piperidyl), tiazolyl, tetrahydrothiophenyl, sulfur oxidized tetrahydrotiophenyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl, indolyl, indolenyl, quinolinyl, isoquinolinyl, benzimidazolyl, piperidinyl, 4-piperidonyl, pyrrolidinyl, 2-pyrrolidonyl, pyrrolinyl, tetrahydroturanyl, bis-tetrahydronuranyl,


tetrahydropyranyl, bis-tetrahydropyranyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, octahydroisoquinolinyl, azocinyl, triazinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, thienyl, thianthrenyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxyathinyl, 2H-pyrrolyl, isothiazolyl, isoxazolyl, pyrazinyl, pyridazinyl, indolizinyl, isoindolyl, 3H-indolyl, 1H-indazolyl, purinyl, 4H-quinolizinyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, 4aH-carbazolyl, carbazolyl, beta-carbolinyl, phenanthridinyl, acridinyl, pyrimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, rurazanyl, phenoxyazinyl, isochromanyl, chromanyl, imidazolidinyl, imidazoliny, pyrazolidinyl, pyrazoliny, piperazinyl, indolinyl, isoindolinyl, quimiclidinyl, morpholinyl, oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazoliny, and isatinoyl.

[0064] “Aryl” means a monovalent aromatic hydrocarbon radical of 6-20 carbon atoms derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Typical aryl groups include, but are not limited to, radicals derived from benzene, substituted benzene, naphthalene, anthracene, biphenyl, and the like.

[0065] “Heteroaryl” means a monovalent aromatic radical of one or more carbon atoms and one or more atoms selected from N, O, S, or P, derived by the removal of one hydrogen atom from a single atom of a parent aromatic ring system. Heteroaryl groups may be a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 3 heteroatoms selected from N, O, P, and S) or a bicyclic having 7 to 10 ring members (4 to 9 carbon atoms and 1 to 3 heteroatoms selected from N, O, P, and S). Heteroaryl bicyclics have 7 to 10 ring atoms (6 to 9 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S) arranged as a bicyclo [4,5], [5,5], [5,6], or [6,6] system; or 9 to 10 ring atoms (8 to 9 carbon atoms and 1 to 2 hetero atoms selected from N and S) arranged as a bicyclo [5,6] or [6,6] system. The heteroaryl group may be bonded to the drug scaffold through a carbon, nitrogen, sulfur, phosphorus or other atom by a stable covalent bond. Heteroaryl groups include, but are not limited to, pyridyl, dihydropyridyl isomers, pyridazinyl, pyrimidinyl, pyrazinyl, s-triazinyl, oxazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, furanyl, thiofuranyl, thienyl, and pyrrolyl.

[0066] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Preferred methods and compositions are described, although any methods and compositions similar or equivalent to those described herein can be used in the practice or testing of the present invention.


[0067] In one aspect, the present invention includes a compound of Formula (I):



wherein R₁ and R₂ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, F, Cl, Br, amine, nitrogen dioxide, indole, alkoxy, cycloalkyl, aryl, heterocycloalkyl, heteroaryl, each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, methoxy;

R₃ is selected from the group consisting of phenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, thiazole, benzothiazole, benzoxazole, imidazole, benzimidazole, thiophene, 1-naphthalene, 2-naphthalene, pyridine, quinoline, isoquinoline, 4N-boc-piperidine-3-phenyl, oxazole, benzothiophene, parathiazine, furan, pyran, chromene, benzofuran, pyrrole, pyrazole, pyrazine, pyrimidine, triazine, indole, purine, phthalazine,; each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, alkoxy, phenyl, cycloalkyl, aryl, piperazine, piperidine, pyridine, morpholine, pyrrolidine, pyrazolidine, imidazolidine, and thiomorpholine;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof; with the proviso that the compound is not



[0068] In one embodiment R₁ is selected from the group consisting of methoxy and Cl when R₂ is H; R₂ is selected from the group consisting of Br, and Cl when R₁ is H; and R₃ is selected from the group consisting of thiazole, 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, 4-methyl-2-thiazole, 5-methyl-2-thiazole, 4, 5-methyl-2-thiazole, phenyl, 1-naphthalene, 2-naphthalene, 1, 4-bi-phenyl, 3-piperazine-phenyl, 4-piperidine-phenyl, 4-piperazine-3-pyridine, 6-methyl-3-pyridine, 3-quinoline, 8-isoquinoline, 2-pyridine, 3-pyridine, 3-tertbutyl-phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, 4-phenyl-2-thiazole, 3-morpholine-phenyl, 4N-boc-piperidine-3-phenyl, 3-piperidine-phenyl, 3-isopropyl-phenyl, and 4-bi-phenyl; or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

[0069] In any of the above embodiments of Formula (I), R₁ is methoxy and R₂ is H.

[0070] In any of the above embodiments of Formula (I), the compound is a pro-drug.

[0071] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, 3-quinoline, 8-isoquinoline, phenyl, and 3-isopropyl-phenyl.

[0072] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-naphthalene, 3-quinoline, 8-isoquinoline, 3-tertbutyl-phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, 4-phenyl-2-thiazole, 4N-boc-piperidine-3-phenyl, and 3-isopropyl-phenyl.

[0073] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-naphthalene, 3-quinoline, 8-isoquinoline, 3-tertbutyl-phenyl, 4N-boc-piperidine-3-phenyl, and 3-isopropyl-phenyl.


[0074] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, and 3-isopropyl-phenyl.

[0075] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, and 4-phenyl-2-thiazole.

[0076] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, and phenyl.


[0077] In one embodiment of Formula (I), R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-naphthalene, 3-quinoline, 8-isoquinoline, 2-pyridine, and 3-pyridine.

[0078] In one embodiment the compound of formula (I) is




[0079] or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof. This is **ML355 (compound 35)**.

[0080] In one embodiment the compound of formula (I) is




[0081] or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

[0082] In one embodiment the compound of formula (I) is



[0083] or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

[0084] In another aspect, the present invention includes a compound of Formula (II):




wherein X is selected from the group consisting of O, S, NH, and C;

wherein R₁ and R₂ are independently selected from the group consisting of H, halogen, hydroxyl, alkoxy, and alkyl;

wherein R₃ through R₆ are independently selected from the group consisting of H, halogen, alkoxy, and alkyl;

[0085] or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

[0086] In another aspect, the present invention includes a compound of Formula (III):




wherein R₁ and R₂ are independently selected from the group consisting of H, halogen, and alkoxy, and further wherein R₁ and R₂ are not both H;

wherein R₄ and R₅ are independently selected from the group consisting of H, alkyl, phenyl, and optionally substituted phenyl;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof;

with the proviso that the compound is not



In one embodiment, R_1 is not a halogen.

In one embodiment, R_1 is not F.


In one embodiment, R_1 is not Br.

In one embodiment, when R_1 is H, R_2 is not methoxy.

In one embodiment, R_5 is not phenyl.


In one embodiment, when R_4 is H, R_5 is not phenyl.

In another aspect, the present invention includes a compound of Formula (IV):




wherein R_1 and R_2 are independently selected from the group consisting of H, halogen, and alkoxy;

wherein R_3 and R_4 are independently selected from the group consisting of H, phenyl,




optionally substituted phenyl, tert-butyl, isopropyl, and

wherein X is selected from the group consisting of NH, O, and



or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.




In one embodiment, when R₄ is , X is not NH.

In one embodiment, R₃ is not phenyl.

In one embodiment, when R₄ is H, R₃ is not phenyl.

In another aspect, the present invention includes a compound of Formula (V):



[0087] wherein X is selected from the group consisting of N, C, S, and O;


wherein R₁ through R₆ are independently selected from the group consisting of H, alkyl, and alkoxy;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

In one embodiment, X is C and R₁ through R₆ are H.

In one embodiment, X is N and R₁ through R₆ are H.

In another aspect, the present invention includes a compound of Formula (VI):



wherein X is selected from the group consisting of N, C, S, and O;


wherein R₁ through R₆ are independently selected from the group consisting of H, alkyl, and alkoxy;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

In one embodiment, X is C and R₁ through R₆ are H.

In one embodiment, X is N and R₁ through R₆ are H.

[0088] In another aspect, the present invention includes a compound of Formula (VII):



[0089] wherein X is selected from the group consisting of N, C, S, and O;

wherein R₁ through R₃ are independently selected from the group consisting of H, alkyl, and alkoxy;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

[0090] In one embodiment, X is S and R₁ through R₃ are H.

[0091] In any of the above embodiments, the compound or salt of Formulas (I) - (VII) includes each diastereomer, optical isomer, enantiomer and the racemic mixtures.

[0092] It is further understood that the above compounds and salts may form solvates, or exist in a substantially uncomplexed form, such as the anhydrous form. When the solvent incorporated in the solvate is water, the molecular complex is a hydrate. Pharmaceutically acceptable solvates include hydrates, alcoholates such as methanolates and ethanolates, acetonitrilates and the like. These compounds can also exist in polymorphic forms.

[0093] The present invention is further directed to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one compound of Formula (I) – (VII) described herein.

[0094] Routes of administration and dosages of effective amounts of the pharmaceutical compositions comprising the compounds are also disclosed. The compounds of the present invention can be administered in combination with other pharmaceutical agents in a variety of protocols for effective treatment of disease.

[0095] The pharmaceutical compositions of the inventions can be administered to any animal that can experience the beneficial effects of the compounds of the invention. Such animals include humans and non-humans such as pets and farm animals.

[0096] The pharmaceutical compositions of the present invention are administered to a subject in a manner known in the art. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

[0097] In addition to the compounds disclosed herein, the pharmaceutical compositions of the present invention may further comprise at least one of any suitable auxiliaries including, but not limited to, diluents, binders, stabilizers, buffers, salts, lipophilic solvents, preservatives, adjuvants or the like. Pharmaceutically acceptable auxiliaries are preferred. Examples and methods of preparing such sterile solutions are well known in the art and can be found in well known texts such as, but not limited to, REMINGTON'S PHARMACEUTICAL SCIENCES (Gennaro, Ed., 18th Edition, Mack Publishing Co. (1990)). Pharmaceutically acceptable carriers can be routinely selected that are suitable for the mode of administration, solubility and/or stability of the compound.

[0098] Pharmaceutical excipients and additives useful in the present invention can also include, but are not limited to, proteins, peptides, amino acids, lipids, and carbohydrates (e.g., sugars, including monosaccharides, di-, tri-, terra-, and oligosaccharides; derivatized sugars such as alditols, aldonic acids, esterified sugars and the like; and polysaccharides or sugar polymers), which

can be present singly or in combination, comprising alone or in combination in ranges of 1-99.99% by weight or volume. Exemplary protein excipients include serum albumin such as human serum albumin (HSA), recombinant human albumin (rHA), gelatin, casein, and the like. Representative amino acid components, which can also function in a buffering capacity, include alanine, glycine, arginine, betaine, histidine, glutamic acid, aspartic acid, cysteine, lysine, leucine, isoleucine, valine, methionine, phenylalanine, aspartame, and the like.

[0099] Carbohydrate excipients suitable for use in the present invention include monosaccharides such as fructose, maltose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, sucrose, trehalose, cellobiose, and the like; polysaccharides, such as raffinose, melezitose, maltodextrins, dextrans, starches, and the like; and alditols, such as mannitol, xylitol, maltitol, lactitol, xylitol, sorbitol (glucitol), myoinositol and the like.

[0100] The composition further can contain, but is not limited to pharmaceutically acceptable carriers such as coloring agents, emulsifying agents, suspending agents, ethanol, EDTA, citrate buffer, flavoring, and water.

[0101] The composition of the invention also can contain the preservatives methylparaben (also known as 4-hydroxybenzoic acid methyl ester, methyl *p*-hydroxybenzoate; or METHYL CHEMOSEPT), ethylparaben (also known as 4-hydroxybenzoic acid ethyl ester; ethyl *p*-hydroxybenzoate; or ETHYL PARASEPT), propylparaben (also known as 4-hydroxybenzoic acid propyl ester; propyl *p*-hydroxybenzoate; NIPASOL; or PROPYL CHEMOSEPT) and/or butylparaben (also known as 4-hydroxybenzoic acid propyl ester; propyl *p*-hydroxybenzoate; or BUTYL CHEMOSEPT). In some embodiments, the composition contains methylparaben and/or propylparaben.

[0102] Emulsifiers of the invention include, but are not limited to ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

[0103] The pharmaceutical compositions comprising the compounds of the present invention can also include a buffer or a pH adjusting agent. Typically, the buffer is a salt prepared from an organic acid or base. Representative buffers include organic acid salts such as salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid; Tris, tromethamine hydrochloride, or phosphate buffers.

[0104] Additionally, the pharmaceutical compositions of the invention can include polymeric excipients/additives such as polyvinylpyrrolidones, ficolls (a polymeric sugar), dextrans (e.g.; cyclodextrins, such as 2-hydroxypropyl-.beta.-cyclodextrin), polyethylene glycols, flavoring agents,

anti-microbial agents, sweeteners, antioxidants, anti-static agents, surfactants (e.g., polysorbates such as "TWEEN 20" and "TWEEN 80"), lipids (e.g., phospholipids, fatty acids), steroids (e.g., cholesterol), and chelating agents (e.g., EDTA or EGTA). These and additional known pharmaceutical excipients and/or additives suitable for use in the present invention are known in the art, e.g., as listed in REMINGTON: THE SCIENCE & PRACTICE OF PHARMACY (19th ed., Williams & Williams (1995)) and PHYSICIAN'S DESK REFERENCE (52nd ed., Medical Economics (1998)), the disclosures of which are expressly entirely incorporated herein by reference.

[0105] The present invention provides stable pharmaceutical compositions as well as preserved solutions and compositions containing a preservative, as well as multi-use preserved compositions suitable for pharmaceutical or veterinary use, comprising at least one compound disclosed herein in a pharmaceutically acceptable composition. Pharmaceutical compositions in accordance with the present invention may optionally contain at least one known preservative. Preservatives include, but are not limited to, phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, phenylmercuric nitrite, phenoxyethanol, formaldehyde, chlorobutanol, magnesium chloride (e.g., hexahydrate), alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent. Any suitable concentration or mixture can be used as known in the art, such as 0.001-5%, or any range or value therein. Non-limiting examples include, no preservative, 0.1-2% m-cresol, 0.1-3% benzyl alcohol, 0.001-0.5% thimerosal, 0.001-2.0% pheno, 0.0005-1.0% alkylparaben(s), and the like.

[0106] Other excipients, e.g., isotonicity agents, buffers, antioxidants, preservative enhancers, can be optionally added to the diluent. An isotonicity agent such as glycerin, is commonly used at known concentrations. A physiologically tolerated buffer is preferably added to provide improved pH control. The pharmaceutical compositions can cover a wide range of pHs, such as from about pH 4 to about pH 10, specifically, a range from about pH 5 to about pH 9, and more specifically, a range of about 6.0 to about 8.0. In one aspect, the formulations of the present invention have pH between about 6.8 and about 7.8. Suitable buffers include phosphate buffers, sodium phosphate and phosphate buffered saline (PBS).

[0107] Other additives, such as a pharmaceutically acceptable solubilizers like Tween 20 (polyoxyethylene (20) sorbitan monolaurate), TWEEN 40 (polyoxyethylene (20) sorbitan monopalmitate), TWEEN 80 (polyoxyethylene (20) sorbitan monooleate), Pluronic F68 (polyoxyethylene polyoxypropylene block copolymers), and PEG (polyethylene glycol) or non-ionic surfactants such as polysorbate 20 or 80 or poloxamer 184 or 188, PLURONIC® polyols, other block co-polymers, and chelators such as EDTA and EGTA can optionally be added to the

pharmaceutical compositions to reduce aggregation. These additives are particularly useful if a pump or plastic container is used to administer the pharmaceutical composition. The presence of pharmaceutically acceptable surfactant mitigates the propensity for the composition to aggregate.

[0108] The composition of the invention also can contain the preservatives methylparaben (also known as 4-hydroxybenzoic acid methyl ester; methyl p-hydroxybenzoate; or METHYL CHEMOSEPT), ethylparaben (also known as 4-hydroxybenzoic acid ethyl ester; ethyl p-hydroxybenzoate; or ETHYL PARASEPT), propylparaben (also known as 4-hydroxybenzoic acid propyl ester; propyl p-hydroxybenzoate; NIPASOL; or PROPYL CHEMOSEPT) and/or butylparaben (also known as 4-hydroxybenzoic acid propyl ester, propyl p-hydroxybenzoate; or BUTYL CHEMOSEPT). In some embodiments, the composition contains methylparaben and/or propylparaben.

[0109] The compositions of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to the compounds of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art (see Prescott, ed., *METH. CELL BIOL.* 14:33 (1976)). Liposomes, methods of making and methods of use are described in U.S. Patent Nos. 4,089,8091 (process for the preparation of liposomes), 4,233,871 (methods regarding biologically active materials in lipid vesicles), 4,438,052 (process for producing mixed miselles), 4,485,054 (large multilamellar vesicles), 4,532,089 (giant-sized liposomes and methods thereof), 4,897,269 (liposomal drug delivery system), 5,820,880 (liposomal formulations), and so forth.

[0110] During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *PROTECTIVE GROUPS IN ORGANIC CHEMISTRY* (1973); and *GREENE AND WUTS, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS* (1991). The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

[0111] The compound of the invention can be solubilized or suspended in a preconcentrate (before dilutions with a diluent), added to the preconcentrate prior to dilution, added to the diluted preconcentrate, or added to a diluent prior to mixing with the preconcentrate. The compound of the

invention can also be co-administered as part of an independent dosage form, for therapeutic effect. Optionally, the compound of the invention can be present in a first, solubilized amount, and a second, non-solubilized (suspended) amount

[0112] The pharmaceutical formulation can also contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations that can be administered to animals, as described herein.

[0113] For oral administration in the form of a tablet or capsule, a compound may be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents may also be incorporated into the mixture. Suitable binders include, without limitation, starch; gelatin; natural sugars such as glucose or beta-lactose; corn sweeteners; natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose; polyethylene glycol; waxes and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

[0114] For oral administration, the composition also optionally contains a sweetener. Sweeteners include but are not limited to sucrose, fructose, sodium saccharin, sucralose (SPLENDA®), sorbitol, mannitol, aspartame, sodium cyclamate, and the like and combinations thereof.

[0115] The aqueous suspensions, emulsions and/or elixirs for oral administration of this invention can be combined with various sweetening agents, flavoring agents, such as, but not limited to orange or lemon flavors, coloring agents, such as dye stuffs, natural coloring agents or pigments, in addition to the diluents such as water, glycerin and various combinations, as described herein.

[0116] The pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, dragees, cachets or tablets each containing a predetermined amount of the compound; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion, and as a bolus, etc.

[0117] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the compound in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be

made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may be optionally coated or scored and may be formulated so as to provide a slow or controlled release of the compound therein.

[0118] In addition, the compositions comprising compounds may be incorporated into biodegradable polymers allowing for sustained release of the compound. The biodegradable polymers and their uses are described in detail in Brem et al., 74 J. NEUROSURG. 441-46 (1991). Suitable examples of sustained-release compositions include semipermeable matrices of solid hydrophobic polymers containing a compound of the present invention, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (including poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT® (TAP Pharmaceuticals, Inc., Chicago, Ill.) (injectable microspheres composed of lactic acid glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

[0119] Pharmaceutical compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multi-dose containers, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[0120] For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous, administration is desired. The pharmaceutical compositions may be administered parenterally via injection of a pharmaceutical composition comprising a compound dissolved in an inert liquid carrier. The term "parenteral," as used herein, includes, but is not limited to, subcutaneous injections, intravenous, intramuscular, intraperitoneal injections, or infusion techniques. Acceptable liquid carriers include, vegetable oils such as peanut oil, cotton seed oil, sesame oil and the like, as well as organic solvents such as solketal, glycerol formal and the like. The pharmaceutical compositions may be prepared by dissolving or suspending the compound in the

liquid carrier such that the final formulation contains from about 0.005% to 30% by weight of a compound.

[0121] The composition of the invention can also include additional therapeutic agents such as, but not limited to hydrophilic drugs, hydrophobic drugs, hydrophilic macromolecules, cytokines, peptidomimetics, peptides, proteins, toxoids, sera, antibodies, vaccines, nucleosides, nucleotides, nucleoside analogs, genetic materials and/or combinations thereof.

[0122] The additional therapeutic agent can be solubilized or suspended in a preconcentrate (before dilutions with a diluent), added to the preconcentrate prior to dilution, added to the diluted preconcentrate, or added to a diluent prior to mixing with the preconcentrate. The additional therapeutic agent can also be co-administered as part of an independent dosage form, for therapeutic effect. Optionally, the additional therapeutic agent(s) can be present in a first, solubilized amount, and a second, non-solubilized (suspended) amount. Such additional therapeutic agent(s) can be any agent(s) having therapeutic or other value when administered to an animal, particularly to a mammal, such as drugs, nutrients, and diagnostic agents.

[0123] In addition to the compound and compositions of the invention, and additional pharmaceutically active agents, the pharmaceutical formulation can also contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations that can be administered to animals, as described herein.

[0124] Pharmaceutical formulations useful in the present invention can contain a quantity of a compound(s) according to this invention in an amount effective to treat or prevent the condition, disorder or disease of the subject being treated.

[0125] The invention further relates to the administration of at least one compound disclosed herein by the following routes, including, but not limited to oral, parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitory, intracelial, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, iontophoretic means, or transdermal means.

[0126] It can be sometimes desirable to deliver the compounds of the present invention to the subject over prolonged periods of time, for periods of one week to one year from a single administration. Certain medical devices may be employed to provide a continuous intermittent or on demand dosing of a patient. The devices may be a pump or diffusion apparatus, or other device containing a reservoir of drug and optionally diagnostic or monitoring components to regulate the

delivery of the drug. Various slow-release, depot or implant dosage forms can be utilized. A dosage form can contain a pharmaceutically acceptable non-toxic salt of a compound disclosed herein that has a low degree of solubility in body fluids, (a) an acid addition salt with a polybasic acid such as phosphoric acid, sulfuric acid, citric acid, tartaric acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalene mono- or di-sulfonic acids, polygalacturonic acid, and the like; (b) a salt with a polyvalent metal cation such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium and the like, or with an organic cation formed from e.g., N,N'-dibenzyl-ethylenediamine or ethylenediamine; or (c) combinations of (a) and (b) e.g., a zinc tannate salt. Additionally, the compounds of the present invention or a relatively insoluble salt such as those just described, can be formulated in a gel, an aluminum monostearate gel with, e.g., sesame oil, suitable for injection. Salts include, but are not limited to, zinc salts, zinc tannate salts, pamoate salts, and the like. Another type of slow-release depot formulation for injection would contain the compound of salt dispersed or encapsulated in a slow degrading, non-toxic, non-antigenic polymer such as a polylactic acid/polyglycolic acid polymer, including the formulations as described in U.S. Patent No. 3,773,919. The compounds or relatively insoluble salts thereof such as those described above can also be formulated in cholesterol matrix silastic pellets, particularly for use in animals. Additional slow-release, depot or implant formulations, e.g., gas or liquid liposomes are known in the literature. *See, e.g.,* U.S. Patent No. 5,770,222; SUSTAINED AND CONTROLLED RELEASE DRUG DELIVERY SYSTEMS (1978).

[0127] Other examples include provision of the compounds of the present invention to be administered by sustained release delivery system containing a biodegradable composition. The biodegradable composition may be composed of a biodegradable, water-coagulable, non-polymeric material and a biocompatible, non-toxic organic solvent that is miscible to dispersible in an aqueous medium. The delivery system may be implanted at an implant site causing the solvent to dissipate, disperse or leach from the composition into surrounding tissue fluid through a resulting microporous matrix.

[0128] The term “implant site” is meant to include a site, in or on which the non-polymeric composition is applied. Implantation or implant site can also include the incorporation of the pharmaceutical composition comprising at least one compound of the present invention with a solid device. The pharmaceutical composition can be incorporated into a coating on a stent that is implanted into a subject. Additionally, other solid or biodegradeable materials can be used as a substrate on which the pharmaceutical composition is applied. The coated material, comprising the pharmaceutical composition is then implanted, inserted or is adjacent to the subject or patient. The term “biodegradable” means that the non-polymeric material and/or matrix of the implant will

degrade over time by the action of enzymes, by simple or enzymatically catalyzed hydrolytic action and/or by other similar mechanisms in the human body. By "bioerodible," it is meant that the implant matrix will erode or degrade over time due, at least in part, to contact with substances found in the surrounding tissue fluids, cellular action, and the like. By "bioabsorbable," it is meant that the non-polymeric matrix will be broken down and absorbed within the human body, by a cell, a tissue, and the like.

[0129] Non-polymeric materials that can be used in the composition generally are those that are biocompatible, substantially insoluble in water and body fluids, and biodegradable and/or bioerodible. The non-polymeric material is capable of being at least partially solubilized in a water-soluble organic solvent. The non-polymeric materials are also capable of coagulating or solidifying to form a solid implant matrix. The non-polymeric material is combined with a compatible and suitable organic solvent to form a composition that has the desired consistency ranging from watery to viscous to a spreadable putty or paste.

[0130] Suitable organic solvents are those that are biocompatible, pharmaceutically-acceptable, and will at least partially dissolve the non-polymeric material. The organic solvent has a solubility in water ranging from miscible to dispersible. Optionally, a pore-forming agent can be included in the composition to generate additional pores in the implant matrix. The pore-forming agent can be any organic or inorganic, pharmaceutically-acceptable substance that is substantially soluble in water or body fluid, and will dissipate from the coagulating non-polymeric material and/or the solid matrix of the implant into surrounding body fluid at the implant site.

[0131] The compounds of the present invention are capable of providing a local or systemic biological, physiological or therapeutic effect in the body of an animal. In formulating some pharmaceutical compositions described herein, the compound is preferably soluble or dispersible in the non-polymeric composition to form a homogeneous mixture, and upon implantation, becomes incorporated into the implant matrix. As the solid matrix degrades over time, the compound is capable of being released from the matrix into the adjacent tissue fluid, and to the pertinent body tissue or organ, either adjacent to or distant from the implant site, preferably at a controlled rate. The release of the compound from the matrix may be varied by the solubility of the compound in an aqueous medium, the distribution of the compound within the matrix, the size, shape, porosity, and solubility and biodegradability of the solid matrix. *See e.g.*, U.S. Patent No. 5,888,533. The amounts and concentrations of ingredients in the composition administered to the patient will generally be effective to accomplish the task intended.

[0132] In other embodiments, the compounds of the present invention may be administered by bioactive agent delivery systems containing microparticles suspended in a polymer matrix. The

microparticles may be microcapsules, microspheres or nanospheres currently known in the art. The microparticles should be capable of being entrained intact within a polymer that is or becomes a gel once inside a biological environment. The microparticles can be biodegradable or non-biodegradable. Many microencapsulation techniques used to incorporate a bioactive agent into a microparticle carrier are taught in the art. *See e.g.*, U.S. Patent Nos. 4,652,441; 5,100,669; 4,438,253; and 5,665,428.

[0133] A preferred polymeric matrix will be biodegradable and exhibit water solubility at low temperature and will undergo reversible thermal gelation at physiological mammalian body temperatures. The polymeric matrix is capable of releasing the substance entrained within its matrix over time and in a controlled manner. The polymers are gradually degraded by enzymatic or non-enzymatic hydrolysis in aqueous or physiological environments. *See e.g.*, U.S. Patent No. 6,287,588.

[0134] Methods of preparing various pharmaceutical compositions with a certain amount of active ingredients are known, or will be apparent in light of this disclosure, to those skilled in the art. Methods of preparing said pharmaceutical compositions can incorporate other suitable pharmaceutical excipients and their formulations as described in REMINGTON'S PHARMACEUTICAL SCIENCES, Martin, E.W., ed., Mack Publishing Company, 19th ed. (1995).

[0135] Methods of preparing the pharmaceutical preparations of the present invention are manufactured in a manner that is known, including conventional mixing, dissolving, or lyophilizing processes. Thus, liquid pharmaceutical preparations can be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary.

[0136] One of ordinary skill in the art will appreciate that a method of administering pharmaceutically effective amounts of the compositions of the invention to a patient in need thereof, can be determined empirically, or by standards currently recognized in the medical arts. The agents can be administered to a patient as pharmaceutical compositions in combination with one or more pharmaceutically acceptable excipients. It will be understood that, when administered to a human patient, the total daily usage of the agents of the compositions of the present invention will be decided within the scope of sound medical judgment by the attending physician. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors: the type and degree of the cellular response to be achieved; activity of the specific agent or composition employed; the specific agents or composition employed; the age, body weight, general health, gender and diet of the patient; the time of administration, route of administration, and rate of excretion of the agent; the duration of the treatment; drugs used in combination or coincidental with

the specific agent; and like factors well known in the medical arts. It is well within the skill of the art to start doses of the agents at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosages until the desired effect is achieved.

[0137] Dosaging can also be administered in a patient-specific manner to provide a predetermined concentration of the agents in the blood, as determined by techniques accepted and routine in the art.

[0138] In general, the compounds disclosed herein may be used alone or in concert with other therapeutic agents at appropriate dosages defined by routine testing in order to obtain optimal efficacy while minimizing any potential toxicity. The dosage regimen utilizing a compound of the present invention may be selected in accordance with a variety of factors including type, species, age, weight, sex, medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound employed. A physician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the condition.

[0139] Optimal precision in achieving concentrations of drug within the range that yields maximum efficacy with minimal toxicity may require a regimen based on the kinetics of the compound's availability to one or more target sites. Distribution, equilibrium, and elimination of a drug may be considered when determining the optimal concentration for a treatment regimen. The dosages of a compound disclosed herein may be adjusted when combined to achieve desired effects. On the other hand, dosages of these various therapeutic agents may be independently optimized and combined to achieve a synergistic result wherein the pathology is reduced more than it would be if either agent were used alone.

[0140] In particular, toxicity and therapeutic efficacy of a compound disclosed herein may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effect is the therapeutic index and it may be expressed as the ratio LD₅₀/ED₅₀. Compounds exhibiting large therapeutic indices are preferred except when cytotoxicity of the compound is the activity or therapeutic outcome that is desired. Although compounds that exhibit toxic side effects may be used, a delivery system can target such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects. Generally, the compounds of the present invention may be administered in a manner that maximizes efficacy and minimizes toxicity.

[0141] Data obtained from cell culture assays and animal studies may be used in formulating a range of dosages for use in humans. The dosages of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods of the present invention, the therapeutically effective dose may be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information may be used to accurately determine useful doses in humans. Levels in plasma may be measured, by high performance liquid chromatography.

[0142] Moreover, the dosage administration of the pharmaceutical compositions of the present invention may be optimized using a pharmacokinetic/pharmacodynamic modeling system. One or more dosage regimens may be chosen and a pharmacokinetic/pharmacodynamic model may be used to determine the pharmacokinetic/pharmacodynamic profile of one or more dosage regimens. Next, one of the dosage regimens for administration may be selected which achieves the desired pharmacokinetic/pharmacodynamic response based on the particular pharmacokinetic/pharmacodynamic profile. See U.S. Patent No. 6,747,002, which is entirely expressly incorporated herein by reference.

[0143] Methods are known in the art for determining effective doses for therapeutic and prophylactic purposes for the disclosed pharmaceutical compositions or the disclosed drug combinations, whether or not formulated in the same composition. For therapeutic purposes, the term “jointly effective amount,” as used herein, means that amount of each active compound or pharmaceutical agent, alone or in combination, that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated. For prophylactic purposes (i.e., inhibiting the onset or progression of a disorder), the term “jointly effective amount” refers to that amount of each active compound or pharmaceutical agent, alone or in combination, that inhibits in a subject the onset or progression of a disorder as being sought by a researcher, veterinarian, medical doctor or other clinician. Thus, the present invention further provides combinations of two or more therapeutic agents wherein, (a) each therapeutic agent is administered in an independently therapeutically or prophylactically effective amount; (b) at least one therapeutic agent in the combination is administered in an amount that is sub-therapeutic or subprophylactic if administered alone, but is therapeutic or prophylactic when administered in combination with the second or additional therapeutic agents according to the invention; or (c) both

therapeutic agents are administered in an amount that is subtherapeutic or subprophylactic if administered alone, but are therapeutic or prophylactic when administered together. Combinations of three or more therapeutic agents are analogously possible. Methods of combination therapy include co-administration of a single formulation containing all active agents; essentially contemporaneous administration of more than one formulation; and administration of two or more active agents separately formulated.

[0144] More specifically, the pharmaceutical compositions may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily. Doses maybe administered for one week, one month, or over the course of several months, 3, 6, 9 or 12 months, or intervals known in the art and determined to be clinically relevant. Doses may be continued throughout the life of the patient, or discontinues when clinical judgment warrants. The daily dosage of the compositions may be varied over a wide range from about 0.0001 to about 1,000 mg per patient, per day. The range may more particularly be from about 0.001 mg/kg to 10 mg/kg of body weight per day, about 0.1-100 mg, about 1.0-50 mg or about 1.0-20 mg per day for adults (at about 60 kg). Additionally, the dosages may be about 0.5-10 mg/kg per day, about 1.0-5.0 mg/kg per day, 5.0-10 mg/kg per day, or equivalent doses as determine by a practitioner, to achieve a serum concentration that is clinically relevant.

[0145] In the case of injections, it is usually convenient to give by an intravenous route in an amount of about 0.01-30 mg, about 0.1-20 mg or about 0.1-10 mg per day to adults (at about 60 kg). Intravenous doses may include a bolus or a slow dosing. In the case of other animals, the dose calculated for 60 kg may be administered as well.

[0146] As a non-limiting example, treatment of humans or animals can be provided as a one-time or periodic dosage of a compound of the present invention 0.0001 to about 1,000 mg per patient, per day. The range may more particularly be from about 0.001 mg/kg to 10 mg/kg of body weight per day, about 0.1-100 mg, about 1.0-50 mg or about 1.0-20 mg per day for adults (at about 60 kg). Additionally, the dosages may be about 0.5-10 mg/kg per day, about 1.0-5.0 mg/kg per day, 5.0-10 mg/kg per day, or equivalent doses as determine by a practitioner, to achieve a serum concentration that is clinically relevant.

[0147] Specifically, the pharmaceutical compositions of the present invention may be administered at least once a week over the course of several weeks. In one embodiment, the pharmaceutical compositions are administered at least once a week over several weeks to several months. In another embodiment, the pharmaceutical compositions are administered once a week over four to eight weeks. In yet another embodiment, the pharmaceutical compositions are administered once a week over four weeks.

[0148] The invention further provides a method for treating or preventing a 12-lipoxygenase mediated disease or disorder. The method comprises administering to a mammal a therapeutically or prophylactically effective amount of a compound of the present invention.

[0149] The 12-lipoxygenase mediated disease or disorder to be treated or prevented is typically a disease or disorder wherein the production of 12-hydroperoxyeicosatetraenoic acid (12(S)-HPETE) and/or 12-hydroxyeicosatetraenoic acid (12(S)-HETE) is implicated in the development or progression of the disease or disorder. 12-LOX mediated diseases and disorders includes those where 12-LOX is a direct mediator of the diseases and disorders, as well as those where the inhibition of 12-LOX results in therapeutic value in the treatment or prevention of the diseases and disorders.

[0150] In one embodiment, the invention provides a method of treating or preventing a 12-lipoxygenase mediated disease or disorder, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of any of compounds of Formula (I) – (VII), or a salt, enantiomers, a mixture of enantiomers, or diastereomer thereof.

[0151] In one embodiment, the 12-lipoxygenase is human the 12-lipoxygenase.

[0152] In one embodiment, the 12-lipoxygenase mediated disease or disorder is selected from the group consisting of type 1 diabetes, type 2 diabetes, diabetic kidney disease, diabetic nerve disease, cardiovascular disease, Alzheimer's disease, Non-Alcoholic steatohepatitis, platelet hemostasis, skin diseases, heparin induced thrombocytopenia, thrombosis, and cancer.

[0153] In one embodiment, the cancer is selected from the group consisting of prostate cancer, colorectal cancer, breast cancer, and lung cancer.

[0154] In one embodiment, the cardiovascular disease is selected from the group consisting of congestive heart failure, myocardial infarction and stroke.

[0155] In one embodiment, the invention provides a method of treating or preventing type 1 and/or type 2, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of any of compounds of Formula (I) – (VII), or a salt, enantiomers, a mixture of enantiomers, or diastereomer thereof.

[0156] In one embodiment, the invention provides a method of treating or preventing thrombosis, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of any of compounds of Formula (I) - (VII), a salt, enantiomers, a mixture of enantiomers, or diastereomer thereof.

[0157] In one embodiment, the invention provides a method for reducing PAR4-AP induced platelet aggregation, comprising administering to a mammal thereof a therapeutically or

prophylactically effective amount of any of compounds of Formula (I) – (VII), a salt, enantiomers, a mixture of enantiomers, or diastereomer thereof.

[0158] In one embodiment, the invention provides a method for reducing PAR4-AP induced calcium mobilization, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of any of compounds of Formula (I) – (VII), a salt, enantiomers, a mixture of enantiomers, or diastereomer thereof.

[0159] In one embodiment, 12-LOX inhibitors have utility in transplantation/xenotransplantation scenarios, for example, where islets are treated ex vivo to improve survival prior to transplant.

[0160] The invention is based, in part, on the surprising discovery that platelet 12(S)-lipoxygenase (12-LOX), a highly expressed oxylipin producing enzyme in the human platelet, is an essential component of Fc γ RIIa-mediated platelet activation. For example, **Fig. 11** shows how 12-LOX modulates Fc γ RIIa-mediated platelet aggregation. Without wishing to be bound by theory, platelets are essential in maintaining hemostasis following inflammation or injury to the vasculature. Activation of the Fc γ RIIa receptor leads to immune-mediated platelet activation, which can result in thrombotic complications leading to myocardial infarction and stroke. Inhibiting Fc γ RIIa-mediated activation in platelets has been shown to limit thrombosis and is the principle target for prevention of immune-mediated platelet activation. However, 12-LOX was not known in the art to be implicated in the pathogenesis of Fc γ RIIa-mediated thrombosis until the inventors' discovery. Accordingly, the technology disclosed herein aims to exploit this surprising discovery and provide novel methods for treating or preventing a disease or disorder in which a Fc γ RIIa-mediated pathway is involved. Such a disease or disorder can be an immune-mediated thrombocytopenia and thrombosis disorder. In some aspects of the invention, disclosed herein are novel methods for inhibiting platelet activation and preventing or treating thrombosis.

[0161] One aspect of the invention relates to a method of inhibiting or decreasing platelet activation, the method comprising contacting a platelet with a 12-lipoxygenase inhibitor. In some embodiments, the platelet activation is immune-mediated. In some embodiments, the immune-mediated platelet activation is resulting from the activation of a Fc γ RIIa receptor.

[0162] A compound or agent is considered a 12-LOX inhibitor if it decreases the function or activity of 12-LOX by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95%. A 12-LOX inhibitor can be an organic compound, an inorganic compound, a biological compound (e.g., proteins or fragments thereof, antibodies or fragments thereof, nucleic acids, nucleic acid analogs, saccharides, or peptides), or any combination thereof. A 12-LOX inhibitor can also be synthetic or naturally

occurring. Non-limiting examples of 12-LOX inhibitors include ETYA (CAS No.: 1191-85-1), baicalein (CAS No.: 491-67-8), 15(S)-HETrE (CAS No.: 13222-49-6), caffeic acid (CAS No.: 331-39-5), CDC (CAS No.: 132465-11-3), esculetin (CAS No.: 305-01-1), ethyl 3,4-dihydroxybenzylidenecyanoacetate (CAS No.: 132464-92-7), ETI (CAS No.: 13488-22-7), ferulic acid (CAS No.: 1135-24-6), Gossypol (CAS No.: 303-45-7), 2-TEDC (CAS No.: 132465-10-2), Hinokitiol (CAS No.: 499-44-5), 8,11,14-eicosatriynoic acid (CAS No.: 34262-64-1), daphnodorin A, ML355, and a combination thereof. The chemical structure of ML355 is disclosed in Luci et al. (J Med Chem. 2014, 57, 495-506), the content of which is incorporated by reference in its entirety.

[0163] 12-LOX inhibitors are also disclosed, for example, in US20130096159, WO1990012008, US6217875, US4605669, US4623661, CA1327204, US20070111965, US4897422, US20060193797, US5120752, US5112848, US5574062, EP0416609, US4822811, EP0456760, US4761403, the content of each of which is incorporated by reference for its teachings of 12-LOX inhibitors.

[0164] In some embodiments, a 12-LOX inhibitor used for the methods disclosed herein can be a combination of 12-LOX inhibitors. In some of these embodiments, the mixing ratios of the 12-LOX inhibitors are optimized to provide maximum therapeutic effects.

[0165] At least some of the 12-LOX inhibitors (e.g., ML355) have been found to show low toxicity and good clearance from the body (Luci et al., J Med Chem. 2014, 57, 495-506), demonstrating the therapeutic values of the methods disclosed herein.

[0166] In some embodiments, the 12-LOX inhibitor contacts the platelet *in vitro*. For example, the platelets can be obtained from a subject and then cultured in a container (e.g., 96-well plate or petri dish). The *in vitro* contacting can be used for applications such as evaluation of the efficacy of the 12-LOX inhibitor.

[0167] In some embodiments, the 12-LOX inhibitor contacts the platelet *in vivo*. In these embodiments, the *in vivo* contacting can be in a subject taking heparin. The subject might be in need of treating or preventing an immune-mediated thrombocytopenia and thrombosis disorder.

[0168] Platelet activation can be assessed or measured by a variety of methods. For example, platelet activation can be quantified by factors such as change in shape and a tendency to aggregate. Shape change of platelets can be assessed by flow cytometry or electron microscopy. Platelet aggregation can be measured in a platelet aggregometer. Platelet activation can also be quantified by measuring the blood or urine levels of relevant platelet metabolic products or biomarkers (e.g., the alpha granule components, beta thromboglobulin and platelet factor 4, thromboxane B₂, or the soluble form of the adhesion molecule P-selectin). Some methods for measuring platelet activation

are disclosed in US7011938, WO1996012956, and US6391568, the content of each of which is incorporated by reference in its entirety.

[0169] As 12-LOX inhibitors can inhibit or decrease immune-mediated platelet activation, another aspect of the invention relates to a method of treating or preventing an immune-mediated thrombocytopenia and thrombosis disorder, the method comprising administering to a subject in need thereof an effective amount of a 12-lipoxygenase inhibitor. Examples of immune-mediated thrombocytopenia and thrombosis disorders include, but are not limited to, heparin-induced thrombocytopenia (HIT); anti-phospholipid syndrome; sepsis syndrome; thrombosis associated with therapeutic or diagnostic monoclonal antibodies; and thrombotic thrombocytopenic purpura.

[0170] In some embodiments, an immune-mediated thrombocytopenia and thrombosis disorder can be triggered in the subject by the use of one or more heparin or derivatives thereof. In these embodiments, the immune-mediated thrombocytopenia and thrombosis disorder is heparin-induced thrombocytopenia (HIT).

[0171] HIT is a disorder resulting from the administration of heparin therapy, and is the development of a low platelet count (e.g., <100,000 platelets per milliliter in the blood). As used herein, the terms “HIT” and “heparin-induced thrombocytopenia and thrombosis (HITT)” are used interchangeably. Paradoxically, anticoagulant medication and thrombocytopenia manifest not as bleeding, but rather as and life-threatening thrombosis. The thrombotic events are multifocal, involving for example, veins, arteries and the microvasculature. In some HIT cases, thrombi form in macrovascular beds with overt clinical symptoms, such as DVT/PE, myocardial infarction, stroke, or limb ischemia. In other cases the thrombi are in the microvasculature, with effects that are overt clinically (e.g., adrenal thrombosis followed by hemorrhage, skin necrosis) or remain subclinical.

[0172] It was found that HIT occurred in 1% to 3% of patients who received unfractionated heparin, and in 0.1% of patients who received low-molecular-weight heparin. Currently there is no standard diagnostic method for HIT. Some diagnostic methods for HIT are disclosed, for example, in WO2001004159 and WO2013165969, the content of each of which is incorporated by reference in its entirety. Clinical scoring systems have also been developed to predict the probability of HIT onset in a variety of situations (Elalamy I, et al., Ann. Med. 2000, 32, 60-67; Samama M, et al., Bull Acad Natl Med. 1998, 182, 1517-1533). For example, in one clinical scoring system, scores are given to factors such as platelet count evolution, development of thrombosis, and other causes of thrombocytopenia.

[0173] For example, a therapeutically effective amount can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model is also

used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in other subjects. Generally, the therapeutically effective amount is dependent of the desired therapeutic effect.

[0174] Dosage regimens of a 12-LOX inhibitor can be adjusted to provide the optimum desired response (e.g. a therapeutic or prophylactic response). For example, a single bolus can be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It can be advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage.

[0175] A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of a 12-LOX inhibitor at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

[0176] A physician or veterinarian having ordinary skill in the art can also readily determine when to initiate administering a pharmaceutical composition comprising a 12-LOX inhibitor. It should be noted that the timing for administration generally depends on diagnostic results. For example, the pharmaceutical composition comprising a 12-LOX inhibitor can be administered when one or more symptoms of HIT have occurred (e.g., a drop in platelet count).

[0177] The method disclosed herein can be used in combination with other methods or therapies for treating an immune-mediated thrombocytopenia and thrombosis disorder (e.g., HIT). For example, one therapeutic approach to treating HIT is the removal of heparin treatment and replacement with direct thrombin inhibitors (e.g., danaparoid, argatroban, or lepirudin). It should be noted that simple withdrawal of heparin does not end the immune-mediated thrombocytopenia and thrombosis disorder. In another example, a Factor Xa inhibitor (e.g., fondaparinux) can be used for HIT treatment. Other therapies include, but are not limited to, thrombolytic therapy (e.g., streptokinase or urokinase), thromboembolectomy, plasmapheresis, high-dose IV IgG, and GPIIb/IIIa inhibitors.

[0178] Yet another aspect of the invention relates to a method of treating or preventing a thrombotic event, myocardial infarction, or stroke, the method comprising administering to a subject in need thereof an effective amount of a 12-LOX inhibitor. Immune-mediated platelet activation can lead to thrombi formation, which can clot arteries and result in stroke, myocardial infarction, organ infarction, limb gangrene, or other serious complications.

[0179] In some aspects of all the embodiments, the subject is a mammal. In some aspects of all the embodiments, the subject is a human.

[0180] In some aspects of all the embodiments, the subject can be one who exhibits one or more risk factors for an immune-mediated thrombocytopenia and thrombosis disorder. In some aspects of all the embodiments, the subject is receiving heparin therapy. In some aspects of all the embodiments, the subject undergoes orthopedic surgery. It is known in the art that orthopedic surgery patients are at higher risk for developing HIT than are patients who receive heparin for other medical reasons.

[0181] Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The term “comprises” means “includes.” The abbreviation, “e.g.” is derived from the Latin exempli gratia, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.”

[0182] Although preferred embodiments have been depicted and described in detail herein, it will be apparent to those skilled in the relevant art that various modifications, additions, substitutions, and the like can be made without departing from the spirit of the invention and these are therefore considered to be within the scope of the invention as defined in the claims which follow. Further, to the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various embodiments herein described and illustrated can be further modified to incorporate features shown in any of the other embodiments disclosed herein.

[0183] All patents and other publications; including literature references, issued patents, published patent applications, and co-pending patent applications; cited throughout this application are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the technology described herein. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.

[0184] The description of embodiments of the disclosure is not intended to be exhaustive or to limit the disclosure to the precise form disclosed. While specific embodiments of, and examples for, the disclosure are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the disclosure, as those skilled in the relevant art will recognize.

For example, while method steps or functions are presented in a given order, alternative embodiments may perform functions in a different order, or functions may be performed substantially concurrently. The teachings of the disclosure provided herein can be applied to other procedures or methods as appropriate. The various embodiments described herein can be combined to provide further embodiments. Aspects of the disclosure can be modified, if necessary, to employ the compositions, functions and concepts of the above references and application to provide yet further embodiments of the disclosure.

[0185] Specific elements of any of the foregoing embodiments can be combined or substituted for elements in other embodiments. Furthermore, while advantages associated with certain embodiments of the disclosure have been described in the context of these embodiments, other embodiments may also exhibit such advantages, and not all embodiments need necessarily exhibit such advantages to fall within the scope of the disclosure.

EXAMPLES

[0186] The following examples illustrate some embodiments and aspects of the invention. It will be apparent to those skilled in the relevant art that various modifications, additions, substitutions, and the like can be performed without altering the spirit or scope of the invention, and such modifications and variations are encompassed within the scope of the invention as defined in the claims which follow. The following examples do not in any way limit the invention.

[0187] The technology described herein is further illustrated by the following examples which in no way should be construed as being further limiting.

[0188] The description below provides the composition of matter and method of use around a 4-((2-hydroxy-3-methoxybenzyl)amino)benzenesulfonamide chemotype. The novel chemotypes for 12-LO inhibition disclosed herein are soluble, have favorable ADME properties, and good *in vivo* PK properties.

[0189] The synthesis of compound **1** involved a reductive amination with 4-aminobenzenesulfonamide and 2-hydroxy-3-methoxybenzaldehyde. A step-wise approach involving pre-formation of the imine was required with 4-amino-*N*-(thiazol-2-yl)benzenesulfonamide (**2**) and the requisite benzaldehyde overnight at 100 °C. Subsequent reduction of the imine with sodium borohydride provided the desired compounds **1**, **8-10** and **14-33** (Scheme 1, Method A). While this route worked for a majority of the analogs, some analogs (**11-13** & **34**) required an alternative route in which a Buchwald-Hartwig type coupling was utilized with the commercially available 4-bromo-*N*-(thiazol-2-yl)benzenesulfonamide (**3**) and 2-methoxy substituted benzyl amines to provide the desired products in a yields of 25-40% (Scheme 1, Method B). For modifications of the thiazole portion of the molecule we wanted to introduce diversity in the final step of the synthesis (Scheme

1, Method C). Therefore, reaction of commercially available 4-aminobenzenesulfonamide (**4**) with 2-hydroxy-3-methoxybenzaldehyde in ethanol at reflux for 18 h, followed by treatment with sodium borohydride provided compound **5** in 95% yield. The resulting sulfonamide derivative was then subjected to Cu-catalyzed N-arylation conditions using the appropriate heteroaryl bromides to afford compounds **35**, **38**, **44-48**, **52**, **53**, **56-59**, **61-69**, **71**, **74**, and **75** in good yields. (Wang, X. et al. *Tetrahedron Lett.* **2012**, *53*, 7-10.) Despite the general versatility of this method, there was a few isolated cases where the Cu-catalyzed reaction failed to produce the desired product (e.g. compounds **77-83**), or the heteroaryl bromides were not readily available. For these compounds a less direct route was developed by heating 4-nitrobenzene-1-sulfonyl chloride (**6**) and the required heteroaryl amines to 100 °C for 1.5 to 8 hours depending on the reactivity of the amine to give the 4-nitrophenyl-sulfonamide derivatives **7** (Scheme 1, Method D). Reduction of the nitro group was achieved using the H-Cube® Pro flow reactor with 10% Pd on carbon at 50 °C and a pressure of 50 bar. Alternatively, for less soluble compounds which were not amendable to flow chemistry, a Zn/AcOH reduction was performed at 60 °C in methanol. Once the desired aniline was in hand a reductive amination was carried out with the corresponding benzyl amine derivative to provide compounds **36**, **37**, **39-43**, **49-51**, **54**, **55**, **60**, **70**, **72**, **73**, and **76-83**. The specific conditions for each step in Scheme 1 as shown in Figure 1 are as follows: (i) RCHO (1.5 equiv), EtOH, 3-18 h, 100 oC, NaBH₄ (3.0 equiv), 0.5-0.6h; (ii) RCH₂NH₂ (1.2 equiv), Xantphos (0.06 equiv), Pd₂dba₃ (0.02 equiv), NaOtBu (2.5 equiv), 1,4-dioxane, MW, 30 min, 100 oC. (iii) 2-hydroxy-3-methoxybenzaldehyde (1.2 equiv), EtOH, 6 h, 100 oC, NaBH₄ (1.5 equiv), 30 min, 92%. (iv) R'Br (1.2 equiv), K₂CO₃ (2.5 equiv), 80 oC, 6-8 h. (v) R'NH₂, pyridine, 100 oC, 1.5-18 h. (vi) 10% Pd/C, MeOH/EtOAc, 50 bar, 50 oC or zinc (4.0 equiv), AcOH (4.0 equiv), methanol, 60 oC, 30 min-2 h. (vii) 2-hydroxy-R-benzaldehyde (1.2 equiv), EtOH, 18 h, 100 oC, NaBH₄ (3.0 equiv).

[0190] General Synthetic Procedures. (Scheme 1, Method A): 4-amino-*N*-(thiazol-2-yl)benzene sulfonamide (0.39 mmol) (**2**), and required benzaldehyde (0.67 mmol) were combined in ethanol (2 mL) in a sealed tube and heated to 100 °C for 4-18 h. The solid was allowed to cool to room temperature and sodium borohydride (0.80 mmol) was added and stirred for 30 min, during which time the reaction turned clear and then cloudy. The resulting solids were filtered, washed with ethanol, and purified using a prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% TFA in water w/ 0.1% TFA) to give the desired product.

[0191] (Scheme 1, Method B) A solution of 4-bromo-*N*-(thiazol-2-yl)benzene sulfonamide (0.31 mmol) (**3**) in dioxane (1 mL) was added to a mixture of sodium tert-butoxide (0.78 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos) (0.02 mmol) and tris(dibenzylideneacetone)dipalladium(0) (Pd₂dba₃) (6.27 µmol) in dioxane (1 mL). The resulting

mixture was degassed with argon for 15 min. then the requisite benzylamine (0.38 mmol) was added, the vessel was sealed and heated to 100 °C for 30 min in Biotage® microwave reactor. The reaction mixture was cooled to room temperature and filtered through celite. Silicycle® silica bound thiol was added and stirred overnight, again filtered through a pad of celite, concentrated, and purified by prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% TFA in water w/ 0.1% TFA) to give the desired product.

[0192] (Scheme 1, Method C) 4-aminobenzenesulfonamide (**4**) (1.00 g, 5.81 mmol), 2-hydroxy-3-methoxybenzaldehyde (1.00 g, 7.00 mmol) in EtOH (29 mL) was heated to reflux for 4 h until reaction is an orange turbid mixture. The reaction mixture was cooled to room temperature before sodium borohydride (0.33 g, 8.71 mmol) was added and stirred for an additional 30 min. A white solid forms after 30 min and is collected by filtration and washed with copious amounts of ethanol, dried under vacuum and used as is in subsequent reactions. ^1H NMR (400 MHz, DMSO- d_6) δ 7.60–7.27 (m, 2H), 6.75–6.40 (m, 4H), 6.06 (t, J = 7.63 Hz, 1H), 4.18 (s, 2H), and 3.65 (s, 3H); ^{13}C NMR (101 MHz, DMSO) δ 40.37, 55.32, 108.91, 109.42, 109.55, 111.29, 111.40, 121.05, 125.05, 127.49, 129.92, 150.17, 152.36, and 156.94; LC-MS retention time (Method 1): 2.876 min. General procedure: (**Step iv**) 4-(2-hydroxy-3-methoxybenzylamino)benzenesulfonamide (**5**) (0.58 mmol), arylbromide (0.70 mmol), K_2CO_3 (1.45 mmol), N,N'-dimethylethylenediamine (0.29 mmol), and copper(I)iodide (0.03 mmol) in 1,4-dioxane (1.5 mL) were place under N_2 and sealed in a 5 mL sealed tube. The reaction was heated to 70 °C for 6 to 8 h and monitored by LC/MS analysis. Upon completion the heterogeneous mixture was cooled to room temperature, filtered, and washed with dioxane. The solution was passed through a thiol cartridge (metal scavenging), diluted with AcOEt and washed with NH_4Cl (2X), water, and brine. The crude material was purified using a prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% TFA in water w/ 0.1% TFA) to give the desired product.

[0193] (Scheme 1, Method D: *Example*) **N-(benzo[d]thiazol-2-yl)-4-nitrobenzenesulfonamide (7):** (Step V) To a stirred solution of benzo[d]thiazol-2-amine (0.50 g, 3.35 mmol) in pyridine (1.60 mL, 20.08 mmol) was added 4-nitrobenzene sulfonyl chloride (**6**) (0.82 g, 3.68 mmol) in three equal parts. The reaction mixture was heated for 75 min at 100 °C, cooled to room temperature, after which time a yellow precipitate formed. The reaction mixture was allowed to sit at room temperature for 2 h then the yellow solid was collected by filtration, washed with ethanol, and dried under reduced pressure overnight to give 1.10 of the desired product. ^1H NMR (400 MHz, DMSO- d_6) δ 8.90 (ddt, J = 0.75, 1.63, and 5.59 Hz, 1H), 8.54, 8.41–8.28 (m, 1H), 8.13–7.96 (m, 2H), 7.82 (dq, J = 0.80, and 7.96 Hz, 1H), 7.45–7.34 (m, 1H), 7.31–

7.19 (m, 1H); LC-MS retention time (Method 1): 3.272 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₃H₁₀N₃O₄S₂, 336.0107) found 336.0107.


[0194] 4-amino-N-(benzo[d]thiazol-2-yl)benzenesulfonamide: (Step VI) N-(benzo[d]thiazol-2-yl)-4-nitrobenzenesulfonamide (0.20 g, 0.60 mmol), zinc (0.16 g, 2.39 mmol), acetic acid (0.14 mL, 2.39 mmol) was dissolved in MeOH (3 mL), and the mixture heated to 60 °C for 2 h. The heterogeneous mixture was filtered through a pad of celite washed with hot methanol, concentrated and purified using prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% NH₄OH in water w/ 0.1% NH₄OH) to give the desired product. Alternate nitro reduction: *N*-(benzo[d]thiazol-2-yl)-4-nitrobenzenesulfonamide was dissolved in MeOH/EtOAc/THF (1:1:1) to give a 0.05 M solution passed through the H-Cube Pro[®] flow reactor using a 10% Pd/C, 70 mm CatCart[®] at 50 bar and 50 °C for two cycles at 0.9 mL/min. The solution was concentrated to give a pale yellow solid in a quantitative yield. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.49–7.34 (m, 3H), 7.18 (ddd, *J* = 0.56, 1.21, and 8.00 Hz, 1H), 7.03 (ddd, *J* = 1.33, 7.24, and 7.97 Hz, 1H), 6.83 (ddd, *J* = 1.20, 7.25, and 7.68 Hz, 1H), 6.51–6.38 (m, 2H), 5.42 (s, 2H); LC-MS retention time (Method 2): 3.933 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₃H₁₂N₃O₂S₂, 306.0365) found 306.0360.

[0195] *N*-(benzo[d]thiazol-2-yl)-4-((2-hydroxy-3-methoxybenzyl)amino)benzenesulfonamide (Step VII, representative example) (35, ML355): 4-amino-N-(benzo[d]thiazol-2-yl)benzenesulfonamide (0.10 g, 0.33 mmol), 2-hydroxy-3-methoxybenzaldehyde (0.075 g, 0.491 mmol) were heated in EtOH (1.5 mL) at 90 °C for 18 h. The reaction mixture was allowed to cool to room temperature before the addition of NaBH₄ (0.04 g, 0.98 mmol), and stirred for an additional 6 h. After this time, the reaction mixture was quenched with methanol and water then stirred for 20 min, the solids were filtered through celite, the filtrate collected, and concentrated under reduced pressure to provide a yellow solid. The crude material was purified using a prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% TFA in water w/ 0.1% TFA) to give the desired product. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.86 (s, 1H), 8.73 (d, *J* = 0.5 Hz, 1H), 7.75 (ddd, *J* = 0.6, 1.2, and 7.9 Hz, 1H), 7.54–7.46 (m, 2H), 7.40–7.31 (m, 1H), 7.28–7.16 (m, 2H), 6.93–6.79 (m, 2H), 6.78–6.55 (m, 4H), 4.23 (d, *J* = 5.8 Hz, 2H) and 3.78 (s, 3H); ¹³C NMR (DMSO-*d*₆) δ ppm 152.4, 147.7, 144.3, 128.2, 125.7, 122.9, 120.4, 119.0, 111.4, 110.9, 56.2 and 40.6; LC-MS retention time (Method 1): 2.260 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₂₁H₁₉N₃O₄S₂, 441.0817) found 441.0819.

[0196] General Methods for Chemistry. All air or moisture sensitive reactions were performed under positive pressure of nitrogen with oven-dried glassware. Chemical reagents and anhydrous solvents were obtained from commercial sources and used as-is. Preparative purification was performed on a Waters semi-preparative HPLC. The column used was a


Phenomenex Luna C18 (5 micron, 30 x 75 mm) at a flow rate of 45 mL/min. The mobile phase consisted of acetonitrile and water (each containing 0.1% trifluoroacetic acid). A gradient of 10% to 50% acetonitrile over 8 minutes was used during the purification. Fraction collection was triggered by UV detection (220 nm). Analytical analysis for purity was determined by two different methods denoted as Final QC Methods 1 and 2. Method 1: Analysis was performed on an Agilent 1290 Infinity Series HPLC. UHPLC Long Gradient Equivalent 4% to 100% acetonitrile (0.05% trifluoroacetic acid) in water over 3.5 minutes run time of 4 minutes with a flow rate of 0.8 mL/min. A Phenomenex Kinetex 1.7 micron C18 column (2.1 x 100 mm) was used at a temperature of 50 °C. Method 2: analysis was performed on an Agilent 1260 with a 7 minute gradient of 4% to 100% acetonitrile (containing 0.025% trifluoroacetic acid) in water (containing 0.05% trifluoroacetic acid) over 8 minute run time at a flow rate of 1 mL/min. A Phenomenex Luna C18 column (3 micron, 3 x 75 mm) was used at a temperature of 50 °C. Purity determination was performed using an Agilent Diode Array Detector for both Method 1 and Method 2. Mass determination was performed using an Agilent 6130 mass spectrometer with electrospray ionization in the positive mode. All of the analogs for assay have purity greater than 95% based on both analytical methods. ¹H and ¹³C NMR spectra were recorded on a Varian 400 (100) MHz spectrometer. High resolution mass spectrometry was recorded on Agilent 6210 Time-of-Flight LC/MS system.

[0197] Description of Embodiments of the Present Invention:




[0198] 4-(2-hydroxy-3-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (1):


Method A: using 2-hydroxy-3-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.67 (s, 1H), 7.44–7.33 (m, 2H), 7.00 (d, *J* = 4.20 Hz, 1H), 6.85–6.62 (m, 3H), 6.60–6.45 (m, 4H), 4.19 (d, *J* = 5.91 Hz, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 2): 4.394 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₄S₂ 392.0733) found, 392.0726.




[0199] 4-(benzylamino)-N-(thiazol-2-yl)benzene sulfonamide TFA (8): Method A: using benzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 12.39 (s, 1H), 7.48–7.39 (m, 2H), 7.35–7.13 (m, 6H), 6.99 (t, J = 6.00 Hz, 1H), 6.71 (d, J = 4.60 Hz, 1H), 6.62–6.54 (m, 2H), and 4.29 (d, J = 5.61 Hz, 2H); LC-MS retention time (Method 2): 4.780 min; HRMS: m/z (M+H) $^+$ = (Calculated for C₁₆H₁₆N₃O₂S₂ 346.0678) found, 346.0672.



[0200] 4-(2-hydroxybenzylamino)-N-(thiazol-2-yl)benzene sulfonamide (9): Method A: using 2-hydroxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 9.49 (s, 1H), 7.44–7.35 (m, 2H), 7.13 (dd, J = 1.65, and 7.55 Hz, 1H), 7.03 (ddd, J = 1.75, 7.32, and 8.01 Hz, 1H), 6.92–6.67 (m, 4H), 6.52–6.41 (m, 2H), 6.38–6.29 (m, 2H), and 4.17 (d, J = 5.68 Hz, 2H); LC-MS retention time (Method 2): 4.213 min; HRMS: m/z (M+H) $^+$ = (Calculated for C₁₆H₁₆N₃O₃S₂ 362.0628) found, 362.0620.




[0201] 4-(3-methoxybenzylamino)-N-(thiazol-2-yl)benzene sulfonamide (10): Method B: using (3-dimethoxyphenyl)methanamine; ^1H NMR (400 MHz, DMSO- d_6) δ 7.48–7.41 (m, 2H), 7.27–7.19 (m, 1H), 7.18 (d, J = 4.63 Hz, 1H), 6.99 (t, J = 6.06 Hz, 1H), 6.93–6.85 (m, 2H), 6.79 (dd, J = 1.10, and 2.48 Hz, 1H), 6.73 (d, J = 4.60 Hz, 1H), 6.63–6.55 (m, 2H), 4.27 (d, J = 5.92 Hz, 2H), and 3.72 (s, 3H); LC-MS retention time (Method 2): 4.777 min; HRMS: m/z (M+Na) $^+$ = (Calculated for C₁₇H₁₇N₃NaO₃S₂ 398.0604) found, 398.0584.




[0202] 4-(2-methoxybenzylamino)-N-(thiazol-2-yl)benzene sulfonamide TFA (11): Method B: using 2-dimethoxyphenyl)methanamine; ^1H NMR (400 MHz, DMSO- d_6) δ 12.39 (s, 1H), 7.48–7.39 (m, 2H), 7.26–7.12 (m, 3H), 6.98 (dd, J = 1.03, and 8.26 Hz, 1H), 6.90–6.76 (m, 2H), 6.71 (d, J = 4.63 Hz, 1H), 6.59–6.50 (m, 2H), 4.23 (d, J = 5.89 Hz, 2H), and 3.81 (s, 3H); LC-MS retention


time (Method 2): 4.888 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₃S₂ 376.0784) found, 376.0765.




[0203] 4-(2,3-dimethoxybenzylamino)-N-(thiazol-2-yl)benzene sulfonamide (12): Method B: using 2,3-dimethoxyphenyl)methanamine; ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.46 (d, *J* = 8.78 Hz, 2H), 7.18 (d, *J* = 4.63 Hz, 1H), 7.06–6.90 (m, 2H), 6.87–6.79 (m, 2H), 6.73 (d, *J* = 4.58 Hz, 1H), 6.59 (d, *J* = 8.84 Hz, 2H), 4.28 (d, *J* = 5.90 Hz, 2H), and 3.78 (d, *J* = 13.99 Hz, 6H); LC-MS retention time (Method 2): 4.756 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₈H₂₀N₃O₄S₂ 406.0890) found, 406.0885.



[0204] 4-(2-aminobenzylamino)-N-(thiazol-2-yl)benzene sulfonamide (13): Method A: using tert-butyl (2-formylphenyl)carbamate and removing the subsequent carbamate with 4 M HCl in dioxane over 30 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.54–7.37 (m, 3H), 7.24–7.06 (m, 3H), 6.96 (d, *J* = 7.86 Hz, 1H), 6.86 (h, *J* = 7.06 Hz, 1H), 6.74 (d, *J* = 4.61 Hz, 1H), 6.65–6.50 (m, 3H), and 4.23 (s, 2H); LC-MS retention time (Method 1): 1.863 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₆H₁₇N₄O₂S₂ 361.0787) found, 361.0784.




[0205] 4-(3-hydroxybenzylamino)-N-(thiazol-2-yl)benzene sulfonamide (14): Method A: using 3-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.31 (s, 1H), 7.51–7.37 (m, 2H), 7.18 (d, *J* = 4.62 Hz, 1H), 7.10 (t, *J* = 7.77 Hz, 1H), 6.97 (t, *J* = 5.90 Hz, 1H), 6.79–6.68 (m, 3H), 6.66–6.49 (m, 3H), and 4.31–4.14 (m, 2H); LC-MS retention time (Method 1): 1.775 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₆H₁₆N₃O₃S₂ 362.0628) found, 362.0614.




[0206] 4-(2-amino-3-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide TFA (15):

Method A: using 3-methoxy-2-nitrobenzaldehyde; A heterogeneous solution of 4-(3-methoxy-2-nitrobenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (0.075 g, 0.178 mmol) in MeOH (1.8 mL), AcOH (0.102 mL, 1.784 mmol) and zinc (0.023 g, 0.357 mmol) were stirred for 30 min, filtered through celite, and washed with MeOH. The filtrate was concentrated and purified using a prep-HPLC (gradient 10-100% acetonitrile w/ 0.1% TFA in water w/ 0.1% TFA) to give the desired product; ^1H NMR (400 MHz, DMSO- d_6) δ 7.44 (d, J = 8.78 Hz, 2H), 7.17 (d, J = 4.61 Hz, 1H), 6.78 (dd, J = 1.23, and 7.81 Hz, 1H), 6.75–6.67 (m, 3H), 6.56 (d, J = 8.82 Hz, 2H), 4.16 (s, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 1): 2.775 min; HRMS: m/z (M+H) $^+$ = (Calculated for $\text{C}_{17}\text{H}_{19}\text{N}_4\text{O}_3\text{S}_2$ 391.0893) found, 391.0874.




[0207] 4-(3-hydroxy-4-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (16):


Method A: using 3-hydroxy-4-methoxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 8.86 (s, 1H), 7.52–7.32 (m, 2H), 7.15 (d, J = 4.61 Hz, 1H), 6.95–6.74 (m, 2H), 6.77–6.62 (m, 4H), 6.62–6.49 (m, 2H), 4.13 (d, J = 5.92 Hz, 2H), and 3.70 (s, 3H); LC-MS retention time (Method 2): 4.123 min; HRMS: m/z (M+H) $^+$ = (Calculated for $\text{C}_{17}\text{H}_{18}\text{N}_3\text{O}_4\text{S}_2$ 392.0733) found, 392.0719.



[0208] 4-((1H-indol-7-yl)methylamino)-N-(thiazol-2-yl)benzenesulfonamide (17): Method A: using 1H-indole-7-carbaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 7.51–7.38 (m, 4H), 7.36–7.31 (m, 1H), 7.16 (t, J = 4.53 Hz, 1H), 6.95–6.85 (m, 2H), 6.70 (d, J = 4.62 Hz, 1H), 6.67–6.57 (m, 2H), 6.44 (dd, J = 1.83, and 3.08 Hz, 1H), and 4.53 (d, J = 5.67 Hz, 2H); LC-MS retention time (Method 2): 4.899 min; HRMS: m/z (M+H) $^+$ = (Calculated for $\text{C}_{18}\text{H}_{17}\text{N}_4\text{O}_2\text{S}_2$ 385.0787) found, 385.0774.




[0209] 4-(2,3-dichlorobenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (18): Method A: using 2,3-dichlorobenzaldehyde; LC-MS retention time (Method 2): 5.786 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₄Cl₂N₃O₂S₂ 413.9899) found, 413.9907.




[0210] 4-(3-chloro-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (19):

Method A: using 3-chloro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.48–7.35 (m, 2H), 7.21 (dd, *J* = 1.63, and 7.94 Hz, 1H), 7.13–7.02 (m, 1H), 6.87 (d, *J* = 3.82 Hz, 1H), 6.76 (t, *J* = 7.77 Hz, 1H), 6.51–6.43 (m, 2H), 6.35 (d, *J* = 3.82 Hz, 1H), and 4.25 (s, 2H); LC-MS retention time (Method 1): 2.076 min; HRMS: *m/z* (M+Na)⁺ = (Calculated for C₁₆H₁₄Cl N₃NaO₃S₂ 419.0085) found, 419.0047.



[0211] 4-(3-fluoro-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (20):


Method A: using 3-fluoro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.63 (s, 1H), 7.49–7.33 (m, 2H), 7.08–6.93 (m, 2H), 6.91–6.80 (m, 1H), 6.72 (t, *J* = 7.94 Hz, 1H), 6.51–6.38 (m, 3H), 6.34 (d, *J* = 3.80 Hz, 1H), and 4.23 (d, *J* = 3.52 Hz, 2H); LC-MS retention time (Method 1): 2.076 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₅FN₃O₃S₂ 380.0533) found, 380.0521.



[0212] 4-(3-bromo-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide, (21):


Method A: using 3-bromo-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.64 (s, 1H), 8.16–8.02 (m, 1H), 7.86 (d, *J* = 8.75 Hz, 2H), 7.79 (dd, *J* = 1.59, and 7.90 Hz, 1H), 7.58 (d, *J* = 4.61 Hz, 1H), 7.52 (dd, *J* = 1.56, and 7.59 Hz, 1H), 7.28 (s, 1H), 7.14 (t, *J* = 7.78 Hz, 1H), 6.96 (d,

J = 8.89 Hz, 2H), and 4.69 (d, *J* = 5.79 Hz, 2H); LC-MS retention time (Method 2): 4.777 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₅BrN₃O₃S₂ 441.9712) found, 441.9705.




[0213] 4-(4-bromo-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (22):

Method A: using 4-bromo-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.06 (s, 1H), 7.47–7.34 (m, 2H), 7.03 (d, *J* = 8.13 Hz, 1H), 6.99–6.92 (m, 2H), 6.88 (dd, *J* = 1.97, and 8.10 Hz, 1H), 6.65–6.31 (m, 4H), and 4.12 (d, *J* = 5.91 Hz, 2H); LC-MS retention time (Method 1): 2.938 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₅BrN₃O₃S₂ 441.9712) found, 441.9704.




[0214] 4-(2-hydroxy-3-methylbenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (23):

Method A: using 2-hydroxy-3-methylbenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.25 (s, 1H), 7.34–7.21 (m, 2H), 7.00 (d, *J* = 4.62 Hz, 1H), 6.84–6.75 (m, 2H), 6.64–6.48 (m, 3H), 6.44–6.31 (m, 2H), 4.08 (d, *J* = 5.76 Hz, 2H), and 2.01 (s, 3H); LC-MS retention time (Method 1): 2.083 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₃S₂ 376.0784) found, 376.0779.




[0215] 4-(3-amino-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (24):

Method A: using 2-hydroxy-3-nitrobenzaldehyde and reducing the nitro-group with zinc and acetic acid conditions (see compound 15 for details). ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.49–7.39 (m, 2H), 7.17 (d, *J* = 4.65 Hz, 1H), 6.92 (t, *J* = 8.63 Hz, 2H), 6.85–6.69 (m, 3H), 6.60–6.50 (m, 2H), and 4.28 (s, 2H); LC-MS retention time (Method 1): 2.263 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₇N₄O₃S₂ 377.0737) found, 377.0730.




[0216] 4-(2-hydroxy-3-nitrobenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (25):

Method 1A: using 2-hydroxy-3-nitrobenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.88 (dd, J = 1.65, and 8.43 Hz, 1H), 7.58–7.24 (m, 3H), 7.17 (d, J = 4.66 Hz, 1H), 7.05–6.88 (m, 2H), 6.72 (d, J = 4.61 Hz, 1H), 6.67–6.34 (m, 2H), and 4.33 (d, J = 4.46 Hz, 2H); LC-MS retention time (Method 2): 4.742 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ = (Calculated for $\text{C}_{16}\text{H}_{15}\text{N}_4\text{O}_5\text{S}_2$ 407.0478) found, 407.0465.




[0217] 4-(3-allyl-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide TFA (26):

Method A: using 3-allyl-2-hydroxybenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.41 (s, 1H), 8.47 (s, 1H), 7.49–7.38 (m, 2H), 7.17 (d, J = 4.61 Hz, 1H), 6.96 (ddd, J = 1.73, 7.52, and 22.52 Hz, 2H), 6.80–6.65 (m, 3H), 6.61–6.50 (m, 2H), 5.93 (ddt, J = 6.63, 10.04, and 16.79 Hz, 1H), 5.08–4.95 (m, 2H), 4.25 (s, 2H), and 3.34 (dt, J = 1.44, and 6.58 Hz, 2H); LC-MS retention time (Method 1): 3.155 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ = (Calculated for $\text{C}_{19}\text{H}_{20}\text{N}_3\text{O}_3\text{S}_2$ 402.0941) found, 402.0926.



[0218] 4-(4-chloro-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (27):


Method A: using 4-chloro-2-hydroxybenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.38 (d, J = 8.6 Hz, 2 H), 7.07 (d, J = 8.2 Hz, 1 H), 6.92–6.61 (m, 3 H), 6.49–6.20 (m, 4 H), and 4.12 (s, 2 H); LC-MS retention time (Method 2): 4.700 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ = (Calculated for $\text{C}_{16}\text{H}_{15}\text{ClN}_3\text{O}_3\text{S}_2$ 396.0238) found, 396.0220.



[0219] 4-(2-hydroxy-4-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (28):


Method A: using 2-hydroxy-4-methoxybenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.51 (s, 1H), 7.41–7.32 (m, 2H), 7.00 (d, J = 8.37 Hz, 1H), 6.87–6.80 (m, 1H), 6.49–6.39 (m, 2H), 6.36 (d, J = 2.47 Hz, 1H), 6.34–6.26 (m, 2H), 6.21 (t, J = 5.86 Hz, 1H), 4.07 (d, J = 5.77 Hz, 2H), and 3.63

(s, 3H); LC-MS retention time (Method 1): 3.155 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₄S₂ 392.0733) found, 392.0715.




[0220] 4-(5-chloro-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (29):

Method A: using 5-chloro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.00 (s, 1H), 7.48–7.37 (m, 2H), 7.13–7.00 (m, 3H), 6.89–6.76 (m, 2H), 6.64 (d, *J* = 4.44 Hz, 1H), 6.58–6.48 (m, 2H), and 4.17 (d, *J* = 5.97 Hz, 2H); LC-MS retention time (Method 2): 4.610 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₆H₁₅ClN₃O₃S₂ 396.0238) found, 396.0233.




[0221] 4-(2-hydroxy-5-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (30):

Method A: using 2-hydroxy-5-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.12 (s, 1H), 7.47–7.38 (m, 2H), 7.08 (d, *J* = 4.44 Hz, 1H), 6.77–6.65 (m, 3H), 6.65–6.48 (m, 4H), 4.15 (d, *J* = 5.91 Hz, 2H), and 3.56 (s, 3H); LC-MS retention time (Method 2): 4.137 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₄S₂ 392.0733) found, 392.0725.




[0222] 4-(5-amino-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (31):

Method A: using 2-hydroxy-5-nitrobenzaldehyde and reducing the nitro-group with zinc and acetic acid conditions (see compound 15 for details). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.92 (s, 1H), 7.46 (d, *J* = 8.83 Hz, 2H), 7.18 (d, *J* = 4.64 Hz, 1H), 7.04–6.92 (m, 2H), 6.86 (d, *J* = 8.23 Hz, 1H), 6.73 (d, *J* = 4.66 Hz, 1H), 6.53 (d, *J* = 8.83 Hz, 2H), and 4.22 (d, *J* = 5.77 Hz, 2H); LC-MS retention time (Method 1): 2.165 min; HRMS: m/z (M+H)⁺ = (Calculated for C₁₆H₁₇N₄O₃S₂ 377.0737) found, 377.0719.




[0223] 4-(5-fluoro-2-hydroxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (32):

Method A: using 5-fluoro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.63–9.54 (m, 1H), 7.44–7.33 (m, 2H), 6.92–6.73 (m, 4H), 6.51–6.36 (m, 3H), 6.36–6.29 (m, 1H), and 4.14 (d, *J* = 6.00 Hz, 2H); LC-MS retention time (Method 1): 3.017 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₅FN₃O₃S₂ 380.0533) found, 380.0526.




[0224] 4-(2-hydroxy-5-nitrobenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (33):

Method A: using 2-hydroxy-5-nitrobenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.92–7.76 (m, 2 H), 7.41 (d, *J* = 9.0 Hz, 2 H), 6.97 (d, *J* = 4.3 Hz, 1 H), 6.56–6.40 (m, 4 H), 5.74 (s, 1 H), and 4.10 (s, 2 H); LC-MS retention time (Method 2): 4.273 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₆H₁₅N₄O₅S₂ 407.0478) found, 407.0470.




[0225] 4-(2-hydroxy-6-methoxybenzylamino)-N-(thiazol-2-yl)benzenesulfonamide (34):

Method B: using 2-hydroxy-6-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.59 (s, 1H), 7.74–7.54 (m, 1H), 7.42 (d, *J* = 8.79 Hz, 2H), 7.24–7.05 (m, 1H), 6.68 (d, *J* = 8.87 Hz, 2H), 6.46 (ddd, *J* = 0.93, 7.08, and 8.22 Hz, 2H), 6.22 (d, *J* = 5.51 Hz, 1H), 4.12 (d, *J* = 5.21 Hz, 2H), and 3.74 (s, 3H); LC-MS retention time (Method 1): 2.960 min; HRMS: *m/z* (M+H)⁺ = (Calculated for C₁₇H₁₈N₃O₄S₂ 392.0733) found, 392.0716.




[0226] *N*-(benzo[d]oxazol-2-yl)-4-((2-hydroxy-3-


methoxybenzyl)amino)benzenesulfonamide (36): Method C: using 2-bromobenzoxazazole; ^1H NMR (600 MHz, DMSO- d_6) δ 12.40 (s, 1H), 8.71 (s, 1H), 7.55 (dd, J = 2.60, and 9.29 Hz, 2H), 7.43 (d, J = 7.93 Hz, 1H), 7.31–7.25 (m, 1H), 7.26–7.20 (m, 1H), 7.16 (td, J = 1.42, and 7.82 Hz, 1H), 6.86 (t, J = 5.64 Hz, 1H), 6.81 (dd, J = 1.72, and 7.88 Hz, 1H), 6.75–6.69 (m, 1H), 6.66 (t, J = 7.85 Hz, 1H), 6.61–6.54 (m, 2H), 4.21 (d, J = 5.58 Hz, 2H), and 3.76 (d, J = 2.79 Hz, 3H); ^{13}C NMR (151 MHz, DMSO- d_6) δ 156.01, 152.37, 147.74, 144.28, 128.27, 125.73, 125.45, 123.67, 120.40, 119.06, 112.19, 111.31, 110.88, 110.59, 56.23, and 41.15; LC-MS retention time (Method 2): 4.848 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_5\text{S}$ 426.1118) found, 426.1098.

[0227] *N*-(1H-benzo[d]imidazol-2-yl)-4-((2-hydroxy-3-


methoxybenzyl)amino)benzenesulfonamide (37): Method D: using 2-aminobenzimidazole and 2-hydroxy-3-methoxybenzaldehyde; ^1H NMR (600 MHz, DMSO- d_6) δ 11.68 (s, 2H), 7.56–7.49 (m, 2H), 7.22 (dt, J = 3.37, and 5.83 Hz, 2H), 7.06 (dt, J = 3.34, and 5.73 Hz, 2H), 6.79 (dd, J = 1.71, and 7.99 Hz, 1H), 6.71 (dd, J = 1.59, and 7.73 Hz, 1H), 6.65 (t, J = 7.83 Hz, 1H), 6.58–6.50 (m, 2H), 4.19 (s, 2H), and 3.75 (s, 3H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 151.47, 150.22, 147.72, 144.26, 130.61, 129.89, 127.61, 125.79, 122.59, 120.44, 119.05, 111.46, 111.09, 110.86, 56.23, and 41.32; LC-MS retention time (Method 2): 4.454min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}_4\text{S}$, 424.1205) found 424.1202.

[0228] **4-(2-hydroxy-3-methoxybenzylamino)-*N*-(thiophen-2-yl)benzenesulfonamide (38):**


Method C: using 2-bromothiophene; ^1H NMR (400 MHz, DMSO- d_6) δ 8.73 (s, 1H), 7.38–7.28 (m, 1H), 7.05–6.98 (m, 1H), 6.91 (t, J = 6.1 Hz, 1H), 6.83 (dd, J = 2.0, and 7.8 Hz, 1H), 6.78–6.64 (m, 3H), 6.61–6.52 (m, 2H), 6.51–6.43 (m, 1H), 4.25 (s, 2H), and 3.80 (s, 3H); LC-MS retention time (Method 1): 2.435 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}_4\text{S}_2$, 391.0781) found 391.0773.




[0229] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(4-methylbenzo[d]thiazol-2-yl)benzenesulfonamide TFA (39): Method D: using 2-amino-4-methylbenzthiazole and 2-hydroxy-3-methoxybenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 12.76 (s, 1H), 8.71 (s, 1H), 7.58–7.45 (m, 3H), 7.18–7.05 (m, 2H), 6.81 (dd, J = 1.73, and 7.89 Hz, 1H), 6.76–6.54 (m, 4H), 4.21 (s, 2H), 3.76 (s, 3H), and 2.34 (s, 3H); LC-MS retention time (Method 2): 5.522 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{22}\text{H}_{22}\text{N}_3\text{O}_4\text{S}_2$, 456.1046) found 456.1037.



[0230] 4-(2-hydroxy-3-methoxybenzylamino)-N-(4-methylthiazol-2-yl)benzenesulfonamide TFA (40): Method C: using 2-bromo-4-methylthiazole; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 12.31 (s, 1 H), 8.70 (s, 1 H), 7.48–7.32 (m, 3 H), 6.89–6.72 (m, 6 H), 6.61–6.47 (m, 3 H), 6.27 (s, 1 H), 4.20 (d, J = 5.90 Hz, 2 H), 3.76 (s, 3 H), and 1.95 (s, 3 H); LC-MS retention time (Method 1): 1.962 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_4\text{S}_2$, 406.0890) found 406.0875.




[0231] 4-(2-hydroxy-3-methoxybenzylamino)-N-(5-methylthiazol-2-yl)benzenesulfonamide TFA (41): Method D: using 2-amino-5-methylthiazole and 2-hydroxy-3-methoxybenzaldehyde; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 12.05 (s, 1H), 8.73 (s, 1H), 7.46–7.38 (m, 2H), 6.91–6.63 (m, 5H), 6.59–6.51 (m, 2H), 4.20 (d, J = 5.86 Hz, 2H), 3.77 (s, 3H), and 2.13 (d, J = 1.39 Hz, 3H); LC-MS retention time (Method 1): 3.182 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_4\text{S}_2$, 406.0890) found 406.0889.




[0232] **4-((2-hydroxy-3-methoxybenzyl)amino)-N-(5-phenylthiazol-2-yl)benzenesulfonamide TFA (42):**


Method D: using 2-amino-5-phenylthiazole and 2-hydroxy-3-methoxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 12.59 (s, 1H), 8.70 (s, 1H), 7.68 (s, 1H), 7.56–7.25 (m, 8H), 6.87–6.78 (m, 2H), 6.79–6.62 (m, 2H), 6.61–6.53 (m, 2H), 4.21 (d, J = 5.8 Hz, 2H), and 3.76 (s, 3H); LC-MS retention time (Method 2): 5.417 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{23}\text{H}_{22}\text{N}_3\text{O}_4\text{S}_2$, 468.1046) found 468.1028.

[0233] ***N*-(4,5-dimethylthiazol-2-yl)-4-(2-hydroxy-3-**

methoxybenzylamino)benzenesulfonamide TFA (43): Method C: using 2-bromo-4,5-dimethylthiazole; ^1H NMR (400 MHz, DMSO- d_6) δ 12.25 (s, 1H), 8.94 (d, J = 0.71 Hz, 1H), 7.73–7.45 (m, 2H), 7.13–6.80 (m, 4H), 6.83–6.62 (m, 2H), 4.53–4.25 (m, 2H), 3.98 (d, J = 0.60 Hz, 3H), ad 2.35–1.95 (m, 6H); LC-MS retention time (Method 2): 4.859 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{19}\text{H}_{21}\text{N}_3\text{O}_4\text{S}_2$, 420.1046) found 420.1037.


[0234] **4-(2-hydroxy-3-methoxybenzylamino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide TFA (44):**

Method D: using 3-amino-5-methylisoxazole and 2-hydroxy-3-methoxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 10.91 (d, J = 1.25 Hz, 1H), 8.76 (d, J = 3.84 Hz, 1H), 7.77–7.55 (m, 1H), 7.52–7.28 (m, 2H), 7.02 (t, J = 5.41 Hz, 1H), 6.77–6.64 (m, 2H), 6.63–6.51 (m, 2H), 6.11–6.02 (m, 1H), 4.27–4.14 (m, 2H), 3.77 (d, J = 1.18 Hz, 3H), and 2.26 (s, 3H); LC-MS retention time (Method 2): 4.804 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_5\text{S}$, 390.1118) found 390.1109.


[0235] **4-((2-hydroxy-3-methoxybenzyl)amino)-N-(3-methoxyphenyl)benzenesulfonamide (45):**

Method C: using 1-bromo-3-methoxybenzene; ^1H NMR (400 MHz, DMSO- d_6) δ 9.87 (s, 1H), 8.73 (d, J = 0.5 Hz, 1H), 7.48–7.36 (m, 2H), 7.13–7.03 (m, 1H), 6.95–6.80 (m, 2H), 6.77–6.47 (m,


7H), 4.20 (d, J = 5.8 Hz, 2H), 3.78 (s, 3H), and 3.64 (s, 3H); LC-MS retention time (Method 2): 5.437 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₁H₂₃N₂O₅S, 415.1322) found 415.1302.




[0236] 4-(2-hydroxy-3-methoxybenzylamino)-N-(quinolin-3-yl)benzenesulfonamide TFA (46): Method C: using 3-bromoquinoline; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.99 (d, J = 1.27 Hz, 1H), 8.20 (dd, J = 0.78, and 2.71 Hz, 1H), 7.60–7.33 (m, 3H), 7.17 (dd, J = 1.31, 6.90, 8.19, and 31.70 Hz, 2H), 7.09–7.04 (m, 2H), 6.41 (dd, J = 1.94, 7.69 Hz, 1H), 6.30–6.20 (m, 2H), 6.17–6.10 (m, 2H), 3.76 (s, 2H), and 3.36 (s, 3H); LC-MS retention time (Method 1): 2.304 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₃H₂₂N₃O₄S, 436.1326) found 436.1316.



[0237] 4-(2-hydroxy-3-methoxybenzylamino)-N-(isoquinolin-8-yl)benzenesulfonamide (47): Method C: using 8-bromoisoquinoline; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.34–10.24 (s, 1H), 9.56 (s, 1H), 8.74 (s, 1H), 8.54 (d, J = 5.95 Hz, 1H), 8.02 (d, J = 5.96 Hz, 1H), 7.91–7.73 (m, 2H), 7.44–7.29 (m, 3H), 6.85 (t, J = 4.80 Hz, 1H), 6.76–6.62 (m, 2H), 6.61–6.44 (m, 2H), 4.19 (s, 2H), and 3.79 (s, 3H); LC-MS retention time (Method 1): 1.914 min; HRMS: m/z (M+Na)⁺ (Calculated for C₂₃H₂₁NaN₃O₄S, 458.1145) found 458.1129.




[0238] 4-(2-hydroxy-3-methoxybenzylamino)-N-phenylbenzenesulfonamide TFA (48): Method C: using bromobenzene; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.85 (s, 1H), 8.73 (d, J = 2.23 Hz, 1H), 7.42–7.34 (m, 2H), 7.21–7.12 (m, 2H), 7.06–6.99 (m, 2H), 6.98–6.91 (m, 1H), 6.82 (dd, J = 1.86, and 7.72 Hz, 1H), 6.74–6.64 (m, 2H), 6.57–6.49 (m, 2H), 4.31–3.99 (m, 2H), and 3.76 (s, 3H); LC-MS retention time (Method 1): 2.750 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₀H₂₁N₂O₄S, 385.1217) found 385.1223.



[0239] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(naphthalen-1-yl)benzenesulfonamide


(49): Method D: using naphthalen-1-amine and 2-hydroxy-3-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.74 (s, 1H), 8.73 (s, 1H), 8.06 (ddd, *J* = 0.74, 1.48, and 8.45 Hz, 1H), 7.89–7.82 (m, 1H), 7.72 (d, *J* = 8.15 Hz, 1H), 7.54–7.27 (m, 5H), 7.15 (dd, *J* = 1.05, and 7.45 Hz, 1H), 6.85 (q, *J* = 4.87 Hz, 2H), 6.75–6.66 (m, 2H), 6.57–6.47 (m, 2H), 4.19 (d, *J* = 5.65 Hz, 2H), and 3.79 (s, 3H); LC-MS retention time (Method 1): 2.744 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₄H₂₃N₂O₄S, 435.1373) found 435.1392.



[0240] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(naphthalen-2-yl)benzenesulfonamide


(50):

[0241] Method D: using naphthalen-2-amine and 2-hydroxy-3-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.11 (d, *J* = 1.4 Hz, 1H), 8.71 (s, 1H), 7.81–7.67 (m, 3H), 7.54–7.30 (m, 5H), 7.27 (dt, *J* = 1.9, and 8.8 Hz, 1H), 6.94–6.76 (m, 2H), 6.76–6.61 (m, 2H), 6.53 (dd, *J* = 8.8, 1.7 Hz, 2H), 4.16 (s, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 1): 3.281 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₄H₂₃N₂O₄S, 435.1373) found 435.1373.




[0242] N-([1,1'-biphenyl]-4-yl)-4-((2-hydroxy-3-

methoxybenzyl)amino)benzenesulfonamide (51): Method D: using biphenyl-4-amine and 2-hydroxy-3-methoxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.99 (s, 1H), 8.72 (s, 1H), 7.60–7.35 (m, 8H), 7.34–7.24 (m, 1H), 7.18–7.09 (m, 2H), 6.90 (t, *J* = 5.9, Hz, 1H), 6.82 (dd, *J* = 1.7, and 7.8 Hz, 1H), 6.76–6.62 (m, 2H), 6.61–6.52 (m, 2H), 4.19 (d, *J* = 5.8 Hz, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 2): 6.433 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₆H₂₅N₂O₄S, 461.1530) found 461.1529.




[0243] *N*-([1,1'-biphenyl]-3-yl)-4-((2-hydroxy-3-

methoxybenzyl)amino)benzenesulfonamide (52): Method C: using 3-bromo-1,1'-biphenyl; ^1H NMR (400 MHz, DMSO- d_6) δ 9.98 (s, 1H), 8.72 (s, 1H), 7.55–7.19 (m, 11H), 7.06 (ddd, J = 1.5, 2.2, and 7.6 Hz, 1H), 6.96–6.73 (m, 2H), 6.75–6.49 (m, 4H), 4.20 (d, J = 5.8 Hz, 2H), and 3.78 (s, 3H); LC-MS retention time (Method 2): 6.131 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{26}\text{H}_{25}\text{N}_2\text{O}_4\text{S}$, 461.1530) found 461.1521.



[0244] 4-((2-hydroxy-3-methoxybenzyl)amino)-*N*-(3-(piperazin-1-


yl)phenyl)benzenesulfonamide (53): Method 1C: using tert-butyl 4-(3-bromophenyl)piperazine-1-carboxylate and the Boc group was removed after the reductive amination with 4 M HCl in dioxanes over a 30 min period. ^1H NMR (400 MHz, DMSO- d_6) δ 9.80 (s, 1H), 8.75 (s, 1H), 7.47–7.38 (m, 2H), 7.05 (t, J = 8.12 Hz, 1H), 6.95–6.81 (m, 2H), 6.77–6.65 (m, 3H), 6.64–6.51 (m, 4H), 4.20 (d, J = 5.58 Hz, 2H), 3.79 (s, 3H), 3.21 (s, 8H), and 0.88–0.76 (m, 1H); LC-MS retention time (Method 1): 1.660 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{24}\text{H}_{29}\text{N}_4\text{O}_4\text{S}$, 469.1904) found 469.1897.




[0245] 4-(2-hydroxy-3-methoxybenzylamino)-*N*-(4-(piperazin-1-

yl)phenyl)benzenesulfonamide (54): Method D: using tert-butyl 4-(4-aminophenyl)piperidine-1-carboxylate and 2-hydroxy-3-methoxybenzaldehyde. The Boc group was removed after the reductive amination with 4 M HCl/dioxanes for 1 h at rt. ^1H NMR (400 MHz, DMSO- d_6) δ 9.50 (s, 1H), 8.74 (d, J = 0.47 Hz, 1H), 7.39–7.30 (m, 2H), 6.98–6.79 (m, 6H), 6.78–6.65 (m, 2H), 6.59–6.50 (m, 2H), 4.20 (d, J = 5.63 Hz, 2H), 3.79 (s, 3H), and 3.19 (s, 8H); LC-MS retention time


(Method 1): 1.648 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₄H₂₉N₄O₄S, 469.1904) found 469.1900.



[0246] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(4-(piperidin-4-yl)phenyl)benzenesulfonamide (55): Method D: using tert-butyl 4-(4-aminophenyl)piperidine-1-carboxylate and 2-hydroxy-3-methoxybenzaldehyde. The Boc group was removed after the reductive amination with 4 M HCl in dioxane for 1 h at rt. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.03 (s, 1H), 8.92 (d, *J* = 0.5 Hz, 1H), 7.63–7.56 (m, 2H), 7.32–7.15 (m, 6H), 6.95–6.82 (m, 2H), 6.78–6.68 (m, 2H), 4.37 (d, *J* = 5.7 Hz, 2H), 3.96 (s, 3H), 3.57–3.42 (m, 4H), 3.14–3.01 (m, 4H), and 2.90 (d, *J* = 11.4 Hz, 1H); LC-MS retention time (Method 1): 1.710 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₅H₃₀N₃O₄S, 468.1952) found 468.1935.




[0247] 4-(2-hydroxy-3-methoxybenzylamino)-N-(6-(piperazin-1-yl)pyridin-3-yl)benzenesulfonamide (56): Method C: using tert-butyl 4-(5-bromopyridin-2-yl)piperazine-1-carboxylate, and the Boc group was removed after the reductive amination with 4 M HCl/dioxanes over a 1 h. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.57 (s, 1H), 9.12 (d, *J* = 7.21 Hz, 2H), 7.74 (d, *J* = 2.65 Hz, 1H), 7.31 (dd, *J* = 4.87, and 7.36 Hz, 2H), 7.07 (s, 1H), 6.89–6.80 (m, 2H), 6.76–6.59 (m, 2H), 6.60–6.51 (m, 2H), 4.19 (s, 2H), 3.77 (s, 3H), 3.75–3.53 (m, 4H), and 3.17–3.07 (m, 4H); LC-MS retention time (Method 1): 1.993 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₃H₂₈N₅O₄S, 470.1857) found 470.1848.




[0248] 4-(2-hydroxy-3-methoxybenzylamino)-N-(6-methylpyridin-3-yl)benzenesulfonamide (57): Method C: using 5-bromo-2-methylpyridine; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.10 (s, 1H), 8.61 (s, 1H), 8.02 (dd, *J* = 0.68, 2.63 Hz, 1H), 7.47–7.39 (m, 1H), 7.32–7.23 (m, 2H), 7.18 (d, *J* = 8.47 Hz, 1H), 6.70 (dd, *J* = 1.92, and 7.68 Hz, 1H), 6.63–6.50 (m, 2H),

6.48–6.40 (m, 2H), 4.06 (s, 2H), 3.64 (s, 3H), and 2.28 (s, 3H); LC-MS retention time (Method 1): 1.840 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₀H₂₂N₃O₄S, 400.1326) found 400.1315.




[0249] 4-(2-hydroxy-3-methoxybenzylamino)-N-(pyridin-2-yl)benzenesulfonamide (58):

Method C: using 2-bromopyridine; ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.71 (d, *J* = 0.54 Hz, 1H), 8.11–7.96 (m, 1H), 7.65–7.58 (m, 1H), 7.52 (d, *J* = 8.70 Hz, 2H), 7.09–6.98 (m, 1H), 6.92–6.79 (m, 3H), 6.76–6.63 (m, 2H), 6.58–6.50 (m, 2H), 4.19 (d, *J* = 5.83 Hz, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 2): 4.540 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₁₉H₂₀N₃O₄S, 386.1169) found 386.1158.




[0250] 4-(2-hydroxy-3-methoxybenzylamino)-N-(pyridin-3-yl)benzenesulfonamide (59):

Method C: using 3-bromopyridine; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.30 (s, 1H), 8.73 (s, 1H), 8.31–8.21 (m, 2H), 7.57 (ddd, *J* = 1.38, 2.63, and 8.37 Hz, 1H), 7.46–7.32 (m, 3H), 6.83 (dd, *J* = 1.92, and 7.71 Hz, 1H), 6.75–6.62 (m, 2H), 6.61–6.51 (m, 2H), 4.19 (s, 2H), 3.77 (s, 3H); LC-MS retention time (Method 1): 1.804 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₁₉H₂₀N₃O₄S, 386.1169) found 386.1166.




[0251] 4-(2-hydroxy-3-methoxybenzylamino)-N-(pyrimidin-2-yl)benzenesulfonamide TFA (60): Method C: using 2-bromopyrimidine; ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.25 (s, 1H), 8.77 (s, 1H), 8.48 (d, *J* = 4.83 Hz, 2H), 7.72–7.50 (m, 2H), 6.99 (dt, *J* = 5.43, and 15.84 Hz, 2H), 6.85 (dd, *J* = 1.69, and 7.85 Hz, 1H), 6.77–6.66 (m, 2H), 6.62–6.56 (m, 2H), 4.23 (d, *J* = 5.65 Hz, 2H), and 3.79 (s, 3H); LC-MS retention time (Method 2): 4.266 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₁₈H₁₉N₄O₄S, 387.1122) found 387.1115.




[0252] *N*-(3-(tert-butyl)phenyl)-4-((2-hydroxy-3-

methoxybenzyl)amino)benzenesulfonamide, TFA (61): Method C: using 1-bromo-3-(tert-butyl)benzene; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.71 (s, 1H), 8.71 (s, 1H), 7.42–7.32 (m, 2H), 7.14–6.93 (m, 3H), 6.93–6.78 (m, 3H), 6.73–6.60 (m, 2H), 6.58–6.47 (m, 2H), 4.18 (s, 2H), 3.76 (s, 3H), and 1.14 (s, 9H); LC-MS retention time (Method 2): 6.184 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{24}\text{H}_{29}\text{N}_2\text{O}_4\text{S}$, 441.0843) found 441.1844.




[0253] *4*-((2-hydroxy-3-methoxybenzyl)amino)-*N*-(6-methoxybenzo[d]thiazol-2-

yl)benzenesulfonamide, TFA (62): Method C: using 2-bromo-6-methoxybenzo[d]thiazole; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.70 (s, 1H), 8.71 (d, $J = 0.47$ Hz, 1H), 7.51–7.41 (m, 2H), 7.38 (d, $J = 2.54$ Hz, 1H), 7.14 (d, $J = 8.76$ Hz, 1H), 6.96–6.78 (m, 3H), 6.76–6.62 (m, 2H), 6.61–6.53 (m, 2H), 4.21 (d, $J = 5.89$ Hz, 2H), 3.75 (s, 3H), and 3.73 (s, 3H); LC-MS retention time (Method 2): 5.278 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{22}\text{H}_{22}\text{N}_3\text{O}_5\text{S}_2$, 472.0995) found 472.0998.




[0254] *4*-((2-hydroxy-3-methoxybenzyl)amino)-*N*-(4-phenylthiazol-2-

yl)benzenesulfonamide, TFA (63): Method C: using 2-bromo-4-phenylthiazole; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.72 (s, 1H), 7.67 (d, $J = 7.47$ Hz, 2H), 7.53–7.24 (m, 6H), 7.08 (d, $J = 19.97$ Hz, 1H), 6.92–6.45 (m, 6H), 4.21 (d, $J = 5.55$ Hz, 2H), and 3.77 (s, 3H); LC-MS retention time (Method 2): 5.200 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{23}\text{H}_{22}\text{N}_3\text{O}_4\text{S}$, 468.1046) found 468.1066.




[0255] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(3-

morpholinophenyl)benzenesulfonamide, TFA (64): Method C: using 4-(3-bromophenyl)morpholine; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.70 (s, 1H), 8.72 (s, 1H), 7.46–7.34 (m, 2H), 7.00 (t, J = 8.09 Hz, 1H), 6.82 (dd, J = 1.86, and 7.75 Hz, 2H), 6.76–6.43 (m, 7H), 4.19 (s, 2H), 3.77 (s, 3H), 3.71–3.59 (m, 4H), and 2.98–2.86 (m, 4H); LC-MS retention time (Method 2): 4.961 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{24}\text{H}_{28}\text{N}_3\text{O}_5\text{S}$, 470.1744) found 470.1753.



[0256] *tert*-butyl4-(3-(4-((2-hydroxy-3-


methoxybenzyl)amino)phenylsulfamido)phenyl)piperidine-1-carboxylate (65): Method C: using *tert*-butyl-4-(3-bromophenyl)piperidine-1-carboxylate; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.71 (s, 1H), 7.46–7.28 (m, 2H), 7.08 (t, J = 7.74 Hz, 1H), 6.94–6.75 (m, 6H), 6.75–6.59 (m, 2H), 6.59–6.44 (m, 2H), 4.18 (d, J = 5.82 Hz, 2H), 4.00 (d, J = 12.83 Hz, 2H), 3.76 (s, 3H), 2.59–2.50 (m, 1H), 1.62 (d, J = 12.87 Hz, 2H), 1.39 (s, 9H), and 1.31 (m, 4H); LC-MS retention time (Method 2): 6.322 min; HRMS: m/z ($\text{M}+\text{H}$) $^+$ (Calculated for $\text{C}_{30}\text{H}_{38}\text{N}_3\text{O}_6\text{S}$, 568.2476) found 568.2461.



[0257] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(3-(piperidin-4-


yl)phenyl)benzenesulfonamide, TFA (66): Method 1C: using *tert*-butyl-4-(3-bromophenyl)piperidine-1-carboxylate the Boc group was removed after the reductive amination with 4 M HCl/dioxanes over a 30 min period. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.73 (s, 1H), 7.53–

7.24 (m, 2H), 7.12 (t, J = 7.85 Hz, 1H), 6.99 (t, J = 1.96 Hz, 1H), 6.93–6.77 (m, 4H), 6.73–6.61 (m, 2H), 6.57–6.48 (m, 2H), 4.33–4.01 (m, 2H), 3.77 (s, 3H), 3.33 (s, 1H), 2.95 (t, J = 12.62 Hz, 2H), 2.83–2.55 (m, 1H), and 1.94–1.43 (m, 4H); LC-MS retention time (Method 2): 4.079 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₅H₃₀N₄O₄S, 468.1952) found 468.1948.




[0258] 4-((2-hydroxy-3-methoxybenzyl)amino)-N-(3-isopropylphenyl)benzenesulfonamide, TFA (67):

Method C: using 1-bromo-3-isopropylbenzene; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.73 (s, 1H), 8.70 (s, 1H), 7.42–7.32 (m, 2H), 7.07 (t, J = 7.77 Hz, 1H), 6.98–6.75 (m, 4H), 6.73–6.61 (m, 2H), 6.58–6.45 (m, 2H), 4.18 (s, 2H), 3.76 (d, J = 0.87 Hz, 3H), 2.72 (h, J = 6.82 Hz, 1H), and 1.07 (dd, J = 0.91, and 6.94 Hz, 6H); LC-MS retention time (Method 2): 6.040 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₃H₂₇N₂O₄S, 427.1686) found 427.1680.



[0259] N-(6-fluorobenzo[d]thiazol-2-yl)-4-((2-hydroxy-3-methoxybenzyl)amino)benzenesulfonamide, TFA (68):


Method C: using 2-bromo-6-fluorobenzo[d]thiazole; ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.72 (s, 1H), 7.69 (dd, J = 2.43, and 8.51 Hz, 1H), 7.54–7.39 (m, 2H), 7.19 (qd, J = 3.40, 8.25, and 8.85 Hz, 3H), 6.88 (t, J = 5.68 Hz, 1H), 6.81 (dd, J = 1.71, and 7.87 Hz, 1H), 6.74–6.63 (m, 2H), 6.60–6.54 (m, 2H), 4.21 (d, J = 5.90 Hz, 2H), and 3.76 (s, 3H); LC-MS retention time (Method 2): 5.164 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₁H₁₉FN₃O₄S₂, 460.0796) found 460.0797.




[0260] 4-(3-chloro-2-hydroxybenzylamino)-N-phenylbenzenesulfonamide, TFA (69):

Method C: using 3-chloro-2-hydroxybenzaldehyde and bromobenzene; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.88 (s, 1H), 9.38 (s, 1H), 7.47–7.34 (m, 2H), 7.28–7.14 (m, 3H), 7.11–7.01 (m, 3H), 7.01–6.88 (m, 2H), 6.79 (t, J = 7.78 Hz, 1H), 6.60–6.50 (m, 2H), and 4.29–4.23 (m, 2H); LC-MS


retention time (Method 2): 5.442 min; HRMS: m/z (M+H)⁺ (Calculated for C₁₉H₁₈ClN₂O₃S, 389.0721) found 389.0702.




[0261] 4-(3-chloro-2-hydroxybenzylamino)-N-(naphthalen-1-yl)benzenesulfonamide, TFA (70): Method D: using naphthalen-1-amine and 3-chloro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.75 (s, 1H), 9.36 (s, 1H), 8.03 (dd, *J* = 1.20, and 8.53 Hz, 1H), 7.90–7.80 (m, 1H), 7.70 (d, *J* = 8.22 Hz, 1H), 7.54–7.28 (m, 3H), 7.23 (dd, *J* = 1.58, and 8.01 Hz, 1H), 7.15 (dd, *J* = 1.08, and 7.40 Hz, 1H), 7.02 (dd, *J* = 1.58, and 7.65 Hz, 1H), 6.90 (t, *J* = 5.82 Hz, 1H), 6.78 (t, *J* = 7.79 Hz, 1H), 6.57–6.42 (m, 2H), and 4.24 (d, *J* = 4.97 Hz, 2H); LC-MS retention time (Method 1): 2.755 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₃H₂₀ClN₂O₃S, 439.0878) found 439.0862.




[0262] N-(benzo[d]thiazol-2-yl)-4-(3-chloro-2-hydroxybenzylamino)benzenesulfonamide, TFA (71): Method C: using 3-chloro-2-hydroxybenzaldehyde and 2-bromobenzthiazole; ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.88 (s, 1H), 9.38 (s, 1H), 7.80–7.69 (m, 1H), 7.59–7.46 (m, 2H), 7.35 (ddd, *J* = 1.24, 7.44, and 8.26 Hz, 1H), 7.31–7.13 (m, 3H), 7.12–7.02 (m, 1H), 6.95 (t, *J* = 5.93 Hz, 1H), 6.78 (t, *J* = 7.79 Hz, 1H), 6.69–6.52 (m, 2H), and 4.28 (d, *J* = 5.72 Hz, 2H); LC-MS retention time (Method 2): 5.258 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₀H₁₇ClN₃O₃S₂, 446.0390) found 446.0379.




[0263] 4-(3-chloro-2-hydroxybenzylamino)-N-(naphthalen-2-yl)benzenesulfonamide, TFA (72): Method D: using naphthalen-2-amine and 3-chloro-2-hydroxybenzaldehyde; LC-MS retention time (Method 1): 2.637 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₃H₂₀ClN₂O₃S, 439.0878) found 439.0867.




[0264] *N*-([1,1'-biphenyl]-4-yl)-4-((3-chloro-2-hydroxybenzyl)amino)benzenesulfonamide, TFA (73): Method D: using biphenyl-4-amine and 3-chloro-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.03 (s, 1H), 9.37 (s, 1H), 7.61–7.36 (m, 9H), 7.35–7.04 (m, 5H), 6.97 (t, *J* = 5.8 Hz, 1H), 6.78 (t, *J* = 7.8 Hz, 1H), 6.62–6.52 (m, 2H), and 4.26 (d, *J* = 5.7 Hz, 2H); LC-MS retention time (Method 2): 6.592 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₅H₂₂ClN₂O₃S, 465.1034) found 465.1021.



[0265] 4-((3-chloro-2-hydroxybenzyl)amino)-N-(isoquinolin-8-yl)benzenesulfonamide, TFA (74): Method C: using 3-chloro-2-hydroxybenzaldehyde and 8-bromoisoquinoline; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.60 (s, 1H), 8.55 (d, *J* = 6.0 Hz, 1H), 8.09 (d, *J* = 6.0 Hz, 1H), 7.99–7.71 (m, 2H), 7.46–7.29 (m, 3H), 7.23 (dd, *J* = 1.6, and 7.9 Hz, 1H), 7.11–6.93 (m, 1H), 6.78 (t, *J* = 7.8 Hz, 1H), 6.64–6.39 (m, 2H), and 4.24 (s, 2H); LC-MS retention time (Method 1): 1.669 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₂H₁₉ClN₃O₃S, 440.0830) found 440.0811.




[0266] 4-((3-chloro-2-hydroxybenzyl)amino)-N-(quinolin-3-yl)benzenesulfonamide, TFA (75): Method C: using 3-chloro-2-hydroxybenzaldehyde and 3-bromoquinoline; LC-MS retention time (Method 1): 2.346 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₂H₁₉ClN₃O₃S, 440.0830) found 440.0824.




[0267] 4-((3-chloro-2-hydroxybenzyl)amino)-N-(4-(piperidin-4-yl)phenyl)benzenesulfonamide, TFA (76): Method D: using *tert*-butyl-4-(4-

aminophenyl)piperidine-1-carboxylate and 3-chloro-2-hydroxybenzaldehyde. The Boc group was removed after the reductive amination with 4 M HCl in dioxane over a 30 min period. ^1H NMR (400 MHz, DMSO- d_6) δ 10.20 (s, 1H), 9.71 (s, 1H), 7.87–7.59 (m, 2H), 7.57 (dd, J = 1.64, and 7.96 Hz, 1H), 7.48–7.15 (m, 6H), 7.21–7.02 (m, 1H), 6.96–6.69 (m, 2H), 4.59 (d, J = 5.77 Hz, 2H), 3.64 (s, 2H), 3.27 (t, J = 12.85 Hz, 3H), 3.10–2.93 (m, 1H), 2.18 (d, J = 13.73 Hz, 2H), and 1.98 (qd, J = 4.00, and 13.16 Hz, 2H); LC-MS retention time (Method 1): 2.884 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{23}\text{H}_{20}\text{ClN}_2\text{O}_3\text{S}$, 472.1456) found 472.1442.




[0268] 4-(4-bromo-2-hydroxybenzylamino)-N-phenylbenzenesulfonamide, TFA (77):

Method D: using 4-bromo-2-hydroxybenzaldehyde and benzylamine; ^1H NMR (400 MHz, DMSO- d_6) δ 10.09 (s, 1H), 9.85 (s, 1H), 7.52–7.28 (m, 2H), 7.28–7.12 (m, 2H), 7.12–6.78 (m, 7H), 6.63–6.41 (m, 2H), 4.12 (d, J = 2.80 Hz, 2H); LC-MS retention time (Method 2): 5.501 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{19}\text{H}_{18}\text{BrN}_2\text{O}_3\text{S}$, 434.0246) found 434.0239.




[0269] 4-(4-bromo-2-hydroxybenzylamino)-N-(naphthalen-1-yl)benzenesulfonamide (78):

Method D: using 1-amino-naphthalene and 4-bromo-2-hydroxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 10.09 (d, J = 1.50 Hz, 1H), 9.74 (s, 1H), 8.03 (dd, J = 1.23, and 8.56 Hz, 1H), 7.87–7.80 (m, 1H), 7.70 (d, J = 8.24 Hz, 1H), 7.45 (ddd, J = 1.23, 6.81, and 8.11 Hz, 1H), 7.41–7.28 (m, 4H), 7.17–7.09 (m, 1H), 7.01–6.94 (m, 2H), 6.92–6.81 (m, 2H), 6.53–6.42 (m, 2H), and 4.12 (s, 2H); LC-MS retention time (Method 2): 6.282 min; HRMS: m/z (M+H) $^+$ (Calculated for $\text{C}_{23}\text{H}_{20}\text{BrN}_2\text{O}_3\text{S}$, 483.0373) found 483.0367.




[0270] N-(benzo[d]thiazol-2-yl)-4-(4-bromo-2-hydroxybenzylamino)benzenesulfonamide, TFA (79): Method D: using 2-aminobenzothiazole and 4-bromo-2-hydroxybenzaldehyde; ^1H NMR (400 MHz, DMSO- d_6) δ 13.15 (s, 1H), 10.38 (s, 1H), 8.07–7.95 (m, 1H), 7.85–7.68 (m, 2H), 7.62


(ddd, $J = 1.22, 7.36$, and 8.30 Hz, 1H), 7.54–7.40 (m, 2H), 7.37–7.04 (m, 5H), 6.93–6.67 (m, 2H), and 4.43 (d, $J = 5.75$ Hz, 2H); LC-MS retention time (Method 1): 2.505 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₀H₁₇BrN₃O₃S₂, 491.9869) found 491.9855.




[0271] 4-(4-bromo-2-hydroxybenzylamino)-N-(quinolin-3-yl)benzenesulfonamide, TFA (80): Method D: using 3-amino-quinoline and 4-bromo-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-d₆) δ 10.41 (d, $J = 2.1$ Hz, 1H), 10.08 (s, 1H), 8.59 (d, $J = 2.6$ Hz, 1H), 7.94–7.83 (m, 3H), 7.61 (tt, $J = 2.43$, and 6.74 Hz, 1H), 7.59–7.41 (m, 3H), 7.01–6.88 (m, 2H), 6.85 (dt, $J = 2.11$, and 8.10 Hz, 1H), 6.53 (dd, $J = 2.22$, and 9.01 Hz, 2H), and 4.10 (s, 2H); LC-MS retention time (Method 1): 2.166 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₂H₁₉BrN₃O₃S, 485.0355) found 485.0331.



[0272] 4-(4-bromo-2-hydroxybenzylamino)-N-(naphthalen-2-yl)benzenesulfonamide, TFA (81): Method D: using 2-amino-naphthalene and 4-bromo-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-d₆) δ 10.09 (s, 1H), 10.07 (s, 1H), 7.79–7.66 (m, 3H), 7.51–7.30 (m, 5H), 7.25 (dd, $J = 2.19$, and 8.81 Hz, 1H), 7.00–6.92 (m, 2H), 6.91–6.81 (m, 2H), 6.61–6.25 (m, 2H), and 4.09 (d, $J = 2.95$ Hz, 2H); LC-MS retention time (Method 1): 2.856 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₃H₂₀BrN₂O₃S, 483.0373) found 483.0372.



[0273] N-([1,1'-biphenyl]-4-yl)-4-((4-bromo-2-hydroxybenzyl)amino)benzenesulfonamide, TFA (82): Method D: using 4-amino-biphenyl and 4-bromo-2-hydroxybenzaldehyde; ¹H NMR (400 MHz, DMSO-d₆) δ 10.03 (s, 1H), 9.37 (s, 1H), 7.61–7.36 (m, 8H), 7.35–7.04 (m, 5H), 6.97 (t, $J = 5.8$ Hz, 1H), 6.78 (t, $J = 7.8$ Hz, 1H), 6.62–6.52 (m, 2H), 4.26 (d, $J = 5.7$ Hz, 2H); LC-MS retention time (Method 2): 6.592 min; HRMS: m/z (M+H)⁺ (Calculated for C₂₅H₂₂BrN₂O₃S, 510.0560) found 510.0549.



[0274] 4-(4-bromo-2-hydroxybenzylamino)-N-(quinolin-8-yl)benzenesulfonamide, TFA (83): Method D: using 8-amino-isoquinoline and 4-bromo-2-hydroxybenzaldehyde; LC-MS retention time (Method 1): 2.524 min; HRMS: *m/z* (M+H)⁺ (Calculated for C₂₂H₁₉BrN₃O₃S, 485.0355) found 485.0345.

[0275] Systematic Structure-Activity Relationship (SAR) explorations as shown in Tables 1-3 were conducted. As shown in Table 1, the removal of the phenolic groups (**8**), 3-OMe group (**9**), or the 2-OH group (**10**) resulted in a complete loss of activity. Protecting the 2-OH moiety as methyl ether (compounds **11-12**) or replacing with an amine (compounds **13** and **15**) also negated all 12-LOX inhibitory activity. A bioisosteric replacement of the catechol moiety with an indole (**17**) was also inactive. The 2,3-dichloro derivative (**18**) had comparable activity to compound **1**. Also, the 3-Cl (**19**, IC₅₀ = 6.1 μM) had comparable activity while the 3-Fluoro and 3-Bromo (**20**, IC₅₀ = 19 μM and **21**, IC₅₀ = 13 μM) was ~4 to ~2.5-fold fold less active, respectively. An interesting and unexpected result was the 2-fold improvement in activity observed for the 4-bromo (**22**, IC₅₀ = 2.2 μM) and 4-chloro (**27**) derivatives. Other 4-substituted derivatives such as 4-methoxy (**28**) had reduced activity compared to compound **1**. The 2-OH was important for activity and the 3-position is most optimal for the methoxy group. This preliminary SAR suggested that the 3-OMe could be replaced with a chloro group (**19**) and the 4-Bromo (**22**) while maintaining comparable, if not improved activity to compound **1**.

[0276] Table 1. 12-LOX Inhibition of Analogue 1, 8-34^a



compd	R	IC ₅₀ [± SD] (μM)
1	2-OH, 3-OMe-Ph	5.1 [0.5]
8	Ph	>40
9	2-OH-Ph	>40
10	3-OMe-Ph	>40
11	2-OMe-Ph	>40
12	2,3-OMe-Ph	>40
13	2-NH ₂ -Ph	>40

14	3-OH-Ph	>40
15	2-NH ₂ , 3-OMe-Ph	>40
16	3-OH, 4-OMe-Ph	>40
17	7-indole	>40
18	2,3-Cl-Ph	>40
19	3-Cl	6.2 [2.0]
20	3-F	19 [4]
21	3-Br	13 [1]
22	4-Br	2.2 [0.5]
23	3-Me	>40
24	3-NH ₂	>40
25	3-N0 ₂	>40
26	3-allyl	>40
27	4-Cl	6.3 [1.0]
28	4-OMe	22 [11]
29	S-Cl	>40
30	S-OMe	>40
31	5-NH ₂	>40
32	5-F	>40
33	5-N0 ₂	>40
34	6-OMe	>40


^aIC₅₀ values represent the half maximal (50%) inhibitory concentration as determined in the UV-vis cuvette-based assay in triplicate.

[0277] The next SAR profile focused on modification of the thiazole group as shown in Table 2. Changes to this region of the molecule led to analogs with improved potency. Replacing the thiazole moiety with a 2-benzothiazole (**35**, **ML355**) resulted in an 18-fold improvement in 12-LOX activity while retaining selectivity. The benzoxazole (**36**) and benzimidazole (**37**) possessed good to excellent activity, and introduction of a methyl group at the 4-position of the benzothiazole ring (**39**), maintained excellent 12-LOX activity (IC₅₀ = 0.24 μM). Potency against 12-LOX was also improved over 10-fold when the thiazole was replaced with a thiophene (**38**), but the substituted thiazole/isoxazole derivatives (**40-43**) did not show this increased potency. This was also the case with the phenyl derivative (**48**), a known thiophene bioisostere, which had potent activity against 12-LOX (IC₅₀ = 0.5 μM). Generally, larger aromatic [1-naphthalene (**49**) and 2-naphthalene (**50**)] and heteroaromatic compounds [3-quinoline (**46**), 8-isoquinoline (**47**), 2-pyridine (**58**) and 3-pyridine (**59**)] were well tolerated and had better potency than the thiazole derivative (**1**). In an effort to improve solubility by adding solubilizing functionality, a few phenyl derivatives substituted with a piperazine moiety at different positions around the aryl ring were synthesized (**53-55**). While these changes were tolerated, they had reduced activity compared to the top actives (e.g. entry **35**, **ML355**).

[0278] Many of these analogs were also tested for activity against 15-LOX-1 to ensure that selectivity was maintained. These compounds were initially tested against 15-LOX-1 at a single

concentration (25 μ M) and an IC₅₀ was determined only on compounds of interest. These studies showed that replacement of the thiazole with a benzothiazole, and its derivatives maintained selectivity with 15/12-LOX ratios of 29-fold (**35**), 18- (**39**), 19- (**62**) and 20- (**68**). The 15-LOX/12-LOX selectivity ratio improved to over 100 with conversion of the benzothiazole to a benzoxazole (**36**), benzimidazole (**37**) and *m*-iPr substituted phenyl (**67**). The phenyl substitution (**48**) only had a selectivity ratio of 15. A wide-range of selectivity was observed despite the compounds bearing comparable structures, ranging from almost complete inhibition of 15-LOX-1 (e.g. **38** and **62**) to minimal inhibition (e.g. **55**, **58**, and **66**). The dramatic effect on selectivity between 12-LOX and 15-LOX-1 in this portion of the molecule, suggests that the bottom of the LOX active site is involved in binding, since it has been shown to control substrate specificity previously.

[0279] **Table 2. 12-LOX Inhibition of Analogue 35–68^a**



compd	R	12 - LOX	15 - LOX - 1	% inh ^b
		IC50 [+ SD] (μ M)	IC50 [+ SD] (μ M)	
35	2-benzothiazole	0.34 [0.04]	9.7 [0.1]	
36	2-benzoxazole	0.79 [0.1]	>100	
37	2-benzimidazole	0.57 [0.04]	>70	
38	2-thiophene	0.35 [0.02]		100
39	4-Me-2-benzothiazole	0.24 [0.01]	0.69 [0.1)	
40	4-Me-2-thiazole	2.7 [0.2]		14
41	5-Me-2-thiazole	3.0 [0.4]		75
42	5-Ph-2-thiazole	91%b		77
43	4,5-Me-2-thiazole	1.4 [0.3]		37
44	5-Me-3-isoxazole	11 [1.2]		
45	3-OMe-Ph	85%b		73
46	3-quinoline	0.48 [0.1]		77
47	8-isoquinoline	0.70 [0.2]		70
48	Ph	0.50 [0.05]	7.6 [1.0]	
49	1-naphthalene	0.51 [0.06]		73
50	2-naphthalene	0.33 [0.05]		54
51	1,4-bi-Ph	1.3 [0.2]		60
52	1,3-bi-Ph	82%b		70
53	3-piperazine-Ph	3.5 [0.5]		31
54	4-piperazine-Ph	11 [2.7]		
55	4-piperidine-Ph	3.7 [0.6]		8
56	4-piperazine-3-pyr	4.0 [0.6]		12


57	6-methyl-3-pyr	5.0 [0.5]	18
58	2-pyr	5.0 [0.5]	3
59	3-pyr	7.0 [0.5]	
60	2-pyrimidine	12 [1.0]	
61	3- ^t Bu-Ph	0.39 [0.8]	66
62	6-OMe-2-benzothiazole	0.26 [0.3]	5.1 [0.6]
63	4-Ph-2-thiazole	0.18 [0.03]	87
64	3-morpholine-Ph	3.8 [0.3]	50
65	4N-boc-piperidine-3-Ph	0.76 [0.05]	34
66	3-piperidine-Ph	1.1 [0.3]	4.7
67	3- <i>i</i> Pr-Ph	0.16 [0.02]	>100
68	6-F-2-benzothiazole	0.22 [0.05)	4.5 [0.4)

^aIC₅₀ values represent the half maximal (50%) inhibitory concentration as determined in the UV-vis cuvette-based assay in triplicate.

^bRepresents inhibition at 25 μM .

[0280] As noted above (Table 1), replacement of the “right-hand” portion of the molecule with a 3-chloro-2-phenol (**19**) or 4-bromo-2-phenol (**22**) resulted in comparable potency to **1** with IC_{50} values of 6.2 and 2.2 μ M respectively. These groups were combined with some of the sulfonamide derivatives discovered as part of the initial SAR efforts (see Table 3, compounds **69-83**). None of the compounds had improved potency and in almost all cases, had reduced activity. Generally, the 2-benzothiazole moiety gave the best activity with both the 3-Chloro-2-phenol (**71**; IC_{50} = 1.6 μ M) and 4-bromo-2-phenol (**79**; IC_{50} = 1.7 μ M), although the 1-naphthalene derivative **78** had comparable potency (IC_{50} = 1.3 μ M).

[0281] Table 3. 12-LOX Inhibition of Analogs 69-83^a



compd	R	12 - LOX	15 - LOX - 1	% inh ^b
		IC50 [+ SD] (μ M)	IC50 [+ SD] (μ M)	
69	Ph	4.5 [0.4]		49
70	1-naphthalene	1.6 [0.2]		70
71	2-benzothiazole	1.3 [0.2]	1.3 [0.2]	
72	2-naphthalene	2.3 [0.3]		75
73	4-bi-Ph	4.2 [1.0]		73
74	8-isoquinoline	4.5 [0.8]		63
75	3-quinoline	5.3 [0.7]		58
76	4-piperidine-Ph	6.3 [3.0]		4
77	Ph	2.9 [0.4]		82

78	1-naphthalene	1.3 [0.3]	83
79	2-benzothiazole	1.7 [0.8]	100
80	3-quinoline	2.3 [0.5]	74
81	2-naphthalene	2.2 [0.3]	78
82	4-bi-Ph	2.5 [0.4]	80
83	8-quinoline	5.6 [2.0]	55

^aIC₅₀ values represent the half maximal (50%) inhibitory concentration as determined in the UV-vis cuvette-based assay in triplicate.

^bRepresents inhibition at 25 μ M.

[0282] Further studies focusing on the selectivity of analogs (**35**, **36**, and **37**) against other human LOX isozymes (5-LOX and 15-LOX-2) were conducted. In addition, the compounds were tested against cyclooxygenase-1 (COX-1) and/or COX-2. As shown in Table 4, there was no significant inhibition against any of these related enzymes, with exception to **ML355** which has modest potency (29-fold less active) towards 15-LOX-1. Few compounds reported in the literature have achieved both nM potency towards 12-LOX and selectivity against other isozymes.

Table 4. Selectivity and Redox Activity of Representative Analogues^a

analogue	12-LOX ^b	15-LOX-1 ^b	15-LOX-2 ^b	5-LOX ^b	COX-1/-2 ^c	redox activity ^d
1	5.1	>50	>40	>200	NT	NT
35	0.34	9.7	>100	>100	NI	NI
36	0.79	>100	>100	>100	NI	NI
37	0.57	>100	>100	>35	NI	NT

^aSelectivity profiling of 1, 35, 36, and 37. ^bIC₅₀ values are reported in μ M. ^cCompounds were tested at 15 μ M and none of the compound exhibited inhibition above 10%. ^dUV-vis pseudoperoxidase activity assay was performed on **35** and **36**, and no degradation of the hydroperoxide product was observed at 234 nm, indicating a nonreductive inhibitory mechanism; NI = no inhibition and NT = not tested.

[0283] LOX inhibitors are known to exhibit a variety of inhibitory mechanisms; therefore the UV pseudoperoxidase assay is used to investigate if the inhibition is reductive in nature. The assay was performed on **35** and **36** with 12-LOX and no degradation of the hydroperoxide product was observed at 234 nm, indicating a nonreductive inhibitory mechanism (Table 4). To investigate the inhibition mechanism further, steady-state kinetics were performed using both **35** and **36** by monitoring the formation of 12-HPETE as a function of substrate and inhibitor concentration in the presence of 0.01% Triton X-100. Replots of K_m/V_{max} and 1/V_{max} versus inhibitor concentration yielded linear trends for both **35** and **36**. The K_i equaled 0.35 \pm 0.08 μ M for **35** and 0.53 \pm 0.2 μ M for **36**, from the K_m/V_{max} graphs (Figures 4 and 6). The K_{i'} equaled 0.72 \pm 0.1 μ M for **35** and 0.62 \pm 0.1 μ M for **36**, from the 1/V_{max} graphs (Figures 5 and 7). The data for both **35** and **36** correlate with their IC₅₀ values (Table 2) and indicate that both molecules are mixed inhibitors, which is a common property of both 12-LOX and 15-LOX-1 inhibitors in general.

[0284] The activity of **ML355** in relevant cell-based systems was explored. As noted previously, 12-LOX has been linked to platelet activation, which plays a central role in the regulation of primary hemostasis and arterial thrombosis. Consequently, failure to attenuate platelet activation results in excessive clot formation leading to adverse cardiovascular events such as myocardial infarction and stroke. Previous studies have shown that 12-LOX in human platelets is highly activated following stimulation of the protease-activated receptor-4 (PAR4) by the PAR4-activating peptide (PAR4-AP). Moreover, the bioactive metabolite of 12-LOX (12-HETE), resulting from the stereospecific oxidation of arachidonic acid (AA) and reduction by peroxidases, demonstrates pro-thrombotic effects in human platelets. Therefore, treatment of PAR4-AP-induced human platelets with a small molecule 12-LOX inhibitor attenuates the platelet aggregation in a dose dependent manner. To confirm this, washed human platelets (1×10^6 platelets/ml) were stimulated with 200 μ M PAR4-AP in the absence or presence of increasing concentrations of **ML355** (Figure 2A). Calcium mobilization was decreased as the concentration of **ML355** was increased. Calcium was measured in real time using a C6 Accuri flow cytometer. The experiments were done in triplicate. In addition, platelet aggregation of human platelets (3×10^8 platelets/ml) was measured in real-time using a Chronolog Lumi-Aggregometer (model 700D) following addition of PAR4-AP (Figure 2B). While 10 μ M **ML355** did not inhibit platelet aggregation, 25 μ M **ML355** inhibited 80% platelet aggregation in washed human platelets. The results in Figure 2B show that **ML355** does in fact significantly reduce PAR4-AP induced platelet aggregation. Further, 12-LOX has been shown to play a role in calcium mobilization in human platelets and inhibition of 12-LOX leads to a reduced concentration of free calcium in the platelet. To study this, human platelets were stimulated with 200 μ M PAR4-AP and the free calcium in the platelet was measured, using a C6 Accuri flow cytometer with various concentrations of **ML355** (Figure 2A). These data demonstrate that at concentrations as low as 250 nM of **ML355** calcium mobilization is reduced significantly (measured as fold change), with complete inhibition of calcium mobilization occurring at \sim 5 μ M. Comparable results were obtained with another top active (compound **36**) shown in Figure 8.

[0285] The ability of **ML355** to inhibit 12-LOX in cell-based models relevant for diabetic disease was also assessed. As discussed above, 12-LOX is expressed in pancreatic β cells and its metabolic product, 12-HETE, is implicated in cytokine-induced cell death. Specifically, 12(S)-HETE has been shown to reduce metabolic activity inhibit insulin secretion and ultimately induce cell death in human islets. **ML355** was tested in both a mouse derived β -cell line (BTC3) and human primary donor islets to determine its ability to inhibit AA/calcium ionophore induced stimulation of 12-HETE. **ML355** was able to potently inhibit 12-HETE in BTC3 cells with an

approximate IC₅₀ of 1 μ M, as measured by ELISA (Figure 3A). Given the difficulty in obtaining primary human islets from donated tissues, the activity in human islets was performed at a single concentration. The data presented in Figure 3B demonstrates significant inhibition of AA/IONO-induced 12-HETE production, at 10 μ M of **ML355**. Comparable results were obtained with another top active (compound **36**) shown in Figure 9.

[0286] Mouse beta cells (BTC3) were treated with arachidonic acid and calcium ionophore (AA/IONO) alone or in the presence of **ML355** (Figure 3A). Graphed are the levels of 12-HETE expressed as a percentage of that detected in cells stimulated with AA/IONO alone. In Figure 3B the graph represents the increase (above control/unstimulated) in 12-HETE for human primary donor islets stimulated with arachidonic acid and calcium ionophore (AA/IONO) alone or in the presence of 10 μ M of **ML355**. 12-HETE was measured by ELISA. In Figure 3C the graph represents the levels of 12-HETE detected when human donor islets were treated with arachidonic acid and calcium ionophore (AA/IONO) alone or in the presence of different concentrations of compound **35** or **36**. The data graphed in figure 3C is a representative experiment with each plotted data point being performed in triplicate. These plotted data are representative of three separate experimental determinations each performed on separate human donor islet preparations. The graphed data are mean \pm SEM, n=3. Error bars for some points are masked by the symbol.

[0287] Figures 3D and 3E show results from glucose-stimulated insulin secretion in human donor islet preparations in response to 3 mM glucose (Low) or 18 mM glucose (High) for control (untreated) islets. The islets were incubated with Cytokines (TNF α , IL-1 β , IFN γ) and islets co-incubated with cytokines and compounds **35** or **36**. Figures 3D and 3E represent results obtained in different human donor islet preparations. Uncoupling of glucose-stimulated insulin secretion by cytokines is inhibited by compounds **35** or **36**.

[0288] While the above data demonstrates activity in disease relevant cell-based models, in order to validate the potential use of **ML355** in proof of concept animal models, both its *in vitro* ADME and *in vivo* PK properties were determined. These data are summarized in Tables 5 and 6 respectively. **ML355** demonstrated excellent microsomal stability with both rat (T_{1/2} >30 minutes) and mouse (T_{1/2} >300 minutes) and was found to be stable to mouse plasma over a 2 hour period (100% remaining). Moreover, **ML355** showed no degradation over various aqueous buffers (pH 2-9) and was stable to 5 mM glutathione suggesting excellent stability (Supplemental Figure S5). Improved solubility is observed in assay buffer (qualitative analysis). In addition the benzoxazole derivative (**36**) has much improved solubility in buffer (>10-fold) albeit with slightly weaker potency towards 12-LOX.

Table 5. In Vitro ADME Profile for 35

<u>Microsomal stability</u>								
	PBS buffer (pH 7.4)	solubility (µg/mL)	T _{1/2} (min) ^{a,b}	T _{1/2} (min) ^{a,b}	permeability (1 × 10 ⁻⁶ cm/s)	efflux ratio	mouse plasma stability remaining at 2 h (%)	PBS buffer (pH 7.4)
compd 35	<5	>30 (rat)	>300 ^a (mouse)	1.5 (A→B) ^a	1.8	100	100	100
36	>60	>30 (rat)	NT	168 ^c	NT	100	100	100

^aThese experiments were conducted at Pharmaron Inc. All other studies were conducted at NCATS.

^bRepresents the stability in the presence of NADPH. 35 and 36 showed no degradation without NADPH present over a 1 h period. ^cRepresents permeability in the Parallel Artificial Membrane Permeability Assay (PAMPA) at pH 7.4.

Table 6. In Vivo PK (Mouse) at 3 mpk (IV) and 30 mpk PO for 35^a

compd	route ^b	T _{1/2} (h)	T _{max} (h)	C _{max} (µM)	AUC _{inf} (µM•h)	V _d (L/kg)	Cl (mL/min/kg)	%F	MRT ^c
35	IV	3.4	NA	112	34	0.55	3.4	NA	2.44 h
	PO	2.9	0.25	38	67	NA	NA	19.8	NA

^aAll experiments were conducted at Pharmaron Inc. using male CD1 mice (6-8 weeks of age). Data was collected in triplicate at 8 time points over a 24 h period. ^bFormulated as a solution (5% DMSO, 10% Solutol, 20% PEG400, 65% water). ^cRepresents the time for elimination of 63.2% of the IV dose.

[0289] **ML355** showed moderate cell permeability in the Caco-2 assay (1.5×10^{-6} cm/s) and does not appear to be a substrate for Pgp given the efflux ratio of <2. The *in vivo* PK (mouse) properties of the molecule were also explored (Table 6). Exploratory formulation studies led to an appropriate vehicle (DMSO:Solutol:PEG400:water; 5/10/20/65 v/v/v/v) in which **ML355** was administered as a solution via IV (3 mpk) and PO (30 mpk). These studies demonstrate that **ML355** is orally bioavailable (%F = ~20), with a moderate half-life ($T_{1/2} = 2.9$ hours). At 30 mpk dosing, **ML355** achieves a C_{max} of over 135 times the *in vitro* IC_{50} and remains over IC_{50} value for over 12 hours. The compound has low clearance (3.4 mL/min/kg) and good overall exposure (AUC_{inf}) of 38 µM. The volume of distribution (V_d) is 0.55 L/kg, which is low but suggests a reasonable distribution between tissue and blood. Given the favorable microsomal stability (phase I metabolism) yet modest *in vivo* $T_{1/2}$ it was believed that the phenolic moiety could be glucuronidated (phase II metabolism) leading to higher clearance than anticipated. In fact, incubation with UDPGA co-factor instead of NADPH led to a $T_{1/2}$ of ~8 minutes (vs. >30 minutes with NADPH). It was further believed that the sterically hindered environment of the 2-OH would possibly prevent glucuronidation from occurring, yet this data suggests otherwise. Another strategy which has been used to obviate a glucuronidation is to introduce electron-withdrawing groups next to the phenolic moiety of interest. Using this strategy analog **71** (2-OH, 3-Cl) was synthesized,

however, this did not change the rate of glucuronidation. In addition to introducing other electron-withdrawing groups to the ring, another approach is to modify the phenolic hydroxyl using a pro-drug approach. Ideally, the pro-drug would be slowly hydrolyzed to the free phenol, after it has bypassed first-pass metabolism. This approach has been used successfully in several marketed drugs, which contain phenolic groups. (Ettmayer, P.; Amidon, G. L.; Clement B.; Testa, B. Lessons learned from marketed and investigational prodrugs. *J. Med. Chem.* **2004**, *47*, 2393-2404.)

[0290] As stated above, previously reported inhibitors of 12-LOX, such as baicalein and nor-dihydroguaiaretic acid (NDGA), “bromo-phenols” or “pyrazole derivatives” (see Figure 1) are not only less potent and selective, but are also not easily amendable to further optimization. Our previously described 12-LOX inhibitor (**ML127**) does demonstrate potent inhibition (<500 nM) and excellent selectivity, but was found to exhibit tight flat SAR thus providing little opportunity for further modification. A structurally distinct, new chemotype that is different from all previously reported inhibitors, and possesses a drug-like scaffold was discovered. **ML355**, and related top analogs demonstrate potent (<500 nM) inhibition towards 12-LOX and excellent selectivity against related enzymes (15-LOX-1, 5-LOX, 15-LOX-2, COX 1/2). This series is readily amendable to structural modifications and displays clear and tractable SAR. **ML355** exhibits a favorable *in vitro* ADME and *in vivo* PK profile with activity in disease relevant cell-based systems, such thrombosis (platelet aggregation and calcium mobilization) and diabetes (12-HETE reducing in β -cells).

[0291] Biological Reagents: All commercial fatty acids (Sigma-Aldrich Chemical Company) were re-purified using a Higgins HAIsil Semi-Preparative (5 μ m, 250 \times 10mm) C-18 column. Solution A was 99.9% MeOH and 0.1% acetic acid; solution B was 99.9% H₂O and 0.1% acetic acid. An isocratic elution of 85% A:15% B was used to purify all fatty acids, which were stored at -80 °C for a maximum of 6 months.

[0292] Human Platelets: Human platelets were obtained from healthy volunteers within the Thomas Jefferson University community and the Philadelphia area. These studies were approved by the Thomas Jefferson University Institutional Review Board, and informed consent was obtained from all donors before blood draw. Blood was centrifuged at 200 g for 13 min at room temperature. Platelet-rich plasma was transferred into a conical tube containing a 10% acid citrate dextrose solution (39 mM citric acid, 75 mM sodium citrate, and 135 mM glucose, pH 7.4) and centrifuged at 2000 g for 15 min at room temperature. Platelets were resuspended in Tyrode's buffer (12 mM NaHCO₃, 127 mM NaCl, 5 mM KCl, 0.5 mM NaH₂PO₄, 1 mM MgCl₂, 5 mM glucose, and 10 mM HEPES), and the final platelet concentration was adjusted to 3 X 10⁸ platelets/mL after counting with a ZI Coulter particle counter (Beckman Coulter, Fullerton, CA).

Reported results are the data obtained using platelets from at least three different subjects. Agonists and inhibitors were used at concentrations indicated in the figures and figure legends.

[0293] Over expression and Purification of 12-Human Lipoxygenase, 5-Human Lipoxygenase, 12/15-Mouse Lipoxygenase and the 15-Human Lipoxygenases: Human platelet 12-lipoxygenase (12-LOX), human reticulocyte 15-lipoxygenase-1 (15-LOX-1), and human epithelial 15-lipoxygenase-2 (15-LOX-2), were expressed as N-terminally, His₆-tagged proteins and purified to greater than 90% purity, as evaluated by SDS-PAGE analysis. Human 5-lipoxygenase was expressed as a non-tagged protein and used as a crude ammonium sulfate protein fraction, as published previously. Iron content of 12-LOX was determined with a Finnigan inductively coupled plasma mass spectrometer (ICP-MS), using cobalt-EDTA as an internal standard. Iron concentrations were compared to standardized iron solutions and used to normalize enzyme concentrations.

[0294] High-throughput Screen Materials: Dimethyl sulfoxide (DMSO) ACS grade was from Fisher, while ferrous ammonium sulfate, Xylenol Orange (XO), sulfuric acid, and Triton X-100 were obtained from Sigma-Aldrich.

[0295] 12-Lipoxygenase qHTS Assay (AID: 1452): All screening operations were performed on a fully integrated robotic system (Kalypsys Inc, San Diego, CA). (Inglese, J. et al. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 11473–11478.) Three μ L of enzyme (approximately 80 nM 12-LOX, final concentration) was dispensed into 1536-well Greiner black clear-bottom assay plates. Compounds and controls (23 nL) were transferred via Kalypsys PinTool equipped with 1536-pin array. The plate was incubated for 15 min at room temperature, and then a 1 μ L aliquot of substrate solution (50 μ M arachidonic acid final concentration) was added to start the reaction. The reaction was stopped after 6.5 min by the addition of 4 μ L FeXO solution (final concentrations of 200 μ M Xylenol Orange (XO) and 300 μ M ferrous ammonium sulfate in 50 mM sulfuric acid). After a short spin (1000 rpm, 15 sec), the assay plate was incubated at room temperature for 30 minutes. The absorbances at 405 and 573 nm were recorded using ViewLux high throughput CCD imager (Perkin-Elmer, Waltham, MA) using standard absorbance protocol settings. During dispense, enzyme and substrate bottles were kept submerged into a +4 °C recirculating chiller bath to minimize degradation. Plates containing DMSO only (instead of compound solutions) were included approximately every 50 plates throughout the screen to monitor any systematic trend in the assay signal associated with reagent dispenser variation or decrease in enzyme specific activity. Data were normalized to controls, and plate-based data corrections were applied to filter out background noise. Further details on data analysis are provided herein.

[0296] Lipoxygenase UV-Vis Assay: The inhibitor compounds were screened initially using one concentration point at 25 μ M on a Perkin-Elmer Lambda 40 UV/Vis spectrophotometer. The percent inhibition was determined by comparing the enzyme rates of the control (DMSO solvent) and the inhibitor sample by following the formation of the conjugated diene product at 234 nm ($\epsilon = 25,000 \text{ M}^{-1} \text{ cm}^{-1}$). The reactions were initiated by adding either of 30 nM 12-LOX, 40 nM 15-LOX-1, 200 nM 15-LOX-2 or 5-10 μ L of 5-LOX crude extract to a cuvette with a 2 mL reaction buffer constantly stirred using a magnetic stir bar at room temperature (22 °C). Reaction buffers used for various lipoxygenase were as follows: 25 mM HEPES (pH 7.3), 0.3 mM CaCl₂, 0.1 mM EDTA, 0.2 mM ATP, 0.01% Triton X-100, 10 μ M AA for the crude, ammonium sulfate precipitated 5-LOX; and 25 mM HEPES (pH 7.5), 0.01% Triton X-100, 10 μ M AA for 12-LOX, 15-LOX-1 and 15-LOX-2. The substrate concentration was quantitatively determined by allowing the enzymatic reaction to go to completion in the presence of 15-LOX-2. For the inhibitors that showed more than 50% inhibition at the one point screens, IC₅₀ values were obtained by determining the % inhibition, relative to solvent vehicle only, at various inhibitor concentrations. The data was then plotted against inhibitor concentration, followed by a hyperbolic saturation curve fit (assuming total enzyme concentration [E] << K_i^{app}, so IC₅₀ ~ K_i^{app}). It should be noted that all of the potent inhibitors displayed greater than 80% maximal inhibition, unless stated in the tables. Inhibitors were stored at -20 °C in DMSO

[0297] Steady State Inhibition Kinetics: The steady-state kinetics experiments were performed with **ML355** to determine the mode of inhibition. The inhibitor concentrations of 0, 0.1, 0.2 and 0.35 μ M were used. Reactions were initiated by adding substrate (range 1 – 5 μ M AA) to approximately 30 nM 12-LOX in a constantly stirring 2 mL cuvette containing 25 mM HEPES buffer (pH 7.5), in the presence of 0.01% Triton X-100. Lipoxygenase rates were determined by monitoring the formation of the conjugated product, 12-HPETE, at 234 nm ($\epsilon = 25\,000 \text{ M}^{-1} \text{ cm}^{-1}$) with a Perkin-Elmer Lambda 45 UV/Vis spectrophotometer. It should be noted that 12-LOX displays higher error in the K_M values at low substrate concentrations (< 1 μ M) due to the limits of the spectrophotometer. To minimize this inherent error, it is best to add 12-LOX first, and then quickly initiate the reaction with the addition of the appropriate amount of substrate, which yielded significantly more reproducible results. The substrate concentration was quantitatively determined by allowing the enzymatic reaction to proceed to completion, using 15-LOX-2. Kinetic data were obtained by recording initial enzymatic rates, at varied substrate and inhibitor concentrations, and subsequently fitted to the Henri-Michaelis-Menten equation, using KaleidaGraph (Synergy) to determine the microscopic rate constants, V_{max} ($\mu\text{mol}/\text{min}/\text{mg}$) and V_{max}/K_M ($\mu\text{mol}/\text{min}/\text{mg}/\mu\text{M}$).

The kinetic rate constants were subsequently replotted with $1/V_{max}$, K_M/V_{max} and $1/K_M$ versus inhibitor concentration, yielding K_i .

[0298] Pseudoperoxidase Assay: The pseudo-peroxidase activity rates were determined with BWb70c as the positive control, 13-(S)-HPODE as the oxidizing product and 12-LOX or 15-LOX-1 on a Perkin-Elmer Lambda 40 UV/Vis spectrophotometer, as described previously.²⁵ Activity was determined by monitoring the decrease at 234 nm (product degradation) in buffer (50 mM Sodium Phosphate (pH 7.4), 0.3 mM CaCl₂, 0.1 mM EDTA, 0.01% Triton X100, and 20 μ M 13-(S)-HPODE). About 60 nM 12-LOX was added to 2 mL buffer containing 20 μ M 13-(S)-HPODE, constantly stirred with a rotating stir bar (22 °C). Reaction was initiated by addition of 20 μ M inhibitor (1:1 ratio to product). The percent consumption of 13-(S)-HPODE was recorded and loss of product less than 20% was not considered as viable redox activity. Individual controls were conducted consisting of enzyme alone with product and **ML355** alone with enzyme. These negative controls formed the baseline for the assay, reflecting non-pseudo-peroxidase dependent hydroperoxide product decomposition. To rule out the auto-inactivation of the enzyme from pseudo-peroxidase cycling, the 12-LOX residual activity was measured after the assay was complete. 20 μ M AA was added to the reaction mixture and the residual activity was determined by comparing the initial rates with inhibitor and 13-(S)-HPODE versus inhibitor alone, since the inhibitor by itself inherently lowers the rate of the oxygenation. Activity is characterized by direct measurement of the product formation with the increase of absorbance at 234 nm.

[0299] Cyclooxygenase assay: Roughly 2-5 μ g of either COX-1 or COX-2 were added to buffer containing 0.1 M Tris-HCl buffer (pH 8.0), 5 mM EDTA, 2 mM phenol and 1 μ M hematin at 37 °C. The selected inhibitors were added to the reaction cell, followed by an incubation of 5 minutes with either of the COX enzymes. The reaction was then initiated by adding 100 μ M AA in the reaction cell. Data was collected using a Hansatech DW1 oxygen electrode and the consumption of oxygen was recorded. Indomethasin and the solvent DMSO, were used as positive and negative controls, respectively and the percent inhibition of the enzyme was calculated by comparing the rates from samples and the controls

[0300] Platelet Aggregation: Washed platelets were adjusted to a final concentration of 3 X 10⁸ platelets/mL. Where indicated, platelets were pretreated with **ML355** for 10 min at the indicated concentrations for 1 min. The aggregation response to PAR4-AP was measured using an aggregometer with stirring at 1100 RPM at 37 °C.

[0301] Calcium Mobilization: Platelets were recalcified to a final concentration of 1 mM followed by pre-incubation with Fluo-4 AM for 10 min. The platelets were then treated with

ML355 for 10 min at the indicated concentrations before stimulation with the indicated agonist. Calcium mobilization was measured using the Accuri C6 flow cytometer.

[0302] Mouse Beta cells (12-HETE Inhibition) Assay: Cells were grown to 90% confluence in 24 well plates in DMEM (Cat# 11885092, Life Technologies Grand Island, NY) +10% FBS. Cells were pre-treated with **ML355** and stimulated as for human islets. After four hours, the media was removed and spun at 1000 RPM for 5 minutes. The cleared supernatant was stored at -80 °C prior to analysis. For analysis, supernatants were extracted on SepPak c18 SPE column (Cat# WAT054945, Waters Corporation, Milford, MA) and dried under nitrogen gas before reconstitution in 500 µL of 12-HETE ELISA buffer and analysis following manufacturers recommendations (Cat# 901-050, Enzo Life Sciences, Plymouth Meeting, PA).

[0303] Human Islet (12-HETE Inhibition) Assay: Human donor islets obtained from integrated islet distribution program (www.iidp.coh.org) were incubated overnight in CMRL media (Cat# 15-110-CV MediaTech, Inc. Manassas, VA) containing 10% Fetal Bovine Serum, 1U penicillin 1 µg streptomycin (pen/strep). Islets were equilibrated in serum free media, (CMRL containing pen/Strep and 1% fatty acid free human serum albumin (Cat# A1887 Sigma, St. Louis, MO)), for 1 hour prior to pretreatment with 10µM **ML355** for 30 mins. For 12-HETE induction, islets were treated with 100 µM arachachidonic acid (Cat# BML-FA003-0100, Enzo Life Sciences Plymouth Meeting, PA), and 5 µM A23187 (Cat# C7522, Sigma, St. Louis, MO), for 4 hours at 37 °C. Islets were harvested, centrifuged at 1000 RPM for 5 minutes with cleared supernatant and islet pellet being stored at -80 °C. For extraction of the supernatants, samples were acidified to pH 3 with 1N HCl for 30 minutes and spun at 1000 RPM for five minutes. Samples were added to a prepared column (prewashed with 3 mL EtOH, followed by 3 mL of H₂O) and washed with 3 mL H₂O, followed by 3 mL 15% EtOH, and 3 mL Hexane. The samples were eluted with 3 mL of ethyl acetate and dried under nitrogen gas before, reconstitution in 500 mL of 12-HETE ELISA sample buffer (Enzo Life Sciences, Plymouth Meeting, PA). Cell pellets were extracted using CHCl₃ / MeOH and samples dried under nitrogen gas before reconstitution in 250 µL of ELISA sample buffer. 12-HETE levels in samples were determined using a 12-HETE ELISA kit (Cat# 901-050, Enzo Life Sciences, Plymouth Meeting, PA).

Examples Related To The Use of 12-Lipoxygenase Inhibitors in Platelet Activation And Treatment Of Thrombosis

Example: Platelet 12-LOX is essential for FcγRIIa-mediated platelet activation

[0304] Pharmacological inhibition of 12-LOX in human platelets resulted in significant attenuation of FcγRIIa-mediated aggregation. 12-LOX was shown to be essential for FcγRIIa-

induced PLC γ 2 activity leading to activation of calcium mobilization, Rap1 and PKC activation, and subsequent activation of the integrin α IIb β 3. Additionally, platelets from transgenic mice expressing human Fc γ RIIa but deficient in platelet 12-LOX failed to form normal platelet aggregates and exhibited deficiencies in Rap1 and α IIb β 3 activation following stimulation of the Fc γ RIIa receptor. These results support an essential role for 12-LOX in regulating Fc γ RIIa-mediated platelet function and identify 12-LOX as a therapeutic target to limit immune-mediated thrombosis.

[0305] Platelet 12(S)-lipoxygenase (12-LOX), an oxygenase highly expressed in platelets, has been shown to potentiate the activation of select signaling pathways including protease-activated receptor 4 (PAR4) and an ITAM-containing receptor complex (GPVI-FcR γ). The most well understood function of 12-LOX is production of oxylipins, most notable being the conversion of arachidonic acid (AA) to 12-hydroxyeicosatetraenoic acid (12-HETE) upon agonist stimulation of platelets through both GPCR- and non-GPCR-mediated pathways. 12-HETE is an oxylipin that has been shown to be pro-thrombotic in platelets. In carotid endarterectomy patients, intra-operative heparin increases the level of plasma 12-HETE which correlates with the patient's platelet reactivity to ADP *ex vivo*. While the mechanism by which 12-LOX regulates platelet activity remains to be fully elucidated, previous publications have demonstrated the ability of 12-LOX activity to augment key signaling components of platelet activation including Rap1, Ca^{2+} mobilization, α IIb β 3 activation, and dense granule secretion. Since 12-LOX activity was recently shown to be required for normal GPVI-mediated platelet activation and Fc γ RIIa shares common signaling components with GPVI, the inventors sought to identify if 12-LOX is a vital component of the Fc γ RIIa signaling pathway.

[0306] In this study human platelets were treated with a 12-LOX inhibitor, ML355, or vehicle control prior to Fc γ RIIa stimulation to determine if 12-LOX plays a role in the Fc γ RIIa signaling pathway. Pharmacological inhibition of 12-LOX activity in human platelets attenuated Fc γ RIIa-mediated platelet aggregation. Consistent with the human studies, murine platelets isolated from hFcR transgenic mice deficient in 12-LOX had an attenuated aggregation in response to Fc γ RIIa stimulation compared to wild-type hFcR transgenic mice. The activity of 12-LOX was further demonstrated to be essential for early PLC γ 2 activation as well as other downstream effectors, such as calcium mobilization, Rap1, PKC, α IIb β 3 activation, and dense granule secretion, while not directly impinging on the phosphorylation of Fc γ RIIa itself. This study is the first to identify 12-LOX activity as an essential component of normal Fc γ RIIa signaling in platelets. Further, the results of this study suggest for the first time that 12-LOX may represent a novel therapeutic target to treat immune-mediated thrombocytopenia and thrombosis.

[0307] *12-LOX regulates Fc γ RIIa-mediated platelet aggregation*

[0308] 12-LOX was previously shown to regulate GPVI-mediated platelet aggregation. In platelets, Fc γ RIIa utilizes many of the same downstream signaling effectors as GPVI-FcR γ (4). To determine if 12-LOX plays a role in Fc γ RIIa-mediated platelet aggregation, washed human platelets were treated with ML355, a selective 12-LOX inhibitor or DMSO (vehicle control) followed by Fc γ RIIa-induced aggregation. Inhibition of platelet 12-LOX activity attenuated Fc γ RIIa-mediated platelet aggregation via anti-CD9 stimulation (Fig. 11A) or Fc γ RIIa antibody cross-linking (Fig. 11B). To confirm the decrease in Fc γ RIIa-mediated platelet aggregation was due to pharmacological inhibition of 12-LOX and not an off-target effect of the ML355, ex vivo platelet aggregation was measured following anti-CD9 stimulation in humanized Fc γ RIIa (hFcR) transgenic mice expressing 12-LOX (ALOX12 $^{+/+}$) or deficient in 12-LOX (ALOX12 $^{-/-}$). Platelets from mice deficient in 12-LOX (hFcR/ALOX12 $^{-/-}$) showed a delayed (Fig. 12B) and attenuated (Fig. 12C) aggregation in response to anti-CD9 stimulation (.125 and .25 μ g/mL) when compared to platelets from mice expressing functional 12-LOX (hFcR/ALOX12 $^{+/+}$). These data suggest that platelets lacking 12-LOX activity through pharmacological or genetic ablation exhibits a significantly attenuated aggregation response to Fc γ RIIa activation.

[0309] *12-LOX regulates α IIb β 3 and Rap1 activity in Fc γ RIIa stimulated platelets*

[0310] Figs. 11 and 12 suggest 12-LOX is essential for normal Fc γ RIIa-mediated platelet aggregation; however, the component(s) in the Fc γ RIIa pathway regulated by 12-LOX remain unclear. As the activation of integrin α IIb β 3 is required for platelet aggregation, the potential role of 12-LOX in mediating α IIb β 3 activation in Fc γ RIIa-stimulated platelets was investigated, α IIb β 3 activation was measured by flow cytometry in Fc γ RIIa-stimulated platelets treated with ML355. Rap1 is a small GTPase whose activation is essential for activation of α IIb β 3. Therefore, Rap1 activation was measured following Fc γ RIIa stimulation in platelets following treatment with or without ML355. As shown in Fig. 13B, while stimulation of platelets in the absence of pre-treatment (control) resulted in a significant increase in Rap1-GTP, platelets treated with ML355 failed to activate Rap1. Platelets treated with DMSO showed no attenuation in active Rap1 compared to control.

[0311] *12-LOX regulates dense but not alpha granule secretion in Fc γ RIIa activated platelets*

[0312] The release of alpha and dense granules from activated platelets serves to amplify platelet aggregation. Surface expression of P-selectin was used as a surrogate marker to measure alpha granule secretion in Fc γ RIIa-stimulated platelets. Treatment with ML355 did not result in attenuation of agonist-induced surface expression of P-selectin compared to control conditions (Fig. 14A). However, Fc γ RIIa-stimulated platelets treated with ML355 showed a significant

decrease in ATP release compared with DMSO treated platelets (Fig. 14B). These data suggest 12-LOX is a regulator of dense secretion downstream of Fc γ RIIa stimulation in platelets.

[0313] *12-LOX modulates proximal signaling components of the Fc γ RIIa pathway in human platelets*

[0314] As shown in Fig. 13, 12-LOX activity is required for the normal activation of distal signaling components (Rap1, α IIb β 3, and platelet aggregation) of the Fc γ RIIa pathway. Crosslinking of Fc γ RIIa with immunoglobulins also initiates proximal signaling components' activation that eventually leads to platelet aggregation. Fc γ RIIa initiates ITAM phosphorylation resulting in Syk activation. Syk activation leads to activation of phospholipase gamma 2 (PLC γ 2) resulting in calcium release and PKC activation, both of which are critical for platelet activation. To determine where in this complex signaling milieu 12-LOX plays an essential role in Fc γ RIIa signaling in the platelet, the individual signaling components directly downstream of Fc γ RIIa initiation were assessed in the presence or absence of ML355.

[0315] To investigate whether 12-LOX directly regulated Fc γ RIIa phosphorylation, Fc γ RIIa was immunoprecipitated from IV.3+GAM-stimulated platelets treated with ML355 or DMSO and immunoblotted for phosphorylation of Fc γ RIIa. No difference in Fc γ RIIa phosphorylation was detected in Fc γ RIIa-stimulated platelets in control or ML355 treated conditions (Fig. 15A). This data suggests that 12-LOX activity is not required for phosphorylation of Fc γ RIIa itself.

[0316] Phosphorylation of Fc γ RIIa results in recruitment and activation of the tyrosine kinase Syk which is crucial for the formation of the LAT signalosome. The phosphorylation of Syk was measured following Fc γ RIIa stimulation of platelets in the presence or absence of ML355. The degree of Syk phosphorylation in response to Fc γ RIIa cross-linking in the presence of ML355 was varied in the subjects tested and may be due to inter-individual variability. Hence, while the observed attenuation of Syk phosphorylation in the presence of ML355 was minor, the number of subjects tested was too small to definitively identify Syk as a node of regulation by 12-LOX (Fig. 15B). To assess if 12-LOX regulates PLC γ 2 activation downstream of Fc γ RIIa, PLC γ 2 phosphorylation was measured in washed human platelets in the presence or absence of ML355. Platelets stimulated by Fc γ RIIa antibody cross-linking were phosphorylated within 15 seconds following stimulation. Platelets treated with ML355 however showed attenuated PLC γ 2 phosphorylation at 15 and 30 seconds compared to control conditions (Fig. 16A).

[0317] Since 12-LOX attenuated Fc γ RIIa-mediated PLC γ 2 activation, its effects on intracellular calcium release were measured. Washed human platelets were stimulated by Fc γ RIIa-mediated antibody cross-linking in the presence of ML355 or vehicle control and calcium release was measured. Platelets treated with ML355 exhibited a decrease in intracellular Ca $^{2+}$ following

platelet stimulation compared to DMSO treated platelets (Fig. 16B). As PLC γ 2 activation leads to activation of PKC in platelets in part through calcium mobilization, the potential regulation of Fc γ RIIa-mediated PKC activity by 12-LOX was also evaluated. A significant decrease in PKC activation in platelets treated with ML355 at 30 and 60 seconds was observed compared to platelets treated with the vehicle control (Fig. 16C). This regulation of PKC was determined to be indirect since PKC activation by PMA, a direct PKC activator, showed no difference in its ability to activate PKC either in control or ML355 treated platelets.

Discussion

[0318] 12-LOX has recently been shown to be an important regulator of GPVI-mediated platelet activation. As Fc γ RIIa and GPVI are purported to signal via a conserved pathway, 12-LOX may play an essential role in the regulation of Fc γ RIIa signaling in human platelets. This study is the first to demonstrate that 12-LOX is an essential component of Fc γ RIIa immune mediated platelet activation. Human platelets treated with a 12-LOX inhibitor, ML355, or Fc γ RIIa transgenic mouse platelets deficient in 12-LOX, showed significantly attenuated aggregation in response to Fc γ RIIa-mediated activation. To investigate the underlying mechanism by which 12-LOX regulates Fc γ RIIa-mediated platelet activation, the activity of multiple signaling intermediates in the Fc γ RIIa pathway were assessed in the presence of the 12-LOX inhibitor ML355. Following stimulation, platelets treated with ML355 were significant attenuated along multiple signaling steps in the immune-mediated Fc γ RIIa activation pathway including α IIb β 3, Rap1, Ca²⁺, PLC γ 2, PKC, and dense granule secretion (Fig. 17).

[0319] The primary known function of 12-LOX is to produce oxylipins (such as 12(S)-HETE) from the fatty acid substrate AA, the predominant fatty acid in the platelet, following platelet activation. Previous publications have shown selective 12-LOX inhibitors significantly reduced 12-HETE production. Though blocking 12-HETE production has been associated with reduced platelet-mediated reactivity, the direct role of 12-HETE on platelet activation remains unclear. Oxylipins generated from COX-1, another oxygenase in platelets, generally act as signaling mediators through GPCRs on the platelet surface. In the case of 12-HETE, published work has demonstrated it signals through GPR31, a GPCR; however, while recent genetic screens have shown the expression of GPR31 in human platelets, there is no direct evidence of GPR31 protein expression on the surface of platelets. In addition to potential autocrine or paracrine signaling through GPR31, 12-LOX oxylipins produced in the platelet may incorporate back into the membrane to induce their regulatory effects. There is a precedent for oxylipin integration into the membrane as 12-HETE has been shown to be esterified into phospholipids in the plasma membrane upon platelet activation. A third potential explanation of the mechanism by which 12-LOX

regulates Fc γ RIIa-mediated signaling may be through direct or indirect 12-LOX scaffolding within the cell. 12-LOX complex formation as a means to regulate downstream signaling has previously been implicated in regulation of cell growth and tumorigenesis. While all of these regulatory mechanisms are plausible, the rapid speed by which 12-LOX impinges on the Fc γ RIIa pathway leaves favors the possibility that 12-LOX and not a 12-LOX oxylipin may act as a direct regulatory component of the Fc γ RIIa pathway.

[0320] Although 12-LOX activity is required for normal Fc γ RIIa-mediated platelet activation, the direct molecular component by which 12-LOX activity is required has yet to be determined. 12-LOX was not required for Fc γ RIIa phosphorylation and only partially affected Syk phosphorylation, suggesting 12-LOX activity is not directly regulating Src family kinase activity. However, 12-LOX activity was shown to be important for early PLCy2 activation, indicating 12-LOX may be an important regulator in the kinetics of PLCy2 activation affecting downstream effectors. The delay in PLCy2 activation due to 12-LOX inhibition may be attributed to direct regulation of PLCy2 or regulation of upstream effectors such as LAT or BTK. The data presented here narrows the scope of where 12-LOX impinges on the Fc γ RIIa pathway to a proximal point in the signaling pathway between the receptor and PLCy2.

Methods

[0321] Mice and platelet preparation: Fc γ RIIA transgenic mice (hFcR/ALOX12⁺⁺) were bred with platelet 12-lipoxygenase knockout (ALOX12^{-/-}) mice to generate Fc γ RIIA transgenic mice deficient in platelet 12-lipoxygenase (hFcR/ALOX12^{-/-}). All mice were housed in the mouse facility of Thomas Jefferson University (TJU). Experimental procedures were approved by the Animal Care and Use Committee of TJU. Blood was drawn from the inferior vena cava of 12 week old anesthetized mice using a syringe containing sodium citrate. Mouse platelet preparation was prepared as described in (Yeung et al., 2012). Murine platelets were resuspended in tyrodes buffer containing fibrinogen (75 μ g/mL) and CaCl₂ (1 mM).

[0322] Preparation of washed human platelets: Prior to blood collection written informed consent was obtained from all volunteers. Washed platelets were isolated from the blood of healthy volunteers as previously described (14). Washed platelets were resuspended in tyrodes buffer at 3.0 $\times 10^8$ platelets/mL unless otherwise indicated. All studies performed with human platelets were approved by the Thomas Jefferson University Institutional Review Board.

[0323] Fc γ RIIa-mediated platelet activation: Fc γ RIIa-mediated platelet activation was initiated by two distinct models; either 1) Fc γ RIIa antibody cross-linking or 2) CD9 monoclonal antibody stimulation. To cross-link Fc γ RIIa, washed platelets were incubated with IV.3, an Fc γ RIIa mouse monoclonal antibody, for one minute followed by the addition of a goat anti-mouse (GAM) IgG

antibody to cross-link Fc γ RIIa. The concentration of Fc γ RIIa cross-linking antibodies used for each experiment is indicated in the text. Alternatively, washed human platelets were stimulated with an anti-CD9 monoclonal antibody to activate Fc γ RIIa mediated signaling. Due to inter-individual variability in anti-CD9 response, a range of anti-CD9 concentration (.25 – 1 ug/ml) was used to achieve an aggregation of EC80. In studies using the 12-LOX inhibitor (ML355), washed platelets were incubated with either ML355 (20 μ M), or DMSO (vehicle control) for 15 minutes prior to Fc γ RIIa stimulation.

[0324] Platelet aggregation: Platelet aggregation was measured with a lumi-aggregometer (Chronolog Corp, Model 700D) under stirring conditions at 37°C.

[0325] PLC γ 2 phosphorylation: Washed human platelets were adjusted to 5×10^8 platelets/mL and stimulated in an aggregometer by antibody cross-linking of Fc γ RIIa and lysed at designated times with 5X reducing buffer to stop the reaction. Samples were separated on a 7.5% SDS-PAGE gel. Antibodies to PLC γ 2 and phospho-Y759 PLC γ 2, a marker of PLC γ 2 activation, were used to evaluate the relative levels of total and active PLC γ 2, respectively.

[0326] Calcium mobilization: Intracellular calcium release was measured as previously described. Briefly, washed human platelets were resuspended at 1.0×10^6 platelets/mL in tyrodes buffer containing 1 mM calcium. Platelets were incubated with Fluo-4 am, a cell permeable calcium sensitive dye, for ten minutes prior to stimulation. Platelets were stimulated by Fc γ RIIa antibody cross-linking and fluorescence intensity was measured in real-time by flow cytometry. Data is reported as the fold change in the fluorescence intensity comparing maximum fluorescence intensity relative to fluorescence intensity prior to platelet stimulation.

[0327] Rap1 activation: Washed human platelets were stimulated by Fc γ RIIa antibody cross-linking for 5 minutes and aggregation was stopped with 2X platelet lysis buffer. RalGDS-RBD was used to selectively precipitate the active conformation of Rap1 from the platelet lysate as previously described. Total platelet lysate and Rap1 pull-down samples were run on a SDS-PAGE gel and identified by Western blot with a Rap1 antibody. The levels of active Rap1 were normalized to the amount of total Rap1 contained in each sample.

[0328] PKC activation assay: Washed platelets were stimulated by Fc γ RIIa antibody cross-linking under stirring conditions (1100 rpm) in an aggregometer at 37°C. Reactions were stopped by the addition of 5X Laemmli sample buffer at the indicated times. As a positive control platelets were treated with PMA (1 mM), a direct PKC agonist, for one minute. Samples were run on an SDS-PAGE gel and Western blots were performed using antibodies specific for PKC substrate phosphorylation and pleckstrin.

[0329] Dense granule secretion: ATP release was measured from washed platelets as surrogate for dense granule secretion. Prior to activation washed platelets were incubated with Chronolume Reagent, an ATP sensitive dye, for one minute. Platelets were stimulated with Fc γ RII antibody cross-linking under stirring conditions and fluorescence was measured in real-time using a lumi-aggregometer.

[0330] Alpha granule release and α IIb β 3 activation: Prior to stimulation, washed platelets were pre-incubated with either FITC-conjugated P-selectin antibody or FITC-conjugated PAC-1, an antibody specific for the active conformation of α IIb β 3. Platelets were stimulated with Fc γ RIIa antibody cross-linking and reactions were stopped by the addition of 2% formaldehyde at indicated times. Fluorescence intensity was measured by flow cytometry. Results are reported as mean fluorescence intensity.

[0331] Western Blotting: Standard Western blots for Rap1 activation, PKC substrates, and PLC γ 2 phosphorylation were used and band intensity were quantified with the Odyssey Infrared Imaging System (LIC-OR Biosystems).

[0332] Statistical analysis: Where applicable, data represents the mean \pm S.E.M. Statistical significance was determined using GraphPad Prism software. A P value less than 0.05 was considered significant.

References

1. Aharony, D.; Smith, J. B.; Silver, M. J. Regulation of arachidonate-induced platelet aggregation by the lipoxygenase product, 12-hydroperoxyeicosatetraenoic acid. *Biochim. Biophys. Acta* 1982, 718, 193-200.
2. Amagata, T.; Whitman, S.; Johnson, T.; Stessmann, C. C.; Carroll, J.; Loo, C.; Clardy, J.; Lobkovsky, E.; Crews, P.; Holman, T. R. Sponge Derived Terpenoids with Selectivity towards Human 15-Lipoxygenase versus Human 12-Lipoxygenase. *J. Nat. Prod.* 2003, 66, 230-235.
3. Arai M, Yamamoto N, Moroi M, Akamatsu N, Fukutake K, and Tanoue K. Platelets with 10% of the normal amount of glycoprotein VI have an impaired response to collagen that results in a mild bleeding tendency. *British journal of haematology*. 1995;89(1):124-30.
4. Asselin J, Gibbins JM, Achison M, Lee YH, Morton LF, Farndale RW, Barnes MJ, and Watson SP. A collagen-like peptide stimulates tyrosine phosphorylation of syk and phospholipase C gamma2 in platelets independent of the integrin alpha2beta1. *Blood*. 1997;89(4):1235-42.

5. Berger, W.; De Chandt, M. T.; Cairns, C. B. Zileuton: Clinical Implications of 5-Lipoxygenase Inhibition in Severe Airway Disease. *Int. J. Clin. Pract.* 2007, 61, 663–676.
6. Bergmeier W, and Stefanini L. Platelet ITAM signaling. *Current opinion in hematology.* 2013;20(5):445-50.
7. Bertoni A, Tadokoro S, Eto K, Pampori N, Parise LV, White GC, and Shattil SJ. Relationships between Rap1b, affinity modulation of integrin alpha IIbbeta 3, and the actin cytoskeleton. *The Journal of biological chemistry.* 2002;277(28):25715-21.
8. Blake RA, Schieven GL, and Watson SP. Collagen stimulates tyrosine phosphorylation of phospholipase C-gamma 2 but not phospholipase C-gamma 1 in human platelets. *FEBS letters.* 1994;353(2):212-6.
9. Bledzka K, Smyth SS, and Plow EF. Integrin alphaIIbbeta3: from discovery to efficacious therapeutic target. *Circulation research.* 2013;112(8):1189-200.
10. Bleich, D.; Chen, S.; Zipser, B.; Sun, D.; Funk, C. D.; Nadler, J. L. Resistance to type 1 diabetes induction in 12-lipoxygenase knockout mice. *J. Clin. Invest.* 1999, 103, 1431-1436.
11. Brash, A. R. A review of possible roles of the platelet 12-lipoxygenase. *Circulation* 1985, 72, 702-707.
12. Brass LF, Manning DR, and Shattil SJ. GTP-binding proteins and platelet activation. *Progress in hemostasis and thrombosis.* 1991;10(127-174)
13. Catalano, A.; Procopio, A. New aspects on the role of lipoxygenases in cancer progression. *Histol. Histopathol.* 2005, 20, 969-975.
14. Chang, J.; Blazek, E.; Kreft, A. F.; Lewis, A. J. Inhibition of platelet and neutrophil phospholipase A2 by hydroxyeicosatetraenoic acids (HETES). A novel pharmacological mechanism for regulating free fatty acid release. *Biochem. Pharmacol.* 1985, 34, 1571-1575.
15. Chen, M.; Yang, Z. D.; Smith, K. M.; Carter, J. D.; Nadler, J. L. Activation of 12-Lipoxygenase in proinflammatory cytokine-mediated β -cell toxicity. *Diabetologia* 2005, 48, 486-495.
16. Chen, X.-S.; Brash, A.; and Funk, C. Purification and characterization of recombinant histidine-tagged human platelet 12-lipoxygenase expressed in a baculovirus/insect cell system. *Eur. J. Biochem.* 1993, 214, 845-852.

17. Chrzanowska-Wodnicka M, Smyth SS, Schoenwaelder SM, Fischer TH, and White GC, 2nd. Rap1b is required for normal platelet function and hemostasis in mice. *The Journal of clinical investigation*. 2005;115(3):680-7.
18. Cichewicz, R. H.; Kenyon, V. A.; Whitman, S.; Morales, N. M.; Arguello, J. F.; Holman, T. R.; Crews, P. Redox inactivation of human 15-lipoxygenase by marine-derived meroterpenes and synthetic chromanes: archetypes for a unique class of selective and recyclable inhibitors. *J. Am. Chem. Soc.* 2004, 126, 14910–14920.
19. Coffey MJ, Jarvis GE, Gibbins JM, Coles B, Barrett NE, Wylie OR, and O'Donnell VB. Platelet 12-lipoxygenase activation via glycoprotein VI: involvement of multiple signaling pathways in agonist control of H(P)ETE synthesis. *Circulation research*. 2004;94(12):1598-605.
20. Cyrus, T.; Witztum, J. L.; Rader, D. J.; Tangirala, R.; Fazio, S.; Linton, M. F.; Funk, C.D. Disruption of the 12/15-lipoxygenase gene diminishes atherosclerosis in apo E-deficient mice *J. Clin. Invest.* 1999, 103, 1597-1604.
21. Deschamps, J. D.; Gautschi, J. T.; Whitman, S.; Johnson, T. A.; Gassner, N. C.; Crews, P.; Holman, T. R. Discovery of platelet-type 12-human lipoxygenase selective inhibitors by high-throughput screening of structurally diverse libraries, *Bioorg. Med. Chem.* 2007, 15, 6900-6908.
22. Deschamps, J. D.; Kenyon, V. A.; Holman, T. R. Baicalein is a potent in vitro inhibitor against both reticulocyte 15-human and platelet 12-human lipoxygenases. *Bioorg. Med. Chem.* 2006, 14, 4295–4301.
23. Ettmayer, P.; Amidon, G. L.; Clement B.; Testa, B. Lessons learned from marketed and investigational prodrugs. *J. Med. Chem.* 2004, 47, 2393-2404.
24. Fukami MH, and Salganicoff L. Human platelet storage organelles. A review. *Thrombosis and haemostasis*. 1977;38(4):963-70.
25. Fulp, A. B.; Johnson, M. S.; Markworth, J.; Marron, B. E.; Seconi, D. C.; West, C. W.; Wang, X.; Zhou, S. Sodium Channel Blockers. US Patent WO/2009/012242, Jul. 14, 2008.
26. Gan, Q.-F.; Browner, M. F.; Sloane, D. L.; Sigal, E. Defining the arachidonic acid binding site of human 15-lipoxygenase. *J. Biol. Chem.* 1996, 271, 25412-25418
27. Ghosh, J. Inhibition of Arachidonate 5-Lipoxygenase Triggers Prostate Cancer Cell Death through Rapid Activation of C-Jun N-Terminal Kinase. *Biochem. Biophys. Res. Commun.* 2003, 307, 342-349.

28. Ghosh, J., Myers, C. E. Inhibition of Arachidonate 5-Lipoxygenase Triggers Massive Apoptosis in Human Prostate Cancer Cells. *Proc. Natl. Acad. Sci. U. S. A.* 1998, 95, 13182-13187.

29. Gibbins J, Asselin J, Farndale R, Barnes M, Law CL, and Watson SP. Tyrosine phosphorylation of the Fc receptor gamma-chain in collagen-stimulated platelets. *The Journal of biological chemistry.* 1996;271(30):18095-9.

30. Greinacher A, Althaus K, Krauel K, and Selleng S. Heparin-induced thrombocytopenia. *Hamostaseologie.* 2010;30(1):17-8, 20-8.

31. Guo Y, Zhang W, Giroux C, Cai Y, Ekambaram P, Dilly AK, Hsu A, Zhou S, Maddipati KR, Liu J, et al. Identification of the orphan G protein-coupled receptor GPR31 as a receptor for 12-(S)-hydroxyeicosatetraenoic acid. *The Journal of biological chemistry.* 2011;286(39):33832-40.

32. Harats, D.; Shaish, A.; George, J.; Mulkins, M.; Kurihara, H.; Levkovitz, H.; Sigal, E. Overexpression of 15-lipoxygenase in vascular endothelium accelerates early atherosclerosis in LDL receptor-deficient mice. *Arteriosler. Thromb. Vasc. Biol.* 2000, 20, 2100-2105.

33. Holmsen H, and Weiss HJ. Secretable storage pools in platelets. *Annual review of medicine.* 1979;30(119-34).

34. Honn, K.V.; Timár, J.; Rozhin, J.; Bazaz, R.; Sameni, M.; Ziegler, G.; Sloane, B. A lipoxygenase metabolite, 12-(S)-HETE, stimulates protein kinase C-mediated release of cathepsin B from malignant cells. *Exp. Cell. Res.* 1994, 214, 120–130.

35. Hoobler, E. K.; Holz, C.; Holman, T. R. Pseudoperoxidase investigations of hydroperoxides and inhibitors with human lipoxygenases. *Bioorg. Med. Chem.* 2013, 21, 3894-3899.

36. Ikei, K. N.; Yeung, J.; Apopa, P. L.; Ceja, J.; Vesci, J.; Holman, T. R., Holinstat, M. Investigations of human platelet-type 12-Lipoxygenase:role of lipoxygenase products in platelet activation. *J. Lipid Res.* 2012, 53, 2546-2559.

37. Inglese, J.; Auld, D. S.; Jadhav, A.; Johnson, R. L.; Simeonov, A.; Yasgar, A.; Zheng, W.; Austin, C. P. Quantitative High-Throughput Screening: A Titration-Based Approach That Efficiently Identifies Biological Activities in Large Chemical Libraries. *Proc. Natl. Acad. Sci. U.S.A.* 2006, 103, 11473–11478.

38. Israels SJ, McNicol A, Robertson C, and Gerrard JM. Platelet storage pool deficiency: diagnosis in patients with prolonged bleeding times and normal platelet aggregation. *British journal of haematology.* 1990;75(1):118-21.

39. Jankun J, Aleem AM, Malgorzewicz S, Szkudlarek M, Zavodszky MI, Dewitt DL, Feig M, Selman SH, and Skrzypczak-Jankun E. Synthetic curcuminoids modulate the arachidonic acid metabolism of human platelet 12-lipoxygenase and reduce sprout formation of human endothelial cells. *Molecular cancer therapeutics*. 2006;5(5):1371-82.

40. Kamitani, H.; Geller, M.; Eling, T. The possible involvement of 15-lipoxygenase/leukocyte type 12-lipoxygenase in colorectal carcinogenesis. *Adv. Exp. Med. Biol.* 1999, 469, 593–598.

41. Kandouz M, Nie D, Pidgeon GP, Krishnamoorthy S, Maddipati KR, and Honn KV. Platelet-type 12-lipoxygenase activates NF- κ B in prostate cancer cells. *Prostaglandins & other lipid mediators*. 2003;71(3-4):189-204.

42. Kasirer-Friede A, Kahn ML, and Shattil SJ. Platelet integrins and immunoreceptors. *Immunological reviews*. 2007;218(247-64).

43. Katoh A, Ikeda H, Murohara T, Haramaki N, Ito H, and Imaizumi T. Platelet-derived 12-hydroxyeicosatetraenoic acid plays an important role in mediating canine coronary thrombosis by regulating platelet glycoprotein IIb/IIIa activation. *Circulation*. 1998;98(25):2891-8.

44. Kenyon, V.; Chorny, I.; Carvajal, W. J.; Holman, T. R.; Jacobson, M. P. Novel human lipoxygenase inhibitors discovered using virtual screening with homology models. *J. Med. Chem.* 2006, 49, 1356–1363.

45. Kenyon, V.; Rai, G.; Jadhav, A.; Schultz, L.; Armstrong, M.; Jameson, B. J.; Perry, S.; Joshi, N.; Bougie, J. M.; Leister, W.; Taylor-Fishwick D. A.; Nadler, J. L.; Holinstat, M.; Simeonov, A.; Maloney, D. J.; Holman, T. R. Discovery of Potent and Selective Inhibitors of Human Platelet-Type 12-Lipoxygenase. *J. Med. Chem.* 2011, 54, 5485-5497.

46. Lee, Y. K.; Parks, D. J.; Lu, T.; Thieu, T. V.; Markotan, T.; Pan, W.; McComsey, D. F.; Milkiewicz, K. L.; Crysler, C. S.; Ninan, N.; Abad, M. C.; Giardino, E. C.; Maryanoff, B. E.; Damiano, B. P.; Player, M. R. *J. Med. Chem.* 2008, 51, 282-297.

47. Luci DK, Jameson JB, 2nd, Yasgar A, Diaz G, Joshi N, Kantz A, Markham K, Perry S, Kuhn N, Yeung J, et al. Synthesis and structure-activity relationship studies of 4-((2-hydroxy-3-methoxybenzyl)amino)benzenesulfonamide derivatives as potent and selective inhibitors of 12-lipoxygenase. *Journal of medicinal chemistry*. 2014;57(2):495-506.

48. Ma, K.; Nunemaker, C. S.; Wu, R.; Chakrabarti, S. K.; Taylor-Fishwick, D. A.; Nadler, J. L. 12-Lipoxygenase Products Reduce Insulin Secretion and β -cell Viability in Human Islets. *J. Clin. Endocrinol. Metab.* 2010, 95, 887-893.

49. Madsen, P. et al. Optimization of Alkylidene Hydrazide Based Humna Glucagon Receptor Antagonists. Discovery of the Highly Potent and Orally Available 3-Cyano-4-hydroxybenzoic Acid [1-(2,3,5,6-Tetramethylbenzyl)-1H-indol-4-ylmethylene]hydrazide. *J. Med. Chem.* 2002, 45, 5755-5775.

50. Malterud, K. E.; Rydland, K. M. Inhibitors of 15-lipoxygenase from orange peel. *J. Agric. Food Chem.* 2000, 48, 5576-5580. (c) Moreau, R. A.; Agnew, J.; Hicks, K. B.; Powell, M. J. Modulation of lipoxygenase activity by bacterial hopanoids. *J. Nat. Prod.* 1997, 60, 397-398.

51. McDuffie, M.; Maybee, N. A.; Keller, S. R.; Stevens, B. K.; Garmey, J. C.; Morris, M. A.; Kropf, E.; Rival, C.; Ma, K.; Carter, J. D. Tersey, S. A.; Nunemaker, C. S.; Nadler, J. L. Nonobese diabetic (NOD) mice congenic for a targeted deletion of 12/15-Lipoxygenase are protected from autoimmune diabetes. *Diabetes* 2008, 57, 199-209.

52. McMahon GS, Jones CI, Hayes PD, Naylor AR, and Goodall AH. Transient heparin-induced platelet activation linked to generation of platelet 12-lipoxygenase. Findings from a randomised controlled trial. *Thrombosis and haemostasis.* 2013;109(6):1099-107.

53. McNicol A, and Israels SJ. Platelet dense granules: structure, function and implications for haemostasis. *Thrombosis research.* 1999;95(1):1-18.

54. Morgan LT, Thomas CP, Kuhn H, and O'Donnell VB. Thrombin-activated human platelets acutely generate oxidized docosahexaenoic-acid-containing phospholipids via 12-lipoxygenase. *The Biochemical journal.* 2010;431(1):141-8.

55. Moroi M, Jung SM, Okuma M, and Shinmyozu K. A patient with platelets deficient in glycoprotein VI that lack both collagen-induced aggregation and adhesion. *The Journal of clinical investigation.* 1989;84(5):1440-5.

56. Nakano, H., Inoue, T., Kawasaki, N., Miyatake, H., Matsumoto, H., Taguchi, T., Inagaki, N., Nagai, H., Satoh, T. Synthesis and Biological Activities of Novel Antiallergic Agents with 5-Lipoxygenase Inhibiting Action. *Bioorg. Med. Chem.* 2000, 8, 373-280.

57. Nappez, C.; Liagre, B.; Beneytout, J. L. Changes in lipoxygenase activities in human erythroleukemia (HEL) cells during diosgenin-induced differentiation. *Cancer Lett.* 1995, 96, 133-140.

58. Natarajan, R.; Esworthy, R.; Bai, W.; Gu, J. L.; Wilczynski, S.; Nadler, J. L. Increase 12-lipoxygenase expression in breast cancer cells and tissues. Regulation by epidermal growth factor. *J. Clin. Endocrinol. Metab.* 1997, 82, 1790-1798.

59. Nie D, Krishnamoorthy S, Jin R, Tang K, Chen Y, Qiao Y, Zacharek A, Guo Y, Milanini J, Pages G, et al. Mechanisms regulating tumor angiogenesis by 12-lipoxygenase in prostate cancer cells. *The Journal of biological chemistry*. 2006;281(27):18601-9.

60. Nie, D.; Hillman, G. G.; Geddes, T.; Tang, K.; Pierson, C.; Grignon, D. J.; Honn, K.V. Platelet-type 12-lipoxygenase in a human prostate carcinoma stimulates angiogenesis and tumor growth. *Cancer Res.* 1998, 58, 4047-4051.

61. Nie, D.; Tang, K.; Diglio, C.; Honn, K. Eicosanoid regulation of angiogenesis: role of endothelial arachidonate 12-lipoxygenase. *Blood* 2000, 95, 2304-2311.

62. Nyby, M. D.; Sasaki, M.; Ideguchi, Y.; Wynne, H. E.; Hori, M. T.; Berger, M. E.; Golub, M. S.; Brickman, A. S.; Tuck, M. L. Platelet lipoxygenase inhibitors attenuate thrombin-and thromboxane mimetic-induced intracellular calcium mobilization and platelet aggregation. *J. Pharmacol. Exp. Ther.* 1996, 278, 503-509.

63. Ohri, R. V.; Radosevich, A. T.; Hrovat, K. J.; Musich, C.; Huang, D.; Holman, T. R.; Toste, F. D. A Re(V)-catalyzed C-N bond-forming route to human lipoxygenase inhibitors, *Org. lett.* 2005, 7, 2501-2504.

64. Phillips DR, and Agin PP. Platelet membrane defects in Glanzmann's thrombasthenia. Evidence for decreased amounts of two major glycoproteins. *The Journal of clinical investigation*. 1977;60(3):535-45.

65. Pratico, D.; Zhukareva, V.; Yao, Y.; Uryu, K.; Funk, C. D.; Lawson, J. A.; Trojanowski, J. Q.; Lee, V. M. 12/15-Lipoxygenase is increased in Alzheimer's disease: possible involvement in brain oxidative stress. *Am. J. Pathol.* 2004, 164, 1655-1662.

66. Radmark, O., Samuelsson, B. 5-Lipoxygenase: Regulation and Possible Involvement in Atherosclerosis. *Prostaglandins Other Lipid Mediat.* 2007, 83, 162-174.

67. Reilly MP, Sinha U, Andre P, Taylor SM, Pak Y, Deguzman FR, Nanda N, Pandey A, Stolla M, Bergmeier W, et al. PRT-060318, a novel Syk inhibitor, prevents heparin-induced thrombocytopenia and thrombosis in a transgenic mouse model. *Blood*. 2011;117(7):2241-6.

68. Robinson, S. J.; Hoobler, E. K.; Riener, M.; Loveridge, S. T.; Tenney, K.; Valeriote, F. A.; Holman, T. R.; Crews, P. Using enzyme assays to evaluate the structure and bioactivity of sponge-derived meroterpenes. *J. Nat. Prod.* 2009, 72, 1857-1863.

69. Rosenfeld SI, Looney RJ, Leddy JP, Phipps DC, Abraham GN, and Anderson CL. Human platelet Fc receptor for immunoglobulin G. Identification as a 40,000-molecular-weight membrane protein shared by monocytes. *The Journal of clinical investigation*. 1985;76(6):2317-22.

70. Sailer, E. R.; Schweizer, S.; Boden, S. E.; Ammon, H. P. T.; Safayhi, H. Characterization of an acetyl-11-keto-B-boswellic acid and arachidonate-binding regulatory site of 5-lipoxygenase using photoaffinity labeling. *Eur. J. Biochem.* 1998, 256, 364–368.

71. Segraves, E. N.; Shah, R. R.; Segraves, N. L.; Johnson, T. A.; Whitman, S.; Sui, J. K.; Kenyon, V. A.; Cichewicz, R. H.; Crews, P.; Holman, T. R. Probing the activity differences of simple and complex brominated aryl compounds against 15-soybean, 15-human, and 12-human lipoxygenase. *J. Med. Chem.* 2004, 47, 4060–4065.

72. Serhan, C. N.; Petasis, N. A. Resolvins and Protectins in Inflammation Resolution. *Chem. Rev.* 2011, 111, 5922-5943.

73. Solomon, E. I.; Zhou, J.; Neese, F.; Pavel, E. G. New insights from spectroscopy into the structure/function relationships of lipoxygenases. *Chem. Biol.* 1997, 4, 795–808. (b) Brash, A. R. Lipoxygenases: Occurrence, Functions, Catalysis and Acquisition of Substrate. *J. Biol. Chem.* 1999, 274, 23679–23682.

74. Soriano, A. F.; Helfrich, B.; Chan, D. C.; Heasley, L. E.; Bunn, P. A.; Chou, T. C. Synergistic effects of new chemopreventive agents and conventional cytotoxic agents against human lung cancer cell lines. *Cancer Res.* 1999, 59, 6178–6184.

75. Strehl A, Munnix IC, Kuijpers MJ, van der Meijden PE, Cosemans JM, Feijge MA, Nieswandt B, and Heemskerk JW. Dual role of platelet protein kinase C in thrombus formation: stimulation of pro-aggregatory and suppression of procoagulant activity in platelets. *The Journal of biological chemistry*. 2007;282(10):7046-55.

76. Sugiyama T, Okuma M, Ushikubi F, Sensaki S, Kanaji K, and Uchino H. A novel platelet aggregating factor found in a patient with defective collagen-induced platelet aggregation and autoimmune thrombocytopenia. *Blood*. 1987;69(6):1712-20.

77. Suzuki-Inoue K, Fuller GL, Garcia A, Eble JA, Pohlmann S, Inoue O, Gartner TK, Hughan SC, Pearce AC, Laing GD, et al. A novel Syk-dependent mechanism of platelet activation by the C-type lectin receptor CLEC-2. *Blood*. 2006;107(2):542-9.

78. Suzuki-Inoue K, Inoue O, and Ozaki Y. Novel platelet activation receptor CLEC-2: from discovery to prospects. *Journal of thrombosis and haemostasis : JTH*. 2011;9 Suppl 1(44-55)

79. Takai T. Roles of Fc receptors in autoimmunity. *Nature reviews Immunology*. 2002;2(8):580-92.

80. Thomas CP, Morgan LT, Maskrey BH, Murphy RC, Kuhn H, Hazen SL, Goodall AH, Hamali HA, Collins PW, and O'Donnell VB. Phospholipid-esterified eicosanoids are generated in agonist-activated human platelets and enhance tissue factor-dependent thrombin generation. *The Journal of biological chemistry*. 2010;285(10):6891-903.

81. Thomas, C. P.; Morgan, L. T.; Maskrey, B. H.; Murphy, R. C; Kuhn, H.; Hazen, S. L.; Goodall, A. H.; Hamali, H. A.; Collins, P. W.; O'Donnell, V. D. Phospholipid-esterified eicosanoids are generated in agonist-activated human platelets and enhance tissue factor-dependant thrombin generation. *J. Biol. Chem.* 2010, 285, 6891-6903.

82. Timár, J.; Silletti, S.; Bazaz, R.; Raz, A.; Honn, K.V. Regulation of melanoma-cell motility by the lipoxygenase metabolite 12-(S)-HETE. *Int. J. Cancer*, 1993, 55, 1003–1010.

83. Tsuji M, Ezumi Y, Arai M, and Takayama H. A novel association of Fc receptor gamma-chain with glycoprotein VI and their co-expression as a collagen receptor in human platelets. *The Journal of biological chemistry*. 1997;272(38):23528-31.

84. van Leyen, K.; Arai, K.; Jin, G.; Kenyon, V.; Gerstner, B.; Rosenberg, P. A.; Holman, T. R.; Lo, E. H. Novel lipoxygenase inhibitors as neuroprotective reagents. *J. Neurosci. Res.* 2008, 86, 904-909.

85. van Leyen, K.; Kim, H. Y.; Lee, S. R.; Jin, G.; Arai, K.; Lo, E. H. Baicalein and 12/15-lipoxygenase in the ischemic brain. *Stroke*. 2006, 37, 3014-3018.

86. Vasquez-Martinez, Y.; Ohri, R. V.; Kenyon, V.; Holman, T. R.; Sepulveda-Boza, S. Structure-activity relationship studies of flavonoids as potent inhibitors of human platelet 12-hLO, reticulocyte 15-hLO-1, and prostate epithelial 15-hLO-2. *Bioorg. Med. Chem.* 2007, 15, 7408–7425.

87. Wang, X.; Guram, A.; Ronk, M.; Milne, J. E.; Tedrow, J. S.; Faul, M. M. Copper-catalyzed N-arylation of sulfonamides with aryl bromides. *Tetrahedron Lett.* 2012, 53, 7-10.

88. Watanabe N, Bodin L, Pandey M, Krause M, Coughlin S, Boussiotis VA, Ginsberg MH, and Shattil SJ. Mechanisms and consequences of agonist-induced talin recruitment to platelet integrin alphaIIbbeta3. *The Journal of cell biology*. 2008;181(7):1211-22.

89. Whitman, S.; Gezginci, M.; Timmermann, B. N.; Holman, T. R. Structure-activity relationship studies of nordihydroguaiaretic acid inhibitors toward soybean, 12-human, and 15-human lipoxygenase. *J. Med. Chem.* 2002, 45, 2659–2661.

90. Yacoub D, Theoret JF, Villeneuve L, Abou-Saleh H, Mourad W, Allen BG, and Merhi Y. Essential role of protein kinase C delta in platelet signaling, alpha IIb beta 3 activation, and thromboxane A2 release. *The Journal of biological chemistry.* 2006;281(40):30024-35.

91. Yamamoto, S. Mammalian lipoxygenases: Molecular Structures and functions. *Biochim. Biophys. Acta.* 1992, 1128, 117-131. (b) Funk, C.D.; Keeney, D.S.; Oliw, E. H.; Vogel, S.; Muller-Decker, K.; Mincheva, A.; Lichten, P.; Marks, F.; Krieg, P. Murine epidermal lipoxygenase (Aloxe) encodes a 12-lipoxygenase isoform. *FEBS Lett.* 1997, 402, 162-166.

92. Yanaga F, Poole A, Asselin J, Blake R, Schieven GL, Clark EA, Law CL, and Watson SP. Syk interacts with tyrosine-phosphorylated proteins in human platelets activated by collagen and cross-linking of the Fc gamma-IIA receptor. *The Biochemical journal.* 1995;311 (Pt 2)(471-8.


93. Yeung J, Apopa PL, Vesci J, Stolla M, Rai G, Simeonov A, Jadhav A, Fernandez-Perez P, Maloney DJ, Boutaud O, et al. 12-lipoxygenase activity plays an important role in PAR4 and GPVI-mediated platelet reactivity. *Thrombosis and haemostasis.* 2013;110(3):569-81.

94. Yeung, J.; Apopa, P. L.; Vesci, J.; Kenyon, V.; Rai, G.; Jadhav, A.; Simeonov, A.; Holman, T. R.; Maloney, D. J.; Boutaud, O.; Holinstat. Protein Kinase C Regulation of 12-Lipoxygenase-Mediated Human Platelet Activation. *Mol. Pharmacol.* 2012, 81, 420-430.

95. Zhi H, Rauova L, Hayes V, Gao C, Boylan B, Newman DK, McKenzie SE, Cooley BC, Poncz M, and Newman PJ. Cooperative integrin/ITAM signaling in platelets enhances thrombus formation in vitro and in vivo. *Blood.* 2013;121(10):1858-67.

CLAIMS


1. A compound of Formula (I)



wherein R₁ and R₂ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, F, Cl, Br, amine, nitrogen dioxide, indole, alkoxy, cycloalkyl, aryl, heterocycloalkyl, heteroaryl, each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, methoxy;

R₃ is selected from the group consisting of phenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, thiazole, benzothiazole, benzoxazole, imidazole, benzimidazole, thiophene, 1-naphthalene, 2-naphthalene, pyridine, quinoline, isoquinoline, 4N-boc-piperidine-3-phenyl, oxazole, benzothiophene, parathiazine, furan, pyran, chromene, benzofuran, pyrrole, pyrazole, pyrazine, pyrimidine, triazine, indole, purine, phthalazine,; each optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, F, Cl, Br, hydroxyl, amine, alkoxy, phenyl, cycloalkyl, aryl, piperazine, piperidine, pyridine, morpholine, pyrrolidine, pyrazolidine, imidazolidine, and thiomorpholine;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof; with the proviso that the compound is not



2. The compound of claim 1, wherein R₁ is selected from the group consisting of methoxy and Cl when R₂ is H;

R₂ is selected from the group consisting of Br, and Cl when R₁ is H; and

R₃ is selected from the group consisting of thiazole, 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, 4-methyl-2-thiazole, 5-methyl-2-thiazole, 4, 5-methyl-2-thiazole, phenyl, 1-naphthalene, 2-naphthalene, 1, 4-bi-phenyl, 3-piperazine-phenyl, 4-piperidine-phenyl, 4-piperazine-3-pyridine, 6-methyl-3-pyridine, 3-quinoline, 8-isoquinoline, 2-pyridine, 3-pyridine, 3-tertbutyl-phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, 4-phenyl-2-thiazole, 3-morpholine-phenyl, 4N-boc-piperidine-3-phenyl, 3-piperidine-phenyl, 3-isopropyl-phenyl, and 4-bi-phenyl;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

3. The compound of claim 2, wherein R₁ is methoxy and R₂ is H.

4. The compound of claim 1, wherein the compound is a pro-drug.

5. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, 3-quinoline, 8-isoquinoline, phenyl, and 3-isopropyl-phenyl.

6. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-

naphthalene, 3-quinoline, 8-isoquinoline, 3-tertbutyl-phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, 4-phenyl-2-thiazole, 4N-boc-piperidine-3-phenyl, and 3-isopropyl-phenyl.

7. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-naphthalene, 3-quinoline, 8-isoquinoline, 3-tertbutyl-phenyl, 4N-boc-piperidine-3-phenyl, and 3-isopropyl-phenyl.


8. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, and 3-isopropyl-phenyl.

9. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 6-methoxy-2-benzothiazole, 6-fluro-2-benzothiazole, and 4-phenyl-2-thiazole.

10. The compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, and phenyl.


11. A compound of claim 3, wherein R₃ is selected from the group consisting of 2-benzothiazole, 2-benzoxazole, 2-benzimidazole, 4-methyl-2-benzothiazole, thiophene, phenyl, 1-naphthalene, 2-naphthalene, 3-quinoline, 8-isoquinoline, 2-pyridine, and 3-pyridine.

12. A compound that is




or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

13. A compound that is




or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

14. A compound that is



or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

15. A compound of formula (II)




wherein X is selected from the group consisting of O, S, NH, and C;

wherein R₁ and R₂ are independently selected from the group consisting of H, halogen, hydroxyl, alkoxy, and alkyl;

wherein R₃ through R₆ are independently selected from the group consisting of H, halogen, alkoxy, and alkyl;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.


16. A compound of formula (III)




wherein R₁ and R₂ are independently selected from the group consisting of H, halogen, and alkoxy, and further wherein R₁ and R₂ are not both H;

wherein R₄ and R₅ are independently selected from the group consisting of H, alkyl, phenyl, and optionally substituted phenyl;

or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof; with the proviso that the compound is not




17. A compound of formula (IV)




wherein R₁ and R₂ are independently selected from the group consisting of H, halogen, and alkoxy;

wherein R₃ and R₄ are independently selected from the group consisting of H, phenyl,



optionally substituted phenyl, tert-butyl, isopropyl, and

wherein X is selected from the group consisting of NH, O, and



or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

18. A method for treating or preventing a 12-lipoxygenase mediated disease or disorder, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of a compound of claim 6, salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

19. The method of claim 18, wherein the 12-lipoxygenase is human the 12-lipoxygenase.

20. The method of claim 18, wherein the 12-lipoxygenase mediated disease or disorder is selected from the group consisting of type 1 diabetes, type 2 diabetes, diabetic kidney disease, diabetic nerve disease, cardiovascular disease, Alzheimer's disease, Non-Alcoholic steatohepatitis, platelet hemostasis, skin diseases, heparin induced thrombocytopenia, thrombosis, anti-phospholipid syndrome; sepsis syndrome; thrombosis associated with therapeutic or diagnostic monoclonal antibodies; and thrombotic thrombocytopenic purpura, and cancer.


21. The method of claim 20, wherein the cancer is selected from the group consisting of prostate cancer, colorectal cancer, breast cancer, and lung cancer.

22. The method of claim 20, wherein the cardiovascular disease is selected from the group consisting of congestive heart failure, myocardial infarction and stroke.

23. The method of claim 20, wherein the 12-lipoxygenase mediated disease or disorder is selected from the group consisting of type 1 and type 2 diabetes.


24. The method of claim 20, wherein the 12-lipoxygenase mediated disease or disorder is selected from the group consisting of thrombosis, heparin-induced thrombocytopenia; anti-phospholipid syndrome; sepsis syndrome; thrombosis associated with therapeutic or diagnostic monoclonal antibodies; and thrombotic thrombocytopenic purpura.

25. The method of claim 18, wherein the compound is




or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

26. The method of claim 20, wherein the compound is




or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

27. The method of claim 20, wherein the compound is



or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

28. The method of claim 20, wherein the compound is




or a pharmaceutically acceptable salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.


29. A method for reducing PAR4-AP induced platelet aggregation, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of a compound of claim 6, salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.

30. A method for reducing PAR4-AP induced calcium mobilization, comprising administering to a mammal thereof a therapeutically or prophylactically effective amount of a compound of claim 6, salt thereof, enantiomers thereof, a mixture of enantiomers thereof, or diastereomers thereof.


Scheme 1. Synthesis of analogs 1-83.



Figures 1A – 1D



Figures 2A - 2B



Figures 3A – 3E




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8







Figure 9

Figures 10A – 10E







Figures 11A-11B




Figures 12A-12C




Figures 13A-13B



Figures 14A-14B



Figures 15A-15B



Figures 16A-16C





Figure 17



Figures 18A – 18B





Figure 19



Figures 20A-20B




Figure 21




Figure 22A-22B




Figure 23

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/60174

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/045, 31/10, 31/18; C12Q 1/26 (2014.01)

CPC - C07D 239/42, 263/50, 277/52, 417/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8)- A61K 31/00, 31/045, 31/10, 31/18, 31/137, 31/506; C07D 209/14, 239/42; C12Q 1/26 (2014.01); CPC - C07D 239/42, 263/50, 277/52, 401/12, 403/12, 417/12, 417/14; 435/7.4, 25; 436/63, 71:514/158, 415, 555, 562, 576, 866

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 435/7.4, 25; 436/63, 71:514/158, 415, 555, 562, 576, 866

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MicroPatent (US-G, US-A, EP-A, EP-B, WO, JP-bib, DE-C,B, DE-A, DE-T, DE-U, GB-A, FR-A); Google Scholar; PubMed; PubChem; IP.com; ProQuest; SureChem; 12-lipoxygenase inhibitors, 12-LOX, benzenesulfonamide, cancer, diabetes, thrombosis, Alzheimer's, PAR4-AP, protease-activated receptor-4, 2-benzothiazole, 2-benzimidazole, phenyl, thiophene, cardiovascular, methoxybenzylamino

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---	PUBCHEM, CID 70701426. February 22, 2013, pp 1-10 [online], [retrieved on 2014-12-10] Retrieved from the internet <URL https://pubchem.ncbi.nlm.nih.gov/compound/70701426#section=Top ; page 3, structure	1-3, 5-12, 15
X	PUBCHEM, CID 70701432. February 22, 2013, pp 1-9 [online], [retrieved on 2014-12-10] Retrieved from the internet <URL https://pubchem.ncbi.nlm.nih.gov/compound/70701432#section=2D-Structure ; page 3, structure	13
X	PUBCHEM, CID 70701440. February 22, 2013, pp 1-8 [online], [retrieved on 2014-12-10] Retrieved from the internet <URL https://pubchem.ncbi.nlm.nih.gov/compound/1079729#section=Top ; page 2, structure	16
X	PUBCHEM, CID 70701358. February 22, 2013, pp 1-8 [online], [retrieved on 2014-12-10] Retrieved from the internet <URL https://pubchem.ncbi.nlm.nih.gov/compound/1079729#section=Top ; page 2, structure	17
A	US 5,571,821 A (CHAN, MF et al.) November 05, 1996; table 1; column 13, lines 33-35; column 16, lines 3-6	4, 14
A	US 6,767,909 B2 (JOUTSAMO, T et al.) July 27, 2004; column 3, line 65 through column 5, line 28	14
A	US 2006/0173075 A1 (KRAUSS, NE et al.) August 03, 2006; pages 5-7	4, 14

Further documents are listed in the continuation of Box C.



• Special categories of cited documents:	
“A” document defining the general state of the art which is not considered to be of particular relevance	“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
“E” earlier application or patent but published on or after the international filing date	“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
“O” document referring to an oral disclosure, use, exhibition or other means	“&” document member of the same patent family
“P” document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 December 2014 (16.12.2014)

Date of mailing of the international search report

31 DEC 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450
Facsimile No. 571-273-3201

Authorized officer:

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/60174

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2011/0077305 A1 (JACKSON, WP) March 31, 2011; paragraphs [0021], [0102]-[0125]	18-28
A	US 2013/0096159 A1 (MALONEY, DJ et al.) April 18, 2013; paragraphs [0015], [0081]	18-30
A	US 2013/0131140 A1 (NADLER, JL et al.) May 23, 2013; claim 1; paragraph [0012]	18-28
A	WO 1999/043809 A2 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) September 02, 1999; page 8, lines 13-16; page 17, lines 5-6; page 25, lines 10-15	29-30
A	WO 2013/050441 A1 (INSERM) April 11, 2013; page 20, lines 14-16; page 22, lines 6-7; page 26, lines 13-20	29-30