

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Purification of polypeptides

(51)⁶ International Patent Classification(s)
C12P 21/02 (2006.01) 20060101AFI2005100
C12P 21/02 8BMEP
PCT/US2004/000499

(21) Application No: 2004230670 (22) Application Date: 2004.01.08

(87) WIPO No: WO04/092393

(30) Priority Data

(31) Number (32) Date (33) Country
60/439,418 2003.01.09 US

(43) Publication Date : 2004.10.28

(71) Applicant(s)
Genentech, Inc.

(72) Inventor(s)
Persson, Josefina, Lester, Philip M.

(74) Agent/Attorney
Griffith Hack, Level 3 509 St Kilda Road, Melbourne, VIC, 3004

(56) Related Art
US 3903254 A
Tchernov et al. Journal of Biotechnology 1999, Vol. 69 No. 1, pages 69-73
JP 60-258127 A
US 5861263 A
Azegami et al. Microbios 1978, Vol. 23 No. 92, pages 73-82

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
28 October 2004 (28.10.2004)

PCT

(10) International Publication Number
WO 2004/092393 A1

(51) International Patent Classification⁷: C12P 21/02 (81) Designated States (unless otherwise indicated, for every kind of national protection available): AF, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CI, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GI, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SI, SG, SK, SI, SY, TI, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/US2004/000499

(22) International Filing Date: 8 January 2004 (08.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/439,418 9 January 2003 (09.01.2003) US

(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LESTER, Philip, M. [US/US]; 15766 Via Colusa, San Lorenzo, CA 94580 (US). PERSSON, Josefine [SE/SE]; Östervägen 16A, S-169 52 Solna (SE).


(74) Agent: HASAK, Janet E., C/O Genentech, Inc., MS 49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PURIFICATION OF POLYPEPTIDES



(57) Abstract: A method for purifying a desired heterologous polypeptide from microbial fermentation broth or homogenate in which it is produced and solubilized is described. This method involves adding to the broth or homogenate an effective amount of a solution of 6,9-diamino-2-ethoxyacridine lactate (ethacridine lactate) to precipitate host cell impurities under conditions wherein the majority of polypeptide remains soluble, and separating the desired polypeptide from the broth or homogenate. The broth or homogenate containing the ethacridine lactate and polypeptide is also disclosed.

WO 2004/092393 A1

PURIFICATION OF POLYPEPTIDES

5

Background of the Invention1. Field of the Invention

This invention relates to a process for purifying polypeptides of interest from microbial fermentation broth or homogenate. More particularly, a precipitation agent is introduced to the broth or homogenate to effect, for example, protein, DNA, and cell debris removal.

10 2. Description of Related Art

The advent of recombinant technology now allows for the production of high levels of proteins within suitably transformed host cells. As a result, there is increased demand for fast, robust, and efficient purification methods to recover the recombinantly produced proteins. Generally, proteins are produced by 15 culturing cells, such as mammalian, insect, fungal, and bacterial cell lines, engineered to produce the protein of interest by insertion of a recombinant plasmid containing the gene for that protein. Since the cell lines used are living organisms, they must be fed with a complex growth medium, containing sugars, amino acids, and growth factors, usually supplied from preparations of animal serum. Separation of the desired protein from the mixture of compounds fed to the cells and from the by-products of the cells themselves to a purity 20 sufficient for use as a human therapeutic poses a formidable challenge.

Procedures for purification of proteins from cell debris initially depend on the site of expression of the protein. Some proteins can be secreted directly from the cell into the surrounding growth media; others are made intracellularly. For polypeptides produced in mammalian cells, the purification scheme is significantly easier than for polypeptides produced in other types of host cells. Mammalian cells export the 25 polypeptides so that they can be collected from the growth media, where they are present in relatively pure form. However, if the polypeptide is produced in a non-mammalian cell, *e.g.*, a microorganism such as fungi or *E. coli*, the polypeptide will be recovered inside the cell or in the periplasmic space (Kipriyanov and Little, *Molecular Biotechnology*, 12: 173-201 (1999); Skerra and Pluckthum, *Science*, 240: 1038-1040 (1988)). Hence, it is necessary to release the protein from the cells to the extracellular medium by extraction 30 such as cell lysis. Such disruption releases the entire contents of the cell into the homogenate, and in addition produces subcellular fragments that are difficult to remove due to their small size. These are generally removed by differential centrifugation or by filtration.

Cell lysis is typically accomplished using mechanical disruption techniques such as homogenization or bead milling. While the protein of interest is generally effectively liberated, such techniques have several 35 disadvantages (Engler, *Protein Purification Process Engineering*, Harrison eds., 37-55 (1994)). Temperature increases, which often occur during processing, may result in inactivation of the protein. Moreover, the resulting suspension contains a broad spectrum of contaminating proteins, nucleic acids, and polysaccharides. Nucleic acids and polysaccharides increase solution viscosity, potentially complicating subsequent processing by centrifugation, cross-flow filtration, or chromatography. Complex associations of 40 these contaminants with the protein of interest can complicate the purification process and result in unacceptably low yields.

As such, more selective means of releasing intracellular proteins facilitates further downstream processing. Several techniques have been reported to permeabilize cells and/or to extract intracellular proteins. These methods include the use of solvents, detergents, chaotropic agents, antibiotics, enzymes, and chelating agents to enhance cell permeability and/or promote extraction. Additions of certain compounds, such as glycine, to the fermentation medium during culture growth have also been reported to promote release of certain intracellular enzymes. Finally, techniques such as freeze-thaw treatment or osmotic shock have also been shown to release subsets of intracellular proteins.

However, these techniques are not necessarily applicable to all intracellular microbial proteins, and all have limited application for large-scale processing, and/or other disadvantages. For example, while solvents such as toluene and chloroform promote release of intracellular proteins, these substances are known to be toxic and/or carcinogenic (Windholtz *et al.*, The Merck Index 10th Edition: 300 and 1364 (1983)). Ionic detergents, such as SDS, often irreversibly denature isolated proteins. Although non-ionic detergents are not normally denaturing, the recovered proteins are often associated with detergent micelles that can require additional processing to yield detergent-free protein. Chaotropic agents, such as urea and guanidine hydrochloride, can be denaturing at the concentrations required for complete release, and their effectiveness may be dependent on the growth phase of the culture. The use of lysozyme, which provides for a relatively gentle means of protein release, is limited because of its relatively high cost and because of the subsequent need to purify the protein of interest from the enzyme reagent. In addition, chelating agents, often used to enhance the effectiveness of other permeabilizing/release techniques such as lysozyme or toluene extraction, suffer from the disadvantage of non-specific release of host proteins.

Other methods for protein release also have disadvantages. For example, osmotic shock, in which cells are suspended in a high osmolarity medium, recovered, and subsequently placed in a low osmolarity buffer, requires additional processing steps with respect to other extraction alternatives (Moir *et al.*, Separation Processes in Biotechnology, Ascenjo eds: 67-94 (1990)) or necessitates the handling of large liquid volumes at low temperatures. This renders the method unattractive for large-scale processing.

Freeze-thaw treatment also releases intracellular proteins, although relatively low yields often result in multiple cycles or additional processing requirements. In addition, cell paste freezing is an added non-trivial processing requirement compared with other extraction alternatives.

Finally, reagents, such as glycine, have been added during fermentation to promote protein release to the extracellular medium (Aristidou *et al.*, Biotechnology Letters 15: 331-336 (1993)). While partial release of several intracellular proteins has been reported, this approach requires direct coupling of fermentation and release strategies and subsequent separation of the protein of interest from a potentially complex extracellular broth.

Once the polypeptide of interest is released from the host cell, purification thereof from other cell components is required. Unfortunately, most extraction approaches, such as cell lysis, not only expose the protein to potential degradation by host cell proteases, but also make isolation of the protein from other elements of the resulting suspension more difficult. For example, the presence of negatively charged molecules, such as DNA, RNA, phospholipids, and lipopolysaccharides (LPS), often requires the use of anion-exchange chromatography (Sassenfeld, TIBTECH, 8: 88-93 (1990); Spars, Biotechnology, vol. 3-Bioprocessing, Rehm eds:40-51 (1993)) and/or precipitation with polycations, such as protamine sulfate

(Kelley *et al.*, Bioseparation, 1: 333-349 (1991); Scopes, Protein Purification Principles and Practice, 2nd edition, Cantor eds., pp. 21-71 (1987)), streptomycin sulfate (Wang *et al.*, eds, Fermentation and Enzyme Technology, 253-256 (1979)), polyethylenimine (PEI) (Kelley *et al.*, *supra*; Sassenfeld, *supra*; Cumming *et al.*, Bioseparation, 6: 17-23 (1996); Jendrisak, The use of polyethylenimine in protein purification, Protein purification: micro to macro, ed. Alan R., Liss, Inc, 75-97 (1987); Salt *et al.*, Enzyme and Microbial Technology, 17: 107-113 (1995)), and/or aqueous two-phase extraction with immiscible polymer systems such as polyethylene glycol (PEG)/phosphate or PEG/dextran (Kelley *et al.*, *supra*, Strandberg *et al.*, Process Biochemistry 26: 225-234 (1991)).

Alternatively, the protein of interest may be precipitated away from non-proteinaceous polyanionic contaminants through the addition of a neutral salt such as ammonium sulfate or potassium chloride (Wheelwright, Protein Purification: Design and Scale up of Downstream Processing: 87-98 (1991); Englund *et al.*, Methods in Enzymology, Volume 182, Deutscher eds.: 285-300 (1990)) and/or a polymer such as PEG or dextran sulfate (Wang *et al.*, *supra*; Wheelwright, *supra*). Where the protein of interest is positively charged, it will tend to bind to any negatively charged molecules present thereby, making purification of the protein virtually impossible.

Typically, researchers have utilized the initial fractionation steps, described above, to separate the offending polyanions from the protein of interest. Unfortunately, each of these initial separation methods suffers from severe disadvantages, especially when used in the manufacture of pharmaceutical reagents. For example, the large quantities of non-proteinaceous polyanionic contaminants found in bacterial lysates tend to reduce the binding capacities of anion-exchange chromatography resins. In addition, regeneration protocols are often rendered ineffective due to tenacious binding of the polyanions to the resins (Spears, *supra*). Finally, the low ionic strength conditions that favor protein binding are ineffective at disrupting polyanion-protein interactions and result in a lack of separation (Scopes, Protein Purification Principles and Practice, 3rd edition, Cantor eds., p. 171 (1994)). Protamine sulfate preparations are plagued by concerns over protease and viral contaminations. Moreover, unwanted protein precipitation can occur using this reagent (Scopes, Protein Purification Principles and Practice, 2nd edition, Cantor eds., 21-71 (1987)).

In the processing of pharmaceutical proteins, streptomycin sulfate is generally not used due to general apprehension over the use of antibiotics as process reagents (Scawen *et al.*, Handbook of Enzyme Biotechnology 2nd edition, Wiseman eds.: 15-53 (1985)). PEI preparations are often contaminated with varying amounts of the ethylenimine monomer, a suspected cancer agent (Scawen *et al.*, *supra*). PEI also tends to bind irreversibly to many chromatography resins, thereby limiting their effectiveness and the number of potential chromatography resins available for use post-PEI clarification. In general, aqueous two-phase extractions systems are difficult to predict and often require an empirical approach for determining conditions that move the protein of interest into the appropriate aqueous phase (Kelley *et al.*, *supra*).

Techniques that specifically precipitate the protein of interest often result in the entrapment of the non-proteinaceous contaminants in the precipitate, rendering the separation ineffective (Scopes, *supra*; Wheelwright, *supra*).

Examples of patents describing protein recovery and purification include the following:

U.S. Pat. No. 5,665,866 discloses a process for obtaining antibodies in soluble and correctly folded and assembled form. It comprises a step to raise the operating temperature to from 34 to 60°C at a time in

the process selected to facilitate the subsequent isolation of soluble, correctly folded and assembled antibody, substantially free of other antibody-related material.

U.S. Pat. No. 5,760,189 discloses a method for releasing a thioredoxin-like fusion protein from *E. coli*, including negatively charged non-proteinaceous material, into a solution by adding chelator to the solution, and precipitating the negatively charged non-proteinaceous material from the solution by adding a divalent cation/alcohol solution to the solution to form a first soluble fraction containing the protein and a first insoluble fraction containing unwanted contaminants. Optionally, the temperature prior to the addition of chelator may be substantially cooler than after the addition of chelator. The divalent cation includes, for example, magnesium, manganese, and calcium, alone or in combination.

US Pat. No. 5,714,583 discloses methods for the purification of factor IX in a solution comprising the steps of applying the solution containing factor IX to an anion-exchange resin, washing the anion-exchange resin with a solution having a conductivity that is less than required to elute factor IX from the resin, eluting the anion-exchange resin with a first eluant to form a first eluate, applying the eluate to a heparin or heparin-like (e.g., negatively charged matrix) resin, eluting the heparin or heparin-like resin with a second eluant to form a second eluate, applying the second eluate to an hydroxyapatite resin, and then eluting the hydroxyapatite resin with a third eluant to form a third eluate containing the purified factor IX.

US Pat. No. 6,322,997 discloses a method for recovering a polypeptide comprising exposing a composition comprising a polypeptide to a reagent that binds to, or modifies, the polypeptide, wherein the reagent is immobilized on a solid phase; and then passing the composition through a filter bearing a charge that is opposite to the charge of the reagent in the composition, so as to remove leached reagent from the composition.

US Pat. No. 6,214,984 discloses low-pH hydrophobic interaction chromatography (LPHIC) for antibody purification. In particular, the patent provides a process for purifying an antibody from a contaminant that comprises loading a mixture containing the antibody and the contaminant on a hydrophobic interaction chromatography column and eluting the antibody from the column with a buffer having a pH of about 2.5-4.5. Usually, the mixture loaded onto the column is at about the same pH as the elution buffer.

US Pat. No. 6,121,428 provides a method for recovering a polypeptide comprising exposing a composition comprising a polypeptide to a reagent that binds to, or modifies, the polypeptide, wherein the reagent is immobilized on a solid phase; and then passing the composition through a filter bearing a charge that is opposite to the charge of the reagent in the composition, so as to remove leached reagent from the composition.

US Pat. No. 5,641,870 provides a process for purifying an antibody is provided, wherein a mixture containing the antibody and contaminant is subjected to LPHIC optionally at low salt concentration. The antibody is eluted from the column in the fraction that does not bind thereto. In the extraction step, frozen cell pellets are re-suspended at room temperature in 20 mM MES buffer, pH 6.0 containing 5 mM EDTA and 20 mM 4,4'-DTP previously dissolved in ethanol (3 liters of buffer/kg of cell pellet). The suspended cells are disrupted by two passages through a Manton Gaulin homogenizer at 5500 to 6500 PSI. The homogenate is adjusted to 0.25% (v/v) with polyethyleneimine (PEI) and diluted with an equal volume of 2-8°C. purified water. The diluted homogenate is then centrifuged. The antibody fragment is found in the supernatant.

2004230670 28 Oct 2010

Historically, immunoglobulin G (IgG) has been purified from human serum and plasma (Putnam, ed., *The Plasma Proteins*, vol. 1 (Academic Press, 1975)). The purification process has often contained one or more precipitation steps. The most commonly used precipitation scheme for recovering IgG is the Cohn fractionation (Cohn *et al.*, *J. Amer. Chem. Soc.*, **72**: 465 (1950)). However, other precipitation techniques have been reported (Niederauer and Glatz, *Advances in Biochemical Engineering Biotechnology*, v. 47 (Springer-Verlag Berlin Heidelberg, 1992); Steinberg and Hershberger, *Biochim. et Biophys. Acta*, **342**: 195-206 (1974)). The pioneering work of purifying IgG from plasma using 6,9-diamino-2-ethoxyacridine lactate (USAN name and herein called ethacridine lactate and also known by the names ETHODIN™ or RIVANOL™), a highly aromatic cationic dye, is reported by Horsjisi and Smetana, *Acta Med. Scand.*, **155**: 65 (1956). The following decade produced a number of publications showing the capability of 6,9-diamino-2-ethoxyacridine lactate to purify IgG and other proteins (Miller, *Nature*, **184**: 450 (1959); Steinbuch and Niewiarowski, *Nature*, **186**: 87 (1960); Neurath and Brunner, *Experientia*, **25**: 668 (1969)) from biological materials, *e.g.*, plasma and growth media. Use of ethacridine lactate to recover antibodies and other proteins from other sources has been reported. See Tchemrov *et al.*, *J. Biotechnol.*, **69**: 69-73 (1999); SU 944580 published 28 July 1982; Franek and Dolnikova, *Biotech-Forum-Eur.*, **7**: 468-470 (1990); EP 250288 published 23 Dec. 1987; DE3604947 published 20 August 1987; Rothwell *et al.*, *Anal. Biochem.*, **149**: 197-201 (1985); Lutsik and Antonyuk, *Biokhimiya*, **47**: 1710-1715 (1982); and Aizenman *et al.*, *Mikrobiol-Zh.*, **44**: 69-72 (1982).

The primary step of recovering polypeptides from microorganisms is most often concerned with removing solid material, *e.g.*, cells and cellular debris. It is important to recognize the need to separate the desired product from components present in conditioned medium with which it specifically interacts. Where the protein of interest is positively charged, it will tend to bind to any negatively charged molecules present, thereby making purification of the protein by traditional methods very difficult. Additional removal of contaminating soluble protein from crude microbial extracts, *e.g.*, *E. coli* homogenate, during this step would simplify subsequent chromatography steps. Such additional removal would be especially valuable for industrial-scale production, resulting in decreased chromatography column size and production times.

All references, including any patents or patent applications, cited in this specification are hereby incorporated by reference. No admission is made that any reference constitutes prior art. The discussion of the references states what their authors assert, and the applicants reserve the right to challenge the accuracy and pertinency of the cited documents. It will be clearly understood that, although a number of prior art publications are referred to herein, this reference does not constitute an admission that any of these documents forms part of the common general knowledge in the art, in Australia or in any other country.

Summary of the Invention

35 The invention involving purification is as claimed.

Specifically, in one aspect, the invention provides a method for purifying a desired heterologous polypeptide from microbial fermentation broth or homogenate in which it is produced and solubilized comprising adding to the broth or homogenate an effective amount of a solution of 6,9-diamino-2-ethoxyacridine lactate (ethacridine lactate) to precipitate host cell impurities under conditions wherein the

2004230670 28 Oct 2010

majority of the polypeptide remains soluble, and separating the desired polypeptide from the broth or homogenate, wherein the temperature of the broth or homogenate after addition of the ethacridine lactate is from about 50 to 65°C held for about 1 to 60 minutes.

5 A second aspect provides a microbial cell fermentation broth or homogenate comprising ethacridine lactate of greater than 0.1% (weight/volume) and a polypeptide heterologous to the cells, wherein the pI of the polypeptide is greater than the average pI of the host proteins contained in the cells, wherein the broth or homogenate is at a temperature of from about 50 to 65°C, and wherein the polypeptide is an antibody or antibody fragment.

10 Addition of ethacridine lactate as a precipitation agent unexpectedly results in a dramatic removal of host debris including host proteins. In this process, the majority of host proteins will be recovered in a precipitate together with the cell debris, and the polypeptide is recovered in the clarified supernatant. The

improved purity of the clarified extract when using ethacridine lactate results in reduction in the volume of chromatographic media or resin required for the columns, thereby reducing the scale needed for subsequent purification. It also results in elimination of some chromatographic step(s), which improves processing time and cost. In addition, the process herein results in a stable feedstock and can be operated at a neutral pH.

5

Brief Description of the Drawings

Figure 1 is a schematic representation of the construction of the antiCD18 F(ab')₂(-leucine zipper) plasmids pS1130 (single promoter) and pxCd18-7T3 (dual-promoter).

10 Figure 2 depicts the inserted nucleic acid sequence (designated as Anti-CD18-7T3.DNA; SEQ ID NO:1) of the dual-promoter construct pxCd18-7T3.

Figures 3A and 3B depict the amino acid sequences (designated in combination as Anti-CD18-7T3.Protein) encoded by the two translational units within the construct pxCd18-7T3 (SEQ ID NOS:2 and 3), designated as STII + Anti-CD18 light chain (Fig. 3A) and STII + Anti-CD18 heavy chain (Fig. 3B), respectively. N-terminal STII secretion signal sequences are underlined.

15 Figure 4 is a schematic of the anti-Tissue Factor IgG1 plasmids paTF130 (*phoA/phoA* promoters) and pxtF-7T3FL (*phoA/tacII*-promoters).

Figure 5 depicts the inserted nucleic acid sequence (designated as Anti-TF-7T3FL.DNA; SEQ ID NO:4) of the *phoA/tacII*-promoter construct pxtF-7T3FL.

20 Figures 6A and 6B depict the amino acid sequences (designated in combination as Anti-TF-7T3FL.Protein) encoded by the two translational units within the construct pxtF-7T3FL (SEQ ID NOS:5 and 6) designated as STII + Anti-TF light chain (Fig. 6A) and STII + Anti-TF heavy chain (Fig. 6B), respectively. N-terminal STII secretion signal sequences are underlined.

Figure 7 depicts the chemical structure of ethacridine lactate.

25 Figures 8A-8C show a non-reduced SDS-PAGE Coomassie-blue-stained gel analysis of three supernatants after precipitation with ethacridine lactate. The precipitation was performed at different pH values, as indicated at each lane. The lanes indicated with an X are the clarified supernatant of the respective *E. coli* homogenate, *i.e.*, anti-CD18 F(ab')₂, anti-TF F(ab')₂, and full-length anti-TF (Figs. 8A, 8B, and 8C, respectively). The homogenates were diluted 4-fold with a 0.8% ethacridine lactate solution, *i.e.*, a final concentration of 0.6% ethacridine lactate in each experiment. All samples were volume compensated before loading on to the gel. Hence, the intensity of the bands should be comparable to the extract (X) if a 100% recovery is obtained. The arrows indicate product band.

30 Figures 9A-9C show a non-reduced SDS-PAGE Coomassie-blue-stained gel analysis of supernatants after precipitation with ethacridine lactate. The precipitation was performed with different ethacridine lactate concentrations, as indicated at each lane. The lanes indicated with an X are the clarified supernatant of the respective *E. coli* homogenate, *i.e.*, anti-CD18F(ab')₂, anti-TF F(ab')₂, and full-length anti-TF (Figs. 9A, 9B, and 9C, respectively). The pH of the anti-CD18 F(ab')₂, anti-TF F(ab')₂, and full-length anti-TF was 8.5, 7.5, and 6.0, respectively. The conductivity in the samples was 3.2 ± 0.2 mS/cm. All samples were volume compensated before loading on to the gel. Hence, the intensity of the bands should be comparable to the extract (X) if a 100% recovery is obtained. The arrows indicate product band.

Figures 10A and 10B show a non-reduced SDS-PAGE Coomassie-blue-stained gel analysis of two supernatants after dilution with water or with ethacridine lactate, respectively. The precipitation was performed at different conductivity levels. *E. coli* homogenate containing anti-CD18 F(ab')₂ was used for this study. The homogenate was diluted 4-fold with either water (Fig. 10A) or a 0.8% ethacridine lactate solution, *i.e.*, a final ethacridine lactate concentration of 0.6% in each experiment (Fig. 10B), and the pH was adjusted to 8.3. To alter the conductivity, NaCl was added to the samples in different concentrations of 0-400 mM (as indicated in the figures). The arrows indicate product band.

Figure 11 shows a graph of solubility of ethacridine lactate at increasing sodium chloride concentrations. The samples were incubated for three hours at room temperature before the concentration of soluble ethacridine lactate was determined. The open keys symbolize the 1.2% ethacridine lactate solution and the closed keys the 0.6% solution. The solid line is the 0.6% ethacridine lactate solution at pH 6.0, the broken line is the 1.2% ethacridine lactate solution at pH 6, the dotted line is the 0.6% ethacridine lactate solution at pH 9, and the broken line with dots is the 1.2% ethacridine lactate solution at pH 9.

Figures 12A-12C show a non-reduced SDS-PAGE Coomassie-blue-stained gel analysis of three supernatants after precipitation with ethacridine lactate. The precipitation was performed at elevated temperatures. The lanes indicated with an X are the clarified supernatant of the respective *E. coli* homogenate, *i.e.*, anti-CD18 (F(ab')₂), anti-TF (F(ab')₂), and full-length anti-TF (Figs. 12A, 12B, and 12C, respectively). The homogenate was diluted 4-fold to a final ethacridine lactate concentration of 0.6% and the pH was adjusted to 8.5, 7.5, and 6.0 for anti-CD18 (F(ab')₂), anti-TF (F(ab')₂), and full-length anti-TF, respectively. The temperature and time for incubation are indicated in the figures. The arrows indicate product band.

Figure 13 shows a graph of turbidity as a function of time for three different supernatants. Supernatants from anti-CD18 homogenate treated with 0.6% ethacridine lactate are symbolized by solid circles (4°C) or open circles (21°C), and 0.2% PEI-treated sample is shown as solid squares (4°C) and open squares (21°C). A supernatant recovered from a clarified anti-CD18 homogenate that had been diluted with water before concentration is shown as solid triangles (4°C) and open triangles (21°C). In all cases the anti-CD18 homogenate had been diluted 4-fold and the pH was 7.2.

Detailed Description of the Preferred Embodiments

30 Definitions

In the claims which follow and in the description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, *i.e.* to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

The expression "microbial fermentation broth or homogenate" refers to broth, paste, or extract, preferably resuspended, obtained from microorganisms, including yeast, fungi, and prokaryotes such as bacteria, that are cultured and consuming nutrients, no matter what culturing vessel is utilized, for example, a shake flask or fermentor. Preferably, the broth or homogenate is from yeast or prokaryotes. More preferably, the broth or homogenate is from bacteria. Homogenate is preferred herein. In some cases if the

28 Oct 2010

2004230670

5 solution has very high conductivity it may be preferred to harvest the cells and re-suspend them, but otherwise, it is preferred to use the homogenate as it is directly from the fermentor. The components of the broth or homogenate include cell debris, host cell protein, DNA, RNA, etc. Thus, the addition of the lactate herein leads to selective precipitation of host cell proteins, etc., giving better purification power than not using the lactate.

The expression "under conditions wherein the majority of the polypeptide remains soluble" refers to addition of ethacridine lactate to the broth or homogenate in amounts and at a temperature and conductivity level that prevent the majority of the target polypeptide from precipitating from the broth or homogenate. Preferably, such conditions prevent more than about 60% of the polypeptide from precipitating, more

5 preferably more than about 70%, still more preferably more than about 75%, even more preferably more than about 80%, even still more preferably more than about 85%, and still more preferably more than about 85%, even still more preferably more than about 90%, and most preferably more than about 95%. This degree of solubility is measured by an appropriate assay, such as, for example, RP-HPLC, affinity chromatography, ELISAs, RIAs, and a combination of SDS-PAGE and high-performance affinity

10 chromatography (HPAC). The choice of assay depends on such factors as the type of host cell used and polypeptide being produced.

The "bacteria" for purposes herein include eubacteria and archaeabacteria. Preferred of these are eubacteria, including gram-positive and gram-negative bacteria. More preferred are gram-negative bacteria. One preferred type of bacteria is *Enterobacteriaceae*. Examples of bacteria belonging to *Enterobacteriaceae* 15 include *Escherichia*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, *Serratia*, and *Shigella*. Other types of suitable bacteria include *Azotobacter*, *Pseudomonas*, *Rhizobia*, *Vitreoscilla*, and *Paracoccus*. *E. coli* is preferred herein. Suitable *E. coli* hosts include *E. coli* W3110 (ATCC 27,325), *E. coli* 294 (ATCC 31,446), *E. coli* B, and *E. coli* X1776 (ATCC 31,537). These examples are illustrative rather than limiting, and W3110 is preferred. Mutant cells of any of the above-mentioned bacteria may also be employed. It is, 20 of course, necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E. coli*, *Serratia*, or *Salmonella* species can be suitably used as the host when well-known plasmids such as pBR322, pBR325, pACYC177, or pKN410 are used to supply the replicon. See further below regarding examples of suitable bacterial host cells.

As used herein, the expressions "cell," "cell line," "strain," and "cell culture" are used 25 interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it 30 will be clear from the context.

As used herein, "polypeptide" refers generally to peptides and proteins from any cell source having 35 more than about ten amino acids. "Heterologous" polypeptides are those polypeptides foreign to the host cell being utilized, such as a human protein produced by *E. coli*. While the heterologous polypeptide may be prokaryotic or eukaryotic, preferably it is eukaryotic, more preferably mammalian, and most preferably human. Preferably, it is a recombinantly produced, or recombinant polypeptide.

The polypeptide is produced and solubilized in the fermentation broth or homogenate, meaning that 40 it is made in such broth or homogenate and is either already in a soluble fraction resulting from production, or is in an insoluble fraction or form or phase that is treated or contacted with a solubilizing agent such as a chaotropic (e.g., urea or guanidine) or detergent (such as sodium dodecyl sulfate (SDS)), with or without a reducing agent (such as dithiothreitol or beta-mercaptoethanol) so as to be solubilized. "Soluble,"

"solubilized," "solubilization," "dissolved," or "dissolution" in the sense used herein means that the polypeptide is in the supernatant rather than in the solids fraction after centrifugation. Precipitation or degree of solubility can be determined, for example, by the appropriate assays as noted above.

Examples of mammalian polypeptides include molecules such as, e.g., renin, a growth hormone, including human growth hormone; bovine growth hormone; growth hormone releasing factor; parathyroid hormone; thyroid stimulating hormone; lipoproteins; 1-antitrypsin; insulin A-chain; insulin B-chain; proinsulin; thrombopoietin; follicle stimulating hormone; calcitonin; luteinizing hormone; glucagon; clotting factors such as factor VIIIIC, factor IX, tissue factor, and von Willebrands factor; anti-clotting factors such as Protein C; atrial natriuretic factor; lung surfactant; a plasminogen activator, such as urokinase or human urine or tissue-type plasminogen activator (t-PA) and variants thereof such as RETEVATM and TNKASETM; bombesin; thrombin; hemopoietic growth factor; tumor necrosis factor-alpha and -beta; antibodies to ErbB2 domain(s) such as 2C4 (WO 01/00245; hybridoma ATCC HB-12697), which binds to a region in the extracellular domain of ErbB2 (e.g., any one or more residues in the region from about residue 22 to about residue 584 of ErbB2, inclusive), enkephalinase; a serum albumin such as human serum albumin; Muellerian-inhibiting substance; relaxin A-chain; relaxin B-chain; prorelaxin; mouse gonadotropin-associated peptide; a microbial protein, such as beta-lactamase; DNase; inhibin; activin; vascular endothelial growth factor (VEGF); receptors for hormones or growth factors; integrin; protein A or D; rheumatoid factors; a neurotrophic factor such as brain-derived neurotrophic factor (BDNF), neurotrophin-3, -4, -5, or -6 (NT-3, NT-4, NT-5, or NT-6), or a nerve growth factor such as NGF; cardiotrophins (cardiac hypertrophy factor) such as cardiotrophin-1 (CT-1); platelet-derived growth factor (PDGF); fibroblast growth factor such as aFGF and bFGF; epidermal growth factor (EGF); transforming growth factor (TGF) such as TGF-alpha and TGF-beta, including TGF-1, TGF-2, TGF-3, TGF-4, or TGF-5; insulin-like growth factor-I and -II (IGF-I and IGF-II); des(1-3)-IGF-I (brain IGF-I), insulin-like growth factor binding proteins; CD proteins such as CD-3, CD-4, CD-8, and CD-19; erythropoietin; osteoinductive factors; immunotoxins; a bone morphogenetic protein (BMP); an interferon such as interferon-alpha, -beta, and -gamma; serum albumin, such as human serum albumin (HSA) or bovine serum albumin (BSA); colony stimulating factors (CSFs), e.g., M-CSF, GM-CSF, and G-CSF; interleukins (ILs), e.g., IL-1 to IL-10; anti-HER-2 antibody; Apo2 ligand; superoxide dismutase; T-cell receptors; surface membrane proteins; decay accelerating factor; viral antigen such as, for example, a portion of the AIDS envelope; transport proteins; homing receptors; addressing; regulatory proteins; antibodies; and fragments of any of the above-listed polypeptides.

Preferred polypeptides herein include human serum albumin (HSA), 2C4, tissue factor, anti-tissue factor, anti-CD20, anti-HER-2, heregulin, anti-IgE, anti-CD11a, anti-CD18, VEGF and receptors and antibodies thereto such as rhuFab V2 and AVASTINTM, growth hormone and its variants, such as hGH, growth hormone receptors, growth hormone releasing protein (GHRP), LIV-1 (EP 1,263,780), TRAIL, tumor necrosis factor (TNF) and antibodies thereto, TNF receptor and related antibodies, TNF-receptor-IgG, TNF receptor associated factors (TRAFs) and inhibitors thereof, Factor VIII, Factor VIII B domain, interferons such as interferon-gamma, transforming growth factors (TGFs) such as TGF-beta, anti-TGF such as anti-TGF-beta, activin, inhibin, anti-activin, anti-inhibin, tissue-plasminogen activators and their variants such as t-PA, RETEPLASETM, and TNKase, anti-Fas antibodies, Apo-2 ligand; Apo-2 ligand inhibitor; Apo-2 receptor, Apo-3, apoptotic factors, Ced-4, DcR3, death receptor and agonist antibodies (DR4, DR5),

lymphotoxin (LT), prolactin, prolactin receptor, SOB proteins, WISP (wnt-induced secreted proteins), neurotoxin-3 (NT-3), nerve growth factor (NGF) and anti-NGF, DNase, hepatitis antigen, herpes simplex antigen, leptin, insulin-like growth factors (IGFs) such as IGF-1 and IGF-2 and their binding proteins and receptors such as IGFBP-1-IGFBP-6, insulin, fibroblast growth factors (FGFs) such as FGF-17, Toll protein, 5 TIE ligands, CD40 and anti-CD40, immunoadhesins, subtilisin, hepatocyte growth factor (HGF), thrombopoietin (TPO), interleukins such as IL-2, IL-12, IL-17, IL-22, IL-8, IL-9, and antibodies thereto, and prostate-specific cancer antigen (PSCA).

10 Examples of antibodies that bind HER2 include 4D5, 7C2, 7F3 and 2C4, as well as humanized variants thereof, including huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and huMAb4D5-8 as described in Table 3 of U.S. Patent 5,821,337; and humanized 2C4 mutant nos. 560, 561, 562, 568, 569, 570, 571, 574, or 56869 as described in WO01/00245. 7C2 and 7F3 and humanized variants thereof are described in WO98/17797.

15 Examples of antibodies that bind the CD20 antigen include: "C2B8" which is now called "Rituximab" ("RITUXAN®") (US Patent No. 5,736,137); the yttrium-[90]-labeled 2B8 murine antibody designated "Y2B8" (US Patent No. 5,736,137); murine IgG2a "B1" optionally labeled with ^{131}I to generate the " ^{131}I -B1" antibody (BEXXAR™) (US Patent No. 5,595,721); murine monoclonal antibody "1F5" (Press *et al.*, *Blood*, 69(2): 584-591 (1987)); "chimeric 2H7" antibody (US Patent No. 5,677,180); and monoclonal antibodies L27, G28-2, 93-1B3, B-C1 or NU-B2 available from the International Leukocyte Typing Workshop (Valentine *et al.*, In: *Leukocyte Typing III* (McMichael, Ed., p. 440, Oxford University Press 20 (1987)).

20 More preferred polypeptides are 2C4, anti-tissue factor, anti-CD20, anti-HER-2, heregulin, anti-IgE, anti-CD11a, anti-CD18, anti-VEGF such as rhuFab V2, hGH, GHRP, LIV-1, TRAIL, antibodies to TNF and TNF receptor and related antibodies, inhibitors of TRAF, TNF-receptor-IgG, Factor VIII, Factor VIII B domain, interferon-gamma, TGF-beta and anti-TGF-beta, activin, inhibin, anti-activin, anti-inhibin, t-PA, TNKase, anti-Fas antibodies, Apo-2 ligand; Apo-2 ligand inhibitor; Apo-2 receptor, Apo-3, DcR3, death receptor and agonist antibodies (DR4, DR5), lymphotoxin (LT), prolactin prolactin receptor, WISP, anti-NGF, NGF, NT-3, anti-IL-8, anti-IL-9, IL-17, IL-22, DNase, GHRP, hepatitis antigen, herpes simplex antigen, leptin, IGF-1 and IGFBP1-6, insulin, FGF-17, Toll protein, TIE ligands, CD40, immunoadhesins, subtilisin, HGF, and TPO.

25 30 Still more preferred polypeptides are 2C4, anti-tissue factor, anti-CD20, anti-HER-2, anti-IgE, anti-CD11a, anti-CD18, anti-VEGF such as rhuFab V2, hGH, LIV-1, TRAIL, antibodies to TNF and TNF receptor and related antibodies, TNF-receptor-IgG, Factor VIII, Factor VIII B domain, interferon-gamma, TGF-beta, activin, inhibin, anti-activin, anti-inhibin, t-PA, TNKase, Apo-2 ligand; Apo-2 ligand inhibitor; Apo-2 receptor, Apo-3, DcR3, death receptor and agonist antibodies (DR4, DR5), WISP, inhibitors of 35 TRAF, anti-NGF, NGF, NT-3, anti-IL-8, anti-IL-9, IL-17, IL-22, anti-TGFs, DNase, GHRP, hepatitis antigen, herpes simplex antigen, leptin, IGF-1 and IGFBP1-6, insulin, FGF-17, Toll protein, TIE ligands, anti-CD40, HGF, and TPO.

40 Particularly preferred polypeptides are recombinant polypeptides, more preferably antibodies, which include monoclonal antibodies and humanized antibodies. Such antibodies may be full-length antibodies or antibody fragments. More preferably, these antibodies are human or humanized antibodies.

These include, *e.g.*, the particularly preferred polypeptides 2C4, anti-tissue factor Fab'2 and full-length, anti-CD20, anti-HER-2, anti-IgE, anti-CD11a, anti-CD18 Fab'2 and full-length, anti-VEGF full-length and rhuFab V2, L1V-1, DR4, DR5, and TRAIL.

Still more preferably, the antibody is an anti-IgE, anti-CD18, anti-VEGF, anti-tissue factor, 2C4, 5 anti-Her-2, anti-CD20, anti-CD40, or anti-CD11a antibody. Antibody fragments encompassed within the definition of polypeptide preferably comprise a light chain, more preferably a kappa light chain. Such preferred fragments include, for example, a Fab, Fab', F(ab')₂, or F(ab')₂-leucine zipper (LZ) fusion, and most preferably are F(ab')₂. The most preferred antibodies are anti-CD18 F(ab')₂, anti-tissue factor F(ab')₂, full-length anti-tissue factor antibody, and anti-VEGF antibody.

10 The term "antibody" herein is used in the broadest sense and specifically covers intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.* bispecific antibodies) formed from at least two intact antibodies, and antibody fragments, so long as they exhibit the desired biological activity.

15 The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations that include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized

20 uncontaminated by other antibodies. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Koehler *et al.*, *Nature*, **256**: 495 (1975), or may be made by recombinant DNA methods (*see, e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, **352**: 624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991), for example.

30 The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, **81**: 6851-6855 (1984)). Chimeric antibodies of interest herein include "primatized" 35 antibodies comprising variable-domain antigen-binding sequences derived from a non-human primate (*e.g.* Old World Monkey, Ape etc) and human constant-region sequences.

40 "Antibody fragments" comprise a portion of an intact antibody, preferably comprising the antigen-binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragment(s).

An "intact" antibody is one that comprises an antigen-binding variable region as well as a light chain constant domain (C_L) and heavy chain constant domains, C_H1 , C_H2 and C_H3 . The constant domains may be native sequence constant domains (e.g. human native-sequence constant domains) or amino acid sequence variant thereof. Preferably, the intact antibody has one or more effector functions.

5 Antibody "effector functions" refer to those biological activities attributable to the Fc region (a native-sequence Fc region or Fc region with amino acid sequence variation) of an antibody. Examples of antibody effector functions include C1q binding, complement dependent cytotoxicity, Fc receptor binding, antibody-dependent cell-mediated cytotoxicity (ADCC), phagocytosis, down-regulation of cell-surface receptors (e.g. B cell receptor; BCR), etc.

10 Depending on the amino acid sequence of the constant domain of their heavy chains, intact antibodies can be assigned to different "classes". There are five classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, having heavy chains designated α , δ , ϵ , γ , and μ , respectively. The γ and α classes are further divided into subclasses on the basis of relatively minor differences in C_H sequence and function, e.g., humans express the following subclasses: IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2. The 15 subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

20 "Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express FcRIII only, whereas monocytes express FcRI, FcRII and FcRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.*, 9: 457-492 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear 25 cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes *et al.* *Proc. Natl. Acad. Sci. USA*, 95: 652-656 (1998).

30 "Human effector cells" are leukocytes that express one or more FcRs and perform effector functions. Preferably, the cells express at least FcRIII and perform ADCC effector function. Examples of human leukocytes that mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells, and neutrophils, with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, e.g., from blood or PBMCs as described herein.

35 "Native antibodies" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the

light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light-chain and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a beta-sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming, part of the beta-sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular cytotoxicity (ADCC).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody that are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a "complementarity-determining region" or "CDR" (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light-chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy-chain variable domain; Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light-chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy-chain variable domain; Chothia and Lesk, *J. Mol. Biol.*, **196**: 901-917 (1987)). "Framework Region" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy-chain and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxyl terminus of the heavy-chain CH1 domain including one or more cysteines from the antibody hinge

region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

5 The "light chains" of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

10 "Single-chain Fv" or "scFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the scFv to form the desired structure for antigen binding. For a review of scFv, see Plückthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenburg and Moore eds. (Springer-Verlag, New York, 1994), pp. 269-315. Anti-ErbB2 antibody scFv fragments are described in WO93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458.

15 The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a variable heavy domain (V_H) connected to a variable light domain (V_L) in the same polypeptide chain (V_H - V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, **90**: 6444-6448 (1993).

20 "Humanized" forms of non-human (*e.g.*, rodent) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody), such as 25 mouse, rat, rabbit, or non-human primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and 30 typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin, and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, **321**: 522-525 (1986); Riechmann *et al.*, Nature, **332**: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., **2**: 593-596 (1992).

35 An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody 40 will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most

2004230670 28 Oct 2010

preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells, since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

A "leucine zipper" is a peptide (often about 20-40 amino acid residues long) having several repeating amino acids, in which every seventh amino acid is a leucine residue. Such leucine zipper sequences form amphipathic alpha-helices, with the leucine residues lined up on the hydrophobic side for dimer formation. Examples of leucine zippers herein include the Fos-Jun leucine zipper (O'Shea *et al.*, *Science*, **245**: 646 (1989)), which may be used for forming heterodimers (e.g., bispecific antibodies); the GCN4 leucine zipper from yeast (Landschulz *et al.*, *Science*, **240**: 1759-1764 (1988)) which may be used for forming homodimers (e.g., monospecific antibodies); and leucine zippers found in other DNA-binding proteins, such as C/EBP and c-myc, as well as variants of any of these.

The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for bacteria include a promoter, optionally an operator sequence, and a ribosome-binding site.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. Linking is accomplished, for example, by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "recovery" of a polypeptide generally means obtaining the polypeptide free from the cells in which it was produced.

"Host cell impurities" means contaminating host proteins and other biomolecular impurities such as DNA and cell debris in the fermentation broth or homogenate.

Modes for Carrying Out the Invention

Disclosed herein is a method for purifying a desired heterologous polypeptide from a microbial fermentation broth or homogenate in which it is produced and solubilized. The polypeptide may already be produced in a soluble fraction, or it may be insoluble (e.g., produced in an insoluble fraction, phase, or form) and therefore contacted or treated so as to dissolve the polypeptide. If the polypeptide is produced in an insoluble state, it is solubilized by exposure to or contact with a solubilizing agent (as noted above) before the ethacridine lactate is added, for example, adding such agent to a fraction containing the insoluble polypeptide. It is preferred that the polypeptide is already in the soluble fraction. The method herein involves adding to the broth or homogenate an effective amount of a solution of ethacridine lactate to precipitate host cell impurities contained in the broth or homogenate. Such addition takes place under

conditions wherein the majority of the polypeptide remains soluble. In the next step the desired polypeptide is separated from the broth or homogenate, including cell debris, host cell protein, DNA, RNA, etc.

Because most host proteins precipitated by the ethacridine lactate have a negative charge while the target polypeptide has a positive surface charge, it is preferred that the target polypeptide have a higher pI than the average pI of host proteins contained in the host cell impurities, so that it can be recovered in the supernatant from the precipitated host proteins. Such average pI can be determined by a 2-D gel of the host proteins, wherein, for example, the band of pI ranges from about 7.5 to 5.0, the average being 6.25. Alternatively, isoelectrofocusing alone (which is the first dimension on a 2-D gel) can be used for this determination, as well as chromatofocusing, and calculations by amino acid composition. The more preferred polypeptides are those having a pI of at least about 7, and preferably about 7-10.

The preferred polypeptides to be employed are set forth above.

The concentration of ethacridine lactate employed is dependent, for example, on the amount of negative charges in solution, which are on the surface of most host cell impurities present in the solution. Hence, the ethacridine lactate concentration depends at least on the amount of host cell impurities such as the DNA and host protein concentration in the solution. The higher the concentration of host protein and DNA in the homogenate, the higher the amount of ethacridine lactate required. Hence, the more negatively charged components available for the ethacridine lactate to complex with and thus precipitate, the higher amount of ethacridine lactate needed to maximize precipitation. The preferred concentration of ethacridine lactate is generally more than about 0.1% weight/volume. More preferred is an ethacridine lactate concentration of about 0.1-5%, still more preferred about 0.4-5%, and most preferably about 0.6-5% weight/volume.

In general, the lower the conductivity of the solution when performing the precipitation, the more efficient the purification of the polypeptide from the cell debris and DNA. The conductivity can be controlled, for example, by the amount of salt in the homogenate or broth or by diluting the homogenate or broth with water or other suitable solvent. It is preferred that the conductivity of the broth or homogenate after addition of the ethacridine lactate is less than about 16 millisiemens per centimetre (mS/cm), more preferably about 1-15 mS/cm, still more preferably about 1-10 mS/cm, and most preferably about 1-5 mS/cm.

The conductivity of the solution during the precipitation will depend at least in part on the type of salt present therein. Halides (e.g., chlorides or bromides) are not preferred anions for the salts, but if they are present, they are preferably at a concentration of less than about 100 mM in solution before adding the ethacridine lactate, and below about 50 mM once the ethacridine lactate is added. Some exemplary salts to employ herein include buffer salts, TRIS, MES, MOPS, acetate, and citrate. The concentration of salts present must not be above an amount that would precipitate the ethacridine lactate. The exact amount is dependent mainly on the type of salt and the stoichiometry between the salt and ethacridine lactate, and the limit is at the low end of the stoichiometry, i.e., the low end means more salt relative to the ethacridine lactate.

The pH of the broth or homogenate after addition of the ethacridine lactate depends, for example, on the pI of the polypeptide, the amount of negative surface charges on the polypeptide, the amount of host cell impurities in the solution, and the concentration of ethacridine lactate. The pH is preferably not higher

28 Oct 2010

2004230670

than the pI of the polypeptide. Generally, the pH range is about 4-10; however, for efficient host cell impurity precipitation, the pH of the broth or homogenate after addition of the ethacridine lactate is preferably no greater than about 9, since the ethacridine lactate becomes less charged above this pH, with the preferred range of about 4-9. More preferably the pH of the broth or homogenate after addition of the ethacridine lactate is about 5-9, and still more preferably, about 6-9. The more negative surface charges on the polypeptide, the lower the pH within this range, with a preferred range for such polypeptides of about pH 6-7.

The broth or homogenate after addition of the ethacridine lactate optionally is incubated at an elevated temperature for a period of time. Whether to raise the temperature and for how long depends on many factors, including the type of polypeptide of interest, what, if any, modifications to the polypeptide of interest occur if exposed to elevated temperature during this period in the process, etc. For example, for the anti-tissue factor F(ab')₂ purifications, elevated temperatures are preferred, whereas for full-length antibody, it is preferred to have no heat or have the temperature no higher than about 25°C. With these factors in mind, in general, the temperature of the broth or homogenate after addition of the ethacridine lactate ranges from about room temperature to about 70°C, more preferably, from about room temperature to about 65°C held for about 1-60 minutes. If the temperature should be elevated, one preferred range is from about 50 to 65°C held for about 1-60 minutes.

Disclosed herein is a composition of matter that is a fermentation broth or homogenate from microbial cells comprising ethacridine lactate and heterologous polypeptide. Preferably, the polypeptide is dissolved in such broth or homogenate. The cells, polypeptide, concentration, and conditions for the broth or homogenate are as indicated above. The dissolution degree of the polypeptide can be determined by an appropriate assay such as the assays as noted above. Culturing parameters are used and polypeptide production is conducted in a conventional manner, such as those procedures described below.

A. Selection of Nucleic Acid and Modifications Thereof

While the polypeptide herein, such as an antibody, may be produced from any source (e.g., peptic cleavage of intact antibodies), preferably it is made recombinantly. The nucleic acid encoding the polypeptide of interest is suitably RNA, cDNA, or genomic DNA from any source, provided it encodes the polypeptide(s) of interest. Methods are well known for selecting the appropriate nucleic acid for expression of heterologous polypeptides (including variants thereof) in microbial hosts. Selection of appropriate nucleic acid to prepare non-antibody polypeptides in microbial cell culture is well known in the art.

If monoclonal antibodies are being produced, DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transformed into the microbial host cells herein to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5: 256-262 (1993) and Plückthun, *Immunol. Revs.*, 130: 151-188 (1992).

Methods for humanizing non-human antibodies have been described in the art. Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human.

These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, **321**: 522-525 (1986); Rechmann *et al.*, *Nature*, **332**: 323-327 (1988); Verhoeyen *et al.*, *Science*, **239**: 1534-1536 (1988)), by substituting hypervariable region sequences

5 for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

10 The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence that is closest to that of the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, **151**: 2296 (1993);

15 Chothia *et al.*, *J. Mol. Biol.*, **196**: 901 (1987)). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, **89**: 4285 (1992); Presta *et al.*, *J. Immunol.*, **151**: 2623 (1993)).

20 It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available that illustrate and display probable three-dimensional conformational structures of 25 selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

30 Various forms of the humanized antibody or affinity-matured antibody are contemplated. For example, the humanized antibody or affinity-matured antibody may be an antibody fragment, such as a Fab, that is optionally conjugated with one or more targeting agent(s) in order to generate an immunoconjugate. Alternatively, the humanized antibody or affinity-matured antibody may be an intact antibody, such as an intact IgG1 antibody.

35 Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology*, **10**: 163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice 40 is a single-chain Fv fragment (scFv) (WO 93/16185; U.S. Pat. Nos. 5,571,894 and 5,587,458). The antibody

fragment may also be a "linear antibody", e.g., as described in U.S. Pat. No. 5,641,870. Such linear antibody fragments may be monospecific or bispecific.

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the Dkk-1 protein.

5 Bispecific antibodies can be prepared as full-length antibodies or antibody fragments (e.g., F(ab'), bispecific antibodies).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, 10 CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable bacterial host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when 15 unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid 20 immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this

asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690.

25 For further details of generating bispecific antibodies see, for example, Suresh *et al.*, Methods in Enzymology, 121: 210 (1986).

According to another approach described in U.S. Pat. No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody

30 constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products 35 such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies

may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed, e.g., in U.S. Pat. No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, 5 *Science*, **229**: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies (Shalaby *et al.*, *J. Exp. Med.*, **175**: 217-225 (1992)).

Various techniques for making and isolating bispecific antibody fragments directly from 15 recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers (Kostelnik *et al.*, *J. Immunol.*, **148**: 1547-1553 (1992)). The leucine zipper peptides from the Fos and Jun proteins are linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers are reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The 20 "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**: 6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker that is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby 25 forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported (Gruber *et al.*, *J. Immunol.*, **152**: 5368 (1994)).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared (Tutt *et al.*, *J. Immunol.*, **147**: 60 (1991)).

Nucleic acid molecules encoding polypeptide variants are prepared by a variety of methods known 30 in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, or cassette mutagenesis of an earlier prepared variant or a non-variant version of the polypeptide.

It may be desirable to modify the antibody disclosed herein with respect to effector function, e.g., 35 so as to enhance Fc receptor binding. This may be achieved by introducing one or more amino acid substitutions into an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region.

To increase the serum half-life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Pat. 5,739,277, for example.

40 As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG

molecule (e.g., IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

Other modifications of the antibody are contemplated herein. For example, the antibody may be linked to one of a variety of non-proteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol.

B. Insertion of Nucleic Acid Into a Replicable Vector

The heterologous nucleic acid (e.g., cDNA or genomic DNA) is suitably inserted into a replicable vector for expression in the microorganism under the control of a suitable promoter. Many vectors are available for this purpose, and selection of the appropriate vector will depend mainly on the size of the nucleic acid to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on the particular host cell with which it is compatible. Depending on the particular type of host, the vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, a promoter, and a transcription termination sequence.

In general, plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell are used in connection with microbial hosts. The vector ordinarily carries a replication site, as well as marking sequences that are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species (see, e.g., Bolivar *et al.*, *Gene*, 2: 95 (1977)). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, or other bacterial plasmid or phage, also generally contains, or is modified to contain, promoters that can be used by the host for expression of the selectable marker genes.

(i) Signal Sequence Component

The DNA encoding the polypeptide of interest herein may be expressed not only directly, but also as a fusion with another polypeptide, preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the polypeptide DNA that is inserted into the vector. The heterologous signal sequence selected should be one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell.

For prokaryotic host cells that do not recognize and process the native or a eukaryotic polypeptide signal sequence, the signal sequence may be substituted by a prokaryotic signal sequence selected, for example, from the group consisting of the *lamB*, *ompF*, alkaline phosphatase, penicillinase, *lpp*, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990.

(ii) Origin of Replication Component

Expression vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of microbes. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria such as *E. coli*.

(iii) Selection Gene Component

Expression vectors generally contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. This selectable marker is separate from the genetic markers as utilized and defined by this invention. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies other than those caused by the presence of the genetic marker(s), or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacillus*.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. In this case, those cells that are successfully transformed with the nucleic acid of interest produce a polypeptide conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin (Southern *et al.*, *J. Molec. Appl. Genet.*, **1**: 327 (1982)), mycophenolic acid (Mulligan *et al.*, *Science*, **209**: 1422 (1980)) or hygromycin (Sugden *et al.*, *Mol. Cell. Biol.*, **5**: 410-413 (1985)). The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid), or hygromycin, respectively.

(iv) Promoter Component

The expression vector for producing the polypeptide of interest contains a suitable promoter that is recognized by the host organism and is operably linked to the nucleic acid encoding the polypeptide of interest. Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems (Chang *et al.*, *Nature*, **275**: 615 (1978); Goeddel *et al.*, *Nature*, **281**: 544 (1979)), the arabinose promoter system (Guzman *et al.*, *J. Bacteriol.*, **174**: 7716-7728 (1992)), alkaline phosphatase, a tryptophan (*tsp*) promoter system (Goeddel, *Nucleic Acids Res.*, **8**: 4057 (1980) and EP 36,776) and hybrid promoters such as the *tac* promoter (deBoer *et al.*, *Proc. Natl. Acad. Sci. USA*, **80**: 21-25 (1983)). However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to DNA encoding the polypeptide of interest (Siebenlist *et al.*, *Cell*, **20**: 269 (1980)) using linkers or adaptors to supply any required restriction sites.

Promoters for use in bacterial systems also generally contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the polypeptide of interest. The promoter can be removed from the bacterial source DNA by restriction enzyme digestion and inserted into the vector containing the desired DNA.

Promoters suitable for use in yeast are well known in the art. Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, *J. Biol. Chem.*, **255**: 2073 (1980)) or other glycolytic enzymes (Hess *et al.*, *J. Adv. Enzyme Reg.*, **7**: 149 (1968); Holland, *Biochemistry*, **17**: 4900 (1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucoisomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2,

isocytchrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

5 (v) Construction and Analysis of Vectors

Construction of suitable vectors containing one or more of the above-listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and re-ligated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to 10 transform *E. coli* K12 strain 294 (ATCC 31,446) or other strains, and successful transformants are selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced by the method of Sanger *et al.*, *Proc. Natl. Acad. Sci. USA*, 74: 5463-5467 (1977) or Messing *et al.*, *Nucleic Acids Res.*, 9: 309 (1981), or by the method of Maxam *et al.*, *Methods in Enzymology*, 65: 499 (1980).

15 C. Selection and Transformation of Host Cells

Suitable host cells for cloning or expressing the DNA in the vectors herein are any microbial cells, including prokaryotes and fungal cells, including yeast. Suitable prokaryotes for this purpose include bacteria as defined above, preferably eubacteria, such as Gram-negative or Gram-positive organisms.

Examples include *Enterobacteriaceae* such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, 20 *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescens*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting. 25 Mutant cells of any of the above-mentioned strains may also be employed as the starting hosts that are then further mutated to contain at least the minimum genotype required herein.

E. coli strain W3110 is a preferred parental *E. coli* host because it is a common host strain for recombinant DNA product fermentations. Examples of starting *E. coli* hosts to be used as parent hosts, along with their genotypes, are included in the table below:

30

Strain	Genotype
W3110	K-12 F lambda ⁺ IN(rrnD-rrnE)1
1A2	W3110 Δ <i>huA</i>
9E4	W3110 Δ <i>huA</i> <i>ptr3</i>
27A7	W3110 Δ <i>huA</i> <i>ptr3</i> <i>phoAΔE15</i> Δ(<i>argF-lac</i>)169
27C6	W3110 Δ <i>huA</i> <i>ptr3</i> <i>phoAΔE15</i> Δ(<i>argF-lac</i>)169 Δ <ompt></ompt>
27C7	W3110 Δ <i>huA</i> <i>ptr3</i> <i>phoAΔE15</i> Δ(<i>argF-lac</i>)169 Δ <ompt>degP4I::kan^R</ompt>
33D3	W3110 Δ <i>huA</i> <i>ptr3</i> <i>lacIq</i> <i>lacL8</i> Δ <ompt>degP4I::kan^R</ompt>

36F8	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $ptr3$ $degP4I::kan^R$ $ilvG2096$
41H1	W3110 $\Delta fhuA$ $phoS^*$ $(T104)$ $\Delta(argF-lac)169$ $degP4I::kan^R$ $ptr3$ $ilvG2096(Vaf^r)$
43D3	W3110 $\Delta fhuA$ $ptr3$ $phoA\Delta E15$ $\Delta(argF-lac)169$ \DeltaompT $degP4I$ kan^R $ilvG2096$
43H1	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $degP4I$ $ilvG2096(Vaf^r)$ $ptr3$ \DeltaompT $prc::kan^R$ $sprW148R$
43E7	W3110 $\Delta fhuA$ $\Delta(argF-lac)169$ \DeltaompT $ptr3$ $phoA\Delta E15$ $degP4I$ $ilvG2096$
44D6	W3110 $\Delta fhuA$ $ptr3$ $\Delta(argF-lac)169$ $degP4I::kan^R$ \DeltaompT $ilvG2096$
45F8	W3110 $\Delta fhuA$ $ptr3$ $\Delta(argF-lac)169$ $degP4I$ \DeltaompT $phoS^*$ $(T10Y)$ $ilvG2096$
45F9	W3110 $\Delta fhuA$ $ptr3$ $\Delta(argF-lac)169$ $degP4I$ \DeltaompT $ilvG2096$ $phoS^*$ $(T10Y)$ $\Delta cyo::kan^R$
49A5	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $deoC2$ $degP4I$ $ilvG2096(Vaf^r)$ $\Delta fucP$ $\Delta malE$
58B3	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $deoC$ $degP4I$ $ilvG2096(Vaf^r)$ Δprc
58H2	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $degP4I$ $ilvG2096(Vaf^r)$ $ptr3$ \DeltaompT $sprW148R$
58H7	W3110 $\Delta fhuA$ $(\Delta tonA)$ $\Delta ptr3$ \DeltaompT $\Delta degP$ $lacIq$ $\Delta lacY$
59A7	W3110 $\Delta fhuA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $deoC$ $degP4I$ $ilvG2096(Vaf^r)$ Δprc $sprW148R$
60H4	W3110 $\Delta fhuA$ $\Delta manA$ $phoA\Delta E15$ $\Delta(argF-lac)169$ $deoC2$ $degP4I$ $ilvG2096(Vaf^r)$ Δprc $prc-suppressor$
61D6	W3110 $\Delta fhuA$ $ptr3$ $lacIq$ $lacL8$ \DeltaompT $(nmprc-fepE)$ $degP4I$
62A7	W3110 $\Delta fhuA$ $ptr3$ $lacIq$ $lacL8$ \DeltaompT $(nmprc-fepE)$ $degP4I$ $ilvG2096$

Also suitable are the intermediates in making strain 36F8, *i.e.*, 27B4 (U.S. Pat. No. 5,304,472) and 35E7 (a spontaneous temperature-resistant colony isolate growing better than 27B4). An additional suitable strain is the *E. coli* strain having the mutant periplasmic protease(s) disclosed in U.S. Pat. No. 4,946,783 issued

5 August 7, 1990.

The above strains may be produced by chromosomal integration of the parental strain or other techniques, including those set forth in the Examples below.

Full-length antibodies may be made in *E. coli* in accordance with the teachings of WO 02/061090 published August 8, 2002.

10 In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable expression hosts for polypeptide-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, **290**: 140 (1981); EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer *et al.*, *Bio/Technology*, **9**: 968-975 (1991)) such as, *e.g.*, *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt *et al.*, *J. Bacteriol.*, **73** (1983)), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickeramii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906; Van den Berg *et al.*, *Bio/Technology*, **8**: 135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna *et al.*, *J. Basic Microbiol.*, **28**: 265-278 (1988)); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case *et al.*, *Proc. Natl. Acad. Sci. USA*, **76**: 5259-5263 (1979)); 20 *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and

2004230670 28 Oct 2010

filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, *Biochem. Biophys. Res. Commun.*, **112**: 284-289 (1983); Tilburn *et al.*, *Gene*, **26**: 205-221 (1983); Yelton *et al.*, *Proc. Natl. Acad. Sci. USA*, **81**: 1470-1474 (1984)) and *A. niger* (Kelly and Hynes, *EMBO J.*, **4**: 475-479 (1985)).

5 Methylotrophic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylotrophs*, 269 (1982).

10 The nucleic acid encoding the polypeptide is inserted into the host cells. Preferably, this is accomplished by transforming the host cells with the above-described expression vectors and culturing in conventional nutrient media modified as appropriate for inducing the various promoters.

15 Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989), is generally used for prokaryotic cells or other cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO, as described in Chung and Miller, *Nucleic Acids Res.*, **16**: 3580 (1988). Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, *J. Bact.*, **130**:946 (1977) and Hsiao *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **76**:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, 20 bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used.

D. Culturing the Host Cells

25 Prokaryotic cells used to produce the polypeptides disclosed herein are grown in media known in the art and suitable for culture of the selected host cells, including the media generally described by Sambrook *et al.*, *supra*. Media that are suitable for bacteria include, but are not limited to, AP5 medium, nutrient broth, Luria-Bertani (LB) broth, Neidhardt's minimal medium, and C.R.A.P. minimal or complete medium, plus necessary nutrient supplements. In preferred embodiments, the media also contains a selection agent, chosen based on the construction of the expression vector, to selectively permit growth of prokaryotic cells containing the expression vector. For example, ampicillin is added to media for growth of cells expressing ampicillin resistant gene. Any necessary supplements besides carbon, nitrogen, and inorganic phosphate sources may also be included at appropriate concentrations introduced alone or as a mixture with another supplement or medium such as a complex nitrogen source. Optionally the culture medium may contain one or more reducing agents selected from the group consisting of glutathione, cysteine, cystamine, thioglycollate, dithioerythritol, and dithiothreitol.

30 Examples of suitable media are given in U.S. Pat. Nos. 5,304,472 and 5,342,763. C.R.A.P. phosphate-limiting media consists of 3.57g (NH₄)₂(SO₄), 0.71g NaCitrate-2H₂O, 1.07g KCl, 5.36g Yeast Extract (certified), 5.36g HycaseSFTM-Sheffield, adjusted pH with KOH to 7.3, qs to 872 ml with SQ H₂O and autoclaved; cooled to 55° C and supplemented with 110 ml 1 M MOPS pH 7.3, 11 ml 50% glucose, 7 ml 1M MgSO₄). Carbenicillin may then be added to the induction culture at a concentration of 50ug/ml

35 The prokaryotic host cells are cultured at suitable temperatures. For *E. coli* growth, for example, the preferred temperature ranges from about 20°C to about 39°C, more preferably from about 25°C to about

2004230670 28 Oct 2010

37°C, even more preferably at about 30°C.

Where the alkaline phosphatase promoter is employed, *E. coli* cells used to produce the polypeptide of interest are cultured in suitable media in which the alkaline phosphatase promoter can be partially or completely induced as described generally, e.g., in Sambrook *et al.*, *supra*. The culturing need never take place in the absence of inorganic phosphate or at phosphate starvation levels. At first, the medium contains inorganic phosphate in an amount above the level of induction of protein synthesis and sufficient for the growth of the bacterium. As the cells grow and utilize phosphate, they decrease the level of phosphate in the medium, thereby causing induction of synthesis of the polypeptide.

If the promoter is an inducible promoter, for induction to occur, typically the cells are cultured until a certain optical density is achieved, e.g., a A_{550} of about 200 using a high cell density process, at which point induction is initiated (e.g., by addition of an inducer, by depletion of a medium component, etc.), to induce expression of the gene encoding the polypeptide of interest.

Any necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art, introduced alone or as a mixture with another supplement or medium such as a complex nitrogen source. The pH of the medium may be any pH from about 5-9, depending mainly on the host organism. For *E. coli*, the pH is preferably from about 6.8 to about 7.4, and more preferably about 7.0.

One selective media that can be used for culturing yeast is a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser *et al.*, *Methods in Yeast Genetics* (Cold Spring Harbor Press, Cold Spring Harbor, NY, 1994), p. 208-210.

E. Detecting Expression

Gene expression may be measured in a sample directly, for example, by conventional Southern blotting, northern blotting to quantitate the transcription of mRNA (Thomas, *Proc. Natl. Acad. Sci. USA*, **77**: 5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences of the polypeptide. Various labels may be employed, most commonly radioisotopes, particularly ^{32}P . However, other techniques may also be employed, such as using biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, assays or gels may be employed for detection of protein.

F. Purification of Polypeptides

When using recombinant techniques, the polypeptide herein is produced intracellularly or in the periplasmic space. If the polypeptide is produced intracellularly, as a first step, the particulate debris, either host cells or lysed cells (e.g. resulting from homogenization), is removed, for example, by centrifugation or ultrafiltration, to produce a cell broth or homogenate.

Then, in accordance with this disclosure, the cellular impurities as defined above are removed from the homogenate or broth by precipitation using the ethacridine lactate under the conditions as set forth above, and the resulting mixture is treated so that the polypeptide of interest, in soluble form, is recovered.

The separation of the target polypeptide from the broth or homogenate may be accomplished by any suitable means, including those well known in the art such as centrifugation and filtration. Preferably, the

separation is performed using centrifugation or tangential flow filtration, for example, using a filter of about 300 kiloDaltons to 1 micron.

After the polypeptide is separated from the broth or homogenate, it may be purified by any known means, including chromatography or filtration such as ultrafiltration/diafiltration or tangential flow filtration.

5 In particular, the following procedures, individually or in combination, are exemplary of suitable purification procedures, with the specific method(s) used being dependent on the type of polypeptide: immobilized metal affinity chromatography (IMAC), aqueous two-phase separation (ATPS), fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reversed-phase HPLC; hydrophobic-interaction chromatography (HIC); chromatography on silica; chromatography on an ion-exchange resin such as S-
10 SEPHAROSE™ and DEAE; chromatofocusing; SDS-PAGE; ammonium-sulfate precipitation; ultrafiltration/diafiltration, tangential flow filtration, and gel filtration using, for example, SEPHADEX™ G-75.

For example, as part of the overall recovery process for the protein, the polypeptide may be exposed to an immobilized reagent that binds to or modifies the polypeptide. Thus, the polypeptide may be 15 subjected to affinity chromatography wherein an immobilized reagent that binds specifically to the polypeptide, such as an antibody, captures the antibody and impurities pass through the affinity chromatography column. The polypeptide can be subsequently eluted from the column by changing the conditions such that the polypeptide no longer binds to the immobilized reagent. The immobilized reagent may also be an enzyme such as a protease that modifies the polypeptide (Sahni *et al.*, *Anal. Biochem.*, **193**: 178-185 (1991) and Voyksner *et al.*, *Anal. Biochem.*, **188**: 72-81 (1990)).

Another type of purification process is filtration. Filtration of fine particle size contaminants from fluids has been accomplished by the use of various porous filter media through which a contaminated composition is passed such that the filter retains the contaminant. Retention of the contaminant may occur by mechanical straining or electrokinetic particle capture and adsorption. In mechanical straining, a particle 25 is retained by physical entrapment when it attempts to pass through a pore smaller than itself. In the case of electrokinetic capture mechanisms, the particle collides with a surface within the porous filter and is retained on the surface by short-range attractive forces. To achieve electrokinetic capture, charge-modifying systems can be used to alter the surface charge characteristics of a filter (see, e.g., WO 90/11814). For example, where the contaminant to be removed is anionic, a cationic charge modifier can be used to alter the charge 30 characteristics of the filter such that the contaminant is retained by the filter.

Monoclonal antibodies may be suitably separated from the precipitants by conventional antibody purification procedures such as, for example, gel filtration or electrophoresis, dialysis, HIC, affinity chromatography, e.g. protein-A SEPHAROSE™, protein-G, antigen-affinity or anti-IgG affinity chromatography, homogenization, clarification by filtration of centrifugation, precipitation, e.g. by treatment 35 with ammonium sulfate, polyethylene glycol or caprylic acid, ion-exchange chromatography, e.g. using resins such as hydroxyapatite, e.g., resins containing calcium-phosphate such as ceramic-hydroxyapatite and BIOGEL H™, and anion-exchange resins including those having a positively charged moiety (at neutral pH), such as diethylaminoethane (DEAE), polyethyleneimine (PEI), and quaternary aminoethane (QAE), for example, Q-SEPHAROSE FAST FLOW™ resin (Pharmacia), DEAE-SEPHAROSE FAST FLOW™ resin, 40 DEAE-TOYOPEARL™ resin, QAE-TOYOPEARL™ resin, POROS-Q™ resin, FRACTOGEL-DMAE™

resin, FRACTOGEL EMD-TMAETM resin, MATREX CELLUFINE DEAETM, and the like. Methods for isolating and purifying antibodies are further described in Antibodies: A Laboratory Manual", Harlow and Lane, eds. (Cold Spring Harbor Laboratories, New York: 1988).

In one specific embodiment, the recovery step involves exposing the solubilized polypeptide to a solid phase to which is immobilized a reagent that binds to, or modifies, the polypeptide. In one embodiment, the solid phase is packed in a column and the immobilized reagent captures the polypeptide. In another embodiment, the reagent chemically and/or physically modifies the polypeptide and is immobilized on the solid phase that is, e.g., packed in a column, and the composition is passed through the column. For example, the polypeptide may comprise a precursor domain that the immobilized reagent removes as part of the recovery process, e.g., the precursor polypeptide is an antibody with a leucine zipper dimerization domain, which is removed by immobilized pepsin in the recovery process.

In this embodiment, the composition comprising the polypeptide and leached reagent (and optionally one or more further contaminants) is then passed through a filter bearing a charge that is opposite to the charge of the reagent at the pH of the composition, so as to remove leached reagent from the composition. The filter may be positively charged to remove contaminants that are negatively charged at the pH of the composition, such as acidic proteases, protein A, protein G or other reagents that can leach from affinity columns. Alternatively, the filter may be negatively charged to remove contaminants that are positively charged at the pH of the composition, such as basic proteases. Preferably, the charge characteristics of the polypeptide of interest in the composition passed through the filter are such that the polypeptide is not significantly retained by the filter and passes therethrough. The filter may be placed "in line" with the effluent treated as in the previous step (i.e., the effluent flows directly through the filter). This can be achieved by connecting the filter directly to the column effluent port, before the effluent is collected into a pool tank. The filter may be regenerated using techniques applicable to the type of filter used.

HIC has also been used for purifying antibody fragments. See, for example, Inouye *et al.*, Protein Engineering, pp. 6, 8 and 1018-1019 (1993); Inouye *et al.*, Animal Cell Technology: Basic & Applied Aspects, 5: 609-616 (1993); Inouye *et al.*, Journal of Biochemical and Biophysical Methods, 26: 27-39 (1993); Morimoto *et al.*, Journal of Biochemical and Biophysical Methods, 24: 107-117 (1992); and Rea *et al.*, Journal of Cell Biochem., Suppl. 0, Abstract No. X1-206 (17 Part A), p. 50 (1993). HIC columns normally comprise a base matrix (e.g., cross-linked agarose or synthetic copolymer material) to which hydrophobic ligands (e.g., alkyl or aryl groups) are coupled. Many HIC columns are available commercially. Examples include, but are not limited to, Phenyl SEPHAROSE 6 FAST FLOWTM column with low or high substitution (Pharmacia LKB Biotechnology, AB, Sweden); Phenyl SEPHAROSETM High Performance column (Pharmacia LKB Biotechnology, AB, Sweden); Octyl SEPHAROSETM High Performance column (Pharmacia LKB Biotechnology, AB, Sweden); FRACTOGELTM EMD Propyl or FRACTOGELTM EMD Phenyl columns (E. Merck, Germany); MACRO-PREPTM Methyl or MACRO-PREPTM t-Butyl Supports (Bio-Rad, California); WP HI-Propyl (C 3)TM column (J. T. Baker, New Jersey); and TOYOPEARLTM ether, phenyl or butyl columns (TosoHaas, PA).

Examples of batch hydrophobic chromatography matrices are well known in the art and include C₁₈ alkyl chains linked to a support matrix such as SEPHAROSETM, agarose, or silica, e.g., butyl, phenyl, or octyl SEPHAROSETM, or polymers such as cellulose or polystyrene. US Pat. No. 6,214,984 describes use of

low-pH hydrophobic interaction chromatography (LPHIC) for antibody and antibody fragment purification. This method is particularly useful for purifying antibody fragments, especially correctly folded and disulfide linked antibody fragments (e.g., Fab fragments) from contaminating antibody fragments that are not correctly folded and/or disulfide linked. Prior to LPHIC, the antibody composition prepared from the cells 5 is preferably subjected to at least one purification step, with examples including hydroxyapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on certain human heavy chains (Lindmark *et al.*, *J. Immunol. Meth.*, **62**: 1-13 (1983)). Protein G is recommended for all mouse isotypes 10 and for one human isotype (Guss *et al.*, *EMBO J.*, **5**: 1567-1575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrene/divinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a CH 3 domain, the BAKERBOND ABXTM resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein 15 purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse-phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™, chromatography on an anion- or cation-exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

G. Uses of Polypeptides

20 The polypeptide thus recovered may be formulated in a pharmaceutically acceptable carrier and is used for various diagnostic, therapeutic, or other uses known for such molecules. For example, antibodies described herein can be used in immunoassays, such as enzyme immunoassays.

25 Therapeutic uses for the polypeptides purified using the method described herein are also contemplated. For example, a growth factor or hormone can be used to enhance growth as desired, and an antibody can be used for redirected cytotoxicity (e.g., to kill tumor cells), as a vaccine adjuvant, for delivering thrombolytic agents to clots, for delivering immunotoxins to tumor cells, for converting enzyme activated prodrugs at a target site (e.g., a tumor), for treating infectious diseases, or for targeting immune complexes to cell surface receptors.

30 Therapeutic formulations of the polypeptide are prepared for storage by mixing the polypeptide having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences*, 16th edition, Osol, A., Ed., (1980)), in the form of lyophilized cake or aqueous solutions.

35 Acceptable carriers, excipients, or stabilizers are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, PLURONICS™, or polyethylene glycol (PEG).

The polypeptide also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions.

5 Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, *supra*.

The polypeptide to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. The antibody ordinarily will be stored in lyophilized form or in solution.

10 Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

15 The route of polypeptide administration is in accord with known methods, *e.g.*, injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial, or intralesional routes, or by sustained-release systems as noted below. The polypeptide is administered continuously by infusion or by bolus injection.

20 Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the polypeptide, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (*e.g.*, poly(2-hydroxyethyl-methacrylate) as described by Langer *et al.*, *J. Biomed. Mater. Res.*, **15**: 167-277 (1981) and Langer, *Chem. Tech.*, **12**: 98-105 (1982) or poly(vinylalcohol)), polyactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, *Biopolymers*, **22**: 547-556 (1983)), non-degradable ethylene-vinyl acetate (Langer *et al.*, *supra*), degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(*-*)-3-hydroxybutyric acid (EP 133,988).

25 While polymers such as ethylenc-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When, for example, encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for antibody stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulphydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

30 Sustained-release polypeptide compositions also include liposomally entrapped polypeptides.

35 Liposomes containing the antibody are prepared by methods known per se: DE 3,218,121; Epstein *et al.*, *Proc. Natl. Acad. Sci. USA*, **82**: 3688-3692 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA*, **77**: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the most effective therapy with polypeptide.

An effective amount of polypeptide to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to tier the dosage and modify the route of administration as required to obtain the most beneficial therapeutic effect. A typical daily dosage might range from about 1 μ g/kg to up to 10 mg/kg or more, depending on the factors mentioned above. Typically, the clinician will administer polypeptide until a dosage is reached that achieves the desired effect. The progress of this therapy is easily monitored by conventional assays.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention. All literature and patent citations herein are incorporated by reference.

EXAMPLE 1

Materials and Methods

A. Plasmids, Transformation, Fermentation15 1. Production of rhuFab'2 (xCD18)a. Plasmid Construction

The control plasmid, pS1130, was designed for the dicistronic expression of anti-CD18 F(ab')₂ and it was based on the vector described by Carter *et al.*, *Bio/Technology*, **10**: 163-167 (1992). This design places transcription of the genes for both light chain and the heavy chain fragment with a C-terminal leucine zipper under the control of a single *phoA* promoter. Transcription ends with a λt_0 transcriptional terminator located downstream of the coding sequence for the heavy chain-leucine zipper (Scholtissek and Grossé, *Nucleic Acids Res.*, **15**(7): 3185 (1987)). The heat stable enterotoxin II signal sequence (STII) precedes the coding sequence for each chain and directs the secretion of the polypeptide into the periplasm (Lee *et al.*, *Infect. Immun.*, **42**: 264-268 (1983); Picken *et al.*, *Infect. Immun.*, **42**: 269-275 (1983)). Leucine zipper was attached to the C-terminal end of heavy chain fragment to promote the dimerization of the two Fab' arms.

The dual-promoter plasmid containing two separate translational units, pxCd18-7T3, temporally separates the transcription of light chain from the transcription of heavy chain. As in pS1130, light chain remains under the control of the *phoA* promoter. However, in pxCd18-7T3, a λt_0 transcriptional terminator follows the light chain coding sequence. Downstream of this terminator, the *tacII* promoter was added to control the transcription of the heavy chain fragment/C-terminal leucine zipper (DeBoer *et al.*, *Proc. Natl. Acad. Sci. USA*, **80**: 21-25 (1983)). A second λt_0 transcriptional terminator follows this coding sequence. Silent codon variants of the STII signal sequence were used to direct the secretion of both chains (Simmons and Yansura, *Nature Biotechnology*, **14**: 629-634 (1996)).

A schematic comparison of the single promoter control plasmid vs. the dual-promoter plasmid is depicted in Figure 1. The expression cassette sequence of pxCd18-7T3 is provided in Figure 2 (SEQ ID NO: 1), and the amino acid sequences (SEQ ID NOS:2 and 3) from the two translational units are shown in Figures 3A (light chain) and 3B (heavy chain), respectively.

b. Fermentation

The host strain used in fermentation was a derivative of *E. coli* W3110, designated 59A7. The complete genotype of 59A7 is W3110 Δ *lhuA* *phoA* Δ *E15* Δ (*argF-lac*) Δ 69 *deoC* *degP41* *ilvG2096*(*Yaf*) Δ *prc*

spvW148R. The 59A7 host cells were transformed with pxD18-7T3 plasmid and successful transformants were selected and grown in culture. An additional plasmid, pMS421, was co-transformed along with pxD18-7T3. This additional plasmid, pMS421, is a pSC101-based plasmid which provides lacIq to improve control of the *tacII* promoter, and which also confers spectinomycin and streptomycin resistance.

5 For each 10-liter fermentation, a single vial containing 1.5 ml of culture in 10-15% DMSO was thawed into a 1-L shake flask containing 500 ml of LB medium supplemented with 0.5 ml of tetracycline solution (5 mg/ml) and 2.5 ml 1M sodium phosphate solution. This seed culture was grown for approximately 16 hours at 30°C and was then used to inoculate a 10-liter fermentor.

10 The fermentor initially started with approximately 6.5 liters of medium containing about 4.4 g of glucose, 100 ml of 1M magnesium sulfate, 10 ml of a trace element solution (100 ml hydrochloric acid, 27 g ferric chloride hexahydrate, 8 g zinc sulfate heptahydrate, 7 g cobalt chloride hexahydrate, 7 g sodium molybdate dihydrate, 8 g cupric sulfate pentahydrate, 2 g boric acid, 5 g manganese sulfate monohydrate, in a final volume of 1 liter), 20 ml of a tetracycline solution (5 mg/ml in ethanol), 10 ml of FERMAX Adjuvant 27TM (or some equivalent anti-foam), 1 bag of HCD salts (37.5 g ammonium sulfate, 19.5 g potassium phosphate dibasic, 9.75 g sodium phosphate monobasic dihydrate, 7.5 g sodium citrate dihydrate, 11.3 g potassium phosphate monobasic), and 200 g of NZ Amine A (a protein hydrolysate). Fermentations were performed at 30 °C with 10 slpm of air flow and were controlled at a pH of 7.0 ± 0.2 (although occasional excursions beyond this range occurred in some cases). The back pressure of the fermentor and agitation rate was varied to manipulate the oxygen transfer rate in the fermentor, and, consequently, control the cellular respiration rate.

15 Following inoculation of the fermentor with the cell-containing medium from the shake flask, the culture was grown in the fermentor to high cell densities using a computer-based algorithm to feed a concentrated glucose solution to the fermentor. Ammonium hydroxide (58% solution) and sulfuric acid (24% solution) were also fed to the fermentor as needed to control pH. Further additions of anti-foam were 20 also used in some cases to control foaming. When the culture reached a cell density of approximately 40 OD550, an additional 100 ml of 1M magnesium sulfate was added to the fermentor. Additionally, a concentrated salt feed (consisting of approximately 10 g ammonium sulfate, 26 g dibasic potassium phosphate, 13 g monobasic sodium phosphate dihydrate, 2 g sodium citrate dihydrate and 15 g monobasic potassium phosphate in 1L of water) to the fermentor was started at a rate of 2.5 ml/min when the culture 25 reached approximately 20 OD550 and continued until approximately 1250 ml were added to the fermentor. Fermentations were typically continued for 72-80 hours.

30 During the fermentation, once the dissolved oxygen setpoint for the fermentation was reached, the concentrated glucose solution was fed based on the dissolved oxygen probe signal in order to control the dissolved oxygen concentration at the setpoint. Consequently, in this control scheme, manipulations of fermentor operating parameters such as the agitation rate or back pressure, which affect the oxygen transfer 35 capacity in the fermentation, correspondingly manipulated the oxygen uptake rate or metabolic rate of the cells.

A mass spectrometer was used to monitor the composition of the off-gas from the fermentations and enabled the calculation of the oxygen uptake and carbon dioxide evolution rates in the fermentations.

When the culture reached a cell density of approximately 220 OD550, the agitation was decreased from an initial rate of 1000 rpm to approximately 725 rpm over approximately 12 hours. Fifty ml of 200 mM of isopropyl β -D-thiogalactopyranoside (IPTG) was added to induce heavy chain synthesis approximately 12 hours after the culture reached a cell density of 220 OD550.

5 2. Production of Anti-Tissue Factor F(ab')₂

 a. Plasmid Construction

A dual promoter plasmid, pxTF7T3, was created similar to the dual promoter plasmid pXCD18-7T3 set forth above, and used to enable temporal separation of anti-Tissue Factor light-chain and heavy-chain expression. The *lacI* sequence from the plasmid pMS421 was also incorporated onto pxTF7T3 to create a new dual-promoter plasmid pJVG3IL.

10 b. Fermentation

The host strain used in these fermentations was a derivative of *E. coli* W3110, designated 60H4. The complete genotype of 60H4 is: W3110 Δ *flhA* Δ *manA* *phoA* Δ *E15* Δ (*argF-lac*)*169* *deoC2* *degP41* *ilvG2096*(*Val*) *Aprc* *pre-suppressor*. The 60H4 host cells were transformed with pJVG3IL and successful transformants were selected and grown in culture.

Fermentations were run under conditions similar to those for anti-CD18 F(ab')₂ as described above, with the principal exceptions that the run length varied between approximately 72 and 114 hours, and heavy chain was induced using IPTG from approximately 4 to 12 hours following the attainment of a culture OD550 of 220.

20 3. Production of full-length anti-TF antibodies

 a. Plasmid Construction

The expression cassette for the plasmid pxTF-7T3FL comprises, from 5' to 3': (1) a *phoA* promoter (Kikuchi *et al.*, *Nucleic Acids Res.*, **9**(21): 5671-5678 (1981)); (2) *trp* Shine-Dalgarno (Yanofsky *et al.*, *Nucleic Acids Res.*, **9**: 6647-6668 (1981)); (3) a silent codon variant of the STII signal sequence (TIR relative strength about 7) (Simmons and Yansura, *Nature Biotechnology*, **14**: 629-634 (1996)); (4) coding sequence for anti-tissue factor light chain; (5) λ_t terminator (Scholtissek and Gross, *Nucleic Acids Res.*, **15**: 3185 (1987)); (6) a *tacII* promoter ((DeBoer *et. al.*, *Proc. Natl. Acad. Sci. USA*, **80**: 21-25 (1983)); (7) a second *trp* Shine-Dalgarno; (8) a second silent codon variant of the STII signal sequence (TIR relative strength about 3); (9) coding sequence for anti-tissue factor full-length heavy chain; and (10) a second λ_t terminator. This expression cassette was cloned into the framework of the *E. coli* plasmid pBR322 (Sutcliffe, *Cold Spring Harbor Symp. Quant. Biol.*, **43**: 77-90 (1978)).

Thus, the vector design of pxTF-7T3FL allows for the temporally separate expression of each chain by using two different, rather than two identical, promoters. In this plasmid, light chain is under the control of the *phoA* promoter. However, the *tacII* promoter is used to control the transcription of heavy chain. As is known in the art, *phoA* and *tacII* promoters are induced under substantially different conditions. A schematic comparison of a single-promoter plasmid paTF130 and pxTF-7T3FL is depicted in Figure 4. The nucleic acid sequence of the expression cassette of pxTF-7T3FL (SEQ ID NO:4) is provided in Figure 5, and the polypeptide sequences it encodes (SEQ ID NOS:5 and 6) are provided in Figures 6A (light chain) and 6B (heavy chain, respectively. The host cells below were co-transformed with pxTF-7T3FL and pJJ247.

pJJ247 encodes a *tacII* promoter driving the expression of both DsbA and DsbC, with DsbA first in the series, and its construction is described in WO02/061090.

b. Fermentation

For small-scale expression *E. coli* strain 61D6, with genotype W3110 *ΔfhuA* (*ΔtonA*) *ptr3 lacIq* 5 *lacL8 ΔompTΔ(nmpc-fepE) degP41*, was used as host cells. Following transformation, selected transformant picks were inoculated into 5 ml of Luria-Bertani medium supplemented with carbenicillin (50 µg/ml) and kanamycin (50 µg/ml) grown at 30°C on a culture wheel overnight. A 10-liter fermentation was conducted using the media as described in WO02/061090 and the basic fermentation conditions were also as described in WO02/061090, except that the following modifications were made to the fermentation process: 300 mL of 10 1M NaPO₄, pH 7.0 was added at approximately 40 hours to give a concentration of about 30 mM. 100 mL of a 200-mM solution of IPTG was added at approximately 44 hours to give a concentration of about 2 mM. The fermentation was harvested 80 hours after inoculation.

B. Protein Identification

The one-dimensional SDS-PAGE gel electrophoresis was carried out in a 4-12% linear acrylamide 15 gradient from Novex. Specifically, the system used was the NOVEX® NUPAGE™ System, consisting of NUPAGE™ Bis-TRIS Pre-Cast Gels (for low- to mid-molecular weight proteins).

C. Chemicals

The precipitation agent ethacridine lactate was 98% pure and purchased from Sigma (St. Louis, 20 MO, USA). Ethacridine lactate has a molecular weight of 361.4Da. All other chemicals were of analytical grade.

D. Precipitation

The antibody- and F(ab')₂-containing *E. coli* materials were homogenized using a microfluidizer 25 from Watts Fluidair Inc. (model B12-04DJC, Kittery, MN, USA). The cells were passed three times through the microfluidizer at 4-bar pressure. To avoid heat degradation of the proteins, the material was passed through an ice-water bath during each pass through the microfluidizer. The total protein concentration in the F(ab')₂ homogenates was 30 mg/ml. The full-length anti-TF homogenate, which was derived from re-suspended paste, had a total protein concentration of 18 mg/ml. The anti-TF paste was re-suspended in 25 mM TRIS-HCl buffer, pH 7.5.

The ethacridine lactate precipitation agent was dissolved in water to the desired final concentration 30 (w/v).

1. pH study:

The precipitation experiments were performed with a constant ethacridine lactate concentration of 0.6% (w/v). A 0.8% ethacridine lactate solution was prepared and mixed with the *E. coli* homogenate in a 3:1 ratio, e.g., 3 ml of ethacridine lactate and 1 ml of *E. coli* homogenate. The pH was adjusted using HCl or 35 NaOH depending on the pH desired.

2. Ethacridine lactate concentration study:

The homogenate was diluted 4-fold with ethacridine lactate stock solutions (1:3 as in the pH study) and the pH was kept at a set value for each of the target proteins. For anti-CD18, the pH was 8.5 and for anti-TF pH 7.5. The final ethacridine lactate concentrations in the precipitation systems were 0.15, 0.30, 0.45, 0.60, 0.75, and 0.9% (w/v). As a reference a set of experiments with 0% ethacridine lactate was performed.

5 3. Conductivity/dilution study:

Various concentrations of NaCl were added to anti-CD18 homogenate to evaluate the effect of conductivity on protein precipitation. The NaCl concentrations studied were 0, 50, 100, 150, 200, and 400 10 mM. A reference series without ethacridine lactate was also performed to determine if any protein was precipitated due to the high salt concentration. The pH in this salt spike study was 8.5 and the anti-CD18 homogenate was diluted 4-fold. The ethacridine lactate concentration in the samples was 0.6% and 0% for the reference experiments.

15 To change the conductivity of the sample the homogenates were diluted in increasing amounts, the pH was kept constant, pH 8.5 and 7.5 for anti-CD18 and anti-TF, respectively. All experiments had a final ethacridine lactate concentration of 0.6% (w/v). The homogenates were diluted 2, 3, 4, 5, 6, and 7-fold.

15 4. Temperature study:

20 Some experiments were performed at elevated temperatures. The *E. coli* homogenate was diluted 4-fold and the final ethacridine lactate concentration was 0.6%. The pH was 8.5 or 7.5 for anti-CD18 and anti-TF, respectively. The samples were incubated in a thermostated water bath at a desired temperature, *i.e.*, 50, 60, and 70°C. The samples were incubated for 20-120 minutes at elevated temperatures. One long-time incubation, 16 hours, was performed at 50°C.

25 After mixing the precipitation agent and *E. coli* homogenate together and adjustment of pH, the samples were incubated under agitating conditions for 30-60 minutes. The precipitation experiments were performed in glass tubes, at 4-ml scale. All experiments were performed in duplicate and average values were reported.

25 E. coli Protein Assay

30 Ethacridine lactate interacts with most commonly used protein measurement assays, *e.g.*, Bradford, BCA, and spectrophotometric absorption measured at 280 nm. Thus, the total protein concentrations were measured using a generic *E. coli* protein ELISA. The samples were diluted in a fish gelatin-containing buffer (0.15 M NaCl, 0.1 M NaPO₄, 0.1% fish gelatin, 0.05% TWEEN 20TM, 0.05% PROCLINTM 300) to reduce the unspecified binding to the anti-*E. coli* protein antibodies. The coating antibody was Goat anti-Whole ECP. The conjugate antibody was an anti-antibody Whole ECP, attached to horseradish peroxidase. The absorption at 405 nm was monitored using a plate reader from Molecular Devices model SPECTRA 35 MAX PLUSTM (Sunnyvale, CA, USA).

35 F. Protein G Assay

40 To measure the recovered F(ab')₂ and antibody concentrations, a protein G affinity chromatography assay was used. An IMMUNO DETECTIONTM protein G column was purchased from PerSeptive Biosystems (Framingham, MA, USA). The column was equilibrated with phosphate-buffered saline (PBS) and eluted with PBS that had been pH adjusted to 2.2 with HCl. To minimize the interference from

ethacridine lactate, samples were treated on exclusion spin columns (BIO-SPIN® 6 Tris columns (Bio-Rad Laboratories, Hercules, CA, USA)) before assay. The spin columns were used as recommended by the vendor. A tetramethylammonium chloride (TMAC) wash step was introduced (Fahrner *et al.*, *Biotechnology and Genetic Engineering Reviews*, 18: 302-327 (2001)) to the chromatography method to

5 minimize any interference of residual ethacridine lactate in the sample. The assay was performed using a HPLC (HP 1090™ liquid chromatograph) from Hewlett Packard (Mountain View, CA, USA). The samples were diluted with PBS. Standard curves were prepared for each of the proteins using purified protein (from Genentech, Inc.).

G. DNA Assay

10 The DNA concentration in the supernatants after precipitation was measured using a Pico Green Kit from Molecular Probes (Eugene, OR, USA). It is a fluorescence assay where the fluorescent reagent (Pico green) binds to double-stranded DNA. The Pico green reagent is excited at 502 nm and the emission at 523 nm is recorded. The assay was performed using a fluorescence plate reader. SPECTRA MAX GENINI XS™, from Molecular Devices (Sunnyvale, CA, USA). Ethacridine lactate interacts with the Pico green assay and thus the precipitation agent was removed from the solution prior to analysis. Ethacridine lactate was removed from the sample using BIO-SPIN® 6 TRIS columns, described in the protein G affinity chromatography assay section.

H. SDS-PAGE

20 Supernatants obtained after ethacridine lactate precipitation were analyzed by SDS-PAGE. Non-reduced 4-12% NUPAGE™ gels from Novex (San Diego, CA, USA) were used to visualize the purification and recovery of anti-CD18 and anti-TF. Pre-cast gels were used and the running buffer was MOPS (pre-made concentrate purchased from Novex). The gels were stained with a filtered solution of COOMASSIE 25 BRILLIANT BLUE R250™. The supernatants were volume compensated in regard to respective clarified *E. coli* extract. In this way, the intensity of the protein bands in the *E. coli* extract and samples should be identical if a 100% yield is obtained in the supernatants after precipitation with ethacridine lactate. Thus, the gels can be used to indicate accurately the extent of purification obtained from the precipitation.

I. Ethacridine lactate Solubility

25 Two ethacridine lactate solutions, *i.e.*, 0.6 and 1.2%, were studied. Each solution was divided into two aliquots and pH adjusted to 6.0 and 9.0, respectively. To obtain a slight buffer capacity in the system, the ethacridine lactate was dissolved in 10 mM Tris-HCl buffer. Each ethacridine lactate solution was exposed to increasing amounts of NaCl, *i.e.*, 0, 50, 100, 150, 200, 300, and 600 mM. The samples were incubated for 3 hours, then centrifuged for 20 minutes at 12000 g in a microfuge (SORVALL MC12V™, DuPont, Wilmington, DE, USA). The supernatants were assayed for ethacridine lactate by measuring the absorption at 270 nm. The spectrophotometer used was a HP8453 UV-VIS™ from Hewlett Packard (Wilmington, DE), now Agilent Technologies (Palo Alto, CA) and known as the AGILENT 8453 UV-VIS™ spectrophotometer. A standard curve was derived from a solution with known ethacridine lactate concentration.

J. Turbidity

To measure the stability of supernatants as a function of time and temperature, the turbidity was monitored. The turbidity meter used was from HACH (model 2100N, Ames, Iowa, USA). The samples were measured at room temperature and without dilution of the sample.

5 The anti-CD18 homogenate was treated either with 0.6% ethacridine lactate, 0.2% PEI, or only water. In all of the three samples the anti-CD18 homogenate was diluted 4-fold and the pH was 7.2 ± 0.2 . After centrifugation for one hour at 4000 g the supernatants were recovered and divided into two aliquots. One part of each sample was incubated at room temperature (21°C) and the other one at 4°C.

10 **Results and Discussion**Effect of pH

The ethacridine lactate molecule is positively charged over most of the pH interval (Miller, *supra*; Neurath and Brunner, *supra*; Franek, *Methods in Enzymology*, ed. Langone, J.J., Van Vunakis, H., 121: 631-638 (1986)). However, since pH changes affect the charge on polypeptides and there is a correlation 15 between the pI of the polypeptide and the pH at which it precipitates when exposed to ethacridine lactate (Neurath and Brunner, *supra*), the effect of pH on degree of purification in the method of this invention was investigated.

20 Homogenates containing anti-CD18 F(ab')₂, anti-TF F(ab')₂, and full-length anti-TF, respectively, were exposed to a 0.6% ethacridine lactate solution adjusted to cover the pH range 4-10. Figs. 8A-8C show the clarified phases after ethacridine lactate treatment and centrifugation of each of these proteins, respectively.

25 In Table 1, anti-TF was studied both as a full-length antibody and as a F(ab')₂. *E. coli* homogenates were treated at a 1:3 ratio with a 0.8% (w/v) ethacridine lactate solution, *i.e.*, a final ethacridine lactate concentration of 0.6% in the sample. The pH was adjusted with HCl and NaOH respectively to obtain desired pH. The yields and purification factors are calculated in respect to each of the clarified cell homogenates. The DNA concentration in the recovered supernatants is also reported in the table.

Table 1

The pH effect on the purification and yield of α CD18 and α TF when treated with ethacridine lactate

pH	anti-CD18 F(ab') ₂		anti-TF F(ab') ₂		anti-TF full-length Ab		DNA conc. (μ g/ml)
	Purification factor*	Yield (%)	Purification factor*	Yield (%)	Purification Factor*	Yield (%)	
4	3.2	100	1.8	93	2.4	100	0.2
5	5.3	100	5.1	93	4.4	100	0.1
6	5.6	100	5.1	67	6.7	100	≤ 0.001
7	5.6	100	5.4	70	7.1	86	≤ 0.001
8	5.6	87	5.0	42	6.2	65	≤ 0.001
9	5.4	84	4.7	22	5.3	24	≤ 0.001
10	4.2	85	4.7	18	0.9	7	0.3

30 * A value of 1 is the same purification as is obtained in the system not treated with ethacridine lactate.

It can be seen that for all three proteins a bell-shaped curve was obtained with respect to extent of purification and DNA concentration over the pH range of 4 to 10. At the middle pH values, *i.e.*, pH 5.9, an approximately 5-fold purification was obtained. The yield of anti-CD18 $\Gamma(ab')_2$ and anti-TF antibody full-length and $F(ab')_2$ decreased with higher pH, and anti-TF antibody full-length and $F(ab')_2$ had a stronger pH dependence than anti-CD18 $F(ab')_2$. The yield of anti-CD18 $F(ab')_2$ is decreased from 100 to 85% and of anti-TF $F(ab')_2$ from 93 to 18% in the pII range 4-10. Without being limited to any one theory, this could partly be due to the lower pI of anti-TF compared to anti-CD18, *e.g.*, pI 7.5 and 8.9, respectively; however, these are theoretically calculated pI values.

The full-length anti-TF protein has an even stronger pH dependence than the $F(ab')_2$ version of the protein. The purity of full-length anti-TF is highest at pH 7.0. At pH above 8 significant yield losses are observed (Fig. 8C). For the full-length anti-TF a pH of about 7.0 is preferred, *i.e.*, a 7.1-fold purification and 86% yield. One possible explanation for the higher losses of anti-TF compared to anti-CD18, without being limited to any one theory, is that anti-TF has more negative surface charges than anti-CD18 when incubated above its pI. Analogous to this, without being limited to any one theory, the larger full-length anti-TF might have more negative surface charges than the corresponding $F(ab')_2$ and thus a significantly higher yield loss is observed when increasing the pH.

However, components other than cell debris and host protein have to be removed from the target polypeptide. One such component is DNA. The major disadvantage of having high DNA concentration together with the target polypeptide is that the viscosity of the solution increases. This will have a negative impact on the further downstream processing. In addition, if an anion-exchange column is used as the first capture column, the negatively charged DNA will bind to the resin and thus reduce the protein capacity of the column.

Hence, the DNA concentration in the supernatants after precipitation with ethacridine lactate was determined. The results show that the DNA concentration in the supernatants after ethacridine lactate precipitation is significantly reduced compared to the initial DNA concentration obtained in the *E. coli* homogenate. However, the DNA concentration in the supernatant increased as the pH thereof decreased, *i.e.*, 0.1 and 0.2 μ g/ml at pH 5.0 and 4.0 respectively. Without being limited to any one theory, this could be due to the fact that the phosphates on the DNA become less negatively charged at lower pII. At very high pII, *e.g.*, pH 10.0, the DNA concentration is significantly increased (0.3 μ g/ml) and the protein purity also was decreased. Without limitation to any one theory, this is partly due to the fact that the pH is above the pI of the antibody and $F(ab')_2$; however, it can also partly be due to the fact that ethacridine lactate is less charged at this pH.

Effect of ethacridine lactate concentration

The *E. coli* homogenates were mixed with ethacridine lactate solutions in a (1:3) ratio. The ethacridine lactate concentration in the samples was increased in 0.15% increments from 0 to 0.9% (w/v). This study was performed at pH 8.5, 7.5, and 6.0 for anti-CD18, anti-TF $F(ab')_2$, and full-length antibody, respectively.

Table 2 shows the effect of ethacridine lactate concentration on purification and yield. In Table 2, anti-TF was studied both as a full-length antibody and as a $F(ab')_2$. *E. coli* homogenates were treated at a 1:3

ratio with different concentrations of ethacridine lactate solutions; the final ethacridine lactate concentration in the sample is reported. The pH of the anti-CD18, anti-TF(F(ab')₂), and full-length anti-TF was 8.5, 7.5 and 6.0, respectively. The DNA concentration in the recovered supernatants is also reported in the table.

5 Table 2
The effect on purity and yield of anti-CD18 and anti-TF when treated with increasing ethacridine lactate concentration

Ethacridine lactate (% w/v)	anti-CD18 F(ab') ₂		anti-TF F(ab') ₂		anti-TF Full-length Ab		DNA conc. (µg/ml)
	Purification factor*	Yield (%)	Purification factor*	Yield (%)	Purification Factor*	Yield (%)	
0	1.0	100	1.0	100	1.0	100	77.9
0.15	1.1	100	1.3	97	3.4	100	3.9
0.30	1.7	100	2.2	93	6.7	100	0.4
0.45	4.0	100	5.2	96	6.5	100	0.2
0.60	5.8	100	5.6	98	6.5	100	≤ 0.001
0.75	5.9	100	5.6	78	6.3	100	≤ 0.001
0.90	5.9	100	5.4	70	6.3	100	≤ 0.001

* A value of 1 is the same purification as is obtained in the system not treated with ethacridine lactate.

10 The purity of the antibodies was shown to be strongly correlated with the concentration of ethacridine lactate (Figs. 9A-9C and Table 2). At ethacridine lactate concentrations above about 0.6% the effect of increased ethacridine lactate concentration on increased purification was not dramatic. However, at lower concentrations of ethacridine lactate, *i.e.* when ethacridine lactate is deficient, every slight addition of the precipitation agent resulted in a substantial additional purification of the F(ab')₂.

15 The yield of anti-CD18 F(ab')₂ was not affected by the addition of different concentrations of ethacridine lactate. The step recovery for both of the F(ab')₂ was about 90% for all the experiments. However, it appears easier to obtain quantitative recovery of the F(ab')₂ anti-CD18 than the anti-TF F(ab')₂. Without being limited to any one theory, this could be because there are more negative surface charges on the anti-TF F(ab')₂ compared to the anti-CD18 F(ab')₂ at the pH studied. The full-length anti-TF reached its maximum purification at a lower ethacridine lactate concentration than the corresponding F(ab')₂ protein, 0.3% and 0.6%, respectively. Without being limited to any one theory, this is probably due to a lower overall protein concentration in the anti-TF full-length homogenate, *i.e.*, 18 and 30 mg/ml for the full-length and F(ab')₂ anti-TF, respectively. The full-length anti-TF is also derived from re-suspended paste and the F(ab')₂ is taken as broth directly from the fermentor. Hence, soluble culture media components present in the *E. coli* broth will not be present in the re-suspended full-length anti-TF material, which could partly explain the differences seen, without limitation to any one theory.

20 The DNA concentration in the supernatants is strongly correlated with the data obtained for the protein purification. At 0.6% or higher ethacridine lactate addition, no DNA was detected in the

supernatants. There is also a clear trend of decreasing DNA concentration, *i.e.*, 78 to 0 $\mu\text{g/ml}$, in the supernatants when the ethacridine lactate concentration is increased from 0 to 0.6%.

Effect of Conductivity

5 As ethacridine lactate precipitates proteins partly due to the charged characteristics of the molecule (Neurath and Brunner, *supra*), the conductivity of the sample may have an effect on the purity of the antibody and $\text{F}(\text{ab}')_2$ after precipitation. Hence, if the sample has high salt concentration, *i.e.*, high conductivity, the salts could shield the proteins from the ethacridine lactate and thus reduce the purification effect.

10 . Anti-CD18 homogenate was subjected to two sets of experiments to separate the effect of the conductivity from the protein concentration in the sample. To both sets of experiments NaCl was added (0-400mM). One set of the experiments contained 0.6% ethacridine lactate and in the other set water was used. The water-containing systems were used as controls, so that if the NaCl would give rise to any precipitation this could be distinguished from the precipitation derived from the ethacridine lactate. In the systems without 15 ethacridine lactate, *i.e.* the water systems, no protein precipitation was observed in the 0-400mM NaCl concentration range (Fig. 10A). The ethacridine lactate-containing experiments showed a strong increase in the purification of anti-CD18 when the conductivity was decreased (Fig. 10B).

Without being limited to any one theory, one reason for the improved purification of anti-CD18 at lower conductivity could be the low shielding capacity of the charged ethacridine lactate at low salt 20 concentration. Analogous shielding effects have been seen when PEI is used for protein purification at increasing salt concentration (Jendrisak, *supra*). However, an even more important factor is the low solubility of ethacridine lactate at higher salt concentrations (Miller, *supra*; Neurath and Brunner, *supra*; Franek, *supra*). At 100mM NaCl precipitation of ethacridine lactate was observed in the extraction system, and as the salt concentration was increased, more of the ethacridine lactate was precipitated. Hence, less 25 ethacridine lactate is soluble in the system and available to precipitate proteins and other biomolecules.

When the solubility of ethacridine lactate as a function of NaCl concentration was studied, a pH dependence was observed (Fig. 11). At pH 6 no significant difference in the solubility between the 0.6 and 1.2% ethacridine lactate solution was observed. Both solutions were soluble at 50mM NaCl but nearly fully precipitated at 100mM. For the pH 9 solutions, where the ethacridine lactate was less charged, a significant 30 difference in solubility was seen between the two concentrations of ethacridine lactate. The 0.6% ethacridine lactate solution precipitated at lower salt concentration than the more concentrated ethacridine lactate solution (Fig. 11). Chlorides have been shown to precipitate ethacridine lactate especially efficiently (Franek, *supra*).

In practice, the conductivity of a precipitation experiment will be determined by the dilution factor. 35 Hence, a set of experiments was performed where the impact of the dilution factor of the *E. coli* homogenate was studied. The overall ethacridine lactate concentration was kept constant, *i.e.*, 0.6% (w/v), but the conductivity of the sample was decreased with increased dilution. The results are shown in Table 3. In Table 3, anti-TF was studied both as a full length antibody and as a $\text{F}(\text{ab}')_2$. The concentration of ethacridine lactate in each experiment was 0.6% (w/v). The pH of the anti-CD18, anti-TF($\text{F}(\text{ab}')_2$), and full-length anti-40 TF was 8.5, 7.5 and 6.0, respectively. The yields and purification factors are calculated in respect to each of

the clarified cell homogenates. The DNA concentration in the recovered supernatants is also reported in the table.

5 Table 3
The effect on the purity and yield of anti-CD18 and anti-TF upon dilution of the *E. coli* homogenates in different amounts

		anti-CD18 F(ab') ₂		anti-TF F(ab') ₂		anti-TF Full-length Ab		
Dilution (times)	Cond. (mS)	Purification factor*	Yield (%)	Purification factor*	Yield (%)	Purification Factor*	Yield (%)	DNA conc. (μ g/ml)
2	5.0	2.4	72	2.3	77	3.4	70	20.7
3	4.0	4.1	76	4.0	60	6.7	82	0.5
4	3.2	5.5	71	4.9	95	6.7	95	≤ 0.001
5	2.7	6.1	88	5.2	100	6.5	100	≤ 0.001
6	2.4	6.1	92	5.0	100	6.0	100	≤ 0.001
7	2.1	6.3	100	5.1	100	6.0	100	≤ 0.001

* A value of 1 is the same purification as is obtained in the system not treated with ethacridine lactate.

10 The results showed that the purification of F(ab')₂ was enhanced if the conductivity was lowered by increasing dilution. However, at a conductivity of 3.5mS or lower the effect of conductivity was minor. For the *E. coli* homogenate used in this Example a 4-fold dilution had to be performed to obtain conductivity below 3.5mS. For the full-length anti-TF antibody a slightly lower dilution can be performed than for the F(ab')₂ homogenates. Without limitation to any one theory, this can be due to the fact that the protein 15 concentration in the full-length homogenate is lower. Furthermore, as the full-length material is derived from re-suspended paste, some of the media components that might affect the precipitation have been removed prior to the ethacridine lactate precipitation.

20 In these experiments the protein concentration was decreased with increased dilution (lowering of the conductivity). Thus, even if 0.6% ethacridine lactate, which was found to be an excess concentration in the ethacridine lactate concentration study, was added, maximum removal of host protein and DNA was not obtained in the less diluted sample. This is due to the increased homogenate concentration, *i.e.*, overall protein and DNA, in these samples compared to the sample performed in the ethacridine lactate concentration study. Hence, the overall concentration of protein, DNA, and other components will affect the ethacridine lactate concentration or alternatively the dilution required. The yield of anti-CD18 F(ab')₂ and 25 full-length and F(ab')₂ anti-TF increased with the decrease in conductivity.

The effect on the DNA removal was also found to be conductivity dependent. At a two-fold dilution, *i.e.*, 5.0 mS, a high DNA concentration was obtained in the supernatant, 20.7 μ g/ml. However, if a 3-fold dilution was performed, *i.e.*, 4.0 mS, the concentration was decreased to 0.5 μ g/ml, and at even higher dilution no DNA was detected. As pointed out earlier, in this case the total amount of ethacridine lactate to

protein and DNA increases with the higher dilutions, *i.e.*, at lower conductivity. Thus, the effect seen in this example is both of lowering conductivity and of increasing amounts of ethacridine lactate.

Effect of temperature

5 Temperature is a factor that is known to be important when performing precipitation experiments. Thus, some elevated temperatures were studied in combination with ethacridine lactate. The effect of incubation time at elevated temperature was also investigated.

10 Incubation at elevated temperature, *i.e.*, 50-70°C, had a positive effect on the purity of the two F(ab')₂ proteins (Figs. 12A and 12B). The higher the temperature, the more efficient the purification. An incubation at 70°C significantly improved the purity of the F(ab')₂ proteins. Incubating the sample for longer time, *i.e.*, 40 min compared to 20 min, at 70°C did not improve the purity of the F(ab')₂ (Figs. 12A and 12B), but an approximately 10% loss in yield was observed. However, when the sample was incubated above 70°C, no F(ab')₂ was recovered, which is due to temperature precipitation of F(ab')₂ as well as other *E. coli* proteins.

15 To investigate the effect of the incubation time more closely, an experiment was performed where the samples were incubated for 16 h and 30 min at 50°C. The results showed that there was no significant purification improvement for the sample that had been incubated for 16 h compared to 30 min at this temperature. This indicates that temperature precipitation is a rapid phenomenon, which suggests that a fast heating to appropriate temperature would be more suitable than a long incubation time.

20 The full-length anti-TF was also studied at elevated temperatures. A 15-min incubation at 50°C had a slightly positive effect on the purity of the antibody and no losses in recovered material were observed compared to a sample incubated at room temperature (Fig. 12C). However, if the temperature was increased to 60°C, nearly all anti-TF precipitated. The data indicate that the full-length anti-TF has lower stability at elevated temperature than the F(ab')₂ anti-TF protein.

25 Since an increase in temperature could give rise to modifications of the target polypeptide, the stability of the particular polypeptide of interest at elevated temperature would need to be evaluated before implementation.

Stability of feed stream

30 It is important to recover the cleanest feed streams possible. However, another important characteristic of the feed stream is its stability over time. Hence, the stability of supernatants after ethacridine lactate or PEI treatment and a supernatant after plain centrifugation was compared. Each of the supernatants was incubated at two temperatures, *i.e.*, room temperature (21°C) and 4°C. The stability was monitored by measuring the turbidity of the respective sample.

35 In Fig. 13 the change in turbidity over time is shown for the three different supernatants, *i.e.*, ethacridine lactate, PEI, and non-treated clarified supernatant, respectively. It can clearly be seen that the polyelectrolytes, *i.e.*, ethacridine lactate and PEI, significantly reduced the turbidity of the supernatant. Directly after clarification the non-treated clarified supernatant had a turbidity of about 700 NTU, whereas the PEI-treated sample only had half that turbidity, *i.e.*, 327 NTU, and the ethacridine lactate supernatant had a turbidity of 1 NTU. There was not a major difference seen, either over time or with incubation

temperatures, when monitoring the turbidity of the PEI-treated supernatant. For the supernatant that had only been centrifuged there was a trend of lower turbidity in the room temperature sample during the first 48 hours. However, the room temperature sample at 72 hours could not be measured due to high turbidity of the sample. The ethacridine lactate-treated supernatant had very low turbidity that was not significantly increased if incubated at 4°C. When the sample was incubated at 21°C the turbidity increased significantly, *i.e.*, from 1 to 100 NTU over 72 hours. However, 100 NTU is still less than the turbidity obtained for the other two supernatants directly after clarification. Hence, it can be concluded that the supernatant recovered after ethacridine lactate treatment at pH 7 is a stable feed stream.

10 **Conclusion**

Ethacridine lactate can successfully be used as precipitation agent for the primary recovery of heterologous polypeptides from culture broth or homogenate. When ethacridine lactate is used as precipitation agent, the ideal target polypeptide preferably has a higher pI than the average host proteins. Hence, most proteins can be negatively charged and become precipitated by ethacridine lactate at the same time as the target protein is positively charged and thus recovered in the supernatant.

15 The preferred concentration of ethacridine lactate for precipitation is highly dependent on the concentration of host protein and DNA in the broth or homogenate. The higher the protein and DNA concentration in the broth or homogenate, the higher is the amount of ethacridine lactate required. Hence, the more negatively charged components available for ethacridine lactate to complex with, and thus precipitate, 20 the higher is the preferred amount of ethacridine lactate for precipitation. The lower the conductivity of the solution when performing the precipitation, the more efficient is the purification of the polypeptide. The precipitation step will give significant polypeptide purification and DNA reduction at the same time as cell debris is removed.

25 For efficient host protein and DNA precipitation, the pH is generally between about 4 to 10, and preferably no more than about pH 9, as the molecule becomes less charged above this pH. Ethacridine lactate should more preferably be used in the about pH 5-9 range when purifying polypeptide. To improve the purification, a short incubation at elevated temperature can be performed. However, stability at elevated temperatures has to be determined for the particular polypeptide of interest to avoid precipitation of the target polypeptide. Also the quality of the recovered target polypeptide has to be investigated to confirm that 30 no modifications of the target polypeptide take place.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for purifying a desired heterologous polypeptide from microbial fermentation broth or homogenate in which it is produced and solubilized comprising adding to the broth or homogenate an effective amount of a solution of ethacridine lactate to precipitate host cell impurities under conditions wherein the majority of the polypeptide remains soluble, and separating the desired polypeptide from the broth or homogenate, wherein the temperature of the broth or homogenate after addition of the ethacridine lactate is from about 50 to 65°C held for about 1 to 60 minutes.
5
2. The method of claim 1 wherein the broth or homogenate is from yeast or prokaryotes.
3. The method of claim 1 or claim 2 wherein the broth or homogenate is from bacteria.
- 10 4. The method of any one of claims 1 to 3 wherein the broth or homogenate is from eubacteria.
5. The method of any one of claims 1 to 4 wherein the broth or homogenate is from gram-negative bacteria.
6. The method of any one of claims 1 to 5 wherein the broth or homogenate is from *E. coli*.
7. The method of any one of claims 1 to 6 wherein the polypeptide is separated from homogenate.
- 15 8. The method of any one of claims 1 to 7 wherein the polypeptide has a higher pI than the average pI of host proteins contained in the host cell impurities.
9. The method of any one of claims 1 to 8 wherein the polypeptide has a pI of at least about 7.
10. The method of any one of claims 1 to 9 wherein the polypeptide is a recombinant polypeptide.
11. The method of any one of claims 1 to 10 wherein the polypeptide is an antibody.
- 20 12. The method of any one of claims 1 to 11 wherein the polypeptide is a humanized antibody.
13. The method of any one of claims 1 to 12 wherein the polypeptide is a full-length antibody.
14. The method of any one of claims 1 to 12 wherein the polypeptide is an antibody fragment.
15. The method of claim 14 wherein the polypeptide is an antibody fragment comprising a light chain.
16. The method of claim 15 wherein the polypeptide is an antibody fragment comprising a kappa light chain.
25
17. The method of any one of claims 14 to 16 wherein the polypeptide is a Fab, Fab', F(ab')₂, or F(ab')₂-leucine zipper fusion.
18. The method of any one of claims 14 to 17 wherein the polypeptide is F(ab')₂.
19. The method of any one of claims 1 to 11 wherein the polypeptide is an anti-IgE, anti-CD18, anti-VEGF,
30 anti-tissue factor, 2C4, anti-Her-2, anti-CD20, anti-CD40, or anti-CD11a antibody or antibody fragment.
20. The method of any one of claims 1 to 11 or 19 wherein the polypeptide is anti-CD18 F(ab')₂, anti-tissue factor F(ab')₂, full-length anti-tissue factor antibody, or anti-VEGF antibody.
35
21. The method of any one of claims 1 to 20 wherein the concentration of ethacridine lactate is about 0.1-5% weight/volume.
22. The method of any one of claims 1 to 21 wherein the concentration of the ethacridine lactate is about 0.4-5% weight/volume.
23. The method of any one of claims 1 to 22 wherein the concentration of the ethacridine lactate is about 0.6-5% weight/volume.

2004230670 28 Oct 2010


24. The method of any one of claims 1 to 23 wherein the conductivity of the broth or homogenate after addition of the ethacridine lactate is about 1-15 mS/cm.
25. The method of any one of claims 1 to 24 wherein the pH of the broth or homogenate after addition of the ethacridine lactate is about 5 to 9.
- 5 26. The method of any one of claims 1 to 25 wherein the pH of the broth or homogenate after addition of the ethacridine lactate is about 6 to 9.
27. The method of any one of claims 1 to 26 wherein the separation is accomplished by centrifugation or filtration.
- 10 28. The method of any one of claims 1 to 27 wherein after the polypeptide is separated from the broth or homogenate, it is further purified by subjection to chromatography or filtration.
29. The method of any one of claims 1 to 28 wherein the polypeptide is produced in a soluble fraction prior to addition of the ethacridine lactate.
- 15 30. The method of any one of claims 1 to 28 wherein the polypeptide is insoluble and is dissolved by contacting it with a solubilizing agent prior to addition of the ethacridine lactate.
31. A microbial cell fermentation broth or homogenate comprising ethacridine lactate of greater than 0.1% (weight/volume) and a polypeptide heterologous to the cells, wherein the pI of the polypeptide is greater than the average pI of the host proteins contained in the cells, wherein the broth or homogenate is at a temperature of from about 50 to 65°C, and wherein the polypeptide is an antibody or antibody fragment.
32. The broth or homogenate of claim 31 that is from bacterial cells.
- 20 33. The broth or homogenate of claim 31 or claim 32 that is from *E. coli*.
34. The broth or homogenate of any one of claims 31 to 33 wherein the polypeptide is a recombinant polypeptide.
35. The broth or homogenate of any one of claims 31 to 34 wherein the polypeptide is F(ab')₂.
36. The broth or homogenate of any one of claims 31 to 34 wherein the polypeptide is anti-IgE, anti-CD18,
- 25 anti-VEGF, anti-tissue factor, 2C4, anti-Her-2, anti-CD20, anti-CD40, or anti-CD11a antibody.
37. The broth or homogenate of any one of claims 31 to 34 or claim 36 wherein the polypeptide is anti-CD18 F(ab')₂, anti-tissue factor F(ab')₂, full-length anti-tissue factor antibody, or anti-VEGF antibody.
38. The broth or homogenate of any one of claims 31 to 37 wherein the polypeptide is soluble in the broth or homogenate.
- 30 39. The broth or homogenate of any one of claims 31 to 38 wherein the concentration of the ethacridine lactate is about 0.1-5% weight/volume.
40. The broth or homogenate of any one of claims 31 to 39, wherein the concentration of ethacridine lactate is about 0.4-5%.
41. The broth or homogenate of any one of claims 31 to 40, wherein the concentration of ethacridine lactate is about 0.6-5%.
- 35 42. The broth or homogenate of any one of claims 31 to 41, wherein the conductivity is less than 16 millisiemens (mS/cm).
43. The broth or homogenate of claim 42, wherein the conductivity is about 1-15 mS/cm.
44. The broth or homogenate of claim 43, wherein the conductivity is about 1-10 mS/cm.
- 40 45. The broth or homogenate of claim 44, wherein the conductivity is about 1-5 mS/cm.

46. The broth or homogenate of any one of claims 31 to 45, wherein the pH ranges from about 4-10.
47. The broth or homogenate of claim 46, wherein the pH is about 4-9.
48. The broth or homogenate of claim 47, wherein the pH is about 5-9.
49. The broth or homogenate of claim 48, wherein the pH is about 6-9.
50. The broth or homogenate of claim 49, wherein the pH is about 6-7.

5 51. A method according to claim 1 or a microbial cell fermentation broth or homogenate according to claim 31, substantially as hereinbefore described with reference to any one of the examples or figures.

Figure 1: Schematic of the anti-CD18 Fab'2 (-leucine zipper) Plasmids

pS1130



pxCD18-7T3




Figure 2
Anti-CD18-7T3.DNA

GAATTCAACTTCCATACTTGGATAAGGAAATACAGACATGAAAATCTCATTCGCTGA
GTGTTTATTTAACGCTGCCAAAAAGAAGAGCTGAACTGTGCGCAGGTAGA
AGCTTGGAGAATTATCGTCACTCAATGCTGCCAATATGCCAAATGACCAACAGG
GTGATTGATCAGGTAGAGGGGGCTGCTACCGAGTAAAGCCCGATGCCAGGATTCTGTA
CGACGATACGGAGCTGCTGCCGATTACGTAAGAGTTATTGAAGCATTCTCGTCAGTA
AAAACTTAATCTTCAACACCTGTCATAAAAGTTGTCACGCCAGACTTATAGTCGCTT
TGTGTTTATTTTAAATGTTACGTAAGTCACTGTAAGGTTACGTTAAAGGTTATCTA
GAATTATGAAAGAATATCGCATTCTCTCGCATCTAATGTCGTTTTTCTATIGCTA
CAAACGCCATACCGTATATCCAGTGACCCAGTCCCGAGCTCCCTGTCGCCCTGTC
GCAGTAGGGTACCATCTGCTGCTGAGGACATCAACAAATTATCTGAACCTG
ATCACACAGAAACCGAAAAGCTCCGAAACTACTGATTACTATACCTCCACCCCTCCACT
CTGGAGTCCTCTCGCTCTGTTCTGGGACGGATTACACTCTGACCATCA
GCACGACCCGAGGAGACTTCGCAACTTAACTGTCAGCAAGGTTAAACTCTGCC
CGAGCTTGGAGAGGAGATCAAACGAACTGTGCTGCCACCATCTGTC
TCATCTCCGCCATCTGAGACGACTCTGAACTCTGAACTGCTCTGTC
TGAATTAACGCTTCCAGGAGGCTGAGCTGAGGAGGCTGAGCTGCT
CGGTTAACCTCCAGGAGGAGTGTACAGAGCAGGACAGGACACCTACACCTCA
GCACGACCCGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGG
TCAACCATCAGGGCGACGCCATGTCGCGACCTCGTGTACCGGGATCTAGGCC
TAAATCTCTGAGGCCGACGCCATGTCGCGACCTCGTGTACCGGGATCTAGGCC
CTCGGTTGCCGCCGGCGTTTTTATTGTTGCCGAGGCCATCTGACTGACGGTGGC
CAATGCTTCTGCTGAGGCCATGCCGAGGCCATGCCGAGGCCATGCCG
CGGTTAACCTCCAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGG
CACTGCTAATTCTGCTGCTCAAGGGCCACTCCGTTCTGGATAATGTTTTGCGCG
ACATCAGTAAACGCTCTGGCAAATTCGAAATGAGCTGAGCTGAGCTGAGCT
AGTTAAATGTTGAGGATAACAAATTAAAGCTTAGGATCTAGAATTATGAG
AAGAATATGCTCTTACTGCTCTATGTTGTTCTATGTTGCTATAGCTAACACCG
GCTGAGGTTCACTGCTGGAGCTGGCGTGGCTGAGGCCAGGGGCTACTCCG
TTGTCCTGCAACTTCTGCTCACCTTTACCGAATACACTATGCACTGGATGCGTC
GCCCGGGTAAGGCCTGGAATGGGTTGCAAGGGATAATCTAAACGGTGGTACACGG
CACACCCAGAGGTTATGAGGACCTTACTATAACGGTAAGATAAAATCCACAGGTAACGG
TACATGCAAATGAAAGCCTGCTGCTGAGGACACTGCGCTCTATTATTGTC
CGACCCCTGAACTACGGCTTGAACGCTCGTATTGAGCTGCTGGGTCAGGAACCC
GTACCCGCTCTCCCGCCCTCACCAAGGGCCATCGGTCTCCCCCTGGCACCCCTCC
AAGAGCACCTCTGGGGCAACGGGCCCTGGCTCTGCTCAAGGACTACTCTCC
CCGGTGAAGGTCTGGAGACTCAGGGGCCCTGACCAAGGGCGTGCACACCTCTCC
GTCTACAGTCTCAGGACTCTACCTCCCTCAGCACGGTGGTGAACGGTGGCCCTCAGCAGC
TTGGGCAACCCACACTCATCTCAACCGTGAATCACAAGCCAGCAACACCAAGGTC
AAGAAAGTTGAGCCAAATCTGACAACACATGCCGCCGTGCCAGC
GAACCTGCTGGGGCGCCGCTGAACACGCTAGGGAGACAAGGTCGAAGAGCTACTCTCAAG
AACTACACCTAGAGAATGAAGTGGCAAGACTCACAACGCTCTGGGGAGCGCTAAGCA
TGGGACGCCCTAAGCTCCCTAAGCTCGGTTGCCGCCGGCTTTTTTATGTTAATC
ATGTTGACGTTATCATGATAAGCTTTAATGCGGTAGTTTACACAGTTAAATGCT
AACGCACTGAGGCCACCGCTATGAAATCTAACAAATGCGCTCATGTCATCTCG
TCACCCCTGAGTCGTAGGCATAGGCTTGGTTATGCCGTACTGCCGGCTCTGGCGG
ATATCGTCCATTCCGACAGCATGCCAGTCATACTGGCTGCTAGCGTATATGCGT

(SEQ ID NO:1)

Figure 3
Anti-CD18-7T3.Protein

FIG. 3A: STII + Anti-CD18 light chain

MKKNIAFLLASMFVFSIATNAYADIONTOSPSSLASASVGDRVTITCRASQDINNYLNWY
QQKPGKAFKLLIYYTSTLHSGVPSRFSGSGSGTDTYTLTISSLQPEDFATYYCQQQGNTLP
PTFGQQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWVKVDNAL
QSGNSQESVTEQDSKDKDSTSLSSTLTLASKADYEKKVYACEVTHQGLSSPVTKSFNRGE
C

(SEQ ID NO:2)


FIG. 3B: STII + Anti-CD18 heavy chain

MKKNIAFLLASMFVFSIATNAYAEVQLVESGGGLVQPGGSLRLSCATSGYIPTTEYTMHW
MRQAPGKGKLEWVAGINPKNGGTSHNQRFMDRFTISVDKSTSTAYMQMNSLRAEDTAVYY
CARWRGLNYGFDVRYFDVWGQGTIVTVSSASTKGPSPVFLAPSSKSTSGGTAA^{IL}GCLVK
DYFPEPVITWSNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLGGRRMKQDEDKVEELLSKNYHLENEVARLKK
LVGER

(SEQ ID NO:3)

Figure 4: Schematic of the anti-Tissue Factor IgG1 Plasmids

paTF130



pxTF-7T3FL




Figure 5
Anti-TF-7T3FL.DNA

GAATTCAACTCTCCATACCTTGGTAAGGAAATACAGACATGAAAAATCTCATTCGCTGA
GTGTTATTAATTCGCCCCAAAAGAACAGAGCTCGAATGACTGTGCGCAGGTAGA
ACGTTTGGAGATATTCTGCATCTGCAATGCTTCGCAATATGGCCAAAATGACCAACAGG
GTGATTGATACAGCTAGGGGGCGCTGTACAGGTAAGGCCGATGCGCAGCATTCCTGGA
CGACGATACAGGACTCTGGCGGATACAGTAAAGAGTATTGGAAAGCTATCTTCCTGACTA
AAAGATTAATCTTCAACAGCTGATCAAGATGCTACGGGAGACATATTAGCTGTT
TGTGTTTATTTTAAATGTTGTAACTAGTACGCAAGTTCACGTAAAAGGGTATCTA
GAATTATGAAAAGAATATCCGATTCCTCTTCATCTCATCTATGTTGCTTTTCTCTGTA
CAAAACCGCTACCGCTGATATCCGATCACCGAGCTCCCGAGCTCTGGCTCCGCTCTGTTG
GCGATAGGCCTCACCATCACCTGCAGGCCAGTCCGAGCATCAAGACCTATCTGACTGTT
ATCACAGAACAGGACAAAGCTCAGTACTGTTATTAATCTGCTACTAGTCTGCGTGTG
AAGGGACTCTCTCGCTTCTGGATCTGGCTTCTGGAGCGGATACACTACGACCATCA
GCAGCTCTGCGAGCAAGAACCTTCGCAACTTATTAATCTGTCAGCAGCGAGAGTCCTCAT
GCACATTTGCAGRCGGGTACCAAGGTGGAGCATAAACGAACTCTGGCTCAGCATCTGCT
TCATCTTCGGCCATCTGATGAGCAGTCTAAATCTGGAACTCTCTGTTCTGCTGCTTC
TGAATTAACCTCTATCCCGAGGAGGCCCCAAAGTACAGTGGAGGTGGATAACGCCCTCCAAAT
CGGTTACCTCTCCAGGAGGTGTCAGAGCAGCACGAGCAACGACACTACAGCTCA
GCAGCACCTGAGCTGAGCAAAGCACACTACAGAACACAAACTCTACGGCTCGAAG
TCACCCATCTGGCCCTGAGCTCGGCGCTAACAAAGGACTTCACAGGGGAGAGGTTAAAT
TAATCTCTACGGCCGAGCAGCTCGGCGAGCTCGGAGCTGGTACCCGGGGATCTGGCTAACG
CTCGGTTGCCGCCGGCTTTTTATGTTGCTGGAGCAGCTCGTACCTGACCTGACCGTC
CAATGCTCTTCGGCTGAGGCCATCGGAGCTGGTGTGTTGCTGAGGTGCTGAAAT
CACTCTCAATTCTGGCTGCTGCTCAAGGCCACTCTGGTGTGATAATTGTTTCTGGCCG
ACATCATAACGGTTCTGGCAAATATTCTGAAATGAGCTGTTGCAAACTTATCATCGAAT
AGTTTAAATGTTGAGGAAATGGTGGAGCCTAACTAATTAAGCTTGTGAGGTTCTAGAAATTATGAG
AAGAATATTGCTGCCCTTACTTGGCTCTATGTTGTTCTGTTCTATGCTCAAAACGGCTG
GCTGAGGTCTCAGTGGTGGAGCTGGCGGTGGCTGGTCAGGCCAGGGGCTACTCCGT
TTGCTCTGTCAGCTTCTGGCTTAATTAATTAGGACTACTACATCTACGGTCTGGCTCAG
GCCGCCGGTAAGGGCTGGTGAATGGTTGATTGATTGATTCAGGACAGCAAGGCAACGATC
TATGACCCCAAGTTCAGGAGCCTGCCACTATTAAGCGCTGACAAATTCCAAAACACAGCA
TACCTGAGATGAAACAGCTCTGGCTTGAGGACACTCTGGCTTATTAATTGTTGTCGAGAC
ACGGCCGCTTACTTCGACTCTGGGTCAAGGGACCTCTGGTCTACGGCTCTCCCTTC
ACCAAGGGGCCATCGGTTCTCCCCCTGGCACCTCTCCAGGACCTCTGGGGGACA
GCCGCCCTGGCTCGCTGGCTGAAGGACTACCTTCCCGAACGGCTGAGCTGCTGGTGAAC
TCAGGCCGCTTACCAAGCGGCCGTCACACTCTGGCTGCTACAGTCTCAGGACT
TACTCTCTGAGCAGCGCTGGTGTGACTGCTCCCTAGAGCTTGGCAGGCCAGACCTACATC
TCAGAACGCTGAACTACACAGCCCAACACAGGTTGGAAGAAGTTGAGGCCAAATCT
TGTGACAAAATCTACACATGCCAGCTGGCCACCTGCAACTCTGACTCTGGGGGAGGCTTC
GCTCTCTCTTCTCCCCAAAAGACCGAGACACCTCTGGTCAATGCTCCGGACCTGGAGGT
ACATGGCTGGTGTGGAGCTGGAGCTGGACAGAACAGGACCTGAGGTCAAGTCTAACCTG
GACGGCGTGGAGGTGTCATAATGCCAGAACAAAGCCGGGGAGGACACTACACACAGC
TACCTGCTGGTGTGAGCTCTCCACGGCTCTGACAGGACTCTGGTGAATGGCAAGGACTAC
AGATGCAAGGTCTTACACAGGCTCTCCAGGCTCTGGTGTGAGGACTCTGGTGAATGGCAAGGACTAC
AAAGGGCAGCCGAGAACCAACGGCTACACCTGGCCCTACGGGAAAGAGTACGCC
AAGAACAGGACTCAGCGCTGAGCTGGTGTGAAAGGCTCTTATCCGGGACATGGAC
GAGTGGAGGAGCAATGGCAGGCCGGAGAACAACTACAGAACGACCCCTGGCTGGAC
TCCGACGGCTCTTCTCTGGTGTGAGCTGGTCAAGTGGCAACGAGCAGGGTGGCAG
GGAAAGCTGTTCTCTGGTGTGAGCTGGTCAAGTGGCAACGACCAACCTGGCAG
AGGCTCTCTCTCTGGTGTGAGCTGGTCAAGTGGCAACGACCAACCTGGCAG
TTGCGCGGGCTTTTATGTTAATCTGAACTCATGTTGAGCTTACATGCTTACAGCT
AATGGCTGGTGTGTTTACACATTAATGCTAACGGAGCTGGTGTGAGGCTTACAGCT
ACATGCGCTTACATGCTGCTTACAGTGGTCAAGTGGCAACGACCAACCTGGCAG

(SEQ. TD. NO. 4)

Figure 6
Anti-TF-7T3FL Protein

FIG. 6A: STII + Anti-TF light chain

```

MKKNIAFLLASMFVFSIATNAYADIQMTQSPSSLSASVGDRVTITCRASRDIKSYLNWY
QQKPGKAPKVLIIYYATSLAEGVPSRFSGSGSTDTLTISSLQPEDFAIYYCLOQHGESP
WTFQCGTKVEIKRTVAAAPSVFIFPPSDEQLKSGTASVVCLLNNFPREAKVQWKVDNAL
QSGNSQESVTEQDSKDKSTYSLSSTLTLSKADYEHKVKVYACEVTHQGLSSPVTKSFNRGC
C

```

(SEQ ID NO:5)

FIG. 6B: STII + Anti-TF heavy chain

```

MKKNIAFLLASMFVFSIATNAYAEVQLVESGGGLVQPGGSLRLSCAASGFNIKEYYMHW
VHQAPGKGLEVGLIDPEQGNTIYDPKFQDRATISADNSKNTAYLQMINSLRAEDITAVYY
CARDTAAYFDYWGQGTIVTVSSASTKGEPSVFLAPPSKSITSGGTAAALGCLVKDYPEPV
TVSNMSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHHKESNTVKDK
KVEPKSCDKTHICPPCPAPELLGGPSVFLFPPKPKDTLMSRTFETVTCVVVDVSHEDPE
VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAP
IEKTIISAKAKGQFRREPQVYTLPPSEEMTKNQVSLTCLVKGFYPSDIIAVEWESNGQFENN
YKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHALHNHYTQKSLSLSPGK

```

(SEQ ID NO:6)

Fig. 7





Fig. 8



anti-CD18 F(ab')₂
FIG 8A

anti-TF F(ab')₂
FIG 8B

anti-TF (Ab)
FIG 8C

Fig. 9




Fig. 10




Fig. 10A

Fig. 10B

Fig. 11




Fig. 12





Fig. 13



2004230670 15 May 2007

P1941R1PCT.txt
Sequence Listing

<110> GENENTECH, INC.
<120> PURIFICATION OF POLYPEPTIDES
<130> P1941R1 PCT
<140> PCT/US2004/000499
<141> 2004-01-08
<150> US 60/439,418
<151> 2003-01-09
<160> 6
<210> 1
<211> 2700
<212> DNA
<213> Artificial sequence
<220>
<223> Sequence is synthesized
<400> 1
gaattcaact tctccatact ttggataagg aaatacagac atgaaaaatc 50
tcattgctga gttgttattt aagcttgccc aaaaagaaga agagtcgaat 100
gaactgtgtc cgccggatgt a gctttggag attatcgta ctgcaatgct 150
tcgcaatatg gcgaaaaatg accaacacgc gttgattgtat caggttaggg 200
gggcgtgtta cgaggtaaag cccgatgcca gcattcctga cgacgatacg 250
gagctgtgc gcgattacgt aaagaagtta ttgaagcatac ctcgtcagta 300
aaaagttaat ctttcaaca gctgtcataa agttgtcactg gcccggactt 350
atagtcgtttt tttttatgtt tttttaatgtt tttttaatgtt gtacgcaagt 400
tcacgtaaaa agggtatcta gaattatgaa aaagaatatac gcatttccttc 450
ttgcatttatg tttttttttt tttttaatgtt tttttaatgtt gtacgcaagt 500
cagatgaccc agtccccgag ctccctgtcc gcctctgtgg gcgatagggt 550
caccatcacc tttttttttt tttttaatgtt tttttaatgtt gtacgcaagt 600
atcaacagaa accaggaaaa gctccgaaac tactgattta ctataccctcc 650
accctccact ctggagtcctt ttctcgcttc tctgggttctg gttctgggac 700
ggattacact ctgaccatca gcagtctgca accggaggac ttgcactt 750
attactgtca gcaaggtaat actctgcccgc cgacggttgc acagggcacg 800
aagggtggaga tcaaacgaac tttttttttt tttttaatgtt tttttaatgtt gtacgcaagt 850
gcattctgtat gacgtttttt tttttttttt tttttaatgtt tttttaatgtt gtacgcaagt 900
tgaataactt ctatcccaaga gaggccaaag tacagtggaa ggtggataac 950
gcctccaat cgggttaactc ccaggagatgt gtcacagagc aggacagcaa 1000
ggacagcacc tacagcctca gcagcaccct gacgctgagc aaagcagact 1050
acgagaaaca caaagtctac gcctgcaag tcacccatca gggcctgagc 1100

Page 1

2004230670 15 May 2007

P1941R1PCT.txt

tcgccccgtca caaagagctt caacaggggg gagtgttaat taaatccctct 1150
acgccggacg catcggtggcg agctcggtac ccggggatct aggcctaacg 1200
ctcggttgcc gccggggcggtt ttttattgtt gccgacgcgc atctcgactg 1250
cacgggtcac caatgcttct ggcgtcaggc agccatcgga agctgtggta 1300
tggctgtgca ggtcgtaaat cactgcataa ttctgtcgcc tcaaggcgca 1350
ctccccgttct ggataatgtt ttttgcggcg acatcataac gggttctggca 1400
aatattctga aatgagctgt tgacaattaa tcatcgaact agttaatgt 1450
gtggaattgt gaggcgataa caatthaagct taggatctag aattatgaag 1500
aagaatattg cgttcctact tgccctatgt tttgtcttt ctatagctac 1550
aaacgcgtac gctgagggttc agctgggtga gtctggcggt ggcctggc 1600
agccaggggg ctcactccgt ttgtccctgtg caacttctgg ctacacccctt 1650
accgaataaca ctagcactg gatgcgtcag gccccggta agggcctggta 1700
atgggttgca gggattaatc ctaaaaaacgg tggtaccagc cacaaccaga 1750
ggttcatgga ccgtttcact ataagcgtag ataaatccac cagtacagcc 1800
tacatgcaaa tgaacagcct gcgctgctgag gacactgccc tctattattg 1850
tgcttagatgg cgaggcctga actacggctt tgacgtccgt tattttgacg 1900
tctggggtca aggaaccctg gtcaccgtct cctcggccctc caccaggc 1950
ccatcggtct tccccctggc acccccttcc aagagcacct ctgggggcac 2000
agcggccctg ggctgctgg tcaaggacta cttcccccga ccggtgacgg 2050
tgtcgtggaa ctcaggcgcc ctgaccagcg gctgtcacac cttcccgct 2100
gtcctacagt cctcaggact ctactccctc agcagcgtgg tgaccgtgcc 2150
ctccagcagc ttgggcaccc agacctacat ctgcaacgtg aatcacaagc 2200
ccagcaacac caaggtcgac aagaaaagtt agcccaaatac ttgtgacaaa 2250
actcacacat gccccgggtg cccagcacca gaactgctgg gcggccgcat 2300
gaaacagcta gaggacaagg tcgaagagct actctccaag aactaccacc 2350
tagagaatga agtggcaaga ctcaaaaagc ttgtcgggga gcgctaagca 2400
tgcgacggcc ctagagttccc taacgctcggtt tgccgcccgg gcgttttta 2450
ttgttaactc atgtttgaca gcttatcatc gataagctt aatgcggtag 2500
tttatcacag ttaaattgtct aacgcgtca ggcaccgtgt atgaaatcta 2550
acaatgcgtct catcgtcatc ctggcaccgg tcacccctggta tgctgttaggc 2600
ataggcttgg ttatgcccgt actgcccggc ctcttgcggg atatcgcca 2650
ttccgacagc atcgccagtc actatggcgt gctgctagcg ctatatgcgt 2700

<210> 2
<211> 237
<212> PRT

2004230670 15 May 2007

P1941R1PCT.txt

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 2

Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15
Ser Ile Ala Thr Asn Ala Tyr Ala Asp Ile Gln Met Thr Gln Ser
20 25 30
Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr
35 40 45
Cys Arg Ala Ser Gln Asp Ile Asn Asn Tyr Leu Asn Trp Tyr Gln
50 55 60
Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Tyr Thr Ser
65 70 75
Thr Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
80 85 90
Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp
95 100 105
Phe Ala Thr Tyr Tyr Cys Gln Gln Gly Asn Thr Leu Pro Pro Thr
110 115 120
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
125 130 135
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
140 145 150
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
155 160 165
Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
170 175 180
Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
185 190 195
Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
200 205 210
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
215 220 225
Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
230 235

<210> 3

<211> 300

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 3

Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15
Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Glu Ser
20 25 30

Page 3

2004230670 15 May 2007

P1941R1PCT.txt

Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
35 40 45
Ala Thr Ser Gly Tyr Thr Phe Thr Glu Tyr Thr Met His Trp Met
50 55 60
Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Gly Ile Asn
65 70 75
Pro Lys Asn Gly Gly Thr Ser His Asn Gln Arg Phe Met Asp Arg
80 85 90
Phe Thr Ile Ser Val Asp Lys Ser Thr Ser Thr Ala Tyr Met Gln
95 100 105
Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
110 115 120
Arg Trp Arg Gly Leu Asn Tyr Gly Phe Asp Val Arg Tyr Phe Asp
125 130 135
Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr
140 145 150
Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr
155 160 165
Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe
170 175 180
Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
185 190 195
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr
200 205 210
Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr
215 220 225
Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys
230 235 240
Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr
245 250 255
Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Arg Met Lys
260 265 270
Gln Leu Glu Asp Lys Val Glu Glu Leu Leu Ser Lys Asn Tyr His
275 280 285
Leu Glu Asn Glu Val Ala Arg Leu Lys Lys Leu Val Gly Glu Arg
290 295 300

<210> 4
<211> 3100
<212> DNA
<213> Artificial sequence

<220>
<223> Sequence is synthesized

<400> 4
gaattcaact tctccatact ttggataagg aaatacagac atgaaaaatc 50
tcattgctga gttgttattt aagcttgccc aaaaagaaga agagtcgaat 100

2004230670 15 May 2007

P1941R1PCT.txt
gaactgtgtg cgccaggtaga agctttggag attatcgta ctgcaatgct 150
tcgcaatatg ggcggaaaatg accaacagcg gttgattgat caggttaggg 200
gggcgcgtgtc cgaggtaaag cccgatgcca gcattcctga cgacgatacg 250
gagctgctgc gcgattacgt aaagaagttt ttaaagcattt ctcgtcgtt 300
aaaaggtaat ctttcaaca gctgtcataa agttgtcagc gcccggactt 350
atagtcgttt tggttttatt ttttaatgtt tttgtacta gtacgcaagt 400
tcacgtaaaa agggtatcta gaattatcaa aaagaatatc gcatttccttc 450
ttgcatttatg ttctattgtt cttttttttt cttttttttt 500
cagatgaccc agtccccgag ctccctgtcc gcctctgtgg gcgatagggt 550
caccatcacc tgcagagcca gtcgcgacat caagagctat ctgaaactgg 600
atcaacagaa accaggaaaa gctccgaaag tactgattt ctatgctact 650
agtctcgctg aaggagtccc ttctcgcttc tctggatccg gttctggac 700
ggatttactt ctgaccatca gcagtctgca gccagaagac ttgcacactt 750
attactgtct tcagcacgga gagtctccat ggacatttgg acagggtacc 800
aagggtggaga tcaaaacgaac tttttttttt tttttttttt 850
gcctctgtat gagcagtttta aatctggaaac tttttttttt tttttttttt 900
ttaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac 950
gcctccaaat cgggttaactc ccaggaggt gtcacagagc aggacagcaa 1000
ggacagcacc tacagcctca gcagcacctt gacgctgagc aaagcagact 1050
acgagaaaca caaagtctac gcctgcgaag tcacccatca gggcctgagc 1100
tcgcccgtca caaagagttt caacagggga gagtgttaat taaatccctct 1150
acgcccggacg catcggtggcg agctcggtac ccggggatct aggcctaact 1200
ctcggttgcc gcccggcggtt tttttttttt gcccggcggtt atctcgactt 1250
cacgggtgcac caatgttttctt ggcgtcaggc agccatcgga agctgtggta 1300
tggctgtgca ggtcgtaat cactgcataa ttctgtgtcgc tcaaggcgca 1350
ctccctgttctt ggataatgtt ttttgcggc acatcataac ggttctggca 1400
aatattctgtt aatgagctgt tgacaattaa tcatcgactt agtttaatgtt 1450
gttggattgtt gacggataa caatggatctt taggatctttagt aattatgtt 1500
aagaatatttgcgtt cttttttttt tttttttttt tttttttttt 1550
aaacgcgtac gctgagggttc agctgggtggaa gtctggcggtt ggcctgggtc 1600
agccagggggg cttttttttt tttttttttt tttttttttt 1650
aaggagttttttttt tttttttttt tttttttttt tttttttttt 1700
atgggttggaa ttgttggatc cagagcaagg caacacgatc tatgacccgaa 1750
agttccagga ccgtgcactt ataagcgctg acaattccaa aaacacagca 1800

Page 5

2004230670 15 May 2007

P1941R1PCT.txt
tacctgcaga tgaacagcct gcgtgcttag gacactgccc tctattattg 1850
tgctcgagac acggccgctt acttcgacta ctggggtaa ggaaccctgg 1900
tcaccgtctc ctcggctcc accaaggccc catcggtctt cccccggca 1950
ccctccctca agagcaccc tggggcaca gcggccctgg gctgcttggt 2000
caaggactac ttccccgaac cggtgacggt gtcgtggac tcaggcgccc 2050
tgaccagcg cggtcacacc ttcccgctg tcctacagtc ctcaggactc 2100
tactccctca gcagcgtggt gactgtgccc tctagcagct tgggcaccca 2150
gacctacatc tgcaacgtga atcacaagcc cagcaacacc aagggtggaca 2200
agaaaatgtga gcccaaatct tgtgacaaa ctcacacatg cccaccgtgc 2250
ccagcacctg aactccctggg gggaccgtca gtctccctt tccccccaaa 2300
acccaaggac accctcatga tctcccgac ccctgaggtc acatgcgtgg 2350
tggtggacgt gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg 2400
gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta 2450
caacagcacg taccgtgtgg tcagcgtctt caccgtctg caccaggact 2500
ggctgaatgg caaggagtag aagtgcagg tctccaacaa agccctccca 2550
gcccccatcg agaaaaccat ctccaaagcc aaagggcagc cccgagaacc 2600
acaggtgtac accctgcccc catcccggga agagatgacc aagaaccagg 2650
tcagcctgac ctgcctggtc aaaggcttct atcccagcga catcgccgtg 2700
gagtggaga gcaatggca gccggagaac aactacaaga ccacgcctcc 2750
cgtgctggac tccgacggct ccttcttctt ctacagcaag ctcaccgtgg 2800
acaagagcag gtggcagcag gggAACGTCT tctcatgctc cgtatgtcat 2850
gaggctctgc acaaccacta cacgcagaag agcctctccc tgcgtccggg 2900
taaataagca tgcgacggcc cttagtccc taacgctgg ttgcgcgg 2950
gcgtttttta ttgttaactc atgtttgaca gcttatcatc gataagcttt 3000
aatgcggtag ttatcacag tttaattgtct aacgcagtca ggcaccgtgt 3050
atgaaatcta acaatgcgtc catcgatc ctcggcaccg tcaccctgga 3100
<210> 5
<211> 237
<212> PRT
<213> Artificial sequence
<220>
<223> Sequence is synthesized
<400> 5
Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15
Ser Ile Ala Thr Asn Ala Tyr Ala Asp Ile Gln Met Thr Gln Ser
20 25 30
Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr
Page 6

2004230670 15 May 2007

P1941R1PCT.txt

35

40

45

Cys Arg Ala Ser Arg Asp Ile Lys Ser Tyr Leu Asn Trp Tyr Gln
50 55 60
Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Tyr Ala Thr
65 70 75
Ser Leu Ala Glu Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
80 85 90
Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp
95 100 105
Phe Ala Thr Tyr Tyr Cys Leu Gln His Gly Glu Ser Pro Trp Thr
110 115 120
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
125 130 135
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
140 145 150
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
155 160 165
Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
170 175 180
Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
185 190 195
Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
200 205 210
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
215 220 225
Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
230 235

<210> 6

<211> 470

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized

<400> 6

Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15

Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Glu Ser
20 25 30

Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
35 40 45

Ala Ala Ser Gly Phe Asn Ile Lys Glu Tyr Tyr Met His Trp Val
50 55 60

Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Leu Ile Asp
65 70 75

Pro Glu Gln Gly Asn Thr Ile Tyr Asp Pro Lys Phe Gln Asp Arg
80 85 90

2004230670 15 May 2007

P1941R1PCT.txt
Ala Thr Ile Ser Ala Asp Asn Ser Lys Asn Thr Ala Tyr Leu Gln
95 100 105
Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
110 115 120
Arg Asp Thr Ala Ala Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu
125 130 135
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
140 145 150
Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
155 160 165
Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
170 175 180
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
185 190 195
Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
200 205 210
Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val
215 220 225
Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro
230 235 240
Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro
245 250 255
Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
260 265 270
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
275 280 285
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
290 295 300
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg
305 310 315
Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
320 325 330
Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
335 340 345
Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser
350 355 360
Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
365 370 375
Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
380 385 390
Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
395 400 405
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
410 415 420
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
425 430 435

Page 8

2004230670 15 May 2007

P1941R1PCT.txt

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
440 445 450
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
455 460 465
Leu Ser Pro Gly Lys
470