



HU000030236T2



(19) **HU**

(11) Lajstromszám: **E 030 236**

(13) **T2**

MAGYARORSZÁG
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM **SZÖVEGÉNEK FORDÍTÁSA**

(21) Magyar ügyszám: **E 12 756223**

(51) Int. Cl.: **A01N 37/42**

(2006.01)

(22) A bejelentés napja: **2012. 09. 10.**

A01N 43/50

(2006.01)

A01P 13/00

(2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20120756223

(86) A nemzetközi (PCT) bejelentési szám:

PCT/EP 12/067652

(97) Az európai bejelentés közzétételi adatai:

EP 2755479 A1 2013. 03. 21.

(87) A nemzetközi közzétételi szám:

WO 13037735

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2755479 B1 2016. 07. 27.

(30) Elsőbbségi adatok:

11181041

2011. 09. 13.

EP

(73) Jogosult(ak):

BASF Agrochemical Products, B.V., 6835 EA Arnhem (NL)

(72) Feltaláló(k):

PFENNING, Matthias, 67365 Schwegenheim (DE)

BREMER, Hagen, 67346 Speyer (DE)

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest

(54) **Eljárás gyomok irtására imazamoxot és növényi növekedést szabályozó anyagokat tartalmazó keverékekkel**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 755 479 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
27.07.2016 Bulletin 2016/30

(51) Int Cl.:
A01N 37/42 (2006.01) **A01N 43/50** (2006.01)
A01P 13/00 (2006.01)

(21) Application number: **12756223.9**(86) International application number:
PCT/EP2012/067652(22) Date of filing: **10.09.2012**(87) International publication number:
WO 2013/037735 (21.03.2013 Gazette 2013/12)(54) **METHOD OF CONTROLLING PARASITIC WEEDS WITH MIXTURES COMPRISING IMAZAMOX AND PLANT GROWTH REGULATORS**

VERFAHREN ZUR BEKÄMPFUNG VON PARASITÄREM UNKRAUT MIT IMAZAMOX UND PFLANZENWACHSTUMSREGULATOREN

PROCÉDÉ DE CONTRÔLE DES HERBES PARASITES AVEC DES MÉLANGES CONTENANT D'IMAZAMOX ET RÉGULATEURS DE CROISSANCE DES PLANTES

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

US-A1- 2010 210 467 US-A1- 2010 304 971
US-A1- 2011 028 324

(30) Priority: **13.09.2011 EP 11181041**

- BEAM J B ET AL: "Interaction of Prohexadione Calcium with Agrichemicals Applied to Peanut (*Arachis hypogaea L.*)", PEANUT SCIENCE, RALEIGH, NC, US, vol. 29, no. 1, 1 January 2002 (2002-01-01), pages 29-35, XP009156683, ISSN: 0095-3679

(43) Date of publication of application:
23.07.2014 Bulletin 2014/30

- WOJCIECH MIZINIAK ET AL: "EFFECT OF SULFOSULFURON + RETARDANTS MIXTURES ON GROWTH AND YIELD OF WINTER WHEAT AND APERA SPICA-VENTI CONTROL", PROGRESS IN PLANT PROTECTION, vol. 48, no. 2, 1 January 2008 (2008-01-01), pages 635-639, XP055019649, DOI: <http://www.progress.plantprotection.pl/pliki/2008/48-2-36.pdf>

(73) Proprietor: **BASF Agrochemical Products, B.V. 6835 EA Arnhem (NL)**

- ROMAN KRAWCZYK: "STUDIES ON EFFECTIVITY ON APPLICATION OF TANK-MIXTURE FLORASULAM WITH GROWTH REGULATORS IN THE WEED CONTROL IN WINTER WHEAT", PROGRESS IN PLANT PROTECTION, vol. 46, no. 2, 1 January 2006 (2006-01-01), pages 200-204, XP055019647, DOI: <http://www.progress.plantprotection.pl/pliki/2006/2/p200-204.pdf>

(72) Inventors:

- PFENNING, Matthias
67365 Schwegenheim (DE)
- BREMER, Hagen
67346 Speyer (DE)

(74) Representative: **BASF IP Association
BASF SE
ZRX-C6
67056 Ludwigshafen (DE)**

(56) References cited:

EP-A1- 0 224 441	EP-A1- 1 088 480
WO-A1-2008/124431	DE-A1- 19 834 627
US-A1- 2005 044 587	US-A1- 2008 269 058

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- Z-W FAN ET AL: "Prohexadione-calcium induces sunflower (*Helianthus annuus*) resistance against the root parasitic weed *Orobanche cumana*", WEED RESEARCH, vol. 47, no. 1, 1 February 2007 (2007-02-01), pages 34-43, XP055019658, ISSN: 0043-1737, DOI: 10.1111/j.1365-3180.2007.00536.x cited in the application

Description

[0001] The present invention relates to a method for controlling parasitic weeds, comprising applying to the host plant, the weeds and/or their habitat herbicidal mixtures or compositions comprising A) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters and B) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl, in a synergistically effective amount. The present invention also relates to a method for improving the yield of the crop plant comprising applying to the host plant, and/or their habitat mixtures comprising components as defined herein. The present invention also relates to herbicidal mixtures comprising imazamox and one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl, compositions comprising said mixtures and their use for the control of parasitic weeds.

[0002] Parasitic weeds derive some or all of their sustenance from another plant by penetrating the host's tissue and drawing nutrients from the host plant (W. Koch, M. Kunisch, PLITS 1989/7 (2), *Principles of Weed Management*, 22-26). This occurs either through the stem, e.g. in *Cuscuta* spp., or from the root tissue, e.g. *Striga* spp. (witchweeds) and *Orobanche* spp. (broomrape). Plants parasitizing on roots cause significant damage in crops before emergence and thus lead to high economical losses. Therefore handpulling or mechanical weeding may prevent the parasite from seed setting, however, these measures come too late to prevent crop losses. The longevity of the seeds of parasitic species is relatively high. Therefore crop rotation or fallow is often not practical since intervals between the cultivation of susceptible crops have to be 5 to 10 years depending on the specific system. Conferring resistance against parasitic weeds on crop plants through selection or genetic modification is only partly successful as parasitic weeds are able to adapt to such changes quickly (Höniges, A., Wegmann, K.* and Ardelean, A. HELIA, 31, Nr. 49, p.p. 1-12, (2008)).

[0003] Plant growth regulators (PGRs) interact with the hormonal system of the treated plants and regulate the growth of a plant or parts of a plant. They affect developmental processes and differentiation in plants at low dosages without having a nutritive value or being phytotoxic. More specifically, various PGRs can, for example, reduce plant height, stimulate seed germination, induce flowering, darken leaf coloring, minimize lodging of cereals, change the rate of plant growth and modify the timing and efficiency of flowering, fruit formation, ripening, fruit drop, defoliation or quality traits.

[0004] There are several different classes of plant growth regulators. Known classes include azoles (such as uniconazole and paclobutrazol), cyclohexane carboxylates (such as trinexapac, trinexapac-ethyl, prohexadione and prohexadione-calcium), pyrimidinyl carbinols (such as flurprimidol and auncyrrnidol), quarternary ammoniums (such as chlormequat-chloride and mepiquat-chloride) and sulphonyl-amino phenyl-acetamides (such as mefluidide).

[0005] Plant growth regulators operate by various modes of action (W. Rademacher, Ann. Rev. Plant Physiol. Plant Biol. 2000, 51, 501-531). Prohexadione, prohexadione-calcium, trinexapac and trinexapac-ethyl act as inhibitors of aminocyclopropane carboxylic acid (ACC) oxidase and hence inhibit the biosynthesis of ethylene.

[0006] Ethylene serves as a hormone in plants (<http://en.wikipedia.org/wiki/Ethylene>, April 2011; Z. Lin et al., "Recent advances in ethylene research", J. Exp. Bot., 2009, 60, 3311-3336). It acts at trace levels within the plant by stimulating or regulating the ripening of fruit as well as senescence of vegetative tissues. Ethylene biosynthesis can be induced or inhibited by plant growth regulators (S. F. Yang et al. "Ethylene biosynthesis and its regulation in higher plants". Ann. Rev. Plant Physiol. 1984, 35, 155-89). As opposed to inhibitors of ethylene biosynthesis, inhibitors of ethylene perception include compounds that have a similar shape to ethylene, but do not elicit the ethylene response. One example of an ethylene perception inhibitor is 1-methylcyclopropene (1-MCP). WO 2008124431 (A1) relates to synergistic mixtures comprising 1-MCP for improving yield, health and/or vigor of plants.

[0007] It has been reported in EP 0123001 (A1) that prohexadione-calcium induces sunflower resistance against infection with the root parasitic weed *Orobanche cumana* (Z.W. Fan et al., Weed Research 2007, 47, 34-43).

[0008] US 2011/0028324, Beam J.B. et al. (Peanut Science, 2002, 29, 29-35), Miziniak W. et al. (Progress in Plant Protection, 2008, 48, 635-839), Krawczyk R. (Progress in Plant Protection, 2006, 46, 200-204), US 2010/304971 (A1), US 2010/210467 (A1), US 2008/0269058 (A1), EP 0 224 441 (A1), DE 198 34 627 (A1) disclose compositions containing acetolactate synthase (ALS) inhibitors and plant growth regulators (PGR) such as prohexadione-calcium, trinexapac or trinexapac-ethyl for combating weeds and for increasing crop yield.

[0009] Acetolactate synthase (ALS) inhibitors are herbicidally active compounds, which inhibit the branched chain amino acid biosynthesis. They belong to the group B of the HRAC classification system. WO 2003/012115 (A2) and WO 2008/124431 (A1) relate to herbicide-resistant sunflower plants and methods for controlling broomrape comprising applying, inter alia, acetolactate synthase inhibitors such as imidazolinones or sulfonylureas. EP 1088480 (A1) relates to the selective use of imazetapyr in varieties of sunflowers that are resistant or tolerant to said herbicide for the control of *Orobanche cernua* or *Orobanche cumana*. US 2005/44587 (A1) relates to sunflower lines and hybrids that are tolerant to sulfonylurea herbicides and to a method for selective control of undesired vegetation, including parasitic weeds, preferably *Orobanche* spp., by applying sulfonylurea herbicides to sulfonylurea-tolerant sunflower crops.

[0010] As to the given mechanisms of action and classification of the active substances, see e.g. "HRAC, Classification of Herbicides According to Mode of Action", <http://www.plantprotection.org/hrac/MOA.html>, April 2011).

[0011] There are only few selective herbicides that control root parasitic weeds season long while they are still underground. In order to prevent damage to the host plant a herbicide that can translocate within the host plant must be applied in concentrations which are non-toxic to the host. For this reason there is a strong need in the agrochemical industry to provide new methods and chemical compositions for the efficient control of root parasitic weeds like, for example, Orobanche spp. and Striga spp..

[0012] It is an objective of the present invention to provide herbicidal mixtures or compositions, which are highly active against parasitic weeds, especially root parasitic weeds. At the same time, the compositions should have good compatibility with useful plants. In addition, the compositions according to the invention should have a broad spectrum of activity and ultimately help increase the yield of the crop plant.

[0013] It was found that herbicidally active mixtures or compositions comprising A) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters and B) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl are very useful for controlling parasitic weeds.

[0014] Surprisingly, the mixtures and compositions according to the invention have better herbicidal activity, i.e. better activity against parasitic weeds, than would have been expected based on the herbicidal activity observed for the individual components. In addition to the improved herbicidal activity the mixtures and compositions according to the present invention also provide for significantly improved yields of the crop plant.

[0015] The herbicidal activity to be expected for mixtures and compositions based on the individual component can be calculated using Colby's formula (see below). If the observed activity exceeds the expected additive activity of the individual components, this effect is referred to as "synergism".

[0016] The present invention relates to a method for controlling parasitic weeds, comprising applying to the host plant, the weeds and/or their habitat a herbicidal mixture comprising components

A) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters, and

B) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl,

in a synergistically effective amount or a composition comprising said mixture.

[0017] In the method according to the invention, the mixtures or compositions comprising said mixtures are preferably applied in a synergistically effective amount, which is defined by a weight ratio of A) to B) from 1 : 300 to 300 : 1, preferably from 1 : 100 to 100 : 1, preferably from 1 : 20 to 20 : 1, preferably from 1 : 3 to 3 : 1, preferably from 1 : 2 to 2 : 1 and also preferably 1 : 1.

[0018] According to one embodiment, the inventive method comprises the application of herbicidal mixtures comprising components A) and B) and as a further ingredient a herbicidal, fungicidal or insecticidal component C).

[0019] Furthermore the present invention also relates to mixtures comprising components

A) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters, and

B) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl.

[0020] Preferably the mixtures comprise components A) and B) in a synergistically effective amount, which is defined by a weight ratio of A) to B) from 1 : 300 to 300 : 1, preferably from 1 : 100 to 100 : 1, preferably from 1 : 20 to 20 : 1, preferably from 1 : 3 to 3 : 1, preferably from 1 : 2 to 2 : 1 and also preferably 1 : 1.

[0021] According to one embodiment of the invention the mixture comprises as a further ingredient a fungicidal, herbicidal or insecticidal component C).

[0022] The term mixture herein relates to combinations of components A), B) and, optionally, C), i.e. active ingredients, which can be applied separately within a time frame that allows simultaneous action of the active ingredients on the plants, preferably within a time-frame of eight weeks, in particular at most 7 days, more particular at most 1 day.

[0023] Examples for suitable components C) which can be used according to the present invention are, for example, strobilurin fungicides, for example, azoxystrobin, coumethoxystrobin, coumoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyribencarb, trifloxystrobin, 2-[2-(2,5-dimethyl-phenoxymethyl)-phenyl]-3-methoxy-acrylic acid methyl ester and 2-(2-(3-(2,6-di-chlorophenyl)-1-methyl-allylideneaminoxyethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide; or azole fungicides such as tebuconazole, metconazole, prothioconazole, difenoconazole, paclobutrazole, flusilazole, cyproconazole, propiconazole, fluquinazole; or succinate dehydrogenase inhibitors such as boscalid, bixafen, fluopyram, penthiopyrad isopyrazam, fluxapyroxad; or carbendazim, thiophanatemethyl, iprodione, chlorothalonil, mancozeb, procymidone, vinclozolin, famoxadone, triadimenol, fenpropimorph, cymoxanil, prochloraz; or herbicides selected

from acetochlor, aclonifen, cycloxydim, tepraloxydim, clethodim, fenoxyaprop, propaquizafop, quizalafop, haloxyfop, pendimethalin, clomazone, metazachlor, Metolachlor, quinmerac, dimethenamid, ethametsulfuron, napropamide, pethoxamide, glyphosate, gluphosinate, sulfentrazone, carfentrazone, trifluralin, flurochloridone, oxadiazone, linuron.

[0024] The active components A), B) and C) that are listed above are known in the art, see, for example, The Compendium of Pesticide Common Names (<http://www.alanwood.net/pesticides/>); Farm Chemicals Handbook 2000 volume 86, Meister Publishing Company, 2000; B. Hock, C. Fedtke, R. R. Schmidt, *Herbicide [Herbicides]*, Georg Thieme Verlag, Stuttgart 1995; W. H. Ahrens, *Herbicide Handbook*, 7th edition, Weed Science Society of America, 1994; and K. K. Hatzios, *Herbicide Handbook, Supplement for the 7th edition*, Weed Science Society of America, 1998.

[0025] In the context of the present invention the following synergistic mixtures as well as the application of said mixtures in the methods according to the invention are particularly preferred. The preferred embodiments of the invention mentioned hereinafter are preferred either independently from each other or in combination with one another.

[0026] According to a preferred embodiment of the invention component A) is imazamox.

[0027] According to a preferred embodiment of the invention component B) is prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl or mixtures thereof.

[0028] According to a preferred embodiment of the invention component B) is prohexadione or prohexadione-calcium or mixtures thereof.

[0029] According to a preferred embodiment of the invention component B) is prohexadione-calcium.

[0030] According to a preferred embodiment of the invention component B) is trinexapac.

[0031] According to a preferred embodiment of the invention component B) is trinexapac-ethyl.

[0032] According to a preferred embodiment of the invention component C) is a fungicidal compound selected from the group of strobilurins such as dimoxystrobin, pyraclostrobin, azoxystrobin, trifloxystrobin, kresoxim-methyl or azole fungicides such as tebuconazole, metconazole, prothioconazole, difenoconazole, paclobutrazole, flusilazole, cyproconazole, propiconazole, fluquinazole; or succinate dehydrogenase inhibitors such as boscalid, bixafen, fluopyram, penthiopyrad isopyrazam, fluxapyroxad; or carbendazim, thiophanatemethyl, iprodione, chlorothalonil, mancozeb, procymidone, vinclozolin, famoxadone, triadimenol, fenpropimorph, cymoxanil, prochloraz.

[0033] According to a preferred embodiment of the invention component C) is a fungicidal compound selected from the group of strobilurins.

[0034] According to a preferred embodiment of the invention component C) is a compound selected from the group of pyraclostrobin and kresoxim-methyl.

[0035] According to a particularly preferred embodiment of the invention component C) is pyraclostrobin.

[0036] According to a preferred embodiment of the invention component C) is a herbicidal compound such as acetochlor, aclonifen, cycloxydim, tepraloxydim, clethodim, fenoxyaprop, propaquizafop, quizalafop, haloxyfop, pendimethalin, clomazone, metazachlor, Metolachlor, quinmerac, dimethenamid, ethametsulfuron, napropamide, pethoxamide, glyphosate, gluphosinate, sulfentrazone, carfentrazone, trifluralin, flurochloridone, oxadiazone, linuron.

[0037] If the components A) and/or B) and/or C) are capable of forming geometrical isomers, for example E/Z isomers, both the pure isomers and mixtures thereof may be used in the compositions according to the invention. If the components A) and/or B) and/or C) have one or more centers of chirality and are thus present as enantiomers or diastereomers, both the pure enantiomers and diastereomers and mixtures thereof may be used in the compositions according to the invention.

[0038] If the components A) and/or B) and/or C) have ionizable functional groups, they can also be employed in the form of their agriculturally acceptable salts. Suitable are, in general, the salts of those cations and the acid addition salts of those acids whose cations and anions, respectively, have no adverse effect on the activity of the active compounds.

[0039] Preferred cations are the ions of the alkali metals, preferably of lithium, sodium and potassium, of the alkaline earth metals, preferably of calcium and magnesium, and of the transition metals, preferably of manganese, copper, zinc and iron, further ammonium and substituted ammonium in which one to four hydrogen atoms are replaced by C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl or benzyl, preferably ammonium or mono-C₁-C₄-alkyl, di-C₁-C₄-alkyl, tri-C₁-C₄-alkyl, tetra-C₁-C₄-alkylammonium that can be substituted by hydroxy or phenyl, for example, methylammonium, isopropylammonium, dimethylammonium, diisopropylammonium, trimethylammonium, tetramethylammonium, tetraethylammonium, tetrabutylammonium, 2-hydroxyethylammonium, 2-(2-hydroxyethyl-1-oxy)ethyl-1-ylammonium, di(2-hydroxyethyl-1-yl)ammonium, benzyltrimethylammonium, benzyltriethylammonium, furthermore phosphonium ions, sulfonium ions, preferably tri(C₁-C₄-alkyl)sulfonium, such as trimethylsulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium.

[0040] Anions of useful acid addition salts are primarily chloride, bromide, fluoride, iodide, hydrogensulfate, methylsulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate and also the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate.

[0041] Active components A) and/or B) and/or C) having a carboxyl group can be employed in the form of the acid, in the form of an agriculturally suitable salt as mentioned above or else in the form of an agriculturally acceptable derivative in the compositions according to the invention, for example as amides, such as mono- and di-C₁-C₆-alkylamides or

arylamides, as esters, for example as allyl esters, propargyl esters, C₁-C₁₀-alkyl esters, alkoxyalkyl esters and also as thioesters, for example as C₁-C₁₀-alkylthio esters. Preferred mono- and di-C₁-C₆-alkylamides are the methyl and the dimethylamides. Preferred arylamides are, for example, the anilides and the 2-chloroanilides. Preferred alkyl esters are, for example, the methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, methyl (1-methylhexyl) or isoctyl (2-ethylhexyl) esters. Preferred C₁-C₄-alkoxy-C₁-C₄-alkyl esters are the straight-chain or branched C₁-C₄-alkoxy ethyl esters, for example the methoxyethyl, ethoxyethyl or butoxyethyl ester. An example of a straight-chain or branched C₁-C₁₀-alkylthio ester is the ethylthio ester.

[0042] In the context of the present invention the following mixtures as well as the application of said mixtures in the methods according to the invention are particularly preferred.

[0043] Mixtures comprising exactly one PGR.

[0044] Mixtures comprising exactly two PGRs.

[0045] Further preferred embodiments relate to ternary mixtures which correspond to the binary mixtures mentioned above and additionally comprise a component C).

[0046] Here and below, the term "binary mixtures" includes mixtures comprising imazamox, including its agriculturally acceptable salts or amides, esters and thioesters and either 1, 2 or 3, PGR components B). Correspondingly, the term "ternary compositions" includes mixtures comprising imazamox, including its agriculturally acceptable salts or amides, esters and thioesters, one or more, for example 1, 2 or 3, components B) and one or more, for example 1, 2 or 3 components C).

[0047] In binary mixtures comprising imazamox, including its agriculturally acceptable salts or amides, esters and thioesters and 1, 2 or 3 components B), the weight ratio of the active compounds A:B is generally in the range from 1:300 to 300:1, in particular in the range from 1:100 to 100:1 and preferably in the range from 1:3 to 3:1, in particular in the range from 1:2 to 2:1 and even more preferably 1:1.

[0048] In ternary mixtures the relative proportions by weight of the components A:B are generally in the range from 1:300 to 300:1, preferably in the range from 1:100 to 100:1 and preferably in the range from 1:3 to 3:1, in particular in the range from 1:2 to 2:1 and even more preferably 1:1. The weight ratio of the components A:C is generally in the range from 1:3000 to 3000:1, preferably in the range from 1:1000 to 1000:1, in particular in the range from 1:100 to 100:1 and particularly preferably in the range from 1:20 to 20:1 and even more preferably in the range from 1:3 to 3:1. The weight ratio of the components B:C is generally in the range from 1:1000 to 1000:1, preferably in the range from 1:500 to 500:1, in particular in the range from 1:250 to 250:1 and particularly preferably in the range from 1:75 to 75:1 and even more preferably in the range from 1:20 to 20:1. The weight ratio of components A + B to component C is preferably in the range from 1:1000 to 1000:1, in particular in the range from 1:100 to 100:1 and particularly preferably in the range from 1:5 to 5:1 and even more preferably in the range from 1:3 to 3:1.

[0049] Particularly preferred embodiments of the present invention relate to methods applying combinations: C.3: A) imazamox and B) prohexadione-calcium; C.10: A) imazamox and B) prohexadione; C.17: A) imazamox and B) trinexapac; C.24: A) imazamox and B) trinexapac-ethyl.

[0050] Further particularly preferred embodiments of the present invention relate to methods comprising applying mixtures C.3, C.10, C.17 or C.24 each in combination with pyraclostrobin.

[0051] Further particularly preferred embodiments of the present invention relate to methods comprising applying mixtures C.3, C.10, C.17 or C.24 each in combination with kresoxim-methyl.

[0052] The mixtures according to the invention are suitable as herbicides. They are suitable as such or as an appropriately formulated composition. They act against broad-leaved weeds and grass weeds, particularly against Orobanche spp. and Striga spp., particularly Orobanche spp. in crops such as sunflower, oilseed rape, wheat, rice, corn, soybeans and cotton without causing any significant damage to the crop plants.

[0053] Another preferred embodiment of the invention relates to a method for controlling parasitic weeds in useful vegetation selected from the group of sunflower, oilseed rape, canola, soybeans, corn, peas, beans and alfalfa.

[0054] Another preferred embodiment of the invention relates to a method for improving the yield of the crop plant in useful vegetation selected from the group of sunflower, oilseed rape, canola, soybeans, corn, peas, beans and alfalfa.

[0055] Another preferred embodiment of the invention relates to a method for controlling parasitic weeds in sunflower and oilseed rape, particularly sunflower.

[0056] Another preferred embodiment of the invention relates to a method for improving the yield of the crop plant in sunflower and oilseed rape, particularly sunflower.

[0057] The components of the mixtures may be applied either simultaneously or sequentially or as premix. If administered sequentially, the components may be applied in any order and combination in a suitable time scale, for example, with up to 8 weeks between the time of applying the first component or combination and the time of applying the last component or combination. According to one embodiment the components are applied within 24 hours. More suitably, the components are applied within a few hours, preferably within one hour. In a preferred embodiment of the invention, the herbicide is applied first in case of sequential application. Preferably the components are applied simultaneously or in sequence. Particularly preferred application sequences D.1 to D.28 of components A), B) and, optionally, C) are listed

below in table IV.

table IV

	1st application	2nd application	3rd application
D.1	A	B	
D.2	A	B	C
D.3	A	C	B
D.4	A	B+C	
D.5	B	A	
D.6	B	A	C
D.7	B	C	A
D.8	B	A+C	
D.9	C	A	B
D.10	C	B	A
D.11	C	A+B	A+B
D.12	C	A+B	
D.13	A+C	B	
D.14	B+C	A	
D.15	B+C	A+B	
D.16	A+B		
D.17	A+B	C	
D.18	A+B	A+B	
D.19	A+B	A+B	A+B
D.20	A+B	A+B	C
D.21	A+B	A+B	B+C
D.22	A+B	C	C
D.23	A+B	B+C	
D.24	A+B+C		
D.25	A+B+C	A+B+C	
D.26	A+B+C	A+B+C	A+B+C
D.27	A+C	B	B
D.28	A+C	A+C	B

[0058] Further preferred embodiments of the present invention relate to a method for controlling parasitic weeds, comprising applying to the host plant, the weeds and/or their habitat, in application sequences defined in rows D.1, D5, D16 or D19, wherein A) is imazamox, B) is prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl or in application sequences defined in rows D2-D4, D6-D15, D17, D18, D20-D28, wherein A) is imazamox, B) is prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl and C) is pyraclostrobin or kresoxim-methyl.

[0059] The invention further relates to methods for improving the yield of the crop plant comprising applying herbicidal mixtures in application sequences as described above.

[0060] If the components are administered simultaneously, they may be administered as a tank mix or as a pre-formulated mixture of all components or as a pre-formulated mixture of some components tank mixed with the remaining components.

[0061] The simultaneous or sequential application of components according to the invention can be done repeatedly.

A preferred embodiment of the invention is a single application. Another preferred embodiment of the invention is a double application. Another preferred embodiment of the invention is a triple application. Another preferred embodiment of the invention is a quadruple application.

[0062] The mixtures and compositions according to the invention can also be used in genetically modified plants. The term "genetically modified plants" herein referred to are plants whose genetic material has been modified by the use of recombinant DNA techniques to include an inserted sequence of DNA that is not native to that plant species' genome or to exhibit a deletion of DNA that was native to that species' genome, wherein the modification(s) cannot readily be obtained by cross breeding, mutagenesis or natural recombination alone. Often, a particular genetically modified plant will be one that has obtained its genetic modification(s) by inheritance through a natural breeding or propagation process from an ancestral plant whose genome was the one directly treated by use of a recombinant DNA technique. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to improve certain properties of the plant. Such genetic modifications also include but are not limited to targeted post-translational modification of protein(s), oligo- or polypeptides. e. g., by inclusion therein of amino acid mutation(s) that permit, decrease, or promote glycosylation or polymer additions such as prenylation, acetylation farnesylation, or PEG moiety attachment.

[0063] Plants that have been modified by breeding, mutagenesis or genetic engineering, e.g. have been rendered tolerant to applications of specific classes of herbicides, such as auxinic herbicides such as dicamba or 2,4-D; bleacher herbicides such as 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors or phytoene desaturase (PDS) inhibitors; acetolactate synthase (ALS) inhibitors such as sulfonylureas or imidazolinones; enolpyruvyl shikimate 3-phosphate synthase (EPSP) inhibitors such as glyphosate; glutamine synthetase (GS) inhibitors such as glufosinate; protoporphyrinogen-IX oxidase inhibitors; lipid biosynthesis inhibitors such as acetylCoA carboxylase (ACCase) inhibitors; or oxynil (i. e. bromoxynil or ioxynil) herbicides as a result of conventional methods of breeding or genetic engineering; furthermore, plants have been made resistant to multiple classes of herbicides through multiple genetic modifications, such as resistance to both glyphosate and glufosinate or to both glyphosate and a herbicide from another class such as ALS inhibitors, HPPD inhibitors, auxinic herbicides, or ACCase inhibitors. These herbicide resistance technologies are, for example, described in Pest Management Science 61, 2005, 246; 61, 2005, 258; 61, 2005, 277; 61, 2005, 269; 61, 2005, 286; 64, 2008, 326; 64, 2008, 332; Weed Science 57, 2009, 108; Australian Journal of Agricultural Research 58, 2007, 708; Science 316, 2007, 1185; and references quoted therein. Several cultivated plants have been rendered tolerant to herbicides by mutagenesis and conventional methods of breeding, e. g., Clearfield® oil seed rape (summer rape or winter rape) or Clearfield® sunflower (BASF SE, Germany) being tolerant to imidazolinones, e. g., imazamox, or ExpressSun® sunflowers (DuPont, USA) or SURES-1 and SURES-2 (USDA-ARS, USA) being tolerant to sulfonyl ureas, e. g., tribenuron. Genetic engineering methods have been used to render cultivated plants such as soybean, cotton, corn, beets and rape, tolerant to herbicides such as glyphosate, imidazolinones and glufosinate, some of which are under development or commercially available under the brands or trade names RoundupReady® (glyphosate tolerant, Monsanto, USA), Cultivance® (imidazolinone tolerant, BASF SE, Germany) and LibertyLink® (glufosinate tolerant, Bayer Crop-Science, Germany).

[0064] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly from *Bacillus thuringiensis*, such as delta-endotoxins, e. g., Cry1A(b), Cry1A(c), Cry1F, Cry1F(a2), Cry1IA(b), Cry1IIA, Cry1IIB(b1) or Cry9c; vegetative insecticidal proteins (VIP), e. g., VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e. g., *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such as Streptomyces toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxy-steroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilbene synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins or toxins are to be understood expressly also as including pre-toxins, hybrid proteins, truncated or otherwise modified proteins. Hybrid proteins are characterized by a new combination of protein domains, (see, e. g., WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e. g., in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 und WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g., in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of arthropods, especially to beetles (Coleoptera), two-winged insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda). Genetically modified plants capable to synthesize one or more insecticidal proteins are, e. g., described in the publications mentioned above, and some of which are commercially available such as YieldGard® (corn cultivars producing the Cry1Ab toxin), YieldGard® Plus (corn cultivars producing Cry1Ab and Cry3Bb1 toxins), Starlink® (corn cultivars producing the Cry9c

toxin), Herculex® RW (corn cultivars producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothrinicin-N-Acetyltransferase [PAT]); NuCOTN® 33B (cotton cultivars producing the Cry1Ac toxin), Bollgard® I (cotton cultivars producing the Cry1Ac toxin), Bollgard® II (cotton cultivars producing Cry1Ac and Cry2Ab2 toxins); VIPCOT® (cotton cultivars producing a VIP-toxin); NewLeaf® (potato cultivars producing the Cry3A toxin); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, 5 Protecta®, Bt11 (e. g., Agrisure® CB) and Bt176 from Syngenta Seeds SAS, France, (corn cultivars producing the Cry1Ab toxin and PAT enzyme), MIR604 from Syngenta Seeds SAS, France (corn cultivars producing a modified version of the Cry3A toxin, c.f. WO 03/018810), MON 863 from Monsanto Europe S.A., Belgium (corn cultivars producing the Cry3Bb1 toxin), IPC 531 from Monsanto Europe S.A., Belgium (cotton cultivars producing a modified version of the Cry1Ac toxin) and 1507 from Pioneer Overseas Corporation, Belgium (corn cultivars producing the Cry1F toxin and PAT 10 enzyme).

[0065] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. Examples of such proteins are the so-called "pathogenesis-related proteins" (PR proteins, see, e.g., EP-A 392 225), plant disease resistance genes (e. g., potato culti-vars, which express resistance genes acting against Phytophthora 15 infestans derived from the Mexican wild potato, Solanum bulbocastanum) or T4-lyso-zym (e.g., potato cultivars capable of synthesizing these proteins with increased resistance against bacteria such as Erwinia amylovora). The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e.g., in the publications mentioned above.

[0066] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize 20 one or more proteins to increase the productivity (e.g., bio-mass production, grain yield, starch content, oil content or protein content), tolerance to drought, salinity or other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral pathogens of those plants.

[0067] Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount 25 of ingredients or new ingredients, specifically to improve human or animal nutrition, e. g., oil crops that produce health-promoting long-chain omega-3 fatty acids or unsaturated omega-9 fatty acids (e. g., Nexera® rape, Dow AgroSciences, Canada).

[0068] Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount 30 of ingredients or new ingredients, specifically to improve raw material production, e.g., potatoes that produce increased amounts of amylopectin (e.g. Amflora® potato, BASF SE, Germany).

[0069] The term seed comprises seed of all types, such as, for example, corns, seeds, fruits, tubers, seedlings and 35 similar forms. Here, preferably, the term seed describes corns and seeds. The seed used can be seed of the useful plants mentioned above, but also the seed of transgenic plants or plants obtained by customary breeding methods.

[0070] The components A), B) and C) according to the present invention, their N-oxides and salts can be converted 40 into customary types of agrochemical compositions, e. g. solutions, emulsions, suspensions, dusts, powders, pastes, granules, pressings, capsules, and mixtures thereof. Examples for composition types are suspensions (e.g. SC, OD, FS), emulsifiable concentrates (e.g. EC), emulsions (e.g. EW, EO, ES, ME), capsules (e.g. CS, ZC), pastes, pastilles, 45 wettable powders or dusts (e.g. WP, SP, WS, DP, DS), pressings (e.g. BR, TB, DT), granules (e.g. WG, SG, GR, FG, GG, MG), insecticidal articles (e.g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e.g. GF). These and further compositions types are defined in the "Catalogue of pesticide formulation types and international coding system", Technical Monograph No. 2, 6th Ed. May 2008, CropLife International. The compositions are prepared in a known manner, such as described by Mollet and Grubemann, Formulation technology, Wiley VCH, Weinheim, 2001; or Knowles, New developments in crop protection product formulation, Agrow Reports DS243, T&F Informa, London, 2005.

[0071] The invention relates in particular to herbicidally active mixtures comprising a component A) and component 45 B) and, optionally, a component C), as defined above, and also, in case of compositions, at least one liquid and/or solid carrier and/or one or more surfactants and, if desired, one or more further auxiliaries customary for crop protection compositions.

[0072] The invention also relates to a mixture formulated as a 1-component composition comprising a component A) and a component B) and, optionally, a component C), and at least one solid or liquid carrier and/or one or more surfactants 50 and, if desired, one or more further auxiliaries customary for crop protection compositions.

[0073] The invention also relates to a mixture formulated as a 2-component composition comprising a first component A), a solid or liquid carrier and/or one or more surfactants, and a second component B) and, optionally, a component C), a solid or liquid carrier and/or one or more surfactants, where additionally both components may also comprise 55 further auxiliaries customary for crop protection compositions.

[0074] Examples for suitable auxiliaries are solvents, liquid carriers, solid carriers or fillers, surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents, colorants, tackifiers and binders.

[0075] Suitable solvents and liquid carriers are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e.g. kerosene, diesel oil; oils of vegetable or animal origin; aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e.g. ethanol, propanol, butanol, benzylalcohol, cyclohexanol; glycols; DMSO; ketones, e.g. cyclohexanone; esters, e.g. lactates, carbonates, fatty acid esters, gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e.g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof.

[0076] Suitable solid carriers or fillers are mineral earths, e.g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharide powders, e.g. cellulose, starch; fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas; products of vegetable origin, e.g. cereal meal, tree bark meal, wood meal, nutshell meal, and mixtures thereof.

[0077] Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, block polymers, polyelectrolytes, and mixtures thereof. Such surfactants can be used as emulsifier, dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant. Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

[0078] Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylarylsulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignine sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxylated arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkylnaphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

[0079] Suitable nonionic surfactants are alkoxylates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxylates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxylated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. Examples of N-substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric surfactants are home- or copolymers of vinylpyrrolidone, vinylalcohols, or vinylacetate.

[0080] Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbetsains and imidazolines. Suitable block polymers are block polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol, polyethylene oxide and polypropylene oxide. Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinylamines or polyethyleneamines.

[0081] Suitable adjuvants are compounds, which have a neglectable or even no pesticidal activity themselves, and which improve the biological performance of the compound I on the target. Examples are surfactants, mineral or vegetable oils, and other auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

[0082] In a preferred embodiment citric acid is used as an adjuvant.

[0083] In another preferred embodiment ammonium sulfate is used as an adjuvant.

[0084] Suitable thickeners are polysaccharides (e.g. xanthan gum, carboxymethylcellulose), anorganic clays (organically modified or unmodified), polycarboxylates, and silicates.

[0085] Suitable bactericides are bronopol and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones.

[0086] Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

[0087] Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

[0088] Suitable colorants (e.g. in red, blue, or green) are pigments of low water solubility and water-soluble dyes. Examples are inorganic colorants (e.g. iron oxide, titan oxide, iron hexacyanoferrate) and organic colorants (e.g. alizarin-, azo- and phthalocyanine colorants).

[0089] Suitable tackifiers or binders are polyvinylpyrrolidone, polyvinylacetates, polyvinyl alcohols, polyacrylates, biological or synthetic waxes, and cellulose ethers.

[0090] The components A), B) and, optionally, C) and the compositions of the invention can for example be formulated as follows:

i) Water-soluble concentrates (SL, LS)

[0091] 10-60 wt% of active components A) and/or B) and, optionally, component C) according to the invention and 5-15 wt% wetting agent (e.g. alcohol alkoxylates) are dissolved in water and/or in a water-soluble solvent (e.g. alcohols) up to 100 wt%. The active substances dissolve upon dilution with water.

ii) Dispersible concentrates (DC)

[0092] 5-25 wt% of active components A) and/or B) and, optionally, component C) according to the invention and 1-10 wt% dispersant (e.g. polyvinylpyrrolidone) are dissolved in up to 100 wt% organic solvent (e.g. cyclohexanone). Dilution with water gives a dispersion.

iii) Emulsifiable concentrates (EC)

[0093] 15-70 wt% of active components A) and/or B) and, optionally, component C) according to the invention and 5-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in up to 100 wt% water-insoluble organic solvent (e.g. aromatic hydrocarbon). Dilution with water gives an emulsion.

iv) Emulsions (EW, EO, ES)

[0094] 5-40 wt% of active components A) and/or B) and, optionally, component C) according to the invention and 1-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in 20-40 wt% water-insoluble organic solvent (e.g. aromatic hydrocarbon). This mixture is introduced into up to 100 wt% water by means of an emulsifying machine and made into a homogeneous emulsion. Dilution with water gives an emulsion.

v) Suspensions (SC, OD, FS)

[0095] In an agitated ball mill, 20-60 wt% of active components A) and/or B) and, optionally, component C) according to the invention are comminuted with addition of 2-10 wt% dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate), 0,1-2 wt% thickener (e.g. xanthan gum) and up to 100 wt% water to give a fine active substance suspension. Dilution with water gives a stable suspension of the active substances. For FS type composition up to 40 wt% binder (e.g. polyvinylalcohol) is added.

vi) Water-dispersible granules and water-soluble granules (WG, SG)

[0096] 50-80 wt% of active components A) and/or B) and, optionally, component C) according to the invention are ground finely with addition of up to 100 wt% dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate) and prepared as water-dispersible or water-soluble granules by means of technical appliances (e.g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active substances.

vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)

[0097] 50-80 wt% of active components A) and/or B) and, optionally, component C) according to the invention are ground in a rotor-stator mill with addition of 1-5 wt% dispersants (e.g. sodium lignosulfonate), 1-3 wt% wetting agents (e.g. alcohol ethoxylate) and up to 100 wt% solid carrier, e.g. silica gel. Dilution with water gives a stable dispersion or solution of the active substances.

viii) Gel (GW, GF)

[0098] In an agitated ball mill, 5-25 wt% of active components A) and/or B) and, optionally, component C) according to the invention are comminuted with addition of 3-10 wt% dispersants (e.g. sodium lignosulfonate), 1-5 wt% thickener (e.g. carboxymethylcellulose) and up to 100 wt% water to give a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substances.

55 ix) Microemulsion (ME)

[0099] 5-20 wt% of active components A) and/or B) and, optionally, component C) according to the invention are added to 5-30 wt% organic solvent blend (e.g. fatty acid dimethylamide and cyclohexanone), 10-25 wt% surfactant blend

(e.g. alkohol ethoxylate and arylphenol ethoxylate), and water up to 100 %. This mixture is stirred for 1 h to produce spontaneously a thermodynamically stable microemulsion.

5 x) Microcapsules (CS)

[0100] An oil phase comprising 5-50 wt% of active components A) and/or B) and, optionally, component C) according to the invention, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), 2-15 wt% acrylic monomers (e.g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 wt% of a compound I according to the invention, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), and an isocyanate monomer (e.g. diphenylmethene-4,4'-diisocyanate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). The addition of a polyamine (e.g. hexamethylenediamine) results in the formation of a polyurea microcapsules. The monomers amount to 1-10 wt%. The wt% relate to the total CS composition.

15 xi) Dustable powders (DP, DS)

[0101] 1-10 wt% of active components A) and/or B) and, optionally, component C) according to the invention are ground finely and mixed intimately with up to 100 wt% solid carrier, e.g. finely divided kaolin.

20 xii) Granules (GR, FG)

[0102] 0.5-30 wt% of active components A) and/or B) and, optionally, component C) according to the invention is ground finely and associated with up to 100 wt% solid carrier (e.g. silicate). Granulation is achieved by extrusion, spray-drying or the fluidized bed.

25 xiii) Ultra-low volume liquids (UL)

[0103] 1-50 wt% of active components A) and/or B) and, optionally, component C) according to the invention are dissolved in up to 100 wt% organic solvent, e.g. aromatic hydrocarbon.

[0104] The compositions types i) to xiii) may optionally comprise further auxiliaries, such as 0,1-1 wt% bactericides, 5-15 wt% anti-freezing agents, 0,1-1 wt% anti-foaming agents, and 0,1-1 wt% colorants.

[0105] The agrochemical compositions generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, and most preferably between 0.5 and 75%, by weight of active substance. The active substances are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

[0106] Water-soluble concentrates (LS), Suspoemulsions (SE), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders (SS), emulsions (ES), emulsifiable concentrates (EC) and gels (GF) are usually employed for the purposes of treatment of plant propagation materials, particularly seeds. The compositions in question give, after two-to-tenfold dilution, active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40% by weight, in the ready-to-use preparations. Application can be carried out before or during sowing. Methods for applying or treating compound I and compositions thereof, respectively, on to plant propagation material, especially seeds include dressing, coating, pelleting, dusting, soaking and in-furrow application methods of the propagation material. Preferably, compound I or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

[0107] When employed in plant protection, the amounts of active substances applied are, depending on the kind of effect desired, from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, more preferably from 0.05 to 0.9 kg per ha, in particular from 0.1 to 0.75 kg per ha.

In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.1 to 1000 g, preferably from 1 to 1000 g, more preferably from 1 to 100 g and most preferably from 5 to 100 g, per 100 kilogram of plant propagation material (preferably seed) are generally required. When used in the protection of materials or stored products, the amount of active substance applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per hectare.

[0108] The required application rates of components A) according to the present invention are generally in the range from 0.0005 kg/ha to 2.5 kg/ha and preferably in the range from 0.005 kg/ha to 2 kg/ha or 0.01 kg/ha to 1.5 kg/ha of active substance.

[0109] The required application rates of components B) according to the present invention are generally in the range

from 0.0005 kg/ha to 2.5 kg/ha and preferably in the range from 0.005 kg/ha to 2 kg/ha or 0.01 kg/ha to 1.5 kg/ha of active substance.

[0110] The required application rates of components C) according to the present invention are generally in the range from 0.0005 kg/ha to 2.5 kg/ha and preferably in the range from 0.005 kg/ha to 2 kg/ha or 0.01 kg/ha to 1.5 kg/ha of active substance.

[0111] Various types of oils, wetters, adjuvants, fertilizer, or micronutrients, and other pesticides (e.g. herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the active substances or the compositions comprising them as premix or, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the compositions according to the invention in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1 and 1:1.

10 The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition according to the invention is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

15 [0112] According to one embodiment, individual components of the composition according to the invention such as parts of a kit or parts of a binary or ternary mixture may be mixed by the user himself in a spray tank and further auxiliaries may be added, if appropriate. In a further embodiment, either individual components of the composition according to the invention or partially premixed components, e. g. comprising components A) and/or components B) and/or components C), may be mixed by the user in a spray tank and further auxiliaries and additives may be added, if appropriate.

20 In a further embodiment, either individual components of the composition according to the invention or partially premixed components, e. g. comprising components A) and/or components B) and/or components C), can be applied jointly (e.g. after tank mix) or consecutively.

25 [0113] Application of the herbicidal compositions according to the present invention can be done before, during and/or after, preferably during and/or after, the emergence or penetration to the host roots by the undesirable plants. The herbicidal compositions according to the present invention can be applied pre- or post-emergence or together with the seed of a crop plant. It is also possible to apply the compounds and compositions by applying to seeds of the host plant, pretreated with a composition of the invention. If the active components A and B and, if appropriate C, are less well tolerated by certain crop plants, application techniques may be used in which the herbicidal compositions are sprayed, with the aid of the spraying equipment, in such a way that as far as possible they do not come into contact with the leaves of the sensitive crop plants, while the active compounds reach the leaves of undesirable plants growing underneath, or the bare soil surface (post-directed, lay-by).

30 [0114] Moreover, it may be advantageous to apply the mixtures of the present invention on their own or jointly in combination with other crop protection agents, for example with agents for controlling pests or phytopathogenic fungi or bacteria or with groups of active compounds which regulate growth. Also of interest is the miscibility with mineral salt solutions which are employed for treating nutritional and trace element deficiencies. Non-phytotoxic oils and oil concentrates can also be added.

Examples

40 [0115] The herbicidal action of the mixtures according to the invention was demonstrated by the following field experiments:

[0116] Seeds of sunflower (*Helianthus annuus* L.) were sown as host test plants using small scale field experiment equipment. Host test plants were, uniform and homogeneous with respect to age, cultivar, and row spacing and distance in the row. Plot size 2,8 m x 10 m - 4 rows per plot. Test area was kept weed free of conventional weeds (monocotyledon and dicotyledonous weeds) by hand hoeing.

[0117] Natural infestations of parasitic weeds were present in the soil of the test areas.

[0118] Postemergence treatments were performed with up to 4 application timings regarding the growth stage of the host plant. Application timings are specified with M, N, O, and P and related to the following growth stages of the host test plants according to the BBCH-scale" Lancashire, P. D., H. Bleiholder, P. Langelüddecke, R. Stauss, T. van den Boom, E. Weber and A. Witzenberger, 1991: A uniform decimal code for growth stages of crops and weeds. Ann. appl. Biol. 119, 561-601", where M = 2 - 4 true leaves (BBCH 12-14), N = 6 - 7 true leaves (BBCH 16-17), O = 8 to 10 leaves (BBCH 18 - 32) and P = 10 leaves up to inflorescence emergence (BBCH 32-51).

[0119] Sequential application were performed with the solo active components and compared to the respective mixture. The respectively stated components A) and B) were used as commercially available formulations and, with addition of a solvent system, introduced into the spray medium used for applying the active composition. In the examples, the spray medium used was water. Imazamox was used as commercial aqueous solution (SL) having an active ingredient concentration of 40 g/liter (Pulsar 40 ®). Prohexadione-calcium was used as commercial water dispersible granules (WG) having an active ingredient concentration of 100 g/kg (Regalis ®).

[0120] The test period extended over 1 to 4 month. During this time, the plants were tended, and their response to the individual treatments was evaluated.

5 [0121] Evaluation was carried out using a scale from 0 to 100 to express activity (in %). 100% means no emergence of the parasitic plants, or complete destruction of at least the above-ground parts, and 0% means no damage or normal course of growth of the parasitic plants. Good herbicidal activity is given at values of at least 70, and very good herbicidal activity is given at values of at least 85.

[0122] In the examples below, using the method of S. R. Colby (1967)" Calculating synergistic and antagonistic responses of herbicide combinations", Weeds 15, p. 22ff., the value E, which is expected if the activity of the individual active compounds is only additive, was calculated.

10

$$E = X + Y - (X \cdot Y / 100)$$

15 where

X = percent activity using active component A) at an application rate a;
 Y = percent activity using active component B) at an application rate b;
 E = expected activity (in %) by A + B) at application rates a + b.

20 [0123] If the value found experimentally is higher than the value E calculated according to Colby, a synergistic effect is present.

[0124] The herbicidal activity of the tested compositions against Orobanche cumana (ORACE) according to the intervention is compiled in table V.

25

30

35

40

45

50

55

table V

Weed	timing [BBCH growth stage]	g a.i./ha total		activity in %			expected activity in % (Colby)	synergistic activity in %
		Imazamox*	Prohexadione-Ca**	Imazamox	Prohexadione-Ca	Imazamox + Prohexadione-Ca		
ORACE	M,N,O,P	50	100	69	2	100	70	+30
ORACE	M,O	50	100	67	5	100	69	+31
ORACE	N,P	50	100	68	8	100	71	+29

* commercial product: Pulsar®

** commercial product: Regalis®

[0125] The delay of weed emergence and associated yield effects observed with the compositions according to the invention when tested against Orobanche sp. (ORASS) in sunflower is compiled in table VI. The number of Orobanche plants per host plant (imiazamox tolerant sunflower) and treatment were counted. The first visible emergence of Orobanche was assessed by counting the days from emergence of the sunflower till the first Orobanche appeared on the host. A later emergence is expressing a depression of the parasite as a result of the application of the active compounds. Yield of sunflower seeds was taken and compared to the control. A delay of emergence of Orobanche plants as well as a reduced number of Orobanche plants per host correlates with higher yield.

table VI

Trtm.	product	a.i. rate g /ha	timing [BBCH growth stage]	first visible emergence of orobanche (DAT***)	no. of orobanche plants per host plant)	yield of sunflower seeds kg/ha
1	untreated			22		1127
2	Imazamox*	30	14-15	81	15	2055
	Imazamox*	20	16-18			
3	Prohexadione- Ca**	60	14-15	63	55	961
	Prohexadione- Ca**	40	16-18			
4	Imazamox*	30	14-15	88	11	2712
	Prohexadione- Ca**	60	14-15			
	Imazamox*	20	16-18			
	Prohexadione- Ca**	40	16-18			
*** DAT = days after treatment						

[0126] The impact of imazamox in a tank mixture with prohexadione-Ca when compared to a single application of either component against Orobanche sp. (ORASS; application at BBCH 16-18) is demonstrated in table VII. The impact on infestation of the host plant (imiazamox tolerant sunflower) was measured 21 days after treatment as a value between 1 and 10 where 1 is equal to no infestation and 10 is equal to high negative impact of the Orobanche on the host plant that eventually leads to the host plant to die off.

table VII

Product	a.i. rate	a.i. unit	impact of host infestation with ORASS
Imazamox* solo treatment	15	g/ha	4
Prohexadione-Ca** solo treatment	75	g/ha	6
Imazamox* tank mixture	15	g/ha	2
Prohexadione-Ca**	75	g/ha	

Claims

1. A method for controlling parasitic weeds, comprising applying to the host plant, the weeds and/or their habitat, a herbicidal mixture of components
 - 1) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters, and
 - 2) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl,

wherein A) and B) are present in a synergistically effective amount or a composition comprising said mixture.

2. The method according to claim 1, wherein the synergistically effective amount is defined by a weight ratio of A) to B) from 1 : 300 to 300 : 1.
3. The method according to claim 1 or 2, wherein the parasitic weed that is controlled is from the genus Orobanche or from the genus Striga.
4. The method according to any one of claims 1 to 3, wherein the mixture or composition comprises as compound C) pyraclostrobin.

10 5. A mixture comprising components

- A) imazamox, including its agriculturally acceptable salts or amides, esters and thioesters, and
- B) one, two or three plant growth regulator(s) (PGR) selected from prohexadione, prohexadione-calcium, trinexapac or trinexapac-ethyl, wherein A) and B) are present in synergistically effective amount.

15 6. The mixture according to claim 5, wherein the synergistically effective amount is defined by a weight ratio of A) to B) from 1 : 300 to 300 : 1.

20 7. The mixture according to claim 5 or 6, wherein component B) is prohexadione or prohexadione-calcium.

8. The mixture according to any one of claims 5 to 7, which comprises as additional compound C) pyraclostrobin.

9. Use of a mixture as defined in any one of claims 5 to 8 for the control of parasitic weeds.

25 10. Compositions comprising mixtures as defined in any one of claims 5 to 8, at least one inert liquid and/or solid carriers and, if desired, at least one surfactant.

Patentansprüche

1. Verfahren zum Bekämpfen von parasitären Unkräutern, bei dem man eine herbizide Mischung der Komponenten A) Imazamox einschließlich seiner landwirtschaftlich unbedenklichen Salze oder Amide, Ester und Thioester, und B) einen, zwei oder drei Pflanzenwachstumsregulator(en) (PGR), ausgewählt aus Prohexadion, Prohexadion-Calcium, Trinexapac oder Trinexapac-Ethyl,

wobei A) und B) in einer synergistisch wirksamen Menge vorliegen, oder eine Zusammensetzung, die diese Mischung umfasst, auf die Wirtspflanze, die Unkräuter und/oder ihren Lebensraum aus bringt.

2. Verfahren nach Anspruch 1, wobei die synergistisch wirksame Menge durch ein Gewichtsverhältnis von A) zu B) von 1:300 bis 300:1 definiert ist.

3. Verfahren nach Anspruch 1 oder 2, wobei das parasitäre Unkraut, das bekämpft wird, aus der Gattung Orobanche oder aus der Gattung Striga stammt.

4. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Mischung oder Zusammensetzung Pyraclostrobin als Verbindung C) umfasst.

5. Mischung, umfassend die Komponenten

- A) Imazamox einschließlich seiner landwirtschaftlich unbedenklichen Salze oder Amide, Ester und Thioester, und
- B) einen, zwei oder drei Pflanzenwachstumsregulator(en) (PGR), ausgewählt aus Prohexadion, Prohexadion-Calcium, Trinexapac oder Trinexapac-Ethyl,

wobei A) und B) in einer synergistisch wirksamen Menge vorliegen.

6. Mischung nach Anspruch 5, wobei die synergistisch wirksame Menge durch ein Gewichtsverhältnis von A) zu B) von 1:300 bis 300:1 definiert ist.
- 5 7. Mischung nach Anspruch 5 oder 6, wobei es sich bei Komponente B) um Prohexadion oder Prohexadion-Calcium handelt.
8. Mischung nach einem der Ansprüche 5 bis 7, die als zusätzliche Verbindung C) Pyraclostrobin umfasst.
- 10 9. Verwendung einer Mischung wie in einem der Ansprüche 5 bis 8 definiert für die Bekämpfung von parasitären Unkräutern.
- 15 10. Zusammensetzungen umfassend Mischungen wie in einem der Ansprüche 5 bis 8 definiert, mindestens einen inerten flüssigen und/oder festen Träger und gegebenenfalls mindestens ein Tensid.

15 **Revendications**

1. Méthode de contrôle d'adventices parasites, comprenant l'application à la plante hôte, aux adventices et/ou à leur habitat, d'un mélange herbicide de composants
- 20 A) d'imazamox, y compris ses sels acceptables sur le plan agricole ou ses amides, esters et thioesters, et B) d'une, deux ou trois substances de croissance (PGR) choisies parmi la prohexadione, la prohexadione-calcium, le trinéxapac ou le trinéxapac-éthyle,
- 25 dans laquelle A) et B) sont présents selon une quantité efficace sur le plan synergique ou une composition comprenant ledit mélange.
2. Méthode selon la revendication 1, dans laquelle la quantité efficace sur le plan synergique est définie par un rapport pondéral de A) à B) allant de 1:300 à 300:1.
- 30 3. Méthode selon la revendication 1 ou 2, dans laquelle l'adventice parasite qui est contrôlée est issue du genre Orobanche ou issue du genre Striga.
- 35 4. Méthode selon l'une quelconque des revendications 1 à 3, dans laquelle le mélange ou la composition comprend, comme composé C), de la pyraclostrobine.
5. Mélange comprenant les composants
- 40 A) d'imazamox, y compris ses sels acceptables sur le plan agricole ou ses amides, esters et thioesters, et B) d'une, deux ou trois substances de croissance (PGR) choisies parmi la prohexadione, la prohexadione-calcium, le trinéxapac ou le trinéxapac-éthyle,
- 65 dans lequel A) et B) sont présents selon une quantité efficace sur le plan synergique.
- 45 6. Mélange selon la revendication 5, dans lequel la quantité efficace sur le plan synergique est définie par un rapport pondéral de A) à B) allant de 1:300 à 300:1.
7. Mélange selon la revendication 5 ou 6, dans lequel le composant B) est la prohexadione ou la prohexadione-calcium.
- 50 8. Méthode selon l'une quelconque des revendications 5 à 7, comprenant, comme composé C) supplémentaire, de la pyraclostrobine.
9. Utilisation d'un mélange tel que défini selon l'une quelconque des revendications 5 à 8, pour le contrôle d'adventices parasites.
- 55 10. Compositions comprenant des mélanges tels que définis selon l'une quelconque des revendications 5 à 8, au moins un véhicule liquide et/ou solide inerte, et, si on le souhaite, au moins un agent tensioactif.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2008124431 A1 [0006] [0009]
- EP 0123001 A1 [0007]
- US 20110028324 A [0008]
- US 2010304971 A1 [0008]
- US 2010210467 A1 [0008]
- US 20080269058 A1 [0008]
- EP 0224441 A1 [0008]
- DE 19834627 A1 [0008]
- WO 2003012115 A2 [0009]
- EP 1088480 A1 [0009]
- US 200544587 A1 [0009]
- WO 02015701 A [0064]
- EP 374753 A [0064]
- WO 93007278 A [0064]
- WO 9534656 A [0064]
- EP 427529 A [0064]
- EP 451878 A [0064]
- WO 0318810 A [0064]
- WO 0352073 A [0064]
- WO 03018810 A [0064]
- EP 392225 A [0065]

Non-patent literature cited in the description

- **HÖNIGES, A. ; WEGMANN, K.* ; ARDELEAN, A.** HELIA, 2008, vol. 31 (49), 1-12 [0002]
- **W. RADEMACHER.** *Ann. Rev. Plant Physiol. Plant Biol.*, 2000, vol. 51, 501-531 [0005]
- **Z. LIN et al.** Recent advances in ethylene research. *J. Exp. Bot.*, 2009, vol. 60, 3311-3336 [0006]
- **S. F. YANG et al.** Ethylene biosynthesis and its regulation in higher plants. *Ann. Rev. Plant Physiol.*, 1984, vol. 35, 155-89 [0006]
- **Z.W. FAN et al.** *Weed Research*, 2007, vol. 47, 34-43 [0007]
- **BEAM J.B. et al.** *Peanut Science*, 2002, vol. 29, 29-35 [0008]
- **MIZINIAK W. et al.** *Progress in Plant Protection*, 2008, vol. 48, 635-839 [0008]
- **KRAWCZYK R.** *Progress in Plant Protection*, 2006, vol. 46, 200-204 [0008]
- Farm Chemicals Handbook 2000. Meister Publishing Company, 2000, vol. 86 [0024]
- **B. HOCK ; C. FEDTKE ; R. R. SCHMIDT.** *Herbizide [Herbicides]*. Georg Thieme Verlag, 1995 [0024]
- **W. H. AHRENS.** *Herbicide Handbook*. Weed Science Society of America, 1994 [0024]
- **K. K. HATZIOS.** *Herbicide Handbook*. Weed Science Society of America, 1998 [0024]
- Catalogue of pesticide formulation types and international coding system. Technical Monograph. CropLife International, May 2008 [0070]
- **MOLLET ; GRUBEMANN.** *Formulation technology*. Wiley VCH, 2001 [0070]
- New developments in crop protection product formulation. **KNOWLES**. Agrow Reports DS243. T&F Informa, 2005 [0070]
- **MCCUTCHEON.** *Emulsifiers & Detergents*. McCutcheon's Directories, 2008, vol. 1 [0077]

Eljárás gyomok írtására imazamoxot és növényi növekedést szabályozó anyagokat tartalmazó keverékekkel

SZABADALMI IGÉNYPONTOK

1. Eljárás gyomok írtására, amely magában foglalja

A) imazamox, ideértve ennek mezőgazdaságilag elfogadható sóit vagy amidjait, észtereit és tioésztereit is, valamint
B) a prohexadion, a prohexadion-kalcium, a trinexapac vagy a trinexapac-etyl közül kiválasztott egy, két vagy három növényi növekedést szabályozó anyag (PGR) komponenseiből álló herbicid hatású keveréknak vagy a keveréket tartalmazó készítménynek a gazdanövényre, a gyomokra és/vagy ezek életterére való kijuttatását, ahol az A) és B) szinergisztikusan hatásos mennyiségben vannak jelen.

2. Az 1. igénypont szerinti eljárás, amelyben a szinergisztikusan hatásos mennyiség egy, az 1:300 és 300:1 közötti értékű A)/B) tömegaránytal van meghatározva.

3. Az 1. vagy 2. igénypont szerinti eljárás, amelyben az írtott gyom az Orobanche nemzetségből vagy a Stringa nemzetségből való.

4. Az 1-3. igénypontok bármelyike szerinti eljárás, amelyben a keverék vagy készítmény C) vegyületként pyraclostrobint tartalmaz.

5. Keverék, amely tartalmazza az alábbi komponenseket:

A) imazamox, ideértve ennek mezőgazdaságilag elfogadható sóit vagy amidjait, észtereit és tioésztereit is, valamint
B) a prohexadion, a prohexadion-kalcium, a trinexapac vagy a trinexapac-etyl közül kiválasztott egy, két vagy három növényi növekedést szabályozó anyag (PGR), ahol az A) és B) szinergisztikusan hatásos mennyiségben vannak jelen.

6. Az 5. igénypont szerinti keverék, ahol a szinergisztikusan hatásos mennyiség egy, az 1:300 és 300:1 közötti értékű A)/B) tömegaránytal van meghatározva.


7. Az 5. vagy 6. igénypont szerinti keverék, amelyben a B) komponens prohexadion vagy prohexadion-kalcium.

8. Az 5-7. igénypontok bármelyike szerinti keverék, amely további C) vegyületként pyraclostrobint tartalmaz.

9. Az 5-8. igénypontok bármelyike szerint meghatározott keverék alkalmazása gyomok írtására.

10. Készítmények, amelyek az 5-8. igénypontok bármelyike szerint meghatározott keverékeket, legalább egy inerit folyékony és/vagy szilárd hordozóanyagot, és amennyiben megkívánt, legalább egy felületaktiv anyagot tartalmaznak.

A meghatalmazott:


Dr. Szabolcs Ábrahám
szabolcs@medika.hu
Szent László Kórház
H-1182 Budapest, Kerepesi út 112.
Telefon: +36 1-3600 1000 Fax: +36 1-3600
E-mail: medika@medika.hu