



(86) Date de dépôt PCT/PCT Filing Date: 2003/08/15  
 (87) Date publication PCT/PCT Publication Date: 2004/02/26  
 (45) Date de délivrance/Issue Date: 2014/07/22  
 (85) Entrée phase nationale/National Entry: 2005/02/02  
 (86) N° demande PCT/PCT Application No.: IB 2003/004502  
 (87) N° publication PCT/PCT Publication No.: 2004/016286  
 (30) Priorité/Priority: 2002/08/16 (US10/222,140)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),  
*A61K 47/26* (2006.01), *A61K 9/19* (2006.01)  
 (72) Inventeurs/Inventors:  
KRAUSE, HANS-JUERGEN, DE;  
BAUST, LISA, DE;  
DICKES, MICHAEL, DE  
 (73) Propriétaire/Owner:  
ABBVIE BIOTECHNOLOGY LTD., BM  
 (74) Agent: TORYS LLP

(54) Titre : ANTICORPS ANTI-TNF ALPHA EN SOLUTION ET LEURS UTILISATIONS  
 (54) Title: ANTI-TNF-ALPHA ANTIBODIES IN SOLUTION AND USES THEREOF

(57) Abrégé/Abstract:

A liquid aqueous pharmaceutical formulation is described which has a high protein concentration, a pH of between about 4 and about 8, and enhanced stability.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
26 February 2004 (26.02.2004)

PCT

(10) International Publication Number  
**WO 2004/016286 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 39/395**,  
9/19, 47/26

(21) International Application Number:  
PCT/IB2003/004502

(22) International Filing Date: 15 August 2003 (15.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
10/222,140 16 August 2002 (16.08.2002) US

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES (BERMUDA) LTD.** [—/—]; Clarendon House, 2 Church Street, Hamilton HM 11 (BM).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KRAUSE, Hans-Juergen** [DE/DE]; Dieselstrasse 33, 67551 Worms (DE). **BAUST, Lisa** [DE/DE]; Stamitzstrasse 8, 68167 Mannheim (DE). **DICKES, Michael** [DE/DE]; Schafergasse 58, 67127 Rodersheim-Gronau (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
22 July 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL ANTI-TNF-ALPHA ANTIBODY FORMULATION

(57) Abstract: A liquid aqueous pharmaceutical formulation is described which has a high protein concentration, a pH of between about 4 and about 8, and enhanced stability.

WO 2004/016286 A3

## ANTI-TNF-ALPHA ANTIBODIES IN SOLUTION AND USES THEREOF

5 Background of the Invention

Tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) is a cytokine produced by numerous cell types, including monocytes and macrophages, that was originally identified based on its capacity to induce the necrosis of certain mouse tumors (see e.g., Old, L. (1985) *Science* 10 230:630-632). Subsequently, a factor termed cachectin, associated with cachexia, was shown to be the same molecule as TNF $\alpha$ . TNF $\alpha$  has been implicated in mediating shock (see e.g., Beutler, B. and Cerami, A. (1988) *Annu. Rev. Biochem.* 57:505-518; Beutler, B. and Cerami, A. (1989) *Annu. Rev. Immunol.* 7:625-655). Furthermore, TNF $\alpha$  has been implicated in the pathophysiology of a variety of other human diseases 15 and disorders, including sepsis, infections, autoimmune diseases, transplant rejection and graft-versus-host disease (see e.g., Moeller, A., et al. (1990) *Cytokine* 2:162-169; U.S. Patent No. 5,231,024 to Moeller et al.; European Patent Publication No. 260 610 B1 by Moeller, A., et al. Vasilli, P. (1992) *Annu. Rev. Immunol.* 10:411-452; Tracey, K.J. and Cerami, A. (1994) *Annu. Rev. Med.* 45:491-503).

20 Because of the harmful role of human TNF $\alpha$  (hTNF $\alpha$ ) in a variety of human disorders, therapeutic strategies have been designed to inhibit or counteract hTNF $\alpha$  activity. In particular, antibodies that bind to, and neutralize, hTNF $\alpha$  have been sought as a means to inhibit hTNF $\alpha$  activity. Some of the earliest of such antibodies were mouse monoclonal antibodies (mAbs), secreted by hybridomas prepared from 25 lymphocytes of mice immunized with hTNF $\alpha$  (see e.g., Hahn T; et al., (1985) *Proc Natl Acad Sci USA* 82: 3814-3818; Liang, C-M., et al. (1986) *Biochem. Biophys. Res. Commun.* 137:847-854; Hirai, M., et al. (1987) *J. Immunol. Methods* 96:57-62; Fendly, B.M., et al. (1987) *Hybridoma* 6:359-370; Moeller, A., et al. (1990) *Cytokine* 2:162-169; U.S. Patent No. 5,231,024 to Moeller et al.; European Patent Publication No. 186 30 833 B1 by Wallach, D.; European Patent Application Publication No. 218 868 A1 by Old et al.; European Patent Publication No. 260 610 B1 by Moeller, A., et al.). While these mouse anti-hTNF $\alpha$  antibodies often displayed high affinity for hTNF $\alpha$  (e.g.,  $K_d \leq 10^{-9}M$ ) and were able to neutralize hTNF $\alpha$  activity, their use *in vivo* may be limited by problems associated with administration of mouse antibodies to humans, such as short 35 serum half life, an inability to trigger certain human effector functions and elicitation of an unwanted immune response against the mouse antibody in a human (the "human anti-mouse antibody" (HAMA) reaction).



- 2 -

In an attempt to overcome the problems associated with use of fully-murine antibodies in humans, murine anti-hTNF $\alpha$  antibodies have been genetically engineered to be more "human-like." For example, chimeric antibodies, in which the variable regions of the antibody chains are murine-derived and the constant regions of the antibody chains are human-derived, have been prepared (Knight, D.M, *et al.* (1993) *Mol. Immunol.* 30:1443-1453; PCT Publication No. WO 92/16553 by Daddona, P.E., *et al.*). Additionally, humanized antibodies, in which the hypervariable domains of the antibody variable regions are murine-derived but the remainder of the variable regions and the antibody constant regions are human-derived, have also been prepared (PCT Publication No. WO 92/11383 by Adair, J.R., *et al.*). However, because these chimeric and humanized antibodies still retain some murine sequences, they still may elicit an unwanted immune reaction, the human anti-chimeric antibody (HACA) reaction, especially when administered for prolonged periods, *e.g.*, for chronic indications, such as rheumatoid arthritis (see *e.g.*, Elliott, M.J., *et al.* (1994) *Lancet* 344:1125-1127; Elliot, M.J., *et al.* (1994) *Lancet* 344:1105-1110).

A preferred hTNF $\alpha$  inhibitory agent to murine mAbs or derivatives thereof (*e.g.*, chimeric or humanized antibodies) would be an entirely human anti-hTNF $\alpha$  antibody, since such an agent should not elicit the HAMA reaction, even if used for prolonged periods. Human monoclonal autoantibodies against hTNF $\alpha$  have been prepared using human hybridoma techniques (Boyle, P., *et al.* (1993) *Cell. Immunol.* 152:556-568; Boyle, P., *et al.* (1993) *Cell. Immunol.* 152:569-581; European Patent Application Publication No. 614 984 A2 by Boyle, *et al.*). However, these hybridoma-derived monoclonal autoantibodies were reported to have an affinity for hTNF $\alpha$  that was too low to calculate by conventional methods, were unable to bind soluble hTNF $\alpha$  and were unable to neutralize hTNF $\alpha$ -induced cytotoxicity (see Boyle, *et al.*; *supra*). Moreover, the success of the human hybridoma technique depends upon the natural presence in human peripheral blood of lymphocytes producing autoantibodies specific for hTNF $\alpha$ . Certain studies have detected serum autoantibodies against hTNF $\alpha$  in human subjects (Fomsgaard, A., *et al.* (1989) *Scand. J. Immunol.* 30:219-223; Bendtzen, K., *et al.* (1990) *Prog. Leukocyte Biol.* 10B:447-452), whereas others have not (Leusch, H-G., *et al.* (1991) *J. Immunol. Methods* 139:145-147).

Alternative to naturally-occurring human anti-hTNF $\alpha$  antibodies would be a recombinant hTNF $\alpha$  antibody. Recombinant human antibodies that bind hTNF $\alpha$  with relatively low affinity (*i.e.*,  $K_D \sim 10^{-7}M$ ) and a fast off rate (*i.e.*,  $K_{off} \sim 10^{-2} \text{ sec}^{-1}$ ) have been described (Griffiths, A.D., *et al.* (1993) *EMBO J.* 12:725-734). However, because of their relatively fast dissociation kinetics, these antibodies may not be suitable for therapeutic use. Additionally, a recombinant human anti-hTNF $\alpha$  has been described

WO 2004/016286

PCT/IB2003/004502

- 3 -

that does not neutralize hTNF $\alpha$  activity, but rather enhances binding of hTNF $\alpha$  to the surface of cells and enhances internalization of hTNF $\alpha$  (Lidbury, A., *et al.* (1994) *Biotechnol. Ther.* 5:27-45; PCT Publication No. WO 92/03145 by Aston, R. *et al.*)

Recombinant human antibodies that bind soluble hTNF $\alpha$  with high affinity and slow dissociation kinetics and that have the capacity to neutralize hTNF $\alpha$  activity, including hTNF $\alpha$ -induced cytotoxicity (*in vitro* and *in vivo*) and hTNF $\alpha$ -induced cell activation, have also been described (see U.S. Patent No. 6,090,382).

### Summary of the Invention

There is a need for a stable aqueous pharmaceutical formulation with an extended shelf life, comprising an antibody which is suitable for therapeutic use to inhibit or counteract detrimental hTNF $\alpha$  activity. There is also a need for a stable aqueous pharmaceutical formulation with an extended shelf life, comprising an antibody suitable for therapeutic use which is easily administered and contains a high protein concentration.

This invention relates to:

- <1> A liquid aqueous pharmaceutical formulation comprising a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody at a concentration from 20 to 150 mg/ml, a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, wherein said formulation has a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
- <2> A liquid aqueous pharmaceutical formulation comprising
- (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody,
  - (b) from 5 to 20 mg/ml of mannitol,
  - (c) from 0.1 to 10 mg/ml of polysorbate 80, and
  - (d) a buffer system comprising citrate and phosphate, with a pH from 4 to 8,



WO 2004/016286

PCT/IB2003/004502

- 4 -

wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

- <3> The liquid aqueous pharmaceutical formulation of <1>, wherein the sugar alcohol is mannitol.
- 5 <4> The liquid aqueous pharmaceutical formulation of <1>, wherein the polysorbate is polysorbate 80.
- <5> The liquid aqueous pharmaceutical formulation of <2>, wherein the antibody is antibody D2E7.
- <6> The liquid aqueous pharmaceutical formulation of <1>, <3> or <4>, wherein  
10 the antibody is antibody D2E7.
- <7> The liquid aqueous pharmaceutical formulation of any one of <1>-<6>, comprising from 30 to 120 mg/ml of the antibody.
- <8> The liquid aqueous pharmaceutical formulation of any one of <1>-<7>, comprising from 35 to 115 mg/ml of the antibody.
- 15 <9> The liquid aqueous pharmaceutical formulation of <8>, comprising from 40 to 110 mg/ml of the antibody.
- <10> The liquid aqueous pharmaceutical formulation of <9>, comprising from 45 to 105 mg/ml of the antibody.
- <11> The liquid aqueous pharmaceutical formulation of any one of <1>-<10>,  
20 wherein the pH is from 4.5 to 6.0.
- <12> The liquid aqueous pharmaceutical formulation of <11>, wherein the pH is from 4.8 to 5.5.
- <13> The liquid aqueous pharmaceutical formulation of <2> or <3>, comprising from 10 to 14 mg/ml of mannitol.
- 25 <14> The liquid aqueous pharmaceutical formulation of <2> or <4>, comprising from 0.5 to 5 mg/ml of polysorbate 80.
- <15> The liquid aqueous pharmaceutical formulation of <2>, which contains
- (a) from 40 to 100 mg/ml of the antibody,
  - (b) from 7.5 to 15 mg/ml of mannitol, and
  - 30 (c) from 0.5 to 5 mg/ml of polysorbate 80.

WO 2004/016286

PCT/IB2003/004502

- 5 -

- <16> The liquid aqueous pharmaceutical formulation of any one of <1>-<4> and <13>-<15>, wherein the antibody comprises an IgG<sub>1</sub> heavy chain constant region.
- <17> The liquid aqueous pharmaceutical formulation of any one of <1>-<4> and <13>-<15>, wherein the antibody comprises an IgG<sub>4</sub> heavy chain constant region.
- 5 <18> A liquid aqueous pharmaceutical formulation having a pH from 4 to 8 and comprising
- (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG<sub>1</sub> antibody comprising a D2E7 light chain variable region and a D2E7 heavy chain variable region;
  - 10 (b) from 10 to 14 mg/ml of mannitol,
  - (c) from 0.1 to 5 mg/ml of polysorbate 80,
  - (d) from 1 to 1.5 mg/ml of citric acid monohydrate,
  - (e) from 0.25 to 0.5 mg/ml of sodium citrate,
  - (f) from 1.25 to 1.75 mg/ml of disodium phosphate dihydrate,
  - 15 (g) from 0.7 to 1.1 mg/ml of sodium dihydrogen phosphate dihydrate, and
  - (h) from 6.0 to 6.4 mg/ml sodium chloride.
- <19> A liquid aqueous pharmaceutical formulation having a pH from 4 to 8 and comprising
- (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha
  - 20 (TNF $\alpha$ ) IgG<sub>4</sub> antibody comprising a D2E7 light chain variable region and a D2E7 heavy chain variable region;
  - (b) from 10 to 14 mg/ml of mannitol,
  - (c) from 0.1 to 5 mg/ml of polysorbate 80,
  - (d) from 1 to 1.5 mg/ml of citric acid monohydrate,
  - 25 (e) from 0.25 to 0.5 mg/ml of sodium citrate,
  - (f) from 1.25 to 1.75 mg/ml of disodium phosphate dihydrate,
  - (g) from 0.7 to 1.1 mg/ml of sodium dihydrogen phosphate dihydrate, and
  - (h) from 6.0 to 6.4 mg/ml sodium chloride.
- <20> Use of the formulation of any one of <1>-<19> in the preparation of a
- 30 medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an autoimmune disease.



WO 2004/016286

PCT/IB2003/004502

- 6 -

- <21> The use of <20>, wherein the autoimmune disease is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis, autoimmune hepatitis, or nephrotic syndrome.
- 5 <22> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an infectious disease.
- <23> The use of <22>, wherein the infectious disease is meningitis, malaria, acquired immune deficiency syndrome, or cytomegalovirus infection secondary to  
10 transplantation.
- <24> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is a pulmonary disorder.
- <25> The use of <24>, wherein the pulmonary disorder is adult respiratory distress  
15 syndrome, shock lung, a chronic pulmonary inflammatory disease, an obstructive pulmonary disease, a chronic obstructive pulmonary disease, pulmonary sarcoidosis, pulmonary fibrosis, or silicosis.
- <26> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human  
20 subject, wherein the disorder is an inflammatory bowel disease.
- <27> The use of <26>, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.
- <28> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human  
25 subject, wherein the disorder is ankylosing spondylitis.
- <29> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating juvenile rheumatoid arthritis in a human subject.
- <30> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating psoriasis in a human subject.
- 30 <31> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating psoriatic arthritis in a human subject.



WO 2004/016286

PCT/IB2003/004502

- 7 -

- <32> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating sarcoidosis in a human subject.
- <33> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating scleroderma in a human subject.
- 5 <34> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a spondyloarthropathy in a human subject.
- <35> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating Still's disease in a human subject.
- <36> Use of the formulation of any one of <1>-<19> in the preparation of a  
10 medicament for treating Sjogren's syndrome in a human subject.
- <37> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating atherosclerosis in a human subject.
- <38> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating uveitis in a human subject.
- 15 <39> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is sepsis, a graft versus host disease, a malignancy, cachexia, an inflammatory bone disorder, a bone resorption disease, hepatitis, a coagulation disturbance, burn, a reperfusion injury, keloid formation, scar tissue  
20 formation, pyrexia, a periodontal disease, obesity, radiation, Alzheimer's disease, asthma, chronic fatigue syndrome, liver failure, chronic liver failure, dermatopolymyositis, diabetic macrovasculopathy, endometriosis, familial periodic fevers, fibrosis, hemodialysis, Jarisch-Herxheimer reaction, Kawasaki syndrome, myelodysplastic syndrome, panciaticular vulgaris, peripheral neuropathy,  
25 polyarticular, polymyositis, progressive renal failure, Reiter's syndrome, stroke, therapy associated syndrome, therapy induced inflammatory syndrome, inflammatory syndrome following IL-2 administration, thoracoabdominal aortic aneurysm repair (TAAA), Vasulo-Behcet's disease, Yellow Fever vaccination, type 1 diabetes mellitus, type 2 diabetes mellitus, neuropathic pain, sciatica, cerebral edema, edema  
30 in or around the spinal cord, vasculitide, Wegener's granulomatosis, temporal arteritis, polymyalgia rheumatica, Takayasu's arteritis, polyarteritis nodosa, microscopic polyangiitis, Churg-Strauss syndrome, Felty's syndrome, mixed

WO 2004/016286

PCT/IB2003/004502

- 7a -

- connective tissue disorder, relapsing polychondritis, pseudogout, loosening of prostheses, sclerosing cholangitis, acute pancreatitis, chronic pancreatitis, glomerulonephritides, post-streptococcal glomerulonephritis or IgA nephropathy, rheumatic heart disease, orchitis, pyoderma gangrenosum, multiple myeloma, TNF
- 5 receptor associated periodic syndrome [TRAPS], atherosclerosis, steroid dependent giant cell arteritismyostitis, or a cardiac disorder.
- <40> Use of the formulation of any one of <1>-<19> for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an autoimmune disease.
- 10 <41> The use of <40>, wherein the autoimmune disease is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis, autoimmune hepatitis, or nephrotic syndrome.
- <42> Use of the formulation of any one of <1>-<19> for treating a disorder in
- 15 which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an infectious disease.
- <43> The use of <42>, wherein the infectious disease is meningitis, malaria, acquired immune deficiency syndrome, or cytomegalovirus infection secondary to transplantation.
- 20 <44> Use of the formulation of any one of <1>-<19> for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is a pulmonary disorder.
- <45> The use of <44>, wherein the pulmonary disorder is respiratory distress syndrome, shock lung, a chronic pulmonary inflammatory disease, an obstructive
- 25 pulmonary disease, a chronic obstructive pulmonary disease, pulmonary sarcoidosis, pulmonary fibrosis, or silicosis.
- <46> Use of the formulation of any one of <1>-<19> for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an inflammatory bowel disease.
- 30 <47> The use of <46>, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.



WO 2004/016286

PCT/IB2003/004502

- 7b -

- <48> Use of the formulation of any one of <1>-<19> for treating ankylosing spondylitis in a human subject.
- <49> Use of the formulation of any one of <1>-<19> for treating juvenile rheumatoid arthritis (JRA) in a human subject.
- 5 <50> Use of the formulation of any one of <1>-<19> for treating psoriasis in a human subject.
- <51> Use of the formulation of any one of <1>-<19> for treating psoriatic arthritis in a human subject.
- <52> Use of the formulation of any one of <1>-<19> for treating sarcoidosis in a  
10 human subject.
- <53> Use of the formulation of any one of <1>-<19> for treating scleroderma in a human subject.
- <54> Use of the formulation of any one of <1>-<19> for treating a spondyloarthropathy in a human subject.
- 15 <55> Use of the formulation of any one of <1>-<19> for treating Still's disease in a human subject.
- <56> Use of the formulation of any one of <1>-<19> for treating Sjogren's syndrome in a human subject.
- <57> Use of the formulation of any one of <1>-<19> for treating atherosclerosis in  
20 a human subject.
- <58> Use of the formulation of any one of <1>-<19> for treating uveitis in a human subject.
- <59> Use of the formulation of any one of <1>-<19> for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is sepsis,  
25 a graft versus host disease, a malignancy, cachexia, an inflammatory bone disorder, a bone resorption disease, hepatitis, a coagulation disturbance, burn, a reperfusion injury, keloid formation, scar tissue formation, pyrexia, a periodontal disease, obesity, radiation, Alzheimer's disease, asthma, chronic fatigue syndrome, liver failure, chronic liver failure, dermatopolymyositis, diabetic macrovasculopathy,  
30 endometriosis, familial periodic fevers, fibrosis, hemodialysis, Jarisch-Herxheimer

- 7c -

reaction, Kawasaki syndrome, myelodysplastic syndrome, panciaticular vulgaris,  
peripheral neuropathy, polyarticular, polymyositis, progressive renal failure, Reiter's  
syndrome, stroke, therapy associated syndrome, therapy induced inflammatory  
syndrome, inflammatory syndrome following IL-2 administration, thoracoabdominal  
5 aortic aneurysm repair (TAAA), Vasulo-Behect's disease, Yellow Fever vaccination,  
type 1 diabetes mellitus, type 2 diabetes mellitus, neuropathic pain, sciatica, cerebral  
edema, edema in or around the spinal cord, vasculitide, Wegener's granulomatosis,  
temporal arteritis, polymyalgia rheumatica, Takayasu's arteritis, polyarteritis nodosa,  
microscopic polyangiitis, Churg-Strauss syndrome, Felty's syndrome, mixed  
10 connective tissue disorder, relapsing polychondritis, pseudogout, loosening of  
prostheses, sclerosing cholangitis, acute pancreatitis, chronic pancreatitis,  
glomerulonephritides, post-streptococcal glomerulonephritis or IgA nephropathy,  
rheumatic heart disease, orchitis, pyoderma gangrenosum, multiple myeloma, TNF  
receptor associated periodic syndrome [TRAPS], atherosclerosis, steroid dependent  
15 giant cell arteritismyositis, or a cardiac disorder.

<60> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that  
specifically binds TNF $\alpha$  to treat psoriasis, wherein the antibody is at a concentration  
from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from  
5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer  
20 system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the  
antibody comprises a light chain variable region and a heavy chain variable region of  
antibody D2E7.

<61> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that  
specifically binds TNF $\alpha$  to treat psoriatic arthritis, wherein the antibody is at a  
25 concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a  
concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10  
mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and  
wherein the antibody comprises a light chain variable region and a heavy chain  
variable region of antibody D2E7.

<62> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that  
specifically binds TNF $\alpha$  to treat ankylosing spondylitis, wherein the antibody is at a  
30 concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a



- 7d -

concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

5 <63> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat juvenile rheumatoid arthritis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and  
10 wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

<64> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat rheumatoid arthritis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a  
15 concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

<65> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that  
20 specifically binds TNF $\alpha$  to treat Crohn's disease, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain  
25 variable region of antibody D2E7.

<66> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat ulcerative colitis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10  
30 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

WO 2004/016286

PCT/IB2003/004502

- 7e -

- 5 <67> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat uveitis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
- 10 <68> Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat Vasulo-Behcet's disease, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
- 15 <69> The use of any one of <60>-<68>, wherein the sugar alcohol is mannitol.
- <70> The use of <69>, wherein the mannitol is at a concentration from 10 to 14 mg/ml.
- <71> The use of any one of <60>-<70>, wherein the polysorbate is polysorbate 80.
- 20 <72> The use of <71>, wherein the polysorbate 80 is at a concentration from 0.5 to 5 mg/ml.
- <73> The use of any one of <60>-<72>, wherein the antibody is at a concentration from 30 to 120 mg/ml.
- <74> The use of <73>, wherein the antibody is at a concentration from 35 to 115 mg/ml.
- 25 <75> The use of <73>, wherein the antibody is at a concentration from 40 to 110 mg/ml.
- <76> The use of <73>, wherein the antibody is at a concentration from 45 to 105 mg/ml.
- <77> The use of any one of <60>-<76>, wherein the pH is from 4.5 to 6.0.
- 30 <78> The use of <77>, wherein the pH is from 4.8 to 5.5.



WO 2004/016286

PCT/IB2003/004502

- 7f -

- <79> The use of any one of <60>-<78>, wherein the antibody is antibody D2E7, the sugar alcohol is mannitol at a concentration from 10 to 14 mg/ml, and the polysorbate is polysorbate 80 at a concentration from 0.5 to 5 mg/ml.
- <80> The use of any one of <60>-<78>, wherein the antibody comprises an IgG<sub>1</sub> or  
5 IgG<sub>4</sub> heavy chain constant region.
- <81> The use of any one of <60>-<78>, wherein the antibody is antibody D2E7.
- <82> The use of any one of <79>-<81>, wherein the use is subcutaneous use.
- <83> A liquid aqueous pharmaceutical formulation comprising a human  
10 anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG<sub>1</sub> antibody at a concentration from 20 to 150 mg/ml, a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate,  
wherein said formulation has a pH from 4 to 8, and  
wherein the antibody comprises a light chain variable region and a heavy  
15 chain variable region of antibody D2E7.
- <84> A liquid aqueous pharmaceutical formulation comprising a human  
anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody at a concentration from 40 to 110 mg/ml, citric acid monohydrate, dibasic sodium phosphate dihydrate, mannitol, monobasic sodium phosphate dihydrate, polysorbate 80, sodium citrate, and sodium  
20 chloride,  
wherein said formulation has a pH from 4 to 8, and  
wherein the antibody comprises a light chain variable region and a heavy  
chain variable region of antibody D2E7.
- <85> The liquid aqueous pharmaceutical formulation of <83> or <84>, wherein the  
25 pH is from 4.5 to 6.0.
- <86> The liquid aqueous pharmaceutical formulation of <83> or <84>, wherein the pH is from 4.8 to 5.5.
- <87> The liquid aqueous pharmaceutical formulation of any one of <83> to <86>, wherein the concentration of the antibody is 50 mg/ml.
- 30 <88> The liquid aqueous pharmaceutical formulation of <83>, wherein the antibody is antibody D2E7, the sugar alcohol is mannitol at a concentration from 10 to

WO 2004/016286

PCT/IB2003/004502

- 7g -

14 mg/ml, and the polysorbate is polysorbate 80 at a concentration from 0.5 to 5 mg/ml.

5 <89> The liquid aqueous pharmaceutical formulation of <84>, wherein the antibody is antibody D2E7, mannitol is at a concentration from 10 to 14 mg/ml, and polysorbate 80 is at a concentration from 0.5 to 5 mg/ml.

10 <90> The liquid aqueous pharmaceutical formulation of <83>, wherein the antibody is antibody D2E7 at a concentration of about 50 mg/ml, the sugar alcohol is mannitol at a concentration of about 12 mg/ml, the polysorbate is polysorbate 80 at a concentration of about 1 mg/ml, and the pH of the buffer system is from about 4.8 to about 5.5.

<91> The liquid aqueous pharmaceutical formulation of <84>, wherein the antibody is antibody D2E7 at a concentration of about 50 mg/ml, mannitol is at a concentration of about 12 mg/ml, polysorbate 80 is at a concentration of about 1 mg/ml, and the pH of the buffer system is from about 5.0 to about 5.2.

15 <92> Use of the formulation of any one of <1>-<19> in the preparation of a medicament for treating rheumatoid arthritis in a human subject.

<93> Use of the formulation of any one of <1>-<19> for treating rheumatoid arthritis in a human subject.



### Detailed Description of the Invention

This invention pertains to a liquid aqueous pharmaceutical formulation with a pH of about 4 to about 8 which contains a high protein concentration, including an antibody concentration ranging from about 1 to about 150 mg/ml, and has enhanced stability. This invention also pertains to a liquid aqueous pharmaceutical formulation for therapeutic use in a subject suffering from a condition characterized by detrimental TNF $\alpha$  activity. The formulation of the invention comprises the following constituents: an antibody which binds to human TNF $\alpha$  with high affinity, a low off rate and high neutralizing capacity; a buffer, which includes citric acid, sodium citrate, disodium phosphate dihydrate, and sodium dihydrogen phosphate dihydrate; tonicity agents, which include mannitol and sodium chloride; a detergent, including polysorbate 80; and sodium hydroxide, for pH adjustment.

### 15 Definitions

In order that the present invention may be more readily understood, certain terms are first defined.

The term "subject" is intended to include living organisms, *e.g.*, prokaryotes and eukaryotes. Examples of subjects include mammals, *e.g.*, humans, dogs, cows, horses, pigs, sheep, goats, cats, mice, rabbits, rats, and transgenic non-human animals. In specific embodiments of the invention, the subject is a human.

The term "pharmaceutical formulation" refers to preparations which are in such form as to permit the biological activity of the active ingredients to be unequivocally effective, and which contain no additional components which are significantly toxic to the subjects to which the formulation would be administered. "Pharmaceutically acceptable" excipients (vehicles, additives) are those which can reasonably be administered to a subject mammal to provide an effective dose of the active ingredient employed.

A "stable" formulation is one in which the antibody therein essentially retains its physical stability and/or chemical stability and/or biological activity upon storage. Various analytical techniques for measuring protein stability are available in the art and are reviewed in Peptide and Protein Drug Delivery, 247-301, Vincent Lee Ed., Marcel Dekker, Inc., New York, N.Y., Pubs. (1991) and Jones, A. Adv. Drug Delivery Rev. 10: 29-90 (1993), for example. Stability can be measured at a selected temperature for a selected time period. Preferably, the formulation is stable at room temperature (about 30°C) or at 40°C for at least 1 month and/or stable at about 2-8°C. for at least 1 year for



- 9 -

at least 2 years. Furthermore, the formulation is preferably stable following freezing (to, *e.g.*, -70°C) and thawing of the formulation, hereinafter referred to as a "freeze/thaw cycle."

An antibody "retains its physical stability" in a pharmaceutical formulation if it shows substantially no signs of aggregation, precipitation and/or denaturation upon visual examination of color and/or clarity, or as measured by UV light scattering or by size exclusion chromatography.

An antibody "retains its chemical stability" in a pharmaceutical formulation, if the chemical stability at a given time is such that the antibody is considered to still retain its biological activity as defined below. Chemical stability can be assessed by detecting and quantifying chemically altered forms of the antibody. Chemical alteration may involve size modification (*e.g.* clipping) which can be evaluated using size exclusion chromatography, SDS-PAGE and/or matrix-assisted laser desorption ionization/time-of-flight mass spectrometry (MALDI/TOF MS), for example. Other types of chemical alteration include charge alteration (*e.g.* occurring as a result of deamidation) which can be evaluated by ion-exchange chromatography, for example.

An antibody "retains its biological activity" in a pharmaceutical formulation, if the antibody in a pharmaceutical formulation is biologically active for its intended purpose. For example, biological activity is retained if the biological activity of the antibody in the pharmaceutical formulation is within about 30%, about 20%, or about 10% (within the errors of the assay) of the biological activity exhibited at the time the pharmaceutical formulation was prepared (*e.g.*, as determined in an antigen binding assay).

"Isotonic" is a term recognized in the art. Isotonic can mean, for example, that the formulation of interest has essentially the same osmotic pressure as human blood. Isotonic formulations will generally have an osmotic pressure from about 250 to 350 mOsm. Isotonicity can be measured using a vapor pressure or ice-freezing type osmometer, for example. A "tonicity agent" is a compound which renders the formulation isotonic.

A "polyol" is a substance with multiple hydroxyl groups, and includes sugars (reducing and nonreducing sugars), sugar alcohols and sugar acids. Preferred polyols herein have a molecular weight which is less than about 600 kD (*e.g.* in the range from about 120 to about 400 kD). A "reducing sugar" is one which contains a hemiacetal group that can reduce metal ions or react covalently with lysine and other amino groups in proteins and a "nonreducing sugar" is one which does not have these properties of a reducing sugar. Examples of reducing sugars are fructose, mannose, maltose, lactose, arabinose, xylose, ribose, rhamnose, galactose and glucose. Nonreducing sugars include



- 10 -

sucrose, trehalose, sorbose, melezitose and raffinose. Mannitol, xylitol, erythritol, threitol, sorbitol and glycerol are examples of sugar alcohols. As to sugar acids, these include L-gluconate and metallic salts thereof. Where it desired that the formulation is freeze-thaw stable, the polyol is preferably one which does not crystallize at freezing  
5 temperatures (*e.g.* -20°C) such that it destabilizes the antibody in the formulation. The polyol may also act as a tonicity agent. In one embodiment of the invention, one ingredient of the formulation is mannitol in a concentration of 5 to 20 mg/ml. In a preferred embodiment of the invention, the concentration of mannitol is 7.5 to 15 mg/ml. In a more preferred embodiment of the invention, the concentration of mannitol  
10 is 10-14 mg/ml.

As used herein, "buffer" refers to a buffered solution that resists changes in pH by the action of its acid-base conjugate components. The buffer of this invention has a pH in the range from about 4 to about 8; preferably from about 4.5 to about 7; and most preferably has a pH in the range from about 5.0 to about 6.5. Examples of buffers that  
15 will control the pH in this range include acetate (*e.g.* sodium acetate), succinate (such as sodium succinate), gluconate, histidine, citrate and other organic acid buffers.

In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" or "effective amount" of an antibody refers to an amount effective in the prevention or treatment of a disorder for the treatment of which  
20 the antibody is effective. A "disorder" is any condition that would benefit from treatment with the antibody. This includes chronic and acute disorders or diseases including those pathological conditions which predisposes the subject to the disorder in question.

A "preservative" is a compound which can be included in the formulation to  
25 essentially reduce bacterial action therein, thus facilitating the production of a multi-use formulation, for example. Examples of potential preservatives include octadecyldimethylbenzyl ammonium chloride, hexamethonium chloride, benzalkonium chloride (a mixture of alkylbenzyltrimethylammonium chlorides in which the alkyl groups are long-chain compounds), and benzethonium chloride. Other types of  
30 preservatives include aromatic alcohols such as phenol, butyl and benzyl alcohol, alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, 3-pentanol, and m-cresol.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as  
35 those in which the disorder is to be prevented.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration,



- 11 -

usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

5           The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

10           The phrase "pharmaceutically acceptable carrier" is art recognized and includes a pharmaceutically acceptable material, composition or vehicle, suitable for administration to mammals. The carriers include liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject agent from one organ, or portion of the body, to another organ, or portion of the body. Each  
15 carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient.

          The term "human TNF $\alpha$ " (abbreviated herein as hTNF $\alpha$ , or simply hTNF), as used herein, is intended to refer to a human cytokine that exists as a 17 kD secreted form and a 26 kD membrane associated form, the biologically active form of which is  
20 composed of a trimer of noncovalently bound 17 kD molecules. The structure of hTNF $\alpha$  is described further in, for example, Pennica, D., *et al.* (1984) *Nature* 312:724-729; Davis, J.M., *et al.* (1987) *Biochemistry* 26:1322-1326; and Jones, E.Y., *et al.* (1989) *Nature* 338:225-228. The term human TNF $\alpha$  is intended to include recombinant human TNF $\alpha$  (rhTNF $\alpha$ ), which can be prepared by standard recombinant expression  
25 methods or purchased commercially (R & D Systems, Catalog No. 210-TA, Minneapolis, MN).

          The term "antibody", as used herein, is intended to refer to immunoglobulin molecules comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy  
30 chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into  
35 regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-



- 12 -

terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In one embodiment of the invention, the formulation contains an antibody with CDR1, CDR2, and CDR3 sequences like those described in U.S. Patent Nos. 6,090,382 and 6,258,562.

5           The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (*e.g.*, hTNF $\alpha$ ). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')<sub>2</sub> fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123). In one embodiment of the invention, the formulation contains an antigen-binding portions described in U.S. Patent Nos. 6,090,382 and 6,258,562, each incorporated by reference herein.

35           Still further, an antibody or antigen-binding portion thereof may be part of a larger immunoadhesion molecules, formed by covalent or noncovalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov, S.M., *et al.* (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal



- 13 -

polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S.M., *et al.* (1994) *Mol. Immunol.* 31:1047-1058). Antibody portions, such as Fab and F(ab')<sub>2</sub> fragments, can be prepared from whole antibodies using conventional techniques, such as papain or pepsin digestion, respectively, of whole antibodies. Moreover, antibodies, antibody portions and immunoadhesion molecules can be obtained using standard recombinant DNA techniques, as described herein.

The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (*e.g.*, mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further in Section II, below), antibodies isolated from a recombinant, combinatorial human antibody library (described further in Section III, below), antibodies isolated from an animal (*e.g.*, a mouse) that is transgenic for human immunoglobulin genes (see *e.g.*, Taylor, L.D., *et al.* (1992) *Nucl. Acids Res.* 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

An "isolated antibody", as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (*e.g.*, an isolated antibody that specifically binds hTNF $\alpha$  is substantially free of antibodies that specifically bind antigens other than hTNF $\alpha$ ). An isolated antibody that specifically binds hTNF $\alpha$  may, however, have cross-reactivity to other antigens, such as TNF $\alpha$



- 14 -

molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

A "neutralizing antibody", as used herein (or an "antibody that neutralized hTNF $\alpha$  activity"), is intended to refer to an antibody whose binding to hTNF $\alpha$  results in inhibition of the biological activity of hTNF $\alpha$ . This inhibition of the biological activity of hTNF $\alpha$  can be assessed by measuring one or more indicators of hTNF $\alpha$  biological activity, such as hTNF $\alpha$ -induced cytotoxicity (either *in vitro* or *in vivo*), hTNF $\alpha$ -induced cellular activation and hTNF $\alpha$  binding to hTNF $\alpha$  receptors. These indicators of hTNF $\alpha$  biological activity can be assessed by one or more of several standard *in vitro* or *in vivo* assays known in the art, and described in U.S. Patent Nos. 6,090,382 and 6,258,562. Preferably, the ability of an antibody to neutralize hTNF $\alpha$  activity is assessed by inhibition of hTNF $\alpha$ -induced cytotoxicity of L929 cells. As an additional or alternative parameter of hTNF $\alpha$  activity, the ability of an antibody to inhibit hTNF $\alpha$ -induced expression of ELAM-1 on HUVEC, as a measure of hTNF $\alpha$ -induced cellular activation, can be assessed.

The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, NJ). For further descriptions, see Jönsson, U., *et al.* (1993) *Ann. Biol. Clin.* 51:19-26; Jönsson, U., *et al.* (1991) *Biotechniques* 11:620-627; Johnson, B., *et al.* (1995) *J. Mol. Recognit.* 8:125-131; and Johnson, B., *et al.* (1991) *Anal. Biochem.* 198:268-277.

The term " $K_{off}$ ", as used herein, is intended to refer to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

The term " $K_d$ ", as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction.

## II. Antibodies of the Formulation

The invention is directed to a liquid aqueous pharmaceutical formulation comprising a therapeutically effective amount of an antibody in a buffered solution forming a formulation having a pH between about 4 and about 8 and having an extended shelf life, preferably of at least about 18 months. In another embodiment, the liquid aqueous pharmaceutical formulation of the invention has enhanced stability. In a further embodiment of the invention, the formulation is not light sensitive. In yet another embodiment of the invention, the claimed formulation remains stable following at least



- 15 -

3 freeze/thaw cycles. In still another embodiment, the pharmaceutical formulation of the invention is suitable for single use sc injection.

Antibodies that can be used in the formulation include polyclonal, monoclonal, recombinant antibodies, single chain antibodies, hybrid antibodies, chimeric antibodies, 5 humanized antibodies, or fragments thereof. Antibody-like molecules containing one or two binding sites for an antigen and a Fc-part of an immunoglobulin can also be used. An example of an antibody-like molecule is the active ingredient etanercept or infliximab. Preferred antibodies used in the formulation are human antibodies which are cloned from human cells or from gene-archives representing the human antibody- 10 reservoir. Especially preferred among the human antibodies are antibodies directed against the antigen TNF $\alpha$ , including human TNF $\alpha$  (or hTNF $\alpha$ ).

In one embodiment, the formulation of the invention includes a combination of antibodies (two or more), or a "cocktail" of antibodies. For example, the formulation can include the antibody D2E7 and one or more additional antibodies.

15 In a preferred embodiment of the invention, the formulation contains an antibody, or antigen-binding portion thereof, dissociates from human TNF $\alpha$  with a  $K_d$  of  $1 \times 10^{-8}$  M or less and a  $K_{off}$  rate constant of  $1 \times 10^{-3}$  s $^{-1}$  or less, both determined by surface plasmon resonance, and neutralizes human TNF $\alpha$  cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$  of  $1 \times 10^{-7}$  M or less. In another preferred embodiment, 20 the formulation of the invention contains an antibody, or antigen-binding portion thereof, like those described in U.S. Patent Nos. 6,090,382 and 6,258,562.

In one aspect, the formulation of the invention contains D2E7 antibodies and antibody portions, D2E7-related antibodies and antibody portions, and other human 25 antibodies and antibody portions with equivalent properties to D2E7, such as high affinity binding to hTNF $\alpha$  with low dissociation kinetics and high neutralizing capacity. In another one embodiment, the formulation of the invention contains an isolated human antibody, or an antigen-binding portion thereof, that dissociates from human TNF $\alpha$  with a  $K_d$  of  $1 \times 10^{-8}$  M or less and a  $K_{off}$  rate constant of  $1 \times 10^{-3}$  s $^{-1}$  or less, both 30 determined by surface plasmon resonance, and neutralizes human TNF $\alpha$  cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$  of  $1 \times 10^{-7}$  M or less. More preferably, the isolated human antibody, or antigen-binding portion thereof, dissociates from human TNF $\alpha$  with a  $K_{off}$  of  $5 \times 10^{-4}$  s $^{-1}$  or less, or even more preferably, with a  $K_{off}$  of  $1 \times 10^{-4}$  s $^{-1}$  or less. More preferably, the isolated human antibody, or antigen-binding portion 35 thereof, neutralizes human TNF $\alpha$  cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$  of  $1 \times 10^{-8}$  M or less, even more preferably with an IC $_{50}$  of  $1 \times 10^{-9}$  M or less and still more preferably with an IC $_{50}$  of  $5 \times 10^{-10}$  M or less. In a preferred embodiment, the



- 16 -

formulation contains an antibody which is an isolated human recombinant antibody, or an antigen-binding portion thereof. In another preferred embodiment, the formulation contains an antibody which also neutralizes TNF $\alpha$ -induced cellular activation, as assessed using a standard *in vitro* assay for TNF $\alpha$ -induced ELAM-1 expression on  
5 human umbilical vein endothelial cells (HUVEC).

### III. Preparation of Formulation

10 The present invention features formulations (*e.g.*, protein formulations and/or antibody formulations) having improved properties as compared to art-recognized formulations. For example, the formulations of the invention have an improved shelf life and/or stability as compared to art recognized formulations. In a preferred aspect, the formulations of the invention comprise a high protein concentration, including, for  
15 example, a protein concentration greater than about 45 mg/ml, a protein concentration greater than about 50 mg/ml, a protein concentration greater than about 100 mg/ml, or a protein concentration greater than about 150 mg/ml. In a preferred embodiment of the invention, the protein is an antibody. In another preferred embodiment, the antibody is D2E7. The invention also provides an aqueous pharmaceutical composition comprising  
20 a polyol, a surfactant, and a buffer system comprising citrate and/or phosphate with a pH of about 4 to 8, in amounts sufficient to formulate an antibody for therapeutic use at a concentration of greater than about, for example, 45 mg/ml.

Preparation of the antibody of interest is performed according to standard methods known in the art. In a preferred embodiment of the invention, the antibody  
25 used in the formulation is expressed in CHO cells and purified by a standard series of chromatography steps. In a further preferred embodiment, the antibody is directed to hTNF $\alpha$ , and is prepared according to the methods described in U.S. Patent Nos. 6,090,382 and 6,258,562.

After preparation of the antibody of interest, the pharmaceutical formulation  
30 comprising the antibody is prepared. The therapeutically effective amount of antibody present in the formulation is determined, for example, by taking into account the desired dose volumes and mode(s) of administration. In one embodiment of the invention, the concentration of the antibody in the formulation is between about 1 to about 150 mg of antibody per ml of liquid formulation. In a preferred embodiment, the concentration of  
35 the antibody in the formulation is between about 5 to about 80 mg per ml. In another preferred embodiment, the concentration of the antibody in the formulation is between about 25 to about 50 mg/ml. The formulation is especially suitable for large antibody



- 17 -

dosages of more than 15 mg/ml. In a preferred embodiment, the concentration of the antibody is 50 mg/ml.

In another embodiment of the invention, the concentration of the antibody in the formulation is about 1-150 mg/ml, about 5-145 mg/ml, about 10-140 mg/ml, about 15-  
5 135 mg/ml, about 20-130 mg/ml, about 25-125 mg/ml, about 30-120 mg/ml, about 35-  
115 mg/ml, about 40-110 mg/ml, about 45-105 mg/ml, about 50-100 mg/ml, about 55-  
95 mg/ml, about 60-90 mg/ml, about 65-85 mg/ml, about 70-80 mg/ml, or about 75  
mg/ml. Ranges intermediate to the above recited concentrations, *e.g.*, about 6-144  
10 mg/ml, are also intended to be part of this invention. For example, ranges of values  
using a combination of any of the above recited values as upper and/or lower limits are  
intended to be included.

In one embodiment, the invention provides a formulation with an extended shelf  
life comprising of an active ingredient, preferably an antibody, in combination with  
mannitol, citric acid monohydrate, sodium citrate, disodium phosphate dihydrate,  
15 sodium dihydrogen phosphate dihydrate, sodium chloride, polysorbate 80, water, and  
sodium hydroxide. In a further embodiment, the formulation of the invention has an  
extended shelf life of at least about 18 months in the liquid state. Freezing the  
formulation of the invention can also be used to further extend its shelf life.

An aqueous formulation is prepared comprising the antibody in a pH-buffered  
20 solution. The buffer of this invention has a pH ranging from about 4 to about 8,  
preferably from about 4.5 to about 6.0, more preferably from about 4.8 to about 5.5, and  
most preferably has a pH of about 5.0 to about 5.2. Ranges intermediate to the above  
recited pH's are also intended to be part of this invention. For example, ranges of values  
using a combination of any of the above recited values as upper and/or lower limits are  
25 intended to be included. Examples of buffers that will control the pH within this range  
include acetate (*e.g.* sodium acetate), succinate (such as sodium succinate), gluconate,  
histidine, citrate and other organic acid buffers.

In a preferred embodiment of the invention, the formulation comprises a buffer  
system which contains citrate and phosphate to maintain the pH in a range of about 4 to  
30 about 8. In a further preferred embodiment the pH range is from about 4.5 to about 6.0,  
more preferably from about pH 4.8 to about 5.5, and most preferably in a pH range of  
about 5.0 to about 5.2. In another preferred embodiment, the buffer system includes  
citric acid monohydrate, sodium citrate, disodium phosphate dihydrate, and/or sodium  
dihydrogen phosphate dihydrate. In a further preferred embodiment, the buffer system  
35 includes about 1.3 mg/ml of citric acid (*e.g.*, 1.305 mg/ml), about 0.3 mg/ml of sodium  
citrate (*e.g.*, 0.305 mg/ml), about 1.5 mg/ml of disodium phosphate dihydrate (*e.g.* 1.53  
mg/ml), about 0.9 mg/ml of sodium dihydrogen phosphate dihydrate (*e.g.*, 0.86), and



- 18 -

about 6.2 mg/ml of sodium chloride (*e.g.*, 6.165 mg/ml). In additional preferred  
embodiments, the buffer system includes 1-1.5 mg/ml of citric acid, 0.25 to 0.5 mg/ml  
of sodium citrate, 1.25 to 1.75 mg/ml of of disodium phosphate dihydrate, 0.7 to 1.1  
mg/ml of sodium dihydrogen phosphate dihydrate, and 6.0 to 6.4 mg/ml of sodium  
5 chloride. In a further embodiment, the pH of the formulation is adjusted with sodium  
hydroxide.

A polyol, which acts as a tonicifier and may stabilize the antibody, is also  
included in the formulation. The polyol is added to the formulation in an amount which  
may vary with respect to the desired isotonicity of the formulation. Preferably the  
10 aqueous formulation is isotonic. The amount of polyol added may also alter with  
respect to the molecular weight of the polyol. For example, a lower amount of a  
monosaccharide (*e.g.* mannitol) may be added, compared to a disaccharide (such as  
trehalose). In a preferred embodiment of the invention, the polyol which is used in the  
formulation as a tonicity agent is mannitol. In a preferred embodiment of the invention,  
15 the mannitol concentration is about 5 to 20 mg/ml. In another preferred embodiment of  
the invention, the concentration of mannitol is about 7.5 to 15 mg/ml. In a more  
preferred embodiment of the formulation of the invention, the concentration of mannitol  
is about 10-14 mg/ml. In the most preferred embodiment, the concentration of mannitol  
is about 12 mg/ml. In another embodiment of the invention, the polyol sorbitol is  
20 included in the formulation.

A detergent or surfactant is also added to the antibody formulation. Exemplary  
detergents include nonionic detergents such as polysorbates (*e.g.* polysorbates 20, 80  
etc) or poloxamers (*e.g.* poloxamer 188). The amount of detergent added is such that it  
reduces aggregation of the formulated antibody and/or minimizes the formation of  
25 particulates in the formulation and/or reduces adsorption. In a preferred embodiment of  
the invention, the formulation includes a surfactant which is a polysorbate. In another  
preferred embodiment of the invention, the formulation contains the detergent  
polysorbate 80 or Tween 80. Tween 80 is a term used to describe polyoxyethylene (20)  
sorbitanmonooleate (see Fiedler, Lexikon der Hifsstoffe, Editio Cantor Verlag  
30 Aulendorf, 4th edi., 1996). In one preferred embodiment, the formulation contains  
between about 0.1 and about 10 mg/ml of polysorbate 80, more preferably between  
about 0.5 and about 5 mg/ml. In another preferred embodiment, about 0.1% polysorbate  
80 is found in the formulation of the invention.

In a preferred embodiment of the invention, the formulation is a 0.8 mL solution  
35 in a vial containing the ingredients shown below in Table 1.

- 19 -

**Table 1****1 vial with 0.8 mL solution for injection<sup>1)</sup> contains:**

<b>Name of ingredient</b>	<b>Quantity</b>	<b>Function</b>
<i>Active substance:</i> Antibody (D2E7) <sup>2)</sup>	40.0 mg	Active substance
<i>Excipients:</i>		
Mannitol	9.6 mg	Tonicity agent
Citric acid monohydrate Citric acid	1.044 mg	Buffer
Sodium citrate Sodium citrate	0.244 mg	Buffer
Disodium phosphate dihydrate Dibasic sodium phosphate dihydrate	1.224 mg	Buffer
Sodium dihydrogen phosphate dihydrate Monobasic sodium phosphate dihydrate	0.688 mg	Buffer
Sodium chloride	4.932 mg	Tonicity agent
Polysorbate 80	0.8 mg	Detergent
Water for injections Water for injection	759.028 – 759.048 mg	Solvent
Sodium hydroxide <sup>3)</sup>	0.02 - 0.04 mg	pH adjustment
<b>Total</b>	817.6 mg	

5 <sup>1)</sup> Density of the solution: 1.022 g/mL<sup>2)</sup> Is used as concentrate<sup>3)</sup> Addition as 1M solution

10 In one embodiment, the formulation contains the above-identified agents (i.e. antibody, buffer, polyol and detergent) and is essentially free of one or more preservatives, such as benzyl alcohol, phenol, m-cresol, chlorobutanol and benzethonium Cl. In another embodiment, a preservative may be included in the formulation, particularly where the formulation is a multidose formulation. One or more other pharmaceutically acceptable carriers, excipients or stabilizers such as those

15 described in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980) may be included in the formulation provided that they do not significantly adversely affect the desired characteristics of the formulation. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed and include; additional buffering agents; co-solvents; antioxidants including ascorbic acid

20 and methionine; chelating agents such as EDTA; metal complexes (e.g. Zn-protein



- 20 -

complexes); biodegradable polymers such as polyesters; and/or salt-forming counterions such as sodium.

The formulation herein may also be combined with one or more other therapeutic agents as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect the antibody of the formulation. Such therapeutic agents are suitably present in combination in amounts that are effective for the purpose intended. Additional therapeutic agents which can be combined with the formulation of the invention are further described in U.S. Patent Nos. 6,090,382 and 6,258,562, each of which is incorporated herein by reference.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to, or following, preparation of the formulation.

#### IV. Administration of Formulation

15

The formulation of the invention can be used in similar indications as those described in U.S. Patent Nos. 6,090,382 and 6,258,562, each is further detailed below.

The language "effective amount" of the formulation is that amount necessary or sufficient to inhibit TNF $\alpha$  activity, *e.g.*, prevent the various morphological and somatic symptoms of a detrimental TNF $\alpha$  activity-associated state. In another embodiment, the effective amount of the formulation is the amount necessary to achieve the desired result. In one example, an effective amount of the formulation is the amount sufficient to inhibit detrimental TNF $\alpha$  activity. In another example, an effective amount of the formulation is 0.8 mL of the formulation containing 40 mg of antibody, as described in table 1. The effective amount can vary depending on such factors as the size and weight of the subject, or the type of illness. For example, the choice of a TNF $\alpha$  activity-inhibiting formulation can affect what constitutes an "effective amount". One of ordinary skill in the art would be able to study the aforementioned factors and make the determination regarding the effective amount of the TNF $\alpha$  activity inhibiting formulation without undue experimentation.

The regimen of administration can affect what constitutes an effective amount. The TNF $\alpha$  activity-inhibiting formulation can be administered to the subject either prior to or after the onset of detrimental TNF $\alpha$  activity. Further, several divided dosages, as well as staggered dosages, can be administered daily or sequentially, or the dose can be continuously infused, or can be a bolus injection. Further, the dosages of the TNF $\alpha$  activity-inhibiting formulation can be proportionally increased or decreased as indicated



- 21 -

by the exigencies of the therapeutic or prophylactic situation.

The term "treated," "treating" or "treatment" includes the diminishment or alleviation of at least one symptom associated or caused by the state, disorder or disease being treated. For example, treatment can be diminishment of one or several symptoms  
5 of a disorder or complete eradication of a disorder.

Actual dosage levels of the active ingredients (antibody) in the pharmaceutical formulation of this invention may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

10 The selected dosage level will depend upon a variety of factors including the activity of the antibody found in the formulation, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health  
15 and prior medical history of the patient being treated, and like factors well known in the medical arts.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition of the present invention required. For example, the physician or veterinarian could start doses of the  
20 compounds of the invention employed in the pharmaceutical formulation at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In general, a suitable daily dose of a formulation of the invention will be that amount of the formulation that is the lowest dose effective to produce a therapeutic  
25 effect. Such an effective dose will generally depend upon the factors described above. An effective amount of the formulation of the present invention is an amount that inhibits TNF $\alpha$  activity in a subject suffering from a disorder in which TNF $\alpha$  activity is detrimental. In a preferred embodiment, the formulation provides an effective dose of 40 mg per injection of the active ingredient, the antibody. In another embodiment, the  
30 formulation provides an effective dose which ranges from about 1 to 150 mg of antibody. If desired, the effective daily dose of the pharmaceutical formulation may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms.

In one embodiment of the invention, the dosage of the antibody in the  
35 formulation is between about 5 to about 80 mg. In another embodiment, the dosage of the antibody in the formulation is between about 25 to about 50 mg. The formulation is especially suitable for large antibody dosages of more than 15 mg. In a preferred



- 22 -

embodiment of the invention, the formulation provides an antibody at a dose of about 40 mg. In another preferred embodiment, the antibody is directed to TNF $\alpha$ . In the most preferred embodiment, the antibody is D2E7.

In one embodiment of the invention, the dosage of the antibody in the  
5 formulation is between about 1-150 mg, about 5-145 mg, about 10-140 mg, about 15-135 mg, about 20-130 mg, about 25-125 mg, about 30-120 mg, about 35-115 mg, about 40-110 mg, about 45-105 mg, about 50-100 mg, about 55-95 mg, about 60-90 mg, about 65-85 mg, about 70-80 mg, or about 75 mg. In a preferred embodiment, the dosage of the antibody is 40 mg. In a further preferred embodiment, the antibody is directed to  
10 TNF $\alpha$ . In the most preferred embodiment, the antibody is D2E7. Ranges intermediate to the above recited dosages, *e.g.*, about 2-149 mg, are also intended to be part of this invention. For example, ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included.

It is to be noted that dosage values may vary with the severity of the condition to  
15 be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

20 The invention provides a pharmaceutical formulation with an extended shelf life, which, in one embodiment, is used to inhibit TNF $\alpha$  activity in a subject suffering from a disorder in which TNF $\alpha$  activity is detrimental, comprising administering to the subject an antibody or antibody portion of the invention such that TNF $\alpha$  activity in the subject is inhibited. Preferably, the TNF $\alpha$  is human TNF $\alpha$  and the subject is a human subject.  
25 Alternatively, the subject can be a mammal expressing a TNF $\alpha$  with which an antibody of the invention cross-reacts. Still further the subject can be a mammal into which has been introduced hTNF $\alpha$  (*e.g.*, by administration of hTNF $\alpha$  or by expression of an hTNF $\alpha$  transgene). A formulation of the invention can be administered to a human subject for therapeutic purposes (discussed further below). In one embodiment of the  
30 invention, the liquid pharmaceutical formulation is easily administratable, which includes, for example, a formulation which is self-administered by the patient. In a preferred embodiment, the formulation of the invention is administered through sc injection, preferably single use. Moreover, a formulation of the invention can be administered to a non-human mammal expressing a TNF $\alpha$  with which the antibody  
35 cross-reacts (*e.g.*, a primate, pig or mouse) for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal models may be useful for evaluating the therapeutic efficacy of antibodies of the invention (*e.g.*, testing of dosages



- 23 -

and time courses of administration).

As used herein, the term "a disorder in which TNF $\alpha$  activity is detrimental" is intended to include diseases and other disorders in which the presence of TNF $\alpha$  in a subject suffering from the disorder has been shown to be or is suspected of being either  
5 responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which TNF $\alpha$  activity is detrimental is a disorder in which inhibition of TNF $\alpha$  activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of TNF $\alpha$  in a biological fluid of a subject  
10 suffering from the disorder (*e.g.*, an increase in the concentration of TNF $\alpha$  in serum, plasma, synovial fluid, *etc.* of the subject), which can be detected, for example, using an anti-TNF $\alpha$  antibody as described above.

There are numerous examples of disorders in which TNF $\alpha$  activity is detrimental. Examples of disorders in which TNF $\alpha$  activity is detrimental are described  
15 in PCT Publication No. WO 2004/009776. Examples in which TNF $\alpha$  activity is detrimental are also described in U.S. Patent Nos. 6,015,557, 6,177,077, 6,379,666, 6,419,934, 6,419,944, 6,423,321, and 6,428,787; U.S. Patent Publ. Nos. US2001/0016195, US2001/0004456 and US2001/026801; PCT Publ. Nos. WO 00/50079 and WO 01/49321.

20 The use of the antibodies and antibody portions of the invention in the treatment of specific disorders is discussed further below:

#### A. Sepsis

Tumor necrosis factor has an established role in the pathophysiology of sepsis,  
25 with biological effects that include hypotension, myocardial suppression, vascular leakage syndrome, organ necrosis, stimulation of the release of toxic secondary mediators and activation of the clotting cascade (see *e.g.*, Tracey, K.J. and Cerami, A. (1994) *Annu. Rev. Med.* 45:491-503; Russell, D and Thompson, R.C. (1993) *Curr. Opin. Biotech.* 4:714-721). Accordingly, the formulation of the invention can be used to treat  
30 sepsis in any of its clinical settings, including septic shock, endotoxic shock, gram negative sepsis and toxic shock syndrome.

Furthermore, to treat sepsis, the formulation of the invention can be coadministered with one or more additional therapeutic agents that may further alleviate sepsis, such as an interleukin-1 inhibitor (such as those described in PCT Publication  
35 Nos. WO 92/16221 and WO 92/17583), the cytokine interleukin-6 (see *e.g.*, PCT Publication No. WO 93/11793) or an antagonist of platelet activating factor (see *e.g.*, European Patent Application Publication No. EP 374 510).



- 24 -

Additionally, in a preferred embodiment, the formulation of the invention is administered to a human subject within a subgroup of sepsis patients having a serum or plasma concentration of IL-6 above 500 pg/ml, and more preferably 1000 pg/ml, at the time of treatment (see PCT Publication No. WO 95/20978 by Daum, L., *et al.*).

5

### B. Autoimmune Diseases

Tumor necrosis factor has been implicated in playing a role in the pathophysiology of a variety of autoimmune diseases. For example, TNF $\alpha$  has been implicated in activating tissue inflammation and causing joint destruction in rheumatoid arthritis (see *e.g.*, Tracey and Cerami, *supra*; Arend, W.P. and Dayer, J-M. (1995) *Arth. Rheum.* 38:151-160; Fava, R.A., *et al.* (1993) *Clin. Exp. Immunol.* 94:261-266). TNF $\alpha$  also has been implicated in promoting the death of islet cells and in mediating insulin resistance in diabetes (see *e.g.*, Tracey and Cerami, *supra*; PCT Publication No. WO 94/08609). TNF $\alpha$  also has been implicated in mediating cytotoxicity to

10 : oligodendrocytes and induction of inflammatory plaques in multiple sclerosis (see *e.g.*, Tracey and Cerami, *supra*). Chimeric and humanized murine anti-hTNF $\alpha$  antibodies have undergone clinical testing for treatment of rheumatoid arthritis (see *e.g.*, Elliott, M.J., *et al.* (1994) *Lancet* 344:1125-1127; Elliot, M.J., *et al.* (1994) *Lancet* 344:1105-1110; Rankin, E.C., *et al.* (1995) *Br. J. Rheumatol.* 34:334-342).

20

The formulation of the invention can be used to treat autoimmune diseases, in particular those associated with inflammation, including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis and nephrotic syndrome. Typically, the formulation is administered systemically, although for certain disorders, local

25 : administration of the antibody or antibody portion at a site of inflammation may be beneficial (*e.g.*, local administration in the joints in rheumatoid arthritis or topical application to diabetic ulcers, alone or in combination with a cyclohexane-ylidene derivative as described in PCT Publication No. WO 93/19751).

30

### C. Infectious Diseases

Tumor necrosis factor has been implicated in mediating biological effects observed in a variety of infectious diseases. For example, TNF $\alpha$  has been implicated in mediating brain inflammation and capillary thrombosis and infarction in malaria (see *e.g.*, Tracey and Cerami, *supra*). TNF $\alpha$  also has been implicated in mediating brain

35 : inflammation, inducing breakdown of the blood-brain barrier, triggering septic shock syndrome and activating venous infarction in meningitis (see *e.g.*, Tracey and Cerami, *supra*). TNF $\alpha$  also has been implicated in inducing cachexia, stimulating viral



- 25 -

proliferation and mediating central nervous system injury in acquired immune deficiency syndrome (AIDS) (see *e.g.*, Tracey and Cerami, *supra*). Accordingly, the antibodies, and antibody portions, of the invention, can be used in the treatment of infectious diseases, including bacterial meningitis (see *e.g.*, European Patent Application  
5 Publication No. EP 585 705), cerebral malaria, AIDS and AIDS-related complex (ARC) (see *e.g.*, European Patent Application Publication No. EP 230 574), as well as cytomegalovirus infection secondary to transplantation (see *e.g.*, Fietze, E., *et al.* (1994) *Transplantation* 58:675-680). The formulation of the invention, also can be used to alleviate symptoms associated with infectious diseases, including fever and myalgias  
10 due to infection (such as influenza) and cachexia secondary to infection (*e.g.*, secondary to AIDS or ARC).

#### D. Transplantation

Tumor necrosis factor has been implicated as a key mediator of allograft  
15 rejection and graft versus host disease (GVHD) and in mediating an adverse reaction that has been observed when the rat antibody OKT3, directed against the T cell receptor CD3 complex, is used to inhibit rejection of renal transplants (see *e.g.*, Tracey and Cerami, *supra*; Eason, J.D., *et al.* (1995) *Transplantation* 59:300-305; Suthanthiran, M. and Strom, T.B. (1994) *New Engl. J. Med.* 331:365-375). Accordingly, the formulation  
20 of the invention, can be used to inhibit transplant rejection, including rejections of allografts and xenografts and to inhibit GVHD. Although the antibody or antibody portion may be used alone, more preferably it is used in combination with one or more other agents that inhibit the immune response against the allograft or inhibit GVHD. For example, in one embodiment, the formulation of the invention is used in  
25 combination with OKT3 to inhibit OKT3-induced reactions. In another embodiment, the formulation of the invention is used in combination with one or more antibodies directed at other targets involved in regulating immune responses, such as the cell surface molecules CD25 (interleukin-2 receptor- $\alpha$ ), CD11a (LFA-1), CD54 (ICAM-1), CD4, CD45, CD28/CTLA4, CD80 (B7-1) and/or CD86 (B7-2). In yet another  
30 embodiment, the formulation of the invention is used in combination with one or more general immunosuppressive agents, such as cyclosporin A or FK506.

#### E. Malignancy

Tumor necrosis factor has been implicated in inducing cachexia, stimulating  
35 tumor growth, enhancing metastatic potential and mediating cytotoxicity in malignancies (see *e.g.*, Tracey and Cerami, *supra*). Accordingly, the formulation of the invention, can be used in the treatment of malignancies, to inhibit tumor growth or



- 26 -

metastasis and/or to alleviate cachexia secondary to malignancy. The formulation may be administered systemically or locally to the tumor site.

#### F. Pulmonary Disorders

5 Tumor necrosis factor has been implicated in the pathophysiology of adult respiratory distress syndrome, including stimulating leukocyte-endothelial activation, directing cytotoxicity to pneumocytes and inducing vascular leakage syndrome (see *e.g.*, Tracey and Cerami, *supra*). Accordingly, the formulation of the invention, can be used to treat various pulmonary disorders, including adult respiratory distress syndrome (see  
10 *e.g.*, PCT Publication No. WO 91/04054), shock lung, chronic pulmonary inflammatory disease, pulmonary sarcoidosis, pulmonary fibrosis and silicosis. The formulation may be administered systemically or locally to the lung surface, for example as an aerosol.

#### G. Intestinal Disorders

15 Tumor necrosis factor has been implicated in the pathophysiology of inflammatory bowel disorders (see *e.g.*, Tracy, K.J., *et al.* (1986) *Science* 234:470-474; Sun, X-M., *et al.* (1988) *J. Clin. Invest.* 81:1328-1331; MacDonald, T.T., *et al.* (1990) *Clin. Exp. Immunol.* 81:301-305). Chimeric murine anti-hTNF $\alpha$  antibodies have undergone clinical testing for treatment of Crohn's disease (van Dullemen, H.M., *et al.*  
20 (1995) *Gastroenterology* 109:129-135). The formulation of the invention, also can be used to treat intestinal disorders, such as idiopathic inflammatory bowel disease, which includes two syndromes, Crohn's disease and ulcerative colitis.

#### H. Cardiac Disorders

25 The formulation of the invention, also can be used to treat various cardiac disorders, including ischemia of the heart (see *e.g.*, European Patent Application Publication No. EP 453 898) and heart insufficiency (weakness of the heart muscle)(see *e.g.*, PCT Publication No. WO 94/20139).

#### I. Others

30 The pharmaceutical formulation of the invention, also can be used to treat various other disorders in which TNF $\alpha$  activity is detrimental. Examples of other diseases and disorders in which TNF $\alpha$  activity has been implicated in the pathophysiology, and thus which can be treated using the formulation of the invention,  
35 include inflammatory bone disorders and bone resorption disease (see *e.g.*, Bertolini, D.R., *et al.* (1986) *Nature* 319:516-518; Konig, A., *et al.* (1988) *J. Bone Miner. Res.* 3:621-627; Lerner, U.H. and Ohlin, A. (1993) *J. Bone Miner. Res.* 8:147-155; and



- 27 -

Shankar, G. and Stern, P.H. (1993) *Bone* 14:871-876), hepatitis, including alcoholic hepatitis (see *e.g.*, McClain, C.J. and Cohen, D.A. (1989) *Hepatology* 9:349-351; Felver, M.E., *et al.* (1990) *Alcohol. Clin. Exp. Res.* 14:255-259; and Hansen, J., *et al.* (1994) *Hepatology* 20:461-474) and viral hepatitis (Sheron, N., *et al.* (1991) *J. Hepatol.* 12:241-245; and Hussain, M.J., *et al.* (1994) *J. Clin. Pathol.* 47:1112-1115), coagulation disturbances (see *e.g.*, van der Poll, T., *et al.* (1990) *N. Engl. J. Med.* 322:1622-1627; and van der Poll, T., *et al.* (1991) *Prog. Clin. Biol. Res.* 367:55-60), burns (see *e.g.*, Giroir, B.P., *et al.* (1994) *Am. J. Physiol.* 267:H118-124; and Liu, X.S., *et al.* (1994) *Burns* 20:40-44), reperfusion injury (see *e.g.*, Scales, W.E., *et al.* (1994) *Am. J. Physiol.* 267:G1122-1127; Serrick, C., *et al.* (1994) *Transplantation* 58:1158-1162; and Yao, Y.M., *et al.* (1995) *Resuscitation* 29:157-168), keloid formation (see *e.g.*, McCauley, R.L., *et al.* (1992) *J. Clin. Immunol.* 12:300-308), scar tissue formation; pyrexia; periodontal disease; obesity and radiation toxicity.

Other disorders in which TNF $\alpha$  activity is detrimental include, but are not limited to, adult Still's disease, Alzheimer's disease, ankylosing spondylitis, asthma, cancer and cachexia, atherosclerosis, chronic atherosclerosis, chronic fatigue syndrome, liver failure, chronic liver failure, obstructive pulmonary disease, chronic obstructive pulmonary disease, congestive heart failure, dermatopolymyositis, diabetic macrovasculopathy, endometriosis, familial periodic fevers, fibrosis, hemodialysis, Jarisch-Herxheimer reaction, juvenile RA, Kawasaki syndrome, myelo dysplastic syndrome, myocardial infarction, panciaticular vulgaris, periodontal disease, peripheral neuropathy, polyarticular, polymyositis, progressive renal failure, psoriasis, psoriatic arthritis, Reiter's syndrome, sarcoidosis, scleroderma, spondyloarthropathies, Still's disease, stroke, therapy associated syndrome, therapy induced inflammatory syndrome, inflammatory syndrome following IL-2 administration, thoracoabdominal aortic aneurysm repair (TAAA), Vasulo-Behcet's disease, Yellow Fever vaccination, type 1 diabetes mellitus, type 2 diabetes mellitus, neuropathic pain, sciatica, cerebral edema, edema in and/or around the spinal cord, vasculitide, Wegener's granulomatosis, temporal arteritis, polymyalgia rheumatica, Takayasu's arteritis, polyarteritis nodosa, microscopic polyangiitis, Churg-Strauss syndrome, Felty's syndrome, Sjogren's syndrome, mixed connective tissue disorder, relapsing polychondritis, pseudogout, loosening of prostheses,, autoimmune hepatitis, sclerosing cholangitis, acute pancreatitis, chronic pancreatitis, glomerulonephritides, post-streptococcal glomerulonephritis or IgA nephropathy, rheumatic heart disease, cardiomyopathy, orchitis, pyoderma gangrenosum, multiple myeloma, TNF receptor associated periodic syndrome [TRAPS], atherosclerosis, steroid dependent giant cell arteritismyositis, uveitis, and drug reactions.



The invention is further illustrated in the following examples, which should not be construed as further limiting.

5

### Examples

#### 10 EXAMPLE 1:      **Preparation of the Formulation**

The pharmaceutical formulation of the invention was made according to the following protocol.

15            Materials which were used in the formulation include: mannitol, citric acid monohydrate (citric acid), sodium citrate, disodium phosphate dihydrate (dibasic sodium phosphate dihydrate), sodium dihydrogen phosphate dihydrate (monobasic sodium phosphate dihydrate), sodium chloride, polysorbate 80, water for the injections, sodium hydroxide, which was used as a 1M solution to adjust the pH, and protein concentrate  
20 (e.g., antibody concentrate).

#### Preparation of 20L of buffer (equivalent to 20.180kg - density of the solution : 1.009 g/ml)

Ingredients were weighed out as follows: 240.0 g mannitol, 26.1 g citric acid  
25 monohydrate, 6.1 g sodium citrate, 30.6 g disodium phosphate dihydrate, 17.2 g sodium dihydrogen phosphate dihydrate, 123.3 g sodium chloride, 20.0 g polysorbate 80, and 19,715.7 to 19,716.1 g of water.

A sodium hydroxide solution was prepared by combining 40.0 g of sodium hydroxide with 1000.8 g of water for injections.

30            Next, a buffer was prepared by dissolving the following pre-weighed ingredients (described above) in about 90% of the water for injections: mannitol, citric acid monohydrate, sodium citrate, disodium phosphate dihydrate, sodium dihydrogen phosphate, sodium chloride, and polysorbate 80. It was determined that the sequence of the addition of the buffer constituents was not important and can, therefore, be chosen at  
35 will.

Following addition of all of the buffer constituents, the pH of the solution was adjusted with 1M sodium hydroxide which was prepared as described above. After the addition of the sodium hydroxide, the final weight of the water was added. The buffer

- 29 -

solution was then filtered through a sterilized filter (hydrophilic polyvinylidene difluoride, 0.22  $\mu\text{m}$  pore size) into a sterilized receptacle. The filtration medium used was filtration sterilized nitrogen.

5 Preparation of 40L of formulation (equivalent to 40.88 kg)

The filtered buffer solution was then added to the thawed and pooled antibody concentrate (the active ingredient of the pharmaceutical formulation), prepared as follows. The antibody (concentrate) was thawed in a water bath prior to the preparation of the pharmaceutical formulation. 34.207 g of antibody concentrate was used, which is  
10 equivalent to 2.0 kg of protein with 60 mg protein/mL protein concentrate. The density of the concentrate was 1.0262 g/mL. Any protein concentrate ranging from 25.655 to 37.316, which is equivalent to a protein concentration in the protein concentrate of 55 to 80 mg/mL, can be used. The buffer was added while stirring, until the final weight of the bulk solution was reached.

15 The formulation, with all of its ingredients included, was then sterilized by filtration as described above, except the formulation was filtered through two sterile 0.22  $\mu\text{m}$  membrane filters. Following sterilization, the formulation was packaged for use in either a vial or a pre-filled syringe.

The skilled artisan will also appreciate that the weight quantities and/or weight-to-volume ratios recited herein, can be converted to moles and/or molarities using the  
20 art-recognized molecular weights of the recited ingredients. Weight quantities exemplified herein (*e.g.*, g or kg) are for the volumes (*e.g.*, of buffer or pharmaceutical formulation) recited. The skilled artisan will appreciate that the weight quantities can be proportionally adjusted when different formulation volumes are desired. For example,  
25 32L, 20L, 10L, 5L, or 1L formulations would include 80%, 50%, 25%, 12.5%, or 2.5%, respectively, of the exemplified weight quantities.

30 **EXAMPLE 2: Freeze/Thaw Studies**

After the formulation buffer for the D2E7 antibody was selected the drug substance was formulated in the same matrix as the finished product.

Freeze thaw behavior of the D2E7 antibody drug substance at a protein concentration of 63 mg/mL was evaluated by cycling drug substance 3 times from the  
35 frozen state to the liquid state. Table N shows the results of an experiment evaluating the effect of three fast and slow freeze-thaw cycles in the presence and absence of 0.1 % polysorbate 80 starting from  $-80^{\circ}\text{C}$  or  $-30^{\circ}\text{C}$ , respectively.



- 30 -

Table 2 shows that the D2E7 antibody drug substance can be thawed/frozen at least 3 times without any detrimental effect on either chemical (cation exchange HPLC, size exclusion HPLC, colour, pH), physicochemical properties (subvisible particles, clarity) or biological activity (in vitro TNF neutralization assay). Also table 2 shows that the inclusion of polysorbate 80 improved the physicochemical properties of the D2E7 antibody drug substance as evidenced by the lower number of subvisible particles regardless whether a slow or fast freeze/thaw cycle was being used (see shaded areas in table 2).

10 Table 2: Effect of freeze thaw on the D2E7 antibody drug substance with/without polysorbate 80

Test criteria	Poly-sorbate (0.1%) <sup>1)</sup>	No freeze/thaw	Slow thaw -30°C in refrigerator	Fast thaw -30°C in water bath	Slow thaw -80° in refrigerator	Fast thaw -80°C in water bath
Clarity	-	25.0	22.5	25.3	25.8	25.6
	+	27.8	28.1	28.2	28.0	28.1
Colour	-	≤ B9	≤ B9	≤ B9	≤ B9	≤ B9
	+	≤ B9	≤ B9	≤ B9	≤ B9	≤ B9
pH	-	5.01	5.02	5.02	5.02	5.02
	+	5.02	5.02	5.02	5.02	5.02
Subvisible particles	-	42	600	303	1891	303
		2	4	5	8	0
	+	0	5	1	0	8
		0	0	0	0	1
Size exclusion HPLC	-	99.8	99.8	99.8	99.8	99.8
	+	99.8	99.8	99.8	99.8	99.8
Cation exchange HPLC	-	87.1	87.0	87.2	86.9	86.9
	+	86.8	87.0	87.1	87.3	86.8
In vitro TNF neutralization test	-	118.0	123.8	118.0	103.3	120.5
	+	111.8	96.2	100.9	96.7	95.8

1) + = formulation with 0.1 % polysorbate 80; - = formulation without 0.1 % polysorbate 80

- 31 -

**EXAMPLE 3: Microbial Studies**

Tests were performed to determine if the formulation can support microbial growth. The results from these experiments showed that the formulation does not support microbial growth if stored at 20 to 25 °C for 14 days. This result was determined by directly inoculating the sterile formulation with microorganisms (*e.g.*, *Staphylococcus aureus*, ATCC-No.: 6538P, *Candida albicans*, ATCC-No.: 10231, *Aspergillus niger*, ATCCC-No.: 16404, *Pseudomonas aeruginosa*, ATCC-No.: 9027, an environmental isolate) at low level (NMT 100 cfu/mL). Inoculated formulations were then examined for overall microbial growth, *e.g.*, for changes in turbidity. A lack of turbidity was an indication of no overall growth, and was detected in the inoculated containers after 14 days. Further, no organisms could be reisolated from these containers. Thus it was concluded that the formulation does not support microbial growth under these conditions.

15

20

**Equivalents**

The scope of the claims should not be limited by the preferred embodiments set forth in the examples, but should be given the broadest interpretation consistent with the description as a whole.

25



page 1

## SEQUENCE LISTING

&lt;110&gt; Abbott Biotechnology Ltd.

&lt;120&gt; PHARMACEUTICAL ANTI-TNF-ALPHA ANTIBODY FORMULATION

&lt;130&gt; 35352-2019

&lt;140&gt; CA 2,494,756

&lt;141&gt; 2003-08-16

&lt;150&gt; 10/222,140

&lt;151&gt; 2002-08-16

&lt;160&gt; 34

&lt;170&gt; PatentIn version 3.5

&lt;210&gt; 1

&lt;211&gt; 107

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Synthetic polypeptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 1

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5				10						15	

Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Arg	Asn	Tyr
			20					25					30		

Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
		35				40						45			

Tyr	Ala	Ala	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				

Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
65					70					75					80

Glu	Asp	Val	Ala	Thr	Tyr	Tyr	Cys	Gln	Arg	Tyr	Asn	Arg	Ala	Pro	Tyr
				85					90					95	

Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

page 2

100

105

<210> 2  
 <211> 121  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic polypeptide

<220>  
 <223> Mutated human antibody

<400> 2  
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr  
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val  
 50 55 60

Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr  
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly  
 100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> 3  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic peptide



page 3

<220>  
<223> Mutated human antibody

<220>  
<221> MOD\_RES  
<222> (9)..(9)  
<223> Thr or Ala

<400> 3  
Gln Arg Tyr Asn Arg Ala Pro Tyr Xaa  
1 5

<210> 4  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<223> Mutated human antibody

<220>  
<221> MOD\_RES  
<222> (12)..(12)  
<223> Tyr or Asn

<400> 4  
Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Xaa  
1 5 10

<210> 5  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<223> Mutated human antibody

<400> 5  
Ala Ala Ser Thr Leu Gln Ser  
1 5

<210> 6  
<211> 17  
<212> PRT

page 4

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Synthetic peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 6

Ala	Ile	Thr	Trp	Asn	Ser	Gly	His	Ile	Asp	Tyr	Ala	Asp	Ser	Val	Glu
1				5					10					15	

Gly

&lt;210&gt; 7

&lt;211&gt; 11

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Synthetic peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 7

Arg	Ala	Ser	Gln	Gly	Ile	Arg	Asn	Tyr	Leu	Ala
1				5					10	

&lt;210&gt; 8

&lt;211&gt; 5

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Synthetic peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 8

Asp	Tyr	Ala	Met	His
1				5

&lt;210&gt; 9

&lt;400&gt; 9



000

<210> 10

<400> 10

000

<210> 11

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 11

Gln Lys Tyr Asn Ser Ala Pro Tyr Ala  
1 5

<210> 12

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 12

Gln Lys Tyr Asn Arg Ala Pro Tyr Ala  
1 5

<210> 13

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

page 6

&lt;400&gt; 13

Gln Lys Tyr Gln Arg Ala Pro Tyr Thr  
1 5

&lt;210&gt; 14

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 14

Gln Lys Tyr Ser Ser Ala Pro Tyr Thr  
1 5

&lt;210&gt; 15

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 15

Gln Lys Tyr Asn Ser Ala Pro Tyr Thr  
1 5

&lt;210&gt; 16

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
peptide

&lt;220&gt;

&lt;223&gt; Mutated human antibody

&lt;400&gt; 16

Gln Lys Tyr Asn Arg Ala Pro Tyr Thr  
1 5



page 7

<210> 17  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<223> Mutated human antibody

<400> 17  
Gln Lys Tyr Asn Ser Ala Pro Tyr Tyr  
1 5

<210> 18  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<223> Mutated human antibody

<400> 18  
Gln Lys Tyr Asn Ser Ala Pro Tyr Asn  
1 5

<210> 19  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<223> Mutated human antibody

<400> 19  
Gln Lys Tyr Thr Ser Ala Pro Tyr Thr  
1 5

<210> 20  
<211> 9  
<212> PRT

page 8

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 20

Gln Lys Tyr Asn Arg Ala Pro Tyr Asn  
1 5

<210> 21

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 21

Gln Lys Tyr Asn Ser Ala Ala Tyr Ser  
1 5

<210> 22

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 22

Gln Gln Tyr Asn Ser Ala Pro Asp Thr  
1 5

<210> 23

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic



peptide

<220>

<223> Mutated human antibody

<400> 23

Gln Lys Tyr Asn Ser Asp Pro Tyr Thr  
1 5

<210> 24

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<220>

<223> Mutated human antibody

<400> 24

Gln Lys Tyr Ile Ser Ala Pro Tyr Thr  
1 5

<210> 25

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<220>

<223> Mutated human antibody

<400> 25

Gln Lys Tyr Asn Arg Pro Pro Tyr Thr  
1 5

<210> 26

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<220>

<223> Mutated human antibody

<400> 26

Gln Arg Tyr Asn Arg Ala Pro Tyr Ala  
1 5

<210> 27

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 27

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Asn  
1 5 10

<210> 28

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 28

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Lys  
1 5 10

<210> 29

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<223> Mutated human antibody

<400> 29

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Tyr









## Claims:

1. A liquid aqueous pharmaceutical formulation comprising  
a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody at a concentration from 20 to 150 mg/ml,  
a sugar alcohol at a concentration from 5 to 20 mg/ml,  
a polysorbate at a concentration from 0.1 to 10 mg/ml, and  
a buffer system comprising citrate and phosphate,  
wherein said formulation has a pH from 4 to 8, and  
wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
2. A liquid aqueous pharmaceutical formulation comprising
  - (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody,
  - (b) from 5 to 20 mg/ml of mannitol,
  - (c) from 0.1 to 10 mg/ml of polysorbate 80, and
  - (d) a buffer system comprising citrate and phosphate, with a pH from 4 to 8,wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
3. The liquid aqueous pharmaceutical formulation of claim 1, wherein the sugar alcohol is mannitol.
4. The liquid aqueous pharmaceutical formulation of claim 1, wherein the polysorbate is polysorbate 80.
5. The liquid aqueous pharmaceutical formulation of claim 2, wherein the antibody is antibody D2E7.

- 33 -

6. The liquid aqueous pharmaceutical formulation of claim 1, 3 or 4, wherein the antibody is antibody D2E7.
7. The liquid aqueous pharmaceutical formulation of any one of claims 1-6, comprising from 30 to 120 mg/ml of the antibody.
8. The liquid aqueous pharmaceutical formulation of any one of claims 1-7, comprising from 35 to 115 mg/ml of the antibody.
9. The liquid aqueous pharmaceutical formulation of claim 8, comprising from 40 to 110 mg/ml of the antibody.
10. The liquid aqueous pharmaceutical formulation of claim 9, comprising from 45 to 105 mg/ml of the antibody.
11. The liquid aqueous pharmaceutical formulation of any one of claims 1-10, wherein the pH is from 4.5 to 6.0.
12. The liquid aqueous pharmaceutical formulation of claim 11, wherein the pH is from 4.8 to 5.5.
13. The liquid aqueous pharmaceutical formulation of claim 2 or 3, comprising from 10 to 14 mg/ml of mannitol.
14. The liquid aqueous pharmaceutical formulation of claim 2 or 4, comprising from 0.5 to 5 mg/ml of polysorbate 80.
15. The liquid aqueous pharmaceutical formulation of claim 2, which contains
  - (a) from 40 to 100 mg/ml of the antibody,
  - (b) from 7.5 to 15 mg/ml of mannitol, and
  - (c) from 0.5 to 5 mg/ml of polysorbate 80.



- 34 -

16. The liquid aqueous pharmaceutical formulation of any one of claims 1-4 and 13-15, wherein the antibody comprises an IgG<sub>1</sub> heavy chain constant region.
17. The liquid aqueous pharmaceutical formulation of any one of claims 1-4 and 13-15, wherein the antibody comprises an IgG<sub>4</sub> heavy chain constant region.
18. A liquid aqueous pharmaceutical formulation having a pH from 4 to 8 and comprising
  - (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG<sub>1</sub> antibody comprising a D2E7 light chain variable region and a D2E7 heavy chain variable region;
  - (b) from 10 to 14 mg/ml of mannitol,
  - (c) from 0.1 to 5 mg/ml of polysorbate 80,
  - (d) from 1 to 1.5 mg/ml of citric acid monohydrate,
  - (e) from 0.25 to 0.5 mg/ml of sodium citrate,
  - (f) from 1.25 to 1.75 mg/ml of disodium phosphate dihydrate,
  - (g) from 0.7 to 1.1 mg/ml of sodium dihydrogen phosphate dihydrate, and
  - (h) from 6.0 to 6.4 mg/ml sodium chloride.
19. A liquid aqueous pharmaceutical formulation having a pH from 4 to 8 and comprising
  - (a) from 20 to 150 mg/ml of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG<sub>4</sub> antibody comprising a D2E7 light chain variable region and a D2E7 heavy chain variable region;
  - (b) from 10 to 14 mg/ml of mannitol,
  - (c) from 0.1 to 5 mg/ml of polysorbate 80,
  - (d) from 1 to 1.5 mg/ml of citric acid monohydrate,
  - (e) from 0.25 to 0.5 mg/ml of sodium citrate,
  - (f) from 1.25 to 1.75 mg/ml of disodium phosphate dihydrate,
  - (g) from 0.7 to 1.1 mg/ml of sodium dihydrogen phosphate dihydrate, and

- 35 -

(h) from 6.0 to 6.4 mg/ml sodium chloride.

20. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an autoimmune disease.
21. The use of claim 20, wherein the autoimmune disease is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis, autoimmune hepatitis, or nephrotic syndrome.
22. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an infectious disease.
23. The use of claim 22, wherein the infectious disease is meningitis, malaria, acquired immune deficiency syndrome, or cytomegalovirus infection secondary to transplantation.
24. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is a pulmonary disorder.
25. The use of claim 24, wherein the pulmonary disorder is adult respiratory distress syndrome, shock lung, a chronic pulmonary inflammatory disease, an obstructive pulmonary disease, a chronic obstructive pulmonary disease, pulmonary sarcoidosis, pulmonary fibrosis, or silicosis.
26. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an inflammatory bowel disease.



- 36 -

27. The use of claim 26, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.
28. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is ankylosing spondylitis.
29. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating juvenile rheumatoid arthritis in a human subject.
30. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating psoriasis in a human subject.
31. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating psoriatic arthritis in a human subject.
32. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating sarcoidosis in a human subject.
33. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating scleroderma in a human subject.
34. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a spondyloarthropathy in a human subject.
35. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating Still's disease in a human subject.
36. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating Sjogren's syndrome in a human subject.

37. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating atherosclerosis in a human subject.
38. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating uveitis in a human subject.
39. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is sepsis, a graft versus host disease, a malignancy, cachexia, an inflammatory bone disorder, a bone resorption disease, hepatitis, a coagulation disturbance, burn, a reperfusion injury, keloid formation, scar tissue formation, pyrexia, a periodontal disease, obesity, radiation, Alzheimer's disease, asthma, chronic fatigue syndrome, liver failure, chronic liver failure, dermatopolymyositis, diabetic macrovasculopathy, endometriosis, familial periodic fevers, fibrosis, hemodialysis, Jarisch-Herxheimer reaction, Kawasaki syndrome, myelodysplastic syndrome, panciaticular vulgaris, peripheral neuropathy, polyarticular, polymyositis, progressive renal failure, Reiter's syndrome, stroke, therapy associated syndrome, therapy induced inflammatory syndrome, inflammatory syndrome following IL-2 administration, thoracoabdominal aortic aneurysm repair (TAAA), Vasulo-Behcet's disease, Yellow Fever vaccination, type 1 diabetes mellitus, type 2 diabetes mellitus, neuropathic pain, sciatica, cerebral edema, edema in or around the spinal cord, vasculitide, Wegener's granulomatosis, temporal arteritis, polymyalgia rheumatica, Takayasu's arteritis, polyarteritis nodosa, microscopic polyangiitis, Churg-Strauss syndrome, Felty's syndrome, mixed connective tissue disorder, relapsing polychondritis, pseudogout, loosening of prostheses, sclerosing cholangitis, acute pancreatitis, chronic pancreatitis, glomerulonephritides, post-streptococcal glomerulonephritis or IgA nephropathy, rheumatic heart disease, orchitis, pyoderma gangrenosum, multiple myeloma, TNF receptor associated periodic syndrome [TRAPS], atherosclerosis, steroid dependent giant cell arteritismyositis, or a cardiac disorder.



- 38 -

40. Use of the formulation of any one of claims 1-19 for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an autoimmune disease.

41. The use of claim 40, wherein the autoimmune disease is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis, autoimmune hepatitis, or nephrotic syndrome.

42. Use of the formulation of any one of claims 1-19 for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an infectious disease.

43. The use of claim 42, wherein the infectious disease is meningitis, malaria, acquired immune deficiency syndrome, or cytomegalovirus infection secondary to transplantation.

44. Use of the formulation of any one of claims 1-19 for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is a pulmonary disorder.

45. The use of claim 44, wherein the pulmonary disorder is respiratory distress syndrome, shock lung, a chronic pulmonary inflammatory disease, an obstructive pulmonary disease, a chronic obstructive pulmonary disease, pulmonary sarcoidosis, pulmonary fibrosis, or silicosis.

46. Use of the formulation of any one of claims 1-19 for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is an inflammatory bowel disease.

- 39 -

47. The use of claim 46, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.
48. Use of the formulation of any one of claims 1-19 for treating ankylosing spondylitis in a human subject.
49. Use of the formulation of any one of claims 1-19 for treating juvenile rheumatoid arthritis (JRA) in a human subject.
50. Use of the formulation of any one of claims 1-19 for treating psoriasis in a human subject.
51. Use of the formulation of any one of claims 1-19 for treating psoriatic arthritis in a human subject.
52. Use of the formulation of any one of claims 1-19 for treating sarcoidosis in a human subject.
53. Use of the formulation of any one of claims 1-19 for treating scleroderma in a human subject.
54. Use of the formulation of any one of claims 1-19 for treating a spondyloarthropathy in a human subject.
55. Use of the formulation of any one of claims 1-19 for treating Still's disease in a human subject.
56. Use of the formulation of any one of claims 1-19 for treating Sjogren's syndrome in a human subject.



- 40 -

57. Use of the formulation of any one of claims 1-19 for treating atherosclerosis in a human subject.

58. Use of the formulation of any one of claims 1-19 for treating uveitis in a human subject.

59. Use of the formulation of any one of claims 1-19 for treating a disorder in which TNF $\alpha$  activity is detrimental in a human subject, wherein the disorder is sepsis, a graft versus host disease, a malignancy, cachexia, an inflammatory bone disorder, a bone resorption disease, hepatitis, a coagulation disturbance, burn, a reperfusion injury, keloid formation, scar tissue formation, pyrexia, a periodontal disease, obesity, radiation, Alzheimer's disease, asthma, chronic fatigue syndrome, liver failure, chronic liver failure, dermatopolymyositis, diabetic macrovasculopathy, endometriosis, familial periodic fevers, fibrosis, hemodialysis, Jarisch-Herxheimer reaction, Kawasaki syndrome, myelodysplastic syndrome, paronychia, peripheral neuropathy, polyarticular, polymyositis, progressive renal failure, Reiter's syndrome, stroke, therapy associated syndrome, therapy induced inflammatory syndrome, inflammatory syndrome following IL-2 administration, thoracoabdominal aortic aneurysm repair (TAAA), Vasculo-Behcet's disease, Yellow Fever vaccination, type 1 diabetes mellitus, type 2 diabetes mellitus, neuropathic pain, sciatica, cerebral edema, edema in or around the spinal cord, vasculitide, Wegener's granulomatosis, temporal arteritis, polymyalgia rheumatica, Takayasu's arteritis, polyarteritis nodosa, microscopic polyangiitis, Churg-Strauss syndrome, Felty's syndrome, mixed connective tissue disorder, relapsing polychondritis, pseudogout, loosening of prostheses, sclerosing cholangitis, acute pancreatitis, chronic pancreatitis, glomerulonephritides, post-streptococcal glomerulonephritis or IgA nephropathy, rheumatic heart disease, orchitis, pyoderma gangrenosum, multiple myeloma, TNF receptor associated periodic syndrome [TRAPS], atherosclerosis, steroid dependent giant cell arteritis/myositis, or a cardiac disorder.

60. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat psoriasis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

61. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat psoriatic arthritis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

62. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat ankylosing spondylitis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

63. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat juvenile rheumatoid arthritis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.



- 42 -

64. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat rheumatoid arthritis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

65. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat Crohn's disease, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

66. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat ulcerative colitis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

67. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat uveitis, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody

- 43 -

comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

68. Use of a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody that specifically binds TNF $\alpha$  to treat Vasulo-Behcet's disease, wherein the antibody is at a concentration from 20 to 150 mg/ml in a solution comprising a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate, at a pH from 4 to 8, and wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

69. The use of any one of claims 60-68, wherein the sugar alcohol is mannitol.

70. The use of claim 69, wherein the mannitol is at a concentration from 10 to 14 mg/ml.

71. The use of any one of claims 60-70, wherein the polysorbate is polysorbate 80.

72. The use of claim 71, wherein the polysorbate 80 is at a concentration from 0.5 to 5 mg/ml.

73. The use of any one of claims 60-72, wherein the antibody is at a concentration from 30 to 120 mg/ml.

74. The use of claim 73, wherein the antibody is at a concentration from 35 to 115 mg/ml.

75. The use of claim 73, wherein the antibody is at a concentration from 40 to 110 mg/ml.



- 44 -

76. The use of claim 73, wherein the antibody is at a concentration from 45 to 105 mg/ml.
77. The use of any one of claims 60-76, wherein the pH is from 4.5 to 6.0.
78. The use of claim 77, wherein the pH is from 4.8 to 5.5.
79. The use of any one of claims 60-78, wherein the antibody is antibody D2E7, the sugar alcohol is mannitol at a concentration from 10 to 14 mg/ml, and the polysorbate is polysorbate 80 at a concentration from 0.5 to 5 mg/ml.
80. The use of any one of claims 60-78, wherein the antibody comprises an IgG<sub>1</sub> or IgG<sub>4</sub> heavy chain constant region.
81. The use of any one of claims 60-78, wherein the antibody is antibody D2E7.
82. The use of any one of claims 79-81, wherein the use is subcutaneous use.
83. A liquid aqueous pharmaceutical formulation comprising a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG<sub>1</sub> antibody at a concentration from 20 to 150 mg/ml, a sugar alcohol at a concentration from 5 to 20 mg/ml, a polysorbate at a concentration from 0.1 to 10 mg/ml, and a buffer system comprising citrate and phosphate,  
wherein said formulation has a pH from 4 to 8, and  
wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.
84. A liquid aqueous pharmaceutical formulation comprising a human anti-Tumor Necrosis Factor alpha (TNF $\alpha$ ) IgG antibody at a concentration from 40 to 110 mg/ml, citric acid monohydrate, dibasic sodium phosphate dihydrate, mannitol, monobasic sodium phosphate dihydrate, polysorbate 80, sodium citrate, and sodium chloride,  
wherein said formulation has a pH from 4 to 8, and

- 45 -

wherein the antibody comprises a light chain variable region and a heavy chain variable region of antibody D2E7.

85. The liquid aqueous pharmaceutical formulation of claim 83 or 84, wherein the pH is from 4.5 to 6.0.

86. The liquid aqueous pharmaceutical formulation of claim 83 or 84, wherein the pH is from 4.8 to 5.5.

87. The liquid aqueous pharmaceutical formulation of any one of claims 83 to 86, wherein the concentration of the antibody is 50 mg/ml.

88. The liquid aqueous pharmaceutical formulation of claim 83, wherein the antibody is antibody D2E7, the sugar alcohol is mannitol at a concentration from 10 to 14 mg/ml, and the polysorbate is polysorbate 80 at a concentration from 0.5 to 5 mg/ml.

89. The liquid aqueous pharmaceutical formulation of claim 84, wherein the antibody is antibody D2E7, mannitol is at a concentration from 10 to 14 mg/ml, and polysorbate 80 is at a concentration from 0.5 to 5 mg/ml.

90. The liquid aqueous pharmaceutical formulation of claim 83, wherein the antibody is antibody D2E7 at a concentration of about 50 mg/ml, the sugar alcohol is mannitol at a concentration of about 12 mg/ml, the polysorbate is polysorbate 80 at a concentration of about 1 mg/ml, and the pH of the buffer system is from about 4.8 to about 5.5.

91. The liquid aqueous pharmaceutical formulation of claim 84, wherein the antibody is antibody D2E7 at a concentration of about 50 mg/ml, mannitol is at a concentration of about 12 mg/ml, polysorbate 80 is at a concentration of about 1 mg/ml, and the pH of the buffer system is from about 5.0 to about 5.2.



- 46 -

92. Use of the formulation of any one of claims 1-19 in the preparation of a medicament for treating rheumatoid arthritis in a human subject.
93. Use of the formulation of any one of claims 1-19 for treating rheumatoid arthritis in a human subject.