



(12)

Oversættelse af europæisk patent

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **A 01 N 63/00 (2006.01)** **A 61 K 35/14 (2006.01)** **A 61 K 39/00 (2006.01)**
A 61 K 45/06 (2006.01) **C 12 N 5/0783 (2010.01)** **C 12 N 15/85 (2006.01)**

(45) Oversættelsen bekendtgjort den: 2015-01-05

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: 2014-11-12

(86) Europæisk ansøgning nr.: **12179377.2**

(86) Europæisk indleveringsdag: **2008-03-31**

(87) Den europæiske ansøgnings publiceringsdag: **2012-12-26**

(30) Prioritet: **2007-03-30 US 921144 P**

(62) Stamansøgningsnr: **08742463.6**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**

(73) Patenthaver: **MEMORIAL SLOAN-KETTERING CANCER CENTER, 1275 York Avenue, New York, NY 10021, USA**

(72) Opfinder: **STEPHAN, Matthias, 111 Queensberry St., Apt. 14, Boston, MA 02215, USA**
SADELAIN, Michel, 444 Central Park West, Apt. 8h, New York, NY 10025, USA

(74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**

(54) Benævnelse: **KONSTITUTIV EKSPRESSION AF CO-STIMULATORISKE LIGANDER PÅ ADOPTIVT OVERFØRTE T-LYMFOCYTER**

(56) Fremdragne publikationer:
KOWOLIK CLAUDIA M ET AL: "CD28-costimulation provided through a CD19-specific chimeric immunoreceptor enhances in vivo persistence and anti-tumor efficacy of adoptively transferred T cells.", BLOOD, vol. 106, no. 11, November 2005 (2005-11), page 372A, XP009130374, & 47TH ANNUAL MEETING OF THE AMERICAN-SOCIETY-OF-HEMATOLOGY; ATLANTA, GA, USA; DECEMBER 10 -13, 2005 ISSN: 0006-4971
GADE TERENCE P F ET AL: "Targeted elimination of prostate cancer by genetically directed human T lymphocytes", CANCER RESEARCH, vol. 65, no. 19, October 2005 (2005-10), page 9080, XP002571428, ISSN: 0008-5472
MELERO I ET AL: "AMPLIFICATION OF TUMOR IMMUNITY BY GENE TRANSFER OF THE CO-STIMULATORY 4-1BB LIGAND: SYNERGY WITH THE CD28 CO-STIMULATORY PATHWAY", EUROPEAN JOURNAL OF IMMUNOLOGY, WILEY - V C H VERLAG GMBH & CO. KGAA, DE, vol. 28, no. 3, 1 January 1998 (1998-01-01), pages 1116-1121, XP000914703, ISSN: 0014-2980
GROSENBACH D W ET AL: "A recombinant vector expressing transgenes for four T-cell costimulatory molecules (OX40L, B7-1, ICAM-1, LFA-3) induces sustained CD4+ and CD8+ T-cell activation, protection from apoptosis, and enhanced cytokine production", CELLULAR IMMUNOLOGY, ACADEMIC PRESS, SAN DIEGO, CA, US, vol. 222, 1 March 2003 (2003-03-01), pages 45-57, XP002995005, ISSN: 0008-8749
BRENTJENS RENIER J ET AL: "Eradication of systemic B-cell tumors by genetically targeted human T lymphocytes co-stimulated by CD80 and interleukin-15", NATURE MEDICINE, NATURE PUBLISHING GROUP, 2003-03-01, pages 45-57, XP002995005, ISSN: 1078-8961

NEW YORK, NY, US, vol. 9, no. 3, 1 March 2003 (2003-03-01), pages 279-286, XP002389127, ISSN: 1078-8956
STEPHAN MATTHIAS T ET AL: "T cell-encoded CD80 and 4-1BBL induce auto- and transcostimulation, resulting in potent tumor rejection", NATURE MEDICINE, vol. 13, no. 12, December 2007 (2007-12), pages 1440-1449, XP002571429, ISSN: 1078-8956
MAUS M V ET AL: "Ex vivo expansion of polyclonal and antigen-specific cytotoxic T lymphocytes by artificial APCs expressing ligands for the T-cell receptor, CD28 and 4-1 BB", NATURE BIOTECHNOLOGY, NATURE PUBLISHING GROUP, NEW YORK, NY, US, vol. 20, no. 2, 1 February 2002 (2002-02-01), pages 143-148, XP002225278, ISSN: 1087-0156
SADELAIN M ET AL: "TARGETING TUMOURS WITH GENETICALLY ENHANCED T LYMPHOCYTES", NATURE REVIEWS. CANCER, NATUR PUBLISHING GROUP, LONDON, GB, vol. 3, no. 1, 1 January 2003 (2003-01-01), pages 35-45, XP009026020, ISSN: 1474-175X
ZHONG ET AL: "Integrated CD28 and 4-1BB Signals Strongly Potentiate CD8+ T Cell Mediated eradication of Metastatic Prostate Cancer", MOLECULAR THERAPY, ACADEMIC PRESS, SAN DIEGO, CA, US, vol. 13, 1 January 2006 (2006-01-01), page S103, XP005675376, ISSN: 1525-0016
C. M. KOWOLIK ET AL: "CD28 Costimulation Provided through a CD19-Specific Chimeric Antigen Receptor Enhances In vivo Persistence and Antitumor Efficacy of Adoptively Transferred T Cells", CANCER RESEARCH, vol. 66, no. 22, 15 November 2006 (2006-11-15), pages 10995-11004, XP055044266, ISSN: 0008-5472, DOI: 10.1158/0008-5472.CAN-06-0160
XIAO-SONG ZHONG ET AL: "Chimeric Antigen Receptors Combining 4-1BB and CD28 Signaling Domains Augment PI3kinase/AKT/Bcl-XL Activation and CD8+ T Cell-mediated Tumor Eradication", MOLECULAR THERAPY, vol. 18, no. 2, 1 February 2010 (2010-02-01), pages 413-420, XP055023623, ISSN: 1525-0016, DOI: 10.1038/mt.2009.210

Description**BACKGROUND OF THE INVENTION**

[0001] Prostate cancer is the most frequent cancer in males in the United States and the cause of nearly 31,000 deaths per year. When diagnosed early, cancer can be effectively treated by surgery or radiation. Postsurgical residual disease requires radiation and/or hormonal therapy, which may prevent tumor progression and metastasis. At present, there is no curative treatment for hormone refractory, metastatic prostate cancer. Immunotherapy is a targeted therapy that in principle provides for the treatment of such cancers. Obstacles remain to induce tumor immunity, which requires the expansion of cytotoxic T lymphocytes to numbers sufficient to mediate tumor rejection. Among the mechanisms limiting efficient T cell priming and tumor rejection is the inherent absence of costimulatory ligands on many malignancies.

SUMMARY OF THE INVENTION

[0002] The disclosure generally provides immunoresponsive cells, including T cells and Natural Killer (NK) cells, expressing at least one of an antigen-recognizing receptor and a costimulatory ligand and methods of use therefore for the treatment of neoplasia, infectious disease, and other pathologies.

[0003] In one aspect, the disclosure generally provides an immunoresponsive cell comprising a receptor that binds an antigen and an exogenous co-stimulatory ligand.

[0004] In another aspect, the disclosure provides a virus specific T cell expressing a vector (e.g., an expression vector) encoding a polypeptide selected from any one or more of CD80, 4-1BBL, OX40L, CD70 and CD30L. In one embodiment, the virus specific T cell recognizes a virus selected from any one or more of Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus antigens.

[0005] In yet another aspect, the disclosure provides a tumor antigen-specific T cell expressing a vector encoding a polypeptide selected from the group consisting of CD80, 4-1BBL, OX40L, CD70 and CD30L. In one embodiment, the cell expresses CD80 and 4-1BBL. In another embodiment, the vector is a retroviral vector (e.g., gamma-retroviral or lentiviral); it may also be non-viral.

[0006] In yet another aspect, the disclosure provides a method of modulating an immune response in a subject, the method comprising administering an effective amount of an immunoresponsive cell of any previous aspect. In one embodiment, the method increases or reduces an immune response. In another embodiment, the method increases self-tolerance or increases tolerance to an organ transplant.

[0007] In yet another aspect, the invention provides a T cell comprising a receptor that

binds a tumor antigen, and exogenous co-stimulatory ligands 4-1BBL and CD80 for use in treating or preventing a neoplasia in a subject. In one embodiment, the neoplasia is selected from the any one or more of prostate cancer, colon cancer, breast cancer, and glioblastoma. In another embodiment, the tumor antigen is prostate-specific membrane antigen, CD19, NY-ESO-1, WT-1, hTERT, or mesothelin.

[0008] In another aspect, the disclosure provides a method of enforcing tolerance in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds an antigen and a vector encoding a co-stimulatory ligand. In one embodiment, the method prevents or reduces an autoimmune disease or a disease associated with allogeneic transplantation.

[0009] In yet another aspect, the disclosure provides a method of treating or preventing a pathogen infection in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds a viral antigen and a vector encoding a co-stimulatory ligand. In one embodiment, the pathogen is a virus, bacteria, fungus, protozoa or parasite. In another embodiment, the virus is selected from any one or more of Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus. In yet another embodiment, the cell is a T cell, a Natural Killer (NK) cell, or a cytotoxic T lymphocyte (CTL).

[0010] In still another aspect, the disclosure provides method for producing an antigen-specific immunoresponsive cell, the method comprising introducing into the immunoresponsive cell a nucleic acid sequence that encodes a chimeric antigen receptor, wherein the chimeric antigen receptor comprises an antigen-binding domain coupled to an intracellular signaling domain that activates an immunoresponsive cell. In one embodiment, the immunoresponsive cell is a T cell, CTL, or NK cell. In another embodiment, the antigen-binding domain is a tumor antigen-binding domain. In yet another embodiment, the tumor antigen is prostate specific membrane antigen (PSMA). In yet another embodiment, the intracellular signaling domain activates a T cell, CTL cell, or NK cell. In yet another embodiment, the intracellular signaling domain is the ξ -chain signaling domain.

[0011] In another aspect, the disclosure provides a method of treating a neoplasia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a T cell comprising a tumor antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from any one or more of a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating cancer in the subject.

[0012] In another aspect, the disclosure provides a method of treating a neoplasia in a subject, the method comprising administering to the subject a therapeutically effective amount of a Natural Killer (NK) cell comprising a tumor antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from any one or more of a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating cancer in the subject. In one embodiment, the TNF ligand is selected from any one

or more of 4-1BBL, OX40L, CD70, CD30L, and LIGHT. In another embodiment, the Ig superfamily ligand is selected from CD80 and CD86. In yet another embodiment, the cell expresses at least two co-stimulatory ligands, where one is a TNF ligand (e.g., 4-1BBL) and the other is an Ig superfamily ligand (e.g., CD80).

[0013] In another aspect, the disclosure provides a method of treating an infectious disease in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a T cell comprising a receptor specific for a viral antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from any one or more of a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating the infectious disease in the subject.

[0014] In yet another aspect, the disclosure provides method of treating an infectious disease in a subject, comprising administering to the subject a therapeutically effective amount of a Natural Killer (NK) cell comprising a receptor specific for a viral antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from any one or more of a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating the infectious disease in the subject. In one embodiment, the subject is an immunocompromised subject. In another embodiment, the TNF ligand is selected from any one or more of 4-1BBL, OX40L, CD70, LIGHT and CD30L. In yet another embodiment, the Ig superfamily ligand is selected from any one or more of CD80 and CD86. In yet another embodiment, the at least two co-stimulatory ligands are a TNF ligand and an Ig superfamily ligand. In another embodiment, the TNF ligand is 4-1BBL and the Ig superfamily ligand is CD80. In another embodiment, the antigen recognition complex is constitutively expressed on the surface of the cell. In another embodiment, the viral antigen is an antigen specific for Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), or influenza virus. In another embodiment, the at least two co-stimulatory ligands are constitutively expressed on the surface of the cell.

[0015] In another aspect, the disclosure provides a pharmaceutical composition comprising an effective amount of an immunoresponsive cell of any previous aspect in a pharmaceutically acceptable excipient.

[0016] In another aspect, the invention provides a pharmaceutical composition for use in the treatment of a neoplasia comprising an effective amount of a tumor antigen-specific T cell comprising a receptor that binds a tumor antigen, and co-stimulatory ligands 4-1BBL and CD80 in a pharmaceutically acceptable excipient.

[0017] In another aspect, the disclosure provides a pharmaceutical composition for the treatment of a pathogen infection comprising an effective amount of a viral-specific T cell of the previous aspect in a pharmaceutically acceptable excipient. In one embodiment, the composition further comprising a cytokine selected from any one or more of IL-2, IL-3, IL-6, IL-11, IL7, IL12, IL15, IL21, granulocyte macrophage colony stimulating factor, alpha, beta or gamma interferon and erythropoietin.

[0018] In another aspect, the disclosure provides a kit comprising an immunoresponsive cell comprising a receptor that binds an antigen and an exogenous co-stimulatory ligand. In one embodiment, the kit further comprises written directions for using said cell for the treatment of a subject having neoplasia, a pathogen infection, an autoimmune disorder, or an allogeneic transplant.

[0019] In various embodiments of any previous aspect, the method further comprises the step of obtaining the immunoresponsive cell or co-stimulatory ligand. In still other embodiments of the previous aspects, the co-stimulatory ligand is constitutively or inducibly expressed. In yet other embodiments of the previous aspects, at least two co-stimulatory ligands are constitutively expressed. In various embodiments of any previous aspect, the cell is selected from any one or more of a T cell, a Natural Killer (NK) cell, a cytotoxic T lymphocyte (CTL), and a regulatory T cell. In still other embodiments of the previous aspects, the antigen is a tumor or pathogen antigen, e.g., any one or more of prostate-specific membrane antigen (PSMA), Carcinoembryonic Antigen (CEA), IL13Ralpha, her-2, CD19, NY-ESO-1, HIV-1 Gag, Lewis Y, Mart-1, gp100, tyrosinase, WT-1, hTERT, mesothelin. In still other embodiments of the previous aspects, the cell expresses a recombinant and/or an endogenous antigen receptor. In still other embodiments of the previous aspects, the co-stimulatory ligand is a tumor necrosis factor (TNF) ligand (e.g., 4-1BBL, OX40L, CD70, LIGHT, and CD30L) or an immunoglobulin (Ig) superfamily ligand (e.g., CD80 and CD86). In still other embodiments of the previous aspects, the immunoresponsive cell expresses at least one TNF ligand and at least one Ig superfamily ligand. In still other embodiments of the previous aspects, the cell expresses 4-1BBL and CD80. In still other embodiments of the previous aspects, an antigen presenting complex and/or co-stimulatory ligands are constitutively or inducibly expressed on the surface of the T cell. In still other embodiments of the previous aspects, the co-stimulatory ligand is expressed in a retroviral vector. In another embodiment, the tumor antigen is prostate-specific membrane antigen, CD19, NY-ESO-1, WT-1 or hTERT. In still other embodiments, the cell expresses a recombinant or an endogenous receptor for the antigen. In still other embodiments, the co-stimulatory ligand is a tumor necrosis factor (TNF) ligand or an immunoglobulin (Ig) superfamily ligand. In still other embodiments, the TNF ligand is selected from any one or more of 4-1BBL, OX40L, CD70, LIGHT, and CD30L. In various embodiments, the intracellular signaling domain is a ζ -chain signaling domain. In other embodiments, in the one costimulatory signal is provided through an engineered antigen receptor and the other by overexpressing a costimulatory ligand. In still other embodiments, the Ig superfamily ligand is selected from the group consisting of CD80 and CD86.

Definitions

[0020] By "CD80 polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: NP_005182 or a fragment thereof that acts as an Ig superfamily ligand.

[0021] By "CD80 nucleic acid molecule" is meant any polynucleotide encoding a CD80 polypeptide. An exemplary CD80 nucleic acid molecule is NM_005191.

[0022] By "4-1BBL polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: P41273 or NP_001552 or a fragment thereof that acts as a tumor necrosis factor (TNF) ligand.

[0023] By "4-1BBL nucleic acid molecule" is meant a polynucleotide encoding a 4-1BBL polypeptide.

[0024] By "OX40L polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: BAB 18304 or NP_003317 or a fragment thereof that is a tumor necrosis factor (TNF) ligand.

[0025] By "OX40L nucleic acid molecule" is meant a polynucleotide encoding a OX40L polypeptide.

[0026] By "CD70 polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: NP_001243 or a fragment thereof that acts as a tumor necrosis factor (TNF) ligand.

[0027] By "CD70 nucleic acid molecule" is meant a polynucleotide encoding a CD70 polypeptide.

[0028] By "Light polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: NP_742011 or a fragment thereof that acts as a tumor necrosis factor (TNF) ligand.

[0029] By "Light nucleic acid molecule" is meant a polynucleotide encoding a Light polypeptide.

[0030] By "CD30L polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: AAB97877 or a fragment thereof that acts as a tumor necrosis factor (TNF) ligand.

[0031] By "CD30L nucleic acid molecule" is meant a polynucleotide encoding a CD30L polypeptide.

[0032] By "CD86 polypeptide" is meant a protein having at least 85% identity to NCBI Reference No: P42081 or a fragment thereof that acts as an Ig superfamily ligand.

[0033] By "CD86 nucleic acid molecule" is meant a polynucleotide encoding a CD86 polypeptide.

[0034] By "Pz1 polypeptide" is meant a protein having at least 85% identity to the protein described by Gong et al., *Neoplasia* 1:123-7, 1999 or a fragment thereof.

[0035] By "P28z polypeptide" is meant a protein having at least 85% identity to the protein described by Maher et al. *Nature Biotechnology*, Vol 20, Jan 2002, 70-75 or a fragment

thereof. The ζ -chain signaling domain is also described by Maher et al. *Nature Biotechnology*, Vol 20, Jan 2002, 70-75.

[0036] Nucleic acid molecules useful in the methods of the invention include any nucleic acid molecule that encodes a polypeptide of the invention or a fragment thereof. Such nucleic acid molecules need not be 100% identical with an endogenous nucleic acid sequence, but will typically exhibit substantial identity. Polynucleotides having "substantial identity" to an endogenous sequence are typically capable of hybridizing with at least one strand of a double-stranded nucleic acid molecule. By "hybridize" is meant pair to form a double-stranded molecule between complementary polynucleotide sequences (e.g., a gene described herein), or portions thereof, under various conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507).

[0037] For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and more preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and more preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30° C, more preferably of at least about 37° C, and most preferably of at least about 42° C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred embodiment, hybridization will occur at 30° C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37° C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 μ g/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42° C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50% formamide, and 200 μ g/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

[0038] For most applications, washing steps that follow hybridization will also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include a temperature of at least about 25° C, more preferably of at least about 42° C, and even more preferably of at least about 68° C. In a preferred embodiment, wash steps will occur at 25° C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42° C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 68° C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art. Hybridization techniques are well known to those skilled in the art and are described, for example, in Benton and Davis (*Science* 196:180, 1977); Grunstein

and Hogness (Proc. Natl. Acad. Sci., USA 72:3961, 1975); Ausubel et al. (Current Protocols in Molecular Biology, Wiley Interscience, New York, 2001); Berger and Kimmel (Guide to Molecular Cloning Techniques, 1987, Academic Press, New York); and Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York.

[0039] By "substantially identical" is meant a polypeptide or nucleic acid molecule exhibiting at least 50% identity to a reference amino acid sequence (for example, any one of the amino acid sequences described herein) or nucleic acid sequence (for example, any one of the nucleic acid sequences described herein). Preferably, such a sequence is at least 60%, more preferably 80% or 85%, and more preferably 90%, 95% or even 99% identical at the amino acid level or nucleic acid to the sequence used for comparison. Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, BLAST, BESTFIT, GAP, or PILEUP/Prettybox programs). Such software matches identical or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. In an exemplary approach to determining the degree of identity, a BLAST program may be used, with a probability score between e-3 and e-100 indicating a closely related sequence.

[0040] By "analog" is meant a structurally related polypeptide or nucleic acid molecule having the function of a reference polypeptide or nucleic acid molecule.

[0041] The term "ligand" as used herein refers to a molecule that binds to a receptor. In particular, the ligand binds a receptor on another cell, allowing for cell-to-cell recognition.

[0042] The term "constitutive expression" as used herein refers to expression under all physiological conditions.

[0043] The term "chimeric antigen receptor" (CAR) as used herein refers to a tumor antigen-binding domain that is fused to an intracellular signaling domain capable of activating T cells. Most commonly, the CAR's extracellular binding domain is derived from a murine or humanized monoclonal antibody.

[0044] By "disease" is meant any condition or disorder that damages or interferes with the normal function of a cell, tissue, or organ. Examples of diseases include neoplasia or pathogen infection of cell.

[0045] By "effective amount" is meant an amount sufficient to arrest, ameliorate, or inhibit the continued proliferation, growth, or metastasis (e.g., invasion, or migration) of a neoplasia.

[0046] By "enforcing tolerance" is meant preventing the activity of self-reactive cells or immunoresponsive cells that target transplanted organs or tissues.

[0047] By "exogenous" is meant a nucleic acid molecule or polypeptide that is not endogenously present in the cell, or not present at a level sufficient to achieve the functional effects obtained when over-expressed. The term "exogenous" would therefore encompass any recombinant nucleic acid molecule or polypeptide expressed in a cell, such as foreign, heterologous, and over-expressed nucleic acid molecules and polypeptides.

[0048] By a "heterologous nucleic acid molecule or polypeptide" is meant a nucleic acid molecule (e.g., a cDNA, DNA or RNA molecule) or polypeptide that is not normally present in a cell or sample obtained from a cell. This nucleic acid may be from another organism, or it may be, for example, an mRNA molecule that is not normally expressed in a cell or sample.

[0049] By "immunoresponsive cell" is meant a cell that functions in an immune response or a progenitor, or progeny thereof.

[0050] By "isolated cell" is meant a cell that is separated from the molecular and/or cellular components that naturally accompany the cell.

[0051] The term "tumor antigen-binding domain" as used herein refers to a domain capable of specifically binding a particular antigenic determinant or set of antigenic determinants present on a tumor.

[0052] The term "obtaining" as in "obtaining the agent" is intended to include purchasing, synthesizing or otherwise acquiring the agent (or indicated substance or material).

[0053] By "modulate" is meant positively or negatively alter. Exemplary modulations include a 1%, 2%, 5%, 10%, 25%, 50%, 75%, or 100% change.

[0054] By "neoplasia" is meant a disease characterized by the pathological proliferation of a cell or tissue and its subsequent migration to or invasion of other tissues or organs. Neoplasia growth is typically uncontrolled and progressive, and occurs under conditions that would not elicit, or would cause cessation of, multiplication of normal cells. Neoplasias can affect a variety of cell types, tissues, or organs, including but not limited to an organ selected from the group consisting of bladder, bone, brain, breast, cartilage, glia, esophagus, fallopian tube, gallbladder, heart, intestines, kidney, liver, lung, lymph node, nervous tissue, ovaries, pancreas, prostate, skeletal muscle, skin, spinal cord, spleen, stomach, testes, thymus, thyroid, trachea, urogenital tract, ureter, urethra, uterus, and vagina, or a tissue or cell type thereof. Neoplasias include cancers, such as sarcomas, carcinomas, or plasmacytomas (malignant tumor of the plasma cells). Illustrative neoplasms for which the invention can be used include, but are not limited to leukemias (e.g., acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia), polycythemia vera, lymphoma (Hodgkin's disease, non-Hodgkin's disease), Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors such as sarcomas and carcinomas (e.g., fibrosarcoma, myxosarcoma,

liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endothelirosarcoma, lymphangiosarcoma, lymphangioendothelirosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, nile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodenrolioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma). In one embodiment, screening methods of the disclosure identify compositions that are useful for treating breast or lung cancer.

[0055] By "receptor" is meant a polypeptide, or portion thereof, present on a cell membrane that selectively binds one or more ligand.

[0056] By "recognize" is meant selectively binds a target. A T cell that recognizes a virus typically expresses a receptor that binds an antigen expressed by the virus.

[0057] By "pathogen" is meant a virus, bacteria, fungi, parasite or protozoa capable of causing disease.

[0058] Exemplary viruses include, but are not limited to, *Retroviridae* (e.g. human immunodeficiency viruses, such as HIV-1 (also referred to as HDTV-III, LAVE or HTLV-III/LAV, or HIV-III; and other isolates, such as HIV-LP; *Picornaviridae* (e.g. polio viruses, hepatitis A virus; enteroviruses, human Coxsackie viruses, rhinoviruses, echoviruses); *Caliciviridae* (e.g. strains that cause gastroenteritis); *Togaviridae* (e.g. equine encephalitis viruses, rubella viruses); *Flaviridae* (e.g. dengue viruses, encephalitis viruses, yellow fever viruses); *Coronoviridae* (e.g. coronaviruses); *Rhabdoviridae* (e.g. vesicular stomatitis viruses, rabies viruses); *Filoviridae* (e.g. ebola viruses); *Paramyxoviridae* (e.g. parainfluenza viruses, mumps virus, measles virus, respiratory syncytial virus); *Orthomyxoviridae* (e.g. influenza viruses); *Bunyaviridae* (e.g. Hantaan viruses, bunga viruses, phleboviruses and Nairo viruses); *Arena viridae* (hemorrhagic fever viruses); *Reoviridae* (e.g. reoviruses, orbiviruses and rotaviruses); *Birnaviridae*; *Hepadnaviridae* (Hepatitis B virus); *Parvovirida* (parvoviruses); *Papovaviridae* (papilloma viruses, polyoma viruses); *Adenoviridae* (most adenoviruses); *Herpesviridae* (herpes simplex virus (HSV) 1 and 2, varicella zoster virus, cytomegalovirus (CMV), herpes virus; *Poxviridae* (variola viruses, vaccinia viruses, pox viruses); and *Iridoviridae* (e.g. African swine fever virus); and unclassified viruses (e.g. the agent of delta hepatitis (thought to be a defective satellite of hepatitis B virus), the agents of non-A, non-B hepatitis (class 1 = internally transmitted; class 2 = parenterally transmitted (i.e. Hepatitis C); Norwalk and related viruses, and astroviruses).

[0059] Exemplary bacteria include, but are not limited to, *Pasteurella*, *Staphylococci*, *Streptococcus*, *Escherichia coli*, *Pseudomonas* species, and *Salmonella* species. Specific examples of infectious bacteria include but are not limited to, *Helicobacter pyloris*, *Borelia*

burgdorferi, Legionella pneumophila, Mycobacteria ssp (e.g. M tuberculosis, M. avium, M. intracellulare, M. kansaii, M. gordonae), Staphylococcus aureus, Neisseria gonorrhoeae, Neisseria meningitidis, Listeria monocytogenes, Streptococcus pyogenes (Group A Streptococcus), Streptococcus agalactiae (Group B Streptococcus), Streptococcus (viridans group), Streptococcus faecalis, Streptococcus bovis, Streptococcus (anaerobic ssp.), Streptococcus pneumoniae, pathogenic Campylobacter sp., Enterococcus sp., Haemophilus influenzae, Bacillus antracis, corynebacterium diphtheriae, corynebacterium sp., Erysipelothrrix rhusiopathiae, Clostridium perfringers, Clostridium tetani, Enterobacter aerogenes, Klebsiella pneumoniae, Pasteurella multocida, Bacteroides sp., Fusobacterium nucleatum, Streptobacillus moniliformis, Treponema pallidum, Treponema pertenue, Leptospira, Rickettsia, and Actinomyces israelii.

[0060] By "specifically binds" is meant a polypeptide or fragment thereof that recognizes and binds a polypeptide of interest, but which does not substantially recognize and bind other molecules in a sample, for example, a biological sample, which naturally includes a polypeptide of the disclosure.

[0061] The term "tumor antigen" as used herein refers to any polypeptide expressed by a tumor that is capable of inducing an immune response.

[0062] By "virus antigen" is meant a polypeptide expressed by a virus that is capable of inducing an immune response.

[0063] The terms "comprises", "comprising", and are intended to have the broad meaning ascribed to them in U.S. Patent Law and can mean "includes", "including" and the like.

[0064] As used herein, "treatment" refers to clinical intervention in an attempt to alter the disease course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Therapeutic effects of treatment include, without limitation, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastases, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. By preventing progression of a disease or disorder, a treatment can prevent deterioration due to a disorder in an affected or diagnosed subject or a subject suspected of having the disorder, but also a treatment may prevent the onset of the disorder or a symptom of the disorder in a subject at risk for the disorder or suspected of having the disorder.

[0065] The term "subject" as used herein refers to a vertebrate, preferably a mammal, more preferably a human.

[0066] The term "immunocompromised" as used herein refers to a subject who has an immunodeficiency. The subject is very vulnerable to opportunistic infections, infections caused by organisms that usually do not cause disease in a person with a healthy immune system, but can affect people with a poorly functioning or suppressed immune system.

[0067] Other aspects of the invention are described in the following disclosure and are within the ambit of the invention.

BRIEF DESCRIPTION OF THE FIGURES

[0068] The following Detailed Description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying drawings.

[0069] Figures 1a-1f show that CD80 and 4-1BBL co-expression in human T-cells elicits robust proliferative responses after cyclic stimulations through their endogenous T cell receptor or through a chimeric antigen receptor without antigen presenting cell (APC)-provided costimulation. 0.4×10^6 purified human T lymphocytes was activated with antibody to CD3 (OKT3) and quantified the expansion of T lymphocytes retrovirally transduced with CD80, 4-1BBL, the combination of the two, or a control vector (Figure 1a,b). Transduction efficiencies were assessed by FACS analysis 48 h after gene transfer (Figure 1a). The absolute count of CD8⁺ T cells transduced with the indicated costimulatory ligand and stimulated weekly with 10 µg/ml plate-bound OKT3 is graphed in Figure 1b. The investigation was extended to Cytomegalovirus (CMV)-specific memory donor T cells (Figure 1c,d) and genetically redirected autologous T cells (Figure 1e,f) CMV- pp65-specific T lymphocytes, briefly expanded on HLA-A*0201⁺pp65⁺ artificial APCs and transduced with CD80, 4-1BBL, or a combination of the two (Figure 1c). Absolute counts of CD8⁺pp65⁺ T cells enriched and transduced as described in c after weekly restimulation with HLA-A*0201⁺ pp65-presenting Caco-2 tumor cells are shown in Figure 1d. To rapidly generate prostate cancer-reactive human T lymphocytes (Figure 1e,f), peripheral blood T cells were retrovirally transduced with the chimeric antigen receptor Pz1 (Gade et al., Cancer Res. 65:9080-9088, 2005), a non-HLA restricted antigen receptor specific for the tumor antigen PSMA. The Pz1 receptor comprises a PSMA-binding single-chain antibody fragment fused to the human CD3 ζ signaling domain. In Figure 1e, primary human CD3⁺ T lymphocytes was retrovirally transduced with Pz1 alone or in combination with CD80 and 4-1BBL. Transduction efficiencies were assessed by FACS analysis. The T cell expansion of Pz1-transduced human T lymphocytes cocultured weekly with LNCaP tumor monolayers is shown in Figure 1f. Costimulatory ligands were expressed on the T cell (top) or the tumor cell (bottom). Respective activation condition are depicted in cartoons shown in the right panels of b,d and f. Data are representative of three independent experiments. Each point in b, d and f represents the mean \pm s.e.m. of three randomly picked wells. * indicates $< 10^4$ cells.

[0070] Figures 2a-2c show the eradication of established - prostate-specific membrane antigen (PSMA) prostate carcinoma tumors in Scid/beige mice by Pz1⁺ T-cells transduced with CD80 and 4-1BBL. Figure 2a shows *in vivo* bioluminescent imaging and corresponding coronal MRI scans of firefly luciferase⁺PC3-PSMA tumors in Scid/beige mice four weeks after systemic inoculation with 4×10^6 tumor cells (day 0 of T-cell treatment), and eighteen days after adoptive transfer of 4×10^6 CD8⁺Pz1⁺ or Pz1⁺CD80⁺4-1BBL⁺ T lymphocytes. An equal number of T-cells bearing the human CD19-targeting chimeric antigen receptor 19z were injected in the control group. Pseudocolor images superimposed on conventional photographs are shown. The same animals imaged before and after treatment using bioluminescent imaging and MRI are shown. The two mice represent a total of n=10. Figure 2b shows three graphs. Bioluminescent tumor signal

quantified per animal every two days over a time period of 28 days. Acquisitions with saturated pixels -although included in the figure to allow direct visual comparison between groups- were not used for photon quantification but repeated at a shorter acquisition time. The graph shows photons/second/cm²/surface radius (sr) versus days after T cell injection. Every line corresponds to one animal with each dot representing the average photon count of the ventral and dorsal acquisition per animal at any given time point. Survival is illustrated in the Kaplan-Meier curve in Figure 2c.

[0071] Figures 3a-3d show robust, yet tumor antigen dependent, *in vivo* proliferation of CD80⁺4-1BBL⁺ T lymphocytes. Figure 3a shows comparative *in vivo* bioluminescent imaging of adoptively transferred T cells in PC3-PSMA tumor bearing Scid/beige mice on days 0, 8 and 18 after the injection of 4 x 10⁶ CD8⁺ clickbeetle luciferase (click-luc)-expressing Pz1-transduced or Pz1⁺CD80⁺4-1BBL⁺ transduced T-lymphocytes. As an antigen specificity control, an equal number of 19z⁺CD80⁺4-1BBL⁺ T cells were infused. T-cell treatment started as in Figure 2 four weeks after the systemic injection of 4 X10⁶ PC3-PSMA tumor cells. The five mice per group shown in each panel represent a total of n=8/group. Acquisitions with saturated pixels -although shown in the figure to allow direct visual comparison were not used for photon quantification but repeated at a shorter acquisition time. Figure 3b shows three graphs. Clickbeetle luciferase signal intensities from sequential bioluminescence imaging were collected every 2 days after T-cell transfer over a sixteen day time period. Every line represents one animal with each dot showing the average photon count of the ventral and dorsal acquisition per animal at any given time point. Figure 3c shows a multicolor FACS analysis of a lung single-cell suspension prepared from a representative animal infused either with Pz1⁺ or Pz1⁺CD80⁺4-1BBL⁺ T lymphocytes, 6 d after T cell transfer. Cells were stained with idiotypic antiserum specific for the chimeric antigen receptor Pz1. In Figure 3d absolute pulmonary Pz1⁺T cell numbers (total cell counts of viable Trypan blue-negative cells x percentages of Pz1⁺T cells). Shown bar graphs represent the mean ± s.e.m. of three mice. Figures 4a-4f show costimulation of T cells in cis as a results of the **colocalization of** CD80, 4-1BBL and their receptors CD28 and 4-1BB into the immunological synapse after induced T-cell/tumor cell cluster formation. In Figure 4a-d, peripheral blood human T lymphocytes were transduced with Pz1 and restimulated with LNCaP-CD80 cells before a second gene transfer with the bicistronic vector encoding the dsRed-monomer-4-1BBL fusion protein and CD80. After negative magnetic CD8⁺ isolation, T cells were labeled with FITC-Choleratoxin B (CTB), and incubated with unmodified LNCaP tumor cells or alone. Fixed conjugates were permeabilized, stained with the indicated antisera and visualized by confocal microscopy. T cell-LNCaP cell clusters from three independent experiments were randomly chosen. Scale bars, 10 µm. The numbers of clusters with clear concentrations of the indicated costimulatory ligand or receptor at the T cell-APC junction over the total number of analyzed clusters is shown in the lower right panels of Figure 4a and b. Figure 4a is a set of ten confocal micrographs exemplifying the polarization of 4-1BBL - expressed as a ds-Red fusion protein in combination with CD80 on CD8⁺ Pz1 transduced T cells - into the immunological synapse. T-cell/LnCaP tumor clusters were incubated for 50 minutes before fixation, permeabilization and incubation with anti 4-1BB antiserum. In Figure 4b, fixed T cell-tumor cell clusters were incubated with anti-CD80 and anti-CD28 antisera. Again, colocalization of CD80 with its receptor CD28 in the immunological synapse after tumor antigen encounter is visualized. Figure 4c illustrates the augmented accumulation of granzyme-B at the immunological synapse. This accumulation is dependent on a

functional engagement of 4-1BB by its ligand 4-1BBL, which is expressed on the same T cell surface. Primary human T lymphocytes were genetically modified as in Figure 4a, b. As indicated, retroviral vectors encoding Pz1 also express control shRNA or 4-1BB targeting shRNA under the control of the U6 promoter in their 3'LTR as described herein below. Cell conjugates in the top (Pz1⁺control shRNA) and middle (and Pz1⁺CD80⁺dsred 4-1BBL⁺control shRNA) row represent CD80⁺dsred 4-1BBL- untransduced and transduced T-lymphocytes, respectively, cultured in the same well and conjugated to LnCaP on the same glass slide. Figure 4d shows the relative recruitment index (RRI) and the relative intensity, calculated as described herein below, of Granzyme-B-Alexa 647 at the T cell - antigen presenting cell interphase. Data points in each group show the calculated value of 35 analyzed conjugates (symbols) and their mean (-) of three independent experiments. *P = 0.0001; **P < 0.0001. In Figure 4e,f, show NF- κ B-luciferase assays in isolated single T cell clones. These results corroborate autocostimulation as an operant mechanism of delivering costimulatory signals. A CD3⁺CD2S⁺4-1BB- Jurkat T cell clone stably transfected with with an NF- κ B-luciferase reporter was retrovirally transduced with a tricistronic vector coexpressing 4-1BB, 4-1BBL and CD80 or a control vector. To preclude bystander costimulation, transduced T lymphocytes were subcloned into OKT-3-coated 96-well plates before the first detectable surface expression of encoded proteins. Twelve hours after activation, the presence of single T cells in the wells was microscopically confirmed and the bioluminescent signal was measured on a single-cell level. Figure 4e shows 36 bioluminescence acquisitions of transduced single T cells assembled with Adobe Illustrator software. Respective bioluminescent single-cell signals were quantified, normalized to background bioluminescence, and plotted in Figure 4f.

[0072] Figures 5a-5e show that CD80 and 4-1BBL-displaying T lymphocytes trans-costimulate unmodified, antigen specific bystander T-cells through physical contact. Figure 5a shows a set of 4 confocal micrographs of a Pz1⁺ T cell engaging an LNCAp tumor cell while in physical T cell-Tcell contact with a bystander Pz1⁺ T lymphocyte transduced with CD80 and dsRed-4-1BBL. Cell clusters were induced and analyzed as described in Figure 4a,b. Scale bars, 10 μ m. In Figure 5b-d, peripheral T lymphocytes of a cytomegalovirus (CMV)-seropositive HLA A2.1⁺ donor were transduced with Pz1 or co-transduced with Pz1, CD80 and 4-1BBL. In parallel, CMV-specific, genetically unmodified cytotoxic T lymphocytes (CTLs) from the same donor were enriched by artificial antigen presenting cell (AAPC) co-culture as described herein below. These cells were labeled with carboxyfluorescein succinimidyl ester (CFSE). Bead-sorted Pz1⁺ (Figure 5b) or Pz1⁺CD80⁺4-1BBL⁺ (Figure 5c,) T lymphocytes were admixed to expanded CMV-reactive pp65⁺ CTLs at a 1:1 ratio and exposed to irradiated Caco-2 tumor cells retrovirally transduced to present surface pp65 in an HLA A2.1-dependent, as well as PSMA in an HLA-independent manner. Alternatively, a transwell membrane separated Pz1⁺CD80⁺4-1BBL⁺ T-cells from pp65⁺T-cells, both engaging Caco-2 tumor (Figure 5d). Respective co-culture conditions are depicted in cartoons shown on the left. Two days after tumor antigen contact intracellular Granzyme B levels gated on CD8⁺ T-cells were quantified by FACS analysis (middle column). Granzyme B-antigen presenting cell (APC) mean fluorescent intensities (MFI) of the CFSE- (Pz1⁺) and CFSE⁺ (pp65⁺) T cell sub-populations are summarized on top of each profile. On day 7 CFSE dilutions and Pz1⁺ compared to pp65⁺ T-cell fractions were analyzed by flow cytometry after pp65 tetramer and CD8 staining (right column)). CFSE-MFI indicated on top of each profile are based on pp65⁺- populations after CD8⁺-gating. Total cell counts of CD8⁺ pp65⁺ T-cells after the 7 day

cultures are graphed in (Figure 5e). Each bar graph represents the mean \pm s.e.m of three randomly chosen wells. Data are representative of two independent experiments. No exogenous cytokines were added at any point of the co-culture.

[0073] Figures 6a and 6b show that the accumulation of adoptively transferred PSMA-redirected T lymphocytes at tumor sites is augmented by Pz1⁺CD80⁺4-1BBL⁺, but not 19z⁺CD80⁺4-1BBL⁺ T-cells in Scid/beige mice. Figure 6a shows dual *in vivo* bioluminescence imaging of external Gaussia luciferase (x-gaus-luc) in RM1.PGLS and Raji tumors in addition to Clickbeetle luciferase (Click-luc) in tumor targeting T-cells. Two weeks after the systemic injection of 1×10^6 CD19⁺x-gaus-luc⁺ Raji tumors and two days after the subsequent infusion of 5×10^5 PSMA⁺x-gaus-luc⁺RM1.PGLS tumor cells into the same animals, established tumors in the bone marrow (Raji) and lung (RM1) were treated with a combination of three cell populations of T cells transduced as indicated. Each animal received a total of 12×10^6 CD8⁺chimeric antigen receptor⁺ T-cells (4×10^6 T-cells/transduction condition). Notably, the T-cell population listed third (Pz1⁺, Pz1⁺CD80⁺4-1BBL⁺, 19z⁺CD80⁺4-1BBL⁺, left, middle, right row, respectively) were injected twelve hours after the combined injection of T-cells listed first and second to avoid T-cell/T-cell interaction in the lung due to crowding and not as a result of selective tumor antigen binding. At the indicated time points x-gaus-luc⁺ tumor cells or click-luc⁺ T-cells were monitored by bioluminescent imaging. On day 0 and day 4 a time period of at least 4 hours between tumor and T-cell imaging ensured the bioluminescent signal to return to background levels. A total number of n=5 Scid/beige mice were imaged per treatment group. Figure 5b shows a series of 6 graphs quantitating clickbeetle luciferase signal intensities from sequential bioluminescence imaging every day after T-cell transfer for a four day time period. Every line represents one animal with each dot showing the average photon count measured over the pulmonary area (top) or both femurs (bottom), respectively at any given time point.

[0074] Figure 7 shows that dual T cell-expressed CD80 and 4-1BBL elicits superior T cell expansion compared to CD28 and/or 4-1BB signaling elements fused in series with the Pz1 chimeric antigen receptor ζ signaling domain. Primary T cells were transduced with Pz1, P28z (Maher et al. Nature Biotechnology, Vol 20, Jan 2002, 70-75), which contains the CD28 signaling domain in series with the ζ chain, or P284-1BBz which includes both, CD28 and 4-1BB signaling regions. Alternatively, T lymphocytes were co-transduced with P28z and 4-1BBL (noted P28Z + 4-1BBL). Pz1 + CD80 + 4-1BBL refers to Pz1+ T cells that co-express both ligands, CD80 and 4-1BBL. Transduced T cells were stimulated weekly (indicated by arrows) on LNCaP tumor monolayers under conditions outlined in detail under Figure 1. The fold expansion of the CD8⁺ Pz1-transduced human T lymphocytes population is graphed.

[0075] Figure 8 provides amino acid sequences for CD80, 4-1BBL, OX40L, CD70, LIGHT, and CD30L.

DETAILED DESCRIPTION OF THE INVENTION

[0076] The present disclosure generally provides cells, including genetically modified

immunoresponsive cells (e.g., T cells, Natural Killer (NK) cells, cytotoxic T lymphocytes (CTL) cells) expressing at least one of an antigen-recognizing receptor and a co-stimulatory ligand and methods of use therefore for the treatment of neoplasia and other pathologies where an increase in an antigen-specific immune response is desired. The invention is based, at least in part, on the discovery that the constitutive retroviral expression of CD80 and 4-1BBL in co-transduced human T cells targeting prostate specific membrane antigen mounted a robust tumor-antigen-dependent T-cell proliferation coupled with a profound *in vivo* rejection of disseminated well-established prostate carcinoma tumors. Furthermore, CD80 and 4-1BBL expressing T cells provided costimulation of bystander T cells in trans in a contact dependent and antigen-specific manner at the tumor site. Taken together, the concept of genetically modified T cells as a constitutive pool of costimulatory ligands to optimally costimulate themselves in addition to enhancing the immunogenicity within the tumor microenvironment represents a significant advance over conventional adoptive T cell therapy. Furthermore, as demonstrated *ex vivo* using enriched CMV-specific T lymphocytes, this approach is not limited to the treatment of neoplasias, but is amenable to a wide range of applications where an increase in an antigen-specific immune response is desired, such applications include not only the treatment of neoplasias, but also for the enhancement of an immune response against a pathogen infection or an infectious disease and to reinforce immune tolerance in regulatory T cells in the context of autoimmunity or allogeneic transplantation.

Hematopoietic Cell Lineages

[0077] Mammalian hematopoietic (blood) cells provide a diverse range of physiologic activities. Hematopoietic cells are divided into lymphoid, myeloid and erythroid lineages. The lymphoid lineage, comprising B, T and natural killer (NK) cells, provides for the production of antibodies, regulation of the cellular immune system, detection of foreign agents in the blood, detection of cells foreign to the host, and the like. The term "T cells" as used herein refers to lymphocytes that mature in the thymus and are chiefly responsible for cell-mediated immunity. T cells are involved in the adaptive immune system. The term "natural killer (NK) cells" as used herein refers to lymphocytes that are part of cell-mediated immunity and act during the innate immune response. They do not require prior activation in order to perform their cytotoxic effect on target cells. Cytotoxic T cells (CTL or killer T cells) are a subset of T lymphocytes capable of inducing the death of infected somatic or tumor cells.

Cells for Use in the Methods

[0078] The present disclosure provides cells expressing at least one of an antigen-recognizing receptor and a co-stimulatory ligand and methods of using such cells for the treatment of a disease that requires an enhanced immune response. In one approach, tumor antigen-specific T cells, NK cells, CTL cells or other immunoresponsive cells are used as shuttles for the selective enrichment of one or more co-stimulatory ligands for the treatment or prevention of neoplasia. For example, a T cell expressing a co-stimulatory

ligands 4-1BBL and CD80 are constitutively co-expressed in a T cell that expresses a chimeric antigen receptor PZ1 that recognizes and binds Prostate Specific Membrane Antigen (PSMA). Such cells are administered to a human subject in need thereof for the treatment or prevention of prostate cancer. In another approach, viral antigen-specific T cells, NK cells, CTL cells can be used for the treatment of viral diseases. For example, CD80 and 4-1BBL are expressed in cytomegalovirus (CMV)-specific cytotoxic T lymphocytes for the treatment of CMV.

Tumor antigen-specific T lymphocytes (and NK cells)

[0079] Types of tumor antigen-specific human lymphocytes that can be used in the methods, without limitation, peripheral donor lymphocytes genetically modified to express chimeric antigen receptors (CARs) (Sadelain, M., et al. 2003 *Nat Rev Cancer* 3:35-45), peripheral donor lymphocytes genetically modified to express a full-length tumor antigen-recognizing T cell receptor complex comprising the α and β heterodimer (Morgan, R.A., et al. 2006 *Science* 314:126-129), lymphocyte cultures derived from tumor infiltrating lymphocytes (TILs) in tumor biopsies (Panelli, M.C., et al. 2000 *J Immunol* 164:495-504; Panelli, M.C., et al. 2000 *J Immunol* 164:4382-4392), and selectively *in vitro*-expanded antigen-specific peripheral blood leukocytes employing artificial antigen-presenting cells (AAPCs) or pulsed dendritic cells (Dupont, J., et al. 2005 *Cancer Res* 65:5417-5427; Papanicolaou, G.A., et al. 2003 *Blood* 102:2498-2505). The T cells may be autologous, allogeneic, or derived *in vitro* from engineered progenitor or stem cells.

[0080] Any suitable tumor antigen (antigenic peptide) is suitable for use in the tumor-related embodiments described herein. Sources of antigen include, but are not limited to cancer proteins. The antigen can be expressed as a peptide or as an intact protein or portion thereof. The intact protein or a portion thereof can be native or mutagenized. One suitable antigen is prostate specific membrane antigen (PSMA).

Viral antigen-specific T lymphocytes (and NK cells)

[0081] Suitable antigens for use in the treatment of pathogen infection or other infectious disease, for example, in an immunocompromised subject include, without limitation, viral antigens present in Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus.

[0082] The unpurified source of CTLs may be any known in the art, such as the bone marrow, fetal, neonate or adult or other hematopoietic cell source, e.g., fetal liver, peripheral blood or umbilical cord blood. Various techniques can be employed to separate the cells. For instance, negative selection methods can remove non-CTLs initially. mAbs are particularly useful for identifying markers associated with particular cell lineages and/or stages of differentiation for both positive and negative selections.

[0083] A large proportion of terminally differentiated cells can be initially removed by a

relatively crude separation. For example, magnetic bead separations can be used initially to remove large numbers of irrelevant cells. Preferably, at least about 80%, usually at least 70% of the total hematopoietic cells will be removed prior to cell isolation.

[0084] Procedures for separation include, but are not limited to, density gradient centrifugation; resetting; coupling to particles that modify cell density; magnetic separation with antibody-coated magnetic beads; affinity chromatography; cytotoxic agents joined to or used in conjunction with a mAb, including, but not limited to, complement and cytotoxins; and panning with antibody attached to a solid matrix, e.g. plate, chip, elutriation or any other convenient technique.

[0085] Techniques for separation and analysis include, but are not limited to, flow cytometry, which can have varying degrees of sophistication, e.g., a plurality of color channels, low angle and obtuse light scattering detecting channels, impedance channels.

[0086] The cells can be selected against dead cells, by employing dyes associated with dead cells such as propidium iodide (PI). Preferably, the cells are collected in a medium comprising 2% fetal calf serum (FCS) or 0.2% bovine serum albumin (BSA) or any other suitable, preferably sterile, isotonic medium.

[0087] Accordingly, the disclosure generally provides an immunoresponsive cell, such as a virus or tumor specific T cell comprising a receptor that binds an antigen and an exogenous co-stimulatory ligand (e.g., CD80, 4-1BBL, OX40L, CD70 and CD30L).

Vectors

[0088] Genetic modification of immunoresponsive cells (e.g., T cells, CTL cells, NK cells) can be accomplished by transducing a substantially homogeneous cell composition with a recombinant DNA construct. Preferably, a retroviral vector (either gamma-retroviral or lentiviral) is employed for the introduction of the DNA construct into the cell. For example, a polynucleotide encoding a co-stimulatory ligand protein (e.g., tumor necrosis factor (TNF) ligand, such as 4-1BBL, OX40L, CD70, LIGHT, and CD30L, or an Ig superfamily ligand, such as CD80 and CD86), or a receptor that binds an antigen, or a variant, or a fragment thereof, can be cloned into a retroviral vector and expression can be driven from its endogenous promoter, from the retroviral long terminal repeat, or from a promoter specific for a target cell type of interest. Non-viral vectors may be used as well.

Co-stimulatory ligands

[0089] The interaction with at least one co-stimulatory ligand provides a non-antigen-specific signal required for full activation of a T cell. Co-stimulatory ligands include, without limitation, tumor necrosis factor (TNF) ligands, cytokines (such as IL-2, IL-12, IL-15 or IL21), and immunoglobulin (Ig) superfamily ligands.

TNF ligands

[0090] Tumor necrosis factor (TNF) is a cytokine involved in systemic inflammation and stimulates the acute phase reaction. Its primary role is in the regulation of immune cells. Tumor necrosis factor (TNF) ligands share a number of common features. The majority of the ligands are synthesized as type II transmembrane proteins (extracellular C-terminus) containing a short cytoplasmic segment and a relatively long extracellular region. TNF ligands include, without limitation, nerve growth factor (NGF), CD40L (CD40L)/CD154, CD137L/4-1BBL, tumor necrosis factor alpha (TNF α), CD134L/OX40L/CD252, CD27L/CD70, Fas ligand (FasL), CD30L/CD153, tumor necrosis factor beta (TNF β)/lymphotoxin-alpha (LT α), lymphotoxin-beta (LT β), CD257/B cell-activating factor (BAFF)/Blys/THANK/Tall-1, glucocorticoid-induced TNF Receptor ligand (GITRL), and TNF-related apoptosis-inducing ligand (TRAIL), LIGHT (TNFSF14).

Ig Superfamily ligands

[0091] The immunoglobulin (Ig) superfamily is a large group of cell surface and soluble proteins that are involved in the recognition, binding, or adhesion processes of cells. These proteins share structural features with immunoglobulins -- they possess an immunoglobulin domain (fold). Immunoglobulin superfamily ligands include, without limitation, CD80 and CD86, both ligands for CD28.

[0092] For initial genetic modification of the cells to provide tumor or viral antigen-specific cells, a retroviral vector is generally employed for transduction, however any other suitable viral vector or delivery system can be used. For subsequent genetic modification of the cells to provide cells comprising an antigen presenting complex comprising at least two co-stimulatory ligands, retroviral gene transfer (transduction) likewise proves effective. Combinations of retroviruses and an appropriate packaging line are also suitable, where the capsid proteins will be functional for infecting human cells. Various amphotropic virus-producing cell lines are known, including, but not limited to, PA12 (Miller, et al. (1985) Mol. Cell. Biol. 5:431-437); PA317 (Miller, et al. (1986) Mol. Cell. Biol. 6:2895-2902); and CRIP (Danos, et al. (1988) Proc. Natl. Acad. Sci. USA 85:6460-6464). Non-amphotropic particles are suitable too, e.g., particles pseudotyped with VSVG, RD114 or GALV envelope and any other known in the art.

[0093] Possible methods of transduction also include direct co-culture of the cells with producer cells, e.g., by the method of Bregni, et al. (1992) Blood 80:1418-1422, or culturing with viral supernatant alone or concentrated vector stocks with or without appropriate growth factors and polycations, e.g., by the method of Xu, et al. (1994) Exp. Hemat. 22:223-230; and Hughes, et al. (1992) J. Clin. Invest. 89:1817.

[0094] Other transducing viral vectors can be used to express a co-stimulatory ligand of the disclosure in an immunoresponsive cell. Preferably, the chosen vector exhibits high efficiency of infection and stable integration and expression (see, e.g., Cayouette et al.,

Human Gene Therapy 8:423-430, 1997; Kido et al., Current Eye Research 15:833-844, 1996; Bloomer et al., Journal of Virology 71:6641-6649, 1997; Naldini et al., Science 272:263-267, 1996; and Miyoshi et al., Proc. Natl. Acad. Sci. U.S.A. 94:10319, 1997). Other viral vectors that can be used include, for example, adenoviral, lentiviral, and adeno-associated viral vectors, vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus (also see, for example, the vectors of Miller, Human Gene Therapy 15:14, 1990; Friedman, Science 244:1275-1281, 1989; Eglitis et al., BioTechniques 6:608-614, 1988; Tolstoshev et al., Current Opinion in Biotechnology 1:55-61, 1990; Sharp, The Lancet 337:1277-1278, 1991; Cornetta et al., Nucleic Acid Research and Molecular Biology 36:311-322, 1987; Anderson, Science 226:401-409, 1984; Moen, Blood Cells 17:407-416, 1991; Miller et al., Biotechnology 7:980-990, 1989; Le Gal La Salle et al., Science 259:988-990, 1993; and Johnson, Chest 107:77S-83S, 1995). Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg et al., N. Engl. J. Med 323:370, 1990; Anderson et al., U.S. Pat. No. 5,399,346).

[0095] Non-viral approaches can also be employed for the expression of a protein in cell. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner et al., Proc. Natl. Acad. Sci. U.S.A. 84:7413, 1987; Ono et al., Neuroscience Letters 17:259, 1990; Brigham et al., Am. J. Med. Sci. 298:278, 1989; Staubinger et al., Methods in Enzymology 101:512, 1983), asialoorosomucoid-polylysine conjugation (Wu et al., Journal of Biological Chemistry 263:14621, 1988; Wu et al., Journal of Biological Chemistry 264:16985, 1989), or by micro-injection under surgical conditions (Wolff et al., Science 247:1465, 1990). Other non-viral means for gene transfer include transfection *in vitro* using calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transplantation of normal genes into the affected tissues of a subject can also be accomplished by transferring a normal nucleic acid into a cultivatable cell type *ex vivo* (e.g., an autologous or heterologous primary cell or progeny thereof), after which the cell (or its descendants) are injected into a targeted tissue or are injected systemically.

[0096] cDNA expression for use in polynucleotide therapy methods can be directed from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element. For example, if desired, enhancers known to preferentially direct gene expression in specific cell types can be used to direct the expression of a nucleic acid. The enhancers used can include, without limitation, those that are characterized as tissue- or cell-specific enhancers. Alternatively, if a genomic clone is used as a therapeutic construct, regulation can be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

[0097] The resulting cells can then be grown under conditions similar to those for unmodified cells, whereby the modified cells can be expanded and used for a variety of purposes.

Polypeptides and Analogs

[0098] Also included in the disclosure are PZ1, P28z, 4-1BBL, OX40L, CD70, LIGHT, and CD30L polypeptides or fragments thereof that are modified in ways that enhance their anti-neoplastic activity when expressed in an immunoresponsive cell. The disclosure provides methods for optimizing an amino acid sequence or nucleic acid sequence by producing an alteration in the sequence. Such alterations may include certain mutations, deletions, insertions, or post-translational modifications. The disclosure further includes analogs of any naturally-occurring polypeptide of the disclosure. Analogs can differ from a naturally-occurring polypeptide by amino acid sequence differences, by post-translational modifications, or by both. Analogs will generally exhibit at least 85%, more preferably 90%, and most preferably 95% or even 99% identity with all or part of a naturally-occurring amino acid sequence. The length of sequence comparison is at least 5, 10, 15 or 20 amino acid residues, preferably at least 25, 50, or 75 amino acid residues, and more preferably more than 100 amino acid residues. Again, in an exemplary approach to determining the degree of identity, a BLAST program may be used, with a probability score between e-3 and e-100 indicating a closely related sequence. Modifications include in vivo and in vitro chemical derivatization of polypeptides, e.g., acetylation, carboxylation, phosphorylation, or glycosylation; such modifications may occur during polypeptide synthesis or processing or following treatment with isolated modifying enzymes. Analogs can also differ from the naturally-occurring polypeptides by alterations in primary sequence. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethanemethylsulfate or by site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, Molecular Cloning: A Laboratory Manual (2d ed.), CSH Press, 1989, or Ausubel et al., *supra*). Also included are cyclized peptides, molecules, and analogs which contain residues other than L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g., β .beta. or γ .gamma. amino acids.

[0099] In addition to full-length polypeptides, the disclosure also provides fragments of any one of the polypeptides or peptide domains. As used herein, the term "a fragment" means at least 5, 10, 13, or 15 amino acids. In other embodiments a fragment is at least 20 contiguous amino acids, at least 30 contiguous amino acids, or at least 50 contiguous amino acids, and in other embodiments at least 60 to 80, 100, 200, 300 or more contiguous amino acids. Fragments can be generated by methods known to those skilled in the art or may result from normal protein processing (e.g., removal of amino acids from the nascent polypeptide that are not required for biological activity or removal of amino acids by alternative mRNA splicing or alternative protein processing events).

[0100] Non-protein analogs have a chemical structure designed to mimic the functional activity of a protein. Such analogs are administered according to methods of the disclosure. Such analogs may exceed the physiological activity of the original polypeptide. Methods of analog design are well known in the art, and synthesis of analogs can be carried out according to such methods by modifying the chemical structures such that the resultant analogs increase the anti-neoplastic activity of the original polypeptide when expressed in an immunoresponsive cell. These chemical modifications include, but are not limited to, substituting alternative R groups and varying the degree of saturation at specific carbon atoms of a reference polypeptide. Preferably, the protein analogs are relatively

resistant to in vivo degradation, resulting in a more prolonged therapeutic effect upon administration. Assays for measuring functional activity include, but are not limited to, those described in the Examples below.

Administration

[0101] Compositions comprising genetically modified immunoresponsive cells (e.g., T cells, NK cells, CTL cells, or their progenitors) can be provided systemically or directly to a subject for the treatment of a neoplasia, pathogen infection, or infectious disease. In one embodiment, cells are directly injected into an organ of interest (e.g., an organ affected by a neoplasia). Alternatively, compositions comprising genetically modified immunoresponsive cells are provided indirectly to the organ of interest, for example, by administration into the circulatory system (e.g., the tumor vasculature). Expansion and differentiation agents can be provided prior to, during or after administration of the cells to increase production of T cells, NK cells, or CTL cells *in vitro* or *in vivo*.

[0102] The modified cells can be administered in any physiologically acceptable vehicle, normally intravascularly, although they may also be introduced into bone or other convenient site where the cells may find an appropriate site for regeneration and differentiation (e.g., thymus). Usually, at least 1×10^5 cells will be administered, eventually reaching 1×10^{10} , or more. Genetically modified immunoresponsive cells can comprise a purified population of cells. Those skilled in the art can readily determine the percentage of genetically modified immunoresponsive cells in a population using various well-known methods, such as fluorescence activated cell sorting (FACS). Preferable ranges of purity in populations comprising genetically modified immunoresponsive cells are about 50 to about 55%, about 55 to about 60%, and about 65 to about 70%. More preferably the purity is about 70 to about 75%, about 75 to about 80%, about 80 to about 85%; and still more preferably the purity is about 85 to about 90%, about 90 to about 95%, and about 95 to about 100%. Dosages can be readily adjusted by those skilled in the art (e.g., a decrease in purity may require an increase in dosage). The cells can be introduced by injection, catheter, or the like. If desired, factors can also be included, including, but not limited to, interleukins, e.g. IL-2, IL-3, IL-6, and IL-11, as well as the other interleukins, the colony stimulating factors, such as G-, M- and GM-CSF, interferons, e.g..gamma.-interferon and erythropoietin.

[0103] Compositions include pharmaceutical compositions comprising genetically modified immunoresponsive cells or their progenitors and a pharmaceutically acceptable carrier. Administration can be autologous or heterologous. For example, immunoresponsive cells, or progenitors can be obtained from one subject, and administered to the same subject or a different, compatible subject.

Peripheral blood derived immunoresponsive cells or their progeny (e.g., *in vivo*, *ex vivo* or *in vitro* derived) can be administered via localized injection, including catheter administration, systemic injection, localized injection, intravenous injection, or parenteral administration. When administering a therapeutic composition (e.g., a pharmaceutical composition containing a genetically modified immunoresponsive cell), it will generally be formulated in a unit dosage injectable form (solution, suspension, emulsion).

Formulations

[0104] Compositions of the disclosure comprising genetically modified immunoresponsive cells can be conveniently provided as sterile liquid preparations, e.g., isotonic aqueous solutions, suspensions, emulsions, dispersions, or viscous compositions, which may be buffered to a selected pH. Liquid preparations are normally easier to prepare than gels, other viscous compositions, and solid compositions. Additionally, liquid compositions are somewhat more convenient to administer, especially by injection. Viscous compositions, on the other hand, can be formulated within the appropriate viscosity range to provide longer contact periods with specific tissues. Liquid or viscous compositions can comprise carriers, which can be a solvent or dispersing medium containing, for example, water, saline, phosphate buffered saline, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like) and suitable mixtures thereof.

[0105] Sterile injectable solutions can be prepared by incorporating the genetically modified immunoresponsive cells in the required amount of the appropriate solvent with various amounts of the other ingredients, as desired. Such compositions may be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose, dextrose, or the like. The compositions can also be lyophilized. The compositions can contain auxiliary substances such as wetting, dispersing, or emulsifying agents (e.g., methylcellulose), pH buffering agents, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard texts, such as "REMINGTON'S PHARMACEUTICAL SCIENCE", 17th edition, 1985, may be consulted to prepare suitable preparations, without undue experimentation.

[0106] Various additives which enhance the stability and sterility of the compositions, including antimicrobial preservatives, antioxidants, chelating agents, and buffers, can be added. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin. However, any vehicle, diluent, or additive used would have to be compatible with the genetically modified immunoresponsive cells or their progenitors.

[0107] The compositions can be isotonic, i.e., they can have the same osmotic pressure as blood and lacrimal fluid. The desired isotonicity of the compositions may be accomplished using sodium chloride, or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

[0108] Viscosity of the compositions, if desired, can be maintained at the selected level using a pharmaceutically acceptable thickening agent. Methylcellulose is preferred because it is readily and economically available and is easy to work with. Other suitable thickening agents include, for example, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, carbomer, and the like. The preferred concentration of the

thickener will depend upon the agent selected. The important point is to use an amount that will achieve the selected viscosity. Obviously, the choice of suitable carriers and other additives will depend on the exact route of administration and the nature of the particular dosage form, e.g., liquid dosage form (e.g., whether the composition is to be formulated into a solution, a suspension, gel or another liquid form, such as a time release form or liquid-filled form).

[0109] Those skilled in the art will recognize that the components of the compositions should be selected to be chemically inert and will not affect the viability or efficacy of the genetically modified immunoresponsive cells as described herein. This will present no problem to those skilled in chemical and pharmaceutical principles, or problems can be readily avoided by reference to standard texts or by simple experiments (not involving undue experimentation), from this disclosure and the documents cited herein.

[0110] One consideration concerning the therapeutic use of genetically modified immunoresponsive cells is the quantity of cells necessary to achieve an optimal effect. The quantity of cells to be administered will vary for the subject being treated. In a one embodiment, between 10^4 to 10^{10} , between 10^5 to 10^9 , or between 10^6 and 10^8 genetically modified immunoresponsive cells are administered to a human subject. In preferred embodiments, at least about 1×10^8 , 2×10^8 , 3×10^8 , 4×10^8 , and 5×10^8 genetically modified immunoresponsive cells are administered to a human subject. The precise determination of what would be considered an effective dose may be based on factors individual to each subject, including their size, age, sex, weight, and condition of the particular subject. Dosages can be readily ascertained by those skilled in the art from this disclosure and the knowledge in the art.

[0111] The skilled artisan can readily determine the amount of cells and optional additives, vehicles, and/or carrier in compositions and to be administered in methods. Typically, any additives (in addition to the active stem cell(s) and/or agent(s)) are present in an amount of 0.001 to 50 % (weight) solution in phosphate buffered saline, and the active ingredient is present in the order of micrograms to milligrams, such as about 0.0001 to about 5 wt %, preferably about 0.0001 to about 1 wt %, still more preferably about 0.0001 to about 0.05 wt % or about 0.001 to about 20 wt %, preferably about 0.01 to about 10 wt %, and still more preferably about 0.05 to about 5 wt %. Of course, for any composition to be administered to an animal or human, and for any particular method of administration, it is preferred to determine therefore: toxicity, such as by determining the lethal dose (LD) and LD50 in a suitable animal model e.g., rodent such as mouse; and, the dosage of the composition(s), concentration of components therein and timing of administering the composition(s), which elicit a suitable response. Such determinations do not require undue experimentation from the knowledge of the skilled artisan, this disclosure and the documents cited herein. And, the time for sequential administrations can be ascertained without undue experimentation.

Methods of Treatment

[0112] Provided herein are methods for treating neoplasia in a subject. Also contemplated

herein are methods for treating a pathogen infection or other infectious disease in a subject, such as an immunocompromised human subject. The methods comprise administering a T cell, NK cell, or CTL cell in an amount effective to achieve the desired effect, be it palliation of an existing condition or prevention of recurrence. For treatment, the amount administered is an amount effective in producing the desired effect. An effective amount can be provided in one or a series of administrations. An effective amount can be provided in a bolus or by continuous perfusion.

[0113] An "effective amount" (or, "therapeutically effective amount") is an amount sufficient to effect a beneficial or desired clinical result upon treatment. An effective amount can be administered to a subject in one or more doses. In terms of treatment, an effective amount is an amount that is sufficient to palliate, ameliorate, stabilize, reverse or slow the progression of the disease, or otherwise reduce the pathological consequences of the disease. The effective amount is generally determined by the physician on a case-by-case basis and is within the skill of one in the art. Several factors are typically taken into account when determining an appropriate dosage to achieve an effective amount. These factors include age, sex and weight of the subject, the condition being treated, the severity of the condition and the form and effective concentration of the antigen-binding fragment administered.

[0114] For adoptive immunotherapy using antigen-specific T cells, cell doses in the range of 10^9 are typically infused. Upon administration of the genetically modified cells into the host and subsequent differentiation, T cells are induced that are specifically directed against the specific antigen. "Induction" of T cells can include inactivation of antigen-specific T cells such as by deletion or anergy. Inactivation is particularly useful to establish or reestablish tolerance such as in autoimmune disorders. The modified cells can be administered by any method known in the art including, but not limited to, intravenous, subcutaneous, intranodal, intratumoral, intrathecal, intrapleural, intraperitoneal and directly to the thymus.

Therapeutic Methods

[0115] The disclosure provides methods for increasing an immune response in a subject in need thereof. In one embodiment, the disclosure provides methods for treating or preventing a neoplasia in a subject. The disclosure provides therapies that are particularly useful for the treatment of subjects having prostate cancer, or metastatic prostate cancer that is not amenable to conventional therapeutic interventions. Suitable human subjects for therapy typically comprise two treatment groups that can be distinguished by clinical criteria. Subjects with "advanced disease" or "high tumor burden" are those who bear a clinically measurable tumor. A clinically measurable tumor is one that can be detected on the basis of tumor mass (e.g., by palpation, CAT scan, sonogram, mammogram or X-ray; positive biochemical or histopathologic markers on their own are insufficient to identify this population). A pharmaceutical composition embodied in this disclosure is administered to these subjects to elicit an anti-tumor response, with the objective of palliating their condition. Ideally, reduction in tumor mass occurs as a result, but any clinical improvement

constitutes a benefit. Clinical improvement includes decreased risk or rate of progression or reduction in pathological consequences of the tumor.

[0116] A second group of suitable subjects is known in the art as the "adjuvant group." These are individuals who have had a history of neoplasia, but have been responsive to another mode of therapy. The prior therapy can have included (but is not restricted to, surgical resection, radiotherapy, and traditional chemotherapy. As a result, these individuals have no clinically measurable tumor. However, they are suspected of being at risk for progression of the disease, either near the original tumor site, or by metastases. This group can be further subdivided into high-risk and low-risk individuals. The subdivision is made on the basis of features observed before or after the initial treatment. These features are known in the clinical arts, and are suitably defined for each different neoplasia. Features typical of high-risk subgroups are those in which the tumor has invaded neighboring tissues, or who show involvement of lymph nodes.

[0117] Another group have a genetic predisposition to neoplasia but have not yet evidenced clinical signs of neoplasia. For instance, women testing positive for a genetic mutation associated with breast cancer, but still of childbearing age, can wish to receive one or more of the antigen-binding fragments described herein in treatment prophylactically to prevent the occurrence of neoplasia until it is suitable to perform preventive surgery.

[0118] Human neoplasia subjects having any of the following neoplasias: glioblastoma, melanoma, neuroblastoma, adenocarcinoma, glioma, soft tissue sarcoma, and various carcinomas (including prostate and small cell lung cancer) are especially appropriate subjects. Suitable carcinomas further include any known in the field of oncology, including, but not limited to, astrocytoma, fibrosarcoma, myxosarcoma, liposarcoma, oligodendrogloma, ependymoma, medulloblastoma, primitive neural ectodermal tumor (PNET), chondrosarcoma, osteogenic sarcoma, pancreatic ductal adenocarcinoma, small and large cell lung adenocarcinomas, chordoma, angiosarcoma, endotheliosarcoma, squamous cell carcinoma, bronchoalveolar carcinoma, epithelial adenocarcinoma, and liver metastases thereof, lymphangiosarcoma, lymphangioendotheliosarcoma, hepatoma, cholangiocarcinoma, synovioma, mesothelioma, Ewing's tumor, rhabdomyosarcoma, colon carcinoma, basal cell carcinoma, sweat gland carcinoma, papillary carcinoma, sebaceous gland carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, testicular tumor, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendrogloma, meningioma, neuroblastoma, retinoblastoma, leukemia, multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease, breast tumors such as ductal and lobular adenocarcinoma, squamous and adenocarcinomas of the uterine cervix, uterine and ovarian epithelial carcinomas, prostatic adenocarcinomas, transitional squamous cell carcinoma of the bladder, B and T cell lymphomas (nodular and diffuse) plasmacytoma, acute and chronic leukemias, malignant melanoma, soft tissue sarcomas and leiomyosarcomas.

[0119] The subjects can have an advanced form of disease, in which case the treatment objective can include mitigation or reversal of disease progression, and/or amelioration of

side effects. The subjects can have a history of the condition, for which they have already been treated, in which case the therapeutic objective will typically include a decrease or delay in the risk of recurrence.

[0120] Accordingly, the disclosure provides a method of treating or preventing a neoplasia in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds a tumor antigen and a vector encoding a co-stimulatory ligand. In one embodiment, the neoplasia is selected from the group consisting of prostate cancer, colon cancer, breast cancer, and glioblastoma. In another embodiment, the tumor antigen is prostate-specific membrane antigen, CD19, NY-ESO-1, WT-1 or hTERT.

[0121] In another approach, the disclosure provides a method of enforcing tolerance in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds an antigen and a vector encoding a co-stimulatory ligand. In one embodiment, the method prevents or reduces an autoimmune disease or a disease associated with allogeneic transplantation.

[0122] As a consequence of constitutive surface expression of co-stimulatory ligands, adoptively transferred human T or NK cells are endowed with augmented proliferative, cytolytic, and survival capacities in an intrinsically poorly immunogenic tumor or immunodeficient environment devoid of co-stimulatory ligands. Furthermore, subsequent to their localization to tumor or viral infection and their proliferation, co-stimulatory ligand-expressing T cells turn the tumor or viral infection site into a highly conductive environment for a wide range of immune cells involved in the physiological anti-tumor or antiviral response (tumor infiltrating lymphocytes, NK-, NKT- cells, dendritic cells, and macrophages).

[0123] In other embodiments, the disclosure provides methods for treating subjects with a pathogen infection (e.g., viral infection, bacterial infection, fungal infection, parasite infection, or protozoal infection. The disclosure is particularly useful for enhancing an immune response in an immunocompromised subject. Exemplary viral infections susceptible to treatment using a method of the disclosure include, but are not limited to, Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus infections.

[0124] Accordingly, the disclosure provides a method of treating or preventing a pathogen infection in a subject, the method comprising administering an effective amount of an immunoresponsive cell as described herein.

Kits

[0125] The disclosure provides kits for the treatment or prevention of a neoplasia, pathogen infection, immune disorder or allogeneic transplant. In one embodiment, the kit includes a therapeutic or prophylactic composition containing an effective amount of an immunoresponsive cell comprising one or more co-stimulatory ligands in unit dosage form.

In some embodiments, the kit comprises a sterile container which contains a therapeutic or prophylactic vaccine; such containers can be boxes, ampules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments.

[0126] If desired the immunoresponsive cell is provided together with instructions for administering the cell to a subject having or at risk of developing a neoplasia, pathogen infection, immune disorder or allogeneic transplant. The instructions will generally include information about the use of the composition for the treatment or prevention of neoplasia, pathogen infection, immune disorder or allogeneic transplant. In other embodiments, the instructions include at least one of the following: description of the therapeutic agent; dosage schedule and administration for treatment or prevention of a neoplasia, pathogen infection, immune disorder or allogeneic transplant or symptoms thereof; precautions; warnings; indications; counter-indications; overdosage information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container.

Recombinant methods are well known in the art. The practice of the disclosure employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", second edition (Sambrook et al., 1989); "Oligonucleotide Synthesis" (Gait, ed., 1984); "Animal Cell Culture" (Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Handbook of Experimental Immunology" (Wei & Blackwell, eds.); "Gene Transfer Vectors for Mammalian Cells" (Miller & Calos, eds., 1987); "Current Protocols in Molecular Biology" (Ausubel et al., eds., 1987); "PCR: The Polymerase Chain Reaction", (Mullis et al., eds., 1994); and "Current Protocols in Immunology" (Coligan et al., eds., 1991). These techniques are applicable to the production of the polynucleotides and polypeptides, and, as such, can be considered in making and practicing the invention. Particularly useful techniques for are discussed in the sections that follow.

EXAMPLES

[0127] The following examples are provided as a further description of the invention, and to illustrate but not limit the invention.

Example 1

T cells co-expressing CD80 and 4-1BBL elicit robust proliferative responses after cyclic stimulations through their endogenous T cell receptor or through a chimeric antigen receptor without antigen presenting cell (APC)-provided costimulation

[0128] To assess whether constitutive expression of costimulatory ligands in T cells could substitute for APC-mediated costimulation, the T cell responses of human primary T cells were first investigated in three experimental systems. Using anti-CD3 (OKT3)-mediated T cell activation, the expansion of peripheral blood T lymphocytes transduced with CD80 and 4-1BBL was quantified (Figure 1a), which were compared to T cells transduced with either ligand alone or none. Recurrent T cell receptor (TCR)-stimulation alone in the absence of costimulatory ligands failed to expand T cells and rapidly induced a decline in T cell number following the first restimulation (Figure 1b). In sharp contrast, OKT3-stimulated CD80⁺ 4-1BBL⁺ T cells triggered a mean 237-fold greater proliferation over 21 days. In comparison, T cells transduced with either ligand alone exhibited a mean 8.1-fold reduced proliferation ($p < 0.0001$). Based on these observations, the concept of T-cell mediated costimulation was extended to two clinically relevant applications of adoptive T cell therapy, using cytomegalovirus (CMV)-specific memory donor T cells and differentiation antigen-specific, genetically redirected autologous T cells. CMV pp65-specific T lymphocytes briefly expanded on HLA-A*0201⁺pp65⁺ artificial APCs²⁴ were readily transduced with CD80 and 4-1BBL (Figure 1c). Following exposure to the HLA-A*0201⁺ pp65-transduced colonic tumor cell line Caco-2, T cells equipped with the costimulatory ligand pair CD80 and 4-1BBL exhibited a significantly greater (209-fold, $p < 0.0001$) expansion, compared to a continuously declining T cell number in the control groups (Figure 1d).

[0129] To rapidly generate tumor-reactive human T lymphocytes, peripheral blood T cells were retrovirally transduced with the chimeric antigen receptor Pz1 (Gade et al., Cancer Res. 65:9080-9088, 2005), a non-HLA-restricted antigen receptor specific for the tumor antigen PSMA. The Pz1 receptor comprises a PSMA-binding single chain antibody fragment fused to the human CD3 ζ signaling domain and is analogous in structure to other chimeric antigen receptors currently in use in clinical trials. Pz1⁺ T cells coexpressing CD80 and 4-1BBL (Figure 1e) mounted a robust proliferative response following three weekly stimulations with PSMA⁺, CD80⁻, CD86⁻, 4-1BBL⁻ LNCaP cells (mean 1042-fold enrichment, Figure 1f, upper panel). This expansion was 9-fold, greater ($p < 0.0001$) than that obtained when expressing CD80 and 4-1BBL in the tumor cells rather than in the T cells (Figure 1f, lower panel). Further analyses documented the greater induction of IL-2 and IFN- γ by exposure to PSMA in T cells coexpressing CD80 and 4-1BBL, as well as their greater antigen-specific cytolytic potential and decreased susceptibility to apoptosis, when compared to conventionally stimulated T lymphocytes (data not shown). In parallel studies, other members of TNF ligand family⁵, including OX40L, CD27L (CD70) or CD30L, with or without CD80, were investigated and the combination of CD80 and 4-1BBL was found to be the most potent (data not shown).

[0130] In aggregate, these *in vitro* studies demonstrate the ability of T lymphocytes coexpressing CD80 and 4-1BBL cells to strongly potentiate suboptimal TCR-activation and, furthermore, to substitute for the lack of costimulation provided by the APC.

Example 2

T cells co-expressing CD80 and 4-1BBL eradicate established, systemic tumors.

[0131] To investigate the potency of our CD80⁺4-1BBL⁺ T cells *in vivo*, a model of multifocal, established prostate cancer utilizing PSMA⁺ PC-3 tumor cells (Gong et al., *Neoplasia* 1:123-7, 1999) was developed. Using dual-modality bioluminescence and magnetic resonance imaging, tumors were visualized four weeks after intravenous inoculation, prior to initiating adoptive T cell therapy. The lungs, cervical lymph nodes, bone marrow, and liver were identified as the main sites of disease (Figure 2a). In this model, animals were treated four weeks after tumor inoculation with a single intravenous infusion of 8×10^6 PSMA-targeted T cells, expressing either CD80, 4-1BBL, both, or neither.

[0132] In control mice treated with 8×10^6 CD 19-targeted T cells, which, like untransduced T cells, fail to lyse PSMA⁺ tumor targets *in vitro* tumor burden steadily progressed until mice had to be sacrificed (Figure 2b,c). Treatment with Pz1⁺ T cells resulted in a short-term reduction of tumor burden, followed by terminal tumor progression (Figure 2b), yielding a modest 12-day survival advantage ($p = 0.0001$, Figure 2c). Constitutive expression of either CD80 or 4-1BBL alone in PSMA-targeted T cells only marginally augmented this therapeutic response, extending the median survival to 63 and 66 days, respectively ($p = 0.077$, $p = 0.056$, respectively). T cells coexpressing CD80 and 4-1BBL induced major responses and reduced the tumor burden 3.3-fold ($p = 0.0028$) four days after adoptive T cell transfer. A 1375-fold reduction was obtained after sixteen days, relative to the Pz1⁺ T cell treatment group ($p = 0.0002$). Seven of ten treated animals remained tumor-free 200 days after infusion of Pz1⁺CD80⁺4-1BBL⁺ T lymphocytes, and none in all other treatment or control groups (Figure 2c). The three mice that failed therapy initially showed marked tumor regression before relapsing (Figure 2b) and survived for 100 days (Figure 2c).

Example 3

In vivo T cell expansion is robust and antigen-specific.

[0133] To track and quantify *in vivo* T-cell migration and accumulation in relation to tumor localization and tumor burden, adoptively transferred T cells were additionally marked with Click Beetle Red-luciferase (Ponomarev et al., *Eur J Nucl Med Mol Imaging*. 2004 May;31(5):740-51) (CBR-luc, Figure 3a). Serial imaging of mice treated with Pz1⁺ T cells showed a progressive increase in signal that reached a peak four days after T cell injection (Figure 3b). A low-level signal remained detectable up to day 18. In the case of

Pz1⁺CD80⁺4-1BBL⁺ T cells, peak signal was detected on day 8 (41-fold higher nadir photon count than Pz1⁺ T cells, $p = 0.0009$), which was followed by a gradual signal decline, although bioluminescence could still be detected until day 100 in some animals (Figure 3b). Importantly, the effect of CD80 and 4-1BBL co-expression was abrogated in 19z1⁺ T lymphocytes (Figure 3b), consistent with the need for antigen stimulation for expansion to occur. Flow cytometric analyses and T cell counts of lung single cell suspensions in three treated mice per group determined the actual T cell number to be highly concordant with acquired bioluminescent signal intensities (Figure 3c,d). These studies, therefore, demonstrate that T cells coexpressing CD80 and 4-1BBL expand in an antigen-dependent manner before eventually entering a contraction phase resulting in substantial, if not complete, T cell clearance.

Example 4

Coalescence of the costimulatory ligands CD80 and 4-1BBL with their respective receptors CD28 and 4-1BB in the immunological synapse precedes functional T cell auto-costimulation.

[0134] To address whether constitutively expressed costimulatory ligands activate T cells in *cis*, the question of whether these ligands colocalize with their cognate receptors during T-cell activation was first examined. Both CD28 and 4-1BB amplify T cell receptor (TCR) signaling after recruitment into central membrane compartments of the immunological synapse³¹⁻³³ Therefore, colocalization of CD80 and 4-1BBL to the T-cell tumor cell contact area is likely a prerequisite for auto-costimulation. To visualize 4-1BBL distribution by confocal microscopy, the cytoplasmic domain was fused to monomeric dsRed and this protein was coexpressed with CD80 in Pz1-transduced CD8⁺ T lymphocytes. Immediately prior to admixing with unmodified CD28⁺, 4-1BB⁺ LNCaP cells, T cells were labeled with fluorescein isothiocyanate-conjugated cholera toxin β subunit (FITC-CTB) to visualize lipid raft clusters after synapse formation. Following LNCaP tumor engagement, 4-1BBL and CD80, as well as 4-1BB and CD28, mobilized into cholera toxin-FITC-positive T-cell-tumor cell contact areas (Figure 4a,b). To investigate the functional consequence of this interaction, the colocalization of Granzyme B (GRB. Defined, albeit sparse, GRB condensation was localized near the contact zone in clusters of Pz1-expressing CD8⁺ T cells and their cognate tumor targets (Figure 4c,d)) was examined and quantified. The median GRB recruitment to T-cell tumor cell junctions was amplified 2.2-fold in CD80⁺4-1BBL⁺ T cells ($p = 0.0001$). To confirm that these observed differences were indeed a result of 4-1BB engagement with 4-1BBL within the same cell, 4-1BB expression was knocked down by stable coexpression of 4-1BBL and 4-1BB-specific shRNA (data not shown). Knock-down of 4-1BB significantly diminished GRB density near the synapse (2.02-fold reduction, $p < 0.0001$), despite the presence of 4-1BBL at the T-cell junction in all imaged cell clusters. These findings strongly suggest that the costimulatory ligand functionally engages its receptor on the same T-cell surface after antigen-induced coalescence in the immunological synapse.

[0135] To further demonstrate auto-costimulation, a single cell assay was devised in which the endogenous 4-1BB/4-1BBL interaction could be analyzed following the earliest expression of 4-1BBL in recently transduced T cells. To achieve this, firefly luciferase was first expressed under the transcriptional control of NF- κ B, a key downstream effector of 4-1BB signaling, in a clone of CD3 $^+$ CD28 $^+$ 4-1BB $^+$ Jurkat T cells (JNL). To preclude NF- κ B induction by sporadic bystander costimulation, JNL cells were exposed to retroviral vectors encoding 4-1BB, 4-1BBL and CD80, and subcloned by limiting dilution 4 hours thereafter, well before the first detectable surface gene expression (data not shown), and immediately stimulated by plate-bound OKT3. As illustrated in Figure 4e, isolated CD80 $^+$ CD28 $^+$ 4-1BBL $^+$ 4-1BB $^+$ T cells markedly up-regulated NF- κ B, in contrast to JNL cells transduced with control vector. The luminescence signal was acquired in 36 wells containing a single, multiply transduced cell as described in Methods. Coexpression of the two ligand pairs increased NF- κ B-dependent signal 3.8-fold (relative to control JNL; $p < 0.0001$; Figure 4f). Since at any point of this assay the sole source of costimulatory ligands was the isolated T cell's own surface, this NF- κ B upregulation could only reflect the impact of auto-costimulation.

Example 5

In vitro *trans*-costimulation of antigen-specific bystander T cells

[0136] The constitutive expression of costimulatory ligands may also enable genetically modified T cells to costimulate T cells in *trans*. Notably, three-cell clusters comprised of CD80 $^+$ 4-1BBL $^+$ and CD80 $^-$ 4-1BBL $^-$ T cells were occasionally noted in the confocal studies shown in Fig. 4 (Figure 5a). To provide functional evidence for *trans*-costimulation, a coculture system was devised in which PSMA-specific T cells expressing CD80 and 4-1BBL were admixed with carboxyfluoroscein succinimidyl ester (CFSE)-labeled T cells that were not transduced with CD80 or 4-1BBL. It was found that these CFSE-labeled, CMV pp65-specific T cells were effectively costimulated by autologous, bystander Pz1 $^+$ CD80 $^+$ 4-1BBL $^+$ T lymphocytes (Figure 5b,c). Physical contact between the two T cell populations was a prerequisite for the CMV-reactive cytotoxic T lymphocytes to expand, as their separation by a transwell membrane greatly reduced the strong induction of GRB and the robust T-cell expansion mounted in pp65 responder cells, consistent with a cell contact-dependent mechanism (Figure 5d).

Example 6

In vivo trans-costimulation of tumor-infiltrating T cells

[0137] These observations prompted us to investigate whether T cell-mediated *trans*-costimulation is operative *in vivo*. To this end, two previously described animal models were combined: the RM1-PSMA tumor model in which tumors are confined to the lungs (Gade et al., Cancer Res. 65:9080-9088, 2005), and the Raji tumor model, in which tumor cells selectively colonize bone marrow (Brentjens et al., Nat Med. 2003 Mar;9(3):279-86). All mice were treated with Pz1-transduced T cells, which also expressed CBR-luc (Figure 3). Luciferase-negative Pz1⁺ T cells, either expressing or lacking CD80 and 4-1BBL, were subsequently infused into the animals. The expression of Gaussia-luciferase (Gau-Luc) in the tumor cells, and of CBR-luc in Pz1⁺ T cells, allowed the use of dual bioluminescence imaging to simultaneously monitor tumor progression and the spatial and temporal accumulation of T cells. Comparable tissue distributions and bioluminescent signals of CBR-luc⁺ lymphocytes were observed six hours (day 0) after adoptive transfer in all treatment groups (Figure 6a). On day 2, the effect of Pz1⁺CD80⁺4-1BBL⁺ T cells on CBR-luc⁺Pz1⁺ T cells was still modest (median 1.2-fold, relative to Pz1⁺ T cells as bystander, p = 0.0947). The CBR-luc signal subsequently decayed in mice given control Pz1⁺ T cells (Figure 6a). In contrast, this photon count recorded over the thoracic area augmented 6.5-fold on average in mice given Pz1⁺CD80⁺4-1BBL⁺ T lymphocytes (p = 0.0122, Figure 6a,b). Importantly, this effect was selective for the PSMA-targeted T cells, since co-injected CBR-luc⁺19z1-transduced T cells, which infiltrated the established Raji tumors in both femurs, did not significantly enhance thoracic bioluminescent signal at any time point (median 1.5-fold increase on day 4, p = 0.0947, Figure 6a,b). Collectively, these *in vitro* and *in vivo* data indicate that CD80⁺4-1BBL⁺ T cells locally enhance T-cell responses by providing costimulation in *trans*.

[0138] The constitutive display of CD80 on T lymphocytes serves as a costimulatory ligand for CD28 but could also engage negative regulators of T cell expansion, such as CTLA-4 at high affinities (Hodi et al. Clin Cancer Res. 2007 Sep 15:13:5238-42). To bypass therapeutically undesirable T cell auto-inhibition mediated by T cell-expressed CD80, T lymphocytes were co-transduced with the chimeric antigen receptor P28z (Maher et al. Nature Biotechnology, Vol 20, Jan 2002, P 70-75) and expressed 4-1BBL on the T cell surface (Figure 7). The dual fusion receptor P28z contains both TCR and CD28 signaling moieties. Whereas signaling through P28z failed to invoke a sustained T cell expansion, the co-expression of 4-1BBL markedly enhanced the proliferative response more than 10-fold by day 14. The synergistic CD80-4-1BBL costimulatory signal, however, was still strongest when both full length ligands were expressed on the surface of Pz1-transduced T lymphocytes (approximately two-fold higher than P28Z + 4-1BBL, Figure 7). CD28 signals relayed by a CD28 signaling element fused into the chimeric antigen receptor construct can therefore endow Pz1⁺4-1BBL⁺T lymphocytes with enhanced proliferative properties, although it is not as robust a T cell proliferation than observed in Pz1⁺4-1BBL⁺CD80⁺ T cells.

[0139] A major goal of cancer immunotherapy is to provide safe and effective costimulation to tumor-reactive T lymphocytes. Using a genetic approach, it was demonstrated that T cells themselves provide potent costimulation. Constitutive expression of costimulatory ligands in human primary CMV-specific cells and PSMA-targeted T cells not only compensated for the absence of these ligands on APCs, but also induced a proliferative response surpassing that elicited by conventional APCs (Figure 1). The survey of a panel of costimulatory ligands indicated that CD80 and 4-1BBL provided the strongest T-cell activation under our experimental conditions, enabling robust T cell expansion following repeated weekly antigenic stimulation. CD80⁺4-1BBL⁺ T cells exhibited superior proliferation, cytokine secretion, in vitro survival and in vivo expansion and persistence, resulting in a 40- to 50-fold greater T cell biomass one week after infusion in tumor-bearing mice, when compared to PSMA-specific T cells that were not transduced with CD80 and 4-1BBL (Figure 3). Whereas animals treated with non-CD80/4-1BBL-transduced PSMA-specific T cells uniformly succumbed to disease, infusion of the same T cells rendered CD80⁺4-1BBL⁺ effectively rejected established, systemic PC-3 tumors in the majority of treated mice (Figure 2). These findings, obtained in a very challenging tumor model, underscore the biological activity and remarkable potency of constitutive, high-level expression of costimulatory ligands in T cells.

[0140] To investigate the mechanism underlying this potentiated response, the spatial distribution of CD80 and 4-1BBL and their receptors on T cells before and after tumor engagement were first examined. CD28 and 4-1BB have been shown to accumulate in close proximity to the TCR within the central supramolecular activation cluster of immunological synapses. Costimulatory ligands expressed by APCs, including CD80, CD40L and CD70, polarize to the synapse, supporting the notion that the immunological synapse provides an ordered contact area to facilitate and propagate costimulatory ligand-receptor interactions. Consistent with this model, T-cell-encoded CD80 and 4-1BBL polarized towards the contact zone between T cells and LNCaP tumor cells with, together with CD28 and 4-1BB (Figure 4). The functional consequence of this colocalization is illustrated by the accumulation of GRB, which is ferried towards the T cell-APC interface along microtubules following antigen contact in synapses containing 4-1BB⁺4-1BBL⁺ foci but not in T cells lacking 4-1BB (Figure 4). The immunological synapse is thus apparently well suited to orchestrate auto-costimulation.

[0141] Auto-costimulation was demonstrated in a single cell assay in which de novo expression of 4-1BB and 4-1BBL in recently transduced T cells activated NF- κ B without any possible contribution from a neighboring T cell. NF- κ B expression was visualized in real-time, after physical separation of single clones shortly after transduction, several hours before costimulatory ligand expression could be detected (Figure 4). Auto-costimulation may thus allow T lymphocytes to override the poor costimulatory capacity of suboptimal APCs, including tumor cells. Furthermore, constitutive cell surface expression of 4-1BBL in T cells ensures costimulation 'on demand' by ensuring that the ligand remains ready to engage transiently upregulated receptors such as 4-1BB. One might speculate that the physical interaction of 4-1BBL with its receptor may occur during early raft reorganization or between opposing plasma membrane folds at the center of the synapse. This effect appears to be self-regulated as CD80⁺4-1BBL⁺ T-cell expansion was gradually attenuated in response to repeated in vitro antigen exposure (Figure 1). In vivo T-cell

expansion was also self-limited (Figure 3), which may be due to tumor elimination that deprives the T cells of antigenic stimulation and attenuated costimulatory signaling.

[0142] Without wishing to be bound by theory, our studies also support trans-costimulation as a mechanism for enhancing immunity through high-level expression of CD80 and 4-1BBL in T lymphocytes. The proliferative and effector functions of T cells engaging APC-presented antigen in the absence of CD80 or 4-1BBL were thus amplified by *in vitro* coculture or *in vivo* coadministration of T cells displaying both ligands. *In vitro* cell mixing and transwell studies indicated that trans-costimulation required cell-cell contact (Figure 5), in agreement with *trans* effects reported for other cell types such as fibroblasts or bystander tumor cells. Trans-costimulation was also induced *in vivo*, in organ-specific fashion, as illustrated in mice bearing two different tumors segregating to different sites (Figure 6).

[0143] Trans-costimulation mediated by T cells paves the way for exciting new therapeutic approaches since costimulation may be delivered to neighboring T cells within the tumor microenvironment. This feature is especially valuable as dendritic cells frequently fail to fully upregulate costimulatory ligands within the tumor microenvironment. Although soluble factors such as interleukin-2 may contribute to T cell aid, the dependence on cell-cell contact effectively restricts trans-costimulation to other tumor-infiltrating T cells. Without wishing to be bound by theory, the effect of T cell competition on the *in vivo* frequency of trans-costimulation, *in trans* may ultimately broaden the anti-tumor immune response via the recruitment of a diverse population of endogenous tumor-infiltrating lymphocytes and thus help to prevent tumor antigen escape.

[0144] In summary, it was shown that CD80 and 4-1BBL expression in human T lymphocytes is a biologically efficacious means to circumvent the lack of conventional APC-mediated costimulation in the tumor microenvironment. This approach is applicable to T cells activated through a transduced antigen receptor as well as their endogenous TCR. T cell-mediated costimulation, whether *in auto*, *trans* or both, may thus be useful in a wide range of malignancies and infectious diseases that are being treated by adoptive T cell therapy.

Embodiments of the Invention

[0145] From the foregoing description, it will be apparent that variations and modifications may be made to the invention described herein to adopt it to various usages and conditions. Such embodiments are also within the scope of the following claims.

[0146] The recitation of a listing of elements in any definition of a variable herein includes definitions of that variable as any single element or combination (or subcombination) of listed elements. The recitation of an embodiment herein includes that embodiment as any single embodiment or in combination with any other embodiments or portions thereof. The disclosure may be further characterized by any of the following items:

1. An immunoresponsive cell comprising a receptor that binds an antigen and an exogenous co-stimulatory ligand.
2. The method of item 1 , wherein the co-stimulatory ligand is constitutively or inducibly expressed.
3. The method of item 1, wherein at least two co-stimulatory ligands are constitutively expressed.
4. The immunoresponsive cell of item 1, wherein the cell is selected from the group consisting of a T cell, a Natural Killer (NK) cell, a cytotoxic T lymphocyte (CTL), and a regulatory T cell.
5. The immunoresponsive cell of item 1, wherein the antigen is a tumor or pathogen antigen.
6. The immunoresponsive cell of item 3, wherein the antigen is selected from the group consisting of prostate-specific membrane antigen (PSMA), Carcinoembryonic Antigen (CEA), IL13Ralpha, her-2, CD19, NY-ESO-1, HIV-1 Gag, Lewis Y, Mart-1, gp100, tyrosinase, WT-1, hTERT, and mesothelin.
7. The immunoresponsive cell of item 1, wherein the cell expresses a recombinant or an endogenous antigen receptor that is Pz1 or P28z.
8. The immunoresponsive cell of item 1, wherein the co-stimulatory ligand is a tumor necrosis factor (TNF) ligand or an immunoglobulin (Ig) superfamily ligand.
9. The immunoresponsive cell of item 8, wherein the TNF ligand is selected from the group consisting of 4-1BBL, OX40L, CD70, LIGHT, and CD30L.
10. The immunoresponsive cell of item 8, wherein the Ig superfamily ligand is selected from the group consisting of CD80 and CD86.
11. The immunoresponsive cell of any one of items 1- 10, wherein the immunoresponsive cell expresses a TNF ligand and an Ig superfamily ligand.
12. The immunoresponsive cell of item 11, wherein the TNF ligand is 4-1BBL and the Ig superfamily ligand is CD80.
13. The immunoresponsive cell of any one of items 1-12, wherein an endogenous or recombinant antigen receptor or antigen presenting complex is constitutively expressed on the surface of the T cell.
14. The cell of any one of items 1-12, wherein the co-stimulatory ligands are constitutively expressed on the surface of the T cell.
15. The immunoresponsive cell of any one of items 1-12, wherein the co-stimulatory ligand is expressed in a retroviral vector.

16. A virus specific T cell expressing a vector encoding a polypeptide selected from the group consisting of CD80, 4-1BBL, OX40L, CD70 and CD30L.
17. The virus specific T cell of item 16, wherein the T cell recognizes a virus selected from the group consisting of Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus antigens.
18. A tumor antigen-specific T cell expressing a vector encoding a polypeptide selected from the group consisting of CD80, 4-1BBL, OX40L, CD70 and CD30L.
19. The tumor antigen- specific T cell of item 15, wherein the cell expresses CD80 and 4-1BBL.
20. The tumor antigen- specific T cell of item 15, wherein the vector is a retroviral vector.
21. A method of modulating an immune response in a subject, the method comprising administering an effective amount of an immunoresponsive cell of any one of items 1-17.
22. The method of item 21, wherein the method increases or reduces an immune response.
23. The method of item 21, wherein the method increases self-tolerance or increases tolerance to an organ transplant.
24. A method of treating or preventing a neoplasia in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds a tumor antigen and a vector encoding a co-stimulatory ligand.
25. The method of item 18, wherein the neoplasia is selected from the group consisting of prostate cancer, colon cancer, breast cancer, and glioblastoma.
26. The method of item 18, wherein the tumor antigen is prostate-specific membrane antigen, CD 19, NY-ESO-1, WT-1, hTERT and mesothelin.
27. A method of enforcing tolerance in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds an antigen and a vector encoding a co-stimulatory ligand.
28. The method of item 27, wherein the method prevents or reduces an autoimmune disease or a disease associated with allogeneic transplantation.
29. A method of treating or preventing a pathogen infection in a subject, the method comprising administering an effective amount of an immunoresponsive cell comprising a receptor that binds a viral antigen and a vector encoding a co-stimulatory ligand.

30. The method of item 29, wherein the pathogen is a virus, bacteria, fungus, protozoa or parasite.
31. The method of item 29, wherein the virus is selected from the group consisting of Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), and influenza virus.
32. The method of any one of items 21-30, wherein the cell is a T cell, a Natural Killer (NK) cell, or a cytotoxic T lymphocyte (CTL), whether collected from an autologous or allogeneic donor, or generated in vitro from an engineered progenitor or stem cell.
33. The method of any one of items 21-30, wherein the cell expresses a recombinant or an endogenous receptor for the antigen.
34. The method of any one of items 21-30, wherein the co-stimulatory ligand is a tumor necrosis factor (TNF) ligand or an immunoglobulin (Ig) superfamily ligand.
35. The method of any one of items 21-31, wherein the TNF ligand is selected from the group consisting of 4-1BBL, OX40L, CD70, LIGHT, and CD30L.
36. The method of any one of items 21-31, wherein the Ig superfamily ligand is selected from the group consisting of CD80 and CD86.
37. The method of any one of items 21-31, wherein the immunoresponsive cell express a TNF ligand and an Ig superfamily ligand.
38. The method of any one of items 21-31, wherein the TNF ligand is 4-1BBL and the Ig superfamily ligand is CD80.
39. The method of any one of items 21-31, wherein a recombinant or an endogenous antigen receptor or an antigen presenting complex is constitutively expressed on the surface of the T cell.
40. The method of any one of items 21-31, wherein the co-stimulatory ligands are constitutively expressed on the surface of the T cell.
41. The method of any one of items 21-31, wherein the vector is a retroviral vector.
42. A method for producing an antigen-specific immunoresponsive cell, the method comprising introducing into the immunoresponsive cell a nucleic acid sequence that encodes a chimeric antigen receptor, wherein the chimeric antigen receptor comprises an antigen-binding domain coupled to an intracellular signaling domain that activates an immunoresponsive cell.
43. The method of item 42, wherein the immunoresponsive cells is a T cell, CTL, or NK cell.


44. The method of item 42, wherein the antigen-binding domain is a tumor antigen-binding domain.
45. The method of item 42, wherein the tumor antigen is prostate specific membrane antigen (PSMA).
46. The method of item 42, wherein the intracellular signaling domain activates a T cell, CTL cell, or NK cell.
47. The method of item 42, wherein the intracellular signaling domain is the [zeta]-chain signaling domain.
48. The method of item 42, wherein the one costimulatory signal is provided through an engineered antigen receptor and the other by overexpressing a costimulatory ligand.
49. A method of treating a neoplasia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a T cell comprising a tumor antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from the group consisting of: a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating cancer in the subject.
50. A method of treating a neoplasia in a subject, the method comprising administering to the subject a therapeutically effective amount of a Natural Killer (NK) cell comprising a tumor antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from the group consisting of: a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating cancer in the subject.
51. The method of items 48-50, wherein the TNF ligand is selected from the group consisting of: 4-1BBL, OX40L, CD70, and CD30L, and LIGHT.
52. The method of items 48-50, wherein the Ig superfamily ligand is selected from the group consisting of: CD80 and CD86.
53. The method of items 48-50, wherein the at least two co-stimulatory ligands are a TNF ligand and an Ig superfamily ligand.
54. The method of item 52, wherein the TNF ligand is 4-1BBL and the Ig superfamily ligand is CD80.
55. The method of items 48-50, wherein the antigen recognition complex is constitutively expressed on the surface of the cell.

56. The method of items 48-50, wherein the tumor antigen is prostate specific membrane antigen (PSMA).
57. The method of items 48-50, wherein the at least two co-stimulatory ligands are constitutively expressed on the surface of the cell.
58. A method of treating an infectious disease in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a T cell comprising a receptor specific for a viral antigen and an antigen presenting complex comprising at least two co-stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from the group consisting of: a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating the infectious disease in the subject.
59. A method of treating an infectious disease in a subject, comprising administering to the subject a therapeutically effective amount of a Natural Killer (NK.) cell comprising a receptor specific for a viral antigen and an antigen presenting complex comprising at least two co- stimulatory ligands, wherein at least one of the two co-stimulatory ligands is selected from the group consisting of: a tumor necrosis factor (TNF) ligand and an immunoglobulin (Ig) superfamily ligand and combinations thereof, thereby treating the infectious disease in the subject.
60. The method of item 58 or 59, wherein the subject is an immunocompromised subject.
61. The method of item 57 or 58, wherein the TNF ligand is selected from the group consisting of: 4-1BBL, OX40L, CD70, LIGHT and CD30L.
62. The method of item 57 or 58, wherein the Ig superfamily ligand is selected from the group consisting of: CD80 and CD86.
63. The method of item 57 or 58, wherein the at least two co-stimulatory ligands are a TNF ligand and an Ig superfamily ligand.
64. The method of item 57 or 58, wherein the TNF ligand is 4-1BBL and the Ig superfamily ligand is CD80.
65. The method of item 57 or 58, wherein the antigen recognition complex is constitutively expressed on the surface of the cell.
66. The method of item 57 or 58, wherein the viral antigen is an antigen specific for Cytomegalovirus (CMV), Epstein Barr Virus (EBV), Human Immunodeficiency Virus (HIV), or influenza virus .
67. The method of item 57 or 58, wherein the at least two co-stimulatory ligands are constitutively expressed on the surface of the cell.

68. A pharmaceutical composition comprising an effective amount of an immunoresponsive cell of any one of items 1-20 in a pharmaceutically acceptable excipient.
69. A pharmaceutical composition for the treatment of a neoplasia comprising an effective amount of a tumor antigen-specific T cell of any one of item 15-17 in a pharmaceutically acceptable excipient.
70. A pharmaceutical composition for the treatment of a pathogen infection comprising an effective amount of a viral-specific T cell of item 14 in a pharmaceutically acceptable excipient.
71. The pharmaceutical composition of any one of items 67-69, further comprising a cytokine selected from the group consisting of IL-2, IL-3, IL-6, IL-11, IL-12, IL7, IL15, IL21, granulocyte macrophage colony stimulating factor, alpha, beta or gamma interferon and erythropoietin.
72. A kit for treatment of a neoplasia, pathogen infection, an autoimmune disorder, or an allogeneic transplant, the kit comprising an immunoresponsive cell comprising a receptor that binds an antigen and an exogenous co-stimulatory ligand.
73. The kit of item 71 , wherein the kit further comprises written instructions for using said cell for the treatment of a subject having a neoplasia, a pathogen infection, an autoimmune disorder, or an allogeneic transplant.
74. The method of any one of items 21 -67, further comprising the step of an immunoresponsive obtaining the immunoresponsive cell or co-stimulatory ligand.

Patentkrav

- 5 1. T-celle, omfattende en receptor, der binder et tumorantigen og exogene co-stimulatoriske ligander 4-1BBL og CD80 til anvendelse ved behandling eller forebyggelse af en neopla*si* i et individ.
- 10 2. T-celle ifølge krav 1, hvor neoplasien er udvalgt fra gruppen bestående af prostatacancer, coloncancer, brystcancer og glioblastom.
- 15 3. T-celle ifølge krav 1, hvor tumorantigenet er prostataspecifikt membran-antigen, CD19, NY-ESO-1, WT-1, hTERT og mesothelin.
- 20 4. Farmaceutisk sammensætning til anvendelse ved behandling af en neopla*si*, omfattende en virksom mængde af en tumorantigen-specifik T-celle, omfattende en receptor, der binder et tumorantigen, og co-stimulatoriske ligander 4-1BBL og CD80 i en farmaceutisk acceptabel excipiens.
- 25 5. Farmaceutisk sammensætning til anvendelse ifølge krav 4, yderligere omfattende et cytokin, der er udvalgt fra gruppen bestående af IL-2, IL-3, IL-6, IL-11, IL-12, IL7, IL15, IL21, granulocyt-makrofag-kolonistimulerende faktor, alfa-, beta- eller gammainterferon og erythropoietin.






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

NCBI Protein Sign In | Regis

Search Protein for

Limits Preview/Index History Clipboard Details

Display GenPept Show 5 Send to

Range: from to Features: CDD

1: NP_005182. Reports CD80 antigen prec...[gi:4885123] BLINK, Conserved Domains, Links

Comment Features Sequence

LOCUS NP_005182 288 aa linear PRI 24-FEB-2008
 DEFINITION CD80 antigen precursor (Homo sapiens).
 ACCESSION NP_005182
 VERSION NP_005182.1 GI:4885123
 DBSOURCE REFSEQ: accession NM_005191.3
 KEYWORDS
 SOURCE Homo sapiens (human)
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.
 1 (residues 1 to 288)
 REFERENCE
 AUTHORS Nolan,A., Weiden,M., Kelly,A., Hoshino,Y., Hoshino,S., Mehta,N. and Gold,J.A.
 TITLE CD40 and CD80/86 act synergistically to regulate inflammation and mortality in polymicrobial sepsis
 JOURNAL Am. J. Respir. Crit. Care Med. 177 (3), 301-308 (2008)
 PUBMED 17989345
 REMARK GeneRIF: CD80 expression was up-regulated on the circulating monocytes in septic subjects on Day 1
 2 (residues 1 to 288)
 REFERENCE
 AUTHORS Stephan,M.T., Ponomarev,V., Brentjens,R.J., Chang,A.H., Dobrenkov,K.V., Heller,G. and Sadelain,M.
 TITLE T cell-encoded CD80 and 4-1BBL induce auto- and transcostimulation, resulting in potent tumor rejection
 JOURNAL Nat. Med. 13 (12), 1440-1449 (2007)
 PUBMED 18026115
 REMARK GeneRIF: CD80 and 4-1BBL induce auto- and transcostimulation in tumor cells
 3 (residues 1 to 288)
 REFERENCE
 AUTHORS Habib-Agahi,M., Phan,T.T. and Searle,P.F.
 TITLE Co-stimulation with 4-1BB ligand allows extended T-cell proliferation, synergizes with CD80/CD86 and can reactivate anergic T cells
 JOURNAL Int. Immunol. 19 (12), 1383-1394 (2007)
 PUBMED 17977894
 REMARK GeneRIF: T cells that had become non-responsive to anti-CD3 could be reactivated to proliferate when costimulated with 4-1BB, either alone or combined with CD80/CD86.
 4 (residues 1 to 288)
 REFERENCE
 AUTHORS Kakoulidou,M., Giscombe,R., Zhao,X., Lefvert,A.K. and Wang,X.
 TITLE Human Soluble CD80 is generated by alternative splicing, and

recombinant soluble CD80 binds to CD28 and CD152 influencing T-cell activation
 Scand. J. Immunol. 66 (5), 529-537 (2007)
 17953528
 REMARK GeneRIF: sCD80 in human serum adds a new member to the family of soluble receptors, implying a network of soluble costimulatory factors with functional relevance
 5 (residues 1 to 288)
 AUTHORS Dopheide,J.F., Sester,U., Schlitt,A., Horstick,G., Rupprecht,H.J., Munzel,T. and Blankenberg,S.
 TITLE Monocyte-derived dendritic cells of patients with coronary artery disease show an increased expression of costimulatory molecules CD40, CD80 and CD86 in vitro
 Coron. Artery Dis. 18 (7), 523-531 (2007)
 17925605
 REMARK GeneRIF: CD40, CD80 and CD86 are upregulated in cultured monocyte-derived dendritic cells of patients with coronary artery disease
 6 (residues 1 to 288)
 AUTHORS Weiskirchen,R., Pino,J.D., Macalma,T., Bister,K. and Beckerle,M.C.
 TITLE The cysteine-rich protein family of highly related LIM domain proteins
 J. Biol. Chem. 270 (48), 28946-28954 (1995)
 7499425
 REFERENCE 7 (residues 1 to 288)
 AUTHORS Freeman,G.J., Disteche,C.M., Gribben,J.G., Adler,D.A., Freedman,A.S., Dougery,J. and Nadler,L.M.
 TITLE The gene for B7, a costimulatory signal for T-cell activation, maps to chromosomal region 3q13.3-3q21
 Blood 79 (2), 489-494 (1992)
 1370389
 REFERENCE 8 (residues 1 to 288)
 AUTHORS Selvakumar,A., Mohanraj,B.K., Eddy,R.L., Shows,T.B., White,P.C. and Dupont,B.
 TITLE Genomic organization and chromosomal location of the human gene encoding the B-lymphocyte activation antigen B7
 JOURNAL Immunogenetics 36 (3), 175-181 (1992)
 1377173
 REFERENCE 9 (residues 1 to 288)
 AUTHORS Freeman,G.J., Gray,G.S., Gimmi,C.D., Lombard,D.B., Zhou,L.J., White,M., Fingerroth,J.D., Gribben,J.G. and Nadler,L.M.
 TITLE Structure, expression, and T cell costimulatory activity of the murine homologue of the human B lymphocyte activation antigen B7
 J. Exp. Med. 174 (3), 625-631 (1991)
 1714935
 REFERENCE 10 (residues 1 to 288)
 AUTHORS Freeman,G.J., Freedman,A.S., Segil,J.M., Lee,G., Whitman,J.F. and Nadler,L.M.
 TITLE B7, a new member of the Ig superfamily with unique expression on activated and neoplastic B cells
 J. Immunol. 143 (8), 2714-2722 (1989)
 2794510
 COMMENT VALIDATED_REFSEQ: This record has undergone validation or preliminary review. The reference sequence was derived from BP225173.1, BC042665.1 and AC073352.22.
 Summary: The B-lymphocyte activation antigen B7-1 (formerly referred to as B7) provides regulatory signals for T lymphocytes as a consequence of binding to the CD28 (MIM 186760) and CTLA4 (MIM 123890) ligands of T cells. [supplied by OMIM].

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

FEATURES Location/Qualifiers

source 1..288
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="3"
 /map="3q13.3-q21"

Protein 1..288
 /product="CD80 antigen precursor"
 /note="costimulatory molecule variant IgV-CD80; CD80 antigen (CD28 antigen ligand 1, B7-1 antigen); costimulatory factor CD80; B-lymphocyte activation antigen B7"

sig_peptide 1..34
 /calculated_mol_wt=3754

mat_peptide 35..288
 /product="CD80 antigen"
 /calculated_mol_wt=29312

Region 37..139
 /region_name="IG"
 /note="Immunoglobulin; smart00409"
 /db_xref="CDD:47718"

Region 37..135
 /region_name="V-set"
 /note="Immunoglobulin V-set domain. This domain is found in antibodies as well as neural protein P0 and CTL4 amongst others; pfam07686"
 /db_xref="CDD:87333"

Region 143..227
 /region_name="C2-set_2"
 /note="CD80-like C2-set immunoglobulin domain. These domains belong to the immunoglobulin superfamily; pfam08205"
 /db_xref="CDD:71639"

CDS 1..288
 /gene="CD80"
 /coded_by="NM_005191.3:396..1262"
 /db_xref="CCDS:CCDS2989.1"
 /db_xref="GeneID:941"
 /db_xref="HGNC:1700"
 /db_xref="HPRD:00202"
 /db_xref="MIM:112203"

ORIGIN

1 mghtrrqgts pskcpylnff qlvlaglsh fcsgvihvtk evkevatlsc ghnvsveela
 61 qtriywqkek kmvltnmsgd mniwpeyknr tifdittnls ivilalrpsd egtyecvvlk
 121 yekdafkreh laevtlsvka dfptpsisdf eiptsnirri icstsgggfpe phlswlenge
 181 elnainttvs qdpetelyav sskldfmmtt nhsfmcliky ghrlrvnqtfn wnttkqehfp
 241 dnllpswait lisvngifvi cclycfaapr crerrrnerl xresvrpv

//

NCBI Protein Search My NCBI Sign In Register

Search Protein for

Limits Preview/Index History Clipboard Details

Display GenPept Show 5 Send to

Range: from begin to end Features: CDD Refresh

1: P41273. Reports Tumor necrosis fa...[gi:728739]

BLINK, Conserved Domains, Links

Comment	Features	Sequence
LOCUS	P41273	254 aa linear PRI 04-DEC-2007
DEFINITION	Tumor necrosis factor ligand superfamily member 9 (4-1BB ligand)	(4-1BBL).
ACCESSION	P41273	
VERSION	P41273.1	GI:728739
DBSOURCE	swissprot: locus TNFL9_HUMAN, accession P41273;	class: standard.
		extra accessions:Q2M3S2
		created: Feb 1, 1995.
		sequence updated: Feb 1, 1995.
		annotation updated: Dec 4, 2007.
		xrefs: U03398.1, AA053134.1, BC104805.1, AA104806.1, BC104807.1, AA104808.1, I38427
		xrefs (non-sequence databases): RefSeq:NP_003802.1, UniGene:Hs_1524, DIP:DIP:3020N, Ensembl:ENSG00000125657, GeneID:8744, KEGG:hsa:8744, H-InvDB:HIX0040144, HGNC:11939, MIM:606182, PharmGKB:PA36629, ArrayExpress:P41273, CleanEx:HS_TNFSF9, GermOnline:ENSG00000125657, GO:0005102, GO:0006915, GO:0008283, GO:0007267, GO:0007165, InterPro:PF00229, SMART:SM00207, PROSITE:PS00251, PROSITE:PS50049
KEYWORDS	Cytokine; Membrane; Polymorphism; Signal-anchor; Transmembrane.	
SOURCE	Homo sapiens (human)	
ORGANISM	Homo sapiens	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.
REFERENCE	1 (residues 1 to 254)	
AUTHORS	Alderson,M.R., Smith,C.A., Tough,T.W., Davis-Smith,T., Armitage,R.J., Falk,B., Roux,E., Baker,E., Sutherland,G.R., Din,W.S. and Goodwin,R.G.	
TITLE	Molecular and biological characterization of human 4-1BB and its ligand	
JOURNAL	Eur. J. Immunol. 24 (9), 2219-2227 (1994)	
PUBMED	8088337	
REMARK	NUCLEOTIDE SEQUENCE [mRNA].	
REFERENCE	2 (residues 1 to 254)	
AUTHORS	Gerhard,D.S., Wagner,L., Feingold,E.A., Shenmen,C.M., Grouse,L.H., Schuler,G., Klein,S.L., Old,S., Rasooly,R., Good,P., Guyer,M., Peck,A.M., Derge,J.G., Lipman,D., Collins,F.S., Jang,W., Sherry,S., Feolo,M., Misquitta,L., Lee,E., Rotmistrovsky,K., Greenhut,S.F., Schaefer,C.F., Buetow,K., Bonner,T.I., Haussler,D., Kent,J., Kiekhaus,M., Furey,T., Brent,M., Prange,C., Schreiber,K..	


Shapiro, N., Bhat, N.K., Hopkins, R.F., Hsieh, F., Driscoll, T., Soares, M.B., Casavant, T.L., Scheetz, T.E., Brown-Stein, M.J., Usdin, T.B., Toshiyuki, S., Carninci, P., Piao, Y., Dudekula, D.B., Ko, M.S., Kawakami, K., Suzuki, Y., Sugano, S., Gruber, C.E., Smith, M.R., Simmons, B., Moore, T., Waterman, R., Johnson, S.L., Ruan, Y., Wei, C.L., Mathavan, S., Gunaratne, P.H., Wu, J., Garcia, A.M., Hulyk, S.W., Fuh, E., Yuan, Y., Snead, A., Kowis, C., Hodgson, A., Muzny, D.M., McPherson, J., Gibbs, R.A., Fahey, J., Helton, E., Ketteman, M., Madan, A., Rodrigues, S., Sanchez, A., Whiting, M., Madari, A., Young, A.C., Wetherby, K.D., Granite, S.J., Kwong, P.N., Brinkley, C.P., Pearson, R.L., Bouffard, G.G., Blakesly, R.W., Green, E.D., Dickson, M.C., Rodriguez, A.C., Grimwood, J., Schmutz, J., Myers, R.M., Butterfield, Y.S., Griffith, M., Griffith, O.L., Krzywinski, M.I., Liao, N., Morin, R., Palmquist, D., Petrescu, A.S., Skalska, U., Smailus, D.E., Stott, J.M., Schnurch, A., Schein, J.E., Jones, S.J., Holt, R.A., Baross, A., Marra, M.A., Clifton, S., Makowski, K.A., Bosak, S. and Malek, J.

CONSRM MGC Project Team
TITLE The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC)
JOURNAL Genome Res. 14 (10B), 2121-2127 (2004)
PUBMED 15489334
REMARK NUCLEOTIDE SEQUENCE [LARGE SCALE MRNA].
TISSUE:Lung
 Erratum:[Genome Res. 2006 Jun;16(6):804. Morrin, Ryan [corrected to Morin, Ryan]]
COMMENT On or before Feb 16, 2007 this sequence version replaced gi:121941506, gi:7512264.
 [FUNCTION] Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.
 [SUBUNIT] Homotrimer (Potential).
 [SUBCELLULAR LOCATION] Membrane; Single-pass type II membrane protein.
 [TISSUE SPECIFICITY] Expressed in brain, placenta, lung, skeletal muscle and kidney.
 [SIMILARITY] Belongs to the tumor necrosis factor family.
FEATURES

source	Location/Qualifiers
	1..254
	/organism="Homo sapiens"
	/db_xref="taxon:9606"
gene	1..254
	/gene="TNFSF9"
Protein	1..254
	/gene="TNFSF9"
	/product="Tumor necrosis factor ligand superfamily member 9"
Region	1..254
	/gene="TNFSF9"
	/region_name="Mature chain"
	/experiment="experimental evidence, no additional details recorded"
	/note="Tumor necrosis factor ligand superfamily member 9. /FTId=PRO_0000185501."
Region	1..28
	/gene="TNFSF9"
	/region_name="Topological domain"
	/inference="non-experimental evidence, no additional details recorded"

```

Region          /note="Cytoplasmic (Potential)."
17
/gene="TNFSF9"
/region_name="Variant"
/experiment="experimental evidence, no additional details
recorded"
/note="P -> A (in dbSNP:rs442511). /FTId=VAR_011928."
Region          29..49
/gene="TNFSF9"
/region_name="Transmembrane region"
/inference="non-experimental evidence, no additional
details recorded"
/note="Signal-anchor for type II membrane protein
(Potential)."
Region          35..41
/gene="TNFSF9"
/region_name="Compositionally biased region"
/experiment="experimental evidence, no additional details
recorded"
/note="Poly-Leu."
Region          50..254
/gene="TNFSF9"
/region_name="Topological domain"
/inference="non-experimental evidence, no additional
details recorded"
/note="Extracellular (Potential)."
Region          91..238
/gene="TNFSF9"
/region_name="TNF"
/note="Tumor Necrosis Factor; TNF superfamily members
include the cytokines: TNF (TNF-alpha), LT
(lymphotoxin-alpha, TNF-beta), CD40 ligand, Apo2L (TRAIL),
Fas ligand, and osteoprotegerin (OPG) ligand; cd00184"
/db_xref="CDD:29146"
Site            order(94,142,144,199,204,234,238)
/gene="TNFSF9"
/site_type="other"
/note="trimer interface"
/db_xref="CDD:29146"
Site            order(110..111,116,162,169,176)
/gene="TNFSF9"
/site_type="other"
/note="receptor binding sites"
/db_xref="CDD:29146"
ORIGIN
1 meyasdasld peapwppapr aracrvlpw aivagllllll laaacavfla cpwawsgara
61 spgsaasprl regpelspdd paglldlrqg mfaqlvagnv llidgplswy sdpglagvsl
121 tgglsykedt kelvvakagv yyvffqleir rvvagegsgs vslalhlqpl rsaagaaala
181 ltvdlppass earnsaafgfq grllhlsagg rlgvhhltea rarhawqltq gatvlgfrrv
241 tpeipaglps prse
//
```



Display GenPept Show 5 Send to

Range: from begin to end Features: + Refresh

1: BAB18304. Reports OX40 ligand [Homo...[gi:11275538]

BLINK, Links

Features Sequence

LOCUS BAB18304 67 aa linear PRI 19-OCT-2004
 DEFINITION OX40 ligand [Homo sapiens].
 ACCESSION BAB18304
 VERSION BAB18304.1 GI:11275538
 DBSOURCE accession AB042988.2
 KEYWORDS .
 SOURCE Homo sapiens (human)
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;
 Catarrhini; Hominidae; Homo.
 REFERENCE 1
 AUTHORS Hikami,K., Tsuchiya,N. and Tokunaga,K.
 TITLE New variations in human OX40 ligand (CD134L) gene
 JOURNAL Genes Immun. 1 (8), 521-522 (2000)
 PUBMED 11197696
 REFERENCE 2 (residues 1 to 67)
 AUTHORS Hikami,K., Tsuchiya,N. and Tokunaga,K.
 TITLE Direct Submission
 JOURNAL Submitted (19-MAY-2000) Koki Hikami, University of Tokyo,
 Department of Human Genetics; 7-3-1 Hongo, Bunkyo-ku, Tokyo
 113-0033, Japan (E-mail:kokihikami@mail.goo.ne.jp),
 Tel:81-3-5841-3693, Fax:81-3-5802-8619
 FEATURES Location/Qualifiers
 source 1..67
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="1"
 /map="1q25"
Protein 1..67
 /product="OX40 ligand"
CDS 1..67
 /gene="OX40L"
 /coded_by="join(AB042988.2:55..207,AB042988.2:462..>510)"
 ORIGIN
 1 mervqpleen vgnaarprfe rnkllllvasv iqqlglllcf tyiclhfsal qvshrypriq
 61 sikvqft
 //

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

NCBI  Protein [Sign In] [Regis]

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search Protein for

Limits Preview/Index History Clipboard Details

Display GenPept

Range: from to Features: CDD

1: NP_001243. Reports tumor necrosis fa...[gi:4507605]

Link, Conserved Domains, Links

Comment Features Sequence

LOCUS NP_001243 193 aa linear PRI 11-FEB-2008
 DEFINITION tumor necrosis factor ligand superfamily, member 7 [Homo sapiens].
 ACCESSION NP_001243
 VERSION NP_001243.1 GI:4507605
 DBSOURCE REFSEQ: accession NM_001252.3
 KEYWORDS
 SOURCE Homo sapiens (human)
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (residues 1 to 193)
 AUTHORS Yang, Z.Z., Novak, A.J., Ziesmer, S.C., Witzig, T.E. and Ansell, S.M.
 TITLE CD70+ non-Hodgkin lymphoma B cells induce Foxp3 expression and regulatory function in intratumoral CD4+CD25 T cells
 JOURNAL Blood 110 (7), 2537-2544 (2007)
 PUBMED 17615291
 REMARK GeneRIF: reveal a novel role for non-Hodgkin lymphoma B cells in the development of intratumoral regulatory T cells
 REFERENCE 2 (residues 1 to 193)
 AUTHORS van Oosterwijk, M.F., Juwana, H., Arens, R., Tesselaar, K., van Oers, M.H., Eldering, E. and van Lier, R.A.
 TITLE CD27-CD70 interactions sensitise naive CD4+ T cells for IL-12-induced Th1 cell development
 JOURNAL Int. Immunol. 19 (6), 713-718 (2007)
 PUBMED 17548342
 REMARK GeneRIF: CD27-CD70 interactions may promote Th1 cell formation by permitting naive T cells to respond to differentiation signals and by promoting survival of activated effector T cells.
 REFERENCE 3 (residues 1 to 193)
 AUTHORS Diegmann, J., Junker, K., Loncarevic, I.F., Michel, S., Schimmel, B. and von Eggeling, F.
 TITLE Immune escape for renal cell carcinoma: CD70 mediates apoptosis in lymphocytes
 JOURNAL Neoplasia 8 (11), 933-938 (2006)
 PUBMED 17132225
 REMARK GeneRIF: Apoptosis mediated by exposure to the CD70 secreted by tumor cells may contribute to the failure of renal cell carcinoma patients to develop an effective lymphocyte-mediated antitumor response
 REFERENCE 4 (residues 1 to 193)
 AUTHORS Adam, P.J., Terrett, J.A., Steers, G., Stockwin, L., Loader, J.A.,

TITLE Fletcher,G.C., Lu,L.S., Leach,B.I., Mason,S., Stamps,A.C.,
 Boyd,R.S., Pezzella,F., Gatter,K.C. and Harris,A.L.
JOURNAL CD70 (TNFSF7) is expressed at high prevalence in renal cell
 carcinomas and is rapidly internalised on antibody binding
PUBMED Br. J. Cancer 95 (3), 298-306 (2006)
16892042
REMARK GeneRIF: Immunocytochemical analysis demonstrated that binding of
 an anti-CD70 antibody to CD70(TNFSF7), endogenously expressed on
 the surface of A498 and 786-O cell lines resulted in the rapid
 internalisation of the antibody-receptor complex.
REFERENCE 5 (residues 1 to 193)
AUTHORS Huang,J., Kerstann,K.W., Ahmadzadeh,M., Li,Y.F., El~Gamil,M.,
 Rosenberg,S.A. and Robbins,P.F.
TITLE Modulation by IL-2 of CD70 and CD27 expression on CD8+ T cells:
 importance for the therapeutic effectiveness of cell transfer
JOURNAL J. Immunol. 176 (12), 7726-7735 (2006)
PUBMED 16751420
REMARK GeneRIF: Interaction of CD70 with CD27 plays a direct role in T
 cell activation mediated by IL-2.
REFERENCE 6 (residues 1 to 193)
AUTHORS Kobata,T., Jacquot,S., Kozlowski,S., Agematsu,K., Schlossman,S.F.
 and Morimoto,C.
TITLE CD27-CD70 interactions regulate B-cell activation by T cells
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 92 (24), 11249-11253 (1995)
PUBMED 7479974
REFERENCE 7 (residues 1 to 193)
AUTHORS Brown,G.R., Meek,K., Nishioka,Y. and Thiele,D.L.
TITLE CD27-CD70 ligand/CD70 interactions enhance alloantigen-induced
 proliferation and cytolytic activity in CD8+ T lymphocytes
JOURNAL J. Immunol. 154 (8), 3686-3695 (1995)
PUBMED 7706711
REFERENCE 8 (residues 1 to 193)
AUTHORS Hintzen,R.Q., Lens,S.M., Koopman,G., Pals,S.T., Spits,H. and van
 Lier,R.A.
TITLE CD70 represents the human ligand for CD27
JOURNAL Int. Immunol. 6 (3), 477-480 (1994)
PUBMED 8186199
REFERENCE 9 (residues 1 to 193)
AUTHORS Hintzen,R.Q., Lens,S.M., Beckmann,M.P., Goodwin,R.G., Lynch,D. and
 van Lier,R.A.
TITLE Characterization of the human CD27 ligand, a novel member of the
 TNF gene family
JOURNAL J. Immunol. 152 (4), 1762-1773 (1994)
PUBMED 8120385
REFERENCE 10 (residues 1 to 193)
AUTHORS Bowman,M.R., Crimmins,M.A., Yetz-Aldape,J., Kriz,R., Kelleher,K.
 and Herrmann,S.
TITLE The cloning of CD70 and its identification as the ligand for CD27
JOURNAL J. Immunol. 152 (4), 1756-1761 (1994)
PUBMED 8120384
COMMENT REVIEWED REFSEQ: This record has been curated by NCBI staff. The
 reference sequence was derived from AC008760.7.

 Summary: The protein encoded by this gene is a cytokine that
 belongs to the tumor necrosis factor (TNF) ligand family. This
 cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on
 activated, but not on resting, T and B lymphocytes. It induces
 proliferation of costimulated T cells, enhances the generation of
 cytolytic T cells, and contributes to T cell activation. This

cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

FEATURES Location/Qualifiers

source 1..193
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="19"
 /map="19p13"

Protein 1..193
 /product="tumor necrosis factor ligand superfamily, member 7"
 /note="CD70 antigen; CD27 ligand; surface antigen CD70; Ki-24 antigen; tumor necrosis factor (ligand) superfamily, member 7"
 /calculated_mol_wt=20987

Region 58..189
 /region_name="TNF"
 /note="Tumor Necrosis Factor; TNF superfamily members include the cytokines: TNF (TNF-alpha), LT (lymphotoxin-alpha, TNF-beta), CD40 ligand, Apo2L (TRAIL), Fas ligand, and osteoprotegerin (OPG) ligand; cd00184"
 /db_xref="CDD:29146"

Site order(61,105,107,154,159,185,189)
 /site_type="other"
 /note="trimer interface"
 /db_xref="CDD:29146"

Site order(76..77,84,119,126,129)
 /site_type="other"
 /note="receptor binding sites"
 /db_xref="CDD:29146"


CDS 1..193
 /gene="CD70"
 /coded_by="NM_001252.3:151..732"
 /db_xref="CCDS:CCDS12170.1"
 /db_xref="GeneID:970"
 /db_xref="HGNC:11937"
 /db_xref="HPRD:18515"
 /db_xref="MIM:602840"

ORIGIN

```

1 mpeegsgcsv rrrpygcvlr aalvplvagl viclvvcigr faqaqqqlpl eslgwdvael
61 qlnhtgpqgd prlywggpa lgrsflhgpe ldkgqlrihr dglymwhiqv tlaicsstta
121 srhhpttlav gicspasrsi sllrlsfhgg ctiasqrtp largdtlctn ltgtllpsrn
181 tddetffgvqw vrp
//
```

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)



Display GenPept Show 5 Send to
 Range: from to Features: SNP CDD Refresh

1: NP_742011. Reports tumor necrosis fa...[gi:25952147] BLINK, Conserved Domains, Links

Comment Features Sequence

LOCUS NP_742011 204 aa linear PRI 16-MAR-2008
DEFINITION tumor necrosis factor ligand superfamily, member 14 isoform 2 [Homo sapiens].
ACCESSION NP_742011
VERSION NP_742011.1 GI:25952147
DBSOURCE REFSEQ: accession NM_172014.1
KEYWORDS
SOURCE Homo sapiens (human)
ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.
REFERENCE 1 (residues 1 to 204)
AUTHORS Pawlak,K., Pawlak,D. and Mysliwiec,M.
TITLE LIGHT--a new member of the TNF superfamily in the plasma, dialysate and urine of uremic patients; the impact of residual diuresis and presence of viral hepatitis
JOURNAL Clin. Biochem. 40 (16-17), 1240-1244 (2007)
PUBMED 17826757
REMARK GeneRIF: The plasma levels of LIGHT seem to be similar in hemodialysis (HD) patients and healthy subjects and were not affected by gender, age, the mean period of HD history, disease etiology, type of medication and type of using dialysis membrane.
REFERENCE 2 (residues 1 to 204)
AUTHORS Celik,S., Langer,H., Stellos,K., May,A.E., Shankar,V., Kurz,K., Katus,H.A., Gawaz,M.P. and Dengler,T.J.
TITLE Platelet-associated LIGHT (TNFSF14) mediates adhesion of platelets to human vascular endothelium
JOURNAL Thromb. Haemost. 98 (4), 798-805 (2007)
PUBMED 17938804
REMARK GeneRIF: platelet-associated LIGHT is involved in adhesion of platelets to endothelium while soluble LIGHT induces a pro-inflammatory state in vascular endothelial cells
REFERENCE 3 (residues 1 to 204)
AUTHORS Loeffler,M., Le'Negrat,G., Krajewska,M. and Reed,J.C.
TITLE Attenuated Salmonella engineered to produce human cytokine LIGHT inhibit tumor growth
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 104 (31), 12879-12883 (2007)
PUBMED 17652173
REMARK GeneRIF: attenuated Salmonella typhimurium expressing LIGHT inhibited growth of primary tumors, as well as the dissemination of pulmonary metastases, in various mouse tumor models employing

REFERENCE 4 (residues 1 to 204)
 AUTHORS Nadiminty,N., Chun,J.Y., Hu,Y., Dutt,S., Lin,X. and Gao,A.C.
 TITLE LIGHT, a member of the TNF superfamily, activates Stat3 mediated by NIK pathway
 JOURNAL Biochem. Biophys. Res. Commun. 359 (2), 379-384 (2007)
 PUBMED 17543278
 REMARK GeneRIF: in addition to activating NF-kappaB/p52, LIGHT also activates Stat3 through the NIK pathway

REFERENCE 5 (residues 1 to 204)
 AUTHORS Gill,R.M., Coleman,N.M. and Hunt,J.S.
 TITLE Differential cellular expression of LIGHT and its receptors in early gestation human placentas
 JOURNAL J. Reprod. Immunol. 74 (1-2), 1-6 (2007)
 PUBMED 17010447
 REMARK GeneRIF: LIGHT system may regulate early to middle stages of placental development via cell-specific, temporally programmed expression of the ligand and its receptors, and may also assist in preserving placental immune privilege.

REFERENCE 6 (residues 1 to 204)
 AUTHORS Yu,K.Y., Kwon,B., Ni,J., Zhai,Y., Ebner,R. and Kwon,B.S.
 TITLE A newly identified member of tumor necrosis factor receptor superfamily (TR6) suppresses LIGHT-mediated apoptosis
 JOURNAL J. Biol. Chem. 274 (20), 13733-13736 (1999)
 PUBMED 10318773

REFERENCE 7 (residues 1 to 204)
 AUTHORS Harrop,J.A., McDonnell,P.C., Brigham-Burke,M., Lyn,S.D., Minton,J., Tan,K.B., Dede,K., Spampinato,J., Silverman,C., Hensley,P., DiPrinzio,R., Emery,J.G., Deen,K., Eichman,C., Chabot-Fletcher,M., Truneh,A. and Young,P.R.
 TITLE Herpesvirus entry mediator ligand (HVEM-L), a novel ligand for HVEM/TR2, stimulates proliferation of T cells and inhibits HT29 cell growth
 JOURNAL J. Biol. Chem. 273 (42), 27548-27556 (1998)
 PUBMED 9765287

REFERENCE 8 (residues 1 to 204)
 AUTHORS Zhai,Y., Guo,R., Hsu,T.L., Yu,G.L., Ni,J., Kwon,B.S., Jiang,G.W., Lu,J., Tan,J., Ugustus,M., Carter,K., Rojas,L., Zhu,F., Lincoln,C., Endress,C., Xing,L., Wang,S., Oh,K.O., Gentz,R., Ruben,S., Lippman,M.E., Hsieh,S.L. and Yang,D.
 TITLE LIGHT, a novel ligand for lymphotoxin beta receptor and TR2/HVEM induces apoptosis and suppresses in vivo tumor formation via gene transfer
 JOURNAL J. Clin. Invest. 102 (6), 1142-1151 (1998)
 PUBMED 9739048

REFERENCE 9 (residues 1 to 204)
 AUTHORS Marsters,S.A., Sheridan,J.P., Pitti,R.M., Brush,J., Goddard,A. and Ashkenazi,A.
 TITLE Identification of a ligand for the death-domain-containing receptor Apo3
 JOURNAL Curr. Biol. 8 (9), 525-528 (1998)
 PUBMED 9560343

REFERENCE 10 (residues 1 to 204)
 AUTHORS Mauri,D.N., Ebner,R., Montgomery,R.I., Kochel,K.D., Cheung,T.C., Yu,G.L., Ruben,S., Murphy,M., Eisenberg,R.J., Cohen,G.H., Spear,P.G. and Ware,C.F.
 TITLE LIGHT, a new member of the TNF superfamily, and lymphotoxin alpha are ligands for herpesvirus entry mediator
 JOURNAL Immunity 8 (1), 21-30 (1998)
 PUBMED 9462508

COMMENT

REVIEWED REFSEQ: This record has been curated by NCBI staff. The reference sequence was derived from AY028261.1 and AF064090.1.

Summary: The protein encoded by this gene is a member of the tumor necrosis factor (TNF) ligand family. This protein is a ligand for TNFRSF14, which is a member of the tumor necrosis factor receptor superfamily, and which is also known as a herpesvirus entry mediator (HVEM). This protein may function as a costimulatory factor for the activation of lymphoid cells and as a deterrent to infection by herpesvirus. This protein has been shown to stimulate the proliferation of T cells, and trigger apoptosis of various tumor cells. This protein is also reported to prevent tumor necrosis factor alpha mediated apoptosis in primary hepatocyte. Two alternatively spliced transcript variant encoding distinct isoforms have been reported.

Transcript Variant: This variant (2) lacks an in-frame coding segment compared to variant 1, resulting an isoform (2) that lacks an internal region, as compared to isoform 1.

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

FEATURES

	Location/Qualifiers
<u>source</u>	1..204 /organism="Homo sapiens" /db_xref="taxon:9606" /chromosome="19" /map="19p13.3"
<u>Protein</u>	1..204 /product="tumor necrosis factor ligand superfamily, member 14 isoform 2" /note="delta transmembrane LIGHT; tumor necrosis factor superfamily member LIGHT; herpesvirus entry mediator A; tumor necrosis factor receptor-like 2; ligand for herpesvirus entry mediator" /calculated_mol_wt=22265
<u>Region</u>	58..202 /region_name="TNF" /note="Tumor Necrosis Factor; TNF superfamily members include the cytokines: TNF (TNF-alpha), LT (lymphotoxin-alpha, TNF-beta), CD40 ligand, Apo2L (TRAIL), Fas ligand, and osteoprotegerin (OPG) ligand; cd00184" /db_xref="CDD:29146"
<u>Site</u>	order(61,106,108,166,171,198,202) /site_type="other" /note="trimer interface" /db_xref="CDD:29146"
<u>Site</u>	order(79,80,85,127,134,139) /site_type="other" /note="receptor binding sites" /db_xref="CDD:29146"
<u>CDS</u>	1..204 /gene="TNFSF14" /coded_by="NM_172014.1:383..997" /note="isoform 2 is encoded by transcript variant 2" /db_xref="GeneID:8740" /db_xref="HGNC:11930" /db_xref="MIM:604520"

ORIGIN

```

1 meesvvrvpsv fvvdgqtidp ftrlggrshrr qscsvardgp agsweqlige rrshevnpaa
61 hltgansslt gsgggpllwet qlqlafirgl syhdgalvvt kagyyyiyks vqlggvgcpl
121 glastithgl ykrtprypee lellvsqqsp cgratsssrw wwdssflggv vhleageevv
181 vrvlderlvr lrdgtrsyyfg afmv
//
```

NCBI   Protein My NCBI [Sign In] [Regis]

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search Protein for

Limits Preview/Index History Clipboard Details

Display GenPept Show 5 Send to 

Range: from to Features: CDD

 1: [AAB97877](#). Reports CD30L protein [Ho...[gi:2815515]]  Conserved Domains, Links

Features Sequence

LOCUS AAB97877 234 aa linear PRI 28-JAN-1998

DEFINITION CD30L protein [Homo sapiens].

ACCESSION AAB97877

VERSION AAB97877.1 GI:2815515

DBSOURCE locus HSCD30L1 accession AF006381.1
locus HSCD30L2 accession AF006382.1
locus HSCD30L3 accession AF006383.1
locus HSCD30L4 accession AF006384.1

KEYWORDS

SOURCE Homo sapiens (human)

ORGANISM Homo_sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.

REFERENCE 1 (residues 1 to 234)

AUTHORS Croager, E.J. and Abraham, L.J.

TITLE Characterisation of the human CD30 ligand gene structure

JOURNAL Biochim. Biophys. Acta 1353 (3), 231-235 (1997)
9349718

REFERENCE 2 (residues 1 to 234)

AUTHORS Croager, E.J. and Abraham, L.J.

TITLE Direct Submission

JOURNAL Submitted (02-JUN-1997) Biochemistry, University of Western Australia, Entrance 2, Hackett Drive, Nedlands, Western Australia 6907, Australia

COMMENT Method: conceptual translation.

FEATURES Location/Qualifiers

source 1..234
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="9"
/map="9q33"

Protein 1..234
/product="CD30L protein"

Region 104..220
/region_name="TNF"
/note="Tumor Necrosis Factor; TNF superfamily members include the cytokines: TNF (TNF-alpha), LT (lymphotoxin-alpha, TNF-beta), CD40 ligand, Apo2L (TRAIL), Fas ligand, and osteoprotegerin (OPG) ligand; cd00184"
/db_xref="CDD:29146"
order(107,140,142,190,195)

Site

Site
/site_type="other"
/note="trimer interface"
/db_xref="CDD:29146"
order(116..117,157,164,169)
/site_type="other"
/note="receptor binding sites"
/db_xref="CDD:29146"
CDS
1..234
/gene="CD30L"
/coded_by="join(AF006381.1:1599..1793,AF006382.1:188..230,
AF006383.1:534..605,AF006384.1:347..741)"
ORIGIN
1 mdpqlqqaln gmappgdtam hvpagsvash lgtsrsyfy lttatlalcl vftvatinv1
61 vvqrtdsipn spdnvplkgg ncseadllcilm krapfkkswa ylgvakhlnk tklswnkdg1
121 lhgvryqdggn lviqfpglyf iicqlqflvq cpnnsvdlkx ellinkhikk qx1vtvcesg
181 mqtkhvyqnl sqflldy1qv nttisvndt xqyidtstfp lenvlsiflx sns1
//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

NCBI Protein (Sign In) (Regis My NI

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search Protein for

Limits Preview/Index History Clipboard Details

Display GenPept

Range: from to Features: CDD

1: P42081. Reports T-lymphocyte acti...[gi:1168851] BLINK, Conserved Domains, Links

Comment Features Sequence

LOCUS P42081 329 aa linear PRI 15-JAN-2008

DEFINITION T-lymphocyte activation antigen CD86 precursor (Activation B7-2 antigen) (CTLA-4 counter-receptor B7.2) (B70) (FUN-1) (BU63).

ACCESSION P42081

VERSION P42081.1 GI:1168851

DBSOURCE swissprot: locus CD86_HUMAN, accession P42081; class: standard.

extra accessions: A0N0P0, Q13655, Q6FHB1, Q6GTS4, Q7M4L5

created: Nov 1, 1995.

sequence updated: Nov 1, 1995.

annotation updated: Jan 15, 2008.

xrefs: [U04343.1](#), [AAB03814.1](#), [L25259.1](#), [AAA58389.1](#), [CR541844.1](#), [CAG46642.1](#), [EF064748.1](#), [ABK41931.1](#), [BC040261.1](#), [AAH40261.1](#), [U17722.1](#), [AAA86473.1](#), [U17717.1](#), [U17718.1](#), [U17719.1](#), [U17721.1](#), [A48754](#), [JC7605](#), [1I85](#), [1NCN](#)

xrefs (non-sequence databases): UniGene:Hs.171182, PDBsum:1I85, PDBsum:1NCN, IntAct:P42081, Ensembl:ENSG00000114013, KEGG:hsa:942, H-InvDB:[HIX0024331](#), HGNC:1705, HPA:CA004319, MIM:601020, PharmGKB:PA26243, LinkHub:P42081, ArrayExpress:P42081, CleanEx:HS_CD86, GermOnline:ENSG00000114013, GO:[0016021](#), GO:[0005886](#), GO:[0015026](#), GO:[0005515](#), GO:[0016563](#), GO:[0007267](#), GO:[0008284](#), GO:[0045086](#), GO:[0045404](#), GO:[0043017](#), GO:[0045630](#), GO:[0045941](#), InterPro:[IPR015651](#), InterPro:[IPR007110](#), InterPro:[IPR013783](#), InterPro:[IPR003006](#), InterPro:[IPR013106](#), InterPro:[IPR003596](#), Gene3D:G3DSA:2.60.40.10, PANTHER:PTHR13712:SF45, Pfam:[PF07686](#), SMART:SM00406, PROSITE:PS50835, PROSITE:PS00290

KEYWORDS 3D-structure; Alternative splicing; Glycoprotein; Host-virus interaction; Immunoglobulin domain; Membrane; Polymorphism; Receptor; Signal; Transmembrane; Ubl conjugation.

SOURCE Homo sapiens (human)

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.

REFERENCE 1 (residues 1 to 329)

AUTHORS Azuma,M., Ito,D., Yagita,H., Okumura,K., Phillips,J.H., Lanier,L.L. and Somoza,C.

TITLE B70 antigen is a second ligand for CTLA-4 and CD28

JOURNAL Nature 366 (6450), 76-79 (1993)

PUBMED 7694153

REMARK NUCLEOTIDE SEQUENCE [MRNA] (ISOFORM 2).

REFERENCE 2 (residues 1 to 329)
 AUTHORS Freeman,G.J., Gibben,J.G., Boussiotis,V.A., Ng,J.W., Restivo,V.A.
 Jr., Lombard,L.A., Gray,G.S. and Nadler,L.M.
 TITLE Cloning of B7-2: a CTLA-4 counter-receptor that costimulates human
 T cell proliferation
 JOURNAL Science 262 (5135), 909-911 (1993)
 PUBMED 7694363
 REMARK NUCLEOTIDE SEQUENCE [mRNA] (ISOFORM 1).
 REFERENCE 3 (residues 1 to 329)
 AUTHORS Magistrelli,G., Caron,G., Gauchat,J.F., Jeannin,P., Bonnefoy,J.Y.
 and Delneste,Y.
 TITLE Identification of an alternatively spliced variant of human CD86
 mRNA
 JOURNAL Biochem. Biophys. Res. Commun. 280 (5), 1211-1215 (2001)
 PUBMED 11162656
 REMARK NUCLEOTIDE SEQUENCE [mRNA] (ISOFORM 3).
 REFERENCE 4 (residues 1 to 329)
 AUTHORS Halleck,A., Ebert,L., Mkoundinya,M., Schick,M., Eisenstein,S.,
 Neubert,P., Kstrang,K., Schatten,R., Shen,B., Henze,S., Mar,W.,
 Korn,B., Zuo,D., Hu,Y. and LaBaer,J.
 TITLE Direct Submission
 JOURNAL Submitted (??-JUN-2004)
 REMARK NUCLEOTIDE SEQUENCE [LARGE SCALE mRNA] (ISOFORM 2).
 REFERENCE 5 (residues 1 to 329)
 AUTHORS Livingston,R.J., Shaffer,T., McFarland,I., Nguyen,C.P.,
 Stanaway,I.B., Rajkumar,N., Johnson,E.J., da Ponte,S.H., Willa,H.,
 Ahearn,M.O., Bertucci,C., Acklestad,J., Carroll,A., Swanson,J.,
 Gildersleeve,H.I. and Nickerson,D.A.
 TITLE Direct Submission
 JOURNAL Submitted (??-OCT-2006)
 REMARK NUCLEOTIDE SEQUENCE (GENOMIC DNA).
 REFERENCE 6 (residues 1 to 329)
 AUTHORS Gerhard,D.S., Wagner,L., Feingold,E.A., Shenmen,C.M., Grouse,L.H.,
 Schuler,G., Klein,S.L., Old,S., Rasooly,R., Good,P., Guyer,M.,
 Peck,A.M., Derge,J.G., Lipman,D., Collins,F.S., Jang,W., Sherry,S.,
 Feolo,M., Misquitta,L., Lee,E., Rotmistrovsky,K., Greenhut,S.F.,
 Schaefer,C.F., Buetow,K., Bonner,T.I., Haussler,D., Kent,J.,
 Kiekhau,M., Furey,T., Brent,M., Prange,C., Schreiber,K.,
 Shapiro,N., Bhat,N.K., Hopkins,R.F., Hsie,F., Driscoll,T.,
 Soares,M.B., Casavant,T.L., Scheetz,T.E., Brown-stein,M.J.,
 Usdin,T.B., Toshiyuki,S., Carninci,P., Piao,Y., Dudekula,D.B.,
 Ko,M.S., Kawakami,K., Suzuki,Y., Sugano,S., Grubetz,C.E.,
 Smith,M.R., Simmons,B., Moore,T., Waterman,R., Johnson,S.L.,
 Ruan,Y., Wei,C.L., Mathavan,S., Gunaratne,P.H., Wu,J., Garcia,A.M.,
 Hulyk,S.W., Fuh,E., Yuan,Y., Sneed,A., Kowis,C., Hodgson,A.,
 Muzny,D.M., McPherson,J., Gibbs,R.A., Fahey,J., Helton,E.,
 Ketteman,M., Madan,A., Rodrigues,S., Sanchez,A., Whiting,M.,
 Madari,A., Young,A.C., Wetherby,K.D., Granite,S.J., Kwong,P.N.,
 Brinkley,C.P., Pearson,R.L., Bouffard,G.G., Blakesly,R.W.,
 Green,E.D., Dickson,M.C., Rodriguez,A.C., Grimwood,J., Schmutz,J.,
 Myers,R.M., Butterfield,Y.S., Griffith,M., Griffith,O.L.,
 Krzywinski,M.I., Liao,N., Morin,R., Palmquist,D., Petrescu,A.S.,
 Skalska,U., Smailus,D.E., Stott,J.M., Schnurch,A., Schein,J.E.,
 Jones,S.J., Holt,R.A., Baross,A., Marra,M.A., Clifton,S.,
 Makowski,K.A., Bosak,S. and Malek,J.
 CONSRM MGC Project Team
 TITLE The status, quality, and expansion of the NIH full-length cDNA
 project: the Mammalian Gene Collection (MGC)
 JOURNAL Genome Res. 14 (10B), 2121-2127 (2004)
 PUBMED 15489334

REMARK NUCLEOTIDE SEQUENCE [LARGE SCALE MRNA] (ISOFORM 1), AND VARIANT
 THR-310.
 TISSUE=Brain
 Erratum: [Genome Res. 2006 Jun;16(6):804. Morrin, Ryan (corrected to
 Morin, Ryan)]
 REFERENCE 7 (residues 1 to 329)
 AUTHORS Jellis,C.L., Wang,S.S., Rennert,P., Borriello,F., Sharpe,A.H.,
 Green,N.R. and Gray,G.S.
 TITLE Genomic organization of the gene coding for the costimulatory human
 B-lymphocyte antigen B7-2 (CD86)
 JOURNAL Immunogenetics 42 (2), 85-89 (1995)
 PUBMED 7541777
 REMARK NUCLEOTIDE SEQUENCE [GENOMIC DNA] OF 7-329.
 TISSUE=Foreskin
 REFERENCE 8 (residues 1 to 329)
 AUTHORS Lanier,L.L., O'Fallon,S., Somoza,C., Phillips,J.H., Linsley,P.S.,
 Okumura,K., Ito,D. and Azuma,M.
 TITLE CD80 (B7) and CD86 (B70) provide similar costimulatory signals for
 T cell proliferation, cytokine production, and generation of CTL
 J. Immunol. 154 (1), 97-105 (1995)
 JOURNAL 7527824
 PUBMED
 REMARK CHARACTERIZATION.
 REFERENCE 9 (residues 1 to 329)
 AUTHORS Engel,F., Gribben,J.G., Freeman,G.J., Zhou,L.J., Nozawa,Y., Abe,M.,
 Nadler,L.M., Wakasa,H. and Tedder,T.F.
 TITLE The B7-2 (B70) costimulatory molecule expressed by monocytes and
 activated B lymphocytes is the CD86 differentiation antigen
 JOURNAL Blood 84 (5), 1402-1407 (1994)
 PUBMED 7520767
 REMARK IDENTIFICATION AS CD86.
 REFERENCE 10 (residues 1 to 329)
 AUTHORS Goto,E., Ishido,S., Sato,Y., Ohgimoto,S., Ohgimoto,K.,
 Nagano-Fujii,M. and Hotta,H.
 TITLE c-MIR, a human E3 ubiquitin ligase, is a functional homolog of
 herpesvirus proteins MIR1 and MIR2 and has similar activity
 J. Biol. Chem. 278 (17), 14657-14668 (2003)
 JOURNAL 12582153
 PUBMED
 REMARK UBIQUITINATION, AND INTERACTION WITH MARCH8.
 COMMENT On or before Feb 27, 2007 this sequence version replaced
 gi:74721835, gi:74757610, gi:627415, gi:25392201.
 [FUNCTION] Receptor involved in the costimulatory signal essential
 for T-lymphocyte proliferation and interleukin-2 production, by
 binding CD28 or CTLA-4. May play a critical role in the early
 events of T-cell activation and costimulation of naive T-cells,
 such as deciding between immunity and anergy that is made by
 T-cells within 24 hours after activation. Isoform 2 interferes with
 the formation of CD86 clusters, and thus acts as a negative
 regulator of T-cell activation.
 [SUBUNIT] Interacts with MARCH8. Interacts with human herpesvirus 8
 MIR2 protein (Probable).
 [SUBCELLULAR LOCATION] Membrane; Single-pass type I membrane
 protein.
 [ALTERNATIVE PRODUCTS] Event=Alternative splicing; Named
 isoforms=3; Name=1; IsoId=P42081-1; Sequence=Displayed; Name=2;
 IsoId=P42081-3; Sequence=VSP_023124; Name=3; Synonyms=CD86 deltaEC;
 IsoId=P42081-2; Sequence=VSP_009125.
 [TISSUE SPECIFICITY] Expressed by activated B-lymphocytes and
 monocytes.
 [PTM] Polyubiquitinated: which is promoted by MARCH8 and results in
 endocytosis and lysosomal degradation.

[SIMILARITY] Contains 1 Ig-like C2-type (immunoglobulin-like) domain.
 [SIMILARITY] Contains 1 Ig-like V-type (immunoglobulin-like) domain.
 [WEB RESOURCE] Name=Wikipedia; Note=CD86 entry;
 URL='<http://en.wikipedia.org/wiki/CD86>'.

FEATURES Location/Qualifiers
source 1..329
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
gene 1..329
 /gene="CD86"
 /note="synonym: CD28LG2"
Protein 1..329
 /gene="CD86"
 /product="T-lymphocyte activation antigen CD86 precursor"
Region 1..23
 /gene="CD86"
 /region_name="Signal"
 /inference="non-experimental evidence, no additional details recorded"
 /note="Potential."
Region 1..6
 /gene="CD86"
 /region_name="Splicing variant"
 /experiment="experimental evidence, no additional details recorded"
 /note="Missing (in isoform 2). /FTId=VSP_023124."
Region 22..234
 /gene="CD86"
 /region_name="Splicing variant"
 /experiment="experimental evidence, no additional details recorded"
 /note="Missing (in isoform 3). /FTId=VSP_009125."
Region 24..329
 /gene="CD86"
 /region_name="Mature chain"
 /experiment="experimental evidence, no additional details recorded"
 /note="T-lymphocyte activation antigen CD86.
 /FTId=PRO_0000014550."
Region 24..247
 /gene="CD86"
 /region_name="Topological domain"
 /inference="non-experimental evidence, no additional details recorded"
 /note="Extracellular (Potential)."
Region 27..31
 /gene="CD86"
 /region_name="Beta-strand region"
 /experiment="experimental evidence, no additional details recorded"
Region 27
 /gene="CD86"
 /region_name="Conflict"
 /experiment="experimental evidence, no additional details recorded"
 /note="K -> E (in Ref. 7; AAA86473)."
Region 33..131
 /gene="CD86"

```
/region_name="Domain"
/experiment="experimental evidence, no additional details
recorded"
/note="Ig-like V-type."
33
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
35..110
/gene="CD86"
/region_name="IGv"
/note="Immunoglobulin V-Type; smart00406"
/db_xref="CDD:47715"
Region
36..38
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Bond
bond(40,110)
/gene="CD86"
/bond_type="disulfide"
/inference="non-experimental evidence, no additional
details recorded"
/note="Potential."
Site
47
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
Region
54..58
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Region
64..69
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Region
80..84
/gene="CD86"
/region_name="Hydrogen bonded turn"
/experiment="experimental evidence, no additional details
recorded"
Region
85..88
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Region
90..92
/gene="CD86"
/region_name="Hydrogen bonded turn"
/experiment="experimental evidence, no additional details
recorded"
Region
95..97
/gene="CD86"
/region_name="Beta-strand region"
```

```
/experiment="experimental evidence, no additional details
recorded"
Region
106..113
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Region
122..133
/gene="CD86"
/region_name="Beta-strand region"
/experiment="experimental evidence, no additional details
recorded"
Site
135
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
Site
146
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
Region
150..225
/gene="CD86"
/region_name="Domain"
/experiment="experimental evidence, no additional details
recorded"
/note="Ig-like C2-type."
Site
154
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
Bond
bond(157, 218)
/gene="CD86"
/bond_type="disulfide"
/inference="non-experimental evidence, no additional
details recorded"
/note="Potential."
Region
170
/gene="CD86"
/region_name="Variant"
/experiment="experimental evidence, no additional details
recorded"
/note="S -> N (in dbSNP:rs9282642). /FTId=VAR_021916."
Site
177
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
Site
192
/gene="CD86"
/site_type="glycosylation"
/inference="non-experimental evidence, no additional
details recorded"
/note="N-linked (GlcNAc...) (Potential)."
```

Site 213
 /gene="CD86"
 /site_type="glycosylation"
 /inference="non-experimental evidence, no additional details recorded"
 /note="N-linked (GlcNAc...) (Potential)."
Region 248..268
 /gene="CD86"
 /region_name="Transmembrane region"
 /inference="non-experimental evidence, no additional details recorded"
 /note="Potential."
Region 269..329
 /gene="CD86"
 /region_name="Topological domain"
 /inference="non-experimental evidence, no additional details recorded"
 /note="Cytoplasmic (Potential)."
Region 310
 /gene="CD86"
 /region_name="Variant"
 /experiment="experimental evidence, no additional details recorded"
 /note="A -> T (in dbSNP:rs1129055). /FTId=VAR_014650."
Region 323
 /gene="CD86"
 /region_name="Variant"
 /experiment="experimental evidence, no additional details recorded"
 /note="D -> N (in dbSNP:rs9282648). /FTId=VAR_021917."
ORIGIN
 1 mdpqctmgls nilfvmafll sgaaplkiga yfnetadlpc qfansqngsl selvvfwqdq
 61 enlvlnevyl gkekfdsvhs kymgrtsfds dswt1rlh1 qikdkglyqc iihhkkptgm
 121 irihqmmsel svlanfsqpe ivpisniten vyinltcssi hgypepkkms vllrtknsti
 181 eydgimqksq dnvteleydvs islsvsfpdv tsmntifcil etdktrllss pfsieledpq
 241 pppdhipwit avlptviicv mvfc1ilwkw kkkkrprnsy kcgtntmre eseqtkkrek
 301 ihipersdea qrvfksskts scdksdacf
//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)