



- (51) **International Patent Classification:**
C12Q 1/68 (2006.01)
- (21) **International Application Number:**
PCT/EP2012/052913
- (22) **International Filing Date:**
21 February 2012 (21.02.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
11155164.4 21 February 2011 (21.02.2011) EP
- (71) **Applicant (for all designated States except US):** **QIAGEN GMBH** [DE/DE]; Qiagen Str. 1, 40724 Hilden (DE).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **DI PASQUALE, Francesca** [IT/DE]; Qiagen GmbH, Qiagen Str. 1, 40724 Hilden (DE). **ENGEL, Holger** [DE/DE]; Qiagen GmbH, Qiagen Str. 1, 40724 Hilden (DE).
- (74) **Agent:** **KILGER, Christian**; Fanelli Haag & Kilger PLLC, Fasanenstrasse 29, 10719 Berlin (DE).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2012/113780 A1

(54) **Title:** METHOD FOR QUANTIFYING HUMAN DNA

(57) **Abstract:** The invention provides for a method for quantifying one or more nucleic acids of a genome in a sample comprising the steps of, (a) amplifying a first nucleic acid to be quantified, (b) determining the amount of said first nucleic acid by comparison of the amount of amplification product from said first nucleic acid with at least one amplification product from a second template nucleic acid, (c) wherein said second template nucleic acid was generated using whole genome amplification and wherein the starting concentration of the second template nucleic acid is known.

METHOD FOR QUANTIFYING HUMAN DNA

Field of the Invention

5

The present invention is in the field of molecular biology, diagnostics, more particularly in the field of analytical and forensic sciences. The invention is further in the field of nucleic acid amplification and quantification, more particularly in the field of DNA quantification in forensics.

10

Background

The determination of the quantity of DNA recovered from forensic samples as well as other samples is a critical step in the over all DNA typing process, but also in the detection of DNA in various other fields of science. A narrow range of input DNA from 0.5 to 2 ng is often needed to produce optimal results with for example multiplex DNA typing kits. Therefore, in order to ensure that a positive result is a positive result and/or a negative result is a negative result due to the absence of DNA, quantification of DNA is of absolute importance. Furthermore, the quality of standards for forensic DNA testing laboratories requires human-specific DNA quantification. This is due to isolation techniques that can recover human DNA as well as bacterial and other exogenous DNA. A number of procedures have been developed to permit quantification of human-specific DNA including start-blot techniques, liquid based hybridization assays and real-time PCR (polymerase chain reaction). Currently, real-time PCR is the dominant technique due to its wide dynamic range and ease of automation.

The modern STR-Kits have become much more sensitive and can obtain good results even using low amounts of DNA. Therefore, there is a desire for a method, kit and nucleic acid region that allows precise and accurate quantification of human DNA even in low concentrated samples. There are certain quantification and detection kits already available, however, these have serious drawbacks. One such kit is the Quantifiler Human Kit (Applied Biosystems) another kit is Quantifiler Duo Kit (Applied Biosystems) another kit is the Plexor HY Real-Time PCR Quantification Kit (Promega). Both the Quantifiler Duo

30

Kit and the Plexor HY Kit target an autosomal and a gonosomal (Y-chromosome) target on the genome.

Drawbacks for the kits: According to LaSalle *et al.*, (Forensic Science International: Genetics, “Analysis of Single and Multi-copy Methods for DNA Quantification by Real-Time Polymerase Chain Reaction”) the Quantifiler Kits are more accurate in the
5 quantification but have a lower dynamic range as the Plexor HY. The Plexor HY offers a higher dynamic range due to the amplification of a multi-copy target, but a lower accuracy. This lower accuracy can be attributed to the multicopy target. If less than the full set of 20 copies on a genome amplify, because of, for example, instability in the target copy number,
10 than the ratio between the amplification between autosomal and gonosomal (Y) target may vary. The dynamic range of the Plexor HY kit is slightly better than that of the other kit (LaSalle *et al.*, Forensic Science International: Genetics, “Analysis of Single and Multi-copy Methods for DNA Quantification by Real-Time Polymerase Chain Reaction”). In a statistical comparison LaSalle *et al.* demonstrated a significant difference between the two
15 kits.

Another important parameter in forensics is the degradation grade of the DNA, that has to be analyzed. Since the amplicon size of the Quantifiler Human and Plexor HY vary from 62 to 133 base pairs (bp), significant differences might be expected when the kits are
20 applied to degraded DNA.

It should be emphasized that there are currently no official calibration standards or reference materials available especially for the quantification of proteins and nucleic acids (Ellison SL, English CA, Burns MJ, Keer JT: Routes to improving the reliability of low
25 level DNA analysis using real-time PCR. BMC Biotechnol 6: 33, 2006). A further problem arises from the pipetting error during the dilution of DNA quantification standards. In addition, long time stability of standards which is sometimes difficult in the case of natural polymers like nucleic acids and proteins must be guaranteed to achieve reproducibility. This is especially important in the case of low amounts of these substances. Furthermore,
30 the manufacturing of reference substances has to be carried out under strict quality control to avoid secondary contaminations.

The Directive 2004/9/EC of THE EUROPEAN PARLIAMENT and of THE COUNCIL of 11 February 2004 on the inspection and verification of good laboratory practice (GLP)

(Official Journal of the European Union, Vol. L 50, pp. 28-43, 20 February 2004) states in Part B (Revised guidance for the conduct of test facility inspections and study audits) following inspection and audit criteria for test systems and test and reference substances:
5 “Test systems Purpose: to determine whether adequate procedures exist for the handling and control of the variety of test systems required by the studies undertaken in the facility, for example, chemical and physical systems, cellular and microbic systems, plants or animals (...). Test and reference substances Purpose: to determine whether the test facility has procedures designed (i) to ensure that the identity, potency, quantity and composition of test and reference substances are in accordance with their specifications, and (ii) to
10 properly receive and store test and reference substances.....”.

According to DIN EN ISO 9000:2005 certified reference material (CRM) is a reference material, accompanied by a certificate, one or more of whose property values are certified by a procedure that establishes metrological traceability to an accurate realization of the
15 unit in which property values are expressed, and for which each certified value is accompanied by an uncertainty at a stated level of confidence.

Processing of the analytical results into DNA profiles also needs to be carefully controlled, and strict guidelines for the identification of alleles have to be developed and followed, to
20 take account of such things as stutters, artefacts, peak size and morphology, variations in peak ratios, and so on. Three independent scientists, or two scientists and an expert system, are currently used in the UK to confirm the accurate designation of DNA profiles.

A centre piece is the quantification of template DNA. This has so far not been addressed
25 adequately.

Summary of the Invention

The present invention solves the above identified problem and provides for the following
30 solution as outlined below.

Hence the invention relates to a method for quantifying one or more nucleic acids of a genome in a sample comprising the steps of, (a) amplifying a first nucleic acid to be quantified, (b) determining the amount of said first nucleic acid by comparison of the

amount of amplification product from said first nucleic acid with at least one amplification product from a second template nucleic acid, (c) wherein said second template nucleic acid was generated using whole genome amplification and wherein the starting concentration of the second template nucleic acid is known.

5

Herein, whole genome amplification is also referred to as "WGA". It is the amplification of substantially large parts or substantially most parts or all parts of a particular genome by any of numerous methods, such as with random primers. A non limiting list of methods according to the invention includes the following WGA methods:

10

The first whole genome amplification methods were described in 1992, and were based on the principles of the polymerase chain reaction. Zhang and coworkers (Zhang, L., et al. Proc. Natl. Acad. Sci. USA, 1992, 89: 5847-5851) developed the primer extension PCR technique (PEP) and Telenius and collaborators (Telenius et al., Genomics. 1992, 13(3):718-25) designed the degenerate oligonucleotide-primed PCR method (DOP-PCR) Zhang et al., 1992).

15

PEP involves a high number of PCR cycles; using Taq polymerase and 15 base random primers that anneal at a low stringency temperature. The PEP protocol has been improved in different ways

20

DOP-PCR is a method which uses Taq polymerase and semi-degenerate oligonucleotides that bind at a low annealing temperature at approximately one million sites within the human genome. The first cycles are followed by a large number of cycles with a higher annealing temperature, allowing only for the amplification of the fragments that were tagged in the first step.

25

Multiple displacement amplification (MDA, also known as strand displacement amplification; SDA) is a non-PCR-based isothermal method based on the annealing of random hexamers to denatured DNA, followed by strand-displacement synthesis at constant temperature (Blanco et al., 1989, J. Biol. Chem. 264:8935-40). It has been applied to small genomic DNA samples, leading to the synthesis of high molecular weight DNA with limited sequence representation bias (Lizardi et al., Nature Genetics 1998, 19, 225-232; Dean et al., Proc. Natl. Acad. Sci. U.S.A. 2002, 99, 5261-5266). As DNA is

30

synthesized by strand displacement, a gradually increasing number of priming events occur, forming a network of hyper-branched DNA structures. The reaction can be catalyzed by the Phi29 DNA polymerase or by the large fragment of the Bst DNA polymerase. The Phi29 DNA polymerase possesses a proofreading activity resulting in error rates 100 times lower than the Taq polymerase.

US 2010/0035232 discloses a further approach and US2004/0209298 discloses further amplification methods.

Herein, human genomic DNA is a sample comprising at least a fraction of human genomic DNA, wherein at least one copy of each human chromosome is present, i.e. in a male individual the male set and for a female individual the female set.

Herein, a male individual is an individual or a cell line that carries a full male chromosome set.

Herein, a female individual is an individual or a cell line that carries a full female chromosome set.

The invention also relates to the use of a nucleic acid amplification mixture generated by whole genome amplification for quantifying a further nucleic acid composition.

And, the invention relates to a kit for quantifying nucleic acids, comprising one more vials with a nucleic acid composition that has been generated by means of whole genome amplification.

As used herein, a kit is a packaged combination optionally including instructions for use of the combination and/or other reactions and components for such use.

30 **Detailed description of the Invention**

As outlined above, known means of quantifying DNA by making use of a reference DNA and an amplification method have the drawback that the reference DNA may vary from lot to lot.

The invention addresses the need to be able to define test sensitivity, i.e. the amount of DNA that can be detected reliably.

- 5 The invention addresses the need to be able to define the amount of DNA that the laboratory can detect reliably in mixture, wherein there is a presence of another DNA.

The invention addresses the need to be able to define the reproducibility with which the laboratory can detect gender specific loci.

10

The invention addresses the need to be able to calibrate the reproducibility of, e.g. signal intensity, laboratory equipment, chemicals or enzymes.

It addresses the need to be able to understand the influence of certain PCR inhibitors.

15

The inventors address the above mentioned needs by providing for a method for quantifying one or more nucleic acids of a genome in a sample comprising the steps of, (a) amplifying a first nucleic acid to be quantified, (b) determining the amount of said first nucleic acid by comparison of the amount of amplification product from said first nucleic acid with at least one amplification product from a second template nucleic acid, (c) wherein said second template nucleic acid was generated using whole genome amplification and wherein the starting concentration of the second template nucleic acid is known.

20

- 25 The amplification method is either a non-isothermal method or an isothermal method.

The non-isothermal amplification method may be selected from the group of polymerase chain reaction (PCR) (Saiki *et al.* (1985) *Science* 230:1350), quantitative real-time PCR (rtPCR), ligase chain reaction (LCR) (Landegren *et al.* (1988) *Science* 241:1077-1080).

30

Polymerase chain reaction amplification is preferred.

The isothermal amplification method may be selected from the group of helicase-dependent amplification (HDA) (Vincent *et al.* (2004) *EMBO rep* 5(8):795-800), thermostable HDA (tHDA) (An *et al.* (2005) *J Biol Chem* 280(32):28952-28958), strand

displacement amplification (SDA) (Walker *et al.* (1992) *Nucleic Acids Res* 20(7):1691-6), multiple displacement amplification (MDA) (Dean *et al.* (2002) *Proc Natl Acad Sci USA* 99(8): 5261-5266), rolling-circle amplification (RCA) (Liu *et al.* (1996) *J Am Chem Soc* 118:1587-1594), restriction aided RCA (Wang *et al.* (2004) *Genome Res* 14:2357-2366),
5 single primer isothermal amplification (SPIA) (Dafforn *et al.* (2004) *Biotechniques* 37(5):854-7), transcription mediated amplification (TMA) (Vuorinen *et al.* (1995) *J Clin Microbiol* 33: 1856-1859), nicking enzyme amplification reaction (NEAR) (Maples *et al.* US2009017453), exponential amplification reaction (EXPAR) (Van Ness *et al.* (2003) *Proc Natl Acad Sci USA* 100(8):4504-4509), loop-mediated isothermal amplification
10 (LAMP) (Notomi *et al.* (2000) *Nucleic Acids Res* 28(12):e63), recombinase polymerase amplification (RPA) (Piepenburg *et al.* (2006) *PloS Biol* 4(7):1115-1120), nucleic acid sequence based amplification (NASBA) (Kievits *et al.* (1991) *J Virol Methods* 35:273-286), smart-amplification process (SMAP) (Mitani *et al.* (2007) *Nat Methods* 4(3):257-62).

15 By “isothermal amplification reaction” in context of the present invention it is meant that the temperature does not significantly change during the reaction. In a preferred embodiment the temperature of the isothermal amplification reaction does not deviate by more than 10 °C, preferably by not more than 5 °C, even more preferably not more than 2 °C during the main enzymatic reaction step where amplification takes place.

20

Depending on the method of isothermal amplification of nucleic acids different enzymes are required for the amplification reaction. Known isothermal methods for amplification of nucleic acids are the above mentioned, wherein the at least one mesophilic enzyme for amplifying nucleic acids under isothermal conditions is selected from the group consisting
25 of helicase, mesophilic polymerases, mesophilic polymerases having strand displacement activity, nicking enzymes, recombination proteins, ligases, glycosylases and/or nucleases.

“Helicases” are known by those skilled in the art. They are proteins that move directionally along a nucleic acid phosphodiester backbone, separating two annealed nucleic acid
30 strands (e.g. DNA, RNA, or RNA-DNA hybrid) using energy derived from hydrolysis of NTPs or dNTPs. Based on the presence of defined helicase motifs, it is possible to attribute a helicase activity to a given protein. The skilled artisan is able to select suited enzymes with helicase activity for the use in a method according to the present invention. In a preferred embodiment the helicase is selected from the group comprising helicases from

different families: superfamily I helicases (e.g. dda, pcrA, F-plasmid traI protein helicase, uvrD), superfamily II helicases (e.g. recQ, NS3-helicase), superfamily III helicases (e.g. AAV rep Helicase), helicases from DnaB-like superfamily (e.g. T7 phage helicase) or helicases from Rho-like superfamily.

5

The amplification methods will comprise buffers, dNTPs or NTPs in addition to the enzymes required.

As used herein, the term "dNTP" refers to deoxyribonucleoside triphosphates. Non-limiting examples of such dNTPs are dATP, dGTP, dCTP, dTTP, dUTP, which may also be present in the form of labelled derivatives, for instance comprising a fluorescence label, a radioactive label, a biotin label. dNTPs with modified nucleotide bases are also encompassed, wherein the nucleotide bases are for example hypoxanthine, xanthine, 7-methylguanine, inosine, xanthosine, 7-methylguanosine, 5,6-dihydrouracil, 5-methylcytosine, pseudouridine, dihydrouridine, 5-methylcytidine. Furthermore, ddNTPs of the above-described molecules are encompassed in the present invention.

10
15

As used herein, the term "NTP" refers to ribonucleoside triphosphates. Non-limiting examples of such NTPs are ATP, GTP, CTP, TTP, UTP, which may also be present in the form of labelled derivatives, for instance comprising a fluorescent label, a radioactive label, a biotin label.

20

In one embodiment the nucleic acid sequences of the first nucleic acid and second template nucleic acid stem from the same organism.

25

In one embodiment the nucleic acids are DNA.

In one embodiment the nucleic acid sequences of the first and second template nucleic acid are of human origin and genomic DNA.

30

Human genomic DNA is a sample comprising at least a fraction of human genomic DNA. Said sample may additionally comprise other biomolecules such as proteins, mitochondrial DNA or lipids but is ideally free of such biomolecules. In a preferred embodiment the human genomic DNA is isolated from a tissue or a cell line and is in soluble form in water

or in a buffer system. However the human genomic DNA may also be lyophilized when starting the WGA amplification. It is important that the respective individual has a normal chromosomal set including all chromosomes.

- 5 In a preferred embodiment said second template nucleic acid stems from a nucleic acid mixture from various individuals of a given organism or various human individuals. These can be humans, e.g. male and female or only male or only female. Other sources are also possible. Genomic DNA is preferred.
- 10 Ideally, said second template nucleic acid was generated using two or more rounds of whole genome amplification.

Preferably, the comparison is done with two or more amplification products from said second template nucleic acid, wherein the two or more amplification products are
15 generated from different starting concentrations of said second template nucleic acids.

For example, using real-time PCR techniques, fluorescence is detected and measured in the real-time PCR thermocycler, and its geometric increase corresponding to exponential increase of the product is used to determine the threshold cycle (C_T) in each reaction. The
20 unknown and each of the standards (amplification product from a second nucleic acid; the WGA DNA) are amplified in separate tubes. A standard curve (plot of C_T value/crossing point against log of amount of standard) is generated using different dilutions of the standard. The C_T value of the unknown samples is compared with the standard curve, allowing calculation of the initial amount of the target. It is important to select an
25 appropriate standard for the type of nucleic acid to be quantified. To generate a standard curve, ideally at least 5 different amounts of the standard should be quantified, and the amount of unknown target should fall within the range of the standard curve. Hence, in one embodiment also the above quantification steps are performed.

Hence, and preferably, the different starting concentrations of said second template nucleic
30 acid have been adjusted in such a way that they reflect a substantially linear range of concentrations.

In one embodiment the second template nucleic acid comprises primarily human nucleic acid sequences and its origin is selected from one of the following tissue types or cell

culture types respectively, blood, whole blood, or fractions of blood or isolated nucleated blood cells, buccal swabs, saliva, sputum, urine, other human sample types like biopsies, or other parts of the human body upon availability for isolation of DNA.

- 5 In the case of forensic DNA amplification assays a representative set of DNA sources of non human origin can be used to exclude and/or document cross reactivity (see table 1). Hence, in one embodiment, these are the source of the first and second nucleic acid. This may include DNA from phylogenetically closely related species like primates as well as farm animals and pets which live in closeness to men. Non-steril human micro-
- 10 environments which are sources of forensic DNA samples are skin, mouth (e.g. buccal swabs, sputum), oesophagus, stomach, vagina, gut, colon and faeces. Table 1 includes some major microbial species which are frequently found within samples prepared from healthy persons.

15 **Table 1: Collection of species for template DNA.**

Trivial (common) name	Scientific name	Source or location
horse	<i>Equus caballus</i>	farm animal, meat
cattle	<i>Bos taurus</i>	farm animal, meat
cat	<i>Felis catus</i>	pet, laboratory animal
dog	<i>Canis lupus familiaris</i>	pet, laboratory animal
mouse	<i>Mus musculus</i>	pet, laboratory animal
rat	<i>Rattus norvegicus</i>	pet, laboratory animal
rabbit	<i>Oryctolagus cuniculus</i>	farm animal, meat
sheep	<i>Ovis aries</i>	farm animal, meat
golden (Syrian) hamster	<i>Mesocricetus auratus</i>	pet, laboratory animal
guinea pig	<i>Cavia aperea</i>	pet, meat, laboratory animal
pig	<i>Sus scrofa</i>	farm animal, meat
orangutan	<i>Pongo pygmaeus</i>	non-human Ape
chimpanzee	<i>Pan troglodytes</i>	non-human Ape
Western gorilla	<i>Gorilla gorilla</i>	non-human Ape
bonobo	<i>Pan paniscus</i>	non-human Ape
chicken	<i>Gallus gallus</i>	farm animal, meat
	<i>Acinetobacter johnsonii</i>	human normal skin
	<i>Aspergillus niger</i>	soil, air
	<i>Bacillus subtilis</i>	soil, air
	<i>Bacteroides fragilis</i>	human stool, distal gut
	<i>Bifidobacterium longum</i>	human stool, distal gut
	<i>Candida albicans</i>	human skin, vagina
	<i>Corynebacterium tuberculostearicum</i>	human normal skin
	<i>Corynebacterium coyleiae</i>	human normal vaginal microflora
	<i>Corynebacterium singulare</i>	human normal skin
	<i>Enterococcus faecalis</i>	human normal vaginal, faeces
	<i>Escherichia coli</i>	human skin, stool

	<i>Finegoldia</i> AB109769	human normal skin
	<i>Gemella haemolysans</i>	human normal oral mriicroflora
	<i>Granulicatella adiacens</i>	human normal oral mriicroflora
	<i>Lactobacillus crispatus</i>	human normal vaginal microflora
	<i>Lactobacillus panis</i>	human normal vaginal microflora
	<i>Lactobacillus rhamnosus</i>	human intestine, stool and dairy products
	<i>Methanobrevibacter smithii</i>	human stool, distal gut
	<i>Neisseria gonorrhoe</i>	human skin
	<i>Neisseria spp.</i>	human normal oral mriicroflora
	<i>Peptostreptococcus harei</i>	human normal vaginal microflora
	<i>Peptostreptococcus vaginalis</i>	human normal vaginal microflora
	<i>Propionibacterium acnes</i>	human normal skin
	<i>Pseudomonas stuzeri</i>	human normal skin
	<i>Saccharomyces cerevisiae</i>	human skin
	<i>Staphylococcus aureus</i>	human skin, oral flora
	<i>Staphylococcus epidermitis</i>	human normal vaginal and skin microflora
	<i>Streptococcus mitis</i>	human normal oral and skin microflora
	<i>Veillionella pravulum</i>	human normal oral mriicroflora

Microbial type strains deposited at microbial strain collections (e.g. American Type Culture Collections, ATCC; German Collection of Microorganisms and Cell Cultures, DSMZ) or pre-characterized field isolates may be used as sources for the DNA as well.

5 Due to the fact that microbial genomes are 100 (yeast) to 1000 fold (bacteria) smaller than mammalian once, less DNA amounts are needed. Typically, DNA amounts which are equivalent to at least 10⁵ genome copies per PCR are sufficient. In some embodiments mixtures of defined DNA samples from microbials are used within one reaction vessel to safe reagents. DNA from these may also be used.

10

The region amplified from the second nucleic acid template (the WGA generated template) is ideally a region in SEQ ID NO. 1. The inventors have astonishingly found that this sequence and/or sequences that share sequence similarity with it may be found many times in the human genome. Hence, ideally primers and/or probes are used that bind this

15 sequence.

The sequences distributed throughout the genome are not all exactly identical. It is important that the selected primers bind also to the nearly identical sequences. Thus, ideally the locus shares at least 60 %, 70 %, 80 %, 90 % or even 95 % or 98 % sequence identity to a sequence according to SEQ ID NO. 1 over a stretch of 80 base pairs.

20

The determination of percent identity between two sequences is accomplished using the mathematical algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA (1993) 90:

5873-5877). Such an algorithm is the basis of the BLASTN and BLASTP programs of Altschul *et al.* (J. Mol. Biol. (1990) 215: 403-410). BLAST nucleotide searches are performed with the BLASTN program, score = 100, word length = 12, to obtain nucleotide sequences homologous to SEQ ID NO. 1.

5 It is an aspect of the invention that multiple copies are amplified from the second WGA generated nucleic acid. Ideally, at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 copies on various chromosomes are amplified. SEQ ID NO. 1 or sequences very similar thereto are present 29 times in the human genome. This is not just astonishing but provides for the power of the present method. Also, the
10 copies may be found on various chromosomes such as 1, 4, 5, 7, 11, 16.

The primers ideally used for amplification have a nucleotide sequence that differs from SEQ ID NO. 2, 3, 5, 6, 8, 9, 10, 11 and 12 by no more than 5 nucleotides over a stretch of 18 nucleotides.

Oligonucleotide primers may be prepared using any suitable method, such as, for example,
15 the phosphotriester and phosphodiester methods or automated embodiments thereof. In one such automated embodiment diethyl phosphoramidites are used as starting materials and may be synthesized as described by Beaucage *et al.* (1981) Tetrahedron Letters 22:1859-1862. One method for synthesizing oligonucleotides on a modified solid support is described in US Patent No. 4,458,006. It is also possible to use a primer, which has been
20 isolated from a biological source (such as a restriction endonuclease digest). Preferred primers have a length of about 6-100 bases, more preferably about 20-50, most preferably about 20-40 bases.

Additionally, the amplification reaction of the first nucleic acid to be amplified may in one
25 embodiment comprise an internal control (IC) nucleic acid.

The invention also relates to the use of a nucleic acid amplification mixture generated by whole genome amplification for quantifying a further nucleic acid composition.

30 First or second nucleic acid may comprise an inhibitor which has been added. Inhibitors of PCR are a serious problem in forensic case work. Although a variety of efficient DNA extraction and purification kits is commercially available which efficiently remove proteinaceous inhibitors (e.g. heparin) some water-soluble compounds may overcome the purification steps. Furthermore, analysis of forensic stains with low amounts of DNA

requires the concentration of DNA extracts from a huge amount of sample material. Wilson gives an overview of inhibitors (Wilson IG. Inhibition and Facilitation of Nucleic Acid Amplification, Appl Environ Microbiol 63, pp. 3741–3751, 1997). Water-soluble model substances can be used as standards for validation purposes. The following
5 compounds are preferred according to the invention: A potent mixture of inhibitors which is extracted from soil or other environmental sample is humic acid (CAS registry number 1415-93-6); hemin (chloro [3,7,12,17-tetramethyl-8,13-divinylporphyrin-2,18-dipropanoato(2-)] iron(III)) (CAS registry number 14875-96-8) or haematin (CAS registry number 15489-90-4) are degradation products of haemoglobin derived from blood
10 samples. Indigo carmine (CAS registry number 860-22-0) resembles the colour of cotton jeans. Bile salts (equimolar mixture of sodium salts of cholic acid and desoxycholic acid) are water soluble extracts of faeces. Other examples are known to those who are skilled in the art.

15 Extracts of natural and synthetic products or mixtures thereof are attractive inhibitors, too. For example hot aqueous extracts (boiling in deionized water for 20 min, supernatant after centrifugation at 12,000 x g for 10 min at room temperature) of cigarette ash or newspapers (inhibition by printing ink) can be standardized for validation purposes.

20 For producing the kit according to the invention it may be necessary to stabilise the DNA and the other components. This is done with trehalose, saccharose, glycerin, polymers such as ficoll, dextran, polyvinylpyrrolidon, polyacrylamide, other substances such as prolin, ectoin, betain, trimethylammoniumchlorid.

25

Figure Captions

Fig. 1

30 Shows data obtained from an experiment with a quantitative real-time PCR using the DNA isolated from Leucocytes (Leuco DNA) and the WGA DNA. In this experiment, a primer and a scorpion primer specific for human DNA were used to amplify a 146bp fragment of a region present on multiple autosomes (named 4NS1C) (Seq-ID # 13 and Seq ID #14). An internal control was coamplified in the same reaction, with primer and scorpions labelled

with a distinguishable dye. The Ct value remained constant for all conditions, demonstrating the success of the reaction and no inhibition present. Data for the IC assay are shown in Fig.2.

Surprisingly, the CT-values of the 4NS1C target using the Leuco DNA or the WGA DNA are highly similar. This result shows the reliability of WGA for amplifying high quantities of DNA for using in DNA quantification assays.

Fig. 2

Shows data for the internal control coamplified in the same reaction in the setup shown in Fig.1, with primer and scorpions labelled with a distinguishable dye. In order to check for successful PCR we also added a primer and a scorpion primer specific for an artificial target (named IC or Internal Control). The artificial template for the IC was added in the same concentration in all reaction wells. The CT-values of the IC are not dependent on which template DNA was used for the 4NS1C target, demonstrating the success of the reaction and no inhibition present, and showing suitability of WGA DNA as a quantification standard.

Fig. 3

Shows the data of real-time PCR experiments comparing quantification results for three different methods. In order to test whether WGA DNA is suitable for quantification purposes we used the WGA DNA as standard to quantify the NIST reference material 2372 (human DNA quantitation standard). This standard has an allegedly known concentration (Component A 52,4 ng/ μ l, Component B 53,6 ng/ μ l and Component C 54,3 ng/ μ l). These three components were diluted to the theoretical concentration of 0,25 ng/ μ l and 0,025 ng/ μ l. These DNA solutions were then quantified using the WGA DNA as a standard by using a quantitative real-time PCR experiment with an amplification reaction composition comprising, (i) Tris-HCl at a pH of between 8 and 8.8 (at 20°C) and/or, (ii) potassium salt selected from the group of, potassium chloride and potassium sulphate and/or, (iii) an ammonium salt, preferably ammonium chloride or ammonium sulphate and/or, (vi) magnesium chloride and/or, (v) a hot-start polymerase and (vi) the oligonucleotides with SEQ ID NO. 13 and SEQ ID NO. 14 the inventors showed that the DNA obtained by WGA is a good substitute for human DNA.

The measured concentrations are also reported in Table 1. All measured concentrations are within the range of $\pm 25\%$ of the theoretical concentration, this error lies within the normal ranges for real-time qPCR. Standard curves were run in duplicates, Test samples (the NIST DNA dilutions) were run in triplicates.

5 For comparison, accuracy of quantification was also determined using commercially available real-time PCR based DNA quantification kits, the Plexor HY kit from Promega and the Quantifiler Human from Applied Biosystems. For both kits, the control DNA coming with the kit was used as quantification standard and the kit used according to the supplied manual. The same dilutions for the NIST DNAs A,B, and C as above were used
10 as unknown Test samples and quantification results were calculated and are shown in Fig 3. It becomes obvious, that the WGA DNA provides at least as accurate quantification results as the conventional DNA controls coming with the two kits. The obtained quantification values for the NIST DNA vary by up to $\pm 50\%$ from the theoretical value for both methods, the Plexor HY and the Quantifiler Human, but not for the WGA method.

15

Fig. 4

Target Sequences for the oligonucleotides given. The first underlined region is the Ht target, the second is the 4NS1C target.

20

Fig. 5

Table with Oligonucleotide sequences

Examples

The quantification of DNA may be performed using different techniques; herein in one example real-time PCR was added. Different commercial kits are available for this purpose. Especially in the forensic field the quantification of human DNA using real-time PCR is a common method.

Of course, the quality of the control DNA, used for the standard curve in the quantification, has a big influence on the quantification result. The problem with control DNAs in currently available commercial kits is the high variability between different lots. Our solution of the problem is to use WGA (whole genome amplification) to amplify a mixture of human DNAs in order to reduce the variability and provide constant results over a long period of time.

In a quantitative real-time PCR experiment using an amplification reaction composition comprising, (i) Tris-HCl at a pH of between 8 and 8.8 (at 20 °C) and/or, (ii) potassium salt selected from the group of, potassium chloride and potassium sulphate and/or, (iii) an ammonium salt, preferably ammonium chloride or ammonium sulphate and/or, (iv) magnesium chloride and/or, (v) a hot-start polymerase and (vi) the oligonucleotides with SEQ ID NO. 11 and SEQ ID NO. 12 the inventors showed that the DNA obtained by WGA is a good substitute for human DNA.

In the given example, Human DNA from Promega (Cat.Nr. G3041) was used as a template for two rounds of WGA (the product of the first WGA was used as a template for a second WGA). The WGAs were carried out using the REPLI-g Midi Kit following manufacturer's guidelines. The purification after each WGA step was carried out using the QIAamp DNA Blood Maxi Kits following manufacturer's guidelines.

After DNA purification, the obtained quantity of WGA DNA was measured with a photometer (A260). This WGA DNA was compared to a DNA isolated from human leucocytes (QIAamp DNA Blod Kit, Qiagen).

As shown in the experiment below (Fig. 1), the result of a duplex quantitative real-time PCR using the DNA isolated from leucocytes (Leuco DNA) and the WGA DNA was compared. In this experiment, a primer and a scorpion primer specific for human DNA

were used to amplify a 146 bp fragment of a region present on multiple autosomes (named 4NS1C/SEQ ID NO. 1; see Fig. 4). In order to check for successful PCR we also added a primer and a scorpion primer specific for an artificial target (named IC or Internal Control). The artificial template for the IC was added in the same concentration in all reaction wells. The Ct-values of the IC are not dependent on which template DNA was used for the 4NS1C target.

4NS1C (SEQ ID NO. 12)	Primer	5'-CCGGGAAGCAGAAGGTGG-3'
4NS1C (SEQ ID NO. 11)	Scorpion	5'-Fam-CGAGCTCAGTTGTGCCTGTAGAGCTCG -dabcyl-C18-ACCTCTTCCTCTTGGCTGGG-3'

Surprisingly, the Ct-values of the 4NS1C target using the Leuco DNA or the WGA DNA are highly similar (Fig. 1). This result shows the reliability of WGA for amplifying high quantities of DNA for using in DNA quantification assays.

When isolating the DNA from natural samples (buccal swabs, blood, etc) the obtained yield using different methods varies. The typical yield for DNA isolation from healthy whole blood is approximately 4–12 µg using 200 µl blood (source QIAamp DNA Blood Mini Kit). The WGA - typical DNA yields per 50 µl reaction are up to 10 µg (Mini Kit) using 1 to 10 ng of template DNA (source REPLI-g Mini Kit). That means that performing WGA the overall yield of DNA isolation using 200 µl blood can be enhanced over a factor of at least 10,000 from 12 µg to 120,000 µg DNA with a constant DNA quality.

The great advantage of WGA DNA is therefore the possibility to produce big lots of a DNA template with a stable and very constant quality for several years.

The highly stable quality of the control DNA in DNA quantification assays will be a great advantage for scientists using this assay, since current limitations of fluctuations in quality and quantity from production lot to production lot is eliminated (http://marketing.appliedbiosystems.com/images/enews/ForensicNews_Vol5/PDF/01_WhatsNew_Letter.pdf)

In order to test whether WGA DNA is suitable for quantification purposes we used the WGA DNA as standard to quantify the NIST reference material 2372 (human DNA

quantitation standard) (see Fig. 3). This standard has an allegedly known concentration (Component A 52,4 ng/ μ l, Component B 53,6 ng/ μ l and Component C 54,3 ng/ μ l). These three components were diluted to the theoretical concentration of 0,25 ng/ μ l and 0,025 ng/ μ l. These DNA solutions were then quantified using the WGA DNA as a standard. The measured concentrations are reported in Table 1. All measured concentrations are within the range of ± 25 % of the theoretical concentration, this error lies within the normal ranges for real-time qPCR. For comparison, accuracy of quantification was also determined using commercially available real-time PCR based DNA quantification kits (see Fig.3), the Plexor HY kit from Promega and the Quantifiler Human from Applied Biosystems. For both kits, the control DNA coming with the kit was used as quantification standard and the kit used according to the supplied manual. The same dilutions for the NIST DNAs A,B, and C as above were used as unknown Test samples and quantification results were calculated and are shown in Fig 3. It becomes obvious, that the WGA DNA provides at least as accurate quantification results as the conventional DNA controls coming with the two kits. The obtained quantification values for the NIST DNA vary by up to ± 50 % from the theoretical value for both methods, the Plexor HY and the Quantifiler Human, but not for the WGA method.

Table 1 Measured concentrations

	theoretical concentration [ng/ μ l]	measured concentration [ng/ μ l]	Standard Deviation	theoretical concentration [ng/ μ l]	measured concentration [ng/ μ l]	Standard Deviation
Component A	0,25	0,227443	0,020820265	0,025	0,0245599	0,001413008
Component B	0,25	0,236829333	0,022564614	0,025	0,031362133	0,003076462
Component C	0,25	0,270458	0,041526476	0,025	0,028518267	0,005086746

CLAIMS

1. Method for quantifying one or more nucleic acids of a genome in a sample
5 comprising the steps of,
 - a. amplifying a first nucleic acid to be quantified,
 - b. determining the amount of said first nucleic acid by comparison of the amount of amplification product from said first nucleic acid with at least one amplification product from a second template nucleic acid,
 - 10 c. wherein said second template nucleic acid was generated using whole genome amplification and wherein the starting concentration of the second template nucleic acid is known.
2. Method according to claim 1, wherein the nucleic acid sequences of the first
15 nucleic acid and second template nucleic acid stem from the same organism.
3. Method according to claims 1 or 2, wherein the nucleic acids are DNA.
4. Method according to claims 1 to 3, wherein the nucleic acid sequences of the first
20 and second template nucleic acid are of human origin.
5. Method according to claims 1 to 4, wherein said second template nucleic acid stems from a nucleic acid mixture from various individuals of a given organism or various human individuals.
25
6. Method according to any of the preceding claims, wherein said second template nucleic acid was generated using two or more rounds of whole genome amplification.
- 30 7. Method according to any of the preceding claims, wherein the comparison is done with two or more amplification products from said second template nucleic acids, wherein the two or more amplification products are generated from different starting concentrations of said second template nucleic acids.

8. Method according to claim 7, wherein the different starting concentrations of said second template nucleic acid have been adjusted in such a way that they reflect a substantially linear range of concentrations.
- 5 9. Method according to any of the preceding claims, wherein the second template nucleic acid comprises primarily human nucleic acid sequences and its origin is selected from one of the following tissue types or cell culture types respectively, blood, whole blood, or fractions of blood or isolated nucleated blood cells, buccal swabs, saliva, sputum, urine, other human sample types like biopsies, or other parts
10 of the human body upon availability for isolation of DNA.
10. Method according to any of the preceding claims, wherein the amplification reaction of the first nucleic acid to be amplified comprises an internal control nucleic acid.
- 15 11. Method according to any of the preceding claims, wherein the amplification process is a real-time amplification process.
- 20 12. Method according claim 11, wherein the Ct value obtained from the first nucleic acid is compared with the standard curve generated using different dilutions of the second template nucleic acid.
- 25 13. Use of a nucleic acid amplification mixture generated by whole genome amplification for quantifying a further nucleic acid composition.
- 30 14. Kit for quantifying nucleic acids, comprising one or more vials with a nucleic acid composition that has been generated by means of whole genome amplification.

FIGURES

FIG. 1




FIG. 2




FIG. 3




FIG. 4

<p>SEQ ID NO. 1 target sequence</p>	<p>5' - TCAACAGGCCACCGTGAGGGAGGAGCTGGGCCGCACGCGGGCTGCTGG GAGGCAGGCAGGGACTTGGCCCCGAGAGGCCCGCCGTGGGGGCAAGAGC TGGGCCTGGAGAGGCCCTGGGAGGCAAGGGCGGGGCTGCAGAGGCT GTTCTCCAACCAGTGCTAGAACTGTACAGGCCACCAGGAGGCAGGAGG TGGGCCCTCAGAGCTTGGCTGGAGAAAAGTTCGGGGCCTACAAAGGCGG TTGGGAGCTGGGCAGGAGTTGAGCCAAAAGAGCTTGCTTACTTGCTGG GAGGCAGGGCCGGGAGAGCCCGACTTCAGGACAACCTGGGCCTGCGGC AGTCGCCGGGAGGCCAACCTTGGCGTGGAGGAGCCACCGACCGGAG ACCATTTGGGGCTGGAGATGCCATCGGAGGGCAGGAGCTCATCCTGG AGAGGCCACCGTGAGGCCTGACCTGGGCCTGGGGAGCTTGGCTTGAGG AAGCTGTGGGCCGACCAAGGCCGCCAGGAGATGGGTAGGCACTGAGTC CAAAGAGGTTGTTGAGAGGCAGGAATCGGGCCTGGAGACCCAACCAGG AAGAAGAGCTGGGCCCGGAGAGAATGCACGGAGGGTGCAAGTGGGTCT GGAGAGGCCGACTTGAGGAGGTTCTGGGCCCGGAGAGGCCCGCCGGAAG GGAAAACTGGGCCTGGAAAGGCCGTTGTGAGGAATGAGCCCCATGGG CCTGAAGAGGGCCACTGGCAGGCCGGGAGCTGGGCCTGCCGAAGCGGCCG AGAGGCAGGAGCTTTGGACTCGGGAGGCCGCAGTGAAGCAACAGCTAG CTGGGCGTGGAGAGTCCGCTGTGAGGCAGAGGCTGGGCCTGTGCAGGC CTTCGGGAGGCAGGAGGCTGGGCCTTGTGAGGCCTGCAGAGGCCACC GAAAGTCAAAGCGGGCTTGGGAAGGCCCGGGGAGGCATGAGCTGG GCTGGGCCGAAAGAGGCCACTGGGAGGCAGGAGGAGCTGGGCCTGGAG AGGCTGCCAAAAGGCAGGAGCTTCGCCTGAGGATGCCACAGTGAGACA CCATCTGGGTCTGGAGGGTCCACTGTGAGGCAGAGGCTGACCTGTAGA GTCCGACAGTAGACAGAAGTTGGGCAAAAAGCCTGATTTGAGGAAGTTT TGGGCTTCAAGAGTCAGCCACGAGGCAGGCACTAGGCCTGGAAATGGC CTCACAGTCATGAGTTGGGCCTAAATGGGCCACTGTGAGGGAGGAGCT GTGCCTGTTGAGGCTGCTGGCAGGCAGGCAGAAATTTGGCCTGGGGCA GCTGCCATGAGGCAAGAGCTGGGCCTGGAAAAAGCCCTGGGAGGCAA GAGCAGGGCCTGCAGAGGCTGTTCTCAAGTCAAAGCTGGGCCTGTTGA TGCCACCGGAAGCAGAAGGTGGGCCTGGAGAGTTTGAAGTTGAGGAAG TTTTGGGCCTACATTTGGCCGCCATGAGCTGGACAGGAAGTTGGGCCAAA AAAGGCTGTTGTGAGGCAGCAGTTGTGCCTGTAGACCCAGCCAAGAGG AAGAGGTGGGTCTGGAGAAGCCCCCATGAGGCAGAGGTTGGGCCTGTA GACGCTGACAGGAGGCAGGAGCTGGGCCTGGACAGGTCAACTTGAGGA GATTTTGGGCCTTCATAGGCCACCAGGAGGCAGTAGTTGGGACTAGAG AGTCTGACTTGAGTAAGTTTTTGGGCCGGAGATGACGTCCTGGGACAG GAGTTGGGCGTGGAGAGGCCACCGTGAGGCATAAGCTGGATGTAGAGA GGCCAGTGTGAGGCAAGACCTGGGCCTGTCTAGGCTGCTGGGAGACAG GCAGGAATCTGGCCAGGGAAGGTTGCCATGAGACAAAAGTTGGGCCTG GAAAGGCCCTTGTGAAGCATGAGCTTGGCCTAAAGAGGCCACTGGGTG GCAGGAGCTGGGTGTGTAGAAGCTGCTGAAAGGTTGGGAGCTTGGCTT GGGGGTCCACAGTGAGGTAGATGCTGGGCCT-3'</p>
---	--

FIG. 5

SEQ ID NO. 2 hT-For1 (primer)	5' -AGTGGGTCTGGAGAGGCCGACTTG-3'
SEQ ID NO. 3 hT-Rev1 (primer)	5' -TCAGGCCCATGGGGCTCATTCCT-3'
SEQ ID NO. 4 hT-Pro1 (probe)	5' -FAM-TTCTGGGCCCGGAGAGGCCGC-BHQ1-3'
SEQ ID NO. 5 hT-For2 (primer)	5' -GCAGAAGGTGGGCCTGGAGAGTTTGAC-3'
SEQ ID NO. 6 hT-Rev2 (primer)	5' -CCTTTTTTGGCCCAGTTCCTGTCCAGC-3'
SEQ ID NO. 7 hT-Pro2 (probe)	5' -FAM-GGAAGTTTTGGGCCTACATTGGCCGCATG-BHQ1-3'
SEQ ID NO. 8 hT-For3 (primer)	5' -AAGGTGGGCCTGGAGAGTTT-3'
SEQ ID NO. 9 hT-Rev3 (primer)	5' -CCTTTTTTGGCCCAGTTCCTGT-3'
SEQ ID NO. 10 hT-Pro3 (probe)	5' -HEX-AAGTTTTGGGCCTACATTGGCCGC-BHQ1-3'
SEQ ID NO. 11 4NS1C Scorpion	5' -Fam-CGAGCTCAGTTGTGCCTGTAGAGCTCG-dabcyl-C18- ACCTCTTCCTCTTGGCTGGG-3'
SEQ ID NO. 12 4NS1C primer	5' -CCGGAAGCAGAAGGTGG-3'

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/052913

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12Q1/68
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ARRIOLA EDURNE ET AL: "Evaluation of Phi29-based whole-genome amplification for microarray-based comparative genomic hybridisation", LABORATORY INVESTIGATION, vol. 87, no. 1, January 2007 (2007-01), pages 75-83, XP002638241, ISSN: 0023-6837	1-4, 7-10,13, 14
Y	abstract Materials and methods ----- -/--	6

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search 24 May 2012	Date of mailing of the international search report 04/06/2012
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Nurmi, Jussi
--	--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2012/052913

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	RENARD A ET AL: "Application of whole genome amplification and quantitative PCR for detection and quantification of spoilage yeasts in orange juice", INTERNATIONAL JOURNAL OF FOOD MICROBIOLOGY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 126, no. 1-2, 15 August 2008 (2008-08-15), pages 195-201, XP023315616, ISSN: 0168-1605, DOI: DOI:10.1016/J.IJFOODMICRO.2008.05.021 [retrieved on 2008-05-24]	1-3,5,7,8,11-14
Y	abstract; figure 2 page 196, column 2, paragraph 2 - page 197, column 2, paragraph 1	6
X	----- HANSEN HELEN M ET AL: "DNA quantification of whole genome amplified samples for genotyping on a multiplexed bead array platform", CANCER EPIDEMIOLOGY BIOMARKERS & PREVENTION, vol. 16, no. 8, August 2007 (2007-08), pages 1686-1690, XP002638242, ISSN: 1055-9965	14
A	abstract page 1687, column 2, paragraph 2	1-11
A	----- WALKER J A ET AL: "Multiplex polymerase chain reaction for simultaneous quantitation of human nuclear, mitochondrial, and male Y-chromosome DNA: application in human identification", ANALYTICAL BIOCHEMISTRY, ACADEMIC PRESS INC, NEW YORK, vol. 337, no. 1, 1 February 2005 (2005-02-01), pages 89-97, XP004707212, ISSN: 0003-2697, DOI: DOI:10.1016/J.AB.2004.09.036	1-12
X	abstract page 92, column 2, paragraph 2 - page 93, column 1, paragraph 2	
X	----- HANSON ERIN K ET AL: "Whole genome amplification strategy for forensic genetic analysis using single or few cell equivalents of genomic DNA", ANALYTICAL BIOCHEMISTRY, vol. 346, no. 2, November 2005 (2005-11), pages 246-257, XP002638243, ISSN: 0003-2697	14
A	abstract	1-11
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/052913

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VALLONE P M ET AL: "Development and usage of a NIST standard reference material for real time PCR quantitation of human DNA", FORENSIC SCIENCE INTERNATIONAL: GENETICS SUPPLEMENT SERIES, ELSEVIER IRELAND LTD, vol. 1, no. 1, 1 August 2008 (2008-08-01), pages 80-82, XP024526726, ISSN: 1875-1768, DOI: DOI:10.1016/J.FSIGSS.2007.10.215 [retrieved on 2008-06-02] abstract	1-12
X	----- LOVMAR L ET AL: "Quantitative evaluation by minisequencing and microarrays reveals accurate multiplexed SNP genotyping of whole genome amplified DNA", NUCLEIC ACIDS RESEARCH, OXFORD UNIVERSITY PRESS, SURREY, GB, vol. 31, no. 21, 1 January 2003 (2003-01-01), pages (E129)1-9, XP002994728, ISSN: 0305-1048, DOI: DOI:10.1093/NAR/GNG129 abstract	14
A	----- CHIA DENNIS J ET AL: "Distinct Alterations in Chromatin Organization of the Two IGF-I Promoters Precede Growth Hormone-Induced Activation of IGF-I Gene Transcription", MOLECULAR ENDOCRINOLOGY, vol. 24, no. 4, April 2010 (2010-04), pages 779-789, XP002638244, ISSN: 0888-8809	1-11
X	----- page 787, column 2, lines 7-9	14
Y	-----	6