



HU000028121T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 028 121**(13) **T2**

EURÓPAI SZABADALOM

SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 10 786915**
- (22) A bejelentés napja: **2010. 06. 11.**
- (96) Az európai bejelentés bejelentési száma:
EP 20100786915
- (97) Az európai bejelentés közzétételi adatai:
EP 2442650 A1 **2010. 12. 16.**
- (97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2442650 B1 **2015. 08. 26.**
- (51) Int. Cl.: **A61K 314/73** (2006.01)
A01N 43/42 (2006.01)
A61K 47/02 (2006.01)
A61P 25/16 (2006.01)
A61P 15/10 (2006.01)
A61K 31/44 (2006.01)
- (86) A nemzetközi (PCT) bejelentési szám:
PCT/US 10/038336
- (87) A nemzetközi közzétételi szám:
WO 10144817

(30) Elsőbbségi adatok: 186445 P 2009. 06. 12. US	(73) Jogosult(ak): Cynapsus Therapeutics Inc., Toronto, ON M6J 1C9 (CA)
(72) Feltaláló(k): GIOVINAZZO, Anthony, John, Oakville ON L6H 0B9 (CA) HEDDEN, David, Bruce, Ann Arbor MI 48103 (US) DE SOMER, Marc, L., Ann Arbor, Michigan 48103 (US) BRYSON, Nathan, John, Toronto ON M4W 2V8 (CA)	(74) Képviselő: dr. Török Ferenc, DANUBIA Szabadalmi és Jogi Iroda Kft., Budapest

(54) **Szublingvális apomorfin**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 442 650 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
26.08.2015 Bulletin 2015/35

(51) Int Cl.:
A61K 31/473 ^(2006.01) **A61K 31/44** ^(2006.01)
A01N 43/42 ^(2006.01) **A61K 47/02** ^(2006.01)
A61P 15/10 ^(2006.01) **A61P 25/16** ^(2006.01)

(21) Application number: **10786915.8**

(86) International application number:
PCT/US2010/038336

(22) Date of filing: **11.06.2010**

(87) International publication number:
WO 2010/144817 (16.12.2010 Gazette 2010/50)

(54) **SUBLINGUAL APOMORPHINE**
SUBLINGUALES APOMORPHIN
APOMORPHINE SUBLINGUALE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO SE SI SK SM TR
Designated Extension States:
BA

- **DE SOMER, Marc, L.**
Ann Arbor, Michigan 48103 (US)
- **BRYSON, Nathan, John**
Toronto
ON M4W 2V8 (CA)

(30) Priority: **12.06.2009 US 186445 P**

(74) Representative: **Høiberg A/S**
St. Kongensgade 59 A
1264 Copenhagen K (DK)

(43) Date of publication of application:
25.04.2012 Bulletin 2012/17

(56) References cited:
WO-A1-98/26763 **WO-A1-98/48781**
WO-A1-2006/120412 **WO-A2-2007/075422**
US-A1- 2004 028 613 **US-A1- 2004 204 440**
US-A1- 2005 031 677 **US-A1- 2007 149 479**
US-A1- 2008 119 504 **US-B2- 6 566 368**

(60) Divisional application:
15175258.1

(73) Proprietor: **Cynapsus Therapeutics Inc.**
Toronto, ON M6J 1C9 (CA)

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

(72) Inventors:
• **GIOVINAZZO, Anthony, John**
Oakville
ON L6H 0B9 (CA)
• **HEDDEN, David, Bruce**
Ann Arbor
MI 48103 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 442 650 B1

Description

Background of the Invention

[0001] The invention relates to compositions including apomorphine and formulated for sublingual administration and the use of such compositions for the treatment of Parkinson's disease.

[0002] Parkinson's disease (PD) is a progressive degenerative disease of the central nervous system. The risk of developing Parkinson's disease increases with age, and afflicted individuals are usually adults over 40. Parkinson's disease occurs in all parts of the world, and affects more than 1.5 million individuals in the United States alone.

[0003] While the primary cause of Parkinson's disease is not known, it is characterized by degeneration of dopaminergic neurons of the substantia nigra. The substantia nigra is a portion of the lower brain, or brain stem that helps control voluntary movements. The shortage of dopamine in the brain caused by the loss of these neurons is believed to cause the observable disease symptoms.

[0004] The symptoms of PD vary from patient to patient. The most common symptom is a paucity of movement and rigidity, characterized by an increased stiffness of voluntary skeletal muscles. Additional symptoms include resting tremor, bradykinesia (slowness of movement), poor balance, and walking problems. Common secondary symptoms include depression, sleep disturbance, dizziness, stooped posture, dementia, problems with speech, breathing, and swallowing. The symptoms become progressively worse with time and ultimately result in death.

[0005] A variety of therapeutic treatments for PD are available. Perhaps the best known is levodopa, a dopamine precursor. While levodopa administration can result in a dramatic improvement in symptoms, patients can experience serious side-effects, including nausea and vomiting. Concurrent carbidopa administration with levodopa is a significant improvement, with the addition of carbidopa inhibiting levodopa metabolism in the gut, liver and other tissues, thereby allowing more levodopa to reach the brain. Other dopamine agonists, such as bromocriptine, pergolide, pramipexole, and amantadine are also used to treat Parkinson's disease, and can be administered to PD patients either alone or in combination with levodopa.

[0006] Many patients develop involuntary choreiform movements which are the result of excessive activation of dopamine receptors. These movements usually affect the face and limbs and can become very severe. Such movements disappear if the dose of dopamine precursor (e.g., levodopa) or dopamine agonist is reduced, but this typically causes rigidity to return. Moreover, the margin between the beneficial and the unwanted effects appear to become progressively narrower as the period of chemotherapeutic treatment lengthens.

[0007] A further complication of long-term chemotherapeutic treatment with dopamine agonists is the development of rapid fluctuations in clinical state where the patient switches suddenly between mobility and immobility for periods ranging from a few minutes to a few hours. The fluctuations are of several general types. "Wearing-off" phenomena are deteriorations in the relief afforded by a dose of levodopa before the next dose takes effect (Van Laar T., CNS Drugs, 17:475 (2003)). Because they are related to a patient's dose schedule, such periods are often relatively predictable (Dewey RB Jr., Neurology, 62(suppl 4):S3-S7 (2004)). In contrast, "on-off" phenomena are sudden transitions from an "on" period of levodopa benefit to an "off" period of akinesia, rigidity, and tremor that occur in minutes or even seconds, (Swope DM., Neurology, 62(suppl 4):S27-S31 (2004)) with no discernible relation to a patient's dose schedule. Two other phenomena are the delayed "on" effect, in which levodopa's effects are substantially delayed, and dose failure (also known as the no-"on" or skipped-dose effect), in which no effects occur at all. These various "off" states can produce such an abrupt loss of mobility that the patient may suddenly stop while walking or be unable to rise from a chair in which he had sat down normally a few moments earlier.

[0008] Subcutaneous injections of apomorphine have proved to be effective in the treatment of "on-off" fluctuations in Parkinson's disease within 5 to 15 minutes, and last for 45 to 90 minutes. Trials have shown consistent reversal of "off" period akinesia, a decrease in daily levodopa requirements and consequently a decrease in the amount of "on" period dyskinesias. Advantages over other dopamine agonists include a quick onset of action and lower incidence of psychological complications. For a "rescue therapy" in patients with "on-off" fluctuations, apomorphine also has the advantage over other dopamine agonists that it has a relatively short half-life.

[0009] Numerous formulations and routes of administration for apomorphine have been studied and apomorphine therapy has been found to be hampered by various complications. For example, oral administration of apomorphine tablets has required high doses to achieve the necessary therapeutic effect because apomorphine administered by this route undergoes extensive metabolism in the small intestine and/or, upon absorption, in the liver; sublingual administration of apomorphine tablets caused severe stomatitis on prolonged use with buccal mucosal ulceration in half the patients treated (see Deffond et al., J. Neurol. Neurosurg. Psychiatry 56:101 (1993)); and intranasal administration produced transient nasal blockage, burning sensation and swollen nose and lips (see Koller et al., Neurology 62:S22 (2004)). While subcutaneous injections of apomorphine have proven effective, an injection by needle is difficult for Parkinson's patients because of impaired motor function. Furthermore, a common side effect of subcutaneous injection is the development of nodules, which often become infected, necessitating antibiotic treatment or surgical debridement.

(see Prietz et al., J. Neurol. Neurosurg. Psychiatry 65:709 (1998)).

[0010] WO 2006/120412 (Amarin Pharmaceuticals Ireland) teaches a two-component kit including (a) an acid addition salt of apomorphine, and (b) a liquid solution including a pH modifying agent. The components of the kit are combined at the point of use to form a composition for buccal administration including apomorphine at a mildly alkaline pH. The document teaches neutralization at the point of use to (i) aid in the sublingual absorption of apomorphine, while (ii) allowing the apomorphine to be kept in an acidified form to prevent oxidation prior to use.

[0011] WO 98/26763 (Lohmann Therapie Systems LTS) teaches film compositions comprising apomorphine which are adapted for buccal delivery. The exemplified compositions include significant quantities (16-19% by weight) of "acidifying agent", and so might be expected to exacerbate acid-induced stomatitis in subjects receiving repeated administrations. There is a need for new formulations of apomorphine and apomorphine prodrugs which are safe, effective, and easy for a Parkinson's patient to use.

Summary of the Invention

[0012] The invention relates to pharmaceutical compositions and uses as described in the claims. The compositions can be useful for the treatment of Parkinson's disease, sexual dysfunction, and depressive disorders therewith.

[0013] In a first aspect, the invention features a pharmaceutical composition in unit dosage form (a film, or a strip) formulated for sublingual administration, the unit dosage form having a first portion including an acid addition salt of apomorphine and a second portion including a pH neutralizing agent (e.g., polyamines, calcium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide, calcium carbonate, iron carbonate, magnesium carbonate, zinc carbonate, sodium acetate, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium phosphate tribasic, and mixtures thereof, or any other suitable base).

[0014] In certain embodiments, the unit dosage form is a film or a strip and wherein the unit dosage form includes a mucoadhesive polymer. The mucoadhesive polymer can be, without limitation, any mucoadhesive polymer described herein.

[0015] In one particular embodiment, the acid addition salt of apomorphine is apomorphine hydrochloride. In another particular embodiment, the acid addition salt of apomorphine is protonated apomorphine complexed to an anionic polyelectrolyte (e.g., alginates, carrageenan, xanthan gum, polyacrylate, or carboxymethylcellulose).

[0016] In still another embodiment, the pharmaceutical

composition is a film or a strip, wherein the first portion is a first layer and the second portion is a second layer, the first layer being acidic and including the acid addition salt of apomorphine and the second layer including the pH neutralizing agent.

[0017] In certain embodiments, the sublingual formulation includes an antioxidant. The antioxidant can be, without limitation, any antioxidant described herein.

[0018] In particular embodiments, the sublingual formulation includes a first portion that is a film including a solid solution of an acid addition salt of apomorphine and includes a second portion that is a particulate base on or within the unit dosage form. The particulate base can include, for example, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium phosphate tribasic, or a mixture thereof.

[0019] In still another embodiment, the first portion of the unit dosage form is separated from the second portion of the unit dosage form by a barrier (e.g., a film separating an acidic layer from a basic layer in a multi-layered film, or a coating on a particulate base or apomorphine particle contained within the unit dosage form). The barrier can be neutral in pH (e.g., between 6 and 7.8), separating the acidic first portion from the basic second portion of the unit dosage form.

[0020] In a related aspect, the pharmaceutical composition formulated for sublingual administration includes apomorphine particles having an effective particle size of from 20 nm to 10 μm , wherein the apomorphine particles include an acid addition salt of apomorphine.

[0021] In certain embodiments, the pharmaceutical composition is in a unit dosage form selected from a film, or a strip.

[0022] The sublingual formulations can include apomorphine particles having an effective particle size of from 1 μm to 10 μm (e.g., an effective particle size of from 1 μm to 9 μm , from 1 μm to 8 μm , from 1 μm to 7 μm , from 1 μm to 6 μm , from 1 μm to 5 μm , from 2 μm to 10 μm , from 3 μm to 10 μm , from 4 μm to 10 μm , from 2 μm to 7 μm , or from 2 μm to 6 μm).

[0023] In certain other embodiments, the sublingual formulations can include apomorphine particles having an effective particle size of from 20 nm to 1 μm (e.g., an effective particle size of from 20 nm to 1 μm , from 40 nm to 1 μm , from 60 nm to 1 μm , from 80 nm to 1 μm , from 100 nm to 1 μm , from 20 nm to 800 nm, from 20 nm to 700 nm, from 50 nm to 700 nm, from 40 nm to 800 nm, from 60 nm to 800 nm, from 100 nm to 800 nm, from 60 nm to 700 nm, from 60 nm to 600 nm, from 100 nm to 600 nm, from 150 nm to 800 nm, or from 150 nm to 600 nm).

[0024] In still other embodiments, the sublingual formulation includes a mucoadhesive polymer. The mucoadhesive polymer can be, without limitation, any mucoadhesive polymer described herein.

[0025] In certain embodiments, the sublingual formulation includes apomorphine particle and the apomorphine particle include an acid addition salt of apomorphine. The acid addition salt can be apomorphine hydrochloride or any acid addition salt described herein.

[0026] In a particular embodiment, the sublingual formulation in unit dosage form is a film or a strip including a mucoadhesive polymer. The mucoadhesive polymer can be, without limitation, any mucoadhesive polymer described herein. For example, the film or strip can include a first layer and a second layer, the first layer being acidic and including the apomorphine particles and the second layer including a pH neutralizing agent (e.g., polyamines, calcium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide, calcium carbonate, iron carbonate, magnesium carbonate, zinc carbonate, sodium acetate, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium phosphate tribasic, and mixtures thereof, or any other suitable base).

[0027] In certain embodiments, the sublingual formulation includes an antioxidant. The antioxidant can be, without limitation, any antioxidant described herein.

[0028] In another aspect, the invention features a pharmaceutical composition formulated for sublingual administration including protonated apomorphine complexed to an anionic polyelectrolyte (e.g., alginates, carrageenan, xanthan gum, polyacrylate, or carboxymethylcellulose). In certain embodiments, the sublingual formulation includes an antioxidant. The antioxidant can be, without limitation, any antioxidant described herein.

[0029] In an embodiment of any of the above pharmaceutical compositions, the pharmaceutical composition is in a unit dosage form including from 2 to 40 mg of an acid addition salt of apomorphine (e.g., from 2 to 5 mg, 4 to 10 mg, 6 to 15 mg, 8 to 20 mg, 10 to 25 mg, 12 to 30 mg, 20 to 35 mg, 25 to 40 mg, or 30 to 40 mg of an acid addition salt of apomorphine). For example, each unit dosage form can contain 3 ± 1 mg, 4 ± 1 mg, 5 ± 1 mg, 8 ± 2 mg, 10 ± 3 mg, 12 ± 3 mg, 15 ± 3 mg, 22 ± 4 mg, 27 ± 4 mg, 30 ± 5 mg, 35 ± 5 mg, or 40 ± 5 mg of an acid addition salt of apomorphine.

[0030] In another embodiment of any of the above pharmaceutical compositions, the unit dosage form when placed in 1 mL of unbuffered water at pH 7 results in a solution having a pH of between 7.4 and 9.1 (e.g., a pH of between 7.4 and 8.8, 7.4 and 8.3, 7.6 and 8.8, 7.6 and 8.5, 8.2 and 8.5, 8.4 and 8.7, 8.6 and 8.8, or 8.7 and 9.1).

[0031] In still another embodiment of any of the above pharmaceutical compositions, following sublingual administration to a subject the unit dosage form produces an average circulating concentration of at least 3 ng/mL within a period of from 5 to 15 minutes following the administration. For example, the unit dosage form can produce an average circulating concentration of from 3 to 6

ng/mL within 7 to 10 minutes, from 5 to 10 ng/mL within 5 to 10 minutes, from 7 to 12 ng/mL within 5 to 10 minutes, from 10 to 16 ng/mL within 5 to 10 minutes, from 3 to 6 ng/mL within 7 to 15 minutes, from 5 to 10 ng/mL within 7 to 15 minutes, from 7 to 12 ng/mL within 7 to 15 minutes, from 10 to 16 ng/mL within 7 to 15 minutes, from 3 to 6 ng/mL within 15 to 20 minutes, from 5 to 10 ng/mL within 15 to 20 minutes, from 7 to 12 ng/mL within 15 to 20 minutes, or from 10 to 16 ng/mL within 15 to 20 minutes following the administration.

[0032] The invention further features pharmaceutical compositions for use in the treatment of Parkinson's disease in a mammal by sublingually administering a pharmaceutical composition of the invention to the mammal in an amount effective to treat the mammal.

[0033] The invention also features pharmaceutical compositions for use in alleviating dyskinesia in a mammal afflicted with Parkinson's disease by sublingually administering a pharmaceutical composition of the invention to the mammal in an amount effective to alleviate the dyskinesia.

[0034] The invention also features pharmaceutical compositions for use in alleviating akinesia in a mammal afflicted with Parkinson's disease by sublingually administering a pharmaceutical composition of the invention to the mammal in an amount effective to alleviate the akinesia.

[0035] The invention features pharmaceutical compositions for use in the treatment of sexual dysfunction in a mammal by sublingually administering a pharmaceutical composition of the invention to the mammal in an amount effective to treat the mammal.

[0036] The invention also features pharmaceutical compositions for use in the treatment of a depressive disorder in a mammal by sublingually administering a pharmaceutical composition of the invention to the mammal in an amount effective to treat the mammal.

[0037] In one embodiment of any of the above uses, the uses further includes administration of an effective amount of an anti-emetic agent (e.g., nicotine, lobeline sulfate, pipamazine, oxypendyl hydrochloride, ondansetron, buclizine hydrochloride, cyclizine hydrochloride, dimenhydrinate, scopolamine, metopimazine, benzaunamine hydrochloride, or diphenidol hydrochloride).

[0038] In an embodiment of any of the above methods and compositions, the apomorphine salt is a racemic mixture of R and S isomers, or enriched in R isomer (i.e., the ratio of R to S isomer for all of the apomorphine in the composition, or all the apomorphine being administered, is from 5:1 to 1,000:1, from 10:1 to 10,000:1, or from 100:1 to 100,000:1, or over all apomorphine isomers in the composition is at least 98% R isomer, 99% R isomer, 99.5% R isomer, 99.9% R isomer, or is free of any observable amount of S isomer.

[0039] The term "administration" or "administering" refers to a method of giving a sublingual dosage of apomorphine to a patient.

[0040] As used herein, the term "apomorphine particle"

refers to microparticles or nanoparticles containing acid addition salts of apomorphine.

[0041] As used herein, the term "average circulating concentration" refers to the average plasma concentration of apomorphine at time t observed for a group of subjects following sublingual administration of a particular unit dosage form of the invention. For example, among 20 subjects the average circulating concentration of apomorphine 10 minutes following sublingual administration of the unit dosage form can be at least 3 ng/mL, 5 ng/mL, 7 ng/mL, 9 ng/mL, 11 ng/mL, 13 ng/mL, or 15 ng/mL, depending upon the amount of apomorphine in the unit dosage.

[0042] By "depressive disorder" is meant any psychological or psychiatric disorder associated with symptoms of depressed mood. Treatable depressive disorders may be characterized by an inhibition or reduction of dopaminergic function in the nucleus accumbens, e.g., major depression, dysthymia, bipolar disorder (manic depression), and post-traumatic stress disorder. As used herein, the terms "effective particle size" and "particle size" are used interchangeably and refer to a mixture of particles having a distribution in which 50% of the particles are below and 50% of the particles are above a defined measurement. The "effective particle size" refers to the volume-weighted median diameter as measured by a laser/light scattering method or equivalent, wherein 50% of the particles, by volume, have a smaller diameter, while 50% by volume have a larger diameter. The effective particle size can be measured by conventional particle size measuring techniques well known to those skilled in the art. Such techniques include, for example, sedimentation field flow fractionation, photon correlation spectroscopy, light scattering (e.g., with a Microtrac UPA 150), laser diffraction, and disc centrifugation.

[0043] As used herein, "pH neutralizing agent" refers to any basic component present in the unit dosage forms of the invention. The pH neutralizing agents which can be used in the unit dosage forms of the invention include organic bases (e.g., amines), inorganic bases (e.g., oxides, hydroxides, carbonates, or phosphates), and mixtures thereof. The pH neutralizing agent is typically present in an amount sufficient to produce a solution having a pH of between 7.4 and 9.1 when the unit dosage form is placed in 1 mL of unbuffered water at pH 7.

[0044] As used herein, "sexual dysfunction" refers to disorders of orgasm, response timing, ejaculation, nociception, congestive arousal and erection, vasculogenic impairment, or desire. In males, the form of sexual dysfunction is typically erectile dysfunction, the inability to achieve and sustain an erection sufficient for intercourse. Females also can have sexual dysfunctions of arousal and orgasm that increase with age and are associated with the presence of vascular risk factors and onset of menopause. Some of the vascular and muscular mechanisms that contribute to penile erection in the male are believed to involve similar vasculogenic factors in female genital responses. Female sexual dysfunction includes

a failure to attain or maintain vaginal lubrication-swelling responses of sexual excitement until completion of the sexual activity.

[0045] As used herein, the term "treating" refers to administering a pharmaceutical composition for prophylactic and/or therapeutic purposes. To "prevent disease" refers to prophylactic treatment of a patient who is not yet ill, but who is susceptible to, or otherwise at risk of, a particular disease. To "treat disease" or use for "therapeutic treatment" refers to administering treatment to a patient already suffering from a disease to ameliorate the disease and improve the patient's condition. Thus, in the claims and embodiments, treating is the administration to a mammal either for therapeutic or prophylactic purposes.

[0046] Other features and advantages of the invention will be apparent from the following detailed description and the claims.

Detailed Description

[0047] The invention features sublingual formulations of apomorphine acid addition salts. The formulations can be useful for the treatment of Parkinson's disease, sexual dysfunction, and depressive disorders therewith. In certain aspects, the invention features (i) sublingual formulations in unit dosage form having a first portion including an acid addition salt of apomorphine and a second portion including a pH neutralizing agent.

[0048] Fluctuations in motor disability and dyskinesias are a significant problem in the long-term treatment of Parkinson's disease. In the later stages of Parkinson's disease, many patients develop severe "off" episodes where, despite continuing to take their medication, they experience periods when they lose the ability to move (e.g., the patients develop bradykinesia (slowed movement) or akinesia (inability to move)). These "off" episodes typically occur 3 to 4 times per day.

[0049] Apomorphine has a rapid onset of action which is ideal for use as a rescue therapy for intractable "off" periods in Parkinson's disease.

[0050] Using the sublingual formulations of the invention, a subject suffering from the effects of middle stage or late stage Parkinson's disease may be able to recognize the onset of their "off" symptoms and be capable of administering a sublingual dose of a formulation of the invention to alleviate the dyskinesia associated with such "off" episodes. The sublingual formulations are easy for a subject with compromised motor skills to administer and can relieve a Parkinson's patient from the need for a caregiver, who might otherwise be needed to administer an injectable dosage form of apomorphine at the onset of an "off" episode.

[0051] The sublingual formulations of the invention can increase the bioavailability of apomorphine, prolong the stability of apomorphine, and/or improve the safety and efficacy of apomorphine therapy. The formulations can produce a rapid uptake of apomorphine into the subject,

allowing dyskinesia episodes to be self-treated. Furthermore, the convenience with which these sublingual formulations can be self administered provides a significant advantage to severely ill patients, such as those with middle stage or late stage Parkinson's disease.

[0052] Additional details of how to make and use the sublingual formulations of the invention are provided below and in the Examples.

Monolayer and Bilayer Films

[0053] The films of the invention are not dissimilar to the films used, for example, to make the Listerine® PocketPak® mouth fresheners. In PocketPak films the polymers used are typically polysaccharide-based or polysaccharide and glycoprotein-based gums such as pullulan, pectin, locust bean gum, xanthan gum, sodium alginate, gum Arabic and the like. These same polymers can be used in the films of the invention.

[0054] The films can include one layer, two layers, or more. If in two layers, the one adapted to adhere to mucosal tissue may be referred to as the "adhesive layer." With two layers, the outer layer can be less adhesive or non-adhesive, and can provide protection against mechanical agitation, such as agitation by a user's tongue. The components of the outer layer might be, of themselves, less dissolvable than the components of an adhesive layer. However, in the aggregate, the film shall dissolve in that it will transition to fully dissolved parts or parts that will be carried away by normal cleaning processes at the mucosal tissue in question. In forming two layers, diffusion or the forming process itself may provide a gradient in component amounts in the transition between the two layers. The two layers can be utilized to separate components (e.g., an apomorphine-containing acidic layer and a buffered pH neutralizing layer), which together enhance absorption of the apomorphine but are otherwise incompatible in a formulation requiring long term stability (i.e., shelf life). Alternatively, the unit dosage form of the invention can be a monolayer film that is an apomorphine-containing acidic layer which is coated with or impregnated with a particulate base. The particulate base can be incorporated into the monolayer film using the methods described in PCT Publication No. WO/2009/052421, U.S. Patent Publication No. 20060210610. The film of the invention can include an effervescent particulate (i.e., a particulate carbonate base). Such effervescent films can be prepared as described in U.S. Patent Publication No. 20010006677.

[0055] The polymers used in the films of the invention can be polymers that affect the rate of hydration or mucosal adhesion properties of an adhesive layer. Such polymers can be, for example, carboxymethylcellulose, cellulose acetate, ethylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose (HPMC, such as Pharmacoat 606™, Shin-Etsu Chemical Company Ltd., Japan), nitrocellulose, polyoxyethylene/polyoxypropylene polymers, copolymers or block copolymers, polyvinylpyrro-

lidone polymers or derivatives, and gums. The average molecular weight of the polymer can be selected based on the swelling and dissolution profile sought. Mixtures of less soluble and/or less swellable polymers with more soluble or swellable polymers can help transition the film to a sufficiently dissolved form. For example, the film can include carbamer, polyethylene oxide, ethylcellulose, titanium oxide and colorant (such as F, D and C blue lake colorant). Often the film is formed using a pharmaceutically appropriate solvent such as ethanol, water, mixtures, or the like. Such solvents are typically largely evaporated away prior to use. Optionally, the films comprise a blend of more than one polymers or more than one molecular weight of a given set of polymers in order to control the rate of hydration, physical properties and mechanical properties.

Basic Layers

[0056] The multi-layered films of the invention can include a film formed from a basic polymer. Polyamines which can be used in the unit dosage forms of the invention include homo and copolymers of dimethylaminoethyl-acrylate, dimethylaminoethyl-methacrylate, dimethylaminopropyl-acrylate, dimethylaminopropyl-methacrylate, or other similar amino-functionalized acrylate, chitosan or partially hydrolyzed chitin in a substantially basic form, homo and co polymers of polyethyleimine, polylysine, polyvinylimidazole, or polyvinylamine. In certain embodiments the polyamine is Eudragit E100.

Other Components

[0057] Plasticizers, penetration enhancers, flavoring agents, preservatives, odorants, coloring agents, and the like can be included in the unit dosage forms of the invention.

[0058] Plasticizers will generally modify the feel, softness, flexibility (in an un-wetted state) of the unit dosage forms of the invention. Penetration enhancers may, in some cases, act as plasticizers. Examples of plasticizers include, without limitation, glycerol, propylene glycol, fatty acid esters, such as glyceryl oleate, polyalcohols, sorbitan esters, citric acid esters, polyethylene glycol (e.g., PEG 400), polyvinyl alcohol, polyvinyl methyl ether, triacetin; mannitol, xylitol, and sorbitol. The plasticizer can be present in any suitable range, including, for example about 0.5% to 30%, 10% to 20%, or 15% to 18% by weight of the dry film.

[0059] Permeation enhancers can be used to improve the permeability of the apomorphine at the mucosal membrane in the unit dosage forms of the invention. One or more permeation enhancers may be used to modulate the rate of mucosal absorption of the apomorphine. Any effective permeation enhancers may be used including, for example, bile salts, such as sodium cholate, sodium glycocholate, sodium glycodeoxycholate, taurodeoxycholate, sodium deoxycholate, sodium lithocholate chen-

ocholate, chenodeoxycholate, ursocholate, ursodeoxycholate, hyodeoxycholate, dehydrocholate, glycochenocholate, taurochenocholate, and taurochenodeoxycholate; sodium dodecyl sulfate (SDS), dimethyl sulfoxide (DMSO), N-lauroyl sacrosine, sorbitan monolaurate, stearyl methacrylate, N-dodecylazacycloheptan-2-one, N-dodecyl-2-pyrrolidinone, N-dodecyl-2-piperidinone, 2-(1-nonyl)-1,3-dioxolane, N-(2-methoxymethyl) dodecylamine, N-dodecylethanolamine, N-dodecyl-N-(2-methoxymethyl)acetamide, 1-N-dodecyl-2-pyrrolidone-5-carboxylic acid, 2-pentyl-2-oxo-pyrrolidineacetic acid, 2-dodecyl-2-oxo-1-pyrrolidineacetic acid, 2-dodecyl-2-oxo-1-pyrrolidineacetic acid, 1-azacycloheptan-2-one-dodecylacetic acid, menthol, propylene glycol, glycerol monostearate, sorbitol monolaurate, glycerol dilaurate, tocopherol acetate, phosphatidyl choline, glycerol, polyethyleneglycol, monoglycerides, diglycerides, triglycerides, lecithin, tween surfactants, sorbitan surfactants, sodium lauryl sulfate; salts and other derivatives of saturated and unsaturated fatty acids, surfactants, bile salt analogs, derivatives of bile salts, or such synthetic permeation enhancers as described in U.S. Pat. No. 4,746,508.

[0060] A flavoring agent and/or odorant can be added to the unit dosage forms of the invention to make them more palatable. At least one flavoring agent or odorant composition may be used. Any effective flavor or odor may be rendered. The flavoring agents may be natural, artificial, or a mixture thereof. The flavoring agent gives a flavor that is attractive to the user. In one embodiment, the flavoring agent may give the flavor of mint, honey lemon, orange, lemon lime, grape, cranberry, vanilla berry, Magnasweet™, bubble gum, or cherry. The flavoring agent can be natural or artificial sweetener, such as sucrose, xylitol, sodium saccharin, cyclamate, aspartame, acesulfame, and salts thereof.

[0061] Apomorphine is susceptible to oxidative degradation. To minimize oxidative degradation it is desirable that the formulations of the invention contain one or more antioxidants. Antioxidants that can be used in the films of the invention can be selected from thiols (e.g., aurothioglucose, dihydrolipoic acid, propylthiouracil, thioredoxin, glutathione, cysteine, cystine, cystamine, thiodipropionic acid), sulphoximines (e.g., buthionine-sulphoximines, homo-cysteine-sulphoximine, buthionine-sulphones, and penta-, hexa- and heptathionine-sulphoximine), metal chelators (e.g., α -hydroxy-fatty acids, palmitic acid, phytic acid, lactoferrin, citric acid, lactic acid, and malic acid, humic acid, bile acid, bile extracts, bilirubin, biliverdin, EDTA, EGTA, and DTPA), sodium metabisulfite, vitamins (e.g., vitamin E, vitamin C, ascorbyl palmitate, Mg ascorbyl phosphate, and ascorbyl acetate), phenols (e.g., butylhydroxytoluene, butylhydroxyanisole, ubiquinol, nordihydroguaiaretic acid, trihydroxybutyrophenone), benzoates (e.g., coniferyl benzoate), uric acid, mannose, propyl gallate, selenium (e.g., selenium-methionine), stilbenes (e.g., stilbene oxide and trans-stilbene oxide), and combinations thereof. The total amount of antioxidant included in the films can be from

0.001% to 3% by weight, preferably 0.01 % to 1% by weight, in particular 0.05% to 0.5% by weight, based on the total weight of the formulation.

[0062] In certain embodiments, the various components (e.g., plasticizers, penetration enhancers, flavoring agents, preservatives, odorants, coloring agents, particulate base, and apomorphine particles) included in the unit dosage forms of the invention can be combined and incorporated into a first portion that is acidic and includes the apomorphine or combined and incorporated into a second portion that is basic and includes a pH neutralizing component, or the components may be divided between the two portions. In some instances it may be desirable to minimize interaction between the acidic portion of the unit dosage form and the basic portion of the unit dosage form by including a barrier between the two. For example, a barrier can be included in the unit dosage forms of the invention as a third layer interposed between the acidic layer and the basic layer of a multilayer sublingual dosage form. Alternatively, the barrier can be a rapidly dissolving coating on the surface of a particulate component in the unit dosage form, such as a coated particulate base coated onto, or embedded within, an acidic portion of the unit dosage form. In still another approach, the barrier can be a rapidly dissolving coating on the surface of apomorphine particles in the unit dosage form, which further includes a basic portion. These approaches can be utilized to ensure that the apomorphine-containing acidic portion of the unit dosage form is not neutralized prior to the administration to a subject.

Apomorphine Particles

[0063] The pharmaceutical formulations described herein can include apomorphine particles having an effective particle size of from about 1 micron to about 10 microns. The starting apomorphine composition can be predominantly crystalline, predominantly amorphous, or a mixture thereof, and can include unmodified apomorphine.

[0064] In an alternative approach, the pharmaceutical formulations described herein can include apomorphine particles having an effective particle size of less than about 1 micron (i.e., nanoparticulate formulations). The starting apomorphine composition can be predominantly crystalline, predominantly amorphous, or a mixture thereof, and can include unmodified apomorphine.

[0065] These apomorphine particles can be made by using any method known in the art for achieving the desired particle sizes. Useful methods include, for example, milling, homogenization, supercritical fluid fracture, or precipitation techniques. Exemplary methods are described in U.S. Patent Nos. 4,540,602; 5,145,684; 5,518,187; 5,718,388; 5,862,999; 5,665,331; 5,662,883; 5,560,932; 5,543,133; 5,534,270; and 5,510,118; 5,470,583.

Milling to Obtain Submicron Apomorphine Particles

[0066] In one approach, the apomorphine or a salt thereof, is milled in order to obtain micron or submicron particles. The milling process can be a dry process, e.g., a dry roller milling process, or a wet process, i.e., wet-grinding. A wet-grinding process is described in U.S. Pat. Nos. 4,540,602, 5,145,684, 6,976,647 and EPO 498,482. Thus, the wet grinding process can be practiced in conjunction with a liquid dispersion medium and dispersing or wetting agents such as described in these publications. Useful liquid dispersion media include safflower oil, ethanol, n-butanol, hexane, or glycol, among other liquids selected from known organic pharmaceutical excipients (see U.S. Patent Nos. 4,540,602 and 5,145,684), and can be present in an amount of 2.0-70%, 3-50%, or 5-25% by weight based on the total weight of the apomorphine in the formulation.

[0067] The grinding media for the particle size reduction step can be selected from rigid media, typically spherical in shape, though non-spherical grinding media could also be used. The grinding media preferably can have a mean particle size from 1 mm to about 500 microns. For fine grinding, the grinding media particles can have a mean particle size from about 0.05 to about 0.6 mm. Smaller size grinding media will result in smaller size apomorphine particles as compared to the same conditions using larger sized grinding media. In selecting material, grinding media with higher density, e.g., glass (2.6 g/cm³), zirconium silicate (3.7 g/cm³), and zirconium oxide (5.4 g/cm³) and 95% zirconium oxide stabilized with yttrium, can be utilized for more efficient milling. Alternatively, polymeric grinding media can be used. Polymeric resins suitable for use herein are chemically and physically inert, substantially free of metals, solvent and monomers, and of sufficient hardness and friability to enable them to avoid being chipped or crushed during grinding. Suitable polymeric resins include, without limitation, crosslinked polystyrenes, such as polystyrene crosslinked with divinylbenzene, styrene copolymers, polycarbonates, polyacetals, such as Delrin™, vinyl chloride polymers and copolymers, polyurethanes, polyamides, poly(tetrafluoroethylenes), e.g., Teflon™, and other fluoropolymers, high density polyethylenes, polypropylenes, cellulose ethers and esters such as cellulose acetate, polyhydroxymethacrylate, polyhydroxyethyl acrylate, and silicone containing polymers such as polysiloxanes.

[0068] Grinding can take place in any suitable grinding mill. Suitable mills include an airjet mill, a roller mill, a ball mill, an attritor mill, a vibratory mill, a planetary mill, a sand mill and a bead mill. A high energy media mill is preferred when small particles are desired. The mill can contain a rotating shaft.

[0069] The preferred proportions of the grinding media, apomorphine the optional liquid dispersion medium, and dispersing, wetting or other particle stabilizing agents present in the grinding vessel can vary within wide limits

and depend on, for example, the size and density of the grinding media, the type of mill selected, the time of milling, etc. The process can be carried out in a continuous, batch or semi-batch mode. In high energy media mills, it can be desirable to fill 80-95% of the volume of the grinding chamber with grinding media. On the other hand, in roller mills, it frequently is desirable to leave the grinding vessel up to half filled with air, the remaining volume comprising the grinding media and the liquid dispersion media, if present. This permits a cascading effect within the vessel on the rollers which permits efficient grinding. However, when foaming is a problem during wet grinding, the vessel can be completely filled with the liquid dispersion medium or an anti-foaming agent may be added to the liquid dispersion.

[0070] The attrition time can vary widely and depends primarily upon the mechanical means and residence conditions selected, the initial and desired final particle size, among other factors. For roller mills, processing times from several days to weeks may be required. On the other hand, milling residence times of less than about 2 hours are generally required using high energy media mills. After attrition is completed, the grinding media is separated from the milled apomorphine particulate product (in either a dry or liquid dispersion form) using conventional separation techniques, such as by filtration, or sieving through a mesh screen.

[0071] To produce apomorphine particles having an effective particle size of less than about 1 micron, the grinding media can be made from beads having a size ranging from 0.05mm to 4 mm. For example, high energy milling of apomorphine with yttrium stabilized zirconium oxide 0.4 mm beads for a milling residence time of 25 minutes to 1.5 hours in recirculation mode at 1200 to 3000 RPM. In another approach, high energy milling of apomorphine with 0.1 mm zirconium oxide balls for a milling residence time of 2 hours in batch mode can be used. The milling concentration can be from about 10% to about 30% apomorphine by weight in comparison to the milling slurry weight, which can contain a wetting and/or dispersing agent to coat the initial suspension so a uniform feed rate may be applied in continuous milling mode. Alternatively, batch milling mode is utilized with a milling media containing an agent to adjust viscosity and/or provide a wetting effect so that the apomorphine is well dispersed amongst the grinding media.

Microprecipitation to Obtain Apomorphine Nanoparticles

[0072] Apomorphine particles can also be prepared by homogeneous nucleation and precipitation in the presence of a wetting agent or dispersing agent using methods analogous to those described in U.S. Patent Nos. 5,560,932 and 5,665,331. Such a method can include the steps of: (1) dispersing apomorphine in a suitable liquid media; (2) adding the mixture from step (1) to a mixture including at least one dispersing agent or wetting agent such that at the appropriate temperature, the apo-

morphine is dissolved; and (3) precipitating the formulation from step (2) using an appropriate anti-solvent. The method can be followed by removal of any formed salt, if present, by dialysis or filtration and concentration of the dispersion by conventional means. In one embodiment, the apomorphine particles are present in an essentially pure form and dispersed in a suitable liquid dispersion media. In this approach the apomorphine particles are a discrete phase within the resulting mixture. Useful dispersing agents, wetting agents, solvents, and anti-solvents can be experimentally determined.

Homogenization to Obtain Apomorphine Nanoparticles

[0073] Apomorphine particles can also be prepared by high pressure homogenization (see U.S. Patent No. 5,510,118). In this approach apomorphine particles are dispersed in a liquid dispersion medium and subjected to repeated homogenization to reduce the particle size of the apomorphine particles to the desired effective average particle size. The apomorphine particles can be reduced in size in the presence of at least one or more dispersing agents or wetting agents. Alternatively, the apomorphine particles can be contacted with one or more dispersing agents or wetting agents either before or after attrition. Other materials, such as a diluent, can be added to the apomorphine/dispersing agent mixture before, during, or after the size reduction process. For example, unprocessed apomorphine can be added to a liquid medium in which it is essentially insoluble to form a premix (i.e., about 0.1-60% w/w apomorphine and about 20-60% w/w dispersing agents or wetting agents). The apparent viscosity of the premix suspension is preferably less than about 1000 centipoise. The premix can then be transferred to a microfluidizer and circulated continuously first at low pressures, and then at maximum capacity (i.e., 3,000 to 30,000 psi) until the desired particle size reduction is achieved. The resulting dispersion of apomorphine particles can be spray coated onto a sublingual pharmaceutical formulation of the invention using techniques well known in the art.

Milling with Simethicone

[0074] Foaming during the nanosizing can present formulation issues and can have negative consequences for particle size reduction. For example, high levels of foam or air bubbles in the mill can cause a drastic increase in viscosity rendering the milling process inoperable. Even a very low level of air presence can dramatically reduce milling efficiency causing the desired particle size unachievable. This may be due to the resultant air in the mill cushioning the milling balls and limiting grinding efficiency. The air also can form a microemulsion with the milled ingredients which presents many issues with respect to the delivery of an accurate dose and palatability. Addition of a small amount of simethicone is a very effective anti-foaming agent which minimizes milling

variability or special handling techniques to avoid the introduction of air into the milling process.

The Use of Wetting and Dispersing Agents

[0075] The apomorphine particles can be prepared with the use of one or more wetting and/or dispersing agents, which are, e.g., adsorbed on the surface of the apomorphine particle. The apomorphine particles can be contacted with wetting and/or dispersing agents either before, during or after size reduction. Generally, wetting and/or dispersing agents fall into two categories: non-ionic agents and ionic agents. The most common non-ionic agents are excipients which are contained in classes known as binders, fillers, surfactants and wetting agents. Limited examples of non-ionic surface stabilizers are hydroxypropylmethylcellulose, polyvinylpyrrolidone, Pladone, polyvinyl alcohol, Pluronic, Tweens and polyethylene glycols (PEGs). Ionic agents are typically organic molecules bearing an ionic bond such that the molecule is charged in the formulation, such as long chain sulfonic acid salts (e.g., sodium lauryl sulfate and dioctyl sodium sulfosuccinate).

[0076] Excipients, such as wetting and dispersing agents, can be applied to the surface of the apomorphine nanoparticulate via spray drying, spray granulation, or spray layering process. These procedures are well known in those skilled in the art. It is also common to add additional excipients prior to removal of solvent in the nanoparticulate suspension to aid in the dispersion of the solid composition in medium in which the solid composition will be exposed (e.g. saliva) to further prevent agglomeration and/or particle size growth of the small apomorphine particles. An example of such an additional excipient is a redispersing agent. Suitable redispersing agents include, without limitation, sugars, polyethylene glycols, urea and quaternary ammonium salts.

Therapy

[0077] Representative examples of diseases and conditions treatable using the sublingual formulations of the invention are as listed hereinabove, and include, but are not limited to, Parkinson's disease, sexual dysfunction, and depressive disorders, such as major depression and bipolar disorder.

[0078] Sublingual formulations of the invention include rapidly disintegrating or dissolving dosage forms, also known as fast dissolve, fast or rapid melt, and quick disintegrating dosage forms. These dosage forms dissolve or disintegrate rapidly in the patient's mouth without chewing or the need for water within a short time frame. Because of their ease of administration, such compositions are particularly useful for the specific needs of patients with compromised motor skills.

[0079] The apomorphine may be administered in its free base form or as a pharmaceutically acceptable salt, such as a non-toxic acid addition salts or metal complex-

es that are commonly used in the pharmaceutical industry. Examples of acid addition salts include organic acids such as acetic, lactic, pamoic, maleic, citric, malic, ascorbic, succinic, benzoic, palmitic, suberic, salicylic, tartaric, methanesulfonic, toluenesulfonic, or trifluoroacetic acids or the like; polymeric acids such as tannic acid, carboxymethyl cellulose, alginic acid, or the like; and inorganic acid such as hydrochloric acid, hydrobromic acid, sulfuric acid phosphoric acid, or the like. Metal complexes include calcium, zinc, iron, and the like. In certain instances the formulation of the invention includes apomorphine hydrochloride.

[0080] The formulations can be administered to patients in therapeutically effective amounts. For example, an amount is administered which prevents, reduces, or eliminates the symptoms of Parkinson's disease, sexual dysfunction, or depression, respectively. Typical dose ranges are from about 2 mg to about 30 mg of apomorphine, or a salt thereof, given up to five times per day. The exemplary dosage of apomorphine to be administered is likely to depend on such variables as the type and extent of the condition, the overall health status of the particular patient, the particular form of apomorphine being administered, and the particular sublingual formulation being used.

[0081] Potential adverse effects can be ameliorated by administering apomorphine in combination with an anti-emetic agent, such as nicotine, lobeline sulfate, pipamazine, oxypendyl hydrochloride, ondansetron, buclizine hydrochloride, cyclizine hydrochloride, dimenhydrinate, scopolamine, metopimazine, benzaunamine hydrochloride or diphenidol hydrochloride. In certain instances it may be desirable to incorporate the anti-emetic into the sublingual formulation for simultaneous administration in combination with apomorphine.

[0082] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the methods and compounds claimed herein are performed, made, and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention.

EXAMPLE 1 - Two Layer Apomorphine Strip

Preparation of the first layer:

[0083] Gelatin and mannitol are dispersed in purified water and mixed thoroughly (i.e., using a vacuum mixer) and homogenized. Apomorphine hydrochloride is added and the mixture was again homogenized to ensure complete dissolution of the apomorphine hydrochloride. The pH of the solution is adjusted to about 3.0 (i.e., by addition of a suitable acid, such as citric acid). The solution is then poured onto a sheet and dried in a heated oven.

Preparation of the second layer:

[0084] Ethyl cellulose, poly(ethylene oxide), and hydroxypropylcellulose are dissolved in anhydrous ethanol.

To the resulting solution is added a pH modifying agent (i.e., calcium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide, calcium carbonate, iron carbonate, magnesium carbonate, zinc carbonate, sodium acetate, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium phosphate tribasic, and mixtures thereof). Alternatively, the pH modifying agent can be a polyamine, such as Eudragit E100.

[0085] A solvent-cast mucoadhesive film is prepared by casting a thin film of the solution onto the first layer. Evaporation of the solvent (ethanol) can be accomplished by drying at 60 °C for 30 minutes.

[0086] The resultant dry film includes (i) a first acidic layer containing apomorphine in a stable acid addition salt form (i.e., the hydrochloride salt), and (ii) a second basic layer (the adhesive layer) capable of neutralizing some of the apomorphine at the time of sublingual administration. The two layer film can enhance the bioavailability of the apomorphine, as absorption is enhanced when apomorphine hydrochloride is neutralized, without compromising the shelf life stability of the film.

[0087] The two-layer film is cut into strips, each strip containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The strips can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

EXAMPLE 2 (Comparative)- Single Layer Nanoparticulate Apomorphine Strip

[0088] Ethyl cellulose, poly(ethylene oxide), and hydroxypropylcellulose are dissolved in anhydrous ethanol to form a solution.

[0089] Nanoparticulate apomorphine hydrochloride is prepared by milling solid apomorphine hydrochloride as described herein.

[0090] The particulate apomorphine is suspended in the solution and a solvent-cast mucoadhesive film is prepared by casting a thin film of the mixture onto a sheet. Evaporation of the solvent (ethanol) can be accomplished by drying at 60 °C for 30 minutes.

[0091] The resultant dry film includes a single adhesive layer that releases nanoparticulate apomorphine. The apomorphine can penetrate the mucosal tissue in its nanoparticulate form, thus enhancing the bioavailability of the apomorphine in the film.

[0092] The single-layer nanoparticulate film is cut into strips, each strip containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The strips can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

orders.

EXAMPLE 3 (Comparative) - Single Layer Apomorphine Free Base Strip

[0093] All materials are degassed prior to use and all steps are carried out under a nitrogen atmosphere.

[0094] Ethyl cellulose, poly(ethylene oxide), and hydroxypropylcellulose are dissolved in anhydrous ethanol to form a polymer solution.

[0095] The apomorphine hydrochloride and an antioxidant (e.g., sodium metabisulfite) are dissolved in a minimal amount of water and added to the polymer solution. The pH of the resulting mixture is adjusted to about 9.0 (i.e., by addition of a suitable base, such as sodium hydroxide).

[0096] A solvent-cast mucoadhesive film is prepared by casting a thin film of the mixture onto a sheet. Evaporation of the solvent (ethanol/water) can be accomplished by drying at 60 °C for 30 minutes, and/or drying under reduced pressure.

[0097] The resultant dry film includes a single adhesive layer that releases free base apomorphine. The apomorphine can penetrate the mucosal tissue in its nanoparticulate form, thus enhancing the bioavailability of the apomorphine in the film.

[0098] The single-layer film is cut into strips, each strip containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The strips can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

EXAMPLE 4 - Two-Layer Apomorphine-Anionic Polyelectrolyte Complex

Preparation of the first layer:

[0099] Apomorphine (free base) is combined with alginic acid to form an apomorphine-alginate complex. To the complex is added gelatin, mannitol, and purified water. The mixture is mixed thoroughly (i.e., using a vacuum mixer) and homogenized. The pH of the solution is adjusted to about 3.0. The solution is then poured onto a sheet and dried in a heated oven.

Preparation of the second layer:

[0100] Ethyl cellulose, poly(ethylene oxide), and hydroxypropyl cellulose are dissolved in anhydrous ethanol. To the resulting solution is added a pH modifying agent (i.e., calcium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide, calcium carbonate, iron carbonate, magnesium carbonate, zinc carbonate, sodium acetate, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium

phosphate tribasic, and mixtures thereof). Alternatively, the pH modifying agent can be a polyamine, such as Eudragit E100.

[0101] A solvent-cast mucoadhesive film is prepared by casting a thin film of the solution onto the first layer. Evaporation of the solvent (ethanol) can be accomplished by drying at 60 °C for 30 minutes.

[0102] The resultant dry film includes (i) a first acidic layer containing apomorphine in a stable acid addition salt complex with alginic acid, and (ii) a second basic layer (the adhesive layer) capable of neutralizing some of the apomorphine at the time of sublingual administration. The two layer film can enhance the bioavailability of the apomorphine, as absorption is enhanced when apomorphine is neutralized, without compromising the shelf life stability of the film.

[0103] The two-layer film is cut into strips, each strip containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The strips can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

EXAMPLE 5 (Comparative)- Nanoparticulate Apomorphine Gel

[0104] Nanoparticulate apomorphine hydrochloride is prepared by milling solid apomorphine hydrochloride as described herein.

[0105] The particulate apomorphine is combined with sodium carboxymethylcellulose and glycerol. The resulting mixture is mixed thoroughly (i.e., using a vacuum mixer) and homogenized. Water is added to the mixture (with extensive mixing) to form a hydrogel containing suspended nanoparticles of apomorphine hydrochloride.

[0106] The nanoparticulate gel can be dispensed under the tongue in amounts containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The gel can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

EXAMPLE 6 (Comparative)- Apomorphine-Anionic Polyelectrolyte Complex Gel

[0107] Apomorphine (free base) is combined with alginic acid to form an apomorphine-alginate complex.

[0108] The apomorphine-alginate complex is combined with sodium carboxymethylcellulose, glycerol, and an antioxidant (e.g., sodium metabisulfite). The resulting mixture is mixed thoroughly (i.e., using a vacuum mixer) and homogenized. Water is added to the mixture (with extensive mixing) to form a hydrogel containing apomorphine-alginate complex.

[0109] The gel can be dispensed under the tongue in amounts containing the equivalent of from 2 mg to 20 mg of apomorphine in its free base form. The gel can be administered to a subject for the treatment of Parkinson's disease, sexual dysfunction, or depressive disorders.

EXAMPLE 7 - Bilayer Apomorphine with Polymer-Based Neutralizer

[0110] The following ingredients are mixed with 200 parts 2-1 water-ethanol solvent in an oxygen free environment: 40 parts apomorphine hydrochloride, 5 parts citric acid, 7 parts Methocel E5, 18 parts Methocel E50, 3 parts Klucel JF, 6 parts Sucralose, 3 parts PEG400, 3 parts sorbitol, 1 Prosweet G, 4 parts maltodextrin M180, 4 parts IPC B792, and 6 parts spearmint. The mixture is spread on a thin plastic liner and dried to produce a film of ca. 40 μm thickness.

[0111] Separately, Eudragit E100 is dissolved to form a viscous mixture in ethanol and acetone 1:1. The mixture is spread on a thin plastic liner to produce a film of ca. 24 μm thickness.

[0112] The apomorphine layer is fused to the Eudragit layer using heat (60 °C) and pressure to create a bilayer film. Individual dosing units of 40 mg apomorphine hydrochloride are obtained by cutting the film to 2.5×1 cm.

EXAMPLE 8 (Comparative)- Tablet with Apomorphine and Basic Agent

[0113] The following ingredients are blended: 40 parts apomorphine hydrochloride jet-milled to 10 μm (D95), 100 parts lactose, 100 parts microcrystalline cellulose, 5 parts sodium phosphate dibasic, 25 parts crosslinked povidone, 18 parts sucralose, 2 parts colloidal silica, 5 parts mint flavoring, and 5 parts magnesium stearate. Tablets of 300mg are pressed to provide tablets containing 40 mg of apomorphine.

[0114] Alternatively, the present ingredients can be segregated into an apomorphine containing mixture and second, sodium phosphate mixture, which are pressed into a bilayer tablet.

EXAMPLE 9 - Dispersed milled Apomorphine in Bilayer Thin Film

[0115] According to methods of a previous example, jet-milled powder of apomorphine hydrochloride (D95 < 10 μm) is added to a mixture of polyethyleneglycol, polyvinylpyrrolidone, sucralose, sorbitol and xylitol in ethanol-ethylacetate to create a homogeneous dispersion of the active ingredient. The mixture is spread on a thin plastic liner and dried to produce a film of ca. 40 μm thickness. This film can be administered as is or combined with a neutralizing layer as per previous examples. Also contemplated, is the addition of jet-milled sodium carbonate to the dispersion of ingredient prior to drying to create a single layer wherein both active apomorphine hydrochloride and a neutralizing agent are dispersed as solid agents within a single layer.

EXAMPLE 10 (Comparative) - Hard Candy Lozenge

[0116] A candy matrix is formed by mixing one cup

(240 grams) of sugar, 1/3 cup (81 cc) of light corn syrup, and slightly more than 1 cup (240 ml) of water. The matrix mixture is heated to a temperature of at least 285 °F, taking care to avoid stirring the mixture at temperatures greater than 200 °F to prevent uncontrolled crystallization of the sugar mixture. The matrix mixture is allowed to cool to 260 °F, and 4 ml of a flavoring agent and 1/8 teaspoon (0.625 cc) of citric acid are added, followed by the addition of 900 mg of milled apomorphine HCl and 1800 mg sodium phosphate dibasic. These ingredients are stirred thoroughly into the matrix, and the resulting mixture is poured into molds which have been sprayed with an anti-stick coating (e.g., sprayed with the anti-stick coating known under the trademark PAM). Sticks are inserted 2 minutes after pouring into the molds. Desirably, the hard candy lozenge is prepared under an inert atmosphere to minimize oxidation of the apomorphine.

[0117] It should be noted that the mixture contains no die, no alcohol, no synthetic flavor agents and no preservatives, and is completely natural.

EXAMPLE 11 - Monolayer Apomorphine with Particulate Base Neutralizer

[0118] The following ingredients are mixed with 200 parts 2-1 water-ethanol solvent in an oxygen free environment: 40 parts apomorphine hydrochloride, 5 parts citric acid, 7 parts Methocel E5, 18 parts Methocel E50, 3 parts Klucel JF, 6 parts Sucralose, 3 parts PEG400, 3 parts sorbitol, 1 Prosweet G, 4 parts maltodextrin M180, 4 parts IPC B792, and 6 parts spearmint. The mixture is spread on a thin plastic liner and dried to produce a film of ca. 40 μm thickness.

[0119] Separately, a pH neutralizing solid (e.g., sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium phosphate monobasic, sodium phosphate dibasic, sodium phosphate tribasic, potassium phosphate monobasic, potassium phosphate dibasic, potassium phosphate tribasic, or mixtures thereof) is milled to produce a microparticulate base. That the neutralizing solid is optionally coated with a neutral barrier, such as polyethylene glycol.

[0120] The apomorphine layer is coated with the microparticulate base, for example, by electrospray or static spray application. Optionally, the layer may be heated (ca. 60 °C) and the microparticulate base pressed into the apomorphine layer to form a monolayer film containing a solid solution of apomorphine hydrochloride with a microparticulate base dispersed within the film. Individual dosing units of 40 mg apomorphine hydrochloride are obtained by cutting the film to 2.5×1 cm.

Claims

1. A pharmaceutical composition in unit dosage form formulated for sublingual administration, wherein said unit dosage form is a film or a strip having a first

- portion comprising apomorphine particles comprising an acid addition salt of apomorphine and a second portion comprising a pH neutralizing agent.
2. The pharmaceutical composition of claim 1, wherein said film form comprises a polysaccharide. 5
 3. The pharmaceutical composition of any of claims 1-2, wherein said unit dosage form comprises a mucoadhesive polymer. 10
 4. The pharmaceutical composition of claim 3, wherein said mucoadhesive polymer comprises carboxymethylcellulose, cellulose acetate, ethylcellulose, hydroxyethylcellulose, or hydroxypropylmethylcellulose. 15
 5. The pharmaceutical composition of any of the preceding claims, further comprising an antioxidant. 20
 6. The pharmaceutical composition of any of the preceding claims, further comprising a permeation enhancer.
 7. The pharmaceutical composition of claim 6, wherein said permeation enhancer is glycerol monostearate. 25
 8. The pharmaceutical composition of any of the preceding claims, wherein said pH neutralizing agent is an organic base. 30
 9. The pharmaceutical composition of any of the preceding claims, wherein said first portion comprises apomorphine particles having an effective particle size of from 1 μm to 10 μm . 35
 10. The pharmaceutical composition of any of the preceding claims, wherein said acid addition salt of apomorphine is apomorphine hydrochloride. 40
 11. The pharmaceutical composition of any of claims 1-10, wherein said unit dosage form comprises from 2 to 40 mg of apomorphine, or an acid addition salt thereof. 45
 12. The pharmaceutical composition of any of claims 1-11 for use in the treatment of Parkinson's disease, dyskinesia, akinesia in a mammal afflicted with Parkinson's Disease. 50
 13. The pharmaceutical composition of any of claims 1-12 for use in the treatment of sexual dysfunction, or depressive disorder. 55

Patentansprüche

1. Pharmazeutische Zusammensetzung in Einheitsdo-

sierungsform, die zur sublingualen Verabreichung formuliert ist, wobei die Einheitsdosierungsform ein Film oder ein Streifen mit einem ersten Abschnitt ist, der Apomorphinpartikel umfasst, die ein Säureadditionssalz von Apomorphin umfassen, und einem zweiten Abschnitt, der ein pH-Neutralisationsmittel umfasst.

2. Pharmazeutische Zusammensetzung nach Anspruch 1, wobei die Filmform ein Polysaccharid umfasst.
3. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1-2, wobei die Einheitsdosierungsform ein mukoadhäsives Polymer umfasst.
4. Pharmazeutische Zusammensetzung nach Anspruch 3, wobei das mukoadhäsive Polymer Carboxymethylcellulose, Celluloseacetat, Ethylcellulose, Hydroxyethylcellulose oder Hydroxypropylmethylcellulose umfasst.
5. Pharmazeutische Zusammensetzung nach einem der vorhergehenden Ansprüche, die weiter ein Antioxidans umfasst.
6. Pharmazeutische Zusammensetzung nach einem der vorhergehenden Ansprüche, die weiter einen Permeationsverstärker umfasst.
7. Pharmazeutische Zusammensetzung nach Anspruch 6, wobei der Permeationsverstärker Glycerinmonostearat ist.
8. Pharmazeutische Zusammensetzung nach einem der vorhergehenden Ansprüche, wobei das pH-Neutralisationsmittel eine organische Base ist.
9. Pharmazeutische Zusammensetzung nach einem der vorhergehenden Ansprüche, wobei der erste Abschnitt Apomorphinpartikel mit einer effektiven Partikelgröße von 1 μm bis 10 μm umfasst.
10. Pharmazeutische Zusammensetzung nach einem der vorhergehenden Ansprüche, wobei das Säureadditionssalz von Apomorphin Apomorphinhydrochlorid ist.
11. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1-10, wobei die Einheitsdosierungsform von 2 bis 40 mg Apomorphin oder ein Säureadditionssalz davon umfasst.
12. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1-11 zur Verwendung bei der Behandlung von Parkinson-Krankheit, Dyskinesie, Akinesie bei einem Säugler, der unter der Parkinson-Krankheit leidet.

13. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1-12 zur Verwendung bei der Behandlung von sexueller Dysfunktion oder von depressiver Störung.

Revendications

1. Composition pharmaceutique sous la forme de dose unitaire formulée pour l'administration sublinguale, dans laquelle ladite forme de dose unitaire est un film ou une bande présentant une première partie comprenant des particules d'apomorphine comprenant un sel d'addition d'acide d'apomorphine et une seconde partie comprenant un agent de neutralisation du pH. 10
2. Composition pharmaceutique selon la revendication 1, dans laquelle ladite forme de film comprend un polysaccharide. 20
3. Composition pharmaceutique selon l'une quelconque des revendications 1 à 2, dans laquelle ladite forme de dose unitaire comprend un polymère muco-adhésif. 25
4. Composition pharmaceutique selon la revendication 3, dans laquelle ledit polymère muco-adhésif comprend de la carboxyméthylcellulose, de l'acétate de cellulose, de l'éthylcellulose, de l'hydroxyéthylcellulose ou de l'hydroxypropylméthylcellulose. 30
5. Composition pharmaceutique selon l'une quelconque des revendications précédentes, comprenant en outre un antioxydant. 35
6. Composition pharmaceutique selon l'une quelconque des revendications précédentes, comprenant un améliorateur de pénétration. 40
7. Composition pharmaceutique selon la revendication 6, dans laquelle ledit améliorateur de pénétration est un monostéarate de glycérol.
8. Composition pharmaceutique selon l'une quelconque des revendications précédentes, dans laquelle ledit agent de neutralisation du pH est une base organique. 45
9. Composition pharmaceutique selon l'une quelconque des revendications précédentes, dans laquelle ladite première partie comprend des particules d'apomorphine présentant une taille de particule effective d' μm à 10 μm . 50
10. Composition pharmaceutique selon l'une quelconque des revendications précédentes, dans laquelle ledit sel d'addition d'acide d'apomorphine est du

chlorhydrate d'apomorphine.

11. Composition pharmaceutique selon l'une quelconque des revendications 1 à 10, dans laquelle ladite forme de dose unitaire comprend de 2 à 40 mg d'apomorphine ou un sel d'addition d'acide de celle-ci. 5
12. Composition pharmaceutique selon l'une quelconque des revendications 1 à 11, destinée à être utilisée dans le traitement de la maladie de Parkinson, la dyskinésie, l'akinésie d'un mammifère atteint par la maladie de Parkinson.
13. Composition pharmaceutique selon l'une quelconque des revendications 1 à 12, destinée à être utilisée dans le traitement du dysfonctionnement sexuel ou de trouble dépressif. 15

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2006120412 A [0010]
- WO 9826763 A [0011]
- WO 2009052421 A [0054]
- US 20060210610 A [0054]
- US 20010006677 A [0054]
- US 4746508 A [0059]
- US 4540602 A [0065] [0066]
- US 5145684 A [0065] [0066]
- US 5518187 A [0065]
- US 5718388 A [0065]
- US 5862999 A [0065]
- US 5665331 A [0065] [0072]
- US 5662883 A [0065]
- US 5560932 A [0065] [0072]
- US 5543133 A [0065]
- US 5534270 A [0065]
- US 5510118 A [0065] [0073]
- US 5470583 A [0065]
- US 6976647 B [0066]
- EP 0498482 A [0066]

Non-patent literature cited in the description

- VAN LAAR T. *CNS Drugs*, 2003, vol. 17, 475 [0007]
- DEWEY RB JR. *Neurology*, 2004, vol. 62 (4), S3-S7 [0007]
- SWOPE DM. *Neurology*, 2004, vol. 62 (4), S27-S31 [0007]
- DEFFOND et al. *J. Neurol. Neurosurg. Psychiatry*, 1993, vol. 56, 101 [0009]
- KOLLER et al. *Neurology*, 2004, vol. 62, 22 [0009]
- PRIETZ et al. *J. Neurol. Neurosurg. Psychiatry*, 1998, vol. 65, 709 [0009]

SZABADALMI IGÉNYPONTOK

1. Egy szublingvális bevételeire tervezett, egységnyi adagolási formájú gyógyszerészeti készítmény, ahol az említett egységnyi adagolási forma egy film vagy egy csik, aminek az első adagja az apomorfin egy savaddíciós sóját tartalmazó apomorfin szemcsékből áll, a második adagja pedig egy pH-semlegesítő szerből.
2. Az 1. igénypont szerinti gyógyszerészeti készítmény, ahol az említett film forma egy poliszacharidot tartalmaz.
3. Az 1-2. igénypontok bármelyike szerinti gyógyszerészeti készítmény, ahol az említett egységnyi adagolási forma egy mukoadhezív polimert tartalmaz.
4. A 3. igénypont szerinti gyógyszerészeti készítmény, ahol az említett mukoadhezív polimer karboxi-metil-cellulózt, cellulóz-acetátot, etil-cellulózt, hidroxietil-cellulózt vagy hidroxipropilmetil-cellulózt tartalmaz.
5. A korábbi igénypontok bármelyike szerinti gyógyszerészeti készítmény, amely még egy antioxidánst is tartalmaz.
6. A korábbi igénypontok bármelyike szerinti gyógyszerészeti készítmény, amely még egy permeációt javító szert is tartalmaz.
7. A 6. igénypont szerinti gyógyszerészeti készítmény, ahol az említett permeációt javító szer a glicerín-monosztearát.
8. A korábbi igénypontok bármelyike szerinti gyógyszerészeti készítmény, ahol az említett pH-semlegesítő szer egy szerves bázis.
9. A korábbi igénypontok bármelyike szerinti gyógyszerészeti készítmény, ahol az említett első adag 1-10 μm effektív szemcseméretű apomorfin részecskékből áll.
10. A korábbi igénypontok bármelyike szerinti gyógyszerészeti készítmény, ahol az apomorfin említett savaddíciós sója az apomorfin-hidroklorid.
11. Az 1-10. igénypontok bármelyike szerinti gyógyszerészeti készítmény, ahol az említett egységnyi adagolási forma 2-40 mg apomorfinból vagy ennek egy savaddíciós sójából áll.
12. Az 1-11. igénypontok bármelyike szerinti gyógyszerészeti készítmény a Parkinson kór, dyskinesia, akinesia kezelésében való felhasználásra Parkinson-kórban szenvedő emlősen.
13. Az 1-12. igénypontok bármelyike szerinti gyógyszerészeti készítmény szexuális zavar vagy depresszív rendellenesség kezelésében való felhasználásra.