

(19) AUSTRALIAN PATENT OFFICE

(54) Title
N- (2-benzyl) -2-phenylbutanamides as androgen receptor modulators

(51)⁶ International Patent Classification(s)

| | | |
|-----------------------------|--------------------|-------------------|
| C07D 213/61 | 20060101ALI2007022 | |
| (2006.01) | 2BMEP | C07C |
| A61K 31/00 (2006.01) | 233/13 | |
| C07C 233/13 | 20060101ALI2006072 | |
| (2006.01) | 2BMEP | C07C |
| C07C 235/34 | 235/34 | |
| (2006.01) | 20060101ALI2006072 | |
| C07C 271/12 | 271/12 | |
| (2006.01) | 20060101ALI2006072 | |
| C07D 295/205 | 295/205 | |
| (2006.01) | 20060101ALI2006072 | |
| C07D 213/61 | 20060101ALI2006072 | |
| 2BMEP | A61K | 2BMEP |
| 31/00 | | PCT/US2005/019554 |

(21) Application No: 2005251766 (22) Application Date: 2005.06.03

(87) WIPO No: W005/120477

(30) Priority Data

| | | |
|-------------|------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/577,698 | 2004.06.07 | US |
| | | 20090528 |

(43) Publication Date: 2005.12.22

(71) Applicant(s)
Merck & Co., Inc.

(72) Inventor(s)
Hanney, Barbara, Meissner, Robert S., Wang, Jiabing, Krout, Michael R., Perkins, James J., Mitchell, Helen J., Kim, Yuntae, Musselman, Jeffrey

(74) Agent/Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 December 2005 (22.12.2005)

PCT

(10) International Publication Number
WO 2005/120477 A3

(51) International Patent Classification⁷: C07D 213/61, C07C 235/06, 233/13, A61K 31/165, A61P 5/26

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number:
PCT/US2005/019554

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 3 June 2005 (03.06.2005)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CE, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

Published:

— with international search report

(26) Publication Language: English

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(30) Priority Data:

60/577,698 7 June 2004 (07.06.2004) US

(88) Date of publication of the international search report:

2 March 2006

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

For two-letter codes and other abbreviations, refer to the "Guidelines Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors; and

(75) Inventors/Applicants (for US only): HANNEY, Barbara [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). KIM, Yuntae [KR/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). KROUT, Michael, R. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). MEISSNER, Robert, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). MITCHELL, Helen, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). MUSSelman, Jeffrey [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). PERKINS, James, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). WANG, Jiabing [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(54) Title: N- (2-BENZYL) -2-PHENYLBUTANAMIDES AS ANDROGEN RECEPTOR MODULATORS

(57) Abstract: Compounds of structural formula I are modulators of the androgen receptor (AR) in a tissue selective manner. These compounds are useful in the enhancement of weakened muscle tone and the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration, including osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, benign prostatic hyperplasia (BPH), abdominal adiposity, metabolic syndrome, type II diabetes, cancer cachexia, Alzheimer's disease, muscular dystrophies, cognitive decline, sexual dysfunction, sleep apnea, depression, premature ovarian failure, and autoimmune disease, alone or in combination with other active agents.

WO 2005/120477 A3

TITLE OF THE INVENTION

N-(2-BENZYL)-2-PHENYLBUTANAMIDES AS ANDROGEN RECEPTOR MODULATORS

FIELD OF THE INVENTION

5 The present invention relates to *N*-(2-benzyl)-2-phenylbutanamide derivatives, their synthesis, and their use as androgen receptor modulators. More particularly, the compounds of the present invention are tissue-selective androgen receptor modulators (SARMs) and are thereby useful for the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration, such as osteoporosis, periodontal disease, bone fracture, frailty, and sarcopenia.

10 Additionally, the SARMs of the present invention can be used to treat mental disorders associated with low testosterone, such as depression, sexual dysfunction, and cognitive decline. SARMs, being antagonists in specific tissues, are also useful in conditions where elevated androgen tone or activity causes symptoms, such as benign prostate hyperplasia and sleep apnea.

15 BACKGROUND OF THE INVENTION

The androgen receptor (AR) belongs to the superfamily of steroid/thyroid hormone nuclear receptors, whose other members include the estrogen receptor, the progesterone receptor, the glucocorticoid receptor, and the mineralocorticoid receptor. The AR is expressed in numerous tissues of the body and is the receptor through which the physiological as well as the pathophysiological effects of androgens, such as testosterone (T) and dihydrotestosterone (DHT), are mediated. Structurally, the AR is composed of three functional domains: the ligand binding domain (LBD), the DNA-binding domain, and amino-terminal domain. A compound that binds to the AR and mimics the effects of an endogenous AR ligand is referred to as an AR agonist, whereas a compound that inhibits the effects of an endogenous AR ligand is termed an AR antagonist.

25 Androgen ligand binding to the AR induces a ligand/receptor complex, which, after translocation into the nucleus of the cell, binds to regulatory DNA sequences (referred to as androgen response elements) within the promoter or enhancer regions of the target genes present in the nucleus. Other proteins termed cofactors are next recruited, which bind to the receptor leading to gene transcription.

30 Androgen therapy has been to treat a variety of male disorders such as reproductive disorders and primary or secondary male hypogonadism. Moreover, a number of natural or synthetic AR agonists have been investigated for the treatment of musculoskeletal disorders, such as bone disease, hematopoietic disorders, neuromuscular disease, rheumatological disease, wasting disease, and for hormone replacement therapy (HRT), such as female androgen deficiency. In addition, AR antagonists,

such as flutamide and bicalutamide, are used to treat prostate cancer. It would therefore be useful to have available compounds that can activate ("agonize") the function of the AR in a tissue-selective manner that would produce the desired osteo- and myoanabolic effects of androgens without the negative androgenic properties, such as virilization and repression of high density lipoprotein cholesterol (HDL).

5 The beneficial effects of androgens on bone in postmenopausal osteoporosis were documented in recent studies using combined testosterone and estrogen administration [Hofbauer, et al., *Eur. J. Edocrinol.*, 140: 271-286 (1999)]. In a large 2-year, double-blind comparison study, oral conjugated estrogen (CEE) and methyltestosterone combinations were demonstrated to be effective in promoting accrual of bone mass in the spine and hip, while conjugated estrogen therapy alone prevented 10 bone loss [*J. Reprod. Med.*, 44: 1012-1020 (1999)].

15 Additionally, there is evidence that hot flushes decrease in women treated with CEE and methyltestosterone; however, 30% of the treated women suffered from significant increases in acne and facial hair, a complication of all current androgen pharmacotherapies [Watts, et al., *Obstet. Gynecol.*, 85: 529-537 (1995)]. It was also found that the addition of methyltestosterone to CEE decreased HDL levels, as seen in other studies. Thus, the virilizing potential and effects on lipid profile of current 15 androgen therapies provide a rationale for developing tissue-selective androgen receptor agonists.

20 Androgens play an important role in bone metabolism in men [Anderson, et al., "Androgen supplementation in eugonadal men with osteoporosis – effects of six months of treatment on bone mineral density and cardiovascular risk factors," *Bone*, 18: 171-177 (1996)]. Even in eugonadal men with osteoporosis, the therapeutic response to testosterone treatment reveals that androgens exert important osteoanabolic effects. Mean lumbar BMD increased from 0.799 gm/cm² to 0.839 g/cm², in 5 to 6 months in response to 250 mg of testosterone ester administered intramuscularly. SARMs can thus be used to treat osteoporosis in men.

25 Androgen deficiency occurs in men with stage D prostate cancer (metastatic) who undergo androgen deprivation therapy (ADT). Endocrine orchiectomy is achieved by long acting GnRH agonists, while androgen receptor blockade is implemented with AR antagonists. In response to hormonal deprivation, these men suffered from hot flushes, significant bone loss, weakness, and fatigue. In a pilot study of men with stage D prostate cancer, osteopenia (50% vs. 38%) and osteoporosis (38% vs. 25%) were more common in men who had undergone ADT for greater than one year than the patients 30 who did not undergo ADT [Wei, et al., *Urology*, 54: 607-611 (1999)]. Lumbar spine BMD was significantly lower in men who had undergone ADT. Thus tissue selective AR antagonists in the prostate that lack antagonistic action in bone and muscle can be useful agents for the treatment of prostate cancer, either alone or as an adjunct to traditional ADT [See also A. Stoch, et al., *J. Clin. Endocrin. Metab.*, 86: 2787-2791 (2001)].

Tissue-selective AR antagonists can also treat polycystic ovarian syndrome in postmenopausal women. See C.A. Eagleson, et al., "Polycystic ovarian syndrome: evidence that flutamide restores sensitivity of the gonadotropin-releasing hormone pulse generator to inhibition by estradiol and progesterone," *J. Clin. Endocrinol. Metab.*, 85: 4047-4052 (2000).

5 SARMs can also treat certain hematopoietic disorders as androgens stimulate renal hypertrophy and erythropoietin (EPO) production. Prior to the introduction of recombinant human EPO, androgens were employed to treat anemia caused by chronic renal failure. In addition, androgens increase serum EPO levels in anemic patients with non-severe aplastic anemia and myelodysplastic syndromes. Treatment for anemia will require selective action such as can be provided by SARMs.

10 SARMs can also have clinical value as an adjunct to the treatment of obesity. This approach to lowering body fat is supported by published observations that androgen administration reduced subcutaneous and visceral fat in obese patients [J.C. Lovejoy, et al., "Oral anabolic steroid treatment, but not parenteral androgen treatment, decreases abdominal fat in obese, older men," *Int. J. Obesity*, 19: 614-624 (1995)], [J.C. Lovejoy, et al., "Exogenous Androgens Influence Body Composition 15 and Regional Body Fat Distribution in Obese Postmenopausal Women - A Clinical Research Center Study," *J. Clin. Endocrinol. Metab.*, 81: 2198-2203 (1996)]. Therefore, SARMs devoid of unwanted androgenic effects can be beneficial in the treatment of obesity.

20 Androgen receptor agonists can also have therapeutic value against metabolic syndrome (insulin resistance syndrome, syndrome X), particularly in men. Low levels of total and free testosterone and sex hormone-binding globulin (SHBG) in men have been associated with type 2 diabetes, visceral obesity, insulin resistance (hyperinsulinemia, dyslipidemia) and metabolic syndrome. D. Laaksonen, et al., *Diabetes Care*, 27 (5): 1036-1041(2004); see also D. Laaksonen, et al. *Euro. J. Endocrin.*, 149: 601-608 (2003); P. Márín, et al. *Int. J. Obesity*, 16: 991-997 (1992), and P. Márín, et al. *Obesity Res.*, 1(4): 245-251 (1993).

25 Androgen receptor agonists can also have therapeutic value against neurodegenerative diseases such as Alzheimer's disease (AD). The ability of androgens to induce neuroprotection through the androgen receptor was reported by J. Hammond, et al., "Testosterone-mediated neuroprotection through the androgen receptor in human primary neurons," *J. Neurochem.*, 77: 1319-1326 (2001). Gouras et al. reported that testosterone reduces secretion of Alzheimer's β -amyloid peptides and can 30 therefore be used in the treatment of AD [*Proc. Nat. Acad. Sci.*, 97: 1202-1205 (2000)]. A mechanism via inhibition of hyperphosphorylation of proteins implicated in the progression AD has also been described [S. Papasozomenos, "Testosterone prevents the heat shock-induced over activation of glycogen synthase kinase-3 β but not of cyclin-dependent kinase 5 and c-Jun NH₂-terminal kinase and

concomitantly abolishes hyperphosphorylation of τ : Implications for Alzheimer's disease," Proc. Natl. Acad. Sci., 99: 1140-1145 (2002)].


Androgen receptor agonists can also have a beneficial effect on muscle tone and strength. Recent studies have demonstrated that "physiologic androgen replacement in healthy, 5 hypogonadal men is associated with significant gains in fat-free mass, muscle size and maximal voluntary strength," [S. Bhagat, et al., J. Endocrin., 170: 27-38 (2001)].

Androgen receptor modulators can be useful in treating decreased libido in both men and women. Androgen deficiency in men is related to diminished libido. S. Howell et al., Br. J. Cancer, 82: 158-161. Low androgen levels contribute to the decline in sexual interest in many women during their 10 later reproductive years. S. Davis, J. Clin. Endocrinol. Metab., 84: 1886-1891 (1999). In one study, circulating free testosterone was positively correlated with sexual desire. Id. In another study, women with primary or secondary adrenal insufficiency were provided physiological DHEA replacement (50 mg/day). Compared with women taking placebo, DHEA-administered women showed an increase in the frequency of sexual thoughts, interest, and satisfaction. W. Arlt, et al., N Engl. J. Med., 341:1013-1020 15 (1999), see also, K. Miller, J. Clin. Endocrinol. Metab., 86: 2395-2401 (2001).

Additionally, androgen receptor modulators may also be useful in treating cognitive impairment. In a recent study, high-dose oral estrogen either alone or in combination with high-dose oral methyltestosterone was given to postmenopausal women for a four-month period. Cognitive tests were administered before and after the four-month hormone treatment. The investigation found that women 20 receiving a combination of estrogen (1.25 mg) and methyltestosterone (2.50 mg) maintained a steady level of performance on the Building Memory task, but the women receiving estrogen (1.25 mg) alone exhibited decreased performance. A. Wisniewski, Horm. Res., 58:150-155 (2002).

SUMMARY OF THE INVENTION

25 The present invention relates to compounds of structural formula I:



or a pharmaceutically acceptable salt or stereoisomer thereof, their uses, and pharmaceutical compositions.

These compounds are effective as androgen receptor agonists and are particularly effective as SARMs. They are therefore useful for the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration.

The present invention also relates to pharmaceutical compositions comprising the 5 compounds of the present invention and a pharmaceutically acceptable carrier.

In this invention, we have identified compounds that function as SARMs using a series of in vitro cell-assays that profile ligand mediated activation of AR, such as (i) N-C interaction, (ii) transcriptional repression, and (iii) transcriptional activation. SARM compounds in this invention, identified with the methods listed above, exhibit tissue selective AR agonism in vivo, i.e. agonism in 10 bone (stimulation of bone formation in a rodent model of osteoporosis) and antagonism in prostate (minimal effects on prostate growth in castrated rodents and antagonism of prostate growth induced by AR agonists).

The compounds of the present invention identified as SARMs are useful to treat diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration.

15 Such compounds are ideal for the treatment of osteoporosis in women and men as a monotherapy or in combination with inhibitors of bone resorption, such as bisphosphonates, estrogens, SERMs, cathepsin K inhibitors, $\alpha v \beta 3$ integrin receptor antagonists, calcitonin, and proton pump inhibitors. They can also be used with agents that stimulate bone formation, such as parathyroid hormone or analogs thereof. The SARM compounds of the present invention can also be employed for treatment of prostate disease, such 20 as prostate cancer and benign prostatic hyperplasia (BPH). Moreover, compounds of this invention exhibit minimal effects on skin (acne and facial hair growth) and can be useful for treatment of hirsutism. Additionally, compounds of this invention can stimulate muscle growth and can be useful for treatment of sarcopenia and frailty. They can be employed to reduce visceral fat in the treatment of obesity. Moreover, compounds of this invention can exhibit androgen agonism in the central nervous system and 25 can be useful to treat vasomotor symptoms (hot flush) and to increase energy and libido. They can be used in the treatment of Alzheimer's disease.

The compounds of the present invention can also be used in the treatment of prostate cancer, either alone or as an adjunct to GnRH agonist/antagonist therapy, for their ability to restore bone, or as a replacement for antiandrogen therapy because of their ability to antagonize androgen in the 30 prostate, and minimize bone depletion. Further, the compounds of the present invention can be used for their ability to restore bone in the treatment of pancreatic cancer as an adjunct to treatment with antiandrogen, or as monotherapy for their antiandrogenic properties, offering the advantage over traditional antiandrogens of being bone-sparing. Additionally, compounds of this invention can increase the number of blood cells, such as red blood cells and platelets, and can be useful for the treatment of


hematopoietic disorders, such as aplastic anemia. Thus, considering their tissue selective androgen receptor agonism listed above, the compounds of this invention are ideal for hormone replacement therapy in hypogonadic (androgen deficient) men.

This invention is also concerned with safely and specifically treating a male subject with

5 abdominal adiposity, metabolic syndrome (also known as the 'insulin resistance syndrome', and 'Syndrome X'), and type II diabetes.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds that are useful as androgen receptor modulators, in particular, as selective androgen receptor modulators (SARMs). Compounds of the present invention are described by structural formula I:



a pharmaceutically acceptable salt or a stereoisomer thereof, wherein:

X is -CH-, or -N-;

15 n is 0, 1, 2, or 3;

m is 0, 1, or 2;

R¹, R⁴, and R⁵ are each independently chosen from

hydrogen,

halogen,

20 cyano,

perfluoroC₁₋₆alkyl,

perfluoroC₁₋₆alkoxy,

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

25 C₂₋₁₀ alkynyl,

C₁₋₁₀ alkylthio,

aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,

C₃₋₈ heterocyclyl C₂₋₁₀ alkyl,

30 C₃₋₈ heterocycloalkyl C₂₋₁₀ alkyl,

(C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
(aryl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
(C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
5 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonylamino C₀₋₁₀ alkyl,
10 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonylamino C₀₋₁₀ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₁₀ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
15 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
20 aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
25 aryl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
30 aryl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₁₋₁₀ alkoxy (carbonyl)0-1C₀₋₁₀ alkyl,

C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
 hydroxycarbonyl C₁₋₁₀ alkyl,
 hydroxycarbonyl C₂₋₁₀ alkenyl,
 hydroxycarbonyl C₂₋₁₀ alkynyl,
 5 C₁₋₁₀ alkoxy,
 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 aryloxy C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyloxy C₀₋₁₀ alkyl,
 C₃₋₈ heterocycl C₂₋₁₀alkyl oxy C₀₋₁₀ alkyl,
 10 C₃₋₈ heterocycloalkylC₂₋₁₀alkyloxy C₀₋₁₀ alkyl,
 C₁₋₁₀ alkylcarbonyloxy C₀₋₁₀ alkyl,
 (C₀₋₁₀ alkyl)1-2aminosulfonyl C₀₋₁₀ alkyl,
 (aryl C₀₋₁₀ alkyl)1-2aminosulfonyl C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminosulfonyl C₀₋₁₀ alkyl,
 15 C₃₋₈ heterocycl C₀₋₁₀ alkyl aminosulfonyl C₀₋₁₀ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminosulfonyl C₀₋₁₀ alkyl,
 C₀₋₁₀ alkyl sulfonylamino C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl sulfonylamino C₀₋₁₀ alkyl,
 C₃₋₈ heterocycl C₀₋₁₀ alkyl sulfonylamino C₀₋₁₀ alkyl,
 20 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl sulfonylamino C₀₋₁₀ alkyl,
 aryl C₀₋₁₀ alkyl sulfonylamino C₀₋₁₀ alkyl,
 C₁₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
 25 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
 (C₀₋₁₀ alkyl)1-2aminocarbonyloxy,
 (aryl C₀₋₁₀ alkyl)1-2aminocarbonyloxy,
 (C₃₋₈ heterocycl C₀₋₁₀ alkyl)1-2aminocarbonyloxy,
 30 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy,
 (C₃₋₈ cycloalkyl C₀₋₁₀alkyl)1-2aminocarbonyloxy, and


hydroxy C₀₋₁₀alkyl,
 and provided that when X is -N- then R₅ is other than a moiety chosen from (C₀₋₁₀alkyl)1-2amino, C₀₋₁₀ alkyloxy carbonylamino, C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonylamino, aryl C₀₋₁₀ alkyloxy carbonylamino, C₁₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino, C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino, and aryl C₀₋₁₀ alkyloxy(carbonyl)0-1C₀₋₁₀ alkylamino;

5 R² and R³ are each independently chosen from
 hydrogen,
 10 halogen,
 cyano,
 amino,
 hydroxy C₀₋₁₀alkyl,
 perfluoroC₁₋₆alkyl,
 15 perfluoroC₁₋₆alkoxy,
 C₁₋₁₀ alkyl,
 C₂₋₁₀ alkenyl,
 C₂₋₁₀ alkynyl,
 aryl C₀₋₁₀ alkyl,
 20 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl,
 (C₀₋₁₀ alkyl)1-2 amino C₀₋₁₀ alkyl,
 (aryl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
 25 (C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₁₀ alkyl,
 (C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₁₀ alkyl,
 (aryl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₁₀ alkyl,
 30 (C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₁₀ alkyl,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₁₀ alkyl,

(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₁₀ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
5 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₁₀ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₁₀ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₁₀ alkyl,
10 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
15 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
20 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
aryl C₀₋₁₀ alkyloxy carbonylamino C₀₋₁₀ alkyl,
arylcarbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl carbonyloxy C₀₋₁₀ alkyl,
25 C₃₋₈ heterocyclyl carbonyloxy C₀₋₁₀ alkyl,
C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
30 aryl C₀₋₁₀ alkyloxy carbonyloxy C₀₋₁₀ alkyl,
C₁₋₁₀ alkoxy (carbonyl)0-1C₀₋₁₀ alkyl,

C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
aryloxy C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyloxy C₀₋₁₀ alkyl,
5 C₃₋₈ heterocyclyloxy C₀₋₁₀ alkyl,
C₃₋₈ heterocyclylC₀₋₁₀alkyloxy C₀₋₁₀ alkyl,
C₁₋₁₀ alkylcarbonyloxy C₀₋₁₀ alkyl,
C₁₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
C₃₋₈ heterocyclyl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
10 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino; and
aryl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino, and
wherein in R₁, R₂, R₃, R₄, and R₅, said alkyl, alkenyl, alkynyl, aryl, heterocyclyl, heterocycloalkyl, and
15 cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, halogen, CO₂H, cyano, O(C=O)C_{1-C6} alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl, C₀₋₁₀ alkylaminocarbonylamino, C₁₋₁₀ alkyloxycarbonylamino, C₁₋₁₀ alkylcarbonylamino, C₀₋₁₀ alkylaminosulfonylamino, C₁₋₁₀ alkylsulfonylamino, C₁₋₁₀ alkylsulfonyl, C₀₋₁₀ alkylaminosulfonyl, C₀₋₁₀ alkylaminocarbonyl and NH₂.

20 In another embodiment of the invention, compounds are described by structural formula
II:



a pharmaceutically acceptable salt or a stereoisomer thereof, wherein:

25 R⁶ and R⁷ are each independently chosen from
hydrogen,
halogen,
cyano,

amino,
hydroxy C₀₋₉alkyl;
perfluoroC₁₋₆alkyl,
perfluoroC₁₋₆alkoxy,
5 C₁₋₉ alkyl,
C₂₋₉ alkenyl,
C₂₋₉ alkynyl,
aryl C₀₋₉ alkyl,
C₃₋₈ cycloalkyl C₀₋₉ alkyl,
10 C₃₋₈ heterocyclyl C₀₋₉ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₉ alkyl,
(C₀₋₁₀ alkyl)1-2 amino C₀₋₉ alkyl,
(aryl C₀₋₁₀ alkyl)1-2amino C₀₋₉ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₉ alkyl,
15 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2amino C₀₋₉ alkyl,
(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2amino C₀₋₉ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₉ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₉ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₉ alkyl,
20 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₉ alkyl,
(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonyloxy C₀₋₉ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonylaminoC₀₋₉ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₉ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₉ alkyl,
25 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₉ alkyl,
(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)1-2aminocarbonylamino C₀₋₉ alkyl,
(C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₉ alkyl,
(aryl C₀₋₁₀ alkyl)1-2aminocarbonyl C₀₋₉ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₉ alkyl,
30 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₉ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₉ alkyl,

C₀₋₁₀ alkyl carbonylamino C₀₋₉ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₉ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₉ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₉ alkyl,
 5 aryl C₀₋₁₀ alkyl carbonylamino C₀₋₉ alkyl,
 C₀₋₁₀ alkyloxy carbonylamino C₀₋₉ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₉ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₉ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carbonylamino C₀₋₉ alkyl,
 10 aryl C₀₋₁₀ alkyloxy carbonylamino C₀₋₉ alkyl,
 C₀₋₁₀ alkyloxy carboxyloxy C₀₋₉ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy carboxyloxy C₀₋₉ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy carboxyloxy C₀₋₉ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy carboxyloxy C₀₋₉ alkyl,
 15 aryl C₀₋₁₀ alkyloxy carboxyloxy C₀₋₉ alkyl,
 C₁₋₁₀ alkoxy (carbonyl)C₀₋₁C₀₋₉ alkyl,
 C₁₋₁₀ alkyloxy C₀₋₉ alkyl,
 aryloxy C₀₋₉ alkyl,
 C₃₋₈ cycloalkyloxy C₀₋₉ alkyl,
 20 C₃₋₈ heterocyclyoxy C₀₋₉ alkyl,
 C₃₋₈ heterocyclylC₀₋₁₀ alkyloxy C₀₋₉ alkyl, and
 C₁₋₁₀ alkylcarboxyloxy C₀₋₉ alkyl, and
 wherein in R⁶, and R⁷, said alkyl, alkenyl, alkynyl, aryl, heterocyclyl, heterocycloalkyl, and cycloalkyl
 25 are each optionally substituted with one or more groups chosen from hydroxy, C₁₋₆ alkyl, C₁₋₆
 alkoxy, halogen, CO₂H, cyano, O(C=O)C_{1-C6} alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -
 O(0-1)(C₁₋₁₀)perfluoroalkyl, C₀₋₁₀ alkylaminocarbonylamino, C₁₋₁₀ alkyloxycarbonylamino,
 C₁₋₁₀ alkylcarboxylamino, C₀₋₁₀ alkylaminosulfonylamino, C₁₋₁₀ alkylsulfonylamino, C₁₋₁₀
 alkylsulfonyl, C₀₋₁₀ alkylaminosulfonyl, C₀₋₁₀ alkylaminocarbonyl and NH₂; and
 R⁸ is chosen from from hydrogen, hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, halogen, CO₂H, cyano,
 30 O(C=O)C_{1-C6} alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl,
 C₀₋₁₀ alkylaminocarbonylamino, C₁₋₁₀ alkyloxycarbonylamino, C₁₋₁₀ alkylcarboxylamino, C₀₋

10 alkylaminosulfonylamino, C₁-10 alkylsulfonylamino, C₁-10 alkylsulfonyl, C₀-10 alkylaminosulfonyl, C₀-10 alkylaminocarbonyl and NH₂.

Illustrative but nonlimiting examples of compounds of the present invention are the following:

- 5 (S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide;
N-(2-fluoro-5-methylbenzyl)-2-phenylbutanamide;
(S)-N-((2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide;
(S)-N-(5-bromo-2-fluorobenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide;
- 10 10 (S)-N-(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide;
(S)-N-(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide;
N-(5-cyclopropyl-2-fluorobenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-vinylbenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-fluorophenyl)butanamide;
- 15 15 (S)-N-(5-ethyl-2-fluorobenzyl)-2-(4-chlorophenyl)butanamide;
N-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide;
(S)-N-((2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide;
(S)-N-((5-ethyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide;
N-(5-bromo-2-fluorobenzyl)-2-phenylbutanamide;
- 20 20 (S)-N-(5-ethyl-2-fluorobenzyl)-2-(3-chlorophenyl)butanamide;
N-(5-ethyl-2-fluorobenzyl)-2-(3,4-dichlorophenyl)butanamide;
(S)-N-(5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide;
(2R or 2S)-N-[(5-cyclopropyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2R or 2S)-N-[(5-ethyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
- 25 25 (2R or 2S)-N-[(5-methyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2R or 2S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide;
(2R or 2S)-N-(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;
(2R or 2S)-N-(5-cyclopropyl-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;
(2R or 2S)-N-(5-chloro-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
- 30 30 (2R or 2S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide;
(2R or 2S)-N-(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
(2R or 2S)-N-(5-cyclopropyl-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
N-(5-(1,1-difluoroethyl)-2-fluorobenzyl)-2-phenylbutanamide;
(2R or 2S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide;

(2*R* or 2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide;
 (2*R* or 2*S*)-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
 (2*R* or 2*S*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide;
 (2*R* or 2*S*)-*N*-(5-ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
 5 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylpropanamide;
 (2*R* or 2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 10 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-bromobenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-chlorobenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R* or 2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-cyclopropylbenzyl)-2-hydroxy-2-phenylbutanamide;
 (2*R* or 2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylbutanamide;
 15 (2*R*)-3,3,3-trifluoro-*N*-(2,3,5-trifluorobenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R* or 2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;
 (2*R* or 2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;
 20 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-3-cyano-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-4-cyano-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide;
 and pharmaceutically acceptable salts and stereoisomers thereof.


The compounds of the present invention can have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, being included in the present invention.

The term "alkyl" shall mean straight or branched chain alkanes of one to ten total carbon atoms, or any number within this range (i.e., methyl, ethyl, 1-propyl, 2-propyl, n-butyl, s-butyl, t-butyl, etc.). The term "C₀₋₈ alkylaryl" (as in "C₀₋₈ alkylaryl") shall refer to the absence of an alkyl group.

The term "alkenyl" shall mean straight or branched chain alkenes of two to ten total carbon atoms, or any number within this range.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds can be present. Thus, "C₂-C₆ alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. The straight, branched or cyclic portion of the alkynyl group can contain triple bonds and can be substituted if a substituted alkynyl group is indicated.

"Cycloalkyl" as used herein is intended to include non-aromatic cyclic hydrocarbon groups, having the specified number of carbon atoms, which may or may not be bridged or structurally constrained. Examples of such cycloalkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, cycloheptyl, tetrahydro-naphthalene, methylenecyclohexyl, and the like. As used herein, examples of "C₃ - C₁₀ cycloalkyl" can include, but are not limited to:



"Alkoxy" represents either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge. "Alkoxy" therefore encompasses the definitions of alkyl and cycloalkyl above.

"Perfluoroalkyl" represents alkyl chains of up to 10 carbon atoms having exhaustive substitution of their corresponding hydrogens with fluorine atoms.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include, but are not limited to, phenyl, naphthyl, tetrahydro-naphthyl, indanyl, or biphenyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring.

The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms chosen from O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: azabenzimidazole, acridinyl, carbazolyl, cinnolinyl benzimidazolyl, benzofuranyl, benzothiophenyl, benzoxazolyl, benzothiazolyl, benzodihydrofuranyl, 1,3-benzodioxolyl, 2,3-dihydro-1,4-benzodioxinyl,

indolyl, quinolyl, quinoxalinyl, isoquinolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, piridazinyl, tetrahydroquinoliny, thiadiazolyl, oxadiazolyl, triazolyl, imidopyridinyl, tetrazolyl, and indanyl. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heterocaryl. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively.

5 Whenever the term "alkyl" or "aryl" or either of their prefix roots appears in a name of a substituent (e.g., aryl C₀₋₈ alkyl), it shall be interpreted as including those limitations given above for 10 "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C₀₋₈) shall refer independently to the number of carbon atoms in an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro, fluoro, bromo and iodo.

15 The term "heterocycle" or "heterocycl" as used herein is intended to mean a 5- to 14-membered aromatic or nonaromatic ring system containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocycl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrahydro analogs thereof. Further examples of "heterocycl" include, but are not limited to the following: azabenzimidazole, benzimidazolyl, 20 benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indazinyl, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridinyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydropyranyl, tetrazolyl, 25 tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidinyl, aziridinyl, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisoxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, 30 dihydropyrimidinyl, dihydropyrrrolyl, dihydroquinoliny, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidinyl, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof. Attachment of a heterocycl substituent can occur via a carbon atom or via a heteroatom.

The terms "arylalkyl" and "alkylaryl" include an alkyl portion where alkyl is as defined above and include an aryl portion where aryl is as defined above. Examples of arylalkyl include, but are not limited to, benzyl, phenylethyl, phenylpropyl, naphthylmethyl, and naphthylethyl. Examples of alkylaryl include, but are not limited to, toluene, ethylbenzene, propylbenzene, methylpyridine,

5 ethylpyridine, propylpyridine and butylpyridine.

The term "oxy" means an oxygen (O) atom. The term "thio" means a sulfur (S) atom.

The term "oxo" means " $=O$ ". The term "carbonyl" means " $C=O$ ".


The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted

10 compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

When any variable (e.g., R^5 , R^6 , etc.) occurs more than one time in any substituent or in formula I, its definition in each occurrence is independent of its definition at every other occurrence.

15 Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a C_{1-5} alkylcarbonylamino C_{1-6} alkyl substituent is equivalent to



In choosing compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. R^1 , R^2 , R^3 , R^4 , R^5 etc., are to be chosen in conformity with well-known principles of chemical structure connectivity.

25 Lines drawn into the ring systems from substituents indicate that the indicated bond can be attached to any of the substitutable ring atoms. If the ring system is polycyclic, it is intended that the bond be attached to any of the suitable carbon atoms on the proximal ring only.

30 It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups can be on the same carbon or on different carbons, so long as a stable structure results. The phrase "optionally substituted with one or

more substituents" should be taken to be equivalent to the phrase "optionally substituted with at least one substituent" and in such cases one embodiment will have from zero to three substituents.

In one embodiment, X is -CH-. In another embodiment, X is -N-.

In one embodiment of the invention, R¹, R⁴, and R⁵ are each independently chosen from

- 5 hydrogen, halogen, cyano, perfluoroC₁₋₆alkyl, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, aryl C₀₋₁₀ alkyl, C₃₋₈ cycloalkyl C₀₋₁₀ alkyl, C₃₋₈ heterocycl C₂₋₁₀ alkyl, C₃₋₈ heterocycloalkyl C₂₋₁₀ alkyl, C₀₋₁₀ alkylamino C₀₋₁₀ alkyl, C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl, aryl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl, C₃₋₈ heterocycl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl, C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl, C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl, C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
- 10 C₃₋₈ heterocycl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl, C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl, aryl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl, C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl, C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl, C₃₋₈ heterocycl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl, C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl, aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl, C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino, hydroxycarbonyl C₁₋₁₀ alkyl,
- 15 hydroxycarbonyl C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxy, C₁₋₁₀alkyloxy C₀₋₁₀alkyl, aryloxy C₀₋₁₀ alkyl, C₃₋₈ cycloalkyloxy C₀₋₁₀ alkyl, C₃₋₈ heterocycl C₂₋₁₀alkyloxy C₀₋₁₀ alkyl, C₃₋₈ heterocycloalkylC₂₋₁₀alkyloxy C₀₋₁₀ alkyl, C₁₋₁₀ alkylcarboxyloxy C₀₋₁₀ alkyl, C₀₋₁₀ alkyl aminosulfonyl C₀₋₁₀ alkyl, C₀₋₁₀ alkyl sulfonlamino C₀₋₁₀ alkyl, aryl C₀₋₁₀ alkyl sulfonlamino C₀₋₁₀ alkyl, C₁₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino, C₃₋₈ heterocycl C₀₋₁₀ alkylamino, C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino, C₃₋₈ cycloalkyl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino, aryl C₀₋₁₀ alkylloxy(carbonyl)0-1C₀₋₁₀ alkylamino, cyano, and hydroxy C₀₋₁₀alkyl; wherein in R¹, R⁴, and R⁵, said alkyl, alkenyl, alkynyl, aryl, heterocycl, heterocycloalkyl, and cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, halogen, CO₂H,
- 25 cyano, O(C=O)C_{1-C6} alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl, C₀₋₁₀ alkylaminocarbonylamino, C₁₋₁₀ alkylloxycarbonylamino, C₁₋₁₀ alkylcarbonylamino, C₀₋₁₀ alkylaminosulfonyl, C₁₋₁₀ alkylsulfonylamino, C₁₋₁₀ alkylsulfonyl, C₀₋₁₀ alkylaminosulfonyl, C₀₋₁₀ alkylaminocarbonyl and NH₂.

In other embodiment of the invention, R¹, R⁴, and R⁵ are each independently chosen

- 30 from hydrogen, halogen, cyano, perfluoroC₁₋₆alkyl, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, perfluoroC₁₋₆alkoxy, C₂₋₁₀ alkynyl, C₁₋₁₀ alkylthio, (C₁₋₁₀ alkyl)2amino C₀₋₁₀ alkyl, (aryl C₀₋₁₀ alkyl)2amino C₀₋₁₀

alkyl, (C₃-8 cycloalkyl C₀-10 alkyl)2amino C₀-10 alkyl, (C₃-8 heterocyclyl C₀-10 alkyl)2amino C₀-10 alkyl, (C₃-8 heterocycloalkyl C₀-10 alkyl)2amino C₀-10 alkyl, (C₁-10 alkyl)2aminocarbonylamino C₀-10 alkyl, (aryl C₀-10 alkyl)1-2aminocarbonylamino C₀-10 alkyl, C₀-10 alkyl aminocarbonylamino C₀-10 alkyl, C₃-8 cycloalkyl C₀-10 alkyl aminocarbonylamino C₀-10 alkyl, C₃-8 heterocyclyl C₀-10 alkyl 5 aminocarbonylamino C₀-10 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyl aminocarbonylamino C₀-10 alkyl, (C₁-10 alkyl)2aminocarbonyl C₀-10 alkyl, (aryl C₀-10 alkyl)1-2aminocarbonyl C₀-10 alkyl, C₀-10 alkyloxy carbonylamino C₀-10 alkyl, C₃-8 cycloalkyl C₀-10 alkyloxy carbonylamino C₀-10 alkyl, C₃-8 heterocyclyl C₀-10 alkyloxy carbonylamino C₀-10 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyloxy carbonylamino C₀-10 alkyl, aryl C₀-10 alkyloxy carbonylamino C₀-10 alkyl, C₀-10 alkyloxy 10 carbonyloxy C₀-10 alkyl, C₃-8 cycloalkyl C₀-10 alkyloxy carbonyloxy C₀-10 alkyl, C₃-8 heterocyclyl C₀-10 alkyloxy carbonyloxy C₀-10 alkyl hydroxycarbonyl C₂-10 alkynyl, C₃-8 heterocycloalkyl C₀-10 alkyloxy carbonyloxy C₀-10 alkyl, aryl C₀-10 alkyloxy carbonyloxy C₀-10 alkyl, C₁-10 alkoxy (carbonyl)0-1C₀-10 alkyl, (C₁-10 alkyl)2aminosulfonyl C₀-10 alkyl, (aryl C₀-10 alkyl)1-2aminosulfonyl C₀-10 alkyl, C₃-8 cycloalkyl C₀-10 alkyl aminosulfonyl C₀-10 alkyl, C₃-8 heterocyclyl 15 C₀-10 alkyl aminosulfonyl C₀-10 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyl aminosulfonyl C₀-10 alkyl, aryl C₀-10 alkyl aminosulfonyl C₀-10 alkyl, C₃-8 cycloalkyl C₀-10 alkyl sulfonylamino C₀-10 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyl sulfonylamino C₀-10 alkyl, (C₁-10 alkyl)2aminocarbonyloxy, (aryl C₀-10 alkyl)1-2aminocarbonyloxy, (C₃-8 heterocyclyl C₀-10 alkyl)1-2aminocarbonyloxy, (C₃-8 heterocycloalkyl C₀-10 alkyl)1-20 2aminocarbonyloxy, cyano, and (C₃-8 cycloalkyl C₀-10 alkyl)1-2aminocarbonyloxy; wherein in R¹, R⁴, and R⁵, said alkyl, alkenyl, alkynyl, aryl, heterocyclyl, heterocycloalkyl, and cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁-6 alkyl, C₁-6 alkoxy, halogen, CO₂H, cyano, O(C=O)C₁-C₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁-10)perfluoroalkyl, C₀-10 alkylaminocarbonylamino, C₁-10 alkyloxycarbonylamino, C₁-10 25 alkylcarbonylamino, C₀-10 alkylaminosulfonylamino, C₁-10 alkylsulfonylamino, C₁-10 alkylsulfonyl, C₀-10 alkylaminosulfonyl, C₀-10 alkylaminocarbonyl and NH₂.

In one embodiment of the invention, R² and R³ are each independently chosen from hydrogen, halogen, amino, hydroxy C₀-10alkyl, perfluoroC₁-6 alkyl, perfluoroC₁-6 alkoxy, C₁-10 alkyl, C₂-10 alkenyl, aryl C₀-10 alkyl, C₃-8 heterocyclyl C₀-10 alkyl, (C₀-10 alkyl)1-2 amino C₀-10 30 alkyl, (aryl C₀-10 alkyl)1-2amino C₀-10 alkyl, (C₃-8 heterocyclyl C₀-10 alkyl)1-2amino C₀-10 alkyl, (C₀-10 alkyl)1-2aminocarbonyloxy C₀-10 alkyl, (aryl C₀-10 alkyl)1-2arylamino carbonyloxy C₀-10

alkyl, aryl C₀-10 alkyloxy carbonyloxy C₀-10 alkyl, C₁-10 alkoxy (carbonyl)0-1C₀-10 alkyl, C₃-8 cycloalkyloxy, C₃-8 heterocyclyloxy, C₁-10 alkyloxy(carbonyl)0-1C₀-10 alkylamino, C₃-8 heterocycl C₀-10 alkyloxy(carbonyl)0-1C₀-10 alkylamino, C₃-8 heterocycloalkyl C₀-10 alkyloxy(carbonyl)0-1C₀-10 alkylamino, C₃-8 cycloalkyl C₀-10 alkyloxy(carbonyl)0-1C₀-10

5 alkylamino, (C₃-8 heterocycl C₀-10 alkyl)1-2aminocarbonyloxy, (C₃-8 heterocycloalkyl C₀-10 alkyl)1-2aminocarbonyloxy, and (C₃-8 cycloalkyl C₀-10 alkyl)1-2aminocarbonyloxy; arylcarbonyloxy C₀-10 alkyl, C₃-8 heterocycloalkyl carbonyloxy C₀-10 alkyl, C₃-8 cycloalkyl carbonyloxy C₀-10 alkyl, and C₃-8 heterocycl carbonyloxy C₀-10 alkyl, and wherein in R², and R³ said alkyl, alkenyl, alkynyl, aryl, heterocycl, heterocycloalkyl, and

10 cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁-6 alkyl, C₁-6 alkoxy, halogen, CO₂H, cyano, O(C=O)C₁-C₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(O-1)(C₁-10)perfluoroalkyl, C₀-10 alkylaminocarbonylamino, C₁-10 alkyloxy carbonylamino, C₁-10 alkylcarbonylamino, C₀-10 alkylaminosulfonylamino, C₁-10 alkylsulfonylamino, C₁-10 alkylsulfonyl, C₀-10 alkylaminosulfonyl, C₀-10 alkylaminocarbonyl and NH₂.

15 In one embodiment of the compounds of structural formula II, R⁶ and R⁷ are each independently chosen from hydrogen, halogen, cyano, amino, hydroxy C₀-9alkyl, perfluoroC₁-6alkyl, perfluoroC₁-6alkoxy, C₁-9 alkyl, aryl C₀-9 alkyl, C₃-8 cycloalkyl C₀-9 alkyl, C₃-8 heterocycl C₀-9 alkyl, (C₀-10 alkyl)1-2 amino C₀-9 alkyl, (aryl C₀-10 alkyl)1-2amino C₀-9 alkyl, (C₃-8 heterocycl C₀-10 alkyl)1-2amino C₀-9 alkyl, (C₀-10 alkyl)1-2aminocarbonyloxy C₀-9

20 alkyl, (aryl C₀-10 alkyl)1-2aminocarbonyloxy C₀-9 alkyl, (C₀-10 alkyl)1-2aminocarbonylamino C₀-9 alkyl, (aryl C₀-10 alkyl)1-2aminocarbonylamino C₀-9 alkyl, C₀-10 alkyl carbonylamino C₀-9 alkyl, aryl C₀-10 alkyl carbonylamino C₀-9 alkyl, C₀-10 alkyloxy carbonylamino C₀-9 alkyl, C₃-8 heterocycl C₀-10 alkyloxy carbonylamino C₀-9 alkyl, aryl C₀-10 alkyloxy carbonylamino C₀-9 alkyl, C₀-10 alkyloxy carbonyloxy C₀-9 alkyl, C₃-8 heterocycl C₀-10 alkyloxy C₀-9 alkyl, and C₁-10

25 alkylcarbonyloxy C₀-9 alkyl, and wherein in R⁶, and R⁷, said alkyl, alkenyl, alkynyl, aryl, heterocycl, heterocycloalkyl, and cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁-6 alkyl, C₁-6 alkoxy, halogen, CO₂H, cyano, O(C=O)C₁-C₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(O-1)(C₁-10)perfluoroalkyl, C₀-10 alkylaminocarbonylamino, C₁-10 alkyloxy carbonylamino, C₁-10 alkylaminosulfonylamino, C₁-10

30 alkylsulfonylamino, C₁-10 alkylsulfonyl, C₀-10 alkylaminosulfonyl, C₀-10 alkylaminocarbonyl and NH₂.

In another embodiment, R⁶ and R⁷ are each independently chosen from hydrogen, C₂-9 alkenyl, C₂-9 alkynyl, C₃-8 heterocycloalkyl C₀-9 alkyl, (C₃-8 cycloalkyl C₀-10 alkyl)1-2amino C₀-9 alkyl, (C₃-8 heterocycloalkyl C₀-10 alkyl)1-2amino C₀-9 alkyl, (C₃-8 cycloalkyl C₀-10 alkyl)1-2aminocarbonyloxy C₀-9 alkyl, (C₃-8 heterocyclyl C₀-10 alkyl)1-2aminocarbonyloxy C₀-9 alkyl, (C₃-8 heterocycloalkyl C₀-10 alkyl)1-2aminocarbonyloxy C₀-9 alkyl, (C₃-8 cycloalkyl C₀-10 alkyl)1-2aminocarbonylamino C₀-9 alkyl, (C₃-8 heterocyclyl C₀-10 alkyl)1-2aminocarbonylamino C₀-9 alkyl, (C₃-8 heterocycloalkyl C₀-10 alkyl)1-2aminocarbonylamino C₀-9 alkyl, (C₀-10 alkyl)1-2aminocarbonyl C₀-9 alkyl, (aryl C₀-10 alkyl)1-2aminocarbonyl C₀-9 alkyl, C₃-8 cycloalkyl C₀-10 alkyl aminocarbonyl C₀-9 alkyl, C₃-8 heterocyclyl C₀-10 alkyl aminocarbonyl C₀-9 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyl aminocarbonyl C₀-9 alkyl, C₃-8 cycloalkyl C₀-10 alkyl carbonylamino C₀-9 alkyl, C₃-8 heterocyclyl C₀-10 alkyl carbonylamino C₀-9 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyl carbonylamino C₀-9 alkyl, C₃-8 cycloalkyl C₀-10 alkyloxy carbonylamino C₀-9 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyloxy carbonylamino C₀-9 alkyl, C₃-8 cycloalkyl C₀-10 alkyloxy carbonyloxy C₀-9 alkyl, C₃-8 heterocyclyl C₀-10 alkyloxy carbonyloxy C₀-9 alkyl, C₃-8 heterocycloalkyl C₀-10 alkyloxy carbonyloxy C₀-9 alkyl, aryl C₀-10 alkyloxy carbonyloxy C₀-9 alkyl, C₁-10 alkoxy (carbonyl)0-1C₀-9 alkyl, C₁-10alkyloxy C₀-9alkyl, aryloxy C₀-9 alkyl, C₃-8 cycloalkyloxy C₀-9 alkyl, C₃-8 heterocyclyloxy C₀-9 alkyl, C₃-8 heterocyclylC₀-10alkyloxy C₀-9 alkyl, and C₁-10 alkylcarbonyloxy C₀-9 alkyl, and wherein in R⁶ and R⁷, said alkyl, alkenyl, alkynyl, aryl, heterocyclyl, heterocycloalkyl, and cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁-6 alkyl, C₁-6 alkoxy, halogen, CO₂H, cyano, O(C=O)C₁-C₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁-10)perfluoroalkyl, C₀-10 alkylaminocarbonylamino, C₁-10 alkyloxycarbonylamino, C₁-10 alkylcarbonylamino, C₀-10 alkylaminosulfonylamino, C₁-10 alkylsulfonylamino, C₁-10 alkylsulfonyl, C₀-10 alkylaminosulfonyl, C₀-10 alkylaminocarbonyl and NH₂.

In one embodiment, the compounds of the present invention are chosen from (2R)-N-[(5-cyclopropyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide; (2R)-N-[(5-ethyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide; (2R)-N-[(5-methyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide; (2R)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide; (2R)-N-(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide; (2R)-N-(5-(cyclopropyl)-2-fluorobenzyl)-2-(3-bromophenyl)butanamide; (2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide; (2R)-N-(5-chloro-2-fluorobenzyl)-2-(4-

bromophenyl)butanamide; (2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide; (2*R*)-*N*-(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide; (2*R*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(4-bromophenyl)butanamide; (2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide; (2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide; (2*R*)-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*R*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide; (2*R*)-*N*-(5-ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*R*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*R*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide; (2*R*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl dimethylcarbamate; (2*R*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; (2*R*)-3-[(2-fluoro-5-methylpyridin-3-yl)methyl]amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; (2*R*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-1-methyl-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; 3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-2-hydroxy-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; and pharmaceutically acceptable salts and stereoisomers thereof.

In another embodiment the compound of formula I is chosen from (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide, (2*R*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate, and (2*R*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide and pharmaceutically acceptable salts and stereoisomers thereof. In a variant of this embodiment the compound in accordance with this invention is (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide and pharmaceutically acceptable salts and stereoisomers thereof. Another variant is the compound of formula I is (2*R*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide and pharmaceutically acceptable salts and stereoisomers thereof.

In one embodiment, the compounds of the present invention are chosen from (S)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide; (S)-*N*-(2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide; (S)-*N*-(5-bromo-2-fluorobenzyl)-2-phenylbutanamide; (S)-*N*-(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide; (S)-*N*-(5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide; (S)-*N*-(5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-(3,4-dichlorophenyl)butanamide; (2*S*)-*N*-(5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-(3,4-dichlorophenyl)butanamide; (2*S*)-*N*-(5-methyl-2-fluoropyridin-3-yl)methyl)-2-(3,4-

dichlorophenyl)butanamide; (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide; (2*S*)-*N*-(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide; (2*S*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(3-bromophenyl)butanamide; (2*S*)-*N*-(5-chloro-2-fluorobenzyl)-2-(4-bromophenyl)butanamide; (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide; (2*S*)-*N*-(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide; (2*S*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(4-bromophenyl)butanamide; (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide; (2*S*)-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide; (2*S*)-*N*-(5-ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-cyclopropylbenzyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylbutanamide; (2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide; (2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide; (2*S*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl dimethylcarbamate; (2*S*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; (2*S*)-3-[(2-fluoro-5-methylpyridin-3-yl)methyl]amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate; (2*S*)-3-[(2-fluoro-5-methylpyridin-3-yl)methyl]amino]-3-oxo-2-phenylpropyl dimethylcarbamate; and pharmaceutically acceptable salts and stereoisomers thereof.

In another embodiment the compound of formula I is chosen from (2*S*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl dimethylcarbamate, (2*S*)-*N*-(5-methyl-2-fluoropyridin-3-yl)methyl)-2-(3,4-dichlorophenyl)butanamide, and (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide and pharmaceutically acceptable salts and stereoisomers thereof. In a variant of this embodiment the compound in accordance with this invention is (2*S*)-*N*-(5-methyl-2-fluoropyridin-3-yl)methyl)-2-(3,4-dichlorophenyl)butanamide and pharmaceutically acceptable salts and stereoisomers thereof. Another variant is the compound of formula I is (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide and pharmaceutically acceptable salts and stereoisomers thereof.

Compounds of the present invention have been found to be tissue-selective modulators of the androgen receptor (SARMS). In one aspect, compounds of the present invention can be useful to activate the function of the androgen receptor in a mammal, and in particular to activate the function of the androgen receptor in bone and/or muscle tissue and block or inhibit ("antagonize") the function of the androgen receptor in the prostate of a male individual or in the uterus of a female individual.

A further aspect of the present invention is the use of compounds of formula I to attenuate or block the function of the androgen receptor in the prostate of a male individual or in the uterus of a female individual induced by AR agonists, but not in hair-growing skin or vocal cords, and activate the function of the androgen receptor in bone and/or muscle tissue, but not in organs which 5 control blood lipid levels (e.g. liver).

Representative compounds of the present invention typically display submicromolar binding affinity for the androgen receptor. Compounds of this invention are therefore useful in treating mammals suffering from disorders related to androgen receptor function. Therapeutically effective amounts of the compound, including the pharmaceutically acceptable salts thereof, are administered to 10 the mammal, to treat disorders related to androgen receptor function, such as, androgen deficiency, disorders which can be ameliorated by androgen replacement, or which can be improved by androgen replacement, including: enhancement of weakened muscle tone, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture (for example, vertebral and non-vertebral fractures), bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, 15 male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, pancreatic cancer, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, benign prostatic hyperplasia (BPH), cancer cachexia, Alzheimer's disease, muscular dystrophies, cognitive decline, sexual dysfunction, sleep apnea, depression, premature ovarian failure, and autoimmune disease. Treatment is effected by 20 administration of a therapeutically effective amount of a compound of structural formula I to a mammal in need of such treatment. In addition, these compounds are useful as ingredients in pharmaceutical compositions alone or in combination with other active agents.

In one embodiment, the compounds of the present invention can be used to treat conditions in a male individual which are caused by androgen deficiency or which can be ameliorated by 25 androgen replacement, including, but not limited to, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, HIV-wasting, prostate cancer, cancer cachexia, obesity, arthritic conditions, anemias, such as for example, aplastic anemia, muscular dystrophies, and Alzheimer's disease, cognitive decline, sexual dysfunction, sleep apnea, depression, benign prostatic hyperplasia (BPH), abdominal obesity, metabolic syndrome, type II diabetes, and atherosclerosis, alone or in 30 combination with other active agents. Treatment is effected by administration of a therapeutically effective amount of a compound of structural formula I to a male individual in need of such treatment.

"Arthritic condition" or "arthritic conditions" refers to a disease wherein inflammatory lesions are confined to the joints or any inflammatory conditions of the joints, most notably osteoarthritis and rheumatoid arthritis (Academic Press Dictionary of Science Technology; Academic Press; 1st

edition, January 15, 1992). The compounds of Formula I are also useful, alone or in combination, to treat or prevent arthritic conditions, such as Behcet's disease; bursitis and tendinitis; CPPD deposition disease; carpal tunnel syndrome; Ehlers-Danlos syndrome; fibromyalgia; gout; infectious arthritis; inflammatory bowel disease; juvenile arthritis; lupus erythematosus; lyme disease; marfan syndrome; myositis;

5 osteoarthritis; osteogenesis imperfecta; osteonecrosis; polyarteritis; polymyalgia rheumatica; psoriatic arthritis; Raynaud's phenomenon; reflex sympathetic dystrophy syndrome; Reiter's syndrome; rheumatoid arthritis; scleroderma; and Sjogren's syndrome. An embodiment of the invention encompasses the treatment or prevention of an arthritic condition which comprises administering a therapeutically effective amount of a Compound of Formula I. A subembodiment is the treatment or prevention of

10 osteoarthritis, which comprises administering a therapeutically effective amount of a Compound of Formula I. Sec: Cutolo M, Seriolo B, Villaggio B, Pizzorni C, Craviotto C, Sulli A. *Ann. N.Y. Acad. Sci.* 2002 Jun;966:131-42; Cutolo, M. *Rheum Dis Clin North Am* 2000 Nov;26(4):881-95; Bijlsma JW, Van den Brink HR. *Am J Reprod Immunol* 1992 Oct-Dec;28(3-4):231-4; Jansson L, Holmdahl R.; *Arthritis Rheum* 2001 Sep;44(9):2168-75; and Purdie DW. *Br Med Bull* 2000; 56(3):809-23. Also, see

15 Merck Manual, 17th edition, pp. 449-451.

When used in combination to treat arthritic conditions, the compounds of Formula I can be used with any of the drugs disclosed herein as useful for combination therapy, or can be used with drugs known to treat or prevent arthritic conditions, such as corticosteroids, cytotoxic drugs (or other disease modifying or remission inducing drugs), gold treatment, methotrexate, NSAIDs, and COX-2 inhibitors.

20 In another embodiment, the compounds of the present invention can be used to treat conditions in a female individual which are caused by androgen deficiency or which can be ameliorated by androgen replacement, including, but not limited to, osteoporosis, osteopenia, aging skin, glucocorticoid-induced osteoporosis, postmenopausal symptoms, periodontal disease, HIV-wasting,

25 cancer cachexia, obesity, anemias, such as for example, aplastic anemia, muscular dystrophies, Alzheimer's disease, premature ovarian failure, cognitive decline, sexual dysfunction, depression, inflammatory arthritis and joint repair, atherosclerosis, and autoimmune disease, alone or in combination with other active agents. Treatment is effected by administration of a therapeutically effective amount of a compound of structural formula I to a female individual in need of such treatment.

30 The compounds of formula I are also useful in the enhancement of muscle tone in mammals, such as for example, humans. The compounds of structural formula I can also be employed as adjuncts to traditional androgen depletion therapy in the treatment of prostate cancer to restore bone, minimize bone loss, and maintain bone mineral density. In this manner, they can be employed together with traditional androgen deprivation therapy, including GnRH agonists/antagonists, such as those

disclosed in P. Limonta, et al., Exp. Opin. Invest. Drugs, 10: 709-720 (2001); H.J. Stricker, Urology, 58 (Suppl. 2A): 24-27 (2001); R.P. Millar, et al., British Medical Bulletin, 56: 761-772 (2000); and A.V. Schally et al., Advanced Drug Delivery Reviews, 28: 157-169 (1997). The compounds of structural formula I can be used in combination with antiandrogens, such as flutamide, 2-hydroxyflutamide (the active metabolite of flutamide), nilutamide, and bicalutamide (CasodexTM) in the treatment of prostate cancer.

5 Further, the compounds of the present invention can also be employed in the treatment of pancreatic cancer, either for their androgen antagonist properties or as an adjunct to an antiandrogen, such as flutamide, 2-hydroxyflutamide (the active metabolite of flutamide), nilutamide, and bicalutamide (CasodexTM).

10 The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

15 Compounds of structural formula I can minimize the negative effects on lipid metabolism. Therefore, considering their tissue selective androgen agonistic properties, the compounds of this invention exhibit advantages over existing approaches for hormone replacement therapy in hypogonadic (androgen deficient) male individuals.

20 Additionally, compounds of the present invention can increase the number of blood cells, such as red blood cells and platelets, and can be used for treatment of hematopoietic disorders, such as aplastic anemia.

25 In one embodiment of the invention, therapeutically effective amounts of the compound of Formula I, are administered to the mammal, to treat or improve disorders selected from enhancement of weakened muscle tone, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, pancreatic cancer, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, benign prostatic hyperplasia (BPH), cancer cachexia, Alzheimer's disease, muscular dystrophies, cognitive decline, sexual dysfunction, sleep 30 apnea, depression, premature ovarian failure, and autoimmune disease.

30 In another embodiment, therapeutically effective amounts of the compound can be used to treat or improve a disorder selected from weakened muscle tone, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, Alzheimer's disease, and frailty.

In another embodiment, the compound in accordance with the invention can be used to treat or improve a disorder such as male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, pancreatic cancer, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, benign prostatic hyperplasia (BPH), cancer cachexia, muscular dystrophies, cognitive decline, sexual dysfunction, sleep apnea, depression, premature ovarian failure, and autoimmune disease.

5 The compounds of the present invention can be administered in their enantiomerically pure form. Racemic mixtures can be separated into their individual enantiomers by any of a number of conventional methods. These include chiral chromatography, derivatization with a chiral auxiliary
10 followed by separation by chromatography or crystallization, and fractional crystallization of diastereomeric salts.

15 As used herein, a compound of the present invention which functions as an "agonist" of the androgen receptor can bind to the androgen receptor and initiate a physiological or a pharmacological response characteristic of that receptor. The term "tissue-selective androgen receptor modulator" refers to an androgen receptor ligand that mimics the action of a natural ligand in some tissues but not in others. A "partial agonist" is an agonist which is unable to induce maximal activation of the receptor population, regardless of the amount of compound applied. A "full agonist" induces full activation of the androgen receptor population at a given concentration. A compound of the present invention which functions as an "antagonist" of the androgen receptor can bind to the androgen receptor and block or inhibit the
20 androgen-associated responses normally induced by a natural androgen receptor ligand.

25 The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Non-limiting representative salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. In one variant of the invention, the salts are chosen from the ammonium, calcium, lithium, magnesium, potassium, and sodium salts. Non-limiting examples of salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N-dibenzylethylenediamine,
30 diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts can be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Representative acids which can be employed include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, formic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic,

5 mandelic, methanesulfonic, malonic, mucic, nitric, pamoic, pantothenic, phosphoric, propionic, succinic, sulfuric, tartaric, p-toluenesulfonic acid, trifluoroacetic acid, and the like. In one variant, the acids are selected from citric, fumaric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al.*, "Pharmaceutical Salts,"

10 *J. Pharm. Sci.*, 1977:66:1-19.

It would also be noted that the compounds of the present invention are potentially internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary nitrogen atom.

The term "therapeutically effective amount" means the amount the compound of structural formula I that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

15 The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

By "pharmaceutically acceptable" it is meant that the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not be deleterious to the recipient thereof.

20 The terms "administration of a compound" and "administering a compound" should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to the individual in need of treatment.

25 By the term "modulating a function mediated by the androgen receptor in a tissue selective manner" it is meant modulating a function mediated by the androgen receptor selectively (or indiscriminately) in anabolic (bone and/or muscular) tissue (bone and muscular) in the absence of such modulation at androgenic (reproductive) tissue, such as the prostate, testis, seminal vesicles, ovary, uterus, and other sex accessory tissues. In one embodiment, the function of the androgen receptor in anabolic tissue is activated whereas the function of the androgen receptor in androgenic tissue is blocked or suppressed. In another embodiment, the function of the androgen receptor in anabolic tissue is blocked or suppressed whereas the function of the androgen receptor in androgenic tissue is activated.

The administration of a compound of structural formula I in order to practice the present methods of therapy is carried out by administering an effective amount of the compound of structural formula I to the patient in need of such treatment or prophylaxis. The need for a prophylactic administration according to the methods of the present invention is determined via the use of well-known

5 risk factors. The effective amount of an individual compound is determined, in the final analysis, by the physician in charge of the case, but depends on factors such as the exact disease to be treated, the severity of the disease and other diseases or conditions from which the patient suffers, the chosen route of administration, other drugs and treatments which the patient can concomitantly require, and other factors in the physician's judgment.

10 If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent(s) within its approved dosage range. Compounds of the instant invention can alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a combination formulation is inappropriate.

15 Generally, the daily dosage of a compound of structural formula I can be varied over a wide range from about 0.01 to about 1000 mg per adult human per day. For example, dosages range from about 0.1 to about 200 mg/day. For oral administration, the compositions can be provided in the form of tablets containing from about 0.01 to about 1000 mg, such as for example, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 3.0, 5.0, 6.0, 10.0, 15.0, 25.0, 50.0, 75, 100, 125, 150, 175, 180, 200, 225, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the mammal to be treated.

20 The dose can be administered in a single daily dose or the total daily dosage can be administered in divided doses of two, three or four times daily. Furthermore, based on the properties of the individual compound selected for administration, the dose can be administered less frequently, e.g., weekly, twice weekly, monthly, etc. The unit dosage will, of course, be correspondingly larger for the less frequent administration.

25 When administered via intranasal routes, transdermal routes, by rectal or vaginal suppositories, or through an intravenous solution, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

Exemplifying the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Also exemplifying the invention
30 is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Formulations of the tissue-selective androgen receptor modulator employed in the present method for medical use comprise a compound of structural formula I together with an acceptable carrier thereof and optionally other therapeutically active ingredients. The carrier must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the

5 formulation and not being deleterious to the recipient subject of the formulation.

The present invention, therefore, further provides a pharmaceutical formulation comprising a compound of structural formula I together with a pharmaceutically acceptable carrier thereof. The formulations include those suitable for oral, rectal, intravaginal, intranasal, topical and parenteral (including subcutaneous, intramuscular and intravenous administration). In one embodiment, 10 the formulations are those suitable for oral administration.

Suitable topical formulations of a compound of formula I include transdermal devices, aerosols, creams, solutions, ointments, gels, lotions, dusting powders, and the like. The topical pharmaceutical compositions containing the compounds of the present invention ordinarily include about 0.005% to about 5% by weight of the active compound in admixture with a pharmaceutically acceptable 15 vehicle. Transdermal skin patches useful for administering the compounds of the present invention include those well known to those of ordinary skill in that art.

The formulations can be presented in a unit dosage form and can be prepared by any of the methods known in the art of pharmacy. All methods include the step of bringing the active compound in association with a carrier, which constitutes one or more ingredients. In general, the 20 formulations are prepared by uniformly and intimately bringing the active compound in association with a liquid carrier, a waxy solid carrier or a finely divided solid carrier, and then, if needed, shaping the product into the desired dosage form.

Formulations of the present invention suitable for oral administration can be presented as discrete units such as capsules, cachets, tablets or lozenges, each containing a predetermined amount of the active compound; as a powder or granules; or a suspension or solution in an aqueous liquid or non-aqueous liquid, e.g., a syrup, an elixir, or an emulsion.

A tablet can be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine the active compound in a free flowing form, e.g., a powder or granules, optionally mixed with accessory 30 ingredients, e.g., binders, lubricants, inert diluents, disintegrating agents or coloring agents. Molded tablets can be made by molding in a suitable machine a mixture of the active compound, preferably in powdered form, with a suitable carrier. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethyl-cellulose, polyethylene glycol, waxes and the like.

Non-limiting representative lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

5 Oral liquid forms, such as syrups or suspensions in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl cellulose and the like, can be made by adding the active compound to the solution or suspension. Additional dispersing agents which can be employed include glycerin and the like.

Formulations for vaginal or rectal administration can be presented as a suppository with 10 a conventional carrier, i.e., a base that is nontoxic and nonirritating to mucous membranes, compatible with a compound of structural formula I, and is stable in storage and does not bind or interfere with the release of the compound of structural formula I. Suitable bases include: cocoa butter (theobroma oil), polyethylene glycols (such as carbowax and polyglycols), glycol-surfactant combinations, polyoxyl 40 stearate, polyoxyethylene sorbitan fatty acid esters (such as Tween, Myrj, and Arlcel), glycerinated 15 gelatin, and hydrogenated vegetable oils. When glycerinated gelatin suppositories are used, a preservative such as methylparaben or propylparaben can be employed.

Topical preparations containing the active drug component can be admixed with a 20 variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

The compounds of the present invention can also be administered in the form of 25 liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention can also be delivered by the use of monoclonal 30 antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention can also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxy-ethylaspartamidephenol, or polyethylene-oxide polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention can be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polyactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Formulations suitable for parenteral administration include formulations that comprise a sterile aqueous preparation of the active compound which can be isotonic with the blood of the recipient. Such formulations suitably comprise a solution or suspension of a compound that is isotonic with the blood of the recipient subject. Such formulations can contain distilled water, 5% dextrose in distilled

5 water or saline and the active compound. Often it is useful to employ a pharmaceutically and pharmacologically acceptable acid addition salt of the active compound that has appropriate solubility for the solvents employed. Useful formulations also comprise concentrated solutions or solids comprising the active compound which on dilution with an appropriate solvent give a solution suitable for parenteral administration.

10 The pharmaceutical composition and method of the present invention can further comprise other therapeutically active compounds usually applied in the treatment of the above mentioned conditions, including osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, post-menopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, hematopoietic disorders, such as for 15 example, aplastic anemia, pancreatic cancer, Alzheimer's disease, inflammatory arthritis, and joint repair.

For the treatment and prevention of osteoporosis, the compounds of the present invention can be administered in combination with at least one bone-strengthening agent selected from antiresorptive agents, osteoanabolic agents, and other agents beneficial for the skeleton through 20 mechanisms which are not precisely defined, such as calcium supplements, flavonoids, and vitamin D analogs. The conditions of periodontal disease, bone fracture, and bone damage following bone reconstructive surgery can also benefit from these combined treatments. For example, the compounds of the instant invention can be effectively administered in combination with effective amounts of other agents such as estrogens, bisphosphonates, SERMs, cathepsin K inhibitors, $\alpha\beta 3$ integrin receptor 25 antagonists, vacuolar ATPase inhibitors, the polypeptide osteoprotegerin, antagonists of VEGF, thiazolidinediones, calcitonin, protein kinase inhibitors, parathyroid hormone (PTH) and analogs, calcium receptor antagonists, growth hormone secretagogues, growth hormone releasing hormone, insulin-like growth factor, bone morphogenetic protein (BMP), inhibitors of BMP antagonism, prostaglandin derivatives, fibroblast growth factors, vitamin D and derivatives thereof, vitamin K and 30 derivatives thereof, soy isoflavones, calcium salts, and fluoride salts. The conditions of periodontal disease, bone fracture, and bone damage following bone reconstructive surgery can also benefit from these combined treatments.

In one embodiment of the present invention, a compound of the instant invention can be effectively administered in combination with an effective amount of at least one bone-strengthening

agent chosen from estrogen, and estrogen derivatives, alone or in combination with progestin or progestin derivatives; bisphosphonates; antiestrogens or selective estrogen receptor modulators; $\alpha\beta 3$ integrin receptor antagonists; cathepsin K inhibitors; osteoclast vacuolar ATPase inhibitors; calcitonin; and osteoprotegerin.

5 In the treatment of osteoporosis, the activity of the compounds of the present invention are distinct from that of the anti-resorptive agents: estrogens, bisphosphonates, SERMs, calcitonin, cathepsin K inhibitors, vacuolar ATPase inhibitors, agents interfering with the RANK/RANKL/Osteoprotegerin pathway, p38 inhibitors or any other inhibitors of osteoclast generation or osteoclast activation. Rather than inhibiting bone resorption, the compounds of structural formula I
10 aid in the stimulation of bone formation, acting, for example, on cortical bone, which is responsible for a significant part of bone strength. The thickening of cortical bone substantially contributes to a reduction in fracture risk, especially fractures of the hip. The combination of the tissue-SARMs of structural formula I with anti-resorptive agents such as for example estrogen or estrogen derivatives, bisphosphonates, antiestrogens, SERMs, calcitonin, $\alpha\beta 3$ integrin receptor antagonists, HMG-CoA
15 reductase inhibitors, vacuolar ATPase inhibitors, and cathepsin K inhibitors is particularly useful due to the complementary effect of the bone anabolic and antiresorptive actions.

Non-limiting representatives of estrogen and estrogen derivatives include steroidal compounds having estrogenic activity such as, for example, 17 β -estradiol, estrone, conjugated estrogen (PREMARIN \circledR), equine estrogen, 17 β -ethynodiol, and the like. The estrogen or estrogen derivative can be employed alone or in combination with a progestin or progestin derivative. Nonlimiting examples of progestin derivatives are norethindrone and medroxyprogesterone acetate.

Non-limiting examples of bisphosphonate compounds which can also be employed in combination with a compound of the present invention include:

(a) alendronate (also known as alendronic acid, 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid, alendronate sodium, alendronate monosodium trihydrate or 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium trihydrate. Alendronate is described in U.S. Patents 4,922,007, to Kieczykowski et al., issued May 1, 1990; 5,019,651, to Kieczykowski, issued May 28, 1991; 5,510,517, to Dauer et al., issued April 23, 1996; 5,648,491, to Dauer et al., issued July 15, 1997;

25 (b) [(cycloheptylaminomethylene]-bis-phosphonate (incadronate), which is described in U.S. Patent 4,970,335, to Isomura et al., issued November 13, 1990;

30 (c) (dichloromethylene)-bis-phosphonic acid (clodronic acid) and the disodium salt (clodronate), which are described in Belgium Patent 672,205 (1966) and *J. Org. Chem.* 32, 4111 (1967);

- (d) [1-hydroxy-3-(1-pyrrolidinyl)-propylidene]-bis-phosphonate (EB-1053);
- (e) (1-hydroxyethylidene)-bis-phosphonate (etidronate);
- (f) [1-hydroxy-3-(methylpentylamino)propylidene]-bis-phosphonate (ibandronate), which is described in U.S. Patent No. 4,927,814, issued May 22, 1990;
- 5 (g) (6-amino-1-hydroxyhexylidene)-bis-phosphonate (neridronate);
- (h) [3-(dimethylamino)-1-hydroxypropylidene]-bis-phosphonate (olpadronate);
- (i) (3-amino-1-hydroxypropylidene)-bis-phosphonate (pamidronate);
- (j) [2-(2-pyridinyl)ethylidene]-bis-phosphonate (piridronate), which is described in U.S. Patent No. 4,761,406;
- 10 (k) [1-hydroxy-2-(3-pyridinyl)-ethylidene]-bis-phosphonate (risedronate);
- (l) {[[(4-chlorophenyl)thio]methylene]-bis-phosphonate (tiludronate), which is described in U.S. Patent 4,876,248, to Breliere et al., October 24, 1989;
- (m) [1-hydroxy-2-(1H-imidazol-1-yl)ethylidene]-bis-phosphonate (zoledronate); and
- (n) [1-hydroxy-2-imidazopyridin-(1,2-a)-3-ylethylidene]-bis-phosphonate (minodronate).

15 In one embodiment of the methods and compositions of the present invention, the bisphosphonate is chosen from alendronate, clodronate, etidronate, ibandronate, incadronate, minodronate, neridronate, olpadronate, pamidronate, piridronate, risedronate, tiludronate, zoledronate, pharmaceutically acceptable salts of these bisphosphonates, and mixtures thereof. In one variant, the bisphosphonate is selected from alendronate, risedronate, zoledronate, ibandronate, tiludronate, and clodronate. In a subclass of this class, the bisphosphonate is alendronate, pharmaceutically acceptable salts and hydrates thereof, and mixtures thereof. A particular pharmaceutically acceptable salt of alendronate is alendronate monosodium. Pharmaceutically acceptable hydrates of alendronate monosodium include the monohydrate and the trihydrate. A particular pharmaceutically acceptable salt of risedronate is risedronate monosodium. Pharmaceutically acceptable hydrates of risedronate monosodium include the hemi-pentahydrate.

20 Still further, antiestrogenic compounds such as raloxifene (see, e.g., U.S. Patent No. 5,393,763), clomiphene, zucloclomiphene, enclomiphene, nafoxidene, CI-680, CI-628, CN-55,945-27, Mer-25, U-11,555A, U-100A, and salts thereof, and the like (see, e.g., U.S. Patent Nos. 4,729,999 and 4,894,373) can be employed in combination with a compound of structural formula I in the methods and compositions of the present invention. These agents are also known as SERMs, or selective estrogen receptor modulators, agents known in the art to prevent bone loss by inhibiting bone resorption via pathways believed to be similar to those of estrogens.

25 Non-limiting representatives of SERMs include, for example, tamoxifen, raloxifene, lasoxifene, toremifene, azoxifene, EM-800, EM-652, TSE 424, clomiphene, droloxifene, idoxifene,

and levormeloxifene [Goldstein, et al., "A pharmacological review of selective estrogen receptor modulators," *Human Reproduction Update*, 6: 212-224 (2000); Lufkin, et al., *Rheumatic Disease Clinics of North America*, 27: 163-185 (2001), and "Targeting the Estrogen Receptor with SERMs," *Ann. Rep. Med. Chem.* 36: 149-158 (2001)].

5 $\alpha\beta 3$ Integrin receptor antagonists suppress bone resorption and can be employed in combination with the SARMs of structural formula I for the treatment of bone disorders including osteoporosis. Peptidyl as well as peptidomimetic antagonists of the $\alpha\beta 3$ integrin receptor have been described both in the scientific and patent literature. For example, reference is made to W.J. Hoekstra and B.L. Poulter, *Curr. Med. Chem.* 5: 195-204 (1998) and references cited therein; WO 95/32710; WO 10 95/37655; WO 97/01540; WO 97/37655; WO 98/08840; WO 98/18460; WO 98/18461; WO 98/25892; WO 98/31359; WO 98/30542; WO 99/15506; WO 99/15507; WO 00/03973; EP 853084; EP 854140; EP 854145; US Patent Nos. 5,204,350; 5,217,994; 5,639,754; 5,741,796; 5,780,426; 5,929,120; 5,952,341; 6,017,925; and 6,048,861.

15 Other $\alpha\beta 3$ antagonists are described in R.M. Keenan et al., *J. Med. Chem.* 40: 2289-2292 (1997); R.M. Keenan et al., *Bioorg. Med. Chem. Lett.* 8: 3165-3170 (1998); and R.M. Keenan et al., *Bioorg. Med. Chem. Lett.* 8: 3171-3176 (1998).

20 Other non-limiting representative examples of published patent and patent applications that describe various $\alpha\beta 3$ integrin receptor antagonists include: those comprising benzazepine, benzodiazepine and benzocycloheptene—PCT Patent Application Nos. WO 96/00574, WO 96/00730, WO 96/06087, WO 96/26190, WO 97/24119, WO 97/24122, WO 97/24124, WO 98/14192, WO 98/15278, WO 99/05107, WO 99/06049, WO 99/15170, WO 99/15178, WO 97/34865, WO 99/15506, and U.S. Patent No. 6,159,964; those comprising dibenzocycloheptene, and dibenzoxapine—PCT Patent Application Nos. WO 97/01540, WO 98/30542, WO 99/11626, WO 99/15508, and U.S. Patent Nos. 6,008,213 and 6,069,158; those having a phenol constraint—PCT Patent Application Nos. WO 25 98/00395, WO 99/32457, WO 99/37621, WO 99/44994, WO 99/45927, WO 99/52872, WO 99/52879, WO 99/52896, WO 00/06169, European Patent Nos. EP 0 820,988, EP 0 820,991, and U.S. Patent Nos. 5,741,796, 5,773,644, 5,773,646, 5,843,906, 5,852,210, 5,929,120, 5,952,281, 6,028,223 and 6,040,311; those having a monocyclic ring constraint—PCT Patent Application Nos. WO 99/26945, WO 99/30709, WO 99/30713, WO 99/31099, WO 99/59992, WO 00/00486, WO 00/09503, European Patent Nos. EP 0 30 796,855, EP 0 928,790, EP 0 928,793, and U.S. Patent Nos. 5,710,159, 5,723,480, 5,981,546, 6,017,926, and 6,066,648; and those having a bicyclic ring constraint—PCT Patent Application Nos. WO 98/23608, WO 98/35949, and WO 99/33798, European Patent No. EP 0 853,084, and U.S. Patent Nos. 5,760,028, 5,919,792, and 5,925,655.

Cathepsin K, formerly known as cathepsin O2, is a cysteine protease and is described in PCT International Application Publication No. WO 96/13523; U.S. Patent Nos. 5,501,969 and 5,736,357. Cysteine proteases, specifically cathepsins, are linked to a number of disease conditions, such as tumor metastasis, inflammation, arthritis, and bone remodeling. At acidic pH's, cathepsins can degrade type-I collagen. Cathepsin protease inhibitors can inhibit osteoclastic bone resorption by inhibiting the degradation of collagen fibers and are thus useful in the treatment of bone resorption diseases, such as osteoporosis. Non-limiting examples of cathepsin K inhibitors can be found in PCT International Publications WO 01/49288 and WO 01/77073.

Members of the class of HMG-CoA reductase inhibitors, known as the "statins," have been found to trigger the growth of new bone, replacing bone mass lost as a result of osteoporosis (see The Wall Street Journal, Friday, December 3, 1999, page B1). Therefore, the statins hold promise for the treatment of bone resorption. Examples of HMG-CoA reductase inhibitors include statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin (see US Patent No. 4,342,767); simvastatin (see US Patent No. 4,444,784); dihydroxy open-acid simvastatin, particularly the ammonium or calcium salts thereof; pravastatin, particularly the sodium salt thereof (see US Patent No. 4,346,227); fluvastatin, particularly the sodium salt thereof (see US Patent No. 5,354,772); atorvastatin, particularly the calcium salt thereof (see US Patent No. 5,273,995); cerivastatin, particularly the sodium salt thereof (see US Patent No. 5,177,080), rosuvastatin, also known as ZD-4522 (see US Patent No. 5,260,440) and pitavastatin, also referred to as NK-104, itavastatin, or nisvastatin (see PCT international application publication number WO 97/23200).

Osteoclast vacuolar ATPase inhibitors, also called proton pump inhibitors, can be employed together with the SARMs of structural formula I. The proton ATPase which is found on the apical membrane of the osteoclast has been reported to play a significant role in the bone resorption process. Therefore, this proton pump represents an attractive target for the design of inhibitors of bone resorption which are potentially useful for the treatment and prevention of osteoporosis and related metabolic diseases [see C. Farina et al., DDT, 4: 163-172 (1999)].

The angiogenic factor VEGF has been shown to stimulate the bone-resorbing activity of isolated mature rabbit osteoclasts via binding to its receptors on osteoclasts [see M. Nakagawa et al., FEBS Letters, 473: 161-164 (2000)]. Therefore, the development of antagonists of VEGF binding to osteoclast receptors, such as KDR/Flik-1 and Flt-1, can provide yet a further approach to the treatment or prevention of bone resorption.

Activators of the peroxisome proliferator-activated receptor- γ (PPAR γ), such as the thiazolidinediones (TZD's), inhibit osteoclast-like cell formation and bone resorption *in vitro*. Results

reported by R. Okazaki et al. in Endocrinology, 140: 5060-5065 (1999) point to a local mechanism on bone marrow cells as well as a systemic one on glucose metabolism. Nonlimiting examples of PPAR γ , activators include the glitazones, such as troglitazone, pioglitazone, rosiglitazone, and BRL 49653.

Calcitonin can also be employed together with the SARMs of structural formula I.

5 Calcitonin is preferentially employed as salmon nasal spray (Azra et al., Calcitonin. 1996. In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press; and Silverman, "Calcitonin," Rheumatic Disease Clinics of North America, 27: 187-196, 2001)

Protein kinase inhibitors can also be employed together with the SARMs of structural formula I. Kinase inhibitors include those disclosed in WO 01/17562 and are in one embodiment 10 selected from inhibitors of p38. Non-limiting examples of p38 inhibitors useful in the present invention include SB 203580 [Badger et al., J. Pharmacol. Exp. Ther., 279: 1453-1461 (1996)].

Osteoanabolic agents are those agents that are known to build bone by increasing the production of the bone protein matrix. Such osteoanabolic agents include, for example, parathyroid hormone (PTH) and fragments thereof, such as naturally occurring PTH (1-84), PTH (1-34), analogs 15 thereof, native or with substitutions and particularly parathyroid hormone subcutaneous injection. PTH has been found to increase the activity of osteoblasts, the cells that form bone, thereby promoting the synthesis of new bone (Modern Drug Discovery, Vol. 3, No. 8, 2000). An injectable recombinant form of human PTH, Forteo (teriparatide), has received regulatory approval in the U.S. for the treatment of osteoporosis.

20 Also useful in combination with the SARMs of the present invention are calcium receptor antagonists which induce the secretion of PTH as described by Gowen et al., J. Clin. Invest. 105: 1595-604 (2000).

Additional osteoanabolic agents include growth hormone secretagogues, growth hormone, growth hormone releasing hormone and the like can be employed with the compounds 25 according to structural formula I for the treatment of osteoporosis. Representative growth hormone secretagogues are disclosed in U.S. Patent Nos. 3,239,345, 4,036,979, 4,411,890, 5,206,235, 5,283,241, 5,284,841, 5,310,737, 5,317,017, 5,374,721, 5,430,144, 5,434,261, 5,438,136, 5,494,919, 5,494,920, 5,492,916 and 5,536,716; European Patent Pub. Nos. 0,144,230 and 0,513,974; PCT Patent Pub. Nos. WO 94/07486, WO 94/08583, WO 94/11012; WO 94/13696, WO 94/19367, WO 95/03289, WO 30 95/03290, WO 95/09633, WO 95/11029, WO 95/12598, WO 95/13069, WO 95/14666, WO 95/16675, WO 95/16692, WO 95/17422, WO 95/17423, WO 95/34311, and WO 96/02530; articles, Science, 260: 1640-1643 (June 11, 1993); Ann. Rep. Med. Chem., 28: 177-186 (1993); Bioorg. Med. Chem. Lett., 4: 2709-2714 (1994); and Proc. Natl. Acad. Sci. USA, 92: 7001-7005 (1995).

Insulin-like growth factor (IGF) can also be employed together with the SARMs of structural formula I. Insulin-like growth factors can be selected from Insulin-like Growth Factor I, alone or in combination with IGF binding protein 3 and IGF II [See Johannson and Rosen, "The IGFs as potential therapy for metabolic bone diseases," 1996, In: Bilezikian, et al., Ed., Principles of Bone

5 Biology, San Diego: Academic Press; and Ghiron et al., J. Bone Miner. Res. 10: 1844-1852 (1995)].

Bone morphogenetic protein (BMP) can also be employed together with the SARMs of structural formula I. Bone morphogenetic protein includes BMP 2, 3, 5, 6, 7, as well as related molecules TGF beta and GDF 5 [Rosen et al., "Bone morphogenetic proteins," 1996. In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press; and Wang EA, Trends Biotechnol.

10 11: 379-383 (1993)].

Inhibitors of BMP antagonism can also be employed together with the SARMs of structural formula I. In one embodiment, BMP antagonist inhibitors are chosen from inhibitors of the BMP antagonists SOST, noggin, chordin, gremlin, and dan [see Massague and Chen, "Controlling TGF-beta signaling," Genes Dev., 14: 627-644, 2000; Aspenberg et al., J. Bone Miner. Res. 16: 497-500,

15 2001; and Brunkow et al., Am. J. Hum. Genet. 68: 577-89 (2001)].

The tissue-selective androgen receptor modulators of the present invention can also be combined with the polypeptide osteoprotegerin for the treatment of conditions associated with bone loss, such as osteoporosis. The osteoprotegerin can be selected from mammalian osteoprotegerin and human osteoprotegerin. The polypeptide osteoprotegerin, a member of the tumor necrosis factor receptor super-

20 family, is useful to treat bone diseases characterized by increased bone loss, such as osteoporosis.

Reference is made to U.S. Patent No. 6,288,032.

Prostaglandin derivatives can also be employed together with the SARMs of structural formula I. Non-limiting representatives of prostaglandin derivatives are selected from agonists of prostaglandin receptors EP1, EP2, EP4, FP, IP and derivatives thereof [Pilbeam et al., "Prostaglandins

25 and bone metabolism," 1996. In: Bilezikian, et al. Ed. Principles of Bone Biology, San Diego: Academic Press; Weinreb et al., Bone, 28: 275-281 (2001)].

Fibroblast growth factors can also be employed together with the SARMs of structural formula I. Fibroblast growth factors include aFGF, bFGF and related peptides with FGF activity [Hurley Florkiewicz, "Fibroblast growth factor and vascular endothelial growth factor families," 1996. In: J. P.

30 Bilezikian, et al., Ed. Principles of Bone Biology, San Diego: Academic Press].

In addition to bone resorption inhibitors and osteoanabolic agents, there are also other agents known to be beneficial for the skeleton through mechanisms which are not precisely defined. These agents can also be favorably combined with the SARMs of structural formula I.

Vitamin D, vitamin D derivatives and analogs can also be employed together with the SARMs of structural formula I. Vitamin D and vitamin D derivatives include, for example, D₃ (cholecaciferol), D₂ (ergocalciferol), 25-OH-vitamin D₃, 1 α ,25(OH)₂ vitamin D₃, 1 α -OH-vitamin D₃, 1 α -OH-vitamin D₂, dihydrotachysterol, 26,27-F₆-1 α ,25(OH)₂ vitamin D₃, 19-nor-1 α ,25(OH)₂ vitamin D₃, 22-oxacalcitriol, calcipotriol, 1 α ,25(OH)₂-16-ene-23-yne-vitamin D₃ (Ro 23-7553), EB1089, 20-epi-1 α ,25(OH)₂ vitamin D₃, KH1060, ED71, 1 α ,24(S)-(OH)₂ vitamin D₃, 1 α ,24(R)-(OH)₂ vitamin D₃ [See, Jones G., "Pharmacological mechanisms of therapeutics: vitamin D and analogs," 1996. In: J. P. Bilezikian, et al. Ed. Principles of Bone Biology, San Diego: Academic Press].

Vitamin K and Vitamin K derivatives can also be employed together with the SARMs of structural formula I. Vitamin K and vitamin K derivatives include menatetrenone (vitamin K2) [see Shiraki et al., *J. Bone Miner. Res.*, 15: 515-521 (2000)].

Soy isoflavones, including ipriflavone, can be employed together with the SARMs of structural formula I.

Fluoride salts, including sodium fluoride (NaF) and monosodium fluorophosphate (MFP), can also be employed together with the SARMs of structural formula I. Dietary calcium supplements can also be employed together with the SARMs of structural formula I. Dietary calcium supplements include calcium carbonate, calcium citrate, and natural calcium salts (Heaney. Calcium. 1996. In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press).

Daily dosage ranges for bone resorption inhibitors, osteoanabolic agents and other agents which can be used to benefit the skeleton when used in combination with a compound of structural formula I are those which are known in the art. In such combinations, generally the daily dosage range for the SARMs of structural formula I ranges from about 0.01 to about 1000 mg per adult human per day, such as for example, from about 0.1 to about 200 mg/day. However, adjustments to decrease the dose of each agent can be made due to the increased efficacy of the combined agent.

In particular, when a bisphosphonate is employed, dosages from about 2.5 to about 100 mg/day (measured as the free bisphosphonic acid) are appropriate for treatment, such as for example ranging from 5 to 20 mg/day, or about 10 mg/day. Prophylactically, doses of about 2.5 to about 10 mg/day and especially about 5 mg/day should be employed. For reduction in side-effects, it can be desirable to administer the combination of a compound of structural formula I and the bisphosphonate once a week. For once weekly administration, doses ranging from about 15 mg to about 700 mg per week of bisphosphonate and from about 0.07 to about 7000 mg of a compound of structural formula I can be employed, either separately, or in a combined dosage form. A compound of structural formula I can

be favorably administered in a controlled-release delivery device, particularly for once weekly administration.

For the treatment of atherosclerosis, hypercholesterolemia, and hyperlipidemia, the compounds of structural formula I can be effectively administered in combination with one or more

5 additional active agents. The additional active agent or agents can be chosen from lipid-altering compounds such as HMG-CoA reductase inhibitors, agents having other pharmaceutical activities, and agents that have both lipid-altering effects and other pharmaceutical activities. Non-limiting examples of HMG-CoA reductase inhibitors include statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin (see US
10 Patent No. 4,342,767); simvastatin (see US Patent No. 4,444,784); dihydroxy open-acid simvastatin, particularly the ammonium or calcium salts thereof; pravastatin, particularly the sodium salt thereof (see US Patent No. 4,346,227); fluvastatin, particularly the sodium salt thereof (see US Patent No. 5,354,772); atorvastatin, particularly the calcium salt thereof (see US Patent No. 5,273,995); cerivastatin, particularly the sodium salt thereof (see US Patent No. 5,177,080), and nisvastatin, also referred to as
15 NK-104 (see PCT international application publication number WO 97/23200).

Additional active agents which can be employed in combination with a compound of structural formula I include, but are not limited to, HMG-CoA synthase inhibitors; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors), acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as

20 well as dual inhibitors of ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; cholesterol absorption inhibitors, such as SCH-58235, also known as ezetimibe and 1-(4-fluorophenyl)-3(R)-[3(S)-(4-fluorophenyl)-3-hydroxypropyl]-4(S)-(4-hydroxyphenyl)-2-azctidinone, which is described in U.S. Patent Nos. 5,767,115 and 5,846,966; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, for example glycoprotein IIb/IIIa
25 fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR γ), agonists, including the compounds commonly referred to as glitazones, for example troglitazone, pioglitazone and rosiglitazone and, including those compounds included within the structural class known as thiazolidinediones as well as those PPAR γ , agonists outside the thiazolidinedione structural class; PPAR α agonists, such as clofibrate, fenofibrate including micronized
30 fenofibrate, and gemfibrozil; PPAR dual α/γ agonists; vitamin B₆ (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B₁₂ (also known as cyanocobalamin); folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the methylglucarime salt; anti-oxidant vitamins such as vitamin C and E and beta carotene; beta-blockers; angiotensin II antagonists such as losartan; angiotensin converting enzyme inhibitors, such as

enalapril and captopril; calcium channel blockers, such as nifedipine and diltiazem; endothelin antagonists; agents such as LXR ligands that enhance ABC1 gene expression; bisphosphonate compounds, such as alendronate sodium; and cyclooxygenase-2 inhibitors, such as rofecoxib and celecoxib, as well as other agents known to be useful in the treatment of these conditions.

5 Daily dosage ranges for HMG-CoA reductase inhibitors when used in combination with the compounds of structural formula I correspond to those which are known in the art. Similarly, daily dosage ranges for the HMG-CoA synthase inhibitors; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors), acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as well as dual inhibitors of

10 ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; cholesterol absorption inhibitors including ezetimibe; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, including glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR γ) agonists; PPAR α agonists; PPAR dual α/γ agonists; vitamin B₆; vitamin B₁₂; folic acid; anti-oxidant vitamins; beta-

15 blockers; angiotensin II antagonists; angiotensin converting enzyme inhibitors; calcium channel blockers; endothelin antagonists; agents such as LXR ligands that enhance ABC1 gene expression; bisphosphonate compounds; and cyclooxygenase-2 inhibitors also correspond to those which are known in the art, although due to the combined action with the compounds of structural formula I, the dosage can be somewhat lower when administered in combination.

20 One embodiment of the invention is a method for affecting a bone turnover marker in a mammal comprising administering a therapeutically effective amount of a compound according to formula I. Non-limiting examples of bone turnover markers can be selected from urinary C-telopeptide degradation products of type I collagen (CTX), urinary N-telopeptide cross-links of type I collagen (NTX), osteocalcin (bone G1a protein), dual energy x-ray absorptionmetry (DXA), bone specific alkaline

25 phosphatase (BSAP), quantitative ultrasound (QUS), and deoxypyridinoline (DPD) crosslinks.

In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents useful for treating diseases caused by androgen deficiency or that can be ameliorated by addition of androgen.

Abbreviations Used in the Description of the Preparation of the Compounds of the Present Invention:


| | |
|------------------------------------|---|
| AcOH | Acetic acid |
| Dess-Martin | Dess Martin periodinane |
| DHT | Dihydrotestosterone |
| DIPEA | diisopropylethylamine |
| DMAP | 4-Dimethylaminopyridine |
| DMEM | Dulbecco modified eagle media |
| DMSO | Dimethylsulfoxide |
| DMF | N,N-Dimethylformamide |
| EA | Ethyl acetate |
| EDC | 1-(3-Dimethylaminopropyl)3-ethylcarbodiimide HCl |
| EDTA | Ethylenediaminetetraacetic acid |
| EtOH | Ethanol |
| Et ₂ B | triethylborane |
| Et ₃ N | Triethylamine |
| FCS | Fetal calf serum |
| h | hour |
| HEPES | (2-Hydroxyethyl)-1-piperazineethanesulfonic acid |
| HOAt or HOAT | 1-hydroxy-7-azabenzotriazole |
| HPLC | High-performance liquid chromatography |
| KHMDS | Potassium bistrimethylsilylamide |
| LC/MS | Liquid chromatography/mass spectroscopy |
| LDA | Lithium diisopropylamide |
| LG | Leaving group |
| MeOH | Methanol |
| NBS | N-bromosuccinimide |
| n-Bu ₄ NI | Tetra-n-butylammonium iodide |
| Pd(PPh ₃) ₄ | Tetrakis(triphenyl phosphine) palladium(0) |
| PMBCL | p-Methoxybenzyl chloride |
| p-TosCl | p-Toluenesulfonyl chloride |
| PyBop | benzotriazol-1-yloxytritypyrrolidinophosphonium hexafluorophosphate |
| Rt or rt | Room temperature |
| tBuSONH ₂ | t-butylsulfonamide |

| | |
|-----|---------------------------|
| TFA | Trifluoroacetic acid |
| TLC | Thin-layer chromatography |

The compounds of this invention may be prepared by employing reactions as shown in the following schemes, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. The illustrative schemes below, therefore, are not limited by the compounds listed or by any particular substituents employed for illustrative purposes. Substituent 5 numbering as shown in the schemes does not necessarily correlate to that used in the claims and often, for clarity, a single substituent is shown attached to the compound in place of multiple substituents which are allowed under the definitions of Formula I defined previously.


Schemes A-F provides general guidelines for making compounds of Formula I. Scheme A shows the preparations of the amides starting from the commercially available 2-phenylbutanoic 10 chloride or (2S)-phenylbutanoic acid. Scheme B indicates the synthetic routes to functionalize 3-position and to introduce various 5-alkyl groups to the 2-pyridine moiety starting from the commercially available 5-bromo-2-fluoropyridine (**B-1**). Scheme C is the synthetic routes to construct 2-fluoropyridine portion with various R₄ starting from the commercially available 5-substituted-2-aminopyridines (**C-1**). Scheme D shows the synthetic methodology for introducing the difluoromethylene groups at the 5-position of the 15 benzylamine moiety. Chemical transformations shown in Scheme E highlight the preparation of 2-alkyl-2-hydroxyphenylacetic acids (**E-3**) starting from the commercially available benzoylformic acids (**E-1**). Scheme F shows two different synthetic routes to synthesize the 2-hydroxy-2-perfluoroalkyl-2-arylacetic acid (**F-5**) either from **F-1** or **F-3**. Scheme G, indicates that R₃, which can be chosen from halides or any 20 group that can be introduced by the cross-coupling reactions, can be added to the benzylamine portion (**G-1**). Scheme H highlights the synthetic pathways to introduce the aminoacyloxymethyl group at 2-position of the phenylacetic acid moiety starting from the commercially available tropic acid (**H-1**).

Scheme A




5


Scheme B




Scheme C



Scheme D




Scheme E




5

Scheme F




10

Scheme G



5

Scheme H




EXAMPLESEXAMPLE 1-1

Examples 1-1 and 1-2 were synthesized in accordance with Scheme 1.

5


Scheme 1



R = substituted phenyl or substituted 3-pyridyl
 * = racemic or (S)

(S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide (1-1)

10




A solution of (S)-2-phenylbutanoic acid (1-d, 50 mg, 0.30 mmol, Sigma-Aldrich, Milwaukee, Wisconsin) and diisopropylethylamine (98 uL, 0.60 mmol) in *N,N*-dimethylformamide (1 mL) was treated at room temperature with benzotriazol-1-yloxytritypyrrolidinophosphonium hexafluorophosphate (PyBop, 158 mg, 0.30 mmol). After 15 min, (2-fluoro-5-(trifluoromethyl)phenyl)methanamine (1-g, 60 mg, 0.30 mmol, Synthesis, Inc., Wyndham, New Hampshire) was added. The reaction mixture was stirred for 3 h, partitioned between dichloromethane and 0.5N-NaOH. The aqueous layer was removed and the organic layer was washed with 0.5N-HCl. The

- 50 -

aqueous layer was removed by filtering through a plastic frit. The organic layer was concentrated *in vacuo* to give the desired product (**1-1**); **HRMS** ($M+1$) = 340.12948; **¹H NMR** (500 MHz, $CDCl_3$) δ 7.49 (bs, 1 H), 7.41 (d, 1 H, J = 6.5 Hz), 7.34 – 7.26 (m, 5 H), 7.10 (t, 1 H, J = 9.0 Hz), 5.77 (bs, 1 H), 4.47 (d, 1 H, J = 3.0 Hz), 3.30 (t, 2 H, J = 7.5 Hz), 2.22 (m, 1 H), 1.82 (m, 1 H), 0.89 (t, 3 H, J = 7.5 Hz).

5 **EXAMPLE 1-2**

***N*-(2-fluoro-5-methylbenzyl)-2-phenylbutanamide (1-2)**



0 A solution of (2-fluoro-5-methylphenyl)methanamine (**1-f**, 50 mg, 0.26 mmol, Oakwood Products, Inc., West Columbia, South Carolina) and diisopropylethylamine (88 μ L, 0.52 mmol) in dichloromethane (1 mL) was treated at room temperature with 2-phenylbutanoyl chloride (**1-a**, 47 mg, 0.26 mmol). The reaction mixture was stirred for 30 min, partitioned between dichloromethane and 0.1N-
5 NaOH. The aqueous layer was removed and the organic layer was washed with 0.1N-HCl. The aqueous layer was removed by filtering through a plastic frit. The organic layer was evaporated *in vacuo* to give the desired product (**1-2**); **HRMS** ($M+1$) = 286.1586.

5 Additionally, Examples **1-3** through **1-28** in Table 1 below were prepared by the general protocols described in Scheme 1. Specific details of the synthesis of particular compounds are presented
0 below.


Examples **1-6** through **1-9** were obtained by direct introduction of ethyl, vinyl or cyclopropyl group to the corresponding bromide (**1-4** or **1-15**) by a protocol shown in Scheme 2 (**1-6** and **1-7**) or known synthetic methods disclosed in Weichert, A. et al., *Synlett* 1996, 473, and Krolski, M. E., et al. *J. Org. Chem.* 1988, 53, 1171 (**1-8** and **1-9**).

25 The 2-arylbutanoic acid portion in the compounds of Examples **1-10**, **1-11**, **1-16** and **1-17** was prepared by alkylation (lithium diisopropylamide; ethyl iodide) of the corresponding ethyl arylacetate followed by the hydrolysis (KOH in MeOH).

20 The 2-arylbutanoic acid portion in Example **1-19** and **1-20** through **1-28** was prepared according to the procedure described in Myers, A. G., et al.; *J Am Chem. Soc.*, 1994, 116, 9361.

30 The benzylamine portions of the compounds of Examples **1-13**, **1-14** and **1-18** (**2-e**, **2-g** and **2-k**), respectively, were synthesized as shown below in Scheme 2.

Scheme 2

5 Step A. 5-Ethyl-2-fluoropyridine (2-b)

A mixture of 5-bromo-2-fluoropyridine (2-a, 5.03 g, 28.6 mmol, Lancaster Synthesis, Inc., Wyndham, New Hampshire), triethylborane (1M in tetrahydrofuran, 42.8 mL, 42.8 mmol), K_2CO_3 (15.8 g, 114.2 mmol) and $Pd(PPh_3)_4$ (1.65 g, 1.43 mmol) in *N,N*-dimethylformamide was heated at 85 °C for 4 h. The reaction mixture was diluted with water and extracted with hexanes. The organic layer was then washed with water (x2), separated, dried ($MgSO_4$), and concentrated *in vacuo*. Chromatography (50%, CH_2Cl_2 in hexanes) afforded the desired product (2-b); 1H NMR (500 MHz, $CDCl_3$) δ 8.04 (d, 1 H, J = 1.0 Hz), 7.60 (dt, 1 H, J = 8.0, 2.5 Hz), 6.85 (dd, 1 H, J = 8.3, 2.8 Hz), 2.65 (q, 2 H, J = 7.8 Hz), 1.21 (t, 3 H, J = 7.7 Hz).

15

15 Step B. 5-Ethyl-2-fluoropyridine-3-carbaldehyde (2-f)

A solution of diisopropylamine (1.01 mL, 7.19 mmol) in tetrahydrofuran (20 mL) was treated in ice-bath with *n*-butyllithium (2.5M, 2.9 mL, 7.19 mmol) and stirred for 30 min. The resulting solution was reacted at -78 °C with 5-ethyl-2-fluoropyridine (2-b, 0.75 g, 5.99 mmol), stirred for 4 h, and treated with *N,N*-dimethylformamide (482 mg, 6.59 mmol). The reaction mixture was quenched with acetic acid (1 mL), partitioned between ethyl acetate and water. The organic layer was washed with 0.5N-HCl and then with brine, separated, dried ($MgSO_4$) and concentrated *in vacuo*. Chromatography (30% ethyl acetate in hexanes) afforded the desired compound (2-f); 1H NMR (500 MHz, $CDCl_3$) δ 10.3 (s, 1 H), 8.29 (d, 1 H, J = 2.0 Hz), 8.12 (dd, 1 H, J = 9.0, 2.5 Hz), 2.73 (q, 2 H, J = 7.8 Hz), 1.29 (t, 3 H, J = 7.5 Hz).

Step C. (5-Ethyl-2-fluoropyridin-3-yl)methanamine-hydrogen chloride (2-g)

A solution of 5-ethyl-2-fluoropyridine-3-carbaldehyde (2-f, 152 mg, 0.99 mmol), *t*-butylsulfinamide (144 mg, 1.19 mmol) and titanium tetraethoxide (679 mg, 2.98 mmol) in tetrahydrofuran (5 mL) was heated at reflux for 2 h. The reaction mixture was cooled to 0 °C and sodium borohydride (150 mg, 3.97 mmol) was introduced. The resulting mixture was stirred at room temperature for 30 min and quenched with methanol. The thick suspension was filtered through a pad of celite and washed with ethyl acetate. The filtrate solution was washed with brine, dried (MgSO_4) and concentrated *in vacuo* to give the intermediate sulfinamide (148 mg, 58 %); The intermediate (148 mg, 0.57 mmol) was treated with ethanol saturated with HCl (5 mL). The reaction mixture was allowed at room temperature to stir for 30 min and diluted with ethanol (20 mL). All the volatiles were removed *in vacuo* to give (5-ethyl-2-fluoropyridin-3-yl)methanamine-hydrogen chloride (2-g); ^1H NMR (500 MHz, DMSO- d_6) δ 8.49 (bs, 3 H), 8.10 (bs, 1 H), 8.03 (dd, 1 H, J = 9.5, 2.5 Hz), 4.05 (d, 2 H, J = 5.0 Hz), 2.64 (q, 2 H, J = 7.5 Hz), 1.21 (t, 3 H, J = 7.7 Hz).

(2-Fluoro-5-methylpyridin-3-yl)methanamine (2-e)


(2-fluoro-5-methylpyridin-3-yl)methanamine (2-e) was prepared by the same synthetic route as that of (2-g) but utilizing the commercially available 2-fluoro-5-methylpyridine (2-c) as a starting material; ^1H NMR (500 MHz, DMSO- d_6) δ 8.65 (bs, 3 H), 8.07 (d, 1 H, J = 1.0 Hz), 8.01 (dd, 1 H, J = 15.5, 3.0 Hz), 4.02 (s, 2 H), 2.30 (s, 3 H).


EXAMPLE 1-3

Compound 1-3 was synthesized as shown in Scheme 3 and described below.

25


Scheme 3





$R_1 = H, F, Cl, C_{1-6}$ alkyl, C_{1-6} cycloalkyl
 $R_2, R_3 =$ independently H, OH, C_{1-6} alkyl, C_{1-6} cycloalkyl, C_{1-6} perfluoroalkyl
 $R_4 =$ independently H, C_{1-6} alkyl or form 5,6-membered ring together

Step A (S)-N-((2-amino-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide (3-g)




5 A solution of 5-(trifluoromethyl)pyridin-2-amine (3-a with $R_4 = CF_3$, 1.6 g, 9.87 mmol) (Maybridge Chemical company, Cornwall, England) in *N,N*-dimethylformamide (30 mL) was treated at room temperature with silver sulfate (3.1 g, 9.87 mmol) and iodine (2.5 g, 9.87 mol). The reaction mixture was stirred for 14 h and filtered. The filtrated solution was concentrated *in vacuo*. The residue was chromatographed (SiO_2 , 25% ethyl acetate in hexanes) to give 5-(trifluoromethyl)-3-iodopyridin-2-amine (3-b, with $R_4 = CF_3$). The iodide (3-b, with $R_4 = CF_3$, 1.0 g, 3.47 mmol) and cuprous cyanide (CuCN, 78 g, 8.68 mmol) was dissolved in *N,N*-dimethylformamide (6 mL) and heated under the microwave at 100 C for 30 min, cooled to ambient temperature, and diluted with ethyl acetate. The precipitates were removed by filtration. The filtrated solution was partitioned between ethyl acetate and water. The organic layer was washed with brine, separated, dried ($MgSO_4$) and concentrated in vacuo.

10 15 Trituration of the residue with hexanes afforded the desired product, 2-amino-5-(trifluoromethyl)pyridine-3-carbonitrile (3-c, with $R_4 = CF_3$). 2-amino-5-(trifluoromethyl)pyridine-3-carbonitrile (3-c, with $R_4 = CF_3$, 545 mg, 29.2 mmol) was stirred in methanol (10 mL, saturated with ammonia) under hydrogen atmosphere in the presence of Pd/C (200 mg) overnight. The reaction mixture was filtered and the filtrate solution was concentrated *in vacuo* to afford the crude amine (3-d, with $R_4 =$

20 CF₃). A solution of (S)-2-phenylbutanoic acid (300 mg, 18.3 mmol) and diisopropylethylamine (888 uL, 52.4 mmol) in *N,N*-dimethylformamide (5 mL) was treated at room temperature with benzotriazol-1-yloxytrityrrolidinophosphonium hexafluorophosphate (PyBop, 953 mg, 18.3 mmol). After 15 min, 3-(aminomethyl)-5-(trifluoromethyl)pyridin-2-amine (500 mg, 26.2 mmol) was added. The reaction

mixture was stirred for 2 h, partitioned between ethyl acetate and 0.5N-NaOH. The aqueous layer was removed and the organic layer was washed with brine. The organic layer was separated, dried (MgSO_4) and concentrated *in vacuo* to give the desired product (3-g); HRMS ($M+1$) = 338.1405.

5 **Step B (2S)-N-((2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide (1-3)**




To a solution of 3-g (100 mg, 0.30 mmol) in HF-pyridine (2 mL, in a PEG culture tube) was added while in ice-bath sodium nitrite (61.5 mg, 0.89 mmol) in portions. The reaction mixture was stirred for 30 min, poured in ice, partitioned between ethyl acetate and saturated in aqueous sodium bicarbonate. The organic layer was washed with brine, separated (MgSO_4) and concentrated *in vacuo*. Chromatography (SiO_2 , 15% ethyl acetate in hexanes) afforded (S)-N-((2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide (1-3); HRMS ($M+1$) = 341.1283; $^1\text{H NMR}$ (500 MHz, CDCl_3) 88.37 (s, 1 H), 7.76 (d, 1 H, J = 8.5 Hz), 7.37 – 7.34 (m, 2 H), 7.31 – 7.26 (m, 3 H), 5.84 (bs, 1 H), 4.50 (dd, 1 H, J = 16.0, 6.0 Hz), 4.40 (dd, 1 H, J = 16.0, 6.0 Hz), 3.31 (t, 2 H, J = 7.8 Hz), 5. 2.20 (m, 1 H), 1.82 (m, 1 H), 0.88 (t, 3 H, J = 7.5 Hz).

EXAMPLE 1-29


Compound 1-29 was synthesized as shown in Scheme 4 and described below.

Scheme 4



$R_1 = H, F, Cl, C_{1-6}$ alkyl, C_{1-6} cycloalkyl
 $R_2, R_3 =$ independently H, OH, C_{1-6} alkyl, C_{1-6} cycloalkyl, C_{1-6} perfluoroalkyl
 $R_4 =$ independently H, C_{1-6} alkyl or form 5,6-membered ring together

Step A 5-acetyl-2-fluorobenzonitrile (4-d)



To a stirred solution of 2-fluoro-5-formylbenzonitrile (110 mg, 0.74 mmol) in THF (5 mL) was added at -78 °C a solution of methylmagnesium bromide (3M in THF, 0.27 mL, 0.81 mmol). The reaction mixture was stirred at the same temperature for 6 h, warmed up to the room temperature, partitioned between ethyl acetate and saturated NH_4Cl . The organic layer was washed with brine,

10 separated, dried ($MgSO_4$) and concentrated *in vacuo* to give the crude product which was oxidized to the ketone; A solution of the crude alcohol obtained above in CH_2Cl_2 (10 mL) was reacted at room temperature with Dess-Martin periodinane (493 mg, 1.16 mmol) for 2 h. The reaction mixture was partitioned between CH_2Cl_2 and 0.5N-NaOH. The organic layer was separated, dried ($MgSO_4$) and concentrated *in vacuo* to give the crude product which was triturated with hexanes (4-d, 106 mg); 1H


15 NMR (500 MHz, $CDCl_3$) δ 8.26 - 8.21 (m, 2 H), 7.34 (t, 1 H, $J = 8.5$ Hz), 2.63 (s, 3 H).

2005251766 18 Jan 2007

5

0

5

Step B N-[5-(1,1-difluoroethyl)-2-fluorobenzyl]-2-phenylbutanamide (1-29)

1-29










To a stirred solution of the ketone (4-d, 106 mg, 0.65 mmol) in CH_2Cl_2 (5 mL) was added at 0 C dropwise [bis(2-methoxyethyl)amino]sulfur trifluoride (717 mg, 3.25 mmol). The reaction mixture was stirred at room temperature for 1 h, portioned between ethyl acetate and saturated aqueous NaHCO_3 . The organic layer was washed with brine, separated, dried (MgSO_4) and concentrated *in vacuo*. The crude product was diluted in MeOH (10 mL) and stirred under H_2 atmosphere in the presence of Pd/C (50 mg). After 3 h, the reaction mixture was filtered and the filtrate solution was concentrated *in vacuo*. The crude amine was diluted with CH_2Cl_2 (5 mL) and reacted with 2-phenylbutanoyl chloride (119 mg) in the presence of DIEA (0.45 mL, 2.6 mmol). After 3 h, the reaction mixture was concentrated *in vacuo* and chromatographed (SiO_2 , 20% ethyl acetate in hexanes) to give the desired product (1-29); **HRMS** ($M+1$) = 336.1573; **$^1\text{H NMR}$** (500 MHz, CDCl_3) δ 7.27 (m, 7 H), 7.04 (t, 1 H, J = 8.5 Hz), 5.78 (s, 1 H), 4.47 (d, 1 H, J = 6.0 Hz), 3.28 (t, 1 H, J = 7.5 Hz), 2.21 (m, 1 H), 1.82 (t, 1 H, J = 18.0 Hz), 0.89 (t, 1 H, J = 7.5 Hz)

Table 1

| Ex. | Structure | Nomenclature | (LC/MS) or (HRMS) ($M+1$ or $M+1\text{-H}_2\text{O}$) |
|-----|-----------|--|---|
| 1-1 | | (S)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide | 340.1294 |
| 1-2 | | <i>N</i> -(2-fluoro-5-methylbenzyl)-2-phenylbutanamide | 286.1586 |

| | | | |
|------|--|--|----------|
| 1-3 | | (<i>S</i>)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide | 341.1283 |
| 1-4 | | (<i>S</i>)- <i>N</i> -(5-bromo-2-fluorobenzyl)-2-phenylbutanamide | 350.0478 |
| 1-5 | | <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide | 340.1255 |
| 1-6 | | <i>N</i> -(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide | 300.1770 |
| 1-7 | | (<i>S</i>)- <i>N</i> -(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide | 300.1768 |
| 1-8 | | <i>N</i> -(5-cyclopropyl-2-fluorobenzyl)-2-phenylbutanamide | 312.1739 |
| 1-9 | | <i>N</i> -(2-fluoro-5-vinylbenzyl)-2-phenylbutanamide | 298.1585 |
| 1-10 | | <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-fluorophenyl)butanamide | 358.1235 |

| | | | |
|------|--|--|----------|
| 1-11 | | <i>N</i> -(5-ethyl-2-fluorobenzyl)-2-(4-chlorophenyl)butanamide | 334.1373 |
| 1-12 | | <i>N</i> -((2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide | 287.1540 |
| 1-13 | | (S)- <i>N</i> -((2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide | 287.1539 |
| 1-14 | | (S)- <i>N</i> -((5-ethyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide | 301.1725 |
| 1-15 | | <i>N</i> -(5-bromo-2-fluorobenzyl)-2-phenylbutanamide | 350.0481 |
| 1-16 | | <i>N</i> -(5-ethyl-2-fluorobenzyl)-2-(3-chlorophenyl)butanamide | 334.1374 |
| 1-17 | | <i>N</i> -(5-ethyl-2-fluorobenzyl)-2-(3,4-dichlorophenyl)butanamide | 368.0989 |
| 1-18 | | (S)- <i>N</i> -((5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide | 313.1685 |



| | | | |
|------|---|--|------------------|
| 1-19 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -[(5-cyclopropyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide | 381.0927 |
| 1-20 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -[(5-ethyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide | (LC/MS) 369.1 |
| 1-21 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -[(5-methyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide | (LC/MS) 355.0 |
| 1-22 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide | (LC/MS) 418.0 |
| 1-23 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide | (LC/MS) 428.0 |
| 1-24 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -[(5-cyclopropyl-2-fluorobenzyl)-2-(3-bromophenyl)butanamide | (LC/MS) 390.0 |
| 1-25 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(5-chloro-2-fluorobenzyl)-2-(4-bromophenyl)butanamide | (LC/MS) 384.0 |
| 1-26 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide | (LC/MS) 418.0 |
| 1-27 |  | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide | (LC/MS) 428.0 |

| | | | |
|------|--|---|------------------|
| 1-28 | | (2R or 2S)-N-(5-(cyclopropyl)-2-fluorobenzyl)-2-(4-bromophenyl)butanamide | (LC/MS) 390.0 |
| 1-29 | | N-[5-(1,1-difluoroethyl)-2-fluorobenzyl]-2-phenylbutanamide | 336.1573 |

EXAMPLE 2-1

Compound 2-1 was synthesized in accordance with the procedure outlined in Scheme 5
5 and described below.

Scheme 5

(2R or 2S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide (2-1)

10 A solution of 2-oxo-2-phenylacetic acid (10 g, 66.6 mmol) (Acros Organics B.V.B.A., Belgium) in tetrahydrofuran (THF, 200 mL) was treated at room temperature with ethylmagnesium bromide (2.5 M, 80 mL, 200 mmol). The reaction mixture was stirred for 5 h, partitioned between ethyl acetate and 1N-HCl. The organic layer was separated, dried (MgSO_4) and concentrated *in vacuo* to afford 2-hydroxy-2-phenylbutanoic acid. A solution of 2-hydroxy-2-phenylbutanoic acid (100 mg, 0.55

mmol) and diisopropylethylamine (188 μ L, 1.11 mmol) in *N,N*-dimethylformamide (2 mL) was treated at room temperature with benzotriazol-1-yloxytritypyrrolidinophosphonium hexafluorophosphate (PyBop, 290 mg, 0.55 mmol). After 15 min, (3-fluoro-5-(trifluoromethyl)phenyl)methanamine (107 mg, 0.55 mmol) was added. The reaction mixture was stirred for 3 h, partitioned between dichloromethane and 0.5N-NaOH. The aqueous layer was removed and the organic layer was washed with 0.5N-HCl. The aqueous layer was removed by filtering through a plastic frit. The organic layer was evaporated *in vacuo* to give the racemic mixture of the amide which was resolved using the chiral column (CiralPak AD) to give the desired product (**2-1**); HRMS (M+H-H₂O) = 338.1167; ¹H NMR (500 MHz, CDCl₃) 87.59 (d, 1 H, *J* = 7.5 Hz), 7.51 (m, 1 H), 7.42 (d, 1 H, *J* = 5.0 Hz), 7.37 (t, 1 H, *J* = 7.5 Hz), 7.32 (d, 1 H, *J* = 7.0 Hz), 7.13 (t, 1 H, *J* = 9.0 Hz), 7.01 (bs, 1 H), 4.52 (dq, 2 H, *J* = 15.6, 6.4 Hz), 2.83 (s, 1 H), 2.36 (m, 1 H), 2.12 (m, 1 H), 0.94 (t, 3 H, *J* = 7.2 Hz).

Compounds, **2-1** through **2-5**, exemplified in Table 2 were synthesized as shown in Scheme 5. The acid portion of the compound of Example **2-2** was prepared by a known method (Negishi, E., *et al.*, *Tetrahedron Lett.* 1983, 24, 5181) followed by Grignard reaction (EtMgBr) and the hydrolysis of the ester (KOH in aq. EtOH).

Table 2

| Ex. | Structure | Nomenclature | HRMS (M+1 or M+1-H ₂ O) |
|------------|-----------|--|------------------------------------|
| 2-1 | | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide | 338.1167 |
| 2-2 | | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide | 389.0811 |
| 2-3 | | (2 <i>R</i> or 2 <i>S</i>)- <i>N</i> -((2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide | 303.1518 |
| 2-4 | | (2 <i>R</i> or 2 <i>S</i>)-2-cyclopropyl- <i>N</i> -((2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide | 315.1521 |


| | | | |
|-----|--|--|----------|
| 2-5 | | (2R or 2S)-N-((5-ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide | 317.1680 |
|-----|--|--|----------|

EXAMPLE 3-1


Compound **3-1** was synthesized in accord with the general procedure outlined in Scheme 6 below.

5

Scheme 6



$R_1 = H, F, Cl, C_{1-6}$ alkyl, C_{1-6} cycloalkyl
 $R_2 = CF_3$ or C_2F_5
 $R_3 = \text{substituted phenyl, substituted 3-pyridyl}$

(2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide (3-1)

10

5 A solution of (2*R*)-3,3,3-trifluoro-2-hydroxy-2-phenylpropanoic acid (**6-e** with $R = CF_3$ and $R' = H$, 105 mg, 0.48 mmol), the amine (**2-e**, 67 mg, 0.48 mmol) and diisopropylethylamine (415 μ L, 2.4 mmol) in *N,N*-dimethylformamide (1 mL) was treated at room temperature with benzotriazol-1-
yloxytritypyrrolidinophosphonium hexafluorophosphate (PyBop, 274 mg, 0.53 mmol). After 14 h, the
reaction mixture was partitioned between ethyl acetate and 0.5N-NaOH. The aqueous layer was removed
and the organic layer was washed with 0.5N-HCl. The aqueous layer was removed by filtering through a
plastic frit. The organic layer was evaporated *in vacuo* to give the desired product (**3-1**); **HRMS** ($M+H$)
= 343.1059; **¹H NMR** (500 MHz, $CDCl_3$) δ 7.89 (bs, 1 H), 7.63 (m, 2 H), 7.43 (m, 3 H), 7.39 (d, 1 H, J =
7.8 Hz), 6.67 (br, 1 H), 4.72 (m, 1 H), 4.48 (d, 2 H, J = 6.1 Hz), 2.26 (s, 3 H); compound **3-3** [(2*R*)-3,3,3-
trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide] was prepared by the
same procedure (2-fluoro-5-trifluoromethylbenzylamine was purchased from Alfa Aesar); **HRMS**
($M+H$) = 396.0825; **¹H NMR** (500 MHz, $DMSO-d_6$) δ 88.95 (t, 1 H), 7.95 (s, 1 H), 7.65 (m, 3 H), 7.39 (m,
5 H), 4.40 (dq, 2 H).


5 The carboxylic acid moieties of **3-1** through **3-10**, **3-11** in Table 3 were prepared from **6-e**
as shown in Scheme 6 by a known method (Mosher, H. S., *et al.*, *J. Org. Chem.* 1969, **34**, 2543).
Resolution of the enantiomeric mixture of the ester was carried out with the ChiralPack AD [360 nm,
95% Hexanes (0.1% diethylamine) and 5% MeOH/EtOH (1:1)] instead of the fractional crystallization of
the acid as reported in the literature. The absolute configuration of (2*R*)-3,3,3-trifluoro-2-hydroxy-2-
phenylpropanoic acid (**6-e** with $R_1 = H$ and $R_2 = CF_3$) was determined to be *R* (measured $[\alpha]_D +24.8$, c
0.1, MeOH; literature $[\alpha]_D +29.8$, c 0.8, MeOH, Sharpless, K. B. *et al.*, *Tetrahedron: Asymmetry* 1994,
5, 1473). The absolute configuration of resolved 3,3,4,4,4-pentafluoro-2-hydroxy-2-phenylbutanoic acid
has not been established. Under the same resolution condition, the pentafluoroethyl analog (**6-d** with R_1
= H and $R_2 = C_2F_5$) gave two peaks at 18 and 21 minutes, respectively. The acid (**6-e** with $R_1 = H$ and R_2
= C_2F_5) that afforded the active coupled compound (for example, (2*R*)-**3-2** or (2*S*)-**3-2**, Table 3) in the
25 biochemical assays was that of the second peak ($RT = 21$ min, $[\alpha]_D +4.4$, c 0.1, MeOH).

EXAMPLE 3-12

Compound **3-12** was also synthesized in accord with the general procedure outlined in
Scheme 6.

18 Jan 2007
2005251766

(2R or 2S)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide (3-12)




To a stirred solution of the pyruvate (6-c with R = CF₃, 4 g, 23.5 mmol) in tetrahydrofuran (100 mL) was added at -78 °C a solution of the Grignard reagent (6-b with R' = 4-Chloro-3-fluoro, 1M, 24.7 mL, 24.7 mmol). After 1 h, the dry-ice bath was removed. The reaction mixture was stirred overnight, quenched with 1N-HCl, partitioned between diethylether, washed with brine, dried (MgSO₄) and concentrated *in vacuo*. Chromatography (SiO₂, 20% ethyl acetate in hexanes) afforded 4.5 g of the desired ester which was subsequently resolved [6-d, Chiracel AD, 10 cm, 5%-(EtOH/MeOH, 1:1) / hexanes (with 1% diethylamine)], hydrolyzed (6-e, KOH, aqueous ethanol) and coupled to the amine to give 3-12, HRMS (M+H) = 448.0340; ¹H NMR (500 MHz, CDCl₃) δ 7.57 (m, 1H), 7.50 - 7.44 (m, 3H), 7.40 (d, 1H, J = 8.6 Hz), 7.17 (t, 1H, J = 9.0 Hz), 6.67 (bs, 1H), 4.60 (d, 2H, J = 5.8 Hz), 4.58 (d, 1H, J = 2.0 Hz).


5 **EXAMPLES 3-13, 3-14 and 3-15**

Compounds 3-13, 3-14 and 3-15 in Table 3 were prepared as shown in Scheme 7.

Scheme 7




R₁ = H, F, Cl, C₁₋₆ alkyl, C₁₋₆ cycloalkyl
R₂ = CF₃, C₂F₅
R₃ = Br, CN
R₄ = CF₃, Et


Step A:

To a suspension of 2-fluoro-5-trifluoromethylbenzylamine (0.2g, 1.0mmol) and 1,3-dibromo-5,5-dimethylhydantoin (0.15g, 0.5mmol) in 3mL anhydrous CH_2Cl_2 was added triflic acid (0.4g, 2.9mmol). The reaction mixture was shielded from light and stirred at room temperature for overnight. 10mL water was added. The organic layer was separated. The water layer was basified by K_2CO_3 and concentrated to yield a solid which was extracted by ether. The combined ether layer was concentrated to afford the desired product (**7-e**, LC/MS found: 273.9); **7-f** was prepared with the same protocol;

10 LC/MS: 232.03.

Step B:

15 A protocol described for the preparation of **3-1**, from **7-e** or **7-f**, was used to prepare **3-13** and **7-g**, respectively (MS found: 473.9924 for **3-13** and 434.0 for **7-g**).

Step C:

20 A mixture of **3-13** (0.2g, 0.4mmol), $\text{Zn}(\text{CN})_2$ (0.1g, 0.8mmol), $\text{Zn}(0.003\text{g}, 0.04\text{mmol})$, $\text{Pd}_2(\text{dba})_3$ (0.02g, 0.02mmol) and dppf (0.02g, 0.04mmol) in 3mL DMF was bubbled N_2 for 5mins. The mixture was heated in a microwave reactor at 150°C for 30mins. The mixture was filtered and loaded to

Gilson™ reverse phase chromatography (5-85% Acetonitrile over 20mins) to afford the desired product **3-14** (MS found: 421.0780); **3-15** was prepared by the same method from **7-g** (MS found: 381.1216).

Compounds in Table 3 were prepared as shown in Scheme 6 (3-1 through 3-12) or Scheme 7 (3-13 through 3-15).

Table 3


| Ex. | Structure | Nomenclature | (LC/MS) or (HRMS) (M+1 or M+1-H ₂ O) |
|-----|-----------|---|--|
| 3-1 | | (2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide | 343.1059 |
| 3-2 | | (2R or 2S)-3,3,4,4,4-pentafluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylbutanamide | 393.1035 |
| 3-3 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide | 396.0825 |
| 3-4 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide | 362.0562 |
| 3-5 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-5-bromobenzyl)-2-hydroxy-2-phenylpropanamide | 406.0060 |
| 3-6 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-5-chlorobenzyl)-2-hydroxy-2-phenylpropanamide | 362.0562 |
| 3-7 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide | 418.0 (LC/MS) |

| | | | |
|------|--|--|----------|
| 3-8 | | (2R or 2S)-3,3,4,4,4-pentafluoro-N-(2-fluoro-5-cyclopropylbenzyl)-2-hydroxy-2-phenylbutanamide. | 418.1238 |
| 3-9 | | (2R or 2S)-3,3,4,4,4-pentafluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylbutanamide. | 446.0796 |
| 3-10 | | (2R)-3,3,3-trifluoro-N-(2,3,5-trifluorobenzyl)-2-hydroxy-2-phenylpropanamide | 364.7592 |
| 3-11 | | (2R or 2S)-2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide | 386.1138 |
| 3-12 | | (2R or 2S)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide | 448.0340 |
| 3-13 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide | 473.9924 |
| 3-14 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-3-cyano-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide | 421.0780 |
| 3-15 | | (2R)-3,3,3-trifluoro-N-(2-fluoro-4-cyano-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide | 381.1216 |

EXAMPLE 4-1

Compound 4-1 in Table 4 was prepared in accordance with Scheme 8.

Scheme 8




R₁ = H, F, Cl, C₁₋₆ alkyl, C₁₋₆ cycloalkyl

R₂ = substituted phenyl, substituted 3-pyridyl

R₃, R₄ = independently H, C₁₋₆ alkyl or form 5 or 6-membered ring together

5 (2*R* or 2*S*)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl dimethylcarbamate (4-
11)



10 A solution of tropic acid (8-a with R₁ = H, 1.00 g, 6.02 mmol) and (3-fluoro-5-(trifluoromethyl)phenyl)methanamine (1.28 g, 6.62 mmol) in *N,N*-dimethylformamide (10 mL) was treated at room temperature with 1-hydroxy-7-azabenzotriazole (HOAt, 0.98 g, 7.22 mmol) and 1-(3-Dimethylaminopropyl)3-ethylcarbodiimide HCl (EDC, 1.38 g, 7.22 mmol). After 14 h, the reaction mixture was partitioned between dichloromethane and 0.5N-NaOH. The organic layer was dried with sodium sulfate, filtered and concentrated *in vacuo*. Chromatography (SiO₂, 0-80% ethyl acetate in hexanes) afforded 1.7 g of the desired alcohol which was subsequently resolved [Chiracel AD, 10 cm, 15 15% isopropanol / hexanes (with 1% diethylamine)], to give alcohol 8-e. A solution of alcohol (8-e, 53

mg, 0.146 mmol) and DMAP (2.5 mg, 0.015 mmol) in dichloromethane (1 mL) was treated at room temperature with carbonyldiimidazole (CDI, 33 mg, 0.205 mmol.). After 1 h, dimethylamine (0.44 mL, 0.438 mmol, 1 M in THF) was added. After 14 h, the reaction mixture was partitioned between dichloromethane and 0.5N-NaOH. The organic layer was dried with sodium sulfate, filtered and concentrated *in vacuo*. Chromatography (SiO₂, 0-80% ethyl acetate in hexanes) afforded the desired product 4-1; LRMS (M+H) = 413.1491; ¹H NMR (500 MHz, CDCl₃) δ 7.51 (m, 1H), 7.45 (m, 1H), 7.26 (m, 5H), 7.11 (t, 1H, *J* = 9.0 Hz), 6.01 (br, 1H), 4.61 (m, 1H), 4.52 (m, 2H), 4.42 (dd, 1H, *J* = 11.0, 6.1 Hz), 3.86 (m, 1H), 2.76 (s, 3H), 2.70 (s, 3H), 2.60 (bs, 1H).

Additionally, compounds 4-2 through 4-6 in Table 4 were prepared by simple modification of the protocols described above for synthesizing compound 4-1 and in Scheme 8. Simple modification includes the use of different acids, different benzyl or pyridinyl amines for amide formation and different amines in the carbamate formation. The acid used for 4-5 was prepared from ethyl phenylacetacetate by reduction with sodium borohydride and hydrolysis of the ester and the acid used for 4-6 was prepared by a known method disclosed in Wang, Z.-M., *et al.*, *Synlett*. 1993, 8, 603, followed by the hydrolysis to the acid.

Table 4

| Ex. | Structure | Nomenclature | (LC/MS) or (HRMS) (M+1 or M+1-H ₂ O) |
|-----|-----------|---|---|
| 4-1 | | (2R or 2S)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl dimethylcarbamate | 413.1491 |
| 4-2 | | (2R or 2S)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate | 439.1641 |
| 4-3 | | (2R or 2S)-3-[(2-fluoro-5-methylpyridin-3-yl)methyl]amino]-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate | 386.2 |

| | | | |
|-----|--|---|----------|
| 4-4 | | (2R or 2S)-3-[(2-fluoro-5-methylpyridin-3-yl)methyl]amino)-3-oxo-2-phenylpropyl dimethylcarbamate | 360.1 |
| 4-5 | | 3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-1-methyl-3-oxo-2-phenylpropylpyrrolidine-1-carboxylate | 453.1804 |
| 4-6 | | 3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-2-hydroxy-3-oxo-2-phenylpropylpyrrolidine-1-carboxylate | 455.2 |

EXAMPLE 5

Pharmaceutical Composition

As a specific embodiment of this invention, 100 mg of (2R or 2S)-3-[(2-fluoro-5-(trifluoromethyl)benzyl)amino]-3-oxo-2-phenylpropyl, is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size 0, hard gelatin capsule.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it is understood that the practice of the invention encompasses all of the usual variations, adoptions, or modifications, as being within the scope of the following claims and their equivalents.

ASSAYS

In Vitro and In Vivo Assays for SARM Activity Identification of Compounds

The compounds exemplified in the present application exhibited activity in one or more of the following assays.

Hydroxylapatite-based Radioligand Displacement Assay of Compound Affinity for Endogenously Expressed AR

Materials:

Binding Buffer: TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM beta-mercaptoethanol, 10 mM Sodium Molybdate, pH 7.2)

50% HAP Slurry: Calbiochem Hydroxylapatite, Fast Flow, in 10 mM Tris, pH 8.0 and 1 mM EDTA.

5 *Wash Buffer:* 40 mM Tris, pH 7.5, 100 mM KCl, 1 mM EDTA and 1 mM EGTA.

95% EtOH

Methyltrienolone, [17 α -methyl- 3 H], (R1881*); NEN NET590

Methyltrienolone (R1881), NEN NLP005 (dissolve in 95% EtOH)

Dihydrotestosterone (DHT) [1,2,4,5,6,7- 3 H(N)] NEN NET453

10 Hydroxylapatite Fast Flow; Calbiochem Cat#391947

Molybdate = Molybdic Acid (Sigma, M1651)

MDA-MB-453 cell culture media:

RPMI 1640 (Gibco 11835-055) w/23.8

mM NaHCO₃, 2 mM L-glutamine

15 in 500 mL of complete media Final conc.

10 mL (1M Hepes) 20 mM

5 mL (200 mM L-glu) 4 mM

0.5 mL (10 mg/mL human insulin) 10 μ g/mL

in 0.01 N HCl

20 Calbiochem#407694-S)

50 mL FBS (Sigma F2442) 10%

1 mL (10 mg/mL Gentamicin) 20 μ g /mL

Gibco#15710-072)

Cell Passaging

25 Cells (Hall R. E., et al., *European Journal of Cancer*, 30A: 484-490 (1994)) are rinsed twice in PBS, phenol red-free Trypsin-EDTA is diluted in the same PBS 1:10. The cell layers are rinsed with 1X Trypsin, extra Trypsin is poured out, and the cell layers are incubated at 37°C for ~ 2 min. The flask is tapped and checked for signs of cell detachment. Once the cells begin to slide off the flask, the complete media is added to kill the trypsin. The cells are counted at this point, then diluted to the

30 appropriate concentration and split into flasks or dishes for further culturing (Usually 1:3 to 1:6 dilution).

Preparation of MDA-MB-453 Cell Lysate

When the MDA cells are 70 to 85% confluent, they are detached as described above, and collected by centrifuging at 1000 g for 10 minutes at 4°C. The cell pellet is washed twice with TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM beta-mercaptoethanol, 10 mM Sodium Molybdate,

pH 7.2). After the final wash, the cells are resuspended in TEGM at a concentration of 10^7 cells/mL. The cell suspension is snap frozen in liquid nitrogen or ethanol/dry ice bath and transferred to -80°C freezer on dry ice. Before setting up the binding assay, the frozen samples are left on ice-water to just thaw (~ 1 hr). Then the samples are centrifuged at 12,500 g to 20,000 g for 30 min at 4°C . The

5 supernatant is used to set-up assay right away. If using 50 μL of supernatant, the test compound can be prepared in 50 μL of the TEGM buffer.

Procedure for Multiple Compound Screening

1x TEGM buffer is prepared, and the isotope-containing assay mixture is prepared in the following order: EtOH (2% final concentration in reaction), ^3H -R1881 or ^3H -DHT (0.5 nM final Conc.

10 in reaction) and 1x TEGM. [eg. For 100 samples, 200 μL (100 x 2) of EtOH + 4.25 μL of 1:10 ^3H -R1881 stock + 2300 μL (100 x 23) 1x TEGM]. The compound is serially diluted, e.g., if starting final conc. is 1 μM , and the compound is in 25 μL of solution, for duplicate samples, 75 μL of 4x1 μM solution is made and 3 μL of 100 μM is added to 72 μL of buffer, and 1:5 serial dilution.

25 μL of ^3H -R1881 trace and 25 μL compound solution are first mixed together,

15 followed by addition of 50 μL receptor solution. The reaction is gently mixed, spun briefly at about 200 rpm and incubated at 4°C overnight. 100 μL of 50% HAP slurry is prepared and added to the incubated reaction which is then vortexed and incubated on ice for 5 to 10 minutes. The reaction mixture is vortexed twice more to resuspend HAP while incubating reaction. The samples in 96-well format are then washed in wash buffer using The FilterMate™ Universal Harvester plate washer (Packard). The 20 washing process transfers HAP pellet containing ligand-bound expressed receptor to Unifilter-96 GF/B filter plate (Packard). The HAP pellet on the filter plate is incubated with 50 μL of MICROSCINT (Packard) scintillant for 30 minutes before being counted on the TopCount microscintillation counter (Packard). IC₅₀s are calculated using R1881 as a reference.

25 The compounds, Examples 1-1 through 1-19, and Examples 2-1 through 2-15, found in Tables 1 and 2, were tested in the above assay and found to have an IC₅₀ value of 1 micromolar or less.

Mammalian Two-Hybrid Assay for the Ligand-induced Interaction of N-Terminus and C-Terminus Domains of the Androgen Receptor (Agonist Mode: VIRCON)

30 This assay assesses the ability of AR agonists to induce the interaction between the N-terminal domain (NTD) and C-terminal domain (CTD) of rhAR that reflects the *in vivo* virilizing potential mediated by activated androgen receptors. The interaction of NTD and CTD of rhAR is quantified as ligand induced association between a Gal4DBD-rhARCTD fusion protein and a VP16-rhARNTD fusion protein as a mammalian two-hybrid assay in CV-1 monkey kidney cells.

The day before transfection, CV-1 cells are trypsinized and counted, and then plated at 20,000 cells/well in 96-well plates or larger plates (scaled up accordingly) in DMEM + 10% FCS. The next morning, CV-1 cells are cotransfected with pCBB1 (Gal4DBD-rhARLBD fusion construct expressed under the SV40 early promoter), pCBB2 (VP16 -rhAR NTD fusion construct expressed under the SV40 early promoter) and pFR (Gal4 responsive luciferase reporter, Promega) using LIPOFECTAMINE PLUS reagent (GIBCO-BRL) following the procedure recommended by the vendor. Briefly, DNA admixture of 0.05 µg pCBB1, 0.05 µg pCBB2 and 0.1 µg of pFR is mixed in 3.4 µL OPTI-MEM (GIBCO-BRL) mixed with "PLUS Reagent" (1.6 µL, GIBCO-BRL) and incubated at room temperature (RT) for 15 min to form the pre-complexed DNA.

5 For each well, 0.4 µL LIPOFECTAMINE Reagent (GIBCO-BRL) is diluted into 4.6 µL OPTI-MEM in a second tube and mixed to form the diluted LIPOFECTAMINE Reagent. The pre-complexed DNA (above) and the diluted LIPOFECTAMINE Reagent (above) are combined, mixed and incubated for 15 minutes at room temperature. The medium on the cells is replaced with 40 µL /well OPTI-MEM, and 10 µL DNA-lipid complexes are added to each well. The complexes are mixed into the

10 medium gently and incubated at 37°C at 5% CO₂ for 5 hours. Following incubation, 200 µL /well DMEM and 13% charcoal-stripped FCS are added, followed by incubation at 37°C at 5% CO₂. After 24 hours, the test compounds are added at the desired concentration(s) (1 nM – 10 µM). Forty eight hours later, luciferase activity is measured using LUC-Screen system (TROPIX) following the manufacturer's protocol. The assay is conducted directly in the wells by sequential addition of 50 µL each of assay

15 solution 1 followed by assay solution 2. After incubation for 40 minutes at room temperature, luminescence is directly measured with 2-5 second integration.

20 Activity of test compounds is calculated as the E_{max} relative to the activity obtained with 3 nM R1881. Typical tissue-selective androgen receptor modulators of the present invention display weak or no agonist activity in this assay with less than 50% agonist activity at 10 micromolar.

25 See He B, Kempainen JA, Voegel JJ, Gronemeyer H, Wilson EM, "Activation function in the human androgen receptor ligand binding domain mediates inter-domain communication with the NH(2)-terminal domain," *J. Biol. Chem.* 274: 37219-37225 (1999). Trans-Activation Modulation of Androgen Receptor (TAMAR)

30 This assay assesses the ability of test compounds to control transcription from the MMTV-LUC reporter gene in MDA-MB-453 cells, a human breast cancer cell line that naturally expresses the human AR. The assay measures induction of a modified MMTV LTR/promoter linked to the LUC reporter gene.

20,000 to 30,000 cells/well are plated in a white, clear-bottom 96-well plate in "Exponential Growth Medium" which consists of phenol red-free RPMI 1640 containing 10%FBS, 4mM L-glutamine, 20mM HEPES, 10ug/mL human insulin, and 20ug/mL gentamicin. Incubator conditions are 37°C and 5% CO₂. The transfection is done in batch mode. The cells are trypsinized and counted to

5 the right cell number in the proper amount of fresh media, and then gently mixed with the Fugenc/DNA cocktail mix and plated onto the 96-well plate. All the wells receive 200 Tl of medium + lipid/DNA complex and are then incubated at 37°C overnight. The transfection cocktail consists of serum-free Optimem, Fugene6 reagent and DNA. The manufacturer's (Roche Biochemical) protocol for cocktail setup is followed. The lipid (Tl) to DNA (Tg) ratio is approximately 3:2 and the incubation time is 20

10 minutes at room temperature. Sixteen to 24 hrs after transfection, the cells are treated with test compounds such that the final DMSO (vehicle) concentration is <3%. The cells are exposed to the test compounds for 48 hours. After 48 hours, the cells are lysed by a Promega cell culture lysis buffer for 30-60 minutes and then the luciferase activity in the extracts is assayed in the 96-well format luminometer.

Activity of test compounds is calculated as the E_{max} relative to the activity obtained

15 with 100 nM R1881.

See R.E. Hall, et al., "MDA-MB-453, an androgen-responsive human breast carcinoma cell line with high androgen receptor expression," *Eur. J. Cancer*, 30A: 484-490 (1994) and R.E. Hall, et al., "Regulation of androgen receptor gene expression by steroids and retinoic acid in human breast-cancer cells," *Int. J. Cancer*, 52: 778-784 (1992).

20 Activity of test compounds is calculated as the E_{max} relative to the activity obtained with R1881. The exemplified tissue selective androgen receptor modulators of the present invention display partial agonist activity in this assay of greater than 10%.

In Vivo Prostate Assay

Male Sprague-Dawley rats aged 9-10 weeks, the earliest age of sexual maturity, are used

25 in prevention mode. The goal is to measure the degree to which androgen-like compounds delay the rapid deterioration (~85%) of the ventral prostate gland and seminal vesicles that occurs during a seven day period after removal of the testes (orchiectomy [ORX]).

Rats are orchiectomized (ORX). Each rat is weighed, then anesthetized by isoflurane gas that is maintained to effect. A 1.5 cm anteroposterior incision is made in the scrotum. The right testicle

30 is exteriorized. The spermatic artery and vas deferens are ligated with 4.0 silk 0.5cm proximal to the testicle. The testicle is freed by one cut of a small surgical scissors distal to the ligation site. The tissue stump is returned to the scrotum. The same is repeated for the left testicle. When both stumps are returned to the scrotum, the scrotum and overlying skin are sutured closed with 4.0 silk. For Sham-ORX,

all procedures excepting ligation and scissors cutting are completed. The rats fully recover consciousness and full mobility within 10-15 minutes.

A dose of test compound is administered subcutaneously or orally to the rat immediately after the surgical incision is sutured. Treatment continues for an additional six consecutive days.

5 Necropsy and Endpoints

The rat is first weighed, then anesthetized in a CO₂ chamber until near death.

Approximately 5ml whole blood is obtained by cardiac puncture. The rat is then examined for certain signs of death and completeness of ORX. Next, the ventral portion of the prostate gland is located and blunt dissected free in a highly stylized fashion. The ventral prostate is blotted dry for 3-5 seconds and

10 then weighed (VPW). Finally, the seminal vesicle is located and dissected free. The ventral seminal vesicle is blotted dry for 3-5 seconds and then weighed (SVWT).

Primary data for this assay are the weights of the ventral prostate and seminal vesicle. Secondary data include serum LH (luteinizing hormone) and FSH (follicle stimulating hormone), and possible serum markers of bone formation and virilization. Data are analyzed by ANOVA plus Fisher

15 PLSD post-hoc test to identify intergroup differences. The extent to which test compounds inhibit ORX-induced loss of VPW and SVWT is assessed.

In Vivo Bone Formation Assay:

Female Sprague-Dawley rats aged 7-10 months are used in treatment mode to simulate adult human females. The rats have been ovariectomized (OVX) 75-180 days previously, to cause bone

20 loss and simulate estrogen deficient, osteopenic adult human females. Pre-treatment with a low dose of a powerful anti-resorptive, alendronate (0.0028mpk SC, 2X/wk) is begun on Day 0. On Day 15, treatment with test compound is started. Test compound treatment occurs on Days 15-31 with necropsy on Day 32. The goal is to measure the extent to which androgen-like compounds increase the amount of bone formation, shown by increased fluorochrome labeling, at the periosteal surface.

25 In a typical assay, nine groups of seven rats each are studied.

On Days 19 and 29 (fifth and fifteenth days of treatment), a single subcutaneous injection of calcein (8mg/kg) is given to each rat.

Necropsy and Endpoints

The rat is first weighed, then anesthetized in a CO₂ chamber until near death.

30 Approximately 5mL whole blood is obtained by cardiac puncture. The rat is then examined for certain signs of death and completeness of OVX. First, the uterus is located, blunt dissected free in a highly stylized fashion, blotted dry for 3-5 seconds and then weighed (UW). The uterus is placed in 10% neutral-buffered formalin. Next, the right leg is disarticulated at the hip. The femur and tibia are separated at the knee, substantially defleshed, and then placed in 70% ethanol.

A 1-cm segment of the central right femur, with the femoral proximal-distal midpoint at center, is placed in a scintillation vial and dehydrated and defatted in graded alcohols and acetone, then introduced to solutions with increasing concentrations of methyl methacrylate. It is embedded in a mixture of 90% methyl methacrylate: 10% dibutyl phthalate that is allowed to polymerize over a 48-72

5 hours period. The bottle is cracked and the plastic block is trimmed into a shape that conveniently fits the vice-like specimen holder of a Leica 1600 Saw Microtome, with the long axis of the bone prepared for cross-sectioning. Three cross-sections of 85 μ m thickness are prepared and mounted on glass slides. One section from each rat that approximates the midpoint of the bone is selected and blind-coded. The periosteal surface of each section is assessed for total periosteal surface, single fluorochrome label,


10 double fluorochrome label, and interlabel distance.

Primary data for this assay are the percentage of periosteal surface bearing double label and the mineral apposition rate (interlabel distance(μ m)/10d), semi-independent markers of bone formation. Secondary data include uterus weight and histologic features. Tertiary endpoints can include serum markers of bone formation and virilization. Data are analyzed by ANOVA plus Fisher PLSD post-

15 hoc test to identify intergroup differences. The extent to which test compounds increase bone formation endpoint are assessed.

The claims defining the invention are as follows:

1. A compound of structural formula I:



(I)

a pharmaceutically acceptable salt or a stereoisomer thereof, wherein:

5 X is -CH-, or -N-;

n is 0, 1, 2, or 3;

m is 0, 1, or 2;

R¹ and R⁴, are each independently chosen from

hydrogen,

halogen,

cyano,

perfluoroC₁₋₆alkyl,

C₁₋₁₀alkyl,

C₂₋₁₀alkenyl,

15 C₃₋₈cycloalkyl C₀₋₁₀alkyl,

C₃₋₈heterocycloalkyl C₂₋₁₀alkyl,

hydroxy C₀₋₁₀alkyl,

R⁵ is hydrogen or CN;

20 R² and R³ are each independently chosen from

hydrogen,

halogen,

hydroxy C₀₋₁₀alkyl,

perfluoroC₁₋₆alkyl,

perfluoroC₁₋₆alkoxy,

25 C₁₋₁₀alkyl,

C₂₋₁₀alkenyl,

C₃₋₈cycloalkyl C₀₋₁₀alkyl,

(C₀₋₁₀alkyl)₁₋₂aminocarbonyloxy C₀₋₁₀alkyl,

(C₃₋₈heterocyclyl C₀₋₁₀alkyl)₁₋₂aminocarbonyloxy C₀₋₁₀alkyl,

30 C₃₋₈heterocyclcarbonyloxy C₀₋₁₀alkyl, and

C₃₋₈heterocycloalkylcarbonyloxy C₀₋₁₀alkyl,

wherein in R¹, R², R³, and R⁴, said alkyl, alkenyl, alkynyl, aryl, heterocyclyl, heterocycloalkyl, and cycloalkyl are each optionally substituted with one or more groups chosen from hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy, halogen, CO₂H, cyano, O(C=O)C_{1-C₆}alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O₍₀₋₁₎(C₁₋₁₀)perfluoroalkyl, C₀₋₁₀alkylaminocarbonylamino, C₁₋₁₀alkyloxycarbonylamino, C₁₋₁₀alkylcarbonylamino, C₀₋₁₀alkylaminosulfonylamino, C₁₋₁₀alkylsulfonylamino, C₁₋₁₀alkylsulfonyl, C₀₋₁₀alkylaminosulfonyl, C₀₋₁₀alkylaminocarbonyl and NH₂.

2. A compound of Claim 1, chosen from:

(S)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide;
10 N-(2-fluoro-5-methylbenzyl)-2-phenylbutanamide;
(S)-N-(2-fluoro-5-(trifluoromethyl)pyridin-3-yl)methyl)-2-phenylbutanamide;
15 (S)-N-(5-bromo-2-fluorobenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-phenylbutanamide;
N-(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide;
(S)-N-(5-ethyl-2-fluorobenzyl)-2-phenylbutanamide;
N-(5-cyclopropyl-2-fluorobenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-vinylbenzyl)-2-phenylbutanamide;
N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-fluorophenyl)butanamide;
20 N-(5-ethyl-2-fluorobenzyl)-2-(4-chlorophenyl)butanamide;
N-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide;
(S)-N-((2-fluoro-5-methylpyridin-3-yl)methyl)-2-phenylbutanamide;
(S)-N-((5-ethyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide;
N-(5-bromo-2-fluorobenzyl)-2-phenylbutanamide;
25 N-(5-ethyl-2-fluorobenzyl)-2-(3-chlorophenyl)butanamide;
N-(5-ethyl-2-fluorobenzyl)-2-(3,4-dichlorophenyl)butanamide;
(S)-N-((5-cyclopropyl-2-fluoropyridin-3-yl)methyl)-2-phenylbutanamide;
(2R)-N-[(5-cyclopropyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
30 (2R)-N-[(5-ethyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2R)-N-[(5-methyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2R)-N-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide;
35 (2R)-N-(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;

(2*R*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;
(2*R*)-*N*-(5-chloro-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
(2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide;
(2*R*)-*N*-(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
5 (2*R*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
(2*S*)-*N*-(5-(cyclopropyl)-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2*S*)-*N*-(5-ethyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
10 (2*S*)-*N*-(5-methyl-2-fluoropyridin-3-yl)methyl]-2-(3,4-dichlorophenyl)butanamide;
(2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-bromophenyl)butanamide;
(2*S*)-*N*-(5-bromo-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;
15 (2*S*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(3-bromophenyl)butanamide;
(2*S*)-*N*-(5-chloro-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
(2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(4-bromophenyl)butanamide;
20 (2*S*)-*N*-(5-bromo-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
(2*S*)-*N*-(5-(cyclopropyl)-2-fluorobenzyl)-2-(4-bromophenyl)butanamide;
N-[5-(1,1-difluoroethyl)-2-fluorobenzyl]-2-phenylbutanamide;
(2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide;
25 (2*R*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide;
(2*R*)-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-(3-chlorophenyl)-2-hydroxybutanamide;
25 (2*R*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide;
(2*R*)-*N*-(5ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
(2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-hydroxy-2-phenylbutanamide;
30 (2*S*)-*N*-(2-fluoro-5-(trifluoromethyl)benzyl)-2-(3-chlorophenyl)-2-hydroxybutanamide;
(2*S*)-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
(2*S*)-2-cyclopropyl-*N*-(2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide;

(2*S*)-*N*-((5-ethyl-2-fluoropyridin-3-yl)methyl)-2-hydroxy-2-phenylbutanamide;
(2*R*)-3,3,3-trifluoro-*N*-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;
5 (2*R*)-3,3,4,4,4-pentafluoro-*N*-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylbutanamide;
(2*S*)-3,3,4,4,4-pentafluoro-*N*-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylbutanamide;
(2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
10 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide;
(2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-bromobenzyl)-2-hydroxy-2-phenylpropanamide;
15 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-chlorobenzyl)-2-hydroxy-2-phenylpropanamide;
(2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;
20 (2*R*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-cyclopropylbenzyl)-2-hydroxy-2-phenylbutanamide;
(2*R*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylbutanamide;
25 (2*S*)-3,3,4,4,4-pentafluoro-*N*-(2-fluoro-5-cyclopropylbenzyl)-2-hydroxy-2-phenylbutanamide;
(2*S*)-3,3,3-trifluoro-*N*-(2,3,5-trifluorobenzyl)-2-hydroxy-2-phenylpropanamide;
30 (2*R*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;
(2*R*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;
(2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;

(2*S*)-2-(4-chloro-3-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-(trifluoromethyl)benzyl]-2-hydroxypropanamide;

(2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

5 (2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-3-cyano-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

(2*R*)-3,3,3-trifluoro-*N*-(2-fluoro-4-cyano-5-ethylbenzyl)-2-hydroxy-2-phenylpropanamide;

10 (2*R*)-3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-3-oxo-2-phenylpropyl dimethylcarbamate;

(2*R*)-3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate;

(2*R*)-3-{[(2-fluoro-5-methylpyridin-3-yl)methyl]amino}-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate;

15 (2*R*)-3-{[(2-fluoro-5-methylpyridin-3-yl)methyl]amino}-3-oxo-2-phenylpropyl dimethylcarbamate;

(2*S*)-3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-3-oxo-2-phenylpropyl dimethylcarbamate;

(2*S*)-3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate;

20 (2*S*)-3-{[(2-fluoro-5-methylpyridin-3-yl)methyl]amino}-3-oxo-2-phenylpropyl pyrrolidine-1-carboxylate;

(2*S*)-3-{[(2-fluoro-5-methylpyridin-3-yl)methyl]amino}-3-oxo-2-phenylpropyl dimethylcarbamate;

25 3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-1-methyl-3-oxo-2-phenylpropylpyrrolidine-1-carboxylate;

3-{[2-fluoro-5-(trifluoromethyl)benzyl]amino}-2-hydroxy-3-oxo-2-phenylpropylpyrrolidine-1-carboxylate; and pharmaceutically acceptable salts and stereoisomers thereof.

30 3. The use of the compound of any one of Claims 1-2 or a pharmaceutically acceptable salt or stereoisomer thereof in the preparation of a medicament for the treatment or prevention of a condition selected from: weakened muscle tone, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis,

35

hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia, hematopoietic disorders, arthritic condition and joint repair, HIV-wasting, prostate cancer, cancer cachexia, muscular dystrophies, Alzheimer's disease, cognitive decline, sexual dysfunction, sleep apnea, benign prostate hyperplasia, abdominal adiposity, metabolic syndrome, type II diabetes, depression, premature ovarian failure, and autoimmune disease, in a mammal in need thereof.

4. The use of Claim 3, wherein said condition is osteoporosis.

5. A pharmaceutical composition comprising a compound of any one of Claims 1-2 or a pharmaceutically acceptable salt or stereoisomer thereof and a pharmaceutically acceptable carrier.

6. A composition of Claim 5, further comprising an active ingredient selected from: an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative, a bisphosphonate, an antiestrogen or a selective estrogen receptor modulator, an $\alpha\beta\beta$ integrin receptor antagonist, a cathepsin K inhibitor, a HMG-CoA reductase inhibitor, an osteoclast vacuolar ATPase inhibitor, an antagonist of VEGF binding to osteoclast receptors, an activator of peroxisome proliferator-activated receptor γ , calcitonin, a calcium receptor antagonist, parathyroid hormone or analog thereof, a growth hormone secretagogue, human growth hormone, insulin-like growth factor, a p38 protein kinase inhibitor, bone morphogenetic protein, an inhibitor of BMP antagonism, a prostaglandin derivative, vitamin D or vitamin D derivative, vitamin K or vitamin K derivative, ipriflavone, fluoride salts, dietary calcium supplements, and osteoprotegerin.

7. A composition of Claim 6, wherein said bisphosphonate is alendronate.

8. A process for making a pharmaceutical composition comprising combining a compound according to any one of Claims 1 to 2 or a pharmaceutically acceptable salt or stereoisomer thereof and a pharmaceutically acceptable carrier.

9. A use of Claim 3, wherein the arthritic condition is selected from rheumatoid arthritis and osteoarthritis.

10. A compound of Claim 1, wherein when R^2 is hydrogen, R^3 is other than hydrogen.

11. A compound of claim 2, chosen from:

3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;
(2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;

(2S)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;

3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

5 (2R)-3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

(2S)-3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

10 2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide;

(2R)-2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide;

(2S)-2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide;

15 and pharmaceutically acceptable salts thereof.

12. A compound of claim 11, chosen from:

(2R)-2-cyclopropyl-N-((2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide;

(2S)-2-cyclopropyl-N-((2-fluoro-5-methylpyridin-3-yl)methyl)-2-hydroxy-2-phenylacetamide;

20 (2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;

(2R)-3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

25 (2R)-2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide;

(2S)-2-(4-fluorophenyl)-3,3,3-trifluoro-[2-fluoro-5-cyclopropylbenzyl]-2-hydroxypropanamide;

30 and pharmaceutically acceptable salts thereof.

13. A compound of claim 12, chosen from:

(2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide;

(2R)-3,3,3-trifluoro-N-(2-fluoro-3-bromo-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide;

35 and pharmaceutically acceptable salts thereof.

14. A compound of Claim 13, chosen from:
(2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide and pharmaceutically acceptable salts thereof.
15. A compound of (2R)-3,3,3-trifluoro-N-[(2-fluoro-5-methylpyridin-3-yl)methyl]-2-hydroxy-2-phenylpropanamide.
16. A compound of Claim 2, chosen from:
(2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide and pharmaceutically acceptable salts and stereoisomers thereof.
17. A compound of Claim 17, chosen from:
(2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide and pharmaceutically acceptable salts thereof.
18. A compound of (2R)-3,3,3-trifluoro-N-(2-fluoro-5-trifluoromethylbenzyl)-2-hydroxy-2-phenylpropanamide.
19. A method of treating or preventing a condition selected from: weakened muscle tone, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia, hematopoietic disorders, arthritic condition and joint repair, HIV-wasting, prostate cancer, cancer cachexia, muscular dystrophies, Alzheimer's disease, cognitive decline, sexual dysfunction, sleep apnea, benign prostate hyperplasia, abdominal adiposity, metabolic syndrome, type II diabetes, depression, premature ovarian failure, and autoimmune disease, in a mammal in need thereof, said method comprising administering to said mammal a therapeutically effective amount of a compound of any one of Claims 1, 2, 10, 11, 12, 13, 14, 16 or a pharmaceutically acceptable salt or stereoisomer thereof.
20. The method of Claim 19, wherein said condition is osteoporosis, rheumatoid arthritis or osteoarthritis.

Dated 4 December 2008
Merck & Co., Inc.