



(86) Date de dépôt PCT/PCT Filing Date: 1993/10/04
(87) Date publication PCT/PCT Publication Date: 1994/04/28
(45) Date de délivrance/Issue Date: 2004/04/06
(85) Entrée phase nationale/National Entry: 1995/04/11
(86) N° demande PCT/PCT Application No.: EP 1993/002706
(87) N° publication PCT/PCT Publication No.: 1994/008979
(30) Priorité/Priority: 1992/10/15 (P 42 34 801.3) DE



(51) Cl.Int.⁶/Int.Cl.⁶ C07D 249/10, A01N 47/38, A01N 57/32, C07F 9/6518

(72) Inventeurs/Inventors:
FINDEISEN, KURT, DE;
LINKER, KARL-HEINZ, DE;
KLUTH, JOACHIM, DE;
MULLER, KLAUS-HELMUT, DE;
RIEBEL, HANS-JOCHEM, DE;
KONIG, KLAUS, DE;
SANTEL, HANS-JOACHIM, DE;
SCHMIDT, ROBERT R., DE

(73) Propriétaire/Owner:
BAYER AKTIENGESELLSCHAFT, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : SULFONYLAMINOCARBONYLTRIAZOLINONES
(54) Title: SULPHONYL AMINO CARBONYL TRIAZOLINONES



(57) **Abrégé/Abstract:**


The invention relates to new sulphonylaminocarbonyltriazolinones of the formula (I) (See formula I) in which Q represents oxygen, sulphur or one of the following groups (See formulae II, III, IV) (in which the radicals R¹, R², R³ and R⁴ have the meanings given in the description), as well as to salts of compounds of the formula (I), to processes and novel intermediates for the preparation of the novel compounds, and to their use as herbicides.



Sulphonylaminocarbonyltriazolinones


A b s t r a c t

The invention relates to new sulphonylaminocarbonyltriazolinones of the formula (I)



in which

Q represents oxygen, sulphur or one of the following groups



(in which the radicals R¹, R², R³ and R⁴ have the meanings given in the description),

as well as to salts of compounds of the formula (I), to processes and novel intermediates for the preparation of the novel compounds, and to their use as herbicides.

BAYER AKTIENGESELLSCHAFT

51368 Leverkusen

Konzernverwaltung RP


Patente Konzern

FILE. PIN IN THIS ATTACHED
TEXT TRANSLATIONSulphonylaminocarbonyltriazolinones

- 5 The invention relates to new sulphonylaminocarbonyltriazolinones, to a plurality of processes and novel intermediates for their preparation, and to their use as herbicides.


It has already been disclosed that certain sulphonylaminocarbonyltriazolinones, such as, forexample, 4,5-dimethyl-2-(2-chlorophenylsulphonyl-aminocarbonyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, have herbicidal properties (cf. EP-A 341489/LeA 26062; cf. also
10 EP-A 422469/LeA 27154; EP-A 425948/LeA 27155; EP-A 431291/LeA 27156). However, the activity of these compounds is not in all respects satisfactory.

There have now been found the new sulphonylaminocarbonyltriazolinones of the general formula (I)



in which

- 15 Q represents oxygen, sulphur or one of the following groups




- R¹ represents hydrogen, hydroxyl, amino, alkylideneamino, or represents a radical from the series consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aralkyl, aryl, alkoxy, alkenyloxy, alkylamino, cycloalkylamino, dialkylamino, alkanoylamino and alkoxycarbonylamino, each of which is optionally substituted,
- 5 R² represents hydrogen, hydroxyl, mercapto, amino, halogen, or represents a radical from the series consisting of alkyl, cycloalkyl, alkenyl, cycloalkenyl, aralkyl, aryl, alkoxy, alkenyloxy, alkynyloxy, cycloalkyloxy, aryloxy, aralkyloxy, alkylthio, alkylsulphinyl, alkylsulphonyl, alkenylthio, alkynylthio, cycloalkylthio, arylthio, aralkylthio, alkylamino, alkenylamino, arylamino, aralkylamino, 10 dialkylamino, aziridino, pyrrolidino, piperidino or morpholino, each of which is optionally substituted,
- R³ represents a radical from the series consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, alkylsulphonyl, alkylaminosulphonyl, dialkylaminosulphonyl, aryl and aralkyl, each of which is optionally substituted,
- 15 R⁴ represents hydrogen, hydroxyl, amino, cyano, alkoxycarbonyl, or represents alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, aralkoxy, aryloxy or dialkoxy(thio)phosphoryl, each of which is optionally substituted, or the following group Q¹-R⁵, in which
- Q¹ represents -CO- or -SO₂- and
- 20 R⁵ represents alkyl, cycloalkyl or aryl, each of which is optionally substituted,

and salts of compounds of the formula (I).

The new sulphonylaminocarbonyltriazolinones of the formula (I) are obtained when

- (a) chlorosulphonylaminocarbonyltriazolinones of the general formula (II)



in which

R^1 and R^2 have the abovementioned meaning,

are reacted with nucleophilic compounds of the general formula (III)




5 in which

Q and R^3 have the abovementioned meaning,

if appropriate in the presence of an acid acceptor and if appropriate in the presence of a diluent,

or when

10 (b) triazolinones of the general formula (IV)



in which

R^1 and R^2 have the abovementioned meaning,


15 are reacted with chlorosulphonyl isocyanate, if appropriate in the presence of a diluent, and the resulting chlorosulphonylaminocarbonyltriazolinones of the formula (II) - above - are reacted, without intermediate isolation, with

nucleophilic compounds of the general formula (III) - above - if appropriate in the presence of an acid acceptor and if appropriate in the presence of a diluent,


and, if appropriate, the compounds of the formula (I) obtained by process (a) or (b) are converted to salts by customary methods.

- 5 Other methods which are possible for the preparation of the compounds of the formula (I) according to the invention are given below, Q, R¹, R² and R³ in each case having the abovementioned meaning:


- (c) Reaction of isocyanates of the formula (V) with triazolinones of the formula (IV):



- (d) Reaction of aminosulphonyl compounds of the formula (VI) with oxycarbonyltriazolinones of the formula (VII) (R: alkyl, aralkyl, aryl):



- (e) Reaction of sulphonylurethanes of the formula (VIII) with triazolinones of the formula (IV) (R: alkyl, aralkyl, aryl):




- 5 The new sulphonylaminocarbonyltriazolinones of the general formula (I) are distinguished by a powerful herbicidal activity.

Surprisingly, the new compounds of the formula (I) display a considerably more powerful herbicidal activity than 4,5-dimethyl-2-(2-chlorophenylsulphonyl-amino-carbonyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, which is known and which has a similar structure.

10

The invention preferably relates to compounds of the formula (I) in which

Q represents oxygen, sulphur or one of the following groups



- 5 R^1 represents hydrogen, hydroxyl, amino, or represents C_2 - C_{10} -alkylideneamino, or represents C_1 - C_6 -alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylcarbonyl or C_1 - C_4 -alkoxy-carbonyl, or represents C_3 - C_6 -alkenyl or C_3 - C_6 -alkinyl, each of which is optionally substituted by fluorine, chlorine and/or bromine, or represents C_3 - C_6 -cycloalkyl which is optionally substituted by fluorine, chlorine, bromine and/or C_1 - C_4 -alkyl, or represents phenyl- C_1 - C_3 -alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl, C_1 - C_4 -alkoxy and/or C_1 - C_4 -alkoxy-carbonyl, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl, C_1 - C_4 -alkoxy, fluorine- and/or chlorine-substituted C_1 - C_3 -alkoxy, C_1 - C_4 -alkylthio, fluorine- and/or chlorine-substituted C_1 - C_3 -alkylthio, C_1 - C_4 -alkylsulphinyl, C_1 - C_4 -alkylsulphonyl and/or C_1 - C_4 -alkoxy-carbonyl, or represents C_1 - C_6 -alkoxy which is optionally substituted by fluorine, chlorine, cyano, phenyl, C_1 - C_4 -alkoxy or C_1 - C_4 -alkoxy-carbonyl, or represents C_3 - C_6 -alkenyloxy, or represents C_1 - C_4 -alkylamino, C_3 - C_6 -cycloalkylamino or di- $(C_1$ - C_4 -alkyl)-amino, each of which is optionally substituted by fluorine, cyano, C_1 - C_4 -alkoxy or C_1 - C_4 -alkoxy-carbonyl, or represents C_1 - C_6 -alkanoylamino, or represents C_1 - C_6 -alkoxycarbonylamino,
- 20 R^2 represents hydrogen, hydroxyl, mercapto, amino, halogen, or represents C_1 - C_6 -alkyl or C_2 - C_6 -alkenyl, optionally substituted by fluorine, chlorine, bromine, cyano, C_3 - C_6 -cycloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio or C_1 - C_4 -alkoxy-carbonyl, or represents C_3 - C_6 -cycloalkyl which is optionally substituted by fluorine, chlorine, bromine and/or C_1 - C_4 -alkyl, or represents cyclohexenyl, or represents phenyl- C_1 - C_3 -alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl, C_1 - C_4 -alkoxy and/or C_1 - C_4 -alkoxy-carbonyl, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl,

C₁-C₄-alkoxy, fluorine- and/or chlorine-substituted C₁-C₃-alkoxy, C₁-C₄-alkyl-
 thio, fluorine- and/or chlorine-substituted C₁-C₃-alkylthio, C₁-C₄-alkyl-sulphinyl,
 C₁-C₄-alkylsulphonyl and/or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkoxy
 5 which is optionally substituted by fluorine, chlorine, cyano, C₁-C₄-alkoxy or
 C₁-C₄-alkoxy-carbonyl, or represents C₃-C₆-alkenyloxy, C₃-C₆-alkinyloxy or C₃-
 C₆-cycloalkyloxy, or represents phenoxy or benzyloxy, each of which is
 optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl,
 trifluoromethyl, C₁-C₄-alkoxy, difluoromethoxy or trifluoromethoxy, or
 10 represents C₁-C₆-alkylthio, C₁-C₆-alkylsulphinyl, C₁-C₆-alkylsulphonyl, C₃-C₆-
 alkenylthio, C₃-C₆-alkinylthio or C₃-C₆-cycloalkylthio, each of which is
 optionally substituted by fluorine, chlorine, cyano, C₁-C₄-alkoxy or C₁-C₄-
 alkoxy-carbonyl, or represents phenylthio or benzylthio, each of which is
 optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl,
 trifluoromethyl, C₁-C₄-alkoxy, difluoromethoxy or trifluoromethoxy, or
 15 represents C₁-C₆-alkylamino or C₃-C₆-alkenylamino, or represents phenylamino
 or benzylamino, each of which is optionally substituted by fluorine, chlorine,
 bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy,
 difluoromethoxy or trifluoromethoxy, or represents di-(C₁-C₄-alkyl)-amino, or
 represents aziridino, pyrrolidino or morpholino, each of which is optionally
 20 substituted by C₁-C₄-alkyl,

R³ represents C₁-C₆-alkyl, C₃-C₆-alkenyl or C₃-C₆-alkinyl, each of which is
 optionally substituted by halogen, cyano, carboxyl, C₁-C₄-alkoxy, C₁-C₄-
 alkylthio, C₁-C₄-alkylsulphinyl, C₁-C₄-alkylsulphonyl or C₁-C₄-alkoxy-carbonyl,
 or represents C₃-C₆-cycloalkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl, each of which
 25 is optionally substituted by halogen, cyano, carboxyl, C₁-C₄-alkyl, C₁-C₄-
 halogenoalkyl or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkylsulphonyl, C₁-
 C₆-alkylaminosulphonyl or di-(C₁-C₄-alkyl)-aminosulphonyl, each of which is
 optionally substituted by halogen, or represents phenyl, naphthyl, tetralinyl,
 phenyl-C₁-C₄-alkyl, naphthyl-C₁-C₄-alkyl or tetralinyl-C₁-C₄-alkyl, each of which
 30 is optionally substituted by halogen, cyano, carboxyl, nitro, by C₁-C₄-alkyl,
 C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulphinyl or C₁-C₄-alkylsulphonyl (in
 each case optionally substituted by halogen, cyano, C₁-C₄-alkoxy or C₁-C₄-

alkoxy-carbonyl), by C₁-C₄-alkylsulphonyloxy, C₁-C₄-alkylaminosulphonyloxy, di-(C₁-C₄-alkyl)aminosulphonyloxy, C₁-C₄-halogenoalkylsulphonyloxy, di-(C₁-C₄-alkyl)-aminocarbonyl or C₁-C₄-alkylaminocarbonyloxy, by C₃-C₆-cycloalkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl (in each case optionally substituted by halogen or C₁-C₄-alkyl), by phenyl, phenoxy, phenylthio, phenylsulphinyl, phenylsulphonyl, phenylsulphonyloxy, phenylamino, phenylcarbonyl or phenyl-C₁-C₄-alkyl (in each case optionally substituted by halogen, cyano, nitro, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkylthio, C₁-C₄-alkoxy-carbonyl or C₁-C₄-alkoxy-amino), by C₁-C₄-alkyl-carbonyl, C₃-C₆-cycloalkylcarbonyl or C₁-C₄-alkoxy-carbonyl (in each case optionally substituted by halogen, C₃-C₆-cycloalkyl or C₁-C₄-alkoxy),

R⁴ represents hydrogen, hydroxyl, amino, cyano, or represents C₁-C₆-alkyl, C₃-C₆-alkenyl or C₃-C₆-alkinyl, each of which is optionally substituted by halogen, cyano, carboxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulphinyl, C₁-C₄-alkylsulphonyl, C₁-C₄-alkylamino, di-(C₁-C₄-alkyl)-amino or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, benzyloxy, phenoxy or di-(C₁-C₄-alkoxy)-(thio)-phosphoryl, or represents the following group Q¹-R⁵ in which

Q¹ represents -CO- or -SO₂- and

R⁵ represents C₁-C₆-alkyl which is optionally substituted by halogen or C₁-C₄-alkoxy, or represents C₃-C₆-cycloalkyl which is optionally substituted by halogen or C₁-C₄-alkyl, or represents phenyl which is optionally substituted by halogen, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl or C₁-C₄-alkoxy.

The invention furthermore preferably relates to the sodium, potassium, magnesium, calcium, ammonium, C₁-C₄-alkylammonium, di-(C₁-C₄-alkyl)-ammonium, tri-(C₁-C₄-alkyl)-ammonium, C₅- or C₆-cycloalkylammonium and di-(C₁-C₂-alkyl)-benzylammonium salts of compounds of the formula (I) in which Q, R¹, R² and R³ have the

meanings mentioned above as being preferred.

In particular, the invention relates to compounds of the formula (I), in which

Q represents oxygen or the group



- 5 R¹ represents hydrogen, hydroxyl, amino, or represents C₃-C₈-alkylideneamino, or represents methyl, ethyl, n- or i-propyl, n-, i-, s- or t-butyl, dichloromethyl, fluoroethyl, chloroethyl, difluoroethyl, dichloroethyl, trifluoroethyl, trichloroethyl, chlorodifluoroethyl, tetrafluoroethyl, cyanomethyl, cyanoethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, or represents allyl, chloroallyl, dichloroallyl, propargyl, or represents chloropropyl, benzyl or phenyl, or
- 10 represents methoxy, ethoxy, n- or i-propoxy, or represents allyloxy, n- or i- or s-butoxy, or represents methylamino, ethylamino, n- or i-propylamino, or represents cyclopropylamino, dimethylamino, diethylamino, acetylamino, methoxycarbonylamino or ethoxycarbonylamino,
- 15 R² represents hydrogen, hydroxyl, mercapto, amino, fluorine, chlorine, bromine, or represents methyl, ethyl, n- or i-propyl, n-, i-, s- or t-butyl, difluoromethyl, trifluoromethyl, dichloromethyl, trichloromethyl, cyanomethyl, cyanoethyl, cyclopropylmethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl, ethylthiomethyl, methylthioethyl or ethylthioethyl, or represents cyclopropyl, difluorocyclopropyl or dichlorocyclopropyl, or represents
- 20 phenyl or benzyl, or represents methoxy, ethoxy, n- or i-propoxy, or represents methoxymethoxy, ethoxymethoxy, methoxyethoxy or ethoxy-ethoxy, or represents phenoxy or benzyloxy, or represents methylthio, ethylthio, n- or i-propylthio, allylthio, propargylthio, cyclopropylmethylthio, methylsulphinyl, ethylsulphinyl, methylsulphonyl, ethylsulphonyl, phenylthio or benzylthio, or
- 25 represents methylamino, ethylamino, n- or i-propylamino, phenylamino or benzylamino, or represents dimethylamino, diethylamino, dipropylamino, methylethylamino, methylpropylamino, or represents aziridino, or represents

pyrrolidino, piperidino or morpholino, each of which is optionally substituted by methyl or ethyl,

- R^3 represents methyl, ethyl, n- or i-propyl or n-, i-, s- or t-butyl, each of which is optionally substituted by fluorine, chlorine, cyano, methoxy, ethoxy, methoxycarbonyl or ethoxycarbonyl, or represents cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl, cyclopentylethyl or cyclohexylethyl, each of which is optionally substituted by fluorine, chlorine, cyano, methyl, ethyl, trifluoromethyl, methoxycarbonyl or ethoxycarbonyl, or represents methylsulphonyl, ethylsulphonyl, propylsulphonyl or butylsulphonyl, optionally substituted by fluorine or chlorine, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, carboxyl, nitro, by methyl, ethyl, methoxy, ethoxy, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, methylsulphonyl or ethylsulphonyl (in each case optionally substituted by fluorine, chlorine, cyano, methoxy, ethoxy, methoxycarbonyl or ethoxycarbonyl), by methylsulphonyloxy, ethylsulphonyloxy, methylaminosulphonyloxy, ethylaminosulphonyloxy, dimethylaminosulphonyloxy, diethylaminosulphonyloxy, trifluoromethylsulphonyloxy, dimethylaminocarbonyl or diethylaminocarbonyl, by cyclohexyl or cyclohexylmethyl (in each case optionally substituted by chlorine, methyl or ethyl), by phenyl, phenoxy, phenylthio, phenylsulphonyl, phenylsulphonyloxy, phenylamino, phenylcarbonyl, phenylmethyl or phenylethyl (in each case optionally substituted by fluorine, chlorine, cyano, nitro, methyl, ethyl, trifluoromethyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, methylthio, ethylthio, trifluoromethylthio, methoxycarbonyl or ethoxycarbonyl), by acetyl, cyclopropylcarbonyl, methoxycarbonyl or ethoxycarbonyl (optionally substituted by fluorine, chlorine, cyclopentyl, cyclohexyl, methoxy or ethoxy), or represents naphthyl or tetralinyl, each of which is optionally substituted by fluorine, chlorine, cyano, nitro, methyl, ethyl, methoxy or ethoxy, or represents naphthylmethyl, naphthylethyl, tetralinylmethyl or tetralinylethyl, each of which is optionally substituted by fluorine, chlorine, methyl, ethyl, methoxy or ethoxy,

5 R⁴ represents hydrogen, cyano, cyanomethyl, cyanoethyl, difluoromethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylethyl, allyl, propargyl, methoxycarbonyl, ethoxycarbonyl, methoxy, ethoxy, allyloxy or benzyloxy, or represents the group Q¹-R⁵ in which

Q¹ represents -CO- or -SO₂- and

10 R⁵ represents methyl, ethyl or propyl, each of which is optionally substituted by fluorine, chlorine, methoxy or ethoxy, or represents cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is optionally substituted by fluorine, chlorine, methyl or ethyl, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, methoxy or ethoxy.

15 The definitions of radicals mentioned above, in general or where preferred ranges are mentioned, apply to the end products of the formula (I) and, analogously, to the starting materials or intermediates required in each case for the preparation. These definitions of radicals can be combined with each other as desired, that is to say combinations between the preferred ranges indicated are also possible.

20 The hydrocarbon radicals mentioned in the definitions of the radicals, such as alkyl, alkenyl or alkynyl, also in combinations with hetero atoms, such as in alkoxy, alkylthio or alkylamino, are straight-chain or branched, even when this is not expressly stated.

Halogen generally represents fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine, in particular fluorine or chlorine.

25 Examples of the compounds of the formula (I) according to the invention are listed in Table 1 below.




Table 1: Examples of the compounds of the formula (I)






Q	R ¹	R ²	R ³
O	CH ₃	C ₂ H ₅	n-C ₃ H ₇
NH	CH ₃	C ₂ H ₅	i-C ₃ H ₇
O	C ₂ H ₅	C ₂ H ₅	
NH	OCH ₃	n-C ₃ H ₇	i-C ₄ H ₉
NH	OC ₂ H ₅	CH ₃	
NCH ₃	CH ₃	i-C ₃ H ₇	
NH	C ₂ H ₅		

Table 1: (continued)












Q	R ¹	R ²	R ³
NH		C ₂ H ₅	
NH	C ₂ H ₅	SCH ₃	
NH	C ₂ H ₅	OC ₂ H ₅	
NCH ₃	OCH ₃	OCH ₃	n-C ₄ H ₉
NH		SCH ₃	
NH		OC ₂ H ₅	
NH	C ₂ H ₅	OCH ₃	
NCH ₃	CH ₃	OC ₂ H ₅	
NH	C ₂ H ₅	SC ₂ H ₅	

Table 1: (continued)




















Q	R ¹	R ²	R ³
O	OC ₂ H ₅	C ₂ H ₅	
NH	CH ₃	SC ₂ H ₅	
NH	C ₂ H ₅	OC ₂ H ₅	
NH	CH ₃		
NH	CH ₃	CH ₃	
NH	C ₂ H ₅		
NH		OCH ₃	
NH	OCH ₃		-CH ₂ COOC ₂ H ₅


Table 1: (continued)

Q	R ¹	R ²	R ³
NH	C ₂ H ₅	N(CH ₃) ₂	
NH	C ₂ H ₅		
O	C ₂ H ₅	Cl	
NH	CH ₃	Br	
NH	CH ₃	i-OC ₃ H ₇	
O	C ₂ H ₅	N(CH ₃) ₂	
NH	CH ₃	-CH ₂ SC ₂ H ₅	-CH ₂ CH ₂ OC ₂ H ₅

If, for example, 2-chlorosulphonylaminocarbonyl-4-ethoxy-5-ethyl-2,4-dihydro-3H-1,2,4-triazol-3-one and 2-fluoro-phenol are used as starting substances, the course of the reaction in process (a) according to the invention can be outlined by the following equation:



- 5 If, for example, 4-ethyl-5-ethylthio-2,4-dihydro-3H-1,2,4-triazol-3-one and chloro-sulphonyl isocyanate and then methyl 2-amino-benzoate are used as starting substances, the course of the reaction in process (b) according to the invention can be outlined by the following equation:



Formula (II) provides a general definition of the chlorosulphonylaminocarbonyl-triazolinones to be used as starting substances in process (a) according to the invention for the preparation of compounds of the formula (I).


5 In formula (II), R^1 and R^2 preferably, or in particular, have those meanings which have already been mentioned above in connection with the description of the compounds of the formula (I) according to the invention as being preferred, or particularly preferred, for R^1 and R^2 .

The following may be mentioned as examples of the starting substances of the formula (II):

10 4,5-dimethyl-, 4,5-diethyl-, 4-ethyl-5-methyl-, 5-ethyl-4-methyl-, 4-methyl-5-propyl-, 4-ethyl-5-propyl-, 4-cyclopropyl-5-ethyl-, 5-cyclopropyl-4-methyl-, 5-cyclopropyl-4-ethyl-, 4-methyl-5-chloro-, 4-ethyl-5-chloro-, 4-methyl-5-bromo-, 4-ethyl-5-bromo-, 4-cyclopropyl-5-chloro-, 4-cyclopropyl-5-bromo-, 4-methoxy-5-methyl-, 4-ethoxy-5-methyl-, 4-ethoxy-5-ethyl-, 4-methoxy-5-ethyl-, 4-methyl-5-methoxy-, 4-ethyl-5-methoxy-, 4-methyl-5-ethylthio-, 4-methyl-5-ethylthio-, 4-ethyl-5-methylthio-, 4-ethyl-5-ethylthio, 4-cyclopropyl-5-methoxy-, 4-cyclopropyl-5-ethoxy-, 4-cyclopropyl-5-methylthio-, 4-cyclopropyl-5-ethylthio-, 4-methoxy-5-cyclopropyl-, 4-ethoxy-5-cyclopropyl-, 4-methoxy-5-ethoxy-, 4,5-dimethoxy-, 4,5-diethoxy-, 4-methyl-5-dimethylamino-, 4-ethyl-5-dimethylamino- and
20 4-cyclopropyl-5-dimethylamino-2-chlorosulphonylaminocarbonyl-2,4-dihydro-3H-1,2,4-triazol-3-one.

The chlorosulphonylaminocarbonyltriazolinones of the general formula (II) are as yet unknown from the literature and, being new substances, also a subject-matter of the present invention.

25 The new compounds of the formula (II) are obtained when triazolinones of the general formula (IV)



in which

R¹ and R² have the abovementioned meaning

are reacted with chlorosulphonyl isocyanate at temperatures of between -20°C and +50°C, if appropriate in the presence of a diluent such as, for example, methylene chloride, and the product is worked up in the customary manner (cf. the preparation examples).

Formula (III) provides a general definition of the nucleophilic compounds to be used as starting substances in processes (a) and (b) according to the invention for the preparation of compounds of the formula (I).

In formula (III) Q and R³ preferably, or in particular, have those meanings which have already been mentioned above in connection with the description of the compounds of the formula (I) according to the invention as being preferred, or particularly preferred, for Q and R³.

The starting substances of the formula (III) are known chemicals for organic synthesis.

Formula (IV) provides a general definition of the triazolinones to be used as starting substances in processes (a) and (b) according to the invention for the preparation of compounds of the formula (I).

In formula (IV) R¹ and R² preferably, or in particular, have those meanings which have already been mentioned above in connection with the description of the compounds of the formula (I) according to the invention as being preferred, or particularly preferred, for R¹ and R².

The starting substances of the formula (IV) are known and/or can be prepared by processes known per se (cf. EP-A 283876, EP-A 294666, EP-A 301946, EP-A 298371, EP-A 341489, EP-A 399294, EP-A 398096, EP-A 422469, EP-A 425948, EP-A 431291, EP-A 477646, DE-OS 4110795 (German Published Specification)).

5 Processes (a) and (b) according to the invention for the preparation of the new compounds of the formula (I) are preferably carried out using diluents. Suitable diluents are virtually all inert organic solvents. These preferably include aliphatic and aromatic, optionally halogenated hydrocarbons such as pentane, hexane, heptane, cyclohexane, petroleum ether, benzine, ligroin, benzene, toluene, xylene, methylene chloride,
 10 ethylene chloride, chloroform, carbon tetrachloride, chlorobenzene and o-dichlorobenzene, ethers such as diethyl ether, dibutyl ether, glycol dimethyl ether, diglycol dimethyl ether, tetrahydrofuran and dioxane, ketones such as acetone, methyl ethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone, esters such as methyl acetate and ethyl acetate, nitriles such as, for example, acetonitrile and propionitrile,
 15 amides such as, for example, dimethylformamide, dimethylacetamide and N-methylpyrrolidone, and also dimethyl sulphoxide, tetramethylene sulphone and hexamethylphosphoric triamide.

Acid acceptors which can be employed in processes (a) and (b) according to the invention are all acid-binding agents which can conventionally be used for reactions of
 20 this type. The following are preferably suitable: alkali metal hydroxides such as, for example, sodium hydroxide and potassium hydroxide, alkaline earth metal hydroxides such as, for example, calcium hydroxide, alkali metal carbonates and alkali metal alcoholates such as sodium carbonate and potassium carbonate, sodium tert-butylate and potassium tert-butylate, furthermore basic nitrogen compounds such as Trimethylamine,
 25 Triethylamine, Tripropylamine, Tributylamine, Diisobutylamine, Dicyclohexylamine, Ethyldiisopropylamine, Ethyldicyclohexylamine, N,N-Dimethylbenzylamine, N,N-Dimethyl-aniline, Pyridine, 2-Methyl-, 3-Methyl-, 4-Methyl-, 2,4-Dimethyl-, 2,6-Dimethyl-, 2-Ethyl-, 4-Ethyl- and 5-Ethyl-2-methyl-pyridine, 1,5-Diazabicyclo-[4,3,0]-non-5-en (DBN), 1,8-Diazabicyclo-[5,4,0]-undec-7-en (DBU) and 1,4-Diazabicyclo-
 30 [2,2,2]-octane (DABCO).

When carrying out processes (a) and (b) according to the invention, the reaction temperatures can be varied within a substantial range. In general, the processes are carried out at temperatures between -20°C and $+80^{\circ}\text{C}$, preferably at temperatures between -10°C and $+50^{\circ}\text{C}$.

- 5 Processes (a) and (b) according to the invention are generally carried out under atmospheric pressure. However, it is also possible to carry out the processes under elevated or reduced pressure.

To carry out processes (a) and (b) according to the invention, the starting substances required in each case are generally employed in approximately equimolar amounts.

- 10 However, one of the components employed in each case may also be used in a larger excess. In general, the reactions are carried out in a suitable diluent in the presence of an acid acceptor, and the reaction mixture is stirred for several hours at the temperature required in each case. Working-up in processes (a) and (b) according to the invention is carried out in each case by customary methods (cf. the preparation examples).

- 15 If appropriate, salts may be prepared from the compounds of the general formula (I) according to the invention. Such salts are obtained in a simple fashion by customary salt formation methods, for example by dissolving or dispersing a compound of the formula (I) in a suitable solvent such as, for example, methylene chloride, acetone, tert-butyl methyl ether or toluene, and adding a suitable base. The salts can then be isolated
20 by concentration or filtration with suction, if appropriate after prolonged stirring.

- The active compounds according to the invention can be used as defoliants, desiccants, agents for destroying broad-leaved plants and, especially, as weed-killers. By weeds, in the broadest sense, there are to be understood all plants which grow in locations where they are undesired. Whether the substances according to the invention act as total or
25 selective herbicides depends essentially on the amount used.

The active compounds according to the invention can be used, for example, in connection with the following plants:

Dicotyledon weeds of the genera: Sinapis, Lepidium, Galium, Stellaria, Matricaria, Anthemis, Galinsoga, Chenopodium, Urtica, Senecio, Amaranthus, Portulaca, Xanthium, Convolvulus, Ipomoea, Polygonum, Sesbania, Ambrosia, Cirsium, Carduus, Sonchus, Solanum, Rorippa, Rotala, Lindernia, Lamium, Veronica, Abutilon, Emex,
 5 Datura, Viola, Galeopsis, Papaver, Centaurea, Trifolium, Ranunculus and Taraxacum.

Dicotyledon cultures of the genera: Gossypium, Glycine, Beta, Daucus, Phaseolus, Pisum, Solanum, Linum, Ipomoea, Vicia, Nicotiana, Lycopersicon, Arachis, Brassica, Lactuca, Cucumis and Cucurbita.

Monocotyledon weeds of the genera: Echinochloa, Setaria, Panicum, Digitaria,
 10 Phleum, Poa, Festuca, Eleusine, Brachiaria, Lolium, Bromus, Avena, Cyperus, Sorghum, Agropyron, Cynodon, Monochoria, Fimbristylis, Sagittaria, Eleocharis, Scirpus, Paspalum, Ischaemum, Sphenoclea, Dactyloctenium, Agrostis, Alopecurus and Apera.

Monocotyledon cultures of the genera: Oryza, Zea, Triticum, Hordeum, Avena, Secale,
 15 Sorghum, Panicum, Saccharum, Ananas, Asparagus and Allium.

However, the use of the active compounds according to the invention is in no way restricted to these genera, but also extends in the same manner to other plants.

The compounds are suitable, depending on the concentration, for the total combating of weeds, for example on industrial terrain and rail tracks, and on paths and squares
 20 with or without tree plantings. Equally, the compounds can be employed for combating weeds in perennial cultures, for example afforestations, decorative tree plantings, orchards, vineyards, citrus groves, nut orchards, banana plantations, coffee plantations, tea plantations, rubber plantations, oil palm plantations, cocoa plantations, soft fruit plantings and hopfields, in lawns, turf and pasture-land, and for the selective combating
 25 of weeds in annual cultures.

The compounds of the formula (I) according to the invention are particularly suitable for selectively combating dicotyledon weeds in monocotyledon cultures by the pre- as

well as post-emergence methods.

Moreover, the active compounds (I) according to the invention also display interesting secondary actions, i.e. a leaf-acting insecticidal action, as well as fungicidal actions in particular against *Pyricularia oryzae*, for example in rice, and against *Erysiphe graminis*.

The active compounds can be converted into the customary formulations, such as solutions, emulsions, wettable powders, suspensions, powders, dusting agents, pastes, soluble powders, granules, suspension-emulsion concentrates, natural and synthetic materials impregnated with active compound, and very fine capsules in polymeric substances.

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is liquid solvents and/or solid carriers, optionally with the use of surface-active agents, that is emulsifying agents and/or dispersing agents and/or foam-forming agents.

In the case of the use of water as an extender, organic solvents can, for example, also be used as auxiliary solvents. As liquid solvents, there are suitable in the main: aromatics, such as xylene, toluene or alkyl-naphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons, such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons, such as cyclohexane or paraffins, for example petroleum fractions, mineral and vegetable oils, alcohols, such as butanol or glycol as well as their ethers and esters, ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents, such as dimethylformamide and dimethyl sulphoxide, as well as water.

As solid carriers there are suitable:

for example ammonium salts and ground natural minerals, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as highly disperse silica, alumina and silicates, as solid carriers for

- granules there are suitable: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, as well as synthetic granules of inorganic and organic meals, and granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks; as emulsifying and/or foam-forming agents there are suitable: for example non-ionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates as well as albumen hydrolysis products; as dispersing agents there are suitable: for example lignin-sulphite waste liquors and methylcellulose.
- 5
- 10 Adhesives such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latexes, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, as well as natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids, can be used in the formulations. Further additives can be mineral and vegetable oils.
- 15 It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs, such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

20 The formulations in general contain between 0.1 and 95 per cent by weight of active compound, preferably between 0.5 and 90%.

For combating weeds, the active compounds according to the invention, as such or in the form of their formulations, can also be used as mixtures with known herbicides, finished formulations or tank mixes being possible.

25 Suitable herbicides for the mixtures are known herbicides, for example anilides such as, for example, diflufenican and propanil; arylcarboxylic acids such as, for example, dichloropicolinic acid, dicamba and picloram; aryloxyalkanoic acids such as, for example, 2,4 D, 2,4 DB, 2,4 DP, fluroxypyr, MCPA, MCPP and triclopyr; aryloxyphenoxy-alkanoic esters such as, for example, diclofop-methyl, fenoxaprop-ethyl,

fluazifop-butyl, haloxyfop-methyl and quizalofop-ethyl; azinones such as, for example, chloridazon and norflurazon; carbamates such as, for example, chlorpropham, desmedipham, phenmedipham and propham; chloroacetanilides such as, for example, alachlor, acetochlor, butachlor, metazachlor, metolachlor, pretilachlor and propachlor;

5 dinitroanilines such as, for example, oryzalin, pendimethalin and trifluralin; diphenyl ethers such as, for example, acifluorfen, bifenox, fluoroglycofen, fomesafen, halosafen, lactofen and oxyfluorfen; ureas such as, for example, chlortoluron, diuron, fluometuron, isoproturon, linuron and methabenzthiazuron; hydroxylamines such as, for example, alloxydim, clethodim, cycloxydim, sethoxydim and tralkoxydim; imidazolinones such

10 as, for example, imazethapyr, imazamethabenz, imazapyr and imazaquin; nitriles such as, for example, bromoxynil, dichlobenil and ioxynil; oxyacetamides such as, for example, mefenacet; sulphonylureas such as, for example, amidosulfuron, bensulfuron-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, metsulfuron-methyl, nicosulfuron, primisulfuron, pyrazosulfuron-ethyl, thifensulfuron-methyl, triasulfuron

15 and tribenuron-methyl; thiocarbamates such as, for example, butylate, cycloate, di-allate, EPTC, esprocarb, molinate, prosulfocarb, thiobencarb and tri-allate; triazines such as, for example, atrazine, cyanazine, simazine, simetryn, terbutryn and terbutylazine; triazinones such as, for example, hexazinone, metamitron and metribuzin; others such as, for example, aminotriazole, benfuresate, bentazone, cinmethylin,

20 clomazone, clopyralid, difenzoquat, dithiopyr, ethofumesate, fluorochloridone, glufosinate, glyphosate, isoxaben, pyridate, quinchlorac, quinmerac, sulphosate and tridiphan.

A mixture with other known active compounds, such as fungicides, insecticides, acaricides, nematocides, birdrepellants, plant nutrients and agents which improve soil

25 structure, is also possible.


The active compounds can be used as such, in the form of their formulations or in the use forms prepared therefrom by further dilution, such as ready-to-use solutions, suspensions, emulsions, powders, pastes and granules. They are used in the customary manner, for example by watering, spraying, atomizing or scattering.

30 The active compounds according to the invention can be applied either before or after

emergence of the plants. They can also be incorporated into the soil before sowing.


The amount of active compound used can vary within a substantial range. It depends essentially on the nature of the desired effect. In general, the amounts used are between 10 g and 10 kg of active compound per hectare of soil surface, preferably between 50
5 g and 5 kg per ha.

The preparation and use of the active compounds according to the invention can be seen from the following examples.

Preparation Examples:Example 1(Process (a))

2.84 g (0.01 mol) of 2-chlorosulphonylaminocarbonyl-4-methyl-5-ethoxy-2,4-dihydro-3H-1,2,4-triazol-3-one are introduced into 100 ml of methylene chloride, and 1.24 g (0.01 mol) of 2-methoxy-phenol and 2.02 g (0.02 mol) of triethylamine are added to this mixture at 20°C. The reaction mixture is stirred for 60 minutes at 20°C. It is then washed three times using 150 ml of water in each case, dried using sodium sulphate and filtered. The filtrate is concentrated under a water pump vacuum, and the residue is recrystallized from isopropanol.


2.5 g (53 % of theory) of the triethylammonium salt of 2-(2-methoxy-phenoxy)-sulphonylaminocarbonyl-4-methyl-5-ethoxy-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 107°C are obtained.

Example 2(Process (a))

5.69 g (0.02 mol) of 2-chlorosulphonylaminocarbonyl-4-methyl-5-ethoxy-2,4-dihydro-


3H-1,2,4-triazol-3-one are introduced into 150 ml of methylene chloride, and a solution of 6.6 g (0.04 mol) of ethyl 2-amino-benzoate in 20 ml of methylene chloride is added to this mixture at 20°C. The reaction mixture is stirred for 60 minutes at 20°C. It is then washed three times using 150 ml of water in each case, dried using sodium sulphate and filtered. The filtrate is concentrated under a water pump vacuum and the residue is recrystallized from isopropanol.

5.7 g (69 % of theory) of 2-(2-ethoxy-carbonyl-phenylamino)-sulphonylaminocarbonyl-4-methyl-5-ethoxy-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 135°C are obtained.

Example 3(Process (a))

5.69 g (0.02 mol) of 2-chlorosulphonylaminocarbonyl-4-ethyl-5-methoxy-2,4-dihydro-3H-1,2,4-triazol-3-one are introduced into 150 ml of methylene chloride, and a solution of 2.74 g (0.02 mol) of 2-methoxy-N-methylaniline and 2.02 g (0.02 mol) of triethylamine in 20 ml of methylene chloride is added to this mixture at 20°C. The reaction mixture is stirred for 60 minutes at 20°C. It is then washed three times using 150 ml of water in each case, dried using sodium sulphate and filtered. The filtrate is concentrated under a water pump vacuum and the residue is recrystallized from isopropanol.

3.2 g (40 % of theory) of 2-(2-methoxy-N-methyl-phenylamino)-sulphonylaminocarbonyl-4-ethyl-5-methoxy-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 134°C are obtained.

Example 4

(Process (b))

4.3 g (0.03 mol) of 4-methyl-5-ethoxy-2,4-dihydro-3H-1,2,4-triazol-3-one are introduced into 150 ml of methylene chloride, and the mixture is cooled to -10°C . A solution of 4.3 g (0.03 mol) of chlorosulphonyl isocyanate in 20 ml of methylene chloride is then added, and the mixture is stirred for 30 minutes. Then, a solution of 4.53 g (0.03 mol) of methyl 2-amino-benzoate and 6.06 g (0.06 mol) of triethylamine in 20 ml of methylene chloride is added dropwise at 20°C , and the reaction mixture is stirred for a further 30 minutes. It is subsequently washed three times using 150 ml of water in each case, dried using sodium sulphate and filtered. The filtrate is concentrated under a water pump vacuum, the residue is stirred with ethyl acetate, and the product which has been obtained as crystals is isolated by filtration with suction.

11 g (73 % of theory) of the triethylammonium salt of 2-(2-methoxycarbonyl-phenyl-amino)-sulphonylaminocarbonyl-4-methyl-5-ethoxy-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 142°C (decomposition) are obtained.

Other examples of the compounds of the formula (I) which can be prepared analogously to Preparation Examples 1 to 4 and in accordance with the general description of the preparation processes according to the invention are those listed in Table 2 below.




Table 2: Preparation examples of the compounds of the formula (I)



Ex. No.	Q	R ¹	R ²	R ³	Physical data
5	NH	CH ₃	CH ₃	s-C ₄ H ₉	m.p.: 80°C
6	NH	CH ₃	CH ₃		m.p.: 68°C
7	NH	CH ₃	CH ₃	t-C ₄ H ₉	m.p.: 234°C
8	NCH ₃	CH ₃	CH ₃	i-C ₃ H ₇	m.p.: 113°C
9	NCH ₃	CH ₃	OC ₂ H ₅	i-C ₃ H ₇	m.p.: 129°C
10	NH	CH ₃	OC ₂ H ₅		m.p.: 153°C
11	NH	CH ₃	OC ₂ H ₅	t-C ₄ H ₉	m.p.: 148°C
12	NH	CH ₃	OC ₂ H ₅	s-C ₄ H ₉	m.p.: 137°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
13	NH	CH ₃	OC ₂ H ₅		m.p.: 107°C
14	NH	CH ₃	CH ₃		m.p.: 153°C
15	NH	CH ₃	CH ₃		m.p.: 140°C Isolated in the form of the triethylammonium salt
16	NH	C ₂ H ₅	OCH ₃		m.p.: 111°C
17	NH	C ₂ H ₅	OCH ₃		m.p.: 145°C
18	NH	C ₂ H ₅	OCH ₃		m.p.: 175°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
19	NH	C ₂ H ₅	OCH ₃		m.p.: 163°C
20	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 3.60-3.78; 4.10; 5.30 ppm
21	NH	C ₂ H ₅	OCH ₃		m.p.: 148°C
22	NH	C ₂ H ₅	OCH ₃		m.p.: 133°C
23	NH	C ₂ H ₅	OCH ₃		m.p.: 147°C
24	NH	C ₂ H ₅	OCH ₃		m.p.: 144°C
25	NH	C ₂ H ₅	OCH ₃		m.p.: 164°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
26	NH	C ₂ H ₅	OCH ₃		m.p.: 160°C
27	NH	C ₂ H ₅	OCH ₃		m.p.: 183°C
28	NH	C ₂ H ₅	OCH ₃		m.p.: 157°C
29	NH	C ₂ H ₅	OCH ₃		m.p.: 156°C
30	NH	CH ₃	OC ₂ H ₅		m.p.: 160°C
31	NH	CH ₃	OC ₂ H ₅		m.p.: 130°C
32	NH	CH ₃	OC ₂ H ₅		m.p.: 160°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
33	NH	CH ₃	OC ₂ H ₅		m.p.: 149°C
34	NH	CH ₃	OC ₂ H ₅		m.p.: 131°C
35	NH	CH ₃	OC ₂ H ₅		m.p.: 151°C
36	NH	CH ₃	OC ₂ H ₅		m.p.: 155°C
37	NH	CH ₃	OC ₂ H ₅		m.p.: 163°C
38	NH	CH ₃	OC ₂ H ₅		m.p.: 148°C
39	NH	CH ₃	OC ₂ H ₅		m.p.: 145°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
40	NH	CH ₃	OC ₂ H ₅		m.p.: 171°C
41	NH	CH ₃	OC ₂ H ₅		m.p.: 120°C
42	NH	CH ₃	OC ₂ H ₅		m.p.: 165°C
43	NH	CH ₃	OC ₂ H ₅		m.p.: 129°C
44	NH	CH ₃	OC ₂ H ₅		m.p.: 170°C
45	NH	CH ₃	OC ₂ H ₅		m.p.: 139°C
46	NH	CH ₃	OC ₂ H ₅		m.p.: 184°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
47	NH	CH ₃	OC ₂ H ₅		m.p.: 158°C
48	NH	CH ₃	OC ₂ H ₅		m.p.: 158°C
49	NH	CH ₃	OC ₂ H ₅		m.p.: 153°C
50	NH	CH ₃	OC ₂ H ₅		m.p.: 160°C
51	NH	CH ₃	OC ₂ H ₅		m.p.: 144°C
52	NH	CH ₃	OC ₂ H ₅		m.p.: 152°C
53	NH	CH ₃	OC ₂ H ₅		m.p.: 127°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
54	NH	CH ₃	OC ₂ H ₅		m.p.: 170°C
55	NH	CH ₃	OC ₂ H ₅		m.p.: 132°C
56	NH	CH ₃	OC ₂ H ₅		m.p.: 143°C
57	NH	CH ₃	OC ₂ H ₅		m.p.: 152°C
58	NH	CH ₃	OC ₂ H ₅		m.p.: 170°C
59	NH	CH ₃	OC ₂ H ₅		m.p.: 148°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
60	NH	CH ₃	OC ₂ H ₅		¹ H NMR (CDCl ₃ , δ): 2.52-2.58; 3.15; 3.70 ppm
61	NH	CH ₃	OC ₂ H ₅		m.p.: 94°C
62	NH	CH ₃	OC ₂ H ₅		m.p.: 163°C
63	NH	CH ₃	OC ₂ H ₅		m.p.: 124°C
64	NH	CH ₃	OC ₂ H ₅		m.p.: 138°C
65	NCH ₃	CH ₃	OC ₂ H ₅		m.p.: 149°C
66	NH	C ₂ H ₅	OCH ₃		m.p.: 182°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
67	NH	C ₂ H ₅	OCH ₃		m.p.: 172°C
68	NH	C ₂ H ₅	OCH ₃		m.p.: 167°C
69	NH	C ₂ H ₅	OCH ₃		m.p.: 140°C
70		C ₂ H ₅	OCH ₃		m.p.: 130°C
71	NH	C ₂ H ₅	OCH ₃		m.p.: 167°C
72	NH	C ₂ H ₅	OCH ₃		m.p.: 145°C
73	-O-NH-	C ₂ H ₅	OCH ₃		m.p.: 165°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
74	NH	C ₂ H ₅	OCH ₃		m.p.: 100°C
75	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 1.23-1.25; 3.60-3.70; 4.08 ppm
76	NH	C ₂ H ₅	OCH ₃		m.p.: 122°C
77	NH	C ₂ H ₅	OCH ₃		m.p.: 148°C
78	NCH ₃	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 3.58; 3.92; 4.13 ppm
79	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 2.50-2.60; 3.67; 4.05 ppm

Table 2: (continued)









Ex. No.	Q	R ¹	R ²	R ³	Physical data
80	NH	C ₂ H ₅	OCH ₃		m.p.: 118°C
81	NH	C ₂ H ₅	OCH ₃		m.p.: 145°C
82	NCH ₃	C ₂ H ₅	OCH ₃		m.p.: 130°C
83	NH	CH ₃	CH ₃		m.p.: 137°C
84	NH		N(CH ₃) ₂		m.p.: 119°C
85	NH	C ₂ H ₅	OCH ₃		m.p.: 152°C
86	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 3.63-3.70; 4.12 ppm

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
87	NH		N(CH ₃) ₂		m.p.: 140°C
88	NH	C ₂ H ₅	OCH ₃		m.p.: 169°C
89	NH	C ₂ H ₅	OCH ₃		¹ H NMR. (CDCl ₃ , δ): 3.60-3.67; 3.70-3.77; 4.10 ppm
90	NH	C ₂ H ₅	OCH ₃		m.p.: 120°C
91	NH	C ₂ H ₅	OCH ₃		m.p.: 158°C
92	NH	C ₂ H ₅	OCH ₃		m.p.: 160°C
93	NH	C ₂ H ₅	OCH ₃		m.p.: 127°C

Table 2: (continued)









Ex. No.	Q	R ¹	R ²	R ³	Physical data
94	NH	C ₂ H ₅	OCH ₃		m.p.: 150°C
95	NH	C ₂ H ₅	OCH ₃		m.p.: 125°C
96	NH	C ₂ H ₅	OCH ₃		m.p.: 121°C
97	NH	C ₂ H ₅	OCH ₃		m.p.: 150°C
98	NH	C ₂ H ₅	OCH ₃		m.p.: 151°C
99	NH	C ₂ H ₅	OCH ₃		m.p.: 133°C
100	NH	C ₂ H ₅	OCH ₃		m.p.: 165°C
101	NH	CH ₃	n-C ₃ H ₇		¹ H NMR (CDCl ₃ , δ): 2.50-2.55; 3.25; 4.15- 4.20 ppm

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
102	NH	CH ₃	n-C ₃ H ₇		m.p.: 180°C
103	NH	CH ₃	n-C ₃ H ₇		m.p.: 111°C
104	NH	C ₂ H ₅	OCH ₃		m.p.: 170°C
105	NH	C ₂ H ₅	OCH ₃		m.p.: 135°C
106	NH	CH ₃	OC ₆ H ₅		m.p.: 146°C
107	NH	C ₂ H ₅	OCH ₃		m.p.: 158°C
108	NH	C ₂ H ₅	OCH ₃		m.p.: 148°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
109	NH	C ₂ H ₅	OCH ₃		m.p.: 140°C
110	NH	C ₂ H ₅	OCH ₃		m.p.: 170°C
111	NH	CH ₃	OC ₂ H ₅		m.p.: 162°C
112		C ₂ H ₅	OCH ₃		m.p.: 103°C
113	NH	C ₂ H ₅	OCH ₃		m.p.: 125°C
114	NH	C ₂ H ₅	OCH ₃		m.p.: 150°C

Table 2: (continued)






Ex. No.	Q	R ¹	R ²	R ³	Physical data
115	$\begin{array}{c} \text{---N---} \\ \\ \text{C=O} \\ \\ \text{OC}_2\text{H}_5 \end{array}$	C ₂ H ₅	OCH ₃		m.p.: 135°C
116	$\begin{array}{c} \text{---N---} \\ \\ \text{CH}_2 \\ \\ \text{C=O} \\ \\ \text{OC}_2\text{H}_5 \end{array}$	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 2.5; 3.60- 3.70; 4.10 ppm
117	NH	C ₂ H ₅	OCH ₃		m.p.: 140°C
118	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 3.60-3.67; 4.05; 8.30 ppm
119	NH	C ₂ H ₅	OCH ₃		m.p.: 165°C

Table 2: (continued)




Ex. No.	Q	R ¹	R ²	R ³	Physical data
120	NH	C ₂ H ₅	OCH ₃		m.p.: 161°C
121	NH	C ₂ H ₅	OCH ₃		m.p.: 172°C
122	NH	C ₂ H ₅	OCH ₃		m.p.: 190°C

Table 2: (continued)




Ex. No.	Q	R ¹	R ²	R ³	Physical data
123	NH	C ₂ H ₅	OCH ₃		m.p.: 185°C
124	NH	C ₂ H ₅	OCH ₃		m.p.: 147°C
125	NH	C ₂ H ₅	OCH ₃		m.p.: 160°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
126	NH	C ₂ H ₅	OCH ₃		m.p.: 193°C
127	NH	C ₂ H ₅	OCH ₃		m.p.: 190°C
128	NH	C ₂ H ₅	OCH ₃		m.p.: 154°C
129	NH	C ₂ H ₅	OCH ₃		m.p.: 168°C
130	NH	C ₂ H ₅	OCH ₃		m.p.: 154°C
131	NH	C ₂ H ₅	OCH ₃		m.p.: 135°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
132	NH	C ₂ H ₅	OCH ₃		m.p. : 193°C
133	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 138°C
134	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 138°C
135	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 144°C
136	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 130°C
137	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 137°C
138	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 146°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
139	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 141°C
140	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 128°C
141	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 135°C
142	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 165°C
143	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 138°C
144	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 120°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
145	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 138°C
146	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 150°C
147	NH	CH ₃	OC ₃ H ₇ -i		m.p. : 150°C
148	NH	C ₂ H ₅	OCH ₃		m.p. : 141°C
149	NH	C ₂ H ₅	OCH ₃		m.p. : 150°C
150	NH	C ₂ H ₅	OCH ₃		m.p. : 170°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
151	NH	C ₂ H ₅	OCH ₃		m.p. : 143°C
152	NH	C ₂ H ₅	OCH ₃		m.p. : 118°C
153	NH	C ₂ H ₅	OCH ₃		m.p. : 142°C
154	NH	C ₂ H ₅	OCH ₃		m.p. : 192°C
155	NH	C ₂ H ₅	OCH ₃		m.p. : 80°C
156	NH	C ₂ H ₅	OCH ₃		m.p. : 107°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
157	NH	C ₂ H ₅	OCH ₃		m.p.: 159°C
158	NH	C ₂ H ₅	OCH ₃		m.p.: 129°C
159	NH	C ₂ H ₅	OCH ₃		m.p.: 137°C
160	NH	C ₂ H ₅	OCH ₃		m.p.: 183°C
161	NH	C ₂ H ₅	OCH ₃		m.p.: 158°C
162	NH	C ₂ H ₅	OCH ₃		m.p.: 155°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
163	NH	CH ₃			m.p.: 140°C
164	NH	CH ₃			m.p.: 129°C
165	NCH ₃	OCH ₃	OCH ₃	CH ₃	m.p.: 154°C
166	NCH ₃	CH ₃	n-C ₃ H ₇	-SO ₂ CH ₃	m.p.: 93°C
167	NCH ₃	CH ₃	C ₂ H ₅	-SO ₂ CH ₃	m.p.: 124°C
168	NCH ₃	CH ₃	SCH ₃	-SO ₂ CH ₃	m.p.: 160°C
169	NCH ₃	CH ₃	SC ₂ H ₅	-SO ₂ CH ₃	m.p.: 116°C
170	NCH ₃	OCH ₃	C ₂ H ₅	-SO ₂ CH ₃	m.p.: 103°C
171	NCH ₃		Cl	-SO ₂ CH ₃	m.p.: 121°C
172	O	CH ₃	OC ₂ H ₅		m.p.: 135°C
173	O	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 3.65-3.72; 3.85; 4.15 ppm

Table 2: (continued)











Ex. No.	Q	R ¹	R ²	R ³	Physical data
174	O		SCH ₃		m.p.: 110°C
175	O		SC ₂ H ₅		m.p.: 100°C
176	O	CH ₃	C ₂ H ₅		m.p.: 132°C.
177	O	CH ₃	n-C ₃ H ₇		m.p.: 107°C
178	O		Br		m.p.: 137°C
179	O	CH ₃	SCH ₃		m.p.: 116°C
180	O	CH ₃	SC ₂ H ₅		m.p.: 103°C

Table 2: (continued)












Ex. No.	Q	R ¹	R ²	R ³	Physical data
181	O		C ₂ H ₅		m.p.: 119°C
182	O		n-C ₃ H ₇		m.p.: 94°C
183	O		Cl		m.p.: 121°C
184	O	OCH ₃	C ₂ H ₅		m.p.: 96°C
185	O		Cl		m.p.: 128°C
186	O	CH ₃	C ₂ H ₅		m.p.: 91°C
187	O	CH ₃	SCH ₃		m.p.: 139°C

Table 2: (continued)












Ex. No.	Q	R ¹	R ²	R ³	Physical data
188	O	CH ₃	SC ₂ H ₅		m.p.: 97°C
189	O		Br		m.p.: 120°C
190	O	CH ₃	SCH ₃		m.p.: 129°C
191	O	CH ₃	SC ₂ H ₅		m.p.: 146°C
192	O		C ₂ H ₅		m.p.: 108°C
193	O		n-C ₃ H ₇		m.p.: 118°C
194	O		n-C ₃ H ₇		m.p.: 99°C

Table 2: (continued)









Ex. No.	Q	R ¹	R ²	R ³	Physical data
195	O	OCH ₃	C ₂ H ₅		m.p.: 120°C
196	O	OCH ₃	n-C ₃ H ₇		m.p.: 105°C
197	NH	CH ₃	OCH ₃		m.p.: 132°C
198		C ₂ H ₅	OCH ₃		m.p.: 157°C
199	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 139°C
200	NH	C ₂ H ₅	OCH ₃		m.p.: 140°C
201	NH	C ₂ H ₅	OCH ₃		m.p.: 178°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
202	NH	C ₂ H ₅	OCH ₃		m.p.: 190°C
203	NH	CH ₃	N(CH ₃) ₂		m.p.: 160°C
204	NH	CH ₃	N(CH ₃) ₂		m.p.: 161°C
205	NH	CH ₃	N(CH ₃) ₂		m.p.: 151°C
206	NH	CH ₃	N(CH ₃) ₂		m.p.: 140°C
207	NH	CH ₃	N(CH ₃) ₂		m.p.: 135°C
208	NH	CH ₃	N(CH ₃) ₂		m.p.: 124°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
209	NH	CH ₃	N(CH ₃) ₂		m.p.: 131°C
210	NH	CH ₃	N(CH ₃) ₂		m.p.: 172°C
211	NH	CH ₃	N(CH ₃) ₂		m.p.: 140°C
212	NH	CH ₃	N(CH ₃) ₂		m.p.: 161°C
213	NH	C ₂ H ₅	OCH ₃		m.p.: 130°C
214	NH	C ₂ H ₅	OCH ₃		¹ H NMR (CDCl ₃ , δ): 4.05; 4.62- 4.64; 4.80-4.82 ppm

Table 2: (continued)









Ex. No.	Q	R ¹	R ²	R ³	Physical data
215	NH	C ₂ H ₅	OCH ₃		m.p. : 174°C
216	NH	C ₂ H ₅	OCH ₃		m.p. : 169°C
217	NH	CH ₃	N(CH ₃) ₂		m.p. : 156°C
218	NH	CH ₃			m.p. : 106°C
219	NH	CH ₃			m.p. : 148°C
220	NH	CH ₃	OCH ₃		m.p. : 149°C

Table 2: (continued)











Ex. No.	Q	R ¹	R ²	R ³	Physical data
221	NH	CH ₃	SC ₂ H ₅		m.p.: 112°C
222	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 131°C
223	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 156°C
224	NH	CH ₃			m.p.: 144°C
225	NH	CH ₃			m.p.: 157°C
226		CH ₃	OC ₃ H ₇ -i		m.p.: 136°C
227	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 161°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
228	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 149°C
229	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 138°C
230	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 149°C
231	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 115°C
232	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 121°C
233	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 123°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
234	NH	CH ₃	OC ₃ H ₇ -i		¹ H NMR (CDCl ₃ , δ): 1.2-1.25; 3.15; 3.15- 3.25 ppm
235	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 130°C
236	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 131°C
237		CH ₃	OC ₃ H ₇ -i		m.p.: 121°C
238	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 139°C

Table 2: (continued)










Ex. No.	Q	R ¹	R ²	R ³	Physical data
239	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 145°C
240	NH	CH ₃			m.p.: 140°C
241	NH	CH ₃			m.p.: 146°C
242		CH ₃	OC ₃ H ₇ -i		m.p.: 156°C
243	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 141°C
244	NH	CH ₃	i-C ₃ H ₇		m.p.: 148°C
245	NH	CH ₃	i-C ₃ H ₇		m.p.: 127°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
246	NH	CH ₃	i-C ₃ H ₇		m.p.: 142°C
247	NH	CH ₃	i-C ₃ H ₇		m.p.: 155°C
248	NH	CH ₃	i-C ₃ H ₇		m.p.: 148°C.
249	NH	CH ₃	i-C ₃ H ₇		m.p.: 137°C
250	NH	CH ₃	i-C ₃ H ₇		m.p.: 144°C
251	NH	CH ₃	i-C ₃ H ₇		m.p.: 136°C
252	NH	CH ₃	i-C ₃ H ₇		m.p.: 120°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
253	NH	CH ₃	O-C ₃ H ₇ -i		m.p.: 139°C
254	NH	CH ₃	i-C ₃ H ₇		m.p.: 96°C
255	NH	CH ₃	O-C ₃ H ₇ -i		m.p.: 154°C
256	NH	CH ₃	O-C ₂ H ₅		m.p.: 165°C
257	NH	CH ₃	O-C ₂ H ₅		m.p.: 164°C
258	NH	CH ₃			m.p.: 155°C

Table 2: (continued)









Ex. No.	Q	R ¹	R ²	R ³	Physical data
259	NH	CH ₃	i-C ₃ H ₇		m.p.: 165°C
260	NH	CH ₃			m.p.: 167°C
261	NH	CH ₃	OC ₂ H ₅		m.p.: 130°C.
262	NH	CH ₃			m.p.: 139°C
263	NH	CH ₃	O-C ₃ H ₇ -i		m.p.: 109°C
264	NH	CH ₃	OC ₂ H ₅		m.p.: 157°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
265	NH	CH ₃	C ₃ H ₇ -i		m.p.: 145°C
266	NH	CH ₃	OC ₂ H ₅		m.p.: 135°C
267	NH	CH ₃	OC ₂ H ₅		m.p.: 124°C
268	NH	CH ₃	OC ₂ H ₅		m.p.: 89°C
269	NH	CH ₃	OC ₂ H ₅		m.p.: 164°C
270	NH	CH ₃	OCH ₃		m.p.: 157°C
271		CH ₃	OC ₂ H ₅	-SO ₂ CH ₃	m.p.: 145°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
272	NH	CH ₃	OC ₃ H ₇ -i		m.p.: 164°C
273	NH	CH ₃	OC ₂ H ₅		m.p.: 136°C
274	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 101°C
275	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 79°C
276	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 84°C
277	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 123°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
278	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 104°C
279	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 111°C
280	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 121°C
281	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 112°C
282	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 117°C
283	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 92°C

Table 2: (continued)








Ex. No.	Q	R ¹	R ²	R ³	Physical data
284	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 103°C
285	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 106°C
286	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 105°C
287	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 115°C
288	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 108°C
289	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 126°C
290	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 132°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
291	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 135°C
292	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 141°C
293	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 79°C
294	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 102°C
295	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 115°C
296	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 68°C

Table 2: (continued)







Ex. No.	Q	R ¹	R ²	R ³	Physical data
297	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 59°C
298	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 71°C
299	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 85°C
300	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 91°C
301	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 114°C
302	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 109°C

Table 2: (continued)











Ex. No.	Q	R ¹	R ²	R ³	Physical data
303	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i *)		(Oil)
304	NH	CH ₃			m.p.: 109°C
305	NH	CH ₃			m.p.: 123°C
306	NH	CH ₃			m.p.: 132°C
307	NH	CH ₃			m.p.: 129°C
308	NH	CH ₃	OC ₂ H ₄ - OC ₂ H ₅ *)		m.p.: 103°C

Table 2: (continued)























Ex. No.	Q	R ¹	R ²	R ³	Physical data
309	NH	CH ₃	OC ₂ H ₄ - OCH ₃	*) 	m.p.: 85°C
310	NH	CH ₃	OC ₂ H ₄ - OC ₃ H ₇ -i	*) 	m.p.: 91°C
311	NH	CH ₃			m.p.: 112°C
312	NH	CH ₃	OCH ₃		m.p.: 146°C
313	NH	CH ₃	SC ₂ H ₅		m.p.: 126°C
314	NH	CH ₃	OCH ₃		m.p.: 133°C

Table 2: (continued)

Ex. No.	Q	R ¹	R ²	R ³	Physical data
315	m.p.	CH ₃	OCH ₃		m.p.: 163°C
316	NH	CH ₃	OCH ₃		m.p.: 146°C
317	NH	CH ₃	SC ₂ H ₅		m.p.: 152°C
318	NH	CH ₃	Cl		m.p.: 147°C
319	NH	CH ₃	Br		m.p.: 148°C
320	NH		Br		m.p.: 153°C
321	NH		Cl		m.p.: 155°C


Ex. No.	Q	R ¹	R ²	R ³	Physical data
322	NH	CH ₃	SC ₂ H ₅		m.p.: 137°C
323	NH		SCH ₃		m.p.: 166°C
324	NH		SC ₂ H ₅		m.p.: 159°C
325	NH	-NHCOO- C ₂ H ₅ *)	OCH ₃		
326	NH	-NHCOO- C ₂ H ₅ *)	OCH ₃		

*) $\begin{matrix} \text{OC}_2\text{H}_4- \\ \text{OC}_2\text{H}_5 \end{matrix} = -\text{OCH}_2\text{CH}_2\text{OCH}_2\text{CH}_3;$ $\begin{matrix} \text{OC}_2\text{H}_4- \\ \text{OCH}_3 \end{matrix} = -\text{OCH}_2\text{CH}_2\text{OCH}_3;$

$\begin{matrix} \text{OC}_2\text{H}_4- \\ \text{OC}_3\text{H}_7-i \end{matrix} = -\text{OCH}_2\text{CH}_2\text{OCH}(\text{CH}_3)_2;$

$\begin{matrix} -\text{NHCOO}- \\ \text{C}_2\text{H}_5 \end{matrix} = -\text{NHCOOCH}_2\text{CH}_3 .$

The compound shown in Table 2 as Example 183 may, for example, also be prepared as follows [Process (c)]:




2.0 g (12.5 mmol) of 4-cyclopropyl-5-chloro-2,4-dihydro-3H-1,2,4-triazol-3-one are dissolved in 40 ml of acetonitrile, and 4.3 g (26.9 mmol) of (2-ethoxy-phenoxy)-sulphonyl isocyanate are added, with stirring. The reaction mixture is stirred for 6 hours at 20°C. It is then concentrated in vacuo, the residue is stirred with diethyl ether, and the crystalline product is isolated by filtration with suction.

4.9 g (97 % of theory) of 2-(2-ethoxy-phenyloxy)-sulphonylaminocarbonyl-4-cyclopropyl-5-chloro-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 121°C are obtained.

Starting substances of the formula (II)

Example (II-1)




28.8 g (0.20 mol) of 5-ethoxy-4-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one are introduced into 250 ml of methylene chloride, and the mixture is cooled to -10°C. 28.3 g (0.20 mol) of chlorosulphonyl isocyanate are added to this mixture, and the reaction mixture is stirred for 30 minutes without cooling. The solvent is then carefully removed by distillation under water pump vacuum.

53 g (93 % of theory) of 2-chlorosulphonylaminocarbonyl-5-ethoxy-4-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one are obtained as a crystalline residue of melting point

106°C.


Example (II-2)



22.6 g (0.20 mol) of 4,5-dimethyl-2,4-dihydro-3H-1,2,4-triazol-3-one are introduced into 250 ml of methylene chloride and the mixture is cooled to -10°C. Then, 28.3 g (0.20 mol) of chlorosulphonyl isocyanate are added, and the mixture is stirred for 20 minutes at -5°C to -10°C, a clear solution first being formed and the product then separating out as crystals. This product is isolated by filtration with suction.

45 g (88 % of theory) of 2-chlorosulphonylaminocarbonyl-4,5-dimethyl-2,4-dihydro-3H-1,2,4-triazol-3-one of melting point 150°C (with decomposition) are obtained.

10 Example (II-3)




9.9 g (72 mmol) of chlorosulphonyl isocyanate are slowly added at -10°C to 0°C to 10 g (72 mmol) of 5-dimethylamino-4-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one in 250 ml of methylene chloride. After the reaction mixture has been stirred for thirty minutes, it has come to room temperature (20°C). Then, the solvent is removed by distillation under water pump vacuum.

19.1 g (94 % of theory) of 2-chlorosulphonylaminocarbonyl-5-dimethylamino-4-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one are obtained as an amorphous residue.

Use Examples:

In the use examples, compound (A) below is used as comparison substance:



2-(2-Chloro-phenylsulphonylaminocarbonyl)-4,5-dimethyl-2,4-dihydro-3H-1,2,4-triazol-3-one (disclosed in EP-A 341 489).

Example A

Pre-emergence test

Solvent: 5 parts by weight of acetone

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, the stated amount of emulsifier is added and the concentrate is diluted with water to the desired concentration.

10 Seeds of the test plants are sown in normal soil and, after 24 hours, watered with the preparation of the active compound. It is expedient to keep constant the amount of water per unit area. The concentration of the active compound in the preparation is of no importance, only the amount of active compound applied per unit area being decisive. After three weeks, the degree of damage to the plants is rated in % damage in comparison to the development of the untreated control. The figures denote:

0% = no action (like untreated control)
15 100% = total destruction

A clearly superior activity compared with the prior art is shown, in this test, for example by the compounds of the following preparation examples: 2, 4, 13, 33, 39, 41, 43, 45, 58, 64 and 65.

Table A: Pre-emergence test/greenhouse




Active compound	Application rate (g/ha)	Wheat	Amaranthus	Galinsoga	Matri-caria	Stel-laria
	250	0	0	0	0	0
 (A) (known) (13)	250	70	90	95	95	80
 x N(C2H5)3 (4)	250	0	70	70	90	60

Table A: Pre-emergence test/greenhouse (continued)




Active compound	Application rate (g/ha)	Wheat	Amaranthus	Galinsoga	Matri-caria	Stel-laria
 (33)	250	0	95	95	90	90
 (2)	250	20	95	95	90	80
 (39)	250	0	80	95	80	50

Table A: Pre-emergence test/greenhouse (continued)







Active compound	Application rate (g/ha)	Wheat	Amaranthus	Galinsoga	Matri-caria	Stel-laria
 (41)	250	0	30	95	95	90
 (43)	250	0	90	95	95	90
 (45)	250	0	95	95	95	95

Table A: Pre-emergence test/greenhouse (continued)

Active compound	Application rate (g/ha)	Wheat	Amaran- thus	Galini- soga	Matri- caria	Stel- laria
 (58)	250	0	-	95	95	70
 (64)	250	30	95	95	60	60
 (65)	250	0	100	95	95	70

Example B

Post-emergence test

Solvent: 5 parts by weight of acetone

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, the stated amount of emulsifier is added and the concentrate is diluted with water to the desired concentration.

10 Test plants which have a height of 5 - 15 cm are sprayed with the preparation of the active compound in such a way as to apply the particular amounts of active compound desired per unit area. The concentration of the spray liquor is so chosen that the particular amounts of active compound desired are applied in 2,000 l of water/ha. After three weeks, the degree of damage to the plants is rated in % damage in comparison to the development of the untreated control. The figures denote:

0% = no action (like untreated control)

15 100% = total destruction

A clearly superior activity compared with the prior art is shown, in this test, for example by the compounds of the following preparation examples: 2, 4, 13, 14, 33, 39, 41 and 43.

Table B: Post-emergence test/greenhouse
Active compound




Active compound	Appli- cation rate (g/ha)	Wheat	Abu- tilon	Galin- soga	Ipo- moea	Matri- caria	Sola- num	Xan- thium
 (A) (known)	250	0	0	0	0	0	70	0
 (13)	250	0	95	100	95	100	95	100
 (14)	250	0	40	50	40	30	90	95

Table B: Post-emergence test/greenhouse (continued)







Active compound	Appli- cation rate (g/ha)	Wheat	Abu- tilon	Galini- soga	Ipo- moea	Matri- caria	Sola- num	Xan- thium
 $\times N(C_2H_5)_3$ (4)	250	0	70	90	80	30	95	100
 (33)	250	0	90	100	70	90	95	90
 (2)	250	0	80	100	90	90	95	100


Table B: Post-emergence test/greenhouse (continued)
Active compound

Active compound	Appli- cation rate (g/ha)	Wheat	Abu- tilon	Galini- soga	Ipo- moea	Matri- caria	Sola- num	Xan- thium
 (39)	250	0	60	95	60	80	90	100
 (41)	250	0	100	100	80	100	100	100
 (43)	250	0	95	100	80	100	100	100

30517-159


CLAIMS:

1. A sulphonylaminocarbonyltriazolinone of the general formula (I):



wherein:

Q represents an oxygen atom, a sulphur atom, or one of the following groups:



R^1 represents hydrogen, hydroxyl, amino, or represents C_2 - C_{10} -alkylidene-amino, or represents C_1 - C_6 -alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylcarbonyl or C_1 - C_4 -alkoxy-carbonyl, or represents C_3 - C_6 -alkenyl or C_3 - C_6 -alkinyl, each of which is optionally substituted by fluorine, chlorine and/or bromine, or represents C_3 - C_6 -cycloalkyl which is optionally substituted by fluorine, chlorine, bromine and/or C_1 - C_4 -alkyl, or represents phenyl- C_1 - C_3 -alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl, C_1 - C_4 -alkoxy and/or C_1 - C_4 -alkoxy-carbonyl, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C_1 - C_4 -alkyl, trifluoromethyl, C_1 - C_4 -alkoxy, fluorine- and/or chlorine-substituted C_1 - C_3 -alkoxy, C_1 - C_4 -alkylthio, fluorine- and/or chlorine-substituted C_1 - C_3 -alkylthio, C_1 - C_4 -

30517-159

alkylsulphinyl, C₁-C₄-alkylsulphonyl and/or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkoxy which is optionally substituted by fluorine, chlorine, cyano, phenyl, C₁-C₄-alkoxy or C₁-C₄-alkoxy-carbonyl, or represents C₃-C₆-
 5 alkenyloxy, or represents C₁-C₄-alkylamino, C₃-C₆-cycloalkylamino or di-(C₁-C₄-alkyl)-amino, each of which is optionally substituted by fluorine, cyano, C₁-C₄-alkoxy or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkanoylamino, or represents C₁-C₆-alkoxycarbonylamino;

10 R² represents hydrogen, hydroxyl, mercapto, amino, halogen, or represents C₁-C₆-alkyl or C₂-C₆-alkenyl, optionally substituted by fluorine, chlorine, bromine, cyano, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio or C₁-C₄-alkoxy-carbonyl, or represents C₃-C₆-cycloalkyl which
 15 is optionally substituted by fluorine, chlorine, bromine and/or C₁-C₄-alkyl, or represents cyclohexenyl, or represents phenyl-C₁-C₃-alkyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy and/or C₁-C₄-
 20 alkoxy-carbonyl, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy, fluorine- and/or chlorine-substituted C₁-C₃-alkoxy, C₁-C₄-alkylthio, fluorine- and/or chlorine-substituted C₁-C₃-alkylthio, C₁-C₄-alkyl-
 25 sulphinyl, C₁-C₄-alkyl-sulphonyl and/or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkoxy which is optionally substituted by fluorine, chlorine, cyano, C₁-C₄-alkoxy or C₁-C₄-alkoxy-carbonyl, or represents C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy or C₃-C₆-cycloalkyloxy, or represents phenoxy or
 30 benzyloxy, each of which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy, difluoromethoxy or trifluoromethoxy, or represents C₁-C₆-alkylthio, C₁-C₆-

30517-159

alkylsulphinyl, C₁-C₆-alkylsulphonyl, C₃-C₆-alkenylthio, C₃-C₆-alkinylthio, or C₃-C₆-cycloalkylthio, each of which is optionally substituted by fluorine, chlorine, cyano, C₁-C₄-alkoxy or C₁-C₄-alkoxy-carbonyl, or represents phenylthio or benzylthio, each of which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy, difluoromethoxy or trifluoromethoxy, or represents C₁-C₆-alkylamino or C₃-C₆-alkenylamino, or represents phenylamino or benzylamino, each of which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy, difluoromethoxy or trifluoromethoxy, or represents di-(C₁-C₄-alkyl)-amino, or represents aziridino, pyrrolidino or morpholino, each of which is optionally substituted by C₁-C₄-alkyl;

R³ represents C₁-C₆-alkyl, C₃-C₆-alkenyl or C₃-C₆-alkinyl, each of which is optionally substituted by halogen, cyano, carboxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulphinyl, C₁-C₄-alkylsulphonyl or C₁-C₄-alkoxy-carbonyl, or represents C₃-C₆-cycloalkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl, each of which is optionally substituted by halogen, cyano, carboxyl, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₆-alkylsulphonyl, C₁-C₆-alkylaminosulphonyl or di-(C₁-C₄-alkyl)-aminosulphonyl, each of which is optionally substituted by halogen, or represents phenyl, naphthyl, tetralinyl, phenyl-C₁-C₄-alkyl, naphthyl-C₁-C₄-alkyl or tetralinyl-C₁-C₄-alkyl, each of which is optionally substituted by halogen, cyano, carboxyl, nitro, by C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulphinyl or C₁-C₄-alkylsulphonyl (in each case optionally substituted by halogen, cyano, C₁-C₄-alkoxy or C₁-C₄-alkoxy-carbonyl), by C₁-C₄-alkylsulphonyloxy, C₁-C₄-alkylaminosulphonyloxy,

30517-159

di-(C₁-C₄-alkyl)aminosulphonyloxy,
 C₁-C₄-halogenoalkylsulphonyloxy, di-(C₁-C₄-alkyl)aminocarbonyl
 or C₁-C₄-alkylcarbonyloxy, by C₃-C₆-cycloalkyl or
 C₃-C₆-cycloalkyl-C₁-C₄-alkyl (in each case optionally
 5 substituted by halogen or C₁-C₄-alkyl), by phenyl, phenoxy,
 phenylthio, phenylsulphinyl, phenylsulphonyl,
 phenylsulphonyloxy, phenylamino, phenylcarbonyl or phenyl-
 C₁-C₄-alkyl (in each case optionally substituted by halogen,
 cyano, nitro, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy,
 10 C₁-C₄-halogenoalkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkylthio,
 C₁-C₄-alkoxycarbonyl or C₁-C₄-alkoxy-amino), by C₁-C₄-alkyl-
 carbonyl, C₃-C₆-cycloalkylcarbonyl or C₁-C₄-alkoxy-carbonyl
 (in each case optionally substituted by halogen,
 C₃-C₆-cycloalkyl or C₁-C₄-alkoxy); and

15 R⁴ represents hydrogen, hydroxyl, amino, cyano, or
 represents C₁-C₆-alkyl, C₃-C₆-alkenyl or C₃-C₆-alkinyl, each
 of which is optionally substituted by halogen, cyano,
 carboxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulphinyl,
 C₁-C₄-alkylsulphonyl, C₁-C₄-alkylamino, di-(C₁-C₄-alkyl)-amino
 20 or C₁-C₄-alkoxy-carbonyl, or represents C₁-C₄-alkoxy-
 carbonyl, or represents C₁-C₆-alkoxy, C₃-C₆-alkenyloxy,
 benzyloxy, phenoxy or di-(C₁-C₄-alkoxy)-(thio)-phosphoryl,
 or represents the group Q¹-R⁵, wherein:

Q¹ represents -CO- or -SO₂-, and


25 R⁵ represents C₁-C₆-alkyl which is optionally
 substituted by halogen or C₁-C₄-alkoxy, or represents
 C₃-C₆-cycloalkyl which is optionally substituted by halogen
 or C₁-C₄-alkyl, or represents phenyl which is optionally
 substituted by halogen, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl or
 30 C₁-C₄-alkoxy;

and a salt thereof.

30517-159

2. A compound of the general formula (I) according to claim 1, wherein:

Q represents oxygen or the group:



5

R^1 represents hydrogen, hydroxyl, amino, or represents C_3 - C_8 -alkylidene-amino, or represents methyl, ethyl, n- or i-propyl, n-, i-, s- or t-butyl, dichloromethyl, fluoroethyl, chloroethyl, difluoroethyl, dichloroethyl, trifluoroethyl, trichloroethyl, chlorodifluoroethyl, tetrafluoroethyl, cyanomethyl, cyanoethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, or represents allyl, chloroallyl, dichloroallyl, propargyl, or represents chloropropyl, benzyl or phenyl, or represents methoxy, ethoxy, n- or i-propoxy, or represents allyloxy, n- or i- or s-butoxy, or represents methylamino, ethylamino, n- or i-propylamino, or represents cyclopropylamino, dimethylamino, diethylamino, acetylamino, methoxycarbonylamino or ethoxycarbonylamino;

R^2 represents hydrogen, hydroxyl, mercapto, amino, fluorine, chlorine, bromine, or represents methyl, ethyl, n- or i-propyl, n-, i-, s- or t-butyl, difluoromethyl, trifluoromethyl, dichloromethyl, trichloromethyl, cyanomethyl, cyanoethyl, cyclopropylmethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl, ethylthiomethyl, methylthioethyl or ethylthioethyl, or represents cyclopropyl, difluorocyclopropyl or dichlorocyclopropyl, or represents phenyl or benzyl, or represents methoxy, ethoxy, n- or i-propoxy, or represents methoxymethoxy, ethoxymethoxy, methoxyethoxy or ethoxyethoxy, or represents phenoxy or benzyloxy, or

30517-159

represents methylthio, ethylthio, n- or i-propylthio, allylthio, propargylthio, cyclopropylmethylthio, methylsulphinyl, ethylsulphinyl, methylsulphonyl, ethylsulphonyl, phenylthio or benzylthio, or represents
5 methylamino, ethylamino, n- or i-propylamino, phenylamino or benzylamino, or represents dimethylamino, diethylamino, dipropylamino, methylethylamino, methylpropylamino, or represents aziridino, or represents pyrrolidino, piperidino or morpholino, each of which is optionally substituted by
10 methyl or ethyl;

R^3 represents methyl, ethyl, n- or i-propyl or n-, i-, s- or t-butyl, each of which is optionally substituted by fluorine, chlorine, cyano, methoxy, ethoxy, methoxycarbonyl or ethoxycarbonyl, or represents
15 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl, cyclopentylethyl or cyclohexylethyl, each of which is optionally substituted by fluorine, chlorine, cyano, methyl,
20 ethyl, trifluoromethyl, methoxycarbonyl or ethoxycarbonyl, or represents methylsulphonyl, ethylsulphonyl, propylsulphonyl or butylsulphonyl, optionally substituted by fluorine or chlorine, or represents phenyl which is optionally substituted by fluorine, chlorine, bromine,
25 cyano, carboxyl, nitro, by methyl, ethyl, methoxy, ethoxy, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, methylsulphonyl or ethylsulphonyl (in each case optionally substituted by fluorine, chlorine, cyano, methoxy, ethoxy, methoxycarbonyl or ethoxycarbonyl), by methylsulphonyloxy,
30 ethylsulphonyloxy, methylaminosulphonyloxy, ethylaminosulphonyloxy, dimethylaminosulphonyloxy, diethylaminosulphonyloxy, trifluoromethylsulphonyloxy, dimethylaminocarbonyl or diethylaminocarbonyl, by cyclohexyl

30517-159

or cyclohexylmethyl (in each case optionally substituted by chlorine, methyl or ethyl), by phenyl, phenoxy, phenylthio, phenylsulphonyl, phenylsulphonyloxy, phenylamino, phenylcarbonyl, phenylmethyl or phenylethyl (in each case
 5 optionally substituted by fluorine, chlorine, cyano, nitro, methyl, ethyl, trifluoromethyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, methylthio, ethylthio, trifluoromethylthio, methoxycarbonyl or ethoxycarbonyl), by acetyl, cyclopropylcarbonyl, methoxycarbonyl or
 10 ethoxycarbonyl (optionally substituted by fluorine, chlorine, cyclopentyl, cyclohexyl, methoxy or ethoxy), or represents naphthyl or tetralinyl, each of which is optionally substituted by fluorine, chlorine, cyano, nitro, methyl, ethyl, methoxy or ethoxy, or represents
 15 naphthylmethyl, naphthylethyl, tetralinylmethyl or tetralinylethyl, each of which is optionally substituted by fluorine, chlorine, methyl, ethyl, methoxy or ethoxy;

R^4 represents hydrogen, cyano, cyanomethyl, cyanoethyl, difluoromethyl, methoxymethyl, ethoxymethyl,
 20 methoxyethyl, ethoxyethyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylethyl, allyl, propargyl, methoxycarbonyl, ethoxycarbonyl, methoxy, ethoxy, allyloxy or benzyloxy, or represents the group Q^1-R^5 , wherein:

25 Q^1 represents $-CO-$ or $-SO_2-$, and

R^5 represents methyl, ethyl, or propyl, each of which is optionally substituted by fluorine, chlorine, methoxy or ethoxy, or represents cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is optionally
 30 substituted by fluorine, chlorine, methyl or ethyl, or represents phenyl which is optionally substituted by


30517-159

fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, methoxy or ethoxy;

and a salt thereof.

3. The compound:


5



and a salt thereof.

4. The compound:


10



and a salt thereof.

5. The compound:


15



and a salt thereof.

6. The compound:


20



30517-159

and a salt thereof.


7. The compound:



5

and a salt thereof.


8. The compound:



10

and a salt thereof.


9. The compound:



15

and a salt thereof.

10. The compound:



20


and a salt thereof.

30517-159

11. The sodium, potassium, magnesium, calcium, ammonium, C₁-C₄-alkyl-ammonium, di-(C₁-C₄-alkyl)-ammonium, tri-(C₁-C₄-alkyl)-ammonium, C₅- or C₆-cycloalkyl-ammonium or di-(C₁-C₂-alkyl)-benzyl-ammonium salt of a compound
 5 according to any one of claims 1 to 10.

12. A process for the preparation of a sulphonylaminocarbonyltriazolinone of the general formula (I) according to claim 1 or 2, wherein:

(a) a chlorosulphonylaminocarbonyltriazolinone of
 10 the general formula (II):




wherein R¹ and R² are as defined in claim 1 or 2,
 15 is reacted with a nucleophilic compound of the general formula (III):



wherein Q and R³ are as defined in claim 1 or 2, if appropriate, in the presence of an acid acceptor and, if
 20 appropriate, in the presence of diluent; or

(b) a triazolinone of the general formula (IV):



25 wherein R¹ and R² are as defined in claim 1 or 2, is reacted with chlorosulphonyl isocyanate, if appropriate,

30517-159

in the presence of a diluent, and the resulting chlorosulphonylaminocarbonyltriazolinone of the formula (II) is reacted, without intermediate isolation, with a nucleophilic compound of the general formula (III), if
5 appropriate, in the presence of an acid acceptor and, if appropriate, in the presence of a diluent; and

if appropriate, the compound obtained by process (a) or (b) is converted to a salt.

13. A herbicidal composition, comprising at least one
10 compound according to any one of claims 1 to 11, and (i) an extender, (ii) a surface-active agent or (iii) a combination of (i) and (ii).

14. Use of a compound according to any one of claims 1 to 11, for combating undesired plant growth.

15 15. A method of combating weeds, comprising allowing to act on the weeds or their environment a compound according to any one of claims 1 to 11.

16. A process for the preparation of a herbicidal composition, comprising mixing a compound according to any
20 one of claims 1 to 11, with (i) an extender, (ii) a surface-active agent or (iii) a combination of (i) and (ii).

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS

