



(86) Date de dépôt PCT/PCT Filing Date: 2015/05/11
(87) Date publication PCT/PCT Publication Date: 2015/11/19
(45) Date de délivrance/Issue Date: 2019/10/22
(85) Entrée phase nationale/National Entry: 2016/11/09
(86) N° demande PCT/PCT Application No.: EP 2015/060343
(87) N° publication PCT/PCT Publication No.: 2015/173181
(30) Priorité/Priority: 2014/05/15 (US61/993,839)

(51) Cl.Int./Int.Cl. *C07D 519/00* (2006.01),
A61K 31/519 (2006.01), *A61P 21/00* (2006.01),
A61P 25/28 (2006.01)

(72) Inventeurs/Inventors:
RATNI, HASANE, FR;
GREEN, LUKE, CH;
NARYSHKIN, NIKOLAI A., US;
WEETALL, MARLA L., US

(73) Propriétaires/Owners:
F. HOFFMANN-LA ROCHE AG, CH;
PTC THERAPEUTICS INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSES POUR LE TRAITEMENT D'UNE AMYOTROPHIE SPINALE
(54) Title: COMPOUNDS FOR TREATING SPINAL MUSCULAR ATROPHY

(57) **Abrégé/Abstract:**

The present invention provides compounds of formula (I) wherein A, R¹, R² and R³ are as described herein, as well as pharmaceutically acceptable salts thereof. Further the present invention is concerned with the manufacture of the compounds of formula (I), pharmaceutical compositions comprising them and their use as medicaments.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)
(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 November 2015 (19.11.2015)

(10) International Publication Number
WO 2015/173181 A1

- (51) International Patent Classification:
C07D 487/04 (2006.01) *A61K 31/5025* (2006.01)
C07D 519/00 (2006.01) *A61P 21/00* (2006.01)

(21) International Application Number:
PCT/EP2015/060343

(22) International Filing Date:
11 May 2015 (11.05.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/993,839 15 May 2014 (15.05.2014) US

(71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(71) Applicant (for US only): **HOFFMANN-LA ROCHE INC.** [US/US]; Great Notch, 150 Clove Road, 8th Floor, Little Falls, New Jersey 07424 (US).

(71) Applicant: **PTC THERAPEUTICS INC.** [—/US]; 100 Corporate Court, South Plainfield, New Jersey 07080-2449 (US).


(72) Inventors: **RATNI, Hasane**; 4, Louis Pasteur, F-68440 Habsheim (FR). **GREEN, Luke**; Oetlingerstrasse 52, CH-4057 Basel (CH). **NARYSHKIN, Nikolai A.**; 101 Wind-song Circle, East Brunswick, New Jersey 08816 (US). **WEETALL, Marla L.**; 70 Burnham Parkway, Morris-town, New Jersey 07960 (US).
- (74) Agent: **HAEFELE RACIN, Thomas**; Grenzacherstrasse 124, 4070 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— of inventorship (Rule 4.17(iv))

Published:
— with international search report (Art. 21(3))



WO 2015/173181 A1

(54) Title: COMPOUNDS FOR TREATING SPINAL MUSCULAR ATROPHY


(57) Abstract: The present invention provides compounds of formula (I) wherein A, R¹, R² and R³ are as described herein, as well as pharmaceutically acceptable salts thereof. Further the present invention is concerned with the manufacture of the compounds of formula (I), pharmaceutical compositions comprising them and their use as medicaments.

Compounds for treating spinal muscular atrophy

Introduction

The present invention provides compounds which are SMN2 gene splicing modulators, their manufacture, pharmaceutical compositions comprising them and their use as medicaments for the treatment of spinal muscular atrophy (SMA).

5 In particular, the present invention relates to compounds of formula (I)



wherein A, R¹, R² and R³ are as described herein, and pharmaceutically acceptable salts thereof.

Background

10 Spinal muscular atrophy (SMA), in its broadest sense, describes a collection of inherited and acquired central nervous system (CNS) diseases characterized by progressive motor neuron loss in the spinal cord and brainstem causing muscle weakness and muscle atrophy. The most common form of SMA is caused by mutations in the Survival Motor Neuron (SMN) gene and manifests over a wide range of severity affecting infants through adults (*Crawford and Pardo,*
 15 *Neurobiol. Dis.*, 1996, 3:97).

Infantile SMA is the most severe form of this neurodegenerative disorder. Symptoms include muscle weakness, poor muscle tone, weak cry, limpness or a tendency to flop, difficulty sucking or swallowing, accumulation of secretions in the lungs or throat, feeding difficulties, and increased susceptibility to respiratory tract infections. The legs tend to be weaker than the arms
 20 and developmental milestones, such as lifting the head or sitting up, cannot be reached. In general, the earlier the symptoms appear, the shorter the lifespan. As the motor neuron cells deteriorate, symptoms appear shortly afterward. The severe forms of the disease are fatal and all

forms have no known cure. The course of SMA is directly related to the rate of motor neuron cell deterioration and the resulting severity of weakness. Infants with a severe form of SMA frequently succumb to respiratory disease due to weakness in the muscles that support breathing. Children with milder forms of SMA live much longer, although they may need extensive
 5 medical support, especially those at the more severe end of the spectrum. The clinical spectrum of SMA disorders has been divided into the following five groups.

- 10 (a) Type 0 SMA (In Utero SMA) is the most severe form of the disease and begins before birth. Usually, the first symptom of Type 0 SMA is reduced movement of the fetus that can first be observed between 30 and 36 weeks of pregnancy. After birth, these newborns have little movement and have difficulties with swallowing and breathing.
- 15 (b) Type 1 SMA (Infantile SMA or Werdnig-Hoffmann disease) presents symptoms between 0 and 6 months. form of SMA is also very severe. Patients never achieve the ability to sit, and death usually occurs within the first 2 years without ventilatory support.
- (c) Type 2 SMA (Intermediate SMA) has an age of onset at 7-18 months. Patients achieve the ability to sit unsupported, but never stand or walk unaided. Prognosis in this group is largely dependent on the degree of respiratory involvement.
- 20 (d) Type 3 SMA (Juvenile SMA or Kugelberg-Welander disease) is generally diagnosed after 18 months. Type 3 SMA individuals are able to walk independently at some point during their disease course but often become wheelchair-bound during youth or adulthood.
- 25 (e) Type 4 SMA (Adult onset SMA). Weakness usually begins in late adolescence in the tongue, hands, or feet, then progresses to other areas of the body. The course of adult SMA is much slower and has little or no impact on life expectancy.

The SMN gene has been mapped by linkage analysis to a complex region in chromosome 5q. In humans, this region contains an approximately 500 thousand base pairs (kb) inverted duplication resulting in two nearly identical copies of the SMN gene. SMA is caused by an inactivating mutation or deletion of the telomeric copy of the gene (SMN1) in both
 30 chromosomes, resulting in the loss of SMN1 gene function. However, all patients retain the centromeric copy of the gene (SMN2), and the copy number of the SMN2 gene in SMA patients generally correlates inversely with the disease severity; i.e., patients with less severe SMA have more copies of SMN2. Nevertheless, SMN2 is unable to compensate completely for the loss of SMN1 function due to alternative splicing of exon 7 caused by a translationally silent C to T
 35 mutation in exon 7. As a result, the majority of transcripts produced from SMN2 lack exon 7 ($\Delta 7$ SMN2), and encode a truncated SMN protein that has an impaired function and is rapidly degraded.

The SMN protein is thought to play a role in RNA processing and metabolism, having a well characterized function of mediating the assembly of a specific class of RNA-protein complexes termed snRNPs. SMN may have other functions in motor neurons, however its role in preventing the selective degeneration of motor neurons is not well established.

5 In most cases, SMA is diagnosed based on clinical symptoms and by the presence of at least one copy of the SMN1 gene test. However, in approximately 5% of cases SMA is caused by mutation in genes other than the inactivation of SMN 1, some known and others not yet defined. In some cases, when the SMN 1 gene test is not feasible or does not show any abnormality, other tests such as an electromyography (EMG) or muscle biopsy may be indicated.

10 Medical care for SMA patients at present is limited to supportive therapy including respiratory, nutritional and rehabilitation care; there is no drug known to address the underlying cause of the disease. Current treatment for SMA consists of prevention and management of the secondary effects of chronic motor unit loss. The major management issue in Type 1 SMA is the prevention and early treatment of pulmonary problems, which are the cause of death in the
15 majority of the cases. While some infants afflicted with SMA grow to be adults, those with Type 1 SMA have a life expectancy of less than two years.

Several mouse models of SMA have been developed. In particular, the SMN delta exon 7 ($\Delta 7$ SMN) model (*Le et al., Hum. Mol. Genet., 2005, 14:845*) carries both the SMN2 gene and several copies of the $\Delta 7$ SMN2 cDNA and recapitulates many of the phenotypic features of Type
20 1 SMA. The $\Delta 7$ SMN model can be used for both SMN2 expression studies as well as the evaluation of motor function and survival. The C/C-allele mouse model (*Jackson Laboratory strain #008714, The Jackson Laboratory, Bar Harbor, ME*) provides a less severe SMA disease model, with mice having reduced levels of both SMN2 full length (FL SMN2) mRNA and SMN protein. The C/C-allele mouse phenotype has the SMN2 gene and a hybrid mSMN1-SMN2 gene
25 that undergoes alternative splicing, but does not have overt muscle weakness. The C/C-allele mouse model is used for SMN2 expression studies.

As a result of improved understanding of the genetic basis and pathophysiology of SMA, several strategies for treatment have been explored, but none have yet demonstrated success in the clinic.

30 Gene replacement of SMN1, using viral delivery vectors, and cell replacement, using differentiated SMN1^{+/+} stem cells, have demonstrated efficacy in animal models of SMA. More research is needed to determine the safety and immune response and to address the requirement for the initiation of treatment at the neonatal stage before these approaches can be applied to humans.

Correction of alternative splicing of SMN2 in cultured cells has also been achieved using synthetic nucleic acids as therapeutic agents: (i) antisense oligonucleotides that target sequence elements in SMN2 pre-mRNA and shift the outcome of the splicing reaction toward the generation of full length SMN2 mRNA (*Passini et al., Sci. Transl. Med., 2011, 3:72ra18*; and, 5 *Hua et al., Nature, 2011, 478:123*) and (ii) trans-splicing RNA molecules that provide a fully functional RNA sequence that replace the mutant fragment during splicing and generate a full length SMN1 mRNA (*Coady and Lorson, J Neurosci., 2010, 30:126*).

Other approaches under exploration include searching for drugs that increase SMN levels, enhance residual SMN function, or compensate for its loss. Aminoglycosides have been shown 10 to enhance expression of a stabilized SMN protein produced from $\Delta 7$ SMN2 mRNA by promoting the translational read-through of the aberrant stop codon, but have poor central nervous system penetration and are toxic after repeat dosing. Chemotherapeutic agents, such as aclarubicin, have been shown to increase SMN protein in cell culture; however, the toxicity profile of these drugs prohibits long-term use in SMA patients. Some drugs under clinical 15 investigation for the treatment of SMA include transcription activators such as histone deacetylase ("HDAC") inhibitors (e.g., butyrates, valproic acid, and hydroxyurea), and mRNA stabilizers (mRNA decapping inhibitor RG3039 from Repligen), the goal being to increase the amount of total RNA transcribed from the SMN2 gene. However, the use of the HDAC inhibitors or mRNA stabilizers does not address the underlying cause of SMA and may result in 20 a global increase in transcription and gene expression with potential safety problems in humans.


In an alternative approach, neuroprotective agents such as Olesoxime have been chosen for investigation. Such strategies are not aimed at SMN for the treatment of SMA, but instead are being explored to protect the SMN-deficient motor neurons from neurodegeneration.

A system designed for identifying compounds that increase the inclusion of exon 7 of 25 SMN into RNA transcribed from the SMN2 gene and certain benzoxazole and benzoisoxazole compounds identified thereby have been described in *International Patent Application WO2009/151546A1*. A system designed for identifying compounds that cause ribosomal frameshifting to produce a stabilized SMN protein from $\Delta 7$ SMN2 mRNA and certain isoindolinone compounds identified thereby have been described in *International Patent* 30 *Applications WO2010/019236A1 and WO2013/119916A2*.

Despite the progress made in understanding the genetic basis and pathophysiology of SMA, there remains a need to identify compounds that alter the course of spinal muscular atrophy, one of the most devastating childhood neurological diseases.


Summary

In one aspect, there is provided a compound of formula (I)



wherein R¹ is hydrogen or C₁₋₇-alkyl; R² is hydrogen, cyano, C₁₋₇-alkyl, C₁₋₇-haloalkyl or C₃₋₈-cycloalkyl; R³ is hydrogen, C₁₋₇-alkyl, or C₃₋₈-cycloalkyl; A is N-heterocycloalkyl or NR¹²R¹³, wherein N-heterocycloalkyl comprises 1 or 2 nitrogen ring atoms and is optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴; R¹² is heterocycloalkyl comprising 1 nitrogen ring atom, wherein heterocycloalkyl is optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴; R¹³ is hydrogen, C₁₋₇-alkyl or C₃₋₈-cycloalkyl; two R¹⁴ together form C₁₋₇-alkylene, or R¹⁴ is independently selected from the group consisting of hydrogen, C₁₋₇-alkyl, amino, amino-C₁₋₇-alkyl, C₃₋₈-cycloalkyl and heterocycloalkyl; with the proviso that if A is N-heterocycloalkyl comprising only 1 nitrogen ring atom, then at least one R¹⁴ substituent is amino or amino-C₁₋₇-alkyl; or a pharmaceutically acceptable salt thereof.


In another aspect, there is provided a compound which is



7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.


5a

In another aspect, there is provided a compound which is



7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one.

In another aspect, there is provided a compound of formula (VI)



wherein R^1 , R^2 and R^3 are as defined herein; Y is halogen or trifluoromethanesulfonate; or a salt thereof.

In another aspect, there is provided a process for the preparation of a compound of the invention, comprising an aromatic nucleophilic substitution reaction between a compound of formula (VI) as defined herein with a compound of formula M-A by heating in a solvent, wherein A, R^1 , R^2 and R^3 are as defined herein, Y is as defined herein, M is hydrogen, sodium or potassium, and wherein M is linked to A via a nitrogen atom of A.

In another aspect, there is provided a compound of the invention, obtained by a process of the invention.

In another aspect, there is provided a pharmaceutical composition comprising a compound as defined herein or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.

In another aspect, there is provided a compound as defined herein or a pharmaceutically acceptable salt thereof for use as therapeutically active substance.

In another aspect, there is provided a compound as defined herein or a pharmaceutically acceptable salt thereof for use in the treatment or prevention of spinal muscular atrophy (SMA).

In another aspect, there is provided a use of a compound as defined herein or a pharmaceutically acceptable salt thereof for the treatment or prevention of spinal muscular atrophy (SMA).

In another aspect, there is provided a use of a compound as defined herein or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment or prevention of spinal muscular atrophy (SMA).

Detailed description of the invention

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, suitable methods and materials are described below.

The nomenclature used in this Application is based on IUPAC systematic nomenclature, unless indicated otherwise.

Any open valency appearing on a carbon, oxygen, sulfur or nitrogen atom in the structures herein indicates the presence of a hydrogen, unless indicated otherwise.

The definitions described herein apply irrespective of whether the terms in question appear alone or in combination. It is contemplated that the definitions described herein can be appended to form chemically-relevant combinations, such as e.g. “heterocycloalkylaryl”, “haloalkylheteroaryl”, “arylalkylheterocycloalkyl”, or “alkoxyalkyl”. The last member of the combination is the radical which is binding to the rest of the molecule. The other members of the combination are attached to the binding radical in reversed order in respect of the literal sequence, e.g. the combination amino-C₁₋₇-alkyl refers to a C₁₋₇-alkyl which is substituted by amino, or e.g. the combination arylalkylheterocycloalkyl refers to a heterocycloalkyl-radical which is substituted by an alkyl which is substituted by an aryl.

The term “moiety” refers to an atom or group of chemically bonded atoms that is attached to another atom or molecule by one or more chemical bonds thereby forming part of a molecule. For example, the variables A, R¹, R² and R³ of formula (I) refer to moieties that are attached to the core structure of formula (I) by a covalent bond.

When indicating the number of substituents, the term “one or more” refers to the range from one substituent to the highest possible number of substitution, i.e. replacement of one hydrogen up to replacement of all hydrogens by substituents.

The term “optional” or “optionally” denotes that a subsequently described event or circumstance can but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

The term “substituent” denotes an atom or a group of atoms replacing a hydrogen atom on the parent molecule.

The term “substituted” denotes that a specified group bears one or more substituents. Where any group can carry multiple substituents and a variety of possible substituents is provided, the substituents are independently selected and need not to be the same. The term “unsubstituted” means that the specified group bears no substituents. The term “optionally substituted” means that the specified group is unsubstituted or substituted by one or more substituents, independently chosen from the group of possible substituents. When indicating the number of substituents, the term “one or more” means from one substituent to the highest possible number of substitution, i.e. replacement of one hydrogen up to replacement of all hydrogens by substituents.

The terms “compound(s) of this invention” and “compound(s) of the present invention” refer to compounds as disclosed herein and stereoisomers, tautomers, solvates, and salts (e.g., pharmaceutically acceptable salts) thereof.

When the compounds of the invention are solids, it is understood by those skilled in the art that these compounds, and their solvates and salts, may exist in different solid forms, particularly different crystal forms, all of which are intended to be within the scope of the present invention and specified formulae.

The term “pharmaceutically acceptable salts” denotes salts which are not biologically or otherwise undesirable. Pharmaceutically acceptable salts include both acid and base addition salts.

The term “pharmaceutically acceptable acid addition salt” denotes those pharmaceutically acceptable salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, carbonic acid, phosphoric acid, and organic acids selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, gluconic acid, lactic acid, pyruvic acid, oxalic acid, malic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, aspartic acid, ascorbic acid, glutamic acid, anthranilic acid, benzoic acid, cinnamic acid, mandelic acid, embonic acid, phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, and salicylic acid.

The term “pharmaceutically acceptable base addition salt” denotes those pharmaceutically acceptable salts formed with an organic or inorganic base. Examples of acceptable inorganic bases include sodium, potassium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, and aluminum salts. Salts derived from pharmaceutically acceptable organic nontoxic bases includes salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins,

such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, and
5 polyamine resins.

Stereochemical definitions and conventions used herein generally follow S. P. Parker, Ed., McGraw-Hill Dictionary of Chemical Terms (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., "Stereochemistry of Organic Compounds", John Wiley & Sons, Inc., New York, 1994. In describing an optically active compound, the prefixes D and L, or R and S,
10 are used to denote the absolute configuration of the molecule about its chiral center(s). The substituents attached to the chiral center under consideration are ranked in accordance with the Sequence Rule of Cahn, Ingold and Prelog. (Cahn et al. Angew. Chem. Inter. Edit. 1966, 5, 385; errata 511). The prefixes D and L or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with (-) or L designating that the compound is
15 levorotatory. A compound prefixed with (+) or D is dextrorotatory.

The term "chiral center" denotes a carbon atom bonded to four nonidentical substituents. The term "chiral" denotes the ability of non-superimposability with the mirror image, while the term "achiral" refers to embodiments which are superimposable with their mirror image. Chiral molecules are optically active, i.e., they have the ability to rotate the plane of plane-polarized
20 light.

Compounds of the present invention can have one or more chiral centers and can exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates. Whenever a chiral center is present in a chemical
25 structure, it is intended that all stereoisomers associated with that chiral center are encompassed by the present invention.

The terms "halo", "halogen", and "halide" are used interchangeably herein and denote fluoro, chloro, bromo, or iodo. One particular example of halogen is fluoro.

The term "alkyl" denotes a monovalent linear or branched saturated hydrocarbon group of
30 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, or tert-butyl. Particular examples for alkyl are methyl and ethyl.

The term "haloalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of
35 the alkyl group has been replaced by same or different halogen atoms, particularly fluoro atoms.

Examples of haloalkyl include monofluoro-, difluoro- or trifluoro-methyl, -ethyl or -propyl, for example 3,3,3-trifluoropropyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, fluoromethyl, or trifluoromethyl and the like. The term “perhaloalkyl” denotes an alkyl group where all hydrogen atoms of the alkyl group have been replaced by the same or different halogen atoms.

5 The term “bicyclic ring system” denotes two rings which are fused to each other via a common single or double bond (annelated bicyclic ring system), via a sequence of three or more common atoms (bridged bicyclic ring system) or via a common single atom (spiro bicyclic ring system). Bicyclic ring systems can be saturated, partially unsaturated, unsaturated or aromatic. Bicyclic ring systems can comprise heteroatoms selected from N, O and S.

10 The term “cycloalkyl” denotes a saturated monocyclic or bicyclic hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means consisting of two saturated carbocycles having one or more carbon atoms in common. Particular cycloalkyl groups are monocyclic. Examples for monocyclic cycloalkyl are cyclopropyl, cyclobutanyl, cyclopentyl, 15 cyclohexyl or cycloheptyl. Examples for bicyclic cycloalkyl are bicyclo[2.2.1]heptanyl, or bicyclo[2.2.2]octanyl. One particular example of cycloalkyl is cyclopropyl.

 The term “heterocycloalkyl” denotes a saturated or partly unsaturated mono-, bi- or tricyclic ring system of 3 to 9 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. In particular embodiments, heterocycloalkyl is 20 a monovalent saturated monocyclic ring system of 4 to 7 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples for monocyclic saturated heterocycloalkyl are aziridinyl, oxiranyl, azetidiny, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydro-thienyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, 25 thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, or oxazepanyl. Examples for bicyclic saturated heterocycloalkyl are 8-aza-bicyclo[3.2.1]octyl, quinuclidinyl, 8-oxa-3-aza-bicyclo[3.2.1]octyl, 9-aza-bicyclo[3.3.1]nonyl, 3-oxa-9-aza-bicyclo[3.3.1]nonyl, or 3-thia-9-aza-bicyclo[3.3.1]nonyl. Examples of a partly unsaturated heterocycloalkyl are dihydrofuryl, imidazolinyl, dihydro-oxazolyl, tetrahydro-pyridinyl, or 30 dihydropyranyl. Particular examples of heterocycloalkyl are 1,4-diazepanyl, hexahydropyrrolo[1,2-a]pyrazinyl, piperidinyl, piperazinyl and pyrrolidinyl. More particular examples of heterocycloalkyl are hexahydropyrrolo[1,2-a]pyrazinyl and piperazinyl.

 The term “N-heterocycloalkyl” denotes a heterocycloalkyl radical containing at least one nitrogen ring atom and where the point of attachment of the heterocycloalkyl radical to the rest 35 of the molecule is through a nitrogen ring atom. Particular examples of N-heterocycloalkyl are

1,4-diazepanyl, hexahydropyrrolo[1,2-a]pyrazinyl, piperidinyl, piperazinyl and pyrrolidinyl. More particular examples of N-heterocycloalkyl are hexahydropyrrolo[1,2-a]pyrazinyl and piperazinyl.

The term “basicity” in reference to a compound is expressed herein by the negative decadic
5 logarithm of the acidity constant of the conjugate acid ($pK_a = -\log K_a$). The larger the pK_a of the conjugate acid, the stronger the base ($pK_a + pK_b = 14$). In this application, an atom or functional group is denoted “basic” if it is suitable to accept a proton and if the calculated pK_a of its conjugate acid is at least 7, more particularly if the calculated pK_a of its conjugate acid is at least 7.8, most particularly if the calculated pK_a of its conjugate acid is at least 8. pK_a values were
10 calculated in-silico as described in F. Milletti et al., *J. Chem. Inf. Model* (2007) 47:2172-2181.

The term “alkylene” denotes a linear saturated divalent hydrocarbon group of 1 to 7 carbon atoms or a divalent branched saturated hydrocarbon group of 3 to 7 carbon atoms. Examples of alkylene groups include methylene, ethylene, propylene, 2-methylpropylene, butylene, 2-ethylbutylene, pentylene, hexylene. Particular examples for alkylene are ethylene, propylene, and
15 butylene.

The term “amino” denotes a group of the formula $-NR'R''$ wherein R' and R'' are independently hydrogen, alkyl, alkoxy, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or as described herein. Alternatively, R' and R'' , together with the nitrogen to which they are attached, can form a heterocycloalkyl. The term “primary amino” denotes a group wherein both R' and R''
20 are hydrogen. The term “secondary amino” denotes a group wherein R' is hydrogen and R'' is a group other than hydrogen. The term “tertiary amino” denotes a group wherein both R' and R'' are other than hydrogen. Particular secondary and tertiary amines are methylamine, ethylamine, propylamine, isopropylamine, phenylamine, benzylamine dimethylamine, diethylamine, dipropylamine and diisopropylamine.

25 The term “active pharmaceutical ingredient” (or “API”) denotes the compound or molecule in a pharmaceutical composition that has a particular biological activity.

The terms “pharmaceutical composition” and “pharmaceutical formulation” (or “formulation”) are used interchangeably and denote a mixture or solution comprising a therapeutically effective amount of an active pharmaceutical ingredient together with
30 pharmaceutically acceptable excipients to be administered to a mammal, e.g., a human in need thereof.

The term “pharmaceutically acceptable” denotes an attribute of a material which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither

biologically nor otherwise undesirable and is acceptable for veterinary as well as human pharmaceutical use.

The terms “pharmaceutically acceptable excipient”, “pharmaceutically acceptable carrier” and “therapeutically inert excipient” can be used interchangeably and denote any
5 pharmaceutically acceptable ingredient in a pharmaceutical composition having no therapeutic activity and being non-toxic to the subject administered, such as disintegrators, binders, fillers, solvents, buffers, tonicity agents, stabilizers, antioxidants, surfactants, carriers, diluents or lubricants used in formulating pharmaceutical products.

The terms “individual” or “subject” refer to a mammal. Mammals include, but are not
10 limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

The term “therapeutically effective amount” denotes an amount of a compound or molecule of the present invention that, when administered to a subject, (i) treats or prevents the
15 particular disease, condition or disorder, (ii) attenuates, ameliorates or eliminates one or more symptoms of the particular disease, condition, or disorder, or (iii) prevents or delays the onset of one or more symptoms of the particular disease, condition or disorder described herein. The therapeutically effective amount will vary depending on the compound, the disease state being treated, the severity of the disease treated, the age and relative health of the subject, the route and
20 form of administration, the judgement of the attending medical or veterinary practitioner, and other factors.

The terms “treating” or “treatment” of a disease state include inhibiting the disease state, i.e., arresting the development of the disease state or its clinical symptoms, or relieving the disease state, i.e., causing temporary or permanent regression of the disease state or its clinical
25 symptoms.

The term “spinal muscular atrophy” (or SMA) relates to a disease caused by an inactivating mutation or deletion in the SMN1 gene on both chromosomes, resulting in a loss of SMN1 gene function.
Symptoms of SMA include muscle weakness, poor muscle tone, weak cry, weak cough,
30 limpness or a tendency to flop, difficulty sucking or swallowing, difficulty breathing, accumulation of secretions in the lungs or throat, clenched fists with sweaty hand, flickering/vibrating of the tongue, head often tilted to one side, even when lying down, legs that tend to be weaker than the arms, legs frequently assuming a “frog legs” position, feeding difficulties, increased susceptibility to respiratory tract infections, bowel/bladder weakness,
35 lower-than-normal weight, inability to sit without support, failure to walk, failure to crawl, and

hypotonia, areflexia, and multiple congenital contractures (arthrogryposis) associated with loss of anterior horn cells.

The term “treating spinal muscular atrophy (SMA)” or “treatment of spinal muscular atrophy (SMA)” includes one or more of the following effects: (i) reduction or amelioration of the severity of SMA; (ii) delay of the onset of SMA; (iii) inhibition of the progression of SMA; (iv) reduction of hospitalization of a subject; (v) reduction of hospitalization length for a subject; (vi) increase of the survival of a subject; (vii) improvement of the quality of life of a subject; (viii) reduction of the number of symptoms associated with SMA; (ix) reduction of or amelioration of the severity of one or more symptoms associated with SMA; (x) reduction of the duration of a symptom associated with SMA; (xi) prevention of the recurrence of a symptom associated with SMA; (xii) inhibition of the development or onset of a symptom of SMA; and/or (xiii) inhibition of the progression of a symptom associated with SMA.

More particular, the term “treating SMA” denotes one or more of the following beneficial effects: (i) a reduction in the loss of muscle strength; (ii) an increase in muscle strength; (iii) a reduction in muscle atrophy; (iv) a reduction in the loss of motor function; (v) an increase in motor neurons; (vi) a reduction in the loss of motor neurons; (vii) protection of SMN deficient motor neurons from degeneration; (ix) an increase in motor function; (x) an increase in pulmonary function; and/or (xi) a reduction in the loss of pulmonary function.


In further detail, the term “treating SMA” refers to the functional ability or retention of the functional ability for a human infant or a human toddler to sit up unaided or for a human infant, a human toddler, a human child or a human adult to stand up unaided, to walk unaided, to run unaided, to breathe unaided, to turn during sleep unaided, or to swallow unaided.

The term “EC_{1.5x} concentration for production of full length SMN2 minigene mRNA” (or “EC_{1.5x} minigene”) is defined as that concentration of test compound that is effective in increasing the amount of full length SMN2 minigene mRNA to a level 1.5-fold greater relative to that in vehicle-treated cells.

The term “EC_{1.5x} concentration for SMN protein expression” (or “EC_{1.5x} SMN protein”) is defined as that concentration of test compound that is effective in producing 1.5 times the amount of SMN protein in an SMA patient fibroblast cell compared to the amount produced from the vehicle control.

-12-

In detail, the present invention relates to compounds of formula (I)



wherein

- R^1 is hydrogen or C_{1-7} -alkyl;
- 5 R^2 is hydrogen, cyano, C_{1-7} -alkyl, C_{1-7} -haloalkyl or C_{3-8} -cycloalkyl;
- R^3 is hydrogen, C_{1-7} -alkyl, or C_{3-8} -cycloalkyl;
- A is N-heterocycloalkyl or $NR^{12}R^{13}$, wherein N-heterocycloalkyl comprises 1 or 2 nitrogen ring atoms and is optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} ;
- 10 R^{12} is heterocycloalkyl comprising 1 nitrogen ring atom, wherein heterocycloalkyl is optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} ;
- R^{13} is hydrogen, C_{1-7} -alkyl or C_{3-8} -cycloalkyl;
- R^{14} is independently selected from hydrogen, C_{1-7} -alkyl, amino, amino- C_{1-7} -alkyl, C_{3-8} -cycloalkyl and heterocycloalkyl or two R^{14} together form C_{1-7} -alkylene;
- 15 with the proviso that if A is N-heterocycloalkyl comprising only 1 nitrogen ring atom, then at least one R^{14} substituent is amino or amino- C_{1-7} -alkyl;

and pharmaceutically acceptable salts thereof.

Particular embodiments of the present invention are compounds of formula (I) and pharmaceutically acceptable salts thereof.

- 20 Further, it is to be understood that every embodiment relating to a specific A, R^1 , R^2 or R^3 as disclosed herein may be combined with any other embodiment relating to another A, R^1 , R^2 or R^3 as disclosed herein.

A particular embodiment of the present invention relates to compounds of formula (I) wherein

R^1 is hydrogen or C_{1-7} -alkyl;

R^2 is hydrogen, cyano, C_{1-7} -alkyl, C_{1-7} -haloalkyl or C_{3-8} -cycloalkyl;

5 R^3 is hydrogen, C_{1-7} -alkyl, or C_{3-8} -cycloalkyl;

A is N-heterocycloalkyl comprising 1 or 2 nitrogen ring atoms, wherein N-heterocycloalkyl is optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} ;

10 R^{14} is independently selected from hydrogen, C_{1-7} -alkyl, amino, amino- C_{1-7} -alkyl, C_{3-8} -cycloalkyl and heterocycloalkyl or two R^{14} together form C_{1-7} -alkylene;

with the proviso that if A is N-heterocycloalkyl comprising only 1 nitrogen ring atom, then at least one R^{14} substituent is amino or amino- C_{1-7} -alkyl;

and pharmaceutically acceptable salts thereof.

15 A particular embodiment of the present invention relates to compounds of formula (I), wherein R^1 is C_{1-7} -alkyl, particularly methyl.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R^2 is hydrogen or C_{1-7} -alkyl, particularly hydrogen or methyl.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R^3 is hydrogen or C_{1-7} -alkyl, particularly hydrogen or methyl.

20 A particular embodiment of the present invention relates to compounds of formula (I), wherein R^{12} is piperidinyl optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} .

A particular embodiment of the present invention relates to compounds of formula (I), wherein R^{13} is hydrogen or C_{1-7} -alkyl, particularly hydrogen or methyl.

25 A particular embodiment of the present invention relates to compounds of formula (I), wherein R^{14} is independently selected from C_{1-7} -alkyl and heterocycloalkyl or two R^{14} together form C_{1-7} -alkylene.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R^{14} is independently selected from methyl, ethyl and pyrrolidinyl or two R^{14} together form ethylene.


-14-

A particular embodiment of the present invention relates to compounds of formula (I), wherein A is a saturated mono- or bicyclic N-heterocycloalkyl comprising 1 or 2 nitrogen atoms and is optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴.

5 A particular embodiment of the present invention relates to compounds of formula (I), wherein the N-heterocycloalkyl in A or the heterocycloalkyl in R¹² as defined herein are substituted with 1 or 2 substituents selected from R¹⁴.

A particular embodiment of the present invention relates to compounds of formula (I), wherein the N-heterocycloalkyl in A as defined herein is further characterized in that one ring nitrogen atoms is basic.

10 A particular embodiment of the present invention relates to compounds of formula (I),



X is N or CH;

R⁴ is hydrogen, C₁₋₇-alkyl or -(CH₂)_m-NR⁹R¹⁰;

R⁵ is hydrogen or C₁₋₇-alkyl;

15 R⁶ is hydrogen or C₁₋₇-alkyl;

R⁷ is hydrogen or C₁₋₇-alkyl;

R⁸ is hydrogen or C₁₋₇-alkyl;

R⁹ and R¹⁰ are independently selected from hydrogen, C₁₋₇-alkyl and C₃₋₈-cycloalkyl;

R¹³ is hydrogen, C₁₋₇-alkyl or C₃₋₈-cycloalkyl;

20 n is 0, 1 or 2;

m is 0, 1, 2 or 3;

or R⁴ and R⁵ together form a C₁₋₇-alkylene;

or R⁴ and R⁷ together form a C₁₋₇-alkylene;

-15-

or R⁵ and R⁶ together form a C₂₋₇-alkylene;

or R⁵ and R⁷ together form a C₁₋₇-alkylene;

or R⁵ and R⁹ together form a C₁₋₇-alkylene;

or R⁷ and R⁸ together form a C₂₋₇-alkylene;


5 or R⁷ and R⁹ together form a C₁₋₇-alkylene;

or R⁹ and R¹⁰ together form a C₂₋₇-alkylene;

with the proviso that if X is CH then R⁴ is -(CH₂)_m-NR⁹R¹⁰; and

with the proviso that if X is N and R⁴ is -(CH₂)_m-NR⁹R¹⁰ then m is 2 or 3.

A particular embodiment of the present invention relates to compounds of formula (I),



X is N or CH;

R⁴ is hydrogen, C₁₋₇-alkyl or -(CH₂)_m-NR⁹R¹⁰;

R⁵ is hydrogen or C₁₋₇-alkyl;

R⁶ is hydrogen or C₁₋₇-alkyl;

15 R⁷ is hydrogen or C₁₋₇-alkyl;

R⁸ is hydrogen or C₁₋₇-alkyl;

R⁹ and R¹⁰ are independently selected from hydrogen, C₁₋₇-alkyl and C₃₋₈-cycloalkyl;

n is 0, 1 or 2;

m is 0, 1, 2 or 3;

20 or R⁴ and R⁵ together form C₁₋₇-alkylene;

or R⁴ and R⁷ together form C₁₋₇-alkylene;

-16-

or R⁵ and R⁶ together form C₂₋₇-alkylene;

or R⁵ and R⁷ together form C₁₋₇-alkylene;

or R⁵ and R⁹ together form C₁₋₇-alkylene;

or R⁷ and R⁸ together form C₂₋₇-alkylene;

5 or R⁷ and R⁹ together form C₁₋₇-alkylene;

or R⁹ and R¹⁰ together form C₂₋₇-alkylene;

with the proviso that if X is CH then R⁴ is -(CH₂)_m-NR⁹R¹⁰; and

with the proviso that if X is N and R⁴ is -(CH₂)_m-NR⁹R¹⁰ then m is 2 or 3.

10 It has been found that brain penetration is improved when at least one of R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

In a particular embodiment of the invention at least one of R⁴, R⁵, R⁶, R⁷ and R⁸ is other than hydrogen.

A particular embodiment of the present invention relates to compounds of formula (I), wherein X is N.

15 A particular embodiment of the present invention relates to compounds of formula (I), wherein n is 1.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁴ is hydrogen, methyl or -(CH₂)_m-NR⁹R¹⁰, more particularly hydrogen.

20 A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁵ is hydrogen, methyl or ethyl, more particularly methyl.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁶ is hydrogen or methyl, more particularly hydrogen.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁷ is hydrogen or methyl.

25 A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁸ is hydrogen.

-17-


A particular embodiment of the present invention relates to compounds of formula (I), wherein m is 0.

A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁴ and R⁵ together form propylene.

5 A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁵ and R⁶ together form ethylene;

A particular embodiment of the present invention relates to compounds of formula (I), wherein R⁹ and R¹⁰ together form butylene.


10 A particular embodiment of the present invention relates to compounds of formula (I), wherein A is selected from the group of:



15 wherein R⁴, R⁵, R⁶, R⁷, R⁸ and R¹³ are as defined herein and wherein R¹¹ is hydrogen or C₁₋₇-alkyl.

A particular embodiment of the present invention relates to compounds of formula (I), wherein A is selected from the group of:

-18-




wherein R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined herein and wherein R¹¹ is hydrogen or C₁₋₇-alkyl.

- 5 A particular embodiment of the present invention relates to compounds of formula (I), wherein A is selected from the group of piperazinyl, diazepanyl, pyrrolidinyl and hexahydropyrrolo[1,2-a]pyrazinyl, each optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴ as defined herein.

- 10 A particular embodiment of the present invention relates to compounds of formula (I), wherein A is selected from the group of piperazin-1-yl, 1,4-diazepan-1-yl, pyrrolidin-1-yl and hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, each optionally substituted with 1 or 2 substituents selected from R¹⁴ as defined herein.

A particular embodiment of the present invention relates to compounds of formula (I), wherein A is NR¹²R¹³, wherein R¹² and R¹³ are as described herein.


A particular embodiment of the present invention relates to compounds of formula (I),



wherein A is
herein.

, wherein R^4 , R^5 , R^6 , R^7 , R^8 and R^{13} are as described











A particular embodiment of the present invention relates to compounds of formula (I),




5 wherein A is

, wherein R^{13} is hydrogen or methyl.


A particular embodiment of the present invention relates to compounds of formula (I),
wherein A is selected from the group of:



A particular embodiment of the present invention relates to compounds of formula (I), wherein A is selected from the group of:



A particular embodiment of the present invention relates to a compound of formula (I),
 5 wherein R^1 is methyl, R^2 is hydrogen or methyl, R^3 is hydrogen, and A is



Particular compounds of formula (I) of the present invention are those selected from the group consisting of:

- 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-(4-methylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one;
- 5 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 10 b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 15 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5R)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 20 a]pyrimidin-4-one;
- 7-(1,4-diazepan-1-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 25 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 7-(1,4-diazepan-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 30 7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 35 b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-

- a]pyrimidin-4-one;
 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 5 a]pyrimidin-4-one;
 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one;
 7-(3,3-dimethylpiperazin-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 10 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-
 15 pyrido[1,2-a]pyrimidin-4-one;
 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
 20 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 25 a]pyrimidin-4-one;
 7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 30 9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 7-(3,3-dimethylpiperazin-1-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 35 a]pyrimidin-4-one;
 7-(4,7-diazaspiro[2.5]octan-7-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
 7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-

yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(3R)-3-ethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

and pharmaceutically acceptable salts thereof.

5 Particular compounds of formula (I) of the present invention are those selected from the group consisting of:

7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5R)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;

15 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;

25 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methylpyrido[1,2-a]pyrimidin-4-one;


7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-(4,7-diazaspiro[2.5]octan-7-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

and pharmaceutically acceptable salts thereof.

Compounds of formula (VI) are suitable as intermediates in the manufacture of compounds of formula (I).

Another embodiment of the invention relates to compounds of formula (VI)



5 wherein R^1 , R^2 and R^3 are as described herein;

Y is halogen or trifluoromethanesulfonate;

and salts thereof.

A particular embodiment of the present invention relates to compounds of formula (VI), wherein Y is fluoro, chloro, bromo, iodo or trifluoromethanesulfonate, particularly fluoro.

10 Particular compounds of formula (VI) of the present invention are those selected from the group consisting of:

7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one;

7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

15 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
and salts thereof.

Manufacturing Processes


Compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above can be prepared following standard methods known in the art.

As illustrated in Scheme 1, the commercially available amino-pyridine of formula (II) can be reacted with a malonic ester to afford the intermediate of formula (III), wherein Y and R³ are as described herein and R is C₁₋₂-alkyl, particularly methyl. The compound of formula (III) is then treated with a chlorinating reagent (such as POCl₃ and the like) to provide a compound of formula (IV). The compound of formula (IV) is then reacted in a Suzuki cross-coupling reaction with a compound of formula (V), wherein R¹ and R² are as described herein and Z is B(OH)₂ or an C₁₋₇-alkyl boronic acid ester such as 4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl, in the presence of a catalyst (such as (1,1'-bis(diphenylphosphino)ferrocene)palladium(II) dichloride (Pd(dppf)Cl₂) and the like) and a base (such as K₂CO₃ and the like) in a suitable solvent (such as DMF and the like), to afford the compound of formula (VI). Finally, the compound of formula (VI) is reacted with a compound M-A either in:

- a) an aromatic nucleophilic substitution reaction (particularly if Y is fluoro) by heating at a temperature from 80°C to 200°C; or
- b) a Buchwald-Hartwig amination reaction in the presence of a palladium catalyst (e.g. tetrakis(triphenylphosphine)palladium (Pd(PPh₃)₄) or bis(dibenzylideneacetone)palladium (Pd(dba)₂) by heating at a temperature from 20°C to 100°C;

in a solvent (e.g. dimethyl sulfoxide (DMSO), N-methylpyrrolidone (NMP), or dimethylformamide (DMF)) to give a compound of formula (I), wherein A is as defined herein, M is hydrogen, sodium or potassium, particularly hydrogen, and wherein M is linked to A via a nitrogen atom of A.

-26-

Scheme 1.

In one embodiment, the invention relates to a process for the manufacture of compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above, comprising the


5 reaction of a compound of formula (VI) with a compound M-A either in:

- an aromatic nucleophilic substitution reaction (particularly if Y is fluoro) by heating at a temperature from 80°C to 200°C; or
 - a Buchwald-Hartwig amination reaction in the presence of a palladium catalyst (e.g. tetrakis(triphenylphosphine)palladium (Pd(PPh₃)₄) or
- 10 bis(dibenzylideneacetone)palladium Pd(dba)₂) by heating at a temperature from 20°C to 100°C;

in a solvent (e.g. dimethyl sulfoxide (DMSO), N-methylpyrrolidone (NMP), or dimethylformamide (DMF)), wherein A, Y, R¹, R² and R³ are as defined herein, M is hydrogen, sodium or potassium, particularly hydrogen, and wherein M is linked to A via a nitrogen atom of

15 A.

-27-



A particular embodiment of the invention relates to a process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above, comprising an aromatic nucleophilic substitution reaction between a compound of formula (VI) as described above with a compound of formula M-A by heating in a solvent, wherein A, R¹, R², R³ and Y are as defined above, M is hydrogen, sodium or potassium, and wherein M is linked to A via a nitrogen atom of A.

A particular embodiment of the invention relates to a process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above, wherein the aromatic nucleophilic substitution reaction is performed at a temperature from 80°C to 200°C.

A particular embodiment of the invention relates to a process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above, wherein the solvent of the aromatic nucleophilic substitution reaction is selected from dimethyl sulfoxide (DMSO), N-methylpyrrolidone (NMP), and dimethylformamide (DMF).

A particular embodiment of the invention relates to a process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts thereof as defined above, wherein M is hydrogen.

Particularly, compounds of formula (I) and pharmaceutically acceptable salts thereof can be prepared in accordance to the methods described in the examples herein.

Pharmaceutical Compositions

Another embodiment provides pharmaceutical compositions or medicaments comprising the compounds of the invention and a therapeutically inert carrier, diluent or pharmaceutically acceptable excipient, as well as methods of using the compounds of the invention to prepare such compositions and medicaments.

Compositions are formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the
5 scheduling of administration, and other factors known to medical practitioners.

The compounds of the invention may be administered by any suitable means, including oral, topical (including buccal and sublingual), rectal, vaginal, transdermal, parenteral, subcutaneous, intraperitoneal, intrapulmonary, intradermal, intrathecal and epidural and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions
10 include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration.

The compounds of the present invention may be administered in any convenient administrative form, e.g., tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches, etc. Such compositions may comprise components conventional in pharmaceutical preparations, e.g., diluents, carriers, pH modifiers,
15 preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents, antioxidants, and further active agents. They can also comprise still other therapeutically valuable substances.

A typical formulation is prepared by mixing a compound of the present invention and a carrier or excipient. Suitable carriers and excipients are well known to those skilled in the art and
20 are described in detail in, e.g., *Ansel H.C. et al., Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems (2004) Lippincott, Williams & Wilkins, Philadelphia; Gennaro A.R. et al., Remington: The Science and Practice of Pharmacy (2000) Lippincott, Williams & Wilkins, Philadelphia; and Rowe R.C, Handbook of Pharmaceutical Excipients (2005) Pharmaceutical Press, Chicago*. The formulations may also include one or more buffers, stabilizing agents,
25 surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants, opaquing agents, glidants, processing aids, colorants, sweeteners, perfuming agents, flavoring agents, diluents and other known additives to provide an elegant presentation of the drug (i.e., a compound of the present invention or pharmaceutical composition thereof) or aid in the manufacturing of the pharmaceutical product (i.e., medicament).

30 The dosage at which compounds of the invention can be administered can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 0.01 to 1000 mg per person of a compound of general formula (I) should be appropriate, although the above upper limit can also be exceeded when necessary.

An example of a suitable oral dosage form is a tablet comprising about 100 mg to 500 mg of the compound of the invention compounded with about 30 to 90 mg anhydrous lactose, about 5 to 40 mg sodium croscarmellose, about 5 to 30 mg polyvinylpyrrolidone (PVP) K30, and about 1 to 10 mg magnesium stearate. The powdered ingredients are first mixed together and then
5 mixed with a solution of the PVP. The resulting composition can be dried, granulated, mixed with the magnesium stearate and compressed to tablet form using conventional equipment.

An example of an aerosol formulation can be prepared by dissolving the compound, for example 10 to 100 mg, of the invention in a suitable buffer solution, e.g. a phosphate buffer, adding a tonicifier, e.g. a salt such as sodium chloride, if desired. The solution may be filtered,
10 e.g., using a 0.2 μm filter, to remove impurities and contaminants.

Uses

As described above, the compounds of formula (I) and their pharmaceutically acceptable
15 salts possess valuable pharmacological properties and have been found to enhance inclusion of exon 7 of SMN1 and/or SMN2 into mRNA transcribed from the SMN1 and/or SMN2 gene, thereby increasing expression of SMN protein in a human subject in need thereof.

The compounds of the present invention can be used, either alone or in combination with other drugs, for the treatment or prevention of diseases caused by an inactivating mutation or
20 deletion in the SMN1 gene and/or associated with loss or defect of SMN1 gene function. These diseases include, but are not limited to spinal muscular atrophy (SMA).

A particular embodiment of the present invention relates to pharmaceutical compositions comprising compounds of formula (I) as defined above or their pharmaceutically acceptable salts as defined above and one or more pharmaceutically acceptable excipients.

25 A particular embodiment of the present invention relates to pharmaceutical compositions comprising compounds of formula (I) or their pharmaceutically acceptable salts as defined above and one or more pharmaceutically acceptable excipients for the treatment or prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or associated with loss or defect of SMN1 gene function, particularly for the treatment or prevention of SMA.

30 A particular embodiment of the present invention relates to compounds of formula (I) or their pharmaceutically acceptable salts as defined above for use as therapeutically active substances, especially for use as therapeutically active substances for the treatment or prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or associated

-30-

with loss or defect of SMN1 gene function, particularly for the treatment or prevention of spinal muscular atrophy (SMA).

A particular embodiment of the present invention relates to compounds of formula (I) or their pharmaceutically acceptable salts as defined above for the use in the treatment or
5 prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or associated with loss or defect of SMN1 gene function, particularly for use in the treatment or prevention of spinal muscular atrophy (SMA).

A particular embodiment of the present invention relates to a method for the treatment or prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or
10 associated with loss or defect of SMN1 gene function, particularly for the treatment or prevention of spinal muscular atrophy (SMA), which method comprises administering compounds of formula (I) or their pharmaceutically acceptable salts as defined above to a subject.

A particular embodiment of the present invention relates to the use of compounds of formula (I) or their pharmaceutically acceptable salts as defined above for the treatment or
15 prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or associated with loss or defect of SMN1 gene function, particularly for the treatment or prevention of spinal muscular atrophy (SMA).

A particular embodiment of the present invention relates to the use of compounds of formula (I) or their pharmaceutically acceptable salts as defined above for the preparation of
20 medicaments for the treatment or prevention of diseases caused by an inactivating mutation or deletion in the SMN1 gene and/or associated with loss or defect of SMN1 gene function, particularly for the treatment or prevention of spinal muscular atrophy (SMA). Such medicaments comprise compounds of formula (I) or their pharmaceutically acceptable salts as defined above.

25

Examples

The invention will be more fully understood by reference to the following examples. They should however not be construed as limiting the scope of the invention.

5 Abbreviations used


ACN: Acetonitrile; CH₂Cl₂: dichloromethane (DCM); DIPEA: diisopropyl ethylamine; DMA: dimethyl acetamide; TEA: triethylamine; RT: room temperature; B₂(pin)₂: bis(pinacolato)diboron; Pd(dppf)Cl₂: (1,1'-Bis(diphenylphosphino)ferrocene)palladium(II) dichloride; PPTS: Pyridinium p-toluenesulfonate.

10

Intermediate 1

7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one

a) 2-chloro-7-fluoro-pyrido[1,2-a]pyrimidin-4-one




15 A mixture of 2-amino-5-fluoropyridine (11.20 g, 0.10 mol) and dimethyl malonate (57.0 mL, 0.50 mol) was heated at 230 °C for 1.5 h. After cooling to room temperature, the precipitate was filtered and washed with ACN (3x) to give 7-fluoro-2-hydroxy-4H-pyrido[1,2-a]pyrimidin-4-one as a dark solid (14 g), which was used directly in the next step. MS *m/z* 181.3 [M+H]⁺.


20 A dark mixture of crude 7-fluoro-2-hydroxy-4H-pyrido[1,2-a]pyrimidin-4-one (14g, ~77 mmol) in POCl₃ (50 mL) and DIPEA (13.3 mL, 77 mmol) was heated at 110 °C for 15 hours. The solvent was removed and the dark residue was treated with ice-water, washed with water (3x) and dried to give a brown solid. The crude brown solid was chromatographed (5% MeOH in CH₂Cl₂) to give 2-chloro-7-fluoro-4H-pyrido[1,2-a]pyrimidin-4-one as a yellow solid (9.84 g, 50%, 2 steps), MS *m/z* 199.2 [M+H]⁺.

25

-32-

b) 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine

A mixture of 6-chloro-2-methylimidazo[1,2-b]pyridazine (900 mg, 5.37 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (1.36 g, 5.37 mmol, 1.0 eq.), KOAc (1.05 g, 10.7 mmol) and Pd(dppf)Cl₂•CH₂Cl₂ (393 mg, 0.54 mmol) in dioxane (50 mL) was
 5 degassed and heated under N₂ at 95 °C. After 15 hours, the mixture was diluted with EtOAc, filtered through celite and concentrated under vacuum to give 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine which was used directly in the next step.

c) 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one

10


To a solution of 2-chloro-7-fluoro-4H-pyrido[1,2-a]pyrimidin-4-one (750 mg, 3.78 mmol) in ACN (36 mL) was added 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine (1.17 g, 4.53 mmol, Eq: 1.2), Pd(PPh₃)₄ (218 mg, 0.189 mmol, 0.05 eq.) and an aqueous solution of K₂CO₃ (3.78 mL, 7.55 mmol, 2.0 eq.). The mixture was degassed
 15 and heated under argon at 105 °C overnight. The reaction was cooled to RT, and filtered. The precipitate was washed with Et₂O and then water, dried *in vacuo* to give 250 mg (22%) of 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one as a light brown solid. MS *m/z* 296.1 [M+H]⁺.

20

Intermediate 2

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one

a) 2,8-dimethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine




5 In a sealed flask, 3,6-dichloro-4-methylpyridazine (27 g, 161 mmol) was suspended in aqueous ammonia (25%, 300 mL). The reaction mixture was heated at 110 °C for 48 hours (turned into solution after 1 hour). After cooling to room temperature, the reaction was poured into CH₂Cl₂, and the organic phase was separated, dried over Na₂SO₄, and concentrated under vacuum, to give 22.4 g of 6-chloro-4-methyl-pyridazin-3-amine and 6-chloro-5-methyl-
10 pyridazin-3-amine as a mixture of regioisomers which were used directly in the next step.

The mixture of regioisomers 6-chloro-4-methyl-pyridazin-3-amine and 6-chloro-5-methyl-pyridazin-3-amine (22.4 g) was suspended in 2-propanol (300 mL). 1-bromo-2,2-dimethoxypropane (36.0 g, 26.6 mL, 193 mmol, 1.2 eq.) and PPTS (2.96 g, 11.6 mmol, 0.0725 eq.) were added, and the resulting solution was heated at 105 °C overnight. The solvent was
15 removed *in vacuo* and the residue was taken up in CH₂Cl₂ and washed with NaHCO₃. The organic phases were dried over Na₂SO₄, concentrated *in vacuo* and the crude light brown solid was chromatographed (EtOAc / Heptane 1/2 -1/1) to give separately 6.1 g of 6-chloro-2,8-dimethyl-imidazo[1,2-b]pyridazine MS *m/z* 182.1 [M+H]⁺ (21%) as a white solid and 5.9 g of 6-chloro-2,7-dimethyl-imidazo[1,2-b]pyridazine MS *m/z* 182.1 [M+H]⁺ (20%) as a white solid.

20 A mixture of 6-chloro-2,8-dimethylimidazo[1,2-b]pyridazine (0.9 g, 4.96 mmol), 4,4,4',4',5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (1.26 g, 4.96 mmol, 1.0 eq.), KOAc (0.97 g, 9.91 mmol) and Pd(dppf)Cl₂•CH₂Cl₂ (363 mg, 0.49 mmol) in dioxane (50 mL) was degassed and heated under N₂ at 110 °C. After 15 hours, the mixture was diluted with EtOAc, filtered through celite and concentrated under vacuum to give 2,8-dimethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine which was used directly in the
25 next step.

-34-

b) 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one




To a solution of 2-chloro-7-fluoro-4H-pyrido[1,2-a]pyrimidin-4-one (750 mg, 3.78 mmol, described herein above) in ACN (36 mL) was added 2,8-dimethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine (1.24 g, 4.53 mmol, 1.2 eq.), Pd(Ph₃P)₄ (218 mg, 0.189 mmol, 0.05 eq.) and an aqueous solution of K₂CO₃ (3.78 mL, 7.55 mmol, 2.0 eq.). The mixture was degassed and heated under argon at 100 °C for 6 hours. The reaction was cooled to RT, and filtered. The precipitate was washed with Et₂O and then water, dried *in vacuo* to give 700 mg (60%) of 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one as a light brown solid. MS *m/z* 310.1 [M+H]⁺.

Intermediate 3

7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one

a) 2-chloro-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one



15


A mixture of 5-fluoro-3-methylpyridin-2-amine (3.3 g, 26.2 mmol) and dimethyl malonate (15.0 mL, 0.13 mol, 5.0 eq.) was heated at 210 °C for 1.5 hours. After cooling to room temperature, the precipitate was filtered and washed with ACN (3x) to give 7-fluoro-2-hydroxy-9-methyl-pyrido[1,2-a]pyrimidin-4-one as a dark solid (2.3 g), which was used directly in the next step. MS *m/z* 195.1 [M+H]⁺.

20

A mixture of crude 7-fluoro-2-hydroxy-9-methyl-pyrido[1,2-a]pyrimidin-4-one (2.3 g, 11.8 mmol) in POCl₃ (7.7 mL, 82.9 mmol) and DIEA (2.07 mL, 11.8 mmol) was heated at 110 °C for 15 hours. The solvent was removed and the residue was treated with ice-water, washed with water (3x) and dried to give a brown solid. The crude brown solid was

chromatographed (5% MeOH in CH₂Cl₂) to give 2-chloro-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one as a yellow solid (1.77 g, 70% over 2 steps), MS *m/z* 213.1 [M+H]⁺.

b) 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




To a solution of 2-chloro-7-fluoro-9-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (2.2 g, 10.3 mmol) in ACN (80 mL) was added 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine (3.22 g, 12.4 mmol, 1.2 eq., described herein above), Pd(PPh₃)₄ (1.20 g, 1.03 mmol, 0.1 eq.) and an aqueous solution of K₂CO₃ (10.3 mL, 20.7 mmol, 2.0 eq.).

10 The mixture was degassed and heated under argon at 100 °C for 6 hours. The reaction was cooled to RT, and filtered. The precipitate was washed with Et₂O and then water, dried *in vacuo* to give 1.80 g (56%) of 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one as a light brown solid. MS *m/z* 310.1 [M+H]⁺.

15

Intermediate 4

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one



To a solution of 2-chloro-7-fluoro-9-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (0.98 g, 4.61 mmol, described herein above) in ACN (50 mL) was added 2,8-dimethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine (1.51 g, 5.53 mmol, 1.2 eq., described herein above), Pd(PPh₃)₄ (0.32 g, 0.277 mmol, 0.06 eq.) and an aqueous solution of K₂CO₃ (4.61 mL, 9.22 mmol, 2.0 eq.). The mixture was degassed and heated under argon at 100 °C for 6 hours. The reaction was cooled to RT, and filtered. The precipitate was washed with Et₂O and water, then dried *in vacuo* to give 0.89 g (60%) of 2-(2,8-dimethylimidazo[1,2-


20

25

b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one as a light brown solid. MS m/z 324.4 $[M+H]^+$.

Example 1


5 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-(4-methylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 35 mg, 0.119 mmol) and 1-methylpiperazine (47.5 mg, 0.474 mmol, 4 eq.) were stirred in DMSO (1 mL) at 120°C overnight. LC-MS showed total
10 conversion. The solvent was removed under high vacuum. The crude product was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to 9/1) to afford the title product (25 mg, 56%) as a light yellow solid. MS m/z 376.3 $[M+H]^+$.

Example 2


15 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 125 mg, 0.426 mmol) and (R)-octahydropyrrolo-[1,2-a]pyrazine (160 mg, 1.27 mmol, 3 eq.) were stirred in DMSO (5 mL) at 125°C overnight. The
20 solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 ,
25 $\text{CH}_2\text{Cl}_2/\text{MeOH}=98/2$ to 95/5) to afford the title product (65 mg, 38%) as a light yellow solid. MS m/z 402.5 $[M+H]^+$.

Example 3

7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




5

In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 200 mg, 0.647 mmol) and (S)-octahydropyrrolo-[1,2-a]pyrazine (286 mg, 2.26 mmol, 3.5 eq.) were stirred in DMSO (5 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=98/2 to 95/5) to afford the title product (115 mg, 43%) as a light yellow solid. MS *m/z* 416.3 [M+H⁺].

15

Example 4

7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 200 mg, 0.647 mmol), DIPEA (0.113 mL, 0.67 mmol, 1 eq.) and (R)-octahydropyrrolo-[1,2-a]pyrazine (245 mg, 1.95 mmol, 3.0 eq.) were stirred in DMSO (2.5 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic

20


-38-

layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=98/2$ to $95/5$) to afford the title product (132 mg, 49%) as a light yellow solid. MS m/z 416.3 $[\text{M}+\text{H}^+]$.

5

Example 5

7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 90 mg, 0.291 mmol), DIPEA (0.05 mL, 0.29 mmol, 1eq.) and (S)-8a-methyloctahydropyrrolo[1,2-a]pyrazine (81 mg, 0.58 mmol, 2.0 eq.) were stirred in DMSO (2.5 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $90/10$) to afford the title product (55 mg, 44%) as a light yellow solid. MS m/z 430.3 $[\text{M}+\text{H}^+]$.

20

Example 6

7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 90 mg, 0.291 mmol), DIPEA (0.05 mL, 0.29 mmol, 1eq.)

-39-

and (R)-8a-methyloctahydropyrrolo[1,2-a]pyrazine (81 mg, 0.58 mmol, 2.0 eq.) were stirred in DMSO (2.5 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by
 5 column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (50 mg, 40%) as a light yellow solid. MS *m/z* 430.4 [M+H⁺].

Example 7


10 **2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5R)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one**



In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 50 mg, 0.162 mmol), and cis-2,6-dimethylpiperazine (74 mg, 0.647 mmol, 4.0 eq.) were stirred in DMSO (1.5 mL) at 110°C overnight. The solvent was
 15 removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (32 mg, 49%) as a light yellow solid. MS *m/z* 404.4 [M+H⁺].

20


-40-

Example 8**2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one**

- 5 In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 33 mg, 0.107 mmol), and (S)-2-methylpiperazine (43 mg, 0.427 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 120°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and
- 10 concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (18 mg, 43%) as a light yellow solid. MS *m/z* 390.3 [M+H⁺].

Example 9


- 15 **2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one**



- In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 85 mg, 0.275 mmol), and (R)-2-methylpiperazine (110 mg, 1.10 mmol, 4.0 eq.) were stirred in DMSO (5 mL) at 120°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (35 mg, 33%) as a light yellow solid. MS *m/z* 390.3 [M+H⁺].
- 20

Example 10

7-(1,4-diazepan-1-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




5

In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 33 mg, 0.107 mmol), and 1,4-diazepane (32 mg, 0.320 mmol, 3.0 eq.) were stirred in DMSO (2 mL) at 120°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (20 mg, 48%) as a light yellow solid. MS *m/z* 390.3 [M+H⁺].

10

Example 11

2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one



15


20

In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), and (S)-2-methylpiperazine (68 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 110°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (40 mg, 63%) as a light yellow solid.

-42-

MS m/z 376.2 $[M+H]^+$.**Example 12**

5 **2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one**




In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), and (R)-2-methylpiperazine (68 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 110°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of $NaHCO_3$. The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $CH_2Cl_2/MeOH=95/5$ to 90/10) to afford the title product (48 mg, 75%) as a light yellow solid. MS m/z 376.3 $[M+H]^+$.

15

Example 13

7-(1,4-diazepan-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




20 In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), and 1,4-diazepane (68 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 110°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of $NaHCO_3$. The organic layer was separated and dried over Na_2SO_4 and concentrated

-43-

in vacuo. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (41 mg, 65%) as a light yellow solid. MS *m/z* 376.2 [M+H⁺].

Example 14


5 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), and *cis*-2,6-dimethylpiperazine (77
10 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 110°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (41 mg, 62%) as a light yellow solid.
15 MS *m/z* 390.3 [M+H⁺].

Example 15

7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



20

In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), and (S)-octahydropyrrolo[1,2-a]pyrazine (85 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The


-44-

solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $90/10$) to afford the title product (36 mg, 53%) as a light yellow solid.

5 MS m/z 402.3 $[\text{M}+\text{H}^+]$.

Example 16

7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




10

In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol) and (S)-8a-methyloctahydropyrrolo[1,2-a]pyrazine (95 mg, 0.677 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $90/10$) to afford the title product (45 mg, 64%) as a light yellow solid. MS m/z 416.3 $[\text{M}+\text{H}^+]$.

15

Example 17

7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-


20

-45-

a]pyrimidin-4-one (**Intermediate 1**; 100 mg, 0.339 mmol) and (R)-8a-methyloctahydropyrrolo[1,2-a]pyrazine (190 mg, 1.35 mmol, 4.0 eq.) were stirred in DMSO (4 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was
 5 separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (45 mg, 64%) as a light yellow solid. MS *m/z* 416.3 [M+H⁺].

Example 18


10 **2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one**



In a microwave reactor, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 45 mg, 0.145 mmol), (R)-1,3'-bipyrrolidine dihydrochloride
 15 (62 mg, 0.291 mmol, 2.0 eq.) and DIPEA (0.20 mL, 1.16 mmol, 8 eq.) were stirred in NMP (3 mL) at 220°C for 1 hour. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=98/2 to 90/10) to afford the title product (25 mg, 40%) as
 20 a light yellow solid. MS *m/z* 430.3 [M+H⁺].

Example 19


7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



5 In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 50 mg, 0.169 mmol), DIPEA (0.24 mL, 1.35 mmol, 8 eq.) and 4,7-diazaspiro[2.5]octane dihydrochloride (62.7 mg, 0.339 mmol, 2.0 eq.) were stirred in DMSO (2 mL) at 125°C for 2 days. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (22 mg, 33%) as a light yellow solid. MS *m/z* 388.3 [M+H⁺].

Example 20

7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




20 In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 50 mg, 0.162 mmol), DIPEA (0.22 mL, 1.29 mmol, 4 eq.) and 4,7-diazaspiro[2.5]octane dihydrochloride (32 mg, 0.320 mmol, 3.0 eq.) were stirred in DMSO (2 mL) at 130°C for 48 hours. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=98/2 to 95/5) to afford the title product (12 mg,

-47-

18%) as a light yellow solid. MS m/z 402.3 $[M+H^+]$.

Example 21

5 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one




10 In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 40 mg, 0.135 mmol), DIPEA (0.19 mL, 1.08 mmol, 8 eq.) and (R)-1,3'-bipyrrolidine dihydrochloride (58 mg, 0.271 mmol, 2.0 eq.) were stirred in DMSO (4 mL) and heated at 220°C for 40 minutes in a microwave. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of $NaHCO_3$. The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $CH_2Cl_2/MeOH=98/2$ to 90/10) to afford the title product (30 mg, 53%) as a light yellow solid. MS m/z 416.3 $[M+H^+]$.

15

Example 22

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one




20 In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 40 mg, 0.129 mmol) and 2,2-dimethylpiperazine (59 mg, 0.517 mmol, 4.0 eq.) were stirred in DMSO (1.6 mL) at 130°C overnight. The solvent was

-48-

removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $9/1$) to afford the title product (29 mg, 55%) as a light yellow solid. MS m/z 404.3 $[\text{M}+\text{H}^+]$.

Example 23

7-(3,3-dimethylpiperazin-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




10

In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 40 mg, 0.135 mmol) and 2,2-dimethylpiperazine (62 mg, 0.542 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $90/10$) to afford the title product (26 mg, 49%) as a light yellow solid. MS m/z 390.3 $[\text{M}+\text{H}^+]$.

20

Example 24

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one




-49-

In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 4**; 50 mg, 0.155 mmol) and (S)-2-methylpiperazine (62 mg, 0.619 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (45 mg, 72%) as a light yellow solid. MS *m/z* 404.3 [M+H⁺].

10


Example 25

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 4**; 50 mg, 0.155 mmol) and (R)-2-methylpiperazine (62 mg, 0.619 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (40 mg, 70%) as a light yellow solid. MS *m/z* 404.3 [M+H⁺].

-50-

Example 26**2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-pyrido[1,2-a]pyrimidin-4-one**

5 In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 4**; 50 mg, 0.155 mmol) and *cis*-2,6-dimethylpiperazine (70 mg, 0.619 mmol, 4.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and
 10 dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (26 mg, 40%) as a light yellow solid. MS *m/z* 418.3 [M+H⁺].

Example 27**2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one**


In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 4**; 50 mg, 0.155 mmol) and 2,2-dimethylpiperazine
 20 (35 mg, 0.309 mmol, 2.0 eq.) were stirred in DMSO (2 mL) at 125°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂,

-51-

CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (36 mg, 56%) as a light yellow solid. MS *m/z* 418.3 [M+H⁺].

Example 28


5 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 4**; 50 mg, 0.155 mmol), DIPEA (0.21 mL, 1.24 mmol, 8 eq.) and 4,7-diazaspiro[2.5]octane dihydrochloride (57 mg, 0.309 mmol, 2.0 eq.) were stirred in DMSO (2 mL) at 125°C for 2 days. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (17 mg, 26%) as a light yellow solid. MS *m/z* 416.3 [M+H⁺].

Example 29

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 50 mg, 0.162 mmol), TEA (0.18 mL, 1.29 mmol, 8 eq.) and

-52-


(2S,6S)-2,6-dimethylpiperazine dihydrochloride (90 mg, 0.485 mmol, 3.0 eq.) were stirred in DMSO (2 mL) at 140°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 9/1) to afford the title product (20 mg, 30%) as a light yellow solid. MS *m/z* 404.3 [M+H⁺].

Example 30

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 2**; 50 mg, 0.162 mmol), DIPEA (0.22 mL, 1.29 mmol, 8 eq.) and (S)-1,3'-bipyrrolidine dihydrochloride (103 mg, 0.485 mmol, 3.0 eq.) were stirred in NMP (2 mL) at 140°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 9/1) to afford the title product (22 mg, 32%) as a light yellow solid. MS *m/z* 430.3 [M+H⁺].

Example 31**2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one**

- 5 In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 75 mg, 0.254 mmol), TEA (0.28 mL, 2.03 mmol, 8 eq.) and (S)-1,3'-bipyrrolidine dihydrochloride (162 mg, 0.762 mmol, 3.0 eq.) were stirred in NMP (4 mL) and heated at 220°C for 1 hour in a microwave. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃.
- 10 The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (12 mg, 11%) as a light yellow solid. MS *m/z* 416.2 [M+H⁺].

Example 32

- 15 **7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one**




- In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 75 mg, 0.254 mmol), TEA (0.28 mL, 2.03 mmol, 8 eq.) and
- 20 (2S,6S)-2,6-dimethylpiperazine dihydrochloride (143 mg, 0.762 mmol, 3.0 eq.) were stirred in DMSO (3 mL) and heated at 140°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude

-54-

was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (10 mg, 10%) as a light yellow solid. MS *m/z* 390.3 [M+H⁺].

Example 33


5 9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one



In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one (**Intermediate 3**; 250 mg, 0.808 mmol), and (S)-2-methylpiperazine (405 mg, 4.04 mmol, 5.0 eq.) were stirred in DMSO (6 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 85/15) to afford the title product (135 mg, 43%) as a light yellow solid. MS *m/z* 390.3 [M+H⁺].

Example 34

9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one



20


In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one (**Intermediate 3**; 250 mg, 0.808 mmol), and (R)-2-methylpiperazine (405 mg, 4.04 mmol, 5.0 eq.) were stirred in DMSO (6 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an

aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $85/15$) to afford the title product (100 mg, 32%) as a light yellow solid. MS m/z 390.3 $[\text{M}+\text{H}^+]$.

5

Example 35


7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



- 10 In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one (**Intermediate 3**; 250 mg, 0.808 mmol), and (2S,6R)-2,6-dimethylpiperazine (461 mg, 4.04 mmol, 5.0 eq.) were stirred in DMSO (6 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH_2Cl_2 and washed with an aqueous saturated solution of NaHCO_3 . The organic layer was separated and dried over
- 15 Na_2SO_4 and concentrated *in vacuo*. The crude was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}=95/5$ to $85/15$) to afford the title product (101 mg, 31%) as a light yellow solid. MS m/z 404.3 $[\text{M}+\text{H}^+]$.

Example 36

- 20 7-(3,3-dimethylpiperazin-1-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one




In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-

-56-

a]pyrimidin-4-one (**Intermediate 3**; 250 mg, 0.808 mmol), and 2,2-dimethylpiperazine (461 mg, 4.04 mmol, 5.0 eq.) were stirred in DMSO (6 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 85/15) to afford the title product (120 mg, 36%) as a light yellow solid. MS *m/z* 404.3 [M+H⁺].

Example 37

10 **7-(4,7-diazaspiro[2.5]octan-7-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one**




In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one (**Intermediate 3**; 125 mg, 0.404 mmol), K₂CO₃ (223 mg, 1.62 mmol, 4 eq.) and 4,7-diazaspiro[2.5]octane dihydrochloride (112 mg, 0.606 mmol, 1.5 eq.) were stirred in DMA (2 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃. The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (75 mg, 46%) as a light yellow solid. MS *m/z* 402.2 [M+H⁺].

-57-

Example 38


7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



- 5 In a sealed tube, 7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one (**Intermediate 3**; 125 mg, 0.404 mmol), K₂CO₃ (223 mg, 1.62 mmol, 4 eq.) and (2S,6S)-2,6-dimethylpiperazine dihydrochloride (113 mg, 0.606 mmol, 1.5 eq.) were stirred in DMA (2 mL) and heated at 130°C overnight. The solvent was removed under high vacuum. The residue was taken up in CH₂Cl₂ and washed with an aqueous saturated solution of NaHCO₃.
- 10 The organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 90/10) to afford the title product (50 mg, 31%) as a light yellow solid. MS *m/z* 404.3 [M+H⁺].

Example 39

- 15 7-[(3R)-3-ethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one



- In a sealed tube, 7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**Intermediate 1**; 200 mg, 0.677 mmol), K₂CO₃ (374 mg, 2.71 mmol, 4 eq.) and (R)-2-ethylpiperazine dihydrochloride (238 mg, 0.606 mmol, 1.5 eq.) were stirred in DMA (3 mL) at 100°C for 4 days. The solvent was removed under high vacuum. The crude was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH=95/5 to 8/2) to afford the title product (168 mg, 64%) as a light yellow solid. MS *m/z* 390.2 [M+H⁺].
- 20

Biological Assays

To describe in more detail and assist in understanding the present description, the following non-limiting biological examples are offered to more fully illustrate the scope of the description and are not to be construed as specifically limiting the scope thereof. Such variations of the present description that may be now known or later developed, which would be within the purview of one skilled in the art to ascertain, are considered to fall within the scope of the present description and as hereinafter claimed. These examples illustrate the testing of certain compounds described herein in vitro and/or in vivo and demonstrate the usefulness of the compounds for treating of SMA by enhancing the inclusion of exon 7 of SMN2 into mRNA transcribed from the SMN2 gene. Compounds of formula (I) enhance inclusion of exon 7 of SMN2 into mRNA transcribed from the SMN2 gene and increase levels of SMN protein produced from the SMN2 gene, and thus can be used to treat SMA in a human subject in need thereof. These examples further illustrate the testing of certain compounds described herein in vitro and/or in vivo and demonstrate the usefulness of the compounds for enhancing the inclusion of exon 7 of SMN1 into mRNA transcribed from the SMN1 gene. Accordingly, compounds of formula (I) also enhance the inclusion of exon 7 of SMN1 into mRNA transcribed from the SMN1 gene and increase levels of SMN protein produced from the SMN1 gene.

Assay 1**SMN2 minigene mRNA splicing RT-qPCR assay in cultured cells**

The reverse transcription-quantitative PCR-based (RT-qPCR) assay is used to quantify the level of the full length SMN2 minigene (referred to herein by the term "FL SMN2mini") mRNA containing SMN2 exon 7 in a HEK293H cell line stably transfected with said minigene and treated with a test compound. Materials used and respective sources are listed below in Table 1.

Material	Source
HEK293H cells	Life Technologies, Inc. (formerly Invitrogen) Catalog No. 11631-017
Cells-To-Ct lysis buffer	Life Technologies, Inc. (formerly Applied Biosystems) part No. 4399002
DMEM	Life Technologies, Inc. (formerly Invitrogen) Catalog No. 11960-044
96-well flat-bottom plates	Becton Dickinson Catalog No. 353072

RT-PCR Enzyme Mix	Life Technologies, Inc. (formerly Applied Biosystems) part No. 4388520
RT-PCR buffer	Life Technologies, Inc. (formerly Applied Biosystems) part No. 4388519
AgPath-ID One-Step RT-PCR kit	Life Technologies, Inc. (formerly Applied Biosystems) part No. 4387391
Thermocycler	Life Technologies, Inc. (formerly Applied Biosystems) 7900HT

Table 1. Materials and their respective sources used in the SMN2 minigene mRNA splicing RT-qPCR assay in cultured cells.

The SMN2-A minigene construct was prepared as described in International Patent Application WO2009/151546A1 page 145 paragraph [00400] to page 147 paragraph [00412] (incl. Figure 1 and Figure 3 therein).

HEK293H cells stably transfected with the SMN2-A minigene construct (10,000 cells/well) are seeded in 200 μ L of cell culture medium (DMEM plus 10% FBS, with 200 μ g/mL hygromycin) in 96-well flat-bottom plates and the plate is immediately swirled to ensure proper dispersal of cells and the formation of an even monolayer of cells. Cells are allowed to attach for 6 hours. Test compounds are serially diluted 3.16-fold in 100% DMSO to generate a 7-point concentration curve. A solution of test compound (1 μ L, 200x in DMSO) is added to each cell-containing well and the plate is incubated for 24 hours in a cell culture incubator (37°C, 5% CO₂, 100% relative humidity). 2 replicates are prepared for each test compound concentration. The cells are then lysed in the Cells-To-Ct lysis buffer and the lysate is stored at -80°C.

Full length SMN2-A minigene and GAPDH mRNA are quantified using the primers and probes referenced in Table 2. Primer SMN Forward A (SEQ ID NO.1) hybridizes to a nucleotide sequence in exon 7 (nucleotide 22 to nucleotide 40), primer SMN Reverse A (SEQ ID NO.2) hybridizes to a nucleotide sequence in the coding sequence of Firefly luciferase, SMN Probe A (SEQ ID NO.3) hybridizes to a nucleotide sequence in exon 7 (nucleotide 50 to nucleotide 54) and exon 8 (nucleotide 1 to nucleotide 21). The combination of these three oligonucleotides detects only SMN1 or SMN2 minigenes (RT-qPCR) and will not detect endogenous SMN1 or SMN2 genes.

Primers/Probes	Sequences	Source
SMN Forward Primer A	SEQ ID NO. 1: GAAGGAAGGTGCTCACATT	PTC ¹
SMN Reverse Primer A	SEQ ID NO. 2: TCTTTATGTTTTTGGCGTCTTC	PTC ¹

SMN Forward Probe A	SEQ ID NO. 3: 6FAM- AAGGAGAAATGCTGGCAT AGAGCAGC-TAMRA	PTC ¹
hGAPDH Forward Probe	SEQ ID NO. 4: VIC-CGCCTGGTCACCAGGGCTGCT- TAMRA	LTI ²
hGAPDH Forward Primer	SEQ ID NO. 5: CAACGGATTTGGTCGTATTGG	LTI ²
hGAPDH Reverse Primer	SEQ ID NO. 6: TGATGGCAACAATATCCACTTTACC	LTI ²

Table 2. ¹ Primers and probes designed by PTC Therapeutics, Inc.; ² Commercially available from Life Technologies, Inc. (formerly Invitrogen).

The SMN forward and reverse primers are used at final concentrations of 0.4 μ M. The SMN probe is used at a final concentration of 0.15 μ M. The GAPDH primers are used at final concentrations of 0.2 μ M and the probe at 0.15 μ M.

The SMN2-minigene GAPDH mix (15 μ L total volume) is prepared by combining 7.5 μ L of 2x RT-PCR buffer, 0.4 μ L of 25x RT-PCR enzyme mix, 0.75 μ L of 20x GAPDH primer-probe mix, 4.0075 μ L of water, 2 μ L of 10-fold diluted cell lysate, 0.06 μ L of 100 μ M SMN forward primer, 0.06 μ L of 100 μ M SMN reverse primer, and 0.225 μ L of 100 μ M SMN probe.

PCR is carried out at the following temperatures for the indicated time: Step 1: 48°C (15 min); Step 2: 95°C (10 min); Step 3: 95°C (15 sec); Step 4: 60°C (1 min); then repeat Steps 3 and 4 for a total of 40 cycles.

Each reaction mixture contains both SMN2-A minigene and GAPDH primers/probe sets (multiplex design), allowing simultaneous measurement of the levels of two transcripts.

The increase in the abundance of the FL SMN2mini mRNA relative to that in cells treated with vehicle control is determined from real-time PCR data using a modified $\Delta\Delta C_t$ method (as described in *Livak and Schmittgen, Methods, 2001, 25:402-8*). The amplification efficiency E is calculated from the slope of the amplification curve for FL SMN2mini and GAPDH individually. The abundance of FL SMN2mini and GAPDH mRNA are then calculated as $(1 + E)^{-C_t}$, where C_t is the threshold value for each amplicon. The abundance of FL SMN2mini mRNA is normalized to GAPDH mRNA abundance. The normalized FL SMN2mini mRNA abundance from test compound-treated samples is then divided by normalized FL SMN2mini mRNA abundance from vehicle-treated cells to determine the level of FL SMN2mini mRNA relative to vehicle control.

Table 3 provides EC_{1.5x} concentrations for production of full length SMN2 minigene mRNA that was obtained from the 7-point concentration data generated according to the above procedure for particular compounds of the present invention.

Particular compounds of the present invention exhibit an EC_{1.5x} concentration for
5 production of full length SMN2 minigene mRNA $\leq 1 \mu\text{M}$.

More particular compounds of the present invention exhibit an EC_{1.5x} concentration for production of full length SMN2 minigene mRNA $\leq 0.1 \mu\text{M}$.

Most particular compounds of the present invention exhibit an EC_{1.5x} concentration for production of full length SMN2 minigene mRNA $\leq 0.02 \mu\text{M}$.

10

Example	EC _{1.5x} minigene (nM)		Example	EC _{1.5x} minigene (nM)		Example	EC _{1.5x} minigene (nM)
1	3.5		14	4.1		27	39.9
2	3.8		15	4		28	5
3	3.2		16	1.1		29	0.3
4	1.8		17	6.4		30	3
5	0.6		18	3.6		31	6.7
6	2.8		19	10.2		32	1.6
7	3.7		20	4.3		33	0.5
8	0.3		21	9.6		34	0.9
9	0.1		22	0.9		35	4.7
10	6.4		23	3.4		36	5
11	1.4		24	0.4		37	4.4
12	1.2		25	0.5		38	0.3
13	5		26	327		39	0.9

Table 3. EC_{1.5x} concentrations for production of full length SMN2 minigene mRNA.

Assay 2

SMN protein assay in cultured cells

The SMN HTRF (homogeneous time resolved fluorescence) assay is used to quantify the level of SMN protein in SMA patient fibroblast cells treated with test compounds. Materials used
5 and respective sources are listed below in Table 4.

Material	Source
SMA Type 1 human cells	GM03813 (Coriell Institute)
Protease inhibitor cocktail	Roche Applied Science Catalog No. 11836145001
Anti-SMN d2	Blue cap Cisbio Catalog No. 63IDC002-SMN
Anti-SMN kryptate	Red cap Cisbio Catalog No. 63IDC002-SMN
SMN reconstitution buffer	Cisbio Catalog No. 63IDC002-SMN-Buffer
DMEM	Life Technologies (formerly Invitrogen) Catalog No. 11960-044
RIPA Lysis Buffer	20 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Thermo Scientific NP-40 Surfact-Amps Detergent Solution (Fisher Scientific, Pittsburgh/PA), 1% Sodium deoxycholate
Diluent Buffer	20 mM Tris-HCl pH 7.5, 150 mM NaCl
Envision Plate Reader	Perkin Elmer Model # 2103

Table 4. Materials and their respective sources used in the SMN protein assay in cultured cells.

Cells are thawed and cultured in DMEM-10% FBS for 72 hours. Cells are trypsinized, counted and re-suspended to a concentration of 25,000 cells/mL in DMEM-10% FBS. The cell
10 suspensions are plated at 5,000 cells per well in a 96 well microtiter plate and incubated for 3 to 5 hours. Test compounds are serially diluted 3.16-fold in 100% DMSO to generate a 7-point concentration curve. 1 μ L of test compound solution is transferred to cell-containing wells and cells are incubated for 48 hours in a cell culture incubator (37°C, 5% CO₂, 100% relative humidity). Triplicate samples are set up for each test compound concentration. After 48 hours,
15 the supernatant is removed from the wells and 25 μ L of the RIPA lysis buffer, containing protease inhibitors, is added to the wells and incubated with shaking at room temperature for 1 hour. 25 μ L of the diluent is added and then 35 μ L of the resulting lysate is transferred to a 384-

well plate, where each well contains 5 μ L of the antibody solution (1:100 dilution of anti-SMN d2 and anti-SMN kryptate in SMN reconstitution buffer). The plate is centrifuged for 1 minute to bring the solution to the bottom of the wells, then incubated overnight at room temperature. Fluorescence for each well of the plate at 665 nm and 620 nm is measured on an EnVision
5 multilabel plate reader (Perkin-Elmer).

The normalized fluorescence signal is calculated for each sample, Blank and vehicle control well by dividing the signal at 665 nm by the signal at 620 nm. Normalizing the signal accounts for possible fluorescence quenching due to the matrix effect of the lysate. The ΔF value (a measurement of SMN protein abundance as a percent value) for each sample well is
10 calculated by subtracting the normalized average fluorescence for the Blank control wells from the normalized fluorescence for each sample well, then dividing this difference by the normalized average fluorescence for the Blank control wells and multiplying the resulting value by 100. The ΔF value for each sample well represents the SMN protein abundance from test compound-treated samples. The ΔF value for each sample well is divided by the ΔF value for the
15 vehicle control wells to calculate the fold increase in SMN protein abundance relative to the vehicle control. Table 5 provides $EC_{1.5x}$ concentrations for SMN protein expression that was obtained from the 7-point concentration data generated according to the above procedure for particular compounds of the present invention.

Particular compounds of the present invention exhibit an $EC_{1.5x}$ concentration for SMN
20 protein expression $\leq 1 \mu M$.

More particular compounds of the present invention exhibit an $EC_{1.5x}$ concentration for SMN protein expression ≤ 100 nM.

Most particular compounds of the present invention exhibit an $EC_{1.5x}$ concentration for SMN protein expression ≤ 30 nM.

25 Table 6 provides the maximum fold increase of SMN protein that was obtained from the 7-point concentration data generated according to the above procedure for particular compounds of the present invention

Particular compounds of the present invention exhibit a maximum fold increase > 1.5 .

More particular compounds of the present invention exhibit a maximum fold increase > 1.7 .

30 Most particular compounds of the present invention exhibit a maximum fold increase > 1.8 .

Example	EC1.5x SMN protein (nM)		Example	EC1.5x SMN protein (nM)		Example	EC1.5x SMN protein (nM)
1	10.8		14	17.6		27	126.5
2	19.8		15	21.2		28	49.7
3	25.6		16	3		29	2.1
4	15.7		17	20.2		30	13.6
5	4.1		18	25		31	27.7
6	11		19	29.8		32	4
7	15.5		20	37		33	4
8	5.9		21	68.7		34	4.4
9	2.5		22	13.8		35	19.5
10	22.8		23	23.9		36	34.4
11	7		24	4.7		37	45
12	7.5		25	11.9		38	3.1
13	3		26	1230		39	15.8


Table 5. EC_{1.5x} concentrations for SMN protein expression.

Example	max. fold increase		Example	max. fold increase		Example	max. fold increase
1	1.84		14	1.86		27	1.57
2	1.76		15	1.94		28	1.72
3	1.81		16	1.83		29	1.81
4	1.76		17	1.98		30	1.84
5	1.71		18	1.75		31	1.65
6	1.84		19	1.83		32	1.88
7	1.76		20	1.72		33	1.82
8	1.85		21	1.54		34	1.89
9	1.92		22	1.69		35	1.79
10	1.95		23	1.63		36	1.77
11	1.9		24	1.77		37	1.87
12	1.77		25	1.79		38	1.85
13	1.91		26	1.52		39	1.81

Table 6. Maximum fold increase of SMN protein.

Claims

1. A compound of formula (I)



wherein

- 5 R¹ is hydrogen or C₁₋₇-alkyl;
- R² is hydrogen, cyano, C₁₋₇-alkyl, C₁₋₇-haloalkyl or C₃₋₈-cycloalkyl;
- R³ is hydrogen, C₁₋₇-alkyl, or C₃₋₈-cycloalkyl;
- 10 A is N-heterocycloalkyl or NR¹²R¹³, wherein N-heterocycloalkyl comprises 1 or 2
 nitrogen ring atoms and is optionally substituted with 1, 2, 3 or 4 substituents
 selected from R¹⁴;
- R¹² is heterocycloalkyl comprising 1 nitrogen ring atom, wherein heterocycloalkyl is
 optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴;
- R¹³ is hydrogen, C₁₋₇-alkyl or C₃₋₈-cycloalkyl;
- 15 two R¹⁴ together form C₁₋₇-alkylene, or R¹⁴ is independently selected from the group
 consisting of hydrogen, C₁₋₇-alkyl, amino, amino-C₁₋₇-alkyl, C₃₋₈-cycloalkyl and
 heterocycloalkyl;
- with the proviso that if A is N-heterocycloalkyl comprising only 1 nitrogen ring atom,
 then at least one R¹⁴ substituent is amino or amino-C₁₋₇-alkyl;

or a pharmaceutically acceptable salt thereof.

2. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

R¹ is hydrogen or C₁₋₇-alkyl;

R² is hydrogen, cyano, C₁₋₇-alkyl, C₁₋₇-haloalkyl or C₃₋₈-cycloalkyl;

5 R³ is hydrogen, C₁₋₇-alkyl, or C₃₋₈-cycloalkyl;

A is N-heterocycloalkyl comprising 1 or 2 nitrogen ring atoms, wherein N-heterocycloalkyl is optionally substituted with 1, 2, 3 or 4 substituents selected from R¹⁴;

10 two R¹⁴ together form C₁₋₇-alkylene, or R¹⁴ is independently selected from the group consisting of hydrogen, C₁₋₇-alkyl, amino, amino-C₁₋₇-alkyl, C₃₋₈-cycloalkyl and heterocycloalkyl;

with the proviso that if A is N-heterocycloalkyl comprising only 1 nitrogen ring atom, then at least one R¹⁴ substituent is amino or amino-C₁₋₇-alkyl.

- 15 3. The compound or pharmaceutically acceptable salt thereof according to claim 1 or 2, wherein R¹ is C₁₋₇-alkyl.

4. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 3, wherein R¹ is methyl.


5. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein R² is hydrogen or C₁₋₇-alkyl.

- 20 6. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 5, wherein R² is hydrogen or methyl.

7. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 6, wherein R³ is hydrogen or C₁₋₇-alkyl.

- 25 8. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 7, wherein R³ is hydrogen or methyl.

9. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 8, wherein R^{12} is piperidinyl optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} .
10. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 9, wherein R^{13} is hydrogen or C_{1-7} -alkyl.
11. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 10, wherein R^{13} is hydrogen or methyl.
12. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 11, wherein two R^{14} together form C_{1-7} -alkylene, or R^{14} is independently selected from the group consisting of C_{1-7} -alkyl and heterocycloalkyl.
13. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 12, wherein two R^{14} together form ethylene, or R^{14} is independently selected from the group consisting of methyl, ethyl and pyrrolidinyl.
14. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 13, wherein the N-heterocycloalkyl in A or the heterocycloalkyl in R^{12} as defined in Claim 1 are further characterized in that one ring nitrogen atom is basic.
15. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 14, wherein



X is N or CH;

R^4 is hydrogen, C_{1-7} -alkyl or $-(CH_2)_m-NR^9R^{10}$;

R^5 is hydrogen or C_{1-7} -alkyl;

R^6 is hydrogen or C_{1-7} -alkyl;

R^7 is hydrogen or C_{1-7} -alkyl;

R^8 is hydrogen or C_{1-7} -alkyl;

5 R^9 and R^{10} are independently selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{3-8} -cycloalkyl;

R^{13} is hydrogen, C_{1-7} -alkyl or C_{3-8} -cycloalkyl;

n is 0, 1 or 2;

m is 0, 1, 2 or 3;

or R^4 and R^5 together form C_{1-7} -alkylene;

10 or R^4 and R^7 together form C_{1-7} -alkylene;

or R^5 and R^6 together form C_{2-7} -alkylene;

or R^5 and R^7 together form C_{1-7} -alkylene;

or R^5 and R^9 together form C_{1-7} -alkylene;

or R^7 and R^8 together form C_{2-7} -alkylene;


15 or R^7 and R^9 together form C_{1-7} -alkylene;

or R^9 and R^{10} together form C_{2-7} -alkylene;

with the proviso that if X is CH then R^4 is $-(CH_2)_m-NR^9R^{10}$; and

with the proviso that if X is N and R^4 is $-(CH_2)_m-NR^9R^{10}$ then m is 2 or 3.

20 16. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 15, wherein



X is N or CH;

R⁴ is hydrogen, C₁₋₇-alkyl or -(CH₂)_m-NR⁹R¹⁰;

R⁵ is hydrogen or C₁₋₇-alkyl;

5 R⁶ is hydrogen or C₁₋₇-alkyl;

R⁷ is hydrogen or C₁₋₇-alkyl;

R⁸ is hydrogen or C₁₋₇-alkyl;

R⁹ and R¹⁰ are independently selected from the group consisting of hydrogen, C₁₋₇-alkyl and C₃₋₈-cycloalkyl;

10 n is 0, 1 or 2;

m is 0, 1, 2 or 3;

or R⁴ and R⁵ together form C₁₋₇-alkylene;

or R⁴ and R⁷ together form C₁₋₇-alkylene;

or R⁵ and R⁶ together form C₂₋₇-alkylene;

15 or R⁵ and R⁷ together form C₁₋₇-alkylene;

or R⁵ and R⁹ together form C₁₋₇-alkylene;

or R⁷ and R⁸ together form C₂₋₇-alkylene;

or R⁷ and R⁹ together form C₁₋₇-alkylene;


or R⁹ and R¹⁰ together form C₂₋₇-alkylene;

with the proviso that if X is CH then R⁴ is -(CH₂)_m-NR⁹R¹⁰; and


with the proviso that if X is N and R⁴ is -(CH₂)_m-NR⁹R¹⁰ then m is 2 or 3.

- 5 17. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 16, wherein X is N.
18. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 17, wherein n is 1.
19. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 18, wherein R⁴ is hydrogen, methyl or -(CH₂)_m-NR⁹R¹⁰.
- 10 20. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 19, wherein R⁴ is hydrogen.
21. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 20, wherein R⁵ is hydrogen, methyl or ethyl.
- 15 22. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 21, wherein R⁵ is methyl.
23. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 22, wherein R⁶ is hydrogen or methyl.
24. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 23, wherein R⁶ is hydrogen.
- 20 25. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 24, wherein R⁷ is hydrogen or methyl.
26. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 25, wherein R⁸ is hydrogen.
- 25 27. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 19, wherein m is 0.

28. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 18, wherein R^4 and R^5 together form propylene.
29. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 18, wherein R^5 and R^6 together form ethylene.
- 5 30. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 18, wherein R^9 and R^{10} together form butylene.
31. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 30, wherein A is selected from the group consisting of:



32. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 31, wherein A is selected from the group consisting of:




wherein R^4 , R^5 , R^6 , R^7 and R^8 are as defined in any one of claims 1 to 31 and wherein R^{11} is hydrogen or C_{1-7} -alkyl.

33. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 32, wherein A is selected from the group consisting of piperazinyl, diazepanyl, pyrrolidinyl and hexahydropyrrolo[1,2-a]pyrazinyl, each optionally substituted with 1, 2, 3 or 4 substituents selected from R^{14} as defined in any one of claims 1 to 32.
34. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 33, wherein A is selected from the group consisting of piperazin-1-yl, 1,4-diazepan-1-yl, pyrrolidin-1-yl and hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, each optionally substituted with 1 or 2 substituents selected from R^{14} as defined in any one of claims 1 to 33.


35. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 31, wherein A is $\text{NR}^{12}\text{R}^{13}$, wherein R^{12} and R^{13} are as defined in any one of claims 1 to 31.

36. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 31, wherein




A is , wherein R^4 , R^5 , R^6 , R^7 , R^8 and R^{13} are as defined in any one of claims 1 to 31.

37. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 34, wherein A is selected from the group consisting of:



38. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 34, wherein A is selected from the group consisting of:



39. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 38, selected from the group consisting of:

2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-(4-methylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

- 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 5 7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5R)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 10 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 7-(1,4-diazepan-1-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 15 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 7-(1,4-diazepan-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 20 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 25 7-[(8aS)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-[(8aR)-8a-methyl-1,3,4,6,7,8-hexahydropyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
- 30 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
- 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-

- yl)pyrido[1,2-a]pyrimidin-4-one;
2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)pyrido[1,2-a]pyrimidin-4-one;
5 7-(3,3-dimethylpiperazin-1-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
10 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-(3,3-dimethylpiperazin-1-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
15 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
20 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-pyrrolidin-1-ylpyrrolidin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
25 9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)-7-[(3R)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
30 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-(3,3-dimethylpiperazin-1-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

7-(4,7-diazaspiro[2.5]octan-7-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-[(3S,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
5 7-[(3R)-3-ethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
and a pharmaceutically acceptable salt thereof.

40. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 39, selected from the group consisting of:

- 10 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
15 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-[(3S,5R)-3,5-dimethylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
20 7-[(8aS)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
25 2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-7-[(3S)-3-methylpiperazin-1-yl]pyrido[1,2-a]pyrimidin-4-one;
7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methyl-pyrido[1,2-a]pyrimidin-4-one;
30 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;
7-(4,7-diazaspiro[2.5]octan-7-yl)-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-

yl)pyrido[1,2-a]pyrimidin-4-one;
and a pharmaceutically acceptable salt thereof.


41. The compound according to claim 1 or 2, wherein the compound is 7-[(8aR)-3,4,6,7,8,8a-hexahydro-1H-pyrrolo[1,2-a]pyrazin-2-yl]-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.

42. The compound according to claim 1 or 2, wherein the compound is 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.

43. The compound according to claim 1 or 2, wherein the compound is 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-9-methylpyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.


44. The compound according to claim 1 or 2, wherein the compound is 7-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.

45. A compound which is




7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one, or a pharmaceutically acceptable salt thereof.

46. A compound which is



7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one.

47. A compound of formula (VI)



(VI)

wherein R¹, R² and R³ are as defined in any one of claims 1 to 7;

Y is halogen or trifluoromethanesulfonate;

or a salt thereof.

48. The compound of formula (VI) or salt thereof according to claim 47, wherein Y is fluoro, chloro, bromo, iodo or trifluoromethanesulfonate.

49. The compound of formula (VI) or salt thereof according to claim 47, wherein Y is fluoro.

50. The compound of formula (VI) or salt thereof according to any one of claims 47 to 49, selected from the group consisting of:

7-fluoro-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-pyrido[1,2-a]pyrimidin-4-one;

7-fluoro-9-methyl-2-(2-methylimidazo[1,2-b]pyridazin-6-yl)pyrido[1,2-a]pyrimidin-4-one;

2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)-7-fluoro-9-methyl-pyrido[1,2-a]pyrimidin-4-one;

and a salt thereof.

51. A process for the preparation of a compound as defined in any one of claims 1 to 46, comprising an aromatic nucleophilic substitution reaction between a compound of formula (VI) as defined in any one of claims 47 to 50 with a compound of formula M-A by heating in a solvent, wherein A, R¹, R² and R³ are as defined in any one of claims 1 to 46, Y is as defined in any one of claims 47 to 50, M is hydrogen, sodium or potassium, and wherein M is linked to A via a nitrogen atom of A.
52. The process according to claim 51, wherein the aromatic nucleophilic substitution reaction is performed at a temperature from 80°C to 200°C.
53. The process according to claim 51 or 52, wherein the solvent of the aromatic nucleophilic substitution reaction is selected from the group consisting of dimethyl sulfoxide, N-methylpyrrolidone, and dimethylformamide.
54. The process according to any one of claims 51 to 53, wherein M is hydrogen.
55. A compound as defined in any one of claims 1 to 46, obtained by a process according to any one of claims 51 to 54.
56. A pharmaceutical composition comprising a compound as defined in any one of claims 1 to 46 or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.
57. A compound as defined in any one of claims 1 to 46 or a pharmaceutically acceptable salt thereof for use as therapeutically active substance.
58. A compound as defined in any one of claims 1 to 46 or a pharmaceutically acceptable salt thereof for use in the treatment or prevention of spinal muscular atrophy (SMA).
59. A use of a compound as defined in any one of claims 1 to 46 or a pharmaceutically acceptable salt thereof for the treatment or prevention of spinal muscular atrophy (SMA).
60. A use of a compound as defined in any one of claims 1 to 46 or a pharmaceutically acceptable salt thereof in the preparation of a medicament for the treatment or prevention of spinal muscular atrophy (SMA).

61. The compound or a pharmaceutically acceptable salt thereof for use of claim 58, in the treatment of SMA.
62. The use of claim 59 or 60, for the treatment of SMA.