

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 August 2008 (28.08.2008)

PCT

(10) International Publication Number
WO 2008/102198 A1

(51) International Patent Classification:
C12N 9/22 (2006.01)

(21) International Application Number:
PCT/IB2007/001527

(22) International Filing Date:
19 February 2007 (19.02.2007)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **CEL-LECTIS** [FR/FR]; 102 route de Noisy, F-93235 Romainville Cedex (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MONTOYA, Guillermo** [ES/ES]; Alejandro Rubio 93, Guadalix de la Sierra, E-28794 Madrid (ES). **BLANCO, Francisco** [ES/ES]; Marqués de Corbera 62, 8°-A, E-28017 Madrid (ES). **PRIETO, Jesus** [ES/ES]; Calle José Bergamin n° 38, 3°-C, E-28030 Madrid (ES).

(74) Agent: **CABINET ORES**; 36 rue de St Pétersbourg, F-75008 Paris (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: LAGLIDADG HOMING ENDONUCLEASE VARIANTS HAVING NOVEL SUBSTRATE SPECIFICITY AND USE THEREOF




Figure 1

(57) Abstract: A LAGLIDADG homing endonuclease variant having novel substrate specificity, said variant being obtainable by a method comprising : (a) the mutation of at least one amino acid residue of the final C-terminal loop of a parent LAGLIDADG homing endonuclease, with the exclusion of the threonine 140 of I-CreI, b) the selection and/or screening of the variants from step (a) having a pattern of cleaved DNA targets that is different from that of the parent LAGLIDADG homing endonuclease.

LAGLIDADG HOMING ENDONUCLEASE VARIANTS HAVING NOVEL SUBSTRATE SPECIFICITY AND USE THEREOF

The invention relates to a method for engineering LAGLIDADG homing endonuclease variants having novel substrate specificity. The invention relates also to a variant obtainable by said method, to a vector encoding said variant, to a cell, an animal or a plant modified by said vector and to the use of said homing endonuclease variant and derived products for genetic engineering, genome therapy and antiviral therapy.

Meganucleases are by definition sequence-specific endonucleases with large (12-45 bp) cleavage sites that can deliver DNA double-strand breaks (DSBs) at specific loci in living cells (Thierry, A. and Dujon B., *Nucleic Acids Res.*, 1992, 20, 5625-5631). Meganucleases have been used to stimulate homologous recombination in the vicinity of their target sequences in cultured cells and plants (Rouet *et al.*, *Mol. Cell. Biol.*, 1994, 14, 8096-8106; Choulika *et al.*, *Mol. Cell. Biol.*, 1995, 15, 1968-1973; Donoho *et al.*, *Mol. Cell. Biol.*, 1998, 18, 4070-4078; Elliott *et al.*, *Mol. Cell. Biol.*, 1998, 18, 93-101; Sargent *et al.*, *Mol. Cell. Biol.*, 1997, 17, 267-277; Puchta *et al.*, *Proc. Natl. Acad. Sci. USA*, 1996, 93, 5055-5060; Chiurazzi *et al.*, *Plant Cell*, 1996, 8, 2057-2066), making meganuclease-induced recombination an efficient and robust method for genome engineering.

The use of meganuclease-induced recombination has long been limited by the repertoire of natural meganucleases, and the major limitation of the current technology is the requirement for the prior introduction of a meganuclease cleavage site in the locus of interest. Thus, the engineering of redesigned meganucleases cleaving chosen targets is under intense investigation.

Such proteins could be used to cleave genuine chromosomal sequences and open new perspectives for genome engineering in wide range of applications. For example, meganucleases could be used to knock-out endogenous genes or knock-in exogenous sequences in the chromosome. It can as well be used for the precise *in situ* correction of mutations linked with monogenic diseases and thereby bypass the risk due to the randomly inserted transgenes encountered with current gene therapy approaches (Hacein-Bey-Abina *et al.*, *Science*, 2003, 302, 415-419).

Recently, Zinc-Finger DNA binding domains of Cys2-His2 type Zinc-Finger Proteins (ZFPs) were fused with the catalytic domain of the *FokI* endonuclease, to induce recombination in various cell types: mammalian cultured cells including human lymphoid cells, plants and insects (Smith *et al.*, Nucleic Acids Res, 1999, 27, 674-81; Pabo *et al.*, Annu. Rev. Biochem, 2001, 70, 313-40; Porteus, M.H. and Baltimore, D., Science, 2003, 300, 763; Urnov *et al.*, Nature, 2005, 435, 646-651; Bibikova *et al.*, Science, 2003, 300, 764; Durai *et al.*, Nucleic Acids Res., 2005, 33, 5978-5990; Porteus M.H., Mol. Ther., 2006, 13, 438-446). The binding specificity of ZFPs is relatively easy to manipulate, and a repertoire of novel artificial ZFPs, able to bind many (g/a)nn(g/a)nn(g/a)nn sequences is now available (Pabo *et al.*, precited; Segal, D.J. and Barbas, C.F., Curr. Opin. Biotechnol., 2001, 12, 632-637; Isalan *et al.*, Nat. Biotechnol., 2001, 19, 656-660). However, preserving a very narrow specificity is one of the major issues for genome engineering applications, and presently it is unclear whether ZFPs would fulfill the very strict requirements for therapeutic applications. Furthermore, these fusion proteins have demonstrated high toxicity in *Drosophila* (Bibikova *et al.*, Science, 2003, 300, 764; Bibikova *et al.*, Genetics, 2002, 161, 1169-1175) and mammalian NIH3T3 cells (Alwin *et al.*, Mol. Ther., 2005, 12, 610-617; Porteus, M.H. and Baltimore, D., Science, 2003, 300, 763; Porteus, M.H. and Carroll, D., Nat. Biotechnol., 2005, 23, 967-973), a genotoxic effect that is probably due to frequent off-site cleavage (Porteus, M.H., Mol. Ther., 2006, 13, 438-446).

In nature, meganucleases are essentially represented by homing endonucleases (HEs), a family of endonucleases encoded by mobile genetic elements, whose function is to initiate DNA double-strand break (DSB)-induced recombination events in a process referred to as homing (Chevalier, B.S. and Stoddard, B.L., Nucleic Acids Res., 2001, 29, 3757-3774; Kostriken *et al.*, Cell, 1983, 35, 167-174; Jacquier, A. and Dujon, B., Cell, 1985, 41, 383-394). Several hundreds of HEs have been identified in bacteria, eukaryotes, and archaea (Chevalier, B.S. and Stoddard, B.L., Nucleic Acids Res., 2001, 29, 3757-3774); however the probability of finding a HE cleavage site in a chosen gene is very low.

Given their biological function and their exceptional cleavage properties in terms of efficacy and specificity, HEs provide ideal scaffolds to derive

novel endonucleases for genome engineering. Furthermore, in addition to their exquisite specificity, homing endonuclease have shown to be less toxic than ZFPs, probably because of better specificity (Alwin *et al.*, Mol. Ther., 2005, 12, 610-617; Porteus, M.H. and Baltimore, D., Science, 2003, 300, 763; Porteus, M.H. and Carroll, D., Nat. Biotechnol., 2005, 23, 967-973), two features that become essential when engaging into therapeutic applications.

Data have accumulated over the last decade, allowing a relatively good characterization of the LAGLIDADG family, the largest of the four HE families (Chevalier, B.S. and Stoddard, B.L., Nucleic Acids Res., 2001, 29, 3757-74).

LAGLIDADG refers to the only sequence actually conserved throughout the family, and is found in one or (more often) two copies in the protein. Proteins with a single motif, such as I-CreI (Wang *et al.*, Nucleic Acids Res., 1997, 25, 3767-3776) form homodimers and cleave palindromic or pseudo-palindromic DNA sequences, whereas the larger, double motif proteins, such as I-SceI (Jacquier, A. and Dujon, B., Cell., 1985, 41, 383-394) or I-DmoI (Dalgaard *et al.*, Proc. Natl. Acad. Sci. USA, 1993, 90, 5417-5417) are monomers and cleave non palindromic targets. Nine different LAGLIDADG proteins have been crystallized with or without bound DNA, showing a very striking core structure conservation that contrasts with the lack of similarity at the primary sequence level (Heath *et al.*, Nature Struct. Biol., 1997, 4, 468-476; Duan *et al.*, Cell., 1997, 89, 555-564; Silva *et al.*, J. Mol. Biol., 2003, 286, 1123-1136; Chevalier *et al.*, Nat. Struct. Biol., 2001, 8, 312-316; Jurica *et al.*, Mol. Cell., 1998, 2, 469-476; Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269; Moure *et al.*, J. Mol. Biol., 2003, 334, 685-695; Moure *et al.*, Nat. Struct. Biol., 2002, 9, 764-770; Ichiyanagi *et al.*, J. Mol. Biol., 2000, 300, 889-901; Gimble *et al.*, J. Biol. Chem., 1998, 273, 30524-30529; Bolduc *et al.*, Genes Dev. 2003, 17, 2875-2888; Silva *et al.*, J. Mol. Biol., 1999, 286, 1123-1136; Nakayama *et al.*, J. Mol. Biol., Epub 29 septembre 2006, Spiegel *et al.*, Structure, 2006, 14, 869-880). In contrast with its DNA bound crystal structure (Jurica *et al.*, Mol. Cell., 1998, 2, 469-476; Chevalier *et al.*, Nat. Struct. Biol., 2001, 8, 312-316; Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269), the structure of I-CreI without bound DNA (Heath *et al.*, Nature Struct. Biol., 1997, 4, 468-476), showed only one I-CreI monomer in the asymmetric unit.

Structural comparisons indicate that LAGLIDADG proteins adopt a similar active conformation and their self-association forms two packed α -helices which separate two monomers or apparent domains. In this core structure (Figure 1), two characteristic $\alpha\beta\beta\alpha\beta\beta\alpha$ folds, contributed by two monomers, or two domains in double LAGLIDADG proteins, are facing each other with a two-fold symmetry. On either side of the LAGLIDADG α -helices, a four stranded β -sheet, forming a saddle on the DNA helix major groove, provides a DNA binding interface that drives the interaction of the protein with a half site of the target DNA sequence (Chevalier *et al.*, Nat. Struct. Biol., 2001, 8, 312-316; Jurica *et al.*, Mol. Cell., 1998, 2, 469-476). The catalytic site is central, formed with contributions from helices of both monomers. Just above the catalytic site, the two LAGLIDADG α -helices play also an essential role in the dimerization interface. In addition to this core structure, other domains can be found, for instance, PI-SceI, an intein, has a protein splicing domain, and an additional DNA-binding domain (Moure *et al.*, Nat. Struct. Biol., 2002, 9, 764-770; Pingoud *et al.*, Biochemistry, 1998, 37, 8233-8243).

Modifying the substrate specificity of DNA binding proteins by mutagenesis and screening/selection is a difficult task (Lanio *et al.*, Protein Eng., 2000, 13, 275-281; Voziyanov *et al.*, J. Mol. Biol., 2003, 326, 65-76; Santoro *et al.*, P.N.A.S., 2002, 99, 4185-4190; Buchholz, F. and Stewart, A.F., Nat. Biotechnol., 2001, 19, 1047-1052). This is even harder in the case of HEs whose main characteristic is their large DNA recognition sites.

Analysis of the I-CreI/DNA crystal structure indicates that in each monomer, nine residues (S32, Y33, Q38, N30, K28, Q26, Q44, R68 and R70) establish direct interaction with eight bases at positions $\pm 3, 4, 5, 6, 7, 9, 10$ and 11 of the homing site (Jurica *et al.*, Mol. Cell., 1998, 2, 469-76; Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269), which randomization would result in 20^9 combinations, a number beyond any screening capacity today. In addition, a total of 28 (Left-monomer) or 24 (Right-monomer) water molecules mediate additional contacts between nucleotides and protein side-chains in the protein/DNA interface (Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269).

Therefore, several laboratories have relied on a semi-rational approach (Chica *et al.*, Curr. Opin. Biotechnol., 2005, 16, 378-384) to limit the

diversity of the mutant libraries to be handled, choosing a small set of relevant amino acid residues according to structural data. This set is generally composed of amino acid residues of the four stranded β -sheet that, in the HE/DNA complex structure, make direct or water-mediated contacts with the nucleotide bases of the homing site.

5 This semi-rational approach was used to locally alter the specificity of the I-CreI (Seligman *et al.*, Genetics, 1997, 147, 1653-64; Seligman *et al.*, Nucleic Acids Res., 2002, 30, 3870-3879; Sussman *et al.*, J. Mol. Biol., 2004, 342, 31-41; Rosen *et al.*, Nucleic Acids Res., 2006, 34, 4791-4800; Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; International PCT Applications WO 2006/097853 and WO
10 2006/097784; Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006), I-SceI (Doyon *et al.*, J. Am. Chem. Soc., 2006, 128, 2477-2484), PI-SceI (Gimble *et al.*, J. Mol. Biol., 2003, 334, 993-1008) and I-MsoI (Ashworth *et al.*, Nature, 2006, 441, 656-659) proteins.

By combining the semi-rational approach and high throughput
15 screening (HTS; Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; International PCT Applications WO 2006/097853 and WO 2006/097784; Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006), it was possible to obtain large number of locally altered variants of the I-CreI meganuclease that recognize a variety of targets, and to assemble them by a combinatorial process, to obtain entirely redesigned mutants with
20 chosen specificity.

However, this approach is not easy since the HEs DNA binding interface is very compact and the two different $\beta\beta$ hairpins which are responsible for virtually all base-specific interactions are part of a single fold. Thus, the mutation of several amino acids placed in close vicinity which is required for binding a target
25 mutated at several positions may disrupt the structure of the binding interface.

Therefore, to reach a larger number of sequences, it would be extremely valuable to be able to identify other regions in the LAGLIDAG endonucleases, which can be engineered to generate novel substrate specificity.

In addition, since homing endonucleases can sometimes be harmful
30 at very high doses (Gouble *et al.*, J. Gene Med., 2006, 8, 616-622), it would be extremely valuable to engineer LAGLIDAG endonucleases which are less toxic.

The inventors have solved the structure of the I-CreI dimer without DNA; its comparison with the DNA bound crystal structure (PDB code 1gz9; Chevalier *et al.*, Nat. Struct. Biol., 2001, 8, 312-316) depicts a different conformation of the C-terminal loop and the final helix α_6 , which suggests its implication in DNA binding. A site-directed mutagenesis study in this region demonstrates that whereas the C-terminal helix is negligible for DNA binding, the final C-terminal loop which is well conserved among homodimeric proteins from the LAGLIDADG family (Figure 2) and makes a number of nonspecific contacts to the DNA phosphate backbone (Jurica *et al.*, Mol. Cell., 1998, 2, 469-76; Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269), plays an important role not only in binding and cleavage but also in target specificity. In addition, some of the mutants in the C-terminal loop were significantly less toxic than wild-type I-CreI.

This region open new possibilities to engineer new homing endonucleases having novel substrate specificities and thereby increases the number of DNA sequences that can be targeted with meganucleases. Thus, redesigned meganucleases cleaving chosen genomic targets from genes of interest can be engineered by combining previously identified mutations as defined above (Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; International PCT Applications WO 2006/097853, WO 2006/097854 and WO 2006/097784; Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006), with mutations in the final C-terminal loop.

In addition, this region allows also the engineering of homing endonucleases which are less toxic.

Potential applications include genetic engineering, genome engineering, gene therapy and antiviral therapy.

The invention relates to a method for engineering a LAGLIDADG homing endonuclease variant having novel substrate specificity, comprising at least the following steps:

(a) the mutation of at least one amino acid residue of the final C-terminal loop of a parent LAGLIDADG homing endonuclease, with the exclusion of the threonine 140 (T140) of I-CreI, and

(b) the selection and/or screening of the variants from step (a) having a pattern of cleaved DNA targets that is different from that of the parent LAGLIDADG homing endonuclease.

Definitions

5 - Amino acid residues in a polypeptide sequence are designated herein according to the one-letter code, in which, for example, Q means Gln or Glutamine residue, R means Arg or Arginine residue and D means Asp or Aspartic acid residue.

 - hydrophobic amino acid refers to leucine (L), valine (V), isoleucine
10 (I), alanine (A), methionine (M), phenylalanine (F), tryptophane (W) and tyrosine (Y).

 - Nucleotides are designated as follows: one-letter code is used for designating the base of a nucleoside: a is adenine, t is thymine, c is cytosine, and g is guanine. For the degenerated nucleotides, r represents g or a (purine nucleotides), k represents g or t, s represents g or c, w represents a or t, m represents a or c, y represents t or c (pyrimidine nucleotides), d represents g, a or t, v represents g, a or c, b
15 represents g, t or c, h represents a, t or c, and n represents g, a, t or c.

 - by "meganuclease" is intended an endonuclease having a double-stranded DNA target sequence of 12 to 45 pb.

 - by "parent LAGLIDADG homing endonuclease" is intended a wild-
20 type LAGLIDADG homing endonuclease or a functional variant thereof. Said parent LAGLIDADG homing endonuclease may be a monomer, a dimer (homodimer or heterodimer) comprising two LAGLIDADG homing endonuclease core domains which are associated in a functional endonuclease able to cleave a double-stranded DNA target of 22 to 24 bp.

25 - by "homodimeric LAGLIDADG homing endonuclease" is intended a wild-type homodimeric LAGLIDADG homing endonuclease having a single LAGLIDADG motif and cleaving palindromic DNA target sequences, such as I-CreI or I-MsoI or a functional variant thereof.

 - by "LAGLIDADG homing endonuclease variant" or "variant" is
30 intended a protein obtained by replacing at least one amino acid of a LAGLIDADG homing endonuclease sequence, with a different amino acid.

- by "functional variant" is intended a LAGLIDADG homing endonuclease variant which is able to cleave a DNA target, preferably a new DNA target which is not cleaved by a wild-type LAGLIDADG homing endonuclease. For example, such variants have amino acid variation at positions contacting the DNA target sequence or interacting directly or indirectly with said DNA target.

- by "homing endonuclease variant with novel specificity" is intended a variant having a pattern of cleaved targets (cleavage profile) different from that of the parent homing endonuclease. The variants may cleave less targets (restricted profile) or more targets than the parent homing endonuclease. Preferably, the variant is able to cleave at least one target that is not cleaved by the parent homing endonuclease.

The terms "novel specificity", "modified specificity", "novel cleavage specificity", "novel substrate specificity" which are equivalent and used indifferently, refer to the specificity of the variant towards the nucleotides of the DNA target sequence.

- by "I-CreI" is intended the wild-type I-CreI having the sequence SWISSPROT P05725 or pdb accession code 1g9y.

- by "domain" or "core domain" is intended the "LAGLIDADG homing endonuclease core domain" which is the characteristic $\alpha_1\beta_1\beta_2\alpha_2\beta_3\beta_4\alpha_3$ fold of the homing endonucleases of the LAGLIDADG family, corresponding to a sequence of about one hundred amino acid residues. Said domain comprises four beta-strands ($\beta_1, \beta_2, \beta_3, \beta_4$) folded in an antiparallel beta-sheet which interacts with one half of the DNA target. This domain is able to associate with another LAGLIDADG homing endonuclease core domain which interacts with the other half of the DNA target to form a functional endonuclease able to cleave said DNA target. For example, in the case of the dimeric homing endonuclease I-CreI (163 amino acids), the LAGLIDADG homing endonuclease core domain corresponds to the residues 6 to 94. In the case of monomeric homing endonucleases, two such domains are found in the sequence of the endonuclease; for example in I-DmoI (194 amino acids), the first domain (residues 7 to 99) and the second domain (residues 104 to 194) are separated by a short linker (residues 100 to 103).

- by "subdomain" is intended the region of a LAGLIDADG homing endonuclease core domain which interacts with a distinct part of a homing endonuclease DNA target half-site. Two different subdomains behave independently and the mutation in one subdomain does not alter the binding and cleavage properties of the other subdomain. Therefore, two subdomains bind distinct part of a homing endonuclease DNA target half-site.

- by "beta-hairpin" is intended two consecutive beta-strands of the antiparallel beta-sheet of a LAGLIDADG homing endonuclease core domain ($\beta_1\beta_2$ or, $\beta_3\beta_4$) which are connected by a loop or a turn,

- by "DNA target", "DNA target sequence", "target sequence", "target-site", "target", "site"; "recognition site", "recognition sequence", "homing recognition site", "homing site", "cleavage site" is intended a 22 to 24 bp double-stranded palindromic, partially palindromic (pseudo-palindromic) or non-palindromic polynucleotide sequence that is recognized and cleaved by a LAGLIDADG homing endonuclease. These terms refer to a distinct DNA location, preferably a genomic location, at which a double stranded break (cleavage) is to be induced by the endonuclease. The DNA target is defined by the 5' to 3' sequence of one strand of the double-stranded polynucleotide. For example, the palindromic DNA target sequence cleaved by wild-type I-CreI presented in figure 8 is defined by the sequence 5'- t₁₂c₁₁a₁₀a₉a₈a₇c₆g₅t₄c₃g₂t₁a₊1c₊2g₊3a₊4c₊5g₊6t₊7t₊8t₊9t₊10g₊11a₊12 (SEQ ID NO :1). Cleavage of the DNA target occurs at the nucleotides in positions +2 and -2, respectively for the sense and the antisense strand. Unless otherwise indicated, the position at which cleavage of the DNA target by an I-Cre I meganuclease variant occurs, corresponds to the cleavage site on the sense strand of the DNA target.

- by "DNA target half-site", "half cleavage site" or "half-site" is intended the portion of the DNA target which is bound by each LAGLIDADG homing endonuclease core domain.

- by "chimeric DNA target" or "hybrid DNA target" is intended the fusion of a different half of two parent meganuclease target sequences. In addition at least one half of said target may comprise the combination of nucleotides which are bound by separate subdomains (combined DNA target).

- by "vector" is intended a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked.

- by "mutation" is intended the substitution, the deletion, and/or the addition of one or more nucleotides/amino acids in a nucleic acid/amino acid sequence.

- by "homologous" is intended a sequence with enough identity to another one to lead to a homologous recombination between sequences, more particularly having at least 95 % identity, preferably 97 % identity and more preferably 99 %.

- "Identity" refers to sequence identity between two nucleic acid molecules or polypeptides. Identity can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base, then the molecules are identical at that position. A degree of similarity or identity between nucleic acid or amino acid sequences is a function of the number of identical or matching nucleotides at positions shared by the nucleic acid sequences. Various alignment algorithms and/or programs may be used to calculate the identity between two sequences, including FASTA, or BLAST which are available as a part of the GCG sequence analysis package (University of Wisconsin, Madison, Wis.), and can be used with, e.g., default settings.

- "individual" includes mammals, as well as other vertebrates (e.g., birds, fish and reptiles). The terms "mammal" and "mammalian", as used herein, refer to any vertebrate animal, including monotremes, marsupials and placental, that suckle their young and either give birth to living young (eutharian or placental mammals) or are egg-laying (metatharian or nonplacental mammals). Examples of mammalian species include humans and other primates (e.g., monkeys, chimpanzees), rodents (e.g., rats, mice, guinea pigs) and ruminants (e.g., cows, pigs, horses).

- "genetic disease" refers to any disease, partially or completely, directly or indirectly, due to an abnormality in one or several genes. Said abnormality can be a mutation, an insertion or a deletion. Said mutation can be a punctual mutation. Said abnormality can affect the coding sequence of the gene or its regulatory sequence. Said abnormality can affect the structure of the genomic sequence or the structure or stability of the encoded mRNA. Said genetic disease can be recessive or

dominant. Such genetic disease could be, but are not limited to, cystic fibrosis, Huntington's chorea, familial hypercholesterolemia (LDL receptor defect), hepatoblastoma, Wilson's disease, congenital hepatic porphyrias, inherited disorders of hepatic metabolism, Lesch Nyhan syndrome, sickle cell anemia, thalassaemias, xeroderma pigmentosum, Fanconi's anemia, retinitis pigmentosa, ataxia telangiectasia, Bloom's syndrome, retinoblastoma, Duchenne's muscular dystrophy, and Tay-Sachs disease.

According to the invention, the amino acids of the final C-terminal loop correspond to positions 137 to 143 in I-*CreI* amino acid sequence SEQ ID NO: 2 or Swissprot P05725. Knowing the positions of the final C-terminal loop in I-*CreI*, one skilled in the art can easily deduce the corresponding positions in another homodimeric LAGLIDADG homing endonuclease, using well-known protein structure analyses softwares such as Pymol. For example, for I-*MsoI*, the final C-terminal loop corresponds to positions 143 to 149. According to an advantageous embodiment of said method, step (a) comprises the mutation of amino acid residue(s) of the final C-terminal loop that are contacting the phosphate backbone of the parent LAGLIDADG endonuclease DNA cleavage site (wild-type LAGLIDADG endonuclease homing site). Preferably said residues are involved in binding and cleavage of said DNA cleavage site. More preferably, said residues are in positions 138, 139, 142 or 143, by reference to the numbering of I-*CreI* amino acid sequence (SEQ ID NO: 2; figure 2). Two residues may be mutated in one variant provided that each mutation is in a different pair of residues chosen from the pair of residues in positions 138 and 139 and the pair of residues in positions 142 and 143.

According to the method of the invention, the mutations which are introduced modify the interaction(s) of said amino acid(s) of the final C-terminal loop with the phosphate backbone of the parent LAGLIDADG endonuclease DNA cleavage site.

According to another advantageous embodiment of said method, the mutation in step (a) is a substitution of at least one amino acid of said final C-terminal loop, with a different amino acid.

Preferably, the residue in position 138 or 139 is substituted by an hydrophobic amino acid to avoid the formation of hydrogen bonds with the phosphate

backbone of the DNA cleavage site. For example, the residue in position 138 is substituted by an alanine or the residue in position 139 is substituted by a methionine.

The residue in position 142 or 143 is advantageously substituted by a small amino acid, for example a glycine, to decrease the size of the side chains of these amino acid residues.

According to the method of the invention, the mutation(s) in step (a) are introduced in either a wild-type LAGLIDADG homing endonuclease or a functional variant thereof.

The wild-type LAGLIDADG homing endonuclease is advantageously homodimeric. Examples of wild-type homodimeric LAGLIDADG homing endonucleases are presented in Table 1 of Lucas *et al.*, Nucleic Acids Res., 2001, 29, 960-969. The wild-type homodimeric LAGLIDADG homing endonuclease may be advantageously selected from the group consisting of: I-*CreI*, I-*CeuI*, I-*MsoI* and I-*CpaI*, preferably I-*CreI*.

The functional variant comprises additional mutations outside the final C-terminal loop, preferably in positions of amino acid residues which interact with a DNA target half-site. The LAGLIDADG homing endonucleases DNA interacting residues are well-known in the art. The residues which are mutated may interact with the DNA backbone or with the nucleotide bases, directly or via a water molecule. Preferably said mutations modify the cleavage specificity of the meganuclease and result in a meganuclease with novel specificity, which is able to cleave a DNA target from a gene of interest. More preferably, said mutations are substitutions of one or more amino acids in a first functional subdomain corresponding to that situated from positions 26 to 40 of I-*CreI* amino acid sequence, that alter the specificity towards the nucleotide in positions ± 8 to 10 of the DNA target, and/or substitutions in a second functional subdomain corresponding to that situated from positions 44 to 77 of I-*CreI* amino acid sequence, that alter the specificity towards the nucleotide in positions ± 3 to 5 of the DNA target, as described previously (International PCT Applications WO 2006/097784 and WO 2006/097853; Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; Smith *et al.*, Nucleic Acids Res., 2006). The substitutions correspond advantageously to positions 26, 28, 30, 32, 33, 38, and/or 40, 44, 68, 70, 75 and/or 77 of I-*CreI* amino acid sequence. For cleaving a

DNA target, wherein n_{-4} is t or n_{+4} is a, said variant has advantageously a glutamine (Q) in position 44; for cleaving a DNA target, wherein n_{-4} is a or n_{+4} is t, said variant has an alanine (A) or an asparagine in position 44, and for cleaving a DNA target, wherein n_{-9} is g or n_{+9} is c, said variant has advantageously an arginine (R) or a lysine (K) in position 38.

According to a most preferred embodiment of said method, the parent LAGLIDADG homing endonuclease is an I-CreI variant having mutations in positions 26 to 40 and 44 to 77 of I-CreI and cleaving a palindromic DNA sequence, wherein at least the nucleotides in positions +3 to +5 and +8 to +10 or -10 to -8 and -5 to -3 of one half of said DNA sequence correspond to the nucleotides in positions +3 to +5 and +8 to +10 or -10 to -8 and -5 to -3 of one half of a DNA target from a gene of interest.

The mutations in step (a) are introduced according to standard mutagenesis methods which are well-known in the art and commercially available. They may be advantageously produced by amplifying overlapping fragments comprising the mutated position(s), as defined above, according to well-known overlapping PCR techniques. Libraries of variants having amino acid variation in the final C-terminal loop may be generated according to standard methods.

Step (a) may comprise the introduction of additional mutations at other positions contacting the DNA target sequence or interacting directly or indirectly with said DNA target, as defined above. This step may be performed by generating combinatorial libraries as described in the International PCT Application WO 2004/067736, Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458 and Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006 and eventually, combining said mutations intramolecularly, by amplifying overlapping fragments comprising each of the mutations, according to well-known overlapping PCR techniques.

Furthermore, random mutations may also be introduced on the whole variant or in part of the variant, in particular the C-terminal half of the variant (positions 80 to 163 of I-CreI amino acid sequence SEQ ID NO:2) in order to improve the binding and/or cleavage properties of the variant towards a DNA target from a gene of interest.

The additional mutations (random or site-specific) and the mutation(s) in the final C-terminal loop may be introduced simultaneously or subsequently.

5 In addition, one or more residues may be inserted at the NH₂ terminus and/or COOH terminus of the variant monomer(s)/domain(s). For example, a methionine residue is introduced at the NH₂ terminus, a tag (epitope or polyhistidine sequence) is introduced at the NH₂ terminus and/or COOH terminus; said tag is useful for the detection and/or the purification of the meganuclease.

10 The selection and/or screening in step (b) may be performed by using a cleavage assay *in vitro* or *in vivo*, as described in the International PCT Application WO 2004/067736.

15 According to another advantageous embodiment of said method, step (b) is performed *in vivo*, under conditions where the double-strand break in a mutated DNA target sequence which is generated by said variant leads to the activation of a positive selection marker or a reporter gene, or the inactivation of a negative selection marker or a reporter gene, by recombination-mediated repair of said DNA double-strand break.

For example, the cleavage activity of the variant of the invention may be measured by a direct repeat recombination assay, in yeast or mammalian cells, 20 using a reporter vector, as described in the PCT Application WO 2004/067736; Epinat *et al.*, Nucleic Acids Res., 2003, 31, 2952-2962; Chames *et al.*, Nucleic Acids Res., 2005, 33, e178, and Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458. The reporter vector comprises two truncated, non-functional copies of a reporter gene (direct repeats) and a chimeric DNA target sequence within the intervening sequence, cloned 25 in a yeast or a mammalian expression vector. The DNA target sequence is derived from the parent homing endonuclease cleavage site by replacement of at least one nucleotide by a different nucleotide. Preferably a panel of palindromic or non-palindromic DNA targets representing the different combinations of the 4 bases (g, a, c, t) at one or more positions of the DNA cleavage site is tested (4ⁿ palindromic targets 30 for n mutated positions). Expression of the variant results in a functional endonuclease which is able to cleave the DNA target sequence. This cleavage induces homologous

recombination between the direct repeats, resulting in a functional reporter gene, whose expression can be monitored by appropriate assay.

According to another advantageous embodiment of said method, step (b) comprises the selection and/or screening of the variants from step (a) which are
5 able to cleave at least one DNA target sequence that is not cleaved by said parent LAGLIDADG homing endonuclease, said DNA target sequence being derived from the parent LAGLIDADG homing endonuclease cleavage site, by the replacement of at least one nucleotide of one half of said cleavage site, with a different nucleotide.

According to the method of the invention, the parent DNA target may
10 be palindromic, non-palindromic or pseudo-palindromic. Preferably, said DNA target sequence is derived from the I-CreI palindromic site having the sequence SEQ ID NO: 1. More preferably, said DNA target has nucleotide mutation(s) in positions ± 1 to 2, ± 6 to 7, ± 8 to 10 and/or ± 11 to 12, still more preferably in positions ± 1 to 2, ± 6 to 7 and/or ± 11 to 12.

According to another advantageous embodiment of said method, it
15 comprises a further step (c) of expressing one variant obtained in step (b), so as to allow the formation of homodimers. Said homodimers are able to cleave a palindromic or pseudo-palindromic target sequences.

According to another advantageous embodiment of said method, it
20 comprises a further step (c') of co-expressing one variant obtained in step (b) and a wild-type LAGLIDADG homing endonuclease or a functional variant thereof, so as to allow the formation of heterodimers. The assembly of functional heterodimers by co-expression of two different LAGLIDADG endonucleases monomers, has been described previously in Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; International
25 PCT Applications WO 2006/097853, WO 2006/097854 and WO 2006/097784; Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006. Preferably, two different variants obtained in step (b) are co-expressed. Said heterodimers are able to cleave a non-palindromic chimeric target.

For example, host cells may be modified by one or two recombinant
30 expression vector(s) encoding said variant(s). The cells are then cultured under conditions allowing the expression of the variant(s) and the homodimers/heterodimers which are formed are then recovered from the cell culture.

According to the method of the invention, single-chain chimeric meganucleases may be constructed by the fusion of one variant obtained in step (b) with a homing endonuclease domain/monomer. Said domain/monomer may be from a wild-type LAGLIDADG homing endonuclease or a functional variant thereof.

5 Preferably, the two domain(s)/monomer(s) are connected by a peptidic linker. More preferably, the single-chain meganuclease comprises two different variants obtained in step (b); said single-chain meganuclease is able cleave a non-palindromic chimeric target comprising one different half of each variant DNA target.

Methods for constructing single-chain chimeric meganucleases derived from homing endonucleases are well-known in the art (Epinat et al., Nucleic Acids Res., 2003, 31, 2952-62; Chevalier et al., Mol. Cell., 2002, 10, 895-905; Steuer et al., Chembiochem., 2004, 5, 206-13; International PCT Applications WO 03/078619 and WO 2004/031346). Any of such methods, may be applied for constructing single-chain chimeric meganucleases derived from the variants as

10 defined in the present invention.

The invention relates also to an homodimeric or heterodimeric LAGLIDADG homing endonuclease variant obtainable by the method as defined above, with the exclusion of the homodimeric variants of SEQ ID NO: 3 and 4 and the homodimeric or heterodimeric variants comprising a monomer of SEQ ID NO: 5; the

20 LAGLIDADG homing endonuclease variant of the invention is also named as variant, meganuclease variant or meganuclease.

According to an advantageous embodiment of said variant, it is an heterodimer comprising monomers from two different variants obtainable by the method as defined above.

25 According to another advantageous embodiment of said variant, it is an I-CreI variant having one or two mutations, each one from a different pair of mutations selected from the group consisting of the pair S138A and K139M and the pair K142G and T143G. Examples of such variants include SEQ ID NO: 6 to 9.

More preferably, said I-CreI variant is an heterodimer, comprising

30 two monomers, each one further comprising different mutations in positions 26 to 40 and 44 to 77 of I-CreI and being able to cleave a genomic DNA target from a gene of interest.

The subject-matter of the present invention is also a single-chain chimeric meganuclease derived from the variant as defined above; the single-chain chimeric meganuclease of the invention is also named as single-chain derivative, single-chain meganuclease, single-chain meganuclease derivative or meganuclease.

5 The meganuclease of the invention includes both the meganuclease variant and the single-chain meganuclease derivative.

 The subject-matter of the present invention is also a polynucleotide fragment encoding a variant or a single-chain derivative as defined above; said polynucleotide may encode one monomer of an homodimeric or heterodimeric
10 variant, or two domains/monomers of a single-chain derivative.

 The subject-matter of the present invention is also a recombinant vector for the expression of a variant or a single-chain derivative according to the invention. The recombinant vector comprises at least one polynucleotide fragment encoding a variant or a single-chain meganuclease, as defined above. In a preferred
15 embodiment, said vector comprises two different polynucleotide fragments, each encoding one of the monomers of an heterodimeric variant.

 A vector which can be used in the present invention includes, but is not limited to, a viral vector, a plasmid, a RNA vector or a linear or circular DNA or RNA molecule which may consists of a chromosomal, non chromosomal, semi-
20 synthetic or synthetic nucleic acids. Preferred vectors are those capable of autonomous replication (episomal vector) and/or expression of nucleic acids to which they are linked (expression vectors). Large numbers of suitable vectors are known to those of skill in the art and commercially available.

 Viral vectors include retrovirus, adenovirus, parvovirus (e. g. adeno-
25 associated viruses), coronavirus, negative strand RNA viruses such as orthomyxovirus (e. g., influenza virus), rhabdovirus (e. g., rabies and vesicular stomatitis virus), paramyxovirus (e. g. measles and Sendai), positive strand RNA viruses such as picornavirus and alphavirus, and double-stranded DNA viruses including adenovirus, herpesvirus (e. g., Herpes Simplex virus types 1 and 2, Epstein-Barr virus, cytomegalovirus), and poxvirus (e. g., vaccinia, fowlpox and canarypox). Other viruses include
30 Norwalk virus, togavirus, flavivirus, reoviruses, papovavirus, hepadnavirus, and hepatitis virus, for example. Examples of retroviruses include: avian leukosis-

sarcoma, mammalian C-type, B-type viruses, D type viruses, HTLV-BLV group, lentivirus, spumavirus (Coffin, J. M., Retroviridae: The viruses and their replication, In Fundamental Virology, Third Edition, B. N. Fields, et al., Eds., Lippincott-Raven Publishers, Philadelphia, 1996).

5 Preferred vectors include lentiviral vectors, and particularly self inactivating lentiviral vectors.

Vectors can comprise selectable markers, for example: neomycin phosphotransferase, histidinol dehydrogenase, dihydrofolate reductase, hygromycin phosphotransferase, herpes simplex virus thymidine kinase, adenosine deaminase,
10 glutamine synthetase, and hypoxanthine-guanine phosphoribosyl transferase for eukaryotic cell culture; TRP1 for *S. cerevisiae*; tetracycline, rifampicin or ampicillin resistance in *E. coli*.

Preferably said vectors are expression vectors, wherein the sequence(s) encoding the variant/single-chain derivative of the invention is placed
15 under control of appropriate transcriptional and translational control elements to permit production or synthesis of said meganuclease. Therefore, said polynucleotide is comprised in an expression cassette. More particularly, the vector comprises a replication origin, a promoter operatively linked to said encoding polynucleotide, a ribosome-binding site, an RNA-splicing site (when genomic DNA is used), a
20 polyadenylation site and a transcription termination site. It also can comprise an enhancer. Selection of the promoter will depend upon the cell in which the polypeptide is expressed. Preferably, when said variant is an heterodimer, the two polynucleotides encoding each of the monomers are included in one vector which is able to drive the expression of both polynucleotides, simultaneously. Suitable promoters
25 include tissue specific and/or inducible promoters. Examples of inducible promoters are: eukaryotic metallothionine promoter which is induced by increased levels of heavy metals, prokaryotic lacZ promoter which is induced in response to isopropyl- β -D-thiogalacto-pyranoside (IPTG) and eukaryotic heat shock promoter which is induced by increased temperature. Examples of tissue specific promoters are skeletal
30 muscle creatine kinase, prostate-specific antigen (PSA), α -antitrypsin protease, human surfactant (SP) A and B proteins, β -casein and acidic whey protein genes.

According to another advantageous embodiment of said vector, it includes a targeting DNA construct comprising sequences sharing homologies with the region surrounding the genomic DNA target cleavage site as defined above.

Alternatively, the vector coding for the meganuclease and the vector comprising the targeting DNA construct are different vectors.

More preferably, the targeting DNA construct comprises:

a) sequences sharing homologies with the region surrounding the genomic DNA cleavage site as defined above, and

b) a sequence to be introduced flanked by sequences as in a).

Preferably, homologous sequences of at least 50 bp, preferably more than 100 bp and more preferably more than 200 bp are used. Indeed, shared DNA homologies are located in regions flanking upstream and downstream the site of the break and the DNA sequence to be introduced should be located between the two arms. The sequence to be introduced is preferably a sequence which repairs a mutation in the gene of interest (gene correction or recovery of a functional gene), for the purpose of genome therapy. Alternatively, it can be any other sequence used to alter the chromosomal DNA in some specific way including a sequence used to modify a specific sequence, to attenuate or activate the endogenous gene of interest, to inactivate or delete the endogenous gene of interest or part thereof, to introduce a mutation into a site of interest or to introduce an exogenous gene or part thereof.

The invention also concerns a prokaryotic or eukaryotic host cell which is modified by a polynucleotide or a vector as defined above, preferably an expression vector.

The invention also concerns a non-human transgenic animal or a transgenic plant, characterized in that all or part of their cells are modified by a polynucleotide or a vector as defined above.

As used herein, a cell refers to a prokaryotic cell, such as a bacterial cell, or eukaryotic cell, such as an animal, plant or yeast cell.

The subject-matter of the present invention is further the use of a meganuclease with the exclusion of SEQ ID NO: 5, one or two derived polynucleotide(s), preferably included in expression vector(s), a cell, a transgenic plant, a non-human transgenic mammal, as defined above, for molecular biology, for

in vivo or *in vitro* genetic engineering, and for *in vivo* or *in vitro* genome engineering, for non-therapeutic purposes.

Non therapeutic purposes include for example (i) gene targeting of specific loci in cell packaging lines for protein production, (ii) gene targeting of
5 specific loci in crop plants, for strain improvements and metabolic engineering, (iii) targeted recombination for the removal of markers in genetically modified crop plants, (iv) targeted recombination for the removal of markers in genetically modified microorganism strains (for antibiotic production for example).

According to an advantageous embodiment of said use, it is for
10 inducing a double-strand break in a site of interest comprising a DNA target sequence, thereby inducing a DNA recombination event, a DNA loss or cell death.

According to the invention, said double-strand break is for: repairing a specific sequence, modifying a specific sequence, restoring a functional gene in place of a mutated one, attenuating or activating an endogenous gene of interest,
15 introducing a mutation into a site of interest, introducing an exogenous gene or a part thereof, inactivating or detecting an endogenous gene or a part thereof, translocating a chromosomal arm, or leaving the DNA unrepaired and degraded.

The subject-matter of the present invention is also a method of genetic engineering, characterized in that it comprises a step of double-strand nucleic
20 acid breaking in a site of interest located on a vector comprising a DNA target as defined hereabove, by contacting said vector with a meganuclease as defined above, with the exclusion of SEQ ID NO: 5, thereby inducing an homologous recombination with another vector presenting homology with the sequence surrounding the cleavage site of said meganuclease.

25 The subject-matter of the present invention is also a method of genome engineering, characterized in that it comprises the following steps: 1) double-strand breaking a genomic locus comprising at least one DNA target of a meganuclease as defined above, by contacting said target with said meganuclease, with the exclusion of SEQ ID NO: 5 ; 2) maintaining said broken genomic locus under
30 conditions appropriate for homologous recombination with a targeting DNA construct comprising the sequence to be introduced in said locus, flanked by sequences sharing homologies with the targeted locus.

The subject-matter of the present invention is also a method of genome engineering, characterized in that it comprises the following steps: 1) double-strand breaking a genomic locus comprising at least one DNA target of a meganuclease as defined above, by contacting said cleavage site with said
5 meganuclease, with the exclusion of SEQ ID NO: 5; 2) maintaining said broken genomic locus under conditions appropriate for homologous recombination with chromosomal DNA sharing homologies to regions surrounding the cleavage site.

The subject-matter of the present invention is also the use of at least one meganuclease as defined above, with the exclusion of SEQ ID NO: 5, one or two
10 derived polynucleotide(s), preferably included in expression vector(s), as defined above, for the preparation of a medicament for preventing, improving or curing a genetic disease in an individual in need thereof, said medicament being administrated by any means to said individual.

The subject-matter of the present invention is also a method for
15 preventing, improving or curing a genetic disease in an individual in need thereof, said method comprising the step of administering to said individual a composition comprising at least a meganuclease as defined above, by any means.

In this case, the use of the meganuclease as defined above, comprises at least the step of (a) inducing in somatic tissue(s) of the individual a
20 double stranded cleavage at a site of interest of a gene comprising at least one recognition and cleavage site of said meganuclease, and (b) introducing into the individual a targeting DNA, wherein said targeting DNA comprises (1) DNA sharing homologies to the region surrounding the cleavage site and (2) DNA which repairs the site of interest upon recombination between the targeting DNA and the chromosomal
25 DNA. The targeting DNA is introduced into the individual under conditions appropriate for introduction of the targeting DNA into the site of interest.

According to the present invention, said double-stranded cleavage is induced, either *in toto* by administration of said meganuclease to an individual, or *ex vivo* by introduction of said meganuclease into somatic cells removed from an
30 individual and returned into the individual after modification.

In a preferred embodiment of said use, the meganuclease is combined with a targeting DNA construct comprising a sequence which repairs a

mutation in the gene flanked by sequences sharing homologies with the regions of the gene surrounding the genomic DNA cleavage site of said meganuclease, as defined above. The sequence which repairs the mutation is either a fragment of the gene with the correct sequence or an exon knock-in construct.

5 For correcting a gene, cleavage of the gene occurs in the vicinity of the mutation, preferably, within 500 bp of the mutation. The targeting construct comprises a gene fragment which has at least 200 bp of homologous sequence flanking the genomic DNA cleavage site (minimal repair matrix) for repairing the cleavage, and includes the correct sequence of the gene for repairing the mutation.
10 Consequently, the targeting construct for gene correction comprises or consists of the minimal repair matrix; it is preferably from 200 pb to 6000 pb, more preferably from 1000 pb to 2000 pb.

For restoring a functional gene, cleavage of the gene occurs upstream of a mutation. Preferably said mutation is the first known mutation in the
15 sequence of the gene, so that all the downstream mutations of the gene can be corrected simultaneously. The targeting construct comprises the exons downstream of the genomic DNA cleavage site fused in frame (as in the cDNA) and with a polyadenylation site to stop transcription in 3'. The sequence to be introduced (exon knock-in construct) is flanked by introns or exons sequences surrounding the cleavage
20 site, so as to allow the transcription of the engineered gene (exon knock-in gene) into a mRNA able to code for a functional protein. For example, the exon knock-in construct is flanked by sequences upstream and downstream.

The subject-matter of the present invention is also the use of at least one meganuclease as defined above, with the exclusion of SEQ ID NO: 5, one or or
25 two derived polynucleotide(s), preferably included in expression vector(s), as defined above for the preparation of a medicament for preventing, improving or curing a disease caused by an infectious agent that presents a DNA intermediate, in an individual in need thereof, said medicament being administrated by any means to said individual.

30 The subject-matter of the present invention is also a method for preventing, improving or curing a disease caused by an infectious agent that presents a DNA intermediate, in an individual in need thereof, said method comprising at least

the step of administering to said individual a composition as defined above, by any means.

The subject-matter of the present invention is also the use of at least one meganuclease as defined above, one or two polynucleotide(s), preferably included
5 in expression vector(s), as defined above, *in vitro*, for inhibiting the propagation, inactivating or deleting an infectious agent that presents a DNA intermediate, in biological derived products or products intended for biological uses or for disinfecting an object.

The subject-matter of the present invention is also a method for
10 decontaminating a product or a material from an infectious agent that presents a DNA intermediate, said method comprising at least the step of contacting a biological derived product, a product intended for biological use or an object, with a composition as defined above, for a time sufficient to inhibit the propagation, inactivate or delete said infectious agent.

15 In a particular embodiment, said infectious agent is a virus. For example said virus is an adenovirus (Ad11, Ad21), herpesvirus (HSV, VZV, EBV, CMV, herpesvirus 6, 7 or 8), hepadnavirus (HBV), papovavirus (HPV), poxvirus or retrovirus (HTLV, HIV).

The subject-matter of the present invention is also a composition
20 characterized in that it comprises at least one meganuclease with the exclusion of SEQ ID NO:5, one or two derived polynucleotide(s), preferably included in expression vector(s), as defined above.

In a preferred embodiment of said composition, it comprises a
25 targeting DNA construct comprising the sequence which repairs the site of interest flanked by sequences sharing homologies with the targeted locus as defined above. Preferably, said targeting DNA construct is either included in a recombinant vector or it is included in an expression vector comprising the polynucleotide(s) encoding the meganuclease, as defined in the present invention.

The subject-matter of the present invention is also products
30 containing at least a meganuclease with the exclusion of SEQ ID NO: 5, or one or two expression vector(s) encoding said meganuclease, and a vector including a targeting

construct, as defined above, as a combined preparation for simultaneous, separate or sequential use in the prevention or the treatment of a genetic disease.

For purposes of therapy, the meganuclease and a pharmaceutically acceptable excipient are administered in a therapeutically effective amount. Such a combination is said to be administered in a "therapeutically effective amount" if the amount administered is physiologically significant. An agent is physiologically significant if its presence results in a detectable change in the physiology of the recipient. In the present context, an agent is physiologically significant if its presence results in a decrease in the severity of one or more symptoms of the targeted disease and in a genome correction of the lesion or abnormality.

In one embodiment of the uses according to the present invention, the meganuclease is substantially non-immunogenic, i.e., engenders little or no adverse immunological response. A variety of methods for ameliorating or eliminating deleterious immunological reactions of this sort can be used in accordance with the invention. In a preferred embodiment, the meganuclease is substantially free of N-formyl methionine. Another way to avoid unwanted immunological reactions is to conjugate meganucleases to polyethylene glycol ("PEG") or polypropylene glycol ("PPG") (preferably of 500 to 20,000 daltons average molecular weight (MW)). Conjugation with PEG or PPG, as described by Davis *et al.* (US 4,179,337) for example, can provide non-immunogenic, physiologically active, water soluble endonuclease conjugates with anti-viral activity. Similar methods also using a polyethylene-polypropylene glycol copolymer are described in Saifer *et al.* (US 5,006,333).

The meganuclease can be used either as a polypeptide or as a polynucleotide construct/vector encoding said polypeptide. It is introduced into cells, *in vitro*, *ex vivo* or *in vivo*, by any convenient means well-known to those in the art, which are appropriate for the particular cell type, alone or in association with either at least an appropriate vehicle or carrier and/or with the targeting DNA. Once in a cell, the meganuclease and if present, the vector comprising targeting DNA and/or nucleic acid encoding a meganuclease are imported or translocated by the cell from the cytoplasm to the site of action in the nucleus.

The meganuclease (polypeptide) may be advantageously associated with: liposomes, polyethyleneimine (PEI), and/or membrane translocating peptides (Bonetta, *The Scientist*, 2002, 16, 38; Ford *et al.*, *Gene Ther.*, 2001, 8, 1-4 ; Wadia and Dowdy, *Curr. Opin. Biotechnol.*, 2002, 13, 52-56); in the latter case, the sequence of the meganuclease fused with the sequence of a membrane translocating peptide (fusion protein).

Vectors comprising targeting DNA and/or nucleic acid encoding a meganuclease can be introduced into a cell by a variety of methods (e.g., injection, direct uptake, projectile bombardment, liposomes, electroporation). Meganucleases can be stably or transiently expressed into cells using expression vectors. Techniques of expression in eukaryotic cells are well known to those in the art. (See *Current Protocols in Human Genetics*: Chapter 12 "Vectors For Gene Therapy" & Chapter 13 "Delivery Systems for Gene Therapy"). Optionally, it may be preferable to incorporate a nuclear localization signal into the recombinant protein to be sure that it is expressed within the nucleus.

The uses of the meganuclease and the methods of using said meganucleases according to the present invention include also the use of the polynucleotide(s), vector(s), cell, transgenic plant or non-human transgenic mammal encoding said meganuclease, as defined above.

According to another advantageous embodiment of the uses and methods according to the present invention, said meganuclease, polynucleotide(s), vector(s), cell, transgenic plant or non-human transgenic mammal are associated with a targeting DNA construct as defined above. Preferably, said vector encoding the monomer(s) of the meganuclease, comprises the targeting DNA construct, as defined above.

The invention concerns also a first method for engineering I-CreI variants able to cleave a genomic DNA target sequence from a gene of interest, comprising at least the steps of:

(a₁) constructing a first series of variants having at least one substitution in a first functional subdomain of the LAGLIDADG core domain situated from positions 26 to 40 of I-CreI,

(b₁) constructing a second series of I-CreI variants having at least one substitution in a second functional subdomain of the LAGLIDADG core domain situated from positions 44 to 77 of I-CreI,

5 (c₁) selecting and/or screening the variants from the first series of step (a₁) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -10 to -8 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions -10 to -8 of said genomic target and (ii) the nucleotide triplet in positions +8 to +10 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions -10 to -8 of said
10 genomic target,

(d₁) selecting and/or screening the variants from the second series of step (b₁) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -5 to -3 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions -5 to -3 of said genomic target and (ii) the nucleotide
15 triplet in positions +3 to +5 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions -5 to -3 of said genomic target,

(e₁) selecting and/or screening the variants from the first series of step (a₁) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet
20 in positions +8 to +10 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions +8 to +10 of said genomic target and (ii) the nucleotide triplet in positions -10 to -8 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions +8 to +10 of said genomic target,

25 (f₁) selecting and/or screening the variants from the second series of step (b) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions +3 to +5 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions +3 to +5 of said genomic target and (ii) the nucleotide triplet in positions -5 to -3 has been replaced with the reverse complementary
30 sequence of the nucleotide triplet which is present in positions +3 to +5 of said genomic target,

(g₁) combining in a single variant, the mutation(s) in positions 26 to 40 and 44 to 77 of two variants from step (c₁) and step (d₁), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions -10 to -8 is identical to the nucleotide triplet which is present in positions -10 to -8 of said genomic target, (ii) the nucleotide triplet in positions +8 to +10 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions -10 to -8 of said genomic target, (iii) the nucleotide triplet in positions -5 to -3 is identical to the nucleotide triplet which is present in positions -5 to -3 of said genomic target and (iv) the nucleotide triplet in positions +3 to +5 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions -5 to -3 of said genomic target,

(h₁) combining in a single variant, the mutation(s) in positions 26 to 40 and 44 to 77 of two variants from step (e₁) and step (f₁), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions +3 to +5 is identical to the nucleotide triplet which is present in positions +3 to +5 of said genomic target, (ii) the nucleotide triplet in positions -5 to -3 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions +3 to +5 of said genomic target, (iii) the nucleotide triplet in positions +8 to +10 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions +8 to +10 of said genomic target and (iv) the nucleotide triplet in positions -10 to -8 is identical to the reverse complementary sequence of the nucleotide triplet in positions +8 to +10 of said genomic target,

(i₁) introducing in the variants from step (g₁) and/or (h₁), at least one mutation in the final C-terminal loop, preferably a substitution in position 138, 139, 142 or 143 of I-*CreI*, as defined above,

(j₁) combining the variants obtained in steps (g₁), (h₁) and/or (i₁) to form heterodimers, and

(k₁) selecting and/or screening the heterodimers from step (j₁) which are able to cleave said genomic DNA target situated in a gene of interest.

Alternatively, the I-*CreI* variant according to the invention may be obtained by a second method for engineering I-*CreI* variants able to cleave a genomic DNA target sequence from a gene of interest, comprising at least the steps of:

(a₂) constructing a first series of I-CreI variants having at least one substitution in a first functional subdomain of the LAGLIDADG core domain situated from positions 26 to 40 of I-CreI and one mutation in the final C-terminal loop, preferably a substitution in position 138, 139, 142 or 143 of I-CreI, as defined above,

5 (b₂) constructing a second series of I-CreI variants having at least one substitution in a second functional subdomain of the LAGLIDADG core domain situated from positions 44 to 77 of I-CreI and one mutation in the final C-terminal loop, preferably a substitution in position 138, 139, 142 or 143 of I-CreI, as defined above, with the proviso that at least one of the two series of I-CreI variants comprise
10 at least one mutation in the final C-terminal loop,

(c₂) selecting and/or screening the variants from the first series of step (a₂) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -10 to -8 and eventually at least one of the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 of the I-CreI site have been replaced,
15 respectively with the nucleotide triplet which is present in positions -10 to -8 and the nucleotide doublet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target (ii) the nucleotide triplet in positions +8 to +10 and eventually at least one of the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 have been replaced with the reverse complementary sequence of respectively, the
20 nucleotide triplet which is present in positions -10 to -8 and the nucleotide doublet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target,

(d₂) selecting and/or screening the variants from the second series of step (b₂) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -5 to -3 and eventually at least one of the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 of the I-CreI site have been replaced
25 respectively with the nucleotide triplet which is present in positions -5 to -3 and the nucleotide doublet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target and (ii) the nucleotide triplet in positions +3 to +5 and eventually at least one of the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 have been replaced with the reverse complementary sequence of respectively the
30 nucleotide triplet which is present in positions -5 to -3 and the nucleotide doublet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target,

with the proviso that at least one of the two mutant *I-CreI* sites in step (c) and (d) have mutation(s) in at least one of the nucleotide doublet(s) in position -12 to -11, -7 to -6 and/or -2 to -1 and at least one of the corresponding nucleotide doublet (s) in positions +1 to +2, +6 to +7, and/or +11 to +12 of the *I-CreI* site,

5 (e₂) selecting and/or screening the variants from the first series of step (a₂) which are able to cleave a mutant *I-CreI* site wherein (i) the nucleotide triplet in positions +8 to +10, and eventually at least one of the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 of the *I-CreI* site have been replaced
10 respectively with the nucleotide triplet which is present in positions +8 to +10, and the nucleotide doublet which is present in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target and (ii) the nucleotide triplet in positions -10 to -8 and eventually the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 have been replaced with the reverse complementary sequence of respectively the nucleotide triplet which is present in positions +8 to +10, and the nucleotide doublet
15 which is present in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target,

(f₂) selecting and/or screening the variants from the second series of step (b₂) which are able to cleave a mutant *I-CreI* site wherein (i) the nucleotide triplet in positions +3 to +5, and eventually the nucleotide doublet(s) in positions +1 to +2,
20 +6 to +7, and/or +11 to +12 of the *I-CreI* site, have been replaced respectively with the nucleotide triplet which is present in positions +3 to +5, and the nucleotide doublet(s) which is present in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target and (ii) the nucleotide triplet in positions -5 to -3, and eventually the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 have been
25 replaced with the reverse complementary sequence of, respectively the nucleotide triplet which is present in positions +3 to +5, and the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target, with the proviso that at least one of the two mutant *I-CreI* sites in step (e) and (f) have mutation(s) in at least one of the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 and
30 at least one of the corresponding nucleotide doublet (s) in positions in positions -12 to -11, -7 to -6 and/or -2 to -1 of the *I-CreI* site,

(g₂) combining in a single variant, the mutation(s) in positions 26 to 40, 44 to 77 and in the final C-terminal loop of two variants from step (c₂) and step (d₂), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions -10 to -8 and the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 are identical, respectively to the nucleotide triplet which is present in positions -10 to -8, and the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target, (ii) the nucleotide triplet in positions +8 to +10 and the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 are identical to the reverse complementary sequence of, respectively the nucleotide triplet which is present in positions -10 to -8 and the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 of said genomic target, (iii) the nucleotide triplet in positions -5 to -3 is identical to the nucleotide triplet which is present in positions -5 to -3 of said genomic target and (iv) the nucleotide triplet in positions +3 to +5 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions -5 to -3 of said genomic target,

(h₂) combining in a single variant, the mutation(s) in positions 26 to 40, 44 to 77 and in the final C-terminal loop of two variants from step (e₂) and step (f₂), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions +3 to +5 and the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 are identical, respectively to the nucleotide triplet which is present in positions +3 to +5 and the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target, (ii) the nucleotide triplet in positions -5 to -3 and the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 are identical to the reverse complementary sequence of, respectively the nucleotide triplet which is present in positions +3 to +5 and the nucleotide doublet(s) present in positions +1 to +2, +6 to +7, and/or +11 to +12 of said genomic target, (iii) the nucleotide triplet in positions +8 to +10 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions +8 to +10 of said genomic target and (iv) the nucleotide triplet in positions -10 to -8 is identical to the reverse complementary sequence of the nucleotide triplet in positions +8 to +10 of said genomic target,

(i₂) combining the variants obtained in steps (g₂) and (h₂) to form heterodimers, and

(j₂) selecting and/or screening the heterodimers from step (i₂) which are able to cleave said genomic DNA target situated in a gene of interest.

5 According to yet another alternative, the I-CreI variant of the invention may be obtained by a third method for engineering I-CreI variants able to cleave a genomic DNA target sequence from a gene of interest, comprising at least the steps of:

10 (a₃) constructing a first series of variants having at least one substitution in a first functional subdomain of the LAGLIDADG core domain situated from positions 26 to 40 of I-CreI,

(b₃) constructing a second series of I-CreI variants having at least one substitution in a second functional subdomain of the LAGLIDADG core domain situated from positions 44 to 77 of I-CreI,

15 (c₃) constructing a third series of variants having at least one mutation in the final C-terminal loop, preferably a substitution in position 138, 139, 142 or 143 of I-CreI, as defined above,

(d₃) selecting and/or screening the variants from the first series of step (a₃) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -10 to -8 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions -10 to -8 of said genomic target and (ii) the nucleotide triplet in positions +8 to +10 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions -10 to -8 of said genomic target,

25 (e₃) selecting and/or screening the variants from the second series of step (b₃) which are able to cleave a mutant I-CreI site wherein (i) the nucleotide triplet in positions -5 to -3 of the I-CreI site has been replaced with the nucleotide triplet which is present in positions -5 to -3 of said genomic target and (ii) the nucleotide triplet in positions +3 to +5 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions -5 to -3 of said genomic target,

30

(f₃) selecting and/or screening the variants from the third series of step (c₃) which are able to cleave a mutant I-*CreI* site wherein (i) the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1, respectively, of said genomic target and (ii) the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions -2 to -1, -7 to -6, and/or -12 to -11, respectively, of said genomic target,

(g₃) selecting and/or screening the variants from the first series of step (a₃) which are able to cleave a mutant I-*CreI* site wherein (i) the nucleotide triplet in positions +8 to +10 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions +8 to +10 of said genomic target and (ii) the nucleotide triplet in positions -10 to -8 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions +8 to +10 of said genomic target,

(h₃) selecting and/or screening the variants from the second series of step (b₃) which are able to cleave a mutant I-*CreI* site wherein (i) the nucleotide triplet in positions +3 to +5 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions +3 to +5 of said genomic target and (ii) the nucleotide triplet in positions -5 to -3 has been replaced with the reverse complementary sequence of the nucleotide triplet which is present in positions +3 to +5 of said genomic target,

(i₃) selecting and/or screening the variants from the third series of step (c₃) which are able to cleave a mutant I-*CreI* site wherein (i) the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 of the I-*CreI* site has been replaced with the nucleotide doublet(s) which is present in positions +1 to +2, +6 to +7, and/or +11 to +12, respectively, of said genomic target and (ii) the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 has been replaced with the reverse complementary sequence of the nucleotide doublet which is present in positions +11 to +12, +6 to +7, and/or +1 to +2, respectively, of said genomic target,

(j₃) combining in a single variant, the mutation(s) in positions 26 to 40, 44 to 77 and in the final C-terminal loop of three variants from step (d₃) (e₃) and

(f₃), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions -10 to -8 is identical to the nucleotide triplet which is present in positions -10 to -8 of said genomic target, (ii) the nucleotide triplet in positions +8 to +10 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions -10 to -8 of said genomic target, (iii) the nucleotide triplet in positions -5 to -3 is identical to the nucleotide triplet which is present in positions -5 to -3 of said genomic target and (iv) the nucleotide triplet in positions +3 to +5 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions -5 to -3 of said genomic target, (v) the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 is identical to the nucleotide triplet which is present in positions -12 to -11, -7 to -6 and/or -2 to -1, respectively, of said genomic target, (vi) the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 is identical to the reverse complementary sequence of the nucleotide doublet(s) which is present in positions -2 to -1, -7 to -6, and/or -12 to -11, respectively, of said genomic target,

(k₃) combining in a single variant, the mutation(s) in positions 26 to 40, 44 to 77 and in the final C-terminal loop of three variants from step (g₃) (h₃) and (i₃), to obtain a novel homodimeric I-*CreI* variant which cleaves a sequence wherein (i) the nucleotide triplet in positions +3 to +5 is identical to the nucleotide triplet which is present in positions +3 to +5 of said genomic target, (ii) the nucleotide triplet in positions -5 to -3 is identical to the reverse complementary sequence of the nucleotide triplet which is present in positions +3 to +5 of said genomic target, (iii) the nucleotide triplet in positions +8 to +10 of the I-*CreI* site has been replaced with the nucleotide triplet which is present in positions +8 to +10 of said genomic target and (iv) the nucleotide triplet in positions -10 to -8 is identical to the reverse complementary sequence of the nucleotide triplet in positions +8 to +10 of said genomic target, (v) the nucleotide doublet(s) in positions +1 to +2, +6 to +7, and/or +11 to +12 is identical to the nucleotide doublet(s) which is present in positions +1 to +2, +6 to +7, and/or +11 to +12, respectively of said genomic target, (vi) the nucleotide doublet(s) in positions -12 to -11, -7 to -6 and/or -2 to -1 is identical to the reverse complementary sequence of the nucleotide doublet(s) which is present in positions +11 to +12, +6 to +7, and/or +1 to +2, respectively of said genomic target,

(l₃) combining the variants obtained in steps (j₃) and (k₃) to form heterodimers, and

(m₁) selecting and/or screening the heterodimers from step (l₃) which are able to cleave said genomic DNA target situated in a gene of interest.

5 The steps (a₁), (a₂), (b₁), (b₂), (a₃), (b₃), (c₃), (g₁), (g₂), (h₁), (h₂), (i₁), (j₃), and (k₃) may comprise the introduction of additional mutations in order to improve the binding and/or cleavage properties of the mutants, particularly at other positions contacting the DNA target sequence or interacting directly or indirectly with said DNA target. These steps may be performed by generating a combinatorial library
10 as described in the International PCT Application WO 2004/067736, Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458 and Smith *et al.*, Nucleic Acids Research, Epub 27 November 2006.

 Steps (g₁), (g₂), (h₁), (h₂), (i₁), (j₃) and (k₃), may further comprise the introduction of random mutations on the whole variant or in a part of the variant, in
15 particular the C-terminal half of the variant (positions 80 to 163). This may be performed by generating random mutagenesis libraries on a pool of variants, according to standard mutagenesis methods which are well-known in the art and commercially available.

 Step (i₁) may also comprise the selection and/or screening of the
20 homodimers which are able to cleave a sequence wherein the nucleotide doublet in positions +1 to +2, +6 to +7 and/or +11 to +12 is identical to the nucleotide doublet which is present in positions +1 to +2, +6 to +7 and/or +11 to +12, respectively of said genomic target, and the nucleotide doublet in positions -12 to -11, -7 to -6, and/or -2 to -1 is identical to the reverse complementary sequence of the nucleotide triplet
25 which is present in positions +11 to +12, +6 to +7, and/or +1 to +2, respectively of said genomic target.

 The (intramolecular) combination of mutations in steps (g₁), (g₂), (h₁), (h₂), (j₃) and (k₃) may be performed by amplifying overlapping fragments comprising each of the two subdomains, according to well-known overlapping PCR
30 techniques, as described for example in Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006.

The (intermolecular) combination of the variants in step (j₁), (i₂) and (l₃) is performed by co-expressing one variant from step (g₁), (g₂) or (i₁), (j₃) with one variant from step (h₁), (h₂) or (i₁), (k₃), respectively, so as to allow the formation of heterodimers. For example, host cells may be modified by one or two recombinant expression vector(s) encoding said variant(s). The cells are then cultured under conditions allowing the expression of the variant(s), so that heterodimers are formed in the host cells, as described previously in Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458; International PCT Applications WO 2006/097853, WO 2006/097854 and WO 2006/097784; Smith *et al.*, Nucleic Acids Res., Epub 27 November 2006.

The selection and/or screening steps may be performed by using a cleavage assay *in vitro* or *in vivo*, as defined above. Preferably, it is performed *in vivo*, under conditions where the double-strand break in the mutated DNA target sequence which is generated by said variant leads to the activation of a positive selection marker or a reporter gene, or the inactivation of a negative selection marker or a reporter gene, by recombination-mediated repair of said DNA double-strand break, as defined above.

The subject-matter of the present invention is also the use of at least one meganuclease, as defined above, as a scaffold for making other meganucleases. For example other rounds of mutagenesis and selection/screening can be performed on the variant, for the purpose of making novel homing endonucleases.

The subject-matter of the present invention is also a method for decreasing the toxicity of a parent LAGLIDADG homing endonuclease, comprising: the mutation of at least one amino acid of the final C-terminal loop of said parent LAGLIDADG homing endonuclease.

According to an advantageous embodiment of said method the parent endonuclease is I-CreI or a functional variant thereof. Preferably, the K139 and/or T143 residues are mutated. More preferably K139 is mutated in an hydrophobic amino acid such as a methionine (K139M) and/or T143 is mutated in a small amino acid such as a glycine (T143G).

The polynucleotide fragments having the sequence of the targeting DNA construct or the sequence encoding the meganuclease variant or single-chain meganuclease derivative as defined in the present invention, may be prepared by any

method known by the man skilled in the art. For example, they are amplified from a DNA template, by polymerase chain reaction with specific primers. Preferably the codons of the cDNAs encoding the meganuclease variant or single-chain meganuclease derivative are chosen to favour the expression of said proteins in the
5 desired expression system.

The recombinant vector comprising said polynucleotides may be obtained and introduced in a host cell by the well-known recombinant DNA and genetic engineering techniques.

The meganuclease variant or single-chain meganuclease derivative
10 as defined in the present the invention are produced by expressing the polypeptide(s) as defined above; preferably said polypeptide(s) are expressed or co-expressed (in the case of the variant only) in a host cell or a transgenic animal/plant modified by one expression vector or two expression vectors (in the case of the variant only), under conditions suitable for the expression or co-expression of the polypeptide(s), and the
15 meganuclease variant or single-chain meganuclease derivative is recovered from the host cell culture or from the transgenic animal/plant.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are
20 within the skill of the art. Such techniques are explained fully in the literature. See, for example, Current Protocols in Molecular Biology (Frederick M. AUSUBEL, 2000, Wiley and son Inc, Library of Congress, USA); Molecular Cloning: A Laboratory Manual, Third Edition, (Sambrook et al, 2001, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press); Oligonucleotide Synthesis (M. J. Gait ed., 1984);
25 Mullis et al. U.S. Pat. No. 4,683,195; Nucleic Acid Hybridization (B. D. Harries & S. J. Higgins eds. 1984); Transcription And Translation (B. D. Hames & S. J. Higgins eds. 1984); Culture Of Animal Cells (R. I. Freshney, Alan R. Liss, Inc., 1987); Immobilized Cells And Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide To Molecular Cloning (1984); the series, Methods In ENZYMOLOGY (J. Abelson and
30 M. Simon, eds.-in-chief, Academic Press, Inc., New York), specifically, Vols.154 and 155 (Wu et al. eds.) and Vol. 185, "Gene Expression Technology" (D. Goeddel, ed.); Gene Transfer Vectors For Mammalian Cells (J. H. Miller and M. P. Calos eds., 1987,

Cold Spring Harbor Laboratory); Immunochemical Methods In Cell And Molecular Biology (Mayer and Walker, eds., Academic Press, London, 1987); Handbook Of Experimental Immunology, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); and Manipulating the Mouse Embryo, (Cold Spring Harbor Laboratory Press, 5 Cold Spring Harbor, N.Y., 1986).

In addition to the preceding features, the invention further comprises other features which will emerge from the description which follows, which refers to examples illustrating I-CreI meganuclease variants and their uses according to the invention, as well as to the appended drawings in which:

10 - figure 1 represents the superposition of the C α ribbon representation of the I-CreI and I-CreI-DNA structures. DNA has been omitted for clarity.

 -figure 2 represents the sequence alignment of the C-terminal region from members of the I-CreI family (Lucas *et al.*, Nucleic Acids Res., 2001, 29, 960-15 969). The position of the mutated residues in the SKTRKTT motif is indicated with a grey triangle (<http://esprict.ibcp.fr/ESPript/cgi-bin/ESPript.cgi>).

 - figure 3 represents a detailed view of S138, K139, K142 and T143 contacts with the DNA backbone (a) and the comparison of the positions of S138, K139, K142 and T143 between the bound and unbound DNA structures (b).

20 - figure 4 illustrates the biophysical characterization of the I-CreI C-terminal region mutants. a) Circular dichroism thermal denaturation. b) Monodimensional H-H NMR spectra.

 - figure 5 illustrates dimer formation by the I-CreI C-terminal region mutants, measured by analytical ultracentrifugation. Sedimentation velocity 25 distribution of the I-CreI proteins (1 mg/ml in PBS buffer) at 42,000 rpm and 20°C. Inset, sedimentation equilibrium gradient of I-CreI proteins (4 mg/ml in PBS buffer) at 11,000 rpm and 20 °C. Open circles represent the experimental data, the two solid lines represent the theoretical gradients of a I-CreI monomer (20,045) and dimer (41,000).

30 - figure 6 represents electrophoretic mobility shift assays of the C-terminal truncated, double and single mutants in the presence of Mg²⁺ and Ca²⁺.

- figure 7 is a summary of the gel *in vitro* cleavage assay of the C-terminal truncated, double and single mutants.

- figure 8 illustrates the *in vivo* cleavage assay used for profiling the single mutants and the 10NNN_P DNA target cleavage profile of the single mutants.

5 a) Yeast screening assay principle. A strain harboring the expression vector encoding a single mutant is mated with a strain harboring a reporter plasmid. In the reporter plasmid, a LacZ reporter gene is interrupted with an insert containing one of the target sites of interest, flanked by two direct repeats. Upon mating, the meganuclease (grey oval) generates a double-strand break at the site of interest, allowing restoration of a
10 functional LacZ gene by single-strand annealing (SSA) between the two flanking direct repeats. The functional LacZ gene is visualized by a blue staining. b) DNA targets. The C1221 target (top) is a palindromic target cleaved by I-CreI. All targets used in this study are palindromic targets derived from C1221 by substitution of six nucleotides in ± 8 , ± 9 and ± 10 (SEQ ID NO: 1 and 10 to 16). A few examples are
15 shown (bottom). The 10GGG_P target differs from the C1221 target by the GGG triplet in -10, -9, -8 and CCC in +8, +9 and +10. c) Mutant target profiles. Each mutant was profiled in yeast on a series of 64 palindromic targets (10NNN_P). An example of cleavage activity in yeast for a single mutant (K139M) compared to I-CreI D75N is presented. Blue staining indicates cleavage. Additionally a representation of
20 the 10NNN_P cleavage profile of all single mutants compared to I-CreI D75N and I-CreI. Grey levels reflect the intensity of the signal. I-CreI is toxic in yeast and profiles have been established at 30 °C instead of 37° C. All other mutants were studied at 37°C.

25 - figure 9 illustrates the 5NNN_P DNA target cleavage profile of the single mutants. The targets (64) are palindromic targets with variations in positions ± 3 to 5).

- figure 10 illustrates the 2NN DNA target cleavage profile of the single mutants. The targets (16 $\times$ 16) are non-palindromic targets with variations in positions ± 1 to 2.

30 - figure 11 illustrates the 12NN_P DNA target (A) and 7NN_P DNA target (B) cleavage profiles of the single mutants. The targets in A (16) and B (16) are palindromic targets with variations in positions ± 11 to 12 and ± 6 to 7, respectively.

Example 1: Structural differences between the bound and unbound I-CreI DNA structures

1) Materials and Methods

a) Protein expression, purification and crystallization

5 Protein expression and purification was performed as in (Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458). An initial screening for I-CreI crystallization conditions was performed in 96 well plates by vapour-diffusion methods using the Hampton crystal screening using drops containing 1 µl protein solution (7 mg/ml in 20 mM HEPES, pH 7.5) and 1 µl precipitant solution equilibrated against 50 µl of
10 reservoir solution at 20° C. Crystals were obtained under several conditions (Crystal Screen 1 conditions 10, 22, 33, 40, 41 and Crystal Screen 2 condition 32). Crystal was made by hanging-drop vapour-diffusion methods using VDX plates; optimization experiments led to the following conditions for crystallization: 1 µl protein at 7 mg/ml in 20 mM HEPES pH 7.5 and 1 µl precipitating buffer containing 20 % PEG 4000, 0.1
15 M HEPES pH 7.5, 10 % Iso-propanol, 10 % Ethylene glycol and 0.01 M Magnesium acetate equilibrated against 500 µl precipitating buffer at 20° C. Rod-shaped crystals grown in 4-8 days and were directly collected and frozen in liquid nitrogen.

b) Data collection, structure solution, model building and refinement

All data were collected at cryogenic temperatures using synchrotron
20 radiation at 100K. I-CreI crystals were mounted and cryoprotected. The data sets were collected using synchrotron radiation at the ID14-4 beamline at the ESRF (Grenoble), and at the PX beamline at the SLS (Villigen). Diffraction data were recorded on an ADSC-Q4 or Mar225 CCD detectors depending on the beamline. Processing and scaling were accomplished with HKL2000 (Otwinowski, Z. and
25 Minor, W.: *Processing of X-ray Diffraction Data Collected in Oscillation Mode, In Methods in Enzymology*, 1997, Academic Press, New York). The structure was solved using the molecular replacement method as implemented in the program MOLREP (Vagin, A. and Teplyakov, A. Acta Crystallogr. D Biol. Crystallogr., 2000, 56 Pt 12, 1622-1624).

30 2) Results

The structure of the I-CreI was solved by molecular replacement and refinement to 2.0Å resolution. The best data set (Table I) was collected using a

$\Delta\phi = 1^\circ$ and a wavelength of 0.97 Å. Statistics for the crystallographic data are summarized in Table I. The search model was based on a poly-alanine backbone derived from the PDB 1gz9 found in the Protein Data Bank. The coordinates from the DNA were deleted in the search model. A refined 2Fo-Fc map showed clear and contiguous electron density for the protein backbone and for many of the side-chains. ARP/wARP and REFMAC5 were applied for automatic model building and refinement to 2.0 Å (Table I).

Table I: Data collection and refinement statistics

Data Collection	
Space group	P4 ₃
Number of crystals	1
Temperature (K)	100
Wavelength (Å)	0.97
Cell dimensions (Å, °),	a=b=69.088, c=93.040 $\alpha\beta\gamma=90^\circ$
No. mol ASU	2
Data collection environment, beamline	ADSC-Q4, ID14-4 ESRF
Completeness (%)	93.2
Multiplicity	5.3
Rsym(%)	6
Refinement	
No Reflections	25943
Resolution range (Å)	34.54-2.00
R-factor / R-free (%)	18/23
No protein atoms (Average B, Å ²) ^c	
No water molecules (Average B, Å ²) ^c	
r.m.s bond length (Å)	0.029
r.m.s. bond angle (°)	2.137
Ramachandran plot outliers (number) ^d	0

The dimer without DNA allowed the observation of the protein conformational changes upon DNA binding after comparison with the protein-DNA complex (PDB code 1gz9) (Figure 1). The most striking differences are in the C-terminal region conformation. Whereas in the DNA bound structure the C-helix and the C-loop are aligned with the DNA, in the unbound structure both elements are located on top of the cavity where the DNA binds, suggesting that the loop and the C-helix could work as a lock opening and closing the DNA binding groove. This region was not observed in a previous structure of I-CreI with only one monomer in the asymmetric unit (Heath *et al.*, Nat Struct Biol, 1997, 4, 468-476). Besides, the C-terminal domain of I-CreI is well conserved among homodimeric proteins from the

LAGLIDADG family (Lucas *et al.*, Nucleic Acids Res., 2001, 29, 960-969) indicating its important role in this meganuclease group working mechanism (Figure 2). A detailed view of the protein-DNA interactions in the C-terminal area showed that Ser138, Lys139, Lys142 and Thr143 at the SKTRKTT motif are involved in hydrogen bonds with the DNA backbone (Figure 3a). The position of these residues is completely different in the unbound DNA state (Figure 3b), indicating that a conformational change is needed to bind the nucleic acid. Although these interactions were described before (Chevalier *et al.*, J. Mol. Biol., 2003, 329, 253-269) and the amino acids are conserved, there is no information about their role during meganuclease action.

Example 2: Biophysical analysis

1) Materials and Methods

a) Construction of the I-CreI mutants

The I-CreI deletion mutants ($\Delta 1$ and $\Delta 2$) were amplified by PCR on the wild-type I-CreI (I-CreI D75) cDNA template, with the forward primer 5' gatataccatggccaataccaaatataac 3' (SEQ ID NO: 18) for both mutants and the reverse primer ICreI deltaCter-R: 5' ttatcagtcggccgcgcgttcagagctgcaatctgatccaccagg 3' (SEQ ID NO: 19) for the $\Delta 1$ mutant or Creh2: 5' gagtgcggccgcagtggtttacgcgtcttagaatcg 3' (SEQ ID NO: 20) for the $\Delta 2$ mutant.

The I-CreI single and double mutants were amplified by round-the-world PCR with a Quickchange® kit (STRATAGENE # 200518), appropriate mutagenizing oligos and the wild-type I-CreI (I-CreI D75) cDNA as template.

b) Circular Dichroism thermal analysis

Data were acquired with a Jasco 810 model dichrograph, previously calibrated with *d*-10-camphorsulphonic acid, and equipped with a Jasco Peltier thermoelectric temperature controller CDF-426S model. Experiments were performed in PBS at 1° C/min intervals. The protein concentration was 10 μ M. The ellipticity at 222 nm was followed from 5 to 95 °C in a 2 mm Hellma 110-QS cell.

c) Analytical ultracentrifugation

Sedimentation equilibrium experiments were performed at 20 °C in an Optima XL-A (Beckman-Coulter) analytical ultracentrifuge equipped with UV-visible optics, using an An50Ti rotor, with 3,mm double sector centerpieces of Epon

charcoal. Protein concentration was 200 μ M in PBS buffer. Short column (23 μ l), low speed sedimentation equilibrium was performed at three successive speeds (11,000, 13,000, and 15,000 rpm), the system was assumed to be at equilibrium when successive scans overlaid and the equilibrium scans were obtained at wavelength of 280 nm. The base-line signal was measured after high speed centrifugation (5 h at 42,000 rpm). Whole-cell apparent molecular weight of the protein was obtained using the program EQASSOC (Minton, A.P., In: *Modern Analytical Ultracentrifugation*, 1994, Birkhauser Boston, Inc., Cambridge, MA). The partial specific volume of I-CreI was 0.7436 ml/g at 20 °C, calculated from the amino acid composition with the program SEDNTERP (retrieved from the RASMB server; Laue, T.M.S., B.D., Ridgeway, T.M., Pelletier, S.L., In: *Computer-aided interpretation of analytical sedimentation data for proteins*, 1992, Royal Society of Chemistry, Cambridge, UK). The sedimentation velocity experiment was carried out in an XL-A analytical ultracentrifuge (Beckman-Coulter Inc.) at 42,000 rpm and 20° C, using an An50Ti rotor and 1.2mm double-sector centerpieces. Absorbance scans were taken at 280 nm. The protein concentration was 50 μ M in PBS. The sedimentation coefficients were calculated by continuous distribution c(s) Lamm equation model (Schuck, P., Biophys. J., 2000, 78, 1606-1619) as implemented in the SEDFIT program. These experimental sedimentation values were corrected to standard conditions to get the corresponding $s_{20,w}$ values using the SEDNTERP program (Laue, T.M.S., B.D., Ridgeway, T.M., Pelletier, S.L., In: *Computer-aided interpretation of analytical sedimentation data for proteins*, 1992, Royal Society of Chemistry, Cambridge, UK). Further hydrodynamic analysis (i.e. calculation of frictional coefficient ratio) was performed with the SEDFIT program to obtain de c(M) distribution (Schuck, P., Biophys. J., 2000, 78, 1606-1619).

d) NMR data acquisition

NMR spectra were recorded at 25 °C in a Bruker AVANCE 600 spectrometer equipped with a cryoprobe. Protein samples were 500 μ M in PBS buffer (137 mM NaCl, 10 mM Na₂HPO₄-2H₂O, 2.7 mM KCl, 2 mM KH₂PO₄, pH 7.4) plus 5 % ²H₂O. DSS (2,2-Dimethyl-2-silapentane-5-sulfonate sodium salt) was used as internal proton chemical shift reference.

2) Results

To unravel the role of the C-terminal domain of I-CreI, a series of trimmed, double and single mutants were designed based on the structural differences between the bound and unbound DNA structures. The two truncated mutants were designed to clarify the role of the C-terminal region. I-CreI $\Delta 1$ (amino acid number 1-137) lacked both the C-loop and the C-Helix whereas I-CreI $\Delta 2$ (amino acid number 1-144) contained the C-loop. Based on the contacts with the DNA backbone in the SKTRKTT motif, the double mutants I-CreI AM (S138A, K139M) and I-CreI GG (K142G, T143G) were produced, as well as their single variants I-CreI S138A, I-CreI K139M, I-CreI K142G, I-CreI T143G. To demonstrate that the effect in meganuclease activity was due to the mutations, their effect in the protein stability, structure and oligomerization state, was studied. Thermal denaturation circular dichroism was performed to confirm that all the mutants were folded. Indeed, all the mutants displayed a sigmoidal curve similar to the wild type (Figure 4a) with different T_m depending on the mutation. In addition, monodimensional H-H NMR confirmed the thermal denaturation experiments, a well defined dispersion of peaks in the amide region demonstrated that all the mutants were folded (Figure 4b). It is well known that the I-CreI family of meganucleases binds DNA as homodimers, therefore to analyze the oligomerization state of the mutants they were subjected to analytical ultracentrifugation. The experiment showed that all the mutants behaved as dimers independently of the mutation, with only little variations corresponding to their molecular weights (Figure 5).

Altogether these experiments indicate that the mutants are folded and conserve the I-CreI scaffold involved in meganuclease activity.

Example 3: DNA-binding activity the C-terminal mutants

1) Materials and Methods

Band shift assay conditions

Band shift assays were performed in 10 mM Tris-HCl pH 8, 50 mM NaCl, 10 mM CaCl₂ or MgCl₂, 1 mM DTT incubated 1 h at room temperature using 5 μ M (0.0793 μ g/ μ l) 6-FAM duplex (SEQ ID NO: 21; see figure 6) and 20 μ M (0.463 μ g/ μ l) protein and electrophoresed in a 15 % Acrylamide-TBE gel.

2) Results

Electrophoretic mobility shift assays (EMSA) in the presence of Mg^{2+} and Ca^{2+} were used to analyze the behavior of the C-terminal mutants in DNA binding (Figure 6). Whereas the presence of Ca^{2+} allows DNA binding, Mg^{2+} is indispensable to bind and cleave DNA (Chevalier *et al.*, Biochemistry, 2004, 43, 14015-14026). Even though the binding capability of I-CreI was abolished in the $\Delta 1$ mutant, the $\Delta 2$ was able to bind the labeled DNA probe demonstrating that the C-loop is essential in DNA binding. In addition, binding was detected in the presence of both cations as in the wild type I-CreI.

On the other hand, both I-CreI AM and I-CreI GG double mutants were severely affected in their DNA binding properties independently of the cation present, indicating that Ser138, Lys139, Lys142 and Thr143 contacts with the DNA backbone are crucial to bind the nucleic acid. Therefore, these residues in the SKTRKTT motif constitute two new hot-spots essential for I-CreI DNA binding.

To define the distinct properties of each site in the C-loop, the single mutants were assayed by EMSA in the same conditions. In contrast with the double mutants all the single ones were able to bind the labeled probe; however they displayed differences depending of the cation present in the assay. Whereas a clear dependence of Mg could be observed in the Ser138-Lys139 site, the single mutants in the Lys142-Thr143 site could bind DNA notwithstanding the cation present in the mobility assay.

Thus, the mutation of both residues in each site is needed to abolish DNA binding, indicating that a synergy between the two residues in each hot-spot is essential for DNA binding.

Example 4: DNA-cleavage activity of the C-terminal mutants, *in vitro*

1) Materials and Methods

In vitro cleavage assay conditions

Cleavage assays were performed at 37 °C in 10 mM Tris-HCl (pH 8), 50 mM NaCl, 10 mM $MgCl_2$ (or $CaCl_2$) and 1 mM DTT. Concentrations were: 100 ng for the *XmnI* linearized target substrate (pGEM-T Easy C1221 GTC) and 40-0.25ng dilutions for I-CreI and helix mutant proteins, in 25 μ l final volume reaction. The linearized target plasmid has 3 kb and after cleavage yields two smaller bands of 2 kb

and 1 kb. Reactions were stopped after 1 hour by addition of 5 μ l of 45 % Glycerol, 95 mM EDTA (pH 8), 1.5 % (w/v) SDS, 1.5 mg/ml Proteinase K and 0.048 % (w/v) Bromophenol blue (6x Buffer Stop), incubated at 37 °C for 30 minutes and electrophoresed in a 1 % agarose gel. The fragments were quantified using SYBR Safe DNA gel staining (IN VITROGEN). Gels were analysed using the ImageJ software (<http://rsb.info.nih.gov/ij/>) to calculate the percentage of cleavage according to $(2\text{kb}+1\text{kb})/(3\text{kb}+2\text{kb}+1\text{kb}) \times 100$ formula.

2) Results

The analysis of the distinct mutants in the DNA binding assays has clear implications for DNA cleavage activity, consequently an examination of their cleavage properties on a wild type DNA sequence was carried out. Figure 7 displays a graph representing the percentage of cleavage against the amount of HE (Gels with raw data are available as supporting information). The mutants can be divided in two groups based on the comparison of their cleavage properties to the wild type HE; the first is composed of the truncated mutants I-CreI $\Delta 1$ and I-CreI $\Delta 2$ and the double mutants I-CreI AM and I-CreI GG which are , whereas the single mutants I-CreI S138A, I-CreI K139M, I-CreI K142G, I-CreI T143G form the second. Members the first group displayed a reduced cleavage activity when compared to the wild type I-CreI. Although I-CreI $\Delta 1$ and I-CreI GG cleavage properties are completely abolished, I-CreI $\Delta 2$ and I-CreI AM showed a reduced activity that is increased when higher HE amounts are used. However the cleavage properties of the single mutants that composed the second group are not only similar to the wild type, but enhanced in some cases (Figure 7).

These results indicate that the trimmed and double mutants whose DNA binding is abolished or severely affected do not cleave DNA or they need higher amounts of HE to cleave the plasmid. Noteworthy is the case of the I-Cre $\Delta 2$, the mutant that conserves the wild type amino acids in the C-loop but lacks the $\alpha 6$ helix, even though its cleavage activity is affected the activity profile is the more similar to the I-CreI wild type.

On the other hand the single mutants depict a slightly enhanced activity with respect to the wild type in all them. The activity assays confirm the DNA binding studies, indicating that the double mutants act in a concerted manner, however

the effect of these mutations have implications not only in nucleic acid binding but also in DNA cleavage as we have shown.

Example 5: DNA-cleavage activity of the C-terminal mutants, *in vivo*

1) Materials and Methods

5 The *in vivo* cleavage assay (Figure 8a) has been described previously in PCT Application WO 2004/067736; Epinat *et al.*, Nucleic Acids Res., 2003, 31, 2952-2962; Chames *et al.*, Nucleic Acids Res., 2005, 33, e178, and Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458.

a) Construction of target clones

10 The C1221 twenty-four bp target sequence (5'-tcaaaacgtcgtacgacgtttga-3': SEQ ID NO: 1) is a palindrome of a half-site of the natural I-CreI target (5'-tcaaaacgtcgtgagacagtttg-3': SEQ ID NO: 17). C1221 is cleaved as efficiently as the I-CreI natural target *in vitro* and *ex vivo* in both yeast and mammalian cells. The palindromic targets, derived from C1221, were cloned as
15 previously described (Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458) using the Gateway protocol (Invitrogen) into the reporter vectors: the yeast pFL39-ADH-LACURAZ and the mammalian vector pcDNA3.1-LACURAZ-ΔURA, both described previously (Epinat *et al.*, Nucleic Acids Res., 2003, 31, 2952-2962) and containing a I-SceI target site as control. Yeast reporter vectors were transformed into *S. cerevisiae*
20 strain FYBL2-7B (*MAT* a, *ura3*Δ851, *trp1*Δ63, *leu2*Δ1, *lys2*Δ202).

b) Screening in yeast

 The protocol for screening homodimer mutants is as described previously (PCT Application WO 2004/067736; Epinat *et al.*, Nucleic Acids Res., 2003, 31, 2952-2962; Chames *et al.*, Nucleic Acids Res., 2005, 33, e178; Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458).
25

c) Mating of meganuclease expressing clones and screening in yeast

 Mating was performed using a colony gridder (QpixII, Genetix). Mutants were gridded on nylon filters covering YPD plates, using a high gridding density (about 20 spots/cm²). A second gridding process was performed on the same
30 filters to spot a second layer consisting of 64 or 75 different reporter-harboring yeast strains for each variant. Membranes were placed on solid agar YPD rich medium, and incubated at 30 °C for one night, to allow mating. Next, filters were transferred to

synthetic medium, lacking leucine and tryptophan, with galactose (2 %) as a carbon source, and incubated for five days at 37 °C (30 °C for I-CreI), to select for diploids carrying the expression and target vectors. After 5 days, filters were placed on solid agarose medium with 0.02 % X-Gal in 0.5 M sodium phosphate buffer, pH 7.0, 0.1 % SDS, 6 % dimethyl formamide (DMF), 7 mM β -mercaptoethanol, 1 % agarose, and incubated at 37 °C, to monitor β -galactosidase activity. Results were analyzed by scanning and quantification was performed using appropriate software.

2) Results

To confirm the results *in vivo* cleavage assays were performed with all the mutants, as well as with the I-CreI and I-CreI D75N proteins, as described previously (Arnould *et al.*, J. Mol. Biol., 2006, 355, 443-458).

None of the double mutants or truncated mutants, whose binding and cleavage activity were affected by the mutations *in vitro*, displayed activity on any of the targets. As an example the I-CreI AM and I-CreI GG are shown (Figure 8c upper panel). By contrast, all the single mutants showed high activity on the wild-type target (C1221).

- 10NNN_P target profiling (figure 8c)

All the single mutants showed high activity on the wild-type target (C1221), 10AAG_P and 10AAT_P. Lower levels of cleavage could also be observed with these four mutants with 10TCG_P and 10AAC_P. In addition the I-CreI K139M mutant was also able to cleave seven additional targets (10AGT_P, 10GAG_P, 10GAA_P, 10GAT_P, 10CAG_P, 10CAA_P, 10CAT_P) as it can be observed in figure 8c. The profile of the I-CreI K139M mutant is very similar to I-CreI (without its toxicity), while the three other single mutants are closer to I-CreI D75N.

- 5NNN_P profiling (figure 9)

The profile of S138A and K139M is similar to the profile of I-CreI D75N, whereas the profile of K142G and T143G is more restricted than the profile of I-CreI D75N.

- 2NN_P profiling (figure 10)

The profile of K142G and S138A is more restricted than the profile of I-CreI D75N. Compared to D75N, T143G and K139M cleave 6 and 10 additional targets, respectively, 6 of which are in common. In addition, at least 8 targets are

cleaved more efficiently by K139M than by D75N. Five targets (2TT_2TG, 2TG_2TT, 2TA_2CT, 2TC_2TC, 2CT_2CT) are not cleaved by K139M; these targets are cleaved by D75N, although less efficiently than by I-CreI.

- 12NN_P profiling (Figure 11A)

5 The profile of K142G and S138A is more restricted than the profile of I-CreI D75N, with the profile of S138A being more restricted than the profile of K142G.

 The profile of T143G is similar to the profile of I-CreI D75N.

 The profile of K139M is similar to the profile of I-CreI but without
10 its toxicity; 7 additional targets are cleaved by K139M as compared to D75N.

- 7NN_P profiling (Figure 11B)

 The profile of K142G and S138A is similar to the profile of I-CreI D75N.

 K139M and T143G cleave 2 additional targets (7CG_P and 7TT_P)
15 as compared to D75N; however the cleavage profile of K139M and T143G is more restricted than the profile of I-CreI.

 These results demonstrate that the C-terminal region of I-CreI is essential for HE activity. Moreover the mutations in the flanking residues of the SKTRKTT region demonstrate that they control not only nucleic acid binding, but
20 also target specificity.

CLAIMS

1°) A method for engineering a LAGLIDADG homing endonuclease variant having novel substrate specificity, comprising at least the following steps:

5 (a) the mutation of at least one amino acid residue of the final C-terminal loop of a parent LAGLIDADG homing endonuclease, with the exclusion of the threonine 140 of I-*CreI*, and

b) the selection and/or screening of the variants from step (a) having a pattern of cleaved DNA targets that is different from that of the parent LAGLIDADG homing endonuclease.

10 2°) The method of claim 1, wherein said mutation(s) are in positions of amino acid residue(s) of the final C-terminal loop which are contacting the phosphate backbone of the parent LAGLIDADG homing endonuclease DNA cleavage site.

15 3°) The method of claim 2, wherein said mutation(s) modify the interaction between said amino acid residue(s) of the final C-terminal loop and the phosphate backbone of the parent LAGLIDADG homing endonuclease DNA cleavage site.

20 4°) The method of claim 2 or claim 3, wherein said mutation(s) are in positions 138, 139, 142 and/or 143, by reference to I-*CreI* amino acid sequence numbering.

5°) The method of claim 4, wherein the residues in positions 138 and/or 139 are substituted by an hydrophobic amino acid and/or the residues in positions 142 and/or 143 are substituted by a small amino acid.

25 6°) The method of claim 5, wherein the residue in position 138 is substituted by an alanine, the residue in position 139 is substituted by a methionine, and/or the residues in positions 142 and/or 143 are substituted by glycines.

7°) The method of anyone of claims 4 to 6, wherein step (a) comprises the mutation of two residues, each one from a different pair chosen from the residues in positions 138 and 139 and the residues in positions 142 and 143.

30 8°) The method of anyone of claims 1 to 7, wherein said parent LAGLIDADG homing endonuclease is an homodimeric LAGLIDADG homing endonuclease.

9°) The method of claim 8, wherein said homodimeric LAGLIDADG homing endonuclease is I-CreI.

10°) The method of claim 8, wherein said homodimeric LAGLIDADG homing endonuclease is an I-CreI variant having mutations in positions 26 to 40 and 44 to 77 of I-CreI and cleaving a palindromic DNA sequence, wherein at least the nucleotides in positions + 3 to + 5 and + 8 to +10 or - 10 to - 8 and -5 to -3 of one half of said DNA sequence correspond to the nucleotides in positions + 3 to + 5 and + 8 to +10 or - 10 to - 8 and -5 to -3 of one half of a genomic DNA target from a gene of interest.

11°) The method of anyone of claims 1 to 10, wherein step (a) comprises, simultaneously or subsequently, the mutation of at least one amino acid residue in a first functional subdomain corresponding to that situated from positions 26 to 40 of I-CreI amino acid sequence, that alter the specificity towards the nucleotide in positions ± 8 to 10 of the DNA target, and/or the mutation of at least amino acid residue in a second functional subdomain corresponding to that situated from positions 44 to 77 of I-CreI amino acid sequence, that alter the specificity towards the nucleotide in positions ± 3 to 5 of the DNA target.

12°) The method of anyone of claims 1 to 11, wherein step (a) comprises, simultaneously or subsequently, the random mutation of the whole or the C-terminal half of said LAGLIDADG homing endonuclease/variant amino acid sequence.

13°) The method of anyone of claims 1 to 12, wherein step (b) comprises the selection and/or screening of the variants from step (a) which are able to cleave at least one DNA target sequence that is not cleaved by said parent LAGLIDADG homing endonuclease, said DNA target sequence being derived from the parent LAGLIDADG homing endonuclease cleavage site, by the replacement of at least one nucleotide of one half of said cleavage site, with a different nucleotide.

14°) The method of claim 13, wherein said DNA target sequence is derived from the I-CreI palindromic site having the sequence SEQ ID NO: 1.

15°) The method of claim 14, wherein said DNA target has mutation(s) in the nucleotide(s) in positions ± 1 to 2, ± 6 to 7, ± 8 to 10 and/or ± 11 to 12.

16°) The method of anyone of claims 11 to 15, wherein said DNA target sequence is a genomic sequence which is present in a gene of interest.

17°) An homodimeric or heterodimeric LAGLIDADG homing endonuclease variant which is obtainable by the method of anyone of claims 1 to 16,
5 with the exclusion of the homodimeric variants of SEQ ID NO: 3 and 4 and the homo- and hetero-dimeric variants comprising a monomer of SEQ ID NO: 5.

18°) The variant of claim 17, which is an heterodimer comprising the monomers of two different variants obtainable by said method.

19°) The variant of claim 17 or claim 18, which is an I-*CreI* variant
10 having one or two mutations, each one from a different pair of mutations selected from the group consisting of the pair S138A and K139M and the pair K142G and T143G.

20°) The variant of claim 19, which is of the sequence SEQ ID NO: 6 to 9.

21°) The variant of claim 19 or claim 20, which is an heterodimeric I-*CreI* variant consisting of two monomers, each monomer further comprising different mutations in positions 26 to 40 and 44 to 77 of I-*CreI*, said variant being able to cleave a genomic DNA target from a gene of interest.

22°) A single-chain chimeric meganuclease comprising two
20 monomers or core domains of one or two variants of anyone of claims 17 to 21, or a combination of both.

23°) A polynucleotide fragment encoding one monomer of the variant of anyone of claims 17 to 21 or the single-chain meganuclease of claim 22.

24°) A recombinant vector comprising at least one polynucleotide
25 fragment of claim 23.

25°) An expression vector comprising two polynucleotide fragments each encoding one of the two monomers of an heterodimeric variant of anyone of claims 17 to 21, said fragment(s) being operatively linked to regulatory sequences allowing the production of the two monomers.

26°) An expression vector comprising a polynucleotide fragment
30 encoding the single-chain meganuclease of claim 22, said fragment being operatively

linked to regulatory sequences allowing the production of said single-chain meganuclease.

27°) The vector of claim 25 or claim 26, which includes a targeting DNA construct comprising sequences sharing homologies with the region surrounding the genomic DNA target sequence as defined in anyone of claims 10, 16 and 21.

28°) The vector of claim 27, wherein said targeting DNA construct comprises : a) sequences sharing homologies with the region surrounding the genomic DNA target sequence as defined in anyone of claims 10, 16 and 21, and b) sequences to be introduced flanked by sequence as in a).

29°) A host cell comprising one or two polynucleotide fragments as defined in claim 23 or claim 25 or a vector according to anyone of claims 24 to 28.

30°) A non-human transgenic animal comprising one or two polynucleotide fragments as defined in claim 23 or claim 25.

31°) A transgenic plant comprising one or two polynucleotide fragments as defined in claim 23 or claim 25.

32°) A pharmaceutical composition comprising at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 25 to 28.

33°) The composition of claim 32, further comprising a targeting DNA construct comprising the sequence which repairs the genomic site of interest flanked by sequences sharing homologies with the targeted locus.

34°) Use of at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 25 to 28, a host cell of claim 29, a transgenic plant of claim 31, a non-human transgenic mammal of claim 30, for molecular biology, for *in vivo* or *in vitro* genetic engineering, and for *in vivo* or *in vitro* genome engineering, for non therapeutic purposes.

35°) Use of at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 25 to 28, for the preparation of a

medicament for preventing, improving or curing a genetic disease in an individual in need thereof, said medicament being intended to be administrated by any means to said individual.

36°) Use of at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 25 to 28, for the preparation of a medicament for preventing, improving or curing a disease caused by an infectious agent that presents a DNA intermediate, in an individual in need thereof, said medicament being intended to be administrated by any means to said individual.

37°) Use of at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 25 to 28, *in vitro*, for inhibiting the propagation, inactivating or deleting an infectious agent that presents a DNA intermediate, in biological derived products or products intended for biological uses or for disinfecting an object.

38°) The use of claim 36 or claim 37, wherein said infectious agent is a virus.

39°) The use of anyone of claims 34 to 37, wherein said variant, single-chain meganuclease, polynucleotides, vector, cell, transgenic plant or non-human transgenic mammal are associated with a targeting DNA construct as defined in claims 27, 28 or 33.

40°) Use of at least a variant of anyone of claims 17 to 21, a single-chain meganuclease of claim 22, one or two polynucleotide fragments as defined in claim 23 or claim 25, a vector of anyone of claims 24 to 28, as a scaffold for engineering other meganucleases.

41°) A method for decreasing the toxicity of a parent LAGLIDADG homing endonuclease, comprising : the mutation of at least one amino acid of the final C-terminal loop of said parent LAGLIDADG homing endonuclease, as defined in anyone of claims 1 to 7.

42°) The method of claim 41, wherein the parent LAGLIDADG homing endonuclease is as defined in anyone of claims 8 to 10.

43°) The method of claim 42, wherein the mutation is K139M and/or T143G.

1/12




Figure 1

3/12




Figure 3

4/12




Figure 4

5/12





Figure 5



Molar ratio [6-FAM:Prot] 1:2

[6FAM]GCAAAACGTCGTGAGACAGTTTGC 3' (SEQ ID NO:21)
CGTTTTCAGCACTCTGTCAAACG 5'

Figure 6




Figure 7




FIGURE 8

9/12




Figure 8 (continuation)

10/12




FIGURE 9

11/12




FIGURE 10




FIGURE 11

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2007/001527

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N9/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, Sequence Search, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/097784 A (COLLECTIS [FR]; PAQUES FREDERIC [FR]) 21 September 2006 (2006-09-21) cited in the application the whole document	
A	SMITH JULIANNE ET AL: "A combinatorial approach to create artificial homing endonucleases cleaving chosen sequences" NUCLEIC ACIDS RESEARCH, vol. 34, no. 22, December 2006 (2006-12), XP009091809 ISSN: 0305-1048 the whole document	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

7 November 2007

Date of mailing of the international search report

16/11/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schneider, Patrick

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2007/001527

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEVALIER BRETT ET AL: "Metal-dependent DNA cleavage mechanism of the I-CreI LAGLIDADG homing endonuclease" BIOCHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, PA, US, vol. 43, no. 44, 9 November 2004 (2004-11-09), pages 14015-14026, XP002412602 ISSN: 0006-2960 figure 1	
A	CHEVALIER B ET AL: "Flexible DNA Target Site Recognition by Divergent Homing Endonuclease Isoschizomers I-CreI and I-MsoI" JOURNAL OF MOLECULAR BIOLOGY, LONDON, GB, vol. 329, no. 2, 30 May 2003 (2003-05-30), pages 253-269, XP004454255 ISSN: 0022-2836 cited in the application the whole document	
E	WO 2007/047859 A (PREC BIOSCIENCES; HELLINGA HOMME W [US]; SMITH JAMES JEFFERSON [US]; J) 26 April 2007 (2007-04-26) page 1 - page 8	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2007/001527

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2006097784	A	21-09-2006	NONE	
WO 2007047859	A	26-04-2007	NONE	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2007/001527

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. type of material
 - ☒ a sequence listing
 - ☐ table(s) related to the sequence listing
 - b. format of material
 - ☒ on paper
 - ☒ in electronic form
 - c. time of filing/furnishing
 - ☒ contained in the international application as filed
 - ☒ filed together with the international application in electronic form
 - ☐ furnished subsequently to this Authority for the purpose of search
2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments: